

A handbook of useful drugs : a selected list of important drugs suggested for the use of teachers of materia medica and therapeutics and to serve as a basis for the examination in therapeutics by state medical examining and licensing boards prepared under the direction and supervision of the Council on Pharmacy and Chemistry of the American Medical Association.

Contributors

Council on Drugs (American Medical Association)

Publication/Creation

Chicago : American Medical Association, 1913.

Persistent URL

<https://wellcomecollection.org/works/w6qr9vqs>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



278
AMERICAN MEDICAL
ASSOCIATION.



22400016936

Med
K13524

42a1

52,000



Digitized by the Internet Archive
in 2016

<https://archive.org/details/b28068245>

A HANDBOOK OF USEFUL DRUGS

A Selected List of Important Drugs Suggested for the Use of
Teachers of Materia Medica and Therapeutics and to
Serve as a Basis for the Examination in Therapeutics
by State Medical Examining and Licensing Boards

PREPARED UNDER THE DIRECTION AND
SUPERVISION OF THE

COUNCIL ON PHARMACY AND
CHEMISTRY

OF THE

AMERICAN MEDICAL
ASSOCIATION

PRESS OF THE AMERICAN MEDICAL ASSOCIATION
535 North Dearborn Street, Chicago
1913

[AUTHORITY TO USE FOR COMMENT THE PHARMACOPEIA OF THE UNITED STATES OF AMERICA, EIGHTH DECENNIAL REVISION, IN THIS VOLUME, HAS BEEN GRANTED BY THE BOARD OF TRUSTEES OF THE UNITED STATES PHARMACOPEIAL CONVENTION, WHICH BOARD OF TRUSTEES IS IN NO WAY RESPONSIBLE FOR THE ACCURACY OF ANY TRANSLATION OF THE OFFICIAL WEIGHTS AND MEASURES, OR FOR ANY STATEMENT AS TO THE STRENGTH OF OFFICIAL PREPARATIONS.]

WELLCOME INSTITUTE LIBRARY	
Coll.	welMOMec
Call	
No.	QV



14796 309

PREFACE

It is generally recognized that a considerable proportion of the articles in the Pharmacopeia of the United States and in the National Formulary are worthless or superfluous. Repeated efforts have been made to eliminate at least the more objectionable of these articles. These efforts, however, have failed because they have uniformly encountered the objection that the articles or preparations are used by some physicians and therefore should be recognized and authoritatively defined.

The Council on Medical Education and the Confederation of State Examining and Licensing Boards have been trying to restrict instruction and examination in materia medica to the more important drugs. These efforts have suggested the desirability of selecting a fundamental list of drugs with which all medical students and practitioners might be expected to be familiar and to which, therefore, state examining and licensing boards might largely or entirely confine their examinations in materia medica.

HOW THE LIST WAS PREPARED

A committee of the Council on Medical Education of the American Medical Association prepared a list of the more important medicaments which was submitted to the members of the National Confederation of State Medical Examining and Licensing Boards. The confederation endorsed the principle and appointed a committee to compile a list which would answer the needs of the state examining and licensing boards.

In pursuance of its fundamental aims to secure the development of a more scientific and rational system of therapeutics, the Council on Pharmacy and Chemistry took up this matter. Largely basing its selection on the conclusions of the committee of the Confederation of State Licensing and Examining Boards, it compiled a preliminary list which it submitted for general discussion. This list was sent to teachers of pharmacology and therapeutics, to deans of medical

schools, to the secretaries and members of state medical examining and licensing boards, and to others presumed to be interested in the subject, with a request for criticism and suggestions. The replies received were compiled and analyzed and a revised list prepared and again submitted. The list was further considered and revised by the Council on Pharmacy and Chemistry and was then published in a preliminary form under the title "Useful Remedies." The object of presenting the book in a tentative form was to obtain further opinions regarding the list and especially suggestions for making the permanent book of the most practical value to all concerned.

The present book has been prepared on the basis of information, advice and suggestions brought out by the preliminary work above referred to. It presents a brief but practical discussion, from the modern viewpoint, of the properties, pharmacologic action, therapeutic uses and dosage of the drugs in the list. As it contains products the value of which is generally recognized it is hoped that the book may serve as a text on which teachers of materia medica and therapeutics may base their instruction, and state examining boards their examinations.

It is confidently predicted that an intelligent and critical use of these selected drugs will prove their general sufficiency and show that many drugs now discussed in text-books are superfluous and that many newly discovered or widely exploited proprietary preparations have no advantages over those contained in this book.

In discussing pharmacologic action the endeavor has been to present the essentials, giving the details of physiologic action only when they have an evident bearing on the therapeutic uses. The latter have been given concisely, but it is hoped in sufficient detail so that no important uses have been overlooked. In dosage the average doses of the pharmacopeia have been given except in those instances in which the importance of the drug makes it necessary to enter more into detail.

Especial attention has been paid to the various methods of administration and it is hoped that the occasional suggestions for the choice of vehicle will be of service to beginners in prescription writing. For the same reason the strength of local applications has been carefully indicated. In giving apothecary and metric doses the attempt to give exact equivalents has not been made. So far as possible, the use of round numbers has been encouraged.

In spelling the book conforms to the style of THE JOURNAL of the American Medical Association, except that the official names of the U. S. Pharmacopeia have been retained in the titles.

ABBREVIATIONS

The following abbreviations occur in the text:

U. S. P.—The Pharmacopeia of the United States of America, Eighth Revision.

N. F.—The National Formulary of Unofficial Preparations, Third Edition.

N. N. R.—New and Nonofficial Remedies, 1913.

STATEMENT OF SOLUBILITY

For ease of reference the solubility of official articles is indicated in approximate terms in accordance with the following equivalents:

Substances that are soluble in less than

1 part of solvent = very soluble.

From 1 to 10 parts of solvent = freely soluble.

From 10 to 100 parts of solvent = soluble.

From 100 to 1,000 parts of solvent = slightly soluble.

From 1,000 to 10,000 parts of solvent = very slightly soluble.

From 10,000 to 100,000 parts of solvent = nearly insoluble.

More than 100,000 parts of solvent = practically insoluble.

USEFUL DRUGS

Acacia.—Acacia, U. S. P., Gum Arabic.

A gummy exudation from *Acacia senegal* and other species of acacia.

PROPERTIES: Acacia occurs in colorless or pale yellowish, opaque, brittle, inodorous tears or fragments which are completely soluble in water but practically insoluble in alcohol.

Mucilago Acaciae.—Mucilage of Acacia, U. S. P.

A 34 per cent. solution of Acacia in a mixture of water and lime-water.

ACTION AND USES: Acacia and its mucilage are used as demulcents and suspending agents in the making of emulsions and mixtures.

Acetanilidum.—Acetanilid, U. S. P., $C_6H_5NH(CH_3CO)$.

Acetanilid is the monacetyl derivative of anilin, $C_6H_5NH_2$.

PROPERTIES: Acetanilid is an odorless, crystalline powder, having a slightly burning taste. It is only slightly soluble in water (1:180), but freely soluble in alcohol (1:2.5).

INCOMPATIBILITIES: Acetanilid is incompatible with spirit of nitrous ether. It forms a semiliquid mass when triturated with chloral or antipyrin.

ACTION AND USES: Acetanilid is analgesic, antipyretic and, in excessive doses, a cardiac depressant. These effects are probably due to para-aminophenol, into which it is converted in the body. The pulse is at first quickened and later slowed by a direct action on the heart muscle. Moderate doses have little effect on the temperature of normal animals and men, but such doses cause a marked reduction of the temperature in fever. Large doses, or small doses taken habitually, convert hemoglobin into met-hemoglobin and may destroy the red blood-corpuscles. In poisonous doses acetanilid produces cyanosis, abnormal reduction of temperature, coldness of the extremities and profuse sweating. In individuals with an idiosyncrasy toward the drug similar symptoms may be produced by small doses. Its use should be avoided in patients who are debilitated from any cause.

Acetanilid is effective for the relief of headache and neuralgic pain, but is not suited to the treatment of pain caused by inflammation.

DOSAGE: 0.20 gm. or 3 grains. It is well to begin with 0.10 gm. or about 2 grains and to repeat cautiously. Formerly mixtures of acetanilid with caffein or ammonium

salts were advised on the supposition that the cardiac depression would thus be avoided, but this does not seem to be the case. Investigation has shown that acetanilid is rendered somewhat more toxic by caffein, but sodium bicarbonate renders it less poisonous. The drug should be used cautiously and only for definite indications. Acetanilid has been widely exploited in the form of varying mixtures under different names as a cure for all pain. Many so-called headache powders contain it, but its indiscriminate use in this way is dangerous. It may be administered dry in the form of powders, cachets or capsules; because of its slight solubility it should not be massed in pills or compressed into tablets unless the tablet is crushed with the teeth before swallowing or unless the tablet will disintegrate rapidly in the stomach.

Acetphenetidinum. — Acetphenetidin, U. S. P., Phenacetin, $C_6H_4(OC_2H_5).NH(CH_3CO)$.

Acetphenetidin differs from acetanilid in containing the ethoxyl group C_2H_5O .

PROPERTIES: Acetphenetidin occurs as white, crystalline scales or a crystalline powder. It is odorless and tasteless. It is only slightly soluble in water (1:925) but soluble in alcohol (1:12).

INCOMPATIBILITIES: The same as for acetanilid.

ACTION AND USES: These are similar to those of acetanilid, but it is supposed to be somewhat safer. Its analgesic, antipyretic and cardiac depressant effects, like those of acetanilid, are due to the formation of para-aminophenol, and its possible advantage over acetanilid is probably due to the fact that this decomposition occurs more slowly. It is best administered in the form of powders, cachets or capsules.

Since the enactment of the Food and Drugs Act, June 30, 1906, acetphenetidin has frequently displaced acetanilid as the active agent in proprietary mixtures for the relief of headache and other pain. Its relation to acetanilid suggests similar caution in its use.

DOSAGE: A full dose is 0.50 gm. or $7\frac{1}{2}$ grains. It is well to begin with 0.30 gm. or 5 grains, and repeat every three hours if needed for a few doses. When small doses fail to relieve headache, larger doses are also usually ineffective.

Acidum Aceticum.—Acetic Acid, U. S. P.

A solution containing 36 per cent. by weight of absolute acetic acid ($H.C_2H_3O_2$) = CH_3COOH . (In some European countries a preparation comparable to glacial acetic acid, U. S. P., containing 99 per cent. of absolute acetic acid, is known as acetic acid, and the article that is official in the U. S. P. as "diluted acetic acid" is sometimes described as acetum or vinegar.)

Diluted Acetic Acid, U. S. P., contains 6 per cent. of absolute acetic acid.

PROPERTIES: Acetic acid is a clear, colorless solution. In all of its forms it is freely miscible with water.

ACTION AND USES: Acetic acid, as such, is not ordinarily used internally, but when administered in the diluted form it is mildly diaphoretic and diuretic. Externally it is a caustic, rubefacient and parasiticide.

Acidum Acetylsalicylicum.—See under Aspirin.

Acidum Benzoicum.—Benzoic Acid, U. S. P.

An organic acid, $(\text{HC}_7\text{H}_5\text{O}_2) = \text{C}_6\text{H}_5\text{COOH}$, obtained from benzoin, by sublimation, or prepared artificially.

PROPERTIES: Benzoic acid occurs as colorless, or almost colorless, lustrous scales or needles having an odor resembling benzoin, when obtained from the latter, and a pungent, acid taste. It is only slightly soluble in water (1:281), but is soluble in alcohol (1:1.8). It reacts with alkali hydroxids, and carbonates to form water-soluble benzoates.

ACTION AND USES: Benzoic acid is a mild antiseptic and diuretic. It is excreted in the urine in the form of hippuric acid (benzoyl glyccoll). Clinically it is of little value.

DOSAGE: 0.5 gm. or $7\frac{1}{2}$ grains. It is preferably dispensed in the form of powder and may be enclosed, dry, in capsules or cachets. It is more frequently used in the form of soluble compounds (see Sodii Benzoas).

Acidum Boricum.—Boric Acid, U. S. P. Boracic Acid (obsolete) $(\text{H}_3\text{BO}_3) = \text{B}(\text{OH})_3$.

PROPERTIES: Boric acid forms transparent, colorless scales or a light, unctuous, very fine powder. It is odorless, has a faintly bitter taste and is slowly soluble in water (1:18) and soluble in alcohol (1:15).

ACTION AND USES: Boric acid is a mild antiseptic. It has been occasionally administered internally, but with little effect, in cystitis. Externally it is frequently used as a dusting powder, either alone or combined with diluents such as starch or talcum, or with active substances such as acetanilid, salicylic acid or iodoform. It is also widely used as a wash or lotion, especially for catarrh of the mucous membranes, cystitis, conjunctivitis, pharyngitis, etc., usually in simple aqueous solutions containing from 2 to 4 per cent. of boric acid. This is one of the most frequently used lotions for conjunctivitis. It is also very useful for irrigating the bladder in cystitis. It is used in the form of glycerite of boroglycerin in washes and injections. The ointment is mildly antiseptic but is chiefly used as a protective dressing.

Glyceritum Boroglycerini.—Glycerite of Boroglycerin, U. S. P.
A glycerin solution representing 30 per cent. of boric acid.

Unguentum Acidi Borici.—Ointment of Boric Acid, U. S. P.

A 10 per cent. mixture of boric acid with paraffin and white petrolatum.

Acidum Citricum.—Citric Acid, U. S. P.

A tribasic organic acid, $\text{H}_3\text{C}_6\text{H}_5\text{O}_7 \cdot \text{H}_2\text{O}$, usually prepared from the juice of limes or lemons.

PROPERTIES: Citric acid forms colorless, transparent crystals; odorless and having an agreeable purely acid taste. It is very soluble in water (1:0.54) and freely soluble in alcohol (1:1.5).

ACTION AND USES: Citric acid may be used as a substitute for lemon-juice. Like the other organic acids, it is oxidized in the system to carbonic acid, so that its acid effects are much reduced. If an alkaline carbonate be added an effervescing mixture is obtained which gives the therapeutic effects of the alkalies.

DOSAGE: 0.5 gm. or $7\frac{1}{2}$ grains. It may be prescribed in the form of syrupus acidi citrici or as lemonade.

Acidum Diaethylbarbituricum.—See under Veronal.

Acidum Hydrochloricum.—Hydrochloric Acid, U. S. P., Muriatic Acid.

A fuming corrosive liquid containing 31.9 per cent. of hydrogen chlorid, HCl.

Acidum Hydrochloricum Dilutum.—Diluted Hydrochloric Acid, U. S. P.

A solution containing 10 per cent. of hydrogen chlorid, HCl.

PROPERTIES: Diluted hydrochloric acid is a colorless, odorless, strongly acid aqueous solution; freely miscible in all proportions with water or alcohol.

INCOMPATIBILITIES: It is incompatible with alkalies, carbonates and oxids, with which it reacts to form chlorids, and with the soluble salts of silver and of lead, forming insoluble silver chlorid and lead chlorid.

ACTION AND USES: Hydrochloric acid is the acid of the gastric juice, the average normal stomach contents containing approximately 0.2 per cent. It is necessary to the digestive action of pepsin and acts as an antiseptic in the stomach. By checking fermentation and putrefaction in the stomach it tends to prevent these processes in the intestine.

Diluted hydrochloric acid is used for the treatment of diseases of the stomach characterized by a deficiency of acid in the gastric juice on the theory that it replaces the acid lacking in the secretion. To restore the acidity of the stomach contents to the normal average would require much larger doses than are commonly given. It seems probable, therefore, that the acid as ordinarily given acts mainly as an appetizer and tonic. The utility of hydrochloric acid in achylia gastrica is more manifest in the nervous forms

and in the earlier stages of the organic variety. In some cases it causes distress and should be discontinued. There is some evidence to show that the continued administration of the acid is capable of increasing the gastric secretion. Hydrochloric acid also exerts a favorable influence on the secretion of the pancreatic and intestinal juices.

Hydrochloric acid is also of service in intestinal putrefaction, especially when the gastric digestion is impaired. It is indicated in achylia gastrica for the diarrhea caused by the irritant action of undigested meat and the putrefaction of proteins which have escaped gastric digestion.

DOSAGE: 1 c.c. or 15 minims in about half a glass of water. It should be given after meals and the dose repeated at the end of an hour. Five drops in a wineglassful of water after meals are often sufficient.

Acidum Hydrocyanicum Dilutum.—Diluted Hydrocyanic Acid, U. S. P.

This preparation in all the leading pharmacopeias of the world, is now uniformly directed to contain 2 per cent. of hydrogen cyanid, HCN.

PROPERTIES: It occurs as a colorless liquid of a characteristic odor resembling that of bitter almonds. On account of its poisonous character it should be tasted only with great caution. Hydrocyanic acid is freely miscible with water and with alcohol.

INCOMPATIBILITIES: It gradually decomposes on exposure to light and is incompatible with salts of iron, silver and mercury.

ACTION AND USES: Hydrocyanic acid is said to be sedative in small doses and has been widely used as an addition to cough mixtures, and as an anti-emetic. On account of its poisonous properties and the readiness with which it decomposes, it is no longer used to the same extent as formerly.

DOSAGE: 0.1 c.c. or 1.5 minims (not drops).

Acidum Nitricum.—Nitric Acid, U. S. P.

A liquid containing about 68 per cent., by weight, of hydrogen nitrate, HNO_3 .

PROPERTIES: Nitric acid is colorless, fuming, very caustic and corrosive and has a peculiar, somewhat suffocating odor. It is miscible with water in all proportions, dissolves mercury and most other metals with evolution of red fumes and stains woolen fabrics and animal tissues a bright yellow.

INCOMPATIBILITIES: Like other inorganic acids, it is incompatible with alkalies, the alkali carbonates, many of the salts of organic acids, and, because of its oxidizing properties, with all readily oxidizable substances. Its addition to organic liquids (alcohol, etc.), is apt to give rise to explosive reactions.

ACTION AND USES: Nitric acid is a powerful caustic, used for removing warts and small nevi and for cauterizing

chaneroids and other sores and bites of rabid animals, but its action is very painful and not readily controlled. When nitric acid is used as a caustic, the surrounding healthy tissue should be coated with petrolatum and the acid applied on the end of a rod of glass or wood.

Acidum Salicylicum.—Salicylic Acid, U. S. P.

An organic acid, $\text{HC}_7\text{H}_5\text{O}_3 = \text{C}_6\text{H}_4.\text{OH}.\text{COOH}$, generally prepared synthetically from phenol.

PROPERTIES: It occurs as fine, white needles or a bulky, white crystalline, odorless powder, possessing a sweetish, subsequently acid taste. Salicylic acid is only slightly soluble in water but freely soluble in alcohol or in ether. Salicylic acid reacts with alkali hydroxids and carbonates to form water-soluble salts. With solution of ferric chlorid it gives a deep purple color.

INCOMPATIBILITIES: It is incompatible with salts of iron and with spirit of nitrous ether.

ACTION AND USES: Salicylic acid is an antiseptic. It is quite irritant to mucous membranes and somewhat corrosive. Internally it has the actions described under sodium salicylate, in which form it is commonly employed. Externally it has been used as an application in pruritus, urticaria, bromidrosis and in some forms of eczema; also in the form of ointments and collodions to soften and remove corns and warts.

DOSAGE: Internally it is best given in the form of soluble salicylates. (See Sodium Salicylate.) Externally it is applied as an astringent in from 1 to 2 per cent. alcoholic solution or ointment; as an antiseptic, antiparasitic and keratolytic agent, in 2 to 5 per cent. dusting-powder, or ointment, and as a strong keratolytic in proportions up to 20 per cent., best dissolved in collodion. Continuous application to the skin may occasion slight corrosion.

Acidum Tannicum.—Tannic Acid, U. S. P., Tannin.

An organic acid, $\text{HC}_{14}\text{H}_9\text{O}_9$, obtained from nutgall.

PROPERTIES: Tannic acid occurs as a light yellowish, amorphous powder, gradually turning darker when exposed to air and light. It has a faint, characteristic odor and a strongly astringent taste. Tannic acid is very soluble in water, alcohol and glycerin.

INCOMPATIBILITIES: It is incompatible with alkalies, alkaloids, salts of iron and of most other metals, albumin and gelatin. With all these substances it reacts to form insoluble compounds.

ACTION AND USES: Tannic acid is used as an astringent and hemostatic. Internally it has been chiefly employed in the treatment of diarrhea. It is but little used with infants or children. It should be employed not as the principal curative agent, but as an occasional adjunct to proper dietetic and physical remedies when the discharges are unduly profuse. Its astringent action, however, often causes nausea and vomiting, and hence some one of the protein

combinations is better for action on the intestine. Local applications of tannic acid are frequently made to inflamed mucous membranes, especially in pharyngitis. It is frequently employed as a local application in the treatment of hemorrhoids in the form of a 20 per cent. ointment or as a suppository containing 0.3 gm. or 5 grains.

DOSAGE: 0.3 gm. or 5 grains.

Glyceritum Acidi Tannici.—Glycerite of Tannic Acid, U. S. P.

A 20 per cent. solution of tannic acid in glycerin.

DOSAGE: 1 c.c. or 15 minims corresponding to 0.2 gm. or 3 grains of tannic acid. This preparation is chiefly used externally and affords a convenient agent for making dilute solutions for local use. For local applications solutions containing from 0.5 to 2 per cent. of tannic acid are appropriate.

To prevent the astringent action of tannic acid on the gastric mucosa, various organic combinations of tannic acid have been prepared which resist the action of the gastric juice to a greater or less extent and escaping solution in the stomach, do not become active until they reach the intestine.

Tannalbin.—Tannalbin, N. N. R., Albumin Tannate.

It is insoluble in gastric juice and becomes effective when it reaches the intestine.

DOSAGE: 2 gm. or 30 grains.

Aconitum.—Aconite, U. S. P.

The tuberous root of *Aconitum napellus* L.

ACTION AND USES: Aconite slows the pulse and thus lowers the blood-pressure. The nervous system is first stimulated and then depressed. Death occurs from respiratory paralysis. Locally applied, aconite causes paralysis of the sensory nerve endings.

Aconite is used internally as a cardiac depressant, antipyretic and diaphoretic, especially in sthenic fevers of short duration or in the initial stage only of other diseases. It should be avoided in fevers accompanied by cardiac weakness. The danger from the depressing action of aconite on the heart makes it an undesirable agent to use as a cardiac depressant. It is being used less and less.

Tinctura Aconiti.—Tincture of Aconite, U. S. P.

One hundred c.c. represent 10 gm. of the drug in approximately 65 per cent. of alcohol.

Externally it is useful as a local application for the relief of neuralgic pain.

DOSAGE: 0.2 c.c. or 3 minims should be given hourly until the desired effect on the pulse is secured. Locally the tincture may be applied over the affected nerve, or a

liniment containing 10 per cent. of the tincture may be prescribed. Application to mucous membranes should be carefully avoided. The local application of the tincture is dangerous, as too much absorption may occur.

Adeps.—Lard, U. S. P.

True fats, chiefly those of animal origin, form an important class of ointment bases. They are especially valuable when penetrating ointments are desired for softening the skin for incisions, etc. Lard is the chief fat thus used. Like all animal fats, it is very liable to become rancid. To mitigate its odor and to hinder the occurrence of rancidity the following preparation is used:

Adeps Benzoinatus.—Benzoinated Lard, U. S. P.

Benzoinated lard is made by incorporating with lard 2 per cent. of benzoin and straining.

Adeps Lanae Hydrosus.—Hydrous Wool Fat, U. S. P., Lanolin.

This preparation is made by mixing the purified fat of sheep's wool with water in a proportion not to exceed 3 parts of the latter to 7 parts of the former.

Lanolin is used as an ointment base. It has the advantage that it does not become rancid, but its odor and consistence are disagreeable. The old statements that it is more readily absorbed than other ointments has not been borne out by experience.

Adnephtrin, Adrenalin.—See Epinephrin.

Aether.—Ether, U. S. P.

A liquid composed of about 96 per cent., by weight, of absolute ether, $(C_2H_5)_2O$, and about 4 per cent. of alcohol, C_2H_5OH .

PROPERTIES: It is volatile, inflammable, and the vapors, which are about two and one half times as heavy as air, are dangerously explosive when mixed with air. Ether is soluble in water (1:10), readily soluble in alcohol, and when administered internally is usually directed to be dispensed in the form of an alcoholic solution.

ACTION AND USES: Ether is used mainly by inhalation for the production of anesthesia. It depresses all parts of the central nervous system, causing loss of sensation, loss of consciousness and abolition of the reflexes. The vital centers of the medulla are involved very late in the poisoning, a fact which enhances the safety of this anesthetic. The respiration is affected first. Later there is depression of the vasomotor center and consequent fall of blood-pressure. Ether does not produce a marked effect on the heart, but its first action is a moderate reflex stimulation, while in poisonous doses it depresses the heart. In the administration of ether as an anesthetic at night, caution should be exercised to have the ether at a distance and, if possible, below any fire or flame, to avoid setting fire to the heavy

inflammable vapors. For anesthesia a pure ether, preferably anhydrous, should always be used. Administered internally it is an anodyne, sedative, carminative and antispasmodic.

DOSAGE: 1 c.c. or 15 minims.

Spiritus Aetheris, U. S. P.

A 32.5 per cent. alcoholic solution.

DOSAGE: 4 c.c. or 1 fluidram well diluted or on cracked ice.

Spiritus Aetheris Compositus, U. S. P.

One hundred c.c. contain ether, 32.5 c.c.; alcohol, 65 c.c., and ethereal oil, 2.5 c.c.

DOSAGE: 4 c.c. or 1 fluidram, best given on a lump of sugar. On account of its transient action the dose may be repeated once in half an hour if needed.

Aether Nitrosus.—Used only in the form of:

Spiritus Aetheris Nitrosi.—Spirit of Nitrous Ether, U. S. P., Sweet Spirits of Niter.

A solution of not less than 4 per cent. of ethyl nitrite, $C_2H_5NO_2$, in alcohol.

PROPERTIES: A pale yellowish liquid having a fragrant ethereal and pungent odor and a sharp burning taste. It is miscible with alcohol or water. It rapidly deteriorates.

INCOMPATIBILITIES: It is incompatible with acetanilid, antipyrin, potassium iodid and sodium salicylate.

ACTION AND USES: Spirit of nitrous ether is popularly used as a weak diuretic and diaphoretic. Its efficiency is doubtful. It is frequently prescribed in mixtures with other diaphoretics, notably the solution of ammonium acetate.

DOSAGE: 2 c.c. or 30 minims or half a teaspoonful.

Aethylis Chloridum.—Ethyl Chlorid, U. S. P.

PROPERTIES: It is a colorless and very volatile liquid having an agreeable odor and a sweetish, burning taste. It should be preserved in hermetically sealed glass tubes and kept in a cool place remote from light or fire.

ACTION AND USES: Ethyl chlorid is widely used in the form of spray to produce a local anesthesia for minor operations. When inhaled it produces prompt anesthesia, suitable for very short operations, but even then not without danger of producing accidents similar to those of chloroform. Because of these dangers and the difficulty of handling, it is now rarely used for general anesthesia.

Aethyl-Morphinae Hydrochloridum.—Ethyl-Morphin Hydrochlorid, N. N. R., Dionin.

PROPERTIES: Ethyl-morphin is an artificial base obtained by the action of ethyl iodid on morphin in the presence of an alkali. It occurs as a white, microscopically crystalline powder, odorless and having only a slight bitter taste. Ethyl-morphin hydro-

chlorid is freely soluble in water and in alcohol, but practically insoluble in ether and in chloroform.

ACTION AND USES: When administered internally, its action is intermediate between those of morphin and codein, but it is claimed that it does not produce constipation, nausea or lassitude and that it promotes healing after operations and injuries. It is the conclusion of careful observers, however, that, for internal use, it possesses no advantage over codein.

When applied to the eye this drug causes a local vasodilation, terminating in acute conjunctival edema. The chemosis thus produced is employed for its analgesic and curative effects in conjunctivitis, corneal ulcer, acute glaucoma, iritis, scleritis and other inflammatory diseases of the uveal tract. The greater the edema of the conjunctiva, the more decided is its analgesic action.

DOSAGE: 0.15 gm. or $\frac{1}{4}$ grain. Externally it is commonly employed in a collyrium in strength varying from 5 to 10 per cent. The strength may be increased to 20 per cent., and it is sometimes used in powder form. The ophthalmologist should make the first application and determine the minimum strength of solution which will produce the necessary chemosis. This dose should not be increased until it loses its effect. The action of the remedy may be regarded as sufficient as long as its application is followed by chemosis, redness and burning sensations for from one to two minutes afterward. Under these circumstances it may be instilled once a day. The patient should be instructed that the swelling of the conjunctiva is necessary to the therapeutic action of the remedy and that no harm to the eye will ensue from its use.

In cases of corneal opacity ethyl-morphin hydrochlorid has been applied to the eye in the form of powder. It may also be used as an ointment in strength varying from 1.5 to 5 per cent.

Alcohol.—Alcohol, U. S. P.

In European pharmacopeias it is usually designated as Spiritus, and varies considerably in strength.

PROPERTIES: The official U. S. P. alcohol is a colorless volatile liquid containing about 94.9 per cent. by volume of absolute ethyl alcohol, C_2H_5OH , and 5.1 per cent. by volume of water. It has a characteristic odor and burning taste and is miscible in all proportions with water, ether or chloroform. In addition to alcohol, the U. S. P. also describes absolute alcohol, used as a laboratory reagent, etc., and also diluted alcohol (approximately 50 per cent.) used in pharmacy as a menstruum.

ACTION AND USES: *Externally*, alcohol is a rubefacient and astringent, and by its evaporation, a refrigerant. It is used to harden and cleanse the skin; as a mild counter-irritant, (soap liniment), etc. In the concentration of 70 per cent. it is markedly antiseptic and is employed in surgery especially as Tincture of Green Soap, to cleanse the skin of patient and operator.

Internally, it is a narcotic, excessive doses depressing and paralyzing the central nervous system. Small doses produce euphoria, stimulate respiration, moderately dilate the cutaneous and splanchnic vessels, and modify the circulation. It is burned in the body and thus serves to a restricted extent as a source of energy.

Alcohol is employed as a diffusible stimulant, diuretic, diaphoretic and hypnotic. In well-selected cases, especially in patients accustomed to its use, it may be very valuable; otherwise it is apt to do more harm than good. In practice it is usually administered in the form of whisky, brandy, wine or other alcohol-containing beverages. It is generally accepted, however, that the aromatic principles in these several articles are even more toxic than is alcohol itself, and these beverages are therefore more poisonous than equal amounts of alcohol diluted with water.

In pharmacy alcohol is used as a solvent and, for administering medicines, is largely used as a vehicle in the form of:

Elixir Aromaticum.—Aromatic Elixir, U. S. P.

An aromatic and sweetened liquid containing about 25 per cent. of alcohol, by volume.

Aloes.—Aloes, U. S. P.

The inspissated juice of various species of *Aloe* is included in all the pharmacopeias. It is used in its original form, as a watery extract or as a purified extract known as aloin.

PROPERTIES: Aloes differs considerably in color and appearance, but in all its forms it has a rather characteristic odor and a nauseous, very bitter taste. Aloes is partially soluble in water.

ACTION AND USES: Aloes belongs to the emodin group of cathartics acting on the large intestine. It is believed to cause pelvic congestion and to have an emmenagogue action. Its action is said to be enhanced by the addition of soap and iron. Its purgative action is slow, evacuation occurring only after some hours, the stools being soft, seldom watery.

Aloes is adapted to the treatment of constipation by daily laxative action, but should rarely be used as a purgative, because of its tendency to cause griping. Because of the pelvic congestion it produces, aloes is contra-indicated in pregnancy and in the presence of hemorrhoids. It is an appropriate remedy in functional amenorrhea if the production of pelvic congestion is deemed advantageous.

DOSAGE: The purgative dose of aloes is from 0.15 to 0.3 gm. or from 2 to 5 grains. For the treatment of chronic constipation smaller doses, 0.03 to 0.05 gm. or $\frac{1}{2}$ to 1 grain, should be used. A preparation of belladonna is usually combined with it.

Extractum Aloes.—Extract of Aloes, U. S. P.

DOSAGE: 0.10 gm. or 2 grains.

Aloinum.—Aloin, U. S. P.

DOSAGE: Purgative dose 0.05 gm. or 1 grain. In the treatment of chronic constipation aloin is frequently given in doses of from 0.005 to 0.02 gm. or 1/10 to 1/3 grain in combination with extract of belladonna and strychnin.

	gm.	or c.c.	
R Strychninae sulphatis	0015		gr. 1/40
Aloinae	02		gr. 1/3
Extracti belladonnae	006		gr. 1/10

Mix and make 1 pill or tablet.

Take 1 after supper or at bedtime; if at bedtime, with plenty of water and a cracker, or at least never on an empty stomach.

Alumen.—Alum, U. S. P.

Crystallized potassium aluminum sulphate, $\text{AlK}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$.

PROPERTIES: Alum occurs as colorless crystals or a white powder without odor, but having a strongly astringent taste. It is freely soluble in water, practically insoluble in alcohol. It contains about 45 per cent. of water of hydration, which can be removed by heat, the product being exsiccated alum.

INCOMPATIBILITIES: Alum is incompatible with alkalies and carbonates, which react with it with the formation of insoluble aluminum hydroxid. It is also incompatible with salts of lead because it precipitates the insoluble lead sulphate.

ACTION AND USES: Alum is astringent, styptic and hemostatic, but is seldom administered internally.

DOSAGE: Alum solution may be used as a gargle (from 1 to 5 per cent.) but it is somewhat injurious to the teeth; it may be given as an injection in gonorrhea (from 0.5 to 1 per cent.) and as a lotion in skin diseases (1 per cent.).

Alumen Exsiccatum.—Exsiccated Alum, U. S. P.

A powder representing about twice its weight of the crystallized alum; it is sometimes used externally as a dusting powder or in the form of ointments.

Alumini Acetas.—Aluminum Acetate.

Used principally in the form of:

Liquor Alumini Acetatis.—Solution of Aluminum Acetate, N. F.

PROPERTIES: A clear, colorless solution, containing from 7.5 to 8 per cent. of basic aluminum acetate, having an acetous odor and a sweetish, astringent taste.

ACTIONS AND USES: Solution of aluminum acetate is used extensively at the present time as a mild astringent and antiseptic. Because of the readiness with which this solution is decomposed it is advisable to dilute it with water only. Applied to the skin it should be diluted from four to nine times.

AMMONIUM COMPOUNDS

Ammonium compounds are divisible into two classes. In the first class the ammonium is combined with a strong acid such as hydrochloric or sulphuric. These form stable, neutral salts which act largely by their salt action. In the second class, the ammonium is combined with a weak acid radical such as hydroxyl or the radical of carbonates. These compounds are unstable, decomposing readily with liberation of ammonia. Such compounds produce reflex effects by the irritating action of the ammonia evolved. All ammonium compounds used in medicine are soluble in water and the carbonate and hydroxid have a strong alkaline reaction.

ACTION AND USES: *Ammonium salts*, when injected into the circulation, stimulate the central nervous system, but they are so rapidly excreted or converted into urea that they cannot produce systemic action when taken by mouth, even though they are absorbed very readily. Their effects are, therefore, entirely local. Ammonium acetate was believed to be diaphoretic, but probably has little value.

The neutral salts are rather irritant and thus cause a mild stimulation of the mucous membranes, explaining the use of ammonium chlorid as an expectorant. With ammonium carbonate, this effect is reinforced by its alkaline reaction, through which it liquefies and dissolves mucus.

Ammonia (NH_3).

A gas which is very soluble in alcohol and water with formation to some extent of NH_4OH . Water of ammonia and the several preparations containing it are strongly alkaline.

INCOMPATIBILITIES: It is incompatible with acids, neutralizing them and forming the salts of ammonium. It is also incompatible with the soluble salts of many metals because it precipitates from these solutions the hydroxid of the metals. Thus ammonia water with solution of ferric chlorid produces an insoluble precipitate of ferric hydroxid. Solutions of ammonia are also incompatible with the salts of alkaloids from which they liberate the alkaloid. Thus ammonia water added to a solution of strychnin sulphate produces a precipitate of the insoluble alkaloid strychnin.

ACTIONS AND USES: Ammonia internally in the form of water of ammonia or of aromatic spirit of ammonia is stimulant, because the ammonia escaping irritates the mucous membranes of the nose and of the stomach and causes a reflex increase in the force of the heart and in the blood-pressure. Little, if any, of the gas is absorbed by the respiratory tract.

Externally, ammonia is used as a counterirritant. It reddens the skin but does not blister, unless applied in concentrated solution.

Aqua Ammoniae.—Ammonia Water, U. S. P.

An aqueous solution, containing about 10 per cent. by weight of ammonia (NH_3). On standing, this solution loses ammonia and hence should be kept in well-stoppered bottles and should frequently be tested by the pharmacist to insure its complying with the pharmacopeial requirements. Ammonia water is a colorless liquid, having a very pungent, characteristic odor and a caustic, soapy taste. It is freely miscible with water and alcohol.

Ammonia water is used chiefly for counterirritation in the form of:

Linimentum Ammoniae.—Ammonia Liniment, U. S. P.

A mixture of ammonia water 35 parts, alcohol 5 parts and a fatty oil 60 parts.

Ammonii Acetas.—Ammonium Acetate.

Used chiefly in the form of:

Liquor Ammonii Acetatis.—Solution of Ammonium Acetate, U. S. P.

An aqueous solution containing about 7 per cent. of ammonium acetate. It is made by dissolving ammonium carbonate in diluted acetic acid. It is a clear, colorless liquid, having a mildly saline, acidulous taste and an acid reaction. It is incompatible with alkaline hydroxids and carbonates, which liberate ammonia.

ACTIONS AND USES: Solution of ammonium acetate was formerly credited with diaphoretic and diuretic powers, but they are very weak. It is used in fevers and infections to promote sweating.

DOSAGE: 15 c.c. or 4 fluidrams, which can be repeated once in two or three hours.

Ammonii Carbonas.—Ammonium carbonate, U. S. P.

Ammonium carbonate consists of a mixture of ammonium bicarbonate and ammonium carbamate.

PROPERTIES: It occurs as white, hard masses, having a strong odor of ammonia and a sharp, saline taste. On exposure to the air the salt loses both ammonia and carbon dioxid. Ammonium carbonate is slowly but freely soluble in water, the ammonium carbamate being thereby converted into normal ammonium carbonate. Alcohol dissolves the carbamate and leaves the bicarbonate.

INCOMPATIBILITIES: Ammonium carbonate is incompatible with acids, which decompose it, forming salts of ammonium and evolving carbon dioxid (CO_2). It precipitates the carbonate or the hydroxid of most metals and the insoluble alkaloids from solutions of their salts.

ACTION AND USES: Ammonium carbonate is largely decomposed (hydrolyzed) when dissolved in water, and its solutions are irritant to mucous membranes from the action of the ammonia set free. It is used by inhalation or in

solutions as a reflex or diffusible stimulant in syncope, or arrest of respiration, and as a liquefying expectorant in bronchitis.

DOSAGE: 0.25 gm. or 4 grains, dissolved in sufficient water to avoid too great irritation, which may result in nausea and vomiting. On the other hand, as the action of the remedy depends on its irritating qualities, it should not be too greatly diluted.

Spiritus Ammoniae Aromaticus. — Aromatic Spirit of Ammonia, U. S. P.

A solution of ammonium carbonate with some free ammonia and aromatic oils in alcohol.

ACTION AND USES: The same as those of Ammonium Carbonate.

DOSAGE: From 1 to 5 c.c. or 15 to 60 minims, suitably diluted with water. As the stimulating action is of short duration a moderate dose may be repeated in from fifteen minutes to half an hour.

Ammonii Chloridum.—Ammonium Chlorid, U. S. P., NH_4Cl .

PROPERTIES: Ammonium chlorid usually occurs as a white, crystalline powder, without odor, having a cooling, saline taste. It is freely soluble in water (1:2), and soluble in alcohol (1:50), its aqueous solutions being neutral or very slightly acid to litmus.

INCOMPATIBILITIES: Ammonium chlorid is incompatible with alkaline hydroxids and carbonates, which liberate ammonia. It precipitates the insoluble chlorids of silver and of lead from solutions of the salts of those metals.

ACTION AND USES: Ammonium chlorid is said to be absorbed more quickly than any other salt, and in general has saline properties. Its principal activity is as an expectorant, though it is slightly diuretic and diaphoretic. Its vapors have been used for inhalation in cases of nasopharyngeal catarrh and as an expectorant in bronchitis. For this purpose it may also be generated in a special apparatus by the union of the vapors of strong hydrochloric acid and ammonia water.

DOSAGE: From 0.30 to 1 gm. (from 5 to 15 grains), repeated every two or three hours, or less frequently, depending on the size of the dose.

Its taste is best concealed or modified by administering it in a sour mixture as:

	gm. or c.c.	
R Ammonii chloridi	5	3 iss
Syrupi acidi citrici	50	or fl. ʒ ii
Aquae q.s.ad	100	fl. ʒ iv

M. et Sig.: A teaspoonful, in water, every two hours.

To the preceding prescription codein sulphate may be added, if desired. The amount of citric acid should be diminished when the prescription is for a child, and after the child has taken a dose of it he could be given a piece of chocolate or a simple peppermint or wintergreen lozenge.

NITRITES

The Nitrites used in medicine are certain salts or esters of nitrous acid and possess a common pharmacologic action. This group also includes certain organic nitrates which are reduced to nitrites in the organism. The chief members are amyl nitrite, sodium nitrite and glycerol nitrate (glyceryl trinitrate or nitroglycerin).

The characteristic action of this group is vasodilatation with a fall of the blood-pressure. The members differ chiefly in the rapidity and duration of their effects, amyl nitrite being the quickest, though its action is of correspondingly short duration.

Amylis Nitris.—Amyl Nitrite, U. S. P.

PROPERTIES: A liquid containing about 80 per cent. of amyl (chiefly iso-amyl) nitrite. It should be kept in hermetically sealed glass bulbs or in dark, amber-colored, glass-stoppered vials, in a cool, dark place. Amyl nitrite is a clear, yellowish liquid of a peculiar, ethereal, fruity odor and a pungent, aromatic taste. It is very volatile even at low temperatures, and is inflammable. It is practically insoluble in water, but miscible in all proportions with alcohol or ether.

ACTION AND USES: When given by inhalation, amyl nitrite produces an almost instantaneous dilatation of the peripheral blood-vessels, shown by redness of the skin beginning in the head and neck, rapidly spreading over the body and sometimes extending to the lower extremities. This is promptly followed by a dilatation of the splanchnic and other vessels so that the blood-pressure soon falls. The lowered pressure increases the heart-rate. There is a feeling of fulness in the head, often accompanied by headache. The breathing is rapid. With excessive doses unconsciousness may supervene, and convulsions may occasionally occur after toxic doses. Large doses kill by respiratory paralysis. It may produce methemoglobin in the blood and cause the excretion of sugar in the urine. As much as 0.6 c.c. may be inhaled without dangerous results.

The effects of the medicinal administration of amyl nitrite are very transient. It is employed to relieve spasm in epilepsy, sometimes averting an expected paroxysm. It is also used to relax the spasm of the blood-vessels in angina pectoris and in other painful affections in which there is reason to believe that the pain depends on arterial spasm.

Amyl nitrite is given to reduce the blood-pressure in cases in which hemorrhage is due to the rupture of a blood-vessel in the lungs, brain or other organ in consequence of elevated blood pressure, but in hemorrhage with normal blood-pressure it may do harm. It has been used with favorable results in bronchial asthma.

DOSAGE: 0.2 c.c. or 3 minims, by inhalation. It is conveniently carried in the form of glass pearls, each pearl containing a dose. When required the pearl is crushed in the handkerchief and the content inhaled.

Amylum.—Starch—Corn-Starch, U. S. P.

The starch grains obtained from the fruit of *Zea mays* occur in the form of powder or irregular, angular, white masses, insoluble in both water and alcohol, but swelling into a colloidal "paste" when boiled in water.

ACTION AND USES: Dry starch is used as a dusting and drying powder and also as a diluent for other more active substances. The mucilage produced on boiling, with water or glycerol is employed as an emollient and protective; as a cataplasm, and as an antidote to iodine poisoning. The starches from other cereals, from cassava and the potato have practically the same medicinal properties as corn-starch.

Antimonii et Potassii Tartras.—Antimony and Potassium Tartrate, U. S. P., Tartar Emetic.

Antimony and potassium tartrate is a double salt of antimony and potassium with the radical of tartaric acid.

PROPERTIES: It occurs as a white powder or as colorless transparent crystals, becoming white and opaque on exposure to the air. It is soluble in water, but practically insoluble in alcohol.

INCOMPATIBILITIES: It is incompatible with alkalies and their carbonates, tannic acid and astringent preparations generally.

ACTION AND USES: Antimony and Potassium Tartrate, when applied to the skin, gives rise slowly to inflammatory changes, with pustules and ulceration, which is somewhat difficult to limit. The ointment formerly employed as a pustulant counterirritant is now rarely used.

Internally it produces local irritation of the gastrointestinal tract, and thereby nausea and vomiting with marked prostration. If absorbed, symptoms very similar to those produced by poisonous doses of arsenic are observed.

The therapeutic uses of tartar emetic are almost entirely confined to the treatment of the first stage of acute laryngitis and bronchitis. It should be avoided in cases marked by depression. When it is given, the object should be to increase secretion and facilitate the expulsion of sputum. The administration of tartar emetic should not be carried beyond the production of slight nausea. For the production of vomiting other agents are preferable.

DOSAGE: As an expectorant small doses should be used, beginning with 0.001 gm. or 1/60 grain, which may be repeated hourly, taking care to avoid too great depression. The emetic dose is 0.03 gm. or 1/2 grain.

Vinum Antimonii.—Wine of antimony, U. S. P.

One hundred c.c. contain 0.4 gm. of antimony and potassium tartrate in white wine fortified by alcohol. This preparation is practically identical with the antimonial wine official in other countries.

DOSAGE: 1 c.c. or 15 minims, equivalent to 0.004 gm. or 1/15 grain of antimonium and potassium tartrate. As an expectorant smaller doses such as 0.2 c.c. or 3 minims should be employed.

Apomorphinae Hydrochloridum.—Apomorphine Hydrochlorid, U. S. P.

The hydrochlorid of an artificial alkaloid prepared from morphin by the abstraction of one molecule of water.

PROPERTIES: Apomorphin hydrochlorid occurs as minute, grayish-white prisms, having a slightly bitter taste and acquiring a greenish tint on exposure to light and air. It is soluble in both water and alcohol. If the salt imparts at once an emerald-green color to 100 parts of water it should be rejected. The amorphous form may contain dangerous impurities.

INCOMPATIBILITIES: It is precipitated by alkalies and the other alkaloidal reagents. Solutions decompose rather readily.

ACTION AND USES: The chief action of apomorphin is the production of vomiting with its usual accompanying symptoms, including nausea with increase of saliva and other secretions, depression of the circulation, sweating, etc. It sometimes produces respiratory paralysis, even in small doses.

The drug is used chiefly as an emetic. For this purpose it has the advantage that it acts on the vomiting center, independently of a local action on the stomach. It produces a prompt emptying of the stomach with little subsequent sickness, unless the dose is large. It is therefore a useful emetic in poisoning, if the stomach-tube cannot be employed. It has been advised for the expulsion of foreign bodies from the air-passages. As an expectorant it is inferior to other nauseants.

It is said to be sometimes useful in asthma. Small doses (0.002 gm., 1/30 grain) are hypnotic, especially in acute alcoholism.

DOSAGE: The emetic dose is 0.005 gm. or 1/10 grain, given preferably by hypodermic injection. This may be repeated at ten-minute intervals until effective, but it should be remembered that in some cases apomorphin produces toxic effects without causing vomiting, and a dose of 0.004 gm. or 1/15 grain is said to have produced death in a person enfeebled by chronic bronchitis; 0.012 or 1/5 grain may be given to robust patients at the first dose if the urgency of the case demands it, but care should be exercised in the use of these doses.

As an expectorant the proper dose is from 0.001 gm. to 0.002 gm. or from 1/60 to 1/30 grain, repeated once an hour or once in two hours, with the avoidance of more than slight nausea.

Antipyrina.—Antipyrine, U. S. P.

This substance is described in the British Pharmacopeia as phenazonum and in the German Pharmacopeia and some

other European pharmacopeias as pyrazolonum phenyldimethylicum.

PROPERTIES: Antipyrin occurs as a colorless, almost odorless, crystalline powder or tabular crystals having a slightly bitter taste. It is very soluble in water and freely soluble in alcohol.

INCOMPATIBILITIES: Antipyrin is incompatible with spirit of nitrous ether, or other nitrites, tannic acid and tannic-acid containing preparations. Mixed dry with sodium salicylate, it liquefies on standing.

ACTION AND USES: Antipyrin is an antipyretic and analgesic, acting similarly to acetanilid.

It is used for the relief of pain, chiefly when of a neuralgic character. It is not suited to the treatment of the pain caused by inflammation.

Antipyrin is now seldom used as an antipyretic. Locally, it is used sometimes as a hemostatic. It is a valuable antispasmodic for use with children for pertussis, bronchitis, etc. It is dangerous, however, in preparations sold directly to the laity.

DOSAGE: 0.25 gm. or 4 grains, given with even greater caution than acetanilid and acetphenetidin. It is best administered by itself in simple solution, or in powders, capsules or cachets.

Aqua.—Water, U. S. P.

Water is described by the Pharmacopeia as potable water in its purest obtainable state. Water used for medicines should be sterile; it is preferable to use:

Aqua Destillata.—Distilled Water, U. S. P.

Distilled water is described as a colorless, limpid liquid, without odor or taste, and perfectly neutral to litmus paper. For some purposes (in making solutions of salvarsan, for example), it is essential that it be freshly distilled. Ordinary distilled water frequently is not sterile.

AQUAE—WATERS

The official medicated waters are aqueous solutions of volatile substances. The aromatic waters are saturated solutions of volatile oils. They are used as vehicles for the more active, water-soluble drugs. For preparations included in this list see:

Aqua Ammoniae, under Ammonia.

Aqua Camphorae, under Camphora.

Aqua Chloroformi, under Chloroformum.

Aqua Cinnamomi, under Cinnamomum.

Aqua Destillata, under Aqua.

Aqua Menthae Piperitae, under Mentha Piperita.

Aqua Rosae, under Rosa.

Argenti Nitras.—Silver Nitrate, U. S. P., Lunar Caustic.

PROPERTIES: Silver nitrate, AgNO_3 , occurs in colorless, crystalline tables, becoming gray, or grayish black on exposure to light

in the presence of organic matter. It is odorless and has a bitter, caustic, and strongly metallic taste. It is very soluble in water and soluble in alcohol.

INCOMPATIBILITIES: Silver nitrate is incompatible with soluble chlorids, bromids and iodids, with which it forms the corresponding very insoluble salts of silver. It is also incompatible with soluble carbonates and hydroxids, which precipitate the oxid of silver, and with all organic drugs and reducing agents.

ACTION AND USES: Silver nitrate is an antiseptic and germicide, destroying many micro-organisms in solutions of a strength of 1 : 1,000 and preventing their growth in a strength of 1 : 10,000. Weak solutions are astringent to mucous membranes and strong solutions are caustic when applied to mucous membranes, denuded surfaces, and, in some cases, to the normal skin. When taken internally silver nitrate is believed to hinder the secretion of the gastric juice, especially when the degree of acidity is excessive.

Silver nitrate is used as a mild caustic to wounds, ulcers and exuberant granulations. It is applied as an astringent and antiseptic in catarrhal infections of the mucous membranes.

Internally it is sometimes employed as an astringent in diarrhea, but its use for this purpose is not to be recommended. It has been advised for the treatment of gastric ulcer and to reduce the gastric secretion in hyperchlorhydria. Its internal use for some time may be followed by its deposition in the skin, producing the condition known as argyria.

DOSAGE: As a caustic, silver nitrate is used in the form of fused silver nitrate. This should be moistened before use and should be held with forceps or in a suitable holder. Care should be taken to limit its action to the affected part.

In applications to mucous membranes the following strengths of solution in water are most suitable:

To the conjunctiva a strength of 4 per cent. may be applied in small quantity and the action stopped by addition of solution of sodium chlorid. This method of use is applicable to cases of severe conjunctivitis, especially gonorrheal.

For the prevention of gonorrheal conjunctivitis in the new-born a drop of 2 per cent. solution should be instilled into the eye as soon as practicable after delivery.

For other diseases of the conjunctiva, solutions varying in strength from 0.2 to 2 per cent. are sometimes used, but it is probable that other astringents are safer and quite as useful.

To the larynx, application is made of solutions containing from 2 to 10 per cent. of silver nitrate.

For the urethra, it is employed in the strength of from 1 : 10,000 to 1 : 2,000, and a solution of the strength of 1 : 5,000 may be injected into the bladder. A solution of

from 1:2,000 to 1:1,000 is used as a gastric douche in hyperchlorhydria and gastric ulcer, followed after two minutes by a solution of sodium chlorid which is thoroughly mixed with the silver solution and then washed out of the stomach.

Solutions of silver nitrate should always be made with distilled water, and the mucous membranes to which they are to be applied should receive a preliminary cleansing to remove mucus, pus, food, etc., which might interfere with its action. The action may be stopped by sodium chlorid solution.

Internally the salt may be given in pill form, preferably mixed with kaolin and massed with petrolatum. The average dose is 0.01 gm. or 1/5 grain. This dose may also be given in solution when the direct effect on the stomach is desired. If the stomach contains acid it would be converted into silver chlorid; hence it should be given on an empty stomach.

Argenti Nitras Fusus.—Molded Silver Nitrate, U. S. P.

A white, hard solid generally in the form of pencils or cones. A small amount of silver chlorid is added in the process for the purpose of toughening the mass.

Argenti Proteinus.—Silver Proteinate; see Protargol, N. N. R.

Aristol.—See Thymolis Iodidum.

Arseni Trioxidum.—Arsenic Trioxid, U. S. P., As_2O_3 . Formerly official as arsenous acid.

PROPERTIES: Arsenic trioxid occurs as an opaque, white powder or in irregular masses of two varieties, one amorphous, the other crystalline, opaque and colorless like glass. Arsenic trioxid is very slowly soluble in water and practically insoluble in alcohol (1:30 to 1:100). It dissolves quite readily in solutions of acids or alkalies.

INCOMPATIBILITIES: Solutions of arsenic are incompatible with salts of iron and of magnesium, lime-water and vegetable astringents.

ACTION AND USES: Local External Effects and Uses: Arsenic trioxid applied to denuded or ulcerated tissue has a mildly caustic action which is quite painful. It has been used as a caustic, especially to malignant growths, but the painful character of the applications, the danger of absorption, and the uncertain extent of the destructive action have justly limited its use.

Internal Actions: Arsenic trioxid irritates the mucous membrane of the stomach and intestines. Toxic doses cause nausea and vomiting and colicky pains in the abdomen. It produces a diarrhea of a watery character resembling that of cholera. In small doses it produces fatty degeneration of the liver and other internal organs; large doses produce great depression, and collapse may ensue.

Arsenic in therapeutic doses is employed in the treatment of neuralgia. It is thought to be especially adapted to

cases of a periodic character. It is also useful in the treatment of chorea. Larger doses, especially if long continued, may cause peripheral neuritis. Arsenic stimulates the action of the blood-forming organs, especially the bone-marrow. It is especially useful in the treatment of pernicious anemia, leukemia and Hodgkin's disease. In these diseases it must be used in as large doses as can be borne. While improvement occurs in these conditions under the use of arsenic it is usually only temporary.

Many skin-diseases are favorably influenced by proper doses of arsenic. It acts by stimulating the skin, in such cases as usually require external stimulating applications. It is also of service in lesions due to disturbances of innervation in which the skin is usually poorly nourished. The following skin affections may be mentioned as likely to be benefited by arsenic: psoriasis, lichen planus, chronic eczema, pemphigus, dermatitis herpetiformis, chronic urticaria and disturbances of the sweat function. On the other hand, acute inflammatory conditions of the skin are made worse by arsenic.

Arsenic is recommended in chronic bronchitis and emphysema. It has been used with apparent good results in tuberculosis. Too great dependence, however, should not be placed on it in the latter affection. It is used in asthma by way of the stomach and in the form of cigarettes.

Arsenic is a powerful remedy against protozoal affections. These include malaria, syphilis and relapsing fever. In the form of arsenic trioxid it has been used successfully in malaria, particularly in the chronic form and in malarial cachexia. See Salvarsan.

DOSAGE: For ordinary affections of metabolism or for a tonic influence the dose may vary from 0.001 to 0.002 gm. or 1/60 to 1/30 grain. In diseases of the blood the dosage should be regulated according to the effect, but it is well to use as large doses as the patient will tolerate. Arsenic may be used in increasing doses until symptoms of mild intoxication appear. One may begin with 0.005 gm. or 1/20 grain of arsenic trioxid three times daily, and increase by 0.001 gm. or 1/60 grain three times daily. In using Fowler's solution the initial dose may be 3 minims three times daily and increase by 1 minim three times daily. A slight toxic action is indicated by nausea, colicky pains or a puffiness under the eyes. The presence of albumin in the urine may also be observed. Such symptoms may make advisable the temporary withdrawal of the remedy.

Liquor Potassii Arsenitis.—Solution of Potassium Arsenite, U. S. P., Fowler's Solution.

An aqueous solution containing potassium arsenite, corresponding to 1 per cent. of arsenic trioxid.

DOSAGE: 0.2 c.c. or 3 minims.

This solution is often somewhat alkaline and is therefore incompatible with alkaloidal salts. This is avoided in the following:

Liquor Acidi Arsenosi.—Solution of Arsenous Acid., U. S. P.

An aqueous solution containing the equivalent of 1 per cent. of arsenic trioxid with 5 per cent. of hydrochloric acid.

DOSAGE: 0.2 c.c. or 3 minims.

Liquor Arseni et Hydrargyri Iodidi.—Solution of Arsenous and Mercuric Iodids, U. S. P.

Commonly designated as Donovan's solution; represents 1 per cent. of arsenous iodid and 1 per cent. of mercuric iodid.

DOSAGE: 0.1 c.c. or 1½ minims.

Asafoetida.—Asafetida, U. S. P.

A gum resin obtained from the roots of *Ferula foetida* and probably other species of *Ferula*.

PROPERTIES: A good quality of asafetida should contain not less than 50 per cent. of matter soluble in alcohol and should yield not more than 10 per cent. of ash. The gum of asafetida is freely soluble in cold or hot water and is present in sufficient amount to suspend in the form of a permanent emulsion the accompanying resin and volatile oil.

ACTION AND USES: Asafetida is used in the treatment of hysteria, acting probably by its odor. It is also carminative.

DOSAGE: 0.25 gm. or 4 grains—preferably in pill form. In tympanites an emulsion may be used as an enema. The emulsion is made by triturating 4 gm. of a good quality of the drug with 100 c.c. of water until a uniform emulsion results. The strength may be varied to meet individual requirements.

Aspidium.—Aspidium, U. S. P., Male Fern.

The dried rhizome of several species of *Dryopteris*; used only in the form of:

Oleoresina Aspidii.—Oleoresin of Aspidium, U. S. P.

An oleoresin prepared by extracting aspidium with acetone, evaporating and recovering the solvent.

PROPERTIES: The most important constituent of male fern is fillic acid.

ACTION AND USES: Ordinarily the active constituents of aspidium are not absorbed and produce no symptoms except some nausea. If absorption occurs, violent symptoms of poisoning may ensue. The symptoms are excessive vomiting and purging, great weakness, spasms in the extremities, convulsions, stupor deepening into coma and collapse. Disturbances of sight and hearing may occur and permanent blindness sometimes follows. Jaundice has been observed.

Oleoresin of aspidium is used as a teniacide against ordinary tapeworm, but it is said to be more efficient against the *Bothriocephalus* than against ordinary tenias.

DOSAGE: 2 gm. or 30 grains is given by the Pharmacopeia, but most authorities recommend a larger dose. The dose should be proportioned to the strength and health of the patient. Robust males can take as much as 8 gm. or 2 fluidrams, while women should be given smaller doses and special caution should be exercised in administering the drug to anemic or debilitated persons. Children of 4 years may take 4 gm. or 60 grains.

Before this remedy is given, the alimentary canal should be emptied by a light diet or fasting for twenty-four hours and the administration of a saline cathartic in the morning before the anthelmintic is given. The drug should be given early in the morning and is best directed to be taken in capsules containing 0.3 gm. or 5 grains, or in the form of emulsion. The total quantity may be given in divided doses at intervals of fifteen minutes and the last dose followed in three hours by a saline laxative. Castor oil or other fixed oils should not be given, because they favor the absorption of the active principle.

A dose of 8 gm. or 2 drams has been fatal to a child and 25 gm. or 6 drams have several times proved fatal to adults.

Aspirin.—Aspirin, N. N. R., Acetylsalicylic Acid, $C_6H_4O(CH_3CO)COOH$.

The acetic acid ester of salicylic acid.

PROPERTIES: Aspirin occurs as a crystalline, odorless powder with a faintly acidulous taste. It is slightly soluble in water (1:100) and freely soluble in alcohol.

ACTION AND USES: Aspirin acts like salicylic acid and the salicylates, but is claimed to be less disturbing to the stomach. It is used as an antipyretic, analgesic and anti-rheumatic. It is much used for the relief of headache and other painful affections.

DOSAGE: 0.3 to 1 gm. or 5 to 15 grains, repeated once in 3 hours until symptoms of salicylism (ringing in the ears, etc.) are noted.

Atoxyl.—See Sodii Arsanilas.

Atropina.—Atropine, U. S. P.

An alkaloid obtained from *Atropa belladonna* and from other solanaceous plants.

PROPERTIES: The alkaloid itself is only slightly soluble in water and is usually prescribed in the form of one of its salts, which are readily soluble.

INCOMPATIBILITIES: Solutions of atropin are stated to be incompatible with alkalies. While no precipitate of the alkaloid would form, unless the solution were more concentrated than those ordinarily used in medicine, the alkaloid in the presence of alkalies is likely to be decomposed.

Solutions of the alkaloid are also incompatible with tannic acid, with Lugol's solution, and with potassiomeric iodid, which precipitate compounds of the alkaloid, and with salts of mercury, such as the chlorid, which decomposes it.

ACTION AND USES: Atropin and the related alkaloids act on the sympathetic and especially on the autonomic nervous system. If the ordinary stimulation of these nerves causes motor or secretory activity, the use of atropin will inhibit it, and if the normal action of the nerve is inhibitory, atropin removes the inhibition.

The alkaloid produces a cerebral stimulation which shows itself by quickened thought and speech, eventuating in a peculiar talkative delirium with hallucinations. Larger doses produce unconsciousness and occasional convulsions.

Atropin checks the secretion of saliva so that the mouth and throat become dry. This dryness is due to some extent to a similar effect on the mucous secretions of the mouth, throat and nose. As an application of this action, the drug is occasionally used to check excessive secretion of saliva in ptyalism, mercurial salivation, etc. It is also used in coryza, especially in the first stage, to lessen the congestion and excessive secretion of the nasal mucus. It is held by some to be very useful in sore throat. It is used in case of excessive expectoration in bronchitis, bronchorrhea, etc. It also checks the secretion of saliva and mucus during anesthesia in operations on the throat, larynx, etc.

Atropin lessens the secretion of hydrochloric acid by the stomach. It is given for this purpose in hyperchlorhydria, gastric ulcer, etc. While it has a decided effect on the secretion, its use should not be continued for a long time. It lessens the secretion of the pancreatic juice, or at least prevents the increase that follows the ingestion of physostigmin and other drugs, but does not prevent the action of secretin. It is doubtful whether it has any effect on bile. It is said not to affect the secretion of intestinal mucus. It relaxes spasm of the intestinal musculature and in small doses favors the normal peristalsis. It is therefore of great service in spasmodic affections of the stomach and intestine. It may be prescribed in colic, painful spasms due to gastric, duodenal or intestinal ulcers, spastic constipation, etc. It is a serviceable anodyne in gall-stone colic and may render the use of morphin unnecessary.

In small or moderate doses it acts as a respiratory stimulant, but large doses cause respiratory paralysis. It may be employed with good effect in cases in which the respiration is embarrassed from other than mechanical agents. It is used for this purpose in morphin poisoning, but it should be administered with great care on account of the respiratory depression caused by large doses. To obviate the effect of morphin on the respiration it is given with the latter drug in hypodermic injections. It has also been given with morphin as a preliminary to anesthesia by ether; in such cases it also serves to lessen the salivary and bronchial secretions.

By paralyzing the vagus endings atropin increases the rapidity of the heart-beat. Its depressant action on the vagus is made use of in the diagnosis of certain disturbances of the cardiac rhythm, particularly bradycardia. If the slow pulse is due to an organic lesion of the conducting mechanism (heart-block), it will persist in spite of the action of atropin, but if it is due to vagal stimulation, an increased rate usually results from an effective dose.

Atropin in moderate doses relaxes the blood-vessels of the skin so that the skin, especially of the face and upper extremities, becomes red, sometimes showing an eruption closely resembling that of scarlet fever. In larger doses it contracts the vessels of the splanchnic area and raises the blood-pressure. In still larger doses a general fall of blood-pressure occurs, accompanied by a very rapid and feeble pulse. It is not, however, an essential heart tonic.

The secretion of sweat is reduced by atropin. It is used for the suppression of night sweats, especially in pulmonary tuberculosis. A single dose given at night may be followed the next night by a larger dose if the first was not successful. The use of atropin for this purpose should be deferred as long as practicable and discontinued as soon as may be on account of disturbing influence on digestion.

Atropin produces dilatation of the pupil, paralysis of the accommodation and consequent disturbance of vision by a local action on the oculo-motor nerve endings in the iris. When the drug is taken internally the effect is due to the atropin circulating in the blood, and is bilateral. When the drug is applied locally the action is unilateral unless some general absorption takes place or some of the solution is introduced into the other eye by accident.

To produce these actions, atropin is employed in solution dropped into the conjunctival sac for the purpose of facilitating the examination of the eye with the ophthalmoscope. The dilatation of the pupil also serves as a diagnostic measure in case of iritis as the pupil dilates irregularly in this disease. The wide dilatation of the pupil also tends to prevent its adhesion to the cornea or lens.

Sufficient absorption may take place from lotions dropped into the conjunctiva to produce general symptoms and even to cause toxic effects.

Atropin is sometimes used externally in the form of ointment of belladonna for the relief of certain forms of neuralgia, especially those in which pain results from local conditions of the nerve or surrounding tissues. It has been advised for local use in the rectum to relieve the pain of hemorrhoids or fissure. It is an old remedy for enuresis, but must be regarded in most cases as a purely empirical treatment to be used after investigation has shown the absence of organic lesions which may be remedied in other ways. It is used for vesical spasm due to irritable neck of the bladder.

While the action of atropin on the milk is not entirely established, it is quite generally used locally in mammitis,

galactorrhea and when it is desired to check the secretion of milk.

DOSAGE: Unpleasant symptoms indicating the beginning of its physiologic effects are produced in some people by 0.0005 gm. or 1/125 grain. It is best, therefore, to begin with half this dose, 0.00025 gm. or 1/250 grain. Doses can be repeated once in two hours until distinct physiologic effects are produced. For the treatment of the stomach the remedy should be given in solution about fifteen minutes before the meal. For action on the bowels it is commonly given at bedtime. The same time is also chosen when giving it for night sweats.

When atropin is being administered the patient or his friends should be warned of the possible appearance of slight toxic symptoms. The first indication is usually dryness of the throat. Some dimness of vision is also likely to be experienced.

Serious poisoning usually begins with rapid pulse, flushing of the skin, talkative delirium, marked dilatation of the pupils, some elevation of the temperature and dryness of the throat and skin. This is followed more or less quickly by unconsciousness, prostration, paralysis of the voluntary muscles and marked vasomotor paralysis.

Atropinae Sulphas.—Atropine Sulphate, U. S. P.

PROPERTIES: Atropin sulphate occurs as a white, crystalline powder or microscopic crystals which are odorless, but have a very bitter, nauseating taste. Atropin sulphate is very soluble in water (1:0.38) and in alcohol (1:3.7).

INCOMPATIBILITIES: Atropin sulphate is incompatible with alkalies and their carbonates and with the precipitants of alkaloids generally.

DOSAGE: 0.4 mg. or 1/160 grain. See Atropin.

Bacterial Vaccines.—See Vaccina.

Balsamum Peruvianum.—Balsam of Peru, U. S. P.

A balsam obtained from a tree, *Toluifera Pereirae*; it contains traces of cinnamic and benzoic acids and their esters and resins.

ACTION AND USES: Balsam of Peru is not used internally, but is sometimes applied in ointment or in alcoholic solution or mixed with castor oil as a stimulant to indolent wounds and ulcers to promote the formation of granulations and the process of cicatrization. Balsam of Peru is much used as a parasiticide in diseases of the skin, such as scabies.

In scabies a 4 per cent. petrolatum ointment of Balsam of Peru, with the same amount of sulphur, chalk and green soap may be used. It should be applied freely at night and in the morning over all of the affected areas. The strength of this mixture can be doubled in obstinate cases. Balsam of Peru can be painted over portions of the body at night,

followed by a bath in the morning. In rare instances, however, Balsam of Peru produces violent dermatitis.

Balsamum Tolutanum.—Balsam of Tolu, U. S. P.

A balsam obtained from *Toluwifera Balsamum*, a tree indigenous to New Granada.

PROPERTIES: Balsam of Tolu occurs as a yellowish-brown, plastic solid, becoming brittle when old, dried or exposed to cold. It is very soluble in alcohol, but nearly insoluble in water.

ACTION AND USES: Balsam of Tolu like balsam of Peru is expectorant and stimulant; it is not used externally at the present time. The use of balsam of Tolu in medicine is largely restricted. It is used as one of the ingredients of compound tincture of benzoin and as a flavor or pleasant vehicle in the form of:

Syrupus Tolutanus.—Syrup of Tolu, U. S. P.

A saturated solution of the aromatic constituents of balsam of Tolu in syrup.

ACTION AND USES: Syrup of Tolu is much used as a vehicle for expectorant mixtures.

DOSAGE: 16 c.c. or 4 fluidrams.

Belladonnae Folia.—Belladonna Leaves, U. S. P.

The dried leaves of *Atropa Belladonna*, yielding by the process outlined in the pharmacopeia, not less than 0.3 per cent. of mydriatic alkaloids, chiefly hyoscyamin.

ACTION AND USES: The active principle of belladonna acts like atropin. Belladonna leaves are narcotic, anodyne, antispasmodic and mydriatic, the medicinal value being due to the contained mydriatic alkaloids.

Tinctura Belladonnae Foliorum.—Tincture of Belladonna Leaves, U. S. P.

One hundred c.c. represent 10 gm. of the drug.

INCOMPATIBILITIES: The tincture should not be prescribed with alkalies.

DOSAGE: 0.5 c.c. or 8 minims, representing approximately 0.00015 gm. or 1/400 grain of mydriatic alkaloids.

Extractum Belladonnae Foliorum.—Extract of Belladonna Leaves, U. S. P.

A hydro-alcoholic extract of belladonna leaves.

DOSAGE: 0.01 gm. or 1/5 grain, corresponding approximately to 0.00015 gm. or 1/400 grain of mydriatic alkaloids.

Emplastrum Belladonnae.—Belladonna Plaster, U. S. P.

This represents approximately 30 per cent. of extract of belladonna. The commercial and widely used plasters are not identical with that described in the pharmacopeia in other than alkaloidal content.

ACTION AND USES: The application of extract of belladonna to the skin secures a local anodyne effect which is employed for the relief of rheumatic and neuralgic pains and soreness. It is a serviceable application in acute inflammatory conditions. Belladonna plaster is often used to relieve palpitation of the heart.

DOSAGE: Plasters are not commonly made by pharmacists, but are supplied already spread by the manufacturers. They are usually prescribed by the size according to the area of skin to be covered.

Unguentum Belladonnae.—Belladonna Ointment, U. S. P.

An ointment containing 10 per cent. of extract of belladonna in a mixture of hydrous wool fat and benzoinated lard.

Benzoinum.—Benzoin, U. S. P.

A balsamic resin obtained from several species of *Styrax* trees indigenous to Siam, Sumatra and Java.

PROPERTIES: Benzoin contains benzoic acid, cinnamic acid and resins which are soluble in alcohol and to some extent at least in oils and fats.

Tinctura Benzoinae Composita.—Compound Tincture of Benzoin, U. S. P.

A mixture of the alcohol-soluble constituents of benzoin, 10 gm.; aloes, 2 gm.; storax, 8 gm, and balsam of Tolu, 4 gm., in sufficient alcohol to make 100 c.c.

ACTION AND USES: Compound tincture of benzoin is used as an application to the inflamed mucous membrane of the throat and bronchi by inhalation. It is soothing and acts as a stimulating expectorant. It is frequently administered by adding a teaspoonful to a glassful of boiling water and inhaling the vapor. It is also used as an addition to lotions of glycerol and water. It is useful as a stimulant and protective for ulcers, bedsores, cracked nipples and fissures of the lips, anus, etc.

Benzosulphinidum.—Benzolsulphinid, Saccharin, U. S. P.

The anhydrid of ortho-sulphamid-benzoic acid (Benzo-sulphonic-imid).

PROPERTIES: Benzosulphinimid is a white, crystalline powder, nearly odorless, having an intensely sweet taste even in dilute solutions. It is soluble in alcohol (1:25), but only slightly soluble in water (1:250). The addition of an alkali materially increases the solubility of saccharin in water by formation of a salt, and it is often prescribed mixed with an equal quantity of sodium bicarbonate or in the form of a sodium salt known as "soluble saccharin." Saccharin is about 500 times as sweet as sugar, but has a slightly different taste which is generally objected to after continued use.

ACTION AND USES: Saccharin has no pronounced pharmacologic effect. It slightly retards the action of the digestive ferments and in very large doses intravenously produces some depression and stupor. The saccharin administered is excreted almost unchanged in the urine.

It is used as a substitute for sugar in the food of diabetics. It must be remembered that saccharin has no food value. Some observers believe it to be always harmful.

DOSAGE: 0.2 gm. or 3 grains. It may be mixed with an equal weight of sodium bicarbonate to insure its solubility. A $\frac{1}{2}$ -grain tablet is the most useful.

Betanaphthol.—Betanaphthol, U. S. P. (Naphthol, U. S. P. 1890).

Betanaphthol is a phenol found in coal-tar, but usually manufactured from naphthalene.

PROPERTIES: It occurs as colorless, or pale buff-colored, crystalline laminae or a crystalline powder, having a faint phenol-like odor and a sharp, pungent, but not persistent taste. It is only very slightly soluble in water, 1:950, but is very soluble in alcohol (1:0.61).

ACTION AND USES: Betanaphthol is irritating to the skin or mucous membranes when applied in solution. If absorbed in considerable amount it may cause nephritis by irritation of the kidneys. It tends to destroy the red blood-corpuscles and has been known to produce changes in the retina and opacity of the lens. It is a powerful antiseptic, several times stronger than phenol.

Betanaphthol is applied externally as a parasiticide and antiseptic. Internally it has been much used as an intestinal antiseptic, but its utility is limited because of the danger of poisonous consequences from its possible absorption. It has also been used as an anthelmintic for the treatment of hookworm disease.

DOSAGE: 0.1 to 0.3 gm. or 2 to 5 grains. It is best given in powder or capsules. Externally it may be used in the form of ointment.

Bismuthi Subcarbonas.—Bismuth Subcarbonate, U. S. P.

A basic carbonate of bismuth of somewhat varying composition, some specimens being more basic than others.

PROPERTIES: Bismuth subcarbonate is practically insoluble in water or alcohol. It is decomposed by hydrochloric or nitric acid, giving a copious effervescence of carbon dioxide and producing a chlorid or nitrate of bismuth which enters into solution. When such a solution is diluted with water the insoluble basic chlorid (oxychlorid or subchlorid) or basic nitrate of bismuth is precipitated. Bismuth subcarbonate is readily decomposed by the acid of the gastric juice, but in this case while the acid is neutralized, an insoluble oxychlorid of bismuth remains, coating the stomach and thus acting therapeutically in the same manner as the original subcarbonate would.

INCOMPATIBILITIES: It is incompatible with sulphids, sulphur, acids and acid salts.

ACTION AND USES: Bismuth subcarbonate acts like other insoluble salts of bismuth. For its uses see Bismuth Subnitrate. In roentgenography it is to be preferred to bismuth subnitrate because of the fact that it cannot give rise to

nitrite poisoning, which has occasionally happened when large doses of the subnitrate were used.

DOSAGE: 0.5 gm. or $7\frac{1}{2}$ grains.

Bismuthi Subgallas.—Bismuth Subgallate, U. S. P., Dermatol.

A basic gallate of bismuth of somewhat varying composition, containing bismuth equivalent to from 52 to 57 per cent. of bismuth oxid.

PROPERTIES: Bismuth subgallate occurs as an amorphous, bright yellow powder without odor or taste. It is practically insoluble in water and in alcohol, but is decomposed by hydrochloric, nitric or sulphuric acid if heated. It is also decomposed by alkali hydroxids.

INCOMPATIBILITIES: Bismuth subgallate is incompatible with acids, alkalies, sulphids and sulphur.

ACTION AND USES: Bismuth subgallate was introduced under the name of "Dermatol" for treatment of skin disease. Its action and uses are similar to those of Bismuth Subnitrate, which see.

DOSAGE: 0.25 gm. or 4 grains.

Bismuth Subnitrates.—Bismuth subnitrate, U. S. P.

Bismuth subnitrate is a basic bismuth nitrate of somewhat varying chemical composition.

PROPERTIES: Bismuth subnitrate occurs as a heavy, white, odorless and almost tasteless powder. It is practically insoluble in water and in alcohol and is little affected by weak acid solutions. Hydrochloric acid of the usual strength of the gastric juice decomposes only a small amount in the course of several hours. Hydrochloric or nitric acid, if not too dilute, decomposes it, producing the chlorid or nitrate, which enters into solution; but when this solution is diluted with water the insoluble basic chlorid (oxychlorid or subchlorid) or basic nitrate (subnitrate) is precipitated.

Basic bismuth nitrate is rendered more basic by the action of alkali until finally it is converted into bismuth oxid, a nitrate of the alkali metal being formed. Under some circumstances the nitrate radical may be reduced to nitrite, especially by the putrefactive bacteria of the large intestine.

INCOMPATIBILITIES: It is incompatible with acids, tannins, sulphids and sulphur. With soluble carbonates and bicarbonates in the presence of water there is a liberation of carbon dioxid, a formation of insoluble bismuth compound, and the nitrate of the alkali metal. With iodids a double decomposition has been noted with the formation of the red basic iodid of bismuth and the nitrate of the metal whose iodid was used.

ACTION AND USES: Soluble bismuth compounds, as a rule, become converted in the presence of water into insoluble basic compounds. Most of the preparations used in medicine are already in the basic form. (As the action of bismuth preparations depends on the action of an insol-

uble powder, there is no object in prescribing a soluble salt.) The salt most frequently used is bismuth subnitrate; but since it sometimes produces poisonous effects, the subcarbonate is the preferable salt. All the compounds of bismuth used in medicine produce essentially the same effects.

Bismuth subnitrate is not appreciably affected by the gastric juice and does not materially lessen its acidity. When given in considerable doses it coats the mucous membrane and acts as a mechanical protective. It thus prevents the action of the digestive secretions and of irritating foods or other substances on the mucous membrane. The same is true of its effect on the mucous membrane of the intestine. It is not absorbed in the stomach. It undergoes chemical changes in the intestine and is probably absorbed there to some extent though seldom in sufficient quantities to produce symptoms of poisoning. It is excreted almost entirely by the cecum and other parts of the large intestine. It is turned black in the large intestine, probably from contact with sulphids. It seems to exert an astringent effect on the gastro-intestinal mucous membrane. When applied to the skin it acts mechanically, but on wounds and ulcers, as on mucous membranes, it acts as a protective, astringent and antiseptic. It is absorbed from wounds to a larger extent than from mucous membranes. A number of cases of poisoning have been so caused.

Bismuth subnitrate and other insoluble salts of bismuth are used in irritation of the stomach and intestines for their protective and astringent powers. They are useful to allay vomiting from gastric irritation. In the same manner they serve to check diarrhea, especially that arising from the ingestion of irritating foods. They are useful in hyperacidity and ulcer of the stomach by coating and protecting the mucous membrane. They are also employed in catarrh of the stomach and intestines. They may be given for the same purpose in ulcerative enteritis. Externally the subnitrate and subcarbonate are used as protective and antiseptic applications in skin diseases and as applications to ulcers or suppurating wounds and to promote the healing of old sinuses and fistulous tracts. In the latter case the bismuth is used in the form of a paste, combined with petrolatum and wax. A number of cases of poisoning have been reported due to the absorption of the bismuth; hence care must be exercised in its use and on the appearance of toxic symptoms, such as a blue line on the gums, headache, nausea and stomatitis, the bismuth should be removed from the fistula by the injection of warm olive oil.

DOSAGE: 1 gm. or 15 grains. For the treatment of ulcer much larger doses are used. Externally it is used freely as a dusting powder or in ointment. Very large quantities are used for the purpose of outlining the stomach and intestines by roentgenography and by the fluoroscope, but the occasional occurrence of nitrite poisoning has led to a preference of the subcarbonate or oxychlorid for

Roentgen-ray work. Poisoning by nitrites is indicated by vasomotor paralysis, tachycardia and asphyxia due to the formation of methemoglobin.

Bismuthi Subsalicylas.—Bismuth Subsalicylate, U. S. P.

PROPERTIES: Bismuth subsalicylate occurs as a white or nearly white amorphous crystalline powder, odorless and tasteless. It is practically insoluble in cold water; on prolonged boiling with water a portion of the salicylic acid passes into solution with the formation of a more basic bismuth salicylate.

INCOMPATIBILITIES: It is decomposed by hydrochloric acid or nitric acid; a more basic insoluble compound remaining. Alcohol also decomposes bismuth subsalicylate with accompanying solution of the liberated salicylic acid. The compound is also incompatible with alkali carbonates and bicarbonates, tannin and salts of iron.

ACTION AND USES: Bismuth subsalicylate, like other salts of bismuth, is protective and astringent; it has also been recommended to some extent as an intestinal antiseptic. Because of the ease with which it is decomposed it is best prescribed in the form of powders, capsules or cachets.

DOSAGE: 0.25 gm. or 4 grains.

CACHETAE—CACHETS

Cachets, or wafers, are made by pressing a mixture of flour, or starch and water between hot iron plates, and are used much in the same way as capsules for enclosing dry and powdered drugs, but are usually of much greater capacity. They have the advantage over capsules of being more readily dissolved and despite their comparatively large size are easily swallowed if previously dipped into water or if floated on a tablespoonful of water.

Caffeina.—Caffein, U. S. P.

A feebly basic alkaloid obtained from the dried leaves of *Thea sinensis* or from the dried seeds of *Coffea arabica* and also found in other plants. Tea contains from 1 to 4 per cent. of caffein; coffee from 1 to 2 per cent.

PROPERTIES: Caffein is trimethyl-xanthin. It occurs as white silky, glistening needles, usually matted together in fleecy masses, odorless and having a bitter taste. It is soluble in water (1:50) and in alcohol (1:50). The solubility in water is materially increased by the addition of sodium benzoate or sodium salicylate.

ACTION AND USES: Small doses of caffein act on the nervous system, stimulating the psychic centers, the respiratory and vasomotor centers and the reflexes. It modifies the circulation by stimulating the heart, and relaxing the vessels by direct action. The flow of urine is increased. Muscular contraction is facilitated and fatigue lessened. Excessive doses produce insomnia, nervousness, headache, palpitation and nausea or vomiting, especially in susceptible

persons. They lessen the capacity for mental or muscular work. Toxic doses may produce tetanic convulsions and cardiac dilatation.

CIRCULATION: Caffein has a rather complex and, therefore, somewhat inconstant action. In therapeutic doses the pulse may be quickened or slowed. The blood-vessels tend to dilate by the peripheral action and to contract by the central action. The dilatation probably predominates in most cases, but the blood-pressure rises slightly by increased force and output of the heart. This increased output and lessened resistance tend to produce a more rapid flow of blood, and this results in an increased flow of urine. These effects make caffein especially efficient in some cases of cardiac dropsy, although it is generally inferior to digitalis. The cardiac stimulation is also useful in temporary cardiac weakness.

By the Germans, and by many physicians in this country, caffein is looked on as the most valuable drug for the treatment of circulatory failure in acute infectious processes, such as pneumonia, peritonitis, scarlet fever, etc.

A disadvantage in the use of large doses is the cerebral stimulation produced, which often prevents sleep. Some authors do not approve of the use of caffein as a cardiac remedy, but believe that its utilities are confined to its diuretic action.

Caffein is used as a nervous stimulant in cases of nerve exhaustion. It is useful in collapse by causing rise of blood-pressure and stimulating the respiration. It may be used in narcotic poisoning in the form of hot coffee, or by itself, for its effect on the respiratory system. It is especially useful in opium poisoning, and it may be used in alcoholic poisoning on the same principle. It relieves some forms of headache, but in the congestive form it may increase the difficulty. It is excreted by the kidney partly under its own form, partly as mono- or dimethyl-xanthin. It does not increase the amount of uric acid in the urine.

DOSAGE: The dose of caffein varies from 0.06 gm. to 0.3 gm., or about 1 to 5 grains. When given in the form of coffee a cup made from a tablespoonful (15 gm.) would contain from 0.1 to 0.2 gm. or from 1½ to 3 grains.

The alkaloid may be given in the form of powder or in capsules or cachets. In combination with equal parts of sodium benzoate or sodium salicylate it dissolves readily, and may be given hypodermically or administered by mouth.

Caffeina Citrata.—Citrated Caffein, U. S. P.

A mixture of equal parts of caffein and citric acid.

PROPERTIES: Citrated caffein is a white odorless powder having a slightly bitter taste and an acid reaction. It is more readily soluble than the alkaloid but is not adapted for hypodermic use.

DOSAGE: 0.1 gm. or about 2 grains.

Caffeinae Sodio-Benzoas.—Caffein Sodio-Benzoate, N. F.

A mixture of equal parts of caffein and sodium benzoate.

PROPERTIES: It is freely soluble (1 : 2) in water and is well adapted for administration in solution either by mouth or hypodermically.

DOSAGE: 0.10 gm., or about 2 grains.

Calci Carbonas Praecipitatus.—Precipitated Calcium Carbonate, U. S. P., Precipitated Chalk, CaCO_3 .

Calcium carbonate, obtained by precipitating a solution of a salt of calcium by a soluble carbonate, collecting and washing the precipitate and drying.

PROPERTIES: Precipitated calcium carbonate is a fine white powder without odor or taste. It is practically insoluble in water, but dissolves to a considerable extent in water containing carbon dioxid. It is decomposed by acids, forming a salt of calcium and giving off carbon dioxid with effervescence.

ACTION AND USES: Calcium carbonate neutralizes the acid of the gastric juice, chlorid of calcium being formed and absorbed to some extent. If the stomach contains no acid it may escape solution and absorption. When absorbed the action is that of the soluble salts of calcium, which will be described under Calcium Chlorid.

Calcium carbonate is used chiefly as an antacid. For this purpose it is used as the base of many tooth-powders. It is given in the form of powder as an antacid in acid gastritis, hyperchlorhydria and gastric ulcer. In prescribing it for such affections its power to check diarrhea and the astringent power with which it is credited should be borne in mind. It is especially appropriate to cases in which there is hyperacidity with accompanying diarrhea and in diarrhea with acid fermentation. It may be used in skin-diseases as a dusting-powder to neutralize acid secretions and to protect the skin.

Calcium carbonate is the appropriate antidote to oxalic acid as it neutralizes the acid with the formation of the insoluble oxalate of calcium. It is also a suitable antidote to other corrosive acids and has the advantage that an excess can be given without doing harm.

DOSAGE: From 1 to 3 gm. or from 15 to 45 grains. The fine powder can be given to adults in teaspoonful doses as no harm is to be feared from moderate excess. One gram will neutralize 0.730 gm. of absolute HCl, or approximately the amount of free HCl in 500 c.c., or 1 pint of stomach contents having 40 degrees of free acidity.

Calci Chloridum.—Calcium Chlorid, U. S. P., CaCl_2 .

PROPERTIES: Calcium chlorid occurs as white, translucent fragments, which are odorless and have a sharp saline taste. It is very deliquescent and should be kept in well-stoppered bottles. It is freely soluble in water (1 : 1.3) and also in alcohol (1 : 8).

INCOMPATIBILITIES: Soluble carbonates, phosphates and sulphates produce a precipitate of the corresponding insoluble salts of calcium.

ACTION AND USES: The chlorid is more irritating to the stomach than other salts of calcium and it should always be given well diluted. It is absorbed slowly and imperfectly; in the intestine it may be converted into the insoluble carbonate or phosphate, and the greater part of the calcium is excreted in this form. Calcium salts are sedatives to muscle and nerve action when absorbed. A lack of calcium in the tissues is believed to be a factor in the development of nervous irritability, tetany and other forms of convulsions. Large intravenous doses of calcium act somewhat like digitalis. It is also believed to increase the coagulability of the blood under certain conditions, but not often. These effects, however, are not produced when calcium is given by the mouth. There is no lack of calcium in the system under ordinary diet. It is therefore doubtful whether calcium chlorid has much if any systemic action when administered by mouth. It is used, however, in hemophilia, typhoid fever and other hemorrhagic conditions, with the idea of increasing the coagulability of the blood. Direct observations of the coagulation time have given contradictory, but generally negative results. The clinical evidence is not very strong, but the drug is at least harmless. Calcium chlorid is also administered, apparently with benefit, against urticaria and serum rashes.

When injected directly into the blood, calcium salts depress the nervous and muscular systems, while the inactivation of calcium by the injection of oxalates and citrates produces convulsions. This is the basis of an as yet unproved hypothesis that certain forms of nervous irritability are due to disturbed calcium metabolism.

DOSAGE: 0.5 gm. or $7\frac{1}{2}$ grains. Calcium chlorid is best administered in dilute solution sweetened with syrup or elixir.

Calcii Hypophosphis.—Calcium Hypophosphite, U. S. P., $\text{Ca}(\text{PH}_2\text{O}_2)_2$.

PROPERTIES: Calcium hypophosphite occurs as colorless, transparent prisms, small lustrous scales or a white crystalline powder, which is odorless and has a nauseous, bitter taste. It is freely soluble in water (1:65) and practically insoluble in alcohol.

INCOMPATIBILITIES: It is incompatible with iodids and oxidizing agents. Caution should be observed in dispensing calcium hypophosphite as explosion is liable to occur when it is triturated or heated with nitrates, chlorates or oxidizing agents.

ACTION AND USES: It was formerly thought that hypophosphites were of special value because the phosphorus was in a less highly oxidized form than in the phosphates. There is no experimental and no convincing clinical evi-

dence for this belief. Calcium hypophosphite has merely the action of other soluble salts of calcium. (See calcium chlorid.)

DOSAGE: 0.5 gm. or $7\frac{1}{2}$ grains.

Calcii Lactas.—Calcium Lactate, N. N. R.

PROPERTIES: Calcium lactate occurs in white crystalline masses or powder, and is odorless and tasteless. It is soluble in water (1:10) but very slightly soluble in alcohol.

INCOMPATIBILITIES: It is incompatible with carbonates, sulphates and other compounds forming insoluble calcium salts.

ACTION AND USES: Calcium lactate has the pharmacologic action of other soluble calcium salts (see Calcium Chlorid) but is less irritating than the chlorid and, therefore, especially suitable for hypodermic use.

DOSAGE: 0.5 gm. or $7\frac{1}{2}$ grains.

Calcii Phosphas Praecipitatus.—Precipitated Calcium Phosphate, U. S. P., $\text{Ca}_3(\text{PO}_4)_2$.

PROPERTIES: Precipitated calcium phosphate occurs as a bulky, white, amorphous powder that is odorless and tasteless. It is practically insoluble in water and in alcohol, but is soluble in hydrochloric or nitric acid.

ACTION AND USES: Because of the fact that calcium phosphate forms much the greater portion of the mineral matter of the human body, it has been supposed to be indicated in a variety of conditions. There is little or no evidence that it is of special value or has any advantages over calcium carbonate. It may be administered in the form of powder or in a complex mixture by dissolving it in a suitable acid.

DOSAGE: 1 gm. or 15 grains.

Calx.—Lime, Calcium Oxid, U. S. P., CaO .

PROPERTIES: Calcium oxid occurs as hard, white or grayish-white masses, which, in contact with the air, gradually attract moisture and carbon dioxid and fall to a white powder; it is odorless and has a caustic taste. Calcium oxid is converted into calcium hydroxid on the addition of water, and this is slightly soluble in cold water (1:760), less soluble in hot water and practically insoluble in alcohol.

Milk of lime is calcium hydroxid mixed with water in the proportion of 1 part of lime to 4 of water.

ACTION AND USES: Lime as calcium hydroxid is a fairly active germicide. It will neutralize acids, forming soluble salts of calcium.

On account of its cheapness lime is much used as a disinfectant. In the form of milk of lime or whitewash, it is a serviceable application to privies or to the infected walls of rooms. It is especially useful for the disinfection of excreta. Freshly prepared milk of lime should be added

in volume equal to that of the material to be disinfected, the mass thoroughly mixed and allowed to stand for two hours before disposal.

Liquor Calcis.—Lime-water, Solution of Calcium Hydroxide, U. S. P.

A saturated solution of calcium hydroxid containing about 0.14 per cent. of $\text{Ca}(\text{OH})_2$.

PROPERTIES: Lime-water is a clear, colorless liquid without odor, and has an alkaline, bitter taste. Lime-water absorbs carbon dioxid from the air, readily forming the insoluble calcium carbonate, and leaving the solution weaker in consequence. The Pharmacopeia gives a convenient method of assay, and there is no excuse for the pharmacist who dispenses an inferior preparation. The strength of lime-water should be preserved by keeping an excess of lime in the bottom of the container. When the lime-water is to be used the clear supernatant liquid should be decanted.

INCOMPATIBILITIES: Lime-water is incompatible with acids and with carbonated, or ordinary hard water.

ACTION AND USES: Lime-water is antacid and astringent and is often used as an addition to milk for both adults and children. The usual proportion is 1 of lime-water to 4 of milk, but a mixture of equal parts may be given. This is administered in small doses to allay nausea and vomiting. It is also used for diarrhea. Lime-water is used externally in the treatment of burns.

DOSAGE: 15 c.c. or about 4 fluidrams, containing approximately 0.02 gm. or $\frac{1}{3}$ grain of calcium hydroxid.

Linimentum Calcis.—Lime Liniment, U. S. P. This is more popularly known as Carron oil from the name of the iron works, in England, where it originated.

PROPERTIES: It is essentially a mixture of equal parts of a bland fixed oil and lime-water of official strength. The Pharmacopeia at present directs the use of linseed oil, but this has no advantages over some of the more limpid, and less odorous fixed oils, and by some is thought to be less desirable.

ACTION AND USES: Lime Liniment is applied to burns by spreading it on cloths and applying to the surface. It should be renewed frequently in order to prevent the dressing from adhering.

Calx Chlorinata.—Chlorinated Lime, Chlorinated Calcium Oxid, U. S. P., Bleaching Powder.

Bleaching powder, often improperly called chlorid of lime, is a variable compound resulting from the action of chlorin on calcium hydroxid. It should contain not less than 30 per cent. of available chlorin, that is, chlorin which is set free by the action of an acid.

PROPERTIES: Chlorinated lime occurs as a white or grayish-white granular powder, having a chlorin-like odor, and a repulsive saline taste. It is only partially soluble in water or alcohol.

ACTION AND USES: Chlorinated lime is a disinfectant of about equal value with lime and is used in the same way.

A 5 per cent. solution is a convenient strength for ordinary use. It is being used quite extensively as the source of chlorin for treating contaminated drinking-waters. A fresh solution of about 0.5 per cent. strength should be added to the water to be treated, in the proportion of about 250 gallons per million gallons of water.

Liquor Sodae Chlorinatae.—Solution of Chlorinated Soda, U. S. P., Labarraque's Solution.

An aqueous solution of sodium hypochlorite and sodium chlorid containing at least 2.4 per cent. of available chlorin. It is made by decomposing a solution of chlorinated lime with sodium carbonate and removing the insoluble calcium carbonate formed.

ACTION AND USES: Solution of chlorinated soda, like chlorinated lime, is used chiefly as a disinfectant and antiseptic. Diluted with from 15 to 20 parts of water it may be used as a spray, gargle or wash.

DOSAGE: 1 c.c. or 15 minims.

Camphora.—Camphor, U. S. P.

A ketone $C_9H_{16}CO$ obtained from *Cinnamomum Camphora*, purified by sublimation.

PROPERTIES: Camphor occurs as white translucent masses of a tough consistence and a crystalline structure, readily pulverizable in the presence of a little alcohol, ether or chloroform; it has a characteristic odor and a pungent taste. It is very slightly soluble in water but freely soluble in alcohol, ether, chloroform and in fixed and volatile oils.

ACTION AND USES: Camphor stimulates the central nervous system, especially the medullary centers, and the circulation; locally it is mildly irritant and antiseptic. In toxic doses it produces delirium followed by unconsciousness with epileptiform convulsions. It has been used as a nerve depressant in epilepsy and convulsions, but owing to its stimulating action this use seems irrational.

Camphor tends to increase the blood-pressure. The effects on animals are complex and rather inconstant, but clinically large doses of camphor improve the pulse in impending cardiac collapse, probably by cardiac stimulation. For this reason it is used as a circulatory and respiratory stimulant in cases of collapse, syncope, cardiac failure, etc. It should be administered hypodermically, 0.5 c.c. of a 10 per cent. sterile solution in olive oil being injected every fifteen minutes for four doses if needed.

The antiseptic action of camphor is utilized in gargles and mouth washes. It is also given in dyspepsia as a carminative, and is used in the first stage of colds and other infections of the respiratory tract. It acts internally as a diaphoretic.

Locally camphor causes a loss of tone of the peripheral blood-vessels, and thus produces marked redness of the

skin. It has also a local anesthetic action. It is employed, for its mild rubefacient and counterirritant effects, as an ingredient of liniments.

DOSAGE: 0.10 gm. or about 2 grains.

The average dose of camphor is commonly given as 2 grains, but when a prompt cardiac or central action is demanded, much larger doses may be required. It may be given in an oily or alcoholic solution or in pills, capsules or cachets.

Aqua Camphorae.—Camphor Water, U. S. P.

A saturated solution of camphor in distilled water. It is a very weak preparation. Camphor water is often used as a vehicle for collyria.

DOSAGE: 10 c.c. or about 2 fluidrams (containing 0.04 gm. or $\frac{1}{2}$ grain of camphor).

Spiritus Camphorae.—Spirit of Camphor, U. S. P.

One hundred c.c. contain 10 gm. camphor in alcohol.

DOSAGE: 1 c.c. or 15 minims.

Linimentum Camphorae.—Camphor Liniment, U. S. P.
Camphorated Oil.

One hundred c.c. contain 20 gm. camphor in cottonseed oil, but a more limpid oil may be preferred, particularly if the preparation is to be used for hypodermic injection.

Cannabis Indica.—Indian Cannabis, U. S. P.

The dried flowering tops of the pistillate plants of *Cannabis sativa*, grown in the East Indies and gathered while the fruits are yet undeveloped, and carrying the whole of their natural resin. Much, if not all, of the available drug fails to comply with the official requirements.

ACTION AND USES: Cannabis Indica is a narcotic poison which stimulates the psychic functions and produces delirium of a pleasant character and a rapid succession of ideas which cause the time to seem very long. From this condition the patient passes into melancholy and then into a deep sleep. An aphrodisiac effect is reported in some cases. In many cases the primary stimulating stage is quite short and often absent. On account of the sleep resulting from this drug, it has been recommended as a hypnotic, but in general the use of cannabis indica as a hypnotic is not advisable. While it has not been known to produce fatal results, the effect on dogs indicates that it is not entirely safe, and the possibility of formation of a habit should be borne in mind.

DOSAGE: The preparations are best given by themselves, the extract in the form of pills and the tincture diluted or mixed with water at the time of administration.

Extractum Cannabis Indicae.—Extract of Indian Cannabis, U. S. P.

DOSAGE: 0.01 gm. or 1/5 grain.

Tinctura Cannabis Indicae.—Tincture of Indian Cannabis, U. S. P. One hundred c.c. represent 10 gm. Indian cannabis in alcohol.

DOSAGE: 0.5 c.c. or about 8 minims.

Cantharis.—Cantharides, U. S. P.

The beetle *Cantharis vesicatoria*, Spanish fly, thoroughly dried.

ACTION AND USES: Cantharides is very irritating to the intestinal canal, producing hyperemia of the mouth and throat and vomiting. It is readily absorbed from the intestinal canal and produces marked irritation of the kidneys. The nephritis affects, at first, the glomeruli and subsequently the urinary tubules. In its passage through the urinary channels, it irritates the mucous membranes of the bladder and urethra and produces a desire to urinate, sometimes amounting to strangury. Cantharides is also a local irritant to the skin and produces blisters. It may be absorbed from the skin in sufficient quantities to cause nephritis. Cantharides has been used for treatment of chronic nephritis and incontinence of urine. Only minute doses should ever be given for these affections, and as soon as any evidence of vesical irritation arises the remedy should be suspended. Many physicians believe that cantharides should never be used internally.

The local irritant action of cantharides is the basis of its use for the treatment of baldness, but it is of little benefit in that condition. It is used for baldness in the form of tincture greatly diluted with alcohol (from 1:15 to 1:30) or in ointments. The chief use of cantharides is as a vesicant. It is contra-indicated in nephritis, and when vesication is desired in nephritis, another agent such as ammonia or chloroform should be selected. It may be used to produce redness and counterirritation in quantities not sufficient to blister. When the irritation is carried just to the point of beginning vesication, the result is known as a flying blister. The counterirritation may be rendered continuous by a succession of such "flying" blisters.

Ceratum Cantharidis.—Cantharides cerate, U. S. P.

This contains 32 per cent. of cantharides.

Ceratum cantharidis may be used for the purpose of vesication, but the blistering plasters commonly put up by manufacturers have a slightly different base. When cantharides is applied for the purpose of vesication a blister will usually be formed in the course of six hours. Vesication can sometimes be hastened by removing the cantharides plaster after a few hours and applying hot poultices. Blisters should not be applied directly over an inflamed part.

The following application has been suggested for baldness:

R	Tr. cantharid.	8		3ij
	Tr. capsici	1		℥xv
	Alcoholis	45		3jss
	Aq. ros.ad	150		3v
M.				

Capsicum.—Capsicum, U. S. P.

The dried, ripe fruit of *Capsicum fastigiatum*.

ACTION AND USES: Capsicum is carminative, stimulant and rubefacient. It is frequently prescribed in atonic dyspepsia, especially in cases due to chronic alcoholism. Such use should be cautious and not long continued.

DOSAGE: 0.05 gm. or about 1 grain. Externally it is frequently used in the form of liniment, preferably a simple mixture of the tincture with the official soap liniment. The tincture mixed with an equal quantity of glycerol is used as a gargle in tonsillitis, pharyngitis, etc.

Capsicum plaster is a suitable preparation for the relief of neuralgia, muscular rheumatism, etc.

Tinctura Capsici.—Tincture of capsicum, U. S. P.

One hundred c.c. represent 10 gm. of the drug in approximately 90 per cent. alcohol.

DOSAGE: 0.5 c.c. or 7½ minims.

CAPSULAE—CAPSULES

Capsules, either hard or soft, are made of gelatin molded over a suitable object. Hard capsules, made in different sizes, have long been used for administering liquids as well as solids, while soft capsules, containing an admixture of glycerol to the gelatin, are used extensively for oils and oleoresins or solutions of active drugs in oils.

Carbo Ligni.—Charcoal, U. S. P.

Prepared from soft wood and very finely powdered.

PROPERTIES: Charcoal is a black, odorless and tasteless powder, free from gritty matter.

ACTIONS AND USES: Locally charcoal is employed as a deodorant for fetid ulcers, etc., usually as a poultice.

Because of its well-known property of absorbing and condensing gases, charcoal has long been administered in various forms of gastric disturbance, the prescribers evidently losing sight of the fact that when thoroughly wet it almost entirely loses its property of absorbing gases. It is used to indicate the length of time food remains in the alimentary tract. Charcoal administered with the test diet causes the feces formed from that diet to have a black color. It is also commonly used to distinguish the periods of diet in metabolism experiments.

DOSAGE: 1 gm. or 15 grains. Preferably administered in cachets or capsules.

Cardamomum.—Cardamom, U. S. P.

The dried nearly ripe fruit of *Elettaria repens*.

ACTION AND USES: Cardamom is used as an aromatic, carminative and stomachic.

Tinctura Cardamomi.—Tincture of Cardamom.

Represents 20 per cent. of cardamom in diluted alcohol. This preparation should not be confounded with compound tincture of cardamom, which is a comparatively weak solution of the soluble constituents of cardamom, cinnamon and caraway in diluted alcohol, colored red with cochineal.

DOSAGE: 5 c.c. or 1 fluidram.

Caryophyllus.—Cloves, U. S. P.

The dried flower buds of *Eugenia aromatica*.

Oleum Caryophylli.—Oil of Cloves, U. S. P.

A volatile oil distilled from cloves consisting largely (80 per cent.) of eugenol.

PROPERTIES: Oil of cloves occurs as a colorless or pale yellow, thin liquid, becoming darker and thicker by age. Oil of cloves is freely soluble in alcohol, but nearly insoluble in water.

ACTION AND USES: Oil of cloves is antiseptic and aromatic. It is frequently used as a carminative and externally as counterirritant. In dental practice it has long been used as an anodyne.

DOSAGE: 0.2 c.c. or 3 minims on granulated sugar or in some emulsion, on shaved ice or in capsules. To relieve toothache it should be dropped on a small piece of cotton and inserted into the cavity in the tooth, if one is present. For external use it is usually diluted with 2 or 3 parts of fatty oil.

CATAPLASMATA—POULTICES

Poultices are soft solid preparations used for the purpose of applying heat and moisture to localized areas of the body. This application of heat induces a superficial hyperemia which is believed to influence the circulation of underlying parts. The poultices may also be made the means of applying counterirritant drugs. A long-continued application tends to cause relaxation of the skin and to render it flabby.

Cera Alba.—White Wax, U. S. P., is the bleached form of:

Cera Flava.—Yellow Wax, U. S. P.

A solid substance prepared from the honeycomb of the bee, *Apis mellifera*. In medicine wax is chiefly used to stiffen ointments.

CERATA—CERATES

Cerates are solid ointment-like preparations containing sufficient wax to prevent them from melting at the temperature of the body.

For the preparation included in this list see:
Ceratum Cantharidis, under Cantharis.

CHARTAE—PAPERS

Medicated papers are pieces of absorbent paper that have been treated with medicinal substances or suitably sized paper coated with a layer of the medicinal substance.

For the preparation included in this list see:
Charta Sinapis, under Sinapis.

Chloralum Hydratum.—Hydrated Chloral, U. S. P. Chloral Hydrate.

A crystalline solid, $\text{CCl}_3\text{CH}(\text{OH})_2$, the hydrate of trichloroacetaldehyd (chloral), CCl_3CHO .

PROPERTIES: Hydrated chloral occurs as colorless and transparent crystals, having an aromatic penetrating odor, and a bitterish, caustic taste. It is very soluble in water, alcohol or ether.

INCOMPATIBILITIES: Hydrated Chloral is incompatible with alkalis and alkali carbonates, which cause the formation of chloroform. In mixtures containing hydrated chloral, an alkali, bromid and alcohol, a compound of chloral believed to be chloral alcoholate sometimes separates.

ACTION AND USES: Hydrated chloral acts on the central nervous system, producing a feeling of weariness followed by quiet sleep in which the pulse and respiration are slowed in the same manner as in normal sleep and the reflexes are not abolished. From this sleep the person can readily be awakened. Larger doses produce more rapid and deeper unconsciousness and abolition of reflexes. There is slowing of the respiration and fall of blood-pressure. The cutaneous vessels are dilated and a marked fall of temperature occurs. With fatal doses death occurs ordinarily by paralysis of the respiratory center but sometimes by paralysis of a weakened heart. Hydrated chloral is therefore to be used with special caution in cases of a weakened heart or blood-vessels.

Hydrated chloral is a very reliable hypnotic in insomnia due to nervous excitation. It is not so valuable when the sleeplessness is primarily due to a painful affection. In such cases morphin or codein is preferable. At times it may be combined with opium or morphin. It should not be used when there is a degeneration of the heart-muscle. In other cases of heart disease and in arteriosclerosis it may be given cautiously.

Hydrated chloral may be given in moderate doses as a nervous sedative, especially in conjunction with the bromids.

Hydrated chloral is also a valuable remedy to relax spasm. For this purpose it is often used in the first stage of labor to overcome rigidity of the os uteri. It is valuable in tetanus, in which it must be given boldly to overcome the convulsions. In delirium in fever it may be administered in small doses. It should not be given, however, if the delirium can be controlled by other means. It is useful as an antispasmodic in obstinate cases of hiccup. Hydrated chloral is a good antidote to strychnin.

DOSAGE: From 0.30 to 1.30 gm. or from 5 to 20 grains dissolved in water and given on cracked ice. The dose may be repeated in three hours, if needed, but the condition of the pulse and respiration should be carefully watched when large doses are given.

While hydrated chloral has the reputation of being especially dangerous, this refers to large doses. Smaller doses, 0.3 gm. or 5 grains, are about as effective as the ordinary doses of other hypnotics and relatively safe. Habit formation is quite common, perhaps more so than with some other hypnotics.

Chloroform.—Chloroform, U. S. P.

A liquid consisting of from 99 to 99.4 per cent. of chloroform, CH_2Cl_3 , and from 0.6 to 1 per cent. of alcohol, $\text{C}_2\text{H}_5\text{OH}$, added as a preservative.

Chloroform is now largely made by the decomposition of chloral by alkalies.

PROPERTIES: Chloroform occurs as a heavy clear, colorless and mobile liquid, of a characteristic odor, and a burning sweet taste. It is but slightly soluble in water (1 : 200), but is miscible, in all proportions, with alcohol, ether and the fixed and volatile oils.

INCOMPATIBILITIES: Chloroform should be protected from the light by storing in a dark place or in dark well-stoppered bottles. It readily deteriorates under the influence of heat, light and air, and the decomposition products must be avoided in the use of this product in general anesthesia. For this reason the vapors should not be allowed to come in contact with a flame.

ACTION AND USES: Chloroform acts locally as a penetrating and fairly powerful irritant, which may blister if its evaporation is prevented. It is used in liniments. Taken by mouth, small doses are carminative, anodyne and antiseptic; it is therefore used in gastric fermentation and colic. Large doses are sometimes employed as a vermifuge, but are rather dangerous. Excessive doses produce unconsciousness and coma, similarly to the results of its inhalation.

Its main use is by inhalation, for the production of general anesthesia. The excitement stage is similar to that with ether, but of shorter duration and therefore less unpleasant. It is much more dangerous, however, most acute fatalities occurring by stoppage of the heart early in the administration. This danger is lessened by atropin.

The anesthetic stage is also more dangerous than with ether, there being a gradual but progressive fall of blood-pressure, even if the administration is carefully managed. The fall is due to depression of both the cardiac muscle and vasomotor center. The respiratory center is also more depressed. If an excessive concentration is given, death occurs, in this stage usually by stoppage of respiration; but since the heart and vasomotor center are also greatly weakened, recovery is more difficult than with ether. Sometimes, especially in cardiac disease, the heart may be the first to give out.

The irritant actions on the kidneys and respiratory tract are probably about the same as with ether.

Prolonged administration is especially dangerous, often producing death after several days by so-called delayed chloroform poisoning. This is characterized by general fatty degeneration, especially marked in the liver, which may pass into a condition analogous to acute yellow atrophy.

Chloroform is distinctly less safe as an anesthetic than ether and should be employed only when ether is unavailable or its use inadmissible for some reason. Chloroform is held by many physicians as specially suitable for anesthesia in children and during childbirth.

It is frequently given to relieve pain in the stomach or bowels, especially pain of a colicky nature.

DOSAGE: From 0.05 to 0.3 c.c. or from 1 to 5 minims.

Chloroform may be prescribed in a variety of forms. The Pharmacopeia includes:

Aqua Chloroformi.—Chloroform water, U. S. P.

A saturated aqueous solution of chloroform containing approximately 1 part in 200.

DOSAGE: It may be made extemporaneously by adding a few drops of chloroform to a glass of water, shaking well and allowing the excess to subside. A teaspoonful contains approximately 0.02 c.c. or $\frac{1}{3}$ minim of chloroform. The average dose of chloroform water is 15 c.c. or 4 fluidrams.

Spiritus Chloroformi.—Spirit of Chloroform, U. S. P.

One hundred c.c. contain 6 c.c. of chloroform in alcohol.

DOSAGE: 2 c.c. or 30 minims containing 0.12 c.c. or 2 minims of chloroform.

Linimentum Chloroformi.—Chloroform Liniment, U. S. P.

A 30 per cent. solution of chloroform in soap liniment.

Chromii Trioxidum.—Chromium Trioxid, U. S. P., CrO_3 .

This is more popularly, though incorrectly, termed chromic acid and should contain not less than 90 per cent. of CrO_3 .

PROPERTIES: Chromium trioxid occurs as needle-shaped crystals or prisms of dark purplish-red color and metallic luster; it is odorless, destructive to animal and vegetable tissue, deliquescent in air and very soluble in water (1:0.5).

INCOMPATIBILITIES: Because of its powerful oxidizing properties chromium trioxid should not be brought in contact with alcohol, glycerol or other oxidizable substances or explosion may result.

ACTION AND USES: In medicine chromium trioxid is used only as a caustic either in the solid form or in aqueous solution. Holt recommends in nasal hemorrhage from ulcer of septum to touch the ulcer with chromium trioxid.

Chrysarobinum.—Chrysarobin, U. S. P.

A neutral principle extracted from Goa powder, a substance found deposited in the wood of *Vouacapoua Araroba*.

PROPERTIES: Chrysarobin is a pale orange-yellow, microcrystalline powder, tasteless, odorless and irritating to the mucous membrane. It is very slightly soluble in water, but rather more soluble in alcohol and partially soluble in fats.

ACTION AND USES: Chrysarobin is antiseptic and antiparasitic and a powerful irritant to the skin. It is a more powerful stimulant than tar and must be used with correspondingly greater caution, except in dermatoses in which the production of an acute inflammatory reaction is not especially to be dreaded. It is employed in the treatment of fungus diseases of the skin but is also of especial use in chronic inflammatory dermatoses to produce a healthy reaction.

Chrysarobin is used externally in ointment or solution in the strength of from 2 to 10 or even 20 per cent. In the weaker proportions it is often quite as effective as in the stronger and does not excite acute dermatitis.

Chrysarobin stains the skin brownish, the hair greenish-yellow, the nails reddish-brown. Its use about the head should be avoided. It also stains clothing a walnut-brown. This stain can be removed by dilute solution of caustic soda or solution of chlorinated soda.

Unguentum Chrysarobini.—Chrysarobin Ointment, U. S. P.

Representing a solution of about 5 per cent. of chrysarobin in benzoinated lard.

Cinchona.—Cinchona, U. S. P.

The dried bark of different species, principally hybrids, of *Cinchona* yielding not less than 5 per cent. of anhydrous cinchona alkaloids, the chief of which is quinin.

ACTION AND USES: The preparations of cinchona are seldom used to obtain the systemic effects of quinin. They are mostly employed as bitter tonics. (See Quinin.)

Tinctura Cinchonae.—Tincture of Cinchona, U. S. P.

One hundred c.c. represent about 20 gm. cinchona in a mixture of alcohol, water and glycerol.

DOSAGE: 4 c.c. or 1 fluidram.

Tinctura Cinchonae Composita.—Compound Tincture of Cinchona, U. S. P.

One hundred c.c. represent 10 gm. red cinchona with bitter orange peel and serpentaria, as aromatics, in a mixture of alcohol, water and glycerol.

DOSAGE: 4 c.c. or 1 fluidram.

Cinnamomum.—Cinnamon, U. S. P.

The commercial drug is the dried bark or inner bark of different species of Cinnamomum.

ACTION AND USES: In medicine it is largely used as an aromatic or carminative preferably in the form of:

Oleum Cinnamomi.—Oil of Cinnamon, U. S. P.

A volatile oil containing about 75 per cent. of cinnamic aldehyd and occurring as a yellowish liquid having the characteristic odor of cinnamon and a sweetish spicy and burning taste. It is freely soluble in alcohol but only very slightly soluble in water.

DOSAGE: 0.05 c.c. or 1 minim.

Aqua Cinnamomi.—Cinnamon Water, U. S. P.

A saturated solution of oil of cinnamon in distilled water; it is largely used as a vehicle.

DOSAGE: 15 c.c. or 4 fluidrams.

Cocaina.—Cocaine, U. S. P.

An alkaloid obtained from several varieties of Coca.

PROPERTIES: Cocain forms large colorless prisms, having a slightly bitter taste and producing on the tongue a temporary numbness. It is only slightly soluble in water (1 : 600), soluble in alcohol (1 : 5), and also soluble in fixed oils, but insoluble in petrolatum and lard. The hydrochlorid is freely soluble in water.

INCOMPATIBILITIES: Solutions of cocain or of any of its salts after being kept a long time, or on boiling, are hydrolyzed more or less completely into ecgonin, benzoic acid and methyl alcohol. Solutions of cocain cannot be sterilized by boiling without some loss, but the amount of decomposition is so small as to be insignificant.

ACTION AND USES: Cocain paralyzes the peripheral nerves when applied locally, causing loss of sensation, and also contracts the blood-vessels. The mucous membrane becomes pale from local constriction of the vessels. When injected into a nerve or around the nerve sheath, anesthesia of the region to which the sensory fibers of the nerve are distributed is produced. By injecting a solution of cocain into the spinal canal anesthesia of the greater part of the body can be produced, but this method has often resulted in serious accidents or fatalities and is not to be recommended, despite the reported successful use by individuals who have become expert in recognizing and combating untoward symptoms.

Cocain produces first stimulation and then depression of the different segments of the central nervous system, beginning with the brain and extending to the spinal cord and the medulla. The stimulation of the brain has generally passed into depression before that of the spinal cord has begun so that when considerable doses have been given the symptoms may consist of a mixture of stimulation and depression. Stimulation of the brain is shown by increased psychic activity, loss of sense of fatigue, insomnia and muscular irritability. Depression is shown by somnolence, stupor and finally coma.

Convulsions frequently occur owing to cerebral action.

Respiration is at first quickened, it may have a Cheyne-Stokes character and later symptoms of respiratory paralysis appear. The heart may be temporarily stimulated, but is later paralyzed. The pulse is accelerated by nervous stimulation. The blood-pressure is raised by small doses but later falls from vasomotor paralysis. The temperature may be increased by cocain.

Cocain is a mydriatic acting both locally and centrally. The dilatation is not so complete as that produced by atropin, and reaction to light is not abolished.

The therapeutic use of cocain is derived almost exclusively from its anesthetic properties. It is much used to secure bloodlessness and anesthesia of the mucous membranes of the conjunctiva, nose and pharynx. In the eye it is said to increase the action of other alkaloids. It has an injurious action on the cornea in certain cases, producing desquamation of the epithelium and causing a keratitis. It is said that the danger of such injury can be minimized by closing the eyes for some time after cocain instillations. A moist compress may also be applied when strong solutions are used. The possible danger of absorption of the drug from the conjunctiva should not be forgotten. Cocain may be used as an anesthetic in operations on the eye and is considered better than any of its substitutes by many ophthalmologists.

In the nose cocain is used for astringent purposes in early stages of coryza and other inflammations in which it renders nose breathing easier by reducing the swelling of the turbinated bodies. It is also employed for the checking of hemorrhage. For anesthetic purposes it is advantageously combined with solution of an epinephrin salt. Hemorrhage of capillary origin is particularly amenable to the use of cocain.

The painful deglutition and other inconveniences of laryngeal tuberculosis may be alleviated by local applications of cocain solutions to the ulcerated areas. A more effectual method is the injection of cocain solution into the sheath of the superior laryngeal nerve. Cocain has been much used by local application to the nasal mucous membrane or by insufflation for the relief of asthma. Swabbing

the pharynx with a solution of cocain is sometimes employed to facilitate the swallowing of the stomach tube. Such an application, however, is seldom needed.

Cocain has been employed as an anti-emetic when vomiting depends on local irritation, but it should not be used for this purpose. It may be applied as a local anesthetic to hemorrhoids, but should be used with caution.

DOSAGE: For internal use the dose is 0.03 gm. or $\frac{1}{2}$ grain. As a local application to mucous membranes it is usually employed in the form of cocain hydrochlorid in a solution of the strength of from 2 to 10 per cent., the average being about 4 per cent. In the stronger solutions it should be used cautiously for fear of poisoning from the alkaloid absorbed. In the vagina and rectum 10 per cent. solutions may be used, but it is not safe to use solutions stronger than from 1 to 2 per cent. in the urethra.

Cocainae Hydrochloridum.—Cocaine Hydrochloride, U. S. P.

The neutral hydrochlorid of the alkaloid cocain.

INCOMPATIBILITIES: It is incompatible with alkalies, sodium borate and zinc sulphate.

DOSAGE: 0.03 gm. or $\frac{1}{2}$ grain.

Codeina.—Codeine, U. S. P.

An alkaloid obtained from opium or prepared from morphin by methylation.

PROPERTIES: Codein occurs as white, nearly translucent crystals or a crystalline powder, odorless and having a faintly bitter taste. Codein is soluble in water (1:88), and freely soluble in alcohol 1:12.

ACTION AND USES: Codein is analgesic, hypnotic and sedative. During recent years it has come to be used widely in place of morphin, particularly as a sedative. As an analgesic it is of value chiefly against abdominal and pelvic pain; it is largely used for cough. Codein is recommended by some authorities in place of opium in the treatment of diabetes.

DOSAGE: 0.03 gm. or $\frac{1}{2}$ grain.

Codeinae Phosphas.—Codeine Phosphate, U. S. P.

A salt that is freely soluble in water (1:2.25) and only slightly soluble in alcohol (1:261). It is preferred for hypodermic use.

DOSAGE: 0.03 gm. or $\frac{1}{2}$ grain.

Codeinae Sulphas.—Codeine Sulphate, U. S. P.

Soluble in water (1:30) and very slightly soluble in alcohol (1:1,035).

DOSAGE: 0.03 gm. or $\frac{1}{2}$ grain.

Colchici Semen.—*Colchicum Seed*, U. S. P.

The seed of *Colchicum autumnale*, assaying not less than 0.55 per cent. of colchicin.

ACTION AND USES: *Colchicum* produces marked irritation of the intestines, leading to looseness of the bowels with much pain and watery stools. It may produce severe inflammation and collapse. The collapse is believed to be due to the intestinal irritation and not to a central action. It also produces marked irritation of the kidney, which may lead to severe nephritis. *Colchicum* seed is said to be antineuralgic and analgesic. By many it is considered to be a specific in acute gout, controlling the pain and cutting short the attack. It may be given to prevent the occurrence of gouty attacks, and it is recommended by some to continue it in smaller doses after the attack.

Tinctura Colchici Seminis.—*Tincture of Colchicum Seed*, U. S. P.

One hundred c.c. represent 10 gm. *colchicum* seed in approximately 55 per cent. alcohol; it should assay approximately 0.05 per cent. of colchicin. It is similar to but not identical with the international standard tincture of *colchicum* seed.

DOSAGE: 2 c.c. or 30 minims.

In acute gout the dose is from 10 to 30 minims of the tincture once every four hours until some decided evidence of its action, such as nausea or slight purging, is induced. Severe purging should be avoided. *Colchicum* is of little value in rheumatism.

COLLODIA—COLLODIONS

Collodions are solutions of pyroxylin (guncotton) in a mixture of ether and alcohol, or of acetone, and are intended for external application.

For the preparations included in this list see the following:

Collodium.—*Collodion*, U. S. P.

A solution containing, in 100 c.c., 4 gm. of pyroxylin, or guncotton, in a mixture of 3 volumes of ether and 1 volume of alcohol. It is used as a protective and a vehicle chiefly in the form of:

Collodium Flexile.—*Flexible Collodion*, U. S. P.

A mixture of collodion with Canada turpentine (5 per cent.), and castor oil (3 per cent.). The addition of the small proportion of castor oil makes the resulting film elastic and more tenacious. The turpentine is superfluous and is probably retained for sentimental reasons.

Colocynthis.—Colocynth, U. S. P. Bitter Apple.

The peeled dried fruit of *Citrullus colocynthis*, from which the seeds should be separated and rejected in making pharmacopeial preparations.

Colocynth belongs to the class of one-time popular hydragogue cathartics whose use appears to be on the decline. It is used in making:

Extractum Colocynthisidis.—Extract of Colocynth, U. S. P.

DOSAGE: 0.03 gm. or $\frac{1}{2}$ grain.

Extractum Colocynthisidis Compositum.—Compound Extract of Colocynth, U. S. P.

A complex mixture of extract of colocynth with aloes and other drugs.

DOSAGE: 0.5 gm. or $7\frac{1}{2}$ grains.

The latter is one of the four preparations used in making the compound cathartic pills of the U. S. P.

CONFECTIONES—CONFECTIONS

Confections, conserves or electuaries were formerly used extensively, and usually occur as soft, pasty solids consisting of active drugs mixed with sugar or honey.

Copaiba.—Copaiba, U. S. P.

An oleoresin derived from one or more South American species of Copaiba.

PROPERTIES: Copaiba is a pale yellow to brownish-yellow liquid, having a peculiar, aromatic odor, and a persistent, bitter and acrid taste. It is practically insoluble in water, partially soluble in alcohol, and freely soluble in fixed and volatile oils.

ACTION AND USES: Copaiba is a mild stimulant, laxative and diuretic. It is chiefly excreted by the kidneys, partly by the bronchial mucous membrane, both of which it stimulates. It is also excreted by the skin and sometimes produces an eruption closely resembling that of measles. It is slightly antiseptic.

Copaiba is sometimes used in chronic bronchitis as a stimulating expectorant. It is chiefly used in gonorrheal urethritis, being applicable only in the chronic form of the disease. It may be given in conjunction with oil of sandal or the oleoresin of cubeb.

DOSAGE: 1 c.c. or 15 minims in capsule or emulsion, four or five times a day.

Creosotum.—Creosote, U. S. P.

A mixture of phenols and phenol derivatives, chiefly guaiacol and creosol, obtained during the distillation of wood-tar, preferably that derived from the beech.

PROPERTIES: It occurs as a colorless or slightly yellowish, highly refractive, oily liquid, having a penetrating smoky odor and a burning, caustic taste. Creosote is slightly, but not com-

pletely soluble in water (1:140), and miscible in all proportions with absolute alcohol; owing to its disagreeable odor and taste, it is seldom administered in the form of solution or mixture.

ACTION AND USES: When given internally creosote acts similarly to phenol. It is antiseptic and is one of the few drugs which appear to have a just claim to be useful as intestinal antiseptics. It is used to some extent externally for its antiseptic power. It is sometimes applied locally for leukorrhea and other infections of the mucous membrane. It has been given as a stimulant expectorant in chronic bronchitis and in tuberculosis. Experiments show that it does not affect the viability of the tubercle bacilli in the lungs. Some observers assert that its favorable action in tuberculosis is due to the production of intestinal antiseptics. It must be remembered, however, that its value in tuberculosis has not been determined beyond doubt. Less reliance is placed on it than formerly.

DOSAGE: 0.2 c.c. or 3 minims three times daily. It is preferably administered in the form of pills or capsules. If it impairs the appetite and disturbs digestion its use should be abandoned.

Cresol.—Cresol, U. S. P.

A mixture of the three isomeric cresols obtained from coal-tar, freed from phenol, hydrocarbons and water.

Cresol occurs as a straw-colored, refractive liquid, having a phenol-like odor and turning yellowish-brown on exposure to the air. It is soluble in water (1:60) and miscible in all proportions with alcohol, petroleum benzin, ether and glycerol; it is miscible with soap solutions and with solutions of alkali hydroxids, yielding an economical form of disinfectant.

ACTION AND USES: Cresol is an active poison resembling phenol in its effects. Its germicidal power is approximately four times as great as that of phenol.

DOSAGE: 0.05 c.c. or 1 minim. As a disinfectant it may be used in solutions varying in strength from $\frac{1}{4}$ to 1 per cent. It is usually employed in the form of:

Liquor Cresolis Compositus.—Compound Solution of Cresol, U. S. P.

A mixture of equal parts of cresol and a solution of a potassium soap.

ACTION AND USES: Compound solution of cresol has about twice the germicidal power of pure phenol. On account of its saponaceous character it is much used for the disinfection of the skin, for lubricating the hands, and for vaginal douches.

DOSAGE: As a douche and for washing wounds, solutions containing from 1 to 5 per cent. are appropriate.

Cupri Sulphas.—Copper Sulphate, U. S. P., $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$.

PROPERTIES: Copper sulphate forms large, transparent, deep-blue crystals, odorless, having a nauseous, metallic taste. Copper sulphate is freely soluble in water (1:2.2) but only slightly soluble in alcohol (1:400).

INCOMPATIBILITIES: Copper sulphate is incompatible with soluble salts of lead, which precipitate the insoluble sulphate of lead; with fixed alkalies and alkaline carbonates, which precipitate copper hydroxid or copper carbonate; with iodids, which form insoluble cuprous iodid with liberation of iodine, and with vegetable astringents containing tannin.

ACTION AND USES: Copper sulphate is astringent in small doses and irritant in large doses, producing nausea and vomiting. Copper sulphate in small amounts exerts a germicidal action in water containing algae, fungi or bacteria of the colon group; but when organic matter is abundantly present the germicidal action is greatly weakened. Externally copper sulphate acts as an astringent, stimulant or mild caustic according to the strength of the application.

Copper sulphate is used as a mild caustic in trachoma. It was formerly much used as an astringent in conjunctivitis. It is sometimes used as an emetic but is not to be recommended except in phosphorus poisoning, in which it acts by precipitating an insoluble compound of phosphorus and copper. It is occasionally prescribed for chronic diarrhea.

DOSAGE: As an astringent in diarrhea, 0.01 gm. or 1/5 grain; as an emetic, 0.3 gm. or 5 grains, not to be repeated.

As a caustic it is applied as the solid crystal or in pencils made by fusing 1 part of potassium alum and 2 parts of copper sulphate. When applications are made to trachomatous lids the affected parts of the everted lid should be touched lightly with the copper stick and the eye washed out afterward with lukewarm water. Collyria containing from 1 part in 1,000 to 1 in 100 may be used.

DECOCTA—DECOCTIONS

Decoctions are aqueous preparations made by boiling vegetable substances in water and then straining.

Diacetyl-Morphinae Hydrochloridum. — Heroin hydrochlorid, N. N. R.

The hydrochloric acid salt of diacetyl-morphin, a base formed by the action of acetic anhydrid on anhydrous morphin.

PROPERTIES: Heroin hydrochlorid occurs as a fine crystalline powder, odorless and having a bitter taste. Diacetyl-morphin hydrochlorid is freely soluble in water and in alcohol.

INCOMPATIBILITIES: It is incompatible with alkaline carbonates and hydroxids and also with strong acids, being quite readily decomposed.

ACTION AND USES: Diacetyl-morphin hydrochlorid resembles morphin, but it affects the brain less and depresses the respiration more. It is only slightly constipating. Repeated

use is likely to produce a habit, as with morphin. The sudden withdrawal of the drug from habitués may occasion dangerous respiratory failure. The use of heroin is resorted to by some cocain habitués who cannot procure their favorite drug.

Diacetyl-morphin hydrochlorid is used as a respiratory sedative to relieve cough. It is perhaps somewhat more effective, but has no important advantage over morphin. It is not so valuable as morphin for the relief of pain.

DOSAGE: 3 mg. or 1/20 grain.

DIGITALIS SUBSTANCES

This group includes as its most important members digitalis, strophanthus and squill. Others of the group are of minor importance. The drugs of the group increase the tone of the heart and arterial muscles and stimulate the vagus mechanism. In this way they slow and regulate the heart-beat, increase the cardiac output, and thus improve the circulation without affecting the blood-pressure directly. They are employed to secure compensation in valvular lesions, relieving the congestion, dyspnea, edema and other distressing symptoms and increasing the flow of urine. When administered by mouth, their absorption is slow and somewhat uncertain and their effects are correspondingly delayed and cumulative, so that they must be carefully watched. Overdoses produce nausea, vomiting, diarrhea, headache, cardiac irregularities and heart-block. The emetic action of ordinary doses is not due to local irritation, as commonly supposed, but is central and therefore cannot be avoided by rectal or intravenous administrations, or by the use of special preparations or isolated principles.

Digitalis.—Digitalis, U. S. P.

The dried leaves of *Digitalis purpurea*. Digitalis contains a number of glucosidal principles, the most important of which are digitoxin, digitalin and digitalein, the actions of which are essentially similar. A number of preparations of these glucosids are on the market, but many are of uncertain composition, and since they have not yet demonstrated any superiority over good preparations of the whole drug, the latter are to be preferred. Digitalis is a fairly stable drug, and those preparations that are made with high percentages of alcohol retain their full activity with little alteration for several years. Aqueous solutions deteriorate very rapidly.

ACTION AND USES: Digitalis, either in substance, or as one of the preparations referred to hereafter, is a cardiac tonic and diuretic. (See the preceding.)

Digitalis is useful whenever the systole of the heart is insufficient on account of incomplete exertion of its muscular power. It causes the heart to empty itself more completely and prevents it from dilating excessively during

diastole. It is useful in decompensated valvular disease, in dropsy from weak heart-action and in constantly irregular pulse, which is due to fibrillation of the auricle.

DOSAGE: 0.065 gm. or 1 grain, in powder or pill. Digitalis is a drug of variable strength, and hence a physiologically standardized preparation should be used. No perceptible effect is to be expected in less than twenty-four hours after oral administration. A slowing of the pulse indicates the beginning of its physiologic action; nausea and vomiting coming on during the administration of the drug are usually due to a commencing toxic action. If the vomiting is due to other causes, the administration of the digitalis may be continued best by a different channel, such as the rectum or the veins, or by hypodermic injection. For hypodermic and intravenous injections special preparations must be used, for which N. N. R. should be consulted.

OVERDOSAGE: As an overdose of digitalis or cumulative effects are most readily detected by a study of the pulse and heart-action, a patient who has been given digitalis should be kept under close observation. As a rule, the conditions which require digitalis also require that the patient stay in bed.

Symptoms of digitalis poisoning are shown by nausea and vomiting, sometimes great gastro-intestinal irritation, with pain and diarrhea, a very slow pulse, followed by a rapid and feeble one, or heart-block and marked prostration. The heart may become irregular and sudden changes in position may result in great aggravation of the condition and sometimes in sudden death. On the occurrence of any of these symptoms, the administration should be suspended.

Aside from its use in powder or pills, digitalis is used largely as:

Infusum Digitalis.—Infusion of Digitalis, U. S. P.

One hundred c.c. represent the water-soluble constituents of 1.5 gm digitalis partially preserved by alcohol (10 per cent.). It should be freshly prepared.

DOSAGE: 8 c.c. or 2 fluidrams.

Tinctura Digitalis.—Tincture of Digitalis, U. S. P.

One hundred c.c. represent 10 gm. digitalis in diluted alcohol.

DOSAGE: 1 c.c. or 15 minims.

Diphtheria Antitoxin.—See Serum Antidiphthericum.

Elaterinum.—Elaterin, U. S. P.

A neutral principle obtained from elaterium, a substance deposited by the juice of the fruit of *Ecballium Elaterium*.

PROPERTIES: Elaterin occurs as minute, white hexagonal scales or prismatic crystals, without odor and having a slightly acid, bitter taste. It is practically insoluble in water and only slightly soluble in alcohol (1:262).

Recent investigations would appear to indicate that the commercial substance is extremely variable in composition, and this is in keeping with clinical observations, many lots of the drug being found to be practically inert. Elaterin should not be confounded with elaterium.

ACTION AND USES: Elaterin is a powerful hydragogue cathartic, causing profuse watery evacuations with comparatively little pain.

It is used in dropsy, convulsions, puerperal eclampsia, etc. If used too freely or in debilitated persons it may produce dangerous weakness.

DOSAGE: 0.005 gm. or 1/10 grain.

ELIXIRIA—ELIXIRS

Elixirs are sweetened, aromatic, alcoholic liquids similar to cordials and having probably the same origin.

For preparations included in this list see:

Elixir Adjuvans, under Glycyrrhiza.

Elixir Aromaticum, under Alcohol.

EMPLASTRA—PLASTERS

Plasters are a class of solid preparations for external use and serve either as simple adhesives or as counter-irritants. The official plasters are practically never used but are replaced by the commercial plasters made on a large scale by machinery, which have as a base a mixture of rubber, with solvents or diluents.

For preparation included in this list see:

Emplastrum Belladonnae, under Belladonnæ Folia.

EMULSA—EMULSIONES—EMULSIONS

Emulsions are aqueous preparations in which oils or resins are suspended by means of mucilaginous substances.

ENEMATA—CLYSTERS

An enema, or clyster, is a liquid preparation intended to be injected into the rectum.

Epinephrina.—Epinephrin, N. N. R.

The blood-pressure-raising principle of the suprarenal gland, also produced synthetically, is official in the French, Italian and Belgian Pharmacopeias as adrenalin and in the German Pharmacopeia and the supplement to the Netherlands Pharmacopeia as suprarenin. In N. N. R. the following synonyms are enumerated: adnephtrin, adrenalin, supracapsulin, suprarenalin and 1-suprarenin synthetic.

PROPERTIES: Chemically epinephrin is described as 1, 2-dihydroxy-4²-methylamino-ethyl-4¹-ol benzene, $C_6H_3(CHOH.CH_2NH-CH_2)$, a substance with feeble basic properties, occurring in the suprarenal gland of the sheep or other animal. As commercially obtained it is a finely crystalline white or yellowish powder, odorless and slightly bitter. The free base is practically insoluble.

ble in water and is usually dispensed in the form of an aqueous solution, 1:1,000, of one of its salts. Epinephrin is oxidized readily and is thus destroyed in dilute alkaline solution.

ACTION AND USES: Epinephrin excites the action of the sympathetic nerves in such a way as to produce a variety of effects according to the function of the part supplied by the nerve. It produces a sudden rise of blood-pressure by contraction of the arterioles. The pulse is slowed by an action on the vagi. The heart is stimulated directly, but the resistance offered by the contraction of the blood-vessels is such that at times the heart is unable to overcome it and suffers passive dilatation. The rise of blood-pressure which results from the action of this drug is very transient, lasting, as a rule, not more than five minutes. When given by the mouth it produces no evident effect on the circulation, but it is readily absorbed from the mucous membranes of the nose, mouth, urethra, vagina and rectum, producing local contraction of the blood-vessels. Very large doses are tolerated when given hypodermically on account of the slow absorption due to the constriction of the blood-vessels of the part into which it is injected. It acts promptly after intravenous injection, but it appears to be rapidly eliminated or destroyed or its action is neutralized by antagonistic influences on the sympathetic ganglia. Epinephrin dilates the pupil, and this dilatation was employed at one time as a measure of the amount of epinephrin present in the blood, serum or other liquid. It inhibits the peristaltic movements of the intestine and increases the secretion of saliva and other glands which receive their nerve-supply from the sympathetic. Epinephrin may produce hyperglycemia and glycosuria, and its continued use may cause a degeneration of the internal coats of the arteries.

The chief therapeutic use of epinephrin is to constrict the peripheral blood-vessels by local application. In this way it may be used to diminish hyperemia of the conjunctiva, to reduce swelling of the turbinated bodies and to arrest hemorrhage from the mucosa of the upper respiratory tract. It is successful only in capillary or small arterial bleeding, as it cannot stop a large vessel hemorrhage. It is used to prevent hemorrhage in operations on the eye, nose, ear, etc.

For the arrest of hemorrhage it must be applied directly to the bleeding vessels or congested area. If the blood washes it away the application may fail because it has not time to act. It may be swallowed to check hemorrhage from the stomach, but the chances of success are small because a quantity of liquid is usually present in the stomach which dilutes the remedy so that it is useless. It should never be given for internal, concealed hemorrhage, because it is never desirable to raise blood-pressure in internal hemorrhage.

Epinephrin is employed in conjunction with local anesthetics, especially cocain, to limit the absorption of the

anesthetic and secure a more efficient local action. It has been used in asthma both by applying a spray to the nose or throat, or by the absorption of a tablet, powder or solution from the tongue, and also by hypodermic injection.

DOSAGE: Epinephrin or one of its salts is employed in solutions of a strength of from 1 : 10,000 to 1 : 1,000. For internal administration the dose of a 1 : 1,000 solution is from 5 to 10 drops. When an oily vehicle is to be used the base itself is prescribed, but when aqueous solutions are wanted one of the salts should be employed.

Ergota.—Ergot, U. S. P.

The sclerotium of *Claviceps purpurea*, replacing the grain of rye. Ergot appears to owe its activity to two active principles, ergotoxin (hydroergotin) and hydroxyphenylethylamin. The latter substance is closely related to epinephrin both in composition and in pharmacologic action.

ACTION AND USES: The several active principles of ergot have somewhat different actions, but the total effect is as follows:

Ergot causes powerful tonic, sometimes tetanic contractions of the uterus. It slows the pulse by stimulating the cardio-inhibitory centers. It also produces contraction of other involuntary muscles such as those of the blood-vessels, stomach and intestines, the bladder, etc. It seems to have a specially beneficial effect on acute congestions of the central nervous system.

Extreme and long-continued contraction of the blood-vessels, especially of the extremities, may lead to gangrene. Some animals do not show this action, while others are very susceptible.

The most common use of ergot is to prevent postpartum hemorrhage. For this purpose a full dose is often given as soon as the second stage of labor terminates, but it is much safer to give it after the placenta has been expelled. Its use during labor should be avoided as it may cause rupture of the uterus or asphyxia of the child. It is an effective remedy for "after-pains." Ergot is much used for hemorrhage from the uterus in menorrhagia and metrorrhagia. Its use for hemorrhage from other internal organs is not rational, as it increases blood-pressure and tends to prolong rather than check the bleeding.

It is asserted to be a good remedy for congestions in various regions. It has been recommended for the treatment of the early stage of acute pneumonia and also for pulmonary congestion in typhoid fever. It is at times useful in diabetes insipidus. In colliquative night-sweats due to relaxation of the blood-vessels, ergot is an efficient remedy.

DOSAGE: 2 gm. or 30 grains. It is sometimes administered in the form of powder.

Fluidextractum Ergotæ.—Fluidextract of Ergot, U. S. P.
(Extractum Ergotæ Fluidum, Pharm. 1890).

A hydro-alcoholic extract of ergot similar to, but not identical with, the international standard fluidextract of ergot.

DOSAGE: 2 c.c. or 30 minims.

Eucalyptus.—Eucalyptus, U. S. P.

The dried leaves of *Eucalyptus Globulus* collected from the older parts of the tree.

Eucalyptol.—Eucalyptol, U. S. P.

An organic oxid (cineol), obtained from the volatile oil of *Eucalyptus Globulus* and from other sources.

PROPERTIES: Eucalyptol occurs as a colorless liquid, having a distinctly camphoraceous odor, and a pungent, spicy and cooling taste. Eucalyptol is soluble in all proportions in alcohol, but is practically insoluble in water.

DOSAGE: 0.3 c.c. or 5 minims.

Oleum Eucalypti.—Oil of Eucalyptus, U. S. P.

The oil distilled from the fresh leaves of eucalyptus, and yielding not less than 50 per cent. of eucalyptol (cineol).

PROPERTIES: Oil of eucalyptus occurs as a colorless or pale yellow liquid, having a somewhat camphoraceous odor, and a pungent, spicy and cooling taste. It is miscible in all proportions with alcohol, but practically insoluble in water.

ACTION AND USES: Eucalyptol and oil of eucalyptus are antiseptic and expectorant. For internal use they are preferably administered in the form of capsules, and for local application in the throat or nose are used either in the form of oil sprays or are directed to be inhaled in the form of vapor from boiling water. Oil of eucalyptus, 2 c.c.; chloroform, 3 c.c. and castor oil, 40 c.c., mixed and taken in two portions at half-hour intervals, is being recommended as an anthelmintic for uncinariasis.

DOSAGE: 0.5 c.c. or 8 minims.

EXTRACTA—EXTRACTS

Extracts are soft solid or powdered preparations made by evaporating a solution of the soluble ingredients of vegetable drugs at a low temperature.

For preparations included in this list see:

Extractum Aloes, under Aloe.

Extractum Belladonnae Foliorum, under Belladonnae Folia.

Extractum Cannabis Indicae, under Cannabis Indica.

Extractum Colocynthis and Extractum Colocynthis

Compositum, under Colocynthis.

Extractum Gentianæ, under Gentiana.

Extractum Nux Vomicae, under Nux Vomica.

Extractum Opii, under Opium.

Extractum Rhamni Purshianae, under Rhamnus Purshiana.

Extractum Rhei, under Rheum.

Fel Bovis.—Oxgall, U. S. P.

The fresh bile of *Bos taurus*.

Fel Bovis Purificatum.—Purified Oxgall, U. S. P.

A yellowish-green, soft solid, or at the present time more frequently a yellow or greenish-yellow powder, having a peculiar odor and a bitter taste.

ACTION AND USES: Purified oxgall is reputed to be an intestinal antiseptic, cholagogue and laxative. The bile salts are held to be the most powerful stimulants to the secretion of bile.

DOSAGE: 0.5 gm. or $7\frac{1}{2}$ grains. It is best administered in the form of pills or in gelatin capsules which have been treated with a solution of formaldehyd.

Ferri Carbonas.—Ferrous Carbonate.

PROPERTIES: Ferrous carbonate is made by precipitating a solution of a soluble ferrous salt by a soluble carbonate. Such a precipitate tends to give off carbon dioxid, absorb oxygen and change rapidly into a basic carbonate, and finally into ferric hydroxid. Various pharmaceutical processes have been devised to prevent such a change. The principal of these are the preparations described below as Vallet's mass (*massa ferri carbonatis*) and Bland's pills (*pilulae ferri carbonatis*).

ACTION AND USES: The action of all forms of iron is essentially the same so far as the action of the iron ion is concerned. When an iron salt is received into the stomach it may be converted into a chlorid, but this is further changed during the process of digestion. The original form in which the iron was combined seems to make little or no difference in regard to the extent or the form in which it is absorbed. A large part of the iron ingested passes through the intestines without being absorbed. A smaller portion is absorbed, mainly through the lymph, and is deposited for a time in the blood-making organs, chiefly the spleen, where it is retained for an indefinite time as "reserve iron." Some of this supply is used in forming hemoglobin, which enters into the red blood-corpuscles. The rest is eliminated by the mucous membrane of the large intestine and only traces by the kidneys. Iron is not eliminated by the bile. The presence of iron in the blood in the amount resulting from medicinal administration produces no recognizable changes in normal individuals. Its salts with the stronger acids may act as gastro-intestinal irritants and astringents.

The only therapeutic action attributable to the iron ion is the improvement in the number of red blood-cells and in the amount of hemoglobin in them. For this purpose it is

indicated in anemia and in diseases of the blood in which anemia is a factor, such as leukemia. It is chiefly of value in anemia following hemorrhage, in chlorosis and in secondary anemias. In pernicious anemia it seems to be useless, but may be given along with arsenic.

Massa Ferri Carbonatis.—Mass of Ferrous Carbonate, U. S. P.

It is better known as Vallet's mass. It contains ferrous carbonate, with honey and sugar as preservatives, and sodium sulphate resulting from the process of manufacture.

DOSAGE: 0.25 gm. or 4 grains.

Pilulae Ferri Carbonatis.—Pills of Ferrous Carbonate, U. S. P.

These are popularly known as Blaud's pills. They consist of ferrous carbonate, potassium sulphate and sugar, with a smaller proportion of tragacanth and althea to make a mass. Each pill represents approximately 0.06 gm. or 1 grain of ferrous carbonate. They should be made fresh when wanted.

DOSAGE: 2 pills.

Ferri Chloridum.—Ferric Chloride, U. S. P. FeCl_3 .

Used in medicine principally in the form of:

Tinctura Ferri Chloridi.—Tincture of Ferric Chloride, U. S. P.

It is a hydro-alcoholic solution of FeCl_3 , containing not less than 13.28 per cent. of the anhydrous salt, corresponding to about 4.5 per cent. of metallic iron.

PROPERTIES: Tincture of ferric chlorid is a bright brownish liquid, having a slightly ethereal odor, a very astringent taste and an acid reaction, miscible in all proportions with either water or alcohol.

INCOMPATIBILITIES: It is incompatible with alkalis, alkali carbonates and with tannin, and the vegetable astringents. These give an inky color to the mixture. Tincture of ferric chlorid is also incompatible with iodids, from which it liberates iodine.

ACTION AND USES: Tincture of ferric chlorid is an astringent and is used in applications to the throat. It may be employed as a hematinic. It is also sometimes given in infections like erysipelas, but it is doubtful if it has any special value in this disease.

DOSAGE: 0.5 c.c. or 8 minims, freely diluted. Care should be taken to prevent injury to the teeth. Equal parts of the tincture, glycerol and water form a useful local application in acute tonsillitis.

Ferri et Ammonii Citras.—Iron and Ammonium Citrate, U. S. P.

It should contain the equivalent of 16 per cent. metallic iron.

PROPERTIES: It forms thin, transparent, garnet-red scales, without odor, having a saline, mildly ferruginous taste; deliquescent in moist air. It is freely and readily soluble in water, but practically insoluble in alcohol.

ACTION AND USES: Iron and ammonium citrate is one of the more widely used of the soluble preparations of iron. It may be directed to be dissolved in water, aromatic elixir, wine or syrup. It has been given hypodermically.

DOSAGE: 0.25 gm. or 4 grains.

Ferri Iodidum.—Ferrous Iodide, FeI_2 .

Ferrous iodid is a very unstable compound, easily undergoing oxidation. In order to preserve it in the ferrous condition it is commonly used in medicine in the form of syrup.

Syrupus Ferri Iodidi.—Syrup of Ferrous Iodide, U. S. P.

This is a syrup liquid containing about 5 per cent. by weight of FeI_2 and practically identical with the international standard syrup of ferrous iodid.

PROPERTIES: Syrup of ferrous iodid occurs as a transparent pale green or yellowish green liquid, having a sweet, strong, ferruginous taste and an acid reaction.

INCOMPATIBILITIES: The syrup is very susceptible to oxidation, and is incompatible with alkali carbonates, acid salts and vegetable astringents. On exposure to light the cane-sugar of the syrup undergoes gradual inversion.

ACTION AND USES: Syrup of ferrous iodid has the general properties of both iron and iodine.

DOSAGE: 1 c.c. or 15 minims, containing approximately 0.008 gm. or $\frac{1}{8}$ grain of iron and 0.04 gm. or $\frac{5}{8}$ grain of iodine.

Ferri Phosphas Solubilis.—Soluble Ferric Phosphate, U. S. P.

PROPERTIES: Soluble ferric phosphate should contain the equivalent of 12 per cent. of metallic iron and occurs in thin, bright-green transparent scales without odor, and having an acidulous, slightly saline taste. It is freely soluble in water, but practically insoluble in alcohol.

INCOMPATIBILITIES: Strong acids decompose it with formation of the astringent ferric compound of the acid used. Like other soluble salts of iron it is incompatible with alkalies, alkali carbonates and vegetable astringents.

ACTION AND USES: Soluble ferric phosphate has been recommended for the administration of iron in soluble form. It may be dissolved in water and flavored with simple elixir or sweetened with syrup.

DOSAGE: 0.25 gm. or 4 grains.

Ferri Sulphas.—Ferrous Sulphate, U. S. P., $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, Green Vitriol.

PROPERTIES: Ferrous sulphate occurs as large, pale bluish-green crystals, without odor, having a saline, styptic taste and efflorescent in dry air. It is freely soluble in water (1:0.9), but practically insoluble in alcohol.

ACTION AND USES: Ferrous sulphate is seldom administered in medicine as such, but is used chiefly for pharmaceutical purposes, in the making of ferrous carbonate. The sulphate was formerly used extensively as a disinfectant and deodorant but is now seldom so employed.

DOSAGE: 0.2 gm. or 3 grains.

Ferri Sulphas Exsiccatus.—Exsiccated Ferrous Sulphate, U. S. P.

One hundred parts represent approximately 150 parts of the crystalline substance.

PROPERTIES: Exsiccated ferrous sulphate occurs as grayish-white powder, having the chemical properties of ferrous sulphate and being slowly but completely soluble in water.

Ferrum.—Iron, U. S. P.

Metallic iron is used pharmaceutically in the production of preparations of iron, but in medicine is used chiefly in the form of:

Ferrum Reductum.—Reduced Iron, U. S. P.

PROPERTIES: Reduced iron should contain not less than 90 per cent. of iron. It occurs as a very fine, grayish-black powder, without odor or taste, and permanent in dry air. It is insoluble in water or alcohol.

DOSAGE: 0.065 gm. or 1 grain. Reduced iron is still widely used as a hematinic and is given preferably in the form of pills or powders, enclosed in capsules or cachets.

FLUIDEXTRACTA—FLUIDEXTRACTS

Fluidextracts are liquid preparations of uniform and definite strength made so that each cubic centimeter represents the soluble ingredients of 1 gm. of the crude drug.

For preparations included in this list see:

Fluidextractum Ergotae, under Ergota.

Fluidextractum Glycyrrhizae, under Glycyrrhiza.

Fluidextractum Hydrastis, under Hydrastis.

Fluidextractum Ipecacuanhae, under Ipecacuanha.

Fluidextractum Rhamni Purshianae, under Rhamnus Purshiana.

Fluidextractum Rhamni Purshianae Aromaticum, under Rhamnus Purshiana.

Fluidextractum Sennae, under Senna.

Fluidextractum Viburni Prunifolii, under Viburnum Prunifolium.

Formaldehydum.—Formaldehyd, CH_2O .

Formaldehyd is the aldehyd of formic acid, HCOOH . Formaldehyd is commonly obtained by oxidation of methyl alcohol. It is used in medicine in the form of:

Liquor Formaldehydi.—Solution of Formaldehyde, U. S. P., often referred to in literature under the proprietary name, formalin.

PROPERTIES: Solution of formaldehyd is an aqueous solution containing not less than 37 per cent. of formaldehyd, CH_2O . It is a clear, colorless liquid, having a pungent odor and caustic taste. Solution of formaldehyd is miscible in all proportions with water and alcohol. On standing it sometimes loses its transparency, its cloudiness being due to the separation of paraformaldehyd, a polymerization product of formaldehyd. Paraformaldehyd is also frequently formed on evaporation of the solution. Paraformaldehyd is a solid which is largely changed again into formaldehyd on heating.

INCOMPATIBILITIES: Solution of formaldehyd is incompatible with oxidizing agents and with alkalies. With ammonia it forms hexamethylenamin.

ACTION AND USES: Formaldehyd is a powerful germicide, especially valuable in the form of gas because of its penetrating power, but it is active only in the presence of an abundance of moisture. The solution is germicidal in the strength of from 1 to 2 per cent. (percentages refer to amounts of absolute formaldehyd, HCOH), but it may require from twenty to thirty minutes for it to act. In a strength of 1 : 5,000 it restrains the growth of many organisms, and in many cases a strength of 1 : 20,000 or 1 : 30,000 is sufficient to prevent the multiplication of bacteria. It is useful as a preservative of urine, although its reducing properties interfere with copper tests for sugar, and it is likely after the urine has stood some time to cause a precipitation of albumin, if present.

It hardens tissues and is used in histology for this purpose. It has a similar hardening effect on the living skin; it is very irritating and if repeatedly or continuously applied produces reddening, inflammation and necrosis. It is applied to the skin to restrain unilateral and excessive sweating. From 1 to 10 per cent. solutions in alcohol are appropriate for this purpose. It is sometimes used for the disinfection of the hands, in connection with a solution of soap.

The use of formaldehyd for the preservation of food has been quite commonly condemned on account of the disturbance of digestion which often follows its ingestion.

The principal application of formaldehyd is in room disinfection. For this purpose the vapor must be generated in a tightly closed room, containing plenty of moisture. Several methods have been described for generating the vapor, the most convenient being by the use of potassium perman-

ganate which, when added to the solution, by decomposing a part of the formaldehyd, generates sufficient heat to vaporize the remainder. For an ordinary-sized room 2 pounds of potassium permanganate are placed in a vessel of at least 25 quarts' capacity and a mixture of 1 quart of formaldehyd solution and 1 quart of water poured on it. Intense heat is generated by the reaction of the two chemicals, and by this heat the formaldehyd is vaporized. The heat is so great as sometimes to cause fire, against which due precautions should be taken. When the mixture has been made the operator should leave the room instantly. After the disinfection is complete the irritating fumes can be neutralized by ammonia.

Gelatinum.—Gelatin, U. S. P.

The purified air-dried product of the hydrolysis of certain animal tissues, as skin, ligaments and bones, by treatment with boiling water.

An amorphous, more or less transparent solid, usually shredded or in thin sheets; colorless or with a slight yellowish tint, inodorous, and having a slight, characteristic, almost insipid taste. Unalterable in the air when dry, but putrefying rapidly when moist or in solution. Gelatin is practically insoluble in cold water, but swells and softens when immersed in it, gradually absorbing from five to ten times its weight of water. It is soluble in boiling water, acetic acid and glycerin, but is practically insoluble in alcohol, ether or chloroform.

INCOMPATIBILITIES: Gelatin is coagulated by tannin, chlorin, bromin, and mercuric chlorid. If a solution of gelatin be mixed with formaldehyd, the gelatin is rendered hard and insoluble after evaporation and drying of the residue.

ACTION AND USES: Gelatin is largely used as a food product, though its full value in this respect has probably been exaggerated. It has also been used to some extent in solution for hypodermic injection to promote the formation of clot in aneurysms, and to arrest hemorrhage. There is, however, serious danger of infection from its use. Even boiling will not insure sterility as it may contain tetanus spores which are not destroyed by simple boiling. In pharmacy gelatin is used for the coating of pills, the making of gelatin capsules, gelatin disks, glycogelatin pastilles and for the making of glycerinated gelatin used as a base for suppositories.

Gentiana.—Gentian, U. S. P.

The dried rhizome and roots of *Gentiana lutea*.

ACTION AND USES: Gentian is one of a large class of substances with a bitter taste which have the power of stimulating the appetite and may possibly stimulate the secretion of the gastric juice. Experiments show that this effect is not due to an appreciable extent to a direct action on the mucous membrane of the stomach, but arises reflexly from the action of the medicine on the nerves in the mouth, chiefly those of taste.

Gentian and other bitters in moderate doses given a short time (five to fifteen minutes) before meals are useful in the treatment of loss of appetite, deficiency of the gastric secretions, chronic or subacute catarrhal gastritis, etc.

Tinctura Gentianae Composita.—Compound Tincture of Gentian, U. S. P.

A hydro-alcoholic tincture representing 10 per cent. of gentian with bitter orange peel and cardamom used for flavor.

DOSAGE: 4 c.c. or 1 fluidram.

Extractum Gentianae.—Extract of Gentian, U. S. P.

An aqueous extract of gentian evaporated to pilular consistence.

DOSAGE: 0.25 gm. or 4 grains.

Glandulae Thyroideae Siccae.—Desiccated Thyroid Glands, U. S. P.

The thyroid glands of the sheep, freed from fat, and cleaned, dried and powdered.

PROPERTIES: Desiccated thyroid gland is a yellowish, amorphous powder, having a slight, peculiar odor, and containing the active constituent of the thyroid tissue. It is partially soluble in water. The preparation should be standardized to contain 0.2 per cent. of iodine in organic combination.

ACTION AND USES: Dried thyroid gland acts chiefly if not entirely through a compound of iodine contained in it; the substance known commercially as "iodothyron" seems sometimes, but not always, to represent the full activity of the gland. When given in therapeutically active doses thyroid causes an increase of the nitrogen of the urine and a decrease in weight; it usually increases the absorption of oxygen and the elimination of carbon dioxide. It is one of the very few drugs which can properly be called stimulants of metabolism. The loss of weight is at first due to loss of water, then to increased metabolism of adipose tissue, although there is also an increased breaking down of protein unless the diet contains an abundance of protein. With larger, or long-continued doses there is a very rapid action of the heart, nervousness, tremors, headache, flushing of the surface, sweating and much more pronounced loss of weight.

Thyroid gland is of service in cases marked by deficient action of the gland. The most striking results are obtained in cretinism and myxedema and in the condition known as cachexia thyreopriva, due to the removal of the thyroid gland. The beneficial effects are seen in the improved condition of the skin, the reestablishment of perspiration and of a normal temperature, increased diuresis and loss of weight, improvement in the mental condition and, in young subjects, renewed growth, especially of the long bones and of the hair. In many cases after the more severe symptoms of hypothyroidism have disappeared, remarkably small doses

suffice to keep the patient in an almost normal state; it is often necessary, however, to continue such small doses indefinitely.

Thyroid is efficacious in many cases of milder degrees of hypothyroidism; but these are often difficult to diagnose.

In some forms of goiter the function of the thyroid is defective and the administration of the dried gland is indicated; but in most cases of goiter its use is condemned. Thyroid has been much used in obesity, but it is indicated in only a small proportion of cases and it should be given in moderate amounts so as not to do harm by the destruction of proteins. The effects, which are marked at first, are not permanent. Thyroid gland has been used with reported success in various skin diseases, such as psoriasis and eczema, and also in certain cases of amenorrhea.

DOSAGE: 0.05 gm. or 1 grain should be given as the initial dose three times daily, increasing gradually until improvement is noted; its administration should be discontinued if toxic symptoms appear. The patient should be careful of exertion and should take sufficient protein in his diet to compensate for increased loss of nitrogen from the action of the drug. The remedy may be given in powder, cachets or capsules. A dose of 0.6 gm. or 10 grains should rarely be exceeded.

Glycerinum.—Glycerin, U. S. P., Glycerol, $C_3H_5(OH)_3$.

A liquid obtained by the decomposition of vegetable or animal fats or fixed oils.

Glycerin occurs as a clear colorless liquid, of a thick, syrupy consistence, smooth to the touch, odorless, sweet to the taste and producing a sensation of warmth in the mouth. It is readily miscible with water or alcohol.

PROPERTIES AND USES: Glycerin is used in medicine chiefly as a solvent in preparing glycerites, and as a sweetening agent or vehicle in place of syrups.

Suppositoria Glycerini.—Suppositories of Glycerin, U. S. P.

Each suppository contains approximately 3 gm. or 45 grains of glycerin gelatinized by means of hard soap.

ACTIONS AND USES: A glycerin suppository, or glycerin itself, when introduced into the rectum tends to absorb water from the surrounding tissues and to cause increased peristalsis by reflex action, thus acting as a prompt laxative.

GLYCERITA—GLYCERITES

Glycerites are solutions of medicinal substances in glycerin.

For preparations of this type included in this list see:

Glyceritum Acidi Tannici, under Acidum Tannicum.

Glyceritum Amyli, under Amylum.

Glyceritum Boroglycerini, under Acidum Boricum.

Gelatinum Glycerinatum, under Gelatinum.

Glycerylis Nitras.—Glyceryl Trinitrate, Nitroglycerin, Trinitrin. Glonoin.

A compound of glyceryl with nitric acid obtained by the action of nitric acid on glycerol in the presence of sulphuric acid.

PROPERTIES: Nitroglycerin is a moderately volatile explosive liquid slightly soluble in water, but quite soluble in alcohol.

ACTION AND USES: Although a nitrate, nitroglycerin has the physiologic action of nitrites but acts more slowly than amyl nitrite. It may be given when it is desired to effect a steady reduction of the arterial pressure, but it gradually becomes inefficient. For this purpose it may be prescribed in arteriosclerosis, and in nephritis in which a high blood-pressure is a prominent symptom. It is used with success in some cases of angina pectoris. It is sometimes combined with digitalis in order to neutralize the contraction of the arterioles by the digitalis under the belief that therapeutic doses of digitalis exert a pronounced vasoconstrictor action.

Spiritus Glycerylis Nitratis.—Spirit of Glyceryl Trinitrate, U. S. P., Spirit of Nitroglycerin (Spiritus Glonoini, Pharm. 1890).

An alcoholic solution containing 1 per cent. by weight of glyceryl trinitrate.

PROPERTIES: Spirit of Glyceryl Nitrate is a clear, colorless liquid, having the odor and taste of alcohol. Caution should be exercised in tasting it, since even a small quantity of it is liable to produce violent headache.

DOSAGE: 0.05 c.c. or 1 minim two or three times a day.

Glycyrrhiza.—Glycyrrhiza, Licorice Root, U. S. P.

The dried rhizome and root of *Glycyrrhiza glabra* (Spanish licorice), or of *Glycyrrhiza glandulifera* (Russian licorice). Used in medicine chiefly as a vehicle and principally in the form of:

Fluidextractum Glycyrrhizæ.—Fluidextract of Glycyrrhiza, U. S. P. [Extractum Glycyrrhizæ Fluidum, Pharm., 1890].

An aqueous extract of glycyrrhiza preserved by means of glycerin and alcohol.

AVERAGE DOSE: 2 c.c. or 30 minims.

A simple mixture of 12 c.c. of fluidextract of glycyrrhiza and 88 c.c. of aromatic elixir is official in the U. S. P. as "elixir adjuvans."

Glycyrrhiza in the form of powder is used as a flavoring agent and vehicle in connection with:

Pulvis Glycyrrhizæ Compositus.—Compound Powder of Glycyrrhiza, U. S. P.

A mixture of senna (18 gm.), glycyrrhiza (23.6 gm.), sulphur (8 gm.), oil of fennel (0.4 gm.), and sugar (50 gm.).

ACTION AND USES: Pulvis Glycyrrhizae Compositus has been widely used as a laxative.

DOSAGE: 4 gm. or 60 grains, administered mixed with a suitable liquid, preferably milk.

Guaiacol.—Guaiacol, U. S. P.

Guaiacol is one of the chief constituents of beechwood creosote.

PROPERTIES: Guaiacol is a colorless or nearly colorless crystalline solid, melting at 28.5 C. (83.3 F.) or a refractive liquid, having an agreeable aromatic odor. It is soluble in water (1:53), and miscible with alcohol in all proportions.

ACTION AND USES: Guaiacol is antiseptic and germicidal, but to a less degree than creosote. When rubbed on the skin it is absorbed. When given in this way it is an active antipyretic, but its action is not easily controlled. It is an intestinal antiseptic and also an expectorant. It is not excreted by the lungs, however.

DOSAGE: Guaiacol may be given in emulsion or in the form of capsules. The initial dose should be 0.1 c.c. or about 1½ minims, gradually increased to 0.6 c.c. or 10 minims, if well borne. It is sometimes applied to the pharynx as a 50 per cent. solution in glycerin. The local use of guaiacol as an antipyretic is not to be recommended on account of the symptoms of collapse which sometimes follow these applications.

Guaiacolis Carbonas.—Guaiacol Carbonate, U. S. P. $(C_7H_7O)_2CO_3$.

Guaiacol carbonate is a white crystalline powder of neutral reaction, almost tasteless and odorless. It is practically insoluble in water, but is soluble in alcohol. It is decomposed readily with the liberation of guaiacol.

INCOMPATIBILITIES: It is incompatible with alkaline hydroxids.

ACTION AND USES: Guaiacol carbonate is inactive until decomposed with the liberation of guaiacol. This occurs only in the presence of putrefactive organisms so that in the intestine it acts only when an antiseptic is needed and the excess fails to be absorbed and is excreted with the feces. Consequently it is ordinarily not poisonous. It is used internally as a tasteless and non-poisonous substitute for guaiacol.

DOSAGE: 1 gm. or 15 grains. It may be given in powders, capsules or cachets.

Heroin Hydrochlorid.—See Diacetylmorphinae Hydrochloridum.

Hexamethylenamina.—Hexamethylenamin, U. S. P.

This preparation is also known under several trade names, of which aminoform, formin and urotropin are included in N. N. R.

Hexamethylenamin is hexamethylenetetramin, $(\text{CH}_2)_6\text{N}_4$, a compound produced by condensation of ammonia and formaldehyd with the elimination of water.

PROPERTIES: Hexamethylenamin forms colorless, lustrous, odorless crystals, freely soluble in water (1:1.5) and alcohol (1:10). The aqueous solution has an alkaline reaction. Hexamethylenamin is a base which combines with acids to form salts. These salts tend to lose formaldehyd, and most acids decompose the base completely with the liberation of formaldehyd. Even in aqueous solution a slow separation of formaldehyd occurs. The basic properties of hexamethylenamin are so pronounced that it displaces ammonia and other weak bases from their combinations.

INCOMPATIBILITIES: Hexamethylenamin is incompatible with acids which liberate formaldehyd and with salts of ammonium from which it separates ammonia. Acid salts like acid sodium phosphate and combined acids like acetylsalicylic acid (aspirin) react with it like other acids. It is also incompatible with tannin and mercuric chlorid, which precipitate hexamethylenamin.

ACTION AND USES: Hexamethylenamin produces no marked physiologic effects except those of formaldehyd, to which it gives rise. It is excreted in the urine, the bile, the cerebrospinal fluid and other serous fluids and by the mucous membranes of the respiratory tract and of the middle ear. When the urine is acid hexamethylenamin is decomposed, yielding formaldehyd. This product exercises a strong antiseptic action on the mucous membrane of the genito-urinary tract. When the urine is alkaline the decomposition does not occur. Formaldehyd is an excellent solvent of uric acid, and hexamethylenamin has been used in the hope of dissolving uric acid calculi, tophi, gravel, etc., and has been thought to increase the elimination of uric acid. Its use for this purpose, however, has practically been abandoned.

The chief use of hexamethylenamin is as a urinary antiseptic. It tends to free the urine from micro-organisms, and in many cases causes pus to disappear. It is of great value as a prophylactic in operations on the urinary organs. It is valuable in cystitis, pyelitis, etc. It is employed as a prophylactic in the bacilluria of typhoid fever. Some advocate its routine use to prevent the occurrence of this complication. It has been recommended to prevent the onset of nephritis in scarlet fever. As it has produced albuminuria and hematuria in some cases, it should be used with caution in cases in which inflammation of the kidney is present or anticipated.

In view of its excretion into the spinal canal, it has been recommended in cerebrospinal meningitis and in poliomyelitis. There is a tendency to extend its employment to infections of the respiratory tract and to the treatment of catarrh of the intestine and biliary passages. Favorable reports have been made of its use in rhinitis, otitis, bronchitis, etc., but further investigations are necessary to establish its efficiency. There is no good reason to expect a therapeutic action from it in alkaline mediums.

DOSAGE: 0.3 gm. or 5 grains three or four times a day in half a glass of water.

Homatropinae Hydrobromidum.—Homatropine Hydrobromide, U. S. P.

The hydrobromid of an alkaloid, tropin mandelate, produced synthetically.

PROPERTIES: It usually occurs as a white, odorless, crystalline powder, having a bitter taste. It is freely soluble in water (1:5.7) and soluble in alcohol (1:32.5).

INCOMPATIBILITIES: Homatropin hydrobromid has the ordinary incompatibilities of the salts of alkaloids.

ACTION AND USES: The actions of homatropin are identical with those of atropin, except for the fact that the mydriasis produced by it occurs more promptly and disappears in about eighteen hours. When applied freely to the conjunctiva it may be absorbed so that the bitter taste can be perceived, but the throat does not become dry as after atropin. The effect of this drug is increased by mixture with cocain.

DOSAGE: 0.0005 gm. or 1/125 grain. Homatropin is used chiefly as a mydriatic in place of atropin. It may be used in aqueous solution of 2 per cent. strength, or a drop of a 1:500 solution may be introduced into the conjunctival sac every five minutes for five times to produce a maximum dilatation in three-quarters of an hour. This will return to normal in from fourteen to eighteen hours.

Hydrargyri Chloridum Corrosivum.—Corrosive Mercuric Chloride, U. S. P., Corrosive Sublimate, HgCl_2 .

PROPERTIES: Mercuric chlorid occurs in the form of heavy colorless crystals or a heavy white powder having an acrid and persistent metallic taste. It is permanent in the air. It is very slowly soluble in water (1:13), and freely soluble in alcohol (1:3). Ammonium chlorid, sodium chlorid, tartaric acid and citric acid enhance its solubility in water.

INCOMPATIBILITIES: Mercuric chlorid is precipitated from its solutions by albumin, but redissolves in an excess of the albumin solution. Albumin in the form of egg-white forms the most useful antidote to corrosive sublimate, but a great excess should be avoided and the antidote should be followed by an emetic.

Mercuric chlorid is incompatible with soluble carbonates and hydroxids, forming insoluble mercuric oxid, and with iodids, forming mercuric iodid or complex mercuric iodids. It is incompatible with many alkaloids and other organic compounds. It is reduced to calomel or metallic mercury by iron, zinc and reducing agents in general. It dulls and tarnishes surgical instruments.

ACTION AND USES: Mercuric chlorid is chiefly used as a germicide and an antiseptic. It is also sometimes used as a specific antisyphilitic agent. In a proportion of 1:20,000 it kills non-spore bearing bacilli and in the proportion of

1:300,000 inhibits the growth of many bacteria. Spores of *Bacillus anthracis* are killed by a solution of 1:1,000. Its disinfectant action is limited by its deficient penetration and by the fact that it is greatly reduced by combination with organic matter. Mercuric chlorid is irritant to the skin, setting up a dermatitis. A sufficient amount may be absorbed from the skin to produce serious poisoning. The effects of the poison when absorbed from the skin or mucous membranes are seen in gastro-intestinal irritation, diarrhea, frequent foul-smelling and bloody passages resembling those of dysentery, various nervous symptoms, irritation of the kidneys with albumin and casts in the urine, marked weakness, etc. When taken in poisonous doses by the mouth it produces, in addition, irritation and ulceration of the mouth and throat, vomiting and corrosion of the mucosa of the stomach and intestines. Salivation and swelling and ulceration of the gums sometimes occur.

Dilute solutions of mercuric chlorid are used by hypodermic or intramuscular injection in the treatment of syphilis. The injections must be repeated daily and have the disadvantage of causing considerable pain.

Mercuric chlorid is used as a local application to the skin in some forms of skin disease, sometimes as an antiseptic, but also for the purpose of producing exfoliation of the epidermis. It is sometimes used in 1 per cent. alcoholic solution as an application to corneal ulcers. In the proportion of 1:5,000 it may be added to collyria to prevent fungus growths.

DOSAGE: From 0.002 to 0.01, or from 1/30 to 1/6 grain, in solution or in pill form. As an antiseptic application it may be used in solutions varying in strength from 1:20,000 to 1:2,000. For disinfection of clothing a solution of 1:1,000 may be used. To excite dermatitis and exfoliation, solutions varying in strength from 1:1,000 to 1:200 may be used, but caution should always be exercised in employing the stronger solutions for fear of absorption of the poison. The injection of mercuric chlorid solutions into the body cavities should be undertaken only with the greatest caution.

Hydrargyri Chloridum Mite.—Mild Mercurous Chloride, U. S. P., Calomel, HgCl .

PROPERTIES: Mild mercurous chlorid is a white impalpable powder, becoming yellowish-white on trituration with strong pressure, odorless, tasteless and permanent in the air. It is practically insoluble in water, alcohol or ether. It undergoes changes when exposed to the action of light or under the influence of alkaline chlorids, bromids or iodids, by which mercuric salts are more or less rapidly formed. The mercuric salt enters into solution in combination with the salt of the alkali metal present. Alkaline hydroxids convert it into mercurous oxid; ammonia forms with it a mixture of mercury and mercuric ammonium chlorid.

INCOMPATIBILITIES: Calomel is incompatible with alkalies, with oxidizing acids like nitric acid and also with soluble bromids and iodids. The fear that non-oxidizing acids like

hydrochloric acid will form mercuric chlorid from it is unfounded. Calomel is not incompatible with such acids.

ACTION AND USES: Mild mercurous chlorid is not irritating to the mucous membrane of the mouth, esophagus and stomach, but it provokes bowel movements by a slow action. This action has been thought to be due to a partial change into a mercuric salt or a protein compound. The absorption of the mercuric salt may produce symptoms of subacute mercurial poisoning. The chief indications of toxic effects are pain in the abdomen, loose passages, salivation, loosening of the teeth, swelling, soreness and ulceration of the gums, foul breath and general malaise.

Calomel was formerly supposed to have a cholagogue action, but it does not increase the quantity of bile secreted, although by its cathartic action it may increase temporarily the amount poured into the intestine. The stools resulting from the action of calomel are frequently greenish, resulting from changes in the bilirubin; this is partly because the bilirubin in the intestinal contents, being hurried through the colon, fails to undergo the change into urobilin which normally occurs. The change in color may also be due in part to the formation of mercuric sulphid.

Calomel is used for a special effect on the gastro-intestinal tract which is sometimes attributed to an antiseptic action. It is thought to be of special value as a cathartic in gastro-enteritis and at the beginning of mild catarrhs of the stomach and intestines. It is also frequently used to empty the bowels in conjunction with the more active salines in cases of infection, or toxemia. It is sometimes useful in small doses to check vomiting and is frequently administered as a laxative when the stomach is irritable because it is retained better than other cathartics. It is useful as an intestinal antiseptic.

Calomel is sometimes an excellent diuretic in cardiac dropsy. It is of much less value in other forms of dropsy.

It is sometimes applied externally to sluggish ulcers, and is used by insufflation on the cornea for ulceration or opacities, phlyctenular conjunctivitis, etc.

DOSAGE: Only a small portion of the calomel is absorbed, so that minute doses are generally effective. From 0.005 to 0.02 gm., or from 1/10 to 1/3 grain may be given every half hour or hour until from 0.1 to 0.2 gm., or from 1 to 3 grains have been given. The calomel should be followed in a few hours or the next morning by a saline cathartic. When calomel is used externally, care should be taken that no iodids are administered internally at the same time, because the presence of iodids in the secretions, for example, tears, may cause the formation of a mercuric salt and induce great irritation.

Hydrargyri Iodidum Flavum.—Yellow Mercurous Iodide, U. S. P., "Protiodid of Mercury," HgI , formerly also called "Green Iodid of Mercury."

PROPERTIES: Mercurous iodid is a bright yellow, amorphous powder, odorless and tasteless. By exposure to light it becomes darker, in proportion as it undergoes decomposition into mercuric iodid and metallic mercury. It is almost insoluble in water and wholly insoluble in alcohol.

INCOMPATIBILITIES: Mercurous iodid is incompatible with oxidizing agents, alkalies and the haloid salts of the alkali metals (chlorids, bromids or iodids), which tend to decompose it with the formation of mercuric salts. Alkali iodids decompose mercurous iodid with formation of metallic mercury and soluble alkali mercuric iodid.

ACTION AND USES: Mercurous iodid is used to secure the constitutional effects of mercury, especially in the treatment of syphilis. Its effects are much the same as those of calomel, although it is not so likely to produce diarrhea or salivation as the latter preparation.

DOSAGE: In syphilis mercurous iodid is best given in pill form in doses gradually increasing until slight toxic symptoms appear in the form of pain in the abdomen, diarrhea, salivation, etc. During its administration careful attention should be given to cleanliness of the mouth and skin. The teeth should be kept in good order, and soreness of the gums should be the signal for interrupting temporarily the administration of the remedy. The initial dose should not exceed 0.015 gm. or $\frac{1}{4}$ grain in the form of tablets, pill or powder, three times a day after meals, and this should be increased by the addition of 0.008 or $\frac{1}{8}$ grain daily to the limit of tolerance.

Hydrargyri Iodidum Rubrum.—Red Mercuric Iodide, U. S. P., "Biniodid of Mercury," HgI_2 .

PROPERTIES: Mercuric iodid is a scarlet-red powder, odorless and tasteless, and permanent in the air. It is nearly insoluble in water, but slightly soluble in alcohol (1:116). It is rendered soluble in the presence of soluble iodids, such as an equal weight of potassium iodid with formation of potassium mercuric iodid.

INCOMPATIBILITIES: Red mercuric iodid is incompatible with alkalies and their carbonates and alkaloids and their salts.

ACTION AND USES: Mercuric iodid is used as an antiseptic and germicide. It is more powerfully antiseptic than mercuric chlorid. It may be applied in potassium iodid solution. It is also used for the internal administration of mercury. A solution of mercuric iodid in sodium iodid (mercuric iodid 1 gm., sodium iodid 3 gm., water to make 100 c.c.) has also been used as a means of administering mercury by intramuscular injection.

DOSAGE: 0.003 or $\frac{1}{20}$ grain three times a day. It is frequently formed for internal administration by the prescription of a mixture of corrosive chlorid of mercury and potassium iodid.

	Gm. or c.c.
℞ Hydrargyri chloridi corrosivi.....	0 060
Potassii iodidi	6 00
Aquae	100
[M.]	

Hydrargyri Oxidum Flavum.—Yellow Mercuric Oxide, U. S. P., HgO .

PROPERTIES: Yellow mercuric oxid is a light orange-yellow, amorphous powder, odorless and having a somewhat metallic taste. It is permanent in the air, but turns darker on exposure to light. It is nearly insoluble in water and in alcohol.

INCOMPATIBILITIES: Yellow mercuric oxid is incompatible with acids, which dissolve it, forming mercuric salts.

ACTION AND USES: Yellow mercuric oxid is employed externally as a stimulant and parasiticide. Its chief use is in blepharitis, phlyctenular conjunctivis, etc., in which it acts as a stimulant.

DOSAGE: Yellow mercuric oxid should be used in the strength necessary to stimulate without producing unnecessary irritation. An ointment varying in strength from 0.5 to 2 per cent. should be prescribed. Some ophthalmologists recommend even weaker preparations.

Unguentum Hydrargyri Oxidi Flavi.—Ointment of Yellow Mercuric Oxide, U. S. P.

A 10 per cent. mixture of yellow mercuric oxid, with water, hydrous wool-fat and petrolatum.

DOSAGE: For use in the eye it should be diluted with from 10 to 100 parts of petrolatum so as to reduce the percentage of mercuric oxid to from 1 to 0.1 per cent.

Hydrargyri Salicylas.—Mercuric Salicylate, N. N. R.

A mercuric salt of salicylic acid in which one atom of mercury is combined with one molecule of salicylic acid.

PROPERTIES: Mercuric salicylate is a white amorphous powder, tasteless, odorless and neutral to litmus paper. It is nearly insoluble in water or alcohol, but soluble at the ordinary temperature in solutions of sodium hydroxid or sodium carbonate with the formation of a double salt.

INCOMPATIBILITIES: Mercuric salicylate is incompatible with iodids and (probably) other halogen salts.

ACTION AND USES: Mercuric salicylate is used as an antiseptic and for producing the internal actions of mercury. It is much used by hypodermic or intramuscular injection in the treatment of syphilis.

DOSAGE: Mercuric salicylate is given by intramuscular injection in a 10 per cent. suspension in liquid paraffin; 0.6 c.c. or 10 minims of this suspension are injected once in four days. It is held by some clinicians that the introduction of sufficient mercury by the mouth to eradicate syphilis thoroughly is practically impossible.

Hydrargyrum.—Mercury, U. S. P.

PROPERTIES: Metallic mercury in its ordinary form is a shining, silver-white, volatile and fluid metal without odor or taste. In its massive form it has comparatively little physiologic action and is not poisonous even in large quantities. When it has been reduced to a fine state of subdivision it is capable of absorption

either by the skin or mucous membranes, and then produces the ordinary pharmacologic effects of its salts. It is also very active in the form of vapor.

Mercury is widely used in the form of the following mixtures:

Hydrargyrum Cum Creta.—Mercury with Chalk, U. S. P., "Gray Powder."

A powder containing 38 per cent. of mercury with clarified honey and prepared chalk.

PROPERTIES: By long shaking and trituration of the mercury with the other ingredients it is reduced to so fine a state of subdivision that distinct globules cannot be seen with a lens magnifying 4 diameters. The preparation then forms a light gray, slightly adhesive powder, with little odor, and a slightly sweetish taste.

ACTION AND USES: This preparation is used in general in the same way as calomel, but it is less irritating to the intestines and in some cases does not move the bowels because of the antagonizing influence of the chalk. It is often used to secure the constitutional effects of mercury in cases of infantile and hereditary syphilis.

DOSAGE: 0.250 gm. or 4 grains contain approximately 0.1 gm. or 1½ grains of metallic mercury.

Massa Hydrargyri.—Mass of Mercury, U. S. P., "Blue Mass."

A mixture of 33 per cent. of mercury with glycyrrhiza, althea, glycerin and honey of rose.

The mercury is reduced to such fine division that globules are not visible under a magnification of 10 diameters.

ACTION AND USES: The effects and indications for this preparation are much the same as those of calomel. It may be used for laxative effects.

DOSAGE: 0.250 gm. or 4 grains.

Unguentum Hydrargyri.—Mercurial Ointment, U. S. P.

A mixture of 50 per cent. of mercury with prepared chalk, benzoinated lard and 2 per cent. of oleate of mercury.

The mercury is reduced by trituration to such fineness that globules are no longer visible under a lens magnifying 10 diameters.

Unguentum Hydrargyri Dilutum.—Blue Ointment, U. S. P.

Blue ointment is made by mixing approximately 2 parts of mercurial ointment with 1 part of petrolatum. It represents approximately 33 per cent. of metallic mercury and conforms to the strength, though not to the composition of the international standard for mercurial ointment.

ACTION AND USES: Mercurial ointment is a parasiticide; it is employed for the destruction of lice, but other measures are more cleanly and less calculated to cause irritation. The rubbing of mercurial ointment into the skin allows the absorption of a part of the mercury, thus securing its con-

stitutional effects. This is regarded by some authorities as the most efficacious method of treating syphilis except the intravenous or hypodermic method. Inunction has the great advantage of leaving the digestive tract undisturbed.

DOSAGE: 2 gm. or 30 grains of the ointment should be rubbed into a chosen area of the skin at night, and the part anointed should be cleansed by washing in the morning. A new area of the skin should be chosen for inunction the next evening. The ointment may be spread on cloths and bound on a part of the skin liable to some friction. If dermatitis is excited by this method of application it will subside rapidly as a rule on washing with a warm alkaline lotion and dusting with some bland powder. If irritation seems likely to arise the ointment may be diluted with an equal part of hydrous wool fat.

Hydrargyrum Ammoniatum.—Ammoniated Mercury, U. S. P.

A mercuric ammonium chlorid produced by the precipitation of a solution of mercuric chlorid by a solution of ammonia; it should represent not less than 78 nor more than 80 per cent. of metallic mercury.

PROPERTIES: Ammoniated mercury forms white pulverulent pieces or a white amorphous powder, having an earthy, afterward styptic and metallic taste. It is practically insoluble in water or in alcohol, but is gradually decomposed by washing with water. It is readily soluble in warm acids with decomposition. It also dissolves in cold solution of ammonium carbonate.

Ammoniated mercury is chiefly used in the form of:

Unguentum Hydrargyri Ammoniatum.—Ointment of Ammoniated Mercury, U. S. P.

A mixture of 10 per cent. of ammoniated mercury with white petrolatum and hydrous wool-fat.

ACTION AND USES: Ammoniated mercury in the form of an ointment of from 2 to 10 per cent. is much used as an antiseptic and local stimulant. In the strength of from 3 to 5 per cent. it is an efficient and non-irritating application for small areas of suppurating dermatitis. In stronger proportions (from 8 to 12 per cent.) it is a useful stimulating ointment for exciting a healthy inflammatory reaction, as in psoriasis. In seborrhea it may be applied after removal of crusts, in the form of a 2 per cent. ointment.

Hydrastis.—Hydrastis, U. S. P., Golden Seal.

The dried rhizome and roots of *Hydrastis canadensis*, yielding when assayed by the process given in the U. S. Pharmacopeia not less than 2.5 per cent. of hydrastin.

ACTION AND USES: Hydrastis is bitter and possesses the tonic and appetizing properties of other bitters. In addition it has been asserted that it has a special influence on mucous membranes, promoting their nutrition and favoring their return to a normal condition. For this property there

is no experimental evidence, and in the opinion of some its existence is doubtful. The action of hydrastis on the circulation is not well understood, but it is believed that the drug stimulates the circulation and raises the blood-pressure. It has a stimulating influence on the uterus, and has been employed to check menorrhagia and metrorrhagia.

Fluidextractum Hydrastis.—Fluidextract of Hydrastis, U. S. P.

A solution of the soluble constituents of hydrastis in a mixture of glycerin, alcohol and water. One hundred c.c. should contain 2 gm. of hydrastin.

Hydrastis contains also berberine, an alkaloid having the properties of a simple bitter.

Hydrastinin is an artificial alkaloid derived from hydrastin. It has marked influence in raising blood-pressure and a stimulating influence on the uterus much greater than hydrastin, from which it is derived. It is chiefly employed to check uterine hemorrhage.

DOSAGE: The dose of fluidextract of hydrastis is 2 c.c. or 30 minims; hydrastin can be given in doses of 0.01 gm. or 1/5 grain. Hydrastinin hydrochlorid is given in doses of 0.03 gm. or 1/2 grain.

Hydrogenii Dioxidum.—Hydrogen Dioxide, Hydrogen Peroxid.

Hydrogen dioxid or peroxid is a compound of hydrogen and oxygen, H_2O_2 , of syrupy consistency, which has a strong tendency to decompose into water and oxygen in the presence of oxidizable substances and of ferments capable of carrying oxygen. It is soluble in water, alcohol and ether. It decomposes readily when heated. Strong solutions are much more stable.

Aqua Hydrogenii Dioxidi.—Solution of Hydrogen Dioxide, U. S. P.

A slightly acid, aqueous solution of H_2O_2 (approximately 3 per cent.) corresponding to about 10 volumes of available oxygen.

PROPERTIES: Solution of hydrogen dioxid is a colorless liquid without odor, but having a slightly acidulous taste and producing a peculiar sensation and thick froth in the mouth.

ACTION AND USES: By its power of giving off oxygen it becomes a germicide and is employed as a non-toxic antiseptic application to suppurating wounds, inflamed mucous membranes, etc. It is employed as a topical application in the throat in diphtheria and other inflammatory exudates.

DOSAGE: Solution of hydrogen dioxid is usually applied diluted with from 1 to 4 volumes of water. It should be used with care in deep cavities. A free opening for the escape of foam and gas should be provided.

Hyoscyamus.—Hyoscyamus, U. S. P., Henbane.

The dried leaves and flowering tops of *Hyoscyamus niger*.

Tinctura Hyoscyami.—Tincture of Hyoscyamus, U. S. P.

One hundred c.c. represent the soluble constituents of 10 gm. of Hyoscyamus in diluted alcohol, and contain approximately 0.007 gm. of mydriatic alkaloids, chiefly hyoscyamin and scopolamin.

ACTION AND USES: The uses of tincture of hyoscyamus are similar to those of tincture of belladonna. It is thought to be more sedative to the nervous system.

DOSAGE: From 0.6 to 2 c.c. or from 10 to 30 minims.

Ichthyol.—Ichthyol, N. N. R., Ammonium Sulpho-ichthyolate.

Ichthyol is an aqueous solution of the ammonium salts of sulphonic acids prepared from the distillate from bituminous shales found in the Tyrol and elsewhere. These shales contain the fossil remains of fishes.

PROPERTIES: Ichthyol is a reddish brown to brown-black syrupy liquid having a characteristic empyreumatic odor and burning taste. It is miscible in all proportions with water, glycerol or oils, but is incompletely soluble in alcohol. Its aqueous solution has a faintly acid reaction.

ACTION AND USES: The actions of ichthyol are not well understood. It penetrates the unbroken skin to a certain extent. It is said to act as a vasoconstrictor on mucous surfaces. It has a mildly antiseptic action. On account of its content of sulphur it is credited with alterative properties.

It is applied to the skin in inflammatory conditions such as erysipelas, acne vulgaris, rosacea, lupus erythematosus, etc., to lessen hyperemia. It has also been used internally, but it is impossible at present to determine its true value. It has no specific action in tuberculosis, although it has seemed to increase appetite and strength in some cases. It has been tried with some apparent success in angioneurotic edema. It is used as a local remedy in gynecologic affections to relieve hyperemia and pain.

DOSAGE: The application of pure ichthyol to the skin is somewhat irritating, but a solution of 25 per cent. strength is generally not irritating. In chronic rheumatism a 50 per cent. ointment has been used locally. In gynecology it is combined with glycerol 1:10: Internally it may be given in solution or in the form of pills. The dose is from 0.2 to 2 c.c. or from 3 to 30 minims.

INFUSA—INFUSIONS

Infusions are aqueous preparations made by pouring hot or cold water over a vegetable drug and allowing the mixture to stand for a definite period and then straining.

For the preparation included in this list see:

Infusum Digitalis, under Digitalis.

Iodoformum.—Iodoform, U. S. P.

Iodoform is tri-iodomethane, CHI_3 , and is usually prepared by the action of iodine on alcohol or acetone in the presence of alkali or alkali carbonate.

PROPERTIES: Iodoform forms a fine, lemon-yellow powder or lustrous crystals having a peculiar, very penetrating and persistent odor, and an unpleasant, slightly sweetish and iodine-like taste. It is very slightly soluble in water (1:10,000), soluble in alcohol (1:50), and very soluble in ether (1:5 approximately). It is also somewhat soluble in fixed oils.

INCOMPATIBILITIES: It is incompatible with calomel, silver salts, chlorates and nitrites.

ACTION AND USES: Iodoform is a local anesthetic and an antiseptic. When absorbed through the skin or from denuded surfaces it produces intoxication, which is not evident until after the lapse of some time. When swallowed it is partly decomposed with the production of iodides, which produce their ordinary effects. Part of the drug is absorbed, however, in a form of combination not yet understood, and produces symptoms that are different from those ordinarily caused by iodine. Iodoform is slowly excreted, iodine compounds appearing in the urine for several days after a single dose.

The symptoms are restlessness, anesthesia, sometimes unconsciousness, occasional convulsions, hallucinations and delusions of persecution, rapid pulse and elevated temperature; in many cases collapse, coma and death may follow.

The physiologic actions of iodoform afford no rational basis for its internal use. Externally it is used as an antiseptic and stimulant in surgical tuberculosis. Several odorless organic compounds of iodine have been devised as substitutes for iodoform, but they are uniformly less actively antiseptic.

DOSAGE: 0.25 gm. or 4 grains. It is usually applied externally in the form of a dusting-powder, but may be used in the form of emulsion, as an ointment or as a surgical dressing in the form of gauze. For the relief of hemorrhoids it should be given in the form of suppositories.

Iodum.—Iodine, U. S. P.

PROPERTIES: Iodine is a heavy, bluish-black, dry and friable solid crystallizing in rhombic plates, having a metallic luster, a distinctive odor and a sharp and acrid taste. It is readily volatile. It is very slightly soluble in water (1:5,000), but soluble in alcohol (1:10). It is also soluble in solutions of iodides.

INCOMPATIBILITIES: Iodine is incompatible with alkalies and alkali carbonates, the alkaloids, with tannin and other vegetable astringents and with most volatile oils, particularly the terpene-containing oils.

ACTION AND USES: Cutaneous: Iodine irritates the skin, causing a sensation of heat and itching. In concentrated solutions it may cause blistering, or even corrosion, but it acts more slowly than many other irritants. It penetrates

into the deeper layers of the skin and small quantities are absorbed.

Iodin is applied to the skin for the purpose of exciting congestion of the underlying tissues. This congestion is supposed to cause the absorption of exudates. Its action is probably overrated. It is also used by surgeons for the disinfection of the skin, for which it is considered to be the most desirable agent. The application is made by painting the tincture over the part to be disinfected. The skin must be dry; wet applications should not be used previously. It is also used in various skin diseases for the purpose of producing an acute inflammatory reaction in the skin, and to cause the destruction of bacteria. Its effect on bacteria below the epidermis is probably due to the inflammatory reaction which it excites rather than to any direct action on the bacteria. For the prevention of tetanus it is recommended to apply to the skin about the wound a 3 per cent. alcoholic solution of iodine and to the wound pieces of gauze soaked in the same solution.

INTERNAL.—Iodin is more irritating to mucous membranes than to the skin. It is seldom used internally because of the irritating action on the stomach and intestines. This irritation may be so great as to excite a suppurative gastritis. The irritating action on the intestines may cause diarrhea. Small quantities of iodine are converted in the intestines into iodides and absorbed in this form. In some cases there is formed, also, a protein compound of iodine. After absorption iodine acts like the iodides (see Potassium Iodide). In cases of iodism, produced by the use of elementary iodine, there is tachycardia and irritation of the nervous system, but not so much affection of the skin and respiratory tract as is seen after the administration of the iodides.

Iodin should rarely be used internally; in treatment of obstinate vomiting the tincture has been recommended in doses of from 0.03 to 0.05 c.c. or from $\frac{1}{2}$ to 1 minim combined with the same quantity of phenol, given in a little water.

LOCAL USES: In the diseases of the eye iodine is sometimes used as a caustic agent and germicide to corneal ulcers of the simple type. It should be applied by means of a pointed toothpick soaked in the solution and used very cautiously.

For the treatment of chronic granular pharyngitis, in acute follicular tonsillitis, and in cases of middle ear catarrh associated with granular pharyngitis, it may be applied mixed with glycerol and combined with other remedies. The following formulas may be used:

		c.c.	
R	Tinct. iodi	3	℥ ℥℥℥
	Glycerini	30	℥ j-℥
R	Tinct. iodi		
	Tinct. ferri chlor.		
	Glycerini āā	q. s.	

In gynecology the tincture is often applied directly to the interior of the cervix and painted over the mucous membrane

of the vagina. This application is especially recommended in acute gonorrheal endocervicitis. Solutions of iodine have been much used in surgery. The tincture is often injected into cysts to cause the adhesion of their walls. Such applications should be made with caution. It is applied in a similar way to fistulous canals. A diluted solution is useful as a stimulant to ulcers.

Tinctura Iodi.—Tincture of Iodine, U. S. P.

One hundred c.c. contain 7 gm. iodine and 5 gm. potassium iodide dissolved in alcohol. This preparation is quite dissimilar to that formerly official in the U. S. P. or the international standard preparation generally prescribed in Europe, as it contains potassium iodide. The U. S. P. tincture of iodine has the advantage of being more stable so far as iodine content is concerned and of being miscible with water.

DOSAGE: 0.1 c.c. or 1½ minims.

Ipecacuanha.—Ipecac, U. S. P.

The dried root of *Cephaelis Ipecacuanha*, commercially known, as Rio, Brazilian or Para ipecac, or of *C. acuminata*, commercially known as Cartagena ipecac.

When assayed according to the method in the U. S. P., it should contain not less than 1.75 per cent. of ipecac alkaloids.

ACTION AND USES: When given by mouth in rather large doses, ipecac causes nausea and vomiting, chiefly through its local irritant action. It is, however, neither a very rapidly acting nor trustworthy emetic. Its use as such is almost entirely confined to pediatric practice.

In smaller doses it is nauseant and is used to promote the secretions of the respiratory tract.

Still smaller doses may act as stomachics through mild irritation of the gastric mucosa. When combined with opium, in the form of Dover's powder, ipecac is a diaphoretic.

Both ipecac and one of its principal alkaloids, emetin, are believed to be specific against amebic dysentery. In the treatment of this disease large doses of ipecac are required and opium or some other depressant drug often has to be given to prevent the occurrence of vomiting. Emetin, in the form of the hydrochloride can, however, be given hypodermically in doses which correspond to very large amounts of the crude drug, and its administration does not cause nausea or vomiting. It is probably this alkaloid alone to which the specific action of ipecac in amebic dysentery is due.

DOSAGE: The expectorant dose of ipecac is 0.05 gm. or 1 grain. As an emetic 1 gm. or 15 grains may be given. For use in dysentery it may be given in salol-coated pills. The coating should not be too thick, lest too large a dose of salol be given. Ipecac may also be given, suspended in mucilage of acacia, by a duodenal catheter. In dysentery an initial dose of 2 gm. or 30 grains may be given and vomit-

ing should be prevented by a previous hypodermic injection of morphin. The dose of emetin hydrochlorid for this purpose is 0.03 gm. or $\frac{1}{2}$ grain.

Fluidextractum Ipecacuanhae.—Fluidextract of Ipecac, U. S. P. (Extractum Ipecacuanhae Fluidum, Pharm. 1890).

One hundred c.c. represent 100 gm. ipecac in approximately 75 per cent. alcohol.

DOSAGE: As an emetic, 1 c.c. or 15 minims; as an expectorant, 0.05 c.c. or 1 minim.

Syrupus Ipecacuanhae.—Syrup of Ipecac, U. S. P.

One hundred c.c. represent 7 c.c. fluidextract of ipecac; it is approximately seven times the strength of the international standard syrup of ipecac.

DOSAGE: As an expectorant, 0.25 c.c. or 4 minims given every two or three hours; as an emetic, 15 c.c. or 4 fluidrams.

Jalap.—Jalap, U. S. P.

The dried tuberous root of *Exogonium Purga*. Frequently used in the form of powder.

ACTION AND USES: Jalap is a powerful purgative, producing copious watery evacuations. It is used for the purpose of removing water from the tissues in the treatment of dropsy.

DOSAGE: 1 gm. or 15 grains. It is best administered as:

Pulvis Jalapae Compositus.—Compound Powder of Jalap, U. S. P.

A mixture of jalap (35 gm.) and potassium bitartrate (65 gm.).

DOSAGE: 2 gm. or 30 grains.

LINIMENTA—LINIMENTS

Liniments are liquid preparations intended for external application.

For preparations included in this list see:

Linimentum Ammoniae, under Ammonia.

Linimentum Calcis, under Calx.

Linimentum Camphorae, under Camphora.

Linimentum Chloroformi, under Chloroformum.

Linimentum Saponis, under Sapo.

Linum.—Linseed, Flaxseed, U. S. P.

The ripe seed of *Linum usitatissimum*. Used extensively in the ground form for making cataplasms.

Oleum Lini.—Linseed Oil, U. S. P.

A fixed oil, expressed from linseed.

PROPERTIES: Linseed oil occurs as a yellowish, oily liquid, having a peculiar odor and a bland taste. It is soluble in absolute alcohol, but practically insoluble in water.

DOSAGE: 30 c.c. or 1 fluidounce.

ACTION AND USES: Linseed oil has the properties of other bland oils, but is seldom administered internally. For pharmaceutical purposes it is used in the making of lime liniment, and of soft soap.

LIQUORES—SOLUTIONS

Pharmaceutically, solutions are aqueous liquid preparations in which one or more substances are completely dissolved.

For preparations included in this list see:

Liquor Acidi Arsenosi, under Arseni Trioxidum.

Liquor Alumini Acetatis, under Alumini Acetas.

Liquor Ammonii Acetatis, under Ammonii Acetas.

Liquor Arseni et Hydrargyri Iodidi, under Arseni Trioxidum.

Liquor Calcis, under Calx.

Liquor Cresolis Compositus, under Cresol.

Liquor Formaldehydi, under Formaldehydum.

Liquor Magnesii Citratis, under Magnesii Citras.

Liquor Plumbi Subacetatis, under Plumbi Acetas.

Liquor Potassii Arsenitis, under Arseni Trioxidum.

Liquor Potassii Hydroxidi, under Potassii Hydroxidum.

Liquor Sodae Chlorinatae, under Calx Chlorinata.

Liquor Sodii Hydroxidi, under Sodii Hydroxidum.

Liquor Zinci Chloridi, under Zinci Chloridum.

Lobelia.—Lobelia, U. S. P.

The dried leaves and tops of *Lobelia inflata*. Occasionally administered in the form of powder.

ACTION AND USES: Lobelia has almost the same action as nicotine. Small doses stimulate, and large doses paralyze, the respiratory center. The vagus endings in the bronchial muscles are also depressed and the bronchi are relaxed. Small doses taken continuously cause a persistent quickening of the pulse; moderate doses produce nausea and large doses produce vomiting with marked and dangerous collapse.

Lobelia is no longer used as an emetic. As an expectorant it may be employed to increase bronchial secretion, but it must be used with caution. It is chiefly used for the treatment of bronchial asthma, in which it often renders distinct service. It should never be employed in asthma due to or associated with heart disease.

Tincture Lobeliae.—Tincture of Lobelia, U. S. P.

One hundred c.c. represent 10 gm. lobelia in diluted alcohol.

Dosage: From 0.5 c.c. to 1.5 c.c. or from 10 to 20 minims.

Ten minims may be given every fifteen minutes until distinct nausea is felt or relief is obtained.

Magnesii Carbonas.—Magnesium Carbonate, U. S. P.

A mixture of magnesium carbonate and magnesium hydroxid, approximately $(\text{MgCO}_3)_4 \cdot \text{Mg}(\text{OH})_2 \cdot 5\text{H}_2\text{O}$.

PROPERTIES: Magnesium carbonate is a light, white, friable mass, or a bulky, white powder, without odor, and having a slightly earthy taste. It is practically insoluble in water and in alcohol, but soluble with effervescence in dilute acids.

INCOMPATIBILITIES: It is incompatible with acids which form salts of magnesium.

ACTION AND USES: Magnesium carbonate when taken internally neutralizes the acids in the stomach. It may be used in cases of hyperacidity or acid gastritis, but it is sometimes objectionable on account of the carbon dioxid gas evolved. The salt formed is laxative. If magnesium carbonate passes the stomach without neutralization, it may escape solution in the intestines and not act as a laxative. Large doses sometimes produce an accumulation of the insoluble carbonate and may lead to intestinal obstruction. Magnesium carbonate is largely used as a dusting powder in intertrigo and in similar conditions, and as a cosmetic.

DOSAGE: 3 gm. or 45 grains.

Magnesii Citras.—Magnesium Citrate.

Liquor Magnesii Citratis.—Solution of Magnesium Citrate, U. S. P.

A solution containing 10 per cent. of a mixture of neutral and acid magnesium citrate.

The materials for making 360 c.c. or 12 ounces of the preparation are placed in a strong bottle, potassium bicarbonate being added last, and the bottle properly corked.

ACTION AND USES: Solution of magnesium citrate is given as a laxative.

DOSAGE: The average dose is one bottle containing 360 c.c. or 12 ounces, often given in divided doses.

Magnesii Oxidum.—Magnesium Oxid, U. S. P., Calcined Magnesia, MgO .

PROPERTIES: Magnesium oxid is a white, very bulky and very fine powder, without odor, and having an earthy but not a saline taste. It is almost insoluble in water and alcohol, but dissolves very readily in acids, forming salts of magnesium.

ACTION AND USES: Magnesium oxid is used mostly as an antacid to neutralize excessive acidity in the gastric juice. It is especially suitable for cases of hyperacidity accompanied by constipation. It is preferable in most cases to the carbonates or bicarbonates because it yields no gas on being neutralized. It is sometimes given in diarrhea with excessive acidity in children. It is an efficient antidote to the corrosive acids.

DOSAGE: From 0.6 to 3 gm. or from 10 to 45 grains; 2.5 gm. or 40 grains are equivalent to the average laxative dose of magnesium sulphate (15 gm. or 240 grains).

Magnesii Sulphas.—Magnesium Sulphate, U. S. P., $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$. "Epsom Salts."

PROPERTIES: Magnesium sulphate occurs as small, colorless prismatic needles or rhombic prisms, without odor, and having a cooling, saline and bitter taste. It is freely soluble in water, but practically insoluble in alcohol.

ACTION AND USES: Magnesium sulphate is one of the most active of the saline cathartics. When injected intravenously or intramuscularly it depresses the nervous and muscular structures, but the absorption from the alimentary canal is too slight to produce these effects. It has been injected for the purpose of producing spinal anesthesia, but its action is too uncertain and dangerous for use in man. It has been used in a few cases by injections in the subarachnoid space for the relief of tetanus.

Concentrated solutions of magnesium sulphate have been widely used as local applications in various inflammations, such as sprains, burns, erysipelas and the like, with asserted beneficial results.

DOSAGE: 15 gm. or 240 grains. Magnesium sulphate may be dissolved so that 1 c.c. of the solution contains 1 gm. of the salt, and of this solution 5 c.c. are given hourly until a laxative action is secured. This dose should be followed by sufficient water to dilute the salt.

Magnesii Sulphas Effervescens.—Effervescent Magnesium Sulphate, U. S. P.

A mixture containing magnesium sulphate (50 per cent.) with sodium bicarbonate, tartaric acid and citric acid.

DOSAGE: 15 gm. or 240 grains.

MASSAE—MASSES

Masses are soft solid preparations of such consistency that they can be made into pills readily.

For preparations included in this list see:

Massa Ferri Carbonatis, under Ferri Carbonas.

Massa Hydrargyri, under Hydrargyrum.

MELLITA—HONEYES

Honeys are sweet liquids having honey as a base; they formerly were much used as vehicles.

Mentha Piperita.—Peppermint, U. S. P.

The dried leaves and flowering tops of *Mentha piperita*.

Oleum Menthae Piperitae.—Oil of Peppermint, U. S. P.

A volatile oil distilled from the fresh or partly dried leaves and flowering tops of peppermint.

PROPERTIES: Oil of peppermint occurs as a colorless liquid, with the characteristic peppermint odor and a strongly aromatic, pungent taste, followed by a sensation of cold when air is drawn into the mouth.

ACTION AND USES: Oil of peppermint is used as an aromatic stimulant and carminative.

DOSAGE: 0.2 c.c. or 3 minims.

Spiritus Menthae Piperitae.—Spirit of Peppermint, U. S. P.

One hundred c.c. contain 10 c.c. oil of peppermint in alcohol, colored with the chlorophyll extracted from peppermint.

ACTION AND USES: See the oil.

DOSAGE: 2 c.c. or 30 minims.

Aqua Menthae Piperitae.—Peppermint water, U. S. P.

A saturated solution of oil of peppermint in distilled water.

ACTION AND USES: Peppermint water is frequently used as a vehicle or flavoring.

DOSAGE: 16 c.c. or 4 fluidrams.

Menthol.—Menthol, U. S. P.

A secondary alcohol, obtained from oil or peppermint, closely allied to camphor.

PROPERTIES: Menthol occurs as colorless, acicular or prismatic crystals, having a strong and pure odor of peppermint and a warm, aromatic taste, followed by a sensation of cold when air is drawn into the mouth. It is only slightly soluble in water, but freely soluble in alcohol, ether, liquid petrolatum, etc.

ACTION AND USES: Menthol has been used internally for the relief of gastric pain. Externally, it is applied on the skin as an anesthetic and cooling application. It is frequently used in the solid form known as Menthol pencils. One of these is rubbed over the painful part in neuralgia or headache.

It may also be applied in solution or ointment for the relief of itching, in a strength of from 1 to 2 per cent. It is used, dissolved in liquid petrolatum, as an application to inflamed mucous membranes, especially in the nose and throat. For this purpose a strength of 1 per cent. is suitable.

DOSAGE: 0.065 gm. or 1 grain.

Methylis Salicylas.—Methyl Salicylate, U. S. P. (Methyl Salicylas, Pharm. 1890).

An ester $C_6H_4OHCOOCH_3$, occurring in oil of birch and in oil of wintergreen, and also produced synthetically.

Oleum Betulae.—Oil of Betula, U. S. P., Oil of Birch.

Oleum Gaultheriae.—Oil of Gaultheria, U. S. P., Oil of Wintergreen.

Oil of birch is also sold commercially as oil of wintergreen.

PROPERTIES: Pure methyl salicylate is a colorless liquid, having a characteristic, strongly aromatic, wintergreen odor and a sweetish, warm and aromatic taste. It is nearly insoluble in water, but miscible in all proportions with alcohol. The oils obtained from natural sources frequently have a pinkish tint, because of the contamination with traces of iron, but in general have the same properties as methyl salicylate, to the requirements and tests for which they conform.

ACTION AND USES: Methyl salicylate and the oils containing it are antiseptic, and when absorbed by the skin or taken internally have the actions of salicylic acid or the salicylates (see sodium salicylate). When rubbed on the skin it is rapidly absorbed. It is frequently applied for the relief of pain in local rheumatic swellings or over inflamed nerves.

DOSAGE: 1 c.c. or 15 minims. When given internally this substance is preferably administered in the form of capsules, care being taken that the stomach is not empty and that plenty of water is taken. Locally, the oil may be applied pure or added to liniments.

MISTURAE—MIXTURES

Mixtures are liquid preparations containing insoluble or partly soluble medicinal substances suspended in them.

Morphina.—Morphine, U. S. P.

An alkaloid obtained from opium, the chief therapeutic constituent of the drug.

PROPERTIES: Morphin occurs as colorless or white crystals or a crystalline powder, odorless and having a bitter taste. It is very slightly soluble in water (1:3,300), but slightly soluble in alcohol (1:168).

INCOMPATIBILITIES: Morphin solutions are incompatible with alkalies, tannic acid, iodids and other precipitants of alkaloids.

ACTION AND USES: There are essentially three actions of morphin: a specific central analgetic action, a depressant action on the entire central nervous system, of a descending type, and a constipating effect resulting from a combination of central and local actions.

Morphin is practically devoid of local action, except on the gastro-intestinal tract. This local action is the subject of much debate, but it seems certain that it plays a part in the causation of the constipation which results from the administration of the drug. The drug has no local analgetic action and its use locally for the relief of pain is irrational.

The systemic actions of morphin are greatly dependent on the dose used. The smallest doses producing therapeutic effects result in the relief of pain; somewhat larger doses cause definite cerebral depression leading to more or less profound and prolonged sleep.

Some persons react peculiarly to morphin, showing one or more of the following symptoms: Cerebral excitation is,

perhaps, more common in women than in men, but it is usually mild and of short duration, soon giving place to the depressant action of the drug. Nausea and even vomiting not infrequently result from the systemic administration of a small dose. In some persons nausea is a very pronounced after-effect of the drug, lasting, at times, for hours.

The drug probably exerts a decided effect on the heart, through the vagus mechanism, chiefly influencing the rhythm, which may become irregular. The rate may be slowed considerably after large doses, but morphin does not endanger life through its cardiac action.

Morphin causes a marked constriction of the pupil when given in moderate doses, and this phenomenon is often used as a gage for the cessation of its administration in cases in which large doses are necessary. It has no local miotic action when dropped into the eyes.

The respiratory center is depressed by relatively small doses of morphin—such as are too small to be hypnotic. Use is made of this action in the treatment of persistent and troublesome cough, but it should be remembered that if the cough is "productive," the depression of the cough reflex may lead to a dangerous retention of the secretions of the inflamed mucosa.

Morphin is used chiefly as an analgetic in conditions of severe acute pain, and its use should be very guarded on account of the great danger of the formation of the morphin habit. In surgical conditions in which the alleviation of severe pain may obscure the course of the disease and lead to the unwarranted postponement of an operation, morphin should not be used, or only in very small doses and with circumspection. In chronic conditions associated with pain morphin should not be used, as the formation of the habit is almost certain to result from its prolonged administration. Exceptions to this generalization are to be found in such conditions as inoperable cancer, etc., in which the condition is hopeless and at the same time the cause of much suffering. Morphin should not be used for the relief of pain in persons of a neurotic or hysteric temperament, unless absolutely unavoidable. In general, it may be said that morphin should not be used for the relief of pain when any satisfactory relief can be obtained by the use of other drugs. Morphin, in moderate doses, is very useful in pain of cardiac origin. Morphin may be used to relieve the attacks of asthma and to lessen dyspnea from other causes, but caution should be exercised that the slowing of the respiration does not embarrass the heart. It should be used cautiously in the pain and dyspnea of uremia, as it interferes with elimination by the kidney and bowel.

Since the introduction of the coal-tar hypnotics and those of the chloral group, the use of morphin as a pure hypnotic has become exceptional.

Morphin, in the form of opium, is often given as a diaphoretic, the well-known Dover's powder being the preparation commonly used.

Habituation to morphin is readily established, and this habit is one of the most difficult to break. The indiscriminate use of morphin and preparations containing it by physicians is a common cause of the habit, and too great discrimination in its use can hardly be urged.

Overdoses of morphin lead to intoxication which may result fatally. The symptoms begin with the usual depression which deepens into sleep. The pupils become extremely constricted. Respiration becomes slow; the sleep deepens into coma from which the patient can be aroused with difficulty at first; later he cannot be aroused at all, and the respiration sinks to as low as three or four per minute. The heart is somewhat weakened and its rate is slowed. Death results from respiratory failure.

The treatment of morphin poisoning is not germane to this work, but its proper execution will often save a patient who is deemed hopeless.

Morphin is excreted largely through the alimentary tract, including the stomach. Some of the morphin thus excreted may be reabsorbed into the circulation; hence, in cases of poisoning, the use of frequent gastric lavage with permanganate of potassium, to destroy the morphin, is an important measure. Even after the hypodermic administration of the drug it is excreted by way of the gastro-intestinal tract.

Morphin is also used to lessen secretion and check peristalsis in diarrhea. For this purpose opium appears to be more efficient than morphin.

In the administration of morphin the danger of forming a habit should never be forgotten.

DOSAGE: 0.01 gm. or 1/6 grain. Smaller doses from 0.005 to 0.008, 1/12 to 1/8 grain, are often sufficient. The minimum fatal dose is 0.06 gm. or 1 grain.

Morphinae Hydrochloridum.—Morphine Hydrochloride, U. S. P.

PROPERTIES: Morphin hydrochlorid occurs as white crystals or as a crystalline powder, odorless and having a bitter taste. It is soluble in water (1:17) and in alcohol (1:42).

DOSAGE: 0.01 gm. or 1/6 grain.

Morphinae Sulphas.—Morphine Sulphate, U. S. P.

PROPERTIES: Morphin sulphate occurs as white, feathery crystals or in cubical masses, odorless, permanent in the air and having a bitter taste. It is soluble in water (1:15.3), but only slightly soluble in alcohol (1:465).

DOSAGE: 0.01 gm. or 1/6 grain.

Morrhuae Oleum.—Cod-Liver Oil, U. S. P.

A fixed oil obtained from the fresh livers of *Gadus morrhua*.

PROPERTIES: Cod-liver oil occurs as a pale yellow, thin, oily liquid having a peculiar, slightly fishy, but not rancid odor, and a bland, fishy taste.

ACTION AND USES: Cod-liver oil is generally considered to be tonic and nutrient, but its value probably depends entirely on its nutritive property. It is preferably admin-

istered in the form of a recently prepared emulsion, though many persons are able to take the oil as food in the same way that olive oil is used. Alcoholic preparations of the so-called active principles of cod-liver oil have no demonstrated therapeutic action.

MUCILAGINES—MUCILAGES

Mucilages are aqueous solutions of gums or the mucilaginous principles of vegetable substances.

For the preparations included in this list see:

Mucilago Acaciae, under Acacia.

Myrrha.—Myrrh, U. S. P.

A gum resin obtained from *Commiphora Myrrha*. Used occasionally in the form of powder, and also used as a carminative, in connection with aloes, in the form of pills.

ACTION AND USES: Myrrh is astringent and carminative. It is used largely as an ingredient in mouth washes and gargles for relaxed throat and spongy gums.

DOSAGE: 0.5 gm. or 7½ grains.

Tinctura Myrrhae.—Tincture of Myrrh, U. S. P.

One hundred c.c. represent 20 gm. myrrh in alcohol.

DOSAGE: 1 c.c. or 15 minims. For use in the mouth dilute with an equal volume of water.

Nux Vomica.—Nux Vomica, U. S. P.

The dried, ripe seed of *Strychnos Nux-vomica*, containing, when assayed by the process given in the U. S. P., not less than 1.25 per cent. of strychnin.

ACTION AND USES: The pharmacologic action of nux vomica is essentially the same as that of the strychnin which it contains. The preparations of nux vomica are used as stomachic tonics, and occasionally as respiratory and nerve stimulants. For the latter purpose the salts of strychnin are generally preferred. (See Strychnin Sulphate.)

Extractum Nucis Vomicae.—Extract of Nux Vomica, U. S. P.

An acetic acid extract of nux vomica purified by precipitating with alcohol and containing, when assayed by the process given in the U. S. P., 5 per cent. of strychnin.

DOSAGE: 0.015 gm. or ¼ grain, preferably given in the form of pills; 0.020 gm. or ⅓ grain contains 0.001 gm. or 1/60 grain of strychnin.

Tinctura Nucis Vomicae.—Tincture of Nux Vomica, U. S. P.

A solution of extract of nux vomica in a mixture of alcohol and water; it should assay 0.1 gm. of strychnin in 100 c.c.

DOSAGE: 0.6 c.c. or 10 minims.

When the tincture is used as an appetizer, its effect depends largely on the psychic stimulation produced by the bitter taste, and the dose may vary from 0.05 c.c. to 0.6 c.c. or from 1 to 10 minims. If it is desired to obtain the effect of strychnin, it is to be remembered that 0.6 c.c. or 10 minims contain approximately 0.6 mg. or 1/100 grain of strychnin.

OLEATA—OLEATES

The official oleates are solutions of alkaloids or metallic oxids in oleic acid.

OLEA PINGUA—FIXED OILS AND FATS

Fixed oils and fats are neutral esters of vegetable or animal derivation being compounds of acids (chiefly oleic, palmitic and stearic) with glycerol.

For fixed oils included in this list see:

Oleum Lini, under Linum.

Oleum Morrhuæ, under Morrhuæ, Oleum.

Oleum Ricini, under Ricini, Oleum.

Oleum Theobromatis, under Theobromatis, Oleum.

Oleum Tiglii, under Tiglii, Oleum.

OLEA VOLATILIA—VOLATILE OR ESSENTIAL OILS

Volatile or essential oils are liquids derived from plants and may contain or consist of neutral principles, aldehyds, ketones, phenols, esters or compound ethers.

For volatile oils included in this list see:

Oleum Betulæ, under Methylis Salicylas.

Oleum Caryophylli, under Caryophyllus.

Oleum Cinnamomi, under Cinnamomum.

Oleum Eucalypti, under Eucalyptus.

Oleum Gaultheriæ, under Methylis Salicylas.

Oleum Menthæ Piperitæ, under Mentha Piperita.

Oleum Rosæ, under Rosa.

Oleum Santali, under Santali, Oleum.

Oleum Sinapis Volatile, under Sinapis.

Oleum Terebinthinae, under Terebinthina.

OLEORESINAE—OLEORESINS

Oleoresins are thick liquid preparations consisting of volatile oils and resins extracted from vegetable substances by ether, acetone or alcohol.

For the preparation included in this list see:

Oleoresina Aspidii, under Aspidium.

Opium.—Opium, U. S. P.

The concrete, milky exudation obtained by incising the unripe capsules of *Papaver somniferum*.

ACTION AND USES: The action of opium closely resembles that of morphin. It, however, is absorbed more slowly and is more constipating and hence is preferred in the treatment

of the gastro-intestinal tract, especially for the purpose of checking diarrhea.

Opii Pulvis.—Powdered Opium, U. S. P.

Yielding, when assayed by the official process, not less than 12.5 per cent. of crystallized morphin.

DOSAGE: 0.065 gm. or 1 grain, containing about 0.008 gm. or about $\frac{1}{8}$ grain of morphin. Suppositories of opium have no advantage on account of local action.

Extractum Opii.—Extract of Opium, U. S. P.

Yielding, when assayed by the official process, 20 per cent. of crystallized morphin.

DOSAGE: 0.03 gm. or $\frac{1}{2}$ grain.

Tinctura Opii.—Tincture of Opium, Laudanum, U. S. P.

A solution of the soluble constituents of opium in diluted alcohol. One hundred c.c. contain not less than 1.2 and not more than 1.25 gm. crystallizable morphin.

DOSAGE: 0.5 c.c. or 8 minims.

Tinctura Opii Deodorati.—Tincture of Deodorized Opium, U. S. P.

Essentially a tincture of opium that has been deodorized by means of purified petroleum benzin.

DOSAGE: 0.5 c.c. or 8 minims.

Tinctura Opii Camphorata.—Camphorated Tincture of Opium, Paregoric, U. S. P.

Each 100 c.c. represent powdered opium (0.4 gm.), benzoic acid (0.4 gm.), camphor (0.4 gm.), oil of anise (0.4 c.c.), glycerol (4 c.c.), in diluted alcohol (to make 100 c.c.).

DOSAGE: 8. c.c. or 2 fluidrams, containing 0.006 gm. or $\frac{1}{10}$ grain of morphin.

Pulvis Ipecacuanhae et Opii.—Powder of Ipecac and Opium, U. S. P., Dover's Powder.

A mixture of ipecac (10 gm.), powdered opium (10 gm.) and sugar of milk (80 gm.).

ACTION AND USES: Dover's Powder is used chiefly for its diaphoretic effect.

DOSAGE: 0.5 gm. or $7\frac{1}{2}$ grains, containing 0.006 gm. or $\frac{1}{10}$ grain of morphin.

Oxygenium Compressum.—Compressed Oxygen, N. N. R.

Gaseous oxygen, O₂, in a compressed state.

PROPERTIES: Compressed oxygen occurs as a colorless, odorless and tasteless gas, slightly soluble in water and neutral to ordinary indicators. This gas is not inflammable, but supports combustion much more vigorously than does air.

ACTION AND USES: Compressed oxygen is administered for the purpose of relieving difficult respiration in cases of mechanical hindrance to the ingress of air to the lungs. It is also mixed with nitrogen monoxid as an anesthetic agent.

Pancreatinum.—Pancreatin, U. S. P.

A mixture of the enzymes naturally existing in the pancreas of warm-blooded animals, usually obtained from the fresh pancreas of the hog, *Sus scrofa*, or the ox, *Bos taurus*. Samples on the market contain much extraneous matter.

PROPERTIES: Pancreatin occurs as a cream-colored, amorphous powder, having a faint, peculiar, not unpleasant odor, and a somewhat meat-like taste. It is partially soluble in water and should contain not more than 10 per cent. of insoluble matter. It is practically insoluble in alcohol. Commercial samples commonly contain no steapsin, and other ferments may be present in traces only.

ACTION AND USES: Pancreatin is used chiefly for the predigestion of protein and starchy foods. Since it is destroyed by the action of the gastric juice, its use for the digestion of food in the stomach is illogical, if the stomach contains any acid. In cases of achylia gastrica in which no hydrochloric acid is secreted, it may be given to secure the digestion of the food in the stomach. In such cases it is well to administer a small amount of an alkali, such as sodium bicarbonate, in order to neutralize any acidity that may be present. In view of the poor quality of pancreatin on the market its use is not to be recommended.

The attempt is sometimes made to further the digestion of protein in the intestines by the administration of pancreatin in pills or capsules coated so as to prevent the action of the gastric juice. The drug may be used in this manner in cases in which it is believed that the secretion of the pancreas is lacking or deficient, but this method is not usually very successful.

DOSAGE: 0.5 gm. or 7½ grains.

Pancreatin may be administered internally in the form of salol-coated pills or in gelatin capsules that have been treated with formaldehyd.

Paraffinum.—Paraffin, U. S. P.

A mixture of solid hydrocarbons, obtained by chilling and pressing the higher distillates from petroleum and purifying the solid press-cake so obtained.

It is practically insoluble in water or in alcohol.

ACTION AND USES: Paraffin is used in surgery for prosthetic purposes. It is injected into the tissue to fill the place of parts which have been destroyed by trauma or pathologic processes. When used for this purpose, the melting-point of the product is important. A preparation melting at from 41 to 45 C. (105.8 to 113 F.) is most suitable. Its use for this purpose is not without danger.

Paraldehydum.—Paraldehyd, U. S. P.

Paraldehyd, $C_6H_{12}O_3$, is a polymer of acetaldehyd, CH_3CHO .

PROPERTIES: Paraldehyd is a colorless, transparent liquid, having a strong characteristic but not unpleasant or pungent odor, and a burning or cooling taste. It is soluble in water (1:8) and miscible in all proportions with alcohol.

ACTION AND USES: Paraldehyd is hypnotic and antispasmodic. It has no direct action on the circulation, except that it dilates the blood-vessels. It acts rapidly and produces a sleep which closely resembles natural sleep. It is very valuable when a rapid action is desired. The disagreeable odor of paraldehyd, persisting in the breath of the patient, interferes with its use in many cases. It may cause a habit very similar to that induced by alcohol.

DOSAGE: 2 c.c. or 30 minims. Preferably administered with cracked ice or ice-water.

Pelletierinae Tannas.—Pelletierine Tannate, U. S. P.

A mixture of the tannates of alkaloids obtained from *Punica Granatum*.

PROPERTIES: Pelletierin tannate occurs as a light yellow, odorless, amorphous powder, having an astringent taste and a weak acid reaction. It is only slightly soluble in water (1:235), but soluble in alcohol (1:12.6).

ACTION AND USES: Pelletierin tannate is used as an anthelmintic and teniafuge, especially in cases of cestode infection.

DOSAGE: 0.25 gm. or 4 grains. Preferably administered in the form of capsules. The alimentary canal should be emptied as completely as practicable by a mild purge (castor oil or a saline), a light diet on the previous day and fasting on the morning on which the anthelmintic is given. This remedy should be followed in two hours by a purge (castor oil). Not more than 0.3 gm. or 5 grains of pelletierin tannate should be given, as this amount has produced paralysis.

Pepsinum.—Pepsin, U. S. P.

A proteolytic ferment or enzyme obtained from the glandular layer of the fresh stomach of the hog, *Sus scrofa*.

PROPERTIES: Pepsin occurs as pale yellow, transparent or translucent scales or grains, or as a cream-colored, amorphous powder, free from any offensive odor and having a slightly acid or saline taste. It is soluble, or almost entirely soluble in water, the solution having more or less opalescence; it is practically insoluble in alcohol.

ACTION AND USES: Pepsin acts only in an acid medium. It is useful to secure the digestion of protein food in the stomach. It is seldom indicated, because the gastric juice usually contains sufficient pepsin to perform gastric digestion. It may be given in conjunction with hydrochloric acid in those cases of acute dyspepsia in which there is an absence of free hydrochloric acid in the stomach contents. In chronic cases it should be given only when the acid and pepsin are

both lacking. As a rule, even when acid is absent, the pepsin is still secreted and digestion will occur normally if hydrochloric acid is given without pepsin. In achylia gastrica, pepsin is a valuable addition to the hydrochloric acid, which is usually prescribed.

DOSAGE: 0.25 gm. or 4 grains.

Petrolatum.—Petrolatum, U. S. P.

A mixture of hydrocarbons, obtained by distilling off the lighter and more volatile portions from petroleum, and purifying the residue.

PROPERTIES: Petrolatum occurs as an unctuous mass, of about the consistence of an ointment, varying in color from white to dark amber.

ACTION AND USES: Petrolatum is used chiefly as a base for ointments. It is sometimes employed as a lubricant, for which purpose only a sterile product should be used.

Petrolatum Liquidum.—Liquid Petrolatum, U. S. P.

A mixture of hydrocarbons, obtained by distilling off most of the lighter and more volatile portions from petroleum, and purifying the liquid residue.

PROPERTIES: It occurs as a colorless or very slightly yellowish, oily, transparent liquid, without odor or taste, but giving off, when heated, a faint odor of petroleum. It is practically insoluble in water and only slightly more soluble in alcohol.

ACTION AND USES: Liquid petrolatum is used as a vehicle for medicinal agents for external application. It is also given internally, largely for its mechanical action as an emollient in constipation. It is not absorbed by the intestine and has no nutritive properties.

Phenol.—Phenol, U. S. P. (Acidum Carbolicum, Pharm. 1890)
 C_6H_5OH .

Hydroxybenzene, obtained from coal-tar by fractional distillation and subsequent purification or made synthetically.

PROPERTIES: Phenol should contain not less than 98 per cent. of C_6H_5OH . It occurs as colorless, interlaced or separate, needle-shaped crystals, having a characteristic somewhat aromatic odor. When copiously diluted with water it has a sweetish taste, with a slightly burning aftertaste, and, when undiluted, cauterizes and whitens the skin and mucous membranes. Phenol is soluble in water (1:20) and miscible with alcohol in all proportions.

ACTION AND USES: Phenol is antiseptic and germicide. A solution of 1:850 will prevent the multiplication of bacteria. A 1 per cent. solution will usually destroy non-sporulating bacteria in a few minutes at ordinary temperature, but a 5 per cent. solution fails to destroy anthrax spores after twenty-four hours' exposure.

Phenol is taken as the type or standard for comparing the activity of disinfectants. The phenol coefficient means the relative strength of a disinfectant, as compared with a solution of phenol acting on the same organism and for the same length of time.

Phenol is escharotic when applied to the skin, turning the skin and tissues white. If a finger or other extremity is surrounded with dressings wet even with a dilute solution of phenol, gangrene is liable to occur. Phenol acts as a local anesthetic. In consequence of this property it is employed to relieve itching. It may be used in the strength of 1 per cent. in solution or in ointment.

Internally, phenol has been used as an antiseptic in mouth-washes, gargles and sprays. It was formerly employed as an antiseptic in fermentation in the stomach, but it should not be used for this purpose. It is sometimes given with success to check obstinate vomiting. It is not so useful as an intestinal antiseptic as some other remedies, particularly salol.

The injection of phenol into the rectum for the destruction of parasites is dangerous and has sometimes resulted fatally.

Superficial burns from the action of phenol should be treated by the application of alcohol, glycerol, ether or oils to remove the poison.

In phenol poisoning the stomach may be washed out by diluted alcohol, which should be completely removed by washing with water. If left in the stomach the alcohol may favor the absorption of phenol.

Phenol Liquefactum.—Liquefied Phenol, U. S. P.

A liquid composed of not less than 86.4 per cent., by weight, of C_6H_5OH , and about 13 per cent., by weight, of water.

PROPERTIES: Liquefied phenol is a colorless liquid which may develop a slight reddish tint on keeping, having a characteristic, somewhat aromatic odor and the general chemical and physical characteristics of phenol.

DOSAGE: 0.05 c.c. or 1 minim.

Phenolphthalein.—Phenolphthalein, N. N. R.

A product of the interaction of phenol and phthalic anhydrid.

PROPERTIES: Phenolphthalein occurs as white, or grayish-white crystals or as an amorphous powder; slightly soluble in water and freely soluble in alcohol. Its solutions in acid liquids are colorless, but turn red when the liquid is made alkaline.

ACTION AND USES: Phenolphthalein acts as a purgative, but appears to possess no further physiologic action, except that it may cause some irritation of the rectum and lower bowel.

DOSAGE: The dosage must vary according to its effect, from 0.05 gm. to 0.5 gm., or from 1 to 8 grains. A case of mild poisoning is reported from taking 1 gm. (15 grains).

Phenylis Salicylas.—Phenyl Salicylate, U. S. P. (Salol, Pharm. 1890).

The salicylic ester of phenyl.

PROPERTIES: Phenylsalicylate occurs as a white crystalline powder, having a faint aromatic odor and a slight but characteristic taste. It is very slightly soluble in water (1:2,333) and freely soluble in alcohol (1:5).

ACTION AND USES: Phenyl salicylate is slightly antiseptic, but its antiseptic action is greatly increased when it is decomposed into its constituents, phenol and salicylic acid. It has very slight action in the mouth or stomach, but in the intestines it is decomposed into its constituents by the action of the fat-splitting ferment of the pancreas. After absorption, it produces the effect of salicylates, but if larger doses are given the phenyl may produce toxic effects. The urine is colored dark by the phenol excreted by the kidneys. If salol is to be used for its constitutional effects as a salicylate, it should always be remembered that about 40 per cent. of it consists of phenol.

Phenyl salicylate is commonly used as an intestinal antiseptic. It is regarded by many as the best available drug for this purpose, but it is the opinion of many physicians that little benefit is to be secured by the use of intestinal antiseptics. It is impossible in practice to check all putrefaction in the intestines, but the use of phenyl salicylate has been shown to lessen the products of putrefaction excreted in the urine.

Phenyl salicylate is used in diarrhea, enteritis and in the diseases believed to be due to intoxication by the products of intestinal putrefaction.

Salol is used in pharmacy for the purpose of coating enteric pills. It must be remembered that the coating of pills of moderate size entails the administration of a considerable dose of salol.

DOSAGE: From 0.2 to 0.5 gm., or from 3 to 8 grains.

It is preferably administered in the form of powder and may be inclosed dry in capsules or cachets. Pills and compressed tablets are objectionable, because of the tendency of the substance to fuse into a hard insoluble mass.

Phosphorus.—Phosphorus, U. S. P.

PROPERTIES: Phosphorus is a translucent, nearly colorless solid, of a waxy luster, having, at ordinary temperatures, about the consistency of beeswax. It has a distinctive and disagreeable odor and taste and should not be exposed to air. It is practically insoluble in water but slightly soluble in absolute alcohol (1:350). It takes fire readily when exposed to the air. Great care should be used in handling elementary phosphorus. It should be carefully kept under water in a moderately cool place. It should be cut, or otherwise divided, under water. It may be secured in the form of small particles by cautiously melting under water and shaking until cool.

ACTION AND USES: In small quantities phosphorus stimulates the growth of bone and has been asserted to have a stimulating action on the nervous tissues; the latter action, however, is doubtful. Somewhat larger doses produce a fatty degeneration of the various organs which is followed by a proliferation of the connective tissue.

Minute doses increase the number of red blood-cells, but do not increase the amount of hemoglobin. In larger doses phosphorus is an irritant poison, causing nausea, vomiting and sometimes diarrhea. If absorbed, the poison produces a wide-spread, fatty degeneration of the organs, but fre-

quently there are no symptoms for several days. This period may be followed by vomiting of blood, drowsiness, enlarged and painful liver and scanty urine. There are usually, in the latter stages, severe nervous symptoms, consisting of delirium, somnolence, coma and occasionally convulsions. In workers in phosphorus, chronic poisoning is shown by a necrosis of the lower jaw. It is believed that this necrosis is due to infection through carious teeth, which is favored by previous changes in the bone due to phosphorus.

Phosphorus was formerly used as a tonic or stimulant to the nervous system, but there is no evidence that it is of service in this way. Its use is gradually being abandoned. Its action on the blood is not regarded as justifying its use in anemia. The principal use of phosphorus is to further the deposition of calcium in growing bone, or in bones undergoing repair. For this purpose it may be given in rickets.

DOSAGE: 0.5 mg. or 1/125 grain. It may be administered either in the form of pills or as phosphorated oil (1:100).

Physostigma.—*Physostigma*, U. S. P. (Calabar bean).

The ripe seed of *Physostigma venenosum*. Used commonly in the form of salts of its principal alkaloid, physostigmin (eserin).

ACTION AND USES: *Physostigma* stimulates the peripheral endings of the autonomous nerves and probably acts on the musculature to enhance its response to normal stimulation. It also depresses the action of the spinal cord. It greatly increases intestinal peristalsis. It slows the pulse and after large doses increases the blood-pressure. It produces contraction of the pupil by local action. When instilled into one eye it causes contraction of the pupil on that side and not on the other.

Physostigma is used for its depressing action on the spinal cord in tetanus, strychnin poisoning and other forms of convulsions. It should be used as an adjuvant to more powerful measures and not relied on as the sole remedy.

Physostigma is used to stimulate peristalsis in the paralytic form of colic and some cases of chronic constipation. It may be employed to overcome acute obstipation, provided there is no inflammation and no mechanical obstruction.

Physostigmin is used in treatment of eye diseases for the purpose of contracting the pupil and also to reduce intra-ocular tension. For the latter purpose it is serviceable in glaucoma. It is a useful remedy in peripheral ulcer of the cornea. It may be employed to hasten recovery from the effect of mydriatics, such as homatropin.

Physostigminae Salicylas.—Physostigmine Salicylate, U. S. P., Eserin Salicylate.

PROPERTIES: Physostigmin salicylate occurs as colorless or faintly yellowish crystals, odorless and having a slightly bitter taste. It is soluble in water and in alcohol.

DOSAGE: 1 mg. or 1/60 grain.

It is used in the eye in solution of the strength of from 0.1 to 1 per cent. To avoid the irritation that it sometimes causes, it is preferable to give it in oily solution, or in conjunction with cocain. The following mixture may be used:

	gm.	or	c.c.	
R Cocainae hydrochlor.....	0		02	gr. $\frac{1}{4}$
Physostigminae sulph.....	0		01	gr. $\frac{1}{8}$
Aquae dest.	10		00	3 ij

For hastening recovery from a miotic a drop of 0.1 per cent. solution in castor oil may be employed.

Physostigminae Sulphas.—Physostigmine Sulphate, U. S. P., Eserin Sulphate.

PROPERTIES: This salt is rather more freely soluble than the salicylate, but is also readily decomposed by exposure to light and air.

DOSAGE: 1 mg. or 1/60 grain.

Pilocarpus.—Pilocarpus, U. S. P., Jaborandi.

The leaflets of *Pilocarpus Jaborandi* or of *Pilocarpus Microphyllus*.

Used chiefly in the form of a salt of its principal alkaloid, pilocarpin.

ACTION AND USES: Pilocarpin stimulates the oculomotor and other autonomous nerves. It produces excessive secretion of the salivary glands and also of the sweat glands. It stimulates the unstriated muscles of the body generally and the motor system of the intestines, and causes a partial spasm of the bronchial muscles. It causes a marked slowing of the pulse and a fall of blood-pressure due to the lessened rate of the heart, but the vagus stimulation is soon followed by depression with an accelerated pulse-rate.

It contracts the pupil and causes spasm of the muscles of accommodation by a peripheral action.

Pilocarpus is administered internally chiefly for its diaphoretic effect. In this action it is serviceable in certain diseases of the skin. Under its continuous use there may be a stimulation of the growth of hair in favorable cases. In doses just short of producing free diaphoresis it is sometimes of great benefit to relieve itching in generalized, acute eczema, urticaria, pruritus, etc.

In diseases of the eye such as glaucoma, corneal ulcer, etc., pilocarpin is employed as a weak miotic.

Pilocarpinae Hydrochloridum.—Pilocarpine Hydrochloride, U. S. P.

PROPERTIES: Pilocarpin hydrochlorid occurs as colorless or white transparent crystals, odorless and having a faintly bitter taste. It is very soluble in water and freely soluble in alcohol.

DOSAGE: From 0.001 to 0.01 gm. or from 1/60 to 1/6 grain.

Pilocarpinae Nitras.—Pilocarpine Nitrate, U. S. P.

PROPERTIES: This salt is less soluble in both water and alcohol than is the hydrochlorid and has no evident advantages over that salt.

DOSAGE: 0.01 gm. or 1/5 grain. Smaller doses of 1 mg. or 1/60 grain are frequently preferable.

Pilulae.—Pills.

Pills are globular, oval or flattened bodies, of such size and consistency that they can be swallowed whole.

For the preparations included in this list see:

Pilula Ferri Carbonatis, under Ferri Carbonas.

Pix Liquida.—Tar, U. S. P.

A product obtained by the destructive distillation of the wood of *Pinus palustris*.

PROPERTIES: Tar occurs as a semiliquid, viscid, blackish-brown product having an empyreumatic odor and a sharp and empyreumatic taste. Tar is miscible with alcohol and fixed or volatile oils but is only partially soluble in water.

ACTION AND USES: Tar is a mild irritant to the skin, and when taken internally stimulates the bronchial mucous membrane. It is employed in the form of syrup as a remedy for subacute bronchitis.

Externally it is employed as a mild stimulant application to the skin. It should be used as the first remedy in changing the treatment from sedative to stimulating applications. It is very useful as an antipruritic for the relief of itching in pruritis, eczema, etc.

DOSAGE: In beginning treatment with tar a dilute preparation should be used. The strength should be from 2 to 4 per cent., which may be tentatively increased. As an antipruritic in acute inflammatory conditions a strength of from 2 to 4 per cent. may be all that can well be used. In chronic cases it is used in the strength of from 6 to 20 per cent., and sometimes the pure tar is applied. For use as an antiseptic in skin affections the following formula may be used:

R	Picis liquidae	2	3 ℥j
	Potassii hydroxidi	1	3 j
	Aquae	8	f℥j

This must be diluted for use.

Unguentum Picis Liquidae.—Tar Ointment, U. S. P.

A mixture of tar (50 gm.) with yellow wax and lard sufficient to make 100 gm.

This should be diluted from 10 to 20 times before application. In chronic conditions, scaly eruptions, psoriasis and ringworm it may be applied in full strength.

Plumbi Acetas.—Lead Acetate, U. S. P., $\text{Pb}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 3\text{H}_2\text{O}$ (Sugar of Lead).

PROPERTIES: Lead acetate occurs as colorless or heavy white crystalline masses or granular crystals, having a faintly acetous odor and a sweetish, astringent, afterward metallic taste. It is freely soluble in water (1:2) and soluble in alcohol (1:30).

INCOMPATIBILITIES: Solutions of lead salts are incompatible with carbonates, hydroxids, iodids, chlorids and sulphates.

ACTION AND USES: The chief internal use of lead acetate is as an astringent in diarrhea. It is rarely used for this purpose for fear of lead poisoning. It may be combined with opium in the form of a lead and opium pill.

DOSAGE: 0.065 gm. or 1 grain.

Liquor Plumbi Subacetatis.—Solution of Lead Subacetate, U. S. P.

An aqueous liquid which contains in solution not less than 25 per cent. of lead subacetate, made by boiling a mixture of lead acetate, lead oxid and distilled water for half an hour.

PROPERTIES: Solution of lead subacetate occurs as a clear, colorless liquid, odorless, having a sweetish, astringent taste and an alkaline reaction.

DOSAGE: This preparation should be diluted from fifteen to thirty times before application. It is employed as an astringent and antipruritic in inflammatory conditions of the skin and as an application to sprains and bruises. It should not be applied to denuded surfaces and should be used with caution on the face.

Podophyllum.—Podophyllum, U. S. P.

The dried rhizome of *Podophyllum peltatum*. Used in medicine chiefly in the form of:

Resina Podophylli.—Resin of Podophyllum, U. S. P.

PROPERTIES: This preparation, popularly known as podophyllin, occurs as an amorphous powder, varying in color from grayish-white to pale greenish-yellow, turning darker when subjected to heat exceeding 35 C., or when exposed to light.

ACTION AND USES: Resin of podophyllum is very irritating to the mucous membrane, especially to that of the eyes. It has a slight, peculiar odor and a faintly bitter taste. It is very soluble in alcohol, but practically insoluble in water. It is used as a laxative and hydragogue cathartic, chiefly in the form of pills. It was formerly thought to exercise a special influence on the liver, but this idea has been abandoned. It is chiefly used in small doses for the treatment of chronic constipation.

DOSAGE: As a laxative in constipation small doses should be used, from 0.003 to 0.006 gm. or from 1/20 to 1/10 grain, once or twice daily. For a cathartic effect it may be given in doses of from 0.008 to 0.03 or from 1/8 to 1/2 grain.

Potassii Acetas.—Potassium Acetate, U. S. P., $KC_2H_3O_2$.

PROPERTIES: Potassium acetate occurs as a white powder or in crystalline masses, odorless, and having a saline taste; it is deliquescent on exposure to air. It is very soluble in water (1:0.4), and is freely soluble in alcohol.

ACTION AND USES: Potassium acetate is oxidized in the organism to potassium carbonate or bicarbonate. This tends to make the blood alkaline, and the salts excreted by the kidney render the urine less acid and, after large doses, alkaline. It is a marked diuretic, increasing the amount of urine and the solids of the urine without irritation of the kidney; consequently it is preferable to other diuretics in the treatment of nephritis.

An alkaline reaction favors oxidation; hence the potassium salts which yield the carbonate on oxidation increase the processes of metabolism. An increase in alkalinity is also believed to favor the production of immunity in infections, and potassium acetate and other salts of potassium which increase the alkalinity of the urine, are useful in infections, especially such as pneumonia and rheumatism. Whenever it is desirable to make the urine neutral or alkaline, potassium acetate affords a convenient means of obtaining the desired result. It is useful as a diuretic in acute nephritis.

Potassium salts are poisonous in excessive doses, but not enough can be administered by mouth to cause toxic effects.

DOSAGE: 2 gm. or 30 grains once in three hours, for a time.

The effect obtained can be determined by examining the reaction of the urine and regulating the dosage accordingly.

Potassii Bicarbonas.—Potassium Bicarbonate, U. S. P., KHCO_3 .

PROPERTIES: Potassium bicarbonate occurs as colorless, transparent crystals, odorless, having a saline and slightly alkaline taste; it is permanent in the air. It is freely soluble in water (1:3), but is practically insoluble in alcohol.

INCOMPATIBILITIES: It is incompatible with acids.

ACTION AND USES: Potassium bicarbonate is sometimes used to neutralize the acidity of the stomach, but sodium bicarbonate is usually preferred. It may be employed for the extemporaneous preparations of potassium acetate or potassium citrate. If a solution of acetic or citric acid or lemon juice is neutralized with potassium bicarbonate, an equivalent dose of potassium acetate or citrate is formed.

Potassium bicarbonate, even when neutralized in the stomach has a tendency to render the blood alkaline and to lessen the acid of the urine. It may be given to secure the same alkaline effects for which the acetates or citrates are usually employed.

Externally, potassium bicarbonate may be employed for its alkaline effects, for the purpose of softening the epidermis.

DOSAGE: 2 gm. or 30 grains.

Potassii Bitartras.—Potassium Bitartrate, U. S. P., $\text{KHC}_4\text{H}_4\text{O}_6$.

PROPERTIES: Potassium bitartrate occurs as a colorless or slightly opaque or white, somewhat gritty, powder, odorless and having a pleasant acidulous taste. It is slightly soluble in water (1:200) and nearly insoluble in alcohol.

ACTION AND USES: Potassium bitartrate is diuretic and aperient. It is used chiefly in combination with jalap as a hydragogue cathartic. See *Pulvis Jalapae Compositus* under Jalap.

DOSAGE: 2 gm. or 30 grains. Preferably administered in the form of a powder or as a "shake" mixture.

Potassii Bromidum.—Potassium Bromide, U. S. P., KBr.

PROPERTIES: Potassium bromid occurs as colorless or white cubical crystals or granular powder, odorless and having a very strongly saline taste. It is freely soluble in water (1:1½ parts), but only slightly soluble in alcohol (1:180 parts).

INCOMPATIBILITIES: It is incompatible with mineral acids or oxidizing agents and with salts of silver or lead.

ACTION AND USES: Potassium bromid is a nerve sedative. It diminishes reflex excitability and depresses the motor area of the cortex. In large doses it is depressant to the circulation. When continued long it disturbs the nutrition and may produce an irritation of the skin similar to that produced by the use of iodin.

Potassium bromid is used to relieve convulsions, either of cerebral or of spinal origin. For this reason it is given in epilepsy. Large doses are also given to relieve the spasms of tetanus.

Potassium bromid is also useful to quiet nervous excitability in neurasthenia and hysteria. It may be given as an adjunct to hypnotics, such as chloral. It is said to be of value for the prevention of seasickness.

DOSAGE: 1 gm. or 15 grains, preferably administered by itself in simple dilute solution.

The dose of potassium bromid may be increased to 5 gm. or more if a powerful action is indicated. When the remedy is long continued it should be interrupted occasionally in order that no accumulation in the system may occur.

Potassii Carbonas.—Potassium Carbonate, U. S. P., K_2CO_3 .

PROPERTIES: Potassium carbonate occurs as a white, granular powder, odorless and having a strongly alkaline taste. It is very soluble in water (1:0.9), but practically insoluble in alcohol.

ACTION AND USES: Potassium carbonate is antacid, but because of its strongly alkaline and caustic properties is rarely used internally unless largely diluted.

Potassii Chloras.—Potassium Chlorate, U. S. P., $KClO_3$.

PROPERTIES: Potassium chlorate is explosive when mixed with oxidizable matter, and caution should be observed in manipulating it. It occurs as colorless crystals or a white granular powder, odorless and having a cooling and characteristic taste. It is soluble in water (1:16), but nearly insoluble in alcohol.

ACTION AND USES: Potassium chlorate is astringent. Large doses are actively poisonous, causing disintegration of the blood-corpuscles. Potassium chlorate is much used as a mouth wash in various forms of stomatitis. It is also

employed as a gargle in the treatment of pharyngitis. Its value in these conditions, however, is uncertain. Its internal use is not to be recommended.

DOSAGE: A saturated solution may be used as a mouth wash or gargle.

Potassii Citras.—Potassium Citrate, U. S. P., $K_3C_6H_5O_7 \cdot H_2O$.

PROPERTIES: Potassium citrate occurs as transparent, prismatic crystals or a white granular powder, odorless, and having a cooling, saline taste. It is very soluble in water (1:0.5), but very slightly soluble in alcohol.

ACTION AND USES: Potassium citrate has the same action as potassium acetate. It is less readily absorbed and consequently, in large doses, is more laxative.

DOSAGE: 1 gm. or 15 grains.

Potassii Citras Effervescens.—Effervescent Potassium Citrate, U. S. P.

A mixture of potassium citrate (20 per cent.) with sodium bicarbonate, tartaric acid and citric acid.

DOSAGE: 4 gm. or 60 grains.

Potassii et Sodii Tartras.—Potassium and Sodium Tartrate, U. S. P., $KNaC_4H_4O_6 \cdot H_2O$.

Popularly known as Rochelle Salt.

PROPERTIES: Potassium and sodium tartrate occurs as colorless, transparent crystals or white powder, odorless and having a cooling saline taste. It is freely soluble in water (1:1.2), but practically insoluble in alcohol.

ACTION AND USES: Potassium and sodium tartrate is used as a saline cathartic.

DOSAGE: 8 gm. or 120 grains.

Pulvis Effervescens Compositus.—Compound Effervescing Powder, U. S. P.

Popularly known as Seidlitz powder. A mixture of sodium bicarbonate (31 gm.), potassium and sodium tartrate (93 gm.) and tartaric acid (27 gm.).

The sodium bicarbonate is mixed with the potassium and sodium tartrate and the mixture divided into twelve equal parts and each part is wrapped separately in a blue paper. Each blue paper contains $7\frac{3}{4}$ gm. of potassium and sodium tartrate. The tartaric acid is also divided into twelve parts and each part wrapped in a separate white paper.

DOSAGE: One set of two papers.

Potassii Hydroxidum.—Potassium Hydroxide, U. S. P., KOH.

PROPERTIES: Potassium hydroxid should contain not less than 85 per cent. of KOH. It occurs as dry, white or nearly white flakes, fused masses, or in pencils, hard and brittle, showing a crystalline fracture, odorless and having a very acrid and caustic taste. It rapidly destroys organic tissues and great caution is necessary in handling. It is very soluble in water and freely soluble in alcohol.

Liquor Potassii Hydroxidi.—Solution of Potassium Hydroxide, U. S. P.

One hundred c.c. contain approximately 5 gm. of potassium hydroxid.

ACTION AND USES: Solution of potassium hydroxid is antacid, but is seldom used internally, and when so employed should be largely diluted. Its action on the urine is similar to that of potassium carbonate, acetate or citrate. It is not regarded as so good a remedy as these.

Externally it may be used to soften the epidermis for the removal of warts, corns, etc.

DOSAGE: 1 c.c. or 15 minims.

Potassii Iodidum.—Potassium Iodide, U. S. P., KI.

PROPERTIES: Potassium iodid occurs as colorless or opaque white, cubical crystals, or granulated powder, having a peculiar, faint, iodine-like odor, and a pungent, saline, afterward bitter taste. It is very soluble in water (1:0.1) and soluble in alcohol (1:12).

INCOMPATIBILITIES: Potassium iodid is incompatible with mineral acids and oxidizing agents and should not be prescribed in solution with alkaloids or alkaloid-containing drugs.

ACTION AND USES: Potassium iodid is slightly irritating to the gastro-intestinal canal, especially the stomach. After absorption ordinary doses produce no symptoms. It probably undergoes some decomposition with the liberation of iodine. In large or moderate doses, long continued, it frequently produces symptoms of iodism. These are due to irritation of the nasal passages, the bronchi and the skin. The irritation of the nose is shown by coryza, and by pain in the region of the frontal sinus.

There is considerable increase of the bronchial secretions.

On the skin it produces various eruptions, generally of a papular character, which rarely become pustular. Eruption and inflammation of the skin may sometimes be so extensive as to produce constitutional depression.

In iodism there are often symptoms resembling those of exophthalmic goiter—tachycardia, tremor, nervous irritability, etc.

Potassium iodid is regarded as an alterative. It is used as an expectorant to increase secretion in the bronchi. It is of value in subacute and chronic bronchitis. The most remarkable action of iodids is the absorption of gummatous exudates in tertiary syphilis. The iodid does not destroy the spirochetes, but possibly causes the formation of a proteolytic ferment that has a selective action on the round cells composing the gumma. By promoting the absorption of cellular exudates in the walls of the blood-vessels, it exerts a beneficial influence in many cases of arteriosclerosis. There is evidence that the forms of arterial disease which are improved by iodids are of syphilitic origin. A similar absorptive action, almost specific, is seen after administration of

large doses of iodids in the granulomatous lesions of actinomycosis, sporotrichosis and blastomycosis.

Potassium iodid is used to promote the elimination of poisons, especially the poisonous metals such as mercury and lead.

DOSAGE: The dosage varies in non-syphilitic cases from 0.3 to 2 gm. or from 5 to 30 grains, but in cases of syphilis, especially in the tertiary form, much larger doses up to 5 gm. and more may be necessary. It is often advisable to begin with 5 drops of a saturated solution three times a day after meals and increase by 1 drop per day added to each dose. It should not be given on an empty stomach, but with some food, preferably liquid, such as milk. On the occurrence of symptoms of iodism, such as coryza, papular or pustular eruptions, rapid pulse, etc., the remedy should be discontinued and resumed in smaller doses after the disappearance of toxic symptoms.

Potassii Permanganas.—Potassium Permanganate, U. S. P., KMnO_4 .

PROPERTIES: Potassium permanganate occurs as slender prisms, of a dark purple color, almost opaque by transmitted light and of a blue metallic luster by reflected light, odorless and having a taste which is at first sweet but afterward disagreeable and astringent. It is soluble in water (1:15) and decomposes when brought into contact with alcohol.

ACTION AND USES: Potassium permanganate is deodorant, disinfectant and possibly emmenagogue. As an emmenagogue, potassium permanganate may be given in doses of from 0.03 to 0.06 gm. (from $\frac{1}{2}$ to 1 grain) three times a day.

It has marked oxidizing properties and is used for this purpose in the treatment of certain forms of poisoning when the poison is still in the stomach. In morphin poisoning the stomach should be washed with a solution of potassium permanganate (1:2,000). The same treatment is useful in phosphorus poisoning. In cases of snake bite it may be given by hypodermic injections of a solution (1:500) in the vicinity of the wound.

Potassium permanganate has been used to disinfect the hands in surgery. The hand is dipped into the disinfectant solution until it is a mahogany brown, and then dipped into a warm saturated solution of oxalic acid and rinsed in sterile water.

It is applied externally for excessive sweating of the feet.

DOSAGE: It is usually administered in the form of pills, being mixed with kaolin and massed with petrolatum. For application to the skin a solution of 1:500 may be used.

Protargol.—Protargol, N. N. R., Silver Proteinate.

This is a compound of albumin and silver containing 8.3 per cent. of silver in organic combination.

PROPERTIES: Protargol occurs as a light brown powder which is slowly but freely soluble in water (1:2). Protargol should be

protected from the light and its solutions should be made as required. It is precipitated by cocain hydrochlorid, but this may be prevented by the addition of boric acid.

ACTION AND USES: Protargol is a non-irritant bactericide and antiseptic. It may be employed in chronic gonorrhea as a non-irritant substitute for silver nitrate, and in diseases of the mucous membranes of the eye, ear, nose and throat as an antiseptic and astringent. It is largely used in the treatment of conjunctivitis.

DOSAGE: It is used chiefly in the form of from 0.25 to 1 per cent. solution for instillations or injections and 1:1,000 to 1:2,000 solution as irrigations. Also used in the form of tampons and bougies. Solutions are preferably made by sprinkling the protargol over the surface of the required amount of cold water and setting aside until solution occurs.

Prunus Virginiana.—Wild Cherry, U. S. P.

The bark of *Prunus serotina*, used in medicine principally in the form of:

Syrupus Pruni Virginianae.—Syrup of Wild Cherry, U. S. P.

Representing an aqueous extract of 15 per cent. of wild cherry bark.

INCOMPATIBILITIES: As it contains tannins it is incompatible with salts of iron.

ACTION AND USES: Syrup of wild cherry is tonic and sedative and largely used as a vehicle for cough medicines.

DOSAGE: 5 c.c. or 1 fluidram.

PULVERES—POWDERS

Pharmaceutically, powders are combinations of several substances in powdered form or powdered vegetable drugs. The term "powders" is also applied to single dose quantities of a drug or mixture of drugs in powdered form wrapped separately in "powder papers."

For the preparations included in this list see:

Pulvis Effervescens Compositus, under Potassii et Sodii Tartras.

Pulvis Glycyrrhizae Compositus, under Glycyrrhiza.

Pulvis Ipecacuanhae et Opii, under Opium.

Pulvis Jalapae Compositus, under Jalapa.

Pulvis Opii, under Opium.

Quinina.—Quinine, U. S. P.

An alkaloid obtained from the bark of various species of *Cinchona*.

PROPERTIES: Quinin occurs as white, flaky or microcrystalline powder, odorless and having a bitter taste. It is very slightly soluble in water (1:1,750), but very soluble in alcohol (1:0.6).

ACTION AND USES: Quinin is antimalarial, antipyretic, ecboic and is also used as a bitter tonic. Quinin is a protoplasmic poison, especially affecting the protozoa more

than bacteria. It is somewhat irritant to the stomach and intestines and when absorbed it causes ringing in the ears, but in moderate doses produces no other marked effects in healthy persons. In patients with fever it is antipyretic. Its solutions produce local anesthesia, especially the solution of quinin and urea hydrochlorid. In large doses it produces depression of the heart and respiration, and collapse. In toxic doses quinin may produce more or less complete hemianopia terminating in permanent loss of sight. Moderately large doses of quinin act as a stimulant to the uterine muscles, but do not produce such spasmodic contractions as ergot.

Quinin may be used as a tonic, like the simple bitters, for the improvement of digestion and nutrition.

Its chief use is as an antiperiodic in malaria. In this disease it should be given in large doses several hours before the time of the expected chill.

DOSAGE: 0.25 gm. or 4 grains. In malaria 0.6 gm. or 10 grains may be given at a dose. For ordinary use it is preferably administered in the form of capsules. For use as a bitter tonic it is given in solution. For its use as a local anesthetic see quinin and urea hydrochlorid. In medicine quinin is customarily used in the form of one of the official or unofficial salts.

The intensely bitter taste of quinin and its soluble salts, which constitutes the greatest objection to its use, especially with children, may be masked by administering the alkaloid or the insoluble tannate in syrup of glycyrrhiza or syrup of yerba santa.

Quininae Bisulphas.—Quinine Bisulphate, U. S. P.

The acid sulphate of the alkaloid quinin.

PROPERTIES: Quinin bisulphate occurs in colorless, transparent crystals or as small whitish needles, odorless and having a very bitter taste. It is freely soluble in water (1:8.5) and soluble in alcohol (1:18).

DOSAGE: 0.25 gm. or 4 grains.

Quininae Hydrochloridum.—Quinine Hydrochloride, U. S. P.

PROPERTIES: Quinin hydrochlorid occurs as white, silky, glistening needles, odorless and having a very bitter taste. It is soluble in water (1:18) and very soluble in alcohol (1:06).

DOSAGE: 0.25 gm. or 4 grains.

Quininae Sulphas.—Quinine Sulphate, U. S. P.

PROPERTIES: Quinin sulphate occurs as white, glistening crystals of prismatic needles, odorless and having a very bitter taste. It is only slightly soluble in water (1:720), but soluble in alcohol (1:86).

DOSAGE: 0.25 gm. or 4 grains.

Quininae Tannas.—Quinine Tannate, N. N. R.

The tannate of the alkaloid quinin, containing from 30 to 35 per cent. of quinin.

PROPERTIES: Quinin tannate occurs as an amorphous, pale lemon-yellow, odorless powder without taste, or at most slightly bitter, with scarcely an astringency. It is only slightly soluble in water, but freely soluble in alcohol (1:3).

DOSAGE: 0.5 gm. or 7½ grains.

Quininae et Ureae Hydrochloridum.—Quinine and Urea Hydrochlorid, N. N. R.

The compound of quinin hydrochlorid and urea hydrochlorid, containing approximately 60 per cent. of anhydrous quinin.

PROPERTIES: Quinin and urea hydrochlorid occurs as white, interlaced prismatic crystals, odorless and having a very bitter taste. It is freely soluble in water (1:1).

ACTIONS AND USES: Quinin and urea hydrochlorid has the actions of quinin. It is non-irritating when injected hypodermically. Recent investigations have shown that when injected hypodermically or when applied locally to mucous membranes, it exerts an anesthetic action similar to that of cocain. It is reported that the anesthesia is in some cases prolonged for several days.

Quinin and urea hydrochlorid is especially useful in the treatment of malaria by hypodermic injections. It has also been applied as a substitute for cocain in the production of local anesthesia.

DOSAGE: The same as quinin. For the production of local anesthesia, injection of a solution of from 0.25 to 1 per cent. strength is said to be free from the risk of producing fibrous indurations, which sometimes occur with the stronger solution. For application to mucous membranes solutions varying in strength from 10 to 20 per cent. should be used.

Resina.—Rosin, U. S. P.

The residue left after distilling off the volatile oil from turpentine.

PROPERTIES: Rosin occurs in sharp, angular fragments, amber-colored, usually covered with a yellowish dust and at ordinary temperatures readily pulverized. It is practically insoluble in water, but soluble in alcohol, ether and the fixed or volatile oils.

ACTION AND USES: Rosin is used as an ingredient of ointments and plasters designed to serve chiefly as stimulating applications to the skin, the simple rosin plaster being used as an adhesive plaster in minor surgery.

RESINAE—RESINS

Resins are usually made by distilling the volatile oil from natural oleoresins or by precipitating resins from alcoholic solutions by the addition of water.

For preparations included in this list see.

Resina, under Resina.

Resina Podophylli, under Podophyllum.

Resorcinol.—Resorcinol, U. S. P. (Resorcinum, Pharmacopeia, 1890).

The diatomic phenol, metadihydroxybenzene, $C_6H_4(OH)_2$.

PROPERTIES: Resorcinol occurs as colorless, needle-shaped crystals, having a faint, peculiar odor and a sweetish, followed by a bitter taste. It is very soluble in water (1:0.5) and in alcohol (1:0.4).

ACTION AND USES: Resorcinol is antiseptic, antizymotic and antipyretic. On account of the readiness with which it causes the formation of methemoglobin and the resulting danger of collapse it is seldom administered internally. It is sometimes given to check fermentation in the stomach. Applied externally it is astringent in from 1 to 3 per cent. solutions, and keratolytic in strong proportions, from 10 to 20 per cent.

DOSAGE: 0.125 gm. or 2 grains.

Rhamnus Purshiana.—Cascara Sagrada, U. S. P.

The dried bark of *Rhamnus Purshiana*.

ACTION AND USES: The preparations of cascara sagrada are laxative, acting mainly on the colon, and are widely used for habitual constipation. The dose can be gradually reduced without constipation following.

Fluidextractum Rhamni Purshianae.—Fluidextract of Cascara Sagrada, U. S. P.

One hundred c.c. represent 100 gm. of the drug in approximately 40 per cent. alcohol. The presence of a bitter principle renders this preparation useful as a tonic as well as a laxative.

DOSAGE: 1 c.c. or 15 minims, three times daily; smaller doses should be used if efficient for laxative effect.

Fluidextractum Rhamni Purshianae Aromaticum.—Aromatic Fluidextract of Cascara Sagrada, U. S. P.

One hundred c.c. represent 100 gm. of the drug, which has been deprived of its bitter principle and aromatized with glycyrrhiza and compound spirit of orange. This preparation is preferred as a laxative on account of its pleasant taste.

DOSAGE: From 0.6 to 2 c.c. or from 10 to 30 minims. The smaller dose may be given several times a day, the larger once daily at bedtime.

Extractum Rhamni Purshianae.—Extract of Cascara Sagrada, U. S. P.

A powdered extract representing four times its weight of the drug.

DOSAGE: From 0.1 to 0.5 gm. or from 2 to 8 grains.

Rheum.—Rhubarb, U. S. P.

The dried rhizome of *Rheum officinale* is occasionally administered either in the form of powder or in the form of "cubes" or "fingers."

ACTION AND USES: Rhubarb and the extract are cathartics, bitter tonics and stomachics, while the aromatic tincture and the aromatic syrup are laxative and to a slight extent astringent. As cathartics they act chiefly on the colon and have a tendency to produce constipation after the initial laxative effect. Hence they are an appropriate remedy in the beginning of diarrhea as they cause the expulsion of irritating substances and promote a return to normal by their constipating influence.

DOSAGE: 1 gm. or 15 grains.

Extractum Rhei.—Extract of Rhubarb, U. S. P.

A hydro-alcoholic extract of rhubarb evaporated to a pilular consistency.

DOSAGE: 0.25 gm. or 4 grains.

Tinctura Rhei Aromatica.—Aromatic Tincture of Rhubarb, U. S. P.

Aromatic tincture of rhubarb represents rhubarb (20 gm.), cinnamon (4 gm.), cloves (4 gm.) and myristica (2 gm.), in a mixture of glycerol, alcohol and water to make 100 c.c.

DOSAGE: 2 c.c. or 30 minims.

Syrupus Rhei Aromaticus.—Aromatic syrup of Rhubarb, U. S. P.

For all practical purposes this represents aromatic tincture of rhubarb (15 c.c.) with sufficient syrup to make 100 c.c.

DOSAGE: 8 c.c. or 2 fluidrams.

Ricini Oleum (Oleum Ricini).—Castor Oil, U. S. P.

A fixed oil expressed from the seed of *Ricinus communis*.

PROPERTIES: Castor oil occurs as a pale yellowish or almost colorless viscid liquid, having a faint, mild odor and a bland afterward slightly acrid and generally offensive taste. It is practically insoluble in water, but freely soluble in alcohol.

ACTION AND USES: Castor oil is used as a cathartic, irritating both the small and the large intestine, and locally, particularly in the eye, as a demulcent.

DOSAGE: 16 c.c. or 4 fluidrams.

Castor oil may be administered in the form of emulsion, Emulsum Olei Ricini, N. F., a 33 $\frac{1}{3}$ per cent. emulsion, or it may be given after wetting the mouth with some hot liquid and giving the oil floating in the liquid. For this purpose milk or tea may be used. Preparations are on the market in which the taste is largely concealed by the addition of saccharin and essential oils, the castor oil itself being made more limpid by the addition of alcohol. Suspended in soda water it is easily given to children.

Rosa.—Rose.

Rose is used in pharmacy chiefly in the form of:

Oleum Rosae.—Oil of Rose, U. S. P.

A volatile oil distilled from the fresh flowers of *Rosa damascena*.

Aqua Rosae.—Rose-Water, U. S. P.

A solution of the volatile oil of rose in water.

ACTION AND USES: Oil of rose and rose-water are used for flavoring.

Saccharum.—Sugar, U. S. P.

Refined sucrose obtained from various sources.

Official in the form of white, dry, hard, distinctly crystalline granules, odorless and having a purely sweet taste. Sugar is very soluble in water (1:0.5) and soluble in alcohol (1:138).

ACTION AND USES: In medicine it is used as a diluent, either as sugar or in the form of:

Syrupus.—Syrup, U. S. P.

An aqueous solution of sugar containing approximately 85 gm. of sugar in 100 c.c. of the liquid.

Saccharum Lactis.—Sugar of Milk, U. S. P., Milk-Sugar. Lactose.

A peculiar crystalline sugar obtained from the whey of cow's milk by evaporation, and purified by recrystallization.

PROPERTIES: Sugar of milk occurs as white, hard, crystalline masses or a white powder feeling gritty on the tongue, odorless and having a faintly sweet taste. It is freely soluble in water (1:5), but practically insoluble in absolute alcohol.

ACTION AND USES: Sugar of milk is used as a diluent, and also as a food, particularly in connection with modified milk for infants. It is also diuretic when given in large quantities.

SALES EFFERVESCENTES—EFFERVESCENT SALTS

Effervescent salts are mixtures of soluble salts with sodium bicarbonate and citric or tartaric acid, or a mixture of the two acids, designed to yield effervescent drafts when added to water.

For preparations included in this list see:

Magnesii Sulphas Effervescens, under Magnesii Sulphas.

Potassii Citras Effervescens, under Potassii Citras.

Pulvis Effervescens, under Potassii et Sodii Tartras.

Sodii Phosphas Effervescens, under Sodii Phosphas.

Salvarsan.—Salvarsan, N. N. R.

Also known as arsenobenzol, "606", and arsenophenolamin hydrochlorid, is 3-diamino-4-dihydroxyl-1-arseno benzene hydrochlorid. Corresponds to 31.57 per cent. arsenic (As).

PROPERTIES: Salvarsan occurs as a yellow, crystalline, hygroscopic powder, very unstable in air. It is soluble in water, yielding a solution with an acid reaction.

ACTION AND USES: Salvarsan is useful as a specific remedy for syphilis in all stages, but is the more efficient the more recent the infection. It is especially indicated in the primary stage; in the later stages it should be given in repeated doses, in conjunction with mercurial courses. In malignant syphilis, which resists mercury, it is often efficient.

Salvarsan is efficient in the various spirillar diseases such as relapsing fever, Vincent's angina, etc. In Vincent's angina local applications of the powder have been found useful, in addition to the intravenous administration. The drug administered intravenously cannot reach the spirilla embedded in the necrotic tissue of the throat.

It has been recommended as a substitute for arsenic in anemia, particularly pernicious anemia, and in diseases of the skin which are amenable to the action of arsenic. Its use in these conditions must be regarded as still experimental.

In certain cases salvarsan has produced toxic results which are equivalent to poisoning by arsenic. These have occurred more commonly after the intramuscular injections. The intramuscular injection is painful and is usually followed by a tender, inflammatory nodule, which persists for some time.

After intravenous injections certain nervous symptoms have frequently arisen which have received the name of neurorecidiv (nervous relapse). The evidence seems to show that these nervous conditions are due not to the action of salvarsan but to the increased activity of the spirochetes. They are best treated, therefore, by a specific remedy: another dose of salvarsan or a compound of mercury.

The optic neuritis which is so frequently produced by other preparations of arsenic has occurred very rarely in connection with salvarsan. The drug should be employed with great caution, however, if at all, in the presence of eye disease even when caused by syphilis.

DOSAGE: 0.5 gm. or $7\frac{1}{2}$ grains.

For subcutaneous and intramuscular injection a suspension in a neutral fluid is commonly employed. For intravenous injection a clear alkaline solution is used. The contents of a tube should be used at once after opening and under no conditions should the contents of a tube, damaged in transportation, or any remnants of the powder from previously used tubes, be employed.

The intravenous method is now recognized as the most suitable. Special care should be taken to see that the water used in making the solutions is freshly distilled and completely sterile. Various forms of apparatus have been

devised to facilitate intravenous injections. Of these, some form of graduated gravity pipet connected with a tube fitted with a three-way stop-cock seems most suitable.

Neosalvarsan is the name applied to a mixture of sodium 3-diamino-4-dihydroxy-1-arsenobenzene-methanal-sulphoxylate with inert, inorganic salts. The arsenic content of 3 parts of neosalvarsan is approximately equal to 2 parts of salvarsan. Solutions of neosalvarsan are more readily prepared; otherwise its actions and uses, as well as the precautions to be observed, are the same as for salvarsan.

Santali Oleum (Oleum Santali).—Oil of Santal, U. S. P.

A volatile oil distilled from the wood of *Santalum album*.

PROPERTIES: Oil of santal occurs as a pale yellow, somewhat thick liquid, having a peculiar aromatic odor, and a pungent, spicy taste. Readily soluble in alcohol, but practically insoluble in water.

ACTION AND USES: Oil of santal is a stimulant and antiseptic to the bronchial mucous membrane and especially to the mucous membranes of the genito-urinary tract. It is not employed in the treatment of respiratory diseases, but is used extensively in the treatment of gonorrhea. It is not suitable to the acute stages, but should be used in the subacute and chronic stages.

DOSAGE: 0.5 c.c. or 8 minims.

Santoninum.—Santonin, U. S. P.

The inner anhydrid or lactone of santonic acid obtained from *santonica*.

PROPERTIES: Santonin occurs as colorless, rhombic prisms, odorless and nearly tasteless when first put into the mouth, but afterward developing a bitter taste. It is very insoluble in water (1:5,300), and soluble in alcohol (1:35).

ACTION AND USES: Santonin is used for its poisonous action on intestinal parasites. It is seldom absorbed to a sufficient extent to produce symptoms. When this occurs, the chief effects are yellow vision and epileptiform convulsions.

It is used especially for its effect on roundworms. In proper doses it is a reasonably safe drug. It is frequently administered with calomel.

DOSAGE: 0.065 gm. or 1 grain.

As absorption is to be avoided the remedy is preferably administered in the form of coarse crystals, either inclosed in a capsule or made up into a lozenge rather than in finely powdered form. The soluble sodium santoninate should be avoided as a vermifuge.

Sapo.—Soap, U. S. P.

Officially directed to be prepared from sodium hydroxid and olive-oil (Castile soap).

PROPERTIES: Soap occurs as a white or whitish solid, hard yet easily cut when fresh, having a faint peculiar odor free from

rancidity, a disagreeable alkaline taste and an alkaline reaction. It is soluble in water and in alcohol.

Linimentum Saponis.—Soap Liniment, U. S. P.

A solution of soap, camphor and oil of rosemary in a mixture of alcohol and water.

ACTION AND USES: Soap liniment applied externally is stimulant and rubefacient, but is used more frequently as a vehicle for the local application of other more active counterirritants.

Sapo Mollis.—Soft Soap, U. S. P.

A soap prepared from potassium hydroxid and linseed oil.

PROPERTIES: Soft soap occurs as a soft, unctuous, yellowish-brown mass, having a characteristic odor and an alkaline taste.

ACTION AND USES: Soft soap is used principally as a detergent; being soluble in alcohol, and in mixtures of alcohol and water it is frequently used as a liquid soap. It has also been used to some extent as a vehicle for other more active medicaments to be applied in the form of an ointment.

Sarsaparilla.—Sarsaparilla, U. S. P.

The dried root of different species of smilax.

ACTION AND USES: Sarsaparilla has long been in use as an "alterative." Practically its only active ingredient appears to be a saponin-like body, and this has been taken advantage of quite extensively in connection with the compound syrup of sarsaparilla for the administration of castor oil at soda-water fountains and as a vehicle for other active medicaments.

Syrupus Sarsaparillae Compositus.—Compound Syrup of Sarsaparilla, U. S. P.

A mixture of fluidextract of sarsaparilla (20 c.c.), fluidextract of glycyrrhiza (1.5 c.c.), fluidextract of senna, (1.5 c.c.), and aromatics with syrup (sufficient to make 100 c.c.).

ACTION AND USES: Compound syrup of sarsaparilla is used as a vehicle.

DOSAGE: 16 c.c. or 4 fluidrams.

Scilla.—Squill, U. S. P.

The bulb of *Urginea maritima*. It is occasionally administered in the form of powder.

ACTION AND USES: The pharmacologic actions of squill resemble those of digitalis qualitatively, but the emetic action appears to be more prominent with squill, and this has led to its use as an emetic in the past, and it is still used as a nauseant, but it seems unwise to use a drug which affects the heart so powerfully when we have other nauseants, such as ipecac, which probably have less undesirable side actions.

Squill is used frequently to promote diuresis, which it probably does after the manner of digitalis, through its action on the heart.

Squill appears to be absorbed slowly from the gastrointestinal tract, but its emetic effects appear to be due to its action on vomiting center in the medulla, and its diuretic effects to be secondary to its action on the heart. Hence it is probable that digitalis could always be used in its place to induce an increased secretion of urine. It is quite possible that the acetic acid plays some part in the usefulness of syrup of squill in expectorant mixtures. Squill is much used as a nauseant expectorant in the first stages of laryngitis, bronchitis, etc.

DOSAGE: 0.125 gm. or 2 grains.

Tinctura Scillae.—Tincture of Squill, U. S. P.

One hundred c.c. represent 10 gm. of squill in approximately 70 per cent. alcohol.

DOSAGE: 1 c.c. or 15 minims.

Syrupus Scillae.—Syrup of Squill, U. S. P.

A mixture of a diluted acetic acid extract of 5 per cent. squill with syrup.

DOSAGE: 2 c.c. or 30 minims.

Scopolaminae Hydrobromidum.—Scopolamine Hydrobromide, U. S. P., Hyoscin Hydrobromid.

The hydrobromid of an alkaloid obtained from plants of the *Solanaceae*.

PROPERTIES: Scopolamin hydrobromid forms colorless crystals, odorless, having an acid, slightly bitter taste, freely soluble in water (1:15) and in alcohol (1:16).

INCOMPATIBILITIES: Scopolamin hydrobromid is incompatible with alkalies and other precipitants of alkaloids.

ACTION AND USES: Scopolamin resembles atropin in its influence on the nerve endings, but differs from it in having a sedative instead of a stimulating effect on the brain. It is used as a cerebral sedative in cases of mania and other forms of insanity, but must be employed with caution, as it sometimes induces a rapid fall in blood-pressure and collapse. It has been extensively used in conjunction with morphin for the production of surgical anesthesia, either as a preliminary to the use of ether or chloroform, or as the sole anesthetic. It is liable to produce dangerous depression of the respiration. Experience in these methods of anesthesia has not been satisfactory. It has been employed as a partial anesthetic in labor, but experience shows that the effect on the fetus is sometimes disastrous, many children being born dead or asphyxiated.

It is frequently used as a mydriatic and is regarded by some ophthalmologists as preferable to atropin because it

is less irritating, and produces a brief and complete cycloplegia.

DOSAGE: 0.5 mg. or 1/125 grain.

Senna.—Senna, U. S. P.

The dried leaflets of *Cassia acutifolia* (Alexandria senna), or of *Cassia angustifolia* (Indian or Tinnevely senna).

ACTION AND USES: Senna belongs to the anthraquinone-containing group of vegetable purgatives and is considered to be one of the more efficient drugs of this class. It is largely used for the treatment of chronic constipation.

DOSAGE: 4 gm. or 60 grains, administered either in the form of powder or as an infusion.

Fluidextractum Sennae.—Fluidextract of Senna, U. S. P.

A hydro-alcoholic extract, 100 c.c. of which represent 100 gm. of senna, previously treated with strong alcohol to remove the resin.

DOSAGE: 2 c.c. or 30 minims.

Syrupus Sennae.—Syrup of Senna, U. S. P.

Represents a mixture of fluidextract of senna (25 c.c.) with syrup (sufficient to make 100 c.c.).

DOSAGE: 4 c.c. or 1 fluidram.

See also Pulvis Glycyrrhizae Compositus, under Glycyrrhiza.

SERA ET VACCINA—SERUMS AND VACCINES, N. N. R.

The vaccines, viruses and serums constitute one of the most important groups of drugs with which the physician has to deal. Some preparations of this group are specific cures for certain diseases; others are invaluable in prophylaxis and diagnosis. The supervision of these drugs is in charge of the United States Public Health Service, which periodically makes inspections of laboratories licensed in accordance with the law passed by Congress in 1902. Antidiphtheric and antitetanic serums are required to conform strictly to the standards which have been established by the United States Public Health Service. There being no established standard for the various other products, they are not examined for their therapeutic value in the laboratory but are tested for the amount of preservative and freedom from bacterial and toxic contaminations. The preparations of this class most widely used are the following, which will be found in their respective alphabetical positions in this book:

Antidiphtheric Serum, under Serum Antidiphthericum.

Antitetanic Serum, under Serum Antitetanicum.

Bacterial Vaccines, under Vaccine.

Tuberculin, under Tuberculinum.

Vaccine Virus, under Virus Vaccinum.

Serum Antidiphthericum.—Antidiphtheric Serum, Diphtheria Antitoxin, U. S. P.

A fluid separated from the coagulated blood of the horse immunized through the inoculation of diphtheria toxin.

PROPERTIES: Antidiphtheric serum occurs as a yellowish or yellowish-brown, transparent or slightly turbid liquid, odorless or having a slight odor due to the presence of an antiseptic used as a preservative. The standard of strength, expressed in units of antitoxin power, must be that approved or established by the United States Public Health Service.

A modification of antidiphtheric serum in concentrated form, known as antidiphtheric globulins, is now being used extensively. This is made by the removal, by precipitation with neutral salts, of most of the constituents of the serum except that fraction of the globulins bearing antitoxic potency.

ACTION AND USES: Antidiphtheric serum neutralizes the toxin of diphtheria and is employed both as a curative and as a prophylactic agent in that disease.

DOSAGE: The dose of diphtheria antitoxin is measured by antitoxic units. From 500 to 1,000 units are commonly given as an immunizing or prophylactic dose. For curative treatment from 3,000 to 10,000 units may be given as an initial dose and repeated at intervals of from six to twenty-four hours, if no improvement results. In urgent cases it may be given intravenously.

Serum Antitetanicum.—Antitetanic Serum, N. N. R.

Antitetanic serum is the blood-serum of horses immunized to the toxin of the tetanus bacillus.

PROPERTIES: Antitetanic serum is marketed in both liquid or dry forms. Some manufacturers prepare an antitetanic globulin; this contains a solution of the globulins of the blood, which are soluble in a saturated sodium chlorid solution, together with the antitoxin, and contains the latter in concentrated form. Antitetanic serum must conform to the standard established by the United States Public Health Service.

ACTION AND USES: It is used both as a prophylactic and as a curative agent in tetanus. The dried product may be used as a dusting-powder in suspicious wounds.

DOSAGE: Immunizing: 1,500 units; in tetanus; from 3,000 to 20,000 units every four to eight hours. Single large doses by the vein are sometimes more effective.

Sinapis.—Mustard.

ACTION AND USES: Mustard is used extensively as a counterirritant. When mustard is mixed with water, the volatile oil is generated by the action of a ferment, myrosin, on a principle called sinigrin, contained in black mustard.

Sinapis Nigra.—Black Mustard, U. S. P.

The seed of *Brassica nigra*.

DOSAGE: As an emetic, 8 gm. or 120 grains.

Charta Sinapis.—Mustard Paper, U. S. P.

A coating of black mustard on rather thick, well-sized paper used in place of the domestic mustard poultice.

Oleum Sinapis Volatile.—Volatile Oil of Mustard, U. S. P.

A volatile oil obtained from black mustard by maceration with water and subsequent distillation, yielding, when assayed by the process given in the U. S. Pharmacopeia, not less than 92 per cent. of allyl iso-thiocyanate.

PROPERTIES: Volatile oil of mustard occurs as a colorless or pale yellow, limpid liquid, having a very pungent and acrid odor. Great caution should be exercised when smelling this oil, and it should not be tasted without being highly diluted. It is miscible with alcohol in all proportions, and is generally soluble in liquids containing alcohol.

ACTION AND USES: See Sinapis.

DOSAGE: 0.008 c.c. or $\frac{1}{8}$ minim.

Sodii Arsanilas.—Sodium Arsanilate, N. N. R., Atoxyl.

Sodium arsanilate is the sodium salt of arsanilic acid, prepared by condensing anilin and arsenic acid.

PROPERTIES: Sodium arsanilate occurs as white, odorless crystals, having a faint salty taste and being freely soluble in water (1:6). The aqueous solution, on standing, assumes a yellowish tint.

ACTION AND USES: Sodium arsanilate has been recommended for the conditions which are favorably influenced by arsenic. The arsenic of the acid is liberated very slowly in the system, producing the therapeutic effects of arsenic with a more continuous and less toxic action. Toxic effects have been frequently noted, however, and the use of all forms of sodium arsanilate has been followed by degeneration of the optic nerve, leading to blindness. This substance has been used extensively in the treatment of trypanosomiasis, but the frequent cases of blindness resulting have caused its use to be almost entirely abandoned.

DOSAGE: 0.08 gm. or $\frac{1}{2}$ grain, gradually increased.

Sodii Arsenas.—Sodium Arsenate, U. S. P., $\text{Na}_2\text{HAsO}_4 \cdot 7\text{H}_2\text{O}$.

PROPERTIES: Sodium arsenate occurs as colorless, transparent monoclinic prisms, odorless, and having a mild alkaline taste. It is freely soluble in water (1:12) and nearly insoluble in alcohol.

ACTION AND USES: Sodium arsenate has an arsenic action and has been employed in chronic skin diseases and in parasitic diseases of the blood, usually given in the form of solution or pill.

DOSAGE: 5 mg. or $\frac{1}{10}$ grain.

Sodii Benzoas.—Sodium Benzoate, U. S. P., $\text{NaC}_7\text{H}_5\text{O}_2$.

PROPERTIES: Sodium benzoate occurs as a white, amorphous powder, odorless, and having a sweetish astringent taste. It is freely soluble in water (1:1.6) and soluble in alcohol (1:43).

INCOMPATIBILITIES: It is incompatible with mineral acids and with ferric salts.

ACTION AND USES: Sodium benzoate has an internal action similar to that of sodium salicylate, but very much weaker, so that the latter salt is generally preferred.

DOSAGE: 1 gm. or 15 grains. It is usually administered in the form of solution.

Sodii Bicarbonas.—Sodium Bicarbonate, U. S. P., NaHCO_3 .

PROPERTIES: Sodium bicarbonate occurs as a white, opaque powder, odorless and having a cooling, mildly alkaline taste. It is soluble in water (1:12) at 15 C., practically without decomposition, but above this temperature the solution gradually loses carbon dioxide, and at the boiling point of water the salt is converted into normal carbonate; it is practically insoluble in alcohol.

INCOMPATIBILITIES: Sodium bicarbonate is incompatible with acids and acid salts, and the solution should be made with cold water, which tends to prevent its conversion into the more alkaline carbonate.

ACTION AND USES: Sodium bicarbonate is used to neutralize the acid of the gastric juice in hyperacidity, acid dyspepsia, etc. For this purpose it is to be preferred in cases which are not accompanied by constipation or in which there is diarrhea. It is sometimes used to render the urine alkaline in gravel, etc., but the citrate is preferable.

Large doses are given by the mouth in cases of acidosis in diabetes, fevers and similar conditions. Sodium bicarbonate may be used intravenously in cases of extreme acidosis. It may also be employed with sodium chlorid for saline infusions in cases accompanied by edema, although edema produced by large doses of alkali has been reported. It should never be injected hypodermically or intramuscularly as it may cause sloughing.

Externally it is used as a mild alkaline wash. Solutions of sodium bicarbonate are antipruritic. A weak solution is employed to soften impacted cerumen. Sodium bicarbonate may be employed to neutralize acetic or citric acids for the extemporaneous preparation of sodium acetate or citrate.

DOSAGE: 1 gm. or 15 grains. Much larger doses may be given in acidosis. As much as 60 gm. or 2 ounces may be given daily. It may be administered by a "drop enema" of a 4 per cent. solution. For intravenous injection a 6 per cent. solution sterilized by boiling and hence partly converted into the carbonate has been recommended. 1,000 c.c. of such a solution may be injected, but great care must be taken that none of the liquid gets outside the vein lest necrosis of the tissues occur.

Sodii Boras.—Sodium Borate, U. S. P., $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$.

Popularly known as borax and properly designated as sodium tetraborate.

PROPERTIES: Sodium borate occurs as colorless, transparent crystals or a white powder, inodorous, and having a sweetish alkaline taste. It is soluble in water, but practically insoluble in alcohol.

INCOMPATIBILITIES: Sodium borate is decomposed by mineral salts and is incompatible with mucilage of acacia, with the metallic salts of the mineral acids and with the salts of most alkaloids.

ACTION AND USES: Sodium borate is antiseptic, astringent and detergent and is used both externally and sometimes internally in the form of solution. The continuous ingestion of small doses produces a deleterious effect. Moderate to large doses may cause nephritis. Very large doses cause gastro-enteritis, nephritis, skin eruptions, visual disturbances, fall of temperature, collapse and a wide-spread fatty degeneration. It is used in from 1 to 2 per cent. solution as an eye-wash in hyperemia of the conjunctiva and catarrhal conjunctivitis.

For this purpose the following formula is very serviceable:

R	Sod. boratis	5	gr. x
	Aq. camph.	10	℥ ii
	Aq. dest. ad.	25	℥ j

It is sometimes used as an antipruritic solution:

R	Sodii boratis	1	gr. xv
	Glycerini	0	℥ x
	Aquae	30	℥ j

DOSAGE: 0.5 gm. or 7½ grains. It may be used as a gargle.

Sodii Bromidum.—Sodium Bromide, U. S. P., NaBr.

PROPERTIES: Sodium bromid occurs as colorless or white, cubical crystals, or a white, granular powder, odorless, and having a saline, slightly bitter taste. It is freely soluble in water (1:2) and soluble in alcohol (1:12.5).

ACTION AND USES: Sodium bromid is used as a nerve sedative and cerebral depressant and in ordinary doses its action and uses are identical with potassium bromid, but it is said to be less irritating. See Potassium Bromid.

DOSAGE: 1 gm. or 15 grains. It is administered in dilute aqueous solution.

Sodii Cacodylas.—Sodium Cacodylate, N. N. R.

Also known as sodium dimethylarsenate. It is the sodium salt of cacodylic acid (dimethyl arsenic acid).

PROPERTIES: Sodium cacodylate occurs as a white powder, very soluble in water. The aqueous solution is alkaline toward litmus paper, but nearly neutral toward phenolphthalein.

ACTION AND USES: The action of sodium cacodylate is similar to other arsenic compounds, but it is said to be much less toxic than the ordinary preparations of arsenic and is also less apt to cause undesirable side effects. This superiority is due to the slow liberation of the arsenous acid in the body.

The cacodylate has been particularly recommended in obstinate psoriasis, pseudoleukemia, diabetes, anemia, chlorosis, tuberculosis, malarial cachexia, etc.

As it is sometimes decomposed in the stomach, it is preferable to give it hypodermically.

DOSAGE: 0.03 gm. or $\frac{1}{2}$ grain. It may be administered hypodermically in aqueous solution or given by mouth in elixir or in the form of pills.

Sodii Carbonas Monohydratus.—Monohydrated Sodium Carbonate, U. S. P., $\text{Na}_2\text{CO}_3\text{H}_2\text{O}$.

The form in which sodium carbonate is now official.

PROPERTIES: Monohydrated sodium carbonate should contain not less than 85 per cent. of anhydrous sodium carbonate (Na_2CO_3). It occurs as a white, crystalline, granular powder, odorless and having a strongly alkaline taste. It is freely soluble in water (1:2.9) and practically insoluble in alcohol. Sodium carbonate, as now official, contains but one molecule of water of crystallization, and is nearly twice as strong in alkaline power as the ordinary crystalline carbonate, $\text{Na}_2\text{CO}_3\cdot 12\text{H}_2\text{O}$, commonly known as "sal soda."

INCOMPATIBILITIES: Sodium carbonate is incompatible with acids and acid salts and with the salts of the heavy metals and of alkaloids.

ACTION AND USES: Sodium carbonate is an antacid and is employed in medicine chiefly in the preparation of alkaline baths. For this purpose it may be used in the proportion of from 2 to 6 ounces for 30 gallons of water.

DOSAGE: 0.25 gm. or 4 grains.

Sodii Chloridum.—Sodium Chloride, U. S. P., Common Salt, NaCl .

PROPERTIES: Sodium chlorid occurs as a colorless or white crystalline powder, odorless and having a purely saline taste. It is freely soluble in water (1:2.8) and nearly insoluble in alcohol.

INCOMPATIBILITIES: Sodium chlorid is incompatible with salts of silver and lead.

ACTION AND USES: Sodium chlorid is used for preparing physiologic salt solution: 9 gm. to 1,000 c.c. of sterile water. This solution is frequently referred to as normal salt solution, but should not be confused with the chemical normal solution of sodium chlorid. When given by mouth in large doses sodium chlorid is emetic, and, in proper dilution, a prompt laxative.

DOSAGE: As an emetic, 16 gm. or 240 grains. As a laxative it is most conveniently given in 1 per cent. solution. The dose, 4 gm. or 60 grains, should be dissolved in from 0.5 to 1 liter (from 1 to 2 pints) of water and drunk on an empty stomach.

Sodii Hydroxidum.—Sodium Hydroxide, U. S. P., NaOH .

PROPERTIES: Sodium hydroxid should contain not less than 95 per cent. of NaOH and not more than 2 per cent. of other inorganic substances, with the exception of water. It occurs as dry, white or nearly white flakes, fused masses or translucent or

opaque white pencils, odorless, and having a caustic taste. Great caution is necessary in tasting and handling it, as it rapidly destroys organic tissue. It is very soluble in water (1:1) and in alcohol.

ACTION AND USES: Sodium hydroxid has properties closely resembling those of potassium hydroxid. It is used chiefly for pharmaceutical purposes, though occasionally it is used locally as a caustic or in dilute solution as an alkaline wash. Solution of sodium hydroxid largely diluted has also been administered internally.

Liquor Sodii Hydroxidi.—Solution of Sodium Hydroxide, U. S. P.

An aqueous solution containing about 5 per cent. of sodium hydroxid, NaOH.

DOSAGE: 1 c.c. or 15 minims.

Sodii Iodidum.—Sodium Iodide, U. S. P., NaI.

PROPERTIES: Sodium iodid occurs as colorless, cubic crystals, or as a white, crystalline powder, odorless and having a saline and slightly bitter taste. It is very soluble in water (1:0.5) and freely soluble in alcohol (1:3).

INCOMPATIBILITIES: Sodium iodid is incompatible with spirit of nitrous ether, bismuth salts, ferric salts, and the salts of many alkaloids.

ACTION AND USES: Sodium iodid has properties closely resembling those of potassium iodid, and like sodium bromid is perhaps less irritating than the potassium salt.

DOSAGE: 0.5 gm. or $7\frac{1}{2}$ grains. For ordinary cases the dose may range from 0.3 to 1.25 gm. or from 5 to 20 grains. When employed for the energetic treatment of tertiary syphilis it is sometimes necessary to increase the dose to from 2 to 5 gm. or from 30 to 75 grains or more. It should not be given on an empty stomach but should be administered with some form of food, preferably liquid, such as milk.

Sodii Nitris.—Sodium Nitrite, U. S. P., NaNO_2

PROPERTIES: Sodium nitrite should contain not less than 90 per cent. of NaNO_2 . It occurs as white, or nearly white, opaque, fused masses or pencils, or colorless, transparent crystals which are odorless and have a mild saline taste. It is freely soluble in water (1:1.1) and very slightly soluble in alcohol.

INCOMPATIBILITIES: It is incompatible with oxidizing agents generally. It must be protected from contact with the air on account of its tendency to oxidation.

ACTION AND USES: Sodium nitrite has the characteristic properties of the nitrites, and resembles nitroglycerin in its action, though its effect is manifested more slowly and is somewhat more lasting.

DOSAGE: 0.065 gm. or 1 grain. It should be administered in solution.

Sodii Phosphas.—Sodium Phosphate, U. S. P., $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$.

PROPERTIES: Sodium phosphate occurs as large, colorless, monoclinic prisms, or a granular, crystalline salt, odorless, and having a cooling saline taste. It is freely soluble in water (1:5.5), but practically insoluble in alcohol.

ACTION AND USES: Sodium phosphate is used as a saline cathartic, and its mode of action has much in common with magnesium sulphate and sodium sulphate. It is less disagreeable, but is less active. The acid phosphate NaH_2PO_4 is used for the purpose of rendering the urine acid. When the official phosphate is used for this purpose the equivalent amount of an acid, such as dilute hydrochloric or phosphoric, may be administered at the same time to convert it into the acid sodium phosphate.

DOSAGE: 2 gm. or 30 grains dissolved in warm water. To render urine acid, from 1 to 2 gm. or from 15 to 30 grains of acid sodium phosphate may be given every two or three hours, for five or ten doses until the urine is acid.

It may also be administered in liquid form by adding 4 parts of sodium nitrate, 13 parts of citric acid and a little water to 100 parts of sodium phosphate.

Sodii Phosphas Effervescens.—Effervescent Sodium Phosphate, U. S. P.

A mixture of exsiccated sodium phosphate, sodium bicarbonate, tartaric acid and citric acid, representing approximately 50 per cent. of sodium phosphate described above.

DOSAGE: 8 gm. or 120 grains.

Sodii Salicylas.—Sodium Salicylate, U. S. P., $\text{NaC}_7\text{H}_5\text{O}_3$.

PROPERTIES: Sodium salicylate occurs as white microcrystalline powder or scales, or as an amorphous, white powder, having not more than a faint pink tinge, odorless, and having a sweetish saline taste. It is very soluble in water (1:0.8) and freely soluble in alcohol (1:5.5).

INCOMPATIBILITIES: Sodium salicylate is incompatible with acids and acid salts and with solutions of most alkaloids, notably quinin, which precipitates as the salicylate.

ACTION AND USES: Sodium salicylate is the salt which is usually employed to secure the constitutional effects of salicylic acid. It is slightly antiseptic, but not so much as the free acid. It is irritant to mucous membranes, and on an empty stomach may cause pain and in large doses may produce vomiting. Large therapeutic doses produce a ringing in the ears, nausea, sometimes vomiting, occasional sweating and an increase in the amount of urine. It increases nitrogenous metabolism, and an increased amount of uric acid is excreted in the urine. In very large doses it may produce depression of the central nervous system, rarely convulsions, a slowing and depression of the respiration, and collapse from depression of the circulation. Large doses may produce abortion, and hence the drug is contra-indicated in pregnancy.

Sodium salicylate is an analgetic and is sometimes administered for the relief of headache or of neuralgic pains. It is chiefly used for its effect in articular rheumatism in which

it is highly efficacious. It promptly relieves all the local joint symptoms and the fever, but does not affect the endocarditis. Its effects last only while the medication is kept up. It is useful in tonsillitis and other catarrhal inflammations, but has not the decided action in the ordinary infections that it has in rheumatic fever. It is used in chorea. It stimulates the secretion of bile.

It produces much benefit in some forms of eye diseases, such as iritis, keratitis or glaucoma. It is of no value in gonorrheal arthritis or in arthritis deformans. It is of little value in gout.

DOSAGE: 1 gm. or 15 grains. The more efficient method is to repeat this dose every hour until salicylism occurs and then three times daily. It should be given in solution, but is sometimes administered in the form of powder inclosed in capsules or cachets, and followed by a sufficient amount of water to dilute it well in the stomach.

Sodii Sulphas.—Sodium Sulphate, U. S. P., $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$.

Popularly known as Glauber's Salt.

PROPERTIES: Sodium sulphate occurs as large, colorless prisms or granular crystals, odorless and having a bitter saline taste. It is freely soluble in water (1:3) and practically insoluble in alcohol.

ACTION AND USES: Sodium sulphate has long been used as a saline cathartic, but is less popular than magnesium sulphate or sodium phosphate.

DOSAGE: 16 gm. or 240 grains.

Sodii Sulphis.—Sodium Sulphite, U. S. P., $\text{Na}_2\text{SO}_3 \cdot 7\text{H}_2\text{O}$.

PROPERTIES: Sodium sulphite occurs as colorless, transparent crystals, odorless and having a cooling, saline, sulphurous taste. It is freely soluble in water (1:2), but nearly insoluble in alcohol.

INCOMPATIBILITIES: It is incompatible with mineral acids.

ACTION AND USES: Sodium sulphite is antiseptic and is principally used externally as a wash.

DOSAGE: It may be applied to the skin in parasitic skin diseases like scabies and favus in a solution of the strength of 1:10 or 1 dram to the ounce. A similar solution may be used as a mouth wash in aphthous stomatitis, mucous patches, or ulcers of the tonsils.

Sodii Thiosulphas.—Sodium Thiosulphate, U. S. P. (Sodii Hyposulphis, Pharm. 1890), $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$.

PROPERTIES: Sodium thiosulphate occurs as colorless, transparent monoclinic prisms, odorless and having a cooling, afterward bitter taste. It is very soluble in water (1:05) and practically insoluble in alcohol.

INCOMPATIBILITIES: Sodium thiosulphate is incompatible with acids, iodine and chlorine. It dissolves insoluble salts of silver.

ACTION AND USES: Sodium thiosulphate is used externally, in the form of a lotion, as an application for ring-worm and other parasitic skin diseases. It is extensively used in photography and has been given internally in place of sodium sulphite. It may be used to remove stains of iodine, silver nitrate, etc.

DOSAGE: 1 gm. or 15 grains. It may be used in aqueous solution or ointment of 10 per cent. strength or 1 dram to the ounce.

Sparteinae Sulphas.—Sparteine Sulphate, U. S. P.

The sulphate of an alkaloid obtained from *scoparius*.

PROPERTIES: Spartein sulphate occurs as colorless crystals, or a crystalline powder, odorless, and having a slightly saline and somewhat bitter taste. It is freely soluble in water (1:1) and in alcohol (1:2.5).

ACTION AND USES: Spartein has very little action on the central nervous system and experimentally, at least, has no pronounced action on the heart. It slows and weakens the heart. It is not a substitute for digitalis and apparently is of little use. It is not diuretic, but *scoparius* possesses this action from the scoparin which it contains.

DOSAGE: The dosage is variously given. The dose given by the Pharmacopeia is 0.01 gm. or 1/5 grain. Others recommend doses as high as 0.1 gm. or 1½ grains. It may be administered in the form of solution, powder or pill.

SPIRITUS—SPIRITS

Spirits are alcoholic solutions of volatile substances; either gases, liquids or solids.

For preparations included in this list see:

Spiritus Aetheris, under Aether.

Spiritus Aetheris Compositus, under Aether.

Spiritus Aetheris Nitrosi, under Aether Nitrosus.

Spiritus Ammoniae Aromaticus, under Ammonia.

Spiritus Camphorae, under Camphora.

Spiritus Chloroformi, under Chloroformum.

Spiritus Glycerylis Nitratis, under Glycerylis Nitrates.

Spiritus Menthae Piperitae, under Mentha Piperita.

Stramonium.—Stramonium, U. S. P.

The dried leaves of *Datura Stramonium*. Now seldom used in medicine except as an ingredient in so-called asthma powders for smoking. Its constituents and action are similar to those of belladonna.

Strophanthinum.—Strophanthin, U. S. P.

A glucosid, or mixture of glucosids, obtained from *strophanthus*.

PROPERTIES: Strophanthin occurs as a white or faintly yellowish powder, having an intensely bitter taste. Because of its

toxicity great caution should be used in tasting it. It is very soluble in water and in diluted alcohol, but less soluble in absolute alcohol.

ACTION AND USES: Strophanthin is used like digitalis. It acts more rapidly and is excreted sooner and hence is regarded as preferable for administration in acute conditions and in emergencies. It is not readily absorbed from the gastro-intestinal tract; hence its oral use is not recommended.

DOSAGE: 0.0003 gm. or 1/200 grain. It is best administered intramuscularly or intravenously, only a single dose being given daily, as a rule.

Strophanthus.—*Strophanthus*, U. S. P.

The ripe seed of *Strophanthus Kombé*. Occasionally administered in the form of powder.

ACTION AND USES: *Strophanthus* and tincture of *strophanthus* have properties similar to those of the glucosid *strophanthin*, though the therapeutic results from the administration of *strophanthus* by mouth are more variable than the results obtained from the hypodermic or intravenous administration of *strophanthin*.

Tinctura Strophanthi.—Tincture of *Strophanthus*, U. S. P.

One hundred c.c. represent 10 gm. *strophanthus* in approximately 65 per cent. of alcohol. It is nearly identical with the international standard tincture of *strophanthus*.

DOSAGE: 0.5 c.c. or 8 minims. For intramuscular or intravenous injection a much smaller dose should be given, not more than 0.12 c.c. or 2 minims, as a rule.

Strychnina.—Strychnine, U. S. P.

An alkaloid obtained from *nux vomica*, and also obtainable from other plants of the Loganiaceae.

PROPERTIES: Strychnin occurs as colorless, transparent crystals or a white crystalline powder, odorless and having an intensely bitter taste. It is very slightly soluble in water (1:6400) and soluble in alcohol (1:110).

INCOMPATIBILITIES: The salts of strychnin are incompatible with alkalies, alkali carbonates, iodids, bromids, arsenates and arsenites.

ACTION AND USES: Strychnin stimulates the reflex activity of the spinal cord, but produces little or no effect on the higher nervous centers. Both the sensory and motor centers are affected. Small doses increase the acuity of both sight and hearing. Large doses raise the blood-pressure by causing a constriction of the arterioles. It seems to have little direct effect on the heart. It stimulates the respiratory center, increasing the rapidity of the respirations. In poisonous doses it produces tonic convulsions similar to those of tetanus, but the trismus is less marked. Between the attacks there is usually complete relaxation of the affected

muscles. The convulsions are spinal, but not cerebral in origin. Death may occur during a convulsion from fixation of the chest by spasm of the respiratory muscles or during the interval from medullary paralysis.

Strychnin is used as a bitter tonic, generally in the form of a preparation of *nux vomica*. It is also serviceable as a tonic to the muscular system. It is chiefly employed as a respiratory stimulant in cases of depression of the respiratory center by narcotic or other poisoning. It is used for this purpose in several diseases of the respiratory organs, such as bronchitis, especially senile, pneumonia, etc.

Strychnin is much employed as a heart tonic, but in many cases improperly. It acts as a heart tonic chiefly by raising the blood-pressure. It should be used for its cardiac effect, if at all, in conditions of low blood-pressure. Thus it may be employed in pneumonia, diphtheria and other infections in which death results from the lowered blood-pressure. Even in these conditions its utility has been overestimated. It has been recommended in conditions of shock and collapse, but in the experience of some it is not effective in these conditions (Crile).

Strychnin is serviceable in some forms of paralysis. It is of no value when the paralysis results from an organic lesion and should not be used in conditions of inflammation of the nerve centers. It may be employed in paralysis due to poisons, such as lead, and in postdiphtheric paralysis. It is also of value in paralysis due to the cortical lesions, if used in moderate doses, to maintain the nutrition of the paralyzed muscles. In incontinence of urine, due to paresis of the vesical sphincter, it is useful, but if the incontinence depends on spasm atropin is more serviceable. Strychnin is useful in amblyopia, acting best in disturbance of vision unattended by changes visible with the ophthalmoscope, especially hysteric and neurasthenic forms. In lesions of the optic nerve it frequently produces an improvement in vision which, however, is only temporary. It is used in acute and chronic alcoholism. It is added to cathartics in the treatment of chronic constipation.

DOSAGE: From 0.0005 to 0.005 gm. or from 1/100 to 1/10 grain, three times a day or even as often as once in three hours.

In threatening respiratory or cardiac paralysis, strychnin must sometimes be used in quite large doses. In these cases it has been advised to push the remedy until twitching of the finger-tips occurs. Too large doses, however, should not be given, as it is possible to bring about a strychnin paralysis of the respiratory center. In such cases the strychnin should always be given hypodermically. In cardiac failure, strychnin, if given too frequently or in too large doses, may do more harm than good by causing irritability of the heart. In toxic amblyopia it is sometimes desirable to use large doses. It has been recommended to give as much as 1/13 grain per day (Nagel).

Strychninae Nitras.—Strychnine Nitrate, U. S. P.

PROPERTIES: Strychnin nitrate occurs as colorless, needle-shaped crystals, odorless, and having an intensely bitter taste. It is soluble in water (1:42) and slightly soluble in alcohol (1:120).

DOSAGE: 0.001 gm. or 1/60 grain.

Strychninae Sulphas.—Strychnine Sulphate, U. S. P.

PROPERTIES: Strychnin sulphate occurs as colorless or white prismatic crystals, or a white crystalline powder, odorless, and having an intensely bitter taste. Soluble in water (1:31) and in alcohol (1:65).

DOSAGE: 0.001 gm. or 1/60 grain.

Sulphonal.—See under Sulphonmethanum.**Sulphonethylmethanum.**—Sulphonethylmethane, U. S. P., Trional.

Official in most foreign pharmacopeias, and usually prescribed in this country as trional. Chemically, it is diethylsulphonemethylethylmethane.

PROPERTIES: Sulphonethylmethane occurs as colorless, odorless, crystalline scales, which have a bitter taste in aqueous solution. It is slightly soluble in water (1:195) and freely soluble in alcohol.

ACTION AND USES: Sulphonethylmethane is a hypnotic, producing, in ordinary doses, no other symptoms than sleep. The sleep comes on in about an hour after the medicine is taken, and is usually quiet. In some cases the sleep is not secured until the next day.

Sulphonethylmethane, when repeatedly taken, may produce poisoning in which hematoporphyrin appears in the urine, giving it a pinkish tinge or red color. The continued use of the remedy may lead to the formation of a habit. In addition to the excretion of hematoporphyrin in the urine there are lassitude, weakness, nausea and gastrointestinal disturbance as shown by diarrhea and constipation. More serious symptoms then develop, including abdominal tenderness, violent vomiting, ataxia, paresis of various muscles, loss of reflexes, and finally a condition of profound collapse. This condition ends in death in about 75 per cent. of the cases. There is a nephritis which involves the glomeruli and may be hemorrhagic in character.

Trional is used as a hypnotic in insomnia, but is of little use when the insomnia is accompanied by severe pain. It is also recommended as an antispasmodic in epilepsy, hiccough, chorea, etc.

DOSAGE: 1 gm. or 15 grains. Owing to its sparing solubility, it should be given with large quantities of hot liquids. It should not be used for more than two or three days consecutively. After this time recourse should be had to other hypnotics, if necessary. The possible appearance of hematoporphyrin, as indicated by pink color in the urine, should be watched for and the remedy suspended on its occurrence, but it is then often too late.

Sulphonmethanum.—Sulphonmethane, U. S. P.

Usually prescribed in this country as sulphonal. Chemically, it is diethylsulphonedimethylmethane.

PROPERTIES: Sulphonmethane occurs as colorless, inodorous and nearly tasteless crystals or a crystalline powder, slightly soluble in water (1:360), but soluble in alcohol (1:47).

ACTION AND USES: Sulphonmethane, or sulphonal, is hypnotic and sedative. See Sulphonethylmethane. It acts more slowly than trional and is best given about 5 p. m.

DOSAGE: 1 gm. or 15 grains, given in a hot liquid.

Sulphur.—Sulphur.

Sulphur is official in the U. S. Pharmacopeia under three headings:

Sulphur Lotum.—Washed Sulphur, U. S. P.

Sulphur Praecipitatum.—Precipitated Sulphur, U. S. P.

Sulphur Sublimatum.—Sublimed Sulphur, U. S. P.

PROPERTIES: Sulphur in any of the foregoing forms should contain from 99 to 99.5 per cent. of pure sulphur. It is a fine yellow powder, the sublimed variety having a slight odor and a faintly acid taste. The other varieties are without odor or taste and all are practically insoluble in water or in alcohol.

ACTION AND USES: Sulphur becomes active in the intestines and on the skin by a partial conversion into hydrogen sulphid or other sulphids. These products are slightly antiseptic and are irritant, so that sulphur becomes a mild purgative.

Sulphur is used externally as a parasiticide in cases of scabies. It is used to produce sulphur dioxid for room disinfection by burning it in the proportion of 3 pounds of sulphur to each thousand cubic feet of air space. All surfaces and articles to be disinfected should be wet.

DOSAGE: Each form, 4 gm. or 60 grains.

Unguentum Sulphuris.—Sulphur Ointment, U. S. P.

A mixture of washed sulphur (15 gm.) with benzoinated lard (85 gm.).

SUPPOSITORIA—SUPPOSITORIES

Suppositories are solid bodies intended to be introduced into the several natural orifices of the body for the purpose of producing systemic or local effects.

For the preparation included in this list see:

Suppositoria Glycerini, under Glycerinum.

Suppositories are usually made with oil of theobroma as a base. Rectal suppositories should be cone-shaped and should weight about 2 gm. or 30 grains.

Supracapsulin, Suprarenalin, L-Suprarenin Synthetic.—See Epinephrin.

SYRUPI—SYRUPS

Syrups are strong solutions of sugar and water with or without the addition of active medicaments.

For preparations included in this list see:

Syrupus, under Saccharum.

Syrupus Ferri Iodidi, under Ferri-Iodidum.

Syrupus Ipecacuanhae, under Ipecacuanha.

Syrupus Pruni Virginianae, under Prunus Virginiana.

Syrupus Rhei Aromaticus, under Rheum.

Syrupus Sarsaparillae Compositus, under Sarsaparilla.

Syrupus Scillae, under Scilla.

Syrupus Sennae, under Senna.

Syrupus Tolutanus, under Balsamum Tolutanum.

TABELLAE—TABLET-TRITURATES—COMPRESSED
TABLETS

Tablet-triturates are small disks made by diluting powdered medicaments with powdered sugar of milk or with powdered sugar, moistening the powder with sufficient alcohol to make a paste, and pressing into suitable molds. Compressed tablets are medicinal substances or mixtures of substances compressed to the form of disks.

Tannalbin.—See under Acidum Tannicum.

Terebinthina.—Turpentine, U. S. P.

A concrete oleoresin obtained from *Pinus palustris*.

Oleum Terebinthinae.—Oil of Turpentine, U. S. P.

A volatile oil recently distilled from turpentine.

PROPERTIES: Oil of turpentine occurs as a thin colorless liquid, having a characteristic odor and taste, both of which become stronger and less pleasant by age and exposure to air. Oil of turpentine is practically insoluble in water, but freely soluble in alcohol (1:3), and in all proportions of oil. For internal use the rectified oil of turpentine (oleum terebinthinae rectificatum) should be used.

ACTION AND USES: Turpentine is antiseptic, anthelmintic and diuretic. Applied externally it is rubefacient and counterirritant.

Turpentine has been used as an expectorant in cases of bronchitis characterized by free secretion. For this purpose it is now generally replaced by terpin hydrate. It is also given for the relief of flatulence and a small amount (from $\frac{1}{2}$ to 1 teaspoonful) may be added to enemas to increase their effectiveness.

Turpentine has been thought to be efficient in cases of internal hemorrhages, but this opinion is not well founded.

DOSAGE: 1 c.c. or 15 minims. It may be administered in the form of emulsion or in capsules.

Terpini Hydras.—Terpin Hydrate, U. S. P.

The hydrate of the diatomic alcohol terpin.

PROPERTIES: Terpin hydrate occurs as colorless, lustrous, rhombic prisms, nearly odorless, and having a slightly aromatic and somewhat bitter taste. It is slightly soluble in water (1:200) and freely soluble in alcohol (1:10).

ACTION AND USES: Terpin hydrate is antiseptic, diaphoretic and diuretic. It is largely used as an expectorant in cases accompanied by free bronchial secretion.

DOSAGE: 0.125 gm. or 2 grains. Terpin hydrate may be administered in the form of powder or in capsules. As a solution, usually in the form of elixir, it requires an excessive amount of alcohol.

Tetanus Antitoxin.—See under Serum Antitetanicum.

Theobromatis Oleum (Oleum Theobromatis).—Oil of Theobroma, U. S. P.

Popularly known as cacao butter, a fixed oil expressed from the roasted seeds of *theobroma cacao*.

PROPERTIES: Oil of theobroma occurs as a yellowish-white solid, having a faint, agreeable odor and a bland chocolate-like taste. It is freely soluble in ether, chloroform and benzene, soluble in absolute alcohol and insoluble in water.

ACTION AND USES: Oil of theobroma is used in pharmacy chiefly for the making of suppositories. Also used as a lubricant in massage, and as an application to sore nipples.

Theobromina.—Theobromin, N. N. R.

3,7-dimethyl-xanthin, a base occurring in *Theobroma cacao*, and other drugs, and also made synthetically. It is closely related to caffein (1,3,7-trimethyl-xanthin).

PROPERTIES: Theobromin occurs as a white, crystalline powder, odorless, and having a bitter taste. It is very slightly soluble in water, but is soluble in alcohol (1:100).

ACTION AND USES: Theobromin has an action on the kidneys and heart similar to that of caffein, but its effect on the central nervous system is less marked.

Theobromin is used as a diuretic in all forms of dropsy, but it is especially useful in renal dropsy. It is non-irritating to the kidney.

DOSAGE: 0.3 gm. or 5 grains.

Theobrominae Sodio-Salicylas.—Theobromin Sodium Salicylate, N. N. R., Diuretin.

A double salt of theobromin-sodium and sodium salicylate.

PROPERTIES: Theobromin sodium salicylate contains 50 per cent of theobromin and occurs as a white powder, odorless and having a saline taste. It is freely soluble in water (1:1), but is readily decomposed on exposure to carbon dioxide or by the action of acids, and must therefore be preserved in well-stoppered bottles.

INCOMPATIBILITIES: Theobromin sodium salicylate is incompatible with acids and has the other incompatibilities of salicylates.

DOSAGE: 0.5 gm. or 7½ grains, three times a day.

Thymol.—Thymol, U. S. P.

A phenol occurring in the volatile oil of *Thymus vulgaris* and in some other volatile oils.

PROPERTIES: Thymol occurs as large, colorless prisms, having an aromatic, thyme-like odor, and a pungent aromatic taste. It is only very slightly soluble in water (1:1,100), but very soluble in alcohol (1:0.9).

ACTION AND USES: Thymol is antiseptic and anthelmintic. As an anthelmintic it is used chiefly for the destruction of hookworm.

DOSAGE: 0.1 gm. or 2 grains.

In the treatment of hookworm disease it should be given in as finely divided state as possible in dosage of from 0.5 to 4 gm. or from 7½ to 60 grains. No fats, oils or alcohols should be given at the same time, for fear of absorption of the drug.

The dosage may be regulated according to age as follows: Up to 5 years of age, 0.5 gm.; up to 10, 1 gm.; up to 15, 1.5 gm.; up to 20, 2 gm.; above 20, 3 to 4 gm.

Thymolis Iodidum.—Thymol Iodide, U. S. P., Aristol.

Dithymol-di-iodid, popularly known as aristol, is obtained by the condensation of two molecules of thymol and the introduction of two atoms of iodine into the phenolic groups of the thymol.

PROPERTIES: Thymol iodid contains, when dried over sulphuric acid, 45 per cent. of iodine. It occurs as a bright, chocolate-colored, or reddish-yellow, bulky powder, with a very slight aromatic odor. It is practically insoluble in water and nearly insoluble in alcohol.

ACTION AND USES: Thymol iodid, or aristol, is antiseptic and is used in place of iodoform, chiefly as a dusting-powder. It is inferior to iodoform in antiseptic action and experience with it has been disappointing.

Tiglii Oleum (Oleum Tiglii).—Croton Oil, U. S. P.

A fixed oil expressed from the seeds of *Croton tiglium*.

PROPERTIES: Croton oil occurs as a pale yellow or brownish-yellow, somewhat viscid, and slightly fluorescent liquid, having a slight, fatty odor, and a mild, oily, afterward acrid and burning taste. It is practically insoluble in water, but when fresh is soluble in alcohol.

ACTION AND USES: Croton oil is a drastic cathartic and applied externally is rubefacient and vesicant. It is used mostly to procure prompt evacuation of the bowels, especially in coma.

DOSAGE: 0.05 c.c. or 1 minim, given in olive oil, butter, etc. It may be placed on the tongue in granulated sugar if the patient cannot swallow.

TINCTURAE—TINCTURES

Tinctures, with a few exceptions, are alcoholic or hydro-alcoholic extractive preparations of vegetable drugs; the

tinctures of potent drugs represent uniformly 10 gm. of drug in 100 c.c. of the preparation, while tinctures of less potent drugs vary in strength, but represent usually 20 gm.

For preparations included in this list see:

Tinctura Aconiti, under Aconitum.
 Tinctura Belladonnae Foliorum, under Belladonnae Folia.
 Tinctura Benzoini Composita, under Benzoinum.
 Tinctura Cannabis Indicae, under Cannabis Indica.
 Tinctura Capsici, under Capsicum.
 Tinctura Cardamomi, under Cardamomum.
 Tinctura Cinchonae, under Cinchona.
 Tinctura Cinchonae Composita, under Cinchona.
 Tinctura Colchici Seminis, under Colchici Semen.
 Tinctura Digitalis, under Digitalis.
 Tinctura Ferri Chloridi, under Ferri Chloridum.
 Tinctura Gentianae Composita, under Gentiana.
 Tinctura Hyoseyami, under Hyoseyamus.
 Tinctura Iodi, under Iodum.
 Tinctura Lobeliae, under Lobelia.
 Tinctura Myrrhae, under Myrrha.
 Tinctura Nucis Vomicae, under Nux Vomica.
 Tinctura Opii, under Opium.
 Tinctura Opii Camphorata, under Opium.
 Tinctura Opii Deodorati, under Opium.
 Tinctura Rhei Aromatica, under Rheum.
 Tinctura Scillae, under Scilla.
 Tinctura Strophanthi, under Strophanthus.
 Tinctura Valerianae Ammoniata, under Valeriana.
 Tinctura Zingiberis, under Zingiber.

Tragacantha.—Tragacanth, U. S. P.

A gummy exudation from *Astragalus gummifer*.

ACTION AND USES: Tragacanth swells on addition of water and gradually forms a cloudy gelatinous mass. On further addition of water it forms a mucilage which is occasionally used in pharmacy in the making of emulsions and is widely used as a basis for a greaseless catheter lubricant and an application for chapped skin.

TRITURATIONES—TRITURATIONS

Pharmacopœial triturations are active remedies diluted usually with 10 parts of sugar of milk.

Trional.—See under Sulphonethylmethanum.

TROSCISCI—TROCHES

Troches, or lozenges, are flat solid bodies intended to be dissolved in the mouth for their local effect on the mucous membrane of the mouth and the throat.

Tuberculinum.—Tuberculin, N. N. R.

This represents the toxins of the tubercle bacillus.

PROPERTIES: Tuberculin is marketed in a variety of forms, either as a filtered extract of the bacilli or as the pulverized insoluble substance of the dead bacilli themselves. In the latter or emulsified form tuberculin is known as tubercle vaccine, and it is closely related to bacterial vaccines mentioned under *Vaccina*.

ACTION AND USES: Tuberculin is used principally as a diagnostic agent and the characteristic properties of the trade product to be used should be studied closely before it is employed.

UNGUENTA—OINTMENTS

Ointments are soft, fatty solids of such consistency that they are readily spread at ordinary temperatures. When intended for systemic effect they are applied by inunction; ordinarily they are used as simple protectives.

For preparations included in this list see:

Unguentum Acidi Borici, under Acidum Boricum.

Unguentum Belladonnae, under Belladonnae Folia.

Unguentum Chrysarobini, under Chrysarobinum.

Unguentum Hydrargyri, under Hydrargyrum.

Unguentum Hydrargyri Ammoniatum, under Hydrargyrum Ammoniatum.

Unguentum Hydrargyri Dilutum, under Hydrargyrum.

Unguentum Hydrargyri Oxidi Flavi, under Hydrargyri Oxidum Flavum.

Unguentum Picis Liquidae, under Pix Liquida.

Unguentum Sulphuris, under Sulphur.

Unguentum Zinci Oxidi, under Zinci Oxidum.

Urotropin.—See under Hexamethylenamina.

VACCINA—BACTERIAL VACCINES, N. N. R.

Bacterial vaccines are suspensions of the killed bacteria in physiologic salt solution. Cresol is usually added as a preservative.

The use of many of these vaccines is in the experimental stage. They are often prepared from cultures obtained from the individual to be treated (autogenous vaccines); these usually give the best results, and some authors maintain that "stock" vaccines should be used only when it is impracticable to secure the autogenous agent. Bacterial vaccines are used to secure the production of an active immunity. Great care and skill are necessary for their proper use, and no definite statements as to dosage, etc., can be given.

Vaccine Virus.—See under Virus Vaccinum.

Valeriana.—Valerian, U. S. P.

The dried rhizome and roots of *Valeriana officinalis*.

ACTION AND USES: Valerian is thought to be antispasmodic and nerve sedative, but its influence is largely psychic, and in the ammoniated tincture the stimulating effects of the aromatic spirit of ammonia no doubt predominate.

Tinctura Valerianae, Ammoniata.—Ammoniated Tincture of Valerian, U. S. P.

One hundred c.c. represent the soluble constituents of 20 gm. of valerian in aromatic spirit of ammonia.

DOSAGE: 2 c.c. or 30 minims.

Veronal.—N. N. R.

Also known as diethyl-barbituric acid, diethyl malonyl urea and malourea, is 2,4,6-trioxy-5-diethyl pyrimidin, a ureid derived from diethylmalonic acid.

PROPERTIES: Veronal occurs as a white, crystalline powder, odorless, and having a faint bitter taste. It is slightly soluble in water (1:150) and freely soluble in alcohol (1:8).

ACTION AND USES: Veronal is quickly absorbed, especially when it is given in solution. In smaller doses it induces sleep, apparently without any other effect. It is a rather slow-acting hypnotic, the hypnotic action beginning in about half an hour after its administration. In larger doses the temperature falls and animals show marked trembling and restlessness in their sleep. A number of fatalities have followed its indiscriminate use.

DOSAGE: 0.3 to 0.6 gm. or 5 to 10 grains. It should never be given in the form of pills or tablets lest they fail to be absorbed.

Sodii Diaethyl-Barbituras.—Sodium diethyl-barbiturate, N. N. R.

Also known as medinal and veronal-sodium, is the monosodium salt of diethyl-barbituric acid.

PROPERTIES: Sodium diethyl-barbiturate occurs as a white, crystalline powder, odorless, and having an objectionably bitter alkaline taste. It is freely soluble in water (1:5).

ACTION AND USES: Sodium diethyl-barbiturate, veronal-sodium or medinal has the same properties as veronal, but acts more rapidly on account of its greater solubility.

DOSAGE: 0.3 to 0.6 gm. or 5 to 10 grains.

Viburnum Prunifolium.—*Viburnum Prunifolium*, U. S. P.

The dried bark of the root of *Viburnum prunifolium*.

ACTION AND USES: *Viburnum prunifolium* is thought to be a uterine sedative and tonic. Like valerian, however, its action, largely psychic, is probably due to its disagreeable odor and taste.

Fluidextractum Viburni Prunifolii.—Fluidextract of *Viburnum Prunifolium*, U. S. P.

One hundred c.c. represent the soluble constituents of 100 gm. *Viburnum prunifolium* in approximately 60 per cent. alcohol.

DOSAGE: 2 c.c. or 30 minims.

VINA—WINES

Medicated wines are solutions of active medicaments in wine or extractive preparations of vegetable drugs made with wine as a menstruum. Because of the great variation in the composition of wine it is unsatisfactory as a menstruum for extracting vegetable drugs, and medicated wines are in fact becoming obsolete.

Vinum Antimonii.—See under Antimonii et Potassii Tartras.

Virus Vaccinum.—Vaccine Virus, N. N. R.

Vaccine virus is the material obtained from skin eruptions of calves having vaccinia. The "pulp" is ground and mixed with varying percentages of glycerol. It is usually marketed in capillary tubes or as glycerinated points.

ACTION AND USES: Vaccine virus is used as prophylactic against small-pox.

Zinci Acetas.—Zinc Acetate, U. S. P., $\text{Zn}(\text{C}_2\text{H}_3\text{O}_2)_2, \text{H}_2\text{O}$.

PROPERTIES: Zinc acetate occurs as soft, white, pearly crystals, having a faintly acetous odor, and in dilute solutions an astringent, metallic taste. It is freely soluble in water (1:2.5) and soluble in alcohol (1:36).

INCOMPATIBILITIES: Zinc acetate is incompatible with alkali carbonates or sodium borate.

ACTION AND USES: Zinc acetate is astringent and antiseptic. It is used principally in the form of solution as an external application, for injection, solution or collyria; rarely used internally.

DOSAGE: 0.125 gm. or 2 grains.

It may be used in collyria in the proportion of from 0.1 to 0.5 per cent. or from $\frac{1}{2}$ to 2 grains per ounce.

Zinci Chloridum.—Zinc Chloride, U. S. P., ZnCl_2 .

PROPERTIES: Zinc chlorid occurs as a white or nearly white granular powder or in porcelain-like masses, irregular or molded into pencils, odorless and of such intensely caustic properties as to make tasting dangerous, unless freely diluted with water. The dilute solution has an astringent metallic taste. It is very soluble in water (1:0.5) and in alcohol.

INCOMPATIBILITIES: Zinc chlorid is incompatible with sodium borate and with alkali carbonates.

ACTION AND USES: Zinc chlorid is used as an antiseptic, astringent and escharotic.

Liquor Zinci Chloridi.—Solution of Zinc Chloride, U. S. P.

An aqueous solution containing about 50 per cent., by weight, of zinc chlorid made by dissolving granulated zinc in hydrochloric acid.

Zinci Oxidum.—Zinc Oxide, U. S. P., ZnO .

PROPERTIES: Zinc oxid occurs as a very fine, amorphous, white or yellowish-white powder, free from gritty particles, without odor or taste. It is practically insoluble in water and in alcohol.

INCOMPATIBILITIES: Zinc oxid is incompatible with acids.

ACTION AND USES: Zinc oxid is antiseptic and astringent and is widely used either alone or in combination with boric acid, bismuth subnitrate and starch as a dusting-powder and as a sedative in ointment for a variety of skin diseases. By some it is thought to be a nervine, but this is probably incorrect. It is now rarely used internally.

DOSAGE: 0.25 gm. or 4 grains.

Unguentum Zinci Oxidi.—Ointment of Zinc Oxid.

A mixture of zinc oxid (20 gm.) with benzoinated lard (80 gm.).

Zinci Stearas.—Zinc Stearate, U. S. P.

PROPERTIES: Zinc stearate occurs as a very fine, white powder, tasteless, and having a very faint odor, resembling that of fat. It is practically insoluble in water and alcohol.

ACTION AND USES: Zinc stearate is used as a dusting-powder, but appears to have little or no advantage over Zinc Oxid.

Zinci Sulphas.—Zinc Sulphate, U. S. P., $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$.

PROPERTIES: Zinc sulphate occurs as colorless, transparent, rhombic crystals, or granular crystalline powder, without odor, and having an astringent, metallic taste. It is very soluble in water, but practically insoluble in alcohol.

INCOMPATIBILITIES: Alkali carbonates, sodium borate, tannin and vegetable astringents.

ACTION AND USES: Zinc sulphate is astringent, styptic and emetic. It is much used in collyria in conjunctivitis and is especially effective in that form caused by Morax-Axenfeld bacillus. It is employed as an injection in the treatment of the chronic stages of gonorrhea.

DOSAGE: As an emetic, 2 gm. or 30 grains.

In collyria it may be used in the strength of from 0.1 to 1.0 per cent. or from 1 to 5 grains per ounce. As an injection in gonorrhea solutions varying in strength, from 0.5 to 4 per cent. may be used.

Zingiber.—Ginger, U. S. P.

The dried rhizome of *Zingiber officinale*. Occasionally administered in the form of powder.

ACTION AND USES: Ginger is aromatic, stimulant and a stomachic tonic. It is a servicable carminative in colic.

DOSAGE: 1 gm. or 15 grains.

Tinctura Zingiberis.—Tincture of Ginger, U. S. P.

One hundred c.c. represent the soluble constituents of 20 gm. ginger in alcohol.

DOSAGE: 2 c.c. or 30 minims.

TABLE SHOWING THE RECORDED SOLUBILITY OF SUBSTANCES INCLUDED IN THE LIST OF IMPORTANT MEDICAMENTS

Abbreviations and signs used:

dec. = decomposed; 00 = sparingly soluble or slightly soluble; 000 = insoluble; v. s. = very soluble; misc. = miscible in all proportions; — = unrecorded or uncertain.

The solubility values are for distilled water at approximately 25 C. and for the official U. S. P. alcohol at the same temperature and indicate the number of parts of the solvent required to dissolve 1 part of the substance.

Substance Soluble in—	Parts of Cold Water.	Parts of Cold Alcohol.
Acetanilidum	179	2.5
Acetphenetidinum	925	12.0
Acidum Benzoicum	281	1.8
Acidum Boricum	18	15.3
Acidum Citricum	0.54	1.55
Acidum Hydrochloricum	misc.
Acidum Hydrocyanicum Dilutum.	misc.
Acidum Nitricum	misc.
Acidum Salicylicum	308	2.0
Acidum Tannicum	0.34	0.23
Acidum Tartaricum	0.71	1.67
Adeps	000	00
Æther	10	misc.
Ætheris Nitrosi, Spiritus	misc.	misc.
Æthylis Chloridum	00	misc.
Æthyl-Morphinae Hydrochloridum	7	2
Aloinum	120	10.50
Alumen	9	000
Alumen Exsiccatum	17	000
Alumini Acetatis, Liquor	misc.
Ammonii Acetatis, Liquor	misc.	misc.
Ammonii Carbonas	4	dec.
Ammonii Chloridum	3	80
Amylis Nitris	00	misc.
Antimonii et Potassii Tartras...	15.5	000
Antipyrina	v.s.	1
Apomorphinae Hydrochloridum...	39.5	38.2
Argenti Nitras	0.54	24.0
Argenti Proteinæ	2
Arseni Trioxidum	100	00
Aspirinum	300	5
Atropina	450	1.46
Benzosulphinidum	250	25
Betanaphthol	950	0.61
Bismuthi Subcarbonas	000	000
Bismuthi Subgallas	000	000
Bismuthi Subnitras	00	000
Bismuthi Subsaliçylas	00
Caffeina	45.6	53.2
Caffeina Citrata	25.0	20.0
Caffeinae Sodio-Benzoas	2	30
Calci Carbonas Præcipitatus...	00	000
Calx	760	000
Camphora	700	v.s.
Chloralum Hydratum	0.3	0.2
Chromii Trioxidum	0.5	dec.

Substance Soluble in—	Parts of Cold Water.	Parts of Cold Alcohol.
Chrysarobinum	4,812	308
Codeina	120	1.6
Codeinæ Phosphas	2.25	261
Codeinæ Sulphas	30	1,035
Creosotum	140	misc.
Cresol	60	misc.
Cresolis Compositus, Liquor....	misc.	misc.
Creta Præparata	00	000
Cupri Sulphas	2.2	400
Diacetylmorphinæ Hydrochloridum	2.0	11
Elaterinum	000	262
Epinephrina	000	00
Ferri Chloridum	v.s.	v.s.
Ferri Chloridi, Tinctura.....	misc.	misc.
Ferri Phosphas Solubilis.....	v.s.	000
Ferri Sulphas	0.9	000
Ferri et Ammonii Citras.....	v.s.	000
Ferrum	000	000
Formaldehydi, Liquor.....	misc.	misc.
Gelatinum	000	000
Glycerinum	misc.	misc.
Glycerylis Nitras	000	v.s.
Guaiacol	53	v.s.
Guaiacolis Carbonas	000	48
Hexamethylenamina	1.5	10
Homatropinæ Hydrobromidum...	5.7	32.5
Hydrargyri Chloridum Corrosivum	13	3
Hydrargyri Chloridum Mite.....	000	000
Hydrargyri Iodidum Flavum....	00	000
Hydrargyri Iodidum Rubrum....	00	116
Hydrargyri Oxidum Flavum....	00	000
Hydrargyri Salicylas	000	000
Hydrargyrum Ammoniatum.....	000	000
Hydrastina	00	135
Hydrogenii Dioxidum, Aqua.....	misc.
Ichthyol	misc.
Iodoformum	9,391	46.7
Iodum	5,000	10
Magnesii Carbonas	000	000
Magnesii Oxidum	00	000
Magnesii Sulphas	1.1	000
Menthol	00	v.s.
Methylis Salicylas	00	v.s.
Morphina	3,330	168
Morphinæ Hydrochloridum.....	17.2	42
Morphinæ Sulphas	15.3	465
Pancreatinum	partly sol.	000
Paraffinum	000	000
Paraldehydum	8	misc.
Pelletierinæ Tannas	235	12.6
Pepsinum	50	000
Petrolatum	000	00
Petrolatum Liquidum	000	00
Phenol	19.6	v.s.
Phenol Liquefactum	12	misc.
Phenolphthaleinum	600	10
Phenylis Salicylas	2,333	5
Phosphorus	000	350
Physostigminæ Salicylas	72.5	12.7
Physostigminæ Sulphas	v.s.	v.s.
Pilocarpinæ Hydrochloridum....	0.3	2.3
Pilocarpinæ Nitras	4	60
Pix Liquida	00	1
Plumbi Acetas	2	30
Potassii Acetas	0.4	2
Potassii Bicarbonas	3	00
Potassii Bitartras	200	00
Potassii Bromidum	1.5	180
Potassii Carbonas	0.91	000

Substance Soluble in—	Parts of Cold Water.	Parts of Cold Alcohol.
Potassii Chloras	16	000
Potassii Citras	0.5	00
Potassii et Sodii Tartras.....	1.2	00
Potassii Hydroxidum	0.4	2
Potassii Iodidum	0.7	12
Potassii Nitras	3.6	00
Potassii Permanganas	15	dec.
Quinina	1,550	0.6
Quininæ Bisulphas	8.5	18
Quininæ Hydrochloridum	18	0.6
Quininæ Sulphas	720	86
Quininæ et Ureæ Hydrochloridum	1
Quininæ Tannas	800	3
Resorcinol	0.5	v.s.
Saccharum	0.46	137.2
Saccharum Lactis	4.79	000
Santoninum	5,300	34
Sapo	20	25
Sapo Mollis	4	1
Scopolaminæ Hydrobromidum....	1.5	16
Salvarsan	vs.
Sodii Arsanilas	6	125
Sodii Arsenas	1.2	00
Sodii Benzoas	1.6	43
Sodii Bicarbonas	12	000
Sodii Boras	17	000
Sodii Bromidum	1.7	12.5
Sodii Cacodylas	0.5	1
Sodii Carbonas Monohydratus...	2.9	000
Sodii Chloridum	2.8	00
Sodii Hydroxidum	1	v.s.
Sodii Iodidum	0.5	3
Sodii Nitris	1.4	50
Sodii Phosphas	5.5	000
Sodii Salicylas	0.8	5.5
Sodii Sulphas	2.8	000
Sodii Sulphis	2	00
Sodii Thiosulphas	0.35	000
Sparteinae Sulphas	1.1	2.4
Strophanthinum	v.s.	v.s.
Strychnina	6,400	110
Strychninæ Nitras	42	120
Strychninæ Sulphas	31	65
Sulphonethylmethanum	195	v.s.
Sulphonmethanum	360	47
Sulphur	000	00
Terpini Hydras	200	10
Theobromina	1,700	50
Theobrominæ Sodio-Salicylas....	1	28
Thymol	1,100	v.s.
Thymolis Iodidum	000	00
Veronal	145	28
Veronal-Sodium	5
Zinci Acetas	2.5	36
Zinci Chloridum	0.4	v.s.
Zinci Oxidum	000	000
Zinci Stearas	000	000
Zinci Sulphas	0.53	000

PHARMACOLOGIC INDEX

The following classification of drugs included in the list of useful remedies according to their therapeutic use has been adapted from the text-book on pharmacology and therapeutics by Arthur R. Cushny. This index serves to indicate the comprehensiveness of the list and should assist in suggesting to practitioners and teachers possible additions and deletions that might be made so that the list will include all of the fairly well-established remedies that are really of use.

For ready reference the titles used are those under which the drug or preparation has been included and described in the list of useful remedies.

I. DRUGS APPLIED FOR THEIR LOCAL ACTION TO THE SKIN, WOUNDS OR VISIBLE MUCOUS MEMBRANES

Corrosives or Caustics :

- Acidum Aceticum
- Acidum Nitricum
- Alumen Exsiccatum
- Argenti Nitras
- Arseni Trioxidum
- Chromii Trioxidum
- Hydrargyrum Ammoniatum
- Iodum
- Phenol
- Potassii Carbonas
- Potassii Hydroxidum
- Sodii Carbonas
- Sodii Hydroxidum
- Zinci Chloridum

Disinfectants and Antiseptics :

- Acidum Benzoicum
- Acidum Boricum
- Acidum Salicylicum
- Argenti Nitras
- Calx
- Calx Chlorinata
- Camphora
- Cresol
- Eucalyptol
- Formaldehydi, Liquor
- Hydrargyri Chloridum Corrosivum
- Hydrargyri Iodidum Rubrum
- Hydrogenii Dioxidum, Aqua
- Iodoformum
- Iodum
- Olea Volatilia
- Phenol
- Pix Liquida
- Potassii Permanganas
- Sulphur
- Thymol
- Zinci Chloridum

Astringents :

- Acidum Tannicum
- Alumen

Alumini Acetatis, Liquor
Argenti Nitras
Bismuthi Subcarbonas
Bismuthi Subgallas
Bismuthi Subnitras
Cupri Sulphas
Ferri Chloridum
Ferri Sulphas
Plumbi Acetas
Zinci Acetas
Zinci Oxidum
Zinci Sulphas

Styptics :

See also Soluble Astringents.
Ferri Chloridum
Alumen Exsiccatum

To Contract Vessels and Reduce Hemorrhage and Swelling :

Cocaina
Epinephrina

Emollients : Powders :

Amylum
Bismuthi Subcarbonas
Bismuthi Subnitras
Magnesii Carbonas
Talcum
Zinci Oxidum

Protectives :

Adeps
Adeps Lanæ
Cera
Olea Pingua
Petrolatum

Local Anodynes and Analgesics for Pain and Itching :

Aconitum
Ammonia, Aqua
Atropina
Chloroformum
Cocaina
Phenol
Sodii Bicarbonas

Local Anesthetics :

Æther
Æthylis Chloridum
Cocaina
Menthol

II. DRUGS USED FOR AFFECTIONS OF THE ALIMENTARY TRACT
Mouth and Throat. (See also Section 1) :

Demulcent :

Acacia
Ammonii Chloridum
Glycerinum
Potassii Chloras

To Lessen Salivation :

Atropina

Flavoring Substances :

Acidum Citricum
Olea Volatilia
Saccharum
Syrupi
Syrupus
Syrupus Pruni Virginianæ
Syrupus Tolutanus

Stomach:

Digestives :

Acidum Hydrochloricum
Pepsinum

Emetics :

Antimonii et Potassii Tartras
Apomorphinæ Hydrochloridum
Cupri Sulphas
Ipecacuanha
Sinapis
Sodii Chloridum
Zinci Sulphas

To Lessen Irritation and Vomiting :

Bismuthi Subcarbonas
Bismuthi Subnitrates
Chloralum Hydratum
Chloroformum
Cocaina
Codeina
Diacetylmorphinæ Hydrochloridum
Liquor Calcis
Menthol
Morphina
Opium

To Lessen Acidity, Antacids :

Calcii Carbonas
Calx (Liquor Calcis)
Magnesii Carbonas
Magnesii Oxidum
Potassii Carbonas
Sodii Bicarbonas
Sodii Hydroxidum

To Increase Secretion, Bitters :

Cinchona
Gentiana
Nux Vomica
Quinina
Strychnina

Carminatives :

See also Bitters.

Alcohol
Camphora
Capsicum
Cardamomum
Caryophyllus
Myrrha
Olea Volatilla
Sinapis
Zingiber

Intestine:

To Promote Digestion :

Pancreatinum

To Promote Evacuation, Purgatives :

Vegetable Purgatives :

Aloe
Aloinum
Colocynthis
Elaterinum
Jalapa
Podophyllum
Rhamnus Purshiana
Rheum

Ricini, Oleum
Senna
Tiglii, Oleum

Saline Purgatives :

Magnesii Carbonas
Magnesii Citratis, Liquor
Magnesii Oxidum
Magnesii Sulphas
Potassii Bitartras
Potassii Citras
Potassii et Sodii Tartras
Sodii Phosphas
Sodii Sulphas

Mercurial Purgatives :

Hydrargyri Chloridum Mite
Hydrargyrum

Miscellaneous :

Fel Bovis
Glycerinum
Sulphur

To Lessen Movement and Relax Spasm :

Acidum Tannicum
Argenti Nitras
Atropina
Belladonna
Bismuthi Subcarbonas
Bismuthi Subgallas
Bismuthi Subnitrates
Calx (Liquor Calcis)
Morphina
Opium
Plumbi Acetas

To Destroy Parasites, Anthelmintics :

Aspidium
Chloroformum
Hydrargyri Chloridum Mite
Pelletierinæ Tannas
Phenylis Salicylas
Santoninum
Terebinthinæ, Oleum
Thymol

Disinfectants and Antiseptics .

See also, Vegetable, Saline and Mercurial Purgatives.
Phenylis Salicylas

III. DRUGS USED FOR THEIR EFFECTS ON THE CIRCULATION

Heart:

To Strengthen Contraction :

Digitalis
Strophanthinum
Strophanthus

To Accelerate Pulse :

Atropina
Caffeina

To Slow Pulse :

Aconitum
Digitalis
Strophanthinum
Strophanthus

Vessels:

To Contract Caliber and Raise Blood-Pressure :

Epinephrina
Ergota

To Relax Vessels and Lower Blood-Pressure :

Amylis Nitris
 Glycerylis Nitratis, Spiritus
 Sodii Nitris

To Arrest Internal Hemorrhage, Styptics :

Ergota
 Hydrastis
 Morphina } To allay restlessness
 Opium }

To Remove Fluid (Dropsy, Anasarca) :

See also Diuretics (Kidney), Diaphoretics (Skin), Vegetable
 and Saline Purgatives (Intestine)
 Digitalis
 Hydrargyri Chloridum Mite.
 Scilla
 Strophanthinum
 Strophanthus

IV. DRUGS USED FOR THEIR EFFECTS ON THE GENITO-URINARY SYSTEM

To Increase the Flow of Urine (Diuretics) :

Ætheris Nitrosi, Spiritus
 Ammonii Acetatis, Liquor
 Caffaina
 Cubeba
 Digitalis
 Hydrargyri Chloridum Mite
 Potassii Acetas
 Potassii Citras
 Potassii Nitrates
 Santali, Oleum
 Scilla
 Sodii Nitris
 Sparteinæ Sulphas
 Strophanthinum
 Strophanthus
 Theobromina

To Render the Urine Less Acid :

Potassii Acetas
 Potassii Bicarbonas
 Potassii Citras
 Sodii Bicarbonas
 Sodii Carbonas

To Render the Urine More Acid :

Acid Sodium Phosphate, see Sodii Phosphas
 Mineral Acids

To Render the Urine Antiseptic :

Acidum Benzoicum
 Acidum Salicylicum
 Aspirinum
 Hexamethylenamina
 Phenylis Salicylas
 Sodii Benzoas
 Sodii Boras
 Sodii Salicylas
 Local antiseptics, etc.

To Promote Menstruation, Emmenagogues :

See also Vegetable Purgatives.
 Aloe
 Aloinum
 Myrrha

V. DRUGS USED FOR THEIR EFFECTS ON THE RESPIRATORY SYSTEM

To Stimulate the Respiratory Center :

Atropina
Caffeina
Camphora
Strychnina

To Reduce the Irritability of the Center in Cough :

Chloralum Hydratum
Chloroformum
Codeina
Diacetylmorphinæ Hydrochloridum
Morphina
Opium

To Increase and Liquefy the Bronchial Secretion :

Ammonii Carbonas
Antimonii et Potassii Tartras
Apomorphinæ Hydrochloridum
Ipecacuanha
Lobelia
Potassii Iodidum
Scilla
Sodii Iodidum

To Lessen the Secretion of the Bronchi (?)

Terpini Hydras

To Relax Bronchial Spasm in Asthma :

Amylis Nitris
Atropina
Belladonna
Glycerylis Nitratis, Spiritus
Sodii Iodidum
Sodii Nitris

VI. DRUGS USED FOR THEIR EFFECTS ON THE CENTRAL NERVOUS SYSTEM

Stimulants :

(a) The spinal cord :

Strychnina

(b) The brain and medulla oblongata :

Atropina
Caffeina
Camphora

Depressants :

(a) To paralyze sensation (general anesthetics) :

Æther
Æthylis Chloridum
Chloroformum

(b) To induce sleep and rest (hypnotics or narcotics) :

Alcohol
Chloralum Hydratum
Codeina
Morphina
Opium
Paraldehydum
Scopolaminæ Hydrobromidum
Sulphonethylmethanum
Sulphonmethanum
Veronal

(c) To relieve pain (analgetics or anodynes) :

Acetanilidum
Acetphenetidinum
Acidum Salicylicum
Alcohol

Antipyrina
Aspirin
Chloralum Hydratum
Codeina

Methylis Salicylas
Morphina
Sodii Salicylas

VII. DRUGS USED TO REDUCE FEVER TEMPERATURE

Acetanilidum
Acetphenetidinum
Acidum Salicylicum
Aconitum
Ammonii Acetatis, Liqueur
Antipyrinum
Aspirin
Gualacol
Phenol
Quinina
Sodii Salicylas

VIII. DRUGS USED FOR THEIR EFFECTS ON THE LIVER

To Increase the Secretion of Bile, Cholagogues :

Acidum Salicylicum
Fel Bovis

IX. DRUGS USED FOR THEIR EFFECTS ON THE BLOOD

To Increase the Hemoglobin :

Arseni Trioxidum
Ferri Carbonas
Ferri Iodidum
Ferri Phosphas Solubilis
Ferri et Ammonii Citras
Sodii Arsanilas
Sodii Cacodylas

To Render the Blood Alkaline :

Potassii Acetas
Potassii Bicarbonas
Potassii Citras
Sodii Bicarbonas
Sodii Carbonas
Sodii Hydroxidum

To Increase the Coagulability (?) :

Calcii Chloridum
Calcii Hypophosphis
Calcii Lactas

X. DRUGS USED FOR SPECIFIED DISEASES

In Malaria :

Arseni Trioxidum
Quinina

In Syphilis :

Hydrargyri Chloridum Corrosivum
Hydrargyri Chloridum Mite
Hydrargyri Iodidum Flavum
Hydrargyri Iodidum Rubrum
Hydrargyri Salicylas
Hydrargyrum
Potassii Iodidum
Salvarsan
Sodii Iodidum

In Rheumatic Fever :

Acidum Salicylicum
Aspirin

Methylis Salicylas
Sodii Salicylas

In Myxedema and Some Other Thyroid Diseases :
Glandulæ Thyroideæ Siccæ

In Diphtheria :
Serum Antidiphthericum

In Tetanus
Serum Antitetanicum

In Trypanosomiasis :
Antimonii et Potassii Tartras
Sodii Arsanilas

In Gout :
Colchici Semen

XI. DRUGS USED FOR THEIR EFFECTS ON THE SKIN

Corrosives or Caustics
Emollients and Protectives
Local Anodynes and Anesthetics

Irritants :

Aconitum
Alcohol
Ammonia
Camphora
Cantharis
Capsicum
Iodum
Menthol
Sinapis
Terebinthinæ, Oleum
Tiglii, Oleum

Disinfectants or Irritants Used Chiefly in the Form of Ointments in Parasitic Skin Diseases :

Balsamum Peruvianum
Benzoinum
Camphora
Chrysarobinum
Hydrargyrum
Ichthyol
Iodum
Pix Liquida
Resorcinum
Sulphur
Thymol

Arsenic, Potassium, Iodid, etc., May Be Used Internally in Skin Diseases.

Drugs Administered Internally to Increase the Secretion of Perspiration, Diaphoretic or Sudorifics :

Ætheris Nitrosi, Spiritus
Antimonii et Potassii Tartras
Camphora
Ipecacuanha
Opium
Pulvis Ipecacuanhæ et Opii
Pillocarpinæ Hydrochloridum

Drugs Administered Internally to Lessen Secretion of Perspiration :

Atropina
Belladonna

XII. DRUGS USED LOCALLY FOR THEIR EFFECTS ON THE EYE

Drugs Dilating the Pupil and Relaxing the Accommodation,
Mydriatics :

Atropina
Cocaina
Homatropinæ Hydrobromidum
Scopolaminæ Hydrobromidum

Drugs Contracting the Pupil and the Ciliary Muscle, Myotics :

Physostigminæ Salicylas
Pilocarpinæ Hydrochloras

OTHER PROPERTIES OF WELL-KNOWN DRUGS

The following classification is taken from "Introduction to Materia Medica and Pharmacology" by Oliver T. Osborne.

Drugs and Preparations Which May Cause an Eruption on,
or Itching of, the Skin :

Antitoxin
Arsenic
Belladonna
Bromids
Chloral
Copaiba
Iodids
Opium
Quinin
Salicylic Acid
Synthetic Compounds
Volatile Oils, and drugs containing them.

Drugs Which May Change the Color of the Urine :

Drugs that increase its amount cause it to be lighter.
Drugs that irritate the kidneys cause it to be darker.
Methylene-blue causes it to be green, if acid.
Phenol may cause it to be brown (same appearance as bile).
Santonin causes it to be yellow, if acid ; purple, if alkaline.
Senna may cause it to be red, if acid ; yellow, if alkaline.
Sulphonals may cause it to be very dark.

Drugs Which Color the Feces :

Bismuth salts color them black or dark gray.
Colchicum colors them greenish.
Iron colors them black.
Mercury colors them green.
Purgatives cause them to be darker.

Drugs Which Are Excreted with the Milk :

Arsenic
Bromids
Hexamethylenamin
Iodids
Lead
Mercury
Opium
Quinin
Sulphur
Vegetable Cathartics
Volatile Oils

INDEX

	PAGE		PAGE
Acacia	7	Ammoniated mercury (hy-	
mucilage of	7	drargyrum ammoni-	
Acetanilid	7	atum)	84
Acetic acid	8	mercury ointment of....	84
Acetphenetidin	8	Ammonium acetate	20
Acetylsalicylic acid (aspirin)	30	acetate, solution of....	20
Achylia gastrica		carbonate	20
diluted hydrochloric acid		chlorid	21
in	10	compounds	19
Acid, acetic	8	sulpho-ichthyolate	86
acetylsalicylic (aspirin)	30	Amyl nitrite	22
arsenous, solution of....	29	Amylum	23
benzoic	9	Analgesics and anodynes, 151, 155	
boric	9	Anemia	
carbolic (phenol)	103	ferrous carbonate in....	67
citric	10	salvarsan in	121
diethylbarbituric (see		sodium cacodylate in... 129	
veronal)	144	Anesthetics, general	155
hydrochloric	10	local	151
hydrochloric, diluted ..	10	Anodynes and analgesics, 151, 155	
hydrocyanic, diluted....	11	Antacids	152
nitric	11	Anthelmintics	153
salicylic	12	Antidiphtheric serum.....	126
tannic	12	Antimony and potassium tar-	
tannic, glycerite of....	13	trate	23
Acidity, to lessen	152	wine of	23
Acne vulgaris		Antipyrin	24
ichthyl in	86	Antiseptics and disinfectants	
Aconite	13150, 153	
tincture of	13	Antitetanic serum.....	126
Actinomycosis		Antitoxin, diphtheria	126
potassium iodid in....	114	Apomorphin hydrochlorid... 24	
Adeps	14	Aqua	25
benzoinatus	14	ammoniae	20
lanæ hydrosus	14	camphoræ	46
Adnephryn, adrenalin (epi-		chloroformi	52
nephryn)	63	destillata	25
Aether	14	hydrogeni dioxidi	85
nitrosus	15	menthae piperitæ	94
nitrosus spiritus	15	rose	120
spiritus	15	Aquæ	25
spiritus compositus....	15	Argenti nitras	25
Aethylis chloridum	15	nitras fusus	27
Aethyl-morphinae hydro-		Argenti proteinas (see pro-	
chloridum	15	targol)	114
Albumin tannate (tannalbin)	13	Aristol (thymol iodid) 141	
Alcohol	16	Aromatic elixir	17
Alcoholic poisoning		Arsenic trioxid	27
cafein in	39	Arsenobenzol (salvarsan) .. 120	
Alcoholism		Arsenous acid (arsenic tri-	
strychnin in	136	oxid)	29
Aloes	17	acid, solution of	29
extract of	17	and mercuric iodids,	
Aloin	18	solution of	29
Alum	18	Asafetida	29
exsiccated	18	Aspidium	29
Aluminum acetate	18	oleoresin of	29
acetate, solution of....	18	Aspirin	30
Amebic dysentery		Asthma	
ipecac in	89	drugs to relax bronchial	
Amenorrhœa		spasm in	155
desiccated thyroid glands		epinephrin in	63
in	73	Astringents	150
Ammonia	19	Atoxyl (sodii arsanilas).... 127	
aromatic spirit of....	21	Atropin	30
liniment	20	sulphate	33
water	20	Bacterial vaccines (vaccina) 143	
		Balsam of Peru	33

	PAGE		PAGE
Balsam of Tolu	34	Calomel (hydrargyri chlori- dum mite)	79
Bed-sores		Calx	43
compound tincture of		chlorinata	44
benzoin in	35	Camphor	45
Belladonna leaves	34	liniment	46
leaves, extract of	34	spirit of	46
leaves, tincture of	34	water	46
ointment	35	Cannabis indica	46
plaster	34	extract	46
Benzoic acid	9	tincture	47
Benzoin	35	Cantharides	47
tincture of, compound..	35	cerate	47
Benzosulphinid	35	Capsicum	48
Betanaphthol	36	tincture	48
Betula, oil of	94	Capsules	48
Biniodid of mercury	81	Carbo ligni	48
Birch, oil of	95	Carbolic acid (phenol)	103
Bismuth subcarbonate	36	Cardamom	49
subgallate	37	tincture of	49
subnitrate	37	Carminatives	152
subsaliolate	39	Carron oil	44
Bitters	152	Caryophyllus	49
Blastomycosis	114	oil of	49
Blood, alkaline, drugs to ren- der	156	Cascara sagrada (rhamnus purshiana)	118
Blood, coagulability, drugs to increase	156	aromatic fluid extract of	118
Blood, drugs, effects on the..	156	extract of	118
Blood-pressure		fluid extract of	118
amyl nitrate, effect on..	22	Castor oil (oleum ricini) ..	119
drugs to lower	154	Cataplasmata	49
strychnin, effect on	135	Caustics or corrosives	150
Blue mass	83	Cera alba	49
ointment	83	flava	49
Borax (sodium borate).....	128	Cerates	50
Boric acid	9	Ceratum cantharidis	47
acid, ointment of	10	Chalk, precipitated	41
Boroglycerin, glycerite of...	9	Chancroids	12
Bothriocephalus	30	Charcoal (carbo ligni)	48
Bronchial secretion, to in- crease and liquefy the..	155	Charta sinapis	127
secretion, to lessen	155	Chartae	50
Bronchitis		Cherry, wild	115
antimony and potassium		Chloral, hydrated	50
tartrate in	23	Chlorinated calcium oxid (chlorinated lime) ..	44
antipyrin in	24	soda, solution of	45
arsenic trioxid in	27	Chloroform	51
strychnin in	135	liniment	52
Bruises		spirit of	52
solution of lead subace- tate in	109	water	52
Burns		Chlorosis	
lime-water in	44	sodium cacodylate in...	129
Cacao butter (oleum theo- bromatis)	140	Cholagogues	156
Cachets	39	Chromium trioxid (chromic acid)	52
Caffein	39	Chrysarobin	53
citratd	40	ointment	53
sodio-benzoate	41	Cinchona	53
Calabar bean (physostigma)..	106	tincture of	53
Calcined magnesia	92	tincture of, compound..	54
Calcium carbonate, precipi- tated	41	Cinnamon	54
chlorid	41	oil of	54
hydroxid, solution of...	44	water	54
hypophosphite	42	Circulation, drugs used for their effects on the..	153
lactate	43	Citratd caffein	40
oxid	43	Citric acid	10
oxid, chlorinated	44	Cloves (caryophyllus)	49
phosphate, precipitated.	43	oil of	49
		Clysters (enemata)	63

	PAGE		PAGE
Cocain	54	Disinfectants and anti-septics	150, 153
hydrochlorid	56	or irritants used in parasitic skin diseases ...	157
Codein	56	Diuretics	154
phosphate	56	Diuretin (theobrominae sodio-salicylas)	140
sulphate	56	Donovan's solution (solution of arsenous and mercuric iodid)	29
Cod-liver oil (morruae oleum)	97	Dover's powder (pulvis ipecacuanhae et opii) ..	100
Colchicum seed	57	Dropsy, renal	140
seed, tincture of	57	Dyspepsia	45
Colds			
camphor for	45	Ear, catarrh of	87
Collodion	57	iodin in	87
flexible	57	Eczema	
Colocynth	58	arsenic trioxid in	27
extract of	58	desiccated thyroid gland in	73
extract of, compound ..	58	Effervescent salts	120
Confections	58	Elaterin	62
Conjunctivitis		Elixir, aromatic	17
boric acid in	9	Elixirs	63
ethyl-morphin hydrochlorid	15	Emetics	152
sodium borate	128	Emmenagogues	154
Conjunctivitis, gonorrheal, ..		Emollients	151
silver nitrate in	25	Emphysema	
Copaiba	58	arsenic trioxid in	27
Copper sulphate	59	Emplastra	63
Corn-starch (amylum)	23	Emulsions	63
Corneal ulcer	16	Endocervicitis	
Corrosive sublimate	78	iodin in	87
Corrosives or caustics	150	Enemata (clysters)	63
Coryza		Enteritis, ulcerative	
atropin in	30	bismuth subnitrate in ...	37
codein in	54	Epinephrin	63
Cough		Epsom Salts (magnesium sulphas)	93
codein for	56	Ergot	65
to reduce nerve center irritability in	155	fluidextract of	66
Creosote	58	Eruption on the skin, drugs which may cause an... 158	
Cresol	59	Erysipelas	
solution of, compound ..	59	ichthyol in	86
Croton oil (oleum tiglli) ...	141	Eserin salicylate (physostigminae salicylas) ..	106
Cupri sulphas	59	sulphate (physostigminae sulphas)	107
Cystitis		Ether (aether)	14
boric acid in	9	nitrous	15
hexamethylenamin in ...	76	nitrous, spirit of	15
Decoctions	60	spirit of	15
Demulcents	151	spirit of, compound....	15
Depressants, central nervous system	155	Ethyl chlorid (aethylis chloridum)	15
Dermatitis herpetiformis ...	28	Ethyl-morphin hydrochlorid..	15
Dermatol (bismuth subgalate)	37	Eucalyptus	66
Diabetes		oil of	66
sodium cacodylate in ...	129	Extract of aloes	17
Diabetes insipidus		of belladonna leaves....	34
ergot in	65	of cannabis indica	46
Diarrhea		of cascara sagrada	118
lime-water in	44	of colocynth	58
Dicetyl-morphinae hydrochloridum	60	of colocynth, compound..	58
Digestives	152	of gentian	73
Digitalis	61	of Indian cannabis	46
infusion of	62	of nux vomica	98
tincture of	62		
Dionin (aethyl-morphinae hydrochloridum)	15		
Diphtheria			
antitoxin (see serum antidiphthericum)	126		
drugs used in	157		
strychnin in	136		

	PAGE		PAGE
of opium	100	Glonoïn	75
of rhamnus purshiana..	118	Glycerin	74
of rhubarb	119	suppositories of	74
Extracts	66	Glyceryl nitrate, spirit of...	75
Eye, drugs used locally for		Glycerite of boroglycerin....	9
their effects on the..	158	Glycerites	74
Fats and oils, fixed.....	99	Glycerol: see Glycerin.....	74
Favus		Glyceryl trinitrate	75
sodium sulphite in	133	Glycyrrhiza	75
Feces, drugs which color the	158	fluidextract of	75
Fel bovis	67	powder of, compound...	75
bovis purificatum	67	Goiter	
Ferri et ammonii citras....	68	desiccated thyroid glands	
Ferric chlorid	68	in	74
chlorid, tincture of	68	Golden seal	84
phosphate	69	Gout, drugs used in	157
Ferrous carbonate	67	Gonorrhea	
carbonate, mass of	68	protargol in	114
carbonate, pills of	68	Gray powder	83
iodid	69	Green vitriol	70
iodid, syrup of	69	Guaiacol	76
sulphate	70	carbonate	76
sulphate, exsiccated ...	70	Gum arabic	7
Ferrum	70	Headache	
reductum	70	acetanilid in	7
Fever, drugs used to reduce		sodium salicylate in....	132
temperature	156	Hemoglobin, drugs used to	
Fevers		increase	156
solution of ammonium		Hemophilia	
acetate	20	calcium chlorid in	42
Flatulence		Hemorrhages	
oil of turpentine in	139	epinephrin in	63
Flavoring substances	151	internal, to arrest in...	154
Flaxseed (linum)	91	Hemorrhoids	
Fluid, to remove	154	iodoform in	87
Fluidextract of cascara sa-		Henbane	86
grada	118	Heroin hydrochlorid (dicetyl-	
of cascara sagrada, aro-		morphinae hydrochlor-	
matic	118	idum)	60
of ergot	66	Hexamethylenamin	76
of glycyrrhiza	75	Homatropin hydrobromid ...	78
of hydrastis.....	85	Honeys (mellita)	93
of ipecac	90	Hookworm disease	
of rhamnus purshiana..	118	thymol in	141
of rhamnus purshiana,		Hydrargyri ammoniati un-	
aromatic	118	guentum	84
of senna	125	chloridum corrosivum...	78
of viburnum prunifolium	144	chloridum mite	79
Fluidextracts	70	dilutum unguentum ...	83
Formaldehyd	71	iodidum flavum	80
solution of.....	71	iodidum rubrum	81
Fowler's solution (potassium		oxidum flavum	82
arsenite)	28	salicylas	82
Galactorrhea		unguentum	83
atropin in	33	Hydrargyrum	82
Gaultheria, oil of	94	ammoniatum	84
Genito-urinary system, drugs		cum creta	83
used for their effects		Hydrastis	84
on the	154	fluidextract of	85
Gentian	72	Hydrated chloral	50
extract of	73	Hydrochloric acid.....	10
tincture of, compound..	73	Hydrocyanic acid, diluted...	11
Ginger (zingiber)	146	Hydrogen dioxide	85
tincture of	146	dioxide, solution of	85
Glauber's salt (sodii sulphas)	133	peroxid	85
Glaucoma		Hydrous wool fat (adeps	
ethyl-morphin hydro-		lanæ)	14
chlorid in	15	Hyoscin hydrobromid	124
sodium salicylate in...	132	Hyoscyamus	86
		tincture of	86

	PAGE		PAGE
Hyperacidity		Linseed oil	91
bismuth subnitrate in..	38	Linum	91
Hypnotics or narcotics	155	Liquor: See Solution.	
Ichthyol	86	Liver, drugs for their effects	
Indian cannabis	46	on the	156
cannabis, extract of	46	Lobelia	91
cannabis, tincture of	47	tincture of	91
Infections		Lunar caustic (argenti	
solution of ammonium		nitras)	25
acetate in	20	Lupus erythematosus	
Infusion of digitalis	62	ichthyol in	86
Infusions	86	Magnesia, calcined	92
Insomnia		Magnesium carbonate	92
hydrated chloral in	50	citrate	92
Iodin	87	citrate, solution of	92
tincture of	89	oxid	92
Iodoform	87	sulphate	93
Ipecac	89	sulphate, effervescent	93
and opium, powder of..	100	Male fern (aspidium)	29
fluidextract of	90	Malaria	
syrup of	90	drugs used in	156
Iritis		quinin and urea hydro-	
ethyl-morphin hydro-		chlorid in	117
chlorid in	15	quinin tannate in	116
Iron (ferrum)	70	Mammitis	
and ammonium citrate..	68	atropin in	32
reduced	70	Mass, blue	83
See also ferric and fer-		of ferrous carbonate	68
rous compounds		of mercury (massa hy-	
Irritants	157	drargyri)	83
or disinfectants used in		Masses	93
parasitic diseases	157	Mellita	93
Itching of the skin, drugs		Menorrhagia	
which may cause an..	158	hydrastis in	84
Jaborandi (pilocarpus)	107	Menstruation, to promote...	154
Jalap	90	Mentha piperita (peppermint)	93
powder of, compound	90	Menthae piperitae, aqua....	94
Keratitis		piperitae oleum	93
sodium salicylate in....	132	piperitae spiritus	94
Labarraque's solution (liquor		Menthol	94
sodae chlorinatae)	45	Mercurial ointment	83
Lanolin (adeps lanae hydro-		Mercuric and arsenous iodids,	
sus)	14	solution of	29
Lard (adeps)	14	chlorid, corrosive	78
benzoinated	14	iodid, red	81
Laryngitis		oxid, yellow	82
antimony and potassium		oxid, yellow, ointment	
tartrate in	23	of	82
squill in	123	salicylate	82
Laudanum (tinctura opii)	100	Mercurous chlorid, mild	79
Lead acetate (plumbi acetas)	108	iodid, yellow	80
subacetate, solution of..	109	Mercury (hydrargyrum)	82
sugar of	108	ammoniated	84
Leukemia		biniodid of	81
ferrous carbonate in....	67	green iodid of	80
Lichen planus		mass of	83
arsenous trioxid in....	27	protiodid	80
Licorice root (glycyrrhiza)	75	with chalk	83
Lime	43	Methyl salicylate	94
chlorinated	44	Metrorrhagia	
liniment	44	hydrastis in	84
water	44	Milk, drugs which are ex-	
Liniment, ammonia	20	creted with the.....	158
camphor	46	Milk-sugar (saccharum lac-	
chloroform	52	tis)	120
lime (calcis)	44	Mixtures	95
soap	123	Morphin	95
Liniments	90	hydrochlorid	97
		sulphate	97
		Morrhuae oleum	97

	PAGE		PAGE
Mucilage of acacia	7	chrysarobin	53
Mucilages	98	mercurial (unguentum	
Mustard (sinapis)	126	hydrargyri)	83
black	126	of ammoniated mercury..	84
oil of, volatile.....	127	of boric acid	10
paper	127	of yellow mercuric oxid..	82
Myrrh	98	of zinc oxid.....	146
tincture of	98	sulphur	138
Myxedema and some other		tar	108
thyroid diseases, drugs		Ointments	143
used in	157	Oleates	99
Naphthol (betanaphthol)....	36	Oleoresin of aspidium.....	29
Narcotics or hypnotics.....	155	Oleoresins	99
Neosalvarsan	122	Oleum: See Oil	
Nephritis		Opium	99
potassium acetate in... 110		and ipecac, powder of..	100
Nerve exhaustion		extract of	100
cafein in	39	powdered	100
Nervous system, central,		tincture of	100
drugs used for their		tincture of, camphorated	100
effects on the.....	155	tincture of, deodorized..	100
Neuralgia		Opium poisoning	
acetanilid in	7	cafein in	39
arsenic trioxid in.....	27	Oxgall (fel bovis).....	67
Nevi		purified	67
nitric acid in.....	11	Oxygen, compressed	100
Nipples, cracked		Pancreatin	101
compound tincture of		Papers (chartae)	50
benzoin in	35	Paraffin	101
Nitric acid	11	Paraldehyd	102
Nitrites	22	Paralysis	
Nitroglycerin	75	strychnin in	135
spirit of	75	Parasites, to destroy.....	153
Nitrous ether, spirit of....	15	Paregoric (tinctura opii cam-	
Nux vomica	98	phorata)	100
extract of	98	Pelletierine tannate	102
tincture of	98	Pemphigus	
Obesity		arsenic trioxid in.....	27
desiccated thyroid glands		Peppermint	93
in	74	oil of	93
Oil, carron (linimentum cal-		spirit of	94
cis)	44	water	94
castor (oleum ricini)..	119	Pepsin	102
croton (oleum tiglli)... 141		Perspiration, drugs adminis-	
linseed (oleum lini)....	91	tered to increase the	
of birch (oleum betulae)	95	secretion of	157
of cinnamon (oleum cin-		Pertussis	
namomi)	54	antipyrin in	24
of cloves (oleum caryo-		Peru, balsam of.....	33
phylli)	49	Petrolatum	102
of eucalyptus (oleum		liquid	102
eucalypti)	66	Pharyngitis	
of gaultheria (oleum		boric acid in.....	9
gaultheriae)	94	capsicum in	48
of mustard, volatile (ol-		iodin in	87
eum sinapis volatile)..	127	Phenacetin (acetphenetidin).	8
of peppermint (oleum		Phenol	103
menthae piperitae)... 93		liquefied	104
of rose (oleum rosae)..	120	Phenolphthalein	104
of santal (oleum santali)	122	Phenyl salicylate	104
of theobroma (oleum the-		Phosphorus	105
obromatis)	140	Physostigma	106
of turpentine (oleum te-		Physostigmine salicylate	106
rebinthinae)	139	sulphate	107
of wintergreen	94	Pills	108
Oils and fats, fixed.....	99	of ferrous carbonate	
volatile or essential....	99	(Blaud's)	68
Ointment, belladonna	35	Pilocarpin hydrochlorid	107
blue (hydrargyri dilutum		nitrate	108
unguentum)	83	Pilocarpus	107

	PAGE		PAGE
Pix liquida	108	Resins	117
unguentum	108	Resorcinol	117
Plaster, belladonna	34	Respiratory system, drugs	
Plasters (emplastra)	63	used for their effects	
Plumbi acetate	108	on the	155
Pneumonia		Rhamnus purshiana	118
potassium acetate in...	109	extract of	118
strychnin in	135	fluidextract	118
Podophyllum	109	fluidextract, aromatic ..	118
resin of	109	Rheumatic fever, drugs used	
Potassium acetate	109	in	156
and sodium tartrate....	112	Rheumatism	
arsenite, solution of....	28	potassium acetate in...	109
bicarbonate	110	sodium salicylate in....	132
bitartrate	110	Rhubarb (rheum)	118
bromid	110	extract of	119
carbonate	111	syrup of, aromatic....	119
chlorate	111	tincture of, aromatic....	119
citrate	112	Ricini oleum	119
citrate, effervescent	112	Rochelle salts (potassii et	
hydroxid	112	sodii tartrates)	112
hydroxid, solution of....	113	Rosacea	
iodid	113	ichthyol in	86
permanganate	114	Rose	119
tartrate and antimony. 23		oil of	120
Poultices (cataplasmata)....	49	water	120
Powder, compound efferves-		Rosin	117
cing	112	Saccharin (benzosulphinid)..	35
Dover's	100	Saccharum	120
of glycyrrhiza, compound 75		lactis	120
of ipecac and opium....	100	Salicylic acid	12
of jalap, compound....	90	Salivation, to lessen	151
Seidlitz	112	Salol (phenyl salicylate)....	104
Powdered opium	100	Salt, common (sodium chlo-	
Powders	115, 151	rid)	130
Precipitated calcium carbon-		Salts, effervescent	120
ate	41	Epsom (magnesi sul-	
calcium phosphate	43	phas)	93
chalk	41	Glauber's	133
Protargol	114	Rochelle (potassii et	
Protectives	151	sodii tartrate)	112
Protiodid of mercury.....	80	Salvarsan	120
Prunus virginiana	115	Santal, oil of.....	122
virginianae syrupus ...	115	Santonin	122
Pseudoleukemia		Sapo	122
sodium cacodylate in...	129	linimentum	123
Psoriasis		mollis	123
arsenic trioxid in.....	27	Sarsaparilla	123
desiccated thyroid glands		syrup of, compound....	123
in	73	Scabies	
sodium cacodylate in...	129	balsam of Peru in.....	33
Pulse, to accelerate.....	153	sodium sulphite in....	133
to slow	153	Scilla	123
Pulveres, pulvis; see powder,		syrupus	124
powders	115	tinctura	124
Purgatives, mercurial	153	Scleritis	
saline	153	ethyl-morphin hydrochlo-	
vegetable	152	rid in	15
Pyelitis		Scopolamin hydrobromid ...	124
hexamethylenamin in ..	76	Seborrhea	
Quinin	115	ointment of ammoniated	
and urea hydrochlorid..	117	mercury in	84
bisulphate	116	Secretion, to increase.....	152
hydrochlorid	116	Seidlitz powder (pulvis effer-	
sulphate	116	vescens compositus) .	112
tannate	116	Senna	125
Relapsing fever		fluidextract of	125
salvarsan in	120	syrup of	125
Resin of podophyllum.....	109	Serum, antidiphtheric	126
		antitetanic	126

	PAGE		PAGE
Serums and vaccines.....	125	of ether	15
Silver nitrate (argenti nitras) ..	25	of ether, compound.....	15
molded	26	of glyceryl nitrate....	75
proteinate (protargol) ..	114	of nitroglycerin	75
Sinapis	126	of nitrous ether.....	15
charta	127	of peppermint (menthae	
nigra	126	piperitae)	94
volatile oleum	127	Sprains	
Skin disease		solution of lead subace-	
arsenic trioxid in.....	27	tate in	109
bismuth subgallate in..	37	Squill (scilla)	123
Skin, drugs used for their ef-		syrup of	124
fects on the.....	157	tincture of	124
Soap	122	Starch (amylum)	23
liniment	123	Stimulants	155
soft	123	Stramonium	134
Sodium arsanilate	127	Strophanthin	134
arsenate	127	Strophanthus	135
benzoate	127	tincture of	135
bicarbonate	128	Strychnin	135
borate	128	nitrate	136
bromid	129	sulphate	137
cacodylate	129	Styptics	151, 154
carbonate, monohydrated	130	Sugar (saccharum)	120
chlorid	130	of lead	108
diethyl-barbiturate	144	of milk	120
hydroxid	130	Sulphonal (sulphonmetha-	
hydroxid, solution of... 131		num)	138
iodid	131	Sulphonethylmethane	137
nitrite	131	Sulphonmethane	138
phosphate	131	Sulphur	138
phosphate, effervescent.	132	ointment	138
salicylate	132	precipitated	138
sulphate	133	sublimed	138
sulphite	133	washed	138
tartrate and potassium.	112	Suppositories	138
thiosulphate	133	of glycerin	74
Solution, Donovan's (arsen-		Supracapsulin, suprarenalin,	
ous and mercuric		l-suprarenin synthetic	
iodids)	29	(epinephrin)	63
Fowler's (potassium ar-		Syrup	120
senite)	28	of ferrous iodid	69
Labarraque's (chlorin-		of ipecac	90
ated soda)	45	of rhubarb, aromatic... 119	
of aluminum acetate....	18	of sarsaparilla, compound 123	
of ammonium acetate....	20	of senna	125
of arsenous acid	29	of squill	124
of arsenous and mercuric		of Tolu	34
iodids	29	of wild cherry	115
of calcium hydroxid....	44	Syphilis, drugs used in.... 156	
of chlorinated soda....	45	Tablets, compressed	139
of cresol, compound....	59	Tablet-triturations	139
of formaldehyd	71	Tannalbin	13
of hydrogen dioxid	85	Tannic acid	12
of lead subacetate.....	109	glycerite of	13
of magnesium citrate..	92	Tar (pix liquida)	108
of potassium arsenite..	28	Tartar emetic (antimony and	
of potassium hydroxid..	113	potassium tartrate) ..	23
of sodium hydroxid.....	131	Terebinthina	139
of zinc chlorid.....	145	oleum	139
Solutions	91	Terpin hydrate	139
Sparteïn sulphate	134	Tetanus antitoxin (serum	
Sporotrichosis		antitetanicum)	126
potassium iodid in.....	113	drugs used in	157
Spasm		Theobroma, oil of	140
hydrated chloral in.....	50	Theobromin	140
to lessen movement and		sodium salicylate	140
relax	153	Thymol	141
Spirit of ammonia, aromatic.	21	iodid	141
of camphor	46		
of chloroform	52		

	PAGE		PAGE
Thyroid glands, desiccated..	73	Urine, drugs to render urine	
Tiglli oleum	141	less acid	158
Tincture of aconite	13	drugs which may change	
of ammoniated valerian..	144	color of	158
of belladonna leaves...	34	Urotropin (hexamethylena-	
of benzoin, compound...	35	mina)	76
of cannabis indica	47	Urticaria	
of capsicum	48	arsenic trioxid in.....	27
of cardamom	49	Uterine hemorrhage	
of cinchona	53	fluidextract of hydrastis	
of cinchona, compound..	54	in	85
of colchicum seed	57	Vaccina	143
of digitalis	62	Vaccine virus	145
of ferric chlorid	68	Vaccines and serums	125
of gentian, compound...	73	Valerian	143
of ginger	146	ammoniated, tincture of	144
of hyoseyamus	86	Vallet's mass	68
of Indian cannabis	47	Vasoconstriction, drugs for..	151
of iodin	89	Veronal	144
of lobelia	91	Viburnum prunifolium	144
of myrrh	98	prunifolium, fluidextract	
of nux vomica.....	98	of	144
of opium	100	Vina	145
of opium, camphorated..	100	Vincent's angina	
of opium, deodorized....	100	salvarsan in	121
of rhubarb, aromatic....	119	Vinum antimonii	23
of squill	124	Virus vaccinum	145
of strophanthus	135	Vitriol, green	70
of valerian, ammoniated	144	Volatile or essential oils	99
Tinctures	141	Vomiting, drugs to lessen	
Tolu, balsam of	34	irritation and	152
syrup of	34	Warts	
Tonsillitis		nitric acid for	11
capsicum in.....	48	Water (aqua)	25
Trachomatous lids		ammonia	20
copper sulphate in.....	59	camphor	46
Tragacanth	142	chloroform	52
Trinitrin	75	cinnamon	54
Trional (sulphonethylme-		peppermint	94
thanum)	137	rose	120
Triturations	142	Waters (aquae)	25
Troches	142	Wax, white (cera alba).....	49
Trypanosomiasis, drugs used		yellow (cera flava)	49
in	157	Wild cherry (prunus virgini-	
Tuberculin	142	ana)	115
Tuberculosis		syrup of	115
sodium cacodylate in... 129		Wine of antimony	23
Turpentine (terebinthina).. 139		Wines (vina)	145
Typhoid fever		Wintergreen, oil of.....	94
calcium chlorid in	41	Wool fat (adeps lanæ	
ergot in	65	hydrated)	14
Ulcer of the stomach		Zinc acetate	145
bismuth subnitrate in.. 37		chlorid	145
Ulcers		oxid	145
compound tincture of		oxid, ointment of	146
benzoin in	35	solution of	145
Unguentum: See Ointment.		stearate	146
Urea and quinin hydro-		sulphate	146
chlorid	117	Zingiber	146
Urethritis, gonorrhea		tinctura	146
copaiba in	58		











