Dental surgery notes.

Contributors

Dowsett, Ernest B.

Publication/Creation

London: Dental Manufacturing Co., 1912.

Persistent URL

https://wellcomecollection.org/works/r4vn7ryb

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org

DENTAL SURCERY NOTES

ev Marker b. dowsett




3/6 peus 100/0



22101902465

Med K48800







GLAUDIUS ASH, SONS & Cº LIMITED. 5 TO 12. BROAD ST., GOLDEN SQUARE, W.1.

DENTAL SURGERY NOTES



DENTAL SURGERY NOTES

BY

ERNEST B. DOWSETT, L.R.C.P., M.R.C.S., L.D.S.

ASSISTANT DENTAL SURGEON AND DEMONSTRATOR OF DENTAL HISTOLOGY TO GUY'S HOSPITAL

THIRD EDITION

COPYRIGHT

LONDON
THE DENTAL MANUFACTURING CO. LIMITED
ALSTON HOUSE, NEWMAN STREET, W.
1912



WELLCOME INSTITUTE LIBRARY							
Coll.	welMOmec						
Call							
No.	WU						

PREFACE

TO THE THIRD EDITION

The Second Edition being now exhausted, I have, in revising for this Third Edition, followed the same lines as before, and again amplified in many places. A few portions have been re-written, and a new chapter has been added on some notes of Hygiene of the Mouth.

ERNEST B. DOWSETT.

August 1912.

4381

PREFACE

TO THE SECOND EDITION

The Second Edition of this little book having been called for, I have, in compiling the same, adhered strictly to the general scheme of its predecessor, but have amplified and given further explanatory notes in many places. I would like to express my thanks to Mr. W. A. Maggs for kindly reading through the manuscript of this edition, and making some valuable suggestions.

ERNEST B. DOWSETT.

February 1910.

FILE COPY

CLAUDIUS ASH, SO 18 & C° LIMITED.

5 TO 12. BROAD ST., GOLDEN SQUARE, W.1.

PREFACE

TO THE FIRST EDITION

I have no hesitation whatever in adding to our list of Dental Literature what is intended to be a condensation of Dental Surgery, but I feel that the production of such a book on such a subject necessitates a slight explanation.

Although there may be many who object theoretically to the principle of presenting any subject in condensed book form, nevertheless practically, as a teacher of Revision Classes, and as a tutor for the Final L.D.S. Examination for the last nine years, I have long felt the want of such a book as I have endeavoured to compile. And the many students that have passed through my hands have so frequently asked me to produce in book form the notes I have been in the habit of giving them, that it is principally with the object of assisting them in their much stocked curriculum, that I venture to present this note-book. I have always believed in strict classification and tabulation as an aid to learning, and therefore I have endeavoured to make them prominent.

I have not attempted to detail the ordinary routine

of Conservative Dentistry, as that does not lend itself to any condensation.

Apart from the intended utility to the Dental Student, I hope also that the book may be of some service to the practitioner at times for easy revision or reference.

ERNEST B. DOWSETT.

January 1909.

CONTENTS

CHAP.			PAGE.
I.	DEVELOPMENT OF THE JAWS		I
II.	ERUPTION OF THE TEETH		6
III.	IRREGULARITIES OF THE TEETH		13
IV.	BACTERIOLOGY OF THE MOUTH		27
V.	CARIES		32
VI.	DISEASES OF THE PULP		42
VII.	DISEASES OF THE PERIODONTAL MEMBRANE		50
VIII.	Injuries to the Teeth		66
IX.	DISEASES OF THE GUMS AND MUCOUS MEN	1-	
	BRANE OF THE MOUTH		71
X.	SALIVARY CALCULUS		83
XI.	SALIVARY FISTULA		87
XII.	ODONTOMES		89
XIII.	Odontalgia and Neuralgia		94
XIV.	SECONDARY AFFECTIONS DUE TO DISEASE	D	
	TEETH		97
XV.	EMPYEMA OF THE ANTRUM		99
XVI.	NECROSIS OF THE JAWS		103
XVII.	FRACTURES OF THE JAWS		107
VIII.	DISLOCATION OF THE TEMPORO-MAXILLAR	Y	
	ARTICULATION		III
XIX.	INABILITY TO OPEN THE MOUTH		113
XX.	SWELLINGS ABOUT THE JAWS		115
XXI.	CLEFT PALATE		119

CONTENTS

CHAP.		PAGE.
XXII. DANGERS ATTENDING THE ADMINISTRATION	OF	
$ m N_2O~Gas$		123
XXIII. HÆMORRHAGE AFTER TOOTH EXTRACTION		125
XXIV. DRUGS COMMONLY USED IN DENTAL SURGE	RY	129
XXV. TREATMENT OF HYPERÆSTHESIA OF DENTIN	E-	
DEVITALISATION OF THE PULP WI	ТН	
Arsenious acid—Bleaching of Tee	TH	
DISCOLOURED BY DEATH OF PULP.		143
XXVI. THE USE OF X-RAYS IN DENTAL SURGERY		148
XXVII. HYGIENE OF THE MOUTH		150
INDEX		151

DENTAL SURGERY NOTES

CHAPTER I

DEVELOPMENT OF THE JAWS

MAXILLA

From 7 centres—5 Primary.
2 Secondary.

Primary :-

- I. Facial surface.
- 2. Posterior part of alveolar process. 7th week.
- 3. Orbital plate.
- 4. Palatal process.5. Intermaxillary bone. Few days later.

Secondary :-

- 1. Nasal process.
- 2. Malar process.

All join up by 10th week, when bone consists of 2 portions:—

- I. Larger part made up of 6 centres.
- 2. Premaxilla.

These separated by suture, which closes about time of birth.

MANDIBLE

Developed principally from *Membrane*, but partly from *Cartilage*.

Meckel's Cartilage.

Atransitory cartilage formed in the first Branchial Arch. It disappears as ossification progresses, but small piece persists as the *malleus*, except which it has entirely gone by end of 7th month.

Centres :-

- "Dentary"—on outer surface of Meckel's cartilage, just below future mental foramen.—
 For body of the bone—5th week.
- "Splenial"—in membrane on inner surface.
 —For inner walls of sockets.
- 3. "Mento-Meckelian" -- for chin.
- 4. For Condyle.
- 5. For Coronoid process.
- 6. For Angle.

Condition of Jaws at Birth:

- Halves of mandible separated by fibrocartilage, allowing free movement.
- Alveolar depressions show wide open spaces, from which teeth drop out in dry specimens.
- At posterior walls of sockets of centrals are shallow depressions for pulps of successors.
- 4. Condyle on level with alveolar margin.
- 5. Coronoid process makes angle of 135° with body of jaw.
- 6. Inferior dental nerve and artery seen at bottom of sockets of lower molars.
- 7. Large shallow depression for 1st and 2nd temporary molars in common.
- 8. Shallow depression extending backwards for 1st permanent molar germ.

- Angle little below level of inferior margin of jaw.
- Antrum is mere depression on outer wall of nasal fossa.
- II. Only thin plate of bone between orbit and mouth.

Teeth at Birth :-

- 1. 3 crown of central temporary incisors.
- 2. $\frac{1}{2}$ crown of lateral temporary incisors.
- 3. Tip of temporary canine.
- 4. Grinding surface of 1st temporary molar.
- 5. Cusps in form of a ring of 2nd temporary molar.
- Anterior cusp of 1st permanent molar commencing.

3 Months after Birth:

Teeth do not fall out in dry specimens, as orifices of crypts are closing over.

8 Months after Birth:

- Eruption of central incisors by absorption of anterior wall of process.
- 2. Bony union of halves of mandible.
- 3. Condyle rising.
- 4. Antrum occupies inner \(^2\) of floor of orbit, and thus considerable space between orbit and mouth.

After 8 Months of Age:

Age of jaw ascertained by position of teeth.

Dates of eruption approximately:-

Centrals . . 7 months.

Laterals . . 9 ,,

Canines . . 18 . ,,

ıst molar . . 12 ,,

2nd molar . . 24 ,,

Lower teeth generally erupt little before upper ones, except perhaps the laterals.

METHOD OF GROWTH OF THE MANDIBLE

1. In Length.

Alveolar process and jaw made longer by absorption in front of coronoid process, and deposition of bone on posterior aspect of vertical ramus, from the condyle to angle of the jaw, actual observation of which can be seen.

2. In Height.

Jaw deepens vertically by alveolar process growing upwards with the teeth:—

a. With temporary teeth (some of this absorbed).

b. With permanent teeth.

Also before age of 7 years, jaw deepens slightly by deposition of bone on base of jaw.

Above can be proved by comparing sagittal sections of jaws of varying ages. Taking the inferior dental canal as the fixed point, it will be seen that up to 7 years the bone is deposited above and below the level of the canal, and after that age only above the canal, which remains after 7 years at an unvarying distance from the base of the jaw.

3. In Thickness.

Posterior plate of mandible remains unchanged, growth going on by bone thickening from within outwards. This is known by the following measurements, &c.:—

a. If a jaw of a 9 months' fœtus be placed over an adult jaw, it will be seen that the inside arches of both are the same.
 (This is a rough-and-ready method of showing that the jaw grew in thickness by deposition of bone on the outside.)

- b. Line joining septum between 1st and 2nd temporary molars on either side remains constant at all ages.
- c. Line joining middle point of the above line with superior genial tubercles remains constant also.
- d. If an adult jaw be shaved down from the outside in the region of the mental foramen, until it is the thickness of a jaw at birth, it will be found, by measuring with a pair of "dividers," that the two mental foramina are the same distance apart and looking in the same direction (forwards) as in the jaw at birth. As the bone is deposited upon the outside of the young jaw, the mental foramina and canal change their direction, passing outwards, upwards, and backwards.
- All the above measurements tend to prove that bone was deposited on the outside of the jaw.

CHAPTER II

ERUPTION OF THE TEETH

- A. Temporary Dentition.
- B. Permanent Dentition.

A. ERUPTION OF THE TEMPORARY TEETH

When a tooth is about to be erupted, active absorption of its crypt begins, particularly on its anterior surface, bone behind remaining for part of crypt for successional tooth. As soon as the crown has passed through the free orifice so formed, bone is deposited and loosely surrounds the neck of the tooth, keeping pace with the elongation of root.

Front teeth are erupted first and then back ones; for if all were erupted at same time, teeth could only be closed at the back of mouth. Hence "proper mutual antagonism."

About 6th year temporary teeth all slightly separated from each other and pushed forward, *possibly* by increased size of crypts for permanent teeth.

Theories of Eruption: -

- 1. Lengthening of Root—evidence against:
 - a. Tooth with fully formed root may not be erupted.
 - b. Canine travels greater distance than length of its roots.
- 2. Growth of Bone at Bottom of Socket—none demonstrable.

- 3. Constant's Theory—increased vascularity of membrane at growing base; thus engorged vessels force up tooth by blood pressure.
- 4. Coleman's Theory—"Bone Currents"—i.e. presence of a tooth causing bone to grow in its vicinity.
- 5. Contraction of Fibres of Dental Ligament—can only be an adjunct to some other force.
- 6. General Growth of Bone—carrying teeth to the surface.
- Delabarre's Theory—analogous to a Gubernaculum; the fibrous remains of the neck of of the enamel organ contracting and pulling tooth out of crypt.

DATES OF ERUPTION AND FORMATION

			Erupt.		Roots complete.		Absorption begins.			
Central			7	7 months.		3 years.		$4\frac{1}{2}$ years.		
Lateral			9	11	3	,,	$4\frac{1}{2}$,,		
Canine			18	1)	5	,,	$6\frac{1}{2}$,,		
1st molar			12	,,,	4	,,	$5\frac{1}{2}$,,		
2nd molar			24	,,	5	,,,	$6\frac{1}{2}$,,		

- Erupt early in—
 a. Syphilitic children (some may erupt before birth).
 - b. Tuberculous children.

Erupt late in-

- a. Rickety children.
- b. Unfavourable hygienic surroundings.
- c. Cretinism.

NORMAL SIGNS OF APPROACHING DENTITION

- I. Increased flow of saliva.
- 2. Tendency of child to bite.
- 3. Gums tense and shining.
- 4. Mouth hot.

COMPLICATIONS OF ERUPTION OF TEMPORARY DENTITION

Local :-

- Acute inflammation of gums, especially over tooth.
- 2. Ulceration; if extensive, called "Odontitis Infantum."
- Cyst over erupting tooth, especially in rickety children.

Treatment-

Lance gums.

Astringent to ulcer.

Purge.

Potassium chlorate gr. j., t.d.s.

Incise cyst.

General :-

I. Nervous System-

Restlessness.

Sleeplessness.

Paralysis.

Convulsions.

Treatment-

Warm mustard bath.

Purge.

Brandy (few drops every half-hour)

2. Respiratory System-

Cough.

Bronchitis.

3. Alimentary System—

Diarrhœa.

Constipation.

Vomiting.

4. Integumentary System-

Erythema.

Eczema.

Herpes.

Impetigo.

5. Genito-urinary System-

Increased or diminished urine.

Urethritis.

6. Coryza.

It may be noted that "every ill that flesh is heir to" in a baby has been attributed to teething. But with the advance in the scientific feeding of infants, there is much evidence to prove that many of the affections that have been put down to teething are really largely due to errors of diet, the teething process only acting possibly as a pre-disposing factor in these cases by the pain and consequent exhaustion, causing a reduction in the general vitality of the child. The nervous complications are certainly more likely to be directly due to the teething, by the stimulus of the erupting tooth acting reflexly upon the nervous system.

ABSORPTION OF ROOTS OF TEMPORARY TEETH

Dates of absorption given above.

Process.

Root at or near its end becomes excavated by cupshaped depressions (*Howship's Lacunæ*), which deepen and coalesce, and thus root is eaten away.

Generally begins on side next the successional tooth, but not always—hence independent of pressure.

Cementum is attacked first, then dentine, and sometimes then enamel.

That part of dentine next pulp is more resistant than other parts, and hence often persists for some time as a hollow column.

Absorption depends upon the integrity of the pulp. If dead, root is only slightly absorbed or not at all. Hence so-called "necrosis of roots of temporary teeth" and protrusion through the gum.

Rate of absorption not constant. May stop for a time and give rise to deposition of cementum, and these processes may go on alternating, the absorption finally exceeding the deposition.

Theories of Absorption.

I. Absorbent Organ of Tomes.

Very vascular soft tissue applied to excavations caused by absorption. Surface may have large cells, somewhat like myeloid or giant cells, each of which fits into an indentation of excavated surface.

- 3 layers of organ
 - a. Large giant cells.
 - b. Ordinary nucleated cells.
 - c. Fibrous tissue.

Action of organ. Theories:-

- a. Cells put forth amœboid processes.
- b. Cells secrete an acid fluid.
- c. Inflammatory process.
- 2. Pressure Theory.

Compare absorption of lower 2nd molar by advancing 3rd molar. Really independent of pressure, as given above.

3. Cannibal Theory.

Eaten up or assimilated by advancing tooth. This cannot apply to absorption of permanent teeth.

B. ERUPTION OF THE PERMANENT TEETH

Analogous to that of milk teeth, an orifice very much larger than crown of tooth being opened up. Hence slightest force directs the rising crown; e.g.:—

- a. Fragment of root of temporary tooth.
- b. Action of lips and tongue.

Temporary teeth stood vertically, but permanent teeth stand obliquely forwards, thus making room for canine, which was out of line during development.

When crown has erupted and reformed bone is enclosing neck, little of root is complete, the *short* and *widely open* root occupies the whole depth of socket, and so as root gets longer its socket is deepened by additions to its *free edges*, and not by the root burrowing deeper into the alveolar process. Thus the sockets grow up with and are moulded round the teeth as they elongate, and therefore the whole of the alveolar process is subservient to the position of the teeth.

A symmetrical arch is moulded by the lips and tongue, giving constant and equal pressure on each side.

DATES OF ERUPTION AND FORMATION

Central incisor		Erupt. 7 years.		Roots Complete.		
		8	,,	II	"	
		12	,,	13	,,	
First premolar		IO	,,	12	11	
Second ,,		II	"	12	,,	
First molar		6	,,	9	, ,	
Second ,,		13	,,	16	,,,	
Third ,,	. I	8-24	"	20	,, .	

RETENTION OF TEMPORARY TEETH WITH PERMANENT TEETH

In order of frequency:—
Upper canines.
Lower 2nd molars.
Upper 2nd molars.
Lower incisors.

May remain till quite late in life.

Results :-

May remain at original level.

May rise to occlusion.

Gum may grow partially over.

May be wedged between two permanent teeth.

Caries from wedging.

Irregularities of permanent teeth.

Treatment :-

Take skiagraph to ascertain position of successor, if unerupted.

a. If in a position to erupt, extract temporary tooth.

b. If absent or much out of position, retain temporary tooth if healthy.

CHAPTER III

IRREGULARITIES OF THE TEETH

A. Temporary Dentition.

B. Permanent Dentition.

Abnormalities in-

- I. Size.
- 2. Number.
- 3. Structure.
- 4. Position.

A. TEMPORARY DENTITION

1. Abnormalities in Size:

Upper canine.
2nd temporary molar. Often larger.

2. Abnormalities in Number:

Supernumerary and supplemental teeth. Usually in incisor region.
Cases on record of total absence.

3. Abnormalities in Structure:

Additional cusps.

Additional roots—2nd molars and lower canine. Honeycombed (rare).

Geminated. Most common in lower incisors and canines. (See below.)

4. Abnormalities in Position:

Teeth—overlapping and twisting. Jaws—protruding and open bite.

B. PERMANENT DENTITION

I. ABNORMALITIES IN SIZE

Abnormally large teeth are generally:— Upper central.

Lower 2nd premolar.

Lower 2nd molar.

Abnormally small teeth are generally:—
Upper lateral.
Upper 3rd molar.

2. Abnormalities in Number

- a. Supernumerary teeth.
- b. Supplemental teeth.
- c. Absence of some or all.

a. SUPERNUMERARY TEETH

Most common in-

- (i) Upper incisor region.
- (ii) Upper 3rd molar region.

Characteristics :-

- (i) Generally single root.
- (ii) Enamel ends straight round middle of tooth.
- (iii) Any form from simple conical tooth to small molar.—With the latter the cusps and roots are rather convergent than divergent.

Theories of Formation:

- (i) Dichotomy—i.e. complete unequal division of tooth germ (Sutton).
- (ii) From masses of epithelium or epithelial pearls (Mallassez).
- (iii) Additional tooth germ from tooth band (? Atavism).

b. SUPPLEMENTAL TEETH

Most common in-

- (i) Upper and lower lateral incisor region.
- (ii) Premolar region.

Characteristic.

Simulate the teeth next to which they stand.

Theory of Formation.

Dichotomy, with complete equal division of tooth germ.

c. ABSENCE OF TEETH

Most common in-

- (i) Upper lateral incisor.
- (ii) Lower 2nd premolar.
- (iii) Upper 3rd molar.

Occasionally seen-

- (i) Absence of several teeth.
- (ii) Absence of all teeth. Rarely seen.

These often associated with other integumentary abnormalities.

3. Abnormalities in Structure

Forms:

- a. Syphilitic teeth.
- b. Stomatitic teeth.
- c. Geminated teeth.
- d. Enamel nodules.

a. Syphilitic Teeth

Found occasionally in the subjects of congenital syphilis.

Characteristics:

Permanent Dentition-

- (i) Narrowing of tooth from gum to cutting edge.
- (ii) Semilunar notch at cutting edge of upper incisors, occupied on eruption by frail enamel, which soon wears away, leaving the notch. Laterals are not always affected.
- (iii) Margins of upper incisors are rounded and illdefined, and distal edges turned outwards.
- (iv) Peg-top shape to lower incisors.
- (v) All teeth small, and spaced as a rule.
- (vi) Implantation very weak.
- (vii) Dome-shaped 1st molars with dwarfed, rounded cusps, and sunken occlusal surfaces.
- (viii) Enamel, normal in colour and thickness.
 - (ix) Lime salts deficient—hence soft and prone to caries.

Temporary Dentition.

Not as a rule affected. May be small and dusky hue, indicating bad calcification.

Coles reports a case where incisors had characteristic markings, as in Permanent Dentition.

Causation.

Very little understood.

Hutchinson says it is due to specific stomatitis occurring about time of or just before birth. This would affect all teeth developing at that time, but this apparently is not found.

Dentine is much more affected than enamel.

b. STOMATITIC TEETH

Also called-

Hypoplastic teeth.

Honeycombed ,,

Mercurial

Rocky

Craggy

Teeth affected.

Generally—

Incisors.

Canine (tip).

ist molar.

Sometimes—

Premolars.

and molars.

All teeth.

Characteristics :-

- (i) Lines of dots or pits transversely across teeth, the number varying.
- (ii) Shallow grooves transversely across teeth may be found.
- (iii) Molars—the contoured portion presents a marked shoulder or rim, and cusps are merely spinous projections.
- (iv) Brown stain when enamel is worn.

Microscopical appearances:

- (i) Enamel thin or absent in grooves and pits.
- (ii) Prisms and "striæ of Retzius" well marked.
- (iii) Prisms may be granular.
- (iv) Some parts—enamel is homogeneous brown mass.
- (v) Interglobular spaces well marked.

Causes :-

(i) Any illness that interferes with calcification, while it is in progress.

Especially exanthematous fevers, as those are the commonest severe illnesses from which a child suffers.

(ii) Bad feeding at same periods.

(iii) Hutchinson says it is due to mercury given during first two years of life.

c. GEMINATED TEETH

Forms :-

- I. True or Developmental Gemination.
- 2. False or Pathological Gemination.

1. True Gemination.

Causes:

- (i) Close apposition and fusion of tooth germs—giving rise to gemination of two normal teeth of the arch.
- (ii) Incomplete dichotomy—giving rise to gemination of one normal tooth with a supernumerary or supplemental.

Varieties in order of frequency :-

Fusion of two incisors—especially in temporary dentition.

Fusion of incisor and supernumerary.

- molar and supernumerary.
- ,, two molars.
- ,, lateral and canine
- ,, canine and premolar.
- ,, two premolars. V

Union :-

(i) By whole length of tooth (most common).

There is then continuity of enamel, dentine, and cementum, with either one or two pulp chambers.

(ii) By crowns only.
There is then continuity of enamel and dentine, with either one or two pulp chambers.

(iii) By roots only.

There is then continuity of dentine and cementum, with either one or two pulp chambers.

2. False Gemination.

Cause.

Chronic inflammation of periodontal membrane, leading to organisation, i.e. productive periodontitis.

Varieties.

Any adjacent teeth may be united.

Common in molars—especially upper 2nd and 3rd.

Union.

By cementum only.

d. ENAMEL NODULES

Enamel excrescences on roots of teeth.

Site.

Generally upon molars between roots.

May be found occasionally on single-rooted tooth.

Structure.

Generally core of dentine connected with that of the root, covered with thick enamel.

Occasionally small pulp cavity found.

May be enamel only, and outside the cementum.

Origin.

If enamel only—formed by portion of "sheath of Hertwig" developing enamel.

If with dentine core—formed by dichotomy of tooth germ. Therefore transition from enamel nodules to supernumerary teeth is only one of degree.

e. Abnormalities in Position

Types described:-

- a. Superior Protrusion.
- b. Inferior Protrusion.
- c. Open Bite.
- d. Misplaced Lower 3rd Molar.

a. Superior Protrusion

Causes :-

(i) Connected with Maxilla.

Excessive development—

Alveolar process.

Whole bone.

Overcrowding of teeth.

Supernumerary teeth.

Narrow or small arch.

Large teeth.

(ii) Connected with Mandible.

Deficient development.

Crowding of lower incisors-fan-shape.

Retardation of eruption of molars, causing mastication to take place on front teeth.

Early extraction of first permanent molars doing the same as above.

(iii) Connected with Soft Parts.

Attached frænum.

Short upper lip, and hence absence of anterior support to upper incisors.

(iv) Habits of Children.

Thumb, finger, toe, lip, and tongue sucking. Rubber teat.

(v) Obstructed Respiration.

Theories of action:-

- a. Tension at sides of mouth by being open.
- β . Absence of tongue support to upper.
- γ. Under development of nasal fossæ; their function being lost from want of use. Hence floor of nose, i.e. palate, is higher and narrower than normal, and so sides are drawn together and anterior part protruded.
- δ. A tmospheric pressure greater in the mouth than in the nose with inspiration. Hence palate forced up.
- (vi) Fault in Temporo-maxillary Articulation. Causing posterior bite (?).

Treatment.

Observe :-

- (i) Facial contour and profile: to determine which parts of arches at fault, and therefore which, if possible, to be corrected.
- (ii) Molar occlusion: to determine whether lower first molar is in normal or distal occlusion to first upper.
- (iii) Incisor occlusion :-
 - Class A. Plenty of space between upper and lower incisors, for the former to be retracted.
 - Class B. Lower incisors biting in such a situation that the upper ones would be bitten out again if merely retracted.

Class A.

Methods available:-

- a. Retract upper incisors, when-
 - (i) Normal molar occlusion.
 - (ii) Only slight distal molar occlusion.

Appliances used:

- (i) Angle's wire arch or modification, with or without skull-cap.
- (ii) Badcock's screw plate and spring wire arch.
- (iii) Rubber band with plate.
- (iv) Wire bows on plate.

(v) T-piece on plate, &c.

If more space than already exists in upper is necessary for this treatment, decide "expansion" or "extraction" according to:—

- (i) Facial contour—expansion broadens face.
- (ii) Amount of space wanted—expansion gives less than extraction.
- (iii) Time at patient's disposal—expansion takes longer.
- (iv) Direction of roots—expansion may tilt teeth too much.

With Expansion—generally necessary to expand lower jaw as well.

Appliances used :-

- (i) Coffin's spring plate.
- (ii) Read's screw plate.
- (iii) Badcock's screw plate.
- (iv) Angle's arch.
- (v) Jack-screw.

With Extraction, be guided by:--

- (i) Which teeth carious.
- (ii) Amount of space wanted.

(iii) Distance from irregularity.

(iv) Possibility of space left after retraction.

b. Jump the bite, when-

Distal molar occlusion is great.

Appliances used :-

- (i) Reciprocal traction with rubber bands. (Baker anchorage, &c.)
- (ii) Anterior inclined plane.

Means of retention :-

- (i) Interlocking wedges on buccal surface of molars.
- (ii) Linked chain anchorage.
- (iii) Inclined plane.

Class B .- Preliminary Treatment.

Methods available:-

- a. Insert bite plate in upper, capping premolars, leaving molars free to rise and occlude, and which also tends to depress lower incisors.—Generally the most satisfactory.
- b. Grind down and polish lower incisors, if only slight amount necessary.—Temporary benefit.

c. Extract one of lower centrals and inser bite plate to depress the others.

d. Extract unerupted premolars, if case seen early enough, and insert bite plate.

After any of these methods of *preliminary* treatment, the case is resolved into *Class A*, and is then further treated as such.

Causes: b. Inferior Protrusion

(i) Connected with Mandible.

Excessive development.

Oblique ramus, causing horizontal part to be pushed forward.

(ii) Connected with Maxilla.

Deficient development.

Tardy eruption of upper incisors.

Instanding upper incisors, from late retention of temporary incisors.

(iii) Habits of Children.

Hooking fingers over lower incisors. Protruding lower jaw.

Treatment:

- (i) Retract lower incisors, extracting premolars if necessary—Generally the most satisfactory.
- (ii) Expand upper jaw.—Only when upper jaw is at fault.

(iii) Separate all upper teeth (Kingsley).

(iv) Skull-cap and chin bandage, if due to habit. —Must be done very early, if done at all, and then the result is doubtful.

c. OPEN BITE

Causes :-

(i) Elongation of molars.

a. Keeping mouth open—
Obstructed respiration.
Cicatrix.

 β . Not capped on regulation plate.

(ii) Congenital syphilis— Small incisors.

Small alveolar process.

- (iii) Thumb and finger sucking, &c.
- (iv) Rubber teat.
- (v) Imperfect development of maxilla and premaxilla.
- (vi) Small or oblique ramus.
- (vii) After fracture—malunion.
- (viii) Hypertrophied tongue.

- (ix) Changes in joint-Dislocation. Osteo-arthritis.
- (x) Scar tissue in front part of neck in child, by contraction, moulding anterior part of mandible downwards.

Treatment:

- (i) Extraction of occluding teeth.—Generally the most satisfactory.
- (ii) Grind down occluding teeth; several sittings with two or three weeks' intervals.-Expedient only when open bite is slight.
- (iii) Chin bandage and skull-cap to force occluding teeth into sockets.—Very tedious and not as as a rule successful.
- (iv) Elongation of upper incisors by banding and pulling to wire bow.—Can only be elongated short distance.
- (v) Treat obstructed respiration.—If case recognised early enough.

d. MISPLACED LOWER THIRD MOLAR

Directions :-

- (i) Obliquely forwards against second molar.

 (ii) Directly forwards against "Impacted."
- second molar.
- (iii) Inwards.
- (iv) Upwards (may reach sigmoid notch).
- (v) Downwards (inverted).

Symptoms and Results.

- (i) Pain
 - a. In tooth—throbbing, jumping, or continuous.

 β In gum—worse on eating.

γ. In another tooth—referred.

c. In second molar.

From pressure.

From absorption.

- ε. Neuralgia—especially in ear—Due to the common origin of the auriculo-temporal and inferior dental nerves from the posterior division of the inferior maxillary nerve.
- (ii) Swelling of gum-painful to bite on.
- (iii) Ulceration of gum.
- (iv) Suppuration in tissues surrounding crown of tooth leading to-

a. Swelling of face. d. Swelling of fauces.

β. Swelling at angle. ε. Swelling of pharynx.

y. Swelling of neck.

Any of which swellings may suppurate.

- (v) Lymphatic glands in neck swollen.
- (vi) Trismus.
- (vii) Constitutional symptoms—

a. Pyrexia.

y. Malaise.

β. Rapid pulse. δ. Constipation, &c.

Treatment: -

- (i) Take skiagraph, if position uncertain.
- (ii) Lance gum by crucial incision, if plenty of room and gum only involved.
- (iii) Extract offending tooth with straight elevator, if symptoms warrant.
- (iv) Extract second molar, if impossible to remove third
 - a. To relieve pressure;
 - β . As preliminary to removal of latter, if necessary.

CHAPTER IV

BACTERIOLOGY OF THE MOUTH

Bacteria, according to shape and arrangement, are thus classified:—

- Bacilli—rods.
- 2. Cocci-round bodies :
 - a. Streptococci—chains.
 - b. Staphylococci—little packets.
 - c. Diplococci—pairs.
 - d. Sarcinæ—four in a square.
- 3. Spirilla—spirals.
- 4. Leptothrix—thread forms.
- 5. Streptothrix—fine threads with lateral branches.

MOUTH BACTERIA

In the human mouth Miller found and cultivated nearly 100 different forms—some pathogenic (disease giving) and some non-pathogenic.

But six forms he described as almost constant, and all having the same peculiarity—that no culture medium was found on which they would grow. These are:—

- I. Leptothrix innominata. Found in soft white deposit on teeth. Stains faint yellow with iodine.
- 2. Bacillus buccalis maximus. Rods and threads 2–10 μ long and 1–1.3 μ wide. Stains deep violet with iodine.

- 3. Leptothrix buccalis maximus. Same form as above, but no iodine reaction.
- Iodococcus vaginatus. Common in unclean mouths. Chains of cells in sheath. Cells stain violet and sheath very little with iodine.
- 5. Spirillum sputigeneum. Found on inflamed gums. Form is like a comma.
- Spirochete dentium. Found on inflamed gums. Long spirals, 8-25 μ long.

To these may be added:-

- Leptothrix Racemosa (Vicentini). Found in soft white deposit. Felted masses of entwined threads, with "fruitful heads," i.e. spore-like areas.
- 8. Strepothrix buccalis (Goadby). Found in soft white deposit, and pyorrhœa. Filamentous forms with lateral branches, whose ends are often club-shaped.

Amongst the many others found in the mouth the following ordinary pathogenic bacteria are frequently observed:—

- 1. Pneumococcus.
- 2. Diphtheria bacillus (Klebs-Læffler).
- 3. Bacillus Coli Commune.
- 4. Streptococcus pyogenes.
- 5. Streptococcus brevis.
- 6. Bacillus pyocyaneus.
- 7. Micrococcus tetragenous.
- 8. Tubercle bacillus.

Miller estimated in one very unclean mouth as many as 1,140,000,000 organisms.

BACTERIOLOGY OF CARIES

Some of mouth bacteria can ferment grape sugar into lactic acid.

Others of mouth bacteria can peptonise, i.e. render soluble, albumen, gelatin, and fibrin.

Hence present view of cause of Caries:-

Certain mouth bacteria are capable of generating an acid which can dissolve the salts out of the tooth substance; while others are capable of dissolving the decalcified matrix—i.e. the organic portion remaining after the salts have gone.

Acid forming Bacteria.

Streptococcus brevis.

Staphylococcus necrodentalis. In deep layers.

Staphylococcus albus.

Streptococcus brevis.

Sarcina lutea.

Sarcina aurantiosa.

Sarcina alba.

Staphylococcus albus.

Staphylococcus aureus.

In superficial layers.

Bacteria capable of liquefying dentine (decalcified) :-

Bacillus mesentericus ruber.

,, vulgatus. ,, fuscus.

,, furvus.

", gingivæ pyogenes.

,, liquefaciens flourescens motilis.

, subtilis.

Proteus Zenkeri.

Bacillus plexiformis.

All in superficial layers.

BACTERIOLOGY OF PULPITIS

Forms of Bacteria found (Miller).

- I. Cocci and diplococci (pathogenic).
- 2. Bacilli-curved and often forming threads.
- 3. Short rods with bipolar staining.

Individual Bacteria described.

- 1. Streptococcus (either pathogenic or non-pathogenic).
- 2. Staphylococcus albus and aureus.
- 3. Micrococcus tetragenous.
- 4. Sarcinæ. (Zierler).
- 5. Bacillus gangrenæ pulpæ. (Arkövy).
- 6. Bacillus mesentericus vulgatus 7. ,, ruber. Goadby, and Sieberth.
- 8. ,, fuscus.

Goadby holds that 5 and 8 are identical.

BACTERIOLOGY OF ALVEOLAR ABSCESS

Cocci mostly found, but not as a rule the common pus cocci (staphylococcus aureus), which are rarely found.

Bacteria commonly found.

- 1. Staphylococcus viscosus. (Goadby.)
- 2. Staphylococcus albus.
- 3. Bacillus gangrenæ pulpæ. (Arkövy.)
- 4. Bacilli of mesenteric group.

Bacteria occasionally found.

- 1. Micrococcus tetragenous.
- 2. Streptococci.
- 3. Bacillus Coli.
- 4. Blastomycetes.
- 5. Staphylococcus aureus.

BACTERIOLOGY OF PYORRHŒA ALVEOLARIS

Forms of Bacteria found (Goadby).

- I. Cocci—generally diplococci.
- 2. Bacilli
 - a. Thick and often jointed.
 - b. Thick, with pointed ends.
 - c. Fine, with irregular banded markings.
 - d. Masses, with some jointed in chains.
- 3. Spirilla and spirochæte.
- 4. Yeast forms—various.
- 5. Streptothrix—with well-marked club ends.

Ordinary pus cocci (staphylococci), rarely found.

None of the organisms found have yet been shown to
be especially related to the disease.

CHAPTER V

CARIES

DISINTEGRATION of tooth substance extending from without inwards towards the pulp, caused by external agencies.

MACROSCOPICAL APPEARANCES

Enamel.

If starting in a crevice—a dark spot.

If starting on a plain surface—a white spot.

Starts usually by enamel losing its translucency, whereby a whitish spot appears, becoming darker if process is slow, and remaining white if rapid.

Surface roughened and disintegrated by the disorganised enamel prisms being washed away. If undermined by caries it appears bluish-white.

Dentine.

This is now reached.

It is first softened and then disintegrated and washed away, forming a cavity; rapidity of process depending upon:—

- a. Structure of tooth.
- b. Virulence of micro-organisms.
- c. Intensity of ferments.

Rapid or Spreading Caries (Caries acuta).—If calcification bad, or many interglobular spaces, caries spreads rapidly sideways, undermining the enamel, which is then attacked from the dentinal side, and is called "secondary enamel decay" (Miller).

This kind of caries is also generally moist (Caries humida), and light yellow or white (Caries

alba).

Slow or Penetrating Caries (Caries chronica).—If calcification is good, caries only extends in directions of dentinal tubes. The shape of the cavity is then conical, with apex towards pulp.

This kind of caries is also generally dry (Caries sicca), and brown or black (Caries pigmenta), which is probably produced by chromogenic

bacteria.

Translucent Zone.

A more or less translucent condition of the dentine, preceding the advance of the caries, and found between the caries and the pulp cavity.

Forms-

- a. Regular zone.
- b. Irregular cone—apex to pulp.
- c. Stripes and spots.

Resembles roots of teeth of aged persons.

Not a phenomenon of caries alone, because it is seen when only enamel is worn.

Has never been produced artificially.

Due to-

Some change that renders the refractive indices of matrix, sheaths, and fibrils very nearly alike, *i.e.*—

- a. Calcification of contents of tubes.
- b. Partial decalcification of matrix.

Arguments in Favour of Calcification of Contents of Tubes:—

- Diameter of tubes is lessened in translucent zone, and not permeated by Charters White's celloidin stain.
- Takes stain with difficulty, whereas even very slightly decalcified dentine stains readily.
- Cannot be produced artificially, which should be easy.
- 4. Not found in carious, pulpless teeth.
- Not found in natural teeth worn as artificial substitutes, when caries supervenes.
- 6. Occurs under worn enamel.
- 7. Decalcification advances in a regular line, whereas this may be very irregular.

Cementum.

Caries may attack any portion exposed to mouth. At first softened and later wide, shallow cavities formed.

May be deep cavity at angle between roots.

MICROSCOPICAL APPEARANCES

Enamel.

Felt-like mass of micro-organisms on surface. Interprismatic substance dissolving away from surface inwards.

Prisms therefore loosened, partially separated, and unduly distinct.

Prisms may appear granular and pigmented if process slow.

Bacteria between prisms.

Dentine.

Mass of debris on surface.

Irregular spaces (liquefaction foci) in dentine, near surface, full of broken-down pieces of dentine and micro-organisms, which have entered the dentinal tubes, bulged them, and finally broken through them by liquefaction of the matrix, which has been previously decalcified by acids that passed down the tubes.

Further in, dentinal tubes full of micro-organisms and bulged, but still intact. Also, tubes normal in calibre with micro-organisms.

These latter, and also many other tubes that contain no micro-organisms, present the "tobacco-pipe" appearance (seen on tranverse section of tubes).

This appearance is due to the partial decalcification of that portion of the matrix immediately surrounding the "sheath of Neuman," thereby rendering the sheath unduly distinct.

In translucent zone—rod-shaped elements and rows of shining granules in tubes, which may be looked upon as consolidated portions of fibrils.

Cementum.

Destruction takes place exactly in same manner as in dentine.

Micro-organisms penetrate principally along Sharpey's fibres, but also along canaliculi, and form liquefaction foci.

CAUSES OF CARIES

- 1. Predisposing
 - a. Structural.
 - b. Local conditions.
 - c. General conditions.
- 2. Exciting.

1. Predisposing.

- a. Structural Defects.
 - (i) Deficiency of lime salts:—
 Females.
 Rickets.
 Temporary teeth.
 - (ii) Presence of pits and crevices.
 - (iii) Increased number of interglobular spaces.
 - (iv) Passage of dentinal tubes into enamel— communicating with enamel spindles.

b. Local Conditions in the Mouth.

- (i) Irregularities of teeth, being inaccessible for cleansing.
- (ii) Crowded arch for same reason.
- (iii) Want of cleansing.
- (iv) Insufficient mastication—which, if properly done, tends to cleanse the teeth.
- (v) Wearing of dentures—collecting debris.
- (vi) Recession of gum.
- (vii) Mouth breathing—natural cleansing by saliva and lips lessened.
- (viii) Class of food:-

Carbohydrates (starches and sugars) especially deleterious.

Glucose, which is used very largely in most sweetened articles of commerce instead of cane or beet sugar is much more easily converted in the mouth into acid than the latter (see below).

Proteids and fats have little or no effect, e.g. Esquimaux—fat eaters; no caries.

(ix) Physical properties of food:—

Sticky foods and those from which fibrous material has been softened or eliminated are deleterious, e.g.:—

Sweets.

All cooked food.

Roller-milled flour.

These all tend to cling, and may be fermented into acid *in situ* on teeth.

(x) Abnormal secretions of mouth:—
Acid saliva in dyspeptics.
Sticky mucoid saliva in
Pregnancy.

Fevers.

Most illnesses.

(xi) Medicines :-

Mineral acids.

Iron salts.

c. General Conditions.

Besides those mentioned incidentally above under "Local Conditions":—

- (i) Hereditary influence.
- (ii) Race.
- (iii) Civilisation.
- (iv) Occupation :-

Millers. Bakers. Flour.

Fur-dyers, &c.:—
Acid fumes.

(v) Artificial feeding of infants.

(vi) Children brought up in districts poor in lime salts in food and water.

Immunity.

Besides all the above predisposing causes there is still some further undiscovered factor which is not understood at all, to explain the facts that sometimes perfect teeth under apparently perfect conditions will decay, and also that sometimes badly formed teeth, even under what are considered bad conditions, will escape caries for many years. Miller looked upon this immunity as due to some bacteriacidal property of the saliva, which, however, he could not determine. Also, it may be that certain saliva has the property of attracting leucocytes in the presence of bacteria.

2. Exciting Causes (in historical order).

a. Inflammatory Theory (Abbot).

That caries was set up by inflammation of living tooth substance internally.

Evidence against:-

- (i) If right, filling would promote caries.
- (ii) No caries without breach of enamel.
- (iii) Caries goes on in dead tooth.
- b. Electrical Theory (Bridgeman).

Crown—positive pole.
Root—negative pole.
Saliva—electrolyte.

Action set up by abnormal saliva.

c. Chemical Theory.

Due to acids of mouth (see below).

d. Parasitic Theory.

Due to micro-organisms alone.

e. Chemico-parasitic Theory (present accepted theory).

Due to action of acids of mouth and micro-organisms.

Decalcification by acids.

Dissolution or peptonising of the remaining organic substance by bacteria acting in an acid medium.

Origin of Acids of Mouth: -

- a. By fermentation.
- b. Introduced with food.
- c. Regurgitated from stomach.
- d. Secreted from mucous glands.
 - a. By Fermentation.

Partly by ptyalin of saliva.

Partly by bacteria themselves.

Sugar and other carbohydrates ferment into-

Lactic acid (principally).

Butyric acid.

Propionic acid.

Valeric acid.

Albumen may ferment into-

Butyric acid.

Valeric acid.

For fermentation process (see below).

b. Introduced with Food or Medicines.

Fruit, vegetables, drinks, &c., contain-

Citric acid.

Malic acid.

Oxalic acid.

Acid oxalates.

Acetic acid.

Tartaric acid.

Acid tartrates.

Alum.

Chloride of iron.

Perchloride of iron.

Sulphate of iron.

All of which have a greater or less dissolving action upon the dental tissues.

c. Regurgitated from Stomach.

Small quantities of-

Lactic acid.

Hydrochloric acid.

d. Secreted from Mucous Glands.

May be very small quantities in certain affections.

Fermentation of Carbohydrates.

Polysaccharides. Disaccharides. Monosaccharides.

 $(C_6H_{10}O_5)n$ $C_{12}H_{22}O_{11}$ $C_6H_{12}O_6$ Starch. Maltose. Dextrose. Dextrin. Lactose. Levulose.

Cellulose. Cane Sugar. Galactose.

Beet Sugar. Glucose.

Grape Sugar.

I. Polysaccharides.

Converted by Ptyalin of Saliva into Maltose thus:—

 $\begin{array}{c} \mathbf{10}(\mathsf{C}_{6}\mathsf{H}_{10}\mathsf{O}_{5})n + 4n\mathsf{H}_{2}\mathsf{O} = 4n(\mathsf{C}_{12}\mathsf{H}_{22}\mathsf{O}_{11}) + \\ \mathsf{Starch.} & \mathsf{maltose} + \mathsf{achroo-dextrin} + \mathsf{erythro-dextrin}. \end{array}$

These gradually change to maltose.

And thence into dextrose and later into lactic acid (see below).

N.B.—Cellulose (coating of starch grains) is not acted upon, and therefore uncooked starch is unaltered by saliva.

2. Disaccharides.

Converted into monosaccharides by hydrolytic ferment produced by some bacteria in mouth thus:—

 $C_{12}H_{22}O_{11}+H_{2}O = C_{6}H_{12}O_{6}+C_{6}H_{12}O_{6}$

(i) Maltose —dextrose —dextrose

(ii) Lactose =dextrose+galactose

(iii) Cane or beet sugar=dextrose+levulose and thence into lactic acid (see below).

Monosaccharides (directly fermentable sugars)
 Converted directly into lactic acid by bacteria thus:—
 C₆H₁₂O₆=2C₃H₆O₃ (lactic acid).

Artificial Production of Caries.

All phenomena can be reproduced except translucent zone.

Method-

Keep teeth or portions of teeth in mixture of *bread* and *saliva* for three months at 98° F., and renew it every two or three weeks. Otherwise putrefaction sets in and the mixture becomes alkaline, and then bacteria cannot act.

Resultant caries is *white*, but by allowing putrefaction at certain stages or by exposure to air or various coloured articles of food, all sorts of coloration appear.

In the mixture—

Bread is to supply the carbohydrates to ferment.

Saliva is to supply bacteria, ptyalin, and moisture.

CHAPTER VI

DISEASES OF THE PULP

In describing the inflammations of the Pulp, it is as well to point out clearly that there are no hard-and-fast lines between the different types of inflammation described, but that they are only thus classified for purposes of description. The different kinds merge into one another, and it is quite impossible to determine where one ends and another begins.

The types of inflammation of the pulp described below are:—

- Hyperæmia, or mild inflammation, extending over short period only, or intermittent with healthy intervals.
- Acute inflammation, or severe condition, extending over short period.
- Chronic inflammation, or variable condition, extending over long period.

HYPERÆMIA OF PULP (Pulpitis)

Causes: -

- 1. Thermal changes in mouth.
- 2. Advancing caries, erosion, attrition, abrasion.
- 3. Exposure of cementum.
- 4. Sweet and acid substances acting through 2 and 3.
- 5. Filling too near pulp.
- 6. Fracture of tooth.

- 7. Incautious use of As₂O₃, ZnCl₂, carbolic acid, &c.
- 8. Susceptibility on part of patient.

Symptoms :-

- I. Intermittent pain— Paroxysmal. Jumping.
- 2. Thermal shocks.
- 3. May be slightly painful to mastication (when hyperæmia has extended to periodontal membrane).

Treatment :-

- I. Remove cause.
- 2. Oil of cloves and temporary filling (oxysulphate).
- 3. Non-conducting filling.
- 4. Counter-irritant to gum.
- 5. Removal of pulp, if persistent.

Microscopical Appearances: --

- 1. Enlarged and tortuous vessels.
- Great increase of pulp cells due to proliferation of pre-existing cells and to escape of leucocytes.
- Increase of odontoblasts, which are arranging themselves in "sheaves" near region of irritation.
- Cylindrical calco-globulin formations in neighbourhood of vessels.

Pathological Results:

- I. If cause removed
 - a. Generally ends in resolution.
 - b. More rarely passes on to acute or chronic inflammation.
- 2. If cause persists—
- Passes on to acute or chronic inflammation.

ACUTE INFLAMMATION OF PULP (Acute Pulpitis) Causes:—

Same set of causes as for Hyperæmia (see above), only in a more pronounced degree or extending over longer period, especially if the pulp has become exposed, allowing irritation from foreign matter.

Also As₂O₃ for devitalising.

Symptoms:

1. Sharp throbbing pain, worse in recumbent position and more or less continuous.

2. Thermal changes increase pain, though cold may relieve in early stage.

If exposure of pulp—
 Wool in cavity causes pain.
 Jet of warm water causes pain.

4. Tooth may be little tender to pressure, it inflammation extended to periosteum.

Treatment:

Anæsthetise with cocaine, or Devitalise with arsenic.

- 2. If in early stage, may be saved by:
 - a. Dressings with—
 Obtundents.
 Astringents.
 Sedatives.
 - b. Capping with a non-irritant and non-con ducting filling. This method should only be adopted in a young and perfectly healthy subject, otherwise degeneration and death of pulp are likely to follow. It is a doubtful procedure, even under favourable circumstances.
 - c. Extract tooth if not saveable.

Microscopical Appearances:

 All the appearances as in Hyperæmia (see above), only more exaggerated.

2. Altered odontoblasts—shrunken or enlarged.

3. Nerve bundles lost their definite structure.

4. Minute abscesses with cells broken down, appearing as little blurred patches with broken centres in stained section.

5. Micro-organisms if exposed pulp or only thin layer of dentine covering.

Pathological Results (in the Pulp) :-

1. Resolution—rare.

2. Suppuration, common because of-

a. Delicate structure.

b. Inability to expand.

3. Passing into chronic inflammation.

4. Death *en masse* (gangrene) from strangulation of vessels at apex.

Changes during Resolution.

Cause ceases, or leucocytes overcome bacteria.

Bacteria if present and toxins absorbed by leucocytes.

Leucocytes and plasm pass back into capillaries.

Blood flow again becomes active.

New vessels formed, when some quite thrombosed, by budding off from endothelium.

Some leucocytes become fibricated and then converted into connective tissue.

Vessels return to normal size.

Changes during Suppuration.

Bacteria and toxins overcome leucocytes, which die.

They liberate proteolytic ferment, which dissolves cells and tissues.

This occurs in small patches, especially near exposure. These extend and coalesce, and finally whole pulp liquefies.

CHRONIC INFLAMMATION OF PULP

(Chronic Pulpitis)

Causes :-

Following Hyperæmia or acute inflammation of pulp from any of their causes as given above.

Symptoms:

Very variable.

Generally pain of intermittent and neuralgic character, but may be any degree of severity. May be painful to mastication.

May give rise to no symptons at all.

Treatment:

Removal of pulp.

Removal of tooth if not saveable.

Pathological Results (in the pulp):—

- I. Suppurative.
 - a. On surface—ulceration of pulp.
 - b. Complete—death of pulp.
- Destructive. Absorption of dentine of pulp cavity.
- 3. Productive.
 - a. Growth of pulp into absorbed dentine.
 - b. Growth of pulp into carious cavity (*Poly-pus*).
 - c. Calcification of above growth (see below).
- 4. Degenerative.
 - a. Fibroid changes.
 - b. Calcareous changes. (see below).
 - c. Fatty changes.

POLYPUS OF PULP (Hypertrophic Pulpitis)

Cause: —Chronic Pulpitis (see above).

Symptoms:

Fleshy mass in carious cavity. Bleeds easily. Insensitive.

Treatment :-

If tooth saveable, remove polypus and fill or crown tooth. (Apical foramen may be too large by absorption.)

If tooth not saveable, extract. Great care must be exercised, as tooth is often very fragile on account of absorption of pulp cavity.

Pathology: -

Pulp hypertrophied-

- a. In carious cavity by growth of granulation tissue.
- b. In pulp chamber by absorbing surrounding dentine.

Microscopical appearances-

Granulation tissue. Fibrous tissue.

Squamous epithelium on surface occasionally (formed by kind of skin grafting).

Abundance of vessels.

Scarcity of nerves.

May undergo-

a. Calcification. b. Any degeneration.

FIBROID DEGENERATION OF PULP

Sometimes described as—

- I. Reticular Atrophy. (Wedl.)
- 2. Areolation of Pulp. (Black.)

Microscopical Appearances.

Presents reticular network throughout pulp. Cells, vessels, nerves, all undergoing fibrification.

Odontoblasts shrunken, converted into fibrous cords or disappeared.

Pulp cells disappeared.

Vessels fewer in number, thin walled and larger, and they have constrictions at intervals by contractions of fibrous tissue.

Nuclei of sheaths have gone, and arteries and veins appear alike.

Nerves may be fatty.

Spaces formed by contraction of fibrous tissue. May be slight calcareous formations.

CALCAREOUS FORMATIONS IN THE PULP

- 1. Adventitious or secondary dentine.
- 2. Pulp stones.
- 3. General calcareous degeneration.

1 Adventitious or Secondary Dentine.

Causes :-

Chronic inflammation of pulp from any of its causes.

Microscopical Appearances:-

Generally definite line of demarcation between the normal and adventitious dentine.

Abrupt bend of tubes where the two join.

First part formed is often very irregular, but becomes more regular towards pulp, if extensive.

Varies very much in arrangement of tissues
—tubes, spaces, cells, calcified connective
tissue fibres may all be mixed up in a
hyaline matrix, or one or other of these
elements may predominate. Thus the
following varieties are described:—

- a. Tubular. b. Areolar.
- c. Lacunar. d. Cellular.
 - e. Fibrillar.

2. Pulp Stones.

Causes :-

- a. Chronic inflammation of pulp
- b. Gout.
- c. Old age.

Symptoms:-

Often no pain at all.

If pain, generally neuralgic.

Diagnosis :-

By process of elimination.

Macroscopical Appearances :-

Occur as varying sized calcareous masses in the bodyof the pulp or attached to the wall.

Microscopical Appearances :-

Generally has nucleus, which is of doubtful origin; supposed to be some small irritant spot in the pulp, around which a barrier of calcareous material has been deposited, e.g. a dead cell.

Often laminated concentrically about the nucleus.

May contain tubes radiating out from centre.

May have no apparent structure at all-

3. General Calcareous Degeneration.

Causes :-

Same as for pulp stones.

Macroscopical Appearances :-

Occurs as-

- a. Gritty small masses throughout pulp-
- b. Trabeculæ.
- c. General calcification.

Microscopical Appearances:—
Vary as with pulp stones.

CHAPTER VII

DISEASES OF THE PERIODONTAL MEMBRANE

AGAIN it is as well to point out that there is no hardand-fast line between the different kinds of inflammation described below, as one may at any stage pass into another.

The types of inflammation described below are:

- 1. Acute local periodontitis.
- 2. Chronic local periodontitis.
- General suppurative cervical periodontitis (Pyorrhœa Alveolaris).
- 4. General periodontitis.

ACUTE LOCAL PERIODONTITIS

(i.e. acute inflammation at the root of one tooth).

If running its full course, it will end in acute alveolar abscess.

But it may at any stage cease of its own accord or as the result of treatment, and then resolve altogether or pass into chronic periodontitis.

Causes :-

- 1. Septic matter passing through apical foramen.
- 2. Arsenic passing through apical foramen.
- 3. Extension of inflammation from pulp.
- 4. Blow on tooth (generally accompanied by death of pulp).

51

Symptoms, in order of their appearance (=Clinical History).

- 4 Stages, which rapidly pass from one to the other:—
 - 1. Dilatation Stage.

Tooth uneasy.

Disposition to grind teeth together, giving relief at first stage.

Tooth raised in socket, due to dilatation of vessels.

2. Exudation Stage.

Gum swollen and painful, with red margin.

Tooth more raised in socket.

Great pain on biting, due to exudation of lymph and leucocytes.

Increased flow of saliva.

3. Pus-forming Stage.

Great throbbing and continuous pain.
Increased swelling of gum.

Inability to eat.

Loss of appetite.

Furred tongue.

Pyrexia.

4. Pointing Stage.

Face œdematous, according to situation of bad tooth.

Pain less, as pressure relieved by pus burrowing through bone.

Lymphatic glands of neck swollen.

Salivary glands may be swollen.

Constitutional disturbance.

Abscess points.

Duration.

Average, 7 or 8 days.

Treatment:

Stages 1 and 2.

Clear out and dress pulp cavity.

Counter-irritants to gum. (See Chapter XXIV.)

Local depletion—

Scarification.

Leeches.

Stages 3 and 4.

If saveable.

Clear out and dress pulp cavity.

If not pointing—

Hot fomentations in mouth.

Poultice with hot fig, &c.

Local depletion.

If pointing—

Open abscess.

Rhizodontrophy—as temporary measure, if tooth too tender to manipulate further.

Purgative.

If about to point externally, cause it to open inside mouth and paint collodium outside to prevent breach of skin.

If not saveable.

Extract.

Pathological Course of Acute Alveolar Abscess:

- Passage of micro-organisms through apex of tooth.
- Action of these micro-organisms on the healthy or perhaps previously inflamed periodontal membrane, leading to severe inflammation around apex.

3. Progress of this inflammation to suppuration of membrane, and pus forming, strips the

membrane from the root.

N.B.—No sac formed when process is acute, but membrane involved breaks up into pus or forms granulation tissue.

- 4. Pus continuing to form makes a cavity for itself in the bone, around the apex of the tooth, by the pressure of pus formation and the inflammation absorbing the bone away. Eventually bone is absorbed right through at one spot, and then-
- 5. Abscess points.

Situations in which an Abscess may point:

1. Usually through outer plate. All the

2. Neck of tooth, along course of root. \ teeth.

3. Hard palate. | 6 · 5 · 2 | 2 · 5 · 6

4. Soft palate.

5. Nasal cavity. I I farely

7 . 6 . 5 | 5 . 6 . 7 6. Antrum.

7. Along margin of mandible on face—

8. Fauces

9. Pharynx. 8

10. Angle.

8 . 7 . 6 6 . 7 . 8 II. Neck.

Complications of Alveolar Abscess:

- I. Forced closure of jaw-by exudation of inflammatory products mechanically fixing the jaw.
- 2. Necrosis of jaw-especially alveolar process.
- 3. Cellulitis of neck.
- 4. Inflammation and suppuration of neighbouring lymphatic glands.

- 5. Empyema of antrum.
- 6. Sinuses on face and depressed cicatrix later.
- 7. Trismus.
- 8. Œdema glottidis.
- 9. Septicæmia and pyæmia.
- 10. Post-pharyngeal abscess.
- Pterygoid Plexus.
 Cavernous Sinus.

CHRONIC LOCAL PERIODONTITIS

According to the cause, this condition may be-

- 1. Confined to apex.
- 2. Confined to neck.
- 3. Affecting whole root.

And either 1 or 2 may become 3 if unchecked.

Also according to virulence of irritation it may be-

- 1. Non-suppurative.
- 2. Suppurative.

Causes :-

- 1. Any one of the causes of Acute Local Periodontitis.
- 2. Traumatic effects:
 - a. Filling left too high.
 - b. Crown left too high.
 - c. Too rapid separating.
 - d. Too rapid regulating.
 - e. Impacted third molar.
 - f. Denture.
 - g. Oblique contact.
 - h. Blow.
 - i. Biting hard substance.
 - j. Gold plugging.
 - k. Toothpick.
 - 1. Rubber dam clamp.

- 3. Foreign bodies:
 - a. Food, wedged interstitially.
 - b. Tartar.
 - c. Fish bone, &c.
 - d. Toothbrush bristle.
 - e. Edge of ill-fitting crown.
 - f. Overhanging edge of filling.
 - g. Denture clasp.
 - h. Ligature.
 - i. Elastic bands.
- 4. Perforation of side of root.
- 5. Tooth rising in socket from loss of antagonist.
- 6. Arsenic dressing leaking on to gum.

Symptoms:

With both non-suppurative and suppurative varieties:-

- I. Loosening and lengthening of tooth in socket.
- 2. Tenderness on percussion.
- 3. Dull pain.
- 4. Gum congested.
- 5. Remission and recurrence.

With suppurative variety only:-

- 6. Gum-boil, generally over or near root of affected tooth, i.e. termination of a sinus leading down to the chronic abscess, cavity around the apex. This gum-boil may fill up from time to time and then burst.
- 7. Pus may be continually draining down canal.
- 8. Pus may be oozing up round neck of tooth.
- 9. Deposit of "seruminal calculus" on tooth from pus, if persistent.

Treatment : __

- I. Remove cause.
- 2. If pulp dead, clear out and dress canals.
- 3. Apply counter-irritants to gum (see Chapter XXIV.).

- 4. Remove "seruminal calculus," if confined to neck.
- 5. Antiseptic mouth-wash, if cervical suppuration.
- 6. If gum-boil present, and not cured by dressing canal, syringe forcibly through canal, and out at sinus with carbolic acid (1 in 20).
- If above does not cure, trephine alveolus and cut off apex.

Pathological Results (directly connected with root).

Productive :-

- a. Abscess Sac.
- b. Granuloma.
- c. Dental Cyst.
- d. Exostosis and Inostosis.

Destructive :-

e. Absorption of Root.

a. Abscess Sac.

Origin :-

When suppuration occurs on the inner surface of the Periodontal Membrane, if remaining chronic, it may gradually expand the membrane and absorb the bone, and thereby form a bag or sac, which communicates with the apical foramen and sometimes with the surface, by the pus which it contains burrowing through the bone and gum.

The inner surface of sac is lined with granulation tissue, and as time goes on its wall thickens and becomes more fibrous, and the pus from time to time discharges down the canal or through the sinus.

The sac is generally pyriform, but may be any shape. It often remains attached to the apex on removal of tooth.

b. Granuloma.

Origin :--

This is a mass of granulation tissue formed from the Periodontal Membrane and possibly partly from the bone. It may be formed primarily without any suppuration, but more frequently as the result of long-standing suppuration, where the pus has continually found an exit through a sinus or down the canal.

Sometimes small aggregations of epithelial cells are found in the substance of the granuloma. Occasionally these are seen in large numbers, and they are then called Epithelial Root Tumours.

c. Dental Cyst.

Origin :-

This is formed from one of the aggregations of epithelial cells frequently found in the Periodontal Membrane, especially in the neighbourhood of the apex, and which are the remains of the epithelial sheath of Hertwig. These epithelial cells are stimulated to growth by some chemical or bacterial irritant that passes through the apical foramen of a dead tooth, but which is insufficient to give rise to suppuration. They proliferate and form first an Epithelial Root Tumour, as given above, and then. as the peripheral cells are proliferating, the central ones degenerate and break down into fluid. This process is progressive, and eventually a large cyst is formed, which hollows out the jaw by absorbing the bone from the inside, and

at the same time bone is deposited in rather less quantity on the outside, and so the appearance of expansion of the jaw is obtained.

Symptoms:-

Painless as a rule, or slight neuralgia.

May be numbness (interference with nerve trunk).

May affect either jaw.

Generally connected with permanent teeth.

Very rarely connected with temporary
teeth.

May occur at any age.

Always in connection with septic tooth (which may, however, have been removed previously).

Smooth regular swelling.

Expands outer plate of alveolus most.

Progressive slow growth—anything from 6 months to 2 years to cause any disfigurement.

No sign of inflammation.

Hard to the touch, if bone not absorbed away.

Egg-shell cracking, if bone thin.

Fluctuation if bone absorbed.

Diagnosis :-

See below under "Swellings of Jaws."

Contents :-

Viscid, transparent fluid, containing serumalbumen, serum-globulin and crystals of cholesterin (cholesterin probably formed by degeneration of epithelial cells).

Microscopical Appearances:-

Cyst wall lined with epithelium of variable thickness and character.

Rarely epithelium may be ciliated. Fibrous wall outside this.

Treatment :-

Remove tooth.

Open up cyst, via sulcus, by removal of outer wall.

Evacuate contents.

Thoroughly scrape away lining membrane. Swab over with pure carbolic acid.

Pack tightly to arrest hæmorrhage.

Later pack more lightly with antiseptic dressing to promote growth of granulalion tissue from the bottom, and thus by frequent dressing (daily), cavity will fill up.

d. Exostosis (Productive Periodontitis).

Outgrowth of cementum on root of tooth.

Causes :-

Chronic periodontitis—rather of the nonsuppurative variety; especially that which is kept up by extension of inflammation through the apical foramen from a *chronic* pulpitis.

E.g. Large metal filling near pulp. Erosion.

Attrition, &c.

Site:-

Any tooth, but especially molars.

Symptoms:-

Often none.

Slight pain on percussion.

Dull pain.

Neuralgia.

Diagnosis:-

From Pulp Stone.

Take skiagraph.

Microscopical Appearances:

Lacunæ very variable in size and shape. Canaliculi very variable in length and direction.

"Lines of Salter" very prominent.

Vascular canals occasionally present.

"Sharpey's fibres" sometimes well marked.

Varieties :-

- a. Macroscopical-
 - I. Smooth.
- 2. Irregular.
- 3. General.
- 4. Nodular.
- 5. Bulbous (at apex). 6. "Candle grease."
- 7. Fusing two teeth together (false gemination).
- b. Microscopical (according to predominance of one or other of its elements)-
 - I. Granular.
- 2. Lacunar.
- 3. Laminar.
- 1. Vascular.

5. Fibrillar.

Treatment:-

Remove cause if possible.

Counter-irritants to gum. (See Chap. XXIV.) Removal of tooth.

Inostosis.

See below under "Absorption of Root."

e. Absorption of Root (Rarefying Periodontitis).

Strictly analogous to that of the Temporary Dentition. Large giant cells found fitting into Howship's Lacunæ. May be any extent, even opening pulp chamber. According to the way the character of inflammation changes from time to time, so this process of absorption may alternate with the deposition of cementum. And if the absorption has extended sufficiently to attack the dentine, then cementum

may be deposited later in the cavity thus formed in the dentine. This microscopically gives rise to an appearance named *Inostosis*.

Cementum may actually be deposited in the pulp chamber, if that has been opened up by the process.

GENERAL SUPPURATIVE CERVICAL PERIO-DONTITIS (Pyorrhæa Alveolaris).

Causes :-

Local-

- Primary Gingivitis—principally from Local Septic causes (see Chapter IX).
 —extending to Periodontal Membrane.
- 2. Primary osteitis of alveolar process (Hopewell Smith)—Clinical evidence seems against this.
- 3. Specific Germ (for Bacteriology, see above).

General-

- 4. Gout.
- 5. Other diatheses.
- 6. After fevers.

Predisposing-

- 7. Abnormal articulation.
- 8. Regulating teeth.

Site :-

Any teeth, but especially upper and lower incisors. May extend all round mouth.

May be limited to one aspect of teeth, or extend right round them.

Often seems to spread from one to another.

Symptoms:

Early Stage-

- 1. Interdental papillæ of gum shrink.
- 2. No pus.

Later Stage-

- 1. Pain and soreness of gums.
- 2. Congestion of gums at margins.
- Pus discharging round neck of tooth on digital pressure.
- 4. Separation of gum from tooth, forming pockets
- Recession of gum and absorption of alveolar process.
- 6. Teeth loosening.
- Teeth shifting position—generally protruding (alveolar support being lost, teeth are forced out by bite).
- Ring of dark calculus round neck of tooth, under margin of gum (deposit from pus), extending further up root as disease advances.
- 9. Foul breath.
- 10. Bad taste in morning.

Above symptoms vary in different patients, and also from time to time in the same patient, according to both the virulence of the poison and the susceptibility of the patient at the moment. And often with long-standing cases the severity will vary directly with the general health of the patient. It is most often found in a chronic condition.

Results :-

- I. General malaise.
- 2. Chronic indigestion—septic gastritis.
- 3. Glossitis.
- 4. Tonsilitis.
- 5. Eruptions of skin.
- 6. Chronic rheumatism and osteo-arthritis.
- 7. Affections of eye.

Pathology :-

Absorption of alveolar margin (=rarefying osteitis).

Slight thickening of process just in advance of absorption.

Hypercalcification and translucency of roots of teeth, especially at apices.

Microscopically-

Hypertrophied papillæ in gum.

Dilated capillaries.

Enlarged epithelium.

Round celled infiltration into gum and periodontal membrane.

Treatment:

Preliminary Treatment for all Cases.

- 1. Remove any tooth that is hopelessly loose.
- 2. Scale thoroughly below gum margin.
- 3. Syringe out pockets with strong antiseptic. (See Chapter XXIV.)

Types of Treatment Available.

I. Drugs, directly applied to pockets—

Tannic acid, if very sore.

Copper sulphate (powdered), if very chronic.

Protargal, 8 per cent.

Trichloracetic acid—apply gently with wood point.

Tincture of Iodine.

Use either about twice a week.

Any of above tend to give temporary relief rather than permanent cure.

2. Electrolysis with tincture of iodine. Tedious, but beneficial.

- Vaccine treatment to improve the opsonic index (Goadby). In association with local treatment, cures may be effected in some cases.
- Extraction of every affected tooth—especially when general symptoms are established.

Adjuncts to any of the above Treatments.

- Saturated solution of tannic acid in spirit, rubbed on the gums with the finger night and morning, is a useful adjunct.
- 2. Good antiseptic and astringent mouth-wash. (See Chapter XXIV.)
- Wire or splint teeth together, to give periodontal membrane a rest. Temporary benefit if teeth loose.
- 4. Aim at obliteration of pockets by
 - a. Cutting off tags of gum.
 - b. Excising or vertically incising gum margin.
 - c. Burring alveolar process.

GENERAL PERIODONTITIS

Causes :-

1. Various diatheses—

Gout.

Rheumatism.

Syphilis.

Diabetes.

Malaria.

Scurvy.

2. After—

Exanthematous fevers. Influenza.

- 3. Prolonged administration of mercury.
- 4. Fumes of phosphorus.
- 5. Extension from Pyorrhœa Alveolaris.
- 6. Injury—

Severe blow on jaw.

Partial dislocation of series of teeth.

DISEASES OF PERIODONTAL MEMBRANE 65

Symptoms :-

Same as for Chronic Local Periodontitis, except that it occurs on several teeth at once.

Treatment : -

- I. Remove and treat any cause.
- 2. Counter-irritants to gum. (See Chap. XXIV.)
- 3. Strong antiseptic and astringent mouth-wash.
- 4. Purges.
- 5. Tonics.

CHAPTER VIII

INJURIES TO THE TEETH

- I. Concussion.
- 2. Dislocation.
- 3. Fracture.
- 4. Dilaceration.
- 5. Erosion.
- 6. Attrition.
- 7. Abrasion.

I. CONCUSSION

Causes :-

Direct blow.

Knocking teeth together.

Biting hard substance.

Results :-

Periodontitis—

a. Immediate.

b. Later, following death of pulp.

Pulpitis—

From the jarring.

Death of pulp by rupture of apical vessels. (This leads to discoloration of tooth.)

Treatment: -

Counter-irritants to gum.

Remove pulp, if pulpitis persistent or pulp dead.

2. DISLOCATION

Degrees :-

- a. Partial.
- b. Complete.

a. Partial.

Treatment :-

Replace tooth in correct position. Fix with ligature or splint. Counter-irritants to gum.

b. Complete.

Treatment :-

If during extraction—replace at once.

Otherwise remove pulp, fill canal, sterilise in mild antiseptic, and replace tooth and fix with splint.

Counter-irritants to gum.

3. FRACTURE

Causes :-

Blow.

Extraction.

Mastication.

Degrees :-

- a. Not involving pulp.
- b. Involving pulp.

a. Not Involving Pulp.

Symptoms:-

Exposed dentine sensitive.

Hyperæmia of pulp.

Periodontitis later.

Treatment:-

Alcohol or zinc chloride to dentine.

Strike up cap and cement on as temporary measure if sensitiveness persistent, as important to save pulp if root not complete.

Counter-irritants to gum.

If pulp suppurates, remove pulp if root complete; otherwise remove tooth.

b. Involving Pulp.

Treatment:-

If age such that root not complete, remove tooth and let space close up, bearing in mind sex of patient and any evidence of irregularity.

If root complete, remove pulp, and prepare for crowning, if direction of fracture will allow of this. Otherwise extract tooth.

Fracture of tooth high up has been known to unite by growth of secondary dentine and cementum.

4. DILACERATION

A distorted condition of tooth from injury to the same during development.

Characteristics :-

 Common in front of mouth, being more liable to injury.

2. Crown or root or both affected, according to

time and nature of injury.

 Any degree of distortion may be found, from a mere bend to a much distorted tooth. There is often a bulging at the bend.

4. Sometimes marked mobility.

5. Generally erupt abnormally, and may not erupt at all.

Causes:

- 1. Direct blow.
- 2. Injury during extraction of the temporary tooth.
- 3. Injury from abscess on the temporary tooth.

5. EROSION

Characteristics :-

- I. Saucer-shaped cavities, deepest near gum.
- 2. Generally labial aspect near gum.
- 3. Cutting edges sometimes involved.
- 4. Surface hard and polished.
- 5. One or more teeth affected.

Causes :-

- I. Entirely abrasion from toothbrush and powders. Strongly supported by Miller's experiments, and no authentic case yet seen in patient not using a toothbrush, or in a situation not easily reached by a brush. Polished cavities in such a situation are always due to abrasion either from Dentures, persistent use of floss silk, or wooden points.
- Acid secretion from buccal mucous glands, viz. acid sodium phosphate. Evidence inconclusive.
- Potassium and ammonium sulphocyanide in saliva. Evidence inconclusive.
- 4. Associated with gout.

Microscopical Appearances :-

Abrupt termination of dentinal tubes, which are otherwise normal.

Zone of translucency under cavity.

No pigmentation.

Secondary dentine in pulp.

Treatment:

Fill, if convenient, with gold or porcelain.

Softer toothbrush and light powder.

Cauterisation of buccal glands has been suggested, but does not seem justifiable.

If sensitive—silver nitrate applied twice a week.

Milk of magnesia as mouth-wash night and
morning.

Following applied daily:—
Powdered gum mastic 3i.
Zinc chloride gr. v.

Chloroform 3ss.

6. ATTRITION

Wearing down of teeth by mastication.

Characteristics:

Few or all teeth affected.

Any degree up to wearing down level with gum.

Causes :-

Edge-to-edge bite or other irregularity.

Harsh character of food.

Want of density of tooth substance.

Habit of grinding teeth together.

Gouty people are frequent subjects.

Pathology: --

Secondary dentine in pulp cavity.

Pulpitis sometimes.

Exposure of pulp rarely.

7. ABRASION

Wearing of teeth by friction with foreign body.

Causes :--

Denture.

Pipe, especially clay pipe.

Toothbrush (see *Erosion*).

Occupations, where things continually held between the teeth.

CHAPTER IX

DISEASES OF GUMS AND MUCOUS MEMBRANE OF MOUTH

- A. Gingivitis.
- B. Stomatitis.
- C. Tumours.
- D. Hypertrophy of Gums.
- E. Scurvy of Gums.
- F. Addison's Disease.
- G. Purpura.
- H. Pemphigus.
- I. Lead Poisoning.

A. GINGIVITIS (Inflammation of Gums)

Causes :-

I. Septic-

Want of cleansing.

Tartar.

Roots of teeth.

Dentures.

Mouth-breathing—natural cleansing by saliva and lips lessened.

Inefficient mastication.

2. Traumatic-

Toothbrush.

Too rapid regulating.

3. Medicinal-

Mercury.

Potassium iodide.

4. Constitutional—

Gout.

Rheumatism.

Syphilis.

Diabetes.

Malaria.

Indigestion.

Alcoholism.

Influenza.

Varieties :-

- I. Confined to gum margins.
- 2. General.

Symptoms:

Gums red, swollen, and soft, especially margins.

May be hypertrophied between teeth.

Often dark-coloured calculus on teeth under free margin of gum, deposited from pus, which oozes up.

Bleed easily.

Treatment:

Treat cause.

Pack powdered copper sulphate under gum twice a week.

Rub on tannic acid in spirit twice a day. Antiseptic and astringent mouth-wash.

Results :-

Cervical periodontitis, if untreated. Recession of gum, if recurring.

B. STOMATITIS

(Inflammation of Mucous Membrane of Mouth)

Varieties :-

- I. Simple.
- 2. Mercurial.
- 3. Follicular.
- 4. Ulcerative.
- 5. Gangrenous.
- 6. Parasitic.
- 7. Syphilitic.

1. SIMPLE STOMATITIS

Causes :-

Uncleanliness.

Tobacco smoke.

Gastric and intestinal troubles.

Fevers.

Associated with nasal catarrh.

Symptoms :-

Mucous membrane tender and bleeds easily.

At first dry, and then wet with sticky secretion which later becomes purulent.

Congested margin to gums.

Gums and sulcus pale and mottled—the mottling being due to sodden epithelium being rubbed off the tips of enlarged papillæ, leaving red patches.

Fœtid breath.

General malaise.

Treatment : -

Antiseptic and astringent mouth-wash.

Purge.

Tonic—Strychnine and Iron,

2. MERCURIAL STOMATITIS

Cause :-

Mercury given medicinally to excess.

Symptoms:

All symptoms as for Simple Stomatitis.

Also-

Teeth and gums tender.

Pain on eating.

Metallic taste.

Salivation.

Periodontitis.

Ulceration of margin of gums.

Sloughing and necrosis.

Treatment:

Stop course of mercury.

Antiseptic and astringent mouth-wash.

Application of tincture of iodine.

Potassium chlorate and potassium iodide internally.

Purge.

3. FOLLICULAR STOMATITIS

Symptoms:

Inflammation of mucous follicle, turning to small ulcer with sharply cut edges and zone of redness.

Very painful.

May be single or multiple.

Most often occur in sulcus.

Treatment:

Touch with copper sulphate or silver nitrate.

4. ULCERATIVE STOMATITIS

Causes :-

Young children out of health. Uncleanliness.

Symptoms:

Starts in front of mouth and extends as line of ulceration along margin of gum, and later on along cheek from contact, leaving ulcer with sharp, irregular edge and bluish margin.

Unilateral as a rule.

Little pain at first.

Salivation.

Fœtid breath.

Swelling of cervical lymphatic glands.

Loosening of teeth.

Sloughing and necrosis.

Treatment :-

Antiseptic and astringent mouth-wash.

Swab out with-

Pot. Chlor. 3i.

Ac. Hydrochlor. dil. 3ss.

Aq. ad. 3xx.

Apply-

Tannic acid.
Dried alum.

Touch with silver nitrate twice a week.

Potassium chlorate internally in syrup.

(gr. i. for every year of child's age up to gr. x.)

Purge.

Tonic.

Results :-

Loss of teeth.

Necrosis.

Lymphatic abscess.

Septicæmia.

Pyæmia.

Septic gastritis.

Septic pneumonia.

Inanition.

Fibrous bands in cheek, on healing.

5. GANGRENOUS STOMATITIS (Cancrum Oris) (Noma)

Causation :-

Generally in children (2–6 years). After fevers. Idiopathic.

Pathology: -

Due to a diffuse capillary thrombosis, induced by germs.

Symptoms:

Great swelling of face, but very little œdema. Ulcer on mucous membrane inside cheek.

Dark spot outside at same place.

Rapid destruction of cheek, making opening, which may be large, and even whole cheek may be destroyed.

Drowsiness and exhaustion, and possibly death. May undergo temporary arrest.

Treatment:

After drying, apply one of following—

- a. Carbolic acid.
- b. Nitric acid.
- c. Actual cautery.

Excise ulcer and perform plastic operation.

Inject pure carbolic acid around ulcer.

Prevent septic matter from entering air passages, &c., by giving good mouth-washes, and applying antiseptic dressing.

Support general system—

Nourishing food.

Stimulants.

Tonics.

Results:

Same as for Ulcerative Stomatitis.

6. Parasitic Stomatitis—Thrush

Causation :-

Occurs in babies mostly.

Occurs in old people, usually just before death, in wasting diseases.

Due to Oidium albicans = mycelium of fungus.

Symptoms :-

Resembles Simple Stomatitis when it begins, but patches rapidly cover with exudation, forming small white patches.

These may peel after a few days, but cover over again.

Treatment :-

Wipe mouth round after every meal.
Borax gently rubbed on.
Alum gently rubbed on.
Sulphurous acid applied.

7. SYPHILITIC STOMATITIS

a. Primary Ulcer-

May occur anywhere in mouth.

b. Secondary Lesions-

Mucous tubercles, especially on edges of tongue, and angles of mouth, and inside cheeks.

May be shallow ulcers.

c. Tertiary Lesions-

Swelling first (gumma), which breaks down, leaving large, deep ulcer.

C. TUMOURS OF GUMS AND MUCOUS MEMBRANE OF MOUTH

Forms:

- 1. Polypus.
- 2. Epulis.
- 3. Papilloma.
- 4. Nævus.
- 5. Epithelioma.
- 6. Sarcoma.
- 7. Gumma.

I. POLYPUS OF GUM (Local Hypertrophy)

Causes :-

Tartar.

Ragged edge of tooth.

Clasp of denture, &c.

Symptoms:

Generally between two teeth.

May be pedunculated or not.

Very tender.

Bleeds easily.

Diagnosis:

From polypus of pulp.

From epulis.

Treatment:

Cut off.

Burn out with red-hot burnisher or cautery. Cauterise away with ethylate of sodium, &c. Push out with G.P. if small.

2. EPULIS

Symptoms:

Generally between two teeth. Predunculated and lobulated.

Springs from junction of periodontal membrane with periosteum.

Usually slow growing, but may be rapid.

May attain large size.

May ulcerate.

More common in women.

Pathology: -

3 varieties-

- a. Fibrous tissue with small round cells.
- b. Fibrous and elastic tissue.

c. Fibro-sarcoma with myeloid cells.

Any of above may contain besides small pieces of bone or root, which have become detached by absorption.

Treatment :-

Cut off and scrape base freely and cauterise with pure carbolic acid.

Remove tooth from which it springs, if growth large.

3. PAPILLOMA

Pathology :-

Hypertrophy of papillæ of mucous membrane. Analogous to a Wart.

Treatment:

Remove and cauterise.

4. NÆVUS

Symptoms :-

Flat, red, and only slightly raised.

Begins as red pimple.

Seldom larger than half an inch across, but may be large.

Bleeds easily.

Blood can be pressed out, but returns when pressure relaxed.

Pathology:

Hypertrophy and multiplication of capillaries.

Treatment:

Remove and cauterise.

5. EPITHELIOMA 6. SARCOMA 7. GUMMA See General Surgery.

D. HYPERTROPHY OF GUMS

Cause :-

Congenital, starting in quite early life.

Symptoms:

Masses of gum tissue, growing up round teeth, and even covering them, and sometimes protruding from mouth.

Masses may be lobulated or have smooth appearance.

Darker than normal gum.

Very vascular and bleed easily.

Often discharge.

Pus oozing up round teeth.

Fœtid breath.

Treatment:

Cut off, and remove surface of alveolar process as well.

After removal, cauterise with actual cautery, because of extensive bleeding.

Pathology:—

Firm fibrous stroma, springing from alveolar process.

Much glandular tissue.

Large vascular papillæ.

Epithelium often very thick.

Alveolar process generally enlarged too.

TRANSPARENT HYPERTROPHY

Symptoms :-

Pale blue cord, running along gum margin. Teeth loosen and fall out.
Cuts like cartilage, and does not bleed.
Very rare condition.

E. SCURVY OF GUMS

Symptoms: -

Gums over process—red and swollen.
Rest of mucous membrane and lips—white.
Bleeds easily.
Teeth loosen.
Ulceration of gums.
Sloughing and necrosis.

F. ADDISON'S DISEASE

Pigmented patches found in gums.
Associated with pigmentation elsewhere.

G. PURPURA

Hæmorrhages under mucous membrane of mouth. Associated with Purpura elsewhere.

H. PEMPHIGUS

Blebs on mucous membrane of mouth, especially palate and cheeks.

Associated with Pemphigus elsewhere.

I. LEAD POISONING

Symptoms: -

Blue line upon gums, generally below the lower incisors.

Absent in edentulous patient.

Discoloured patches on mucous membrane of cheeks.

Colic.

Constipation.

Metallic taste.

Fœtid breath.

General debility and anæmia.

Wrist drop.

Diagnosis: -

Blue line has to be diagnosed from-

- a. Deposit of white lead, not in but on gums.
- b. Charcoal tooth-powder.
- c. Carbon dust in miners.
- d. Copper and bismuth poisoning (rare).
- e. Black tartar under margin of gums.

Sources of Lead Poisoning: -

- I. Lead and its salts in factories.
- 2. Occupations, &c.
 - a. Painters.
 - b. Potmen.
 - c. Compositors.
 - d. Brussels lace makers.
 - e. Glazed card makers.
- 3. Applications
 - a. Ointments.
 - b. Hair dyes.
- 4. Water—

Leaden cisterns and pipes.

5. Food.

CHAPTER X

SALIVARY CALCULUS

Salivary Calculus is a deposit from the saliva of salts, held in solution by excess of carbon dioxide gas, which is liberated on exposure to the air, and thus the salts are deposited.

Sites :-

- I. In ducts and glands.
- 2. On teeth.

Tartai,

3. On dentures.

I. IN DUCTS AND GLANDS

Results :-

- a. Retention cyst.
- b. Suppuration of cyst.
- c. Salivary fistula.
- d. Ulceration of duct.

2. ON TEETH

Sites :-

Especially—

- a. Lingual aspect of lower incisors.
- b. Buccal aspect of upper molars.

Being opposite salivary ducts.

But may occur anywhere in mouth, and is occasionally sufficient to completely hide the teeth of one side.

Conditions favouring Accumulation of Tartar: -

a. Lack of Mastication.

Painful tooth-

- (i) Exposed pulp.
- (ii) Periodontitis.
- (iii) Polypus of pulp or gum.

Sore gums.

Loss of antagonising teeth.

Habit of using one side only.

With any of above, tartar may collect over all the teeth on one side.

- b. Lack of brushing.
- c. Rough surfaces.
 - (i) Cavities.
 - (ii) Pits.
 - (iii) Bad fillings.
 - (iv) Dentures.
- d. Irregularities in position.
- e. Condition of saliva (varies in individuals).
- f. Condition of health. (?)
- g. Gout. (?)

Effects:

Gingivitis.

Recession of gum.

Periodontitis (Pyorrhœa).

Absorption of alveolus.

Looseness of teeth.

Sore tongue.

Lodgment of food.

Foul breath.

Gastric catarrh.

Composition :-

a. Salts.

Carbonates of Calcium and Magnesium— Chief salts from parotid, e.g. on molars.

Phosphates of Calcium and Magnesium— Chief salts from sublingual, e.g. on lower incisors.

Chlorides of Sodium and Potassium— Small quantities from both glands.

Sulphocyanide of Potassium— Traces only.

b. Animal Matter.

Epithelium.

Food.

Pus cells.

c. Other Matter.

Water.

Mucous.

Ptyalin.

Pigment (smoke, &c.).

Quantitative Analysis shows great variation.

Roughly-

Salts . . 80 per cent.

Water and Mucous 12 ,,

Animal Matter . 7 ,,

Ptyalin . . I ,,

Hard tartar will contain rather more salts. Soft tartar will contain rather less salts.

Treatment:

Removal by "Scaling."

Principles of Scaling: -

- a. Remove every particle and polish.
- b. Chip off in as large pieces as possible leaving less to scrape.
- c. Push towards gum when large amount.
- d. Pull away from gum when small amount.

Precautions in Scaling:-

- a. Instruments to be sterile and sharp.
- b. Avoid injuring gum.
- c. Avoid injuring enamel.
- d. Support loose teeth.
- e. Prevent patient swallowing particles.
- f. Give mouth-wash afterwards.

CHAPTER XI

SALIVARY FISTULA

Varieties :-

- 1. Connecting mouth with exterior.
- 2. Connecting salivary duct with exterior.

1. Connecting Mouth with Exterior.

Causes :-

- a. Alveolar abscess bursting externally and internally.
- b. Necrosis.
- c. Bad fracture.
- d. Slipping of elevator.
- e. Cancrum oris.
- f. Syphilitic ulceration.
 Any of which may be means of saliva finding its way through.

Treatment:-

- a. Treat cause.
- b. Plastic operation.
- c. Pass seton through (old treatment).

2. Connecting Salivary Duct with Exterior.

Causes:-

Blocking of duct by-

- a. Calculus.
- b. Cicatrix.
- c. Growth.

Opening of duct by-

- a. Knite.
- b. Suppuration.
- c. Ulceration of malignant growth.

Treatment :-

Remove calculus or growth. Cause duct to open inside. Plastic operation outside.

CHAPTER XII

ODONTOMES

An odontome is a neoplasm composed of dental tissues in varying proportions and different degrees of development, arising from a tooth germ or a tooth still in its process of growth (Bland Sutton).

Classification (Bland Sutton):-

- I. Aberrations of enamel organ— Epithelial odontome.
- 2. Aberrations of follicle
 - a. Dentigerous or follicular odontome.
 - b. Fibrous odontome.
 - c. Cementoma.
 - d. Compound follicular odontome.
- 3. Aberrations of papilla— Radicular odontomes.
- 4. Aberrations of whole tooth germ—
 Composite odontomes.
- N.B.—The above is the accepted classification, but it will be seen that the different forms of odontomes merge so much into one another that it makes a comprehensive classification a very difficult matter.

89

1. ABERRATIONS OF ENAMEL ORGAN

EPITHELIAL ODONTOME

Site:

Generally lower molar region.

Pathology:

On section—numerous cysts of various sizes, separated by fibrous or bony septa.

Contain brown mucoid fluid.

Walls and septa, on microscopical examination, show branching columns of epithelium with fibrous tissue stroma.

2. ABERRATIONS OF DENTAL FOLLICLE

a. Dentigerous Cyst or Follicular ODONTOME

Site:

Lower molar region. Antral region.

Pathology:

Excessive secretion of fluid between the developing tooth and the follicle wall, gradually enlarging its crypt and body of jaw.

Fluid is vellowish in colour, and may contain cholesterin.

Walls are fibrous, and usually have epithelial lining.

Generally contains tooth, either loose or partially fixed in floor, or may be inverted.

May suppurate.

N.B.—The epithelial lining may possibly indicate that the enamel organ may have aberrated after forming the enamel, and therefore that this form is not really an aberration of the follicle.

b. FIBROUS ODONTOME

Pathology: -

Hypertrophy of tooth sac. May be laminated on section. May have tooth in centre.

c. CEMENTOMA

Pathology: -

Calcification en masse of a fibrous odontome.

Formed of cementum, since follicle forms that tissue.

d. COMPOUND FOLLICULAR ODONTOME

Pathology :-

Calcification of a fibrous odontome in a sporadic manner, with sometimes cystic formation.

Made up of masses of cementum and bone, embedded in loose fibrous tissue, or lying loose in fluid, when the follicle has also become cystic.

N.B.—Certain cysticodontomes, containing many denticles of enamel, dentine and cementum, although sometimes classified under this heading, because they are cystic, would perhaps be better classified as "Composite Odontomes."

3. ABERRATIONS OF DENTINE PAPILLA

Site:

Lower jaw.

Pathology: -

Due to an outgrowth of the dentine papilla, after the completion of the crown and during or before the formation of the root, and then partial or complete calcification. Thus, according to the predominance of the tissue formed, there may be—

- a. Radicular dentoma.
- b. Radicular osteo-den!oma.
- c. Radicular cementoma.

The cementum in the last may be formed by the diseased dentine papilla, or by the surrounding follicle, enlarged by the dilatation of the papilla.

N.B.—Many irregularly-shaped teeth with malformed crowns may be classified as this form of odontome, having possibly arisen by an outgrowth of the dentine papilla before the commencement of calcification of the crown.

4. ABERRATIONS OF WHOLE TOOTH GERM

COMPOSITE ODONTOME

Pathology: -

Due to irregular growths of all parts of the tooth germ, forming irregular masses of calcified tissue.

On section they contain enamel, dentine, and cementum, all arranged quite indefinitely, either in one solid mass, or a large number of smaller masses or denticles.

Symptoms of all Odontomes:—

1. Often no pain.

 May be merely sense of uneasiness or neuralgic pain.

 Tumour, generally encroaching upon alveolus, increasing very slowly, and varying in physical characters, according to variety of odontome.

- 4. May become inflamed, and finally suppurate and point, leaving sinuses.
- 5. Generally tooth missing, but not always.
- 6. Although starting to grow at usual time of development of tooth, not as a rule noticed till about 15 to 20 years of age.

Treatment of all Odontomes:-

Removal.

If cystic, lining membrane must be scraped away.

CHAPTER XIII

ODONTALGIA AND NEURALGIA

ODONTALGIA

Odontalgia is pain in or around a tooth.

Forms:

- I. Local.
- 2. Referred.

I. LOCAL

Causes :-

Morbid condition of that particular tooth. (Fully considered in foregoing pages.)

2. Referred

Causes :-

Exactly the same as for neuralgia. (See below.)

N.B.—If the cause is *peripheral*, it is always on the same side as the pain, which is never referred across the middle line.

NEURALGIA OF THE 5th NERVE

Neuralgia is pain in the course of, or in the area of distribution of, one or more of the branches of the nerve.

Causes :-

Predisposing: -

- a. Overwork.
- b. Neurotic temperament.
- c. Reduced vitality.

Exciting:-

- a. Peripheral—
 - (i) Affections of pulp.
 - (ii) ,, periodontal membrane.
 - (iii) ,, periosteum.
 - (iv) ,, gum.
 - (v) ,, mucous membrane of mouth, nose, or antrum,
 - (vi) ,, skin.
 - (vii) ,, eye.
- (viii) ,, viscera, uterus, &c.

b. Course of Nerve-

- (i) Injury.
- (ii) Pressure.

Wisdom tooth. Tumours.

c. Cerebral—

- (i) Degenerations.
- (ii) Tumours.
- (iii) Hysteria.

d. Systemic-

- (i) Gout.
- (ii) Rheumatism.
- (iii) Syphilis.
- (iv) Malaria.
- e. Idiopathic.

Dental Causes :-

- 1. Caries.
- 2. Metal fillings too close to pulp.
- 3. Oxychloride fillings.
- 4. Erosion.
- 5. Attrition.
- 6. Abrasion.
- 7. Fracture of tooth.
- 8. Exposure of cementum.
- 9. Calcification of pulp.
- 10. Degeneration of pulp.
- Periodontitis from any cause, especially the chronic form.
- 12. Exostosis.
- 13. Buried root.
- 14. After extraction—
 Tearing of nerve.
 Healing of tissues.
- 15. Erupting teeth.
- 16. Impacted wisdom teeth.
- 17. Absorption of 2nd molar by 3rd molar.
- 18. Inflammation of gums.
- 19. Growths, odontomes, and cysts.

Diagnosis re Teeth:

Probe all round and under gums.

Suspect any filling.

Test each tooth with heat and cold.

Tap all teeth.

Press teeth in all directions.

Suspect pulp of lately dying.

Note wisdom teeth—whether erupted.

Note position of pain-

Upper teeth—infra- and supra-orbital. Lower teeth—ear, neck, and shoulder.

Upper and lower teeth—parietal region.

Failing the teeth, look for other cause.

CHAPTER XIV

SECONDARY AFFECTIONS DUE TO DISEASED TEETH

1. Direct Septic Infection—i.e. extension of inflammation.

Empyema of antrum.

Necrosis.

Post-pharyngeal abscess.

Œdema glottidis.

Cellulitis of neck.

Cellulitis of orbit and affections of eye.

2. Remote Septic Infection.

Lymphadenitis.

Lymphatic abscess.

Meningitis.

Thrombosis of sinuses.

Via lymphatics.

Via facial vein.

Septic gastritis. By swallowing septic matter. Appendicitis.

Septic pneumonia. By inhaling septic matter.

3. Septic Absorption—by toxins or micro-organisms getting into blood stream.

Sapræmia.

Septicæmia.

Pvæmia.

Septic Anæmia.

Septic Neuritis.

Puerperal fever.

Eruptions of skin.

Chronic Rheumatism.

Osteo-arthritis.

4. Direct Absorption—via exposed pulp, periodontal membrane or mucous membrane.

Phosphorous Necrosis.

 Nervous Irritation—by reflex irritation of sensory, motor, trophic or vaso-motor nerves.

Neuralgia.

Epilepsy.

Hysteria.

Trismus.

Spasm of facial muscles.

Wry neck.

Paralysis.

Herpes.

Various diseases of eye.

6. Tissue Irritation—by continuous irritation from rough surface, combined with sepsis.

Ulcer of tongue.

Epithelioma.

7. Malnutrition—from want of mastication.

Indigestion. Wasting.

CHAPTER XV

EMPYEMA OF ANTRUM

Causes :-

- Extension of inflammation from roots of teeth—any upper tooth, but especially 1st molar.
- Extension of inflammation from nasal cavity, —catarrh, &c.
- 3. Extension of inflammation from—
 Frontal sinus. | Via nose.
- 4. Alveolar abscess bursting into antrum, instead of in sulcus.
- 5. Passage of foreign body
 - a. Root.
 - b. Food.
 - c. Toothpick.
- 6. Necrosed bone; syphilitic, &c.
- 7. Blow on face.
- 8. Operations on face.
- 9. Fracture of maxilla.
- 10. After Influenza.

Symptoms:

Acute condition—when ostium maxillaire will probably be closed by swelling of mucous membrane— Pain over forehead and face.

Infra-orbital neuralgia.

Throbbing—worse when recumbent.

Tender to pressure.

Bulging of antrum, especially on face.

Parchment crackling if bone thinned.

Fluctuation if bone absorbed away.

Pyrexia.

Rigor.

Malaise.

Chronic condition—(more common)—when ostium maxillaire will probably be open—

Pain as above, though less.

Discharge from one nostril if ostium not too high up.

Surging of fluid, heard by patient only.

Smell, noticed by patient only.

Indigestion from swallowing pus.

Pus coughed up in the morning from pharynx.

Diagnosis :-

Means :-

- I. Look for cause.
- Position of patient—head forward and affected side uppermost, pus may trickle from nostril.
- Speculum in nose—pus may be seen in middle meatus.
- 4. Catheterisation of ostium maxillaire by nose.
- Transillumination by electric light in mouth or pharynx. Infra-orbital zone of light if antrum normal.
- Aspiration with needle and syringe—through inferior meatus.

Treatment:

- 1. Open antrum and drain
 - a. Socket of extracted tooth. Buccal root of 1st molar fo: preference.
 - b. Canine fossa.
 - c. Under malar bone.
 - d. Inferior meatus.

Open by means of— Ordinary gimlet.

Antral gimlet with guard.

Burr or drill on engine.

Straight elevator.

Opening to be made *large* (as little finger), if special plug or tube is to be made.

Opening to be made same size as tube, if an already made tube is to be used.

2. Syringe forcibly at once with any mild antiseptic solution by means of antral syringe.

Get the solution and pus to pass out through nose if possible.

3. Drain.

a. By putting in at once a temporary drain,
 e.g. piece of catheter,

piece of rubber tubing,

piece of gauze,

and proceed to make tube or plug to pass through hole into antrum, and retain by a flange if in sulcus, or a denture if in alveolus.

b. By putting in at once a-

Symond's tube, Ackland's tube, McGavin's tube,

and leave in, syringing by means of the

respective special syringes.

Whichever means of drainage is used the antrum must be syringed once or twice a day with any mild antiseptic forcibly, so as to penetrate into all the recesses. Change of antiseptic is often beneficial.

In favourable cases the above should effect a cure in 3 or 4 weeks. But the more chronic cases may go on discharging for months, years, or even be absolutely incurable by this means. In any case the hole should be kept open as long as there is any discharge.

4. Further treatment for chronic cases.

Open more widely and pack with lint dipped in carbolic acid (1 in 20) to destroy diseased mucous membrane and promote growth of fresh one, or swab over with pure carbolic acid.

Pack daily.

May be opened more widely still, and scraped and tightly packed. In this case antrum fills up with granulation tissue and probably becomes obliterated.

Or it may be left open after scraping, with the object of allowing a new mucous membrane to grow over the surface of the antrum.

This latter method of treatment is the one that is gaining more favour every day with modern surgeons as the *primary* measure in treating antral disease.

CHAPTER XVI

NECROSIS OF THE JAWS

Sites :-

Either jaw, but more frequently mandible, because—

- a. More exposed to injury.
- b. More compact bone.
- c. Less blood supply.

Causes :-

Extension of inflammation from alveolar abscess—

By vessels of bone becoming inflamed and thrombosed.

2. Escape of arsenic dressing through apex of tooth or on to gum—

By corrosive action of drug.

3. Blow on jaw-

By periosteal abscess stripping off periosteum.

4. Fracture-

By-

- a. Directly cutting off blood supply.
- b. Becoming septic from mouth.
- 5. Inhalation of fumes of yellow phosphorus— By direct corrosive action.
- 6. Administration of mercur —

By acting through the system.

7. Exanthematous fevers—

By periostitis probably during acute stage

8. Ulcerative stomatitis and cancrum oris— By suppuration of periosteum.

9. Syphilis-

By breaking down gummatous infiltration.

10. Tubercle-

By carious disintegration,

11. Malignant disease—

By extension into bone.

12. Actinomycosis—

By extension into bone.

13. Idiopathic

Symptoms :-

Gums swollen and tender.

Abscess pointing.

Sinuses in mouth and on face.

N.B.—Multiple sinuses on face almost invariably indicate necrosis.

Teeth loose.

Breath fœtid.

Skin red and œdematous.

May end fatally.

Treatment:

Remove cause.

Hot fomentations in mouth.

Antiseptic mouth-wash.

If abscess shows signs of pointing on face, apply hot fomentations, or open externally.

If possible, get free exit for pus in mouth.

Dress external sinuses.

Remove sequestrum when loose and covered only by soft tissues (ascertained by probing sinuses, two at once if possible, with stiff probes, to feel if bone rocks).

Give periodical purges.

PHOSPHOROUS NECROSIS

Cause : -

Fumes of yellow phosphorus enter deep parts by—

a. Breach of periosteum.

b. Socket of recently extracted tooth.

c. Exposed pulp.

N.B.—It is thought by some to be due, not to phosphorus at all, but to the tubercle bacillus, to which those working in phosphorous factories may be specially liable.

Site: -

Body of jaws— Upper and lower.

Symptoms: -

Great pain, beginning often with toothache.

Great swelling.

Great suppuration and several sinuses.

Great constitutional disturbance.

Pathology: -

Sequestrum has upon it a "pumice stone deposit," formed by periosteum.

Sequestrum also generally presents large interlacing Haversian canals running at right angles to direction of bone.

Extensive production of new bone.

EXANTHEMATOUS NECROSIS

Causes :-

Any of the exanthematous fevers, but especially Scarlet fever.

Measles.

Generally occurs about 4th year and never after 10th year.

Site: -

Alveolar portion of jaw.
Usually symmetrical if near front of mouth.

Symptoms:

Breath very fœtid—often the only symptom, till gums observed.

Insidious onset.

No pain as a rule.

No swelling.

Pus oozing up round teeth, which loosen.

Gums strip off alveolar process in a vertical direction.

Pathology:

Sequestrum is a dirty yellowish colour, and may be dark brown if exposed to air.

Margins ragged.

Worm-eaten appearance.

CHAPTER XVII

FRACTURES OF THE JAWS

Causes:

Direct Violence :-

Blows.

Kicks.

Falls.

Gunshots.

Violent extraction.

Indirect Violence :-

Coughing, &c., when jaw weakened by disease.

Sites :-

Mandible—

Anywhere, but especially—

Canine region.

Angle.

Molar region.

Symphysis.

Neck.

Maxilla-

Generally alveolar portion, and especially in front.

Symptoms:

Pain in region of fracture and worse on movement.

Loss of power.

Deformity—line of teeth altered, due to displaced fragment.

Swelling.

Crepitus.

Mobility of fragments.

Causes of Displacement:-

Voluntary muscular action—
Eating, talking, &c.
Involuntary muscular action—tonic spasm.
Swelling.
Obliquity of line of fracture.
Soft tissue between fragments.
Tooth between fragments.
Piece of bone between fragments.
Callus between fragments.

Treatment :-

a. MANDIBLE.

Bandages.

4-tailed.

Hamilton.
Either alone or in conjunction with chin boot of some sort.

To be used—

When fracture single, simple, and no displacement, and only very slightly movable.

2. Hammond's wire splint or modification, surrounding all teeth and wired to teeth with binding wire.

To be used—

When fracture single and anywhere in front of or in premolar region. When little displacement and easily rectified. When most of teeth are standing and firm.

- 3. Tomes' Splint.---Metal cap covering all the teeth, and pressed home with cement or G.P.
- 4. Hayward's Splint.—Vulcanite cap over all the teeth with wire wings prejectingoutofmouth along cheek. Pressed home with G.P., and bandaged under chin and over wire wings.
- 5. Gunning's Splint or modification.—C a p s of vulcanite or metal covering both upper and lower jaws, fixed together at sides. Bandage also applied.
- 6. Angle's studded bands and wires.—Opposing teeth in upper and lower on each side are banded and wired together over the studs on the bands.

To be used—

When conditions are similar to those for a "Hammond," except that it can be used when fewer teeth are standing, and if they happen to be loose.

Also in children, in preference to a "Hammond," on account of ease of application.

To be used—
As for Tomes' splint.

To be used—

When angle or vertical ramus fractured.

When more than one fracture.

When great difficulty in rectifying displacement.

In edentulous patient, wherever the site of fracture.

With these forms-

Upper jaw forms a fixed point for the lower to be held motionless against it.

7. Surgical wiring of fragments.

To be performed—

When fragments displaced considerably.

When difficulty of retaining splints from any cause.

General directions to patient.

Jaw not to be moved, if bandaged up.

Soft food with Hammond's, Tomes', and Hayward's splints.

Slop diet with Gunning's splint and Angle's methods.

Good mouth-wash and toothbrush frequently used.

If slight— b. MAXILLA

Ligature teeth together.

If extensive—

Small Hammond's or Tomes' splint.

Complications of Fracture of Mandible.

Wounds of face.

Dislocation or fracture of teeth.

Abscess, necrosis, and sinuses.

Salivary fistula.

Injury to base of skull.

Injury to brain, through glenoid cavity.

Dislocation of jaw.

Irregular union.

Ununited fracture.

Splintering of bone.

Nerve injured or involved in callus—Anæsthesia.

Neuralgia.

Paralysis.

Infiltration of blood about base of tongue— Dyspnœa.

Hæmorrhage.

General—

Shock. Septicæmia, &c. Erysipelas. Fat embolism.

CHAPTER XVIII

DISLOCATION OF TEMPORO-MAXILLARY ARTICULATION

Causes :-

Direct Violence-

Blows.

Kicks.

Falls.

Violent extraction.

Muscular Action-

Yawning.

Shouting.

Opening mouth wide to put large things in

Predisposed to by-

Previous dislocation.

Weakly young subjects—loose joint.

Arthritis.

Varieties: -

Unilateral.

Bilateral.

Complicated with fracture.

Symptoms:

Deformity of jaw.

Projection below malar bone—coronoid process.

Projection above zygoma—contracted temporal muscle.

Masseter muscle hard and prominent.

Hollow behind condyle.

Pain and swelling.

Dribbling of saliva.

Loss of power-speech and deglutition altered.

Treatment:

Reduction at once by disengaging condyle from emenentia articularis, and application of 4-tailed bandage for some days.

CHAPTER XIX

INABILITY TO OPEN THE MOUTH

(" FORCED CLOSURE ")

May be-

- I. Complete.
- 2. Incomplete.

Causes :-

- 1. Organic lesions.
- 2. Spasmodic conditions.
- 3. Traumatic conditions.

1. Organic Lesions :-

- a. Infiltration of soft tissue around vertical ramus and angle with inflammatory products—
 - (i) Lower 3rd molar trouble.
 - (ii) Inflammation of lymphatic glands. (Often incorrectly called Trismus.)
- b. Diseases of joint—
 - (i) Acute or chronic arthritis.
 - (ii) Anchylosis—bony or fibrous.
- c. Fibrous bands in cheek, i.e. cicatrices from previous ulceration, &c.—

Burns.

Lupus.

Caustics.

Cancrum oris.

Ulcerative stomatitis, &c.

- d. Parotid and other tumours.
- e. Mumps.
- f. Deep-seated growths.
- g. Exostosis of zygoma.
- h. "Bony bar"—ossification of pterygo-maxillary ligament.

2. Spasmodic Conditions:

a. Trismus-spasm of masticatory muscles.

Causes :-

Inflammatory conditions. Erupting wisdom tooth. Carious tooth. Hysteria.

- b. Tetanus—due to tetanus bacillus.
 Closure of jaw may be first symptom, followed by spasm of other muscles.
- c. Strychnine poisoning.
- d. Hydrophobia.

3. Traumatic Conditions: -

- a. Fracture—especially if severe.
- b. Dislocation-jaw fixed open.

CHAPTER XX

SWELLINGS ABOUT THE JAWS

Diagnosis: -

First Determine-

- Connected with soft parts only—freely movable over jaw.
- 2. Connected with jaw-
 - a. Attacked externally—circumscribed swelling.
 - b. Involving substance—fusiform swelling.

If involving substance of jaw, it is possibly of dental origin, therefore—

Next Determine-

- 1. Fluid swelling.
- 2. Solid swelling
 - a. Innocent.
 - b. Malignant.

FLUID SWELLINGS

Characteristics of all :-

- 1. Regular, smooth, and globular
- 2. May fluctuate.
- 3. May be egg-shell crackling.

- 4. Thinner wall of jaw (outer plate) enlarging first.
- 5. Alveolar border not interfered with.

 Besides the above points common to all fluid swellings in jaw, note also the following individual characters:—

a. Abscess.

Presence of root or dead tooth.

Presence of inflammation somewhere.

Short history (comparatively).

Varies in size, perhaps, or—

Came quickly and remained so.

b. Dental Cyst.

Presence of root or dead tooth.

No inflammation around (unless it suppurates later).

Slow, steady growth.

c. Dentigerous Cyst (Follicular Odontome).

Absence of tooth from arch usually. No inflammation around. Very slow growth.

Age—15 to 20 years.

Skiagraph may show tooth.

d. Multilocular Cyst (Epithelial Odontome).

Absence of tooth from arch usually. No inflammation around. Very slow growth.

Age—15 to 20 years.

Tendency to be nodular.

e. Cyst of Independent Origin.

Mucous cyst of antrum.

No inflammation around.

SOLID INNOCENT SWELLINGS

Characteristics of All:

- 1. Slow growth.
- 2. Solid feel—according to kind and consistency.
- 3. Alveolar border displaced little.
- 4. Affects one wall of jaw as much as another.
- 5. Little pain.

Forms :-

Fibroma.

Enchondroma.

Osteoma (Exostosis).

Myxoma.

Solid odontomes-

Fibrous.

Cementoma.

Composite.

Radicular.

SOLID MALIGNANT SWELLINGS

Characteristics:-

- 1. Quick growth.
- 2. Solid or pulpy feel.
- 3. Alveolar border displaced irregularly.
- 4. Affects one wall more than another, but not necessarily the thinner
- 5. Severe pain.
- 6. Early implication of skin.
- 7. Emaciation.
- 8. Œdema of face.
- 9. Enlarged glands-carcinoma.
- 10. Attacks of hæmorrhage if superficial.

Forms: -

Sarcoma.

Age-young or adults. Tendency to bleed.

Carcinoma.

Age-adults or aged. Glands affected.

CHAPTER XXI

CLEFT PALATE

Causes :-

- 1. Congenital.
- 2. Acquired.

1. CONGENITAL

Due to-

Arrest of development.

N.B.—Not merely an arrest of union, but also arrest of formation of tissue.

Varieties :-

Any extent of cleft, from a bifid uvula up to a complete cleft passing through alveolus on one or both sides. May then be associated with single or double hare-lip.

N.B.—With this latter the premaxilla is deformed and very much protruded.

Relation of teeth to cleft through alveolus:—

a. (Central on mesial side) Most (lateral missing). Canine on distal side. (Central on mesial side.

b. Small tooth—generally called precanine (?lateral) with canine on distal side.

c. Lateral on mesial side.

Effects :--

Deglutition interfered with little.

Speech interfered with great deal—

D pronounced as N. B pronounced as M.

G and K are impossible.

Treatment:

A. Operative.

B. Mechanical.

A. Operative—

Staphylorraphy—union of soft palate. Uranoplasty—union of hard palate.

Very many different plastic operations of the above have been devised. Tendency now with most surgeons is to operate early, i.e. when patient is about 1 to 3 months old.

B. Mechanical—

a. Velum-

A soft, elastic, and movable appliance, fixed to the back of a denture, and which should imitate the natural soft palate.

Forms-

(i) Sercombe's.

Flap of sheet rubber cut out to cover cleft and extend to within quarter of inch of posterior wall of pharynx, and sewn on to a metal flange projecting from the posterior edge of a denture.

(ii) Kingsley's.

Double flap of soft rubber exactly fitting and embracing the sides of the cleft.

b. Obturator-

A hard, rigid plug of vulcanite or metal that fills in the cleft and is fixed to the back of a denture.

Forms-

- (i) Suerson's.Wedge-shaped plug of vulcanite with base upwards.
- (ii) Kingsley's.

 Large hollow bulb of vulcanite.

A velum is the better appliance with which to learn to speak correctly, but when once this is accomplished, an obturator may be substituted with advantage, as the velum wants frequently renewing on account of deterioration of the soft rubber.

When the appliance is first inserted, the patient must actually be *taught* to speak, which before was impossible. The *immediate* effect of the plate is very little as a rule.

2. ACQUIRED

Causes :-

a. Accidents.

Stabs.

Gun shots, &c.

- b. Syphilis.
- c. Removal of growths.

Varieties :-

Usually asymmetrical.
Usually perforation anywhere in palate.
Rarely cleft through posterior edge.

Effects :-

Deglutition. Speech. Both interfered with great deal.

Treatment :-

A. Operative—

Plastic operation if feasible and *not* due to syphilis.

N.B.—Site of previous syphilitic ulceration would probably break down on attempting to close.

B. Mechanical-

- a. Simple denture if hard palate only involved.
- b. Denture with sheet rubber extension if soft palate involved.

CHAPTER XXII.

DANGERS ATTENDING ADMINISTRATION OF NITROUS OXIDE

A. Connected directly with the N₂O

- I. Cyanosis (Patient dark-blue colour)—
 - Obstructed respiration from
 - a. Tongue falling back.
 - b. Tongue swelling.
 - c. Growths in mouth or neck.
 - d. Œdema of mouth or throat.
 - e. Malposition of head.
 - f. Tight clothing.
 - g. Mucous in air passages, from catarrh.
 - h. Vomit in air passages.
 - i. Lung disease.
 - j. Overdose.
- 2. Syncope (Patient pale)
 - a. Heart disease.
 - b. Weak heart.
 - c. Alcoholics (bad arteries).
 - d. Fright.
- 3. Shock
 - a. Action of N₂O.
 - b. Operating before Vagus centre inhibited.
 N.B.—Vaso-motor centre paralysed before Vagus centre.

- 4. Struggling of Patient
 - a. Alcoholics.
 - b. Neurasthenics
- 5. Apoplexy (Patient paralysed somewhere)
 - a. Granular Kidney.
 - b. Syphilis.
 - c. Arterial Sclerosis.
- 6. Miscarriage.

B. Connected with Operation

- I. Foreign Body in Air-passages or Œsophagus— E.g. a. Tooth.
 - b. Piece of forceps.
 - c. Piece of gag.
 - d. Denture.
 - e. Blood clot.
- 2. Tooth Broken or Forced Out with Gag-

Precautions and Remedies for above.

- 1. Avoid operating under two hours after a meal.
- 2. Loosen clothing.
- 3. Remove dentures.
- 4. Avoid placing gag on loose or frail tooth.
- 5. Remove each tooth from mouth as extracted.
- 6. Use cloth to catch upper roots.
- For Cyanosis—a. Open mouth with Mason's gag, or screw gag if necessary.
 - b. Pull out tongue with tongue forceps.
 - c. Artificial respiration in prone position.
 - d. Tracheotomy—as a last resource.
- 8. For Syncope or Shock
 - a. Amyl nitrie capsules.
 - b. Hypodermic injection of alcohol or ether.
 - c. Artificial respiration.

CHAPTER XXIII

HÆMORRHAGE AFTER TOOTH EXTRACTION

Causes : --

- 1. Laceration with tearing of large vessel.
- 2. Vessels not contracting—
 Old age.
 Syphilis.
 Inflammation.
- 3. Blood not clotting—
 Sea scurvy.
 Purpura.
 Hæmophilia.

Menstruation.

Varieties :-

- 1. Primary.
- 2. Recurrent.

I. PRIMARY

Ordinary hæmorrhage should cease in a few minutes, but if not, means must be taken to stop it, starting with the milder remedies:—

- a. Ice-cold water or ice to suck.
- b. Styptics on wool in socket for 15 minutes or more.
 - E.g. Adrenalin chloride.

 Tincture of Hamamelis.

 Tannic acid.

 Perchloride of Iron (strong solution).

c. Pressure with any of above for few minutes. If above fail, then proceed as given below for recurrent hæmorrhage.

2. RECURRENT

This usually comes on some few hours after the extraction. First observe whether blood clotting or not.

If clotting, give one of the following:-

a. Ext. Ergotæ Liq. m. xx. every 3 hours;

b. Injectio Ergotinæ Hypodermica, m. x.

If not clotting, give one of the following:—

a. R. Liq. Ferri Perchlor. m. xx.;
Mag. Sulph. gr. xv.;
Syrupi Limonis m. xxx.;
Aq. ad \(\frac{7}{3}\)i.
Dose, \(\frac{7}{3}\)i. every 3 hours.

b. Calcii Lact. gr. x., in tabloids every 4 hours.

Next observe whether bleeding from-

a. Gums.

b. Socket.

a. Gums.

If capillary oozing-

Pressure with pad of styptic (see above).

If arterial bleeding—

Twist vessel with artery forceps.

If above fail—

Apply pad of styptic and 4-tailed bandage till next morning.

b. Socket.

Wash well out and plug tightly at once with one of following:—

(i) Lint, tape, or wool soaked in styptic.

(ii) The extracted tooth.

(iii) Plug of wood, wrapped in lint and styptic.

Keep in place with one of following:-

- (i) Pad of lint or cork and 4-tailed bandage.
- (ii) Wedge of wood between adjoining teeth.
- (iii) Figure-of-eight ligature around adjoining teeth.

Leave for 24 hours. If still bleeding on removal—replug. This may be done 3 or 4 times if necessary. More drastic methods if above fail:

- (i) Actual cautery.
- (ii) Trephine inferior dental canal (if for lower tooth), and plug with ivory peg.
- (iii) Digital pressure on common carotid artery.
- (iv) Tie common carotid artery.

General Directions :---

Whether bleeding from gums or socket— Avoid all stimulants. Avoid hot drinks. Avoid all excitement. During day, remain sitting with body erect. During night, recline with head raised.

HÆMOPHILIA

It is extremely advisable never to operate at all on a true hæmophilic patient, as the risk of hæmorrhage and the difficulty of control is graver as a rule than that of leaving a diseased tooth. If, however, an extraction is absolutely necessary, it should be done with the consent and under the supervision of the medical attendant, and the patient should be given one of the following for at least 10 days before, and the operation performed in the morning, to give all day for treatment if necessary:-

- a. R. Ext. Ergotæ Liq. m. xxx.
 Liq. Ferri Perchlor. m. xv.
 Ac. Citrici gr. v.
 Aq. Chloroformi ad 3i. misce.
 Dose—3i. three times a day.
- b. R. Calcii Lact., gr. x. (tabloids).

 Dose—One tabloid three times a day.

CHAPTER XXIV

DRUGS COMMONLY USED IN DENTAL SURGERY

ACTIONS OF DRUGS

1. Analgesics—See Sedatives.

2. Anæsthetics (local) and Anodynes—

Act upon and depress nerve endings, thereby rendering conduction of impulses impossible, and therefore preventing or relieving pain.

Used as injections and local applications.

Examples-

Aconite, tincture of.
Arnica, tincture of.
Cloves, oil of.
Cocaine, hydrochloride of.
Creosote.
Eucaine.
Ethyl, chloride of.
Menthol.
Novocaine.

3. Antacids-

Have direct counteraction to acid conditions in the mouth.

Used in mouth-washes and tooth-powders.

Examples—

Magnesium, heavy carbonate of.
Magnesium, hydroxide of.
Sodium, bicarbonate of.
Precipitated chalk.

4. Antiseptics—

Prevent putrefaction by arresting growth of, or destroying, micro-organisms.

Used in mouth-washes, lotions, and applications.

Examples-

Alcohol.

Carbolic acid.

Chinosol.

Cloves, oil of.

Copper, sulphate of.

Creosote.

Formalin.

Hydrogen, peroxide of.

Iodine, tincture of.

Lysol.

Mercury, perchloride of.

Silver, nitrate of.

Thymol.

Zinc, chloride of.

5. Aperients—

Mildly increase the secretions of the intestines and the peristaltic action, and relieve blood pressure.

Examples—

Carbonate of magnesium. Sulphate of magnesium, &c.

6. Astringents—

a. Coagulate protoplasm of upper layers of cells.

b. Constrict vessels and lessen exudation.

Used in mouth-washes, tooth-powders, and applications.

Examples—

Alum.

Potassium, chlorate of.

Iodine, tincture of. Tannic acid.

Copper, sulphate of. Rhatany, tincture of.

Myrrh, tincture of. Zinc, chloride of.

7. Caustics (escharotics)—

Destroy the surface of living tissues. When acting upon ends of dentinal fibrils they are rendered insensitive.

Examples—

Arsenious acid. Carbolic acid. Copper, sulphate of. Silver, nitrate of. Zinc, chloride of.

8. Counter-irritants—

- a. Dilate superficial vessels, thereby causing free flow of blood through the part, and hence blood and plasma drawn more rapidly through the deeper and inflamed area.
- b. Irritate the superficial nerves, and thereby reflexly relieve the pain from the deeper nerves irritated by the inflammation.

Example—

Iodine, strong solution of (Liq. Iodi Fortis).

Iodine, tincture of.

9. Emollients—

Soften, relax, and protect the mucous membrane. Used in mouth-washes and applications.

Examples-Glycerine.

10. Escharotics—See Caustics.

11. Hæmostatics (general)—

Given internally, constrict arteries and lessen frequency of heart or assist the blood to clot, thereby tending to arrest hæmorrhage. Given by mouth or by injection.

Examples—

Ergot, liquid extract of.

Ergotin.

Iron, perchloride of (strong solution).

12. Obtundents—

Applied locally to deaden pain (see Chap. XXV.).

13. Sedatives (nervine)—

Given internally, act upon central nervous system, dulling the conduction and reception of impulses, thereby relieving pain.

Examples-

Antifebrin.

Antipyrin.

Aspirin.

Phenacetin.

14. Styptics (local hæmostatics)—

- a. Hasten coagulation of blood by precipitation of albumen.
- b. Constrict broken vessel.
- c. Coagulate perivascular tissue.

Thereby tending to arrest hæmorrhage.

Used as local applications.

Examples—

Adrenalin, chloride of.

Hamamelis, tincture of.

Tannic acid.

Iron, solution of perchloride of (tends to slough tissues).

USES OF DRUGS

Uses.

Strength or Dose. Action.

Aconite, Liniment of.

Periodontitis . . Pure (B.P.) Anodyne

Pulpitis . . . ,,

Uses.	Strength or Dose.	Action.
Adrenalin Chloride.		
Hæmorrhage .	. I in 1000	Styptic
Pulp extirpation .	. I in 1000	
	(with cocaine	,,
Tooth extraction .		,
	• "	"
Alcohol, Absolute (C ₂ H ₆ O)		01 / 1 /
Sensitive dentine .		Obtundent
Septic pulp cavities	. Pure or	
	with HgCl2	Antiseptic
Mouth-washes .	. about I in 20	"
Cavity drying .	TO TO	
		agent
Alum (Al ₂ (SO ₄) ₃ K ₂ SO ₄).		agone
111111 (1112(004)3112004).	Mouth washes	Astringent
Inflammation of mouth	Mouth-washes,	Astringent
Inflammation of mouth		
Laceration of mouth	1	,,
Ulceration of mouth	. Glyc. Alumi-	
	nis (I in 7)	
Antifebrin (C6H5NHC2H3O)).	
Periodontitis		Sedative
Pulpitis		
Pain from erupting 3rd	• ,,	,,
molar		"
Neuralgia		,,
Antipyrin (C ₆ H ₅ (CH ₃) ₂ C ₃ HI	N ₂ O).	
As for Antifebrin.		
Arnica, Tincture of.		
Laceration of mouth	.) Mouth-wash	Anodyne
Ulceration of mouth	$mvm.x.to \overline{3}i$	
Arsenious Acid (As,O,).		
	or 1	Caustic.
Devitalising pulp .	- 10	Caustic.
Sensitive dentine (fa		
from pulp)		"
Aspirin (CH ₃ COOC ₆ H ₄ COO)	H).	
As for Antifebrin.		

Calcium Carbonate (CaCo ₃) (precipitated chalk).	Action.
Tooth-powders . 3v. or vi. to 3i. clea	
	ent
Carbolic Acid (C ₆ H ₅ OH).	
All septic conditions in [1 in 200 to 1 in]	
mouth 60 in mouth- Ant	tiseptic
After extractions . wash	
Syringing septic socket 1 in 40	"
Syringing abscess . 1 in 40	11
Sterilising instruments I in 20	1)
Tooth-powders . gr. v. to 3i.	**
Mopping out cysts .	
Painting ulcers Ac. Carbolici	
Pain after extraction Lig (BP)	Caustic
Sensitive dentine . —000/	caustic
Kinnig pulps of teni-	
porary teeth	
Cinnamon, Oil of.	
· · · ·	tiseptic
Flav	vouring
	Ourme
	igent
Chinosol.	agent
	agent
Chinosol.	agent
All septic conditions in a in 500 Anti- mouth after extractions in mouth-wash Syringing septic socket	agent
Chinosol. All septic conditions in 1 in 500 Antimouth after extractions in mouth-wash Syringing septic socket	agent
All septic conditions in 1 in 500 Antimouth after extractions in mouth-wash Syringing septic socket	septic
Chinosol. All septic conditions in 1 in 500 Antimouth after extractions in mouth-wash Syringing septic socket Syringing Abscess Cloves, Oil of (Oleum Caryophylli).	septic
Chinosol. All septic conditions in 1 in 500 Antimouth after extractions in mouth-wash Syringing septic socket Syringing Abscess Cloves, Oil of (Oleum Caryophylli).	septic
Chinosol. All septic conditions in 1 in 500 Antimouth after extractions in mouth-wash Syringing septic socket Syringing Abscess Cloves, Oil of (Oleum Caryophylli). Pulpitis Pure And Exposed pulp	septic odyne
Chinosol. All septic conditions in a in 500 Antimouth after extractions in mouth-wash Syringing septic socket Syringing Abscess Cloves, Oil of (Oleum Caryophylli). Pulpitis Pure And Exposed pulp	septic odyne
Chinosol. All septic conditions in a lin 500 Antimouth after extractions in mouth-wash Syringing septic socket Syringing Abscess Cloves, Oil of (Oleum Caryophylli). Pulpitis Pure And Exposed pulp	septic odyne ,,
Chinosol. All septic conditions in a lin 500 Antimouth after extractions in mouth-wash Syringing septic socket Syringing Abscess Cloves, Oil of (Oleum Caryophylli). Pulpitis Pure And Exposed pulp	septic odyne nt anti- eptic

Uses.	Strongth or D	loca Action		
Cocaine Hydrochloride (C ₁₇ H ₂₁ NO ₄ HCl). Strength or Dose. Action.				
Extractions .	171121110 41101).			
Demonal of mouth				
т .	gr. ½ in m. x.) Local		
Manalakama		anæsthetic		
Opening ontown				
	.) .			
1	. Paint, 20 %	200		
TO 1 1 1 1 1	. ,,	,,		
	Paint, 5 %	,,		
	. Saturated	,,		
	solution			
Copper Sulphate (CuSO ₄).				
Syringing abscess, &c.	gr. ij. to \(\frac{7}{3} \)i.	Antiseptic		
		& astringent		
Pyorrhœa	Powdered	"		
	crystals			
Follicular stomatitis .	Touch with	Caustic		
	crystal			
Root dressing . Po	owdered crystals	Antiseptic		
Creosote.				
Root dressing	Pure (with	Antiseptic		
e	equal quantity	and		
	of Formalin)			
Solvent for As ₂ O ₃	. Pure	Anodyne		
Ergot, Liquid extract of.				
	m. xxx. t.d.s.	Hæmostatic		
Ethyl Chloride.				
Extractions	Pure	Local		
(sprayed on)	anæsthetic		
Small operations on	"	,,		
gums				
Opening abscess	"	5		
Sensitive dentine	Pure (on	"		
(small cavities)	cotton wool)			

Uses. Stren	gth or Dose. Action.		
Eucaine (C ₁₉ H ₂₇ NO ₄ HCI). As for Cocaine.			
Formalin (40 percent. solution of Formic Aldehyde in water).			
All septic conditions of			
mouth in mo	uth-wash		
Syringing septic socket 1	in 1000 ,,		
Syringing abscess, &c. 1	in 1000 ,,		
Sterilising instruments I	in 500 ,,		
Root dressing 40% s	solution "		
Glycerine.			
Most mouth-washes .about	1 in 20 Emollient		
Hamamelis, Tincture of.			
Hæmorrhage Pi	ure as Styptic		
	lication		
Inflammation of mouth m. xx			
	uth-wash		
	a car ii casar		
Hydrogen Peroxide (H ₂ O ₂).			
All septic conditions of mouth	vols. Antiseptic		
Syringing abscess, &c. 10			
Pyorrhœa 20			
	about 6%)		
Root dressing 20			
N.B.—" 20 vols." indicates that one volume of the liquid is capable of giving off 20 volumes of oxygen.			
Iodine,			
Periodontitis Lic	a. Iodi Counter-		
	(I in 9) irritant		
Pulpitis	Iodi Antiseptic		
(I in	40) & astringent		
Syringing abscess, &c. Iodi			
	w colour)		

Uses. Strength or Dose. Action. Iron, Perchloride of (strong solution) (Liq. Ferri Perchlor. Fort.). Hæmorrhage . . see Chap. XXIII. Lime, Chloride of (CaOCl₂). Pure Bleaching Discoloured teeth . (see Chap. XXV.) agent Very foul condition of gr. x. to 3i. Antiseptic mouth in mouth wash Lysol. Sterilising instruments I in 50 Antiseptic & lubricant Magnesium Carbonate (Heavy) ((MgCO₃)₃Mg(HO)₂). Oral acidity gr. xx. to 3i. Antacid (mouth-wash) Tooth-powders . . 3ij. to 3i. Antacid & cleansing Magnesium Hydroxide (Mg(HO)₂). Oral acidity .) gr. xxiv. to 3i. Artacid Rampant Caries in mouth wash Menthol (C, H, O). Sensitive dentine . Pure or with Obtundent Carbolic Acid Mercuric Chloride (Hydrargyri Perchloridum) (HgCl₂). All septic conditions of I in 5000 Antiseptic mouth to I in 2000 Syringing abscess, &c. I in 1000 Root dressing . . . I in 50 (alcohol) Syringing pyorrhœa . I in 500 Myrrh, Tincture of. Inflammation of mouth m. xv. to 3i. Astringent (mouth-wash) Laceration. . 27

Uses. Strength or Dose. Action. Novacaine (NH₂C₆H₄COOC₂H₄N(C₂H₅)₂). As for Cocaine. Phenacetin (C10H13NO2). As for Antifebrin. Potassium Chlorate (KClO3). Inflammation of mouth gr. i. to x. inter- Specific nally & gr. x. to 3i. Astringent (mouth-wash) Laceration. Ulceration . . Tooth-powders . . 3i. to 3i. Potassium Permanganate (KMnO₄). All septic conditions of gr. iv. to 3i. Antiseptic mouth Syringing abcess, &c. Rhatany, Tincture of (Tinctura Krameriæ). Inflammation of mouth m. ij. to 3i. Astringent (mouth-wash) Laceration. . . ,, Ulceration. Silver Nitrate (AgNO₃). Sensitive dentine . Pure Caustic Erosion . . Carious dentine. Antiseptic 2.2 Exposed dentine . Roots before crowning Ulceration Caustic All septic conditions of gr. ij. to 3i. Antiseptic (mouth-wash) mouth

Syringing abscess, &c. gr. iv. to 3i.

DRUGS USED IN DENTAL SURGERY 139

Uses.

Strength or Dose. Action.

Soap, Hard (Sapo Durus).

Tooth-powders . . 3i. to 3i. Antacid and cleansing agent

Tannic Acid (C14H10O9).

Hæmorrhage . . Pure Application to pulp after ,, As2O3

Styptic

Inflammation of mouth

Laceration. . Ulceration. .

a. Mouth-wash, Astringent gr. iv. to 3i.

b. Application Glyc. Ac. Tannici

(I in 5)

. . Saturated sol. Pyorrhæa. in spirit.

Tooth-powders . . gr. xx. to 3i.

Thymol (C10H13OH).

Root dressing . . I in 4 Antiseptic with Zn oxide

Zinc Chloride (ZnCl2).

mouth Sensitive dentine . Erosion . . .

All septic conditions of gr. ij. to 3i. Antiseptic (mouth-wash) & astringent Syringing abscess, &c gr. iv. to 3i. ,, Pure Caustic

,,

PRESCRIPTION WRITING

Superscription.

R = recipe = take of.

Inscription.

Names of the drugs, written in Latin in the genitive case, and their quantities in the accusative case. The last constituent or vehicle is followed by "ad," meaning "up to," so that a definite quantity is always made up, and is generally written in the accusative case.

Subscription.

Directions to the dispenser, written in Latin. E.g. Misce=mix.

Fiat collutorium=let a mouth-wash be made.

Fiat dentifricium = let a tooth-powder be made.

Fiat lotio=let a lotion be made. Fiat mistura let a mixture be made.

Signature.

Directions to the patient, generally written in English—preceded by the word Signa.

Names, &c.

Patient's name and prescriber's initials and date.

Abbreviations.

m.—misce. ss.—semi=half.

s.—signa. ft.—fiat=let it be made.

āā.—ana=of each. q.s.—quantum sufficit=a sufficiency.

Examples.

The first is written out in full, but the others are abbreviated, as is the more usual form.

DRUGS USED IN DENTAL SURGERY 141

I. Antiseptic and anodyne mouth-wash.

Fiat collutorium.

Signum.—" Use with an equal quantity of warm water to rinse the mouth frequently."

(Prescriber's initials)

(Patient's name)

(Date)

2. Antiseptic and astringent mouth-wash.

R Ac. Carbolici . 3i.
Tr. Krameriæ . 3ss.
Glyc. . 3ss.
Sp. rect. . ad 3iv.

Fiat collutorium.

S.—" Use a teaspoonful to half a wine-glass of water to rinse the mouth frequently."

(Prescriber's initials)

(Patient's name)

(Date)

3. Tooth-powder.

Fiat pulv. Mitte 3iv.

S.—" The Tooth-powder."

(Prescriber's initials)

(Patient's name)

(Date)

N.B.—The soap may be omitted if objected to, and the essential oil may be replaced by Ol. Cinnamoni, Ol. Caryoph., or Ol. Rosæ, or combinations of these to suit the patient's taste.

4. Liquid Dentifrice.

Misce.

S.—" A few drops to be placed on a moist tooth-brush, and used vigorously."

(Prescriber's initials)

(Patient's name)

(Date)

5. Counter-irritant.

R Lin. Aconiti Liq. Iodi Fort. āā 3ij.

Fiat applic.

S.—Apply to the gum over the affected tooth with a camel's-hair brush 2 or 3 times a day.

(Prescriber's initials)

(Patient's name)

(Date)

6. Aperient.

R Mag. Carb. Lev. . 31.
Mag. Sulph. . . 3ij.
Aq. Menth. Pip. ad 3vj.
Fiat mist.

S.—" A sixth part twice a day."

(Prescriber's initials)

(Patient's name)

(Date)

CHAPTER XXV

TREATMENT OF HYPERÆSTHESIA OF DENTINE

Manipulation.

Sharp and warm instruments.
Purposive movements—no scraping.
Use much tact, sympathy, and forbearance.

Drugs.

1. For Immediate Effect:

a. Dehydration of Fibrils-

Rubber dam.

Blasts of hot air.

Alcohol. Applied on wool.

b. Cauterisation of Fibrils-

Carbolic (liquid).

Menthol.

Carbolic and menthol.

Zinc chloride (deliquesced).

Silver nitrate (20% sol.).

Applied on wool and dried out with hot air.

c. Production of Cold-

Chloroform.

Ether.

Applied on wool

Ethyl chloride.

143

d. Direct Anæsthetic Action-

Oil of cloves.

Oil of Cinnamon.

Oil of peppermint.

Cocaine hydrochloride.

Menthol.

Novocaine.

Eucaine.

Stovaine.

Cataphoresis and cocaine.

Injection of cocaine into dentine by special syringe.

Applied on wool.

Injection into gum of-

Cocaine.

Novocaine.

Stovaine.

Eucaine.

2. For Later Effect: -

Carbolic acid-sealed in for few days.

Oil of cloves—sealed in for few days.

Arsenious acid—sealed in for 24 hours (very shallow cavities).

Paraform paste-sealed in for few days.

Made as follows :-

Paraform—I part.

Zinc oxide—19 parts.

Carbolic and menthol, q.s. to make a paste.

Oxychloride cement, for few weeks.

Oxyphosphate cement, for few weeks.

General Treatment.

Improve general health.

(Teeth more sensitive when patient unwell.)

Milk of magnesia, if exposed cavities sensitive.

DEVITALISATION OF THE PULP WITH ARSENIOUS ACID

Action of As₂O₃ on the Pulp.

Dilates vessels.

Forms compound with hæmaglobin—arsenhæmaglobin.

Increase in size of connective tissue cells—3 or 4 times their normal size.

Increase of nuclei of neurilemma of nerve fibres, and wasting of axis cylinders.

Pulp and dentine stained brown colour.

Contra-indications to its Use:

- When Pressure Anæsthesia can be conveniently used.
- 2. Open apex. (See Chap. II. for dates of formation.)
- 3. When great pain has been caused by previous use.
- 4. Patients susceptible to action of As₂ O₃.

Necrosis.

Suppuration.
As. poisoning.

Told by previous use.

5. Where it cannot be sealed in.

6. Previous or existing periodontitis.

7. When patient cannot be seen for long time.

8. Patients taking arsenic medicinally may possibly be immune to the action on their pulps. (?)

Action Retarded by:

1. Faulty application—

Too little.

Not over the exposure.

Pushed off by filling.

Washed out by leaking, &c.

- 2. Too thick layer of dentine.
- 3. Chronic inflammation of pulp, especially—
 Pulp stones.
 Secondary dentine.
 Fibroid degeneration.
 Polypus.
 Suppuration of surface.

THE BLEACHING OF TEETH DISCOLOURED BY DEATH OF PULP

Preliminary Procedure.

Prepare cavity.

Clear canal and render aseptic.

Plug apex.

Then apply any of following methods, using rubber dam and platinum broaches:—

1. Chlorinated Lime.

Fill canal and pulp cavity with the chloride of lime moistened with 50 per cent. acetic acid. Seal with G.P.
Renew in 2 or 3 days until bleached.

2. Sulphur Dioxide.

Dessicated sodium sulphite, 10 parts.

Dessicated boracic acid, 7 parts.

Pack in mixed powder, moisten with water, and seal at once.

SO₂ is liberated.

3. Pyrozone.

Mop out cavity continually for \(\frac{1}{4} \) hour. Seal little in for few days.

4. Sodium Peroxide.

Put saturated solution on asbestos wool in cavity for 10 minutes.

Wash with warm water.

Seal in pyrozone for few days.

5. Chlorine Water.

Prepare fresh solution of chlorine gas in water and mop out cavity continually for ½ hour.

CHAPTER XXVI

THE USE OF X-RAYS IN DENTAL SURGERY

As a Means of Diagnosis in the following:—

- Unerupted tooth—especially lower 3rd molars.
- 2. Buried root.
- 3. Orthodontia. To indicate
 - a. Presence of teeth:-
 - (i) Whether extracted.
 - (ii) Whether not erupted.
 - (iii) Whether undeveloped.
 - b. Direction of roots.
 - c. If temporary tooth persists, whether permanent tooth is near.
 - d. Whether root of temporary tooth is absorbed.
 - e. Presence of unerupted supernumerary tooth.
- 4. Exostosis of root.
- 5. Fracture of root.
- 6. Absorption of root.
- 7. Inspection of root canals and canal fillings.
- 8. Pulp stones.
- 9. Odontomes and other solid tumours.
- 10. Bony growth of jaw.
- 11. Fracture of jaw.
- 12. Necrosis of jaw—shown as lighter area and fine cancellous structure destroyed.
- 13. Dislocation of jaw.

- 14. Presence of periodontitis—shown by thickening of membrane.
- 15. Presence of abscess cavity, when not apparent clinically—shown as light area around apex of tooth.
- 16. Presence of cyst—shown as light area.
- 17. Foreign bodies in the antrum.
- 18. Salivary calculus in ducts or gland.

As a Therapeutic Measure in the following: -

- 1. Pyorrhœa Alveolaris.
- 2. Empyema of Antrum.
- 3. Malignant disease.

Not sufficiently tried to justify comment.

Methods of Using.

Outside Method-

Plate placed outside cheek of side required.

Tube placed in such a position on the other side of patient that—

- a. For the mandible, the rays pass obliquely upwards through floor of mouth, thus avoiding the side of jaw not required.
- b. For the maxilla, the rays pass between the teeth of the opposite side, with the mouth propped wide open.

Inside Method-

Protected film placed inside mouth and held against the jaw.

Tube placed outside.

Especially useful for incisor region.

A tube has been made that can be used inside the mouth, with the plate outside, but it is necessarily very small, and not of much practical value at present.

CHAPTER XXVII HYGIENE OF THE MOUTH

Natural Means of Cleansing.

Breast feeding for at least 9 months—jaw developes better and teeth less prone to caries. Proper mastication of suitable food to develop jaw, and cleanse the teeth.

Slop diet for children should be avoided.

Meals should never be finished with a sticky carbohydrate, but rather with a cleansing fruit. Prevention or correction of mouth-breathing. Prevention of thumb sucking or use of comforters. Correction of irregularities.

Artificial Means of Cleansing.

Teeth should be brushed in all directions, night and morning.

No food should be taken after brushing at night. Tooth-brush should be of medium hardness with serrated edge.

Judicious use of tooth-powders and pastes. (See Chap. XXIV.)

Floss silk to clean interstitial spaces.

Judicious use of toothpicks.

Mouth-washes-

a. Alkaline—to neutralise acidity.

b. Antiseptic—to kill or inhibit growth of bacteria.

(See Chap. XXIV.)

Periodical scaling and cleansing by the dentist. (Compressed air spray, useful adjunct.)

INDEX

ABNORMAL number of teeth, 13, 14 position of teeth, 13, 20 size of teeth, 13, 14 structure of teeth, 13, 15 Abrasion, 70 Abscess, acute alveolar, 50 et seq. bacteriology of, 30 chronic, 54 et seq., 116 complications of, 53 pointing of, 53 sac, 56 Absence of teeth, 15 Absorbent organ, 10 Absorption of permanent teeth, 46, 60 of temporary teeth, 9 Acids in mouth, 37, 39, 40 Actinomycosis, 104 Adventitious dentine, 48 Alcohol, 133 Alimentary tract, affections of during teething, 8 Anæsthetics, local, 129 dangers of N2O, 123 Analgesics, 129 Anchylosis of jaw, 113 Angle's methods of regulating, 22 Anodynes, 129 Antacids, 129 Antiseptics, 129 Antrum, empyema of, 99 Aperients, 130

Appendicitis and diseased teeth, 97
Arsenious acid, action on pulp, 145
causing necrosis, 103
causing periodontitis, 50
uses of, 133
Arthritis, 24, 113
Astringents, 130
Attrition, 70

Bacterial, classification of,
27
Bacteriology of alveolar abscess, 30
of dental caries, 29, 40
of mouth, 27
of pulpitis, 30
of pyorrhœa alveolaris, 31
Bakers' predisposition to
caries, 37
Bite, open, 24
raising of, 23
underhung, 23
Bleaching of teeth, 146
Blue line on gums, 81

Calcium carbonate, 134 Calcium lactate, 126, 128 Calculus, salivary, 83 seruminal, 55, 61 Cancrum oris, 76 Cane sugar, fermentation of, Capping pulp, 44 Carbohydrates in caries, 36 39, 40 Carcinoma, 118 Caries, artificial production of, 41 bacteria of, 29 causes of, exciting, 38 causes of, predisposing, 36 cementum, of, 34 classes of food and, 36 course of, 32, 33 dentine of, 32 enamel of, 32 microscopical appearances physical properties of tood, and, 37 pigmentation in, 33 Caustics, 131 Cellulitis of neck, 53, 97 Cellulose, 40 Cementoma, 91, 92 Cementum, caries of, 34 Chalk, precipitated, 134, 141 Cholesterine, 58, 90 Cicatrices, deformities from, 24, 113 Cleft palate, congenital, 119 acquired, 121 Closure of jaws, 113 Completion of roots of permanent teeth, II temporary teeth, 7 Complications of teething, 8 Composite odontome, 92 Compound follicular odontome, 91 Concussion of teeth, 66 Convulsions, due to teething, 8

Counter-irritants, 131 Crowding of teeth, 20 Cysts, antral, 116 dental, 57, 116 dentigerous or follicular, 90, 110 Cysts, multilocular, 90, 116 over erupting teeth, 8 DANGERS of No.O, 123 Deciduous teeth, abnormalities of, 13 absorption of, 9 complications of eruption of, 8 dates of formation and eruption, 7 gemination of, 13 persistence of, 12 Degeneration of pulp, 46et seq. Dental cyst, 57, 116 Dentigerous cyst, 90, 116 Dentine, adventitious or secondary, 48 caries of, 32 sensitive, 139 translucency of, 33 Dentition, temporary, 6 et seq. permanent, II et seq. Development of jaws, 1 et seq. Diabetes, 64 Dichotomy, 14, 15, 18 Diet, effect on caries of, 36, 39 Dilaceration, 68 Dislocation of jaw, III of teeth, 67 Drugs, 129 et seq. Dyspepsia, 37, 73, 97, 98 EMOLLIENTS, 131 Empyema of antrum, 99 et Enamel, caries of, 32 erosion of, 69 nodule, 19 organ, aberration of, 90

Epilepsy, 98 Epithelial odontome, 90 pearls, 14, 57 root tumours, 57 Epulis, 78 Ergot, 126, 127 Erosion, 69 Eruption of lower third molar 11, 25 of permanent teeth, 11 of temporary teeth, 6 Escharotics, 131 Esquimaux, 36 Exanthematous fevers, 18, necrosis, 105 Exostosis of jaw, 114, 117 of teeth, 59 Expansion of arch, 22, 24

FERMENTATION, 39, 40
Fibroid degeneration of pulp,
47
Fibroma of gum (epulis), 78
of jaw, 117
Fibrous food, 36
odontome, 91
Fifth nerve, neuralgia of, 94
Follicle, aberrations of, 90
Follicular odontome, 90, 116
stomatitis, 74
Food, effect on teeth of, 36,
39
Forced closure of jaws, 113
Fracture of jaws, 107 et seq.
of teeth, 67

Gastritis, septic, 75, 97 Gemination, 18 General diseases from teeth, 97 Gingivitis, 71 Glands, enlargement of lymphatic, 26, 53, 75, 97 Glucose, 36.
Gout, 49, 61, 64, 69, 71, 84, 95
Granuloma, 57
Gum-boil, 55
Gums, diseases of, 71 et seq.
hypertrophy of, 80
inflammation of, 71
tumours of, 78
Gunning splint, 109

Habits of children and irregularities, 21, 24 Hæmorrhage after extraction Hæmorrhagic diathesis, 127 Hæmostatics, 131, 132 Hammond's splint, 108 Hayward's splint, 109 Heart trouble and N2O, 123 Honeycombed teeth, 13, 17 Howship's lacunæ, 9 Hutchinsonian teeth, 16 Hygiene of Mouth, 150 Hyperæmia of pulp, 42 Hyperæsthesia of dentine, 143 Hypertrophy of gums, 79 Hypoplasia of teeth, 17 Hysteria, 95, 98, 114

Immunity, 37
Impacted wisdom teeth, 25
Inclined plane, 23
Infant feeding, 9, 37
Inferior protrusion, 23
Inflammation of gums, 71
of periodontal membrane, 50 et seq.
of pulp, 42 et seq.
Influenza, 64
Injuries to the teeth, 66
Inostosis, 60
Interglobular spaces, 17, 33
Iron, perchloride of, 125, 126, 128

Irregularities of permanent teeth in number, 14 position, 20 structure, 15 size, 14 of temporary teeth, 13

Jaw, actinomycosis of, 104 closure of, 113 condition of, at birth, 2 development of, 1 et seq. dislocation of, 111 fracture of, 107 et seq. method of growth of lower, 4 necrosis of, 103

Jumping the bite, 23

KINGSLEY'S obturator, 121 velum, 120

Lactic acid, 39, 40
Lancing of gum, 8, 26
Lead poisoning, 82
Lips, action on teeth, 11, 20
Liquefaction foci, 35

MALARIA, 64, 71, 95 Malignant tumours, 104, 117 Malnutrition, 98 Mandible, bending of, 24 deficient development of, development of, 2 excessive development of, fracture of, 107 et seq. method of growth of, 4 Masticatory muscles, spasm of, 26, 98 Maxilla, deficient development of, 24 development of, I excessive development of, fracture of, 107 et seq.

Meat diet, 36 Meckel's cartilage, 2 Meningitis, 97 Menstruation and hæmorrhage, 125 Mercurial necrosis, 103 stomatitis, 73 teeth, 17 Method of growth of mandible, 4 Misplaced lower third molar, Mouth bacteria, 27 breathing, 21 Mucous cysts, 116 glands, 39, 69 Multilocular cysts, 116 Myeloid epulis, 78 Myxoma, 117

Nævi of gums, 79
Necrosis of jaws, 103 et seq.
Nervous irritation from diseased teeth, 98
Nervous system, affections of, during teething, 8
Neuralgia, 94
Neuritis, toxic, 97
Nitrous oxide, dangers of,123
Nodules, enamel, 19
pulp, 48
Noma, 75
Nose, affections of, causing empyema, 99
Notches in teeth, 16

OBTUNDENTS, 132, 143, 144
Obturators, 121
Odontalgia, 94
Odontitis infantum, 8
Odontomes, 89 et seq.
Open bite, 24
Osteo-arthritis, 97, 111, 113
Osteoma, 117

PALATE, abscess in, 53 Papilla, aberrations of, 91 Papilloma of gum, 79 Paralysis, due to dental disease, 8, 98 Pemphigus, 81 Penetrating caries, 33 Perchloride of iron, 125, 126, Perforation of palate, 121 Periodontitis, acute local, 50 chronic local, 54 general, 64 general suppurative cervical, 61 productive, 59 rarefying, 60 Permanent teeth, abnormalities of, 14 dates of formation and eruption of, II Permanent teeth, eruption of, gemination of, 18 Persistent temporary teeth, Phosphorous necrosis, 105 Pigmentation in caries, 33, 41 Polypus of gum, 78 of pulp, 46 Pregnancy and caries, 37 Prescription writing, 140 et Pterygo-maxillary ligament, 114 Puerperal fever, 97 Pulp, bacteriology of, 30 capping of, 44 diseases ofacute inflammation, 44 chronic inflammation, 46 degeneration, 47, 48, 49 hyperæmia, 42 stones, 48 Purpura, 81

Pyæmia, 54, 75, 97 Pyorrhœa alveolaris, 61 Radicular odontomes, 92 Reflex dental irritation, 98 Resolution of pulpitis, 45 Respiratory system, affections of, during teething, Retention of temporary teeth, 12 Reticular atrophy of pulp, 47 Rheumatism, 62, 64, 71, 95, Rickets, 7 Roots, absorption of, 7, 9, 60 completion of, 7, 11 Saliva, bactericidal properties of, 37 deposit from, 83 et seq. physical characters of, 37 Salivary calculus, 83 et seq. fistula, 87 Sapræmia, 54, 75, 97, 110 Sarcoma, 118 Scurvy of gums, 81 Secondary dentine, 48 affections due to diseased teeth, 97 Sedatives, 132 Sensitive dentine, 139 Septic absorption from diseased teeth, 97 infection from diseased teeth, 97 Septicæmia, 54, 75, 97, 110

Septicæmia, 54, 75, 97, 110
Sequestrum, removal of, 104
Shock, 110
Sinuses on face, 54, 104
Skin, affections of, due to
teething, 9
Spasmodic closure of jaw, 114
Splints, interdental, 108, 109

Starch, fermentation of, 40

Stomatitic teeth, 17 Stomatitis, follicular, gangrenous, 76 mercurial, 74 parasitic, 77 simple, 73 syphilitic, 77 ulcerative, 74 Strychnine poisoning, 114 Styptics, 132 Suerson's obturator, 121 Sugar, fermentation of, 39, 40 Superior protrusion, 20 Supernumerary teeth, 14 Supplemental teeth, 15 Suppuration of pulp, changes during, 45 Suppurative periodontitis, 50 et seg. Swellings of jaws, 115 et seq. Syncope, 123 Syphilis, effects on teeth of, 16 effects on jaws of, 24, 104 effects on mouth of, 77 Syphilitic necrosis, 104 stomatitis, 77 teeth, 16

Tartar, 83 et seq.
Teeth, affections from diseased, 97
at birth, 3
bleaching of, 142
dislocation of, 67
eruption of, 6, 11
injuries of, 66 et seq.
stomatitic, 17
syphilitic, 16
Teething, affections arising

Temporary teeth, abnormalities of, 13 absorption of, 9 dates of eruption, 7 dates of formation, 7 eruption of, 6 gemination of, 13 persistence of, 12 Temporo-maxillary articulation, dislocation of, III fixation of, 113 Tetanus, 114 Thrush, 76 Tobacco-pipe appearance in caries, 35 Tomes' splint, 109 Tongue, action of, on teeth, Tooth germ, aberration of, 92 Toothache, causes of, 94 Tooth-powders, 141 Toxic neuritis, 97 Translucent zone, 33, 35 Trismus, 26, 54, 98, 108 Tubercular necrosis, 104 Tumours of gums, 78 of jaws, 115 UNDERHUNG bite, 23 Vaccine treatment for pyorrhœa, 63 Vela, 120 Visceral troubles, causing neuralgia, 95 WISDOM tooth, misplaced lower, 25 Wry neck, 98 X-rays in dentistry, 148 Zone of translucency in caries, 33, 35

Printed by ALABASTER,

from, 8

PASSNORB & SONSVLTD.

TD. London and Maidstone.

