

Illustrations of comparative anatomy, vertebrate and invertebrate : for the use of students in the Museum of Zoology and Comparative Anatomy.

Contributors

Clark, John Willis, 1833-1910.
Bridge, T. W.

Publication/Creation

Cambridge : The University Press, 1875.

Persistent URL

<https://wellcomecollection.org/works/rus9ej63>


License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>






22200068198

Med
K7763

F. Leigh.

Jan. 1891.

Longway.



ILLUSTRATIONS
OF
COMPARATIVE ANATOMY,
VERTEBRATE AND INVERTEBRATE,

FOR THE USE OF STUDENTS IN THE MUSEUM OF
ZOOLOGY AND COMPARATIVE ANATOMY.

SECOND EDITION.

Cambridge:
AT THE UNIVERSITY PRESS.

London: CAMBRIDGE WAREHOUSE, 17, PATERNOSTER ROW,
Cambridge: DEIGHTON, BELL, AND CO.

1875.

7 721 420

Cambridge:

PRINTED BY C. J. CLAY, M.A.
AT THE UNIVERSITY PRESS.

WELLCOME INSTITUTE LIBRARY	
Coll.	weIMOmec
Call	
No.	QS

PREFACE.

THE specimens and dissections described in the following pages have been prepared to illustrate the most important forms in Invertebrate and Vertebrate Anatomy.

When the first edition of these descriptions, entitled 'List of Dissections,' was published in 1871, Invertebrate Anatomy alone was illustrated. We have now included four representative Vertebrate forms—Rat, Pigeon, Frog, and Perch: and have increased the series of Invertebrata from 55 to 90; adding to it several specimens, with detailed descriptions, of the Exoskeleton in those forms where such descriptions are not to be met with in the text-books that are commonly used by English students.

All the descriptions have been carefully revised. As in the former edition, we have not attempted to give a complete account of each dissection, but merely an indication of its leading features, to enable students more readily to recognise the points dwelt upon in the literature of the subject, or in lectures; and to shew them clearly the position and relation of the organs which they will subsequently have to examine when they begin to dissect. For this reason the very language used by Prof. Rolleston, Prof. Huxley, and others has been

reproduced where it has been possible to do so. Our obligations to Prof. Rolleston are however far greater than mere passages of description, for it is to his work that our own owes its existence.

For the sake of brevity we have cited as 'Rolleston,' *Forms of Animal Life*, by G. ROLLESTON, M.D., Oxford, 1870.

J. W. CLARK,
Superintendent.

T. W. BRIDGE,
Demonstrator of Comparative Anatomy.

MUSEUM OF ZOOLOGY AND COMPARATIVE ANATOMY,
CAMBRIDGE, May 20, 1875.

ILLUSTRATIONS OF COMPARATIVE ANATOMY, VERTEBRATE AND INVERTEBRATE.

SUB-KINGDOM PROTOZOA.

1. A piece of the common Freshwater Sponge (*Spongilla fluviatilis*), exhibiting its siliceous skeleton.

Rolleston, p. 163.

SUB-KINGDOM CŒLENTERATA.

CLASS HYDROZOA.

2. A Sea-Fir (*Sertularia abietina*).

It consists of a slender pinnately-branched 'cœnosarc' invested by a chitinous 'periderm.' Both the main stem and the lateral pinnæ are beset by numerous sessile or sub-sessile more or less flask-shaped chitinous 'hydrothecæ,' in which the 'polypites' are lodged. In addition to these, larger but similarly shaped vesicles are to be observed studding the pinnæ. These are the 'gonothecæ' and contain the generative zooids. The animal when mature is non-locomotive and is attached to its base by a 'hydrorhiza.'

3. A Sea-Anemone (*Tealia crassicornis*), from which the base has been removed. The tentacles and the space round the mouth—'peristomial disc'—are too much retracted to be well seen. On the other side of the preparation the internal surface of the digestive cavity is exposed, marked with fine radiating lines, which correspond with the attachments to it of the vertical muscular lamellar plates termed 'mesenteries.' These

radiate from the outer surface of the stomach to the inner wall of the outer integument, and thus divide the space between the two—the body cavity—into a number of wedge-shaped compartments. Some of these ‘mesenteries’ fail by greater or less intervals to reach the outer surface of the stomach, and are therefore called ‘secondary’ or ‘tertiary’ mesenteries, while those that are attached to that organ and to the outer integument are called ‘primary.’ Between them are seen some of the generative glands. The cord-like ‘craspeda’ are also to be seen, attached to the inner edges of the mesenteries.

Rolleston, p. 158.

SUB-KINGDOM ECHINODERMATA.

4. A Common Cross-Fish (*Asterias rubens*).

The animal consists of a discoid body, produced horizontally into five radial prolongations. The ventral surface of the disk is depressed into a cavity having a pentagonal outline, and which is further continuous with the mesial lanceolate grooves extending along the ventral surfaces of each ray. The dorsal surfaces of the disk and rays, on the other hand, are convex and not grooved. In the middle of the ventral surface of the disk is the mouth: and on the corresponding part of the dorsal surface, the anus. Protruding from the radial grooves or avenues are the cæcal tube-feet: arranged in four rows, two on either side of a line running down the centre of each ray. There are two sets of spines bounding each avenue, ‘one of which is placed internally and consists of two rows of long and slender spines: while the more externally placed is made up of three rows of shorter and blunter spines.’ Externally to the latter is a third series of spines, marking the line of junction of the dorsal and lateral surfaces. Furthermore the dorsal surface of both rays and disk is studded with variously shaped spines, some short and obtuse, while others carry a small bunch of calcified setæ at their free extremities. None of the spines are moveable. On the extreme edge of the dorsal surface of the disk, and in one of the inter-radial spaces, there is a porous elevation—the madreporic tubercle—through the medium of which, and of the ‘sand canal’ with which it is in relation internally, the sea-

water finds its way to the circular water-vascular canal surrounding the œsophagus. The tube-feet being restricted to the ventral surface of the rays and disk, that surface is distinguished by the term 'ambulacral' from the dorsal and lateral surfaces, which do not possess tube-feet, and are therefore called ant-ambulacral. Though the radial symmetry of the disk and arms is apparently very evident, yet the animal is really bilaterally symmetrical, thus:—the madreporic tubercle is inter-radial, and as both mouth and anus are central, a vertical plane drawn through tubercle, mouth and anus, will divide the animal into two symmetrical halves, each half including two rays and one half of the bisected ray. The two rays between which the madreporic tubercle lies are distinguished as the 'bivium,' while the remaining rays constitute the 'trivium.'

5. Another specimen, to shew the structure of the investing coriaceous test, and the disposition of the ossicles composing it.

The whole of the disk, with the exception of that portion on which the madreporic tubercle is situated, has been removed:—and, with a single exception, the cavity of each ray has been laid open by a longitudinal incision on the ant-ambulacral surface. Four rays have been divided transversely. Each ray consists of a solid axis and of the calcified arch supported by it. The axis itself is composed of a longitudinal series of ossicles, apposed to each other after the manner of vertebræ, and forming a solid though flexible rod extending along the ventral surface of each ray from the mouth to the tip. Each ossicle is double—that is, it is composed of two lateral halves, closely applied in the mesial line, and retained in position partly by a tendinous investment—and partly also by the longitudinal muscular fibres, which, in the fresh condition, pass from ossicle to ossicle. Each lateral ossicle gives off from its outer side a transverse process, which is slightly sinuous, or emarginate on opposite sides: so that by the apposition of the concavities of contiguous processes two alternating rows of foramina for the exit of the tube-feet are formed on each side of the median ossicles. The external extremity of each of the transverse processes abuts upon a small ad-ambulacral ossicle, which in addition to contributing to form the prominent margin of the ambulacral groove, and

to supporting the innermost row of slender spines described in the preceding preparation, also marks the junction of the ambulacral and ant-ambulacral areas. The ad-ambulacral ossicle is best seen in the transverse section. Externally to the ad-ambulacral ossicles is a series of stout, rib-like rods, fewer in number than the transverse processes, as there is but one of the former to three of the latter. These form the lateral boundaries of the ad-ambulacral region. These 'ribs' are connected with each other by at least two well-marked rows of cross-bars on each side; extending from one 'rib' to the other. The 'rib' is not a single ossicle, but composed of several pieces joined together by oblique surfaces: so as to admit of some amount of sliding motion upon one another, in order to allow the consequent dilatation or contraction of the cavity, the walls of which they contribute to form. Towards the dorsal region, or roof of the ray, the 'rib' breaks up, and the pieces cross in all directions, forming an irregular network of ossicles imbedded in the thick integument. The first axial ossicle in each ray is much larger than the rest: and by their abutment upon each other a strong calcified ring is formed, encircling the mouth. The skeleton of the disk in no way differs from the structure of the rays: as it is in fact formed by the confluence of the ant-ambulacral integument of the five rays: and the axial ossicles themselves with their transverse processes are continued to the margin of the mouth, and only then abut upon each other. In the fresh state the radial ambulacral canal would lie in the ambulacral groove immediately beneath the axial series of ossicles, and still more superficially the radial nerve-cord:—while in the cavity of the ray the digestive and generative systems are contained. In the peristomial depression, and external to the calcified oral ring, the pentagonal nerve-collar would lie. It will thus be seen that from its relation to the nervous system and the radial ambulacral canals, the skeleton is really an internal one, and does not correspond to the ambulacral and inter-ambulacral plates of the *Echinus*, which form a true exoskeleton.

6. A Common Cross-Fish (*Asterias rubens*), dissected to shew its digestive system.

The whole of the integument has been removed from the

upper or anal surface. In the centre is the membranous stomach, which has been stained pink, as also have been the five cæcal tubes—'diverticula'—radiating from it. A portion of the integument has been preserved round the anus, into which a black bristle has been inserted. From the intestiniform portion of the digestive tract arise some tufted cæcal appendages. The tubes which are given off from the stomach in the direction of each of the rays very soon divide into two, each of which gives off a number of arborescent cæca, filling up the whole cavity of each ray. At the base of each ray is placed a pair of generative glands. Each gland is made up of numerous cæca, converging to a duct opening at the base of the ray. The madreporic tubercle is between the first and second rays on the right of the preparation, counting from the top. In each ray, and on each side of the median longitudinal ossicles, may be seen numerous 'ampullæ.'

Rolleston, pp. 141—145.

7. A Common Cross-Fish (*Asterias rubens*), dissected to shew its generative system.

The dorsal portion of the integument, with the whole of the digestive apparatus, has been removed. The generative organs, ovaries or testes (the Echinodermata being 'dioecious'), are seen to consist of bundles of arborescent cæca, two to each ray, each bundle being attached at the base of the cavity which contains the digestive cæca, one on each side of the median longitudinal ossicles. The attached extremity of each gland indicates the position of its duct, which opens at the base of each ray in the inter-radial space.

8. A Common Cross-Fish (*Asterias rubens*), dissected to shew its nervous system.

The tube-feet have been removed. The nervous system lies on the ventral surface of the animal, and consists of a pentagonal ganglionated ring surrounding the mouth, from which a single cord passes down the centre of each ray.

9. A crinoid Echinoderm (*Pentacrinus caput-medusæ*), dredged off Barbadoes, 1869.

10. A Sea-Urchin (*Echinus sphæra*), divested of its spines, and dried.

The exo-skeleton is an immovably articulated, inflexible

shell, composed of numerous—upwards of six hundred—calcified pieces, arranged in a definite pattern. The shape of the test is that of a sphere with flattened poles. One pole however, the oral one, is much more flattened than the other,—the anal one. With the exception of small circum-anal and circum-oral areas the remainder of the body, 'corona' as it is called, is composed of numerous calcified plates, disposed in twenty longitudinal series. These are furthermore arranged in ten double rows, radiating from one pole to the other. The five double rows of plates, which are perforated by the external apertures of small canals, constitute the ambulacra, while the remaining five double rows, alternating with the foregoing, and imperforate, form the inter-ambulacra. The ambulacral and inter-ambulacral plates in converging towards the poles of the shell become much smaller, and do not quite reach the poles, but leave small areas round the mouth and anus, in which the calcified plates have a different arrangement.

In the neighbourhood of the anus this space is occupied by two concentric rows of plates, the anal plates. The much larger circum-oral space—the 'peristoma'—is in the fresh specimen filled up by a leathery membrane strengthened by irregularly shaped calcified plates, in the centre of which is the mouth. Each ambulacral row of plates is separated from the adjacent row of inter-ambulacral plates by a tolerably straight but not very well defined suture. On the other hand the suture separating the two rows which together make up an ambulacrum or an inter-ambulacrum is a zigzag one: being formed by the alternation of the triangular internal extremities of the component plates. There is consequently such a suture running down the middle of each ambulacrum and inter-ambulacrum. The plates composing an ambulacrum are pentagonal, and transversely elongate. Each is perforated by six obliquely disposed apertures or pores, affording a passage to the tube-feet: and each carries on its surface conical tubercles varying in size. Furthermore each plate is subdivided by faintly visible oblique transverse sutures into three primary divisions or pore-plates, each pore-plate containing a pair of obliquely set pores: and the whole six apertures are situated close to the outer or inter-ambulacral edge of the plate. 'There is therefore a pore-plate or subdivision of

the ambulacral plate corresponding to each pair of pores: and this arrangement would appear to hold good for all the *Echinoidea* except the *Clypeasteridae*, where the pores of the locomotive feet are grouped together on a single plate.' (Huxley.)

Each row of ambulacral plates carries two rows of alternating conical tubercles, and in addition there are inner tubercles of various sizes scattered between the larger ones: these tubercles carry the spines, which beset the external surface of the skeleton.

The inter-ambulacral plates, though similar in shape, and carrying tubercles, are larger in size than the ambulacral, and imperforate. If an ambulacrum be traced upwards towards the apical pole of the shell, it will be found to abut upon one of the five smaller ossicles which form the outer row of circum-anal plates. Each of these angular plates is perforated, and supports an eye-spot. Hence they are called 'ocular plates.' The apical extremities of the inter-ambulacra are similarly related to the larger plates which form the inner series of circum-anal ossicles, and alternate with the ocular plates. These plates, of the inner row, are likewise perforated with a small aperture for the passage outwards of the generative products. Hence they have been termed 'genital plates.' One of these is much larger than any of the others, and is marked by a porous elevation, the madreporic tubercle: which internally is in relation with the sand canal.

The small area within the ocular and genital plates is occupied by small calcified tubercles and granules, within which the anus is situated. The growth of the shell in the *Echinidae* is effected in two ways: partly by addition to the circumference of existing plates, and partly by the interposition of new ambulacral and inter-ambulacral plates at the apical end of each series. The tubercles which cover the plates terminate in rounded, ball-like summits, either simple, or marked by a small pit. The spines articulating with these tubercles have a cup-shaped depression at their attached extremity, which receives the rounded end of the tubercle in such a manner as to form a ball and socket joint. The spine is attached to the tubercle partly by a ligament which passes from the bottom of its articular surface to the pit in the tubercle: and partly by the muscular fibres which connect the neck of the spine with

the base of the tubercle, and in addition produce all the motion of which these organs are capable.

It will be noticed that in addition to the absence of any lateral prolongation of the body into rays, an *Echinus* differs from a Starfish in the enormous development of the ambulacra at the expense of the ant-ambulacral region: so that the latter eventually becomes reduced to a very small space round the apex. As in the Starfish, the test is bilaterally symmetrical, for on each side of the vertical plane in which the mouth, anus, and madreporic tubercle lie, there are two and a half ambulacra, and two and a half inter-ambulacra. The disposition of the ambulacra into bivium and trivium is also obvious.

11. Two Sea-Urchins (*Echinus sphaera*), with the spines and tube-feet intact.

The spines cover the whole of the ambulacra and ant-ambulacra. They fail however to be developed for some little distance round the mouth, though they extend up to, and even arch over, the anus. Though they vary much in size, they are all of one shape. Each spine is cylindric, striated longitudinally as well as transversely, and tapers gradually towards its apex, which is tipped with purple. The spines that form a continuous series from one end to the other of an ambulacrum or inter-ambulacrum are distinguished as 'primary' spines, while the smaller and less regularly arranged are, in proportion to their size, termed 'secondary' or 'tertiary.' The five pairs of rows of tube-feet are to be seen protruding from the pores. Each foot is a tubular cæcal process, terminating distally in a slightly expanded, disk-like extremity:—while internally it is in connection with the 'ampullæ' and one of the 'radial trunks' of the water-vascular system. In the peristomial area, and immediately surrounding the mouth, the tube-feet become modified into a circle of more or less plumose appendages, which may be regarded as the aborted rudiments of the tentacular crown of the *Holothuroidea*.

12. A Sea-Urchin (*E. brevispinosus*), bisected horizontally, to shew the internal surface of the shell.

The ambulacral plates are produced into five short truncated processes, so perforated as to form five small vertical arches,

which arch over the ambulacra, and are called 'auriculæ.' Through these apertures the radial nerve-cords and the radial canals of the water-vascular system pass from the nerve-collars and water-vascular ring. From the relation which these 'auriculæ' bear to the nervous and water-vascular systems it would seem as though they represented the ambulacral ossicles of the Starfishes.

13. A *Clypeaster* bisected to shew the great development of the 'auriculæ,' which here are so large as to unite the oral and apical walls of the 'corona.'

14. A *Spatangus* (*S. purpureus*), bisected to shew the total absence of 'auriculæ.'

15. A Sea-Urchin (*Echinus sphaera*), prepared to shew its generative and digestive systems.

The test has been horizontally bisected, and the apical half of the shell suspended above the oral half.

The œsophagus, as it emerges from the arrangement of calcified jaws called Aristotle's lantern, to be described in the next specimen, is of moderate calibre, but subsequently it somewhat abruptly expands into a large and thin-walled intestine, which, after turning twice round the inside of the test, passes along an inter-ambulacral space to the anus. At the commencement of its second and superior convolution round the shell it bends upon itself and runs in an opposite direction to its first circuit, to which however it still remains parallel. The two convolutions are attached by a much-fenestrated mesentery to the inside of the test, and are disposed in festoons. Two pseud-hæmal vessels can be seen in relation with the intestine, one running along its inner, and the other along its outer or mesenteric border.

The ovaries are five in number, and are arranged round the apical area, one in each of the five inter-ambulacral spaces. A short tubular oviduct conveys the ova to the apertures before described as existing in the five genital plates.

The ampullæ are seen on the internal surface of the shell in the ambulacral areas; and mesially between each of the two rows of ampullæ in an ambulacral area one of the five radial nerve-cords can be seen.

Rolleston, p. cl.

16. The oral skeleton, whole and separated into its component parts.

17. The lower half of an Echinus (*E. sphaera*), prepared to shew the mouth, and the muscles that move the jaws.

"The apparatus consists of five hollow wedge-shaped calcareous pieces, the 'alveoli,' each of which is composed of two halves united together in the mesial line, while each half again consists of a superior epiphysis and an inferior principal portion. Each 'alveolus' serves as the socket to a long tooth shaped like the incisor of a rodent, harder externally than internally, so as always to develop a sharp edge with wear, and constantly growing from its upper extremity. The five alveoli, fitted together, form a cone, whose applied surfaces are united by strong transverse muscular fibres, while superiorly the epiphyses of each pair of alveoli are connected by long radial pieces, the 'rotulæ.' To the inner extremity of each 'rotula' finally a slender arcuated rod presenting indications of a division in the middle of its length is articulated, and running outwards parallel to the 'rotula' terminates in a free bifurcated extremity. This is the 'radius' or 'compass' of Valentin. Altogether the 'lantern' consists of twenty principal pieces, five teeth, five 'alveoli,' five 'rotulæ,' or 'falces,' and five 'radii' or 'compasses:' of which the alveoli are again divisible into four pieces each, and the radii into two: making a total of forty pieces. In their normal position it must be remembered that the alveoli and teeth are inter-ambulacral, while the 'radii' and 'rotulæ' are ambulacral. Besides the inter-alveolar muscles this complex apparatus has protractor muscles arising from the inter-ambulacral region of the oral edge of the 'corona,' and inserted into the upper part of the alveoli; slender oblique muscles, with a similar origin, but inserted into the radii; transverse muscles connecting the radii together; and retractor muscles arising from the arches of the 'auriculæ,' and inserted into the oral ends of the 'alveoli.'"

Huxley, *Medical Times and Gazette*, 1856, vol. xiii. p. 587.

The next three preparations illustrate the anatomy of the *Holothuroidea*. In each a blue rod has been placed in the mouth: a red rod in the anus.

18. A Great Sea-Cucumber (*Cucumaria frondosa*), verti-

cally laid open. In this specimen the animal has contracted so much, that the whole of the tract round the mouth—'the peristomial space,'—with the tentacles, has been withdrawn within the body-walls, and hangs down like a bag, which has been laid open so as to expose the tentacles, of a black colour.

Five pairs of longitudinal muscular bands, radiating from mouth to anus, are seen to divide the body-walls into a corresponding number of crescent-shaped divisions, in which the muscular fibres, which encircle the animal transversely, are well seen. In the interval between the two elements of each pair of muscular bands lies the longitudinal water-vascular ambulacral vessel. In this family—the *Dendrochirotae*—each longitudinal muscle gives off a long slip which inserts itself into the corresponding radial ossicle of the calcareous ring encircling the mouth, and acts as a retractor. The water-vascular ring surrounds the muscular pharynx, a little way posteriorly to the calcareous ring. In the centre of the preparation are the muscular pharynx and stomach. On the exterior may be noticed the longitudinal rows of apertures for the tube-feet, which, however, are now partially retracted.

Rolleston, p. 145 sq., and p. cxlvii.

19. A specimen of *Psolus phantapus* with part of the dorsal and ventral surfaces removed. The animal is contracted as in the last preparation. The longitudinal and retractor muscles, and the calcareous ring with its two Polian vesicles, are well seen.

On the ventral side of the preparation is seen the whole of the digestive tract, spread out upon a sheet of mica. The anal end receives the ducts of the so-called respiratory trees, which in the preparation depend downwards from the cloaca, one on each side. These organs consist of hollow stems carrying somewhat scanty ramifications; they reach a considerable distance into the body, to the walls of which they are attached by mesenteries.

On the dorsal surface is the generative gland—a bundle of simple unbranched cæca of various lengths. The mesentery, by which it was attached to the body-walls, has been removed. A single efferent duct passes from these cæca to open inside the

circle of oral tentacles, above the calcareous ring. The ventral surface is covered with calcareous scales, and is destitute of tube-feet: though these latter do exist in other parts of the body.

20. A specimen of *Holothuria tubulosa*, dissected to shew the generative system and 'respiratory tree.' The greater portion of the left side of the body has been removed, together with the whole of the digestive tract, except the pharynx and cloaca. These are well displayed, as also are the mesenteric bands by which the cloaca is attached to the body-walls.

The generative cæca have been spread out on a piece of mica. They are disposed in two groups, but are shorter and much less subdivided than in *Psolus*, and end in dilated cæca. The two ducts, one from each group, shortly unite to form the common efferent duct, appended to which, at about the level of the calcareous ring, is a number of minute pyriform vesicles. The external opening of the generative duct has a black bristle inserted into it. The respiratory tree consists of a hollow stem extending from the cloaca, with which it is in communication, as far forwards as the pharynx. Numerous filiform and arborescent cæcal appendages are given off from it. The mesenteric fibres by which the stem is attached to the body-wall are well seen.

On the left of the preparation is the 'respiratory tree,' attached by mesentery to the body-wall. A green rod has been inserted into the orifice by which it opens into the 'cloaca.' Depending from the calcareous ring are the 'ampullæ' of the water-vascular ring surrounding the mouth.

SUB-KINGDOM VERMES.

DIVISION ANNULATA.

Class I. ANNULATA PROPER.

21. An *Amphinome*. The body, though elongated, is of considerable transverse width, and is divided by numerous transverse constrictions into about 60 segments. With the

exception of the cephalic the whole of the segments differ from each other in little else than their comparative calibre. The integument is not hardened by the deposition of chitinous or calcareous matter, and the appendages attached to the different 'somites' are soft, and devoid of segments or of joints. A typical 'somite,' from the middle region of the body, may be said to consist of dorsal and ventral arcs—each arc terminates on each side in a conical eminence, constituting with its appendages ('cirri' and 'setæ') a 'notopodium' or 'neuropodium' according to the arc with which it is in relation. The appendages attached to the 'notopodium' are; first, a tuft of branching cæca dorsally placed; secondly, a fasciculus of chitinous setæ; thirdly, a small 'cirrus.' The neuropodial appendages are separated from the notopodial by a well-marked interspace, and consist of, (1) a small fasciculus of setæ, and (2) a cirrus: but no branchiæ. Towards the head and anal extremity the branchiæ become very small and are absent altogether on the first two post-cephalic segments: and the setæ are also insignificant in size. The mouth and anus are terminal: the former is situated on the ventral surface of the third somite.

22. A Common Earthworm (*Lumbricus terrestris*), dissected to shew its digestive system.

The transversely annulated integument has been divided down the medio-dorsal line and pinned out on either side. The digestive tract is seen in the middle line of the preparation, and extends, without any convolutions, direct from mouth to anus.

The mouth (formed by a modification of the first segment) leads into an oval pharynx, whose external surface has a coarsely villous appearance, due to the breaking away of the delicate muscular bands by which it was attached to the body-wall. The œsophagus, in length about five-eighths of an inch, and of much smaller calibre, leads from the pharynx to the large and globular crop. Immediately succeeding this latter cavity, and of slightly smaller size, and lighter colour, is the so-called gizzard. Posteriorly to the gizzard the intestine is continued as a slightly sacculated tube of even diameter to the anus. Attached to the œsophagus immediately anterior to the

crop are three pairs of minute glands of unknown function called œsophageal glands.

The membranous partitions extending from the body-wall to the alimentary canal, and dividing the perivisceral cavity into a series of inter-communicating spaces, are well seen in the middle third of the preparation. In the inter-mesenteric spaces of the middle third of the body are to be seen the 'segmental organs,' consisting in each space of a looped internally ciliated tube, communicating by its internal extremity with the perivisceral cavity, and by its other extremity opening externally.

Rolleston, p. 119.

23. The same, dissected to shew its nervous system.

The integument, divided in the medio-dorsal line, has been pinned out on either side, and the whole of the digestive and generative systems removed. The nervous tract is seen occupying the medio-ventral line, and extending from the oral to the anal extremity of the body. The supra-œsophageal mass is seen to consist of two pear-shaped lobes applied together in the median line by their broad ends, their apices pointing outwards. From the outer end of each of these two lobes a large nerve passes forwards to supply the upper lip, and a bristle has been placed underneath them. A commissural cord on each side proceeds downwards from the supra-œsophageal ganglia, encircling the œsophagus and connecting these ganglia with the ventral nervous cord.

The anterior portion of the cord is of considerable thickness, and, on close inspection, may be seen to be composed of two distinct strands; the ganglionic enlargements are, however, with difficulty recognizable. In the middle third of the body the cord becomes much thinner, but assumes a beaded appearance owing to the greater comparative size of the ganglia and the wider interspaces which separate them. In the posterior third of the body it again becomes of considerable size, but the ganglia are less distinctly visible and more closely apposed.

A pair of nerves is given off from each pair of ganglia, and a single nerve on each side from the segments of the cord interposed between each pair of ganglia.

24. A Sea-Mouse (*Aphrodite aculeata*), dissected to shew its digestive organs.

The animal has been laid open along the ventral surface, the nervous system having been removed.

The mouth, into which a black bristle has been inserted, leads into a thick, cylindrical, muscular sac, called by some a gizzard. The posterior extremity of this opens by a narrow, sigmoid segment, into a funnel-shaped stomach, which gradually diminishes in size, and insensibly passes into an intestine, terminating at the anus. Throughout the whole length of the stomach, cæcal processes are given off on each side to the number of about twenty pairs. They are slender at their origin, but gradually enlarge, and give off lateral 'diverticula,' which again, in their turn, divide and subdivide, terminating finally in fusiform dilatations. These cæca are probably homologous with the gastric cæca of the leech; and the so-called 'gizzard,' which is capable of protrusion, has been considered to be homologous with the projectile proboscis of other Annelids.

The stomach has been filled with red injection, and the cæca partially so. They have been dissected and pinned out on the left side. On the right they remain *in situ*, lying among the transverse muscles. On this side also may be seen the delicate muscular bands, which pass from the base of the bundles of setæ to the integument, and effect their protrusion.

25. The same, dissected to shew the 'elytra.'

A longitudinal strip of the dorsal coating of 'felt' has been removed and the 'elytra' displayed.

The 'elytra' are broad membranous plates developed as appendages from the notopodia, and, arching inwards over the back, form two lateral series on either side of the medio-dorsal line. Beneath the elytra is the membranous partition forming the dorsal boundary of the visceral cavity. The nervous system may be seen on the ventral surface of the body, through the integument.

It is to be observed that the 'Sea Mouse' is much less distinctly annulated than the Earthworm: and that the 'setæ,' which in the Earthworm are almost microscopic, are here of large size and iridescent.

26. A Leech (*Hirudo medicinalis*), dissected to shew its nervous and generative systems.

The nervous system consists of a longitudinal ganglionated

ventral cord containing about twenty-four ganglia, from which delicate nerves radiate on either side. The ganglia are more approximated at both ends than at the middle of the animal. From the foremost, or sub-œsophageal, ganglion pass the nerves to the oral muscles and the cords on either side of the œsophagus. These, meeting above, join the super-œsophageal ganglion from which radiate nerves to the ten ocelli and other parts of the head.

The testes are nine spherical bodies on either side of the nervous cord connected by their ducts with a common canal, or 'vas deferens,' which runs external to them and communicates in front with a receptacle or 'vas deferens.' The two receptacles are connected with the muscular ball from which the penis is continued. The ovaries, smaller than the testes, are situated, one on either side, just behind the 'receptacula seminis,' and are connected by short oviducts with the muscular bag or vagina.

DIVISION ANNULOIDA.

Class II. NEMATELMINTHES.

27. The generative organs of a male and female *Ascaris* (*Ascaris lumbricoides*, L.), taken from the duodenum of a young woman. The male, on the left, is smaller than the female. It—the male—has been opened longitudinally, and the intestine, a straight tube, is seen passing from the mouth to the anus. It has been pinned somewhat to the left of the specimen. The testis is a long curling tube, very attenuated at its commencement. It gradually becomes thicker, and is disposed in numerous coils round the alimentary canal. About three inches from its termination it suddenly dilates into a larger tube or receptacle, which, narrowing again, is continued to the posterior extremity of the animal. Its lowest part, beneath which a slip of blue paper has been placed, admits of protrusion, and is called the penis.

The female has also been laid open longitudinally. The intestine, like that of the male, is a straight tube, extending the whole length of the animal, but it expands much more

near its anal end. The ovarian tubes commence as thin filiform cæca arranged in complicated coils round the intestine. Near the anus however these tubes become suddenly expanded, and extend as far forwards as the anterior third of the body: where they unite to form a short and very narrow oviduct, which opens outwards. This, turning round the intestine and becoming narrow, terminates at the genital aperture one-third from the anterior end of the animal.

Class III. PLATYELMINTHES.

28. A Tape-Worm (*Tænia solium*).

The specimen, a mature worm (*Strobilia*), is a flattened, ribbon-shaped organism, divided by transverse constrictions into numerous well-defined segments—the ‘proglottides,’—which are for the most part of large size, but on approaching the head they become gradually smaller, and the constrictions less evident, till in the vicinity of the head the body becomes reduced to a mere flattened thread, and the constrictions are only represented by faintly marked ‘striæ.’ A short, smooth, faintly-striated neck is succeeded by the head. This is about the size of a pin’s head, and is armed with a double circlet of minute spines, and carries in addition four minute suckers. The ‘proglottides’ in the hinder part of the specimen are of a whitish colour, opaque, and three or four times as long as they are broad. The head, ‘nurse,’ or ‘scolex,’ is asexual. The segments succeeding are sexual, but immature: that is, their generative organs and their products are but imperfectly developed. The segments which succeed and make up the bulk of the organism, are sexually mature, hermaphrodite ‘proglottides.’ The new segments are produced by a process of gemmation from the head: and the segments thus produced are intercalated between the head and the segments already formed. The apertures of the efferent ducts of the male and female organs are situated on small cup-shaped eminences placed on the lateral margins of the ‘proglottides,’ the side alternating in adjoining segments. The specimen is eight feet in length, and was taken from a human intestine.

SUB-KINGDOM ARTHROPODA.

Division TRACHEATA.

Class MYRIAPODA.

29. A Centipede, *Scolopendra*, a Chilopodous Myriapod.

The segments of the exo-skeleton differ from those of Insecta and Crustacea in not being definitely marked out into cephalo-thoracic, abdominal, and post-abdominal regions. There are twenty-three post-cephalic segments, to each of which a single pair of appendages is attached. Any one of these appendages consists of a dorsal and ventral arc, united together laterally by the soft integument from which the appendages take origin. On either side of the mesial line of the ventral arc are two very faint lines, which mark off a median portion, or 'sternum,' from two lateral elements, or 'episterna.' These faint lines are best seen in the sterna of the anterior segments. Imbedded in the lateral integument are one or two chitinous plates or tubercles, representing the rudimentary 'epimera' or lateral elements of a typical 'somite.' Each appendage is many-jointed. The proximal joint is the 'coxa,' and the distal one carries a sharp curved claw. Each segment is to be regarded as being composed of two sub-segments, which at first are equally developed, but as development proceeds, one of them increases in size more rapidly than the other; and moreover develops a pair of appendages which the other does not. This latter, the anterior one, becomes abortive, though in the Chilognatha it develops equally with the posterior one, with which it eventually coalesces, and moreover carries a pair of appendages. Nevertheless in this specimen the remains of the anterior sub-segment are to be seen. Immediately in front of each sternal piece, and on each side of the mesial line, are three small angular chitinous plates. The pair nearest the mesial line are the two uncoalesced halves of the sternum of the aborted anterior sub-segment. The pair external to these are the episterna, and the most external and smaller pair are the epimeral plates. Epimeral plates are also situated, one pair on the hinder edges of the sternum, and two small plates just in front of the coxa, imbedded in the lateral integument. The two sub-segments,

one of which in Chilopoda is abortive, are equally developed in Chilognatha, and each carries a distinct appendage. The terga, however, are confluent, so that the double nature of each body-segment is only evident when the ventral surface is examined, where two pairs of appendages will be found attached to each segment. The segmental appendages are composed of six joints, coxa, tibia, femur, metatarsus, tarsus, and claw. The anterior appendages are the smallest, but they gradually increase in size posteriorly; and the hindmost pairs are directed backwards and not outwards, as is the case with the anterior ones. The last pair is very much larger than any of the rest, curves inwards, and by mutual apposition forms an efficient clasping apparatus. The anus opens on the under surface of the last segment. In several of the segments a pair of small, crescentic, laterally placed slits surrounded by prominent chitinous lips is to be seen a little inferior to the dorsal arc: these are the 'stigmata.'

The head is probably formed by the fusion of at least five of the most anterior segments. The superior surface is formed by a broad, shield-like plate, quite smooth, and devoid of sutures, with the exception of a faint median groove. To the anterior margin of this plate is attached a pair of multiarticulate antennæ consisting of not fewer than seven segments. In this respect the Order Chilopoda, to which this species belongs, differs from the Order Chilognatha, in which the antennæ are composed of fewer than seven segments. At the base of the antennæ, and external to them, are four sessile 'oculi.' On the inferior surface of the head, in addition to the antennæ already described, the following appendages are to be seen. First, a pair of mandibles, toothed on their inner edges, and carrying small rudimentary palpi on their outer edge. Secondly, two pairs of maxillæ, of which the anterior pair is jointed, palpiform, and forms the external elements of the labium, while the posterior pair is composed of two small conical pieces applied together in the mesial line, but not united, and constituting the mesial part of the 'labium.'

The appendages hitherto enumerated belong to the head, the remainder, proceeding backwards, to regions which in the Lobster are termed thoracic, abdominal and post-abdominal.

Succeeding the maxillæ is a pair of jointed appendages armed with a small claw. The next are the large cheliform poison-fangs. These are armed with claws, and the extremity of each claw is perforated by the aperture of a small duct leading from a poison-gland. The basal joints are confluent, so that by this union a second labium is formed posterior to the first. The two last-mentioned appendages may be taken as corresponding to two of the three pairs of maxillipeds in the Lobster, and to the two anterior pairs of legs in Insects.

The segment to which the large 'chelæ' are attached is larger than the segment immediately succeeding it.

30. A Centipede (*Scolopendra*), dissected to shew the digestive and female generative systems.

The animal has been laid open along its dorsal surface, so as to expose the alimentary canal, which is simple, and passes in almost a straight course from mouth to anus, the terminal portion of the intestine alone indicating any approach to a convolution. It is distended with food. The stomach, which occupies the central third of the body, is the widest part of the canal: it communicates anteriorly with the mouth by a narrow œsophagus, and posteriorly is marked off from the intestine by a slight constriction. The intestine, at its junction with the stomach, though of less calibre than the latter, is yet of considerable size; but diminishes as it proceeds, and after describing a sigmoid flexure terminates in an anus.

The ovary is a long and simple tube, attached above to about the middle of the stomach, and gradually diminishing in size till it reaches its vaginal outlet. The ovi-sacs may be seen through its semi-transparent wall throughout its whole extent.

The white cæca pinned on either side of the lower part of the preparation are the 'accessory glands.'

31. A Centipede (*Scolopendra*), dissected to shew the male generative organs.

The animal has been laid open along its dorsal surface, and the digestive tract removed. The testes consist of a series of pairs of fusiform vesicles, each vesicle being applied along its whole length to its fellow, from which however it can be easily separated. They occupy a considerable portion of the abdomen,

and extend forwards as far as the anterior fifth of the body. Each end of each vesicle communicates by a duct with a minute common longitudinal canal. There would thus be twenty pairs of ducts passing from the testes to the central canal. Posteriorly this canal becomes of considerable size, and is disposed in spiral coils before it reaches its external opening. Beneath it, and in the last two segments of the body, are seen the 'accessory glands.' The long dorsal vessel has been pinned on the left of the specimen.

32. A Centipede (*Scolopendra*), dissected to shew its nervous system.

It has been laid open along its ventral surface. A black bristle indicates the position of the œsophagus.

The nerve-system consists of a double cord placed medially in the body. The cephalic ganglion supplies the antennæ and eyes. Posteriorly to this the cords divide to admit of the passage of the œsophagus. Subsequently there are ganglia at regular intervals, corresponding in number to the segments of the body. From each nerves are given off to supply the legs. The first ganglion is oval in form, the others are round.

Class INSECTA.

1. Order *Coleoptera*.

33. A Goliath Beetle (*Cetonia cæcicus*), disarticulated and mounted on wires to shew its segments and their appendages.

34. Digestive system of an Oil-Beetle (*Meloe proscarabæus*).

Nearly the whole of the exoskeleton, together with the other organs, has been removed.

The œsophagus, which is very short, leads into a large elongated oval stomach, occupying the greater part of the abdominal cavity. On leaving the stomach the intestine becomes suddenly extremely narrow, and is slightly convoluted. Its terminal segment is dilated into a rectum.

Near the posterior end of the stomach the hepatic vessels, a number of thread-like cæcal tubules, may be seen.

35. A Stag Beetle (*Lucanus cervus*), dissected to shew its nervous system.

The supra-oesophageal mass consists, as usual, of two cephalic ganglia, from which are derived the nerves which supply the eyes. These ganglia are connected with the ventral cord by a double nervous cord encircling the oesophagus. Three large ganglia supply the thorax. The first pair of abdominal ganglia are very large, but the remaining six, with the exception of the last, are small. This ganglion sends off a bundle of nerves which supply the segments posterior to it.

36. A Common Cockchafer (*Melolontha vulgaris*), female, dissected to shew its generative and digestive systems, which have been arranged on the left and right sides of the tablet respectively.

Generative organs. Each ovary is composed of a single fasciculus of egg-bearing cæca. These cæca, or tubules, at their commencement are small and slender, but as they descend they dilate and assume a granular appearance, eventually fusing to form the oviduct. The cæca composing the right ovary have been partially separated. The two oviducts combine to form the common vaginal canal, which opens externally. On the right of the 'vagina,' and opening into it by a short duct, is the pyriform 'spermatheca:' on the same side and with the same termination is the glandular accessory cæcum.

Digestive system. This is seen to be a canal of great length in proportion to the size of the insect, of tolerably uniform diameter throughout, and much convoluted.

2. Order *Orthoptera*.

37. A Common Cock-Roach (*Periplaneta orientalis*), dissected to shew its digestive system.

The oesophagus opens into a large pear-shaped crop. On either side of the oesophagus may be seen the two-lobed salivary gland. Posterior to the crop, and separated from it by a slight constriction, is the muscular gizzard. The true chylic stomach which succeeds receives at its anterior extremity a whorl of eight cæca; and a cluster of much longer and more slender tubules is similarly arranged round its lower end. The first eight cæca are regarded as analogous to the liver of higher animals, while the tubules are probably renal. The 'colon,'

which is connected with the stomach by a short segment of intestine, of small calibre, is large, and bent upon itself. The terminal portion, or 'rectum,' is marked by longitudinal muscular bands.

Rolleston, p. 86 and p. 199.

38. A Locust (*Locusta*), dissected to shew the female generative system and the digestive system.

The insect has been opened on its dorsal surface, most of which has been removed.

The ovigerous cæca form two bundles, one on each side. The tubules composing them unite to form an oviduct. The two oviducts join in the median line, and form the short and wide vaginal canal which opens externally at the base of the large ovipositor. Communicating with this canal by a short duct is the 'spermatheca' (pinned to the left side), and also a long thread-like accessory cæcum (pinned to the right side).

The digestive system of the insect has been arranged on the right of the preparation. It closely resembles that of the Cock-Roach; but the hepatic cæca are unusually large, and the renal tubes are small.

39. A Locust, dissected to shew the male generative organs.

The insect has been prepared in the same way as the last. The testes are two large oval compressed bundles of sperm-secreting tubules, one on each side of the body. Between them is a mass of accessory glands, which have taken the form of minute, much convoluted tubes.

3. Order *Diptera*.

40. A Common Flesh-Fly (*Sarcophaga carnaria*), dissected to shew its digestive and female generative organs.

The insect has been laid open along its dorsal surface.

The alimentary canal has been pinned to the right. It consists of a long tube, in which scarcely any difference between stomach and intestine can be observed. The ovaries are seen to be clusters of white cæca, arranged spirally. In this species the female is viviparous.

Packard on Insects, p. 408.

4. Order *Lepidoptera*.

41. Larva of Privet Hawk Moth (*Sphinx ligustri*).

Bristles have been inserted into the spiracles of the right side.

Rolleston, p. 73.

42. Pupa of the same.

43. Larva of the same, dissected to shew the nervous system.

It has been laid open on its superior surface, and the digestive system removed, with the exception of a piece of the œsophagus, which has been pinned back, in front of the head.

The nerve-system consists of twelve ganglia, arranged in a linear series along the ventral surface of the larva. The first pair of these are the cerebral ganglia, situated immediately above the œsophagus. They are connected by the usual nerve-collar, embracing the œsophagus, with the first post-oral ganglion. The terminal ganglion of the series is composed of two ganglia, closely opposed to each other, from which a leash of nerves is given off to the terminal segments of the body.

Rolleston, p. 82.

44. Larva of Goat Moth (*Cossus ligniperda*), dissected to shew the internal organs.

The integument has been divided down the medio-dorsal line and pinned out on each side.

Occupying the greater part of the body-cavity and investing the contained viscera is a mass of lobulated adipose tissue, known as the 'fat body' or 'rete,' which disappears almost completely in the 'imago' of those insects whose metamorphosis is complete.

In the median line is the digestive tract, which passes, without forming any convolutions, direct from mouth to anus. The walls of the œsophagus are thin, and thus readily distinguish it from the thick-walled, opaque, corrugated stomach.

A little way below the pylorus the intestine receives the common ducts of the tubular renal organs, the so-called 'Malpighian vessels.' The common duct on each side bifurcates, and

the branches so formed extend anteriorly and form two loops on either side of the stomach; the trunk forming the outermost of the two loops in descending bifurcates again on each side. There are thus three trunks on each side of the stomach, forming an intricate plexus round the pylorus, and finally terminating blindly. The long, convoluted, and much larger tubes seen on each side of the digestive tract are the silk-glands; they commence blindly in the posterior part of the body, and terminate in a common duct opening on the modified 'labium' or 'spinneret.'

On the left side of the stomach is seen a thin-walled, transparent sac, the posterior end of which is connected with a much-convoluted tubular gland, which bends forwards, and in the preparation lies on the outer side and in advance of the sac. This sac is connected anteriorly with the mouth by a wide duct. The tubular gland, bladder, and duct correspond with the smaller salivary glands of the imago.

In the intervals of the fat body fasciculi of tracheæ are seen diverging to distribute themselves upon the viscera.

The following points in which the viscera of the larval form differ from those of the imago may be noticed in this preparation:—the presence of the 'fat body,' the larger calibre and lesser length of the digestive canal, and the absence of convolutions, and the presence and large size of the silk-glands.

Rolleston, p. 79.

Class ARACHNIDA.

45. A Scorpion (*Scorpio afer*), to shew the exo-skeleton.

The exo-skeleton presents a well-marked division into cephalo-thoracic, abdominal, and post-abdominal regions, and with the exception that the abdominal segments are here distinct from each other, these divisional areas approximate closely to the same parts in the lobster. The cephalo-thorax is a broad and but slightly convex plate marked by a median antero-posterior groove, and with a deeply emarginate anterior border. At the commencement of the posterior fourth of its extent there is a shallow transverse groove curving forwards on either side,

dividing the cephalo-thorax into a large anterior and a much smaller posterior division, and bearing some resemblance to the cephalic suture, to be subsequently described, in the lobster. In the centre of the cephalo-thorax, and situated upon a little eminence, is the medio-dorsal pair of 'oculi,' one on each side of the median groove. There are also three additional 'oculi' at each antero-external angle, on each side of the anterior median notch.

The segmental appendages, of which there are six pairs, namely, counting from before backwards, antennæ, mandibles, and four pairs of ambulatory legs, are entirely confined to the cephalo-thorax. Functional antennæ are absent: but the homologues of the antennæ of Insecta are the large and strong cheliform appendages situated in front of the mouth, and so modified from their ordinary structure as to become efficient prehensile organs. The first post-oral pair, the mandibles, are in this species remarkable for the extraordinarily disproportionate development of the mandible proper and its 'palpus.' While the former is a small, internally flattened, slightly serrated plate, the latter, absent in Insects and with the exception of a rudiment, in Myriapods also, is enormously elongated, and carries huge 'chelæ' at its distal extremity. The two succeeding pairs of appendages correspond to the two pairs of 'maxillæ' in the lobster. In the first pair the 'maxillæ' themselves are small, somewhat conical, curved inwards, and form the external lateral elements of the 'labium.' The 'palpi,' on the contrary, are greatly elongated, six-jointed appendages, with two small claws attached to their distal segment. In the second pair a similar arrangement exists. The maxillæ are two triangular processes, applied by their straight internal edges to each other in the median line, and so forming the mesial elements of the 'labium.' The palpi resemble those of the first pair of maxillæ. The next two pairs of feet, representing the two pairs of 'maxillipeds' in the Lobster, and the two anterior pairs of legs in Insects, resemble the maxillary palpi just described, with the exception that they are longer, and have much larger basal segments.

The next segment, which in the Crustacea would carry the third pair of 'maxillipeds,' is, as in the Chilognathous Myriapods, the genital segment. In the centre of the sternal surface of this segment is the opening of the efferent duct of the reproductive

organs. An opercular process, attached in front, and free posteriorly, hangs like a flap over this aperture. This may probably be regarded as the sternal element of this segment. The sternal elements of the two maxillary segments of the 'cephalo-thorax' are aborted, but in the two succeeding segments the sterna, though small, are still sufficiently obvious.

The segments and appendages hitherto described belong to the cephalo-thorax: the five succeeding ones to the abdominal region. The abdominal segments are distinct from each other, and, with the doubtful exception of the first, carry no appendages. Each consists of a broad tergal and sternal arc, united laterally by non-chitinous membrane. The sternal element of each segment is divided into a median division, 'sternum,' and two lateral divisions, 'episterna,' by two well-marked grooves, one on each side of the median line. On each 'episternum,' with the exception of that of the first pair, is a small oblique slit, the external opening of the pulmonary sacs. In addition to the absence of pulmonary apertures, the first abdominal segment further differs from the rest in the small size of the sternum, and in the presence of a pair of pectiniform appendages. Each appendage presents indications of being composed of several segments, and carries on its hinder border a row of close-set filaments.

The six segments which succeed the abdomen are the post-abdominal segments. With the exception of the first, which resembles the abdominal segments in its general shape, each segment is more or less cylindriciform, completely chitinized, convex above and concave below, and marked with longitudinal serrated ridges. Attached to the last post-abdominal segment is a conical, somewhat tumid segment, which carries a strong curved spine upon its distal extremity, and may represent the 'telson' of the Crustacea. On the ventral surface, and in the interspace of the last two segments, is the anus.

46. A Scorpion (*Scorpio*), dissected to shew its digestive tract.

The alimentary canal is seen to be a straight tube, destitute of convolutions, and passing directly from mouth to anus. The liver is very large, occupying four-fifths of the thoracic cavity, and from it numerous ducts pass to the stomach. The anus is situated on the under surface of the last segment

but one. The nervous system is seen at intervals underlying the digestive tract in the abdomen.

47. A gravid female Scorpion, dissected to shew the position of the embryos in an ovi-viparous Arachnid.

The dorsal surface of the animal has been removed. The embryos, differing from the parent only in size, are seen occupying the ovarian tubes, which overlie the liver or are even embedded in its substance.

48. A Spider, dissected to shew its nervous system.

The whole of the post-oral ganglia are seen to be concentrated into a single stellate mass, occupying the larger portion of the ventral surface of the cephalo-thorax. From this mass, which may be taken to represent the confluent thoracic and abdominal ganglia of Insecta, radiate, on each side, five principal nerves to supply the legs and maxillary palpi. Several nerves pass posteriorly from its hinder edge to supply the abdomen. The cephalic ganglia (two in number) are situated on the anterior edge of this stellate mass, between which and the former the œsophagus passes. From this pair of ganglia nerves are seen passing forwards to supply the eyes and mandibles.

Division ARTHROPODA BRANCHIATA.

Class CRUSTACEA.

1. Order *Decapoda*.

49. A Lobster (*Homarus vulgaris*), disarticulated, so as to shew its segmentation.

The different segments with their appendages have been as far as possible separated from each other, and serially arranged in their natural order. The segments have been numbered from the first, or ophthalmic segment, backwards. The carapace has been removed, and fixed on the right of the sub-abdominal segments.

The skeleton consists of a broad, shield-like carapace, 'cephalo-thorax,' and of the six segments, or 'somites,' which enter into the composition of the post-abdomen. Attached to the ventral surface of these different segments are the variously

modified appendages, to the number of twenty pairs. These are named in order from the first segment as follows. In front of the mouth are a pair of eyes, a pair of antennules, and a pair of antennæ. Behind the mouth are a pair of mandibles, two pairs of maxillæ, three pairs of maxillipeds, five pairs of ambulatory legs, and six pairs of swimmerets: while to the last sub-abdominal segment is attached a median azygos element, the 'telson.'

A sub-abdominal 'somite,' which may be regarded as a typical one, is made up as follows. It is nearly semicircular in vertical section, the dorsal wall or 'tergum' being very convex, and the ventral wall or 'sternum' nearly straight or but slightly convex. At the infero-lateral angle both the 'tergum' and the 'sternum' are produced downwards and outwards into a lobate hollow process, termed the 'pleuron.' In the angle between the 'sternum' and the inner surface of the 'pleuron,' on each side, is an articular cavity, which receives the proximal extremity of the appendage attached to this 'somite.' A transverse groove marks off a crescentic area on the anterior surface of the 'tergum,' which is overlapped by the hinder part of the 'somite' in front of it. This area is the tergal facet. A similar flattened surface exists on the anterior portion of each 'pleuron.' This is the pleural facet. Each appendage consists of a basal attached portion (divided longitudinally into two segments), the 'protopodite.' Each 'protopodite' (sometimes called 'basipodite') carries at its distal extremity an external and internal filament, to the former of which the term 'endopodite,' and to the latter 'exopodite,' has been applied.

This description will in the main apply to the four central sub-abdominal segments, but the first and the sixth of these segments have their appendages somewhat modified for special purposes. In the sixth sub-abdominal segment, the 'exopodite' and 'endopodite' are enormously expanded into flat setose plates, and the exopodite is further divided by a transverse joint.

The 'telson' is a flattened conical setose plate, attached to the hinder margin of this 'somite,' and forms, with the expanded 'exopodite' and 'endopodite,' a powerful swimming tail.

In the first sub-abdominal 'somite' each appendage consists of

two elements, the distal one of which is grooved on its inner face, and is capable, by being applied to its fellow, of forming a short cylindrical tube which probably conveys the male semen to the vulva of the female.

The 'cephalo-thorax' is formed by the fusion of the terga of the cephalic, the thoracic, and the abdominal segments. The cephalo-thoracic pleura are well developed, and extend so far downwards as to hide the bases of the appendages.

The sternal portions of the pleura are membranous and are reflected upwards to the roof of the cephalo-thorax, and then, after being deflected, become continuous with the partially calcified 'epimera.' In this specimen this membranous portion has been torn through by the separation of the tergal half of the cephalo-thorax from the coalescent sterna, which form the ventral surface of this part of the animal.

On the dorsal surface the 'cephalo-thorax' does not present any indications of division into distinct terga, corresponding to the sterna and appendages of the ventral surface, but nevertheless contains areas more or less distinctly marked out. A groove commences at the base of the antennæ, and, after passing directly backwards for a short distance, curves upwards and fuses with the corresponding curve of the opposite side at about the middle of the dorsum. The part in front of this suture has been called the 'cephalo-stegite,' and the portion behind the 'homo-stegite.' The 'homo-stegite' is further divided by two lateral linear depressions, one on each side of the median line, called the 'branco-cardiac sutures,' and by a very faint transverse ridge, which is to be seen a little way behind the cervical suture, into a quadrangular area which corresponds externally with the position of the heart internally. In this specimen this area is tolerably well indicated by a boundary line of irregularly shaped white spots. The cervical suture indicates the line of separation between the cephalic and thoracic segments. The area between the cervical suture in front, and the faintly marked ridge behind, represents the confluent terga of the three thoracic segments, while that part of the 'homo-stegite' posterior to this corresponds with the terga of the five abdominal segments. The lateral areas which lie on the side of these median divisions represent the conjoined pleura of the

eight thoracic and abdominal 'somites:' and from the fact that these areas cover in and protect the branchial chamber, they have been called 'branchio-stegites.' Anteriorly the 'cephalo-stegite' is produced into a long serrated rostrum. The sterna corresponding to the 'cephalo-thorax' are, like the terga, confluent, owing to the calcification of the membrane connecting the sterna and epimera of adjacent somites. The sterna which are in relation to the three posterior pairs of ambulatory legs are tolerably wide, but the sterna anterior to these become so narrow as to allow the bases of the appendages to meet in the median line. The calcified inter-epimeral, and inter-sternal membranes send inwards calcified processes, which, after dividing, reunite with each other in such a manner as to divide the floor of the cephalo-thorax into a series of inter-communicating chambers. A median canal is also formed by these processes, called the 'sternal canal.' It contains the thoracic portion of the ventral nervous cord, and is indicated by the insertion into it of a red glass rod.

The abdominal segments which have been already described are tolerably simple, and only slightly modified: but those appended to the 'cephalo-thorax' exhibit much greater diversity of form, and subserve a variety of functions.

Taking the segments in order, from behind forwards, and beginning with the first sub-abdominal or fifteenth body-segment, which has been already described, we find the following cephalo-thoracic appendages.

XIVth pair. (Ambulatory legs.)

Each appendage consists of a right 'protopodite' and a long segmented 'endopodite,' the 'exopodite' being absent. The joints of the appendage are named by Milne-Edwards as follows: 'coxopodite' ['protopodite'], which is attached to the 'somite,' and pierced by the external aperture of the 'vas deferens,' into which a black arrow has been inserted:—'basi-podite,' 'ischiopodite,' 'meropodite,' 'carpopodite,' 'propodite,' and 'dactylopodite.'

XIIIth, XIIth, XIth, Xth pairs, (ambulatory legs,)

are similar to the fourteenth pair, but each possesses a long curved membranous appendage, which ascends from the 'coxopodite,' and in the natural condition penetrated into the branchial chamber. This is the 'epipodite.' Attached to each 'coxopodite' may also be seen a lamellar gill.

In the twelfth, eleventh, and tenth segments the 'propodite' is prolonged outwards into a process which bites against the 'dactylopodites,' and so forms the chelæ, which in the tenth pair are of great size.

IXth, VIIIth, VIIth pairs. (Maxillipeds.)

Each maxilliped is composed of three divisions, articulated to a 'coxopodite.' In the first maxilliped [No. IX.], the outermost of these three divisions, the 'epipodite' is a curved lanceolate lamina. A little in front of it is a gill, likewise attached to the 'coxopodite.' The middle division is a many-jointed palpi-form filament, which corresponds to the 'exopodite.' The innermost element is the seven-jointed 'endopodite.' The second pair of maxillipeds [No. VIII.] much resembles the first, but the 'endopodite' is smaller and somewhat flattened, resembling the foliaceous 'endopodite' of the first pair. In the third pair [No. VII.] the 'endopodite' has been split into an inner flattened plate-like lamella and an outer flattened filament. The 'exopodite' is small, but the 'epipodite' is large, and in the specimen directed backwards.

VIth, Vth pairs. (Maxillæ.)

In the first pair [No. VI.] the 'endopodite' has been split into five divisions, and is in consequence somewhat foliaceous. The 'exopodite' is a flattened filament. The 'epipodite' is curved inwards. The 'endopodite' of the second pair [No. V.] differs from that of the first in having only three divisions. The 'exopodite' and 'epipodite' are absent.

IVth pair. (Mandibles.)

Each mandible carries a jointed palp, which corresponds to the terminal joints of the mandibular 'endopodite' (Huxley). The calcified rod projecting outwards is the tendon of the adductor muscle of the mandibles.

III^d pair. (Antennæ.)

The antennæ are composed of a short basal 'protopodite,' which carries a long multi-articulated 'endopodite' and a short flattened scale-like 'exopodite.' The black arrow is inserted into the orifice of the duct of the 'green gland.'

II^d pair. (Antennules.)

A three-jointed 'protopodite' is succeeded by two long multi-articulated filaments, the 'exopodite' and 'endopodite.' An aperture protected by membrane in the proximal joint, and towards which the arrow points, is the auditory organ.

Ist pair. (Ophthalmic peduncles.)

The eye-peduncles are composed of two segments, the distal one of which carries the eye itself.

On the left of the specimen is a portion of the skeleton of a female, with the ventral surface turned upwards, so as to shew the openings of the oviducts on the 'protopodite' of the third pair of ambulatory legs. The two orifices are indicated by two black arrows. The first pair of post-abdominal appendages differs from those of the male in being simple unmodified filaments.

The structure of the stomach is illustrated by the two specimens marked *A* and *B*. In *B* the greater portion of the cephalothorax has been removed so as to expose the cavity of the thorax, with the stomach in its natural position. This viscus occupies nearly the whole of that part of the cavity which lies in front of the cephalic suture, and its membranous walls are supported and strengthened by a some-

what complicated arrangement of calcified bars and plates. It is divided transversely by a rather deep fissure into an anterior layer or cardiac portion, and a posterior smaller or pyloric division. The structure of the calcified gastric skeleton is as follows.

Crossing the stomach transversely, at the hinder border of the cardiac division, is a broad and slightly arcuated plate, the 'cardiac ossicle.' Attached to each extremity of this 'cardiac ossicle' is a curved triangular 'ptero-cardiac ossicle.' On each side a large elongated 'supero-lateral ossicle,' wider posteriorly than anteriorly, is connected with the lower end of the 'ptero-cardiac ossicle,' and passing upwards and backwards articulates by the anterior part of its hinder extremity with a small transverse rod, which forms the medial part of the anterior boundary of the pyloric portions, and is termed the 'pyloric ossicle.' The posterior part of the superior extremity of the 'supero-lateral ossicle' is connected with a triangularly shaped imperfectly calcified plate, the 'meso-pyloric ossicle.' Anteriorly this ossicle becomes continuous with the hinder edge of the pyloric ossicle. "These pieces, it will be observed, form a sort of hexagonal frame whose anterior and lateral angles are formed by moveable joints, while the posterior angles are united by an elastic meso-pyloric plate." From the middle of the cardiac piece a strong calcified process passes downwards and backwards underneath the 'meso-pyloric ossicle,' terminating in a broad thickened extremity, which may be termed the post-cardiac process, and to which is attached internally a broad and flat chitinous plate. This plate has its posterolateral and internal angles slightly nodular. These nodules represent the cardiac teeth of the Common Cray-fish. A very narrow and uncalcified band connects the end of the post-cardiac process with a broad calcified process, the 'uro-cardiac ossicle.' This ossicle carries on its inferior surface a strong conical tooth, which projects into the cavity of the pyloric divisions of the stomach. Finally, connected in a similar manner with the end of the uro-cardiac piece, is a broad somewhat triangularly shaped ossicle, convex above and concave beneath, the 'pre-pyloric ossicle.' This passes upwards and forwards, and connects the uro-cardiac with the pyloric ossicle. In *A* the greater

part of the meso-pyloric ossicle has been removed, so as better to exhibit the 'pre-pyloric ossicle' which lies beneath it. The post-cardiac process and the uro-cardiac ossicle, with its teeth, are also best seen in this specimen. The supero-lateral ossicle sends inwards from its inferior border a strong calcified process, which carries on its internal face the large lateral gastric tooth. On the infero-lateral parietes of the stomach there are several rod-like ossicles. To the inferior extremity of one of these, the 'lateral cardiac,' stretching downwards and backwards from the 'ptero-cardiac,' is attached a small ossicle, which internally bears a bifid tooth. The under surface of the stomach is strengthened by two nearly parallel rods, which extend from the pylorus forwards to the cardiac division, where they become divergent and more or less membranous. Their posterior extremities, also slightly divergent, are somewhat thickened, and each articulates with two rods, one of which is connected with the ossicle just described as bearing the bifid tooth, while the other joins the 'supero-lateral ossicle.' There are thus five gastric teeth, one median, and two lateral on each side.

In *A*, on looking into the gastric cavity and towards the pylorus, we can see, situated beneath the median gastric tooth, a bilobed valvular process; and on either side, immediately beneath the great lateral teeth, is a similar cushion-like process. These processes are beset with hairs, and by their mutual apposition form an efficient sieve to prevent the undigested food from passing into the intestine.

For an account of the structure and functions of the stomach of *Astacus fluviatilis*, which differ but little from that of the specimen described above, see Huxley, "*Medical Times and Gazette*," 1857, Vol. xiv. p. 255: also Rolleston, p. 90.

50. The cephalo-thorax and the first fourteen segments of a Langouste (*Palinurus vulgaris*.)

The same numbers have been affixed to the same parts as in the preceding specimen: and the description is in the main equally applicable.

51. A Lobster (*Homarus vulgaris*), dissected to shew its nervous system, which is of the type termed 'homogangliate:' that is, it consists of a longitudinal series of ganglia

of various sizes, united by nerve-cords, termed 'commissural cords,' and occupies the ventral surface of the body.

The total number of ganglia is thirteen. Of these, the first is the præ-oral or cephalic ganglion, from which are derived the nerves supplied to the eyes, antennæ, and antennules. Black bristles have been placed under these nerves. The remaining and post-oral twelve are equally divided between the thoracic and abdominal segments. The first post-oral ganglion (formed by the fusion of six embryonic ganglia) is of large size, and supplies six pairs of appendages, viz. the mandibles, two pairs of maxillæ, and three pairs of 'maxillipeds.' The five succeeding ganglia, smaller in size, and comparatively close together, supply the five pairs of ambulatory legs. The abdominal ganglia, still smaller in size and separated by greater intervals, supply the powerful abdominal muscles, and the segmental appendages. The last of these ganglia is of larger size than the rest, and gives off a leash of nerves to supply not only its own segment and appendages, but the 'telson.' The longitudinal commissural cord is double in the segment anterior to the abdomen, but in the latter region it is reduced by fusion to a single thread. Between some of the abdominal ganglia may be seen the 'nervi transversi.' These are small nerves given off from the longitudinal cord, at a point about midway between the ganglia.

52. A Cray-Fish (*Astacus fluviatilis*), dissected to shew the male generative system.

The whole of the heart and the greater part of the liver have been removed. The testes are situated immediately behind the posterior end of the stomach, and consist of two lateral halves, which posteriorly fuse into a long thin mass, situated in the median line. On either side the 'vasa deferentia' quit the gland at the junction of its three apparent lobes, and after describing a mass of convolutions on either side of the hinder part of the thoracic cavity, terminate by a small prominent orifice on the basal joint of the last pair of thoracic legs. A slip of blue paper has been placed under the terminal segment of each 'vas deferens.'

Rolleston, p. 106, sq.

2. Order *Stomapoda*.

53. A Stomapod (*Squilla mantis*), dissected to shew the same parts.

The double 'commissural cord' connecting the præ-oral with the post-oral ganglion is remarkably well seen. The remaining thoracic ganglia supply the thoracic legs. The abdomen contains six ganglia, each of which gives off two or more nerves on each side to supply the muscles of its segment. The longitudinal commissural cord connecting the different ganglia is more or less distinctly double throughout.

Sub-Class CIRRIPIEDIA.

54. A Barnacle (*Lepas anatifera*).

The body of the animal consists of 'capitulum' and 'peduncle.' The peduncle is soft and fleshy, and invested externally with a tough rugose skin, which by its distal, clavate extremity serves to attach the animal. The 'capitulum' is somewhat triangular, consisting of two lateral valves, strengthened externally by several calcified plates, and encloses the major part of the body of the organism. The calcified pieces composing the valves are four in number, two in each valve. The superior or distal plates are called 'terga,' and the lower or proximal plates, also the largest, are called 'scuta.' The two valves are connected with each other along one side, through the intermediation of a mesial curved piece, the 'carina.' Within those valves the body of the animal is to be seen:—its dorsal surface in relation with the inner surface of the 'carina.' The body consists of a 'prosoma' which includes the cephalic segments—of six thoracic segments—and a tubular extension from the hinder edge of the last thoracic segment, which functionally is a penis, but morphologically may be the aborted representative of the abdomen, or post-abdomen, of other crustacea. The mouth is bounded anteriorly by a tumid, transversely extended labrum. Behind this we find, first, a pair of mandibles, each armed with a setose palp, and two pairs of maxillæ. Six pairs of appendages are attached to the thorax, each consisting of a basal 'protopodite' terminating distally in a long multiarticulated 'exopodite' and 'endopodite.' That part of the body which lies in front of the

mouth, together with the whole of the peduncle, is the result of an extraordinary development of the pre-oral 'somites'—and the only appendage to be found in relation with these segments is a pair of sucker-like antennæ, to be seen by careful examination of the attached surface of the peduncle.

55. A Barnacle (*Lepas fascicularis*), dissected to shew its digestive system.

One half of the calcareous carapace has been removed, together with the investing membrane and the left testicular mass. A black bristle has been inserted through the mouth and œsophagus into the stomach, and a white one into the anus. The mouth, encircled by its mandibles and maxillæ, leads by a short œsophagus into a large oval or sub-globular stomach. The intestine is a simple tube of considerable size at its commencement. It bends upon the stomach, and proceeds along the dorsal aspect of the animal, gradually diminishing in size, to the anus, which is situated at the base of the penis.

Upon the stomach is a mass of dark-coloured glandular cæca, probably hepatic in function,—(a portion has been removed with the wall of the stomach,)—which communicate by several large orifices with its cavity. The cuticular lining of the œsophagus is continued for some distance into the stomach, and terminates therein by a well-defined serrated edge.

56. The same, dissected, to shew its hermaphrodite generative system.

In the upper of the two specimens on the tablet the carapace and investing membrane have been removed, and the organs displayed *in situ*, on the left side.

The testis, of irregular shape, is composed of minute branching cæca, and not only overlies the pyloric end of the stomach, but also gives off prolongations which enter the pedicels, and the basal segments of the rami of the cirri.

In the lower specimen a great part of the testis has been removed, so as to shew the 'vas deferens,' or dilated portion of the efferent duct of the testis, ascending parallel to the terminal portion of the intestine. It gradually diminishes in size to the base of the penis, where it joins the corresponding duct of the opposite side to form the common canal, which, after traversing

the penis, opens at its end. A black bristle has been placed under the two ducts.

The ovaria are disposed in the peduncle and in the body. In the peduncle they may be seen as a tangled mass of branching tubes, filled with granular matter and immature ova. From the peduncle two main ducts (not seen in this specimen), one on each side, enter the body at the anterior corner of the rostrum, whence they curve round towards the base of the first pair of cirri. There they form two glandular masses resting on the upper edge of the stomach, of an orange colour, and generally subdivided near the mouth. These masses were thought by Cuvier to be salivary glands, but are now known to be true ovaries. No proper oviduct has yet been discovered. In the uppermost of the two specimens, between the body of the animal and the peduncle may be seen one of the ovigerous lamellæ in which the ova exist previous to their expulsion.

Darwin, "Monograph on the Cirripedia," p. 55, sq.

SUB-KINGDOM MOLLUSCA.

Division I. MOLLUSCOIDEA.

Class TUNICATA.

In the following preparations of Ascidians a *blue* rod has been placed in the inhalent, a *red* rod in the exhalent, aperture.

Rolleston, p. 66.

57. A simple Ascidian (*Cynthia microcosmus*.)

One half of the test, which in this species is of a leathery consistency, has been removed, to shew the animal suspended within it. The mantle is attached to the test at the two apertures, and is very muscular. Fibres may be seen passing in a longitudinal, oblique and transverse direction across its surface. The two apertures are regarded by some as homologous with the inhalent and exhalent siphons of the Siphonate Lamelli-branchiates; and by others, as the mouth and anus.

58. One-half of the test, mantle, and branchial sac has been removed, so as to expose the cavity of the latter. The branchial

sac is of considerable size, with its walls produced into a series of parallel longitudinal folds, radiating from the inhalent orifice to the mouth. A black bristle has been introduced into the aperture leading from the branchial sac to the stomach—an aperture which has been regarded by some as the mouth, while others maintain that the mouth is better represented by the inhalent aperture.

59. To shew the nervous system.

The single nerve-ganglion is situated in the substance of the mantle on one side of the inhalent aperture, between it and the exhalent aperture, and is indicated in the specimen by the insertion of a black bristle behind it. The ganglion is fusiform in shape, and from its anterior and posterior ends nerves are seen passing towards the inhalent and exhalent apertures respectively.

The genital ducts are seen emerging from the lower margin of the first intestinal coil, and following the course of the intestine open into the atrium in close contiguity to the anus.

60. An Ascidian, of a different species, dissected to shew its digestive tract.

The external test has been removed, and the mantle and wall of the branchial sac reflected.

A black bristle has been passed through the inhalent aperture, branchial sac, and mouth, into the stomach, and a red rod through the exhalent aperture and atrial chamber. The mouth is situated at the bottom of the branchial sac, and leads by a short œsophageal canal into an oblong stomach. Proceeding from the stomach the intestine describes two curves, the first of which has its convexity directed towards the nerve-ganglion (marked by a black bristle being placed under it), and its concavity towards the heart. Hence the primary flexure of the intestine is said to be hæmal. The second curve has its concavity towards the ganglion, and terminates in a rectum which has its anal orifice in the atrial chamber. The floor of the stomach is seen to be marked by a longitudinal ridge, which is prolonged into and along the intestine, so as partially to divide the tube into two canals, by which arrangement the absorptive surface is much increased. On the side of the branchial one opposite to the mouth is to be seen a portion of the 'endostyle.'

Family *Salpidae*.

61. A *Salpa maxima*, the asexual form. A green rod has been passed through the body-cavity, so as to protrude from the anterior and posterior orifices. The test is double, being composed of an inner and an outer layer: of which the latter is semi-transparent, uniform in texture, and almost colourless: the former is more opaque, of a yellowish tint, and provided with numerous transverse muscular bands. The two are seen to be united at one point on the right side near the posterior orifice.

Extending across the body-cavity diagonally, from right to left, are the branchiæ. In the left-hand lower corner is the liver, close to which is a young chain of sexual salps.

62. In this specimen the chain of sexual salps is nearly mature, and will shortly separate from the animal.

63. A chain of sexual forms of the same species. In the centre are two specimens, chained together, of *Salpa pinnata*.

Bay of Naples, 1874. F. M. Balfour.

Class POLYZOA.

64. A Sea-Mat (*Flustra foliacea*).

The zooids in the living specimen are contained within the ovoidal corneous cells of which the ramose and lamellar 'cænæcium' or 'polyzoarium' is composed. These cells are arranged on both sides of the frond in parallel longitudinal rows; examined with a lens, from four to eight short and blunt spines are seen to surround the broad end of each cell. The stone to which these fronds are attached is encrusted with these cells.

Class BRACHIOPODA.

65. Two specimens of *Lingula anatina*.

They have been attached to a sheet of mica. The valves are of a horny texture, and of nearly equal size, differing in this respect from the majority of the class, which have equilateral but inequivalve shells. Furthermore, the valves are not lateral,

as in the Lamellibranchiata, but dorsal and ventral in relation to the animal they contain. Between the margins of the umbones the long muscular peduncle by which the animal is attached is protruded. The dorsal valve and mantle of the right-hand specimen have been removed, but the ventral mantle-lobe, with its margin and fringe of setæ, is left intact, adhering to the inner surface of the ventral valve. Little of the body, which occupies but a small part of the shell, can be distinguished, with the exception of a white glandular central mass, the liver, and the various muscles by which the adduction and adjustment of the valves is effected. The lateral margins of the mouth are produced outwards into two muscular ciliated 'arms,' which occupy quite one-half of the shell, and are curved inwards towards each other. The bases of the arms are connected with each other by a vertical membranous partition.

Division II. MOLLUSCA PROPER.

Class LAMELLIBRANCHIATA.

66. A Fresh-water Mussel (*Anodonta cygnea*), removed from its shell, so as to shew the general relation of the mantle and branchiæ. The two strings by which the animal is suspended are attached to the mantle, one to each of its lobes. Bristles are inserted into the mouth and anus.

The mantle consists of two lateral lobes, confluent with each other along the medio-dorsal line, but free throughout the whole of their ventral margins. With the exception of the free, thickened, glandular, and shell-secreting ventral margin, which is homologous with the 'collar' of Gasteropoda, each pallial lobe is comparatively thin, and in the fresh condition nearly transparent. In the vicinity of the anus the free edge of each mantle-lobe is beset with a fringe of minute papillæ. This papillose border corresponds with that part of the mantle which in the siphonate Lamellibranchiata is produced into the cylindrical exhalent and inhalent siphons. There are two pairs of branchiæ on each side of the central body-mass; each gill is a flattened bag composed of longitudinal and transverse fibres,

and richly ciliated. By the fusion of the two inner gill-lamellæ through the hinder third of their extent, the mantle-cavity is divided into a superior or anal chamber, into which the anus opens, and an inferior or pallial chamber. The anal chamber, indicated in the preparation by a white bristle, corresponds to the exhalent siphon. Anterior to this union the two laminae are free, but are united to the body-mass anteriorly to this. On each side of the mouth is a pair of lamellar labial palpi.

67. A vertical section of an *Anodon*. The section has been taken through the heart and intestine, the organ of Bojanus, the visceral mass, and the foot. A white bristle has been inserted into the ventricle, through which the intestine is seen to pass. Immediately beneath the heart a black bristle is inserted into the great dorsal sinus, and right and left of this, a white bristle has been placed in the secretory portion of the organ of Bojanus, which overlies the excretory portion. The preparation also shews the relation of the mantle and the two pairs of gills to the median visceral mass. The three inferior bristles indicate sections of the intestine which have been cut through in the specimen.

68. The same, prepared to shew its digestive system.

The right lobe of the mantle and the right gill have been removed, and part of the muscular foot and liver cut away to shew the convolutions of the alimentary canal, which has been filled with a light blue injection. A blue glass rod has been placed in the mouth.

69. The same, prepared to shew the two muscles which serve to draw the valves of the shell together (the anterior and posterior adductor muscles). The strings by which the proportion is suspended are attached to the posterior extremity of the shell. Rolleston, p. 54.

70. The same, to shew its nervous system.

The animal is suspended on mica with its mouth upwards. Part of the muscular foot has been removed on the left side to shew the pedal ganglion *in situ*. Part also of the organ of Bojanus has been cut away to shew the nervous commissure connecting the cephalic with the parieto-splanchnic ganglion, and posteriorly the two inner lamellæ of the two inner gills have been separated so as to shew the commissure last men-

tioned throughout its entire length. The 'labial' ganglion is seen lying on the anterior retractor muscle of the foot, and immediately below the anterior adductor. Three commissural cords, under each of which a slip of blue paper has been placed pass off from it. The one which is seen passing vertically downwards along the base of the inner gill and through the organ of Bojanus, connects the labial with the parieto-splanchnic or branchial ganglion, which is seen lying on the posterior adductor. A little way to the left may be seen the terminal portion of the corresponding commissural cord of the opposite side. A slip of blue paper has been placed beneath it. A second cord is seen passing obliquely to the left, into the substance of the foot, to join the pedal ganglion. The third cord passes upwards towards the anterior adductor, and connects the labial ganglion with its fellow of the opposite side. From the pedal ganglion several nerves may be seen radiating to supply the muscular foot. A black bristle has been placed beneath a bundle of them. The two parieto-splanchnic ganglia of the two sides of the body are closely apposed, so as to form a transversely oblong mass. Nerves from these ganglia may be noticed passing to the mantle and inner gills.

71. Two Oysters (*Ostrea edulis*).

The upper specimen shews the single adductor muscle; the lower, the hinge-ligament, which causes the valves to open. Each valve is composed of a series of concentric calcareous laminæ, overlapping each other by their free margins.

72. A specimen of *Pholas dactylus*, to illustrate the arrangement of the mantle in a *Siphonate* Gasteropod.

A *red* rod has been placed in the exhalent orifice, and a *blue* rod in the inhalent.

The valves of the shell have been partially opened, so as to shew the union of the ventral margins of the mantle. In front the mantle-lobes are widely separated, so as to allow of the protrusion of the muscular foot between them. In front of the foot they are again united. Inferiorly the mantle-lobes are prolonged downwards into a cylindrical tube, divided into two canals by a median attachment. These canals are the exhalent and inhalent orifices.

Class GASTEROPODA.

Order 1. *Opistho-branchiata*.

73. A Sea-Hare (*Aplysia*), dissected to shew the general disposition of its organs.

In the upper of the two specimens on the tablet the rudimentary shell has been displaced to the right side of the animal, and the left mantle-lobe deflected, thus exposing the heart, branchiæ, and terminal segment of intestine.

The arborescent branchiæ, protected by the rudimentary shell, are situated behind the heart and towards the rear of the body (hence the position of the animal in the order *Opistho-branchiata*), but in front of the anus, into which a black bristle has been inserted. A black bristle has also been placed under the bilocular heart and the rectum.

In the lower preparation, the muscular envelope of the animal has been separated from the foot along the left side, and reflected to the right, together with the heart, branchiæ, and parts of the generative system. The digestive tract has been arranged mainly on the left side.

The ribbon-like salivary glands (two in number) have been pinned to the left of the animal's head. Posteriorly to the buccal mass the œsophagus is seen communicating with the stomach, which is divided by certain constrictions into three compartments, the last of which, from the muscularity of its walls, has been termed a gizzard. Surrounding the gizzard and communicating with it by a large duct is the liver. The intestine, on emerging from the liver, curves round the lower border of the specimen, and passes to the anus. On the exterior of the liver is the hermaphrodite gland, a white, irregularly shaped, granular mass. From it a slightly convoluted hermaphrodite duct passes forwards to the albuminiparous gland; and after this junction and the reception of the duct of a minute vesicle (seen near the anterior margin of the gland), the duct passes upwards as a much larger tube to its external opening, prior to reaching which it communicates with a globular spermatheca. A little to the right of this part are the heart and branchiæ. A bristle has been placed under the former.

The nervous collar, consisting of supra-œsophageal and sub-œsophageal ganglia, connected by commissural strands, is seen surrounding the œsophagus. From the sub-œsophageal ganglia two nervous chords pass obliquely to a ganglion on the right, (indicated by a black bristle), situated immediately in front of the spermatheca: from this ganglion nerves are given off to the genitals, liver, and branchiæ. From the cephalic ganglia nerves are seen passing to the tentacles.

Order 2. *Prosobranchiata*.

74. A Common Whelk (*Buccinum undatum*).

The proboscis has been extended to its full length, and a blue rod inserted in the mouth. The foot and operculum are in the positions which they would assume when the animal is in motion.

75. Another specimen, to shew the structure of the proboscis, which has been laid open. A blue rod has been inserted into the œsophagus. At the end of the proboscis is seen the strap-like tongue, armed with a treble row of horny teeth. At the end of the proboscis nearest to the head it is marked horizontally with numerous striations, indicating the mode in which it is folded upon itself within the body of the animal. Its termination, which is not so marked, remains unfolded when the organ is drawn in. The movements of the tongue are effected by the longitudinal muscles seen beneath the œsophagus.

76. Another specimen, to shew the penis and the sperm-duct.

The penis has been protruded, and a portion of the shell taken away so as to shew the sperm-duct, under which, where it emerges from the body of the animal, a black bristle has been placed. The penis is a very large, flattened organ, and when laid open is found to be composed of interlacing muscular fibres. The sperm-duct, having pursued a tortuous course through the penis, terminates by opening externally at the extremity of a projection near its blunt end.

77. Another specimen, to shew the digestive tract.

The whole of the mantle has been removed, the animal suspended by its penis and proboscis, and the alimentary canal

partially filled with red injection. The proboscis has been unfolded and laid open so as to display the commencement of the alimentary canal, into which a blue rod has been inserted.

The œsophagus is of considerable length. At the base of the proboscis it is clasped by large, coarsely-lobed, irregular, salivary glands whose ducts open into it, as does a long and apparently glandular cæcum, which is coiled round its subsequent portion. Before it reaches the stomach there is a small conical crop. The stomach is somewhat peculiar in shape, of moderate size, and has its œsophageal end produced into a short conical cæcum. A black bristle has been placed under this portion. The intestine, compared with the preceding portion of the digestive tract, is remarkably short. Its primary flexure is hæmal, and its terminal segment, or 'rectum,' very large and capacious. The anus opens into the pallial cavity, and is indicated by the insertion into it of a red rod. Near the pyloric end of the stomach is the bi-ocular heart.

78. A specimen of *Triton rugosus*, prepared to shew the relation of a Branchio-gasteropod to its shell, when it is retracted within its shell, and the mouth closed by the operculum.

The first whorl of the shell, being broken away, shews the broad muscular foot; the tentacles behind; and further still the proboscis, much retracted, which has been made visible by removing a portion of the overlying integument. Behind this again is seen the mantle, a large portion of which has been cut away, so as to expose its cavity, in which are contained the gills, the exhalent siphon, and the intestine. The branchiæ are seen as a fringe, depending from the lower border of the mantle-cavity. A blue rod has been placed in the anus, a red rod in the exhalent siphon. At the termination of the branchiæ the liver commences, and occupies the last two whorls of the shell.

Order 3. *Pulmonifera*.

79. The upper specimen on the tablet, a Garden Snail (*Helix aspersa*), shews the position and relations of the aperture of the pulmonary sac when the animal is in motion. The aperture,

into which a white bristle has been inserted, is situated immediately within the peristomial margin of the shell, a little to the right side of the animal, but to the left of the anus, which is indicated by a black bristle.

In the lower of the two specimens, a Roman Snail (*Helix pomatia*), nearly the whole of the shell has been removed, and a red injection thrown into the auricle and pulmonary vessels. An incision has been made transversely through the roof of the pulmonary sac, and its interior exposed by turning back the posterior flap. A black bristle has been passed through the external orifice into its cavity. Attached to the displaced posterior half of the sac is the pale-coloured and triangular kidney, and close to it on the left side the bilocular heart.

The floor of the respiratory cavity is non-vascular, being composed of muscular fibres arranged transversely and longitudinally: and functionally representing a diaphragm. On the right side of this floor the rectum may be seen passing forwards to the anus, which is situated to the right of, but in close proximity to, the respiratory orifice.

80. A Roman Snail (*Helix pomatia*), dissected to shew the general disposition of its organs.

The shell has been removed. The liver, heart, and pulmonary sac have been arranged on the left side of the preparation, the stomach and salivary glands in the middle, and the generative system mainly on the right side.

Posteriorly to the buccal mass, and on emerging from the nerve-collar through which it passes, the œsophagus expands into a long and fusiform stomach. The salivary glands overlie the stomach about the middle of its length, and the two ducts therefrom pass forwards through the nerve-collar and pierce the buccal mass anteriorly to it. Posteriorly the stomach diminishes in size, and at its point of junction with the intestine receives the single large hepatic duct. The first segment of the intestine ascends towards the heart, and, together with the stomach, makes up a curve having its concavity towards the pedal ganglia. Hence its primary flexure is said to be neural. Below the heart the intestine describes certain convolutions in the substance of the liver, and reappearing on the extreme left of the preparation, proceeds straight to the anus, which is

indicated by a black bristle. The nerve-collar (through which a black bristle has been passed) encircles the œsophagus. From the supra-œsophageal portion delicate nerves pass forwards to supply the tentacles, lips, &c. A little to the left of the left salivary duct may be seen the left parietal nerve, double at its termination, passing from its origin in the parieto-splanchnic ganglion to its peripheral distribution in the body-walls. Between the left salivary duct and the stomach the central portions of three other nerves may be seen in connection with the same ganglia, viz. the two parietal nerves of the right side and the great visceral nerve in connection with the aorta. The second black bristle (the lower one) is inserted into the anterior aorta, which passes through a foramen between the upper or parieto-splanchnic, and the lower or pedal portion of the sub-œsophageal mass.

On the left of the preparation is the pulmonary sac. The greater portion of its diaphragm-like floor has been removed, and its highly vascular roof exposed. At the lower edge of the sac, and on the right side of it, is the bilocular heart. To the left of the heart is the pale triangular kidney.

The hermaphrodite gland is lodged in the concavity of the penultimate and ante-penultimate coils of the liver, which occupy the lowest part of the preparation. From this gland a convoluted hermaphrodite duct passes upwards to a tongue-shaped body, the albuminiparous gland; and after this junction the duct passes upwards again as a thicker and much plicated tube, having closely applied along it a ribbon-like glandular prostatic mass. At about the level of the animal's head this hermaphrodite duct divides into the oviduct and the vas deferens. The vas deferens passes forwards and joins the base of the penis at the point where the long whip-like flagellum also joins it. On the extreme right of the preparation is the globular spermatheca with its duct leading to the oviduct into which it opens. Connected with the vagina is the 'dart sac,' and between the latter and the oviduct are seen the 'multifid vesicles.'

The external generative opening is on the right side of the animal at the base of the right superior tentacle.

81. A Roman Snail (*Helix pomatia*), dissected to shew its nervous system.

The viscera have been removed, with the exception of the buccal mass, œsophagus, and portions of the generative system.

The œsophagus has been drawn through the nerve-collar and pinned in front of the animal's head.

As in other Gasteropoda the central nervous system forms a collar encircling the œsophagus. The supra-œsophageal part of the collar is composed of the two cephalic ganglia and the broad commissural band connecting them. The sub-œsophageal ganglia, which are connected with the former on each side by a double nervous cord, consist of a lower or pedal, and an upper or parieto-splanchnic element. From the former of these elements nerves may be seen passing to the muscular foot, while from the parieto-splanchnic portion three large nerves pass off to supply the parietes. Of these, two pass to the right side, and one, which bifurcates, to the left. Arising from the same source is the great splanchnic nerve. This nerve proceeds posteriorly, in close proximity with the anterior aorta, as far as the albuminiparous gland, at which point it divides into two branches, one of which goes to the liver, and the other to the hermaphrodite gland. The anterior aorta passes forwards through a foramen between the parieto-splanchnic and pedal ganglia to supply the anterior part of the body. This fact, together with the distribution of the nerves of the two ganglia respectively, enables these two elements of the sub-œsophageal mass to be differentiated.

From the cephalic ganglia nerves, under which a slip of blue paper has been placed, may be seen passing forwards to supply the superior or eye-bearing tentacles, together with the upper lip and adjacent sensitive surface.

From each cephalic ganglion a nerve passes to join a small stomato-gastric ganglion, situated on either side of the point at which the œsophagus joins the buccal mass. The two stomato-gastric ganglia are connected by a commissural strand, and from each nerves are given off to supply the buccal mass, salivary glands, and œsophagus.

82. A Grey Slug (*Limax cinereus*), dissected to shew the general distribution of its organs.

The muscular envelope of the animal has been separated from the foot proper along the left side, and turned over to the right, together with the shield-shaped mantle, heart, and pulmonary sac. The digestive tract and the glandular organs in connection with it have been displaced from their natural positions and pinned to the tablet on the left side of the preparation, while the generative organs, similarly displaced, occupy the right side. The superior tentacles have been pinned by their retractor muscles in front of the animal's head.

On the left side of the head are the salivary glands, two lobulated organs connected with the mouth by two ducts which pass forwards through the nerve-collar and pierce the buccal mass anteriorly to it. The glandular mass which encircles the anterior margin of the buccal mass is 'Semper's organ,' supposed by its discoverer to be olfactory.

The œsophagus, emerging from the buccal mass, passes through the nerve-collar, and almost immediately expands into a long fusiform stomach, the posterior segment of which, separated from the anterior three-fourths by a slight constriction, receives the two biliary ducts, one from each of the two principal lobes into which the liver is divided. A convoluted intestine succeeds the stomach, passing to the middle line of the body, where, after a short retrograde course posteriorly, it bends sharply upon itself, ascends, and terminates at an anus situated on the right side of the respiratory aperture.

In the upper portion of the specimen, and on the right side, is seen the reticulated structure of the roof of the pulmonary sac, the floor having been removed.

A little posterior to this point is seen the bilocular heart, together with portions of the two trunks into which the aorta is divided. Beneath the heart is the sub-triangular yellow kidney. In the natural position of these parts the kidney would of course overlie the heart. Immediately posterior to the buccal mass, and encircling the œsophagus, is the nerve-collar, consisting, superiorly, of the two cephalic ganglia connected with each other by a broad commissural band, and inferiorly of the parieto-splanchnic and pedal ganglia. The supra-œsophageal and sub-œsophageal masses are mutually connected by a double commissural strand embracing the gullet.

From the cephalic ganglia nerves may be seen passing anteriorly to supply the tentacles, 'Semper's organ,' and the adjacent parts. From the upper or parieto-splanchnic segment of the sub-œsophageal mass nerves are given off to the parietes and viscera. Of the visceral nerves, the peripheral divisions have been removed, their origins alone remaining. From the lower or pedal segment a leash of nerves passes off to supply the muscular foot. A black bristle (the longest of the two) has been inserted into the foramen which gave passage to the anterior branch of the aorta.

In the angle which the gullet makes with the buccal mass, and situated on the latter, is the right stomato-gastric ganglion of the sympathetic system.

Of the generative organs, the hermaphrodite gland (situated at the bottom of the right side of the tablet) is connected by a single convoluted duct with the large, somewhat irregularly shaped, and coarsely lobulated albuminiparous gland.

The duct, on emerging from the gland, is of much larger size and plicated. It has associated with it throughout the greater portion of its length a ribbon-like granular prostatic mass. Immediately above the pulmonary sac this subdivides into a long thread-like duct, or 'vas deferens,' which joins the root of the penis, and a short, thick oviduct which opens outwards.

In the re-entering angle between the penis and the oviduct, and communicating with the latter, is the globular spermatheca. The penis, pinned by its retractor muscle a little to the right of the right superior tentacle, is seen to communicate by its saccular portion with an aperture common to it and the oviduct. The 'multifid vesicles,' and 'dart sac,' so characteristic of the majority of the European Helicidæ, are absent.

83. A Black Slug (*Arion empiricorum*), dissected to shew its general anatomy.

The structure of this species differs in no very important respect from the last. The most noteworthy points of distinction are the following:—the extremely lobulated and diffuse character of the salivary glands and liver; the large and compact hermaphrodite gland, and the smaller and more finely lobulated albuminiparous gland. The spermatheca also

is larger, and connected by a much shorter duct with the vagina.

On the right side of the preparation the heart is well displayed, and proceeding from it the aorta, divided into its anterior and posterior branches, the former of which passes anteriorly to supply the organs of that part of the body, while the latter passes posteriorly to supply the liver and intestine.

A yellow glass rod has been placed under the cephalic ganglia, and a red rod under the sub-œsophageal mass.

84. A wax model of a Grey Slug (*Limax cinereus*), prepared to shew its general anatomy.

A. The animal entire.

B. Anterior half of the animal. A portion of the mantle has been removed in order to expose the rudimentary shell which would not otherwise be seen.

C. General disposition of the internal organs, in their natural position, on the removal of the integument.

D. The organs have been partially separated in order that the general disposition of the alimentary canal and the generative organs may be seen.

E. Designed to shew the salivary glands, the nervous system, and the pulmonary sac; together with the heart and kidney.

F. The generative system.

G. The buccal mass laid open to shew the odontophore and the nervous system.

The numbers affixed denote the different organs as follows:

- | | |
|--------------------------|---|
| 1. Buccal mass. | 14. Posterior Aorta. |
| 2. Salivary Glands. | 15. Pulmonary vessels lining roof of pulmonary sac. |
| 3. Stomach. | 16. Branches of posterior aorta supplying viscera. |
| 4. Intestine. | 17. Supra-œsophageal ganglion. |
| 5. Liver. | 18. Sub-œsophageal ganglion. |
| 7. Rectum. | 19. Right and left stomatogastric ganglia. |
| 8. Anus. | 20. Buccal mass laid open to show odontophore. |
| 9. Renal organ. | 21. Duct of salivary gland. |
| 10. Duct of Renal organ. | |
| 11. Ventricle. | |
| 12. Auricle. | |
| 13. Anterior Aorta. | |

- | | |
|---------------------------|-------------------------|
| 22. Hermaphrodite gland. | 27. Hermaphrodite duct. |
| 23. Shell. | 28. Vas deferens. |
| 24. Shield-shaped mantle. | 29. Oviduct. |
| 25. Albuminiparous gland. | 30. Spermatheca. |
| 26. Penis. | |

Class CEPHALOPODA.

85. A Cephalopod (*Eledone*).

A *blue* rod has been inserted into the mouth, a *red* rod into the funnel.

The body is symmetrical, the organs being arranged symmetrically with reference to an imaginary vertical plane drawn from before backwards through the centre of the body. The animal is divided into an anterior or cephalic portion, 'prosoma,' and a posterior, or 'metasoma.' In the centre of the 'prosoma' and surrounded radially by the arms, is the mouth, and on the opposite sides of it are the sessile eyes. The eight muscular arms which surround the mouth correspond to the anterior portion of the foot, 'propodium,' of the Gasteropoda, and are formed by the extension of the antero-lateral parts of the foot forwards in front of the head, and their union with each other in front of the mouth: so that this aperture, from its position in front of the anterior margin of the foot, comes to lie in the centre of it. The margins are produced into long muscular processes or arms; which, in this species, are set with a single row of suckers, by means of which the animal can attach itself to foreign bodies, seize its prey, or move from place to place.

The 'metasoma,' which contains the chief organs of vegetative life, is invested externally with a thick muscular coat, or mantle. This terminates in front freely, and so allows the entrance of water into the pallial cavity. On the dorsal surface of the specimen a muscular tube may be seen emerging from the pallial cavity: this is the 'Funnel.' It is formed by an extension of the 'epipodia' upwards, and their subsequent fusion in the medio-dorsal line. The Funnel is broader posteriorly than it is anteriorly, and its base is covered by the superior margins of the mantle.

The *Octopoda*, to which this species belongs, have no internal skeleton of any kind. The specimen was taken at Algiers.

86. A Calamary (*Loligo vulgaris*), dissected to shew its internal skeleton, the 'gladius' or 'pen.' The 'gladius' extends throughout the whole length of the dorsal portions of the mantle. It is of a horny texture, dark in colour, and consists of a median shaft with two lateral alæ. It lies loosely in the cavity, so as easily to fall out when the integument is cut away. This species represents the family *Teuthidæ*.

87. A Sepia (*Sepia officinalis*), dissected to shew the position of the internal skeleton, or 'cuttle-bone.' The 'cuttle-bone' is imbedded in the dorsal integument. It is a light, cellular, calcareous body, peculiarly characteristic of the genus Sepia. Its form is an elongated oval, convex, and marked by a slight median keel on the dorsal surface, and partly convex and partly concave on the ventral surface. The margin of the 'bone' is seen to be more or less horny, and darker in colour. This species represents the family *Sepiadæ*.

88. Portion of 'cuttle-bone' from a Sepia, treated with acid to remove the earthy matter, and shew the laminated structure of the residual animal matter.

89. The lips and jaws of a Sepia (*Sepia officinalis*).

A portion of the lips has been removed, and a black and white bristle passed through the mouth into the œsophagus.

The lip, which surrounds the mouth loosely, is beset with numerous elongated papillæ. The jaws consist of two horny beaks of a brown colour, in shape not unlike those of a parrot. They are embedded in a mass of muscular fibres by which the movements of the jaws are regulated. The margin of the lower one overlaps that of the upper. Between them is seen the tongue with its serrated surface. From the hinder portion of the buccal mass the œsophagus may be seen passing downwards through the cartilaginous ring towards the stomach.

90. A Cuttlefish (*Eledone*), dissected to shew its digestive, respiratory, circulatory, and generative organs.

The bag containing the viscera has been laid open, and the mantle pinned back. An incision has been carried through the cartilaginous ring at the base of the tentacles and the muscular mass surrounding the jaws, so as to expose the origin of the œsophagus. This is a narrow tube, which, after a course of uniform diameter for about an inch, suddenly expands into a

large, easily dilateable, 'crop' (pinned to the right in the preparation). A short passage leads thence to the stomach, a small muscular organ which has been called 'gizzard,' from its muscularity. Close to, and opening into, the stomach is the spiral cæcum; a little beyond this the intestine receives the duct of the liver, and then proceeds in an upward direction to the anus, which is situated near the 'funnel.' The liver is a large globular organ, containing within its substance the inkbag, the duct of which opens into the intestine near the anus. There are two sets of salivary glands. The uppermost are closely attached to the under surface of the buccal mass, and open by a single duct into the œsophagus. The lower pair are larger and have each a duct, which shortly unite to form a common canal, opening into the buccal mass at the base of the tongue.

On either side of the lower part of the preparation are the branchiæ, each with its venous heart attached. The great venous trunk has been removed, but the cut end of one of the two veins that unite to form it has been stitched back in the centre of the preparation. A piece of red glass indicates the aorta. On the left of this is the great pallial ganglion, from which numerous nerves may be seen radiating into the substance of the mantle. Between the two venous hearts, arching over the systemic heart, are the two venæ cavæ, furnished with clusters of 'spongy cellular bodies.' At the bottom of the preparation is the ovary, with its ducts, of which one is pinned out to the right.

91. Circulatory system of the same species. The arterial portion has been injected with blue. The systemic heart is in the centre; on either side of it are the venous dilatations into which the blood is received from the branchiæ, prior to its reception by the ventricle. Pinned on the right and left sides of the preparation, and uninjected, are the pulmonary hearts, with their so-called 'appendages.' From the systemic ventricle the anterior and posterior aortæ may be seen passing respectively upwards and downwards.

From a specimen taken at Weymouth.

SUB-KINGDOM. VERTEBRATA.

Division I. ANALLANTOIDEA.

Sub-division I. ICHTHYOPSIDA.

Class PISCES.

Order TELEOSTEI.

92. A common Perch (*Perca fluviatilis*).

The roof of the cranial cavity and the greater part of the left wall of the abdomen have been removed, exposing the brain and most of the internal viscera. The following external features may be noticed: the absence of distinct divisions between cephalic, thoracic and caudal regions; the imbricated scales of the variety called ctenoid; the paired, and median unpaired fins supported by hard, unbranched, non-striated rays, and by soft, branching, striated rays; the large size of the eyes; the absence of distinct eyelids; the presence of persistent visceral clefts interposed between the visceral arches, and protected partly by a long operculum and partly by a membrane supported on a series of branchio-stegal rays. Bristles are inserted into the paired cæcal nasal sacs, each of which has two apertures. The pectoral fins are situated a little behind the free edge of the operculum; and behind these again are the paired ventral fins. All the rays of the pectoral fins are soft, but the first ray of the ventral fin is hard. The mesial fins are the dorsal, divided into two divisions, the anterior of which is composed entirely of hard rays, as is the rule in Acanthopterygian fishes, the caudal, and the anal. "A series of scales perforated by mucous ducts supplied with nerves forms a lateral line along the trunk, the contour of which corresponds with that of the upper line of the back." (Rolleston.) A portion of the skin has been removed from the lateral surface in the vicinity of the head—thus exposing the lateral branch of the pneumogastric nerve, which passes posteriorly in relation with the lateral line. A slip of blue paper has been placed beneath this nerve. The tendinous intersections dividing the great lateral muscles into a series of segments, may also be noticed.

A blue injection has been thrown into the gills, which are four in number on each side. Each gill carries on its outer or convex edge a double row of short, flattened, vascular filaments, and on its inner or concave side a row of short bony tubercles. In the angle between the converging gills are to be seen the central parts of the circulatory system, consisting of a triangular ventricle, leading into the pulmonary aorta anteriorly, an auricle which slightly overlaps the ventricle, and a 'sinus venosus,' into which the great veins pour their blood. The heart is separated from the subjacent liver by a transverse peritoneal partition. The latter viscus is of large size, rounded in front and above, or divided inferiorly by a deep fissure into a right and left lobe. Depending as it were from beneath the inferior edge of the liver is the cæcal stomach, with the intestine, to which are appended three pyloric cæca passing out of it at right angles. After describing an S-shaped convolution, the intestine expands into a dilated, thin-walled, and transparent rectum. Behind the rectum is the single large, pear-shaped ovary, and still more posteriorly the small oval bladder. The spleen lies in a loop of intestine, and a portion of it, stained with the blue injection, is to be seen between the lower edge of the stomach and the ovary. The large semi-transparent air-sac, closely applied to the hinder wall of the abdomen, has been left intact in the greater part of its extent; but the posterior third has been removed, in order to expose a portion of the kidney, which lies between it and the vertebral column throughout the whole extent of the abdomen. A slip of blue paper has been placed beneath one of the ureters. The anus is situated anteriorly to the openings of the generative and urinary ducts, and not posteriorly, as in the higher Vertebrata.

In the brain the cerebral hemispheres are small, and their surfaces are slightly corrugated. From their anterior extremities the olfactory nerves are given off. The optic lobes, which succeed them, are the largest divisions of the brain: and by their opposition to the hemispheres in front, the optic thalami are hidden from view. The next division is the small, smooth cerebellum, and the fourth ventricle which it bounds anteriorly is widely open. (Rolleston, p. 40.)

Class II. AMPHIBIA.

Order BATRACHIA.

93. A Common Frog (*Rana temporaria*), dissected and injected to shew the reni-portal system, and the chief trunks of the arterial and venous systems.

A white bristle has been placed beneath the crural vein; if this vein be traced upwards from the leg it will be seen that in the groin it divides into two branches, and that one of them, the outer one, passes upwards to the kidney, along the outer side of which it runs, and ultimately breaks up into capillaries in that organ. This is the reni-portal vein. The second branch of the crural curves round the front of the groin, towards the medio-ventral line, where it unites with the corresponding branch of the crural vein of the other leg to form the epigastric or umbilical vein. The latter vein, after receiving one or two small twigs from the large and persistent allantois, passes up the median line of the abdomen till it reaches the liver, where it joins the portal vein, and subsequently ramifies through that organ. It will thus be observed that the whole of the blood in the hinder extremities of the body, after passing through the capillaries proper to these parts, traverses in addition one of two sets of capillaries, either going by the reni-portal vein to those of the kidney, or by the epigastric to those of the liver. The hepatico-portal system is also shewn. A loop of intestine has been pulled out and fastened to the right side of the animal. The veins of the intestine are seen to unite and re-unite, and by successive unions to form the portal vein which passes upwards to the liver. The left superior vena cava is, as usual, formed by the union of the jugular and subclavian veins. A black bristle has been placed beneath it, just before it curves round the base of the heart to reach the right auricle. The large inferior vena cava is formed by the confluence of the veins from the two kidneys, between which it is to be seen. It passes forwards in relation with the posterior surface of the liver, and, after receiving the hepatic veins, enters the right auricle. (The liver has been displaced to the right so that the posterior surface has, in this specimen, become the anterior.)

Arterial System.—On the left side of the heart the three trunks into which the 'bulbus arteriosus' divides on that side are also to be seen. A slip of blue paper has been placed beneath them: the most anterior of the three is the carotid. The central one, which is the systemic aorta, after giving off the subclavian and vertebral arteries (the roots only of these are seen), passes downwards and joins its fellow of the opposite side, to form the abdominal aorta. The mesenteric artery which supplies the intestines is given off immediately below this junction. The third trunk is the pulmo-cutaneous artery—the pulmonary branches, proceeding to the lungs, only being seen, as the cutaneous branch has been removed. After giving off the mesenteric artery, the abdominal aorta disappears behind the kidney, though at some distance inferiorly one of its branches, the crural artery, passes downwards over a black bristle, to supply the right leg.

94. A Frog, dissected to shew its nervous system.

The nervous system consists of a brain and spinal cord. The cerebral hemispheres are much elongated, and are, as in all Vertebrata except fishes, the largest division of the encephalon. Olfactory nerves are given off from their anterior extremities to the nasal sacs. The optic thalami which succeed are largely exposed between the hemispheres and the corpora bigemina. The latter are comparatively small. The fourth ventricle is widely exposed, and the only representative of a cerebellum is the narrow band of transverse fibres which is to be seen crossing over the anterior part of the ventricle. The spinal cord is short and stout, and nerves are given off from it to the number of ten pairs. The pia mater of the cord is continued as a thin sheath, destitute of nervous matter, to the extremity of the neural canal. The last two or three pairs of spinal nerves pass backwards in the spinal canal for some little distance before passing out to their peripheral distribution. The ganglia on the posterior roots at the point where they join the anterior roots are to be seen. A slip of blue paper has been placed under the two ganglia. On the ventral surface of the throat slips of blue paper have been placed beneath two cranial nerves and the first spinal, viz. the glossopharyngeal, pneumogastric and hypoglossal.

95. A Frog, dissected to shew certain features not shewn,

or but imperfectly shewn, in the previous preparations. The greater part of the stomach and intestine has been removed, a portion of the dilated rectum alone remaining. The heart, consisting of two auricles and a ventricle, is seen in the mesial line at the base of the throat. The carotid with its bulb-like carotid gland and the lingual artery which it gives off, the systemic and pulmonary trunks, into which the aortic bulb divides on each side, are perhaps better seen. The same remark will also apply to the left musculo-cutaneous vein, which is to be seen returning the blood from the vascular and respiratory skin to the left superior vena cava. On each side of the heart are the simple sac-like but internally vesicular lungs. Below the heart are two lobes of the liver, the third having been removed to shew the large gall-bladder and the root of the epigastric vein. A little to the right of the darkly stained gall-bladder is the globular spleen. The two oval white bodies seen beneath the liver are the testes. The vasa deferentia emerge from and pass through the substance of the anterior part of the kidney, and finally open into the urogenital duct, which can be traced down the outer edge of the latter organ. To the terminal part of each of these ducts is appended a large lateral dilatation, the vesicula seminalis. The large bifid allantois hangs down in front of the rectum.

Division II. ALLANTOIDEA.

Sub-division. SAUROPSIDA.

Class AVES.

Order CARINATÆ.

96. A Pigeon (*Columba livia*), dissected to shew the chief features by which it differs from a Mammal.

The roof of the cranial cavity has been removed and the brain exposed. The right wall of the chest and abdomen has also been removed. The origins of the well-developed pectoralis major and pectoralis minor muscles are to be seen along the edge of the sternal keel. The skin has been removed from the front of the throat, and the crop, after having been distended with spirit and hardened, has been laid open and its cavity displayed. The cerebral hemispheres are pear-shaped, with a smooth surface,

and differ from the corresponding parts of the Reptilian brain in being in contact with the cerebellum, and also by reason of their much greater size. The cerebellum is large, and divided by numerous transverse 'sulci' into well-marked lobules. It has no lateral lobes. The 'corpora bigemina,' instead of being in contact with each other mesially, as is always the case in other Vertebrates, are thrown down towards the infero-lateral surface of the brain, so that they appear as rounded lateral projections. The other points to be noted about the head, which is supported upon a long neck consisting of more than seven vertebræ, are the presence of feathers; the absence of a 'concha' to the external auditory meatus; a well-developed third eyelid, which can be drawn vertically across the eye; the edentulous character of the mouth; the horny and non-prehensile tongue; and the absence of an epiglottis. The widely dilated œsophagus expanding below into a bi-sacculated crop occupies the front of the flexible neck. The cartilaginous tracheal rings are complete. Between the chest and the abdomen the absence of a diaphragm is to be remarked. In the chest, and occupying a mesial position, is the long, acutely conical heart. The short, thick aorta which it gives off, is seen to divide into three large trunks, two 'arteriæ innominatæ,' and a descending aorta. The aorta curves over the right bronchus and not over the left, as in the Rat. Behind the crop, and lying in close apposition with each other, in a deep groove formed on the anterior surfaces of the cervical vertebræ by the inferior transverse processes, are the two carotids. A piece of blue paper has been placed beneath them. The right lung is also seen in section. It is not divided into lobules, nor is it suspended in a pleural cavity; but is so firmly connected by connective tissue to the back and side-walls of the chest as to be marked by deep grooves, the impressions of the ribs. Each innominate artery divides into three branches—the carotid, sub-clavian, and pectoral arteries. Below the heart is the liver, composed, as in most reptiles, of a large right lobe and a smaller left one, the two being connected by a transverse lobe. The apex of the heart is received into a mesial depression between the two lobes. Overlapped by the left lobe, and partially concealed by it, is the thick-walled and muscular gizzard; the

intestine, with the exception of part of the duodenum and the rectum, has been removed. The duodenum hangs down from the gizzard, and contains within its primary convolution the compact pancreas, with the three ducts by which its secretion is poured into the intestine. In addition, two other ducts, one of considerable size, and the other very small, are seen leaving the liver and joining the duodenum. These are the hepatic ducts. The right testis is also to be seen, as a large ovoid whitish body. A little beneath and behind the testis is the tri-lobed kidney. Each of its lobes lies in a depression in the under surface of the ischium. The ureter emerges from its inner edge, and passes posteriorly to its cloacal outlet. A little to its outer side is the sinuous vas deferens. The rectum hangs down from the bottom of the specimen, and about an inch from the anus two small conical cæca join it. On the dorsal surface of the extreme tip of the tail is the uropygial gland, consisting of two pear-shaped lobes. The biceps muscle of the right leg should be noticed passing downwards from the ischium. It passes through a pulley-like loop of tendon at the back of the knee to its insertion in the tibia.

Class MAMMALIA.

Order RODENTIA.

97. A Rat (*Mus decumanus*), dissected to shew the distinctive features of its class.

In addition to the more important anatomical details, the following external points should be noticed: the hairy skin, the existence of distinct outlets for the urinary and generative organs as well as for the intestine, the vibrissæ on the snout, the distinctly annulated tail, the cleft upper lip, the moveable muffle, and the ungulate digits. To expose the viscera, the greater part of the right side of the abdomen and thorax has been removed. A red injection has been thrown into the arteries and veins. The general visceral cavity is divided by the transverse tendino-muscular diaphragm into a thorax, containing the heart and lungs, and an abdomen, in which the liver, stomach, intestines, &c. are situated. The lungs are freely suspended within a pleural cavity, and are not closely adherent

to the posterior walls of the thorax, as they were in the Pigeon ; the right lung is divided into three lobes, and the left into two. The heart is mesially placed between the two lungs, and its anterior surface is about equally divided between the two ventricles. The aorta is seen curving over to the left side, and from the summit of the arch the three following vessels take origin : first, the right arteria innominata which soon splits into a right subclavian and carotid arteries ; a little to the left of this the left carotid arises directly from the arch ; and still more to the left the left subclavian, but this is hidden from view by the left superior vena cava. On each side of the root of the neck the jugular and subclavian veins are to be seen converging ; and subsequently by their union they form the right and left superior venæ cavæ. In order that these vessels might be better displayed, the thymus, or hibernating gland, has been displaced, and turned forwards over the front of the heart. Descending the neck, in close relation with the carotid artery, is the pneumogastric nerve, and a little to the outer side of the same artery the cervical portion of the sympathetic may be seen ; while in close proximity with the trachea is the delicate recurrent laryngeal. Subsequently to giving off the arteries to the head and fore limbs, the aorta passes downwards through the chest just in front of the vertebral column, and gives off numerous intercostal arteries in its course : eventually it pierces the diaphragm and becomes the abdominal aorta. A little to the outer side of the thoracic aorta a rather large vein passes upwards to the left superior vena cava, receiving during its ascending course small veins from the intercostal spaces ; this is the azygos vein. The phrenic nerves in their course downwards from their origins in the cervical portion of the spinal cord to their distribution in the diaphragm, of which they are the motor nerves, pass a little to the outer side of the superior vena cava, and afterwards in front of the roots of the lungs ; the right phrenic nerve is in close relation with the inferior vena cava. Behind, and a little to the left of the last-mentioned vein, is the thoracic portion of the œsophagus. Beneath the concavo-convex diaphragm, and attached to it by a peritoneal process—the suspensory ligament—is the five-lobed liver. This again is rounded above and adapted by its convex upper surface to the concave inferior

surface of the diaphragm, while inferiorly it more or less conceals the shrunken stomach. The stomach has a large cardiac pouch with a thin wall, and a pyloric division with much thicker and glandular walls: the œsophagus opens into it about midway between the two divisions. The epiploon, a special process of peritoneum peculiar to this class, depends from the inferior border of the stomach, and in its natural position would hang in front of the small intestines. To the left of the stomach, and attached to it by the gastro-splenic omentum, is the spleen, while below the stomach the great mass of the large and small intestines lie. Of the large intestines, the darkly stained and dilated cæcum is the most prominent, though smaller in the Rat than in most Rodentia. In the first convolution which the duodenum makes after leaving the stomach, the ramifying and diffuse pancreas lies. A black bristle has been placed beneath the hepatic duct near to its opening into the intestine at a point about an inch and a half from the stomach. On each side of the vertebral column, the slightly asymmetrically disposed kidneys are to be seen lying on the psoas muscles. Three vessels may be noticed passing out of the hilum of the right kidney; of these the anterior one is the renal artery and the posterior one the ureter, while the median and largest of the three is the renal vein. The small, buff-coloured and triangular masses attached to the anterior ends of the kidneys are the supra-renal bodies; the left one only is seen, as the right one is hidden by the hepatic lobes. Immediately below the kidney the right ovary can be seen and from it the Fallopian tube passes to the right cornu of the uterus. At a point just behind the bladder the two cornua converge to form an extremely small corpus uteri which opens into the large vagina. 'The vagina, rectum and bladder have each of them separate and independent outlets,' and into each of the three bristles have been inserted.

In the interspace between the gastric and mesenteric veins and close to the aorta from which these vessels are derived is the small, pale-coloured, semilunar ganglion: and from it two small nerves are to be observed passing obliquely upwards across the aorta to connect this ganglion with the chain of subvertebral ganglia which together constitute the sympathetic system.

The following points are to be noticed about the head in

addition to those already mentioned, viz. the large submaxillary salivary glands bounding the angle of the lower jaw, and the much smaller and more diffuse parotid gland placed behind and beneath the external ear. On the right side of the animal's head and situated at a point just in front of the ear, the extra-orbital factor of the right lachrymal gland can be seen; from this portion a duct, with a slip of blue paper beneath it, passes forwards to join a second duct from the intra-orbital part which is situated at the posterior angle of the eye and inside the orbit. Near the inner and anterior angle of the eye there is another small gland about the size of a pin's head; this is the Harderian gland. Of the third eyelid, or *membrana nictitans*, with which this gland is always correlated, a rudiment exists and can be seen covering the anterior third of the eyeball.

A black bristle has been placed beneath one or two of the external branches of the seventh or facial nerve, as they cross the largely developed masseter muscle.

The characteristic Rodent dentition should be noted: the large, scalpriform, incisor teeth, of which there are two in each jaw, growing from persistent pulps; the absence of canines, and the large interval which separates the incisors from the molars.

The roof of the skull as well as the neural arches of the cervical vertebræ have been removed in order to expose the brain and spinal cord. The hemispheres are pear-shaped, thin, and tapering in front, but broad behind; their surfaces are smooth. In the median longitudinal fissure which separates the hemispheres, the longitudinal sinus, filled with red injection, can be seen. The olfactory lobes are very large and are altogether uncovered by any anterior extension of the hemispheres. The cerebellum has a large median and two smaller lateral lobes. In the transverse slit between the hemispheres and the cerebellum the corpora quadrigemina are partially exposed. The medulla oblongata is wide and passes behind into the spinal cord.




In the Press.

A CATALOGUE
OF
EROTYLIDÆ

BY THE LATE
GEORGE ROBERT CROTCH, M.A.,
ST JOHN'S COLLEGE, CAMBRIDGE.

CAMBRIDGE:
AT THE UNIVERSITY PRESS.



UNIVERSITY PRESS, CAMBRIDGE

January, 1890

CATALOGUE OF

WORKS

PUBLISHED FOR THE SYNDICS

OF THE

Cambridge University Press.



London: C. J. CLAY AND SONS,
CAMBRIDGE UNIVERSITY PRESS WAREHOUSE,
AVE MARIA LANE.

GLASGOW: 263, ARGYLE STREET.

Cambridge: DEIGHTON, BELL AND CO.

Leipzig: F. A. BROCKHAUS.

PUBLICATIONS OF
The Cambridge University Press.

THE REVISED VERSION
OF THE
OLD AND NEW TESTAMENTS.

*The Revised Version is the Joint Property of the Universities of
Oxford and Cambridge.*

(The Cambridge & Oxford Editions are uniform in Type, Size, & Price.)

The following Editions of the Revised Version of the Holy Bible and New Testament have been already published and may be had in a great variety of cloth and leather bindings of all booksellers.

THE HOLY BIBLE.

N.B. *The Pearl 16mo., the Ruby 16mo., and the Minion 8vo., are
facsimile editions and correspond page for page with each other.*

Cheap edition for use in Schools.

- | | | |
|----------------------|-------------|-----|
| 1. Pearl type, 16mo. | prices from | 1/6 |
| 2. Ruby type, 16mo. | do. | 4/6 |
| 3. Minion type, 8vo. | do. | 7/6 |
-

A large type edition in one volume.

4. Small Pica type, Imperial 8vo. prices from 18/-
-

LIBRARY EDITIONS.

In five vols., or the Old Testament only, in four volumes.

- | | | |
|---|----------------------|-------------|
| 5. Pica type, Demy 8vo. | 5 vols., prices from | £2 |
| 6. Pica type, Demy 8vo. Old Testament only, 4 vols., | do. | £1. 12s. |
| 7. Pica type, Royal 8vo. | 5 vols., do. | £3. 2s. 6d. |
| 8. Pica type, Royal 8vo. Old Testament only, 4 vols., | do. | £2. 10s. |
-

THE PARALLEL BIBLE.

Being the AUTHORISED VERSION arranged in Parallel columns with
the REVISED VERSION.

9. Minion type, Crown 4to. prices from £1. 6s.
-

*London: C. F. CLAY & SONS, Cambridge University Press Warehouse,
Ave Maria Lane.*

THE REVISED VERSION OF THE NEW TESTAMENT.

Cheap editions for use in Schools.

- | | | |
|---------------------------|-------------|-----|
| 1. Nonpareil type, 32mo. | prices from | -/6 |
| 2. Brevier type, 16mo. | do. | 1/- |
| 3. Long Primer type, 8vo. | do. | 1/6 |

LIBRARY EDITIONS.

- | | | |
|--------------------------|-------------|------|
| 4. Pica type, Demy 8vo. | prices from | 8/- |
| 5. Pica type, Royal 8vo. | do. | 12/6 |

THE PARALLEL NEW TESTAMENT.

Giving the Authorised and Revised Versions side by side.

- | | | |
|---------------------------------------|-------------|-----|
| 6. Pearl type, 16mo. (Pocket Edition) | prices from | 1/6 |
| 7. Minion type, 8vo. | do. | 4/6 |
| 8. Long Primer type, 4to. | do. | 7/6 |

STUDENT'S LARGE PAPER EDITION.

- | | | |
|----------------------------|-------------|------|
| 9. Minion type, Crown 4to. | prices from | 10/6 |
|----------------------------|-------------|------|

All Editions of the Parallel New Testament correspond page for page with each other.

THE PARALLEL NEW TESTAMENT, Greek and English. The Greek Text edited by the Rev. F. H. A. SCRIVENER, M.A., LL.D., and printed on alternate pages with the English Parallel Minion Edition of the Revised Version.

Minion type, Crown 8vo. prices from 12/6

THE NEW TESTAMENT IN GREEK, according to the Text followed in the Authorised Version with the variations adopted in the Revised Version. Edited by the Rev. F. H. A. SCRIVENER, M.A., LL.D.

Crown 8vo. prices from 6/-

Specimens of type and size of pages with prices sent on application.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse,
Ave Maria Lane.

THE HOLY SCRIPTURES, &c.

THE CAMBRIDGE PARAGRAPH BIBLE of the Authorized English Version, with the Text Revised by a Collation of its Early and other Principal Editions, the Use of the Italic Type made uniform, the Marginal References remodelled, and a Critical Introduction prefixed, by F. H. A. SCRIVENER, M.A., LL.D., Editor of the Greek Testament, Codex Augiensis, &c., and one of the Revisers of the Authorized Version. Crown 4to. gilt. 21s.

From the *Times*.

"Students of the Bible should be particularly grateful (to the Cambridge University Press) for having produced, with the able assistance of Dr Scrivener, a complete critical edition of the Authorized Version of the English Bible, an edition such as, to use the words of the Editor, 'would have been executed long ago had this version been nothing more than the greatest and best known of English classics.' Falling at a time when the formal revision of this version has been undertaken by a distinguished company of scholars and divines, the publication of this edition must be considered most opportune."

From the *Athenæum*.

"Apart from its religious importance, the English Bible has the glory, which but few sister versions indeed can claim, of being the chief classic of the language, of having, in conjunction with Shakspeare, and in an immeasurable degree more than he, fixed the language beyond any possibility of important change. Thus the recent contributions to the

literature of the subject, by such workers as Mr Francis Fry and Canon Westcott, appeal to a wide range of sympathies; and to these may now be added Dr Scrivener, well known for his labours in the cause of the Greek Testament criticism, who has brought out, for the Syndics of the Cambridge University Press, an edition of the English Bible, according to the text of 1611, revised by a comparison with later issues on principles stated by him in his Introduction. Here he enters at length into the history of the chief editions of the version, and of such features as the marginal notes, the use of italic type, and the changes of orthography, as well as into the most interesting question as to the original texts from which our translation is produced."

From the *London Quarterly Review*.

"The work is worthy in every respect of the editor's fame, and of the Cambridge University Press. The noble English Version, to which our country and religion owe so much, was probably never presented before in so perfect a form."

THE CAMBRIDGE PARAGRAPH BIBLE. STUDENT'S EDITION, on *good writing paper*, with one column of print and wide margin to each page for MS. notes. This edition will be found of great use to those who are engaged in the task of Biblical criticism. Two Vols. Crown 4to. gilt. 31s. 6d.

THE AUTHORIZED EDITION OF THE ENGLISH BIBLE (1611), ITS SUBSEQUENT REPRINTS AND MODERN REPRESENTATIVES. Being the Introduction to the Cambridge Paragraph Bible (1873), re-edited with corrections and additions. By F. H. A. SCRIVENER, M.A., D.C.L., LL.D., Prebendary of Exeter and Vicar of Hendon. Crown 8vo. 7s. 6d.

THE LECTIONARY BIBLE, WITH APOCRYPHA, divided into Sections adapted to the Calendar and Tables of Lessons of 1871. Crown 8vo. 3s. 6d.

THE NEW TESTAMENT IN GREEK according to the text followed in the **Authorised Version**, with the Variations adopted in the **Revised Version**. Edited by F. H. A. SCRIVENER, M.A., D.C.L., LL.D. Crown 8vo. 6s. Morocco boards or limp. 12s.

The Revised Version is the Joint Property of the Universities of Cambridge and Oxford.

PREVIARIUM ROMANUM a FRANCISCO CARDINALI QUIGNONIO editum et recognitum iuxta editionem Venetiis A.D. 1535 impressam curante JOHANNES WICKHAM LEGG Societatis Antiquariorum atque Coll. Reg. Medicorum Londin. Socio. Demy 8vo. 12s.

London: C. J. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

BREVIARIUM AD USUM INSIGNIS ECCLESIAE SARUM. Juxta Editionem maximam pro CLAUDIO CHEVALLON ET FRANCISCO REGNAULT A.D. MDXXXI. in Alma Parisiorum Academia impressam: labore ac studio FRANCISCI PROCTER, A.M., ET CHRISTOPHORI WORDSWORTH, A.M.

FASCICULUS I. In quo continentur KALENDARIUM, et ORDO TEMPORALIS sive PROPRIUM DE TEMPORE TOTIUS ANNI, una cum ordinali suo quod usitato vocabulo dicitur PICA SIVE DIRECTORIUM SACERDOTUM. Demy 8vo. 18s.

"The value of this reprint is considerable to liturgical students, who will now be able to consult in their own libraries a work absolutely indispensable to a right understanding of the history of the Prayer-Book, but which till now

usually necessitated a visit to some public library, since the rarity of the volume made its cost prohibitory to all but a few."—*Literary Churchman*.

FASCICULUS II. In quo continentur PSALTERIUM, cum ordinario Officii totius hebdomadae juxta Horas Canonicas, et proprio Completorii, LITANIA, COMMUNE SANCTORUM, ORDINARIUM MISSAE CUM CANONE ET XIII MISSIS, &c. &c. Demy 8vo. 12s.

"Not only experts in liturgiology, but all persons interested in the history of the Anglican Book of Common Prayer, will be grateful to the Syndicate of the Cambridge University Press for forwarding the publication of the volume which bears the above title."—*Notes and Queries*.

"Cambridge has worthily taken the lead with the Breviary, which is of especial value for that part of the reform of the Prayer-Book which will fit it for the wants of our time."—*Church Quarterly Review*.

FASCICULUS III. In quo continetur PROPRIUM SANCTORUM quod et sanctorale dicitur, una cum accentuario. Demy 8vo. 15s.

. An Introduction of 130 pages, prefixed to this volume, contains (besides other interesting information as to the Breviary and its contents) Mr BRADSHAW'S exhaustive lists of editions and copies of the Breviary and allied liturgical books.

FASCICULI I. II. III. complete, £2. 2s.

GREEK AND ENGLISH TESTAMENT, in parallel Columns on the same page. Edited by J. SCHOLEFIELD, M.A. Small Octavo. New Edition, with the Marginal References as arranged and revised by Dr SCRIVENER. Cloth, red edges. 7s. 6d.

GREEK AND ENGLISH TESTAMENT. THE STUDENT'S EDITION of the above, on *large writing paper*. 4to. 12s.

GREEK TESTAMENT, ex editione Stephani tertia, 1550. Small 8vo. 3s. 6d.

THE PARALLEL NEW TESTAMENT, GREEK AND ENGLISH, being the **Authorised Version** set forth in 1611 arranged in Parallel Columns with the **Revised Version** of 1881, and with the original Greek, as edited by F. H. A. SCRIVENER, M.A., D.C.L., LL.D. Crown 8vo. 12s. 6d. *The Revised Version is the Joint Property of the Universities of Cambridge and Oxford.*

THE BOOK OF ECCLESIASTES, with Notes and Introduction. By the Very Rev. E. H. PLUMPTRE, D.D., Dean of Wells. Large Paper Edition. Demy 8vo. 7s. 6d.

London: C. J. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

THE OLD TESTAMENT IN GREEK ACCORDING TO THE SEPTUAGINT. Edited by H. B. SWETE, D.D., Honorary Fellow of Gonville and Caius College. Vol. I. Genesis—IV Kings. Crown 8vo. 7s. 6d.

Volume II. By the same Editor.

[*In the Press.*]

"Der Zweck dieser Ausgabe, den ganzen in den erwähnten Hss. vorliegenden kritischen Stoff übersichtlich zusammenzustellen und dem Benützer das Nachschlagen in den Separatausgaben jener Codices zu ersparen, ist hier in compendiösester Weise vortrefflich erreicht. Bezüglich der Klarheit, Schönheit und Correctheit des Drucks gebührt der Ausgabe das höchste Lob. Da zugleich der Preis sehr niedrig gestellt ist, so ist zu hoffen und zu wünschen, dass sie auch außerhalb des englischen Sprachkreises ihre Verbreitung finden werde.

Bezüglich der Accente und Spiritus der Eigennamen sind die Herausg. ihre eigenen Wege gegangen."—*Deutsche Literaturzeitung.*

"The Edition has been executed in the very best style of Cambridge accuracy, which has no superior anywhere, and this is enough to put it at the head of the list of editions for manual use."—*Academy.*

"An edition, which for ordinary purposes will probably henceforth be that in use by readers of the Septuagint."—*Guardian.*

THE BOOK OF PSALMS IN GREEK ACCORDING TO THE SEPTUAGINT, being a portion of Vol. II. of the above. Crown 8vo. 2s. 6d.

THE GOSPEL ACCORDING TO ST MATTHEW in Anglo-Saxon and Northumbrian Versions, synoptically arranged: with Collations exhibiting all the Readings of all the MSS. Edited by the Rev. W. W. SKEAT, Litt.D., Elrington and Bosworth Professor of Anglo-Saxon. **New Edition.** Demy 4to. 10s.

"By the publication of the present volume Prof. Skeat has brought to its conclusion a work planned more than a half century ago by the late J. M. Kemble... Students of English have every reason to be grateful to Prof. Skeat

for the scholarly and accurate way in which he has performed his laborious task. Thanks to him we now possess a reliable edition of all the existing MSS. of the old English Gospels."—*Academy.*

THE GOSPEL ACCORDING TO ST MARK, uniform with the preceding, by the same Editor. Demy 4to. 10s.

THE GOSPEL ACCORDING TO ST LUKE, uniform with the preceding, by the same Editor. Demy 4to. 10s.

THE GOSPEL ACCORDING TO ST JOHN, uniform with the preceding, by the same Editor. Demy 4to. 10s.

"The Gospel according to St John, in Anglo-Saxon and Northumbrian Versions: completes an undertaking designed and commenced by that distinguished scholar, J. M. Kemble, some forty years ago. Of the par-

ticular volume now before us, we can only say it is worthy of its two predecessors. We repeat that the service rendered to the study of Anglo-Saxon by this Synoptic collection cannot easily be overstated."—*Contemporary Review.*

THE FOUR GOSPELS (as above) bound in one volume, price 30s.

THE POINTED PRAYER BOOK, being the Book of Common Prayer with the Psalter or Psalms of David, pointed as they are to be sung or said in Churches. Royal 24mo. 1s. 6d.
The same in square 32mo. cloth. 6d.

THE CAMBRIDGE PSALTER, for the use of Choirs and Organists. Specially adapted for Congregations in which the "Cambridge Pointed Prayer Book" is used. Demy 8vo. cloth extra, 3s. 6d. cloth limp, cut flush. 2s. 6d.

London: C. J. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

THE PARAGRAPH PSALTER, arranged for the use of Choirs by BROOKE FOSS WESTCOTT, D.D., Regius Professor of Divinity in the University of Cambridge. Fcap. 4to. 5s.

The same in royal 32mo. Cloth 1s. Leather 1s. 6d.

THE MISSING FRAGMENT OF THE LATIN TRANSLATION OF THE FOURTH BOOK OF EZRA, discovered, and edited with an Introduction and Notes, and a facsimile of the MS., by ROBERT L. BENSLY, M.A., Lord Almoner's Professor of Arabic. Demy 4to. 10s.

"It has been said of this book that it has added a new chapter to the Bible, and, startling as the statement may at first sight appear, it is no exaggeration of the actual fact, if by the

Bible we understand that of the larger size which contains the Apocrypha, and if the Second Book of Esdras can be fairly called a part of the Apocrypha."—*Saturday Review*.

THE HARKLEAN VERSION OF THE EPISTLE TO THE HEBREWS, Chap. xi. 28—xiii. 25. Now edited for the first time with Introduction and Notes on this Version of the Epistle. By ROBERT L. BENSLY, M.A. Demy 8vo. 5s.

NOTITIA CODICIS QUATTUOR EVANGELIORUM Græci Membranacei viris doctis hucusque incogniti quem in museo suo asservat EDUARDUS REUSS Argentoratensis. 2s.

THE ORIGIN OF THE LEICESTER CODEX OF THE NEW TESTAMENT. By J. RENDEL HARRIS, M.A. With 3 plates. Demy 4to. 10s. 6d.

THE REST OF THE WORDS OF BARUCH: A Christian Apocalypse of the Year 136 A.D. The Text revised with an Introduction. By J. RENDEL HARRIS, M.A. Royal 8vo. 5s.

CODEx S. CEADDAE LATINUS. Evangelia SSS. Matthæi, Marci, Lucae ad cap. III. 9 complectens, circa septimum vel octavum saeculum scriptus, in Ecclesia Cathedrali Lichfieldiensi servatus. Cum codice versionis Vulgatae Amiatino contulit, prolegomena conscripsit, F. H. A. SCRIVENER, A.M., D.C.L., LL.D., With 3 plates. £1. 1s.

THEOLOGY—(ANCIENT).

THE GREEK LITURGIES. Chiefly from original Authorities. By C. A. SWAINSON, D.D., late Master of Christ's College, Cambridge. Crown 4to. Paper covers. 15s.

"Jeder folgende Forscher wird dankbar anerkennen, dass Swainson das Fundament zu einer historisch-kritischen Geschichte der

Griechischen Liturgien sicher gelegt hat."—ADOLPH HARNACK, *Theologische Literatur-Zeitung*.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

THEODORE OF MOPSUESTIA'S COMMENTARY ON THE MINOR EPISTLES OF S. PAUL. The Latin Version with the Greek Fragments, edited from the MSS. with Notes and an Introduction, by H. B. SWETE, D.D. In Two Volumes. Volume I., containing the Introduction, with Facsimiles of the MSS., and the Commentary upon Galatians—Colossians. Demy 8vo. 12s.

"It is the result of thorough, careful, and patient investigation of all the points bearing on the subject, and the results are presented with admirable good sense and modesty."—*Guardian*.

"In dem oben verzeichneten Buche liegt uns die erste Hälfte einer vollständigen, ebenso sorgfältig gearbeiteten wie schön ausgestatteten Ausgabe des Commentars mit ausführlichen Prolegomena und reichhaltigen kritischen und erläuternden Anmerkungen vor."—*Literarisches Centralblatt*.

"Auf Grund dieser Quellen ist der Text bei Swete mit musterhafter Akribie hergestellt. Aber auch sonst hat der Herausgeber mit unermüdlichem Fleisse und eingehend-

ster Sachkenntniss sein Werk mit allen denjenigen Zugaben ausgerüstet, welche bei einer solchen Text-Ausgabe nur irgend erwartet werden können. . . . Von den drei Haupt-handschriften . . . sind vortreffliche photographische Facsimile's beigegeben, wie überhaupt das ganze Werk von der *University Press* zu Cambridge mit bekannter Eleganz ausgestattet ist."—*Theologische Literaturzeitung*.

"Herrn Swete's Leistung ist eine so tüchtige dass wir das Werk in keinen besseren Händen wissen möchten, und mit den sichersten Erwartungen auf das Gelingen der Fortsetzung entgegen sehen."—*Göttingische gelehrte Anzeigen* (Sept. 1881).

VOLUME II., containing the Commentary on 1 Thessalonians—Philemon, Appendices and Indices. 12s.

"Eine Ausgabe . . . für welche alle zugänglichen Hülfsmittel in musterhafter Weise benutzt wurden . . . eine reife Frucht siebenjährigen Fleisses."—*Theologische Literaturzeitung*

(Sept. 23, 1882).

"Mit derselben Sorgfalt bearbeitet die wir bei dem ersten Theile gerühmt haben."—*Literarisches Centralblatt* (July 29, 1882).

SAYINGS OF THE JEWISH FATHERS, comprising Pirke Aboth and Pereq R. Meir in Hebrew and English, with Critical and Illustrative Notes. By CHARLES TAYLOR, D.D., Master of St John's College, Cambridge. Demy 8vo. 10s.

"The 'Masseketh Aboth' stands at the head of Hebrew non-canonical writings. It is of ancient date, claiming to contain the dicta of teachers who flourished from B.C. 200 to the same year of our era. Mr Taylor's explanatory and illustrative commentary is very full and satisfactory."—*Spectator*.

"A careful and thorough edition which does credit to English scholarship, of a short treatise from the Mishna, containing a series of sentences or maxims ascribed mostly to Jewish teachers immediately preceding, or immediately following the Christian era. . . ."—*Contemporary Review*.

A COLLATION OF THE ATHOS CODEX OF THE SHEPHERD OF HERMAS. Together with an Introduction by SPYR. P. LAMBROS, PH. D., translated and edited with a Preface and Appendices by J. ARMITAGE ROBINSON, M.A., Fellow and Dean of Christ's College, Cambridge. Demy 8vo. 3s. 6d.

THE PALESTINIAN MISHNA. By W. H. LOWE, M.A., Lecturer in Hebrew at Christ's College, Cambridge. Royal 8vo. 21s.

SANCTI IRENÆI EPISCOPI LUGDUNENSIS libros quinque adversus Hæreses, versione Latina cum Codicibus Claromontano ac Arundeliano denuo collata, præmissa de placitis Gnosticatorum prolusione, fragmenta necnon Græce, Syriace, Armeniace, commentatione perpetua et indicibus variis edidit W. WIGAN HARVEY, S.T.B. Collegii Regalis olim Socius. 2 Vols. 8vo. 18s.

M. MINUCII FELICIS OCTAVIUS. The text revised from the original MS., with an English Commentary, Analysis, Introduction, and Copious Indices. Edited by H. A. HOLDEN, LL.D. Examiner in Greek to the University of London. Crown 8vo. 7s. 6d.

London: C. J. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

- THEOPHILI EPISCOPI ANTIOCHENSIS LIBRI
TRES AD AUTOLYCUM edidit, Prolegomenis Versione Notulis
Indicibus instruxit G. G. HUMPHRY, S.T.B. Post 8vo. 5s.
- THEOPHYLACTI IN EVANGELIUM S. MATTHÆI
COMMENTARIUS, edited by W. G. HUMPHRY, B.D. Prebendary
of St Paul's, late Fellow of Trinity College. Demy 8vo. 7s. 6d.
- TERTULLIANUS DE CORONA MILITIS, DE SPEC-
TACULIS, DE IDOLOLATRIA, with Analysis and English Notes,
by GEORGE CURREY, D.D. Preacher at the Charter House, late
Fellow and Tutor of St John's College. Crown 8vo. 5s.
- FRAGMENTS OF PHILO AND JOSEPHUS. Newly
edited by J. RENDEL HARRIS, M.A., Fellow of Clare College,
Cambridge. With two Facsimiles. Demy 4to. 12s. 6d.
- THE TEACHING OF THE APOSTLES. Newly edited,
with Facsimile Text and Commentary, by J. RENDEL HARRIS, M.A.
Demy 4to. £1. 1s.

THEOLOGY—(ENGLISH).

- WORKS OF ISAAC BARROW, compared with the Ori-
ginal MSS., enlarged with Materials hitherto unpublished. A new
Edition, by A. NAPIER, M.A. 9 Vols. Demy 8vo. £3. 3s.
- TREATISE OF THE POPE'S SUPREMACY, and a
Discourse concerning the Unity of the Church, by ISAAC BARROW.
Demy 8vo. 7s. 6d.
- PEARSON'S EXPOSITION OF THE CREED, edited
by TEMPLE CHEVALLIER, B.D. New Edition. Revised by R. SINKER,
B.D., Librarian of Trinity College. Demy 8vo. 12s.
- "A new edition of Bishop Pearson's famous
work *On the Creed* has just been issued by the
Cambridge University Press. It is the well-
known edition of Temple Chevallier, thoroughly
overhauled by the Rev. R. Sinker, of Trinity
College. Altogether this appears to be the
most complete and convenient edition as yet
published of a work which has long been re-
cognised in all quarters as a standard one."—
Guardian.
- AN ANALYSIS OF THE EXPOSITION OF THE
CREED written by the Right Rev. JOHN PEARSON, D.D. late Lord
Bishop of Chester, by W. H. MILL, D.D. Demy 8vo. 5s.
- WHEATLY ON THE COMMON PRAYER, edited by
G. E. CORRIE, D.D. late Master of Jesus College. Demy 8vo. 7s. 6d.
- TWO FORMS OF PRAYER OF THE TIME OF QUEEN
ELIZABETH. Now First Reprinted. Demy 8vo. 6d.
- CÆSAR MORGAN'S INVESTIGATION OF THE
TRINITY OF PLATO, and of Philo Judæus, and of the effects
which an attachment to their writings had upon the principles and
reasonings of the Fathers of the Christian Church. Revised by H. A.
HOLDEN, LL.D. Crown 8vo. 4s.

SELECT DISCOURSES, by JOHN SMITH, late Fellow of Queens' College, Cambridge. Edited by H. G. WILLIAMS, B.D. late Professor of Arabic. Royal 8vo. 7s. 6d.

"The 'Select Discourses' of John Smith, collected and published from his papers after his death, are, in my opinion, much the most considerable work left to us by this Cambridge School [the Cambridge Platonists]. They have a right to a place in English literary history."—Mr MATTHEW ARNOLD, in the *Contemporary Review*.

"Of all the products of the Cambridge School, the 'Select Discourses' are perhaps the highest, as they are the most accessible and the most widely appreciated...and indeed

no spiritually thoughtful mind can read them unmoved. They carry us so directly into an atmosphere of divine philosophy, luminous with the richest lights of meditative genius... He was one of those rare thinkers in whom largeness of view, and depth, and wealth of poetic and speculative insight, only served to evoke more fully the religious spirit, and while he drew the mould of his thought from Plotinus, he vivified the substance of it from St Paul."—Principal TULLOCH, *Rational Theology in England in the 17th Century*.

THE HOMILIES, with Various Readings, and the Quotations from the Fathers given at length in the Original Languages. Edited by the late G. E. CORRIE, D.D. Demy 8vo. 7s. 6d.

DE OBLIGATIONE CONSCIENTIÆ PRÆLECTIONES decem Oxonii in Schola Theologica habitæ a ROBERTO SANDERSON, SS. Theologiæ ibidem Professore Regio. With English Notes, including an abridged Translation, by W. WHEWELL, D.D. late Master of Trinity College. Demy 8vo. 7s. 6d.

ARCHBISHOP USHER'S ANSWER TO A JESUIT, with other Tracts on Popery. Edited by J. SCHOLEFIELD, M.A. late Regius Professor of Greek in the University. Demy 8vo. 7s. 6d.

WILSON'S ILLUSTRATION OF THE METHOD OF explaining the New Testament, by the early opinions of Jews and Christians concerning Christ. Edited by T. TURTON, D.D. 8vo. 5s.

LECTURES ON DIVINITY delivered in the University of Cambridge, by JOHN HEY, D.D. Third Edition, revised by T. TURTON, D.D. late Lord Bishop of Ely. 2 vols. Demy 8vo. 15s.

S. AUSTIN AND HIS PLACE IN THE HISTORY OF CHRISTIAN THOUGHT. Being the Hulsean Lectures for 1885. By W. CUNNINGHAM, D.D. Demy 8vo. Buckram, 12s. 6d.

THE GOSPEL HISTORY OF OUR LORD JESUS CHRIST IN THE LANGUAGE OF THE REVISED VERSION, arranged in a Connected Narrative, especially for the use of Teachers and Preachers. By Rev. C. C. JAMES, M.A., Rector of Wortham, Suffolk, and late Fellow of King's College. Crown 8vo. [Nearly ready.]

ARABIC, SANSKRIT, SYRIAC, &c.

THE DIVYÂVADÂNA, a Collection of Early Buddhist Legends, now first edited from the Nepalese Sanskrit MSS. in Cambridge and Paris. By E. B. COWELL, M.A., Professor of Sanskrit in the University of Cambridge, and R. A. NEIL, M.A., Fellow and Lecturer of Pembroke College. Demy 8vo. 18s.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

POEMS OF BEHA ED DIN ZOHEIR OF EGYPT.

With a Metrical Translation, Notes and Introduction, by E. H. PALMER, M.A., Barrister-at-Law of the Middle Temple, late Lord Almoner's Professor of Arabic, formerly Fellow of St John's College, Cambridge. 2 vols. Crown 4to.

Vol. I. The ARABIC TEXT. 10s. 6d.

Vol. II. ENGLISH TRANSLATION. 10s. 6d.

"We have no hesitation in saying that in both Prof. Palmer has made an addition to Oriental literature for which scholars should be grateful; and that, while his knowledge of Arabic is a sufficient guarantee for his mastery of the original, his English compositions are distinguished by versatility, command of language, rhythmical cadence, and, as we have

remarked, by not unskilful imitations of the styles of several of our own favourite poets, living and dead."—*Saturday Review*.

"This sumptuous edition of the poems of Behá-ed-dín Zoheir is a very welcome addition to the small series of Eastern poets accessible to readers who are not Orientalists."—*Academy*.

THE CHRONICLE OF JOSHUA THE STYLITE, com-

posed in Syriac A.D. 507, with an English translation and notes, by the late W. WRIGHT, LL.D., Professor of Arabic. Demy 8vo. 10s. 6d.

"Die lehrreiche kleine Chronik Josuas hat nach Assemani und Martin in Wright einen dritten Bearbeiter gefunden, der sich um die Emendation des Textes wie um die Erklärung der Realien wesentlich verdient gemacht hat . . . Ws. Josua-Ausgabe ist eine sehr dankenswerte Gabe und besonders empfehlenswert als

ein Lehrmittel für den syrischen Unterricht; es erscheint auch gerade zur rechten Zeit, da die zweite Ausgabe von Roedigers syrischer Chrestomathie im Buchhandel vollständig vergriffen und diejenige von Kirsch-Bernstein nur noch in wenigen Exemplaren vorhanden ist."—*Deutsche Literaturzeitung*.

KALĪLAH AND DIMNAH, OR, THE FABLES OF

BIDPAI; being an account of their literary history, together with an English Translation of the same, with Notes, by I. G. N. KEITH-FALCONER, M.A., late Lord Almoner's Professor of Arabic in the University of Cambridge. Demy 8vo. 7s. 6d.

NALOPĀKHYĀNAM, OR, THE TALE OF NALA; containing the Sanskrit Text in Roman Characters, followed by a Vocabulary and a sketch of Sanskrit Grammar. By the late Rev. THOMAS JARRETT, M.A. Trinity College, Regius Professor of Hebrew. Demy 8vo. 10s.

NOTES ON THE TALE OF NALA, for the use of Classical Students, by J. PEILE, Litt.D., Master of Christ's College. Demy 8vo. 12s.

CATALOGUE OF THE BUDDHIST SANSKRIT

MANUSCRIPTS in the University Library, Cambridge. Edited by C. BENDALL, M.A., Fellow of Gonville and Caius College. Demy 8vo. 12s.

"It is unnecessary to state how the compilation of the present catalogue came to be placed in Mr Bendall's hands; from the character of his work it is evident the selection was judicious, and we may fairly congratulate

those concerned in it on the result . . . Mr Bendall has entitled himself to the thanks of all Oriental scholars, and we hope he may have before him a long course of successful labour in the field he has chosen."—*Athenaeum*.

THE HISTORY OF ALEXANDER THE GREAT,

being the Syriac version of the Pseudo-Callisthenes. Edited from Five Manuscripts, with an English Translation and Notes, by E. A. W. BUDGE, M.A., Assistant in the Department of Egyptian Antiquities, British Museum. Demy 8vo. 25s. (*The Edition is limited to 250 copies.*)

London: C. J. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

GREEK AND LATIN CLASSICS, &c.

SOPHOCLES: The Plays and Fragments, with Critical Notes, Commentary, and Translation in English Prose, by R. C. JEBB, Litt.D., LL.D., Regius Professor of Greek in the University of Cambridge.

Part I. Oedipus Tyrannus. Demy 8vo. *New Edition.* 12s. 6d.

Part II. Oedipus Coloneus. Demy 8vo. *New Edition.* 12s. 6d.

Part III. Antigone. Demy 8vo. 12s. 6d.

Part IV. Philoctetes.

[*In the Press.*]

"Of his explanatory and critical notes we can only speak with admiration. Thorough scholarship combines with taste, erudition, and boundless industry to make this first volume a pattern of editing. The work is made complete by a prose translation, upon pages alternating with the text, of which we may say shortly that it displays sound judgment and taste, without sacrificing precision to poetry of expression."—*The Times*.

"Professor Jebb's edition of Sophocles is already so fully established, and has received such appreciation in these columns and elsewhere, that we have judged this third volume when we have said that it is of a piece with the others. The whole edition so far exhibits perhaps the most complete and elaborate editorial work which has ever appeared."—*Saturday Review*.

"Prof. Jebb's keen and profound sympathy, not only with Sophocles and all the best of ancient Hellenic life and thought, but also with modern European culture, constitutes him an ideal interpreter between the ancient writer and the modern reader."—*Athenæum*.

"It would be difficult to praise this third instalment of Professor Jebb's unequalled edition of Sophocles too warmly, and it is almost a work of supererogation to praise it at all. It is equal, at least, and perhaps superior, in merit, to either of his previous instalments; and when this is said, all is said. Yet we cannot refrain from formally recognising once more the consummate Greek scholarship of the editor, and from once more doing grateful homage to his masterly tact and literary skill, and to his unwearied and marvellous industry."—*Spectator*.

AESCHYLI FABULAE.—ΙΚΕΤΙΔΕΣ ΧΟΗΦΟΡΟΙ IN LIBRO MEDICEO MENDOSE SCRIPTAE EX VV. DD. CONIECTURIS EMENDATIUS EDITAE cum Scholiis Graecis et brevi adnotatione critica, curante F. A. PALEY, M.A., LL.D. Demy 8vo. 7s. 6d.

THE AGAMEMNON OF AESCHYLUS. With a Translation in English Rhythm, and Notes Critical and Explanatory. **New Edition Revised.** By the late BENJAMIN HALL KENNEDY, D.D., Regius Professor of Greek. Crown 8vo. 6s.

"One of the best editions of the masterpiece of Greek tragedy."—*Athenæum*.

THE THEÆTETUS OF PLATO with a Translation and Notes by the same Editor. Crown 8vo. 7s. 6d.

ARISTOTLE.—ΠΕΡΙ ΨΥΧΗΣ. ARISTOTLE'S PSYCHOLOGY, in Greek and English, with Introduction and Notes, by EDWIN WALLACE, M.A., late Fellow and Tutor of Worcester College, Oxford. Demy 8vo. 18s.

"The notes are exactly what such notes ought to be,—helps to the student, not mere displays of learning. By far the more valuable parts of the notes are neither critical nor literary, but philosophical and expository of the thought, and of the connection of thought, in the treatise itself. In this relation the notes are invaluable. Of the translation, it may be said that an English reader may fairly master by means of it this great treatise of Aristotle."—*Spectator*.

"Wallace's Bearbeitung der Aristotelischen Psychologie ist das Werk eines denkenden und in allen Schriften des Aristoteles und grösstenteils auch in der neueren Litteratur zu denselben belesenen Mannes... Der schwächste Teil der Arbeit ist der kritische... Aber in allen diesen Dingen liegt auch nach der Absicht des Verfassers nicht der Schwerpunkt seiner Arbeit, sondern."—Prof. Susemihl in *Philologische Wochenschrift*.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse
Ave Maria Lane.

ARISTOTLE.—ΠΕΡΙ ΔΙΚΑΙΟΣΤΗΗΣ. THE FIFTH BOOK OF THE NICOMACHEAN ETHICS OF ARISTOTLE. Edited by HENRY JACKSON, Litt.D., Fellow of Trinity College, Cambridge. Demy 8vo. 6s.

"It is not too much to say that some of the points he discusses have never had so much light thrown upon them before. . . . Scholars

will hope that this is not the only portion of the Aristotelian writings which he is likely to edit."—*Athenæum*.

ARISTOTLE. THE RHETORIC. With a Commentary by the late E. M. COPE, Fellow of Trinity College, Cambridge, revised and edited by J. E. SANDYS, Litt.D. With a biographical Memoir by the late H. A. J. MUNRO, Litt.D. 3 Vols., Demy 8vo.

Now reduced to 21s. (originally published at 31s. 6d.)

"This work is in many ways creditable to the University of Cambridge. If an English student wishes to have a full conception of what is contained in the *Rhetoric* of Aristotle, to Mr Cope's edition he must go."—*Academy*.

"Mr Sandys has performed his arduous duties with marked ability and admirable tact. . . . In every part of his work—revising, supplementing, and completing—he has done exceedingly well."—*Examiner*.

PINDAR. OLYMPIAN AND PYTHIAN ODES. With Notes Explanatory and Critical, Introductions and Introductory Essays. Edited by C. A. M. FENNELL, Litt.D., late Fellow of Jesus College. Crown 8vo. 9s.

"Mr Fennell deserves the thanks of all classical students for his careful and scholarly edition of the Olympian and Pythian odes. He brings to his task the necessary enthusiasm for

his author, great industry, a sound judgment, and, in particular, copious and minute learning in comparative philology."—*Athenæum*.

— THE ISTHMIAN AND NEMEAN ODES. By the same Editor. Crown 8vo. 9s.

" . . . As a handy and instructive edition of a difficult classic no work of recent years surpasses Mr Fennell's 'Pindar.'"—*Athenæum*.

"This work is in no way inferior to the previous volume. The commentary affords

valuable help to the study of the most difficult of Greek authors, and is enriched with notes on points of scholarship and etymology which could only have been written by a scholar of very high attainments."—*Saturday Review*.

DEMOSTHENES. PRIVATE ORATIONS OF, with Introductions and English Notes, by the late F. A. PALEY, M.A. and J. E. SANDYS, Litt.D. Fellow and Tutor of St John's College, and Public Orator in the University of Cambridge.

PART I. Contra Phormionem, Lacritum, Pantaenetum, Boeotum de Nomine, Boeotum de Dote, Dionysodorum. **New Edition.** Crown 8vo. 6s.

"Mr Paley's scholarship is sound and accurate, his experience of editing wide, and if he is content to devote his learning and abilities to the production of such manuals as these, they will be received with gratitude throughout the higher schools of the country. Mr Sandys is deeply read in the German

literature which bears upon his author, and the elucidation of matters of daily life, in the delineation of which Demosthenes is so rich, obtains full justice at his hands. . . . We hope this edition may lead the way to a more general study of these speeches in schools than has hitherto been possible."—*Academy*.

PART II. Pro Phormione, Contra Stephanum I. II.; Nicostratum, Cononem, Calliclem. **New Edition.** Crown 8vo. 7s. 6d.

"It is long since we have come upon a work evincing more pains, scholarship, and varied research and illustration than Mr Sandys's contribution to the 'Private Orations of De-

mosthenes'."—*Saturday Review*.

" . . . the edition reflects credit on Cambridge scholarship, and ought to be extensively used."—*Athenæum*.

DEMOSTHENES. LEPTINES. With Introductory Essay and Critical and Explanatory Notes. Edited by J. E. SANDYS, Litt.D. Demy 8vo. [Nearly ready.]

London: C. F. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

DEMOSTHENES AGAINST ANDROTION AND AGAINST TIMOCRATES, with Introductions and English Commentary, by WILLIAM WAYTE, M.A., late Professor of Greek, University College, London. Crown 8vo. 7s. 6d.

"These speeches are highly interesting, as illustrating Attic Law, as that law was influenced by the exigences of politics . . . As vigorous examples of the great orator's style,

they are worthy of all admiration . . . Besides a most lucid and interesting introduction, Mr Wayte has given the student effective help in his running commentary."—*Spectator*.

PLATO'S PHÆDO, literally translated, by the late E. M. COPE, Fellow of Trinity College, Cambridge, revised by HENRY JACKSON, Litt.D., Fellow of Trinity College. Demy 8vo. 5s.

P. VERGILI MARONIS OPERA, cum Prolegomenis et Commentario Critico edidit B. H. KENNEDY, S.T.P., Extra Fcap. 8vo. 3s. 6d.

THE BACCHAE OF EURIPIDES. With Introduction, Critical Notes, and Archæological Illustrations, by J. E. SANDYS, Litt.D. New and Enlarged Edition. Crown 8vo. 12s. 6d.

"Of the present edition of the *Bacchæ* by Mr Sandys we may safely say that never before has a Greek play, in England at least, had fuller justice done to its criticism, interpretation, and archæological illustration, whether for the young student or the more advanced scholar. The Cambridge Public Orator may be said to have taken the lead in issuing a complete edition of a Greek play, which is destined perhaps to gain redoubled favour now that the study of ancient monuments has been applied to its illustration."—*Saturday Review*.

"The volume is interspersed with well-executed woodcuts, and its general attractiveness of form reflects great credit on the University Press. In the notes Mr Sandys has more than sustained his well-earned reputation as a careful and learned editor, and shows considerable advance in freedom and lightness of style. . . . Under such circumstances it is superfluous to say that for the purposes of teachers and advanced students this handsome edition far surpasses all its predecessors."—*Athenæum*.

THE TYPES OF GREEK COINS. By PERCY GARDNER, Litt. D., F.S.A. With 16 Autotype plates, containing photographs of Coins of all parts of the Greek World. Impl. 4to. Cloth extra, £1. 11s. 6d.; Roxburgh (Morocco back), £2. 2s.

"Professor Gardner's book is written with such lucidity and in a manner so straightforward that it may well win converts, and it may

be distinctly recommended to that omnivorous class of readers—'men in the schools'."—*Saturday Review*.

ESSAYS ON THE ART OF PHEIDIAS. By C. WALDSTEIN, Litt. D., Phil. D., Reader in Classical Archæology in the University of Cambridge. Royal 8vo. With numerous Illustrations. 16 Plates. Buckram, 30s.

"His book will be universally welcomed as a very valuable contribution towards a more thorough knowledge of the style of Pheidias."—*The Academy*.

"'Essays on the Art of Pheidias' form an extremely valuable and important piece of work. . . . Taking it for the illustrations alone, it is an exceedingly fascinating book."—*Times*.

AN INTRODUCTION TO GREEK EPIGRAPHY. Part I. The Archaic Inscriptions and the Greek Alphabet by E. S. ROBERTS, M.A., Fellow and Tutor of Gonville and Caius College. Demy 8vo. With illustrations. 18s.

"We will say at once that Mr Roberts appears to have done his work very well. The book is clearly and conveniently arranged. The inscriptions are naturally divided according to the places to which they belong. Under each head are given illustrations sufficient to show the characteristics of the writing, one copy in letters of the original form (sometimes a facsimile) being followed by another in the usual cursive. References, which must have cost great labour, are given to the scattered

notices bearing on each document. Explanatory remarks either accompany the text or are added in an appendix. To the whole is prefixed a sketch of the history of the alphabet up to the terminal date. At the end the result is resumed in general tables of all the alphabets, classified according to their connexions; and a separate table illustrates the alphabet of Athens. The volume contains about five hundred inscriptions, and forms a moderate octavo of about four hundred pages."—*Saturday Review*.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

M. TULLI CICERONIS AD M. BRUTUM ORATOR.

A revised text edited with Introductory Essays and with critical and explanatory notes, by J. E. SANDYS, Litt.D. Demy 8vo. 16s.

"This volume, which is adorned with several good woodcuts, forms a handsome and welcome addition to the Cambridge editions of Cicero's works."—*Athenæum*.

"A model edition."—*Spectator*.

"The commentary is in every way worthy of the editor's high reputation."—*Academy*.

M. TULLI CICERONIS DE FINIBUS BONORUM ET MALORUM LIBRI QUINQUE. The text revised and explained; with a Translation by JAMES S. REID, Litt.D., Fellow and Tutor of Gonville and Caius College. 3 Vols. [In the Press.]

VOL. III. Containing the Translation. Demy 8vo. 8s.

M. T. CICERONIS DE OFFICIIS LIBRI TRES, with Marginal Analysis, English Commentary, and copious Indices, by H. A. HOLDEN, LL.D. Sixth Edition, Revised and Enlarged. Cr. 8vo. 9s.

"Few editions of a classic have found so much favour as Dr Holden's *De Officiis*, and the present revision (sixth edition) makes the

position of the work secure."—*American Journal of Philology*.

M. T. CICERONIS DE OFFICIIS LIBER TERTIUS, with Introduction, Analysis and Commentary, by H. A. HOLDEN, LL.D. Crown 8vo. 2s.

M. TULLI CICERONIS PRO C. RABIRIO [PERDVLLIONIS REO] ORATIO AD QVIRITES, with Notes, Introduction and Appendices by W. E. HEITLAND, M.A., Fellow and Tutor of St John's College, Cambridge. Demy 8vo. 7s. 6d.

M. TULLII CICERONIS DE NATURA DEORUM Libri Tres, with Introduction and Commentary by JOSEPH B. MAYOR, M.A., together with a new collation of several of the English MSS. by J. H. SWAINSON, M.A.

Vol. I. Demy 8vo. 10s. 6d. Vol. II. 12s. 6d. Vol. III. 10s.

"Such editions as that of which Prof. Mayor has given us the first instalment will doubtless do much to remedy this undeserved neglect. It is one on which great pains and much learning have evidently been expended, and is in every

way admirably suited to meet the needs of the student . . . The notes of the editor are all that could be expected from his well-known learning and scholarship."—*Academy*.

See also Pitt Press Series, pp. 30—34.

MATHEMATICS, PHYSICAL SCIENCE, &c.

MATHEMATICAL AND PHYSICAL PAPERS. By Sir W. THOMSON, LL.D., D.C.L., F.R.S., Professor of Natural Philosophy in the University of Glasgow. Collected from different Scientific Periodicals from May 1841, to the present time. Vol. I. Demy 8vo. 18s. Vol. II. 15s. [Volume III. In the Press.]

MATHEMATICAL AND PHYSICAL PAPERS, by Sir G. G. STOKES, Sc.D., LL.D., F.R.S., Lucasian Professor of Mathematics in the University of Cambridge. Reprinted from the Original Journals and Transactions, with Additional Notes by the Author. Vol. I. Demy 8vo. 15s. Vol. II. 15s. [Vol. III. In the Press.]

London: C. J. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

- A HISTORY OF THE THEORY OF ELASTICITY AND OF THE STRENGTH OF MATERIALS, from Galilei to the present time. VOL. I. Galilei to Saint-Venant, 1639-1850. By the late I. TODHUNTER, Sc.D., F.R.S., edited and completed by Professor KARL PEARSON, M.A. Demy 8vo. 25s.
- Vol. II. By the same Editor. *[In the Press.]*
- THE ELASTICAL RESEARCHES OF BARRÉ DE SAINT-VENANT (Extract from Vol. II. of TODHUNTER'S History of the Theory of Elasticity), edited by Professor KARL PEARSON, M.A. Demy 8vo. 9s.
- A TREATISE ON GEOMETRICAL OPTICS. By R. S. HEATH, M.A., Professor of Mathematics in Mason Science College, Birmingham. Demy 8vo. 12s. 6d.
- AN ELEMENTARY TREATISE ON GEOMETRICAL OPTICS. By R. S. HEATH, M.A. Crown 8vo. 5s.
- A TREATISE ON ELEMENTARY DYNAMICS. By S. L. LONEY, M.A., Fellow of Sidney Sussex College. Crown 8vo. 7s. 6d.
- A TREATISE ON PLANE TRIGONOMETRY. By E. W. HOBSON, M.A., Fellow and Lecturer of Christ's College, Cambridge. Demy 8vo. *[In the Press.]*
- CATALOGUE OF SCIENTIFIC PAPERS COMPILED BY THE ROYAL SOCIETY OF LONDON: Vols. 1-6 for the years 1800-1863, Royal 4to. cloth (vol. 1 in half morocco) £4 (net); half morocco £5. 5s. (net). Vols. 7-8 for the years 1864-1873, cloth £1. 11s. 6d. (net); half morocco £2. 5s. (net). Single volumes cloth 20s. or half-morocco 28s. (net). New Series for the years 1874-1883. *[In the Press.]*
- THE COLLECTED MATHEMATICAL PAPERS OF ARTHUR CAYLEY, Sc.D., F.R.S., Sadlerian Professor of Pure Mathematics in the University of Cambridge. Demy 4to. 10 vols. Vols. I. and II. 25s. each. *[Vol. III. In the Press.]*
- THE SCIENTIFIC PAPERS OF THE LATE PROF. J. CLERK MAXWELL. Edited by W. D. NIVEN, M.A. In 2 vols. Royal 4to. *[Nearly ready.]*
- A HISTORY OF THE STUDY OF MATHEMATICS AT CAMBRIDGE. By W. W. ROUSE BALL, M.A., Fellow and Lecturer on Mathematics of Trinity College, Cambridge. Crown 8vo. 6s.
- A TREATISE ON ANALYTICAL STATICS, by E. J. ROUTH, Sc.D., F.R.S., Fellow of the University of London, Honorary Fellow of St Peter's College, Cambridge. *[In the Press.]*
- A CATALOGUE OF THE PORTSMOUTH COLLECTION OF BOOKS AND PAPERS written by or belonging to SIR ISAAC NEWTON. Demy 8vo. 5s.

A TREATISE ON NATURAL PHILOSOPHY. By Sir W. THOMSON, LL.D., D.C.L., F.R.S., and P. G. TAIT, M.A.
Part I. Demy 8vo. 16s. **Part II.** Demy 8vo. 18s.

ELEMENTS OF NATURAL PHILOSOPHY. By Professors Sir W. THOMSON and P. G. TAIT. Demy 8vo. 9s.

AN ATTEMPT TO TEST THE THEORIES OF CAPILLARY ACTION, by FRANCIS BASHFORTH, B.D., and J. C. ADAMS, M.A., F.R.S. Demy 4to. £1. 1s.

A TREATISE ON THE THEORY OF DETERMINANTS and their applications in Analysis and Geometry, by R. F. SCOTT, M.A., Fellow of St John's College. Demy 8vo. 12s.

HYDRODYNAMICS, a Treatise on the Mathematical Theory of the Motion of Fluids, by H. LAMB, M.A. Demy 8vo. 12s.

THE ANALYTICAL THEORY OF HEAT, by JOSEPH FOURIER. Translated, with Notes, by A. FREEMAN, M.A., formerly Fellow of St John's College, Cambridge. Demy 8vo. 12s.

PRACTICAL WORK AT THE CAVENDISH LABORATORY. HEAT. Edited by W. N. SHAW, M.A. Demy 8vo. 3s.

THE ELECTRICAL RESEARCHES OF THE Hon. H. CAVENDISH, F.R.S. Written between 1771 and 1781. Edited from the original MSS. in the possession of the Duke of Devonshire, K. G., by the late J. CLERK MAXWELL, F.R.S. Demy 8vo. 18s.

AN ELEMENTARY TREATISE ON QUATERNIONS. By P. G. TAIT, M.A. 3rd Edition. Enlarged. Demy 8vo. 18s.

COUNTERPOINT. A Practical Course of Study, by the late Professor Sir G. A. MACFARREN, M.A., Mus. Doc. New Edition, revised. Crown 4to. 7s. 6d.

A TREATISE ON THE GENERAL PRINCIPLES OF CHEMISTRY, by M. M. PATTISON MUIR, M.A. **Second Edition.** Demy 8vo. 15s.

"The value of the book as a digest of the historical developments of chemical thought is immense."—*Academy*.

"Theoretical Chemistry has moved so rapidly of late years that most of our ordinary text books have been left far behind. German students, to be sure, possess an excellent guide to the present state of the science in 'Die Modernen Theorien der Chemie' of Prof.

Lothar Meyer; but in this country the student has had to content himself with such works as Dr Tilden's 'Introduction to Chemical Philosophy', an admirable book in its way, but rather slender. Mr Pattison Muir having aimed at a more comprehensive scheme, has produced a systematic treatise on the principles of chemical philosophy which stands far in advance of any kindred work in our language."—*Athenaeum*.

ELEMENTARY CHEMISTRY. By M. M. PATTISON MUIR, M.A., and CHARLES SLATER, M.A., M.B. Crown 8vo. 4s. 6d.

PRACTICAL CHEMISTRY. A Course of Laboratory Work. By M. M. PATTISON MUIR, M.A., and D. J. CARNEGIE, B.A. Crown 8vo. 3s.

NOTES ON QUALITATIVE ANALYSIS. Concise and Explanatory. By H. J. H. FENTON, M.A., F.I.C., Demonstrator of Chemistry in the University of Cambridge. Cr. 4to. *New Edition.* 6s.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse,
Ave Maria Lane.

LECTURES ON THE PHYSIOLOGY OF PLANTS.

by S. H. VINES, Sc.D., Professor of Botany in the University of Oxford. Demy 8vo. With Illustrations. 21s.

"To say that Dr Vines' book is a most valuable addition to our own botanical literature is but a narrow meed of praise: it is a work which will take its place as cosmopolitan: no more clear or concise discussion of the difficult chemistry of metabolism has appeared....

In erudition it stands alone among English books, and will compare favourably with any foreign competitors."—*Nature*.

"The work forms an important contribution to the literature of the subject....It will be eagerly welcomed by all students."—*Academy*.

A SHORT HISTORY OF GREEK MATHEMATICS.

By J. GOW, Litt.D., Fellow of Trinity College. Demy 8vo. 10s. 6d.

DIOPHANTOS OF ALEXANDRIA; a Study in the

History of Greek Algebra. By T. L. HEATH, M.A., Fellow of Trinity College, Cambridge. Demy 8vo. 7s. 6d.

"This study in the history of Greek Algebra is an exceedingly valuable contribution to the history of mathematics."—*Academy*.

"The most thorough account extant of Diophantus's place, work, and critics."—*Athenæum*.

THE MATHEMATICAL WORKS OF ISAAC BAR-

ROW, D.D. Edited by W. WHEWELL, D.D. Demy 8vo. 7s. 6d.

THE FOSSILS AND PALÆONTOLOGICAL AFFIN-

ITIES OF THE NEOCOMIAN DEPOSITS OF UPWARE AND BRICKHILL with Plates, being the Sedgwick Prize Essay for 1879. By the late W. KEEPING, M.A. Demy 8vo. 10s. 6d.

THE BALA VOLCANIC SERIES OF CAERNAR-

VONSHIRE AND ASSOCIATED ROCKS, being the Sedgwick Prize Essay for 1888 by A. HARKER, M.A., F.R.S., Fellow of St John's College. Demy 8vo. 7s. 6d.

A CATALOGUE OF BOOKS AND PAPERS ON PRO-

TOZOA, CŒLENTERATES, WORMS, and certain smaller groups of animals, published during the years 1861—1883, by D'ARCY W. THOMPSON, M.A. Demy 8vo. 12s. 6d.

ASTRONOMICAL OBSERVATIONS made at the Obser-

vatory of Cambridge by the late Rev. J. CHALLIS, M.A. from 1846 to 1860.

ASTRONOMICAL OBSERVATIONS from 1861 to 1865.

Vol. XXI. Royal 4to. 15s. From 1866 to 1869. Vol. XXII. Royal 4to. 15s.

A CATALOGUE OF THE COLLECTION OF BIRDS

formed by the late H. E. STRICKLAND, now in the possession of the University of Cambridge. By O. SALVIN, M.A. Demy 8vo. £1. 1s.

A CATALOGUE OF AUSTRALIAN FOSSILS, Strati-

graphically and Zoologically arranged, by R. ETHERIDGE, Jun., F.G.S. Demy 8vo. 10s. 6d.

ILLUSTRATIONS OF COMPARATIVE ANATOMY,

VERTEBRATE AND INVERTEBRATE, for the Use of Students in the Museum of Zoology and Comparative Anatomy. Second Edition. Demy 8vo. 2s. 6d.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse,
Ave Maria Lane.

A CATALOGUE OF THE COLLECTION OF CAMBRIAN AND SILURIAN FOSSILS contained in the Geological Museum of the University of Cambridge, by J. W. SALTER, F.G.S. With a Portrait of PROFESSOR SEDGWICK. Royal 4to. 7s. 6d.

CATALOGUE OF OSTEOLOGICAL SPECIMENS contained in the Anatomical Museum of the University of Cambridge. Demy 8vo. 2s. 6d.

LAW.

ELEMENTS OF THE LAW OF TORTS. A Text-book for Students. By MELVILLE M. BIGELOW, Ph.D., Lecturer in the Law School of the University of Boston, U.S.A. Crown 8vo. 10s. 6d.

"It is based on the original American edition, but it is an English Text-book with English authorities and statutes and illustrations substituted very generally for the American... The style is easy and lucid, though condensed,

showing great grasp of subject... A very full index enhances the value of this book, which should take a prominent place among the really trustworthy text-books for the use of students."—*Law Times*.

A SELECTION OF CASES ON THE ENGLISH LAW OF CONTRACT. By GERARD BROWN FINCH, M.A., of Lincoln's Inn, Barrister at Law. Royal 8vo. 28s.

"An invaluable guide towards the best method of legal study."—*Law Quarterly Review*.

THE INFLUENCE OF THE ROMAN LAW ON THE LAW OF ENGLAND. Being the Yorke Prize Essay for 1884. By T. E. SCRUTTON, M.A. Demy 8vo. 10s. 6d.

"Legal work of just the kind that a learned University should promote by its prizes."—*Law Quarterly Review*.

LAND IN FETTERS. Being the Yorke Prize Essay for 1885. By T. E. SCRUTTON, M.A. Demy 8vo. 7s. 6d.

COMMONS AND COMMON FIELDS, OR THE HISTORY AND POLICY OF THE LAWS RELATING TO COMMONS AND ENCLOSURES IN ENGLAND. Being the Yorke Prize Essay for 1886. By T. E. SCRUTTON, M.A. 10s. 6d.

HISTORY OF THE LAW OF TITHES IN ENGLAND. Being the Yorke Prize Essay for 1887. By W. EASTERBY, B.A., LL.B., St John's College and the Middle Temple. Demy 8vo. 7s. 6d.

HISTORY OF LAND TENURE IN IRELAND. Being the Yorke Prize Essay for 1888. By W. E. MONTGOMERY, M.A., LL.M. Demy 8vo. 10s. 6d.

AN ANALYSIS OF CRIMINAL LIABILITY. By E. C. CLARK, LL.D., Regius Professor of Civil Law in the University of Cambridge, also of Lincoln's Inn, Barrister-at-Law. Crown 8vo. 7s. 6d.

PRACTICAL JURISPRUDENCE, a Comment on AUSTIN. By E. C. CLARK, LL.D. Crown 8vo. 9s.

"Damit schliesst dieses inhaltreiche und nach allen Seiten anregende Buch über Prac-

tical Jurisprudence."—König. *Centralblatt für Rechtswissenschaft*.

London: C. J. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

A SELECTION OF THE STATE TRIALS. By J. W. WILLIS-BUND, M.A., LL.B., Professor of Constitutional Law and History, University College, London. Crown 8vo. Vols. I. and II. In 3 parts. **Now reduced to 30s.** (*originally published at 46s.*)

"This work is a very useful contribution to that important branch of the constitutional history of England which is concerned with the

growth and development of the law of treason, as it may be gathered from trials before the ordinary courts."—*The Academy*.

THE FRAGMENTS OF THE PERPETUAL EDICT OF SALVIUS JULIANUS, collected, arranged, and annotated by BRYAN WALKER, M.A., LL.D., late Law Lecturer of St John's College, and Fellow of Corpus Christi College, Cambridge. Crown 8vo. 6s.

"In the present book we have the fruits of the same kind of thorough and well-ordered study which was brought to bear upon the notes to the Commentaries and the Institutes. . . . Hitherto the Edict has been almost inaccessible to the ordinary English student, and

such a student will be interested as well as perhaps surprised to find how abundantly the extant fragments illustrate and clear up points which have attracted his attention in the Commentaries, or the Institutes, or the Digest."—*Law Times*.

BRACTON'S NOTE BOOK. A Collection of Cases decided in the King's Courts during the reign of Henry the Third, annotated by a Lawyer of that time, seemingly by Henry of Bratton. Edited by F. W. MAITLAND of Lincoln's Inn, Barrister at Law, Downing Professor of the Laws of England. 3 vols. Demy 8vo. Buckram. £3. 3s. *Net*.

AN INTRODUCTION TO THE STUDY OF JUSTINIAN'S DIGEST. Containing an account of its composition and of the Jurists used or referred to therein. By HENRY JOHN ROBY, M.A., formerly Prof. of Jurisprudence, University College, London. Demy 8vo. 9s.

JUSTINIAN'S DIGEST. Lib. VII., Tit. I. De Usufructu, with a Legal and Philological Commentary. By H. J. ROBY, M.A. Demy 8vo. 9s.

Or the Two Parts complete in One Volume. Demy 8vo. 18s.

"Not an obscurity, philological, historical, or legal, has been left unsifted. More informing aid still has been supplied to the student of the Digest at large by a preliminary account, covering nearly 300 pages, of the mode of composition of the Digest, and of the jurists

whose decisions and arguments constitute its substance. Nowhere else can a clearer view be obtained of the personal succession by which the tradition of Roman legal science was sustained and developed."—*The Times*.

THE COMMENTARIES OF GAIUS AND RULES OF ULPIAN. With a Translation and Notes, by J. T. ABDY, LL.D., Judge of County Courts, late Regius Professor of Laws in the University of Cambridge, and the late BRYAN WALKER, M.A., LL.D., New Edition by BRYAN WALKER. Crown 8vo. 16s.

"As scholars and as editors Messrs Abdy and Walker have done their work well. . . . For one thing the editors deserve special commendation. They have presented Gaius to the reader with few notes and those merely by

way of reference or necessary explanation. Thus the Roman jurist is allowed to speak for himself, and the reader feels that he is really studying Roman law in the original, and not a fanciful representation of it."—*Athenæum*.

THE INSTITUTES OF JUSTINIAN, translated with Notes by J. T. ABDY, LL.D., and the late BRYAN WALKER, M.A., LL.D. Crown 8vo. 16s.

"We welcome here a valuable contribution to the study of jurisprudence. The text of the *Institutes* is occasionally perplexing, even to practised scholars, whose knowledge of classical models does not always avail them in dealing with the technicalities of legal phraseology. Nor can the ordinary dictionaries be expected to furnish all the help that is wanted. This translation will then be of great use. To

the ordinary student, whose attention is distracted from the subject-matter by the difficulty of struggling through the language in which it is contained, it will be almost indispensable."—*Spectator*.

"The notes are learned and carefully compiled, and this edition will be found useful to students."—*Law Times*.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

SELECTED TITLES FROM THE DIGEST, annotated by the late B. WALKER, M.A., LL.D. Part I. Mandati vel Contra. Digest XVII. 1. Crown 8vo. 5s.

— Part II. De Adquirendo rerum dominio and De Adquirenda vel amittenda possessione. Digest XLI. 1 and 11. Crown 8vo. 6s.

— Part III. De Conditionibus. Digest XII. 1 and 4—7 and Digest XIII. 1—3. Crown 8vo. 6s.

GROTIUS DE JURE BELLII ET PACIS, with the Notes of Barbeyrac and others; accompanied by an abridged Translation of the Text, by W. WHEWELL, D.D. late Master of Trinity College. 3 Vols. Demy 8vo. 12s. The translation separate, 6s.

HISTORICAL WORKS, &c.

THE LIFE AND LETTERS OF THE REVEREND ADAM SEDGWICK, LL.D., F.R.S., Fellow of Trinity College, Cambridge, and Woodwardian Professor of Geology from 1818 to 1873. (Dedicated, by special permission, to Her Majesty the Queen.) By JOHN WILLIS CLARK, M.A., F.S.A., formerly Fellow of Trinity College, and THOMAS M^cKENNY HUGHES, M.A., Woodwardian Professor of Geology. 2 vols. Demy 8vo. [Nearly ready.]

THE DESPATCHES OF EARL GOWER, English Ambassador at the court of Versailles from June 1790 to August 1792, to which are added the Despatches of Mr Lindsay and Mr Munro, and the Diary of Lord Palmerston in France during July and August 1791. Edited by OSCAR BROWNING, M.A. Demy 8vo. 15s.

LIFE AND TIMES OF STEIN, OR GERMANY AND PRUSSIA IN THE NAPOLEONIC AGE, by J. R. SEELEY, M.A., Regius Professor of Modern History in the University of Cambridge, with Portraits and Maps. 3 Vols. Demy 8vo. 30s.

"DR BUSCH's volume has made people think and talk even more than usual of Prince Bismarck, and Professor Seeley's very learned work on Stein will turn attention to an earlier and an almost equally eminent German statesman.... He was one, perhaps the chief, of the illustrious group of strangers who came to the rescue of Prussia in her darkest hour, about the time of the inglorious Peace of Tilsit, and who laboured to put life and order into her dispirited army, her impoverished finances, and her inefficient Civil Service. Englishmen will feel very pardonable pride at seeing one of their countrymen undertake to write the history of a period from the investigation of

which even laborious Germans are apt to shrink."—*Times*.

"In a notice of this kind scant justice can be done to a work like the one before us; no short *résumé* can give even the most meagre notion of the contents of these volumes, which contain no page that is superfluous, and none that is uninteresting.... To understand the Germany of to-day one must study the Germany of many yesterdays, and now that study has been made easy by this work, to which no one can hesitate to assign a very high place among those recent histories which have aimed at original research."—*Athenæum*.

THE GROWTH OF ENGLISH INDUSTRY AND COMMERCE DURING THE EARLY AND MIDDLE AGES.

By W. CUNNINGHAM, D.D., University Lecturer. Demy 8vo. 16s.

CHRONOLOGICAL TABLES OF GREEK HISTORY.

Accompanied by a short narrative of events, with references to the sources of information and extracts from the ancient authorities, by CARL PETER. Translated from the German by G. CHAWNER, M.A., Fellow of King's College, Cambridge. Demy 4to. 10s.

London: C. J. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

THE ARCHITECTURAL HISTORY OF THE UNIVERSITY OF CAMBRIDGE AND OF THE COLLEGES OF CAMBRIDGE AND ETON, by the late ROBERT WILLIS, M.A. F.R.S., Jacksonian Professor in the University of Cambridge. Edited with large Additions and brought up to the present time by JOHN WILLIS CLARK, M.A., formerly Fellow of Trinity College, Cambridge. Four Vols. Super Royal 8vo. £6. 6s.

Also a limited Edition of the same, consisting of 120 numbered Copies only, large paper Quarto; the woodcuts and steel engravings mounted on India paper; price Twenty-five Guineas *net* each set.

THE UNIVERSITY OF CAMBRIDGE FROM THE EARLIEST TIMES TO THE ROYAL INJUNCTIONS OF 1535, by J. B. MULLINGER, M.A., Lecturer on History and Librarian to St John's College. Part I. Demy 8vo. (734 pp.), 12s.

Part II. From the Royal Injunctions of 1535 to the Accession of Charles the First. Demy 8vo. 18s.

"He shews in the statutes of the Colleges, the internal organization of the University, its connection with national problems, its studies, its social life. All this he combines in a form which is eminently readable."—PROF. CREIGHTON in *Cont. Review*.

"Mr Mullinger displays an admirable thoroughness in his work. Nothing could be more exhaustive and conscientious than his method: and his style...is picturesque and elevated."—*Times*.

SCHOLAE ACADEMICAE: some Account of the Studies at the English Universities in the Eighteenth Century. By C. WORDSWORTH, M.A., Fellow of Peterhouse. Demy 8vo. 10s. 6d.

"Mr Wordsworth has collected a great quantity of minute and curious information about the working of Cambridge institutions in the last century, with an occasional comparison of the corresponding state of things at Oxford.

... To a great extent it is purely a book of reference, and as such it will be of permanent value for the historical knowledge of English education and learning."—*Saturday Review*.

HISTORY OF THE COLLEGE OF ST JOHN THE EVANGELIST, by THOMAS BAKER, B.D., Ejected Fellow. Edited by JOHN E. B. MAYOR, M.A. Two Vols. Demy 8vo. 24s.

HISTORY OF NEPĀL, translated by MUNSHĪ SHEW SHUNKER SINGH and PANDIT SHRĪ GUNĀNAND; edited with an Introductory Sketch of the Country and People by Dr D. WRIGHT, late Residency Surgeon at Kāthmāndū, and with facsimiles of native drawings, and portraits of Sir JUNG BAHĀDUR, the KING OF NEPĀL, &c. Super-royal 8vo. 10s. 6d.

KINSHIP AND MARRIAGE IN EARLY ARABIA, by W. ROBERTSON SMITH, M.A., LL.D., Professor of Arabic and Fellow of Christ's College. Crown 8vo. 7s. 6d.

"It would be superfluous to praise a book so learned and masterly as Professor Robertson Smith's; it is enough to say that no student of

early history can afford to be without *Kinship in Early Arabia*."—*Nature*.

TRAVELS IN ARABIA DESERTA IN 1876 AND 1877. BY CHARLES M. DOUGHTY, of Gonville and Caius College. With Illustrations and a Map. 2 vols. Demy 8vo. £3. 3s.

"This is in several respects a remarkable book. It records the ten years' travels of the author throughout Northern Arabia, in the Hejas and Nejd, from Syria to Mecca. No doubt this region has been visited by previous travellers, but none, we venture to think, have done their work with so much thoroughness or with more enthusiasm and love."—*Times*.

"We judge this book to be the most remarkable record of adventure and research which has been published to this generation."—*Spectator*.

"Its value as a storehouse of knowledge simply cannot be exaggerated."—*Saturday Review*.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

A JOURNEY OF LITERARY AND ARCHÆOLOGICAL RESEARCH IN NEPAL AND NORTHERN INDIA, during the Winter of 1884-5. By CECIL BENDALL, M.A., Professor of Sanskrit in University College, London. Demy 8vo. 10s.

THE CONSTITUTION OF CANADA. By J. E. C. MUNRO, LL.M., Professor of Law and Political Economy at Victoria University, Manchester. Demy 8vo. 10s.

CAMBRIDGE HISTORICAL ESSAYS.

POLITICAL PARTIES IN ATHENS DURING THE PELOPONNESIAN WAR, by L. WHIBLEY, M.A., Fellow of Pembroke College, Cambridge. (Prince Consort Dissertation, 1888.) *Second Edition.* Crown 8vo. 2s. 6d.

POPE GREGORY THE GREAT AND HIS RELATIONS WITH GAUL, by F. W. KELLETT, M.A., Sidney Sussex College. (Prince Consort Dissertation, 1888.) Crown 8vo. 2s. 6d.

THE CONSTITUTIONAL EXPERIMENTS OF THE COMMONWEALTH, being the Thirlwall Prize Essay for 1889, by E. JENKS, B.A., LL.B. Crown 8vo. [In the Press.]

MISCELLANEOUS.

THE LITERARY REMAINS OF ALBRECHT DÜRER, by W. M. CONWAY. With Transcripts from the British Museum MSS., and Notes by LINA ECKENSTEIN. Royal 8vo. 21s. (*The Edition is limited to 500 copies.*)

A LATIN-ENGLISH DICTIONARY. Printed from the (Incomplete) MS. of the late T. H. KEY, M.A., F.R.S. Cr. 4to. 31s. 6d.

THE COLLECTED PAPERS OF HENRY BRADSHAW, including his Memoranda and Communications read before the Cambridge Antiquarian Society. *With 13 fac-similes.* Edited by F. J. H. JENKINSON, M.A., Fellow of Trinity College. Demy 8vo. 16s.

THE LATIN HEPTATEUCH. Published piecemeal by the French printer WILLIAM MOREL (1560) and the French Benedictines E. MARTÈNE (1733) and J. B. PITRA (1852-88). Critically reviewed by JOHN E. B. MAYOR, M.A., Professor of Latin in the University of Cambridge. Demy 8vo. 10s. 6d.

A CATALOGUE OF ANCIENT MARBLES IN GREAT BRITAIN, by Prof. ADOLF MICHAELIS. Translated by C. A. M. FENNELL, Litt. D. Royal 8vo. Roxburgh (Morocco back), £2. 2s.

"The book is beautifully executed, and with its few handsome plates, and excellent indexes, does much credit to the Cambridge Press. All lovers of true art and of good work should be

grateful to the Syndics of the University Press for the liberal facilities afforded by them towards the production of this important volume by Professor Michaelis."—*Saturday Review.*

London: C. F. CLAY & SONS, Cambridge University Press Warehouse,
Ave Maria Lane.

CONTRIBUTIONS TO THE TEXTUAL CRITICISM

OF THE DIVINA COMMEDIA. Including the complete collation throughout the *Inferno* of all the MSS. at Oxford and Cambridge. By the Rev. EDWARD MOORE, D.D. Demy 8vo. 21s.

RHODES IN ANCIENT TIMES. By CECIL TORR, M.A.

With six plates. Demy 8vo. 10s. 6d.

RHODES IN MODERN TIMES. By the same Author.

With three plates. Demy 8vo. 8s.

THE WOODCUTTERS OF THE NETHERLANDS

during the last quarter of the Fifteenth Century. In 3 parts. I. History of the Woodcutters. II. Catalogue of their Woodcuts. III. List of Books containing Woodcuts. By W. M. CONWAY. Demy 8vo. 10s. 6d.

THE LITERATURE OF THE FRENCH RENAISSANCE.

An Introductory Essay. By A. A. TILLEY, M.A. Cr. 8vo. 6s.

FROM SHAKESPEARE TO POPE: an Inquiry into

the causes and phenomena of the rise of Classical Poetry in England. By EDMUND GOSSE, M.A. Crown 8vo. 6s.

CHAPTERS ON ENGLISH METRE. By Rev. JOSEPH

B. MAYOR, M.A. Demy 8vo. 7s. 6d.

A GRAMMAR OF THE IRISH LANGUAGE. By Prof.

WINDISCH. Translated by Dr NORMAN MOORE. Crown 8vo. 7s. 6d.

LECTURES ON TEACHING, delivered in the University

of Cambridge in the Lent Term, 1880. By J. G. FITCH, M.A., LL.D. Her Majesty's Inspector of Training Colleges. Cr. 8vo. New Edit. 5s.

"Mr Fitch's book covers so wide a field and touches on so many burning questions that we must be content to recommend it as the best existing *vade mecum* for the teacher."—*Pall Mall Gazette*.

LECTURES ON THE SCIENCE OF EDUCATION.

By FRANCIS WARNER, M.D., F.R.C.P.

[*In the Press.*]

OCCASIONAL ADDRESSES ON EDUCATIONAL

SUBJECTS. By S. S. LAURIE, M.A., LL.D. Crown 8vo. 5s.

A MANUAL OF CURSIVE SHORTHAND. By H. L.

CALLENDAR, M.A., Fellow of Trinity College. Ex. Fcap. 8vo. 2s.

A SYSTEM OF PHONETIC SPELLING ADAPTED

TO ENGLISH. By H. L. CALLENDAR, M.A. Ex. Fcap. 8vo. 6d.

A PRIMER OF CURSIVE SHORTHAND. By H. L.

CALLENDAR, M.A. Ex. Fcap. 8vo. 6d.

READING PRACTICE IN CURSIVE SHORTHAND.

Easy Extracts for Beginners. The Gospel according to St Mark, (First half). The Vicar of Wakefield. Chaps. I.—V. Alice in Wonderland. Chap. VII. 3d. each.

For other books on Education, see Pitt Press Series, p. 39.

London: C. J. CLAY & SONS, Cambridge University Press Warehouse,
Ave Maria Lane.

STUDIES IN THE LITERARY RELATIONS OF ENGLAND WITH GERMANY IN THE SIXTEENTH CENTURY. By C. H. HERFORD, M.A. Crown 8vo. 9s.

ADMISSIONS TO GONVILLE AND CAIUS COLLEGE in the University of Cambridge March 1558—9 to Jan. 1678—9 Edited by J. VENN, Sc.D., and S. C. VENN. Demy 8vo. 10s.

ECCLESIAE LONDINO—BATAVAE ARCHIVVM. TOMVS PRIMVS. ABRAHAMI ORTELII et virorum eruditorum ad eundem et ad JACOBVM COLIVM ORTELIANVM Epistulae, 1524—1628. TOMVS SECVNDVS. EPISTVLAE ET TRACTATVS cum Reformationis tum Ecclesiae Londino-Batavae Historiam Illustrantes 1544—1622. Ex autographis mandante Ecclesia Londino-Batava edidit JOANNES HENRICVS HESSELS. Demy 4to. Each volume, separately, £3. 10s. Taken together £5. 5s. *Net*.

CATALOGUE OF THE HEBREW MANUSCRIPTS preserved in the University Library, Cambridge. By Dr S. M. SCHILLER-SZINESSY. Volume I. containing Section I. *The Holy Scriptures*; Section II. *Commentaries on the Bible*. Demy 8vo. 9s.

A CATALOGUE OF THE MANUSCRIPTS preserved in the Library of the University of Cambridge. Demy 8vo. 5 Vols. 10s. each. INDEX TO THE CATALOGUE. Demy 8vo. 10s.

A CATALOGUE OF ADVERSARIA and printed books preserved in the Library of the University of Cambridge. 3s. 6d.

THE ILLUMINATED MANUSCRIPTS IN THE LIBRARY of the Fitzwilliam Museum, Catalogued with Descriptions, and an Introduction, by W. G. SEARLE, M.A. Demy 8vo. 7s. 6d.

A CHRONOLOGICAL LIST OF THE GRACES, Documents, and other Papers in the University Registry which concern the University Library. Demy 8vo. 2s. 6d.

CATALOGUS BIBLIOTHECÆ BURCKHARDTIANÆ. Demy 4to. 5s.

GRADUATI CANTABRIGIENSES: SIVE CATALOGUS exhibens nomina eorum quos gradu quocunque ornavit Academia Cantabrigiensis (1800—1884). Cura H. R. LUARD S. T. P. Demy 8vo. 12s. 6d.

STATUTES OF THE UNIVERSITY OF CAMBRIDGE and for the Colleges therein, made, published and approved (1878—1882) under the Universities of Oxford and Cambridge Act, 1877. With an Appendix. Demy 8vo. 16s.

STATUTES OF THE UNIVERSITY OF CAMBRIDGE. With Acts of Parliament relating to the University. 8vo. 3s. 6d.

ORDINANCES OF THE UNIVERSITY OF CAMBRIDGE. Demy 8vo., cloth. 7s. 6d.

TRUSTS, STATUTES AND DIRECTIONS affecting (1) The Professorships of the University. (2) The Scholarships and Prizes. (3) Other Gifts and Endowments. Demy 8vo. 5s.

COMPENDIUM of UNIVERSITY REGULATIONS. 6d.

London: C. J. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

The Cambridge Bible for Schools and Colleges.

GENERAL EDITOR: THE VERY REVEREND J. J. S. PEROWNE, D.D.,
DEAN OF PETERBOROUGH.

"It is difficult to commend too highly this excellent series."—*Guardian*.

"The modesty of the general title of this series has, we believe, led many to misunderstand its character and underrate its value. The books are well suited for study in the upper forms of our best schools, but not the less are they adapted to the wants of all Bible students who are not specialists. We doubt, indeed, whether any of the numerous popular commentaries recently issued in this country will be found more serviceable for general use."—*Academy*.

"One of the most popular and useful literary enterprises of the nineteenth century."—*Baptist Magazine*.

"Of great value. The whole series of comments for schools is highly esteemed by students capable of forming a judgment. The books are scholarly without being pretentious: information is so given as to be easily understood."—*Sword and Trowel*.

The Very Reverend J. J. S. PEROWNE, D.D., Dean of Peterborough, has undertaken the general editorial supervision of the work, assisted by a staff of eminent coadjutors. Some of the books have been already edited or undertaken by the following gentlemen:

- Rev. A. CARR, M.A., *late Assistant Master at Wellington College.*
 Rev. T. K. CHEYNE, M.A., D.D., *Canon of Rochester.*
 Rev. S. COX, *Nottingham.*
 Rev. A. B. DAVIDSON, D.D., *Professor of Hebrew, Edinburgh.*
 The Ven. F. W. FARRAR, D.D., *Archdeacon of Westminster.*
 Rev. C. D. GINSBURG, LL.D.
 Rev. A. E. HUMPHREYS, M.A., *late Fellow of Trinity College, Cambridge.*
 Rev. A. F. KIRKPATRICK, B.D., *Fellow of Trinity College, Regius Professor of Hebrew.*
 Rev. J. J. LIAS, M.A., *late Professor at St David's College, Lampeter.*
 Rev. J. R. LUMBY, D.D., *Norrisian Professor of Divinity.*
 Rev. G. F. MACLEAR, D.D., *Warden of St Augustine's College, Canterbury.*
 Rev. H. C. G. MOULE, M.A., *late Fellow of Trinity College, Principal of Ridley Hall, Cambridge.*
 Rev. E. H. PEROWNE, D.D., *Master of Corpus Christi College, Cambridge.*
 The Ven. T. T. PEROWNE, B.D., *Archdeacon of Norwich.*
 Rev. A. PLUMMER, M.A., D.D., *Master of University College, Durham.*
 The Very Rev. E. H. PLUMPTRE, D.D., *Dean of Wells.*
 Rev. H. E. RYLE, M.A., *Hulsean Professor of Divinity.*
 Rev. W. SIMCOX, M.A., *late Rector of Weyhill, Hants.*
 W. ROBERTSON SMITH, M.A., *Professor of Arabic and Fellow of Christ's College.*
 The Very Rev. H. D. M. SPENCE, M.A., *Dean of Gloucester.*
 Rev. A. W. STREANE, M.A., *Fellow of Corpus Christi College, Cambridge.*

London: C. F. CLAY & SONS, Cambridge University Press Warehouse,
Ave Maria Lane.

THE CAMBRIDGE BIBLE FOR SCHOOLS & COLLEGES. *Cont.**Now Ready. Cloth, Extra Fcap. 8vo.*

- THE BOOK OF JOSHUA. By the Rev. G. F. MACLEAR, D.D.
With 2 Maps. 2s. 6d.
- THE BOOK OF JUDGES. By the Rev. J. J. LIAS, M.A.
With Map. 3s. 6d.
- THE FIRST BOOK OF SAMUEL. By the Rev. Professor
KIRKPATRICK, B.D. With Map. 3s. 6d.
- THE SECOND BOOK OF SAMUEL. By the Rev. Professor
KIRKPATRICK, B.D. With 2 Maps. 3s. 6d.
- THE FIRST BOOK OF KINGS. By Rev. Prof. LUMBY, D.D. 3s. 6d.
- THE SECOND BOOK OF KINGS. By the same Editor. 3s. 6d.
- THE BOOK OF JOB. By the Rev. A. B. DAVIDSON, D.D. 5s.
- THE BOOK OF ECCLESIASTES. By the Very Rev. E. H.
PLUMPTRE, D.D. 5s.
- THE BOOK OF JEREMIAH. By the Rev. A. W. STREANE,
M.A. With Map. 4s. 6d.
- THE BOOK OF HOSEA. By Rev. T. K. CHEYNE, M.A., D.D. 3s.
- THE BOOKS OF OBADIAH AND JONAH. By Archdeacon
PEROWNE. 2s. 6d.
- THE BOOK OF MICAH. By Rev. T. K. CHEYNE, D.D. 1s. 6d.
- THE BOOKS OF HAGGAI AND ZECHARIAH. By Arch-
deacon PEROWNE. 3s.
- THE GOSPEL ACCORDING TO ST MATTHEW. By the
Rev. A. CARR, M.A. With 2 Maps. 2s. 6d.
- THE GOSPEL ACCORDING TO ST MARK. By the Rev.
G. F. MACLEAR, D.D. With 4 Maps. 2s. 6d.
- THE GOSPEL ACCORDING TO ST LUKE. By Archdeacon
F. W. FARRAR. With 4 Maps. 4s. 6d.
- THE GOSPEL ACCORDING TO ST JOHN. By the Rev.
A. PLUMMER, M.A., D.D. With 4 Maps. 4s. 6d.
- THE ACTS OF THE APOSTLES. By the Rev. Professor
LUMBY, D.D. With 4 Maps. 4s. 6d.
- THE EPISTLE TO THE ROMANS. By the Rev. H. C. G.
MOULE, M.A. 3s. 6d.
- THE FIRST EPISTLE TO THE CORINTHIANS. By the Rev.
J. J. LIAS, M.A. With a Map and Plan. 2s.
- THE SECOND EPISTLE TO THE CORINTHIANS. By the
Rev. J. J. LIAS, M.A. 2s.
- THE EPISTLE TO THE EPHESIANS. By the Rev. H. C. G.
MOULE, M.A. 2s. 6d.
- THE EPISTLE TO THE PHILIPPIANS. By the Rev. H. C. G.
MOULE, M.A. 2s. 6d.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse,
Ave Maria Lane.

THE CAMBRIDGE BIBLE FOR SCHOOLS & COLLEGES. *Cont.*

THE EPISTLE TO THE HEBREWS. By Arch. FARRAR. 3s. 6d.

THE GENERAL EPISTLE OF ST JAMES. By the Very Rev.
E. H. PLUMPTRE, D.D. 1s. 6d.THE EPISTLES OF ST PETER AND ST JUDE. By the
same Editor. 2s. 6d.THE EPISTLES OF ST JOHN. By the Rev. A. PLUMMER,
M.A., D.D. 3s. 6d.*Preparing.*THE BOOK OF GENESIS. By the Very Rev. the DEAN OF
PETERBOROUGH.THE BOOKS OF EXODUS, NUMBERS AND DEUTERO-
NOMY. By the Rev. C. D. GINSBURG, LL.D.THE BOOKS OF EZRA AND NEHEMIAH. By the Rev.
Prof. RYLE, M.A.

THE BOOK OF PSALMS. By the Rev. Prof. KIRKPATRICK, B.D.

THE BOOK OF ISAIAH. By Prof. W. ROBERTSON SMITH, M.A.

THE BOOK OF EZEKIEL. By the Rev. A. B. DAVIDSON, D.D.

THE BOOK OF MALACHI. By Archdeacon PEROWNE.

THE EPISTLE TO THE GALATIANS. By the Rev. E. H.
PEROWNE, D.D.THE EPISTLES TO THE COLOSSIANS AND PHILEMON.
By the Rev. H. C. G. MOULE, M.A.THE EPISTLES TO TIMOTHY AND TITUS. By the Rev.
A. E. HUMPHREYS, M.A.

THE BOOK OF REVELATION. By the Rev. W. SIMCOX, M.A.

The Smaller Cambridge Bible for Schools.

The Smaller Cambridge Bible for Schools will form an entirely new series of commentaries on some selected books of the Bible. It is expected that they will be prepared for the most part by the Editors of the larger series (The Cambridge Bible for Schools and Colleges). The volumes will be issued at a low price, and will be suitable to the requirements of preparatory and elementary schools.

*Now ready. Price 1s. each.*THE FIRST AND SECOND BOOKS OF SAMUEL. By
Rev. Prof. KIRKPATRICK, B.D.THE GOSPEL ACCORDING TO ST MATTHEW. By Rev.
A. CARR, M.A.THE GOSPEL ACCORDING TO ST MARK. By Rev.
G. F. MACLEAR, D.D.*Nearly ready.*THE GOSPEL ACCORDING TO ST LUKE. By ARCHDEACON
FARRAR, D.D.

*London: C. F. CLAY & SONS, Cambridge University Press Warehouse
Ave Maria Lane.*

The Cambridge Greek Testament for Schools and Colleges,

with a Revised Text, based on the most recent critical authorities, and
English Notes, prepared under the direction of the General Editor,
THE VERY REVEREND J. J. S. PEROWNE, D.D.

Now Ready.

THE GOSPEL ACCORDING TO ST MATTHEW. By the
Rev. A. CARR, M.A. With 4 Maps. 4s. 6d.

"Copious illustrations, gathered from a great variety of sources, make his notes a very valuable aid to the student. They are indeed remarkably interesting, while all explanations on meanings, applications, and the like are distinguished by their lucidity and good sense."—*Pall Mall Gazette*.

THE GOSPEL ACCORDING TO ST MARK. By the Rev.
G. F. MACLEAR, D.D. With 3 Maps. 4s. 6d.

"The Cambridge Greek Testament, of which Dr Maclear's edition of the Gospel according to St Mark is a volume, certainly supplies a want. Without pretending to compete with the leading commentaries, or to embody very much original research, it forms a most satisfactory introduction to the study of the New Testament in the original . . . Dr Maclear's introduction contains all that is known of St Mark's life, an account of the circumstances in which the Gospel was composed, an excellent sketch of the special characteristics of this Gospel; an analysis, and a chapter on the text of the New Testament generally . . . The work is completed by three good maps."—*Saturday Review*.

THE GOSPEL ACCORDING TO ST LUKE. By Archdeacon
FARRAR. With 4 Maps. 6s.

THE GOSPEL ACCORDING TO ST JOHN. By the Rev. A.
PLUMMER, M.A., D.D. With 4 Maps. 6s.

"A valuable addition has also been made to 'The Cambridge Greek Testament for Schools,' Dr Plummer's notes on 'the Gospel according to St John' are scholarly, concise, and instructive, and embody the results of much thought and wide reading."—*Expositor*.

THE ACTS OF THE APOSTLES. By the Rev. Prof. LUMBY, D.D.,
with 4 Maps. 6s.

THE FIRST EPISTLE TO THE CORINTHIANS. By the
Rev. J. J. LIAS, M.A. 3s.

THE SECOND EPISTLE TO THE CORINTHIANS. By the
Rev. J. J. LIAS, M.A. [Preparing.]

THE EPISTLE TO THE HEBREWS. By Arch. FARRAR, D.D.
3s. 6d.

THE EPISTLES OF ST JOHN. By the Rev. A. PLUMMER,
M.A., D.D. 4s.

London: C. J. CLAY & SONS, Cambridge University Press Warehouse,
Ave Maria Lane.

THE PITT PRESS SERIES.

[Copies of the Pitt Press Series may generally be obtained bound in two parts for Class use, the text and notes in separate volumes.]

I. GREEK.

ARISTOPHANES—AVES—PLUTUS—RANAE. With English Notes and Introduction by W. C. GREEN, M.A., late Assistant Master at Rugby School. 3s. 6d. each.

EURIPIDES. HERACLEIDÆ. With Introduction and Explanatory Notes by E. A. BECK, M.A., Fellow of Trinity Hall. 3s. 6d.

EURIPIDES. HERCULES FURENS. With Introductions, Notes and Analysis. By A. GRAY, M.A., Fellow of Jesus College, and J. T. HUTCHINSON, M.A., Christ's College. New Edition. 2s.

EURIPIDES. HIPPOLYTUS. By W. S. HADLEY, M.A. Fellow of Pembroke College. 2s.

EURIPIDES. IPHIGENEIA IN AULIS. By C. E. S. HEADLAM, B.A., Fellow of Trinity Hall. 2s. 6d.

HERODOTUS, BOOK V. Edited with Notes, Introduction and Maps by E. S. SHUCKBURGH, M.A., late Fellow of Emmanuel College. 3s.

HERODOTUS, BOOK VI. By the same Editor. 4s.

HERODOTUS, BOOK VIII., CHAPS. 1—90. By the same Editor. 3s. 6d.

"We could not wish for a better introduction to Herodotus."—*Journal of Education*.

HERODOTUS, BOOK IX., CHAPS. 1—89. By the same Editor. 3s. 6d.

HOMER—ODYSSEY, BOOKS IX. X. With Introduction, Notes and Appendices. By G. M. EDWARDS, M.A., Fellow and Classical Lecturer of Sidney Sussex College. 2s. 6d. each.

HOMER—ODYSSEY, BOOK XXI. By the same Editor. 2s.

LUCIANI SOMNIUM CHARON PISCATOR ET DE LUCTU, with English Notes by W. E. HEITLAND, M.A., Fellow of St John's College, Cambridge. New Edition, with Appendix. 3s. 6d.

PLATONIS APOLOGIA SOCRATIS. With Introduction, Notes and Appendices by J. ADAM, M.A., Fellow and Classical Lecturer of Emmanuel College. 3s. 6d.

"A worthy representative of English Scholarship."—*Classical Review*.

— CRITO. With Introduction, Notes and Appendix. By the same Editor. 2s. 6d.

"Mr Adam, already known as the author of a careful and scholarly edition of the Apology of Plato, will, we think, add to his reputation by his work upon the Crito."—*Academy*.

"A scholarly edition of a dialogue which has never been really well edited in English."—*Guardian*.

— EUTHYPHRO. By the same Editor. [In the Press.]

London: C. F. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

- PLUTARCH. LIVES OF THE GRACCHI. With Introduction, Notes and Lexicon by Rev. HUBERT A. HOLDEN, M.A., LL.D. 6s.
- PLUTARCH. LIFE OF NICIAS. With Introduction and Notes. By Rev. HUBERT A. HOLDEN, M.A., LL.D. 5s.
 "This edition is as careful and thorough as Dr Holden's work always is."—*Spectator*.
- PLUTARCH. LIFE OF SULLA. With Introduction, Notes, and Lexicon. By the Rev. HUBERT A. HOLDEN, M.A., LL.D. 6s.
- PLUTARCH. LIFE OF TIMOLEON. With Introduction, Notes and Lexicon. By Rev. HUBERT A. HOLDEN, M.A., LL.D. 6s.
- SOPHOCLES.—OEDIPUS TYRANNUS. School Edition, with Introduction and Commentary, by R. C. JEBB, Litt. D., LL.D., Regius Professor of Greek in the University of Cambridge. 4s. 6d.
- THUCYDIDES. BOOK VII. With Notes and Introduction By H. R. TOTTENHAM, M.A., Fellow of St John's College. [*In the Press*].
- XENOPHON.—AGESILAUS. The Text revised with Critical and Explanatory Notes, Introduction, Analysis, and Indices. By H. HAILSTONE, M.A., late Scholar of Peterhouse. 2s. 6d.
- XENOPHON.—ANABASIS, BOOKS I. III. IV. and V. With a Map and English Notes by ALFRED PRETOR, M.A., Fellow of St Catharine's College, Cambridge. 2s. each.
 "Mr Pretor's 'Anabasis of Xenophon, Book IV.' displays a union of accurate Cambridge scholarship, with experience of what is required by learners gained in examining middle-class schools. The text is large and clearly printed, and the notes explain all difficulties. . . . Mr Pretor's notes seem to be all that could be wished as regards grammar, geography, and other matters."—*The Academy*.
- — BOOKS II. VI. and VII. By the same. 2s. 6d. each.
 "Had we to introduce a young Greek scholar to Xenophon, we should esteem ourselves fortunate in having Pretor's text-book as our chart and guide."—*Contemporary Review*.
- XENOPHON.—ANABASIS. By A. PRETOR, M.A., Text and Notes, complete in two Volumes. 7s. 6d.
- XENOPHON.—CYROPAEDEIA. BOOKS I. II. With Introduction, Notes and Map. By Rev. H. A. HOLDEN, M.A., LL.D. 2 vols. Vol. I. Text. Vol. II. Notes. 6s.
 "The work is worthy of the editor's well-earned reputation for scholarship and industry."—*Athenæum*.
- — BOOKS III., IV., V. By the same Editor. 5s.
 "Dr Holden's Commentary is equally good in history and in scholarship."—*Saturday Review*.

II. LATIN.

- BEDA'S ECCLESIASTICAL HISTORY, BOOKS III., IV., the Text from the very ancient MS. in the Cambridge University Library, collated with six other MSS. Edited, with a life from the German of EBERT, and with Notes, &c. by J. E. B. MAYOR, M.A., Professor of Latin, and J. R. LUMBY, D.D., Norrisian Professor of Divinity. Revised edition. 7s. 6d.

BOOKS I. and II. *In the Press*.

"In Bede's works Englishmen can go back to *origines* of their history, unequalled for form and matter by any modern European nation. Prof. Mayor has done good service in rendering a part of Bede's greatest work accessible to those who can read Latin with ease. He has adorned this edition of the third and fourth books of the 'Ecclesiastical History' with that amazing erudition for which he is unrivalled among Englishmen and rarely equalled by Germans. And however interesting and valuable the text may be, we can certainly apply to his notes the expression, *La sauce vaut mieux que le poisson*. They are literally crammed with interesting information about early English life. For though ecclesiastical in name, Bede's history treats of all parts of the national life, since the Church had points of contact with all."—*Examiner*.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse,
 Ave Maria Lane.

CAESAR. DE BELLO GALLICO COMMENT. I. With Maps and English Notes by A. G. PESKETT, M.A., Fellow of Magdalene College, Cambridge. 1s. 6d.

CAESAR. DE BELLO GALLICO COMMENT. II. III. By the same Editor. 2s.

CAESAR. DE BELLO GALLICO COMMENT. I. II. III. by the same Editor. 3s.

CAESAR. DE BELLO GALLICO COMMENT. IV. AND V. and COMMENT. VII. by the same Editor. 2s. each.

CAESAR. DE BELLO GALLICO COMMENT. VI. AND COMMENT. VIII. by the same Editor. 1s. 6d. each.

CICERO. ACTIO PRIMA IN C. VERREM. With Introduction and Notes. By H. COWIE, M.A., Fellow of St John's College, Cambridge. 1s. 6d.

CICERO. DE AMICITIA. Edited by J. S. REID, Litt.D., Fellow and Tutor of Gonville and Caius College. New Edition. 3s. 6d.

"Mr Reid has decidedly attained his aim, namely, 'a thorough examination of the Latinity of the dialogue.' . . . The revision of the text is most valuable, and comprehends sundry acute corrections. . . . This volume, like Mr Reid's other editions, is a solid gain to the scholarship of the country."—*Athenæum*.

"A more distinct gain to scholarship is Mr Reid's able and thorough edition of the *De Amicitia* of Cicero, a work of which, whether we regard the exhaustive introduction or the instructive and most suggestive commentary, it would be difficult to speak too highly. . . . When we come to the commentary, we are only amazed by its fulness in proportion to its bulk. Nothing is overlooked which can tend to enlarge the learner's general knowledge of Ciceronian Latin or to elucidate the text."—*Saturday Review*.

CICERO. DE SENECTUTE. Edited by J. S. REID, Litt.D. Revised Edition. 3s. 6d.

"The notes are excellent and scholarlike, adapted for the upper forms of public schools, and likely to be useful even to more advanced students."—*Guardian*.

CICERO. DIVINATIO IN Q. CAECILIUM ET ACTIO PRIMA IN C. VERREM. With Introduction and Notes by W. E. HEITLAND, M.A., and HERBERT COWIE, M.A., Fellows of St John's College, Cambridge. 3s.

CICERO. PHILIPPICA SECUNDA. With Introduction and Notes by A. G. PESKETT, M.A., Fellow of Magdalene College. 3s. 6d.

CICERO. PRO ARCHIA POETA. Edited by J. S. REID, Litt.D. Revised Edition. 2s.

"It is an admirable specimen of careful editing. An Introduction tells us everything we could wish to know about Archias, about Cicero's connexion with him, about the merits of the trial, and the genuineness of the speech. The text is well and carefully printed. The notes are clear and scholar-like. . . . No boy can master this little volume without feeling that he has advanced a long step in scholarship."—*The Academy*.

CICERO. PRO BALBO. Edited by J. S. REID, Litt.D. 1s. 6d.

"We are bound to recognize the pains devoted in the annotation of these two orations to the minute and thorough study of their Latinity, both in the ordinary notes and in the textual appendices."—*Saturday Review*.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

CICERO. PRO MILONE, with a Translation of Asconius' Introduction, Marginal Analysis and English Notes. Edited by the Rev. JOHN SMYTH PURTON, B.D., late President and Tutor of St Catharine's College. 2s. 6d.

"The editorial work is excellently done."—*The Academy*.

CICERO. PRO MURENA. With English Introduction and Notes. By W. E. HEITLAND, M.A., Fellow and Classical Lecturer of St John's College, Cambridge. Second Edition, carefully revised. 3s.

"Those students are to be deemed fortunate who have to read Cicero's lively and brilliant oration for L. Murena with Mr Heitland's handy edition, which may be pronounced 'four-square' in point of equipment, and which has, not without good reason, attained the honours of a second edition."—*Saturday Review*.

CICERO. PRO PLANCIO. Edited by H. A. HOLDEN, LL.D., Examiner in Greek to the University of London. Second Edition. 4s. 6d.

CICERO. PRO SULLA. Edited by J. S. REID, Litt.D. 3s. 6d.

"Mr Reid is so well known to scholars as a commentator on Cicero that a new work from him scarcely needs any commendation of ours. His edition of the speech *Pro Sulla* is fully equal in merit to the volumes which he has already published. . . It would be difficult to speak too highly of the notes. There could be no better way of gaining an insight into the characteristics of Cicero's style and the Latinity of his period than by making a careful study of this speech with the aid of Mr Reid's commentary. . . Mr Reid's intimate knowledge of the minutest details of scholarship enables him to detect and explain the slightest points of distinction between the usages of different authors and different periods. . . The notes are followed by a valuable appendix on the text, and another on points of orthography; an excellent index brings the work to a close."—*Saturday Review*.

CICERO. SOMNIUM SCIPIONIS. With Introduction and Notes. By W. D. PEARMAN, M.A., Head Master of Potsdam School, Jamaica. 2s.

HORACE. EPISTLES, BOOK I. With Notes and Introduction by E. S. SHUCKBURGH, M.A. 2s. 6d.

LIVY. BOOK IV. With Notes and Introduction, by Rev. H. M. STEPHENSON, M.A. 2s. 6d.

LIVY. BOOK V. With Notes and Introduction by L. WHIBLEY, M.A., Fellow of Pembroke College. 2s. 6d. [Shortly.]

LIVY. BOOKS XXI., XXII. With Notes, Introduction and Maps. By M. S. DIMSDALE, M.A., Fellow of King's College. 2s. 6d. each.

LUCAN. PHARSALIA LIBER PRIMUS. Edited with English Introduction and Notes by W. E. HEITLAND, M.A. and C. E. HASKINS, M.A., Fellows and Lecturers of St John's College, Cambridge. 1s. 6d.

"A careful and scholarlike production."—*Times*.

"In nice parallels of Lucan from Latin poets and from Shakspeare, Mr Haskins and Mr Heitland deserve praise."—*Saturday Review*.

LUCRETIUS. BOOK V. With Notes and Introduction by J. D. DUFF, M.A., Fellow of Trinity College. 2s.

OVID. FASTI. LIBER VI. With a Plan of Rome and Notes by A. SIDGWICK, M.A., Tutor of Corpus Christi College, Oxford. 1s. 6d.

"Mr Sidgwick's editing of the Sixth Book of Ovid's *Fasti* furnishes a careful and serviceable volume for average students. It eschews 'construes' which supersede the use of the dictionary, but gives full explanation of grammatical usages and historical and mythical allusions, besides illustrating peculiarities of style, true and false derivations, and the more remarkable variations of the text."—*Saturday Review*.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse,
Ave Maria Lane.

QUINTUS CURTIUS. A Portion of the History.

(ALEXANDER IN INDIA.) By W. E. HEITLAND, M.A., Fellow and Lecturer of St John's College, Cambridge, and T. E. RAVEN, B.A., Assistant Master in Sherborne School. 3s. 6d.

"Equally commendable as a genuine addition to the existing stock of school-books is *Alexander in India*, a compilation from the eighth and ninth books of Q. Curtius, edited for the Pitt Press by Messrs Heitland and Raven. . . . The work of Curtius has merits of its own, which, in former generations, made it a favourite with English scholars, and which still make it a popular text-book in Continental schools. . . . The reputation of Mr Heitland is a sufficient guarantee for the scholarship of the notes, which are ample without being excessive, and the book is well furnished with all that is needful in the nature of maps, indices, and appendices."—*Academy*.

VERGIL. AENEID. LIBRI I., II., III., IV., V., VI., VII. VIII., IX., X., XI., XII. Edited with Notes by A. SIDGWICK, M.A., Tutor of Corpus Christi College, Oxford. 1s. 6d. each.

"Mr Sidgwick's Vergil is . . . we believe, the best school edition of the poet."—*Guardian*.

"Mr Arthur Sidgwick's 'Vergil, Aeneid, Book XII.' is worthy of his reputation, and is distinguished by the same acuteness and accuracy of knowledge, appreciation of a boy's difficulties and ingenuity and resource in meeting them, which we have on other occasions had reason to praise in these pages."—*The Academy*.

"As masterly in its clearly divided preface and appendices as in the sound and independent character of its annotations. . . . There is a great deal more in the notes than mere compilation and suggestion. . . . No difficulty is left unnoticed or unhandled."—*Saturday Review*.

VERGIL. AENEID. LIBRI IX. X. in one volume. 3s.

VERGIL. AENEID. LIBRI X., XI., XII. in one volume. 3s. 6d.

VERGIL. BUCOLICS. With Introduction and Notes, by the same Editor. 1s. 6d.

VERGIL. GEORGICS. LIBRI I. II. By the same Editor. 2s. LIBRI III. IV. 2s.

"This volume, which completes the Pitt Press edition of Virgil's Georgics, is distinguished by the same admirable judgment and first-rate scholarship as are conspicuous in the former volume and in the 'Aeneid' by the same talented editor."—*Athenæum*.

VERGIL. The Complete Works, edited with Notes, by A. SIDGWICK, M.A., Two vols. Vol. I. containing the Text. Vol. II. The Notes. [Nearly ready.]

III. FRENCH.

CORNEILLE. LA SUITE DU MENTEUR. A Comedy in Five Acts. Edited with Fontenelle's Memoir of the Author, Voltaire's Critical Remarks, and Notes Philological and Historical. By the late GUSTAVE MASSON. 2s.

DE BONNECHOSE. LAZARE HOCHÉ. With Four Maps, Introduction and Commentary, by C. COLBECK, M.A., late Fellow of Trinity College, Cambridge. Revised Edition. 2s.

D'HARLEVILLE. LE VIEUX CÉLIBATAIRE. A Comedy. With a Biographical Memoir, and Grammatical, Literary and Historical Notes. By GUSTAVE MASSON. 2s.

London: C. J. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

DE LAMARTINE. JEANNE D'ARC. With a Map and Notes Historical and Philological and a Vocabulary by Rev. A. C. CLAPIN, M.A., St John's College, Cambridge, and Bachelier-ès-Lettres of the University of France. Enlarged Edition. 2s.

DE VIGNY. LA CANNE DE JONC. Edited with Notes by Rev. H. A. BULL, M.A. 2s.

ERCKMANN-CHATRIAN. LA GUERRE. With Map, Introduction and Commentary by the Rev. A. C. CLAPIN, M.A. 3s.

LA BARONNE DE STAËL-HOLSTEIN. LE DIRECTOIRE. (Considérations sur la Révolution Française. Troisième et quatrième parties.) With a Critical Notice of the Author, a Chronological Table, and Notes Historical and Philological, by G. MASSON, B.A., and G. W. PROTHERO, M.A. Revised and enlarged Edition. 2s.

"Prussia under Frederick the Great, and France under the Directory, bring us face to face respectively with periods of history which it is right should be known thoroughly, and which are well treated in the Pitt Press volumes. The latter in particular, an extract from the world-known work of Madame de Staël on the French Revolution, is beyond all praise for the excellence both of its style and of its matter."—*Times*.

LA BARONNE DE STAËL-HOLSTEIN. DIX ANNÉES D'EXIL. LIVRE II. CHAPITRES 1—8. With a Biographical Sketch of the Author, a Selection of Poetical Fragments by Madame de Staël's Contemporaries, and Notes Historical and Philological. By GUSTAVE MASSON and G. W. PROTHERO, M.A. Revised and enlarged edition. 2s.

LEMERCIER. FRÉDÉGONDE ET BRUNEHAUT. A Tragedy in Five Acts. Edited with Notes, Genealogical and Chronological Tables, a Critical Introduction and a Biographical Notice. By GUSTAVE MASSON. 2s.

MOLIÈRE. LE BOURGEOIS GENTILHOMME, Comédie-Ballet en Cinq Actes. (1670.) With a life of Molière and Grammatical and Philological Notes. By Rev. A. C. CLAPIN. Revised Edition. 1s. 6d.

MOLIÈRE. L'ÉCOLE DES FEMMES. Edited with Introduction and Notes by GEORGE SAINTSBURY, M.A. 2s. 6d.

"Mr Saintsbury's clear and scholarly notes are rich in illustration of the valuable kind that vivifies textual comment and criticism."—*Saturday Review*.

MOLIÈRE. LES PRÉCIEUSES RIDICULES. With Introduction and Notes by E. G. W. BRAUNHOLTZ, M.A., Ph.D. University Lecturer in French. 2s.

PIRON. LA METROMANIE, A Comedy, with a Biographical Memoir, and Grammatical, Literary and Historical Notes. By G. MASSON. 2s.

RACINE. LES PLAIDEURS. With Introduction and Notes by E. G. W. BRAUNHOLTZ, M.A., Ph.D. 2s.

SAINTE-BEUVE. M. DARU (Causeries du Lundi, Vol. IX.). With Biographical Sketch of the Author, and Notes Philological and Historical. By GUSTAVE MASSON. 2s.

SAINTINE. LA PICCIOLA. The Text, with Introduction, Notes and Map, by Rev. A. C. CLAPIN. 2s.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse,
Ave Maria Lane.

SCRIBE AND LEGOUVÉ. BATAILLE DE DAMES.

Edited by Rev. H. A. BULL, M.A. 2s.

SCRIBE. LE VERRE D'EAU. With a Biographical Memoir, and Grammatical, Literary and Historical Notes. By C. COLBECK, M.A. 2s.

"It may be national prejudice, but we consider this edition far superior to any of the series which hitherto have been edited exclusively by foreigners. Mr Colbeck seems better to understand the wants and difficulties of an English boy. The etymological notes especially are admirable. . . . The historical notes and introduction are a piece of thorough honest work."—*Journal of Education*.

SÉDAINE. LE PHILOSOPHE SANS LE SAVOIR.

Edited with Notes by Rev. H. A. BULL, M.A., late Master at Wellington College. 2s.

THIERRY. LETTRES SUR L'HISTOIRE DE FRANCE

(XIII.—XXIV.). By GUSTAVE MASSON, B.A. and G. W. PROTHERO, M.A. With Map. 2s. 6d.

THIERRY. RÉCITS DES TEMPS MÉROVINGIENS

I—III. Edited by GUSTAVE MASSON, B.A. Univ. Gallic., and A. R. ROPES, M.A. With Map. 3s.

VILLEMAIN. LASCARIS, OU LES GRECS DU XV^E.

SIÈCLE, Nouvelle Historique, with a Biographical Sketch of the Author, a Selection of Poems on Greece, and Notes Historical and Philological. By GUSTAVE MASSON, B.A. 2s.

VOLTAIRE. HISTOIRE DU SIÈCLE DE LOUIS XIV.

Part I. Chaps. I.—XIII. Edited with Notes Philological and Historical, Biographical and Geographical Indices, etc. by G. MASSON, B.A. Univ. Gallic., and G. W. PROTHERO, M.A., Fellow of King's College, Cambridge. 2s. 6d.

— Part II. Chaps. XIV.—XXIV. With Three Maps of the Period. By the same Editors. 2s. 6d.

— Part III. Chap. XXV. to the end. By the same Editors. 2s. 6d.

XAVIER DE MAISTRE. LA JEUNE SIBÉRIENNE.

LE LÉPREUX DE LA CITÉ D'AOSTE. With Biographical Notice, Critical Appreciations, and Notes. By G. MASSON, B.A. 1s. 6d.

IV. GERMAN.

BALLADS ON GERMAN HISTORY. Arranged and

Annotated by W. WAGNER, Ph.D., late Professor at the Johanneum, Hamburg. 2s.

"It carries the reader rapidly through some of the most important incidents connected with the German race and name, from the invasion of Italy by the Visigoths under their King Alaric, down to the Franco-German War and the installation of the present Emperor. The notes supply very well the connecting links between the successive periods, and exhibit in its various phases of growth and progress, or the reverse, the vast unwieldy mass which constitutes modern Germany."—*Times*.

London: C. J. CLAY & SONS, Cambridge University Press Warehouse,
Ave Maria Lane.

BENEDIX. DOCTOR WESPE. Lustspiel in fünf Aufzügen. Edited with Notes by KARL HERMANN BREUL, M.A. 3s.

FREYTAG. DER STAAT FRIEDRICHS DES GROSSEN. With Notes. By WILHELM WAGNER, Ph.D. 2s.

GERMAN DACTYLIC POETRY. Arranged and Annotated by the same Editor. 3s.

Goethe's Knabenjahre. (1749—1759.) GOETHE'S BOYHOOD: being the First Three Books of his Autobiography. Arranged and Annotated by the same Editor. 2s.

GOETHE'S HERMANN AND DOROTHEA. With an Introduction and Notes. By the same Editor. Revised edition by J. W. CARTMELL, M.A. 3s. 6d.

"The notes are among the best that we know, with the reservation that they are often too abundant."—*Academy*.

GUTZKOW. ZOPF UND SCHWERT. Lustspiel in fünf Aufzügen von. With a Biographical and Historical Introduction, English Notes, and an Index. By H. J. WOLSTENHOLME, B.A. (Lond.). 3s. 6d.

"We are glad to be able to notice a careful edition of K. Gutzkow's amusing comedy 'Zopf und Schwert' by Mr H. J. Wolstenholme. . . . These notes are abundant and contain references to standard grammatical works."—*Academy*.

HAUFF. DAS BILD DES KAISERS. Edited by KARL HERMANN BREUL, M.A., Ph.D. 3s.

HAUFF. DAS WIRTHSHAUS IM SPESSART. Edited by A. SCHLOTTMANN, Ph.D., late Assistant Master at Uppingham School. 3s. 6d.

HAUFF. DIE KARAVANE. Edited with Notes by A. SCHLOTTMANN, Ph.D. 3s. 6d.

IMMERMANN. DER OBERHOF. A Tale of Westphalian Life. With a Life of Immermann and English Notes, by WILHELM WAGNER, Ph.D., late Professor at the Johanneum, Hamburg. 3s.

KOHLRAUSCH. Das Jahr 1813 (THE YEAR 1813). With English Notes. By W. WAGNER. 2s.

LESSING AND GELLERT. SELECTED FABLES. Edited with Notes by KARL HERMANN BREUL, M.A., Lecturer in German at the University of Cambridge. 3s.

MENDELSSOHN'S LETTERS. Selections from. Edited by JAMES SIME, M.A. 3s.

RAUMER. Der erste Kreuzzug (THE FIRST CRUSADE). Condensed from the Author's 'History of the Hohenstaufen', with a life of RAUMER, two Plans and English Notes. By W. WAGNER. 2s.

"Certainly no more interesting book could be made the subject of examinations. The story of the First Crusade has an undying interest. The notes are, on the whole, good."—*Educational Times*.

RIEHL. CULTURGESCHICHTLICHE NOVELLEN. With Grammatical, Philological, and Historical Notes, and a Complete Index, by H. J. WOLSTENHOLME, B.A. (Lond.). 3s. 6d.

SCHILLER. WILHELM TELL. Edited with Introduction and Notes by KARL HERMANN BREUL, M.A., University Lecturer in German. 2s. 6d.

UHLAND. ERNST, HERZOG VON SCHWABEN. With Introduction and Notes. By H. J. WOLSTENHOLME, B.A. (Lond.), Lecturer in German at Newnham College, Cambridge. 3s. 6d.

V. ENGLISH.

ANCIENT PHILOSOPHY. A SKETCH OF, FROM THALES TO CICERO, by JOSEPH B. MAYOR, M.A. 3s. 6d.

"Professor Mayor contributes to the Pitt Press Series *A Sketch of Ancient Philosophy* in which he has endeavoured to give a general view of the philosophical systems illustrated by the genius of the masters of metaphysical and ethical science from Thales to Cicero. In the course of his sketch he takes occasion to give concise analyses of Plato's Republic, and of the Ethics and Politics of Aristotle; and these abstracts will be to some readers not the least useful portions of the book."—*The Guardian*.

ARISTOTLE. OUTLINES OF THE PHILOSOPHY OF. Compiled by EDWIN WALLACE, M.A., LL.D. (St Andrews), late Fellow of Worcester College, Oxford. Third Edition Enlarged. 4s. 6d.

"A judicious selection of characteristic passages, arranged in paragraphs, each of which is preceded by a masterly and perspicuous English analysis."—*Scotsman*.

"Gives in a comparatively small compass a very good sketch of Aristotle's teaching."—*Sat. Review*.

BACON'S HISTORY OF THE REIGN OF KING HENRY VII. With Notes by the Rev. J. RAWSON LUMBY, D.D. 3s.

COWLEY'S ESSAYS. With Introduction and Notes. By the Rev. J. RAWSON LUMBY, D.D., Norrisian Professor of Divinity; Fellow of St Catharine's College. 4s.

MORE'S HISTORY OF KING RICHARD III. Edited with Notes, Glossary and Index of Names. By J. RAWSON LUMBY, D.D. to which is added the conclusion of the History of King Richard III. as given in the continuation of Hardyng's Chronicle, London, 1543. 3s. 6d.

MORE'S UTOPIA. With Notes by the Rev. J. RAWSON LUMBY, D.D. 3s. 6d.

"It was originally written in Latin and does not find a place on ordinary bookshelves. A very great boon has therefore been conferred on the general English reader by the managers of the *Pitt Press Series*, in the issue of a convenient little volume of *More's Utopia* not in the original Latin, but in the quaint *English Translation thereof made by Raphe Robynson*, which adds a linguistic interest to the intrinsic merit of the work. . . . All this has been edited in a most complete and scholarly fashion by Dr J. R. Lumby, the Norrisian Professor of Divinity, whose name alone is a sufficient warrant for its accuracy. It is a real addition to the modern stock of classical English literature."—*Guardian*.

THE TWO NOBLE KINSMEN, edited with Introduction and Notes by the Rev. Professor SKEAT, Litt.D., formerly Fellow of Christ's College, Cambridge. 3s. 6d.

"This edition of a play that is well worth study, for more reasons than one, by so careful a scholar as Mr Skeat, deserves a hearty welcome."—*Athenæum*.

"Mr Skeat is a conscientious editor, and has left no difficulty unexplained."—*Times*.

VI. EDUCATIONAL SCIENCE.

COMENIUS. JOHN AMOS, Bishop of the Moravians. His Life and Educational Works, by S. S. LAURIE, M.A., F.R.S.E., Professor of the Institutes and History of Education in the University of Edinburgh. New Edition, revised. 3s. 6d.

London: C. F. CLAY & SONS, Cambridge University Press Warehouse, Ave Maria Lane.

EDUCATION. THREE LECTURES ON THE PRACTICE OF. I. On Marking, by H. W. EVE, M.A. II. On Stimulus, by A. SIDGWICK, M.A. III. On the Teaching of Latin Verse Composition, by E. A. ABBOTT, D.D. 2s.

LOCKE ON EDUCATION. With Introduction and Notes by the Rev. R. H. QUICK, M.A. 3s. 6d.

"The work before us leaves nothing to be desired. It is of convenient form and reasonable price, accurately printed, and accompanied by notes which are admirable. There is no teacher too young to find this book interesting; there is no teacher too old to find it profitable."—*The School Bulletin, New York.*

MILTON'S TRACTATE ON EDUCATION. A facsimile reprint from the Edition of 1673. Edited, with Introduction and Notes, by OSCAR BROWNING, M.A. 2s.

"A separate reprint of Milton's famous letter to Master Samuel Hartlib was a desideratum, and we are grateful to Mr Browning for his elegant and scholarly edition, to which is prefixed the careful *résumé* of the work given in his 'History of Educational Theories.'"—*Journal of Education.*

MODERN LANGUAGES. LECTURES ON THE TEACHING OF, delivered in the University of Cambridge in the Lent Term, 1887. By C. COLBECK, M.A., Assistant Master of Harrow School. 2s.

ON STIMULUS. A Lecture delivered for the Teachers' Training Syndicate at Cambridge, May 1882, by A. SIDGWICK, M.A. 1s.

TEACHER. GENERAL AIMS OF THE, AND FORM MANAGEMENT. Two Lectures delivered in the University of Cambridge in the Lent Term, 1883, by Archdeacon FARRAR, D.D., and R. B. POOLE, B.D. Head Master of Bedford Modern School. 1s. 6d.

TEACHING. THEORY AND PRACTICE OF. By the Rev. EDWARD THRING, M.A., late Head Master of Uppingham School and Fellow of King's College, Cambridge. New Edition. 4s. 6d.

"Any attempt to summarize the contents of the volume would fail to give our readers a taste of the pleasure that its perusal has given us."—*Journal of Education.*

BRITISH INDIA, A SHORT HISTORY OF. By Rev. E. S. CARLOS, M.A., late Head Master of Exeter Grammar School. 1s.

GEOGRAPHY, ELEMENTARY COMMERCIAL. A Sketch of the Commodities and the Countries of the World. By H. R. MILL, Sc.D., F.R.S.E., Lecturer on Commercial Geography in the Heriot-Watt College, Edinburgh. 1s.

AN ATLAS OF COMMERCIAL GEOGRAPHY. Intended as a Companion to the above. By J. G. BARTHOLOMEW, F.R.G.S. With an Introduction by Dr H. R. MILL. 3s.

VII. MATHEMATICS.

EUCLID'S ELEMENTS OF GEOMETRY. BOOKS I. & II. Edited by H. M. TAYLOR, M.A., Fellow and formerly Tutor of Trinity College, Cambridge. 1s. 6d.

[Other Volumes are in preparation.]

London: C. F. CLAY & SONS, Cambridge University Press Warehouse,
Ave Maria Lane.

University of Cambridge.

LOCAL EXAMINATIONS.

Examination Papers, for various years, with the *Regulations for the Examination*. Demy 8vo. 2s. each, or by Post 2s. 2d.

Class Lists, for various years, Boys 1s., Girls 6d.

Annual Reports of the Syndicate, with Supplementary Tables showing the success and failure of the Candidates. 2s. each, by Post 2s. 3d.

HIGHER LOCAL EXAMINATIONS.

Examination Papers for various years, *to which are added the Regulations for the Examination*. Demy 8vo. 2s. each, by Post 2s. 2d.

Class Lists, for various years. 1s. each. By Post 1s. 2d.

Reports of the Syndicate. Demy 8vo. 1s., by Post 1s. 2d.

LOCAL LECTURES SYNDICATE.

Calendar for the years 1875—80. Fcap. 8vo. cloth. 2s.; for 1880—81. 1s.

TEACHERS' TRAINING SYNDICATE.

Examination Papers for various years, *to which are added the Regulations for the Examination*. Demy 8vo. 6d., by Post 7d.

CAMBRIDGE UNIVERSITY REPORTER.

Published by Authority.

Containing all the Official Notices of the University, Reports of Discussions in the Schools, and Proceedings of the Cambridge Philosophical, Antiquarian and Philological Societies. 3d. weekly.

CAMBRIDGE UNIVERSITY EXAMINATION PAPERS.

These Papers are published in occasional numbers every Term, and in volumes for the Academical year.

VOL. XVI. Parts 44 to 65. PAPERS for the Year 1886—87, 15s. cloth.

VOL. XVII. „ 65 to 86. „ „ 1887—88, 15s. cloth.

VOL. XVIII. „ 87 to 107. „ „ 1888—89, 15s. cloth.

Oxford and Cambridge Schools Examinations.

Papers set in the Examination for Certificates, July, 1888. 2s. 6d.

List of Candidates who obtained Certificates at the Examination held in 1889; and Supplementary Tables. 6d.

Regulations of the Board for 1890. 9d.

Regulations for the Commercial Certificate, 1890. 3d.

Report of the Board for the year ending Oct. 31, 1889. 1s.

Studies from the Morphological Laboratory in the Uni-

versity of Cambridge. Edited by ADAM SEDGWICK, M.A., Fellow and Lecturer of Trinity College, Cambridge. Vol. II. Part I. Royal 8vo. 10s.

Vol. II. Part II. 7s. 6d. Vol. III. Part I. 7s. 6d. Vol. III. Part II. 7s. 6d.

Vol. IV. Part I. 12s. 6d. Vol. IV. Part II. 10s. Vol. IV. Part III. 5s.

London: C. J. CLAY AND SONS,

CAMBRIDGE UNIVERSITY PRESS WAREHOUSE,

AVE MARIA LANE.

GLASGOW: 263, ARGYLE STREET.

