

**How to feed the horse, avoid disease and save money / by George Armatage.**

**Contributors**

Armatage, George.

**Publication/Creation**

London and New York : F. Warne and co., 1890.

**Persistent URL**

<https://wellcomecollection.org/works/n24zej6b>

**License and attribution**

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.


You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>

# How to Feed THE HORSE, avoid Disease and Save Money.

BY  
George  
Armatage



LONDON & NEW YORK  
FREDERICK WARNE & CO





*Price 1s. each.*

## WARNE'S USEFUL BOOKS.

A Series of Handy and Practical Guides on Popular Subjects.

*In fcap. or crown 8vo, cloth, or picture boards.*

HANDBOOKS FOR THE COUNTRY AND SEASIDE.

**OUT-DOOR COMMON BIRDS:** Their Habits and General Characteristics.

By HENRY STANNARD. With Eighty Original Illustrations.

"The habits, appearance and peculiarities of the birds are graphically and minutely described."—*Graphic*.

**THE HORSE:** Its Varieties and Management in Health and Disease.

"The whole duty of man as regards horses may be found in these pages."—

*Sporting Life.*

**CATTLE:** Their Varieties and Management in Health and Disease. By

GEORGE ARMATAGE, M.R.C.V.S. With Illustrations.

"Cheap, portable, neatly got-up, and full of varied information, and contains useful facts as to habits, training, breeding, &c."—*Sporting Gazette*.

**THE SHEEP:** Its Varieties and Management in Health and Disease. By

GEORGE ARMATAGE, M.R.C.V.S. With Illustrations.

"This handbook is thoroughly trustworthy and can be recommended to any farmer with perfect confidence."—*The Times*.

**THE BRITISH BIRD PRESERVER.** With Practical Instructions on How

to Skin, Stuff, and Mount Birds and Animals, with a Chapter on their Localities, Habits, and How to Obtain Them; also Instructions in Moth and Butterfly Catching, Setting and Preserving. By SAMUEL WOOD, Naturalist and Professional Bird Preserver.

"A more complete, handy little manual, and one more thoroughly adapted to the purposes it has in view, we are not acquainted with."—*Canterbury Gazette*.

"The instructions are detailed and minute."—*Scotsman*.

**BIRD-KEEPING:** A Practical Guide for the Management of Cage and Singing Birds. By C. E. DYSON. With Thirty Illustrations.

"This book will be heartily welcomed by owners of cage pets. Almost every bird suitable for an aviary is fully described, with simple instructions for their management in sickness as well as health."—*Pall Mall Gazette*.

**THE DOG:** Its Varieties and Management in Health and Disease. By STONEHENGE. With numerous and Choice Wood Engravings.

"There is no handier volume issued on the subject than this excellent little manual."—*Field*.

**HAPPY PLANTS FOR LITTLE FRONT GARDENS.** By S. STACKHOUSE.

save the cultivator  
und in the present

se persons who, re-  
for gardening under

tructions on the  
VATTS.

to become growers of flowers."—*Weekly Times*.



22101886615



Med

K51314

ICAL HANDBOOKS.

2

ice 1s. each.

SEFUL BOOKS—*continued*.

TO GROW THEM: A Guide to the Kitchen  
Garden. By E. WATTS.

This Book is the property of

THE WELLCOME PHYSIOLOGICAL  
RESEARCH LABORATORIES,  
BROCKWELL HALL,  
HERNE HILL, LONDON, S.E.

Anyone finding and returning it to the above  
address will be handsomely rewarded.

*The Standard Gardener* is a simple and handy guide to the various everyday  
operations required to ensure success in gardening."

THE STANDARD MANUAL ON SINGING.  
ADVICE TO SINGERS. By F. J. CROWEST. Fourth Edition.

"We cordially recommend a new and enlarged edition of 'Advice to Singers' as  
a little book deserving to be in the hands of all to whom it is addressed. The general  
advice is very sensible."—*The Guardian*.



M. xx 19/a

868



1. D. 7875-

# HOW TO FEED THE HORSE



HOW TO FEED THE HORSE



# HOW TO FEED THE HORSE

*AVOID DISEASE AND SAVE MONEY*

BY

GEORGE ARMATAGE

FORMERLY LECTURER IN VETERINARY SCIENCE IN THE ALBERT AND GLASGOW  
VETERINARY COLLEGES

AUTHOR OF

"THE HORSE DOCTOR," "THE CATTLE DOCTOR," "THE HORSEOWNER  
AND STABLEMAN'S COMPANION," ETC.

Third Edition, Revised



LONDON AND NEW YORK  
FREDERICK WARNE AND CO.

1890

7 425348



WELLCOME INSTITUTE LIBRARY	
Coll.	welMOmec
Call No.	
	✓



## PREFACE TO THE THIRD EDITION.



SINCE the appearance of this work in 1868, much satisfaction and encouragement has been gained from the numerous instances in which horse proprietors have been led to look into the subject of feeding. While the immediate object has been that of reducing actual cost, the system, when faithfully carried out, has yielded more than was sought for. It has increased the working ability of the animals treated by conferring perfect health, and, as an inevitable outcome, common diseases have been abolished, and the natural term of life considerably lengthened. This is acting on the wiser side of prevention, always

safe and more satisfactory than courting disease and gaining credit by its cure.

The interest which the subject has awakened is seen in various ways. There is scarcely a retail corn chandler who has not his special "Horse Mixture" for sale. This must be a great benefit to many small proprietors, when such is compounded with honest material and in honest proportions. Several who have adopted our advice in this respect have earned good reputation and lucrative returns. In addition, firms of millers have extended their premises and machinery, and, by the aid of our little manual, they have worked up a large connection, sending annually hundreds of tons of "Horse Mixtures" over the entire country.

Happening to make a call in a country town on one occasion, and being in the company of one of the leading tradesmen, the subject of feeding was introduced by himself. We preserved



---

silence, and allowed him to enlighten us. At first he claimed originality of thought and design, but added there were points he had not been quite able to clear up. We ventured to state that his line of investigation seemed to be on parallel lines with those of the writer, naming the present work. We also pointed to his show-cards, announcing various foods for pheasants, poultry, pigs, &c. &c., and the prizes obtained at certain agricultural shows. At first he was disinclined to acknowledge the source of his information, but on our claiming the authorship of the manual, he brought it from the secret recesses of his desk and made humble confession, which was duly acknowledged by leaving with him a few more hints. We sincerely hope his popularity as a benefactor to winged bipeds, as well as porkers, may bring its desired pecuniary reward.

These are a few of the means of usefulness

permitted to previous editions of the manual. The Third Edition is issued with the desire that similar results may follow. Let men grow rich, but let animal suffering grow less.

GEORGE ARMATAGE.

*5th Month, 1890.*



# CONTENTS.

## PART I.

	PAGE
MISMANAGEMENT . . . . .	1
DEVELOPMENT AND MAINTENANCE:	
Immediate Object of Food—Development—Maintenance—Waste or Decay—Importance of Good Food for Young Animals—Evils of an Insufficiency . . .	8
ORGANS OF DIGESTION :	
Enumeration—Prehension—Incisor Teeth—Tongue—Cheeks—Molars or Grinders—Pharynx—Gullet—Deglutition . . . . .	14
INSALIVATION :	
Saliva—Salivary Glands—Solvent Action—Chemical Action—Quantity of Saliva—Uses and Importance of Saliva . . . . .	16
STOMACH :	
Peculiarities in the Horse, Ox, and Man—Broken Wind, and Chronic Cough—Rapid Digestion in the Horse—Small Stomach—Necessity for Regular Feeding . . .	20
INTESTINES :	
Enumeration and Division—Length—Small Intestines—Digestion in—Secretions—Chyle—Lacteals—Absorption and Assimilation—Capacity—Large Intestines—Division—Absorption—Capacity . . . . .	23



## THE DIGESTIVE PROCESS:

PAGE

- A Complex Action—Gastric Digestion—Gastric Juice  
—Intestinal Digestion—Chyle—Formation of Blood 26

## ELEMENTARY PRINCIPLES OF FOOD:

- Enumeration—Flesh Formers—Heat Producers—Salts  
—Acids—Fæces or Dung . . . . . 28

## ESSENTIAL CHARACTERS OF FOOD:

- Nutritious Principles—Their Identity from all Sources  
—Non-nutritious Principles—Animal Heat—Neces-  
sity for Substances of a Mixed Character—Bulk or  
Volume . . . . . 29

## ERRORS TO BE AVOIDED:

- Fluids—Cooked Food—Injurious Effect of Common  
Salt — Diabetes and Albuminuria — Farcy and  
Glanders . . . . . 40

## ADVANTAGES OF PROPER FOOD AND SYSTEM . . . 46

## REGULAR FEEDING . . . . . 50

## CALCULI OR STONES:

- Their Origin—Different Kinds—Frequently Prove Fatal 51

## OBJECTIONS TO THE USE OF DRY FOOD:

- Do Oats Pass Out Unchanged?—Proofs—Necessity for  
Healthy Condition—Care Required in Feeding—  
Importance of a Superintendent . . . . . 54

## PART II.

## VARIETIES OF FOOD:

- Nutrition in Each—Maize—Oats—Cost of Feeding  
Upon—Measure and Weight should be Combined 63

## SELECTION AND PURCHASE OF GRAIN . . . . . 68

## ECONOMY OF FOOD:

- Indian Corn Injurious — Linseed — Tares — Mixtures  
Necessary . . . . . 70



---

SYSTEMS ADOPTED ON VARIOUS COLLIERIES :	PAGE
Hunting on Cut Food—Hetton System . . .	74
WHAT CONSTITUTES A CHEAP FOOD . . .	81
OBJECTIONS TO A CHANGE OF GRAIN . . .	82
BRAN, ITS NATURE, USES, AND ABUSES . . .	83
LONDONDERRY COLLIERY SYSTEM :	
Mixture—Steamed Food—Waste, &c. . . .	85
FORMS OF ADMIXTURE :	
Low Rates—High Rates . . . . .	89
GREEN FOOD, ITS BENEFIT AND INJURY . . .	97
ECONOMY IN USING CHAFF AND BRUISED CORN .	99
PEA AND BEAN STRAW . . . . .	101
SAVING TO BE EFFECTED . . . . .	102
PRACTICAL TESTS OF THE SYSTEM . . . .	104
WORK . . . . .	116
IMPORTANCE OF GROOMING . . . . .	123





# HOW TO FEED THE HORSE.



## PART I.

### MISMANAGEMENT.

WE trace to this source the rapid deterioration, prevalence of disease, and short life of the animals comprising the studs of many public companies, the numerous losses among farm horses, and those of large proprietors. It is the bane which robs the poor man of his only animal.

Mismanagement assumes a variety of forms. It will be sufficient for present purposes if it is simply stated with respect to Food and Work.

The character of food in many instances decides the infliction of actual cruelty in more ways than one. Horses are injudiciously fed. The system either takes the form of a lavish abundance or niggardly parsimony in kind or quantity of the aliment, with the view of bringing the actual cost within certain limits, without reference to the nutritious value of the food. In some instances no respect is paid to the quality, and sadly inferior articles are supposed to make



up for all deficiencies by an extra allowance. Even when good food is supplied, and due care is lacking in other respects, the results are not always of a satisfactory nature, as the prevalence of indigestion in its many forms so abundantly proves. Veterinary surgeons in many districts testify that the bulk of their cases arise from this cause alone, not a few sudden deaths being caused by rupture of the diaphragm, stomach, or intestines. At one period in the experience of the writer, the number of deaths among horses, from the presence of calculi or stones, the evidence of a life-long injudicious mode of feeding, was almost incredible. It had also escaped observation that the attendant evils of unwise systems created the necessity for medicines, and these, combined with the operation of unsanitary conditions, so impaired the system as to cause grease, farcy, and glanders to be constantly present in many stables.

And the category is far from complete. The London brewers suffer the loss of their plump dray-horses from disease and rupture of the liver, due to engorgement, the result of overfeeding. In Scotland, the exclusive use of boiled mixtures of food, besides sometimes inducing similar results, augment the casualties by the number of deaths from colic, and a form of albuminuria still more rapid and fatal; while grease, farcy, and glanders are sometimes fearfully prevalent.



Among the large studs employed in coal-mines, notwithstanding the animals are in many cases more advantageously situated, the deaths from various causes are the result of mainly preventible diseases, accidents being of course excepted.

The character of work has also its share in the multiplication of evils which cut short the life of working animals. As they are fed according to a fallacious method, which recognises only actual cost, the same means are employed to decide how much work should be done without reference to health or condition. When this is added to "penny wise and pound foolish" ways of feeding, the result is a catastrophe, representing so much financial loss. The close connection that exists between food and work is not generally understood as it should be. It is often altogether ignored, and even deplorable results fail to secure a change to improved systems. We have known frequent deaths in draught horses to arise from rupture of the diaphragm taking place during severe exertion after a full meal. Notwithstanding the fullest reports as to cause and prevention, the evil was not abated. The same has occurred in many instances with regard to other preventible diseases. The errors of feeding and work have been pointed out, yet the evils continue, the proprietors in some way settling down to the practice of considering the matter fully ac-



counted for when the value of the dead creature is posted to the debit of "P. and L. account."

Perhaps the most egregious form of mismanagement is to be witnessed in connection with some large establishments, where "grandeeism" rides roughshod over common-sense and practicability in beings of a supposed lower grade. If the secret history of the downward progress of many defunct public enterprises could be written, the first cause would be traced to some form of mismanagement, generally of an intentional character and designed for personal benefit. In one instance out of several we could name, a whole stud of horses, purchased for and by the "promoter" of a public company, were mostly inferior and unsound animals; yet the company paid an exorbitant price, the "promoter" pocketing a considerable sum. The disadvantages arising from working a number of animals which soon became confirmed and incurable cripples were so apparent that "grandeeism" determined to recoup the failing exchequer by taking the feeding into his own hands. From a daily allowance of twenty pounds of rich grain mixture, the creatures were reduced to one of twelve, containing not more than a third of the nutritious value of the former. It also consisted chiefly of maize, to which the horses were not accustomed. This they refused for weeks; meanwhile they lost flesh under the infliction of the hardest work,



and became very weak. The orders were that they *should be starved*, and thus forced to eat the maize or go without. The small proportion of oats was consumed with the chaff, but being sadly insufficient to sustain the vital powers, disease set in, and accomplished its work most grievously. Several animals fell dead as they were taken from the stables to work, and others contracted farcy and glanders, and were destroyed; others were seized with a malignant form of purpura hæmorrhagica and died. For two years previously, during which the concern had been in existence, not a case of either farcy or glanders had been seen in the whole of the stables; from that time to the present they have not been free. During one period of that time the company paid no dividend: it afterwards yielded to the demands of the shareholders, and paid a small acknowledgment. Somewhat later the shareholders looked more closely into existing matters, and the result was that "grandeeism" was dethroned, the paternal care of the State being graciously extended for a time, and since his liberation he has found it more convenient to reside abroad.

In a paper read before the members of a northern counties farmers' club, we gave, among numerous instances, the highly successful results in feeding certain colliery horses under the superintendence of the resident veterinary surgeon, the annual saving being many hundreds



of pounds. The local newspapers gave full reports, at which "grandeeism" grew jealous, and immediately took the feeding into his own hands, returning to the old as well as unsuitable and more expensive system.

The attention of thinking men had, however, been drawn to the question, and Mr. Hunting, M.R.C.V.S. of South Heston, was invited by the Newcastle Farmers' Club to give the results of his splendid system. Again the usual feeling of malice prevailed, for some person sent a parcel of hay-seeds and rubbish to the chairman, with the intimation that the sample had been taken from the kind of provender used by Mr. Hunting, and which he would recommend to the meeting.

Thus during many years it was impossible to make desirable progress in the spread of information on the subject of systematic and profitable feeding hard-working animals. In many instances the feeding was in the hands of those who were averse to change, and to whom the purchase of grain meant so much gain. Although it came within the scope of duties imposed upon the veterinary surgeon, to report the need and advantage of change in matter, intimately affecting the well being of so many animals, and, at the same time, promising in a twofold degree, a certain financial success, he must remain silent. If he ventured to report or suggest to "grandeeism," he was told to mind his own business; if he went to head quarters, he was deposed,



iniquity triumphed, and ruinously expensive systems prevailed.

The principles upon which working animals should be fed are those which rest upon scientific data. When we recognise this, the system is elevated to the dignity of an art, abrogating the uncertainties of the "rule of thumb," and ensuring results of a definite and useful character in the maintenance of the general health of the stock, and a corresponding increase over former states in the power of endurance. "Wear and tear" being also greatly reduced, the average life of each animal is augmented; the cost per head is correspondingly reduced, and thus combined advantages are derived.

Ample opportunity of verifying these statements will be found wherever the plan is adopted, whether the stud be large or small. The results are proportionate to the faithfulness with which the rules are carried out. The system propounded in the following pages is essentially one of prevention. The mistake of the past has consisted in our seeking for cures and ignoring causes. We should aim at conserving the forces at our command. While we maintain these in their fulness of vigour, the animal is in the enjoyment of perfect health and spirits. He possesses the greatest aptitude for work, and in a state most likely to resist the more common modes of attack by disease.



---

## DEVELOPMENT AND MAINTENANCE.

A discussion of the subject of food and its application to the animal fabric, involves a consideration of scientific principles which reveal the exact nature of the substances employed, their transformation within the organism, and the offices they perform at their destination.

The *immediate object of food* is the development and maintenance of the animal body.

*Development* may be briefly stated to be that process by which the various parts or organs assume their relative form, size, and capabilities for functional activity—*e.g.*, secretion, excretion, and the like. Development is principally referable to intra-uterine life, or that period during which the future animal lies within the womb of its mother, when the various organs, as the brain, heart, lungs, liver, and in fact all parts of the body, are acquiring their peculiar and characteristic form.

Each primitive portion gradually and constantly receives fresh additions, and each stage constitutes also a farther elaboration and assumption of higher powers.

Development continues also throughout a period after birth. The long-legged, weedy-looking foal is an example excellently suitable for the purpose of illustration. Although as far as internal organs are concerned, development may be said to be in a measure complete, yet there



are important changes to be otherwise effected. The muscles of the body are small and pale, and the bones (so called) are very deficient in osseous material (earthy salts). A few teeth only are through the gums, and others are lying in different stages beneath in the form of a highly vascular pulp, quite unlike its future self.

In each there are important changes to take place.

The bundles of cartilage in the centre of the limbs and beneath the muscles of the body, &c., gradually become harder, and lose their characteristic elasticity. Their structure is altered by the deposition of bony material, and by and bye we have the osseous framework or skeleton completed.

The pale muscular fibres assume greater dimensions, and at the same time a darker colour, and power to act more forcibly on the individual bones.

The tooth pulp within the gums, already provided with innumerable blood vessels and nerves, gradually acquires the shape of a tooth, incisor or molar, its different layers of hard substance variously termed dentine, enamel, &c., and only at a specified time will it appear above the surface.

At length the animal is "filled up," "made up," or "furnished," in stable phraseology, and the period of *youth* gives way to that which is known as the *adult stage*.



*Maintenance* has been going on throughout the whole of this time. Functional activity is attended by a process of *waste* or *decay* of the parts implicated. This process will be hereafter expressed as *metamorphosis of tissues*.

In no period of life can this process be said to be dormant or non-existent, but in none so slow or so little required as in early youth. At this time the building up of the animal body by the accumulation of blood, bone, muscle, hoof, horn, teeth, hair, &c., &c., is actively carried on, and their waste or decay comparatively small. In adult life, however, when development has gone on to completion, and great exertion is called forth, metamorphosis or change in the constitution of the tissues above mentioned is great.

This requires the process of maintenance or repair to restore, during repose, that which has been constantly lost during activity or exercise.

The movements of the arms or legs in walking, the tongue and jaws in speaking, lungs during respiration, heart in its beating, and intestines during contraction,—in short, any act, voluntary or involuntary, which calls forth muscular movements in any part of the animal body, is attended with the expenditure of vital force, as shown in the change or alteration of the condition, waste, decay, or metamorphosis of the composing material.

During metamorphosis of tissue chemical



action is instituted, and new compounds are produced which are no longer useful to the body. They are called *effete* materials, and after being collected by appropriate vessels termed *lymphatics*, are by them conveyed to special organs for the purpose of being expelled. If retained, they would prove highly detrimental to the body, and even destructive to life.

The *waste of muscular tissue* from work and ordinary exercise is considerable, that substance forming a great portion of the animal body. It is constantly in need of fresh material to restore or replace those parts which have been removed by waste, and the elements necessary are provided by the food, after undergoing important changes hereafter to be mentioned under digestion.

Such facts, briefly as they are detailed, and divested, as far as practicable, of technicalities, present most important points to view, and enable us to direct particular attention to the application and economising of material (food) for the production of the greatest amount of actual force—strength and vigour in working animals, or of fat in those destined for the butcher.

With these before us, the effects of some of the modes in which animals are kept, will be clearly understood.

Under the subject of development it has been stated the process is confined to intra-uterine



life and early periods of youth. It is then we find the great demands upon the system in order to complete the formation and building up of the whole body. Every one almost knows that much more food than ordinarily is consumed by the pregnant mare or cow, and that the young animal consumes much food of a nutritious quality without the appearance being in all cases obviously benefited thereby.

These illustrations serve to show the great demands for blood, bone, muscle, skin, hair, hoof, tendons, &c. &c., and a moment's reflection will cause one to pause and wonder how some persons can really expect to make a horse, cow, or other animal upon the quantities of miserable rubbish which are put in the poor creatures' way when young.

How they can witness their colts and calves standing without cover in an empty straw-yard or bare pasture in the cold and rain during the winter, and expect the grass of the coming summer will make up for the previous starvation, is indeed a paradox. It appears quite sufficient with some that a horse should taste corn only when he is able to earn it, and cattle when they commence to give milk, or there are prospects of their being useful to the butcher.

All this is mistaken policy. The young horse or ox requires ample food for the building up of the frame, and we discern the wise provision of nature in furnishing the rich elements in the



milk of the mother. Both need highly nutritious food, and when this is denied, the owner finds he is on the sure road towards spoiling them. They always suffer acutely when good food is afterwards given; the change serves only to engender serious, if not fatal, disease. Instances will be found in the maladies *black-leg* or *quarter-evil*, *splenic apoplexy*, *blain*, &c., of cattle and sheep.

In the horse we find, when put to work for the first time, he is "unable to stand the corn;" the legs swell, coat stares, he performs his work badly, and there are other indications of a disordered state within.

*Colic*, or *purpura hæmorrhagica*, speedily kills him, or he is left to dole out a miserable existence with protracted suffering from *chronic disease of the lungs* or *liver*, *diabetes*, *canker*, *laminitis* (founder), constant attacks of *lymphangitis* (weed), and *farcy*, terminating, after effectually propagating the contagion, in *glanders* and *death*.

As life and development proceeds, food then serves a different purpose. While it is expended in supplying fresh elements to the blood, that fluid has now to furnish elaborate materials to replace the constituents of the body lost during waste of muscle and other tissues, which are rendered visible in the shape of the products of respiration, perspiration, the urine, and fæces. What formerly went to form



muscle, bone, &c., now goes to repair them. The blood in all cases is the fluid which furnishes the *pabulum* whence tissues derive their support. That fluid is maintained by regular additions of elements obtained from the food. The process which prepares them we have to consider shortly.

### ORGANS OF DIGESTION.

The organs of digestion comprise the *mouth*, *teeth*, *tongue*, *salivary glands*, *pharynx*, *æso-phagus* or *gullet*, *stomach*, *intestines*, *liver*, and *pancreas* or *sweetbread*.

In connexion with the mouth we find appropriate muscles forming the bulk of the lips and sides for the *prehension* or gathering of food, and *incisor teeth* or *nippers*—six above and below in the horse, and eight only below in the ox, sheep, deer, &c.—for the purpose of cutting off the herbage. Within the mouth the *tongue* performs the important office of distinguishing by the peculiar sense of taste, the difference in each variety of substances introduced, and moves them from side to side in order to bring them within the pressure of the *molar teeth* or *grinders*. The *tongue* at length, by a contraction upon itself backwards, carries the bolus of food with it, which is then seized by the *pharynx*, and passes immediately along the *gullet* to the stomach.

The *cheeks* on each side being formed of



muscles, assist materially in keeping the food effectually between the molar teeth. Through their substance pass ducts or tubes which convey necessary lubricating and solvent fluids to the mouth.

The *molars* or *grinders* are twenty-four in number, six of which are situated above and below upon each side of the jaws in the back part of the mouth. In order to observe them particularly the mouth must be held wide open.

They are large and rough upon the wearing surface, and form most perfect agents in crushing and grinding, or masticating the hard grain and kernels which this animal requires as natural food.

The *pharynx* is a musculo-membranous bag or pouch which opens at the back of the mouth, and grasps the bolus of food as it is presented by the tongue. Arrived at this point, its appropriate muscles successively contract and force onwards the pellet to the further end opening into the gullet. The food is now beyond the control of the will, and cannot be returned by any effort of a natural character.

The *æso-phagus* or *gullet* is also composed partly of muscles and membrane. It forms an elastic tube reaching from the pharynx above to the stomach below, and occupies a position in the neck upon the left side of the lower portion. A knowledge of this fact enables us to trace the passage of substances throughout its visible



course. The remaining part of the gullet passes through the chest, and is called the *thoracic* portion. The passage of food through the pharynx and gullet is termed *deglutition* or swallowing.

### INSALIVATION.

The mouth is abundantly supplied with fluid for the purpose of lubricating its surfaces during motion in mastication. It also possesses peculiar solvent and chemical actions upon alimentary matters introduced, and thus effects special changes in their constitution, of vital importance to digestion and assimilation.

*Saliva* is the name given to this fluid, and is the produce of glands in the vicinity, the largest, the parotid gland, being placed below the ear behind the angle of the lower jaw. In addition there are others of a smaller character situated beneath the tongue, between the sides of the jaw, inside the lips and other parts; all of which separately contribute important properties to the fluids which mix in the mouth, whence they are conveyed by appropriate tubes or ducts.

The *solvent action* of *saliva* depends upon the presence of peculiar salts and a principle called *ptyalin* by scientific men, and is easily demonstrated by placing food within the mouth. It is also found to act efficiently when collected in suitable vessels and brought into contact with



certain alimentary matters. Its power of altering the constitution of compounds is purely a *chemical action*, and is most peculiarly well marked upon those of a starchy character, abundance of which are found in the food upon which horses and cattle subsist. Through this influence they are converted into a principle called *dextrine*, and subsequently into *grape sugar*.

The salivary glands are under the control of the nervous system. In the human subject, the sight or smell of food of an agreeable character is sufficient to cause a copious flow to the mouth; hence the common expression "the mouth waters."

*Saliva is secreted abundantly.* The exact amount of this fluid which is formed is not easily obtained. A large horse is supposed to secrete 84lbs. per day, an ox 102lbs. Mr. Hunting obtained half a gallon, or about five pounds in half an hour.

The *uses of saliva* are important and manifold. First, we have an abundant flow; second, its peculiar solvent power; third, its chemical action in converting starchy matters into sugar; fourth, its lubricating qualities during the presence of absorbing, or coarse and rough, substances; fifth, *it is secreted in major part during mastication, and continues as long as that act is carried on.*

This category of properties in a simple-looking fluid like saliva must have been conferred for a beneficial purpose.



We find such to be a law throughout nature. That purpose is essentially the primary step in rendering the component parts of the food assimilable, *i.e.*, converting them into constituent parts of the blood.

Being secreted at the time when the food is undergoing comminution between the teeth, it is presented at a period when it will be most effective. Food, therefore, which is swallowed quickly or greedily, without proper mastication, obtains little saliva, and the necessary transformations are not carried out.

*It is impossible to supply a fluid artificially which can take the place of saliva.* Mere saturation of the food by water does not promote digestibility as a consequence.

This has been satisfactorily proved many times. If the salivary glands are prevented from discharging their contents into the mouth during mastication, and water is thrown among the mass within the stomach, digestion is retarded. We have positive evidence of this in daily practice in those establishments where owners persist in using boiled mixtures of food, which is done in utter neglect of the most important fact that the horse has perfect grinders to crush and break down everything which comes in the way of *natural* food, with the best solvent immediately at hand, and in unlimited quantity, to effect a primary transformation towards the production of blood, bone, muscle, hoof, hair, &c. &c.



The practice is an attempt to set aside nature, and might be excused if none of the above appliances are present, or their capabilities in part destroyed. We are so accustomed to treat the horse by analogy, thus convicting him in the exercise of irregular practices, errors of body and mind in common with ourselves, that we naturally prescribed a mode of treatment based upon conditions supposed to exist.

Even a moderate acquaintance with the organized fabric of man and the higher animals leads to a different conclusion. It proves the practice unscientific, unsystematic, foreign, and *unnatural*.

I shall have more to offer upon the question of boiled food when a description of other organs has been given.

After the food has undergone the necessary processes of mastication and insalivation, and, as it passes backwards in the act of swallowing or *deglutition*, it becomes coated with a thick mucous or viscous secretion, thrown out from glands on the inner surface of the pharynx and gullet. It accumulates as the mass descends, and forms a thick greasy kind of coating, the proceeding having for its object mainly the avoidance of aggregation and stoppage in the gullet, a condition which under aggravated states constitutes choking.\*

---

\* It not uncommonly happens after hard work and long fasts the horse, returning weary and hungry, ravenously swallows his



---

THE STOMACH.

Descending the gullet the bolus of food at length reaches the stomach, a pouch or dilatation of that tube. In the stomach further important changes are executed in the constitution of the food.

In man this organ performs the greater portion of the process of digestion. In the horse and ox but little analogy exists, and the essential points of difference will repay even a cursory examination.

The ox is endowed with four stomachs, or what is more in keeping with anatomical description, a quadrisect stomach, *i.e.*, containing four distinct compartments, each of which possesses different functions.

The first compartment is one of immense capacity. It occupies three-fourths of the abdomen, and is able to accommodate a great quantity of ingesta.

The second is small, and contains more fluid

---

food without due mastication and insalivation. The secretions are deficient by reason of fatigue or nervous depression and other causes. The bolus is speedily despatched, almost dry, and choking is apt to occur in its worst forms. If the food reaches the stomach aggravated colic is almost certain.

See my Prize Essay "On the Diseases of Farm Horses," published in the Transactions of the Highland and Agricultural Society of Scotland, Article—"Choking."



than the first, but both act in common in maccrating and securing successive changes in the food.

The third exerts a degree of pressure, by means of its peculiar leaves, upon the alimentary matters arrested by it.

The fourth is the true digestive compartment, and resembles the stomach of man and the dog principally.

The ox is essentially an animal capable only of limited and not rapid exertion. The structure and arrangement of the digestive organs are clearly intended to accommodate large quantities of herbage of a mixed character, for which the functions are admirably fitted.

The horse, on the other hand, being made and required for rapid exertion, could not fulfil those ends with the digestive organs of the ox, and, therefore, like man and the carnivora, but one stomach or compartment is provided, which is both comparatively and relatively smaller.

The process of digestion is also wonderfully effective.

Whilst the ox has been known to retain food for upwards of six weeks within his stomach, little more than as many hours will suffice to ensure digestion in the horse. The process is, therefore, also wonderfully rapid.

The stomach of the horse, replete with food, acts as a great impediment to the action of the lungs, hence the great number of cases of *broken*



*wind* and *chronic cough*, which are known to occur from irregular principles of feeding.

The horse cannot occupy himself in rumination or digestion almost entirely as the ox. He is required to undertake the removal of heavy loads, or otherwise engaged in rapid exertion. A rapid digestion in his stomach is therefore a very wise provision to relieve the lungs during action.

I have purposely administered substances of different degrees of solubility, and possessing a strong odour, to animals about to be slaughtered, and have been particularly struck with the fact that, in the short space of time which has elapsed between the administration of the medicine and opening the stomach, not a trace beyond the odour of the substance could be detected. The paper in which it was wrapped was usually found in the stomach, but the medicine had passed several yards along the gut, frequently within one hour.

Similar facts are observable with regard to food. I have known horses and ponies suddenly killed in coal mines by a fall of portions of the roof within one or two hours after a meal, and on examining the stomach it has been observed that digestion has progressed rapidly and effectively in that organ, only a small portion, the most indigestible, remaining behind. The more easily digestible portion had usually passed along the intestines in distances varying from ten to twenty feet or more.



It is a rule that all portions of greater solubility in the secretions, *i.e.*, more digestible, first pass out from the stomach in health.

*Fluids also pass rapidly.* They are usually carried to the cæcum, a large gut situate about twenty yards from the stomach in large horses, twenty minutes being usually sufficient.

These illustrations definitely explain why a horse occupies so much time consuming food. Put him into the stable after a day's work, and he will be found eating the greater part of the night. A small stomach, capable of effective and rapid action, is quickly emptied of its contents, and the desire for food, therefore, becomes almost constant.

We should gather from this also the absolute necessity of a regular supply of food, with abolition of long fasts and full racks and mangers, when the animal returns to the stable.

### THE INTESTINES.

The intestines are of two kinds, small and large. In the horse they occupy the greater portion of the cavity of the abdomen, and constitute the largest part of the alimentary track.

The whole length may be estimated at about ninety feet, or ranging from twenty-six to thirty yards.



The *small intestines* commence at the stomach, and in their course receive the several names *duodenum*, *jejunum*, and *ilium*, the divisions being purely imaginary.

About five or six inches below the stomach are the orifices which discharge the secretions of the liver (bile) and pancreas (a fluid resembling saliva). Besides these, other special fluids are poured from the walls (coats) throughout the length of the canal.

The walls are strong and provided with muscular fibres, as the gullet and stomach, to produce the necessary contractions (vermicular or worm-like) in order to subject the contents (ingesta) to the action of the various fluids, as well as cause it to pass onwards.

Throughout the inner surfaces of the small intestines a number of peculiar vessels are seen to enter. Their office is to abstract the nutritious elements of the food, which after meeting with the various secretions in the tube, assume a whitish creamy consistence, and is termed *chyle*. The vessels here spoken of are termed *lacteals*. They communicate with other vessels and glands. In them the chyle as it passes onwards alters its constitution, and by successive stages assumes the character of the blood with which it is afterwards mixed. This constitutes the process of *assimilation*.

The small intestines are from fifty to sixty feet in length, and will accommodate from eight to eleventh gallons of fluid.



The *large intestines* of the horse are very capacious organs. The first, the *cæcum*, is a large conical pouch or bag, usually known as the *blind gut*. In it are collected principally the fluids drank, together with solid matters. The former occupy the extremity, the latter remain at the base. Absorption of fluids takes place from the *cæcum* in large proportion, and its contents are usually alkaline from the changes which take place in starchy constituents.

The *colon* is continued next in order to the *cæcum*. It is doubled upon itself, both parts throughout its entire length being united by intervening tissue, and traverses both sides and front of the abdomen twice, terminating in the rectum near the point from which it arises from the *cæcum* on the right side.

The *cæcum* and *colon* do not possess such thick muscular walls as the small intestines, but strong bands run longitudinally on four sides. These being considerably shorter than the intestines, have the effect of drawing them into puckers or folds, forming on the inner side a number of pouches, which assist in giving the characteristic form to the *fæces* or dung in the horse.

The *rectum* is very strong and muscular, but not so capacious, tolerably cylindrical, and terminates the alimentary track at the *anus* or fundament. Absorption from the large intestines is principally confined to the *cæcum*. The *colon*



and rectum minister but little to that process. The *capacity of the large intestines* greatly exceeds that of the small. The cæcum will contain about four gallons of fluid, the colon about twelve, and the rectum about three gallons.

### THE DIGESTIVE PROCESS.

The process of digestion is most complex and important, and deservedly receives an extended notice in all authentic works on animal physiology. It is impossible here to do more than briefly notice the leading stages, which are indispensable, but sufficient towards explaining generally the object for which these pages are written.

*Gastric digestion.*—In the stomach the food, already incorporated with the salivary and other secretions, is subjected to peculiar movements or contractions of the muscular walls, described usually as a “churning action.” It is thus moved from one part to another and further incorporated with secretions (gastric juice) derived from glands (gastric glands) situate in the walls of the organ.

The *gastric juice* possesses acid properties, and a peculiar principle termed *pepsine*, which with the action of the stomach effects a reduction of the food to a uniform mass. The particles are in a fine state of division, and albuminous prin-



ciples chiefly suffer chemical decomposition. At length the whole becomes a thick fluid and passes through the *pyloric* orifice of the stomach to the small intestines.

This fluid is called *chyme*. It contains nutritious matters in a state of mechanical suspension, others in chemical and simple solution, while a third variety are unacted upon from various causes.

Chyme proceeds along the intestines by virtue of the peculiar action of the organs. It first meets with fluids from special glands in the vicinity of the outlet from the stomach. Next with those from the liver and pancreas (sweetbread). By constant motion perfect incorporation is effected, and the following outline of changes may be observed.

- 1st. The acid character of chyme is neutralized—it is now alkaline.
- 2nd. Albuminous matters escaping from the stomach unchanged are transformed.
- 3rd. Starchy matters, unacted upon by saliva, are now effectually converted into sugar.
- 4th. Fatty bodies are emulsified or converted into a kind of soap.
- 5th. and lastly, all the nutritious principles have conferred upon them properties which facilitate their absorption and passage through the lacteals towards the blood, and the fluid mass now receives the name of *chyle*.



Chyle passes through the lacteals from the intestines. Its constituents gradually form a granular-looking mass, in which float a number of small vesicles or cells. By passage through an assemblage of bodies called *mesenteric glands*, the cells increase in number, and at length acquire colour, and eventually are poured into blood vessels near the heart. It is thus the blood becomes the pabulum whence all tissues are nourished. In the elements of food are found the materials which, after assimilation, partake of the characters of the tissues of the body; and the circulation of the blood explains how each kind receives its share.

#### ELEMENTARY PRINCIPLES OF FOOD.

The chyle contains the whole of the elements of the food. These are of a mixed character. First, we have *nitrogenous*, so-called because they contain largely the gas nitrogen, or *albuminous* principles which are strictly the elements from which muscle is developed—hence the term “flesh formers” which is applied to them. Second, we have *fatty bodies* and *saccharine* or sugary principles—starch, gum, sugar, &c.—*heat-producers*. Third, there are essential constituents in the form of *salts*—of lime, potash, soda, magnesia, &c. Fourth, various *acids*, as hydrochloric (muriatic acid or spirits of salt), nitric



(aquafortis), sulphuric (oil of vitriol), lactic and phosphoric, &c.

In a chemical point of view the acids are most powerful agents, and by union with various substances form compounds of vital importance.

Lastly, there are compounds of no service whatever to the system. These are principally indigestible bodies, such as ligneous or fibrous parts of plants and foreign bodies which have gained access to the digestive organs, cells, earthy matter, and portions of undigested food which have been present in excess. Liebig has termed these the incombustible and unburned parts of food. They are the refuse from the digestive process, and together with *effete* or useless parts thrown off from the system, constitute the *fæces* or *dung*.

#### ESSENTIAL CHARACTERS OF FOOD.

Certain conditions are essential for the proper digestion and assimilation of food. When properly understood, they indicate the principles of an economical system of feeding animals with a view towards the preservation of health and vigour under continued laborious exertion, as well as preventing an undue rate of mortality resulting from it, and at a low rate of cost.

These conditions are, first, that food should be nutritious ; second, that it contains elements of



a mixed character ; third, that it possesses proportionate bulk ; fourth, it should be regularly supplied ; and lastly, the digestive organs generally must be in a state of healthy action.

It is imperative that food possess *nutritious principles*. This must be apparent from the fact that from it all parts of the body are built up and being constantly renovated. In order to possess this property, bodies rich in nitrogen are requisite, which are denominated *nitrogenous*, *azotised*, *nutritious*, or *flesh-forming compounds*. Examples of these are *albumen*, *fibrine*, and *caseine*—terms which would indicate different substances. Chemically, however, they are alike in composition, but exhibit physically different appearances.

If wheaten flour be placed under a stream of cold water a sticky paste is left behind. This is called *gluten*, and is identical in chemical composition with the flesh of man and animals. It is the nitrogenous or albuminous principle of wheat flour.

If an egg be broken a glairy fluid escapes from the shell, which becomes white and coagulates into a solid mass when heated. This is *albumen*—gluten in another form, identical in composition, and is the nitrogenous principle of the egg from which the flesh, feathers, claws, &c., of the bird are formed.

If milk is taken and an acid added, a solid mass of curds is speedily produced. This is



*caseine*—or gluten, albumen, and fibrine in another form, supplied in the milk of the mother to young animals, and in peas, beans, &c., to older ones; from which also hoof, horn, hair, wool, skin, flesh, and feathers, &c. &c., are formed, developed, and repaired.

Lastly, if blood be coagulated, the mass which separates proves itself an identical compound to flesh itself—*fibrine*.

The whole then are the various forms in which the elements are conveyed to the blood, as found in the food upon which the animal subsists. By the action of vital processes within the system, each assumes those conditions which in themselves are vital. Their supply to the system must be constant in order to keep pace with the waste. Hence they are found abundantly in the food upon which man and animals live. Horses and cattle meet with them in the grass and corn which they daily consume, and man and carnivora obtain it directly by using the flesh and blood (*fibrine*) of animals as food.

The following table shows how these substances resemble each other in chemical composition:—

	Gluten from flour. Boussingault.	Caseine from peas. Scherer.	Albumen from eggs. Jones.	Ox-blood. Playfair.	Ox-flesh. Playfair.
Carbon . .	54.2	54.138	55.000	54.35	54.12
Hydrogen . .	7.5	7.156	7.073	7.50	7.89
Nitrogen . .	13.9	15.672	15.920	15.76	15.67
Oxygen . .	24.4	23.034	22.007	22.39	22.32
	<hr/> 100.0	<hr/> 100.000	<hr/> 100.000	<hr/> 100.000	<hr/> 100.000



### NON-NUTRITIOUS PRINCIPLES.

In addition to the flesh-forming constituents of food, there are also needed others known as *non-nitrogenous*, *non-azotised*, or *non-nutritious*, so named in contradistinction to those which contain nitrogen. Examples are found in starch, sugar, gum, and fat itself. They are composed of carbon, hydrogen, and oxygen, *minus* nitrogen, and are found abundantly in the different varieties of corn and vegetables used as food.

Their services are required in the system equally with the albuminous compounds, but for a different purpose, *viz.*, the production of animal heat, and formation and storing up of fat within the system.

### ANIMAL HEAT.

In order to render somewhat intelligible the principles upon which heat is developed, and maintained in the body, and the part which food plays in that process, attention must again be directed to facts.

Here it must be understood that although the term "heat producer" is applied to the saccharine principles of food, it by no means establishes an isolated fact. They are *not* the sole agents in the production of animal heat. I will attempt to explain.



To those who have studied chemistry, even but a little, the rapid development of heat unbearable in water previously cold is familiar. It is a good illustration for our purpose.

A quantity of cold water is taken in a tumbler or other thin glass vessel, and held in the hand, taking care to grasp that part in contact with the water. Sulphuric acid (oil of vitriol) is then added in nearly equal quantity, when by chemical action, which immediately takes place between the acid and water, intense heat is developed, rendering it an impossibility to hold the vessel in the manner adopted at the commencement.

The rubbing together of certain substances, or beating of metals upon hard surfaces, also produces great heat. Here the molecular disturbance—or the alteration of position of the ultimate particles composing the mass—which ensues from the concussion may not be unlike that which accompanies chemical action, although the means adopted to bring about each may be different.

It is, however, an undoubted fact that heat is developed by the chemical union of substances outside the body, and similar conditions are now found to apply to substances which are contained within it.

There are always met with elements and substances which, in contact with vital organs and their secretions assume the power of union by chemical action. They pervade the whole of the



tissues of the body. All the actions essential to life are carried on at the expense of the materials introduced, and those which are present as ready formed tissue. Their formation and development are attended with the union and disunion of the elements of compounds present, and their waste or decay is likewise attended with the same results. Thus we have a general chemical action, and this accounts for the equal state of temperature throughout the body.

Under ordinary circumstances the saccharine principles of food are being constantly caused to assume the form of fat, which under passive states, or where little exertion is carried on, is stored up beneath the skin, between muscles and around various organs, giving the animal that appearance of rotundity so much prized when intended for the butcher.

If an animal in such a condition were caused to exert himself for any length of time, the fat is absorbed and consumed. Its consumption is an essential act, not only in maintaining animal heat for the purposes of warmth, but as an agent which facilitates the decomposition of other bodies intended for the use of the system, in a different manner. Such an animal becomes lean. His muscles are distinctly observed to be well mapped out beneath the skin. The same appearance is also brought out by illness or disorder, and is the process generally understood as "wasting of the body."



There is a number of animals, as the hedgehog, (hybernating,) which during the summer become enormously fat, and sleep throughout the whole of the winter. In this case, as no food is taken, the body maintains its animal heat entirely by the consumption of fat which has been stored up in the system.

The bear, too, disposes of himself nearly in the same manner. It is also a notable fact that our cattle lay on, during the summer, a great quantity of fat, which is derived from the plentiful herbage of the period. In both these instances we also perceive the provision which is ensured towards obtaining the necessary amount of warmth during cold weather, and when food is either withheld or not very abundant.

*Animal heat is not maintained by saccharine bodies or fat alone.* In carnivorous animals, as the lion, tiger and wolf, whose diet is confined to flesh, also the wild hunters of some countries who occupy their time chiefly in the saddle, sufficient fat cannot be obtained, nor are saccharine compounds forthcoming. Natural warmth then must be procured from another source.

Wild animals in their natural state, and the hunters of uncivilised nations, are on an equality. The habits of both necessitate much exertion—activity of muscle—and consequently waste or decay.

Flesh, upon which they live, being purely a nitrogenous compound, would furnish none of the



materials for warmth under other circumstances. But under active work or movements, albuminous compounds are broken up, and the carbonaceous elements rendered available for the purpose.

A reference to the table at page 31 will explain how this can be permitted. In farther explanation it may be stated, that fatty or saccharine compounds are serviceable *only* for the production of heat as an adjunct to vital force ; but nitrogenous compounds are capable, by peculiar action, of becoming not only useful for the manufacture of flesh, but also for the production of the necessary animal heat during exercise or work.

This explains why animals in high condition appear fresh and excited. There is a necessity for movement created in order to consume the highly nutritious material within the body. Wild animals confined to cages are seen to move about incessantly when awake. Here is another mode in which the system endeavours to appropriate the food, and bring about essential vital actions. Were it not thus, disease or death would be an early result, from an accumulation of deleterious principles in the blood, by which it is no longer able to support the body.

Man and animals consume little food when the body is covered by fat, and exercise but sparingly taken. The muscles are not developed because they are not used, in fact such a body is not capable of much exertion, by reason of that want of muscular development. It is neither so



---

healthy, and is prone to disease in consequence of the absence of movements which promote vital action.

These facts are well known to practical horse owners, and they in consequence always avoid fat animals for immediate active work, and delay until the place is occupied by muscle.

We thus perceive that in feeding fattening animals and working horses two opposite principles must be carried out. It would be as absurd to feed a hunter or draught horse on the materials given to the cow or ox in the feeding house, as to expect they should replace each other in the scale of usefulness to man.

These facts dispose of the first and second propositions. We find that food is nutritious when it can minister to the formation and development of the body, and maintenance against waste ; while substances of a mixed character are needed in order to keep pace with the requirements of the body in the production of a necessary temperature, and assistance in the accumulation of vital force.

Without materials specially provided for the latter process our working animals would be reduced to the condition of flesh-eating or caged wild creatures, with this serious disadvantage, the work imposed might act prejudicially, as it would not, in all cases, be so nicely proportioned as to suit the wants of the system, or when taken in accordance with the promptings of instinct in the



creature itself. In short, food possessing elements exclusively of one kind, too rich in nitrogen, or too rich in carbon, at once proves insufficient to support life in a proper manner for any length of time. The experiments of Majendie and others who fed animals exclusively on one of the substances known as sugar, gum, starch, albumen, fibrine, or caseine, set this matter at rest for ever, and we are thereby taught that the animal economy can live and thrive only upon food provided *naturally*, and which contains *all* the elements calculated to minister to the tissues and functions of the body.

The poor inhabitants of Ireland, as well as the negroes of the Indies, also establish the truth of the principle. The former, who consume potatoes in large quantity, would exist in a poor degree of capability for exertion, were they not to add to this expensive and innutritious article of diet one of the compounds very rich in nitrogen, *viz.*, milk.

The coolies, who eat impure sugar, receive with it also nitrogenous compounds, gathered from the vegetable kingdom, and all the eaters of maize and rice resort to milk for the azotised principle, caseine.

Food, rich in mixed characters, supplies the necessary elements without disturbing the balance of the functions, which occurs when too much of one kind is given indiscriminately. All vegetable food is of a mixed character, but each kind differs in the richness of its constituents. A



---

knowledge of this is of great value to all concerned in the keeping of animals.

### BULK OR VOLUME.

Food should always possess an amount of bulk. Nature has not been unmindful of this when providing the nutritious principles of grain. The grain, or kernel, contains the nutrition in a concentrated form, and bulky material is to be found in the husk or stem.

*Proportionate bulk is requisite to ensure digestion.* The stomach cannot abstract nutrition from small quantities of concentrated food with benefit. The digestion and solution is not efficiently performed, as the stomach lacks the stimulus of contact, so essential to healthy action and secretion.

Grass, straw, and hay contain but little nutrition, and to ensure its abstraction, bulk is given to it consisting of water, ligneous matter (woody fibre), and salts.

The people of uncivilised nations have exemplified this from time remote.

The Kamschatdales mix earth and sawdust with the train oil they use as food, and in other northern regions a kind of bread is made from sawdust.

The natives of Ceylon use scraps of decayed wood with the honey consumed as food. Among



animals the wolf is known to appease the sensations of hunger by taking into the stomach a great quantity of mud ; the dog thrives best when he obtains his food from the ground, when it is mixed with grit and dirt, and in birds, small stones or sand is constantly being swallowed.

Food thus accompanied is fully compressed by the walls of the stomach, but when adventitious matters are not present, however nutritive, it does not fulfil the wants of the economy by virtue of its elements not being extracted.

#### ERRORS TO BE AVOIDED.

One of the great evils attendant upon the feeding of horses, and even cattle, is the use of *too bulky material*. By it the wants of the system are not satisfied, the stomach is over distended, and the process of chymification is retarded by the powers of the stomach being destroyed.

The walls are reduced in thickness, and rupture frequently takes place from the effects.

Fluids, however nutritious, as a rule, are not so easily appropriated as solid food. No better evidence is found than where cooked food is served to horses. The soft watery mass is too rapidly swallowed, and becomes as unnatural as it is innutritious.

*It is an acknowledged fact that no process of cooking or preparation will render the food more*



*nutritious, and there is positive proof that a mixture of substances, boiled to a pulp, are not so digestible as when given in a natural condition to working horses.*

I do not expect that all who read this statement will be converted to the truth it proclaims. I am, however, certain that in the many places where the cooked system is carried out, there will be found persons desirous of successful reform and amelioration. To such I have great pleasure in addressing these pages. Others there are whom no amount of argument would convince, or practical demonstration convert. Bigotry, prejudice, and a stupid adherence to old customs have blinded them.

To change is considered unmanly, and, as existing affairs have probably prevailed for many years, unguided by either the light of reason or science, and having tradition only for their adoption and continuance, alterations would amount to sacrilege or disrespect to the blundering system which is worshipped with such folly and stupidity.

To resume. The stomach and intestines of the horse are not intended for sloppy food. The whole arrangement forms an assemblage of perfect organs eminently fitted for bruising, insalivating, digesting, and appropriating *natural* food, and unnatural slop-food—trash concocted by the device of man is attended with disease and mortality. Among horses, if we go no further



than colic alone, the number of cases which occur where boiled food is used exceed those where attention is paid to the selection and supply of proper diet by *ninety per cent.*

*Cooked food* is open to grave objections. It weakens the digestive organs. It is swallowed rapidly, and the stomach becomes greatly distended, by which secretions are prevented or altogether stopped. Little or no insalivation takes place, and the food does not undergo those important and preliminary changes which have already been insisted upon. Secretions, otherwise necessary, are of no use with such an excess of fluid food, and if poured out are too far diluted. The stomach acquires in time an immense capacity and the muscular powers are weakened. The liver becomes diseased and the natural secretions very limited or absent. The intestines now suffer from this combination of results, and colic becomes of periodical occurrence, eventually ending in death.

The horses of many firms with which I am acquainted in Scotland, to whom boiled food is given, suffer very frequently from colic, and deaths are common.

Where proper systems are carried out, I have known three hundred animals belonging to one firm, doing the hardest work, kept in the best of health, and for a whole year not a single case occurs.

Mr. Hunting states that 120 pit animals under



his care, all in regular work, continued for six years without a single case of colic.

The bulk given to boiled food is looked upon by some as an advantage, and in illustration of the belief, a gentleman remarked recently that the food thus supplied to his horses must be more nutritious than other kinds, as it is softer, partially digested to begin with, and every twelve pounds put into the copper are increased to forty-eight. It must, however, be remembered that *thirty-six pounds of this is only water*. I would like to know who can conscientiously expect a horse to work well and continue in health on food which contains three hundred parts of water for every hundred of spoiled grain. It is an injustice to treat an animal in such a manner, which deserves more consideration on account of his usefulness, and whose better judgment would enable him to take water with greater comfort and benefit than can possibly accrue from deceiving him to swallow unlimited quantities in the form of a mess presenting such indescribable qualities and disproportionate quantities.

Cooked food for horses is a form which certainly has no analogy in nature, and wherever dictated, must inevitably arise from neglect or total ignorance of the anatomy of the digestive organs, with their physiology and the laws which govern assimilation.

To another objection raised against the cooked meat system, it is urged that horses so fed usually



drink as much water as those confined to the dry meat or manger system. I believe there is truth in this as a rule, but the fact is not favourable to the plan, and great reasons may be assigned in the large quantities of common salt used in the mess, which occasions an unnatural thirst. Excess of common salt taken into the system proves highly injurious and predisposes the animal to disease, particularly of a congestive or low form, a condition of unnatural plethora being established.

The continued presence of sloppy food, besides acting—like bran mashes—as a foreign body, deranges the balance and harmony of vital functions generally, those of digestion primarily and particularly, and the large quantities of nutritious matter which may be present, having no admixture with natural secretions are not rendered assimilable. *It therefore proves an expensive mode of feeding.*

Nutritious matters not having undergone those necessary chemical and vital changes which are ensured by the secretions of the digestive organs, are not in a fit state to enter the blood. They may be taken up, but will act as a foreign body there as they do to the intestines, and must be expelled or communicate disease.

If they remain in the intestines, diarrhœa is induced by irritative action, and horses so fed void their dung much like a cow. Such a state in this animal is not proper order, and if allowed to go on, disease of some kind supervenes.



On the other hand, if the unassimilated principle gains access to the blood, it is as soon as possible carried to the kidneys, and by them expelled. The animal frequently voids his urine. It will be frequently found to contain modified albumen, and even blood, and this accounts for the excessive number of cases of *diabetes* or *profuse staling* and *albuminuria* which have come under my notice during my residence in Glasgow.

Such cases are so common that they are considered trivial, and no doubt are, primarily; but when the errors of diet are allowed to proceed, they become marked by such characters as in the man wine-bibber and gourmand, or epicure, are modified, and appear as dyspepsia, biliousness, severe headache, piles, &c., a tolerable bloating of the countenance, with enlargement of the abdomen, which generally signifies organic disease.

The horse when suffering from these conditions is usually well drugged while at work, and as the same kind of feeding is persisted in, medicines repeatedly follow the aliment, the animal loses condition, and we may trace numerous instances of *farcy* and *glanders* to this as an undoubted cause.

Other terminations are dilated stomach, broken wind, congested, or schirrus liver, calculi (or stones) in the intestines, recurrent colic, organic disease of the kidneys and bladder, or probably speedy death from over-distention and fermentation of the contained food, causing rupture of the



stomach, some part of the intestines, or diaphragm (midriff).

#### ADVANTAGES OF PROPER FOOD AND SYSTEM.

My experience, and that of others who have devoted attention to the conditions discussed in the preceding section, clearly shows that the secret lies in *prevention*. This is comprised in cleanliness, ventilation, care and attention to the quality, quantity, and regularity of feeding, and due proportion of work.

The author is old enough to remember the effect of a journey from London to the north by stage coach, the character of animals selected for the work, and the amount they were required to perform. With such experiences he has frequently paused to enquire how the facts have failed to carry conviction in analogous cases at the present day.

In many of the coaches which ran between London, York, and Leeds, the horses were known to "do their fourteen miles out and the same number in" six days in the week. Their work was testified by the wear of the shoes, which, made of the toughest metal, and not unfrequently having an admixture of steel, were worn out by the fore feet in three weeks, and replaced on the hind by new ones nearly every fourteen days. Notwithstanding this, by proper feeding, care,



and ventilation of the stables, these animals retained their health and usefulness for years. Among them was a celebrated mare, "Old Sal." She ran in one of the above coaches as "off wheeler" for years, was known by all upon the road, had never been sick a single day, and when railways revolutionized the system of transit was over twenty years of age, and even then "as fresh as paint."

When the fly-boats plied between Glasgow and Edinbro' on the Forth and Clyde Canal, an old mare named "Maggie Lauder," was stationed to run between Port Dundas and Glasgow Bridge, a distance of eight miles, the time allowed being one hour. After a rest of one hour the return journey to Port Dundas was made, and in the afternoon she performed the whole distance over again, thus travelling and drawing the boat thirty-two miles per day.

The person from whom I obtained my information rode the animal daily for seven years, during which time she was doing the work alluded to six days in the week, "was never sick nor sorry a single day, nor ever had a day's rest in addition to the usual Sunday."

When the boats were superseded by railways, "Maggie Lauder" was sold at the age of *twenty-nine years*.

A similar instance is related of a horse employed in like manner on the Paisley canal. He was sold at the age of twenty-seven when the



boats were discontinued, and, being "fresh as a lark" at the time, suddenly fractured the bone of one of his legs in his gambols while being led home by his purchaser.

I am aware that breeding will in a measure account for "pluck" and disposition to work in animals as well as in man, but it will not stand in the place of *ability* under any circumstances. The willingness or pluck may be always present, but ability will depend upon a condition of strength.

One thorough-bred horse will resemble another very much in disposition, but differ widely in ability from mode of living. Take the first from the green pasture and run him alongside that brought direct from hard dry corn and sound hay, upon which he has subsisted months, and performed daily exertion. The effect is easily perceived; weight or distance is scarcely an object to him, but the first is blown or lamed before half a dozen fences are crossed, or has received his death summons from various causes.

Harness horses, and horses used in draught, require similar treatment to produce strength and endurance.

While pluck is derived from breeding, strength is derived from food and a healthy digestion. Corn and hay, transmuted within the penetralia of the living organism, becomes muscle. Manure, the refuse of digestion and the animal body, the



agriculturist knows, nourishes his land, without which he can expect no crop. In the production of artificial light, all depends upon the supply of combustive material and agents which support or promote the process, which, in proportion to quantity and quality, afford a good or bad kind of illumination. In the warming of buildings the maximum temperature can only be obtained by instituting an operation of the laws of combustion upon materials capable of undergoing that process, and, likewise, by the operation of laws within the animal organism, if we need strength (force), it must first be supplied in the shape of sound, dry provender.

There is a very erroneous idea, at least to my perception, entertained by many, that where a draught horse is required for moving extreme weights he should be large and ponderous. If the dealer is inquired of, "You want weight," says he. If a friend be appealed to, a similar recommendation is given. It is in this way many useless, heavy-legged, unsightly, lugubrious, and slovenly animals are tolerated. What these are supposed to gain by superior capabilities in moving weights they lose in speed, and hence are seen creeping along the streets, and, as occurs in some towns, creating quite an obstruction to general traffic.

It appears to me that strength is required, not absolute weight, but a good development of



muscle. If weight only is wanted, it would not be so frequently remarked that horses "will not pull a sitting hen off her nest."

And, again, if nothing but weight will suffice, then a lump of lead or any inanimate object would answer equally as well.

I would refer my readers to Youatt's excellent treatise on draught, and they will perceive that other objections are against tall animals for moving loads, particularly on four-wheeled carriages.

#### REGULAR FEEDING.

It is one of the essentials of good management that horses should receive their food as regularly as possible. Without regularity, and especially with long fasts, the digestive organs are prostrated or weakened, and food, which would otherwise be nutritious, brings about those changes already described.

In coal mines, where feeding and work is dictated by those who know nothing about it, broken wind, colic, diabetes, organic diseases of various kinds, calculi and death by rupture, are common. When, on the contrary, attention is given, these fatalities are rare, and when they occur are usually traced to other causes.\*

---

\* Much valuable information has from time to time been detailed to me by my friends, Mr. Charles Hunting, M.R.C.V.S., and Mr. Luke Scott, M.R.C.V.S., whose experience in the management of pit animals is of the most extensive character in Britain, and therefore to be received with respect.



## CALCULI OR STONES IN THE INTESTINES.

The nature and aggregation of the particles which compose these bodies are not without interest, as affording information on the conditions which are present in the stomach and intestines. *Calculi* are very common in the horses and ponies of some coal mines, and also among the horses of millers, general carters, and those used in large establishments where the system of feeding and work is defective.

Their origin is usually considered to arise from water impregnated by the salts of lime, which are precipitated in the same manner from solution as occurs in the steam boiler or tea kettle.

It does not follow that calculi form on the use of such kind of water. It is also a most notorious fact that from water of precisely the same character supplied to the animals in two coal pits, different results are manifest. In one, where proper care in feeding and work is observed, there is the greatest amount of health, but in the other, where systems are the reverse, and especially when work is excessive and irregular, calculi exist.

That the water draining from the magnesian limestone, and holding lime in solution, has not much to do with their formation, is proved by several circumstances.

First: Many calculi which I have found in



such animals have little or no lime in their composition, being mainly composed of mucus, hair, dirt, coals, &c., closely matted together, and known as "*dust balls*."

Second: The miners consume large quantities of the same water, but are not known to suffer from intestinal calculi.

Third: From personal experience and residence in districts where calcareous or hard waters only can be had, I have ascertained that calculi are not more prevalent when system is observed, and

Fourth: I have found calculi to exist more abundantly where water is of the purest character. Some of the largest specimens which have been obtained were from horses using soft water alone, and therefore must have another origin.

To attribute their formation to any special kind of water is a false theory, and negatives the power of the intestines to remove useless matter.

The presence of a nail, piece of wood, stone, or other hard substance, is also said to insure the formation of calculi. But horses and cattle meet with these things constantly among their food, and, while many are doubtless rejected by the sensitive lips, others are swallowed, as proved by their presence in the dung during life, or intestines after death.

The *origin* of the material which forms the substance of a stone or calculus is undoubtedly from the *food*. Hard waters may assist under certain circumstances.



They may contain a large quantity of lime, but the food relatively contains a much larger proportion. It is a substance largely in demand for the wants of the system.

If the animal is enfeebled by overtaxing work, long fasts, and supplied with food of an inferior quality, a condition known by the term *bulimia*, is established. This is known by an irregular, capricious, and morbid appetite, irregular bowels, staring coat, leanness, inaptitude for work, and a desire to lick the walls. Sometimes, however, these symptoms are either not well marked, or escape observation.

Under such circumstances the animal swallows sticks, stones, and rubbish of all kinds, to appease the ravenous desire within. Food, yet the most nutritious, is not properly digested, the secretions are vitiated, or altogether deficient, and act imperfectly. Further derangement occurs, in which the liver particularly suffers, and affairs now assume an aggravated character.

During these conditions assimilation and absorption are not carried on perfectly, and the mineral matters of food and water are deposited in the solid form, and aggregate around any rough surface or object which may be present.

As the mass increases in size corresponding derangement is continued, and thus secures material for its development, or the formation of others—as many as fifty having been found in one animal.



Some time may elapse before acute symptoms are observed. Suddenly abdominal pain arises. Symptoms are continued, and become aggravated, admitting of no relief, and the animal dies.

Upon making a *post-mortem* examination, one or more of these stones are found to have passed into a narrow part of the gut, and become imprisoned by spasmodic action of the muscular walls which tightly enclose it on all sides. Sometimes destruction of the gut has occurred, and the stone is partially or wholly liberated along with the contents of the intestines.\*

#### OBJECTIONS TO THE USE OF DRY FOOD.

It is urged by many persons enthusiastic in the feeding and management of horses, that a change

---

\* I have succeeded in obtaining a great number of these interesting specimens during the past sixteen years. In less than two years twenty-six were obtained from pit animals, having caused death, and many more were found in animals dying from other causes.

Their composition was principally mucus, the felted down from the oat, silicious matter and carbonate of lime. The last-named substance usually formed the outer portion, while the interior was filled by pieces of coal mixed with the other ingredients. Many of these were presented to the museum of the Albert Veterinary College, London.

Several fine specimens have recently been forwarded to me by my friend, Mr. Thomas Foreman, M.R.C.V.S., Leadgate, Durham, exhibiting similar peculiarities. Of these he has collected a great number from pit animals.



to a system of bruising and cutting of food gains only one advantage, *viz.*, the animal fills his stomach quicker, and is thereby enabled to take more rest.

This property is pre-eminently claimed in favour of cooked food, and in addition, that it is partially digested for the animal. Experience and systematic inquiry prove the fallacy of these tenets.

I have shown that digestion, when properly carried on, is wonderfully rapid and effective.

The stomach, being small, is rapidly emptied again. This obtains in all horses, including those fed on the *manger system*.

Every one knows the effects of bran mashes. How much more, then, must constant supplies of food, supposed to be half digested at commencement, containing much water, and being an unnatural mixture, act like a foreign body?

The laws of the animal economy render such results inevitable, for as quickly, in proportion, as the stomach is voiding the digestible portions, the sensation of hunger arises, and desire for food is appeased by taking in more. Thus it will be found, in opposition to the argument in favour of a half-digested food, that the process of digestion is accelerated, and more is required. Such food always fails to give up the whole of its nutrition; *the animal eats more, costs also more, and gains the least by the method.*

It is thus that an equally long time is occupied



in the consumption of food, and visit him at any hour of the night, he will be found having an appetite, and like Oliver Twist "looking for more."

*Rapidity of digestion* is a provision established by nature. If the stomach had been from the first intended to receive the large quantities frequently placed before horses, or to perform functions assigned to the teeth and salivary glands in addition to its own, the logical inference is that, as nature, in her development of all things, has not studied ornament merely, the stomach would have been endowed with greater capacity and powers, the teeth and salivary glands, in all probability, absent entirely.

It is also urged against the dry meat or manger system, that horses fail to masticate or grind the whole of their corn, that much in a state capable of germination or growth passes out in the excrement, and, of course, the cooked meat system supplies this deficiency.

During a season of extreme scarcity in India, it has been stated,\* the famine-hunted wretches followed the English camp, and drew their principal nourishment from the grains of corn extracted from the excrement of horses.

I well remember an extensive firm employing many horses, whose manure was objected to by

---

\* Letter from an Indian officer to J. Curwen, M.P., quoted in Blaine's (fifth) edition of "The Veterinary Art."



several farmers because they obtained crops of oats in places where they were not required, after using it on the land. I have also observed the heap of manure literally covered with green blades from the growing oats. How was this, and why has such a simple circumstance had no better effect than to induce a system of porridge feeding for an animal eminently unfitted for it?

It was in this wise. The remedy was sought for in a manner too superficially; and in the absence of an acquaintance with the principles of management, the horse has been compelled to suffer for the omissions of others.

In such cases I have known carpenters, and even more unlikely men appealed to for information and a remedy. The system resembles very much that so commonly adopted under certain Acts of Parliament, where discretion and power is given to tailors, joiners, bailiffs, policemen *et hoc genus omne*, to pronounce in a magisterial court what is cruelty to animals in a medical point of view, and to descant upon the pathological signs which prove a carcass to be diseased, and all this in the teeth of an educated scientific witness.

Such matters require dealing with philosophically. They are not within the comprehension of every clodhopper, groom, and coachman, who fancies he knows all about a horse because he cleans him. Their duties lie in the practical administration of the brush and broom, and faithful



execution of all orders entrusted to them. Management and dictation form no part of these.

The arrangement and functions of the internal organs of the animal body are matters beyond the comprehension of illiterate brains. The mutual relations of the horse and his master will exist in proportion as this truth is observed.

It is important to us to enquire whether oats or any kind of grain are capable of growing after having passed through from eighty to ninety feet of intestines subjected to healthy acid, alkaline, and other peculiar juices which convert their starch into sugar, separate and modify their albumen and caseine, and dissolve out salts, or to a temperature varying from 98° to 102° F., occupying time varying from ten to twenty-four hours?

I venture to predict not, and to assert that very few grains will pass out unacted upon. If such were the case the whole processes of mastication, insalivation, digestion, and that of making malt are but a farce, and cannot be accounted for by the present deductions of science.

The solution of the growing oats, however, was soon made apparent.

The manger system had been imperfectly carried on for some time without any alteration in the size or character of the mangers themselves. These were shallow, narrow, and not provided with cross-bars. In this way much was



wasted by the horses throwing out the corn when searching for better portions.

In addition, I entered the stables on one occasion when the premises were supposed to be deserted, and observed a number of those young urchins who are always a kind of *sine qua non* about lorries and stables, carrying on a mimic warfare, the missiles in the case being oats, obtained from the open corn-bin or granary. It was afterwards ascertained this had been a common practice. The successful remedy soon followed in the shape of deep and wide mangers, with iron cross-bars, padlocks to the corn-bin and granary door, and institution of a proper room for mixing the corn and cut food.

That the half-famished Indians picked something out of the excrements of the horses I have no doubt, and they most probably resembled oats in appearance. Birds, especially sparrows and common fowls, are also captivated to make such a selection for their crop. But only in outward appearance would these abstracted matters resemble the grain of oats. Few have trusted themselves and their judgment in troubling in this matter farther than making the broad assertion on the strength of appearances. A superficial mode, and very delusive proposition to be circulated. I have tested the matter and found it to be an error.

If the stomach and digestive organs generally are in a state of health and order, very few



grains escape their action, and the common inference is that if the horses of the Indian campaign referred to really parted with so much corn as to support such a number of followers, they must have been fed in a very imperfect manner, their digestive organs in a state of disorder, and from the loss of the food, in a condition certainly not desirable in an enemy's country.

In France the subject met with the attention of Boussingault and Papin, who instituted experiments, and ascertained correctly that the passage of undigested grain from the bowels of horses in health is merely nominal.

Out of eight pounds allowed per diem only one ounce of unchanged grain could be detected, and this weight included moisture.

*The digestive organs generally should be in a state of healthy action in order to secure the proper benefit from food.*

Under this head there requires little to be written. I have fully shown how disease originates by false management, and how waste occurs in food supplied under such circumstances. It must, therefore, be apparent that those organs specially set apart for providing the fabric of the animal body should not be overtaxed or enfeebled. If such occurs the whole body suffers more or less.

*Care is especially needed after protracted hard work and long fasts, avoiding the use of large*



quantities of cold water upon the contents of a well-filled stomach, or supplying too much food at one time. In fact these precautions would repay, if followed, at all times ; neglect of them is a frequent source of derangement, and from which fatal consequences usually succeed.

Where a great number of horses are kept it is far more profitable to employ a person whose sole occupation is confined to the feeding department. Corn, &c., should be transferred to his possession under proper rules for mixture and use, the quantities being accurately stated in documents handed to him. In return he should furnish evidence of having carefully complied with the terms, and his papers should also afford particular information as to consumption under all circumstances, regular or variable.

He should be a man who will carefully notice the condition of all horses when brought to the stable, and willing to minister to their comfort in properly regulating the quantity of food each may receive without injury.

The details of such management require special attention in order to be properly applied to the various circumstances which obtain in different establishments. There is, however, no particular difficulty beyond a willingness that need be encountered, in order to make the whole safe, satisfactory, and profitable.

In the absence of particulars it would be an



impossibility to attempt to lay down a plan suitable for any particular establishment. An outline must be framed on the principles here given, the work, size, and nature of animal, &c., being points of importance for consideration.



## PART II.

## VARIETIES OF FOOD.

THE food used for working horses are those cereals and leguminous seeds which, usually denominated corn, consist of oats, barley, Indian corn or maize, peas, beans, and tares, together with hay, straw, bran, and linseed.

These vary much in their properties as well as nutrition, and on that ground alone arises the question "What kinds are best, and what proportion should be given?"

An answer will greatly depend upon circumstances, such as the character of work imposed, together with the market price and condition of the provender.

The relative proportions of nutritious matter contained in different kinds of food have been ascertained from time to time by analysis, an outline of which is as follows:—

In 100 pounds.	Nutritious matter.	Fat, or heat producers.
Turnips . . . . .	1	9
Red beet . . . . .	1½	8½
Carrots . . . . .	2	10
Potatoes . . . . .	2	25
Hay . . . . .	8	68½



In 100 pounds.	Nutritious matter.	Fat, or heat producers
Maize, or Indian corn . . . . .	12	67
Oats . . . . .	14	68
Barley . . . . .	18	68½
Bran . . . . .	18	4
Linseed . . . . .	24	64¾
Beans . . . . .	31	51½
Peas . . . . .	32	51½
Tares, or lentils . . . . .	33	48

*Maize or Indian Corn.*—By the above table it will be observed that maize among corn contains the least percentage of nutritious matter, and from it we gather that it is not suited as a principal article of diet for working animals. It contains a great proportion of water and starchy matters, and forms in consequence an excellent food for pigs and fattening animals, given with other varieties.

*Oats* stand next in proportion, and by custom have been most commonly selected as a principal article of diet for horses, but prove very expensive as sometimes used. This is very apparent on calculation, and results are demonstrative in practice.

A horse doing little work, or what actually only amounts to exercise, would be economically and advantageously fed upon hay and oats, as the demand for muscular power not being great, such articles form a very suitable diet, without causing him to become overloaded with fat and a burden to himself.



Estimating oats to weigh 42lbs. per bushel, and costing twenty-six shillings per quarter (eight bushels), the cost per week for different allowances, together with the amount of nutritious matter contained, would be somewhat as follows :—

Pounds per day.	Pounds per week.	Cost.		Azotised or nutritious matter in pounds.
		s.	d.	
10	70	5	5	9·8
12	84	6	6	11·76
14	98	7	7	13·52
16	112	8	8	15·68
18	126	9	9	17·64
20	140	10	10	19·60

In addition, sixpence must be added for each stone (14lbs.) of hay consumed, which yields 11·20 oz. of nutritious matter.

Oats should be heavy in the hand, devoid of thick husks, and short and plump in the kernel. Good dry potato or Scotch oats, weighing 39 to 42 pounds per bushel, are undoubtedly cheapest to purchase at all times, and to be preferred to other kinds weighing from 33 to 37 pounds, even when four or five shillings extra per quarter is paid.

It is no saving to purchase corn, especially oats, *made up* to a certain weight. Some dealers guarantee four bushels of oats for a stated price to weigh, say 160 pounds, and when measured



the quantity considerably exceeds the four bushels.

Under such circumstances the purchase cannot be as beneficial as if the oats were 160 lbs. natural weight. The inference is, the grain is of an inferior quality, the kernel being light and the husks disproportionately heavy. *It is always more profitable to obtain standard measure and weight*, as this proves an important item in twelve months, and particularly when animals are doing variable work.

The lower priced oats may answer well for moderate exercise or work, but as soon as the labour is increased, or a change is made from standard weighing oats to others of a lower nutritious percentage, animals suddenly fall away in condition, and become liable to disease, and in ignorance of the real causes, are not unfrequently impregnated with a course of useless and even dangerous drugs. It is thus the expenses of feeding are obviously increased, as to them are to be added less ability for work, and an additional incurring of expense, while the original cost of the provender is really but a fraction below the price of good material.

When the kernel of oats is small the husk preponderates, the former containing as low as eight per cent. only of nutritious matter, and the latter no better than straw itself, but for which the price of oats is paid.

Mr. Hunting proved this by a set of tedious



but ingenious experiments. He says,\* "One ounce of foreign oats, 39lbs. per bushel, was weighed, the same weight of old Scotch, 40lbs. per bushel, and the same weight of Tartar oats 36lbs. per bushel. The foreign oats contained 1,112 kernels, the Scotch oats, 1,084, and the Tartars, 1,144. The husks were weighed. The husks of the 1,144 grains of Tartars weighed 120 grains; of the 1,112 kernels of foreign oats, 126 grains; while the husk of the 1,084 kernels of Scotch only weighed 96 grains."

Under these circumstances, horses fed upon three bushels of oats per week, weighing 42lbs. per bushel, would receive 126lbs. total weight, of which 25lbs. would be husks, having a value only of four per cent., in nutritious matter.

Change this to Tartar oats at 39lbs. per bushel, exactly 39lbs. must be extracted for husks, and if the foreign oats are used, 36lbs. would be the amount of husk in the same quantity.

It must be borne in mind that the inferior kinds of oats are not only remarkable for a preponderance of husk, but their nutritious matter, and consequently their feeding qualities, are considerably below the standard weighing Scotch oats.

Similar rules should be applied to other kinds of grain or corn, and their standard qualities

---

\* "On the Feeding and Management of our Domestic Animals." p 14.



obtained from analyses upon which reliance can be placed, and information will thus be easily obtained whereby to institute a system of mixtures for any purpose of feeding.

### SELECTION AND PURCHASE OF GRAIN.

Most persons are aware that corn should be thoroughly dry for feeding purposes, otherwise inconveniences occur in the form of indigestion, colic, weed, grease, or loss of condition, &c.

Besides, in purchasing that which is not dry, or seasoned, a decided loss occurs, which buyers should avoid by securing a reduced price in accordance.

Owing to this, large feeders take advantage of the markets, and purchase oats, peas, beans, barley, &c., as they are aware that in good corn the loss by a reasonable evaporation of water is an ample payment for the construction of granaries for spreading out and constantly turning; added to which, animals are kept in better condition, and work is less interfered with by illness on that account.

I have known hundreds of quarters of corn purchased under such conditions, and with the observance of other measures, to be noticed, a balance has been shown which has cleared the cost of keeping the animals for a great portion of the time, as compared with the previous cost from another mode of feeding.



The purchase of grain should not be referred to week, month, or year, in fact, to no period if it can be avoided. It is much better to store up corn and ensure it being thoroughly dry, than to depend upon purchase when it is immediately required. Many good bargains are put in the way of owners who are open to purchase, and the profit obtained helps to improve the appearance of the credit side of the balance sheet.

Where roomy grain floors cannot be had, it becomes a matter of greater necessity that the provender should be dry, and of *guaranteed weight and measure*. Of a necessity also, a higher price must be paid for it, and even under those circumstances it answers far better than the use of inferior kinds. To constant purchasers, dealers are to be found who will endeavour to provide what is required in the shape of a regular sample, and thus difficulties are greatly overcome.

These principles have been more understood of late by those who feed large numbers of horses, and whose competitive kind of work requires them to study every item of expenditure. In some instances animals are fed inexpensively, and preserve their health and condition with remarkable benefit from the system, even under much harder work.



## ECONOMY OF FOOD.

The method of economising food consists in using *a variety of grain* instead of one kind, and that exclusively oats. Some have tried the different leguminous kinds also separately, but found them inefficient in economy, and even injurious.

A farmer of my acquaintance having about thirty horses, purchased, at a cheap rate, a quantity of Indian corn, under the idea that it would effect a saving in the cost of feeding. On the contrary, the health and condition were greatly sacrificed.

Others, again, give a preponderance of beans, acting upon the well known fact that, being more nutritive than oats, they must be more economical. In the mode adopted, however, they have proved highly expensive as well as injurious.

*Linseed* is also added, and with pernicious results. It is too laxative for general use in quantity, but judiciously administered proves assimilative, hastens the process of assimilation, and assists in the formation of fat and flesh. As a nutritious body it is very highly expensive.

*Tares*, which are the most nutritious diet we can use for horses, are objectionable as being unpalatable in quantity. Similar objections may be urged against other varieties of food when exclusively used, particularly in their liability to produce disorder of the digestive organs.



A mixture, therefore, should be regulated by special conditions. Due consideration is to be given to the nutritious matter contained in each ingredient, and as a whole, the amount and character of the work, quantity allowed to each animal, and the price paid at the time. Where the consumer is compelled to go to the dealer for his week's, fortnight's, or month's supply, this is particularly needed, as there is no room for speculation on the rise or fall of corn, upon which frequently large sums are to be saved.

In estimating the amount of nutritious matter contained in food of different kinds, and how much is required to preserve horses in health under their work, the table given in pages 63, 64 will be found of great service. Practically, it is answered by observing the quantity of oats of standard nutrition, or other kind of food, which may be required to keep the animals in condition for work and perfect health. The amount of nutritious matter is then determined according to the per centage stated opposite each kind by simple rule of three. This done, the estimate of nutrition contained in other kinds is observed, and mixtures arranged, forms of which will shortly be submitted.

It is to be next inquired, will this ensure a less cost than is incurred by the principal use of oats? The answer is emphatically, yes, and with a greater supply of nutrition.

I am indebted to a gentleman of great ex-



perience among horses for valuable information in the feeding of his animals.

They are employed in the town of Sheffield, which is not unlike Glasgow for its hills.

The work is severe, and consists in removing the heavy manufactures of steel, stoves, hardware, &c., to and from the railway station. Drays or lorries are generally used, and three tons is a common load.

In a letter to me dated April 4th, 1864, he says:—

“Our horses’ diet is as follows:—

Hay . . . .	16 pounds per day,	8 stones per week.
Oats . . . .	10   ”   ”	5   ”   ”
Beans . . . .	5   ”   ”	2½   ”   ”
Maize . . . .	4   ”   ”	2   ”   ”
Bran . . . .	2   ”   ”	1   ”   ”
<hr/>		
Total .	37   ”   ”	18½   ”   ”

No loose hay, all is chopped; oats, beans, Indian corn all crushed separately, then the whole is mixed with the chop. The hay costs 4s. per cwt., oats, 1s. per stone,\* maize, 11d., beans, 1s. 2d., and bran 7½d. per stone.

“I always buy of the best quality without regard to price, as I find that the cheapest, all points considered. As a rule, I greatly prefer English oats.† The harvest of 1862 was an

\* The hundredweight is 112 lbs., the stone 14 lbs.

† I think the term English is here applied without prejudice



exception, and I bought foreign oats principally, being in a superior condition to English. In addition to the above—which is the winter scale, the summer is somewhat reduced—we every Saturday night give each horse a mash of linseed mixed with a small proportion of bran, boiled altogether and given warm.

“This serves to lubricate and clear the bowels, and comforts the animals. I do not know whether that is the cause; but since its adoption we have had few, if any, cases of colic, or severe inflammation; prior to the adoption of that system these cases were of great occurrence.”

It must be borne in mind that these are heavy horses, and we have here also an evidence from an unprejudiced source, that a considerable amount of nutritious matter is required to keep up the health and condition of the animals, since when the inferior kinds of mixture were supplied, in the shape of oats principally, disease was common. This I can testify, having been in professional attendance.

We are also taught practically that a loss of condition and proneness to disease is brought about by food containing too little nutrition, when the demands upon the system are excessive.

---

to Scotch oats, but as a distinction to foreign and inferior kinds. The writer is keenly alive to the qualities of the various kinds of corn, having had unlimited experience in the feeding and management of the best draught horses in large numbers, and doing the hardest work.



Suppose these animals had been fed exclusively upon oats and hay, they would require to consume weekly *four-and-a-half bushels* of the former, and about *ten stones* of the latter, in order to receive the same amount of nutrition, viz.  $31\frac{1}{2}$  pounds derived from the mixture quoted, and which would cost at the same prices about *eighteen shillings and sixpence*.

Here is also a decided saving of three shillings and sixpence per horse per week, by the use of the mixture, which only costs *fifteen shillings*, and with fifty horses would realize £8 15s., and for a year £455, besides almost an immunity from disease and death, to which they were before exposed.

#### SYSTEMS ADOPTED ON VARIOUS COLLIERY ESTATES.

It is by paying particular attention to these matters that so much has been effected in the saving over former expenditure, on many colliery establishments, where the supply and mixing of grain is under the management of the veterinary surgeon.

Mr. Hunting, in his pamphlet already referred to, has shown that with the number of horses and ponies employed at the various collieries of the South Hetton Coal Company—all doing the hardest work—the saving effected in ten years amounted to no less than the enormous sum of



£31,876 2s. 7½d. by cutting the hay into chaff and mixing with small quantities of straw, and substituting superior kinds of beans, peas, barley, tares, &c., in lieu of oats. He also states that out of 225 horses employed in two pits, during six weeks between 15th December, 1850, and 24th January, 1851, when the old system was pursued, there died of colic and its consequences *three horses and ponies*; while with the same number of animals under the improved *régime* during seven years, there were *less than three cases per annum*.

#### CUT FOOD AS AN AGENT IN PROMOTING PROPER INSALIVATION.

The value of this kind of provender as an agent in causing a proper insalivation of food is undoubted. On this subject Mr. Hunting is as explicit as he is full of information regarding it. He found that a number of animals, all selected as near as possible with an average age, height, and common appetite, required much longer time to consume the same weight and measure of food cut and mixed in the manger than when it was given in the shape of long hay in the rack, and oats only in the manger, from ten to thirteen minutes more being required.

After what has been said in reference to the objects of the teeth and salivary glands and their



secretions, it must be obvious that in giving food in its naturally dry state, it serves more important uses in the animal economy. And as the *quantity* of saliva plays also an equally important part, the use of cut food is here unmistakably shown to be a very desirable proceeding.

At the *Hetton Colliery*, Mr. Luke Scott, M.R.C.V.S. has followed out most useful principles. His attention had been arrested in 1851 by the inefficiency of the existing system of feeding, and proneness to mortality which was constant among the horses and ponies employed.

The quantities allowed per pay\* to each animal employed in connection with the estate were as follows:—

	Oats in	Cost.		Hay in	Cost.	Total.			
	Bushels.	£	s.	d.	stones.	s.	£	s.	d.
Waggon Horse .	8	1	6	0	28	14	2	0	0
Farm Horse . .	6		19	6	20	10	1	9	6
Pit Horse . .	6		19	6	24	12	1	11	6
Pit Pony . .	3		9	9	12	6		15	9

It is necessary to state, the whole of the hay here allowed was not consumed. Much is constantly wasted in collieries by being carelessly sent down the pits, or conveyed to the stables in large trusses, which come in contact with water, dirt, and grease. The animals consequently refuse it, and generally place it beneath them, sufficient being

---

\* Colliery accounts in the county of Durham are made up once a fortnight, when wages are paid; hence the term "pay."



frequently gathered to make a very good bed. I have seen a week's supply of hay and corn lying exposed to the dust flying from the pit mouth, as well as steam, condensing as it falls from the boilers, and, in common instances, to the rain of one or two nights in addition, before being sent down. The former kind of unsystematic arrangement was of frequent occurrence at one pit, but no kind of remonstrance succeeded in effecting an alteration.

Pit animals are usually kept *twelve hours* at work, and in many cases, *without food and water*,\* or at least without a satisfactory amount of either, and seldom periods of rest.

I have known animals kept from their stable thirty-six hours at a time, and when complaints have been made they have been met with a stout denial from the powers that be; or, where denial has been useless, it has been assigned as a cause, that the veterinary surgeon has not provided a sufficient number of animals for the working of the pit, an office which he does not include among his duties without express orders.†

Such long fasts act very prejudicially upon

---

\* The practice is not so common as formerly. In some pits it is abolished, but slumbers in others.

† Lord Kinnaird, in his letter to the Home Secretary on the Earndale Colliery explosion, dated 15th November, 1867, has fully shown the aspect of colliery affairs when he quotes the words of the miner, who says, "If peoples as knows dared to speak, these things would soon be stopped."



the animal. When he returns to the stable the food is seized and ravenously swallowed, and, as a consequence, it does the least good, but is more frequently productive of the greatest harm.

By a reference to previous remarks on the estimate of nutritious matter in the various articles of food, it will be found that the Hetton Colliery "old plan" of feeding was doubly expensive and extravagant. It was innutritious as it yielded to the waggon horse (and others proportionately) one fourth less nutriment than the mixture given hereafter. It was expensive because it cost nineteen shillings per week, from its use much was wasted, the animals were in bad condition, and mortality great.

It was decided at length, to allow Mr. Scott to regulate the system, which he did, in the following way:—

His first act was to have the hay cut in trusses of sufficient size to fit a coal tub,\* in which they are sent down the pit, and thus totally prevented from collecting dirt and grease as before. This resulted in a saving of the amount wasted, and also supply in a better condition, which would be turned to account in the animal body.

The varieties of grain fixed upon to be used were bruised separately, and mixed in the following proportions:—

---

\* This plan is also carried out at all the collieries under Mr. Hunting's superintendence.



Grain.	Weight per bush. lbs.	Price per qr. s.	Actual weight. lbs.	Cost. s.
Barley, 4 bushels . .	56	30	224	15
Oats, 4 „ . .	42	26	168	13
Peas, 2 „ . .	66	40	132	10
			<hr/> 524	<hr/> £1 18

In order to arrive at an average number of animals to be fed, the ponies and horses in the pits are classed as follows :—

All above 15 hands are called horses.

Three above 13 hands, and under 15 hands,  
equal to two horses.

Two above 11 hands, and under 13 hands,  
equal to one horse.

Under this arrangement the whole were reduced to an average of 130 horses.

The superiority of the system is at once apparent by a reference to the subjoined analysis, in which the two plans are contrasted.

*Old System, for One Pit Horse.*

6 bushels of oats per pay, at 3s. 3d. . .	19s. 6d.
24 stones of hay, at 6d. . . . .	12s. 0d.

---

£1 11s. 6d.

---

Or for 26 pays (one year) . . . £40 19s. 0d.

---

Nutritious matter contained . . . 52·08 lbs.



*New System, for One Pit Horse.*

176 lbs. of the mixture . . . . .	14s. 4d.
18 stones of hay, at 6d. . . . .	9s. 0d.
21 pounds of bran . . . . .	11d.

---

Total per pay . . . £1 4s. 3d.

---

Or for 26 pays (one year) . . . £31 10s. 6d.  
 Saving effected for one horse for  
 one year . . . . . 9 8s. 6d.

---

Cost per horse under old plan . . £40 19s. 0d.

---

Saving in 130 horses for one year £1225 5s. 0d.

---

Nutritious matter contained . . . 48·18 lbs.

In the old system it will be observed that 52·08lbs. of N.M. (nutritious matter) are contained in the food supplied, against 48·18 lbs. in the new. The greater part, however, was not obtained by the animals on account of the excessive waste already mentioned in the hay, and from the greedy manner in which the corn was devoured.

The additional 4 lbs. of N.M. supplied also proved expensive, besides useless, as it was unavailable. For it the sum of 7s. 3d. per pay was being paid, as forming part of a system which engendered disease, and gave the animals even less support, although professedly, a mode which furnished a large amount.

It is now a custom to allow the pit animals a portion of the hay and corn during the day,



instead of causing them to fast twelve hours, as before.

The result is, Mr. Scott has found that the food is more leisurely taken, masticated, and thoroughly digested.

Besides the saving effected in actual expenditure, the reduction of disease and losses by death, is an important item. In repeated visits to the animals in the pits fed upon Mr. Scott's principle, it is due from me to state that I never saw a greater uniformity in condition while the hardest work was being imposed, and cases of indigestion, colic, and death in consequence, were the exception, and of extremely rare occurrence.

#### WHAT CONSTITUTES A CHEAP FOOD.

It is usually considered a cheap mode of feeding, when material can be supplied for one penny per pound, and I find several owners base their calculations of cost at this rate. But it must be borne in mind that food costing only one penny per pound is not inevitably an economical food. We must look farther than mere cost. Economy does not consist in price alone, to such must be added the veterinary surgeon's account, whose services in the main will be found to have been occasioned by the supposed economical food, and in addition, the value of the amount wasted by refusal, fermentation, or that which is hastened



through the bowels in the state of "partial digestion," so much insisted upon.

Again, the amount of nutritious matter contained must be accurately calculated, or no reliable estimate of cheapness can be made. I am aware of several large firms in Glasgow who believe they are feeding economically upon one penny per pound rates, while their food yields ten per cent. less nutrition than the preceding mixtures, and cases of acute indigestion, &c., &c., are constant and numerous. These form a good comparison with the systems of Messrs. Hunting and Scott, the mixture proposed by the latter costing only ninety-eight parts of a penny per pound, with a high scale of nutritive value.

#### OBJECTIONS TO A CHANGE OF GRAIN.

It is frequently urged that to adopt a total change in the kind of grain used, is to produce serious evils and fatal disorder, as exemplified by animals gaining access to the open corn bin, or to a heap of wheat or barley, when either rupture of the diaphragm, stomach, or intestines takes place, and death speedily ensues, while at other times founders (laminitis) occurs.

These certainly appear grave objections at first sight, but in reality are difficulties of no moment. We are speaking of *systematic* feeding, *not* deliberate engorgement. Death or disease results in



one case by repletion and fermentation consequent upon the halting of an unnatural quantity of food, which, if taken under proper principles, would in the other prove nutritious and life supporting. The death of horses by these means is fully carried out in analogy among mankind, with the exception that in the latter gluttony generally exerts itself a little more slowly.

*In no case, with proper caution, will evil consequences ensue by a change to the dry, or manger system, from any other. It need therefore cause no apprehension.*

#### NATURE, USES, AND ABUSES OF BRAN.

Bran will be found by analysis to contain from 14 to 18 per cent. of azotised matter, equal in fact to oats or barley. In nutrition, however, it is inferior to the straw of wheat or barley, a property which appears to obtain with the husk of grain generally. Owing to some peculiar form of combination, the azotised matter is not liberated by the process of digestion; hence those who consume bread in which the bran of wheat has been retained, under the idea that such is more nutritious, are greatly mistaken. The fæces or excrement are loaded with bran which has passed out almost unchanged, and horses fed upon it exhibit the same conditions.

That an admixture of bran with wheaten flour



or with the provender of horses, is useful and even profitable, there can be no doubt. This is to be attributed to the mechanical action set up by the siliceous particles, which obviates constipation, and preserves the proper action of the digestive organs when given in a judicious manner. Messrs. Hunting and Scott take advantage of this property, and use it *daily* with their superior mixtures of corn.

As a *laxative*, bran is justly called into requisition periodically as a warm mash for animals in whom there exists an innate disposition to constipation. I place the action of a bran mash, *given occasionally*, as one of the safest, most natural, and acceptable adjuncts towards the preservation of health; which effect is produced with more benefit and less deterioration to the system than by any other means. There are few horses that will not take it when offered as a change, and I would recommend, especially in winter, that it be given at the temperature of new milk, *not cold*, and the use of it should not be insisted upon indiscriminately, or ill effects are speedily shown.

Nothing can be more anomalous than the opinion entertained on the use of bran, as it obtains in many quarters. Believed to be non-nutritious, it is given largely during disease, to ward off critical inflammation, which a diet of corn might increase, but why it is persisted in with animals suffering from general prostration and weakening complaints is quite paradoxical.



It is a very common practice in some places to leave the bran mash, even for days, before the sick animal. The mixture ferments, fouls the woodwork, and nauseates the sufferer, who shows his disgust by standing as far away as his chain will allow.

#### SYSTEM PURSUED AT THE LONDONDERRY COLLIERIES.

The horses and ponies employed at the collieries of the late Marchioness of Londonderry (now Earl Vane's), during my appointment as Veterinary Surgeon to her Ladyship, and subsequently for a short period to the Earl, were fed upon oats and peas—five parts of the former to one of the latter. The mixture was, however, never regulated by weighing. It was quite of an extemporaneous character, and entire guess work, a dash of peas being hastily put into the bottom of a "*poke*," and afterwards covered to the top with bruised oats, and the *whole* then weighed. In addition, hay, and green food in summer, was allowed, the cost of which I find by returns in my possession, amounted to £1 3s. 10½*d.* per horse per pay, all ponies being classed as two equal to one horse.

Under this arrangement 106 horses and 322 ponies exclusively employed in producing coal, and of course below ground principally, would in round numbers give 267 horses, the cost of feeding each being £31 0s. 9*d.* per annum.



At first sight this appears, and is really believed to be a very economical rate of feeding, being lower than Mr. Scott's expenditure at Hetton-le-hole. Quality, however, is the test of cheapness, not the price alone at which the food is supplied. This allowance yields not more than 37·52 pounds of nutritious matter—assuming the one part of peas are carefully added, which I have grave reasons to doubt—against 48·18 in Mr. Scott's feed, in which 10·66 lbs. extra are supplied each pay per horse during the year, for which he incurs a cost only of 9s. 9d., the difference per horse per year.

But, as in the case of the old system at Hetton-le-hole, the animals do not derive the whole of even this limited amount of benefit from the corn allowed. In winter *steamed food*, consisting of hay, linseed, and the one-fifth part of peas abstracted from the mixture, are supplied. This is sent down the pits during the afternoon, and frequently before the animals reach the stable in which the mess has been deposited by the horse-keeper, I have found it undergoing fermentation. Some, in fact many, of the animals refused it even when fresh, and from these causes the most nutritious portions were lost to them.

The process of steaming was conducted at separate places contiguous to a number of collieries, and the food conveyed in boxes or coal-tubs placed in carts, and throughout the distance a dark-coloured fluid drained in profusion, of



course carrying with it some of the most useful soluble principles of the food.\*

The process of steaming food may answer well where inferior food and hay abound, cattle are to be fed, and aged dependants require some light employment. But as applied to good food, and carried out in the manner alluded to, it is a complete farce, a useless addition to expenditure—which, by the bye, is not made to appear in connection with the feeding—and deprivation to the hard-working animals. In fact, the whole system of feeding is no better, hence the number of cases of colic and indigestion, rapid falling away in condition, diabetes, &c., &c., which occur, and not being required to be reported at headquarters nothing is known of them. But if the gentlemen who attempt to feed horses without visiting them at their work or in their stables, and others who rule with pens and ink in a comfortable office, had to ride off at all hours in all kinds of weather three, six, eight, or ten miles, descend a coal-mine, remain in the heated atmosphere several hours, or take the place of their coachman *outside* the conveyance in the cold night air, after influences equal to a severe

---

\* In a visit to a large estate owned by a noble earl, where feeding is supposed to be conducted upon scientific principles, steamed food forms a large item for cattle. After the process is carried sufficiently far the food is removed, and the fluid accumulated from the condensed steam, containing useful soluble matters, is allowed to drain away in the gutter, while the animals are supplied with water for drinking. *Cui bono?*



vapour bath, each would perceive the advisability of reform. As such irregularities are in a coalpit hundreds of feet below the surface, they *do not see*, and it is very useful to be *determined not to see*.

With the exercise of common knowledge a man may be expert at purchasing corn, and make good bargains, but it requires a little more than that to adapt a proper system to the constantly varying wants of a large number of animals. If the principles of dieting and management require no more philosophy, and can be conducted with success without presence or information, the sooner such gentlemen take up also the treatment of disease by a system of telegraphy the better, and probably they may then make both ends meet still more satisfactorily to themselves.

From a strict examination of all phases and conditions, as they exist on that estate in common with many others, a more highly nutritious diet can be guaranteed—a reduction of disease and mortality therefrom to a minimum by the exercise of care and watchfulness, and these with a saving of *hundreds of pounds annually*.

The principle is of easy calculation. In large numbers, say three or four hundred animals, if two or three shillings only per week can be saved, it is a large sum at the end of the year. With other agencies it may also be increased. But on collieries there are objections to figures, except when they are used by the powers that be, and



reports, although desired, are as waste paper. Like the phoenix, however, these may rise from their ashes, and, under another name, with an official envelope, appear as new and *original* creatures altogether.

In leaving this part of the subject, I would enquire how it is possible for a pit horse of fifteen or sixteen hands, with his large muscular system—doing from twenty to thirty miles a day, drawing a train of coal tubs, one way empty the other laden with many tons, in a highly heated and dusty atmosphere, and perspiring freely—to obtain sufficient nutriment from an allowance of twelve pounds of oats per day. The same is allowed to many carriage horses, and those of our cavalry regiments receive it within a fraction, for which exertion amounting to healthy and necessary exercise only, compared with the work of a pit horse, is exacted.

#### OTHER FORMS OF ADMIXTURE.

I have shown that nutritious mixtures of corn may be used with great advantage at a low cost. Their use and application should be guided by existing circumstances.

Animals should be selected as much as possible from an average class, character, and age; but an intelligent overlooker would soon be able to perceive, even among a mixed class where any differences existed, and modification is required.



Some would need a little more food than others—particularly aged ones—while there are others whose assimilative powers being more active, would derive more nutrition even from a little less food, and thus spare a trifle to their less endowed neighbours.

In the studs under the care of Mr. Hunting at South Hetton, Murton, Rhyhope, Trimdon Grange, Seaton Delaval, &c., &c., and Mr. Scott, in the old Hetton Collieries, animals of all ages are to be found, and, by the system carried out, are equally well provided for.

If, owing to the severity of the work, a more nutritious diet is called for, it will be found in the forms as follow, which provide it at an equally cheap rate.

*Mixture No. 1 (Low rates).*

				Weight.	Cost.		
				lbs.	£	s.	d.
Peas	4 bushels	.	.	264	1	0	0
Barley	4	„	.	224		15	0
Oats	3	„	.	126		9	9
Bran*	.	.	.	98		4	1
				<hr/> 712	<hr/> £2	<hr/> 8	<hr/> 10

This mixture, divided among seven horses during one week, would allow each over  $14\frac{1}{2}$  pounds per day, at a cost of 6s.  $11\frac{3}{4}d.$ , out of which 20·14 lbs. of N.M. (nutritious matter)

---

\* The nitrogenous principle of bran is not calculated in these mixtures for reasons stated on page 83.



would be obtained. Divided among six horses each would cost 8s.  $1\frac{1}{2}d.$ , and obtain 23·63 lbs. of N.M.

Divided among five horses, each would cost 9s. 9d., and obtain 28·36 lbs. of N.M.

*Mixture No. 2.*

				Weight.			Cost.		
				lbs.			£	s.	d.
Peas	4 bushels	.	.	264			1	0	0
Barley	4	"	.	224				15	0
Tares	1	"	.	67				5	$4\frac{1}{2}$
Oats	1	"	.	42				3	9
Bran		"	.	98				4	1
				<hr/>			<hr/>		
				695			£2	8	$2\frac{1}{2}$

Divided among seven horses, each would receive over 99 lbs. per week, costing 6s.  $10\frac{1}{2}d.$ , and obtain 21·65 lbs. N.M.

Six horses would receive  $115\frac{1}{2}$  lbs. each, costing 8s., and obtain 25·26 lbs. N.M.

Five horses would receive 139 lbs. each, costing 9s.  $7\frac{1}{2}d.$ , and obtain 30·31 lbs. N.M.

*Mixture No. 3.*

				Weight.			Cost.		
				lbs.			£	s.	d.
Peas	4 bushels	.	.	264			1	0	0
Barley	4	"	.	224				15	0
Tares	2	"	.	134				10	9
Bran	.	.	.	98				4	1
				<hr/>			<hr/>		
				720			£2	9	10



Divided among seven horses, each would receive over 102 lbs. per week, costing 7s. 1 $\frac{1}{4}$ d., and obtain 23·86 lbs. N.M.

Six horses would receive 120 lbs., costing 8s. 3 $\frac{1}{2}$ d., and obtain 27·84 lbs. N.M.

Five horses would receive 124 lbs., costing 9s. 11 $\frac{1}{2}$ d., and obtain 33·40 lbs. N.M.

It must be borne in mind that good hay yields five per cent. of N.M. which must be added to the above for the quantity used.

It will also be observed that peas are named in the foregoing mixtures to the exclusion of beans. They are not so productive of constipation as beans, which on that account enables us to use them with greater freedom. Beans are, however, unobjectionable when used with oats and bran, chaff, &c., in sufficient quantities. Prices must regulate these mixtures from time to time, due consideration being paid to the N.M. contained in the various kinds of grain.

When the prices of grain are much increased, such mixtures as the following must be taken, and will be found adequate at an equally low cost.

From the *Field* of 21st September, 1867, we learn the prices current at Mark Lane were, for oats, 35s.; barley, 43s.; peas, 43s.; beans, 44s. per quarter of eight bushels, and tares, 7s. per bushel. Bran cost at that time 6s. per cwt. of eight bushels.



*Mixture No. 4 (High rates).*

				Weight.	Cost.		
				lbs.	£	s.	d.
Peas	6 bushels	.	.	396	1	12	3
Beans	4	„	.	264	1	2	0
Tares	2	„	.	134		14	0
Barley	1	„	.	56		5	6
Bran	.	.	.	112		6	0
				962	£3	19	9

Divided among twelve horses, each would receive 80 lbs. per week, costing 6s. 7 $\frac{3}{4}$ d., furnishing 21 $\frac{1}{2}$  lbs. N.M.

Eleven horses would receive 87 $\frac{1}{2}$  lbs. each, costing 7s. 3d., and obtain 23 $\frac{1}{2}$  lbs. N.M.

Ten horses would receive 96 lbs. each, costing 7s. 11 $\frac{3}{4}$ d., and obtain 25 $\frac{3}{4}$  lbs. N.M.

Nine horses would receive 106 $\frac{1}{2}$  lbs., costing 8s. 9 $\frac{1}{4}$ d., and obtain 28 $\frac{1}{2}$  lbs. of N.M.

Eight horses would receive 120 lbs. each, costing 9s. 11 $\frac{1}{2}$ d., and obtain 32 $\frac{1}{8}$  lbs. of N.M.

*Mixture No. 5.*

				Weight.	Cost.		
				lbs.	£	s.	d.
Beans	4 bushels	.	.	264	1	2	0
Peas	4	„	.	264	1	1	6
Tares	2	„	.	134		14	0
Oats	1	„	.	42		4	4 $\frac{1}{2}$
Bran	.	.	.	112		6	0
				816	£3	7	10 $\frac{1}{2}$



Ten horses would each receive  $81\frac{1}{2}$  lbs., costing 6s.  $9\frac{1}{2}d.$ , and obtain 21 lbs. N.M.

Nine horses would each receive  $90\frac{1}{2}$  lbs., costing 7s.  $6\frac{1}{2}d.$ , and obtain  $23\frac{1}{2}$  lbs. of N.M.

Eight horses would each receive 102 lbs., costing 8s.  $6d.$ , and obtain  $26\frac{1}{2}$  lbs. of N.M.

Seven horses would each receive  $106\frac{1}{2}$  lbs., costing 9s.  $8\frac{1}{4}d.$ , and obtain  $30\frac{1}{2}$  lbs. of N.M.

*Mixture No. 6.*

					Weight.			Cost.		
					lbs.			£ s. d.		
Beans	4	bushels	.	.	264			1	2	0
Peas	4	"	.	.	264			1	1	6
Barley	3	"	.	.	168			0	16	$1\frac{1}{2}$
Tares	2	"	.	.	134			0	14	0
Bran	.	.	.	.	112			0	6	0
					<hr/>			<hr/>		
					942			£3	19	$7\frac{1}{2}$

Twelve horses would receive per week each,  $78\frac{1}{2}$  lbs., costing 6s.  $7\frac{1}{2}d.$ , and obtain  $19\frac{3}{4}$  lbs. N.M.

Eleven horses would receive each  $85\frac{3}{4}$  lbs., costing 7s.  $2\frac{3}{4}d.$ , and obtain  $21\frac{1}{2}$  lbs. N.M.

Ten horses would receive each 94 lbs., costing 7s.  $11\frac{1}{2}d.$ , and obtain  $23\frac{3}{4}$  lbs. N.M.

Nine horses would receive each 104 lbs., costing 8s.  $9d.$ , and obtain 26 lbs. N.M.

Eight horses would receive each  $117\frac{1}{4}$  lbs., costing 9s.  $11\frac{3}{4}d.$ , and obtain  $29\frac{1}{4}$  lbs. of N.M.

Seven horses would receive each  $134\frac{1}{2}$  lbs., costing 11s.  $4\frac{1}{4}d.$ , and obtain  $33\frac{3}{4}$  lbs. of N.M.



The present prices of grain, &c. (February, 1871), afford the means of suitable contrast with the foregoing. By using the following mixtures, actual cost need not be increased.

*Mixture No. 7.*

				Weight.			Cost.		
				lbs.			£	s.	d.
Peas	8 bushels	.	.	504			2	0	0
Beans	4	„	.	252			1	6	0
Barley	4	„	.	200			0	15	0
Oats	4	„	.	152			0	12	0
				<hr/>			<hr/>		
				1108			£4	13	0

Fourteen horses would receive 79 lbs. per week, at a cost of 6s.  $7\frac{3}{4}d.$  each, yielding 21 lbs. of nutritious matter.

Thirteen horses would receive 85 lbs., at a cost of 7s.  $2d.$ , yielding  $22\frac{3}{4}$  lbs. N.M.

Twelve horses would receive  $92\frac{1}{4}$  lbs., at a cost of 7s.  $9d.$ , yielding  $24\frac{3}{4}$  lbs. of N.M.

Eleven horses would receive 100 lbs., at a cost of 8s.  $5\frac{1}{2}d.$ , yielding 27 lbs. N.M.

Ten horses would receive  $110\frac{3}{4}$  lbs., at a cost of 9s.  $3\frac{1}{2}d.$ , yielding  $29\frac{3}{4}$  lbs. N.M.

*Mixture No. 8.*

				Weight.			Cost.		
				lbs.			£	s.	d.
Peas	8 bushels	.	.	504			2	0	0
Barley	4	„	.	200			0	15	0
Oats	4	„	.	152			0	12	0
				<hr/>			<hr/>		
				856			£3	7	0



Ten horses would receive 85 lbs. per week, at a cost of 6s. 8 $\frac{1}{4}$ d. each, yielding 21 lbs. N.M.

Nine horses would receive 95 lbs., at a cost of 7s. 5 $\frac{1}{4}$ d., yielding 24 lbs. N.M.

Eight horses would receive 107 lbs., at a cost of 8s. 4 $\frac{1}{2}$ d., yielding 27 lbs. N.M.\*

It remains only to be added that by a careful estimate of the prices of grain and leguminous seeds, together with the relative amounts of nutritious matter contained in each, it is possible to vary the mixture in such manner as the foregoing, that, notwithstanding the excessive fluctuations of markets, the cost of each animal per week for corn will not vary throughout the entire year more than a few pence.

The bulk will, however, be somewhat less, and requires to be made up by the use of hay, or hay and straw chaff.

As these mixtures are designed for hard-working animals only, it will be almost unnecessary after what has been said when speaking of animal heat, &c., to state that such food will prove pre-eminently injurious to idle animals, except given in very small quantities. To those doing no work, or only exercise, oats are decidedly the safest article of diet.

---

\* In feeding horses upon oats only, weighing 38 lbs. per bushel, costing 24s. per quarter, each must consume 5 bushels weekly, at a cost of 15s., in order to derive the same nutrition as here afforded by a mixture of 27 lbs. per day, at a cost of 8s. 4 $\frac{1}{2}$ d. per week.



---

GREEN FOOD.

Among hard-working horses, I have found the use of green food too indiscriminately adopted, and it frequently proves as injurious as a mass of cooked food.

That it is useful and beneficial I fully believe, but this occurs under proper management. When given, it should by no means take the place of corn. This is, however, too commonly done, the animal being allowed too much.

He then perspires freely, becomes weak and liable to disease. I have known for years that among town cart and cab horses to whom grass has been allowed as described, sore throats and influenza have appeared almost simultaneously with its use, besides colic, weed, &c., &c.

Two or three weeks' feeding while the plants are young may answer for easily wrought horses, but in my opinion that state of the animal's body which we term condition should not be so trifled with. Aptitude for work is not a condition which is obtained in a few days or hours. It is only obtained during weeks, and at considerable expense. Most persons know, or at least presume to know, what training the racer requires, and how long a hunter is in being got ready for the field; yet we find the same people sacrifice the condition of a draught horse during a season



when he requires the greatest strength, by the supply of an innutritious and watery food.

I maintain, if horses are in condition and required for work, that green food is an interruption to the formation and maintenance of muscle, and it should be avoided. If he requires rest, green food will be serviceable ; and, like his more favoured master, he should be allowed a trip to the country, where, in a roomy loose box or covered yard, he can enjoy the green fruits of the earth, which are regularly mown and brought to him, protected from the rays of a scorching summer sun, or the pelting rains that fall, and flies which torment at this season.

When these conditions are not fulfilled, green food proves expensive and unprofitable. The animal loses condition and is below the standard for work, while he becomes peculiarly liable to disease, and exhibits the greatest difficulty in taking on flesh, appearing lean, hide bound, and generally betokening want of power and activity.

Similar remarks apply to the use of turnips or carrots. Moderately used raw in winter they prove serviceable—in excess they are positively injurious.

The use of green food, in accordance with common-sense principles, answers among pit animals. Although it favours perspiration, the temperature being uniform, the results are not so pernicious as among animals so fed and worked above ground.



---

## ECONOMY OF USING CHAFF AND BRUISED CORN.

With regard to the bruising of corn and cutting hay into chaff, I find the practice attended with good results. It suits a mixed class of animals better, and causes a more perfect mixture of each kind than would be derived from the food in the whole state.

It is an important proceeding where boys are employed ; for where they have access to unbruised corn, they often take out beans, peas, or tares for pigs, rabbits, poultry, or pigeons at home.

The cutting of food, as already shown, causes a proper and continued flow of saliva—an important process, in order to ensure digestion of the numerous starchy elements of food upon which horses and cattle live. The cost\* incurred is very little with proper machinery, and amply repays for the outlay in a very short time.

By the addition of *oat* or *wheat-straw*, a saving is not only effected, but the food is further aided in digestion. The usual proportions are one-fourth straw and three-fourths hay, but many persons use equal parts.

These are strictly non-nutritious agents as used. They are given purposely for providing

---

\* It is frequently estimated that hay costs 2s. per ton cutting, and the bruising of corn one penny per bushel. Mr. Hunting, however, informs me that the work is performed at the South Hetton colliery for half these sums.



necessary bulk, and assisting in grinding down the other portions of food within the digestive organs. As an agent of nutrition hay is very expensive, and in that respect little superior to straw, which, with other food, I have known given regularly to animals, no hay being allowed, while the condition has been all that could be desired.

In all the cereals and leguminosæ ripeness of the plant is evidenced by the development and perfection of the seed in the various receptacles; prior to that stage the stem contains the nutrition. Notwithstanding this fact, so apparent as it must be to every farmer that the analogy exists in each, hay is allowed to stand for days, and even weeks, before being cut, when it must be an utter impossibility for any increase to take place. Growth is completed, the plant ripens, as indicated by the seeds becoming plump. They contain all the nutriment, and in the main are shed upon the ground by every wind that blows, leaving the hay nothing better than a fine sample of straw.

The higher price asked for this fine sample, we may infer, is to cover the loss occasioned by this species of neglect.

I wish it to be understood that I am not prejudiced to the exclusive use of cut hay and bruised corn. When given whole these substances may be effectively masticated, and the animals will appear excellent and pay well for



the proper selection and mixture of food. I, nevertheless, consider that a larger profit accrues by *avoiding the waste* which always happens when hay is given whole (or long), besides suiting for general purposes much better.

Food so prepared will require *wide and deep mangers* fitted with cross bars, to prevent the animals throwing it out in their search for the more tasty portions.

On this subject many investigations have taken place at home and abroad, and the results are, without exception, favourable.

In France, Leblanc found that animals fed upon boiled food fattened, but lost vigour and became affected with disease; while dry food had a contrary effect. M. Charlier says, with cooked food the animals are very subject to disease, but since the regular adoption, by omnibus proprietors, of dry, bruised, and cut food, founder, colic, and illnesses generally, which each week were numerous, had become quite exceptional.

#### PEA AND BEAN STRAW.

The question has frequently been put to me, "Is pea or bean-straw useful for feeding horses?" To this an affirmative answer may be given, when the article has been carefully gathered and housed. It should always be cut up small, and mixed with



the bruised mixtures of corn, and large quantities should be avoided by using with it cut hay. Equal parts of hay and pea-straw, or three parts of hay to one of bean-straw, will answer very well for working horses, but I deprecate filling the racks with either.

The many fatal cases of disease which occur from their use, is owing to the inferior condition of the straw, coupled with being in undue quantities and other irregularities, which too often exist in the farm stable.

#### SAVING TO BE EFFECTED.

Carefully carried out, these principles will effect an enormous saving, and I have no hesitation in stating that, among the numbers of animals employed in Glasgow and other large towns of Great Britain, it may be raised to *thousands of pounds annually*, and, in addition, a corresponding sum by the avoidance of disease.

The farmer need not grudge his horse corn, since it can be shown that he can be fed at less cost than many now incur on inferior hay, straw, and provender generally, and derive greater aptitude for work.

The poor man may also feed better and save a few shillings weekly, which would be well applied to the formation of a fund wherewith he could replace his animal in the event of death, without



resorting to the subscription system over his immediate district, which is too commonly done.

The principles are important to large owners, who might also raise a fund with the savings for the same purpose, and to remunerate, in some cases I could name, those who are able, and study to apply effectually the system calculated to bring about the change.

*All food should be of the best quality and kind.* If inferior kinds are purchased, and require disguising by some process to make them palatable, the sooner he who purchases such is removed the better.

No mixture or process of doctoring can render such available as nutritive food when the elements are not present, nor can any admixture of condimental nonsense effect it either. The experiments of Mr. J. B. Lawes have definitely proved that when animals improve their condition with the use of condiments—which in themselves have no nutrition scarcely—the result occurs from an *increased consumption of corn*. The proprietors of such compounds may state what they please in puffing their wares. But against their puffing it can be positively asserted, as an undoubted fact, that condiments, spicy foods, or by whatever term they are known, do not effect any improvement upon inferior food with which they may be mixed, and are not economical, as set against common salt, which is far superior at one-fortieth the price. On the other hand, they



are expensive articles of diet, and can only be looked upon as substances quite dispensable, and of no great service in the feeding of animals.

#### PRACTICAL TESTS OF THE SYSTEM.

We have endeavoured to show, from the testimony of others, to what extent the system advocated in these pages may be productive of financial success, as well as securing general health and ability for work. It remains for us to notice a few, from many similar instances of equal success which have come within direct personal observation.

Shortly after the first edition of this present work left the press, we were consulted by the proprietor of a single horse in reference to his animal, which was suffering from severe colic. The case was pronounced hopeless from the first, and as his plan of feeding was known to be the principal cause, from which four other animals had died, we pressed upon him the necessity for reform. The usual plan of cooking sound food had been long practised, and *the weekly cost averaged twenty-one shillings.*

Having become the possessor of the volume he proceeded seriously to study its pages, and, although nearly convinced, he hesitated to adopt the plan, fearing that he might not succeed. The men he met with on the road dissuaded him from it. They declared the system to be false;



and when he produced the book, that was said to be a tissue of lies. A friend, who advanced money for the purchase of another horse, also strongly supported our advice, and the man consented. Instructions were given for making up a mixture, which was compounded and paid for each week out of his earnings, the benefit of which was rapidly apparent. After two or three weeks the animal, having improved so much, he made four extra journeys per week to the colliery, distant about three miles, *realising one pound, at the same time the weekly cost of feeding had been reduced from twenty-one shillings, the cost of the cooked food, to fourteen shillings, making in all a clear gain of twenty-seven shillings per week, or seventy pounds four shillings per annum, in respect of that one animal.*

This man was under our observation about three years. He became the proprietor of other horses, which were engaged in the same work—that of carting coals from a colliery outside Glasgow to some ironworks. His animals remained in splendid condition, and, during the time above stated, he had neither lameness, illness, nor death.

A firm of coal merchants at Leighton Buzzard were induced to give the plan a trial, and four horses were made the subject of the experiment. The report, within a month, was to the effect that the health of the horses was all that could be desired. They were doing 20 per cent. more



work, with less fatigue—equal to a saving of one pound per week per horse; while in the feeding, the saving per horse per week, the cost was seven shillings and sixpence less than on the old system. The aggregate annual result was a clear saving of £286 on the four horses.

The following communication is dated from

“HERTFORD,  
3rd August, 1878.

“MR. GEORGE ARMATAGE

“DEAR SIR,—

“I have been working two broken-winded horses, which I bought cheap some months ago. They have been in constant work (*very hard work*) up to the present time, and seem to do their work with ease. I have been feeding them on a mixture of grain, according to your system, with wetted chaff. The object of this note is to ask you whether it would be safe to give them dry food?

“And may I trouble you to let me know if broken-winded horses are worse (*cæteris paribus*) in summer or during the winter season?

“I must apologise for troubling you, which I do only after a fruitless search for the information in your very valuable work.

“Believe me,

“Yours truly,

“GEORGE R. DURRANT.

“P.S.—You will feel interested in the fact



that we have been saving £1 a week by the adoption of your really scientific method of feeding. The saving is really very much larger than stated, as our *four horses* (costing 9s. 8d. each per week) have done 10 per cent. more work than *seven horses* (costing 14s. 6d. each per week). Further, the horses we work this year cost £54 the four. Last year the seven cost £190."

It should be stated, for the information of the reader, that the writer of the foregoing communication was at the time engaged in the manufacture of mineral waters, which constitutes a summer trade. The animals are, therefore, purchased each year, and used only during that time, being sold at the close of the season. A lower class horse answers the purpose, constituting a saving in the outlay; yet, with the system adopted, although defective in wind, and, as we are informed, they were doing "very hard work," a considerable and satisfactory economy was pursued.

With regard to the suitability of the system for broken-winded horses, what is conducive to the comfort and well-being of the healthy horse is eminently suited for those suffering in the form alluded to. The old and defective system is the bane of animals sound in wind. That advocated by us is preventive of the disease, and palliative to those diseased.



A later communication brings the information up to date :

“ HERTFORD,  
3rd May, 1890.

“ MR. GEORGE ARMATAGE.

“ DEAR SIR,—

“ I regret that, owing to absence from business, I have only to-day seen your letter of the 30th ult.

“ Until I used your method of feeding I was always in trouble with my horses, but after reading your book and thoroughly mastering its contents, and carefully carrying out the suggestions your book contains, my horses improved rapidly, and I saved at least 10 per cent. on the cost of feeding stuff, and the horses did more work.

“ You are welcome to make any use you like of this.

“ Both of your books (“ Stable Management ” and “ The Horse ”) are the best treatises on the respective subjects I have ever read.

“ Yours faithfully,

“ GEORGE DURRANT.

“ P.S.—I have sold my business, and have no horses at the present time.—G. D.”

In an article, entitled “ Tramway Traction,” which appeared in the *Railway and Tramway*



*Express*, p. 219, April 4, 1885, by C. Challenger, Traffic Manager of the Bristol Tramways Company, Limited, the following observations are made :

“ Mayhew, and Youatt, and Stonehenge have been excellent authorities in their time, but other and better authorities have stepped on their heels and forced them off the stage. The veterinary profession has of late years made rapid strides since the above-named wrote, and some of its members have been entrusted not only with the medical care of hundreds of horses of large firms, such as are to be found at the extensive North of England collieries, but they have had, at the same time, the arrangements for the feeding of these horses as well. They have carried science into their principles of feeding, and have ascertained with exactness the requirements of the horse under every condition ; and they have done the world the service of publishing an account of the results of their scientific experiments and practice.

\* \* \* \* \*

“ With regard to the feeding and general care of horses, applying especially to tramway horses, the most modern, and I believe the only one that has reduced the feeding of horses to a science, is a book, entitled ‘ The Horseowner and Stableman’s Companion,’ by George Arma-



tage. It was written in 1878,\* and published by F. Warne & Company, Bedford Street, Strand. Price 5s.

“When the teachings of this book are put more into general practice, I should expect to see the condition of our useful friend, the tramway horse, raised above its present crestfallen state.”

Referring to the duration of life in horses drawing public conveyances, the writer observes :

“When a ‘big man’ makes an assertion he is generally believed. Thus it behoves all who come under this appellation, when they make an assertion, especially before a public tribunal, to be correct. A statement was made some few years ago, before a Committee of the House of Commons, by, I believe, Dr. Wood, who was then Managing Director of the Edinburgh Tramways, to the effect that the life of a tramway horse was only four years ; and I find that no longer ago than November last, in the sketch of the North Metropolitan Tramways appearing in the *Railway and Tramway Express*, it was stated

---

\* The work originally appeared in 1868, as, at the present, in the form of two manuals, one expressly devoted to the principles of feeding, under the title “How to Feed the Horse, Avoid Disease, and Save Money ;” the second, “The Horseowner and Stableman’s Companion,” dealing with Stable Management.



that the average life of a tramway horse was from four to five years.\*

“ I consider these statements are not reliable. I allow seven years. A horse is five years growing to maturity, and philosophers state that the laws of Nature allow every animal, including man, if it has fair-play, to live six times longer than the time it takes to pass through youth and arrive at maturity. This would give to the horse a life of thirty years, and to man one hundred and twenty. The horse is fit for tramway service at five, and not a week before. He should be able, if cared for in a proper manner, and fed on nutritious food—food that would impart muscle and bone—to draw a tramway car until twelve years old. His knees would probably be “ gone over,” more or less, after seven years’ trotting on the paved road, although there are able-bodied horses at the present time in the service of the Bristol Tramways which work side by side with six and seven years’ old horses, and which have been working regularly ever since the opening of the first section, in August, 1875, nearly ten years. They are neither over at the knee nor ‘ out at the elbow.’ What is there about the

---

\* This estimate had its origin in respect of horses used in omnibuses. Many died as the result of hard work, defective feeding and management, during the first and second years of their introduction.



work, taking into consideration the long rests, to use up a horse in four years?

"In Armatage's book, life-sustaining principles of feeding are treated of in a masterly manner.

"The weekly amount of corn given to a tramway horse should yield, at least, 22 lbs. of nutritious matter. Bran, hay, and straw are given as bulk, to bring about a healthy assimilation, and are not taken into account when ascertaining the amount of nutritious matter.

"He says, at page 82: 'The amount of nutritious matter must be accurately calculated, or no reliable estimate of cheapness can be made.'

"Again, on page 73: 'Considerable amount of nutritious matter is required to keep up the health and condition of animals. . . . We are also taught practically that a loss of condition and proneness to disease are brought about by food containing too little nutrition, when the demands upon the system are excessive.' And, page 70: 'The method of economising food consists in using a variety of grain, instead of one kind.'

"The horses of the Bristol Tramways have been fed, during the last few years, on the following:



	Per day. lbs.	Per week. lbs.	N.M. Per cent.	N.M. Weekly.
Barley . . .	5½	38½	18	6·54
Beans . . .	3	21	31	6·72
Oats . . .	3	21	14	2·94
Peas . . .	2	14	32	4·48
Maize . . .	2	14	12	1·68
Hay . . .	11	77	„	„
Straw . . .	5½	38½	„	„
Bran . . .	1	7	„	„
<hr/>				
Daily . . .	33			22·36

“The amount of nutritious matter is ascertained by simple rule of three.

“The cost per horse per week, at the present price of corn, is 10s. 2¾d.; peat bedding, 30 lbs., 6d.; total, 10s. 8¾d. Horses do well on the above. The mortality of the stud speaks well for the health-ensuring properties of the food. Since November 30, 1883, out of 372 horses, only four deaths have occurred.

“The corn given by the Glasgow Tramways Company, according to the article, ‘Our Tramways,’ yields, per week, 15·54 lbs. of nutritious matter, and that of the Dublin Tramways 15·26 lbs. of nutritious matter. Their horses may be smaller than those of the Bristol Tramways, as they have fewer hills to contend with. If it were otherwise, experience teaches one that they would be underfed.”

Some years ago we were entrusted with the



management of a large stud of horses, the property of a tramway company. At the time, all kinds of provender were realising the highest prices. Sainfoin was £7 per load of 36 trusses; peas and beans, 42s. per quarter; oats, 36s.; straw, 40s. per load; other kinds of food in proportion. The works not being completed, advantage was taken of the available time—some two or three weeks—to get the animals into condition. They were then costing, on short allowances, eighteen shillings per week. We had no machinery, nor any convenience for food preparation during this time, but we supplied a more nutritious kind at a cost of about ten shillings, and upon this the animals thrived rapidly. They also entered upon their work in surprising condition, doing first twelve, and, as further extension of the route took place, eighteen miles per day.

Owing to the delay in putting down machinery and adapting the buildings, &c., the system could not be pursued as we intended; therefore, the cost of food increased, as a natural consequence, as soon as the work commenced. No bruising could be done, and we considered the feeding was not so complete nor so cheaply carried out, the cost being about fifteen or sixteen shillings. Shortly, however, the requisite conveniences were supplied, and we could cut up hay and straw into chaff, instead of purchasing it so prepared. The grain was also bruised and transferred to



the floors, where thorough incorporation was performed. The cost, under the rates already quoted, never exceeded *thirteen shillings and sixpence, including straw for bedding*. During eighteen months the animals continued to do well; not a single case of colic occurred, nor indeed any serious affection. The sole cause of idleness was lameness arising from over-driving and accidents due to the ignorance of drivers not previously accustomed to horses. At the time also influenza prevailed largely in the metropolis, and created appalling havoc, many proprietors and public firms suffering absolute paralysis of work; their animals were suffering and dying in large numbers. On the other hand, our horses continued at work nearly throughout, suffering slightly, but we had no deaths.

After we had pulled through these difficulties, "big men" interfered, and we threw up the responsibility. Had we been permitted to see the prices of grain, &c., reduced to usual rates, the saving would have been enormous, the total cost not being more than nine or ten shillings weekly per horse, with a corresponding healthy condition and immunity from disease. The result of the change was serious. At the expiration of two years we were twice invited to resume the management, as the losses from disease were numerous and the ability of the stud at the lowest. This we could not see our way to



accept, as no guarantee from unwarranted interference could be ensured.

### WORK.

The subject of equal importance with feeding is work. It is of little consequence what system is adopted—the quantity or quality of the food or the monetary saving—if there is no care exercised in the imposition of work. It is possible for men, either ignorant of or indifferent to the powers of animal endurance, to frustrate every good effort and intention of those who have charge of the feeding department, and such results have been shamefully accomplished in notable instances.

In the management of horses the all-important object is the avoiding of loss. To keep animals capable only of limited exertion, constantly out of condition or ability, and sickening after every unusual effort, is one source of loss; the other lies in overwork, from whatever cause. It is, therefore, of paramount importance to be able, first, to create the true ability for exertion—in other words, to establish the best of health—and while that is duly maintained by an intelligent system of feeding, to apportion the work which shall lie within the ability of the animal. It is far from true economy to push exertion to the extent of exhaustion. The horse should come in



from his labour, in a sense, "fresh." When he is tired—that is, suffering from muscular exhaustion—every organ in his frame is equally exhausted. In this state the natural secretions are perverted—altered in constitution as well as quantity; they fail to act suitably on any food he may take. Nature often takes away the appetite under such circumstances, and the animal suffers less. But when exhausted conditions are accompanied by a ravenous appetite—which, by the way, is an unhealthy condition—then follows various forms of indigestion, which saps the source of profit.

Work should be so apportioned that general health shall not be interfered with. When a good system of feeding is adopted, the regular imposition of labour secures a full digestion and appropriation of the elements of nutrition. Blood, bone, and muscle are fed and sustained in such degrees of tone and vigour as to indicate the highest ability for exertion. This we term "condition," or the standard of excellent health, and should not be trifled with on any consideration. As long as this state is ensured by food, the imposition of regular and properly apportioned labour will maintain the balance of healthy function, and secure the evidences of ability by the comparative ease with which every effort is accomplished. Overwork destroys all this. It is betokened by loss of tone, marked inability, failing condition, consumption of a



larger quantity of food, inferior digestion, liability to disease, and inability to perform the daily portion of even moderate work. These indications, although happening in the order given, are not generally interpreted as soon as they should be by many to whom the responsibility of management is entrusted. Many are wilfully blind, and seek to make capital out of the malady.

The great barrier which often exists and stops the way of progress is an objection to allow other persons the claim to special knowledge. The desire to be first swamps every principle of right, often regardless of consequences, which exhibit themselves in the shape of enormous expenses, if not paralysis and ultimate failure of the concern.

If public companies could see their way to commission the veterinary surgeon to undertake the feeding and management of their studs on the principle of preventing disease, they would find it to be the most profitable undertaking. The traffic manager and road inspectors should act in concert with the veterinary surgeon; otherwise, all the latter may do may be speedily undone by the two first. This has been painfully apparent in repeated instances which have come beneath the writer's notice. One example may be taken. The veterinary surgeon to a large establishment was informed that an appointment had been made of the character



referred to, and he was requested to supply horses as they were wanted, without having the liberty of controlling the amount or nature of their work. He remonstrated, but could not obtain a hearing; all he could do was to quietly submit. The officers appointed were ex-police-men, having no knowledge of matters pertaining to the horse. The end of things soon came to pass. They imposed additional tasks to the already overworked animals, often taking them out twice during the same day, and always working seven days each week. In addition, they were stationed too far from their work, having to proceed nearly two miles to and from the place of starting. From that time the success of the system was totally frustrated, ending in absolute failure. Those who formerly had the management and responsibility wisely retired, but the eyes of the powers above were so securely closed that they only acquired full and complete vision when the concern was verging upon general collapse.

In judging of the amount of work to be imposed, some consideration must be given to the pace or rate at which the animals must travel, the kind of carriage which contains the load, the size of wheels, nature—especially gradients—of the road, and last, but not least, the kind of anti-friction material used, commonly called grease. The slowest pace, it will be naturally conceded, involves the least exertion; therefore,



conversely, the fast pace, if with it are combined defective carriages, small wheels, bad grease, heavy gradients, frequent stoppages, and an inhuman driver, reduce the best animals to the worst condition. In level towns the difficulties are proportionately lessened, and a faster pace is tolerated. Tramway horses have not only the exhausting effects of frequent startings, the moving of a dead, enormous weight, with the disadvantage of short levers in the shape of small wheels, but additional drawbacks in the friction involved in passing the numerous junctions and injudicious use of the break, as well as the interposition of dirt upon the metals. We have known the condition of these animals to be seriously damaged by the wheels being too wide for the metals, a stream of fire in dry weather issuing from the flanges as the wheels ground against the sides of the groove.

The difference in the severity of the exertion required of tramway horses and those used in omnibuses is very apparent. The larger wheels of the latter, as well as being close together, greatly reduce the severity of effort in starting. The horses also often work somewhat longer hours, and maintain their condition better, because the efforts on all hands, including the pace, are less severe. Notwithstanding some of these doing not more, on the average, than twelve miles per day, their duration of service extends only to four years, spare horses and invalids



amounting to 7 per cent. On the other hand, we know of tramway horses doing fifteen or sixteen miles daily, averaging in pace six miles an hour on indifferent roads. This we consider severe and expensive. One company imposed eighteen miles, and, when underhorsed, put on an extra journey of six miles; but the false economy soon made itself evident in the shape of lameness and loss of condition.

In overworked establishments the number of idle or really incapacitated horses is always great, and proportionate to the extent of undue labour imposed. The weekly returns of a certain company showed that 20 per cent. of the whole number, consisting chiefly of lameness, with loss of condition, arose from this cause. Reports sent to the board of directors failed to draw attention to the defect, and the results became serious. It was then discovered that he who had imposed the work had also altered the figures in the returns, showing a false number of workable animals, and, to meet the case, he ordered the sound animals extra journeys. Overwork causes severe depression, which no amount of food, however excellent in quality, can overcome. This has already been stated in various ways, and cannot be insisted upon too strongly.

With regard to spare or relief horses, the number required must very much depend upon the kind and extent of the work. We have known excellent work to be accomplished by



having one extra animal for every twelve; thus the weekly expense of each has not been raised more than one shilling. With more severe work, one to eight may be required. By these means every animal may obtain an uninterrupted rest of twenty-four hours—a proceeding which will tell on the right side in course of years.

Under present circumstances the rest allowed to many studs is obtained by a species of smuggling or “ringing the changes,” somewhat after the following manner. We will suppose eight horses are set apart to work one conveyance, four journeys being made daily throughout seven days. The first pair going out on Monday morning takes the last journey on Tuesday evening, and possibly the first or second on Wednesday, so that another pair obtains a rest of similar duration as the first. All, however, do their allotted work in the week, and the rest exists only in name, not from work. Mr. Hunting states that in one colliery he found, as a result of overwork, one-sixth of the animals were incapacitated. In another, where system was instituted, the same state of inefficiency having previously existed, spare animals amounted only to one-fiftieth of the whole number.\*

To compare the duration of life as well as the nature of work of pit animals with those above

---

\* Paper “On the Feeding and Management of Draught Horses,” read before the Newcastle Farmers’ Club, 1874.



ground may seem irrelevant. It is not, however, a useless comparison. The work is of the hardest kind, and often extends to ten or twelve hours; yet some of these creatures, as a result of good management, remain useful to the age of ten or fifteen years. As a rule, the duration of life, which is largely influenced by fatal accidents, extends to an average of over eight years, where a proper system of feeding and general management is pursued. With all these disadvantages, the pit animal is more highly favoured in having an atmosphere of fairly uniform temperature. Other conditions being equal, this is great gain and safety to his constitution, while his neighbours above ground suffer the vicissitudes of a changeable and often severe climate. Notwithstanding, the estimate of four years as the average duration of life seems to us to be very low, and one which a proper system of food and work should speedily extend at least one-half, and possibly three-fourths, as already shown in the case of the horses owned by the Bristol Tramway Company.

#### IMPORTANCE OF GROOMING.

There is one particular in the management of horses which so materially influences the effects of food upon the system that it will not admit of being unnoticed, although the attention be exceedingly brief. This is *grooming*.



How far this principle is carried with many of our town, dray, and cab horses can be clearly ascertained by merely passing within a few yards as they stand in the street. After they have spent hours in the open air, the odour of the stable is yet strongly upon them.

There are, doubtless, good reasons to be assigned for this in some instances, the most common being that of absolute laziness. There are, however, cases where a groom or horsekeeper is expected to execute multifarious duties in addition to those of the stable, and in consequence the horse is neglected. Another fruitful source of neglect is to be found in low wages, and the men endeavour to make up the deficiency by looking after percentages.

Therefore it proves to his interest to be careless and incur expense upon all sides, which is done to an alarming pitch in some instances. A third cause is the unreasonable number of horses which are placed under the care of horsekeepers. This is a most monstrous practice in some of the coal-mines. I have found it to occur invariably, that where the animals are working in the greatest amount of heat and dust—two or even three miles from the bottom of the shaft—where they perspire most freely, and endure every vicissitude inimical to general health, there they have had the least attention.

All this arose from an excess of duties.

In such cases I found men having the care of



too many animals, to which they could not in any sense render needful attention.

One horsekeeper had under his care fifteen horses and nineteen ponies; a second, four horses and twenty-four ponies; a third, twenty-three ponies; a fourth, two horses and thirty-eight ponies; a fifth, nine horses and twenty-eight ponies.

Three men had the care of five horses and eighty-nine ponies in a sixth instance, and three others had the care of twenty-two horses and fifty-one ponies.

These animals are to be fed, harnessed, and cleaned to go out to work at 5 A.M. by the men, who descend several hours previously in order to attempt an impossibility.

If we take the mean of these numbers, we shall find that each man had an average of thirty animals to attend to. The arrangements of the stables are not always suited towards reducing the labours of the men, being frequently in a continuous line of stalls arranged on one side, which necessitates several journeys to each animal for turning to water, tying-up, supplying with corn and hay when they cannot be fed at the head. Estimating the trouble which these men have, and the risk they run for a few shillings a week, one cannot be surprised that the horses and ponies do not receive their proper share of attention.

The idea of cleaning them is estimated as



the boy did the operation of washing his hands and face, when he refused on the score of their becoming as bad again shortly. Such an estimate, however, does not render the fact as it exists less flagrant.

Let each man be limited to a specified time for the execution of the various details expected from him, and it will be more apparent how absurd must be the belief that these animals can obtain one-fourth the necessary attention under their peculiar and extreme conditions.

Feeding with corn, each one minute	=	30	minutes.
"      hay,      "	=	30	"
Harnessing, with repairs, two minutes	=	60	"
Turning to water and tying-up, one minute	=	30	"
Brushing down, each five minutes	=	150	"
		<hr/>	
Which occupies for thirty animals		800	"
or five hours in all.			

The men descend at 2 A.M., which only leaves *three hours* to perform a number of duties, the time for which, estimated as low as possible, is not sufficient to allow of their being effected in any other than a most slovenly manner—really occupies two hours more than the men can actually bestow on them. At night they are merely stripped, watered, and fed, in order to allow them rest.

To occupy more time with them would be



also prejudicial, as the hours of rest would be materially interfered with ; and consequently an average of thirty animals, estimated at about £300, are thought no more of than to value their care and management at the rate of fifteen shillings a week, or sixpence per head—the amount paid to the horsekeepers.

Here, where the truth of the old maxim, “A good cleaning is equal to a feed of corn,” would be faithfully realised, it cannot be carried out.

When complaints were made, and exact conditions represented, all was pooh-pooh'd, and any alterations, it was said, would cost the estate £300 per annum, as ascertained by figures, which, emanating from the executive, were reliable, but worth nothing when they arose from the mental calculations of one who was not a “Viewer.”

Such is the connection which exists between the skin and digestive organs, that if these poor creatures could receive a guaranteed dressing of fifteen minutes only, each night and morning, the effects would be marvellous.

In a solitary instance, one man had fourteen animals under his care. As he was infirm and contented himself solely with the employment as horsekeeper, he could devote more attention to them than was possible in the other pits.

Every morning and night each was well dressed with a coarse brush, and left in a much more comfortable state.



His horses and ponies would have caused many owners of animals above ground to blush with shame at their superior condition and clean shining skins. They were always doing equal work with the animals of other collieries, but *suffered the least of all, and consumed the least corn.*

I have preferred to recite these facts rather than dictate a philosophical explanation of the uses of a brush and comb to the body of the horse. Those who are so pertinaciously obstinate as to deny him the influence of this luxury, I am afraid are not in a condition to explain the benefits of such an application from any experience of the adoption of similar means to themselves. However, these are considered to be measures not truly essential, and therefore, while the attempt is on the one hand to save, as these *pseudo savants* suppose, the sum of £300 per annum, they have not the philosophy to see, or the honesty to allow another to show, that four times that amount could be saved in the feeding in twelve months, and as much more in a variety of ways of which they cannot form any conception.

For further details on this subject, see "The Horseowner and Stableman's Companion." \*

---

\* London : F. Warne & Co. Price 1s.



# INDEX.

---

**ABILITY**, 48  
**Absorption**, 24  
**Acids of digestion**, 29  
**Advantages of proper food and system**, 42, 46  
**Albumin**, 30  
**Albuminous principles of food**, 28-30  
**Albuminuria**, 45  
**Animal heat**, 32  
     not maintained by fat alone, 35  
     produced in part from nitrogenous compounds, 32  
**Animals, hybernating**, 35  
**Assimilation**, 24

**BRAN and pea straw**, 101  
**Bran: its nature, uses, and abuses**, 83  
     as a laxative, 84  
**Breeding or pluck**, 48  
**Broken wind, how caused**, 21  
**Bruised corn, economy of**, 99  
**Boussingault and Papin, experiments of**, 60  
**Bulk or volume**, 39

**CALCULI or stones in the intestines**, 51  
     their origin, 51, 52  
     usual composition of, 52  
     presented by Mr. Foreman, 54

**Capacity of large intestines**, 26  
     small ditto, 24  
**Care required in feeding after work**, 60  
**Carrots and turnips**, 98  
**Caseine**, 30  
**Chaff, economy of**, 99  
**Change of grain, objections to**, 82  
**Changes in chyme**, 27  
**Cheap food, what constitutes a**, 81  
**Cheeks**, 14  
**Choking, causes of**, 19  
**Chronic cough, causes of**, 22  
**Chyle**, 24  
**Chyme**, 27  
     changes in, 27  
**Coal mines, feeding of horses and ponies in**, 2, 79, 85  
     neglect of horses and ponies in, 77, 105  
**Cæcum**, 25  
     principally contains fluid, 23  
**Colon**, 25  
**Colliery estates, systems of feeding upon**, 74, 76, 79, 85  
**Condiments, objections to**, 103  
**Consequences of imperfect feeding**, 2, 75, 76  
**Cooked food**, 18, 42, 101  
     disease arising from, 45, 101  
     not economical, 55  
     not more digestible or nutritious, 41



- Cooked food produces liability to disease, 45, 101  
 Corn, mixture of, 89  
 Cost of feeding horses in Sheffield, 72  
     feeding on oats, 65  
 Cough, chronic, how caused, 22  
 Cut food promotes insalivation, 75  
  
**DEVELOPMENT** and maintenance, 8  
 Deglutition, 17, 19  
 Diabetes, 13, 42  
 Digestion, organs of, 14, 108  
     should be in healthy condition, 60  
     acids of, 29  
     in intestines, 27  
     in stomach, 22  
     rapid in the horse, 22, 56  
 Digestive process, 26  
 Diseases arising from cooked food, 45, 101  
 Do horses masticate the whole of the grain? 56  
 Dry food, objections to, 54  
 Dung or fæces, 29  
 Duodenum, 24  
  
**ECONOMY** of food, 71  
     using chaff and bruised corn, 99  
 Effete or useless matter, 11, 29  
 Elementary principles of food, 28  
 Errors to be avoided, 40  
 Essential characters of food, 29  
     qualities of oats, 65  
 Evil effects of insufficient food, 12  
 Experiments of Boussingault and Papin, 60  
  
**FÆCES** or dung, 29  
 Fallacious ideas as to passage of unchanged grain, 56  
 Fat or heat producers, 28, 32  
 Feeding in coal-mines, 7, 76, 79, 85  
     regular, 23, 46, 50  
  
 Feeding, saving effected by Mr. Hunting, 74  
     saving effected by Mr. Scott, 80  
     system of, at Hetton Colliery, 76  
     system of, at Londonderry Collieries, 85  
     the manger system, 55  
 Fibrine, 30, 31  
 Flesh-formers, 28  
 Fluids, passage of, 23  
 Food, advantages of good, 12, 42, 46  
     economy of, 71  
     elementary principles of, 28, 29  
     essential characters of, 29  
 Food, green, 97  
     immediate object of, 8  
     indigestible parts of, 29  
     insufficient, effects of, 12  
     relative proportion of nutritious and starchy matters in, 63  
     requires bulk, 39  
     steamed, objectionable, 87  
     varieties of, 63  
     when cooked, injurious, &c., 18, 42, 55, 101  
 Foreman, Mr. Thos., furnishes calculi, 54  
 Forms of grain mixtures, 89  
  
**GASTRIC** juice, 26  
     digestion, 26  
 Gluten, 30, 31  
 Grain, does it pass unchanged? 57  
     economy of storing, 68  
     objections to a change of, 62  
     selection and purchase of, 68  
 Green food, 97  
 Grinders, 15  
 Grooming, importance of, 123  
 Gullet, 15  
  
**HARD** work, care required in feeding after, 23, 60  
 Healthy condition of digestive organs necessary, 60



Heat, animal, 32  
 producers, 28  
 relative proportion of, in food, 63  
 Heavy draught horses, 49  
 Hetton Colliery system of feeding, 76  
 Horse, digestive organs and cooked food, 41  
 Hours of work of pit animals, 77  
 Hunting, C., on the proportion of husk in grain, 67  
 his saving in feeding, 74  
 Husks in grain, proportion of, 67  
 Hybernating animals, 35  
 IDENTITY of nitrogenous compounds, 31  
 Ilium, 24  
 Ill effects of maize, 70  
 Immediate objects of food, 8  
 Importance of grooming, 123  
 prevention, 4, 46  
 Incisor teeth, 14  
 Indian corn or maize, 70  
 Indigestible parts of food, 29  
 Injurious effects of cooked food, 18, 45, 87, 101  
 Insalivation, 16  
 Insufficient food ill effects of, 12  
 Intestines, capacity of, 26  
 division of, 23  
 stones or calculi in, 51  
 JEJUNUM, 24  
 LACTEALS or absorbents, 24  
 Large intestines, 25  
 number of horses, importance of a special superintendent for, 61  
 Laxative, bran as a, 84  
 Linseed, 71  
 Londonderry Collieries, system of feeding at, 85  
 Long fasts prejudicial, 23, 78  
 Longevity promoted by good food, 46

MAINTENANCE, 10  
 Maize or Indian corn, 70  
 Manger system, 55  
 Mastication, 15  
 Mesenteric glands, 28  
 Metamorphosis of tissue, 10, 11  
 Mismanagement, 1  
 Mixtures of grain for horses, 89  
 Molar teeth, 15  
 Mortality from mismanagement, 2-7  
 avoided by Mr. Hunting's system, 75  
 avoided by Mr. Scott's system, 81  
 NATURE, &c., of bran, 83  
 Neglect of grooming, results of, 123  
 Nitrogenous compounds, 30  
 principles of food, 30, 63  
 Non-nitrogenous principles of food, 32  
 relative proportions in food, 63  
 OATS, 64  
 cost of feeding with, 65  
 essential qualities of, 65  
 straw, 99  
 Objections to a change of grain, 83  
 to green food, 97  
 to the use of dry food, 54  
 Œsophagus, 15  
 Organs of digestion, 14  
 Origin of calculi, 52  
 PEA and bean straw, 101  
 Pharynx, 15  
 Pit animals, neglect of, 77  
 their hours of work, 77  
 horsekeepers, excessive duties of, 124  
 Pluck or breeding, 48  
 Practical tests of the system, 104  
 Prehension, 14  
 Prejudice against system on colliery estates, 77, 87, 127  
 Prevention, importance of, 4, 46  
 Process of digestion, 26



- Profuse staling, 42  
 Proportion of husk in grain, 67  
 Provision for maintaining  
     warmth in the body, 35  
 Ptyalin, 16  
 RAPIDITY of digestion, 22  
 Rectum, 25  
 Regular feeding, 23, 46, 50  
 Relative proportion of heat pro-  
     ducers in food, 63  
     of nutritious matter in food,  
     31, 63  
 SACCHARINE principles of food,  
     17, 28, 32, 63  
 Saliva, 16  
     cannot be replaced, 18  
     chemical action of, 17  
     flow of, promoted by cut  
     food, 75, 99  
     is secreted abundantly, 17  
     solvent action of, 16  
     quantity of, important in di-  
     gestion, 17, 99  
     uses of, 17, 18  
 Saving effected by Mr. Hunt-  
     ing, 74  
     by systematic feeding, 102  
 Scott, Mr. L., his system and  
     saving by feeding, 79  
 Selection and purchase of grain,  
     68  
 Sheffield, cost of feeding at, 72  
 Small intestines, 24  
 Spicy foods or condiments, 103  
 Staling, profuse, 42  
 Steamed food objectionable, 86  
 Stomach of horse, ox, and man,  
     20  
     of horse, small, 21  
     of ox, divided into four parts,  
     20  
 Stones in the intestines, 51  
     their origin, 52  
 Storing of grain, 68  
 Straw of beans, 101  
 Straw of oats, 99  
     peas, 101  
     wheat, 99  
 Strength or ability derived from  
     food, 48  
     required for work, 49  
 Superintendent, importance of  
     a, 61  
 Swallowing, 17, 19  
 System of feeding on collieries,  
     74, 85  
     of saving to be effected by a,  
     102  
     practical tests of the, 104  
 TARES, 71  
 Teeth, incisor, 14  
     molar, 15  
 Tissue, metamorphosis of, 10,  
     11  
 Tongue, 14  
 Turnips and carrots, 98  
 UNDIGESTED food, 29  
     grain, passage of, 57  
 Urination, profuse, 42  
 Useless or effete matters, 11  
 Use and abuse of bran, 83  
     of dry food, objections to the,  
     54  
     of saccharine principles of  
     food, 17, 28, 32  
     of saliva, 17  
 VARIETIES of food, 63  
 Various mixtures of grain, 89  
 WANT of condition, effects of a,  
     13, 97  
 Waste of animal tissues, 10  
 Wasting of the body, causes of,  
     34  
 What constitutes a cheap food,  
     81  
 Work, 116  
 YOUNG animals require good  
     food, 12







*Price 1s. each.*WARNE'S USEFUL BOOKS—*continued.*

## LETTER-WRITERS.

**LADIES' AND GENTLEMEN'S MODEL LETTER WRITER:** A Complete Guide to Correspondence on all Subjects, with Household and Commercial Forms.

"Every possible subject appears to have been dealt with, and a table of contents serves to bring those who are in difficulties to the object of their search."—

*Army and Navy Gazette.*

**THE COMPANION LETTER WRITER:** A Guide to Correspondence on all Subjects relating to Friendship, Love and Business, with Commercial Forms, &c.

"This volume has been prepared with great care, and will be found most useful to persons requiring such assistance."

## ATHLETIC, SCIENTIFIC AND HOUSEHOLD MANUALS.

**THE MODERN GYMNAST.** Practical Instructions on the Horizontal Bar, Parallel Bars, Flying Trapeze, &c. &c. With One Hundred and Twenty Practical Illustrations. By CHARLES SPENCER.

"The directions given in this work are short and simple; the book is well worth having, and we recommend it to all who may affect gymnastic exercises."—*The Field.*

"By means of one hundred and twenty illustrations it sets forth *advanced* gymnastics in a very practical manner."—*Builder.*

**THE MODERN FENCER.** With the most Recent Means of Attack and Defence when Engaged with an Adversary. With Practical Instructions. By Captain T. GRIFFITHS.

"The author's aim has been to convey a practical knowledge of fencing in the simplest manner possible, and in the shortest time."

**MAGIC AND ITS MYSTERIES.** Explained by Professor J. D. THEOBALD.

"By following out with care the hints and instructions given in these pages, even those most ignorant of conjuring may, in a few evenings, commence to delight their friends and acquaintances with simple and effective amusements."

**THE MAGIC LANTERN MANUAL.** By W. J. CHADWICK, Hon. Sec. Manchester Photographic Society. Illustrated by 105 Practical Woodcuts.

"Although forming a book of instruction to the learner, it will take its place as a handbook to the expert lanternist."

"Will be very useful to all teachers who wish to make themselves master of the art of managing a magic lantern."—*National Schoolmaster.*

**VENTRILOQUISM (THE ART OF).** With Full Directions for Learners. How to Acquire a Pleasing Vocalisation; with Amusing Dialogues. By FREDERICK MACCABE.

"That master of the art, Frederick Maccabe, clearly explains what ventriloquism is, describes how the art might be acquired, gives several funny dialogues for beginners, &c."—*Plymouth Mercury.*

**ONE THOUSAND OBJECTS FOR THE MICROSCOPE.** By M. C. COOKE. With 500 Figures Illustrated. Twelve pages of Tinted Plates.

"A valuable and entertaining companion."—*Sunday Times.*

"The thousand objects are all clearly described."—*Public Opinion.*

**WASHING AND CLEANING:** A Manual for Domestic Use. By BESSIE TREMAINE.

"This little manual will be found of great service in setting up washing at home, and also useful in that it contains many hints as to the various kinds of washing, starching, and getting up lace, linen, &c."—*Bazaar.*



*Price 1s. each.*WARNE'S USEFUL BOOKS—*continued.*

**CARPENTRY AND JOINERY:** A Useful Manual for the Many. By S. T. AVELING. With numerous Original and Practical Illustrations, and Full Instructions as to Choice of Materials, Tools, and how to make a variety of Useful Articles.

"This is an excellent little handbook on a most important branch of industry, and no apprentice who wishes to excel as a workman should be without a copy. The illustrations are practical, showing the right way of doing a thing."—*Weekly Times.*

**THE MONEY MARKET.** What it Is, What it Does, and How it is Managed. An Introduction to Financial Science.

"It imparts a good deal of information on such themes as banking, exchange, the stock and share markets, discount, &c."—*Nonconformist.*

**HOW TO SEND A BOY TO SEA.** By FRANKLIN FOX, Captain P. and O. Service.

"'How to Send a Boy to Sea,' by Franklin Fox, will be of benefit to the parents of the legion of youths aspiring to a nautical career, as it satisfactorily explains the branches and advantages of the mercantile marine service."—*Lloyd's.*

"His advice is sound and practical. There is a great deal of information in his book which cannot fail to be of use to those who wish to send a boy to sea, but have a limited acquaintance with naval matters."—*Broad Arrow.*

*Price 1s. each.***WARNE'S LEGAL HANDBOOKS.**

CLEAR, CONCISE AND TRUSTWORTHY.

By CHARLES E. BAKER, Solicitor of the Supreme Court.

*In small crown 8vo, cloth.*

**THE LOCAL GOVERNMENT ACT, 1888.** With Introduction and Notes.

**WILLS:** How to Make and How to Prove Them, and the Law of Administration. With nearly 150 Forms and Index. Fourth Edition. Revised.

**LANDLORDS, TENANTS AND LODGERS.** Embodying the Law of Distress Amendment Act, 1888, and numerous Forms and Index. Fourth Edition, Revised.

**HUSBAND AND WIFE,** and the Married Woman's Property Act. New Edition.

**THE LAW OF MASTER AND SERVANT.** With Appendix and Index.

*Also Uniform with the above by J. RENWICK SEAGER.*

**THE REPRESENTATION OF THE PEOPLE ACT, 1884,** and the Redistribution Act, 1885. With Introduction, Appendix of Forms, and full Index. Third Edition.

**THE MUNICIPAL ELECTIONS ACT, 1884.** With Introduction and Full Index.



