#### The pharmacopoeia of the Royal London Ophthalmic Hospital.

#### **Contributors**

Royal London Ophthalmic Hospital.

#### **Publication/Creation**

[London]: [n.p.], [1882] [(London]: [Spottiswoode.])

#### **Persistent URL**

https://wellcomecollection.org/works/gqj5p89y

#### License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



# PHARMACOPEIA OF THE ROYAL LONDON OPHTHALMIC HOSPITAL

K50271



# Med K50271

PHARMACOPŒIA.

LONDON: PRINTED BY

SPOTTISWOODE AND CO., NEW-STREET SQUARE

AND PARLIAMENT STREET

#### THE

# PHARMACOPŒIA

OF THE

ROYAL LONDON

# OPHTHALMIC HOSPITAL.

SEVENTH EDITION.

LONDON: 1882.

17 694 +88

WELLCOME INSTITUTE LIBRARY		
Coll.	WelMOmec	
Coll.		
No.	WW	

Consulting Physician.
FREDERICK JOHN FARRE, M.D.

Physician.

ROBERT MARTIN, M.D.

Consulting Surgeons.

JAMES DIXON.

CHORGE CRITCHITT.

WILLIAM BOWMAN, F.R.S. JONATHAN HUTCHINSON.

Surgeons.

JOHN CAWOOD WORDSWORTH.

JOHN FREMLYN STREATFFILD.

JOHN WHITAKER HULKE, F.R.S.
GEORGE LAWSON.

JOHN COUPER.

Assistant Surgeons.

WARREN TAY.

JAMES EDWARD ADAMS.

JOHN TWEEDY.

ROBERT LYELL.

LOUIS PRESENT.

# PHARMACOPŒIA.

CAUSTICA — CAUSTICS.

# CAUSTICUM ZINCI CHLORIDI.

(MIDDLESEX HOSPITAL.)

# Take of

Chloride of Zin	nc		480 grains
Wheat Flour			180 grains
Water .	KNE IST	Us 18	1 fluid ounce

Mix, and apply the heat of a water bath for a few minutes only, until a paste is formed.

Or-

Take of

Chloride of Zinc . . . } equal parts

Mix thoroughly with pestle and mortar, and preserve in a stoppered bottle.

#### CONFECTIO \_\_\_\_ CONFECTION.

#### CONFECTIO SULPHURIS SIMPLEX.

#### Take of

Sublimed Sulphur . . 4 ounces
Acid Tartrate of Potash . 1 ounce
Treacle, by weight . . 4 ounces

Rub well together.

Dose, one or two teaspoonfuls for an adult.

#### DECOCTA - DECOCTIONS.

#### DECOCTUM PAPAVERIS.

# Take of

Poppy Capsules, bruised (without seeds) . . 1 ounce

Distilled Water . . 1 pint

Boil for a quarter of an hour in a covered vessel, and strain.

#### DECOCTUM QUERCUS.

# Take of

Oak Bark, bruised . . 1 ounce Distilled Water . . 1 pint

Boil for a quarter of an hour in a covered vessel, and strain.

#### FOMENTUM \_\_\_ FOMENTATION.

#### FOMENTUM BELLADONNÆ.

#### Take of

Extract of Belladonna . 60 grains Distilled Water . . . 1 pint

Rub together till dissolved.

To be made hot and used as a fomentation.

#### GARGARISMATA — GARGLES.

# GARGARISMA ACIDI CARBOLICI.

# Take of

Glycerine of Carbolic Acid

B.P. . .  $2\frac{1}{2}$  fluid drachms

Glycerine . . . . 3 fluid drachms

Water . . . up to 20 fluid ounces

#### Mix.

#### GARGARISMA ACIDI TANNICI.

# Take of

Diluted Hydrochloric Acid 2 fluid drachms Glycerine of Tannin B.P. . 3 fluid drachms Water . . up to 20 fluid ounces

#### GARGARISMA ALUMINIS.

# Take of

Powdered Alum . . . 180 grains . . 1 fluid ounce Glycerine .

Distilled Water . up to 20 fluid ounces

Dissolve and mix.

# GARGARISMA HYDRARGYRI PERCHLORIDI.

# Take of

Perchloride of Mercury . 2 grains

Tincture of Cochineal . 2 fluid drachms

Water . . up to 20 fluid ounces

Dissolve and mix.

#### GARGARISMA POTASSÆ CHLORATIS.

# Take of

Chlorate of Potash . . . 90 grains
Hydrochloric Acid . . . 40 minims

Glycerine . . . 1 fluid ounce

Distilled Water . . up to 20 fluid ounces

Mix the acid and chlorate in a bottle. Cork and let it stand for three minutes. Then add the water gradually with much shaking. Lastly add the glycerine.

#### GARGARISMA SODÆ CHLORATÆ.

Take of

Solution of Chlorinated Soda . . . . 1 ounce Distilled Water . . . 19 ounces

Mix.

GLYCERINA ---- GLYCERINES.

#### GLYCERINUM ACIDI TANNICI P.B.

Take of

Tannic Acid . . . 1 ounce
Glycerine . . 4 fluid ou . 4 fluid ounces Glycerine

Dissolve with gentle heat.

#### GLYCERINUM BORACIS P.B.

Take of

Borax, in powder . . 1 ounce Glycerine . . 4 fluid ounces

Rub together till dissolved.

#### GLYCERINUM BELLADONNÆ.

# Take of

Extract of Belladonna	. 1 ounce
Distilled Water	. 4 fluid drachms
Glycerine	. 3½ fluid ounces

Mix the extract with the water, add the glycerine, and rub together till dissolved.

#### GLYCERINUM PLUMBI SUBACETATIS.

# Take of

Acetate of Lead .	5 ounces
Oxide of Lead, in powder	$3\frac{1}{2}$ ounces
Glycerine	20 fluid ounces
Distilled Water	12 fluid ounces

Mix together and boil for a quarter of an hour; then filter and evaporate to one pint.

#### GUTTÆ \_\_\_ DROPS.

# GUTTÆ ARGENTI NITRATIS.

# Take of

Nitrate of Silver		1 grain
Distilled Water		1 fluid ounce

#### GUTTÆ ARGENTI NITRATIS FORTIORES.

Take of

Nitrate of Silver · . 2 grains

Distilled Water . . 1 fluid ounce

Dissolve.

#### GUTTÆ ATROPIÆ SULPHATIS.

Take of

Sulphate of Atropia . . 1 grain

Distilled Water . . 1 fluid ounce

Dissolve.

#### GUTTÆ ATROPIÆ SULPHATIS FORTIORES.

Take of

Sulphate of Atropia . . 2 grains

Distilled Water . . 1 fluid ounce

Dissolve.

# GUTTÆ ATROPIÆ CUM ALUMINE.

Take of

. 1 grain Sulphate of Atropia

Alum . . . . 3 grains
Distilled Water . . . 1 fluid o

. 1 fluid ounce

# GUTTÆ ATROPIÆ CUM ZINCO.

# Take of

Sulphate of Atropia . . 1 grain

Sulphate of Zinc . . 1 grain
Distilled Water . . 1 fluid ounce

Dissolve.

#### GUTTÆ CUPRI SULPHATIS.

# Take of

Sulphate of Copper . . 2 grains

Distilled Water . . 1 fluid ounce

Dissolve.

#### GUTTÆ DATURIÆ.

# Take of

Sulphate of Daturine . . 2 grains

Distilled Water . . 1 fluid ounce

Dissolve.

#### GUTTÆ ESERIÆ.

#### Take of

Sulphate of Eserine . . 2 grains

Distilled Water . . 1 fluid ounce

#### GUTTÆ OPII.

# Take of

Wine of Opium . . 2 fluid drachms Distilled Water . . 6 fluid drachms

Mix.

#### GUTTÆ PILOCARPIÆ.

# Take of

Nitrate of Pilocarpine . 2 grains

Distilled Water . . 1 fluid ounce

Dissolve.

#### GUTTÆ POTASSII IODIDI.

# Take of

Iodide of Potassium . . 4 grains

Distilled Water . 1 fluid ounce

Dissolve.

#### GUTTÆ TEREBINTHINÆ.

# Take of

Oil of Turpentine . . 1 fluid drachm Olive Oil . . . 7 fluid drachms

#### GUTTÆ ZINCI CHLORIDI.

Take of

Chloride of Zinc . . 1 grain

Distilled Water . . 1 fluid ounce

Dissolve.

#### GUTTÆ ZINCI SULPHATIS.

Take of

Sulphate of Zinc . . 2 grains

Distilled Water . . 1 fluid ounce

Dissolve.

#### GUTTÆ ZINCI SULPHATIS ET ALUMINIS.

Take of

Sulphate of Zinc . . 1 grain

Alum . . . . 3 grains

Distilled Water . . 1 fluid ounce

Dissolve.

# INJECTIONES \_\_\_\_ INJECTIONS.

#### INJECTIO ALKALINA.

Take of

Carbonate of Potash · 120 grains

Distilled Water . . 1 pint

#### INJECTIO ASTRINGENS.

# Take of

Alum in powder . . 80 grains Sulphate of Zinc . . 80 grains Tannic Acid . . . 30 grains Distilled Water . . 1 pint

Dissolve.

#### INJECTIO PLUMBI.

#### Take of

Diluted Solution of Sub-

acetate of Lead .  $1\frac{1}{2}$  fluid drachm Rectified Spirit . 2 fluid drachms

Distilled Water . up to 1 pint

Mix.

# INJECTIONES HYPODERMICE —— HYPO-DERMIC INJECTIONS.

#### INJECTIO ATROPIÆ SULPHATIS.

# Take of

Sulphate of Atropia . . ½ grain

Distilled Water . . 1 fluid drachm

#### Mik.

Dose, by subcutaneous injection, two to four minims. (Three minims contain th of a grain of the sulphate of atropia.)

# INJECTIO MORPHIÆ HYPODERMICA P.B.

A Solution of Acetate of Morphia containing one grain of acetate in twelve minims of the injection.

Dose, one to six minims.

#### INJECTIO MORPHIÆ ET ATROPIÆ.

#### Take of

Sulphate of Atropia . . 1 grain

Hypodermic Injection of

Morphia B.P. . . 6 fluid drachms

#### Mix.

Dose, one to four minims. Three minims contain  $\frac{1}{4}$ th of a grain of acetate of morphia and  $\frac{1}{120}$ th of sulphate of atropia.

#### LINCTUS.

# Take of

Treacle		2 fluid ounces
Powdered Tragaca	inth	40 grains
Tincture of Opiun	n .	30 minims
Water		2 fluid ounces
Diluted Sulphuric	Acid	 30 minims
Vinegar of Squills	BEILDIG	3 fluid drachms

#### Mix.

Dose, one or two teaspoonfuls.

#### LINIMENTA — LINIMENTS.

#### LINIMENTUM ACONITI ET SAPONIS.

# Take of

Liniment of Aconite B.P 1 part Liniment of Soap . . 3 parts

Mix.

#### LINIMENTUM AMMONIÆ P.B.

#### Take of

Solution of Ammonia . 1 part Olive Oil . . . 3 parts

Mix.

# LINIMENTUM BELLADONNÆ COM-POSITUM.

# Take of

Extract of Belladonna . 1 fluid drachm Distilled Water . . 1 fluid drachm Liniment of Soap . . 6 fluid drachms

Mix.

#### LINIMENTUM CROTONIS

# Take of

Croton Oil . . . 1 fluid drachm
Oil of Turpentine . . 3 fluid drachms
Olive Oil . . . 4 fluid drachms

#### LINIMENTUM HYDRARGYRI OLEATIS

(10 per cent.)

(UNIVERSITY COLLEGE HOSPITAL.)

#### Take of

Yellow Oxide of Mercury 1 drachm Oleic Acid . . . . 10 fluid drachms

To the oleic acid, kept agitated in a mortar, sprinkle in the oxide of mercury gradually; triturate the mixture frequently during twenty-four hours until the oxide is dissolved and a viscid solution is formed.

# LINIMENTUM HYDRARGYRI OLEATIS CUM MORPHIA.

(University College Hospital.)

#### Take of

Morphia (pure alkaloid) . 10 grains
Oleic Acid . . . 5 fluid drachms

# Dissolve and add

Liniment of Oleate of Mercury (10 per cent.) . 5 drachms

# LIQUORES \_\_\_\_ SOLUTIONS.

# LIQUOR ARGENTI NITRATIS

# Take of

Nitrate of Silver . 5, 10, 15, or 20 grains 1 fluid ounce Distilled Water

# Dissolve.

N.B.—These preparations are intended only for application to the surface of the conjunctiva of the lids with a camel's-hair brush, the excess to be afterwards neutralised with a solution of salt and water five grains to the ounce.

# LIQUOR ATROPIÆ SULPHATIS P.B.

# Take of

Sulphate of Atropia . . 4 grains Distilled Water . 1 fluid ounce

#### LIQUOR EPISPASTICUS.

# Take of

Cantharides, in powder . 8 ounces
Glacial Acetic Acid . 2 fluid ounces
Ether . . . a sufficiency

Mix the cantharides with the acid and two fluid ounces of ether; pack in a percolator, and after twenty-four hours pass ether slowly through until twenty fluid ounces are obtained. Keep in a stoppered bottle.

# LOTIONES \_\_\_ LOTIONS.

#### LOTIO ACETI.

#### Take of

Diluted Acetic Acid . . 30 minims
Distilled Water . up to 1 fluid ounce
Mix.

#### LOTIO ACIDI HYDROCYANICI.

# Take of

Diluted Hydrocyanic Acid 5 minims
Distilled Water . . 1 fluid ounce

# LOTIO ACIDI HYDROCYANICI CUM SODA.

# Take of

Bicarbonate of Soda. . 5 grains Borax . . . 5 grains Diluted Hydrocyanic Acid 5 minims Distilled Water . . 1 fluid ounce

Dissolve and mix.

#### LOTIO ALUMINIS.

# Take of

Alum, in powder . . 6 grains Distilled Water . . 1 fluid ounce

Dissolve.

#### LOTIO ALUMINIS MITIS.

# Take of

Alum, in powder . . 3 grains . 1 fluid ounce Distilled Water

Dissolve.

#### LOTIO ALUMINIS COMPOSITA.

# Take of

Alum, in powder . . 3 grains Sulphate of Zinc . . 1 grain

. . 1 fluid ounce Distilled Water

#### LOTIO ARNICÆ.

# Take of

Tincture of Arnica . . 20 minims

Distilled Water . . 1 fluid ounce

Mix.

#### LOTIO BELLADONNÆ CUM ALUMINE.

# Take of

Extract of Belladonna . 5 grains

Alum, in powder . . 3 grains

Distilled Water . . 1 fluid ounce

Dissolve.

#### LOTIO BORACIS.

#### Take of

Glycerine of Borax . . 1 fluid drachm

Distilled Water . . 1 fluid ounce

Dissolve.

#### LOTIO FERRI SULPHATIS.

# Take of

Sulphate of Iron, in powder 5 grains
Distilled Water . . 1 fluid ounce

#### LOTIO HYDRARGYRI PERCHLORIDI.

# Take of

Perchloride of Mercury . 1/8 grain Distilled Water . . 1 fluid ounce

Dissolve.

# LOTIO PLUMBI ACETATIS.

# Take of

Acetate of Lead, in powder 2 grains Diluted Acetic Acid . . 2 minims Distilled Water . . 1 fluid ounce

Dissolve and mix.

#### LOTIO PLUMBI FORTIS.

# Take of

Solution of Subacetate of Distilled Water . up to 1 fluid ounce

#### LOTIO POTASSÆ CHLORATIS.

# Take of

Chlorate of Potash . . 5 grains
Diluted Hydrochloric Acid 2 minims
Distilled Water . . 1 fluid ounce

Dissolve and mix.

#### LOTIO RUBRA.

# Take of

Sulphate of Zinc . . 1 grain
Compound Tincture of
Lavender . . . 15 minims
Spirit of Rosemary . . . 15 minims
Distilled Water . up to 1 fluid ounce

Dissolve and mix.

# LOTIO ZINCI SULPHATIS.

# Take of

Sulphate of Zinc . . 1 grain

Distilled Water . . 1 fluid ounce

#### MISTURÆ --- MIXTURES.

#### MISTURA ACIDI HYDROCHLORICI.

# Take of

Diluted Nitro-hydrochloric

. . . . 10 minims Acid

Tincture of Chiretta . . 15 minims

Water , . . up to 1 fluid ounce

Mix.

# MISTURA ACIDI SULPHURICI CUM MAGNESIA.

# Take of

Sulphate of Magnesia . 60 grains
Diluted Sulphuric Acid . 10 minims

Glycerine . . . 30 minims

Peppermint Water . up to 1 fluid ounce

Mix.

#### MISTURA ÆTHERIS COMPOSITA.

# Take of

. 30 minims Spirit of Ether.

Aromatic Spirit of Am-

monia . . . 30 minims
Syrup of Tolu . . 1 fluid drachm
Distilled Water . . 1 fluid ounce

<sup>&</sup>lt;sup>1</sup> The ordinary dose of all the mixtures is one ounce for an adult.

#### MISTURA AMMONIÆ ACETATIS.

# Take of

Solution of Acetate of Am-

monia . . . . 2 fluid drachms

Distilled Water . up to 1 fluid ounce

Mix.

# MISTURA AMMONIÆ ACETATIS CUM MAGNESIA

(VEL MISTURA SALINA CATHARTICA).

#### Take of

Sulphate of Magnesia . 60 grains

Mixture of Acetate of Am-

monia . . . 1 fluid ounce

Mix.

#### MISTURA AMMONIÆ CUM IPECACUANHA.

#### Take of

Carbonate of Ammonia . 3 grains

Wine of Ipecacuanha . 10 minims

Glycerine . . . 30 minims

Distilled Water . up to 1 fluid ounce

Dissolve and mix.

#### MISTURA BISMUTHI.

# Take of

Subnitrate of Bismuth . 10 grains Powdered Tragacanth . 3 grains

Distilled Water . . 1 fluid ounce

Mix.

# MISTURA CHLOROFORMI CUM AMMONIA.

# Take of

Carbonate of Ammonia . 3 grains Spirit of Chloroform . 15 minims Infusion of Gentian . up to 1 fluid ounce

Dissolve and mix.

# MISTURA COLCHICI.

# Take of

Wine of Colchicum . . 20 minims Carbonate of Magnesia . 10 grains Pimento Water . up to 1 fluid ounce

Mix.

#### MISTURA COPAIBÆ.

# Take of

Balsam of Copaiva . . 20 minims Solution of Potash . . 10 minims Pimento Water . up to 1 fluid ounce

#### MISTURA CRETÆ AROMATICA.

# Take of

Prepared Chalk . . 10 grains Compound Cinnamon Pow-

der . . . . 10 grains
Gum Acacia . . . 10 grains

Distilled Water . . 1 fluid ounce

Mix.

#### MISTURA CRETÆ AROMATICA ANODYNA.

# Take of

Aromatic Chalk Mixture . 1 fluid ounce Tincture of Opium . 5 minims

Mix.

#### MISTURA FERRI APERIENS.

# Take of

Sulphate of Iron . . . 1 grain
Sulphate of Magnesia . . 30 grains
Diluted Sulphuric Acid . 5 minims
Distilled Water . . 1 fluid ounce

#### MISTURA FERRI COMPOSITA P.B.

Make according to the instructions given in the British Pharmacopæia, using distilled water in place of rose water.

#### MISTURA FERRI PERCHLORIDI.

# Take of

Tincture of Perchloride of

· . . 10 minims 

Distilled Water . up to 1 fluid ounce

Mix.

#### MISTURA FILICIS.

#### Take of

Liquid Extract of Male Fern 30 minims

Powdered Tragacanth . . 10 grains

Peppermint Water . up to 1 fluid ounce

Rub the liquid extract and the tragacanth together, and then add the peppermint water and mix well.

# MISTURA GENTIANÆ CUM AMMONIA.

# Take of

Carbonate of Ammonia . 4 grains

Bicarbonate of Soda . . 10 grains

Compound Infusion of

. . 1 fluid ounce Gentian . .

Dissolve and mix.

#### MISTURA GENTIANÆ CATHARTICA.

# Take of

Sulphate of Magnesia . 60 grains
Compound Infusion of
Gentian . . . 1 fluid ounce

Dissolve.

#### MISTURA GUAIACI COMPOSITA.

# Take of

Ammoniated Tincture of
Guaiacum . . . 30 minims
Mucilage of Gum Acacia . 1 fluid drachm
Distilled Water . up to 1 fluid ounce

Mix the tincture with the mucilage; then add the water and shake well.

#### MISTURA HYDRARGYRI PERCHLORIDI.

#### Take of

Perchloride of Mercury .  $\frac{1}{16}$  grain Tincture of Cochineal . 10 minims Distilled Water . up to 1 fluid ounce

#### MISTURA HYDRARGYRI IODIDI.

# Take of

Iodide of Potassium . . 3 grains Solution of Perchloride of

Mercury . . . 30 minims
Distilled Water . up to 1 fluid ounce

Mix.

#### MISTURA MAGNESIÆ CUM RHEO.

# Take of

Carbonate of Magnesia . 15 grains Rhubarb, in powder . . 5 grains
Cinnamon Water . . 1 fluid ounce

Mix.

#### MISTURA NUCIS VOMICÆ.

# Take of

Tincture of Nux Vomica . 10 minims

Compound Infusion of

Gentian . . . 1 fluid ounce

## MISTURA POTASSÆ TARTRATIS EFFERVESCENS.

#### Take of

Bicarbonate of Potash . 15 grains

Distilled Water . . 4 fluid drachms

# Dissolve, and add

Tartaric Acid . . . 12 grains

Distilled Water . . 4 fluid drachms

To be taken while effervescing.

#### MISTURA POTASSII BROMIDI.

## Take of

Bromide of Potassium . 10 grains Carbonate of Ammonia . 3 grains

Tincture of Calumba . . 30 minims

Distilled Water . . . 1 fluid ounce

Dissolve.

# MISTURA POTASSII IODIDI.

## Take of

Iodide of Potassium . . 5 grains

Carbonate of Ammonia . 3 grains

Tincture of Calumba . 30 minims

Distilled Water. . up to 1 fluid ounce

Dissolve and mix.

#### MISTURA QUINIÆ.'

# Take of

Sulphate of Quinia . . 1 grain

Dilute Sulphuric Acid . 5 minims

Distilled Water . . 1 fluid ounce

Dissolve.

#### MISTURA QUINIÆ CATHARTICA

# Take of

Sulphate of Magnesia . 60 grains
Quinine Mixture . 1 fluid ounce

Dissolve.

#### MISTURA QUINIÆ CUM FERRO.

# Take of

Sulphate of Iron . . 1 grain

Quinine Mixture . . 1 fluid ounce

# Mix.

<sup>&</sup>lt;sup>1</sup> For out-patients sulphate of cinchonidine and quinidine may be substituted for quinia.

#### MISTURA SCILLÆ COMPOSITA.

## Take of

Tincture of Squills . . 15 minims Spirit of Nitrous Ether . 30 minims

Solution of Acetate of Am-

monia . . . 1 fluid drachm-Distilled Water . . up to 1 fluid ounce

Mix.

#### MISTURA SENNÆ COMPOSITA.

# Take of

Sulphate of Magnesia . 120 grains

Tincture of Senna . . 2 fluid drachms

Compound Tincture of Car-

damoms . . . 30 minims

Infusion of Senna . up to 1 fluid ounce

Dissolve and mix.

### PASTA IODI ET AMYLI.

(University College Hospital.)

# Take of

Starch, in powder . . 1 ounce

Glycerine . . . 2 fluid ounces

. 6 fluid ounces Water .

Boil together, and when nearly cold add

Solution of Iodine B. P. . 1 fluid ounce

Mix well.

#### PILULÆ \_\_\_ PILLS.

#### PILULA ACONITI ET BELLADONNÆ

## Take of

Extract of Aconite . . 1 grain Extract of Belladonna . 1/6 grain . 2 grains Sugar of Milk . . . . . a sufficiency Treacle .

Mix, and make a pill.

#### PILULA APERIENS.

# Take of

Barbadoes Aloes . 1 grain Powdered Scammony Powdered Scammony . 1 grain
Powdered Jalap . 1 grain
Powdered Ginger . 1 grain
Treacle . . a suffici . 1 grain . a sufficiency

Mix, and make a pill.

# PILULA BELLADONNÆ CUM ZINCI SULPHATE.

# Take of

Extract of Belladonna . 1 grain Powdered Sulphate of Zinc 1 grain Sugar of Milk . . . 1 grain Treacle . . . a sufficiency

# PILULA CALOMELANOS CUM COLO-CYNTHIDE.

# Take of

Subchloride of Mercury . 1 grain

Compound Extract of Co-

locynth . . . 3 grains

Extract of Henbane . . 1 grain

Mix, and make a pill.

#### PILULA CALOMELANOS CUM OPIO.

## Take of

Subchloride of Mercury . 1 grain

Powder of Opium . . 4 grain

Wheaten Flour . . . 1 grain

Confection of Roses . . a sufficiency

Mix, and make a pill.

# PILULA CALOMELANOS ET PULVERIS IPE-CACUANHÆ COMPOSITI.

#### Take of

Subchloride of Mercury . 1 grain

Compound Powder of Ipe-

cacuanha . . 4 grains

Mucilage of Acacia . . a sufficiency

# PILULA CANNABIS INDICÆ.

## Take of

Extract of Indian Hemp . ½ grain
Sugar of Milk . . . 2 grains
Treacle . . . a sufficiency

Mix, and make a pill.

## PILULA COLCHICI ET QUINIÆ.

# Take of

Acetic Extract of Colchicum . . . . 1 grain Sulphate of Quinia . . 1 grain

Mix, and make a pill.

# PILULA HYDRARGYRI CUM COLO-CYNTHIDE.

# Take of

Mercurial Pill . . . 2 grains
Compound Pill of Colocynth . . . 3 grains

# PILULA HYDRARGYRI CUM CRETA ET PULVERIS IPECACUANHÆ COMPOSITI.

Take of

Mercury with Chalk . . 2 grains

Compound Powder of Ipe-

cacuanha . . . 3 grains

Mucilage of Acacia . . a sufficiency

Mix, and make a pill.

### PILULA HYDRARGYRI CUM OPIO.

Take of

Mercurial Fill . . . 5 grains

Powder of Opium . . ½ grain

Mix, and make a pill.

#### PILULA HYDRARGYRI CUM RHEO.

Take of

Mercurial Pill . . . 2½ grains

Compound Rhubarb Pill . 21 grains

Mix, and make a pill.

#### PILULA SODÆ ARSENIATIS.

Take of

Arseniate of Soda .  $\frac{1}{16}$  grain

Sugar of Milk, in powder . 1 grain

Extract of Hop . . 3 grains

## PULVERES --- POWDERS.

## PULVIS ALTERATIVUS.

Take of

Mercury with Chalk . . 1 part Bicarbonate of Soda . . 2 parts

Powdered Rhubarb . . 3 parts

Mix.

Dose, 6 to 12 grains.

# PULVIS CALOMELANOS CUM JALAPA.

Take of

Mix.

Dose, 12 grains.

## PULVIS CALOMELANOS CUM RHEO.

Take of

Subchloride of Mercury
Ginger, in powder . . . } each 1 part
Rhubarb, in powder . . 4 parts

Mix.

Dose, 12 grains.

## PULVIS CALOMELANOS CUM SCAMMONIA.

Take of

Subchloride of Mercury
Ginger, in powder . . . } each 1 part
Scammony, in powder . . 4 parts

Mix.

Dose, 12 grains.

#### PULVIS CINCHONÆ ET SODÆ.

Take of

Powdered Cinchona Bark
Bicarbonate of Soda . . . } equal parts

Mix.

Dose, 5 to 10 grains.

# PULVIS HYDRARGYRI CUM CRETA ET BELLADONNÆ.

Take of

Mercury with Chalk . 2 parts

Powdered Belladonna

Leaves . . . 1 part

Powdered Sugar . . 2 parts

Mix.

Dose, 5 grains.

# PULVIS HYDRARGYRI CUM CRETA ET RHEI.

Take of

Mercury with Chalk . . 1 part Rhubarb, in powder . . 2 parts

Mix.

Dose, 3 to 6 grains.

## PULVIS JALAPÆ ET RHEI.

Take of

Jalap, in powder . . 2 parts Rhubarb, in powder . . 1 part Mix.

Dose, 3 to 15 grains.

#### PULVIS RHEI ET MAGNESIÆ.

Take of

Rhubarb, in powder . . 1 part Carbonate of Magnesia . 2 parts Mix.

Dose, 15 to 20 grains.

#### PULVIS RHEI ET SODÆ.

Take of

Rhubarb, in powder . Bicarbonate of Soda . } equal parts Mix.

Dose, 10 to 20 grains.

#### PULVIS SENNÆ COMPOSITUS

(VEL PULVIS LIQUIRITIÆ COMPOSITUS PHARM. GERM.).

## Take of

Senna Leaves, in powder . 2 parts
Liquorice Root, in powder . 2 parts
Fennel Fruit, in powder . 1 part
Sublimed Sulphur . 1 part

White Sugar, in powder . 6 parts

# Mix.

Dose, one teaspoonful for an adult.

#### UNGUENTA1 \_\_\_OINTMENTS.

# UNGUENTUM BELLADONNÆ COMPO-SITUM.

# Take of

Extract of Belladonna . 80 grains
Distilled Water . a sufficiency

Ammoniated Mercury, in

powder . . . 40 grains
Benzoated Lard . . . 360 grains

Rub the extract with water to form a soft paste; then add the ammoniated mercury, and lastly the lard, and mix well.

N.B.—All the ointments may be made with Paraffin Ointment as a basis. This is a mixture of paraffins of a

# UNGUENTUM HYDRARGYRI AMMONIATI DILUTUM.

Take of

Ammoniated Mercury, in

powder . . . 8 grains
Simple Ointment . . 1 ounce

Mix.

# UNGUENTUM HYDRARGYRI AMMONIATI CUM SULPHURE.

Take of

Ammoniated Mercury, in

powder . . . 6 grains
Sublimed Sulphur . . 30 grains

Benzoated Lard . . 1 ounce

Mix.

# UNGUENTUM HYDRARGYRI NITRATIS MITE.

Take of

Ointment of Nitrate of

Mercury . . . 1 part
Paraffin Ointment . . 7 parts

Rub them well together.

soft unctuous consistency, having a melting point of 95° to 100° F. It is odourless, has a bland taste, and may be of a white opal or pale yellow colour. It has a minute crystalline structure not visible to the naked eye. It does not become rancid, and is not affected by boiling with a caustic alkaline solution. In commerce the substances known as Vaseline and Cosmoline (unguentum petrolei) may be taken as representing it.

# UNGUENTUM HYDRARGYRI NITRATIS MITIUS.

Take of

Ointment of Nitrate of

Mercury . . . 1 part

Paraffin Ointment . . . 11 parts

Mix.

## UNGUENTUM HYDRARGYRI OXIDI FLAVI.

Take of

Yellow Oxide of Mercury . 8 grains

Paraffin Ointment . . 1 ounce

Mix.

# UNGUENTUM HYDRARGYRI OXIDI RUBRI DILUTUM.

Take of

Ointment of Red Oxide of

Mercury . . . 1 part

Paraffin Ointment . . 7 parts

Rub them well together.

## UNGUENTUM HYDRARGYRI CUM BELLA-DONNA.

Take of

Extract of Belladonna . 2 parts

Distilled Water . . a sufficiency

Ointment of Mercury. . 6 parts

Rub the extract with water to form a soft paste; add the ointment, and mix well.

# UNGUENTUM PLUMBI SUBACETATIS CUM PARAFFINE.

Take of

Glycerole of Subacetate of

. 4½ fluid ounces

Lead . . Paraffin Ointment . 18 ounces

Paraffin Wax . . . 6 ounces

Melt the paraffin wax and ointment together, add the glycerole of lead, and stir till cold.

### UNGUENTUM SIMPLEX.

Take of

White Wax . . . 1 part

Prepared Lard . . . 2 parts

Melt them together.

# UNGUENTUM STAPHISAGRIÆ.

Take of

Stavesacre, in powder . 1 part Olive Oil . . . 1 part Prepared Lard . . . 2 parts

Melt the lard in a water bath, add the oil and stavesacre, and maintain the heat for a quarter of an hour, then remove and stir till cold.

#### UNGUENTUM ZINCI COMPOSITUM.

Take of

Compound Ointment of Subacetate of Lead B.P.

Rub them well together.

#### UNGUENTUM ZINCI OLEATIS.

## Take of

Oxide of Zinc . . 1 ounce
Oleic Acid . . 8 fluid ounces

Mix, and set aside for six hours; then apply heat to liquefy and form a perfect solution.

To this add

Paraffin Ointment . . 9 ounces

Dissolve, and stir till cold.

LONDON: PRINTED BY
SPOTTISWOODE AND CO., NEW-STREET SQUARE
AND PARLIAMENT STREET







