

A pharmacopœia : including the outlines of materia medica and therapeutics for the use of practitioners and students of veterinary medicine / by the late Richard V. Tuson.

Contributors

Tuson, Richard V.
Bayne, James

Publication/Creation

1895

Persistent URL

<https://wellcomecollection.org/works/pxfze425>

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>





22200197021

Med
K53601


L. G. Mrs. E. Holland

18th "The King's" Hussars


5/6

72m 2 Holland

XV Hussars



VETERINARY PHARMACOPŒIA



A
P H A R M A C O P Œ I A

INCLUDING THE OUTLINES OF
MATERIA MEDICA AND THERAPEUTICS

FOR THE USE OF
PRACTITIONERS AND STUDENTS
OF
Veterinary Medicine

BY THE LATE RICHARD V. TUSON

FIFTH EDITION
REVISED AND EDITED BY
JAMES BAYNE, F.C.S.

PROFESSOR OF CHEMISTRY AND TOXICOLOGY AT THE ROYAL VETERINARY COLLEGE



LONDON
J. & A. CHURCHILL
11 NEW BURLINGTON STREET

1895

31275689

WELLCOME INSTITUTE LIBRARY	
Coll.	welMOmec
Call	
No.	V

PREFACE
TO
THE FIFTH EDITION


IN THIS EDITION the original arrangement is retained. Several important preparations, which have been introduced since the last issue, have been added, and the whole work carefully revised and brought up to date. Reference has again been made to Finlay Dun's 'Veterinary Medicines,' and to Shoemaker's 'Materia Medica and Therapeutics.'

My thanks are due to my colleagues Professors MACQUEEN and HOBDAY, for the valuable assistance and advice which they have rendered me in the selection of new preparations, and the revision of doses. I am also indebted to my assistant, Mr. H. E. LINDLEY, F.C.S., for his help in the correction of proofs.

JAMES BAYNE.

ROYAL VETERINARY COLLEGE, N.W.

Jan. 14th, 1895.



PREFACE TO THE FIRST EDITION

THIS work is designed for the use of practitioners and students of veterinary medicine.

The agents are arranged in alphabetical order, according to the plan adopted in the 'British Pharmacopœia,' and are treated of under the following heads :

1. Latin Pharmaceutic Name.
2. English Pharmaceutic Name.
3. Synonyms.
4. Natural Order.
5. Composition.
6. Mode of Preparation.
7. Characters and Tests.
8. Actions and Uses.
9. Doses.
10. Modes of Application.
11. Incompatibles.
12. Antidotes.
13. Preparations.

Many medicines bearing old and familiar names have, for the sake of order, received new titles ; thus, the 'Diuretic Mass' of Morton is, in the present work, termed *Massa Resinæ Composita*. By whatever name, however, preparations may be known to the reader, they can be readily found by reference to the General Index.

The modes of preparation, characters, and tests are in most cases the same as those given in the 'British Pharmacopœia.'

The composition of bodies is expressed in modern atomic weights, a table of which, together with a list of the old ones, will be found in the Appendix.

In the directions for preparing chemico-pharmaceutic substances old chemical names are, for the most part, employed ; whereas, in explaining reactions and decompositions, that which the author believes to be the best of the new systems of nomenclature, supplemented where necessary by the old, is adopted.

The incompatibles mentioned are those recorded as such by the most eminent authorities ; although, no doubt, in many instances the incompatibility indicated is based upon hypothesis rather than upon the results of observation or experiment.

For a more detailed account of the actions and uses of veterinary medicines the reader is referred to the books of Dun and Morton, also to the forthcoming work on 'Veterinary Therapeutics' by Professor George Brown.

To the last-named gentleman the author is deeply indebted for his kind assistance in revising those portions of the work which are comprised under the sub-headings 'Doses' and 'Actions and Uses,' as well as for many valuable suggestions made during the progress of the volume through the press.


The principal sources, in addition to the 'British Pharmacopœia' of 1867, from which the author has obtained much valuable information, are Morton's 'Veterinary Pharmacy,' Dun's 'Veterinary Medicines,' Macnamara's edition of 'Neligan's Medicines,' and Garrod's 'Materia Medica and Therapeutics.'

ROYAL VETERINARY COLLEGE :

October 1869

CONTENTS

	PAGE
PREFACE TO THE FIFTH EDITION	v
PREFACE TO THE FIRST EDITION	vii
VETERINARY PHARMACOPŒIA	1
ADDENDA	325
APPENDIX:	
1. SYMBOLS AND ATOMIC WEIGHTS OF THE ELEMENTARY BODIES MENTIONED IN THIS WORK	331
2. WEIGHTS AND MEASURES OF THE BRITISH PHARMA- COPŒIA	332
3. WEIGHTS AND MEASURES OF THE METRICAL SYSTEM	333
4. RELATION OF THE WEIGHTS AND MEASURES OF THE BRITISH PHARMACOPŒIA TO THOSE OF THE METRI- CAL SYSTEM	333-34
INDEX TO VETERINARY MEDICINES, ARRANGED ACCORDING TO THEIR ACTIONS AND USES	335
GENERAL INDEX	349



A

VETERINARY PHARMACOPŒIA

ACACIA GUMMI

GUM ACACIA

Synonym.—Gum Arabic.

Composition.—An exudation from certain species of acacia, belonging to the natural order *Leguminosæ*, and consisting, when pure, of arabin, $C_{12}H_{22}O_{11}$.

Characters and Tests.—Spheroidal fragments, with shining surfaces ; nearly colourless ; bland and mucilaginous taste ; soluble in water, insoluble in alcohol, ether, and oils. Its cold aqueous solution gives a white gelatinous precipitate with lead subacetate, but no blue colour with iodine water, unless it be adulterated with starch.

Actions and Uses.—See Mucilago Acaciæ.

ACETANILIDE

PHENYL ACETAMIDE

Synonym.—Antifebrin.

Composition.—A derivative of aniline.

Characters and Tests.—A white powder, of neutral reaction, slightly pungent, slightly soluble in water, and

B

freely soluble in ether and alcohol. Not changed by acids or alkalies.

Actions and Uses.—Febrifuge, anti-spasmodic, anodyne, used to relieve pain in nervous affections and rheumatism.



Doses.—CATTLE, 1 drachm dissolved in ether.—DOGS, 5 grains dissolved in ether. These doses repeated every four hours, according to Mr. J. A. Nunn, Army Veterinary Department, produced marked lowering of temperature in fever, unaccompanied by nausea or other untoward effects.

ACETUM

VINEGAR

Composition.—Water holding in solution about 5·41 per cent. of real acetic acid, $\text{HC}_2\text{H}_3\text{O}_2$, together with colouring and flavouring matters.

Mode of Preparation.—Weak alcoholic liquids, *e.g.* beer, poor wines, &c., are exposed to the air and suffered to undergo the acetous fermentation, whereby the alcohol is oxidised first into aldehyd, and finally into acetic acid, thus :—



Characters and Tests.—A liquid of a brown colour, acetous odour, and sour taste. Sp. gr. 1·017 to 1·019. With barium chloride it should give no, or a very slight, precipitate, showing the absence, or the presence of only a minute quantity, of sulphuric acid.¹

¹ The addition of 1 part of sulphuric acid to 999 of vinegar is permitted by law.

Actions and uses.—*Internally* :—Refrigerant to allay febrile heat ; styptic, astringent, and diuretic ; also litholytic for calculi and urinary deposits consisting of calcium or magnesium phosphate or carbonate. Likewise employed as an antidote to poisoning by the alkalies and their carbonates. Very seldom, however, used internally.

Doses.—HORSE, 1 to 2 fluid ounces.—CATTLE, 1 to 2 fluid ounces.—SHEEP, 1 to 2 fluid drachms.—PIG, 1 to 2 fluid drachms.—DOG, 10 to 20 minims.

Incompatibles.—Alkalies ; earths ; and all carbonates.

Antidotes.—Potassium, sodium, calcium, or magnesium carbonate or bicarbonate.

ACETUM CANTHARIDIS

VINEGAR OF CANTHARIDES

Composition. — An impure solution of cantharidin, $C_8H_6O_2$, in acetic acid.

Mode of Preparation.—Take of

Cantharides in powder	.	.	2 ounces.
Glacial Acetic Acid	.	.	2 fluid ounces.
Acetic Acid	.	.	{ 18 fluid ounces, or a sufficiency.

Mix thirteen fluid ounces of the acetic acid with the glacial acetic acid, and digest the cantharides in this mixture for two hours at a temperature of 200° F. Cool the ingredients, transfer them to a percolator, and when the liquid ceases to flow, pour five fluid ounces of acetic acid over the residue in the apparatus. When the percolation is complete, press the contents of the percolator, filter the product, mix the liquids, and add sufficient acetic acid to make one pint.


Actions and Uses.—*Externally*:—Counter-irritant. Vesicant and rubefacient in local congestion, and in irritation of important organs.

ACIDUM ACETICUM

ACETIC ACID

Composition.—Water mixed with 33 per cent. of real acetic acid, $\text{HC}_2\text{H}_3\text{O}_2$.

Mode of Preparation.—By distilling a mixture of crystallised sodium acetate and sulphuric acid; sodium sulphate remains in the retort, while water and acetic acid, in the proportions above named, pass over into the receiver.



Characters and Tests.—A colourless, sour liquid, having a pungent odour. Sp. gr. 1.044. Should leave no residue when evaporated, and will give no precipitate with hydro-sulphuric acid, barium chloride, or silver nitrate, if free from lead, copper, sulphuric acid, and chlorine. Put a fluid drachm of the acid, mixed with half an ounce of distilled water and half a drachm of pure hydrochloric acid, also a few pieces of granulated zinc, into a flask. While effervescence continues suspend a slip of white blotting-paper moistened with a solution of subacetate of lead in the upper part of the flask above the liquid for about five minutes. The paper should not become discoloured, thus indicating the absence of sulphurous acid.

Actions and Uses.—*Externally*:—Stimulant, astringent, styptic, rubefacient, and vesicant. As a caustic for the removal of warts and similar growths. For the removal of

scurf in ringworm, mallenders and sallenders, scab, and mange.¹

Preparations.—Acetum Cantharidis ; Acidum Aceticum Dilutum.

ACIDUM ACETICUM DILUTUM

DILUTED ACETIC ACID

Composition.—Water mixed with 4·27 per cent. of real acetic acid, $\text{HC}_2\text{H}_3\text{O}_2$.

Mode of Preparation.—Take of

Acetic Acid	1 pint.
Distilled Water	7 pints.

Mix.

Characters and Tests.—Colourless, sour liquid. Sp. gr. 1·006.

Actions and Uses.

Doses.

Incompatibles.

Antidotes.

} Similar to vinegar.

} See Acetum.

ACIDUM ACETICUM GLACIALE

GLACIAL ACETIC ACID

Synonyms.—Acetic Hydrate ; Real Acetic Acid.

Composition.— $\text{HC}_2\text{H}_3\text{O}_2$.

Mode of Preparation.—By distilling a mixture of concentrated sulphuric acid and sodium acetate deprived of its water of crystallisation by fusion ; sodium sulphate remains in the retort, and glacial acetic acid passes over

¹ For these purposes, and as an astringent and styptic, impure pyroligneous acid is preferred to Acidum Aceticum, on account of the former containing carbolic acid and similar compounds.

into the receiver. See *Acidum Aceticum* for an explanation of the decomposition.

Characters and Tests.—At ordinary temperatures a colourless liquid, with a pungent odour and sour taste. Sp. gr. 1.065 to 1.066. Cooled to 34°F. it crystallises and remains solid unless heated to 48°F. Tested for sulphurous acid, as directed under *Acidum Aceticum*, it should give no indication of this impurity.

Actions and Uses.—*Internally* :—Irritant poison. *Externally* :—Rubefacient, vesicant, and caustic. Used to destroy warts.

Antidotes.—See *Acetum*.

Preparations.—*Acetum Cantharidis* ; *Mistura Creasoti*.

ACIDUM ARSENIOSUM

ARSENIOUS ACID

Synonyms.—Arsenious Anhydride ; White Arsenic.

Composition.— As_2O_3 .

Mode of Preparation.—By roasting arsenical ores, *e.g.* arsenical pyrites, FeSAs , in a current of air ; the arsenicum combines with oxygen and becomes arsenious anhydride, which is purified by sublimation.

Characters and Tests.—Heavy, white, glistening powder or in translucent masses resembling porcelain. Completely volatile at 400°F. Sublimed in a test-tube, it condenses in brilliant octahedral or triangular crystals. Soluble in water, 1 in 80 parts. Its aqueous solution furnishes a yellow precipitate (silver arsenite) with silver ammonio-nitrate, and a green precipitate (copper arsenite) with copper ammonio-sulphate. These precipitates are soluble in ammonia and in nitric acid.

Actions and Uses.—*Internally* : In excessive doses, an irritant poison ; in medicinal doses, alterative and tonic in chronic rheumatism, paralysis, epilepsy, chorea, farcy, and mange. *Externally* :—Used in the cure of mange, scab, and other skin diseases ; also as a caustic to eradicate warts, and to produce the sloughing and effect the removal of malignant tumours. When used externally, especially for the latter purpose, arsenious acid must be employed with great caution, as it is liable to be absorbed and to kill the patient.

Doses.—CATTLE, 5 to 10 grains.—HORSE, 5 to 10 grains. SHEEP, 1 to 2 grains.—PIG, $\frac{1}{2}$ to 2 grains.—DOG, $\frac{1}{15}$ to $\frac{1}{10}$ of a grain.

Modes of Application.—*Internally* :—The powder sprinkled over, or mixed with, the animal's food.¹ *Externally* :—Liquor Arsenicalis ; Unguentum Arsenici.

Antidotes.—Freshly made moist hydrated peroxide of iron ; magnesia ; cold affusions ; inhalation of ammonia.

Preparations.—Liquor Arsenicalis ; Unguentum Arsenici.

ACIDUM BORICUM

BORIC ACID

Synonym.—Boracic Acid.

Composition.— H_3BO_3 .

Mode of Preparation.—Found naturally in the volcanic districts of North Italy. See Borax.

Characters and Tests.—A transparent, colourless solid which crystallises in six-sided plates. It has a feebly acid taste. It dissolves in 25 parts of water or 15 of alcohol,

¹ See Liquor Arsenicalis, a better form of arsenical medicine for internal use.

and in 3 parts of boiling water or 5 of boiling alcohol. An alcoholic solution burns with a green flame.

Actions and Uses.—*Externally* :—Antiseptic, used as a dressing for wounds. The saturated solution is used in the treatment of scaly and parasitic skin eruptions.

Preparations.—Boroglyceride, a combination of boric acid (62 parts) and glycerin (92 parts). This diluted with an equal quantity of glycerin makes the fifty per cent. boroglyceride, in which form it is used as an antiseptic.

Lister's antiseptic ointment for dressing wounds is made by adding 1 part each of boric acid and white wax to 2 parts each of paraffin and almond oil. Borated lint or borated cotton-wool is made by steeping the substance in a saturated solution of the acid and allowing it to dry.


ACIDUM CARBOLICUM

CARBOLIC ACID

Synonyms.—Carbolic Hydrate ; Phenic Acid ; Phenol ; Phenylic Alcohol.

Composition.— $\text{HC}_6\text{H}_5\text{O}$.

Mode of Preparation.—Those portions of coal-tar oil which distil between 300°F . and 400°F . are mixed with a strong and boiling solution of potassium hydrate. The resulting potassium carbolate is decomposed by hydrochloric acid, and the liberated carbolic acid rectified by distillation off calcium chloride. The calcium chloride deprives the acid of moisture.



Characters and Tests.—Acicular crystals, which, when freshly prepared and quite pure, are colourless, but are often brown or turn so by keeping. At 95°F. they become an oily liquid, having an odour and taste like creasote. Sp. gr. 1·065 ; boiling-point, 370°F. Exposed to the air, the crystals absorb moisture and liquefy. The acid is slightly soluble in water, 1 in 15 parts, but freely soluble in glycerin, alcohol, and ether. About a grain of hypochlorite of calcium added to a little aqueous solution of carbolic acid, placed in a test-tube, produces, after agitation, the addition of a few drops of ammonia, and the application of a gentle heat, a bright blue colour with a tinge of green. An aqueous solution of carbolic acid will also give a blue colour with a few drops of ferric chloride. One drachm of the acid, if pure, completely dissolves on being shaken with half a pint of warm water.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses it is occasionally employed as an anthelmintic ; also given to dogs to allay vomition. *Externally* :—Caustic, disinfectant, astringent, and styptic ; applied to cancerous sores and ulcers which emit a fetid odour, *e.g.* canker and thrush in the foot of the horse, and foot-rot in sheep. It modifies suppuration, and facilitates cicatrisation when applied to wounds. Said to render inert the virus of cholera, cattle plague, and other contagious diseases.

Doses.—*Crystals* :—HORSE, 20 to 80 grains.—CATTLE, 20 to 80 grains.—SHEEP, 5 to 15 grains.—PIG, 5 to 15 grains.—DOG, 1 to 5 grains.

Fluid : HORSE, 20 to 80 minims.—CATTLE, 20 to 80 minims.—SHEEP, 5 to 15 minims.—PIG, 5 to 15 minims.—DOG, 1 to 5 minims.

Modes of Application.—*Internally* :—Dissolved in dilute spirits of wine, or made into a bolus with linseed meal or

common mass. *Externally* :—As a caustic, the undiluted acid ; for other purposes, in the form of lotion, liniment, or ointment.

Buildings may be disinfected (?) and the atmosphere impregnated with the acid by steeping rags in it and then suspending them in various parts of the room, stable, &c. ; or the acid, in a diluted form,¹ may be sprinkled over floors and walls.

Antidotes.—Albumen ; soap ; demulcent drinks ; oil.

Preparations

Glycerinum Acidi Carbolici	Unguentum Acidi Carbolici.
Linimentum Acidi Carbolici.	Collodium Hæmostaticum.
Lotio Acidi Carbolici.	Sodii Sulphocarbolas.

ACIDUM CHROMICUM

CHROMIC ACID

Synonym.—Chromic Anhydride.

Composition.— CrO_3 .

Mode of Preparation.—By adding to one measure of a saturated solution of potassium dichromate one and a half measures of concentrated sulphuric acid, by small portions at a time, and allowing the solution to cool, when chromic anhydride crystallises out.

Characters and Tests.—Crystalline in fine crimson needles, which are deliquescent, and very soluble in water. They decompose sometimes with explosion when mixed

¹ Pour an ounce or two of the fluid acid into a gallon bottle, nearly fill the bottle with water, shake, and set aside for twenty-four hours. From the undissolved acid decant the aqueous solution for use.

with alcohol. An explosion occurs when 1 part of chromic anhydride is mixed with 2 parts of glycerin.

Actions and Uses.—*Externally* in solution (from 5 to 20 per cent.) it acts as a caustic. In the solid form it rapidly destroys tissue (on account of its affinity for water), forming an eschar. It is a powerful disinfectant.

ACIDUM GALLICUM

GALLIC ACID

Composition.— $\text{H}_3\text{C}_7\text{H}_3\text{O}_5, \text{H}_2\text{O}$.

Mode of Preparation.—Take of

Galls, in coarse powder	.	.	.	1 pound.
Distilled Water	.	.	.	A sufficiency.

Place the powdered galls in a porcelain dish, pour on as much water as will convert it into a thick paste, and keep it in this moist condition for six weeks, at a temperature between 60° and 70° ; at the end of that time boil the paste for twenty-minutes with forty-five ounces of water, strain through calico, and when the fluid has cooled collect the crystalline deposit which has formed, and let it drain. If the gallic acid is required very pure, it should be recrystallised.

Characters and Tests.—Crystalline in acicular prisms or silky needles of a fawn colour; it is soluble 1 in 100 parts of cold water, but dissolves in 3 parts of boiling water. Soluble also in rectified spirit 1 in 8, glycerin 1 in 20. It leaves no residue on ignition with free access of air, it gives a bluish-black precipitate with a ferric salt, but no precipitate with a ferrous salt or a solution of gelatin, showing the absence of tannic acid.

Actions and Uses.—The same as Tannic Acid.


ACIDUM HYDROCHLORICUM

HYDROCHLORIC ACID

Synonyms.—Muriatic Acid ; Spirit of Salt.

Composition.—Water holding in solution 31·8 per cent. of its weight of hydrochloric acid gas, HCl.

Mode of Preparation.—By distilling a mixture of sodium chloride, sulphuric acid, and water, and allowing the distillate to pass into a receiver containing distilled water.



Sodium sulphate remains in the retort, while hydrochloric acid gas passes over and is absorbed by the water in the receiver.

Characters and Tests.—Colourless, strongly acid liquid, which emits white vapours and has a pungent odour. Sp. gr. 1·16. With silver nitrate it gives a white curdy precipitate (silver chloride), which is soluble in ammonia but insoluble in boiling nitric acid. Should entirely evaporate on the application of heat, and, after being diluted with four times its volume of distilled water, should give no precipitate with hydrosulphuric acid or barium chloride, proving the absence of lead, copper, and sulphuric acid ; neither should a strip of polished copper tarnish when boiled with the acid thus diluted, showing the absence of arsenic and antimony.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, tonic. Said to act as a litholytic on earthy phosphates and carbonates ; probably it possesses this property only when injected into the

bladder. If employed for this purpose, it must be highly diluted. See *Acidum Hydrochloricum Dilutum*. *Externally* :—Caustic, astringent, antiseptic.

Doses.—For internal use, *Acidum Hydrochloricum Dilutum* is employed.

Modes of Application.—For external use the part is painted with the acid.

Antidotes.—Carbonate of lime (chalk or whiting); magnesia; carbonate of magnesia; demulcent drinks; large draughts of water.

Preparation.—*Acidum Hydrochloricum Dilutum*.

ACIDUM HYDROCHLORICUM DILUTUM

DILUTE HYDROCHLORIC ACID

Composition.—Six fluid ounces of this preparation consist of water holding in solution 36·5 grains (one molecular weight) of hydrochloric acid gas, HCl.

Mode of Preparation.—Take of

Hydrochloric Acid 8 fluid ounces.

Distilled Water A sufficiency.

Mix the hydrochloric acid with sixteen fluid ounces of the distilled water, and add more water, so that at 60°F. the mixture shall measure twenty-six and a half fluid ounces.

Characters and Tests.—Sp. gr. 1·052. For tests, see *Acidum Hydrochloricum*.

Actions and Uses.—Tonic and litholytic internally.

Doses.—HORSE, 1 to 4 fluid drachms.—CATTLE, 1 to 4 fluid drachms.—SHEEP, 10 to 20 minims.—PIG, 10 to 20 minims.—DOG, 2 to 10 minims.

Modes of Application.—Diluted with about a hundred times its bulk of water, or combined with infusions of vegetable tonics. As a litholytic, see *Acidum Hydrochloricum*.

Incompatibles.—Carbonates ; sulphides ; salts of silver.

Antidotes.—Same as to *Acidum Hydrochloricum*.


ACIDUM HYDROCYANICUM DILUTUM

DILUTED HYDROCYANIC ACID

Synonym.—Prussic Acid.

Composition.—Water containing 2 per cent.¹ by weight of real hydrocyanic acid, HCN or HCy.

Mode of Preparation.—By distilling a mixture of potassium ferrocyanide, sulphuric acid, and water.



Potassium-hydrogen sulphate and potassium-ferrous ferrocyanide remain in the retort, hydrocyanic acid accompanied by water passes over into the receiver, into which a quantity of water has been previously placed. By the addition of water the distillate is made of such a strength that 100 grains or 110 minims of it precipitated with silver nitrate shall yield ten grains of silver cyanide.

Characters and Tests.—Colourless liquid, with a peculiar

¹ Scheele's Prussic acid, as met with in the shops, is a preparation of uncertain strength, the amount of real hydrocyanic acid contained in it varying from one to four per cent.

odour.¹ Sp. gr. 0.997. With potassium hydrate and a mixture of ferrous and ferric sulphates it yields, after being acidulated with hydrochloric acid, a precipitate of Prussian blue. Silver nitrate furnishes a white curdy precipitate of silver cyanide which is soluble in ammonia and in boiling nitric acid. Barium chloride should give no precipitate, proving the absence of sulphuric acid and sulphates.

Actions and Uses.—*Internally* :—In excessive doses, narcotic poison ; in medicinal doses, sedative, antispasmodic, and anodyne. Used in chronic cough, chorea, epilepsy, chronic vomiting, carditis, palpitation of the heart, rheumatism, and tetanus.

Doses.—HORSE, 20 to 30 minims.—CATTLE, 20 to 30 minims.—SHEEP, 5 to 10 minims.—PIG, 5 to 10 minims.—DOG, 1 to 3 minims.

To prevent accidents, the smaller doses should be given at first, and afterwards gradually increased.

Modes of Application.—*Internally* :—Diluted with water. In troublesome cough in horses, twenty minims of the acid with a drachm each of camphor and extract of belladonna, made into a ball with powdered liquorice or linseed meal, and given two or three times a day. To the dog, in obstinate vomiting, two grains of the acid with ten grains of carbonate of soda and one ounce of water may be administered every hour. One drachm of the acid with about a quart of water, employed slowly as an enema two or three times a day, lessens muscular contractions in tetanus. *Externally* :—To allay pain and irritation in chronic skin affections, especially in dogs, two to three drachms of the acid are mixed with a pint of distilled or rain water.

¹ This character should be very cautiously observed, as the vapour of the acid, when inhaled, is highly dangerous.

Incompatibles.—Nitrate of silver ; sulphates and chlorides of iron with an alkali.

Antidotes.—Fresh air ; affusions of cold water ; inhalation of ammonia or chlorine ; freshly precipitated hydrated peroxide of iron mixed with an alkaline carbonate.


ACIDUM NITRICUM

NITRIC ACID

Synonym.—Aqua fortis.

Composition.—Water containing 70 per cent. of real nitric acid, HNO_3 .

Mode of Preparation.—By distilling a mixture of potassium (or sodium) nitrate and concentrated sulphuric acid ; potassium-hydrogen sulphate remains in the retort, and nitric acid distils over.



Characters and Tests.—Colourless liquid. Sp. gr. 1.42 ; boiling-point, 250°F . Exposed to air, it emits acrid and corrosive fumes. Copper dropped into a mixture of equal parts of the acid and water causes the evolution of a colourless gas (nitric oxide), which, as it mixes with the oxygen of the air, forms ruddy fumes (nitrous anhydride, N_2O_3 , and nitric peroxide, NO_2). A crystal of ferrous sulphate dropped into the acid, diluted with an equal bulk of water, becomes surrounded by a brown cloud if the mixture be cold. The acid should leave no residue on evaporation. Diluted with six times its volume of distilled water, it should give no precipitate with barium nitrate or silver nitrate, showing the absence of sulphuric and hydrochloric acids.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, astringent and tonic. *Externally* :—Deodoriser and disinfectant ; also used as a caustic for removing warts and fungous growths, and in canker ; for improving the condition and destroying the fœtor of unhealthy wounds, caries, foul, and foot-rot.

Doses.—See Acidum Nitricum Dilutum, the preparation of nitric acid for internal use.

Modes of Application.—*Internally* : see Acidum Nitricum Dilutum. *Externally* :—As a caustic, by means of a pledget of tow tied to a stick, or tow saturated with the acid laid on the part. As an excitant and deodoriser to indolent and sphacelated wounds and ulcers, a mixture of two to four drachms of the acid and one pint of water may be employed. Tow thoroughly saturated with a mixture of four parts of tar and one part of nitric acid is applied to the feet of horses affected with canker.

Antidotes.—Copious draughts of water ; demulcent drinks ; albumen ; milk ; soap ; alkaline carbonates or bicarbonates ; carbonate of calcium (chalk or whiting) ; magnesia ; carbonate of magnesia.

Preparation.—Acidum Nitricum Dilutum.

ACIDUM NITRICUM DILUTUM

DILUTED NITRIC ACID

Composition.—Water containing 17·44 per cent. of real nitric acid, HNO_3 .

Mode of Preparation.—Take of

Nitric Acid	6 fluid ounces.
Distilled Water	A sufficiency.

Dilute the acid with twenty-four ounces of the water,

then add more water, so that at 60°F. the mixture shall measure thirty-one fluid ounces.

Tests.—Colourless, sour. Sp. gr. 1·101.

Actions and Uses.—*Internally*:—Tonic, astringent; during convalescence from debilitating disorders, in chronic skin diseases, and in chronic enlargement of the liver in cattle.

Doses.—HORSE, 1 to 3 fluid drachms.—CATTLE, 2 to 4 fluid drachms.—SHEEP, 10 to 20 minims.—PIG, 10 to 20 minims.—DOG, 2 to 10 minims.

Mode of Application.—Diluted with from fifty to a hundred times its bulk of water.

Incompatibles.—Alkalies, earths, and other oxides; carbonates; bicarbonates; sulphides; sulphate of iron; and acetate of lead.

Antidotes.—See Acidum Nitricum.

ACIDUM NITRO-HYDROCHLORICUM DILUTUM

DILUTED NITRO-HYDROCHLORIC ACID

Synonym.—Diluted Aqua Regia.

Composition.—An aqueous solution of chlorine, chloro-nitric and chloro-nitrous gases.



Mode of Preparation.—Take of

Nitric Acid	3 fluid ounces.
Hydrochloric Acid	4 fluid ounces.
Distilled Water	25 fluid ounces.

Mix the acids and allow them to remain for at least twenty-four hours in a bottle, the mouth of which is

partially closed ; then add the water in successive portions, shake the bottle after each addition, and preserve the mixture in a stoppered bottle.

On mixing the acids they gradually suffer mutual decomposition.



Characters and Tests.—Colourless liquid. Sp. gr. 1·074.
Dissolves gold leaf.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, alterative and tonic. For the latter purposes it is preferred to either nitric or hydrochloric acid separately.

Doses.—HORSE, 1 to 3 fluid drachms.—CATTLE, 2 to 4 fluid drachms.—SHEEP, 10 to 20 minims.—PIG, 10 to 20 minims.—DOG, 2 to 10 minims.

Mode of Application.—Diluted with not less than a hundred times its bulk of water.

Incompatibles.	}	Same as Acidum Nitricum Dilutum.
Antidotes.		

ACIDUM PHOSPHORICUM CONCENTRATUM

PHOSPHORIC ACID

Synonym.—Orthophosphoric acid.

Composition.— H_3PO_4 with 33 per cent. of water.

Mode of Preparation.—By heating phosphorus with dilute nitric acid until nitrous fumes cease to form and diluting with water until the sp. gr. is 1·5.

Characters and Tests.—A colourless, sour, syrupy liquid, with an acid reaction. In diluted solution it gives a yellow precipitate with silver nitrate.

Actions and Uses.—*Externally* :—An irritant and escharotic.

ACIDUM PHOSPHORICUM DILUTUM

DILUTE PHOSPHORIC ACID

Composition.—Water containing 13·8 per cent. of H_3PO_4 .

Mode of Preparation.—By mixing three parts of concentrated phosphoric acid with twenty of water.

Actions and Uses.—In debilitated conditions of the system, has the reputation of checking tuberculosis. It is a good tonic, and useful in bronchial catarrh.

Doses.—HORSE, 1 to 2 drachms.—DOG, 5 to 10 minims.

Preparations.—Phosphates of ammonia ; iron and soda ; Syrupus ferri quiniæ et strychninæ phosphatum ; syrup of the phosphates of iron, quinine, and strychnine (Easton's syrup) contains iron phosphate 1·33, quinine 1·33, strychnine 0·04, phosphoric acid 8, sugar 60, water q.s. ad 100 parts : each drachm contains about $\frac{1}{32}$ gr. of strychnine.

ACIDUM SALICYLICUM

SALICYLIC ACID

Synonym.—Hydroxybenzoic acid.

Composition.— $\text{HC}_7\text{H}_5\text{O}_3$ or $\text{C}_6\text{H}_4\cdot\text{OH}\cdot\text{COOH}$.

Mode of Preparation.—By passing carbonic anhydride through sodium phenol for several hours, heating to 482°F ., and treating the residue with dilute hydrochloric acid.

Characters and Tests.—A colourless powder, usually crystallising in minute acicular crystals. Taste at first sweet, afterwards bitter. Soluble in water, 1 in 500 to 700 ; more soluble in hot water, alcohol, and ether. Ferric chloride gives a violet colour both with the acid and its salts. Copper sulphate gives an emerald green colour.

Actions and Uses.—Antiseptic, antipyretic, both the acid, and sodium salicylate are employed in the treatment of rheumatism. *Externally* :—It is used as a dressing for wounds, and to relieve the itching and discharge of eczema and psoriasis. Also in the treatment of open joints.

Doses.—HORSES and CATTLE, 1 to 2 drachms ; SHEEP and DOGS, from 10 to 15 grains.

Preparations.—Solution of one part each of salicylic acid and borax to thirty or fifty parts of water, used for soaking lint, &c., as an antiseptic covering for wounds.

Ointment.—Salicylic acid, one part, mixed warm with twenty to twenty-five of vaselin.

The B.P. ointment contains one part salicylic acid, eighteen soft paraffin, and nine hard paraffin.


ACIDUM SULPHURICUM

SULPHURIC ACID


Synonym.—Oil of Vitriol.

Composition.—96·8 per cent. by weight of sulphuric hydrate, H_2SO_4 , with 3·2 per cent. of water.


Mode of Preparation.—Sulphurous anhydride, vapour of nitric acid, steam, and air are conducted into a leaden chamber, where they react upon one another in the following manner :



The resulting nitric oxide abstracts oxygen from the air, and becomes nitric peroxide :



The nitric peroxide thus formed oxidises further quantities of sulphurous anhydride, and, with the assistance of steam, converts it into sulphuric acid :



The acid produced in the leaden chambers is very dilute ; it is concentrated by evaporation, first in leaden basins, and finally in glass or platinum retorts.

Characters and Tests.—Colourless, heavy, oily, intensely acid liquid. Sp. gr. 1·843. Mixed with water, much heat is evolved. Diluted with five or six times its volume of water, it gives, with barium chloride, a white precipitate (barium sulphate), which is insoluble in nitric acid ; it should yield no precipitate or darkening of colour with hydrosulphuric acid, indicating the absence of lead. When a solution of ferrous sulphate is carefully poured upon the surface of the acid, there should be no brown colour developed where the two liquids unite, showing the absence of nitric acid and oxides of nitrogen. The acid should leave no residue on evaporation.

Actions and Uses.—*Internally* :—Irritant and corrosive poison ; see Acidum Sulphuricum Dilutum. *Externally* :—Employed as a caustic for the removal of warts and cancerous growths.

Mode of Application.—Made into a paste with flowers of sulphur, sawdust, finely powdered charcoal, or dried alum.

Antidotes.—Same as for Acidum Nitricum

ACIDUM SULPHURICUM DILUTUM**DILUTED SULPHURIC ACID**

Composition.—Water containing 12·42 per cent. of real sulphuric acid, H_2SO_4 .

Mode of Preparation.—Take of

Sulphuric Acid	7 fluid ounces.
Distilled Water	A sufficiency.

Place seventy-seven fluid ounces of distilled water in a thin glass or earthenware vessel, and then pour into it the seven fluid ounces of acid. Stir the mixture, and, when cooled to 60°F ., add more water, so that it shall measure eighty-three and a half fluid ounces.

Characters and Tests.—Colourless, sour liquid. Sp. gr. 1·094.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, astringent ; also tonic during convalescence, and in other cases in which mineral tonics are indicated. *Externally* :—Astringent.

Doses.—HORSE, 2 to 4 fluid drachms.—CATTLE, 2 to 4 fluid drachms.—SHEEP, 10 to 20 minims.—PIG, 10 to 20 minims.—DOG, 2 to 10 minims.

Modes of Application.—*Internally* :—As a tonic, it is given diluted with from fifty to a hundred times its bulk of water. In diarrhœa and dysentery, accompanied by alkaline discharges, two fluid drachms of acid, with one fluid ounce of laudanum, may be administered twice a day in gruel or some mucilaginous drink. One fluid drachm of this acid, with one ounce each of sulphuric ether and powdered cinchona bark, is sometimes given in gruel or

ale to horses suffering from typhoid influenza. In cases of relaxed and ulcerated sore throat in horses, the acid, highly diluted with water, may be slowly given.

Externally :—As an astringent, thirty minims of acid in one fluid ounce of water.

Incompatibles.	} Same as Acidum Nitricum.
Antidotes.	

ACIDUM SULPHUROSUM

SULPHUROUS ACID

Composition.—Water containing 9·2 per cent. by weight of sulphurous anhydride (sulphurous acid gas), SO_2 .¹


Mode of Preparation.²—Take of

Sulphuric Acid	4 fluid ounces.
Wood Charcoal, in coarse powder	1 ounce.
Common Water	2 fluid ounces.
Distilled Water	20 fluid ounces.


Put the charcoal and sulphuric acid into a glass flask, connected by a glass tube with a wash-bottle containing the two fluid ounces of common water, whence a second tube leads into a pint bottle containing the distilled water, to the bottom of which the gas-delivery tube should pass. Apply heat to the flask until gas is evolved, which is to be conducted through the water in the wash-bottle, and then into the distilled water, the latter being kept cold, and the

¹ Equal to 11·78 per cent. of hypothetical sulphurous acid, H_2SO_3 .


² For fumigating or disinfecting purposes, sulphurous anhydride may be easily evolved by setting fire to a few fragments of roll sulphur (brimstone) placed in a saucer or other suitable vessel.



process being continued until the bubbles of gas pass through the liquid undiminished in size.



The product should be preserved in a stoppered bottle, to prevent the conversion of the sulphurous anhydride, by absorption of oxygen, into sulphuric acid ; thus :—



It should be kept in a cool place, to prevent loss of gas.

Characters and Tests.—A colourless liquid, with an odour like that emitted from burning sulphur. Sp. gr. 1.04. Should leave no residue on evaporation, and should give no precipitate, or but a very slight one, with barium chloride, showing the absence, or nearly so, of sulphuric acid.

Actions and Uses.—Antiseptic, deodoriser, disinfectant.¹ Applied externally to kill fungi that cause or accompany certain skin affections.

Mode of Application.—One part of the acid mixed with three parts of water, or two of glycerin, applied to the diseased surface with a sponge.

ACIDUM TANNICUM

TANNIC ACID

Synonym.—Tannin.

Composition.—Pure tannic acid has the formula $\text{C}_{13}\text{H}_9\text{O}_7\cdot\text{CO}_2\text{H}$, but that obtained by the method described

¹ Active principles of Tuson's disinfectant, sulphurous acid and chloride of zinc. See Zinci Chloridum.

below, and taken from the 'British Pharmacopœia' of 1867, is mixed with gallic acid, $C_6H_2(OH)_3CO_2H$.¹

Mode of Preparation.—Take of

Oak Galls in powder	} Of each a sufficiency.
Ether	

Expose the powdered galls to a damp atmosphere for two or three days, and afterwards add sufficient ether to form a soft paste. Let this stand in a well-closed vessel for twenty-four hours, then, having quickly enveloped it in a linen cloth, submit it to a strong pressure in a suitable press so as to separate the liquid portion. Reduce the pressed cake to powder, mix it with sufficient ether, to which one sixteenth of its bulk of water has been added, to form again a soft paste, and press this as before. Mix the expressed liquids, and expose the mixture to a spontaneous evaporation until, by the aid subsequently of a little heat, it has acquired the consistence of a soft extract; then place it on earthen plates or dishes, and dry it in a hot-air chamber at a temperature not exceeding 212°F.

Under the combined influence of moisture and atmospheric oxygen tannic acid is, according to one view, converted into gallic acid, water, and carbonic anhydride, and, according to another view, into gallic acid and glucose.

Characters and Tests.—Pale yellow vesicular masses or thin glistening scales, strongly astringent taste, and acid reaction. Readily soluble in water, ten in eight parts, and rectified spirit, sparingly soluble in ether. With an aqueous solution gelatin gives a yellowish-white precipitate (gelatin tannate), and ferric chloride furnishes a bluish-black

¹ By some practitioners gallic acid is regarded as being a more effective astringent and styptic than tannic acid, and is prescribed in cases where hæmorrhage has to be reached through the circulation. The doses are the same as those of tannic acid.

precipitate (ferric tannate). The acid should leave no residue when burned with free access of air.

Actions and Uses.—*Internally and Externally* :—Astringent and styptic. Employed to diminish mucous discharges, as in diarrhœa and dysentery ; also used as a vermicide. Antidote to tartar emetic ; also to strychnia, morphia, and other alkaloidal poisons. Tannic acid is an ingredient of styptic collodion.

Doses.—HORSE, 20 to 60 grains.—CATTLE, 20 to 60 grains.—SHEEP, 4 to 12 grains.—PIG, 4 to 12 grains.—DOG, 1 to 3 grains.

Modes of Application.—*Internally* :—Either in the form of bolus, pill, or solution in water or spirit of wine. *Externally* :—One part of acid dissolved in six parts water ; or Unguentum Acidi Tannici.

Incompatibles.—Mineral acids ; alkalies ; salts of lead, silver, iron, antimony ; alkaloids ; gelatin, and emulsions.

Preparations.—Glycerinum Acidi Tannici ; Unguentum Acidi Tannici.

ACONITI FOLIA

ACONITE LEAVES

The fresh leaves and flowering tops of *Aconitum Napellus* (Monkshood, Wolfsbane, Blue Rocket), gathered when about one third of the flowers are expanded, from plants cultivated in Britain.

Natural Order.—*Ranunculaceæ*.

Characters.—Leaves smooth, palmate, divided into five deeply cut wedge-shaped segments, which are further

incised in a pinnatifid manner, dark green above, paler beneath ; slowly exciting, when *cautiously* chewed, a sensation of tingling. Flowers numerous, irregular, deep blue in dense racemes.

Actions and Uses.—*Internally*:—In excessive doses, cerebro-spinal poison ; in medicinal doses, sedative and anodyne
Externally :—Occasionally applied to allay neuralgic and rheumatic pains ; for this purpose, however, Linimentum Aconiti or Tinctura Aconiti is usually preferred.

Antidotes.—Emetics, when they can be employed ; stimulants, internally and externally.

Preparations.—Extractum Aconiti ; Linimentum Aconiti ; Tinctura Aconiti.

ACONITI RADIX

ACONITE ROOT

The dried root of *Aconitum Napellus*. Imported from Germany or cultivated in Britain, and collected in the winter or early spring, before the leaves have appeared.

Natural Order.—*Ranunculaceæ*.

Characters.—Usually from one to three inches long, not more than about three quarters of an inch in thickness at the crown, tapering ; externally, blackish-brown ; internally, whitish ; earthy odour. A minute portion *cautiously* chewed causes prolonged tingling and numbness.

Antidotes.—See Aconiti Folia,

Preparations.—Aconitia ; Linimentum Aconiti ; and Tinctura Aconiti,

ACONITINA**ACONITINE**

Synonym.—Aconitia.

A highly poisonous alkaloid, $C_{33}H_{43}NO_{12}$, residing in, and forming the active principle of, the root, leaves, and other parts of several species of aconite. Not used in veterinary practice, as it is very costly.

ADEPS PRÆPARATUS**PREPARED LARD**

Obtained from *Sus scrofa*.

Natural Order.—*Pachydermata*.

Synonyms.—Axungia ; Hog's Lard.

Composition.—A mixture of 38 per cent. of margarin and stearin and palmitin, with 62 per cent. of olein.

Preparation.—The fat of the hog (*Sus scrofa*), taken from the loins, omentum, and mesentery, is deprived of its membranes and purified by being kneaded with cold water, melted, strained, dried at a little above 212°F ., again strained through flannel, and finally run into bladders or casks and allowed to solidify.

Characters and Tests.—Soft, white, fatty substance, melting at about 100°F . Should not have a rancid odour, and should dissolve entirely in ether. If boiled with distilled water and filtered, the filtrate should give no precipitate with silver nitrate, showing the absence of common salt ; and, when quite cold, no blue colour with solution of iodine, proving the absence of starch and flour.

Actions and Uses.—Chiefly employed externally as a simple dressing to ulcers and blistered surfaces, which it softens and protects from the irritating action of the atmosphere and of acrid discharges. Occasionally to kill the acarus in mange, scab, and similar complaints. Extensively used in the preparation of ointments.

ÆTHER

ETHER

Synonyms.—Ethylic Oxide ; Ethylic Ether ; Sulphuric Ether.

Composition.—About 92 per cent. of pure ether (C_2H_5)₂O or $C_4H_{10}O$, with about 8 per cent. of impurities, chiefly water and alcohol.

Preparation.—Take of



Rectified Spirit	.	.	.	50 fluid ounces.
Sulphuric Acid	.	.	.	10 fluid ounces.
Chloride of Calcium	.	.	.	10 ounces.
Slaked Lime	.	.	.	$\frac{1}{2}$ ounce.
Distilled Water	.	.	.	13 fluid ounces.

Mix the sulphuric acid with twelve fluid ounces of the spirit in a glass retort capable of holding at least two pints ; connect the retort with a Liebig's condenser to which a receiver has been attached, and distil with a heat sufficient to maintain the liquid in brisk ebullition. Allow the remainder of the spirit to slowly flow in a continuous stream into the retort. For this purpose use a tube furnished with a stopcock to regulate the supply, connecting one end of the tube with a vessel containing the spirit raised above the level of the retort, and passing the other end through a cork fitted into the neck of the retort. When the whole of the spirit has been added, and forty-two

fluid ounces of impure ether have distilled over, the process may be stopped.

Dissolve the chloride of calcium in the distilled water, add the lime, and agitate the mixture in a bottle with the impure ether ; the chloride of calcium absorbs water, and the lime sulphurous acid. Leave the mixture at rest for ten minutes, pour off the light supernatant fluid, and distil it with a gentle heat until a glass bead of sp. gr. 0·735 placed in the receiver begins to float. The product is ether of the composition already stated.

In the foregoing process the decomposition takes place in two stages :



The ether distils over, and the sulphuric acid remains in the retort.

Characters and Tests.—Colourless, very volatile, and inflammable liquid, emitting a strong and characteristic odour, and boiling below 105°F. Sp. gr. 0·735. Fifty measures shaken with an equal volume of water are reduced to forty-five, by an absorption of 10 per cent. It should evaporate without residue.

Actions and Uses.—*Internally* :—Stimulant, narcotic, and antispasmodic.—*Externally* :—Refrigerant. Inhaled, it acts as an anæsthetic. Used also in the production of local anæsthesia.

Doses.—HORSE, 1 to 2 fluid ounces.—CATTLE, 2 to 3 fluid ounces.—SHEEP, 2 to 4 fluid drachms.—PIG, 2 to 4 fluid drachms.—DOG, $\frac{1}{2}$ to 2 fluid drachms.

Modes of Application.—As a stimulant and antispasmodic, it is given with ten parts cold water ; as a refrigerant, it is applied to the part either alone or mixed with a cooling lotion ; as a general anæsthetic, a sponge saturated with it is placed in the bottom of a nose-bag attached to the patient's head, and perforated so as to ensure the ether vapour being mixed with air. For the production of local anæsthesia, Dr. Richardson's spray-diffuser must be employed.

Preparations.—Collodium ; Collodium Flexile ; Spiritus *Ætheris*.

ALBUMEN OVI

EGG ALBUMEN

Liquid white of the egg of the *Gallus Bankiva*, var. *domesticus*.

Natural Order.—*Gallinæ*.

Composition.—Consists of carbon, hydrogen, oxygen, nitrogen, sulphur, and phosphorus, but the molecular weight has not yet been determined.

Actions and Uses.—As an antidote to corrosive poisons.¹

Doses.—Ad libitum.

Modes of Application.—Before administration it should be whisked, either alone or with water or demulcent drinks.

ALOE BARBADENSIS

BARBADOES ALOES

The inspissated juice of the leaf of *Aloe vulgaris*, belonging to the natural order *Liliaceæ*. Imported from Barbadoes, usually in gourds.

¹ Serum from blood, emulsions of bean or pea meal, or milk, may be used in the absence of egg albumen.

Composition.—Aloin, $C_{17}H_{18}O_7$, mixed with resin and other impurities.

Characters.—Yellowish-brown or dark brown opaque masses ; breaks with a dull conchoidal fracture ; bitter nauseous taste, and a strong disagreeable odour ; dissolves almost entirely in proof spirit, and during solution exhibits under the microscope numerous crystals.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, purgative, nauseant, tonic, diuretic, vermifuge. *Externally* :—Traumatic.

Doses.—*Purgative.*—HORSE, 2 to 8 drachms.—CATTLE, 1 to 2 ounces.—SHEEP, $\frac{1}{2}$ to 1 ounce.—PIG, 1 to 4 drachms.—DOG, 20 grains to 2 drachms.

Modes of Application.—*Internally* :—Massa Aloes, Liquor Aloes, and Liquor Aloes cum Sodæ Carbonate. *Externally* :—As a traumatic, the powder is sprinkled over wounds ; Tinctura Aloes Composita.

Preparations.—Liquor Aloes ; Liquor Aloes cum Sodæ Carbonate ; Massa Aloes ; Tinctura Aloes Composita.

ALOE CAPENSIS

CAPE ALOES ¹

The inspissated juice of the leaf of *Aloe spicata* and several other allied species belonging to the natural order *Liliaceæ*. Imported from the Cape of Good Hope in skins and in chests.

Composition.—Similar to that of Aloe Barbadensis.

¹ Barbadoes aloes is usually preferred to this preparation, as it is considered by most veterinary practitioners to be more certain in its therapeutic action.

Characters.—Glossy, resinous appearance ; dark brown colour, with a strong, greenish-yellow shade, especially when in small fragments ; strong, disagreeable odour, increased by breathing on it ; acrid, bitter taste ; very brittle and readily reduced to powder, which has a shining greenish-yellow colour.

Actions and Uses.

Doses.

Modes of Application.

} Similar to those of *Aloe Bar-*
badensis.

ALOE SOCOTRINA

SOCOTRINE ALOES

The inspissated juice of the leaf of one or more undetermined species of *Aloe*, probably from *Aloe socotrina* and *Aloe purpurascens*, belonging to the natural order *Liliaceæ*. Imported from the island of Socotra, also from Zanzibar. It comes over in skins, casks, kegs, and chests.

Composition.—Similar to *Aloe Barbadensis*.

Characters.—Occurs in masses, varying in colour in different parts of the same mass, sometimes garnet red, at other times much paler, and when quite dry golden red, yielding a golden-yellow powder. The colour is deepened by exposure to air ; fine pieces break with a smooth, glassy, conchoidal fracture, but specimens of good quality often break with a roughish fracture. The taste is very bitter, and the odour of fresh-broken pieces (especially when breathed on) is very fragrant ; it dissolves entirely in proof spirit, and during solution exhibits under the microscope numerous minute crystals.

Actions and Uses.

Doses.

Modes of Application.

} Similar to those of *Aloe Bar-*
badensis.

ALOIN**ALOIN**

Synonym.—Aloine.

Composition.— $C_{17}H_{18}O_7$

This is the principal constituent of aloes, of which it forms about 60 per cent. It is contained in the cold infusion of aloes, and also in a decoction which has cooled ; it may be obtained from either by evaporation. Thus procured it is a brown bitter mass, readily soluble in water but with difficulty in alcohol. Its insolubility in ether distinguishes it from the bitter principle of rhubarb.


Doses.—HORSE, 1 to $1\frac{1}{2}$ drachm.—DOG, 3 to 20 grains.

ALUMEN**ALUM¹**

Synonyms.—Potassium-aluminum Sulphate ; Potassium Alum.


Composition.— $KAl_2SO_4.12Aq$, or $K_2SO_4.Al_2(SO_4)_3.24Aq$.

Mode of Preparation.—There are various modes, but the following is the simplest :—Cornish or pipe clay is boiled with sulphuric acid, so as to form aluminum sulphate and silica.



¹ In the last 'British Pharmacopœia' ammonium alum, $NH_4Al_2SO_4.12Aq$, is substituted for potassium alum. When solution of ammonium alum is mixed with potassium hydrate, ammoniacal gas is evolved and aluminum hydrate is precipitated, which dissolves in excess of the alkali.

The aluminum sulphate, after being separated from the silica, is mixed with solution of potassium sulphate; the two salts unite, and on evaporation crystallise out as alum:



Characters and Tests.—Colourless transparent crystalline masses, exhibiting the faces of the regular octahedron, and having an astringent taste.

Its aqueous solution gives a white gelatinous precipitate (aluminum hydrate) with potassium hydrate soluble in excess of this reagent, indicating the presence of aluminum. A platinum wire moistened with the solution, held in the reducing flame of the blowpipe, communicates a violet colour to the flame, showing the presence of potassium. Barium chloride added to the aqueous solution gives a white precipitate (barium sulphate), insoluble in nitric acid, proving that the compound is a sulphate. Potassium ferrocyanide should not produce either a pale or dark blue colour, showing the absence of iron.

Actions and Uses.—*Internally*:—In excessive doses, irritant poison; in medicinal doses, astringent in diarrhœa, dysentery, and obstinate diabetes; as an antidote to poisoning by lead, and to arrest the secretion of milk in cows. *Externally*:—Astringent to open joints and wounded thecæ; as a collyrium in chronic conjunctivitis; also styptic.

Doses.—HORSE, 2 to 4 drachms.—CATTLE, 2 to 4 drachms.—SHEEP, $\frac{1}{2}$ to 2 drachms.—PIG, $\frac{1}{2}$ to 2 drachms.—DOG, 10 to 30 grains.

Modes of Application.—*Internally*:—Dissolved in water, or in the form of bolus, either alone or in conjunction with opium and carminatives. *Externally*:—Pulvis Aluminis Compositus sprinkled over the wound; otherwise apply Liquor Aluminis, or Unguentum Aluminis Com-

positum, or, in cases of open joints, one part of finely powdered alum with two or three parts of flour may be dusted over the opening.

Incompatibles.—Tannic acid and vegetable astringents containing it ; alkalies, earths, and their carbonates ; acetate of lead.

Preparations.—Alumen Exsiccatum ; Liquor Aluminis Compositus ; Unguentum Aluminis ; Unguentum Aluminis Compositum.

ALUMEN EXSICCATUM

DRIED ALUM

Synonym.—Burned Alum.

Composition.—Alum deprived of its water of crystallisation by heat.

Mode of Preparation.—Heat alum in a porcelain dish till aqueous vapour ceases to be given off, taking care that temperature never exceeds 400°F. Pulverise the residue, and preserve in a well-stoppered bottle.

Actions and Uses.—Caustic and astringent ; employed externally for the same purposes as alum, than which it is more powerful.

Preparation.—Pulvis Aluminis Compositus.

AMMONIACUM

AMMONIACUM

Synonym.—Gum Ammoniacum.

A gum-resinous exudation from *Dorema Ammoniacum*. Collected in Persia and the Punjaub.

Natural Order.—*Umbelliferae*.

Composition.— $C_{40}H_{30}O_9$. When distilled yields umbelliferone $C_9H_6O_3$.

Characters.—In tears or masses; the tears from one-fifth to four-fifths of an inch in diameter, pale cinnamon-brown, breaking with a smooth, shining, opaque, white surface; the masses composed of agglutinated tears, hard and brittle when cold, but readily softening with heat. Faint odour, and a bitter, acrid, nauseous taste. Rubbed with water, it forms a nearly white emulsion.

Actions and Uses.—*Internally*:—Stimulant and antispasmodic in coughs, spasmodic nervous diseases, *e.g.* chorea, epilepsy, &c.; also vermifuge.

Doses.—HORSE, 2 to 4 drachms.—CATTLE, 2 to 4 drachms.—SHEEP, $\frac{1}{2}$ to $1\frac{1}{2}$ drachm.—PIG, $\frac{1}{2}$ to $1\frac{1}{2}$ drachm.—DOG, 10 to 20 grains.

Modes of Application.—Administered in the form of bolus or emulsion. As a vermifuge it may be given either by the mouth or the rectum.

AMMONIÆ CARBONAS


CARBONATE OF AMMONIA

Synonym.—Commercial Carbonate of Ammonia.

Composition.—Probably $2NH_4HCO_3 + NH_4NH_2CO_2$, *i.e.* a mixture or compound of two molecules of ammonium-hydrogen carbonate and one molecule of ammonium carbonate.

Preparation.—By heating to redness a mixture of one part of ammonium chloride and two parts of calcium carbonate (chalk), in an iron or earthenware retort, to which is luted an earthenware or leaden receiver. When the receiver has become filled with the 'carbonate of ammonia'

by repeated distillations, it is broken or cut in two, and its contents removed.




The ammoniacal gas and water are first disengaged, then the ‘carbonate of ammonia’ distils over, and the calcium chloride remains in the retort.

Characters and Tests.—Translucent crystalline masses, with a strong ammoniacal odour¹ and alkaline reaction; soluble in cold water,² sparingly so in spirit of wine. Volatilised entirely by heat, and dissolved by acids with effervescence. If dissolved in slight excess of dilute nitric acid and boiled, the solution should give no white precipitate with silver nitrate, showing the absence of chlorine, and no blue colour with potassium ferrocyanide, proving the absence of iron.

Actions and Uses.—*Internally*:—In excessive doses, irritant and narcotic poison; in medicinal doses, stimulant, antacid, and resolvent. Given in influenza, scarlatina,

¹ The odour is due to the escape of ammoniacal gas. By exposure to air the preparation also loses carbonic anhydride and becomes ammonium-hydrogen carbonate, which is opaque, odourless, and antacid, but not stimulant; hence, ‘carbonate of ammonia’ should be preserved in well-stoppered bottles.

² When commercial or pharmaceutical ‘carbonate of ammonium’ is dissolved in water, the ammonium carbamate which it contains is soon transformed into normal ammonium carbonate; thus:



erysipelas, and other typhoid affections; also in inflammation of the lungs, pleurisy, and similar complaints.

Doses.—HORSE, 1 to 2 drachms.—CATTLE, 2 to 4 drachms.—SHEEP, 20 grains to 1 drachm.—PIG, 20 grains to 1 drachm.—DOG, 3 to 10 grains.

Modes of Application.—In a bolus, pill, or in cold gruel.

Incompatibles.—Acids; alkalies; lime-water; and most metallic salts.

Antidotes.—Copious draughts of water or mucilaginous drinks, oil, or highly diluted acids.

Preparations.—Liquor Ammoniae Acetatis; Spiritus Ammoniae Aromaticus.

AMMONII CHLORIDUM

CHLORIDE OF AMMONIUM

Synonyms.—Ammonium Chloride; Hydrochlorate of Ammonia; Muriate of Ammonia; Sal Ammoniac.

Composition.— NH_4Cl .

Preparation.—‘Gas liquor,’ which is water holding in solution carbonate and other salts of ammonium, is acidulated with hydrochloric acid and evaporated to dryness. The crude ammonium chloride, constituting the residue, is purified by sublimation.

Characters and Tests.—Colourless, odourless, translucent, fibrous masses, tough and difficult to powder; soluble in water and in rectified spirit. Its aqueous solution, heated with potassium hydrate, evolves ammoniacal gas, which may be recognised by its odour; with silver nitrate it gives a white curdy precipitate (silver chloride), insoluble in boiling nitric acid, soluble in ammonia.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses,¹ stimulant, resolvent, diuretic. Given in mucous fevers after subsidence of acute inflammatory symptoms ; in the milder forms of pneumonia and inflammations of serous membranes ; in mucus diarrhœa, chronic rheumatism, and passive dropsies. *Externally* :—Stimulant to bruises, sprains, and inflammatory swellings.

Doses.—HORSE, $\frac{1}{2}$ to $1\frac{1}{2}$ ounce.—CATTLE, $\frac{1}{2}$ to $1\frac{1}{2}$ ounce.—SHEEP, $\frac{1}{2}$ to 2 drachms.—PIG, $\frac{1}{2}$ to 2 drachms.—DOG, 5 to 20 grains.

Modes of Application.—*Internally* :—Bolus, pill, or dissolved in water. *Externally* :—Lotio Ammonii Chloridi et Camphoræ ; Lotio Ammonii Chloridi et Potassæ Nitratis.

Incompatibles.—Sulphuric and nitric acids ; potash, soda, lime, and their carbonates ; and most metallic salts.

Antidotes.—Large draughts of water ; and, when possible, vomiting should be promoted by warm demulcent and mucilaginous drinks.

Preparations.—Lotio Ammonii Chloridi et Camphoræ ; Lotio Ammonii Chloridi et Potassæ Nitratis.

AMYLUM

STARCH

Synonym.—Fecula.

Composition.— $C_6H_{10}O_5$, or more probably $C_{18}H_{30}O_{15}$.

Mode of Preparation.—Chiefly obtained by grinding wheat or rice, or rasping potatoes, making the product into

¹ Seldom employed internally in this country, but on the Continent extensively used as a resolvent in inflammatory diseases, it is said, with the greatest advantage.

a pulp with water, and washing it in a sieve. The starch passes through the sieve, subsides, and is collected and dried at a gentle heat.

Characters and Tests.—White columnar masses. Rubbed in a mortar with a little cold water, it should be neither acid nor alkaline to test-paper. Mixed with boiling water and cooled, it gives a deep blue colour (starch iodide) with solution or tincture of iodine.

Actions and Uses.—*Internally* :—Demulcent and emollient, in the form of gruel. *Externally* :—Applied in the state of dry powder, it diminishes the discharge from wounds, and, mixed with about one-eighth of its weight of alum, it is employed to arrest the flow of synovia from open joints. Also used in the preparation of starch bandages.

AMYL NITRIS

NITRITE OF AMYL

Synonym.—Amyl Nitrite.

Composition.— $C_5H_{11}NO_2$.

Mode of Preparation.—By passing nitrous acid into amyl alcohol.

Characters and Tests.—Nitrite of amyl is a yellowish ethereal liquid ; sp. gr. of liquid, 0·877 ; boiling-point, 205°F. ; soluble in alcohol, insoluble in water ; converted by fused potassium hydrate into valerianate of potassium ($KC_5H_9O_2$).

Actions and Uses.—Sedative, antispasmodic, and anæsthetic. Used in the treatment of tetanus.

Doses.—As an anæsthetic the vapour of 1 to 3 drachms for the HORSE ; 10 to 20 minims for the DOG.

ANISI FRUCTUS**ANISE FRUIT**

Synonym.—Aniseed.

The fruit of *Pimpinella Anisum*. Cultivated in China, Japan, and Tartary.

Natural Order.—*Umbelliferæ*.

Composition.—The active principle is a volatile oil.¹

Characters.—Ovoid ; composed of two mericarps with five primary ridges ; slightly hairy ; yellowish-brown colour, peculiar sweet aromatic odour, and a warm sweetish taste.

Actions and Uses.—Stomachic, carminative, aromatic. Used internally in indigestion and flatulency, to mask the disagreeable flavour of medicines, and to diminish the nauseating and griping effects of purgatives.

Doses.—HORSE, $\frac{1}{2}$ to 2 ounces.—CATTLE, 1 to 2 ounces.—SHEEP, 2 to 3 drachms.—PIG, 2 to 3 drachms.—DOG, 20 grains to 1 drachm.

Modes of Application.—The powdered fruit is administered alone, or in conjunction with ginger or some other aromatic, in ale or in spirit and water.

ANTHEMIDIS FLORES**CHAMOMILE FLOWERS**

The dried single and double flower-heads of the common chamomile, *Anthemis nobilis*. Found in a wild state in all the temperate parts of Europe. Cultivated at Mitcham, Derbyshire, and other parts of England.

¹ The volatile oil of anise, Oleum Anisi, is commonly added to masses as a flavouring ingredient, and is likewise employed to destroy pediculi on dogs and other small animals.

Natural Order.—*Compositæ.*

Composition.—The active principles are a bitter substance of unknown composition and a volatile oil.

Characters.—The single variety consists of both yellow tubular and white strap-shaped florets, the double of white strap-shaped florets only ; all arising from a conical scaly receptacle. Both varieties, but especially the single, are bitter and very aromatic.

Actions and Uses.—*Internally* :—Tonic and carminative in indigestion, and during convalescence after inflammatory attacks.

Doses.—HORSE, 2 to 6 drachms.—CATTLE, 2 to 6 drachms.—SHEEP, $\frac{1}{2}$ to 2 drachms.—PIG, $\frac{1}{2}$ to 2 drachms.—DOG, 10 to 60 grains.

Mode of Application.—The dried flowers powdered, in combination with ginger or some other aromatic, once or twice a day.

Incompatibles.—Mineral acids and most metallic salts.

ANTIMONII OXIDUM

OXIDE OF ANTIMONY

Synonyms.—Antimonious Oxide ; Antimony Trioxide ; Teroxide of Antimony.

Composition.— Sb_2O_3 .


Mode of Preparation.—Take of

Solution of Chloride of Antimony .	16 fluid ounces.
Carbonate of Soda	6 ounces.
Common Water	2 gallons.
Distilled Water	A sufficiency.

Pour the chloride of antimony into the common water, and mix thoroughly ; let the precipitate (antimony oxy-

chloride) which forms settle, and, after removing the supernatant liquid by a siphon, add to it one gallon of distilled water, agitate well, let the precipitate subside ; again withdraw the fluid and repeat the processes of affusion of distilled water, agitation, and subsidence. Now add the carbonate of soda, previously dissolved in two pints of distilled water ; leave them in contact for half an hour, stirring frequently ; collect the deposit on a calico filter, and wash it with boiling distilled water until the washings, after being acidulated by nitric acid, cease to give a precipitate with solution of silver nitrate. Dry the product (oxide of antimony) at a temperature not exceeding 212°F.

The oxide of antimony thus prepared is formed in two stages :



Characters and Tests.—Greyish-white powder, fusible at a low red heat ; insoluble in water, but readily dissolved by hydrochloric acid. The hydrochloric solution, dropped into distilled water, gives a white deposit (antimony oxychloride), which is immediately changed to orange (antimonious sulphide) by sulphuretted hydrogen. Should entirely dissolve when boiled with water and excess of potassium-hydrogen tartrate.

Actions and Uses.—Diaphoretic and febrifuge for the dog. Chiefly used in the preparation of tartarated antimony.

Preparations.—Pulvis Antimonialis ; Antimonium Tartaratum.

ANTIMONIUM NIGRUM**BLACK ANTIMONY**

Synonyms.—Antimonious Sulphide ; Antimony Trisulphide ; Tersulphide of Antimony.

Composition.— Sb_2S_3 .

Mode of Preparation.—Native antimonious sulphide is fused in a perforated crucible placed over another ; the siliceous impurities are retained by the upper crucible, and the molten sulphide which flows into the lower one is solidified by cooling, and reduced to a fine powder.

Characters and Tests.—Greyish-black crystalline powder. It dissolves entirely in boiling hydrochloric acid, evolving sulphuretted hydrogen and furnishing a solution (antimonious chloride) which throws down a white precipitate (antimony oxychloride) when poured into water.

Actions and Uses.—Alterative, anthelmintic, and emetic. Its action is uncertain and irregular, and it should, therefore, not be employed as a remedial agent. Frequently given in combination with sulphur and nitrate of potash as an alterative to horses. Also used in the preparation of *Liquor Antimonii Chloridi*.

ANTIMONIUM TARTARATUM**TARTARATED ANTIMONY**


Synonyms.—Potassium-Antimony Tartrate ; Tartrate of Potassium and Antimony ; Tartar Emetic ; Emetic Tartar ; Tartarised Antimony.

Composition.— $\text{K}(\text{SbO})\text{C}_4\text{H}_4\text{O}_6 \cdot \text{H}_2\text{O}$.

Mode of Preparation.—Take of

Oxide of Antimony	5 ounces.
Acid Tartrate of Potash, in fine powder	} . 6 ounces.
Distilled Water	
	2 pints.

Mix the oxide of antimony and acid tartrate of potash with sufficient distilled water to form a paste, and set aside for twenty-four hours ; then add the remainder of the water, and boil for a quarter of an hour, stirring frequently. Filter, and set aside the clear liquid which contains the tartarated antimony, to crystallise. Pour off the mother liquor, evaporate to one-third, and set aside, that more crystals may form. Dry the crystals (tartarated antimony) on filtering paper at the temperature of the air.



Characters and Tests.—Colourless, transparent crystals, exhibiting triangular facets, soluble in water, and less so in proof spirit. Decrepitates, blackens, and emits an odour characteristic of tartrates when heated on platinum foil. Its solution in water gives, with hydrochloric acid, a white precipitate (antimony oxychloride) soluble in excess ; if tartaric acid be previously added, no precipitate is formed by hydrochloric acid.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in small doses it promotes the secretion of mucous membranes, skin, liver, pancreas, and kidneys, likewise stimulates the activity of the absorbent system ; in larger doses it causes nausea, relaxes muscular fibres depresses the nervous system, and, in animals capable of vomition, acts as an emetic. Used internally as an alterative, antiphlogistic, nauseant, sedative, and emetic. Given

in the early stages of inflammatory diseases, in pneumonia, bronchitis, articular rheumatism, &c.; also employed as a vermifuge. *Externally*:—Vesicant, counter-irritant. Sometimes added to ordinary blistering ointments to increase their activity.

Doses.—*Alterative*:—HORSE, $\frac{1}{2}$ to 1 drachm.—CATTLE, $\frac{1}{2}$ to 1 drachm.

Antiphlogistic, Nauseant, Sedative:—HORSE, 1 to 2 drachms.—CATTLE, 2 to 6 drachms.—DOG, $\frac{1}{8}$ to 2 grains.

Emetic:—PIG, 4 to 16 grains.—DOG, 1 to 4 grains.

Vermifuge:—HORSE, 1 to 2 drachms.¹

Modes of Application.—In the form of bolus, pill, or dissolved in water. As an emetic to dogs it may be given rolled up in a piece of meat.

Externally:—Liquor Antimonii Tartarati; Unguentum Antimonii Tartarati.

Incompatibles.—Gallic and tannic acids; alkalies; and lead salts.

Antidotes.—Tannic acid, or agents containing it, *e.g.* powdered gall-nuts, catechu, &c.

Preparations.—Liquor Antimonii Tartarati; Pulvis Antimonialis; Unguentum Antimonii Tartarati.

ANTIPYRIN

DI-METHYL-OXY-QUINIZINE

Composition.— $C_{11}H_{12}N_2O$.

Preparation.—A synthetical product, belonging to the Aniline series.

Characters and Tests.—A white crystalline powder, of a slightly bitter taste, freely soluble in water, alcohol and

¹ Should be given at night and followed by an aloetic purge next morning.

chloroform, less readily soluble in ether. Ferric chloride gives a dark brown colour. Spirits of nitrous ether give a green colour.

Actions and Uses.—Antipyretic ; it causes a decided lowering of temperature in fever. Anodyne in neuralgic and rheumatic affections. Used sometimes in chorea.

Doses.—Dogs, 15 to 30 grains.

AQUA DESTILLATA

DISTILLED WATER

Composition.— H_2O .

Mode of Preparation.—Take of

Common Water	.	.	.	10 gallons.
--------------	---	---	---	-------------

Distil from a copper still, connected with a block-tin worm ; reject the first half-gallon, and preserve the next eight gallons.

Tests.—A fluid ounce evaporated in a clean glass capsule should leave scarcely any visible residue, indicating the absence of solid non-volatile impurities. Should not be affected by hydrosulphuric acid, ammonium oxalate, silver nitrate, barium chloride, or lime water, proving the absence of lead and other metals precipitable by sulphuretted hydrogen, calcium salts, chlorides, sulphates, carbonates, and carbonic acid respectively.

Uses.—In the preparation of, and for dissolving, many medicinal agents ; also for making all aqueous solutions of tests. When distilled water cannot be obtained, its best substitute is boiled and filtered rain water.

AQUA MENTHÆ PIPERITÆ**PEPPERMINT WATER**

Mode of Preparation.—Take of

Oil of Peppermint $1\frac{1}{2}$ fluid drachm.

Water $1\frac{1}{2}$ gallon.

Distil one gallon.

Actions and Uses.—Carminative and stimulant. Used to relieve flatulence, and as a vehicle for other medicines.

Doses.—HORSE, 8 to 10 fluid ounces.—CATTLE, 8 to 12 fluid ounces.—DOG, 1 to 2 fluid ounces.

ARECÆ SEMINA**ARECA SEEDS**

Synonyms.—Nux Areca Catechu ; Areca Nut ; Betel Nut.

The seeds or kernels of the fruit of the catechu or betel-nut palm, *Areca Catechu*.

Natural Order.—*Palmaceæ*.

Composition.—Their medicinal properties depend upon tannic and gallic acids.

Characters.—Spheroidal, about three-quarters of an inch in diameter ; hard ; colour, reddish brown ; taste, astringent.

Actions and Uses.—*Internally* :—Astringent ; employed as an anthelmintic, especially for dogs.

Doses.—HORSE, 4 to 6 drachms.—CATTLE, 4 to 8 drachms.—DOG, 30 grains to 2 drachms.

Modes of Application.—The powdered seeds made into a bolus, or suspended in milk or gruel.

Incompatibles.—Same as Acidum Tannicum.

ARGENTI NITRAS

NITRATE OF SILVER


Synonyms.—Silver Nitrate ; Lunar Caustic.

Composition.— AgNO_3 .

Mode of Preparation.—Take of

Purified Silver	.	.	.	3 ounces.
Nitric Acid	.	.	.	$2\frac{1}{2}$ fluid ounces.
Distilled Water	.	.	.	3 fluid ounces.

Put the silver into a flask, pour on to it the nitric acid and water, and apply a gentle heat until the metal is dissolved. Decant the clear liquid from any black powder (gold) which may be present into a porcelain dish, evaporate, and set aside to crystallise ; pour off the liquor, and again evaporate and crystallise. Let the crystals (nitrate of silver) drain in a glass funnel, and dry them by exposure to the air, carefully avoiding the contact of all organic substances. To obtain the nitrate in rods, fuse the crystals in a platinum or thin porcelain capsule, and pour the melted salt into proper moulds. Nitrate of silver should be preserved in carefully stoppered bottles.



Characters and Tests.—Colourless tubular crystals, the primal form of which is the right rhombic prism ; or in white cylindrical rods. Soluble in water and rectified spirit. A fragment heated on charcoal with the blowpipe melts,

deflagrates, and leaves a white metallic coating of silver. Its aqueous solution, mixed with excess of dilute hydrochloric acid, yields a white precipitate (silver chloride), which darkens by exposure to light, and which should entirely dissolve in ammonia. If the white precipitate, with its accompanying liquid, be thrown on a filter and thoroughly washed with boiling distilled water, the filtrate should leave no residue on evaporation, proving the absence of potassium nitrate, lead, zinc, and copper.

Actions and Uses.—*Internally* :—In excessive doses, irritant and corrosive poison ; in medicinal doses, tonic and antispasmodic. It is sometimes given to the dog in chorea, epilepsy, and other nervous affections ; and has also been found of service to the dog in diarrhœa, dysentery, and cholera. *Externally* :—Stimulant, astringent, and caustic. Used as a collyrium in conjunctivitis and similiar superficial inflammatory complaints ; for the improvement of indolent sores, mange, ringworm, and other chronic skin diseases ; as a caustic for the eradication of warts, fungous and other growths. Applied also to poisoned wounds, and to the bites of rabid and venomous animals.

Doses.—Dog :—In chorea, epilepsy, &c., $\frac{1}{8}$ to $\frac{1}{2}$ grain ; in diarrhœa, dysentery, and cholera, $\frac{1}{4}$ to 1 grain.

These doses should be repeated two or three times a day.

Modes of Application :—*Internally* :—In the form of bolus or pill. *Externally* :—In the form of stick ; Liquor Argenti Nitratis ; Unguentum Argenti Nitratis.

Incompatibles.—Sulphuric, hydrochloric, and hydrocyanic acids ; alkalies ; and soluble chlorides.

Antidotes.—Solution of common salt in some demulcent drink.

Preparations.—Liquor Argenti Nitratis ; Unguentum Argenti Nitratis.

ARNICÆ RADIX**ARNICA ROOT**

The dried rhizome and rootlets of *Arnica montana*, or Leopard's Bane. Collected in the mountains of middle and southern Europe.

Natural Order.—*Compositæ*.

Composition.—The most important, because probably the active, constituents are resin containing cytosin and arnicin and volatile oil.

Characters.—Rhizome from one to three inches long, and from one-tenth to three-tenths of an inch thick, cylindrical, contorted, rough from the scars of the coriaceous leaves, and furnished with numerous long, slender fibres; has a peppery taste and peculiar odour.

Preparation.—*Tinctura Arnicæ*.

ASSAFŒTIDA**ASSAFŒTIDA**

A gum resin obtained by incision from the living root of *Narthex Assafoetida*.

Natural Order.—*Umbelliferae*.

Composition.—Its active constituents are resinous matter and a volatile oil containing allyl sulphide $(C_3H_5)_2S$.

Characters.—Irregular masses, partly composed of tears, moist or dry. The colour of a freshly cut or broken piece is opaque white, but gradually becomes purplish pink, and ultimately dull yellowish or pinkish brown. Taste, bitter, acrid; odour (due to allyl sulphide), fetid, alliaceous, and

persistent. Dissolves almost entirely in rectified spirit. Powdered with difficulty, unless triturated with carbonate of potash.

Actions and Uses.—A moderate stimulant, powerful antispasmodic, expectorant, and vermifuge. Chiefly used in treating colic and coughs. Given also to dogs suffering from chorea and other nervous affections.

Doses.	}	Similar to Ammoniacum.
Modes of Application.		

Preparations.—Enema Assafœtidæ ; Spiritus Ammoniæ foetidus.

ATROPINA

ATROPINE

Composition.— $C_{17}H_{23}NO_3$.

An extremely poisonous crystallisable alkaloid, obtained from belladonna root, and constituting the active principle of Belladonnæ Folia, Belladonnæ Radix, Extractum Belladonnæ, and Tinctura Belladonnæ.

Preparations.—Atropine sulphate. The B. P. Liquor atropinæ sulphatis consists of one part atropine sulphate and ninety-nine of camphor water ; it contains four grains in one fluid ounce.

Doses of the Liquor.—HORSES and CATTLE, from 1 to 3 fluid drachms ; DOGS, 1 to 3 minims.

One-fifth of these doses suffices when used hypodermically. Equal proportions of atropine sulphate and acetate of morphine solutions give prompt antispasmodic and anodyne effects when used conjointly.

Ointment.—Atropine dissolved in three and a half parts rectified spirit and fifty-five parts benzoated lard.

BALSAMUM PERUVIANUM**BALSAM OF PERU**

A balsam¹ obtained from *Myroxylon Pereiræ*, belonging to the natural order *Leguminosæ*. It exudes from the trunk of the tree after the bark has been scorched and removed. From Salvador in Central America.

Composition.—A mixture of volatile oil called cinnamoin, $C_{27}H_{26}O_4$, cinnamic acid, $HC_9H_7O_2$, and resinous compounds.

Characters.—Reddish-brown or nearly black liquid, translucent in thin films; syrupy consistence, balsamic odour, and an acrid slightly bitter taste; soluble in five parts of rectified spirit. Undergoes no diminution when mixed with water.

Actions and Uses.—Stimulant and antispasmodic. Used in the preparation of *Tinctura Benzoini composita* (Friar's Balsam).

BELLADONNÆ FOLIA**BELLADONNA LEAVES**

The fresh leaves, with the branches to which they are attached, of Deadly Nightshade, *Atropia Belladonna*; also the leaves separated from the branches, and carefully dried and finely powdered; gathered from wild or cultivated British plants when the fruit has begun to form.

Natural Order.—*Solanaceæ*.

Characters.—Leaves alternate, three to six inches long, ovate, acute, entire, smooth, the uppermost in pairs, and

¹ A balsam is a natural mixture of essential oil and resinous substances.

unequal. The expressed juice, or an infusion, dropped into the eye, dilates the pupil.

Actions and Uses.—*Internally* :—In excessive doses, a narcotic-acrid poison ; in medicinal doses, sedative, anodyne, and antispasmodic. In whatever doses it is administered, or by whatever channel it enters the circulation, it causes dilatation of the pupil. Given in colic, acute and chronic rheumatism, bronchitis, influenza, and in cases of sore throat and cough which often accompany or succeed this last-named disease.

Externally :—As a dressing for painful and irritable tumours ; for tender enlarged glands ; for cases of garget and sore throat ; and, in the form of injection, for allaying irritation of the bladder or rectum, and counteracting spasmodic contractions of the uterus.

Doses.¹—HORSE, 1 to 3 ounces.—CATTLE, 1 to 3 ounces.—SHEEP, 10 to 20 grains.—PIG, 8 to 12 grains.—DOG, 2 to 5 grains.

Modes of Application.—*Internally* :—In the form of bolus or pill. See also Extractum Belladonnæ ; Tinctura Belladonnæ ; Massa Belladonnæ composita. *Externally* :—Emplastrum Belladonnæ ; Unguentum Belladonnæ. As an injection, see Extractum Belladonnæ.

Antidotes.—An emetic of sulphate of copper, when practicable ; purgatives to empty the alimentary canal ; after vomition or purgation, administer opium in sufficient doses to counteract the effects of the belladonna. Move the animal about ; affusions of cold water over the head and chest ; artificial respiration ; galvanism ; ether ; brandy ; inhalations of ammonia.

Preparations.—Extractum Belladonnæ ; Tinctura Belladonnæ.

¹ Although the doses of the dried and powdered leaves are here given, Extractum Belladonnæ or Tinctura Belladonnæ is to be preferred for internal use.

BELLADONNÆ RADIX**BELLADONNA ROOT**

The dried root of *Atropa Belladonna*, or Deadly Nightshade. Cultivated in Britain or imported from Germany.

Natural Order.—*Solanaceæ*.

Characters.—From one to two feet long, and from half an inch to two inches thick ; branched and wrinkled, brownish white. An infusion dilates the pupil.

Preparations.—Atropia, and Linimentum Belladonnæ.

BENZOINUM**BENZOIN**

Synonym.—Gum Benzoin.

A balsamic resin, obtained from *Styrax Benzoin*, belonging to the natural order *Styracææ*, by making incisions in the bark of the tree, and allowing the liquid that exudes to concrete by exposure to the air. Imported from Siam and Sumatra.

Composition.—Contains resins, benzoic acid ($\text{HC}_7\text{H}_5\text{O}_2$), volatile oil, and aromatic extract.

Characters.—Lumps consisting of agglutinated tears, or a brownish mottled mass with or without tears imbedded in it ; has little taste, but an agreeable odour ; gives off, when heated, fumes of benzoic acid ; soluble in rectified spirit and in solution of potash.

Actions and Uses.—*Internally* :—Stimulant and antispasmodic ; but seldom or never given at the present day. *Externally* :—Excitant to wounds and contusions in the form of Tinctura Benzoini composita (Friar's Balsam).

BOLI**BALLS**


See Massæ (Masses).

BORAX**BORAX**

Synonyms.—Sodium Diborate ; Biborate of Soda.

Composition.— $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{Aq.}$ Found native in the waters of certain lakes in Thibet and Persia. Also imported in the crude state from the East Indies under the name of tincal.

Mode of Preparation.—Jets of steam (*fumerolles* or *suffioni*) charged with boracic acid, HBO_3 , which issue from the earth in Tuscany, are conducted into *lagoni* (little lakes), in which the acid condenses. The solution thus formed, after being concentrated by heat, is neutralised with sodium carbonate, and, on evaporation, yields crystals of borax.



In the act of crystallising, the sodium diborate unites with ten molecules of water, and becomes $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{Aq.}$

Characters and Uses.—Transparent colourless crystals, sometimes slightly effloresced, with a weak alkaline reaction ; insoluble in rectified spirit. Solubility in water, 1 in 22 ; boiling water, 1 in 2. By the aid of 1 of glycerin, 1 part of borax will dissolve in 12 of water. A hot saturated solution, when acidulated with any of the mineral acids,

lets fall, as it cools, a scaly crystalline deposit of boracic acid, HBO_3 , the solution of which in spirits of wine burns with a green flame.

Actions and Uses.—*Externally*:—Detergent and astringent in aphthous affections of the mouth.

Modes of Application.—A mixture of one part of finely powdered borax and six parts of honey or glycerin is applied to the part affected.

BUCHU FOLIA

BUCHU LEAVES

The dried leaves of *Barosma betulina*, *B. crenulata*, and *B. serratifolia*.

Natural Order.—*Rutaceæ*.

Characters.—The leaves are smooth, dull yellow-green, with a strong penetrating odour, a bitter, aromatic taste, and varying in different species from half an inch to an inch and a half in length. They contain a volatile oil, which is the most active constituent. A camphoraceous substance, barosma camphor, is deposited in the oil in the cold. They also contain mucilage.

Actions and Uses.—A mild tonic, diuretic, diaphoretic, but the principal use of this agent is for disorders of a catarrhal character affecting mucous membranes and diseases of the genito-urinary organs.

Doses.—HORSES or CATTLE, 1 to 4 ounces ; for DOGS, 10 to 30 grains infused in a covered vessel with 20 parts of water for half an hour ; this infusion is given with linseed tea or barley water. It may be advantageously combined with belladonna, opium, hyoscyamus, or potassium bromide.

CALCII CHLORIDUM


CHLORIDE OF CALCIUM

Synonym.—Calcium Chloride.


Composition.— CaCl_2 .

Mode of Preparation.—By neutralising hydrochloric acid with marble (calcium carbonate), adding a little solution of calcium hypochlorite and calcium hydrate to the solution, filtering, evaporating until it becomes solid, and finally drying the salt at about 400°F .


When calcium carbonate dissolves in hydrochloric acid, calcium chloride, water, and carbonic anhydride are produced, thus :



Any iron that the marble may contain in the state of carbonate passes into solution as ferrous chloride, and is converted into ferric chloride and ferric hydrate by the calcium hypochlorite.



The calcium hydrate transforms the ferric chloride produced in the last reaction into ferric hydrate.



In this way the iron is precipitated, and by filtration it is completely removed.

Characters and Tests.—White agglutinated masses, dry, but very deliquescent. If hydrochloric acid be poured on it, no chlorine or hypochlorous acid should be evolved, proving the absence of calcium hypochlorite. Should dissolve entirely in water and alcohol. Lime water added to its aqueous solution should give no precipitate, indicating its freedom from iron.

Uses.—In consequence of its strong affinity for water, it is employed to withdraw this impurity from ether, and various other liquids, as well as from gases.

CALCIS HYDRAS

SLAKED LIME

Synonyms.—Calcium Hydrate ; Hydrate of Lime.

Composition.— $\text{Ca}(\text{HO})_2$.


Preparation.—Take of

Freshly burned Lime	.	.	.	2 pounds.
Distilled Water ¹	.	.	.	1 pint.

Place the lime in a metal or earthenware vessel furnished with a lid ; pour the water upon it, and when vapour ceases to be disengaged, cover the vessel with its lid, and set it aside to cool. When the temperature has fallen to that of the atmosphere, put the slaked lime on an iron-wire sieve, and, by gentle agitation, cause the fine powder to pass through the sieve, and reject what is left. Put the powder into a well-stoppered bottle, and keep it excluded as much as possible from the air, to prevent its absorbing carbonic

¹ Although distilled water is directed to be employed in this preparation by the *British Pharmacopæia*, common water may be used in its stead.

anhydride. In the process of slaking, lime unites chemically with water.



Characters and Tests.—A white caustic and strongly alkaline powder. It is slightly soluble in water (1 in 800), more so in cold than hot. Should not effervesce on the addition of an acid, showing its freedom from carbonate.

Actions and Uses.—*Internally* :—Irritant, corrosive, and antacid. Given chiefly to cattle, as an antacid in indigestion, diarrhœa, and tympanitis. Sometimes employed as an antidote to poisoning by arsenic and irritant acids.

Doses.—HORSE, 1 to 3 drachms.—CATTLE, 1 to 3 drachms.—SHEEP, 20 grains to 1 drachm. PIG, 20 grains to 1 drachm.—DOG, 5 to 20 grains.

Modes of Application.—*Internally* :—In the form of bolus or pill, or else suspended in water, milk, or some mucilaginous drink. Also administered as *Liquor Calcis* and *Liquor Calcis Saccharatus*.

Incompatibles.—Mineral and vegetable acids ; metallic and ammoniacal salts ; tartarated antimony.

Preparations.—*Linimentum Calcis* ; *Liquor Calcis* ; *Liquor Calcis Saccharatus* ; *Lotio Hydrargyri flava* ; *Lotio Hydrargyri nigra*.

CALUMBÆ RADIX

CALUMBA ROOT

Synonym.—Calumba Root.

The root of the *Jateorhiza Calumba* and *Cocculus palmatus*, sliced transversely and dried.

Natural Order.—*Menispermaceæ*.

Characters and Tests.—The root consists of several fasciculated, fusiform, fleshy tubercles, which are brown externally and deep yellow internally, odourless, and very bitter. Moistened with a solution of iodine, it becomes blue, indicating presence of starch. A decoction is not blackened by the persalts of iron, indicating absence of astringent matter.

Composition.—The principal constituents of calumba are calumbin ($C_{21}H_{22}O_7$), and the alkaloid berberia ($C_{20}H_{17}NO_4$) combined with calumbic acid ($C_{22}H_{24}O_7$) and starch.

Actions and Uses.—Calumba root is an excellent tonic, promoting the appetite, assisting the digestive process, and improving the quality of the secretions from the gastrointestinal mucous membranes. It does not appear either to constipate or relax the bowels.

Doses.—(Of the powder) HORSE, 2 to 4 drachms.—CATTLE, 2 to 6 drachms.—SHEEP, 1 to 2 drachms.—PIG, $\frac{1}{2}$ to 1 drachm.—DOG, 10 to 20 grains.

Modes of Application.—The root is best administered in the form of powder or infusion.

Preparation.—Infusum Calumbæ.

CALX


LIME

Synonyms.—Calcium Oxide ; Oxide of Calcium ; Quick-lime.

Composition.—CaO.

Mode of Preparation.—Some form of calcium carbonate, *e.g.* chalk, limestone, or marble, is calcined so as to expel

carbonic anhydride ; lime, mixed with a small amount of impurity, remains.



Characters and Tests.—Compact masses of a whitish colour, which readily absorb water, and which, when rather less than their weight of water is added, crack, and fall into powder with the development of much heat. The powder thus obtained, agitated with distilled water and filtered, yields a clear solution, having an alkaline reaction, and giving a white precipitate (calcium oxalate) with ammonium oxalate.

Actions and Uses.—*Externally* :—As a caustic and desiccant, dusted over the part affected.

Preparation.—*Calcis Hydras*.


CALX CHLORATA

CHLORINATED LIME

Synonyms.—Calcium Hypochlorite ; Hypochlorite of Lime ; Chloride of Lime.

Composition.—Not well established. By some authorities it is regarded as lime chloride CaOCl_2 , by others as calcium hypochlorite $\text{Ca}(\text{ClO})_2$; mixed with calcium chloride and variable quantities of calcium hydrate.

Preparation.—Slaked lime is exposed in stone chambers to the action of chlorine gas as long as the latter is absorbed. Assuming that the activity of chlorinated lime depends upon its containing calcium hypochlorite, its production may be thus explained :



Characters and Tests.—Dull white powder, with a feeble odour of chlorine ; partially soluble in water. It evolves chlorine on being mixed with an acid. Its aqueous solution, mixed with a dilute mineral acid, bleaches indigo.

Actions and Uses.—*Internally* :—In tympanites and hoven to absorb the carbonic anhydride and decompose the sulphuretted hydrogen which are developed in these affections. *Externally* :—Disinfectant, deodoriser, and stimulant. Applied to gangrenous wounds, fistulæ, phagedenic ulcers, thrush, canker, grease, and to the treatment of mange and other skin diseases. Also employed as a lotion for checking conjunctival ophthalmia and other circumscribed and superficial inflammations.

Doses.—HORSE, 2 to 4 drachms.—CATTLE, 2 to 4 drachms.—SHEEP, 1 to 2 drachms.

Modes of Application.—*Internally* :—Suspended in water, or, preferably, as Liquor Calcis Chloratæ. *Externally* :—Liquor Calcis Chloratæ ; Unguentum Calcis Chloratæ.

Incompatibles.—Acids and most metallic salts.

Preparations.—Liquor Calcis Chloratæ ; Unguentum Calcis Chloratæ ; Vapor Chlori.

CAMBOGIA

GAMBOGE

A gum-resin obtained from *Garcinia Morella*, var. *pedicellata* belonging to the natural order *Guttiferæ*. Imported from Siam.

Composition.—From sixty-eight to seventy-five per cent. of resin (active principle), $C_{20}H_{23}O_4$, gum, and a minute quantity of woody fibre.

Characters and Tests.—Cylindrical pieces, breaking easily with a smooth, conchoidal, glistening fracture; colour, tawny, changing to yellow when rubbed with water; taste, acrid.

If adulterated with starch, an emulsion, made with boiling water and cooled, becomes green if treated with solution of iodine.

Actions and Uses.—In large doses, irritant poison; in medicinal doses, purgative. Should not be given to horses or dogs, on account of the uncertainty and violence of its action. To cattle and sheep it is sometimes administered in indigestion; also, combined with aloes or sulphate of magnesia, in impaction of the omasum (*fardel bound*) and similar disorders.

Doses.—CATTLE, 4 to 8 drachms.—SHEEP, 15 to 30 grains.

Mode of Application.—Made into an emulsion with water.

Antidotes.—Demulcent drinks, and small, but repeated, doses of opium.

CAMPHORA

CAMPHOR

A concrete volatile oil (stearopten), obtained in China and Japan from the wood and other parts of a species of laurel, *Camphora officinarum*, belonging to the natural order *Lauraceæ*.

Composition.— $C_{10}H_{16}O$.

Mode of Preparation.—The small branches, leaves, wood, and root of the tree are cut into small pieces and boiled with water in an iron vessel, to which an earthen hood is luted; the camphor sublimes and condenses on straws placed within the hood. The crude product thus obtained, after its importation into Europe, is mixed with quicklime to retain impurities, and sublimed in glass vessels.

Characters and Tests.—White, translucent, tough,¹ and crystalline; has a powerful penetrating odour, and a pungent taste, followed by a sensation of cold; floats on water; volatilises slowly at ordinary temperatures; slightly soluble in water, readily soluble in rectified spirit and in ether; sublimes entirely when heated.

Actions and Uses.—*Internally*:—In large doses, irritant and narcotic poison; in medicinal doses, stimulant, sedative, antispasmodic; also, after long exhibition, diuretic. Occasionally given in tympanites, and, combined with opium or digitalis, in chronic cough. *Externally*:—Discutient and anodyne for chronic sprains, bruises, and tumours, also for infiltration into the cellular tissue. Added to oil or ointment of cantharides, or sprinkled in fine powder over linseed poultices, it allays irritation, although for this purpose extract of belladonna is usually preferred.

Doses.—HORSE, 1 to 2 drachms.—CATTLE, 2 to 4 drachms.—SHEEP, 10 to 30 grains.—PIG, 5 to 20 grains.—DOG, 3 to 10 grains.

Modes of Application.—*Internally*:—Spiritus Camphoræ; Oleum Camphoratum, or in the form of ball or pill. *Externally*:—Linimentum Camphoræ; Linimentum Terebinthinæ; Linimentum Terebinthinæ Aceticum; Linimentum Saponis; Tinctura Camphoræ.

¹ Camphor is difficult to pulverise unless a little spirit of wine be added to it.

Incompatibles.—Camphor is not likely to be administered with anything that would counteract its effects.

Preparations.

Linimentum Aconiti.	Linimentum Saponis com-
„ Camphoræ.	positum.
„ Iodi composi-	„ Terebinthinæ.
tum.	Spiritus Camphoræ.
„ Opii	

CANTHARIS

CANTHARIDES

Synonyms.—*Cantharis Vesicatoria* ; *Lytta Vesicatoria* ; Blistering Fly ; Spanish Fly.

Natural Order.—*Coleoptera*.

Composition.—Contains an active principle called cantharidin, $C_{10}H_{12}O_4$.

The beetles are collected, chiefly in Hungary, at night time, by shaking the trees on which they feed, and so causing them to fall into cloths placed beneath the trees for their reception. After being transferred to sieves, they are killed by the vapour of vinegar or turpentine, or by being placed in a vacuum, and finally dried.

Characters and Tests.—From eight-tenths to an inch in length ; furnished with two wing-covers of a shining metallic green colour, under which are two membranous transparent wings ; odour, strong and disagreeable ; powder, greyish brown, containing shining green particles. Should be free from mites.

Actions and Uses.—*Internally* :—In large doses, irritant poison ; in medicinal doses, stimulant, diuretic. With vegetable bitters, sometimes used as a stimulating tonic in cases of debility, accompanied or not with anasarca ; also

in farcy, glanders, and some other affections. As cantharides are liable to inflame the neck of the bladder and cause strangury, they are seldom employed as a diuretic. *Externally* :—Rubefacient, vesicant, counter-irritant, stimulant. Employed as a blister, and for charging materials used for setons ; also to promote the adhesion of unhealthy wounds and fistulæ ; to stimulate weak and callous ulcers ; and to induce an improved condition of skin in mallenders, ringworm, and inveterate mange ; likewise to promote the growth of hair.

Doses.—HORSE, 5 to 10 grains.—CATTLE, 10 to 20 grains.—SHEEP, 2 to 8 grains.—PIG, 1 to 4 grains.—DOG, $\frac{1}{4}$ to 1 grain.

Modes of Application.—*Internally* :—In a bolus, or as Tinctura Cantharidis. *Externally* :—To keep up discharges, also to increase the activity of mustard poultices and similar applications, the powdered insect is used. To promote the growth of hair, a solution of one part of cantharides in twenty parts of acetic acid, or an ointment consisting of one part of cantharides and twenty parts of lard, is to be applied with friction. For the other applications of cantharides, see the ‘Preparations’ enumerated below.

Preparations.

Acetum Cantharidis.	Oleum Cantharidis.
Linimentum Cantharidis.	Tinctura Cantharidis.
Liquor Cantharidis Tere-	„ „ fortior.
binthinatus.	Unguentum Cantharidis.

CAPSICI FRUCTUS

CAPSICUM FRUIT

The dried ripe fruit of *Capsicum fastigiatum*. Imported from Zanzibar, and, in a pulverised state, in gourds from the West Indies. It is distinguished in commerce as

Guinea Pepper and Pod Pepper ; when powdered, and mixed with a fourth part of common salt, it forms the well-known condiment 'Cayenne Pepper.'

Natural Order.—*Solanaceæ*.

Composition.—The fruit contains an alkaloid and a crystallisable solid oil, to both of which the name capsicum has been applied ; the latter body is very acrid, and constitutes the active principle of capsicum.

Characters and Tests.—Pod membranous, from five- to eight-tenths of an inch long, two-tenths of an inch broad ; straight, conical, pointed, smooth, shining, but somewhat corrugated ; orange-red ; intensely hot taste.

Actions and Uses.—*Internally* :—In large doses, irritant poison ; in medicinal doses, stimulant and carminative. *Externally* :—Rubefacient and vesicant.

Doses.—HORSE, 5 to 20 grains.—CATTLE, 10 to 30 grains.—SHEEP, 5 to 10 grains.—PIG, 2 to 8 grains.—DOG, 1 to 5 grains.

Modes of Application.—*Internally* :—The powdered fruit in a bolus, or, preferably, suspended in gruel. *Externally* :—In the form of ointment.

CARBO LIGNI

WOOD CHARCOAL

Composition.—Carbon mixed with from one to five per cent. of ash.

Mode of Preparation.—By exposing various kinds of wood to a red heat without access of air.

Characters.—Black, brittle, porous masses, tasteless, odourless, very light ; retains the shape and texture of the wood from which it was obtained. When burned with free

access of air it should not leave more than two per cent. of ash.

Actions and Uses.—Deodoriser, disinfectant, desiccant. *Internally* :—In dyspepsia, accompanied by flatus, to absorb the gases causing gastric distension ; in diarrhœa and dysentery, to correct the fœtor of the evacuations ; as an antidote to arsenic, aconite, strychnia, and probably to most vegetable poisons.¹ *Externally* :—As a desiccant and deodoriser to unhealthy wounds and phagedenic ulcers, especially when accompanied by fetid and irritating discharges ; also in cases of mange, scab, and grease.

Wood charcoal, particularly when strongly heated and subsequently cooled just before being used, is a valuable deodoriser if strewn over the floors of a stable, &c., or sprinkled over putrefying substances.

Doses.—HORSE, $\frac{1}{2}$ to 1 ounce.—CATTLE, 1 to 2 ounces.—SHEEP, 1 to 3 drachms.—PIG, 1 to 3 drachms.—DOG, 20 to 60 grains.

Modes of Application.—*Internally* :—Pulverised and mixed with gruel or some mucilaginous drink. *Externally* : Dusted over the part in the state of powder, or as Cataplasma Carbonis.

Preparation.—Cataplasma Carbonis.

CARDAMOMUM

CARDAMOMS

The dried capsules of the Malabar cardamom, *Elettaria Cardamomum*. Cultivated in Malabar. The seeds are best kept in their pericarps, from which they should be sepa-

¹ Animal charcoal (*Carbo animalis*) is to be preferred to wood charcoal in consequence of its greater absorptive power for the poisons above referred to.

rated when required for use, the pericarpal coats being rejected.

Natural Order.—*Zingiberaceæ*.

Composition.—The therapeutic activity of cardamoms is due to a volatile oil contained in the seeds.

Characters.—Seeds obtusely angular, corrugated, reddish brown ; internally white, with a warm, aromatic agreeable taste and odour ; contained in ovate-oblong, triangular, pale brown, coriaceous, ribbed pericarps.

Actions and Uses.

Doses.

Modes of Application.

} Same as Anisi Fructus.

CARUI FRUCTUS

CARAWAY FRUIT

Synonym.—Caraway Seeds.

The dried fruit of *Carum Carui*. Cultivated in England and Germany.

Natural Order.—*Umbelliferae*.

Composition.—The active principle is a volatile oil.

Characters.—Fruit usually separating into two parts, which are about two-tenths of an inch long, curved, tapering at each end, brown, with five paler longitudinal ridges ; agreeable aromatic odour and spicy taste.

Actions and Uses.

Doses.

Modes of Application.

} Same as Anisi Fructus.

CARYOPHYLLUM**CLOVES**

The dried unexpanded flower-buds of *Caryophyllus aromaticus*. Cultivated in Penang, Bencoolen, and Amboyna.

Natural Order.—*Myrtaceæ*.

Composition.—The active principle is a volatile oil.

Characters and Test.—About six-tenths of an inch long ; dark reddish brown ; plump and heavy ; consisting of a nearly cylindrical body surmounted by four teeth and a globular head ; strong fragrant odour, and a bitter, spicy, pungent taste. Emits oil when indented with the finger-nail.

Actions and Uses.

Doses.

Modes of Application.

} Same as Anisi Fructus.

Preparation.—Oleum Caryophylli.

CASCARILLÆ CORTEX**CASCARILLA BARK**

The bark of *Croton Eluteria*. From the island of Eluthera, one of the Bahamas.

Natural Order.—*Euphorbiaceæ*.

Composition.—Its therapeutic actions depend upon a crystalline bitter principle, *cascarillin*, and a volatile oil.

Characters.—Quills, two or three inches long, and from two- to five-tenths of an inch in diameter ; dull brown, but more or less coated with white crustaceous lichens ; breaks with a short resinous fracture ; warm and bitter taste, and emits a fragrant odour when burned.

Actions and Uses.—Aromatic tonic and slightly astringent. Sometimes given in indigestion, diarrhœa, chronic typhous affections, and during convalescence from debilitating diseases.

Doses.—HORSE, 2 to 4 drachms.—CATTLE, 3 to 6 drachms.—SHEEP, 1 to 2 drachms.—PIG, 1 to 2 drachms.—DOG, $\frac{1}{4}$ to 1 drachm.

Mode of Application.—In the form of ball.

CATAPLASMA CALCIS CHLORATÆ

CHLORINATED LIME POULTICE

Take of

Solution of Chlorinated Lime	.	2 fluid ounces.
Linseed Meal	.	4 ounces.
Boiling Water	.	8 fluid ounces.

Mix the linseed meal gradually with the water, and add the solution of chlorinated lime, with constant stirring.

Use.—Deodoriser, antiseptic, and excitant to unhealthy wounds and ulcers.

CATAPLASMA CARBONIS

CHARCOAL POULTICE

Take of

Wood Charcoal, in powder	.	$\frac{1}{2}$ ounce.
Linseed Meal	.	$3\frac{1}{2}$ ounces.
Boiling Water.	.	10 fluid ounces.

Add the linseed meal to the water, and stir them together so that a soft poultice may be formed. Mix with this half the charcoal, and sprinkle the remainder on the surface of the poultice.

Use.—Applied while warm to absorb the fœtor of foul ulcers.

CATAPLASMA FERMENTI**YEAST POULTICE**

Take of

Beer Yeast	6 fluid ounces.
Wheaten Flour	14 ounces.
Water heated to 100°F.	6 fluid ounces.

Mix the yeast with the water, and stir in the flour.
Place the mass near the fire till it rises.

Use.—Stimulant and antiseptic to indolent ulcers.

CATAPLASMA FURFURIS**BRAN POULTICE**

Take of

Bran	2 to 3 parts.
Linseed meal	1 part.
Boiling Water	A sufficiency.

Mix the bran and linseed meal together, and then add the water, with constant stirring, so as to form a soft poultice.

Use.—Emollient; relaxes the vessels of the part to which it is applied, and thereby allays pain and inflammation.

CATAPLASMA LINI**LINSEED POULTICE**

Take of

Linseed Meal	4 ounces.
Olive Oil	$\frac{1}{2}$ fluid ounce.
Boiling Water	10 fluid ounces.

Mix the linseed meal gradually with the water, and then add the oil, with constant stirring.

Use.—Same as Cataplasma Furfuris. Opium, belladonna, or subacetate of lead may be added to this poultice when it is desired to allay irritation.

CATAPLASMA SINAPIS

MUSTARD POULTICE

Take of

Mustard, ¹ in powder	2½ ounces.
Linseed Meal	2½ ounces.
Warm ² Water	10 fluid ounces.

Mix the linseed meal gradually with the water, and add the mustard, with constant stirring.

Use.—Rubefacient to allay inflammations of serous and mucous surfaces ; to relieve congestion of various organs, and to alleviate neuralgic and other pains and spasms.

CATAPLASMA SINAPIS AMMONIATUM

AMMONIATED MUSTARD POULTICE

Take of

Mustard, in powder	} of each a sufficiency.
Solution of Ammonia	

Mix so as to form a poultice.³

Use.—Similar to, but much more powerful than, Cataplasma Sinapis.

¹ See note to *Sinapis*.

² Boiling water should not be used, as a temperature of 212°F. prevents the formation of the vesicating principle, allyl sulphocyanide.

³ Oil of turpentine is sometimes added to this poultice, but such addition should be omitted in treating affections of the kidneys.

CATECHU PALLIDUM**PALE CATECHU**

Synonym.—Terra Japonica.

An extract of the leaves and young shoots of *Uncaria Gambir*, belonging to the natural order *Cinchonaceæ*. Prepared at Singapore and in other places in the Eastern Archipelago.

Composition.—Mainly consists of catechin, $C_{13}H_{12}O_5$, and tannic acid.

Mode of Preparation.—The leaves of *Uncaria Gambir* are boiled in water immediately after they are pulled from the tree, and the decoction concentrated and run into square or moulds, to form the commercial cubes of catechu.

Characters.—Cubes, or masses formed of coherent cubes; the former about an inch in diameter, externally brown, internally ochrey yellow or pale brick-red, breaks easily with a dull earthy fracture. Taste, bitter, very astringent, and mucilaginous, succeeded by slight sweetness. Entirely soluble in boiling water. The decoction, when cool, should not be rendered blue by iodine, indicating the absence of starch.

Actions and Uses.—Astringent. *Internally*:—Given in atony and relaxation, and in excessive mucous discharges, especially from the alimentary canal; also in diarrhoea. *Externally*:—To sluggish sores and ulcerations, for excoriation on the udder, &c.

Doses.—HORSE, 2 to 4 drachms.—CATTLE, 2 to 4 drachms.—SHEEP, $\frac{1}{2}$ to 1 drachm.—PIG, 10 to 20 grains.—DOG, 1 to 10 grains. Three or four times a day.

Modes of Application.—*Internally* :—Massa Catechu Composita ; Infusum Catechu ; Tinctura Catechu, or the powder in sufficient gruel or mucilage to mask its astringent flavour. Catechu is likewise given with aromatics to remove flatulency, with opium to allay irritability, and with magnesia or an alkaline carbonate to correct acidity. *Externally* :—The powder ; Infusum Catechu ; and, occasionally, Unguentum Catechu.

Incompatibles.—Alkalies (?) ; most metallic salts ; gelatin.

Preparations.—Infusum Catechu ; Massa Catechu composita. Tinctura Catechu.

CERA FLAVA

YELLOW WAX

Synonym.—Unbleached Beeswax.

A secretion of certain glands (*wax pockets*) situated on the abdomen of the common bee, *Apis mellifica*, and used by the insect for constructing the cells of the honeycomb.

Composition.—A mixture of myricin, cein, and cerolein.

Mode of Preparation.—The comb, after being freed from honey by dripping and pressure, is melted in hot water to free it from impurities ; the residue constitutes yellow wax. White wax (*Cera Alba*) is obtained by agitating molten yellow wax with water, and bleaching in the open air.

Characters.—Firm, breaking with a granular fracture ; yellowish, having an agreeable honeylike odour. Not unctuous to the touch ; does not melt at 140°F. ; yields nothing to rectified spirit, but is entirely soluble in oil of turpentine. Boiling water, in which it has been agitated,

when cooled, should not be rendered blue by iodine, showing its freedom from starch.

Actions and Uses.—Emollient and to increase consistency of ointments.

Preparation.—Emplastrum Picis.

CEREVISIÆ FERMENTUM

BEER YEAST

The ferment obtained in brewing beer.

Characters.—Viscid, semifluid, frothy, exhibiting under the microscope numerous round or oval confervoid cells, *Torula cerevisiæ*.

Preparation.—Cataplasma Fermenti.

CERII OXALAS

OXALATE OF CERIUM

Composition.— CeC_2O_4 .

Mode of Preparation.—By adding oxalic acid to a solution of a cerium salt, cerium oxalate is precipitated.

Characters and Tests.—A white granular powder, odourless and tasteless, insoluble in water and alcohol, soluble in hydrochloric acid.

Actions and Uses.—*Internally*:—It is sometimes given in the treatment of chronic bronchitis, and chorea.

(Chloral Hydrate 3T)



CHLORAL HYDRAS

HYDRATE OF CHLORAL

Synonyms.—Chloral Hydrate ; Trichloraldehyd.

Composition.— $C_2HCl_3O.H_2O$ or $CCl_3.CO.H_2O$.

Mode of Preparation.¹—Dry chlorine gas is passed for several days through absolute alcohol, which is at first kept cool to prevent explosion, but which, towards the end of the operation, is heated to nearly its boiling-point to complete the decomposition. The resulting liquid (impure chloral, C_2HCl_3O) is exposed to the air for a day or two, when it absorbs water and becomes converted into a solid mass of crude chloral hydrate. This is purified by agitating it with four times its bulk of concentrated sulphuric acid, separating the anhydrous chloral which floats on the surface, and subjecting it to fractional distillation. The anhydrous chloral thus purified is placed in a still with eleven per cent. of water, and distilled off chalk to remove any hydrochloric acid that may be present. The solid distillate (chloral hydrate) obtained in this manner is fused and cast in shallow vessels into cakes.



Characters and Tests.—White, opaque, crystalline solid ; pungent odour, resembling that of a ripe melon ; soluble in water, alcohol, and glycerin. One hundred grains dissolved in $\frac{1}{2}$ oz. of distilled water, well shaken with 1 oz. of Liquor Potassæ, and allowed to stand several hours in a

¹ Dr. Squire's process.

well-stoppered bottle, will, if pure, yield a layer of at least 46 grain measures of chloroform.

Actions and Uses.—*Internally* :—In excessive doses, narcotic poison ; in medicinal doses, stimulant, sedative, anodyne, antispasmodic, and narcotic. Said to possess the good, but none of the objectionable, qualities of opium ; successfully used in milk fever in cattle.¹

Doses.—HORSE, $\frac{1}{2}$ to 1 ounce.—CATTLE, 1 to 2 ounces.—SHEEP, 1 to 2 drachms.—PIG, 1 to 2 drachms.—DOG, 10 to 30 grains.

Mode of Application.—Dissolved in a large quantity of water.

Preparation.—Syrupus Chloralis.

CHLOROFORMI MISTURA HYDROCYANATA

COMPOUND CHLOROFORM MIXTURE

Synonym.—Chlorodyne.

Composition.—The two following formulæ may be used in the preparation of chlorodyne :

1. Morphine sulphate 4 grains, oil of peppermint 8 grains, ether 1 drachm, alcohol 1 drachm, hydrocyanic acid (dilute) 5 drachms, chloroform 3 ounces, syrup to 17 ounces.

2. Ten parts each of chloroform, ether, Indian hemp, and morphine ; two parts of tincture of capsicum and prussic acid ; three parts of tinctures of aconite and hyoscyamus ; one part oil of peppermint, five parts of hydrochloric acid, and fifty parts of simple syrup.

Actions and Uses.—Anodyne, antispasmodic, and to relieve gastro-intestinal and bronchial irritation.

Doses.—HORSE, 2 to 4 drachms.—DOG, 20 to 40 minims.

¹ Dr. Take, in speaking of its effects on maniacal patients, says : 'Its advantages over other hypnotics are as follows: it is more uniform in its action and its effects are more lasting; it has no depressing influence; it does not constipate nor produce nausea.'

CHLOROFORMUM

CHLOROFORM

Synonyms.—Formyl Trichloride ; Terchloride of Formyl.

Composition.— CHCl_3 .




Mode of Preparation.—Take of

Chlorinated Lime	10 pounds.
Rectified Spirit	30 fluid ounces.
Slaked Lime	A sufficiency.
Water	3 gallons.
Sulphuric Acid	A sufficiency.
Chloride of Calcium, in small fragments	2 ounces.
Distilled Water	9 fluid ounces.

Place the water and the spirit in a capacious still, and raise the mixture to the temperature of 100°F . Add the chlorinated lime and five pounds of the slaked lime, and mix thoroughly. Connect the still with a condenser terminating in a narrow-necked receiver, and apply heat so as to cause distillation, taking care to withdraw the fire the moment that the process is well established. When the distilled product measures fifty ounces, the receiver is to be withdrawn. Pour its contents into a gallon bottle half filled with water, mix well by shaking, and set at rest for a few minutes, when the mixture will separate into two layers of different densities. Let the lower layer, which constitutes crude chloroform, be washed by agitating it in a bottle with three ounces of the distilled water. Allow the chloroform to subside, withdraw the water, and repeat the washing, with the rest of the distilled water, in successive quantities of three ounces at a time. Agitate the washed chloroform for five minutes in a bottle with an equal

volume of sulphuric acid, allow the mixture to settle, and transfer the upper layer of liquid to a flask containing the chloride of calcium mixed with half an ounce of perfectly dry slaked lime, and mix well by agitation. After the lapse of an hour connect the flask with a Liebig's condenser, and distil over the pure chloroform by means of a water-bath. Preserve the product in a cool place in well-stoppered bottles. The lighter liquid which floats on the crude chloroform after its agitation with water, and the washings with distilled water, should be preserved and employed in a subsequent operation.

The exact nature of the chemical changes which occur in the production of chloroform by the foregoing process is still undetermined, but the following equations represent the probable reactions which take place :



Characters and Tests.—Limpid, colourless liquid, of an agreeable ethereal odour and sweet taste. Soluble in alcohol and ether in all proportions; dissolves slightly in water, communicating to it a sweetish taste. Burns, though not readily, with a green and smoky flame. Sp. gr. 1.49. Should not be coloured by agitation with sulphuric acid, and should leave no residue nor unpleasant odour after evaporation.

Actions and Uses.—In excessive doses, whether the liquid be swallowed or the vapour inhaled, narcotic poison ;

in medicinal doses, stimulant and antispasmodic ; breathed in admixture with air, anæsthetic.

Doses.—See Spiritus Chloroformi.

Modes of Application.—As an antispasmodic: Spiritus Chloroformi. As an anæsthetic, a sponge saturated with pure chloroform is placed in the bottom of a nose-bag attached to the patient's head, and perforated so as to insure the chloroform vapour being mixed with a sufficiency of air.

Antidote.—Fresh air.

Preparations.—Spiritus Chloroformi ; Tinctura Chloroformi Composita.

CINCHONÆ FLAVÆ CORTEX

YELLOW CINCHONA BARK

The bark of *Cinchona Calisaya*. Collected in Bolivia and Southern Peru.

Composition.—Its therapeutic value depends upon the presence of quinine ($C_{20}H_{24}N_2O_2$) and cinchona ($C_{20}H_{24}N_2O$), which, with several other alkaloids, exist in the bark in combination with kinic, kinovic, and tannic acids.

Mode of Preparation.—The trees are generally cut down, and the bark, after removal from the stem and branches, is carefully dried, so that it may retain its bright colour ; the larger and thicker portions are dried so as to form flat pieces, while the smaller are allowed to curl into quills. Unless very coarse and injured, the epidermis, with the lichens which grow on it, is carefully preserved on the bark.

Characters.—Flat pieces, uncoated or deprived of the periderm, rarely in coated quills, from six to eighteen

inches long, one to three inches wide, and two- to four-tenths of an inch thick, compact and heavy ; outer surface brown, marked by broad, shallow, irregular, longitudinal depressions ; inner surface tawny yellow and fibrous ; transverse fracture shortly and finely fibrous. Powder cinnamon brown, somewhat aromatic ; persistently bitter.

Test.—Boil one hundred grains of the finely powdered bark, for a quarter of an hour, in a fluid ounce of distilled water, acidulated with ten minims of hydrochloric acid, and allow it to macerate for twenty-four hours. Transfer the whole to a small percolator, and, after the fluid has ceased to drop, add at intervals about an ounce and a half of similarly acidulated water, or until the fluid which passes through is free from colour. Add to the percolated fluid solution of subacetate of lead until the whole of the colouring matter has been removed, taking care that the fluid remains acid in reaction. Filter and wash with a little distilled water. To the filtrate add about thirty-five grains of caustic potash, or as much as will cause the precipitate which is at first formed to be nearly re-dissolved, and afterwards six fluid drachms of pure ether. Then shake briskly, and, having removed the ether, repeat the process twice with three fluid drachms of ether or until a drop of the ether employed leaves, on evaporation, scarcely any perceptible residue. Lastly, evaporate the mixed ethereal solutions in a capsule at the temperature of boiling water. The residue, which consists of nearly pure quinia, when dry, should weigh not less than two grains, and should be readily soluble in dilute sulphuric acid.

Actions and Uses.—*Internally* :—Tonic and astringent. Given in debility ; in periodic and intermittent diseases, *e.g.* spasm, neuralgia, and periodical hæmorrhage ; in continued fever, rheumatism, and erysipelas ; in chronic atonic affections of the alimentary canal, *e.g.* dyspepsia ;

in chorea, passive hæmorrhages, profuse mucous discharges, leucorrhœa, and chronic diarrhœa ; in enlargements and indurations of the absorbent glands in scrofula ; in convalescence after either acute or lingering maladies, and during debility after surgical operations. *Externally* : Astringent and antiseptic in leucorrhœa and to fetid ulcers.

Doses.—HORSE, 2 to 4 drachms.—CATTLE, 2 to 4 drachms.—SHEEP, 1 to 2 drachms.—PIG, 1 to 2 drachms. DOG, $\frac{1}{2}$ to 1 drachm. Twice or thrice a day. Sometimes the administration of cinchona to dogs is followed by nausea or vomition ; in such cases the dose should be greatly diminished.

Modes of Application.—*Internally* :—The powdered bark, in the form of ball ; Infusum Cinchonæ Flavæ ; Tinctura Cinchonæ Flavæ. *Externally* :—Infusum Cinchonæ Flavæ ; Tinctura Cinchonæ Flavæ.

Incompatibles.—Alkalies ; alkaline carbonates ; metallic salts ; gelatin.

Preparations.—Infusum Cinchonæ Flavæ ; Tinctura Cinchonæ Flavæ.

CINCHONÆ PALLIDÆ CORTEX

PALE CINCHONA BARK

Therapeutic properties, uses, and doses, similar to those of Cinchonæ Flavæ Cortex.

CINCHONÆ RUBRÆ CORTEX

RED CINCHONA BARK

Therapeutic properties, uses, and doses, similar to those of Cinchonæ Flavæ Cortex.

CINNAMOMUM

CINNAMON

The inner bark of the shoots of *Cinnamomum zeylanicum* (Ceylon cinnamon), or of one or more undetermined species of Chinese cinnamon, belonging to the natural order *Lauraceæ*.

Composition.—Contains a volatile oil, which is used for flavouring purposes, tannic acid, mucilage, colouring matter, an acid, and lignin.

Actions and Uses.—A hæmostatic, and stomachic and carminative, useful in the treatment of diarrhœa. Used in conjunction with purgatives to prevent griping.

Doses : of the bark.—HORSES, $\frac{1}{2}$ to 1 ounce.

Of the oil. $\left\{ \begin{array}{l} \text{DOGS, } \frac{1}{2} \text{ to 1 drachm.} \\ \text{HORSES, 20 minims to 1 drachm.} \\ \text{DOGS, 1 to 4 minims.} \end{array} \right.$

COCAINE

COCAINE

An alkaloid obtained from the leaves of the *Erythroxylon Coca*, a small tree of Peru and Bolivia, belonging to the natural order *Erythroxylaceæ*.

Composition.— $\text{C}_{17}\text{H}_{21}\text{NO}_4$.

Characters and Tests.—Crystallises in colourless prisms requiring for solution 704 parts of water. The Hydrochlorate $\text{C}_{17}\text{H}_{21}\text{NO}_4\text{HCl}$ is readily soluble in water and alcohol, and is therefore usually employed.

Actions and Uses.—*Internally* :—Stimulant and tonic in treatment of sore throat. *Externally* :—As a local anæsthetic. Principally employed in veterinary practice in

diseases of the eye. Twenty minims of a 4 per cent. solution of the hydrochlorate are dropped into the eye, for lessening irritability and pain in cases of conjunctivitis, ophthalmia, and ulceration of the cornea, for stitching torn eyelids, or for treatment of other injuries to the eye. In conjunction with Atropine Sulphate, it is employed in ophthalmoscopic examinations.

Professor Hobday has found that it is unwise to inject more than two grains of hydrochlorate of cocaine subcutaneously to produce local anæsthesia in small dogs, such as fox terriers, or four grains in the case of large dogs, collies, &c. That is, one drachm of a 4 per cent. solution for small dogs, and two drachms of a 4 per cent. solution for large dogs. Local anæsthesia will be manifested in from three to five minutes, and extends for about half an inch to one inch around the seat of injection, and lasts for about twenty minutes.

COLCHICI CORMUS

COLCHICUM CORM

The fresh corm of Meadow Saffron, *Colchicum autumnale*, belonging to the natural order *Melanthaceæ*; collected about the end of June, stripped of its coats, sliced transversely, and dried at a temperature not exceeding 150°F.

Composition.—Contains a poisonous crystallisable alkaloid called cholchicine ($C_{17}H_{19}NO_5$).

Characters.—Fresh corm about the size of a chestnut, flattened where it has an undeveloped bud; furnished with an outer brown and an inner yellow coat; internally white, solid, and fleshy, yielding, when cut, a milky, acrid, and bitter juice. Dried slices about a tenth of an inch thick, moderately indented on one, rarely on both, sides; firm, flat, whitish, amylaceous.

Actions and Uses.—In excessive doses, irritant poison ; in medicinal doses, cathartic, emetic, and sedative ; it also increases the secreting action of the skin, kidneys, and particularly that of the intestinal mucous membrane. Occasionally prescribed on account of its sedative and diuretic qualities, in small and frequently repeated doses, in rheumatism and rheumatic influenza, also in deep-seated or constitutional ophthalmia.

Doses.—HORSE, $\frac{1}{2}$ to 2 drachms.—CATTLE, $\frac{1}{2}$ to 2 drachms. SHEEP, 5 to 30 grains.—DOG, 2 to 8 grains.

Modes of Administration.—In powder, ball, or Tinctura Colchici Seminum.

Incompatibles.—Tincture of Iodine ; Guaiacum ; and all astringents.

Antidotes.—Emetics, where practicable, followed by demulcent drinks ; and, if coma be present, brandy, ammonia, and other powerful stimulants.

COLCHICI SEMINA

COLCHICUM SEEDS

The fully ripe seeds of *Colchicum autumnale*.

Characters.—About the size of white mustard seed ; very hard, and of a reddish-brown colour.

Actions and Uses. { Similar to those of the corm, but
by some considered milder and
more certain.

Doses.—Same as of corm.

Preparation.—Tinctura Colchici Seminum.

COLLODIUM**COLLODION**

Take of

Pyroxylin	1 ounce.
Ether	36 fluid ounces.
Rectified Spirit	12 fluid ounces.

Mix the ether and the spirit, and add the pyroxylin. Set aside for a few days, and, should there be any sediment, decant the clear solution. Keep in a well-corked or stoppered bottle.

Characters.—Colourless, highly inflammable liquid, with ethereal odour, which dries rapidly on exposure to the air, and leaves a thin transparent film, insoluble in water or rectified spirit.

Uses.—As an adhesive to excoriations, ulcers, burns, wounds, &c. ; likewise in erysipelas.

Mode of Application.—Brushed over the part. The applications are repeated, at intervals of a few minutes, until the film has become sufficiently thick to protect the affected part from atmospheric and other causes of irritation.

COLLODIUM FLEXILE**FLEXIBLE COLLODION**

Take of

Collodion	6 fluid ounces.
Canada Balsam	120 grains.
Castor Oil	1 fluid drachm.

Mix, and keep in a well-corked bottle.

Uses and mode of application similar to those of Collodion, to which, however, this preparation is superior on account of its elasticity.

COLLODIUM HÆMOSTATICUM

HÆMOSTATIC COLLODION

Take of

Collodion	10 fluid ounces
Carbolic Acid	1 ounce
Tannic Acid	$\frac{1}{2}$ ounce.

Dissolve.

Action and Use.—*Externally* :—Styptic

Mode of Application.—Same as Collodion.

COLOCYNTHIS

COLOCYNTH

Synonym.—Bitter Cucumber.

The fruit of *Citrullus Colocynthis* deprived of its rind, a native of Western Asia, belonging to the natural order *Cucurbitaceæ*.

Composition.—It contains a bitter alkaloidal principle Colocynthin, a resin Colocynthein, and a tasteless crystalline body Colocynthitin.

Actions and Uses.—A hydragogue purgative in chronic constipation. Its griping tendency may be overcome by combining it with aromatics, or a small proportion of hyoscyamus or belladonna.

Doses.—Of the extract for Dogs, 1 to 2 grains in combination with other purgatives.

CONFECTIO ROSÆ GALLICÆ

CONFECTION OF ROSES

Take of

Fresh Red Rose Petals	.	1 pound.
Refined Sugar	. . .	3 pounds.

Beat the petals to a pulp in a stone mortar, add the sugar, and rub them well together.

Use.—In making *Pilula Hydrargyri cum Ferro*.

CORIANDRI FRUCTUS

CORIANDER FRUIT

Synonym.—Coriander Seed.

The dried fruit of *Coriandrum sativum*, belonging to the natural order *Umbelliferae*. Cultivated in Britain.

Composition.—The active principle is a volatile oil ($C_{10}H_{16}H_2O$).

Characters.—Globular, nearly as large as white pepper, beaked, finely ribbed, yellowish brown; has an agreeable aromatic odour and flavour.

Actions and Uses.

Doses.

Modes of Application.

} Same as *Anisi Fructus*.

CREASOTUM

CREASOTE

A product of the distillation of wood tar.

Composition.—A mixture of carbolic and kresylic acids with other bodies.

Characters and Tests.—Liquid, colourless or with a yellowish tinge, and a strong empyreumatic odour. Sparingly dissolved by water, but freely by alcohol, ether, and glacial acetic acid. Sp. gr. 1.071. Coagulates albumen. A slip of deal dipped into it, and afterwards into hydrochloric acid, acquires on exposure for a short time to the air a greenish-blue colour. Dropped on white filtering paper and exposed to a heat of 212°F., it leaves no translucent stain. It turns the plane of polarisation of a ray of polarised light to the right. Is not solidified by the cold produced by a mixture of hydrochloric acid and sodium sulphate.

Actions and Use.

Doses.

Modes of Application.

Antidotes.

} Similar to those of Acidum
Carbolicum.

Preparations.—Linimentum Creasoti Compositum ; Mistura Creasoti ; Unguentum Creasoti.

CRETA

CHALK

Native friable calcium carbonate (carbonate of lime), CaCO_3 , with small quantities of siliceous and other impurities.

Preparation.—Creta Præparata.

CRETA PRÆPARATA

PREPARED CHALK

Chalk freed from most of its impurities by elutriation, and afterwards dried in small, usually conical, masses.

Characters and Tests.—White amorphous powder ; dissolves with effervescence (escape of carbonic anhydride) in dilute hydrochloric acid, indicating that it is a carbonate, and leaving only a small residue. The solution thus formed gives with ammonium oxalate a white precipitate (calcium oxalate), showing the presence of calcium. The salt (calcium chloride) formed by dissolving prepared chalk in hydrochloric acid, if rendered neutral by evaporation to dryness, and re-dissolved in water, gives only a very scanty precipitate on the addition of saccharated solution of lime, showing the absence of calcium phosphate.

Actions and Uses.—*Internally* :—Antacid in indigestion, chronic diarrhœa, and dysentery ; antidote to oxalic and most other acids. *Externally* :—Desiccant for absorbing irritating discharges ; for protecting wounds, burns, and scalds from the air ; also in erysipelas.

Doses.—HORSE, $\frac{1}{2}$ to 1 ounce.—CATTLE, 1 to 2 ounces.—SHEEP, 2 to 4 drachms.—PIG, 1 to 2 drachms.—DOG, 5 to 15 grains.

Modes of Application.—*Internally* :—In the form of bolus, or suspended in gruel, mucilage, or milk. Given in diarrhœa and dysentery ; seldom administered alone, but usually in combination with opium, ginger, and catechu ; the Mistura Cretæ Composita. *Externally* :—As a desiccant it is used in the form of powder ; to scalds and burns it may be applied mixed with water into a stiff paste.

Incompatibles.—All acids.

Preparations.—Hydrargyrum cum Creta ; Mistura Cretæ Composita

CROCUS**SAFFRON**

The dried stigma and part of style of *Crocus sativus*.
Imported from Spain, France, and Italy.

Natural Order.—*Iridaceæ*.

Composition.—Contains an active principle called saffranin ($C_{48}O_{60}H_{18}$), a volatile oil, and also a colouring matter, crocin ($C_{16}H_{18}O_{16}$).

Characters and Tests.—Hay saffron consists of the dried parts of the styles, with their attached stigmas entangled together. The odour of saffron is penetrating, aromatic, and, in large quantities, narcotic. The taste is bitter and somewhat aromatic. When chewed, saffron tinges the mouth and saliva yellow; and when rubbed on the moistened finger it produces an intensely orange stain.

Actions and Uses.—Seldom employed internally; principally used as a colouring and flavouring ingredient.

CROTONIS SEMINA**CROTON SEEDS**

Synonyms.—*Crotonis Tiglii Semina*, Croton Beans.

Natural Order.—*Euphorbiaceæ*.

Composition.—Contains, in addition to oil, an acrid purgative principle, the chemical nature of which is still unknown.

Characters.—About the size of a coffee grain, oblong, rounded at the extremities, with two faces, the external more convex than the internal. If the shelly covering of

the seeds is partially removed, they present a mottled appearance ; but if entirely so, they are brownish black.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, cathartic in obstinate constipation, and when it is necessary to effect a speedy evacuation of the contents of the intestines.

Doses.—HORSE, 30 to 36 grains.¹—CATTLE, 30 to 80 grains.¹—SHEEP, 5 to 15 grains.¹—DOG, 3 to 10 grains.¹

Modes of Application.—The seed, crushed to powder, made into a bolus, mixed with the animal's food, or diluted with linseed oil.

Preparation.—Oleum Crotonis.

CUMINI FRUCTUS

CUMIN FRUIT

Synonym.—Cumin Seeds.

The dried fruit of *Cuminum Cyminum*.

Natural Order.—*Umbelliferae*.

Composition.—The properties of cumin depend upon a volatile oil, which is of a pale yellow colour and limpid. It is very acrid to the taste, and of a disagreeable odour.

Characters.—The fruit commonly termed *cumin seed* is larger than anise, and of a light brown or greyish-yellow colour. It has some resemblance to, though it is larger than, caraway. Each mericarp has five primary ridges, which are filiform, and furnished with very fine prickles, The four secondary ridges are prominent and prickly, and under each there is one vitta. The odour of the fruit is

¹ One seed weighs about three grains.

strong and aromatic. Both in odour and taste it resembles caraway, though less agreeable.

Actions and Uses.

Doses.

Modes of Application.

} Same as Anisi Fructus.

CUPRI AMMONIO-SULPHAS

AMMONIO-SULPHATE OF COPPER

Composition.—Probably $\text{CuSO}_4 \cdot 4\text{NH}_3 \cdot \text{H}_2\text{O}$.

Mode of Preparation.—Take of

Sulphate of Copper, in powder . 2 ounces.

Carbonate of Ammonia, in powder 3 ounces.

Rub them together in a porcelain mortar until effervescence has ceased ; then roll up the product in bibulous paper, and place it on a porous brick. When dry, preserve it in a well-stoppered bottle.

Characters.—Azure-blue coloured powder ; emits an ammoniacal odour, and has an astringent metallic taste.

Actions and Uses.—Tonic, stimulant, antispasmodic, astringent. Given to horses and cattle in influenza, pleuropneumonia, consumption, and other debilitating maladies ; to dogs it is administered in chorea and other nervous complaints.

Doses.—HORSE, 1 to 2 drachms.—CATTLE, 1 to 2 drachms.—SHEEP, 5 to 20 grains.—DOG, 1 to 4 grains.

Modes of Application.—In the form of ball, or suspended in cold gruel or mucilaginous drink.

Incompatibles.—Acids ; fixed alkalies.

Antidotes.—Albumen ; bean or pea meal suspended in water.

CUPRI IODIDUM CUM IODO

IODIDE OF COPPER WITH IODINE


Synonym.—Improperly, Iodide of Copper.

Composition.—A mixture of cuprous iodide, Cu_2I_2 , and free Iodine.

Mode of Preparation.—Take of

Iodide of Potassium	2 ounces.
Sulphate of Copper	4 ounces.
Boiling Distilled Water	$1\frac{1}{2}$ pint.

Dissolve the sulphate of copper in one pint of the distilled water, and the iodide of potassium in the remaining half-pint. Mix the solutions ; collect the precipitate which forms (mixture of cuprous iodide and iodine) on a filter, wash it with distilled water, and, when dried over sulphuric acid, pulverise it, and preserve it in a stoppered bottle.



Characters and Tests.—Light fawn-coloured powder, which changes colour by exposure to air ; metallic taste. The presence of free iodine is recognised by the preparation emitting the odour peculiar to this element, and by distilled water, after having been shaken with it and filtered, giving a blue colour (starch iodide) with cold aqueous solution of starch. The presence of combined copper and iodine may be proved by dissolving the solid in nitric acid, and testing the resulting solution for these elements.

Actions and Uses.—*Internally* :—Tonic and alterative ; also stimulant to the absorbents. Has been employed in the treatment of diabetes insipidus, glanders, farcy, and nasal gleet. *Externally* :—Stimulant and astringent.

Applied to chronic œdematous enlargements of the legs, ill-conditioned ulcers, and in inveterate grease.

It is doubtful, however, whether the internal or external applications of this agent are attended with any greater benefits than those derived from the use, in similar cases, of sulphate of copper.

Dose.—HORSE, 1 to 3 drachms, daily, combined with gentian and pimento, or some other carminative.

Modes of Application.—*Internally*:—In the form of bolus. *Externally*:—The powder, either alone or made into an ointment with 4 parts of lard.

Incompatibles.—Acids; acidulous salts; alkalies and their carbonates; lime water; vegetable astringents.

CUPRI SUBACETAS

SUBACETATE OF COPPER

Synonyms.—Copper Subacetate; Diacetate of Copper
Ærugo; Verdigris.

Composition.—The blue variety chiefly consists of the compound, $\text{Cu}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot \text{CuO} \cdot 6\text{H}_2\text{O}$; the green variety contains, in addition to this compound, variable quantities of several basic copper acetates.

Mode of Preparation.—Plates of copper are subjected for several weeks to the united action of atmospheric oxygen and acetic acid. Sometimes the acid is derived from refuse grape husks undergoing the acetous fermentation; at other times, cloths soaked in vinegar are arranged alternately with the plates of copper. When a sufficiency of the salt has formed, it is scraped off from any copper that may have remained unacted upon.

Characters and Tests.—Coarse masses or powder, either of a rich blue colour (blue verdigris) or of a pale bluish-

green colour (green verdigris). Astringent and metallic taste. Partially soluble in water, completely dissolved by hydrochloric acid. Evolves an acetous odour, which is increased on the application of heat or of strong sulphuric acid, showing the compound to be an acetate. Boiled with water and a few drops of acetic acid and filtered, a solution is obtained which furnishes a maroon-red precipitate (cupric ferrocyanide) with solution of potassium ferrocyanide, indicating the presence of copper.

Actions and Uses.

Doses.

Modes of Application.

Incompatibles.

Antidotes.

} Same as Cupri Sulphas.

Preparations.—Linimentum Cupri Subacetatis ; Unguentum Cupri Subacetatis.

CUPRI SULPHAS


SULPHATE OF COPPER

Synonyms.—Cupric Sulphate ; Blue Vitriol ; Blue Stone.

Composition.— $\text{CuSO}_4 \cdot 5\text{Aq}$.

Mode of Preparation.—Scraps of metallic copper are heated in a current of air so as to convert them superficially into cupric oxide, CuO , which forms in flakes and is known in commerce as 'copper scales.' The oxide thus produced, digested with dilute sulphuric acid, is converted into sulphate.¹ By filtration and evaporation the salt is obtained in the crystalline state.


¹ Cupric sulphate is also prepared by boiling concentrated sul-



Characters and Tests.—Blue crystalline salt, in oblique prisms, soluble in water, forming a pale blue solution which strongly reddens blue litmus. The aqueous solution gives with barium chloride a white precipitate (barium sulphate) insoluble in hydrochloric acid, showing the salt to be a sulphate ; and with potassium ferrocyanide a maroon-red precipitate, indicating the presence of copper (cupric ferrocyanide). If an aqueous solution of the salt be mixed with twice its volume of chlorine water, to peroxidise any iron that may be present, and solution of ammonia be added, the precipitate (cupric hydrate) formed by the first addition of the ammonia will be dissolved by a further and sufficient addition of the alkali, and a violet-blue solution (copper ammonio-sulphate) will be produced, leaving nothing undissolved unless iron be present, in which case a reddish-brown precipitate will be left.

Actions and Uses.—*Internally* :—In large doses, irritant poison ; in medicinal doses, astringent and tonic. Given in debility, scrofula, dysentery, obstinate diarrhoea, glanders, and farcy ; also, combined or alternated with Liquor Ammoniae or Ammoniae Carbonas, in purpura, typhoid fever, and other debilitating disorders. Likewise administered as an emetic to the dog. *Externally* :—Stimulant astringent, styptic, mild escharotic, detergent. Applied to

phuric acid with metallic copper, when the following decomposition takes place :



phagedenic ulcers, *e.g.* in canker, thrush, and foot-rot, also to farcy buds ; sometimes a weak solution is injected up the vessels or into the frontal or maxillary sinuses in ulcerated conditions of the Schneiderian membranes ; to the skin in scurvy ; to arrest hæmorrhages ; as a collyrium in chronic ophthalmia.

Doses.—*Tonic and Astringent* :—HORSE, 1 to 5 drachms.—CATTLE, 1 to 3 drachms.—SHEEP, 20 to 30 grains.—PIG, 5 to 10 grains.—DOG, 1 to 3 grains.

Modes of Application.—*Internally* :—As a tonic and astringent, in the form of *Massa Cupri Sulphatis*, or suspended in gruel or linseed tea. As an emetic, dissolved in warm water. *Externally* :—In the state of powder ; *Liquor Cupri Sulphatis* ; *Liquor Cupri Sulphatis Compositus* ; *Linimentum Cupri Sulphatis*.

Incompatibles.—Alkalies ; alkaline carbonates ; vegetable astringents.

Antidotes.—Albumen ; milk ; bean or pea meal suspended in water ; mucilaginous drinks.

Preparations.—*Linimentum Cupri Sulphatis* ; *Liquor Cupri Sulphatis* ; *Liquor Cupri Sulphatis Compositus* ; *Massa Cupri Sulphatis*.

CUSSO

KOUSSO

The flowering panicles of *Brayera anthelmintica*, gathered before the seeds are quite ripe, dried in the sun, and reduced to coarse powder. Imported from Abyssinia.

Natural Order.—*Rosaceæ*.

Composition.—Contains two varieties of tannin, a bitter acrid and a tasteless resin, a fixed and a volatile oil, a

crystalline body termed kosing, chlorophyll, sugar, gum, &c.

Characters and Tests.—The branches of flowers are of a greenish-yellow colour, the edges of the petals having a purplish hue; balsamic odour, when freshly opened, compared by Pereira to the combined odour of tea, hops, and senna leaves; slightly acrid and unpleasant taste. On the addition of a ferric salt to an infusion or decoction of kousso, a dark green colour develops.

Actions and Uses.—Anthelmintic, especially for tænia.

Doses.—HORSE, $\frac{1}{2}$ to 1 pound.—DOG, 2 to 6 drachms.

Modes of Application.—In the form of Infusum Cusso.

Preparation.—Infusum Cusso.

DECOCTUM TABACI

DECOCTION OF TOBACCO

Take of

Tobacco 1 ounce.

Water 1 pint.

Boil the water with the tobacco for a few seconds (for ten minutes, if the time will permit), strain, and add cold water to the strained liquid until it measures two pints.

Use.—Antidote to Nux Vomica; also to Strychnia and its preparations.

Doses.—HORSE, 1 to 3 pints.—CATTLE, 1 to 3 pints.—SHEEP, 5 to 15 fluid ounces.—PIG, 5 to 15 fluid ounces.—DOG, 2 to 5 fluid ounces.

Should the spasms continue after the administration of the tobacco, repeat the dose, at intervals to be determined by the effect produced, until complete muscular relaxation ensues. The effects of an overdose of tobacco may be counteracted by the exhibition of diffusible stimulants.

DIGITALINUM**DIGITALIN**

Composition.— $C_{54}H_{84}O_{27}$

The difficultly crystallisable sedative principle of *Digitalis purpurea*. Too expensive for veterinary use.

DIGITALIS FOLIA**DIGITALIS LEAF**

The dried leaf of *Digitalis purpurea*, Purple Foxglove.

Natural Order.—*Scrophulariaceæ*.

Composition.—Contains a neutral active principle, termed digitalin. See Digitalinum.

Mode of Preparation.—The leaves are gathered from the wild indigenous plant, in June, and July, when about two-thirds of the flowers are expanded, and dried by the heat of a stove in a dark place.

Characters.—Ovate-lanceolate, shortly petiolate, rugose, downy, crenate. Bright green colour, which is paler on the under than on the upper surface ; scarce any odour ; acrid and nauseous taste.

Actions and Uses.—*Internally* :—In large doses, irritant and sedative poison ; in medicinal doses, sedative and diuretic. Liable to accumulate in the system and irritate the intestines ; its action should, therefore, be carefully watched. Chiefly given as a sedative in the secondary stages of inflammatory affections, functional diseases of the heart, and chronic rheumatism ; also recommended in pneumonia

and chronic cough. Prescribed as a diuretic, either alone or in combination with nitrate of potash, in dropsical complaints. *Externally* :—Irritant, diuretic, sedative, anodyne. Occasionally employed as a collyrium in ophthalmia.

Doses.—HORSE, 3 grains to 1 drachm.—CATTLE, $\frac{1}{2}$ to 1 drachm.—SHEEP, 5 to 15 grains.—PIG, 2 to 10 grains.—DOG, 1 to 3 grains.

Modes of Application.—*Internally* :—The powdered leaf made into a ball ; Extractum Digitalis. For coughs, Massa Digitalis Composita. *Externally* :—Infusum Digitalis.

Incompatibles.—Sulphate and chloride of iron, the acetates of lead, and the preparations of cinchona bark.

Antidotes.—Empty the alimentary canal, and apply stimulants internally and externally.

Preparations.—Extractum Digitalis ; Massa Digitalis Composita.

EMPLASTRUM PICIS

PITCH PLASTER

Take of

Burgundy Pitch	.	.	.	20 ounces.
Resin .	.	} of each .	.	2 $\frac{1}{4}$ ounces.
Yellow Wax	.		.	
Olive Oil	1 $\frac{1}{2}$ fluid ounce.
Water	.	.	.	1 fluid ounce.

Add the oil and the water to the Burgundy pitch, resin, and wax, previously melted together ; then, constantly stirring, evaporate to a proper consistency.

Use.—Adhesive to wounds when spread on leather.

ENEMA ALOES**ENEMA OF ALOES**

Take of

Aloes	4 drachms.
Carbonate of Potassium	30 grains.
Warm Mucilage of Starch or Gruel	} 1 pint.

Mix and rub together.

ENEMA ASSAFŒTIDÆ**ENEMA OF ASSAFŒTIDA**

Take of

Assafœtida	2½ drachms.
Warm Water	1 pint.

ENEMA MAGNESIÆ SULPHATIS**ENEMA OF SULPHATE OF MAGNESIA**

Take of

Sulphate of Magnesia	2 ounces.
Linseed or Castor Oil	2 fluid ounces.
Mucilage of Starch or Gruel	1 pint.

Dissolve the sulphate of magnesia in the mucilage of starch or gruel, add the oil, and mix.

ENEMA OPII**ENEMA OF OPIUM**

Take of

Tincture of Opium . . .	5 fluid drachms.
Warm Mucilage of Starch or Gruel	1 pint.

Mix.

ENEMA TABACI**ENEMA OF TOBACCO**

Take of

Leaf Tobacco . . .	1 drachm.
Boiling Water . . .	20 fluid ounces (1 pint).

Infuse in a covered vessel for half an hour, and strain.

ENEMA TEREBINTHINÆ**ENEMA OF TURPENTINE**

Take of

Oil of Turpentine . . .	2 fluid ounces.
Mucilage of Starch or Gruel . . .	1 pint.

ERGOTA**ERGOT****Synonym.**—Ergot of Rye.**Natural Order.**—*Fungi*.

The sclerotium (compact mycelium or spawn) of a fungus, *Claviceps purpurea*, produced within the paleæ of the common rye, *Secale cereale*.

Composition.—Complicated, and not well known. Its parturient effects are probably due to a mixture of an alkaloid Cornutin and an acid (Sphacelinic Acid).

Characters.—Subtriangular, curved with a longitudinal furrow on the concave side, obtuse at the ends ; from one-third of an inch to an inch and a half in length ; violet-brown colour superficially, pinkish internally ; solid, frangible, fracture short ; faint odour, which may be increased by trituration with solution of potash.

Actions and Uses.—*Internally* :—In large doses, irritant poison ; in medicinal doses, styptic in hæmorrhages of the lungs, kidneys, and other organs, but principally used as a parturient, *i.e.* to excite the uterus to contract and expel its contents during parturition. Its efficacy in this respect is, however, doubtful on the lower animals. *Externally* :—Employed as a styptic, to arrest hæmorrhages, *e.g.* of the vagina, uterus, &c.

Doses as an Ecboic.—HORSE, 2 to 4 drachms.—CATTLE, 2 to 4 drachms.—SHEEP, 20 to 40 grains.—PIG, 15 to 30 grains.—DOG, 2 to 10 grains.

Modes of Application.—*Internally* :—The powder, in ale or porter, with a carminative ; Extractum Ergotæ Liquidum,¹ Infusum Ergotæ, Tinctura Ergotæ. *Externally* :—Extractum Ergotæ Liquidum ; Infusum Ergotæ ; Tinctura Ergotæ.

Preparations.—Extractum Ergotæ Liquidum ; Infusum Ergotæ ; Tinctura Ergotæ.

¹ Until we are in possession of more complete and accurate information respecting the therapeutic actions of the oil, said to be the only poisonous principle of ergot, Extractum Ergotæ Liquidum, which is freed from this constituent, should be preferred to its other preparation as a parturient.

EUCALYPTUS**EUCALYPTUS**

Synonym.—Australian Blue Gum-tree.

The leaves of the *Eucalyptus globulus*, belonging to the natural order *Myrtaceæ* ; from the leaves an oil is obtained by distillation. (See *Oleum Eucalypti*.)

EUPHORBIIUM**EUPHORBIIUM**

Synonym.—Gum Euphorbium.

An acrid resin, obtained from *Euphorbia officinarum*, and probably other species of euphorbia, belonging to the natural order *Euphorbiaceæ*.

Characters.—Irregular, slightly friable tears, usually pierced with one or two holes. These tears are almost odourless, but their dust is very irritating when applied to the mucous lining of the nose. Taste at first slight, afterwards acrid and burning. On heating, euphorbium melts, swells up imperfectly, evolves an odour like benzoic acid, takes fire, and burns with a pale flame. Alcohol, ether, and oil of turpentine are its best solvents ; water only dissolves a small portion of it.

Composition.—Contains about 60 per cent. of a reddish-brown, hard, brittle resin, which is its active constituent.

Actions and Uses.—Sometimes it is employed as an errhine in chronic affections of the eyes, ears, or brain ; but its local action is so violent that it can only be applied when largely diluted with starch or flour. Mixed with turpen-

tine, Burgundy pitch, or resin, it is employed in the form of plaster, as a rubefacient, in chronic affections of the joints. Sometimes used as a vesicant.

EXTRACTUM ACONITI

EXTRACT OF ACONITE

Mode of Preparation.—Take of

The fresh leaves and flowering tops of Aconite	} 112 pounds.
---	---------------

Bruise in a stone mortar, and press out the juice ; heat it gradually to 130°F., and separate the green colouring matter by a calico filter. Heat the strained liquor to 200°F. to coagulate albumen, and filter. Evaporate the filtrate on a water-bath to the consistence of a thin syrup ; then add to it the green colouring matter previously separated, and, stirring the whole together assiduously, continue the evaporation at a temperature not exceeding 140°F., until the extract is of suitable consistence for forming balls or pills.

Doses.—HORSE, 10 to 15 grains.—CATTLE, 15 to 30 grains.—SHEEP, 2 to 5 grains.—PIG, 2 to 5 grains.—DOG, $\frac{1}{4}$ to 2 grains.

Modes of Application.—In the form of bolus, or rubbed up with cold water.

EXTRACTUM BELLADONNÆ

EXTRACT OF BELLADONNA

Mode of Preparation.—Take of

The fresh leaves and young branches of Belladonna	} 112 pounds.
--	---------------

Bruise in a stone mortar, and press out the juice ; heat it gradually to 130°F., and separate the green colouring matter by a calico filter. Heat the strained liquor to 200°F. to coagulate albumen, and filter. Evaporate the filtrate on a water-bath to the consistence of a thin syrup ; then add to it the green colouring matter previously separated, and, stirring the whole together assiduously, continue the evaporation at a temperature not exceeding 140°F., until the extract is of a suitable consistence for forming balls or pills.

Doses.—HORSE, 1 to 2 drachms.—CATTLE, 1 to 2 drachms.—SHEEP, 3 to 10 grains.—PIG, 2 to 5 grains.—DOG, $\frac{1}{2}$ to 3 grains.

Modes of Application.—In the form of bolus, or rubbed up with cold water.

EXTRACTUM CANNABIS INDICÆ

EXTRACT OF INDIAN HEMP

Preparation.—Take 1 lb. of the dried flowering or fruiting tops of the female plants of *Cannabis sativa*, from which the resin has not been removed ; powder coarsely, macerate for seven days, press out the tincture, distil off the spirit, and evaporate to the consistency of a soft extract.

The *Cannabis sativa* grows in India, where it is known as Gunjah or Ganga ; it is rough to the feel, brittle, of a greenish colour, and with a peculiar characteristic narcotic odour.

Characteristics.—A soft extract, of a greenish colour, and with the peculiar characteristic narcotic odour of the plant.

Natural Order.—*Cannabinaceæ*.

Actions and Uses.—Antispasmodic, hypnotic, and sedative. Used internally in colic and inflammation of the bowels ; in chronic cough ; in tetanus ; and in chorea of dogs it has been observed to lessen the involuntary movements. It has not the effect of producing constipation or loss of appetite when repeatedly given, and is very convenient to administer as bolus or pill.

Doses.—HORSE, 2 to 4 drachms.—DOG, 1 to 3 grains.

Preparation.—Tinctura Cannabis Indicæ.

EXTRACTUM DIGITALIS

EXTRACT OF DIGITALIS

Mode of Preparation.—Take of

Fresh leaves of Digitalis . 112 pounds.

Bruise in a stone mortar, and press out the juice ; heat it gradually to 130°F., and separate the green colouring matter by a calico filter. Heat the strained liquor to 200°F. to coagulate albumen, and filter. Evaporate the filtrate on a water-bath to the consistence of a thin syrup ; then add to it the green colouring matter previously separated, and, stirring the whole together assiduously, continue the evaporation at a temperature not exceeding 140°F., until the extract is of a suitable consistence for forming balls or pills.

Doses.—HORSE, 5 to 10 grains.—CATTLE, 5 to 10 grains.—SHEEP, 1 to 5 grains.—PIG, 1 to 5 grains.—DOG, $\frac{1}{4}$ to 1 grain.

Modes of Application.—In the form of bolus, or rubbed up with cold water.

EXTRACTUM ERGOTÆ LIQUIDUM

LIQUID EXTRACT OF ERGOT

Mode of Preparation.—Take of

Ergot, in coarse powder	. 1 pound.
Ether 1 pint, or a sufficiency.
Distilled Water 3½ pints.
Rectified Spirit 8 fluid ounces.

Shake the ether in a bottle with half a pint of the water, and after the two liquids have separated from one another on standing, decant the ether. Place the ergot in a percolator, and free it from its oil by passing the washed ether slowly through it. Remove the marc, and digest it in three pints of the water at 160°F. for twelve hours. Press out, strain, and evaporate the liquor by the heat of a water-bath to nine fluid ounces ; when cold add the spirit. Allow it to stand for half an hour to coagulate, then filter. The product should measure sixteen fluid ounces.

Actions and Uses.—See Ergota.

Doses.—HORSE, 1 to 4 fluid drachms.—CATTLE, 2 to 4 fluid drachms.—SHEEP, 10 to 40 minims.—PIG, 10 to 20 minims.—DOG, 10 to 30 minims.

Mode of Application.—Diluted with water.

EXTRACTUM FILICIS LIQUIDUM

LIQUID EXTRACT OF MALE FERN

Mode of Preparation.—Take of

Fern Root, in coarse powder	2 pounds.
Ether. 4 pints or a sufficiency

Mix the fern root with two pints of the ether, pack closely in a percolator, and add the remainder of the ether at intervals, until it passes through colourless. Evaporate off the ether over a water-bath, or recover it by distillation, and preserve the oily extract which remains.

Doses.—HORSES, $\frac{1}{2}$ to 1 ounce.—DOG, $\frac{1}{2}$ to 1 drachm.

EXTRACTUM HYOSCYAMI

EXTRACT OF HYOSCYAMUS

Mode of Preparation.—Take of

The fresh leaves and young branches of Hyoscyamus	} 112 pounds.
--	---------------

Bruise in a stone mortar, and press out the juice ; heat it gradually to 130°F., and separate the green colouring matter by a calico filter. Heat the strained liquor to 200°F., and filter. Evaporate the filtrate on a water-bath to the consistence of a thin syrup ; then add to it the green colouring matter previously separated, and, stirring the whole assiduously, continue the evaporation at a temperature not exceeding 140°, until the extract is of a suitable consistence for forming balls or pills.

Doses.—HORSE, 1 to 2 drachms.—CATTLE, 1 to 2 drachms. SHEEP, 15 to 30 grains.—PIG, 10 to 20 grains.—DOG, 2 to 8 grains.

Modes of Application.—In the form of bolus, or rubbed up with cold water.


FERRI CARBONAS SACCHARATA**SACCHARATED CARBONATE OF IRON**

Composition.—Ferrous carbonate (carbonate of iron), FeCO_3 , mixed with ferric oxide, Fe_2O_3 , and sugar, the carbonate forming at least fifty-seven per cent. of the mixture.

Mode of Preparation.—Take of

Sulphate of Iron	.	.	.	2 ounces.
Carbonate of Ammonia	.	.	.	$1\frac{1}{4}$ ounce.
Boiling Distilled Water.	.	.	.	2 gallons.
Refined Sugar	.	.	.	1 ounce.

Dissolve the sulphate of iron and the carbonate of ammonia separately in half a gallon of the water, and mix the two solutions with brisk stirring in a deep cylindrical vessel, which is then to be covered as accurately as possible. Set the mixture by for twenty-four hours, and from the precipitate (carbonate of iron) which has subsided separate the supernatant liquid by a siphon. Pour on the remainder of the water, stir well, and after subsidence again remove the clear liquid. Collect the resulting carbonate on a calico filter, and, having first subjected it to expression, rub it with the sugar in a porcelain mortar. Finally, dry the mixture at a temperature not exceeding 212°F .



The sugar is added to prevent the absorption of atmospheric oxygen, the loss of carbonic anhydride, and the con-

sequent conversion of the ferrous carbonate into ferric oxide.¹


Characters and Tests.—Small coherent lumps of a grey colour, with a sweet very feeble chalybeate taste. Dissolves with effervescence in warm hydrochloric acid diluted with half its volume of water, indicating that it contains a carbonate, and the solution gives but a very slight precipitate with barium chloride, showing its almost total freedom from sulphate; with potassium ferrocyanide the solution gives a pale blue precipitate, which, on exposure to air, changes to a dark blue, indicating the presence of a ferrous salt.

Actions and Uses. }
Doses. } Same as Ferri Sulphas.


Modes of Application.—In bolus, pill, or suspended in some mucilaginous drink.

Incompatibles.—Acids; acidulous salts; vegetable astringents.

¹ When commercial or pharmaceutical carbonate of ammonia is dissolved in water, the ammonium carbamate which it contains is rapidly converted into normal ammonium carbonate $(\text{NH}_4)_2\text{CO}_3$, thus:



This ammonium carbonate decomposes the ferrous sulphate in the manner indicated above, while the ammonium-hydrogen carbonate in the commercial salt acts upon ferrous sulphate in the way explained by the following equation:



FERRI IODIDUM

IODIDE OF IRON


Composition.—Ferrous iodide, FeI_2 , with about eighteen per cent. of water of crystallisation and a little ferric oxide.

Mode of Preparation.—Take of

Fine Iron Wire	.	.	$1\frac{1}{2}$ ounce.
Iodine	.	.	3 ounces.
Distilled Water	.	.	15 fluid ounces.

Put the iodine, iron, and twelve ounces of the water into a flask, and, having heated the mixture gently for about ten minutes, raise the heat and boil till the froth becomes white. Pass the solution as quickly as possible through a wetted calico filter into a dish of polished iron, washing the filter with the remainder of the water, and boil down until a drop of the solution taken out on the end of an iron wire solidifies on cooling. The liquid should now be poured out on a porcelain dish, and, as soon as it has solidified, should be broken into fragments, and enclosed in a well-stoppered bottle.

In this process iron and iodine combine chemically, thus :



The iodide acquires five molecules of water of crystallisation, and becomes $\text{FeI}_2.5\text{Aq}$.

Characters and Tests.—Crystalline ; green, with a tinge of brown ; inodorous ; deliquescent ; almost entirely soluble in water, forming a slightly green solution, which gradually deposits a rust-coloured sediment (ferric oxide), and

acquires a red colour. Its aqueous solution gives a copious blue precipitate with potassium ferricyanide, showing the presence of iron. Mixed with mucilage or starch, it acquires a blue colour on the addition of a minute quantity of solution of chlorine or nitric acid, indicating that the compound is an iodide.

Actions and Uses.—Tonic, hæmatinic, alterative, astringent. Given in *diabetes insipidus* and nasal gleet.

Doses.—HORSE, $\frac{1}{2}$ to 2 drachms.—CATTLE, 1 to 2 drachms.—SHEEP, 15 to 30 grains.—PIG, 10 to 20 grains.—DOG, 1 to 8 grains.

Modes of Application.—In bolus, pill, or solution in water.

Incompatibles.—Acids; acidulous salts; alkalies; alkaline carbonates; lime water; vegetable astringents.

FERRI PEROXIDUM HUMIDUM

MOIST PEROXIDE OF IRON

Synonyms.—Moist Ferric Hydrate ; Ferri Peroxidum Hydratum ; Moist Hydrated Peroxide of Iron.


Composition.—Ferric hydrate, $\text{Fe}_2(\text{HO})_6$, with about eighty-six per cent. of uncombined water.

Mode of Preparation.—Take of

Solution of Persulphate of Iron .	4 fluid ounces.
Solution of Soda . . .	33 fluid ounces.
Distilled Water . . .	A sufficiency.

Mix the solution of persulphate of iron with a pint of the distilled water, and add this gradually to the solution of soda, stirring constantly and briskly. Let the mixture stand for two hours, stirring it occasionally, then put it on a calico filter, and, when the liquid has drained away,

wash the precipitate (moist peroxide of iron) with distilled water until the filtrate ceases to give a precipitate with barium chloride, which shows that the precipitate is freed from sulphate. Lastly, enclose the precipitate, without drying it, in a stoppered bottle from which evaporation cannot take place. This preparation, when used, should be recently made.



Characters and Tests.—Soft pasty mass, of a reddish-brown colour. Dissolves readily in diluted hydrochloric acid without the aid of heat, and the solution gives a copious blue precipitate [ferric ferrocyanide, Prussian blue, $\text{Fe}_4(\text{FeCy}_6)_3$] with potassium ferrocyanide, but none with potassium ferricyanide. A little of it dried at 212°F . until it ceases to lose weight gives off water of hydration when heated to dull redness in a test-tube.

Actions and Uses.—Antidote to poisoning by arsenic, in which case it forms an insoluble and inert ferrous sub-arsenate.

Doses.—The administration should be frequently repeated until at least twelve times the supposed quantity of the arsenic has been given, or until it appears to have been effective in allaying the action of the poison.

Mode of Application.—Mixed with water or some demulcent drink.

FERRI SULPHAS

SULPHATE OF IRON

Synonyms.—Ferrous Sulphate ; Protosulphate of Iron ; Green Vitriol ; Copperas.

Composition.— $\text{FeSO}_4 \cdot 7\text{Aq}$.

no precipitate with hydrosulphuric acid, showing its freedom from lead and other metals precipitable by this reagent.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, astringent, tonic, and hæmatinic. Given in, and during convalescence from, debilitating diseases ; in dysentery, internal hæmorrhages, hæmaturia, diabetes, chorea, and occasionally in epilepsy. *Externally* :—Astringent to ulcers, to check capillary hæmorrhages, and to diminish profuse discharges.

Doses.—HORSE, 1 to 2 drachms.—CATTLE, 2 to 3 drachms.—SHEEP, 15 to 30 grains.—PIG, 10 to 20 grains.—DOG, 1 to 10 grains.

Modes of Application.—*Internally* :—Massa Ferri Sulphatis, or dissolved in water. *Externally* :—In powder or strong aqueous solution.

Incompatibles.—Alkalies ; alkaline carbonates ; lime-water ; vegetable astringents.

Preparation.—Massa Ferri Sulphatis.

FERRUM

IRON

Wrought iron, in the form of wire or nails, free from oxide.

Composition.—An element ; symbol Fe.

Preparations.—Ferri Carbonas Saccharata ; Ferri Iodidum ; Ferri Peroxidum Humidum ; Ferri Sulphas ; Liquor Ferri Perchloridi Fortior ; Liquor Ferri Persulphatis.

FILIX MAS**MALE FERN**

Synonyms.—Male Shield Fern ; Fern Root.

The dried rhizome, with the bases of the foot-stalks and portions of the root-fibres, of *Aspidium* or *Nephrodium Filix-mas*, belonging to the natural order *Filices*, collected in summer. Indigenous.

Composition.—According to Bock the root contains 0·04 per cent. of a volatile oil, to which are attributed its anthelmintic properties. It also contains 4 per cent. of resin, a green fatty oil, tannic acid, and filicic acid.

Mode of Preparation.—The rhizome should be dug up in summer, cleared of root-fibres, &c., but not washed, and dried quickly and thoroughly in the open air, in the shade, or in a hot-air press at a temperature not above 140°F.; the tufts and those parts of the root-stock which are greenish internally are alone to be kept ; they should be reduced to powder immediately, and preserved in well-stoppered bottles ; the stock should be renewed annually, as in two years the plant loses its medicinal qualities.

Characters.—Tufted, scaly, greenish brown ; powder greenish yellow, with a disagreeable odour, and a nauseous, bitter, somewhat astringent taste.

Action and Use.—Anthelmintic for tænia.

Doses.—HORSE, 8 to 16 ounces.—CATTLE, 8 to 16 ounces.—SHEEP, 3 to 4 ounces.—DOG, 1 to 3 ounces.

Mode of Application.—The powder made into a bolus or mixed with gruel ; *Extractum Filicis Liquidum*.¹

Preparation. *Extractum Filicis Liquidum*.

¹ This preparation is to be preferred to the powder, as the latter is uncertain in its action, no matter how well it may be kept.

GALBANUM

GALBANUM

A gum-resin, derived from *Ferula galbaniflua* and probably other allied umbelliferous plants. Imported from India and the Levant.

Composition.—Its medicinal properties depend upon the presence of a resinous body and a volatile oil.

Characters.—Irregular tears, about the size of a pea, usually agglutinated into masses; greenish-yellow colour, translucent, having a strong disagreeable odour, and an acrid bitter taste.

Actions and Uses.	} Similar to Ammoniacum and Assa-
Doses.	

GALLA

GALLS

Synonyms.—Oak-galls; Nut-galls.

Excrescences formed on the young branches of the oak, *Quercus infectoria*, belonging to the natural order *Cupuliferæ*, as a consequence of the irritation produced by the puncture and deposited ova of the female of a hymenopterous insect, *Cynips Gallæ tinctoriæ*.

Composition.—Their therapeutic properties are chiefly due to their containing about 40 to 75 per cent. of tannic acid, $C_{13}H_9O_7CO_2H$, and 3 per cent. of gallic acid $C_6H_2(HO)_3CO_2H$.

Characters.—Hard, heavy, globular bodies, varying in size from half an inch to three-fourths of an inch in

diameter ; tuberculated on the surface, the tubercles and intervening spaces smooth ; bluish-green colour superficially, yellowish white within, with a small central cavity ; intensely astringent.

Actions and Uses.—*Internally and Externally* :—Astringent and styptic. Employed in same cases as Acidum Tannicum.

Doses.—HORSE, 1 to 3 drachms.—CATTLE, 2 to 4 drachms.—SHEEP, 20 to 40 grains.—PIG, 10 to 20 grains.—DOG, 2 to 10 grains.

Modes of Application.—*Internally* :—In the form of bolus or Tinctura Gallæ. *Externally* :—Tinctura Gallæ ; Unguentum Gallæ ; Unguentum Gallæ cum Opio.

Incompatibles.—Mineral acids ; soluble salts of iron and lead ; sulphate of copper ; nitrate of silver ; tartar emetic ; lime-water ; alkaline carbonates and bicarbonates, and many vegetable agents containing alkaloids.

Preparations.—Acidum Gallicum ; Acidum Tannicum ; Unguentum Gallæ ; Unguentum Gallæ cum Opio.

GENTIANÆ RADIX

GENTIAN ROOT

The dried root of *Gentiana lutea*. Collected in the mountain districts of central and southern Europe.

Natural Order.—*Gentianaceæ*.

Composition.—Gentian root consists for the most part of a volatile, odorous, and butyraceous oily matter, a bitter crystalline body consisting of an acid (gentisic acid), and a bitter crystalline principle (gentiopicroin), with gum, sugar, pectin, wax, caoutchouc, a yellow colouring matter, and woody fibre.

Characters.—From half an inch to one inch in thickness, several inches in length, often twisted, much wrinkled or marked with close transverse rings ; brown externally, yellow internally ; tough and spongy ; taste at first sweetish, but afterwards very bitter.

Actions and Uses.—*Internally* :—Tonic and stomachic. *Externally* :—Occasionally employed as a mild excitant and antiseptic.

Doses.—HORSE, 2 to 4 drachms.—CATTLE, 2 to 4 drachms.—SHEEP, 1 to 2 drachms.—PIG, $\frac{1}{2}$ to 1 drachm.—DOG, 10 to 20 grains.

Modes of Application.—In the form of bolus ; Infusum Gentianæ Compositum ; Tinctura Gentianæ Composita.

Preparations.—Infusum Gentianæ Compositum.

GLYCERINUM

GLYCERIN

Synonym.—Propenyl Alcohol.

Composition.— $C_3H_8O_3$, or $C_3H_5(HO)_3$, with a small amount of water.

Mode of Preparation.—Price's glycerin is preferred to that of other makers. Molten fat is resolved by the action of high-pressure steam, at a temperature of from 500° to $600^{\circ}F.$, into glycerin and a mixture of stearic, palmitic, and oleic acids ; the former dissolves in the water formed by the condensation of the steam, while the latter floats on the surface of the solution of glycerin. After the removal of the fatty acids the glycerin is deprived of nearly the whole of its water by evaporation.

Characters.—Clear colourless fluid, oily to the touch. Sp. gr. 1.25. Slight odour ; sweet taste ; freely soluble

in water and in alcohol. Strongly heated, it decomposes and evolves intensely irritating vapours (acrolein, C_3H_4O).

Actions and Uses.—Chiefly employed externally as an emollient in skin affections accompanied by dryness and irritation ; to excoriations, cracked heels, burns, scalds, and freshly blistered surfaces. When used to allay irritability it may be advantageously mixed with an equal bulk of Liquor Plumbi Subacetatis. Also employed as a vehicle for many medicines, and, in consequence of its non-drying properties, it is often added to masses to prevent their hardening.

GLYCERINUM ACIDI CARBOLICI

GLYCERIN OF CARBOLIC ACID

Take of

Carbolic Acid	1 part.
Glycerin	4 parts.

Rub together in a mortar until the acid is dissolved.

Use.—Astringent and stimulant to indolent wounds and ulcers, also to abrasions, sore backs, cracked heels, &c.

GOSSYPIUM

COTTON WOOL

The carded hairs of the seeds of various species of *Gossypium*.

Natural Order.—*Malvaceæ*.

Composition.—Nearly pure cellulose or woody fibre, $C_6H_{10}O_5$.

Uses.—To blistered surfaces, as in burns, scalds, erysipelas, &c. ; and to wounds for the purpose of protecting them from the irritating effects of the atmosphere.

Preparations.—Collodium ; Collodium Flexile.

HYDRARGYRI IODIDUM RUBRUM

RED IODIDE OF MERCURY


Synonyms.—Mercuric Iodide ; Biniodide of Mercury.

Composition.— HgI_2 .

Mode of Preparation.—Take of

Perchloride of Mercury	.	.	4 ounces.
Iodide of Potassium	.	.	5 ounces.
Boiling Distilled Water	.	.	2 quarts.

Dissolve the perchloride of mercury in three pints, and the iodide of potassium in the remainder, of the water, and mix the solutions. When the temperature has fallen to that of the atmosphere, decant the supernatant liquor from the precipitate (red iodide of mercury), and, having collected the latter on a filter, wash it twice with cold distilled water, and dry it at a temperature not exceeding 212°F .



Characters and Tests.—Crystalline powder of a vermilion colour, becoming yellow when gently heated over a lamp on a sheet of paper ; almost insoluble in water, dissolves sparingly in alcohol, but freely in ether or in an aqueous solution of iodide of potassium. Should be completely soluble in ether and entirely volatilised at a high temperature. Heated in a dry test-tube with dry sodium

carbonate, a sublimate, consisting of bluish-white liquid metallic globules, is obtained, indicating the presence of mercury ; the residue remaining in the bottom of the test-tube dissolved in water, acidified with nitric acid, and mixed with solution of starch, produces a blue colour (starch iodide), proving the compound to be an iodide.

Actions and Uses.—*Externally* :—In the form of Unguentum Hydrargyri Iodidi Rubri, as a stimulant detergent and caustic to unhealthy sores and farcy ulcers ; also as a counter-irritant to sore throats, splints, curbs, spavins, enlarged bursæ, thickening of the integument, and similar affections.

Preparation.—Unguentum Hydrargyri Iodidi Rubri.

HYDRARGYRI OXIDUM RUBRUM

RED OXIDE OF MERCURY

Synonyms.—Mercuric Oxide ; Red Precipitate.


Composition.— HgO .

Mode of Preparation.—Take of

Mercury, by weight	.	.	.	8 ounces.
Nitric Acid	.	.	.	$4\frac{1}{2}$ fluid ounces.
Water	.	.	.	2 fluid ounces.

Dissolve half the mercury in the nitric acid diluted with the water, evaporate the solution to dryness, and, with the dry salt (mercuric nitrate) thus obtained, triturate the remainder of the mercury until the two are uniformly blended together. Heat the mixture in a porcelain dish, with repeated stirring, until acid vapours (nitric peroxide) cease to be evolved, and, when cold, enclose the product (red oxide of mercury) in a bottle. The chemical changes

in the foregoing process may be regarded as taking place in two stages, thus :



Characters and Tests.—Orange-red powder, readily dissolved by hydrochloric acid, yielding a solution of mercuric chloride, which gives a yellow precipitate (yellow mercuric oxide) with excess of caustic potash, and a white precipitate (mercuric ammonium chloride, NH_2HgCl) with ammonia. Strongly heated in a test-tube, it should be entirely converted into mercury, which condenses in the upper and cooler parts of the tube, and into oxygen, which may be recognised by its rekindling an incandescent match. During the heating of the oxide no red vapours should be evolved, showing its freedom from nitrate.

Actions and Uses.—*Externally* :—As an escharotic, the powder is sprinkled over wounds ; in the form of Unguentum Hydrargyri Oxidi Rubri it is used as a stimulant and detergent.

Preparation.—Unguentum Hydrargyri Oxidi Rubri. Hydrargyri Oleatum (Oleate of Mercury) is prepared by mixing oleic acid with 10 to 20 per cent. of mercuric

¹ Mercuric nitrate heated alone furnishes mercuric oxide, nitric peroxide, and free oxygen [$2\text{Hg}(\text{NO}_3)_2 = 2\text{HgO} + 4\text{NO}_2 + \text{O}_2$], the latter product escaping into the air and becoming wasted. In the 'British Pharmacopœia' process, above described, this oxygen combines with the additional mercury, and is thereby economised.

oxide. It is used in articular inflammation, exostoses, indurations of the udder, and other glandular and indolent swellings, and as an antiparasitic.

HYDRARGYRI PERCHLORIDUM

PERCHLORIDE OF MERCURY


Synonyms.—Mercuric Chloride ; Bichloride of Mercury ; Corrosive Sublimate ; Mercury Ball.

Composition.— HgCl_2 .

Mode of Preparation.—Take of

Sulphate of Mercury, dried . . .	20 ounces.
Chloride of Sodium, dried . . .	16 ounces.
Black Oxide of Manganese ¹ . . .	1 ounce.

Reduce the sulphate of mercury and the chloride of sodium separately to fine powder, and having mixed them and the oxide of manganese thoroughly by trituration in a mortar, put the mixture into a subliming apparatus, and apply sufficient heat to cause the vapours of perchloride of mercury to rise into the less heated part of the apparatus.



Characters and Tests.—Heavy colourless masses of prismatic crystals, possessing a highly acrid metallic taste ; more soluble in alcohol, and still more so in ether, than in water. Heated, it liquefies, and completely sublimes with-

¹ If the mercuric sulphate (sulphate of mercury) employed be free from mercurous sulphate, the addition of the oxide of manganese may be dispensed with, as in that case the production of calomel need not be apprehended.

out decomposing. Its aqueous solution gives a yellow precipitate (yellow mercuric oxide) with caustic potash, a white precipitate (mercuric-ammonium chloride, NH_2HgCl) with ammonia, and a curdy white precipitate (silver chloride) with silver nitrate ; the first two reactions are indicative of the salt being a mercuric compound, and the third reaction shows that it is a chloride.

Actions and Uses.—*Internally* :—In large doses, irritant and corrosive poison ; seldom administered medicinally, although, in doses of from five to ten grains, it has been recommended in glanders and farcy. *Externally* :—A most powerful antiseptic and disinfectant.¹ Caustic to indolent ulcers, fistulæ, and bony tumours ; as a stimulant in scab, mange, and similar affections ; also as an antiparasitic for the destruction of pediculi, acari, &c. With suitable precautions watery solutions are used for disinfecting infected premises, and carcasses of animals which have died of contagious diseases.

Modes of Application.—As a caustic, the powder ; Liquor Hydrargyri Perchloridi ; Unguentum Hydrargyri Perchloridi. As a stimulant and antiparasitic, Spiritus Hydrargyri Perchloridi Compositus.

Antidotes.—Albumen ; wheaten flour ; bean or pea meal ; gruel or other demulcent drinks, followed by vegetable astringents.

Preparations.—Liquor Hydrargyri Perchloridi ; Spiritus Hydrargyri Perchloridi Compositus ; Unguentum Hydrargyri Perchloridi.

¹ Koch, in 1880, demonstrated that mercuric chloride in a solution of 1 to 1,000 would soon destroy the spores of anthrax. The addition of a small quantity of hydrochloric or tartaric acid is an improvement, as it prevents the coagulation of albumen.

HYDRARGYRI SUBCHLORIDUM

SUBCHLORIDE OF MERCURY


Synonyms.—Mercurous Chloride ; Protochloride of Mercury ; Calomel.

Composition.— HgCl , or probably Hg_2Cl_2 .

Mode of Preparation.—Take of

Sulphate of Mercury . . .	10 ounces.
Mercury	7 ounces.
Chloride of Sodium, dried . .	5 ounces
Boiling Distilled Water . .	A sufficiency.

Moisten the sulphate of mercury with some of the water, and rub it and the mercury together until globules are no longer visible ; add the chloride of sodium, and thoroughly mix the whole by continued trituration. Sublime by a suitable apparatus into a chamber of such a size that the calomel, instead of adhering to its sides as a crystalline crust, shall fall as a fine powder on its floor. Wash this powder with boiling distilled water until the washings cease to be darkened by a drop of ammonium sulphide, showing that any corrosive sublimate that may have been formed in the above process has been removed from the calomel.



Characters and Tests.—Dull white, heavy, and nearly tasteless powder, rendered yellowish by trituration in a mortar ; insoluble in water, spirit, and ether. Digested with solution of potash, it becomes black (mercurous oxide), indicating that it is a mercurous compound ; and

the clear solution obtained by decantation, or filtering the mixture, acidulated with nitric acid, gives a copious white precipitate (silver chloride) with silver nitrate, showing it to be a chloride. Should be entirely volatilised by a sufficient heat, indicating the absence of fixed impurity ; and ether, which has been shaken with it in a bottle, should leave no residue on evaporation, proving the calomel to be free from corrosive sublimate.

Actions and Uses.—*Internally* :—In continued doses it produces ptyalism ; in excessive doses, irritant poison ; in medicinal doses, cathartic, stimulant, sedative, resolvent, cholagogue, diuretic, sialagogue, and, on some animals, diaphoretic, alterative, antiphlogistic, anthelmintic. Given in diseases of the liver ; combined with opium or belladonna, in enteritis and dysentery ; in febrile affections, in inflammation of serous membranes, in typhous inflammation, in lingering inflammation of the intestines, and in chronic glandular enlargements. As a cathartic it should never be given alone to the horse, as its action on this animal is both uncertain and violent. Calomel is seldom given to cattle or sheep, except in cases of obstinate dysentery.

Doses.—HORSE, 20 to 60 grains.—CATTLE, 10 to 30 grains.—SHEEP, 4 to 8 grains.—PIG, 1 to 4 grains.—DOG, 1 to 4 grains.

Modes of Application.—In the form of bolus, or suspended in some mucilaginous drink.

Incompatibles.—Lime-water ; alkalies ; alkaline carbonates.

Antidotes.—Albumen ; wheaten flour ; bean or pea-meal.

Preparation.—Lotio Hydrargyri Nigra.

HYDRARGYRI SULPHAS

SULPHATE OF MERCURY


Synonyms.—Mercuric Sulphate ; Persulphate of Mercury.

Composition.— HgSO_4 .



Mode of Preparation.—Take of

Mercury, by weight . . .	20 ounces.
Strong Sulphuric Acid . . .	12 fluid ounces.

Heat the mercury with the sulphuric acid in a porcelain vessel, stirring constantly until the metal disappears ; then continue the heat until a dry white salt (sulphate of mercury) remains.



Or (in two stages)—



Characters.—White crystalline heavy powder, rendered yellow (mercuric oxysulphate, $\text{HgSO}_4.2\text{HgO}$) by affusion of water. Entirely volatilised by heat.

Uses.—In the preparation of Hydrargyri Perchloridum and Hydrargyri Subchloridum.


HYDRARGYRUM

MERCURY

Synonym.—Quicksilver.

Composition.—An element, symbol Hg.

Mode of Preparation.—Although it is met with in the native metallic state in the quicksilver mines of South America, it is chiefly imported into this country from Spain, Illyria, Bavaria, and several other parts of the world, where it is extracted from a native sulphide termed cinnabar (mercuric sulphide). In order that the cinnabar may be made to yield the mercury it contains, it is either roasted or heated with lime or iron in a suitable distillatory apparatus. The following equation represents the changes which take place in the simpler of these processes :



The mercury distils over, and the ferrous sulphide remains in the retort.

Characters and Tests.—A metal, liquid at ordinary temperature, brilliantly lustrous, and easily divisible into spherical globules. Volatilises at a heat below that of visible redness, and should leave no residue.

Preparations.

Hydrargyri Iodidum Ru-	Hydrargyrum cum Cretâ.
brum.	Lotio Hydrargyri Flava.
„ Oxidum Rubrum.	„ Hydrargyri Nigra.
„ Perchloridum.	Liquor Hydrargyri Nitratis
„ Subchloridum.	Acidus.
„ Sulphas.	Liquor Hydrargyri Perchlo-
Hydrargyrum Ammoniatum.	ridi.

Pilula Hydrargyri Subchloridi et Ferri.	Unguentum Hydrargyri Nitratis.
Unguentum Hydrargyri.	Unguentum Hydrargyri Oxidi Rubri.
„ Hydrargyri Ammoniati.	Unguentum Hydrargyri Iodidi Rubri.
Unguentum Hydrargyri Compositum.	

HYDRARGYRUM AMMONIATUM

AMMONIATED MERCURY


Synonyms.—Hydrargyri Ammonio-chloridum ; Hydrargyri Precipitatum Album ; Mercuric-ammonium Chloride ; Chloride of Mercuric-ammonium ; Amidochloride of Mercury ; White Precipitate.

Composition.—Various views have been adopted respecting the constitution of this body ; it may, however, be regarded as the ammonium chloride, in which two atoms of hydrogen are replaced by a bivalent atom of mercury ; its formula then becomes NH_2HgCl .

Mode of Preparation.—Take of

Perchloride of Mercury	.	.	3 ounces.
Solution of Ammonia	.	.	4 fluid ounces.
Distilled Water	.	.	3 pints.

Dissolve the perchloride of mercury in the water, with the aid of a moderate heat. Mix the solution with the ammonia, constantly stirring. Collect the precipitate (ammoniated mercury) on a filter, and wash it well with cold distilled water until the liquid which passes through ceases to give a precipitate when dropped into a solution of silver nitrate acidulated by nitric acid. Lastly, dry the product at a temperature not exceeding 212°F .



Characters and Tests.—Opaque white powder, on which cold water, alcohol, and ether have no action. Digested with caustic potash, it evolves ammonia and acquires a pale yellow colour (mercuric oxide). The clear liquid obtained by filtering this mixture, after being acidulated with nitric acid, gives a white precipitate (silver chloride) with silver nitrate. Boiled with stannous chloride, it becomes grey and affords globules of metallic mercury. Entirely volatilises at a heat under redness.

Actions and Uses.—*Internally* :—Irritant poison. *Externally* :—The powder, or an ointment consisting of one part of white precipitate and eight parts of lard, is used for the destruction of pediculi and acari.

HYDRARGYRUM CUM CRETA

MERCURY WITH CHALK

Synonym.—Grey powder.

Composition.—An intimate mixture of metallic mercury, mercurous oxide, and calcic carbonate. The proportion of mercurous oxide, which is probably the active ingredient of this preparation, to the other constituents has not yet been ascertained, but it is most likely variable.

Mode of Preparation.—Take of

Mercury, by weight	,	,	1 ounce.
Prepared Chalk	,	,	2 ounces.

Rub the mercury and chalk together in a porcelain mortar until metallic globules cease to be visible to the naked eye, and the mixture acquires a uniform grey colour.

Characters and Tests.—Powder of a light grey colour, free from grittiness, insoluble in water, partly dissolved with effervescence by dilute hydrochloric acid, leaving the mercury in a finely divided state. The solution formed with hydrochloric acid is not precipitated by the addition of stannous chloride, showing that the preparation contains no mercuric oxide.

Actions and Uses.—Chiefly given to the dog as an alterative.

Dose.—Dog, from 3 to 10 grains.

Mode of Application.—In the form of pill or powder.

Incompatibles.—Acids and acidulous salts.

HYOSCYAMI FOLIA

HYOSCYAMUS LEAVES

Synonym.—Henbane Leaves.

Natural Order.—*Solanaceæ*.

The fresh leaves, with the branches to which they are attached, of *Hyoscyamus niger* ; also the leaves separated from the branches and carefully dried, gathered from wild or cultivated British biennial plants, when about two thirds of the flowers are expanded.

Characters.—Leaves sinuated, clammy, and hairy. The fresh herb has a strong unpleasant odour, and a slightly acrid taste, which nearly disappear on drying. The fresh juice dropped into the eye dilates the pupil.

Preparations.—Extractum Hyoscyami ; Tinctura Hyoscyami.

INFUSUM CALUMBÆ

INFUSION OF CALUMBA

Take of

Calumba, in coarse powder . . . 1 ounce.
Cold water 1 pint.

Macerate one hour, and strain.

Doses.—HORSE, 6 to 8 fluid ounces.—CATTLE, 8 to 10 fluid ounces.—SHEEP, 2 to 4 fluid ounces.—PIG, 1 to 2 fluid ounces.—DOG, $\frac{1}{2}$ to 1 fluid ounce.

INFUSUM CATECHU

INFUSION OF CATECHU

Take of

Pale Catechu, in coarse powder . . 6 drachms.
Cinnamon Bark, bruised . . . 1 drachm.
Boiling Distilled Water . . . 1 pint.

Infuse in a covered vessel for half an hour, and strain.

Doses.—HORSE, 4 to 6 fluid ounces.—CATTLE, 4 to 6 fluid ounces.—SHEEP, 1 to 2 fluid ounces.—PIG, 1 to 2 fluid ounces.—DOG, $\frac{1}{4}$ to 1 fluid ounce.

INFUSUM CINCHONÆ FLAVÆ

INFUSION OF YELLOW CINCHONA

Take of

Yellow Cinchona Bark, in coarse }
powder } 1 ounce.
Boiling Distilled Water 1 pint.

Infuse in a covered vessel for two hours, and strain.

Doses.—HORSE, 4 to 6 fluid ounces.—CATTLE, 4 to 6 fluid ounces.—SHEEP, 1 to 2 fluid ounces.—PIG, 1 to 2 fluid ounces.—DOG, $\frac{1}{4}$ to 1 fluid ounce.

INFUSUM CUSSO

INFUSION OF KOUSSO

Take of

Kousso, in coarse powder	.	.	$\frac{1}{2}$ ounce
Boiling Distilled Water	.	.	8 fluid ounces.

Infuse in a covered vessel for fifteen minutes. Not to be strained.

Doses.—HORSE, 20 to 30 fluid ounces.—CATTLE, 20 to 40 fluid ounces.—SHEEP, 6 to 10 fluid ounces.—PIG, 3 to 5 fluid ounces.—DOG, 1 to 3 fluid ounces.

INFUSUM ERGOTÆ

INFUSION OF ERGOT

Take of

Ergot, in coarse powder	.	.	$\frac{1}{2}$ ounce.
Boiling Distilled Water	.	.	1 pint.

Infuse in a covered vessel for half an hour, and strain.

Doses.—HORSE, 6 to 8 fluid ounces.—CATTLE, 8 to 10 fluid ounces.—SHEEP, 4 to 6 fluid ounces.—PIG, 2 to 4 fluid ounces.—DOG, $\frac{1}{2}$ to 1 fluid ounce.

INFUSUM GENTIANÆ COMPOSITUM

COMPOUND INFUSION OF GENTIAN

Take of

Gentian Root, sliced	.	.	4 drachms.
Pimento Berries, bruised	,	,	2 drachms.
Boiling Distilled Water	.	,	1 pint.

Infuse in a covered vessel for four hours, and strain.

Doses.—HORSE, 6 to 8 fluid ounces.—CATTLE, 8 to 10 fluid ounces.—SHEEP, 2 to 4 fluid ounces.—PIG, 1 to 2 fluid ounces.—DOG, $\frac{1}{2}$ to 1 fluid ounce.

INFUSUM LINI

INFUSION OF LINSEED

Synonym.—Linseed Tea.

Take of

Linseed, crushed	.	.	.	1 ounce.
Boiling Distilled Water.	.	.	.	1 pint.

Infuse in a covered vessel for four hours, and strain.

Dose.—*Ad libitum*.

INFUSUM TABACI

INFUSION OF TOBACCO

Take of

Tobacco, in leaf or shreds	.	.	.	1 drachm.
Boiling Distilled Water	.	.	.	1 pint.

Infuse in a covered vessel for one hour, and strain.

Use.—*Externally* :—As a parasiticide.

IODIFORMUM

IODIFORM

Composition.— CHI_3 .

Mode of Preparation.—By mixing together hot alcoholic solutions of Iodine and Potassium Carbonate.

Characters and Tests.—It occurs in yellow, scaly crystals having a persistent, disagreeable odour and taste ;

it is scarcely soluble in cold water, but dissolves in ether and chloroform ; it is very volatile, and should be kept in well-stoppered bottles.

Actions and Uses.—Antiseptic and deodoriser, it is devoid of local irritant effect, but is a local anæsthetic. A few grains dusted over ulcers or malignant growths deodorise and promote their healing. Also used as a dressing for ordinary wounds.

IODOL

TETRA-IODOPYROL

Composition.— C_4I_4NH .

Mode of Preparation.—By the action of iodine upon pyrol, a constituent of mineral oil.

Characters and Tests.—A grayish white powder, free from the disagreeable odour of iodoform.


Actions and Uses.—Iodol may be dusted over wounds in the form of a powder, or it may be used as an ointment.

IODUM

IODINE

Composition.—An elementary body, having the symbol I.

Mode of Preparation.—Sea-weed is collected, dried, and burned ; the semi-vitreous ash (kelp) which remains is lixiviated with water, and the aqueous solution of alkaline sulphates, carbonates, chlorides, and iodides thus formed is evaporated. Sodium sulphate and carbonate, and potassium and sodium chloride, crystallise out and leave a 'mother liquor' rich in iodides, chiefly sodium iodide. **This** 'mother liquor' is mixed with sulphuric acid and manganese peroxide and subjected to distillation.



The iodine distils over and is collected in a series of glass receivers ; the other products remain in the retort.

Iodine thus obtained is purified by being re-sublimed ; it should be preserved in a well-stoppered bottle.

Characters and Tests.—Laminar crystals, of a peculiar odour, bluish-black colour, and metallic lustre, which, when heated, yield a beautiful violet-coloured vapour ; very sparingly soluble in water, but freely dissolved by alcohol, ether, and a solution of potassium iodide. Its aqueous solution strikes a deep blue colour (starch iodide) with cold solution of starch. It should sublime without leaving any residue, and the portion which first comes over should not include any pungent, slender, colourless prisms (cyanogen iodide). If shaken in a dry bottle, iodine will not adhere to the sides if it be free from moisture.

Actions and Uses.—*Internally*:—In excessive doses, irritant and corrosive poison ; in medicinal doses, tonic, alterative, deobstruent, resolvent ; if exhibited for a sufficient length of time, it occasions debility and gives rise to a disturbance of the constitution termed *iodism*. Given as a tonic and alterative in inflammatory affections after the subsidence of fever and acute inflammation ; as a resolvent and stimulant in ascites, hydrothorax, and similar complaints in which effusion is present ; as a deobstruent in chronic enlargement of the liver, and the mammary and other glands, also in indurations of mucous membranes. Likewise given in rheumatism, in scrofulous diseases, and

to arrest the secretion of urine in diabetes insipidus. Recommended as an antidote to poisoning by mercury and the vegetable alkaloids. *Externally* :—Stimulant, counter-irritant, resolvent in sore throat, bursal enlargements, swellings of joints, thickening of the periosteum, tumours of various kinds, strains of tendons ; also in scab, mange, ringworm, mallenders and sallenders, and other skin diseases.

Doses.—HORSE, 10 to 20 grains.—CATTLE, 20 to 30 grains.—SHEEP, 5 to 10 grains.—PIG, 2 to 5 grains.—DOG, 1 to 2 grains.

Modes of Application.—*Internally* :—In the form of bolus, preferably as Liquor Iodi, sometimes as Tinctura Iodi. *Externally* :—Linimentum Iodi ; Tinctura Iodi ; Unguentum Iodi ; Unguentum Sulphuris Iodidi.

Incompatibles.—Ammonia ; metallic salts ; mineral acids ; vegetable alkaloids.

Antidotes.—Emetics, when applicable, aided by demulcent drinks ; starch, flour, and similar amylaceous meals, suspended in water. If the starch-containing remedies be boiled with water and cooled before administration, their efficiency will be increased.

Preparations.

Ferri Iodidum.	Potassii Iodidum.
Hydrargyri Iodidum Rubrum.	Sulphuris Iodidum.
Linimentum Iodi.	Tinctura Iodi.
Liquor Iodi et Potassii Iodidi.	Unguentum Iodi.
	Unguentum Sulphuris Iodidi.

IPECACUANHA**IPECACUANHA**

The dried root of *Cephaëlis Ipecacuanha*, powdered.
Imported from Brazil.

Natural Order.—*Cinchonaceæ*.

Composition.—Its medicinal activity is ascribed to an alkaloid, *emetina*. The formula of this body has not yet been well established.

Characters.—Pieces three or four inches long, about the diameter of a small quill, contorted and irregularly annulated. Colour, brown of various shades. It consists of two parts—the cortical or active portion, which is brittle, and a slender, tough, white, woody centre. Powder, pale brown, with a faint nauseous odour and a somewhat acrid and bitter taste.

Actions and Uses.—*Internally*:—In excessive doses, irritant poison ; in medicinal doses, emetic and sedative ; also mildly cathartic. Chiefly used in dog practice as an emetic, and in the treatment of coughs and colds.

Doses.—As an emetic to the Dog ; from 15 to 30 grains are given in tepid water, either alone or with 1 grain of tartar emetic.

Preparations.—*Pulvis Ipecacuanhæ Compositus* ; *Vinum Ipecacuanhæ*.

JALAPA**JALAP**

The dried tubercles of *Exogonium Purga*, powdered.
Imported from Mexico, from a town of which, Xalapa, its name is derived.

Natural Order.—*Convolvulaceæ*.

Composition.—Its cathartic principle is a resinous body, soluble in alcohol, insoluble in water.

Characters.—Varying from the size of a nut to that of an orange, ovoid, the larger tubercles frequently incised, covered with a thin, brown, wrinkled cuticle ; presenting, when cut, a yellowish-grey colour, with dark brown concentric circles.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, it is sometimes given to sheep and pigs, but usually to the dog, as a cathartic and vermifuge.

Doses.—PIG, 1 to 2 drachms.—DOG, $\frac{1}{2}$ to 3 drachms.

Modes of Application.—In the form of bolus, or rubbed up with some mucilaginous drink.

Preparation.—Extractum Jalapæ, which is a constituent of purgative pills.

KAMALA

KAMALA

Synonym.—Kamela.

Minute glands, which are found adhering to the capsules of *Rottlera tinctoria* ; imported from India.

Natural Order.—*Euphorbiaceæ*.

Composition.—It contains a yellow crystalline principle called Rottlerin.

Characters and Tests.—A fine, granular, mobile, orange-red powder ; scarcely miscible with water, but soluble for the most part in alcohol and ether, forming red-coloured solutions. The residue insoluble in ether should consist principally of tufted hairs, and be free from sand and earthy impurities.

Actions and Uses.—Purgative and anthelmintic for tænia.

Doses.—HORSE, 1 to 2 ounces.—DOG, 1 to 2 drachms.

Modes of Application.—In the form of bolus or suspended in gruel.

KINO

KINO

The inspissated juice obtained from incisions made in the trunk of *Pterocarpus Marsupium*. Imported from Malabar.

Natural Order.—*Leguminosæ*.

Composition.—Kino contains (Kino-tannic acid), Kinoin, Pyrocatechin, Pectin, &c.

Characters and Tests.—Small, angular, brittle, glistening reddish-black fragments. When entire they are opaque, but in thin laminæ are translucent and ruby-red. They are inodorous, but possess a very astringent taste. Soluble in alcohol ; insoluble in ether. The mineral acids, and solutions of gelatine, tartarated antimony, acetate of lead, perchloride of iron, and nitrate of silver, produce precipitates with the watery infusion.

Actions and Uses.

Doses.

Modes of Application.

} Same as Catechu Pallidum.

LINI FARINA

LINSEED MEAL

Linseed, or the cake from which the greater part of the oil has been expressed, reduced to powder.

Natural Order.—*Linacææ*.

Composition.—Chiefly valuable in medicine on account of the mucilage it contains.

Uses.—As an excipient for powders.

Preparations.—Cataplasma Calcis Chloratæ ; Cataplasma Carbonis ; Cataplasma Lini ; Cataplasma Sinapis.

LINI SEMINA

LINSEED

The seeds of the common flax, *Linum usitatissimum*. Cultivated in Britain.

Natural Order.—*Linaceæ*.

Composition.—Its value in medicine depends upon the mucilage it contains, and the oil that is obtained from it by expression.

Characters.—Small, oval, pointed, flat, with acute edges ; smooth, shining, brown externally, yellowish-white internally ; mucilaginous oily taste.

Preparations.—Infusum Lini ; Lini Farina ; Oleum Lini.

LINIMENTUM ACIDI CARBOLICI

LINIMENT OF CARBOLIC ACID

This is synonymous with Glycerinum Acidi Carbolici, which see.

LINIMENTUM ACONITI

LINIMENT OF ACONITE

Take of

Aconite Root, in coarse powder	. 20 ounces.
Camphor 1 ounce.
Rectified Spirit A sufficiency

Moisten the root for three days, then pack in a percolator, and pour sufficient rectified spirit upon it to produce a pint.

Use.—Applied with a camel-hair pencil alone, or mixed, in equal proportions, with soap liniment or compound camphor liniment, and rubbed on the part, it relieves acute rheumatism.

LINIMENTUM AMMONIÆ

LINIMENT OF AMMONIA

Take of

Solution of Ammonia B.P. .	. 1 fluid ounce.
Olive Oil 2 fluid ounces.

Mix together with agitation.

Use.—Counter-irritant in sore throat, rheumatism, sprains, chronic tumours, bronchitis, &c.

LINIMENTUM AMMONIÆ COMPOSITUM

COMPOUND LINIMENT OF AMMONIA

Take of

Solution of Ammonia B.P. .	} Equal parts.
Olive Oil	
Oil of Turpentine	

Shake the solution of ammonia and the olive oil together, then add the oil of turpentine, and shake again until the ingredients are thoroughly mixed.

Use.—Same as Linimentum Ammoniaë, but more active.

LINIMENTUM CALCIS**LINIMENT OF LIME****Synonym.**—Carron Oil.¹

Take of

Solution of Lime	}	Of each, equal parts.
Olive Oil ²		

Mix together with agitation.

Use.—Applied to burns and scalds.**LINIMENTUM CAMPHORÆ****LINIMENT OF CAMPHOR**

Take of

Camphor	1 ounce.
Olive Oil ³	4 fluid ounces.

Dissolve the camphor in the oil.

Uses.—A stimulating embrocation for deep-seated inflammations, glandular swellings, &c.**LINIMENTUM CAMPHORÆ COMPOSITUM****COMPOUND LINIMENT OF CAMPHOR****Synonym.**—White Oil.

Take of

Camphor	1 ounce.
Rectified Spirit	4 fluid ounces.
Olive Oil	1 pint.
Solution of Ammonia	2 fluid ounces.

¹ Carron Oil is usually made with Linseed Oil, as it is cheaper.—ED.² Sometimes one part of oil of turpentine is added to this liniment to increase its activity.³ Oil of turpentine is sometimes added to increase the activity of this liniment.

Dissolve the camphor in the spirit, shake the olive oil with the solution of ammonia, and mix the whole together, so as to form a liniment.

Uses.—Same as *Linimentum Camphoræ* and *Linimentum Ammoniaë*, but more active.

LINIMENTUM CANTHARIDIS

LINIMENT OF CANTHARIDES

Synonyms.—*Oleum Cantharidis* ; Oil of Cantharides.

Take of

Cantharides, in powder	1 part.
Olive Oil ¹ ²	8 parts.

Digest in a water-bath for two or more hours, and strain.

Use.—Vesicant.

LINIMENTUM CREASOTI COMPOSITUM

COMPOUND LINIMENT OF CREASOTE

Take of

Creasote	2 parts.
Oil of Turpentine	4 parts.
Olive Oil	4 parts.

Mix together with agitation.

Uses.—Similar to those of *Glycerinum Acidi Carbolici*.

¹ Or six to ten parts of rape or linseed oil.—ED.

² Oil of turpentine is sometimes added to increase the activity of this liniment.

LINIMENTUM CROTONIS**LINIMENT OF CROTON OIL**

Take of

Croton Oil	1 part.
Oil of Turpentine, or Soap Liniment	6 to 8 parts

Mix together with agitation.

Use.—Vesicant.

LINIMENTUM CUPRI SUBACETATIS**LINIMENT OF SUBACETATE OF COPPER¹**

Synonyms.—Linimentum *Æruginis* ; Liniment of Verdigris.

Take of

Subacetate of Copper, ² in fine powder	} 9 ounces.
Alum, in fine powder	
Treacle	1½ pound.

Boil and stir together until the mixture assumes a brown colour.

Uses.—Erodent and digestive in canker, thrush, and foot-rot.

¹ A mixture of one part of subacetate of copper and eight parts of lard is generally used instead of this liniment.

² The less expensive sulphate of copper may be substituted for the subacetate in this preparation.

LINIMENTUM CUPRI SULPHATIS**LINIMENT OF SULPHATE OF COPPER**

Take of

Sulphate of Copper, in fine powder .	1 part.
Tar or Treacle	4 parts.

Mix, stir, and heat together until the mass assumes a reddish-brown colour.

Uses.—Same as Linimentum Cupri Subacetatis.

LINIMENTUM HYDRARGYRI**COMPOUND LINIMENT OF MERCURY**

Take of

Ointment of Mercury	3 ounces.
Solution of Ammonia	3 ounces.
Liniment of Camphor	3 ounces.

Melt the ointment in the liniment, add the ammonia, and shake them together.

Uses.—A stimulating liniment, applied to indolent ulcers, &c.

LINIMENTUM IODI COMPOSITUM**COMPOUND LINIMENT OF IODINE**

Synonym.—Linimentum Iodi, B.P.

Take of

Iodine	1 $\frac{1}{4}$ ounce.
Iodide of Potassium	$\frac{1}{2}$ ounce.
Camphor	$\frac{1}{4}$ ounce.
Rectified Spirit	10 fluid ounces.

Dissolve the iodine, iodide of potassium, and camphor in the spirit.

Uses.—Stimulant and deobstruent to bursæ and enlarged glands.

LINIMENTUM PICIS LIQUIDÆ

LINIMENT OF TAR

Take of

Oil of Tar	}	Of each, equal parts.
Rape Oil		

Mix together with agitation.

Use.—In skin affections.

LINIMENTUM OLEI PICIS LIQUIDÆ

COMPOSITUM

COMPOUND LINIMENT OF TAR

Take of

Oil of Turpentine	}	Of each, equal parts.
Oil of Tar		
Rape Oil		

Mix together with agitation.

Use.—In skin affections.

LINIMENTUM OPII

LINIMENT OF OPIUM

Take of

Tincture of Opium, by measure	. 1 part.
Compound Liniment of Soap,	} 3 parts.
by measure	

Mix.

Use.—Anodyne to local pains and sprains ; in rheumatism, &c.

LINIMENTUM PLUMBI SUBACETATIS**LINIMENT OF SUBACETATE OF LEAD**

Take of

Solution of Subacetate of Lead	. 1 part.
Olive Oil	. 4 parts.

Mix.

Uses.—Sedative to excoriated surfaces, or after the application of a blister or the actual cautery.

LINIMENTUM SAPONIS COMPOSITUM**COMPOUND LINIMENT OF SOAP**

Synonym.—Opodeldoc.

Take of

Soft Soap	. 4 ounces.
Camphor	. 1 ounce.
Proof Spirit	. 2 pints.
Solution of Ammonia	. $\frac{1}{2}$ pint.

Dissolve the soap and the camphor in the spirit, then add the solution of ammonia, and filter for use.

Uses.—Stimulant and anodyne, after the subsidence of local inflammation ; to sprains and bruises ; to tumours, slight attacks of sore throat, &c.

LINIMENTUM TEREBINTHINÆ**LINIMENT OF TURPENTINE**

Take of

Soft Soap	. 2 ounces.
Camphor	. 1 ounce.
Oil of Turpentine	. 16 fluid ounces.

Dissolve the camphor in the oil of turpentine, then add the soap, rubbing them together until they are thoroughly mixed.

Uses.—Same as *Linimentum Saponis Compositum*.

LINIMENTUM TEREBINTHINÆ COMPOSITUM

COMPOUND LINIMENT OF TURPENTINE

Take of

Croton Oil	2 fluid drachms.
Oil of Turpentine	4 fluid ounces.
Olive or Rape Oil	25 fluid ounces.

Mix.

Uses.—Rubefacient and vesicant.

LIQUOR ACIDI CARBOLICI

SOLUTION OF CARBOLIC ACID

Take of

Carbolic Acid	1 ounce.
Water	40 ounces.

Shake well together.

Uses.—Lotion for extensive wounds ; after operations, as a styptic, and as an antiseptic to prevent the decomposition of any blood that may accumulate from hæmorrhage after the wound has been sewed up. For the destruction of pediculi and acari, it is applied as hot as the animal can bear it.

LIQUOR ALOES**SOLUTION OF ALOES**

Take of

Barbadoes Aloes, ¹ in small pieces	1 pound.
Distilled Water	5 pints.
Proof Spirit	16 fluid ounces.

Digest the aloes and water together over a water-bath until the former is dissolved, remove the solution from the water-bath, add the spirit, and thoroughly mix.

Uses.—Laxative and cathartic.

Doses.—HORSE, 2 to 4 fluid ounces.—CATTLE, 2 to 6 fluid ounces.—SHEEP, 1 to 2 fluid ounces.

LIQUOR ALOES ET SODÆ CARBONATIS**SOLUTION OF ALOES AND CARBONATE OF SODA**

Take of

Barbadoes Aloes, in small pieces	1 pound.
Carbonate of Soda ²	$\frac{1}{2}$ pound.
Water	1 gallon.

Dissolve, with the aid of a water-bath, the aloes in seven pints of the water, and, when cold, pour off the clear solution. Dissolve the carbonate of soda in the remaining pint of water, previously made hot; allow the solution to cool, add it to the solution of aloes, and preserve the mixture in a well-corked or stoppered bottle.

¹ Spiked aloes is frequently substituted for Barbadoes aloes in this preparation, on account of the more ready solubility of the former.

² Four ounces of Potassium Carbonate may be substituted for this.

Uses.—Laxative and cathartic.

Doses.—HORSE, 8 to 16 fluid ounces.—CATTLE, 8 to 16 fluid ounces.—SHEEP, 2 to 6 fluid ounces.

LIQUOR ALUMINIS

SOLUTION OF ALUM

Take of

Alum in Powder	.	.	.	1 ounce.
Distilled Water	.	.	.	16 fluid ounces.

Dissolve the alum in the water.

Uses.—*Externally* :—As an astringent or styptic.

LIQUOR AMMONIÆ

SOLUTION OF AMMONIA

Mode of Preparation.—Take of

Strong Solution of Ammonia	.	.	.	1 pint.
Distilled Water	.	.	.	2 pints.

Mix, and preserve in a stoppered bottle.

Tests.—Sp. gr. 0·959. Other characters similar to Liquor Ammonia Fortior.

Actions and Uses.—*Internally* :—In excessive doses and undiluted, irritant and narcotic poison ; in medicinal doses, antacid, stimulant, antispasmodic, resolvent, and diuretic. Given in tympanitis, hoven, and in many inflammatory and debilitating complaints. *Externally* :—Counter-irritant, rubefacient, vesicant.

Doses.—HORSE, $\frac{1}{2}$ to 1 fluid ounce.—CATTLE, 1 to 4 fluid ounces.—SHEEP, 1 to 3 fluid drachms.—PIG, 1 to 3 fluid drachms.—DOG, 5 to 20 minims.

Modes of Application.—*Internally* :—Diluted with from twenty to thirty parts of water, with cold gruel, or other mucilaginous drink. As a stimulant, sometimes administered to horses and cattle in the form of *Mistura Ammoniaë*; in indigestion and colic, one fluid ounce of *Liquor Ammoniaë* mixed with three to five drachms of aloes, rubbed down in water, may be given to the same animals. *Externally* :—In the form of *Linimentum Ammoniaë*, *Linimentum Ammoniaë Compositum*, *Linimentum Camphoræ Compositum*.

Incompatibles.—Acids and metallic salts, except those of potassium, sodium, barium, and calcium.

Antidotes.—Dilute vinegar, and solutions of the non-poisonous vegetable acids.

Preparations.—*Linimentum Ammoniaë*, *Linimentum Ammoniaë Compositum*, *Linimentum Camphoræ Compositum*.

LIQUOR AMMONIÆ ACETATIS

SOLUTION OF ACETATE OF AMMONIA

Synonym.—Mindererus's Spirit.

Composition.—Ammonium acetate (acetate of ammonia) $(\text{NH}_4)\text{C}_2\text{H}_3\text{O}_2$, dissolved in water.


Mode of Preparation.—Take of

Acetic Acid	10 fluid ounces.
Carbonate of Ammonia	$3\frac{1}{4}$ ounces, or a sufficiency.
Distilled Water	$2\frac{1}{2}$ pints.

Powder the carbonate of ammonia, and add it gradually to the acetic acid until a neutral solution is formed, then add the water.

Assuming, for simplicity's sake, the carbonate of am-

monia to have the formula $(\text{NH}_4)_2\text{CO}_3$, the following equation expresses the decomposition which takes place during its conversion into acetate :



Characters and Tests.—Transparent, colourless, and nearly odourless liquid, having a cooling, saline, and unpleasant taste. Mixed with solution of potash it evolves ammoniacal gas, and warmed with strong sulphuric acid it emits the vapour of acetic acid. The ammoniacal gas and the vapour of acetic acid may each be recognised by their peculiar odour.

Actions and Uses.—*Internally* :—Mild stimulant, diaphoretic, and diuretic. Given as a febrifuge in inflammation of the lungs, influenza, and other febrile affections, usually in combination with Potassæ Nitras, Spiritus Ætheris Nitrosi, or with Spiritus Ætheris Nitrosi and Extractum Belladonnæ. *Externally* :—As a discutient.

Doses.—HORSE, 1 to 4 fluid ounces.—CATTLE, 4 to 8 fluid ounces.—SHEEP, 1 to 3 fluid ounces.—PIG, 1 to 3 fluid ounces.—DOG, 2 to 6 drachms.

Modes of Application.—*Internally* :—Diluted with about three times its bulk of water. *Externally* :—Lotio Ammoniae Acetatis.


Incompatibles.—Acids ; fixed alkalies and their carbonates ; lime-water ; nitrate of silver ; acetate of lead ; and metallic sulphates.

LIQUOR AMMONIÆ FORTIOR

STRONG SOLUTION OF AMMONIA

Composition.—Water holding in solution 32·5 per cent. of ammoniacal gas, NH_3 .

Mode of Preparation.—A mixture of calcium hydrate and coarsely powdered ammonium chloride is heated in an iron bottle connected by a tube with a series of Woulffe's bottles. The first two Woulffe's bottles are empty. Two retain impurities that may pass over from the iron bottle, and the others contain water, by which the ammoniacal gas, set free from the ammonium chloride, is absorbed. The aqueous solution of ammonia thus formed constitutes Liquor Ammonia Fortior.



Characters and Tests.—Colourless liquid, with a characteristic and very pungent odour and strong alkaline reaction. Sp. gr. 0·891.

When diluted with four times its volume of distilled water, it should give no precipitate with lime-water, showing the absence of ammonium carbonate; no precipitate with ammonium oxalate, indicating the absence of calcium; no precipitate with ammonium sulphide, showing the absence of iron; and no precipitate with copper ammonio-sulphate, proving the absence of ammonium sulphide. Liquor Ammonia Fortior acidified with nitric acid should not be rendered turbid by silver nitrate, indicating the absence of chlorine, or by barium chloride, showing that it is free from sulphate.¹

¹ When the boracic acid of Tuscany is saturated with soda, considerable quantities of pure ammonia are evolved. The Liquor

Preparations.—Linimentum Camphoræ Compositum ; Liquor Ammoniaë ; Spiritus Ammoniaë Aromaticus.

LIQUOR ANTIMONII CHLORIDI

SOLUTION OF CHLORIDE OF ANTIMONY


Synonyms.—Solutions of Antimonious Chloride ; Solution of Terechloride of Antimony ; Butter of Antimony.

Composition.—Water holding in solution antimonious chloride, SbCl_3 .

Mode of Preparation.—Take of

Black Antimony (Native Antimonious Sulphide), in powder	}	1 pound.
Hydrochloric Acid		
		4 pints.

Place the black antimony in a porcelain vessel, pour upon it the hydrochloric acid, and, constantly stirring, apply to the mixture, beneath a flue with a good draught, a gentle heat, which must be gradually augmented as the evolution of sulphuretted hydrogen gas begins to slacken, until the liquid boils. Maintain it at this temperature for fifteen minutes ; then remove the vessel from the fire, and filter the liquid (solution of antimonious chloride) through calico into another vessel, returning what passes through first, that a perfectly clear solution may be obtained. Evaporate this to the bulk of two pints, and preserve it in a stoppered bottle.



Ammoniaë and Ammoniaë Carbonas obtained from this source are known in commerce as 'volcanic ammonia,' and are to be preferred to the same preparations procured in any other way on account of their greater purity.

Characters and Tests.—Heavy liquid, sp. gr. 1·47, usually of a yellowish-red colour.¹ A little of it dropped into water gives a white precipitate (antimonious oxychloride, SbOCl), and the filtered solution gives a white precipitate (silver chloride) with silver nitrate, indicating the presence of chlorine. If the white precipitate formed by water be treated with sulphuretted hydrogen, it becomes orange-coloured (antimonious sulphide, Sb_2S_3).

Actions and Uses.—*Internally* :—Irritant and corrosive poison. *Externally* :—Either alone or mixed with four parts of Tinctura Myrrhæ Composita, it is employed as a caustic in the treatment of fistulæ, thrush, canker, luxuriant granulations, foul and foot-rot. It occasions but little pain.

Antidotes.—Chalk ; magnesia or its carbonates ; and demulcent drinks.

Preparation.—Antimonii Oxidum.

LIQUOR ANTIMONII TARTARATI

SOLUTION OF TARTARATED ANTIMONY

Take of

Tartarated Antimony, in powder 1 ounce.

Distilled Water 4 to 8 ounces

Dissolve the tartarated antimony in the water.

Use.—Counter-irritant in deep-seated inflammations and chronic rheumatism of the joints.

¹ The colour is usually darker than here described, owing to the presence of ferric chloride, derived from the iron vessels used in its preparation.

LIQUOR ARGENTI NITRATIS**SOLUTION OF NITRATE OF SILVER**

Take of

Nitrate of Silver	. . .	10 to 15 grains,
Distilled Water	. . .	1 fluid ounce.

Dissolve the nitrate of silver in the water.

Uses.—As an excitant to wounds, &c. One or two drops of the solution, mixed with an equal bulk of water, are sometimes introduced into the eye to remove opacity of the cornea.

LIQUOR ARSENICALIS**ARSENICAL SOLUTION**

Synonyms.—Liquor Potassæ Arsenitis ; Fowler's Solution.

Composition.—By some authorities it is regarded as arsenious anhydride dissolved in a solution of potassium carbonate ; by others as an aqueous solution of potassium arsenite, $KAsO_2$, and carbonate.¹ It contains an amount of arsenical compound corresponding to four grains of arsenious anhydride in one fluid ounce.

Mode of Preparation.—Take of


Arsenious Acid, in powder	} of each, 80 grains.
Carbonate of Potash	
Distilled Water	. . . A sufficiency.

Place the arsenious acid and the carbonate of potash in a flask with ten ounces of water, and apply heat until

¹ Some potassium arsenite is probably formed, which has the power of dissolving arsenious anhydride more readily than water can.—ED.

a clear solution is obtained. Allow this to cool, and then add as much distilled water as will make the bulk one pint.

If the former of the above views respecting the composition of this solution be correct, no decomposition takes place during its preparation ; but if the latter be adopted, part of the carbonate of potash must be decomposed by the arsenious acid in the manner indicated by the following equation :



Characters and Tests.—Clear, colourless liquid, alkaline to test-paper. Sp. gr. 1.009. After being acidulated with hydrochloric acid it gives, with sulphuretted hydrogen, a yellow precipitate (arsenious sulphide, As_2S_3). Soluble in ammonium carbonate, which is brightest when the arsenical solution has been previously diluted.

Actions and Uses.—*Internally* :—Similar to, but more certain in its action and less liable to produce local irritation than, arsenious acid. *Externally* :—In obstinate skin affections, and for the destruction of pediculi, acari, and other external parasites.¹

Doses.—HORSE, $\frac{1}{2}$ to 1 fluid ounce.—CATTLE, $\frac{1}{2}$ to 1 fluid ounce.—SHEEP, $\frac{1}{2}$ to 3 drachms.—DOG, 2 to 10 minims.

¹ Several 'sheep-washes' and 'dips' have a composition similar to that of Liquor Arsenicalis. The simplest is thus prepared :

Take of Arsenious Acid, in powder	} of each, 8 to 10 ounces.
Carbonate of Potash	
Water	
	20 gallons.

Boil together for half an hour. This quantity is sufficient for twenty sheep. Sometimes $8\frac{1}{2}$ ounces of soft soap and 6 ounces of flowers of sulphur are added to the foregoing mixture before it is boiled.

Tuson's sheep dip is a solution of arsenic acid (H_3AsO_4) in water, mixed with a little paraffin oil.

Modes of Application. — *Internally* :—Diluted with sufficient water to make a draught. *Externally* :—Diluted with from two to three times its bulk of water.

Antidote.—A mixture of an alkaline carbonate and Ferri Peroxidum Humidum.

LIQUOR ARSENICI HYDROCHLORICUS

HYDROCHLORIC SOLUTION OF ARSENIC

Synonym.—Liquor Arsenici Chloridi.

Composition.—A solution of arsenious anhydride (arsenious acid) in dilute hydrochloric acid. One ounce of the solution contains four grains of arsenious anhydride.

Mode of Preparation.—Take of

Arsenious Acid, in powder	.	80 grains.
Hydrochloric Acid	.	2 fluid drachms.
Distilled Water	.	A sufficiency.

Boil the arsenious acid with the hydrochloric acid and four ounces of the water, then add distilled water to make the bulk up to one pint.

No decomposition takes place in this process ; the arsenious acid is simply dissolved by the dilute acid.

Characters and Tests.—Colourless liquid, having an acid reaction. Sp. gr. 1·009. Sulphuretted hydrogen gives at once a bright yellow precipitate (arsenious sulphide). Soluble in ammonium carbonate.

Actions and Uses.

Mode of Application.

Doses.

} Same as Liquor Arsenicalis.

Antidote.—A mixture of an alkaline carbonate and Ferri Peroxidum Humidum

Preparation.—Liquor Arsenici et Hydrargyri Iodidi (the B.P. imitation of Donovan's solution) contains about one per cent. by weight of arsenious iodide and of mercuric iodide; it is useful in chronic skin and rheumatic complaints.

Dose.—For HORSE and CATTLE, 1 to 2 ounces.

LIQUOR CALCIS

SOLUTION OF LIME

Synonyms.—Aqua Calcis ; Lime-Water.

Composition.—An aqueous solution of calcium hydrate, $\text{Ca}(\text{HO})_2$. Ten fluid ounces of the solution should contain 7·4 grains of calcium hydrate.

Mode of Preparation.—Take of

Slaked Lime, freshly prepared	. 2 ounces
Distilled Water 1 gallon.

Put the lime into a stoppered bottle containing the water, and shake well for two or three minutes. After twelve hours the excess of lime will have subsided, and the clear solution may be drawn off with a siphon as it is required for use, or transferred to a green glass bottle¹ furnished with a well-ground stopper.

Characters and Tests.—Transparent, colourless, and odourless liquid, having a disagreeable alkaline taste. Feebly alkaline to test-paper. Exposed to the air, a white pellicle or film (calcium carbonate) forms upon its surface,² owing to the absorption of carbonic anhydride by the calcium hydrate. With solution of oxalic acid it gives a white precipitate (calcium oxalate), insoluble in acetic acid.

¹ Lime-water should not be kept in white glass bottles, as it attacks the lead of which they are partly composed.

² This reaction takes place much more rapidly if air from the lungs be blown through a tube into the lime-water.

Actions and Uses.—Chiefly as an antacid ; occasionally as an antidote for poisoning by acids.

Doses.—HORSE, 4 to 6 fluid ounces.—CATTLE, 4 to 6 fluid ounces.—SHEEP, $\frac{1}{2}$ to 2 fluid ounces.—PIG, $\frac{1}{2}$ to 2 fluid ounces.—DOG, 1 to 4 fluid drachms.

Mode of Application.—Mixed with sufficient water to form a draught.

Incompatibles.—Acids, and most metallic salts.

Preparations.—Linimentum Calcis ; Lotio Hydrargyri Flava ; Lotio Hydrargyri Nigra.

LIQUOR CALCIS CHLORATÆ

SOLUTION OF CHLORINATED LIME

Synonyms.—Solution of Chloride of Lime ; Solution of Hypochlorite of Lime.

Composition.—Water holding in solution an amount of calcium hypochlorite equivalent to thirteen grains of available chlorine in one ounce of the fluid.

Mode of Preparation.—Take of

Chlorinated Lime	.	.	.	1 pound.
Distilled Water	.	.	.	1 gallon.

Well mix the water and the chlorinated lime by trituration in a mortar, and, having transferred the mixture to a stoppered bottle, shake it occasionally during three hours ; now pour out the contents of the bottle on a calico filter, and preserve the solution, which passes through, in a stoppered bottle.

Characters and Tests.—Transparent, nearly colourless liquid. Sp. gr. 1.035. Exposed to the atmosphere, the calcium hypochlorite is decomposed by carbonic anhydride ; the calcium carbonate produced is deposited as a white

precipitate, and the chlorine which is set free may be recognised by its odour.

Actions and Uses.—Stimulant, mild erodent, and deodoriser to ill-conditioned and fetid wounds and fistulous sores. Sprinkled about stables and other buildings, it acts as a deodoriser and disinfectant.

In cases of grease, exfoliating bone, &c., it should be diluted with from ten to fifteen times its bulk of water.

Preparation.—Cataplasma Calcis Chloratæ.

LIQUOR CANTHARIDIS TEREBINTHINATUS

TEREBINTHINATED SOLUTION OF CANTHARIDES

Take of

Cantharides, in powder	1 part.
Oil of turpentine	8 parts.

Digest, at a gentle heat, for three or four days, with frequent agitation, and filter. To every ounce of the filtrate add an equal quantity of Canada balsam, and thoroughly mix them together.

Use.—For medicating materials used for setons. The tape, cotton, cord, or other material is immersed in the liniment until it ceases to imbibe any more of the liquid ; it is then to be removed and drawn between the finger and thumb to deprive it of any superfluous fluid, and finally hung up to dry.

LIQUOR CUPRI SULPHATIS

SOLUTION OF SULPHATE OF COPPER

Take of

Sulphate of Copper, in powder . .	5 ounces.
Boiling Distilled Water	1 pint.

Dissolve and filter.

Uses.—For injection into sinuses, one part of this solution is mixed with from two to three times its volume of water. See also Cupri Sulphas.

LIQUOR CUPRI SULPHATIS COMPOSITUS

COMPOUND SOLUTION OF SULPHATE OF COPPER

Take of

Sulphate of Copper	.	.	}	of each
Alum	.	.	}	3 ounces.
Water	.	.	.	1½ pint.
Sulphuric Acid	.	.	.	1½ ounce.

Uses.—See Cupri Sulphas.

LIQUOR FERRI PERCHLORIDI FORTIOR

STRONG SOLUTION OF PERCHLORIDE OF IRON

Composition.—One fluid drachm contains 35·47 grains of perchloride of iron (ferric chloride, Fe_2Cl_6), dissolved in water.


This preparation also contains free hydrochloric acid.

Mode of Preparation.—Take of

Iron Wire	.	.	.	2 ounces.
Hydrochloric Acid	.	.	.	12 fluid ounces.
Nitric Acid	.	.	.	9 fluid drachms.
Distilled Water	.	.	.	8 fluid ounces.

Mix eight fluid ounces of the hydrochloric acid with the distilled water, and in this dissolve the iron at a gentle heat. Filter the solution, add it to the remainder of the hydrochloric acid and the nitric acid, heat the mixture briskly

until on the sudden evolution of red fumes the liquid becomes of an orange-brown colour, then evaporate by the heat of a water-bath until it is reduced to ten fluid ounces. The production of perchloride of iron takes place in two stages :



Characters and Tests.—Orange-brown solution, with a strong astringent and inky taste ; miscible with water and rectified spirit in all proportions. Sp. gr. 1.338. Diluted with water, it gives a white precipitate (silver chloride) with silver nitrate, showing that it contains chlorine, and a dark blue precipitate (Prussian blue) with potassium ferrocyanide, indicating the presence of a ferric salt.

Uses.—In preparing Tinctura Ferri Perchloridi (1 of liquor to 3 of spirit) and in preparing Liquor Ferri Perchloridi (1 of Liquor to 3 of water.)

LIQUOR FERRI PERSULPHATIS


SOLUTION OF PERSULPHATE OF IRON

Take of

Sulphate of Iron	.	.	.	8 ounces.
Sulphuric Acid	.	.	.	$\frac{3}{4}$ fluid ounce.
Nitric Acid	.	.	.	$\frac{3}{4}$ fluid ounce.
Distilled Water	.	.	.	12 fluid ounces.

Add the sulphuric acid to ten ounces of the water, and

dissolve the sulphate of iron in the mixture with the aid of heat. Mix the nitric acid with the remaining two ounces of water, and add the diluted acid to the solution of sulphate of iron. Concentrate the whole by boiling, until, by the sudden disengagement of ruddy vapours, the liquid ceases to be black and acquires a red colour. Test a drop of the solution with potassium ferrocyanide; if a blue precipitate be produced, indicating the presence of some unaltered ferrous salt, a few additional drops of nitric acid are to be added and the boiling renewed in order that the whole may become converted into ferric salt.¹ When the solution is cold, make up the quantity to eleven fluid ounces by the addition, if necessary, of distilled water.



Uses.—Styptic; also in the preparation of Ferri Peroxidum Humidum.

LIQUOR HYDRARGYRI PERCHLORIDI

SOLUTION OF PERCHLORIDE OF MERCURY

Take of

Perchloride of Mercury	3 to 6 grains.
Water	1 fluid ounce.

Uses.—*Externally*:—In the treatment of scab and mange, also for the destruction of vermin infesting the skin.²

¹ The boiling, with small additional quantities of nitric acid, must be repeated until a drop of the solution ceases to yield a blue precipitate with potassium ferrocyanide.

² 2 minims of hydrocyanic acid added to 2 grains of perchloride of mercury dissolved in 1 fluid ounce of water forms a lotion which is sometimes applied to the skin, especially of dogs, to relieve itching,

LIQUOR IODI ET POTASSII IODIDI
SOLUTION OF IODINE AND
IODIDE OF POTASSIUM

Synonyms.—Liquor Iodi ; Lugol's Solution.

Take of

Iodine	22 grains.
Iodide of Potassium	33 grains.
Distilled Water	1 fluid ounce.

Mix and dissolve.

Uses.—Injected into cavities to cause adhesion of their sides ; one fluid drachm diluted with three fluid ounces of water injected into the bladder, to arrest hæmorrhage from that organ ; applied with friction to the skin, when depilated, it is said to promote the return of hair.

Internally :—Antiseptic, and in treatment of Bots.

Doses.—HORSE, 2 to 4 drachms.

LIQUOR MORPHINÆ ACETATIS
SOLUTION OF ACETATE OF MORPHIA

Take of

Acetate of Morphia	9 grains.
Dilute Acetic Acid	18 minims.
Rectified Spirit	$\frac{1}{2}$ ounce.
Distilled Water	$1\frac{1}{2}$ ounce.

Dissolve the acetate of morphia in the mixed liquids
The strength is about 1 in 100.

Actions and Uses.—Stimulant, sedative, and narcotic.

Doses.—HORSE, 2 to 6 fluid drachms.—CATTLE, 2 to 6 fluid drachms.—DOG, 5 to 15 minims.

Preparation.—Injectio morphinæ hypodermica. A solution of acetate of morphine, containing one grain of the acetate in 10 minims of the injection. Use from 30 to 60 minims for the horse in abdominal affections.

LIQUOR MORPHINÆ HYDROCHLORATIS

SOLUTION OF HYDROCHLORATE OF MORPHIA

Take of

Hydrochlorate of Morphia	9 grains.
Diluted Hydrochloric Acid	. 18 minims.
Rectified Spirit	. . . $\frac{1}{2}$ ounce.
Distilled Water	. . . $1\frac{1}{2}$ ounce.

Mix the hydrochloric acid, the spirit, and the water, and dissolve the hydrochlorate of morphia in the mixture. The strength is about 1 in 100.

Uses.—As a sedative, soporific, antispasmodic, &c.

Doses.—See Liquor Morphiæ Acetatis.

LIQUOR PLUMBI SUBACETATIS

SOLUTION OF SUBACETATE OF LEAD

Synonyms.—Liquor Plumbi Diacetatis ; Solution of Diacetate of Lead ; Goulard's Extract.

Composition.—Not accurately known ; probably a mixture of two or more basic acetates of lead.

Mode of Preparation.—Take of

Acetate of Lead	. . . 5 ounces.
Oxide of Lead (Litharge),	} $3\frac{1}{2}$ ounces.
in powder	
Distilled Water	. . . 1 pint, or a sufficiency.

Boil the acetate and oxide of lead in the water for half an hour, constantly stirring ; then filter, and when the liquid is cold add to it more distilled water until the product

measures twenty fluid ounces. Keep the clear solution in well-stoppered bottles, to prevent its being decomposed by atmospheric carbonic anhydride. In this process the acetate of lead unites with the oxide of lead, and is transformed into the mixture of basic salts above referred to.

Characters and Tests.—Heavy, clear, colourless liquid. Sp. gr. 1.26. Alkaline reaction, sweet astringent taste. Exposed to the air it becomes turbid,¹ from the absorption of carbonic anhydride and consequent formation of lead carbonate. The addition of strong sulphuric acid throws down a white precipitate (lead sulphate), and liberates the vapour of acetic acid, which can be identified by its odour. The former reaction indicates the presence of lead; the latter that the salt is an acetate.

Use.—In preparing Linimentum Plumbi Subacetatis.

LIQUOR PLUMBI SUBACETATIS DILUTUS

DILUTED SOLUTION OF SUBACETATE OF LEAD

Synonym.—Goulard's Water.

Mode of Preparation.—Take of

Solution of Subacetate of Lead	} of each, 2 fluid drachms.
Rectified Spirit . . .	
Distilled Water ² . . .	
	19½ fluid ounces.

Mix, and filter through paper. Keep the clear solution in a stoppered bottle.

¹ Air from the lungs blown through a tube into a solution produces the turbidity much more rapidly. Compare with Plumbi Acetas.

² Common water should never be employed for diluting subacetate of lead, as it precipitates the lead as sulphate and carbonate, and thus destroys or diminishes the activity of the preparation.

Actions and Uses.—*Externally* :—Astringent and sedative. Applied, by means of rags wetted with it, to the skin to allay pain and irritation, and to subdue superficial inflammation. See also Linimentum Plumbi Subacetatis. As a collyrium in conjunctival ophthalmia, one part of this preparation, diluted with an equal bulk of water, is to be used.

LIQUOR PLUMBI SUBACETATIS ET OPII

SOLUTION OF SUBACETATE OF LEAD AND OPIUM

Take of

Solution of Subacetate of Lead	1 fluid drachm.
Tincture of Opium	. . . 1 fluid drachm.
Distilled Water	. . . 6 fluid ounces.

Mix.

Uses.—Anodyne to wounds, to blistered and cauterised surfaces, and in superficial inflammation of the eye, skin, or joints.

LIQUOR POTASSÆ

SOLUTION OF POTASH


Composition.—Twenty-seven grains of potassium hydrate, KHO, in one fluid ounce of water.

Mode of Preparation.—Take of

Carbonate of Potash	. . . 1 pound.
Slaked Lime	. . . 12 ounces.
Distilled Water	. . . 1 gallon.

Dissolve the carbonate of potash in the water, and having heated the solution to the boiling-point in a clean iron vessel, gradually mix with it the slaked lime, and continue the ebullition for ten minutes with constant stirring. Then remove the vessel from the fire ; and when, by the

subsidence of the insoluble matter (calcium carbonate), the supernatant liquor (solution of potassium hydrate) has become perfectly clear, transfer it by means of a syphon to a green glass bottle¹ furnished with an air-tight stopper, and add distilled water, if necessary, to make it correspond with the test of specific gravity and neutralising power.²



Characters and Tests.—Transparent, colourless, and odourless liquid. Sp. gr. 1.058. Alkaline to test-paper; acrid taste; feels soapy when rubbed between the finger and thumb. Mixed with excess of solution of tartaric acid, and stirred, it yields a white granular precipitate (hydrogen-potassium tartrate), showing the presence of potassium. Added to excess of diluted hydrochloric acid, no effervescence should be occasioned, indicating its freedom from carbonate. Treated with slight excess of nitric acid, and evaporated to dryness, the residue dissolves in water and yields a nearly clear solution, which should give no precipitate, or a very slight one, when separate portions of it are mixed with silver nitrate, barium chloride, and solution of ammonia; these reactions prove the absence or nearly so, of chlorides, sulphates, and iron respectively.

Use.—In preparing Potassa Caustica.

¹ As it slowly acts upon and dissolves the lead of white bottles.

² See the 'British Pharmacopœia' published in 1867, if it be required to make this test.

LIQUOR POTASSÆ NITRATIS**SOLUTION OF NITRATE OF POTASH**

Take of

Nitrate of Potash, in powder	. 1 ounce.
Water	7 fluid ounces.

Dissolve, and apply immediately after the solution of the salt has been effected.

Uses.—Stimulant to gangrenous and other wounds, also to superficial inflammations and slight sprains.

LIQUOR POTASSÆ PERMANGANATIS**SOLUTION OF PERMANGANATE OF POTASH**

Take of

Permanganate of Potash	4 grains.
Water	1 fluid ounce.

Dissolve.

Use.—Cleansing wash for diseased surfaces.

LIQUOR POTASSÆ SULPHURATÆ**SOLUTION OF SULPHURATED POTASH**

Take of

Sulphurated Potash	1 ounce.
Water	8 fluid ounces.

Dissolve.

Use.—Applied to the skin in scab, mange, and other skin affections.

LIQUOR SODÆ


SOLUTION OF SODA

Composition.—18·8 grains of sodium hydrate, NaHO, in every fluid ounce of water.

Mode of Preparation.—Take of

Carbonate of Soda	.	.	.	28 ounces.
Slaked Lime	.	.	.	12 ounces.
Distilled Water	.	.	.	1 gallon.

Dissolve the carbonate of soda in the water, and, having heated the solution to the boiling-point in a clean iron vessel, gradually mix with it the slaked lime, and continue the ebullition for ten minutes with constant stirring. Then remove the vessel from the fire, and when, by the subsidence of the insoluble matter (calcium carbonate), the supernatant liquid (solution of sodium hydrate) has become perfectly clear, transfer it by means of a siphon to a green glass¹ bottle furnished with an air-tight stopper, and add distilled water, if necessary, to make it correspond with the tests of specific gravity and neutralising power.²



Characters and Tests.—Transparent, colourless, and odourless liquid. Sp. gr. 1047. Alkaline to test-paper; acrid taste; feels soapy when rubbed between the finger and thumb. Mixed with excess of solution of tartaric

¹ As it slowly acts upon and dissolves the lead of white ones.

² See the 'British Pharmacopœia' for 1867, if it be desired to apply this test.

acid, and stirred, no precipitate is thrown down ; thus it is distinguished from Liquor Potassæ. If a platinum wire, moistened with the solution, be heated before the blowpipe, a yellow colour is imparted to the flame, indicating the presence of sodium. It should not effervesce when added to an excess of diluted hydrochloric acid, proving the absence of carbonate. Chlorides, sulphates, and iron are detected in the manner described under Liquor Potassæ.

Preparation.—Soda Caustica.

LIQUOR STRYCHNIÆ

SOLUTION OF STRYCHNINE

Take of

Strychnia, in crystals	. . .	10 grains.
Diluted Hydrochloric Acid	. . .	15 minims.
Rectified Spirit	. . .	5 fluid drachms.
Distilled Water	. . .	15 fluid drachms.

Mix the hydrochloric acid with ten drachms of the water, and dissolve the strychnia in the mixture by the aid of heat ; then add the spirit and the remainder of the water.¹

The strychnine, during its solution in the hydrochloric acid, is converted into hydrochlorate, $C_{21}H_{22}N_2O_2.HCl$.

Actions and Uses.—Stimulant and tonic.

Doses.¹—HORSE, 2 to 6 fluid drachms.—CATTLE, 2 to 6 fluid drachms.—SHEEP, $\frac{1}{2}$ to 2 fluid drachms.—DOG, 3 to 10 minims.²

¹ Two fluid drachms of Liquor Strychniæ contain one grain of strychnia.

² Liquor Strychniæ is the safest form of administering strychnine to dogs.—ED.

LIQUOR ZINCI CHLORIDI

SOLUTION OF CHLORIDE OF ZINC


Synonym.—Butter of Zinc.

Composition.—Prepared in the manner described in the 'British Pharmacopœia' for 1867, one fluid ounce contains 366 grains of zinc chloride, ZnCl_2 . Sir William Burnett's Disinfecting Fluid is a very similar preparation, and is the form of chloride of zinc most commonly employed in veterinary practice; one fluid ounce of it contains 200 grains of zinc chloride. A solution of 'British Pharmacopœia' strength, but containing small quantities of iron and other impurities, may be obtained by the following

Mode of Preparation.—Take of

Granulated Zinc . . .	1 pound.
Hydrochloric Acid . . .	44 fluid ounces.
Distilled Water . . .	1 pint.

Mix the hydrochloric acid and water in a porcelain dish, add the zinc, and apply a gentle heat to promote the action until gas (hydrogen) is no longer evolved. Boil for half an hour, supplying the water lost by evaporation, and allow the product to cool. Filter the liquid (zinc chloride) into a porcelain basin, and evaporate until it is reduced to the bulk of two pints.



Characters and Tests.—Heavy, oily, colourless liquid, having a powerfully astringent and metallic taste. Diluted with water, it will give no precipitate with sulphuretted hydrogen unless lead be present, when a black precipitate or brown coloration (lead sulphide) is produced. If

solution of ammonia be slowly added to the preparation diluted with water, a white, or, if iron be present, a brownish-white, precipitate (zinc hydrate) will be thrown down, which dissolves in excess of the precipitant. On saturating this ammoniacal solution, after filtration, if necessary, with sulphuretted hydrogen, a white precipitate (zinc sulphide) is produced, which readily dissolves in dilute hydrochloric acid ; this reaction is indicative of the presence of zinc. Another portion of the diluted preparation, acidified with nitric acid, and treated with silver nitrate, furnishes a white precipitate (silver chloride), showing that the salt is a chloride.

Actions and Uses.—Chiefly employed as a deodoriser and disinfectant ; for this purpose one fluid ounce of chloride of zinc may be diluted with one gallon of water and distributed about the apartment in shallow plates, or it may be sprinkled over the floor, &c. Fifty minims of Burnett's Fluid, diluted with one pint of water, forms a lotion which may be applied, by means of a sponge saturated with it, to open joints.

In the concentrated state it is a powerful astringent and caustic, and used in the treatment of fetid ulcers and wounds, fistulæ, luxuriant growths, &c. See also *Zinci Chloridum*.

Preparation.¹—*Zinci Chloridum*.

LIQUOR ZINCI SULPHATIS

SOLUTION OF ZINC SULPHATE

Take of

Sulphate of Zinc . . .	5 ounces.
Distilled Water . . .	20 ounces.

¹ Tuson's Liquid Disinfectant consists of a solution of zinc chloride saturated with sulphurous anhydride.

Mix the sulphate of zinc with the water, and dissolve by the aid of heat.

Actions and Uses.—Similar to *Liquor Zinci Chloridum*.

LOTIO ACIDI CARBOLICI

LOTION OF CARBOLIC ACID

Take of

Carbolic Acid	1 fluid drachm.
Water	6 to 12 fluid ounces.

Mix and well shake.

Uses.—Dressing for unhealthy sores, indolent ulcers, foot-rot in sheep, the feet of cattle suffering from foul ; in the treatment of eczema, grease, mange, and scab ; also to allay itching in skin affections.¹

LOTIO AMMONIÆ ACETATIS

LOTION OF ACETATE OF AMMONIA

Take of

Solution of Acetate of Ammonia	} Of each,
Rectified Spirit	
Water	4 fluid ounces.
	1 pint.

Mix.

Use.—Discutient. The part is to be kept constantly moist with the lotion.

¹ One drachm of carbolic acid shaken with a quart of water forms an efficient wash for the mouth, feet, and udders of animals suffering from aphthous epizootic.

LOTIO AMMONII CHLORIDI ET CAMPHORÆ

LOTION OF CHLORIDE OF AMMONIUM AND CAMPHOR

Synonyms.—Lotio Discutiens ; Discutient Lotion.

Mode of Preparation.—Take of

Chloride of Ammonium, in powder	1 ounce.
Diluted Acetic Acid	8 fluid ounces.
Spirit of Camphor.	1 fluid ounce.

Dissolve the chloride of ammonium in the diluted acetic acid, and add the spirit of camphor.

Use.—Discutient to indolent tumours and chronic sprains.

It should be well shaken before use, and applied with friction.

LOTIO AMMONII CHLORIDI ET POTASSÆ NITRATIS

LOTION OF CHLORIDE OF AMMONIUM AND NITRATE OF POTASH

Synonyms.—Lotio Refrigerans ; Cooling Lotion.

Take of

Chloride of Ammonium, in powder	} Of each
Nitrate of Potash, in powder.	
Water	
	2½ ounces.
	1 pint.

Add the chloride of ammonium and nitrate of potash to the water, and, directly they are dissolved, dip cloths in the solution and lay them on the inflamed part.

LOTIO HYDRARGYRI FLAVA

YELLOW MERCURIAL LOTION

Synonym.—Yellow Wash.

Take of

Perchloride of Mercury . . .	18 grains.
Solution of Lime . . .	10 fluid ounces.

Mix.

Use.—Stimulant to unhealthy sores and ulcers.

LOTIO HYDRARGYRI NIGRA

BLACK MERCURIAL LOTION

Synonym.—Black Wash.

Take of

Subchloride of Mercury . . .	30 grains.
Solution of Lime . . .	13 fluid ounces.

Mix.

Use.—Stimulant to unhealthy sores and ulcers.

MAGNESIA ¹

MAGNESIA

Synonyms.—Magnesium Oxide ; Magnesia Usta.

Composition.—MgO.

Mode of Preparation.—Take of


Carbonate of Magnesia ² . . .	4 ounces.
--	-----------

Put it into a Cornish or Hessian crucible closed loosely by a lid, and expose it to a low red heat until a small

¹ Magnesia Levis (Light Magnesia) is a bulky white powder, possessing the same chemical and therapeutic properties as this preparation.

² Really a mixture of carbonate and hydrate. See *Magnesiae Carbonas*.

quantity taken from the centre of the crucible, when it has cooled, and dropped into dilute sulphuric acid, causes no effervescence, showing that it contains no carbonate.



Characters.—White powder, very slightly soluble in water, but readily dissolves in acids without effervescence. Its solution in hydrochloric acid, when neutralised by a mixed solution of ammonium chloride and ammonia, gives a copious crystalline precipitate (ammonia-magnesium phosphate) when hydrogen-di-sodium phosphate is added to it. Dissolved in nitric acid, it should give no precipitate with barium chloride, showing the absence of sulphate, and when neutralised with a mixture of ammonium chloride and ammonia, it should yield no precipitate with ammonium oxalate, indicating its freedom from calcium.

Actions and Uses.—Antacid and laxative. Only given by the veterinarian to foals or calves suffering from indigestion.

Doses.—From 2 to 8 drachms, combined with a carminative, are administered, suspended in milk or gruel.

MAGNESIÆ CARBONAS¹

CARBONATE OF MAGNESIA

Synonym.—Magnesia Alba.


Composition.—A mixture of magnesium carbonate and hydrate, to which the formula $3\text{MgCO}_3.\text{Mg}(\text{HO})_2.4\text{H}_2\text{O}$ has been assigned.

¹ Magnesiæ Carbonas Levis (Light Carbonate of Magnesia) is a bulky white powder, possessing the same chemical and therapeutic properties as this preparation.

Mode of Preparation.—Take of

Sulphate of Magnesia . . .	10 ounces.
Carbonate of Soda . . .	12 ounces.
Boiling Distilled Water . . .	a sufficiency.

Dissolve the sulphate of magnesia and the carbonate of soda each in a pint of the water, mix the two solutions, and evaporate the whole to perfect dryness by means of a sand-bath. Digest the residue for half an hour with two pints of the water, and having collected the insoluble matter (official carbonate of magnesia) on a calico filter, wash it repeatedly with distilled water (to remove sodium sulphate) until the washings cease to give a precipitate with barium chloride. Finally, dry the product at a temperature not exceeding 212° F.



Characters and Tests.—White granular powder, which dissolves with effervescence in diluted nitric acid, showing that the preparation contains a carbonate, and yielding a solution which, when treated with ammonium chloride, ammonia, and hydrogen-di-sodium phosphate, gives a copious white crystalline precipitate (ammonio-magnesium phosphate), showing the presence of magnesium. With slight excess of hydrochloric acid it forms a clear solution, in which barium chloride will cause no precipitate if sulphates be absent. Another portion of the solution supersaturated with ammonia gives no precipitate with ammonium oxalate or sulphuretted hydrogen, unless calcium and iron are respectively present.

Actions and Uses.	} Same as Magnesia.
Doses.	


MAGNESIÆ SULPHAS

SULPHATE OF MAGNESIA

Synonyms.—Magnesium Sulphate ; Epsom Salts.

Composition.— $\text{MgSO}_4 \cdot 7\text{Aq.}$


Mode of Preparation.—Various methods are adopted, but the simplest consists in dissolving *magnesite* (native magnesium carbonate, MgCO_3) in dilute sulphuric acid, heating the solution formed (magnesium sulphate) until carbonic anhydride ceases to be evolved, filtering and evaporating the filtrate, so that the resulting magnesium sulphate may crystallise out on cooling and standing.



In assuming the solid state the salt acquires seven molecules of water of crystallisation.¹

Character and Tests.—Very small, colourless, transparent rhombic prisms, possessing a bitter taste. Readily dissolves in water, and the solution gives a copious white crystalline precipitate (ammonio-magnesium phosphate) with ammonium chloride, ammonia, and hydrogen-disodium phosphate ; and a white precipitate, insoluble in nitric acid, with barium chloride, showing that the salt is

¹ Magnesium sulphate is also prepared by decomposing magnesian limestone ($\text{MgCO}_3 + \text{CaCO}_3$) with sulphuric acid.



The magnesium sulphate is separated from the comparatively insoluble calcium sulphate by solution in water, and is purified by crystallisation.

a sulphate. Should give no blue precipitate with potassium ferrocyanide, indicating its freedom from iron.

Actions and Uses.—Cathartic, diuretic, and febrifuge. Chiefly employed, combined with ginger, as a purgative for cattle and sheep. Used also as a febrifuge for the horse, and as an antidote to poisoning by lead.

Doses.—*Cathartic* :—CATTLE, 1 to 1½ pound. —SHEEP, 1 to 2 ounces. —DOG, 1 to 4 drachms.

Febrifuge :—HORSE, 2 to 4 ounces, dissolved in water twice a day.

Mode of Application.—Dissolved in from ten to twenty times its weight of water. Sometimes, as in cases of obstinate constipation and torpidity of the bowels in cattle, a drachm of calomel, or ten to fifteen croton beans, are added to the ordinary dose of the salt, to increase its cathartic activity.

Incompatibles.—Alkaline carbonates ; lime-water ; acetate of lead.

MANGANESII OXIDUM NIGRUM

BLACK OXIDE OF MANGANESE

Synonyms.—Manganic Peroxide ; Manganese Dioxide ; Peroxide of Manganese ; Binoxide of Manganese.

Composition.— MnO_2 . Found native.

Character and Tests.—Heavy black powder, which dissolves almost entirely in hydrochloric acid, with evolution of chlorine, and gives off oxygen when heated to bright redness.

Use.—Chiefly for the production of chlorine, and in the preparation of Potassium Permanganate.

MARMOR ALBUM**WHITE MARBLE**

Composition.—Naturally crystallised calcium carbonate, CaCO_3 .

Characters and Tests.—Heavy, white crystalline, solid. Dissolves almost completely in hydrochloric acid, with evolution of carbonic anhydride.

Use.—Chiefly in the production of carbonic anhydride.

MASSA ALOES**MASS OF ALOES**

Synonym.—Cathartic Mass.

Take of

Barbadoes Aloes, in small pieces	. 8 ounces.
Glycerin	2 ounces.
Ginger, ¹ in powder	1 ounce.
Rape Oil	1 ounce.

Melt together in a water-bath, and thoroughly incorporate by frequent stirring.

Use.—Cathartic for the horse.

Dose.—From 6 to 8 drachms.

¹ In cases in which the use of ginger may be considered objectionable substitute an equal weight of powdered gentian.

MASSA ALOES COMPOSITA**COMPOUND MASS OF ALOES**

Synonym.—Alterative Mass.

Take of

Barbadoes Aloes, in powder . . .	1 ounce.
Soft Soap	1 ounce.
Common Mass	6 ounces.

Thoroughly incorporate by beating in a mortar, so as to form a mass.

Use.—Alterative for the horse.

Dose.—1 ounce.

**MASSA ANTIMONII TARTARATI
COMPOSITA****COMPOUND MASS OF TARTARATED
ANTIMONY**

Synonym.—Fever Ball.

Take of

Tartarated Antimony, in powder . . .	$\frac{1}{2}$ drachm.
Camphor, in powder	$\frac{1}{2}$ drachm.
Nitrate of Potash, in powder	2 drachms.
Common Mass	a sufficiency.

Mix so as to form a bolus.

Use.—Febrifuge for the horse.

Dose.—The above mixture constitutes one dose.

MASSA BELLADONNÆ COMPOSITA

COMPOUND MASS OF BELLADONNA

Synonym.—Cough Ball.**Take of**

Extract of Belladonna . . .	$\frac{1}{2}$ to 1 drachm.
Barbadoes Aloes, in powder . .	1 drachm.
Nitrate of Potash, in powder . .	2 drachms.
Common Mass	a sufficiency.

Mix so as to form a bolus.**Use.**—For the horse in chronic cough.**Dose.**—The above mixture constitutes one dose.**MASSA CATECHU COMPOSITA**

COMPOUND MASS OF CATECHU

Synonym.—Astringent Mass.**Take of**

Extract of Catechu, in fine powder . .	1 ounce.
Cinnamon Bark, in fine powder . . .	1 ounce.
Common Mass	6 ounces.

Mix.**Use.**—Astringent for the horse.**Dose.**—1 ounce, in the form of bolus.**MASSA COMMUNIS**

COMMON MASS

Take of

Linseed, finely ground	} of each, equal parts.
Treacle	

Mix together so as to form a mass.

Use.—An excipient for medicinal agents when they are to be administered in the form of bolus.

MASSA CUPRI SULPHATIS

MASS OF SULPHATE OF COPPER

Synonym.—Tonic Mass.

Take of

Sulphate of Copper, finely powdered	. 1 ounce.
Ginger, in powder 1 ounce.
Common Mass 6 ounces.

Mix.

Use.—Tonic for the horse.

Dose.—6 to 8 drachms.

MASSA DIGITALIS COMPOSITA

COMPOUND MASS OF DIGITALIS

Synonym.—Cough Ball.

Take of

Barbadoes Aloes, in powder 2 ounces.
Digitalis 1 ounce.
Common Mass 13 ounces.

Mix.

Use.—For the horse in chronic cough.

Dose.—1 ounce, once or twice a day.

MASSA FERRI SULPHATIS**MASS OF SULPHATE OF IRON**

Synonym.—Tonic Mass.

Take of

Sulphate of Iron, in powder	.	.	.	2 ounces.
Ginger, in powder	.	.	.	1 ounce.
Common Mass	.	.	.	5 ounces.

Mix.

Use.—Tonic for the horse.

Dose.—6 to 8 drachms.

MASSA RESINÆ COMPOSITA**COMPOUND MASS OF RESIN**

Synonym.—Diuretic Mass.

Take of

Resin, in powder	.	.	.	} of each, equal parts.
Nitrate of Potash, in powder	.	.	.	
Hard Soap	.	.	.	

Mix.

Use.—Diuretic for the horse.

Dose.—1 ounce.

MASSA ZINGIBERIS COMPOSITA**COMPOUND MASS OF GINGER****Synonym.**—Cordial Mass.**Take of**

Ginger, in powder	}	of each, equal parts.
Gentian Root, in powder		
Treacle		a sufficiency.

Mix so as to form a mass.**Use.**—Stomachic for the horse.**Dose.**—1 ounce.**MISTURA AMMONIÆ****AMMONIA MIXTURE****Take of**

Solution of Ammonia	}	of each, equal parts.
Spirit of Nitrous Ether		
Compound Tincture of Gentian		

Actions and Uses.—Stimulant and antispasmodic.**Doses.**—HORSE, 1 to 2 fluid ounces.—CATTLE, 2 to 4 fluid ounces.—SHEEP, 4 to 6 fluid drachms.**MISTURA CREASOTI****CREASOTE MIXTURE****Take of**

Creasote	16 minims.
Acetic Acid	16 minims.

Spirit of Juniper	.	.	.	$\frac{1}{2}$ drachm.
Syrup	.	.	.	1 ounce.
Distilled Water.	.	.	.	15 ounces.

Mix.

Actions and Uses.—Astringent, styptic, and antiseptic.

Doses.—HORSE, 7 to 10 ounces.—DOG, $\frac{1}{2}$ to 1 ounce.

MISTURA CRETÆ COMPOSITA

COMPOUND CHALK MIXTURE

Synonym.—Sheep and Calves' Cordial.

Take of

Prepared Chalk	.	.	.	2 ounces.
Catechu, in powder	.	.	.	1 ounce.
Ginger	.	.	.	$\frac{1}{2}$ ounce.
Opium	.	.	.	1 drachm.
Peppermint Water	.	.	.	1 pint.

Use.—To check diarrhœa in calves and sheep.

Doses.—CALF, $\frac{1}{2}$ to 1 fluid ounce.—SHEEP, 4 to 8 fluid drachms.

MISTURA RICINI

CASTOR-OIL MIXTURE

Take of

Castor Oil	.	.	.	3 ounces
Syrup of Buckthorn	.	.	.	2 ounces
Syrup of Poppies	.	.	.	1 ounce.

Mix.

Use.—Cathartic for the dog.

Dose.—4 to 8 fluid drachms.

MORPHIÆ ACETAS**ACETATE OF MORPHIA**

Composition.— $C_{17}H_{19}NO_3C_2H_4O_2$.

Preparation.—Acetate of morphia is prepared by decomposing a solution of hydrochlorate of morphia by ammonia solution, adding dilute acetic acid to the precipitated morphia, and drying at a gentle heat.

Characters and Tests.—It closely resembles the alkaloid, from which it is distinguished by the evolution of an acetous odour on the addition of sulphuric acid.

Actions and Uses.

Doses.

Modes of Application.

} See Morphicæ
Hydrochloras.

Preparation.—Liquor Morphicæ Acetatis.

MORPHIÆ HYDROCHLORAS**HYDROCHLORATE OF MORPHIA**

Synonyms.—Morphicæ Murias ; Muriate of Morphia.

Composition.— $C_{17}H_{19}NO_3.HCl.3H_2O$.

Mode of Preparation.—Take of

Opium sliced	1 pound.
Chloride of Calcium	$\frac{3}{4}$ ounce.
Purified Animal Charcoal	$\frac{1}{4}$ ounce.
Diluted Hydrochloric Acid	{ 2 fluid ounces, or a sufficiency.
Solution of Ammonia	
Distilled Water	{ of each, a sufficiency.

Macerate the opium for twenty-four hours with two pints of the water, and decant. Macerate the residue for twelve hours with two pints of the water, decant, and repeat the process with the same quantity of the water, subjecting the insoluble residue to strong pressure. Unite the liquors, evaporate in a water-bath to the bulk of one pint, and strain through calico. Pour in now the chloride of calcium, previously dissolved in four fluid ounces of distilled water, and evaporate until the solution is so far concentrated that upon cooling it becomes solid. Envelop the mass in a double fold of strong calico, and subject it to powerful pressure, preserving the dark fluid which exudes. Triturate the squeezed cake with about half a pint of boiling distilled water, and, the whole being thrown upon a paper filter, wash the residue well with boiling distilled water. The filtered fluids having been evaporated as before, cooled and solidified, again subject the mass to pressure ; and, if it be still much coloured, repeat the process a third time, the expressed liquids being always preserved. Dissolve the pressed cake in six fluid ounces of boiling distilled water ; add the animal charcoal, and digest for twenty minutes ; filter, wash the filter and charcoal with boiling distilled water, and to the solution thus obtained add the solution of ammonia in slight excess.

Let the pure crystalline morphia, which separates as the liquid cools, be collected on a paper filter and washed with cold distilled water until the washings cease to give a precipitate with solution of nitrate of silver acidulated by nitric acid.

From the dark liquids expressed in the above process an additional product may be obtained by diluting them with distilled water, precipitating with solution of potash added in considerable excess, filtering, and supersaturating the filtrate with hydrochloric acid. This acid liquid, digested

with a little animal charcoal, and again filtered, gives, upon the addition of ammonia, a small quantity of pure morphia.

Diffuse the pure morphine, obtained as above, through two fluid ounces of boiling distilled water placed in a porcelain capsule, kept hot, and add, constantly stirring, the diluted hydrochloric acid, proceeding with caution, so that the morphine may be entirely dissolved and a neutral solution obtained. Set aside to cool and crystallise. Drain the crystals and dry them on filtering-paper. By further evaporating the mother liquor, and again cooling additional crystals are obtained. Water extracts from opium the meconate and sulphate of morphine and codeine ; a part of the narcotine, of the meconine, of the narceine, and of the thebaine ; the brown acid extractive, and a part of the resin and of the fat oil. When chloride of calcium is added to infusion of opium, meconate, with a little sulphate of lime, and some resinous colouring matter, are precipitated, while the hydrochlorates of morphine and of codeine are left in solution. A watery solution of the impure crystals obtained by evaporation is then decomposed by ammonia by which the morphia is precipitated, while codeine and hydrochlorate of ammonia are left in solution. The morphine is dissolved in hydrochloric acid, and the solution of the hydrochloric decolorised by charcoal.

Characters and Tests.—White, flexible, acicular prisms of a silky lustre, not changed by exposure to the air, soluble in water and alcohol. The aqueous solution gives a white curdy precipitate with nitrate of silver, and a white one with potash, which is redissolved when an excess of the alkali is added. Moistened with strong nitric acid it becomes orange-red, and with solution of perchloride of iron greenish-blue. Entirely destructible by heat, leaving no residue. Twenty grains of the salt dissolved in half an ounce of warm water, with ammonia added in the slightest possible excess, give on cooling a crystalline precipitate

which, when washed with a little cold water and dried by exposure to the air, weighs 15 to 18 grains.

Actions and Uses.—*Internally*:—Anodyne, sedative, soporific, and antispasmodic. *Externally*:—To relieve neuralgia, tetanus, acute rheumatism, and enteritis in horses.

Doses.¹—HORSE, 3 to 10 grains.—CATTLE, 3 to 10 grains. SHEEP, $\frac{1}{2}$ to 2 grains.—PIG, $\frac{1}{2}$ to 2 grains.—DOG, $\frac{1}{8}$ to $\frac{1}{2}$ grain.

Modes of Application.—*Internally*:—In the form of bolus, or dissolved in water acidulated with a little hydrochloric acid. *Externally*:—By hypodermic injection. See Liquor Morphiae Acetatis.

Antidotes.—See Opium.

MORPHINA

MORPHINE

An alkaloid, $C_{17}H_{19}NO_3$, obtained from opium. Nearly insoluble in cold water. Readily soluble in hydrochloric or acetic acid. Used in the form of acetate or hydrochlorate of morphia.

APOMORPHINE

APOMORPHINE

Composition.— $C_{17}H_{17}NO_2$.

Mode of Preparation.—By heating morphine for some hours with an excess of hydrochloric acid, and precipitating the hydrochloride thus formed with sodium carbonate.

¹ For Liquor Morphiae Hydrochloratis, see p. 174.

Characters and Tests.—An amorphous white powder, which rapidly turns green in air, and dissolves in ether with a pink colour.

Actions and Uses.—A powerful emetic.

Dose.—DOGS, $\frac{1}{10}$ to $\frac{1}{8}$ of a grain.

MUCILAGO ACACIÆ

MUCILAGE OF GUM ACACIA

Take of

Gum Acacia, ¹ in small pieces	. 13 ounces.
Distilled Water	. . . 1 pint.

Put the gum and water into a covered earthenware jar and stir them frequently until the gum is dissolved. If necessary, strain the solution through muslin.

Actions and Uses.—Demulcent and emollient to allay irritation of the alimentary canal, whether caused by disease or poison ; also injected into the bladder in inflammation of that organ and of the kidneys.

Dose.—*Ad libitum*.

MUCILAGO AMYLI²

MUCILAGE OF STARCH

Take of

Starch, in powder $\frac{1}{2}$ ounce.
Water 1 pint.

¹ The much cheaper and equally efficacious dextrin, or British gum, may be substituted for gum acacia in this preparation.

² For most veterinary purposes ordinary gruel may be used instead of this preparation.

Triturate the starch with the water, gradually added ; then boil for a few minutes, constantly stirring.

Use.—In preparing enemas.

MYLABRIS

MYLABRIS

Synonyms.—Mylabris Cichorii ; Chinese Blistering Fly.

Natural Order.—*Coleoptera*.

Composition.—Its vesicant properties are due to the presence of cantharidin.

Characters.—An insect, found on the flowers of the succory plants in India and China. About an inch and a quarter in length ; sheath-wings black, each presenting anteriorly two almost quadrate brownish-yellow spots ; behind these two brownish-yellow bands, each of which equals about one-sixth of the length of the sheath-wings.

Actions and Uses.—Its physiological actions are the same as those of cantharides, except that it is said not to affect the kidneys when topically applied. Employed in the form of Unguentum Mylabridis as a counter-irritant and vesicant.

Preparation.—Unguentum Mylabridis.

MYRRHA

MYRRH

A gum-resinous exudation from the stem of *Balsamodendron Myrrha*. Collected in Arabia Felix and Abyssinia.

Natural Order.—*Amyridaceæ* or *Burseraceæ*.

Composition.—Its medicinal activity depends upon the resin and volatile oil which it contains ; they are both per-

fectly soluble in rectified spirit, but only partially so in proof spirit.

Characters.—In irregularly shaped tears or masses, varying much in size, slightly translucent, of a reddish-yellow or reddish-brown colour ; fractured surface, irregular, and somewhat oily ; odour agreeable and aromatic ; taste acrid and bitter.

Actions and Uses.—*Internally* :—Stimulant and tonic. It is sometimes given in dyspepsia depending upon debility, but recommended chiefly for cattle in cases of chronic cough ; for the latter purpose it is usually combined with opium. *Externally* :—As an excitant and deodoriser to wounds and indolent and fœtid ulcers.

Doses.—HORSE, 2 to 4 drachms.—CATTLE, $\frac{1}{2}$ to 1 ounce.—SHEEP, 1 to 2 drachms.—PIG, 1 to 2 drachms.—DOG, 10 to 30 grains.

Modes of Application.—*Internally* :—In the form of bolus ; pill ; Tinctura Myrrhæ, Tinctura Aloës Composita. *Externally* :—Powder ; Tinctura Myrrhæ ; Tinctura Aloës Composita.

Preparations.—Tinctura Aloës Composita ; Tinctura Myrrhæ.

NUX VOMICA

NUX VOMICA

The seeds of *Strychnos Nux-vomica*. Imported from the East Indies.

Natural Order.—*Loganiaceæ*.

Composition.—Contains the highly poisonous alkaloids strychnine, $C_{21}H_{22}N_2O_2$, and Igasurin, and an almost inert alkaloid, brucine, in combination with igasuric acid.

Characters.—Nearly circular and flat, about an inch in diameter, umbilicated and slightly convex on one side ; externally of an ash-grey colour, thickly covered with short satiny hairs ; internally translucent, tough, and horny ; taste intensely bitter ; inodorous.

Actions and Uses.—In excessive doses, highly poisonous ; in medicinal doses, it is given as a nervous stimulant in general paralysis, paraplegia, and amaurosis ; also as a tonic in chorea and affections of the alimentary canal, *e.g.* dyspepsia, dysentery, colic arising from lead poisoning, &c.

Doses.—HORSE, 30 to 60 grains.—CATTLE, 60 to 120 grains.—SHEEP, 5 to 15 grains.—PIG, 5 to 15 grains.—DOG, $\frac{1}{2}$ to 3 grains.¹

Modes of Application.—The powdered seed made into a bolus or pill ; Extractum Nucis Vomicae, or Tinctura Nucis Vomicae.

Antidotes.—Decoctum Tabaci.

Preparations.—Strychnia ; Tinctura Nucis Vomicae.

OLEUM ANISI

OIL OF ANISE

Composition.—A mixture of two distinct volatile oils, one of which solidifies at temperatures below 50° F., and is named *anethol* or *anise camphor*, $C_{10}H_{12}O$, while the other remains fluid at all temperatures. The constitution of the latter oil is not known, but it is believed to be isomeric with oil of turpentine.

¹ As the powdered drug varies very much in alkaloidal strength it should be avoided in dog practice, and Liquor Strychninae Hydrochloratis used in its place.—ED.

Mode of Preparation.—In Europe by distilling the fruit of *Pimpinella Anisum*, and in China the fruit of *Illicium anisatum*, with water.

Characters.—Colourless or pale yellow, with the odour of anise, and a warm sweetish taste. Completely volatilised by heat; congeals at temperatures between 50° and 60°F. (10° to 15·5° C.).

Uses.—For flavouring medicines, especially masses; also for destroying pediculi on pet dogs and other small animals.

OLEUM CAJUPUTI

CAJUPUT OIL

Composition.—According to Blanchet, the composition is $C_{10}H_{18}O$.

Mode of Preparation.—By distilling the leaves of *Melaleuca Cajuputi* (*Myrtaceæ*). Imported from Batavia and Singapore.

Characters.—Pale bluish or myrtle green colour. It is transparent, limpid, of a strong penetrating agreeable odour resembling that of camphor, rosemary, and cardamoms combined, and of a warm aromatic camphoraceous taste, succeeded by a sensation of coldness in the mouth. Specific gravity varies from 0·914 to 0·930. Oil of Cajuput is soluble in alcohol.

Actions and Uses.—*Internally* :—A powerful diffusible stimulant, antispasmodic, and diaphoretic. Useful in painful spasmodic affections of the stomach, and in flatulent colic. *Externally* :—As an application in chronic rheumatism.

Doses.—Same as *Oleum Terebinthinæ*.

Modes of Application.—*Internally* :—In the form of an Emulsion. *Externally* :—Mixed with olive oil

OLEUM CAMPHORATUM

CAMPHORATED OIL

Take of

Camphor	1 part.
Olive Oil	4 parts.

Dissolve the camphor in the oil.

Uses.—Discutient and anodyne for tumours, bruises, and chronic sprains.

OLEUM CANTHARIDIS

OIL OF CANTHARIDES

Take of

Cantharides, in powder	1 ounce.
Olive Oil	8 fluid ounces.

Digest over a water-bath for two or three hours, and filter for use.

Use.—To maintain the action of blisters.

OLEUM CARYOPHYLLI

OIL OF CLOVES

The oil of the clove (*Eugenia caryophyllata*) distilled in Britain.

Characters.—Colourless or pale yellow when fresh, becomes reddish brown on keeping. Aromatic odour and pungent taste.

Uses.¹—*Internal*:—Carminative, antispasmodic, stoma-
chic, and corrective. Frequently given with purgatives
to prevent griping. *External*:—Used to arrest the pain
and check the flow of synovia in open joints and as a dress-
ing for wounds.

Doses.—HORSE, 20 to 30 minims.—DOG, 1 to 4 minims.

¹ Used in the preparation of the 10 to 20 per cent. solution of Cocaine, for local anæsthesia.

OLEUM CROTONIS

CROTON OIL

Composition.—Principally stearin, palmitin, and two compounds allied to olein ; myristic, lauric, crotonic, and angelic acids have also been obtained from it. Its irritant effects upon the skin are attributed to a body called crotonal, $C_9H_{14}O_2$; but the exact nature of its purgative principle is still unknown.

Mode of Preparation.—By pressure from the kernels of the seeds of *Croton Tiglium*, belonging to the natural order *Euphorbiaceæ*.

Characters.—Fixed, viscid ; colour brownish-yellow ; taste acrid ; odour faintly nauseous. The oil expressed in England is soluble in an equal volume of rectified spirit of wine. Indian croton oil, agitated with cold rectified spirit, forms a milky-looking emulsion, which becomes transparent on the application of heat, but which, on cooling and standing, allows the oil to separate and subside. Croton oil is readily dissolved by sulphuric ether and by the fixed and volatile oils.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, drastic purgative. Given in obstinate constipation ; in torpidity of the bowels dependent on a disordered state of the nervous system, as in tetanus and parturient apoplexy ; to produce copious fluid evacuations and excite extensive counter-irritation in passive dropsies and in local inflammation in parts remote from the alimentary canal ; in anasarca swellings and effusions into cavities, when purgatives are admissible ; and in affections of the kidneys when it is desirable to avoid irritating them. Also valuable for the horse in cases in which

neither a bolus nor a draught can be given, as when the animal is unmanageable or unable to swallow. *Externally* : Vesicant and counter-irritant. Applied to all animals, but especially to cattle, in bronchitis, pleurisy, pneumonia, chronic rheumatism, and glandular and other indolent swellings.

Doses.—HORSE, 10 to 20 minims.—CATTLE, 30 to 60 minims.—SHEEP, 2 to 5 minims.—PIG, 2 to 5 minims.—DOG, $\frac{1}{2}$ to 3 minims, usually given with other purgatives.

Modes of Application.—*Internally* : Mixed with linseed meal so as to form a bolus, dissolved in olive or linseed oil or put into the animal's mash or food. The oil may be placed upon the tongue of horses that refuse their provender, or that are vicious or incapable of swallowing. To remove constipation in cattle, the oil is frequently given in conjunction with sulphate of magnesia or calomel. *Externally* :—One part of croton oil mixed with from four to eight parts of soap liniment, or Linimentum Terebinthinæ et Olei Crotonis, or in oil or gruel.

Antidotes.—Where available, an emetic of sulphate of copper, demulcent drinks, and opium to check the purgation.

Preparation.—Linimentum Terebinthinæ Compositum.

OLEUM EUCALYPTI

EUCALYPTUS OIL

Composition.—A volatile oil containing Eucalyptol, Terpene, and Cymol.

Actions and Uses.—A good disinfectant ; applied to the skin it is an irritant. The vapour of the oil is useful in the treatment of influenza and bronchitis.

OLEUM JUNIPERI

OIL OF JUNIPER

Composition.— $C_{10}H_{16}$.

Mode of Preparation.—In Britain, by distilling the unripe fruit of *Juniperus communis*, belonging to the natural order *Coniferae*, with water.

Characters.—Colourless or pale greenish-yellow, sweetish odour, and warm aromatic taste. Completely volatilised by heat.

Actions and Uses.—Stomachic and diuretic. Given as a diuretic.

Doses.—HORSE, 1 drachm.—CATTLE, 2 drachms.—SHEEP, 15 minims.—PIG, 10 minims.—DOG, 2 to 5 minims.

Mode of Application.—Suspended in water or some mucilaginous drink.

OLEUM LINI

LINSEED OIL

Composition.—Linolein, palmitin, and possibly stearin.

Mode of Preparation.—By expression, without heat, from linseed, belonging to the natural order *Linaceae*.

Characters.—A fixed oil, viscid, yellow, faint odour, and oleaginous taste.

Actions and Uses.—*Internally* :—Nutritive, cathartic, and emollient ; chiefly employed, either alone or conjoined with sulphate of magnesia, as a laxative for cattle and sheep. Its activity may be increased, if desirable, by the

addition of croton oil. Given in cases of intestinal irritability arising from natural causes or from the action of poison ; in colic and diarrhœa, where saline and other vegetable cathartics have proved inactive, or where their repetition is inexpedient ; to give relief in cases of choking ; also injected into the bladder or rectum to allay irritation. *Externally* :—In skin affections, as an emollient to hard and dry surfaces, and as a soothing application in cases of cutaneous and other superficial irritability. Likewise often used in the preparation of liniments instead of the non-siccative, and consequently preferable, olive oil.

Doses.—HORSE, 1 to 2 pints.—CATTLE, 1 to 2 pints.—SHEEP, 3 to 6 fluid ounces.—PIG, 3 to 6 fluid ounces.—DOG, 1 to 3 fluid ounces.

OLEUM MENTHÆ PIPERITÆ

. OIL OF PEPPERMINT.

Composition.—The oil contains two bodies, the fluid Menthene, $C_{10}H_{18}$, and the solid crystalline Menthol, $C_{10}H_{20}O$.

Characters and Tests.—A colourless or pale yellow oil, which has a warm aromatic taste.

Actions and Uses.—Antiseptic, carminative, and parasiticide. Used to prevent the griping effects of purgatives and to flavour medicines.

Doses.—HORSES and CATTLE, 20 to 30 minims.—DOGS, 3 to 5 minims.

The solid menthol is used in the form of a pencil to relieve neuralgic and other pains.

OLEUM MORRHUÆ**COD-LIVER OIL**

Synonym.—Oleum Jecoris Aselli.

Composition.—Chiefly consists of olein, palmitin, and stearin, with small quantities of free butyric and acetic acids, biliary constituents, gaduine, trimethylamine and other peculiar substances, iodine, bromine, sulphur, phosphorus, and about one per cent. of mineral salts.

Mode of Preparation.—Although directed by the 'British Pharmacopœia' to be obtained from the cod *Gadus Morrhua*, it is also procured from other allied species, *e.g.* ling, *Gadus lota*; the dorse, *Gadus callarias*; the torsk, *Gadus Brosme*, &c., belonging to the natural order *Acipenser*. The livers are exposed to the sun and allowed to putrefy; the oil runs from them, and is collected in vessels placed for its reception. Imported principally from Newfoundland and the north of England.

Characters.—Fixed, viscid, yellow colour, faint odour, and oleaginous taste.

Actions and Uses.—Nutritive and alterative in debility accompanied with emaciation, chronic rheumatism, scrofula, phthisis pulmonalis, and other diseases affecting the respiratory organs, chronic skin affections, and distemper in dogs.

Doses.—HORSE, 8 fluid ounces.—CATTLE, 8 fluid ounces.—SHEEP, 2 to 4 fluid ounces.—PIG, 2 to 4 fluid ounces.—DOG, $\frac{1}{4}$ to 1 fluid ounce.

Twice a day for some considerable period, and gradually increased to double the dose originally given.

Modes of Application.—Administered alone or in milk or gruel, flavoured (if necessary) with an aromatic. In dog practice it may be given in capsules, or in biscuits.

OLEUM OLIVÆ¹

OLIVE OIL

Composition.—Chiefly olein and palmitin.

Mode of Preparation.—In the south of Europe by pressing the fleshy portion or pericarp of the ripe fruit of the olive, *Olea europæa*, belonging to the natural order *Oleaceæ*, in a mill.

Characters.—Fixed, pale-yellow colour; scarcely any odour; and a bland oleaginous taste. It partially congeals at about 36°F.

Actions and Uses.	} Same as Oleum Lini.
Doses.	

OLEUM PALMÆ

PALM OIL

Synonym.—Palm Butter.

Composition.—Tripalmitin, $C_3H_5(C_{16}H_{31}O_2)_3$, with a small quantity of olein, $C_3H_5(C_{18}H_{33}O_2)_3$.

Mode of Preparation.—The kernels of the fruit of certain kinds of palm (*Cocos butyracea* or *Arvira Elais*, or both) are crushed and heated with water. The oil rises to the surface in the liquid condition, and is removed.

Characters.—In this country it usually has the consistency of butter. The fresh oil melts at about 76 ° F. Colour usually yellow, which can be removed by bleach-

¹ Linseed oil, rape oil, and whale oil are frequently used as substitutes for olive oil in veterinary pharmacy.

ing ; odour peculiar and agreeable, said to resemble that of violets.

Actions and Uses.—Substitute for lard as an emollient, and in the preparation of ointments.

OLEUM RICINI

CASTOR OIL

Composition.—Principally ricinoleate of glyceryl with other fixed oils, a resin, and possibly an alkaloid, ricinine, and a drastic purgative principle.

Mode of Preparation.—That imported from the West Indies, and the finer qualities from the East Indies, are procured by submitting the seeds of *Ricinus communis*, belonging to the natural order *Euphorbiaceae*, after the removal of their coats, to pressure, with or without heat. Most of the East Indian oil, however, is obtained by boiling the seeds in water, drying and crushing them, and again boiling them until the oil separates and floats on the surface. Cold-drawn castor oil is considered to possess the finest quality.

Actions and Uses.—Cathartic, chiefly for the dog. For other animals linseed oil is usually substituted as an oleaginous purgative.

Doses.—HORSE, 1 to 1½ pint.—CATTLE, ¼ to 1½ pint.—SHEEP, 2 to 3 fluid ounces.—PIG, 2 to 3 fluid ounces.—DOG, ½ to 2 fluid ounces.

Modes of Application.—Usually administered alone. It is, however, sometimes given with linseed or olive oil, or gruel and aromatics, to diminish its irritant effects ; to increase its cathartic activity it is combined with small doses of oil of turpentine or of croton.

Preparations.—Collodium Flexile ; Mistura Ricini.

OLEUM TEREBINTHINÆ**OIL OF TURPENTINE**

Synonyms.—Spirits of Turpentine, Terebene.

Composition.— $C_{10}H_{16}$.

Mode of Preparation.—The outer bark near the root is removed from *Pinus palustris*, *Pinus tæda*, *Pinus pinaster*, as well as from other varieties of pines. Incisions are then made through the inner bark into the wood, from which a mixture of volatile oil and resin (crude and common turpentine) flows into pits dug in the earth, whence it is transferred to casks. By distillation, with or without water, the volatile oil (oil or spirit of turpentine) is separated from the resin.

Characters.—Volatile, limpid, colourless ; strong peculiar odour ; pungent and bitter taste. Imported from France and America.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses astringent, stimulant, cathartic, and diaphoretic. Chiefly employed as a diuretic, antispasmodic, and anthelmintic. *Externally* :—Vesicant, counterirritant, stimulant, and digestive. Principally applied to cattle, in conjunction with mustard or ammonia, in the treatment of inflammation of the bowels or lungs, and in chronic rheumatism.

Doses.—*Diuretic* :—HORSE, 4 to 8 fluid drachms.—CATTLE, 4 to 8 fluid drachms.—SHEEP, 10 to 15 minims.—PIG, 10 to 15 minims.

Antispasmodic :—HORSE, 1 to 2 fluid ounces.—CATTLE, 1 to 2 fluid ounces.—SHEEP, 20 to 60 minims.—PIG, 20 to 60 minims.

Anthelmintic :—HORSE,¹ 2 to 3 fluid ounces.—CATTLE,¹ 2 to 3 fluid ounces.—SHEEP,¹ 2 to 4 fluid drachms.—PIG,¹ 2 to 4 fluid drachms.—DOG,¹ 1 to 1½ drachm.

Modes of Application.—*Internally* :—Dissolved in linseed, olive, or other fixed oil ; beaten into an emulsion with yolk of egg (one yolk to every two drachms of oil of turpentine) ; or Enema Terebinthinæ. *Externally* :—As a rubefacient or digestive, the oil alone ; Linimentum Terebinthinæ ; Linimentum Terebinthinæ Compositum. As a counterirritant for cattle, Cataplasma Terebinthinæ Compositum ; for the dog, Cataplasma Terebinthinæ.

Preparations.—Cataplasma Terebinthinæ ; Cataplasma Terebinthinæ Compositum ; Enema Terebinthinæ ; Linimentum Terebinthinæ ; Linimentum Terebinthinæ Compositum.

OPIUM

OPIUM

Composition of Opium.—Opium contains about 25 per cent. of a gummy substance, 20 per cent. of ill-defined organic matters, a little caoutchouc, resin, oil, and water, and variable proportions of a large number of alkaloids, of which morphine, narcotine, and Narceine are the most abundant. Subjoined is a list of the opium alkaloids :

Hydrocotarnine, $C_{12}H_{15}NO_3$	Papaverine, $C_{21}H_{21}NO_4$
Morphine, $C_{17}H_{19}NO_3$	Meconidine, $C_{21}H_{23}NO_4$
Oxymorphine, $C_{17}H_{19}NO_4$	Laudanosine, $C_{21}H_{27}NO_4$
Codeine, $C_{18}H_{21}NO_3$	Cryptopine, $C_{21}H_{23}NO_5$
Thebaine, $C_{19}H_{21}NO_3$	Narcotine, $C_{22}H_{23}NO_7$
Codamine } $C_{20}H_{25}NO_4$	Lanthopine, $C_{23}H_{25}NO_4$
Laudanine }	Narceine, $C_{23}H_{29}NO_9$
Protopine, $C_{30}H_{19}NO_5$	

¹ Combined with a small dose of castor oil, linseed oil, or solution of aloes.

These alkaloids are present principally as meconates and sulphates.

Mode of Preparation.—Incisions, by means of an instrument consisting of four or five heart-shaped blades tied together with thread, are made in the unripe capsules of the poppy *Papaver somniferum*, belonging to the natural order *Papaveraceæ*. A white milky juice exudes from the incisions in drops, and is allowed to remain on the poppy head for twenty-four hours in order that it may thicken. The inspissated exudation is then scraped off, and the different collections stirred together. The opium is finally dried without the aid of artificial heat, usually by exposure to the sun, and made into cakes or masses, which are in most places wrapped in poppy leaves to prevent their adhering to one another. Chiefly imported from Turkey, Egypt, and India.

Characters.—Irregular lumps, weighing from four ounces to two pounds ; enveloped in the remains of poppy leaves, and generally covered with the chaffy fruits of a species of *Rumex* ; when fresh, plastic, tearing with an irregular, slightly moist, chestnut-brown surface, shining when rubbed smooth with the finger ; peculiar odour and bitter taste.

Actions and Uses.—*Internally* :—In excessive doses, narcotic poison ; in medicinal doses, stimulant, sedative, narcotic, anodyne, and antispasmodic. Given in gastritis, diarrhœa, dysentery, enteritis, colic, peritonitis, pleurisy, bronchitis, pneumonia, tetanus, rheumatism, and very many other maladies. *Externally* :—Anodyne to wounds and blistered and cauterised surfaces ; in cases of superficial inflammation of the eye, skin, or joints ; and in the treatment of hæmorrhoids.

Doses.—HORSE, 1 to 2 drachms.—CATTLE, 2 to 4 drachms.—SHEEP, 10 to 40 grains.—PIG, 10 to 40 grains.—DOG, $\frac{1}{2}$ to 3 grains.

Modes of Application.—*Internally* :—In the form of bolus, Tinctura Opii, or Tinctura Opii Ætherea ; also Enema Opii. Although frequently given alone, it is often combined with belladonna, aconite, tartar emetic, or calomel when required to act as a sedative ; and with chloroform, sulphuric ether, or nitrous ether when employed as a stimulant or antispasmodic. As an injection, as well as by the mouth,¹ in enteritis and inflammation of the bladder and kidneys. *Externally* :—Liquor Plumbi Subacetatis et Opii ; Linimentum Opii ; Tinctura Opii.

Incompatibles.—Lime-water ; alkalies, and their carbonates ; the acetate of lead ; sulphates of iron, zinc, and copper ; perchloride of mercury ; arsenites of potash and soda ; and all vegetable astringents.

Antidotes.—Where possible, empty the stomach by means of emetics or the stomach-pump ; administer brandy, ammonia, carbonate of ammonia, or some other stimulant ; apply ammonia or strong acetic acid to the nostrils ; dash cold water over the body, and endeavour to keep the patient moving about.

Preparations.—Enema Opii ; Linimentum Opii ; Liquor Plumbi Subacetatis et Opii ; Pulvis Ipecacuanhæ Compositus ; Pulvis Opii Compositus ; Tinctura Opii.

PARAFFINUM

PARAFFIN

Synonym.—Tar-oil Stearin.

Composition.—A mixture of several solid hydrocarbons.

Mode of Preparation.—Paraffin is found native in the coal-measures and other bituminous strata constituting

¹ Two or three times as much opium may be administered *per rectum* as would be given by the mouth.

the minerals known as fossil wax, ozocerite, &c. It exists also in the state of solution in many petroleum springs, and may be separated by distilling the more volatile portion and exposing the remainder to a low temperature.

Characters and Tests.—A colourless, crystalline, fatty substance. Melting at about 113°F. Soluble in ether in all proportions. When melted should dissolve entirely in fixed and volatile oils. Sulphuric acid, chlorine, and nitric acid, below the temperature of 212°F., have no action upon it.

Actions and Uses.—Employed externally. Occasionally used in the manufacture of ointments.

PHENACETINE

PARA-ACETPHENITIDIN

Composition.—A coal-tar product, analogous in composition to acetanilid.

Characters and Tests.—A white crystalline powder, of slightly bitter taste, without odour; soluble in alcohol and glycerin, and sparingly soluble in water.

Actions and Uses.—A nervous sedative and antipyretic, useful in influenza, rheumatism, and pneumonia. Dissolved in glycerine and water it has been found useful in distemper.

Doses.—Dogs, 2 to 8 grains.

PHYSOSTIGMATIS SEMEN

CALABAR BEAN

The seed of the *Physostigma venenosum*.

Natural Order.—*Leguminosae*.

Composition.—It contains the alkaloid physostigmine,

also known as eserine, $C_{15}H_{21}N_3O_2$; it also contains calabrine.

Actions and Uses.—Calabar bean stimulates voluntary and involuntary muscular fibres and paralyses nerve centres. It is very poisonous, causing tetanic convulsions. It contracts the pupil of the eye. Principally employed in veterinary medicine in the treatment of obstinate constipation. A solution of the alkaloid physostigmine is usually employed hypodermically for this purpose.

Doses—of the bean.—HORSES and CATTLE, 15 to 30 grains.—DOGS, $\frac{1}{4}$ to $\frac{1}{2}$ a grain.

Preparations.—Extractum Physostigmatis, Physostigmine or Eserine.¹

PILULA HYDRARGYRI ET FERRI

PILL OF MERCURY AND IRON

Synonyms.—Mercurial Pill with Iron ; Ferruginate Blue Pill.

Preparation.—Take of

Mercury, pure	2 parts.
Hydrated Peroxide of Iron	1 part.
Confection of Roses	3 parts.

Rub together until the globules of mercury are no longer visible.

Use.—Alterative for the horse.²

Dose.— $\frac{1}{2}$ to 1 drachm.

PIMENTA

PIMENTO

Synonym.—Allspice.

The dried unripe berries of the allspice, *Eugenia Pimenta*. Imported from the West Indies.

¹ For actions and uses &c. see Addenda.

² If there be any apprehension of this preparation exciting catharsis, it should be combined with opium.

Natural Order.—*Myrtaceæ*.

Composition.—Therapeutic value dependent on the presence of a volatile oil.

Characters.—Of the size of a small pea, rough, crowned with the teeth of the calyx; contains two dark-brown seeds; externally brown, internally yellowish; odour and taste hot, aromatic, and peculiar.

Actions and Uses.—Aromatic, carminative, stomachic, and antispasmodic. Given in indigestion, relaxed stomach, flatulency, and in colic. Employed also to disguise the flavour of nauseous medicines.

Doses.—HORSE, 2 to 4 drachms.—CATTLE, 2 to 6 drachms.—SHEEP, $\frac{1}{2}$ to 1 drachm.—PIG, $\frac{1}{2}$ to 1 drachm.—DOG, 10 to 30 grains.

Modes of Application.—In the form of powder, bolus, or Tinctura Pimentæ.

Preparation.—Tinctura Pimentæ.

PIPER NIGRUM

BLACK PEPPER

Natural Order.—*Piperaceæ*.

Composition.—Its medicinal properties are due chiefly to a volatile oil, and an alkaloid piperine.

Mode of Preparation.—The berries of *Piper nigrum*, before they have all changed to red, are collected, dried, and ground to powder.

Characters.—Small roundish wrinkled bodies; tegument brownish black, containing a greyish-yellow globular seed (white pepper).

Actions and Uses.—Principally employed as a stomachic and carminative in simple indigestion, and as an aromatic to mask the unpleasant flavour of many medicines.

Doses.—HORSE, 2 drachms.—CATTLE, 3 drachms.—SHEEP, 20 to 40 grains.—Repeated two or three times a day.

Modes of Application.—In the form of bolus, or, preferably, suspended in gruel.

PIX BURGUNDICA

BURGUNDY PITCH

A resinous exudation from the stem of the spruce fir, *Abies excelsa*, belonging to the natural order *Coniferae*, melted and strained. Imported from Switzerland.

Characters.—Hard and brittle, yet gradually taking the form of the vessel in which it is kept ; opaque, varying in colour, but generally dull reddish brown ; of a peculiar, somewhat empyreumatic perfumed odour, and aromatic taste, without bitterness ; free from vesicles ; gives off no water when heated.

Actions and Uses.—*Internally* :—Irritant and diuretic. *Externally* :—Rubefacient and digestive ; stimulant to sprains ; to give support to fractures ; and as an adhesive to wounds.

Modes of Application.—*Externally* :—Melted and applied, after the hair of the part has been loosened by the currycomb, to fractures and sprains. As an adhesive, Emplastrum Picis is used.

PIX LIQUIDA

TAR

Synonym.—Wood Tar.

Composition.—Hydrocarbons, methylic alcohol, carbolic acid, acetic acid, cresylic acid, resinous bodies, pitch, and many other substances.

Mode of Preparation.—By submitting to destructive distillation the wood of *Pinus sylvestris* and other pines.

Characters.—Thick, viscid, brownish-black liquid, of a well-known peculiar aromatic odour. Water agitated with it acquires a pale-brown colour, sharp empyreumatic taste, and acid reaction.

Actions and Uses.—*Externally* :—As an excitant, rubefacient, antiseptic, and deodoriser in the treatment of thrush in the horse, and of broken horns and punctured wounds of the belly and chest of cattle ; foot-rot in sheep ; and in mange, eczema, impetigo, and other skin affections.¹

Modes of Application.—In thrush and foot-rot, tar alone is applied to the diseased parts ; for broken horns and punctured wounds, the tar should be spread over coarse cloth ; for skin affections, Linimentum Olei Picis Liquidæ Compositum or Linimentum Picis Liquidæ is used ; and as a common foot ointment for all domesticated animals, Unguentum Picis Liquidæ is employed.

Preparations.—Linimentum Olei Picis Liquidæ Compositum ; Linimentum Picis Liquidæ ; Unguentum Picis Liquidæ.

¹ A ' *Stopping for Feet* ' is made by mixing together—

Tar	2 parts.
Soft Soap	1 part.
Linseed Meal	{ a sufficiency to impart tenacity to the mixture.

PIX NIGRA

PITCH

Synonym.—Black Pitch.

The black, resinoid, and highly carbonaceous residue which is left when the volatile constituents of tar are removed by distillation.

Uses.—Mild stimulant in thrush, canker, and sand-crack in horses, and foot-rot in sheep. Also used to impart consistency and adhesiveness to plasters and ‘charges.’

PLUMBI ACETAS

ACETATE OF LEAD

Synonyms.—Lead Acetate ; Sugar of Lead.

Composition.— $\text{Pb}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 3\text{Aq.}$


Mode of Preparation.—Take of

Oxide of Lead (Litharge, } 24 ounces.
in fine powder . . . }

Acetic Acid . . . 2 pints, or a sufficiency.

Distilled Water . . . 1 pint.

Mix the acetic acid and the water, add the oxide of lead, and dissolve with the aid of a gentle heat. Filter, evaporate till a pellicle forms, and set aside to crystallise, first adding a little acetic acid should the fluid not have a distinctly acid reaction. Drain, and dry the crystals on filtering paper without heat.



In the act of crystallising the salt unites with three molecules of water.

Characters and Tests.—White crystalline masses, slightly efflorescent ; acetous odour, and sweet astringent taste. It dissolves in distilled water, and forms a clear solution, or one which has only a slight milkiness, which disappears on the addition of acetic acid. Its solution in water slightly reddens litmus, gives a yellow precipitate (lead iodide) with potassium iodide, and a white precipitate (lead sulphate) with dilute sulphuric acid, acetic acid being set free and recognisable by its odour ; the precipitates by potassium iodide and sulphuric acid indicate the presence of lead, and the elimination of acetic acid shows that the salt is an acetate. Air blown from the lungs through its aqueous solution should not occasion any turbidity, showing its freedom from subacetate.

Actions and Uses.—Sedative and astringent. Seldom employed in veterinary practice ; it is, however, occasionally administered in from 1 to 2 drachm doses to the horse in diarrhoea and diabetes insipidus. Principally used in the production of *Liquor Plumbi Subacetatis*.

Preparations.¹—*Liquor Plumbi Subacetatis*.


PLUMBI OXIDUM

OXIDE OF LEAD

Synonyms.—Lead Monoxide ; Protoxide of Lead ; Litharge.

Composition.— PbO .

Mode of Preparation.—Metallic lead is oxidised by being heated in a current of air.



¹ *Plumbi acetas* and *zinci sulphas*, dissolved in water, form the well known 'white lotion.'

Characters and Tests.—Heavy scales, pale brick-red colour, completely soluble without effervescence in diluted nitric and acetic acids, either solution, when neutral, giving a yellow precipitate (lead iodide) with potassium iodide, indicating the presence of lead. Its solution in diluted nitric acid, when saturated with ammonia and filtered, does not exhibit a blue colour, showing that it is free from copper.

Uses.—In preparing Plumbi Acetas and Liquor Plumbi Subacetatis.

PODOPHYLLI RHIZOMA

PODOPHYLLUM RHIZOME

Synonym.—Podophylli Radix.

The dried rhizome of *Podophyllum peltatum*, belonging to the natural order *Ranunculaceae*. Imported from North America.

Characters.—In pieces of variable length, about two inches thick ; mostly wrinkled longitudinally ; dark reddish-brown externally, whitish within ; breaking with a short fracture ; accompanied with pale brown rootlets. Powder yellowish-grey, sweetish in odour, bitterish, subacid, and nauseous in taste.

Preparation.—Resina Podophylli.

Actions and Uses.—Cholagogue, purgative ; used as a substitute for calomel. Experiments have shown that podophyllin is not likely to be serviceable as a purgative for the domesticated animals. It is given to lower the heart's action in acute diseases of the respiratory organs ; rheumatism, laminitis, and other inflammatory disorders.

Doses.—As a sedative. HORSE, 1 to 2 drachms.—CATTLE, 1 to 2 drachms.—DOG, 1 to 2 grains.

Modes of Application.—United with calomel, or potassium nitrate, or magnesium sulphate. For dogs it may be administered with calomel, grey powder, or ipecacuanha.

PODOPHYLLI RESINA

RESIN OF PODOPHYLLUM

Take of

Podophyllum Root, in coarse powder .	1 pound.
Rectified Spirit	{ 3 pints, or a sufficiency.
Distilled Water a sufficiency.
Hydrochloric Acid a sufficiency.

Exhaust the podophyllum by percolation with the spirit ; distil over the spirit ; slowly pour the liquid remaining after the distillation of the tincture into three times its volume of water, acidulated with one twenty-fourth part of its weight of hydrochloric acid, constantly stirring ; let it stand twenty-four hours ; collect the resin which falls, wash on a filter with distilled water, and dry in a stove.

Characters and Tests.—A pale, greenish-brown, amorphous powder, soluble in rectified spirit and in ammonia ; precipitated from the former solution by water, from the latter by acids. Almost entirely soluble in pure ether.

POTASSA CAUSTICA

CAUSTIC POTASH

Synonyms.—Potassæ Hydras ; Potassa Fusa ; Potassium Hydrate ; Hydrate of Potash.

Composition.—KOH.

Mode of Preparation.—Take of

Solution of Potash	2 pints.
------------------------------	----------

Boil down the solution of potash rapidly in a silver or clean iron vessel until there remains a fluid of oily consistence, a drop of which, when removed on a warm glass rod, solidifies on cooling. Pour this into proper moulds, and when it has solidified, and while it is still warm, put it into a stoppered bottle.

Characters and Tests.—Hard white pencils, very deliquescent, powerfully alkaline, and corrosive. A watery solution acidulated by hydrochloric acid gives a yellow precipitate (potassium-platinic chloride) with platinic chloride, indicating the presence of potassium. It should dissolve in diluted nitric acid without effervescence, showing the absence of carbonate, and give but scanty white precipitates with barium chloride and silver nitrate, showing the presence of traces only of sulphate and chloride respectively.

Actions and Uses.—Principally applied as an escharotic to wounds produced by the bites of rabid animals, also for stimulating unhealthy ulcers and inducing suppuration. In consequence of the highly deliquescent nature of this preparation, it has an objectionable tendency to spread beyond the part upon which it is required to act. The use of Soda Caustica is not open to this objection, and, as its escharotic effects rival those of Potassa Caustica, it may be advantageously substituted for the latter agent.

POTASSA SULPHURATA

SULPHURATED POTASH

Synonyms.—Potassii Sulphuratum ; Hepar Sulphuris ; Sulphide of Potassium ; Liver of Sulphur.

Composition.—A mixture of several sulphides, sulphates, and other compounds of potassium.

Mode of Preparation.—Take of

Carbonate of Potash . . .	10 ounces.
Sublimed Sulphur . . .	5 ounces.

Mix the carbonate of potash and the sulphur in a warm mortar, and having introduced them into a Cornish or Hessian crucible, let them be heated, first gradually, until effervescence has ceased, and finally to dull redness, so as to produce perfect fusion. Let the liquid contents of the crucible be then poured out on a clean flagstone, and covered quickly with an inverted porcelain basin, so as to exclude the air as completely as possible while solidification is taking place. The solid product thus obtained should, when cold, be broken into fragments, and immediately enclosed in a green glass bottle furnished with an air-tight stopper.

Characters and Tests.—Solid greenish fragments, liver-brown when recently broken, alkaline and acrid to the taste, readily forms with water a yellow solution which has the odour of sulphuretted hydrogen, and evolves it freely when excess of hydrochloric acid is dropped into it, sulphur being at the same time deposited. The acid fluid, when boiled and filtered, is precipitated yellow (potassium-platinic chloride) by platinic chloride, showing the presence of potassium, and white by barium chloride, indicating the presence of sulphate. About three-fourths of its weight is dissolved by rectified spirit.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in doses of from half to one drachm to the horse it acts as a stimulant and diaphoretic. Very seldom employed as an internal remedy in this country. *Externally* :—In chronic skin diseases, especially those of a scaly character.

Modes of Application.—*Externally* :—Liquor Potassæ Sulphuratæ ; or Unguentum Potassæ Sulphuratæ.

Antidotes.—Solution of chlorinated lime, with an emollient drink.

Preparations.—Liquor Potassæ Sulphuratæ ; Unguentum Potassæ Sulphuratæ.

POTASSÆ BICARBONAS

BICARBONATE OF POTASH

Synonym.—Potassium-hydrogen Carbonate.

Composition.— KHCO_3 .


Mode of Preparation.—Take of

Carbonate of Potash . . .	1 pound.
Distilled Water . . .	2 pints.
Hydrochloric Acid . . .	$1\frac{1}{2}$ pint.
Common Water . . .	3 pints.
White Marble, in fragments	{ 1 pound, or a sufficiency.

Dissolve the carbonate of potash in the distilled water, and filter the solution into a three-pint bottle capable of being tightly closed by a cork traversed by a glass tube sufficiently long to pass to the bottom of the fluid. Introduce the marble into another bottle, in the bottom of which a few small holes have been drilled, and the mouth of which is closed by a cork also traversed by a glass tube, and place the bottle in a jar of the same height as itself. Join the two glass tubes, so that the connection is air-tight, by a caoutchouc tube. The cork of the bottle containing the carbonate of potash having been placed loosely, and that of the bottle containing the marble tightly, in its mouth, pour into the jar surrounding the latter bottle the hydrochloric acid, previously diluted with the common water.

When carbonic anhydride has passed through the carbonate of potash solution for two minutes, so as to expel the

whole of the air of the apparatus, fix the cork tightly in the neck of the bottle, and let the process go on for a week. At the end of this time numerous crystals of bicarbonate of potash will have formed, which are to be removed, shaken with twice their bulk of cold distilled water, and, after decantation of the water, drained, and dried on filtering-paper by exposure to the air. The mother-liquor, filtered if necessary, and concentrated to one-half, at a temperature not exceeding 110°F., will yield more crystals. The tube immersed in the solution of carbonate of potash, which should have as large a diameter as possible, may require the occasional removal of the crystals formed within it, in order that the process may not be interrupted.



Characters and Tests.—Colourless right rhombic prisms, not deliquescent ; saline, feebly alkaline, taste ; not corrosive. The addition of diluted hydrochloric acid causes strong effervescence (escape of carbonic anhydride), and forms a solution which gives a yellow precipitate (potassium-platinic chloride). An aqueous solution gives no precipitate with magnesium sulphate, owing to the formation of soluble magnesium-hydrogen carbonate.

Actions and Uses.—In excessive doses, irritant poison ; in medicinal doses, antacid and diuretic.

Doses.—HORSE, 2 to 6 drachms.—CATTLE, 2 to 6 drachms. SHEEP, 40 grains to 2 drachms.—PIG, 40 grains to 2 drachms.—DOG, 10 to 30 grains.

Modes of Application.—In the form of bolus or dissolved in water.

Incompatibles.—Acids ; lime-water ; acetate and chloride of ammonium ; most metallic salts, but not sulphate of magnesia.

Antidotes.—Highly diluted acetic acid or vinegar, and linseed or olive oil.

POTASSÆ CARBONAS

CARBONATE OF POTASH

Synonym.—Potassium Carbonate.

Composition.— K_2CO_3 , with about 16 per cent. of water of crystallisation.

Mode of Preparation.—In the United States, Canada, Russia, and on the shores of the Baltic, various kinds of wood are burned, and their ashes collected and lixiviated. By lixiviation various potassium compounds—*e.g.* carbonate, sulphate, chloride, phosphate, and silicate—are dissolved, while silica and certain salts of calcium, magnesium, iron, and aluminum remain insoluble. The soluble portions of the ash are evaporated to dryness in iron pots, and deprived of their volatile impurities by being heated in an oven-shaped furnace, whose flame is made to play over the mass. The product thus obtained is impure carbonate of potash, and is commercially termed pearlash. In order to obtain medicinal carbonate of potash, pearlash (to dissolve the carbonate and leave undissolved the other potassium salts) is treated with its own weight of distilled water, and the solution so formed evaporated to dryness, while it is kept briskly agitated.

Characters and Tests.—White crystalline powder, alkaline and caustic to the taste, very deliquescent, readily soluble in water, but insoluble in spirit of wine. It effervesces with dilute hydrochloric acid, showing that it is a

carbonate, and forms a solution which gives a yellow precipitate (potassium-platinic chloride) with platinic chloride. Loses about 16 per cent. of its weight (water) when exposed to a red heat. Its aqueous solution gives a white precipitate (magnesium carbonate) with solution of magnesium sulphate ; it is thus distinguished from Potassæ Bicarbonas. If dissolved in water and mixed with excess of nitric acid, it should give but a faint precipitate on the addition of barium chloride and silver nitrate, indicating its almost perfect freedom from sulphates and chlorides.

Actions and Uses.—Same as Potassæ Bicarbonas, but more liable to irritate the stomach and intestines than that agent.

Doses.—HORSE, 2 to 4 drachms.—CATTLE, 2 to 4 drachms. SHEEP, 30 grains to $1\frac{1}{2}$ drachm.—PIG, 30 grains to $1\frac{1}{2}$ drachm.—DOG, 5 to 20 grains.

Modes of Application.—In the form of bolus, or dissolved in water.

Incompatibles.—Same as with Potassæ Bicarbonas.

Antidotes.—Same as for Potassæ Bicarbonas.

POTASSII BROMIDUM

BROMIDE OF POTASSIUM

Synonym.—Potassium Bromide.

Composition.—KBr.


Mode of Preparation.—Take of

Solution of Potash	.	.	.	2 pints.
Bromine	.	.	.	4 fluid ounces.
Wood Charcoal, in fine powder	.	.	.	2 ounces.
Boiling Distilled Water	.	.	.	$1\frac{1}{2}$ pint.

Put the solution of potash into a porcelain vessel, and add the bromine in successive portions, with agitation, until the mixture has acquired a brown tint. Evaporate to dryness, reduce the residue to a fine powder, and mix this intimately with the charcoal. Throw the mixture, in small quantities at a time, into a red-hot crucible, and when the whole has been brought to a state of fusion remove the crucible from the fire and turn out its contents.

When the fused mass has cooled, dissolve it in the water, filter the solution through paper, and set it aside to crystallise. The crystals should be dried at a gentle heat.

The chemical reaction occurring in this process takes place in two stages :



Characters and Tests.—In colourless cubical crystals with no odour, but a pungent, saline taste. Soluble in water, 1 in 2 ; less soluble in spirit. Its aqueous solution gives a white crystalline precipitate with tartaric acid (KHC₄H₄O₆), proving the presence of potassium. When its solution in water is mixed with a little chlorine, chloroform agitated with it, on falling to the bottom, exhibits a red colour (bromine). Silver nitrate will also give a yellowish-white precipitate, insoluble in dilute nitric acid ; dissolving sparingly in ammonium hydrate, but readily in potassium cyanide.

Actions and Uses.—As a sedative. It exerts a powerful influence on the generative organs, lowering their functions in a marked degree.

Doses.—HORSE, 2 to 4 drachms.—CATTLE, 2 to 5 drachms.—DOGS, 5 to 20 grains. Repeated when required.

POTASSÆ CHLORAS

CHLORATE OF POTASH


Synonym.—Potassium Chlorate.


Composition.— KClO_3 .

Mode of Preparation.—Take of

Carbonate of Potash	.	.	20 ounces.
Slaked Lime	.	.	53 ounces.
Distilled Water	.	.	a sufficiency.
Black Oxide of Manganese	.	.	83 ounces.
Hydrochloric Acid	.	.	24 pints.

Mix the lime with the carbonate of potash, and triturate them with a few ounces of the water, so as to make the mixture slightly moist. Place the oxide of manganese in a large retort or flask, and, having poured upon it the hydrochloric acid, diluted with six pints of water, apply a gentle heat, and conduct the chlorine as it comes over, first through a bottle containing six ounces of water, and then into a large carboy containing the mixture of carbonate of potash and slaked lime. When the whole of the chlorine has come over, remove the contents of the carboy, and boil them for twenty minutes with seven pints of the water, filter and evaporate till a film forms on the surface, and set aside to cool and crystallise. The crystals thus obtained are to be purified by dissolving them in three times their weight of boiling distilled water and again allowing the solution to crystallise.





The potassium chloride crystallises out, while the calcium chloride remains in solution.

Characters and Tests.—Colourless rhomboidal crystalline plates, cool saline taste, soluble in cold water 1 in 12, boiling water 1 in 2. Its aqueous solution is not affected by silver nitrate or ammonium oxalate. Strongly heated, it fuses, gives off oxygen gas, and leaves a white residue (potassium chloride), which readily dissolves in water, and produces a solution which yields a white precipitate (silver chloride) with silver nitrate, indicating the presence of chlorine, and a yellow precipitate (potassium-platinic chloride) with platinum chloride, showing the presence of potassium. It explodes when triturated in a mortar with sulphur.¹

Actions and Uses.—*Internally* :—Mild stimulant, diuretic, and refrigerant. Given principally in tympanites and hoven, from its having the power, it is said, of causing a condensation of the gases developed in these complaints. *Externally* :—Stimulant and refrigerant to cancerous and other ill-conditioned ulcers.

Doses.—HORSE, 1 to 4 drachms.—CATTLE, 2 to 6 drachms.—SHEEP, 20 to 40 grains.—PIG, 20 to 40 grains. DOG, 5 to 15 grains.

Modes of Application.—*Internally* :—Dissolved in water. *Externally* :—In the form of Liquor Potassæ Chloratis Compositus.

Preparation.—Liquor Potassæ Chloratis Compositus.

¹ To avoid accidents, the particles used in performing this experiment should not be larger than a pin's head.


POTASSÆ NITRAS

NITRATE OF POTASH



Synonyms.—Potassium Nitrate ; Nitre ; Saltpetre.

Composition.— KNO_3 .


Mode of Preparation.—In India and other hot climates putrefying organic matter is naturally, and in some parts of Europe it is artificially, associated with alkaline and earthy carbonates, together with other materials which render the mass porous. Under these conditions the ammonia arising from the substances undergoing putrefaction is transformed into nitric acid at the expense of atmospheric oxygen.



The nitric acid thus formed is converted into calcium and potassium nitrates by contact with calcium and potassium carbonates.



Soil containing nitrates is known as *nitre-* or *saltpetre-earth*, and is the source of large quantities of the nitrate of potash used in medicine and in the arts. The nitre-earth is lixiviated, and the watery solution obtained is mixed with solution of potassium carbonate, to convert any calcium nitrate into the corresponding potassium salt.



By decantation or filtration a clear solution is procured, which, on evaporation, deposits nitrate of potash in a crystalline state. The salt is purified by solution and careful recrystallisation.

Characters and Tests.—White crystalline masses or fragments of striated six-sided prisms, colourless ; peculiar cool saline taste. Soluble in cold water 1 in 4 parts, boiling water 1 in $2\frac{1}{2}$ parts. Thrown on the fire or on red-hot charcoal, deflagration takes place ; warmed in a test-tube with sulphuric acid and metallic copper, red fumes (nitric peroxide) are evolved ; these two reactions show that the salt is a nitrate. Its aqueous solution, acidulated with hydrochloric acid, gives a yellow precipitate (potassium-platinic chloride) with platinic chloride, indicating the presence of potassium. Its aqueous solution should not be affected by barium chloride or silver nitrate, proving its freedom from sulphates and chlorides respectively.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, diuretic, febrifuge, and refrigerant. *Externally* :—Stimulant to unhealthy wounds.

Doses.—HORSE, 2 to 4 drachms.—CATTLE, 2 to 4 drachms.—SHEEP, $\frac{1}{2}$ to 1 drachm.—PIG, $\frac{1}{2}$ to 1 drachm.—DOG, 5 to 10 grains.

Modes of Application.—*Internally* :—As a febrifuge, it is best given, when possible, in the form of bolus ; as a diuretic, it is administered in solution in water. Where nitre is used as a refrigerant, the solution should not be made until just before it is to be given.—*Externally* :—In the form of *Liquor Potassæ Nitratis*.

Incompatibles.—Sulphuric acid and sulphates.

Preparations.—Liquor Potassæ Nitratis ; Liquor Ammonii Chloridi et Potassæ Nitratis.

POTASSÆ PERMANGANAS

PERMANGANATE OF POTASH

Synonym.—Potassium Permanganate.

Composition.— $K_2Mn_2O_8$.

Mode of Preparation.—Take of


Caustic Potash	5 ounces.
Black Oxide of Manganese, in fine Powder	} 4 ounces.
Chlorate of Potash	
Diluted Sulphuric Acid	a sufficiency.
Distilled Water	$2\frac{1}{4}$ pints.

Finely pulverise the chlorate of potash in a perfectly clean mortar, and mix it with the oxide of manganese ;¹ put the mixture into a porcelain basin, and add to it the caustic potash, previously dissolved in four ounces of the water. Evaporate to dryness, stirring diligently to prevent spurt-ing. Pulverise the residual mass, put it into a Hessian or Cornish crucible, and expose it to a dull red heat for an hour, or till it has assumed the condition of a semi-fused green mass (potassium manganate). When cool, pulverise it, and boil it with a pint and a half of the water, for the purpose of converting it into permanganate. Let the insoluble matter (manganese dioxide) subside, decant the fluid, boil again with half a pint of the water, again decant,

¹ The mixture should be effected by stirring with a spatula ; if the manganese and chlorate be ground together with a pestle, a dangerous explosion may occur.

neutralise the united liquors accurately with the diluted sulphuric acid, and evaporate till a pellicle forms. Set aside to cool and crystallise. Drain the crystalline mass, boil it in six ounces of the water, and strain through a funnel, the throat of which is slightly obstructed by a little asbestos. Allow the fluid to cool and crystallise, drain the crystals, and dry them by placing them under a bell-jar over a vessel containing sulphuric acid.

The chemical reactions occurring in this process take place in two stages :



The diluted sulphuric acid is added to neutralise the potassium hydrate liberated in the second reaction.

Characters and Tests.—Dark purple, prismatic crystals, with a sweet astringent taste. Completely dissolved by cold water, forming a rich purple solution.

Actions and Uses.—Applied as an excitant and deodoriser to cancerous and ill-conditioned wounds, and in the treatment of certain skin affections. Principally employed as a deodoriser and disinfectant.

Modes of Application.—To fœtid wounds and ulcers in the form of *Liquor Potassæ Permanganatis*. For ordinary

deodorising and disinfecting purposes, mix a wineglassful of Condyl's fluid (commercial potassium permanganate) with a pailful of water, and well sprinkle it, or place it in dishes, about the room, stable, &c. In the case of an offensively smelling drain the above mixture should be poured down it.

Preparation.—Liquor Potassæ Permanganatis.

POTASSÆ PRUSSIÆ FLAVA

YELLOW PRUSSIAN OF POTASH

Synonyms.—Potassium Ferrocyanide ; Ferrocyanide of Potassium.

Composition.— $K_4FeC_6N_6 \cdot 3Aq$, or $K_4FeCy_6 \cdot 3Aq$.

Mode of Preparation.—Cuttings of horns, skins, hoofs, and dry refuse animal matter of various kinds are fused at a red heat in an iron vessel with potassium carbonate and scraps of iron. The resulting mass is treated with hot water and filtered. On evaporation the filtrate deposits the ferrocyanide, which is purified by recrystallisation. Carbon and nitrogen from the animal matter and potassium from the potassium carbonate unite with the iron to form potassium ferrocyanide.

Characters and Tests.—Yellow crystals derived from the octahedron, permanent in the air, soluble in water, insoluble in alcohol. The aqueous solution gives a deep blue precipitate (ferric ferrocyanide ¹) with a ferric salt ; a precipitate which is at first white (potassio-ferrous ferrocyanide ²), then light blue, and finally, by exposure to the

¹ Fe_7Cy_{18} or $Fe_4Fe_3Cy_{18}$. This compound, plus $18Aq$, constitutes Prussian Blue.

² $K_2Fe_2Cy_6$.

air, dark blue,¹ with a ferrous salt ; and a reddish-brown precipitate (cupric ferrocyanide) with cupric sulphate. Heated with diluted sulphuric acid, hydrocyanic acid vapours are evolved.

Use.—For preparing Acidum Hydrocyanicum Dilutum.


POTASSÆ SULPHAS

SULPHATE OF POTASH

Synonym.—Potassium Sulphate.

Composition.— K_2SO_4 .

Mode of Preparation.—Hydrogen-potassium sulphate, which is left in the retort in the preparation of nitric acid, is dissolved in water and neutralised by potassium carbonate. By evaporating the solution the salt is thrown down, and by recrystallisation it is purified.



Use.—In the preparation of Pulvis Ipecacuanhæ Compositus, to facilitate the intimate mixture, and to prevent the agglutination, of the ingredients.

POTASSII IODIDUM

IODIDE OF POTASSIUM

Synonym.—Potassium Iodide.

Composition.—KI.

Mode of Preparation.—Take of



Solution of Potash 1 gallon.

¹ Contact with air causes the ferrous to become a ferric salt by oxidation and the consequent transformation of white potassio-ferrous ferrocyanide into dark blue ferric ferrocyanide.

Iodine	{ 29 ounces, or a sufficiency.
Wood Charcoal, in fine powder	3 ounces.

Put the solution of potash into a glass or porcelain vessel, and add the iodine in small quantities at a time, with constant agitation, until the solution acquires a permanent brown tint. Evaporate the whole to dryness in a porcelain dish, pulverise the residue (a mixture of potassium iodide and iodate), and mix it intimately with the charcoal. Throw the mixture, in small quantities at a time, into a red-hot iron crucible, and, when the whole has been brought to a state of fusion, remove the crucible from the fire and pour out its contents.

When the fused mass (impure potassium iodide) has cooled, dissolve it in two pints of boiling distilled water, filter through paper, wash the filter with a little boiling distilled water, unite the liquids, evaporate the whole till a film forms on the surface, and set it aside to cool and crystallise. Drain the crystals, and dry them quickly with a gentle heat. More crystals may be obtained by evaporating the mother-liquor and cooling. The salt should be kept in a stoppered bottle. Two decompositions take place in this process :



Characters and Tests.—Colourless, generally opaque, cubic crystals, readily soluble in water, less so in spirit of wine. It commonly has a feeble alkaline reaction. Its solution mixed with mucilage of starch gives a blue colour on the addition of a minute quantity of solution of chlorine

(to set free iodine), thus proving that the salt is an iodide. Its aqueous solution mixed with solution of tartaric acid, or hydrogen-sodium tartrate, furnishes a crystalline precipitate (hydrogen-potassium tartrate), indicating the presence of potassium.

Silver nitrate gives, with an aqueous solution of the salt, a yellowish-white precipitate (silver iodide), which, when agitated with ammonia, yields by subsidence a clear liquid, in which excess of nitric acid should cause no turbidity, showing the absence of chlorides.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, alterative, deobstruent, and diuretic. Given in scrofula, enlargement of the lymphatic glands, periostitis, articular rheumatism, and dropsy.¹ *Externally* :—To promote the absorption of tumours and other abnormal growths. Also employed as a solvent for iodine, which it strongly resembles in therapeutic action.

Doses.—HORSE, 1 to 2 drachms.—CATTLE, 1 to 3 drachms.—SHEEP, 20 grains to 1 drachm.—PIG, 20 grains to 1 drachm.—DOG, 1 to 5 grains. Repeated two or three times a day.

Modes of Application.—*Internally* :—In the form of bolus, or dissolved in water. *Externally* :—Unguentum Potassii Iodidi ; Unguentum Iodidi Compositum.

Incompatibles.—Acids, and Spiritus Ætheris Nitrosi.

Preparations.

Linimentum Iodi Comp.	Tinctura Iodi.
Liquor Iodi et Potassii Iodidi.	Unguentum Iodi.
	Unguentum Iodi Comp.

¹ Potassium Iodide is now largely used in the treatment of actinomycosis and other tumours due to discomyces.

PULVIS ANTIMONIALIS

ANTIMONIAL POWDER

Synonym.—Factitious James's Powder.

Take of

Oxide of Antimony	1 ounce.
Phosphate of Lime	2 ounces.

Actions and Uses.—Similar to those of tartarated antimony, but not so likely to cause local irritation. Employed as a febrifuge for the dog.

Doses.—3 to 10 grains.

PULVIS FERRI SULPHATIS COMPOSITUS

COMPOUND POWDER OF SULPHATE OF IRON

Take of

Sulphate of Iron, in powder .	1 to 1½ drachm.
Gentian, in powder	4 drachms. ¹

Mix.

This mixture forms one dose for the horse, to which animal it is given as a tonic, sprinkled over his food.

PULVIS IPECACUANHÆ COMPOSITUS

COMPOUND POWDER OF IPECACUANHA

Synonym.—Factitious Dover's Powder.

Take of

Ipecacuanha, in powder . . .	1 ounce.
Opium, in powder	1 ounce.
Sulphate of Potash, in powder .	8 ounces.

¹ If it be thought desirable, from 30 to 40 grains of ground ginger may be added to this powder.

Mix thoroughly, pass the powder through a fine sieve, and finally rub it lightly in a mortar. Keep it in a stoppered bottle.

Actions and Uses.—Given to the dog as a nauseant, sedative, in from 5- to 15-grain doses, repeated four or five times a day.

PULVIS OPII COMPOSITUS

COMPOUND POWDER OF OPIUM

Take of

Opium	}	In powder, of each equal parts.
Galls		
Pimento		
Carbonate of Soda		

Mix.

Use.—In diarrhœa occasioned, or accompanied, by a relaxed or irritable condition of the intestines.

Doses.—HORSE, 4 to 6 drachms.—CATTLE, 4 to 6 drachms.—SHEEP, 1 to 1½ drachm.—PIG, 1 to 1½ drachm. Dog, 10 to 30 grains.

Modes of Application.—In the form of bolus, or mixed with gruel.

PULVIS RHEI COMPOSITUS

COMPOUND POWDER OF RHUBARB

Synonym.—Gregory's Powder, or Gregory's Mixture.

Take of

Rhubarb Root, in powder	2 ounces.
Ginger, in powder	1 ounce.
Light Magnesia	6 ounces.

Mix thoroughly, and pass the powder through a fine sieve.

Actions and Uses.—Antacid and stomachic in indigestion.

Doses.—HORSE, 2 to 3 ounces.—CATTLE, 2 to 4 ounces. SHEEP, 1 to 2 drachms.—PIG, 1 to 2 drachms.—DOG, 10 to 30 grains.

Modes of Application.—In the form of bolus, or mixed with gruel.

PULVIS ZINCI CARBONATIS COMPOSITUS

COMPOUND POWDER OF CARBONATE OF ZINC

Take of

Carbonate of Zinc, in powder .	. 4 ounces.
Alum, in powder 1 ounce.

Mix.

Uses.—Excitant and astringent when sprinkled over abrasions, ulcerated heels, open joints, &c.

PYROXYLIN

GUN COTTON

Synonym.—Dinitro-cellulose.

Composition.— $C_{18}H_{21}O_6(OH)(NO_3)_8$.

Mode of Preparation.—Take of

Cotton Wool	1 ounce.
Sulphuric Acid	} of each, 5 fluid ounces.
Nitric Acid	

Mix the acids in a porcelain mortar, immerse the cotton in

the mixture, and stir it for three minutes with a glass rod until it is thoroughly wetted with the acids. Transfer the cotton to a vessel containing water, stir it well with a glass rod, decant the liquid, pour more water upon the mass, agitate again, and repeat the affusion, agitation, and decantation until the washings cease to give a precipitate with barium chloride, indicating that the acids have been completely removed. Drain the product on filtering-paper, and dry it in a water-bath.

Tests.—Readily soluble in a mixture of ether and rectified spirit ; should leave no residue when exploded by heat.

Preparations.—Collodium ; Collodium Flexile.

QUERCUS CORTEX

OAK BARK

The dried bark of the small branches and young stems of *Quercus pedunculata*. Collected in spring from trees growing in Britain.

Natural Order.—*Cupuliferæ*.

Composition.—Its medicinal activity depends upon its containing from fifteen to twenty per cent. of tannic acid with a small quantity of gallic acid.

Characters.—Covered with a greenish shining epidermis ; cinnamon-coloured on the surface ; fibrous, brittle, and strongly astringent.

Actions and Uses.—*Internally* :—Astringent and tonic. Employed to arrest diarrhœa, dysentery, and similar mucous discharges. *Externally* :—Astringent, styptic, and excitant. Occasionally used for stimulating unhealthy wounds and arresting gangrene ; also in prolapsus recti and prolapsus uteri.

Doses.—HORSE, 2 to 6 drachms.—CATTLE, 2 to 6 drachms.—SHEEP, $\frac{1}{2}$ to 2 drachms.—PIG, $\frac{1}{2}$ to 2 drachms. DOG, 10 to 30 grains.

Modes of Application.—*Internally* :—The powdered bark made into a bolus, or mixed with gruel, but preferably in the form of Decoctum Quercûs. *Externally* :—The powder, or Decoctum Quercûs.

Preparation.—Decoctum Quercûs.

QUINIÆ SULPHAS

SULPHATE OF QUININE

Synonyms.—Quinia Sulphate ; Sulphate of Quinine ; Disulphate of Quinine.

Composition.— $(C_{20}H_{24}N_2O_2)H_2SO_4.7Aq.$

Mode of Preparation.—Take of

Yellow Cinchona Bark, in	}	1 pound.
coarse powder		
Hydrochloric Acid		3 fluid ounces.
Distilled Water		a sufficiency.
Solution of Soda		4 pints.
Diluted Sulphuric Acid . .		a sufficiency.

Dilute the hydrochloric acid with ten pints of the water. Place the cinchona bark in a porcelain basin, and add to it as much of the diluted hydrochloric acid as will render it thoroughly moist. After maceration, with occasional stirring, for twenty-four hours, place the bark in a displacement apparatus, and percolate with the diluted hydrochloric acid until the solution which drops through is nearly destitute of bitter taste. Into this liquid pour the solution of soda, agitate well, let the precipitate (impure quinine) completely subside, decant the supernatant fluid, collect the precipi-

tate on a filter, and wash it with cold distilled water until the washings cease to have colour. Transfer the precipitate to a porcelain dish containing a pint of distilled water, and, applying to this the heat of a water-bath, gradually add diluted sulphuric acid until very nearly the whole of the precipitate has been dissolved and a neutral liquid has been obtained.

Filter the solution while hot through paper, wash the filter with boiling distilled water, concentrate the filtrate till a film forms upon its surface, and set it aside to crystallise.

Dry the crystals (quinine sulphate) on filtering-paper without the application of heat.

By digestion with diluted hydrochloric acid the bark is exhausted of quinine, cinchonine, and other alkaloids which are precipitated on the addition of soda to the solution. The precipitated alkaloids, on being dissolved in diluted sulphuric acid, are converted into sulphates, and their aqueous solution, when evaporated till a film forms upon its surface and allowed to cool, deposits the less soluble quinine sulphate, while the more soluble cinchonine sulphate remains in the mother-liquor.

Characters and Tests.—Filiform, silky, snowy-white¹ crystals, of a pure intensely bitter taste, sparingly soluble in water, yet imparting to it a peculiar bluish tint. The solution gives with barium chloride a white precipitate (barium sulphate), insoluble in nitric acid, showing that the salt is a sulphate, and when treated first with solution of chlorine and afterwards with ammonia, it assumes a splendid emerald-green colour, highly characteristic of quinine.

Should dissolve in pure sulphuric acid with a feeble

¹ Herring's 'Hospital Sulphate of Quinine' is less costly than this preparation, owing to its not having been deprived of colour, and may be substituted for it in veterinary medicine.

yellowish tint, and undergo no further change of colour when gently warmed. Ten grains, with ten minims of diluted sulphuric acid and half a fluid ounce of water, form a perfect solution, from which ammonia throws down a white precipitate (quinine). This should redissolve on agitating the whole of it with half a fluid ounce of ether, without leaving any crystalline matter (cinchonine) floating on the lower of the two strata into which the agitated fluid separates on rest. Twenty-five grains of the salt should lose 3·6 grains of water by drying at 212° F.

Actions and Uses.—Tonic and antiperiodic in intermittent and other fevers; in loss of appetite, and weak digestion depending on debility; in acute rheumatism, chorea in dogs, and constitutional or deep-seated ophthalmia in horses; during convalescence from acute complaints; also during the exhaustion which follows influenza and other diseases.

Doses.—HORSE, 20 to 40 grains.—CATTLE, 20 to 60 grains.—SHEEP, 5 to 10 grains.—PIG, 5 to 10 grains.—DOG, 1 to 5 grains.

Modes of Application.—In the form of bolus, or mixed with gruel.

Incompatibles.—Alkalies and their carbonates; infusions of vegetable astringents precipitate quinia tannate especially in the presence of sulphuric acid.

RESINA

RESIN

Synonym.—Yellow Resin.

The residue obtained by distilling the turpentine from various species of *Pinus*, and withdrawing the heat before the whole of the volatile oil is expelled.

Composition.—A mixture of abietic (sylvic) acid, which is crystallisable, and pinic acid, which is amorphous.

Characters.—Translucent, yellowish, brittle, pulverisable ; fracture shining ; odour and taste faintly terebinthinate. When heated it easily fuses, and burns with a dense yellow flame and much smoke. Soluble in rectified spirit ; insoluble in water.

Actions and Uses.—*Internally*:—Diuretic. *Externally*:—Mild excitant, astringent, and styptic. Also employed for imparting consistency and adhesiveness to plasters.

Doses.—HORSE, 4 to 6 drachms.—CATTLE, 4 to 6 drachms.—SHEEP, 1 to 2 drachms. PIG, 1 to 2 drachms.—DOG, 20 to 30 grains.

Modes of Application.—*Internally*:—Resin is principally given to the horse. It is administered in the form of bolus, preferably as *Massa Resinæ Composita*. *Externally*:—The powder sprinkled over wounds. To arrest hæmorrhage after castration a small quantity of the powder is sometimes applied to the ends of the spermatic cord and melted by a hot iron.

Preparations.—*Massa Resinæ Composita* ; *Emplastrum Picis* ; *Unguentum Resinæ*.

RHEI RADIX

RHUBARB ROOT

The dried root of *Rheum officinale* and of other undetermined species of *Rheum*. Grows in China, Chinese Tartary, and Thibet. Imported from Shanghai and Canton, and brought overland by way of Moscow.

Natural Order.—*Polygonaceæ*.

Composition.—The root contains a number of principles, including chrysophanic acid, erythroretin, phæoretin, and aporetin. At present it is not known whether the physiological effects of rhubarb are due to one particular constituent or to the combined action of several of them.

Characters. — Trapezoidal, roundish, cylindrical, or flattish pieces, frequently bored with one hole ; yellow externally, internally marbled with fine waving greyish and reddish lines ; finely gritty under the teeth ; taste bitter, faintly astringent and aromatic ; odour peculiar. Should be free from decay and not worm-eaten. Boracic acid does not turn rhubarb brown unless it is adulterated with turmeric.

Actions and Uses.—But little used in veterinary medicine. Chiefly given as a tonic to young animals with weak digestions, and to the dog as a cathartic.

Doses.—*Tonic* :—HORSE, 1 to $1\frac{1}{2}$ ounce.—CATTLE, 1 to 2 ounces.—SHEEP, 2 to 4 drachms.—PIG, 1 to $1\frac{1}{2}$ drachm. DOG, 10 to 30 grains. *Cathartic* :—DOG, 2 to 3 drachms.

Mode of Application.—The powdered root made into a bolus or mixed with gruel.

Preparations.—Pulvis Rhei Compositus ; Tinctura Rhei.

ROSÆ GALLICÆ PETALA

RED ROSE PETALS

The fresh and dried unexpanded petals of *Rosa gallica*. From plants cultivated in Britain.

Natural Order.—*Rosaceæ*.

Characters.—Colour fine purplish red, retained after drying ; taste bitterish, feebly acid, and astringent ; odour

roseate (owing to presence of volatile oil), developed by drying.

Use.—In preparing Confectio Rosæ Gallicæ.

SABINÆ CACUMINA

SAVIN TOPS

The fresh and dried tops of *Juniperus Sabina*. Collected in spring from plants cultivated in Britain.

Natural Order.—*Coniferæ*.

Composition.—Medicinal properties are due to a volatile oil having the formula $C_{10}H_{16}$.

Characters.—Twigs densely covered with minute imbricated appressed leaves in four rows ; odour strong, peculiar, and unpleasant ; taste acrid, bitter, resinous, and disagreeable.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, anthelmintic, but is very seldom employed for this purpose, on account of its violent and uncertain irritant qualities. *Externally* :—Stimulant to ill-conditioned wounds and indolent sores ; also used for maintaining the action of, and keeping up the discharge from, blisters and setons.

Mode of Application.—*Externally* :—Infusum Sabinæ, and, preferably, Unguentum Sabinæ.

Preparations.—Infusum Sabinæ ; Unguentum Sabinæ.

SACCHARUM PURIFICATUM

REFINED SUGAR

Pure cane sugar prepared from the juice of the stem of *Saccharum officinarum*, cultivated in the West Indies and other tropical countries.

Natural Order.—*Graminaceæ*.

Composition.— $C_{12}H_{22}O_{11}$.

Characters.—Compact crystalline conical loaves, known in commerce as lump sugar.

Preparations.—Confectio Rosæ Gallicæ ; Ferri Carbonas Saccharata.

SANTONINUM

SANTONIN

A crystalline neutral principle obtained from the *Artemisia maritima*, which is not a seed, but the unexpanded flower-heads of a species of *Artemisia* imported from Russia. It is the only so-called worm-seed which yields santonin, a crystalline neutral principle, in quantity worth extracting.

Natural Order.—*Compositæ*.

Composition.— $C_{15}H_{18}O_3$.

Characters and Tests.—In colourless, flat, rhombic prisms, feebly bitter, sparingly soluble in water. Solubility in rectified spirit 1 in 50. Not dissolved by diluted mineral acids. Entirely destructible by a red heat with free access of air.

Actions and Uses.—*Internally* :—Anthelmintic, useful both for tape- and thread-worms.

Doses.—HORSE, 20 to 40 grains.—DOG, 2 to 10 grains.

SAPO DURUS


HARD SOAP

Composition.—Chiefly sodium oleate,¹ $NaC_{18}H_{33}O_2$, with a large percentage of water.

Mode of Preparation.—Olive oil¹ is boiled with caustic

¹ Although the 'British Pharmacopœia' directs hard soap to be made from olive oil, the best 'curd soap' of commerce, which is a

soda (sodium hydrate) ; glycerin passes into solution, and sodium oleate (hard soap) floats on the surface of the liquid. After remaining quiescent for a few hours the soap is skimmed off, transferred to 'frames,' in which it solidifies on cooling, and finally cut by wires into slabs or bars.



Characters.—Greyish white, dry, horny, and pulverisable when kept in dry warm air ; easily moulded when heated. Soluble in rectified spirit, producing a solution which, when evaporated on paper, does not leave an oily stain. Incinerated, it yields an ash which does not deliquesce.

Actions and Uses.—*Internally*:—Cathartic and diuretic, but chiefly employed as an antacid, as an antidote to poisoning by the mineral acids, and in the preparation of clysters. *Externally* :—Stimulant to sprains and contusions ; also applied to burns and scalds to diminish the irritation and pain by excluding air. Likewise employed as an excipient in making up boluses, liniments, and ointments.

Doses.—HORSE, 1 to 2 ounces.—CATTLE, 1 to 2 ounces. SHEEP, 2 to 6 drachms.—PIG, 2 to 6 drachms.—DOG, 20 to 60 grains.

Modes of Application.—In the form of bolus, or beaten up with water.

Incompatibles.—Acids, and most metallic salts.


mixture of sodium stearate, palmitate, and oleate, and which is prepared from various solid and liquid fats, may be substituted for it in practice.

SAPO MOLLIS

SOFT SOAP

Composition.—Principally potassium oleate,¹ $\text{KC}_{18}\text{H}_{33}\text{O}_2$, with glycerin, free potassium hydrate, and a large percentage of water.

Mode of Preparation.—By boiling together olive oil¹ and caustic potash (potassium hydrate) until the soap produced assumes the appearance of a transparent jelly, and until a drop of it, when allowed to fall on a glass plate, remains clear, and does not exhibit a fatty border on cooling.



Use.—In the preparation of Linimentum Saponis Compositum.

Preparation.—Linimentum Saponis Compositum.

SCILLA

SQUILL

The sliced and dried bulb of *Urginea Scilla*.

Natural Order.—*Liliaceæ*.

Composition.—It contains scillipierin, scillitoxin and scillin. The active principle scillitoxin is soluble in water, acetic acid, and alcohols.

¹ Olive oil is directed by the 'British Pharmacopœia' to be employed in the preparation of soft soap; but the commercial article, which is the material used by the veterinary surgeon, is made from tallow, also from whale, linseed, and other oils.

Actions and Uses.—Principally used in catarrhal and bronchial affections.

Doses of the Syrup.—HORSES, $1\frac{1}{2}$ ounce.—DOGS, 10 to 60 minims.

Preparation.—Syrupus Scillæ.

SEVUM PRÆPARATUM

PREPARED SUET

The internal fat of the abdomen of the sheep, *Ovis Aries*, purified by melting and straining.

Composition.—A mixture of stearin, palmitin, and olein.

Characters.—White, smooth, almost scentless ; fusible at 103° F.

Uses.—Sometimes employed as a substitute for, or in conjunction with, lard, in the preparation of ointments.

SINAPIS


MUSTARD

The seeds of *Sinapis nigra* and *Sinapis alba* ; also the seeds of both reduced to powder and mixed.¹

Natural Order.—*Cruciferae*.

Composition.—The vesicating power of mustard, and the pungent odour emitted by it when moistened, depend upon a volatile oil (allyl sulphocyanide, C_3H_5CNS). It does not exist ready formed in the seed, but is developed by the mutual action of two of its proximate principles, viz. myrosin, a kind of ferment, and potassium myronate, in the presence of hot, but not boiling water.¹

¹ Boiling water coagulates myrosin, and destroys its power as a ferment.



Actions and Uses.—Vesicant and counterirritant, in the form of poultice.

Preparations.—Cataplasma Sinapis ; Cataplasma Sinapis Ammoniatum.

SODA CAUSTICA

CAUSTIC SODA

Synonyms.—Sodium Hydrate ; Hydrate of Sodium ; Hydrate of Soda.

Composition.— NaOH .

Mode of Preparation.—Take of

Solution of Soda 2 pints.

Boil down the solution of soda rapidly in a silver or clean iron vessel until there remains a fluid of oily consistency, a drop of which, when removed on a warmed glass rod, solidifies on cooling. Pour the fluid on a clean silver or iron plate, or into moulds, and as soon as it has solidified break it into pieces, and preserve it in stoppered green glass bottles.

Characters and Tests.—Hard and greyish white, very alkaline and corrosive. Does not deliquesce on exposure to the atmosphere. It imparts a yellow colour to flame indicating the presence of sodium, and its solution in water, acidulated by nitric acid, should give only scanty white precipitates with argentum nitrate and barium chloride, showing the presence of traces only of chlorides and sulphates respectively.

Actions and Uses.—*Externally*:—As a caustic. It is equal in activity to Potassa Caustica, but preferable to this agent, as it does not deliquesce and spread.


SODÆ ACETAS

ACETATE OF SODA

Synonym.—Sodium Acetate.

Composition.— $\text{NaC}_2\text{H}_3\text{O}_2 \cdot 3\text{Aq.}$

Mode of Preparation.—By dissolving sodium carbonate in acetic acid and evaporating the solution, so that the sodium acetate produced may crystallise out. The crystals are drained, and subsequently dried by exposure to air on a porous brick.



The salt, in assuming the crystalline state, acquires three molecules of water.

Characters and Tests.—Transparent colourless crystals, soluble in water, forming a solution neutral to test-paper. The salt, or its solution in water, warmed with dilute sulphuric acid, emits the odour of vinegar. The aqueous solution, when dilute, is not precipitated by barium chloride or silver nitrate, showing its freedom from sulphates and chlorides.

Use.—In the preparation of acetic acid.


SODÆ BICARBONAS

BICARBONATE OF SODA

Synonym.—Sodium-Hydrogen Carbonate.

Composition.— NaHCO_3 .

Mode of Preparation.—Carbonic anhydride¹ is passed into a vessel containing a mixture of two parts of crystallised and three parts of dried sodium carbonate until the gas is no longer absorbed. The damp salt which is formed is shaken occasionally during half an hour with half its weight of cold distilled water, for the purpose of dissolving an unaltered carbonate, and the remaining insoluble bicarbonate is dried by exposure to air on filtering-paper placed on porous bricks. The object of using the crystallised carbonate is that it may conveniently furnish the necessary amount of water for the success of the process.



Characters and Tests.—In powder, or small opaque irregular scales, white ; saline unpleasant taste. Imparts a yellow colour to flame, showing the presence of sodium. Soluble in water 1 part in 10. Dissolves with effervescence in diluted hydrochloric acid, showing that it is a carbonate ; and the hydrochloric acid solution gives no precipitate with platinic chloride, indicating that it is neither a potassium nor an ammonium salt. Its aqueous solution acidified with nitric acid should give but faint indications of sulphates and chlorides on the addition of barium chloride and silver nitrate. A solution of the salt in cold water gives a white and not a coloured precipitate with mercuric chloride, unless

¹ For mode of generating this gas, see Potassæ Bicarbonas.

sodium carbonate be present. Distinguished from sodium carbonate by its aqueous solution not giving a precipitate with magnesium sulphate.

Actions and Uses.—Similar to those of Potassæ Bicarbonas. Less alkaline and milder than Sodæ Carbonas.

Doses.—HORSE, 4 to 6 drachms.—CATTLE, 4 to 8 drachms.—SHEEP, 1 to 2 drachms.—PIG, 1 to 2 drachms. Dog, 5 to 40 grains.

Modes of Application. }
Incompatibles. } Same as Potassæ Bicarbonas.


SODÆ CARBONAS

CARBONATE OF SODA


Synonyms.—Sodium Carbonate ; Washing-Soda.

Composition.— $\text{Na}_2\text{CO}_3 \cdot 10\text{Aq.}$

Mode of Preparation.—Sodium chloride (common salt) is decomposed by sulphuric acid, whereby sodium sulphate ('salt cake') and hydrochloric acid are produced ; the former remains on the hearth, while the latter escapes by the chimney, of the furnace in which the operation is conducted.



The sodium sulphate is next heated with a mixture of carbon (coal-dust) and calcium carbonate (chalk or limestone), when the following changes take place :



The resulting mixture of sodium carbonate and calcium sulphide ('black ash' or 'ball soda'), when cold, is broken up into little pieces and lixiviated with water; the sodium carbonate passes into solution, and the calcium sulphide is undissolved. By evaporating to dryness the solution thus formed, 'soda ash' is procured, and by dissolving 'soda ash' in water, filtering, and evaporating the solution, and then allowing it to cool, sodium carbonate is obtained in the crystallised state.

Characters and Tests.—Transparent, colourless, laminar, rhombic crystals, efflorescent; harsh alkaline taste, and strong alkaline reaction. It imparts a yellow colour to flame, showing the presence of sodium, and dissolves with effervescence in diluted hydrochloric acid, indicating that it is a carbonate. Soluble in cold water 1 in 2; insoluble in rectified spirit. The hydrochloric acid solution gives no precipitate with platinic chloride, showing that it is neither a potassium nor an ammonium salt. Its aqueous solution gives a white precipitate (magnesium carbonate) with magnesium sulphate, a reaction by which it is distinguished from Sodæ Bicarbonas. Strongly heated, it undergoes the aqueous fusion, and then dries up, losing water to the extent of sixty-three per cent. of its weight. When its aqueous solution is treated with excess of nitric acid, it should yield but slight precipitates with barium chloride and silver nitrate, indicating the almost total absence of sulphates and chlorides.

Actions and Uses.—Very similar to those of Potassæ Bicarbonas, Potassæ Carbonas, and Sodæ Bicarbonas.

Doses.—Half those of Sodæ Bicarbonas.

Mode of Application.

Incompatibles.

} Same as Potassæ Bicarbonas.

Preparations.—Liquor Sodæ; Liquor Sodæ Chloratæ
Sodæ Bicarbonas; Sodæ Carbonas Exsiccata.

SODÆ CARBONAS EXSICCATA**DRIED CARBONATE OF SODA**

Composition.— Na_2CO_3 .

Mode of Preparation.—Crystallised sodium carbonate is heated in a porcelain capsule until the liquid which first forms is converted into a dry cake. This cake, when pulverised, is to be preserved in a stoppered bottle.

Actions and Uses.—Similar to those of the potassium and sodium carbonates and bicarbonates.

Doses.—About one fourth of those of Sodæ Bicarbonas.

Modes of Application. } Same as Potassæ Bicarbonas.
Incompatibles. }


SODÆ HYPOSULPHIS**HYPOSULPHITE OF SODA**

Synonyms.—Sodium Thiosulphate ; Antichlor.

Composition.— $\text{Na}_2\text{S}_2\text{O}_3 \cdot 10\text{Aq}$.

Mode of Preparation.—An aqueous solution of sodium sulphite is boiled with flowers of sulphur and filtered. On evaporating the filtrate the salt crystallises out.

Assuming the first of the above formulæ to represent the composition of the salt, the following equation explains its formation.



During crystallisation the salt acquires five molecules of water.

Characters and Tests.—Rhombic prisms, soluble in water. Imparts a yellow colour to flame, indicating the presence of sodium. Diluted hydrochloric acid added to its aqueous solution causes effervescence (due to the expulsion of sulphurous anhydride, recognisable by its odour), and the solution is, at the same time, rendered turbid by the precipitation of sulphur. The latter reaction distinguishes this salt from sodium sulphite.

Actions and Uses.

Doses.

Incompatibles.

} Same as Sodæ Sulphis.


SODÆ SULPHAS

SULPHATE OF SODA

Synonyms.—Sodium Sulphate ; Glauber's Salt.

Composition.— $\text{Na}_2\text{SO}_4 \cdot 10\text{Aq.}$

Mode of Preparation.—By neutralising the residue (sodium-hydrogen sulphate) left in the manufacture of hydrochloric acid with sodium carbonate, and crystallising the product from solution in water.



Characters and Tests.—Transparent oblique prisms ; has a saline and bitter taste ; effloresces on exposure to the air ; soluble in water, 1 in 3. Heated in a porcelain crucible, it loses 55·9 per cent of water. It imparts a yellow colour to flame, indicating the presence of sodium. With barium chloride its aqueous solution furnishes a white precipitate (barium sulphate) insoluble in nitric acid, showing

that the salt is a sulphate. Its aqueous solution mixed with diluted hydrochloric acid gives no precipitate with platinic chloride, proving that it is neither a potassium nor an ammonium salt.

Actions and Uses.—Cathartic, diuretic, and resolvent. *Magnesia Sulphas* being more certain in its action than *Sodæ Sulphas*, the latter is not much employed by the veterinarian, except occasionally as a purgative for cattle.

Dose.—Purgative for cattle, from 1 to 1½ pound.

Mode of Application.—Dissolved in water, or mixed with ginger and treacle.


SODÆ SULPHIS

SULPHITE OF SODA


Synonym.—Sodium Sulphite.

Composition.— $\text{Na}_2\text{SO}_3 \cdot 7\text{Aq.}$

Mode of Preparation.—An aqueous solution of sodium carbonate is saturated with sulphurous anhydride; sodium-hydrogen sulphite is formed and remains in solution, while carbonic anhydride escapes in the gaseous state.



The solution of sodium-hydrogen sulphite is mixed while warm with as much sodium carbonate as was used in the first operation; on cooling, sodium sulphite crystallises out. The crystals, after being drained, are dried on filtering paper.



In the act of crystallising the salt takes up seven molecules of water.

Characters and Tests.—Sometimes in transparent prismatic crystals, but usually in white opaque masses. Soluble in water. It imparts a yellow colour to flame, showing the presence of sodium. On adding diluted hydrochloric acid to its aqueous solution, sulphurous anhydride (recognisable by its odour) is evolved with effervescence, and the solution remains bright and transparent.

Actions and Uses.—*Internally* :—Antiseptic, disinfectant, and alterative. Given in indigestion, tympanites, and in so-called blood diseases. *Externally* :—Antiseptic, deodoriser, and disinfectant. Employed in the treatment of ulcerated sore throat and phagedænic wounds and ulcers.

Doses.—HORSE, 1 to $1\frac{1}{2}$ ounce.—CATTLE, 2 to 4 ounces.—SHEEP, 2 to 6 drachms.—PIG, 2 to 6 drachms.—DOG, 20 to 60 grains. Frequently repeated.

Modes of Application.—*Internally* :—Dissolved in water. *Externally* :—Three to four drachms, dissolved in every ounce of water employed.

Incompatibles.—Acids, and most metallic salts.

SODII CHLORIDUM

CHLORIDE OF SODIUM

Synonym.—Sodium Chloride ; Common Salt.

Composition.—NaCl.

Mode of Preparation.—Crude 'rock salt,' or a similar product obtained by evaporating sea-water, or the water of brine springs, is purified by solution and crystallisation.

Characters and Tests.—Small white crystalline grains, or transparent cubic crystals, free from moisture ; purely saline taste ; soluble in water. Its aqueous solution mixed with hydrochloric acid and platinic chloride yields no precipitate, showing that it is neither an ammonium nor a potassium salt. It imparts a yellow colour to flame, indicating that sodium is present ; and its aqueous solution gives with silver nitrate a white precipitate (silver chloride) soluble in ammonia, but insoluble in boiling nitric acid, proving the salt to be a chloride.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, cathartic, anthelmintic, emetic, diuretic, tonic, alterative, and resolvent. To cattle and sheep it is given as an anthelmintic ; also as a purgative in diarrhœa caused by the presence of an irritant or by over-feeding, in fardel-bound, and in distention of the stomach by food. To sheep it is likewise given as a preventive of rot, for the destruction of intestinal worms and as a corrective of the bad effects of damp and badly kept fodder. It is employed as an emetic for the dog, and is given to all animals, in frequently repeated doses, as a tonic and stimulant in impaired digestion. Rock salt should always be kept as a provocative of the appetite in the horse's manger, the ox's crib, and the sheep's trough. *Externally* :—As a stimulant in chronic sprains, many affections of the joints, and in some diseases of the feet, particularly amongst cattle and sheep. Common salt is frequently added to laxative enemata.

Doses.—*Anthelmintic and Cathartic* :—CATTLE, 12 to 24 ounces.—SHEEP, 1 to 3 ounces.

Emetic :—DOG, 1 to 2 drachms.

Tonic :—HORSE, 1 to 2 ounces.—CATTLE, 2 to 4 ounces. SHEEP, 2 to 4 drachms.—DOG, 10 to 30 grains.

Modes of Application.—*Internally* :—As a cathartic and emetic dissolved in tepid water. Sometimes half doses of chloride of sodium and sulphate of magnesia are mixed with ginger, pimento, or some other aromatic and treacle ; such a mixture will often be voluntarily drunk by cattle and sheep. Croton oil, oil of turpentine, and calomel may be added to common salt, to increase its cathartic and anthelmintic action ; and mustard, sulphate of zinc, sulphate of copper, or tartarated antimony may be combined with it to ensure its prompt action as an emetic. As a tonic and stomachic, it should be sprinkled over the animal's food, either in the state of powder or in solution. *Externally* :—A lotion consisting of one pound of common salt dissolved in a gallon of cold water. This lotion should be applied during, or immediately after, the solution of the salt.

SPIRITUS ÆTHERIS

SPIRIT OF ETHER

Synonym.—Spiritus Ætheris Sulphurici.

Composition.—A mixture of sulphuric ether with twice its bulk of rectified spirit.

Mode of Preparation.—Take of

Ether	10 fluid ounces.
Rectified Spirit	1 pint.

Mix.

Test.—Sp. gr. 0·809.

Actions and Uses.—Diffusible stimulant ; antispasmodic.

Doses.—HORSE, 3 to 6 fluid ounces.—CATTLE, 5 to 10 fluid ounces.—SHEEP, 1 to 2 fluid ounces.—PIG, 1 to 2 fluid ounces.—DOG, 1 to 5 drachms.

SPIRITUS ÆTHERIS NITROSI**SPIRIT OF NITROUS ETHER**

Synonyms.—Spiritus Ætheris Nitrici ; Sweet Spirits of Nitre.


Composition.—An alcoholic solution of nitrous ether (ethyl nitrite), $C_2H_5NO_2$.

Mode of Preparation.—Take of

Nitric Acid	3 fluid ounces.
Sulphuric Acid	2 fluid ounces.
Copper in fine wire (about No. 25)	} 2 ounces.
Rectified Spirit	
	a sufficiency.

To one pint of the spirit add gradually the sulphuric acid, and stir them together ; then add, in the same way, two and a half fluid ounces of the nitric acid. Put the mixture into a retort or other suitable apparatus, into which the copper has been introduced, and to which a thermometer has been fitted. Attach an efficient condenser, and, applying a gentle heat let the spirit distil at a temperature, commencing at $170^{\circ}F.$, and rising to 175° , but not exceeding 180° , until twelve fluid ounces have passed over and been collected in a bottle kept cool, if necessary, with ice-cold water ; then withdraw the heat and, having allowed the contents of the retort to cool, introduce the remaining half-ounce of nitric acid, and resume the distillation as before, until the distilled product (spirit of nitrous ether) has been increased to fifteen fluid ounces. Mix this with two pints of the rectified spirit, or as much as will make it correspond to the tests of specific gravity and percentage of ether separated by calcium chloride. Preserve it in well-stoppered bottles.

The following equation explains the principal changes which probably take place in the foregoing process :



The ethyl nitrite distils over, accompanied by alcohol and other bodies of minor importance.

Characters and Tests.—Transparent and nearly colourless, but with a very slight tinge of yellow ; mobile, inflammable ; peculiar penetrating apple-like odour, and sweetish cooling, sharp taste. Sp. gr. 0·845. It effervesces feebly or not at all when shaken with a little sodium-hydrogen carbonate, indicating the presence of a trace or the absence of free acid. When agitated with solution of ferrous sulphate and a few drops of sulphuric acid, it becomes deep olive-brown or black. Agitated with twice its volume of a saturated solution of calcium chloride (to absorb water and alcohol) in a closed graduated tube, two per cent. of its original volume should separate in the form of ethyl nitrite and rise to the surface of the mixture.

Actions and Uses.—In excessive doses narcotic and sedative ; in medicinal doses, stimulant, antispasmodic, diuretic, and diaphoretic. Given in colic, indigestion, tympanites, and local congestions ; during convalescence from debilitating diseases ; also in combination with tincture of opium for diminishing the pains and spasms which sometimes follow parturition in cows.

Doses.—HORSE, 1 to 2 fluid ounces.—CATTLE, 3 to 4 fluid ounces.—SHEEP, 3 to 6 fluid drachms.—PIG, 3 to 6

fluid drachms.—Dog, $\frac{1}{2}$ to 2 fluid drachms. Frequently repeated.

Mode of Application.—Mixed with sufficient cold water to make a draught.

SPIRITUS AMMONIÆ AROMATICUS¹

AROMATIC SPIRIT OF AMMONIA

Take of

Carbonate of Ammonia	. . .	8 ounces.
Strong Solution of Ammonia	. . .	4 fluid ounces.
Volatile Oil of Nutmeg	. . .	4 fluid drachms.
Oil of Lemon	. . .	6 fluid drachms.
Rectified Spirit	. . .	6 pints.
Water	. . .	3 pints.

Mix and distil seven pints.

Test.—Sp. gr. 0·870.

Actions and Uses.—Diffusible stimulant, antacid, and antispasmodic.

Doses.—HORSE, $\frac{1}{2}$ to 1 fluid ounce.—CATTLE, $\frac{1}{2}$ to 2 fluid ounces.—SHEEP, 2 to 4 fluid drachms.—PIG, 2 to 4 fluid drachms.—Dog, 10 to 20 minims. Mixed with water.

SPIRITUS AMMONIÆ FŒTIDUS

FETID SPIRIT OF AMMONIA

Take of

Assafœtida	. . .	12 ounces.
Strong Solution of Ammonia	. . .	2 fluid ounces.
Rectified Spirit	. . .	a sufficiency.

¹ This preparation is stronger in spirit, and about one-half stronger in ammonia, than the Spiritus Ammoniæ Aromaticus of the 'British Pharmacopœia.'

Break the assafœtida into small pieces, and macerate it in a closed vessel, in fifteen fluid ounces of the spirit, for twenty four hours ; then distil off the spirit, mix the product with the solution of ammonia, and add sufficient rectified spirit to make one pint.

Actions and Uses.—Stimulant and antispasmodic.

Doses.—HORSE, $\frac{1}{2}$ to 1 fluid ounce.—CATTLE, $\frac{1}{2}$ to 2 fluid ounces.—SHEEP, 2 to 4 fluid drachms.—PIG, 2 to 4 fluid drachms.—DOG, 10 to 20 minims. Mixed with water.

SPIRITUS CAMPHORÆ

SPIRIT OF CAMPHOR

Synonym.—Tinctura Camphoræ.

Take of

Camphor	1 ounce.
Rectified Spirit	4 fluid ounces.

Dissolve.

Actions and Uses.—*Externally* :—Stimulant, anodyne and discutient.

SPIRITUS CHLOROFORMI

SPIRIT OF CHLOROFORM

Synonym.—Chloric Ether.

Take of

Chloroform	1 fluid ounce.
Rectified Spirit	19 fluid ounces.

Dissolve.

Test.—Sp. gr. 0·871.

Actions and Uses.—Stimulant, antispasmodic, and anodyne.

Doses.—HORSE, 2 to 4 fluid ounces.—CATTLE, 2 to 4 fluid ounces.—SHEEP, $\frac{1}{2}$ to 1 fluid ounce.—PIG, $\frac{1}{2}$ to 1 fluid ounce.—DOG, 1 to 3 fluid drachms. Mixed with water.

SPIRITUS HYDRARGYRI PERCHLORIDI COMPOSITUS

COMPOUND SPIRIT OF PERCHLORIDE OF MERCURY

Synonym.—Liquor Hydrargyri Bichloridi.

Take of

Perchloride of Mercury, in	}	1 ounce.
powder		
Hydrochloric Acid		1 fluid ounce.
Rectified Spirit		7 fluid ounces.

Add the acid to the spirit, and dissolve the perchloride mercury in the mixture.

Uses.—*Externally* :—Occasionally to farcy ulcers, and for arresting synovial discharges.

SPIRITUS METHYLATUS

METHYLATED SPIRIT

Composition.—A mixture consisting of ninety per cent. of spirit of wine and ten per cent. of methylic alcohol, CH_3HO .

Uses.—As a substitute for the more expensive spirit of wine in the preparation of tinctures, &c., for *external use only*. The law forbids methylated spirit, or any preparation containing it, to be used internally.


SPIRITUS RECTIFICATUS

RECTIFIED SPIRIT

Synonym.—Spirit of Wine.

Composition.—Alcohol (absolute alcohol, C_2H_5HO), with sixteen per cent. of water.

Mode of Preparation.—Saccharine fluids are permitted to ferment, whereby the sugar which they contain is transformed into alcohol and carbonic anhydride. The former is separated from the impurities with which it is mixed by distillation, and the latter escapes into the air in the state of gas.



Characters and Tests.—Colourless, transparent, very mobile and inflammable ; peculiar pleasant odour, and a strong spirituous burning taste. Burns with a blue flame, without smoke. Sp. gr. 0.358. Should have a purely alcoholic odour and taste, and should remain clear when diluted with distilled water, indicating its freedom from fusel oil (amylic alcohol, $C_5H_{11}.OH$).

Actions and Uses.—*Internally* :—Stimulant, diuretic, and diaphoretic. Alcohol, in the form of ale, brandy, whisky, or gin, for the larger animals, and in the form of wine for the smaller ones, is employed as a stimulant in colic and tympanites ; in the latter stages of influenza, bronchitis, and other inflammatory diseases ; also in, and during convalescence from, debilitating complaints. *Externally* :—Rectified spirit is added to cooling lotions to increase their refrigerating power. Rectified spirit is likewise employed in the preparation of medicinal spirits, tinctures, and extracts.

Doses.¹—HORSE, 1 to 3 ounces.—CATTLE, 3 to 6 ounces.
SHEEP, $\frac{1}{2}$ to $1\frac{1}{2}$ ounce.—PIG, $\frac{1}{2}$ to $1\frac{1}{2}$ ounce.—DOG, 1 to 3
drachms. Repeated every two hours.

*Table showing the average amounts by measure of absolute
alcohol ($C_2H_5.OH$) contained in 100 volumes of the
following alcoholic beverages :*

Name of Beverage.	Absolute Alcohol in 100 volumes.
Small Beer and Table Ale	1 to 2
Porter	about 5
Ale (various kinds)	5 to 9
Sherry	15 to 17
Port	17 to 18
Brandy (common)	36
Gin	45
Brandy (Cognac)	46
Rum	48
Whisky	50

SPIRITUS TENUIOR

PROOF SPIRIT

Composition.—One hundred parts, at 60°F., consists of
forty-nine parts by weight of absolute alcohol, and fifty-
one parts by weight of water.

Take of

Rectified Spirit	5 pints.
Distilled Water	3 pints.

Mix.

Test.—Sp. gr. 0.92.

Use.—In the preparation of tinctures.

¹ The doses here enumerated refer to rectified spirit; the deter-
mination of those of the ordinary alcoholic beverages must be left
to the judgment of the prescriber.

STAPHISAGRIÆ SEMINA

STAVESACRE SEEDS

The seeds of *Delphinium Staphisagria*.

Natural Order.—*Ranunculaceæ*.

Composition.—The seeds contain several alkaloids, the most important being delphinine, delphisine, and delphinidine. They also contain a fixed oil.

Actions and Uses.—A solution made by boiling the seeds in water, 1 part of seeds to 20 or 30 of water, is used as a parasiticide lotion.

STRYCHNINA

STRYCHNINE

Composition.— $C_{21}H_{22}NO_2$.

Mode of Preparation.—Take of

Nux Vomica	1 pound.
Acetate of Lead	180 grains.
Solution of Ammonia	a sufficiency.
Rectified Spirit	a sufficiency.
Distilled Water	a sufficiency

Soften the nux vomica by subjecting it for two hours to steam in any convenient vessel ; chop or slice it ; dry it in a water-bath or hot-air chamber, and immediately grind it in a coffee-mill. Digest the powder at a gentle heat for twelve hours with two pints of the spirit and one of the water, to dissolve out the igasurates of strychnine and brucine ; strain through linen, express strongly, and repeat the process twice. Distil off the spirit from the mixed fluid, evaporate the watery residue to about sixteen ounces,

and filter when cold. Add now the acetate of lead previously dissolved in distilled water, so long as it occasions any precipitate (lead igasurate) ; filter ; wash the precipitate with ten ounces of cold water, add the washings to the filtrate (containing the acetates of strychnia and brucia) ; evaporate the clear fluid to eight ounces, and when it has cooled add the ammonia in slight excess, stirring thoroughly. Let the mixture stand, at the ordinary temperature, for twelve hours ; collect the precipitate (impure strychnine) on a filter, wash it at once with a few ounces of cold distilled water, dry it in a water-bath or hot-air chamber, and boil it with successive portions of rectified spirit till the fluid scarcely tastes bitter. Distil off most of the spirit, evaporate the residue to the bulk of about half an ounce, and set it aside to cool. Cautiously pour off the yellowish mother-liquor (which contains the brucine of the seeds) from the white crust of strychnine which adheres to the vessel. Throw the crust on a paper filter, wash it with a mixture of two parts of rectified spirit and one of water till the washing ceases to become red on the addition of nitric acid, showing that the brucine has been removed from the strychnia ; finally, dissolve it (the strychnia) by boiling it with an ounce of rectified spirit, and set it aside to crystallise. More crystals of strychnine may be obtained by evaporating the mother-liquor.

Characters and Tests.—Right square octahedrons or prisms, colourless and inodorous ; sparingly soluble in water, but communicating to it an intensely bitter taste ; soluble in boiling rectified spirit, chloroform, and in commercial ether, but not in absolute alcohol or ether. Pure sulphuric acid forms with it a colourless solution, which on the addition of solid potassium bichromate acquires a purple hue, changes to yellowish-red, and ultimately leaves a green residue. Should not be coloured by nitric

acid ; if it turns yellow, it shows that brucine is present. Should leave no ash when burned with free excess of air.

Actions and Uses.—In excessive doses, highly poisonous ; in medicinal doses, stimulant and tonic in paralysis, and in chorea in the dog.

Doses.—HORSE, 1 to 3 grains.—CATTLE, 1 to 3 grains.—SHEEP, $\frac{1}{3}$ to 1 grain.—DOG, $\frac{1}{30}$ to $\frac{1}{10}$ th grain. Once or twice a day.

Modes of Application.—In the form of bolus, pill, or dissolved in water acidulated with acetic or sulphuric acid.

Antidotes.—Same as to *Nux Vomica*.

Preparation.—Liquor Strychniæ.

SUCCUS RHAMNI

BUCKTHORN JUICE

Synonym.—Rhamni Succus.

The recently expressed juice of the ripe berry of the common buckthorn, *Rhamnus catharticus*, belonging to the natural order *Rhamnaceæ*.

Preparation.—Syrupus Rhamni.

SULPHUR SUBLIMATUM

SUBLIMED SULPHUR

Synonym.—Flowers of Sulphur.

Composition.—An elementary body, having the symbol S.

Mode of Preparation.—Crude sulphur, imported from Sicily, Naples, the Roman States, and other volcanic districts, is placed in a retort connected with a brick chamber.

On the application of heat to the retort the sulphur sublimes into the chamber, to the walls of which it attaches itself, while the earthy impurities remain behind. When a sufficient quantity of the sulphur has collected on the walls of the condensing chamber, it is scraped off for use.

Characters and Tests.—A slightly gritty crystalline powder, of a fine greenish-yellow colour, tasteless and odourless, unless heated; burns in open vessels with a blue flame and the evolution of sulphurous anhydride, recognisable by its odour. Entirely volatilised by heat. Insoluble in water, alcohol, and ether, but soluble in carbon disulphide and turpentine. Should not redden moistened litmus paper. Solution of ammonia agitated with it and filtered does not leave any residue on evaporation, showing that it is free from arsenicum trisulphide.

Actions and Uses.—*Internally* :—In excessive doses irritant poison; in medicinal doses laxative, and as an alterative in rheumatism, skin diseases, and many other affections. *Externally* :—In scabies and other skin diseases.

Doses.—*Laxative* :—HORSE, 3 to 4 ounces.—CATTLE, 5 to 6 ounces.—SHEEP, 2 ounces.—PIG, $1\frac{1}{2}$ to 2 ounces.—DOG, 2 to 8 drachms.

Alterative :—HORSE, $\frac{3}{4}$ to $1\frac{1}{2}$ ounce.—CATTLE, 1 to 2 ounces.—SHEEP, 6 drachms.—PIG, 4 to 6 drachms.—DOG, $\frac{1}{2}$ to 2 drachms.

Modes of Application.—*Internally* :—In the form of bolus or pill, mixed with gruel, or dissolved in oil. *Externally* :—Unguentum Sulphuris.

Preparations.—Unguentum Sulphuris. Unguentum Sulphuris Compositum.

SULPHURIS IODIDUM

IODIDE OF SULPHUR

Composition.—Not well established.

Mode of Preparation.—Take of

Iodine	4 ounces.
Sublimed Sulphur	1 ounce.

Rub them together in a Wedgwood mortar until they are thoroughly mixed. Put the mixture into a flask, close the orifice loosely, and apply a gentle heat, so that the colour of the mass shall become gradually darkened. When the colour has become uniformly dark throughout, increase the heat so as to produce liquefaction. Then incline the flask in different directions, in order to return into the liquid any portion of the iodine which may have condensed on the inner surface of the vessel. Lastly, withdraw the heat, and when the liquid has congealed remove the mass by breaking the flask, reduce it to pieces, and keep them in a well-stoppered bottle.

Characters and Tests.—A greyish-black solid, with a radiated crystalline appearance. Resembles iodine in odour and in the property of staining the skin. Soluble in about sixty parts of glycerin ; insoluble in water, but decomposed when boiled with it.

Actions and Uses.—*Externally* :—In scabies, urticaria, and other skin affections ; also applied to farcy ulcers and indolent sores.

Mode of Application.—Unguentum Sulphuris Iodidi.

Preparation.—Unguentum Sulphuris Iodidi.

SYRUPUS CHLORALIS HYDRATIS**SYRUP OF HYDRATE OF CHLORAL**

Preparation.—Take of

Hydrate of Chloral	. . .	80 grains.
Distilled Water	. . .	1½ drachms.

Syrup to make 1 ounce.

Mix together.

The specific gravity should be about 1.32.

Composition.—One drachm contains 10 grains of Chloral Hydrate.

Uses.—See Chloral Hydrate. Used in canine practice.

Dose.—Dog, 30 minims to 2 drachms.

SYRUPUS PAPAVERIS**SYRUP OF POPPIES**

Take of

Poppy Capsules, dried, freed from seeds, and coarsely powdered.	}	36 ounces.
Rectified Spirit		
Refined Sugar		4 pounds.
Boiling Distilled Water		a sufficiency.

Mix the poppy capsules with four pints of the water, and infuse for twenty-four hours, stirring them frequently ; then pack them in a percolator, and, adding more of the water, allow the liquor to pass slowly until about two gallons have been collected, or the poppies are exhausted. Evaporate the liquor by a water-bath until it is reduced to three pints. When quite cold, add the spirit, let the mixture stand for twelve hours, and filter. Distil off the spirit, evaporate the remaining liquor to two pints, and

then add the sugar. The product should weigh six and a half pounds, and should have the sp. gr. 1.32.

Actions and Uses.—Sedative, anodyne, and hypnotic for the dog.

Dose.—2 to 5 fluid drachms.

Preparation.—Mistura Ricini.

SYRUPUS RHAMNI

SYRUP OF BUCKTHORN

Take of

Buckthorn Juice	.	4 pints.
Ginger, sliced	}	. $\frac{3}{4}$ ounce.
Pimento, bruised		
Refined Sugar	.	5 pounds, or a sufficiency.
Rectified Spirit	.	6 fluid ounces.

Evaporate the juice to two and a half pints, add the ginger and pimento, digest at a gentle heat for four hours, and strain. When cold, add the spirit, let the mixture stand for two days, then decant off the clear liquor, and in this dissolve the sugar with a gentle heat, so as to make the sp. gr. 1.32.

Use.—Mild cathartic and laxative for the dog, and in preparing Mistura Ricini.

Dose.—Dog, $\frac{1}{2}$ to 2 ounces.—Cat, 2 to 6 drachms.

TABACI FOLIA¹

LEAF TOBACCO

The dried leaves of Virginian Tobacco, *Nicotiana Tabacum*. Cultivated in America.

Natural Order.—*Solanaceæ*.

¹ In practice the ordinary 'shag' tobacco is usually employed.

Characters.—Large mottled-brown, ovate or lanceolate acuminate leaves, bearing numerous short glandular hairs ; having a peculiar heavy odour and nauseous, bitter, acrid taste ; yielding, when distilled with potassium hydrate, an alkaline fluid, which has the peculiar odour of nicotine (the active principle of tobacco), and precipitates with platonic chloride and tincture of galls.

Preparations.—Decoctum Tabaci ; Enema Tabaci ; Infusum Tabaci.

TARAXACUM

DANDELION

The root of *Taraxacum Dens-leonis* (Natural Order *Compositæ*).

Composition.—It contains asparagin, taraxacin, and taraxacerin.

Actions and Uses.—According to Rutherford it is a feeble hepatic stimulant, it is also a laxative and diuretic and is of service in duodenal dyspepsia. The best preparation is the fresh succus, the dose of which is for the HORSE, one ounce ; DOG, a drachm and a half to two drachms.

TEREBINTHINA CANADENSIS

CANADA BALSAM

The turpentine obtained by incision from the stem of *Abies balsamea*, Balm of Gilead fir. Cultivated in Canada.

Natural Order.—*Coniferæ*.

Characters.—Pale yellow ductile oleo-resin, of the consistence of thin honey ; peculiar agreeable odour, slightly bitter, feebly acrid taste ; by exposure to air it dries very slowly into a transparent adhesive varnish ; solidifies when mixed with a sixth of its weight of magnesia.

Use.—In preparing Collodium Flexile.

THERIACA

TREACLE

Synonyms.—Sacchari Fæx ; Molasses.

Composition.—A mixture chiefly consisting of uncrystallisable sugar, produced by the action of too high a temperature on cane sugar, unaltered cane sugar, and colouring matter.

Characters.—Thick brown fermentable syrup, very sweet, not crystallising by rest or evaporation. Sp. gr. about 1·4. Nearly free from empyreumatic odour and flavour.

Actions and Uses.—Laxative. Employed for hastening the activity of cathartics, and as a vehicle for medicines having a disagreeable flavour, or which occasion nausea. Also used in the preparation of masses.

Doses.—HORSE, 1 to 1½ pound.—CATTLE, 1 to 1½ pound, SHEEP, 3 to 4 ounces.—PIG, 3 to 4 ounces.—DOG, 1 to 2 ounces. Frequently repeated until a laxative effect is produced.

THYMOL

THYMOL

A stearoptin obtained from the volatile oil of thyme, which is extracted from the *Thymus vulgaris*.

Natural Order.—*Labiatae*.

Composition.— $C_{10}H_{13}OH$.

Characters and Tests.—Thymol crystallises in hexagonal forms, nearly or quite colourless, it has an aromatic thyme-like odour, a pungent aromatic taste. It is soluble in about

1,200 parts of cold water, and 900 of boiling water, freely soluble in alcohol, ether, chloroform, and oils.

Actions and Uses.—A powerful disinfectant and antiseptic, also diaphoretic and diuretic, principally employed externally for allaying irritation and removing scales in chronic eczema and psoriasis, either in solution 1 to 2 grains in diluted spirit, or as an ointment 8 to 10 grains to an ounce of vaselin.

TINCTURA ACONITI¹

TINCTURE OF ACONITE

Take of

Aconite Root, in coarse powder	2½ ounces.
Rectified Spirit	1 pint.

Macerate the aconite root for forty-eight hours in fifteen fluid ounces of the spirit in a closed vessel, and agitate occasionally ; transfer to a percolator, and, when the fluid

¹ This is the tincture of the ‘British Pharmacopœia,’ 1867 ; it has one fourth of the strength of Tinctura Aconiti, Dublin, and one third of the strength of Tinctura Aconiti, London. Fleming’s Tincture of Aconite is nearly four times as strong as that of the ‘British Pharmacopœia.’

Squire, in his admirable ‘Companion to the British Pharmacopœia,’ gives the following directions for percolating tinctures :— ‘After the materials have been macerated for forty-eight hours in three fourths of the menstruum ordered, percolation will be most efficiently performed by decanting the liquid, pressing the ingredients in the hand, and carefully packing them, in small portions at a time, in a conical percolator, so that the mass shall be uniformly tight throughout. The decanted liquid may then be poured upon the ingredients and suffered to percolate ; the remainder of the menstruum being afterwards poured upon them in order to chase the strong tincture out. As soon as the liquid ceases to drop, the ingredients are to be removed and pressed. Any deficiency in the product may be made up by adding more of the menstruum and repeating the pressure.’

ceases to pass, continue the percolation with the remaining five ounces of spirit ; then subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient rectified spirit to make one pint.

Uses.—*Internally* :—As a sedative. *Externally* :—As an anodyne.

Doses.—HORSE, 30 to 50 minims.—CATTLE, 40 to 80 minims.—SHEEP, 8 to 12 minims.—PIG, 8 to 12 minims.—DOG, 3 to 10 minims. Given in water, and repeated every two hours.

TINCTURA ALOES COMPOSITA

COMPOUND TINCTURE OF ALOES

Take of

Barbadoes Aloes, in coarse powder	} . 10 drachms.
Myrrh	5 drachms.
Methylated or Rectified Spirit	14 fluid ounces.
Water	6 fluid ounces.

Macerate the aloes and the myrrh in the spirit and the water, previously mixed, for fourteen days in a closed vessel, and shake frequently. Filter and add sufficient proof spirit to make one pint.

Uses.—*Externally* :—Excitant to wounds, obstinate ulcers, and excoriations.

TINCTURA ARNICÆ

TINCTURE OF ARNICA

Take of

Arnica Root, bruised	1 ounce.
Proof Spirit	1 pint.

Macerate the arnica for forty-eight hours in fifteen fluid ounces of the spirit, in a closed vessel, agitating occasionally ; transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Then subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient rectified spirit to make one pint.

Uses.—*Externally* :—Stimulant to sprains and bruises, and in chronic rheumatism.

TINCTURA CAMPHORÆ

See Spiritus Camphoræ.

TINCTURA CANNABIS INDICÆ

TINCTURE OF CANNABIS INDICA

Take of

Extract of Indian Hemp . . .	1 ounce.
Rectified Spirit . . .	20 ounces.

Dissolve.

Uses.—*Internally* :—*See* Extractum Cannabis Indicæ.

Doses.—HORSE, 1 to 2 ounces.—DOG, 10 to 40 minims.

TINCTURA CANTHARIDIS

TINCTURE OF CANTHARIDES

Take of

Cantharides, in coarse powder . . .	$\frac{1}{4}$ ounce.
Proof Spirit	1 pint.

Macerate for seven days in a closed vessel, with occa-

sional agitation ; strain, press, and add sufficient proof spirit to make one pint.

Uses.—*Internally* :—Stimulant and tonic. *Externally* : Tinctura Cantharidis Fortior is used instead.

Doses.—HORSE, 1 to 4 fluid ounces.—CATTLE, 3 to 6 fluid ounces.—SHEEP, $\frac{1}{2}$ to $1\frac{1}{2}$ fluid ounce.—PIG, $\frac{1}{2}$ to $1\frac{1}{2}$ fluid ounce.—DOG, 1 to 3 fluid drachms.

TINCTURA CANTHARIDIS FORTIOR

STRONG TINCTURE OF CANTHARIDES

Take of

Cantharides, in coarse powder .	. 1 ounce.
Proof Spirit 1 pint.

Macerate for seven days in a closed vessel, with occasional agitation ; strain, press, filter, and add sufficient proof spirit to make one pint.

Use.—*Externally* :—Stimulant and rubefacient.

TINCTURA CARDAMOMI COMPOSITA

COMPOUND TINCTURE OF CARDAMOMS

Take of

Cardamom Seeds, freed from their pericarps and bruised	} 1 ounce.
Cinnamon, bruised 2 ounces.
Proof Spirit 80 fluid ounces.

Macerate, with occasional agitation, for forty-eight hours with sixty fluid ounces of the spirit, pack in a percolator, and let it drain ; pour on the remaining spirit, and, when it ceases to drop, press, and wash the mass with sufficient

spirit to make the resulting tincture measure eighty fluid ounces.

Uses.—Carminative and antispasmodic.

Doses.—HORSE, 1 to 4 fluid ounces.—CATTLE, 2 to 6 fluid ounces.—SHEEP, 1 to 2 fluid ounces.—PIG, 1 to 2 fluid ounces.—DOG, 2 to 6 fluid drachms.

Preparation.—Tinctura Chloroformi Composita.

TINCTURA CHLOROFORMI COMPOSITA

COMPOUND TINCTURE OF CHLOROFORM

Take of

Chloroform	1 fluid ounce.
Rectified Spirit	4 fluid ounces.
Compound Tincture of Cardamoms	10 fluid ounces.

Mix.

Uses.—Similar to Spiritus Chloroformi.

Doses.—Same as Spiritus Chloroformi.

TINCTURA CINCHONÆ FLAVÆ

TINCTURE OF YELLOW CINCHONA

Take of

Yellow Cinchona Bark, in moderately fine powder	4 ounces.
Proof Spirit	1 pint.

Macerate the cinchona bark for forty-eight hours in fifteen fluid ounces of the spirit, in a closed vessel, agitating occasionally ; then transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Afterwards subject the

contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient proof spirit to make one pint.

Uses.—*Internally* :—Stimulant, stomachic, and tonic.

Doses.—HORSE, 1 to 3 fluid ounces.—CATTLE, 3 to 6 fluid ounces.—SHEEP, $\frac{1}{2}$ to 1 fluid ounce.—PIG, $\frac{1}{2}$ to 1 fluid ounce.—DOG, 2 to 5 fluid drachms.

TINCTURA COLCHICI SEMINIS

TINCTURE OF COLCHICUM SEEDS

Take of

Colchicum Seeds, bruised	.	.	2 $\frac{1}{2}$ ounces.
Proof Spirit	.	.	1 pint.

Macerate the colchicum for forty-eight hours in fifteen fluid ounces of the spirit, in a closed vessel, agitating occasionally ; then transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Afterwards subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient proof spirit to make one pint.

Uses.—*Internally* :—Sedative. *Externally* :—Anodyne.

Doses.—HORSE, $\frac{1}{2}$ to 2 fluid ounces.—CATTLE, $\frac{1}{2}$ to 2 fluid ounces.—SHEEP, 1 to 4 fluid drachms.—PIG, 1 to 4 fluid drachms.—DOG, 20 to 80 minims.

TINCTURA CONII

TINCTURE OF HEMLOCK

Synonym.—Tinctura Conii Fructus.

Take of

Hemlock Fruit, bruised	.	.	2 $\frac{1}{2}$ ounces.
Proof Spirit	.	.	1 pint.

Macerate the hemlock fruit for forty-eight hours in fifteen ounces of the spirit, in a closed vessel, agitating occasionally ; transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Then subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient proof spirit to make one pint.

Uses.—*Internally* :—Alterative and sedative.

Doses.—HORSE, 1 to 2 fluid ounces.—CATTLE, 1 to 2 fluid ounces.—SHEEP, 3 to 6 fluid drachms.—PIG, 3 to 6 fluid drachms.—DOG, 10 to 40 minims.

TINCTURA CROTONIS

TINCTURE OF CROTON

Take of

Croton Seeds, bruised	.	.	.	1½ ounce.
Rectified Spirit	.	.	.	1 pint.

Macerate for seven days in a closed vessel, with occasional agitation ; strain, press, filter, and add sufficient rectified spirit to make one pint.

Uses.—*Internally* :—Cathartic. *Externally* :—Rube-facient and counterirritant.

Doses.—HORSE, ½ to 1 fluid ounce.—CATTLE, ½ to 1½ fluid ounce.—SHEEP, 1 to 3 fluid drachms.—PIG, 1 to 3 fluid drachms.—DOG, 20 to 50 minims.

TINCTURA ERGOTÆ

TINCTURE OF ERGOT

Take of

Ergot, in coarse powder	.	.	.	5 ounces.
Proof Spirit	.	.	.	1 pint.

Macerate the ergot for forty-eight hours in fifteen fluid ounces of the spirit, in a closed vessel, agitating occasionally ; then transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Afterwards subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient proof spirit to make one pint.

Use.—*Internally* :—Parturient.

Doses.—MARE, 1 to 2 fluid ounces.—Cow, 1 to 3 fluid ounces.—EWE, $1\frac{1}{2}$ to 3 fluid drachms.—Sow, $1\frac{1}{2}$ to 3 fluid drachms.—BITCH, 20 to 60 minims.

TINCTURA FERRI PERCHLORIDI

TINCTURE OF PERCHLORIDE OF IRON

Synonym.—Tinctura Ferri Sesquichloridi.

Take of

Strong Solution of Perchloride of Iron	}	5 fluid ounces.
Rectified Spirit		15 fluid ounces.

Mix, and preserve in a stoppered bottle.

Uses.—*Internally* :—Astringent and tonic in hæmaturia, farcy, many typhoid affections, and indigestion ; also in distemper in dogs. *Externally* :—Astringent, styptic, and caustic.

Doses.—HORSE, 1 to 2 fluid ounces.—CATTLE, 1 to 2 fluid ounces.—SHEEP, 3 to 6 fluid drachms. PIG, 3 to 6 fluid drachms.—DOG, 5 to 20 minims. Diluted with water

TINCTURA IODI¹

TINCTURE OF IODINE

Take of

Iodine	2½ ounces.	1 1/2	35	3 1/2
Iodide of Potassium	1¼ ounce.	30	3 1/2	3 1/2
Rectified Spirit	1 pint.	310	35	3 1/2

Dissolve the iodine and iodide of potassium in the spirit.

Uses.—*Internally*:—Alterative. *Externally*:—Caustic, stimulant, and deobstruent.

Doses.—HORSE, 2 to 6 fluid drachms.—CATTLE, 3 to 8 fluid drachms.—SHEEP, 1 to 2 fluid drachms.—PIG, 20 to 60 minims.—DOG, 10 to 30 minims.

TINCTURA MYRRHÆ

TINCTURE OF MYRRH

Take of

Myrrh, in coarse powder	2½ ounces.
Rectified Spirit	1 pint.

Macerate the myrrh for forty-eight hours in fifteen fluid ounces of the spirit, in a closed vessel, agitating occasionally; then transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Afterwards subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient rectified spirit to make one pint.

Uses.—*Externally*:—Mild astringent and excitant to foul and indolent ulcers, and to canker in the mouth.

¹ This preparation is five times as strong as the Tinctura Iodi of the 'British Pharmacopœia.'

TINCTURA NUCIS VOMICÆ¹**TINCTURE OF NUX VOMICA**

Take of

Nux Vomica	2 ounces.
Rectified Spirit	1 pint.

Apply steam to the nux vomica until it is thoroughly softened, then dry rapidly, and reduce it to fine powder. Macerate the powder for forty-eight hours in fifteen fluid ounces of the spirit, in a closed vessel, agitating occasionally ; then transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Afterwards subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient rectified spirit to make one pint.

Uses.—Stimulant and tonic in paralysis, chorea, and dyspepsia.

Doses.—HORSE, 4 to 8 fluid drachms.—CATTLE, 4 to 8 fluid drachms.—SHEEP, 1 to 2 fluid drachms.—PIG, 1 to 2 fluid drachms.—DOG, 10 to 40 minims.

TINCTURA OPII**TINCTURE OF OPIUM**

Synonym.—Laudanum.

Take of

Opium, in coarse powder	1½ ounce.
Proof Spirit	1 pint.

Macerate for seven days in a closed vessel, with occa-

¹ Morton's Tinctura Nucis Vomicae is two and a half times stronger than this preparation.

sional agitation ; then strain, press, filter, and add sufficient proof spirit to make one pint.

Uses.—Anodyne, hypnotic, sedative, and antispasmodic.

Doses.—HORSE, 1 to 3 fluid ounces.—CATTLE, 1 to 3 fluid ounces.—SHEEP, 2 to 8 fluid drachms.—PIG, 2 to 8 fluid drachms.—DOG, 10 to 30 minims.

Preparations.—Enema Opii ; Linimentum Opii.

TINCTURA PIMENTÆ

TINCTURE OF PIMENTO

Take of

Pimento Berries, bruised	2½ ounces.
Proof Spirit	1 pint.

Macerate for seven days in a closed vessel, with occasional agitation ; then strain, press, filter, and add sufficient proof spirit to make one pint.

Uses.—Antispasmodic and carminative.

Doses.—HORSE, 2 to 4 fluid ounces.—CATTLE, 3 to 6 fluid ounces.—SHEEP, 1 to 3 fluid ounces.—PIG, 1 to 3 fluid ounces.—DOG, 1 to 3 fluid drachms.

TINCTURA SANTALIS RUBRA

TINCTURE OF RED SANDERS WOOD

Take of

Sanders Wood, rasped to powder	½ ounce.
Proof Spirit	1 pint.

Macerate for seven days in a closed vessel, with occasional agitation ; then strain, press, filter, and add sufficient proof spirit to make one pint.

Use.—To impart colour to draughts.

TINCTURA ZINGIBERIS**TINCTURE OF GINGER**

Take of

Ginger, in coarse powder	.	.	.	2½ ounces.
Rectified Spirit	.	.	.	1 pint.

Macerate the ginger for forty-eight hours in fifteen fluid ounces of the spirit, in a closed vessel, agitating occasionally ; then transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Afterwards subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient rectified spirit to make one pint.

Uses.—Carminative, and as an adjunct to tonic, stimulant, and purgative medicines.

Doses.—HORSE, 1 to 2 fluid ounces.—CATTLE, 1 to 3 fluid ounces.—SHEEP, 2 to 6 fluid drachms.—PIG, 2 to 6 fluid drachms.—DOG, ½ to 2 fluid drachms.

UNGUENTUM ACIDI CARBOLICI**OINTMENT OF CARBOLIC ACID**

Take of

Carbolic Acid	1 ounce.
Prepared Lard	6 ounces.

Mix thoroughly.

Uses.—Applied to sloughing and unhealthy wounds.

UNGUENTUM ACIDI CARBOLICI COMPOSITUM

COMPOUND OINTMENT OF CARBOLIC ACID

Take of

Ointment of Carbolic Acid . . . 7 ounces.

Sublimed Sulphur . . . 2 ounces.

Mix thoroughly.

Uses.—In the treatment of scab and mange.

UNGUENTUM ACIDI TANNICI

OINTMENT OF TANNIC ACID

Take of

Tannic Acid . . . 1 part.

Prepared Lard . . . 6 parts.

Mix thoroughly.

Use.—Astringent. Sometimes from $\frac{1}{2}$ to 1 part of opium is added to this ointment.

UNGUENTUM ACONITI ¹

OINTMENT OF ACONITE

Take of

Extract of Aconite . . . 2 ounces.

Prepared Lard . . . 4 ounces.

Mix thoroughly.

Use.—Anodyne.

¹ Unguentum Aconitiæ of the 'British Pharmacopœia' consists of 8 grains of Aconitia, $\frac{1}{2}$ a fluid drachm of Rectified Spirit, and 1 ounce of Prepared Lard.

UNGUENTUM ALUMINIS COMPOSITUM**COMPOUND OINTMENT OF ALUM**

Take of

Alum, in fine powder	.	.	.	1 ounce.
Common Turpentine	.	.	.	1 ounce.
Prepared Lard	.	.	.	3 ounces.

Melt the turpentine and the lard together on a water-bath, add the alum, and stir the mixture while it cools.

Uses.—Astringent and digestive in ‘grease.’

UNGUENTUM ANTIMONII TARTARATI**OINTMENT OF TARTARATED ANTIMONY**

Take of

Tartarated Antimony, in fine powder	1 ounce.
Prepared Lard	4 ounces.

Mix thoroughly.

Uses.—Counterirritant to the chest in pulmonary complaints, and to joints in chronic articular affections.

UNGUENTUM ARGENTI NITRATIS**OINTMENT OF NITRATE OF SILVER**

Take of

Nitrate of Silver, in fine powder	.	5 to 10 grains.
Prepared Lard	.	1 ounce.

Mix thoroughly.

Use.—A piece the size of a pea is placed between the eyelids every third day in chronic ophthalmia.

UNGUENTUM ARSENICI

OINTMENT OF ARSENIC

Take of

Arsenious Acid, in fine powder	. 6 grains.
Prepared Lard	. 1 ounce.

Mix thoroughly.

Use.—Caustic to warts and cancerous growths.

UNGUENTUM CALCIS CHLORATÆ

OINTMENT OF CHLORINATED LIME

Take of

Chlorinated Lime	. 1 ounce.
Prepared Lard	. 4 to 8 ounces.

Mix thoroughly.

Uses.—In grease, especially when accompanied by considerable fœtor.

UNGUENTUM CANTHARIDIS

OINTMENT OF CANTHARIDES

Synonym.—Blistering Ointment.

Take of

Cantharides, in powder	. 1 ounce.
Prepared Lard	. 6 ounces.

Digest the cantharides and lard together over a water-

bath for three hours, with occasional stirring; while hot filter through paper, and allow the clear liquid to cool.

Uses.—Irritant to keep open blisters, and to promote the discharge from setons and ulcers.

UNGUENTUM CREASOTI

OINTMENT OF CREASOTE

Take of

Creasote	1 fluid ounce.
Prepared Lard	8 ounces.

Mix thoroughly.

Uses.—Same as Unguentum Acidi Carbolici.

UNGUENTUM CUPRI SUBACETATIS

OINTMENT OF SUBACETATE OF COPPER

Synonym.—Unguentum *Æruginis*; Ointment of Verdigris.

Take of

Subacetate of Copper, in fine powder	} 1 ounce.
Common Turpentine	
Prepared Lard	12 ounces.

Melt the turpentine and lard together over a water-bath, then withdraw the heat, add the subacetate of copper, and thoroughly mix by stirring the whole until cold.

Use.—Detergent to foul ulcers.

UNGUENTUM GALLÆ

OINTMENT OF GALLS

Take of

Galls, in fine powder	.	.	.	1 ounce.
Prepared Lard	.	.	.	4 ounces

Mix thoroughly.

Uses.—Astringent to ulcers with profuse discharge.

UNGUENTUM GALLÆ ET OPII

OINTMENT OF GALLS AND OPIUM

Take of

Ointment of Galls	.	.	.	5 ounces.
Opium, in fine powder	.	.	.	$\frac{1}{2}$ ounce.

Mix thoroughly.

Uses.—Astringent and anodyne.

UNGUENTUM HELLEBORI NIGRI

OINTMENT OF BLACK HELLEBORE

Take of

Black Hellebore Leaves	.	}	of each, equal parts.
Common Turpentine	.		
Prepared Lard	.		

Melt the turpentine and the lard together in a water-bath, add the hellebore, and digest for one hour; then remove the mixture and express through calico.

Use.—Digestive to wounds of Cattle.

UNGUENTUM HYDRARGYRI

OINTMENT OF MERCURY

Synonym.—Blue Ointment.

Take of

Mercury	1 pound.
Prepared Lard	1 pound.
Prepared Suet	1 ounce.

Rub them together until metallic globules cease to be visible when a small portion of the ointment is rubbed with the finger on brown paper.

Uses.—In scab, mange, and other scurfy cutaneous affections ; to indolent sores and ulcers ; as a resolvent in glandular enlargements ; and applied to the skin with friction to expedite and maintain the constitutional effects produced by the internal administration of mercury. For skin affections, however, the next preparation is generally used.

UNGUENTUM HYDRARGYRI

AMMONIATI

OINTMENT OF AMMONIATED MERCURY

Take of

Ammoniated Mercury	.	.	.	1 ounce.
Prepared Lard	.	.	.	6 to 8 ounces.

Mix thoroughly.

Use.—To destroy pediculi and acari.

UNGUENTUM HYDRARGYRI COMPOSITUM

COMPOUND OINTMENT OF MERCURY

Take of

Ointment of Mercury	1 ounce.
Soft Soap	2 ounces.

Mix thoroughly.

Uses.—*See* Unguentum Hydrargyri.

UNGUENTUM HYDRARGYRI IODIDI RUBRI¹

OINTMENT OF RED IODIDE OF MERCURY

Synonyms.—Unguentum Hydrargyri Biniodidi ; Ointment of Biniodide of Mercury ; Blistering Ointment.

Take of

Red Iodide of Mercury, in fine powder	} 1 ounce. ²
Prepared Lard	
	8 ounces.

Mix thoroughly.

Uses.—Caustic, stimulant, and detergent to unhealthy sores ; counterirritant ; also resolvent in the treatment of curbs, splints, incipient spavins, and enlarged bursæ.

¹ This preparation is nearly three and a half times stronger than Unguentum Hydrargyri Iodidi Rubri of the ' British Pharmacopœia.'

² An ointment containing twice the quantity of iodide of mercury forms a powerful counterirritant when applied to the skin of the ox.

UNGUENTUM HYDRARGYRI NITRATIS


OINTMENT OF NITRATE OF MERCURY

Synonym.—Citrine Ointment.

Take of

Mercury, by weight	.	.	4 ounces.
Nitric Acid	.	.	12 fluid ounces.
Prepared Lard	.	.	15 ounces.
Olive Oil	.	.	32 fluid ounces.

Dissolve the mercury in the nitric acid with the aid of a gentle heat.



Melt the lard in the oil by a steam- or water-bath, in a porcelain vessel capable of holding six times the quantity, and, while the mixture is hot, add the solution of mercuric nitrate, also hot, mixing them thoroughly. If the mixture does not froth up, increase the heat till this occurs. Keep it stirred until it is cold.

Uses.—Stimulant and detergent in grease, ringworm, and other skin affections. One part of this ointment, mixed with one part of lard and one of olive oil, is employed in the treatment of tarsal ophthalmia, and for the removal of nebulæ from the cornea.

UNGUENTUM HYDRARGYRI PERCHLORIDI

OINTMENT OF PERCHLORIDE OF MERCURY

Take of

Perchloride of Mercury, in fine powder	}	1 ounce.
Prepared Lard		
						2 ounces.

Mix thoroughly.

Uses.—Spread on leather for the removal of bony tumours. One part of this ointment, mixed with two parts of lard, is used for dressing Professor Varnell's caustic clams.

UNGUENTUM IODI OINTMENT OF IODINE

Take of

Iodine	1 ounce.
Prepared Lard	8 ounces.
Rectified Spirit or Glycerine	1 drachm.

Handwritten notes:
3 1/2 oz Iodine
8 oz Lard
1 drachm Spirit or Glycerine
Total 12 1/2 oz

Mix thoroughly.

Uses.—Stimulant and resolvent.

UNGUENTUM IODI COMPOSITUM COMPOUND OINTMENT OF IODINE

Take of

Iodine	1/2 ounce.
Iodide of Potassium, in fine powder	1 ounce.
Prepared Lard	8 ounces.
Rectified Spirit or Glycerine	1 drachm.

Handwritten notes:
1/2 oz Iodine
1 oz Iodide of Potassium
8 oz Lard
1 drachm Spirit or Glycerine
Total 10 1/2 oz

Mix thoroughly.

Uses.—Stimulant and resolvent.

UNGUENTUM MYLABRIDIS OINTMENT OF MYLABRIS

Take of

Mylabris, in fine powder	1 ounce.
Prepared Lard	6 ounces.

Digest the mylabris and the lard together over a water-bath for three hours, with occasional stirring; while hot filter through paper, and allow the clear liquid to cool.

Uses.—Same as those of Unguentum Cantharidis, but has no diuretic effect.

UNGUENTUM PICIS LIQUIDÆ

OINTMENT OF TAR

Take of

Common Tar . . .	} of each, equal parts.
Resin . . .	
Prepared Lard . . .	

Melt together with a gentle heat, and stir the mixture while it cools.

Uses.—In foot-rot in sheep, and as a foot ointment for all other domesticated animals.

UNGUENTUM POTASSÆ SULPHURATÆ

OINTMENT OF SULPHURATED POTASH

Take of

Sulphurated Potash, in fine powder	1 ounce.
Prepared Lard	8 ounces.

Triturate the sulphurated potash in a porcelain mortar, and gradually add the lard, rubbing them together until the ointment is perfectly smooth and free from grittiness. This ointment, when used, should be recently prepared.

Uses.—In mange and other skin affections.

UNGUENTUM RESINÆ

OINTMENT OF RESIN

Synonym.—Digestive Ointment.

Take of

Resin, in coarse powder . . .	8 ounces.
Yellow Wax	6 ounces.

Almond Oil	3 ounces.
Prepared Lard	3 ounces.

Melt with a gentle heat, strain the mixture while hot through flannel, and stir constantly while it cools.

Uses.—Stimulant to ulcers, wounds, blistered surfaces, and to impart consistency to other ointments.

UNGUENTUM SABINÆ

OINTMENT OF SAVIN

Take of

Savin Tops, bruised	1 ounce.
Prepared Lard	2 ounces.

Melt the lard and the wax together on a water-bath, add the savin, and digest for twenty minutes. Then remove the mixture, and express through calico.

Uses.—Irritant to blistered surfaces, and for the removal of warts.

UNGUENTUM SULPHURIS

OINTMENT OF SULPHUR

Take of

Sublimed Sulphur	1 ounce.
Prepared Lard	4 ounces.

Mix.

Uses.—For scab in sheep, parasiticide, and in treatment of eczema, mange, &c.

UNGUENTUM SULPHURIS COMPOSITUM**COMPOUND OINTMENT OF SULPHUR**

Synonym.—Sheep Ointment.

Take of

Sublimed Sulphur	.	.	.	1 pound.
Common Turpentine	.	.	.	4 ounces.
Ointment of Mercury	.	.	.	2 ounces.
Linseed Oil.	.	.	.	1 pint.

Melt the oil and turpentine together over a water-bath ; when the mixture is nearly cold, stir in the sulphur, and afterwards the ointment of mercury, until the whole is thoroughly mixed.

Uses.—For eradicating scab in sheep.

UNGUENTUM SULPHURIS IODIDI**OINTMENT OF IODIDE OF SULPHUR**

Take of

Iodide of Sulphur, in fine powder	.	.	.	1 ounce.
Prepared Lard	.	.	.	8 ounces.

Triturate the iodide of sulphur in a porcelain mortar, and gradually add the lard, rubbing them together until the ointment is perfectly smooth and free from grittiness.

Uses.—In mange, urticaria, and other skin diseases.

UNGUENTUM TEREBINTHINÆ**OINTMENT OF TURPENTINE**

Synonym.—Digestive Ointment.

Take of

Common Turpentine . . .	1 ounce.
Prepared Lard	3 ounces.

Melt together over a water-bath, and mix thoroughly by stirring, and cool.

Use.—Stimulant to wounds.

UNGUENTUM VERATRI**OINTMENT OF WHITE HELLEBORE**

Take of

White Hellebore Root, dried, and in } fine powder	1 ounce.
Prepared Lard	4 ounces.

Thoroughly mix.

Uses.—Applied to the skin as a parasiticide, and for charging materials to be used as setons.

UNGUENTUM ZINCI CARBONATIS**OINTMENT OF CARBONATE OF ZINC**

Synonym.—Healing-Ointment.

Take of

Carbonate of Zinc, in fine powder .	1 ounce.
Prepared Lard	6 ounces.

Mix thoroughly.

Uses.—Desiccant and mild astringent to slight excoriations.

UNGUENTUM ZINCI OXIDI

OINTMENT OF OXIDE OF ZINC

Take of

Oxide of Zinc	1 ounce.
Prepared Lard	5 ounces.

Mix.

Uses.—Employed in slight excoriations and ulcerations.


VAPOR CHLORI

CHLORINE GAS

Composition.—An element ; symbol Cl.

Modes of Preparation.—There are two methods commonly resorted to.

1st Method.—Put three or four ounces of powdered manganese peroxide into a flask, basin, or other suitable vessel ; make it into a thin paste with strong hydrochloric acid, and apply a gentle heat to the mixture.



*2nd Method.*¹—Take of


Sodium Chloride (common salt), in fine powder	2½ ounces.
Manganese Peroxide, in fine powder	2 ounces.

¹ Chlorine may also be evolved by pouring diluted sulphuric or hydrochloric acid on chlorinated lime.

Strong Sulphuric Acid	2½ fluid ounces.
Water	5 fluid ounces.

Intimately mix the manganese peroxide and sodium chloride, and place them in a flask, basin, &c. Pour the sulphuric acid into the water contained in porcelain, earthenware, or thin glass vessel. Add the diluted sulphuric acid to the mixture of manganese peroxide and sodium chloride, and thoroughly stir or shake them together.

If necessary, the evolution of chlorine may be promoted by the application of a gentle heat.



Characters and Tests.—Heavy greenish-yellow gas, with a peculiar odour, and producing great irritation when breathed. Bleaches blue litmus paper and develops a blue colour (starch iodide) when brought in contact with paper imbued with a mixture of starch and potassium iodide.

Actions and Uses.—Chiefly employed for disinfecting the air of stables and other buildings. Occasionally used as a therapeutic in glanders, farcy, and similar affections, in which cases it first increases, but ultimately diminishes, the secretion of the nasal membrane. It has also been recommended as an antidote to poisoning by hydrocyanic acid and sulphuretted hydrogen.

Modes of Application.—As an aërial disinfectant, the apparatus is to be placed in the stable, &c. ; the doors and other openings should be closed for several hours during the generation of the gas, and afterwards opened for some time previously to the entrance of any person.

When the gas is to be inhaled the patient should be placed in a loose box, along with the apparatus containing one of the mixtures above described. As soon as the gas begins to exert its irritating effects upon the attendant, he should withdraw, remove the apparatus, and close the door of the box.

Antidotes.—Coughing, and other symptoms of the irritant effects of chlorine, may be allayed by inhaling the vapour of ammonia, ether, alcohol, or hot water.

VASELIN

VASELINE

A substance of the consistence of butter, obtained during the fractional distillation of rock oil.

Composition.—Hexadecane, $C_{16}H_{34}$, and other hydrocarbons.

Actions and Uses.—It dissolves iodine, sulphur iodide, and carbolic acid, it is used as a lubricant, and emollient for irritable, inflamed, or blistered mucous and skin surfaces. It is nearly free from greasiness, and is not oxidisable; it therefore forms a good basis for ointments. It is advisable to use the best varieties of vaseline, as the cheaper preparations, owing to imperfect rectification, frequently cause irritation and even blistering.

VERATRUM

WHITE HELLEBORE

The rhizome of *Veratrum album*. A native of the mountainous regions of Central and Southern Europe.

Natural Order.—*Melanthaceæ*.

Composition.—Its medicinal activity is attributed to an alkaloid named *veratrine*, $C_{37}H_{53}NO_{11}$.

Characters.—Usually consists of the rhizome with the radicles attached ; in pieces of from two to three inches long, about the thickness of the little finger ; covered with a rough dark brown bark ; greyish white internally. In the fresh state it emits a strong disagreeable odour, which it nearly loses by drying, but it retains its acrid, intensely bitter taste.

Actions and Uses.—*Internally* :—In excessive doses an irritant poison ; in medicinal doses nauseant, emetic, sedative, purgative, and anthelmintic. Given sometimes to the horse as a sedative, and as an excitant to promote absorption in œdematous enlargements of the legs. *Externally* :—Applied to the skin as a parasiticide ; also in the cure of mange and other cutaneous diseases.

Doses.—HORSE :—*As a sedative*, 20 to 30 grains every four or six hours, until its depressant action is manifested. *As an excitant*, $\frac{1}{2}$ to 1 ounce.

Modes of Application.—*Internally* :—In the form of bolus, or tincture made with proof spirit. *Externally* :—The powdered rhizome, or Unguentum Veratri.

Antidotes.—Demulcent drinks : also tannic acid, and vegetable astringents containing it. If coma be present, ammonia, or some powerful alcoholic stimulant, should be administered.

Preparation.—Unguentum Veratri.

VINUM FERRI

WINE OF IRON

Composition.—A weak solution of ferrous tartarate, malate, and acetate in sherry.

Take of

Fine Iron Wire (about No. 35)	. 1 ounce.
Sherry 1 pint.

Macerate for thirty days in a closed vessel, the iron being almost, but not quite, wholly immersed in the wine ; frequently shake the vessel, remove the stopper after each shaking, and filter.

Use.—Tonic.

Dose.—Dog, 1 to 4 fluid drachms two or three times a day.

ZINCI ACETAS

ACETATE OF ZINC


Synonym.—Zinc Acetate.

Composition.— $\text{Zn}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 2\text{Aq}$.

Mode of Preparation.—Take of

Carbonate of Zinc	2 ounces.
Acetic Acid	{ 5 ounces,
		{ or a sufficiency.
Distilled Water	6 fluid ounces.

Add the carbonate of zinc in successive portions to three ounces of the acetic acid, previously mixed with the water in a flask ; heat gently ; add by degrees the remainder of the acid till the carbonate is dissolved ; boil for a few minutes, filter while hot, and set it aside for two days to crystallise. Decant the mother-liquor ; evaporate to one half, and again set it aside for two days to crystallise. Place the crystals (zinc acetate) in a funnel to drain, then spread them on filtering-paper on a porous tile, and dry them by exposure to the air at ordinary temperatures.



In the act of crystallising the salt takes up two molecules of water.

Characters and Tests.—Thin, translucent, and colourless crystalline plates, of a pearly lustre, with a sharp unpleasant taste; evolving acetic acid (recognisable by the characteristic odour of its vapour) when decomposed by sulphuric acid; soluble in water, forming a solution which gives a white precipitate (zinc sulphide) with ammonium sulphhydrate. A dilute aqueous solution should not be affected by barium chloride (showing the absence of sulphate) or by silver nitrate (showing the absence of chlorides); and when slightly acidulated with hydrochloric acid, it should furnish no precipitate with sulphuretted hydrogen. After it has been boiled for a few minutes with a little nitric acid (to convert into ferric salt any ferrous compound that may be present), it yields, with ammonia, a white precipitate (zinc hydrate), which is soluble, without colour (if iron be absent), in an excess of the reagent.

Uses.—*Externally*:—Stimulant and astringent to wounds, especially those accompanied by profuse discharges; also to allay superficial inflammation. A wash-leather bandage, saturated with a watery solution of acetate of zinc, is used to reduce the swellings on the legs of hunters.

Modes of Application.—A lotion consisting of twenty-five to fifty grains of the acetate dissolved in one pint of cold water, and applied as soon after the solution of the salt as possible.

ZINCI CARBONAS¹


CARBONATE OF ZINC

Composition.—A mixture or compound of zinc carbonate and zinc hydrate, $\text{ZnCO}_3 \cdot 2\text{Zn}(\text{HO})_2$.

Mode of Preparation.—Take of

Sulphate of Zinc . . .	10 ounces.
Carbonate of Soda . . .	$10\frac{1}{2}$ ounces.
Boiling Distilled Water . .	a sufficiency.

Dissolve the carbonate of soda with a pint of the water in a capacious porcelain vessel, and pour into it the sulphate of zinc, also dissolved in a pint of the water, stirring diligently. Boil for fifteen minutes after effervescence (escape of carbonic anhydride) has ceased, and let the precipitate (carbonate of zinc, B.P.) subside. Decant the supernatant liquid, pour on the precipitate three pints of boiling distilled water, agitating briskly; let the precipitate again subside, and repeat the processes of affusion of hot distilled water and subsidence till the washings are no longer precipitated by barium chloride, showing that sulphate of zinc is completely removed from the carbonate. Collect the precipitate on calico, let it drain, and dry it with a gentle heat.



¹ In veterinary practice the native greyish-brown zinc carbonate (calamine) is generally used, after it has been partially purified by levigation. It is then known as 'Prepared Calamine.'

Characters and Tests.—White, tasteless, inodorous, insoluble in water, soluble with effervescence and without residue in diluted nitric acid. This solution should not be affected by barium chloride, showing the absence of sulphates; or silver nitrate, showing the absence of chlorides; and gives with ammonium carbonate a white precipitate (zinc carbonate and hydrate), entirely soluble, without colour (if iron be absent), in an excess of the reagent, forming a solution which furnishes a white precipitate (zinc sulphide) with ammonium sulphide, showing the presence of zinc.

Actions and Uses.—*Externally* :—Desiccant, stimulant, and astringent.

Mode of Application.—The powder dusted over the wounds, &c., or Unguentum Zinci Carbonatis.

Preparations.—Unguentum Zinci Carbonatis; Zinci Acetas; Zinci Oxidum.

ZINCI CHLORIDUM

CHLORIDE OF ZINC

Synonyms.—Zinc Chloride; Butter of Zinc.¹


Mode of Preparation.—Take of

Granulated Zinc .	. 16 ounces.
Hydrochloric Acid .	. 44 fluid ounces.
Solution of Chlorine .	. a sufficiency.
Carbonate of Zinc .	. $\frac{1}{2}$ ounce, or a sufficiency.
Distilled Water .	. 1 pint.

Put the zinc into a porcelain basin; add by degrees the

¹ Tuson's fluid disinfectant is an aqueous solution of zinc chloride impregnated with sulphurous acid gas (see *Acidum Sulphurosus*).

hydrochloric acid, previously mixed with the water ; and aid the action by gently warming it on a sand-bath until gas (hydrogen) is no longer evolved. Boil for half an hour, supplying the water lost by evaporation, and allow it to stand on a cool part of the sand-bath for twenty-four hours, stirring frequently. Filter the product into a gallon bottle, and pour in the solution of chlorine by degrees, with frequent agitation (to convert any ferrous chloride into ferric chloride), until the fluid acquires a permanent odour of the gas. Add the carbonate of zinc (to precipitate iron and lead, if present) in small quantities at a time, and with renewed agitation, until a brown sediment (ferric hydrate and lead peroxide) appears. Filter through paper into a porcelain basin, and evaporate until a portion of the liquid, withdrawn on the end of a glass rod and cooled, forms an opaque white solid (zinc chloride). Pour it now into proper moulds, and, when the salt has solidified, but before it has cooled, place it in closely stoppered bottles.



Characters and Tests.—Colourless opaque rods or tablets, very deliquescent and caustic ; almost entirely soluble in water, alcohol, and ether. The watery solution gives a white precipitate (zinc sulphide) with ammonium sulphide, showing the presence of zinc ; also a white precipitate (silver chloride) with silver nitrate, indicating that the salt is a chloride ; if first acidulated with hydrochloric acid, it is not affected with sulphuretted hydrogen. Its aqueous solution should give no precipitate with barium chloride, showing the absence of sulphates ; or ammonium oxalate, showing the absence of calcium ; and should not be tinged blue by potassium ferro- or ferric cyanide, showing the absence of iron. Ammonia throws down a white precipi-

tate (zinc hydrate), entirely soluble in an excess of the reagent.

Uses.—In the solid state, or in the form of concentrated solution, as a caustic and deodoriser in canker, quittor, foot-rot in sheep, fistulæ, ill-conditioned wounds, phagedænic ulcers, luxuriant granulations, &c. In the form of Liquor Zinci Chloridi it is applied to wounds as an astringent, stimulant, and antiseptic; to the skin of horses and dogs as a parasiticide, and in eczema; also as a 'dip' or 'wash' for killing ticks and preventing the attacks of the fly on sheep. For deodorising and disinfecting stables, manure heaps, &c.

Preparation.—Liquor Zinci Chloridi.


ZINCI OXIDUM

OXIDE OF ZINC

Synonym.—Zinc Oxide.

Composition.— ZnO .

Mode of Preparation.—Carbonate of zinc is exposed to a dull red heat in a loosely covered Hessian crucible, until a portion taken from the centre of the contents of the crucible and cooled no longer effervesces when dropped into dilute sulphuric acid, showing that the expulsion of carbonic anhydride is complete. Let the crucible cool, and transfer the product (zinc oxide) to a stoppered bottle.



Characters and Tests.—Soft, nearly white, tasteless, and inodorous powder, becoming pale yellow when heated. Dissolves without effervescence in diluted nitric acid, showing absence of carbonate; forming a solution which is

not affected by barium chloride, showing the absence of chlorides ; and gives with ammonium carbonate a white precipitate, zinc carbonate and hydrate, which dissolves entirely without colour (if iron be absent) in an excess of the reagent, forming a solution which gives a white precipitate (zinc sulphide) with ammonium sulphide, indicating the presence of zinc.

Uses.—*Externally* :—Astringent and desiccant to ichorous ulcers, excoriations, &c.

Mode of Application.—In the form of powder, and ointment.

ZINCI SULPHAS

SULPHATE OF ZINC

Synonyms.—Zinc Sulphate ; White Vitriol.

Composition.— $\text{ZnSO}_4 \cdot 7\text{Aq}$.


Mode of Preparation.—Take of

Granulated Zinc . . .	16 ounces.
Sulphuric Acid . . .	12 fluid ounces.
Distilled Water . . .	4 pints.
Solution of Chlorine . .	a sufficiency.
Carbonate of Zinc . . .	$\frac{1}{2}$ ounce, or a sufficiency.

Pour the sulphuric acid, previously mixed with the water on the zinc, contained in a porcelain basin, and, when effervescence (escape of hydrogen) has nearly ceased, aid the action by a gentle heat. Filter the fluid into a gallon bottle, and add gradually, with constant agitation, the solution of chlorine¹ until the fluid acquires a permanent odour of chlorine. Add now, with continued agitation, carbonate of zinc,¹ until a brown precipitate (ferric hy-

¹ The uses of the chlorine water and carbonate of zinc are explained under Zinci Chloridum.

drate and lead peroxide) appears ; let it settle, filter the solution, evaporate till a pellicle forms on the surface, and set aside to crystallise. Dry the crystals (zinc sulphate) by exposure to the air on filtering paper placed on porous tiles. More crystals may be obtained by again evaporating the mother liquor.



In the act of crystallising, the salt acquires seven molecules of water.

Characters and Tests.—Colourless, transparent, prismatic crystals, strongly resembling sulphate of magnesia, but with a metallic astringent taste. Its aqueous solution gives a white precipitate (zinc sulphide) with ammonium sulphide, indicating the presence of zinc, and a white precipitate (barium sulphate) with barium chloride, insoluble in nitric acid, showing that the salt is a sulphate. Its watery solution should give no bluish black precipitate or coloration with tincture of galls, showing the absence of iron, and when acidulated with hydrochloric acid it should not be affected by sulphuretted hydrogen. After it has been boiled for a few minutes with a little nitric acid (to convert into ferric salt any ferrous compound that may be present), it should yield with ammonium hydrate a white precipitate (zinc hydrate), which is entirely soluble, without colour, showing absence of iron, in an excess of the reagent.

Actions and Uses.—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, astringent and tonic. Also emetic when given in sufficient quantity to the pig or dog. *Externally* :—Astringent, excitant, mild escharotic, and desiccant when applied to wounds, injected into sinuses, and in quittor.

Doses.—*Tonic* :—HORSE, 1 to 2 drachms.—CATTLE, 2 to 3 drachms.—SHEEP, 20 to 30 grains.—PIG, 10 to 20 grains. DOG, 2 to 5 grains.

Emetic :—PIG, 30 to 50 grains.—DOG, 8 to 15 grains.

Modes of Application.—*Internally* :—As a tonic, in the form of bolus or dissolved in water ; as an emetic, dissolved in tepid water. *Externally* :—In the state of powder or dissolved in water, the solution varying in strength from half an ounce to seven ounces of the salt to a pint of distilled water, according to the purpose for which the lotion is required.

Incompatibles.—Alkalies and their carbonates ; lime-water ; acetate of lead ; nitrate of silver ; astringent vegetable infusion ; and milk.

Antidotes.—Where possible, warm demulcent drinks, to promote the evacuation of the poison by vomiting.

Preparation.—Zinci Carbonas.

ZINCUM GRANULATUM

GRANULATED ZINC

Composition.—An element, symbol Zn.

Zinc of commerce is heated just sufficiently high to melt it, and the fused metal is poured in a thin stream into a vessel containing about two gallons of cold water. The granulated zinc is to be removed from the water and dried.

Preparations.—Liquor Zinci Chloridi ; Zinci Chloridum ; Zinci Sulphas.

ZINGIBER

GINGER

The scraped and dried rhizome of *Zingiber officinale* obtained from plants cultivated in the West Indies, India, and other countries.

Natural Order.—*Zingiberaceæ*.

Composition.—Its medicinal properties are attributed to the volatile oil, which, according to Thrush, is a complex mixture of hydrocarbons and products of their oxidation.

Characters.—Irregularly lobed decorticated pieces, three or four inches long, subcompressed, yellowish white, but not chalky, on the surface ; has a short mealy fracture, hot taste, and agreeable aroma. Powder yellowish-white.

Actions and Uses.—Stimulant, stomachic, carminative, and tonic. Given in flatulent colic and debility of the stomach and intestines ; also combined with cathartics to increase their activity and to prevent griping.

Doses.—HORSE, 2 to 4 drachms.—CATTLE, 2 to 6 drachms.—SHEEP, $\frac{1}{2}$ to 2 drachms.—PIG, $\frac{1}{2}$ to 1 drachm.—DOG, 10 to 30 grains.

Mode of Application.—In the form of bolus, or as Tinctura Zingiberis.

Preparations.—Massa Aloes ; Syrupus Rhamni ; Tinctura Zingiberis.

WINTER

1880

The first of the season was a very cold one, with a heavy frost on the 1st of December.

The weather was very pleasant on the 2nd of December, with a light frost.

The 3rd of December was a very cold day, with a heavy frost.

The 4th of December was a very cold day, with a heavy frost.

The 5th of December was a very cold day, with a heavy frost.

The 6th of December was a very cold day, with a heavy frost.

The 7th of December was a very cold day, with a heavy frost.

The 8th of December was a very cold day, with a heavy frost.

The 9th of December was a very cold day, with a heavy frost.

The 10th of December was a very cold day, with a heavy frost.

The 11th of December was a very cold day, with a heavy frost.

The 12th of December was a very cold day, with a heavy frost.

The 13th of December was a very cold day, with a heavy frost.

The 14th of December was a very cold day, with a heavy frost.

The 15th of December was a very cold day, with a heavy frost.

The 16th of December was a very cold day, with a heavy frost.


The 17th of December was a very cold day, with a heavy frost.

The 18th of December was a very cold day, with a heavy frost.

The 19th of December was a very cold day, with a heavy frost.

The 20th of December was a very cold day, with a heavy frost.

ADDENDA



PHYSOSTIGMINA

PHYSOSTIGMINE

Synonym.—Eserine.

Composition.— $C_{15}H_{21}N_3O_2$.

Mode of Preparation.—Obtained from the alcoholic extract of Calabar bean, by dissolving the extract in water, adding sodium bicarbonate, shaking the mixture with ether, and evaporating the ethereal liquid.

Characters and Tests.—An alkaloid which occurs in colourless or pinkish crystals, slightly soluble in water, but readily soluble in alcohol and in dilute acids.

The aqueous solution when warmed with, or when shaken with, dilute solution of potash becomes red; and when evaporated to dryness over a water-bath leaves a bluish residue, the acidified solution of which is beautifully dichroic, being blue and red.

Action and Uses.—Hypodermically, as sulphate or salicylate of eserine; the salicylate is preferable, as it keeps for a much longer period.

For therapeutic action see *Physostigmatis Semen*.

Doses.—HORSE, 2 to 3 grains.—DOG, $\frac{1}{20}$ to $\frac{1}{10}$ of a grain.
Hypodermically, one-half these doses.

EXTRACTUM PHYSOSTIGMATIS**EXTRACT OF CALABAR BEAN****Mode of Preparation.—**


Calabar Bean in No. 40 powder	1 pound
Rectified Spirit	4 pints.

Macerate the bean for forty-eight hours with one pint of the spirit in a closed vessel, agitating occasionally ; then transfer to a percolator, and when the fluid ceases to pass add the remainder of the spirit, so that it may slowly percolate through the powder. Subject the residue of the bean to pressure, adding the expressed liquor to the product of the percolation ; filter, distil off most of the spirit, and evaporate what is left in the retort by a water-bath to the consistence of a soft extract.

Preparation.—Physostigmine or Eserine.

Uses.—See *Physostigmatis Semen*.

APPENDIX



SYMBOLS AND ATOMIC WEIGHTS OF THE ELEMENTARY BODIES MENTIONED IN THIS WORK.

Elementary Bodies.	Symbols.	Atomic Weights.
Aluminium	Al	= 27
Antimony (Stibium)	Sb	= 120
Arsenic	As	= 75
Barium	Ba	= 137
Bismuth	Bi	= 219
Boron	B	= 11
Bromine	Br	= 80
Cadmium	Cd	= 112
Calcium	Ca	= 40
Carbon	C	= 12
Chlorine	Cl	= 35.5
Chromium	Cr	= 52.5
Copper (Cuprum)	Cu	= 63.5
Gold (Aurum)	Au	= 196.4
Hydrogen	H	= 1
Iodine	I	= 127
Iron (Ferrum)	Fe	= 56
Lead (Plumbum)	Pb	= 207
Magnesium	Mg	= 24
Manganese	Mn	= 55
Mercury (Hydrargyrum)	Hg	= 200
Nitrogen	N	= 14
Oxygen	O	= 16
Phosphorus	P	= 31
Platinum	Pt	= 195
Potassium (Kalium)	K	= 39
Silver (Argentum)	Ag	= 108
Sodium (Natrium)	Na	= 23
Sulphur	S	= 32
Tin (Stannum)	Sn	= 118
Zinc	Zn	= 65

WEIGHTS AND MEASURES OF THE BRITISH PHARMACOPŒIA ¹

WEIGHTS

1 Grain	gr.			
1 Ounce	oz.		=	437.5 grains
1 Pound	lb.	=	16 ounces	= 7000 „

MEASURES OF CAPACITY

1 Minim	min.			
1 Fluid Drachm	fl. dr.	=	60 minims	
1 Fluid Ounce	fl. oz.	=	8 fluid drachms	
1 Pint	O.	=	20 fluid ounces	
1 Gallon	C.	=	8 pints	

MEASURES OF LENGTH

1 line	=	$\frac{1}{12}$ inch
1 inch	=	$\frac{1}{39.1393}$ second pendulum
12 inches	=	1 foot
36 „	=	3 feet = 1 yard

Length of pendulum vibrating seconds of
 mean time in the latitude of London, } 39.1393 inches
 in a vacuum at the level of the sea . }

RELATION OF MEASURES TO WEIGHTS

1 Minim is the measure of		0.91 grain of water
1 Fluid Drachm „		54.68 grains of water
1 Fluid Ounce „	1 ounce or	437.5 „
1 Pint „	1.25 pound or	8750.0 „
1 Gallon „	10 pounds or	70000.0 „

¹ Although the drachm of sixty grains is discarded by the Editors of the *British Pharmacopœia*, the Author has retained it in the body of his work in consequence of its being a convenient, commonly employed, and well-known quantity in Veterinary Medicine.

WEIGHTS AND MEASURES OF THE METRICAL SYSTEM

WEIGHTS

1 Milligramme	= the thousandth part of one grm. or 0·001 grm.	
1 Centigramme	= the hundredth	0·01 „
1 Decigramme	= the tenth	0·1 „
1 Gramme	= the weight of a cubic centimetre of water at 4°C.	1·0 „
1 Decagramme	= ten grammes	10·0 grms.
1 Hectogramme	= one hundred grammes	100·0 „
1 Kilogramme	= one thousand grammes	1000·0 „

MEASURES OF CAPACITY

1 Millilitre	= 1 cub. centim. or the mea. of 1 grm. of water	
1 Centilitre	= 10	10 grms. „
1 Decilitre	= 100	100 „ „
1 Litre	= 1000	1000 „ (1 kilo.)

MEASURES OF LENGTH

1 Millimetre	= the thousandth part of one metre, or 0·001 metre	
1 Centimetre	= the hundredth	0·01 „
1 Decimetre	= the tenth	0·1 „
1 Metre	= the ten-millionth part of a quarter of the meridian of the earth.	

RELATION OF THE WEIGHTS OF THE BRITISH PHARMACOPŒIA TO THE METRICAL WEIGHTS

1 Pound	= 453·5925 grammes
1 Ounce	= 28·3495 „
1 Grain	= 0·0648 „

RELATION OF MEASURES OF CAPACITY OF THE BRITISH
PHARMACOPŒIA TO THE METRICAL MEASURES

1 Gallon	=	4·543487 litres	
1 Pint	=	0·567936 „	or 567·936 cubic centimetres
1 Fluid Ounce	=	0·028396 „	28·396 „
1 Fluid Drachm	=	0·003549 „	3·549 „
1 Minim	=	0·000059 „	0·059 „


RELATION OF THE METRICAL WEIGHTS TO THE WEIGHTS OF THE
BRITISH PHARMACOPŒIA

1 Milligramme	=	0·015432 grains	
1 Centigramme	=	0·15432 „	
1 Decigramme	=	1·5432 „	
1 Gramme	=	15·432 „	
1 Kilogramme	=	2 lbs. 3 oz. 119·8 grs. or 15432·348 „	

RELATION OF THE METRICAL MEASURES TO THE MEASURES OF
THE BRITISH PHARMACOPŒIA

1 Millimetre	=	0·03937 inches	
1 Centimetre	=	0·39371 „	
1 Decimetre	=	3·93708 „	
1 Metre	=	39·37079 „	or 1 yard 3·7 inches
1 Cubic Centimetre	=	15·432 grain-measures	
1 Litre	=	1 pint 15 oz. 2 drs. 11 m. or 15432·348 grain-measures	

INDEX
TO
VETERINARY MEDICINES
ARRANGED ACCORDING TO THEIR
ACTIONS AND USES



INDEX

TO

VETERINARY MEDICINES

Arranged according to their Actions and Uses.

Adhesives (*adhæreo*, to stick to).—Agents employed to keep the edges of wounds together.

Collodium.

Collodium Flexile.

Emplastrum Picis.

Alteratives (*altero*, to change).—Medicines which gradually change the condition and function of organs from a diseased to a healthy state.

Acidum Arseniosum.

— Nitro-hydrochloricum.

— — dilutum.

Antimonium Sulphuratum.

— Tartaratum.

Argenti Nitras.

Ferri Iodidum.

Hydrargyri Perchloridum.

— Subchloridum.

Hydrargyrum cum Cretâ.

Iodum.

Liquor Arsenicalis.

Oleum Morrhuæ.

Pilula Hydrargyri et Ferri.

Potassa Sulphurata.

Potassii Iodidum.

Pulvis Antimonialis.

Sodæ Bicarbonas.

— Carbonas.

— — exsiccata.

Sodii Chloridum.

Sulphur Sublimatum.

Sulphuris Iodidum.

Tinctura Iodi.

Unguentum Hydrargyri.

— — compositum.

— — Iodidi Rubri.

— — Nitratis.

— Iodi.

— — compositum.

— Sulphuris Iodidi.

Anæsthetics (*a*, *a*, privative, and *αἰσθητικός*, *aisthētikos*, belonging to *αἴσθησις*, *aisthēsis*, or sensation).—Agents which cause insensibility to pain.

Æther.

Chloroform.

Cocaine.

Cold water.

Ice.

Anodynes (*α, α*, privative, and *ὀδύνη, odunē*, pain).—Agents which allay or diminish pain.

Æther.	Spiritus Ætheris Nitrosi.
Enema Opii.	— Camphoræ.
Extractum Aconiti.	— Chloroformi.
— Hyoscyami.	Syrupus Papaveris.
Hyoscyami Folia.	Tinctura Aconiti.
Linimentum Aconiti.	— Belladonnæ.
— Belladonnæ.	— Chloroformi composita.
— Camphoræ.	— Hyoscyami.
— Chloralis Hydratis.	— Opii.
— Chloroformi.	Unguentum Aconiti.
— Opii.	— Gallæ.
Opium.	— — cum Opio.
Spiritus Ætheris.	Unguentum Veratriæ.

Antacids (*ἀντί, anti*, against, and *acidus*, acid).—Agents which neutralise acidity.

Ammoniæ Carbonas.	Mistura Cretæ.
Calcis Carbonas.	Potassæ Bicarbonas.
Calx.	— Carbonas.
Creta Præparata.	Sapo Durus.
Liquor Ammoniæ.	— Mollis.
— Calcis.	Sodæ Bicarbonas.
— — Saccharatus.	— Carbonas.
Magnesia.	— — exsiccata.
Magnesiæ Carbonas.	Spiritus Ammoniæ Aromaticus.

Antemetics (*ἀντί, anti*, against, and *ἐμετικά, emetika*, from *ἐμέω, emeo*, I vomit).—Agents which check or allay vomition.

Acidum Carbolicum.	Creasotum.
— Hydrocyanicum.	Magnesia.
Bismuthi Subnitratis.	Magnesiæ Carbonas.
Caryophyllum.	

Anthelmintics (*ἀντί, anti*, against, and *ἑλμινς, helmins*, a worm).—Agents which kill, expel, or prevent the return of worms.

Aloes.	Enema Sodii Chloridi.
Areçæ Semina.	— Terebinthinæ.
Cambogia.	Filix Mas.
Cusso.	Kamala.
Enema Aloes.	Oleum Terebinthinæ.
— Etheris.	Santonin.
— Quassisë.	Sodii Chloridum.

Antidotes (*ἀντί, anti*, against, and *δίδωμι, didōmi*, I give).—Agents which counteract the injurious effects of poisons.

Antilithics (*ἀντί, anti*, against, and *λίθος, lithos*, a stone).—Agents which dissolve or prevent the formation of calculi and urinary deposits.

Acidum Hydrochloricum dilu-
tum.

— Nitricum dilutum.

Acidum Nitro-hydrochloricum
dilutum.

Antiparasitics (*ἀντί, anti*, against, and *παράσιτος, parasitos*, a parasite).—Agents which prevent the attacks of, or destroy parasites.

Acidum Arseniosum.

— Carbolicum.

— Sulphurosum.

Decoctum Tabaci.

Liquor Calcis Chloratæ.

Oleum Anisi.

Sapo Durus.

— Mollis.

Sodæ Arsenias.

Sulphur Sublimatum.

Sulphuris Iodidum.

Unguentum Hydrargyri Ammo-
niati.

— Sulphuris.

— — Iodidi.

Antiperiodics (*ἀντί, anti*, against, and *περίοδος, periodos*, a period).—Agents which counteract periodicity in disease, *e.g.* in inter-mittent fever.

Acidum Arseniosum.

Cinchonæ Flavæ Cortex.

— Pallidæ Cortex.

— Rubræ Cortex.

Decoctum Cinchonæ Flavæ.

Liquor Arsenicalis.

Liquor Arsenici Hydrochlorici.

— Sodii Arseniatis.

Piper Nigrum.

Quiniæ Sulphas.

Sodii Salicylas.

Antiphlogistics (*ἀντί, anti*, against, and *φλέγω, phlegō*, I burn).—Agents which counteract inflammation. See Sedatives.

Antiseptics (*ἀντί, anti*, against, and *σηπτικός, sēptikos*, putrefying).—Agents which prevent, retard, or arrest putrefaction.

Acidum Arseniosum.

— Aceticum dilutum.

— Carbolicum.

— Sulphurosum.

Cataplasma Fermenti.

Cerevisiæ Fermentum.

Creasotum.

Glycerinum Acidi Carbolic.

Liquor Acidi Carbolic.

Sodii Chloridi.

Zinci Chloridum.

— Sulphas.

Antispasmodics (*ἀντί, anti*, against, and *σπασμός, spasmus*, a spasm).

Agents which prevent or allay the irregular muscular contraction called spasm or cramp.

Æther.	Extractum Hyoscyami.
Assafoetida.	Ipecacuanha.
Belladonnæ Folia.	Spiritus Ætheris.
Cannabis Indica.	Spiritus Ætheris Nitrosi.
Chloralis Hydras.	— Ammoniaë Aromaticus.
Chlorodyne.	— — Foetidus.
Chloroformum.	— Chloroformi.
Cupri Sulphas.	Tinctura Asafoetidæ.
Enema Assafoetidæ.	— Belladonnæ.
— Tabaci.	— Chloroformi composita.
— Terebinthinæ.	— Opii.
Extractum Belladonnæ.	Unguentum Belladonnæ.

Aperients (*aperio*, to open).—Agents which act gently on the bowels, so as to cause mild purgation. See Laxatives.

Aromatics (*ἀρί, ari*, intensely, and *ὀσφ, ozō*, to smell).—Agents possessing an agreeable taste and odour.

Anethi Fructus.	Mistura Cretæ.
Carui Fructus.	Oleum Menthæ Piperitæ.
Cascarillæ Cortex.	— Caryophylli.
Coriandri Fructus.	Pimenta.
Cumini Fructus.	Piper Nigrum.
Enema Assafoetidæ.	Spiritus Ætheris Nitrosi.
Fenugreek.	— Chloroformi.
Infusum Digitalis.	Tinctura Zingiberis.

Arterial Sedatives.—Agents which lower the action of the heart and of the vascular system.

Digitalis and its preparations. | Plumbi Acetas.

Astringents (*ad, to*, and *stringo*, I bind).—Agents which cause a contraction of muscular tissue, promote the coagulation of albuminous fluids, and check secretion.

Acetum.	Calx.
Acidum Carbolicum.	Catechu.
— Gallicum.	— Pallidum.
— Sulphuricum dilutum.	Chlorodyne.
— Tannicum.	Cinchonæ Flavæ Cortex.
Alumen.	Creasotum.
Borax.	Cupri Sulphas.
Calcis Hydras.	Ferri Peroxidum humidum.

Astringents—*continued*.

Ferri Sulphas.	Quercus Cortex.
Galla.	Sodii Chloridum.
Gossypium.	Tinctura Catechu.
Kino.	— Gallæ.
Liquor Ferri Perchloridi fortior.	Unguentum Gallæ.
— Plumbi Subacetatis.	— — et Opii.
— — dilutus.	Zinci Acetas.
Myrrha.	— Carbonas.
Plumbi Acetas.	— Oxidum.
— Carbonas.	— Sulphas.
	Ice and other cold applications.

Carminatives (*carmen*, a verse or charm).—Agents which were formerly supposed to possess the power of *charming* away flatulency and pain in the alimentary canal. See Aromatics.

Cathartics (*καθαίρω*, *kathairo*, I cleanse).—Agents which excite the bowels to increased peristaltic action, and enable them to eject their contents. See Laxatives, Drastics, Hydragogues, and Cholagogues.

Caustics (*καίω*, *kaio*, I burn).—Agents which, by chemical action, destroy the parts to which they are applied. The more powerful caustics produce an eschar (scab or crust), and are hence termed *Escharotics*.

Acidum Aceticum Glaciale.	Hydrargyri Perchloridum.
— Arseniosum.	Liquor Ammoniae fortior.
— Carbolicum.	— Antimonii Chloridi.
— Hydrochloricum.	— Ferri Perchloridi fortior.
— Sulphuricum.	— Hydrargyri Perchloridi.
Alumen exsiccatum.	— Zinci Chloridi.
Argenti Nitras.	Potassa Caustica.
Calx.	Soda Caustica.
Creasotum.	Tinctura Ferri Perchloridi.
Cupri Sulphas.	— Iodidi composita.
Hydrargyri Iodidum Rubrum.	Zinci Chloridum.
— Oxidum Rubrum.	

Cholagogues, *Cholagogue Purgatives* (*χολή*, *cholē*, bile, and *ἀγωγός*, *agōgos*, an expeller).—Agents which have the reputation of in-

creasing the secreting power of the liver and the flow of bile into the intestines. It is probable, however, that they simply cause an emptying of the gall-bladder. They also induce purgation.

Aloes.	Hydrargyrum cum Cretâ.
Hydrargyri Subchloridum.	

Corrosives.—See Caustics.

Counterirritants.—See Irritants.

Demulcents (*demulceo*, to soften).—See Emollients.

Deobstruents (*de*, from, and *obstruo*, I stop up).—Agents which remove any obstruction in the body.

Cambogia.	Magnesiæ Sulphas.
Iodine.	Potassii Iodidum.

Deodorisers (*de*, from, and *odor*, a scent).—Agents which absorb or decompose fetid effluvia. See Antiseptics and Disinfectants.

Desiccants (*desicco*, to dry up).—Agents which dry up mucons discharges and purulent secretions from ulcers and wounds.

Calcis Carbonas.	Plumbi Acetas.
— Hydras.	— Subacetas.
Calx.	Unguentum Plumbi Acetatis.
Catechu.	— — Subacetatis.
Creta Præparata.	— Potassæ Sulphuratæ.
Liquor Calcis.	— Zinci.
Magnesiæ Carbonas.	Zinci Oxidum.

Diaphoretics (*διαφορέω*, *diaphoreo*, to throw off).—Agents which increase the action of the skin, and produce sweating, but in a milder degree than sudorifics.

Disinfectants (*dis*, indicating separation, and *inficio*, I infect).—Agents which render infectious matter inert.

Acidum Arseniosum.	Cataplasma Calcis Chloratæ.
— Carbolicum.	— Carbonis.
— Nitricum.	Liquor Calcis Chloratæ.
— Sulphurosum.	Potassæ Permanganas.
Calx Chlorata.	Vapor Chlori.
Carbo Ligni.	Zinci Chloridum.

Diuretics (*δίᾳ, dia*, through, and *οὐρέω, oureo*, I make water).—
Agents which cause an increased secretion and discharge of urine.

Colchici Cormus.	Potassæ Carbonas.
— Semina.	— Chloras.
Digitalis Folia.	— Nitras.
Extractum Digitalis.	Resina.
Liquor Ammoniae Acetatis.	Sodæ Acetas.
Magnesiæ Sulphas.	— Bicarbonas.
Oleum Terebinthinæ.	— Carbonas.
Pix Burgundica.	Spiritus Ætheris Nitrosi.
Potassæ Acetas.	Terebinthina Canadensis.
— Bicarbonas.	Tinctura Digitalis.

Drastic Purgatives (*δραστικός, drastikos*, effective, powerful).—
Violent cathartics, which produce speedy and copious evacuations.

Cambogia.	Oleum Terebinthinæ.
Jalapa.	Rhamni Succus.
Oleum Crotonis.	Terebinthina Canadensis.

Ecbolics (*ἐκ, ek*, out of, and *βάλλω, ballo*, I throw).—Agents which cause the uterus to contract and expel its contents.

Ergota.	Infusum Ergotæ.
---------	-----------------

Emetics (*ἐμετικά, emetika*, from *ἐμέω, emeo*, I vomit).—Agents which cause vomition.

Antimonium Tartaratum.	Zinci Acetas.
Cupri Sulphas.	— Sulphas.
Ipecacuanha.	

Emollients (*emollio*, to soften).—Agents which soften the part to which they are applied, and diminish irritation. When used internally to protect the mucous membranes of the alimentary canal from the action of irritants, they are termed *Demulcents*.

Acaciæ Gummi.	Glycerinum.
Adeps Præparatus.	Infusum Lini.
Albumen Ovi.	Lini Farina.
Amylum.	— Semina.
Cera Flava.	Mucilago Acaciæ.
Collodium.	Oleum Olivæ.
— Flexile.	Sevum Præparatum.

Epispastics (ἐπισπάω, *epispao*, to draw to).—Agents which, when applied to the skin, produce a vesicle or blister.

Acetum Cantharidis.	Linimentum Crotonis.
Acidum Aceticum Glaciale.	Mylabris.
Cantharis.	Unguentum Cantharidis.
Linimentum Cantharidis.	— Mylabridis.

Escharotics (ἐσχαρόω, *escharoō*, to scab over).—See Caustics.

Excitants (*excitans*, exciting, stimulating).—See Stimulants.

Febrifuges (*febris*, a fever, and *fugo*, to drive away).—Agents which moderate or abate the violence of fevers.

Hæmatinics (αἱματίνα, *hæmatina*, the red colouring matter of the blood).—Preparations of iron, which increase the number of red corpuscles in the blood.

Cambogia.	Ferri Sulphas.
Ferri Carbonas Saccharata.	Manganesii Oxidum Nigrum.
— Iodidum.	Pilula Hydrargyri et Ferri.
— Peroxidum humidum.	

Hæmostatics (αἷμα, *haima*, blood, and στάσις, *stasis*, a standing).—See Styptics.

Hydragogues (ὑδωρ, *hudōr*, water, and ἀγωγός, *agōgos*, expeller).—Cathartics which have the power of causing a very large secretion of fluid from the mucous membrane of the intestines.

Hypnotics (ὑπνος, *hupnos*, sleep).—See Narcotics.

Irritants (*irrito*, to excite).—Agents which stimulate and irritate the skin or other parts to which they are applied. See derivatives, Rubefacients, Epispastics, and Pustulants.

Laxatives (λαξω, to loosen).—Mild cathartics.

Aloes.	Oleum Ricini.
Jalapa.	Potassæ Sulphas.
Magnesia.	Rhamni Succus.
Magnesiae Carbonas.	Rhei Radix.
— Sulphas.	Sodæ Sulphas.
Oleum Lini.	Sulphur Sublimatum.
— Olivæ.	Theriaca.

Lithontriptics (λίθος, *lithos*, a stone, and τρίβω, *tribo*, to rub down).—See Antilithics.

Narcotics (νάρκη, *narkē*, stupor).—Agents which allay pain and produce sleep.

Chloroformum.

Opium.

Tinctura Opii.

Nutritives (*nutrio*, to nourish).—Agents which facilitate the assimilative process, and improve the condition of the tissues.

Oleum Liní.

— Morrhuae.

Oleum Olivæ.

Sevum Præparatum.

Purgatives (*purgo*, to cleanse).—See Cathartics.

Pustulants (πύον, *puon*, pus).—Agents which, when applied to the skin, produce a pustule.

Antimonium Tartaratum.

Argenti Nitras.

Oleum Crotonis.

Refrigerants (*refrigero*, to cool).—Agents which diminish heat, lower the circulation, and quench thirst.

Acetum.

Acidum Hydrochloricum dilu-
tum.

— Nitricum dilutum.

— Sulphuricum dilutum.

Ammonii Chloridum.

Potassæ Chloras.

— Nitras.

— Permanganas.

Liquor Ammoniae Acetatis.

Spiritus Ætheris Nitrosi.

Resolvents (*resolvo*, I unloose).—Agents having the power to dissolve or relax.

Rubefacients (*rubefacio*, to make red).—Agents which, when applied to the skin, irritate and redden it.

Acetum Cantharidis.

Cataplasma Sinapis.

Hydrargyri Perchloridi.

Iodum.

Liquor Ammoniae (weak solu-
tion).

Linimentum Ammoniae.

— — compositum.

Linimentum Camphoræ Compo-
situm.

Oleum Terebinthinæ.

Sedatives (*sedo*, to allay).—Agents which depress nervous power without previously exalting it.

Acidum Carbolicum.	Hydrargyri Subchloridum.
— Hydrocyanicum dilutum.	Hyoscyami Folia.
Aconiti Folia.	Ipecacuanha.
— Radix.	Liquor Plumbi Subacetatis.
Antimonii Oxidum.	Opium.
Antimonium Tartaratum.	Plumbi Acetas.
Belladonnæ Folia.	Spiritus Camphoræ.
Camphora.	Syrupus Papaveris.
Colchici Cormus.	Tinctura Aconiti.
— Semina.	— Camphoræ composita.
Creasote.	— Opii.
Digitalis Folia.	Zinci Oxidum.
Extractum Aconiti.	Ether Spray.
— Belladonnæ.	

Sialagogues (*σίαλον*, *siālon*, saliva, and *ἄγω*, *ago*, I lead).—Agents which promote or increase the secretion of saliva.

Antimonium Tartaratum.	Potassii Iodidum.
Hydrargyri Subchloridum.	Sinapis.
Hydrargyrum cum Cretâ.	Zingiber.
Iodine.	

Soporifics (*sopor*, a heavy sleep, and *fero*, to bring).—See Narcotics.

Stimulants (*stimulo*, to excite).—Agents which rapidly, but transiently, excite the nervous system to increased activity, and which, as a consequence, induce a more energetic performance of the functions of the whole body.

Acidum Aceticum dilutum.	Infusum Gentianæ compositum.
— Hydrochloricum dilutum.	Nux Vomica.
Ammoniæ Carbonas.	Pimenta.
Ammonii Chloridum.	Rhei Radix.
Arnicae Radix.	Sinapis.
Assafoetida.	Spiritus Chloroformi.
Cantharis.	— Rectificatus.
Capsici Fructus.	— — tenuior.
Cardamomum.	Tinctura Aloes composita.
Catechu Pallidum.	— Chloroformi composita.
Chloralis Hydras.	— Cinchonæ Flavæ.
Chloroform.	— Gentianæ composita.
Gentianæ Radix.	Zingiber.
Infusum Cinchonæ Flavæ.	Alcoholic Beverages.

Styptics (στυφω, *stupho*, I restrain).—Agents which arrest bleeding.

Acidum Aceticum dilutum.	Liquor Plumbi Subacetatis.
— Carbolicum.	Myrrha.
— Gallicum.	Plumbi Acetas.
— Tannicum.	— Carbonas.
Alumen.	Quercus Cortex.
Catechu.	Spiritus Rectificatus.
Collodium Hæmostaticum.	Tinctura Ferri Perchloridi.
Cupri Sulphas.	— Gallæ.
Ergota.	— Myrrhæ.
Ferri Sulphas.	Zinci Acetas.
Galla.	— Carbonas.
Gossypium.	— Oxidum.
Infusum Ergotæ.	— Sulphas.
Liquor Calcis.	Cold Applications.
— Ferri Perchloridi fortior.	

Sudorifics (*sudor*, to sweat, and *fiō*, to become).—Agents which increase the exhalant functions of the skin, and produce sweating.

Ammoniæ Carbonas.	Potassæ Nitras.
Antimonii Oxidum.	Pulvis Antimonialis.
Camphora.	— Ipecacuanhæ compositus.
Hyoscyami Folia.	Spiritus Ætheris Nitrosi.
Ipecacuanha.	— Camphoræ.
Liquor Ammoniæ.	Sulphur Præcipitatum.
— — Acetatis.	Terebinthina Canadensis.
Oleum Terebinthinæ.	

Tonics (τείνω, *teino*, to stretch, the quality of muscular fibre in an active state).—Agents which gradually improve the condition and functions of the digestive organs, so as to enable the system to acquire increased vigour.

Acidum Hydrochloricum dilutum.	Cupri Sulphas.
— Nitricum dilutum.	Decoctum Cinchonæ Flavæ.
— Sulphuricum dilutum.	Ferri Carbonas Saccharata.
Aloes.	— Iodidum.
Anthemidis Flores.	— Peroxidum Hydratum.
Argenti Nitras.	— Sulphas.
Cerevisiæ Fermentum.	Gentianæ Radix.
Cinchonæ Flavæ Cortex.	Infusum Cinchonæ Flavæ.
	— Gentianæ compositum.

Tonics—*continued*.

Iodum.	Tinctura Ferri Perchloridi.
Liquor Ferri Perchloridi.	— Gentianæ composita.
Quiniæ Sulphas.	— Myrrhæ.
Rhei Radix.	— Nucis Vomicæ.
Sodii Chloridum.	Vinum Ferri.
Tinctura Cantharidis.	Zinci Acetas.
— Cinchonæ Flavæ.	— Sulphas.

Traumatics (*τραῦμα, trauma*, a wound).—Agents applied to wounds.

Collodium.	Tinctura Aloes composita.
Myrrh.	Unguentum Terebinthinæ.
Resin.	Zinci Oxidum.

Vermicides (*vermis*, a worm, and *cædo*, to kill).—See Anthelmintics.**Vermifuges** (*vermis*, a worm, and *fugo*, to drive away).—See Anthelmintics.**Vesicants** (*vas*, a vessel, a bladder).—See Epispastics.

INDEX

	PAGE		PAGE
ACACIA Gum	1	Acid, Nitric	16
— — Mucilage of	1	— — Diluted	17
Acaciæ Gummi	1	— Nitro-hydrochloric	18
— Mucilago	1	— Phenic	8
Acetanilide	1	— Prussic	14
Acetic Acid	4	— — Scheele's	14
— — Diluted	5	— Phosphoric, concentrated	19
— — Glacial	5	— — Dilute	20
— — Hydrate	5	— Pyroligneous	5
Acetum	2	— Salicyclic	20
— Cantharidis	3	— Sulphuric	21
Acid, Acetic	4	— — Dilute	23
— — Dilute	5	— Sulphurous	24
— — Glacial	5	— Tannic	25
— Arsenious	6	— — Ointment of	297
— Boracic	7	Acidi Carbolicæ Glyceri-	
— Boric	7	num	126
— Carbolic	8	— — Linimentum	156
— — Glycerin of	126	— — Liquor	156
— — Liniment of	156	— — Lotio	183
— — Lotion of	183	— — Unguentum	296
— — Ointment of	296	— — — compositum	297
— — — Compound of	297	— Tannici Unguentum	297
— — Solution of	156	Acidum Aceticum	4
— Chromic	10	— — dilutum	5
— Gallic	11	— Glaciale	5
— Hydrochloric	12	— Arseniosum	6
— — Dilute	13	— Boricum	7
— Hydrocyanic, Dilute	14	— Carbolicum	8
— Muriatic	12	— Chromicum	10

	PAGE		PAGE
Acidum Gallicum	11	Aloe Barbadensis	32
— Hydrochloricum	12	— Capensis	33
— — dilutum	13	— Socotrina	34
— Hydrocyanicum dilutum	14	Aloes, Barbadoes	32
— Nitricum	16	— Cape	33
— — dilutum	17	— Enema of	106
— Nitro-hydrochloricum dilu- tum	18	— Liquor	157
— Phosphoricum concentra- tum	19	— — et Sodæ Carbonatis	157
— — dilutum	20	— Massa	190
— Salicylicum	20	— — composita	191
— Sulphuricum	21	— Socotrine	34
— — dilutum	23	— Solution of	157
— Sulphurosum	24	— — — and Carbonate of Soda	157
— Tannicum	25	— Tinctura, composita	286
Aconite Leaves	27	— Tincture of, Compound	286
— Extract of	110	Aloin	35
— Ointment of	297	Allspice	219
— Root	28	Alterative Mass	191
— Tincture of	285	Alum	35
Aconiti, Extractum	110	— Ammonium	35
— Folia	27	— Burned	37
— Radix	28	— Compound Ointment of	298
— Tinctura	285	— Dried	37
— Unguentum	297	— Ointment of, Compound	298
Aconitia	29	— Potassium	35
Aconitina	29	— Solution of	157
Aconitine	29	Alumen	35
Adeps Præparatus	29	— Exsiccatum	37
Æruginis, Unguentum	300	Aluminis, Liquor	158
— Linimentum	152	— Unguentum compositum	298
Ærugo	99	Amido-chloride of Mercury	136
Æther	30	Ammonia Acetate, Solution of	159
Ætheris, Spiritus	268	— — Lotion of	183
— Nitrici, Spiritus	269	— Aromatic Spirit of	271
— Nitrosi, Spiritus	269	— Carbonate of	38
— Sulphurici, Spiritus	268	— Chloride of	40
Albumen, Egg	32	— Fetid Spirit of	271
— Ovi	32	— Mixture	195
Alcohol Ethylic	274	— Muriate of	40
Phenylic	8	— Solution of	158

	PAGE		PAGE
Ammonia, Strong Solution of	161	Antimonii Tartarati Unguen-	
Ammoniacum	37	tum	298
— Gum	37	— — Liquor	163
Ammoniæ Acetatis, Liquor .	159	Antimonious Chloride, Solu-	
— — Lotio	183	tion of	162
— Aromaticus Spiritus . . .	271	— Oxide	44
— Carbonas	38	— Sulphide	46
— Fœtidus Spiritus	271	Antimonium Nigrum	46
— Fortior Liquor	161	— Tartaratum	46
— Linimentum	149	Antimony, Black	46
— Liquor	158	— Butter of	162
— Mistura	195	— Chloride of, Solution of .	162
Ammoniated Mercury	136	— Ointment of Tartarated .	298
— Mustard Poultice	76	— Oxide of	44
Ammonii Chloridum	40	— Solution of Tartarated .	163
— Chloridi et Camphoræ,		— — Chloride of	162
Lotio	184	— — Terchloride of	162
— — et Potassæ Nitratis		— Tartarated	46
Lotio	184	— — Compound Mass of .	191
Ammonium Alum	35	— — Solution of	163
— Chloride	40	— Terchloride, Solution of .	162
— — Lotion of	184	— Teroxide	46
— — Lotion of, and Potassium		— Tersulphide	46
Nitrate	184	— Trichloride	162
— — and Camphor	184	— Trioxide	44
Amyl Nitris	42	— Trisulphide	46
— Nitrite	42	Antipyrin	48
Amyli Mucilago	201	Apomorphine	200
Amylum	41	Aqua Calcis	167
Anhydride, Arsenious	6	— Destillata	49
— Sulphurous	24	— Fortis	16
Anise Fruit	43	— Menthæ Piperitæ	50
— Oil of	204	— Regia, Diluted	18
Aniseed	43	Areca Nut	50
Anisi Fructus	43	— Seeds	50
— Oleum	204	Arecae Semina	50
Anthemidis Flores	43	Argenti Nitras	51
Antichlor	263	— Nitratis, Liquor	164
Antifebrin	1	— — Unguentum	298
Antimonii Chloridi, Liquor .	162	Arnica Root	53
— Oxidum	44	— Tincture of	286
— Tartarati composita, Massa	191	Aromatic Spirit of Ammonia .	271

	PAGE		PAGE
Arsenic, Hydrochloric Solu-		Black Pepper	220
tion of	166	— Pitch	223
— Ointment of	299	— Wash	188
— White	310	Blistering Fly	68
Arsenical Solution	164	— Chinese Fly	202
Arsenicalis, Liquor	164	— Ointment	299, 303
Arsenici Hydrochloricus, Li-		Blue Ointment	302
quor	166	— Pill, Ferruginate	219
— Unguentum	299	— Stone	100
Arsenious Acid	6	— Vitriol	100
— Anhydride	6	Boli (<i>see</i> Mass)	
Assafoetida	53	Borax	58
— Enema of	106	Bran Poultice	75
Assafoetidæ, Enema	106	British Gum	201
Astringent Mass	192	Buchu Folia	59
Atropia	54	— Leaves	59
Atropina	54	Buckthorn Juice	282
Atropine	54	— Syrup of	282
Axungia	29	Burgundy Pitch	221
		Burnett's, Sir William, Fluid .	181
BALLS (<i>see</i> Mass)		Butter of Antimony	162
Balsam, Canada	283		
— of Peru	55	CALABAR Bean	218
Balsamum Peruvianum	55	Calamine	316
Barbadoes Aloes	32	Calcii Chloridum	60
Bean, Calabar,	218	Calcis, Aqua	167
Belladonna, Extract of	110	— Chloratæ, Liquor	168
— Leaves	55	— — Unguentum	299
Belladonna, Compound Mass		— Hydras	61
of	192	— Linimentum	150
— Root	57	— Liquor	167
Belladonnæ, Extractum	110	Calcium Carbonate	93
— Folia	55	— Chloride	60
— Massa composita	192	— Chloride of	60
— Radix	57	— Hydrate	61
Benzoin	57	— Hypochlorite	64
Benzoinum	57	— Oxide	63
Betel Nut	50	— Oxide of	63
Black Antimony	46	Calomel	132
— Hellebore, Ointment of	301	Calumba Root	62
— Mercurial Lotion	185	Calumbæ Radix	62

	PAGE		PAGE
Calves', Sheep and, Cordial	196	Capsicine	69
Calx	63	Capsicum Fruit	69
Calx Chlorata	64	Caraway Fruit	72
Cambogia	65	— Seeds	72
Camphor	66	Carbolic Acid	8
— Compound Liniment of	150	— — Glycerin of	126
— Liniment of	150	— — Liniment of	156
— Lotion of Chloride of		— — Lotion of	183
Ammonium and Cam-		— — Ointment of	296
phor	184	— — — Compound	297
— Spirit of	272	— — Solution of	156
Camphora	66	Carbo Ligni	70
Camphoræ, Linimentum	150	Cardamomi, Tinctura, compo-	
— — Compositum	150	sita	281
— Lotio Ammonii Chloridi et	184	Cardamoms	71
— Spiritus	272	— Tincture of, Compound	281
— Tinctura	287	Cardamomum	71
Camphorated Oil	206	Carron Oil	150
Canada Balsam	283	Carui Fructus	72
Cantharis	68	Caryophyllum	73
Cantharis Vesicatoria	68	Cascarilla Bark	73
Cantharides	68	Cascarillæ Cortex	73
— Liniment of	151	Cascarillin	73
— Oil of	206	Castor Oil	213
— Ointment of	299	— — Mixture	196
— Solution of, Terebinthi-		Cataplasma Calcis Chloratæ	74
nated	169	— Carbonis	74
— Strong Tincture of	287	— Fermenti	75
— Terebinthinated Solution		— Furfuris	75
of	169	— Lini	75
— Tincture of	287	— Sinapis	76
— — Strong	288	— — Ammoniatum	76
— Vinegar of	3	Catechu composita, Massa	192
Cantharidis Acetum	3	— Infusion of	139
— Linimentum	151	— Infusum	139
— Oleum	206	— Pale	77
— Terebinthinatus, Liquor	169	— Pallidum	77
— Tinctura	287	Cathartic Mass	190
— — Fortior	288	Caustic Potash	176
— Unguentum	299	— Soda	179
Cantharidin	68	Cera Flava	78
Capsici Fructus	69	Cerii Oxalas	79

	PAGE		PAGE
Cerevisiæ Fermentum	79	Colchici Semina	89
Chalk	93	— Seminis Tinctura	290
— Mercury with	137	Colchicum Corm	88
— Mixture, Compound	196	— Seeds	89
— Prepared	93	— — Tincture of	290
Chamomile Flowers	43	Collodion	90
Charcoal Poultice	74	— Flexible	90
— Wood	70	— Hæmostatic	91
Chinese Blistering Fly	202	Collodium	90
Chloral Hydrate	80	— Flexile	90
— Hydras	80	— Hæmostaticum	91
Chloric Ether	272	Colocynth	91
Chlorinated Lime	64	Colocynthis	91
— — Poultice	74	Common Mass	192
Chlorine gas	310	— Salt	266
Chlorodyne	81	Compound Chalk Mixture	196
Chloroform	82	— Infusion of Gentian	140
Chloroform mixture, Com-		— Liniment of Ammonia	149
pound	81	— — of Creasote	151
Chloroform, Spirit of	272	— — of Iodine	153
— Tincture of, Compound	289	— — of Soap	155
Chloroformi, Spiritus	272	— — of Turpentine	155
— Mistura Hydrocyanata	81	— Mass of Aloes	191
— Tinctura, composita	289	— — of Belladonna	192
Chloroformum	82	— — of Catechu	192
Cinchona Bark, Pale	86	— — of Digitalis	193
— — Red	86	— — of Ginger	195
— — Yellow	84	— — of Resin	194
— Infusion of Yellow	139	— — of Tartarated Antimony	191
— Tincture of Yellow	289	— Ointment of Carbolic Acid	297
Cinchonæ Flavæ Cortex	84	— — of Alum	298
— — Infusum	139	— — of Mercury	303
— — Tinctura	289	— — of Iodine	305
— Pallidæ Cortex	86	— — of Sulphur	308
— Rubræ Cortex	86	— Powder of Carbonate of	
Cinnamon	87	Zinc	246
Cinnamomum	87	— — of Ipecacuanha	244
Citrine Ointment	304	— — of Opium	245
Cloves	73	— — of Rhubarb	245
Cocaine	87	— — of Sulphate of Iron	244
Cod Liver Oil	211	— Solution of Sulphate of	
Colchici Cormus	88	Copper	170

	PAGE		PAGE
Compound Spirit of Perchloride of Mercury . . .	273	Creasoti Unguentum . . .	300
— Tincture of Aloes . . .	286	Creasotum	92
Condy's Fluid	238	Creta	93
Confectio Rosæ Gallicæ . . .	92	— Præparata	93
Confection of Roses	92	— Hydrargyrum cum . . .	137
Conii, Tinctura	290	Cretæ composita, Mistura .	196
— Fructus, Tinctura	290	Crocus	95
Cooling Lotion	184	Croton Beans	95
Copper, Ammonio-Sulphate of	98	— Oil	207
— Compound Solution of Sulphate of	170	— — Liniment of	152
— Diacetate of	99	— Seeds	95
— Iodide of	98	— Tincture of	291
— — with Iodine	98	Crotonis, Oleum	207
— Liniment of Subacetate of .	152	— Linimentum	152
— — of Sulphate of	153	— Semina	95
— Mass of Sulphate of	193	— Tigllii Semina	95
— Subacetate of	99	— Tinctura	291
— — Liniment of	152	Cumin Fruit	96
— Sulphate of	100	Cumini Fructus	96
— Sulphate, Compound Solution of	170	Cupri Ammonio-Sulphas . .	97
— — Liniment of	153	— Iodidum cum Iodo . . .	98
— — Mass of	193	— Subacetas	99
— — Solution of	169	— Subacetatis, Linimentum .	152
Copperas	100	— — Unguentum	300
Cordial Mass	195	— Sulphatis Massa	193
— Sheep and Calves'	196	— — Linimentum	152
Coriandri Fructus	92	— — Liquor	169
Coriander Fruit	92	— — Compositus, Liquor .	170
— Seed	92	— Sulphas	100
Corrosive Sublimate	130	Cupric Sulphate	100
Cotton Wool	126	Cusso	102
Cough Ball	192		
Creasote	92	DANDELION	283
— Compound Liniment of . . .	151	Decoctum Tabaci	103
— Mixture	195	Decoction of Tobacco . . .	103
— Ointment of	300	Digestive Ointment	306
Creasoti compositum, Linimentum	151	Digitalin	104
		Digitalinum	104
		Digitalis composita, Massa .	193
		— Extract of	112
		— Extractum	112

	PAGE		PAGE
Digitalis Folia	104	Ethylic Oxide	30
— Leaves	104	Eucalyptus	109
Dinitro-cellulose	246	Euphorbium	109
Dip, Sheep	165	Extract of Aconite	110
Discutient Lotion	184	— of Belladonna	110
Distilled Water	49	— of Digitalis	112
Diuretic Mass	194	— of Indian Hemp	111
Dover's Powder, Factitious	244	— Liquid, of Ergot	113
		— of Male Fern	113
EGG Albumen	32	— Goulard's	174
Emetic, Tartar	46	— of Hyoscyamus	114
Emplastrum Picis	105	— of Physostigmine	328
Enema Aloes	106	Extractum Aconiti	110
— of Aloes	106	— Belladonnæ	110
— Assafœtidæ	106	— Cannabis Indicæ	111
— of Assafœtida	106	— Digitalis	112
— Magnesiæ Sulphatis	106	— Ergotæ Liquidum	113
— Opii	107	— Filicis Liquidum	113
— of Opium	107	— Hyoscyami	114
— of Sulphate of Magnesia	106	— Physostigmatis	328
— Tabaci	107		
— Terebinthinæ	107	FACTITIOUS Dover's Powder	244
— of Tobacco	107	— James's Powder	244
— of Turpentine	107	Fecula	41
Epsom Salts	188	Fern, Male	122
Ergot	107	— — Liquid Extract of	113
— Infusion of	140	— — Shield	122
— Liquid Extract of	113	— Root	122
— of Rye	107	Ferri Carbonas Saccharata	115
— Tincture of	291	— Iodidum	117
Ergota	107	— Perchloridi Fortior, Liquor	170
Ergotæ, Extractum, liquidum	113	— — Tinctura	292
— Infusum	140	— Peroxidum Humidum	118
— Tinctura	291	— — Hydratum	118
Eserine	327	— Persulphatis, Liquor	171
Ether	30	— Pilula Hydrargyri et	219
— Chloric	272	— Protosulphas	119
— Ethylic	30	— Sesquichloridi, Tinctura	292
— Spirit of	268	— Sulphas	119
— — of Nitrous	269	— Sulphatis Compositus, Pulv.	244
— Sulphuric	30	— — Massa	194
Ethylic Ether	30		

	PAGE		PAGE
Ferri Vinum	313	Gossypium	126
Ferric Hydrate, Moist	118	Goulard's Extract	174
Ferruginate Blue Pill	219	— Water	175
Ferrum	121	Granulated Zinc	322
Fetid Spirit of Ammonia	271	Green Vitriol	119
Fever Ball	191	Gregory's Mixture	245
— Mass	191	— Powder	245
Filix Mas	122	Grey Powder	137
Flowers of Sulphur	278	Gum Acacia	1
Fluid, Sir Wm. Burnett's	181	— — Mucilage of	1
— Condyl's	178	— Ammoniacum	37
Fly, Chinese Blistering	202	— Arabic	1
Formyl Trichloride	82	— Benzoin	57
Fowler's Solution	164	— British	201
		— Euphorbium	109
		Gun Cotton	246
GALBANUM	123		
Galla	123	HARD Soap	254
Gallæ, Unguentum	301	Healing Ointment	309
— — et Opii	301	Hellebore, Black, Ointment of	301
Gallic Acid	11	— White	309
Galls	123	— — Ointment of	309
— Nut	123	Hellebori Nigri, Unguentum	301
— Oak	123	Hemlock, Tincture of	290
— Ointment of	301	Henbane Leaves	138
— and Opium, Ointment of	301	Hepar Sulphuris	227
Gamboge	65	Hog's Lard	29
Gentian, Compound Infusion		Hydrargyri Ammoniati, Un-	
of	140	guentum	302
— Radix	124	— Ammonio-chloridum	136
— Root	124	— Bichloridi, Liquor	172
Gentianæ compositum, In-		— Biniodidi, Unguentum	303
fusum	140	— Flava Lotio	185
Ginger	323	— Iodidi Rubri, Unguentum	303
— Compound Mass of	195	— Iodidum Rubrum	127
— Tincture of	296	— Nigra Lotio	188
Glacial Acetic Acid	5	— Nitrico-oxidum	128
Glauber's Salt	264	— Oxidum Rubrum	128
Glycerin	125	— Perchloridi compositus,	
— of Carbolic Acid	126	Spiritus	273
Glycerinum	125	— — Liquor	172
— Acidi Carbolici	126		

	PAGE		PAGE
Hydrargyri Perchloridum	130	Iodi et Potassii Iodidi, Liquor	173
— Pilula, et Ferri	219	— Tinctura	293
— Precipitatum Album	136	— Unguentum	305
— Subchloridum	132	— — compositum	305
— Sulphas	134	Iodine	142
— Unguentum	302	— Compound Liniment of	153
— — compositum	303	— — Ointment of	305
Hydrargyrum	135	— Ointment of	305
— Ammoniatum	136	— — Compound	305
— cum Creta	137	— Solution of, and Iodide of	
Hydrate Acetic	5	Potassium	173
Hydrochloric Acid	12	— Tincture of	293
— — Dilute	13	Iodoform	141
— Solution of Arsenic	166	Iodoformum	141
Hydrocyanic Acid, Diluted	14	Iodol	142
Hydrogen-Potassium Carbon-		Iodum	142
ate	220	Ipecacuanha	145
Hyoscyami Folia	138	— Compound Powder of	244
Hyoscyamus Leaves	138	— Powder of, Compound	244
		Ipecacuanhæ compositus,	
		Pulvis	244
		Iron	121
INFUSION of Calumba	139	— Iodide of	117
— of Catechu	139	— Mercurial Pill with	219
— Compound, of Gentian	140	— Moist Peroxide of	118
— of Ergot	140	— — Hydrated Peroxide of	118
— of Kousso	140	— Perchloride of, Strong So-	
— of Linseed	141	lution of	170
— of Tobacco	141	— — Tincture of	292
— of Yellow Cinchona	139	— Pill of Mercury and	219
Infusum Calumbæ	139	— Protosulphate of	119
— Catechu	139	— Saccharated Carbonate	
— Cinchonæ Flavæ	139	of	115
— Cusso	140	— Solution of Perchloride of,	
— Ergotæ	140	Strong	170
— Gentianæ compositum	140	— Strong Solution of Perchlo-	
— Lini	141	ride of	170
— Tabaci	141	— Sulphate of	119
Iodi, compositum, Linimen-		— — Compound Powder of	244
tum	153	— — Mass of	194
— Linimentum	153	— Tincture of Perchloride of	292
— Liquor	173	— Wine of	118

	PAGE		PAGE
JALAP	145	Lime, Solution of Chlorinated	64
Jalapa	145	— Liniment of	150
James's Powder, Factitious .	244	— Slaked	61
Jecoris Aselli, Oleum	211	— Solution of	167
Juice, Buckthorn	278	— — Chloride of	64
Juniper, Oil of	209	— — Chlorinated	64
Juniperi, Oleum	209	— — Hypochlorite of	64
		— Water	167
		Lini Farina	147
KAMALA	146	— Infusum	141
Kamela	146	— Oleum	209
Kino	147	— Semina	148
Kousso	102	Liniment of Aconite	148
		— of Ammonia	149
		— — Compound	149
		— Camphor	150
LARD, Hog's	29	— — Compound	150
— Prepared	29	— of Cantharides	151
Laudanum	294	— of Carbolic Acid	148
Lead Acetate	223	— Compound, of Ammonia .	149
— Acetate of	223	— — of Soap	155
— Diacetate, Solution of .	174	— — of Turpentine	155
— Diluted Solution of Sub-		— Compound, of Iodine . . .	153
acetate of	175	— of Creasote, Compound .	151
— Liniment of Subacetate of	155	— of Croton Oil	152
— Monoxide	224	— of Iodine, Compound . . .	153
— Protoxide of	224	— of Lime	150
— Subacetate of, Liniment of	155	— of Opium	154
— — Solution of	174	— of Subacetate of Copper .	152
— Subacetate of, Diluted So-		— — Lead	155
lution of	175	— of Sulphate of Copper . .	153
— Solution of Diacetate of .	174	— of Tar	154
— — — Diluted	175	— — Compound	154
— — — and Opium	176	— of Turpentine	155
— Sugar of	223	— of Verdigris	152
Leaf Tobacco	282	Linimentum Acidi Carbolici .	148
Light Magnesia	186	— Aconiti	148
— Carbonate of Magnesia .	186	— Æruginis	152
Lime	63	— Ammonia	149
— Carbonate of	93	— — compositum	149
— Chloride of	64	— Calcis	150
— — Solution of	168	— Camphoræ	150

	PAGE		PAGE
Linimentum Camphoræ com-		Liquor Cupri Sulphatis com-	
positum	150	positus	170
— Cantharidis	151	— Ferri Perchloridi Fortior .	170
— Creasoti compositum	151	— — Persulphatis	171
— Crotonis	152	— Hydrargyri Perchloridi .	172
— Cupri Subacetatis	152	— Iodi	173
— — Sulphatis	153	— et Potassii Iodidi	173
— Hydrargyri compositum	153	— Morphinæ Acetatis	173
— Iodi	153	— — Hydrochloratis	174
— — compositum	153	— Plumbi Diacetatis	174
— Olei Picis Liquidæ compo-		— — Subacetatis	174
situm	154	— — — dilutus	175
— Opii	154	— — — et Opii	176
— Picis Liquidæ	154	— Potassæ	176
— Plumbi Subacetatis	155	— — Arsenitis	164
— Saponis compositum	155	— — Nitratis	178
— Terebinthinæ	155	— — Permanganatis	178
— — compositum	156	— Potassæ Sulphuratae	178
Linseed	148	— Sodæ	179
— Infusion of	141	— Strychniæ	180
— Meal	147	— Zinci Chloridi	181
— Oil	209	— Zinci Sulphatis	182
— Poultice	75	Litharge	224
— Tea	141	Liver of Sulphur	227
Liquor Acidi Carbolici	156	Lotio Acidi Carbolici	183
— Aloes	157	— Ammoniae Acetatis	183
— — et Sodæ Carbonatis	157	— Ammonii Chloridi et Cam-	
— Aluminis	158	phoræ	184
— Ammoniae	158	— — — et Potassæ Nitratis .	184
— — Acetatis	159	— Discutiens	184
— — Fortior	161	— Hydrargyri Flava	185
— Antimonii Chloridi	162	— — Nigra	185
— — Tartarati	163	— Refrigerans	184
— Argenti Nitratis	164	Lotion of Acetate of Ammonia	183
— Arsenicalis	164	— Black Mercurial	185
— Arsenici Chloridi	166	— of Carbolic Acid	183
— — Hydrochloricus	166	— of Chloride of Ammonium	
— Calcis	167	and Camphor	184
— — Chloratae	168	— — — and Nitrate of Potash	184
— Cantharidis Terebinthin-		— Cooling	184
atus	169	— Discutient	184
— Cupri Sulphatis	169	— Yellow Mercurial	185

	PAGE		PAGE
Lugol's Solution	173	Mass of Sulphate of Iron . .	194
Lunar Caustic	51	— of Tartarated Antimony,	
Lytta Vesicatoria	68	Compound	191
		— Tonic	194
MAGNESIA	185	Massa Aloes	190
— Alba	186	— — composita	191
— Carbonate of	186	— Antimonii Tartarati com-	
— Levis	185	posita	191
— Light	185	— Belladonnæ composita . .	192
— — Carbonate of	186	— Catechu composita . .	192
— Sulphate of	188	— Communis	192
— — Enema	106	— Cupri Sulphatis	193
— Usta	185	— Digitalis composita . .	193
Magnesiæ Carbonas	186	— Ferri Sulphatis	194
— — Levis	185	— Resinæ composita	194
— Sulphas	188	— Zingiberis composita . .	195
— Sulphatis, Enema	106	Mercurial Lotion, Black . .	185
Magnesium Oxide	185	— — Yellow	185
— Sulphate	188	— Pill with Iron	219
Manganese, Black Oxide of . .	189	Mercuric-ammonium Chloride .	136
— Binoxide of	189	Mercuric Chloride	130
— Peroxide of	189	Mercuric Iodide	127
— Dioxide	189	— Oxide	128
Manganesii Oxidum Nigrum . .	189	— Sulphate	134
Manganic Peroxide	189	Mercurous Chloride	132
Marble, White	190	Mercury	135
Marmor Album	190	— Amidochloride of	136
Mass of Aloes	190	— Ammoniated	136
— — Compound	191	— Ball	130
— Alterative	191	— Bichloride of	130
— Astringent	192	— Biniodide of	127
— of Belladonna, Compound . .	192	— — Ointment of	303
— of Catechu, Compound . .	192	— with Chalk	137
— Cathartic	190	— Compound Ointment of . .	303
— Common	192	— — Spirit of Perchloride of .	273
— Cordial	195	— Nitrate of, Ointment of . .	304
— of Digitalis, Compound . .	193	— Ointment of	302
— Diuretic	194	— — Compound	303
— of Ginger, Compound . .	195	— — of Biniodide of	303
— of Resin, Compound . .	194	— — of Nitrate of	304
— of Sulphate of Copper . .	193	— — of Perchloride of . .	304
		— — Red Iodide of	303

	PAGE		PAGE
Mercury, Perchloride of	130	Mylabris	202
— — Compound Spirit of	273	— Cichorii	202
— — Ointment of	304	— Ointment of	305
— — Solution of	172	Myrrh	202
— Persulphate of	134	— Tincture of	293
— Pill of, and Iron	219	Myrrha	202
— Protochloride of	132	Myrrhæ, Tinctura	293
— Red Iodide of	127		
— — Ointment of	303	NITRE	236
— — Oxide of	128	— Sweet Spirits of	269
— Spirit of Perchloride of,		Nitric Acid	16
Compound	273	— — Diluted	17
— Subchloride of	132	Nitro-hydrochloric Acid, Diluted	18
— Sulphate of	134	Nitrous Ether, Spirit of	269
Methylated Spirit	273	Nucis Vomicae, Tinctura	294
Mindererus's Spirit	159	Nut-galls	123
Mistura Ammoniaë	195	Nux Vomica	203
— Creasoti	195	— — Tincture of	294
— Cretæ composita	196		
— Ricini	196	OAK-BARK	247
Mixture, Ammonia	195	Oak-galls	123
— Castor Oil	196	Oil of Anise	43, 204
— Chalk, Compound	196	— Cajuput	205
— Compound Chalk	196	— Camphorated	206
— Gregory's	245	— of Cantharides	206
Moist Peroxide of Iron	118	— Carron	150
— Hydrated Peroxide of Iron	118	— Castor	213
— Ferric Hydrate	118	— — Mixture	196
Molasses	284	— Cod-liver	211
Monkshood	27, 28	— Croton	207
Morphiæ Acetas	197	— — Liniment of	152
— Hydrochloras	197	— Eucalyptus	208
Morphina	197	— of Juniper	209
Morrhua, Oleum	211	— Linseed	209
Mucilage of Gum Acacia	201	— Olive	212
— of Starch	201	— Palm	212
Mucilago Acaciæ	201	— Peppermint	210
— Amyli	201	— of Turpentine	214
Muriatic Acid	12	— of Vitriol	21
Mustard	257	— White	150
— Poultice	76		
Mylabridis, Unguentum	305		

	PAGE		PAGE
Ointment of Aconite . . .	297	Oleum Cajuput . . .	205
— Alum, Compound . . .	298	— Camphoratum . . .	206
— of Ammoniated Mercury .	302	— Cantharidis . . .	206
— Arsenic . . .	299	— Caryophylli . . .	206
— Black Hellebore . . .	301	— Crotonis . . .	207
— Biniiodide of Mercury .	302	— Eucalypti . . .	208
— Blistering . . .	299, 302	— Jecoris Aselli . . .	211
— Blue . . .	302	— Juniperi . . .	209
— of Cantharides . . .	299	— Lini . . .	209
— of Carbolic Acid . . .	296	— Menthæ Piperitæ . .	210
— — — Compound . . .	297	— Morrhuæ . . .	211
— of Carbonate of Zinc . .	309	— Olivæ . . .	212
— of Chlorinated Lime . .	299	— Palmæ . . .	212
— Citrine . . .	304	— Ricini . . .	213
— of Creasote . . .	300	— Terebinthinæ . . .	214
— Digestive . . .	306, 309	Olive Oil . . .	212
— of Galls . . .	301	Opii compositus, Pulvis .	245
— — and Opium . . .	301	— Enema . . .	107
— Healing . . .	309	— Linimentum . . .	154
— of Iodide of Sulphur . .	308	— Liquor Plumbi Subaceta-	
— of Iodine . . .	305	tis et . . .	176
— — Compound . . .	305	— Tinctura . . .	294
— of Mercury . . .	302	Opium . . .	215
— — Compound . . .	303	— Compound Powder of .	245
— of Mylabris . . .	305	— Enema of . . .	107
— of Nitrate of Mercury .	304	— Liniment of . . .	154
— — of Silver . . .	298	— Powder of, Compound .	245
— of Perchloride of Mercury	304	— Solution of Subacetate of	
— of Red Iodide of Mercury	302	Lead and . . .	176
— of Resin . . .	306	— Tincture of . . .	294
— of Savin . . .	307	Opodeldoc . . .	155
— Sheep . . .	308		
— of Subacetate of Copper .	300		
— of Sulphur, Compound .	308		
— of Sulphurated Potash .	306	PALM Butter . . .	212
— of Tannic Acid . . .	297	— Oil . . .	212
— of Tar . . .	306	Palmæ, Oleum . . .	212
— of Tartarated Antimony .	297	Papaveris, Syrupus . .	281
— of Turpentine . . .	309	Paraffin . . .	217
— of Verdigris . . .	300	Paraffinum . . .	217
— of White Hellebore . .	309	Pepper, Black . . .	220
Oleum Anisi . . .	43, 204	Peru, Balsam of . . .	55

	PAGE		PAGE
Phenacetine	218	Potash, Hydrate of	226
Phenic Acid	8	— Lotion of Chloride of Am-	
Phenol	8	monium and Nitrate	
Phenylic Alcohol	8	of	184
Physostigmatis extractum	328	— Nitrate of	236
— semen	218	— Permanganate of	238
Physostigmina	327	— Solution of	176
Picis, Emplastrum	105	— Sulphate of	241
— Liquidæ, Unguentum	306	— Sulphurated	227
Pill, Blue, Ferruginate	219	— — Ointment of	306
— Ferruginate Blue	219	— Yellow Prussiate of	240
— Mercurial, with Iron	219	Potassa Caustica	226
— of Mercury and Iron	219	— Fusa	226
Pilula Hydrargyri et Ferri	219	— Sulphurata	227
Pimenta	219	Potassæ Arsenitis, Liquor	164
Pimentæ, Tinctura	286	— Bicarbonas	229
Pimento	219	— Carbonas	231
— Tincture of	296	— Chloras	234
Piper Nigrum	220	— Hydras	226
Pitch Plaster	105	— Liquor	176
— Black	223	— Nitras	236
Pitch, Burgundy	221	— Nitratis, Lotio Ammonii	
Pix Burgundica	221	Chloridi et	184
— Liquida	222	— Permanganas	238
— Nigra	223	— Prussia Flava	240
Plaster, Pitch	105	— Sulphas	241
Plumbi Acetas	223	— Sulphurata, Liquor	178
— Diacetatis, Liquor	174	— — Unguentum	306
— Oxidum	224	Potassii Bromidum	232
— Subacetatis Linimen-		— Iodidum	241
tum	155	Potassium Alum	35
— — Liquor	174	— Aluminum Sulphate	35
— — dilutus, Liquor	175	— Antimony Tartrate	46
— et Opii, Liquor	176	— Bromide	232
Podophylli Rhizoma	225	— Carbonate	231
— resina	226	— Chlorate	234
Poppies, Syrup of	281	— Ferrocyanide	240
Potash, Bicarbonate of	229	— Hydrate	236
— Bromide	232	— Hydrogen Carbonate	229
— Carbonate of	231	— Iodide	241
— Caustic	226	— Nitrate	236
— Chlorate of	234	— Permanganate	238

	PAGE		PAGE
Potassium, Solution of Iodine		QUERCUS Cortex	247
and Iodide of	173	Quicklime	63
— Sulphate	241	Quicksilver	135
— Sulphide of	227	Quinia Sulphate	248
Potassii Iodidi, Liquor Iodi et	173	— Sulphate of	248
— Iodidum	241	Quiniæ Sulphas	248
Poultice, Ammoniated Mus-		Quinine, Disulphate of .	248
tard	76	— Sulphate of	248
— Bran	75		
— Charcoal	74	RECTIFIED Spirit	274
— Chlorinated Lime	74	Red Iodide of Mercury . .	127
— Linseed	75	— Oxide of Mercury . .	128
— Mustard	76	— Precipitate	128
— Yeast	75	— Rose Petals	252
Powder, Antimonial	244	— Sanders Wood, Tincture of	295
Powder of Carbonate of Zinc,		— Refined Sugar	253
Compound	246	Resin	250
— Dover's Factitious	244	— Compound Mass of . .	194
— Gregory's	245	— Ointment of	306
— Grey	137	— Yellow	250
— of Ipecacuanha, Compound	244	Resina	250
— James's Factitious	244	Resinæ composita, Massa .	194
— of Opium, Compound . . .	245	— Unguentum	306
— of Rhubarb, Compound . .	245	Rhamni Succus	278
— of Sulphate of Iron, Com-		— Syrupus	282
pound	244	Rhei compositus, Pulvis .	245
Prepared Chalk	93	— Radix	251
— Lard	29	Rhubarb, Compound Powder	
— Suet	257	of	245
Proof Spirit	275	— Powder of, Compound .	245
Protochloride of Mercury . .	132	— Root	251
Prussic Acid	14	Rosæ Gallicæ Petala . . .	252
— Scheele's	14	Rose Petals, Red	252
Pulvis Antimonialis	244	Roses, Confection of . . .	92
— Ferri Sulphatis compositus	244	Rye, Ergot of	107
— Ipecacuanhæ compositus .	244		
— Opii compositus	245	SABINÆ Cacumina	253
— Rhei compositus	245	— Unguentum	302
— Zinci Carbonatis compositus		Saccharated Carbonate of Iron	115
tus	246	Sacchari Fæx	284
Pyroligneous Acid	4		
Pyroxylin	246		

	PAGE		PAGE
Saccharum Purificatum	253	Soda, Hydrate of	258
Saffron	95	— Hyposulphite of	263
Sal Ammoniac	40	— Solution of	179
Salt, Common	266	— Sulphate of	264
— Glauber's	264	— Sulphite of	265
— Spirit of	12	— Washing	261
Saltpetre	236	Sodæ Acetas	259
Sanders Wood, Tincture of		— Bicarbonas	260
Red	295	— Carbonas	261
Santalidis Rubra, Tinctura	295	— — Exsiccata	263
Santoninum	254	— Carbonatis, Liquor Aloes	
Sapo Durus	254	et	157
— Mollis	256	— Hyposulphis	263
Saponis compositum, Lini-		— Liquor	179
mentum	155	— Sulphas	264
Savin, Ointment of	307	— Sulphis	265
— Tops	253	Sodii Chloridum	266
Scheele's Prussic Acid	14	Sodium Acetate	259
Scillæ	256	— Biborate of	58
Sevum Præparatum	257	— Carbonate	261
Sheep and Calves' Cordial	196	— Chloride	266
— Ointment	308	— Chloride of	266
— Dip or Wash (Arsenical)	165	— Diborate	58
Silver, Nitrate of	51	— Hydrate	258
— — Ointment of	298	— Hydrate of	258
— — Solution of	164	— Hydrogen Carbonate	260
— Ointment of Nitrate of	298	— Hyposulphite	263
— Solution of Nitrate of	164	— Sulphate	264
Sinapis	257	— Sulphite	265
Slaked Lime	61	Soft Soap	256
Soap, Compound Liniment of	155	Solution of Acetate of Am-	
— Hard	254	monia	159
— Soft	256	— of Aloes	157
Soda, Acetate of	259	— — and Carbonate of Soda	157
— Biborate	58	— of Alum	158
— Bicarbonate of	260	— of Ammonia	158
— Carbonate of	261	— — Strong	161
— — Dried	263	— of Antimonious Chloride	162
— — Solution of Aloes and	157	— Arsenical	164
— Caustic	258	— of Arsenic, Hydrochloric	166
— Caustica	258	— of Cantharides, Terebin-	
— Dried Carbonate of	263	thinated	169

	PAGE		PAGE
Solution of Carbolic Acid	156	Spirit of Perchloride of Mer-	
— of Chloride of Antimony	162	cury, Compound	273
— — of Lime	168	— Proof	275
— — of Zinc	181	— Rectified	274
— of Chlorinated Lime	168	— of Salt	12
— of Diacetate of Lead	174	— of Wine	274
— Fowler's	164	Spirits of Nitre, Sweet	269
— of Hypochlorite of Lime	168	— of Turpentine	214
— of Iodine and Iodide of		Spiritus Ætheris	268
Potassium	173	— — Nitrici	269
— of Lime	167	— — Nitrosi	269
— Lugol's	173	— Ætheris Sulphurici	268
— of Permanganate of Potash	178	— Ammoniaë Aromaticus	271
— of Nitrate of Potash	178	— — Fœtidus	271
— — of Silver	164	— Camphoræ	272
— of Perchloride of Iron,		— Chloroformi	272
Strong	170	— Hydrargyri Perchloridi	
— of Perchloride of Mer-		compositus	273
cury	172	— Methylatus	273
— of Persulphate of Iron	171	— Rectificatus	274
— of Potash	176	— Tenuior	275
— of Soda	179	Staphisagriaë Semina	276
— of Subacetate of Lead	174	Starch	41
— — — Diluted	175	— Mucilage of	201
— — — and Opium	176	Stavesacre Seeds	276
— of Sulphate of Copper	169	Strong Solution of Ammonia	161
— — Compound	176	Strychnine	276
— of Strychnia	180	— Solution of	180
— of Sulphurated Potash	178	Strychniaë Liquor	180
— of Terchloride of Anti-		Sublimed Sulphur	278
mony	162	Succus Rhamni	208
— of Tartarated Antimony	163	Suet, Prepared	257
— Sulphate of Zinc	182	Sugar of Lead	223
Spanish Fly	68	Sulphur, Compound Ointment	
Spirit of Ammonia, Aromatic	271	of	308
— — Fetid	271	— Flowers of	278
— of Camphor	272	— Iodide of	280
— of Chloroform	272	— — Ointment of	308
— of Ether	268	— Liver of	306
— Methylated	273	— Ointment of, Compound	308
— Mindererus's	159	— — of Iodide of	280
— of Nitrous Ether	269	— Sublimatum	278

	PAGE		PAGE
Sulphur, Sublimed	280	Terebinthinæ Unguentum	309
Sulphurated Potash	227	Terebinthinated Solution of	
— — Ointment of	306	Cantharides	169
Sulphuric Acid	21	Terra Japonica	77
— — Diluted	23	Theriaca	284
— Ether	30	Thymol	284
Sulphuris Iodidi, Unguentum	308	Tinctura Aconiti	285
— Iodi	305	— Aloes composita	286
— Unguentum compositum	308	— Arnicæ	286
Sulphurous Acid	24	— Camphoræ	287
— Anhydride	24	— Cannabis Indicæ	287
Sweet Spirits of Nitre	269	— Cantharidis	287
Syrup of Buckthorn	282	— — Fortior	288
— of Chloral Hydrate	281	— Cardamomi composita	288
— of Poppies	281	— Chloroformi composita	289
Syrupus Chloralis Hydratis	281	— Cinchonæ Flavæ	289
— Papaveris	281	— Colchici Seminis	290
— Rhamni	282	— Conii	290
		— — Fructus	290
TABACI, Decoctum	103	— Crotonis	291
— Enema	107	— Ergotæ	291
— Folia	282	— Ferri Perchloridi	292
— Infusum	141	— — Sesquichloridi	292
Tannic Acid	25	— Iodi	293
— — Ointment of	297	— Myrrhæ	293
Tannin	25	— Nucis Vomicae	294
Tar	222	— Opii	294
— Ointment of	306	— Pimentæ	295
— Wood	222	— Santalis Rubra	295
Tartar Emetic	46	— Zingiberis	296
Tartarated Antimony	46	Tincture of Aconite	285
— — Compound Mass of	191	— of Aloes, Compound	286
— — Solution of	163	— of Arnica	286
Tartarised Antimony	46	— of Cantharides	287
Taraxacum	283	— — Strong	288
Terebinthina Canadensis	283	— of Cardamoms, Compound	288
Terebinthinæ compositum, Li-		— of Chloroform, Compound	289
nimentum	156	— of Colchicum Seeds	290
— Enema	107	— of Croton	291
— Linimentum	155	— of Ergot	291
— Oleum	214	— of Ginger	296
		— of Hemlock	290

	PAGE		PAGE
Tincture of Iodine	293	Unguentum Hydargyri Bin-	
— of Myrrh	293	iodidi	303
— of Nux Vomica	294	— — compositum	303
— of Opium	294	— — Iodidi Rubri	303
— of Perchloride of Iron	292	— — Nitratis	304
— of Pimento	295	— — Perchloridi	304
— of Red Sanders Wood	295	— Iodi	305
— of Yellow Cinchona	289	— — compositum	305
Tobacco, Decoction of	103	— Mylabridis	305
— Enema of	107	— Picis Liquidæ	306
— Infusion of	141	— Potassæ Sulphuratæ	306
— Leaf	282	— Resinæ	306
Tonic Mass	194	— Sabinæ	307
Treacle	284	— Sulphuris	307
Trichloraldehyd	80	— — compositum	308
Turpentine, Compound Lini-		— — Iodidi	308
ment of	156	— Terebinthinæ	309
— Enema of	107	— Veratri	309
— Liniment of	155	— Zinci Carbonatis	309
— — Compound	156	— — Oxidi	310
— Oil of	214	VAPOR Chlorig	310
— Ointment of	309	Vaselin	312
— Spirits of	214	Vaseline	312
UNGUENTUM Acidi Carbolici	296	Veratri, Unguentum	309
— — — compositum	297	Veratrum	312
— — Tannici	297	Verdigris	99
— Aconiti	297	— Liniment of	152
— Æruginis	300	— Ointment of	300
— Aluminis compositum	298	Vinegar	2
— Antimonii Tartarati	298	— of Cantharides	3
— Argenti Nitratis	298	Vinum Ferri	313
— Arsenici	299	Vitriol, Blue	100
— Calcis Chlorati	299	— Green	119
— Cantharidis	299	— Oil of	21
— Creasoti	300	— White	320
— Cupri Subacetatis	300	WASH, Sheep	165
— Gallæ	301	Washing Soda	261
— et Opii	301	Water, Distilled	49
— Hellebori Nigri	301	Wax, Bees', Unbleached	78
— Hydrargyri	302	— Yellow	78
— — Ammoniati	302		

	PAGE		PAGE
White Hellebore	312	Zinc Carbonate, Ointment of	309
— — Ointment of	309	— Chloride	317
— Marble	190	— Chloride of	317
— Oil	150	— — Solution of	181
— Precipitate	136	— Compound Powder of Car-	
— Vitriol	320	bonate of	246
Whiting	93	— Granulated	322
Wine of Iron	313	— Ointment of Carbonate	
— Spirit of	274	of	309
Wood Charcoal	70	— Oxide	319
— Tar	222	— Oxide of	319
		— Solution of Chloride of . .	81
YEAST	79	— Sulphate	320
— Poultice	75	— Sulphate of	320
Yellow Mercurial Lotion . .	185	Zinci Acetas	314
— Resin	250	— Carbonas	316
— Wash	185	— Carbonatis, Pulvis, compo-	
		situs	246
ZINC Acetate	314	— Carbonatis, Unguentum .	309
— Acetate of	314	— Chloridi, Liquor	181
— Butter of	317	— Chloridum	317
— Carbonate	316	— Oxidum	319
— Carbonate of	316	— Sulphas	320
— Carbonate, Compound		Zincum Granulatum	322
Powder of	246	Zingiber	323
		Zingiberis composita, Massa .	195
		— Tinctura	296

COOLEY'S CYCLOPÆDIA OF PRACTICAL RECEIPTS AND COLLATERAL INFORMATION

*IN THE ARTS, MANUFACTURES, PROFESSIONS,
AND TRADES,*

INCLUDING

MEDICINE, PHARMACY, HYGIENE, AND DOMESTIC
ECONOMY,

DESIGNED AS A COMPREHENSIVE SUPPLEMENT TO THE
PHARMACOPŒIA AND GENERAL BOOK OF REFERENCE.

Edited by W. NORTH, M.A., F.C.S.

With 370 Illustrations. 2 vols. royal 8vo. 42s.

THE acknowledged usefulness of this work is proved by its having now reached its seventh edition.

In the preparation of this edition, the object kept in view has been the revision and amplification of the matter originally contained in the work, rather than the enlargement of its scope.

The more important additions to the work will be found under Photography, Surveying, and Insects Injurious to Crops. The old article on Photography has been entirely re-written and greatly enlarged, and a collection of formulæ and processes, from the very best authorities of the present day, has been included in it, which it is hoped will prove useful, not only to the amateur, but also to the professional.

Surveying appeared to demand a place in a book which is largely used by country gentlemen and colonists, and though the methods given are chiefly those used for military purposes, practical experience of the value of these methods for purely civil purposes,

[TURN OVER.]

has led the editor to deal with the subject from this point of view at some length.

The series of articles on Insects Injurious to Crops are practically a reprint of reports made by Mr. Chas. Whitehead, F.Z.S., to the Agricultural Department of the Privy Council and to the Board of Agriculture. These articles are reproduced by special permission of the Board.

The revision of the Pharmacy, one of the most important divisions of the work, has been carried out very thoroughly by Mr. A. W. Gerrard, Pharmacist to University College Hospital. Great additions have been made, and nothing has been removed in any way likely to be of practical utility. The character of 'Cooley's Cyclopædia' as a pharmaceutical reference-book has always been considerable, and no effort has been spared to keep it well up to the mark in this respect.

The general Chemistry has been most thoroughly revised by various hands, and brought well up to date. Special attention has been paid throughout to all commercial and practically useful methods and processes, whilst at the same time the scientific aspect of the various branches of the subject has not been neglected. The work was begun by Dr. G. McGowan, of Bangor, carried on by Messrs. W. K. Tomkins, B.Sc., E. P. Perman, B.Sc., and C. F. Baker, B.Sc., of University College, London, and completed by Mr. J. T. Norman.

The Veterinary Medicine has been considerably modified and brought more into harmony with modern practice. The Domestic Medicine has been advisedly reduced, and detailed accounts of many diseases and modes of treatment have been removed in all cases in which the supervision of a qualified medical practitioner is necessary or desirable. The gaps thus caused have been more than filled by practical information on first aid to the sick and injured.

The general receipts throughout the work have been carefully revised and largely added to, and it is believed much increased in practical value.

London: J. & A. CHURCHILL, 11 New Burlington Street.

No. 2.

London, 7, Great Marlborough Street,
November, 1901.

A S E L E C T I O N
FROM
J. & A. CHURCHILL'S CATALOGUE,
COMPRISING
MOST OF THE RECENT WORKS PUBLISHED BY THEM.

N.B.—J. & A. Churchill's larger Catalogue, which contains over 600 works with a Complete Index to their Subjects, will be sent on application.

Human Anatomy : a Treatise by various Authors. Edited by HENRY MORRIS, M.A., M.B. Lond., F.R.C.S., Surgeon to, and Lecturer on Surgery at, the Middlesex Hospital. Second Edition. Roy. 8vo, with 790 Illustrations, nearly all original, and many of them in several colours, 36s.

Dissection Outlines for Use with Morris's Anatomy. By WILLIAM A. CAMPBELL, B.S., M.D. Second Edition. 8vo, 3s.

Heath's Practical Anatomy : a Manual of Dissections. Eighth Edition. Edited by WILLIAM ANDERSON, F.R.C.S., Surgeon and Lecturer on Anatomy at St. Thomas's Hospital. Crown 8vo, with 329 Engravings. 15s.

Wilson's Anatomist's Vade-Mecum. Eleventh Edition, by HENRY E. CLARK, M.R.C.S. Eng., F.F.P.S. Glasg., Examiner in Anatomy F.P.S. Crown 8vo, with 492 Engravings and 26 Coloured Plates, 18s.

An Atlas of Human Anatomy. By RICKMAN J. GODLEE, M.S., F.R.C.S., Surgeon and late Demonstrator of Anatomy, University College Hospital. With 48 Imp. 4to Plates (112 figures), and a volume of Explanatory Text. 8vo, £4 14s. 6d.

Human Osteology. By LUTHER HOLDEN. Eighth Edition, edited by CHARLES STEWART, F.R.S., Conservator of the Museum R.C.S., and ROBERT W. REID, M.D., F.R.C.S., Regius Professor of Anatomy in the University of Aberdeen. 8vo, with 59 Lithographic Plates and 74 Engravings, 16s.

By the same Author.

Landmarks, Medical and Surgical. Fourth Edition. 8vo, 3s. 6d.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

A Manual of Practical Anatomy. By the late Professor ALFRED W. HUGHES, M.B., M.C. Edin., Professor of Anatomy, King's College, London, Edited and completed by ARTHUR KEITH, M.D., Lecturer on Anatomy, London Hospital Medical College. In three parts, royal 8vo. Part I., with 38 Coloured Plates, and 116 Figures in the Text, 10s. 6d.

The Essentials of Regional Anatomy. By RICHARD J. BERRY, M.D., F.R.S., F.R.C.S., (Edin.), Lecturer on Anatomy in the New School of Medicine of the Royal Colleges, Edinburgh. Post 8vo, interleaved, 10s. net.

Anatomy of the Joints of Man. By Henry MORRIS, Senior Surgeon to the Middlesex Hospital. With 44 Lithographic Plates (several coloured). 8vo, 16s.

A Manual of General Pathology, for Students and Practitioners. By W. S. LAZARUS-BARLOW, B.A., M.D., Pathologist and Lecturer on Pathology, Westminster Hospital. 8vo, 21s.

Pathological Anatomy of Diseases. Arranged according to the nomenclature of the R.C.P. Lond. By NORMAN MOORE, M.D., F.R.C.P., Assistant Physician and Lecturer on Pathological Anatomy to St. Bartholomew's Hospital. Fcap. 8vo, with 111 Engravings, 8s. 6d.

A Manual of Clinical and Practical Pathology. By W. E. WYNTER, M.D., M.R.C.P., Assistant Physician to the Middlesex Hospital, and F. J. WETHERED, M.D., F.R.C.P., Assistant Physician to the Consumption Hospital, Brompton. With 4 Coloured Plates and 67 Engravings. 8vo, 12s. 6d.

General Pathology (an Introduction to). By JOHN BLAND SUTTON, F.R.C.S., Assistant Surgeon to, and Lecturer on Anatomy at, Middlesex Hospital. 8vo, with 149 Engravings, 14s.

The Pathologist's Handbook: a Manual for the Post-mortem Room. By T. N. KELYNACK, M.D., late Demonstrator in Morbid Anatomy, Owens College, Manchester. With 126 Illustrations, fcap. 8vo, pegamoid, 4s. 6d.

Selected Researches in Pathology. By A. G. AULD, M.D., M.R.C.P. With 14 Illustrations, 8vo, 6s.

The Human Brain: Histological and Coarse Methods of Research. By W. BEVAN LEWIS, L.R.C.P. Lond., Medical Superintendent, West Riding Lunatic Asylum. 8vo, with Wood Engravings and Photographs, 8s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

The Physiology and the Pathology of the

Cerebral Circulation: an Experimental Research. By LEONARD HILL, M.D., Hunterian Professor, R.C.S. With 41 Illustrations. Royal 8vo, 12s.

Elements of Human Physiology. By Ernest

H. STARLING, M.D., F.R.C.P., F.R.S., Jodrell Professor of Physiology in University College, London. Fourth Edition, much enlarged, with 317 Illustrations 8vo, 12s. 6d.

Manual of Physiology: for the Use of Junior

Students of Medicine. By GERALD F. YEO, M.D., F.R.S. Third Edition. Crown 8vo, with 254 Engravings and Plate of Spectra, 14s.

Elementary Practical Physiology, including

Histology, Chemical and Experimental Physiology. By DE BURGH BIRCH, M.D., F.R.S.E., Professor of Physiology in the Yorkshire College of the Victoria University. With 62 Engravings. Crown 8vo, 6s. 6d.

Practical Lessons in Elementary Biology, for

Junior Students. By PEYTON T. B. BEALE, F.R.C.S., Lecturer on Elementary Biology and Demonstrator in Physiology in King's College, London. Crown 8vo, 3s. 6d.

Medical Jurisprudence: its Principles and

Practice. By ALFRED S. TAYLOR, M.D., F.R.C.P., F.R.S. Fourth Edition, by THOMAS STEVENSON, M.D., F.R.C.P., Lecturer on Medical Jurisprudence at Guy's Hospital. 2 vols. 8vo, with 189 Engravings, 31s. 6d.

Lectures on Medical Jurisprudence and Toxi-

cology. By FRED. J. SMITH, M.D., F.R.C.P., Lecturer on Forensic Medicine and Toxicology at the London Hospital. Crown 8vo, 7s. 6d.

The Theory and Practice of Hygiene. (Notter

and Firth). By J. LANE NOTTER, M.D., Examiner in Hygiene and Public Health in the University of Cambridge and the English Conjoint Board, Professor of Hygiene in the Army Medical School; and W. H. HORROCKS, M.B., B.Sc., Assistant Professor of Hygiene in the Army Medical School. Second Edition. With 15 Plates and 134 other Illustrations. Royal 8vo, 25s.

A Manual of Practical Hygiene. By the late

E. A. PARKES, M.D., F.R.S. Eighth Edition, by J. LANE NOTTER, A.M., M.D., Professor of Hygiene in the Army Medical School. 8vo, with 10 Plates and 103 Engravings, 18s.

A Handbook of Hygiene and Sanitary Science.

By GEO. WILSON, M.A., M.D., LL.D., D.P.H. Camb. Medical Officer of Health for Mid-Warwickshire. Eighth Edition. Post 8vo, with Engravings, 12s. 6d.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Hygiene and Public Health: a Treatise by various Authors. Edited by THOMAS STEVENSON, M.D., and SHIRLEY F. MURPHY. In 3 vols., royal 8vo, fully Illustrated. Vol. I., 28s.; Vol. II., 32s.; Vol. III., 20s.

A Simple Method of Water Analysis, especially designed for the use of Medical Officers of Health. By JOHN C. THRESH, M.D.Vic., D.Sc Lond., D.P.H. Camb. Third Edition, enlarged. Fcap. 8vo, 2s. 6d.

Elements of Health: an Introduction to the Study of Hygiene. By LOUIS C. PARKES, M.D., D.P.H. Lond., Medical Officer of Health for Chelsea, Lecturer on Public Health at St. George's Hospital. Post 8vo, with 27 Engravings, 3s. 6d.

Diet and Food considered in relation to Strength and Power of Endurance, Training and Athletics. By ALEXANDER HAIG, M.D., F.R.C.P. Third Edition. Crown 8vo, 2s.

The Prevention of Epidemics and the Construction and Management of Isolation Hospitals. By ROGER MCNEILL, M.D. Edin., D.P.H. Camb., Medical Officer of Health for the County of Argyll. 8vo, with several Hospital Plans, 10s. 6d.

Effects of Borax and Boracic Acid on the Human System. By Dr. OSCAR LIEBREICH, Professor in the University of Berlin. With Plates, post 4to, 2s.

A Manual of Bacteriology, Clinical and Applied. With an Appendix on Bacterial Remedies, &c. By RICHARD T. HEWLETT, M.D., Professor of General Pathology and Bacteriology in King's College, London. With 75 Illustrations, post 8vo, 10s. 6d.

An Introduction to the Bacteriological Examination of Water. By W. H. HORROCKS, M.B., B.Sc., Lond., Assistant Professor of Military Hygiene in the Army Medical School, Netley, Major, Royal Army Medical Corps. With 5 Lithographic Plates, 8vo, 10s. 6d.

Hospitals and Asylums of the World: their Origin, History, Construction, Administration, Management, and Legislation. By Sir H. C. BURDETT, K.C.B. 4 vols., royal 8vo, and Portfolio. Complete, £12 12s. net. Vols. I. and II.—Asylums, £6 15s. net. Vols. III. and IV.—Hospitals, with Plans and Portfolio, £9 net.

Mental Diseases: Clinical Lectures. By T. S. CLOUSTON, M.D., F.R.C.P. Edin., Lecturer on Mental Diseases in the University of Edinburgh. Fifth Edition. Cr. 8vo, with 19 Plates, 14s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

The Insane and the Law: a Plain Guide for

Medical Men, Solicitors, and Others as to the Detention and Treatment, Maintenance, Responsibility, and Capacity either to give evidence or make a will of Persons Mentally Afflicted. With Hints to Medical Witnesses and to Cross-Examining Counsel. By G. PITT-LEWIS, Q.C., R. PERCY SMITH, M.D., F.R.C.P. and J. A. HAWKE, B.A., Barrister-at-Law. 8vo, 14s.

A Dictionary of Psychological Medicine, giving

the Definition, Etymology, and Synonyms of the Terms used in Medical Psychology; with the Symptoms, Treatment, and Pathology of Insanity; and THE LAW OF LUNACY IN GREAT BRITAIN AND IRELAND. Edited by D. HACK TUKE, M.D., LL.D., assisted by nearly 130 Contributors, British, Continental and American. 2 vols., 1,500 pages, royal 8vo, Illustrated, 42s.

Mental Physiology, especially in its Relation

to Mental Disorders. By THEO. B. HYSLOP, M.D., Resident Physician and Medical Superintendent at Bethlem Royal Hospital, Lecturer on Mental Diseases in St. Mary's Hospital Medical School. 8vo, 18s.

The Mental Affections of Children: Idiocy,

Imbecility, and Insanity. By WM. W. IRELAND, M.D. Edin., formerly Medical Superintendent of the Scottish Institution for the Education of Imbecile Children. Second Edition. With 21 Plates, 8vo, 14s.

Mental Affections of Childhood and Youth

(Lettsomian Lectures for 1887, etc.). By J. LANGDON-DOWN, M.D., F.R.C.P., Consulting Physician to the London Hospital. 8vo, 6s.

The Journal of Mental Science. Published

Quarterly, by Authority of the Medico-Psychological Association. 8vo, 5s.

Manual of Midwifery, including all that is

likely to be required by Students and Practitioners. By ALFRED L. GALABIN, M.A., M.D., F.R.C.P., Obstetric Physician and Lecturer on Midwifery and Diseases of Women to Guy's Hospital. Fifth Edition. Post 8vo, with 298 Engravings, 15s.

The Practice of Midwifery: a Guide for Prac-

tioners and Students. By D. LLOYD ROBERTS, M.D., F.R.C.P., Lecturer on Clinical Midwifery and Diseases of Women at the Owens College; Consulting Obstetric Physician to the Manchester Royal Infirmary. Fourth Edition. Fcap. 8vo, with Coloured Plates and Wood (226) Engravings, 10s. 6d.

A Short Practice of Midwifery, embodying the

Treatment adopted in the Rotunda Hospital, Dublin. By HENRY JELLETT, M.D., B.A.O. Dub., late Assistant Master, Rotunda Hospital. Third Edition. With 124 Illustrations. Crown 8vo, 8s. 6d.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Obstetric Aphorisms: for the Use of Students

commencing Midwifery Practice. By JOSEPH G. SWAYNE, M.D.,
Lecturer on Midwifery in the Bristol Medical School. Tenth Edition.
Fcap. 8vo, with 20 Engravings. 3s. 6d.

Economics, Anæsthetics, and Antiseptics in

the Practice of Midwifery. By HAYDN BROWN, L.R.C.P., L.R.C.S.
Edin. Fcap. 8vo, 2s. 6d.

Lectures on Obstetric Operations: including

the Treatment of Hæmorrhage, and forming a Guide to the Manage-
ment of Difficult Labour. By ROBERT BARNES, M.D., F.R.C.P.,
Consulting Obstetric Physician to St. George's Hospital. Fourth
Edition. 8vo, with 121 Engravings, 12s. 6d.

By the same Author.

A Clinical History of Medical and Surgical

Diseases of Women. Second Edition. 8vo, with 181 Engravings, 28s.

Outlines of Gynæcological Pathology and

Morbid Anatomy. By C. HUBERT ROBERTS, M.D. Lond., Physician
to the Samaritan Free Hospital for Women. With 151 Illustrations,
8vo, 21s.

Diseases of Women. (Student's Guide Series.)

By ALFRED L. GALABIN, M.A., M.D., F.R.C.P., Obstetric Phy-
sician to, and Lecturer on Midwifery and Diseases of Women at,
Guy's Hospital. Fifth Edition. Fcap. 8vo, with 142 Engravings, 8s. 6d.

A Short Practice of Gynæcology. By Henry

JELLETT, M.D., B.A.O. Dub., late Assistant Master, Rotunda
Hospital, Dublin With 125 Illustrations, crown 8vo, 7s. 6d.

Manual of the Diseases peculiar to Women.

By JAMES OLIVER, M.D., F.R.S.E., M.R.C.P., Physician to the
Hospital for Diseases of Women, London. Fcap. 8vo, 3s. 6d.

By the same Author.

Abdominal Tumours and Abdominal Dropsy

in Women. Crown 8vo, 7s. 6d.

Sterility. By ROBERT BELL, M.D., F.F.P. & S. Glasg.,

Senior Physician to the Glasgow Hospital for Diseases peculiar to
Women. 8vo, 5s.

A First Series of Fifty-four Consecutive Ovario-

tomies, with Fifty-three Recoveries. By A. C. BUTLER-SMYTHE,
F.R.C.P. Edin., Surgeon to the Samaritan Free Hospital, Senior
Surgeon to the Grosvenor Hospital for Women and Children. 8vo,
6s. 6d.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

A Manual for Hospital Nurses and others engaged in Attending on the Sick. By E. J. DOMVILLE, Surgeon to the Devon and Exeter Hospital. Eighth Edition. Crown 8vo, 2s. 6d.

A Short Manual for Monthly Nurses. By CHARLES J. CULLINGWORTH, M.D., F.R.C.P., Obstetric Physician to St. Thomas's Hospital. Revised by the Author, with the assistance of M. A. ATKINSON, Matron of the General Lying-in Hospital, Lambeth. Fifth Edition. Fcap. 8vo, 1s. 6d.

Notes on Gynæcological Nursing. By John BENJAMIN HELLIER, M.D., Surgeon to the Hospital for Women, etc., Leeds. Crown 8vo, 1s. 6d.

Lectures on Medicine to Nurses. By Herbert E. CUFF, M.D., F.R.C.S., Medical Superintendent, North Eastern Fever Hospital, London. Third Edition. With 29 Illustrations. Crown 8vo, 3s. 6d.

Antiseptic Principles for Nurses. By C. E. RICHMOND, F.R.C.S. Fcap. 8vo, 1s.

A Practical Treatise on Disease in Children. By EUSTACE SMITH, M.D., F.R.C.P., Physician to the King of the Belgians, and to the East London Hospital for Children, etc. Second Edition. 8vo, 22s.

By the same Author.

Clinical Studies of Disease in Children. Second Edition. Post 8vo, 7s. 6d.

Also.

The Wasting Diseases of Infants and Children. Sixth (cheap) Edition. Post 8vo, 6s.

The Diseases of Children. By James F. GOODHART, M.D., F.R.C.P. Sixth Edition, with the assistance of G. F. STILL, M.D., Medical Registrar to the Hospital for Sick Children, Great Ormond Street. Crown 8vo, 10s. 6d.

On the Natural and Artificial Methods of Feeding Infants and Young Children. By EDMUND CAUTLEY, M.D., Physician to the Belgrave Hospital for Children. Crown 8vo, 7s. 6d.

Materia Medica, Pharmacy, Pharmacology, and Therapeutics. By W. HALE WHITE, M.D., F.R.C.P., Physician to, and Lecturer on Pharmacology and Therapeutics at, Guy's Hospital. Sixth Edition, based upon the B.P. of 1898 and the Indian and Colonial Addendum. Fcap. 8vo, 7s. 6d.

Southall's Organic Materia Medica. Sixth Edition, adapted to the B.P. of 1898. Edited by JOHN BARCLAY, B.Sc. Lond. Crown 8vo, 7s. 6d.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

An Introduction to the Study of Materia

Medica, designed for Students of Pharmacy and Medicine. By HENRY G. GREENISH, F.I.C., F.L.S., Professor of Materia Medica and Pharmacy to the Pharmaceutical Society. With 213 Illustrations, 8vo, 15s.

Materia Medica and Therapeutics. By Charles

D. F. PHILLIPS, M.D., LL.D., F.R.S. Edin.

Vegetable Kingdom—Organic Compounds—Animal Kingdom. 8vo, 25s.
Inorganic Substances. Second Edition. 8vo, 21s.

Practical Pharmacy: an Account of the

Methods of Manufacturing and Dispensing Pharmaceutical Preparations; with a chapter on the Analysis of Urine. By E. W. LUCAS, F.C.S., Examiner at the Pharmaceutical Society. With 283 Illustrations. Roy. 8vo, 12s. 6d.

Galenic Pharmacy: a Practical Handbook to

the Processes of the British Pharmacopœia. By R. A CRIPPS, M.P.S. 8vo, with 76 Engravings, 8s. 6d.

Practical Pharmacy. By Barnard S. Proctor.

Third Edition. 8vo, with Engravings and Fac-simile Prescriptions, 14s.

The Galenical Preparations of the British

Pharmacopœia: a Handbook for Medical and Pharmaceutical Students. By C. O. HAWTHORNE, M.D., C.M., late Lecturer on Materia Medica and Therapeutics, Queen Margaret's College, Glasgow. 8vo, 4s. 6d.

The Pharmaceutical Formulary: a Synopsis

of the British and Foreign Pharmacopœias. By HENRY BEASLEY. Twelfth Edition by J. OLDHAM BRAITHWAITE. 18mo, 6s. 6d.

By the same Author.

The Druggist's General Receipt-Book. Tenth

Edition. 18mo, 6s. 6d.

Also.

The Book of Prescriptions: containing upwards

of 3,000 Prescriptions collected from the Practice of the most eminent Physicians and Surgeons, English and Foreign. Seventh Edition, 18mo, 6s. 6d.

A Companion to the British Pharmacopœia.

By PETER SQUIRE. Revised by PETER WYATT SQUIRE, F.L.S., F.C.S. Seventeenth Edition. 8vo, 12s. 6d.

By the same Authors.

The Pharmacopœias of thirty of the London

Hospitals, arranged in Groups for Comparison. Seventh Edition. Fcap. 8vo, 6s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Pereira's Selecta è Prescriptis : containing

Lists of Terms used in Prescriptions, with Explanatory Notes, etc.
Also, a Series of Abbreviated Prescriptions with Translations.
Eighteenth Edition, by JOSEPH INCE. 24mo, 5s.

Year-Book of Pharmacy: containing the Trans-

actions of the British Pharmaceutical Conference. Annually. 8vo, 10s.

Manual of Botany, in two Vols., crown 8vo.

By J. REYNOLDS GREEN, Sc.D., M.A., F.R.S., Professor of Botany to the Pharmaceutical Society.

Vol. I.: Morphology and Anatomy, with 788 Engravings. Second Edition. 7s. 6d.

Vol. II.: Classification and Physiology, with 417 Engravings, 10s.

By the same Author.

An Introduction to Vegetable Physiology.

With 184 Illustrations, 8vo, 10s. 6d.

The Student's Guide to Systematic Botany,

including the Classification of Plants and Descriptive Botany. By ROBERT BENTLEY, late Emeritus Professor of Botany in King's College and to the Pharmaceutical Society. Fcap. 8vo, with 350 Engravings, 3s. 6d.

Medicinal Plants: being Descriptions with

original figures, of the Principal Plants employed in Medicine, and an account of their Properties and Uses. By Prof. BENTLEY and Dr. H. TRIMEN, F.R.S. In 4 vols., large 8vo, with 306 Coloured Plates, bound in Half Morocco, Gilt Edges, £11 11s.

Therapeutic Electricity and Practical Muscle

Testing. By W. S. HEDLEY, M.D., in charge of the Electro-therapeutic Department of the London Hospital. With 110 Illustrations. Roy. 8vo, 8s. 6d.

Practical Therapeutics: a Manual. By

EDWARD J. WARING, C.I.E., M.D., F.R.C.P., and DUDLEY W. BUXTON, M.D., B.S. Lond. Fourth Edition. Crown 8vo, 14s.

By the same Author.

Bazaar Medicines of India, and Common

Medical Plants. With Full Index of Diseases, indicating their Treatment by these and other Agents procurable throughout India, etc. Fifth Edition. Fcap. 8vo, 5s.

7, GREAT MARLBOROUGH STREET.

*

J. & A. Churchill's Recent Works.

Climate and Fevers of India, with a Series
of Cases (Croonian Lectures, 1882). By Sir JOSEPH FAYRER,
K.C.S.I., M.D. 8vo, with 17 Temperature Charts, 12s.

Psilosis or "Sprue": its Nature and Treat-
ment; with Observations on various Forms of Diarrhœa acquired in
the Tropics. By GEORGE THIN, M.D. Second and Enlarged Edition,
with Illustrations. 8vo, 10s.

A Manual of Family Medicine and Hygiene
for India. Published under the Authority of the Government of
India. By Sir WILLIAM J. MOORE, K.C.I.E., M.D., late Surgeon-
General with the Government of Bombay. Sixth Edition. Post 8vo,
with 71 Engravings, 12s.

By the same Author.

A Manual of the Diseases of India: with a
Compendium of Diseases generally. Second Edition. Post 8vo,
10s.

The Prevention of Disease in Tropical and
Sub-Tropical Campaigns. (Parkes Memorial Prize for 1886.) By
Lieut.-Col. ANDREW DUNCAN, M.D., B.S. Lond., F.R.C.S., H.M.
Indian Medical Service. 8vo, 12s. 6d.

A Commentary on the Diseases of India. By
NORMAN CHEVERS, C.I.E., M.D., F.R.C.S., Deputy Surgeon-General
H.M. Indian Army. 8vo, 24s.

Hooper's Physicians' Vade-Mecum: a Manual
of the Principles and Practice of Physic. Tenth Edition. By W. A.
GUY, F.R.C.P., F.R.S., and J. HARLEY, M.D., F.R.C.P. With 118
Engravings. Fcap. 8vo, 12s. 6d.

A Text-Book of Medicine. By the late C.
HILTON FAGGE, M.D., and P. H. PYE-SMITH, M.D., F.R.S., Consulting
Physician to Guy's Hospital. Fourth Edition. 2 vols. royal 8vo.
Vol. I., cloth, 21s.

Manual of the Practice of Medicine. By
FREDERICK TAYLOR, M.D., F.R.C.P., Physician to, and Lecturer
on Medicine at, Guy's Hospital. Sixth Edition. Post 8vo, with
Engravings, 16s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

A Dictionary of Practical Medicine. By various writers. Edited by JAS. KINGSTON FOWLER, M.A., M.D., F.R.C.P., Physician to Middlesex Hospital and the Hospital for Consumption. 8vo, cloth, 21s.; half calf, 25s.

The Practice of Medicine. By M. Charteris, M.D., Professor of Therapeutics and Materia Medica in the University of Glasgow. Eighth Edition. Edited by F. J. CHARTERIS, M.B., Ch. B. Crown 8vo, with Engravings on Copper and Wood, 10s.

A Text-Book of Bacteriology for Students and Practitioners of Medicine. By G. M. STERNBERG, M.D., Surgeon-General, U.S. Army. With 9 Plates and 200 Figures in the Text. 8vo, 24s.

How to Examine the Chest: a Practical Guide for the use of Students. By SAMUEL WEST, M.D., F.R.C.P., Assistant Physician to St. Bartholomew's Hospital. Third Edition. With 46 Engravings. Fcap. 8vo, 5s.

An Atlas of the Pathological Anatomy of the Lungs. By the late WILSON FOX, M.D., F.R.S., F.R.C.P., Physician to H.M. the Queen. With 45 Plates (mostly Coloured) and Engravings. 4to, half-bound in Calf, 70s.

By the same Author.

A Treatise on Diseases of the Lungs and Pleura. Edited by SIDNEY COUPLAND, M.D., F.R.C.P., Physician to Middlesex Hospital. Roy. 8vo, with Engravings; also Portrait and Memoir of the Author, 36s.

The Student's Guide to Diseases of the Chest. By VINCENT D. HARRIS, M.D. Lond., F.R.C.P., Physician to the City of London Hospital for Diseases of the Chest, Victoria Park. Fcap 8vo, with 55 Illustrations (some Coloured), 7s. 6d.

The Schott Methods of the Treatment of Chronic Diseases of the Heart, with an account of the Nauheim Baths, and of the Therapeutic Exercises. By W. BEZLY THORNE, M.D., M.R.C.P. Third Edition. 8vo, with Illustrations, 6s.

Guy's Hospital Reports. By the Medical and Surgical Staff. Vol. XL. Third Series. 8vo, 10s. 6d.

St. Thomas's Hospital Reports. By the Medical and Surgical Staff. Vol. XXVII. New Series. 8vo, 8s. 6d.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Text-Book of Medical Treatment (Diseases and Symptoms). By NESTOR I. C. TIRARD, M.D., F.R.C.P., Professor of the Principles and Practice of Medicine, King's College, London. 8vo, 15s.

Medical Diagnosis. (Student's Guide Series.) By SAMUEL FENWICK, M.D., F.R.C.P., and W. SOLTAU FENWICK, M.D., B.S. Eighth Edition. Crown 8vo, with 135 Engravings, 9s.

By the same Authors.

Outlines of Medical Treatment. Fourth Edition. Crown 8vo, with 35 Engravings, 10s.

Also.

Ulcer of the Stomach and Duodenum. With 55 Illustrations. Roy. 8vo, 10s. 6d.

Also, by Dr. Samuel Fenwick.

Clinical Lectures on some Obscure Diseases of the Abdomen. Delivered at the London Hospital. 8vo, with Engravings, 7s. 6d.

And

The Saliva as a Test for Functional Diseases of the Liver. Crown 8vo, 2s.

The Liver. By LIONEL S. BEALE, M.B., F.R.S., Consulting Physician to King's College Hospital. With 24 Plates (85 Figures), 8vo, 5s.

By the same Author.

On Slight Ailments: and on Treating Disease. Fourth Edition. 8vo, 5s.

The Blood: how to Examine and Diagnose its Diseases. By ALFRED C. COLES, M.D., B.Sc. With 6 Coloured Plates. 8vo, 10s. 6d.

The Physiology of the Carbohydrates; their Application as Food and Relation to Diabetes. By F. W. PAVY, M.D., LL.D., F.R.S., F.R.C.P., Consulting Physician to Guy's Hospital. Royal 8vo, with Plates and Engravings, 10s. 6d.

Medical Lectures and Essays. By Sir G. JOHNSON, M.D., F.R.C.P., F.R.S., Consulting Physician to King's College Hospital. 8vo, with 46 Engravings, 25s.

By the same Author.

An Essay on Asphyxia (Apnœa). 8vo, 3s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Uric Acid as a Factor in the Causation of
Disease. By ALEXANDER HAIG, M.D., F.R.C.P. Physician to the
Metropolitan Hospital and the Royal Hospital for Children and
Women. Fifth Edition. 8vo, with 75 Illustrations, 14s.

Bronchial Asthma: its Pathology and Treat-
ment. By J. B. BERKART, M.D., late Physician to the City of
London Hospital for Diseases of the Chest. Second Edition, with 7
Plates (35 Figures). 8vo, 10s. 6d.

Treatment of Some of the Forms of Valvular
Disease of the Heart. By A. E. SANSOM, M.D., F.R.C.P., Physician
to the London Hospital. Second Edition. Fcap. 8vo, with 26 Engrav-
ings, 4s. 6d.

Medical Ophthalmoscopy: a Manual and Atlas.
By Sir WILLIAM R. GOWERS, M.D., F.R.C.P., F.R.S. Third Edition.
Edited with the assistance of MARCUS GUNN, M.B., F.R.C.S., Surgeon
to the Royal London Ophthalmic Hospital. With Coloured Plates
and Woodcuts. 8vo, 16s.

By the same Author.

A Manual of Diseases of the Nervous System.
VOL. I.—Nerves and Spinal Cord. Third
Edition, by the Author and JAMES TAYLOR, M.D., F.R.C.P.
Roy. 8vo, with 192 Engravings, 15s.

VOL. II.—Brain and Cranial Nerves: General
and Functional Diseases of the Nervous System. Second Edition
Roy. 8vo, with 182 Engravings, 20s.

Also.

Clinical Lectures on Diseases of the Nervous
System. 8vo. 7s. 6d.

Also.

Epilepsy and other Chronic Convulsive
Diseases: their Causes, Symptoms, and Treatment. Second Edition
8vo, 10s. 6d.

Also.

Diagnosis of Diseases of the Brain. Second
Edition. 8vo, with Engravings, 7s. 6d.

Also.

Syphilis and the Nervous System: being a
Revised Reprint of the Lettsomian Lectures for 1890. Delivered before
the Medical Society of London. 8vo, 4s.

The Nervous System, Diseases of. By J. A.
ORMEROD, M.D., F.R.C.P., Physician to the National Hospital for the
Paralysed and Epileptic. With 66 Illustrations. Fcap. 8vo, 8s. 6d.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Text-Book of Nervous Diseases for Students

and Practitioners of Medicine. By CHARLES L. DANA, M.D., Professor of Nervous and Mental Diseases in Bellevue Hospital Medical College, New York. Fourth Edition. With 246 Illustrations. 8vo, 20s.

Diseases of the Nervous System. Lectures

delivered at Guy's Hospital. By Sir SAMUEL WILKS, Bart., M.D., F.R.S. Second Edition. 8vo, 18s.

Handbook of the Diseases of the Nervous

System. By JAMES ROSS, M.D., F.R.C.P., late Professor of Medicine in the Victoria University, and Physician to the Royal Infirmary, Manchester. Roy. 8vo, with 184 Engravings, 18s.

Stammering: its Causes, Treatment, and

Cure. By A. G. BERNARD, M.R.C.S., L.R.C.P. Crown 8vo, 2s.

Secondary Degenerations of the Spinal Cord

(Gulstonian Lectures, 1889). By HOWARD H. TOOTH, M.D., F.R.C.P., Assistant Physician to the National Hospital for the Paralysed and Epileptic. With Plates and Engravings. 8vo, 3s. 6d.

Diseases of the Nervous System. Clinical

Lectures. By THOMAS BUZZARD, M.D., F.R.C.P., Physician to the National Hospital for the Paralysed and Epileptic. With Engravings. 8vo, 15s.

By the same Author.

Some Forms of Paralysis from Peripheral

Neuritis; of Gouty, Alcoholic, Diphtheritic, and other origin. Crown 8vo, 5s.

Also.

On the Simulation of Hysteria by Organic

Disease of the Nervous System. Crown 8vo, 4s. 6d.

On the Typhoid Bacillus and Typhoid Fever,

being the Goulstonian Lectures delivered before the Royal College of Physicians in March, 1900, by P. HORTON-SMITH, M.D., F.R.C.P. With Illustrations, 8vo, 2s. 6d.

Gout in its Clinical Aspects. By J. Mortimer

GRANVILLE, M.D. Crown 8vo, 6s.

Diseases of the Liver: with and without

Jaundice. By GEORGE HARLEY, M.D., F.R.C.P., F.R.S. 8vo, with 2 Plates and 36 Engravings, 21s.

Rheumatic Diseases (Differentiation in). By

HUGH LANE, Surgeon to the Royal Mineral Water Hospital, Bath. Second Edition, much Enlarged, with 8 Plates. Crown 8vo, 3s. 6d.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Diseases of the Abdomen, comprising those of the Stomach and other parts of the Alimentary Canal, Œsophagus, Cæcum, Intestines, and Peritoneum. By S. O. HABERSHON, M.D., F.R.C.P. Fourth Edition. 8vo, with 5 Plates, 21s.

On Gallstones, or Cholelithiasis. By E. M. BROCKBANK, M.D. Vict., M.R.C.P. Lond., Honorary Physician to the Ancoats Hospital, Manchester. Crown 8vo, 7s.

On the Relief of Excessive and Dangerous Tympanites by puncturing the Abdomen. By JOHN W. OGLE, M.D., Consulting Physician to St. George's Hospital. 8vo, 5s. 6d.

Headaches: their Nature, Causes, and Treatment. By W. H. DAY, M.D., Physician to the Samaritan Hospital. Fourth Edition. Crown 8vo, with Engravings, 7s. 6d.

A Handbook of Medical Climatology, embodying its Principles and Therapeutic Application, with Scientific Data of the chief Health Resorts of the World. By S. EDWIN SOLLY, M.D. M.R.C.S., late President of the American Climatological Association. With Engravings and Coloured Plates. 8vo, 16s.

The Mineral Waters of France, and its Wintering Stations (Medical Guide to). With a Special Map. By A. VINTRAS, M.D., Physician to the French Embassy, and to the French Hospital, London. Second Edition. Crown 8vo, 8s.

Surgery: its Theory and Practice. By William J. WALSHAM, F.R.C.S., Surgeon to, and Lecturer on Anatomy at, St. Bartholomew's Hospital. Seventh Edition. Post 8vo, with 483 Engravings (including 28 Skiagrams), 15s.

A Synopsis of Surgery. By R. F. Tobin, Surgeon to St. Vincent's Hospital, Dublin. Crown 8vo, interleaved, leather binding, 6s. 6d.

Surgical Emergencies: together with the Emergencies attendant on Parturition and the Treatment of Poisoning. By PAUL SWAIN, F.R.C.S., Surgeon to the South Devon and East Cornwall Hospital. Fifth Edition. Crown 8vo, with 149 Engravings, 6s.

Illustrated Ambulance Lectures: (to which is added a NURSING LECTURE) in accordance with the Regulations of the St. John's Ambulance Association for Male and Female Classes. By JOHN M. H. MARTIN, M.D., F.R.C.S., Hon. Surgeon to the Blackburn Infirmary. Fourth Edition. Crown 8vo, with 60 Engravings, 2s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Operations on the Brain (a Guide to). By ALEC FRASER, Professor of Anatomy, Royal College of Surgeons in Ireland. Illustrated by 42 life-size Plates in Autotype, and 2 Woodcuts in the text. Folio, 63s.

Abdominal Surgery. By J. Greig Smith, M.A., F.R.S.E. Sixth Edition. Edited by JAMES SWAIN, M.S., M.D. Lond., F.R.C.S. Eng., Assistant-Surgeon to the Bristol Royal Infirmary, Professor of Surgery, University College, Bristol. 2 vols., 8vo, with 224 Engravings, 36s.

The Physiology of Death from Traumatic Fever; a Study in Abdominal Surgery. By JOHN D. MALCOLM, M.B., C.M., F.R.C.S.E., Surgeon to the Samaritan Free Hospital. 8vo, 3s. 6d.

The Surgery of the Alimentary Canal. By ALFRED ERNEST MAYLARD, M.B. Lond. and B.S., Senior Surgeon to the Victoria Infirmary, Glasgow. With 27 Swantype Plates and 89 Figures in the Text, 8vo, 25s.

By the same Author.

A Student's Handbook of the Surgery of the Alimentary Canal. With 97 Illustrations. Crown 8vo, 8s. 6d.

Surgery. By C. W. Mansell Moullin, M.A., M.D. Oxon., F.R.C.S., Surgeon and Lecturer on Physiology to the London Hospital. Large 8vo, with 497 Engravings, 34s.

The Practice of Surgery : a Manual. By THOMAS BRYANT, Consulting Surgeon to Guy's Hospital. Fourth Edition. 2 vols. crown 8vo, with 750 Engravings (many being Coloured), and including 6 chromo plates, 32s.

The Surgeon's Vade-Mecum : a Manual of Modern Surgery. By R. DRUITT, F.R.C.S. Twelfth Edition. By STANLEY BOYD, M.B., F.R.C.S., Assistant Surgeon and Pathologist to Charing Cross Hospital. Crown 8vo, with 373 Engravings, 16s.

The Operations of Surgery : intended for use on the Dead and Living Subject alike. By W. H. A. JACOBSON, M.A., M.B., M.Ch. Oxon., F.R.C.S., Assistant Surgeon to, and Lecturer on Anatomy at, Guy's Hospital. Third Edition. 8vo, with 401 Illustrations, 34s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

A Course of Operative Surgery. By CHRISTOPHER HEATH, Surgeon to University College Hospital. Second Edition. With 20 Coloured Plates (180 figures) from Nature, by M. LÉVEILLÉ, and several Woodcuts. Large 8vo, 30s.

By the same Author.

The Student's Guide to Surgical Diagnosis. Second Edition. Fcap. 8vo, 6s. 6d.

Also.

A Manual of Minor Surgery and Bandaging. For the use of House-Surgeons, Dressers, and Junior Practitioners. Twelfth Edition, revised by BILTON POLLARD, F.R.C.S., Surgeon to University College Hospital. Fcap. 8vo, with 195 Engravings, 6s. 6d.

Also.

Injuries and Diseases of the Jaws. Fourth Edition. Edited by HENRY PERCY DEAN, M.S., F.R.C.S., Assistant Surgeon to the London Hospital. 8vo, with 187 Wood Engravings, 14s.

Ovariectomy and Abdominal Surgery. By HARRISON CRIPPS, F.R.C.S., Surgical Staff, St. Bartholomew's Hospital. With numerous Plates, royal 8vo, 25s.

Diseases of Bones and Joints. By CHARLES MACNAMARA, F.R.C.S., Surgeon to, and Lecturer on Surgery at, the Westminster Hospital. 8vo, with Plates and Engravings, 12s.

Surgical Pathology and Morbid Anatomy. By ANTHONY A. BOWLBY, F.R.C.S., Assistant Surgeon to St. Bartholomew's Hospital. Fourth Edition. Crown 8vo, with 186 Engravings, 10s. 6d.

By the same Author.

Injuries and Diseases of Nerves, and their Surgical Treatment. 8vo, with 20 Plates, 14s.

Chloroform: a Manual for Students and Practitioners. By EDWARD LAWRIE, M.B. Edin., Lieut.-Col. I.M.S., Residency Surgeon, Hyderabad. Illustrated, crown 4to, 5s. net.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Diseases of the Thyroid Gland and their

Surgical Treatment. By JAMES BERRY, B.S. Lond., F.R.C.S., Surgeon to the Royal Free Hospital and Lecturer on Surgery at the London School of Medicine for Women. With 121 Illustrations, 8vo, 14s.

The Human Foot: its Form and Structure,

Functions and Clothing. By THOMAS S. ELLIS, Consulting Surgeon to the Gloucester Infirmary. With 7 Plates and Engravings (50 Figures). 8vo, 7s. 6d.

Short Manual of Orthopædy. By Heather

BIGG, F.R.C.S. Ed., Part I. Deformities and Deficiencies of the Head and Neck. 8vo, 2s. 6d.

Face and Foot Deformities. By Frederick

CHURCHILL, C.M. 8vo, with Plates and Illustrations, 10s. 6d.

Royal London Ophthalmic Hospital Reports.

By the Medical and Surgical Staff. Vol. XIV., Part 2. 8vo, 5s.

Ophthalmological Society of the United King-

dom. Transactions. Vol. XX. 8vo, 12s. 6d.

Manual of Ophthalmic Surgery and Medicine.

By W. H. H. JESSOP, M.A., F.R.C.S., Ophthalmic Surgeon to St. Bartholomew's Hospital. With 5 Coloured Plates and 110 Woodcuts. Crown 8vo, 9s. 6d.

Nettleship's Diseases of the Eye. Sixth Edition.

Revised and Edited by W. T. HOLMES SPICER, M.B., F.R.C.S., Ophthalmic Surgeon to St. Bartholomew's Hospital and the Victoria Hospital for Children. With 161 Engravings and a Coloured Plate illustrating Colour-Blindness. Crown 8vo, 8s. 6d.

Diseases and Refraction of the Eye. By

N. C. MACNAMARA, F.R.C.S., Surgeon to Westminster Hospital, and GUSTAVUS HARTRIDGE, F.R.C.S., Surgeon to the Royal Westminster Ophthalmic Hospital. Fifth Edition. Crown 8vo, with Plate, 156 Engravings, also Test-types, 10s. 6d.

On Diseases and Injuries of the Eye: a Course

of Systematic and Clinical Lectures to Students and Medical Practitioners. By J. R. WOLFE, M.D., F.R.C.S.E. With 10 Coloured Plates and 157 Wood Engravings. 8vo, 21s.

Convergent Strabismus, and its Treatment,

an Essay. By EDWIN HOLTHOUSE, M.A., F.R.C.S., Surgeon to the Western Ophthalmic Hospital. 8vo, 6s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Normal and Pathological Histology of the

Human Eye and Eyelids. By C. FRED. POLLOCK, M.D., F.R.C.S., and F.R.S.E., Surgeon for Diseases of the Eye to Anderson's College Dispensary, Glasgow. Crown 8vo, with 100 Plates (230 drawings), 15s.

Atlas of Ophthalmoscopy. Composed of 12

Chromo-lithographic Plates (59 Figures drawn from nature) and Explanatory Text. By RICHARD LIEBREICH, M.R.C.S. Translated by H. ROSBOROUGH SWANZY, M.B. Third Edition, 4to, 40s.

Refraction of the Eye: a Manual for Students.

By GUSTAVUS HARTRIDGE, F.R.C.S., Surgeon to the Royal Westminster Ophthalmic Hospital. Eleventh Edition. Crown 8vo, with 105 Illustrations, also Test-types, etc., 6s.

By the same Author.

The Ophthalmoscope: a Manual for Students.

Fourth Edition. Crown 8vo, with 65 Illustrations and 4 Plates, 4s. 6d.

Glaucoma: its Pathology and Treatment. By

PRIESTLEY SMITH, Ophthalmic Surgeon to the Queen's Hospital, Birmingham. 8vo, with 64 Engravings and 12 Zinco-photographs. 7s. 6d.

Methods of Operating for Cataract and

Secondary Impairments of Vision, with the results of 500 cases. By Major G. H. FINK, H.M. Indian Medical Service. Crown 8vo, with 15 Engravings, 5s.

Diseases of the Eye: a Practical Handbook

for General Practitioners and Students. By CECIL EDWARD SHAW, M.D., M.Ch., Ophthalmic Surgeon to the Ulster Hospital for Children and Women, Belfast. With a Test-Card for Colour-Blindness. Crown 8vo, 3s. 6d.

Diseases of the Ear, including the Anatomy

and Physiology of the Organ, together with the Treatment of the Affections of the Nose and Pharynx, which conduce to Aural Disease (a Treatise). By T. MARK HOVELL, Senior Aural Surgeon to the London Hospital, and Lecturer on Diseases of the Throat in the College. Second Edition. 8vo, with 128 Engravings, 21s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works

Diseases and Injuries of the Ear. By Sir WILLIAM B. DALBY, F.R.C.S., M.B., Consulting Aural Surgeon to St. George's Hospital. Fourth Edition. Crown 8vo, with 8 Coloured Plates and 38 Wood Engravings. 10s. 6d.

By the same Author.

Short Contributions to Aural Surgery, between 1875 and 1896. Third Edition. 8vo, with Engravings, 5s.

A System of Dental Surgery. By Sir John TOMES, F.R.S., and C. S. TOMES, M.A., F.R.S. Fourth Edition. Post 8vo, with 289 Engravings, 16s.

Dental Anatomy, Human and Comparative : A Manual. By CHARLES S. TOMES, M.A., F.R.S. Fifth Edition. Post 8vo, with 263 Engravings, 14s.

Decay in Teeth : an Investigation into its Cause and Prevention. By J. SIM WALLACE, M.D., B.Sc., L.D.S.R.C.S. 8vo, 5s.

Dental Materia Medica, Pharmacology and Therapeutics. By CHARLES W. GLASSINGTON, M.R.C.S., L.D.S. Edin.; Senior Dental Surgeon, Westminster Hospital; Dental Surgeon, National Dental Hospital, and Lecturer on Dental Materia Medica and Therapeutics to the College. Crown 8vo, 6s.

Dental Medicine : a Manual of Dental Materia Medica and Therapeutics. By FERDINAND J. S. GORGAS, M.D., D.D.S., Professor of the Principles of Dental Science in the University of Maryland. Sixth Edition. 8vo, 18s.

A Manual of Dental Metallurgy. By Ernest A. SMITH, F.I.C., Assistant Instructor in Metallurgy, Royal College of Science, London. With 37 Illustrations, crown 8vo, 6s. 6d.

A Manual of Nitrous Oxide Anæsthesia. By J. FREDERICK W. SILK, M.D. Lond., M.R.C.S., Assistant Anæsthetist to Guy's Hospital, Anæsthetist to the Dental School of Guy's Hospital, and to the Royal Free Hospital. 8vo, with 26 Engravings, 5s.

Practical Treatise on Mechanical Dentistry. By JOSEPH RICHARDSON, M.D., D.D.S. Seventh Edition, revised and edited by GEORGE W. WARREN, D.D.S. Royal 8vo, with 690 Engravings, 22s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

A Handbook on Leprosy. By S. P. Impey, M.D., late Chief and Medical Superintendent, Robben Island Leper and Lunatic Asylums, Cape Colony. With 38 Plates, 8vo, 12s.

Diseases of the Skin (Introduction to the Study of). By P. H. PYE-SMITH, M.D., F.R.S., F.R.C.P., Physician to Guy's Hospital. Crown 8vo, with 26 Engravings, 7s. 6d.

A Manual of Diseases of the Skin, with an Analysis of 20,000 Consecutive Cases and a Formulary. By DUNCAN E. BULKLEY, M.D., New York. Fourth Edition, royal 16mo, 6s. 6d.

Skin Diseases of Children. By Geo. H. Fox, M.D., Clinical Professor of Diseases of the Skin, College of Physicians and Surgeons, New York. With 12 Photogravure and Chromographic Plates and 60 Illustrations in the Text. Royal 8vo, 12s. 6d.

The Operative Surgery of Malignant Disease. By HENRY T. BUTLIN, F.R.C.S., Surgeon to St. Bartholomew's Hospital. Second Edition, with 12 Engravings. 8vo, 14s.

By the same Author.

Malignant Disease (Sarcoma and Carcinoma) of the Larynx. 8vo, with 5 Engravings, 5s.

Also.

Sarcoma and Carcinoma: their Pathology, Diagnosis, and Treatment. 8vo, with 4 Plates, 8s.

Cancers and the Cancer Process: a Treatise, Practical and Theoretic. By HERBERT L. SNOW, M.D., Surgeon to the Cancer Hospital, Brompton. 8vo, with 15 Plates. 15s.

By the same Author.

The Re-appearance (Recurrence) of Cancer after apparent Extirpation. 8vo, 5s. 6d.

Also.

The Palliative Treatment of Incurable Cancer. Crown 8vo, 2s. 6d.

The Diagnosis and Treatment of Syphilis. By TOM ROBINSON, M.D. St. And., Physician to the Western Skin Hospital. Second Edition. Crown 8vo, 3s. 6d.

By the same Author.

The Diagnosis and Treatment of Eczema. Second Edition. Crown 8vo, 3s. 6d.

Also.

Illustrations of Diseases of the Skin and Syphilis, with Remarks. Fasc. I. with 3 Plates. Imp. 4to, 5s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Cancerous Affections of the Skin (Epithelioma and Rodent Ulcer). By GEORGE THIN, M.D. Post 8vo, with 8 Engravings, 5s.
By the same Author.

Pathology and Treatment of Ringworm. 8vo, with 21 Engravings, 5s.

Ringworm, and some other Scalp Affections: their Cause and Cure. By HAYDN BROWN, L.R.C.P. Ed. 8vo, 5s.

Urinary and Renal Derangements and Calculous Disorders. By LIONEL S. BEALE, F.R.C.P., F.R.S., Physician to King's College Hospital. 8vo, 5s.

Chemistry of Urine: a Practical Guide to the Analytical Examination of Diabetic, Albuminous, and Gouty Urine. By ALFRED H. ALLEN, F.I.C., F.C.S., Public Analyst for the West Riding of Yorkshire, &c. 8vo, with Engravings, 7s. 6d.

Clinical Chemistry of Urine (Outlines of the). By C. A. MACMUNN, M.A., M.D. 8vo, with 64 Engravings and Plate of Spectra, 9s.

Diseases of the Male Organs of Generation. By W. H. A. JACOBSON, M.Ch.Oxon., F.R.C.S., Assistant-Surgeon to Guy's Hospital. 8vo, with 88 Engravings, 22s.

Atlas of Electric Cystoscopy. By E. HURRY FENWICK, F.R.C.S., Surgeon to the London Hospital and St. Peter's Hospital for Stone. Royal 8vo, with 34 Coloured Plates, embracing 83 Figures. 21s.

By the same Author.

Electric Illumination of the Bladder and Urethra, as a Means of Diagnosis of Obscure Vesico-Urethral Diseases. By E. HURRY FENWICK, F.R.C.S., Surgeon to London Hospital and St. Peter's Hospital for Stone. Second Edition. 8vo, with 54 Engravings, 6s. 6d.

Also.

Operative and Inoperative Tumours of the Urinary Bladder: a Clinical and Operative Study based on 500 cases. With 39 Illustrations, 8vo, 5s.

Also.

Tumours of the Urinary Bladder. Fas. I. Royal 8vo, 5s.

Also.

Ulceration of the Bladder, Simple, Tuberculous, and Malignant: a Clinical Study. With Illustrations, 8vo, 5s.

Also.

The Cardinal Symptoms of Urinary Disease: their Diagnostic Significance and Treatment. 8vo, with 36 Illustrations, 8s. 6d.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

By SIR HENRY THOMPSON, BART., F.R.C.S.

Diseases of the Urinary Organs. Clinical
Lectures. Eighth Edition. 8vo, with 121 Engravings, 10s. 6d.

Some Important Points connected with the
Surgery of the Urinary Organs. Lectures delivered in the R.C.S.
8vo, with 44 Engravings. Student's Edition, 2s. 6d.

Practical Lithotomy and Lithotrity; or, an
Inquiry into the Best Modes of Removing Stone from the Bladder.
Third Edition. 8vo, with 87 Engravings, 10s.

The Preventive Treatment of Calculous Dis-
ease, and the Use of Solvent Remedies. Third Edition. Cr. 8vo, 2s. 6d.

Tumours of the Bladder: their Nature, Sym-
ptoms, and Surgical Treatment. 8vo, with numerous Illustrations, 5s.

Stricture of the Urethra, and Urinary Fistulæ:
their Pathology and Treatment. Fourth Edition. 8vo, with 74 En-
gravings, 6s.

The Suprapubic Operation of Opening the
Bladder for Stone and for Tumours. 8vo, with Engravings, 3s. 6d.

The Clinical Examination of Urine, with an
Atlas of Urinary Deposits. By LINDLEY SCOTT, M.A., M.D., with
41 original Plates (mostly in colours). Crown 4to, 15s. net.

The Surgical Diseases of the Genito-Urinary
Organs, including Syphilis. By E. L. KEYES, M.D., Professor of
Genito-Urinary Surgery, Syphilology, and Dermatology in Bellevue
Hospital Medical College, New York (a revision of VAN BUREN and
KEYES' Text-book). Roy. 8vo, with 114 Engravings, 21s.

Selected Papers on Stone, Prostate, and other
Urinary Disorders. By REGINALD HARRISON, F.R.C.S., Surgeon to
St. Peter's Hospital. 8vo, with 15 Illustrations, 5s.

Syphilis. By Alfred Cooper, F.R.C.S., Con-
sulting Surgeon to the West London and the Lock Hospitals. Second
Edition. Edited by EDWARD COTTERELL, F.R.C.S., Surgeon (out-
patients) to the London Lock Hospital. 8vo, with 24 Full-page
Plates (12 coloured), 18s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

On Maternal Syphilis, including the presence
and recognition of Syphilitic Pelvic Disease in Women. By JOHN
A. SHAW-MACKENZIE, M.D. With Coloured Plates. 8vo, 10s. 6d.

Diseases of the Rectum and Anus. By Alfred
COOPER, F.R.C.S., Senior Surgeon to St. Mark's Hospital for
Fistula; and F. SWINFORD EDWARDS, F.R.C.S., Senior Assistant
Surgeon to St. Mark's Hospital. Second Edition, with Illustrations.
8vo, 12s.

Diseases of the Rectum and Anus. By
HARRISON CRIPPS, F.R.C.S., Assistant Surgeon to St. Bartholomew's
Hospital, etc. Second Edition. 8vo, with 13 Lithographic Plates and
numerous Wood Engravings, 12s. 6d.

By the same Author.

Cancer of the Rectum. Especially considered
with regard to its Surgical Treatment. Jacksonian Prize Essay.
Third Edition. 8vo, with 13 Plates and several Wood Engravings, 6s.

Also

The Passage of Air and Fæces from the
Urethra. 8vo, 3s. 6d.

A Medical Vocabulary: an Explanation of all
Terms and Phrases used in the various Departments of Medical Science
and Practice, their Derivation, Meaning, Application, and Pronuncia-
tion. By R. G. MAYNE, M.D., LL.D. Sixth Edition, by W. W.
WAGSTAFFE, B.A., F.R.C.S. Crown 8vo, 10s. 6d.

A Short Dictionary of Medical Terms. Being
an Abridgment of Mayne's Vocabulary. 64mo, 2s. 6d.

Dunghlison's Dictionary of Medical Science.
Containing a full Explanation of its various Subjects and Terms,
with their Pronunciation, Accentuation, and Derivation. Twenty-
second Edition. By RICHARD J. DUNGLISON, A.M., M.D. Royal 8vo,
30s.

Terminologia Medica Polyglotta: a Concise
International Dictionary of Medical Terms (French, Latin, English,
German, Italian, Spanish, and Russian). By THEODORE MAXWELL,
M.D., B.Sc., F.R.C.S. Edin. Royal 8vo, 16s.

A German-English Dictionary of Medical
Terms. By Sir FREDERICK TREVES, K.C.V.O., Surgeon to the London
Hospital; and HUGO LANG, B.A. Crown 8vo, half-Persian calf, 12s.

A Handbook of Physics and Chemistry, adapted
to the requirements of the first examination of the Conjoint Board
and for general use. By HERBERT E. CORBIN, B.Sc. Lond., and ARCHI-
BALD M. STEWART, B.Sc. Lond. With 120 Illustrations, crown 8vo.,
6s. 6d.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

A Manual of Chemistry, Theoretical and Practical. By WILLIAM A. TILDEN, D.Sc., F.R.S., Professor of Chemistry in the Royal College of Science, London; Examiner in Chemistry to the Department of Science and Art. With 2 Plates and 143 Woodcuts, crown 8vo, 10s.

Chemistry, Inorganic and Organic. With Experiments. By CHARLES L. BLOXAM. Eighth Edition, by JOHN MILLAR THOMSON, F.R.S., Professor of Chemistry in King's College, London, and ARTHUR G. BLOXAM, Head of the Chemistry Department, the Goldsmiths' Institute, New Cross. 8vo, with 281 Engravings, 18s. 6d.

By the same Author.

Laboratory Teaching; or, Progressive Exercises in Practical Chemistry. Sixth Edition, by ARTHUR G. BLOXAM. Crown 8vo, with 80 Engravings, 6s. 6d.

Watts' Organic Chemistry. Edited by William A. TILDEN, D.Sc., F.R.S., Professor of Chemistry, Royal College of Science, London. Second Edition. Crown 8vo, 10s.

Practical Chemistry, and Qualitative Analysis. By FRANK CLOWES, D.Sc. Lond., Emeritus Professor of Chemistry in the University College, Nottingham. Seventh Edition. Post 8vo, with 101 Engravings and Frontispiece, 8s. 6d.

Quantitative Analysis. By Frank Clowes, D.Sc. Lond., Emeritus Professor of Chemistry in the University College, Nottingham, and J. BERNARD COLEMAN, Assoc. R. C. Sci. Dublin; Professor of Chemistry, South-West London Polytechnic. Fifth Edition. Post 8vo, with 122 Engravings, 10s.

By the same Authors.

Elementary Practical Chemistry and Qualitative Analysis. Third Edition. With 68 Engravings, Post 8vo, 3s. 6d.

Also

Elementary Quantitative Analysis. With 62 Engravings, Post 8vo, 4s. 6d.

Qualitative Analysis. By R. Fresenius. Translated by CHARLES E. GROVES, F.R.S. Tenth Edition. 8vo, with Coloured Plate of Spectra and 46 Engravings, 15s.

By the same Author.

Quantitative Analysis. Seventh Edition.

VOL. I., Translated by A. VACHER. 8vo, with 106 Engravings, 15s.

VOL. II., Translated by C. E. GROVES, F.R.S. 8vo, with 143 Engravings, 20s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Inorganic Chemistry. By Sir Edward Frank-
LAND, K.C.B., D.C.L., LL.D., F.R.S., and FRANCIS R. JAPP, M.A.,
Ph.D., F.I.C., F.R.S., Professor of Chemistry in the University of
Aberdeen. 8vo, with numerous Illustrations on Stone and Wood, 24s.

Inorganic Chemistry (A System of). By
WILLIAM RAMSAY, Ph.D., F.R.S., Professor of Chemistry in the
University College, London. 8vo, with Engravings, 15s.

By the same Author.

Elementary Systematic Chemistry for the Use
of Schools and Colleges. With Engravings. Crown 8vo, 4s. 6d.
Interleaved, 5s. 6d.

Valentin's Practical Chemistry and Qualitative
and Quantitative Analysis. Edited by Dr. W. R. HODGKINSON,
F.R.S.E., Professor of Chemistry and Physics at the Royal Military
Academy, and Artillery College, Woolwich. Ninth Edition. 8vo, with
Engravings and Map of Spectra. 9s. (The Tables separately, 2s. 6d.)

Practical Chemistry, Part I. Qualitative Exer-
cises and Analytical Tables for Students. By J. CAMPBELL BROWN,
Professor of Chemistry in Victoria University and University College,
Liverpool. Fourth Edition. 8vo, 2s. 6d.

The Analyst's Laboratory Companion: a Col-
lection of Tables and Data for Chemists and Students. By ALFRED
E. JOHNSON, A.R.C.S.I., F.I.C. Second Edition. Crown 8vo, cloth,
5s.; leather, 6s. 6d.

Commercial Organic Analysis: a Treatise on
the Properties, Modes of Assaying, Proximate Analytical Examination,
etc., of the various Organic Chemicals and Products employed in the
Arts, Manufactures, Medicine, etc. By ALFRED H. ALLEN, F.I.C.

Third Edition.

VOL. I., 18s.; VOL. II., Part I., 14s.; VOL. II., Part
II., 14s.; VOL. II., Part III., ; VOL. III., Part I., 18s.

Second Edition.

VOL. III., Pt. II., 18s.; VOL. III., Pt. III., 16s.;
VOL. IV., completing the work, 18s.

Volumetric Analysis (A Systematic Hand-
book of); or the Quantitative Estimation of Chemical Substances by
Measure, applied to Liquids, Solids, and Gases. By FRANCIS SUTTON,
F.C.S., F.I.C., Public Analyst for the County of Norfolk. Eighth
Edition. 8vo, with 116 Engravings, 20s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Chemical Technology; or, Chemistry in its

Applications to Arts and Manufactures. Edited by CHARLES E. GROVES, F.R.S., and WILLIAM THORP, B.Sc.

VOL. I.—Fuel and its Applications. By E. J.

MILLS, D.Sc., F.R.S., and F. J. ROWAN, C.E. Royal 8vo, with 606 Engravings, 30s.

VOL. II.—Lighting, Fats and Oils, by W. Y.

DENT. STEARINE INDUSTRY, by J. McARTHUR. CANDLE MANUFACTURE, by L. FIELD and F. A. FIELD. THE PETROLEUM INDUSTRY AND LAMPS, by BOVERTON REDWOOD. MINERS' SAFETY LAMPS, by B. REDWOOD and D. A. LOUIS. Royal 8vo, with 358 Engravings and Map, 20s.

VOL. III.—Gas Lighting. By Charles Hunt.

With 2 Plates and 292 Engravings, 8vo, 18s.

Cooley's Cyclopædia of Practical Receipts,

and Collateral Information in the Arts, Manufactures, Professions, and Trades: including Medicine, Pharmacy, Hygiene, and Domestic Economy. Seventh Edition, by W. NORTH, M.A. Camb., F.C.S. Vols., Roy. 8vo, with 371 Engravings, 42s.

Chemical Technology: a Manual. By Rudolf

VON WAGNER. Translated and Edited by Sir WILLIAM CROOKES, F.R.S., from the Thirteenth Enlarged German Edition as remodelled by Dr. FERDINAND FISCHER. 8vo, with 596 Engravings, 32s.

Technological Handbooks. Edited by John

GARDNER, F.I.C., F.C.S., and JAMES CAMERON, F.I.C.

Brewing, Distilling, and Wine Manu-

facture. Crown 8vo, with Engravings, 6s. 6d.

Oils, Resins, and Varnishes. Crown 8vo,

with Engravings, 7s. 6d.

Soaps and Candles. Crown 8vo, with 54 En-

gravings, 7s.

Chemistry an Exact Mechanical Philosophy.

By FRED. G. EDWARDS. Illustrated, 8vo, 3s. 6d.

The Quarterly Journal of Microscopical Science.

Edited by E. RAY LANKESTER, M.A., LL.D., F.R.S.; with the co-operation of ADAM SEDGWICK, M.A., F.R.S., W.F. R. WELDON, M.A., F.R.S., and SYDNEY J. HICKSON, M.A., F.R.S. Each Number, 10s.

7, GREAT MARLBOROUGH STREET.

J. & A. Churchill's Recent Works.

Methods and Formulæ used in the Preparation

of Animal and Vegetable Tissues for Microscopical Examination, including the Staining of Bacteria. By PETER WYATT SQUIRE, F.L.S. Crown 8vo, 3s. 6d.

The Microscope and its Revelations. By the

late WILLIAM B. CARPENTER, C.B., M.D., LL.D., F.R.S. Eighth Edition, by the Rev. W. H. DALLINGER, LL.D., F.R.S. With 23 Plates and more than 800 Wood Engravings. 8vo, 28s. Half Calf, 32s.

The Microtometist's Vade-Mecum: a Handbook

of the Methods of Microscopic Anatomy. By ARTHUR BOLLES LEE. Fifth Edition, 8vo, 15s.

Photo-Micrography (Guide to the Science of).

By EDWARD C. BOUSFIELD, L.R.C.P. Lond. 8vo, with 34 Engravings and Frontispiece, 6s.

A Treatise on Physics. By Andrew Gray,

LL.D., F.R.S., Professor of Natural Philosophy in the University of Glasgow. Vol. I. Dynamics and Properties of Matter. With 350 Illustrations, 8vo. 15s.

An Introduction to Physical Measurements,

with Appendices on Absolute Electrical Measurements, etc. By Dr. F. KOHLRAUSCH. Third English Edition, by T. H. WALLER, B.A., B.Sc., and H. R. PROCTER, F.I.C., F.C.S. 8vo, with 91 Illustrations, 12s. 6d.

Tuson's Veterinary Pharmacopœia, including

the Outlines of Materia Medica and Therapeutics. Fifth Edition Edited by JAMES BAYNE, F.C.S., Professor of Chemistry and Toxicology in the Royal Veterinary College. Crown 8vo, 7s. 6d.

The Veterinarian's Pocket Remembrancer:

being Concise Directions for the Treatment of Urgent or Rare Cases, By GEORGE ARMATAGE, M.R.C.V.S. Second Edition. Post 8vo, 3s.

Chauveau's Comparative Anatomy of the

Domesticated Animals. Revised and Enlarged, with the Co-operation of S. ARLOING, Director of the Lyons Veterinary School, and Edited by GEORGE FLEMING, C.B., LL.D., F.R.C.V.S., late Principal Veterinary Surgeon of the British Army. Second English Edition. 8vo with 585 Engravings, 31s. 6d.

Human Nature, its Principles and the Principles

of Physiognomy. By PHYSICIST. Part I., Imp. 16mo, 2s. Part II. completing the work), 2s. 6d.

Encyclopædia Medica. Edited by Chalmers

Watson, M.B., M.R.C.P.E. In about 12 Volumes, 20s. each net. Vols. I. to VIII. now ready.

7, GREAT MARLBOROUGH STREET.

INDEX TO J. & A. CHURCHILL'S CATALOGUE.

- Allen's Chemistry of Urine, 22
 ——— Commercial Organic Analysis, 26
 Armatage's Veterinary Pocket Remembrancer, 28
 Auld's Researches in Pathology, 2
 Barnes' (R.) Obstetric Operations, 6
 ——— Diseases of Women, 6
 Beale (L. S.) on Liver, 12
 ——— Slight Ailments, 12
 ——— Urinary and Renal Derangements, 22
 Beale (P. T. B.) on Elementary Biology, 3
 Beasley's Book of Prescriptions, 8
 ——— Druggists' General Receipt Book, 8
 ——— Pharmaceutical Formulary, 8
 Bell on Sterility, 6
 Bentley and Trimen's Medicinal Plants, 9
 Bentley's Systematic Botany, 9
 Berkart's Bronchial Asthma, 13
 Bernard on Stammering, 14
 Berry's (Jas.) Thyroid Gland, 18
 ——— (R. J.) Regional Anatomy, 2
 Bigg's Short Manual of Orthopædy, 18
 Birch's Practical Physiology, 3
 Bloxam's Chemistry, 25
 ——— Laboratory Teaching, 25
 Bousfield's Photo-Micrography, 28
 Bowlby's Injuries and Diseases of Nerves, 17
 ——— Surgical Pathology and Morbid Anatomy, 17
 Brockbank on Gallstones, 15
 Brown's (Haydn) Midwifery, 6
 ——— Ringworm, 22
 Brown's Practical Chemistry, 26
 Bryant's Practice of Surgery, 16
 Bulkley on Skin, 21
 Burdett's Hospitals and Asylums of the World, 4
 Butler-Smythe's Ovariectomies, 6
 Butlin's Operative Surgery of Malignant Disease, 21
 ——— Sarcoma and Carcinoma, 21
 ——— Malignant Disease of the Larynx, 21
 Buzzard's Diseases of the Nervous System, 14
 Buzzard's Peripheral Neuritis, 14
 ——— Simulation of Hysteria, 14
 Cameron's Oils, Resins, and Varnishes, 27
 ——— Soaps and Candles, 27
 Campbell's Dissection Outlines, 1
 Carpenter and Dallinger on the Microscope, 28
 Cautley on Feeding Infants, 7
 Charteris' Practice of Medicine, 11
 Chauveau's Comparative Anatomy, 28
 Chevers' Diseases of India, 10
 Churchill's Face and Foot Deformities, 18
 Clouston's Lectures on Mental Diseases, 4
 Clowes and Coleman's Quantitative Analysis, 25
 Clowes and Coleman's Elementary Practical Chemistry, 25
 Clowes' Practical Chemistry, 25
 Coles on Blood, 12
 Cooley's Cyclopædia of Practical Receipts, 27
 Cooper's Syphilis, 23
 Cooper and Edwards' Diseases of the Rectum, 24
 Corbin and Stewart's Physics and Chemistry, 24
 Cripps' (H.) Ovariectomy and Abdominal Surgery, 17
 ——— Diseases of the Rectum and Anus, 24
 ——— Cancer of Rectum, 24
 ——— Air and Faces in Urethra, 24
 Cripps' (R. A.) Galenic Pharmacy, 8
 Cuff's Lectures to Nurses, 7
 Cullingworth's Monthly Nurses, 7
 Dalby's Diseases and Injuries of the Ear, 20
 ——— Short Contributions, 20
 Dana on Nervous Diseases, 14
 Day on Headaches, 15
 Domville's Manual for Nurses, 7
 Druitt's Surgeon's Vade-Mecum, 16
 Duncan (A.) on Prevention of Diseases in Tropics, 10
 Dunlison's Med. Dictionary, 24
 Edwards' Chemistry, 27
 Ellis's (T. S.) Human Foot, 18
 [Continued on next page]

7, GREAT MARLBOROUGH STREET.

INDEX TO J. & A. CHURCHILL'S CATALOGUE—continued.

- Encyclopædia Medica, 28
 Fagge's Principles and Practice of Medicine, 10
 Fayrer's Climate and Fevers of India, 10
 Fenwick (E. H.), Electric Illumination of Bladder, 22
 ——— Atlas of Electric Cystoscopy, 22
 ——— Symptoms of Urinary Disease, 22
 ——— Tumours of Bladder, 22
 ——— Ulceration of Bladder, 22
 Fenwick's (S.) Medical Diagnosis, 12
 ——— Ulcer of Stomach, 12
 ——— Obscure Diseases of the Abdomen, 12
 ——— Outlines of Medical Treatment, 12
 ——— The Saliva as a Test, 12
 Fink's Operating for Cataract, 19
 Fowler's Dictionary of Practical Medicine, 11
 Fox (G. H.) on Skin Diseases of Children, 21
 Fox (Wilson), Atlas of Pathological Anatomy of the Lungs, 11
 — Treatise on Diseases of the Lungs, 11
 Frankland and Japp's Inorganic Chemistry, 26
 Fraser's Operations on the Brain, 16
 Fresenius' Qualitative Analysis, 25
 ——— Quantitative Analysis, 25
 Galabin's Diseases of Women, 6
 ——— Manual of Midwifery, 5
 Gardner's Brewing, Distilling, and Wine Manufacture, 27
 Glassington's Dental Materia Medica, 20
 Godlee's Atlas of Human Anatomy, 1
 Goodhart's Diseases of Children, 7
 Gorgas' Dental Medicine, 20
 Gowers' Diagnosis of Brain Disease, 13
 ——— Diseases of Nervous System, 13
 ——— Medical Ophthalmoscopy, 13
 ——— Syphilis and the Nervous System, 13
 ——— Epilepsy, 13
 Granville on Gout, 14
 Gray's Treatise on Physics, 28
 Green's Manual of Botany, 9
 ——— Vegetable Physiology, 9
 Greenish's Materia Medica, 8
 Groves and Thorp's Chemical Technology, 27
 Guy's Hospital Reports, 11
 Habershon's Diseases of the Abdomen, 15
 Haig's Uric Acid, 13
 ——— Diet and Food, 4
 Harley on Diseases of the Liver, 14
 Harris's (V. D.) Diseases of Chest, 11
 Harrison's Urinary Organs, 23
 Hartridge's Refraction of the Eye, 19
 ——— Ophthalmoscope, 19
 Hawthorne's Galenical Preparations, 8
 Heath's Injuries and Diseases of the Jaws, 17
 ——— Minor Surgery and Bandaging, 17
 ——— Operative Surgery, 17
 ——— Practical Anatomy, 1
 ——— Surgical Diagnosis, 17
 Hedley's Therapeutic Electricity, 9
 Hellier's Notes on Gynæcological Nursing, 7
 Hewlett's Bacteriology, 4
 Hill on Cerebral Circulation, 3
 Holden's Human Osteology, 1
 ——— Landmarks, 1
 Holthouse on Strabismus, 18
 Hooper's Physicians' Vade Mecum, 10
 Horrocks' Bacteriological Examination of Water, 4
 Horton-Smith on Typhoid, 14
 Hovell's Diseases of the Ear, 19
 Hughes and Keith's Practical Anatomy, 2
 Human Nature and Physiognomy, 28
 Hyslop's Mental Physiology, 5
 Impey on Leprosy, 21
 Ireland's Mental Affections of Children, 5
 Jacobson's Male Organs, 22
 ——— Operations of Surgery, 16
 Jellett's Midwifery, 5
 ——— Gynæcology, 6
 Jessop's Ophthalmic Surgery and Medicine, 18
 Johnson's (Sir G.) Asphyxia, 12
 ——— Medical Lectures and Essays, 12
 ——— (A. E.) Analyst's Companion, 26
 Journal of Mental Science, 5

[Continued on next page.]

7, GREAT MARLBOROUGH STREET.

INDEX TO J. & A. CHURCHILL'S CATALOGUE—continued.

- Kelynack's Pathologist's Handbook, 2
 Keyes' Genito-Urinary Organs and Syphilis, 23
 Kohlrausch's Physical Measurements, 28
 Lane's Rheumatic Diseases, 14
 Langdon-Down's Mental Affections of Childhood, 5
 Lawrie on Chloroform, 17
 Lazarus-Barlow's General Pathology, 2
 Lee's Microtometist's Vade-Mecum, 28
 Lewis (Bevan) on the Human Brain, 2
 Liebreich (O.) on Borax and Boracic Acid, 4
 Liebreich's (R.) Atlas of Ophthalmoscopy, 19
 Lucas's Practical Pharmacy, 8
 MacMunn's Clinical Chemistry of Urine, 22
 Macnamara's Diseases and Refraction of the Eye, 18
 ——— Diseases of Bones and Joints, 17
 McNeill's Isolation Hospitals, 4
 Malcolm's Physiology of Death, 16
 Martin's Ambulance Lectures, 15
 Maxwell's Terminologia Medica Polyglotta, 24
 Maylard's Surgery of Alimentary Canal, 16
 Mayne's Medical Vocabulary, 24
 Microscopical Journal, 27
 Mills and Rowan's Fuel and its Applications, 27
 Moore's (N.) Pathological Anatomy of Diseases, 2
 Moore's (Sir W. J.) Diseases of India, 10
 ——— Family Medicine, etc., for India, 10
 Morris's Human Anatomy, 1
 ——— Anatomy of Joints, 2
 Moullin's (Mansell) Surgery, 16
 Nettleship's Diseases of the Eye, 18
 Notter's Hygiene, 3
 Ogle on Tympanites, 15
 Oliver's Abdominal Tumours, 6
 ——— Diseases of Women, 6
 Ophthalmic (Royal London) Hospital Reports, 18
 Ophthalmological Society's Transactions, 18
 Ormerod's Diseases of the Nervous System, 13
 Owen's (J.) Diseases of Women, 6
 Parkes' (E. A.) Practical Hygiene, 3
 Parkes' (L. C.) Elements of Health, Pavy's Carbohydrates, 12
 Pereira's Selecta à Prescriptis, 9
 Phillips' Materia Medica and Therapeutics, 8
 Pitt-Lewis's Insane and the Law, 5
 Pollock's Histology of the Eye and Eyelids, 19
 Proctor's Practical Pharmacy, 8
 Pye-Smith's Diseases of the Skin, 21
 Ramsay's Elementary Systematic Chemistry, 26
 ——— Inorganic Chemistry, 26
 Richardson's Mechanical Dentistry, 20
 Richmond on Antiseptics, 7
 Roberts' (C. Hubert) Gynaecological Pathology, 6
 Roberts' (D. Lloyd), Practice of Midwifery, 5
 Robinson's (Tom) Eczema, 21
 ——— Illustrations of Skin Diseases, 21
 ——— Syphilis, 21
 Ross's Diseases of the Nervous System, 14
 St. Thomas's Hospital Reports, 11
 Sansom's Valvular Disease of the Heart, 13
 Scott's Atlas of Urinary Deposits, 23
 Shaw's Diseases of the Eye, 19
 Shaw-Mackenzie on Maternal Syphilis, 24
 Short Dictionary of Medical Terms, 24
 Silk's Manual of Nitrous Oxide, 20
 Smith's (Ernest), Dental Metallurgy, 20
 ——— (Eustace) Clinical Studies, 7
 ——— Disease in Children, 7
 ——— Wasting Diseases of Infants and Children, 7
 ——— (Fred. J.) Medical Jurisprudence, 3
 Smith's (J. Greig) Abdominal Surgery, 16
 Smith's (Priestley) Glaucoma, 19
 Suow's Cancers and the Cancer Process, 21
 ——— Palliative Treatment of Cancer, 21
 ——— Reappearance of Cancer, 21
 [Continued on next page]

7, GREAT MARLBOROUGH STREET.


INDEX TO J. & A. CHURCHILL'S CATALOGUE—continued.

- | | |
|--|--|
| Solly's Medical Climatology, 15 | Tirard's Medical Treatment, 12 |
| Southall's Organic Materia Medica, 7 | Tobin's Synopsis of Surgery, 15 |
| Squire's (P.) Companion to the Pharmacopœia, 8 | Tomes' (C. S.) Dental Anatomy, 20 |
| ——— London Hospitals Pharmacopœias, 8 | ——— (J. & C. S.) Dental Surgery, 20 |
| ——— Methods and Formulæ 28 | Tooth's Spinal Cord, 14 |
| Starling's Elements of Human Physiology, 3 | Treves and Lang's German-English Dictionary, 24 |
| Sternberg's Bacteriology, 11 | Tuke's Dictionary of Psychological Medicine, 5 |
| Stevenson and Murphy's Hygiene, 4 | Tuson's Veterinary Pharmacopœia, 28 |
| Sutton's (F.) Volumetric Analysis, 26 | Valentin and Hodgkinson's Practical Chemistry, 26 |
| Sutton's (J. B.) General Pathology, 2 | Vintras on the Mineral Waters, etc., of France, 15 |
| Swain's Surgical Emergencies, 15 | Wagner's Chemical Technology, 27 |
| Swayne's Obstetric Aphorisms, 6 | Wallace on Dental Caries, 20 |
| Taylor's (A. S.) Medical Jurisprudence, 3 | Walsham's Surgery: its Theory and Practice, 15 |
| Taylor's (F.) Practice of Medicine, 10 | Waring's Indian Bazaar Medicines, 9 |
| Thin's Cancerous Affections of the Skin, 22 | ——— Practical Therapeutics, 9 |
| ——— Pathology and Treatment of Ringworm, 22 | Watts' Organic Chemistry, 25 |
| ——— Psilosis or "Sprue," 10 | West's (S.) How to Examine the Chest, 11 |
| Thompson's (Sir H.) Calculous Diseases, 23 | White's (Hale) Materia Medica, Pharmacy, etc., 7 |
| ——— Diseases of the Urinary Organs, 23 | Wilks' Diseases of the Nervous System, 13 |
| ——— Lithotomy and Lithotripsy, 23 | Wilson's (Sir E.) Anatomist's Vademecum, 1 |
| ——— Stricture of the Urethra, 23 | Wilson's (G.) Handbook of Hygiene, 3 |
| ——— Suprapubic Operation, 23 | Wolfe's Diseases and Injuries of the Eye, 18 |
| ——— Surgery of the Urinary Organs, 23 | Wynter and Wethered's Practical Pathology, 2 |
| ——— Tumours of the Bladder, 23 | Year Book of Pharmacy, 9 |
| Thorne's Diseases of the Heart, 11 | Yeo's (G. F.) Manual of Physiology, 3 |
| Thresh on Water Analysis, 4 | |
| Tilden's Chemistry, 25 | |

N.B.—J. & A. Churchill's larger Catalogue of about 600 works on Anatomy, Physiology, Hygiene, Midwifery, Materia Medica, Medicine, Surgery, Chemistry, Botany, etc. etc., with a complete Index to their Subjects, for easy reference, will be forwarded post free on application.


AMERICA.—J. & A. Churchill being in constant communication with various publishing houses in America are able to conduct negotiations favourable to English Authors.

LONDON: 7, GREAT MARLBOROUGH STREET.



72 m of Holland

XV Messers



F. G. m. s. to. Holland

10th "The Kings" Hussars

