**A pharmacopoeia including the outlines of materia medica and therapeutics for the use of practitioners and students of veterinary medicine / by the late Richard V. Tuson.**


**Contributors**

Tuson, Richard V.
Bayne, James.

**Publication/Creation**

London : J. & A. Churchill, 1904.

**Persistent URL**

https://wellcomecollection.org/works/k6ew5dfb

**License and attribution**

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).

wellcome collection

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
https://wellcomecollection.org

155 1



Med
K53602

# VETERINARY PHARMACOPŒIA

3/8/04

A

# PHARMACOPŒIA

INCLUDING THE OUTLINES OF

## MATERIA MEDICA AND THERAPEUTICS

FOR THE USE OF

PRACTITIONERS AND STUDENTS

OF

## Veterinary Medicine

BY THE LATE RICHARD V. TUSON

SIXTH EDITION

REVISED AND EDITED BY

JAMES BAYNE, F.I.C., F.C.S.

LATE PROFESSOR OF CHEMISTRY AND TOXICOLOGY AT THE
ROYAL VETERINARY COLLEGE

LONDON

J. & A. CHURCHILL

7 GREAT MARLBOROUGH STREET

1904

117 041

PRINTED BY
SPOTTISWOODE AND CO. LTD., NEW-STREET SQUARE
LONDON



# PREFACE

TO

# THE SIXTH EDITION

In this, as in the Fifth Edition, the original arrangement is retained. Some new preparations, which are now in general use by veterinary practitioners, have been added. The whole work has been carefully revised, and some alterations have been made to bring it as nearly as possible into line with the 'British Pharmacopœia,' 1898.

In addition to this, reference has been made to Shoemaker's 'Materia Medica and Therapeutics,' Gresswell's 'Veterinary Pharmacopœia,' and Winslow's 'Veterinary Materia Medica and Therapeutics.'

I have especially to express my indebtedness to my former colleague, Professor Macqueen, for the valuable assistance I have derived from the Tenth Edition of Finlay-Dun's 'Veterinary Medicines' (edited by Professor Macqueen) in the correction and bringing up to date the

actions, uses, and doses of medicines. My thanks are also due to my partner, Mr. Hubert E. Lindley, F.C.S., for assistance in the compilation of the work; and to my assistant, Mr. L. Elliott, for help in the correction of proofs.

JAMES BAYNE

THE EUSTON LABORATORY,
4 EUSTON BUILDINGS, LONDON, N.W.:
*July*, 1904

# PREFACE TO THE FIRST EDITION

This work is designed for the use of practitioners and students of veterinary medicine.

The agents are arranged in alphabetical order, according to the plan adopted in the 'British Pharmacopœia,' and are treated of under the following heads:

1. Latin Pharmaceutic Name.
2. English Pharmaceutic Name.
3. Synonyms.
4. Natural Order.
5. Composition.
6. Mode of Preparation.
7. Characters and Tests.
8. Actions and Uses.
9. Doses.
10. Modes of Application.
11. Incompatibles.
12. Antidotes.
13. Preparations.

Many medicines bearing old and familiar names have, for the sake of order, received new titles; thus, the 'Diuretic Mass' of Morton is, in the present work, termed Massa Resinæ Composita. By whatever name, however, preparations may be known to the reader, they can be readily found by reference to the General Index.

The modes of preparation, characters, and tests are in most cases the same as those given in the 'British Pharmacopœia.'

The composition of bodies is expressed in modern atomic weights, a table of which, together with a list of the old ones, will be found in the Appendix.

In the directions for preparing chemico-pharmaceutic substances old chemical names are, for the most part, employed; whereas, in explaining reactions and decompositions, that which the author believes to be the best of the new systems of nomenclature, supplemented where necessary by the old, is adopted.

The incompatibles mentioned are those recorded as such by the most eminent authorities; although, no doubt, in many instances the incompatibility indicated is based upon hypothesis rather than upon the results of observation or experiment.

For a more detailed account of the actions and uses of veterinary medicines the reader is referred to the books of Dun and Morton, also to the forthcoming work on 'Veterinary Therapeutics' by Professor George Brown.

To the last-named gentleman the author is deeply indebted for his kind assistance in revising those portions of the work which are comprised under the sub-headings 'Doses' and 'Actions and Uses,' as well as for many valuable suggestions made during the progress of the volume through the press.

The principal sources, in addition to the 'British Pharmacopœia' of 1867, from which the author has obtained much valuable information, are Morton's 'Veterinary Pharmacy,' Dun's 'Veterinary Medicines,' Macnamara's edition of 'Neligan's Medicines,' and Garrod's 'Materia Medica and Therapeutics.'

ROYAL VETERINARY COLLEGE:
*October* 1869

# CONTENTS


PAGE

PREFACE TO THE SIXTH EDITION . . . . . v

PREFACE TO THE FIRST EDITION . . . . . vii

VETERINARY PHARMACOPŒIA . . . . . 1

ADDENDA . . . . . 331

APPENDIX :

1. SYMBOLS AND ATOMIC WEIGHTS OF THE ELEMENTARY BODIES MENTIONED IN THIS WORK . . 337
2. WEIGHTS AND MEASURES OF THE 'BRITISH PHARMACOPŒIA' . . . . . 338
3. WEIGHTS AND MEASURES OF THE METRICAL SYSTEM . . . . . 339
4. RELATION OF THE WEIGHTS AND MEASURES OF THE 'BRITISH PHARMACOPŒIA' TO THOSE OF THE METRICAL SYSTEM . . . . . 339–340

INDEX TO VETERINARY MEDICINES, ARRANGED ACCORDING TO THEIR ACTIONS AND USES . . . . 341

GENERAL INDEX . . . . . 355

A

# VETERINARY PHARMACOPŒIA

## ACACIA GUMMI

GUM ACACIA

**Synomyn.**—Gum Arabic.

**Composition.**—An exudation from *Acacia Senegal* and other species of *Acacia*, and consisting, when pure, of arabin, $C_{12}H_{22}O_{11}$.

**Natural Order.**—*Leguminosæ.*

**Characters and Tests.**—Spheroidal fragments, with shining surfaces; nearly colourless; bland and mucilaginous taste; soluble in water, insoluble in alcohol, ether, and oils. Its cold aqueous solution gives a white gelatinous precipitate with lead subacetate, but no blue colour with iodine-water, unless it be adulterated with starch.

**Actions and Uses.**—See Mucilago Acaciæ.

## ACETANILIDUM

PHENYL ACETAMIDE

**Synonym.**—Antifebrin.

**Composition.**—$CH_3.CO.NH.C_6H_5$.

**Characters and Tests.**—A colourless, inodorous, and glistening solid, in lamellar crystals, slightly pungent,

slightly soluble in water, and freely soluble in ether, alcohol, benzol, and chloroform. Not changed by acids or alkalies.

**Actions and Uses.**—Antipyretic, diuretic, anti-spasmodic, anodyne. Used to relieve pain in nervous affections and rheumatism.

**Doses.**—For HORSES or CATTLE, from 30 grains to 2 drachms.—DOGS, 2 to 10 grains may be given in powder, pill, or electuary, or in water containing alcohol. These doses repeated every four hours, according to Major J. A. Nunn, Army Veterinary Department, produced marked lowering of temperature in fever, unaccompanied by nausea or ether untoward effects.

## ACETUM

VINEGAR

**Composition.**—Water holding in solution about 5·41 per cent. of real acetic acid, $HC_2H_3O_2$, together with colouring and flavouring matters.

**Mode of Preparation.**—Weak alcoholic liquids—*e.g.*, beer, poor wines, &c.—are exposed to the air and suffered to undergo the acetous fermentation, whereby the alcohol is oxidised first into aldehyd, and finally into acetic acid, thus :—

1. $\overset{\text{Alcohol.}}{2C_2H_6O} + \overset{\text{Oxygen.}}{O_2} = \overset{\text{Aldehyd.}}{2C_2H_4O} + \overset{\text{Water.}}{2H_2O}$

2. $\overset{\text{Aldehyd.}}{2C_2H_4O} + \overset{\text{Oxygen.}}{O_2} = \overset{\text{Acetic Acid.}}{2HC_2H_3O_2}$

Also by the destructive distillation of wood.

**Characters and Tests.**—A liquid of a brown colour, acetous odour, and sour taste. Sp. gr. 1·017 to 1·019. With barium chloride it should give no, or a very slight,

precipitate, showing the absence, or the presence of only a minute quantity, of sulphuric acid.[1]

**Actions and Uses.**—*Internally* :—Refrigerant to allay febrile heat ; styptic, astringent, and diuretic ; also litholytic for calculi and urinary deposits consisting of calcium or magnesium phosphate or carbonate. Likewise employed as an antidote to poisoning by the alkalies and their carbonates. Very seldom, however, used internally.

**Doses.**—HORSE, 1 to 2 fluid ounces.—CATTLE, 1 to 2 fluid ounces.—SHEEP, 1 to 2 fluid drachms.—PIG, 1 to 2 fluid drachms.—DOG, 10 to 20 minims.

**Incompatibles.**—Alkalies, earths, and all carbonates.

**Antidotes.**—Potassium, sodium, calcium, or magnesium carbonate or bicarbonate.

## ACETUM CANTHARIDIS

### VINEGAR OF CANTHARIDES

**Composition.**—An impure solution of cantharidin, $C_5H_6O_2$, in acetic acid.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Cantharides in powder . . | 2 ounces |
| Glacial Acetic Acid . . . | 2 fluid ounces |
| Acetic Acid . . . | 18 fluid ounces, or a sufficiency |

Mix thirteen fluid ounces of the acetic acid with the glacial acetic acid, and digest the cantharides in this mixture for two hours at a temperature of 200° F. Cool the ingredients, transfer them to a percolator, and when the

[1] The addition of 1 part of sulphuric acid to 999 of vinegar is permitted by law.

liquid ceases to flow, pour five fluid ounces of acetic acid over the residue in the apparatus. When the percolation is complete, press the contents of the percolator, filter the product, mix the liquids, and add sufficient acetic acid to make one pint.

**Actions and Uses.**—*Externally* :—Counter-irritant. Vesicant and rubefacient in local congestion, and in irritation of important organs.

## ACIDUM ACETICUM

ACETIC ACID

**Composition.**—Water mixed with 33 per cent. of real acetic acid, $HC_2H_3O_2$.

**Mode of Preparation.**—By distilling a mixture of crystallised sodium acetate and sulphuric acid ; sodium sulphate remains in the retort, while water and acetic acid, in the proportions above named, pass over into the receiver.

| Sodium Acetate. | | Sulphuric Acid. | | Sodium Sulphate. | | Acetic Acid. |
|---|---|---|---|---|---|---|
| $2NaC_2H_3O_2$ | + | $H_2SO_4$ | = | $Na_2SO_4$ | + | $2HC_2H_3O_2$ |

Also by the destructive distillation of wood (pyroligneous acid).

**Characters and Tests.**—A colourless, sour liquid, having a pungent odour. Reddens litmus-paper and gives a red colour with neutral ferric chloride; evolves odour of acetic ether when warmed with alcohol. Sp. gr. 1·044. Should leave no residue when evaporated, and will give no precipitate with hydrosulphuric acid, barium chloride, or silver nitrate if free from lead, copper, sulphuric acid, and chlorine. Put a fluid drachm of the acid, mixed with half an ounce of distilled water and half a drachm of pure hydrochloric acid, also a few pieces of granulated zinc,

into a flask. While effervescence continues suspend a slip of white blotting-paper moistened with a solution of sub-acetate of lead in the upper part of the flask above the liquid for about five minutes. The paper should not become discoloured, thus indicating the absence of sulphurous acid.

**Actions and Uses.**—*Externally* :—Stimulant, astringent, styptic, rubefacient, and vesicant. As a caustic for the removal of warts and similar growths. As it destroys cryptogamic parasites and acari, it is used for the removal of scurf in ringworm, mallenders and sallenders, scab, and mange.[1]

**Preparations.**—Acetum Cantharidis; Acidum Aceticum Dilutum.

## ACIDUM ACETICUM DILUTUM

DILUTE ACETIC ACID

**Composition.**—Water mixed with 4·27 per cent. of real acetic acid, $HC_2H_3O_2$ (hydrogen acetate).

**Mode of Preparation.**—Take of

| | |
|---|---|
| Acetic Acid . . . . . | 1 pint |
| Distilled Water . . . . | 7 pints |

Mix.

**Characters and Tests.**—Colourless, sour liquid. Sp. gr. 1·006.

| | |
|---|---|
| **Actions and Uses.** | Similar to vinegar. |
| **Doses.** | Similar to vinegar. |
| **Incompatibles.** | See Acetum. |
| **Antidotes.** | See Acetum. |

[1] For these purposes, and as an astringent and styptic, impure pyroligneous acid is preferred to acidum aceticum, on account of the former containing carbolic acid and similar compounds.

## ACIDUM ACETICUM GLACIALE

GLACIAL ACETIC ACID

**Synonyms.**—Acetic Hydrate; Hydrogen Acetate.

**Composition.**—$HC_2H_3O_2$. Contains 99 parts of hydrogen acetate.

**Mode of Preparation.**—By distilling a mixture of concentrated sulphuric acid and sodium acetate deprived of its water of crystallisation by fusion; sodium sulphate remains in the retort, and glacial acetic acid passes over into the receiver. See Acidum Aceticum for an explanation of the decomposition.

**Characters and Tests.**—At ordinary temperatures a colourless liquid, with a pungent odour and sour taste. Sp. gr. 1·065 to 1·066. Cooled to 34° F. it crystallises and remains solid unless heated to 48° F. Tested for sulphurous acid, as directed under Acidum Aceticum, it should give no indication of this impurity.

**Actions and Uses.**—*Internally*:—Irritant poison. *Externally*:—Rubefacient, vesicant, and caustic. Used to destroy warts.

**Antidotes.**—See Acetum.

**Preparations.**—Acetum Cantharidis; Mistura Creosoti.

## ACIDUM ARSENIOSUM

ARSENIOUS ACID

**Synonyms.**—Arsenious Anhydride; White Arsenic.

**Composition.**—$As_2O_3$.

**Mode of Preparation.**—By roasting arsenical ores—*e.g.*, arsenical pyrites, FeSAs—in a current of air; the

arsenic combines with oxygen and becomes arsenious anhydride, which is purified by sublimation.

**Characters and Tests.**—Heavy, white, glistening powder or in translucent masses resembling porcelain. Completely volatile at 400° F. Sublimed in a test-tube, it condenses in brilliant octahedral or triangular crystals. Soluble in water, 1 in 80 parts. Its aqueous solution furnishes a yellow precipitate (silver arsenite) with silver ammonio-nitrate, and a green precipitate (copper arsenite) with copper ammonio-sulphate. These precipitates are soluble in ammonium hydrate and in nitric acid.

**Actions and Uses.**—*Internally* :—In excessive doses, an irritant poison; in medicinal doses, alterative and tonic in chronic rheumatism, paralysis, epilepsy, chorea, farcy, and mange. *Externally* :—Used in the cure of mange, scab, and other skin-diseases; also as a caustic to eradicate warts, and to produce the sloughing and effect the removal of malignant tumours. When used externally, especially for the latter purpose, arsenious acid must be employed with great caution, as it is liable to be absorbed and to kill the patient.

**Doses.**—Cattle, 2 to 10 grains.—Horse, 2 to 10 grains.—Sheep, 1 to 2 grains.—Pig, $\frac{1}{2}$ to 2 grains.—Dog, $\frac{1}{15}$ to $\frac{1}{10}$ grain.

**Modes of Application.**—*Internally* :—The powder sprinkled over, or mixed with, the animal's food.[1] *Externally* :—Liquor Arsenicalis; Unguentum Arsenici.

**Antidotes.**—Freshly made moist hydrated peroxide of iron; magnesia; cold affusions; inhalation of ammonia.

**Preparations.** — Liquor Arsenicalis; Unguentum Arsenici.

[1] See Liquor Arsenicalis, a better form of arsenical medicine for internal use.

## ACIDUM BORICUM

BORIC ACID

**Synonyms.**—Boracic Acid ; Hydrogen Borate.

**Composition.**—$H_3BO_3$.

**Mode of Preparation.**—By purifying the acid found naturally in the volcanic districts of North Italy. See Borax.

**Characters and Tests.**—A transparent, colourless solid which crystallises in six-sided plates. It has a feebly acid taste. It dissolves in 25 parts of water or 15 of alcohol, and in 3 parts of boiling water or 5 of boiling alcohol. An alcoholic solution burns with a green flame.

**Actions and Uses.**—*Externally* :—Antiseptic, used as a dressing for wounds. The saturated solution is used in the treatment of scaly and parasitic skin-eruptions.

**Preparations.**—Boroglyceride, a combination of boric acid (62 parts) and glycerin (92 parts). This diluted with an equal quantity of glycerin makes the 50 per cent. boroglyceride, in which form it is used as an antiseptic.

Lister's antiseptic ointment for dressing wounds is made by adding 1 part each of boric acid and white wax to 2 parts each of paraffin and almond oil. Borated lint or borated cotton-wool is made by steeping the substance in a saturated solution of the acid and allowing it to dry.

## ACIDUM CARBOLICUM

CARBOLIC ACID

**Synonym.**—Phenol.

**Composition.**—$C_6H_5OH$.

**Mode of Preparation.**—Those portions of coal-tar oil which distill between 300° and 400° F. are mixed

with a strong and boiling solution of potassium hydrate. The resulting potassium phenol is decomposed by hydrochloric acid, and the liberated carbolic acid rectified by distillation off calcium chloride. The calcium chloride deprives the acid of moisture.

| | Carbolic Acid. | | Potassium Hydrate. | | Potassium Phenol | | Water. |
|---|---|---|---|---|---|---|---|
| 1. | $C_6H_5OH$ | + | KHO | = | $KC_6H_5O$ | + | $H_2O$ |

| | Potassium Phenol. | | Hydrochloric Acid. | | Carbolic Acid. | | Potassium Chloride. |
|---|---|---|---|---|---|---|---|
| 2. | $KC_6H_5O$ | + | HCl | = | $HC_6H_5O$ | + | KCl |

**Characters and Tests.**—Acicular crystals which, when freshly prepared and quite pure, are colourless, but exposed to moist air acquire a pink tinge. At 95° F. they become an oily liquid, having an odour and taste like creosote. Sp. gr. 1·065 ; boiling-point, 370° F. Exposed to the air, the crystals absorb moisture and liquefy. The acid is slightly soluble in water, 1 in 15 parts, but freely soluble in glycerin, alcohol, and ether. About a grain of hypochlorite of calcium added to a little aqueous solution of carbolic acid, placed in a test-tube, produces, after agitation, the addition of a few drops of ammonia, and the application of a gentle heat, a bright blue colour with a tinge of green. An aqueous solution of carbolic acid will also give a purple colour with a few drops of ferric chloride. One drachm of the acid, if pure, completely dissolves on being shaken with half a pint of warm water. Carbolic acid is distinguished by its odour. Bromine-water in aqueous solution gives pale yellow needles of tribromophenol. According to Martindale ('ExtraPharmacopœia'), there are seven commercial varieties in general use : (1) Absolute Phenol. (2) No. 1 Carbolic Acid in crystals, and No. 1 Acid in liquid containing 10 per cent. of added

water; the crystals are soluble in 14 parts of water. (3) No. 2 Carbolic Acid, also in crystals, soluble in 18 parts of water, and No. 2 Acid in liquid. (4) No. 4 Carbolic Acid, a pale straw coloured liquid containing about 10 per cent. of phenol and nearly 90 per cent. of cresols. (5) No. 5 Carbolic Acid, a dark-coloured liquid of uncertain strength.

**Actions and Uses.**—*Internally*:—In excessive doses, narcotic and irritant poison; in medicinal doses it possesses antiseptic and antipyretic properties. It is occasionally employed as an anthelmintic; also given to dogs to allay vomition. *Externally*:—Caustic, disinfectant, astringent, and styptic; applied to cancerous sores and ulcers which emit a fetid odour—*e.g.*, canker and thrush in the foot of the horse, and foot-rot in sheep. It modifies suppuration, and facilitates cicatrisation when applied to wounds. It is administered in various contagious and zymotic diseases, to prevent or arrest the development of micro-organisms.

**Doses.**—*Crystals*:—Horse, 20 to 80 grains.—Cattle, 20 to 80 grains.—Sheep, 5 to 15 grains.—Pig, 5 to 15 grains.—Dog, 1 to 5 grains. *Fluid*:—Horse, 20 to 80 minims.—Cattle, 20 to 80 minims.—Sheep, 5 to 15 minims.—Pig, 5 to 15 minims.—Dog, 1 to 5 minims.

**Modes of Application.**—*Internally*:—Dissolved in glycerin or glycerin and water, or made into a bolus with linseed meal or common mass. *Externally*:—As a caustic, the undiluted acid; for other purposes, in the form of lotion, liniment, or ointment.

Buildings may be disinfected and the atmosphere impregnated with the acid by steeping rags in it and then suspending them in various parts of the room, stable, &c.;

or the acid, in a diluted form,[1] may be sprinkled over floors and walls; or by vapourising the acid on hot metal plates.

**Antidotes.**—Albumen; soap; demulcent drinks; oil.

### Preparations

| | |
|---|---|
| Glycerinum (Linimentum) Acidi Carbolici. | Collodium Hæmostaticum. |
| Lotio Acidi Carbolici. | Sodii Sulphocarbolas. |
| Unguentum Acidi Carbolici. | Carbolic Oil.[2] |
| | Sulphocarbolic Acid. |

## ACIDUM CHROMICUM

CHROMIC ACID

**Synonym.**—Chromic Anhydride.

**Composition.**—$CrO_3$.

**Mode of Preparation.**—By adding to one measure of a saturated solution of potassium dichromate one and a half measure of concentrated sulphuric acid, by small portions at a time, and allowing the solution to cool, when chromic anhydride crystallises out.

**Characters and Tests.**—Crystalline in fine crimson needles, which are deliquescent, and are very soluble in water. They decompose sometimes with explosion when mixed with alcohol. An explosion occurs when 1 part of chromic anhydride is mixed with 2 parts of glycerin.

**Actions and Uses.**—*Externally* in solution (from 5 to 20 per cent.) it acts as a caustic. In the solid form

[1] Pour an ounce or two of the fluid acid into a gallon bottle, nearly fill the bottle with water, shake, and set aside for twenty-four hours. From the undissolved acid decant the aqueous solution for use.

[2] 1 of Phenol, 7 of Castor Oil, and 8 of Almond Oil, used for oiling catheters and instruments after they have been disinfected.

it rapidly destroys tissue (on account of its affinity for water), forming an eschar. It is a powerful disinfectant.

The Liquor Acidi Chromici is made with 1 part of acid to 3 of water.

## ACIDUM GALLICUM

GALLIC ACID

**Synonym.**—Trihydroxy Benzoic Acid.

**Composition.**—$C_6H_2(OH)_3.CO.OH.H_2O$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Galls, in coarse powder . . . | 1 pound |
| Distilled Water . . . . | a sufficiency |

Place the powdered galls in a porcelain dish, pour on as much water as will convert the powder into a thick paste, and keep it in this moist condition for six weeks, at a temperature between 60° and 70°; at the end of that time boil the paste for twenty minutes with forty-five ounces of water, strain through calico, and when the fluid has cooled collect the crystalline deposit which has formed, and let it drain. If the gallic acid is required very pure, it should be recrystallised.

**Characters and Tests.**—Crystalline in acicular prisms or silky needles of a fawn colour; it is soluble 1 in 100 parts of cold water, but dissolves in 3 parts of boiling water. Soluble also in rectified spirit 1 in 8, glycerin 1 in 12. It leaves no residue on ignition with free access of air, it gives a bluish-black precipitate with a ferric salt, but no precipitate with a ferrous salt or a solution of gelatin, showing the absence of tannic acid.

**Actions and Uses.**—The same as Tannic Acid.

## ACIDUM HYDROCHLORICUM

### HYDROCHLORIC ACID

**Synonyms.**—Muriatic Acid ; Spirit of Salt.

**Composition.**—Water holding in solution 31·8 per cent. of its weight of hydrochloric acid gas, HCl.

**Mode of Preparation.**—By distilling a mixture of sodium chloride, sulphuric acid, and water, and allowing the distillate to pass into a receiver containing distilled water.

| Sodium Chloride. | | Sulphuric Acid. | | Hydrochloric Acid. | | Sodium Sulphate. |
|---|---|---|---|---|---|---|
| $2NaCl$ | + | $H_2SO_4$ | = | $2HCl$ | + | $Na_2SO_4$ |

Sodium sulphate remains in the retort, while hydrochloric-acid gas passes over and is absorbed by the water in the receiver.

**Characters and Tests.**—Colourless, strongly acid liquid, which emits white vapours and has a pungent odour. Sp. gr. 1·16. With silver nitrate it gives a white curdy precipitate (silver chloride), which is soluble in ammonium hydrate but insoluble in boiling nitric acid. Should entirely evaporate on the application of heat, and, after being diluted with four times its volume of distilled water, should give no precipitate with hydrosulphuric acid or barium chloride, proving the absence of lead, copper, and sulphuric acid ; neither should a strip of polished copper tarnish when boiled with the acid thus diluted, showing the absence of arsenic and antimony.

**Actions and Uses.**—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, tonic. *Externally* :—Caustic, astringent, antiseptic.

**Modes of Application.**—For internal use, Acidum

Hydrochloricum Dilutum is employed. For external use the part is painted with the acid.

**Antidotes.**—Carbonate of lime (chalk or whiting); magnesia; carbonate of magnesia; demulcent drinks; large draughts of water.

**Preparation.**—Acidum Hydrochloricum Dilutum.

## ACIDUM HYDROCHLORICUM DILUTUM

### DILUTE HYDROCHLORIC ACID

**Composition.**—One hundred parts by weight should contain 10·58 parts of Hydrogen Chloride.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Hydrochloric Acid . . . | 6 fluid ounces |
| Distilled Water . . . . | a sufficiency |

Mix the hydrochloric acid with sixteen fluid ounces of the distilled water, and add more water, so that at 60° F. the mixture shall measure one pint.

**Characters and Tests.**—Sp. gr. 1·052. For tests, see Acidum Hydrochloricum.

**Actions and Uses.**—Tonic internally. Usually given in conjunction with gentian or other bitters.

**Doses.**—Horse, 1 to 2 fluid drachms.—Cattle, 1 to 4 fluid drachms.—Sheep, 10 to 20 minims.—Pig, 10 to 20 minims.—Dog, 2 to 10 minims.

**Modes of Application.**—Diluted with about a hundred times its bulk of water, or combined with infusions of vegetable tonics.

**Incompatibles.**—Carbonates; sulphides; salts of silver.

**Antidotes.**—Same as to Acidum Hydrochloricum.

## ACIDUM HYDROCYANICUM DILUTUM

### DILUTE HYDROCYANIC ACID

**Synonym.**—Prussic Acid.

**Composition.**—Water containing 2 per cent.[1] by weight of real hydrocyanic acid, HCN or HCy (hydrogen cyanide).

**Mode of Preparation.**—By distilling a mixture of potassium ferrocyanide, sulphuric acid, and water.

| Potassium Ferrocyanide. | | Sulphuric Acid. | | Hydrocyanic Acid. | | Potassium-hydrogen Sulphate. | |
|---|---|---|---|---|---|---|---|
| $2K_4FeCy_6$ | + | $6H_2SO_4$ | = | $6HCy$ | + | $6KHSO_4$ | + |

Potassium Ferrous Ferrocyanide
(Everitt's Yellow Salt).
$K_2Fe_2Cy_6$

Potassium-hydrogen sulphate and potassium ferrous ferrocyanide remain in the retort, and hydrocyanic acid, accompanied by water, passes over into the receiver, into which a quantity of water has been previously placed. By the addition of water the distillate is made of such a strength that 100 grains or 110 minims of it precipitated with silver nitrate shall yield ten grains of silver cyanide.

**Characters and Tests.**—Colourless liquid, with a peculiar odour.[2] Sp. gr. 0·997. With potassium hydrate and a mixture of ferrous and ferric sulphates it yields, after being acidulated with hydrochloric acid, a precipitate of Prussian blue. Silver nitrate furnishes a white

[1] Scheele's prussic acid, as met with in the shops, is a preparation of uncertain strength, the amount of real hydrocyanic acid contained in it varying from 1 to 4 per cent.

[2] This character should be very cautiously observed, as the vapour of the acid, when inhaled, is highly dangerous.

curdy precipitate of silver cyanide which is soluble in ammonium hydrate and in boiling nitric acid. Barium chloride should give no precipitate, proving the absence of sulphuric acid and sulphates.

**Actions and Uses.**—*Internally* :—In excessive doses, narcotic and cerebro-spinal poison; in medicinal doses, sedative, antispasmodic, and anodyne. Used in chronic cough, chorea, epilepsy, chronic vomiting, carditis, palpitation of the heart, rheumatism, and tetanus.

**Doses.**—Horse, 20 to 40 minims.—Cattle, 20 to 40 minims.—Sheep, 5 to 10 minims.—Pig, 5 to 10 minims. Dog, 1 to 3 minims.

To prevent accidents, the smaller doses should be given at first, and afterwards gradually increased.

**Modes of Application.**—*Internally* :—Diluted with water. In troublesome cough in horses, twenty minims of the acid with a drachm each of camphor and extract of belladonna, made into a ball with powdered liquorice or linseed meal, and given two or three times a day. To the dog, in obstinate vomiting, two grains of the acid, with ten grains of carbonate of soda and one ounce of water, may be administered every hour. One drachm of the acid with about a quart of water, employed slowly as an enema two or three times a day, lessens muscular contractions in tetanus. *Externally* :—To allay pain and irritation in chronic skin-affections, especially in dogs, two or three drachms of the acid mixed with a pint of distilled or rain water.

**Incompatibles.**—Nitrate of silver; sulphates and chlorides of iron with an alkali.

**Antidotes.**—Fresh air; affusions of cold water; inhalation of ammonia or chlorine; freshly precipitated

hydrated peroxide of iron mixed with an alkaline carbonate.

## ACIDUM NITRICUM

NITRIC ACID

**Synonym.**—Aqua Fortis.

**Composition.**—Water containing 70 per cent. of real nitric acid, $HNO_3$ (hydrogen nitrate).

**Mode of Preparation.**—By distilling a mixture of potassium (or sodium) nitrate and concentrated sulphuric acid; potassium-hydrogen sulphate remains in the retort, and nitric acid distils over.

| Potassium Nitrate. | | Sulphuric Acid. | | Nitric Acid. | | Potassium-hydrogen Sulphate. |
|---|---|---|---|---|---|---|
| $KNO_3$ | + | $H_2SO_4$ | = | $HNO_3$ | + | $KHSO_4$ |

**Characters and Tests.**—Colourless liquid. Sp. gr. 1·42; boiling-point, 250° F. Exposed to air, it emits acrid and corrosive fumes. Copper dropped into a mixture of equal parts of the acid and water causes the evolution of a colourless gas (nitric oxide), which, as it mixes with the oxygen of the air, forms ruddy fumes (nitrous anhydride, $N_2O_3$, and nitric peroxide, $NO_2$). A crystal of ferrous sulphate dropped into the acid, diluted with an equal bulk of water, becomes surrounded by a brown cloud if the mixture be cold. The acid should leave no residue on evaporation. Diluted with six times its volume of distilled water, it should give no precipitate with barium nitrate or silver nitrate, showing the absence of sulphuric and hydrochloric acids.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant poison; in medicinal doses, astringent and tonic,

*Externally* :—Deodoriser and disinfectant; also used as a caustic for removing warts and fungoid growths and in canker; for improving the condition and destroying the fœtor of unhealthy wounds, caries, foul, and foot-rot.

**Modes of Application.**—*Internally* :—See Acidum Nitricum Dilutum. *Externally* :—As a caustic, by means of a pledget of tow tied to a stick, or tow saturated with the acid laid on the part. As an excitant and deodoriser to indolent and sphacelated wounds and ulcers, a mixture of two to four drachms of the acid and one pint of water may be employed. Tow thoroughly saturated with a mixture of four parts of tar and one part of nitric acid is applied to the feet of horses affected with canker.

**Antidotes.**—Copious draughts of water; demulcent drinks; albumen; milk; soap; alkaline carbonates or bicarbonates; carbonate of calcium (chalk or whiting); magnesia; carbonate of magnesia.

**Preparation.**—Acidum Nitricum Dilutum.

## ACIDUM NITRICUM DILUTUM

### DILUTE NITRIC ACID

**Composition.**—Water containing 17·44 per cent. of real nitric acid, $HNO_3$ (hydrogen nitrate).

**Mode of Preparation.**—Take of

| | |
|---|---|
| Nitric Acid . . . . | 3 fluid ounces |
| Distilled Water . . . | a sufficiency |

Dilute the acid with water at 60° F. so that the mixture shall measure one pint.

**Tests.**—Colourless, sour. Sp. gr. 1·101.

**Actions and Uses.**—*Internally*:—Tonic, astringent; during convalescence from debilitating disorders, in chronic skin-diseases, and in chronic enlargement of the liver in cattle.

**Doses.**—HORSE, 1 to 2 fluid drachms. CATTLE, 1 to 2 fluid drachms. SHEEP, 10 to 20 minims.—PIG, 10 to 20 minims.—DOG, 2 to 10 minims.

**Mode of Application.**—Diluted with from fifty to a hundred times its bulk of water.

**Incompatibles.**—Alkalies, earths, and other oxides; carbonates and bicarbonates; sulphides; sulphate of iron; acetate of lead.

**Antidotes.**—See Acidum Nitricum.

## ACIDUM NITRO-HYDROCHLORICUM DILUTUM

### DILUTE NITRO-HYDROCHLORIC ACID

**Synonym.**—Dilute Aqua Regia.

**Composition.**—An aqueous solution of chlorine, hydrochloric, nitrous, and nitric acids.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Nitric Acid . . . . | 3 fluid ounces |
| Hydrochloric Acid . . | 4 fluid ounces |
| Distilled Water . . . | 25 fluid ounces |

Mix the acids and allow them to remain for at least twenty-four hours in a bottle the mouth of which is partly closed; then add the water in successive portions, shake the bottle after each addition, and preserve the mixture in a stoppered bottle.

On mixing the acids they gradually suffer mutual decomposition.

| | Nitric Acid. | | Hydrochloric Acid. | | Chlorine. | | Chloro-nitric Gas. | | Water. |
|---|---|---|---|---|---|---|---|---|---|
| 1. | $2HNO_3$ | + | $6HCl$ | = | $Cl_2$ | + | $2NOCl_2$ | + | $4H_2O$ |

| | Nitric Acid. | | Hydrochloric Acid. | | Chlorine. | | Chloro-nitrous Gas. | | Water. |
|---|---|---|---|---|---|---|---|---|---|
| 2. | $HNO_3$ | + | $3HCl$ | = | $Cl_2$ | + | $NOCl$ | + | $2H_2O$ |

**Characters and Tests.**—Colourless liquid. Sp. gr. 1·074. Dissolves gold leaf.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant poison; in medicinal doses, tonic and stimulant action on the skin, liver, and intestinal glands. For the latter purposes it is preferred to either nitric or hydrochloric acid separately.

**Doses.**—Horse, 1 to 2 fluid drachms.—Cattle, 1 to 2 fluid drachms.—Sheep, 10 to 20 minims.—Pig, 10 to 20 minims.—Dog, 2 to 10 minims.

**Mode of Application.**—Diluted with not less than a hundred times its bulk of water.

**Incompatibles.**
**Antidotes.** } Same as Acidum Nitricum Dilutum.

## ACIDUM PHOSPHORICUM CONCENTRATUM

PHOSPHORIC ACID

**Synonym.**—Orthophosphoric Acid.

**Composition.**—$H_3PO_4$ with 33 per cent. of water.

**Mode of Preparation.**—By heating phosphorus with dilute nitric acid until nitrous fumes cease to form, and diluting with water until the sp. gr. is 1·5.

**Characters and Tests.**—A colourless, sour, syrupy liquid, with an acid reaction. In dilute solution it gives a yellow precipitate with silver nitrate.

**Actions and Uses.**—*Externally*:—An irritant and escharotic.

## ACIDUM PHOSPHORICUM DILUTUM

DILUTE PHOSPHORIC ACID

**Composition.**—Water containing 13·8 per cent. of $H_3PO_4$.

**Mode of Preparation.**—By mixing three parts of concentrated phosphoric acid with twenty of water.

**Actions and Uses.**—In debilitated conditions of the system, has the reputation of checking tuberculosis. It is a good tonic, and useful in bronchial catarrh. It is also useful in the convalescent stage of distemper in the dog.

**Doses.**—Horse, 1 to 2 drachms.—Dog, 5 to 10 minims.

**Preparations.**—Phosphates of ammonia, iron, and soda. Syrupus ferri, quininæ, et strychninæ phosphatis (syrup of the phosphates of iron, quinine, and strychnine) (Easton's syrup) contains iron phosphate 1·33, quinine 1·33, strychnine 0·04, phosphoric acid 8, sugar 60, water to 100 parts: each drachm contains about $\frac{1}{32}$ gr. of strychnine.

## ACIDUM SALICYLICUM

SALICYLIC ACID

**Synonym.**—Hydroxybenzoic Acid.

**Composition.**—$HC_7H_5O_3$ or $C_6H_4.OH.COOH$.

**Mode of Preparation.**—By passing carbonic anhydride through sodium phenol for several hours, heating

to 482° F., and treating the residue with dilute hydrochloric acid.

Also found as salicylate in oils of wintergreen (*Gaultheria procumbens*) and sweet birch (*Betula lenta*).

**Characters and Tests.**—A colourless powder, usually crystallising in minute acicular crystals. Taste at first sweet, afterwards bitter. Soluble in water, 1 in 500 to 700; more soluble in hot water, alcohol, and ether. Ferric chloride gives a violet colour with both the acid and its salts. Copper sulphate gives an emerald green colour.

**Actions and Uses.**—Antiseptic, antipyretic. Both the acid and sodium salicylate are employed in the treatment of rheumatism. *Externally*:—It is used as a dressing for wounds, and to relieve the itching and discharge of eczema and psoriasis. Also in the treatment of open joints.

**Doses.**—Horses and Cattle, 4 to 8 drachms.—Sheep, 1 to 2 drachms.—Dogs, from 10 to 15 grains.

**Preparations.**—Solution of one part each of salicylic acid and borax to thirty or fifty parts of water, used for soaking lint, &c., as an antiseptic covering for wounds.

**Ointment.**—Salicylic acid one part, mixed warm with twenty to twenty-five of vaselin.

The B.P. ointment contains one part of salicylic acid, eighteen of soft paraffin, and nine of hard paraffin.

## ACIDUM SULPHOCARBOLICUM

SULPHOCARBOLIC ACID

**Composition.**—$HC_6H_4O.HSO_3$.

**Mode of Preparation.**—By the action of strong sulphuric acid on phenol.

**Characters and Tests.**—Crystallises slowly in colourless, deliquescent needles, having less odour than carbolic acid; soluble in water, alcohol, ether, and glycerin. Forms salts called sulphocarbolates.

**Actions and Uses.**—Antiseptic.[1]

**Mode of Application.**—A 10-per-cent. aqueous solution is used as a dressing for wounds.

## ACIDUM SULPHURICUM

### SULPHURIC ACID

**Synonym.**—Oil of Vitriol.

**Composition.**—Should contain 98 per cent. by weight of hydrogen sulphate, $H_2SO_4$.

**Mode of Preparation.**—Sulphurous anhydride, vapour of nitric acid, steam, and air are conducted into a leaden chamber, where they react upon one another in the following manner:

| Sulphurous Anhydride. | | Nitric Acid. | | Water. | | Sulphuric Acid. | | Nitric Oxide. |
|---|---|---|---|---|---|---|---|---|
| $3SO_2$ | + | $2HNO_3$ | + | $2H_2O$ | = | $3H_2SO_4$ | + | $2NO$ |

The resulting nitric oxide abstracts oxygen from the air, and becomes nitric peroxide:

| Nitric Oxide. | | Oxygen. | | Nitric Peroxide. |
|---|---|---|---|---|
| $2NO$ | + | $O_2$ | = | $2NO_2$ |

The nitric peroxide thus formed oxidises further quantities of sulphurous anhydride, and, with the assistance of steam, converts it into sulphuric acid:

| Sulphurous Anhydride. | | Nitric Peroxide. | | Water. | | Sulphuric Acid. | | Nitric Oxide. |
|---|---|---|---|---|---|---|---|---|
| $SO_2$ | + | $NO_2$ | + | $H_2O$ | = | $H_2SO_4$ | + | $NO$ |

[1] Aseptol, a syrupy dark liquid, freely soluble in alcohol, water, and glycerin. It contains orthophenol-sulphonic acid (33½ per cent.). Used diluted (1 to 20 of water) as an antiseptic for surgical purposes.

The acid produced in the leaden chambers is very dilute; it is concentrated by evaporation, first in leaden basins, and finally in glass or platinum retorts.

**Characters and Tests.**—Colourless, heavy, oily, intensely acid liquid. Sp. gr. 1·843. Mixed with water, much heat is evolved. Diluted with five or six times its volume of water, it gives, with barium chloride, a white precipitate (barium sulphate), which is insoluble in nitric acid; it should yield no precipitate or darkening of colour with hydrosulphuric acid, indicating the absence of lead. When a solution of ferrous sulphate is carefully poured upon the surface of the acid, there should be no brown colour developed where the two liquids unite, showing the absence of nitric acid and oxides of nitrogen. The acid should leave no residue on evaporation. The commercial acid prepared from pyrites is always contaminated with arsenic.

**Actions and Uses.**—*Internally*:—Irritant and corrosive poison; see Acidum Sulphuricum Dilutum. *Externally*:—Employed as caustic for the removal of warts and cancerous growths.

**Mode of Application.**—Made into a paste with flowers of sulphur, sawdust, finely powdered charcoal, or dried alum.

**Antidotes.**—Same as for Acidum Nitricum.

## ACIDUM SULPHURICUM DILUTUM

DILUTE SULPHURIC ACID

**Composition.**—Water containing 13·65 per cent. of hydrogen sulphate, $H_2SO_4$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Sulphuric Acid . . . . | 7 fluid ounces |
| Distilled Water . . . . | a sufficiency |

Place seventy-seven fluid ounces of distilled water in a thin glass or earthenware vessel, and then pour into it the seven fluid ounces of acid. Stir the mixture, and, when cooled to 60° F., add more water, so that it shall measure eighty-three and a half fluid ounces.

**Characters and Tests.**—Colourless, sour liquid. Sp. gr. 1·094.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant poison; in medicinal doses, astringent; also tonic during convalescence, and in other cases in which mineral tonics are indicated. *Externally*:—Astringent.

**Doses.**—Horse, 2 to 4 fluid drachms.—Cattle, 2 to 4 fluid drachms.—Sheep, 10 to 20 minims.—Pig, 10 to 20 minims.—Dog, 2 to 10 minims.

**Modes of Application.**—*Internally*:—As a tonic, it is given diluted with from fifty to a hundred times its bulk of water. In diarrhœa and dysentery, accompanied by alkaline discharges, two fluid drachms of acid, with one fluid ounce of laudanum, may be administered twice a day in gruel or some mucilaginous drink. One fluid drachm of this acid, with one ounce each of sulphuric ether and powdered cinchona-bark, is sometimes given in gruel or ale to horses suffering from typhoid influenza. In cases of relaxed and ulcerated sore throat in horses, the acid, highly diluted with water, may be slowly given. *Externally*:—As an astringent, thirty minims of acid in one fluid ounce of water.

**Incompatibles.**
**Antidotes.** } Same as Acidum Nitricum.

## ACIDUM SULPHUROSUM

SULPHUROUS ACID

**Composition.**—Water containing 5 per cent. by weight of sulphurous anhydride (sulphurous-acid gas), $SO_2$.[1]

**Mode of Preparation.**[2]—Take of

| | |
|---|---|
| Sulphuric Acid . . . . | 4 fluid ounces |
| Wood Charcoal, in coarse powder | 1 ounce |
| Common Water . . . . | 2 fluid ounces |
| Distilled Water . . . . | 20 fluid ounces |

Put the charcoal and sulphuric acid into a glass flask, connected by a glass tube with a wash-bottle containing the two fluid ounces of common water, whence a second tube leads into a pint bottle containing the distilled water, to the bottom of which the gas-delivery tube should pass. Apply heat to the flask until gas is evolved, which is to be conducted through the water in the wash-bottle, and then into the distilled water, the latter being kept cold, and the process being continued until the bubbles of gas pass through the liquid undiminished in size.

| Sulphuric Acid. | | Carbon. | | Sulphurous Anhydride. | | Carbonic Oxide. | | Water. |
|---|---|---|---|---|---|---|---|---|
| $H_2SO_4$ | + | C | = | $SO_2$ | + | CO | + | $H_2O$ |

The product should be preserved in a stoppered bottle, to prevent the conversion of the sulphurous anhydride, by absorption of oxygen, into sulphuric acid thus:

| Sulphurous Anhydride. | | Water. | | Oxygen from Air. | | Sulphuric Acid. |
|---|---|---|---|---|---|---|
| $SO_2$ | + | $H_2O$ | + | O | = | $H_2SO_4$ |

[1] Equal to 6·4 per cent. of hypothetical sulphurous acid, $H_2SO_3$.

[2] For fumigating or disinfecting purposes, sulphurous anhydride may be easily evolved by setting fire to a few fragments of roll sulphur (brimstone) placed in a saucer or other suitable vessel:

| Sulphur. | | Oxygen from Air. | | Sulphurous Anhydride. |
|---|---|---|---|---|
| S | + | $O_2$ | = | $SO_2$ |

Sulphurous anhydride, $SO_2$, may also be prepared by boiling sulphuric acid with mercury, copper, or sulphur. It should be kept in a cool place, to prevent loss of gas.

**Characters and Tests.**—A colourless liquid, with an odour like that emitted from burning sulphur. Sp. gr. 1·04. Should leave no residue on evaporation, and should give no precipitate, or but a very slight one, with barium chloride, showing the absence, or nearly so, of sulphuric acid.

**Actions and Uses.**—Antiseptic, deodoriser, disinfectant.[1] Applied externally to kill fungi that cause or accompany certain skin-affections. Antiseptic properties are also possessed by the sulphites and hyposulphites.

**Mode of Application.**—One part of the acid mixed with three parts of water, or two of glycerin, applied to the diseased surface with a sponge.

## ACIDUM TANNICUM

### TANNIC ACID

**Synonym.**—Tannin.

**Composition.**—Pure tannic acid has the formula $C_{13}H_9O_7.CO_2H$, but that obtained by the method described below, and taken from the 'British Pharmacopœia' of 1867, is mixed with gallic[2] acid, $C_6H_2(OH)_3$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Oak Galls in powder . . | } of each a sufficiency |
| Ether . . . . | |

[1] Active principles of Tuson's disinfectant, sulphurous acid and chloride of zinc. See Zinci Chloridum.

[2] By some practitioners gallic acid is regarded as being a more effective astringent and styptic than tannic acid, and is prescribed in cases where hæmorrhage has to be reached through the circulation. The doses are the same as those of tannic acid.

Expose the powdered galls to a damp atmosphere for two or three days, and afterwards add sufficient ether to form a soft paste. Let this stand in a well-closed vessel for twenty-four hours, then, having quickly enveloped it in a linen cloth, submit it to a strong pressure in a suitable press so as to separate the liquid portion. Reduce the pressed cake to powder, mix it with sufficient ether, to which one-sixteenth of its bulk of water has been added, to form again a soft paste, and press this as before. Mix the expressed liquids, and expose the mixture to a spontaneous evaporation until, by the aid subsequently of a little heat, it has acquired the consistence of a soft extract; then place it on earthen plates or dishes, and dry it in a hot-air chamber at a temperature not exceeding 212° F.

Under the combined influence of moisture and atmospheric oxygen tannic acid is converted into gallic acid and glucose.

**Characters and Tests.**—Pale yellow vesicular masses, or thin glistening scales, with strongly astringent taste, and acid reaction. Readily soluble in water, ten in eight parts, and rectified spirit, sparingly soluble in ether. With an aqueous solution gelatin gives a yellowish-white precipitate (gelatin tannate), and ferric chloride furnishes a bluish-black precipitate (ferric tannate). The acid should leave no residue when burned with free access of air.

**Actions and Uses.**—*Internally and Externally*:—Astringent and styptic. Employed to diminish mucous discharges, as in diarrhœa and dysentery; also used as a vermicide. Antidote to tartar emetic; also to strychnine, morphine, and other alkaloidal poisons. Tannic acid is an ingredient of styptic collodion.

**Doses.**—Horse, 20 to 60 grains.—Cattle, 20 to 60

grains.—SHEEP, 4 to 12 grains.—PIG, 4 to 12 grains.—DOG, 1 to 3 grains.

**Modes of Application.**—*Internally* :—Either in the form of bolus, pill, or solution in water or spirit of wine, or in glycerin and water. *Externally* :—One part of acid dissolved in six parts of water; or unguentum acidi tannici.

**Incompatibles.**—Mineral acids; alkalies; salts of lead, silver, iron, antimony; alkaloids; gelatin and emulsions.

**Preparations.**—Glycerinum Acidi Tannici; Unguentum Acidi Tannici.

## ACONITI FOLIA

ACONITE-LEAVES

The fresh leaves and flowering tops of *Aconitum Napellus* (Monkshood, Wolfsbane, Blue Rocket), gathered when about one third of the flowers are expanded, from plants cultivated in Britain.

**Natural Order.**—*Ranunculaceæ.*

**Characters.**—Leaves smooth, palmate, divided into five deeply cut wedge-shaped segments, which are further incised in a pinnatifid manner, dark green above, paler beneath; slowly exciting, when *cautiously* chewed, a sensation of tingling.

**Actions and Uses.**—*Internally* :—In excessive doses, cerebro-spinal poison; in medicinal doses, sedative and anodyne, febrifuge, diaphoretic, and diuretic. *Externally* :—Occasionally applied to allay neuralgic and rheumatic pains; for this purpose, however, linimentum aconiti or tinctura aconiti is usually preferred.

**Antidotes.**—Emetics, when they can be employed; stimulants, internally and externally.

**Preparations.**—Extractum Aconiti; Linimentum Aconiti; Tinctura Aconiti.

## ACONITI RADIX

ACONITE-ROOT

The dried root of *Aconitum Napellus*. Imported from Germany or cultivated in Britain, and collected in the winter or early spring, before the leaves have appeared.

**Natural Order.**—*Ranunculaceæ.*

**Characters.**—Usually from one to three inches long, not more than about three-quarters of an inch in thickness at the crown, tapering; externally, blackish-brown; internally, whitish; earthy odour. A minute portion *cautiously* chewed causes prolonged tingling and numbness.

**Antidotes.**—See Aconiti Folia.

**Preparations.** — Aconitina; Linimentum Aconiti; Tinctura Aconiti.

## ACONITINA

ACONITINE

**Synonym.**—Aconitia.

A highly poisonous alkaloid, $C_{33}H_{43}NO_{12}$, forming the active principle of the root, leaves, and other parts of several species of *Aconitum*. Not often used in veterinary practice, as it is very costly.

## ADEPS PRÆPARATUS

PREPARED LARD

**Natural Order.**—*Pachydermata.*

**Synonyms.**—Axungia ; Hog's Lard.

**Composition.**—A mixture of 38 per cent. of stearin and palmitin, with 62 per cent. of olein.

**Preparation.**—The fat of the hog (*Sus scrofa*), taken from the loins, omentum, and mesentery, is deprived of its membranes and purified by being kneaded with cold water, melted, strained, dried a little above 212° F., again strained through flannel, and finally run into bladders or casks and allowed to solidify.

**Characters and Tests.**—Soft, white, fatty substance, melting at about 100° F. Should not have a rancid odour, and should dissolve entirely in ether. If boiled with distilled water and filtered, the filtrate should give no precipitate with silver nitrate, showing the absence of common salt ; and, when quite cold, no blue colour with solution of iodine, proving the absence of starch and flour.

**Actions and Uses.**—Chiefly employed externally as a simple dressing to ulcers and blistered surfaces, which it softens and protects from the irritating action of the atmosphere and of acrid discharges. Occasionally to kill the acarus in mange, scab, and similar complaints. Extensively used in the preparation of ointments.

Benzoated lard, preferable on account of its agreeable odour and diminished liability to rancidity, is made by melting one pound of purified lard over a water-bath, and stirring in 210 grains of benzoin, and afterwards straining.

## ÆTHER

ETHER

**Synonyms.**—Ethyl Oxide; Ethylic Ether; Sulphuric Ether.

**Composition.**—About 92 per cent. of pure Ether $(C_2H_5)_2O$, with about 8 per cent. of impurities, chiefly water and alcohol.

**Preparation.**—Take of

| | | |
|---|---|---|
| Rectified Spirit | 50 | fluid ounces |
| Sulphuric Acid | 10 | fluid ounces |
| Chloride of Calcium | 10 | ounces |
| Slaked Lime | ½ | ounce |
| Distilled Water | 13 | fluid ounces |

Mix the sulphuric acid with twelve fluid ounces of the spirit in a glass retort capable of holding at least two pints; connect the retort with a Liebig's condenser to which a receiver has been attached, and distil with a heat sufficient to maintain the liquid in brisk ebullition. Allow the remainder of the spirit slowly to flow in a continuous stream into the retort. For this purpose use a tube furnished with a stopcock to regulate the supply, connecting one end of the tube with a vessel containing the spirit raised above the level of the retort, and passing the other end through a cork fitted into the neck of the retort. When the whole of the spirit has been added, and forty-two fluid ounces of impure Ether have distilled over, the process may be stopped.

Dissolve the chloride of calcium in the distilled water, add the lime, and agitate the mixture in a bottle with the impure ether; the chloride of calcium absorbs water, and the lime sulphurous acid. Leave the mixture at rest for ten minutes, pour off the light supernatant fluid, and distil

it with a gentle heat until a glass bead of sp. gr. 0·735 placed in the receiver begins to float. The product is ether of the composition already stated.

In the foregoing process the decomposition takes place in two stages:

| | Alcohol. | | Sulphuric Acid. | | Sulphethylic Acid. | | Water. |
|---|---|---|---|---|---|---|---|
| 1. | $C_2H_5OH$ | + | $H_2SO_4$ | = | $C_2H_5HSO_4$ | + | $H_2O$ |

| | Alcohol. | | Sulphethylic Acid. | | Ether. | | Sulphuric Acid. |
|---|---|---|---|---|---|---|---|
| 2. | $C_2H_5OH$ | + | $C_2H_5HSO_4$ | = | $(C_2H_5)_2O$ | + | $H_2SO_4$ |

The ether distils over, and the sulphuric acid remains in the retort.

**Characters and Tests.**—Colourless, very volatile, and inflammable liquid, emitting a strong and characteristic odour, and boiling below 105° F. Sp. gr., pure ether 0·720 to 0·722; B.P. 0·735. Fifty measures shaken with an equal volume of water are reduced to forty-five, by an absorption of 10 per cent. Miscible in all proportions with alcohol and chloroform. It should evaporate without residue.

**Actions and Uses.**—*Internally*:—Stimulant, narcotic, and antispasmodic.—*Externally*:—Refrigerant. Inhaled, it acts as an anæsthetic. Used also in the production of local anæsthesia.

**Doses.**—Horse, 1 to 3 fluid ounces.—Cattle, 2 to 3 fluid ounces.—Sheep, 2 to 4 fluid drachms.—Pig, 2 to 4 fluid drachms.—Dog, $\frac{1}{2}$ to 2 fluid drachms.

As an anæsthetic the larger animals take from 3 to 8 ounces, the smaller 4 drachms to 1 ounce.

**Modes of Application.**—As a stimulant and antispasmodic, it is given with ten parts of cold water; as a refrigerant, it is applied to the part either alone or mixed

with a cooling lotion; as a general anæsthetic, a sponge saturated with it is placed in the bottom of a nose-bag attached to the patient's head, and perforated so as to ensure the ether vapour being mixed with air. For the production of local anæsthesia, Dr. Richardson's spray-diffuser must be employed.

**Preparations.**—Collodium; Collodium Flexile; Spiritus Ætheris.

## ALBUMEN OVI

### EGG-ALBUMEN

Liquid white of the egg of the *Gallus Bankiva* var. *domesticus*.

**Natural Order.**—*Gallinæ*.

**Composition.**—Consists of carbon, hydrogen, oxygen, nitrogen, sulphur, and phosphorus, $C_{72}H_{108}N_{10}O_{22}S$.

**Actions and Uses.**—As an antidote to corrosive poisons.[1]

**Doses.**—Ad libitum.

**Modes of Application.**—Before administration it should be whisked, either alone or with water or demulcent drinks.

## ALOE BARBADENSIS

### BARBADOS ALOES

The inspissated juice of the leaf of *Aloe vera*, and other species. Imported from Barbados, usually in gourds.

**Natural Order**—*Liliaceæ*.

[1] Serum from blood, emulsions of bean or pea meal, or milk may be used in the absence of egg-albumen

**Composition.**—Aloin, a volatile oil, a brown resin, mineral matter, and albumen, $C_{16}H_{16}O_7.3H_2O$.

**Characters.**—Yellowish-brown or dark-brown opaque masses; breaks with a dull conchoidal fracture; bitter nauseous taste, and a strong disagreeable odour; dissolves almost entirely in proof spirit, and during solution exhibits under the microscope numerous crystals.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant poison; in medicinal doses, purgative, nauseant, tonic, diuretic, vermifuge. *Externally*:—Traumatic.

**Doses.**—*Purgative.*—HORSE, 2 to 8 drachms.—CATTLE, 1 to 2 ounces.—SHEEP, ½ to 1 ounce.—PIG, 1 to 4 drachms.—DOG, 20 grains to 2 drachms. As a tonic the dose of Aloes for any of the domesticated animals is about one-eighth to one-tenth of that given as a purgative.

**Modes of Application.**—*Internally*:—Massa Aloes, Liquor Aloes, and Liquor Aloes et Sodii Carbonatis. *Externally*:—As a traumatic, the powder is sprinkled over wounds; Tinctura Aloes Composita.

**Preparations.**—Liquor Aloes; Liquor Aloes et Sodii Carbonatis; Massa Aloes; Tinctura Aloes Composita.

## ALOE CAPENSIS

### CAPE ALOES[1]

The inspissated juice of the leaf of *Aloe spicata* and several other allied species. Imported from the Cape of Good Hope in skins and in chests.

**Natural Order.**—*Liliaceæ.*

[1] Barbados aloes is usually preferred to this preparation, as it is considered by some veterinary practitioners to be more certain in its therapeutic action, but the better qualities of Cape aloes are little, if at all, inferior to Barbados or Socotrine aloes.

**Composition.**—Similar to that of Aloe Barbadensis.

**Characters.**—Glossy, resinous appearance; dark brown colour, with a strong, greenish-yellow shade, especially when in small fragments; strong, disagreeable odour, increased by breathing on it; acrid, bitter taste; very brittle and readily reduced to powder, which has a shining greenish-yellow colour.

**Actions and Uses.**
**Doses.**
**Modes of Application.**
} Similar to those of Aloe Barbadensis.

## ALOE SOCOTRINA

SOCOTRINE ALOES

The inspissated juice of the leaf of one or more undetermined species of *Aloe*, probably *Aloe socotrina* and *Aloe Perryi*, belonging to the natural order *Liliaceæ*. Imported from the island of Socotra, also from Zanzibar. It comes over in skins, casks, kegs, and chests.

**Composition.**—Similar to Aloe Barbadensis.

**Characters.**—Occurs in masses, varying in colour in different parts of the same mass, sometimes garnet red, at other times much paler, and when quite dry golden red, yielding a golden-yellow powder. The colour is deepened by exposure to air; fine pieces break with a smooth, glassy, conchoidal fracture, but specimens of good quality often break with a roughish fracture. The taste is very bitter, and the odour of freshly broken pieces (especially when breathed on) is very fragrant. Dissolves entirely in proof spirit, and during solution exhibits under the microscope numerous minute crystals.

**Actions and Uses.**
**Doses.**
**Modes of Application.**
} Similar to those of Aloe Barbadensis.

## ALOINUM

ALOIN

**Synonym.**—Aloine.

**Composition.**—$C_{16}H_{16}O_7.3H_2O$. Three crystalline principles are known—namely, Aloin, Nataloin, and Socaloin : they are either isomeric or form a homologous series. These are the principal constituents of aloes, of which they form about 60 per cent. They are contained in the cold infusion of aloes, and also in a decoction which has cooled ; they may be obtained from either by evaporation. Thus procured, they are crystalline principles, readily soluble in water but with difficulty in alcohol. Their insolubility in ether distinguishes them from the bitter principle of rhubarb.

**Doses.**—Horse, 1 to 1½ drachm.—Dog, 3 to 20 grains.

## ALUMEN

ALUM

**Synonyms.**—Ammonium-Aluminium Sulphate or Potassium-Aluminium Sulphate ; Potassium Alum.

**Composition.**—$(NH_4)_2SO_4.Al_2(SO_4)_3 24H_2O$, or $K_2SO_4.Al_2(SO_4)_3 24H_2O$.

**Mode of Preparation.**—There are various modes, but the following is the simplest : Cornish or pipe clay is boiled with sulphuric acid, so as to form aluminium sulphate and silica :

| Clay (Aluminium Silicate). | | Sulphuric Acid. | | Aluminium Sulphate. | | Silica (Silicic Anhydride). | | Water |
|---|---|---|---|---|---|---|---|---|
| $Al_2(SiO_3)_3$ | + | $3H_2SO_4$ | = | $Al_2(SO_4)_3$ | + | $3SiO_2$ | + | $3H_2O$ |

The aluminium sulphate, after being separated from the silica, is mixed with solution of potassium sulphate; the two salts unite, and on evaporation crystallise out as alum:

$$Al_2(SO_4)_3.K_2SO_4.24H_2O.$$

**Characters and Tests.**—Colourless transparent crystalline masses, exhibiting the faces of the regular octahedron, and having an astringent taste.

An aqueous solution gives a white gelatinous precipitate (aluminium hydrate) with potassium hydrate, soluble in excess of this reagent, indicating the presence of aluminum. A platinum wire moistened with the solution, held in the reducing flame of the blowpipe, communicates a violet colour to the flame, showing the presence of potassium. Barium chloride added to the aqueous solution gives a white precipitate (barium sulphate), insoluble in nitric acid, proving that the compound is a sulphate. Potassium ferrocyanide should not produce either a pale or a dark blue colour, showing the absence of iron.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant poison; in medicinal doses, astringent in diarrhœa and dysentery; as an antidote to poisoning by lead, and to arrest the secretion of milk in cows. *Externally*:—Astringent to open joints and wounded thecæ; as a collyrium in chronic conjunctivitis; also styptic.

**Doses.**—Horse, 2 to 5 drachms.—Cattle, 2 to 5 drachms.—Sheep, ½ to 2 drachms.—Pig, ½ to 2 drachms. Dog, 10 to 30 grains.

**Modes of Application.**—*Internally*:—Dissolved in water, or in the form of bolus, either alone or in conjunction with opium and carminatives. *Externally*:—Pulvis Aluminis sprinkled over the wound; otherwise apply Liquor Aluminis or Unguentum Aluminis Compositum,

or, in cases of open joints, one part of finely powdered Alum with two or three parts of flour may be dusted over the opening.

**Incompatibles.**—Tannic acid and vegetable astringents containing it; alkalies, earths, and their carbonates; acetate of lead.

**Preparations.**—Alumen Exsiccatum; Liquor Aluminis; Unguentum Aluminis Compositum; Glycerin of Alum.[1]

## ALUMEN EXSICCATUM

DRIED ALUM

**Synonym.**—Burned Alum.

**Composition.**—Alum deprived of its water of crystallisation by heat.

**Mode of Preparation.**—Heat alum in a porcelain dish till aqueous vapour ceases to be given off, taking care that temperature never exceeds 400° F. Pulverise the residue, and preserve in a well-stoppered bottle.

**Actions and Uses.**—Caustic and astringent; employed externally for the same purposes as alum, than which it is more powerful.

**Preparation.**—Pulvis Aluminis.

## AMMONIACUM

AMMONIACUM

**Synonym.**—Gum Ammoniacum.

A gum-resinous exudation from *Dorema Ammoniacum*. Collected in Persia and the Punjaub.

[1] Alum 1 ounce, water 3 fluid drachms, and glycerin 6 ounces. Used for chapped teats and eczema.

**Natural Order.**—*Umbelliferæ.*

**Composition.**—$C_{40}H_{30}O_9$. When distilled yields umbelliferone, $C_9H_6O_3$.

**Characters.**—In tears or masses; the tears from one-fifth to four-fifths of an inch in diameter, pale cinnamon-brown, breaking with a smooth, shining, opaque, white surface; the masses composed of agglutinated tears, hard and brittle when cold, but readily softening with heat. Faint odour, and a bitter, acrid, nauseous taste. Rubbed with water forms a nearly white emulsion.

**Actions and Uses.**—*Internally*: Stimulant and antispasmodic in coughs and spasmodic nervous diseases—*e.g.*, chorea, epilepsy, &c.; also vermifuge.

**Doses.**—Horse, 2 to 4 drachms.—Cattle, 2 to 4 drachms.—Sheep, $\frac{1}{2}$ to $1\frac{1}{2}$ drachm.—Pig, $\frac{1}{2}$ to $1\frac{1}{2}$ drachm.—Dog, 10 to 20 grains.

**Modes of Application.**—Administered in the form of bolus or emulsion. As a vermifuge it may be given either by the mouth or by the rectum.

## AMMONII CARBONAS

### CARBONATE OF AMMONIUM

**Synonym.**—Commercial Carbonate of Ammonia.

**Composition.**—Probably $2NH_4HCO_3NH_4NH_2CO_2$—*i.e.*, a mixture or compound of two molecules of ammonium-hydrogen carbonate and one molecule of ammonium carbamate.

**Preparation.**—By heating to redness a mixture of one part of ammonium chloride and two parts of calcium carbonate (chalk) in an iron or earthenware retort, to which is luted an earthenware or leaden receiver. When

the receiver has become filled with the 'carbonate of ammonia' by repeated distillations, it is broken or cut in two, and its contents removed.

'Carbonate of Ammonia'

| Ammonium Chloride. | Calcium Carbonate. | | Ammonium-Hydrogen Carbonate. | | Ammonium Carbamate. | |
|---|---|---|---|---|---|---|
| $6NH_4Cl$ | + $3CaCO_3$ | = | $[2NH_4HCO_3$ | + | $NH_4NH_2CO_2]$ | + |

| Ammoniacal Gas. | | Calcium Chloride. | | Water. |
|---|---|---|---|---|
| $2NH_3$ | + | $3CaCl_2$ | + | $H_2O$ |

The ammoniacal gas and water are first disengaged, then the 'carbonate of ammonia' distils over, and the calcium chloride remains in the retort.

**Characters and Tests.**—Translucent crystalline masses, with a strong ammoniacal odour[1] and alkaline reaction; soluble in cold water,[2] sparingly so in spirit of wine. Volatilised entirely by heat, and dissolved by acids with effervescence. If dissolved in slight excess of dilute nitric acid and boiled, the solution should give no white precipitate with silver nitrate, showing the absence of chlorine, and no blue colour with potassium ferrocyanide, proving the absence of iron.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant and narcotic poison; in medicinal doses, stimulant, antacid, and resolvent. Given in influenza and typhoid

[1] The odour is due to the escape of ammoniacal gas. By exposure to air the preparation also loses carbonic anhydride and becomes ammonium-hydrogen carbonate, which is opaque, odourless, and antacid, but not stimulant; hence, 'carbonate of ammonia' should be preserved in well-stoppered bottles.

[2] When commercial or pharmaceutical 'carbonate of ammonium' is dissolved in water, the ammonium carbamate which it contains is soon transformed into normal ammonium carbonate; thus:

| Ammonium Carbamate. | | Water. | | Ammonium Carbonate. |
|---|---|---|---|---|
| $NH_4(NH_2CO_2)$ | + | $H_2O$ | = | $(NH_4)_2CO_3$ |

affections; also in inflammation of the lungs, pleurisy, and similar complaints.

**Doses.**—Horse, 2 to 4 drachms.—Cattle, 3 to 6 drachms.—Sheep, 20 grains to 1 drachm.—Pig, 20 grains to 1 drachm.—Dog, 3 to 10 grains.

**Modes of Application.**—In a bolus or pill, or in cold gruel. As a stimulant it is given mixed with alcohol or ether, as a febrifuge with spiritus ætheris nitratis, potassii chloras, and camphor.

**Incompatibles.**—Acids, alkalies, lime-water, and most metallic salts.

**Antidotes.**—Copious draughts of water or mucilaginous drinks, oil, or highly diluted acids.

**Preparations.**—Liquor Ammoniæ Acetatis; Spiritus Ammoniæ Aromaticus.

## AMMONII CHLORIDUM

### CHLORIDE OF AMMONIUM

**Synonyms.**—Ammonium Chloride; Muriate of Ammonia; Sal Ammoniac.

**Composition.**—$NH_4Cl$.

**Preparation.**—'Gas liquor,' which is water holding in solution carbonate and other salts of ammonium, is acidulated with hydrochloric acid and evaporated to dryness. The crude ammonium chloride, constituting the residue, is purified by sublimation.

**Characters and Tests.**—Colourless, odourless, translucent, fibrous masses, tough and difficult to powder; soluble in water and in rectified spirit. Its aqueous solution, heated with potassium hydrate, evolves ammoniacal gas, which may be recognised by its odour; silver

nitrate gives a white curdy precipitate (silver chloride), insoluble in boiling nitric acid, soluble in ammonia.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant poison; in medicinal doses, stimulant, resolvent, diuretic. Given in mucous fevers after subsidence of acute inflammatory symptoms; in the milder forms of pneumonia and inflammations of serous membranes; in mucous diarrhœa, chronic rheumatism, and passive dropsies. *Externally*:—Stimulant to bruises, sprains, and inflammatory swellings.

**Doses.**—Horse, 2 to 15 drachms.—Cattle, 15 drachms to 1 ounce.—Sheep, ½ to 2 drachms.—Pig, ½ to 2 drachms.—Dog, 5 to 20 grains.

**Modes of Application.**—*Internally*:—Bolus, pill, or dissolved in water. *Externally*:—Lotio Ammonii Chloridi et Camphoræ; Lotio Ammonii Chloridi et Potassæ Nitratis.

**Incompatibles.**—Sulphuric and nitric acids; potash, soda, lime, and their carbonates; and most metallic salts.

**Antidotes.**—Large draughts of water; and, when possible, vomiting should be promoted by warm demulcent and mucilaginous drinks.

**Preparations.**—Lotio Ammonii Chloridi et Camphoræ; Lotio Ammonii Chloridi et Potassii Nitratis.

## AMYL NITRIS

### NITRITE OF AMYL

**Synonym.**—Amyl Nitrite.

**Composition.**—$C_5H_{11}NO_2$.

**Mode of Preparation.**—By passing nitrous acid into amyl alcohol.

**Characters and Tests.**—Nitrite of amyl is a yellowish ethereal liquid; sp. gr. of liquid, 0·877; boiling-point, 205° F.; soluble in alcohol, insoluble in water; converted by fused potassium hydrate into valerianate of potassium ($KC_5H_9O_2$).

**Actions and Uses.**—Sedative, antispasmodic, and anæsthetic. Used in the treatment of tetanus, epilepsy, and angina pectoris.

As an inhalation when too much chloroform has been administered.

**Doses.**—As an anæsthetic the vapour of 1 to 3 drachms for the HORSE; 10 to 20 minims for the DOG. Hypodermically or internally, about one-third or one-fourth of the above doses may be used.

## AMYLUM

STARCH

**Synonym.**—Fecula.

**Composition.**—$C_6H_{10}O_5$, or some multiple.

**Mode of Preparation.**—Chiefly obtained by grinding wheat or rice, or rasping potatoes, making the product into a pulp with water, and washing it in a sieve. The starch passes through the sieve, subsides, and is collected and dried at a gentle heat.

**Characters and Tests.**—White columnar masses. Rubbed in a mortar with a little cold water, it should be neither acid nor alkaline to test-paper. Mixed with boiling water and cooled, it gives a deep blue colour (starch iodide) with solution or tincture of iodine.

**Actions and Uses.**—*Internally*:—Demulcent and emollient, in the form of gruel. *Externally*:—Applied in the state of dry powder, it diminishes the discharge from

wounds, and, mixed with about one-eighth of its weight of alum, it is employed to arrest the flow of synovia from open joints. Also used in the preparation of starch bandages.

## ANISI FRUCTUS

ANISE-FRUIT

**Synonym.**—Aniseed.

The fruit of *Pimpinella Anisum*. Cultivated in China, Japan, and Tartary.

**Natural Order.**—*Umbelliferæ*.

**Composition.**—Contains about 5 per cent. of a mixture of a fixed oil, a volatile oil,[1] and a volatile anethol or camphor-like body, common to the *Umbelliferæ*.

**Characters.**—Ovoid; composed of two mericarps with five primary ridges; slightly hairy; yellowish-brown colour, peculiar sweet aromatic odour, and a warm sweetish taste.

**Actions and Uses.**—Stomachic, carminative, aromatic. Used internally in indigestion and flatulency, to mask the disagreeable flavour of medicines, and to diminish the nauseating and griping effects of purgatives.

**Doses.**—Horse, ½ to 2 ounces.—Cattle, 1 to 2 ounces.—Sheep, 2 to 3 drachms.—Pig, 2 to 3 drachms.—Dog, 20 grains to 1 drachm.

**Modes of Application.**—The powdered fruit is administered alone, or, in conjunction with ginger or some other aromatic, in ale or in spirit and water.

[1] The volatile oil of anise, oleum anisi, is commonly added to masses as a flavouring ingredient, and is likewise employed to destroy pediculi on dogs and other small animals.

## ANTHEMIDIS FLORES

CHAMOMILE-FLOWERS

The dried single and double flower-heads of the common chamomile, *Anthemis nobilis*. Found in a wild state in all the temperate parts of Europe. Cultivated at Mitcham, in Derbyshire, and in other parts of England.

**Natural Order.**—*Compositæ.*

**Composition.**—The active principles are a bitter substance of unknown composition and a volatile oil.

**Characters.**—The single variety consists of both yellow tubular and white strap-shaped florets, the double of white strap-shaped florets only; all arising from a conical scaly receptacle. Both varieties, but especially the single, are bitter and very aromatic.

**Actions and Uses.**—*Internally*:—Tonic and carminative in indigestion, and during convalescence after inflammatory attacks. Not used externally.

**Doses.**—Horse, 2 to 6 drachms.—Cattle, 2 to 6 drachms.—Sheep, ½ to 2 drachms.—Pig, ½ to 2 drachms. Dog, 10 to 60 grains.

**Mode of Application.**—The dried flowers powdered, in combination with ginger or some other aromatic, once or twice a day.

**Incompatibles.**—Mineral acids and most metallic salts.

## ANTIMONII OXIDUM

OXIDE OF ANTIMONY

**Synonyms.**—Antimonious Oxide; Antimony Trioxide; Teroxide of Antimony.

**Composition.**—$Sb_2O_3$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Solution of Chloride of Antimony | 16 fluid ounces |
| Carbonate of Sodium . . . | 6 ounces |
| Common Water . . . . | 2 gallons |
| Distilled Water . . . . | a sufficiency |

Pour the chloride of antimony into the common water, and mix thoroughly; let the precipitate (antimony oxychloride) which forms settle, and, after removing the supernatant liquid by a siphon, add to it one gallon of distilled water, agitate well, let the precipitate subside; again withdraw the fluid and repeat the processes of affusion of distilled water, agitation, and subsidence. Now add the carbonate of sodium, previously dissolved in two pints of distilled water; leave in contact for half an hour, stirring frequently; collect the deposit on a calico filter, and wash it with boiling distilled water until the washings, after being acidulated by nitric acid, cease to give a precipitate with solution of silver nitrate. Dry the product (oxide of antimony) at a temperature not exceeding 212° F.

The oxide of antimony thus prepared is formed in two stages:

| | Antimony Trichloride. | | Water. | | Antimony Oxychloride. | | Hydrochloric Acid. |
|---|---|---|---|---|---|---|---|
| 1. | $SbCl_3$ | + | $H_2O$ | = | $SbOCl$ | + | $2HCl$ |

| | Antimony Oxychloride. | | Sodium Carbonate. | | Antimonious Oxide (Oxide of Antimony). | | |
|---|---|---|---|---|---|---|---|
| 2. | $2SbOCl$ | + | $Na_2CO_3$ | = | $Sb_2O_3$ | + | |

| Sodium Chloride. | | Carbonic Anhydride. |
|---|---|---|
| $2NaCl$ | + | $CO_2$ |

**Characters and Tests.**—Greyish-white powder, fusible at a low red heat; insoluble in water, but readily dissolved by hydrochloric acid. The hydrochloric solution,

dropped into distilled water, gives a white deposit (antimony oxychloride), which is immediately changed to orange (antimonious sulphide) by sulphuretted hydrogen. Should entirely dissolve when boiled with water and excess of potassium-hydrogen tartrate.

**Actions and Uses.**—Diaphoretic and febrifuge for the dog. Chiefly used in the preparation of tartarated antimony.

**Preparations.**—Pulvis Antimonialis; Antimonium Tartaratum.

## ANTIMONIUM NIGRUM

BLACK ANTIMONY

**Synonyms.**—Antimonious Sulphide; Antimony Trisulphide; Tersulphide of Antimony.

**Composition.**—$Sb_2S_3$.

**Mode of Preparation.**—Native antimonious sulphide is fused in a perforated crucible placed over another; the siliceous impurities are retained by the upper crucible, and the molten sulphide, which flows into the lower one, is solidified by cooling, and reduced to a fine powder.

**Characters and Tests.**—Greyish-black crystalline powder. It dissolves entirely in boiling hydrochloric acid, evolving sulphuretted hydrogen and furnishing a solution (antimonious chloride) which throws down a white precipitate (antimony oxychloride) when poured into water.

**Actions and Uses.**—Alterative, anthelmintic, and emetic. Its action is uncertain and irregular, and it should, therefore, not be employed as a remedial agent. Was formerly frequently given in combination with sulphur and nitrate of potassium as an alterative to horses. Also used in the preparation of Liquor Antimonii Chloridi.

# ANTIMONIUM TARTARATUM

## TARTARATED ANTIMONY

**Synonyms.**—Potassium-Antimony Tartrate; Tartrate of Potassium and Antimony; Tartar Emetic; Emetic Tartar; Tartarised Antimony.

**Composition.**—$[K(SbO)C_4H_4O_6]_2.H_2O$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Oxide of Antimony . . . . | 5 ounces |
| Acid Tartrate of Potash, in fine powder | 6 ounces |
| Distilled Water . . . . . | 2 pints |

Mix the oxide of antimony and acid tartrate of potash with sufficient distilled water to form a paste, and set aside for twenty-four hours; then add the remainder of the water, and boil for a quarter of an hour, stirring frequently. Filter, and set aside the clear liquid, which contains the tartarated antimony, to crystallise. Pour off the mother-liquor, evaporate to one-third, and set aside that more crystals may form. Dry the crystals (tartarated antimony) on filtering-paper at the temperature of the air.

| Antimonious Oxide. | | Potassium-hydrogen Tartrate. | | Potassium-antimony Tartrate (Tartarated Antimony). | | Water. |
|---|---|---|---|---|---|---|
| $Sb_2O_3$ | + | $2KHC_4H_4O_6$ | = | $2K(SbO)C_4H_4O_6$ | + | $H_2O$ |

**Characters and Tests.**—Colourless, transparent crystals, exhibiting triangular facets, soluble in water, and less so in proof spirit. Decrepitates, blackens, and emits an odour characteristic of tartrates when heated on platinum foil. Its solution in water gives, with hydrochloric acid, a white precipitate (antimony oxychloride) soluble in excess; if tartaric acid be previously added, no precipitate is formed by hydrochloric acid.

**Actions and Uses.**—*Internally* :—In excessive doses, irritant poison; in small doses it promotes the secretion of mucous membranes, skin, liver, pancreas, and kidneys, likewise stimulates the activity of the absorbent system; in larger doses it causes nausea, relaxes muscular fibres, depresses the nervous system, and, in animals capable of vomition, acts as an emetic. Used internally as an alterative, antipyretic, nauseant, sedative, and emetic. Given in the early stages of inflammatory diseases, in pneumonia, bronchitis, articular rheumatism, &c.; also employed as a vermifuge. *Externally* :—Vesicant, counter-irritant. Sometimes added to ordinary blistering ointments to increase their activity.

**Doses.**—*Alterative*:—Horse, ½ to 1 drachm.—Cattle, ½ to 1 drachm. *Antipyretic, Nauseant, Sedative* :—Horse, 1 to 2 drachms.—Cattle, 2 to 6 drachms.—Dog, ⅛ to 2 grains. *Emetic* :—Pig, 4 to 16 grains.—Dog, 1 to 4 grains. *Vermifuge* :—Horse, 1 to 2 drachms.[1]

**Modes of Application.**—*Internally* :—In the form of bolus or pill, or dissolved in water. As an emetic to dogs it may be given rolled up in a piece of meat. *Externally*:—Liquor Antimonii Tartarati; Unguentum Antimonii Tartarati.

**Incompatibles.**—Gallic and tannic acids, alkalies, and lead salts.

**Antidotes.**—Tannic acid or agents containing it—*e.g.*, powdered gall-nuts, catechu, &c.

**Preparations.**—Liquor Antimonii Tartarati; Pulvis Antimonialis; Unguentum Antimonii Tartarati.

[1] Should be given at night and followed by an aloetic purge next morning.

## ANTIPYRINE

ANTIPYRIN

**Synonyms.**—Phenazone; Phenazonum.

**Composition.**—Phenyl - dimethyl - iso - pyrazolone, $C_6H_5(CH_3)_2C_3HN_2O$.

**Preparation.**—Obtained from phenyl-hydrazine by interaction with aceto-acetic ether, and the subsequent interaction of the resulting phenyl-methyl-iso-pyrazolone with methyl iodide.

**Characters and Tests.**—It occurs in colourless, odourless, scaly crystals, with a bitter taste. Soluble in about one and a half part of water, alcohol, or chloroform, and in forty parts of ether. An aqueous solution yields a white precipitate with tannin, and is coloured yellow by concentrated nitric acid. Ferric chloride gives a dark brown colour. Spirit of nitrous ether gives a green colour.

**Actions and Uses.**—Antiseptic, antipyretic, and analgesic; it causes a decided lowering of temperature in fever. Anodyne in neuralgic and rheumatic affections, and in influenza in horses. Used sometimes in chorea.

**Doses.**—Horse, 2 to 4 drachms.—Cattle, 3 to 6 drachms.—Sheep, 1 drachm.—Dog, 15 to 30 grains.

**Modes of Application.**—In bolus, drench, electuary, or in the drinking-water.

## AQUA DESTILLATA

DISTILLED WATER

**Composition.**—$H_2O$.

**Mode of Preparation.**—Take of

Common Water . . . . 10 gallons

Distil from a copper still, connected with a block-tin

worm ; reject the first half-gallon, and preserve the next eight gallons.

**Tests.**—A fluid ounce evaporated in a clean glass capsule should leave scarcely any visible residue, indicating the absence of solid non-volatile impurities. Should not be affected by hydrosulphuric acid, ammonium oxalate, silver nitrate, barium chloride, or lime-water, proving the absence of lead and other metals precipitable by sulphuretted hydrogen, calcium salts, chlorides, sulphates, carbonates, and carbonic acid respectively.

**Uses.**—In the preparation of, and for dissolving, many medicinal agents ; also for making all aqueous solutions of tests. When distilled water cannot be obtained, its best substitute is boiled and filtered rain-water.

## AQUA MENTHÆ PIPERITÆ

PEPPERMINT-WATER

**Mode of Preparation.**—Take of

| | |
|---|---|
| Oil of Peppermint . . . | 1½ fluid drachm |
| Water . . . . . | 1½ gallon |

Distil one gallon.

**Actions and Uses.**—Carminative and stimulant. Used to relieve flatulence, and as a vehicle for other medicines.

**Doses.**—Horse, 8 to 10 fluid ounces.—Cattle, 8 to 12 fluid ounces.—Dog, 1 to 2 fluid ounces.

## ARECÆ SEMINA

ARECA-SEEDS

**Synonyms.**—Nux Areca Catechu ; Areca-nut ; Betel-nut.

The seeds or kernels of the fruit of the catechu or betel-nut palm, *Areca Catechu*.

**Natural Order.**—*Palmaceæ*.

**Composition.**—They contain, besides tannin, the alkaloids arecoline ($C_8H_{13}NO_2$), arecaine, arecaidine, and guacine.

**Characters.**—Spheroidal, about three-quarters of an inch in diameter; hard; colour reddish brown; taste astringent.

**Actions and Uses.**—*Internally*:—Is a powerful sialogogue, diaphoretic, intestinal stimulant, and vermifuge. In colic and fœcal impaction, it is almost as powerful as eserine, and stronger and more rapid than pilocarpine. It is an effective vermicide, especially for dogs, destroying both tape and round worms.

*Tæniline*.—A registered liquid preparation, which contains the tænifuge principle of freshly ground Areca-nut; this is a convenient and efficient vermicide.

**Doses.**—Horse, 4 to 6 drachms.—Cattle, 4 to 8 drachms.—Dog, 10 grains to 2 drachms.

**Modes of Application.**—The powdered seeds made into a bolus, or suspended in milk, gruel, linseed oil, or mucilage.

**Incompatibles.**—Same as Acidum Tannicum.

## ARGENTI NITRAS

### NITRATE OF SILVER

**Synonyms.**—Silver Nitrate; Lunar Caustic.

**Composition.**—$AgNO_3$.

**Mode of Preparation.**—Take of

| | | |
|---|---|---|
| Purified Silver | 3 | ounces |
| Nitric Acid | 2½ | fluid ounces |
| Distilled Water | 3 | fluid ounces |

Put the silver into a flask, pour on to it the nitric acid and water, and apply a gentle heat until the metal is dissolved. Decant the clear liquid from any black powder (gold) which may be present into a porcelain dish, evaporate, and set aside to crystallise; pour off the liquor, and again evaporate and crystallise. Let the crystals (nitrate of silver) drain in a glass funnel, and dry them by exposure to the air, carefully avoiding the contact of all organic substances. To obtain the nitrate in rods, fuse one part of the crystals in a platinum or thin porcelain capsule with two parts of potassium nitrate, and pour the melted salt into proper moulds. Nitrate of silver should be preserved in carefully stoppered bottles.

| Silver. | | Nitric Acid. | | Silver Nitrate. | | Nitric Oxide. | | Water. |
|---|---|---|---|---|---|---|---|---|
| $Ag_3$ | + | $4HNO_3$ | = | $3AgNO_3$ | + | $NO$ | + | $2H_2O$ |

**Characters and Tests.**—Colourless tubular crystals, the primal form of which is the right rhombic prism; or in white cylindrical rods. Soluble in water, rectified spirit, ether, and glycerin. A fragment heated on charcoal with the blowpipe melts, deflagrates, and leaves a white metallic coating of silver. Its aqueous solution, mixed with excess of dilute hydrochloric acid, yields a white precipitate (silver chloride), which darkens by exposure to light, and which should entirely dissolve in ammonium hydrate. If the white precipitate, with its accompanying liquid, be thrown on a filter and thoroughly washed with boiling distilled water, the filtrate should leave no residue on evaporation, proving the absence of potassium nitrate, lead, zinc, and copper.

**Actions and Uses.**—*Internally* :—In excessive doses, irritant and corrosive poison ; in medicinal doses, tonic and antispasmodic. It is sometimes given to the dog in chorea, epilepsy, and other nervous affections ; and has also been found of service in canine diarrhœa, dysentery, and cholera. *Externally* :—Stimulant, astringent, and caustic. Used as a collyrium in conjunctivitis and similar superficial inflammatory complaints ; for the improvement of indolent sores, mange, ringworm, and other chronic skin-diseases ; as a caustic for the eradication of warts, fungous and other growths. Applied also to poisoned wounds, and to the bites of rabid and venomous animals.

**Doses.**—Dog :—In chorea, epilepsy, &c., $\frac{1}{8}$ to $\frac{1}{2}$ grain ; in diarrhœa, dysentery, and cholera, $\frac{1}{4}$ to 1 grain.

These doses should be repeated two or three times a day.

**Modes of Application.**—*Internally* :—In the form of bolus or pill. *Externally* :—In the form of stick ; Liquor Argenti Nitratis ; Unguentum Argenti Nitratis.

**Incompatibles.**—Sulphuric, hydrochloric, and hydrocyanic acids ; alkalies ; and soluble chlorides.

**Antidote.**—Solution of common salt in some demulcent drink.

**Preparations.**—Liquor Argenti Nitratis ; Unguentum Argenti Nitratis.

## ARNICÆ RHIZOMA

ARNICA-RHIZOME

**Synonym.**—Arnicæ Radix.

The dried rhizome and rootlets of *Arnica montana*, or Leopard's Bane. Collected in the mountains of Middle and Southern Europe.

**Natural Order.**—*Compositæ.*

**Composition.**—The most important, because probably the active, constituents are a resin containing cytosin and arnicin, and a volatile oil.

**Characters.**—Rhizome from one to three inches long, and from one-tenth to three-tenths of an inch thick, cylindrical, contorted, rough from the scars of the coriaceous leaves, and furnished with numerous long, slender fibres ; has a peppery taste and peculiar odour.

**Actions and Uses.**—See Tinctura Arnicæ.

**Preparation.**—Tinctura Arnicæ.

## ASAFETIDA

ASAFETIDA

A gum resin obtained by incision from the living root of *Ferula fœtida*, and probably other species.

**Natural Order.**—*Umbelliferæ.*

**Composition.**—Its active constituents are resinous matter and a volatile oil containing allyl sulphide $(C_3H_5)_2S$.

**Characters.**—Irregular masses, partly composed of tears, moist or dry. The colour of a freshly cut or broken piece is opaque white, but gradually becomes purplish pink, and ultimately dull yellowish or pinkish brown. Taste bitter, acrid ; odour (due to allyl sulphide) fetid, alliaceous, and persistent. Dissolves almost entirely in rectified spirit. Powdered with difficulty, unless triturated with carbonate of potash.

**Actions and Uses.**—A moderate stimulant, powerful antispasmodic, expectorant, and vermifuge. Chiefly used in treating colic and coughs. Given also to dogs suffering from chorea and other nervous affections.

**Doses.**
**Modes of Application.** } Similar to Ammoniacum.

**Preparations.**—Enema Asafetidæ; Spiritus Ammoniæ Fœtidus.

## ATROPINA

ATROPINE

**Composition.**—$C_{17}H_{23}NO_3$.

An extremely poisonous crystallisable alkaloid, obtained from belladonna-root, and constituting the active principle of Belladonnæ Folia, Belladonnæ Radix, Extractum Belladonnæ, and Tinctura Belladonnæ.

**Preparations.**—Atropine sulphate. The B.P. liquor atropinæ sulphatis consists of one part atropine sulphate and ninety-nine of camphor-water; it contains four grains in one fluid ounce.

**Doses of the Liquor.**—Horse and Cattle, from 1 to 3 fluid drachms.—Dog, 1 to 3 minims.

One-fifth of these doses suffices when used hypodermically. Equal proportions of atropine sulphate and acetate of morphine solutions give prompt antispasmodic and anodyne effects when used conjointly.

**Ointment.**—Atropine dissolved in three and a half parts rectified spirit and fifty-five parts benzoated lard.

## BALSAMUM PERUVIANUM

BALSAM OF PERU

A balsam [1] obtained from *Myroxylon Pereiræ*, belonging to the natural order *Leguminosæ*. It exudes from

[1] A balsam is a natural mixture of essential oil and resinous substances.

the trunk of the tree after the bark has been scorched and removed. From Salvador in Central America.

**Composition.**—A mixture of volatile oil called cinnamein ($C_{27}H_{26}O_4$), cinnamic acid ($HC_9H_7O_2$), and resinous compounds.

**Characters.**—Reddish-brown or nearly black liquid, translucent in thin films; syrupy consistence, balsamic odour, and an acrid, slightly bitter taste; soluble in five parts of rectified spirit. Undergoes no diminution when mixed with water.

**Actions and Uses.**—Stimulant and antispasmodic. Used in the preparation of Tinctura Benzoini Composita (Friar's Balsam).

## BELLADONNÆ FOLIA

BELLADONNA-LEAVES

The fresh leaves, with the branches to which they are attached, of Deadly Nightshade, *Atropa Belladonna*; also the leaves separated from the branches, and carefully dried and finely powdered; gathered from wild or cultivated British plants when the fruit has begun to form.

**Natural Order.**—*Atropaceæ.*

**Characters.**—Leaves alternate, three to six inches long, ovate-acute, entire, smooth, the uppermost in pairs, and unequal. The expressed juice or an infusion, dropped into the eye, dilates the pupil.

**Actions and Uses.**—*Internally*:—In excessive doses, a narcotic-acrid poison; in medicinal doses, sedative, anodyne, and antispasmodic. In whatever doses it is administered, or by whatever channel it enters the circulation, it causes dilatation of the pupil. Given in colic,

acute and chronic rheumatism, bronchitis, influenza, and in cases of sore throat and cough which often accompany or succeed this last-named disease. *Externally* :—As a dressing for painful and irritable tumours; for tender enlarged glands; for cases of garget and sore throat. In the form of injection, for allaying irritation of the bladder or rectum, and counteracting spasmodic contractions of the uterus.

**Doses.**[1]—Horse and Cattle, 4 drachms to 1 ounce. —Sheep, 10 to 20 grains.—Pig, 8 to 12 grains.—Dog, 2 to 5 grains.

**Modes of Application.**—*Internally* :—In the form of bolus or pill. See also Extractum Belladonnæ; Tinctura Belladonnæ; Massa Belladonnæ Composita. As an injection :—See Extractum Belladonnæ.

**Antidotes.**—An emetic of sulphate of copper, when practicable; purgatives to empty the alimentary canal; after vomition or purgation, administer opium in sufficient doses to counteract the effects of the belladonna. Move the animal about; affusions of cold water over the head and chest; artificial respiration; galvanism; ether; brandy; inhalations of ammonia.

**Preparations.**—Extractum Belladonnæ; Tinctura Belladonnæ.

## BELLADONNÆ RADIX

BELLADONNA-ROOT

The dried root of *Atropa Belladonna*, or Deadly Nightshade. Cultivated in Britain or imported from Germany.

[1] Although the doses of the dried and powdered leaves are here given Extractum Belladonnæ or Tinctura Belladonnæ is to be preferred for internal use.

**Natural Order.**—*Atropaceæ.*

**Characters.**—From one to two feet long, and from half an inch to two inches thick; branched and wrinkled, brownish white. An infusion dilates the pupil.

**Preparation.**—Atropina.

## BENZOINUM

BENZOIN

**Synonym.**—Gum Benzoin.

A balsamic resin, obtained from *Styrax Benzoin*, by making incisions in the bark of the tree and allowing the liquid that exudes to concrete by exposure to the air. Imported from Siam and Sumatra.

**Natural Order.**—*Styraceæ.*

**Composition.**—Contains resins, benzoic acid ($HC_7H_5O_2$), volatile oil, and aromatic extract.

**Characters.**—Lumps consisting of agglutinated tears, or a brownish mottled mass with or without tears embedded in it; has little taste, but an agreeable odour; gives off, when heated, fumes of benzoic acid; soluble in rectified spirit and in solution of potash.

**Actions and Uses.**—*Internally* :—Stimulant and antispasmodic; but seldom or never given at the present day. *Externally* :—Excitant to wounds and contusions in the form of Tinctura Benzoini Composita (Friar's Balsam).

## BENZOL

BENZOL

**Synonym.**—Benzene.

A mixture of homologous hydrocarbons obtained from light coal-tar oil. It contains about 70 per cent. of benzene ($C_6H_6$) and 20 to 30 per cent. of toluene ($C_6H_5CH_3$).

**Characters.**—A colourless, volatile liquid free from opalescence, with a strong characteristic odour. Specific gravity from 0·880 to 0·888. It is inflammable, and burns with a smoky flame.

**Actions and Uses.**—Antiseptic, irritant, and parasiticide for various skin-parasites. It is usually applied after a thorough wash with soap and water, either undiluted or, in cats, dogs and fowls, mixed with one to three parts of oil or vaseline.

## BOLI

BALLS

See Massæ (Masses).

## BORAX

BORAX

**Synonyms.**—Sodium Diborate ; Biborate of Soda.

**Composition.**—Sodium Pyroborate, $Na_2B_4O_7.10H_2O$. Found native in the waters of certain lakes in Thibet and Persia. As crude Borax or Tincal, it is imported from Calcutta.

**Mode of Preparation.**—Jets of steam (*fumerolles* or *suffioni*) charged with boric acid, $H_3BO_3$, which issue from the earth in Tuscany, are conducted into *lagoni* (little lakes), in which the acid condenses. The solution thus formed, after being concentrated by heat, is neutralised with sodium carbonate, and, on evaporation, yields crystals of borax.

In the act of crystallising, the sodium diborate unites with ten molecules of water, and becomes $Na_2B_4O_7.10H_2O$.

**Characters and Tests.**—Transparent colourless

crystals, sometimes slightly effloresced, with a weak alkaline reaction ; insoluble in rectified spirit. Solubility 1 in 22 of cold water, and 1 in 2 of boiling water. By the aid of 1 of glycerin, 1 part of borax will dissolve in 12 of cold water. A hot saturated solution, when acidulated with any of the mineral acids, lets fall, as it cools, a scaly crystalline deposit of boric acid, $H_3BO_3$, the solution of which in spirit of wine burns with a green flame.

**Actions and Uses.**— Antiseptic, parasiticide ; slightly astringent and alkaline. Used to allay irritation of the skin and mucous membrane, and as a detergent and astringent in aphthous affections of the mouth.

**Mode of Application.**—A mixture of one part of finely powdered borax and six parts of honey or glycerin is applied to the part affected.

## BUCHU FOLIA

BUCHU-LEAVES

The dried leaves of *Barosma betulina*, *B. crenulata*, and *B. serratifolia*.

**Natural Order.**—*Rutaceæ*.

**Characters.**—The leaves are smooth, dull yellow-green, with a strong penetrating odour, a bitter, aromatic taste, and varying in different species from half an inch to one and a half inch in length. They contain a volatile oil, which is the most active constituent. A camphoraceous substance, barosma camphor, is deposited in the oil in the cold. They also contain mucilage.

**Actions and Uses.**—A mild tonic, diuretic, diaphoretic, but the principal use of this agent is for disorders of a catarrhal character affecting mucous membranes and diseases of the genito-urinary organs.

**Doses.** — HORSE or CATTLE, 1 to 4 ounces; for DOGS, 10 to 30 grains infused in a covered vessel with twenty parts of water for half an hour; this infusion is given with linseed-tea or barley-water. It may be advantageously combined with belladonna, opium, hyoscyamus, or potassium bromide.

## CALCII HYDRAS

### CALCIUM HYDRATE

**Synonyms.**—Calcium Hydroxide; Hydrate of Lime; Slaked Lime.

**Composition.**—$Ca(HO)_2$.

**Preparation.**—Take of

| | |
|---|---|
| Freshly burned Lime . . . . | 2 pounds |
| Distilled Water [1] . . . . | 1 pint |

Place the lime in a metal or earthenware vessel furnished with a lid; pour the water upon it, and when vapour ceases to be disengaged, cover the vessel with its lid, and set it aside to cool. When the temperature has fallen to that of the atmosphere, put the slaked lime on an iron-wire sieve, and, by gentle agitation, cause the fine powder to pass through the sieve, and reject what is left. Put the powder into a well-stoppered bottle, and keep it excluded as much as possible from the air, to prevent its absorbing carbonic anhydride. In the process of slaking, lime unites chemically with water.

| Calcium Oxide (Lime). | | Water. | | Calcium Hydrate. |
|---|---|---|---|---|
| $CaO$ | + | $H_2O$ | = | $Ca(HO)_2$ |

[1] Although distilled water is directed to be employed in this preparation by the *British Pharmacopœia*, common water may be used in its stead

**Characters and Tests.** — A white, caustic, and strongly alkaline powder. It is slightly soluble in water (1 in 800), more so in cold than in hot. Should not effervesce on the addition of an acid, showing its freedom from carbonate.

**Actions and Uses.**—*Internally* :—Irritant, corrosive, and antacid. Given chiefly to cattle in the form of Lime-water as an antacid in indigestion, diarrhœa, and tympanitis. Sometimes employed as an antidote to poisoning by arsenic and irritant acids. *Externally* :—Occasionally substituted for zinc oxide in cases of eczema.

**Doses.**—Horse, 1 to 3 drachms.—Cattle, 1 to 3 drachms.—Sheep, 20 grains to 1 drachm. Pig, 20 grains to 1 drachm.—Dog, 5 to 20 grains.

**Modes of Application.**—In the form of bolus or pill, or else suspended in water, milk, or some mucilaginous drink. Also administered as Liquor Calcis.

**Incompatibles.**—Mineral and vegetable acids ; metallic and ammoniacal salts ; tartarated antimony.

**Preparations.**—Linimentum Calcis ; Liquor Calcis ; Lotio Hydrargyri Flava ; Lotio Hydrargyri Nigra.

## CALUMBÆ RADIX

### CALUMBA-ROOT

**Synonym.**—Calumba.

The root of *Jateorhiza Columba*, sliced transversely and dried.

**Natural Order.**—*Menispermaceæ.*

**Characters and Tests.**—The root consists of several fasciculated, fusiform, fleshy tubercles, which are brown

externally and deep yellow internally, odourless, and very bitter. Moistened with a solution of iodine, it becomes blue, indicating presence of starch. A decoction is not blackened by the persalts of iron, indicating absence of astringent matter.

**Composition.**—The principal constituents of calumba are calumbin ($C_{21}H_{22}O_7$) and the alkaloid berberia ($C_{20}H_{17}NO_4$) combined with calumbic acid ($C_{22}H_{24}O_7$) and starch.

**Actions and Uses.**—Calumba-root is an excellent tonic promoting the appetite, assisting the digestive process, and improving the quality of the secretions from the gastro-intestinal mucous membranes. It does not appear either to constipate or to relax the bowels. The infusion is used to destroy worms in the horse's rectum.

**Doses of the Powder.**—Horse, 2 to 4 drachms.—Cattle, 2 to 6 drachms.—Sheep, 1 to 2 drachms.—Pig, ½ to 1 drachm.—Dog, 10 to 20 grains.

**Modes of Application.**—The root is best administered in the form of powder or infusion.

**Preparation.**—Infusum Calumbæ.

## CALX

### LIME

**Synonyms.**—Calcium Oxide; Oxide of Calcium; Quicklime.

**Composition.**—CaO.

**Mode of Preparation.**—Some form of calcium carbonate—*e.g.*, chalk, limestone, or marble—is calcined so

as to expel carbonic anhydride; lime, mixed with a small amount of impurity, remains.

| Calcium Carbonate. | | Calcium Oxide (Lime). | | Carbonic Anhydride. |
|---|---|---|---|---|
| $CaCO_3$ | = | $CaO$ | + | $CO_2$ |

**Characters and Tests.** — Compact masses of a whitish colour, which readily absorb water, and which, when rather less than their weight of water is added, crack and fall into powder with the development of much heat. The powder thus obtained, agitated with distilled water and filtered, yields a clear solution, having an alkaline reaction, and giving a white precipitate (calcium oxalate) with ammonium oxalate.

**Actions and Uses.**—*Externally* :—As a caustic and desiccant, dusted over the part affected.

**Preparation.**—Calcii Hydras.

## CALX CHLORINATA

### CHLORINATED LIME

**Synonyms.** — Calcium Hypochlorite ; Hypochlorite of Lime ; Chloride of Lime.

**Mode of Preparation.**—Slaked lime is exposed in stone chambers to the action of chlorine gas as long as the latter is absorbed. Assuming that the activity of chlorinated lime depends upon its containing calcium hypochlorite, its production may be thus explained :

| | | | | Chlorinated Lime. | | | |
|---|---|---|---|---|---|---|---|
| Calcium Hydrate. | | Chlorine. | | Calcium Hypochlorite. | Calcium Chloride. | | Water. |
| $2Ca(HO)_2$ | + | $2Cl_2$ | = | $Ca(ClO)_2$ | $CaCl_2$ | + | $2H_2O$ |

**Characters and Tests.**—Dull white powder, with a

feeble odour of chlorine; partially soluble in water. It evolves chlorine on being mixed with an acid. Its aqueous solution, mixed with a dilute mineral acid, bleaches indigo.

**Actions and Uses.**—*Internally*:—In tympanitis or hoven, to absorb the carbonic anhydride and decompose the sulphuretted hydrogen which are developed in these affections. Seldom given internally. *Externally*:—Disinfectant, deodoriser, and stimulant. Applied to gangrenous wounds, fistulæ, phagedænic ulcers, thrush, canker, grease, and to the treatment of mange and other skin-diseases. Also employed as a lotion for checking conjunctival ophthalmia, and other circumscribed and superficial inflammations.

**Doses.**—Horse, 1 to 2 drachms.—Cattle, 2 to 4 drachms.—Sheep, 1 to 2 drachms.

**Modes of Application.**—*Internally*:—Suspended in water, or, preferably, as Liquor Calcis Chlorinatæ. *Externally*:—Liquor Calcis Chlorinatæ; Unguentum Calcis Chlorinatæ.

**Incompatibles.**—Acids and most metallic salts.

**Preparations.**—Liquor Calcis Chlorinatæ; Unguentum Calcis Chlorinatæ; Vapor Chlori.

## CAMBOGIA

### GAMBOGE

A gum-resin obtained from *Garcinia Hanburii*, belonging to the natural order *Guttiferæ*. Imported from Siam, Singapore, and Ceylon.

**Composition.**—From 15 to 20 per cent. of soluble gum and about 70 per cent. of an active orange-yellow resin.

**Characters and Tests.**—Cylindrical pieces, breaking easily with a smooth, conchoidal, glistening fracture; colour tawny, changing to yellow when rubbed with water; taste acrid.

If adulterated with starch, an emulsion, made with boiling water and cooled, becomes green if treated with solution of iodine.

**Actions and Uses.**—In large doses, irritant poison; in medicinal doses, a drastic hydragogue cathartic. Should not be given to horses or dogs, on account of the uncertainty and violence of its action. To cattle and sheep it is sometimes administered in indigestion; also, combined with aloes or sulphate of magnesia, in impaction of the omasum (*fardel bound*) and similar disorders.

**Doses.**—Cattle, ½ to 1 ounce.—Sheep, 15 to 30 grains.

**Mode of Application.**—Given with other purgatives in solution.

**Antidotes.**—Demulcent drinks, and small but repeated doses of opium.

## CAMPHORA

CAMPHOR

A white crystalline substance, obtained in China and Japan from the wood and other parts of *Cinnamomum Camphora*, purified by sublimation.

**Natural Order.**—*Lauraceæ*.

**Composition.**—$C_{10}H_{16}O$.

**Mode of Preparation.**—The small branches, leaves, wood, and root of the tree are cut into small pieces and boiled with water in an iron vessel to which an earthen

hood is luted; the camphor sublimes and condenses on straws placed within the hood. The crude product thus obtained, after its importation into Europe, is mixed with quicklime to retain impurities, and sublimed in glass vessels.

**Characters and Tests.**—White, transparent, tough,[1] and crystalline; also in rectangular tablets, or in pulverulent masses, known as 'flowers of camphor.' Has a powerful penetrating odour and a pungent taste, followed by a sensation of cold; floats on water; volatilises slowly at ordinary temperatures; slightly soluble in water, readily soluble in rectified spirit and in ether; sublimes entirely when heated.

**Actions and Uses.**—*Internally*:—In large doses, irritant and narcotic poison; in medicinal doses, stimulant, sedative, antispasmodic, anodyne, and aphrodisiac; also, after long exhibition, diuretic. Occasionally given in tympanitis, and, combined with opium or digitalis, in chronic cough. *Externally*:—Discutient and anodyne for chronic sprains, bruises, and tumours, also for infiltration into the cellular tissue. Added to oil or ointment of cantharides, or sprinkled in fine powder over linseed poultices, it allays irritation, although for this purpose extract of belladonna is usually preferred.

**Doses.**—Horse, 1 to 2 drachms.—Cattle, 2 to 4 drachms.—Sheep, 10 to 40 grains.—Pig, 5 to 20 grains.—Dog, 3 to 10 grains.

**Modes of Application.**—*Internally*:—Spiritus Camphoræ; Oleum Camphoratum; or in the form of ball or pill. *Externally*:—Linimentum Camphoræ; Linimentum

[1] Camphor is difficult to pulverise unless a little spirit of wine be added to it.

Terebinthinæ; Linimentum Terebinthinæ Aceticum; Linimentum Saponis; Tinctura Camphoræ.

**Incompatibles.**—Camphor is not likely to be administered with anything that would counteract its effects.

### Preparations

Linimentum Aconiti.
„ Camphoræ.
„ Iodi Compositum.
„ Opii.
Linimentum Saponis Compositum.
„ Terebinthinæ.
Spiritus Camphoræ.

## CANTHARIS

CANTHARIDES

**Synonyms.**—The Dried Beetle; Blistering Fly; Spanish Fly; Cantharis Vesicatoria; Lytta Vesicatoria.

**Class.**—*Insecta.* **Natural Order.**—*Coleoptera.*

**Composition.**—Contains an active principle called cantharidin, $C_{10}H_{12}O_4$, animal matters, acetic and uric acids, and two oils.

The beetles are collected, chiefly in Hungary, at night time, by shaking the trees on which they feed, and so causing them to fall into cloths placed beneath the trees for their reception. After being transferred to sieves, they are killed by the vapour of vinegar or turpentine, or by being placed in a vacuum, and finally dried.

**Characters and Tests.**—From eight-tenths to an inch in length; furnished with two wing-covers of a shining metallic-green colour, under which are two membranous transparent wings; odour, strong and disagreeable; powder greyish brown, containing shining green particles. Should be free from mites.

**Actions and Uses.**—*Internally*:—In large doses, irritant poison; in medicinal doses, stimulant, diuretic. With vegetable bitters, sometimes used as a stimulating tonic in cases of debility, accompanied or not with anasarca; also in farcy, glanders, and some other affections. As cantharides are liable to inflame the neck of the bladder and cause strangury, they are seldom employed as a diuretic. *Externally*:—Rubefacient, vesicant, counter-irritant, stimulant. Employed as a blister, and for charging materials used for setons; also to promote the adhesion of unhealthy wounds and fistulæ; to stimulate weak and callous ulcers; and to induce an improved condition of skin in mallenders, ringworm, and inveterate mange; likewise to promote the growth of hair.

**Doses.**—Horse, 5 to 10 grains.—Cattle, 10 to 20 grains.—Sheep, 2 to 8 grains.—Pig, 1 to 4 grains.—Dog, $\frac{1}{4}$ to 1 grain.

**Modes of Application.**—*Internally*:—In a bolus, or as Tinctura Cantharidis. *Externally*:—To keep up discharges, also to increase the activity of mustard poultices and similar applications, the powdered insect is used. To promote the growth of hair, a solution of one part of cantharides in twenty parts of acetic acid, or an ointment consisting of one part of cantharides and twenty parts of lard, is to be applied with friction. For the other applications of cantharides, see the 'Preparations' enumerated below.

## Preparations

Acetum Cantharidis.
Linimentum Canthardis.
Liquor Cantharidis Terebinthinatus.
Oleum Cantharidis.
Tinctura Cantharidis.
,, ,, Fortior.
Unguentum Cantharidis.

## CAPSICI FRUCTUS

CAPSICUM-FRUIT

The dried ripe fruit of *Capsicum minimum.* Imported from Zanzibar, and, in a pulverised state, in gourds from the West Indies. It is distinguished in commerce as Guinea Pepper and Pod Pepper; when powdered, and mixed with a fourth part of common salt, it forms the well-known condiment 'Cayenne Pepper.'

**Natural Order.**—*Solanaceæ.*

**Composition.**—The fruit contains an alkaloid, capsicine, and a crystallisable solid oil, capsaicin. The latter body is very acrid, and constitutes the active principle of capsicum.

**Characters and Tests.**—Pod membranous, from five- to eight-tenths of an inch long, two-tenths of an inch broad; straight, conical, pointed, smooth, shining, but somewhat corrugated; orange-red; intensely hot taste.

**Actions and Uses.**—*Internally*:—In large doses, irritant poison; in medicinal doses, stimulant and carminative. *Externally*:—Rubefacient and vesicant.

**Doses.**—Horse, 5 to 20 grains.—Cattle, 10 to 30 grains.—Sheep.—5 to 10 grains.—Pig, 2 to 8 grains.—Dog, 1 to 5 grains.

**Modes of Application.**—*Internally*:—The powdered fruit in a bolus, or, preferably, suspended in gruel. *Externally*:—In the form of ointment.

## CARBO LIGNI

WOOD CHARCOAL

**Composition.**—Carbon mixed with from 1 to 5 per cent. of ash.

**Mode of Preparation.**—By exposing various kinds of wood to a red heat without access of air.

**Characters.**—Black, brittle, porous masses, tasteless, odourless, very light; retains the shape and texture of the wood from which it was obtained. When burned with free access of air it should not leave more than 2 per cent. of ash.

**Actions and Uses.**—Deodoriser, disinfectant, desiccant. *Internally*:—In dyspepsia, accompanied by flatus, to absorb the gases causing gastric distension; in diarrhœa and dysentery, to correct the fœtor of the evacuations; as an antidote to arsenic, aconite, strychnine, and probably to most vegetable poisons.[1] *Externally*:—As a desiccant and deodoriser to unhealthy wounds and phagedænic ulcers, especially when accompanied by fetid and irritating discharges; also in cases of mange, scab, and grease.

Wood charcoal, particularly when strongly heated and subsequently cooled just before being used, is a valuable deodoriser if strewn over the floors of a stable, &c., or sprinkled over putrefying substances.

**Doses.**—Horse, ½ to 1 ounce.—Cattle, 1 to 2 ounces. Sheep, 1 to 3 drachms.—Pig, 1 to 3 drachms.—Dog, 20 to 60 grains.

**Modes of Application.** — *Internally*: — Pulverised and mixed with gruel or some mucilaginous drink. *Externally*:—Dusted over the part in the state of powder, or as Cataplasma Carbonis.

**Preparation.**—Cataplasma Carbonis.

[1] Animal charcoal (*carbo animalis*) is to be preferred to wood charcoal in consequence of its greater absorptive power for the poisons above referred to.

## CARDAMOMI SEMINA

CARDAMOM-SEEDS

The dried capsules of the Malabar cardamom, *Elettaria Cardamomum*. Cultivated in Malabar. The seeds are best kept in their pericarps, from which they should be separated when required for use, the pericarpal coats being rejected.

**Natural Order.**—*Zingiberaceæ.*

**Composition.**—The therapeutic activity of cardamoms is due to a volatile oil contained in the seeds.

**Characters.**—Seeds obtusely angular, corrugated, reddish brown, internally white, with a warm, aromatic, agreeable taste and odour; contained in ovate-oblong, triangular, pale brown, coriaceous, ribbed pericarps.

**Actions and Uses.**
**Doses.**
**Modes of Application.**
} Same as Anisi Fructus.

## CARUI FRUCTUS

CARAWAY-FRUIT

**Synonym.**—Caraway-seeds.

The dried fruit of *Carum Carvi*. Cultivated in England and Germany.

**Natural Order.**—*Umbelliferæ.*

**Composition.**—The active principle is a volatile oil.

**Characters.**—Fruit usually separating into two parts, which are about two-tenths of an inch long, curved, tapering at each end, brown, with five paler longitudinal ridges; agreeable aromatic odour and spicy taste.

**Actions and Uses.**
**Doses.**
**Modes of Application.** } Same as Anisi Fructus.

## CARYOPHYLLUM

CLOVES

The dried unexpanded flower-buds of *Eugenia Caryophyllata*. Cultivated in Penang, Bencoolen, and Amboyna.

**Natural Order.**—*Myrtaceæ.*

**Composition.**—The active principle is a volatile oil.

**Characters and Test.**—About six-tenths of an inch long; dark reddish brown; plump and heavy; consisting of a nearly cylindrical body surmounted by four teeth and a globular head; strong fragrant odour, and a bitter, spicy, pungent taste. Emits oil when indented with the finger-nail.

**Actions and Uses.**
**Doses.**
**Modes of Application.** } Same as Anisi Fructus.

**Preparation.**—Oleum Caryophylli.

## CASCARILLÆ CORTEX

CASCARILLA-BARK

The bark of *Croton Eluteria*. From the island of Eluthera, one of the Bahamas.

**Natural Order.**—*Euphorbiaceæ.*

**Composition.**—Its therapeutic actions depend upon a crystalline bitter principle (cascarillin) and a volatile oil.

**Characters.**—Quills, two or three inches long, and from two- to five-tenths of an inch in diameter; dull brown,

but more or less coated with white crustaceous lichens; breaks with a short resinous fracture; warm and bitter taste, and emits a fragrant odour when burned.

**Actions and Uses.**—Aromatic, tonic, and slightly astringent. Sometimes given in indigestion, diarrhœa, chronic typhous affections, and during convalescence from debilitating diseases.

**Doses.**—Horse, 2 to 4 drachms.—Cattle, 3 to 6 drachms.—Sheep, 1 to 2 drachms.—Pig, 1 to 2 drachms. Dog, $\frac{1}{4}$ to 1 drachm.

**Mode of Application.**—In the form of ball.

## CATAPLASMA CALCIS CHLORINATÆ

### CHLORINATED LIME POULTICE

Take of

| | |
|---|---|
| Solution of Chlorinated Lime | 2 fluid ounces |
| Linseed Meal | 4 ounces |
| Boiling Water | 8 fluid ounces |

Mix the linseed meal gradually with the water, and add the solution of chlorinated lime, with constant stirring.

**Use.**—Deodoriser, antiseptic, and excitant to unhealthy wounds and ulcers.

## CATAPLASMA CARBONIS

### CHARCOAL POULTICE

Take of

| | |
|---|---|
| Wood Charcoal, in powder | $\frac{1}{2}$ ounce |
| Linseed Meal | $3\frac{1}{2}$ ounces |
| Boiling Water | 10 fluid ounces |

Add the linseed meal to the water, and stir them together so that a soft poultice may be formed. Mix with this half the charcoal, and sprinkle the remainder on the surface of the poultice.

**Use.**—Applied while warm to absorb the fœtor of foul ulcers.

## CATAPLASMA FERMENTI

YEAST POULTICE

Take of

| | |
|---|---|
| Beer Yeast . . . . | 6 fluid ounces |
| Wheaten Flour . . . | 14 ounces |
| Water heated to 100° F. . . | 6 fluid ounces |

Mix the yeast with the water, and stir in the flour. Place the mass near the fire till it rises.

**Use.**—Stimulant and antiseptic to indolent ulcers.

## CATAPLASMA FURFURIS

BRAN POULTICE

Take of

| | |
|---|---|
| Bran . . . . . | 2 to 3 parts |
| Linseed Meal . . . . | 1 part |
| Boiling Water . . . . | a sufficiency |

Mix the bran and linseed meal together, and then add the water, with constant stirring, so as to form a soft poultice.

**Use.**—Emollient; relaxes the vessels of the part to which it is applied, and thereby allays pain and inflammation.

## CATAPLASMA LINI

### LINSEED POULTICE

Take of

| | |
|---|---|
| Linseed Meal . . . . | 4 ounces |
| Olive Oil . . . . . | ½ fluid ounce |
| Boiling Water . . . . | 10 fluid ounces |

Mix the linseed meal gradually with the water, and then add the oil, with constant stirring.

**Use.**—Same as Cataplasma Furfuris. Opium, belladonna, or subacetate of lead may be added to this poultice when it is desired to allay irritation.

## CATAPLASMA SINAPIS

### MUSTARD POULTICE

Take of

| | |
|---|---|
| Mustard,[1] in powder . . | 2½ ounces |
| Linseed Meal . . . . | 2½ ounces |
| Warm [2] Water . . . . | 10 fluid ounces |

Mix the linseed meal gradually with the water, and add the mustard, with constant stirring.

**Use.**—Rubefacient to allay inflammations of serous and mucous surfaces, to relieve congestion of various organs, and to alleviate neuralgic and other pains and spasms.

[1] See note to *Sinapis*.

[2] Boiling water should not be used, as a temperature of 212° F. prevents the formation of the vesicating principle, allyl sulphocyanide.

## CATAPLASMA SINAPIS AMMONIATUM

### AMMONIATED MUSTARD POULTICE

Take of

Mustard, in powder
Solution of Ammonia } of each a sufficiency

Mix so as to form a poultice.[1]

**Use.**—Similar to, but much more powerful than, Cataplasma Sinapis.

## CATECHU PALLIDUM

### PALE CATECHU

**Synonym.**—Terra Japonica.

An extract of the leaves and young shoots of *Uncaria Gambier*. Prepared at Singapore and in other places in the Eastern Archipelago.

**Natural Order.**—*Rubiaceæ.*

**Composition.** — Mainly consists of catechin, $C_{13}H_{12}O_5$, and tannic acid.

**Mode of Preparation.** — The leaves of *Uncaria Gambier* are boiled in water immediately after they are pulled from the tree, and the decoction concentrated and run into square moulds, to form the commercial cubes of catechu.

**Characters.**—Cubes, or masses formed of coherent cubes; the former about an inch in diameter, externally brown, internally ochrey yellow or pale brick-red. Breaks easily with a dull earthy fracture. Taste bitter, very

[1] Oil of turpentine is sometimes added to this poultice, but such addition should be omitted in treating affections of the kidneys.

astringent, and mucilaginous, succeeded by slight sweetness. Entirely soluble in boiling water. The decoction, when cool, should not be rendered blue by iodine, indicating the absence of starch.

**Actions and Uses.** — Astringent. *Internally*:—Given in atony and relaxation, and in excessive mucous discharges, especially from the alimentary canal; also in diarrhœa. *Externally*:—To sluggish sores and ulcerations, for excoriations on the udder, &c.

**Doses.**—Horse, 1 to 3 drachms.—Cattle, 2 to 6 drachms.—Sheep, 1 to 2 drachms—Pig, 10 to 20 grains.—Dog, 1 to 10 grains. Three or four times a day.

**Modes of Application.** — *Internally*: — Massa Catechu Composita; Infusum Catechu; Tinctura Catechu; or the powder in sufficient gruel or mucilage to mask its astringent flavour. Catechu is likewise given with aromatics to remove flatulency, with opium to allay irritability, and with magnesia or an alkaline carbonate to correct acidity. *Externally*:—The powder; Infusum Catechu; and, occasionally, Unguentum Catechu.

**Incompatibles.**—Alkalies (?); most metallic salts; gelatin.

**Preparations.**—Infusum Catechu; Massa Catechu Composita; Tinctura Catechu.

## CERA FLAVA

### YELLOW WAX

**Synonym.**—Unbleached Beeswax.

A secretion of certain glands (*wax pockets*) situated on the abdomen of the common bee, *Apis mellifica*, and used by the insect for constructing the cells of the honeycomb.

**Composition.**—A mixture of myricin, cerin, and cerolein.

**Mode of Preparation.** — The comb, after being freed from honey by dripping and pressure, is melted in hot water to free it from impurities: the residue constitutes yellow wax. White wax (*Cera Alba*) is obtained by agitating molten yellow wax with water, and bleaching in the open air.

**Characters.**—Firm, breaking with a granular fracture; yellowish, and having an agreeable honeylike odour. Not unctuous to the touch; does not melt at 140° F.; yields nothing to rectified spirit, but is entirely soluble in oil of turpentine. Boiling water, in which it has been agitated, when cooled should not be rendered blue by iodine, showing its freedom from starch.

**Actions and Uses.**—Emollient, and to increase consistency of ointments.

**Preparation.**—Emplastrum Picis.

## CEREVISIÆ FERMENTUM

BEER YEAST

The ferment obtained in brewing beer.

**Characters.** — Viscid, semi-fluid, frothy, exhibiting under the microscope numerous round or oval confervoid cells, *Torula cerevisiæ*.

**Preparation.**—Cataplasma Fermenti.

## CERII OXALAS

OXALATE OF CERIUM

**Composition.**—$CeC_2O_4.9H_2O$.

**Mode of Preparation.**—By adding oxalic acid to a solution of a cerium salt, cerium oxalate is precipitated.

**Characters and Tests.**—A white granular powder, odourless and tasteless, insoluble in water and alcohol, soluble in hydrochloric acid.

**Uses.**—*Internally*:—It is sometimes given in the treatment of chronic bronchitis, and chorea.

## CHLORAL HYDRAS

### CHLORAL HYDRATE

**Synonyms.**—Hydrate of Chloral; Trichloraldehyd; Trichlorethylidine-glycol.

**Composition.**—$CCl_3.COH.(OH)_2$.

**Mode of Preparation.**[1]—Dry chlorine gas is passed for several days through absolute alcohol, which is at first kept cool to prevent explosion, but which, towards the end of the operation, is heated to nearly its boiling-point to complete the decomposition. The resulting liquid (impure chloral, $C_2HCl_3O$) is exposed to the air for a day or two, when it absorbs water and becomes converted into a solid mass of crude chloral hydrate. This is purified by agitating it with four times its bulk of concentrated sulphuric acid, separating the anhydrous chloral which floats on the surface, and subjecting it to fractional distillation. The anhydrous chloral thus purified is placed in a still with 11 per cent. of water, and distilled off chalk to remove any hydrochloric acid that may be present. The solid distillate (chloral hydrate) obtained in this manner is fused and cast in shallow vessels into cakes.

| | Alcohol. | | Chlorine. | | Chloral. | | Hydrochloric Acid. |
|---|---|---|---|---|---|---|---|
| 1. | $C_2H_6O$ | + | $4Cl_2$ | = | $C_2HCl_3O$ | + | $5HCl$. |

| | Chloral. | | Water. | | Chloral Hydrate. |
|---|---|---|---|---|---|
| 2. | $C_2HCl_3O$ | + | $H_2O$ | = | $C_2HCl_3(HO)_2$ |

[1] Dr. Squire's process.

**Characters and Tests.**—White, opaque, crystalline solid; pungent odour, resembling that of a ripe melon; soluble in water, alcohol, and glycerin. One hundred grains dissolved in half an ounce of distilled water, well shaken with one ounce of liquor potassæ, and allowed to stand several hours in a well-stoppered bottle, will, if pure, yield a layer of at least forty-six grain measures of chloroform.

**Actions and Uses.** — *Internally* : — In excessive doses, narcotic poison; in medicinal doses, stimulant, sedative, anodyne, antispasmodic, and narcotic; destroys micro-organisms. Said to possess the good, but none of the objectionable, qualities of opium; successfully used in milk fever in cattle.[1]

**Doses.**—Horse, 1 to 2 ounces.—Cattle, 1 to 2 ounces.—Sheep, 1 to 2 drachms.—Pig, 1 to 2 drachms.—Dog, 10 to 30 grains.

**Mode of Application.**—Dissolved in a large quantity of water.

**Preparation.**—Syrupus Chloralis Hydratis.

## CHLOROFORMI MISTURA HYDROCYANATA

### COMPOUND CHLOROFORM MIXTURE

**Synonym.**—Chlorodyne.

**Composition.**—The two following formulæ may be used in the preparation of chlorodyne:

1. Morphine sulphate 4 grains, oil of peppermint 8 grains, ether 1 drachm, alcohol 1 drachm, hydrocyanic

[1] Dr. Tuke, in speaking of its effects on maniacal patients, says: 'Its advantages over other hypnotics are as follows: it is more uniform in its action and its effects are more lasting; it has no depressing influence; it does not constipate or produce nausea.'

acid (dilute) 5 drachms, chloroform 3 ounces, syrup to 17 ounces.

2. Ten parts each of chloroform, ether, Indian hemp, and morphine; two parts of tincture of capsicum and prussic acid; three parts of tinctures of aconite and hyoscyamus; one part of oil of peppermint, five parts of hydrochloric acid, and fifty parts of simple syrup.

**Actions and Uses.**—Anodyne, antispasmodic, and to relieve gastro-intestinal and bronchial irritation.

**Doses.**—Horse, 2 to 4 drachms.—Dog, 20 to 40 minims.

## CHLOROFORMUM

CHLOROFORM

**Synonyms.**—Formyl Trichloride; Trichloromethane.

**Composition.**—$CHCl_3$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Chlorinated Lime | 10 pounds |
| Rectified Spirit | 30 fluid ounces |
| Slaked Lime | a sufficiency |
| Water | 3 gallons |
| Sulphuric Acid | a sufficiency |
| Chloride of Calcium, in small fragments | 2 ounces |
| Distilled Water | 9 fluid ounces |

Place the water and the spirit in a capacious still, and raise the mixture to the temperature of 100° F. Add the chlorinated lime and five pounds of the slaked lime, and mix thoroughly. Connect the still with a condenser terminating in a narrow-necked receiver, and apply heat so as to cause distillation, taking care to withdraw the fire the moment that the process is well established. When

the distilled product measures fifty ounces, the receiver is to be withdrawn. Pour its contents into a gallon bottle half filled with water, mix well by shaking, and set at rest for a few minutes, when the mixture will separate into two layers of different densities. Let the lower layer, which constitutes crude chloroform, be washed by agitating it in a bottle with three ounces of the distilled water. Allow the chloroform to subside, withdraw the water, and repeat the washing with the rest of the distilled water, in successive quantities of three ounces at a time. Agitate the washed chloroform for five minutes in a bottle with an equal volume of sulphuric acid, allow the mixture to settle, and transfer the upper layer of liquid to a flask containing the chloride of calcium mixed with half an ounce of perfectly dry slaked lime, and mix well by agitation. After the lapse of an hour, connect the flask with a Liebig's condenser, and distil over the pure chloroform by means of a water-bath. Preserve the product in a cool place in well-stoppered bottles. The lighter liquid which floats on the crude chloroform after its agitation with water, and the washings with distilled water, should be preserved and employed in a subsequent operation.

The exact nature of the chemical changes which occur in the production of chloroform by the foregoing process is still undetermined, but the following equations represent the probable reactions which take place :

| | Alcohol. | | Oxygen from the Calcium Hypochlorite (Chlorinated Lime). | | Aldehyd. | | Water. |
|---|---|---|---|---|---|---|---|
| 1. | $2C_2H_6O$ | + | $O_2$ | = | $2C_2H_4O$ | + | $2H_2O$ |
| | Aldehyd. | | Chlorine from the Calcium Hypochlorite (Chlorinated Lime). | | Chloral. | | Hydrochloric Acid. |
| 2. | $C_2H_4O$ | + | $3Cl_2$ | = | $C_2HCl_3O$ | + | $3HCl$ |
| | Chloral. | | Calcium Hydrate. | | Chloroform. | | Calcium Formate. |
| 3. | $3C_2HCl_3O$ | + | $Ca(HO)_2$ | = | $2CHCl_3$ | + | $Ca(CHO_2)_2$ |

**Characters and Tests.**—Limpid, colourless liquid, of an agreeable ethereal odour and sweet taste. Soluble in alcohol and ether in all proportions ; dissolves slightly in water, communicating to it a sweetish taste. Burns, though not readily, with a green and smoky flame. Sp. gr. 1·490 to 1·495. Should not be coloured by agitation with sulphuric acid, and should leave no residue or unpleasant odour after evaporation.

**Actions and Uses.**—In excessive doses, whether the liquid be swallowed or the vapour inhaled, narcotic poison ; in medicinal doses, stimulant and antispasmodic ; breathed in admixture with air, anæsthetic.

**Doses.**—See Spiritus Chloroformi.

**Modes of Application.**—As an antispasmodic : Spiritus Chloroformi. As an anæsthetic, a sponge saturated with *pure* chloroform is placed in the bottom of a nose-bag attached to the patient's head, and perforated so as to insure the chloroform-vapour being mixed with a sufficiency of air.

**Antidote.**—Fresh air.

**Preparations.** — Spiritus Chloroformi; Tinctura Chloroformi Composita.

## CINCHONÆ FLAVÆ CORTEX

YELLOW CINCHONA BARK

The bark of *Cinchona Calisaya*. Collected in Bolivia and Southern Peru.

**Natural Order.**—*Rubiaceæ*.

**Composition.**—Its therapeutic value depends upon the presence of quinine ($C_{20}H_{24}N_2O_2$) and cinchonine

($C_{20}H_{24}N_2O$), which, with several other alkaloids, exist in the bark in combination with kinic, kinovic, and tannic acids.

**Mode of Preparation.**—The trees are generally cut down, and the bark, after removal from the stem and branches, is carefully dried, so that it may retain its bright colour; the larger and thicker portions are dried so as to form flat pieces (chips), while the smaller are allowed to curl into quills.

**Characters.**—Flat pieces, coated with the periderm, and as quills, from six to eighteen inches long, one to three inches wide, and two- to four-tenths of an inch thick, compact and heavy; outer surface brown, marked by broad, shallow, irregular, longitudinal depressions; inner surface tawny yellow and fibrous; transverse fracture shortly and finely fibrous. Powder cinnamon brown, somewhat aromatic, persistently bitter.

**Test.**—Boil one hundred grains of the finely powdered bark, for a quarter of an hour, in a fluid ounce of distilled water, acidulated with ten minims of hydrochloric acid, and allow it to macerate for twenty-four hours. Transfer the whole to a small percolator, and, after the fluid has ceased to drop, add at intervals about an ounce and a half of similarly acidulated water, or until the fluid which passes through is free from colour. Add to the percolated fluid solution of subacetate of lead until the whole of the colouring matter has been removed, taking care that the fluid remains acid in reaction. Filter and wash with a little distilled water. To the filtrate add about thirty-five grains of caustic potash, or as much as will cause the precipitate which is at first formed to be nearly re-dissolved, and afterwards six fluid drachms of pure ether. Then

shake briskly, and, having removed the ether, repeat the process twice with three fluid drachms of ether, or until a drop of the ether employed leaves, on evaporation, scarcely any perceptible residue. Lastly, evaporate the mixed ethereal solutions in a capsule at the temperature of boiling water. The residue, which consists of nearly pure quinia, when dry, should weigh not less than two grains, and should be readily soluble in dilute sulphuric acid.

**Actions and Uses.**—*Internally* :—Tonic and astringent. Given in debility ; in periodic and intermittent diseases—*e.g.*, spasm, neuralgia, and periodical hæmorrhage ; in continued fever, rheumatism, and erysipelas ; in chronic atonic affections of the alimentary canal—*e.g.*, dyspepsia ; in chorea, passive hæmorrhages, profuse mucous discharges, leucorrhœa, and chronic diarrhœa ; in enlargements and indurations of the absorbent glands in scrofula ; in convalescence after either acute or lingering maladies ; and during debility after surgical operations. *Externally* : Astringent and antiseptic in leucorrhœa and to fetid ulcers.

**Doses.**—Horse, 2 to 4 drachms.—Cattle, 1 to 2 drachms.—Sheep, 1 to 2 drachms.—Pig, 1 to 2 drachms. Dog, ½ to 1 drachm. Twice or thrice a day. Sometimes the administration of cinchona to dogs is followed by nausea or vomition ; in such cases the dose should be greatly diminished.

**Modes of Application.**—*Internally* :—The powdered bark, in the form of ball ; Infusum Cinchonæ Flavæ ; Tinctura Cinchonæ Flavæ. *Externally* :—Infusum Cinchonæ Flavæ ; Tinctura Cinchonæ Flavæ.

**Incompatibles.**—Alkalies, alkaline carbonates, metallic salts, gelatin.

**Preparations.**—Infusum Cinchonæ Flavæ; Tinctura Cinchonæ Flavæ.

## CINCHONÆ PALLIDÆ CORTEX

PALE CINCHONA BARK

From the stem and branches of *Cinchona officinalis* and *C. Condaminea.*

Therapeutic properties, uses, and doses, similar to those of Cinchonæ Flavæ Cortex.

## CINCHONÆ RUBRÆ CORTEX

RED CINCHONA BARK

From *Cinchona Succirubra.*

Therapeutic properties, uses, and doses, similar to those of Cinchonæ Flavæ Cortex.

(This is the official Cinchona-bark of the B.P.)

## CINNAMOMI CORTEX

CINNAMON-BARK

The inner bark of the shoots of *Cinnamomum zeylanicum* (Ceylon cinnamon), or of one or more undetermined species of Chinese cinnamon.

**Natural Order.**—*Lauraceæ.*

**Composition.**—Contains a volatile oil, which is used for flavouring-purposes, tannic acid, mucilage, colouring matter, an acid, and lignin.

**Actions and Uses.**—A hæmostatic, and stomachic and carminative, useful in the treatment of diarrhœa. Used in conjunction with purgatives to prevent griping.

**Doses.**—Of the bark: Horse, ½ to 1 ounce.

Of the oil.
- Cattle, ½ to 1 drachm.
- Horse, 20 minims to 1 drachm.
- Dog, 1 to 4 minims.

## COCAINA

COCAINE

An alkaloid obtained from the leaves of *Erythroxylon Coca*, a small tree of Peru and Bolivia, belonging to the natural order *Erythroxylaceæ*.

**Composition.**—$C_{17}H_{21}NO_4$.

**Characters and Tests.**—Crystallises in colourless prisms, requiring for solution 704 parts of water. The hydrochloride ($C_{17}H_{21}NO_4HCl$) is readily soluble in water and alcohol, and is therefore usually employed.

**Actions and Uses.**—*Internally*:—Stimulant and tonic in treatment of sore throat. *Externally*:—As a local anæsthetic. Principally employed in veterinary practice in diseases of the eye. Twenty minims of a 4-per-cent. solution of the hydrochloride are dropped into the eye, for lessening irritability and pain in cases of conjunctivitis, ophthalmia, and ulceration of the cornea, for stitching torn eyelids, or for treatment of other injuries to the eye. In conjunction with atropine sulphate, it is employed in ophthalmoscopic examinations.

Professor Hobday has found that it is unwise to inject more than two grains of hydrochloride of cocaine subcutaneously to produce local anæsthesia in small dogs, such as fox terriers, or four grains in the case of large dogs, collies, &c. That is, one drachm of a 4-per-cent. solution for small dogs, and two drachms of a 4-per-cent. solution for large dogs. Local anæsthesia will be manifested in from

three to five minutes, and extends for about half an inch to one inch around the seat of injection, and lasts for about twenty minutes.

## COLCHICI CORMUS

COLCHICUM-CORM

The fresh corm of Meadow Saffron, *Colchicum autumna* ; collected about the end of June, stripped of its coats, sliced transversely, and dried at a temperature not exceeding 150° F.

**Natural Order.**—*Colchicaceæ.*

**Composition.**—Contains a poisonous crystallisable alkaloid called colchicine ($C_{17}H_{19}NO_5$).

**Characters.**—Fresh corm about the size of a chestnut, flattened where it has an undeveloped bud ; furnished with an outer brown and an inner yellow coat ; internally white, solid, and fleshy, yielding, when cut, a milky, acrid, and bitter juice. Dried slices about a tenth of an inch thick, moderately indented on one side, rarely on both ; firm, flat, whitish, amylaceous.

**Actions and Uses.**—In excessive doses, irritant poison ; in medicinal doses, cathartic, emetic, and sedative ; it also increases the secreting action of the skin, kidneys, and particularly that of the intestinal mucous membrane. Occasionally prescribed on account of its sedative and diuretic qualities, in small and frequently repeated doses, in rheumatism and rheumatic influenza, also in deep-seated or constitutional ophthalmia.

**Doses.**—Horse, ½ to 1 drachm.—Cattle, 1 to 2 drachms.—Sheep, 5 to 30 grains.—Dog, 2 to 8 grains.

**Modes of Administration.**—In powder, ball, or Tinctura Colchici.

**Incompatibles.**—Tincture of Iodine ; Guaiacum ; and all astringents.

**Antidotes.**—Emetics, where practicable, followed by demulcent drinks ; and, if coma be present, brandy, ammonia, and other powerful stimulants.

## COLCHICI SEMINA

COLCHICUM-SEEDS

The fully ripe seeds of *Colchicum autumnale.*

**Natural Order.**—*Colchicaceæ.*

**Characters.**—About the size of white mustard seed ; very hard, and of a reddish-brown colour.

**Actions and Uses.** Similar to those of the corm, but by some considered milder and more certain.

**Doses.**—Same as of corm.

**Preparation.**—Tinctura Colchici Seminum.

## COLLODIUM

COLLODION

**Modes of Preparation.**—Take of

| | |
|---|---|
| Pyroxylin . . . | 1 ounce |
| Ether . . . . | 36 fluid ounces |
| Alcohol (90-per-cent.) . | 12 fluid ounces |

Mix the ether and the spirit, and add the pyroxylin. Set aside for a few days, and, should there be any sediment, decant the clear solution. Keep in a well-corked or stoppered bottle.

**Characters.**—Colourless, highly inflammable liquid, with ethereal odour; dries rapidly on exposure to the air, and leaves a thin transparent film, insoluble in water or rectified spirit.

**Uses.**—As an adhesive to excoriations, ulcers, burns, wounds, &c.; likewise in erysipelas.

**Mode of Application.**—Brushed over the part. The applications are repeated, at intervals of a few minutes, until the film has become sufficiently thick to protect the affected part from atmospheric and other causes of irritation.

## COLLODIUM FLEXILE

FLEXIBLE COLLODION

Take of

| | |
|---|---|
| Collodion . . . | 6 fluid ounces |
| Canada Balsam . . | 120 grains |
| Castor Oil . . . | 1 fluid drachm |

Mix, and keep in a well-corked bottle.

Uses and mode of application similar to those of Collodion, to which, however, this preparation is superior on account of its elasticity.

## COLLODIUM HÆMOSTATICUM

HÆMOSTATIC COLLODION

Take of

| | |
|---|---|
| Collodion . . . | 10 fluid ounces |
| Carbolic Acid . . . | 1 ounce |
| Tannic Acid . . . | ½ ounce |
| Benzoic Acid . . . | ½ ounce |

Dissolve.

**Action and Use.**—*Externally*:—Styptic.

**Mode of Application.**—Same as Collodion.

## COLOCYNTHIDIS PULPA

COLOCYNTH-PULP

**Synonym.**—Bitter Cucumber.

The dried pulp of the fruit of *Citrullus Colocynthis* deprived of its rind, a native of Western Asia.

**Natural Order.**—*Cucurbitaceæ.*

**Composition.**—It contains a bitter alkaloidal principle (colocynthin), a resin (colocynthein), and a tasteless crystalline body (colocynthidin).

**Actions and Uses.**—A hydragogue purgative in chronic constipation. Its griping tendency may be overcome by combining it with aromatics or a small proportion of hyoscyamus or belladonna.

**Dose** (of the extract).—For Dogs, 1 to 2 grains in combination with other purgatives.

## CONFECTIO ROSÆ GALLICÆ

CONFECTION OF ROSES

**Mode of Preparation.**—Take of

| | |
|---|---|
| Fresh Red Rose Petals . . | 1 pound |
| Refined Sugar . . . | 3 pounds |

Beat the petals to a pulp in a stone mortar, add the sugar, and rub them well together.

**Use.**—In making Pilula Hydrargyri cum Ferro.

## CORIANDRI FRUCTUS

CORIANDER-FRUIT

**Synonym.**—Coriander-seed.

The dried fruit of *Coriandrum sativum*, cultivated in Britain.

**Natural Order.**—*Umbelliferæ.*

**Composition.**—The active principle is a volatile oil ($C_{10}H_{16}H_2O$).

**Characters.** — Globular, nearly as large as white pepper, beaked, finely ribbed, yellowish brown; has an agreeable aromatic odour and flavour.

**Actions and Uses.**
**Doses.**
**Modes of Application.**
} Same as Anisi Fructus.

## CREOSOTUM

CREOSOTE

A product of the fractional distillation of wood tar (B.P.)

**Composition.**—A mixture of guaiacol, creosal, and other phenols.

**Characters and Tests.**—Liquid, colourless or with a yellowish tinge, and a strong empyreumatic odour. Sparingly dissolved by water, but freely by alcohol, ether, and glacial acetic acid. Sp. gr. 1·071. Coagulates albumen. A slip of deal dipped into it, and afterwards into hydrochloric acid, acquires on exposure for a short time to the air a greenish-blue colour. Dropped on white filtering-paper and exposed to a heat of 212° F., it leaves no translucent stain. It turns the plane of polarisation of a ray of polarised light to the right (?). Is not solidified by the cold produced by a mixture of hydrochloric acid and sodium sulphate.

**Actions and Use.**
**Doses.**
**Modes of Application.**
**Antidotes.**
} Similar to those of Acidum Carbolicum.

**Preparations.**—Linimentum Creosoti Compositum; Mistura Creosoti; Unguentum Creosoti.

## CRETA PRÆPARATA

### PREPARED CHALK

Chalk freed from most of its impurities by elutriation, and afterwards dried in small, usually conical, masses.

**Characters and Tests.**—White amorphous powder; dissolves with effervescence (escape of carbonic anhydride) in dilute hydrochloric acid, indicating that it is a carbonate, and leaving only a small residue. The solution thus formed gives with ammonium oxalate a white precipitate (calcium chloride), showing the presence of calcium. The salt (calcium chloride) formed by dissolving prepared chalk in hydrochloric acid, if rendered neutral by evaporation to dryness, and re-dissolved in water, gives only a very scanty precipitate on the addition of saccharated solution of lime, showing the absence of calcium phosphate.

**Actions and Uses.**—*Internally*:—Antacid in indigestion, chronic diarrhœa, and dysentery; antidote to oxalic and most other acids. *Externally*:—As a desiccant and for absorbing irritating discharges; for protecting wounds, burns, and scalds from the air; also in erysipelas.

**Doses.** — Horse, 1 to 2 ounces. — Cattle, 1 to 2 ounces.—Sheep, 2 to 4 drachms.—Pig, 1 to 2 drachms. Dog, 5 to 15 grains.

**Modes of Application.**—*Internally*: In the form of bolus, or suspended in gruel, mucilage, or milk. Given in diarrhœa and dysentery; seldom administered alone, but usually in combination with opium, ginger, and catechu; the Mistura Cretæ Composita. *Externally*:—

As a desiccant it is used in the form of powder; to scalds and burns it may be applied mixed with water into a stiff paste.

**Incompatibles.**—All acids.

**Preparations.**—Hydrargyrum cum Cretâ; Mistura Cretæ Composita.

## CROCUS

SAFFRON

The dried stigma and part of style of *Crocus sativus*. Imported from Spain, France, and Italy.

**Natural Order.**—*Iridaceæ.*

**Composition.**—Contains an active principle called saffranin ($C_{48}O_{60}H_{18}$), a volatile oil, and also a colouring matter, crocin ($C_{16}H_{18}O_{16}$).

**Characters and Tests.**—Hay saffron consists of the dried parts of the styles, with their attached stigmas entangled together. The odour of saffron is penetrating, aromatic, and, in large quantities, narcotic. The taste is bitter and somewhat aromatic. When chewed, saffron tinges the mouth and saliva yellow; and when rubbed on the moistened finger it produces an intense orange stain.

**Actions and Uses.**—Seldom employed internally; principally used as a colouring and flavouring ingredient.

## CROTONIS SEMINA

CROTON-SEEDS

**Synonyms.**—Crotonis Tiglii Semina; Croton-beans.

**Natural Order.**—*Euphorbiaceæ.*

**Composition.**—Contains, in addition to oil, crotonic and crotonolic acids, an acrid purgative principle, crotonal ($C_9H_{14}O_2$).

**Characters.**—About the size of a coffee-berry, oblong, rounded at the extremities, with two faces, the external more convex than the internal. If the shelly covering of the seeds is partially removed, they present a mottled appearance; but if entirely so, they are brownish black.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant poison; in medicinal doses, cathartic in obstinate constipation, and when it is necessary to effect a speedy evacuation of the contents of the intestines.

**Doses.**—Horse, 30 to 36 grains.[1]—Cattle, 30 to 80 grains.[1]—Sheep, 5 to 15 grains.[1]—Dog, 3 to 10 grains.[1]

**Modes of Application.**—The seed, crushed to powder, made into a bolus, mixed with the animal's food or diluted with linseed oil.

**Preparation.**—Oleum Crotonis.

## CUMINI FRUCTUS

### CUMIN-FRUIT

**Synonym.**—Cumin-seeds.

The dried fruit of *Cuminum Cyminum.*

**Natural Order.**—*Umbelliferæ.*

**Composition.**—The properties of cumin depend upon a volatile oil, which is of a pale yellow colour and limpid. It is very acrid to the taste, and of a disagreeable odour.

**Characters.**—The fruit, commonly termed *cumin-seed*, is larger than anise, and of a light brown or greyish-

[1] One seed weighs about three grains.

yellow colour. It has some resemblance to, though it is larger than, caraway. Each mericarp has five primary ridges, which are filiform, and furnished with very fine prickles. The four secondary ridges are prominent and prickly, and under each there is one vitta. The odour of the fruit is strong and aromatic. Both in odour and taste it resembles caraway, though less agreeable.

**Actions and Uses.**
**Doses.**
**Modes of Application.**
} Same as Anisi Fructus.

## CUPRI IODIDUM CUM IODO

### IODIDE OF COPPER WITH IODINE

**Synonym.**—Improperly, Iodide of Copper.

**Composition.**—A mixture of cuprous iodide ($Cu_2I_2$) and free iodine.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Iodide of Potassium . . . . | 2 ounces |
| Sulphate of Copper . . . . | 4 ounces |
| Boiling Distilled Water . . . | $1\frac{1}{2}$ pint |

Dissolve the sulphate of copper in one pint of the distilled water, and the iodide of potassium in the remaining half-pint. Mix the solutions ; collect the precipitate which forms (mixture of cuprous iodide and iodine) on a filter, wash it with distilled water, and, when dried over sulphuric acid, pulverise it, and preserve it in a stoppered bottle.

| Cupric Sulphate. | | Potassium Iodide. | | Iodide of Copper with Iodine. | | Potassium Sulphate. |
|---|---|---|---|---|---|---|
| $2CuSO_4$ | + | $4KI$ | = | $Cu_2I_2+I_2$ | + | $2K_2SO_4$ |

**Characters and Tests.**—Light fawn-coloured powder which changes colour by exposure to air ; metallic taste.

The presence of free iodine is recognised by the preparation emitting the odour peculiar to this element, and by distilled water, after having been shaken with it and filtered, giving a blue colour (starch iodide) with cold aqueous solution of starch. The presence of combined copper and iodine may be proved by dissolving the solid in nitric acid, and testing the resulting solution for these elements.

**Actions and Uses.**—*Internally* :—Tonic and alterative ; also stimulant to the absorbents. Has been employed in the treatment of diabetes insipidus, glanders, farcy, and nasal gleet. *Externally* :—Stimulant and astringent. Applied to chronic œdematous enlargements of the legs, ill-conditioned ulcers, and in inveterate grease.

It is doubtful, however, whether the internal or external applications of this agent are attended with any greater benefits than those derived from the use, in similar cases, of sulphate of copper.

**Dose.**—Horse, 1 to 3 drachms daily, combined with gentian and pimento, or some other carminative.

**Modes of Application.**—*Internally* :—In the form of bolus. *Externally* :—The powder, either alone or made into an ointment with four parts of lard.

**Incompatibles.**—Acids ; acidulous salts ; alkalies and their carbonates ; lime-water ; vegetable astringents.

## CUPRI SUBACETAS

### SUBACETATE OF COPPER

**Synonyms.**—Copper Subacetate ; Ærugo ; Verdigris

**Composition.**—The blue variety chiefly consists of the compound $Cu(C_2H_3O_2)_2.CuO.6H_2O$ ; the green variety

contains, in addition to this compound, variable quantities of several basic copper acetates.

**Mode of Preparation.**—Plates of copper are subjected for several weeks to the united action of atmospheric oxygen and acetic acid. Sometimes the acid is derived from refuse grape-husks undergoing the acetous fermentation; at other times, cloths soaked in vinegar are arranged alternately with the plates of copper. When a sufficiency of the salt has formed, it is scraped off from any copper that may have remained unacted upon.

**Characters and Tests.**—Coarse masses or powder, either of a rich blue colour (blue verdigris) or of a pale bluish-green colour (green verdigris). Astringent and metallic taste. Partially soluble in water, completely dissolved by hydrochloric acid. Evolves an acetous odour which is increased on the application of heat or of strong sulphuric acid, showing the compound to be an acetate. Boiled with water and a few drops of acetic acid and filtered, a solution is obtained which furnishes a maroon-red precipitate (cupric ferrocyanide) with solution of potassium ferrocyanide, indicating the presence of copper.

**Actions and Uses.**—See preparations.

**Doses.**
**Modes of Application.**
**Incompatibles.**
**Antidotes.**
} Same as Cupri Sulphas.

**Preparations.**—Linimentum Cupri Subacetatis; Unguentum Cupri Subacetatis.

## CUPRI SULPHAS

SULPHATE OF COPPER

**Synonyms.**—Cupric Sulphate; Blue Vitriol; Blue Stone.

**Composition.**—$CuSO_4.5H_2O$.

**Mode of Preparation.**—Scraps of metallic copper are heated in a current of air so as to convert them superficially into cupric oxide, CuO, which forms in flakes and is known in commerce as 'copper scales.' The oxide thus produced, digested with dilute sulphuric acid, is converted into sulphate.[1] By filtration and evaporation the salt is obtained in the crystalline state.

| | Copper. | | Oxygen. | | Cupric Oxide. |
|---|---|---|---|---|---|
| 1. | $2Cu$ | + | $O_2$ | = | $2CuO$ |

| | Cupric Oxide | | Sulphuric Acid. | | Cupric Sulphate. | | Water. |
|---|---|---|---|---|---|---|---|
| 2. | $CuO$ | + | $H_2SO_4$ | = | $CuSO_4$ | + | $H_2O$ |

**Characters and Tests.**—Blue crystalline salt, in oblique prisms, soluble in water, forming a pale blue solution which strongly reddens blue litmus. The aqueous solution gives with barium chloride a white precipitate (barium sulphate) insoluble in hydrochloric acid, showing the salt to be a sulphate; and with potassium ferrocyanide a maroon-red precipitate, indicating the presence of copper (cupric ferrocyanide). If an aqueous solution of the salt be mixed with twice its volume of chlorine-water, to peroxidise any iron that may be present, and solution of ammonia be added, the precipitate (cupric hydrate) formed by the first addition of the ammonia will be dissolved by a further and sufficient addition of the alkali, and a violet-blue solution (copper ammonio-sulphate) will be produced, leaving nothing undissolved unless iron be present, in which case a reddish-brown precipitate will be left.

[1] Cupric sulphate is also prepared by boiling concentrated sulphuric acid with metallic copper, when the following decomposition takes place:

| Copper. | | Sulphuric Acid. | | Cupric Sulphate. | | Sulphurous Anhydride. | | Water. |
|---|---|---|---|---|---|---|---|---|
| $2Cu$ | + | $4H_2SO_4$ | = | $2CuSO_4$ | + | $2SO_2$ | + | $4H_2O$ |

**Actions and Uses.**—*Internally*:—In large doses, irritant poison; in medicinal doses, astringent and tonic. Given in debility, scrofula, dysentery, obstinate diarrhœa, glanders, and farcy; also, combined or alternated with liquor ammoniæ or ammonii carbonas, in purpura, typhoid fever, and other debilitating disorders. Likewise administered as an emetic to the dog. *Externally*:—Stimulant, astringent, styptic, mild escharotic, detergent. Applied to phagedænic ulcers—*e.g.*, in canker, thrush, and foot-rot, —also to farcy buds; sometimes a weak solution is injected up the vessels or into the frontal or maxillary sinuses in ulcerated conditions of the Schneiderian membranes; to the skin in scurvy; to arrest hæmorrhages; as a collyrium in chronic ophthalmia.

**Doses.** — *Tonic and Astringent*: — HORSE, 1 to 2 drachms.—CATTLE, 1 to 4 drachms.—SHEEP, 20 to 30 grains.—PIG, 5 to 10 grains.—DOG, 1 to 2 grains.

**Modes of Application.**—*Internally*:—As a tonic and astringent, in the form of Massa Cupri Sulphatis, or suspended in gruel or linseed tea. As an emetic, dissolved in warm water. *Externally*:—In the state of powder; Liquor Cupri Sulphatis; Liquor Cupri Sulphatis Compositus; Linimentum Cupri Sulphatis.

**Incompatibles.**—Alkalies; alkaline carbonates; vegetable astringents.

**Antidotes.**—Albumen; milk; bean or pea meal suspended in water; mucilaginous drinks.

**Preparations.**—Linimentum Cupri Sulphatis; Liquor Cupri Sulphatis; Liquor Cupri Sulphatis Compositus; Massa Cupri Sulphatis.

## CUSSO

KOUSSO

The flowering panicles of pistillate flowers of *Brayera anthelmintica*, gathered before the seeds are quite ripe, dried in the sun, and reduced to coarse powder. Imported from Abyssinia.

**Natural Order.**—*Rosaceæ.*

**Composition.**—Contains two varieties of tannin, a bitter acrid and a tasteless resin, a fixed and a volatile oil, a crystalline body termed koussin, chlorophyll, sugar, gum, &c.

**Characters and Tests.**—The branches of flowers are of a greenish-yellow colour, the edges of the petals having a purplish hue; balsamic odour, when freshly opened, compared by Pereira to the combined odour of tea, hops, and senna-leaves; slightly acrid and unpleasant taste. On the addition of a ferric salt to an infusion or decoction of kousso, a dark green colour develops.

**Action and Use.** — Anthelmintic, especially for tænia.

**Doses.**—Horse, ½ to 1 pound.—Sheep, 4 drachms to 1 ounce.—Lamb, 30 to 50 grains.—Dog, 10 grains to 6 drachms.

**Modes of Application.**—In the form of Infusum Cusso.

**Preparation.**—Infusum Cusso.

## DECOCTUM TABACI

DECOCTION OF TOBACCO

Take of

| | |
|---|---|
| Tobacco . . . . . | 1 ounce |
| Water . . . . . | 1 pint |

Boil the water with the tobacco for a few seconds (for ten minutes, if the time will permit), strain, and add cold water to the strained liquid until it measures two pints.

**Use.**—Antidote to nux vomica; also to strychnine and its preparations.

**Doses.**—HORSE, 1 to 3 pints.—CATTLE, 1 to 3 pints. SHEEP, 5 to 15 fluid ounces.—PIG, 5 to 15 fluid ounces.—DOG, 2 to 5 fluid ounces.

Should the spasms continue after the administration of the tobacco, repeat the dose, at intervals to be determined by the effect produced, until complete muscular relaxation ensues. The effects of an overdose of tobacco may be counteracted by the exhibition of diffusible stimulants.

## DIGITALINUM

DIGITALIN

**Composition.**—$C_{54}H_{84}O_{27}$.

The difficultly crystallisable sedative principle of *Digitalis purpurea.* Too expensive for veterinary use.

## DIGITALIS FOLIA

DIGITALIS-LEAF

The dried leaf of *Digitalis purpurea*, Purple Foxglove.

**Natural Order.**—*Scrophulariaceæ.*

**Composition.**—Contains a bitter glucoside, termed digitalin; also digitoxin, digitolein, digitonin, and digitin.

**Mode of Preparation.**—The leaves are gathered from the wild indigenous plant, in June and July, when about two-thirds of the flowers are expanded, and dried by the heat of a stove in a dark place.

**Characters.**—Ovate-lanceolate, shortly petiolate, rugose, downy, crenate; bright green colour, which is paler on the under than on the upper surface; scarce any odour; acrid and nauseous taste.

**Actions and Uses.**—*Internally*:—In large doses, irritant and sedative poison; in medicinal doses, sedative and diuretic. Liable to accumulate in the system and irritate the intestines; its action should therefore be carefully watched. Chiefly given as a sedative in the secondary stages of inflammatory affections, functional diseases of the heart,[1] and chronic rheumatism; also recommended in pneumonia and chronic cough. Prescribed as a diuretic, either alone or in combination with nitrate of potash, in dropsical complaints. *Externally*:—Irritant, diuretic, sedative, anodyne. Occasionally employed as a collyrium in ophthalmia.

**Doses.**—Horse, 15 grains to 1 drachm.—Cattle, ½ to 1 drachm.—Sheep, 5 to 15 grains.—Pig, 5 to 10 grains.—Dog, 1 to 3 grains.

**Modes of Application.** — *Internally*:—The powdered leaf made into a ball; Extractum Digitalis. For coughs, Massa Digitalis Composita. *Externally*:—Infusum Digitalis.

[1] Medicinal doses are vascular and cardiac stimulants and tonics, and are prescribed to increase the force and co-ordinating power of the heart and relieve congestion of veins and capillaries.

**Incompatibles.**—Sulphate and chloride of iron, the acetates of lead, and the preparations of cinchona-bark.

**Antidotes.**—Empty the alimentary canal, and apply stimulants internally and externally.

**Preparations.**—Extractum Digitalis; Massa Digitalis Composita; Tinctura Digitalis.

## EMPLASTRUM PICIS

PITCH PLASTER

Take of

| | |
|---|---|
| Burgundy Pitch | 20 ounces |
| Resin, Yellow Wax | of each $2\frac{1}{4}$ ounces |
| Olive Oil | $1\frac{1}{2}$ fluid ounce |
| Water | 1 fluid ounce |

Add the oil and the water to the Burgundy pitch, resin, and wax, previously melted together; then, constantly stirring, evaporate to a proper consistency.

**Use.**—Adhesive to wounds when spread on leather.

## ENEMA ALOES

ENEMA OF ALOES

Take of

| | |
|---|---|
| Aloes | 4 drachms |
| Carbonate of Potassium | 30 grains |
| Warm Mucilage of Starch or Gruel | 1 pint |

Mix and rub together.

## ENEMA ASAFETIDÆ

ENEMA OF ASAFETIDA

Take of

| | |
|---|---|
| Asafetida | 2½ drachms |
| Warm Water | 1 pint |

## ENEMA MAGNESIÆ SULPHATIS

ENEMA OF SULPHATE OF MAGNESIA

Take of

| | |
|---|---|
| Sulphate of Magnesia | 2 ounces |
| Linseed or Castor Oil | 2 fluid ounces |
| Mucilage of Starch or Gruel | 1 pint |

Dissolve the sulphate of magnesia in the mucilage of starch or gruel, add the oil, and mix.

## ENEMA OPII

ENEMA OF OPIUM

Take of

| | |
|---|---|
| Tincture of Opium | 5 fluid drachms |
| Warm Mucilage of Starch or Gruel | 1 pint |

Mix.

## ENEMA TABACI

ENEMA OF TOBACCO

Take of

| | |
|---|---|
| Leaf Tobacco | 1 drachm |
| Boiling Water | 20 fluid ounces (1 pint) |

Infuse in a covered vessel for half an hour, and strain.

## ENEMA TEREBINTHINÆ

ENEMA OF TURPENTINE

Take of

| | |
|---|---|
| Oil of Turpentine . . . | 2 fluid ounces |
| Mucilage of Starch or Gruel . | 1 pint |

## ERGOTA

ERGOT

**Synonym.**—Ergot of Rye.

The sclerotium (compact mycelium or spawn) of a fungus, *Claviceps purpurea*, produced within the paleæ of the common rye, *Secale cereale.*

**Natural Order.**—*Basidiomycetes.*

**Composition.**—About 30 per cent. of a fixed oil; a sugar (mycose); lactic, acetic, and formic acids; colouring matter; and three active principles, cornutin (an alkaloid) and sphacelinic and ergotinic acids.

**Characters.**—Subtriangular, curved with a longitudinal furrow on the concave side, obtuse at the ends; from one-third of an inch to one and a half inch in length; violet-brown colour superficially, pinkish internally; solid, frangible, fracture short; faint odour, which may be increased by trituration with solution of potash.

**Actions and Uses.**—*Internally*:—In large doses, irritant poison; in medicinal doses, styptic in hæmorrhages of the lungs, kidneys, and other organs, but principally used as a parturient—*i.e.*, to excite the uterus to contract and expel its contents during parturition. Its efficacy in this respect is, however, doubtful on the lower animals. *Externally*:—Employed as a styptic, to arrest hæmorrhages—*e.g.*, of the vagina, uterus, &c.

**Doses as an Ecbolic.** — MARE, ½ ounce to 2 ounces.—Cow, 2 to 4 drachms.—SHEEP, 2 drachms.—Sow, ½ drachm to 1 drachm.—BITCH, 10 grains to 1 drachm, according to size.

**Modes of Application.**—*Internally* :—The powder, in ale or porter, with a carminative ; Extractum Ergotæ Liquidum ; Infusum Ergotæ ; Tinctura Ergotæ. *Externally* :—Extractum Ergotæ Liquidum ; Infusum Ergotæ ; Tinctura Ergotæ.

**Preparations.**—Extractum Ergotæ Liquidum ; Infusum Ergotæ ; Tinctura Ergotæ.

## EUCALYPTUS

EUCALYPTUS

**Synonym.**—Australian Blue Gum.

The leaves of the *Eucalyptus globulus*; from the leaves an oil is obtained by distillation (see Oleum Eucalypti).

**Natural Order.**—*Myrtaceæ.*

**Actions and Uses.**—See Oleum Eucalypti.

## EUPHORBIUM

EUPHORBIUM

**Synonym.**—Gum Euphorbium.

An acrid resin, obtained from *Euphorbia officinarum*, and probably other species of *Euphorbia*.

**Natural Order.**—*Euphorbiaceæ.*

**Characters.**—Irregular, slightly friable tears, usually pierced with one or two holes. These tears are almost odourless, but their dust is very irritating when applied

to the mucous lining of the nose. Taste at first slight, afterwards acrid and burning. On heating, euphorbium melts, swells up imperfectly, evolves an odour like benzoic acid, takes fire, and burns with a pale flame. Alcohol, ether, and oil of turpentine are its best solvents; water only dissolves a small portion of it.

**Composition.**—Contains about 60 per cent. of a reddish-brown, hard, brittle resin, which is its active constituent.

**Actions and Uses.**—Sometimes it is employed as an errhine in chronic affections of the eyes, ears or brain; but its local action is so violent that it can only be applied when largely diluted with starch or flour. Mixed with turpentine, Burgundy pitch, or resin, it is employed in the form of plaster, as a rubefacient, in chronic affections of the joints. Sometimes used as a vesicant.

## EXTRACTUM ACONITI

### EXTRACT OF ACONITE

**Mode of Preparation.**—Take of

| | |
|---|---|
| The fresh leaves and flowering tops of Aconite . . . | 112 pounds |

Bruise in a stone mortar, and press out the juice; heat it gradually to 130° F., and separate the green colouring matter by a calico filter. Heat the strained liquor to 200° F. to coagulate albumen, and filter. Evaporate the filtrate on a water-bath to the consistence of a thin syrup; then add to it the green colouring matter previously separated, and, stirring the whole together assiduously, continue the evaporation at a temperature not

exceeding 140° F., until the extract is of suitable consistence for forming balls or pills.

**Doses.**—Horse, 10 to 15 grains.—Cattle, 15 to 30 grains.—Sheep, 2 to 5 grains.—Pig, 2 to 5 grains.—Dog, $\frac{1}{4}$ to 2 grains.

**Modes of Application.**—In the form of bolus, or rubbed up with cold water.

## EXTRACTUM BELLADONNÆ

EXTRACT OF BELLADONNA

**Mode of Preparation.**—Take of

| | |
|---|---|
| The fresh leaves and young branches of Belladonna . | . 112 pounds |

Bruise in a stone mortar, and press out the juice; heat it gradually to 130° F., and separate the green colouring matter by a calico filter. Heat the strained liquor to 200° F. to coagulate albumen, and filter. Evaporate the filtrate on a water-bath to the consistence of a thin syrup; then add to it the green colouring matter previously separated, and, stirring the whole together assiduously, continue the evaporation, at a temperature not exceeding 140° F., until the extract is of a suitable consistence for forming balls or pills.

**Doses.**—Horse, 1 to 2 drachms.—Cattle, 2 to 4 drachms.—Sheep, 3 to 10 grains.—Pig, 2 to 5 grains.—Dog, 1 to 5 grains.

**Modes of Application.**—In the form of bolus, or rubbed up with cold water.

## EXTRACTUM CANNABIS INDICÆ

EXTRACT OF INDIAN HEMP

**Preparation.**—Take one pound of the dried flowering or fruiting tops of the female plants of *Cannabis sativa*, from which the resin has not been removed; powder coarsely, macerate for seven days, press out the tincture, distil off the spirit, and evaporate to the consistency of a soft extract.

The *Cannabis sativa* grows in India, where it is known as Gunjah or Ganga; it is rough to the feel, brittle, of a greenish colour, and with a peculiar characteristic narcotic odour.

**Characteristics.**—A soft extract, of a greenish colour, and with the peculiar narcotic odour of the plant.

**Natural Order.**—*Cannabinaceæ.*

**Actions and Uses.**—Antispasmodic, hypnotic, and sedative. Used internally in colic and inflammation of the bowels; in chronic cough; in tetanus; and in chorea of dogs it has been observed to lessen the involuntary movements. It has not the effect of producing constipation or loss of appetite when repeatedly given, and is very convenient to administer as bolus or pill.

**Doses.**—Horse, 1 to 4 drachms.—Dog, 1 to 3 grains.

**Preparation.**—Tinctura Cannabis Indicæ.

## EXTRACTUM DIGITALIS

EXTRACT OF DIGITALIS

**Mode of Preparation.**—Take of

Fresh leaves of Digitalis . 112 pounds

Bruise in a stone mortar, and press out the juice; heat it gradually to 130° F., and separate the green colouring matter by a calico filter. Heat the strained liquor to 200° F. to coagulate albumen, and filter. Evaporate the filtrate on a water-bath to the consistence of a thin syrup; then add to it the green colouring matter previously separated, and, stirring the whole together assiduously, continue the evaporation at a temperature not exceeding 140° F., until the extract is of a suitable consistence for forming balls or pills.

**Doses.** — Horse, 5 to 10 grains. — Cattle, 5 to 10 grains.—Sheep, 1 to 5 grains.—Pig, 1 to 5 grains.—Dog, $\frac{1}{4}$ to 1 grain.

**Modes of Application.**—In the form of bolus, or rubbed up with cold water.

## EXTRACTUM ERGOTÆ LIQUIDUM

### LIQUID EXTRACT OF ERGOT

**Mode of Preparation.**—Take of

| | |
|---|---|
| Ergot, in coarse powder . | 1 pound |
| Ether . . . . | 1 pint, or a sufficiency |
| Distilled Water . . | $3\frac{1}{2}$ pints |
| Rectified Spirit . . | 8 fluid ounces |

Shake the ether in a bottle with half a pint of the water, and after the two liquids have separated from one another on standing, decant the ether. Place the ergot in a percolator, and free it from its oil by passing the washed ether slowly through it. Remove the marc, and digest it in three pints of the water at 160° F. for twelve hours. Press out, strain, and evaporate the liquor by the heat of a water-bath to nine fluid ounces; when cold add the spirit. Allow it

to stand for half an hour to coagulate, then filter. The product should measure sixteen fluid ounces.

**Actions and Uses.**—See Ergota.

**Doses.**—Horse, 1 to 4 fluid drachms.—Cattle, 2 to 4 fluid drachms.—Sheep, 10 to 40 minims.—Pig, 10 to 20 minims.—Dog, 10 to 30 minims.

**Mode of Application.**—Diluted with water.

## EXTRACTUM FILICIS LIQUIDUM

LIQUID EXTRACT OF MALE FERN

**Mode of Preparation.**—Take of

| | |
|---|---|
| Fern Root, in coarse powder | 2 pounds |
| Ether . . . . | 4 pints or a sufficiency |

Mix the fern root with two pints of the ether, pack closely in a percolator, and add the remainder of the ether at intervals, until it passes through colourless. Evaporate off the ether over a water-bath, or recover it by distillation, and preserve the oily extract which remains.

**Doses.** — Horse, 2 to 6 drachms. — Dog, ½ to 1 drachm.

## EXTRACTUM HYOSCYAMI

EXTRACT OF HYOSCYAMUS

**Mode of Preparation.**—Take of

| | |
|---|---|
| The fresh leaves and young branches of Hyoscyamus | 112 pounds |

Bruise in a stone mortar, and press out the juice; heat it gradually to 130° F., and separate the green colouring matter by a calico filter. Heat the strained liquor to

200° F., and filter. Evaporate the filtrate on a water-bath to the consistence of a thin syrup; then add to it the green colouring matter previously separated, and, stirring the whole assiduously, continue the evaporation at a temperature not exceeding 140°, until the extract is of a suitable consistence for forming balls or pills.

**Doses.**—Horse, 1 to 2 drachms.—Cattle, 1 to 2 drachms.—Sheep, 15 to 30 grains.—Pig, 10 to 20 grains. Dog, 2 to 8 grains.

**Modes of Application.**—In the form of bolus, or rubbed up with cold water.

## FERRI CARBONAS SACCHARATA

### SACCHARATED CARBONATE OF IRON

**Composition.**—Ferrous oxycarbonate—

$$x\mathrm{FeCO_3},y\mathrm{Fe(OH)_2}$$

—more or less oxidised, mixed with sugar, the carbonate, reckoned as $FeCO_3$, forming about one third of the mixture.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Sulphate of Iron . . . . | 2 ounces |
| Carbonate of Ammonium . . . | 1¼ ounce |
| Boiling Distilled Water . . . | 2 gallons |
| Refined Sugar . . . . . | 1 ounce |

Dissolve the sulphate of iron and the carbonate of ammonium separately in half a gallon of the water, and mix the two solutions with brisk stirring in a deep cylindrical vessel, which is then to be covered as accurately as possible. Set the mixture by for twenty-four hours, and from the precipitate (carbonate of iron) which has sub-

sided separate the supernatant liquid by a siphon. Pour on the remainder of the water, stir well, and after subsidence again remove the clear liquid. Collect the resulting carbonate on a calico filter, and, having first subjected it to expression, rub it with the sugar in a porcelain mortar. Finally, dry the mixture at a temperature not exceeding 212° F.

| Ferrous Sulphate. | | Ammonium Carbonate. | | Ferrous Carbonate (Carbonate of Iron). | | Ammonium Sulphate. |
|---|---|---|---|---|---|---|
| $FeSO_4$ | + | $(NH_4)_2CO_3$ | = | $FeCO_3$ | + | $(NH_4)_2SO_4$ |

The sugar is added to prevent the absorption of atmospheric oxygen, the loss of carbonic anhydride, and the consequent conversion of the ferrous carbonate into ferric oxide.

**Characters and Tests.**—Small coherent lumps of a grey colour, with a sweet very feeble chalybeate taste. Dissolves with effervescence in warm hydrochloric acid diluted with half its volume of water, indicating that it contains a carbonate, and the solution gives but a very slight precipitate with barium chloride, showing its almost total freedom from sulphate; with potassium ferrocyanide the solution gives a pale blue precipitate, which, on exposure to air, changes to a dark blue, indicating the presence of a ferrous salt.

**Actions and Uses.**—Same as Ferri Sulphas.

**Doses.**—Three times larger than Ferri Sulphas.

**Modes of Application.**—In bolus, pill, or suspended in some mucilaginous drink.

**Incompatibles.**—Acids; acidulous salts; vegetable astringents.

## FERRI IODIDUM

IODIDE OF IRON

**Composition.**—Ferrous iodide, $FeI_2$, with about 18 per cent. of water of crystallisation and a little ferric oxide.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Fine Iron Wire . . . | 1½ ounce |
| Iodine . . . . . | 3 ounces |
| Distilled Water . . . | 15 fluid ounces |

Put the iodine, iron, and twelve ounces of the water into a flask, and, having heated the mixture gently for about ten minutes, raise the heat and boil till the froth becomes white. Pass the solution as quickly as possible through a wetted calico filter into a dish of polished iron, washing the filter with the remainder of the water, and boil down until a drop of the solution taken out on the end of an iron wire solidifies on cooling. The liquid should now be poured out on a porcelain dish, and, as soon as it has solidified, should be broken into fragments, and enclosed in a well-stoppered bottle.

In this process iron and iodine combine chemically, thus :

| Iron. | | Iodine. | | Ferrous Iodide (Iodide of Iron). |
|---|---|---|---|---|
| $2Fe$ | + | $2I_2$ | = | $2FeI_2$ |

The iodide acquires five molecules of water of crystallisation, and becomes $FeI_2.5H_2O$.

**Characters and Tests.**—Crystalline ; green, with a tinge of brown ; inodorous ; deliquescent ; almost entirely soluble in water, forming a slightly green solution, which gradually deposits a rust-coloured sediment (ferric oxide)

and acquires a red colour. Its aqueous solution gives a copious blue precipitate with potassium ferricyanide, showing the presence of iron. Mixed with mucilage or starch, it acquires a blue colour on the addition of a minute quantity of solution of chlorine or nitric acid, indicating that the compound is an iodide.

**Actions and Uses.**—Tonic, hæmatinic, alterative, astringent. Given in diabetes insipidus and nasal gleet.

**Doses.**—HORSE, ½ to 2 drachms.—CATTLE, 1 to 2 drachms.—SHEEP, 15 to 30 grains.—PIG, 10 to 20 grains. DOG, 1 to 8 grains.

**Modes of Application.**—In bolus, pill, or solution in water, or as the official Syrup.[1]

**Incompatibles.**—Acids; acidulous salts; alkalies; alkaline carbonates; lime-water; vegetable astringents.

## FERRI PEROXIDUM HUMIDUM

### MOIST PEROXIDE OF IRON

**Synonyms.**—Moist Ferric Hydrate; Ferri Peroxidum Hydratum; Moist Hydrated Peroxide of Iron.

**Composition.**—Ferric hydrate, $Fe_2(HO)_6$, with about 86 per cent. of uncombined water.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Solution of Persulphate of Iron . | 4 fluid ounces |
| Solution of Soda . . . | 33 fluid ounces |
| Distilled Water . . . . | a sufficiency |

Mix the solution of persulphate of iron with a pint of

[1] This syrup contains about 5½ grains of ferrous iodide in each drachm, and is frequently given in bronchitis and canine distemper.

the distilled water, and add this gradually to the solution of soda, stirring constantly and briskly. Let the mixture stand for two hours, stirring it occasionally, then put it on a calico filter, and, when the liquid has drained away, wash the precipitate (moist peroxide of iron) with distilled water until the filtrate ceases to give a precipitate with barium chloride, which shows that the precipitate is freed from sulphate. Lastly, enclose the precipitate, without drying it, in a stoppered bottle from which evaporation cannot take place. This preparation, when used, should be recently made.

| Ferric Sulphate. | | Sodium Hydrate. | | Ferric Hydrate (Moist Peroxide of Iron). | | Sodium Sulphate. |
|---|---|---|---|---|---|---|
| $Fe_2(SO_4)_3$ | + | $6NaHO$ | = | $Fe_2(HO)_6$ | + | $3Na_2SO_4$ |

**Characters and Tests.**—Soft pasty mass, of a reddish-brown colour. Dissolves readily in dilute hydrochloric acid without the aid of heat, and the solution gives a copious blue precipitate [ferric ferrocyanide, Prussian blue, $Fe_4(FeCy_6)_3$] with potassium ferrocyanide, but none with potassium ferricyanide. A little of it dried at 212° F. until it ceases to lose weight gives off water of hydration when heated to dull redness in a test-tube.

**Actions and Uses.**—Antidote to poisoning by arsenic, in which case it forms an insoluble and inert ferrous subarsenite.

**Doses.**—The administration should be frequently repeated until at least twelve times the supposed quantity of the arsenic has been given, or until it appears to have been effective in allaying the action of the poison.

**Mode of Application.**—Mixed with water or some demulcent drink.

# FERRI SULPHAS

## SULPHATE OF IRON

**Synonyms.**—Ferrous Sulphate; Protosulphate of Iron; Green Vitriol; Copperas.

**Composition.**—$FeSO_4.7H_2O$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Iron Wire . . . | 4 ounces |
| Sulphuric Acid . . | 4 fluid ounces |
| Distilled Water . . | 1½ pint |

Pour the water on the iron placed in a porcelain dish, add the sulphuric acid, and, when the disengagement of hydrogen has nearly ceased, boil for ten minutes. Filter through paper, and after the lapse of twenty-four hours separate the crystals (sulphate of iron) which have deposited from the solution. Dry these on filter-paper placed on porous bricks, and preserve them in a stoppered bottle.

| Iron. | | Sulphuric Acid. | | Ferrous Sulphate (Sulphate of Iron). | | Hydrogen. |
|---|---|---|---|---|---|---|
| $Fe_2$ | + | $2H_2SO_4$ | = | $2FeSO_4$ | + | $2H_2$ |

In the act of crystallising, seven molecules of water are taken up by the salt, forming the compound $FeSO_4.7H_2O$.[1]

**Characters and Tests.**—Oblique, rhombic, prismatic crystals; pale greenish-blue colour and astringent taste; soluble in water, insoluble in rectified spirit. Its aqueous solution gives a white precipitate (barium

[1] This salt is most extensively obtained by the oxidation of iron pyrites (iron disulphide), $FeS_2$.

| Iron Disulphide. | | Oxygen (from the air). | | Water. | | Ferrous Sulphate. | | Sulphuric Acid. |
|---|---|---|---|---|---|---|---|---|
| $FeS_2$ | + | $7O_2$ | + | $2H_2O$ | = | $2FeSO_4$ | + | $2H_2SO_4$ |

sulphate) with barium chloride, a dark blue one (ferrous ferricyanide,[1] *Turnbull's Blue*, $Fe_3Fe_2Cy_{12}$) with potassium ferricyanide, and a nearly white or light blue precipitate (ferrous ferrocyanide,[2] $Fe_2FeCy_6$) with potassium ferrocyanide. It should give no precipitate with hydrosulphuric acid, showing its freedom from lead and other metals precipitable by this reagent.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant poison ; in medicinal doses, astringent, tonic, and hæmatinic. Given in, and during convalescence from, debilitating diseases ; in dysentery, internal hæmorrhages, hæmaturia, diabetes, chorea, and occasionally in epilepsy. *Externally*:—Astringent to ulcers, to check capillary hæmorrhages, and to diminish profuse discharges.

**Doses.**—Horse, 1 to 2 drachms.—Cattle, 2 to 3 drachms.—Sheep, 15 to 30 grains.—Pig, 10 to 20 grains. Dog, 1 to 10 grains.

**Modes of Application.**—*Internally*:—Massa Ferri Sulphatis, or dissolved in water. *Externally*:—In powder or strong aqueous solution.

**Incompatibles.**—Alkalies ; alkaline carbonates ; lime-water ; vegetable astringents.

**Preparation.**—Massa Ferri Sulphatis.

## FERRUM

### IRON

Annealed iron wire having a diameter of about 0·005 inch, or wrought-iron nails, free from oxide.

[1] Ferricyanogen is a compound radical, having the formula $Fe_2(CN)_{12}$, and sometimes written $Fe_2Cy_{12}$.

[2] Ferrocyanogen is a compound radical, having the formula $Fe(CN)_6$, and sometimes written $FeCy_6$.

**Composition.**—An element; symbol Fe.

**Preparations.**—Ferri Carbonas Saccharata; Ferri Iodidum; Ferri Peroxidum Humidum; Ferri Sulphas; Liquor Ferri Perchloridi Fortis; Liquor Ferri Persulphatis.

## FILIX MAS

### MALE FERN

**Synonyms.**—Male Shield Fern; Fern Root.

The dried rhizome, with the bases of the foot-stalks and portions of the root-fibres, of *Aspidium Filix-mas*, collected in summer. Indigenous.

**Natural Order.**—*Filices.*

**Composition.**—According to Bock the root contains 0·04 per cent. of a volatile oil, to which are attributed its anthelmintic properties. It also contains 4 per cent. of resin, a green fatty oil, tannic acid, and filicic acid.

**Mode of Preparation.**—The rhizome should be dug up in summer, cleared of root-fibres, &c., but not washed, and dried quickly and thoroughly in the open air, in the shade, or in a hot-air press at a temperature not above 140° F.; the tufts and those parts of the root-stock which are greenish internally are alone to be kept; they should be reduced to powder immediately, and preserved in well-stoppered bottles; the stock should be renewed annually, as in two years the plant loses its medicinal qualities.

**Characters.**—Tufted, scaly, greenish brown; powder greenish yellow, with a disagreeable odour and a nauseous, bitter, somewhat astringent taste.

**Action and Use.**—Anthelmintic for tænia.

**Doses.**—Horse, 8 to 16 ounces.—Cattle, 8 to 16 ounces.—Sheep, 3 to 4 ounces.—Dog, 1 to 3 ounces.

**Mode of Application.**—The powder made into a bolus or mixed with gruel ; Extractum Filicis Liquidum.[1]

**Preparation.**—Extractum Filicis Liquidum.

## GALBANUM

GALBANUM

A gum-resin, derived from *Ferula galbaniflua* and probably other allied umbelliferous plants. Imported from India and the Levant.

**Composition.**—Its medicinal properties depend upon the presence of a resinous body and a volatile oil.

**Characters.**—Irregular tears, about the size of a pea, usually agglutinated into masses ; greenish-yellow colour, translucent, having a strong disagreeable odour, and an acrid bitter taste.

**Actions and Uses.** } Similar to Ammoniacum and
**Doses.** } Asafetida.

## GALLA

GALLS

**Synonyms.**—Oak-galls ; Nut-galls.

Excrescences formed on the young branches of the oak, *Quercus infectoria*, belonging to the natural order

[1] This preparation is to be preferred to the powder, as the latter is uncertain in its action no matter how well it may be kept.

*Cupuliferæ*, as a consequence of the irritation produced by the puncture and deposited ova of the female of a hymenopterous insect, *Cynips Gallæ tinctoriæ*.

**Composition.**—Their therapeutic properties are chiefly due to their containing about 40 to 75 per cent. of tannic acid ($C_{13}H_9O_7CO_2H$) and 3 per cent. of gallic acid [$C_6H_2(HO)_3CO_2H$].

**Characters.**—Hard, heavy, globular bodies, varying in size from half an inch to three-fourths of an inch in diameter; tuberculated on the surface, the tubercles and intervening spaces smooth; bluish-green colour superficially, yellowish white within, with a small central cavity; intensely astringent.

**Actions and Uses.**—*Internally and Externally*:—Astringent and styptic. Employed in same cases as Acidum Tannicum.

**Doses.**—Horse, 1 to 3 drachms.—Cattle, 2 to 4 drachms.—Sheep, 20 to 40 grains.—Pig, 10 to 20 grains. Dog, 2 to 10 grains.

**Modes of Application.**—*Internally*:—In the form of bolus or Tinctura Gallæ. *Externally*:—Tinctura Gallæ; Unguentum Gallæ; Unguentum Gallæ cum Opio.

**Incompatibles.**—Mineral acids; soluble salts of iron and lead; sulphate of copper; nitrate of silver; tartar emetic; lime-water; alkaline carbonates and bicarbonates, and many vegetable agents containing alkaloids.

**Preparations.**—Acidum Gallicum; Acidum Tannicum; Unguentum Gallæ; Unguentum Gallæ cum Opio.

## GENTIANÆ RADIX

GENTIAN-ROOT

The dried root of *Gentiana lutea*. Collected in the mountain districts of Central and Southern Europe.

**Natural Order.**—*Gentianaceæ.*

**Composition.**—Gentian-root consists for the most part of a volatile, odorous, and butyraceous oily matter, a bitter crystalline body consisting of an acid (gentisic acid), and a bitter crystalline principle (gentiopicrin), with gum, sugar, pectin, wax, caoutchouc, a yellow colouring matter, and woody fibre.

**Characters.**—From half an inch to one inch in thickness, several inches in length, often twisted, much wrinkled or marked with close transverse rings; brown externally, yellow internally; tough and spongy; taste at first sweetish, but afterwards very bitter.

**Actions and Uses.**—*Internally*:—Tonic and stomachic. *Externally*:—Occasionally employed as a mild excitant and antiseptic.

**Doses.**—Horse, 2 to 4 drachms.—Cattle, 2 to 4 drachms.—Sheep, 1 to 2 drachms.—Pig, ½ to 1 drachm. Dog, 10 to 20 grains.

**Modes of Application.**—In the form of bolus; Infusum Gentianæ Compositum; Tinctura Gentianæ Composita.

**Preparation.**—Infusum Gentianæ Compositum.

## GLYCERINUM

GLYCERIN

**Synonym.**—Glycerol.

**Composition.**—$C_3H_5(HO)_3$, a trihydric alcohol with a small amount of water.

**Mode of Preparation.**—Price's glycerin is preferred to that of other makers. Molten fat is resolved by the action of high-pressure steam, at a temperature of from 500° to 600° F., into glycerin and a mixture of stearic, palmitic, and oleic acids; the former dissolves in the water formed by the condensation of the steam, while the latter floats on the surface of the solution of glycerin. After the removal of the fatty acids the glycerin is deprived of nearly the whole of its water by evaporation.

**Characters.**—Clear colourless fluid, oily to the touch. Sp. gr. 1·25. Slight odour, sweet taste; freely soluble in water and in alcohol. Strongly heated, it decomposes and evolves intensely irritating vapours (acrolein, $C_3H_4O$).

**Actions and Uses.**—Chiefly employed externally as an emollient in skin-affections accompanied by dryness and irritation; to excoriations, cracked heels, burns, scalds, and freshly blistered surfaces. When used to allay irritability it may be advantageously mixed with an equal bulk of Liquor Plumbi Subacetatis. Also employed as a vehicle for many medicines, and, in consequence of its non-drying properties, it is often added to masses to prevent their hardening.

## GLYCERINUM ACIDI CARBOLICI

GLYCERIN OF CARBOLIC ACID

Take of

| | |
|---|---|
| Carbolic Acid (Phenol) . . . | 1 part |
| Glycerin . . . . . | 4 parts |

Rub together in a mortar until the Phenol is dissolved.

**Use.**—Astringent and stimulant to indolent wounds and ulcers, also to abrasions, sore backs, cracked heels, &c.

## GOSSYPIUM

COTTON WOOL

The carded hairs of the seeds of various species of *Gossypium.*

**Natural Order.**—*Malvaceæ.*

**Composition.**—Nearly pure cellulose or woody fibre, $C_6H_{10}O_5$.

**Uses.**—To blistered surfaces, as in burns, scalds, erysipelas, &c., and to wounds, for the purpose of protecting them from the irritating effects of the atmosphere.

**Preparations.**—Collodium ; Collodium Flexile.

## HYDRARGYRI IODIDUM RUBRUM

RED IODIDE OF MERCURY

**Synonyms.**—Mercuric Iodide ; Biniodide of Mercury.

**Composition.**—$HgI_2$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Perchloride of Mercury . . | 4 ounces |
| Iodide of Potassium . . . | 5 ounces |
| Boiling Distilled Water . . | 2 quarts |

Dissolve the perchloride of mercury in three pints, and the iodide of potassium in the remainder, of the water, and mix the solutions. When the temperature has fallen to that of the atmosphere, decant the supernatant liquor from the precipitate (red iodide of mercury), and, having collected the latter on a filter, wash it twice with cold distilled water, and dry it at a temperature not exceeding 212° F.

| Mercuric Chloride. | | Potassium Iodide. | | Mercuric Iodide (Red Iodide of Mercury). | | Potassium Chloride. |
|---|---|---|---|---|---|---|
| $HgCl_2$ | + | 2KI | = | $HgI_2$ | + | 2KCl |

**Characters and Tests.**—Crystalline powder of a vermilion colour, becoming yellow when gently heated over a lamp on a sheet of paper; almost insoluble in water, dissolves sparingly in alcohol, but freely in ether or in an aqueous solution of iodide of potassium. Should be completely soluble in ether and entirely volatilised at a high temperature. Heated in a dry test-tube with dry sodium carbonate, a sublimate, consisting of bluish-white liquid metallic globules, is obtained, indicating the presence of mercury; the residue remaining in the bottom of the test-tube, dissolved in water acidified with nitric acid, and mixed with solution of starch, produces a blue colour (starch iodide), proving the compound to be an iodide.

**Actions and Uses.**—*Externally*:—In the form of Unguentum Hydrargyri Iodidi Rubri, as a stimulant, detergent, and caustic to unhealthy sores and farcy ulcers; also as a counter-irritant to sore throats, splints, curbs,

spavins, enlarged bursæ, thickening of the integument, and similar affections.

**Preparation.**—Unguentum Hydrargyri Iodidi Rubri.

## HYDRARGYRI OXIDUM RUBRUM

### RED OXIDE OF MERCURY

**Synonyms.**—Mercuric Oxide ; Red Precipitate.

**Composition.**—HgO.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Mercury, by weight . . | 8 ounces |
| Nitric Acid . . . . | 4½ fluid ounces |
| Water . . . . . | 2 fluid ounces |

Dissolve half the mercury in the nitric acid diluted with the water, evaporate the solution to dryness, and, with the dry salt (mercuric nitrate) thus obtained, triturate the remainder of the mercury until the two are uniformly blended together. Heat the mixture in a porcelain dish, with repeated stirring, until acid vapours (nitric peroxide) cease to be evolved, and, when cold, enclose the product (red oxide of mercury) in a bottle. The chemical changes in the foregoing process may be regarded as taking place in two stages, thus :

| | Mercury. | | Nitric Acid. | | Mercuric Nitrate. | | Nitric Oxide. | |
|---|---|---|---|---|---|---|---|---|
| 1. | $3Hg$ | + | $8HNO_3$ | = | $3Hg(NO_3)_2$ | + | $2NO$ | + |

Water.
$4H_2O$

| | Mercuric Nitrate. | | Mercury. | | Mercuric Oxide (Red Oxide of Mercury). | | Nitric Peroxide. |
|---|---|---|---|---|---|---|---|
| 2. | $2Hg(NO_3)_2$ | + | $2Hg$ | = | $4HgO$ | + | $4NO_2$[1] |

[1] Mercuric nitrate heated alone furnishes mercuric oxide, nitric peroxide, and free oxygen [$2Hg(NO_3)_2 = 2HgO + 4NO_2 + O_2$], the latter pro-

**Characters and Tests.**—Orange-red powder, readily dissolved by hydrochloric acid, yielding a solution of mercuric chloride, which gives a yellow precipitate (yellow mercuric oxide) with excess of caustic potash, and a white precipitate (mercuric ammonium chloride, $NH_2HgCl$) with ammonia. Strongly heated in a test-tube, it should be entirely converted into mercury, which condenses in the upper and cooler parts of the tube, and into oxygen, which may be recognised by its rekindling an incandescent match. During the heating of the oxide no red vapours should be evolved, showing its freedom from nitrate.

**Actions and Uses.**—*Externally*:—As an escharotic, the powder is sprinkled over wounds; in the form of Unguentum Hydrargyri Oxidi Rubri it is used as a stimulant and detergent.

**Preparation.**—Unguentum Hydrargyri Oxidi Rubri. Hydrargyri Oleatum (Oleate of Mercury) is prepared by mixing oleic acid with 10 to 20 per cent. of mercuric oxide. It is used in articular inflammation, exostoses, indurations of the udder, and other glandular and indolent swellings, and as an antiparasitic.

## HYDRARGYRI PERCHLORIDUM

### PERCHLORIDE OF MERCURY

**Synonyms.**—Mercuric Chloride; Bichloride of Mercury; Corrosive Sublimate; Mercury Ball.

duct escaping into the air and becoming wasted. In the 'British Pharmacopœia' process, above described, this oxygen combines with the additional mercury, and is thereby economised.

**Composition.**—$HgCl_2$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Sulphate of Mercury, dried . . | 20 ounces |
| Chloride of Sodium, dried . . | 16 ounces |
| Black Oxide of Manganese [1] . . | 1 ounce |

Reduce the sulphate of mercury and the chloride of sodium separately to fine powder, and having mixed them and the oxide of manganese thoroughly by trituration in a mortar, put the mixture into a subliming apparatus, and apply sufficient heat to cause the vapours of perchloride of mercury to rise into the less heated part of the apparatus.

| Mercuric Sulphate. | | Sodium Chloride. | | Mercuric Chloride (Perchloride of Mercury). | | Sodium Sulphate. |
|---|---|---|---|---|---|---|
| $HgSO_4$ | + | $2NaCl$ | = | $HgCl_2$ | + | $Na_2SO_4$ |

**Characters and Tests.**—Heavy colourless masses of prismatic crystals, possessing a highly acrid, metallic taste ; more soluble in alcohol, and still more so in ether, than in water. Heated, it liquefies, and completely sublimes without decomposing. Its aqueous solution gives a yellow precipitate (yellow mercuric oxide) with caustic potash, a white precipitate (mercuric-ammonium chloride, $NH_2HgCl$) with ammonia, and a curdy white precipitate (silver chloride) with silver nitrate ; the first two reactions are indicative of the salt being a mercuric compound, and the third reaction shows that it is a chloride.

**Actions and Uses.**—*Internally* :—In large doses, irritant and corrosive poison ; seldom administered medicinally, although, in doses from five to ten grains, it has been recommended in glanders and farcy. *Externally* :—

[1] If the mercuric sulphate (sulphate of mercury) employed be free from mercurous sulphate, the addition of the oxide of manganese may be dispensed with, as in that case the production of calomel need not be apprehended.

A most powerful antiseptic and disinfectant.[1] Caustic to indolent ulcers, fistulæ, and bony tumours; as a stimulant in scab, mange, and similar affections; also as an antiparasitic for the destruction of pediculi, acari, &c. With suitable precautions watery solutions are used for disinfecting infected premises and carcases of animals which have died of contagious diseases.

**Modes of Application.**—As a caustic, the powder; Liquor Hydrargyri Perchloridi; Unguentum Hydrargyri Perchloridi. As a stimulant and antiparasitic, Spiritus Hydrargyri Perchloridi Compositus.

**Antidotes.**—Albumen; wheaten flour; bean or pea meal; gruel or other demulcent drinks, followed by vegetable astringents.

**Preparations.**—Liquor Hydrargyri Perchloridi; Spiritus Hydrargyri Perchloridi Compositus; Unguentum Hydrargyri Perchloridi.

## HYDRARGYRI SUBCHLORIDUM

### SUBCHLORIDE OF MERCURY

**Synonyms.**—Mercurous Chloride; Protochloride of Mercury; Calomel.

**Composition.**—HgCl, or probably $Hg_2Cl_2$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Sulphate of Mercury . . . | 10 ounces |
| Mercury . . . . . | 7 ounces |
| Chloride of Sodium, dried . . | 5 ounces |
| Boiling Distilled Water . . | a sufficiency |

[1] Koch, in 1880, demonstrated that mercuric chloride in a solution of 1 to 1000 would soon destroy the spores of anthrax. The addition of a small quantity of hydrochloric or tartaric acid is an improvement, as it prevents the coagulation of albumen

Moisten the sulphate of mercury with some of the water, and rub it and the mercury together until globules are no longer visible; add the chloride of sodium, and thoroughly mix the whole by continued trituration. Sublime by a suitable apparatus into a chamber of such a size that the calomel, instead of adhering to its sides as a crystalline crust, shall fall as a fine powder on its floor. Wash this powder with boiling distilled water until the washings cease to be darkened by a drop of ammonium sulphide, showing that any corrosive sublimate that may have been formed in the above process has been removed from the calomel.

| Mercuric Sulphate. | | Mercury. | | Sodium Chloride. | | Mercurous Chloride (Calomel). | | Sodium Sulphate. |
|---|---|---|---|---|---|---|---|---|
| $2HgSO_4$ | + | $Hg_2$ | + | $4NaCl$ | = | $4HgCl$ | + | $2Na_2SO_4$ |

**Characters and Tests.**—Dull white, heavy, and nearly tasteless powder, rendered yellowish by trituration in a mortar; insoluble in water, spirit, and ether. Digested with solution of potash, it becomes black (mercurous oxide), indicating that it is a mercurous compound; and the clear solution obtained by decantation, or filtering the mixture, acidulated with nitric acid, gives a copious white precipitate (silver chloride) with silver nitrate, showing it to be a chloride. Should be entirely volatilised by a sufficient heat, indicating the absence of fixed impurity; and ether, which has been shaken with it in a bottle, should leave no residue on evaporation, proving the calomel to be free from corrosive sublimate.

**Actions and Uses.**—*Internally*:—In continued doses it produces ptyalism; in excessive doses, irritant poison; in medicinal doses, cathartic, stimulant, sedative, resolvent, cholagogue, diuretic, sialagogue, and, on some animals, diaphoretic, alterative, antiphlogistic, anthelmintic. Given

in diseases of the liver; combined with opium or belladonna, in enteritis and dysentery; in febrile affections, in inflammation of serous membranes, in typhous inflammation, in lingering inflammation of the intestines, and in chronic glandular enlargements. As a cathartic it should never be given alone to the horse, as its action on this animal is both uncertain and violent. Calomel is seldom given to cattle or sheep, except in cases of obstinate dysentery.

**Doses.**—Horse, 20 to 60 grains.—Cattle, 10 to 30 grains.—Sheep, 4 to 8 grains.—Pig, 1 to 4 grains.—Dog, 1 to 4 grains.

**Modes of Application.**—In the form of bolus, or suspended in some mucilaginous drink.

**Incompatibles.**—Lime-water; alkalies; alkaline carbonates.

**Antidotes.**—Albumen; wheaten flour; bean or pea meal.

**Preparation.**—Lotio Hydrargyri Nigra.

## HYDRARGYRUM

MERCURY

**Synonym.**—Quicksilver.

**Composition.**—An element, symbol Hg.

**Mode of Preparation.**—Although it is met with in the native metallic state in the quicksilver-mines of South America, it is chiefly imported into this country from Spain, Illyria, Bavaria, and several other parts of the world, where it is extracted from a native sulphide termed cinnabar (mercuric sulphide). In order that the cinnabar

may be made to yield the mercury it contains, it is either roasted or heated with lime or iron in a suitable distillatory apparatus. The following equation represents the changes which take place in the simpler of these processes:

| Mercuric Sulphide. | | Iron. | | Mercury. | | Ferrous Sulphide. |
|---|---|---|---|---|---|---|
| $2HgS$ | + | $Fe_2$ | = | $Hg_2$ | + | $2FeS$ |

The mercury distils over, and the ferrous sulphide remains in the retort.

**Characters and Tests.**—A metal, liquid at ordinary temperature, brilliantly lustrous, and easily divisible into spherical globules. Volatilises at a heat below that of visible redness, and should leave no residue.

## Preparations.

Hydrargyri Iodidum Rubrum.
„ Oxidum Rubrum.
„ Perchloridum.
„ Subchloridum.
„ Sulphas.
Hydrargyrum Ammoniatum
Hydrargyrum cum Cretâ.
Lotio Hydrargyri Flava.
„ Hydrargyri Nigra.
Liquor Hydrargyri Nitratis Acidus.
Liquor Hydrargyri Perchloridi
Pilula Hydrargyri Subchloridi et Ferri.
Unguentum Hydrargyri.
„ Hydrargyri Ammoniati.
Unguentum Hydrargyri Compositum.
Unguentum Hydrargyri Nitratis.
Unguentum Hydrargyri Oxidi Rubri.
Unguentum Hydrargyri Iodidi Rubri.

## HYDRARGYRUM AMMONIATUM

### AMMONIATED MERCURY

**Synonyms.**—Hydrargyri Ammonio-chloridum ; Hydrargyri Præcipitatum Album ; Mercuric-ammonium Chloride ; Chloride of Mercuric Ammonium ; Amidochloride of Mercury ; White Precipitate.

**Composition.** — Various views have been adopted respecting the constitution of this body ; it may, however, be regarded as the ammonium chloride in which two atoms of hydrogen are replaced by a bivalent atom of mercury ; its formula then becomes $NH_2HgCl$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Perchloride of Mercury . . | 3 ounces |
| Solution of Ammonia . . | 4 fluid ounces |
| Distilled Water . . . | 3 pints |

Dissolve the perchloride of mercury in the water, with the aid of a moderate heat. Mix the solution with the ammonia, constantly stirring. Collect the precipitate (ammoniated mercury) on a filter, and wash it well with cold distilled water until the liquid which passes through ceases to give a precipitate when dropped into a solution of silver nitrate acidulated by nitric acid. Lastly, dry the product at a temperature not exceeding 212° F.

| Mercuric Chloride. | | Ammonia. | | Mercuric-ammonium Chloride (Ammoniated Mercury). | |
|---|---|---|---|---|---|
| $HgCl_2$ | + | $2NH_4HO$ | = | $NH_2HgCl$ | + |

| Ammonium Chloride. | | Water. |
|---|---|---|
| $NH_4Cl$ | + | $2H_2O$ |

**Characters and Tests.**—Opaque white powder, on which cold water, alcohol, and ether have no action. Digested with caustic potash, it evolves ammonia and acquires a pale yellow colour (mercuric oxide). The clear liquid obtained by filtering this mixture, after being acidulated with nitric acid, gives a white precipitate (silver chloride) with silver nitrate. Boiled with stannous chloride, it becomes grey and affords globules of metallic mercury. Entirely volatilises at a heat under redness.

**Action and Use.**—*Internally*:—Irritant poison. *Externally*:—The powder, or an ointment consisting of one part of white precipitate and eight parts of lard, is used for the destruction of pediculi and acari.

## HYDRARGYRUM CUM CRETA

MERCURY WITH CHALK

**Synonym.**—Grey powder.

**Composition.**—An intimate mixture of metallic mercury, mercurous oxide, and calcic carbonate. The proportion of mercurous oxide, which is probably the active ingredient of this preparation, to the other constituents has not yet been ascertained, but it is most likely variable.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Mercury, by weight . . | 1 ounce |
| Prepared Chalk . . . | 2 ounces |

Rub the mercury and chalk together in a porcelain mortar until metallic globules cease to be visible to the naked eye, and the mixture acquires a uniform grey colour.

**Characters and Tests.**—Powder of a light-grey colour, free from grittiness, insoluble in water, partly dissolved with effervescence by dilute hydrochloric acid, leaving the mercury in a finely divided state. The solution formed with hydrochloric acid is not precipitated by the addition of stannous chloride, showing that the preparation contains no mercuric oxide.

**Action and Use.**—Chiefly given to the dog as an alterative.

**Dose.**—Dog, from 3 to 10 grains.

**Mode of Application.** — In the form of pill or powder.

**Incompatibles.**—Acids and acidulous salts.

## HYOSCYAMI FOLIA

HYOSCYAMUS-LEAVES

**Synonym.**—Henbane-leaves.

The fresh leaves, with the branches to which they are attached, of *Hyoscyamus niger*; also the leaves separated from the branches and carefully dried, gathered from wild or cultivated British biennial plants, when about two-thirds of the flowers are expanded.

**Natural Order.**—*Solanaceæ.*

**Characters.**—Leaves sinuated, clammy, and hairy. The fresh herb has a strong unpleasant odour, and a slightly acrid taste, which nearly disappear on drying. The fresh juice dropped into the eye dilates the pupil.

**Preparations.** — Extractum Hyoscyami; Tinctura Hyoscyami.

## INFUSUM CALUMBÆ

INFUSION OF CALUMBA

Take of

| | |
|---|---|
| Calumba, in coarse powder | 1 ounce |
| Cold water | 1 pint |

Macerate one hour, and strain.

**Doses.**—Horse, 6 to 8 fluid ounces.—Cattle, 8 to 10 fluid ounces.—Sheep, 2 to 4 fluid ounces.—Pig, 1 to 2 fluid ounces.—Dog, $\frac{1}{2}$ to 1 fluid ounce.

## INFUSUM CATECHU

INFUSION OF CATECHU

Take of

| | |
|---|---|
| Pale Catechu, in coarse powder | 6 drachms |
| Cinnamon-bark, bruised | 1 drachm |
| Boiling Distilled Water | 1 pint |

Infuse in a covered vessel for half a hour, and strain.

**Doses.**—Horse, 4 to 6 fluid ounces.—Cattle, 4 to 6 fluid ounces.—Sheep, 1 to 2 fluid ounces.—Pig, 1 to 2 fluid ounces.—Dog, $\frac{1}{4}$ to 1 fluid ounce.

## INFUSUM CINCHONÆ FLAVÆ

INFUSION OF YELLOW CINCHONA

Take of

| | |
|---|---|
| Yellow Cinchona Bark, in coarse powder | 1 ounce |
| Boiling Distilled Water | 1 pint |

Infuse in a covered vessel for two hours, and strain.

**Doses.**—Horse, 4 to 6 fluid ounces.—Cattle, 4 to 6 fluid ounces.—Sheep, 1 to 2 fluid ounces.—Pig, 1 to 2 fluid ounces.—Dog, $\frac{1}{4}$ to 1 fluid ounce.

## INFUSUM CUSSO

INFUSION OF KOUSSO

Take of

| | |
|---|---|
| Kousso, in coarse powder . . | $\frac{1}{2}$ ounce |
| Boiling Distilled Water . . | 8 fluid ounces |

Infuse in a covered vessel for fifteen minutes. Not to be strained.

**Doses.** — Horse, 20 to 30 fluid ounces. — Cattle, 20 to 40 fluid ounces.—Sheep, 6 to 10 fluid ounces.—Pig, 3 to 5 fluid ounces.—Dog, 1 to 3 fluid ounces.

## INFUSUM ERGOTÆ

INFUSION OF ERGOT

Take of

| | |
|---|---|
| Ergot, in coarse powder . . | $\frac{1}{2}$ ounce |
| Boiling Distilled Water . . | 1 pint |

Infuse in a covered vessel for half an hour, and strain.

**Doses.**—Horse, 6 to 8 fluid ounces.—Cattle, 8 to 10 fluid ounces.—Sheep, 4 to 6 fluid ounces.—Pig, 2 to 4 fluid ounces.—Dog, $\frac{1}{2}$ to 1 fluid ounce.

## INFUSUM GENTIANÆ COMPOSITUM

COMPOUND INFUSION OF GENTIAN

Take of

| | |
|---|---|
| Gentian-root, sliced . . | 4 drachms |
| Pimento-berries, bruised . . | 2 drachms |
| Boiling Distilled Water . . | 1 pint |

Infuse in a covered vessel for four hours, and strain.

**Doses.**—Horse, 6 to 8 fluid ounces.—Cattle, 8 to 10 fluid ounces.—Sheep, 2 to 4 fluid ounces.—Pig, 1 to 2 fluid ounces.—Dog, ½ to 1 fluid ounce.

## INFUSUM LINI

### INFUSION OF LINSEED

**Synonym.**—Linseed Tea.

Take of

| | |
|---|---|
| Linseed, crushed . . . | 1 ounce |
| Boiling Distilled Water . . | 1 pint |

Infuse in a covered vessel for four hours, and strain.

**Dose.**—*Ad libitum.*

## INFUSUM TABACI

### INFUSION OF TOBACCO

Take of

| | |
|---|---|
| Tobacco, in leaf or shreds . . | 1 drachm |
| Boiling Distilled Water . . | 1 pint |

Infuse in a covered vessel for one hour, and strain.

**Use.**—*Externally*:—As a parasiticide.

## IODOFORMUM

### IODOFORM

**Synonym.**—Tri-iodomethane.

**Composition.**—$CHI_3$.

**Mode of Preparation.**—By mixing together hot alcoholic solutions of Iodine and Potassium Carbonate.

**Characters and Tests.**—It occurs in yellow, scaly crystals having a persistent, disagreeable odour and taste;

is scarcely soluble in cold water, but dissolves in ether and chloroform; is very volatile, and should be kept in well-stoppered bottles.

**Actions and Uses.**—Antiseptic and deodoriser, it is devoid of local irritant effect, but is a local anæsthetic. A few grains dusted over ulcers or malignant growths deodorise and promote their healing. Also used as a dressing for ordinary wounds.

## IODOL

TETRA-IODOPYROL

**Composition.**—$C_4I_4NH$.

**Mode of Preparation.**—By the action of iodine upon pyrol, a constituent of mineral oil.

**Characters and Tests.**—A greyish-white powder, free from the disagreeable odour of iodoform.

**Uses.**—Iodol may be dusted over wounds in the form of a powder, or it may be used as an ointment.

## IODUM

IODINE

**Composition.**—A solid non-metallic element, having the symbol I.

**Mode of Preparation.**—Seaweed is collected, dried, and burned; the semi-vitreous ash (kelp) which remains is lixiviated with water, and the aqueous solution of alkaline sulphates, carbonates, chlorides, and iodides thus formed is evaporated. Sodium sulphate and carbonate, and potassium and sodium chloride, crystallise out and leave a 'mother liquor' rich in iodides, chiefly

sodium iodide. This 'mother liquor' is mixed with sulphuric acid and manganese peroxide and subjected to distillation.

| Sodium Iodide. | | Sulphuric Acid. | | Manganese Peroxide. | | Iodine. | | Sodium Sulphate. | |
|---|---|---|---|---|---|---|---|---|---|
| $2NaI$ | + | $2H_2SO_4$ | + | $MnO_2$ | = | $I_2$ | + | $Na_2SO_4$ | + |

| Manganese Sulphate. | | Water. |
|---|---|---|
| $MnSO_4$ | + | $2H_2O$ |

The iodine distils over and is collected in a series of glass receivers; the other products remain in the retort.

Iodine thus obtained is purified by being resublimed; it should be preserved in a well-stoppered bottle.

**Characters and Tests.**—Laminar crystals, of a peculiar odour, bluish-black colour, and metallic lustre, which, when heated, yield a beautiful violet-coloured vapour; very sparingly soluble in water, but freely dissolved by alcohol, ether, and a solution of potassium iodide. Its aqueous solution strikes a deep blue colour (starch iodide) with cold solution of starch. It should sublime without leaving any residue, and the portion which first comes over should not include any pungent, slender, colourless prisms (iodine cyanide). If shaken in a dry bottle, iodine will not adhere to the sides if it be free from moisture.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant and corrosive poison; in medicinal doses, tonic, alterative, deobstruent, resolvent; if exhibited for a sufficient length of time, it occasions debility and gives rise to a disturbance of the constitution termed *iodism*. Given as a tonic and alterative in inflammatory affections after the subsidence of fever and acute inflammation; as a resolvent and stimulant in ascites, hydrothorax, and similar complaints in which effusion is present; as a

deobstruent in chronic enlargement of the liver and the mammary and other glands, also in indurations of mucous membranes. Likewise given in rheumatism, in scrofulous diseases, and to arrest the secretion of urine in diabetes insipidus. Recommended as an antidote to poisoning by mercury and the vegetable alkaloids. *Externally* :— Stimulant, counter-irritant, resolvent in sore throat, bursal enlargements, swellings of joints, thickening of the periosteum, tumours of various kinds, strains of tendons ; also in scab, mange, ringworm, mallenders and sallenders, and other skin-diseases.

**Doses.**—HORSE, 10 to 20 grains.—CATTLE, 20 to 30 grains.—SHEEP, 5 to 10 grains.—PIG, 2 to 5 grains.—DOG, 1 to 2 grains.

**Modes of Application.**—*Internally* :—In the form of bolus, preferably as Liquor Iodi, sometimes as Tinctura Iodi. *Externally* :—Linimentum Iodi ; Tinctura Iodi ; Unguentum Iodi ; Unguentum Sulphuris Iodidi.

**Incompatibles.**—Ammonia ; metallic salts ; mineral acids ; vegetable alkaloids.

**Antidotes.** — Emetics, when applicable, aided by demulcent drinks ; starch, flour, and similar amylaceous meals, suspended in water. If the starch-containing remedies be boiled with water and cooled before administration, their efficiency will be increased.

## Preparations

Ferri Iodidum.
Hydrargyri Iodidum Rubrum.
Linimentum Iodi Compositum.
Liquor Iodi et Potassii Iodidi.
Potassii Iodidum.
Sulphuris Iodidum.
Tinctura Iodi.
Unguentum Iodi.
Unguentum Sulphuris Iodidi.

## IPECACUANHA

IPECACUANHA

The dried root of *Psychotria Ipecacuanha*, powdered. Imported from Brazil.

**Natural Order.**—*Rubiaceæ.*

**Composition.**—Its medicinal activity is ascribed to an alkaloid, *emetine*. The formula of this body has not yet been well established.

**Characters.**—Pieces three or four inches long, about the diameter of a small quill, contorted and irregularly annulated. Colour, brown of various shades. It consists of two parts—the cortical or active portion, which is brittle, and a slender, tough, white, woody centre. Powder pale brown, with a faint nauseous odour and a somewhat acrid and bitter taste.

**Actions and Uses.**—*Internally* :—In excessive doses irritant poison ; in medicinal doses, emetic and sedative ; also mildly cathartic. Chiefly used in dog practice, and in the treatment of coughs and colds.

**Doses.**—As an emetic to the Dog, from 15 to 30 grains are given in tepid water, either alone or with 1 grain of tartar emetic.

**Preparations.**—Pulvis Ipecacuanhæ Compositus ; Vinum Ipecacuanhæ.

## JALAPA

JALAP

The dried tubercles of *Ipomœa Purga*, powdered. Imported from Mexico, from a town of which, Xalapa, its name is derived.

**Natural Order.**—*Convolvulaceæ.*

**Composition.**—Its cathartic principle is a resinous body, soluble in alcohol, insoluble in water.

**Characters.**—Varying from the size of a nut to that of an orange, ovoid, the larger tubercles frequently incised, covered with a thin, brown, wrinkled cuticle; presenting, when cut, a yellowish-grey colour, with dark-brown concentric circles.

**Actions and Uses.**—*Internally* :—In excessive doses, irritant poison; in medicinal doses, it is sometimes given to sheep and pigs, but usually to the dog, as a cathartic and vermifuge.

**Doses.**—Pig, 1 to 2 drachms.—Dog, ½ to 3 drachms.

**Modes of Application.**—In the form of bolus or rubbed up with some mucilaginous drink.

**Preparation.**—Extractum Jalapæ, which is a constituent of purgative pills.

## KAMALA

KAMALA

**Synonym.**—Kamela.

Minute glands, which are found adhering to the capsules of *Rottlera tinctoria*. Imported from India.

**Natural Order.**—*Euphorbiaceæ.*

**Composition.**—It contains a yellow crystalline principle called *rottlerin.*

**Characters and Tests.**—A fine, granular, mobile, orange-red powder; scarcely miscible with water, but soluble for the most part in alcohol and ether, forming

red-coloured solutions. The residue insoluble in ether should consist principally of tufted hairs, and be free from sand and earthy impurities.

**Actions and Uses.**—Purgative and anthelmintic for tænia.

**Doses.**—Horse, 1 to 2 ounces.—Dog, 1 to 2 drachms.

**Modes of Application.**—In the form of bolus or suspended in gruel.

## KINO

KINO

The inspissated juice obtained from incisions made in the trunk of *Pterocarpus Marsupium*. Imported from Malabar.

**Natural Order.**—*Leguminosæ*.

**Composition.** — Kino contains kino-tannic acid, kinoin, pyrocatechin, pectin, &c.

**Characters and Tests.**—Small, angular, brittle, glistening reddish-black fragments. When entire they are opaque, but in thin laminæ are translucent and ruby-red. They are inodorous, but possess a very astringent taste. Soluble in alcohol, insoluble in ether. The mineral acids, the solutions of gelatin, tartarated antimony, acetate of lead, perchloride of iron, and nitrate of silver produce precipitates with the watery infusion.

**Actions and Uses.**
**Doses.**
**Modes of Application.**
} Same as Catechu Pallidum

## LINI FARINA

LINSEED MEAL

Linseed, or the cake from which the greater part of the oil has been expressed, reduced to powder.

**Natural Order.**—*Linaceæ.*

**Composition.**—Chiefly valuable in medicine on account of the mucilage it contains.

**Preparations.**—Cataplasma Calcis Chlorinatæ ; Cataplasma Carbonis ; Cataplasma Lini ; Cataplasma Sinapis.

## LINI SEMINA

LINSEED

The seeds of the common flax, *Linum usitatissimum.* Cultivated in Britain.

**Natural Order.**—*Linaceæ.*

**Composition.**—Its value in medicine depends upon the mucilage it contains and the oil that is obtained from it by expression.

**Characters.**—Small, oval, pointed, flat, with acute edges ; smooth, shining, brown externally, yellowish-white internally ; mucilaginous oily taste.

**Preparations.**—Infusum Lini ; Lini Farina ; Oleum Lini.

## LINIMENTUM ACIDI CARBOLICI

LINIMENT OF CARBOLIC ACID

This is synonymous with Glycerinum Acidi Carbolici, which see.

## LINIMENTUM ACONITI

### LINIMENT OF ACONITE

Take of

| | |
|---|---|
| Aconite-root, in coarse powder | 20 ounces |
| Camphor | 1 ounce |
| Rectified Spirit | a sufficiency |

Moisten the root for three days, then pack in a percolator, and pour sufficient rectified spirit upon it to produce a pint.

**Use.**—Applied with a camel-hair pencil alone, or mixed, in equal proportions, with soap liniment or compound camphor liniment, and rubbed on the part, it relieves acute rheumatism.

## LINIMENTUM AMMONIÆ

### LINIMENT OF AMMONIA

Take of

| | |
|---|---|
| Solution of Ammonia B.P. | 1 fluid ounce |
| Olive Oil | 2 fluid ounces |

Mix together with agitation.

**Use.**—Counter-irritant in sore throat, rheumatism, sprains, chronic tumours, bronchitis, &c.

## LINIMENTUM AMMONIÆ COMPOSITUM

### COMPOUND LINIMENT OF AMMONIA

Take of

| | |
|---|---|
| Solution of Ammonia B.P. | equal parts |
| Olive Oil | |
| Oil of Turpentine | |

Shake the solution of ammonia and the olive oil together, then add the oil of turpentine, and shake again until the ingredients are thoroughly mixed.

**Use.**—Same as Linimentum Ammoniæ, but more active.

## LINIMENTUM CALCIS

### LINIMENT OF LIME

**Synonym.**—Carron Oil.[1]

Take of

| | |
|---|---|
| Solution of Lime . . . . | equal parts |
| Olive Oil[2] . . . . . | |

Mix together with agitation.

**Use.**—Applied to burns and scalds.

## LINIMENTUM CAMPHORÆ

### LINIMENT OF CAMPHOR

Take of

| | |
|---|---|
| Camphor . . . . . | 1 ounce |
| Olive Oil[3]. . . . . | 4 fluid ounces |

Dissolve the camphor in the oil.

**Uses.**—A stimulating embrocation for deep-seated inflammations, glandular swellings, &c.

[1] Carron Oil is usually made with linseed oil, as it is cheaper.—Ed.

[2] Sometimes one part of oil of turpentine is added to this liniment to increase its activity.

[3] Oil of turpentine is sometimes added to increase the activity of this liniment

## LINIMENTUM CAMPHORÆ COMPOSITUM

### COMPOUND LINIMENT OF CAMPHOR

**Synonym.**—White Oil.

Take of

| | |
|---|---|
| Camphor . . . . . | 1 ounce |
| Rectified Spirit . . . | 4 fluid ounces |
| Olive Oil . . . . . | 1 pint |
| Solution of Ammonia . . | 2 fluid ounces |

Dissolve the camphor in the spirit, shake the olive oil with the solution of ammonia, and mix the whole together, so as to form a liniment.

**Uses.**—Same as Linimentum Camphoræ and Linimentum Ammoniæ, but more active.

## LINIMENTUM CANTHARIDIS

### LINIMENT OF CANTHARIDES

**Synonyms.**—Oleum Cantharidis; Oil of Cantharides.

Take of

| | |
|---|---|
| Cantharides, in powder . . . | 1 part |
| Olive Oil[1] [2] . . . . . | 8 parts |

Digest in a water-bath for two or more hours, and strain.

**Use.**—Vesicant.

[1] Or six to ten parts of rape or linseed oil.—Ed.

[2] Oil of turpentine is sometimes added to increase the activity of this liniment.

## LINIMENTUM CREOSOTI COMPOSITUM

COMPOUND LINIMENT OF CREOSOTE

Take of

| | |
|---|---|
| Creosote | 2 parts |
| Oil of Turpentine | 4 parts |
| Olive Oil | 4 parts |

Mix together with agitation.

**Uses.**—Similar to those of Glycerinum Acidi Carbolici.

## LINIMENTUM CROTONIS

LINIMENT OF CROTON OIL

Take of

| | |
|---|---|
| Croton Oil | 1 part |
| Oil of Turpentine, or Soap Liniment | 6 to 8 parts |

Mix together with agitation.

**Use.**—Vesicant.

## LINIMENTUM CUPRI SUBACETATIS

LINIMENT OF SUBACETATE OF COPPER [1]

**Synonyms.**—Linimentum Æruginis; Liniment of Verdigris.

Take of

| | |
|---|---|
| Subacetate of Copper,[2] in fine powder | 9 ounces |
| Alum, in fine powder | 6 ounces |
| Treacle | $1\frac{1}{2}$ pound |

[1] A mixture of one part of subacetate of copper and eight parts of lard is generally used instead of this liniment.

[2] The less expensive sulphate of copper may be substituted for the subacetate in this preparation.

Boil and stir together until the mixture assumes a brown colour.

**Uses.**—Erodent and digestive in canker, thrush, and foot-rot.

## LINIMENTUM CUPRI SULPHATIS

### LINIMENT OF SULPHATE OF COPPER

Take of

| | |
|---|---|
| Sulphate of Copper, in fine powder . | 1 part |
| Tar or Treacle . . . . . | 4 parts |

Mix, stir, and heat together until the mass assumes a reddish-brown colour.

**Uses.**—Same as Linimentum Cupri Subacetatis.

## LINIMENTUM HYDRARGYRI

### COMPOUND LINIMENT OF MERCURY

Take of

| | |
|---|---|
| Ointment of Mercury . . . | 3 ounces |
| Solution of Ammonia . . . | 3 ounces |
| Liniment of Camphor . . . | 3 ounces |

Melt the ointment in the liniment, add the ammonia, and shake them together.

**Uses.**—A stimulating liniment, applied to indolent ulcers, &c.

## LINIMENTUM IODI COMPOSITUM

COMPOUND LINIMENT OF IODINE

**Synonym.**—Linimentum Iodi, B.P.

Take of

| | |
|---|---|
| Iodine | 1¼ ounce |
| Iodide of Potassium | ½ ounce |
| Camphor | ¼ ounce |
| Rectified Spirit | 10 fluid ounces |

Dissolve the iodine, iodide of potassium, and camphor in the spirit.

**Uses.**—Stimulant and deobstruent to bursæ and enlarged glands.

## LINIMENTUM PICIS LIQUIDÆ

LINIMENT OF TAR

Take of

| | |
|---|---|
| Oil of Tar | equal parts |
| Rape Oil | |

Mix together with agitation.

**Use.**—In skin-affections.

## LINIMENTUM OLEI PICIS LIQUIDÆ COMPOSITUM

COMPOUND LINIMENT OF TAR

Take of

| | |
|---|---|
| Oil of Turpentine | equal parts |
| Oil of Tar | |
| Rape Oil | |

Mix together with agitation.

**Use.**—In skin-affections.

## LINIMENTUM OPII

LINIMENT OF OPIUM

Take of

| | |
|---|---|
| Tincture of opium, by measure | 1 part |
| Compound Liniment of Soap, by measure | 3 parts |

Mix.

**Use.**—Anodyne to local pains and sprains; in rheumatism, &c.

## LINIMENTUM PLUMBI SUBACETATIS

LINIMENT OF SUBACETATE OF LEAD

Take of

| | |
|---|---|
| Solution of Subacetate of Lead | 1 part |
| Olive Oil | 4 parts |

Mix.

**Uses.**—Sedative to excoriated surfaces, or after the application of a blister or the actual cautery.

## LINIMENTUM SAPONIS COMPOSITUM

COMPOUND LINIMENT OF SOAP

**Synonym.**—Opodeldoc.

Take of

| | |
|---|---|
| Soft Soap | 4 ounces |
| Camphor | 1 ounce |
| Proof Spirit | 2 pints |
| Solution of Ammonia | ½ pint |

Dissolve the soap and the camphor in the spirit, then add the solution of ammonia, and filter for use.

**Uses.**—Stimulant and anodyne, after the subsidence of local inflammation ; to sprains and bruises ; to tumours, slight attacks of sore throat, &c.

## LINIMENTUM TEREBINTHINÆ

LINIMENT OF TURPENTINE

Take of

| | |
|---|---|
| Soft Soap | 2 ounces |
| Camphor | 1 ounce |
| Oil of Turpentine | 16 fluid ounces |

Dissolve the camphor in the oil of turpentine, then add the soap, rubbing them together until they are thoroughly mixed.

**Uses.**—Same as Linimentum Saponis Compositum.

## LINIMENTUM TEREBINTHINÆ COMPOSITUM

COMPOUND LINIMENT OF TURPENTINE

Take of

| | |
|---|---|
| Croton Oil | 2 fluid drachms |
| Oil of Turpentine | 4 fluid ounces |
| Olive or Rape Oil | 25 fluid ounces |

Mix.

**Uses.**—Rubefacient and vesicant.

## LIQUOR ACIDI CARBOLICI

SOLUTION OF CARBOLIC ACID

Take of

| | |
|---|---|
| Carbolic Acid | 1 ounce |
| Water | 40 ounces |

Shake well together.

**Uses.**—Lotion for extensive wounds; after operations, as a styptic, and an antiseptic to prevent the decomposition of any blood that may accumulate from hæmorrhage after the wound has been sewn up. For the destruction of pediculi and acari, it is applied as hot as the animal can bear it.

## LIQUOR ALOES

SOLUTION OF ALOES

Take of

| | |
|---|---|
| Barbados Aloes,[1] in small pieces | 1 pound |
| Distilled Water . . . . | 5 pints |
| Proof Spirit . . . . . | 16 fluid ounces |

Digest the aloes and water together over a water-bath until the former is dissolved, remove the solution from the water-bath, add the spirit, and thoroughly mix.

**Uses.**—Laxative and cathartic.

**Doses.**—Horse, 2 to 4 fluid ounces.—Cattle, 2 to 6 fluid ounces.—Sheep, 1 to 2 fluid ounces.

## LIQUOR ALOES ET SODII CARBONATIS

SOLUTION OF ALOES AND CARBONATE OF SODA

Take of

| | |
|---|---|
| Barbados Aloes, in small pieces . | 1 pound |
| Carbonate of Soda[2] . . . | ½ pound |
| Water . . . . . | 1 gallon |

Dissolve, with the aid of a water-bath, the aloes in seven pints of the water, and, when cold, pour off the

[1] Spiked aloes is frequently substituted for Barbados aloes in this preparation, on account of the more ready solubility of the former.

[2] Four ounces of Potassium Carbonate may be substituted for this.

clear solution. Dissolve the carbonate of soda in the remaining pint of water, previously made hot; allow the solution to cool, add it to the solution of aloes, and preserve the mixture in a well-corked or stoppered bottle.

**Uses.**—Laxative and cathartic.

**Doses.**— Horse, 8 to 16 fluid ounces.—Cattle, 8 to 16 fluid ounces.—Sheep, 2 to 6 fluid ounces.

## LIQUOR ALUMINIS

SOLUTION OF ALUM

Take of

| | |
|---|---|
| Alum, in powder . . . | 1 ounce |
| Distilled Water . . . | 16 fluid ounces |

Dissolve the alum in the water.

**Uses.**—*Externally* :—As an astringent or styptic.

## LIQUOR AMMONIÆ

SOLUTION OF AMMONIA

**Mode of Preparation.**—Take of

| | |
|---|---|
| Strong Solution of Ammonia . . | 1 pint |
| Distilled Water . . . . | 2 pints |

Mix, and preserve in a stoppered bottle.

**Tests.**—Sp. gr. 0·959. Other characters similar to Liquor Ammoniæ Fortis.

**Actions and Uses.**—*Internally* :—In excessive doses and undiluted, irritant and narcotic poison; in medicinal doses, antacid, stimulant, antispasmodic, resolvent, and diuretic. Given in tympanitis, hoven, and many inflammatory and debilitating complaints. *Externally* :—Counter-irritant, rubefacient, vesicant.

**Doses.**—Horse, ½ to 1 fluid ounce.—Cattle, 1 to 4 fluid ounces.—Sheep, 1 to 3 fluid drachms.—Pig, 1 to 3 fluid drachms.—Dog, 5 to 20 minims.

**Modes of Application.**—*Internally*:—Diluted with from twenty to thirty parts of water, with cold gruel, or other mucilaginous drink. As a stimulant, sometimes administered to horses and cattle in the form of Mistura Ammoniæ; in indigestion and colic, one fluid ounce of Liquor Ammoniæ mixed with three to five drachms of aloes, rubbed down in water, may be given to the same animals. *Externally*:—In the form of Linimentum Ammoniæ, Linimentum Ammoniæ Compositum, Linimentum Camphoræ Compositum.

**Incompatibles.**—Acids, and metallic salts, except those of potassium, sodium, barium, and calcium.

**Antidotes.**—Dilute vinegar, and solutions of the non-poisonous vegetable acids.

**Preparations.**—Linimentum Ammoniæ, Linimentum Ammoniæ Compositum, Linimentum Camphoræ Compositum.

## LIQUOR AMMONII ACETATIS

### SOLUTION OF ACETATE OF AMMONIA

**Synonym.**—Mindererus's Spirit.

**Composition.**—Ammonium acetate (acetate of ammonia), $(NH_4)C_2H_3O_2$, dissolved in water.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Acetic Acid . . . . | 10 fluid ounces |
| Carbonate of Ammonia . . | 3¼ ounces, or a sufficiency |
| Distilled Water . . . | 2½ pints |

Powder the carbonate of ammonia, and add it gradually to the acetic acid until a neutral solution is formed, then add the water.

Assuming, for simplicity's sake, the carbonate of ammonia to have the formula $(NH_4)_2CO_3$, the following equation expresses the decomposition which takes place during its conversion into acetate :

| Ammonium Carbonate. | | Acetic Acid. | | Ammonium Acetate (Acetate of Ammonia). | | |
|---|---|---|---|---|---|---|
| $(NH_4)_2CO_3$ | + | $2HC_2H_3O_2$ | = | $2(NH_4)C_2H_3O_2$ | + | |

| Carbonic Anhydride. | | Water. |
|---|---|---|
| $CO_2$ | + | $H_2O$ |

**Characters and Tests.**—Transparent, colourless, and nearly odourless liquid, having a cooling, saline, and unpleasant taste. Mixed with solution of potash it evolves ammoniacal gas, and warmed with strong sulphuric acid it emits the vapour of acetic acid. The ammoniacal gas and the vapour of acetic acid may each be recognised by its peculiar odour.

**Actions and Uses.**—*Internally* :—Mild stimulant, diaphoretic, and diuretic. Given as a febrifuge in inflammation of the lungs, influenza, and other febrile affections, usually in combination with Potassæ Nitras, Spiritus Ætheris Nitrosi, or Spiritus Ætheris Nitrosi and Extractum Belladonnæ. *Externally* :—As a discutient.

**Doses.**—Horse, 1 to 4 fluid ounces.—Cattle, 4 to 8 fluid ounces.—Sheep, 1 to 3 fluid ounces.—Pig, 1 to 3 fluid ounces.—Dog, 2 to 6 drachms.

**Modes of Application.**—*Internally* :—Diluted with about three times its bulk of water. *Externally* :—Lotio Ammoniæ Acetatis.

**Incompatibles.**—Acids ; fixed alkalies and their carbonates ; lime-water ; nitrate of silver ; acetate of lead ; metallic sulphates.

## LIQUOR AMMONIÆ FORTIS

STRONG SOLUTION OF AMMONIA

**Composition.**—Water holding in solution 32·5 per cent. of ammoniacal gas, $NH_3$.

**Mode of Preparation.**—A mixture of calcium hydrate and coarsely powdered ammonium chloride is heated in an iron bottle connected by a tube with a series of Woulffe's bottles. The first two Woulffe's bottles are empty. Two retain impurities that may pass over from the iron bottle, and the others contain water, by which the ammoniacal gas, set free from the ammonium chloride, is absorbed. The aqueous solution of ammonia thus formed constitutes Liquor Ammoniæ Fortis.

| Ammonium Chloride. | | Calcium Hydrate. | | Ammoniacal Gas | | Calcium Chloride. | | Water |
|---|---|---|---|---|---|---|---|---|
| $2(NH_4)Cl$ | + | $Ca(HO)_2$ | = | $2NH_3$ | + | $CaCl_2$ | + | $2H_2O$ |

**Characters and Tests.**—Colourless liquid, with a characteristic and very pungent odour and strong alkaline reaction. Sp. gr. 0·891.

When diluted with four times its volume of distilled water, it should give no precipitate with lime-water, showing the absence of ammonium carbonate ; no precipitate with ammonium oxalate, indicating the absence of calcium ; no precipitate with ammonium sulphide, showing the absence of iron ; and no precipitate with copper ammonio-sulphate, proving the absence of ammonium sulphide. Liquor ammoniæ fortis acidified with nitric acid should not be rendered turbid by silver nitrate, indicating the

absence of chlorine, or by barium chloride, showing that it is free from sulphate.[1]

**Preparations.**—Linimentum Camphoræ Compositum; Liquor Ammoniæ; Spiritus Ammoniæ Aromaticus.

## LIQUOR ANTIMONII CHLORIDI

### SOLUTION OF CHLORIDE OF ANTIMONY

**Synonyms.**—Solution of Antimonious Chloride; Solution of Terchloride of Antimony; Butter of Antimony.

**Composition.**—Water holding in solution antimonious chloride, $SbCl_3$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Black Antimony (Native Antimonious Sulphide), in powder . . . | 1 pound |
| Hydrochloric Acid . . . . | 4 pints |

Place the black antimony in a porcelain vessel, pour upon it the hydrochloric acid, and, constantly stirring, apply to the mixture, beneath a flue with a good draught, a gentle heat, which must be gradually augmented as the evolution of sulphuretted hydrogen gas begins to slacken, until the liquid boils. Maintain it at this temperature for fifteen minutes; then remove the vessel from the fire, and filter the liquid (solution of antimonious chloride) through calico into another vessel, returning what passes through first, that a perfectly clear solution may be

[1] When the boric acid of Tuscany is saturated with soda, considerable quantities of pure ammonia are evolved. The liquor ammoniæ and ammonii carbonas obtained from this source are known in commerce as 'volcanic ammonia,' and are to be preferred to the same preparations procured in any other way on account of their greater purity.

obtained. Evaporate this to the bulk of two pints, and preserve it in a stoppered bottle.

| Antimonious Sulphide. | | Hydrochloric Acid. | | Antimonious Chloride. | | Sulphuretted Hydrogen. |
|---|---|---|---|---|---|---|
| $Sb_2S_3$ | + | 6HCl | = | $2SbCl_3$ | + | $3H_2S$ |

**Characters and Tests.**—Heavy liquid, sp. gr. 1·47, usually of a yellowish-red colour.[1] A little of it dropped into water gives a white precipitate (antimonious oxychloride, SbOCl), and the filtered solution gives a white precipitate (silver chloride) with silver nitrate, indicating the presence of chlorine. If the white precipitate formed by water be treated with sulphuretted hydrogen, it becomes orange-coloured (antimonious sulphide, $Sb_2S_3$).

**Actions and Uses.**—*Internally*:—Irritant and corrosive poison. *Externally*:—Either alone or mixed with four parts of tinctura myrrhæ composita, it is employed as a caustic in the treatment of fistulæ, thrush, canker, luxuriant granulations, foul, and foot-rot. It occasions but little pain.

**Antidotes.**—Chalk; magnesia or its carbonates; and demulcent drinks.

**Preparation.**—Antimonii Oxidum.

## LIQUOR ANTIMONII TARTARATI

### SOLUTION OF TARTARATED ANTIMONY

Take of

| | |
|---|---|
| Tartarated Antimony, in powder | 1 ounce |
| Distilled Water . . . . | 4 to 8 ounces |

Dissolve the tartarated antimony in the water.

[1] The colour is usually darker than here described, owing to the presence of ferric chloride, derived from the iron vessels used in its preparation.

**Use.**—Counter-irritant in deep-seated inflammations and chronic rheumatism of the joints.

## LIQUOR ARGENTI NITRATIS

SOLUTION OF NITRATE OF SILVER

Take of

| | |
|---|---|
| Nitrate of Silver . . . | 10 to 15 grains |
| Distilled Water . . . | 1 fluid ounce |

Dissolve the nitrate of silver in the water.

**Uses.**—As an excitant to wounds, &c. One or two drops of the solution, mixed with an equal bulk of water, are sometimes introduced into the eye to remove opacity of the cornea.

## LIQUOR ARSENICALIS

ARSENICAL SOLUTION

**Synonyms.**—Liquor Potassæ Arsenitis; Fowler's Solution.

**Composition.**—By some authorities it is regarded as arsenious anhydride dissolved in a solution of potassium carbonate; by others as an aqueous solution of potassium arsenite, $KAsO_2$, and carbonate. It contains an amount of arsenical compound corresponding to four grains of arsenious anhydride in one fluid ounce.

**Mode of Preparation.**[1]—Take of

| | |
|---|---|
| Arsenious Anhydride, in powder | of each |
| Carbonate of Potash . . . | $87\frac{1}{2}$ grains |
| Distilled Water . . . . | a sufficiency |

[1] In the B.P. preparation five fluid drachms of Compound Tincture of Lavender are added

Place the arsenious acid and the carbonate of potash in a flask with ten ounces of water, and apply heat until a clear solution is obtained. Allow this to cool, and then add as much distilled water as will make the bulk one pint.

If the former of the above views respecting the composition of this solution be correct, no decomposition takes place during its preparation; but if the latter be adopted, part of the carbonate of potash must be decomposed by the arsenious acid in the manner indicated by the following equation:

| Potassium Carbonate. | | Arsenious Anhydride. | | Potassium Arsenite. | | Carbonic Anhydride. |
|---|---|---|---|---|---|---|
| $K_2CO_3$ | + | $As_2O_3$ | = | $2KAsO_2$ | + | $CO_2$ |

**Characters and Tests.**—Clear, colourless liquid, alkaline to test-paper. Sp. gr. 1·009. After being acidulated with hydrochloric acid it gives, with sulphuretted hydrogen, a yellow precipitate (arsenious sulphide, $As_2S_3$) soluble in ammonium carbonate, which is brightest when the arsenical solution has been previously diluted.

**Actions and Uses.**—*Internally*:—Similar to, but more certain in its action and less liable to produce local irritation than, arsenious acid. *Externally*:—In obstinate skin-affections, and for the destruction of pediculi, acari, and other external parasites.[1]

[1] Several 'sheep-washes' and 'dips' have a composition similar to that of liquor arsenicalis. The simplest is thus prepared:

| | | |
|---|---|---|
| Take of Arsenious Acid, in powder<br>Carbonate of Potash . | } | of each, 8 to 10 ounces |
| Water . . . . . | | 20 gallons |

Boil together for half an hour. This quantity is sufficient for twenty sheep. Sometimes 8½ ounces of soft soap and 6 ounces of flowers of sulphur are added to the foregoing mixture before it is boiled.

Tuson's sheep-dip is a solution of arsenic acid ($H_3AsO_4$) in water mixed with a little paraffin oil.

**Doses.**—Horse, ½ to 1 fluid ounce.—Cattle, ½ to 1 fluid ounce.—Sheep, ½ to 3 drachms.—Dog, 2 to 10 minims.

**Modes of Application.**—*Internally* :—Diluted with sufficient water to make a draught. *Externally* :—Diluted with from two to three times its bulk of water.

**Antidote.**—A mixture of an alkaline carbonate and Ferri Peroxidum Humidum.

## LIQUOR ARSENICI HYDROCHLORICUS

HYDROCHLORIC SOLUTION OF ARSENIC

**Synonym.**—Liquor Arsenici Chloridi.

**Composition.**—A solution of arsenious anhydride (arsenious acid) in dilute hydrochloric acid. One ounce of the solution contains four grains of arsenious anhydride.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Arsenious Anhydride, in powder | 80 grains |
| Hydrochloric Acid . . . | 2 fluid drachms |
| Distilled Water . . . . | a sufficiency |

Boil the arsenious anhydride with the hydrochloric acid and four ounces of the water, then add distilled water to make the bulk up to one pint.

No decomposition takes place in this process; the anhydride is simply dissolved by the dilute acid.

**Characters and Tests.**—Colourless liquid, having an acid reaction. Sp. gr. 1·009. Sulphuretted hydrogen gives at once a bright yellow precipitate (arsenious sulphide) soluble in ammonium carbonate.

**Actions and Uses.**
**Mode of Application.**
**Doses.**
} Same as Liquor Arsenicalis

**Antidote.**—A mixture of an alkaline carbonate and Ferri Peroxidum Humidum.

**Preparation.**—Liquor Arsenici et Hydrargyri Iodidi (the B.P. imitation of Donovan's solution) contains about 1 per cent. by weight of arsenious iodide and of mercuric iodide; it is useful in chronic skin and rheumatic complaints.

**Dose.**—For HORSE and CATTLE, 1 to 2 ounces.

## LIQUOR CALCIS

### SOLUTION OF LIME

**Synonyms.**—Aqua Calcis; Lime-water.

**Composition.**—An aqueous solution of calcium hydrate, $Ca(HO)_2$. Ten fluid ounces of the solution should contain 7·4 grains of calcium hydrate.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Slaked Lime, freshly prepared | 2 ounces |
| Distilled Water | 1 gallon |

Put the lime into a stoppered bottle containing the water, and shake well for two or three minutes. After twelve hours the excess of lime will have subsided, and the clear solution may be drawn off with a siphon as it is required for use, or transferred to a green-glass bottle [1] furnished with a well-ground stopper.

[1] Lime-water should not be kept in white-glass bottles, as it attacks the lead of which they are partly composed.

**Characters and Tests.**—Transparent, colourless, and odourless liquid, having a disagreeable alkaline taste. Feebly alkaline to test-paper. Exposed to the air, a white pellicle or film (calcium carbonate) forms upon its surface,[1] owing to the absorption of carbonic anhydride by the calcium hydrate. With solution of oxalic acid it gives a white precipitate (calcium oxalate), insoluble in acetic acid.

**Actions and Uses.**—Chiefly as an antacid; occasionally as an antidote for poisoning by acids.

**Doses.**—Horse, 4 to 6 fluid ounces.—Cattle, 4 to 6 fluid ounces.—Sheep, ½ to 2 fluid ounces.—Pig, ½ to 2 fluid ounces.—Dog, 1 to 4 fluid drachms.

**Mode of Application.**—Mixed with sufficient water to form a draught.

**Incompatibles.**—Acids and most metallic salts.

**Preparations.**—Linimentum Calcis; Lotio Hydrargyri Flava; Lotio Hydrargyri Nigra.

## LIQUOR CALCIS CHLORINATÆ

### SOLUTION OF CHLORINATED LIME

**Synonyms.**—Solution of Chloride of Lime; Solution of Hypochlorite of Lime.

**Composition.**—Water holding in solution an amount of calcium hypochlorite equivalent to thirteen grains of available chlorine in one ounce of the fluid.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Chlorinated Lime . . . | 1 pound |
| Distilled Water . . . . | 1 gallon |

[1] This reaction takes place much more rapidly if air from the lungs be blown through a tube into the lime-water.

Well mix the water and the chlorinated lime by trituration in a mortar, and, having transferred the mixture to a stoppered bottle, shake it occasionally during three hours ; now pour out the contents of the bottle on a calico filter, and preserve the solution, which passes through, in a stoppered bottle.

**Characters and Tests.**—Transparent, nearly colourless liquid. Sp. gr. 1·035. Exposed to the atmosphere, the calcium hypochlorite is decomposed by carbonic anhydride ; the calcium carbonate produced is deposited as a white precipitate, and the chlorine which is set free may be recognised by its odour.

**Actions and Uses.**—Stimulant, mild erodent, and deodoriser to ill-conditioned and fœtid wounds and fistulous sores. Sprinkled about stables and other buildings, it acts as a deodoriser and disinfectant.

In cases of grease, exfoliating bone, &c., it should be diluted with from ten to fifteen times its bulk of water.

**Preparation.**—Cataplasma Calcis Chlorinatæ.

## LIQUOR CANTHARIDIS TEREBINTHINATUS

TEREBINTHINATED SOLUTION OF CANTHARIDES

Take of

| | |
|---|---|
| Cantharides, in powder . . . | 1 part |
| Oil of Turpentine . . . . | 8 parts |

Digest, at a gentle heat, for three or four days, with frequent agitation, and filter. To every ounce of the filtrate add an equal quantity of Canada balsam, and thoroughly mix them together.

**Use.**—For medicating materials used for setons. The tape, cotton, cord, or other material is immersed in the liniment until it ceases to imbibe any more of the liquid; it is then to be removed and drawn between the finger and thumb to deprive it of any superfluous fluid, and finally hung up to dry.

## LIQUOR CUPRI SULPHATIS

SOLUTION OF SULPHATE OF COPPER

Take of

| | |
|---|---|
| Sulphate of Copper, in powder . | 5 ounces |
| Boiling Distilled Water . . | 1 pint |

Dissolve and filter.

**Uses.**—For injection into sinuses, one part of this solution is mixed with from two to three times its volume of water. See also Cupri Sulphas.

## LIQUOR CUPRI SULPHATIS COMPOSITUS

COMPOUND SOLUTION OF SULPHATE OF COPPER

Take of

| | |
|---|---|
| Sulphate of Copper . . | of each |
| Alum . . . . . | 3 ounces |
| Water . . . . . | 1½ pint |
| Sulphuric Acid . . . | 1½ ounce |

**Uses.**—See Cupri Sulphas.

## LIQUOR FERRI PERCHLORIDI FORTIS

STRONG SOLUTION OF PERCHLORIDE OF IRON

**Composition.** — One fluid drachm contains 35·47 grains of perchloride of iron (ferric chloride, $Fe_2Cl_6$), dissolved in water.

This preparation also contains free hydrochloric acid.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Iron Wire . . . | 4 ounces |
| Hydrochloric Acid . | 20½ fluid ounces |
| Nitric Acid . . . | 1½ fluid drachm |
| Distilled Water . . | a sufficient quantity |

Mix eight fluid ounces of the hydrochloric acid with the distilled water, and in this dissolve the iron at a gentle heat. Filter the solution, add it to the remainder of the hydrochloric acid and the nitric acid, heat the mixture briskly until on the sudden evolution of red fumes the liquid becomes of an orange-brown colour, then evaporate by the heat of a water-bath until it is reduced to ten fluid ounces. The production of perchloride of iron takes place in two stages :

| | Iron. | | Hydrochloric Acid. | | Ferrous Chloride. | | Hydrogen. |
|---|---|---|---|---|---|---|---|
| 1. | Fe | + | 2HCl | = | $FeCl_2$ | + | $H_2$ |

| | Ferrous Chloride. | | Hydrochloric Acid. | | Nitric Acid. | | Ferric Chloride (Perchloride of Iron). | |
|---|---|---|---|---|---|---|---|---|
| 2. | $6FeCl_2$ | + | 6HCl | + | $2HNO_3$ | = | $3Fe_2Cl_6$ | + |

| | Nitric Oxide. | | Water. |
|---|---|---|---|
| | 2NO | + | $4H_2O$ |

**Characters and Tests.** — Orange-brown solution, with a strong astringent and inky taste ; miscible with

water and rectified spirit in all proportions. Sp. gr. 1·338. Diluted with water, it gives a white precipitate (silver chloride) with silver nitrate, showing that it contains chlorine, and a dark blue precipitate (Prussian blue) with potassium ferrocyanide, indicating the presence of a ferric salt.

**Uses.**—In preparing Tinctura Ferri Perchloridi (1 of liquor to 3 of spirit) and Liquor Ferri Perchloridi (1 of liquor to 3 of water).

## LIQUOR FERRI PERSULPHATIS

### SOLUTION OF PERSULPHATE OF IRON

Take of

| | |
|---|---|
| Sulphate of Iron | 8 ounces |
| Sulphuric Acid | ¾ fluid ounce |
| Nitric Acid | ¾ fluid ounce |
| Distilled Water | a sufficient quantity |

Add the sulphuric acid to ten ounces of the water, and dissolve the sulphate of iron in the mixture with the aid of heat. Mix the nitric acid with the remaining two ounces of water, and add the diluted acid to the solution of sulphate of iron. Concentrate the whole by boiling, until, by the sudden disengagement of ruddy vapours, the liquid ceases to be black and acquires a red colour. Test a drop of the solution with potassium ferrocyanide: if a blue precipitate be produced, indicating the presence of some unaltered ferrous salt, a few additional drops of nitric acid are to be added and the boiling renewed in order that the whole may become converted into ferric salt.[1] When the solution is cold, make up the quantity

[1] The boiling, with small additional quantities of nitric acid, must be repeated until a drop of the solution ceases to yield a blue precipitate with potassium ferrocyanide

to eleven fluid ounces by the addition, if necessary, of distilled water.

| Sulphate of Iron (Ferrous Sulphate). | | Sulphuric Acid. | | Oxygen from Nitric Acid. | | Persulphate of Iron (Ferric Sulphate). | | Water. |
|---|---|---|---|---|---|---|---|---|
| $2FeSO_4$ | + | $2H_2SO_4$ | + | $O_2$ | = | $2Fe_2(SO_4)_3$ | + | $2H_2O$ |

**Uses.**—Styptic ; also in the preparation of Ferri Peroxidum Humidum.

## LIQUOR HYDRARGYRI PERCHLORIDI

### SOLUTION OF PERCHLORIDE OF MERCURY

Take of

| | |
|---|---|
| Perchloride of Mercury . . . | 10 grains |
| Distilled Water . . . . | 1 pint |

**Uses.**—*Externally* :—In the treatment of scab and mange, also for the destruction of vermin infesting the skin.[1]

## LIQUOR IODI ET POTASSII IODIDI

### SOLUTION OF IODINE AND IODIDE OF POTASSIUM

**Synonyms.**—Liquor Iodi ; Lugol's Solution.

Take of

| | |
|---|---|
| Iodine . . . . . | 22 grains |
| Iodide of Potassium . . . | 33 grains |
| Distilled Water . . . . | 1 fluid ounce |

Mix and dissolve.

[1] Two minims of hydrocyanic acid added to 2 grains of perchloride of mercury dissolved in 1 fluid ounce of water forms a lotion which is sometimes applied to the skin, especially of dogs, to relieve itching.

**Uses.**—Injected into cavities to cause adhesion of their sides; one fluid drachm diluted with three fluid ounces of water injected into the bladder, to arrest hæmorrhage from that organ; applied with friction to the skin, when depilated, it is said to promote the return of hair. *Internally*:—Antiseptic, and in treatment of bots.

**Doses.**—Horse, 2 to 4 drachms.

## LIQUOR MORPHINÆ ACETATIS

### SOLUTION OF MORPHINE ACETATE

Take of

| | |
|---|---|
| Morphine Acetate . . | 17½ grains |
| Dilute Acetic Acid . . | 38 minims |
| Alcohol (90-per-cent.) . | 1 fluid ounce |
| Distilled Water . . | a sufficient quantity |

Dissolve the acetate in the mixed liquids. Dilute with sufficient distilled water to produce four fluid ounces of solution.[1]

**Actions and Uses.**—Stimulant, sedative, and narcotic.

**Doses.**—Horse, 2 to 6 fluid drachms.—Cattle, 2 to 6 fluid drachms.—Dog, 5 to 15 minims.

**Preparation.**—Injectio Morphinæ Hypodermica, a solution of acetate of morphine containing one grain of the acetate in ten minims of the injection. Use from thirty to sixty minims for the horse in abdominal affections.

[1] One hundred and ten minims contain 1 grain of morphine acetate

## LIQUOR MORPHINÆ HYDROCHLORIDI

SOLUTION OF MORPHINE HYDROCHLORIDE

Take of

| | |
|---|---|
| Morphine Hydrochloride . | 17½ grains |
| Dilute Hydrochloric Acid . | 38 minims |
| Rectified Spirit . . | 1 fluid ounce |
| Distilled Water . . | a sufficient quantity |

Mix the hydrochloric acid, the spirit, and the water, and dissolve the hydrochloride in the mixture. Dilute with sufficient distilled water to produce four fluid ounces of the solution.

**Uses.**—As a sedative, soporific, antispasmodic, &c.

**Doses.**—See Liquor Morphinæ Acetatis.

## LIQUOR PLUMBI SUBACETATIS

SOLUTION OF SUBACETATE OF LEAD

**Synonym.**—Goulard's Extract.

**Composition.**—Not accurately known; probably a mixture of two or more basic acetates of lead.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Acetate of Lead . . | 5 ounces |
| Oxide of Lead (Litharge), in powder . . . | 3½ ounces |
| Distilled Water . . | 1 pint, or a sufficiency |

Boil the acetate and oxide of lead in the water for half an hour, constantly stirring; then filter, and when the liquid is cold add to it more distilled water until the

product measures twenty fluid ounces. Keep the clear solution in well-stoppered bottles, to prevent its being decomposed by atmospheric carbonic anhydride. In this process the acetate of lead unites with the oxide of lead and is transformed into the mixture of basic salts above referred to.

**Characters and Tests.**—Heavy, clear, colourless liquid. Sp. gr. 1·26. Alkaline reaction, sweet astringent taste. Exposed to the air it becomes turbid,[1] from the absorption of carbonic anhydride and consequent formation of lead carbonate. The addition of strong sulphuric acid throws down a white precipitate (lead sulphate), and liberates the vapour of acetic acid, which can be identified by its odour. The former reaction indicates the presence of lead ; the latter that the salt is an acetate.

**Use.**—In preparing Linimentum Plumbi Subacetatis.

## LIQUOR PLUMBI SUBACETATIS DILUTUS

### DILUTE SOLUTION OF SUBACETATE OF LEAD

**Synonym.**—Goulard's Water.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Solution of Subacetate of Lead | of each 2 fluid drachms |
| Rectified Spirit . . . | |
| Distilled Water [2] . . . | 19½ fluid ounces |

Mix, and filter through paper. Keep the clear solution in a stoppered bottle.

[1] Air from the lungs blown through a tube into a solution produces the turbidity much more rapidly. Compare with Plumbi Acetas.

[2] Common water should never be employed for diluting subacetate of lead, as it precipitates the lead as sulphate and carbonate, and thus destroys or diminishes the activity of the preparation.

**Actions and Uses.** — *Externally*:—Astringent and sedative. Applied, by means of rags wetted with it, to the skin to allay pain and irritation, and to subdue superficial inflammation. See also Linimentum Plumbi Subacetatis. As a collyrium in conjunctival ophthalmia, one part of this preparation, diluted with an equal part of water, is to be used.

## LIQUOR PLUMBI SUBACETATIS ET OPII

SOLUTION OF SUBACETATE OF LEAD AND OPIUM

Take of

| | |
|---|---|
| Solution of Subacetate of Lead | 1 fluid drachm |
| Tincture of Opium . . | 1 fluid drachm |
| Distilled Water . . . | 6 fluid ounces |

Mix.

**Uses.**—Anodyne to wounds, to blistered and cauterised surfaces, and in superficial inflammation of the eye, skin, or joints.

## LIQUOR POTASSÆ

SOLUTION OF POTASH

**Composition.**—Twenty-seven grains of potassium hydroxide, KOH, in one fluid ounce of water.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Carbonate of Potash . . . | 1 pound |
| Slaked Lime . . . . | 12 ounces |
| Distilled Water . . . . | 1 gallon |

Dissolve the carbonate of potash in the water, and, having heated the solution to the boiling-point in a clean

iron vessel, gradually mix with it the slaked lime, and continue the ebullition for ten minutes with constant stirring. Then remove the vessel from the fire; and when, by the subsidence of the insoluble matter (calcium carbonate), the supernatant liquor (solution of potassium hydrate) has become perfectly clear, transfer it by means of a siphon to a green-glass bottle [1] furnished with an air-tight stopper, and add distilled water, if necessary, to make it correspond with the test of specific gravity and neutralising power.[2]

| Potassium Carbonate. | | Calcium Hydrate. | | Potassium Hydrate (Caustic Potash). | | Calcium Carbonate. |
|---|---|---|---|---|---|---|
| $K_2CO_3$ | + | $Ca(HO)_2$ | = | $2KHO$ | + | $CaCO_3$ |

**Characters and Tests.** — Transparent, colourless, and odourless liquid. Sp. gr. 1·058. Alkaline to test-paper; acrid taste; feels soapy when rubbed between the finger and thumb. Mixed with excess of solution of tartaric acid, and stirred, it yields a white granular precipitate (hydrogen-potassium tartrate), showing the presence of potassium. Added to excess of dilute hydrochloric acid, no effervescence should be occasioned, indicating its freedom from carbonate. Treated with slight excess of nitric acid, and evaporated to dryness, the residue dissolves in water and yields a nearly clear solution, which should give no precipitate, or a very slight one, when separate portions of it are mixed with silver nitrate, barium chloride, and solution of ammonia; these reactions prove the absence or nearly so, of chlorides, sulphates, and iron respectively.

**Use.**—In preparing Potassa Caustica.

[1] As it slowly acts upon and dissolves the lead of white bottles.

[2] See the 'British Pharmacopœia' published in 1898, if it be required to make this test.

## LIQUOR POTASSII NITRATIS

SOLUTION OF NITRATE OF POTASH

Take of

| | |
|---|---|
| Nitrate of Potash, in powder . | 1 ounce |
| Water . . . . . | 7 fluid ounces |

Dissolve, and apply immediately after the solution of the salt has been effected.

**Uses.**—Stimulant to gangrenous and other wounds, also to superficial inflammations and slight sprains.

## LIQUOR POTASSII PERMANGANATIS

SOLUTION OF PERMANGANATE OF POTASH [1]

Take of

| | |
|---|---|
| Permanganate of Potash . . | 87½ grains |
| Water . . . . . | 1 pint |

Dissolve.

**Use.**—Cleansing wash for diseased surfaces.

## LIQUOR POTASSÆ SULPHURATÆ

SOLUTION OF SULPHURATED POTASH

Take of

| | |
|---|---|
| Sulphurated Potash . . . | 1 ounce |
| Water . . . . . | 8 fluid ounces |

Dissolve.

**Use.**—Applied to the skin in scab, mange, and other skin-affections.

[1] 110 minims contain 1 grain of Potassium Permanganate.

## LIQUOR SODÆ

SOLUTION OF SODA

**Composition.** — 18·8 grains of sodium hydrate, NaOH, in every fluid ounce of water.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Carbonate of Soda . . . | 28 ounces |
| Slaked Lime . . . . | 12 ounces |
| Distilled Water . . . | 1 gallon |

Dissolve the carbonate of soda in the water, and, having heated the solution to the boiling-point in a clean iron vessel, gradually mix with it the slaked lime, and continue the ebullition for ten minutes with constant stirring. Then remove the vessel from the fire, and when, by the subsidence of the insoluble matter (calcium carbonate), the supernatant liquid (solution of sodium hydrate) has become perfectly clear, transfer it by means of a siphon to a green-glass [1] bottle furnished with an air-tight stopper, and add distilled water, if necessary, to make it correspond with the tests of specific gravity and neutralising power.[2]

| Sodium Carbonate. | | Calcium Hydrate. | | Sodium Hydrate (Caustic Soda). | | Calcium Carbonate. |
|---|---|---|---|---|---|---|
| $Na_2CO_3$ | + | $Ca(HO)_2$ | = | $2NaOH$ | + | $CaCO_3$ |

**Characters and Tests.** — Transparent, colourless, and odourless liquid. Sp. gr. 1·047. Alkaline to test-paper; acrid taste; feels soapy when rubbed between the

[1] As it slowly acts upon and dissolves the lead of white ones.

[2] See the 'British Pharmacopœia,' 1867, if it be desired to apply this test.

finger and thumb. Mixed with excess of solution of tartaric acid, and stirred, no precipitate is thrown down: thus it is distinguished from Liquor Potassæ. If a platinum wire, moistened with the solution, be heated before the blowpipe, a yellow colour is imparted to the flame, indicating the presence of sodium. It should not effervesce when added to an excess of diluted hydrochloric acid, proving the absence of carbonate. Chlorides, sulphates, and iron are detected in the manner described under Liquor Potassæ.

**Preparation.**—Soda Caustica.

## LIQUOR STRYCHNINÆ HYDROCHLORIDI

### SOLUTION OF STRYCHNINE HYDROCHLORIDE [1]

Take of

| | | |
|---|---|---|
| Strychnine Hydrochloride | . | 17½ grains |
| Alcohol (90-per-cent.) | . . | 5 fluid drachms |
| Distilled Water to | . . . | 4 fluid ounces |

Dissolve the strychnine hydrochloride in the alcohol mixed with sufficient distilled water to produce four fluid ounces of solution.

**Actions and Uses.**—Stimulant and tonic.

**Doses.** — Horse, 2 to 6 fluid drachms.—Cattle, 2 to 6 fluid drachms.—Sheep, ½ to 2 fluid drachms.—Dog, 3 to 10 minims.[2]

[1] 110 minims contain 1 grain of Strychnine Hydrochloride.

[2] Liquor Strychninæ Hydrochloridi is the safest form of administering strychnine to dogs.—Ed.

## LIQUOR ZINCI CHLORIDI

SOLUTION OF CHLORIDE OF ZINC

**Synonym.**—Butter of Zinc.

**Composition.**—Prepared in the manner described in the 'British Pharmacopœia,' 1898, one fluid ounce contains 366 grains of zinc chloride, $ZnCl_2$. Sir William Burnett's Disinfecting-fluid is a very similar preparation, and is the form of chloride of zinc most commonly employed in veterinary practice; one fluid ounce of it contains 200 grains of zinc chloride. A solution of 'British Pharmacopœia' strength, but containing small quantities of iron and other impurities, may be obtained by the following

**Mode of Preparation.**—Take of

| | |
|---|---|
| Granulated Zinc . . . | 1 pound |
| Hydrochloric Acid . . | 44 fluid ounces |
| Distilled Water . . . | 1 pint |

Mix the hydrochloric acid and water in a porcelain dish, add the zinc, and apply a gentle heat to promote the action until gas (hydrogen) is no longer evolved. Boil for half an hour, supplying the water lost by evaporation, and allow the product to cool. Filter the liquid (zinc chloride) into a porcelain basin, and evaporate until it is reduced to the bulk of two pints.

| Zinc. | | Hydrochloric Acid. | | Zinc Chloride (Chloride of Zinc). | | Hydrogen. |
|---|---|---|---|---|---|---|
| Zn | + | 2HCl | = | $ZnCl_2$ | + | $H_2$ |

**Characters and Tests.**—Heavy, oily, colourless liquid, having a powerfully astringent and metallic taste. Diluted with water, it will give no precipitate with sulphuretted hydrogen unless lead be present, when a black precipitate or brown coloration (lead sulphide) is produced. If solution of ammonia be slowly added to the preparation diluted with water, a white, or, if iron be present, a brownish-white, precipitate (zinc hydrate) will be thrown down, which dissolves in excess of the precipitant. On saturating this ammoniacal solution, after filtration, if necessary, with sulphuretted hydrogen, a white precipitate (zinc sulphide) is produced, which readily dissolves in dilute hydrochloric acid; this reaction is indicative of the presence of zinc. Another portion of the diluted preparation, acidified with nitric acid, and treated with silver nitrate, furnishes a white precipitate (silver chloride), showing that the salt is a chloride.

**Actions and Uses.**—Chiefly employed as a deodoriser and disinfectant; for this purpose one fluid ounce of chloride of zinc may be diluted with one gallon of water and distributed about the apartment in shallow plates, or it may be sprinkled over the floor, &c.[1] Fifty minims of Burnett's Fluid, diluted with one pint of water, forms a lotion which may be applied, by means of a sponge saturated with it, to open joints.

In the concentrated state it is a powerful astringent and caustic, used in the treatment of fetid ulcers and wounds, fistulæ, luxuriant growths, &c. See also Zinci Chloridum.

[1] Tuson's Liquid Disinfectant consists of a solution of zinc chloride saturated with sulphurous anhydride.

## LIQUOR ZINCI SULPHATIS

SOLUTION OF ZINC SULPHATE

Take of

| | |
|---|---|
| Sulphate of Zinc . . . . | 5 ounces |
| Distilled Water . . . . | 20 ounces |

Mix the sulphate of zinc with the water, and dissolve by the aid of heat.

**Actions and Uses.** — Similar to Liquor Zinci Chloridi.

## LOTIO ACIDI CARBOLICI

LOTION OF CARBOLIC ACID

Take of

| | |
|---|---|
| Carbolic Acid . . . | 1 fluid drachm |
| Water . . . . | 6 to 12 fluid ounces |

Mix and well shake.

**Uses.**—Dressing for unhealthy sores, indolent ulcers, foot-rot in sheep, the feet of cattle suffering from foul; in the treatment of eczema, grease, mange, and scab; also to allay itching in skin-affections.[1]

## LOTIO AMMONII ACETATIS

LOTION OF ACETATE OF AMMONIA

Take of

| | |
|---|---|
| Solution of Acetate of Ammonia | of each |
| Rectified Spirit . . . . | 4 fluid ounces |
| Water . . . . . | 1 pint |

Mix.

[1] One drachm of carbolic acid shaken with a quart of water forms an efficient wash for the mouth, feet, and udders of animals suffering from aphthous epizootic.

**Use.**—Discutient. The part is to be kept constantly moist with the lotion.

## LOTIO AMMONII CHLORIDI ET CAMPHORÆ

### LOTION OF CHLORIDE OF AMMONIUM AND CAMPHOR

**Synonyms.**—Lotio Discutiens ; Discutient Lotion.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Chloride of Ammonium, in powder | 1 ounce |
| Dilute Acetic Acid . . . | 8 fluid ounces |
| Spirit of Camphor . . . | 1 fluid ounce |

Dissolve the chloride of ammonium in the dilute acetic acid, and add the spirit of camphor.

**Use.**—Discutient to indolent tumours and chronic sprains.

It should be well shaken before use, and applied with friction.

## LOTIO AMMONII CHLORIDI ET POTASSII NITRATIS

### LOTION OF CHLORIDE OF AMMONIUM AND NITRATE OF POTASH

**Synonyms.**—Lotio Refrigerans ; Cooling Lotion.

Take of

| | |
|---|---|
| Chloride of Ammonium, in powder | of each |
| Nitrate of Potash, in powder . . | 2½ ounces |
| Water . . . . . | 1 pint |

Add the chloride of ammonium and nitrate of potash to the water.

**Use.**—Directly the salts are dissolved, dip cloths in the solution and lay them on the inflamed part.

## LOTIO HYDRARGYRI FLAVA

YELLOW MERCURIAL LOTION

**Synonym.**—Yellow Wash.

Take of

| | |
|---|---|
| Perchloride of Mercury . . | 18 grains |
| Solution of Lime . . . | 10 fluid ounces |

Mix.

**Use.**—Stimulant to unhealthy sores and ulcers.

## LOTIO HYDRARGYRI NIGRA

BLACK MERCURIAL LOTION

**Synonym.**—Black Wash.

Take of

| | |
|---|---|
| Subchloride of Mercury . . | 30 grains |
| Solution of Lime . . . | 13 fluid ounces |

Mix.

**Use.**—Stimulant to unhealthy sores and ulcers.

## MAGNESIA[1]

MAGNESIA

**Synonyms.**—Magnesium Oxide ; Magnesia Usta.

**Composition.**—MgO.

[1] Magnesia Levis (Light Magnesia) is a bulky white powder, possessing the same chemical and therapeutic properties as this preparation.

**Mode of Preparation.**—Take of

Carbonate of Magnesia[1] . . . 4 ounces

Put it into a Cornish or Hessian crucible closed loosely by a lid, and expose it to a low red heat until a small quantity taken from the centre of the crucible, cooled, and dropped into dilute sulphuric acid, causes no effervescence, showing that it contains no carbonate.

| Official Carbonate of Magnesia. | | Magnesia. | | Carbonic Anhydride. | | Water. |
|---|---|---|---|---|---|---|
| $3MgCO_3.Mg(HO)_2.4H_2O$ | = | $4MgO$ | + | $3CO_2$ | + | $5H_2O$ |

**Characters.**—White powder, very slightly soluble in water, but readily dissolving in acids without effervescence. Its solution in hydrochloric acid, when neutralised by a mixed solution of ammonium chloride and ammonia, gives a copious crystalline precipitate (ammonio-magnesium phosphate) when hydrogen-di-sodium phosphate is added to it. Dissolved in nitric acid, it should give no precipitate with barium chloride, showing the absence of sulphate, and when neutralised with a mixture of ammonium chloride and ammonia, it should yield no precipitate with ammonium oxalate, indicating its freedom from calcium.

**Actions and Uses.**—Antacid and laxative. Only given by the veterinarian to foals or calves suffering from indigestion.

**Doses.**—From 2 to 8 drachms, combined with a carminative, are administered, suspended in milk or gruel.

[1] Really a mixture of carbonate and hydrate. See Magnesii Carbonas.

## MAGNESII CARBONAS PONDEROSA[1]

HEAVY MAGNESIUM CARBONATE

**Synonym.**—Magnesia Alba.

**Composition.**—A mixture of magnesium carbonate and hydrate, to which the formula $3MgCO_3.Mg(HO)_2.4H_2O$ has been assigned.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Magnesium Sulphate . . | 10 ounces |
| Carbonate of Soda . . . | 12 ounces |
| Boiling Distilled Water . . | a sufficiency |

Dissolve the magnesium sulphate and the carbonate of soda each in a pint of the water, mix the two solutions, and evaporate the whole to perfect dryness by means of a sand-bath. Digest the residue for half an hour with two pints of the water, and having collected the insoluble matter (official carbonate of magnesia) on a calico filter, wash it repeatedly with distilled water (to remove sodium sulphate) until the washings cease to give a precipitate with barium chloride. Finally, dry the product at a temperature not exceeding 212° F.

| Magnesium Sulphate. | | Sodium Carbonate. | | Water. |
|---|---|---|---|---|
| $4MgSO_4$ | + | $4Na_2CO_3$ | + | $5H_2O$ |

| | Official Carbonate of Magnesia. | | Sodium Sulphate. | | Carbonic Anhydride. |
|---|---|---|---|---|---|
| = | $3MgCO_3.Mg(HO)_2.4H_2O$ | + | $4Na_2SO_4$ | + | $CO_2$ |

**Characters and Tests.**—White granular powder, which dissolves with effervescence in dilute nitric acid, showing that the preparation contains a carbonate, and

[1] Magnesii Carbonas Levis (Light Carbonate of Magnesia) is a bulky white powder, possessing the same chemical and therapeutic properties as this preparation

yielding a solution which, when treated with ammonium chloride, ammonia, and hydrogen-di-sodium phosphate, gives a copious white crystalline precipitate (ammonio-magnesium phosphate), showing the presence of magnesium. With slight excess of hydrochloric acid it forms a clear solution, in which barium chloride will cause no precipitate if sulphates be absent. Another portion of the solution supersaturated with ammonia gives no precipitate with ammonium oxalate or sulphuretted hydrogen, unless calcium and iron are respectively present.

**Actions and Uses.**
**Doses.** } Same as Magnesia.

## MAGNESII SULPHAS

### MAGNESIUM SULPHATE

**Synonyms.**—Sulphate of Magnesia ; Epsom Salts.

**Composition.**—$MgSO_4.7H_2O$.

**Mode of Preparation.** — Various methods are adopted, but the simplest consists in dissolving *magnesite* (native magnesium carbonate, $MgCO_3$) in dilute sulphuric acid, heating the solution formed (magnesium sulphate) until carbonic anhydride ceases to be evolved, filtering, and evaporating the filtrate, so that the resulting magnesium sulphate may crystallise out on cooling and standing.

| Magnesium Carbonate. | | Sulphuric Acid. | | Magnesium Sulphate (Sulphate of Magnesia). | | Water. | | Carbonic Anhydride. |
|---|---|---|---|---|---|---|---|---|
| $MgCO_3$ | + | $H_2SO_4$ | = | $MgSO_4$ | + | $H_2O$ | + | $CO_2$ |

In assuming the solid state the salt acquires seven molecules of water of crystallisation.

**Characters and Tests.** — Very small, colourless, transparent, rhombic prisms, possessing a bitter taste. Readily dissolves in water, and the solution gives a copious white crystalline precipitate (ammonio-magnesium phosphate) with ammonium chloride, ammonia, and hydrogen-di-sodium phosphate; and a white precipitate, insoluble in nitric acid, with barium chloride, showing that the salt is a sulphate. Should give no blue precipitate with potassium ferrocyanide, indicating its freedom from iron.

**Actions and Uses.**—Cathartic, diuretic, and febrifuge. Chiefly employed, combined with ginger, as a purgative for cattle and sheep. Used also as a febrifuge for the horse, and as an antidote to poisoning by lead.

**Doses.** — *Cathartic*: — CATTLE, 1 to 1½ pound.— SHEEP, 1 to 2 ounces.—DOG, 1 to 4 drachms. *Febrifuge*:—HORSE, 2 to 4 ounces, dissolved in water, twice a day.

**Mode of Application.**—Dissolved in from ten to twenty times its weight of water. Sometimes, as in cases of obstinate constipation and torpidity of the bowels in cattle, a drachm of calomel, or ten to fifteen croton beans, are added to the ordinary dose of the salt, to increase its cathartic activity.

**Incompatibles.**—Alkaline carbonates; lime-water; acetate of lead.

## MANGANESII OXIDUM NIGRUM

BLACK OXIDE OF MANGANESE

**Synonyms.**—Manganic Peroxide; Manganese Dioxide; Peroxide of Manganese; Binoxide of Manganese.

**Composition.**—$MnO_2$. Found native.

**Characters and Tests.** — Heavy black powder, which dissolves almost entirely in hydrochloric acid, with evolution of chlorine, and gives off oxygen when heated to bright redness.

**Use.**—Chiefly for the production of chlorine, and in the preparation of Potassium Permanganate.

## MARMOR ALBUM

WHITE MARBLE

**Composition.**—Naturally crystallised calcium carbonate, $CaCO_3$.

**Characters and Tests.**—Heavy, white, crystalline solid. Dissolves almost completely in hydrochloric acid, with evolution of carbonic anhydride.

**Use.** — Chiefly in the production of carbonic anhydride.

## MASSA ALOES

MASS OF ALOES

**Synonym.**—Cathartic Mass.

Take of

| | |
|---|---|
| Barbados Aloes, in small pieces | 8 ounces |
| Glycerin | 2 ounces |
| Ginger,[1] in powder | 1 ounce |
| Rape Oil | 1 ounce |

Melt together in a water-bath, and thoroughly incorporate by frequent stirring.

[1] In cases in which the use of ginger may be considered objectionable substitute an equal weight of powdered gentian.

**Use.**—Cathartic for the horse.

**Dose.**—From 6 to 8 drachms.

## MASSA ALOES COMPOSITA

COMPOUND MASS OF ALOES

**Synonym.**—Alterative Mass.

Take of

| | |
|---|---|
| Barbados Aloes, in powder . . | 1 ounce |
| Soft Soap . . . . . | 1 ounce |
| Common Mass . . . . | 6 ounces |

Thoroughly incorporate by beating in a mortar, so as to form a mass.

**Use.**—Alterative for the horse.

**Dose.**—1 ounce.

## MASSA ANTIMONII TARTARATI COMPOSITA

COMPOUND MASS OF TARTARATED ANTIMONY

**Synonym.**—Fever-ball.

Take of

| | |
|---|---|
| Tartarated Antimony, in powder | ½ drachm |
| Camphor, in powder . . . | ½ drachm |
| Nitrate of Potash, in powder | 2 drachms |
| Common Mass . . . . | a sufficiency |

Mix so as to form a bolus.

**Use.**—Febrifuge for the horse.

**Dose.**—The above mixture constitutes one dose.

## MASSA BELLADONNÆ COMPOSITA

### COMPOUND MASS OF BELLADONNA

**Synonym.**—Cough-ball.

Take of

| | |
|---|---|
| Extract of Belladonna . | . ½ to 1 drachm |
| Barbados Aloes, in powder | . 1 drachm |
| Nitrate of Potash, in powder | . 2 drachms |
| Common Mass . . | . a sufficiency |

Mix so as to form a bolus.

**Use.**—For the horse in chronic cough.

**Dose.**—The above mixture constitutes one dose.

## MASSA CATECHU COMPOSITA

### COMPOUND MASS OF CATECHU

**Synonym.**—Astringent Mass.

Take of

| | |
|---|---|
| Extract of Catechu, in fine powder | . 1 ounce |
| Cinnamon-bark, in fine powder | . 1 ounce |
| Common Mass . . . . | . 6 ounces |

Mix.

**Use.**—Astringent for the horse.

**Dose.**—1 ounce, in the form of bolus.

## MASSA COMMUNIS

### COMMON MASS

Take of

| | |
|---|---|
| Linseed, finely ground . | } equal parts |
| Treacle . . . . | } |

Mix together so as to form a mass.

**Use.**—An excipient for medicinal agents when they are to be administered in the form of bolus.

## MASSA CUPRI SULPHATIS

### MASS OF SULPHATE OF COPPER

**Synonym.**—Tonic Mass.

Take of

| | |
|---|---|
| Sulphate of Copper, finely powdered | 1 ounce |
| Ginger, in powder | 1 ounce |
| Common Mass | 6 ounces |

Mix.

**Use.**—Tonic for the horse.

**Dose.**—6 to 8 drachms.

## MASSA DIGITALIS COMPOSITA

### COMPOUND MASS OF DIGITALIS

**Synonym.**—Cough-ball.

Take of

| | |
|---|---|
| Barbados Aloes, in powder | 2 ounces |
| Digitalis | 1 ounce |
| Common Mass | 13 ounces |

Mix.

**Use.**—For the horse in chronic cough.

**Dose.**—1 ounce, once or twice a day.

## MASSA FERRI SULPHATIS

### MASS OF SULPHATE OF IRON

**Synonym.**—Tonic Mass.

Take of

| | |
|---|---|
| Sulphate of Iron, in powder . . | 2 ounces |
| Ginger, in powder . . . . | 1 ounce |
| Common Mass . . . . | 5 ounces |

Mix.

**Use.**—Tonic for the horse.

**Dose.**—6 to 8 drachms.

## MASSA RESINÆ COMPOSITA

### COMPOUND MASS OF RESIN

**Synonym.**—Diuretic Mass.

Take of

| | |
|---|---|
| Resin, in powder . . . | equal parts |
| Nitrate of Potash, in powder . | |
| Hard Soap . . . . | |

Mix.

**Use.**—Diuretic for the horse.

**Dose.**—1 ounce.

## MASSA ZINGIBERIS COMPOSITA

### COMPOUND MASS OF GINGER

**Synonym.**—Cordial Mass.

Take of

| | |
|---|---|
| Ginger, in powder . . | } equal parts |
| Gentian-root, in powder . | |
| Treacle . . . . | a sufficiency |

Mix so as to form a mass.

**Use.**—Stomachic for the horse.

**Dose.**—1 ounce.

## MISTURA AMMONIÆ

### AMMONIA MIXTURE

Take of

| | |
|---|---|
| Solution of Ammonia . . | } equal parts |
| Spirit of Nitrous Ether . | |
| Compound Tincture of Gentian | |

**Actions and Uses.**—Stimulant and antispasmodic.

**Doses.**—Horse, 1 to 2 fluid ounces.—Cattle, 2 to 4 fluid ounces.—Sheep, 4 to 6 fluid drachms.

## MISTURA CREOSOTI

### CREOSOTE MIXTURE

Take of

| | |
|---|---|
| Creosote | 16 minims |
| Acetic Acid | 16 minims |
| Spirit of Juniper | ½ drachm |
| Syrup | 1 ounce |
| Distilled Water | 15 ounces |

Mix.

**Actions and Uses.**—Astringent, styptic, and antiseptic.

**Doses.**—Horse, 7 to 10 ounces.—Dog, ½ to 1 ounce.

## MISTURA CRETÆ COMPOSITA

### COMPOUND CHALK MIXTURE

**Synonym.**—Sheep and Calves' Cordial.

Take of

| | |
|---|---|
| Prepared Chalk | 2 ounces |
| Catechu, in powder | 1 ounce |
| Ginger | ½ ounce |
| Opium | 1 drachm |
| Peppermint-water | 1 pint |

**Use.**—To check diarrhœa in calves and sheep.

**Doses.**—Calf, ½ to 1 fluid ounce.—Sheep, 4 to 8 fluid drachms.

## MISTURA RICINI

### CASTOR-OIL MIXTURE

Take of

| | |
|---|---|
| Castor Oil . . . . . | 3 ounces |
| Syrup of Buckthorn . . . | 2 ounces |
| Syrup of Poppies . . . | 1 ounce |

Mix.

**Use.**—Cathartic for the dog.

**Dose.**—4 to 8 fluid drachms.

## MORPHINÆ ACETAS

### MORPHINE ACETATE

**Composition.**—$C_{17}H_{19}NO_3C_2H_4O_2.3H_2O$.

**Preparation.**—Acetate of Morphine is prepared by decomposing a solution of hydrochloride of morphine by ammonia solution, adding dilute acetic acid, to the precipitated morphine, and drying at a gentle heat.

**Characters and Tests.**—It closely resembles the alkaloid, from which it is distinguished by the evolution of an acetous odour on the addition of sulphuric acid.

**Actions and Uses.**
**Doses.**
**Modes of Application.**
} See Morphinæ Hydrochloridum.

**Preparation.**—Liquor Morphinæ Acetatis.

## MORPHINÆ HYDROCHLORIDUM

MORPHINE HYDROCHLORIDE

**Synonyms.**—Morphinæ Murias ; Muriate of Morphia.

**Composition.**—$C_{17}H_{19}NO_3.HCl.3H_2O$.

**Mode of Preparation.**—The carefully dried salt, obtained by neutralising morphine with hydrochloric acid.

**Characters and Tests.**—White, flexible, acicular prisms of a silky lustre, not changed by exposure to the air, soluble in water and alcohol. The aqueous solution gives a white curdy precipitate with nitrate of silver, and a white one with potash, which is redissolved when an excess of the alkali is added. Moistened with strong nitric acid it becomes orange-red, and with solution of perchloride of iron greenish-blue. Entirely destructible by heat, leaving no residue. Twenty grains of the salt dissolved in half an ounce of warm water, with ammonia added in the slightest possible excess, give on cooling a crystalline precipitate which, when washed with a little cold water and dried by exposure to the air, weighs fifteen to eighteen grains.

**Actions and Uses.**—*Internally* :—Anodyne, sedative, soporific, and antispasmodic. *Externally* :—To relieve neuralgia, tetanus, acute rheumatism, and enteritis in horses.

**Doses.**[1]—Horse, 3 to 10 grains.—Cattle, 3 to 10 grains.—Sheep, ½ to 2 grains.—Pig, ½ to 2 grains.—Dog, ⅛ to ½ grain.

[1] For Liquor Morphinæ Hydrochloridi see p. 176.

**Modes of Application.**—*Internally*:—In the form of bolus, or dissolved in water acidulated with a little hydrochloric acid. *Externally*:—By hypodermic injection. See Liquor Morphinæ Acetatis.

**Antidotes.**—See Opium.

## MORPHINA

### MORPHINE

An alkaloid, $C_{17}H_{19}NO_3$, obtained from opium. Nearly insoluble in cold water; readily soluble in hydrochloric or acetic acid. Used in the form of acetate or hydrochloride of morphia.

## APOMORPHINA

### APOMORPHINE

**Composition.**—$C_{17}H_{17}NO_2$.

**Mode of Preparation.**—By heating morphine for some hours with an excess of hydrochloric acid, and precipitating the hydrochloride[1] thus formed with sodium carbonate.

**Characters and Tests.**—An amorphous white powder, which rapidly turns green in air, and dissolves in ether with a pink colour.

**Actions and Uses.**—A powerful emetic.

**Dose.**—Dogs, $\frac{1}{10}$ to $\frac{1}{8}$ grain.

[1] A solution of the hydrochloride (1 grain in 110 minims) is usually employed. Dose.—Dogs: Hypodermically, 5 to 10 minims; by the mouth, 1 grain of the solid salt.

## MUCILAGO ACACIÆ

### MUCILAGE OF GUM ACACIA

Take of

| | |
|---|---|
| Gum Acacia,[1] in small pieces | 13 ounces |
| Distilled Water | 1 pint |

Put the gum and water into a covered earthenware jar and stir them frequently until the gum is dissolved. If necessary, strain the solution through muslin.

**Actions and Uses.**—Demulcent and emollient to allay irritation of the alimentary canal, whether caused by disease or poison; also injected into the bladder in inflammation of that organ and of the kidneys.

**Dose.**—*Ad libitum.*

## MUCILAGO AMYLI[2]

### MUCILAGE OF STARCH

Take of

| | |
|---|---|
| Starch, in powder | ½ ounce |
| Water | 1 pint |

Triturate the starch with the water, gradually added; then boil for a few minutes, constantly stirring.

**Use.**—In preparing enemas.

[1] The much cheaper and equally efficacious dextrin, or British gum, may be substituted for gum acacia in this preparation.

[2] For most veterinary purposes ordinary gruel may be used instead of this preparation.

## MYLABRIS

MYLABRIS

**Synonyms.**—Mylabris Cichorii; Chinese Blistering-fly.

**Natural Order.**—*Coleoptera.*

**Composition.**—Its vesicant properties are due to the presence of cantharidin.

**Characters.**—An insect, found on the flowers of the succory-plant in India and China. About an inch and a quarter in length; sheath-wings black, each presenting anteriorly two almost quadrate brownish-yellow spots; behind these two brownish-yellow bands, each of which equals about one-sixth of the length of the sheath-wings.

**Actions and Uses.**—Its physiological actions are the same as those of cantharides, except that it is said not to affect the kidneys when topically applied. Employed in the form of Unguentum Mylabridis as a counter-irritant and vesicant.

**Preparation.**—Unguentum Mylabridis.

## MYRRHA

MYRRH

A gum-resinous exudation from the stem of *Balsamodendron Myrrha.* Collected in Arabia Felix and Abyssinia.

**Natural Order.**—*Amyridaceæ* or *Burseraceæ.*

**Composition.**—Its medicinal activity depends upon the resin and volatile oil which it contains; they are both

perfectly soluble in rectified spirit, but only partially so in proof spirit.

**Characters.**—In irregularly shaped tears or masses, varying much in size, slightly translucent, of a reddish-yellow or reddish-brown colour; fractured surface irregular, and somewhat oily; odour agreeable and aromatic; taste acrid and bitter.

**Actions and Uses.**—*Internally*:—Stimulant and tonic. It is sometimes given in dyspepsia depending upon debility, but recommended chiefly for cattle in cases of chronic cough; for the latter purpose it is usually combined with opium. *Externally*:—As an excitant and deodoriser to wounds and indolent and fœtid ulcers.

**Doses.**—Horse, 2 to 4 drachms.—Cattle, ½ to 1 ounce.—Sheep, 1 to 2 drachms.—Pig, 1 to 2 drachms. Dog, 10 to 30 grains.

**Modes of Application.**—*Internally*:—In the form of bolus; pill; Tinctura Myrrhæ; Tincture Aloës Composita.—*Externally*:—Powder; Tinctura Myrrhæ; Tinctura Aloës Composita.

**Preparations.**—Tinctura Aloës Composita; Tinctura Myrrhæ.

## NUX VOMICA

### NUX VOMICA

The seeds of *Strychnos Nux-Vomica*. Imported from the East Indies.

**Natural Order.**—*Loganiaceæ*.

**Composition.**—Contains the highly poisonous alkaloids strychnine ($C_{21}H_{22}N_2O_2$) and igasurin, and an almost inert alkaloid, brucine, in combination with igasuric acid.

**Characters.**—Nearly circular and flat, about an inch in diameter, umbilicated and slightly convex on one side; externally of an ash-grey colour, thickly covered with short satiny hairs; internally translucent, tough, and horny; taste intensely bitter; inodorous.

**Actions and Uses.**—In excessive doses, highly poisonous; in medicinal doses, it is given as a nervous stimulant in general paralysis, paraplegia, and amaurosis; also as a tonic in chorea and affections of the alimentary canal—*e.g.*, dyspepsia, dysentery, colic arising from lead-poisoning, &c.

**Doses.**—Horse, 30 to 60 grains.—Cattle, 60 to 120 grains.—Sheep, 5 to 15 grains.—Pig, 5 to 15 grains. Dog, ½ to 3 grains.[1]

**Modes of Application.**—The powdered seed made into a bolus or pill; Extractum Nucis Vomicæ; Tinctura Nucis Vomicæ.

**Antidote.**—Decoctum Tabaci.

**Preparations.**—Strychnina; Tinctura Nucis Vomicæ.

## OLEUM ANISI

### OIL OF ANISE

**Composition.**—A mixture of two distinct volatile oils, one of which solidifies at temperatures below 50° F., and is named *anethol* or *anise camphor*, $C_{10}H_{12}O$, while the other remains fluid at all temperatures. The constitution of the latter oil is not known, but it is believed to be isomeric with oil of turpentine.

[1] As the powdered drug varies very much in alkaloidal strength, it should be avoided in dog practice, and Liquor Strychninæ Hydrochloridi used in its place.—Ed

**Mode of Preparation.**—In Europe by distilling the fruit of *Pimpinella Anisum*, and in China the fruit of *Illicium anisatum*, with water.

**Characters.**—Colourless or pale yellow, with the odour of anise and a warm sweetish taste. Completely volatilised by heat; congeals at temperatures between 50° and 60° F. (10° and 15·5° C.).

**Uses.**—For flavouring medicines, especially masses; also for destroying pediculi on pet dogs and other small animals.

## OLEUM CAJUPUTI

### CAJUPUT OIL

**Composition.**—According to Blanchet, the composition is $C_{10}H_{18}O$.

**Mode of Preparation.**—By distilling the leaves of *Melaleuca Cajuputi*. Imported from Batavia and Singapore.

**Natural Order.**—*Myrtaceæ*.

**Characters.**—Pale bluish or myrtle green colour. It is transparent, limpid, of a strong, penetrating, agreeable odour, resembling that of camphor, rosemary, and cardamoms combined, and of a warm, aromatic, camphoraceous taste, succeeded by a sensation of coldness in the mouth. Specific gravity varies from 0·914 to 0·930. Oil of Cajuput is soluble in alcohol.

**Actions and Uses.**—*Internally*:—A powerful diffusible stimulant, antispasmodic, and diaphoretic. Useful in painful spasmodic affections of the stomach, and in flatulent colic. *Externally*:—As an application in chronic rheumatism.

**Doses.**—Same as Oleum Terebinthinæ.

**Modes of Application.**—*Internally*:—In the form of an emulsion. *Externally*:—Mixed with olive oil.

## OLEUM CAMPHORATUM

CAMPHORATED OIL

Take of

| | |
|---|---|
| Camphor . . . . . | 1 part |
| Olive Oil . . . . . | 4 parts |

Dissolve the camphor in the oil.

**Uses.**—Discutient and anodyne for tumours, bruises, and chronic sprains.

## OLEUM CANTHARIDIS

OIL OF CANTHARIDES

Take of

| | |
|---|---|
| Cantharides, in powder . . . | 1 ounce |
| Olive Oil . . . . . | 8 fluid ounces |

Digest over a water-bath for two or three hours, and filter for use.

**Use.**—To maintain the action of blisters.

## OLEUM CARYOPHYLLI

OIL OF CLOVES

The oil of the clove (*Eugenia caryophyllata*) distilled in Britain.

**Characters.**—Colourless or pale yellow when fresh, becomes reddish brown on keeping. Aromatic odour and pungent taste.

**Uses.** — *Internal* : — Carminative, antispasmodic, stomachic, and corrective. Frequently given with purgatives to prevent griping. *External* :—Used to arrest the pain and check the flow of synovia in open joints and as a dressing for wounds.

**Doses.**—Horse, 20 to 30 minims.—Dog, 1 to 4 minims.

## OLEUM CROTONIS

### CROTON OIL

**Composition.**—Principally stearin, palmitin, and two compounds allied to olein; myristic, lauric, crotonic, and angelic acids have also been obtained from it. Its irritant effects upon the skin are attributed to a body called crotonal, $C_9H_{14}O_2$; but the exact nature of its purgative principle is still unknown.

**Mode of Preparation.**—By pressure from the kernels of the seeds of *Croton Tiglium*, belonging to the natural order *Euphorbiaceæ*.

**Characters.**—Fixed, viscid; colour brownish-yellow; taste acrid; odour faintly nauseous. The oil expressed in England is soluble in an equal volume of rectified spirit of wine. Indian croton oil, agitated with cold rectified spirit, forms a milky-looking emulsion, which becomes transparent on the application of heat, but which on cooling and standing allows the oil to separate and subside. Croton oil is readily dissolved by sulphuric ether and by the fixed and volatile oils.

**Actions and Uses.**—*Internally* :—In excessive doses, irritant poison; in medicinal doses, drastic purgative.

Given in obstinate constipation; in torpidity of the bowels dependent on a disordered state of the nervous system, as in tetanus and parturient apoplexy; to produce copious fluid evacuations and excite extensive counter-irritation in passive dropsies and in local inflammation in parts remote from the alimentary canal; in anasarcous swellings and effusions into cavities, when purgatives are admissible; and in affections of the kidneys when it is desirable to avoid irritating them. Also valuable for the horse in cases in which neither a bolus nor a draught can be given, as when the animal is unmanageable or unable to swallow. *Externally*:—Vesicant and counter-irritant. Applied to all animals, but especially to cattle, in bronchitis, pleurisy, pneumonia, chronic rheumatism, and glandular and other indolent swellings.

**Doses.**—Horse, 10 to 20 minims.—Cattle, 30 to 60 minims.—Sheep, 2 to 5 minims.—Pig, 2 to 5 minims. Dog, ½ to 3 minims. Usually given with other purgatives.

**Modes of Application.**—*Internally*:—Mixed with linseed meal so as to form a bolus, dissolved in olive or linseed oil or put into the animal's mash or food. The oil may be placed upon the tongue of horses that refuse their provender, or that are vicious or incapable of swallowing. To remove constipation in cattle, the oil is frequently given in conjunction with sulphate of magnesia or calomel. *Externally*:—One part of Croton Oil mixed with from four to eight parts of Soap Liniment, or Linimentum Terebinthinæ Compositum.

**Antidotes.**—Where available, an emetic of sulphate of copper, demulcent drinks, and opium to check the purgation.

**Preparation.**—Linimentum Terebinthinæ Compositum.

## OLEUM EUCALYPTI

EUCALYPTUS OIL

The oil distilled from the fresh leaves of *Eucalyptus globulus*.

**Composition.**—A volatile oil containing eucalyptol, terpene, and cymol.

**Natural Order.**—*Myrtaceæ*.

**Actions and Uses.**—A good disinfectant; applied to the skin it is an irritant. The vapour of the oil is useful in the treatment of influenza and bronchitis.

## OLEUM JUNIPERI

OIL OF JUNIPER

**Composition.**—$C_{10}H_{16}$.

**Mode of Preparation.**—In Britain by distilling the unripe fruit of *Juniperus communis*.

**Natural Order.**—*Coniferæ*.

**Characters.**—Colourless or pale greenish-yellow, sweetish odour, and warm aromatic taste. Completely volatilised by heat.

**Actions and Use.**—Stomachic and diuretic. Given as a diuretic.

**Doses.**—Horse, 1 drachm.—Cattle, 2 drachms.—Sheep, 15 minims.—Pig, 10 minims.—Dog, 2 to 5 minims.

**Mode of Application.**—Suspended in water or some mucilaginous drink.

## OLEUM LINI

LINSEED OIL

**Composition.** — Linolein, palmitin, and possibly stearin.

**Mode of Preparation.**—By expression, without heat, from linseed.

**Natural Order.**—*Linaceæ.*

**Characters.**—A fixed oil, viscid, yellow, faint odour, and oleaginous taste.

**Actions and Uses.**—*Internally*:—Nutritive, cathartic, and emollient; chiefly employed, either alone or conjoined with sulphate of magnesia, as a laxative for cattle and sheep. Its activity may be increased, if desirable, by the addition of croton oil. Given in cases of intestinal irritability arising from natural causes or from the action of poison; in colic and diarrhœa, where saline and other vegetable cathartics have proved inactive, or where their repetition is inexpedient; to give relief in cases of choking; also injected into the bladder or rectum to allay irritation. *Externally*:—In skin-affections, as an emollient to hard and dry surfaces, and as a soothing application in cases of cutaneous and other superficial irritability. Likewise often used in the preparation of liniments instead of the non-siccative, and consequently preferable, olive oil.

**Doses.**—Horse, 1 to 2 pints.—Cattle, 1 to 2 pints. Sheep, 3 to 6 fluid ounces.—Pig, 3 to 6 fluid ounces.—Dog, 1 to 3 fluid ounces.

## OLEUM MENTHÆ PIPERITÆ

### OIL OF PEPPERMINT

The oil distilled from fresh flowering peppermint.

**Composition.**—The oil contains two bodies, the fluid menthene, $C_{10}H_{18}$, and the solid crystalline menthol, $C_{10}H_{20}O$.

**Natural Order.**—*Labiatæ.*

**Characters and Tests.**—A colourless or pale yellow oil, which has a warm aromatic taste.

**Actions and Uses.**—Antiseptic, carminative, and parasiticide. Used to prevent the griping effects of purgatives and to flavour medicines.

**Doses.**—Horse and Cattle 20 to 30 minims.—Dog, 3 to 5 minims.

The solid menthol is used in the form of a pencil to relieve neuralgic and other pains.

## OLEUM MORRHUÆ

### COD-LIVER OIL

**Synonym.**—Oleum Jecoris Aselli.

**Composition.**—Chiefly consists of olein, palmitin, and stearin, with small quantities of free butyric and acetic acids, biliary constituents, gaduine, trimethylamine, and other peculiar substances, iodine, bromine, sulphur, phosphorus, and about 1 per cent. of mineral salts.

**Mode of Preparation.**—Although directed by the 'British Pharmacopœia' to be obtained from the cod, *Gadus Morrhua*, it is also procured from other species—

*e.g.*, the ling, *Gadus lota*; the dorse, *Gadus callarias*; the torsk, *Gadus Brosme*, &c.—belonging to the natural order *Acipenser*. The livers are exposed to the sun and allowed to putrefy; the oil runs from them, and is collected in vessels placed for its reception. Imported principally from Newfoundland and the North of Europe.

**Characters.**—Fixed, viscid, yellow colour, faint odour, and oleaginous taste.

**Actions and Uses.**—Nutritive and alterative in debility accompanied with emaciation, chronic rheumatism, scrofula, phthisis pulmonalis, and other diseases affecting the respiratory organs, chronic skin-affections, and distemper in dogs.

**Doses.**—Horse, 8 fluid ounces.—Cattle, 8 fluid ounces.—Sheep, 2 to 4 fluid ounces.—Pig, 2 to 4 fluid ounces.—Dog, $\frac{1}{4}$ to 1 fluid ounce. Twice a day for some considerable period, and gradually increased to double the dose originally given.

**Modes of Application.**—Administered alone or in milk or gruel, flavoured (if necessary) with an aromatic. In dog practice it may be given in capsules or in biscuits.

## OLEUM OLIVÆ [1]

### OLIVE OIL

**Composition.**—Chiefly olein and palmitin.

**Mode of Preparation.**—In the South of Europe by pressing the fleshy portion or pericarp of the ripe fruit of the olive, *Olea europæa*.

[1] Linseed oil, rape oil, and whale oil are frequently used as substitutes for olive oil in veterinary pharmacy.

**Natural Order.**—*Oleaceæ.*

**Characters.**—Fixed, pale-yellow colour; scarcely any odour; and a bland oleaginous taste. It partly congeals at about 36° F.

**Actions and Uses.**
**Doses.** } Same as Oleum Lini.

## OLEUM PALMÆ

PALM OIL

**Synonym.**—Palm Butter.

**Composition.**—Tripalmitin, $C_3H_5(C_{16}H_{31}O_2)_3$, with a small quantity of olein, $C_3H_5(C_{18}H_{33}O_2)_3$.

**Natural Order.**—*Palmaceæ.*

**Mode of Preparation.**—The kernels of the fruit of certain kinds of palm (*Cocos butyracea* or *Elæis guineënsis* or both) are crushed and heated with water. The oil rises to the surface in the liquid condition, and is removed.

**Characters.**—In this country it usually has the consistency of butter. The fresh oil melts at about 76° F. Colour usually yellow, which can be removed by bleaching; odour peculiar and agreeable, said to resemble that of violets.

**Actions and Uses.**—Substitute for lard as an emollient, and in the preparation of ointments.

## OLEUM RICINI

CASTOR OIL

**Composition.**—Principally ricinoleate of glyceryl, with other fixed oils, a resin, and possibly an alkaloid, ricinine, and a drastic purgative principle.

**Mode of Preparation.** — That imported from the West Indies, and the finer qualities from the East Indies, are procured by submitting the seeds of *Ricinus communis*, after the removal of their coats, to pressure, with or without heat. Most of the East Indian oil, however, is obtained by boiling the seeds in water, drying and crushing them, and again boiling them until the oil separates and floats on the surface. Cold-drawn castor oil is considered to possess the finest quality.

**Natural Order.**—*Euphorbiaceæ.*

**Actions and Uses.**—Cathartic, chiefly for the dog. For other animals linseed oil is usually substituted as an oleaginous purgative.

**Doses.** — HORSE, 1 to 1½ pint. — CATTLE, 1 to 1½ pint.—SHEEP, 2 to 3 fluid ounces.—PIG, 2 to 3 fluid ounces.—DOG, ½ to 2 fluid ounces.

**Modes of Application.** — Usually administered alone. It is, however, sometimes given with linseed or olive oil, or gruel and aromatics, to diminish its irritant effects; to increase its cathartic activity it is combined with small doses of oil of turpentine or of croton.

**Preparations.**—Collodium Flexile; Mistura Ricini.

## OLEUM TEREBINTHINÆ

### OIL OF TURPENTINE

**Synonyms.**—Spirit of Turpentine; Terebene.

**Composition.**—$C_{10}H_{16}$.

**Mode of Preparation.**—The outer bark near the root is removed from *Pinus sylvestris* as well as from other species of pines. Incisions are then made through the

inner bark into the wood, from which a mixture of volatile oil and resin (crude or common turpentine) flows into pits dug in the earth, whence it is transferred to casks. By distillation, with or without water, the volatile oil (oil or spirit of turpentine) is separated from the resin.

**Characters.** — Volatile, limpid, colourless; strong peculiar odour; pungent and bitter taste. Imported from France and America.

**Actions and Uses.** — *Internally*: — In excessive doses, irritant poison; in medicinal doses, astringent, stimulant, cathartic, and diaphoretic. Chiefly employed as a diuretic, antispasmodic, and anthelmintic. *Externally* :—Vesicant, counter-irritant, stimulant, and digestive. Principally applied to cattle, in conjunction with mustard or ammonia, in the treatment of inflammation of the bowels or lungs, and in chronic rheumatism.

**Doses.**—*Diuretic*:—Horse, 4 to 8 fluid drachms.—Cattle, 4 to 8 fluid drachms.—Sheep, 10 to 15 minims.—Pig, 10 to 15 minims. *Antispasmodic*:—Horse, 1 to 2 fluid ounces.—Cattle, 1 to 2 fluid ounces.—Sheep, 20 to 60 minims.—Pig, 20 to 60 minims. *Anthelmintic*:—Horse,[1] 2 to 3 fluid ounces.—Cattle,[1] 2 to 3 fluid ounces.—Sheep,[1] 2 to 4 fluid drachms.—Pig,[1] 2 to 4 fluid drachms.—Dog,[1] 1 to 1½ fluid drachm.

**Modes of Application.**—*Internally*:—Dissolved in linseed, olive, or other fixed oil; beaten into an emulsion with yolk of egg (one yolk to every two drachms of oil of turpentine); or Enema Terebinthinæ. *Externally*:—As a rubefacient or digestive, the oil alone; Linimentum Terebinthinæ; Linimentum Terebinthinæ Compositum.

[1] Combined with a small dose of castor oil, linseed oil, or solution of aloes.

As a counter-irritant for cattle, Cataplasma Terebinthinæ Compositum; for the dog, Cataplasma Terebinthinæ.

**Preparations.** — Cataplasma Terebinthinæ; Cataplasma Terebinthinæ Compositum; Enema Terebinthinæ; Linimentum Terebinthinæ; Linimentum Terebinthinæ Compositum.

## OPIUM

OPIUM

**Composition.**—Opium contains about 25 per cent. of a gummy substance, 20 per cent. of ill-defined organic matters, a little caoutchouc, resin, oil, and water, and variable proportions of a large number of alkaloids, of which morphine, narcotine, and narceine are the most abundant. Subjoined is a list of the opium alkaloids:

Hydrocotarnine, $C_{12}H_{15}NO_3$
Morphine, $C_{17}H_{19}NO_3$
Oxymorphine, $C_{17}H_{19}NO_4$
Codeine, $C_{18}H_{21}NO_3$
Thebaine, $C_{19}H_{21}NO_3$
Codamine, Laudanine } $C_{20}H_{25}NO_4$
Protopine, $C_{30}H_{19}NO_5$
Papaverine, $C_{21}H_{21}NO_4$
Meconidine, $C_{21}H_{23}NO_4$
Laudanosine, $C_{21}H_{27}NO_4$
Cryptopine, $C_{21}H_{23}NO_5$
Narcotine, $C_{22}H_{23}NO_7$
Lanthopine, $C_{23}H_{25}NO_4$
Narceine, $C_{23}H_{29}NO_9$

These alkaloids are present principally as meconates and sulphates.

**Mode of Preparation.**—Incisions, by means of an instrument consisting of four or five heart-shaped blades tied together with thread, are made in the unripe capsules of the poppy, *Papaver somniferum*. A white milky juice exudes from the incisions in drops, and is allowed to remain on the poppy head for twenty-four hours in order

that it may thicken. The inspissated exudation is then scraped off, and the different collections stirred together. The opium is finally dried without the aid of artificial heat, usually by exposure to the sun, and made into cakes or masses, which are in most places wrapped in poppy-leaves to prevent their adhering to one another. Chiefly imported from Turkey, Egypt, and India.

**Natural Order.**—*Papaveraceæ.*

**Characters.**—Irregular lumps, weighing from four ounces to two pounds; enveloped in the remains of poppy-leaves, and generally covered with the chaffy fruits of a species of *Rumex*; when fresh, plastic, tearing with an irregular, slightly moist, chestnut-brown surface, shining when rubbed smooth with the finger; peculiar odour and bitter taste.

**Actions and Uses.** — *Internally*: — In excessive doses, narcotic poison; in medicinal doses, stimulant, sedative, narcotic, anodyne, and antispasmodic. Given in gastritis, diarrhœa, dysentery, enteritis, colic, peritonitis, pleurisy, bronchitis, pneumonia, tetanus, rheumatism, and very many other maladies. *Externally*:—Anodyne to wounds and blistered and cauterised surfaces; in cases of superficial inflammation of the eye, skin, or joints; and in the treatment of hæmorrhoids.

**Doses.**—Horse, 1 to 2 drachms.—Cattle, 2 to 4 drachms.—Sheep, 10 to 40 grains.—Pig, 10 to 40 grains.—Dog, $\frac{1}{2}$ to 3 grains.

**Modes of Application.**—*Internally*:—In the form of bolus, Tinctura Opii, or Tinctura Opii Ætherea; also Enema Opii. Although frequently given alone, it is often combined with belladonna, aconite, tartar emetic, or calo-

mel when required to act as a sedative; and with chloroform, sulphuric ether, or nitrous ether when employed as a stimulant or antispasmodic. As an injection, as well as by the mouth,[1] in enteritis and inflammation of the bladder and kidneys. *Externally*:—Liquor Plumbi Subacetatis et Opii; Linimentum Opii; Tinctura Opii.

**Incompatibles.**—Lime-water; alkalies, and their carbonates; acetate of lead; sulphates of iron, zinc, and copper; perchloride of mercury; arsenites of potash and soda; and all vegetable astringents.

**Antidotes.**—Where possible, empty the stomach by means of emetics or the stomach-pump; administer brandy, ammonia, carbonate of ammonia, or some other stimulant; apply ammonia or strong acetic acid to the nostrils; dash cold water over the body, and endeavour to keep the patient moving about.

**Preparations.**—Enema Opii; Linimentum Opii; Liquor Plumbi Subacetatis et Opii; Pulvis Ipecacuanhæ Compositus; Pulvis Opii Compositus; Tinctura Opii.

## PARAFFINUM

### PARAFFIN

**Synonym.**—Tar-oil Stearin.

**Composition.**—A mixture of several solid hydrocarbons of the paraffin series.

**Mode of Preparation.**—Paraffin is found native in the coal-measures and other bituminous strata constituting the minerals known as fossil wax, ozocerite, &c. It

[1] Two or three times as much opium may be administered *per rectum* as would be given by the mouth.

exists also in a state of solution in many petroleum springs, and may be separated by distilling the more volatile portion and exposing the remainder to a low temperature.

**Characters and Tests.**—A colourless, crystalline, fatty substance, melting at about 113° F.; soluble in ether in all proportions. When melted should dissolve entirely in fixed and volatile oils. Sulphuric acid, chlorine, and nitric acid, below the temperature of 212° F., have no action upon it.

**Uses.**—Employed externally. Occasionally used in the manufacture of ointments.

## PHENACETINUM

PHENACETIN

**Composition.**—Para-acet-phenacetin—

$$C_2H_5O,C_6H_4.NHCOCH_3.$$

**Preparation.**—By the interaction of glacial acetic acid and para-phenetidin, a body obtained from para-nitro-phenol.

**Characters and Tests.**—A white crystalline powder, tasteless, without odour; soluble in alcohol and glycerin, and sparingly soluble in water.

**Actions and Uses.**—A nervous sedative and anti-pyretic, useful in influenza, rheumatism, and pneumonia. Dissolved in glycerin and water it has been found useful in distemper.

**Dose.**—Dog, 2 to 8 grains.

## PHYSOSTIGMATIS SEMINA

CALABAR BEANS

The seed of the *Physostigma venenosum*.

**Natural Order.**—*Leguminosæ*.

**Composition.**—It contains the alkaloid physostigmine, also known as eserine, $C_{15}H_{21}N_3O_2$; it also contains calabrine.

**Actions and Uses.**—Calabar bean stimulates voluntary and involuntary muscular fibres and paralyses nerve-centres. It is very poisonous, causing tetanic convulsions. It contracts the pupil of the eye. Principally employed in veterinary medicine in the treatment of obstinate constipation. A solution of the alkaloid physostigmine is usually employed hypodermically for this purpose.

**Doses** (of the bean).—Horse and Cattle, 15 to 30 grains.—Dog, $\frac{1}{4}$ to $\frac{1}{2}$ grain.

**Preparations.**—Extractum Physostigmatis; Physostigmine or Eserine.

## PHYSOSTIGMINA

PHYSOSTIGMINE

**Synonym.**—Eserine.

**Composition.**—$C_{15}H_{21}N_3O_2$.

**Mode of Preparation.**—Obtained from the alcoholic extract of Calabar bean, by dissolving the extract in water, adding sodium bicarbonate, shaking the mixture with ether, and evaporating the ethereal liquid.

**Characters and Tests.**—An alkaloid which occurs in colourless or pinkish crystals, slightly soluble in water but readily soluble in alcohol and in dilute acids.

The aqueous solution when warmed with, or when shaken with, dilute solution of potash becomes red; and when evaporated to dryness over a water-bath leaves a bluish residue, the acidified solution of which is beautifully dichroic, being blue and red.

**Action and Uses.**—Hypodermically, as sulphate or salicylate of eserine; the salicylate is preferable, as it keeps for a much longer period. For therapeutic action see Physostigmatis Semina.

**Doses.**—Horse, 2 to 3 grains.—Dog, $\frac{1}{20}$ to $\frac{1}{10}$ grain. Hypodermically, one-half these doses.

## EXTRACTUM PHYSOSTIGMATIS

### EXTRACT OF CALABAR BEAN

**Mode of Preparation.**—Take of

| | |
|---|---|
| Calabar Bean in No. 40 powder | 1 pound |
| Rectified Spirit | 4 pints |

Macerate the bean for forty-eight hours with one pint of the spirit in a closed vessel, agitating occasionally; then transfer to a percolator, and when the fluid ceases to pass add the remainder of the spirit, so that it may slowly percolate through the powder. Subject the residue of the bean to pressure, adding the expressed liquor to the product of the percolation; filter, distil off most of the spirit, and evaporate what is left in the retort by a water-bath to the consistence of a soft extract.

**Preparation.**—Physostigmine or Eserine.

**Uses.**—See Physostigmatis Semina.

## PILULA HYDRARGYRI ET FERRI

PILL OF MERCURY AND IRON

**Synonyms.**—Mercurial Pill with Iron; Ferruginat Blue Pill.

**Preparation.**—Take of

| | |
|---|---|
| Mercury, pure . . . . | 2 parts |
| Moist hydrated Peroxide of Iron . | 1 part |
| Confection of Roses . . . | 3 parts |

Rub together until the globules of mercury are no longer visible.

**Use.**—Alterative for the horse.[1]

**Dose.**—½ to 1 drachm.

## PIMENTA

PIMENTO

**Synonym.**—Allspice.

The dried unripe berries of the allspice, *Pimenta officinalis*. Imported from the West Indies.

**Natural Order.**—*Myrtaceæ*.

**Composition.**—Therapeutic value dependent on the presence of a volatile oil.

**Characters.**—Of the size of a small pea, rough, crowned with the teeth of the calyx; contains two dark-

[1] If there be any apprehension of this preparation exciting catharsis, it should be combined with opium.

brown seeds; externally brown, internally yellowish; odour and taste hot, aromatic, and peculiar.

**Actions and Uses.**—Aromatic, carminative, stomachic, and antispasmodic. Given in indigestion, relaxed stomach, flatulency, and colic. Employed also to disguise the flavour of nauseous medicines.

**Doses.**—Horse, 2 to 4 drachms.—Cattle, 2 to 6 drachms.—Sheep, ½ to 1 drachm.—Pig, ½ to 1 drachm.—Dog, 10 to 30 grains.

**Modes of Application.**—In the form of powder, bolus, or Tinctura Pimentæ.

**Preparation.**—Tinctura Pimentæ.

## PIPER NIGRUM

BLACK PEPPER

**Natural Order.**—*Piperaceæ.*

**Composition.**—Its medicinal properties are due chiefly to a volatile oil and an alkaloid (piperine).

**Mode of Preparation.**—The berries of *Piper nigrum*, before they have all changed to red, are collected, dried, and ground to powder.

**Characters.**—Small roundish wrinkled bodies; tegument brownish black, containing a greyish-yellow globular seed (white pepper).

**Actions and Uses.** — Principally employed as a stomachic and carminative in simple indigestion, and as an aromatic to mask the unpleasant flavour of many medicines.

**Doses.**—Horse, 2 drachms.—Cattle, 3 drachms.—Sheep, 20 to 40 grains. Repeated two or three times a day.

**Mode of Application.**—In the form of bolus, or, preferably, suspended in gruel.

## PIX BURGUNDICA

### BURGUNDY PITCH

A resinous exudation from the stem of the spruce fir, *Picea excelsa*, melted and strained. Imported from Switzerland.

**Natural Order.**—*Coniferæ.*

**Characters.**—Hard and brittle, yet gradually taking the form of the vessel in which it is kept; opaque, varying in colour, but generally dull reddish brown; of a peculiar, somewhat empyreumatic perfumed odour, and aromatic taste, without bitterness; free from vesicles; gives off no water when heated.

**Actions and Uses.** — *Internally*: — Irritant and diuretic. *Externally*:—Rubefacient and digestive; stimulant to sprains; to give support to fractures; and as an adhesive to wounds.

**Modes of Application.**—*Externally*:—Melted and applied, after the hair of the part has been loosened by the currycomb, to fractures and sprains. As an adhesive, Emplastrum Picis is used.

**Doses.**—*Internally*:—Horse, 1 to 3 ounces.—Cattle, 1 to 3 ounces.—Sheep, 1 to 2 drachms.—Dog, 20 to 40 grains.

## PIX LIQUIDA

TAR

**Synonyms.**—Wood Tar ; Stockholm Tar.

**Composition.**—Hydrocarbons, methylic alcohol, carbolic acid, acetic acid, cresylic acid, resinous bodies, pitch, and many other substances.

**Mode of Preparation.**—By submitting to destructive distillation the wood of *Pinus sylvestris* and other pines.

**Characters.**—Thick, viscid, brownish-black liquid, of a well-known peculiar aromatic odour. Water agitated with it acquires a pale-brown colour, sharp empyreumatic taste, and acid reaction.

**Actions and Uses.**—*Externally* :—As an excitant, rubefacient, antiseptic, and deodoriser in the treatment of thrush in the horse, and of broken horns and punctured wounds of the belly and chest of cattle; in foot-rot in sheep; and in mange, eczema, impetigo, and other skin-affections.[1]

**Modes of Application.**—In thrush and foot-rot, tar alone is applied to the diseased parts; for broken horns and punctured wounds, the tar should be spread over coarse cloth ; for skin-affections, Linimentum Olei Picis Liquidæ Compositum or Linimentum Picis Liquidæ is used ; and as a common foot-ointment for all domesticated animals, Unguentum Picis Liquidæ is employed.

[1] A '*Stopping for Feet*' is made by mixing together—

| | |
|---|---|
| Tar | 2 parts |
| Soft Soap | 1 part |
| Linseed Meal | a sufficiency to impart tenacity to the mixture. |

**Preparations.** — Linimentum Olei Picis Liquidæ Compositum; Linimentum Picis Liquidæ; Unguentum Picis Liquidæ

## PIX NIGRA

PITCH

**Synonym.**—Black Pitch.

The black, resinoid, and highly carbonaceous residue which is left when the volatile constituents of tar are removed by distillation.

**Uses.**—Mild stimulant in thrush, canker, and sand-crack in horses, and foot-rot in sheep. Also used to impart consistency and adhesiveness to plasters and 'charges.'

## PLUMBI ACETAS

ACETATE OF LEAD

**Synonyms.**—Lead Acetate; Sugar of Lead.

**Composition.**—$Pb(C_2H_3O_2)_2.3H_2O$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Oxide of Lead (Litharge), in fine powder . . | 24 ounces |
| Acetic Acid . . . | 2 pints or a sufficiency |
| Distilled Water . . | 1 pint |

Mix the acetic acid and the water, add the oxide of lead, and dissolve with the aid of a gentle heat. Filter, evaporate till a pellicle forms, and set aside to crystallise, first adding a little acetic acid should the fluid

not have a distinctly acid reaction. Drain, and dry the crystals on filtering-paper without heat.

| Lead Monoxide. | | Acetic Acid. | | Lead Acetate (Acetate of Lead). | | Water. |
|---|---|---|---|---|---|---|
| $PbO$ | + | $2HC_2H_3O_2$ | = | $Pb(C_2H_3O_2)_2$ | + | $H_2O$ |

In the act of crystallising the salt unites with three molecules of water.

**Characters and Tests.**—White crystalline masses, slightly efflorescent; acetous odour, and sweet astringent taste. It dissolves in distilled water, and forms a clear solution, or one which has only a slight milkiness, which disappears on the addition of acetic acid. Its solution in water slightly reddens litmus, gives a yellow precipitate (lead iodide) with potassium iodide, and a white precipitate (lead sulphate) with dilute sulphuric acid, acetic acid being set free and recognisable by its odour; the precipitates by potassium iodide and sulphuric acid indicate the presence of lead, and the elimination of acetic acid shows that the salt is an acetate. Air blown from the lungs through its aqueous solution should not occasion any turbidity, showing its freedom from subacetate.

**Actions and Uses.**—Sedative and astringent. Seldom employed in veterinary practice; it is, however, occasionally administered in from 1 to 2 drachm doses to the horse in diarrhœa and diabetes insipidus. Principally used in the production of Liquor Plumbi Subacetatis.

**Preparation.**[1]—Liquor Plumbi Subacetatis.

[1] Plumbi acetas and zinci sulphas, dissolved in water, form the well-known 'white lotion'

## PLUMBI OXIDUM

OXIDE OF LEAD

**Synonyms.**—Lead Monoxide; Protoxide of Lead; Litharge.

**Composition.**—PbO.

**Mode of Preparation.**—Metallic lead is oxodised by being heated in a current of air.

| Lead. | | Oxygen (from Air). | | Lead Monoxide. |
|---|---|---|---|---|
| $2Pb$ | + | $O_2$ | = | $2PbO$ |

**Characters and Tests.**—Heavy scales, pale brick-red colour, completely soluble without effervescence in dilute nitric and acetic acids, either solution, when neutral, giving a yellow precipitate (lead iodide) with potassium iodide, indicating the presence of lead. Its solution in dilute nitric acid, when saturated with ammonia and filtered, does not exhibit a blue colour, showing that it is free from copper.

**Uses.**—In preparing Plumbi Acetas and Liquor Plumbi Subacetatis.

## PODOPHYLLI RESINA

RESIN OF PODOPHYLLUM

Take of

| | |
|---|---|
| Podophyllum-root, in coarse powder . | 1 pound |
| Alcohol (90-per-cent.) . . . | 3 pints, or a sufficiency |
| Distilled Water . . . . . | a sufficiency |
| Hydrochloric Acid . . . . | a sufficiency |

Exhaust the podophyllum by percolation with the spirit; distil over the spirit; slowly pour the liquid

remaining after the distillation of the tincture into three times its volume of water acidulated with one twenty-fourth part of its weight of hydrochloric acid, constantly stirring; let it stand twenty-four hours; collect the resin which falls, wash on a filter with distilled water, and dry in a stove.

**Characters and Tests.**—A pale greenish-brown, amorphous powder, soluble in rectified spirit and in ammonia; precipitated from the former solution by water, from the latter by acids. Almost entirely soluble in pure ether.

**Actions and Uses.**—See Podophylli Rhizoma.

**Doses.**—Horse, 1 to 2 drachms.—Cattle, 1 to 2 drachms.—Dog, 1 to 2 grains.

## PODOPHYLLI RHIZOMA

PODOPHYLLUM-RHIZOME

**Synonyms.**—Podophylli Radix; Podophyllum-root.

The dried rhizome of *Podophyllum peltatum.* Imported from North America.

**Natural Order.**—*Ranunculaceæ.*

**Characters.**—In pieces of variable length, about two inches thick; mostly wrinkled longitudinally; dark reddish brown externally, whitish within; breaking with a short fracture; accompanied with pale brown rootlets. Powder yellowish-grey, sweetish in odour, bitterish, subacrid, and nauseous in taste.

**Preparation.**—Podophylli Resina.

**Actions and Uses.**—Cholagogue, purgative; used as a substitute for calomel. Experiments have shown that podophyllin is not likely to be serviceable as a purgative for the domesticated animals. It is given to lower the

heart's action in acute diseases of the respiratory organs, rheumatism, laminitis, and other inflammatory disorders.

**Doses.**—*As a sedative* :—HORSE, 1 to 2 drachms.—CATTLE, 1 to 2 drachms.—DOG, 1 to 2 grains.

**Modes of Application.**—United with calomel, or potassium nitrate, or magnesium sulphate. For dogs it may be administered with calomel, grey powder, or ipecacuanha.

## POTASSA CAUSTICA

### POTASSIUM HYDROXIDE

**Synonyms.**—Potassii Hydras ; Potassium Hydrate ; Caustic Potash.

**Composition.**—KOH.

**Mode of Preparation.**—Take of

Solution of Potash . . . 2 pints

Boil down the solution of potash rapidly in a silver or clean iron vessel until there remains a fluid of oily consistence, a drop of which, when removed on a warm glass rod, solidifies on cooling. Pour this into proper moulds, and when it has solidified, and while it is still warm, put it into a stoppered bottle.

**Characters and Tests.**—Hard white pencils, very deliquescent, powerfully alkaline, and corrosive. A watery solution acidulated by hydrochloric acid gives a yellow precipitate (potassium-platinic chloride) with platinic chloride, indicating the presence of potassium. It should dissolve in dilute nitric acid without effervescence, showing the absence of carbonate, and give but scanty white precipitates with barium chloride and silver nitrate, showing the presence of traces only of sulphate and chloride respectively.

**Actions and Uses.**—Principally applied as an escharotic to wounds produced by the bites of rabid animals, also for stimulating unhealthy ulcers and inducing suppuration. In consequence of the highly deliquescent nature of this preparation, it has an objectionable tendency to spread beyond the part upon which it is required to act. The use of Soda Caustica is not open to this objection, and as its escharotic effects rival those of Potassa Caustica, it may be advantageously substituted for the latter agent.

## POTASSA SULPHURATA

### SULPHURATED POTASH

**Synonyms.**—Potassii Sulphuratum; Hepar Sulphuris; Sulphide of Potassium; Liver of Sulphur.

**Composition.**—A mixture of several sulphides, sulphates, and other compounds of potassium.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Carbonate of Potash . . . | 10 ounces |
| Sublimed Sulphur . . . | 5 ounces |

Mix the carbonate of potash and the sulphur in a warm mortar, and, having introduced them into a Cornish or Hessian crucible, let them be heated, first gradually, until effervescence has ceased, and finally to dull redness, so as to produce perfect fusion. Let the liquid contents of the crucible be then poured out on a clean flagstone, and covered quickly with an inverted porcelain basin, so as to exclude the air as completely as possible while solidification is taking place. The solid product thus obtained should, when cold, be broken into fragments, and immediately enclosed in a green-glass bottle furnished with an air-tight stopper.

**Characters and Tests.**—Solid greenish fragments liver-brown when recently broken, alkaline and acrid to

the taste; readily forms with water a yellow solution which has the odour of sulphuretted hydrogen, and evolves it freely when excess of hydrochloric acid is dropped into it, sulphur being at the same time deposited. The acid fluid, when boiled and filtered, is precipitated yellow (potassium-platinic chloride) by platinic chloride, showing the presence of potassium, and white by barium chloride, indicating the presence of sulphate. About three-fourths of its weight is dissolved by rectified spirit.

**Actions and Uses.**—*Internally* :—In excessive doses, irritant poison ; in doses of from half to one drachm to the horse it acts as a stimulant and diaphoretic. Very seldom employed as an internal remedy in this country. *Externally* :—In chronic skin-diseases, especially those of a scaly character.

**Modes of Application.**—*Externally* :—Liquor Potassæ Sulphuratæ ; Unguentum Potassæ Sulphuratæ.

**Antidotes.**—Solution of chlorinated lime, with an emollient drink.

**Preparations.**—Liquor Potassæ Sulphuratæ; Unguentum Potassæ Sulphuratæ.

## POTASSII BICARBONAS

### POTASSIUM BICARBONATE

**Synonym.**—Potassium-hydrogen Carbonate.

**Composition.**—$KHCO_3$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Carbonate of Potash . . . | 1 pound |
| Distilled Water . . . . | 2 pints |
| Hydrochloric Acid . . . | 1½ pint |
| Common Water . . . . | 3 pints |
| White Marble, in fragments . | 1 pound, or a sufficiency |

Dissolve the carbonate of potash in the distilled water, and filter the solution into a three-pint bottle capable of being tightly closed by a cork traversed by a glass tube sufficiently long to pass to the bottom of the fluid. Introduce the marble into another bottle, in the bottom of which a few small holes have been drilled, and the mouth of which is closed by a cork also traversed by a glass tube, and place the bottle in a jar of the same height as itself. Join the two glass tubes, so that the connection is air-tight, by a caoutchouc tube. The cork of the bottle containing the carbonate of potash having been placed loosely, and that of the bottle containing the marble tightly, in its mouth, pour into the jar surrounding the latter bottle the hydrochloric acid, previously diluted with the common water.

When carbonic anhydride has passed through the carbonate-of-potash solution for two minutes, so as to expel the whole of the air from the apparatus, fix the cork tightly in the neck of the bottle, and let the process go on for a week. At the end of this time numerous crystals of bicarbonate of potash will have formed, which are to be removed, shaken with twice their bulk of cold distilled water, and, after decantation of the water, drained, and dried on filtering-paper by exposure to the air. The mother-liquor, filtered if necessary, and concentrated to one-half, at a temperature not exceeding 110° F., will yield more crystals. The tube immersed in the solution of carbonate of potash, which should have as large a diameter as possible, may require the occasional removal of the crystals formed within it, in order that the process may not be interrupted.

| Potassium Carbonate. | | Water. | | Carbonic Anhydride. | | Potassium-hydrogen Carbonate (Bicarbonate of Potash). |
|---|---|---|---|---|---|---|
| $K_2CO_3$ | + | $H_2O$ | + | $CO_2$ | = | $2HKCO_3$ |

**Characters and Tests.**—Colourless right rhombic prisms, not deliquescent; saline, feebly alkaline, taste; not corrosive. The addition of dilute hydrochloric acid causes strong effervescence (escape of carbonic anhydride), and forms a solution which gives a yellow precipitate (potassium-platinic chloride). An aqueous solution gives no precipitate with magnesium sulphate, owing to the formation of soluble magnesium-hydrogen carbonate.

**Actions and Uses.**—In excessive doses, irritant poison; in medicinal doses, antacid and diuretic.

**Doses.**—Horse, 2 to 6 drachms.—Cattle, 2 to 6 drachms.—Sheep, 40 grains to 2 drachms.—Pig, 40 grains to 2 drachms.—Dog, 10 to 30 grains.

**Modes of Application.**—In the form of bolus or dissolved in water.

**Incompatibles.**—Acids; lime-water; acetate and chloride of ammonium; most metallic salts, but not sulphate of magnesia.

**Antidotes.**—Highly diluted acetic acid or vinegar, and linseed or olive oil.

## POTASSII BROMIDUM

### POTASSIUM BROMIDE

**Synonym.**—Bromide of Potassium.

**Composition.**—KBr.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Solution of Potash . . . | 2 pints |
| Bromine . . . . . | 4 fluid ounces |
| Wood Charcoal, in fine powder . | 2 ounces |
| Boiling Distilled Water . . | 1½ pint |

Put the solution of potash into a porcelain vessel, and add the bromine in successive portions, with agitation, until the mixture has acquired a brown tint. Evaporate to dryness, reduce the residue to a fine powder, and mix this intimately with the charcoal. Throw the mixture, in small quantities at a time, into a red-hot crucible, and when the whole has been brought to a state of fusion remove the crucible from the fire and turn out its contents.

When the fused mass has cooled, dissolve it in the water, filter the solution through paper, and set it aside to crystallise. The crystals should be dried at a gentle heat.

The chemical reaction occurring in this process takes place in two stages:

| | Potassium Hydrate. | | Bromine. | | Potassium Bromate. | | Potassium Bromide. | | Water. |
|---|---|---|---|---|---|---|---|---|---|
| 1. | $6KHO$ | + | $3Br_2$ | = | $KBrO_3$ | + | $5KBr$ | + | $3H_2O$ |

| | Potassium Bromide. | | Potassium Bromate. | | Carbon. | | Potassium Bromide. | | Carbonic Oxide. |
|---|---|---|---|---|---|---|---|---|---|
| 2. | $10KBr$ | + | $2KBrO_3$ | + | $3C_2$ | = | $12KBr$ | + | $6CO$ |

**Characters and Tests.**—In colourless, cubical crystals, with no odour, but a pungent, saline taste. Soluble in water, 1 in 2; less soluble in spirit. Its aqueous solution gives a white crystalline precipitate with tartaric acid ($KHC_4H_4O_6$), proving the presence of potassium. When its solution in water is mixed with a little chlorine, chloroform agitated with it, on falling to the bottom, exhibits a red colour (bromine). Silver nitrate will also give a yellowish-white precipitate, insoluble in dilute nitric acid; dissolving sparingly in ammonium hydrate, but readily in potassium cyanide.

**Action and Use.**—As a sedative. It exerts a powerful influence on the generative organs, lowering their functions in a marked degree.

**Doses.**—HORSE, 2 to 4 drachms.—CATTLE, 2 to 5 drachms.—DOG, 5 to 20 grains. Repeated when required.

## POTASSII CARBONAS

### POTASSIUM CARBONATE

**Synonyms.**—Carbonate of Potash; Salt of Tartar.

**Composition.**—$K_2CO_3$, with about 16 per cent. of water of crystallisation.

**Mode of Preparation.**—In the United States, Canada, Russia, and on the shores of the Baltic, various kinds of wood are burned, and their ashes collected and lixiviated. By lixiviation various potassium compounds —*e.g.*, carbonate, sulphate, chloride, phosphate, and silicate—are dissolved, while silica and certain salts of calcium, magnesium, iron, and aluminium remain insoluble. The soluble portions of the ash are evaporated to dryness in iron pots, and deprived of their volatile impurities by being heated in an oven-shaped furnace, whose flame is made to play over the mass. The product thus obtained is impure carbonate of potash, and is commercially termed pearlash. In order to obtain medicinal carbonate of potash, pearlash is treated with its own weight of distilled water (to dissolve the carbonate and leave undissolved the other potassium salts), and the solution so formed evaporated to dryness, while it is kept briskly agitated.

**Characters and Tests.**—White crystalline powder, alkaline and caustic to the taste, very deliquescent, readily soluble in water, but insoluble in spirit of wine. It effervesces with dilute hydrochloric acid, showing that it is a carbonate, and forms a solution which gives a yellow precipitate (potassium-platinic chloride) with platinic chloride. Loses about 16 per cent. of its weight (water) when exposed to a red heat. Its aqueous solution gives a white precipitate (magnesium carbonate) with solution of magnesium sulphate: it is thus distinguished from potassii bicarbonas. If dissolved in water and mixed with excess of nitric acid, it should give but a faint precipitate on the addition of barium chloride and silver nitrate, indicating its almost perfect freedom from sulphates and chlorides.

**Actions and Uses.**—Same as Potassii Bicarbonas, but more liable than that agent to irritate the stomach and intestines.

**Doses.**—Horse, 2 to 4 drachms.—Cattle, 2 to 4 drachms.—Sheep, 30 grains to 1½ drachm.—Pig, 30 grains to 1½ drachm.—Dog, 5 to 20 grains.

**Mode of Application.**—In the form of bolus or dissolved in water.

**Incompatibles.**—Same as with Potassii Bicarbonas.

**Antidotes.**—Same as for Potassii Bicarbonas.

## POTASSII CHLORAS

### POTASSIUM CHLORATE

**Synonym.**—Chlorate of Potassium

**Composition.**—$KClO_3$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Carbonate of Potash . . | 20 ounces |
| Slaked Lime . . . . | 53 ounces |
| Distilled Water . . . | a sufficiency |
| Black Oxide of Manganese . | 83 ounces |
| Hydrochloric Acid . . . | 24 pints |

Mix the lime with the carbonate of potash, and triturate them with a few ounces of the water, so as to make the mixture slightly moist. Place the oxide of manganese in a large retort or flask, and, having poured upon it the hydrochloric acid, diluted with six pints of water, apply a gentle heat, and conduct the chlorine as it comes over, first through a bottle containing six ounces of water, and then into a large carboy containing the mixture of carbonate of potash and slaked lime. When the whole of the chlorine has come over, remove the contents of the carboy, and boil them for twenty minutes with seven pints of the water, filter, evaporate till a film forms on the surface, and set aside to cool and crystallise. The crystals thus obtained are to be purified by dissolving them in three times their weight of boiling distilled water and again allowing the solution to crystallise.

| Calcium Hydrate. | | Potassium Carbonate. | | Chlorine. | | Potassium Chlorate (Chlorate of Potash). | |
|---|---|---|---|---|---|---|---|
| $5Ca(HO)_2$ | + | $K_2CO_3$ | + | $6Cl_2$ | = | $2KClO_3$ | + |

| Calcium Chloride. | | Carbonic Anhydride. | | Water. |
|---|---|---|---|---|
| $5CaCl_2$ | + | $CO_2$ | + | $5H_2O$ |

The potassium chloride crystallises out, while the calcium chloride remains in solution.

**Characters and Tests.**—Colourless, rhomboidal, crystalline plates; cool saline taste; soluble in cold water

1 in 12, boiling water 1 in 2. Its aqueous solution is not affected by silver nitrate or ammonium oxalate. Strongly heated, it fuses, gives off oxygen gas, and leaves a white residue (potassium chloride), which readily dissolves in water and produces a solution which yields a white precipitate (silver chloride) with silver nitrate, indicating the presence of chlorine, and a yellow precipitate (potassium-platinic chloride) with platinum chloride, showing the presence of potassium. It explodes when triturated in a mortar with sulphur.[1]

**Actions and Uses.**—*Internally*:—Mild stimulant, diuretic, and refrigerant. Given principally in tympanites and hoven, from its having the power, it is said, of causing a condensation of the gases developed in these complaints. *Externally*:—Stimulant and refrigerant to cancerous and other ill-conditioned ulcers.

**Doses.**—Horse, 1 to 4 drachms.—Cattle, 2 to 6 drachms.—Sheep, 20 to 40 grains.—Pig, 20 to 40 grains. Dog, 5 to 15 grains.

**Modes of Application.**—*Internally*:—Dissolved in water. *Externally*:—In the form of Liquor Potassii Chloratis Compositus.

**Preparation.**—Liquor Potassii Chloratis Compositus.

## POTASSII IODIDUM

### IODIDE OF POTASSIUM

**Synonym.**—Potassium Iodide.

**Composition.**—KI.

[1] To avoid accidents, the particles used in performing this experiment should not be larger than a pin's head

**Mode of Preparation.**—Take of

| | |
|---|---|
| Solution of Potash . . . | 1 gallon |
| Iodine . . . . . | 29 ounces, or a sufficiency |
| Wood Charcoal, in fine powder . | 3 ounces |

Put the solution of potash into a glass or porcelain vessel, and add the iodine in small quantities at a time, with constant agitation, until the solution acquires a permanent brown tint. Evaporate the whole to dryness in a porcelain dish, pulverise the residue (a mixture of potassium iodide and iodate), and mix it intimately with the charcoal. Throw the mixture, in small quantities at a time, into a red-hot iron crucible, and when the whole has been brought to a state of fusion, remove the crucible from the fire and pour out its contents.

When the fused mass (impure potassium iodide) has cooled, dissolve it in two pints of boiling distilled water, filter through paper, wash the filter with a little boiling distilled water, mix the liquids, evaporate the whole till a film forms on the surface, and set it aside to cool and crystallise. Drain the crystals, and dry them quickly with a gentle heat. More crystals may be obtained by evaporating the mother-liquor and cooling. The salt should be kept in a stoppered bottle. Two decompositions take place in this process :

| | Potassium Hydrate. | | Iodine. | | Potassium Iodide. | | Potassium Iodate. | | Water. |
|---|---|---|---|---|---|---|---|---|---|
| 1. | $6HKO$ | + | $3I_2$ | = | $5KI$ | + | $KIO_3$ | + | $3H_2O$ |

| | Potassium Iodide. | | Potassium Iodate. | | Carbon. | | Potassium Iodide. | | Carbonic Oxide. |
|---|---|---|---|---|---|---|---|---|---|
| 2. | $5KI$ | + | $KIO_3$ | + | $C_3$ | = | $6KI$ | + | $3CO$ |

**Characters and Tests.**—Colourless, generally opaque, cubic crystals, readily soluble in water, less so in

spirit of wine. It commonly has a feeble alkaline reaction. Its solution mixed with mucilage of starch gives a blue colour on the addition of a minute quantity of solution of chlorine (to set free iodine), thus proving that the salt is an iodide. Its aqueous solution mixed with solution of tartaric acid or hydrogen-sodium tartrate, furnishes a crystalline precipitate (hydrogen-potassium tartrate), indicating the presence of potassium.

Silver nitrate gives, with an aqueous solution of the salt, a yellowish-white precipitate (silver iodide), which, when agitated with ammonia, yields by subsidence a clear liquid, in which excess of nitric acid should cause no turbidity, showing the absence of chlorides.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant poison; in medicinal doses, alterative, deobstruent, and diuretic. Given in scrofula, enlargement of the lymphatic glands, periostitis, articular rheumatism, and dropsy.[1] *Externally*:—To promote the absorption of tumours and other abnormal growths. Also employed as a solvent for iodine, which it strongly resembles in therapeutic action.

**Doses.**—Horse, 1 to 2 drachms.—Cattle, 1 to 3 drachms.—Sheep, 20 grains to 1 drachm.—Pig, 20 grains to 1 drachm.—Dog, 1 to 5 grains. Repeated two or three times a day.

**Modes of Application.**—*Internally*:—In the form of bolus or dissolved in water. *Externally*:—Unguentum Potassii Iodidi; Unguentum Iodi Compositum.

**Incompatibles.**—Acids and Spiritus Ætheris Nitrosi.

[1] Potassium Iodide is now largely used in the treatment of actinomycosis and other tumours due to discomyces. See also Liq. Potassæ, and Liq. Calcis Iodinatæ (in Addenda).

### Preparations.

| | |
|---|---|
| Linimentum Iodi Comp. | Tinctura Iodi. |
| Liquor Iodi et Potassii Iodidi. | Unguentum Iodi. |
| | Unguentum Iodi Comp. |

## POTASSII NITRAS

POTASSIUM NITRATE

**Synonyms.**—Nitrate of Potash ; Nitre ; Saltpetre.

**Composition.**—$KNO_3$.

**Mode of Preparation.**—In India and other hot climates putrefying organic matter is naturally, and in some parts of Europe it is artificially, associated with alkaline and earthy carbonates, together with other materials which render the mass porous. Under these conditions the ammonia arising from the substances undergoing putrefaction is transformed into nitric acid at the expense of atmospheric oxygen.

| Ammonia. | | Oxygen. | | Nitric Acid. | | Water. |
|---|---|---|---|---|---|---|
| $NH_3$ | + | $2O_2$ | = | $HNO_3$ | + | $H_2O$ |

The nitric acid thus formed is converted into calcium and potassium nitrates by contact with calcium and potassium carbonates.

| | Nitric Acid. | | Calcium Carbonate. | | Calcium Nitrate. | | Carbonic Anhydride. | | Water. |
|---|---|---|---|---|---|---|---|---|---|
| 1. | $2HNO_3$ | + | $CaCO_3$ | = | $Ca(NO_3)_2$ | + | $CO_2$ | + | $H_2O$ |

| | Nitric Acid. | | Potassium Carbonate. | | Potassium Nitrate (Nitrate of Potash). | | Carbonic Anhydride. | | Water. |
|---|---|---|---|---|---|---|---|---|---|
| 2. | $2HNO_3$ | + | $K_2CO_3$ | = | $2KNO_3$ | + | $CO_2$ | + | $H_2O$ |

Soil containing nitrates is known as *nitre-* or *saltpetre-earth*, and is the source of large quantities of the nitrate

of potash used in medicine and in the arts. The nitre-earth is lixiviated, and the watery solution obtained is mixed with solution of potassium carbonate, to convert any calcium nitrate into the corresponding potassium salt.

| Calcium Nitrate. | | Potassium Carbonate. | | Potassium Nitrate (Nitrate of Potash). | | Calcium Carbonate. |
|---|---|---|---|---|---|---|
| $Ca(NO_3)_2$ | + | $K_2CO_3$ | = | $2KNO_3$ | + | $CaCO_3$ |

By decantation or filtration a clear solution is procured, which, on evaporation, deposits nitrate of potash in a crystalline state. The salt is purified by solution and careful recrystallisation.

**Characters and Tests.**—White crystalline masses or fragments of striated six-sided prisms, colourless; peculiar cool saline taste. Soluble in cold water 1 in 4 parts, boiling water 1 in $2\frac{1}{2}$ parts. Thrown on the fire or on red-hot charcoal, deflagration takes place; warmed in a test-tube with sulphuric acid and metallic copper, red fumes (nitric peroxide) are evolved; these two reactions show that the salt is a nitrate. Its aqueous solution, acidulated with hydrochloric acid, gives a yellow precipitate (potassium-platinic chloride) with platinic chloride, indicating the presence of potassium. Its aqueous solution should not be affected by barium chloride or silver nitrate, proving its freedom from sulphates and chlorides respectively.

**Actions and Uses.** — *Internally*:—In excessive doses, irritant poison; in medicinal doses, diuretic, febrifuge, and refrigerant. *Externally*:—Stimulant to unhealthy wounds.

**Doses.**—Horse, 2 to 4 drachms.—Cattle, 2 to 4 drachms.—Sheep, $\frac{1}{2}$ to 1 drachm.—Pig, $\frac{1}{2}$ to 1 drachm.—Dog, 5 to 10 grains.

**Modes of Application.**—*Internally*:—As a febrifuge, it is best given, when possible, in the form of bolus; as a diuretic, it is administered in solution in water. Where nitre is used as a refrigerant, the solution should not be made until just before it is to be given.—*Externally*:—In the form of Liquor Potassii Nitratis.

**Incompatibles.**—Sulphuric acid and sulphates.

**Preparations.** — Liquor Potassii Nitratis; Lotio Ammonii Chloridi et Potassii Nitratis.

## POTASSII PERMANGANAS

### PERMANGANATE OF POTASH

**Synonym.**—Potassium Permanganate.

**Composition.**—$K_2Mn_2O_8$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Caustic Potash . . . . | 5 ounces |
| Black Oxide of Manganese, in fine powder . . . | 4 ounces |
| Chlorate of Potash . . . | $3\frac{1}{2}$ ounces |
| Dilute Sulphuric Acid . . . | a sufficiency |
| Distilled Water . . . . | $2\frac{1}{4}$ pints |

Finely pulverise the chlorate of potash in a perfectly clean mortar, and mix it with the oxide of manganese;[1] put the mixture into a porcelain basin, and add to it the caustic potash, previously dissolved in four ounces of the water. Evaporate to dryness, stirring diligently to prevent spurting. Pulverise the residual mass, put it into a Hessian or Cornish crucible, and expose it to a dull

[1] The mixture should be effected by stirring with a spatula; if the manganese and chlorate be ground together with a pestle, a dangerous explosion may occur.

red heat for an hour, or till it has assumed the condition of a semi-fused green mass (potassium manganate). When cool, pulverise it, and boil it with a pint and a half of the water, for the purpose of converting it into permanganate. Let the insoluble matter (manganese dioxide) subside, decant the fluid, boil again with half a pint of the water, again decant, neutralise the united liquors accurately with the dilute sulphuric acid, and evaporate till a pellicle forms. Set aside to cool and crystallise. Drain the crystalline mass, boil it in six ounces of the water, and strain through a funnel the throat of which is slightly obstructed by a little asbestos. Allow the fluid to cool and crystallise, drain the crystals, and dry them by placing them under a bell-jar over a vessel containing sulphuric acid.

The chemical reactions occurring in this process take place in two stages.

| | Manganese Dioxide. | | Potassium Chlorate. | | Potassium Hydrate. | | Potassium Manganate. | |
|---|---|---|---|---|---|---|---|---|
| 1. | $3MnO_2$ | + | $KClO_3$ | + | $6KHO$ | = | $3K_2MnO_4$ | + |

| | Potassium Chloride. | | Water. |
|---|---|---|---|
| | $KCl$ | + | $3H_2O$ |

| | Potassium Manganate. | | Water. | | Potassium Permanganate (Permanganate of Potash). | | Manganese Dioxide. | |
|---|---|---|---|---|---|---|---|---|
| 2. | $3K_2MnO_4$ | + | $2H_2O$ | = | $K_2Mn_2O_8$ | + | $MnO_2$ | + |

| Potassium Hydrate. |
|---|
| $4KHO$ |

The dilute sulphuric acid is added to neutralise the potassium hydrate liberated in the second reaction.

**Characters and Tests.** — Dark purple, prismatic crystals, with a sweet astringent taste. Completely dissolved by cold water, forming a rich purple solution.

**Actions and Uses.** — Applied as an excitant and deodoriser to cancerous and ill-conditioned wounds, and in the treatment of certain skin-affections. Principally employed as a deodoriser and disinfectant.

**Modes of Application.** — To fœtid wounds and ulcers in the form of Liquor Potassii Permanganatis. This preparation is also used in aphthous affections of the mouth and throat, and as an injection in foul-smelling discharges of the uterus and vagina.

**Preparation.**—Liquor Potassii Permanganatis.

## POTASSII PRUSSIAS FLAVA

### YELLOW PRUSSIATE OF POTASH

**Synonyms.**—Potassium Ferrocyanide ; Ferrocyanide of Potassium.

**Composition.**—$K_4FeC_6N_6.3H_2O$, or $K_4FeCy_6.3H_2O$.

**Mode of Preparation.** — Cuttings of horns, skins, hoofs, and dry refuse animal matter of various kinds are fused at a red heat in an iron vessel with potassium carbonate and scraps of iron. The resulting mass is treated with hot water and filtered. On evaporation the filtrate deposits the ferrocyanide, which is purified by recrystallisation. Carbon and nitrogen from the animal matter and potassium from the potassium carbonate unite with the iron to form potassium ferrocyanide.

**Characters and Tests.** — Yellow crystals derived from the octahedron, permanent in the air, soluble in water, insoluble in alcohol. The aqueous solution gives a

deep blue precipitate (ferric ferrocyanide[1]) with a ferric salt; a precipitate which is at first white (potassio-ferrous ferrocyanide[2]), then light blue, and finally, by exposure to the air, dark blue,[3] with a ferrous salt; and a reddish-brown precipitate (cupric ferrocyanide) with cupric sulphate. Heated with dilute sulphuric acid, hydrocyanic acid vapours are evolved.

**Use.**—For preparing Acidum Hydrocyanicum Dilutum.

## POTASSII SULPHAS

### POTASSIUM SULPHATE

**Synonym.**—Sulphate of Potash.

**Composition.**—$K_2SO_4$.

**Mode of Preparation.**—Hydrogen-potassium sulphate, which is left in the retort in the preparation of nitric acid, is dissolved in water and neutralised by potassium carbonate. By evaporating the solution the salt is thrown down, and by recrystallisation it is purified.

| Hydrogen-potassium Sulphate. | | Potassium Carbonate. | | Potassium Sulphate. | | Carbonic Anhydride. | | Water |
|---|---|---|---|---|---|---|---|---|
| $2HKSO_4$ | + | $K_2CO_3$ | = | $2K_2SO_4$ | + | $CO_2$ | + | $H_2O$ |

**Use.**—In the preparation of Pulvis Ipecacuanhæ Compositus, to facilitate the intimate mixture, and to prevent the agglutination, of the ingredients.

[1] $Fe_7Cy_{18}$ or $Fe_4Fe_3Cy_{18}$. This compound, *plus* $18H_2O$, constitutes Prussian Blue.

[2] $K_2Fe_2Cy_6$.

[3] Contact with air causes the ferrous to become a ferric salt by oxidation and the consequent transformation of white potassio-ferrous ferrocyanide into dark blue ferric ferrocyanide.

## PULVIS ANTIMONIALIS

ANTIMONIAL POWDER

**Synonym.**—Factitious James's Powder.

Take of

| | |
|---|---|
| Oxide of Antimony . . . . | 1 ounce |
| Phosphate of Lime . . . . | 2 ounces |

**Actions and Use.**—Similar to those of tartarated antimony, but not so likely to cause local irritation. Employed as a febrifuge for the dog.

**Dose.**—3 to 10 grains.

## PULVIS FERRI SULPHATIS COMPOSITUS

COMPOUND POWDER OF SULPHATE OF IRON

Take of

| | |
|---|---|
| Sulphate of Iron, in powder . | 1 to 1½ drachm |
| Gentian, in powder . . | 4 drachms.[1] |

Mix.

This mixture forms one dose for the horse, to which animal it is given as a tonic, sprinkled over his food.

## PULVIS IPECACUANHÆ COMPOSITUS

COMPOUND POWDER OF IPECACUANHA

**Synonym.**—Factitious Dover's Powder.

Take of

| | |
|---|---|
| Ipecacuanha, in powder . . . | 1 ounce |
| Opium, in powder . . . . | 1 ounce |
| Sulphate of Potash, in powder . | 8 ounces |

[1] If it be thought desirable, from 30 to 40 grains of ground ginger may be added to this powder.

Mix thoroughly, pass the powder through a fine sieve, and finally rub it lightly in a mortar. Keep it in a stoppered bottle.

**Uses.**—Given to the dog as a nauseant and sedative, in from 5- to 15-grain doses, repeated four or five times a day.

## PULVIS OPII COMPOSITUS

### COMPOUND POWDER OF OPIUM

Take of

| | |
|---|---|
| Opium . . . . . | in powder, equal parts |
| Galls . . . . . | |
| Pimento . . . . . | |
| Carbonate of Soda . . | |

Mix.

**Use.**—In diarrhœa occasioned or accompanied by a relaxed or irritable condition of the intestines.

**Doses.**—Horse, 4 to 6 drachms.—Cattle, 4 to 6 drachms.—Sheep, 1 to 1½ drachm.—Pig, 1 to 1½ drachm. Dog, 10 to 30 grains.

**Mode of Application.**—In the form of bolus or mixed with gruel.

## PULVIS RHEI COMPOSITUS

### COMPOUND POWDER OF RHUBARB

**Synonym.**—Gregory's Powder or Gregory's Mixture.

Take of

| | |
|---|---|
| Rhubarb-root, in powder . . . | 2 ounces |
| Ginger, in powder . . . . | 1 ounce |
| Light Magnesia . . . . . | 6 ounces |

Mix thoroughly, and pass the powder through a fine sieve.

**Uses.**—Antacid and stomachic in indigestion.

**Doses.**—Horse, 2 to 3 ounces.—Cattle, 2 to 4 ounces.—Sheep, 1 to 2 drachms.—Pig, 1 to 2 drachms. Dog, 10 to 30 grains.

**Mode of Application.**—In the form of bolus or mixed with gruel.

## PULVIS ZINCI CARBONATIS COMPOSITUS

COMPOUND POWDER OF CARBONATE OF ZINC

Take of

| | |
|---|---|
| Carbonate of Zinc, in powder . . | 4 ounces |
| Alum, in powder . . . . | 1 ounce |

Mix.

**Uses.**—Excitant and astringent when sprinkled over abrasions, ulcerated heels, open joints, &c.

## PYROXYLINUM

PYROXYLIN

**Synonym.**—Dinitro-cellulose.

**Composition.**—$C_{18}H_{21}O_6(OH)(NO_3)_8$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Cotton Wool . . . | 1 ounce |
| Sulphuric Acid . . | of each, 5 fluid ounces |
| Nitric Acid . . . | |

Mix the acids in a porcelain mortar, immerse the cotton

in the mixture, and stir it for three minutes with a glass rod after it is thoroughly wetted with the acids. Transfer the cotton to a vessel containing water, stir it well with a glass rod, decant the liquid, pour more water upon the mass, agitate again, and repeat the affusion, agitation, and decantation until the washings cease to give a precipitate with barium chloride, indicating that the acids have been completely removed. Drain the product on filtering-paper, and dry it in a water-bath.

**Tests.**—Readily soluble in a mixture of ether and rectified spirit; should leave no residue when exploded by heat.

**Preparations.**—Collodium; Collodium Flexile.

## QUERCUS CORTEX

OAK-BARK

The dried bark of the small branches and young stems of *Quercus pedunculata*. Collected in spring from trees growing in Britain.

**Natural Order.**—*Cupuliferæ*.

**Composition.**—Its medicinal activity depends upon its containing from 15 to 20 per cent. of tannic acid with a small quantity of gallic acid.

**Characters.**—Covered with a greenish shining epidermis; cinnamon-coloured on the surface; fibrous, brittle, and strongly astringent.

**Actions and Uses.**—*Internally*:—Astringent and tonic. Employed to arrest diarrhœa, dysentery, and similar mucous discharges. *Externally*:—Astringent,

stypic, and excitant. Occasionally used for stimulating unhealthy wounds and arresting gangrene; also in prolapsus recti and prolapsus uteri.

**Doses.**—Horse, 2 to 6 drachms.—Cattle, 2 to 6 drachms.—Sheep, ½ to 2 drachms.—Pig, ½ to 2 drachms. Dog, 10 to 30 grains.

**Modes of Application.**—*Internally*:—The powdered bark made into a bolus or mixed with gruel, but preferably in the form of Decoctum Quercûs. *Externally*:—The powder or Decoctum Quercûs.

## QUININÆ SULPHAS

### QUININE SULPHATE

**Synonyms.**—Quinia Sulphate; Disulphate of Quinine.

**Composition.**—$[(C_{20}H_{24}N_2O_2)H_2SO_4]_2 15H_2O$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Yellow Cinchona Bark, in coarse powder | 1 pound |
| Hydrochloric Acid | 3 fluid ounces |
| Distilled Water | a sufficiency |
| Solution of Soda | 4 pints |
| Dilute Sulphuric Acid | a sufficiency |

Dilute the hydrochloric acid with ten pints of the water. Place the cinchona-bark in a porcelain basin, and add to it as much of the dilute hydrochloric acid as will render it thoroughly moist. After maceration, with occasional stirring, for twenty-four hours, place the bark in a displacement apparatus, and percolate with the dilute hydrochloric acid until the solution which drops

through is nearly destitute of bitter taste. Into this liquid pour the solution of soda, agitate well, let the precipitate (impure quinine) completely subside, decant the supernatant fluid, collect the precipitate on a filter, and wash it with cold distilled water until the washings cease to have colour. Transfer the precipitate to a porcelain dish containing a pint of distilled water, and, applying to this the heat of a water-bath, gradually add dilute sulphuric acid until very nearly the whole of the precipitate has been dissolved and a neutral liquid has been obtained.

Filter the solution while hot through paper, wash the filter with boiling distilled water, concentrate the filtrate till a film forms upon its surface, and set it aside to crystallise.

Dry the crystals (quinine sulphate) on filtering-paper without the application of heat.

By digestion with dilute hydrochloric acid the bark is exhausted of quinine, cinchonine, and other alkaloids, which are precipitated on the addition of soda to the solution. The precipitated alkaloids, on being dissolved in dilute sulphuric acid, are converted into sulphates, and their aqueous solution, when evaporated till a film forms upon its surface and allowed to cool, deposits the less soluble quinine sulphate, while the more soluble cinchonine sulphate remains in the mother-liquor.

**Characters and Tests.** — Filiform, silky, snowy-white [1] crystals, of an intensely bitter taste, sparingly soluble in water, yet imparting to it a peculiar bluish fluorescence. The solution gives with barium chloride a white precipitate (barium sulphate) insoluble in nitric

[1] Herring's 'Hospital Sulphate of Quinine' is less costly than this preparation, owing to its not having been deprived of colour, and may be substituted for it in veterinary medicine.

acid, showing that the salt is a sulphate, and when treated first with solution of chlorine and afterwards with ammonia, it assumes a splendid emerald-green colour, highly characteristic of quinine.

Should dissolve in pure sulphuric acid with a feeble yellowish tint, and undergo no further change of colour when gently warmed. Ten grains, with ten minims of dilute sulphuric acid and half a fluid ounce of water, form a perfect solution, from which ammonia throws down a white precipitate (quinine). This should re-dissolve on agitating the whole of it with half a fluid ounce of ether, without leaving any crystalline matter (cinchonine) floating on the lower of the two strata into which the agitated fluid separates on rest. Twenty-five grains of the salt should lose 3·6 grains of water by drying at 212° F.

**Actions and Uses.**—Tonic, antipyretic, and anti-periodic in intermittent and other fevers; in loss of appetite and weak digestion depending on debility; in acute rheumatism, chorea in dogs, and constitutional or deep-seated ophthalmia in horses; during convalescence from acute complaints; also during the exhaustion which follows influenza and other diseases.

**Doses.**—Horse, 20 to 40 grains.—Cattle, 20 to 60 grains.—Sheep, 5 to 10 grains.—Pig, 5 to 10 grains.—Dog, 1 to 5 grains.

**Mode of Application.** — In the form of bolus or mixed with gruel.

**Incompatibles.** — Alkalies and their carbonates; infusions of vegetable astringents precipitate quinia tannate especially in the presence of sulphuric acid.

## RESINA

RESIN

**Synonym.**—Yellow Resin.

The residue obtained by distilling the turpentines from various species of *Pinus*, and withdrawing the heat before the whole of the volatile oil is expelled.

**Composition.**—A mixture of abietic (sylvic) acid, which is crystallisable, and pinic acid, which is amorphous.

**Characters.** — Translucent, yellowish, brittle, pulverisable; fracture shining; odour and taste faintly terebinthinate. When heated it easily fuses, and burns with a dense yellow flame and much smoke. Soluble in rectified spirit, insoluble in water.

**Actions and Uses.** — *Internally*: — Diuretic. *Externally*:—Mild excitant, astringent, and styptic. Also employed for imparting consistency and adhesiveness to plasters.

**Doses.**—Horse, 4 to 6 drachms.—Cattle, 4 to 6 drachms.—Sheep, 1 to 2 drachms. Pig, 1 to 2 drachms.—Dog, 20 to 30 grains.

**Modes of Application.** — *Internally*: — Resin is principally given to the horse. It is administered in the form of bolus, preferably as Massa Resinæ Composita. *Externally*:—The powder sprinkled over wounds. To arrest hæmorrhage after castration a small quantity of the powder is sometimes applied to the ends of the spermatic cord and melted by a hot iron.

**Preparations.**—Massa Resinæ Composita; Emplastrum Picis; Unguentum Resinæ.

## RHEI RADIX

RHUBARB-ROOT

The dried rhizome, or so-called root, of *Rheum officinale*, *Rheum palmatum*, and other undetermined species of *Rheum*. Grows in China, Chinese Tartary, and Thibet. Imported from Shanghai and Canton, and brought overland by way of Moscow.

**Natural Order.**—*Polygonaceæ*.

**Composition.**—The root contains a number of principles, including chrysophanic acid, erythroretin, phæoretin, and aporetin. At present it is not known whether the physiological effects of rhubarb are due to one particular constituent or to the combined action of several of them.

**Characters.**—Trapezoidal, roundish, cylindrical, or flattish pieces, frequently bored with one hole; yellow externally, internally marbled with fine waving greyish and reddish lines; finely gritty under the teeth; taste bitter, faintly astringent, and aromatic; odour peculiar. Should be free from decay and not worm-eaten. Boric acid does not turn rhubarb brown unless it is adulterated with turmeric.

**Actions and Uses.**—But little used in veterinary medicine. Chiefly given as a tonic to young animals with weak digestions, and to the dog as a cathartic.

**Doses.**—*Tonic*:—Horse, 1 to 1½ ounce.—Cattle, 1 to 2 ounces.—Sheep, 2 to 4 drachms.—Pig, 1 to 1½ drachm. Dog, 10 to 30 grains. *Cathartic*:—Dog, 2 to 3 drachms.

**Mode of Application.** — The powdered root made into a bolus or mixed with gruel.

**Preparations.**—Pulvis Rhei Compositus; Tinctura Rhei.

## ROSÆ GALLICÆ PETALA

### RED ROSE PETALS

The fresh and dried unexpanded petals of *Rosa gallica*. From plants cultivated in Britain.

**Natural Order.**—*Rosaceæ.*

**Characters.**—Colour fine purplish red, retained after drying; taste bitterish, feebly acid, and astringent; odour roseate (owing to presence of volatile oil), developed by drying.

**Use.**—In preparing Confectio Rosæ Gallicæ.

## SABINÆ CACUMINA

### SAVIN TOPS

The fresh and dried tops of *Juniperus Sabina*. Collected in spring from plants cultivated in Britain.

**Natural Order.**—*Coniferæ.*

**Composition.**—Medicinal properties are due to a volatile oil having the formula $C_{10}H_{16}$.

**Characters.**—Twigs densely covered with minute imbricated appressed leaves in four rows; odour strong, peculiar, and unpleasant; taste acrid, bitter, resinous, and disagreeable.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant poison; in medicinal doses, anthelmintic, but is

very seldom employed for this purpose, on account of its violent and uncertain irritant qualities. *Externally*:—Stimulant to ill-conditioned wounds and indolent sores; also used for maintaining the action of, and keeping up the discharge from, blisters and setons.

**Modes of Application.**—*Externally*:—Infusum Sabinæ and, preferably, Unguentum Sabinæ.

**Preparation.**—Unguentum Sabinæ.

## SACCHARUM PURIFICATUM

REFINED SUGAR

Pure cane sugar prepared from the juice of the stem of *Saccharum officinarum*, cultivated in the West Indies and other tropical countries.

**Natural Order.**—*Graminaceæ.*

**Composition.**—$C_{12}H_{22}O_{11}$.

**Characters.**—Compact, crystalline, conical loaves, known in commerce as lump sugar.

**Preparations.**—Confectio Rosæ Gallicæ; Ferri Carbonas Saccharata.

## SANTONINUM

SANTONIN

A crystalline neutral principle obtained from santonica, the dried unexpanded flower-heads of *Artemisia Santonica*, imported from Russia. It is the only so-called worm-seed which yields santonin, a crystalline neutral principle, in quantity worth extracting.

**Natural Order.**—*Compositæ.*

**Composition.**—$C_{15}H_{18}O_3$.

**Characters and Tests.**—In colourless, flat, rhombic prisms, feebly bitter, sparingly soluble in water. Solubility in rectified spirit 1 in 50. Not dissolved by dilute mineral acids. Entirely destructible by a red heat with free access of air.

**Actions and Uses.**—*Internally*:—Anthelmintic, useful both for tape- and thread-worms.

**Doses.**—Horse, 20 to 40 grains.—Dog, 2 to 10 grains.

## SAPO DURUS

HARD SOAP

**Composition.**—Chiefly sodium oleate,[1] $NaC_{18}H_{33}O_2$, with a large percentage of water; contains about 30 per cent. of glycerin.

**Mode of Preparation.**—Olive oil[1] is boiled with caustic soda (sodium hydrate); glycerin passes into solution, and sodium oleate (hard soap) floats on the surface of the liquid. After remaining quiescent for a few hours the soap is skimmed off, transferred to 'frames,' in which it solidifies on cooling, and is finally cut by wires into slabs or bars.

| Olive Oil (Triolein). | | Sodium Hydrate. | | Sodium Oleate. | | Glycerin. |
|---|---|---|---|---|---|---|
| $C_3H_5(C_{18}H_{33}O_2)_3$ | + | $3NaHO$ | = | $3NaC_{18}H_{33}O_2$ | + | $H_3C_3H_5O_3$ |

**Characters.**—Greyish white, dry, horny, and pulverisable when kept in dry warm air; easily moulded

[1] Although the 'British Pharmacopœia' directs hard soap to be made from olive oil, the best 'curd soap' of commerce, which is a mixture of sodium stearate, palmitate, and oleate, and which is prepared from various solid and liquid fats, may be substituted for it in practice.

when heated. Soluble in rectified spirit, producing a solution which, when evaporated on paper, does not leave an oily stain. Incinerated, it yields an ash which does not deliquesce.

**Actions and Uses.**—*Internally* :—Cathartic and diuretic, but chiefly employed as an antacid, as an antidote to poisoning by the mineral acids, and in the preparation of clysters. *Externally* :—Stimulant to sprains and contusions; also applied to burns and scalds to diminish the irritation and pain by excluding air. Likewise employed as an excipient in making up boluses, liniments, and ointments.

**Doses.**—Horse, 1 to 2 ounces.—Cattle, 1 to 2 ounces. Sheep, 2 to 6 drachms.—Pig, 2 to 6 drachms.—Dog, 20 to 60 grains.

**Mode of Application.**—In the form of bolus or beaten up with water.

**Incompatibles.**—Acids and most metallic salts.

## SAPO MOLLIS

SOFT SOAP

**Composition.** — Principally potassium oleate,[1] $KC_{18}H_{33}O_2$, with glycerin, free potassium hydrate, and a large percentage of water.

**Mode of Preparation.**—By boiling together olive oil[1] and caustic potash (potassium hydroxide) until the soap produced assumes the appearance of a transparent

[1] Olive oil is directed by the 'British Pharmacopœia' to be employed in the preparation of soft soap; but the commercial article, which is the material used by the veterinary surgeon, is made from tallow, also from whale, linseed, and other oils.

jelly, and until a drop of it, when allowed to fall on a glass plate, remains clear, and does not exhibit a fatty border on cooling.

| Olive Oil (Triolein). | | Potassium Hydrate. | | Potassium Oleate. | | Glycerin. |
|---|---|---|---|---|---|---|
| $C_3H_5(C_{18}H_{33}O_2)_3$ | + | $3KHO$ | = | $3KC_{18}H_{33}O_2$ | + | $H_3C_3H_5O_3$ |

**Use.**—In the preparation of Linimentum Saponis Compositum.

**Preparation.**—Linimentum Saponis Compositum.

## SCILLA

SQUILL

The sliced and dried bulb of *Urginea Scilla.*

**Natural Order.**—*Liliaceæ.*

**Composition.**—It contains scillipicrin, scillitosin, and scillin. The active principle scillitosin is soluble in water, acetic acid, and alcohols.

**Actions and Uses.**—Principally used in catarrhal and bronchial affections.

**Doses** (of the Syrup).—Horse, 1½ ounce.—Dog, 10 to 60 minims.

**Preparation.**—Syrupus Scillæ.

## SEVUM PRÆPARATUM

PREPARED SUET

The internal fat of the abdomen of the sheep, *Ovis Aries,* purified by melting and straining.

**Composition.**—A mixture of stearin, palmitin, and olein.

**Characters.**—White, smooth, and almost scentless: fusible at 103° F.

**Uses.**—Sometimes employed as a substitute for, or in conjunction with, lard, in the preparation of ointments.

## SINAPIS

### MUSTARD

The seeds of *Brassica nigra* and *Brassica alba*; also the seeds of both reduced to powder and mixed.

**Natural Order.**—*Cruciferæ.*

**Composition.**—The vesicating power of mustard, and the pungent odour emitted by it when moistened, depend upon a volatile oil (allyl sulphocyanide, $C_3H_5CNS$). It does not exist ready formed in the seed, but is developed by the mutual action of two of its proximate principles—viz., myrosin, a kind of ferment, and potassium myronate—in the presence of hot, but not boiling, water.[1]

| Potassium Myronate. | | Allyl Sulphocyanide. | | Glucose. | | Potassium Sulphate. |
|---|---|---|---|---|---|---|
| $KC_{10}H_{18}NS_2O_{10}$ | = | $C_3H_5CNS$ | + | $C_6H_{12}O_6$ | + | $KHSO_4$ |

**Actions and Uses.**—Vesicant and counter-irritant, in the form of poultice.

**Preparations.** — Cataplasma Sinapis; Cataplasma Sinapis Ammoniatum.

## SODA CAUSTICA

### CAUSTIC SODA

**Synonyms.**—Sodium Hydrate; Hydrate of Sodium; Hydrate of Soda.

[1] Boiling water coagulates myrosin, and destroys its power as a ferment.

**Composition.**—NaOH.

**Mode of Preparation.**—Take of

Solution of Soda . . . . 2 pints

Boil down the solution of soda rapidly in a silver or clean iron vessel until there remains a fluid of oily consistency, a drop of which, when removed on a warmed glass rod, solidifies on cooling. Pour the fluid on to a clean silver or iron plate, or into moulds, and as soon as it has solidified break it into pieces, and preserve it in stoppered green-glass bottles.

**Characters and Tests.**—Hard and greyish white, very alkaline and corrosive. Does not deliquesce on exposure to the atmosphere. It imparts a yellow colour to flame, indicating the presence of sodium, and its solution in water acidulated by nitric acid should give only scanty white precipitates with silver nitrate and barium chloride, showing the presence of traces only of chlorides and sulphates respectively.

**Action and Use.**—*Externally* :—As a caustic. It is equal in activity to Potassa Caustica, but preferable to this agent, as it does not deliquesce and spread.

## SODII BICARBONAS

### SODIUM BICARBONATE

**Synonym.**—Sodium-hydrogen Carbonate.

**Composition.**—$NaHCO_3$.

**Mode of Preparation.** — Carbonic anhydride [1] is passed into a vessel containing a mixture of two parts of

[1] For mode of generating this gas, see Potassii Bicarbonas.

crystallised and three parts of dried sodium carbonate until the gas is no longer absorbed. The damp salt which is formed is shaken occasionally during half an hour with half its weight of cold distilled water, for the purpose of dissolving any unaltered carbonate, and the remaining insoluble bicarbonate is dried by exposure to air on filtering-paper placed on porous bricks. The object of using the crystallised carbonate is that it may conveniently furnish the necessary amount of water for the success of the process.

| Sodium Carbonate. | | Carbonic Anhydride. | | Water. | | Sodium-hydrogen Carbonate. |
|---|---|---|---|---|---|---|
| $Na_2CO_3$ | + | $CO_2$ | + | $H_2O$ | = | $2NaHCO_3$ |

**Characters and Tests.** — In powder, or small opaque monoclinic crystals, white; saline unpleasant taste. Imparts a yellow colour to flame, showing the presence of sodium. Soluble in water 1 part in 10. Dissolves with effervescence in dilute hydrochloric acid, showing that it is a carbonate; and the hydrochloric-acid solution gives no precipitate with platinic chloride, indicating that it is neither a potassium nor an ammonium salt. Its aqueous solution acidified with nitric acid should give but faint indications of sulphates and chlorides on the addition of barium chloride and silver nitrate. A solution of the salt in cold water gives a white, and not a coloured, precipitate with mercuric chloride, unless sodium carbonate be present. Distinguished from sodium carbonate by its aqueous solution not giving a precipitate with magnesium sulphate.

**Actions and Uses.**—Similar to those of Potassii Bicarbonas. Less alkaline and milder than Sodii Carbonas.

**Doses.**—Horse, 4 to 6 drachms.—Cattle, 4 to 8 drachms.—Sheep, 1 to 2 drachms.—Pig, 1 to 2 drachms. Dog, 5 to 40 grains.

**Modes of Application.** } Same as Potassii Bicar-
**Incompatibles.** } bonas.

## SODII CARBONAS

### SODIUM CARBONATE

**Synonyms.**—Carbonate of Soda ; Washing-soda.

**Composition.**—$Na_2CO_3.10H_2O$.

**Mode of Preparation.**—Sodium chloride (common salt) is decomposed by sulphuric acid, whereby sodium sulphate ('salt cake') and hydrochloric acid are produced; the former remains on the hearth, while the latter escapes by the chimney, of the furnace in which the operation is conducted.

| Sodium Chloride. | | Sulphuric Acid. | | Sodium Sulphate. | | Hydrochloric Acid. |
|---|---|---|---|---|---|---|
| $2NaCl$ | + | $H_2SO_4$ | = | $Na_2SO_4$ | + | $2HCl$ |

The sodium sulphate is next heated with a mixture of carbon (coal-dust) and calcium carbonate (chalk or limestone), when the following changes take place:

| | Sodium Sulphate. | | Carbon. | | Sodium Sulphide. | | Carbonic Oxide. |
|---|---|---|---|---|---|---|---|
| 1. | $Na_2SO_4$ | + | $4C$ | = | $Na_2S$ | + | $4CO$ |

| | Sodium Sulphide. | | Calcium Carbonate. | | Sodium Carbonate. | | Calcium Sulphide. |
|---|---|---|---|---|---|---|---|
| 2. | $Na_2S$ | + | $CaCO_3$ | = | $Na_2CO_3$ | + | $CaS$ |

The resulting mixture of sodium carbonate and calcium sulphide ('black ash' or 'ball soda'), when cold, is broken up into little pieces and lixiviated with water; the sodium carbonate passes into solution, and the

calcium sulphide is undissolved. By evaporating to dryness the solution thus formed, 'soda ash' is procured, and by dissolving 'soda ash' in water, filtering and evaporating the solution, and then allowing it to cool, sodium carbonate is obtained in the crystallised state.

**Characters and Tests** — Transparent, colourless, laminar, rhombic crystals, effloresecent; harsh alkaline taste, and strong alkaline reaction. It imparts a yellow colour to flame, showing the presence of sodium, and dissolves with effervescence in dilute hydrochloric acid, indicating that it is a carbonate. Soluble in cold water 1 in 2, insoluble in rectified spirit. The hydrochloric-acid solution gives no precipitate with platinic chloride, showing that it is neither a potassium nor an ammonium salt. Its aqueous solution gives a white precipitate (magnesium carbonate) with magnesium sulphate and a red precipitate with solution of mercuric chloride, reactions by which it is distinguished from sodii bicarbonas. Strongly heated, it undergoes the aqueous fusion, and then dries up, losing water to the extent of 63 per cent. of its weight. When its aqueous solution is treated with excess of nitric acid, it should yield but slight precipitates with barium chloride and silver nitrate, indicating the almost total absence of sulphates and chlorides.

**Actions and Uses.** — Very similar to those of Potassii Bicarbonas, Potassii Carbonas, and Sodii Bicarbonas.

**Doses.**—Half those of Sodii Bicarbonas.

**Mode of Application.** } Same as Potassii Bicar-
**Incompatibles.** } bonas.

**Preparations.**—Liquor Sodæ; Sodii Bicarbonas; Sodii Carbonas Exsiccata.

## SODII CARBONAS EXSICCATA

DRIED SODIUM CARBONATE

**Composition.**—$Na_2CO_3$.

**Mode of Preparation.**—Crystallised sodium carbonate is heated in a porcelain capsule until the liquid which first forms is converted into a dry cake. This cake, when pulverised, is to be preserved in a stoppered bottle.

**Actions and Uses.**—Similar to those of the potassium and sodium carbonates and bicarbonates.

**Doses.**—About one fourth of those of Sodii Bicarbonas.

**Modes of Application.** } Same as Potassii Bicar-
**Incompatibles.** } bonas.

## SODII HYPOSULPHIS

SODIUM HYPOSULPHITE

**Synonyms.**—Sodium Thiosulphate ; Antichlor.

**Composition.**—$Na_2S_2O_3.10H_2O$.

**Mode of Preparation.**—An aqueous solution of sodium sulphite is boiled with flowers of sulphur and filtered. On evaporating the filtrate the salt crystallises out. The following equation explains its formation :

| Sodium Sulphite. | | Sulphur. | | Sodium Hyposulphite. |
|---|---|---|---|---|
| $2Na_2SO_3$ | + | $S_2$ | = | $2Na_2S_2O_3$ |

During crystallisation the salt acquires five molecules of water.

**Characters and Tests.**—Rhombic prisms, soluble in water. Imparts a yellow colour to flame, indicating

the presence of sodium. Dilute hydrochloric acid added to its aqueous solution causes effervescence (due to the expulsion of sulphurous anhydride, recognisable by its odour), and the solution is, at the same time, rendered turbid by the precipitation of sulphur. The latter reaction distinguishes this salt from sodium sulphite.

**Actions and Uses.**
**Doses.**
**Incompatibles.**
} Same as Sodii Sulphis.

## SODII SALICYLAS

### SODIUM SALICYLATE

**Composition.**—$(C_6H_4.OH.COONa)_2.H_2O$.

**Mode of Preparation.**—Salicylate of Sodium is prepared by the action of salicylic acid on either sodium carbonate or sodium hydroxide.

**Characters and Tests.**—Small, colourless or nearly colourless, crystalline, lustrous scales; inodorous, and has a sweet saline taste. It is soluble in less than its own weight of cold water, and in six parts of alcohol (90-per-cent.); dissolves in cold sulphuric acid without coloration or effervescence.

The solutions are neutral or faintly acid.

Perchloride of iron gives, with a concentrated solution, a reddish-brown colour; in dilute solutions a violet colour.

**Actions and Uses.**—Similar to salicylic acid.[1] It is antiseptic, antiperiodic, antipyretic, irritant, and astrin-

[1] *Dithion* or Di-thio-salicylate of sodium, a greyish-white powder freely soluble in water, is antiseptic and antipyretic. In foot-and-mouth disease it has been given internally, and applied externally as a dusting-powder and as a lotion (2½ to 5 per cent.) to the sores of the feet, mouth and udder ('Lancet,' 1892).

gent, and is especially of value in the treatment of rheumatism.

**Dose.**—Horse, 1 to 5 drachms.—Cattle, 2 to 7 drachms.—Dog, 5 to 20 grains.

## SODII SULPHAS

### SODIUM SULPHATE

**Synonyms.**—Sulphate of Soda; Glauber's Salt.

**Composition.**—$Na_2SO_4.10H_2O$.

**Mode of Preparation.**—By neutralising the residue (sodium-hydrogen sulphate) left in the manufacture of hydrochloric acid with sodium carbonate, and crystallising the product from solution in water.

| Sodium-hydrogen Sulphate. | | Sodium Carbonate. | | Sodium Sulphate. | | Carbonic Anhydride. | | Water. |
|---|---|---|---|---|---|---|---|---|
| $2NaHSO_4$ | + | $Na_2CO_3$ | = | $2Na_2SO_4$ | + | $CO_2$ | + | $H_2O$ |

**Characters and Tests.**—Transparent oblique prisms; has a saline and bitter taste; effloresces on exposure to the air; soluble in water, 1 in 3. Heated in a porcelain crucible, it loses 55·9 per cent of water. It imparts a yellow colour to flame, indicating the presence of sodium. With barium chloride its aqueous solution furnishes a white precipitate (barium sulphate) insoluble in nitric acid, showing that the salt is a sulphate. Its aqueous solution mixed with dilute hydrochloric acid gives no precipitate with platinic chloride, proving that it is neither a potassium nor an ammonium salt.

**Actions and Uses.**—Cathartic, diuretic, and resolvent. Magnesii sulphas being more certain in its action

than sodii sulphas, the latter is not much employed by the veterinarian, except occasionally as a purgative for cattle.

**Dose.**—Purgative for cattle, from 1 to 1½ pound.

**Mode of Application.**—Dissolved in water or mixed with ginger and treacle.

## SODII SULPHIS

### SODIUM SULPHITE

**Synonym.**—Sulphite of Soda.

**Composition.**—$Na_2SO_3.7H_2O$.

**Mode of Preparation.**—An aqueous solution of sodium carbonate is saturated with sulphurous anhydride; sodium-hydrogen sulphite is formed and remains in solution, while carbonic anhydride escapes in the gaseous state.

| Sodium Carbonate. | | Sulphurous Anhydride. | | Water. | | Sodium-hydrogen Sulphite. | | Carbonic Anhydride. |
|---|---|---|---|---|---|---|---|---|
| $Na_2CO_3$ | + | $2SO_2$ | + | $H_2O$ | = | $2NaHSO_3$ | + | $CO_2$ |

The solution of sodium-hydrogen sulphite is mixed while warm with as much sodium carbonate as was used in the first operation; on cooling, sodium sulphite crystallises out. The crystals, after being drained, are dried on filtering-paper.

| Sodium-hydrogen Sulphite. | | Sodium Carbonate. | | Sodium Sulphite. | | Carbonic Anhydride. | | Water. |
|---|---|---|---|---|---|---|---|---|
| $2NaHSO_3$ | + | $Na_2CO_3$ | = | $2Na_2SO_3$ | + | $CO_2$ | + | $H_2O$ |

In the act of crystallising the salt takes up seven molecules of water.

**Characters and Tests.**—Sometimes in transparent prismatic crystals, but usually in white opaque masses. Soluble in water. It imparts a yellow colour to flame, showing the presence of sodium. On adding dilute hydrochloric acid to its aqueous solution, sulphurous anhydride (recognisable by its odour) is evolved with effervescence, and the solution remains bright and transparent.

**Actions and Uses.**—*Internally*:—Antiseptic, disfectant, and alterative. Given in indigestion, tympanites, and in so-called blood-diseases. *Externally*:—Antiseptic, deodoriser, and disinfectant. Employed in the treatment of ulcerated sore throat and phagedænic wounds and ulcers.

**Doses.**—Horse, 1 to 1½ ounce.—Cattle, 2 to 4 ounces. Sheep, 2 to 6 drachms.—Pig, 2 to 6 drachms.—Dog, 20 to 60 grains. Frequently repeated.

**Modes of Application.**—*Internally*:—Dissolved in water. *Externally*:—Three to four drachms dissolved in every ounce of water employed.

**Incompatibles.**—Acids and most metallic salts.

## SODII CHLORIDUM

### CHLORIDE OF SODIUM

**Synonyms.**—Sodium Chloride; Common Salt.

**Composition.**—NaCl.

**Mode of Preparation.**—Crude 'rock salt,' or a similar product obtained by evaporating sea-water or the water of brine springs, is purified by solution and crystallisation.

**Characters and Tests.**—Small white crystalline grains, or transparent cubic crystals, free from moisture; purely saline taste; soluble in water. Its aqueous solution mixed with hydrochloric acid and platinic chloride yields no precipitate, showing that it is neither an ammonium nor a potassium salt. It imparts a yellow colour to flame, indicating that sodium is present; and its aqueous solution gives with silver nitrate a white precipitate (silver chloride) soluble in ammonia, but insoluble in boiling nitric acid, proving the salt to be a chloride.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant poison; in medicinal doses, cathartic, anthelmintic, emetic, diuretic, tonic, alterative, and resolvent. To cattle and sheep it is given as an anthelmintic; also as a purgative in diarrhœa caused by the presence of an irritant or by over-feeding, in fardel-bound, and in distension of the stomach by food. To sheep it is likewise given as a preventive of rot, for the destruction of intestinal worms, and as a corrective of the bad effects of damp and badly kept fodder. It is employed as an emetic for the dog, and is given to all animals, in frequently repeated doses, as a tonic and stimulant in impaired digestion. Rock salt should always be kept as a provocative of the appetite in the horse's manger, the ox's crib, and the sheep's trough. *Externally*:—As a stimulant in chronic sprains, many affections of the joints, and in some diseases of the feet, particularly amongst cattle and sheep. Common salt is frequently added to laxative enemas.

**Doses.**—*Anthelmintic and Cathartic*:—Cattle, 12 to 24 ounces.—Sheep, 1 to 3 ounces. *Emetic*:—Dog, 1 to 2 drachms. *Tonic*:—Horse, 1 to 2 ounces.—Cattle, 2 to 4 ounces.—Sheep, 2 to 4 drachms.—Dog, 10 to 30 grains.

**Modes of Application.**—*Internally* :—As a cathartic and emetic dissolved in tepid water. Sometimes half-doses of chloride of sodium and sulphate of magnesia are mixed with ginger, pimento, or some other aromatic, and treacle; such a mixture will often be voluntarily drunk by cattle and sheep. Croton oil, oil of turpentine, and calomel may be added to common salt to increase its carthartic and anthelmintic action ; and mustard, sulphate of zinc, sulphate of copper, or tartarated antimony may be combined with it to ensure its prompt action as an emetic. As a tonic and stomachic, it should be sprinkled over the animal's food, either in the state of powder or in solution. *Externally* :—A lotion consisting of one pound of common salt dissolved in a gallon of cold water. This lotion should be applied during, or immediately after, the solution of the salt.

## SPIRITUS ÆTHERIS

SPIRIT OF ETHER

**Synonym.**—Spiritus Ætheris Sulphurici.

**Composition.**—A mixture of sulphuric ether with twice its bulk of rectified spirit.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Ether . . . . . | 10 fluid ounces |
| Alcohol (90-per-cent.) . . | 1 pint |

Mix.

**Test.**—Sp. gr. 0·806 to 0·811.

**Actions and Uses.**—Diffusible stimulant; anti-spasmodic.

**Doses.**—Horse, 3 to 6 fluid ounces.—Cattle, 5 to 10 fluid ounces.—Sheep, 1 to 2 fluid ounces.—Pig, 1 to 2 fluid ounces.—Dog, 1 to 5 drachms.

## SPIRITUS ÆTHERIS NITROSI

SPIRIT OF NITROUS ETHER

**Synonym.**—Sweet Spirit of Nitre.

**Composition.**—An alcoholic solution of nitrous ether (ethyl nitrite), $C_2H_5NO_2$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Nitric Acid . . . . | 3 fluid ounces |
| Sulphuric Acid . . . | 2 fluid ounces |
| Copper in fine wire (about No. 25) . . . . | 2 ounces |
| Alcohol (90-per-cent.) . . | a sufficiency |

To one pint of the spirit add gradually the sulphuric acid, and stir them together ; then add, in the same way, two and a half fluid ounces of the nitric acid. Put the mixture into a retort or other suitable apparatus, into which the copper has been introduced, and to which a thermometer has been fitted. Attach an efficient condenser, and, applying a gentle heat, let the spirit distil at a temperature commencing at 170° F., and rising to 175°, but not exceeding 180°, until twelve fluid ounces have passed over and been collected in a bottle kept cool, if necessary, with ice-cold water ; then withdraw the heat and, having allowed the contents of the retort to cool, introduce the remaining half-ounce of nitric acid, and resume the distillation as before, until the distilled product (spirit of nitrous ether) has been increased to fifteen fluid ounces. Mix this with two pints of the rectified spirit, or as much as will make it correspond to the tests of specific gravity and percentage of ether separated by calcium chloride. Preserve it in well-stoppered bottles.

The following equation explains the principal changes which probably take place in the foregoing process:

| Alcohol. | | Sulphuric Acid. | | Nitric Acid. | | Copper. |
|---|---|---|---|---|---|---|
| $2C_2H_5HO$ | + | $2H_2SO_4$ | + | $2HNO_3$ | + | $Cu_2$ |

| | Ethyl Nitrite. | | Cupric Sulphate. | | Water. |
|---|---|---|---|---|---|
| = | $2C_2H_5NO_2$ | + | $2CuSO_4$ | + | $4H_2O$ |

The ethyl nitrite distils over, accompanied by alcohol and other bodies of minor importance.

**Characters and Tests.**—Transparent and nearly colourless, but with a very slight tinge of yellow; mobile, inflammable; peculiar penetrating apple-like odour, and sweetish cooling, sharp taste. Sp. gr. 0·845. It effervesces feebly or not at all when shaken with a little sodium-hydrogen carbonate, indicating the presence of a trace or the absence of free acid. When agitated with solution of ferrous sulphate and a few drops of sulphuric acid, it becomes deep olive-brown or black. Agitated with twice its volume of a saturated solution of calcium chloride (to absorb water and alcohol) in a closed graduated tube, 2 per cent. of its original volume should separate in the form of ethyl nitrite and rise to the surface of the mixture.

**Actions and Uses.**—In excessive doses, narcotic and sedative; in medicinal doses, stimulant, antispasmodic, diuretic, and diaphoretic. Given in colic, indigestion, tympanites, and local congestions; during convalescence from debilitating diseases; also in combination with tincture of opium for diminishing the pains and spasms which sometimes follow parturition in cows.

**Doses.**—Horse, 1 to 2 fluid ounces.—Cattle, 3 to 4 fluid ounces.—Sheep, 3 to 6 fluid drachms.—Pig, 3 to

6 fluid drachms.—Dog, ½ to 2 fluid drachms. Frequently repeated.

**Mode of Application.**—Mixed with sufficient cold water to make a draught.

## SPIRITUS AMMONIÆ AROMATICUS[1]

AROMATIC SPIRIT OF AMMONIA

Take of

| | |
|---|---|
| Carbonate of Ammonia | 8 ounces |
| Strong Solution of Ammonia | 4 fluid ounces |
| Volatile Oil of Nutmeg | 4 fluid drachms |
| Oil of Lemon | 6 fluid drachms |
| Rectified Spirit | 6 pints |
| Water | 3 pints |

Mix and distil seven pints.

**Test.**—Sp. gr. 0·870.

**Actions and Uses.**—Diffusible stimulant, antacid, and antispasmodic.

**Doses.**—Horse, ½ to 1 fluid ounce.—Cattle, ½ to 2 fluid ounces.—Sheep, 2 to 4 fluid drachms.—Pig, 2 to 4 fluid drachms.—Dog, 10 to 20 minims. Mixed with water.

## SPIRITUS AMMONIÆ FŒTIDUS

FETID SPIRIT OF AMMONIA

Take of

| | |
|---|---|
| Asafetida | 1½ ounce |
| Strong Solution of Ammonia | 2 fluid ounces |
| Alcohol (90-per-cent.) | a sufficiency |

[1] This preparation is stronger in spirit, and about one-half stronger in ammonia, than the Spiritus Ammoniæ Aromaticus of the 'British Pharmacopœia.'

Break the asafetida into small pieces, and macerate it in a closed vessel, in fifteen fluid ounces of the spirit, for twenty-four hours; then distil off the spirit, mix the product with the solution of ammonia, and add sufficient rectified spirit to make one pint.

**Actions and Uses.**—Stimulant and antispasmodic.

**Doses.**—Horse, ½ to 1 fluid ounce.—Cattle, ½ to 2 fluid ounces.—Sheep, 2 to 4 fluid drachms.—Pig, 2 to 4 fluid drachms.—Dog, 10 to 20 minims. Mixed with water.

## SPIRITUS CAMPHORÆ

### SPIRIT OF CAMPHOR

**Synonym.**—Tinctura Camphoræ.

Take of

| | |
|---|---|
| Camphor . . . . . | 1 ounce |
| Alcohol (90-per-cent.) enough to make . . . . . | 10 fluid ounces |

Dissolve.

**Actions and Uses.**—*Externally*:—Stimulant, anodyne, and discutient.

## SPIRITUS CHLOROFORMI

### SPIRIT OF CHLOROFORM

**Synonym.**—Chloric Ether.

Take of

| | |
|---|---|
| Chloroform . . . . | 1 fluid ounce |
| Rectified Spirit . . . . | 19 fluid ounces |

Dissolve.

**Test.**—Sp. gr. 0·871.

**Actions and Uses.**—Stimulant, antispasmodic, and anodyne.

**Doses.**—HORSE, 2 to 4 fluid ounces.—CATTLE, 2 to 4 fluid ounces.—SHEEP, ½ to 1 fluid ounce.—PIG, ½ to 1 fluid ounce.—DOG, 1 to 3 fluid drachms. Mixed with water.

## SPIRITUS HYDRARGYRI PERCHLORIDI COMPOSITUS

### COMPOUND SPIRIT OF PERCHLORIDE OF MERCURY

**Synonym.**—Liquor Hydrargyri Bichloridi.

Take of

| | |
|---|---|
| Perchloride of Mercury, in powder | 1 ounce |
| Hydrochloric Acid | 1 fluid ounce |
| Rectified Spirit | 7 fluid ounces |

Add the acid to the spirit, and dissolve the perchloride of mercury in the mixture.

**Uses.**—*Externally*:—Occasionally to farcy ulcers, and for arresting synovial discharges.

## SPIRITUS METHYLATUS

### METHYLATED SPIRIT

**Composition.**—A mixture consisting of 90 per cent. of spirit of wine and 10 per cent. of methylic alcohol, $CH_3HO$.

**Uses.**—As a substitute for the more expensive spirit of wine in the preparation of tinctures, &c., for *external use only*. The law forbids methylated spirit, or any preparation containing it, to be used internally.

## SPIRITUS RECTIFICATUS

RECTIFIED SPIRIT

**Synonym.**—Spirit of Wine.

**Composition.**—Alcohol (absolute alcohol, $C_2H_5OH$) 90 per cent.

**Mode of Preparation.**—Saccharine fluids are permitted to ferment, whereby the sugar which they contain is transformed into alcohol and carbonic anhydride. By distillation the former is separated from the impurities with which it is mixed, and the latter escapes into the air in the state of gas.

| Grape Sugar (Glucose). | | Alcohol. | | Carbonic Anhydride. |
|---|---|---|---|---|
| $C_6H_{12}O_6$ | = | $C_2H_5OH$ | + | $2CO_2$ |

**Characters and Tests.** — Colourless, transparent, very mobile and inflammable; peculiar pleasant odour, and a strong spirituous, burning taste. Burns with a blue flame, without smoke. Sp. gr. 0·340. Should have a purely alcoholic odour and taste, and should remain clear when diluted with distilled water, indicating its freedom from fusel oil (amylic alcohol, $C_5H_{11}.OH$).

**Actions and Uses.** — *Internally* :—Stimulant, diuretic, and diaphoretic. Alcohol, in the form of ale, brandy, whisky, or gin for the larger animals, and in the form of wine for the smaller ones, is employed as a stimulant in colic and tympanites ; in the latter stages of influenza, bronchitis, and other inflammatory diseases ; also in, and during convalescence from, debilitating complaints. *Externally* :—Rectified spirit is added to cooling lotions to increase their refrigerating power. It is likewise employed in the preparation of medicinal spirits, tinctures, and extracts.

**Doses.**[1]—HORSE, 1 to 3 ounces.—CATTLE, 3 to 6 ounces.—SHEEP, ½ to 1½ ounce.—PIG, ½ to 1½ ounce.—DOG, 1 to 3 drachms. Repeated every two hours.

*Table showing the average amounts by measure of absolute alcohol* ($C_2H_5OH$) *contained in* 100 *volumes of the following alcoholic beverages:*

| Name of Beverage | Absolute Alcohol in 100 volumes |
|---|---|
| Small Beer and Table Ale | 1 to 2 |
| Porter | about 5 |
| Ale (various kinds) | 5 to 9 |
| Sherry | 15 to 17 |
| Port | 17 to 18 |
| Brandy (common) | 36 |
| Gin | 45 |
| Brandy (Cognac) | 46 |
| Rum | 48 |
| Whisky | 50 |

## SPIRITUS TENUIOR

### PROOF SPIRIT

**Composition.**—One hundred parts, at 60° F., consist of forty-nine parts by weight of absolute alcohol, and fifty-one parts by weight of water.

Take of

| | |
|---|---|
| Rectified Spirit | 5 pints |
| Distilled Water | 3 pints |

Mix.

**Test.**—Sp. gr. 0·92.

**Use.**—In the preparation of tinctures.

[1] The doses here enumerated refer to rectified spirit; the determination of those of the ordinary alcoholic beverages must be left to the judgment of the prescriber.

## STAPHISAGRIÆ SEMINA

STAVESACRE-SEEDS

The seeds of *Delphinium Staphisagria.*

**Natural Order.**—*Ranunculaceæ.*

**Composition.**—The seeds contain several alkaloids, the most important being delphinine, delphisine, and delphinoidine. They also contain a fixed oil.

**Actions and Uses.**—A solution made by boiling the seeds in water, one part of seeds to twenty or thirty of water, is used as a parasiticide lotion.

## STRYCHNINA

STRYCHNINE

**Composition.**—$C_{21}H_{22}N_2O_2$. An alkaloid obtained from the dried ripe seeds of *Strychnos Nux-Vomica.*

**Natural Order.**—*Loganiaceæ.*

**Mode of Preparation.**—Take of

| | |
|---|---|
| Nux Vomica . . . . . | 1 pound |
| Acetate of Lead . . . . | 180 grains |
| Solution of Ammonia . . . | a sufficiency |
| Rectified Spirit . . . . | a sufficiency |
| Distilled Water . . . . | a sufficiency |

Soften the nux vomica by subjecting it for two hours to steam in any convenient vessel; chop or slice it; dry it in a water-bath or hot-air chamber, and immediately grind it in a coffee-mill. Digest the powder at a gentle heat for twelve hours with two pints of the spirit and one of the water, to dissolve out the igasurates of strychnine and brucine; strain through linen, express strongly, and

repeat the process twice. Distil off the spirit from the mixed fluid, evaporate the watery residue to about sixteen ounces, and filter when cold. Add now the acetate of lead previously dissolved in distilled water, so long as it occasions any precipitate (lead igasurate); filter; wash the precipitate with ten ounces of cold water, add the washings to the filtrate (containing the acetates of strychnia and brucia); evaporate the clear fluid to eight ounces, and when it has cooled add the ammonia in slight excess, stirring thoroughly. Let the mixture stand, at the ordinary temperature, for twelve hours; collect the precipitate (impure strychnine) on a filter, wash it at once with a few ounces of cold distilled water, dry it in a water-bath or hot-air chamber, and boil it with successive portions of rectified spirit till the fluid scarcely tastes bitter. Distil off most of the spirit, evaporate the residue to the bulk of about half an ounce, and set it aside to cool. Cautiously pour off the yellowish mother-liquor (which contains the brucine of the seeds) from the white crust of strychnine which adheres to the vessel. Throw the crust on a paper filter, wash it with a mixture of two parts of rectified spirit and one of water till the washing ceases to become red on the addition of nitric acid, showing that the brucine has been removed from the strychnia; finally, dissolve it (the strychnia) by boiling it with an ounce of rectified spirit, and set it aside to crystallise. More crystals of strychnine may be obtained by evaporating the mother-liquor.

**Characters and Tests.**—Trimetric prisms, colourless and inodorous; sparingly soluble in water, but communicating to it an intensely bitter taste; soluble in boiling rectified spirit, in chloroform, and in commercial ether, but not in absolute alcohol or ether. Pure sulphuric acid forms with it a colourless solution, which on

the addition of solid potassium bichromate acquires a purple hue, changes to yellowish-red, and ultimately leaves a green residue. Should not be coloured by nitric acid; if it turns yellow, it shows that brucine is present. Should leave no ash when burned with free access of air.

**Actions and Uses.**—In excessive doses, highly poisonous; in medicinal doses, stimulant and tonic in paralysis, and in chorea in the dog.

**Doses.**—HORSE, 1 to 3 grains.—CATTLE, 1 to 3 grains.—SHEEP, $\frac{1}{3}$ to 1 grain.—DOG, $\frac{1}{30}$ to $\frac{1}{10}$ grain. Once or twice a day.

**Modes of Application.**—In the form of bolus or pill, or dissolved in water acidulated with acetic or sulphuric acid.

**Antidote.**—Same as to Nux Vomica.

**Preparation.**—Liquor Strychninæ Hydrochloridi.

## SUCCUS RHAMNI

BUCKTHORN-JUICE

**Synonym.**—Rhamni Succus.

The recently expressed juice of the ripe berry of the common buckthorn, *Rhamnus catharticus*, belonging to the natural order *Rhamnaceæ*.

**Uses.**—See Syrupus Rhamni.

**Preparation.**—Syrupus Rhamni.

## SULPHUR SUBLIMATUM

SUBLIMED SULPHUR

**Synonym.**—Flowers of Sulphur.

**Composition.**—An elementary body, having the symbol S.

**Mode of Preparation.**—Crude sulphur, imported from Sicily, Naples, the Roman States, and other volcanic districts, is placed in a retort connected with a brick chamber. On the application of heat to the retort the sulphur sublimes into the chamber, to the walls of which it attaches itself, while the earthy impurities remain behind. When a sufficient quantity of the sulphur has collected on the walls of the condensing-chamber, it is scraped off for use.

**Characters and Tests.**—A slightly gritty, crystalline powder, of a fine greenish-yellow colour, tasteless and odourless, unless heated; burns in open vessels with a blue flame and the evolution of sulphurous anhydride, recognisable by its odour. Entirely volatilised by heat. Insoluble in water, alcohol, and ether, but soluble in carbon disulphide and turpentine. Should not redden moistened litmus-paper. Solution of ammonia agitated with it and filtered does not leave any residue on evaporation, showing that it is free from arsenic trisulphide.

**Actions and Uses.**—*Internally*:—In excessive doses, irritant poison; in medicinal doses, laxative, and as an alterative in rheumatism, skin-diseases, and many other affections. *Externally*:—In scabies and other skin-diseases.

**Doses.**—*Laxative*:—Horse, 3 to 4 ounces.—Cattle, 5 to 6 ounces.—Sheep, 2 ounces.—Pig, 1½ to 2 ounces. Dog, 2 to 8 drachms. *Alterative*:—Horse, ¾ to 1½ ounce.—Cattle, 1 to 2 ounces.—Sheep, 6 drachms.—Pig, 4 to 6 drachms.—Dog, ½ to 2 drachms.

**Modes of Application.**—*Internally*:—In the form of bolus or pill, or mixed with gruel or dissolved in oil. *Externally*:—Unguentum Sulphuris.

**Preparations.**—Unguentum Sulphuris; Unguentum Sulphuris Compositum.

## SULPHURIS IODIDUM

SULPHUR IODIDE

**Composition.**—Not well established.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Iodine . . . . . | 4 ounces |
| Sublimed Sulphur . . . | 1 ounce |

Rub them together in a Wedgwood mortar until they are thoroughly mixed. Put the mixture into a flask, close the orifice loosely, and apply a gentle heat, so that the colour of the mass shall become gradually darkened. When the colour has become uniformly dark throughout, increase the heat so as to produce liquefaction. Then incline the flask in different directions, in order to return into the liquid any portion of the iodine which may have condensed on the inner surface of the vessel. Lastly, withdraw the heat, and when the liquid has congealed remove the mass, by breaking the flask, and reduce to pieces. Keep in a well-stoppered bottle.

**Characters and Tests.**—A greyish-black solid, with a radiated crystalline appearance. Resembles iodine in odour and in the property of staining the skin. Soluble in about sixty parts of glycerin; insoluble in water, but decomposed when boiled with it.

**Actions and Uses.**—*Externally*:—In scabies, urticaria, and other skin-affections; also applied to farcy ulcers and indolent sores.

**Mode of Application.**—Unguentum Sulphuris Iodidi.

**Preparation.**—Unguentum Sulphuris Iodidi.

## SYRUPUS CHLORALIS HYDRATIS

SYRUP OF HYDRATE OF CHLORAL

**Preparation.**—Take of

| | |
|---|---|
| Hydrate of Chloral . . . | 80 grains |
| Distilled Water . . . . | 1½ drachm |
| Syrup to make . . . . | 1 ounce |

Mix together.

The specific gravity should be about 1·32.

**Composition.**—One drachm contains ten grains of Chloral Hydrate.

**Uses.**—See Chloral Hydras. Used in canine practice.

**Dose.**—Dog, 30 minims to 2 drachms.

## SYRUPUS PAPAVERIS

SYRUP OF POPPIES

Take of

| | |
|---|---|
| Poppy-capsules, dried, freed from seeds, and coarsely powdered . . . . | 36 ounces |
| Rectified Spirit . . . | 16 fluid ounces |
| Refined Sugar . . . | 4 pounds |
| Boiling Distilled Water . | a sufficiency |

Mix the poppy-capsules with four pints of the water, and infuse for twenty-four hours, stirring them frequently; then pack them in a percolator, and, adding more of the water, allow the liquor to pass slowly until about two gallons have been collected, or the poppies are exhausted. Evaporate the liquor by a water-bath until it is reduced to three pints. When quite cold, add the spirit, let the mixture stand for twelve hours, and filter. Distil off the spirit, evaporate the remaining liquor to two pints, and

then add the sugar. The product should weigh six and a half pounds, and should have the sp. gr. 1·32.

**Actions and Uses.**—Sedative, anodyne, and hypnotic for the dog.

**Dose.**—2 to 5 fluid drachms.

**Preparation.**—Mistura Ricini.

## SYRUPUS RHAMNI

### SYRUP OF BUCKTHORN

Take of

| | | |
|---|---|---|
| Buckthorn-juice | . | 4 pints |
| Ginger, sliced<br>Pimento, bruised | . | ¾ ounce of each |
| Refined Sugar | . | 5 pounds, or a sufficiency |
| Rectified Spirit | . | 6 fluid ounces |

Evaporate the juice to two and a half pints, add the ginger and pimento, digest at a gentle heat for four hours, and strain. When cold, add the spirit, let the mixture stand for two days, then decant off the clear liquor, and in this dissolve the sugar with a gentle heat, so as to make the sp. gr. 1·32.

**Uses.**—Mild cathartic and laxative for dogs and cats, and in preparing Mistura Ricini.

**Doses.**—Dog, ½ to 2 ounces.—Cat, 2 to 6 drachms.

## TABACI FOLIA[1]

### LEAF TOBACCO

The dried leaves of Virginian Tobacco, *Nicotiana Tabacum*. Cultivated in America.

**Natural Order.**—*Solanaceæ*.

[1] In practice the ordinary 'shag' tobacco is usually employed.

**Characters.**—Large, mottled-brown, ovate or lanceolate, acuminate leaves, bearing numerous short glandular hairs; having a peculiar heavy odour and nauseous, bitter, acrid taste; yielding, when distilled with potassium hydrate, an alkaline fluid, which has the peculiar odour of nicotine (the active principle of tobacco), and precipitates with platinic chloride and tincture of galls.

**Preparations.**—Decoctum Tabaci; Enema Tabaci; Infusum Tabaci.

## TARAXACI RADIX

TARAXACUM-ROOT

The root of *Taraxacum officinale.*

**Natural Order.**—*Compositæ.*

**Composition.**—It contains asparagin, taraxacin, and taraxacerin.

**Actions and Uses.**—According to Rutherford, it is a feeble hepatic stimulant; it is also a laxative and diuretic, and is of service in duodenal dyspepsia. The best preparation is the fresh juice, the dose of which is for the Horse, 1 ounce; Dog, 1½ to 2 drachms.

## TEREBINTHINA CANADENSIS

CANADA BALSAM

The turpentine obtained by incision from the stem of *Abies balsamea,* Balm of Gilead fir. Cultivated in Canada.

**Natural Order.**—*Coniferæ.*

**Characters.**—Pale yellow, ductile oleo-resin, of the consistence of thin honey; peculiar agreeable odour and

slightly bitter, feebly acrid taste; by exposure to air it dries very slowly into a transparent adhesive varnish; solidifies when mixed with a sixth of its weight of magnesia.

**Use.**—In preparing Collodium Flexile.

## THERIACA

### TREACLE

**Synonyms.**—Sacchari Fæx; Molasses.

**Composition.**—A mixture chiefly consisting of uncrystallisable sugar produced by the action of too high a temperature on cane sugar, unaltered cane sugar, and colouring-matter.

**Characters.**—Thick, brown, fermentable syrup, very sweet, not crystallising by rest or evaporation. Sp. gr. about 1·4. Nearly free from empyreumatic odour and flavour.

**Actions and Uses.** — Laxative. Employed for hastening the activity of cathartics, and as a vehicle for medicines having a disagreeable flavour, or which occasion nausea. Also used in the preparation of masses.

**Doses.**—Horse, 1 to 1½ pound.—Cattle, 1 to 1½ pound.—Sheep, 3 to 4 ounces.—Pig, 3 to 4 ounces.—Dog, 1 to 2 ounces. Frequently repeated until a laxative effect is produced.

## THYMOL

### THYMOL

The solid part of the volatile oil of thyme, which is extracted from *Thymus vulgaris*.

**Natural Order.**—*Labiatæ*.

**Composition.**—$C_6H_3.OH.CH_3.C_3H_7$.

**Characters and Tests.**—Thymol has the form of oblique prismatic crystals, nearly or quite colourless; it has an aromatic thyme-like odour and a pungent aromatic taste. It is soluble in about 1200 parts of cold water, and 900 of boiling water, freely soluble in alcohol, ether, chloroform, and oils. A solution of Thymol in half its bulk of glacial acetic acid, warmed with an equal volume of sulphuric acid, assumes a reddish-violet colour.

**Actions and Uses.**—A powerful disinfectant and antiseptic, also diaphoretic and diuretic, principally employed externally for allaying irritation and removing scales in chronic eczema and psoriasis, either in solution (one to two grains in diluted spirit) or as an ointment (eight to ten grains to an ounce of vaseline).

## TINCTURA ACONITI [1]

### TINCTURE OF ACONITE

Take of

| | |
|---|---|
| Aconite-root, in No. 40 powder | 1 ounce |
| Alcohol (70-per-cent.) | a sufficient quantity |

Moisten the powder with four fluid drachms of the alcohol, and complete the percolation process. The resulting tincture should measure one pint.

[1] This is the tincture of the 'British Pharmacopœia,' 1867; it has one fourth of the strength of Tinctura Aconiti, Dublin, and one-third of the strength of Tinctura Aconiti, London. Fleming's Tincture of Aconite is nearly four times as strong as that of the 'British Pharmacopœia.'

Squire, in his admirable 'Companion to the British Pharmacopœia,' gives the following directions for percolating tinctures:—'After the materials have been macerated for forty-eight hours in three-fourths of the menstruum ordered, percolation will be most efficiently performed by decanting the liquid, pressing the ingredients in the hand, and carefully

**Uses.**—*Internally*:—As a sedative. *Externally*:—As an anodyne.

**Doses.**—Horse, 30 to 50 minims.—Cattle, 40 to 80 minims.—Sheep, 8 to 12 minims.—Pig, 8 to 12 minims.—Dog, 3 to 10 minims. Given in water and repeated every two hours.

## TINCTURA ALOES COMPOSITA

### COMPOUND TINCTURE OF ALOES

Take of

| | |
|---|---|
| Barbados Aloes, in coarse powder | 10 drachms |
| Myrrh | 5 drachms |
| Methylated or Rectified Spirit | 14 fluid ounces |
| Water | 6 fluid ounces |

Macerate the aloes and the myrrh in the spirit and the water, previously mixed, for fourteen days in a closed vessel, and shake frequently. Filter and add sufficient proof spirit to make one pint.

**Uses.**—*Externally*:—Excitant to wounds, obstinate ulcers, and excoriations.

packing them, in small portions at a time, in a conical percolator, so that the mass shall be uniformly tight throughout. The decanted liquid may then be poured upon the ingredients and suffered to percolate; the remainder of the menstruum being afterwards poured upon them in order to chase the strong tincture out. As soon as the liquid ceases to drop, the ingredients are to be removed and pressed. Any deficiency in the product may be made up by adding more of the menstruum and repeating the pressure.'

## TINCTURA ARNICÆ

### TINCTURE OF ARNICA

Take of

| | |
|---|---|
| Arnica-rhizome, bruised . . | 1 ounce |
| Alcohol (70-per-cent.) . . . | 1 pint |

Macerate the arnica for forty-eight hours in fifteen fluid ounces of the spirit, in a closed vessel, agitating occasionally; transfer to a percolator, and when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Then subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient rectified spirit to make one pint.

**Uses.** — *Externally*: — Stimulant to sprains and bruises, and in chronic rheumatism.

## TINCTURA BELLADONNÆ

### TINCTURE OF BELLADONNA

Take of

| | |
|---|---|
| Liquid Extract of Belladonna | 2 fluid ounces |
| Alcohol (60-per-cent.) . . | a sufficient quantity |

To the liquid extract of belladonna add enough of the alcohol to form thirty fluid ounces of the tincture; set aside for twenty-four hours; filter.

**Doses.**—For the larger animals, 1 fluid ounce. For the smaller, 5 to 15 minims.

## TINCTURA CAMPHORÆ

See Spiritus Camphoræ.

## TINCTURA CANNABIS INDICÆ

TINCTURE OF INDIAN HEMP

Take of

| | |
|---|---|
| Extract of Indian Hemp . . . | 1 ounce |
| Alcohol (90-per-cent.) . . . | 20 ounces |

Dissolve.

**Uses.**—*Internally*:—See Extractum Cannabis Indicæ.

**Doses.** — Horse, 1 to 2 ounces.—Dog, 10 to 40 minims.

## TINCTURA CANTHARIDIS

TINCTURE OF CANTHARIDES

Take of

| | |
|---|---|
| Cantharides, in coarse powder . . | ¼ ounce |
| Alcohol (90-per-cent.) . . . | 1 pint |

Macerate for seven days in a closed vessel, with occasional agitation; strain, press, and add sufficient proof spirit to make one pint.

**Uses.** — *Internally*: — Stimulant and tonic. *Externally*:—Tinctura Cantharidis Fortior is used instead.

**Doses.**—Horse, 1 to 4 fluid ounces.—Cattle, 3 to 6 fluid ounces.—Sheep, ½ to 1½ fluid ounce.—Pig, ½ to 1½ fluid ounce.—Dog, 1 to 3 fluid drachms.

## TINCTURA CANTHARIDIS FORTIOR

STRONG TINCTURE OF CANTHARIDES

Take of

| | |
|---|---|
| Cantharides, in coarse powder . . | 1 ounce |
| Proof Spirit . . . . . | 1 pint |

Macerate for seven days in a closed vessel, with occasional agitation ; strain, press, filter, and add sufficient proof spirit to make one pint.

**Use.**—*Externally* :—Stimulant and rubefacient.

## TINCTURA CARDAMOMI COMPOSITA

COMPOUND TINCTURE OF CARDAMOMS

Take of

| | |
|---|---|
| Cardamom-seeds, freed from their pericarps and bruised | 1 ounce |
| Cinnamon, bruised . . . | 2 ounces |
| Proof Spirit . . . . | 80 fluid ounces |

Macerate, with occasional agitation, for forty-eight hours with sixty fluid ounces of the spirit, pack in a percolator, and let it drain ; pour on the remaining spirit, and, when it ceases to drop, press, and wash the marc with sufficient spirit to make the resulting tincture measure eighty fluid ounces.

**Uses.**—Carminative and antispasmodic.

**Doses.**—Horse, 1 to 4 fluid ounces.—Cattle, 2 to 6 fluid ounces.—Sheep, 1 to 2 fluid ounces.—Pig, 1 to 2 fluid ounces.—Dog, 2 to 6 fluid drachms.

**Preparation.**—Tinctura Chloroformi Composita.

## TINCTURA CHLOROFORMI COMPOSITA

### COMPOUND TINCTURE OF CHLOROFORM

Take of

| | |
|---|---|
| Chloroform . . . . . | 1 fluid ounce |
| Rectified Spirit . . . . | 4 fluid ounces |
| Compound Tincture of Cardamoms | 10 fluid ounces |

Mix.

**Uses.**—Similar to Spiritus Chloroformi.

**Doses.**—Same as Spiritus Chloroformi.

## TINCTURA CINCHONÆ

### TINCTURE OF CINCHONA

Take of

| | |
|---|---|
| Red Cinchona Bark, in moderately fine powder . . . . | 4 ounces |
| Alcohol (70-per-cent.) . . . | 1 pint |

Macerate the cinchona-bark for forty-eight hours in fifteen fluid ounces of the spirit, in a closed vessel, agitating occasionally; then transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Afterwards subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient proof spirit to make one pint.

**Uses.**—*Internally*:—Stimulant, stomachic, and tonic.

**Doses.**—HORSE, 1 to 3 fluid ounces.—CATTLE, 3 to 6 fluid ounces.—SHEEP, ½ to 1 fluid ounce.—PIG, ½ to 1 fluid ounce.—DOG, 2 to 5 fluid drachms.

## TINCTURA COLCHICI SEMINUM

TINCTURE OF COLCHICUM-SEEDS

Take of

| | |
|---|---|
| Colchicum-seeds, powdered . . | 4 ounces |
| Alcohol (45-per-cent.) . . . | 1 pint |

Macerate the colchicum for forty-eight hours in fifteen fluid ounces of the spirit, in a closed vessel, agitating occasionally; then transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Afterwards subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient proof spirit to make one pint.

**Uses.** —*Internally*: — Sedative. *Externally*: — Anodyne.

**Doses.**—Horse, ½ to 2 fluid ounces.—Cattle, ½ to 2 fluid ounces.—Sheep, 1 to 4 fluid drachms.—Pig, 1 to 4 fluid drachms.—Dog, 20 to 80 minims.

## TINCTURA CONII

TINCTURE OF HEMLOCK

**Synonym.**—Tinctura Conii Fructus.

Take of

| | |
|---|---|
| Hemlock-fruit, bruised . . | 4 ounces |
| Alcohol (70-per-cent.) . . . | 1 pint |

Macerate the hemlock-fruit for forty-eight hours in fifteen ounces of the spirit, in a closed vessel, agitating occasionally; transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Then subject the contents of the percolator to pressure, filter the product, mix

the liquids, and add sufficient proof spirit to make one pint.

**Uses.**—*Internally*:—Alterative and sedative.

**Doses.**—HORSE, 1 to 2 fluid ounces.—CATTLE, 1 to 2 fluid ounces.—SHEEP, 3 to 6 fluid drachms.—PIG, 3 to 6 fluid drachms.—DOG, 10 to 40 minims.

## TINCTURA CROTONIS

### TINCTURE OF CROTON

Take of

| | |
|---|---|
| Croton-seeds, bruised . . . | 1½ ounce |
| Rectified Spirit . . . . | 1 pint |

Macerate for seven days in a closed vessel, with occasional agitation; strain, press, filter, and add sufficient rectified spirit to make one pint.

**Uses.**—*Internally*:—Cathartic. *Externally*:—Rubefacient and counter-irritant.

**Doses.**—HORSE, ½ to 1 fluid ounce.—CATTLE, ½ to 1½ fluid ounce.—SHEEP, 1 to 3 fluid drachms.—PIG, 1 to 3 fluid drachms.—DOG, 20 to 50 minims.

## TINCTURA DIGITALIS

### TINCTURE OF DIGITALIS

Take of

| | |
|---|---|
| Digitalis-leaves, powdered . . | 2½ ounces |
| Alcohol (60-per-cent.) . . . | 1 pint |

Moisten the powder with two fluid ounces of the alcohol, and complete the percolation process. The resulting tincture should measure one pint.

**Doses.**—HORSES and CATTLE, 2 to 4 drachms.—SHEEP, ½ to 1 drachm.—DOGS, 2 to 10 minims.

## TINCTURA ERGOTÆ

TINCTURE OF ERGOT

Take of

| | |
|---|---|
| Ergot, in coarse powder . . . | 5 ounces |
| Proof Spirit . . . . . | 1 pint |

Macerate the ergot for forty-eight hours in fifteen fluid ounces of the spirit, in a closed vessel, agitating occasionally; then transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Afterwards subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient proof spirit to make one pint.

**Use.**—*Internally*:—Parturient.

**Doses.**—MARE, 1 to 2 fluid ounces.—Cow, 1 to 3 fluid ounces.—EWE, 1½ to 3 fluid drachms.—Sow, 1½ to 3 fluid drachms.—BITCH, 20 to 60 minims.

## TINCTURA FERRI PERCHLORIDI[1]

TINCTURE OF PERCHLORIDE OF IRON

**Synonym.**—Tinctura Ferri Sesquichloridi.

Take of

| | |
|---|---|
| Strong Solution of Perchloride of Iron . . . . . | 5 fluid ounces |
| Alcohol (90-per-cent.) . . | 5 fluid ounces |
| Distilled Water to . . . | 1 pint |

Mix and preserve in a stoppered bottle.

[1] The Liquor Ferri Perchloridi Fortis diluted with three parts of distilled water constitutes the medicinal Liquor Ferri Perchloridi, which for ordinary purposes is as effectual as the tincture.

**Uses.**—*Internally*: Astringent and tonic in hæmaturia, farcy, many typhoid affections, and indigestion; also in distemper in dogs. *Externally*:—Astringent, styptic, and caustic.

**Doses.**—Horse, ½ to 1 fluid ounce.—Cattle, ½ to 1 fluid ounce.—Sheep, 20 to 30 minims.—Pig, 10 to 20 minims.—Dog, 1 to 10 minims. Diluted with water.

## TINCTURA IODI[1]

### TINCTURE OF IODINE

Take of

| | |
|---|---|
| Iodine . . . . . | 2½ ounces |
| Iodide of Potassium . . . | 1¼ ounce |
| Alcohol (90-per-cent.) . . | 1 pint |

Dissolve the iodine and iodide of potassium in the spirit.

**Uses.**—*Internally*:— Alterative. *Externally*:— Caustic, stimulant, and deobstruent.

**Doses.**—Horse, 2 to 6 fluid drachms.—Cattle, 3 to 8 fluid drachms.—Sheep, 1 to 2 fluid drachms.—Pig, 20 to 60 minims.—Dog, 10 to 30 minims.

## TINCTURA MYRRHÆ

### TINCTURE OF MYRRH

Take of

| | |
|---|---|
| Myrrh, in coarse powder . . | 4 ounces |
| Alcohol (90-per-cent.) . . . | 1 pint |

Macerate the myrrh for forty-eight hours in fifteen fluid ounces of the spirit, in a closed vessel, agitating

[1] This preparation is five times as strong as the Tinctura Iodi of the 'British Pharmacopœia.'

occasionally; then transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Afterwards subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient rectified spirit to make one pint.

**Uses.**—*Internally*:—Stomachic, carminative, and mild laxative. *Externally*:—Mild astringent and excitant to foul and indolent ulcers, and to canker in the mouth.

**Doses.**—Horse, 2 drachms.—Cattle, 2 drachms.—Sheep, ½ to 1 drachm.—Pig, ½ to 1 drachm.—Dog, 10 to 20 minims.

## TINCTURA NUCIS VOMICÆ

### TINCTURE OF NUX VOMICA

Take of

| | |
|---|---|
| Liquid Extract of Nux Vomica . | 2 fluid ounces |
| Alcohol (90-per-cent.) . . | a sufficiency |
| Distilled Water . . . . | 3 fluid ounces |

Mix the liquid extract with the distilled water; add sufficient alcohol to produce 12 fluid ounces of the tincture; filter.

**Uses.**—Stimulant and tonic in paralysis, chorea, and dyspepsia.

**Doses.**—Horse, 4 to 8 fluid drachms.—Cattle, 4 to 8 fluid drachms.—Sheep, 1 to 2 fluid drachms.—Pig, 1 to 2 fluid drachms.—Dog, 5 to 15 minims.

## TINCTURA OPII

TINCTURE OF OPIUM

**Synonym.**—Laudanum.

Take of

| | |
|---|---|
| Opium, in coarse powder . . | 3 ounces |
| Alcohol (90-per-cent.) . . . | 10 ounces |
| Distilled Water . . . . | 10 ounces |

Macerate for seven days in a closed vessel, with occasional agitation; then strain, press, filter, and add sufficient proof spirit to make one pint.

**Uses.**—Anodyne, hypnotic, sedative, and antispasmodic.

**Doses.**—Horse, 1 to 3 fluid ounces.—Cattle, 1 to 3 fluid ounces.—Sheep, 2 to 8 fluid drachms.—Pig, 2 to 8 fluid drachms.—Dog, 15 to 40 minims.

**Preparations.**—Enema Opii; Linimentum Opii.

## TINCTURA PIMENTÆ

TINCTURE OF PIMENTO

Take of

| | |
|---|---|
| Pimento-berries, bruised . . | 2½ ounces |
| Proof Spirit . . . . | 1 pint |

Macerate for seven days in a closed vessel, with occasional agitation; then strain, press, filter, and add sufficient proof spirit to make one pint.

**Uses.**—Antispasmodic and carminative.

**Doses.**—Horse, 2 to 4 fluid ounces.—Cattle, 2 to 6 fluid ounces.—Sheep, 1 to 3 fluid ounces.—Pig, 1 to 3 fluid ounces.—Dog, 1 to 3 fluid drachms.

## TINCTURA SANTALIS RUBRA

TINCTURE OF RED SANDERS WOOD

Take of

| | |
|---|---|
| Sanders Wood, rasped to powder . | ½ ounce |
| Proof Spirit . . . . . | 1 pint |

Macerate for seven days in a closed vessel, with occasional agitation; then strain, press, filter, and add sufficient proof spirit to make one pint.

**Use.**—To impart colour to draughts.

## TINCTURA ZINGIBERIS

TINCTURE OF GINGER

Take of

| | |
|---|---|
| Ginger, in coarse powder . . | 2½ ounces |
| Alcohol (90-per-cent.) . . . | 1 pint |

Macerate the ginger for forty-eight hours in fifteen fluid ounces of the spirit, in a closed vessel, agitating occasionally; then transfer to a percolator, and, when the fluid ceases to pass, continue the percolation with the remaining five ounces of spirit. Afterwards subject the contents of the percolator to pressure, filter the product, mix the liquids, and add sufficient rectified spirit to make one pint.

**Uses.**—Carminative, and as an adjunct to tonic, stimulant, and purgative medicines.

**Doses.**—Horse, 1 to 2 fluid ounces.—Cattle, 1 to 3 fluid ounces.—Sheep, 2 to 6 fluid drachms.—Pig, 2 to 6 fluid drachms.—Dog, ½ to 2 fluid drachms.

## UNGUENTUM ACIDI CARBOLICI

OINTMENT OF CARBOLIC ACID

Take of

| | |
|---|---|
| Carbolic Acid . . . . | 1 ounce |
| Prepared Lard . . . . | 6 ounces |

Mix thoroughly.

**Uses.**—Applied to sloughing and unhealthy wounds.

## UNGUENTUM ACIDI CARBOLICI COMPOSITUM

COMPOUND OINTMENT OF CARBOLIC ACID

Take of

| | |
|---|---|
| Ointment of Carbolic Acid . . | 7 ounces |
| Sublimed Sulphur . . . | 2 ounces |

Mix thoroughly.

**Uses.**—In the treatment of scab and mange.

## UNGUENTUM ACIDI TANNICI

OINTMENT OF TANNIC ACID

Take of

| | |
|---|---|
| Tannic Acid . . . . . | 1 part |
| Prepared Lard . . . . | 6 parts |

Mix thoroughly.

**Use.**—Astringent. Sometimes from ½ to 1 part of opium is added to this ointment.

## UNGUENTUM ACONITI[1]

OINTMENT OF ACONITE

Take of

| | |
|---|---|
| Extract of Aconite . . . | 2 ounces |
| Prepared Lard . . . . | 4 ounces |

Mix thoroughly.

**Use.**—Anodyne.

## UNGUENTUM ALUMINIS COMPOSITUM

COMPOUND OINTMENT OF ALUM

Take of

| | |
|---|---|
| Alum, in fine powder . . . | 1 ounce |
| Common Turpentine . . . | 1 ounce |
| Prepared Lard . . . . | 3 ounces |

Melt the turpentine and the lard together on a water-bath, add the alum, and stir the mixture while it cools.

**Uses.**—Astringent and digestive in 'grease.'

## UNGUENTUM ANTIMONII TARTARATI

OINTMENT OF TARTARATED ANTIMONY

Take of

| | |
|---|---|
| Tartarated Antimony, in fine powder | 1 ounce |
| Prepared Lard . . . . . | 4 ounces |

Mix thoroughly.

**Uses.**—Counter-irritant to the chest in pulmonary complaints, and to joints in chronic articular affections.

[1] Unguentum Aconitinæ of the 'British Pharmacopœia' 1898 consists of 10 grains of Aconitine, 80 grains of Oleic Acid, and 410 grains of Prepared Lard.

## UNGUENTUM ARGENTI NITRATIS

### OINTMENT OF NITRATE OF SILVER

Take of

| | |
|---|---|
| Nitrate of Silver, in fine powder . | 5 to 10 grains |
| Prepared Lard . . . . | 1 ounce |

Mix thoroughly.

**Use.**—A piece the size of a pea is placed between the eyelids every third day in chronic ophthalmia.

## UNGUENTUM ARSENICI

### OINTMENT OF ARSENIC

Take of

| | |
|---|---|
| Arsenious Acid, in fine powder . | 6 grains |
| Prepared Lard . . . . | 1 ounce |

Mix thoroughly.

**Use.**—Caustic to warts and cancerous growths.

## UNGUENTUM CALCIS CHLORINATÆ

### OINTMENT OF CHLORINATED LIME

Take of

| | |
|---|---|
| Chlorinated Lime . . . . | 1 ounce |
| Prepared Lard . . . . | 4 to 8 ounces |

Mix thoroughly.

**Use.**—In grease, especially when accompanied by considerable fœtor.

## UNGUENTUM CANTHARIDIS

OINTMENT OF CANTHARIDES

**Synonym.**—Blistering-ointment.

Take of

| | |
|---|---|
| Cantharides, in powder . . . | 1 ounce |
| Prepared Lard . . . . | 6 ounces |

Digest the cantharides and lard together over a water-bath for three hours, with occasional stirring; while hot filter through paper, and allow the clear liquid to cool.

**Use.**—Irritant to keep open blisters, and to promote the discharge from setons and ulcers.

## UNGUENTUM CREOSOTI

OINTMENT OF CREOSOTE

Take of

| | |
|---|---|
| Creosote . . . . . | 1 fluid ounce |
| Prepared Lard . . . . | 8 ounces |

Mix thoroughly.

**Uses.**—Same as Unguentum Acidi Carbolici.

## UNGUENTUM CUPRI SUBACETATIS

OINTMENT OF SUBACETATE OF COPPER

**Synonyms.**—Unguentum Æruginis; Ointment of Verdigris.

Take of

| | |
|---|---|
| Subacetate of Copper, in fine powder . . . . . | 1 ounce |
| Common Turpentine . . . | 1 ounce |
| Prepared Lard . . . . | 12 ounces |

Melt the turpentine and lard together over a water-bath, then withdraw the heat, add the subacetate of copper, and thoroughly mix by stirring the whole until cold.

**Use.**—Detergent to foul ulcers.

## UNGUENTUM GALLÆ

### OINTMENT OF GALLS

Take of

| | |
|---|---|
| Galls, in fine powder | 1 ounce |
| Prepared Lard | 4 ounces |

Mix thoroughly.

**Use.**—Astringent to ulcers with profuse discharge.

## UNGUENTUM GALLÆ CUM OPIO

### OINTMENT OF GALLS AND OPIUM

Take of

| | |
|---|---|
| Ointment of Galls | 5 ounces |
| Opium, in fine powder | ½ ounce |

Mix thoroughly.

**Uses.**—Astringent and anodyne.

## UNGUENTUM HELLEBORI NIGRI

### OINTMENT OF BLACK HELLEBORE

Take of

| | |
|---|---|
| Black Hellebore Leaves<br>Common Turpentine<br>Prepared Lard | equal parts |

Melt the turpentine and the lard together in a water-

bath, add the hellebore, and digest for one hour; then remove the mixture and express through calico.

**Use.**—Digestive to wounds of cattle.

## UNGUENTUM HYDRARGYRI

OINTMENT OF MERCURY

**Synonym.**—Blue Ointment.

Take of

| | |
|---|---|
| Mercury . . . . . | 1 pound |
| Prepared Lard . . . . | 1 pound |
| Prepared Suet . . . . . | 1 ounce |

Rub them together until metallic globules cease to be visible when a small portion of the ointment is rubbed with the finger on brown paper.

**Uses.**—In scab, mange, and other scurfy cutaneous affections; to indolent sores and ulcers; as a resolvent in glandular enlargements; and applied to the skin with friction to expedite and maintain the constitutional effects produced by the internal administration of mercury. For skin-affections, however, Unguentum Hydrargyri Ammoniati is generally used.

## UNGUENTUM HYDRARGYRI AMMONIATI

OINTMENT OF AMMONIATED MERCURY

Take of

| | |
|---|---|
| Ammoniated Mercury . . | 1 ounce |
| Prepared Lard . . . | 6 to 8 ounces |

Mix thoroughly.

**Uses.**—To destroy pediculi and acari, and for skin affections.

## UNGUENTUM HYDRARGYRI COMPOSITUM

### COMPOUND OINTMENT OF MERCURY

Take of

| | |
|---|---|
| Ointment of Mercury . . . | 1 ounce |
| Soft Soap . . . . . | 2 ounces |

Mix thoroughly.

**Uses.**—See Unguentum Hydrargyri.

## UNGUENTUM HYDRARGYRI IODIDI RUBRI[1]

### OINTMENT OF RED IODIDE OF MERCURY

**Synonyms.**—Unguentum Hydrargyri Biniodidi; Ointment of Biniodide of Mercury; Blistering-ointment.

Take of

| | |
|---|---|
| Red Iodide of Mercury, in fine powder . . . . . | 1 ounce[2] |
| Prepared Lard . . . . . | 8 ounces |

Mix thoroughly.

**Uses.**—Caustic, stimulant, and detergent to unhealthy sores; counter-irritant; also resolvent in the treatment of curbs, splints, incipient spavins, and enlarged bursæ.

[1] This preparation is nearly three-and-a-half times stronger than Unguentum Hydrargyri Iodidi Rubri of the 'British Pharmacopœia.'

[2] An ointment containing twice the quantity of iodide of mercury forms a powerful counter-irritant when applied to the skin of the ox.

## UNGUENTUM HYDRARGYRI NITRATIS

OINTMENT OF NITRATE OF MERCURY

**Synonym.**—Citrine Ointment.

Take of

| | |
|---|---|
| Mercury, by weight . . | 4 ounces |
| Nitric Acid . . . . | 12 fluid ounces |
| Prepared Lard . . . | 15 ounces |
| Olive Oil . . . . | 32 fluid ounces |

Dissolve the mercury in the nitric acid with the aid of a gentle heat.

| Mercury. | Nitric Acid. | | Mercuric Nitrate. | | Nitric Oxide. | | Water. |
|---|---|---|---|---|---|---|---|
| $Hg_3$ | + $8HNO_3$ | = | $3Hg(NO_3)_2$ | + | $2NO$ | + | $4H_2O$ |

Melt the lard in the oil by a steam- or water-bath, in a porcelain vessel capable of holding six times the quantity, and while the mixture is hot add the solution of mercuric nitrate, also hot, mixing them thoroughly. If the mixture does not froth up, increase the heat till this occurs. Keep the ointment stirred until it is cold.

**Uses.**—Stimulant and detergent in grease, ringworm, and other skin-affections. One part of this ointment, mixed with one part of lard and one of olive oil, is employed in the treatment of tarsal ophthalmia, and for the removal of nebulæ from the cornea.

## UNGUENTUM HYDRARGYRI PERCHLORIDI

OINTMENT OF PERCHLORIDE OF MERCURY

Take of

| | |
|---|---|
| Perchloride of Mercury, in fine powder . | 1 ounce |
| Prepared Lard . . . . . | 2 ounces |

Mix thoroughly.

**Uses.**—Spread on leather for the removal of bony tumours. One part of this ointment, mixed with two parts of lard, is used for dressing Professor Varnell's caustic clams.

## UNGUENTUM IODI

### OINTMENT OF IODINE

Take of

| | |
|---|---|
| Iodine | 1 ounce |
| Prepared Lard | 8 ounces |
| Rectified Spirit or Glycerin | 1 drachm |

Mix thoroughly.

**Uses.**—Stimulant and resolvent.

## UNGUENTUM IODI COMPOSITUM

### COMPOUND OINTMENT OF IODINE

Take of

| | |
|---|---|
| Iodine | ½ ounce |
| Iodide of Potassium, in fine powder | 1 ounce |
| Prepared Lard | 8 ounces |
| Rectified Spirit or Glycerin | 1 drachm |

Mix thoroughly.

**Uses.**—Stimulant and resolvent.

## UNGUENTUM MYLABRIDIS

### OINTMENT OF MYLABRIS

Take of

| | |
|---|---|
| Mylabris, in fine powder | 1 ounce |
| Prepared Lard | 6 ounces |

Digest the mylabris and the lard together over a

water-bath for three hours, with occasional stirring; while hot filter through paper, and allow the clear liquid to cool.

**Uses.**—Same as those of Unguentum Cantharidis, but has no diuretic effect.

## UNGUENTUM PICIS LIQUIDÆ

### OINTMENT OF TAR

Take of

| | |
|---|---|
| Common Tar . . . | equal parts |
| Resin . . . . | |
| Prepared Lard . . . | |

Melt together with a gentle heat, and stir the mixture while it cools.

**Uses.**—In foot-rot in sheep, and as a foot-ointment for all other domesticated animals.

## UNGUENTUM POTASSÆ SULPHURATÆ

### OINTMENT OF SULPHURATED POTASH

Take of

| | |
|---|---|
| Sulphurated Potash, in fine powder | 1 ounce |
| Prepared Lard . . . . | 8 ounces |

Triturate the sulphurated potash in a porcelain mortar, and gradually add the lard, rubbing them together until the ointment is perfectly smooth and free from grittiness. This ointment, when used, should be recently prepared.

**Uses.**—In mange and other skin-affections.

## UNGUENTUM RESINÆ

OINTMENT OF RESIN

**Synonym.**—Digestive Ointment.

Take of

| | |
|---|---|
| Resin, in coarse powder . . | 8 ounces |
| Yellow Wax . . . . | 6 ounces |
| Almond Oil . . . . | 3 ounces |
| Prepared Lard . . . . | 3 ounces |

Melt with a gentle heat, strain the mixture while hot through flannel, and stir constantly while it cools.

**Uses.**—Stimulant to ulcers, wounds, and blistered surfaces, and to impart consistency to other ointments.

## UNGUENTUM SABINÆ

OINTMENT OF SAVIN

Take of

| | |
|---|---|
| Savin-tops, bruised . . . | 1 ounce |
| Prepared Lard . . . . | 2 ounces |

Melt the lard on a water-bath, add the savin, and digest for twenty minutes. Then remove the mixture, and express through calico.

**Uses.**—Irritant to blistered surfaces, and for the removal of warts.

## UNGUENTUM SULPHURIS

OINTMENT OF SULPHUR

Take of

| | |
|---|---|
| Sublimed Sulphur . . . | 1 ounce |
| Prepared Lard . . . . | 4 ounces |

Mix.

**Uses.**—For scab in sheep, parasiticide, and in treatment of eczema, mange, &c.

## UNGUENTUM SULPHURIS COMPOSITUM

### COMPOUND OINTMENT OF SULPHUR

**Synonym.**—Sheep-ointment.

Take of

| | |
|---|---|
| Sublimed Sulphur . . . | 1 pound |
| Common Turpentine . . . | 4 ounces |
| Ointment of Mercury . . . | 2 ounces |
| Linseed Oil . . . . . | 1 pint |

Melt the oil and turpentine together over a water-bath; when the mixture is nearly cold, stir in the sulphur, and afterwards the ointment of mercury, until the whole is thoroughly mixed.

**Uses.**—For eradicating scab in sheep.

## UNGUENTUM SULPHURIS IODIDI

### OINTMENT OF IODIDE OF SULPHUR

Take of

| | |
|---|---|
| Iodide of Sulphur, in fine powder . | 1 ounce |
| Prepared Lard . . . . | 8 ounces |

Triturate the iodide of sulphur in a porcelain mortar, and gradually add the lard, rubbing them together until the ointment is perfectly smooth and free from grittiness.

**Uses.**—In mange, urticaria, and other skin-diseases.

## UNGUENTUM TEREBINTHINÆ

OINTMENT OF TURPENTINE

**Synonym.**—Digestive Ointment.

Take of

| | |
|---|---|
| Common Turpentine . . . | 1 ounce |
| Prepared Lard . . . . | 3 ounces |

Melt together over a water-bath, mix thoroughly by stirring, and cool.

**Use.**—Stimulant to wounds.

## UNGUENTUM VERATRI

OINTMENT OF WHITE HELLEBORE

Take of

| | |
|---|---|
| White Hellebore Root, dried, and in fine powder . . . . | 1 ounce |
| Prepared Lard . . . . | 4 ounces |

Thoroughly mix.

**Uses.**—Applied to the skin as a parasiticide, and for charging materials to be used as setons.

## UNGUENTUM ZINCI CARBONATIS

OINTMENT OF CARBONATE OF ZINC

**Synonym.**—Healing-ointment.

Take of

| | |
|---|---|
| Carbonate of Zinc, in fine powder . | 1 ounce |
| Prepared Lard . . . . | 6 ounces |

Mix thoroughly.

**Uses.** — Desiccant and mild astringent to slight excoriations.

## UNGUENTUM ZINCI OXIDI

OINTMENT OF OXIDE OF ZINC

Take of

| | |
|---|---|
| Oxide of Zinc . . . . | 1 ounce |
| Prepared Lard . . . . | 5 ounces |

Mix.

**Uses.**—Employed in slight excoriations and ulcerations.

## VAPOR CHLORI

CHLORINE GAS

**Composition.**—An element; symbol Cl.

**Modes of Preparation.** — There are two methods commonly resorted to.

*1st Method.*—Put three or four ounces of powdered manganese peroxide into a flask, basin, or other suitable vessel; make it into a thin paste with strong hydrochloric acid, and apply a gentle heat to the mixture.

| Manganese Peroxide. | | Hydrochloric Acid. | | Manganese Chloride. | | Water. | | Chlorine. |
|---|---|---|---|---|---|---|---|---|
| $MnO_2$ | + | $4HCl$ | = | $MnCl_2$ | + | $2H_2O$ | + | $Cl_2$ |

*2nd Method.*[1]—Take of

| | |
|---|---|
| Sodium Chloride (common salt), in fine powder . . . . | $2\frac{1}{2}$ ounces |
| Manganese Peroxide, in fine powder | 2 ounces |
| Strong Sulphuric Acid . . . | $2\frac{1}{2}$ fluid ounces |
| Water . . . . . | 5 fluid ounces |

[1] Chlorine may also be evolved by pouring dilute sulphuric or hydrochloric acid on chlorinated lime.

Intimately mix the manganese peroxide and sodium chloride, and place them in a flask or basin. Pour the sulphuric acid into the water contained in a porcelain, earthenware, or thin glass vessel. Add the dilute sulphuric acid to the mixture of manganese peroxide and sodium chloride, and thoroughly stir or shake them together.

If necessary, the evolution of chlorine may be promoted by the application of a gentle heat.

| Sodium Chloride. | | Manganese Peroxide. | | Sulphuric Acid. | | Chlorine. | | Sodium Sulphate. | |
|---|---|---|---|---|---|---|---|---|---|
| $2NaCl$ | + | $MnO_2$ | + | $2H_2SO_4$ | = | $Cl_2$ | + | $Na_2SO_4$ | + |

| Manganese Sulphate. | | Water. |
|---|---|---|
| $MnSO_4$ | + | $2H_2O$ |

**Characters and Tests.**—Heavy greenish-yellow gas, with a peculiar odour, and producing great irritation when breathed. Bleaches blue litmus-paper, and develops a blue colour (starch iodide) when brought into contact with paper imbued with a mixture of starch and potassium iodide.

**Actions and Uses.**—Chiefly employed for disinfecting the air of stables and other buildings. Occasionally used as a therapeutic in glanders, farcy, and similar affections, in which cases it first increases, but ultimately diminishes, the secretion of the nasal membrane. It has also been recommended as an antidote to poisoning by hydrocyanic acid and sulphuretted hydrogen.

**Modes of Application.**—As an aërial disinfectant, the apparatus is to be placed in the stable; the doors and other openings should be closed for several hours during the generation of the gas, and afterwards opened for some time previously to the entrance of any person.

When the gas is to be inhaled, the patient should be

placed in a loose box, along with the apparatus containing one of the mixtures above described. As soon as the gas begins to exert its irritating effects upon the attendant, he should withdraw, remove the apparatus, and close the door of the box.

**Antidotes.**—Coughing, and other symptoms of the irritant effects of chlorine, may be allayed by inhaling the vapour of ammonia, ether, alcohol, or hot water.

## 'VASELINE' [1]

'VASELINE'

A substance of the consistence of butter, obtained by concentrating petroleum by dry heat and subsequent filtration through animal charcoal.

**Synonyms.**—Petroleum Molle ; Soft Petroleum ; Cosmoline.

**Composition.**—Hexadecane, $C_{16}H_{34}$, and other hydro carbons.

**Actions and Uses.**—It dissolves iodine, sulphur iodide, and carbolic acid ; it is used as a lubricant and an emollient for irritable, inflamed, or blistered mucous and skin surfaces. It is nearly free from greasiness, and is not oxidisable ; it therefore forms a good basis for ointments.

## VERATRUM

WHITE HELLEBORE

The rhizome of *Veratrum album*, a native of the mountainous regions of Central and Southern Europe.

[1] According to the Court of Appeal, there is only one make of Vaseline on the market—that of the Chesebrough Company, who are the owners of the trade-mark 'Vaseline.'

**Natural Order.**—*Melanthaceæ.*

**Composition.**—Its medicinal activity is attributed to an alkaloid named *veratrine*, $C_{37}H_{53}NO_{11}$.

**Characters.**—Usually consists of the rhizome with the radicles attached; in pieces of from two to three inches long, about the thickness of the little finger; covered with a rough, dark-brown bark; greyish white internally. In the fresh state it emits a strong disagreeable odour, which it nearly loses by drying, but it retains its acrid, intensely bitter taste.

**Actions and Uses.**—*Internally*:—In excessive doses, an irritant poison; in medicinal doses, nauseant, emetic, sedative, purgative, and anthelmintic. Given sometimes to the horse as a sedative, and as an excitant to promote absorption in œdematous enlargements of the legs. *Externally*:—Applied to the skin as a parasiticide; also in the cure of mange and other cutaneous diseases.

**Doses.**—Horses and Cattle:—*As a sedative*, ½ to 1 drachm every four or six hours, until its depressant action is manifested. *As an excitant*, ½ to 1 ounce.—Sheep and Pigs, 20 to 30 grains.

**Modes of Application.**—*Internally*:—In the form of bolus, or tincture made with proof spirit. *Externally*:—The powdered rhizome, or Unguentum Veratri.

**Antidotes.**—Demulcent drinks; also tannic acid, and vegetable astringents containing it. If coma be present, ammonia, or some powerful alcoholic stimulant, should be administered.

**Preparation.**—Unguentum Veratri.

## VINUM FERRI

### WINE OF IRON

**Composition.**—A weak solution of ferrous tartrate, malate, and acetate in sherry.

Take of

| | |
|---|---|
| Fine Iron Wire (about No. 35) | 1 ounce |
| Sherry | 1 pint |

Macerate for thirty days in a closed vessel, the iron being almost, but not quite, wholly immersed in the wine; frequently shake the vessel, removing the stopper after each shaking; filter.

**Use.**—Tonic.

**Dose.**—Dog, 1 to 4 fluid drachms two or three times a day.

## ZINCI ACETAS

### ACETATE OF ZINC

**Synonym.**—Zinc Acetate.

**Composition.**—$Zn(C_2H_3O_2)_2.2H_2O$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Carbonate of Zinc | 2 ounces |
| Acetic Acid | 5 ounces, or a sufficiency |
| Distilled Water | 6 fluid ounces |

Add the carbonate of zinc in successive portions to three ounces of the acetic acid previously mixed with the water in a flask; heat gently; add by degrees the remainder of the acid till the carbonate is dissolved; boil

for a few minutes, filter while hot, and set it aside for two days to crystallise. Decant the mother-liquor; evaporate to one-half, and again set it aside for two days to crystallise. Place the crystals (zinc acetate) in a funnel to drain, then spread them on filtering-paper on a porous tile, and dry them by exposure to the air at ordinary temperatures.

| Zinc Carbonate. | | Acetic Acid. | | Zinc Acetate. | | Carbonic Anhydride. | | Water. |
|---|---|---|---|---|---|---|---|---|
| $ZnCO_3$ | + | $2C_2H_4O_2$ | = | $Zn(C_2H_3O_2)_2$ | + | $CO_2$ | + | $H_2O$ |

In the act of crystallising the salt takes up two molecules of water.

**Characters and Tests.**—Thin, translucent, and colourless crystalline plates, of a pearly lustre, with a sharp, unpleasant taste; evolving acetic acid (recognisable by the characteristic odour of its vapour) when decomposed by sulphuric acid; soluble in water, forming a solution which gives a white precipitate (zinc sulphide) with ammonium sulphydrate. A dilute aqueous solution should not be affected by barium chloride (showing the absence of sulphate) or by silver nitrate (showing the absence of chlorides); and when slightly acidulated with hydrochloric acid it should furnish no precipitate with sulphuretted hydrogen. After it has been boiled for a few minutes with a little nitric acid (to convert into ferric salt any ferrous compound that may be present), it yields, with ammonia, a white precipitate (zinc hydrate), which is soluble without colour (if iron be absent) in an excess of the reagent.

**Uses.**—*Externally* :—Stimulant and astringent to wounds, especially those accompanied by profuse discharges; also to allay superficial inflammation. A wash-leather bandage, saturated with a watery solution of

acetate of zinc, is used to reduce the swellings on the legs of hunters.

**Modes of Application.**—A lotion consisting of twenty-five to fifty grains of the acetate dissolved in one pint of cold water, and applied as soon after the solution of the salt as possible.

## ZINCI CARBONAS[1]

### CARBONATE OF ZINC

**Composition.**—A mixture or compound of zinc carbonate and zinc hydrate, $ZnCO_3.2Zn(OH)_2.H_2O$.

**Mode of Preparation.**—Take of

| | |
|---|---|
| Sulphate of Zinc . . . | 10 ounces |
| Carbonate of Soda . . . | 10½ ounces |
| Boiling Distilled Water . . | a sufficiency |

Dissolve the carbonate of soda with a pint of the water in a capacious porcelain vessel, and pour into it the sulphate of zinc, also dissolved in a pint of the water, stirring diligently. Boil for fifteen minutes after effervescence (escape of carbonic anhydride) has ceased, and let the precipitate (carbonate of zinc, B.P.) subside. Decant the supernatant liquid, pour on the precipitate three pints of boiling distilled water, agitating briskly; let the precipitate again subside, and repeat the processes of affusion of hot distilled water and subsidence till the washings are no longer precipitated by barium chloride, showing that sulphate of zinc is completely removed from the carbonate.

[1] In veterinary practice the native greyish-brown zinc carbonate (calamine) is generally used, after it has been partly purified by levigation. It is then known as 'Prepared Calamine.'

Collect the precipitate on calico, let it drain, and dry it with a gentle heat.

| Zinc Sulphate. | | Sodium Carbonate. | | Water. | | Carbonate of Zinc, B.P. | |
|---|---|---|---|---|---|---|---|
| $3ZnSO_4$ | + | $3Na_2CO_2$ | + | $2H_2O$ | = | $ZnCO_3.2Zn(OH)_2$ | + |

| Sodium Sulphate. | | Carbonic Anhydride. |
|---|---|---|
| $3Na_2SO_4$ | + | $2CO_2$ |

**Characters and Tests.**—White, tasteless, inodorous, insoluble in water, soluble with effervescence and without residue in dilute nitric acid. This solution should not be affected by barium chloride, showing the absence of sulphates, or by silver nitrate, showing the absence of chlorides, and gives with ammonium carbonate a white precipitate (zinc carbonate and hydrate) entirely soluble without colour (if iron be absent) in an excess of the reagent, forming a solution which furnishes a white precipitate (zinc sulphide) with ammonium sulphide, showing the presence of zinc.

**Actions and Uses.**—*Externally* :—Desiccant, stimulant, and astringent.

**Mode of Application.**—The powder dusted over wounds, &c., or Unguentum Zinci Carbonatis.

**Preparations.**—Unguentum Zinci Carbonatis ; Zinci Acetas ; Zinci Oxidum.

## ZINCI CHLORIDUM

### CHLORIDE OF ZINC

**Synonyms.**—Zinc Chloride ; Butter of Zinc.[1]

**Composition.**—$ZnCl_2$.

[1] Tuson's Fluid Disinfectant is an aqueous solution of zinc chloride impregnated with sulphurous-acid gas (see Acidum Sulphurosum).

**Mode of Preparation.**—Take of

| | | |
|---|---|---|
| Granulated Zinc | . | 16 ounces |
| Hydrochloric Acid | . | 44 fluid ounces |
| Solution of Chlorine | . | a sufficiency |
| Carbonate of Zinc | . | ½ ounce, or a sufficiency |
| Distilled Water | . | 1 pint |

Put the zinc into a porcelain basin; add by degrees the hydrochloric acid, previously mixed with the water; and aid the action by gently warming it on a sand-bath until gas (hydrogen) is no longer evolved. Boil for half an hour, supplying the water lost by evaporation, and allow it to stand on a cool part of the sand-bath for twenty-four hours, stirring frequently. Filter the product into a gallon bottle, and pour in the solution of chlorine by degrees, with frequent agitation (to convert any ferrous chloride into ferric chloride), until the fluid acquires a permanent odour of the gas. Add the carbonate of zinc (to precipitate iron and lead, if present) in small quantities at a time, and with renewed agitation, until a brown sediment (ferric hydrate and lead peroxide) appears. Filter through paper into a porcelain basin, and evaporate until a portion of the liquid, withdrawn on the end of a glass rod and cooled, forms an opaque white solid (zinc chloride). Pour it now into proper moulds, and, when the salt has solidified, but before it has cooled, place it in closely stoppered bottles.

| Zinc. | | Hydrochloric Acid. | | Zinc Chloride. | | Hydrogen. |
|---|---|---|---|---|---|---|
| Zn | + | 2HCl | = | $ZnCl_2$ | + | $H_2$ |

**Characters and Tests.**—Colourless opaque rods or tablets, very deliquescent and caustic; almost entirely soluble in water, alcohol, and ether. The watery solution gives a white precipitate (zinc sulphide) with ammonium sulphide, showing the presence of zinc; also a white

precipitate (silver chloride) with silver nitrate, indicating that the salt is a chloride; if first acidulated with hydrochloric acid, it is not affected by sulphuretted hydrogen Its aqueous solution should give no precipitate with barium chloride, showing the absence of sulphates, or ammonium oxalate, showing the absence of calcium; and should not be tinged blue by potassium ferro- or ferric cyanide, showing the absence of iron. Ammonia throws down a white precipitate (zinc hydrate) entirely soluble in an excess of the reagent.

**Uses.**—In the solid state, or in the form of concentrated solution, as a caustic and deodoriser in canker, quittor, foot-rot in sheep, fistulæ, ill-conditioned wounds, phagedænic ulcers, luxuriant granulations, &c. In the form of Liquor Zinci Chloridi it is applied to wounds as an astringent, stimulant, and antiseptic; to the skin of horses and dogs as a parasiticide, and in eczema; also as a 'dip' or 'wash' for killing ticks and preventing the attacks of the fly on sheep. For deodorising and disinfecting stables, manure-heaps, &c.

**Preparation.**—Liquor Zinci Chloridi.

## ZINCI OXIDUM

OXIDE OF ZINC

**Synonym.**—Zinc Oxide.

**Composition.**—ZnO.

**Mode of Preparation.**—Carbonate of zinc is exposed to a dull red heat in a loosely covered Hessian crucible, until a portion taken from the centre of the contents of the crucible and cooled no longer effervesces when dropped into dilute sulphuric acid, showing that the expulsion of

carbonic anhydride is complete. Let the crucible cool, and transfer the product (zinc oxide) to a stoppered bottle.

| Zinc Carbonate. | | Zinc Oxide. | | Carbonic Anhydride. |
|---|---|---|---|---|
| $ZnCO_3$ | = | $ZnO$ | + | $CO_2$ |

**Characters and Tests.**—Soft, nearly white, tasteless, and inodorous powder, becoming pale yellow when heated. Dissolves without effervescence in dilute nitric acid, showing absence of carbonate; forms a solution which is not affected by barium chloride, showing the absence of chlorides; and gives with ammonium carbonate a white precipitate (zinc carbonate and hydrate) which dissolves entirely without colour (if iron be absent) in an excess of the reagent, forming a solution which gives a white precipitate (zinc sulphide) with ammonium sulphide, indicating the presence of zinc.

**Uses.**—*Externally*:—Astringent and desiccant to ichorous ulcers, excoriations, &c.

**Modes of Application.**—In the form of powder and of ointment.

## ZINCI SULPHAS

### SULPHATE OF ZINC

**Synonyms.**—Zinc Sulphate; White Vitriol.

**Composition.**—$ZnSO_4.7H_2O$.

**Mode of Preparation.**—Take of

| | | |
|---|---|---|
| Granulated Zinc | 16 | ounces |
| Sulphuric Acid | 12 | fluid ounces |
| Distilled Water | 4 | pints |
| Solution of Chlorine | a | sufficiency |
| Carbonate of Zinc | ½ | ounce, or a sufficiency |

Pour the sulphuric acid, previously mixed with the water, on the zinc, contained in a porcelain basin, and, when effervescence (escape of hydrogen) has nearly ceased, aid the action by a gentle heat. Filter the fluid into a gallon bottle, and add gradually, with constant agitation, the solution of chlorine [1] until the fluid acquires a permanent odour of chlorine. Add now, with continued agitation, carbonate of zinc,[1] until a brown precipitate (ferric hydrate and lead peroxide) appears; let it settle, filter the solution, evaporate till a pellicle forms on the surface, and set aside to crystallise. Dry the crystals (zinc sulphate) by exposure to the air on filtering-paper placed on porous tiles. More crystals may be obtained by again evaporating the mother-liquor.

| Zinc. | | Sulphuric Acid. | | Zinc Sulphate. | | Hydrogen. |
|---|---|---|---|---|---|---|
| $Zn$ | + | $H_2SO_4$ | = | $ZnSO_4$ | + | $H_2$ |

In the act of crystallising, the salt acquires seven molecules of water.

**Characters and Tests.**—Colourless, transparent, prismatic crystals, strongly resembling sulphate of magnesia, but with a metallic astringent taste. Its aqueous solution gives a white precipitate (zinc sulphide) with ammonium sulphide, indicating the presence of zinc, and a white precipitate (barium sulphate) with barium chloride, insoluble in nitric acid, showing that the salt is a sulphate. Its watery solution should give no bluish-black precipitate or coloration with tincture of galls, showing the absence of iron, and when acidulated with hydrochloric acid it should not be affected by sulphuretted hydrogen. After it has been boiled for a few minutes with a little nitric acid

[1] The uses of the chlorine-water and carbonate of zinc are explained under Zinci Chloridum.

(to convert into ferric salt any ferrous compound that may be present), it should yield with ammonium hydrate a white precipitate (zinc hydrate) which is entirely soluble without colour (showing absence of iron) in an excess of the reagent.

**Actions and Uses.**—*Internally* :—In excessive doses, irritant poison ; in medicinal doses, astringent and tonic. Also emetic when given in sufficient quantity to the pig or dog. *Externally* :—Astringent, excitant, mild escharotic, and desiccant when applied to wounds, injected into sinuses, and in quittor.

**Doses.**—*Tonic* :—HORSE, 1 to 2 drachms.—CATTLE, 2 to 3 drachms.—SHEEP, 20 to 30 grains.—PIG, 10 to 20 grains.—DOG, 2 to 5 grains. *Emetic* :—PIG, 30 to 50 grains.—DOG, 8 to 15 grains.

**Modes of Application.**—*Internally* :—As a tonic, in the form of bolus or dissolved in water; as an emetic, dissolved in tepid water. *Externally* :—In the state of powder, or dissolved in water, the solution varying in strength from half an ounce to seven ounces of the salt to a pint of distilled water, according to the purpose for which the lotion is required.

**Incompatibles.**—Alkalies and their carbonates, lime-water, acetate of lead, nitrate of silver, astringent vegetable infusions, and milk.

**Antidotes.**—Where possible, warm demulcent drinks, to promote the evacuation of the poison by vomiting.

**Preparation.**—Zinci Carbonas.

## ZINGIBER

### GINGER

The scraped and dried rhizome of *Zingiber officinale* obtained from plants cultivated in the West Indies, India, and other countries.

**Natural Order.**—*Zingiberaceæ.*

**Composition.**—Its medicinal properties are attributed to the volatile oil, which, according to Thrush, is a complex mixture of hydrocarbons and products of their oxidation.

**Characters.**—Irregularly lobed decorticated pieces, three or four inches long, sub-compressed, yellowish white, but not chalky, on the surface; has a short mealy fracture, hot taste, and agreeable aroma. Powder yellowish white.

**Actions and Uses.** — Stimulant, stomachic, carminative, and tonic. Given in flatulent colic and debility of the stomach and intestines; also combined with cathartics to increase their activity and to prevent griping.

**Doses.**—Horse, 2 to 4 drachms.—Cattle, 2 to 6 drachms.—Sheep, ½ to 2 drachms.—Pig, ½ to 1 drachm.—Dog, 10 to 30 grains.

**Mode of Application.**—In the form of bolus or as Tinctura Zingiberis.

**Preparations.** — Massa Aloes; Syrupus Rhamni; Tinctura Zingiberis.

# ADDENDA

## LIQUOR CALCIS IODINATÆ[1]

### SOLUTION OF HYPOIODITE OF CALCIUM

A solution of hypoiodite of calcium.

**Actions and Uses.**—It has been pointed out by Ringer and others that iodine in this form has better practical results than the iodides when administered internally.

In cases of actinomycosis, the solution should be administered internally and locally applied.

For washing out wounds, injecting sinuses, and purifying sores, the solution does not affect the healthy tissues injuriously, as do most antiseptics, while it acts powerfully upon disease-products and infective matter, completely destroying their noxious qualities. Also useful diluted with water as a lotion, and with oil as a liniment.

**Dose.**—Horse, 2 drachms to 1 ounce.

## LIQUOR POTASSÆ IODINATÆ[1]

### SOLUTION OF HYPOIODITE OF POTASSIUM

A solution of hypoiodite of potassium.

**Actions and Uses.**—Useful in the treatment of parasitic and other skin-diseases. The solution is used internally and externally, and may be employed with

[1] These are registered preparations of Messrs. Willows, Francis Butler, & Thompson, Limited, Aldersgate Street, London.

advantage as a substitute for both potassium iodide and the preparations of free iodine.

Also used as an outward application as a liniment in the proportion of one part to five of oil.

**Dose.**—Horse, 1 drachm to 4 ounces.

## SODII SULPHOCARBOLAS

### SODIUM SULPHOCARBOLATE

**Composition.**—$C_6H_4OH,SO_2ONa,2H_2O$.

**Actions and Uses.**—According to Gresswell,[1] sulphocarbolate of sodium is a safe and useful antiseptic, and has also alterative properties. It has been given in indigestion in horses, in flatulent dyspepsia in dogs, and in tympanitis in cattle. For influenza in horses it is in some cases advised by Mr. J. B. Gresswell, and also for tabes mesenterica in cattle. Professor Macqueen[2] says the sulphocarbolates have certainly not fulfilled the expectations formed of them when they were introduced thirty years ago as antiseptics.

**Doses.**—Horse, ½ to 1½ drachm.—Cattle, 1 to 2½ drachms.—Dog, 3 to 12 grains.

[1] Gresswell's 'Veterinary Pharmacopœia.'

[2] 'Veterinary Medicines.' (Finlay Dunn.) Fifth edition By James Macqueen, F.R.C.V.S.

# APPENDIX

## SYMBOLS AND ATOMIC WEIGHTS OF THE ELEMENTARY BODIES MENTIONED IN THIS WORK.[1]

| Elementary Bodies. | Symbols. | Atomic Weights. |
|---|---|---|
| Aluminium | Al | = 27 |
| Antimony (Stibium) | Sb | = 120 |
| Arsenic | As | = 75 |
| Barium | Ba | = 137 |
| Bismuth | Bi | = 207 |
| Boron | B | = 11 |
| Bromine | Br | = 80 |
| Cadmium | Cd | = 112 |
| Calcium | Ca | = 40 |
| Carbon | C | = 12 |
| Chlorine | Cl | = 35·5 |
| Chromium | Cr | = 52 |
| Copper (Cuprum) | Cu | = 63·1 |
| Gold (Aurum) | Au | = 196 |
| Hydrogen | H | = 1 |
| Iodine | I | = 127 |
| Iron (Ferrum) | Fe | = 56 |
| Lead (Plumbum) | Pb | = 207 |
| Magnesium | Mg | = 24 |
| Manganese | Mn | = 55 |
| Mercury (Hydrargyrum) | Hg | = 200 |
| Nitrogen | N | = 14 |
| Oxygen | O | = 16 |
| Phosphorus | P | = 31 |
| Platinum | Pt | = 195 |
| Potassium (Kalium) | K | = 39 |
| Silver (Argentum) | Ag | = 108 |
| Sodium (Natrium) | Na | = 23 |
| Sulphur | S | = 32 |
| Tin (Stannum) | Sn | = 118 |
| Zinc | Zn | = 65 |

[1] These weights are expressed in round numbers or to the first place of decimals only.

# WEIGHTS AND MEASURES OF THE 'BRITISH PHARMACOPŒIA'[1]

## WEIGHTS

| | | | | |
|---|---|---|---|---|
| 1 Grain | gr. | | | |
| 1 Ounce | oz. | | = | 437·5 grains |
| 1 Pound | lb. | = 16 ounces | = | 7000 ,, |

## MEASURES OF CAPACITY

| | | |
|---|---|---|
| 1 Minim | min. | |
| 1 Fluid Drachm | fl. drm. | = 60 minims |
| 1 Fluid Ounce | fl. oz. | = 8 fluid drachms |
| 1 Pint | O. | = 20 fluid ounces |
| 1 Gallon | C. | = 8 pints |

## MEASURES OF LENGTH

1 line = $\frac{1}{12}$ inch
1 inch = $\frac{1}{39 \cdot 1393}$ second pendulum
12 inches = 1 foot
36 ,, = 3 feet = 1 yard

Length of pendulum vibrating seconds of mean time in the latitude of London, in a vacuum at the level of the sea . } 39·1393 inches

## RELATION OF MEASURES TO WEIGHTS

| | | | | |
|---|---|---|---|---|
| 1 Minim is the measure of | | | 0·91 | grain of water |
| 1 Fluid Drachm | ,, | | 54·68 | grains of water |
| 1 Fluid Ounce | ,, | 1 ounce or | 437·5 | ,, |
| 1 Pint | ,, | 1·25 pound or | 8750·0 | ,, |
| 1 Gallon | ,, | 10 pounds or | 70000·0 | ,, |

[1] Although the drachm of sixty grains is discarded by the compilers of the 'British Pharmacopœia,' the Author has retained it in the body of his work in consequence of its being a convenient, commonly employed, and well-known quantity in Veterinary Medicine.

# WEIGHTS AND MEASURES OF THE METRICAL SYSTEM

## WEIGHTS

| | | | |
|---|---|---|---|
| 1 Milligramme | = the thousandth part of one grm. | or 0·001 | grm. |
| 1 Centigramme | = the hundredth „ | 0·01 | „ |
| 1 Decigramme | = the tenth „ | 0·1 | „ |
| 1 Gramme | = the weight of a cubic centimetre of water at 4° C. | 1·0 | „ |
| 1 Decagramme | = ten grammes | 10·0 | grms. |
| 1 Hectogramme | = one hundred grammes | 100·0 | „ |
| 1 Kilogramme | = one thousand grammes | 1000·0 | „ |

## MEASURES OF CAPACITY

| | | | | |
|---|---|---|---|---|
| 1 Millilitre | = 1 cub. centim. | or the mea. of 1 grm. | of water | |
| 1 Centilitre | = 10 „ | 10 grms. | „ | |
| 1 Decilitre | = 100 „ | 100 „ | „ | |
| 1 Litre | = 1000 „ | 1000 „ | (1 kilo.) | |

## MEASURES OF LENGTH

| | | | |
|---|---|---|---|
| 1 Millimetre | = the thousandth part of one metre, | or 0·001 | metre |
| 1 Centimetre | = the hundredth „ | 0·01 | „ |
| 1 Decimetre | = the tenth „ | 0·1 | „ |
| 1 Metre | = the ten-millionth part of a quarter of the meridian of the earth. | | |

## RELATION OF THE WEIGHTS OF THE 'BRITISH PHARMACOPŒIA' TO THE METRICAL WEIGHTS

| | | | |
|---|---|---|---|
| 1 Pound | = | 493·5925 | grammes |
| 1 Ounce | = | 28·3495 | „ |
| 1 Grain | = | 0·0648 | „ |

### RELATION OF MEASURES OF CAPACITY OF THE 'BRITISH PHARMACOPŒIA' TO THE METRICAL MEASURES

| | | | |
|---|---|---|---|
| 1 Gallon | = 4·543487 litres | | |
| 1 Pint | = 0·567936 „ | or 567·936 | cubic centimetres |
| 1 Fluid Ounce | = 0·028396 „ | 28·396 | „ |
| 1 Fluid Drachm | = 0·003549 „ | 3·549 | „ |
| 1 Minim | = 0·000059 „ | 0·059 | „ |

### RELATION OF THE METRICAL WEIGHTS TO THE WEIGHTS OF THE 'BRITISH PHARMACOPŒIA'

| | | |
|---|---|---|
| 1 Milligramme | = | 0·015432 grains |
| 1 Centigramme | = | 0·15432 „ |
| 1 Decigramme | = | 1·5432 „ |
| 1 Gramme | = | 15·432 „ |
| 1 Kilogramme | = 2 lbs. 3 oz. 119·8 grs. or | 15432·348 „ |

### RELATION OF THE METRICAL MEASURES TO THE MEASURES OF THE 'BRITISH PHARMACOPŒIA'

| | | |
|---|---|---|
| 1 Millimetre | = | 0·03937 inches |
| 1 Centimetre | = | 0·39371 „ |
| 1 Decimetre | = | 3·93708 „ |
| 1 Metre | = | 39·37079 „ or 1 yard 3·7 inches |
| 1 Cubic Centimetre | = | 15·432 grain-measures |

1 Litre = 1 pint 15 oz. 2 drs. 11m. or 15432·341 grain-measures

# INDEX

TO

# VETERINARY MEDICINES

ARRANGED ACCORDING TO THEIR

# ACTIONS AND USES

# INDEX

TO

# VETERINARY MEDICINES

*Arranged according to their Actions and Uses*

---

**Adhesives** (*adhæreo*, to stick to).—Agents employed to keep the edges of wounds together.

Collodium.
Collodium Flexile.
Emplastrum Picis.

**Alteratives** (*altero*, to change).—Medicines which gradually change the condition and function of organs from a diseased to a healthy state.

Acidum Arseniosum.
— Nitro-hydrochloricum.
— — Dilutum.
Antimonium Sulphuratum.
— Tartaratum.
Argenti Nitras.
Ferri Iodidum.
Hydrargyri Perchloridum.
— Subchloridum.
Hydrargyrum cum Cretâ.
Iodum.
Liquor Arsenicalis.
Oleum Morrhuæ.
Pilula Hydrargyri et Ferri.
Potassa Sulphurata.
Potassii Iodidum.
Pulvis Antimonialis.
Sodii Bicarbonas.
— Carbonas.
— — Exsiccata.
— Chloridum.
Sulphur Sublimatum.
Sulphuris Iodidum.
Tinctura Iodi.
Unguentum Hydrargyri.
— — Compositum.
— — Iodidi Rubri.
— — Nitratis.
— Iodi.
— — Compositum.
— Sulphuris Iodidi.

**Anæsthetics** (α, *a*, privative, and αἰσθητικός, *aisthēticos*, belonging to αἴσθησις, *aisthēsis*, or sensation).—Agents which cause insensibility to pain.

Æther.
Chloroformum.
Cocaina.
Cold water.
Ice.

**Anodynes** (α, *a*, privative, and ὀδύνη, *odunē*, pain).—Agents which allay or diminish pain.

Æther.
Enema Opii.
Extractum Aconiti.
— Hyoscyami.
Hyoscyami Folia.
Linimentum Aconiti.
— Belladonnæ.
— Camphoræ.
— Chloralis Hydratis.
— Chloroformi.
— Opii.
Opium.
Spiritus Ætheris.
Spiritus Ætheris Nitrosi.
— Camphoræ.
— Chloroformi.
Syrupus Papaveris.
Tinctura Aconiti.
— Belladonnæ.
— Chloroformi Composita.
— Hyoscyami.
— Opii.
Unguentum Aconiti.
— Gallæ.
— — cum Opio.
— Veratriæ.

**Antacids** (ἀντί, *anti*, against, and *acidus*, acid).—Agents which neutralise acidity.

Ammonii Carbonas.
Calcii Carbonas.
Calx.
Creta Præparata.
Liquor Ammoniæ.
— Calcis.
— — Saccharatus.
Magnesia.
Magnesii Carbonas.
Mistura Cretæ.
Potassii Bicarbonas.
— Carbonas.
Sapo Durus.
— Mollis.
Sodii Bicarbonas.
— Carbonas.
— — Exsiccata.
Spiritus Ammoniæ Aromaticus.

**Antemetics** (ἀντί, *anti*, against, and ἐμετικά, *emetika*, from ἐμέω *emeo*, I vomit).—Agents which check or allay vomition.

Acidum Carbolicum.
— Hydrocyanicum.
Bismuthi Subnitratis.
Caryophyllum.
Creosotum.
Magnesia.
Magnesii Carbonas.

**Anthelmintics** (ἀντί, *anti*, against, and ἕλμινς, *helmins*, a worm).—Agents which kill, expel, or prevent the return of worms.

Aloes.
Arecæ Semina.
Cambogia.
Cusso.
Enema Aloes.
— Ætheris.
— Quassiæ.
Enema Sodii Chloridi.
— Terebinthinæ
Filix Mas.
Kamala.
Oleum Terebinthinæ.
Santonin.
Sodii Chloridum.

**Antidotes** (ἀντί, *anti*, against, and δίδωμι, *didōmi*, I give).—Agents which counteract the injurious effects of poisons.

**Antilithics** (ἀντί, *anti*, against, and λίθος, *lithos*, a stone).—Agents which dissolve or prevent the formation of calculi and urinary deposits.

Acidum Hydrochloricum Dilutum.
— Nitricum Dilutum.
Acidum Nitro-hydrochloricum Dilutum.

**Antiparasitics** (ἀντί, *anti*, against, and παράσιτος, *parasitos*, a parasite).—Agents which prevent the attacks of, or destroy, parasites.

Acidum Arseniosum.
— Carbolicum.
— Sulphurosum.
Decoctum Tabaci.
Liquor Calcis Chlorinatæ.
Oleum Anisi.
Sapo Durus.
Sapo Mollis.
Sulphur Sublimatum.
Sulphuris Iodidum.
Unguentum Hydrargyri Ammoniati.
— Sulphuris.
— — Iodidi.

**Antiperiodics** (ἀντί, *anti*, against, and περίοδος, *periodos*, a period). Agents which counteract periodicity in disease—*e.g.*, in intermittent fever.

Acidum Arseniosum.
Cinchonæ Flavæ Cortex.
— Pallidæ Cortex.
— Rubræ Cortex.
Decoctum Cinchonæ Flavæ.
Liquor Arsenicalis.
— Arsenici Hydrochlorici.
Piper Nigrum.
Quininæ Sulphas.
Sodii Salicylas.

**Antiphlogistics** (ἀντί, *anti*, against, and φλέγω, *phlegō*, I burn).—Agents which counteract inflammation. See Sedatives.

**Antipyretics** lower the temperature of the body in fever.

Acidum Benzoicum.
— Carbolicum.
— Salicylicum.
Aconitum.
Antimonial salts.
Antipyrine.
Camphora.
Cinchona alkaloids.
Digitalis.
Opium.
Salicinum.
Sodii Salicylas.
Spiritus Ætheris Nitrosi.
Veratrina.

**Antiseptics** (*ἀντί, anti*, against, and *σηπτικός, sēptikos*, putrefying) Agents which prevent, retard, or arrest putrefaction.

Acidum Arseniosum.
— Aceticum Dilutum.
— Carbolicum.
— Sulphurosum.
Cataplasma Fermenti.
Cerevisiæ Fermentum.
Creosotum.
Glycerinum Acidi Carbolici.
Liquor Acidi Carbolici.
Sodii Chloridi.
Zinci Chloridum.
— Sulphas.

**Antispasmodics** (*ἀντί, anti*, against, and *σπασμός, spasmos*, a spasm). Agents which prevent or allay the irregular muscular contraction called spasm or cramp.

Æther.
Asafetida.
Belladonnæ Folia.
Cannabis Indica.
Chloralis Hydras.
Chlorodynum.
Chloroformum.
Cupri Sulphas.
Enema Asafetidæ.
— Tabaci.
— Terebinthinæ.
Extractum Belladonnæ.
Extractum Hyoscyami.
Ipecacuanha.
Spiritus Ætheris.
— — Nitrosi.
— Ammoniæ Aromaticus.
— — Fœtidus.
— Chloroformi.
Tinctura Asafetidæ.
— Belladonnæ.
— Chloroformi Composita.
— Opii.
Unguentum Belladonnæ.

**Aperients** (*aperio*, to open).—Agents which act gently on the bowels, so as to cause mild purgation. See Laxatives.

**Aromatics** (*ἀρι, ari*, intensely, and *ὄζω, ozō*, to smell).—Agents possessing an agreeable taste and odour.

Anethi Fructus.
Carui Fructus.
Cascarillæ Cortex.
Coriandri Fructus.
Cumini Fructus.
Enema Asafetidæ.

**Aromatics**—*continued.*

Fenugreek.
Infusum Digitalis.
Mistura Cretæ.
Oleum Menthæ Piperitæ.
— Caryophylli.
Pimenta.
Piper Nigrum.
Spiritus Ætheris Nitrosi.
— Chloroformi.
Tinctura Zingiberis.

**Arterial Sedatives.**—Agents which lower the action of the heart and of the vascular system.

Digitalis and its preparations.
Plumbi Acetas.

**Astringents** (*ad*, to, and *stringo*, I bind).—Agents which cause a contraction of muscular tissue, promote the coagulation of albuminous fluids, and check secretion.

Acetum.
Acidum Carbolicum.
— Gallicum.
— Sulphuricum Dilutum.
— Tannicum.
Alumen.
Borax.
Calcii Hydras.
Calx.
Catechu.
— Pallidum.
Chlorodynum.
Cinchonæ Flavæ Cortex.
Creasotum.
Cupri Sulphas.
Ferri Peroxidum Humidum.
Ferri Sulphas.
Galla.
Gossypium.
Kino.
Liquor Ferri Perchloridi Fortior.
— Plumbi Subacetatis.
— — Dilutus.
Myrrha.
Plumbi Acetas.
— Carbonas.
Quercûs Cortex.
Sodii Chloridum.
Tinctura Catechu.
— Gallæ.
Unguentum Gallæ.
— — et Opii.
Zinci Acetas.
— Carbonas.
— Oxidum.
— Sulphas.
Ice and other cold applications.

**Carminatives** (*carmen*, a verse or charm).—Agents which were formerly supposed to possess the power of *charming* away flatulency and pain in the alimentary canal. See Aromatics.

**Cathartics** (*καθαίρω*, *kathairo*, I cleanse).—Agents which excite the bowels to increased peristaltic action, and enable them to eject their contents. See Laxatives, Drastic Purgatives, Hydragogues, and Cholagogues.

**Caustics** (*καίω, kaio*, I burn).—Agents which, by chemical action, destroy the parts to which they are applied. The more powerful caustics produce an eschar (scab or crust), and are hence termed *Escharotics.*

Acidum Aceticum Glaciale.
— Arseniosum.
— Carbolicum.
— Hydrochloricum.
— Sulphuricum.
Alumen Exsiccatum.
Argenti Nitras.
Calx.
Creosotum.
Cupri Sulphas.
Hydrargyri Iodidum Rubrum.
— Oxidum Rubrum.
Hydrargyri Perchloridum.
Liquor Ammoniæ Fortior.
— Antimonii Chloridi.
— Ferri Perchloridi Fortior.
— Hydrargyri Perchloridi.
— Zinci Chloridi.
Potassa Caustica.
Soda Caustica.
Tinctura Ferri Perchloridi.
— Iodidi Composita.
Zinci Chloridum.

**Cholagogues,** *Cholagogue Purgatives* (*χολή, cholē*, bile, and *ἀγωγός, agōgos*, an expeller). Agents which have the reputation of increasing the secreting power of the liver and the flow of bile into the intestines. It is probable, however, that they simply cause an emptying of the gall-bladder. They also induc purgation.

Aloes.
Hydrargyri Subchloridum.
Hydrargyrum cum Cretâ.

**Corrosives.**—See Caustics.

**Counter-irritants.**—See Epispastics, Pustulants, and Rubefacients.

**Demulcents** (*demulceo*, to soften).—See Emollients.

**Deobstruents** (*de*, from, and *obstruo*, I stop up).—Agents which remove any obstruction in the body.

Cambogia.
Iodum.
Magnesii Sulphas.
Potassii Iodidum.

**Deodorisers** (*de*, from, and *odor*, a scent).—Agents which absorb or decompose fetid effluvia. See Antiseptics and Disinfectants.

**Desiccants** (*desicco*, to dry up).—Agents which dry up mucous discharges and purulent secretions from ulcers and wounds.

Calcii Carbonas.
— Hydras.
Calx.
Catechu.
Creta Præparata.
Liquor Calcis.

**Desiccants**—*continued.*

Magnesii Carbonas.
Plumbi Acetas.
— Subacetas.
Unguentum Plumbi Acetatis.
Unguentum Plumbi Subacetatis.
— Potassæ Sulphuratæ.
— Zinci.
Zinci Oxidum.

**Diaphoretics** (διαφορέω, *diaphoreo*, to throw off).—Agents which increase the action of the skin, and produce sweating, but in a milder degree than sudorifics.

Æther.
Alcohol.
Amyl Nitris.
Antimonium Tartaratum.
Ipecacuanha.
Jaborandi.
Liquor Ammonii Acetatis.
Physostigmine.
Spiritus Ætheris Nitrosi.

**Disinfectants** (*dis*, indicating separation, and *inficio*, I infect).—Agents which render infectious matter inert.

Acidum Arseniosum.
— Carbolicum.
— Nitricum.
— Sulphurosum.
Calx Chlorinata.
Carbo Ligni.
Cataplasma Calcis Chlorinatæ.
— Carbonis.
Liquor Calcis Chlorinatæ.
Potassii Permanganas.
Vapor Chlori.
Zinci Chloridum.

**Diuretics** (διά, *dia*, through, and οὐρέω, *oureo*, I make water).—Agents which cause an increased secretion and discharge of urine.

Colchici Cormus.
— Semina.
Digitalis Folia.
Extractum Digitalis.
Liquor Ammonii Acetatis.
Magnesii Sulphas.
Oleum Terebinthinæ.
Pix Burgundica.
Potassii Acetas.
— Bicarbonas.
Potassii Carbonas.
— Chloras.
— Nitras.
Resina.
Sodii Acetas.
— Bicarbonas.
— Carbonas.
Spiritus Ætheris Nitrosi.
Terebinthina Canadensis.
Tinctura Digitalis.

**Drastic Purgatives** (δραστικός, *drastikos*, effective, powerful).—Violent cathartics, which produce speedy and copious evacuations.

Cambogia.
Jalapa.
Oleum Crotonis.
Oleum Terebinthinæ.
Rhamni Succus.
Terebinthina Canadensis.

**Ecbolics** (ἐκ, *ek*, out of, and βάλλω, *ballo*, I throw). Agents which cause the uterus to contract and expel its contents.

Ergota.
Infusum Ergotæ.

**Emetics** (ἐμετικά, *emetika*, from ἐμέω, *emeo*, I vomit).—Agents which cause vomition.

Antimonium Tartaratum.
Cupri Sulphas.
Ipecacuanha.
Zinci Acetas.
— Sulphas.

**Emollients** (*emollio*, to soften). Agents which soften the part to which they are applied, and diminish irritation. When used internally to protect the mucous membranes of the alimentary canal from the action of irritants, they are termed *Demulcents*.

Acaciæ Gummi.
Adeps Præparatus.
Albumen Ovi.
Amylum.
Cera Flava.
Collodium.
— Flexile.
Glycerinum.
Infusum Lini.
Lini Farina.
— Semina.
Mucilago Acaciæ.
Oleum Olivæ.
Sevum Præparatum.

**Epispastics** (ἐπισπάω, *epispao*, to draw to).—Agents which, when applied to the skin, produce a vesicle or blister.

Acetum Cantharidis.
Acidum Aceticum Glaciale.
Cantharis.
Linimentum Cantharidis.
Linimentum Crotonis.
Mylabris.
Unguentum Cantharidis.
— Mylabridis.

**Escharotics** (ἐσχαρόω, *escharoō*, to scab over).—See Caustics.

**Excitants** (*excitans*, exciting, stimulating).—See Stimulants.

**Febrifuges** (*febris*, a fever, and *fugo*, to drive away).—Agents which moderate or abate the violence of fevers.

**Hæmatinics** (αἱμάτινα, *hæmatina*, the red colouring matter of the blood).—Preparations which increase the number of red corpuscles in the blood.

Cambogia.
Ferri Carbonas Saccharata.
— Iodidum.
— Peroxidum Humidum.
Ferri Sulphas.
Manganesii Oxidum Nigrum.
Pilula Hydrargyri et Ferri.

**Hæmostatics** (αἷμα, *haima*, blood, and στάσις, *stasis*, a standing).—See Styptics.

**Hydragogues** (ὕδωρ, *hudor*, water, and ἀγωγός, *agōgos*, expeller).—Cathartics which have the power of causing a very large secretion of fluid from the mucous membrane of the intestines. See Drastic Purgatives.

**Hypnotics** (ὕπνος, *hupnos*, sleep).—See Narcotics.

**Irritants** (*irrito*, to excite).—Agents which stimulate and irritate the skin or other parts to which they are applied. See derivatives, Rubefacients, Epispastics, and Pustulants.

**Laxatives** (*laxo*, to loosen).—Mild cathartics.

Aloes.
Jalapa.
Magnesia.
Magnesii Carbonas.
— Sulphas.
Oleum Lini.
— Olivæ.
Oleum Ricini.
Potassii Sulphas.
Rhamni Succus.
Rhei Radix.
Sodii Sulphas.
Sulphur Sublimatum.
Theriaca.

**Lithontriptics** (λίθος, *lithos*, a stone, and τρίβω, *tribo*, to rub down).—See Antilithics.

**Narcotics** (νάρκη, *narkē*, stupor).—Agents which allay pain and produce sleep.

Chloroformum.
Opium.
Tinctura Opii.

**Nutritives** (*nutrio*, to nourish).—Agents which facilitate the assimilative process, and improve the condition of the tissues.

Oleum Lini.
— Morrhuæ.
Oleum Olivæ.
Sevum Præparatum.

**Purgatives** (*purgo*, to cleanse).—See Laxatives.

**Pustulants** (πῦον, *puon*, pus).—Agents which, when applied to the skin, produce a pustule.

Antimonium Tartaratum.
Argenti Nitras.
Oleum Crotonis.

**Refrigerants** (*refrigero*, to cool).—Agents which diminish heat lower the circulation, and quench thirst.

Acetum.
Acidum Hydrochloricum Dilutum.
— Nitricum Dilutum.
— Sulphuricum Dilutum.
Ammonii Chloridum.
Potassii Chloras.
— Nitras.
— Permanganas.
— Liquor Ammonii Acetatis.
Spiritus Ætheris Nitrosi.

**Resolvents** (*resolvo*, I unloose).—Agents having the power to dissolve or relax.

**Rubefacients** (*rubefacio*, to make red).—Agents which, when applied to the skin, irritate and redden it.

Acetum Cantharidis.
Cataplasma Sinapis.
Hydrargyri Perchloridum.
Iodum.
Liquor Ammoniæ (weak solution).
Linimentum Ammoniæ.
— — Compositum.
Linimentum Camphoræ Compositum.
Oleum Terebinthinæ.

**Sedatives** (*sedo*, to allay).—Agents which depress nervous power without previously exalting it.

Acidum Carbolicum.
— Hydrocyanicum Dilutum.
Aconiti Folia.
— Radix.
Antimonii Oxidum.
Antimonium Tartaratum.
Belladonnæ Folia.
Camphora.
Colchici Cormus.
— Semina.
Creosote.
Digitalis Folia.
Extractum Aconiti.
— Belladonnæ.
Hydrargyri Subchloridum.
Hyoscyami Folia.
Ipecacuanha.
Liquor Plumbi Subacetatis.
Opium.
Plumbi Acetas.
Spiritus Camphoræ.
Syrupus Papaveris.
Tinctura Aconiti.
— Camphoræ Composita.
— Opii.
Zinci Oxidum.
Ether spray.

**Sialagogues** (σίαλον, *siălon*, saliva, and ἄλω, *ago*, I lead).—Agents which promote or increase the secretion of saliva.

Antimonium Tartaratum.
Hydrargyri Subchloridum.
Hydrargyrum cum Cretâ.
Iodum.
Potassii Iodidum.
Sinapis.
Zingiber.

**Soporifics** (*sopor*, a heavy sleep, and *fero*, to bring).—See Narcotics.

**Stimulants** (*stimulo*, to excite).—Agents which rapidly, but transiently, excite the nervous system to increased activity, and

**Stimulants**—*continued.*

which, as a consequence, induce a more energetic performance of the functions of the whole body.

Ammonii Carbonas.
— Chloridum.
Acidum Aceticum Dilutum.
— Hydrochloricum Dilutum.
Arnicæ Radix.
Asafetida.
Cantharis.
Capsici Fructus.
Cardamomum.
Catechu Pallidum.
Chloralis Hydras.
Chloroformum.
Gentianæ Radix.
Infusum Cinchonæ Flavæ.
Infusum Gentianæ Compositum.
Nux Vomica.
Pimenta.
Rhei Radix.
Sinapis.
Spiritus Chloroformi.
— Rectificatus.
— — Tenuior.
Tinctura Aloes Composita.
— Chloroformi Composita.
— Cinchonæ Flavæ.
— Gentianæ Composita.
Zingiber.
Alcoholic beverages.

**Styptics** (στύφω, *stupho*, I restrain).—Agents which arrest bleeding.

Acidum Aceticum Dilutum.
— Carbolicum.
— Gallicum.
— Tannicum.
Alumen.
Catechu.
Collodium Hæmostaticum.
Cupri Sulphas.
Ergota.
Ferri Sulphas.
Galla.
Gossypium.
Infusum Ergotæ.
Liquor Calcis.
— Ferri Perchloridi Fortior.
Liquor Plumbi Subacetatis.
Myrrha.
Plumbi Acetas.
— Carbonas.
Quercûs Cortex.
Spiritus Rectificatus.
Tinctura Ferri Perchloridi.
— Gallæ.
— Myrrhæ.
Zinci Acetas.
— Carbonas.
— Oxidum.
— Sulphas.
Cold applications.

**Sudorifics** (*sudor*, to sweat, and *fio*, to become).—Agents which increase the exhalant functions of the skin, and produce sweating.

Ammonii Carbonas.
Antimonii Oxidum.
Camphora.
Hyoscyami Folia.
Ipecacuanha.
Liquor Ammoniæ.
— Ammonii Acetatis.
Oleum Terebinthinæ.

**Sudorifics**—*continued.*

Potassii Nitras.
Pulvis Antimonialis.
— Ipecacuanhæ Compositus.
Spiritus Ætheris Nitrosi.
Spiritus Camphoræ.
Sulphur Præcipitatum.
Terebinthina Canadensis.

**Tonics** (τείνω, *teino*, to stretch ; the quality of muscular fibre in an active state).—Agents which gradually improve the condition and functions of the digestive organs, so as to enable the system to acquire increased vigour.

Acidum Hydrochloricum Dilutum.
— Nitricum Dilutum.
— Sulphuricum Dilutum.
Aloes.
Anthemidis Flores.
Argenti Nitras.
Cerevisiæ Fermentum.
Cinchonæ Flavæ Cortex.
Cupri Sulphas.
Decoctum Cinchonæ Flavæ.
Ferri Carbonas Saccharata.
— Iodidum
— Peroxidum Hydratum.
— Sulphas.
Gentianæ Radix.
Infusum Cinchonæ Flavæ.
— Gentianæ Compositum.
Iodum.
Liquor Ferri Perchloridi.
Quininæ Sulphas.
Rhei Radix.
Sodii Chloridum.
Tinctura Cantharidis.
— Cinchonæ Flavæ.
— Ferri Perchloridi.
— Gentianæ Composita.
— Myrrhæ.
— Nucis Vomicæ.
Vinum Ferri.
Zinci Acetas.
— Sulphas.

**Traumatics** (τραῦμα, *trauma*, a wound).—Agents applied to wounds.

Collodium.
Myrrha.
Resina.
Tinctura Aloes Composita.
Unguentum Terebinthinæ.
Zinci Oxidum.

**Vermicides** (*vermis*, a worm, and *cædo*, to kill).—See Anthelmintics.

**Vermifuges** (*vermis*, a worm, and *fugo*, to drive away).—See Anthelmintics.

**Vesicants** (*vas*, a vessel, a bladder).—See Epispastics.

# INDEX


| | PAGE |
|---|---|
| Acacia Gum | 1 |
| — — Mucilage of | 202 |
| Acaciæ Gummi | 1 |
| — Mucilago | 202 |
| Acetanilide | 1 |
| Acetic Acid | 4 |
| — — Diluted | 5 |
| — — Glacial | 6 |
| — — Hydrate | 6 |
| Acetum | 2 |
| — Cantharidis | 3 |
| Acid, Acetic | 4 |
| — — Dilute | 5 |
| — — Glacial | 6 |
| — Arsenious | 6 |
| — Boracic | 8 |
| — Boric | 8 |
| — Carbolic | 8 |
| — — Glycerin of | 128 |
| — — Liniment of | 149 |
| — — Lotion of | 185 |
| — — Ointment of | 304 |
| — — — Compound, of | 304 |
| — — Solution of | 157 |
| — Chromic | 11 |
| — Gallic | 12 |
| — Hydrochloric | 13 |
| — — Dilute | 14 |
| — Hydrocyanic, Dilute | 15 |
| — Muriatic | 13 |
| Acid, Nitric | 17 |
| — — Dilute | 18 |
| — Nitro-hydrochloric | 19 |
| — Phosphoric, Concentrated | 20 |
| — — Dilute | 21 |
| — Prussic | 15 |
| — — Scheele's | 15 |
| — Pyroligneous | 5 |
| — Salicylic | 21 |
| — Sulpho-carbolic | 22 |
| — Sulphuric | 23 |
| — — Dilute | 24 |
| — Sulphurous | 26 |
| — Tannic | 27 |
| — — Ointment of | 304 |
| Acidi Carbolici Glycerinum | 128 |
| — — Linimentum | 149 |
| — — Liquor | 157 |
| — — Lotio | 185 |
| — — Unguentum | 304 |
| — — — Compositum | 304 |
| — Tannici Unguentum | 304 |
| Acidum Aceticum | 4 |
| — — Dilutum | 5 |
| — — Glaciale | 6 |
| — Arseniosum | 6 |
| — Boricum | 8 |
| — Carbolicum | 8 |
| — Chromicum | 11 |
| — Gallicum | 12 |

| | PAGE |
|---|---|
| Acidum Hydrochloricum | 13 |
| — — Dilutum | 14 |
| — Hydrocyanicum Dilutum | 15 |
| — Nitricum | 17 |
| — — Dilutum | 18 |
| — Nitro-hydrochloricum Dilutum | 19 |
| — Phosphoricum Concentratum | 20 |
| — — Dilutum | 21 |
| — Salicylicum | 22 |
| — Sulphuricum | 23 |
| — — Dilutum | 24 |
| — Sulphurosum | 26 |
| — Tannicum | 27 |
| Aconite, Extract of | 111 |
| — Leaves | 29 |
| — Ointment of | 305 |
| — Root | 30 |
| — Tincture of | 291 |
| Aconiti, Extractum | 111 |
| — Folia | 29 |
| — Radix | 30 |
| — Tinctura | 291 |
| — Unguentum | 305 |
| Aconitia | 30 |
| Aconitina | 30 |
| Aconitine | 30 |
| Adeps Præparatus | 31 |
| Æruginis, Unguentum | 307 |
| — Linimentum | 153 |
| Ærugo | 100 |
| Æther | 32 |
| Ætheris, Spiritus | 274 |
| — Nitrosi, Spiritus | 275 |
| — Sulphurici, Spiritus | 274 |
| Albumen, Egg | 34 |
| — Ovi | 34 |
| Alcohol, Ethylic | 280 |
| Allspice | 223 |
| Aloe Barbadensis | 34 |
| Aloe Capensis | 35 |
| — Socotrina | 36 |
| Aloes, Barbados | 34 |
| — Cape | 35 |
| — Enema of | 107 |
| — Liquor | 158 |
| — — et Sodæ Carbonatis | 158 |
| — Massa | 192 |
| — — Composita | 193 |
| — Socotrine | 36 |
| — Solution of | 158 |
| — — and Carbonate of Soda | 158 |
| — Tinctura Composita | 292 |
| — Tincture of, Compound | 292 |
| Aloin | 37 |
| Alterative Mass | 193 |
| Alum | 37 |
| — Ammonium | 37 |
| — Burned | 39 |
| — Compound Ointment of | 305 |
| — Dried | 39 |
| — Ointment of, Compound | 305 |
| — Potassium | 37 |
| — Solution of | 159 |
| Alumen | 37 |
| — Exsiccatum | 39 |
| Aluminis, Liquor | 159 |
| — Unguentum Compositum | 305 |
| Amido-chloride of Mercury | 137 |
| Ammonia Acetate, Solution of | 160 |
| — — Lotion of | 185 |
| — Aromatic Spirit of | 277 |
| — Carbonate of | 40 |
| — Chloride of | 42 |
| — Fetid Spirit of | 277 |
| — Mixture | 197 |
| — Muriate of | 42 |
| — Solution of | 159 |
| — Strong Solution of | 162 |

| | PAGE |
|---|---|
| Ammoniacum | 39 |
| — Gum | 39 |
| Ammoniæ Acetatis, Liquor | 160 |
| — — Lotio | 185 |
| — Aromaticus Spiritus | 277 |
| — Carbonas | 40 |
| — Fœtidus, Spiritus | 277 |
| — Fortior, Liquor | 162 |
| — Linimentum | 150 |
| — Liquor | 159 |
| — Mistura | 197 |
| Ammoniated Mercury | 137 |
| — Mustard Poultice | 78 |
| Ammonii Chloridi et Camphoræ, Lotio | 186 |
| — — et Potassæ Nitratis, Lotio | 186 |
| — Chloridum | 42 |
| Ammonium Alum | 37 |
| — Chloride | 42 |
| — — Lotion of, and Potassium Nitrate | 186 |
| — — and Camphor | 186 |
| Amyl Nitris | 43 |
| — Nitrite | 43 |
| Amyli Mucilago | 202 |
| Amylum | 44 |
| Anhydride, Arsenious | 6 |
| — Sulphurous | 26 |
| Anise Fruit | 45 |
| — Oil of | 205 |
| Aniseed | 45 |
| Anisi Fructus | 45 |
| — Oleum | 205 |
| Anthemidis Flores | 46 |
| Antichlor | 268 |
| Antifebrin | 1 |
| Antimonii Chloridi, Liquor | 163 |
| — Oxidum | 46 |
| — Tartarati Composita, Massa | 193 |
| — — Liquor | 164 |

| | PAGE |
|---|---|
| Antimonii Tartarati, Unguentum | 305 |
| Antimonious Chloride, Solution of | 163 |
| — Oxide | 46 |
| — Sulphide | 48 |
| Antimonium Nigrum | 48 |
| — Tartaratum | 49 |
| Antimony, Black | 48 |
| — Butter of | 163 |
| — Chloride of, Solution of | 163 |
| — Ointment of Tartarated | 305 |
| — Oxide of | 46 |
| — Solution of Chloride of | 163 |
| — — Tartarated | 164 |
| — — Terchloride of | 163 |
| —Tartarated | 49 |
| — — Compound Mass of | 193 |
| — — Solution of | 164 |
| — Terchloride, Solution of | 163 |
| — Teroxide | 46 |
| — Tersulphide | 48 |
| — Trichloride | 163 |
| — Trioxide | 46 |
| — Trisulphide | 48 |
| Antipyrin | 51 |
| Apomorphine | 201 |
| Aqua Calcis | 168 |
| — Destillata | 51 |
| — Fortis | 17 |
| — Menthæ Piperitæ | 52 |
| — Regia, Dilute | 19 |
| Areca Nut | 52 |
| — Seeds | 52 |
| Arecæ Semina | 52 |
| Argenti Nitras | 53 |
| — Nitratis, Liquor | 165 |
| — — Unguentum | 306 |
| Arnica Root | 55 |
| — Tincture of | 293 |
| Aromatic Spirit of Ammonia | 277 |

PAGE

Arsenic, Hydrochloric Solution of . . . 167
— Ointment of . . . 306
— White . . . . 6
Arsenical Solution . . . 165
Arsenicalis, Liquor . . 165
Arsenici Hydrochloricus, Liquor . . . . 167
— Unguentum . . . 306
Arsenious Acid . . . 6
Anhydride . . . . 6
Asafetida . . . . 56
— Enema of . . . . 108
Asafetidæ, Enema . . . 108
Aseptol . . . . . 23
Astringent Mass . . . 194
Atropina . . . . 57
Atropine . . . . 57
Axungia . . . . 31

Balls (*see* Mass)
Balsam, Canada . . . 289
— of Peru . . . . 57
Balsamum Peruvianum . . 57
Barbados Aloes . . . 34
Bean, Calabar . . . 221
Belladonna, Compound Mass of . . . . 194
— Extract of . . . 112
— Leaves . . . . 58
— Root . . . . . 59
Belladonnæ, Extractum . 112
— Folia . . . . . 58
— Massa Composita . . 194
— Radix . . . . 59
Benzoin . . . . . 60
Benzoinum . . . . 60
Benzol . . . . . 60
Betel Nut . . . . 52

PAGE

Black Antimony . . . 48
— Hellebore, Ointment of . 308
— Mercurial Lotion . . 187
— Pepper . . . . 224
— Pitch . . . . 227
— Wash . . . . 187
Blistering Fly . . . 203
— — Chinese . . . 70
— Ointment . . . 307, 310
Blue Ointment . . . 309
— Pill, Ferruginate . . 223
— Stone . . . . 101
— Vitriol . . . . 101
Boli (*see* Mass)
Borax . . . . . 61
Boroglyceride . . . 8
Bran Poultice . . . 77
British Gum . . . . 202
Buchu Folia . . . . 62
— Leaves . . . . 62
Buckthorn Juice . . . 284
— Syrup of . . . . 284
Burgundy Pitch . . . 225
Burnett's, Sir William, Fluid . 163
Butter of Antimony . . 163

Calabar Bean . . . 221
Calamine . . . . 323
Calcii Hydras. . . . 63
Calcis, Aqua . . . . 168
— Chlorinatæ, Liquor . . 169
— — Unguentum . . . 306
— Linimentum . . . 151
— Liquor . . . . 168
Calcium Carbonate . . 96
— Hydrate . . . . 63
— Hypochlorite . . . 66
— Oxide . . . . 65
— Oxide of . . . 65

PAGE

Calomel . . . . . 133
Calumba Root . . . 64
Calumbæ Radix . . . 64
Calves', Sheep and, Cordial . 198
Calx . . . . . 65
— Chlorinata . . . 66
Cambogia . . . . 67
Camphor . . . . 68
— Compound Liniment of . 152
— Liniment of . . . 151
— Lotion of Chloride of Ammonium and . . 186
— Spirit of . . . . 278
Camphora . . . . 68
Camphoræ, Linimentum . 151
— — Compositum . . . 152
— Lotio Ammonii Chloridi et . . . . 186
— Spiritus . . . . 278
— Tinctura . . . . 294
Camphorated Oil . . . 207
Canada Balsam . . . 289
Cantharides . . . . 70
— Liniment of . . . 152
— Oil of . . . . 207
— Ointment of . . . 307
— Solution of, Terebinthinated . . . . 170
— Strong Tincture of . . 295
— Terebinthinated Solution of . . . . 170
— Tincture of . . . 294
— — Strong . . . . 295
— Vinegar of . . . 3
Cantharidin . . . . 70
Cantharidis Acetum . . 3
— Linimentum . . . 152
— Oleum . . . . 207
— Terebinthinatus, Liquor . 170
— Tinctura . . . . 294
— — Fortior . . . . 295

PAGE

Cantharidis Unguentum . 307
Cantharis . . . . 70
— Vesicatoria . . . 70
Capsici Fructus . . . 72
Capsicine . . . . 72
Capsicum Fruit . . . 72
Caraway Fruit . . . 74
— Seeds . . . . 74
Carbo Ligni . . . . 72
Carbolic Acid . . . . 8
— — Glycerin of . . . 128
— — Liniment of . . . 149
— — Lotion of . . . 185
— — Ointment of . . . 304
— — — Compound . . 304
— — Solution of . . . 157
Cardamomi, Tinctura, Composita . . . . 295
Cardamoms . . . . 74
— Tincture of, Compound . 295
Cardamomum . . . 74
Carron Oil . . . . 151
Carui Fructus . . . 74
Caryophyllum . . . 75
Cascarilla Bark . . . 75
Cascarillæ Cortex . . . 75
Cascarillin . . . . 75
Castor Oil . . . . 214
— — Mixture . . . 199
Cataplasma Calcis Chlorinatæ 76
— Carbonis . . . . 76
— Fermenti . . . . 77
— Furfuris . . . . 77
— Lini . . . . . 78
— Sinapis . . . . 78
— — Ammoniatum . . 79
Catechu Composita, Massa . 194
— Infusion of . . . 140
— Infusum . . . . 140
— Pale . . . . . 79
— Pallidum . . . . 79

| | PAGE |
|---|---|
| Cathartic Mass | 192 |
| Caustic Potash | 231 |
| — Soda | 263 |
| Cera Flava | 80 |
| Cerevisiæ Fermentum | 81 |
| Cerii Oxalas | 81 |
| Chalk | 96 |
| — Mercury with | 138 |
| — Mixture, Compound | 198 |
| — Prepared | 96 |
| Chamomile Flowers | 46 |
| Charcoal Poultice | 77 |
| — Wood | 72 |
| Chinese Blistering Fly | 203 |
| Chloral Hydrate | 82 |
| — Hydras | 82 |
| Chloric Ether | 278 |
| Chlorinated Lime | 66 |
| — — Poultice | 76 |
| Chlorine Gas | 317 |
| Chlorodyne | 83 |
| Chloroform | 84 |
| — Mixture, Compound | 83 |
| — Spirit of | 278 |
| — Tincture of, Compound | 296 |
| Chloroformi, Mistura, Hydrocyanata | 83 |
| — Spiritus | 278 |
| — Tinctura, Composita | 296 |
| Chloroformum | 84 |
| Cinchona Bark, Pale | 89 |
| — — Red | 89 |
| — — Yellow | 86 |
| — Infusion of Yellow | 140 |
| Cinchonæ Flavæ Cortex | 86 |
| — — Infusum | 140 |
| — Pallidæ Cortex | 89 |
| — Rubræ Cortex | 89 |
| Cinnamomum | 89 |
| Cinnamon | 89 |
| Citrine Ointment | 311 |
| Cloves | 75 |
| Cocaine | 90 |
| Cod-liver Oil | 212 |
| Colchici Cormus | 91 |
| — Semina | 92 |
| — Seminum, Tinctura | 297 |
| Colchicum Corm | 91 |
| — Seeds | 92 |
| — — Tincture of | 297 |
| Collodion | 92 |
| — Flexible | 93 |
| — Hæmostatic | 93 |
| Collodium | 92 |
| — Flexile | 93 |
| — Hæmostaticum | 93 |
| Colocynth | 94 |
| Colocynthis | 94 |
| Common Mass | 194 |
| — Salt | 272 |
| Compound Chalk Mixture | 198 |
| — Infusion of Gentian | 141 |
| — Liniment of Ammonia | 150 |
| — — of Creosote | 153 |
| — — of Iodine | 155 |
| — — of Soap | 156 |
| — — of Tar | 155 |
| — — of Turpentine | 157 |
| — Mass of Aloes | 193 |
| — — of Belladonna | 194 |
| — — of Catechu | 194 |
| — — of Digitalis | 195 |
| — — of Ginger | 197 |
| — — of Resin | 196 |
| — — of Tartarated Antimony | 193 |
| — Ointment of Alum | 305 |
| — — of Carbolic Acid | 304 |
| — — of Iodine | 312 |
| — — of Mercury | 310 |
| — — of Sulphur | 315 |
| — Powder of Carbonate of Zinc | 251 |

PAGE

Compound Powder of Ipecacuanha . . . 249
— — of Opium . . . 250
— — of Rhubarb . . . 250
— — of Sulphate of Iron . 249
— Solution of Sulphate of Copper . . . 171
— Spirit of Perchloride of Mercury . . . 279
— Tincture of Aloes . . 292
Confectio Rosæ Gallicæ . . 94
Confection of Roses . . 94
Conii Fructus, Tinctura . 297
— Tinctura . . . . 297
Cooling Lotion . . . 186
Copper, Compound Solution of Sulphate of . . 171
— Iodide of . . . . 99
— — with Iodine . . . 99
— Liniment of Subacetate of 153
— — of Sulphate of . . 154
— Mass of Sulphate of . . 195
— Subacetate of . . . 100
— — Liniment of . . . 153
— Sulphate of . . . 101
— — Compound Solution of . 171
— — Liniment of . . . 154
— — Mass of . . . 195
— — Solution of . . . 171
Copperas . . . . 101
Cordial Mass . . . . 197
— Sheep and Calves' . . 198
Coriander Fruit . . . 94
— Seed . . . . . 94
Coriandri Fructus . . . 94
Corrosive Sublimate . . 131
Cotton Wool . . . . 128
Cough Ball . . . . 195
Creosote . . . . . 95
— Compound Liniment of . 153
— Mixture . . . . 198

PAGE

Creosote, Ointment of . . 307
Creosoti Compositum, Linimentum . . . 153
— Unguentum . . . 307
Creosotum . . . . 95
Creta . . . . . 96
— Hydrargyrum cum . . 138
— Præparata . . . . 96
Cretæ Composita, Mistura . 198
Crocus . . . . . 97
Croton Beans . . . 97
— Oil . . . . . 208
— — Liniment of . . . 153
— Seeds . . . . 97
— Tincture of . . . 298
Crotonis, Linimentum . . 153
— Oleum . . . . 208
— Semina . . . . 97
— Tiglii Semina . . . 97
— Tinctura . . . . 298
Cumin Fruit . . . . 98
Cumini Fructus . . . 98
Cupri Iodidum cum Iodo . 99
— Subacetas . . . . 100
— Subacetatis, Linimentum . 153
— — Unguentum . . . 307
— Sulphas . . . . 101
— Sulphatis, Compositus, Liquor . . . . 171
— — Linimentum . . . 153
— — Liquor . . . . 171
— — Massa . . . . 195
Cupric Sulphate . . . 101
Cusso . . . . . 104

Decoction of Tobacco . . 105
Decoctum Tabaci . . . 105
Digestive Ointment . . 314
Digitalin . . . . 105
Digitalinum . . . . 105

| | PAGE |
|---|---|
| Digitalis Composita, Massa | 195 |
| — Extract of | 113 |
| — Extractum | 113 |
| — Folia | 105 |
| — Leaves | 105 |
| Dinitro-cellulose | 251 |
| Dip, Sheep | 166 |
| Discutient Lotion | 186 |
| Distilled Water | 51 |
| Diuretic Mass | 196 |
| Dover's Powder, Factitious | 249 |
| Egg Albumen | 34 |
| Emetic, Tartar | 49 |
| Emplastrum Picis | 107 |
| Enema Aloes | 107 |
| — of Aloes | 107 |
| — of Asafetida | 108 |
| — Asafetidæ | 108 |
| — Magnesiæ Sulphatis | 108 |
| — Opii | 108 |
| — of Opium | 108 |
| — of Sulphate of Magnesia | 108 |
| — Tabaci | 108 |
| — Terebinthinæ | 109 |
| — of Tobacco | 108 |
| — of Turpentine | 109 |
| Epsom Salts | 190 |
| Ergot | 109 |
| — Infusion of | 141 |
| — Liquid Extract of | 114 |
| — of Rye | 109 |
| — Tincture of | 299 |
| Ergota | 109 |
| Ergotæ, Extractum, Liquidum | 114 |
| — Infusum | 141 |
| — Tinctura | 299 |
| Eserine | 221 |
| Ether | 32 |
| — Chloric | 278 |
| Ether, Ethylic | 32 |
| — Spirit of | 274 |
| — — of Nitrous | 275 |
| — Sulphuric | 32 |
| Ethylic Ether | 32 |
| — Oxide | 32 |
| Eucalyptus | 110 |
| Euphorbium | 110 |
| Extract of Aconite | 111 |
| — of Belladonna | 112 |
| — of Digitalis | 113 |
| — of Indian Hemp | 113 |
| — Liquid, of Ergot | 114 |
| — of Male Fern | 115 |
| — Goulard's | 177 |
| — of Hyoscyamus | 115 |
| — of Physostigmine | 221 |
| Extractum Aconiti | 111 |
| — Belladonnæ | 112 |
| — Cannabis Indicæ | 113 |
| — Digitalis | 113 |
| — Ergotæ Liquidum | 114 |
| — Filicis Liquidum | 115 |
| — Hyoscyami | 115 |
| — Physostigmatis | 221 |
| Factitious Dover's Powder | 249 |
| — James's Powder | 249 |
| Fecula | 44 |
| Fern, Male | 123 |
| — — Liquid Extract of | 115 |
| — — Shield | 123 |
| — Root | 123 |
| Ferri Carbonas Saccharata | 116 |
| — Iodidum | 118 |
| — Perchloridi Fortior, Liquor | 172 |
| — — Tinctura | 299 |
| — Peroxidum Humidum | 119 |
| — — Hydratum | 119 |
| — Persulphatis, Liquor | 173 |

| | PAGE |
|---|---|
| Ferri, Pilula Hydrargyri et | 223 |
| — Protosulphas | 121 |
| — Sesquichloridi, Tinctura | 299 |
| — Sulphas | 121 |
| — Sulphatis Compositus, Pulv. | 249 |
| — — Massa | 196 |
| — Vinum | 321 |
| Ferric Hydrate, Moist | 119 |
| Ferruginate Blue Pill | 223 |
| Ferrum | 122 |
| Fetid Spirit of Ammonia | 277 |
| Fever Ball | 193 |
| — Mass | 193 |
| Filix Mas | 123 |
| Flowers of Sulphur | 284 |
| Fluid, Sir Wm. Burnett's | 183 |
| Fly, Chinese Blistering | 203 |
| Formyl Trichloride | 84 |
| Fowler's Solution | 165 |
| Galbanum | 124 |
| Galla | 124 |
| Gallæ, Unguentum | 308 |
| — — cum Opio | 308 |
| Gallic Acid | 12 |
| Galls | 124 |
| — Nut | 124 |
| — Oak | 124 |
| — Ointment of | 308 |
| — and Opium, Ointment of | 308 |
| Gamboge | 67 |
| Gentian, Compound Infusion of | 141 |
| — Root | 126 |
| Gentianæ Compositum, Infusum | 141 |
| — Radix | 126 |
| Ginger | 330 |
| — Compound Mass of | 197 |
| — Tincture of | 303 |
| Glacial Acetic Acid | 6 |
| Glauber's Salt | 270 |
| Glycerin | 127 |
| — of Carbolic Acid | 128 |
| Glycerinum | 127 |
| — Acidi Carbolici | 128 |
| Gossypium | 128 |
| Goulard's Extract | 176 |
| — Water | 177 |
| Green Vitriol | 121 |
| Gregory's Mixture | 250 |
| — Powder | 250 |
| Grey Powder | 138 |
| Gum Acacia | 1 |
| — — Mucilage of | 1 |
| — Ammoniacum | 39 |
| — Arabic | 1 |
| — Benzoin | 60 |
| — British | 202 |
| — Euphorbium | 110 |
| Gun Cotton | 251 |
| Hard Soap | 260 |
| Healing Ointment | 316 |
| Hellebore, Black, Ointment of | 308 |
| — White | 319 |
| — — Ointment of | 316 |
| Hellebori Nigri, Unguentum | 308 |
| Hemlock, Tincture of | 297 |
| Henbane Leaves | 139 |
| Hepar Sulphuris | 232 |
| Hog's Lard | 31 |
| Hydrargyri Ammoniati, Unguentum | 309 |
| — Ammonio-chloridum | 137 |
| — Bichloridi, Liquor | 174 |
| — Biniodidi, Unguentum | 310 |
| — Flava, Lotio | 187 |
| — Iodidi Rubri, Unguentum | 310 |
| — Iodidum Rubrum | 128 |

PAGE

Hydrargyri Nigra, Lotio . 187
— Nitrico-oxidum . . . 130
— Oxidum Rubrum . . 130
— Perchloridi Compositus, Spiritus . . . 279
— — Liquor . . . . 174
— Perchloridum . . . 131
— Pilula, et Ferri . . . 233
— Præcipitatum Album . 137
— Subchloridum . . . 133
— Unguentum . . . 309
— — Compositum . . . 310
Hydrargyrum . . . 135
— Ammoniatum . . . 137
— cum Cretâ . . . . 138
Hydrate, Acetic . . . 6
Hydrochloric Acid . . . 13
— — Dilute . . . . 14
— Solution of Arsenic . . 167
Hydrocyanic Acid, Dilute . 15
Hydrogen Acetate . . . 6
Hydrogen-Potassium Carbonate . . . . 233
Hydroxybenzoic Acid . . 21
Hyoscyami Folia . . . 139
Hyoscyamus Leaves . . 139

Infusion of Calumba . . 140
— of Catechu . . . 140
— Compound, of Gentian . 141
— of Ergot . . . . 141
— of Kousso . . . . 141
— of Linseed . . . . 142
— of Tobacco . . . 142
— of Yellow Cinchona . . 140
Infusum Calumbæ . . . 140
— Catechu . . . . 140
— Cinchonæ Flavæ . . 140
— Cusso . . . . 141
— Ergotæ . . . . 141

PAGE

Infusum Gentianæ Compositum . . . . 141
— Lini . . . . . 142
— Tabaci . . . . 142
Iodi Compositum, Linimentum . . . . 155
— Linimentum . . . 156
— Liquor . . . . 174
— et Potassii Iodidi, Liquor 174
— Tinctura . . . . 300
— Unguentum . . . 312
— — Compositum . . . 312
Iodine . . . . . 143
— Compound Liniment of . 155
— — Ointment of . . . 312
— Ointment of . . . 312
— — Compound . . . 312
— Solution of, and Iodide of Potassium . . . 174
— Tincture of . . . 300
Iodoform . . . . 142
Iodoformum . . . . 142
Iodol . . . . . 143
Iodum . . . . . 143
Ipecacuanha . . . . 146
— Compound Powder of . 249
— Powder of, Compound . 249
Ipecacuanhæ Compositus, Pulvis . . . . 249
Iron . . . . . 122
— Iodide of . . . . 118
— Mercurial Pill with . . 233
— Moist Hydrated Peroxide of 119
— — Peroxide of . . . 119
— Perchloride of, Strong Solution of . . . 172
— — Tincture of . . . 292
— Pill of Mercury and . . 233
— Protosulphate of . . 121
— Saccharated Carbonate of . . . . 116

PAGE

Iron, Solution of Perchloride of, Strong . . . 172
— Strong Solution of Perchloride of . . . . 172
— Sulphate of . . . 121
— — Compound Powder of . 249
— — Mass of . . . 196
— Tincture of Perchloride of 299
— Wine of . . . . 321

Jalap . . . . . 146
Jalapa . . . . . 146
James's Powder, Factitious . 249
Jecoris Aselli, Oleum . . 212
Juice, Buckthorn . . . 284
Juniper, Oil of . . . 210
Juniperi, Oleum . . . 210

Kamala . . . . . 147
Kamela . . . . . 147
Kino . . . . . 148
Kousso . . . . . 104

Lard, Hog's . . . . 31
— Prepared . . . . 31
Laudanum . . . . 302
Lead Acetate . . . . 227
— Acetate of . . . . 227
— Diluted Solution of Subacetate of . . . 177
— Liniment of Subacetate of 156
— Monoxide . . . . 229
— Protoxide of . . . 229
— Solution of Subacetate . 176
— — — Diluted . . . 177
— — — and Opium . . 178
— Subacetate of, Diluted Solution of . . . 177

PAGE

Lead, Subacetate of, Liniment of . . . . . 156
— — Solution of . . . 176
— Sugar of . . . . 227
Leaf Tobacco . . . . 288
Light Carbonate of Magnesia . 189
— Magnesia . . . . 189
Lime . . . . . 65
— Carbonate of . . . 96
— Chloride of . . . 66
— Chlorinated . . . 66
— Hypochlorite of . . 66
— Liniment of . . . 151
— Slaked . . . . 63
— Solution of . . . 168
— — of Chlorinated . . 169
— Water . . . . 168
Lini Farina . . . . 149
— Infusum . . . . 142
— Oleum . . . . 210
— Semina . . . . 149
Liniment of Aconite . . 150
— of Ammonia . . . 150
— — Compound . . . 150
— Camphor . . . . 151
— — Compound . . . 152
— of Cantharides . . . 152
— of Carbolic Acid . . 149
— Compound, of Ammonia . 150
— — of Iodine . . . 155
— — of Soap . . . 156
— — of Turpentine . . 157
— of Creosote, Compound . 153
— of Croton Oil . . . 153
— of Iodine, Compound . 155
— of Lime . . . . 151
— of Opium . . . . 156
— of Subacetate of Copper . 153
— — of Lead . . . 156
— of Sulphate of Copper . 154
— of Tar . . . . 155

PAGE
Liniment of Tar, Compound . 155
— of Turpentine . . . 157
— of Verdigris . . . 153
Linimentum Acidi Carbolici . . . . 149
— Aconiti . . . . 150
— Æruginis . . . . 153
— Ammoniæ . . . . 150
— — Compositum . . . 150
— Calcis . . . . 151
— Camphoræ . . . . 151
— — Compositum . . . 152
— Cantharidis . . . 152
— Creosoti Compositum . 123
— Crotonis . . . . 153
— Cupri Subacetatis . . 153
— — Sulphatis . . 154
— Hydrargyri Compositum . 154
— Iodi . . . . . 174
— — Compositum . . . 155
— Olei Picis Liquidæ Compositum . . . 155
— Opii . . . . . 156
— Picis Liquidæ . . . 155
— Plumbi Subacetatis . . 156
— Saponis Compositum . . 156
— Terebinthinæ . . . 157
— — Compositum . . . 157
Linseed . . . . . 149
— Infusion of . . . 142
— Meal . . . . . 149
— Oil . . . . . 211
— Poultice . . . . 78
— Tea . . . . . 142
Liquor Acidi Carbolici . . 157
— Aloes . . . . . 158
— — et Sodæ Carbonatis . 158
— Aluminis . . . . 159
— Ammoniæ . . . . 159
— — Acetatis . . . 160
— — Fortior . . . . 162
PAGE
Liquor Antimonii Chloridi . 163
— — Tartarati . . . 164
— Argenti Nitratis . . 165
— Arsenicalis . . . 165
— Arsenici Chloridi . . 167
— — Hydrochloricus . . 167
— Calcis . . . . 168
— — Chlorinatæ . . . 169
— — Iodinatæ . . . 333
— Cantharidis Terebinthinatus . . . . 170
— Cupri Sulphatis . . . 171
— — — Compositus . . 171
— Ferri Perchloridi Fortior . 172
— — Persulphatis . . . 173
— Hydrargyri Perchloridi . 174
— Iodi . . . . . 174
— — et Potassii Iodidi . . 174
— Morphinæ Acetatis . . 175
— — Hydrochloridi . . 176
— Plumbi Subacetatis . . 176
— — — Dilutus . . . 177
— — — et Opii . . . 178
— Potassæ . . . . 178
— — Arsenitis . . . 165
— — Iodinatæ . . . 333
— — Sulphuratæ . . . 180
— Potassii Nitratis . . 180
— — Permanganatis . . 180
— Sodæ . . . . . 181
— Strychninæ Hydrochloridi 182
— Zinci Chloridi . . . 183
— — Sulphatis . . . 185
Litharge . . . . . 229
Liver of Sulphur . . . 232
Lotio Acidi Carbolici . . 185
— Ammoniæ Acetatis . . 185
— Ammonii Chloridi et Camphoræ . . . . 186
— — — et Potassii Nitratis . . . . 186

| | PAGE |
|---|---|
| Lotio Discutiens | 186 |
| — Hydrargyri Flava | 187 |
| — — Nigra | 187 |
| — Refrigerans | 186 |
| Lotion of Acetate of Ammonia | 185 |
| — Black Mercurial | 187 |
| — of Carbolic Acid | 185 |
| — of Chloride of Ammonium and Camphor | 186 |
| — — — and Nitrate of Potash | 186 |
| — Cooling | 186 |
| — Discutient | 186 |
| — Yellow Mercurial | 187 |
| Lugol's Solution | 174 |
| Lunar Caustic | 53 |
| Lytta Vesicatoria | 70 |
| Magnesia | 187 |
| — Alba | 189 |
| — Carbonate of | 189 |
| — Levis | 189 |
| — Light | 189 |
| — — Carbonate of | 189 |
| — Sulphate of | 190 |
| — — Enema | 108 |
| — Usta | 187 |
| Magnesiæ Carbonas | 189 |
| — — Levis | 189 |
| — Sulphas | 190 |
| — Sulphatis, Enema | 108 |
| Magnesium Oxide | 187 |
| — Sulphate | 190 |
| Manganese, Binoxide of | 191 |
| — Black Oxide of | 191 |
| — Dioxide | 191 |
| — Peroxide of | 191 |
| Manganesii Oxidum Nigrum | 191 |
| Manganic Peroxide | 191 |
| Marble, White | 192 |
| Marmor Album | 192 |
| Mass of Aloes | 192 |
| — — Compound | 193 |
| — Alterative | 193 |
| — Astringent | 194 |
| — of Belladonna, Compound | 194 |
| — of Catechu, Compound | 194 |
| — Cathartic | 192 |
| — Common | 194 |
| — Cordial | 197 |
| — of Digitalis, Compound | 195 |
| — Diuretic | 196 |
| — of Ginger, Compound | 197 |
| — of Resin, Compound | 196 |
| — of Sulphate of Copper | 195 |
| — — of Iron | 196 |
| — of Tartarated Antimony, Compound | 193 |
| — Tonic | 196 |
| Massa Aloes | 192 |
| — — Composita | 193 |
| — Antimonii Tartarati Composita | 193 |
| — Belladonnæ Composita | 194 |
| — Catechu Composita | 194 |
| — Communis | 194 |
| — Cupri Sulphatis | 195 |
| — Digitalis Composita | 195 |
| — Ferri Sulphatis | 196 |
| — Resinæ Composita | 196 |
| — Zingiberis Composita | 197 |
| Mercurial Lotion, Black | 187 |
| — — Yellow | 187 |
| — Pill with Iron | 223 |
| Mercuric Ammonium Chloride | 137 |
| — Chloride | 131 |
| — Iodide | 128 |
| — Oxide | 130 |
| Mercurous Chloride | 133 |
| Mercury | 135 |
| — Amidochloride of | 137 |
| — Ammoniated | 137 |

| | PAGE |
|---|---|
| Mercury Ball | 131 |
| — Bichloride of | 131 |
| — Biniodide of | 128 |
| — — Ointment of | 310 |
| — with Chalk | 138 |
| — Compound Ointment of | 310 |
| — — Spirit of Perchloride of | 279 |
| — Nitrate of, Ointment of | 311 |
| — Ointment of | 309 |
| — — of Biniodide of | 310 |
| — — Compound | 310 |
| — — of Nitrate of | 311 |
| — — of Perchloride of | 311 |
| — — of Red Iodide of | 310 |
| — Perchloride of | 131 |
| — — Compound Spirit of | 279 |
| — — Ointment of | 311 |
| — — Solution of | 174 |
| — Pill of, and Iron | 223 |
| — Protochloride of | 133 |
| — Red Iodide of | 128 |
| — — — Ointment of | 130 |
| — — Oxide of | 130 |
| — Spirit of Perchloride of, Compound | 279 |
| — Subchloride of | 133 |
| Methylated Spirit | 279 |
| Mindererus's Spirit | 160 |
| Mistura Ammoniæ | 197 |
| — Creosoti | 198 |
| — Cretæ Composita | 198 |
| — Ricini | 199 |
| Mixture, Ammonia | 197 |
| — Castor Oil | 214 |
| — Chalk, Compound | 198 |
| — Compound Chalk | 198 |
| — Gregory's | 256 |
| Moist Ferric Hydrate | 119 |
| — Hydrated Peroxide of Iron | 119 |
| — Peroxide of Iron | 119 |
| Molasses | 290 |
| Monkshood | 29, 30 |
| Morphinæ Acetas | 199 |
| — Hydrochloridum | 200 |
| Morphine | 201 |
| Morrhuæ, Oleum | 211 |
| Mucilage of Gum Acacia | 202 |
| — of Starch | 202 |
| Mucilago Acaciæ | 202 |
| — Amyli | 202 |
| Muriatic Acid | 13 |
| Mustard | 263 |
| — Poultice | 78 |
| Mylabridis, Unguentum | 312 |
| Mylabris | 203 |
| — Cichorii | 203 |
| — Ointment of | 312 |
| Myrrh | 203 |
| — Tincture of | 300 |
| Myrrha | 203 |
| Myrrhæ, Tinctura | 300 |
| Nitre | 243 |
| — Sweet Spirit | 271 |
| Nitric Acid | 17 |
| — — Dilute | 18 |
| Nitro hydrochloric Acid, Dilute | 19 |
| Nitrous Ether, Spirit of | 271 |
| Nucis Vomicæ, Tinctura | 301 |
| Nut-galls | 124 |
| Nux Vomica | 204 |
| — — Tincture of | 301 |
| Oak-bark | 252 |
| Oak-galls | 124 |
| Oil of Anise | 205 |
| — Cajuput | 206 |
| — Camphorated | 207 |
| — of Cantharides | 207 |
| — Carron | 151 |

PAGE

Oil, Castor . . . 214
— — Mixture . . 199
— of Cloves . . . . 207
— Cod-liver . . . . 212
— Croton . . . . 208
— — Liniment of . . . 153
— Eucalyptus . . . 210
— of Juniper . . . . 210
— Linseed . . . . 211
— Olive . . . . . 213
— Palm . . . . . 214
— Peppermint . . . 212
— of Turpentine . . . 215
— of Vitriol . . . . 23
— White . . . . 152
Ointment of Aconite . . 305
— Alum, Compound . . 305
— of Ammoniated Mercury . 309
— Antiseptic, Lister's . . 8
— Arsenic . . . . 306
— Biniodide of Mercury . 310
— Black Hellebore . . 308
— Blistering . . . 307, 310
— Blue . . . . . 309
— of Cantharides . . . 307
— of Carbolic Acid . . 304
— — — Compound . . 304
— of Carbonate of Zinc . . 316
— of Chlorinated Lime . . 306
— Citrine . . . . 311
— of Creosote . . . 307
— Digestive . . . 314, 316
— of Galls . . . . 308
— — and Opium . . . 308
— Healing . . . . 316
— of Iodide of Sulphur . . 315
— of Iodine . . . . 312
— — Compound . . . 312
— of Mercury . . . 309
— — Compound . . . 310
— of Mylabris . . . 312

PAGE

Ointment of Nitrate of Mercury . . . 311
— — of Silver . . . 306
— of Oxide of Zinc . . 317
— of Perchloride of Mercury 311
— of Red Iodide of Mercury . 310
— of Resin . . . . 314
— of Savin . . . . 314
— Sheep . . . . 315
— of Subacetate of Copper . 307
— of Sulphur . . . 314
— — Compound . . . 315
— of Sulphurated Potash . 318
— of Tannic Acid . . . 304
— of Tar . . . . 313
— of Tartarated Antimony . 305
— of Turpentine . . . 316
— of Verdigris . . . 307
— of White Hellebore . . 316
Oleum Anisi . . . . 205
— Cajuput . . . . 206
— Camphoratum . . . 207
— Cantharidis . . . 207
— Caryophylli . . . 207
— Crotonis . . . . 208
— Eucalypti . . . . 210
— Jecoris Aselli . . . 212
— Juniperi . . . . 210
— Lini . . . . . 210
— Menthæ Piperitæ . . 212
— Morrhuæ . . . . 212
— Olivæ . . . . 213
— Palmæ . . . . 214
— Ricini . . . . 214
— Terebinthinæ . . . 215
Olive Oil . . . . 213
Opii Compositus, Pulvis . 250
— Enema . . . . 108
— Linimentum . . . 156
— Liquor Plumbi Subacetatis et . . . . 178

| | PAGE |
|---|---|
| Opii, Tinctura | 302 |
| Opium | 217 |
| — Compound Powder of | 250 |
| — Enema of | 108 |
| — Liniment of | 156 |
| — Powder of, Compound | 250 |
| — Solution of Subacetate of Lead and | 178 |
| — Tincture of | 302 |
| Opodeldoc | 156 |
| Orthophosphoric Acid | 20 |
| Palm Butter | 214 |
| — Oil | 214 |
| Palmæ, Oleum | 214 |
| Papaveris, Syrupus | 287 |
| Paraffin | 219 |
| Paraffinum | 219 |
| Pepper, Black | 224 |
| Peru, Balsam of | 57 |
| Phenacetine | 220 |
| Phenol | 8 |
| Phenyl Acetamide | 1 |
| Phosphoric Acid | 20 |
| Physostigmatis, Extractum | 222 |
| — Semen | 221 |
| Physostigmina | 221 |
| Picis, Emplastrum | 107 |
| — Liquidæ, Unguentum | 313 |
| Pill, Blue, Ferruginate | 223 |
| — Ferruginate Blue | 223 |
| — Mercurial, with Iron | 223 |
| — of Mercury and Iron | 223 |
| Pilula Hydrargyri et Ferri | 223 |
| Pimenta | 223 |
| Pimentæ, Tinctura | 302 |
| Pimento | 223 |
| — Tincture of | 302 |
| Piper Nigrum | 224 |
| Pitch, Black | 227 |
| Pitch, Burgundy | 225 |
| — Plaster | 107 |
| Pix Burgundica | 225 |
| — Liquida | 226 |
| — Nigra | 227 |
| Plaster, Pitch | 107 |
| Plumbi Acetas | 227 |
| — Oxidum | 229 |
| — Subacetatis Dilutus, Liquor | 177 |
| — — Linimentum | 156 |
| — — Liquor | 176 |
| — — et Opii, Liquor | 178 |
| Podophylli Resina | 229 |
| — Rhizoma | 230 |
| Poppies, Syrup of | 287 |
| Potash, Caustic | 231 |
| — Hydroxide | 231 |
| — Lotion of Chloride of Ammonium and Nitrate of | 186 |
| — Nitrate of | 243 |
| — Permanganate of | 245 |
| — Solution of | 178 |
| — Sulphate of | 248 |
| — Sulphurated | 232 |
| — — Ointment of | 313 |
| — Yellow Prussiate of | 247 |
| Potassa Caustica | 231 |
| — Fusa | 231 |
| — Sulphurata | 232 |
| Potassæ Liquor | 178 |
| — Sulphuratæ, Liquor | 180 |
| — — Unguentum | 313 |
| Potassii Arsenitis, Liquor | 165 |
| — Bicarbonas | 233 |
| — Bromidum | 235 |
| — Carbonas | 237 |
| — Chloras | 238 |
| — Hydras | 231 |
| — Iodidi, Liquor Iodi et | 174 |
| — Iodidum | 240 |

PAGE

Potassii Nitras . . . 243
— Nitratis, Lotio Ammonii Chloridi et . . . 186
— Permanganas . . . 245
— Prussia Flava . . . 247
— Sulphas . . . . 248
Potassium Alum . . . 37
— Aluminium Sulphate . 37
— Antimony Tartrate . . 49
— Bicarbonate . . . 233
— Bromide . . . . 235
— Carbonate . . . . 237
— Chlorate . . . . 238
— Ferrocyanide . . . 247
— Hydrogen Carbonate . 233
— Hydroxide . . . 231
— Iodide . . . . 240
— Nitrate . . . . 243
— Permanganate . . . 245
— Solution of Iodine and Iodide of . . . 174
— Sulphate . . . . 248
— Sulphide of . . . 232
Poultice, Ammoniated Mustard . . . . 79
— Bran . . . . . 77
— Charcoal . . . . 76
— Chlorinated Lime . . 76
— Linseed . . . . 78
— Mustard . . . . 78
— Yeast . . . . 77
Powder, Antimonial . . 249
— of Carbonate of Zinc, Compound . . . . 251
— Dover's, Factitious . . 249
— Gregory's . . . . 250
— Grey . . . . . 138
— of Ipecacuanha, Compound 249
— James's, Factitious . . 249
— of Opium, Compound . 250
— of Rhubarb, Compound . 250

PAGE

Powder of Sulphate of Iron, Compound . . . 240
Prepared Chalk . . . 96
— Lard . . . . . 31
— Suet . . . . . 262
Proof Spirit . . . . 281
Protochloride of Mercury . 133
Prussic Acid . . . . 15
— — Scheele's . . . 15
Pulvis Antimonialis . . 249
— Ferri Sulphatis Compositus 249
— Ipecacuanhæ Compositus . 249
— Opii Compositus . . 250
— Rhei Compositus . . 250
— Zinci Carbonatis Compositus . . . . 251
Pyroligneous Acid . . . 5
Pyroxylin . . . . 251

Quercus Cortex . . . 252
Quicklime . . . . 65
Quicksilver . . . . 135
Quininæ Sulphas . . . 253
Quinine, Disulphate of . . 253
— Sulphate . . . . 253
— Sulphate of . . . 253

Rectified Spirit . . . 280
Red Iodide of Mercury . . 128
— Oxide of Mercury . . 130
— Precipitate . . . 137
— Rose Petals . . . 258
— Sanders Wood, Tincture of 303
Refined Sugar . . . 259
Resin . . . . . 256
— Compound Mass of . . 196
— Ointment of . . . 314
— Yellow . . . . 256
Resina . . . . . 256
Resinæ Compositæ, Massa . 196

| | PAGE |
|---|---|
| Resinæ, Unguentum | 314 |
| Rhamni Succus | 284 |
| — Syrupus | 284 |
| Rhei Compositus, Pulvis | 250 |
| — Radix | 257 |
| Rhubarb, Compound Powder of | 250 |
| — Powder of, Compound | 250 |
| — Root | 257 |
| Rosæ Gallicæ Petala | 258 |
| Rose Petals, Red | 258 |
| Roses, Confection of | 94 |
| Rye, Ergot of | 109 |
| Sabinæ Cacumina | 258 |
| — Unguentum | 314 |
| Saccharated Carbonate of Iron | 116 |
| Sacchari Fæx | 290 |
| Saccharum Purificatum | 259 |
| Saffron | 97 |
| Sal Ammoniac | 42 |
| Salt, Common | 272 |
| — Glauber's | 270 |
| — Spirit of | 13 |
| Saltpetre | 243 |
| Sanders Wood, Tincture of Red | 303 |
| Santalis Rubra, Tinctura | 303 |
| Santonin | 259 |
| Santoninum | 259 |
| Sapo Durus | 260 |
| — Mollis | 261 |
| Saponis Compositum, Linimentum | 156 |
| Savin, Ointment of | 314 |
| — Tops | 258 |
| Scheele's Prussic Acid | 151 |
| Scillæ | 262 |
| Sevum Præparatum | 262 |
| Sheep and Calves' Cordial | 198 |
| Sheep Dip or Wash (Arsenical) | 166 |
| — Ointment | 315 |
| Silver, Nitrate of | 53 |
| — — Ointment of | 306 |
| — — Solution of | 165 |
| — Ointment of Nitrate of | 306 |
| — Solution of Nitrate of | 165 |
| Sinapis | 263 |
| Slaked Lime | 63 |
| Soap, Compound Liniment of | 156 |
| — Hard | 260 |
| — Soft | 261 |
| Soda Biborate | 61 |
| — Bicarbonate of | 264 |
| — Carbonate of | 266 |
| — — Dried | 268 |
| — — Solution of Aloes and | 158 |
| — Caustic | 263 |
| — Caustica | 263 |
| — Dried Carbonate of | 268 |
| — Hydrate of | 263 |
| — Hyposulphite of | 268 |
| — Solution of | 181 |
| — Sulphate of | 270 |
| — Sulphite of | 271 |
| — Washing | 266 |
| Sodii Bicarbonas | 264 |
| — Carbonas | 266 |
| — — Exsiccata | 268 |
| — Carbonatis, Liquor Aloes et | 158 |
| — Chloridum | 272 |
| — Hyposulphis | 268 |
| — Liquor | 181 |
| — Sulphas | 270 |
| — Sulphis | 271 |
| — Sulphocarbolas | 334 |
| Sodium, Biborate of | 61 |
| — Carbonate | 266 |
| — Chloride | 272 |
| — Chloride of | 272 |
| — Diborate | 61 |

PAGE

Sodium Hydrate . . . 263
— Hydrate of . . . 263
— Hydrogen Carbonate . . 264
— Hyposulphite . . . 268
— Salicylas . . . . 269
— Sulphate . . . . 270
— Sulphite . . . . 271
— Sulphocarbolate . . 334
Soft Soap . . . . 261
Solution of Acetate of Ammonia . . . . 160
— of Aloes . . . . 158
— — and Carbonate of Soda 158
— of Alum . . . . 159
— of Ammonia . . . 159
— — Strong . . . . 162
— of Antimonious Chloride . 163
— of Arsenic, Hydrochloric . 167
— Arsenical . . . . 165
— of Cantharides, Terebinthinated . . . 170
— of Carbolic Acid . . 157
— of Chloride of Antimony . 163
— — of Lime . . . 169
— — of Zinc . . . . 183
— of Chlorinated Lime . . 169
— Fowler's . . . . 165
— of Hypochlorite of Lime . 169
— of Iodine and Iodide of Potassium . . . 174
— of Lime . . . . 168
— Lugol's . . . . 174
— of Nitrate of Potash . . 180
— — of Silver . . . 165
— of Perchloride of Iron, Strong . . . 172
— — of Mercury . . . 174
— of Permanganate of Potash 180
— of Persulphate of Iron . 173
— of Potash . . . . 178
— of Soda . . . . 181

PAGE

Solution of Strychnine Hydrochloride . . . 182
— of Subacetate of Lead . 176
— — — Dilute . . . 177
— — — and Opium . . 178
— of Sulphate of Copper . 171
— — — Compound . . 171
— — Zinc . . . . 185
— of Sulphurated Potash . 180
— of Tartarated Antimony . 164
— of Terchloride of Antimony . . . . 163
Spanish Fly . . . . 70
Spirit of Ammonia, Aromatic 277
— — Fetid . . . . 277
— of Camphor . . . 278
— of Chloroform . . . 278
— of Ether . . . . 274
— Methylated . . . 279
— Mindererus's . . . 160
— of Nitre, Sweet . . . 275
— of Nitrous Ether . . 275
— of Perchloride of Mercury, Compound . . . 279
— Proof . . . . . 281
— Rectified . . . . 280
— of Salt . . . . 131
— of Turpentine . . . 215
— of Wine . . . . 280
Spiritus Ætheris . . . 274
— — Nitrici . . . . 275
— — Nitrosi . . . . 275
— — Sulphurici . . . 274
— Ammoniæ Aromaticus . 277
— — Fœtidus . . . 277
— Camphoræ . . . 278
— Chloroformi . . . 278
— Hydrargyri Perchloridi Compositus . . . 279
— Methylatus . . . 279
— Rectificatus . . . 280

| | PAGE |
|---|---|
| Spiritus Tenuior | 281 |
| Staphisagriæ Semina | 282 |
| Starch | 44 |
| — Mucilage of | 202 |
| Stavesacre Seeds | 282 |
| Strong Solution of Ammonia | 162 |
| Strychnine | 282 |
| — Hydrochloride, Solution of | 182 |
| Strychninæ Hydrochloridi, Liquor | 182 |
| Sublimed Sulphur | 284 |
| Succus Rhammi | 284 |
| Suet, Prepared | 262 |
| Sugar of Lead | 227 |
| Sulphur, Compound Ointment of | 315 |
| — Flowers of | 284 |
| — Iodide of | 286 |
| — — Ointment of | 315 |
| — Liver of | 232 |
| — Ointment of, Compound | 315 |
| — — of Iodide of | 315 |
| — Sublimatum | 284 |
| — Sublimed | 284 |
| Sulphurated Potash | 232 |
| — — Ointment of | 313 |
| Sulphuric Acid | 23 |
| — — Dilute | 24 |
| — Ether | 32 |
| Sulphuris Iodi | 286 |
| — Iodidi, Unguentum | 315 |
| — Unguentum Compositum | 315 |
| Sulphurous Acid | 26 |
| — Anhydride | 26 |
| Sweet Spirit of Nitre | 275 |
| Syrup of Buckthorn | 288 |
| — of Chloral Hydrate | 287 |
| — of Poppies | 287 |
| Syrupus Chloralis Hydratis | 287 |
| — Papaveris | 287 |
| — Rhamni | 288 |

| | PAGE |
|---|---|
| Tabaci, Decoctum | 10 |
| — Enema | 108 |
| — Folia | 288 |
| — Infusum | 142 |
| Tannic Acid | 27 |
| — — Ointment of | 304 |
| Tannin | 27 |
| Tar | 226 |
| — Ointment of | 313 |
| — Wood | 226 |
| Taraxacum Root | 289 |
| Tartar Emetic | 49 |
| Tartarated Antimony | 49 |
| — — Compound Mass of | 193 |
| — — Solution of | 164 |
| Tartarised Antimony | 49 |
| Terebinthina Canadensis | 289 |
| Terebinthinæ Compositum, Linimentum | 157 |
| — Enema | 109 |
| — Linimentum | 157 |
| — Oleum | 215 |
| — Unguentum | 316 |
| Terebinthinated Solution of Cantharides | 170 |
| Terra Japonica | 79 |
| Theriaca | 290 |
| Thymol | 290 |
| Tinctura Aconiti | 291 |
| — Aloes Composita | 292 |
| — Arnicæ | 293 |
| — Belladonnæ | 293 |
| — Camphoræ | 294 |
| — Cannabis Indicæ | 294 |
| — Cantharidis | 294 |
| — — Fortior | 295 |
| — Cardamomi Composita | 295 |
| — Chloroformi Composita | 296 |
| — Cinchonæ | 296 |
| — Colchici Seminum | 297 |
| — Conii | 297 |

| | PAGE |
|---|---|
| Tinctura Conii Fructus | 290 |
| — Crotonis | 289 |
| — Digitalis | 298 |
| — Ergotæ | 299 |
| — Ferri Perchloridi | 299 |
| — — Sesquichloridi | 299 |
| — Iodi | 300 |
| — Myrrhæ | 300 |
| — Nucis Vomicæ | 301 |
| — Opii | 302 |
| — Pimentæ | 302 |
| — Santalis Rubra | 303 |
| — Zingiberis | 303 |
| Tincture of Aconite | 291 |
| — of Aloes, Compound | 292 |
| — of Arnica | 293 |
| — of Cantharides | 294 |
| — — Strong | 295 |
| — of Cardamoms, Compound | 295 |
| — of Chloroform, Compound | 296 |
| — of Colchicum Seeds | 297 |
| — of Croton | 298 |
| — of Ergot | 299 |
| — of Ginger | 303 |
| — of Hemlock | 297 |
| — of Iodine | 300 |
| — of Myrrh | 300 |
| — of Nux Vomica | 301 |
| — of Opium | 302 |
| — of Perchloride of Iron | 299 |
| — of Pimento | 302 |
| — of Red Sanders Wood | 303 |
| Tobacco, Decoction of | 105 |
| — Enema of | 108 |
| — Infusion of | 142 |
| — Leaf | 288 |
| Tonic Mass | 195 |
| Treacle | 290 |
| Trichloraldehyd | 82 |
| Turpentine, Compound Liniment of | 157 |
| Turpentine, Enema of | 109 |
| — Liniment of | 157 |
| — — Compound | 157 |
| — Oil of | 215 |
| — Ointment of | 316 |
| — Spirit of | 215 |
| Unguentum Acidi Carbolici | 304 |
| — — — Compositum | 304 |
| — — Tannici | 304 |
| — Aconiti | 305 |
| — Æruginis | 307 |
| — Aluminis Compositum | 305 |
| — Antimonii Tartarati | 305 |
| — Argenti Nitratis | 306 |
| — Arsenici | 306 |
| — Calcis Chlorinatæ | 306 |
| — Cantharidis | 307 |
| — Creosoti | 307 |
| — Cupri Subacetatis | 307 |
| — Gallæ | 308 |
| — Opii | 308 |
| — Hellebori Nigri | 308 |
| — Hydrargyri | 309 |
| — — Ammoniati | 309 |
| — — Biniodidi | 310 |
| — — Compositum | 310 |
| — — Iodidi Rubri | 310 |
| — — Nitratis | 311 |
| — — Perchloridi | 311 |
| — Iodi | 312 |
| — — Compositum | 312 |
| — Mylabridis | 312 |
| — Picis Liquidæ | 313 |
| — Potassæ Sulphuratæ | 313 |
| — Resinæ | 314 |
| — Sabinæ | 314 |
| — Sulphuris | 314 |
| — — Compositum | 315 |
| — — Iodidi | 315 |

| | PAGE |
|---|---|
| Unguentum Terebinthinæ | 316 |
| — Veratri | 316 |
| — Zinci Carbonatis | 316 |
| — — Oxidi | 317 |
| Vapor Chlori | 317 |
| Vaseline | 319 |
| Veratri, Unguentum | 316 |
| Veratrum | 319 |
| Verdigris | 100 |
| — Liniment of | 153 |
| — Ointment of | 307 |
| Vinegar | 2 |
| — of Cantharides | 3 |
| Vinum Ferri | 321 |
| Vitriol, Blue | 101 |
| — Green | 121 |
| — Oil of | 23 |
| — White | 327 |
| Wash, Sheep | 166 |
| Washing Soda | 266 |
| Water, Distilled | 51 |
| Wax, Bees', Unbleached | 80 |
| — Yellow | 80 |
| White Hellebore | 319 |
| — — Ointment of | 316 |
| — Marble | 192 |
| — Oil | 152 |
| — Precipitate | 137 |
| — Vitriol | 327 |
| Whiting | 96 |
| Wine of Iron | 321 |
| — Spirit of | 280 |
| Wood Charcoal | 72 |
| — Tar | 226 |
| Yeast | 81 |
| — Poultice | 77 |
| Yellow Mercurial Lotion | 187 |
| — Resin | 256 |
| — Wash | 187 |
| Zinc Acetate | 321 |
| — Acetate of | 321 |
| — Butter of | 324 |
| — Carbonate | 323 |
| — Carbonate of | 323 |
| — — Compound Powder of | 251 |
| — — Ointment of | 216 |
| — Chloride | 324 |
| — Chloride of | 324 |
| — — Solution of | 183 |
| — Compound Powder of Carbonate of | 251 |
| — Ointment of Carbonate of | 316 |
| — Oxide | 326 |
| — Oxide of | 326 |
| — Solution of Chloride of | 183 |
| — Sulphate | 327 |
| — Sulphate of | 327 |
| Zinci Acetas | 321 |
| — Carbonas | 323 |
| — Carbonatis Compositus, Pulvis | 251 |
| — — Unguentum | 316 |
| — Chloridi, Liquor | 183 |
| — Chloridum | 324 |
| — Oxidum | 326 |
| — Sulphas | 327 |
| Zingiber | 330 |
| Zingiberis Composita, Massa | 197 |
| — Tinctura | 303 |


*Spottiswoode & Co. Ltd., Printers, New-street Square, London.*

No. 2.

*London, 7, Great Marlborough Street,*
*April*, 1904.

# A SELECTION
FROM
# J. & A. CHURCHILL'S CATALOGUE,
COMPRISING
MOST OF THE RECENT WORKS PUBLISHED BY THEM.

*N.B.—J. & A. Churchill's larger Catalogue, which contains over* 600 *works, with a Complete Index to their Subjects, will be sent on application.*

**Human Anatomy: a Treatise by various** Authors. Edited by HENRY MORRIS, M.A., M.B. Lond., F.R.C.S., Surgeon to, and Lecturer on Surgery at, the Middlesex Hospital. Third Edition. Roy. 8vo, with 846 Illustrations, of which 267 are in several colours, 30s. net.

**Heath's Practical Anatomy: a Manual of** Dissections. Ninth Edition. Edited by J. ERNEST LANE, F.R.C.S., Surgeon and Lecturer on Anatomy at St. Mary's Hospital. Crown 8vo, with 321 Engravings, of which 32 are coloured, 12s. 6d.

**Wilson's Anatomist's Vade-Mecum.** Eleventh Edition, by HENRY E. CLARK, M.R.C.S. Eng., F.F.P.S. Glasg., Examiner in Anatomy F.P.S. Crown 8vo, with 492 Engravings and 26 Coloured Plates, 18s.

**An Atlas of Human Anatomy. By Rickman** J. GODLEE, M.S., F.R.C.S., Surgeon and late Demonstrator of Anatomy, University College Hospital. With 48 Imp. 4to Plates (112 figures), and a volume of Explanatory Text. £4 14s. 6d. net.

**Human Osteology. By Luther Holden.** Eighth Edition, edited by CHARLES STEWART, F.R.S., Conservator of the Museum R.C.S., and ROBERT W. REID, M.D., F.R.C.S., Regius Professor of Anatomy in the University of Aberdeen. 8vo, with 59 Lithographic Plates and 74 Engravings, 16s.

*By the same Author.*

**Landmarks, Medical and Surgical.** Fourth Edition. 8vo, 3s. 6d.

**A Manual of Practical Anatomy.** By the late Professor ALFRED W. HUGHES, M.B., M.C. Edin., Professor of Anatomy, King's College, London. Edited and completed by ARTHUR KEITH, M.D., Lecturer on Anatomy, London Hospital Medical College. In three parts, royal 8vo. Part I., with 38 Coloured Plates, and 116 Figures in the Text, 10s. 6d. Part II., with 4 Coloured Plates and 151 Figures in the Text, 8s. 6d. Part III., with 12 Coloured Plates and 204 Figures in the Text, 10s. 6d.

**The Essentials of Regional Anatomy.** By RICHARD J. BERRY, M.D., F.R.S., F.R.C.S., (Edin.), Lecturer on Anatomy in the New School of Medicine of the Royal Colleges, Edinburgh. Second Edition. 8vo, interleaved, 3 vols., 5s. each, net.

**Anatomy of the Joints of Man.** By Henry MORRIS, Senior Surgeon to the Middlesex Hospital. With 44 Lithographic Plates (several coloured). 8vo, 16s.

**A Manual of General or Experimental Pathology,** for Students and Practitioners. By W. S. LAZARUS-BARLOW, M.D., F.R.C.P., Director of the Cancer Research Laboratories, Middlesex Hospital. Roy. 8vo, 21s. net.

*By the same Author.*

**The Elements of Pathological Anatomy and Histology** for Students. With 7 Coloured Plates and 171 Figures in the Text. Royal 8vo, 24s. net.

**Pathological Anatomy of Diseases.** Arranged according to the nomenclature of the R.C.P. Lond. By NORMAN MOORE, M.D., F.R.C.P., Assistant Physician to St. Bartholomew's Hospital. Fcap. 8vo, with 111 Engravings, 8s. 6d.

**A Manual of Clinical and Practical Pathology.** By W. E. WYNTER, M.D., M.R.C.P., and F. J. WETHERED, M.D., F.R.C.P. With 4 Coloured Plates and 67 Engravings. 8vo, 12s. 6d.

**The Pathologist's Handbook: a Manual for** the Post-mortem Room. By T. N. KELYNACK, M.D., late Demonstrator in Morbid Anatomy, Owens College, Manchester. With 126 Illustrations, fcap. 8vo, pegamoid, 4s. 6d.

**A Manual of Antenatal Pathology and Hygiene.** The Fœtus. By J. W. BALLANTYNE, M.D., F.R.C.P.E., F.R.S. Edin., Lecturer on Antenatal Pathology and Teratology in the University of Edinburgh. With 69 Illustrations (14 coloured), roy. 8vo, 20s. net.

**The Human Brain: Histological and Coarse** Methods of Research. By W. BEVAN LEWIS, L.R.C.P. Lond. 8vo, with Wood Engravings and Photographs, 8s.

**The Physiology and the Pathology of the** Cerebral Circulation: an Experimental Research. By LEONARD HILL, M.D., Hunterian Professor, R.C.S. With 41 Illustrations. Royal 8vo, 12s.

**Elements of Human Physiology. By Ernest** H. STARLING, M.D., F.R.C.P., F.R S., Jodrell Professor of Physiology in University Colleg, London. Fifth Edition, with 321 Illustrations, 8vo, 12s. 6d.

**Essentials of Human Physiology for Medical** Students. By D. NOËL PATON, M.D., B.Sc., Lecturer on Physiology, School of Medicine of the Royal Colleges, Edinburgh. With 110 Illustrations. 8vo., 12s. net.

**Elementary Practical Physiology, including** Histology, Chemical and Experimental Physiology. By DE BURGH BIRCH, M.D., F.R.S.E., Professor of Physiology in the Yorkshire College of the Victoria University. With 62 Engravings. Crown 8vo, 6s. 6d.

**Practical Lessons in Elementary Biology, for** Junior Students. By PEYTON T. B. BEALE, F.R.C.S., Lecturer on Elementary Biology and Demonstrator in Physiology in King's College, London. Crown 8vo, 3s. 6d.

**Medical Jurisprudence: its Principles and** Practice. By ALFRED S. TAYLOR, M.D., F.R.C.P., F.R.S. Fourth Edition, by THOMAS STEVENSON, M.D., F.R.C.P., Lecturer on Medical Jurisprudence at Guy's Hospital. 2 vols. 8vo, with 189 Engravings, 31s. 6d.

**Lectures on Medical Jurisprudence and Toxi-** cology. By FRED J. SMITH, M.D., F.R.C.P., Lecturer on Forensic Medicine and Toxicology at the London Hospital. Crown 8vo, 7s. 6d.

**The Theory and Practice of Hygiene. (Notter** and Firth). By J. LANE NOTTER, M.D., late Professor of Hygiene in the Army Medical School; and W. H. HORROCKS, M.B., B.Sc., Assistant Professor of Military Hygiene in the Staff Medical College. Second Edition. With 15 Plates and 134 other Illustrations. Royal 8vo, 25s.

**Manual of Hygiene. By W. H. Hamer, M.D.,** D.P.H., F.R.C.P., Lecturer on Public Health, St. Bartholomew's Hospital, Assistant Medical Officer of Health of the Administrative County of London. With 93 Illustrations. 8vo, 12s. 6d. net.

**A Handbook of Hygiene and Sanitary Science.** By GEO. WILSON, M.A., M.D., LL.D., D.P.H. Camb. Medical Officer of Health for Mid-Warwickshire. Eighth Edition. Post 8vo, with Engravings, 12s. 6d.

**Public Health and Preventive Medicine. By** C. J. LEWIS, M.D., D.Sc. Edin., and ANDREW BALFOUR, M.D., B Sc. Edin., D.P.H. Camb. With 11 Plates and 189 Figures in the Text. Imp. 8vo, 25s. net.

**Hygiene and Public Health: a Treatise by** various Authors. Edited by THOMAS STEVENSON, M.D., and SHIRLEY F. MURPHY. In 3 vols., royal 8vo, fully Illustrated. Vol. I., 28s.; Vol. II., 32s.; Vol. III., 20s.

**A Simple Method of Water Analysis, especially** designed for the use of Medical Officers of Health. By JOHN C. THRESH, M.D.Vic., D.Sc. Lond., D.P.H. Camb. Fourth Edition, enlarged. Fcap. 8vo, 2s. 6d.

**Elements of Health: an Introduction to the** Study of Hygiene. By LOUIS C. PARKES, M.D., D.P.H. Lond., Medical Officer of Health for Chelsea, Lecturer on Public Health at St. George's Hospital. Post 8vo, with 27 Engravings, 3s. 6d.

**Diet and Food considered in relation to** Strength and Power of Endurance, Training and Athletics. By ALEXANDER HAIG, M.D., F.R.C.P. Fourth Edition. Crown 8vo, 2s.

**The Prevention of Epidemics and the Con-** struction and Management of Isolation Hospitals. By ROGER MCNEILL, M.D. Edin., D.P.H. Camb., Medical Officer of Health for the County of Argyll. 8vo, with several Hospital Plans, 10s. 6d.

**Second Treatise on the Effects of Borax** and Boric Acid on the Human System. By Dr. OSCAR LIEBREICH, Professor in the University of Berlin. With Plates, post 4to, 2s 6d.

**A Manual of Bacteriology, Clinical and Ap-** plied. With an Appendix on Bacterial Remedies, &c. By RICHARD T. HEWLETT, M.D., Professor of General Pathology and Bacteriology in King's College, London. Second Edition. With 20 Plates, and 66 Figures in the Text, post 8vo, 12s. 6d.

*By the same Author.*

**Serum Therapy, Bacterial Therapeutics, and** Vaccines. With 20 Figures, crown 8vo, 5s. net.

**An Introduction to the Bacteriological Exami-** nation of Water. By W. H. HORROCKS, M.B., B.Sc., Lond., Assistant Professor of Military Hygiene in the Staff Medical College, Major, Royal Army Medical Corps. With 5 Lithographic Plates, 8vo, 10s. 6d.

**Lessons in Disinfection and Sterilisation: an** Elementary Course of Bacteriology. With Experiments. By F. W. ANDREWES, M.D., F.R.C.P., Lecturer on Pathology, St. Bartholomew's Hospital. With 31 Illustrations, crown 8vo, 3s. net.

**Mental Diseases: Clinical Lectures. By T. S.** CLOUSTON, M.D., F.R.C.P. Edin., Lecturer on Mental Diseases in the University of Edinburgh. Fifth Edition. Cr. 8vo, with 19 Plates, 14s.

**Hospitals and Asylums of the World: their** Origin, History, Construction, Administration, Management, and Legislation. By Sir H. C. BURDETT, K.C.B. 4 vols., royal 8vo, and Portfolio. Complete, £12 12s. net. Vols. I. and II.—Asylums, £6 15s. net. Vols. III. and IV.—Hospitals, with Plans and Portfolio, £9 net.

**The Insane and the Law: a Plain Guide for** Medical Men, Solicitors, and Others. By G. PITT-LEWIS, K.C., R. PERCY SMITH, M.D., F.R.C.P., and J. A. HAWKE, B.A., Barrister-at-Law. 8vo, 14s.

**A Dictionary of Psychological Medicine, giving** the Definition, Etymology, and Synonyms of the Terms used in Medical Psychology; with the Symptoms, Treatment, and Pathology of Insanity; and THE LAW OF LUNACY IN GREAT BRITAIN AND IRELAND. Edited by D. HACK TUKE, M.D., LL.D., assisted by nearly 130 Contributors, British, Continental and American. 2 vols., 1,500 pages, royal 8vo, Illustrated, 42s.

**The Force of Mind: or, the Mental Factor** in Medicine. By ALFRED T. SCHOFIELD, M.D., Hon. Physician to Friedenheim Hospital. Second Edition. Crown 8vo, 5s. net.

**The Mental Affections of Children: Idiocy,** Imbecility, and Insanity. By WM. W. IRELAND, M.D. Edin., formerly Medical Superintendent of the Scottish Institution for the Education of Imbecile Children. Second Edition. With 21 Plates, 8vo, 14s.

**The Journal of Mental Science. Published** Quarterly, by Authority of the Medico-Psychological Association. 8vo, 5s.

**Manual of Midwifery, including all that is** likely to be required by Students and Practitioners. By ALFRED L. GALABIN, M.A., M.D., F.R.C.P., Obstetric Physician and Lecturer on Midwifery and Diseases of Women to Guy's Hospital. Fifth Edition. Post 8vo, with 298 Engravings, 15s.

**The Practice of Midwifery: a Guide for Prac-** titioners and Students. By D. LLOYD ROBERTS, M.D., F.R.C.P., Lecturer on Clinical Midwifery and Diseases of Women at the Owens College; Consulting Obstetric Physician to the Manchester Royal Infirmary. Fourth Edition. Fcap. 8vo, with Coloured Plates and Wood (226) Engravings, 10s. 6d.

**Obstetric Aphorisms: for the Use of Students** commencing Midwifery Practice. By JOSEPH G. SWAYNE, M.D., Lecturer on Midwifery in the Bristol Medical School. Tenth Edition. Fcap. 8vo, with 20 Engravings 3s. 6d.

**A Short Practice of Midwifery, embodying the** Treatment adopted in the Rotunda Hospital, Dublin. By HENRY JELLETT, M.D., B.A.O. Dub., Ex-Assistant Master, Rotunda Hospital. Fourth Edition. With 152 Illustrations. Crown 8vo, 8s. 6d.

*By the same Author.*

**A Short Practice of Midwifery for Nurses.** With 67 Illustrations, crown 8vo, 6s.

**Lectures on Obstetric Operations: including** the Treatment of Hæmorrhage, and forming a Guide to the Management of Difficult Labour. By ROBERT BARNES, M.D., F.R.C.P., Consulting Obstetric Physician to St. George's Hospital. Fourth Edition. 8vo, with 121 Engravings, 12s. 6d

*By the same Author.*

**A Clinical History of Medical and Surgical** Diseases of Women. Second Edition. 8vo, with 181 Engravings, 28s.

**Outlines of Gynæcological Pathology and** Morbid Anatomy. By C. HUBERT ROBERTS, M.D. Lond., Physician to the Samaritan Free Hospital for Women. With 151 Illustrations, 8vo, 21s.

**Diseases of Women. By Alfred L. Galabin,** M.A., M.D., F.R.C.P., Consulting Obstetric Physician to Guy's Hospital. Sixth Edition. 8vo, with 284 Engravings, 16s. net.

**A Short Practice of Gynæcology. By Henry** JELLETT, M.D., B.A.O. Dub., Ex-Assistant Master, Rotunda Hospital, Dublin. Second Edition. With 225 Illustrations, 8vo, 10s. 6d.

**Manual of the Diseases peculiar to Women.** By JAMES OLIVER, M.D., F.R.S.E., M.R.C.P., Physician to the Hospital for Diseases of Women, London. Fcap. 8vo, 3s. 6d.

*By the same Author.*

**Abdominal Tumours and Abdominal Dropsy** in Women. Crown 8vo, 7s. 6d.

**Sterility.** By ROBERT BELL, M.D., F.F.P. & S. Glasg., Senior Physician to the Glasgow Hospital for Diseases peculiar to Women. 8vo, 5s.

**Nursing, General, Medical, and Surgical, with** Appendix on Sickroom Cookery. By WILFRED J. HADLEY, M.D., F.R.C.P., Physician to the London Hospital. Crown 8vo, 3s. 6d.

**A Short Manual for Monthly Nurses.** By CHARLES J. CULLINGWORTH, M.D., F.R.C.P., Obstetric Physician to St. Thomas's Hospital. Revised by the Author, with the assistance of M. A. ATKINSON, Matron of the General Lying-in Hospital, Lambeth. Fifth Edition. Fcap. 8vo, 1s. 6d.

**Lectures on Medicine to Nurses.** By Herbert E. CUFF, M.D., F.R.C.S., Medical Superintendent, North Eastern Fever Hospital, London. Fourth Edition. With 29 Illustrations. Crown 8vo, 3s. 6d.

**Antiseptic Principles for Nurses.** By C. E. RICHMOND, F.R.C.S. Fcap. 8vo, 1s.

**A Practical Treatise on Disease in Children.** By EUSTACE SMITH, M.D., F.R.C.P., Physician to the King of the Belgians, and to the East London Hospital for Children, etc. Second Edition. 8vo, 22s.

*By the same Author.*

**Clinical Studies of Disease in Children.** Second Edition. Post 8vo, 7s. 6d.

*Also.*

**The Wasting Diseases of Infants and Children.** Sixth (cheap) Edition. Post 8vo, 6s.

**The Diseases of Children.** By James F. GOODHART, M.D., F.R.C.P. Seventh Edition, with the assistance of G. F. STILL, M.D., F.R.C.P., Assistant Physician to the Hospital for Sick Children, Great Ormond Street. 8vo, 12s. 6d. net.

**On the Natural and Artificial Methods of Feeding** Infants and Young Children. By EDMUND CAUTLEY, M.D., Physician to the Belgrave Hospital for Children. Second Edition. Crown 8vo, 7s. 6d.

**Materia Medica, Pharmacy, Pharmacology,** and Therapeutics. By W. HALE WHITE, M.D., F.R.C.P., Physician to, and Lecturer on Medicine at, Guy's Hospital. Eighth Edition, based upon the B.P. of 1898 and the Indian and Colonial Addendum Fcap. 8vo, 7s. 6d.

**Southall's Organic Materia Medica.** Sixth Edition, adapted to the B.P. of 1898. Edited by JOHN BARCLAY, B.Sc. Lond. Crown 8vo, 7s. 6d.

**An Introduction to the Study of Materia** Medica, designed for Students of Pharmacy and Medicine. By HENRY G. GREENISH, F.I.C., F.L.S., Professor of Pharmaceutics to the Pharmaceutical Society. With 213 Illustrations, 8vo, 15s.

*By the same Author.*

**The Microscopical Examination of Foods and** Drugs, in the Entire, Crushed, and Powdered States. With 168 Illustrations, 8vo, 10s. 6d. net.

*Also, with Mr. Eugene Collin,*

**An Anatomical Atlas of Vegetable Powders,** Designed as an Aid to the Microscopic Analysis of Powdered Foods and Drugs. With 138 Illustrations. 4to, 12s. 6d. net.

**Practical Pharmacy: an Account of the** Methods of Manufacturing and Dispensing Pharmaceutical Preparations; with a chapter on the Analysis of Urine. By E. W. LUCAS, F.C.S., Examiner at the Pharmaceutical Society. With 283 Illustrations. Roy. 8vo, 12s. 6d.

**Galenic Pharmacy: a Practical Handbook to** the Processes of the British Pharmacopœia. By R. A CRIPPS, M.P.S. 8vo, with 76 Engravings, 8s. 6d.

**Practical Pharmacy. By Barnard S. Proctor.** Third Edition. 8vo, with Engravings and Fac-simile Prescriptions, 14s.

**The Galenical Preparations of the British** Pharmacopœia: a Handbook for Medical and Pharmaceutical Students. By C. O. HAWTHORNE, M.D., C.M., late Lecturer on Materia Medica and Therapeutics, Queen Margaret's College, Glasgow. 8vo, 4s. 6d.

**The Pharmaceutical Formulary: a Synopsis** of the British and Foreign Pharmacopœias. By HENRY BEASLEY. Twelfth Edition by J. OLDHAM BRAITHWAITE 18mo, 6s. 6d.

*By the same Author.*

**The Druggist's General Receipt-Book.** Tenth Edition. 18mo, 6s. 6d.

*Also.*

**The Book of Prescriptions: containing upwards** of 3,000 Prescriptions collected from the Practice of the most eminent Physicians and Surgeons, English and Foreign. Seventh Edition, 18mo, 6s. 6d.

**A Companion to the British Pharmacopœia.** By PETER SQUIRE. Revised by PETER WYATT SQUIRE. F.L.S., F.C.S. Seventeenth Edition. 8vo, 12s. 6d.

*By the same Authors.*

**The Pharmacopœias of thirty of the London** Hospitals, arranged in Groups for Comparison. Seventh Edition. Fcap. 8vo, 6s.

**Pereira's Selecta è Prescriptis: containing** Lists of Terms used in Prescriptions, with Explanatory Notes, etc. Also, a Series of Abbreviated Prescriptions with Translations. Eighteenth Edition, by JOSEPH INCE. 24mo, 5s.

**Year-Book of Pharmacy: containing the Trans-** actions of the British Pharmaceutical Conference. Annually. 8vo, 10s.

**Manual of Botany, in two Vols., crown 8vo.** By J. REYNOLDS GREEN, Sc.D., M.A., F.R.S., Professor of Botany to the Pharmaceutical Society.

Vol. I.: Morphology and Anatomy. Third Edition, with 778 Engravings. 7s. 6d.

Vol. II.: Classification and Physiology. Second Edition, with 466 Engravings, 10s.

*By the same Author.*

**An Introduction to Vegetable Physiology.** With 184 Illustrations, 8vo, 10s. 6d.

**The Student's Guide to Systematic Botany,** including the Classification of Plants and Descriptive Botany. By ROBERT BENTLEY, late Emeritus Professor of Botany in King's College and to the Pharmaceutical Society. Fcap. 8vo, with 350 Engravings, 3s. 6d.

**Medicinal Plants: being Descriptions with** original figures, of the Principal Plants employed in Medicine, and an account of their Properties and Uses. By Prof. BENTLEY and Dr. H. TRIMEN, F.R.S. In 4 vols., large 8vo, with 306 Plates (not coloured), cloth binding, £5 5s. net.

**Therapeutic Electricity and Practical Muscle** Testing. By W. S. HEDLEY, M.D., in charge of the Electro-therapeutic Department of the London Hospital. With 110 Illustrations. Roy. 8vo, 8s. 6d.

**Practical Therapeutics: a Manual. By** EDWARD J. WARING, C.I.E., M.D., F.R.C.P., and DUDLEY W. BUXTON, M.D., B.S. Lond. Fourth Edition. Crown 8vo, 14s.

*By the same Author.*

**Bazaar Medicines of India, and Common** Medical Plants. With Full Index of Diseases, indicating their Treatment by these and other Agents procurable throughout India, etc. Fifth Edition. Fcap. 8vo, 5s.

**Climate and Fevers of India, with a Series** of Cases (Croonian Lectures, 1882). By Sir JOSEPH FAYRER, Bart., K.C.S.I., M.D. 8vo, with 17 Temperature Charts, 12s.

**The Malarial Fevers of British Malaya. By** HAMILTON WRIGHT, M.D. (McGILL), Director of the Institute for Medical Research, Federated Malay States. With Map and Charts. Royal 8vo, 3s. net.

*By the same Author.*

**The Etiology and Pathology of Beri-beri.** With Map and Charts. Roy. 8vo, 3s. net.

**Psilosis or "Sprue": its Nature and Treatment**; with Observations on various Forms of Diarrhœa acquired in the Tropics. By GEORGE THIN, M.D. Second and Enlarged Edition, with Illustrations. 8vo, 10s.

**A Manual of Family Medicine and Hygiene** for India. Published under the Authority of the Government of India. By Sir WILLIAM J. MOORE, K.C.I.E., M.D. Seventh Edition revised by Major J. H. TULL WALSH, I.M.S. Post 8vo, with 70 Engravings, 12s.

*By the same Author.*

**A Manual of the Diseases of India: with a** Compendium of Diseases generally. Second Edition. Post 8vo, 10s.

**The Prevention of Disease in Tropical and** Sub-Tropical Campaigns. (Parkes Memorial Prize for 1886.) By Lieut.-Col. ANDREW DUNCAN, M.D., B.S. Lond., F.R.C.S., H.M. Indian Medical Service. 8vo, 12s. 6d.

**A Text-Book of Medicine. Begun by the late** C. HILTON FAGGE, M.D.; completed and re-written by P. H. PYE-SMITH, M.D., F.R.S. Fourth Edition. 2 vols. royal 8vo, 42s.

**Manual of the Practice of Medicine. By** FREDERICK TAYLOR, M.D., F.R.C.P., Physician to, and Lecturer on Medicine at, Guy's Hospital. Sixth Edition. Post 8vo. with Engravings, 16s.

**A System of Clinical Medicine for Practitioners** and Students, giving an account of the Diagnosis, Prognosis, and Treatment of Disease. By THOS. D. SAVILL, M.D., M.R.C.P. In 2 vols., post 8vo, with 4 Coloured Plates, and 200 Figures in the Text.

VOL. I.—**Local Diseases and Pyrexial Disorders.** 12s. 6d. net.

VOL. II.—**Diseases of the Skin, the Nervous System, etc.** (*shortly*).

**A Dictionary of Practical Medicine.** By various writers. Edited by JAS. KINGSTON FOWLER, M.A., M.D. F.R.C.P., Physician to Middlesex Hospital and the Hospital for Consumption. 8vo, cloth, 21s.

**The Practice of Medicine.** By M. Charteris, M.D., Professor of Therapeutics and Materia Medica in the University of Glasgow. Eighth Edition. Edited by F. J. CHARTERIS, M.B., Ch. B. Crown 8vo, with Engravings on Copper and Wood, 10s.

**A Text-Book of Bacteriology for Students and** Practitioners of Medicine. By G. M. STERNBERG, M.D., Surgeon-General, U.S. Army. Second Edition. With 9 Plates and 198 Figures in the Text. 8vo, 26s.

**How to Examine the Chest: a Practical** Guide for the use of Students. By SAMUEL WEST, M.D., F.R.C.P., Assistant Physician to St. Bartholomew's Hospital. Third Edition. With 46 Engravings. Fcap. 8vo, 5s.

**Atlas of the Pathological Anatomy of the Lungs.** By the late WILSON FOX, M.D., F.R.S. With 45 Plates (mostly Coloured) and Engravings. 4to, half-bound in Calf, 70s. net.

*By the same Author.*

**A Treatise on Diseases of the Lungs and** Pleura. Edited by SIDNEY COUPLAND, M.D., F.R.C.P., Physician to Middlesex Hospital. Roy. 8vo, with Engravings; also Portrait and Memoir of the Author, 36s. net.

**Guy's Hospital Reports.** By the Medical and Surgical Staff. Vol. XL. Third Series. 8vo, 10s. 6d.

**St. Thomas's Hospital Reports.** By the Medical and Surgical Staff. Vol. XXX. New Series. 8vo, 8s. 6d.

**Text-Book of Medical Treatment (Diseases** and Symptoms). By NESTOR I. C. TIRARD, M.D., F.R.C.P., Professor of the Principles and Practice of Medicine, King's College, London. 8vo, 15s.

**Student's Guide to Medical Diagnosis.** By SAMUEL FENWICK, M.D., F.R.C.P., and W. SOLTAU FENWICK, M.D., B.S. Ninth Edition. Crown 8vo, with 139 Engravings, 9s.

*By the same Authors.*

**Outlines of Medical Treatment.** Fourth Edition. Crown 8vo, with 35 Engravings, 10s.

*Also.*

**Ulcer of the Stomach and Duodenum.** With 55 Illustrations. Roy. 8vo, 10s. 6d.

*Also.*

**Cancer and other Tumours of the Stomach.** With 70 Illustrations, royal 8vo, 10s. 6d.

**On Syphonage and Hydraulic Pressure in the** Large Intestine, with their Bearing upon the Treatment of Constipation, Appendicitis, etc. By RALPH WINNINGTON LEFTWICH, M.D. Crown 8vo, 3s. net.

**The Schott Methods of the Treatment of** Chronic Diseases of the Heart, with an account of the Nauheim Baths, and of the Therapeutic Exercises. By W. BEZLY THORNE, M.D., M.R.C.P. Fourth Edition. 8vo, with Illustrations, 6s.

**The Liver.** By Lionel S. Beale, M.B., F.R.S., Consulting Physician to King's College Hospital With 24 Plates (85 Figures), 8vo, 5s.

*By the same Author.*

**On Slight Ailments: and on Treating Disease.** Fourth Edition. 8vo, 5s.

**The Blood: how to Examine and Diagnose** its Diseases. By ALFRED C. COLES, M.D., D.Sc., F.R.S. Edin. Second Edition. With 6 Coloured Plates. 8vo, 10s. 6d.

**The Physiology of the Carbohydrates; their** Application as Food and Relation to Diabetes. By F. W. PAVY, M.D., LL.D., F.R.S., F.R.C.P., Consulting Physician to Guy's Hospital. Royal 8vo, with Plates and Engravings, 10s. 6d.

**Uric Acid as a Factor in the Causation of Disease.** By ALEXANDER HAIG, M.D., F.R.C.P. Physician to the Metropolitan Hospital and the Royal Hospital for Children and Women. Sixth Edition. 8vo, with 75 Illustrations, 15s.

**Bronchial Asthma: its Pathology and Treatment.** By J. B. BERKART, M.D., late Physician to the City of London Hospital for Diseases of the Chest. Second Edition, with 7 Plates (35 Figures). 8vo, 10s. 6d.

**Treatment of Some of the Forms of Valvular Disease of the Heart.** By A. E. SANSOM, M.D., F.R.C.P., Physician to the London Hospital. Second Edition. Fcap. 8vo, with 26 Engravings, 4s. 6d.

**Medical Ophthalmoscopy: a Manual and Atlas.** By Sir WILLIAM R. GOWERS, M.D., F.R.C.P., F.R.S. Third Edition. Edited with the assistance of MARCUS GUNN, M.B., F.R.C.S., Surgeon to the Royal London Ophthalmic Hospital. With Coloured Plates and Woodcuts. 8vo, 16s.

*By the same Author.*

**A Manual of Diseases of the Nervous System.**

VOL. I.—**Nerves and Spinal Cord.** Third Edition, by the Author and JAMES TAYLOR, M.D., F.R.C.P. Roy. 8vo, with 192 Engravings, 15s.

VOL. II.—**Brain and Cranial Nerves: General and Functional Diseases of the Nervous System.** Second Edition. Roy. 8vo. with 182 Engravings, 20s.

*Also.*

**Clinical Lectures on Diseases of the Nervous System.** 8vo 7s. 6d.

*Also.*

**Subjective Sensations of Sight and Sound, Abiotrophy, and other Lectures on Diseases of the Nervous System.** With 18 Illustrations. 8vo, 6s. net.

*Also.*

**Epilepsy and other Chronic Convulsive Diseases: their Causes, Symptoms, and Treatment.** Second Edition. 8vo, 10s. 6d.

*Also.*

**Diagnosis of Diseases of the Brain.** Second Edition. 8vo, with Engravings, 7s. 6d.

*Also.*

**Syphilis and the Nervous System: being a Revised Reprint of the Lettsomian Lectures for 1890.** Delivered before the Medical Society of London. 8vo, 4s.

**The Nervous System, Diseases of.** By J. A. Ormerod, M.D., F.R.C.P., Physician to the National Hospital for the Paralysed and Epileptic. With 66 Illustrations. Fcap. 8vo, 8s. 6d.

**Text-Book of Nervous Diseases for Students** and Practitioners of Medicine. By Charles L. Dana, M.D., Professor of Nervous and Mental Diseases in Bellevue Hospital Medical College, New York. Fourth Edition. With 246 Illustrations. 8vo, 20s.

**Diseases of the Nervous System. Lectures** delivered at Guy's Hospital. By Sir Samuel Wilks, Bart., M.D., F.R.S. Second Edition. 8vo, 18s.

**Handbook of the Diseases of the Nervous** System. By James Ross, M.D., F.R.C.P., late Professor of Medicine in the Victoria University, and Physician to the Royal Infirmary, Manchester. Roy. 8vo, with 184 Engravings, 18s.

**Stammering: its Causes, Treatment, and** Cure. By A. G. Bernard, M.R.C.S., L.R.C.P. Crown 8vo, 2s.

**Diseases of the Nervous System. Clinical** Lectures. By Thomas Buzzard, M.D., F.R.C.P., Physician to the National Hospital for the Paralysed and Epileptic. With Engravings. 8vo, 15s.

*By the same Author.*

**Some Forms of Paralysis from Peripheral** Neuritis; of Gouty, Alcoholic, Diphtheritic, and other origin. Crown 8vo, 5s.

*Also.*

**On the Simulation of Hysteria by Organic** Disease of the Nervous System. Crown 8vo, 4s. 6d.

**Gout in its Clinical Aspects.** By J. Mortimer Granville, M.D. Crown 8vo, 6s.

**Rheumatic Diseases (Differentiation in).** By Hugh Lane, Surgeon to the Royal Mineral Water Hospital, Bath. Second Edition, much Enlarged, with 8 Plates. Crown 8vo, 3s. 6d.

**Diseases of the Abdomen, comprising those** of the Stomach and other parts of the Alimentary Canal, Œsophagus, Cæcum, Intestines, and Peritoneum. By S. O. HABERSHON, M.D., F.R.C.P. Fourth Edition. 8vo, with 5 Plates, 21s.

**On Gallstones, or Cholelithiasis.** By E. M. BROCKBANK, M.D. Vict., M.R.C.P. Lond., Honorary Physician to the Ancoats Hospital, Manchester. Crown 8vo, 7s.

**Obstinate Hiccough; the Physiology, Pathology,** and Treatment, based on a collection of over 150 cases from British and Foreign Works. By L. F. B. KNUTHSEN, M.D. Edin. Royal 8vo, 6s.

**Headaches: their Nature, Causes, and Treatment.** By W. H. DAY, M.D., Physician to the Samaritan Hospital. Fourth Edition. Crown 8vo, with Engravings, 7s. 6d.

**A Handbook of Medical Climatology, embodying** its Principles and Therapeutic Application, with Scientific Data of the chief Health Resorts of the World. By S. EDWIN SOLLY, M.D., M.R.C.S., late President of the American Climatological Association. With Engravings and Coloured Plates. 8vo, 16s.

**The Mineral Waters of France, and its** Wintering Stations (Medical Guide to). With a Special Map. By A. VINTRAS, M.D., Physician to the French Embassy, and to the French Hospital, London. Second Edition. Crown 8vo, 8s.

**Surgery: its Theory and Practice. By William** J. WALSHAM, F.R.C.S., Surgeon to St. Bartholomew's Hospital. Eighth Edition, by W. G. SPENCER, F.R.C.S., Surgeon to the Westminster Hospital. 8vo, with 622 Engravings (including 28 Skiagrams), 18s. net.

**A Synopsis of Surgery. By R. F. Tobin,** Surgeon to St. Vincent's Hospital, Dublin. Second Edition. Crown 8vo, interleaved, leather binding, 6s. 6d.

**Surgical Emergencies: together with the** Emergencies attendant on Parturition and the Treatment of Poisoning. By PAUL SWAIN, F.R.C.S., Surgeon to the South Devon and East Cornwall Hospital. Fifth Edition. Crown 8vo, with 149 Engravings, 6s.

**Illustrated Ambulance Lectures: (to which is** added a NURSING LECTURE) in accordance with the Regulations of the St. John's Ambulance Association for Male and Female Classes. By JOHN M. H. MARTIN, M.D., F.R.C.S., Hon. Surgeon to the Blackburn Infirmary. Fourth Edition. Crown 8vo, with 60 Engravings, 2s.

**Operations on the Brain (a Guide to).** By ALEC FRASER, Professor of Anatomy, Royal College of Surgeons in Ireland. Illustrated by 42 life-size Plates in Autotype, and 2 Woodcuts in the text. Folio, 63s. net.

**Abdominal Surgery.** By J. Greig Smith, M.A., F.R.S.E. Sixth Edition. Edited by JAMES SWAIN, M.S., M.D. Lond., F.R.C.S. Eng., Assistant-Surgeon to the Bristol Royal Infirmary, Professor of Surgery, University College, Bristol. 2 vols., 8vo, with 224 Engravings, 36s.

**The Physiology of Death from Traumatic Fever**; a Study in Abdominal Surgery. By JOHN D. MALCOLM, M.B., C.M., F.R.C.S.E., Surgeon to the Samaritan Free Hospital. 8vo, 3s. 6d.

**The Surgery of the Alimentary Canal.** By ALFRED ERNEST MAYLARD, M.B. Lond. and B.S., Senior Surgeon to the Victoria Infirmary, Glasgow. With 27 Swantype Plates and 89 Figures in the Text, 8vo, 25s.

*By the same Author.*

**A Student's Handbook of the Surgery of the Alimentary Canal.** With 97 Illustrations. Crown 8vo, 8s. 6d.

**Hare-Lip and Cleft Palate.** By R. W. MURRAY, F.R.C S., Surgeon, David Lewis Northern Hospital, late Surgeon, Liverpool Infirmary for Children. With 25 Illustrations, 8vo, 3s.

**Modern Bullet-Wounds and Modern Treatment**, with Special Regard to Long Bones and Joints, Field Appliances and First Aid. Part of the Alexander Essay for 1903. By Major F. SMITH, D.S.O., R.A.M.C. Crown 8vo, 3s. net.

**The Practice of Surgery: a Manual.** By THOMAS BRYANT, Consulting Surgeon to Guy's Hospital. Fourth Edition. 2 vols. crown 8vo, with 750 Engravings (many being Coloured), and including 6 chromo plates, 32s.

**The Operations of Surgery: intended for use** on the Dead and Living Subject alike. By W. H. A. JACOBSON, M.Ch. Oxon., F.R.C.S., Surgeon Guy's Hospital, and F. J. STEWARD, M.S. Lond., F.R.C.S., Assistant Surgeon, Guy's Hospital. Fourth Edition. 2 vols., roy. 8vo, with 550 Illustrations, 42s.

**Muco-Membranous Entero-Colitis; Symptoms**, Complications, Etiology, and Treatment. By MAURICE DE LANGENHAGEN, M.D., Consulting Physician at Plombières, Vosges, France. Fcap. 8vo, 3s. 6d.

**A Course of Operative Surgery.** By Christopher Heath, Surgeon to University College Hospital. Second Edition. With 20 Coloured Plates (180 figures). 30s. net.

*By the same Author.*

**A Manual of Minor Surgery and Bandaging.** Twelfth Edition, revised by Bilton Pollard, F.R.C.S., Surgeon to University College Hospital. Fcap. 8vo, with 195 Engravings, 6s. 6d.

*Also.*

**Injuries and Diseases of the Jaws.** Fourth Edition. Edited by Henry Percy Dean, M.S., F.R.C.S., Assistant Surgeon to the London Hospital. 8vo, with 187 Wood Engravings, 14s.

*Also.*

**Clinical Lectures on Surgical Subjects** delivered at University College Hospital. First Series, 6s.; Second Series, 6s.

**Clinical Essays and Lectures.** By Howard Marsh, F.R.C.S., Professor of Surgery in the University of Cambridge, late Surgeon to St. Bartholomew's Hospital. With 26 Illustrations, 8vo, 7s. 6d.

**Ovariotomy and Abdominal Surgery.** By Harrison Cripps, F.R.C.S., Surgical Staff, St. Bartholomew's Hospital. With numerous Plates, royal 8vo, 25s.

**Diseases of Bones and Joints.** By Charles Macnamara, F.R.C.S., Surgeon to, and Lecturer on Surgery at, the Westminster Hospital. 8vo, with Plates and Engravings, 12s.

**Surgical Pathology and Morbid Anatomy.** By Anthony A. Bowlby, F.R.C.S., Assistant Surgeon to St. Bartholomew's Hospital. Fourth Edition. Crown 8vo, with 186 Engravings, 10s. 6d.

*By the same Author.*

**Injuries and Diseases of Nerves, and their** Surgical Treatment. 8vo, with 20 Plates, 14s.

**Chloroform: a Manual for Students and Practitioners.** By Edward Lawrie, M.B. Edin., Lieut.-Col. I.M.S., Residency Surgeon, Hyderabad. Illustrated, crown 4to, 5s. net.

**A Pocket Guide to Anæsthetics for the** Student and General Practitioner. By Thomas D. Luke, M.B., F.R.C.S., Edinburgh. With 43 Engravings, crown 8vo, 5s. net.

**Diseases of the Thyroid Gland and their** Surgical Treatment. By JAMES BERRY, B.S. Lond., F.R.C.S., Surgeon to the Royal Free Hospital and Lecturer on Surgery at the London School of Medicine for Women. With 121 Illustrations, 8vo, 14s.

**The Human Foot: its Form and Structure,** Functions and Clothing. By THOMAS S. ELLIS, Consulting Surgeon to the Gloucester Infirmary. With 7 Plates and Engravings (50 Figures). 8vo, 7s. 6d.

**Deformities and Deficiencies of the Head and** Neck. By HEATHER BIGG, F.R.C.S. Ed. 8vo, 2s. 6d.

**Face and Foot Deformities. By Frederick** CHURCHILL, C.M. 8vo, with Plates and Illustrations, 10s. 6d.

**Royal London Ophthalmic Hospital Reports.** By the Medical and Surgical Staff. Vol. XV., Part 3. 8vo, 5s.

**Ophthalmological Society of the United King-** dom. Transactions. Vol. XXIII. 8vo, 12s. 6d. net.

**Manual of Ophthalmic Surgery and Medicine.** By W. H. H. JESSOP, M.A., F.R.C.S., Ophthalmic Surgeon to St. Bartholomew's Hospital. With 5 Coloured Plates and 110 Woodcuts Crown 8vo, 9s. 6d.

**Nettleship's Diseases of the Eye.** Sixth Edition. Revised and Edited by W. T. HOLMES SPICER, M.B., F.R.C.S., Ophthalmic Surgeon to St. Bartholomew's Hospital Surgeon to the Royal London Ophthalmic Hospital. With 161 Engravings and a Coloured Plate illustrating Colour-Blindness. Crown 8vo, 8s. 6d.

**Diseases and Refraction of the Eye. By** N. C. MACNAMARA, F.R.C.S., Surgeon to Westminster Hospital, and GUSTAVUS HARTRIDGE, F.R.C.S., Surgeon to the Royal Westminster Ophthalmic Hospital. Fifth Edition. Crown 8vo, with Plate, 156 Engravings, also Test-types, 10s. 6d.

**Practical Handbook of Diseases of the Eye.** By D. CHALMERS WATSON, M.B., Ophthalmic Physician, Marshall Street Dispensary, Edinburgh. Second Edition, with 9 Coloured Plates and 31 Figures in the Text, fcap. 8vo, 5s. net.

**Elementary Ophthalmic Optics, including** Ophthalmoscopy and Retinoscopy. By J. HERBERT PARSONS, B.S., B.Sc., F.R.C.S., Curator, Royal London Ophthalmic Hospital. With 66 Illustrations, 8vo, 6s. 6d.

**Ocular Therapeutics according to the most** Recent Discoveries. By Dr. A. Darier. Translated by Sydney Stephenson, M.B., C.M., Ophthalmic Surgeon to the Evelina Hospital and the North-Eastern Hospital for Children. Roy. 8vo, 10s. 6d. net.

**Atlas of Ophthalmoscopy. Composed of 12** Chromo-lithographic Plates (59 Figures drawn from nature) and Explanatory Text. By Richard Liebreich, M.R.C.S. Translated by H. Rosborough Swanzy, M.B. Third Edition, 4to, 40s. net.

**Refraction of the Eye: a Manual for Students.** By Gustavus Hartridge, F.R.C.S., Surgeon to the Royal Westminster Ophthalmic Hospital. Twelfth Edition. Crown 8vo, with 105 Illustrations, also Test-types, etc., 6s.

*By the same Author.*

**The Ophthalmoscope: a Manual for Students.** Fourth Edition. Crown 8vo, with 65 Illustrations and 4 Plates, 4s. 6d.

**Glaucoma: its Pathology and Treatment. By** Priestley Smith, Ophthalmic Surgeon to the Queen's Hospital Birmingham. 8vo, with 64 Engravings and 12 Zinco-photographs. 7s. 6d.

**Methods of Operating for Cataract and** Secondary Impairments of Vision, with the results of 500 cases. By Major G. H. Fink, H.M. Indian Medical Service. Crown 8vo, with 15 Engravings, 5s.

**Diseases of the Eye: a Practical Handbook** for General Practitioners and Students. By Cecil Edward Shaw, M.D., M.Ch., Ophthalmic Surgeon to the Ulster Hospital for Children and Women, Belfast. With a Test-Card for Colour-Blindness. Crown 8vo, 3s. 6d.

**The Accessory Sinuses of the Nose, their** Surgical Anatomy and the Diagnosis and Treatment of their Inflammatory Affections. By A. Logan Turner, M.D. Edin., Surgeon for Diseases of the Ear and Throat, Deaconess Hospital, Edinburgh. With 81 Illustrations, imp. 8vo, 12s. net.

**Diseases of the Ear, including the Anatomy** and Physiology of the Organ, together with the Treatment of the Affections of the Nose and Pharynx, which conduce to Aural Disease (a Treatise). By T. Mark Hovell, Senior Aural Surgeon to the London Hospital, and Lecturer on Diseases of the Throat in the College. Second Edition. 8vo, with 128 Engravings, 21s.

**Diseases and Injuries of the Ear.** By Sir William B. Dalby, F.R.C.S., M.B., Consulting Aural Surgeon to St. George's Hospital. Fourth Edition. Crown 8vo, with 8 Coloured Plates and 38 Wood Engravings. 10s. 6d.

*By the same Author.*

**Short Contributions to Aural Surgery, between** 1875 and 1896. Third Edition. 8vo, with Engravings, 5s.

**A System of Dental Surgery.** By Sir John Tomes, F.R.S., and C. S. Tomes, M.A., F.R.S. Fourth Edition. Post 8vo, with 289 Engravings, 16s.

**Dental Anatomy, Human and Comparative:** A Manual. By Charles S. Tomes, M.A., F.R.S. Fifth Edition. Post 8vo, with 263 Engravings, 14s.

**Decay in Teeth: an Investigation into its** Cause and Prevention. By J. Sim Wallace, M.D., D.Sc., L.D.S.R.C.S. Second Edition. 8vo, 5s.

**Dental Materia Medica, Pharmacology and** Therapeutics. By Charles W. Glassington, M.R.C.S., L.D.S. Edin.; Senior Dental Surgeon, Westminster Hospital; Dental Surgeon, National Dental Hospital, and Lecturer on Dental Materia Medica and Therapeutics to the College. Crown 8vo, 6s.

**Dental Medicine: a Manual of Dental Materia** Medica and Therapeutics. By Ferdinand J. S. Gorgas, M.D., D.D.S., Professor of the Principles of Dental Science in the University of Maryland. Sixth Edition. 8vo, 18s.

**A Manual of Dental Metallurgy.** By Ernest A. Smith, Assay Office, Sheffield. Second Edition. With 38 Illustrations, crown 8vo, 6s. 6d.

**A Manual of Nitrous Oxide Anæsthesia.** By J. Frederick W. Silk, M.D. Lond., M.R.C.S., Assistant Anæsthetist to Guy's Hospital, Anæsthetist to the Dental School of Guy's Hospital, and to the Royal Free Hospital. 8vo, with 26 Engravings, 5s.

**Practical Treatise on Mechanical Dentistry.** By Joseph Richardson, M.D., D.D.S. Seventh Edition, revised and edited by George W. Warren, D.D.S. Royal 8vo, with 690 Engravings, 22s.

**A Handbook on Leprosy.** By S. P. Impey, M.D., late Chief and Medical Superintendent, Robben Island Leper and Lunatic Asylums, Cape Colony. With 38 Plates, 8vo, 12s.

**Diseases of the Skin (Introduction to the Study of).** By P. H. Pye-Smith, M.D., F.R.S., F.R.C.P., Physician to Guy's Hospital. Crown 8vo, with 26 Engravings, 7s. 6d.

**A Manual of Diseases of the Skin, with an Analysis of 20,000 Consecutive Cases and a Formulary.** By Duncan E. Bulkley, M.D., New York. Fourth Edition, royal 16mo, 6s. 6d.

**Skin Diseases of Children.** By Geo. H. Fox, M.D., Clinical Professor of Diseases of the Skin, College of Physicians and Surgeons, New York. With 12 Photogravure and Chromographic Plates and 60 Illustrations in the Text. Royal 8vo, 12s. 6d.

**The Operative Surgery of Malignant Disease.** By Henry T. Butlin, F.R.C.S., Surgeon to St. Bartholomew's Hospital. Second Edition, with 12 Engravings. 8vo, 14s.

*By the same Author.*

**Malignant Disease (Sarcoma and Carcinoma) of the Larynx.** 8vo, with 5 Engravings, 5s.

**Cancers and the Cancer Process: a Treatise,** Practical and Theoretic. By Herbert L. Snow, M.D., Surgeon to the Cancer Hospital, Brompton. 8vo, with 15 Plates. 15s.

*By the same Author.*

**The Re-appearance (Recurrence) of Cancer after apparent Extirpation.** 8vo, 5s. 6d.

*Also.*

**The Palliative Treatment of Incurable Cancer.** Crown 8vo, 2s. 6d.

**The Diagnosis and Treatment of Syphilis.** By Tom Robinson, M.D. St. And., Physician to the Western Skin Hospital. Second Edition. Crown 8vo, 3s. 6d.

*By the same Author.*

**The Diagnosis and Treatment of Eczema.** Second Edition. Crown 8vo, 3s. 6d.

*Also.*

**Illustrations of Diseases of the Skin and Syphilis, with Remarks.** Fasc. I. with 3 Plates. Imp. 4to, 5s. net.

**Cancerous Affections of the Skin (Epithelioma** and Rodent Ulcer). By GEORGE THIN, M.D. Post 8vo, with 8 Engravings, 5s.

*By the same Author.*

**Pathology and Treatment of Ringworm.** 8vo, with 21 Engravings, 5s.

**Ringworm, and some other Scalp Affections:** their Cause and Cure. By HAYDN BROWN, L.R.C.P. Ed. 8vo, 5s.

**Urinary and Renal Derangements and Calculous** Disorders. By LIONEL S. BEALE, F.R.C.P., F.R.S., Physician to King's College Hospital. 8vo, 5s.

**Chemistry of Urine: a Practical Guide to the** Analytical Examination of Diabetic, Albuminous, and Gouty Urine. By ALFRED H. ALLEN, F.I.C., F.C.S., Public Analyst for the West Riding of Yorkshire, &c. 8vo, with Engravings, 7s. 6d.

**Atlas of Electric Cystoscopy. By E. Hurry** FENWICK, F.R.C.S., Surgeon to the London Hospital and St. Peter's Hospital for Stone. Royal 8vo, with 34 Coloured Plates, embracing 83 Figures. 21s. net.

*By the same Author.*

**Electric Illumination of the Bladder and** Urethra, as a Means of Diagnosis of Obscure Vesico-Urethral Diseases. By E. HURRY FENWICK, F.R.C.S., Surgeon to London Hospital and St. Peter's Hospital for Stone. Second Edition. 8vo, with 54 Engravings, 6s. 6d.

*Also.*

**Operative and Inoperative Tumours of the** Urinary Bladder: a Clinical and Operative Study based on 500 cases. With 39 Illustrations, 8vo, 5s. net.

*Also.*

**Tumours of the Urinary Bladder. Fas. I.** Royal 8vo, 5s. net.

*Also.*

**Ulceration of the Bladder, Simple, Tuber-** culous, and Malignant: a Clinical Study. With Illustrations, 8vo, 5s.

*Also.*

**The Cardinal Symptoms of Urinary Disease:** their Diagnostic Significance and Treatment. 8vo, with 36 Illustrations, 8s. 6d.

*Also.*

**Obscure Diseases of the Urethra.** With 63 Illustrations, 8vo, 6s. 6d.

*Also.*

**Obscure Diseases of the Kidney, the Value of** Ureteric Meatoscopy in their Diagnosis and Treatment. Based on 400 Major Operations on the Kidney. With 14 Plates and 43 Figures in the Text. 8vo, 6s. 6d.

*By SIR HENRY THOMPSON, BART., F.R.C.S.*

**Diseases of the Urinary Organs. Clinical** Lectures. Eighth Edition. 8vo, with 121 Engravings, 10s. 6d.

**Practical Lithotomy and Lithotrity; or, an** Inquiry into the Best Modes of Removing Stone from the Bladder. Third Edition. 8vo, with 87 Engravings, 10s.

**The Preventive Treatment of Calculous Dis-** ease, and the Use of Solvent Remedies. Third Edition. Cr. 8vo, 2s. 6d.

**Tumours of the Bladder: their Nature, Sym-** ptoms, and Surgical Treatment. 8vo, with numerous Illustrations, 5s.

**Stricture of the Urethra, and Urinary Fistulæ:** their Pathology and Treatment. Fourth Edition. 8vo, with 74 Engravings, 6s.

---

**The Clinical Examination of Urine, with an** Atlas of Urinary Deposits. By LINDLEY SCOTT, M.A., M.D., with 41 original Plates (mostly in colours). Crown 4to, 15s. net.

**Clinical Chemistry of Urine (Outlines of the).** By C. A. MACMUNN, M.A., M.D. 8vo, with 64 Engravings and Plate of Spectra, 9s.

**The Surgical Diseases of the Genito-Urinary** Organs, including Syphilis. By E. L. KEYES, M.D., Professor of Genito-Urinary Surgery, Syphilology, and Dermatology in Bellevue Hospital Medical College, New York (a revision of VAN BUREN and KEYES' Text-book). Roy. 8vo, with 114 Engravings, 21s.

**Selected Papers on Stone, Prostate, and other** Urinary Disorders. By REGINALD HARRISON, F.R.C.S., Surgeon to St. Peter's Hospital. 8vo, with 15 Illustrations, 5s.

**Syphilis. By Sir Alfred Cooper, F.R.C.S., Con-** sulting Surgeon to the West London and the Lock Hospitals. Second Edition. Edited by EDWARD COTTERELL, F.R.C.S., Surgeon (out-patients) to the London Lock Hospital. 8vo, with 24 Full-page Plates (12 coloured), 18s.

---

**On Maternal Syphilis, including the presence** and recognition of Syphilitic Pelvic Disease in Women. By JOHN A. SHAW-MACKENZIE, M.D. With Coloured Plates. 8vo, 10s. 6d.

**Diseases of the Rectum and Anus. By Sir** ALFRED COOPER, F.R.C.S., Senior Surgeon to St. Mark's Hospital for Fistula; and F. SWINFORD EDWARDS, F.R.C.S., Senior Assistant Surgeon to St. Mark's Hospital. Second Edition, with Illustrations. 8vo, 12s.

**Cancer of the Rectum. Especially considered** with regard to its Surgical Treatment. Jacksonian Prize Essay. By HARRISON CRIPPS, F.R.C.S., Assistant Surgeon to St. Bartholomew's Hospital. Third Edition. 8vo, with 13 Plates and several Wood Engravings, 6s.

*Also*

**The Passage of Air and Fæces from the** Urethra. 8vo, 3s. 6d.

**A Medical Vocabulary: an Explanation of all** Terms and Phrases used in the various Departments of Medical Science and Practice, their Derivation, Meaning, Application, and Pronunciation. By R. G. MAYNE, M.D., LL.D. Seventh Edition, by W. W. WAGSTAFFE, B.A., F.R.C.S., and G. D. PARKER, M.B. Crown 8vo, 12s. 6d.

**A Short Dictionary of Medical Terms. Being** an Abridgment of Mayne's Vocabulary. 64mo, 2s. 6d.

**Dunglison's Dictionary of Medical Science.** Containing a full Explanation of its various Subjects and Terms, with their Pronunciation, Accentuation, and Derivation. Twenty-second Edition. By RICHARD J. DUNGLISON, A.M., M.D. Royal 8vo, 30s.

**Terminologia Medica Polyglotta: a Concise** International Dictionary of Medical Terms (French, Latin, English, German, Italian, Spanish, and Russian). By THEODORE MAXWELL, M.D., B.Sc., F.R.C.S. Edin. Royal 8vo, 16s.

**A German-English Dictionary of Medical** Terms. By Sir FRED. TREVES, Bart, K.C.V.O., Surgeon to the London Hospital; and HUGO LANG, B.A. Crown 8vo, half-Persian calf, 12s.

**A Handbook of Physics and Chemistry, adapted** to the requirements of the first examination of the Conjoint Board and for general use. By HERBERT E. CORBIN, B.Sc. Lond., and ARCHIBALD M. STEWART, B.Sc. Lond. Second Edition. With 126 Illustrations, crown 8vo, 6s. 6d.

**A Manual of Chemistry, Theoretical and Practical.** By WILLIAM A. TILDEN, D.Sc., F.R.S, Professor of Chemistry in the Royal College of Science, London; Examiner in Chemistry to the Department of Science and Art. With 2 Plates and 143 Woodcuts, crown 8vo, 10s.

**Chemistry, Inorganic and Organic. With Experiments.** By CHARLES L. BLOXAM. Ninth Edition, by JOHN MILLAR THOMSON, F.R.S., Professor of Chemistry in King's College, London, and ARTHUR G. BLOXAM, F.I.C., Consulting Chemist and Chartered Patent Agent. 8vo, with 284 Engravings, 18s. net.

*By the same Author.*

**Laboratory Teaching; or, Progressive Exercises in Practical Chemistry.** Sixth Edition, by ARTHUR G. BLOXAM. Crown 8vo, with 80 Engravings, 6s. 6d.

**Watts' Organic Chemistry. Edited by William A. TILDEN, D.Sc., F.R.S., Professor of Chemistry, Royal College of Science, London. Second Edition. Crown 8vo, 10s.**

**Practical Chemistry, and Qualitative Analysis.** By FRANK CLOWES, D.Sc. Lond., Emeritus Professor of Chemistry in the University College, Nottingham. Seventh Edition. Post 8vo, with 101 Engravings and Frontispiece, 8s. 6d.

**Quantitative Analysis. By Frank Clowes, D.Sc. Lond., Emeritus Professor of Chemistry in the University College, Nottingham, and J. BERNARD COLEMAN, Assoc. R. C. Sci. Dublin; Professor of Chemistry, South-West London Polytechnic. Sixth Edition. Post 8vo, with 125 Engravings, 10s.**

*By the same Authors.*

**Elementary Practical Chemistry.** Fourth Edition.

Part I. General Chemistry. With 75 Engravings, 2s. 6d. net.
Part II. Analytical Chemistry, with 20 Engravings. 2s. 6d. net.

*Also*

**Introduction to Quantitative Analysis.** With 62 Engravings, Post 8vo, 4s. 6d.

**Qualitative Analysis. By R. Fresenius. Translated by CHARLES E. GROVES, F.R.S. Tenth Edition. 8vo, with Coloured Plate of Spectra and 46 Engravings, 15s.**

*By the same Author.*

**Quantitative Analysis.** Seventh Edition.

VOL. I., Translated by A. VACHER. 8vo, with 106 Engravings, 15s.

VOL. II., Translated by C. E. GROVES, F.R.S. 8vo with 143 Engravings, 20s.

**Inorganic Chemistry.** By Sir Edward Frankland, K.C.B., D.C.L., LL.D., F.R.S., and Francis R. Japp, M.A., Ph.D., F.I.C., F.R.S. 8vo, with numerous Illustrations, 24s.

**Inorganic Chemistry (A System of).** By Sir William Ramsay, K.C.B., Ph.D., F.R.S., Professor of Chemistry in the University College, London. 8vo, with Engravings, 15s.

*By the same Author.*

**Elementary Systematic Chemistry for the Use** of Schools and Colleges. With Engravings. Crown 8vo, 4s. 6d. Interleaved, 5s. 6d.

**Valentin's Practical Chemistry and Qualitative** and Quantitative Analysis. Edited by Dr. W. R. Hodgkinson, F.R.S.E., Professor of Chemistry and Physics at the Royal Military Academy, and Artillery College, Woolwich. Ninth Edition. 8vo, with Engravings and Map of Spectra. 9s. (The Tables separately, 2s. 6d.)

**Practical Chemistry, Part I.** Qualitative Exercises and Analytical Tables for Students. By J. Campbell Brown, Professor of Chemistry in Victoria University and University College, Liverpool. Fourth Edition. 8vo, 2s. 6d.

**The Analyst's Laboratory Companion: a Col**lection of Tables and Data for Chemists and Students. By Alfred E. Johnson, F.I.C. Second Edition. Cloth, 5s.; leather, 6s. 6d.

**The Chemistry of the Terpenes.** By F. Heusler, Ph.D. Translated and enlarged by F. J. Pond, M.A., Ph.D. 8vo, 17s. net.

**Commercial Organic Analysis: a Treatise on** the Properties, Modes of Assaying, Proximate Analytical Examination, etc., of the various Organic Chemicals and Products employed in the Arts, Manufactures, Medicine, etc. By Alfred H. Allen, F.I.C.

*Third Edition.*

Vol. I., 18s.; Vol. II., Part I., 14s.; Vol. II., Part II., 14s.; Vol. II., Part III., ; Vol. III., Part I., 18s.

*Second Edition.*

Vol. III., Pt. II., 18s.; Vol. III., Pt. III., 16s.

Vol. IV., completing the work, 18s.

**Volumetric Analysis: or the Quantitative** Estimation of Chemical Substances by Measure, applied to Liquids, Solids, and Gases. By Francis Sutton, F.C.S., F.I.C., Public Analyst for the County of Norfolk. Eighth Edition. 8vo, with 116 Engravings, 20s.

**Chemical Technology; or, Chemistry in its** Applications to Arts and Manufactures. Edited by CHARLES E. GROVES, F.R.S., WILLIAM THORP, B.Sc., and W. J. DIBDIN, F.I.C.

**VOL. I.—Fuel and its Applications. By E. J.** MILLS, D.Sc., F.R.S., and F. J. ROWAN, C.E. Royal 8vo, with 606 Engravings, 30s.

**VOL. II.—Lighting.—Fats and Oils, by W. Y.** DENT. STEARINE INDUSTRY, by J. MCARTHUR. CANDLE MANUFACTURE, by L. FIELD and F. A. FIELD. THE PETROLEUM INDUSTRY AND LAMPS, by BOVERTON REDWOOD. MINERS' SAFETY LAMPS, by B. REDWOOD and D. A. LOUIS. Royal 8vo, with 358 Engravings and Map, 20s.

**VOL III.—Gas Lighting. By Charles Hunt.** With 2 Plates and 292 Engravings, 8vo, 18s.

**VOL. IV.—Electric Lighting. By A. G.** COOKE, M.A., A.M.I.E.E. PHOTOMETRY. By W. J. DIBDIN, F.I.C., F.C.S. Royal 8vo., with 10 Plates and 181 Engravings, 20s.

**Cooley's Cyclopædia of Practical Receipts,** and Collateral Information in the Arts, Manufactures, Professions, and Trades: including Medicine, Pharmacy, Hygiene, and Domestic Economy. Seventh Edition, by W. NORTH, M.A. Camb., F.C.S. 2 Vols., Roy. 8vo, with 371 Engravings, 42s.

**Chemical Technology: a Manual. By Rudolf** VON WAGNER. Translated and Edited by Sir WILLIAM CROOKES, F.R.S., from the Thirteenth Enlarged German Edition as remodelled by Dr. FERDINAND FISCHER. 8vo, with 596 Engravings, 32s.

**Technological Handbooks. Edited by John** GARDNER, F.I.C., F.C.S., and JAMES CAMERON, F.I.C.

**Brewing, Distilling, and Wine Manu-** facture. Crown 8vo, with Engravings, 6s. 6d.

**Soaps and Candles.** Crown 8vo, with 54 Engravings, 7s.

**Chemistry an Exact Mechanical Philosophy.** By FRED. G. EDWARDS. Illustrated, 8vo, 3s. 6d.

**The Quarterly Journal of Microscopical Science.** Edited by E. RAY LANKESTER, M.A., LL.D., F.R.S.; with the co-operation of ADAM SEDGWICK, M.A., F.R.S., W.F. R. WELDON, M.A., F.R.S., and SYDNEY J. HICKSON, M.A., F.R.S. Each Number, 10s.

**Methods and Formulæ used in the Preparation** of Animal and Vegetable Tissues for Microscopical Examination, including the Staining of Bacteria. By PETER WYATT SQUIRE, F.L.S. Crown 8vo, 3s. 6d.

**The Microscope and its Revelations. By the** late WILLIAM B. CARPENTER, C.B., M.D., LL.D., F.R.S. Eighth Edition, by the Rev. W. H. DALLINGER, LL.D., F.R.S. With 23 Plates and more than 800 Wood Engravings. 8vo, 28s. Half Calf, 32s.

**The Microtomist's Vade-Mecum: a Handbook** of the Methods of Microscopic Anatomy. By ARTHUR BOLLES LEE. Fifth Edition, 8vo, 15s.

**Photo-Micrography (Guide to the Science of).** By EDWARD C. BOUSFIELD, L.R.C.P. Lond. 8vo, with 34 Engravings and Frontispiece, 6s.

**A Treatise on Physics. By Andrew Gray,** LL.D., F.R.S., Professor of Natural Philosophy in the University of Glasgow. Vol. I. Dynamics and Properties of Matter. With 350 Illustrations, 8vo. 15s.

**An Introduction to Physical Measurements,** with Appendices on Absolute Electrical Measurements, etc. By Dr. F. KOHLRAUSCH. Third English Edition, by T. H. WALLER, B.A., B.Sc., and H. R. PROCTER, F.I.C., F.C.S. 8vo, with 91 Illustrations, 12s. 6d.

**Tuson's Veterinary Pharmacopœia, including** the Outlines of Materia Medica and Therapeutics. Fifth Edition Edited by JAMES BAYNE, F.C.S., Professor of Chemistry and Toxicology in the Royal Veterinary College. Crown 8vo, 7s. 6d.

**The Veterinarian's Pocket Remembrancer:** being Concise Directions for the Treatment of Urgent or Rare Cases, By GEORGE ARMATAGE, M.R.C.V.S. Second Edition. Post 8vo, 3s.

**Chauveau's Comparative Anatomy of the** Domesticated Animals. Revised and Enlarged, with the Co-operation of S. ARLOING, Director of the Lyons Veterinary School, and Edited by GEORGE FLEMING, C.B., LL.D., F.R.C.V.S., late Principal Veterinary Surgeon of the British Army. Second English Edition. 8vo, with 585 Engravings, 31s. 6d.

**Encyclopædia Medica. Edited by Chalmers** Watson, M.B., M.R.C.P.E. In 13 Volumes, £13 net.

Allen's Chemistry of Urine, 22
——— Commercial Organic Analysis, 26
Andrewes on Disinfection, 4
Armatage's Veterinary Pocket Remembrancer, 28

Ballantyne's Antenatal Pathology and Hygiene, 2
Barnes' (R.) Obstetric Operations, 6
——— Diseases of Women, 6
Beale (L. S.) on Liver, 12
——— Slight Ailments, 12
——— Urinary and Renal Derangements, 22
Beale (P. T. B.) on Elementary Biology, 3
Beasley's Book of Prescriptions, 8
——— Druggists' General Receipt Book, 8
——— Pharmaceutical Formulary, 8
Bell on Sterility, 6
Bentley and Trimen's Medicinal Plants, 9
Bentley's Systematic Botany, 9
Berkart's Bronchial Asthma, 13
Bernard on Stammering, 14
Berry's (Jas.) Thyroid Gland, 18
——— (R. J.) Regional Anatomy, 2
Bigg's Short Manual of Orthopædy, 18
Birch's Practical Physiology, 3
Bloxam's Chemistry, 25
——— Laboratory Teaching, 25
Bousfield's Photo-Micrography, 28
Bowlby's Injuries and Diseases of Nerves, 17
——— Surgical Pathology and Morbid Anatomy, 17
Brockbank on Gallstones, 15
Brown's (Haydn) Ringworm, 22
Brown's Practical Chemistry, 26
Bryant's Practice of Surgery, 16
Bulkley on Skin, 21
Burdett's Hospitals and Asylums of the World, 5
Butlin's Operative Surgery of Malignant Disease, 21
——— Malignant Disease of the Larynx, 21
Buzzard's Diseases of the Nervous System, 14
Buzzard's Peripheral Neuritis, 14
——— Simulation of Hysteria, 14

Cameron's Soaps and Candles, 27
Carpenter and Dallinger on the Microscope, 28
Cautley on Feeding Infants, 7
Charteris' Practice of Medicine, 11
Chauveau's Comparative Anatomy, 28
Churchill's Face and Foot Deformities, 18
Clouston's Lectures on Mental Diseases, 4
Clowes and Coleman's Quantitative Analysis, 25
Clowes and Coleman's Elementary Practical Chemistry, 25
Clowes' Practical Chemistry, 25
Coles on Blood, 12
Cooke's Electric Lighting, 27
Cooley's Cyclopædia of Practical Receipts, 27
Cooper's Syphilis, 23
Cooper and Edwards' Diseases of the Rectum, 24
Corbin and Stewart's Physics and Chemistry, 24
Cripps' (H.) Ovariotomy and Abdominal Surgery, 17
——— Cancer of Rectum, 24
——— Air and Fæces in Urethra, 24
Cripps' (R. A.) Galenic Pharmacy, 8
Cuff's Lectures to Nurses, 7
Cullingworth's Monthly Nurses, 7

Dalby's Diseases and Injuries of the Ear, 20
———Short Contributions, 20
Dana on Nervous Diseases, 14
Darier's Ocular Therapeutics, 19
Day on Headaches, 15
De Langenhagen on Entero-Colitis 16
Dibdin's Photometry, 27
Duncan (A.) on Prevention of Diseases in Tropics, 10
Dunglison's Med. Dictionary, 24

Edwards' Chemistry, 27
Ellis's (T. S.) Human Foot, 18
Encyclopædia Medica, 28

Fagge's Principles and Practice of Medicine, 10
Fayrer's Climate and Fevers of India, 10

[*Continued on next page.*

Fenwick (E. H.), Electric Illumination of Bladder, 22
——— Obscure Diseases of the Urethra, 22
——— Atlas of Electric Cystoscopy, 22
——— Kidney, 22
——— Symptoms of Urinary Disease, 22
——— Tumours of Bladder, 22
——— Ulceration of Bladder, 22
Fenwick's (S.) Medical Diagnosis, 12
——— Ulcer of Stomach, 12
——— Cancer of Stomach, 12
——— Outlines of Medical Treatment, 12
Fink's Operating for Cataract, 19
Fowler's Dictionary of Practical Medicine, 11
Fox (G. H.) on Skin Diseases of Children, 21
Fox (Wilson), Atlas of Pathological Anatomy of the Lungs, 11
—— Treatise on Diseases of the Lungs, 11
Frankland and Japp's Inorganic Chemistry, 26
Fraser's Operations on the Brain, 16
Fresenius' Qualitative Analysis, 25
——— Quantitative Analysis, 25

Galabin's Diseases of Women, 6
——— Manual of Midwifery, 5
Gardner's Brewing, Distilling, and Wine Manufacture, 27
Glassington's Dental Materia Medica, 20
Godlee's Atlas of Human Anatomy, 1
Goodhart's Diseases of Children, 7
Gorgas' Dental Medicine, 20
Gowers' Diagnosis of Brain Disease, 13
———Diseases of Nervous System, 13
———Medical Ophthalmoscopy, 13
———Subjective Sensations, 13
———Syphilis and the Nervous System, 13
———Epilepsy, 13
Granville on Gout, 14
Gray's Treatise on Physics, 28
Green's Manual of Botany, 9
———Vegetable Physiology, 9
Greenish's Materia Medica, 8
——— Microscopical Examination of Foods and Drugs, 8
Greenish's and Collin's Vegetable Powders, 8
Groves, Thorp, and Dibdin's Chemical Technology, 27
Guy's Hospital Reports, 11

Habershon's Diseases of the Abdomen, 15
Hadley on Nursing, 6
Haig s Uric Acid, 13
——— Diet and Food, 4
Hamer's Manual of Hygiene, 3
Harrison's Urinary Organs, 23
Hartridge's Refraction of the Eye, 19
———Ophthalmoscope, 19
Hawthorne's Galenical Preparations, 8
Heath's Injuries and Diseases of the Jaws, 17
——— Minor Surgery and Bandaging, 17
——— Operative Surgery, 17
——— Practical Anatomy, 1
——— Clinical Lectures, 17
Hedley's Therapeutic Electricity, 9
Heusler on the Terpenes, 26
Hewlett's Bacteriology, 4
——— Serum Therapy, 4
Hill on Cerebral Circulation, 3
Holden's Human Osteology, 1
——— Landmarks, 1
Horrocks' Bacteriological Examination of Water, 4
Hovell's Diseases of the Ear, 19
Hughes and Keith's Practical Anatomy, 2

Impey on Leprosy, 21
Ireland's Mental Affections of Children, 5

Jacobson's Operations of Surgery, 16
Jellett's Midwifery, 6
——— for Nurses, 6
——— Gynæcology, 6
Jessop's Ophthalmic Surgery and Medicine, 18
Johnson's (A. E.) Analyst's Companion, 26
Journal of Mental Science, 5

Kelynack's Pathologist's Handbook, 2
Keyes' Genito-Urinary Organs and Syphilis, 23

[*Continued on next page.*

Knuthsen on Hiccough, 15
Kohlrausch's Physical Measurements, 28

Lane's Rheumatic Diseases, 14
Lawrie on Chloroform, 17
Lazarus-Barlow's General Pathology, 2
———— Pathological Anatomy, 2
Lee's Microtomist's Vade-Mecum, 28
Leftwich on Syphonage in Large Intestine, 12
Lewis and Balfour's Public Health, 3
Lewis (Bevan) on the Human Brain, 2
Liebreich (O.) on Borax and Boracic Acid, 4
Liebreich's (R.) Atlas of Ophthalmoscopy, 19
Lucas's Practical Pharmacy, 8
Luke's Anæsthetics, 17

MacMunn's Clinical Chemistry of Urine, 23
Macnamara's Diseases and Refraction of the Eye, 18
———— Diseases of Bones and Joints, 17
McNeill's Isolation Hospitals, 4
Malcolm's Physiology of Death, 16
Marsh's Clinical Essays and Lectures, 17
Martin's Ambulance Lectures, 15
Maxwell's Terminologia Medica Polyglotta, 24
Maylard's Surgery of Alimentary Canal, 16
Mayne's Medical Vocabulary, 24
Microscopical Journal, 27
Mills and Rowan's Fuel and its Applications, 27
Moore's (N.) Pathological Anatomy of Diseases, 2
Moore's (Sir W. J.) Diseases of India, 10
———— Family Medicine, etc., for India, 10
Morris's Human Anatomy, 1
———— Anatomy of Joints, 2
Murray's Hare-Lip and Cleft Palate, 16

Nettleship's Diseases of the Eye, 18
Notter's Hygiene, 3

Oliver's Abdominal Tumours, 6
————Diseases of Women, 6
Ophthalmic (Royal London) Hospital Reports, 18
Ophthalmological Society's Transactions, 18
Ormerod's Diseases of the Nervous System, 14

Parkes' (L. C.) Elements of Health, 4
Parsons' Ophthalmic Optics, 18
Paton's (Noël) Physiology, 3
Pavy's Carbohydrates, 12
Pereira's Selecta è Prescriptis, 9
Pitt-Lewis's Insane and the Law, 5
Proctor's Practical Pharmacy, 8
Pye-Smith's Textbook of Medicine, 10
————Diseases of the Skin, 21

Ramsay's Elementary Systematic Chemistry, 26
————Inorganic Chemistry, 26
Richardson's Mechanical Dentistry, 20
Richmond on Antiseptics, 7
Roberts' (C. Hubert) Gynæcological Pathology, 6
Roberts' (D. Lloyd), Practice of Midwifery, 5
Robinson's (Tom) Eczema, 21
————Illustrations of Skin Diseases, 21
————Syphilis, 21
Ross's Diseases of the Nervous System, 14

St. Thomas's Hospital Reports, 11
Sansom's Valvular Disease of the Heart, 13
Savill's Clinical Medicine, 11
Schofield's Force of Mind, 5
Scott's Atlas of Urinary Deposits, 23
Shaw's Diseases of the Eye, 19
Shaw-Mackenzie on Maternal Syphilis, 24
Short Dictionary of Medical Terms, 24
Silk's Manual of Nitrous Oxide, 20
Smith's (Ernest), Dental Metallurgy, 20
———— (Eustace) Clinical Studies, 7
———— Disease in Children, 7
———— Wasting Diseases of Infants and Children, 7
Smith (F.) on Bullet Wounds, 16
———— (Fred. J.) Medical Jurisprudence, 3
Smith's (J. Greig) Abdominal Surgery, 16
Smith's (Priestley) Glaucoma, 19

[*Continued on next page.*

Snow's Cancers and the Cancer Process, 21
——— Palliative Treatment of Cancer, 21
——— Reappearance of Cancer, 21
Solly's Medical Climatology, 15
Southall's Organic Materia Medica, 7
Squire's (P.) Companion to the Pharmacopœia, 8
——— London Hospitals Pharmacopœias, 8
——— Methods and Formulæ 28
Starling's Elements of Human Physiology, 3
Sternberg's Bacteriology, 11
Stevenson and Murphy's Hygiene, 4
Sutton's (F.) Volumetric Analysis, 26
Swain's Surgical Emergencies, 15
Swayne's Obstetric Aphorisms, 5

Taylor's (A. S.) Medical Jurisprudence, 3
Taylor's (F.) Practice of Medicine, 10
Thin's Cancerous Affections of the Skin, 22
——— Pathology and Treatment of Ringworm, 22
——— Psilosis or "Sprue," 10
Thompson's (Sir H.) Calculous Diseases, 23
——— Diseases of the Urinary Organs, 23
——— Lithotomy and Lithotrity, 23
——— Stricture of the Urethra, 23
——— Tumours of the Bladder, 23
Thorne's Diseases of the Heart, 12
Thresh on Water Analysis, 4
Tilden's Chemistry, 25
Tirard's Medical Treatment, 12
Tobin's Synopsis of Surgery, 15
Tomes' (C. S.) Dental Anatomy, 20
——— (J. & C. S.) Dental Surgery, 20
Treves and Lang's German-English Dictionary, 24
Tuke's Dictionary of Psychological Medicine, 5
Turner on Sinuses of the Nose, 19
Tuson's Veterinary Pharmacopœia, 28

Valentin and Hodgkinson's Practical Chemistry, 26
Vintras on the Mineral Waters, etc., of France, 15

Wagner's Chemical Technology, 27
Wallace on Dental Caries, 20
Walsham and Spencer's Surgery, 15
Waring's Indian Bazaar Medicines, 9
——— Practical Therapeutics, 9
Watson (D. C.) on the Eye, 18
Watts' Organic Chemistry, 25
West's (S.) How to Examine the Chest, 11
White's (Hale) Materia Medica, Pharmacy, etc., 7
Wilks' Diseases of the Nervous System, 13
Wilson's (Sir E.) Anatomist's Vade-Mecum, 1
Wilson's (G.) Handbook of Hygiene, 3
Wright (H.) on Fevers of British Malaya, 10
——— Beri-beri, 10
Wynter and Wethered's Practical Pathology, 2

Year Book of Pharmacy, 9

*N.B.—J. & A. Churchill's larger Catalogue of about 600 works on Anatomy, Physiology, Hygiene, Midwifery, Materia Medica, Medicine, Surgery, Chemistry, Botany, etc. etc., with a complete Index to their Subjects, for easy reference, will be forwarded post free on application.*

AMERICA.—*J. & A. Churchill being in constant communication with various publishing houses in America are able to conduct negotiations favourable to English Authors.*

LONDON: 7, GREAT MARLBOROUGH STREET.