Delineations of cutaneous diseases : exhibiting the characteristic appearances of the principal genera and species comprised in the classification of the late Dr. Willan and completing the series of engravings begun by that author.

Contributors

Bateman, Thomas, 1778-1821. Willan, Robert, 1757-1812. University of Leeds. Library

Publication/Creation

London : Longman, Hurst, Rees, Orme, and Brown, 1817.

Persistent URL

https://wellcomecollection.org/works/jvkcay4e

Provider

Leeds University Archive

License and attribution

This material has been provided by This material has been provided by The University of Leeds Library. The original may be consulted at The University of Leeds Library. where the originals may be consulted.


This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org





















Digitized by the Internet Archive in 2015

https://archive.org/details/b28191262







DELINEATIONS

OF

CUTANEOUS DISEASES:

EXHIBITING THE CHARACTERISTIC APPEARANCES

OF THE

PRINCIPAL GENERA AND SPECIES

COMPRISED IN THE CLASSIFICATION OF THE LATE



DR WILLAN;

AND COMPLETING THE SERIES OF ENGRAVINGS BEGUN BY THAT AUTHOR.

BY THOMAS BATEMAN, M.D. F.L.S.

LIBRARIAN OF THE MEDICAL AND CHIRURGICAL SOCIETY OF LONDON, HONORARY MEMBER OF THE LITERARY AND PHILOSOPHICAL SOCIETY OF NEW YORK, AND PHYSICIAN TO THE PUBLIC DISPENSARY, AND TO THE LONDON HOUSE OF RECOVERY.

LONDON : PRINTED FOR LONGMAN, HURST, REES, ORME, AND BROWN, PATERNOSTER-ROW.

1817.

40

JUTANEOUS DISEASES

AND COMPLEXING. FILESERIES OF EX-MAVINES REGIN &V TH

603107

T. DAVISON, Lombard-street, Whitefriars, London.

objects of autoral history, namely, the facility of equicanting them by

In pathology, as well as in other departments of science, the advantages of methodical arrangement and a definite nomenclature, in aiding the progress of inquiry, have been generally acknowledged; but no branch of nosology appears to be so necessary or amenable to practical utility; as the classification of those diseases, which, being external, exhibit palpable differences of character, and are subject, like other objects of natural history, to the examination of our senses. From this circumstance alone, cutaneous diseases are, therefore, peculiarly capable of illustration by nosological arrangement; and, in fact, nothing but the want of such an arrangement can account for the general confusion of language and indiscriminate practice which have heretofore prevailed in this branch of medicine. Among the various attempts which have from time to time been made to class these diseases, the method proposed by Professor Plenck has alone any pretensions to accuracy. It is founded upon the proper basis of external character; and, indeed, appears to have been the model adopted by Dr. Willan for his own classification: but it was not followed out in its various parts with the same correct adherence to the fundamental principles, nor supported by the same extensive practical observation, the same perspicuity of diagnosis, and the same clear detail of symptoms belonging to each

genus and species. Hence it was not calculated, like the more correct system devised by Dr. Willan, to fulfil all the purposes of nosological investigation.

There is another circumstance, however, which renders the diseases of the skin equally capable of methodical classification with the other objects of natural history, namely, the facility of representing them by drawings. This was first attempted by Dr. Willan, and the engravings annexed to his work contributed greatly to the elucidation both of the general principles of the classification and nomenclature, and of the forms and progressive changes of various species of cutaneous disease. It was a source of considerable regret and disappointment, therefore, to the profession, that this work, and especially its excellent illustrations, should have been but half completed by its lamented author. Many intimations of this feeling reached me soon after the publication of my Synopsis, in consequence of its acknowledged deficiency in respect to the plates, which should have accompanied the work as completed by the hands of its author. The publication of the whole system in a compendious and accessible form has, however, probably been the means of diffusing a knowledge of it through a much wider field, and thus, I trust, of contributing to extend this part of pathology beyond the limits which it would have reached under the more costly form in which it originally appeared. For I have the satisfaction to state, that within the space of three years four large impressions of the Synopsis have been called for; and that it has been translated into the French and German languages, (into the latter under the direction and with the annotations of the learned Professor Sprengel,) and republished in America. In order, however, to fulfil the wishes of the profession by the completion, as far as it was in my power, of the series of engravings

iv

begun by Dr. Willan, I have purchased the copy-right of that work, together with the drawings and engravings procured by him, and have now brought that series to a conclusion.

It remains for me to say a few words respecting the mode in which this has been accomplished. The first four Orders, Papula, Squama, Exanthemata, and Bulla, have been illustrated by the reprinting of Dr. Willan's engravings, improved by Mr. Stewart, with the following exceptions. Plate XII, representing the Psoriasis gyrata, was re-engraved. In Plate XIII, which is new, a very accurate portrait of P. inveterata is substituted for Dr. Willan's confused and indistinct representation of that disease; and another variety of P. diffusa is superadded. Plates XVIII and XXX are also new: the former, exhibiting the disease Ichthyosis when it affects the face; and the latter, a variety of Purpura unnoticed by Dr. Willan. The second figure was added to Plate XXXII, from a drawing left by Dr. Willan, illustrating a species of Erythema, which he had omitted to represent. With Plate XXXIV the new series of engravings commences; and the superiority of execution by the able hand of Mr. Stewart will be very evident through the remainder of the series, with the exception of Plates XLVIII, LXIII, LXVII, and LXIX, which were already rather coarsely engraved by another artist for Dr. Willan. I think it but justice to Mr. Stewart to point out this circumstance. With the exception of the representations of Impetigo, Porrigo, Ecthyma cachecticum, Varicella (P. XLVI), Herpes præputialis, Rupia, Eczema impetiginodes, & mercuriale, the carbuncle of the plague, Vitiligo, Molluscum contagiosum, Sycosis barbæ, and Elephantiasis, all the new engravings were made from drawings found in Dr. Willan's collection, with their names written by his own hand, and obviously intended to illustrate his work, had he

lived to complete it. I stated, in the advertisement prefixed to the first Fasciculus of this work, some reasons for omitting the engravings representing those varieties of the different genera, which were reputed syphilitic. Further observation has confirmed the propriety of that omission: for the subject has been rather perplexed than elucidated by the most recent experiments; and it would be presumptuous, in the present state of our knowledge, to attempt to determine the character of the eruptions of true syphilis.

Т. В.

14, Bloomsbury-square, Oct. 1st, 1817.

3.VIII and XXX are shown in the tormer, evoluting the sistent foltheorie when a mirete the first; and the latter, a montr of Pagner atmosted by Dr. Wilton. The scient figure was wided to Flate XXXII; from a deriving tells by Dr. Wilton, illustrating a species of Drythewie which he had omitted to represent. With Flate XXXII that new serve of control on Market will be arresed and through the remainder of control on Market will be arresed and through ethics with the had omitted to represent. With Flate XXXII the remainder of the second, with the exception of the set of the remainder of the second, with the exception of the gravely of the remainder of the second, with the exception of the gravely of the remainder of the second, with the exception of the gravel by mother artic for Dr. Wiles. I thus a that parte as Mit. Typeschart to point out the mireurstande. Write the exception of the steriation of the second, the page. I thus a that parte as Mit. (P. XLVI), Hoyne preparates for Dr. Wiles. A thus a that parte as Mit. (P. XLVI), Hoyne preparates, the page. Finites (Heliarena contegiorum, fracting for the contents of the page. Finites (Heliarena contegiorum, diracting found in the Willing collection, with their names written by drawing found in the Willing collection, with their names written by drawing found in the Willing collection, with their names written by drawing found in the Willing collection, with their names written by INDEX.

soole month -

The first LZ.	Plate. Fig.	and the state of the second	Plate. Fig.
ACNE simplex	LXII.	Lichen simplex .	IV.
punctata		pilaris	V. 1
indurata	LXIII.	circumscriptus .	- 3
rosacea	LXIV.	agrius	IV. 2
Ecthyma vulgare .	XLIII. 1	lividus	V. 2
luridum .	2	Lupus	LXVII.
cachecticum .	XLIV. 1 & 2	Miliaria	LV. 1
Eczema solare	LVI.	Molluscum	LX. 1
	LV. 2	contagiosum .	LXI.
rubrum	LVII.	Nævus L	XX. LXXI-II.
mercuriale	LVIII.	Pityriasis capitis .	XV. 1
Elephantiasis	LXVIII.	versicolor .	2
Ephelis	LXIX.	Plague, Carbuncle of the .	LIX. 1-4
Erythema marginatum .	XXXII. 2	Pompholyx diutinus .	XXXIII. 1
papulatum .	XXXI. I	pruriginosus .	2
tuberculatum .	2	Porrigo larvalis	XXXVII.
nodosum .	XXXII. 1	furfurans	XXXVIII.
Herpes phlyctænodes .	XLIX.	scutulata	XXXIX.
zoster . ,	L.	decalvans	XL.
circinatus	LI. 1	favosa	XLI.
præputialis	- 2	faciei .	XLII.
iris	LII.	Prurigo mitis	VI. 1
Ichthyosis simplex .	XVI.	formicans .	- 2
cruris	XVII.	senilis	- 3
faciei	XVIII.	Psoriasis guttata	IX. 1
Impetigo figurata .	XXXIV.	diffusa	- 2
sparsa	XXXV.	, other varieties	X, XI, XIII. 1
scabida	XXXVI.	gyrata	XII.
Lepra vulgaris	VII.	inveterata	XIII. 2
	VIII. 1	palmaria	XIV.
nigricans	2	Purpura simplex	XXVIII. 1

1117

······· sine catatella

	N	D	16	х	
٠		**			

	Plate. Fig.	A CALL AND A	Plate. Fig.	
Purpura hæmorrhagica .	XXVIII. 2	Scabies purulenta . X	LVI. 1	
urticans	XXIX.	porcina	2	
senilis	XXX.	Scarlatina simplex . X	XII.	
Roseola æstiva	XXV. 1	anginosa . X	XXIII.	
autumnalis	2	Strophulus intertinctus . I.		
annulata	XXVI. 1		I.	
infantilis	2	confertus . Il	II. 1	
variolosa	XXVII. 1-2		- 2	
vaccina	3	candidus	- 3	
Rubeola vulgaris	XIX.	Sycosis menti I	LXV.	
	XX, XXI. 2	capillitii I	XVI.	
nigra	XXI. 1	Varicella XLVI	I, XLVIII.	
Rupia simplex	LIII.	Vitiligo I	X. 2	
prominens	LIV.	Urticaria febrilis X	XIV. 2	
Scabies lymphatica	XLV.	evanida	1	
A CONTRACT OF A				

viii

. .

ADVERTISEMENT

ADVERTISEMENT.

FROM the many urgent representations, that were made to me, while I was preparing my Synopsis of Cutaneous Diseases for the press, respecting the deficiencies of a mere descriptive treatise, unaccompanied by any graphical illustrations,—and from the obvious advantages which an ocular view of these external maladies affords, in conjunction with verbal description,—I am induced to suppose that a publication, like the present, will be acceptable to most of the readers of the Synopsis. With a view to their convenience, these plates will be published separately from the text of that work, of which they are intended as an illustration; a few words of description and diagnosis only being annexed, for the purpose of rendering the plates not altogether unintelligible alone; but which will add nothing to the *cost* of the work.

Upon this subject, I may observe, that it would have been more satisfactory to myself, if the Fasciculi could have been published at a lower price: but it would have ill answered the purpose, either of the editor or the purchasers, to sacrifice accuracy to cheapness; or to incur the risk of misleading, and being misled, by imperfect and ill-executed representations of the diseases attempted to be pourtrayed, merely to facilitate the sale of the work. I trust, however, that the appearance of a Fasciculus four times in the year, for three successive years, will not be deemed too frequent, nor prevent my professional brethren from taking a share in the necessary expenses of such a work.

ADVERTISEMENT.


The copper-plates, belonging to Dr. Willan's Treatise, having come into my possession, together with several drawings, prepared for the subsequent Orders, upon which he had not published any thing; and many original drawings, relative to the whole classification, having also been procured; I propose to publish twelve Fasciculi, containing six plates each, illustrative of all the principal genera and species, described in the Synopsis. The first four Orders, consisting of Pimples, Scales, Rashes, and Blebs or Blains, (Bullæ) will be illustrated chiefly by the republication of Dr. Willan's plates, improved at a considerable expense by the able hands of Mr. Stewart, and tinted with every possible care. The syphilitic eruptions, being a distinct class of diseases, and being, in fact, very insufficiently ascertained or characterized, will be, for the present, omitted; and a few additions, illustrative of other varieties, more immediately belonging to the arrangement, will be made. The last four Orders, comprising the Pustular, Vesicular, Tubercular, and Macular diseases, will be illustrated by a series of new engravings, executed from the original drawings above mentioned. This collection of drawings is nearly completed, and a few of them already engraved; so that punctuality in the completion of the publication may be confidently promised.

14, BLOOMSBURY SQUARE, 20th December, 1814.

T. B.

price : but it would have ill answered the purpose, either of the editor of the purchasers, to sacrifice accuracy to cheaparess ; or to incur the the mideading, and being misled, by imperfect and ill-executed representation of the diseases attempted to be pourtrayed, merely to facilitate the ; de the work. I trust, however, that the appearance of a Facticulas for the in the year, for three successive years, will not be deemed too freque is nor prevent my professional brethrea from taking a share in the necessive expenses of such a work.

ii





EXPLANATION OF THE PLATES.

THIS Fasciculus contains a representation of the principal species of the first Order of Cutaneous Eruptions, which are characterized by PAPULE or *Pimples**. The three genera of this Order are STROPHULUS, LICHEN, and PRURIGO.

PLATE I.

The STROPHULUS intertinctus; popularly termed the Red Gum, or Gown; a pimply eruption, of a vivid red colour, rising sensibly above the level of the skin, and intermixed often with dots and red patches, which have no elevation. It is peculiar to very young infants; and often consistent with good health.

* See Definition 1. Synopsis, p. xx.

EXPLANATION OF THE PLATES.

Tars Passionius contains a representation of the principal species of the first Order of Cataneous Eruptions, which are characterized by Parenes or Pimples *. The three genera of this Order are Stratence, Licensy, and Prentoo.

PLATE I.

The STROPRULT'S interfactus; popularly termed the Ref Curr, of Gome; a pimply cruption, of a vivid red colour, rishig sensibly above the level of the skin, and intermixed often with dots and fiel putckey, whele have no elevation. It is peculiar to very young infinite; and offen colour sistent with good bealth.

", and Definition 1. Synopsis, p. sx. 17







PLATE II.

The STROPHULUS albidus; which has been denominated by some writers the White Gum.⁷ It exhibits numerous, little, hard, whitish pimples, that remain a long time, on the face and neck, and, like the red gum, is frequently unaccompanied by any indisposition, but, like it, is occasionally the indication of disturbance in the alimentary canal.

Thurse call it pin Felon








PLATE III.

Fig. 1.—The STROPHULUS confertus; commonly called the rank Red Gum, and also, from its frequent occurrence in children a little elder, during the commencement of dentition, the Tooth-Rash. It is remarkable for the more extensive crop of papulæ, which are smaller, and more crowded, than in the other species; and, in children of seven or eight months, of a higher colour, exhibiting a general redness in each patch. It spreads along the fore-arm, cheeks, and forehead, which are its most common seats, as represented in the plate.

Fig. 2.—The STROPHULUS volaticus, or Feu volage; a less common eruption, which is distinguished by its small circular clusters of *papula*, each patch containing from three to ten, which, as well as the interstices between them, are of a high red colour. They rapidly appear in succession, turn brown, and exfoliate.

Fig. 3.—The STROPHULUS candidus; differing from the preceding in the larger size of the *papulæ*, which are smooth and shining, and without any surrounding redness, and stand at a considerable distance from each other. Fig. 1.— The Statemuture conjecture commonly called the rank field Gam, and also, from its finquent accurrence in children a field when for the more extensive cop of station, the Toste Kask. It is remarkable crowted, than in the after even of statistic, in children of seven or eight months, of a lighter colour, exhibiting a general reduces in each patch. It spreads along the foresum, circulate and to the date of the second spreads along the foresum, circulate and to the second of the second patch in the patch.

lie. 3.—The Seasenver er expensive e districts from the preceding in the larger size of the popular which are known and shinkle, and schoot are uncounting reduces, and shipt are considerable distance from such




Fig. 1. Lichen simplex V



Fig. 2. Lichen agrius. I

PLATE IV.


Comprises two species of LICHEN, a non-contagious, papular eruption, affecting adults.

Fig. 1.—The LICHEN simplex; which is characterized by a scattered eruption of distinct red *papulæ*, with a very moderate inflammation surrounding their base, terminating with a slight scurfy exfoliation of the cuticle. It is liable to be mistaken for the itch, the eruption of which, however, is usually less red, and more humid, being vesicular or pustular, but seldom purely papular. The distinct and distant papulæ distinguish it from the measles and scarlatina, for which it has been mistaken.

Fig. 2.—The LICHEN agrius; which is a more inflammatory form of the disease, and marked by the highly red hue of some of its patches, the skin of which becomes at length harsh, thickened, chapped, and excoriated. This tendency, and the diffuse redness, connecting the *papulæ*, distinguish this Lichen from the other species.

Fig. 1.—The Licenni simples ; which is characterized by a scattered impless of distinct red popole, with a very moderate information surrounding their base, terminating with a soluble scurfy extellation of the cuticle. It is liable to be mistaken for the itch, the emption of which, however, is usually less red, and more humid; beidg verticular or postular, bot seldom purely papelar. The distinct and distant papular distinguish is from the measles and scaription, for which it has been mistaken.

Fig. 2.—The Litence s agriant which is a more inflammatory farm of the disease, and numbed by the highly-red har of combod its patches, the skin of which becomes at length have, this leased, chapped, and excerimied. This tendency, and the diliuse reduces, connecting the popular, distinguish this fichen from the other species.







PLATE V.

Fig. 1.—The LICHEN *pilaris*; in which the *papulæ* chiefly appear at the roots of the cutaneous hairs.

Fig. 2.—LICHEN lividus; which is in fact a modification of Purpura, consisting of a sort of papular or elevated petechiæ.

Fig. 3.—The LICHEN circumscriptus; an annulated form of the eruption, less common than the two first, in which the *papulæ* are arranged in circular clusters, which have a well defined margin. They are apt to extend themselves by this papulated border, while the central surface becomes even, but continues slightly red and rough. This species was neither figured nor described in the first edition of Dr. Willan's work.






Fig ? Drurigo formicans.



Plate 17.


PLATE VI.


Contains the three species of PRURIGO, the only remaining disease of this Order, which is characterized by nearly colourless *papulæ*, with intense itching.


Fig. 1.—PRURIGO mitis; which is distinguished from LICHEN simplex by its soft and smooth papulæ, of nearly the same colour as the rest of the skin, and which seldom appear red, except when violently rubbed. As in the following species, minute black scabs become the most visible objects, which form upon them by the concretion of a little watery humour and blood, which ooze out, when their tops are abraded.

Fig. 2, and S.—PRURIGO formicans and senilis; which differ little from each other, or from the preceding species, except that the papulæ are larger in the last: these eruptions are more visible from the formation of the little dark scabs just mentioned, than from the appearance of the papulæ themselves, which are apt to escape observation from their hue. The combination of these colourless papulæ, with the little dark central specks upon those which have been abraded and inflamed, as is well represented in the plate, serve to determine the diagnosis, between this disease and the contagious itch.

Fig. 4.—Represents an insect, of which a great number were detected on the skin of an old man, affected with PRURIGO senilis, by Dr. Willan, who never met with a second instance of the same occurrence. Neither the disease nor the insect was communicated to the patient's wife, or to any of his family. It is obviously not a *pediculus*; but, both from the structure of its hind legs, and the rapidity of its jumping motion, it was deemed to belong to the genus *pulex*.







FASCICULUS II.

EXPLANATION OF THE PLATES.

THIS Fasciculus contains delineations of several species of the two principal genera of the second Order of Cutaneous Eruptions, which are distinguished by the formation of Scales*. There are four genera of scaly diseases, which are denominated LEPRA, PSORIASIS, PITYRIASIS, and ICHTHYOSIS.

PLATE VII.

LEPRA vulgaris, the common Leprosy; which is characterized, like all the species of this genus, by "scaly patches, of different sizes, having always nearly a circular form." It appears, first, in small, distinct, red, and shining elevations of the cuticle, on the centres of which thin white scales are speedily formed (as at A). In the course of three or four days, these spots are flattened and dilated, and become more scaly, equalling in size a silver penny (as at B); and they continue to enlarge, still retaining their circular form, to the size of a shilling or half a crown, or sometimes larger (as at D). Occasionally, though rarely, the scales accumulate into a thick crust (as at C). The patches are commonly surrounded by a red and slightly elevated border, which generally remains scaly after the central area has recovered its texture in the progress of cure. The leprous patches often appear first at the elbow, or immediately below the knee (as at E).

* See Definition 2. Synopsis, p. xx.

PASCICULUS II

EXPLANATION OF THE FLATES

I une Fasciculus contains delineations of several species of the two principal genera of the second Order of Cutaneous Eraptions, which are distinguished by the formation of Scales*. There are four genera of acaly diseases, which are denominated LEPHARTER FURALISTS, and ICHTUROSIS.

HV ATA19

Larra valoris, the common Leping; which is characterized, ille all the species of this genus, by " scaly patches, of different sizes, having airays pearly a circular form." It appears, first, in small, distinct, red, and shining clovations of the caticle, on the ceptres of which this white scales are specify fameed (as at A). In the course of three or four days, these spots are flattened and dilated, and become more scaly, equiling in size a silver penny (as at B); and they confinue to enlarge, still retaining their circular form, to the size of a shilling or half a crown, or concluses larger (as at B); and they confinue to enlarge, still retaining their circular form, to the size of a shilling or half a crown, or someticas larger (as at B). Occasionally, though rarely, the scyles acconsolate into a thick creat (as at C). The patches are commonly acrounded by a red and slightly clevated horder, which generally recussing scaly after the central area has recovered in texture in the acrounded by the kneet (as at C). The patches are commonly from a scale of the central area has recovered in texture in the acrounded by a red and sightly clevated horder, which generally recussing scaly after the central area has recovered in texture in the acrounded by a red and sightly clevated horder, which generally area instituted in the central area has recovered in texture in the acrounded by the kare (as at f).

· See Defaition C. Synophis p. zz.




Fig. 1. Lepra Alphoides. V 0 0 Fig. 2. Lepra nigricans. V

PLATE VIII.

Fig. 1.—LEPRA alphoides; which differs from the preceding form of the disease chiefly in the small size of the circular patches, which seldom extend beyond a few lines in diameter, and in the minuteness and greater whiteness of the scales.

Fig. 2.—LEPRA nigricans; which is a more rare form of the disease, differing from the L. vulgaris in the dark and livid colour of its patches, but resembling it in regard to its form and distribution.

PLATE VIII

the disease chiefly in the small size of the circular patches, which seldom extend beyond a few lines in diameter, and in the minuteness and greater whiteness of the scales.

big. 2.—Levn a nigricant y which is a more rate form of the discase, differing from the L. culgarie in the dark and livid colour of its patches, but reaembling it in regard to its form and distribution.






PLATE IX.


Fig. 1.—PSORIASIS guttata; a form of the scaly tetter most nearly allied to the Lepra, and characterized by small, distinct, patches of thin scales, with little or no inflammation round them, and without the regular circular figure, or the elevated border of Lepra. Their circumference is often angular, and sometimes slightly serpentine.

Fig. 2.—PSORIASIS diffusa; which consists of irregular, rough, and reddish patches, of various sizes, interspersed with chops and scales, the surface of which is exceedingly tender and irritable. In this variety, the patches are separate; but they are apt to expand till they become confluent.

PLATE IX.

Fig. 1.—FROMIASIS guillete: a form of the scale terter most meany allied to the Lepta, and characterized by small, distinct, patches of thin scales, with little or no influenzation round them, and without the regular circular figure, or the elevated border of Lepta. Their circumference is often angular, and sometimes sightly setpentine.

reddish patelos, of various sizes, interspersed with shops and scales, the stuface of which is exceedingly tender and imitable. In this variety, the pateles are reparate, but they are apt to expand till they become confluent.





Proviasis diffusa, peculiar to Washer Homen. V

Published April : 105 by Longmon Mart Res Orms & Brown

PLATE X.

Fig. 1.—PSORIASIS diffusa. Another specimen of the same species, in which the scales are more minute, and more readily exfoliate.

Fig. 2.—PSORIASIS diffusa; a variety of the same species, which occurs chiefly about the hands and arms of washerwomen, in consequence of the irritation of soap, and which is generally more highly inflamed, chopped, and tender, than any other variety of the disease. The scales separate in large irregular flakes.

X HLVIL

in which the scales me more minutes and more readily extended.

Fig. 2.—Prontacts diffuse; a variety of the same species, which occurs chiefly about the hands and aims of washerwomen, in concequencest the initation of soap, and which is generally some highly influence, chopped, and tender; than any solver unisty of the discuse. The colles separate in luge irrector flate.






PLATE XI.

PSORIASIS diffusa; a variety of the disease which often affects the back of the hand of bakers, and has been thence called the *Baker's itch*. It is accompanied with numerous chops and fissures about the knuckles, and the surface is a little tumefied, and the scaliness slight.

In his second edition of this plate, Dr. Willan had introduced a figure of the PSORIASIS *inveterata*; but the representation was so totally inadequate to convey any idea of the disease, that it has been deemed necessary to expunge it, and to restore the engraving to its original state. A new representation of this form of Psoriasis will be given in the next Fasciculus, Plate XIII.

PLATE XI.

back of the hand of baiters, and has been themer called the Rate's itch. It is accompanied with numerous chops and fissures about the Lanckles, and the surface is a built tunctied, and the scalarss sight.

In the Propriest of the of this plate, Dr. When had here been a figure of the Propriests invituants, but the representation was so totaly in adequate to convey any idea of the distance, that it has been deened necessary to expange it, and to restare has cograving to its original state. A new representation of this form of Provide's will be given in the treates to Passiculate, Plate XIII.






PLATE XII.

PSORIASIS gyrata; which is chiefly distinguished by the singular form and distribution of the scaly patches. These are tortuous and vermiform, or circular and semicircular with tortuous appendages, and are often arranged in a similar manner on each breast, or on each side of the spine. Slight scales are separated from the surface of the patches.

In consequence of the imperfection of the original drawing, this plate has been re-engraved. PLATE XE.

Provinsen and distribution of the achieventy distinguished by the singular form and distribution of the achieventian muchos. These are torizons and evernitisms or circular and semicircular with torizons appendances, and are often homored in a similar meaner on each breast, or an each side of the spine. Stickers are separated from the surface of the patcher.

consequences at the angenticeton of the original disk inst this white




Plate All Fig. 1. Derivasis diffusa. V Fig 2 Deriases inveterata & Deann by T.B. ent be at Stee

al July 2 2221 In Longman The st. Breach & Brown

FASCICULUS III.

EXPLANATION OF THE PLATES.

THIS Fasciculus comprises the remainder of the specimens of the second Order, or scaly diseases of the skin; namely, two additional varieties of PSORIASIS, and the principal modifications of PITYRIASIS and ICH-THYOSIS.

PLATE XIII.

Fig. 1.—PSORIASIS diffusa; a variety of the eruption, not figured by Dr. Willan, which occurs almost exclusively in young subjects, affecting the whole surface of the body, and producing a considerable degree of itching and irritation, but without any material derangement of the health. The scales and patches are both more numerous and more minute on the trunk than on the extremities. This drawing was taken from a boy about nine years of age.

Fig. 2.—PSOBIASIS inveterata; the most extensive and severe form of the disease, "is characterized by an almost universal scaliness, and a harsh, dry, thickened state of the skin:" it begins in separate patches on the extremities, and rapidly spreads over nearly the whole surface of the body. The scales are quickly produced and thrown off, and the skin is red, deeply furrowed, thickened, and rigid, so that motion is often painful and difficult.

EASOICULUS III.

.1

d non i associate comprises the remainder of the speciment of the specimen of the speciment of the speciment of the specimen

PLATE XIII.

Dr. Wahan, which occurs alarcar a variety of the croption, not referred by the whole surface of the body, and preducing a considerable degree of itching and irritation, but without any material deragements the dealth. The scales and patches are inthe user ounterous and can intain a boy about nice years of are

Fig. 2.-effort and attricted is the most estensive and rear form of the disease, " is chorecterized by an almost universal scalares, and a barsh, dry, thickened state of the skin ." it begins is separate patches on the extremities, and rapidly spreads over nearly the whole sudice of the body, The scale are quickly produced and thrown off, and the skin is rad, deeply furrowed, thickened, and right, so that motion is often painful and difficult







PLATE XIV.

PSORIASIS *palmaria*; an obstinate form of the *scaly tetter*, which is nearly confined to the palm of the hand, but sometimes appears slightly on the inside of the fingers and wrist. The cuticular furrows are deep, and even fissured at the bottom, so as to bleed when the fingers are strongly extended; and the whole surface of the diseased patch is rough, in consequence of the detached and elevated edges of the scaly cuticle. The complaint is troublesome from the heat, pain, itching, and stiffness which attend it.








PLATE XV.

Fig. 1.—PITYRIASIS capitis; which, when it occurs in young infants, as here represented, is called *the Dandriff*. On the upper part of the forehead and temples, it appears as a slight whitish scurf; but on the other parts of the head there are large, flat, semitransparent scales.

Fig. 2.—PITYRIASIS versicolor; a chequered discoloration of the skin, of a brown or yellowish hue, appearing chiefly about the breast, abdomen, shoulders, and upper arms. The discoloured patches are slightly rough with minute scurf or scales, which, however, are not always very conspicuous. Sometimes there is a slight redness in the patches, and then the eruption is accompanied by some degree of itching. Eig. 1.- Per ratatte copitie; which, when it occurs in young infants, as here represented; is colled the Dandriff. On the upper part of the forchead and complex, it appears as a shight whitish scurf; but on the other parts of the head there are large, flat, semistrumsparent scales.

of a brown or visitorish hue, oppraving chiefly about the breast, abdoupen, shoulders, and upper orms. The discoloured patches are slightly rough with minute sourt or scales, which, however, are not always very conspicuous. Sometimes there is a slight reduces in the patches, and then the emptime is necompanied by some degree of itching.








PLATE XVI.

ICHTHYOSIS simplex; the fish-skin disease, which is characterized by a permanently harsh, dry, rough, thickened, and slightly scaly texture of the integuments, generally of a brown or mud-colour. The scales are not deciduous, as in Lepra and Psoriasis.







Published Sulv Matas by Longman Maryt Reve Come & Br.

PLATE XVII.

ICHTHYOSIS cruris; a variety of the disease, which occurs in the lower extremities, in some cases of anasarca, forming a rugged coating, which ultimately cracks, and permits a discharge of the effused lymph through innumerable crevices.

PLATE XVII.

ICHTHYOS IS CRUTHY & VARIETY OF the disease, which occurs in the lower extremities, in some cases of nuesarea, forming a rugged conting, which a ultimately cracks, and permits e discharge of the effused lymph through innumerable crevices.







PLATE XVIII.

ICHTHYOSIS *faciei*. The face is often free from the disease, while the rest of the surface is affected with it; but in the case from which this drawing of the eruption was made, it was confined exclusively to the cheeks, the two patches uniting at one time across the nose. The head is imitated from one of M. Alibert's engravings.

PLATE XVIII

leurn vosts sacist. The face is often fee from the disease, while the fest of the surface is affected with it; but is the case from which this drawing of the eraption was made, it was conflued exclusively to the checks, the two patches pairing at one ting across the nose. The hand is imitated time one of M. Alibert's engravings.





FASCICULUS IV.

EXPLANATION OF THE PLATES.

THIS Fasciculus comprehends the three most important genera of the *third* Order of cutaneous diseases, the *Exanthemata*, or *Rashes**; namely, RUBEOLA, SCARLATINA, and URTICARIA.

PLATE XIX.

RUBEOLA vulgaris; the common form of measles, as it appears on the fifth day of the fever, and the second of the efflorescence, consisting of an intermixture of small, distinct, and nearly circular spots (A); and of little irregular patches, formed by the coalescence of these, which affect somewhat of a semicircular or crescent form (B), and are slightly raised. Upon the neck and arms a few miliary vesicles sometimes appear, during the height of the eruption, as at (C).

* See Definition 3. Synopsis, p. xx.

FASCIOULUS IV.

XPLANATION OF THE PLATES

The Pasciculus comprehends the three most important genera of the third Order of cutançous discasts, the Heanthemata, or Rashes"; unpely, literated a. Beautarrya, and Unreanta.

PLATE XIX.

Represent the first of the compon firm of measles, as it appears on the fifth day of the feren, and the second of the efforescence, consisting of m_1 internalstore of small, distinct, and nearly circular spots (A): and of little precular patches, formed by the contracence of these, which affect score what of a semicircular or crescent form (B), and are slightly maized. Upon the neck and arms a few milincy vesicles sometimes appears, during the bright of the cruption, as at (C).







PLATE XX.

RUBEOLA sine catarrho. This variety has been observed, during an epidemic season, in a few rare instances, in which the eruption of measles went through its stages, unaccompanied by any symptoms of fever, ophthalmia, or catarrh. It is so far important, however, that it does not secure the constitution from a subsequent attack of the febrile measles.

PLATE XX.

RUBROLA sife colurvio. This variety has been observed, during an epidemic season, in a few rare instances, in which the cruption of measles went through its stages, unaccompanied by any symptoms of fever, ophthalmia, or catarrh. It is so int important, however, that it does not secure the constitution from a subsequent attack of the febrile measles.



Rubeola nigra

Fig.1.



Appearance of the Rubeola vulgaris on its decline?

Pig. 2.

PLATE XXI.

Fig. 1.—RUBEOLA nigra. Dr. Willan gave this appellation to an unusual appearance, which he had sometimes seen about the seventh or eighth day of measles, the rash becoming suddenly black, or of a dark purple colour, with a mixture of yellow. It has continued ten days, or longer, without materially disturbing the functions of the patient.

Fig. 2.—Is a representation of the Rubeola vulgaris, about the eighth day, when it is declining, and assumes a slightly yellowish hue.

6

PLATE XXI

Fig. 1.— RUBEOLA migro. Dr. William gave this appellation to an unusual appearance, which he had sometimes seen about the seventh or eighth day of measles, the rash becoming suddenly black, or of a dark purple colour, with a mixture of yellow. It has continued ten days, or longer, without materially disturbing the functions of the patient.

Fig. 2.— Is a representation of the Rabcola valgars, about the eighth day, when it is declining, and assumes a slightly vellowish line.




Plate XXII. Fig. 2. TO SERVE AND DEVELOPED Flg. 5. COMPANY STREET, SALAR Scarlatina simplex?

brutt dell

PLATE XXII.


SCARLATINA simplex; the rash of scarlet fever, when it is not attended by ulceration of the throat. It is at first composed of innumerable red points (as at A, Fig. 2.), which on the following day coalesce, or are extended into an almost continuous efflorescence, especially over the cheeks and limbs; and on these parts, the miliary glands and papillæ of the skin are so much distended with blood, as to be raised into minute papulæ, which produce a roughness of the surface, like the cutis anserina (as at Fig. 1. B); and sometimes into miliary vesicles (as at Fig. 1. A). But there are commonly many interstices of skin-colour, of very irregular forms (as at C.). When the eruption begins to decline, the scarlet hue becomes less vivid, and the interstices are very numerous, as at the commencement (Fig. 2.).

Fig. 3. represents the speckled appearance, which often shews itself in a scurfiness (A), or in little patches of cuticle, which have sometimes been mistaken for vesicles (B).

PLATE XXII.

SCARLATINATINA Simpler; the rash of scarlet fever, when it is not attended by diceration of the threat. It is at first composed of innumerable red points (as at 1, Fig. 2.), which on the following day cosieses, or are extended, into an almost continuous efforescence, especially over the checks and limbs; and on these parts, the miliary glands and papille of the skin are so much distended, with bland, as to be mined into minute (as at Fig. 1 B), and admentines into military colours (as at Fig. 1, A). But there are commonly many intestices of skin-colour, of very irregular forms (as at C.) Wheat the comption begins to decline, the searlet here becomes test rink with the surregiver are reprint scarlet here becomes intuition the surregiver are reprint scarlet here becomes rest rink, and the surregiver here on a fig. 1, A).

Fig. 3. represents the spectical appearance, which often shows itself in a summers $(\Lambda)_{*}$ or in hith patches of calicle, which have sometimes been mistaken for evolvies (B).







PLATE XXIII.

Fig. 1. represents the inflamed and elongated papillæ of the tongue, in Scarlatina anginosa.

Fig. 2. The eruption of Scarlatina *anginosa* on the arm; shewing the goose-skin appearance from minute papulæ (as at A), and the general fulness of the cutaneous vessels (as at B).

Fig. 3. The rash of Scarlatina maligna, of a darker hue, and intermixed with Petechiæ.








Plate XXIV Fig. 1. Urticaria evanida Syd Edwards detin! Eng. 2. Urticaria Febrilis Instances for a state belongen and the stillene dance & Brann I Southiar Sculpt Indekahanika dal ?


PLATE XXIV.

Fig. 1.—URTICARIA evanida, a chronic form of nettle-rash, in which the wheals are not stationary, and are sometimes round, and sometimes oblong, like the strokes of a whip. They are never surrounded by an extensive blush, but sometimes slightly red round the base.

Fig. 2.—URTICARIA febrilis, a species of nettle-rash, which is preceded by some degree of fever, and in which the wheals appear in the midst of irregular patches of a vivid red hue, which fades during the day, but returns in the evening, with a slight febrile paroxysm. The patches are often a little elevated, and give some degree of tension and hardness to the skin, and terminate by slight exfoliations of the cuticle.







FASCICULUS V.


EXPLANATION OF THE PLATES.


THIS Fasciculus contains the fourth and fifth genera of the *third* Order of cutaneous diseases, or *Rashes*; namely, the principal varieties of ROSEOLA and PURPURA.

PLATE XXV.

Fig. 1.—ROSEOLA *astiva*: a rose-coloured rash, which chiefly occurs in summer, and is preceded by slight febrile indisposition for a few days. It differs from the Rubeola in the larger size, and more irregular forms and distribution of its patches; and from Scarlatina it is still more distinguished by the roseate hue, and by its mode of distribution.

Fig. 2.—ROSEOLA autumnalis; consists of distinct circular patches, of a damask hue, occurring principally on the arms of children from five to ten years of age, in autumn, and disappearing in about a week.








PLATE XXVI.

Fig. 1.—ROSEOLA annulata, appears in rose-coloured rings, with central areas of the usual colour of the skin. The rings are at first minute, but gradually dilate, in the order A, B, C, and E, sometimes to nearly half an inch in diameter.

Fig. 2.—ROSEOLA infantilis, is a closer rash than the R. æstiva, leaving very small interstices. It occurs in infants during any irritation occasioned by dentition, fevers, or bowel-complaints, and is very irregular in its duration, sometimes continuing but for a night, and sometimes appearing and disappearing for several successive days. It has been sometimes mistaken for measles and scarlatina. central areas of the usual colour of the skin. The rings, with minute, but gradually dilute, in the order M. D. C. and E. sometimes to ready half an inch in diameter.

prety anall interstees. It corears in fallout data the R. estivo, leaving rety anall interstees. It corears in fallouts during any irritation occaaimoed by dentitions ferens, or bowdecomplaints, and is very irregular in its duration, sometimes configuring the for a cight, and icenchines appending and disappearing far several successive risks. It has been








PLATE XXVII.

Fig. 1. and 2.—ROSEOLA variolosa; a rash which occurs during the eruptive fever of both the inoculated and the casual small-pox, first on the arms, breast, and face, and on the following day it extends over the trunk and extremities. In some cases it forms contiguous arches, (as in Fig. 1.) and in others longitudinal irregular patches, with small distinct dots intermixed (Fig. 2. B.)

Fig. 3.—ROSEOLA vaccina; an efflorescence which commonly appears in a congeries of dots and small patches, as here represented, but is sometimes diffuse, like the variolous rash. It usually occurs at the same time with the areola, and round the inoculated part.








PLATE XXVIII.

Fig. 1.—PURPURA simplex; the least complicated form of the "petechiæ sine febre" of authors, consisting only of minute effusions of dark blood under the cuticle, usually termed petechiæ.

Fig. 2.—PURPURA hæmorrhagica; a more severe modification of the disease, in which the effusions under the cuticle are more extensive, forming ecchymoses and vibices, and are accompanied by actual hæmorrhages from those parts where the cuticle is so delicate as to be easily ruptured, and therefore chiefly from the internal parts. The colour of the spots, in both varieties, is different at different periods of their continuance; being brighter on their first appearance, and becoming subsequently purple or livid, and lastly brownish or yellowish, when the effused blood is nearly absorbed.

PLATE XXVIII

Fig. 1.—Fourvas singler; the least complicated form of the " pelechim size fours" of authors, consisting only of minute effusions of dark "Mood under the cutiele, simily termed petechim.

Lig. 2.-- Ponevas hannevhasicat a more severe modification of the disease, in which the efficients under the cuticle are none extensive, forming evelopmoses and vibices, and are accompanied by actual harmorrhages from those parts where the cuticle is so delicate as to be easily ruprored; and therefore chiefly from the internal parts. The colour of the parts, in both unretice, is different at different periods of their continuance; being brighter on their first appearance, and becoming subservently purple of livid, and hertly brownish or y cllowish, when the efficience into a livid, and hertly brownish or y cllowish, when the






PLATE XXIX.


Fig. 1.—PURPURA urticans, is characterized by commencing with hard, reddish, and rounded elevations of the cuticle, resembling wheals, but not accompanied by any tingling or itching. These little tumours gradually dilate, but within a day or two they subside into livid spots of the same extent. On the extremities, where they most frequently appear, they are often mixed with petechiæ.

Fig. 2. and Fig. 3. are intended to shew the relative characters of petechiæ and flea-bites; the latter being distinguished by a central punctum, which remains under pressure, while the surrounding redness disappears.

PLATE XXIX.

Fig. 1.—Pregren anticana, is characterized by commenting with hard, reddish, and tounded elevations of the cuticle, resembling wheals, but not accompanied by any tangling or itching. These little camous gradually dilate, but within a day or two they subside into livid spots of the same extent. On the extremities, where they most frequently appear, they are often mixed with peterine.

Fig. 2. and Fig. 3. are intended to shew the relative characters of petechiae and flea-bites; the latter being distinguished by a central punctum, which remains under pressure, while the surrounding redness disappears.







PLATE XXX.

PURPURA senilis. This form of the Purpura was not noticed by Dr. Willan, nor in the two first editions of my synopsis. A few cases of it have occurred, with considerable uniformity of character, in elderly women. It appears principally along the outside of the fore-arm, in successive dark purple blotches, of an irregular form, and various magnitude. Each of these continues from a week to ten or twelve days, when the extravasated blood is absorbed. A constant series of these ecchymoses had appeared in one case during ten years, and in others for a considerable period; and in all, the skin of the arms was left of a brown colour. The health did not appear to suffer materially, as in the other forms of Purpura.

PLATE XXX.

Perrovs smills. This form of the Parpura was not noticed by Dr. Willan, nor in the two first editions of my synopsis. A few cases of it have occurred, with classiderable uniformity of character, in elderly women. It appears principally along the outside of the forcanta, in successive dark purple blotches, or an inregular form, and various magnitude. Each of these continues from a week to ten or twelve days, when the extra stried blood is accurate daring ton years, and in others for a considerable period; and to de the string ton years, and in others for a considerable period; and to de the string ton years, and in others brown colour. The health fid not uppear to suffer materially, as in the other forms of Pirroura.



rg.1 Crythema papulatum? Plate XXXI. e like Fig. 2.

Crythema tuberculatum?

Strutt Adars!

J. Seillier Sculpt

Published And said belongman Marst Bees Orme & Brown.

FASCICULUS VI.

EXPLANATION OF THE PLATES.

IN this Fasciculus, the *third* Order of cutaneous eruptions is completed by the representation of the varieties of ERYTHEMA; the *fourth* Order, *Bulla*, is illustrated by two varieties of POMPHOLYX; and the *fifth* Order, *Pustula*, is commenced with the three species of IMPETIGO.

PLATE XXXI.

Fig. 1.—ERYTHEMA papulatum. An eruption, chiefly affecting the arms, neck, and breast, in extensive irregular patches, of a bright red hue; but which consists for the first two days of imperfect papulæ, which disappear as the efflorescence brightens. The skin continues red and furrowed for ten days or a fortnight; and a blueish or livid hue succeeds the decline of the rash.

Fig. 2.—ERYTHEMA tuberculatum. This eruption somewhat resembles the former in its irregular and bright patches, in which, however, there are dispersed small round tumours, slightly elevated, but not so hard as tubercles, which subside in about a week. The erythematous patches turn livid about a week afterwards, and gradually disappear.

FASCICULUS VI.

EXPLANATION OF THE PLATES

In this Fasciculus, the stird Order of cutancous cruptions is completed by the representation of the varieties of Envruences; the fourth Order, Bulle, is illustrated by two varieties of Pournourz; and the fifth Order, Pastale, is commenced with the three species of Invertion.

.FLATE XXXI.

Fig. 1.—Earrurun papulatum. An eruption, chicky affecting the arms, neck, and breast, in extensive irregular patches, of a bright red intes; but which consists for the first two days of imperfect papular, which despipers as the efforence or pightens. The skin continues red and furrowed for ten days or a fortnight; and a bluch or livid bue succeeds the decline of the risk.

Fig. 2.— Larren a. a takerentatean. This cruption source hat reachebles the sourcer, is its urreation and height purcises, in which, however, there are dispensed much runnet turn his, dightly clevated, but not so hard as takendes, which satiside in clout a week. The crythenatous polycluss turn livid about a week after wards, and gradually disappear.




Plate XXXII. Crythema nodosum! Crythema marginatum I Joelliar Scalp!

PLATE XXXII.

Fig. 1.— ERYTHEMA nodosum. This rash usually affects the fore part of the legs, and seems to occur only in women. Many of the patches are of an oval form, the central parts of which are very gradually elevated into hard and painful protuberances on the 6th or 7th day, from which time to the 10th they constantly soften and subside, the red colour turning blueish or livid on the 8th or 9th day.

Fig. 2.—ERYTHEMA marginatum. This eruption was not delineated in Dr. Willan's work; but the little figure, here substituted for an obscure representation of a supposed syphilitic erythema, was found among his drawings. The erythema marginatum occurs in large patches, which are bounded on one side by a hard, elevated, tortuous, red border, in some places obscurely papulated; but have no regular margin on the open side. The duration of the disease is variable, from three to six weeks.

PLATE XXXII.

Fig. 1.— Ex vrn zur nodecum. This rash usually affects the fore part of the legs, and seems to occur oily in women. Many of the patches are of an oval form, the central parts of which are very graduelly devated into hard and painful protuberances on the 6th or 7th day, from which time to the 10th they constantly soften and subside, the red colour turning blassish or hvid on the 6th or 9th day.

Fig. 9.— Eurrunas surginatum. This cruption was not defineated in Dr. Willan's work : but the fittle figure, here substituted for an obsected representation of a supposed symplotic certificans, was found among his drawings. The crytheum margination preuze in large patches, which are bounded on one side by a lard, elevated, justwods, red border, in some places obsecuely papainted: but have no regular margin on the open side. The duration of the discuse is variable, from three to six weeks.




Fig. L. Plate XXXIII. Dompholyx diutinus. D Е Fig. 2 Dompholyx pruriginosus. 童子 thed April 1355 by Longman Burd Res. Orme & Bree W.Dorton delis!

PLATE XXXIII.

It was scarcely necessary to figure so common a disease as Erysipelas; and this representation of POMPHOLYX, therefore, constitutes the only illustration of the *fourth* Order of Cutaneous Diseases, BULLE, * Blebs, or Blains.

Fig. 1.—POMPHOLYX diutinus. "An eruption of bulla, without any inflammation around them, and without fever." The vesications seem to commence from red tingling elevations of the cuticle, nearly resembling papulæ (as at A.); these rise into transparent vesicles (as at B.); which become yellow, and enlarge (as at C and D.); break and inflame (as at E); and at length form a scab on the excoriated surface (as at S.).

Fig. 2.—Represents an appearance of the Pompholyx, which is sometimes connected with the Prurigo formicans.

* See Definition 4. Synopsis, p. xxi.

PLATE XXXIII.

It was scarcely necessary to figure so common a disease as Erysipelas; and this representation of Pournoirs, therefore, constitutes the only illustriction of the fourth Order of Cutanoous Daeases, Burns, Blebs, or Blaim.

B.): which become yellow, and colarge (as at C and D.); hreak and B.): which become yellow, and colarge (as at C and D.); hreak and B.): which become yellow, and colarge (as at C and D.); hreak and inflame (as at E); and at length form a scab on the exterioristic surface (as at S.);

Fig. 9.2 Represents an appearance of the Pompholys, which is some-

ice & See Delinition 4. Symposic, 5. 321.






PLATE XXXIV.

With this Plate the new series of engravings commences, and also the *fifth* Order of Cutaneous Diseases, the PUSTULES, *Pustulæ*; on which subject Dr. Willan had given nothing to the public, his treatise having extended no farther than the fourth Order.

IMPETIGO figurata, the first species of the pustular, or humid tetter, is characterized by circumscribed patches, a little elevated, which at first consist of clusters of psydracious pustules, * closely congregated, and surrounded by a slight inflammatory margin; but after some days, the pustules break, and an irritating humour exudes, which soon concretes into yellowish or greenish-brown scabs. These patches are often situated on the arm and wrist, and extend to the back of the hand, and one of them not unfrequently takes its station between the metacarpal bones of the thumb and forefinger, as here represented.

* See Definition 5, b. Synopsis, p. xxi.

PLATE XXXIV.

With this Plate the new series of moravings commences, and also the fifth Order of Cutaneous Discastes, the Persturss, Pastule; on which subject Dr. Willan had given nothing to the public, his treatise having 'extended as father than the fourth Gener.

In retro four a the first spaces of the postular, or humid tetter, is characterized by commercial pairine, a fittle elevated, which at first counist of chasters of paymanna purtules, * closely congregated, and surrounded by a might influence or margin ; but after some days, the pustices into a distance influence or scale. There patches are often cretes into a dise are and wrist and retread to the back of the hand, and situated on the are and wrist and retread to the back of the hand, and one of the mark and universe the set of the back of the hand, and bones of the think and toronge, as has a set of the set of the back of the hand.

is a support of the stand of the second






PLATE XXXV.

IMPETIGO sparsa; which differs from the preceding species rather in the indeterminate and extensive distribution of the pustules and scabs, than in its general character. It is commonly, however, a more severe disease, and connected with more constitutional derangement than the I. figurata. In the lower extremities, especially, it is often extremely obstinate and severe, continuing for many weeks, and even passing into the next species.

APLATE XXXV.

A second second state of the second s







PLATE XXXVI.

IMPETIGO scabida; which is distinguished by the formation of a thick, yellowish, scabby crust, by which the whole limb or limbs are encased, and which is often separated by many fissures and cracks, from which a thin ichor copiously exudes, and concretes into additional and irregular layers. The whole crust, indeed, is the result of the concretion of this acrimonious humour, which is discharged in great abundance from numerous psydracious pustules, as they successively form, break, and ulcerate over the surface of the limb.

PLATE XXXVI.

Invertee cables, which is distinguished by the formation of a thick, yellowish, scabby creat, by which the whole limb or limbs are encased, and which is often separated by many fissures and cracks, from which a thin icher copionary exceles, and concretes into additional and irreguiar layers. The whole crust, indeed, is the result of the concretion of this acrimonious humour, which is discharged in great abundance from numerous psychology pustoles, as they successively form, break, and ulcerate over the suffice of the limb.





FASCICULUS VII.

EXPLANATION OF THE PLATES.

THIS Fasciculus comprises six specimens of the formidable and unsightly genus PORRIGO, the second in the Order of *Pustules*.

PLATE XXXVII.

PORRIGO larvalis, the Crusta lactea of medical authors, is almost exclusively a disease of infancy, occurring during the period of lactation; whence its popular name. It is commonly confined at first to the forehead and cheeks, but usually extends to the scalp and ears, and sometimes to the breast and extremities. It consists of an eruption of numerous, minute, and whitish pustules, denominated achores,* which are crowded together upon a red surface, upon which, as they break and discharge, thin yellowish or brownish-green scabs are formed by the concretion of the viscid humour. The whole face is thus often enveloped as it were by a mask, whence it has acquired the epithet larvalis. Notwithstanding the extent to which the infant is disfigured, no permanent scar or deformity ever ensues.

* See Definition 5, c. Synopsis, p. xxii.

PASCICULUS VII.

XPLANATION OF THE PLATES.

1 mis l'ascir alus comprases six que ontone of the identidable and ordightly is

PLATE ANAVIL

Pormioo ferrafis, the Crassi hecked of medical makters, is almost exclusively a disease of infance, transmag during the period of lactation; whence its popular sime. It is comparing contined at first to and similarity to the trenut and extremities. It consists of an exaction of aumerous, remute, and which quantities. It consists of an exaction are convict repeters upon a ned surface, upon which, as they back and discharge, this reflects to income it is autoe, upon which, as they back the concetant of the result frame it. Yin make the scale back and discharge, this reflects to income it is a sequired to entry the the concetant of the result frame it. Yin make face is thus then the concetant of the result frame it. The make face is thus then the concetant of the result frame it. The make face is thus then the concetant is write for a next, whence it has acquired the pathet for each discharge, the next is a next, whence it has acquired the pathet for the concetant for a first frame it is a set in the set of the result in the concetant is distanting the extent to a first the infant is disting the permanent year of defining the extent to a first the infant is disting the permanent year of defining the extent to a first the infant is disting the permanent year of defining the extent to a first the infant is disting to the

* men Definition 5, c. Stanues, p. 190







PLATE XXXVIII.

PORRIGO *furfurans*. Instead of concreting into thick scabs, the humour, which exudes in very small quantity, dries speedily, and separates in thin laminated scabs, or scale-like exfoliations. The scalp is usually affected extensively with the furfuraceous porrigo, and becomes tender and sore. The disease is commonly limited to the scalp, but occasionally extends to the external ears, and a little on the forehead and temples, beyond the roots of the hair, as in the case from which this drawing was made. It occurs principally in adults, or in persons attaining to the period of puberty.









PLATE XXXIX.

PORRIGO scutulata; popularly known by the name of Ringworm; an unmanageable and very contagious variety of the disease, which, though as old as the Greeks, has recently been propagated extensively, by the multiplication of boarding-schools and manufactories, in this country. It commences with clusters of small achores, or pimple-like pustules, in distinct small patches, of an irregularly circular figure; which are sometimes limited to one scutulum, and sometimes numerous, and distant. Though the exudation is commonly small, yet, if neglected, it will concrete into thin scabs: the areas of the patches also dilate, and become confluent; and the hair upon them becomes lighter in colour, and at length is destroyed at the roots, leaving patches of baldness.









PLATE XL.

PORRIGO decalvans; a variety, which is also called ringworm, and which may sometimes be the result of the decalvating process of the preceding species. In general, however, there is no manifest appearance of disease in the patches of this species, except complete baldness, of a very white and smooth appearance, and in form more or less circular, while the hair surrounding the patches is as thick as usual.








PLATE XLI.

PORRIGO favosa, the scalled head, consists of an eruption of the large, soft, straw-coloured pustules, denominated Favi.* This Scall, however, may occur on all parts of the body, and not unfrequently affects the face and limbs, and even the trunk. The scab which results from these pustules, wherever they appear, is very characteristic; being generally soft, yellowish, or greenish, elevated, and semi-transparent, with a very irregular surface, indented into hollows; whence the term honey-comb scab has been applied to it. This is particularly conspicuous when it occurs on the face. Considerable ulceration often takes place under these scabs, especially when they are much irritated by picking and scratching them about the edges; and the neighbouring glands usually exhibit marks of irritation.

* See Definition 5, d. Synopsis, p. xxii.

Possano area an and and a second and a second and a second and an area and a second a se






PLATE XLII.

PORRIGO favosa affecting the face. As in this situation the scab, which succeeds the favous pustules, is conspicuously characterized, I deemed it right to give this view of it, in preference to figuring the Porrigo lupinosa, which indeed is of such rare occurrence, that I have not been able to procure a good drawing of it since the commencement of this undertaking. Notwithstanding the extreme deformity occasioned by this scall, it often yields readily (as in the case from which this drawing was taken) to medical treatment, and leaves no scar behind; a circumstance which renders it important to distinguish it from the varieties of Lupus, Scrofula, and other scabby ulcerations.

PLATE XLII

Poneration from a starting the face. As in this stimuted the state of a starting the stimuted the state of a start of a s




Plate XLIII.



FASCICULUS VIII.

EXPLANATION OF THE PLATES.

IN this Fasciculus the fifth Order of cutaneous diseases, PUSTULE, is completed by the representation of four specimens of ECTHYMA, and three of SCABIES; it being deemed unnecessary to figure the varieties of that well known eruption, VARIOLA, of which I am in possession of several excellent drawings, procured by Dr. Willan. The two remaining plates of this Fasciculus exhibit the varieties of the first genus of the sixth Order, which is characterized by the formation of Vesicles,* namely, VARICELLA.

PLATE XLIII.

Fig. 1.—ECTHYMA vulgare: an eruption of the large pustules, denominated phlyzacia, † occurring principally on some part of the extremities, and gradually enlarging and inflaming, for a week or ten days, one after another, when they break, pour out their pus, and afterwards a thinner fluid, which concretes into brownish scabs. In about a week more, the inflammation subsides, and the scabs soon afterwards fall off. The various stages of the pustules are here delineated, in a case of some severity. In the milder cases, they often do not exceed in size the pustule near the wrist.

Fig. 2.—ECTHYMA luridum; is characterized by the darker hue of the bases of the phlyzacious pustules, and by the long and slow succession in which they spread over both the trunk and limbs, for a period of several weeks. When they break, a curdly or sanious discharge issues, and dark hard scabs form by its concretion, and remain surrounded by dark inflamed borders till they are about to separate.

These two figures are from Dr. Willan's collection of drawings.

* See Definition 6. Synopsis, p. xxii. † Definition 5, a. ibid. p. xxi.

PASCICULUS VIII.


Is this Fasciculus the 15th Order of cutanious diseases, Provence, is domplicitie in the representation of four speciments of F. rurats, and there; of Stantas représentation of four speciments of F. rurats, and efficient well known eropaine. Namericas of which I am in possession of sevents exercitant dravinges provinced by Dr. William. The two remaining plates of this Fastinguins calable the two surficients of the first genus of the states rate of this fasting also adding the section of the first genus of the states of the fasting also adding the the formation of fermation of the states of the fasting also an observation of the first genus of the states of the fasting also adding the the first fermation of fermation of the states of the fasting also adding the fasting the fermation of fermation of the states of the fasting and the states of the first genus of the states of the fasting also adding the first fermation of fermation of the states of the fasting also adding the first fermation of fermation of the first the states of the fasting also adding the first fermation of the fermation the states of the fasting also adding the first fermation of the first genus of

PLATE XILIL. PLATE XEUL

Fig. 1.—Ectury a suggerer an emption of the large pustiles, denomilitiged "Elfernie if a coopriner an emption of the large pustile, extroministed physicalitatic conversing architikally on some work on the days, unities and prisedra it here is a coopriner and originally on some work on the days, mutical and prisedra it here they threak open control was controller exacts at the date of addition of the set of the transformer and the origination of the set of the set of the set of the transformer at the date of the set is because and the set of the set of the set of the set is because at the interval of the set of the set of the set is because at the interval of the set of the set of the set is the set of the set is because at ages and the set of the set of the set of the set is set of the mission of the set of the set of the set of the set of the mission of the set of the set of the set of the pustole near the wrist.

Lig. 2.— Ecriptic 1 invidum; is characterized by the darker has of the bases of the phlysacsous postales, and by the long and alow praced sion in which they spread over both the trunk and hatso terms derived of severall weeks in When they orches, an cardly for similar to dark a cases, and dark indich scale reaches this test concretant to a reaches sole rounded by digk inflament backers will the same whose us darks a sole

r nese two ngures are from ils. William's collection of drawings





Ecthyma cachecticum.

PLATE XLIV.

ECTHYMA cachecticum. This species of Ecthyma was not noticed in the first and second editions of my Synopsis, whence it may be necessary to notice it more at large here.

These two figures present specimens of the two most common forms of an extensive eruption of phlyzacious pustules, which not unfrequently occurs, in connection with a state of cachexia, resembling that which is produced by the operation of a morbid poison in the habit, and is often treated as syphilitic. It is usually preceded for two or three days by a febrile paroxysm, after which numerous pustules, with a hard inflamed base, arise in daily succession, going through their successive stages of inflammation, suppuration, scabbing, and desquamation, so that, in the progress of the disease, they are commonly seen under all these conditions at the same time, as represented in the plate. The pustules which occupy the breast and abdomen are generally less prominent, and contain less matter than those on the face and arms.

The febrile symptoms are diminished, but not removed, on the appearance of the eruption; for a constant erethism or hectic continues during the progress of the disease, with great languor and depression of strength and spirits. It is also accompanied by pains in the head and limbs, and commonly by some degree of ophthalmia, and by a slow inflammation, with superficial ulceration, of the fauces.

The duration of these complaints is considerable, extending to eight, ten, or twelve weeks, but yielding at length to the use of sarsaparilla, and cinchona, with the assistance of alteratives. F PLATE XERV

by a febrile parorysin, where which connercts pratifies, with a land inmestales which occupy the breast and abdousen are generally less pro-







PLATE XLV.

SCABIES lymphatica. It is difficult to give a distinct representation of the almost invisible vesicles of scabies; and this engraving expresses rather too strongly the slight inflammatory redness, which accompanies them, having exceeded the drawing in that respect: but it exhibits the distribution of the disease, in a well marked case, pretty distinctly.








PLATE XLVI.

Fig. 1.—Scables purulenta; or pocky itch. The size, prominence, and purulence of the eruption in this form of itch, which is here faithfully represented, frequently mislead the observer, who allows this severe disease to go on for a considerable time without applying the specific, to which it speedily yields. Where the pustules coalesce, they form irregular blotches, which sometimes ulcerate to some extent, and afterwards form hard and dry scabs.

Fig. 2.—SCABIES porcina. It is well known that a severe scabies is sometimes excited by the contact of dogs, cats, pigs, &c. affected with the mange. The vesicles and pustules are generally numerous and rank, and the general surface of the skin becomes rough, and of a dirty or brownish hue. The case, from which this drawing was made, occurred in a man who had attended a mangy hog.

IV.IX MYA.M

Fig. 1.—Starts paralents to pocky itch. The time, prominence, and puralence of the complete in this form of itch, which is here, kathfully, represented, frequently, mishead the direct cours he is college this servere disease to go on for a considerable there without applying the specific, to which it specifily yields. Whete the pustales copies of the form integular blotches, which sometimes derive advantation applying the afterwards first interland dry scales and means of a spine exactly and afterwards first interland dry scales and means of a spine exactly and

Fig. 2.—Sommers percing. It is well known that a screw scables is sometimes excited by the doubtedt of deply's core thint. Not affected of the the mange. The residence of the other phase of the screw fig. a decrement and the and the general surface of the shire being pressed by and of a party brownish have. There are been shire before any single and of a derive of in a man when has half attended any approximation and of a derive of in a man when has half attended any approximation and a second of a course of the screw board of any approximation and a second in a man when half attended any approximation and and of a derive of a course of the screw board of any approximation and a start of the screw by the screw and the screw bar and the screw bar and a start of the screw board of a start of the screw bar and a start of the screw bar and a start of the screw bar and the screw bar and a start of the screw bar and a start of the screw bar and the screw bar and a start of the screw bar and a start of the screw bar and the screw bar and a start of the screw bar and the screw bar and a start of the screw bar and the screw bar and a start of the screw bar and a start of the screw bar and a start of the screw bar and the screw bar and a start of the screw bar and the screw bar and a start of the screw bar and the screw bar and a start of the screw bar and the screw bar and the screw bar and a start of the screw bar and a start of the screw bar and the screw bar and the screw bar and start a screw bar and the screw bar and the







PLATE XLVII.

The first genus of the numerous Order of Vesicles is the VARICELLA, Chicken Pox, or Swine Pox.

The three varieties, which the Varicella assumes, have been introduced into this plate. The vesicles on the upper arm are the common *lenticular* eruption, which is peculiarly called *Chicken-pox*; those on the forearm are the more acuminated *conoidal* vesicles, which have received the appellation of *Swine-pox*; and the hand and wrist exhibit the *globated* vesicles, which, in the north, have been distinguished by the name of *Hives*. The intermixture of rising vesicles, with those that are puckered and subsiding, and others that are drying into scabs, on the fifth or sixth day, as here represented, constitutes a principal point in the diagnosis between this eruption and small pox.

TILATE XLVII

The first yeans of the numerous Order of Fedeles is the Vancri, Chicken For, or Same For. The three varieties, which the Varietiz assumes, have been introduced into this plate. The vesicles of the apper arm are the the teamon kentcalar emption, which is preatinely called Cheken-par; those on the forearm are the more accumulated consulat reaches, which have genered the uppellation of Scher-par; and the hered and which they globated vesicles, which he the north, have been distinctioned by the ratio of the effect. The intermittane of the seconds, which they globated state subsiding; and others that are diving into each, on the foresial between this emprised the seconds, with these that are previous of more subsiding; and others that are diving into each, on the fifth or sials between this emprised and small point in the diag.








PLATE XLVIII.

This plate, from a drawing of Dr. Willan's, represents a full eruption of the lenticular varicella, with some intermixture of conoidal vesicles. The little circumscribed brown scabs, here and there formed, while the rest of the eruption is full, mark the character of the disease, together with the form of the vesicles, although they have assumed a yellowish, or almost purulent appearance.

PLATE XIVIII.

The lasticular varicella, with some intermixture of conoidal vesicles. of the lasticular varicella, with some intermixture of conoidal vesicles. The little circumscribed brown scales, here and there formed, while the sint of the eraption is fall, mark the character of the disease, rogether with the form of the scales, although they fare usuand a yellowish, or shock purariest appearance.





FASCICULUS IX.

EXPLANATION OF THE PLATES.

THIS Fasciculus exhibits the principal species of two genera of the Order Vesiculæ; namely, HERPES and RUPIA.

PLATE XLIX.

HERPES phlyctænodes. In this form of Herpes, the vesicles are usually smaller than in the Shingles, and the successive clusters extend upon the limb longitudinally, and do not encircle it as the latter eruption surrounds the trunk. The progress of the patches downwards is marked in the plate by the incrustation of the upper one, while the smaller vesicles of the lower one are forming.

FASCICULUS IX.

EXPLANATION OF THE PLATES

I ars Fasciculus exhibits the principal species of two genera of the Order Fericular, mimely, Harris and Rurra.

PLATE NILY.

HERE'S physicanodes. In this form of Herpes, the vesicles are usually smaller than in the Shingles, and the succe swe clusters extend upon the limb longitudinally, and do not encircle it as the latter cruption surrously the trunk. The progress of the patches downwards is marked in the plate by the incrustration of the upper one, while the smaller vesicles of the lower one are forming.







PLATE L.

HERPES zoster ; commonly termed the Shingles ; a species of Herpes peculiar to the trunk of the body, and generally spreading half round, by a succession of irregular clusters of the vesicles. As the patches, which first appeared, subside, the vesicles become partially confluent, and assume a livid or blackish hue, and terminate in thin dark scabs. This complaint is generally of little importance, but is occasionally accompanied, especially on the decline of the eruption, by an intense deep-seated pain in the chest, which is not easily allayed by medicine. HEREES zoster ; commonly tarmed the Shingles ; a species of Herpes peculiar to the trunk of the body, and rememby spreading half round, by a succession of irregular charters of the resides. As the patches, which first appeared, subside, the vesicles become partially confident, and assume a hold or blackish here and terminate in thin dark scale, This confplant is generally of fittle importances but is occasionally accompanied, especially on the decline of the cruption, by an intense deep-seated jana at the chest, which is and cerains allayed by medicine.






PLATE LI.


Fig. 1.—HERPES circinatus. The plate represents one of the most extensive varieties of this species of Herpes. The patches are often much smaller, consisting of little rings of vesicles, not much larger than a shilling, and accompanied by very slight redness. In children it is called the *ring-worm*, and sometimes mistaken for the porriginous disease.

Fig. 2 and 3.—Two specimens of the HERPES praputialis, a form of the disease, which, from its situation on the prepuce, is liable to be confounded by the patients themselves, and by ignorant practitioners, with chancre. The drawings represent the progress of the eruption, which generally consists of a single patch, in two cases, and the numbers mark the days on which the drawings were made; namely, in Fig. 2, on the second, fifth, and eighth; and in Fig. 3, on the second, third, fifth, and sixth days.

ILI STALLA

Fig. 1.—If rar as circinates. The plate represents one of the most extensive varieties of this species of Herpes. The patches are often much smaller, consisting of little rings of vesicles, not much larger than a shilling, and accompanied by very slight redness. In children it is called the ring-norm, and sometimes mistaken for the porriginaus disease.

Fig. 2 and 3.—Two specimens of the Huurus preparation, a form of the disease, which, from its situation on the prepuce, is huble to be canfounded by the patients themselves, and hy ignorant, practificners, with chancre. The drawings represent the progress of the emption, which generally consists of a single patch, in two crees, and the anumbers mark the days on which the drawings were muther manuely, in Fig. 2, on the second, fifth, and eighth ; and m Fig. 3, on the second, third, fifth, and







PLATE LII.

HERPES Iris. This rare and curious variety of Herpes occurs principally in the situation in which it is here figured, but sometimes in the palms of the hands, or on the instep. The vesicles are not large or elevated, but sufficiently distinct to characterize the disease. They are disposed in two or three concentric rings, with a central vesicle, which arise in succession, as in the other forms of Herpes.

PLATE MIL

HEREES In This mee and carines variety of Helpes occurs principally in the signatum in which it is here heared, but constitues in the palms of the hands, or on the instep. The concless are not large or elevated, but sufficiently distinct to connected at the discase. They are disposed in two or three concentric rings, with a central vesicle, which arise in succession, as in the other forms of Herpes.

1







PLATE LIII.

RUPIA simplex. All the forms of this eruption are connected with a cachectic state of the constitution, and some of them, like the one here figured, are accompanied by symptoms resembling those produced by a morbific poison. It will be seen that they differ from the Ecthymata in the form and size of the eruption, and in the colour and consistence of the contained fluid, consisting in fact of flattened muddy vesicles, and forming a larger and more circular scab.

HILL GUALDS

Iterra simpler. All the forms of this eruption are connected with a cachectic state of the constitution, and some of them, like the one here figured, are accompanied by a meltoms reambling those produced by a morbilit preserve. It will be seen that they differ from the Eethynanta in the form and size of the croption, and is the colour and consistence of the contained find, consisting in fact of insteaded middly vesicles, and forming a farger and more consistence.








PLATE LIV.

RUPIA prominens. I am indebted to the liberality of a professional friend, distinguished by the union of graphic with anatomical science, for this excellent representation of this form of Rupia, which greatly exceeds in the expression of character the drawings which I had previously procured. It exhibits very distinctly the original vesicular form of the eruption, and the limpet-shell appearance of the scab.







FASCICULUS X.

EXPLANATION OF THE PLATES.

IN this Fasciculus the remaining genera of the Order VESICULE are comprehended; namely, MILIARIA, and ECZEMA: and as a substitute for the genus APHTHA, which required no graphical illustration, I have been enabled to figure the vesicle or carbuncle of the Plague, which I trust will be deemed an acquisition. The last Plate commences the seventh Order of *Tubercles*, including specimens of MOLLUSCUM and VITILIGO.

PLATE LV.

Fig. 1.—MILIARIA. The miliary eruption, when it affects the extremities, as here represented, is less copious than on the back, neck, and breast, where it is liable to occur only as a symptom of fevers, in which much heat of the skin and sweating have been accidentally excited.

Fig. 2.—ECZEMA impetiginodes. This local variety of Eczema chiefly affects the back of the hands, about the knuckles, or in the situation here figured. After a succession of rising vesicles, an inflamed and irritable state of the skin is induced, forming a patch of an irregular circular form, which is hot, itches, and is disposed to crack, but on which neither scabs nor pustules form, as in Impetigo.

PASCIOULUS X.

EXPLANATION OF THE PLAN

Is the formation of the reaction formation of the Order Ventering and comprehensive many property of the second formation of the substitute for the second many of the second of the second of the Player, which have been many is found to accord on the back of the Player, which I trust will "many and an acquisition. The last Plate commences the second Order in Tubersies including second of Motorsector of First 199.

PLATE LV.

Erg. 1.—MITELADIA. The miliary cemptron, when it afford the entremities; as here represented, is less copions than on the mech, nock and breast, where it is liable to occur only as a symptom of trues, in which much heat of the skin and sweating have been accidentally excited

Fig. 2.—Eczena imperiements. The head variety of bereine chiefly effects the back of the hands, about the hunchles or in the silter tion here figured. After a succession of rising vesicles, an inflated and irritable state of the ishin is induced, forming a patch of as foregilar circular form, which is hot, steles, and is disposed to travely, but on which acadier wakes nor postules form, as is imperiate.






PLATE LVI.

ECZEMA solare. This form of the disease may arise from various sources of irritation, but is most common in summer, being the result of the direct irritation of the sun or heated air, and hence occurs on parts most exposed to their influence, more particularly on the back of the hands and fingers, as here represented.

PLATE LVL

Excession where the form of the domast may arise from various sources of irrutation but is most common in summer, being the result of the direct irritation of the ann or heated his, and hence occurs on parts most exposed to their infinences more particularly on the back of the heads and intervents of here represented.






PLATE LVII.

ECZEMA rubrum. This form of Eczema varies somewhat in different cases; but always appears under the form of minute vesicles, accompanied by a diffuse redness, which the roughness and elevation of the vesicles distinguish from simple Erythema. This disease has been observed to follow exposure to cold, and to recur in the same individual at irregular intervals without any manifest cause.

PLATE EVIL

Ecara a rubram. This form of Eczema varies somewhat in different cases ; but nivers appears under the form of minute vesicles, accompanied by a diffuse reduces, which the roughness and dicration of the vesicles distinguish from simple Ergthema. This disease has been observed to follow exposure to cold; and to recut in the same individual at irregular intervals without any manifest cause.







PLATE LVIII.

ÉCZEMA rubrum mercuriale. The most remarkable and common form of Eczema rubrum is that which results, in some constitutions, from the irritation of mercury. The hue of this eruption is commonly less florid than that of the former variety, or of a browner tint; though we have seen it highly florid in one or two instances. The distribution and extent vary infinitely in different cases, and the specimen here figured can give but a general notion of either.

PLATE LIVE

Energy relevant and an exclusion of the constrable and common form of Ecterns and units and which results in some constitutions, from the instation of meetrics. The bins of this compton is commonly less therid that that that do the torrest variable of the compton is commonly less have seen it inguly though is one or two instances. The distribution and extent vary infinitely in different cores, and the specimen here beared can are but a research instance of ether.








PLATE LIX.

Vesicle and Carbuncle of the Plague. The liberality of Dr. Calvert, physician to the Army at Malta, enables me to give a representation of this pestilential eruption, which, as its first appearance is in the form of a vesicle, not much unlike that of the cow-pock, I place here. (See Fig. 1.) The progress of the vesicle to the tumid and fiery ulcer of the carbuncle is delineated by the pencil of Dr. Calvert in four successive stages.









Molluscum pendulum?

PLATE LX.

Fig. 1. and 2. represent the decline of the eruption of VITILIGO, in which the elevations appear to have subsided nearly to the level of the surface, chequering it in a singular manner. I have seen little of this rare disease, and am unacquainted with the whole progress of the case from which this drawing was accurately made by Mr. H. Thomson.

Fig. 3.—MOLLUSCUM pendulum. This drawing was procured by Dr. Willan (a small portion of it only being here figured) from a case which occurred under his own observation. Some of the tubercles were considerably larger than those included in the present figure; yet, like the poor man of Muhlberg, whose portrait was published by Prof. Tilesius, he appeared to suffer neither indisposition nor material inconvenience.







FASCICULUS XI.

EXPLANATION OF THE PLATES.

In this Fasciculus the seventh Order of *Tubercles* is continued. It comprises, besides an undescribed species of MOLLUSCUM, all the species of the common tubercular eruption, ACNE, and of SYCOSIS.

PLATE LXI.

MOLLUSCUM contagiosum. This singular eruption had not been noticed by Dr. Willan, and was unknown to myself till after the publication of two editions of my Synopsis; it may, therefore, be proper to give a fuller account of it than is usual in this work. The peculiar circumstances which distinguish this species of Molluscum from the common excrescences of that name are its contagious quality, and the emission of a milky fluid from an imperceptible aperture in the apex of most of the larger tubercles, upon pressure; by means of which fluid it is probable that the disease is inoculated. The young woman, whose face and neck were disfigured in the manner represented in this plate, had received the eruption from a child whom she nursed, on whose cheek a large tubercle of the same kind existed, who had doubtless been infected by a former nurse, on whose face the same eruption had been observed, and who had communicated it also to two other children in the family. I have subsequently seen this eruption in another child, who had apparently been infected by an older child, who was in the habit of nursing it, and on whose face the same tubercles, with the milky fluid, previously appeared. The tubercles are hard, smooth, and nearly of the colour of the skin, but with a shining surface, and a slight appearance of transparency: they are generally of a globular form, sometimes ovate and sessile upon a contracted base. Their progress is not very rapid; but some of them, on attaining a considerable size, proceed to a slow and curdly suppuration; and, in the case here figured, the cervical glands, under those on the neck, were swollen, and the surface was somewhat discoloured.


FASOICULUS XI

STITE AND AD ADDRENTER

Is this Forthating the strength Order of Yolf release confinently. It comprises here the an undescribed species of Mon russian, all the species of the common tubercular cruption. Access and a Sycons.

PA ATE EXT

Morrisers a sectorious. This sincular emption had not been noticed by Draw the public not very mpid ; but some of them, on attaining a considerable size, pro-






PLATE LXII.

ACNE simplex; ACNE punctata. These two species of Acne so constantly occur together, as in the case here engraved from a drawing of Dr. Willan's, that it was not deemed necessary to figure them separately. The black *puncta*, indeed, sometimes appear on the top of inflamed tubercles, which they have excited, being, in fact, little plugs of hardened mucus or sebaceous matter, irritating and over-distending the little ducts of the sebaceous glands, in which they are lodged. In general, however, these *puncta* are surrounded only by a very slight raised border of cuticle.

PLATE EXH.

A constantly occur tegether, in in the case here engineed from a denving of it: Without, that it was not decoust meressary to fruce them sepatately. Without, that it was not decoust meressary to fruce them sepatately. The battly gassin, indeed, sumetimes appear on the top of inhamed fairendes, which they linke excited, bring in fact, little plugs of hardened much or reburbous matter, instative and over-distending also tittle dueth of the schaceous glands in which they up lodged. In startic however, these panets are surrounded only by query slight matrix there of our here.






PLATE LXIII.

ACNE indurata. This accurate representation of the more violent and indurated species of Acne was procured and already engraved for Dr. Willan's work. The various forms, acuminated, conical, or oblong and conoidal, which the tubercles assume, are depicted in the plate, as well as the livid hue which they put on in their progress, especially those which do not proceed to suppuration.

PLATE LXIII.

Areas indurates (Inis air unate representation of the more violent and innumied sporte of Arm was produced and already engraved for Dr. Willian's vork. The various forms, accombated, conical, or objerty and conoidal, which the tubercles assume, are depicted in the plate, as well as the limit has which they put on in their progress, especially thoses which do not proceed to supply their on the second.






PLATE LXIV.

ACNE rosacea. This species is somewhat different from the preceding, as it is accompanied with general redness, which commences on and spreads from the end of the nose, on both sides, to the cheeks. In the progress of the disease, the cuticle becomes thicker, and its surface uneven, and variegated by reticulations of enlarged cutaneous veins, with smaller red lines, sometimes stretching across the cheeks.

PLATE LXIV.

Acre a remoter. This apecies is somewhat different from the preceding, as it is accompanied with general redness, which commences on and spreads thin, the end of the nose, on both sides, to the checks. In the progress of the disease, the outicle becomes thicker, and its surface uneven, and variegated by redealstons of calarged cataneous venue, with smaller red lime, sometimes strefching across the checks.

.






PLATE LXV.

Sycosis menti. The tubercles in this eruption, which almost exclusively affects men, usually appear first on the under lip, or prominent part of the chin, in an irregular cluster: but other clusters and single tubercles soon follow, often covering nearly the whole surface on which the beard grows, up to the ears, and under the jaw towards the neck. The tubercles are red and smooth, and nearly equal to a pea in magnitude, but of a conoidal form. As they suppurate slowly and partially, and discharge a small quantity of thick matter, by which the hairs of the unshaven beard become matted together, that appearance like the pulp of a fig, which has given origin to the name of the disease, and which is here represented on the chin, is often produced.

PLATE LXV.

Sroosis meant. The tubercles in this cruption, which almost exclusively affects men, isually appear first on the under lip, or prominent part of the chin, in an inegular cluster: but other clusters and single tabercles soon follow, often covering nearly the whole surface on which the beard grows, up to the cars, and under the jaw towards the neck. The tabercles are red and smooth, and nearly equal to a pea in magnitude, but of a conoidal form. As they suppurate slowly and partially, and discharge a small quantity of thick matter, by which the like the pulp of a fig, which has given origin to the mane of the discuse, and which is here represented on the course in the produced.







PLATE LXVI.

Success capillitii. This variety of Sycosis, affecting the hairy scalp, is seated chiefly about the temples, near the external ear, forehead, and occiput, near the margin of the hair. The tubercles arise in clusters like the former, but are softer and more acuminated than those on the chin, and pass more rapidly into suppuration, soon becoming confluent, and producing an elevated, unequal, and ulcerated surface. This is also sometimes granulated, affording some resemblance to the inside pulp of a fig.

PLATE LXVL

Sreests cajulatif. This variety of Specsus adocting the harry scalp, is seated chiefly about the temples, near the external car, forehead, and occiput, near the margin of the bair. The tubercles arise in clusters like the former, but are somer and more neuminated than those on the chin, and pass more impidir into suppurations some becominage confident, and producing an elevated, intequal, and uccrated surface. This is also sometimes granulated, affording some rescublance to the inside pdip of





FASCICULUS XII.

EXPLANATION OF THE PLATES.

IN this concluding Fasciculus are exhibited two formidable diseases of the *tubercular* Order, LUPUS, and ELEPHANTIASIS; and specimens of the principal varieties of the genera EPHELIS and NEVUS, which constitute the eighth Order, *Macula*.

PLATE LXVII.

LUPUS, noli me tangere. This representation of the extensive ulceration of the face, occasioned by this severe and unmanageable disease, was procured and already engraved at the instance of Dr. Willan. It produces even more deformity than Elephantiasis, with which it is sometimes confounded; but from which it differs essentially in its character and progress. It has been sometimes also deemed cancerous; and the circumstance of an occasional cure having been effected by the use of arsenic has probably given rise to the opinion that that medicine has proved a remedy for cancer.

Is this concluding Franculus are eslibited two foundable or costs of the reference Order. Lurges, and Excensions and specimics of the mineipal varieties of the genera Econoris and Surves, which rou-

rtitute the cighth Orlor, Macuke.

PLATE INVIL

Lorves, nois we tanger: This representation of the target to the target of target of the target of the target of the target of t






PLATE LXVIII.

ELEPHANTIASIS. This hideous disease is fortunately never seen in this country, except in a few individuals, natives or residents of more southern climates. This was the first case that had occurred to my notice, and I am indebted to the kindness and liberality of my friend, Mr. Lawrence, for the opportunity of procuring this excellent representation of the disease, as it appeared in the boy, Charles Uncle, a patient in St. Bartholomew's Hospital, whose case has been detailed by Mr. Lawrence, in the sixth volume of the Medico-Chirurgical Transactions, for the year 1815. The engraving here given is an exact portrait of that individual, after the tubercles had been rendered a little smoother than they originally appeared by the application of poultices.

PLATE LXVIII.


Research the set of the fide one disease is boronally never terms and the courts, creept is a greater the first case that half occurred to mansouth methanes. This was the first case that half occurred to my built, and I are articleted to the leadness and liberaity of my friend, when is and I are articleted to the leadness and liberaity of my friend, are another a statement of the leadness and liberaity of my friend, are another a statement of the leadness and liberaity of my friend, are another a statement of the leadness and liberaity of my friend, are another a statement of the superset in the boy. Charles Use, a parent is statement in the article and the boy. Charles I age, are been a statement of the case the subset of the boy for the boy are been as the term the state of the superset in the boy. Charles I age, are been and the boy of the case the subset of the boy for the been state of the tract of the term the case the subset of the boy for the been readered a little





PLATE LXIX.

EPHELIS. This representation of the large dusky patches, which differ from freckles only in their extent, and in occurring also on parts of the skin not exposed to the direct influence of the sun, was already engraved for Dr. Willan's work. These discolorations approximate in their character to the Pityriasis versicolor, but are more diffuse, and generally of a darker brown hue, and never shew any tendency to the minute furfuraceous exfoliations, which often accompany that affection.






Plate TXX Naivus foliaceus

PLATE LXX.

N \pm vus maternus; mother-spots, &c. This and the two following plates exhibit some of the varieties of those discolorations and excrescences of the skin, with which different individuals are marked from their birth. In consequence of the absurd popular notion, that these congenital alterations of the texture of the skin originate from impressions of various external objects on the senses and imagination of the mother during gestation, they have received various appellations, corresponding with the objects which they have been supposed to resemble. The specimens here figured were among the drawings procured by Dr. Willan, with the names here annexed to them. The two on this plate, from their resemblance to a leaf and a cherry, were called N \pm vus foliaceus, and cerasus.

two on this plate, how their resublicary to a by and a clearly, ware called Navra foline as and echoner.




Fig. 2.

Navus Cerasus.



Fig.1.

Navus.

Fig.3.



Navus Arancus.

Engthy J. Dewart

PLATE LXXI.

FIG. 1. This is one of the most common forms of the N_Evus, and is sometimes compared to the stain of port-wine, or claret.

FIG. 2. NÆVUS araneus, being supposed to bear the likeness of a spider.








PLATE LXXII.

This drawing exhibits an unusual number and variety of congenital marks, all occurring in the same individual.

