Fentazin : chiff chaff.

Contributors

Allen & Hanburys

Publication/Creation

London : Allen & Hanburys, 1980.

Persistent URL

https://wellcomecollection.org/works/ecgukgx9

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



Fentazin

(perphenazine BP)

A potent tranquilliser to control agitation, anxiety and tension, and for the treatment of nausea and vomiting.

PRESENTATIONS AND BASIC NHS COSTS

Fintaxin Tablets 2mg, 4mg and 8mg each contain perphenazine BP 2mg, 4mg and 8mg, respectively. In cartons of 100 in 10 foil strips of 10 (5 x 2) tablets. Fentaxin Tablets 2mg – £2:17, Fentazin Tablets 4mg – £2:57, Fentazin Tablets 8mg – £3:22, Fentaxin Injection contains perphenazine BP 5mg in each ampoule of 1ml. Boxes of 5 and 100 ampoules, Basic NHS cost of 5 ampoules 55p.

INDICATIONS

INDICATIONS For severe anxiety and tension states, particularly where agitation is present; anxiety associated with psychosomatic disorders, compulsive or obsessive disorders; psychotic states characterised by CNS hyperactivity, including schizophrenia of various types and senile psychoses; behaviour disorders; nausea and vomiting; and as adjunctive therapy

Some payon as the fail of the second second

DOSAGE AND ADMINISTRATION

The usual oral dosage is 4mg three times daily and this may be adjusted according to response but the total should not exceed 24mg daily. Fentazin Injection is given intramus-cularly in a dose of 5 to 10mg, which may be followed by further doses of 5mg at six hourly intervale. intervals. Children: Not suitable for children under 14 years of age.

CONTRA-INDICATIONS, WARNINGS, ETC.

Fentazin should not be given to patients with leucopenia or with drugs liable to cause bone-marrow depression.

Use instruct expression: It should be used with caution in patients with congestive heart failure or coronary artery disease. Unnecessary administration of drugs during the first trimester of pregnancy is undesirable.

Side effects include extrapyramidal symptoms such as dystonia, torsion spasm and pseudo-Parkinsonism. Persistent oral dyskinesia has been reported occasionally, particu-larly in elderly female patients, after long-term treatment with potent phenothiazine drugs including perphenazine.

SELECTED BIRDS OF BRITAIN

CHIFFCHAFF

Phylloscopus collybita

The Chiffchaff comes to this country from Africa in the early spring and returns south during October. Although it may be found in most parts of Britain it is rare in Scotland. It favours areas containing trees or shrubs, including large gardens. Insects such as midges may be taken on the wing or snatched from leaves. The Chiffchaff is very similar in appearance to the Willow Warbler but may be distinguished by its song, a repetitive 'chiff-chaff' interspersed with a low churring note. The call note is a high 'ho-eet'. During courtship there is a slow display in flight.

The female builds a domed nest of grass, moss and leaves lined with feathers and soft grass. It is usually sited in a low bush or in undergrowth. 5-7 white eggs, finely speckled with purplish-brown, are laid. The sexes are alike in appearance; they reach 41 inches (11 cm) in length.

(Illustration from "A History of British Birds" by F. O. Morris)

LEGAL CATEGORY

80/5/HN

POM

PRODUCT LICENCE NUMBERS Fentazin Tablets 2 mg 0045/5082; Fentazin Tablets 4 mg 0045/5083; Fentazin Tablets 8 mg 0045/5084. Fentazin Injection 5 mg 0045/5055.

> Further information is available on request. Fentazin is a trade mark of ALLEN & HANBURYS LTD, LONDON E2 6LA

