

Propaderm topical steroid therapy in eczemas and dermatoses : Montagu's harrier.

Contributors

Allen & Hanburys

Publication/Creation

London : Allen & Hanburys, 1978.

Persistent URL

<https://wellcomecollection.org/works/sr9k9p4q>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



MONTAGU'S HARRIER.

Propaderm

(Beclomethasone dipropionate BP for use on the skin)

TOPICAL STEROID THERAPY IN ECZEMAS AND DERMATOSES

PRESENTATIONS

Propaderm Cream – preferred for weeping conditions or where the skin is moist or hairy.

Propaderm Ointment – suitable for dry, lichenified or scaly conditions or for use under occlusive dressings.

Propaderm Lotion – primarily intended for lesions involving large areas, skin folds or hairy sites and, for cosmetic reasons, may be preferred for the face or other exposed areas.

AND FOR WHEN INFECTION IS PRESENT OR IS LIKELY TO OCCUR

Propaderm-A Ointment – also containing Chlortetracycline Hydrochloride BP 3.0%.

Propaderm-C Cream & Ointment – also containing Clioquinol BP 3.0%. Particularly suitable where infection is likely to be of fungal origin.

PACKAGE QUANTITIES AND BASIC NHS COST

Propaderm Cream	15g tube	41p	50g tube	£1.06
Propaderm Ointment	15g tube	41p	50g tube	£1.06
Propaderm Lotion	15ml bottle	40p	50ml bottle	£1.06
Propaderm-A Ointment	15g tube	46p	50g tube	£1.15
Propaderm-C Cream	15g tube	44p		
Propaderm-C Ointment	15g tube	44p		

SELECTED BIRDS OF BRITAIN

MONTAGU'S HARRIER

Circus Pygargus

A summer visitor from Africa, found in marshes, heaths and sand dunes in southern England. One of the rarest British breeding species. Keeping close to the ground, it searches for and then pounces on small animals and birds. Also eats eggs of other birds. Silent, except on the breeding ground and possibly at communal roosts. Then there is an interchange of high-pitched, staccato calls. The female builds a nest on the turf of marshes or heaths, using weeds and coarse grasses, lined with finer grass. Lays four or five eggs, white or pale blue (occasionally marked by sparse red-brown spots or streaks). The print illustrates the male. The female is larger, 19 in (48 cm) long, and dark brown with paler streaks on head and neck, and a barred tail.

(Illustration from "A History of British Birds" by F. O. Morris)

SELECTED BIBLIOGRAPHY

- Ashurst, P. J. et al (1970) *Brit. J. clin. Pract.*, **24**, 45.
Caldwell, I. W. et al (1968) *Brit. J. Derm.*, **79**, 487.
Fine, S. et al (1968) *Practitioner*, **201**, 362.
Raffle, E. J. and Frain-Bell, W. (1967) *Brit. J. Derm.*, **79**, 487.
Raffle, E. J. and Frain-Bell, W. (1968) *Brit. J. Derm.*, **80**, 124.

Further information is available on request.

Propaderm is a trade mark of
ALLEN & HANBURY LTD LONDON E2 6LA

