

Sednine : as an example of English pottery there is no improving on this Wedgewood vase and stopper ...

Contributors

Allen & Hanburys

Publication/Creation

London : Allen & Hanburys, 1966.

Persistent URL

<https://wellcomecollection.org/works/tsryzewa>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



Photograph by courtesy of the Victoria and Albert Museum

As an example of English pottery there is no improving on this Wedgwood vase and stopper (circa 1870) in jasper ware. The applied medallions are after the antique and show : Apollo with Citharoedus, Silenus with panther, Apollo and Marsegas, and Ceres with Cupid . . .

. . . and as a cough suppressant there is no improving on Pholcodine ; but the addition of Pseudoephedrine provides the advantage of easier breathing, making

SEDNINE an ideal treatment for all forms of cough.

To the Medical Profession

SEDNINE

a significant advance in the relief of cough
suppresses useless cough
makes productive coughing easier
and
facilitates breathing

for effective control of cough

SEDNINE

combines Pholcodine—the reliable anti-tussive, non-constipating, and safe for adult and child
with

Pseudoephedrine — the effective decongestant and bronchodilator less likely to produce
cardiovascular or CNS side effects than ephedrine.

SEDNINE

for all forms of troublesome cough, either productive or non-productive. Especially indicated where cough is associated with bronchospasm or upper respiratory congestion.

SEDNINE

a significant advance in the relief of cough

DOSAGE

Adults: two teaspoonfuls (10ml) three or four times a day as required.

Children: 6 to 12 years, half to one teaspoonful (2.5 to 5ml) three or four times a day as required.

CAUTION

Pseudoephedrine in clinical doses has very little pressor activity in normotensive persons, but minor side effects similar to those commonly encountered with ephedrine may occur if the recommended dosage is exceeded. Similarly, pseudoephedrine should not be given to patients suffering from advanced hypertension or heart failure.

FORMULA

Each adult dose (10 ml) of Sednine contains:

Pholcodine BP 15 mg

Pseudoephedrine hydrochloride 60 mg

PRESENTATION

Sednine is supplied in bottles of 142 ml (5 fl oz) and 2 litres.

BASIC N.H.S. COST

142 ml 4/8d

SEDNINE

is a Trade Mark of Allen and Hanburys Ltd



Manufactured in England by
Allen & Hanburys Limited London E2

B2 66/531/HN

PRINTED IN ENGLAND