

Sulpenin V widens the antibacterial spectrum of Penicillin V.

Contributors

Allen & Hanburys

Publication/Creation

London : Allen & Hanburys, 1958.

Persistent URL

<https://wellcomecollection.org/works/em65e6yy>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

● BLOTTER

*widens the
antibacterial spectrum
of Penicillin V*

SULPENIN V



Each tablet contains:-

PENICILLIN V 90 mg.
SULPHADIMIDINE 0.5 gramme

Presentation — Tubes of 10 and bottles of 100 tablets

MANUFACTURED IN ENGLAND BY ALLEN & HANBURY LTD LONDON E2

To the Medical Profession

RECOMMENDED DOSAGE

Sulpenin V should be given at intervals of 6 hours

AGE	INITIAL DOSE	MAINTENANCE DOSE
	No. of tablets to be given	No. of tablets to be given
Adult	4	2
11-15 years	3	1
4-10	2	1
Up to 3 years	1-1½	½-1

NOTE

It is recommended the tablets be swallowed with a copious drink of fluid, and adequate fluid intake be maintained during treatment.

IN ACUTE INFECTIONS

In acute meningococcal, streptococcal and pneumococcal infections when high blood concentrations of both penicillin and sulphonamide are necessary, the initial dose may be increased to 6 tablets followed by 2 or 3 tablets every 6 hours.

K58/460/HN Printed in England