

Trandate tablets : sandpiper.

Contributors

Allen & Hanburys

Publication/Creation

London : Allen & Hanburys, 1981.

Persistent URL

<https://wellcomecollection.org/works/rm6sj455>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).

**wellcome
collection**

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



SANDPIPER.

Trandate

(labetalol hydrochloride)

TABLETS

For the management of all grades of hypertension

PRESENTATIONS AND BASIC NHS COSTS

Trandate Tablets 100mg

Film-coated biconvex tablets, each containing labetalol hydrochloride 100mg, marked TRANDATE 100AH on one face. Containers of 50 £4.54 and 250 £21.65.

Trandate Tablets 200mg

Film-coated biconvex tablets, each containing labetalol hydrochloride 200mg, marked TRANDATE 200AH on one face. Containers of 50 £7.32 and 250 £34.30.

Trandate Tablets 400mg

Film-coated biconvex tablets, each containing labetalol hydrochloride 400mg, marked TRANDATE 400AH on one face. Containers of 50 £11.64 and 250 £55.32.

INDICATIONS

For the treatment of all grades of hypertension (mild, moderate and severe).

DOSAGE AND ADMINISTRATION

The usual initial dosage should not exceed 100mg three times daily. Dosage adjustment should be gradual. Most patients are controlled with 600mg daily or less, but severe cases may require up to 2,400mg daily. Administration should be after food. Once dosage is stabilised a twice-daily regimen may be introduced.

Children: Not applicable.

For use with other agents or transfer to Trandate from other drugs, see Data Sheet.

CONTRA-INDICATIONS, WARNINGS, ETC.

Trandate should not be given to patients with uncontrolled or digitalis-resistant heart failure or with atrioventricular block. Caution is necessary in patients with asthma. Patients with severe liver damage will probably require lower doses of Trandate than usual. Unnecessary administration of drugs during the first trimester of pregnancy is undesirable.

Warning: There have been reports of skin rashes and/or dry eyes associated with the use of beta-adrenoceptor blocking drugs. The reported incidence is small and in most cases the symptoms have cleared when the drug was withdrawn. Discontinuance of the drug should be considered if any such reaction is not otherwise explicable. Cessation of therapy with a beta-adrenoceptor blocking drug should be gradual.

Symptoms of postural hypotension may occur if initial dosage is too high or increased too rapidly but are uncommon, except at very high doses, if used as recommended. Mild and usually transient side effects include: headache, tiredness, dizziness, depressed mood, lethargy, difficulty in micturition, epigastric pain, and nausea and vomiting. Tingling in the

SELECTED BIRDS OF BRITAIN

SANDPIPER

Tringa hypoleucos

The Common Sandpiper arrives in Britain during the spring from more Southerly areas. Most pairs nest by streams and lochs in Scotland, and in more Northerly parts of England and Wales. Characteristically it flies low over water, wings bowed, in search of food such as water insects, larvae and tadpoles. Snails, worms and some vegetable matter is also taken. The call note is a high pitched and repeated "twee". During courtship the male performs a spiralling flight and utters a trilling song. The nest, often close to water, is a scrape, lined with grass and leaves, and well concealed in vegetation. 4 buff and brown-marked eggs are laid. The clutch is incubated by both parents. The sexes are of similar appearance; they reach a length of 8 inches (20 cm).

(Illustration from "A History of British Birds" by F. O. Morris)

scalp has occasionally occurred but is usually transient. Rarely, a lichenoid rash has been reported which disappears on drug withdrawal.

LEGAL CATEGORY

[POM]

PRODUCT LICENCE NUMBERS

Trandate Tablets 100mg 0045/0106

Trandate Tablets 200mg 0045/0107

Trandate Tablets 400mg 0045/0109

HC 8195/Gp -
RML/Feb 1981

Further information is available on request
Trandate is a trade mark of
ALLEN & HANBURY LTD, LONDON E2 6LA

