Triptafen: crow.

Contributors

Allen & Hanburys

Publication/Creation

London: Allen & Hanburys, 1978.

Persistent URL

https://wellcomecollection.org/works/hzsbv7zk

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).





(Amitriptyline Hydrochloride BP and Perphenazine BP)

FOR THE COMPREHENSIVE TREATMENT OF DEPRESSION AND ANXIETY NOW IN FOIL STRIP PACKS FOR ADDED CONVENIENCE AND CHILD SAFETY

PRESENTATIONS

Triptafen-DA Tablets – providing Amitriptyline Hydrochloride BP 25mg and Perphenazine BP 2mg – whenever depression predominates.

Triptafen-Minor Tablets - providing Amitriptyline Hydrochloride BP 10mg and Perphenazine BP 2mg - when anxiety is the obvious feature.

Triptafen-Forte Tablets – providing Amitriptyline Hydrochloride BP 25mg and Perphenazine BP 4mg – when agitation complicates the picture.

PACKAGE QUANTITIES AND BASIC NHS COST

All Triptafen preparations are issued as tablets in printed cartons each containing 100 tablets in 10 foil strips of 10 (5 x 2) tablets.

Triptafen-DA Tablets 100's £2.72 £3.02 £2.40 Triptafen-Forte Tablets 100's Triptafen-Minor Tablets 100's

TABLET IDENTIFICATION CODE Triptafen-Forte Tablets Triptafen-DA Tablets AH 1D 40 Triptafen-Minor Tablets

SELECTED BIBLIOGRAPHY Browne, M. W. (1969) Brit. J. Psychiat., 115, 693.

SELECTED BIRDS OF BRITAIN

CARRION CROW

Corvus corone

This crow is found all over England, Wales and the southerly parts of Scotland. In contrast to the gregarious rook, the crow is a comparatively solitary creature which can be sighted almost anywhere, but is most common over arable land or amongst trees. It eats any dead animal matter, also eggs, insects and small mammals. The call is the well-known croaking 'krah-krah-krah'. In courtship, the male spreads his wings and bowing, fans out his tail. The birds then pair for life. The nest is built in the stout fork of a tree or on a cliff ledge. Similar to, but more compact than the rook's nest, it is constructed of sticks and lined with hay and wool. Up to 5 (very rarely 7) eggs may be laid. They have a bluish-green background splashed with brown. The male and female are alike, measuring up to 20 in (50 cm).

(Illustration from "A History of British Birds" by F. O. Morris)

Chacon, C. and Downham, E. T. (1967) Brit. J. Psychiat., 113, 201. Collins, A. D. and Dundas, J. (1967) *Brit. J. Psychiat.*, **113**, 1425. Dabbs, A. R. and Lyons, J. A. (1966) *Clin. Trials J.*, **3**, 573. Haider, I. (1967) *Brit. J. Psychiat.*, **113**, 195.

McKnight, E. and Bodger, J. N. (1971) Brit. J. clin. Pract., 25, 381.

Martins, J. K. (1966) Curr. ther. Res., 8, 17.

78/218/HN

Naftulin, D. H. and Ware, J. E. (1972) *Psychosomatics*, **13**, 125. Rao, S. V. (1967) *Brit. J. clin, Pract.*, **21**, 348.

Full prescribing information is available on request. Triptafen is a trade mark of ALLEN & HANBURYS LTD LONDON E2 6LA

