

Identification and dosage card for the new BRL Penicillins.

Contributors

Beecham Research Laboratories

Publication/Creation

Brentford : Beecham Research Laboratories Ltd, [between 1955 and 1965?]

Persistent URL

<https://wellcomecollection.org/works/sq335jph>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).

**wellcome
collection**

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

Tablets and Capsules Actual Size

BROXIL

Tablets: 125 and 250 mg.
Syrup: 60 ml.
(125 mg./5 ml.)



BROCILLIN

Tablets: 125 and 250 mg.
Syrup: 60 ml.
(62.5 mg./5 ml.)



PENBRITIN

Capsules: 250 mg.
Paediatric
Tablets: 125 mg.



ORBENIN

Capsules: 250 mg.
Vials: 250 mg. to be made up for i/m injection by addition of 1.5 ml. 'Water for Injection' BP



CELBENIN

Vials: 1 G. to be made up for i/m injection by the addition of 1.5 ml. 'Water for Injection' BP



**IDENTIFICATION
AND DOSAGE CARD
FOR THE NEW
BRL PENICILLINS**

BROXIL
(potassium phenethicillin)

Oral Penicillin

BROCILLIN
(potassium propicillin)

Oral Penicillin

PENBRITIN
(ampicillin)

Oral Broad Spectrum Penicillin

ORBENIN
(sodium cloxacillin)
**Oral Penicillinase
Stable Penicillin**

CELBENIN
(sodium methicillin)
**Penicillinase
Stable Penicillin**

BEECHAM RESEARCH LABORATORIES LTD
BRENTFORD * ENGLAND
Telephone : ISLeworth 4111



9898

INDICATIONS

DOSAGE

BROXIL

**BOILS, CARBUNCLES,
OTITIS MEDIA.**
Wound and skin infections.
Tonsillitis and Laryngitis.
Lobar and Bronchopneumonia.

ADULTS:

125-250 mg. 4-6 hourly.

CHILDREN:

62.5-125 mg. 4-6 hourly.

BROCILLIN

TONSILLITIS, LARYNGITIS.
Lobar and Bronchopneumonia.
Otitis Media.
Boils and Carbuncles.
Wound and skin infections.

ADULTS:

125-250 mg. 4-6 hourly.

CHILDREN:

62.5-125 mg. 4-6 hourly.

PENBRITIN

Chronic Bronchitis.

Urinary Tract infections.
Gastro-intestinal infections.

ADULTS:

250 mg. 6 hourly.
(Prophylaxis 250 mg. 12 hourly.)
500 mg. 8 hourly.
500-750 mg. 8 hourly.

CHILDREN:

0-2 years 62.5 mg. 6 hourly.
3-10 years 125 mg. 6 hourly.

ORBENIN

**Resistant staphylococcal in-
fections.**

Mixed Gram-positive infec-
tions.

ADULTS:

Oral

500 mg. 6 hourly.

Intramuscular

250 mg. 6 hourly.

CHILDREN:

0-2 years Quarter adult dose.
3-10 years Half adult dose.

CELBENIN

**Resistant staphylococcal in-
fections.**

ADULTS:

1 G. i/m. 6 hourly.

CHILDREN:

100 mg./kg. body wt./day.