

Prodexin : good living, a series of drawings by the Cruikshanks.

Contributors

Cruikshank, Robert, 1789-1856
C.L. Bencard Ltd.

Publication/Creation

London : C.L. Bencard Ltd., [1977?]

Persistent URL

<https://wellcomecollection.org/works/heqzg2jr>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).

**wellcome
collection**

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



Drawn & Engraved by J. R. & G. Cruikshank

Tom & Jerry larking at a Masquerade Supper, at the Opera House

Good Living

A SERIES OF DRAWINGS BY THE CRUIKSHANKS

No. 4 "*The Masquerade Supper*"

By Robert and George Cruikshank

Our print is the result of collaboration between both Cruikshank brothers, and is an illustration from the book "*Life in London*" by Pierce Egan (1821).

The Ideal Antacid

The tonic effect of conviviality is usually denied to the dyspeptic or ulcer patient. And at times of dietary indiscretion, his means of relief is too often inaccessible just when it is most wanted.

On these occasions he needs rapid, predictable and *discreet* medication such as that provided by Prodexin.

The sucking of a Prodexin tablet supplies to the hyperacidic stomach a continuous flow of efficient and persistent buffer antacid. Prodexin simulates the action of an intragastric drip, but is available at any time, anywhere, in the form of a pleasant-tasting tablet.

For both occasional dyspepsia and active peptic ulceration, rapid, sustained and safe control of acidity is achieved without alkalinity or the risk of constipation.

PRODEXIN

for continuous control of gastric pH

Each Prodexin tablet contains

Aluminium glycinat (dihydroxy aluminium aminoacetate) . . . 0.9 gramme
Light magnesium carbonate 0.1 gramme

DOSAGE

To ensure a steady and prolonged flow of medicament to the stomach the tablet should be allowed to dissolve *slowly* in the mouth.

For hyperacidity

One to two tablets as required.

For prevention of hyperacidic attacks

One tablet every hour. If pain recurs regularly it is best to anticipate each attack by sucking a tablet 15 to 30 minutes before pain is expected.

For peptic ulcer

One to four tablets every hour, depending on the degree of hyperacidity.

PACKAGES

Cartons of 30 individually wrapped tablets and dispensing packs of 240.
Basic N.H.S. cost of 240 tablets: 23s. 4d.



C. L. BENCARD LTD. PARK ROYAL, LONDON N.W. 10

Telephone: ELGar 6681

Telegrams: Bencarlond, Harles, London