

**B.D.H. Androgens : methyl-testosterone B.D.H. Testosterone propionate  
B.D.H.**

**Contributors**

The British Drug Houses Ltd.

**Publication/Creation**

[1950?]

**Persistent URL**

<https://wellcomecollection.org/works/ttpvuh7h>

**License and attribution**

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>

# **B.D.H. ANDROGENS**

*Methyl-testosterone B.D.H.*

*Testosterone Propionate B.D.H.*



*B.D.H. Androgens are today effectively  
employed in gynæcological conditions in  
addition to their well-established uses in  
the male*

THE BRITISH DRUG HOUSES LTD. LONDON N.1

For oral or sublingual administration

## METHYL-TESTOSTERONE *B.D.H.*

| INDICATIONS                                   | DOSAGE   |
|---|--|
| <b>IN THE MALE</b>                            |  |
| Eunuchism complete                            | 25—50 mg. daily  |
| Eunuchoidism                                  | 10—25 mg. daily  |
| Hypogonadism                                  |  |
| Impotence                                     | 15 mg. daily   |
| Climacteric                                   | 10—25 mg. daily  |
| Premature senility                            |  |
| Benign prostatic enlargement                  | 25 mg. daily for 6 weeks, thereafter 10 mg. daily  |
| <b>IN THE FEMALE</b>                          |  |
| Metrorrhagia                                  | 25—50 mg. daily  |
| Chronic mastitis                              | 15 mg. daily   |
| Mastodynia                                    |  |
| Dysmenorrhœa                                  | 5—10 mg. daily continuously till relief is obtained, then in the post-ovulatory phase only |
| Premenstrual tension                          | 5—15 mg. daily, reducing when relief is obtained   |
| Menopause (particularly hæmorrhagic symptoms) | 10—30 mg. daily  |
| Frigidity                                     | 25—50 mg. daily until clitoris is sufficiently enlarged                                    |

The doses quoted above are minimal and may be increased at the discretion of the physician.

For administration by injection and implantation

## TESTOSTERONE PROPIONATE *B.D.H.*

| INDICATIONS                                   | DOSAGE   |
|---|--|
| <b>IN THE MALE</b>                            |  |
| Eunuchism (complete)                          | 25 mg. twice weekly  |
| Eunuchoidism                                  | 5—10 mg. twice weekly  |
| Climacteric                                   |  |
| Premature senility                            | 10—25 mg. twice weekly   |
| Hypogonadism                                  |  |
| Impotence                                     | 10 mg. twice weekly for 6 weeks followed by 5 mg. twice weekly   |
| Benign prostatic enlargement                  | 25 mg. every second to fifth day   |
| Angina pectoris                               |  |
| <b>IN THE FEMALE</b>                          |  |
| Metrorrhagia                                  | 25 mg. daily until bleeding controlled—subsequently 25 mg. every other day for 2 weeks preceding expected bleeding |
| Frigidity                                     | 25 mg. twice weekly until clitoris sufficiently enlarged   |
| Mastitis (chronic)                            | 10 mg. twice weekly  |
| Mastodynia                                    |  |
| Mammary carcinoma                             | 25 mg. daily   |
| Menopause (particularly hæmorrhagic symptoms) | 10—50 mg. daily, decreasing as control of symptoms permits   |
| Dysmenorrhœa                                  | 10—25 mg. once or twice weekly continuously until relief is obtained, then in post-ovulatory phase only            |
| Premenstrual tension                          | 10—25 mg. once or twice during week preceding menstruation, reducing when relief is obtained                       |

The doses quoted above are minimal and may be increased at the discretion of the physician.

Further information is available on request.



## PACKINGS AND PRICES

---

### Methyl-testosterone B.D.H.

|                           |        |
|---------------------------|--------|
| Tablets containing 5 mg.  |        |
| Bottle of 25              | 14/5   |
| Bottle of 100             | 51/9   |
| Bottle of 250             | 123/4  |
| Tablets containing 10 mg. |        |
| Bottle of 25              | 26/4   |
| Bottle of 100             | 94/6   |
| Tablets containing 25 mg. |        |
| Bottle of 25              | 63/-   |
| Bottle of 100             | 226/10 |
| Tablets containing 50 mg. |        |
| Bottle of 25              | 122/5  |
| Bottle of 100             | 439/2  |

### Testosterone Propionate B.D.H.

For parenteral administration

|                                      |       |
|--------------------------------------|-------|
| Ampoules containing 5 mg. per ml.    |       |
| Box of 6 × 1 ml.                     | 11/6  |
| Box of 25 × 1 ml.                    | 45/-  |
| Ampoules containing 10 mg. per ml.   |       |
| Box of 3 × 1 ml.                     | 10/2  |
| Box of 6 × 1 ml.                     | 19/7  |
| Box of 25 × 1 ml.                    | 78/9  |
| Ampoules containing 25 mg. per ml.   |       |
| Box of 1 × 1 ml.                     | 7/10  |
| Box of 3 × 1 ml.                     | 22/3  |
| Box of 12 × 1 ml.                    | 86/7  |
| Box of 25 × 1 ml.                    | 180/- |
| Ampoules containing 50 mg. per ml.   |       |
| Box of 1 × 1 ml.                     | 14/2  |
| Box of 3 × 1 ml.                     | 39/7  |
| Box of 12 × 1 ml.                    | 152/1 |
| Vials containing 50 mg. per ml.      |       |
| Vial of 10 ml.                       | 117/- |
| Ampoules containing 100 mg. in 2 ml. |       |
| Box of 1 × 2 ml.                     | 26/7  |
| Box of 3 × 2 ml.                     | 77/5  |
| Box of 12 × 2 ml.                    | 295/8 |

For implantation

|                            |      |
|----------------------------|------|
| Pellets containing 100 mg. |      |
| Tube of 1 pellet           | 38/3 |

Prices in Great Britain to the medical profession

---

THE BRITISH DRUG HOUSES LTD. LONDON N.1

TORONTO · SYDNEY · BOMBAY · JOHANNESBURG · AUCKLAND

M&TstP/F/BI/5006/M

Printed in Great Britain