

Brontisol : aerosol therapy for generalized airways obstruction.

Contributors

The British Drug Houses Ltd.

Publication/Creation

[1993?]

Persistent URL

<https://wellcomecollection.org/works/cjr3v7jx>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

brontisol*



Aerosol therapy for
generalized airways obstruction

PRESCRIBING INFORMATION

Indications - Chronic airways obstruction due to asthmatic bronchitis, chronic bronchitis or asthma, and all other respiratory conditions where bronchodilation, control of hypersecretion and protection against allergic effects on the respiratory system are advantageous.

Dose

Adults 1-3 inhalations every 4-5 hours as directed by the physician.

Children 1-2 inhalations every 4-5 hours as directed by the physician. For children under 6 years of age, administration should be supervised by a responsible adult. Not more than 12 inhalations in 24 hours.

Side Effects - The side effects which may be encountered are those normally associated with the use of sympathomimetic and anticholinergic drugs: - dizziness, tachycardia, dry mouth and blurred vision. These side effects do not occur frequently and are seldom severe. In practice they will usually lessen or disappear with continued usage. Occasionally dosage adjustment may be necessary.

Special Precautions - Brontisol should be used with caution in the presence of impaired cardiac function, tuberculosis, diabetes, hyperthyroidism and hypertension.

Pack Information - Brontisol is available in the form of a 15 ml. self-propelled metered dose aerosol complete with an anatomically designed plastic mouthpiece.

brontisol*



Aerosol therapy for generalized airways obstruction

Formula - Brontisol is a combination of dectropine citrate 2 mg/ml. and isoprenaline hydrochloride 3 mg/ml. suspended in an inert propellant. The 15 ml cartridge delivers 300 doses each containing:

Deptropine citrate	0.10 mg.
Isoprenaline hydrochloride	0.15 mg.

Presentation - Each complete unit, contained in one carton, consists of the 300 dose metal container, a plastic mouthpiece and a comprehensive instruction leaflet for patients, including simple breathing exercises.

National Health Service Cost

Available only as one complete unit.

Basic N.H.S. price 16/6 d. + purchase tax.



BrocaDes|Great Britain|Ltd.,



Trend House, Pyrford Road, West Byfleet, Surrey.

Sole distributors in the U.K. Camden Chemical Co. Ltd. 61 Gray's Inn Road, London, W.C.1.