Your arse can absorb HIV like a sponge: if you get fucked without a condom HIV can enter the bloodstream through the lining in the arse / GMFA.

Contributors

Gay Men Fighting AIDS (Organization)

Publication/Creation

[between 2005 and 2015?]

Persistent URL

https://wellcomecollection.org/works/cmccmd35

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org

If you get fucked without a condom, the gay men's he HIV can enter the bloodstream through the lining in the arse.



YOUR CAN ABSORB HIV LIKE A SPONGE



- Most gay men with HIV became infected by getting fucked without a condom. This is because the thin lining of your arse is extremely efficient at absorbing HIV.
- Your arse does not have to be damaged or bleeding to be vulnerable to HIV. However, damage to the lining inside your arse does increase the risk of transmission.
- Dildos or fisting can damage the lining of your arse without you noticing, and douching (washing the inside of your arse) can cause irritation. These can make your arse even more vulnerable to infections like HIV.
- Having an STI can also make it easier to catch or pass on HIV. If you fuck or get fucked, using condoms is the most reliable way to prevent the transmission of HIV.

For information on how HIV is transmitted: Visit: www.gmfa.org.uk/cocksure Call: London Lesbian and Gay Switchboard, 020 7837 7324; THT Direct on 0845 12 21 200.

To express your views on HIV services in London, visit: www.ergoclear.com/express

