

You can get HIV through your cock : if you fuck without a condom, HIV can enter the bloodstream through the cock / GMFA.

Contributors

Gay Men Fighting AIDS (Organization)

Publication/Creation

[between 2005 and 2015?]

Persistent URL

<https://wellcomecollection.org/works/s8bpetct>

License and attribution


Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



1102



If you fuck
without a condom,

HIV can enter the
bloodstream
through
the cock.

YOU CAN GET HIV
THROUGH YOUR

COCK

- ❖ Some men with HIV became infected when they were the active partner and fucked without a condom. During anal sex, HIV can pass through the tissue that lines the tube you piss out of, your foreskin or the head of your cock.
- ❖ Your cock does not need to be damaged for HIV to get in. However, damage, like a sore or a cut, does make transmission more likely. When fucking, tiny abrasions can appear on the skin of your cock that then allow HIV in.
- ❖ Anal mucus, which lines the inside of your arse, can contain more HIV than cum. So if you have HIV and get fucked without a condom, you can pass the infection to your partner.
- ❖ Having an STI can also make it easier to catch or pass on HIV. If you fuck or get fucked, using condoms is the most reliable way to prevent the transmission of HIV.

This resource is funded by the Pan London HIV Prevention Programme.

GMFA projects are developed by positive and negative volunteers.
To volunteer or donate, go to www.gmfa.org.uk or call 020 7738 6872.
Charity no. 1076854.