

How HIV transmission works: getting fucked : fluids were a matter of some concern for Howard / GMFA, London Gay Men's HIV Prevention Partnership, MetroM8.

Contributors

London Gay Men's HIV Prevention Partnership
Gay Men Fighting AIDS (Organization)
MetroM8

Publication/Creation

[2009?]

Persistent URL

<https://wellcomecollection.org/works/q9t7uz4u>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

How HIV transmission works: Getting Fucked



**Fluids were a matter of some
concern for Howard**

How HIV transmission works: Getting Fucked

There is HIV in pre-cum but even more in cum. The inside of the arse absorbs liquid like a sponge. If HIV-infected cum or pre-cum gets into the arse, from fucking without a condom, HIV can get directly into the bloodstream.

Being fucked can also cause scratches and tears in the lining of the arse. When this happens, the cells beneath the damaged skin are the ones that are most easily infected by HIV.

For more information on HIV & AIDS call THT Direct Helpline on 0845 122 1200 Mon-Fri 10am-10pm; Sat-Sun 12noon-6pm or visit www.metromate.org.uk

GMFA's campaigns and actions are designed, planned and executed by positive, negative and untested volunteers. To volunteer for GMFA, phone, write or e-mail: 020 7738 6872
Unit 43, The Eurolink Centre, 49 Effra Road, London SW2 1BZ. newvol@gmfa.org.uk
Registered Charity no: 1076854 Part of the London Gay Men's HIV Prevention Partnership.



LONDON GAY MEN'S
HIV Prevention Partnership

