

HIV : now there is a treatment that may prevent HIV infection after the virus has entered the body : PEP post exposure prophylaxis : treatment only works if started within 72 hours of exposure : 1 month course of anti-HIV medication ... / Terrence Higgins Trust, developed as part of CHAPS.

Contributors

Terrence Higgins Trust

Publication/Creation

[2004]

Persistent URL

<https://wellcomecollection.org/works/ncsag3c9>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

**NOW THERE IS A TREATMENT
THAT MAY PREVENT HIV
INFECTION AFTER THE VIRUS
HAS ENTERED THE BODY**

PEP

POST EXPOSURE PROPHYLAXIS

**TREATMENT ONLY WORKS IF STARTED
WITHIN 72 HOURS OF EXPOSURE**

1 MONTH COURSE OF ANTI-HIV MEDICATION

**CAUTION: PEP IS *NOT*
A SUBSTITUTE FOR CONDOMS**

**SIDE EFFECTS CAN INCLUDE DIARRHOEA,
NAUSEA AND SEVERE HEADACHES**

**terrence
higgins
TRUST** 

For information you need to know about
PEP, visit www.chapsonline.org.uk call
THT DIRECT HELPLINE **0845 12 21 200** or pick
up a copy of the leaflet from gay venues.