

Act fast and PEP can stop HIV / Terrence Higgins Trust ; funded by the Pan-London HIV Prevention Programme.

Contributors

Terrence Higgins Trust
Pan London HIV Prevention Programme

Publication/Creation

[approximately 2011?]

Persistent URL


<https://wellcomecollection.org/works/faua4xz2>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



ACT FAST and PEP can STOP HIV

terrence
HIGGINS
TRUST



CAUTION

PEP (post exposure prophylaxis) is a **one month** course of anti-HIV pills. Side effects can include diarrhoea, nausea and severe headaches.

PEP is NOT a substitute for condoms.


STOP HIV

PEP may prevent HIV infection after the virus has entered the body.

Treatment is available at all London sexual health clinics and hospital emergency departments.

ACT FAST

You must start PEP within 72 hours of unsafe sex or a condom breaking.

For more details: www.gmfa.org.uk/pep or call **THT DIRECT** GUY SEXUAL HEALTH  020 7998 4161