

**'Migril' masters migraine : "the toast is 'Migril'".**

**Contributors**

Eccles, W. J.  
Burroughs Wellcome and Company

**Publication/Creation**

[betwen 1962 and 1964?]

**Persistent URL**

<https://wellcomecollection.org/works/g9rjbk7k>

**License and attribution**

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>



**'MIGRIL'**  
masters  
migraine

*“The toast is ‘Migril’ ”*

*‘Migril’ will master all the primary  
symptoms of a migraine attack.*

*‘Migril’ provides the most effective  
means of stopping the migraine  
headache and ocular disturbances  
as well as nausea and vomiting.*

*Carried in the pocket or handbag*

*‘Migril’ gives confidence to the  
migraine sufferer that he possesses  
insurance against another attack.*

*‘Migril’ is the all-round Master  
of Migraine.*

# 'MIGRIL' brand Ergotamine Compound

## Indications

'Migril' is indicated in all forms of migraine.

## Dosage

'Migril' should be taken at the first warning of an attack, 1 to 2 tablets initially, followed if necessary by  $\frac{1}{2}$  to 1 tablet at half-hourly or hourly intervals. The smallest effective dose should always be used and should never exceed 4 tablets for any single attack. Not more than 12 tablets should be taken in the course of a week. Children should be given proportionately less.

## Side-Effects

Rarely there may be sensitivity to ergotamine; if, therefore, a patient feels a sensation of coldness or aching in the extremities, he should take no more tablets. Occasionally abdominal pain may occur.

## Contra-Indications

'Migril' is contra-indicated in the presence of peripheral vascular disease, coronary disease, kidney and liver diseases, pregnancy and febrile conditions.

**'MIGRIL'** brand Ergotamine Compound

the all-round master  
of migraine

Each 'MIGRIL' Tablet contains:

Ergotamine Tartrate            2 mgm.

Cyclizine Hydrochloride    50 mgm.

Caffeine                            100 mgm.

*Basic cost to N.H.S.: 7s.1d. for 10 tablets*



**BURROUGHS WELLCOME & CO., LONDON**

*(The Wellcome Foundation Ltd.)*