'Migril' masters migraine : a first!.

Contributors

Eccles, W. J. Burroughs Wellcome and Company

Publication/Creation

[betwen 1962 and 1964?]

Persistent URL

https://wellcomecollection.org/works/qyxpc5ud

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).





'MIGRIL' masters migraine

A first!

'Migril' so effectively masters all the primary symptoms of migraine that the victim will not live in fear of his complaint or its treatment.

'Migril' rapidly overcomes the headache, ocular disturbances, and nausea and vomiting.

'Migril' Masters Migraine.

'MGRIL' brand Ergotamine Compound

Indications

'Migril' is indicated in all forms of migraine.

Dosage

'Migril' should be taken at the first warning of an attack, 1 to 2 tablets initially, followed if necessary by ½ to 1 tablet at half-hourly or hourly intervals. The smallest effective dose should always be used and should never exceed 4 tablets for any single attack. Not more than 12 tablets should be taken in the course of a week. Children should be given proportionately less.

Side-Effects

Rarely there may be sensitivity to ergotamine; if, therefore, a patient feels a sensation of coldness or aching in the extremities, he should take no more tablets. Occasionally abdominal pain may occur.

Contra-Indications

'Migril' is contra-indicated in the presence of peripheral vascular disease, coronary disease, kidney and liver diseases, pregnancy and febrile conditions.

'MGRIL' brand Ergotamine Compound

the all-round master of migraine

Each 'MIGRIL' Tablet contains:

Ergotamine Tartrate 2 mgm.

Cyclizine Hydrochloride 50 mgm.

100 mgm. Caffeine

Basic cost to N.H.S.: 7s.1d. for 10 tablets



BURROUGHS WELLCOME & CO., LONDON

(The Wellcome Foundation Ltd.)