

**'Pro-Actidil'.**

**Contributors**

Burroughs Wellcome and Company

**Publication/Creation**

1968.

**Persistent URL**

<https://wellcomecollection.org/works/kfgvbj9q>

**License and attribution**

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>



# 'Pro-Actidil'

Triprolidine Hydrochloride Tablets

## Dosage

For adults, and children over 10 years: one tablet should be swallowed whole every 24 hours—preferably between 6 and 7 p.m. (or 4 to 5 hours before retiring) so that, should drowsiness occur, it will do so during normal sleeping time. Very severe cases may require 2 tablets in 24 hours. Children under 10 years should be given 'Actidil' Elixir.

## Side-effects and Precautions

In case drowsiness occurs, patients should be warned to determine their individual response to the drug before driving or doing any mechanical work which requires concentration. Alcohol may enhance any sedative effect of 'Pro-Actidil' therapy.

## Presentation

'Pro-Actidil' brand Triprolidine Hydrochloride, 10 mg. tablets.

N.B. Triprolidine hydrochloride is also available as 'Actidil' 2.5 mg. tablets (useful for daytime use and when a short duration of action is required) and as 'Actidil' Elixir containing 2 mg. in each 5 ml. (ideal for all children).

Basic cost to N.H.S.: 4s. 6d. for 10 tablets.

Additional information is available on request.



**Burroughs Wellcome & Co.** (The Wellcome Foundation Ltd.) **London**

7989a (H) Printed in England Jan 68-32 A