

Childhood protection : the following schedule was suggested at a Symposium on Childhood Immunization held in London, May 1959.

Contributors

Burroughs Wellcome and Company

Publication/Creation

[1961?]

Persistent URL

<https://wellcomecollection.org/works/h5j3v6zz>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).

**wellcome
collection**

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

Childhood Protection

The following Schedule was suggested at a Symposium on Childhood Immunization held in London, May 1959
Brit. Med. J. (1959), 1, 1342

AGE	VISIT	VACCINE	NO. OF INJECTIONS	INTERVAL
2-6 months	1, 2, 3	★ Triple (<i>diphtheria, tetanus, pertussis</i>)	3	4 weeks or more
7-10 months	4, 5	□ Poliomyelitis	2	4 weeks or more
15-18 months	6	★ Triple (<i>diphtheria, tetanus, pertussis</i>)	1	same visit
		□ Poliomyelitis	1	
Smallpox sometime during first 5 years				
School entry	7	⊙ Diphtheria and tetanus	1	
8-9 years	8	⊙ Diphtheria and tetanus (small dose, e.g. 0.25 c.c.) Smallpox (<i>revaccination</i>)	1	same visit
10-15 years	9	B.C.G.	1	

The following vaccines are suitable:

- ★ 'TRIVAX' BRAND Triple Antigen (DTP/vac)
- 'POLIMYLEX' BRAND Poliomyelitis Vaccine (POL/vac)
- ⊙ 'WELLCOME' BRAND Diphtheria-Tetanus Prophylactic (DT/vac)

'TRIVAX' BRAND Triple Antigen The original half volume vaccine for simultaneous immunization against diphtheria, tetanus and whooping cough. Each 0.5 c.c. of 'Trivax' contains 25 Lf of diphtheria toxoid, 5 Lf of tetanus toxoid and 20,000 million *B. pertussis* – an ideal balance of components for triple protection.
Set of 3 × 0.5 c.c. ampoules and rubber-capped bottles of 5 c.c.

'POLIMYLEX' BRAND Poliomyelitis Vaccine A Salk-type vaccine containing formalin-inactivated cultures of the three immunological groups of poliomyelitis virus. The strains used are:
Type 1 – Modified Brunhilde (Enders). Type 2 – M.E.F.-1. Type 3 – Saukett.
Ampoules of 1 c.c. and rubber-capped bottles of 10 c.c.

'WELLCOME' BRAND Diphtheria-Tetanus Prophylactic A mixture of purified diphtheria and tetanus toxoids. Each 0.5 c.c. contains 25 Lf of diphtheria toxoid and 5 Lf of tetanus toxoid – the most effective strengths for reinforcement of immunity in the older child.
Ampoules of 0.5 c.c. and rubber-capped bottles of 5 c.c.



BURROUGHS WELLCOME & CO. (The Wellcome Foundation Ltd.) LONDON

6693

Printed in England

L61.3.35

APR 1961