

Real relief required : for pain of all dimensions prescribe Doxypol first.

Contributors

CAPS

Publication/Creation

[1979?]

Persistent URL

<https://wellcomecollection.org/works/dmab65w9>


License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

REAL RELIEF REQUIRED




FOR PAIN OF ALL DIMENSIONS

- mild to severe
- acute, chronic, persistent, recurrent
- of any etiology

PRESCRIBE DOXYPOL FIRST

the 3-in-1 analgesic



The analgesic properties of d-Propoxyphene are potentiated in combination with paracetamol

DOXYPOL offers these advantages:

- ☐ No codeine: No constipation
- ☐ No aspirin: Does not irritate the gastro-intestinal mucosa
- ☐ No phenacetin
- ☐ Remarkably well tolerated

PP Medication: Only available on doctor's R_x

Dosage: One capsule 3-4 times daily with food


DOXYPOL

PROMPT POTENT PREDICTABLE



A PHARMACEUTICAL HALLMARK FOR QUALITY, EFFICACY AND ECONOMY

d-Propoxyphene 65 mg
Paracetamol 300 mg
Caffeine 30 mg
Reg. No. 32,278/91
DOXYPOL capsules PP



REGISTRATION NUMBER

3/2.9/91

PHARMACOLOGICAL

CLASSIFICATION

2.9 non-narcotic analgesic and
antipyretic

DISTRIBUTION CATEGORY

PP

TRADE NAME

DOXYPOL

COMPOSITION

65 mg dextropropoxyphene
hydrochloride

300 mg paracetamol

30 mg caffeine per capsule

IDENTIFICATION

White powder in red and yellow capsule

PHARMACOLOGICAL ACTION

The analgesic properties of dextropropoxyphene hydrochloride are potentiated in combination with paracetamol (an analgesic and an antipyretic); and caffeine (a central nervous system stimulant) is considered to enhance analgesic action.

INDICATIONS

Doxypol is indicated for the relief of pain of moderate severity including chronic and persistent conditions.

DOSAGE AND DIRECTIONS FOR USE

Adults one capsule when necessary up to a maximum of four in twenty-four hours

Children over 12 years one to two capsules per day after meals

SIDE EFFECTS AND SPECIAL PRECAUTIONS

Doxypol may cause nausea on an empty stomach; and dextropropoxyphene hydrochloride should be given with care to patients suffering from respiratory depression.

CONDITIONS OF REGISTRATION OF THE MEDICINE

None


PRESENTATION

Containers of 20, 100 and 1 000 capsules

STORAGE DIRECTIONS

Keep well closed in a cool dark dry place beyond the reach of children

Further information is available on request



DOXYPOL

FOR REAL RELIEF



A PHARMACEUTICAL HALLMARK FOR QUALITY, EFFICACY AND ECONOMY