For winter chest infections Tetralysal : marked clinical superiority : superior absorption higher blood levels lower dosage improved tolerance.

Contributors

Carlo Erba

Publication/Creation

[approximately 1964]

Persistent URL

https://wellcomecollection.org/works/ujtm6mpa

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



CHEST, decorated with carvings and inlay of the so-called « Carthusian » type. Lombardy. XVth century.

83.5 × 209 × 53 cm; walnut, inlaid in maple ebony and woods of various fruit trees. Inv. no. 17

The left side, lid and lower portion of the arch motif have been restored. The closing plate is missing.

plate is missing. The decoration of the front is divided into seven square panels,inlaid with various geometric, vegetable and architectonic motives on a series of blind arches. It is already of Renaissance style. Typical of the 15th century, it is decorated with sunken squares, while the height of the chest, the small niche-like inlays between the squares, and the twisted frame place this chest among works completed in the transition period. The motif of the row of embossed arches is found both in articles of German workmanship and those from Northern Italy.

Reproduced from « I mobili nelle civiche raccolte artistiche di Milano » - edited by Gilda Rosa and published by Martello. Photographs taken in the «Castello Sforzesco » in Milan.





FOR WINTER CHEST INFECTIONS TETRALYSAL® (Lymecycline)

marked clinical superiority

SUPERIOR ABSORPTION HIGHER BLOOD LEVELS LOWER DOSAGE IMPROVED TOLERANCE

TETRALYSAL®

INDICATIONS: All infections caused by germs sensitive to tetracyclines

PACKINGS

ORAL FOR ADULTS - Bottles of 16, 100 and 1000 capsules. Each capsule contains tetracycline-l-methylene-lysine 204 mg equivalent to 150 mg tetracycline base.

ORAL FOR CHILDREN - Boxes of 10 tablets. Each tablet contains tetracycline-l-methylene-lysine equivalent to 40 mg tetracycline base.

INJECTION FOR ADULTS - Boxes of 1 and 10 vials of 135 mg lyophilized tetracycline-l-methylene-lysine, equivalent to 100 mg tetracycline base with ampoules of solvent, each containing 40 mg procaine.

INJECTION FOR CHILDREN - Boxes of 1 and 10 vials of 67 mg lyophilized tetracycline-1-methylene-lysine, equivalent to 50 mg tetracycline base with ampoules of solvent each containing 20 mg procaine.

DOSAGE

Orally: Adults - 4 capsules daily in divided doses Children - 10 mg/kg/daily in divided doses

Intramuscularly: 2-3 vials daily in divided doses Dosage may be increased in the more serious cases.

NHS COST:

Tetralysal is the least expensive tetracycline available under the NHS.

REFERENCES:

full list of reference may be obtained on request.

Brit. J. Geriat., (1963), 12 Medical Digest, (1964), 9, 41 J. Coll. Gen. Practit., (1964), 8, 140 Brit. Med. J., (1964), 11, 1491 Brit. Med. J., (1964), 11, 1495

CARLO ERBA MILANO

SOLE DISTRIBUTORS FOR U.K.: FRASER CHEMICALS LTD. 88, Lewisham High Street London S.E. 13 Tel. LEEgreen 7485