

For winter chest infections Tetralysal : marked clinical superiority : superior absorption higher blood levels lower dosage improved tolerance.

Contributors

Carlo Erba

Publication/Creation

[approximately 1964]

Persistent URL

<https://wellcomecollection.org/works/qda6dukn>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>




Gilt CHEST. Venice. XVth century.

58 × 166 × 49 cm. poplar-wood. Inv. no. 42

The shape of this exhibit, the absence of supports and the proportions still belong to the fifteenth century. The rich decoration in black and gold, with stylized plant motives is arranged in bands. The upper portion contains two coats-of-arms in typical Venetian shields. Two iron handles are fixed at the sides.

Reproduced from « I mobili nelle civiche raccolte artistiche di Milano » - edited by Gilda Rosa and published by Martello. Photographs taken in the « Castello Sforzesco » in Milan.



FOR WINTER CHEST INFECTIONS

TETRALYSAI[®]

(Lymecycline)

ERBA

marked clinical superiority

SUPERIOR ABSORPTION
HIGHER BLOOD LEVELS
LOWER DOSAGE
IMPROVED TOLERANCE

TETRALYSAL[®]

(Lymecycline)

ERBA

INDICATIONS: All infections caused by germs sensitive to tetracyclines

PACKINGS

ORAL FOR ADULTS - Bottles of 16, 100 and 1000 capsules. Each capsule contains tetracycline-l-methylene-lysine 204 mg equivalent to 150 mg tetracycline base.

ORAL FOR CHILDREN - Boxes of 10 tablets. Each tablet contains tetracycline-l-methylene-lysine equivalent to 40 mg tetracycline base.

INJECTION FOR ADULTS - Boxes of 1 and 10 vials of 135 mg lyophilized tetracycline-l-methylene-lysine, equivalent to 100 mg tetracycline base with ampoules of solvent, each containing 40 mg procaine.

INJECTION FOR CHILDREN - Boxes of 1 and 10 vials of 67 mg lyophilized tetracycline-l-methylene-lysine, equivalent to 50 mg tetracycline base with ampoules of solvent each containing 20 mg procaine.

DOSAGE

Orally: Adults - 4 capsules daily in divided doses
Children - 10 mg/kg/daily in divided doses

Intramuscularly: 2-3 vials daily in divided doses
Dosage may be increased in the more serious cases.

NHS COST:

Tetralysal is the least expensive tetracycline available under the NHS.

REFERENCES:

full list of reference may be obtained on request.

Brit. J. Geriat., (1963), 12

Medical Digest, (1964), 9, 41

J. Coll. Gen. Practit., (1964), 8, 140

Brit. Med. J., (1964), II, 1491

Brit. Med. J., (1964), II, 1495

CARLO ERBA MILANO

SOLE DISTRIBUTORS FOR U.K.: FRASER CHEMICALS LTD. 88, Lewisham High Street London S.E. 13 Tel. LEEgreen 7486