

Argyrol.

Contributors

Fassett & Johnson

Publication/Creation

[between 1904 and 1909?]

Persistent URL

<https://wellcomecollection.org/works/fym7tzmh>

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

ARGYROL

(BRAND) SILVER VITELLIN

NON-IRRITANT - NON-TOXIC

Posological Table

EAR, NOSE AND THROAT.

ANTRUM INFECTION. Instil 10% solution with other suitable measures.

ASTHMA. Dowling Argyrol pack treatment. Snugly fitting cotton wool swabs soaked in 10% solution and inserted high into both nostrils on wire or long probes. Keep in position 20 minutes. After withdrawal thoroughly cleanse the nares.

CHRONIC OTORRHOEA. Swab with 5% sol.

COMMON WINTER COLD. Tampons soaked in 10% to 20% solution.

DEAFNESS in children. Instil few drops of 25% sol., follow by mineral oil spray.

ETHMOIDITIS. As for Asthma.

FURUNCULOSIS (nasal vestibule). Constantly apply nasal pack soaked with 20% solution.

HAY FEVER. Insert pledgets soaked in 10% solution. Retain for 15 to 30 minutes.

LARYNGITIS. Spray with 10% to 20% sol.

NASAL INFECTIONS. Swab or spray nose with 20% solution.

OTITIS MEDIA. Cleanse and gently plug with gauze soaked in 10% to 20% sol. Water and Glycerine in equal parts.

PHARYNGITIS. Swab or spray with 20% to 30% solution every 4 hours.

POST-OPERATIVE Spray 20% solution in throat and nose or as drops in ear.

RHINITIS. Instil 20 minims of 10% sol.

SINUSITIS. Cleanse nose and instil 5 drops of 10% solution.

TONSILITIS. As pharyngitis.

TRACHEITIS. As laryngitis.

OPHTHALMOLOGY.

CONJUNCTIVITIS.

Catarrhal, 10% to 25% solution, repeat frequently; instil also into nose.

Diphtheritic, 25% sol. eyes and nostrils.

Follicular, 10% to 15% solution.

Gonorrhoeal, Gentle, frequent, thorough cleansing and then instillations of 25% to 50% solution.

Ophthalmia Neonatorum, Frequent cleansing and instillations of 25% sol.

PRE- & POST-OPERATIVE. 15% instillations.

TRACHOMA. Remove granulations. Apply 10% to 30% solutions.

UROLOGY & GYNAECOLOGY.

CHANCROID. Cleanse and dry; then apply pure powdered Argyrol freely two or three times a day.

CYSTITIS. Irrigate bladder 1-500 solution; then instil 1-ounce 10% solution.

GONORRHOEA (Male) Small syringe, Inject 3-10% solution T.D.S.

(Female) Inject 5-10% solution in urethra and bladder. Instil similar solution into cervical os. Insert vaginal tampon soaked in 20% solution for several hours each day.

GONORRHOEAL PROCTITIS. 1-500 rectal irrigations followed by injections of 10% solution.

PYELITIS. Inject per ureter catheter 1-500 solution or stronger up to 5%.

VAGINAL DISCHARGE, same as Gonorrhoea.

VULVO-VAGINITIS (children). Swab vulva with and instil into vagina 10% solution.

Supplied in Crystals and Tablets

FASSETT & JOHNSON, LTD.

86 Clerkenwell Road, London, E.C.1.

SYDNEY

CAPE TOWN

WELLINGTON