Something old : poppie (papaver somniferum) : something new in the treatment of cough Extil compound linctus.

Contributors

Duncan, Flockhart & Co.

Publication/Creation

[approximately 1965]

Persistent URL

https://wellcomecollection.org/works/cetc2gw5

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org





something **NEW** in the treatment of cough

Extil Trade Mark compound linctus

Extil compound linctus

Poppie

Of the heades boyled in water, is made a Syrupe, which doth al/o cau/e Sleep, and is very good agayn/t the *f*ubtil Rheumes, and Catharres, that di/till and fal downe from the brayne upon the lunges, and agayn/t the cough, taking his beginning of *f*uch *f*ubtil humours.

Rembert Dodoens, A NIEVVE HERBALL OR HISTORIE OF PLANTES Translated out of French into Engli/h by Henry Lyte E/quyer London 1578

References:

Bickerman, H. A. and Barach, A.L., (1954), Am. J. med. Sci., 228, 156-163. Bickerman, H. A., et al., (1957) Am. J. med. Sci. 234, 191-205. Harris, E. A., (1965) Brit. J. Pharmacol. 24, 532-537.

Bergmann, M., and Stolzer, H. (1956) Wien. med. Wochschr. 106, 232-233. Janschulte, B. (1956) Med. Klin. (Berlin), 50, 1788-1789.

Segal, M. S. et al., (1957) Diseases of Chest, 32, 305-309,

World Health Organisation (1959) Wld. Org. tech. Rep. Ser., 160, 10.

Each adult dose (10 ml) contains: Noscapine 25 mg Pseudoephedrine 49.2 mg (equivalent to 60 mg pseudoephedrine hydrochloride) Carbinoxamine maleate 6 mg

NOSCAPINE

• recommended by the World Health Organisation (1959) as a substitute for codeine for the suppression of cough.

• is an isoquinoline derivative and does not produce addiction.

• unlike codeine it has no narcotic or analgesic effect and, in normal dosage, does not cause constipation.

• unlike codeine it does not depress the respiratory centre. (Harris 1965).

• confirmed antitussive properties in man with experimental cough. (Bickerman and Barach 1954; Bickerman *et al.*, 1957).

• confirmed antitussive properties in man with pathological cough. (Bergmann and Stolzer 1956; Janschulte 1956; Segal *et al.*, 1957).

PSEUDOEPHEDRINE

a powerful decongestant.

 reduces bronchospasm without impairing ciliary action thereby facilitating the removal of mucus.

 resembles ephedrine pharmacologically but causes less central stimulation.

CARBINOXAMINE

a potent antihistamine.

well tolerated—it has only slight sedative properties.

 complements the action of pseudoephedrine to promote decongestion of the WHOLE of the upper respiratory tract.

INDICATIONS:

Troublesome cough, especially cough accompanied by bronchospasm or congestion of the upper respiratory tract.

Cough associated with the common cold; in such conditions EXTIL COMPOUND LINCTUS not only relieves the cough but reduces both nasal and bronchial congestion.

DOSAGE:

Adults and children over 12:

Two teaspoonfuls (10 ml) three or four times a day as required.

Children over 2 years and under 12:

Half to one teaspoonful (2.5 to 5 ml) three or four times a day as required.

CONTRAINDICATIONS:

EXTIL COMPOUND LINCTUS should not be given to patients suffering from advanced hypertension or heart failure.

PRESENTATION:

EXTIL COMPOUND LINCTUS is available in bottles of 5 fluid ounces.

BASIC N.H.S. COST:

The basic N.H.S. cost of a bottle of 5 fluid ounces is 5/- plus P.T.

Manufactured in England by DUNCAN. FLOCKHART and EVANS Ltd., LONDON E2





DUNCAN, FLOCKHART and EVANS Ltd LONDON E2