

Pollenosis source and season : annual meadow (poa annua)... : for the patient suffering from severe hay fever Extil.

Contributors

Duncan, Flockhart & Co.

Publication/Creation

[between 1960 and 1969?]

Persistent URL

<https://wellcomecollection.org/works/pbfezs9t>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).

**wellcome
collection**

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



Pollenosis
source and season



Annual Meadow

(Poa annua)

seems to prefer the waste land, sprouting (often unwanted) from gravel paths and on stony ground. It has a long flowering season, from May right through to August.

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC

THIS CARD IS ONE OF A SERIES ON COMMON CAUSATIVE FACTORS OF HAY FEVER IN GREAT BRITAIN ISSUED BY



TO THE MEDICAL PROFESSION

For the patient suffering
from severe hay fever

Extil

Trade Mark

especially where there is
associated bronchospasm

EXTIL TABLETS

Each EXTIL TABLET contains:

Carbinoxamine maleate 6mg
Pseudoephedrine hydrochloride 60mg

EXTIL SYRUP

Each 10ml (2 teaspoonfuls) contains:

Carbinoxamine maleate 6mg
Pseudoephedrine hydrochloride 60mg

The combination of the potent anti-allergic properties of carbinoxamine with the nasopharyngeal vasoconstrictor properties of pseudoephedrine provide symptomatic control of hay fever, allergic rhinitis, and vasomotor rhinitis. Oral therapy with EXTIL eliminates the need for topical therapy with nasal drops or sprays which, all too frequently, give rise to the risk of abuse and subsequent habituation.

EXTIL TABLETS

Dosage:

Adults and children over 12 years:
One tablet every four to six hours.

EXTIL SYRUP

Dosage:

Adults and children over 12 years:
Two teaspoonfuls every four to six hours.
Children over 2 years and under 12 years:
Half to one teaspoonful every four to six hours.



MADE IN ENGLAND
DUNCAN, FLOCKHART and EVANS Ltd LONDON E2