Furadantin dosage card - tablet of nitrofurantoin 50 mg.

Contributors

Duncan, Flockhart & Co.

Publication/Creation

[between 1950 and 1959?]


Persistent URL

https://wellcomecollection.org/works/xh8vhh7z

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).





11/2

2

NUMBER OF TABLETS -50 mg. - four times daily The above are average doses. The dose should lie in the range 5-10 mg./kg. (2.2-4.5 mg./lb.) It may be reduced towards the lower limit if nausea occurs, and increased towards the upper if the infection is slow to respond.

3

31/2

4

21/2

'FURADANTIN' is an antibacterial agent for the specific treatment of urinary tract infections. It is active against most organisms responsible for such infections, including many strains of B. proteus and some strains of Ps. pyocyanea.

Bacterial resistance to 'Furadantin' develops very slowly if at all. 'FURADANTIN' cannot cause cross-resistance to antibiotics.

'FURADANTIN' neither interferes with the bacterial flora of the large intestine nor favours the development of yeast infections:

INDICATIONS:

Cystitis

Pyelitis (including pyelitis in pregnancy and childhood)

Pyelonephritis

Urethritis Prostatitis Before and after prostatectomy


Infection following

passage of instruments Trichomonas infection

in the male

ADMINISTRATION:

see dosage table overleaf. To avoid nausea the tablets should be taken during meals. In the treatment of chronic infections, 'Furadantin' has been administered continuously for long periods without toxic effects. The longest period reported is 2 years.



Sole distributors in the U.K.