

For the control of pain...just one tablet : Paramol-118.

Contributors

Duncan, Flockhart & Co.

Publication/Creation

[1975]

Persistent URL

<https://wellcomecollection.org/works/fuhuv27x>

License and attribution


Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

For
the control
of PAIN...

just one tablet



Paramol-118

Paramol-118

is an effective combination of two analgesics in low dosage. Although each tablet contains only 500 mg paracetamol fortified with a low dose (10 mg) of dihydrocodeine, **a single tablet every four hours can be relied upon to relieve most of the aches and pains you meet daily in your practice.**

To the Medical Profession

just one tablet

Paramol-118

low dosage, low cost
in the control of PAIN

PARAMOL-118 TABLETS

Presentation White tablets engraved BDH/P each tablet containing:

Paracetamol BP 500 mg
Dihydrocodeine tartrate BP 10 mg

Indications Muscular conditions such as fibrositis, lumbago and other back pain; also in other commonly encountered conditions of persistent pain such as sprains and strains, dysmenorrhoea, severe headache, sinusitis and neuralgia.

Dosage and administration Paramol-118 tablets should, if possible, be taken during or after meals.

Adults and children over 12 years: One tablet every four hours. This may be increased to two tablets every four hours if necessary, or as required. Paramol-118 is not recommended for children under 12.

Contraindications, warnings, etc. There are no contraindications.

Precautions: Paramol-118 should be given with caution to patients with allergic disorders and should not be given during an attack of asthma. Caution should also be observed if there is marked impairment of liver function or advanced kidney disease. Notice should be taken of current views that any drug should be prescribed with caution during pregnancy.

Side Effects: Constipation, if it occurs, is readily treated with a mild laxative. Nausea and giddiness may occur in a few patients.

Overdosage: Conservative management is recommended; gastric lavage should be carried out. Severe respiratory depression can be treated with nalorphine 5-10 mg intravenously, which may be cautiously repeated to a total of 40-100 mg.

Package Quantities Paramol-118 Tablets are available in packs of 100 and 500.

Basic NHS Cost 25 Paramol-118 tablets (dispensed from containers of 500) cost 14p.

Clinical experience shows that

Paramol-118

controls pain in dosage of only
one tablet every four hours



Made in England

Paramol is a Trade Mark of

DUNCAN, FLOCKHART & CO. LTD. LONDON E2 6LA

holder of product licence number 0021/5056

Full information is available on request.

MAR 1975

DF75/3/HN