

Duphalac : suspends the monthly sentence of dysmenorrhoea.

Contributors

Duphar Laboratories

Publication/Creation

1971.

Persistent URL

<https://wellcomecollection.org/works/yfuxtypm>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



A party of young men and women Hals (c. 1580 (?) - 1666)

August


September 1971

October

M	T	W	T	F	S	S	M	T	W	T	F	S	S	M	T	W	T	F	S	S	
						1			1	2	3	4	5						1	2	3
2	3	4	5	6	7	8	6	7	8	9	10	11	12	4	5	6	7	8	9	10	
9	10	11	12	13	14	15	13	14	15	16	17	18	19	11	12	13	14	15	16	17	
16	17	18	19	20	21	22	20	21	22	23	24	25	26	18	19	20	21	22	23	24	
23	24	25	26	27	28	29	27	28	29	30	25	26	27	28	29	30	31				
30	31																				

duphaston
 suspends the monthly sentence of dysmenorrhoea

National Gallery London



duphaston[®]

**puts back the days
that dysmenorrhoea
takes away**

For some women, dysmenorrhoea makes the monthly period a monthly sentence of misery, a few days lost from their normal lives.

Duphaston can suspend this monthly sentence. It is the only orally active non-contraceptive progestogen which protects the woman against the pain and distress of dysmenorrhoea. Duphaston is free from oestrogenic effect; it does not suppress ovulation or have any adverse hormonal effects, and it is extremely well tolerated. Clotting mechanism, hepatic function and basal temperature are unaffected.

Duphaston is supplied as tablets of 10 mg. dydrogesterone, in packs of 80 and 500. The usual dosage to control dysmenorrhoea is 10 mg. twice daily from day 5 to day 25 of the cycle. This may be reduced in subsequent cycles according to response.

Side-effects

Duphaston has no serious side-effects, and no greater incidence of minor effects than with placebo.

Contraindications

There are no known contraindications to the use of Duphaston.

Other indications

Besides its use in the prevention and treatment of dysmenorrhoea, Duphaston is also indicated in the management of threatened and habitual abortion, endometriosis, and infertility due to corpus luteum insufficiency.

Basic NHS cost

80 tablets, £6.70; 500 tablets, £34.80.

Further information available on request.

duphaston[®]

dydrogesterone

duphar DUPHAR LABORATORIES LIMITED BASINGSTOKE HANTS. TEL 0256 26351
PHILIPS-DUPHAR (IRELAND) LIMITED RAINSFORD STREET DUBLIN 8 TEL DUBLIN 754271 

**it is
important
to take
medicines
exactly as
prescribed**

M7 09 87870K