

Chilean girl gathering the leaves of the Boldo Tree : used for the phytotherapeutic treatment of constipation : Boldolaxine.

Contributors

W. Fletcher (Chemists) Ltd.
Mercier, A.

Publication/Creation

[approximately 1936]

Persistent URL

<https://wellcomecollection.org/works/dubv9het>

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).

**wellcome
collection**

Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



Chilean girl gathering the leaves of the BOLDO TREE

used for the PHYTOTHERAPEUTIC TREATMENT OF

CONSTIPATION

BOLDOLAXINE

BOLDOLAXINE

PHYTOTHERAPEUTIC TREATMENT OF CONSTIPATION

D E F I N I T I O N

BOLDOLAXINE derives its action principally from the fresh leaves of the BOLDO TREE (*Peumus Boldus*) a native Chilean Tree. For the last few years BOLDOLAXINE has been the object of important research work.

Physiological and clinical observations from tests carried out warrant our offering this product under the name of BOLDOLAXINE, an entirely new product in which full advantage has been taken of the laxative properties of this native of the Chilean Forest.

A C T I O N

It is not a purgative — it is a true physiological laxative ensuring natural stools. It greatly assists intestinal evacuation by its stimulating action on the liver and intestinal glands ; by its invigorating action on the muscle fibres and its equalizing power over the vegetative nervous system (large sympathetic and pneumo-gastric nerves).

I N D I C A T I O N S

Constipation in every form is relieved by BOLDOLAXINE with the assurance of a gentle effective action without colic. It is not habit forming and has a valuable diuretic action on the kidneys and a stimulating action on the liver.

I N S T R U C T I O N S F O R U S E

One tablet half an hour before each evening meal. (Children : — half a tablet). Two tablets are seldom necessary.

Sole Distributors : **W. FLETCHER (CHEMISTS) LTD.** 5 Rampayne Street, London S.W.I.
