Anatomical dialogues, or, A breviary of anatomy ...: chiefly compiled for the use of the young gentlemen in the navy and army / by a gentleman of the faculty.

Contributors

Gentleman of the faculty.

Publication/Creation

London: Printed for G. Robinson, 1778.

Persistent URL

https://wellcomecollection.org/works/m3s8pnxw

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.




Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org




19.0.3 AL ARMY MEDICAL. TRO RAMAC COLL. 1ANA

22101943607



HOYAL ARMY MEDICAL



ANATOMICAL DIALOGUES; my OR, A Dantsan BREVIARY OF ANATOMY. WHEREIN ALL THE PARTS OF THE HUMAN BODY

CONCISELY AND ACCURATELY DESCRIBED.

THEIR USES EXPLAINED;

BY WHICH

THE YOUNG PRACTITIONER

MAY ATTAIN A RIGHT METHOD OF

TREATING DISEASES, AS FAR AS IT DEPENDS ON ANATOMY.

CHIEFLY COMPILED

FOR THE USE OF THE YOUNG GENTLEMEN

THE NAVY AND ARMY.

BY A GENTLEMAN OF THE FACULTY.

LONDON:

Printed for G. ROBINSON, No. 25, PATERNOSTER-ROW. MDCCLXXVIII,

BIR MEN PROPERTY ON A PROPERTY OF THE PROPERTY AND PROPERTY AND THE . Privillion C. Routs, out, No. 27 Pa can work acon. TO WILLIAM HUNTER, M. D. PHYSICIAN TO HER MAJESTY,

FELLOW OF THE ROYAL COLLEGES

OF PHYSICIANS IN LONDON AND EDINBURGH,

OF

THE ROYAL SOCIETY,

ANDOF

THE SOCIETY OF ANTIQUARIANS,

THE FOLLOWING

ANATOMICAL DIALOGUES

ARE,

WITH THE HIGHEST ADMIRATION OF HIS ABILITIES,


AND WITH THE GREATEST RESPECT FOR HIS VIRTUES,

MOST HUMBLY INSCRIBED,

BY HIS MOST OBEDIENT

AND OBLIGED PUPIL,

THE EDITOR.



COLLEGE LIBRARY

PREFACE.

S the Public will undoubtedly expect I some Reason to be given for intruding on them another Work of this Kind, (when fo many learned and judicious Authors have professedly written on the Subject) I think it necessary to inform them, "That it is neither from Vanity nor Prospect of Gain, that the following Sheets were compiled, but merely to render the Study of Anatomy more agreeable, and to be obtained at a less Price than any System of Anatomy yet extant." Was Ambition the Motive, I should not conceal my Name; and was Profit the Inducement, I should then have made it more voluminous and fystematical: However, I hope I have omitted no effential Part of the Science necessary to be known by the young Practitioner, in order to attain a right Method of treating Diseases, as far as it depends on Anatomy; but how I have fucceeded. I must leave to the impartial and judicious Reader to determine.

This Book was compiled with a view to facilitate the Knowledge of Anatomy to fuch young Gentlemen as are intended for the Service vice of the Army and Navy, as well as domeftic Practice.

A Compendium, containing all the Disco. veries of the modern Anatomists, it is prefumed has long been wanted, as a Pocket-Companion for the young Student, previous to his Examination at Surgeon's Hall, in which he may see at one View, a concise Description of the Parts which compose the human Body, without distracting his Memory by the Perusal of a Number of Books, which often contain physiological Explanations very contradictory to each other.

I have thrown the whole into the Form of Dialogue, which I flatter myself will not only be more pleasing to the younger Part of the Profession, (for whom this Work is principally designed) but make a greater Impression on their Memory—Instruction being most impressive when least incumbered.

Should this Compendium be censured or condemned, by the self-sufficient and ill-natured Part of the Faculty, I doubt not but those of Learning and Liberality of Sentiment, who are an Honour to the Profession, will overlook with Candour any trisling Defects which

which may be found in it. Though this Performance is not by any means intended for the Use of the elder and more experienced Artists; yet even to them, it may supply the Place of a Remembrancer, of what they were before more fully informed. I am not so vain as to think it will improve their Knowledge, since nothing new can reasonably be expected in a Work of this Kind, which is chiefly collected from other Authors: But if it answers the End designed, of assisting the young Student, I shall not think my Labour and Attention in compiling it, though it has cost me much Trouble, ill bestowed.

To render the Book more useful, I have added a copious Index, by which the Reader may readily advert to any particular Subject he may want to be informed of.

London, July, 1778.

CONTENTS.

		The same
DIALOGUE	I. Of Anatomy in General	Page
	II. Of the Head and its Parts	50
	III. Of the Neck and its Parts	110
	IV. Of the Thorax and its Parts	132
	V. Of the Abdomen and its Part	s 177
	VI. Of the Arm and its Parts	342
1 ,4:	VII. Of the Leg and its Parts	285
	VIII. Of the Parts of Genera- tion in both Sexes	325
lat Subject	IX. Of the Embryo, and the Fœtus in Utero	346
	X. Of the Eye	354
	XI. Of the Ear	363

ROYAL ARMY MEDICAL COLLEGE LIBRARY.

ANATOMICAL

DIALOGUES.

DIALOGUE I.

Of Anatomy in general.

A. Anatomy is the accurate dissection, or the cutting to pieces of a human body, in order to discover its structure and uses; in which every physician and surgeon ought to be well skilled, exactly to know, and fully to understand every part, and its proper use.

Q. What is the use of anatomy?

A. It gives you the perfect knowledge of the cause and cure of diseases; without which, neither the physician nor the surgeon can do justice in their professions, but would rather be detrimental, than beneficial to mankind.

Q. How is the body distinguished by anatomists?

A. Into venters and limbs. The venters are the more remarkable cavities of the body, in which some principal part or bowel is contained, viz. the belly, breaft, and head: the limbs are, the arms and legs; the cavity of the breast is called thorax, and that of the belly, abdomen; their sub-contained parts are named as follows, viz. the hollow on the middle of the thorax, under the breafts, is called fcrobiculus cordis; the middle of the abdomen, for about three fingers breadth above and below the navel, is called regio umbilicalis; the middle part above this, epigastrium; under the cartilages of the lower ribs, hypochondrium; and from below the regio umbilicalis to the offa ilia and offa pubis, hypogaftrium.

Q. What are the external parts, and com-

mon integuments?

A. These are, the cuticula, epidermis, or fearf-skin; the corpus reticulare, vel reticulum mucosum; the cutis, or true skin; the glandulæ subaceæ, vel miliares; the membrana adiposa; and the pinguedo, or fat.

Q. What is the cuticula, epidermis, or

feart fkin?

A. It is a very fine, thin, smooth, insensible membrane, closely lying upon the cutis, or true skin, of which it seems a part, and is what rises into a bladder in burns, or when a blister is applied: the colour of it in Europeans is white, but black in many other nations: hav-

ing no blood-vessels or nerves, it is void of sensation.

Q. What is the use of the cuticle?

A. To defend the true skin from injury, dryness, and pain; and to assist, and, at the same time, to moderate the sense of feeling. This membrane appears scaly and porous (through a good microscope); and Lewenhoeck and others say, every scale has about 500 pores; so that a grain of sand (according to this account) will cover 125,000 pores through which we perspire.

Q. What is the corpus reticulare, vel reti-

culum mucofum?

A. A fine net-like membrane, lying immediately under the cuticle, and adhering firmly to it: from its colour, and the colour of the mucus it contains, the skin appears to be either black or white: in Europeans, white; in Africans, black; in the tawny, yellowish.

Q. What are the uses of the corpus reticu-

lare?

A. To preferve the structure of the other parts of the integuments, to give passage to the hairs, papillæ, and excretory ducts, and retain them in their places; and it has some share in preserving the softness of the papillæ, which renders them fit for the sense of feeling.

Q. What is the cutis, or true skin?

A. The true skin is a very compact, thick, strong, and sensible membrane covering the whole body, and so plentifully supplied with nerves for the sense of touching, that the very

B 2

fineft

finest pointed instrument can prick no where, without touching some of them. Immediately above this membrane, is the corpus reticulare and cuticula; and under it, the membrana adiposa, and fat. Its thickness is very different in feveral parts of the body. It is composed of a multitude of tendinous fibres, a vast number of blood-vessels and nerves, which constitute rhe pyramidal papillæ that raise themselves through the pores of the corpus reticulare, and constitute the organs of feeling. The true skin is white in all mankind. Its appearing white, black, or tawny (according to the climate) is owing to the colour of the cuticula and corpus reticulare, and not to the colour of the skin, which is always white in all nations. The red colour of the lips is owing to the blood-veffels in the muscular flesh; and that of the cheeks in white people, to the blood in the minute vessels of the skin.

Q. What are the uses of the skin?

A. To defend the parts underneath from external accidents, to be the organ of feeling, to wrap the parts more firmly together, and to be an universal emunctory to the body, cleansing the blood of its redundancies by sweat and perspiration, which, at the same time, prevents its flaccidity or dryness.

Q. What are the glandulæ sebaceæ?

A. The glandulæ sebaceæ, vel miliares, are small bodies like millet-seeds, seated immediately under the skin in the axillas; and are said to have been found under all other parts of

of the skin when looked for with good microscopes.

Q. What are the uses of the glandulæ fe-

baceæ, vel miliares?

A. These glands are supposed to separate fweat; which fluid was thought to be only the materia perspirabilis, flowing in a greater quantity, and condensed, till Sanctorius affured us that it is not so; and that more of the materia perspirabilis is separated in equal times, than of sweat: of the former he says, there are ufually fifty-two ounces a-day feparated (in Italy, where his experiments were made), but of the latter, not near fo much in the most profuse sweats. But whoever reads Mr. Hales's experiments, will find what Sanctorius accounted for by an imaginary, infensible perspiration, different from that which in the greatest degree produces fweat, is really made by the lungs in respiration, in ten times a greater quantity than all the ordinary perspiration through the cutis, and feems to be but the fame kind of fluid discharged both ways; for whenever it is interrupted through the skin in cold weather, then the lungs are overcharged, and a cough is produced.

Q. What is the membrana adipofa?

A. All that membrane immediately under the skin, containing the fat, in numerous cells, by some named cellulæ, and the substance made up of them, termed the cellular substance: this membrane adheres very closely to the skin, and runs between the muscles in ge-

B 3

neral,

neral, and between their feveral fibres in particular; and communicates with the membrane which lines the infide of the thorax and abdomen; and all its cells communicate throughout the whole body, so that from any one part the whole may be filled with air, as is evident in beafts, by the butcher's blowing up their meat when newly killed; and in an emphysema of the human body, sometimes occafioned by a broken rib, &c. And in these cells the water is contained in an anafarca, which, from its weight, first fills the depending parts; and when these cells are very full, the water frequently passes from them into the abdomen, and, after tapping, though the limbs were ever fo full, they will almost empty themfelves in one night's time. This membrane is also the usual seat of imposthumations, carbuncles, and boils.

Q. What are the uses of the membrana

adipofa?

A. It contributes to keep the inner parts warm and pliant; and by filling the interflices of the muscles, renders the surface of the body smooth.

Q. What is the pinguedo, or fat?

A. It is an oleaginous or butyraceous matter fecreted from the blood (in the little arteries of the adipose membrane) and filling up the cavity of the cells of the membrana adiposa: and that this oleaginous matter has a circulatory motion, or an egress into the veins, is very evident, from the judden consumption

of it in many diseases, and from its vast diminution by exercise and labour:

Q. What are the uses of the fat?

A. To serve as a kind of covering to the body, in order to defend it from cold, and other injuries; to defend the more tender and fensible parts from being too strongly vellicatedby the falts; to preferve, in good order, the flexion of the muscles and cutis, and of the other parts between, and about which it is placed; to facilitate the motions of some parts, as the eyes, jaws, &c. to fill up the intersticial spaces, and by that means to add to the beauty of the parts, as is evident in the face, neck, &c. and to facilitate the diffension of the parts, the spaces between which it thus fills up. There is a manifest use of the fat about the vagina, anus, offa ischia, and pudenda; in the exclusion of the fœtus, and the harder excrements; and in the foles of the feet, the nates, and other the like parts, in all which the fat is very copiously disposed, and ferves in the place of a cushion for the muscular flesh to rest on, and to prevent the painful pressure and attrition of the parts: and finally, there is great reason to suppose, that when the body does not receive nourishment in the usual way, the regress of the fat into the veins supplies that defect.

Q. What are the constituent parts of the

body?

A. The body consists of bones, cartilages, ligaments, muscles, tendons, arteries, veins, nerves,

nerves, lymphæducts, glands, excretory vessels, membranes, fibres, hair and nails, besides the integuments before mentioned.

Q. What are the bones?

A. The bones are hard, brittle, insensible parts (but covered both on their infide and outfide with an exquifitely fenfible, nervous, vafcular membrane, called periosteum, except on the skull, where it is called pericranium) containing more or less marrow: and on the surface of the bones, at the ends, are two kinds of prominences; the one termed apophysis, or processus, and the other epiphysis, or appendage. Befides the common large cavity, most bones have superficial cavities or sinuses, with furrows and holes through which the nutritious and medullary veffels enter (as all the membranes of the bones, both within and without, are supplied with blood-vessels and nerves, as is the marrow). The most considerable of the nutritious vessels enter at the ends of the bones, viz. the artery at one end, and the veins at the other. If the bones had no cavities, they would, if they were strait, sustain the same weight; but being made hollow, their strength to relift breaking transversely is increased in proportion to their diameters, without increafing their weight, which is very evident in the wings and quills of birds: but the bones in the legs of all animals are more folid, being formed to support weight.

Q. What is the apophysis of a bone, and

its ule?

A. The apophysis is a continued part, or excrescence of the bone (as a branch is of a tree) jutting out from it, to make the better articulations, and for the more commodious insertion of the muscles.

Q. What is the epiphysis of a bone, and

its use?

A. The epiphysis is an additional bone growing to another by mere contiguity, being generally more soft and porous than the other, though it mostly degenerates into an apophysis in adults, and therefore of the same use.

Q. What is the medulla, or marrow, con-

tained in the bones?

A. It consists (besides the blood-vessels) of an investing membrane, in which are included membranaceous lobules and bags that fill up the cells of the bones; and in these bags are vesiculæ, or glandulous bladders, very like the vesicular substance of the lungs. The large middle cavity of all cylindrical bones, contains an oily marrow; but the great number of leffer cells, towards their extremities, contain a bloody marrow, or rather a kind of red, fatty, medullary juice; which last is found in all fpongy heads and cells of bones. The marrow in young bones is more red and bloody than in old ones, as the oily marrow would otherwise render their fibres too foft. The medullary veffels, found running here and there, through their proper canals, penetrate into the inner cavity of the bones, and fecrete the the medullary part from the blood, the blood being afterwards returned by the veins: the nerves are distributed to the same places, for the sake of sense and motion: these vessels enter the bones obliquely, that they may not weaken them by dividing too many fibres in the same place.

Q. What is the use of the marrow?

A. The marrow being more or less distributed all over the bones, and transuding through their plates and fibres, makes them tougher and less brittle; but does not nourish them, as was originally believed.

Q. What is the periofteum?

A. An exquisite, sensible, nervous, vascular membrane, which lines and covers all the bones in the body, internally and externally, except a part of the teeth, and the places in bones where the muscles are inferted. The periofteum on the outfide of the bones, is derived from the membranes of the muscles that lie upon it; that on the infide, from the dura mater. The inner superficies of the periosteum flick as close to the bone, as if it were glued to it; and besides, the periosteum has little fibrillæ or threads continued from it, that enter into the substance of the bone, which give them, probably, some internal sense. The periosteum constitutes the first rudiments of the bones in a fœtus in utero. It is every where full of small blood-vessels, which enter the bones for their nourishment; but the internal substance of the larger bones is nourished by the

the vessels which enter obliquely through their middles. The periosteum is of different thickness in different parts; but, in general, the internal is vastly thinner than the external; and though some have supposed it to arise from the dura mater, yet it is evidently formed, at the same time, with the dura mater in the fœtus.

A. It gives fensibility to the bones, which otherwise might be sawed, cut, or burnt, without pain, as their whole sensibility is owing to this membrane: and it also gives the determination and figure to bones; as is evident from this, that when it is wounded, exostoses, tophi, and caries, arise in the part: it is likewise the organ of secretion for the bony matter, as the membrana adiposa is for the fat: and it serves also for the muscles to slide easy upon, and to hinder them from being lacerated by the roughness and hardness of the bones.

Q. What is the substance of the bones?

A. The substance of the bones consists of lamellæ or plates, lying one upon the other, joined together by transverse fibres, and, as it were, archwise: these fibres, when first formed, are very soft, but grow by degrees to the hardness of a cartilage, and afterwards to that of a perfect bone. But the change is neither made in a very short time, nor begun in all the parts of the same bone at once.

Q. What are the uses of the bones?

A. They give strength and shape to the whole body, sustain all its organs, and keep

the machine in proper order to perform its various functions; therefore the exact know-ledge of the bones is the foundation of all anatomy. The bones are connected together various ways, according to the various purposes they are to serve; some being intended for motion, others for rest, and the support of the incumbent parts only.

Q. How many bones are there in the hu-

man body?

A. Two hundred and forty-seven (exclusive of the sesamoidal bones, whose number is uncertain); but the number is various in various subjects; some say 300, or 307; others, 318, but late writers six it at most to 250, and commonly but 247, as follows.

A. The findfance of the cones confile of

Sardyonether by maniveric horrs, and, as in

in a very thort time, not began in all the mark

A TABLE of OSTEOLOGY.

	-		1	10
9.00			Os Parietalia	2.
[Æthmoides	1-
-5	Th	e bones of the crani-	Frentis	I-
2	1 111	m, or skull	Occipitis.	1-
2	-	m, or man	Sphenoides	1 -
2.			Temporum	2-
	A Total Control of the		Os Malarum	2 -
re		State of the state	Maxillaria	2 -
ca			Nafi	
ea		e bones of the fauces	¿ Plana	2
,u	1 2	and face	Palati	2-
th	1		Vomer	1-
The bones of the head are	!		[Ungues	2-
63	T	ne bone of the lower	Os Maxilla Inferior	1-
one		jaw	1 03 IVANATITA THICHOI	228 65
2	1 .		Dent. Canini	4-
The state of	D	entes, the teeth	3 Incifores	8-
F.	1		(Molares	20-
	T	he bones of the os hy-	S Os Hyoides	5-
	oides of the tongue			2
	1	COs Incus		
	T	he bones of the internal	Malleus	2
	1	ear	Orbiculare	2 *
1 Si			L Stapes	2
			No	67
			The state of the s	07
			(Neck	7
1	1	V	ertebræ { Back	13
tru		The spine	Loins	
(Sternum 3				
Line of 1 mil - thouses J		The standard of	ibs	5 2 3 24
5 i C 0s		COs Illian	n I	
nc	;	The pubes	Offa Ischiu	ım I
2	1	The Paoes	Inominata Pubis	1
2	111			
-	4			

14 ANATOMICAL

TABLE continued.

. 6		Ne
es.	The shoulder { Clavicula	2
The bones of the upper extremities	Scapula Scapula	2
rem	The a m Os Humeri	2
Xt	The fore arm { Ulna	2
1	Radius Os Naviculare	2
bbe	CLunare (Lunare	2 2
0	Cuneiforme	2 *
the	Magnum	2
Jo	Carpus Crbiculare	2
60	Trapezium	2
OH	The hand Trapezoides	2
e p	Metacarpus	2
Lh	L Phalanges	8
12	the ways	30
60	C SHANKE STATE OF THE STATE OF	
bones of the lower extremities.	Thatlish O. F. L.	
em	The thigh—Os Femoris Os Patella	2
xtr	The leg { Tibia	2
re	L Fibula	2 2
We	, COs Calcis	2
	Aftragalus	2
th	Tarius Cuboides	2
of	The foot \ Cuneiformia	6
nes	Metatarius Naviculate	2
Pol	Phalanges	10
The	and the second	28
FI	ALL DESCRIPTION OF THE PROPERTY OF THE PROPERT	In all 247
2	most of the second second second second	Q. What
	The state of the s	· · · · · · · · · · · · · · · · · · ·

Q. What are cartilages?

A. Cartilages approach much to the nature of bones, but are smooth, lubricous, slexible, and elastic. There is no sensible cavity, cell, or pore, in any part of the substance of those which cover the ends of bones, except very small passages for the blood-vessels, &c. They are whitish, or pearl-coloured; and with age, sometimes grow so hard, as to become perfect bone.

Q. What are the uses of the cartilages?

A. Their use is to cover the ends of the bones that have motion, to prevent them from being damaged by a continual friction; and to contribute, in a great measure, to the formation of several parts, as the larynx, nose, ears, &c.

Q. What are ligaments?

A. Ligaments are white, fibrous, close, compact substances, more flexible than cartilages, not easily ruptured or torn, and do not yield, or at least but very little, when pulled. They form either narrow cords, broad bands, or thin webs. In all the moveable joints there are capsular ligaments; so called, from a very fluid, mucilaginous liquor (termed synovia) being contained in capsulæ, or bags; which liquor continually moistens the articulations. It is generally agreed, that the ligaments are infensible; but experience shews, that they are capable of very acute pains, nothing being more painful than collections of matter within these

these parts, or sharp medicines applied to them when laid bare.

Q. What are the uses of the ligaments?

A. Their use is either to bind down the tendons, or to tie together such bones as have motion, to prevent them from parting from each other, as happens in luxations.

Q. What are muscles?

A. Muscles are distinct portions of flesh, foft and red; compounded of fibres, tendons, nerves, veins, and arteries, all enclosed in a peculiar membrane: every muscle has one or more tendons at its beginning and end, though fometimes fo very small, as not to be feen without a microscope. Where the nerve is inserted, is the head or beginning of the muscle; the middle, or fleshy part, is its belly; and where the tendons are greatest, and most in number, is called the tail or end; and if the end be broad and membranous, it is called aponeurosis; but in many of the muscles both the extremities are moveable: in those that part, which of the two is least moveable, is generally called the origin, or head; though in the ferrati antici, and some of the muscles of the abdomen, it is the reverse. All the muscles are either rectilinear or penniform, and formed for flexion or extension, and sometimes both. To every flexor is opposed an extensor; to every adductor, an abductor; to every elevator, a depreffor, excepting only in the sphineters, cremasters, and cesophagus. Some muscles are mostly carnous, as all the sphincters and muscles

muscles of the tongue; others, mostly nervous and membranous, as the fascia lata tibia adducens, quadratus vel distortor oris; by Galen named platysma myoides, being first contracted involuntarily in the spasmus cynicus.

Q. What are the uses of the muscles?

A. To promote voluntary motion, as by the contraction of the muscles the several motions of the body are performed. Thus, the foul effects motion in the muscle at pleasure, by transmitting from the brain by the nerves, animal fpirits, by whose copious influx, the muscles to be moved are in an instant inflated, and so contracted, that they pull and move the member to which the tail is fastened: to this end, nature hath inferted a nerve or more into each muscle; so that flesh and nerves are the principal constituents of a muscle; and indeed, some muscles seem to have their bodies made up of nothing else than an intertexture of nervous and fleshy fibres; as the muscles of the forehead, temples, bladder, anus, and all sphincters. The motions of the far greater part of the muscles are voluntary, or dependent on our will; those of a few others involuntary: those which perform the voluntary motions, receive nerves from the brain or ipinal marrow; those which perform their motions involuntarily, have their nerves from the cerebellum; and those whose motion is partly voluntary, and partly involuntary, have theirs in part from the brain, and in part from the cerecerebellum. And as a muscle can no longer act when its nerve is either cut asunder, or tied up, so nearly the same absolute dependance it has on its artery; for on cutting, or tying up the artery, the muscle in the same manner, some hours after, loses its whole power of action, as if the nerve had been cut or tied up.

Q What are tendons?

A. Tendons are white, firm, tenacious parts, composed of the same fibres with the muscles (which see). When the fibres of a tendon expand themselves into a membrane, it is called aponeurosis.

Q. What are the uses of tendons?

A. The same as that of the muscles.

Q. What are arteries?

A. Arteries are conical tubes, which arife from the ventricles of the heart; and thence dividing into branches, are distributed to all parts of the body. They are composed of three membranes, or coats. The external and internal are membranous, but the middle coat is rather muscular; consisting of circular or spiral fibres, which being very elastic, contract themselves with some force, when the power ceases by which they have been stretched out. They have two reciprocal motions, or pulsations, like the pulses of the heart; being a systole and a diastole, keeping opposite times; the systole of the one, answering to the diastole of the other.

Q. What is the use of the arteries?

A. To

the

A. To convey the blood from the heart to all parts of the body.

Q. What are the principal arteries of the

human body?

A. The aorta vel arteria magna, and the arteria pulmonalis: all the other arteries of the body, though diftinguished by particular names, are only branches of these two.

Q. What is the aorta, and how distributed?

A. The aorta vel arteria magna is a large artery which comes out from the left ventricle of the heart in a fingle trunk, above its valves, called femilunares vel figmoides; from this all the other arteries proceed, and by which the whole mass of blood is conveyed to all parts of the body. The aorta is by anatomists generally divided into the aorta afcendens and aorta descendens, though both are but one and the same trunk. It is termed ascendens, from where it leaves the heart to the extremity of the great curvature, or arch. The descendens is that part of the trunk which, after the archlike inflection descends through the thorax and abdomen, down to the os facrum; and is ufually larger in women than in men.

Q. How is the aorta ascendens distributed?

A. The aorta, before it perforates the pericardium, affords to the heart itself the arteriæ coronariæ; and then passing the pericardium, it is termed aorta ascendens; when, after ascending two or three inches upwards, its trunk is bent in manner of an arch, from which arises three ascending branches that form

the carotid and fubclavian arteries. The right carotid and fubclavian proceed first in one trunk, but the left carotid and fubclavian immediately fingle; the left carotid forming the middle branch. From the two fubclavian branches (while yet within the breast) near the uppermost rib, proceeds, 1. arteria intercostalis superior, proper to the four upper ribs; 2. arteria mammaria, proper to the breafts; 3. cervicalis, proper to the muscles of the neck and head, and by communication, partly to the brain; 4. carotis, the external, proper to the larynx, tongue, neck, head, and brain; the internal, chiefly to the brain. When the fubclavian branches have left the cavity of the thorax, they are termed axillares, which are distributed to the outer parts of the breast and arms by thoracica superior & inferior; 5. scapularis; 6. humeralis; then they approach the arm, where they lie under the branches of the axillary vein, and pass to all parts of the arm, bearing the same names with the veins that accompany them.

Q. How is the aorta descendens distri-

buted?

A. The aorta being reflected under the left lobe of the lungs, it commences aorta descendens; which name it keeps through the thorax and abdomen, where it passes on the left side of the spine, till its division into iliac arteries, between the third and fourth vertebræ of the loins. This descendent trunk, which is the greatest, being yet within the capacity of the thorax,

thorax, fends, 1. intercostalis inferior, to the eight lower ribs; 2. bronchiales, to the lungs; 3. phrenicæ, to the diaphragm; 4. cæliaca, whose branches are bestowed upon the liver, pancreas, spleen, stomach, omentum, and duodenum; which are named from the parts they are bestowed on, except two upon the stomach, which are called coronaria ventriculi, fuperior et inferior; and one upon the duodenum, named intestinalis; 5. mesenterica superior, whose branches are bestowed upon all the intestinum jejunum, and ileum, part of the colon, and fometimes one branch upon the liver; 6. emulgentes, to the kidnies; 7. spermaticæ, to the peritonæum, ureters, testicles, and epidydimes; 8. lumbares, to the loins; 9. melenterica inferior, to the lower part of the colon, and the rectum; 10. muscula superior, to the muscles of the belly. As soon as the aorta divides upon the loins, it sends off an artery into the pelvis, upon the os facrum, called arteria facra; and the branches the aorta divides into, are called iliacæ, which, in about two inches space, divide into external and internal. The iliacæ internæ, fend, 1. arteria inferior, to the muscles; 2. umbilicalis, which are collapsed in adult bodies, except at their beginnings, which are kept open for the collateral branches on each fide, one to the bladder, and one to the penis or uterus; 3. hypogastrica: the rest of the branches of the interhal iliac are bestowed upon the buttocks, and upper parts of the thighs. The iliacæ externæ

næ run over the offa pubis into the thighs, sending off, 1. epigastricæ, to the fore-part of the integuments of the abdomen under the recti muscles into the pelvis, and also through the foramina of the offa innominata, to the muscles of those parts; 2. inguinalis, to parts of the groin; 3. cruralis, to the thigh; 4. poplitea, to the ham; 5. tibialis antica medica, et postica, which supply the leg, foot, and toes. Thus you have a description of all the large and finall capital branches of the aorta, which are for the most part disposed in pairs, and are uniform in most bodies; but the lesser branches are distributed, like the branches of trees, in so different a manner in one body from another, that it is highly probable no two bodies are exactly alike, nor the two fides in any one body.

Q. How is the arteria pulmonaris diftri-

buted ?

A. Only through the lungs, but with a vaft number of ramifications. It arises from the right ventricle of the heart, and soon divides into two branches, one to each lobe of the lungs; then they are subdivided into smaller and smaller branches, until they are distributed through every part of the lungs. The extreme branches, both of the arteries and veins, have very numerous communications, like those in the stamina of the leaves of plants; by which communications, the blood that is obstructed in any particular vessel, may pass off by other vessels that are not obstructed, &c. and as many

many of the leffer veffels are more exposed to pressure than any of the large ones, those communications in the leffer veffels are therefore made more numerous. By fuch communications, the blood circulates in a limb that has had part amputated, and the fluids contained in a large inflammation, suppurates into one cavity. It is computed, that each ventricle of the heart holds five ounces of blood (and they are filled and emptied every fystole and diaftole) and that there are commonly eighty pulses in a minute: if so, there then flows twentyfive pounds of blood through each ventricle of the heart in a minute. Dr. Keil has shewn, that the fum of all the fluids in a man exceed the fum of all the folids; and yet the quantity of blood which all the visible arteries of a man will contain, is less than four pounds; and if we may suppose all the visible veins, including the vena portæ, hold four times as much, the whole, then, that the visible vessels contain, is not twenty pounds; but the whole that they do contain, is but very little more than the veins can contain, seeing the arteries are always found almost empty in dead bodies. How much the invisible arteries and veins contain, however (I mean those which contain fuch a compound fluid as is found in the larger veffels) there is no way to judge, unless we knew what proportion these vessels bear to those that carry the nutritious juices and serum (if there are fuch) without the globuli of the blood. It is probable, that in all animals the velocity

velocity of the blood, as well as the necessity of food, is, cæteris paribus, in proportion to their quantity of action: if fo, it appears how those animals which use no exercise, and whose blood moves extremely flow in the winter, can fubfift without any fresh supply of food; while others that use a little more exercise, require a little more food; and those who use equal exercife winter and fummer, require equal quantities of food at all times; the end of eating and drinking being to repair, what exercise and the motion of the blood have destroyed, or made useless: and is not the less velocity of the blood in some animals than in others, the reason why wounds and bruises in those animals do not so soon destroy life, as they do in animals whose blood moves swifter?

Q. What occasions the pulse?

A. When the left ventricle of the heart contracts, and throws its blood into the aorta. or great artery, the blood in the artery is not only thrust forward towards the extremities. but the channel of the artery is likewise dilated; because fluids, when they are pressed, press again to all sides, and their pressure is always perpendicular to the fide of the containing veffels: but the coats of the artery by any small impetus may be distended; therefore upon the contraction of the heart, the blood from the left ventricle will not only press the blood in the artery forwards, but both together will diftend the fides of the artery. When the impetus of the blood against the

the fides of the artery ceases, that is, when the left ventricle ceases to contract, the spiral fibres of the artery, by their natural elasticity, return again to their former state, and contract the channel of the artery till it is again dilated by the systole of the heart. This diaftole of the artery is its pulse, and the time the spiral fibres are returning to their natural state. is the distance between two pulses. The pulse is in all the arteries of the body at the same time; for whilst the blood is thrust out of the heart into the artery, the artery being full, the blood must move in all the arteries at the fame time: and because the arteries are conical, and the blood moves from the basis of the cone to the apex, therefore the blood must strike against the sides of the vessels, and confequently every part of the artery must be dilated at the fame time that the blood is thrown out of the left ventricle of the heart; and as foon as the elafticity of the spiral fibres can overcome the impetus of the blood, the arteries are again contracted. Thus these are two causes, which operate alternately, and keep the blood in a continual motion, viz. the heart, and fibres of the arteries; but because the one is stronger than the other, therefore, though the blood runs continually, yet when the artery is opened, it is feen to move per faltum.

Q. What are veins?

A. Veins are tubes or veffels which carry the blood from all parts of the body to the heart. They are composed principally of a membra-

membranous, a vafcular, and a mufcular tunic, the same with those of the arteries, only the muscular coat is much thinner; as thin in all the veins as it is in the capillary arteries; the pressure of the blood against the sides of the veins being less than that against the sides of the arteries. The veins are only a continuation of the extreme capillary arteries, reflected back again towards the heart, which, uniting their channels as they approach it, at last form three large primary veins, viz. the vena cava, vena portæ, and the vena pulmonaris. In the veins there is no pulse, because the blood is thrown into them with a continued ftream, and because it moves from a narrow channel to a wider. The capillary veins unite with one another, as I have faid of the capil. lary arteries. The veins accompany the arteries in almost every part of the body, and have the same names in the several parts with the arteries which they accompany. The veins of the limbs are more than double the number of the arteries, there being one on each fide each artery, even to the smallest branches that we can trace, besides the veins which lie immediately under the skin. The external veins have frequent communications with the internal, and are always fullest when we use the most exercise; because the blood being expanded by the heat which exercise produces, it requires the vessels to be distended; and the inner veffels being compressed by the actions of the muscles, they cannot dilate enough; but

but these vessels being seated on the outsides of the muscles, are capable of being much dilated. In all the veins which are perpendicular to the horizon, excepting those of the uterus, and the vena portæ, there are small membranes or valves; fometimes there is only one, fometimes there are two, and fometimes three, placed together, like so many half thimbles stuck to the sides of the veins, with their mouths towards the heart. In the motion of the blood towards the heart, they are preffed closely to the fides of the veins; but if the blood should fall back, it must repel the valves, fo that no blood can repais them, or return towards the extreme parts from whence it came. The blood distributed to all parts of the body by the two arteries, the aorta and pulmonary, is returned by the three veins called vena cava, vena portæ, and pulmonary vein. The vena cava carries back to the right auricle of the heart the blood conveyed by the aorta to all the parts of the body, except what goes by the coronary arteries of the heart. It receives all this blood from the arterial ramifications in part directly, and in part indirectly. The vena portæ receives the blood carried to the floating viscera of the abdomen (by the cæliac artery, and the two mesenteric arteries) and conveys it to the hepatic vein, and from thence to the vena cava. The pulmonary vein conveys to the pulmonary finus, or left auricle of the heart, the blood carried to the lungs by the pulmonary artery. Q. How Q. How is the vena cava distributed?

A. The vena cava arises with a larger sinus from the right auricle of the heart, where it first sends out a vein to the heart itself, called the coronary vein, and is then divided into two trunks; one running upward, called vena cava descendens vel superior (because the blood descends through it to the heart) and the other running downward, called vena cava afcendens vel inferior (because the blood ascends through it to the heart) but the antients, respecting the direction of the tubes or veffels only, called them the reverfe, viz. the fuperior ascendens, and the inferior descendens. The moderns name them as here mentioned. according to the flowing of the blood through them. The descending, or superior trunk, is distributed chiefly to the thorax, head, and upper extremities, and but very little to the parts below the diaphragm. The afcending, or inferior trunk, is distributed chiefly to the abdomen and lower extremities, and but very little to the parts above the diaphragm.

Q. How the vena cava superior (called

alfo vena cava descendens) distributed?

A. The vena cava superior (called also vena cava descendens) a little higher than the aorta, as high as the cartilage of the first true rib, terminates by a bifurcation or division into the right and left subclavian branches, which are of unequal lengths, because the trunk of the vena cava lies more towards the right side, where the left subclavian arises as well

well as the right, and is consequently longest. From the heart to the bifurcation of the trunk of the superior cava, arise the following veins, viz. vena (azygos) fine pari, which fend branches to the eight lower ribs, and another to the left emulgent vein; 2. intercostalis superior, one on each fide, which afford branches to the four upper ribs (if the azygos doth not bestow branches on all the interstices of the costæ) the remarkable veins and arteries as well as nerves relating to the coftæ, are couched in a furrow on the under edge of each rib, where they fafely pass; 3. bronchialis, which accompanies the bronchial artery to return the blood to the cava. This vein in some subjects indeed, does not rife separate, but comes from the azygos, and fometimes from the intercostals, and in some is altogether wanting. After fending off some small branches, the cava paffeth to the clavicle, where it divides itself (as before mentioned) into a right and left subclavian branch, then sends off some fmall branches to the muscles of the neck and upper part of the breast; and these four capital branches, viz. jugularis externa et interna, vertebralis, and axillaris; but this last is rather a continuation than a branch of the fubclavian; 1. jugularis externa, chiefly to the external parts of the throat, neck, and head; and its branches receives names from the parts they are distributed to, as frontalis, temporalis, occipitalis, &c. 2. jugularis interna, to the internal parts of the head and neck, which gives

gives ramifications to the larynx, pharynx, muscles of the os hyoides and tongue; and befides these, its truk terminates in a fack, and brings back the blood from the brain and finules of the dura mater; 2. vertebralis, which afcends to the cranium through the transverse apophyses of the vertebræ of the neck; and with the cervicalis (which is generally a branch of this vein) supplies the muscles of the neck, accompanying the vertebral artery through all the transverse apophyses of the vertebræ colli, all the way to the great foramen occipitale, communicating with the occipital veins, and fmall occipital finuses of the dura mater. The fubclavian branch going out of the thorax on each fide, is termed, 4. axillaris, which is rather a continuation of the fubclavian, than a distinct branch : before it leaves the thorax, it fends off venæ thoracicæ, which are fpent on the muscles of the thorax. From the axillaris (after it leaves the capacity of the thorax) branches are fent off to the external and internal muscles of the scapula, &c. and to the axillary glands; and having reached the fide of the head of the os humeri, it fends forth cephalica to the arm (which creeps along between the fkin and the muscles): the axillaris then runs along the arm by the name of vena basilica; but both this, and the cephalica, may be looked upon as two principal branches of the axillaris; both are distributed, by numerous ramifications, to all parts of the arm, fore-arm, and hand. At the bend of the elbow

bow (or flexure of the cubit) they form three capital branches; the uppermost is called cephalica; the middle, mediana, and the next, basilica. The vein which runs over the back of the hand, towards the little finger, is called the salvatella.

Q. How is the vena cava inferior (called

also vena cava ascendens) distributed?

A. The vena cava inferior vel ascendens. is remarkable for its valves, which ferve to prevent the blood from returning towards the extremities: it is distributed thus, I. immediately after it passes out of the pericardium, it perforates the diaphragm, to which it gives the venæ diaphragmaticæ inferiores, or phrenicæ; 2. hepaticæ, to the liver; 3. renales vel emulgentes, to the kidneys; 4. spermaticæ, to the testicles; 5. lumbares, to the loins and their vertebræ: after this, the trunk having reached the os facrum, it there lofes the name of cava; and terminating by a bifurcation like that of the descending aorta, it sends off the vena facra to accompany the artery of the same name, and then divides into the two venæ iliacæ, each of which is divided into two large trunks, or fecondary iliac veins: this fecond bifurcation is about a finger's breadth below that of the iliac arteries. The original iliacs are distinguished into the right and left; and the secondary iliacs (which are four trunks) are named external and internal. or anterior and posterior. The two external trunks are also named simply iliaca, and the

two internal, hypogastricæ; the former seems to be a true continuation of the original iliac trunk, but the latter only a branch, I mean in adult bodies, for in a fœtus there is a manifest difference. From the hypogastrica arise the musculares, hæmorrhoidales externæ, and other branches to the parts of generation, bladder, anus, &c. From the external branches goes the epigastrica to the muscles of the belly and hips; after this, the iliac branches paffing out of the abdomen, are termed cruralis, and from thence arife, 1. faphena, paffing between the muscles and integuments (only covered with the (kin and fat) from the inguen to the knee, ankle, upper part of the foot and great toe; 2. faphena minor, to the back part of the thigh and leg, and to the heel; 3. poplitea, formed of a double crural branch, runs through the ham, on the back of the gastrocnemii to the tendo achillis; 4. tibialis anterior, runs down the fore part of the leg, between the mufculus tibialis anticus, and the extenfor digitorum communis, to the upper part of the foot; 5. tibialis posterior vel suralis, is distributed through the calf and back-part of the leg (as the anterior is on the fore-part) down to the heel and foot; 6. peronæa, runs down on the infide of the fibula as low as the outer ankle, fometimes double, fometimes triple, fending ramifications to the neighbouring portions of the musculi peronæi, and long flexors of the toes.

Q. How is the vena portæ distributed?

A. The

A. The vena portæ, in its structure, has some resemblance to a tree, being divided into innumerable branches, which are dispersed throughout the whole substance of the liver. Where the trunk begins to divide, it constitutes the finus portæ in the liver, and from this proceeds the numberless ramifications as before mentioned. The roots, or inferior branches of this vein, are divided into right and left; but first the trunk sends off, 1. cysticæ gemellæ, to the vesica fellis; 2. gastrica dextra, to the stomach; 3. duodenalis vel intestinalis, to the intestinum duodenum; and from this last often proceeds the pancreaticæ. From the right branch (before-mentioned) arife, 1. mesentericæ, to the mesentery and intestines; 2. hæmorrhoidalis interna, to the rectum; 3. epiploicæ dextræ, to the caul, the right fide of the cæliac artery. From the left branch, which paffeth to the spleen, and is called splenica, arise, 1. gastricæ (which are various); 2. coronaria ventriculi, proceeding from the former, and distributed on the stomach; 3. vafa brevia, which are formed by some of the branches of the coronaria ventriculi and splenica; 4. epiploica sinistra, and 5. gastro-epiploicæ, to the caul and stomach, the left side of the cæliac artery; 6. pancreaticæ, to the pancreas; and fometimes also the hæmorrhoidalis interna. All venal branches may be accounted the roots, in regard that their lesser branches first resorb the residuous blood deposited in any part by the arteries, and carry the the same into their greater branches, and so into the trunk: thus the blood is carried by the veins and their branches to the liver; and from all parts of the body by the branches of the cava into its ascending and descending trunk, which convey it to the heart. The distribution of the veins is so various, that it is rare to see subcutaneous veins in any two persons alike.

Q. How is the vena pulmonaris distributed?

A. The vena pulmonaris arises from the left ventricle of the heart, where it first forms a sinus, then is divided into four branches, and afterwards into innumerable ramifications, which are distributed through the whole substance of the lungs, and accompany the pulmonary artery, to return the blood into the heart.

Q. What are nerves?

A. Nerves are bundles of whitish, cylindrical fibres, arising from the medulla oblongata of the brain, and the medulla spinalis, from which they go out in pairs like so many distinct trunks, and are afterwards divided into branches, ramifications, and filaments, and terminate in all the sensitive parts of the body, being the immediate organs of sensation. They are all wrapt up in the dura and pia mater, which last covers all the nerves in common, and also incloses every fibre (of which they are composed) in particular. To these membranous coats, an infinite number of vessels, both arteries and veins, are distributed. Though the

the nerves do not appear through the best microscope to have any cavity or fluid contained in them, yet it is possible (and indeed it is the general opinion) that there may be fuch cavities, and also highly probable (from the experiments of Bellini and others) that there is fuch a fluid, though too fubtile to be perceived by us. This fluid (named animal spirits) is supposed to be conveyed by the nerves to all parts of the body, being an extreme fubtile fluid, fecreted in the brain and spinal marrow, and is thought to be destined for no less noble a purpose than the sensation, motion, and nutrition of the feveral parts of the human struc-But the existence of this fluid and the cavity of the nerves, are still the subjects of dispute, though I am inclined to favour both. Those who deny animal spirits in the nerves, suppose that the sensation is conveyed by a vibration, which, though it feems to me improbable, yet it is possible it may be conveyed either or both ways (though at present undetermined) and probably some nerves may be fo fmall, as to escape even microscopical obfervation likewife. The harmony and fympathy of the nervous parts is of great use in physic; for without an accurate knowledge of this, many symptoms of difeases can scarcely be explained. There is a wonderful connection, sympathy, and communication of motion as well as fensation, when they are affected by any violent cause: all which is owing to the nerves; for when any violence is offered them.

them, fo as to threaten a folution of union, it creates a pain and stricture of the adjacent, and even of the remote parts, especially of the veffels. The nerves generally run as strait as the parts of the body and their own fafety from external injuries will admit, fending off their branches at very acute angles, and confequently running more parallel than the bloodveffels; and I am inclined to think that every the minutest nerve, terminating in any part, is a distinct cord from its origin, or else I do not fee how they could produce distinct fensations in every part of the body; and the distinct points of sensation throughout the body are to very numerous, that the whole body of nerves (which, taken together, would not make a cord of an inch diameter) must be divided into fuch a number, to afford one for every part that has a distinct sensation, that furely fuch a nerve would be too fmall to be feen by the best microscope.

Q. How are the nerves divided?

A. The nerves are divided into those which come immediately out of the cranium from the brain; and those which come out between the vertebræ from the medulla spinalis.

O. How are the nerves from the brain di-

fiributed?

A. The nerves from the brain, or medulla oblongata, are ten pair, and are thus distributed, viz. 1. the olfactory pair, which passing through the os cribrosum, vel ethmoides, vel cribrisorme, are spread over the membrane of the

the nostrils; 2. the optic pair, which, by their expansion, form the retina of the eye; 3. the motory pair of the eyes, each of which is divided near the orbit into fix parts or branches; the first branch goes to the elevator palpebræ; the second, to the elevator occuli; the third, to the depressor; the fourth, to the adducens; the fifth, to the obliquus inferior; and the fixth, into the tunics of the eye; 4. the pathetic pair, are very small, and run to the obliquus superior, or trochlear muscle of the eye; 5. the gustatory pair, which are very large, and divided within the cranium into three branches, immediately under the dura mater: of these, the first branch, called the opthalmic, runs to various parts of and about the eye, the eye-lids, the muscles of the forehead and nofe, and the integuments of the face; the fecond branch may be called the fuperior maxillary one, as being finally diftributed through all parts of the upper jaw, the lips, nose, palate, uvula, gums and teeth; a branch of it also runs to the ear, and, joining with a branch of the seventh pair, forms the chorda tympani; the third branch may be called the maxillaris inferior, as being diffributed over the feveral parts of the lower jaw, the tongue, and other parts of the mouth; whence the whole pair of nerves has obtained the name of par gustatorium; though a great part of them serves to very different purposes, and is carried to parts that have nothing to do with tasting; 6. the abducent pair (except a 1) 3 branch

branch for the formation of the intercostal nerve) is wholly carried to the abducens occuli; whence its name. The intercostal nerve is formed either of ramifications of the two preceding nerves, or only of those of the fixth pair. It makes its way out of the cranium by the passage of the internal carotid, and defcends near the eighth pair through the neck, and thence through the breast and abdomen, even to the pelvis, making in its way various plexuses and ganglia, and sending branches to almost all the parts contained in the breast and abdomen; 7. the auditory pair arise with two trunks, one of which is called the portio dura, and the other portio mollis: this last enters the foramen of the os petrofum; and thence, through various little apertures, gets into the labyrinth of the ear, where it is expanded over all its parts, and constitutes the primary organ of hearing. The harder portion, paffing through the aquæductus Fallopii, fends back one branch into the cavity of the cranium; it also sends off another branch, which helps to form the chorda tympani, and others, to the muscles of the tympanum; the rest of this pair goes to the external ear, the pericranium, the muscles of the os hyoides, the lips, the eye-lids, and the parotids; 8. the par vagum, with the accessorius Willisii, pass out near the lateral finuses of the dura mater; and descending through the neck and thorax to the abdomen, fend out branches by the way to the larynx, the pharynx, heart, lungs, and especially

to the stomach: it also sends off, from the upper part of the thorax, large branches, which are variously implicated in the neck, thorax, and abdomen; with the linguals, the cervicals, and the intercostals; 9. the lingual pair go immediately to the tongue, and are called by fome, the motory nerves of the tongue; but by others, with more justice, the gustatory nerves; 10. this pair comes out from the beginning of the medulla spinalis, betwixt the os occipitis and first vertebra colli, and is all (except what goes to the ganglion of the intercostal) spent on the musculi obliqui, and extensores capitis. But we are to observe, says Heister, that the pair of nerves which the generality of writers have called the tenth pair of the head, are, for many unanswerable reasons, to be properly called the first pair of nerves of the neck.

Q. How are the nerves from the medulla

spinalis distributed?

A. The nerves which come out between the vertebræ from the medulla spinalis, are generally reckoned thirty pair, viz. the nerves of the neck are eight pair; the nerves of the back, twelve pair; the nerves of the loins, sive pair; and those of the facrum form five or six pair, though not always determinately and regularly so. From the nerves of the neck are innumerable branches distributed through the muscles of the head, neck, scapula, and humerus; from the third, sourth, and fifth pair, are formed the nerves of the diaphragm; and

and the fixth, feventh, and eighth pair, together with the first pair of the back, form the fix robust nerves of the arms and hands: to this division are the accessory spinal nerves of Willis to be referred, which arise about the origin of the third or fourth pair. The nerves of the back, besides the branch they give to the brachial nerves, run entirely in the same furrow along the course of the ribs, and are dispersed over the pleura, the intercostal, pectoral, and abdominal muscles, the breast, and other parts of the thorax. The nerves of the loins are generally dispersed over the loins, the peritonæum, integuments and muscles of the abdomen; and besides this, their first pair often gives, on each fide, a branch to the diaphragm; the fecond pair, after inosculating with the branches of the first, third, and fourth pairs, forms the crural nerves, which are distributed over the anterior part of the thigh; and in the same manner a branch is formed of the conjunctions of the second, third, and fourth pairs, which passes through the great foramen of the os pubis, to the scrotum, testicles, and the adjoining parts; the fourth and fifth pair of the nerves of the loins, joining with the first, fecond, third, and fourth pair of the os facrum, compose the nerve called ischiatic, which is the largest in the body; it descends along the hinder part of the thigh, and its branches are diftributed over the whole leg, the foot, and toes. The nerves of the facrum form five or fix pair, and pass through the foramina of this bone: the

the superior of them, as I have already obferved, compose the ischiatic nerve; and what remains, is dispersed in a multitude of ramisications over the parts contained in the pelvis, the intestinum rectum, the bladder, the parts of generation, and the parts adjacent.

See the Table of the Arteries, Veins, and

Nerves.

Q. What are lymphæducts, or lymphatic veins?

A. Lymphæducts, or lymphatic veins, are fine transparent tubes or vessels, which carry lymph from all parts, especially the glands, which they discharge into the larger veins, and into the vafa lactea. They are not continuations of arteries, but have their origin in all the cavities and cellular substance of the body: they are more numerous in glands than other parts, especially in those glands which separate the viscidest fluids, as may be observed in the liver and testes. The lymphatics have but one difcernible coat, which is very thin, having valves at small and uncertain diftances, to prevent the regress of their fluid; and they have frequent communications, like the veins. The larger trunks are in many places attended with small glands, through which they run, and at the fame time fend communicant branches over them, as a fecurity against obstructions from diseases in those glands: they all ultimately terminate in the receptaculum chyli.

Q. What are the uses of the lymphatics?

A. The

A. The use of the lymphatics is to dilute the chyle and absorb the sluids which are thrown into the several interstices of the body, thereby preventing any morbid accumulation. When a lymphatic vessel bursts, it occasions a dropfy in the cavity into which it opens. The lymphatics are perfectly similar to the lacteals. The lacteals are the absorbents of the intestines, as the lymphatics are of the other parts; there is no difference but the name. The same vessels are called lacteals in the intestines, and lymphatics in the other parts of the body.

Q. What are glands?

A. Glands are small bodies formed by the interweaving of veffels of every kind, covered with a membrane; generally composed of an artery, vein, lymphatic, excretory duct and nerve. The glands are of two kinds, viz. the fimple, called conglobate glands; and the compound, named conglomerate glands. Some glands are confiderably hard and firm; others very foft and tender: of the latter kind in particular are the glands in the articulations of the bones. The glands differ very confiderably in colour, figure, and fize: fome take their names from their peculiar figure; as the glandula pinealis, the miliares, and others; fome, from their use and contents, as falival glands, mucofe and lymphatic, &c. and others take their names from their fituation; fuch are the parotid glands, axillary, inguinal, mesenteric, lingual, &c.

Q. What are the uses of the glands?

A. To fecrete fluids from the mass of blood for divers purposes, such as perfecting the lymph, &c. Their uses are as different as their colours or figures; some of them are salival, mucose, and lymphatic; others are mucilaginous, sebaceous, and waxy; others lachrymal, pituitary, &c. and from these their several contents or secretions, they are termed lachrymal, &c.

Q. What is a conglobate gland?

A. A conglobate gland is a little smooth body, wrapt up in a fine skin or membrane, by which it is separated from all other parts, only admitting an artery and a nerve to pass in, and giving way to a vein and excretory canal or duct, to come out; of which fort are the glands of the brain and testes. Winslow includes, under the name of conglobate glands, the lymphatic glands alone, and calls all the other glands of the body by the name of conglomerate.

Q. What is a conglomerate gland?

A. A conglomerate gland is that which is composed of several little conglobate glands, all tied up together in one common tunicle or membrane. Sometimes all their excretory facts unite, and make one common pipe, through which the liquor of them all runs, as the pancreas and parotides do. Sometimes the ducts uniting, form several pipes, which only communicate with one another by cross canals; and such are the manimæ: others again have several pipes without any communication with

one another; of which forts are the glandulæ lachrymales, and proflatæ; and a fourth fort is, when each little gland has its own excretory duct, through which it transmits its liquor to a common bason; as the kidneys.

Q. What are excretory veffels?

A. Excretory veffels are either tubes from glands to convey the fecreted fluids to their refpective places; or vessels from the small guts to carry the chyle to the blood veffels: thefe last are called vasa lactea, or lacteal vessels.

Q. What are the lacteal veffels?

A. The vafa lactea, or lacteal veffels, are the venæ lactæ, receptaculum chyli, and ductus thoracicus, filled with a white milky fluid, called chyle.

Q. What are the venæ lacteæ?

A. The venæ lacteæ, &c. have the name of veins, because their valves are disposed as the ordinary blood veins, and because the fluid which they contain runs from fmaller into larger tubes or vessels. The lacteal veins are long, slender pipes, whose coats are so thin as to become invisible when they are not distended with chyle. They arise from all the parts of the small guts, by a vast number of fine capillary, pellucid tubes, which, as they run from the fides of the intestines to the glands in the mesentery, frequently unite, and form fewer and larger branches (called venæ lacteæ primi generis) which pass through the glands of the mesentery; and from these arise other lacteals of a larger fize (named lactea fecundi generis)

generis) which carry the chyle immediately into the receptaculum chyli. The mouths of these lacteals, which are open into the cavity of the guts, from whence they receive the chyle, are fo fmall, as not to be feen by the very best microscope: it was necessary they should be smaller than the finest arteries in the body, that nothing might enter which could ftop the circulation of the blood. The fame extremity of the lacteals has likewife communication with the capillary arteries of the guts, by which they receive a lymph that dilutes and propels the chyle forwards, and washes the lacteals and glands, that they may not be obstructed by the chyle's staying in them upon fasting. The other extremity of the lacteals discharges the chyle into the vesicular cells of the mesenteric glands, &c. as I have beforementioned. The lacteal veins have valves at certain distances, which hinder the chyle from returning back into the intestines. The office of these veins is to receive the fluid part of the digested aliment, which is called chyle, and convey it to the receptaculum chyli, that it may be thence carried through the ductus thoracicus into the blood vessels.

Q. What is the receptaculum chyli?

A. The receptaculum chyli is a membranous fomewhat pyriform bag, two-thirds of an inch long, one third of an inch over in its largest part when collapsed; situated on the first vertebra lumborum to the right of the aorta, a little higher than the arteria emulgens dextra,

dextra, under the right inferior muscle of the diaphragm. It is formed by the union of three tubes; one, from under the aorta; the fecond, from the interftice of the aorta and cava; the third, from under the emulgents of the right fide. It lies between the descending trunk of the aorta and the above vertebræ, and is biggeft between the cæliac and emulgent arteries, furrounded by veficular glands, called glandulæ lumbares, which difcharge their lymph into it. The receptacle receives all the fecond order of the lacteals, as well as all the lymphatic veins, both of the legs, and of all the parts of the abdomen. This faccus chyliferus will contain about one ounce of water; at its fuperior part becoming gradually fmaller, it is contracted into a slender, membranous pipe, of about a line diameter, called the ductus thoracicus.

Q. What is the ductus thoracicus?

A. The ductus thoracicus, or thoracic duct, is the superior part of the receptaculum chyli contracted into a slender, membranous pipe, of about a line diameter, which passes between the appendices musculosæ diaphragmatis, on the right of, and somewhat behind the aorta, then lodged in the cellular substance under the pleura: it mounts between this artery and the vena sine pari, or azygos, as far as the sisth vertebra thoracis, where it is hid by the azygos, as this vein rises forward to join the cava descendens; after which, the duct passes obliquely over to the left side under the cesophagus,

gus, aorta descendens, and great curvature of the aorta, until it reaches the left carotid, stretching farther towards the left internal jugular, by a circular turn, whose convex part is uppermost: at the top of this arch it splits into two, for one half line; the superior branch receiving into it a large lymphatic from the cervical gland. This lymphatic appears, by blowing and injections, to have two valves: when the two branches are united, the duct continues its course to the internal jugular, behind which it descends; and immediately at the left fide of the infertion of this vein, enters the fuperior and posterior part of the left subclavian, whose internal membrane duplicated, forms a femilunar externally convex valve, that covers two thirds of the orifice of the duct: immediately below this orifice, a cervical vein, from the musculi scaleni, enters the subclavian. The ductus thoracicus has a thin coat, and valves, at feveral distances (commonly ten or twelve) which hinder the chyle that has once paffed them, from falling back. The diameter of the duct varies in most bodies; and in the same subject is uniform; but frequently fudden enlargements, or facculi of it, are obfervable. The divisions which authors mention of this duct within the thorax, are very uncertain; as is the precise vertebra, where it begins to turn towards the left. Frequently it does not split at its superior arch; in which case, a large saccus is found near its aperture into the fubclavian vein: generally it has but

one orifice, though I have feen two in one body, and three in another; nay, fometimes it divides into two, under the curvature of the great artery, one going to the right, another to the left fubclavian: this however, is very rare. The lymphatic which enters the superior arch, is often fent from the thyroid gland. The thoracic duct receives the lymphæducts from the feveral parts of the thorax, as it passes along to the fubclavian vein: by its running up to the left fide, the chyle receives a new impetus from the pullation of the aorta; whereas, on the right fide, it must have ascended only by the pressure of the diaphragm, and the muscles of the lower belly upon the receptacle, which it equally enjoys in its present situation. The receptaculum chyli is eafily found in living bodies, but with greater difficulty in those that are dead.

Q. What are membranes?

A. Membranes are a pliable texture of fibres interwoven together, and expanded, to cover or line any other part. Every diffinct part of the body is covered, and every cavity is lined with a fingle membrane, whose thickness and strength is as the bulk of the part it belongs to, and as the friction to which it is naturally exposed. The membranes of the body are various, and variously denominated: those which serve as integuments, or covers of vessels, are called tunics or coats; those that cover the brain, meninges; the skull, pericranium; the bones in general, periosteum; that which lines the

the thorax, pleura; the abdomen, peritonæum; and that which includes the heart, pericardium. The muscles too, are each inclosed in a peculiar membrane; as are the bowels, &c.

Q. What are the uses of the membranes?

A. The uses of the membranes are to cover and wrap up the parts, strengthen them, and fave them from external injuries; those that contain distinct parts, keep the parts they contain together, and render their surfaces smooth, and less subject to be lacerated by the actions of the body; and those which line cavities, ferve to render the cavities smooth and fit for the parts they contain to move against. The membranes are not only of use to join one part to another, but also to preserve the natural heat, to fustain small vessels, and the nerves, which run through their duplicatories; to stop the returning of the humours in their vessels (as the valves stop the returning of the blood in the veins and heart) of the chyle in the thoracic duct, and of the lymph in the lymphatic veffels. The membranes of all the cavities that contain folid parts, are studded with glands, or are provided with veffels which separate a mucus to make the parts contained move fmoothly against one another, and not grow together; and those cavities which are exposed to the air, as the nofe, ears, mouth, and the trachea arteria, have their membranes beset with glands which separate matter to defend them from the outer air.

Q. What are fibres?

A. Fibres appear to be fimple, thread-like bodies, which ferve to form other parts; hence fome are very hard, as the bony ones; and others foft, as the fleshy parts.

Q. What is hair composed of?

A. Of five or fix oblong, flender, flexible filaments, inclosed in a tube, and transparent, growing out of the pores, and is either strait or crooked, according to the rectitude or obliquity of the pores.

Q. What are nails?

A. A kind of bony excrescence growing on the fingers and toes; their substance consists of the cutaneous papillæ, elongated and indurated in that form.

DIALOGUE II.

Of the HEAD, and its PARTS.

QWHAT are the parts of the head, and

how distinguished?

A. The hair is termed capillus; that part which it covers is termed the scalp; the crown of the head, vertex; the hinder part, occiput; the fore-part, sinciput; that part between the hair and the eye-brows, frons, the forehead; the sides of it tempora, the temples; the eye-brows, supercilia; the space between the eye-brows, glabella; the eye lids (both upper and lower) palpebræ superior et inferior, and the carti-

cartilaginous edges of both, with the hairs growing thereon, are termed cilia; the hollow of the eye, cavum oculi; the inner angle of the eye, canthus major; the outer corner of the eye, canthus minor; and the eye, oculus; which confifts of proper tunicles and humours. The ridge of the nose is termed spina nasi; the tip of the nose, orbiculus, vel globulus nasi; the nostrils, nares; the partition between them, feptum nasi; the sides of the nostrils, alæ; the hairs growing within, vibrifæ: internally, the maxillary, fphenoidal, and frontal finuses. The outer ear is termed auricula, but more properly auris externa: in this there are a great many eminences and cavities; the upper part is termed pinna; the lower part, lobus, or fibra; the outer circle, helix; the inner circle (or femi-circle) antihelix; the space between the two circles, scapha; the lower end of the antihelix, or femicircle, makes a little prominence, which is called antitragus - because there is another prominence just opposite to it, called tragus; the cavity made by the extremity of the helix, is called concha; the hollow in the middle of the ear, is termed alvearium, and has a hole or passage into the internal ear, named meatus auditorius. The upper jaw is termed maxilla fuperior; the prominent part of the cheek, mala; the lower jaw, maxilla inferior; the hollow, or the cheeks stretched out in blowing, bucca; the chin, mentum; the beard, or place where it grows on the upperlip, mystax; the gutter in the middle under

the nose, is philtrum. The external parts of the mouth are the lips, one upper, and the other under: the internal parts are the gums, termed gingivæ; the teeth, dentes; the tongue, lingua; the roof of the mouth, palatum; with the uvula there suspended for modulating the voice, &c. The other internal parts of the skull, eye, nose, ear, and mouth; such as the brain, glands, membranes, muscles, &c. I shall speak of hereafter in the particular descriptions of these parts.

Q. What are the bones of the head?

A. The bones of the cranium, or skull, which contains the brain, &c. the bones of the face, jaws, teeth, tongue, and internal ear.

Q. What are the bones of the cranium?

A. The bones of the cranium are but eight; though if we reckon the offa petrofa distinctly from the offa temporum, they are ten, viz. os frontis, offa parietalia, two; offa temporum, two; os occipitis, os ethmoides, os sphenoides.

Q. Which is the os frontis?

A. The os frontis is situated in the fore-part of the skull, and forms that part of the face called the forehead; from whence its name. It is joined by suture to the offa parietalia, ungues vel lachryma nasi, maxillaria, malarum, os ethmoides et sphenoides. The os frontis contains the anterior lobes of the brain: in its middle there is generally a ridge, or spine, to strengthen it, and to which the longitudinal sinus of the dura mater adheres; and from the middle

middle of this bone externally, goes a process to fupport the bones of the nose; and the lower parts of the os frontis compose the upper parts of the orbits of the eyes. Immediately above the os ethmoides is a small blind hole, through which runs a vein to the longitudinal finus of the dura mater, and through the upper edge of each orbit, nerves, and an artery, pass to the forehead; and in each orbit, near the os planum, passes a branch of the fifth pair of nerves. In the substance of this bone, near the note, are feveral finuses more in some subjects than in others; in children feldom any. These finuses, and the spine in this bone, render it dangerous, if not impracticable, to apply a trephine on the middle and lower part of the forehead.

. Q. What are the offa parietalia?

A. The offa parietalia (by some named bregmati, or sincipitis) are two bones larger than any other in the skull, forming the superior and lateral parts of it, and to which the temporal muscles are partly fixed.

Q. Which are the offa temporum?

A. The offa temporum, vel squamosa, are two bones situated below the parietal bones, at the middle and lower parts of the sides of the skull, from which proceed the mammillary and zygomatic processes; and it has an exterior sinus lined with a cartilage, which receives the process of the lower jaw.

The offa petrofa lie between the former and the occipital bones, or are truly portions of the

tempo-

temporal bones, being never found separate in adult bodies. In each of these bones there is an external and an internal process; the former is named styliformis; near it is the sixth foramen, through which the carotid arteries pass to the brain; and that on the inside of the skull, leading to the organs of hearing, is the seventh foramen; the latter process is called the os petrosum, which contains the whole meatus auditorius and cavity of the tympanum.

Q. Which is the os occipitis?

A. The os occipitis makes all the back part of the skull: it is bounded by the sphenoidal, temporal, petrofal, and parietal bones. This bone is articulated to the spine; and between its apophyses is the great or tenth foramen, through which the medulla oblongata descends into the spine, the cervical arteries and veins pass: on the inside of this bone is a crucial ipine; and on the outfide a spine, or an apophysis, to strengthen it: the thinner parts of this bone are also defended by the muscles that cover them, as blows here are of worse consequence than on any other part of the skull; because wounds in the cerebellum, which is underneath, are mortal. Near the apophyses of this bone is the ninth foramen of the skull, through which pass the ninth pair of nerves; and behind each apophysis of the occipital bone there is a foramen, or a finus, through which pass sinuses from the lateral sinuses of the external cervical veins, by means of which

(as in all other communications of the finuses) the blood passes from those that happen to be surcharged by any posture of the head, into those that, from the same posture, would have been almost empty.

Q. Which is the os ethmoides?

A. The os ethmoides, vel cribriforme, is a small bone, about two inches in circumference, lying in the fore-part of the skull, and almost surrounded by the os frontis. It is full of holes like a sieve; it is a principal part of the organ of smelling, and gives a very great extent to the pituitary membrane in a small compass: in the middle of this bone is the cristal galli process, and opposite to it a thin one, which in part divides the nose.

Q. Which is the os sphenoides?

A. The os sphenoides is fixed like a wedge in the midst of the os frontis, ethmoides, vomer, occipitis, maxillæ fuperioris, offa parietalia, palati, malarum, temporum, and petrofum: on its infide is a cavity, named fella turcica, vel equina, and the four clinoid processes; under the two foremost of which, pass the internal carotid arteries: opposite to the sella turcica is a process, which makes part of the septum narium: on the outfide of the skull, adjoining to the upper jaw, are the pterygoid proceffes: under the fella turcica, vel equina, in this bone, is the sphenoidal sinus, which is iometimes double, and opens into the nostrils; but fometimes it is totally wanting, especially in children. At the infide of the basis of the two anterior clinoid processes, are the first foramina of the skull through which the optic nerves pass: near this is the second foramina, through which pass nerves and blood vessels into the the orbits of the eyes; and towards the occiput are the third foramina, through which pass nerves to the face; nearer the occiput are the fourth foramina, through which pass the largest branches of the fifth pair of nerves; and a straw's breadth farther are the fifth foramina, through which those branches of the carotid arteries enter, that are bestowed upon the dura mater.

Q. How are the bones of the skull com-

posed?

A. They are of unequal thickness in the several parts, and are composed of two lamellæ, or tables, laid over one another; between which there is a diploe, or meditullium, being a thin, spongeous substance, made of bony fibres detached from each lamina, and full of little cells. The tables are hard and folid, the fibres being close to one another. The diploe is foft, the bony fibres being here at a greater distance (a contrivance of the all-wife Creator's) whereby the skull is not only made lighter, but less liable to fractures. The external lamina of the cranium is smooth, and covered with the pericranium, which in other bones is called by the general name of periosteum, because of its adhering to the bone: it is found connected to the dura mater by fibres transmitted from it to that membrane through the futures. the

the origin of the temporal muscles the coats of the pericranium part; the outer passing over those muscles, and the inner still adhering close to the cranium. The internal lamina of the cranium, or inside of the skull, is likewise smooth, except the furrows made by the pulsation of the arteries of the dura mater before the cranium was arrived at its consistence. All the bones of the cranium are found to be imperfect in new-born infants; the sinus and its meditullium are almost wholly wanting.

Q. What are futures?

A. Sutures are the closing or joining together the bones of the skull, like the teeth of faws set one into another; and these are either common or proper; the proper sutures are diffinguished into the true, and the false or spurious.

Q. Which are called true futures?

A. They are called true futures which are denticulated mutually into each other with a multitude of faw-like teeth, and are most plainly to be feen; such are the coronal, fagittal, and lambdoidal sutures.

Q. Which are the false, or spurious sutures?

A. The false, or spurious sutures, are those squamose ones of the temporal and parietal bones, and of the os frontis and sphenoides in the angle where they unite with the parietals.

Q. Which are the common futures?

A. The common futures are the transversal ones, which join the os frontis with the bones below it; but these are of little moment.

Q. Which

58

Q. Which are the coronal futures?

A. The futura coronalis runs across the skull from one upper edge of the sphenoidal bone to the other, and joins the parietal bone to the frontal.

Q. Which are the fagittal futures?

A. The futura fagittalis joins the parietal bones, beginning at the os occipitis, and is continued to the os frontis; in children down to the nose; the os frontis in them being two bones, and sometimes so in adult bodies.

Q. Which are the lambdoidal futures?

A. The futura lambdoidalis joins the back part of the offa bregmatis, or parietal bones, to the upper part of the occipital.

Q. Which are the fquamofe futures?

A. The futura fquamofa is made by the upper part of the temporal and sphenoidal bones wrapping over the lower edges of the parietal bones.

O. Which are the transversal sutures?

A. They run across the face, through the bottoms of the orbits of the eyes; they join the lower edge of the frontal bone to the os sphenoides, maxillæ superioris, ossa nasi, ungues, palati, and jugalia, or malarum.

O. What are the uses of the futures?

A. Not only to join the bones of the cranium together, but the skull being thus divided into many bones, renders it less apt to be fractured, and, when it is fractured, they prevent its being extended, as it would have been, were it composed of one bone only. They are also

of use to join the dura mater very firmly to the cranium and pericranium in those parts, and for the offisication of the bones; and in infants, that the head may be the more easily extended in its growth, and also that they may give way in the birth: and the openness of the sutures in children have another advantage, viz. that medicinal applications to the external part of the head may penetrate and do service; and lastly, that the transpiration from the brain may be the more free and easy at that time of life in which the bones are open, and in which also the habit is more humid.

Q. What use are the foramina, or holes of

the fkull?

A. To give passage to the spinal marrow, nerves, arteries, and veins.

Q. How are they distinguished?

A. Into external and internal.

Q. Which are the external?

A. The external are meant those which are easily discovered on the outside of the skull.

Q. Which are the internal?

A. By the internal are meant those which are most obvious in the internal surface of the skull.

Of the larger internal foramina there are eleven pair, affording passage to the arteries, veins, and nerves of the brain; besides these, there is one that is single, viz. the great foramen of the occipital bones, that gives passage to the medulla spinalis, accessory spinal nerves, and vertebral arteries: 1st pair, gives passage

to the olfactory nerves; 2d pair, to the optic nerves; 3d pair, to the third and fourth pair of nerves, to the first branch of the fifth pair, and to the fixth pair, as also to the emissary of the receptacles of the dura mater; 4th pair, give passage to the second branch of the fifth pair of nerves, which is distributed to the feveral parts of the upper jaw; 5th pair, to the third branch of the fifth pair of nerves, and to the emissary of the dura mater; 6th pair, give passage to an artery (which is distributed over the dura mater, and is that which forms the tree-like impressions on the inside of the parietal bones); 7th pair, is between the fella equina and the petrofe apophysis, and is shut up by the dura mater; 8th pair, give passage to the carotid arteries; and here the intercostal nerve goes out; 9th pair, to the auditory nerves; 10th pair, to the par vagum, the lateral finuses of the dura mater, and to the spinal nerve; 11th pair, give passage to the lingual nerves: besides these large foramina, there are a number of little ones often very visible.

Q. What external foramina are there?

A. Two of the os frontis, a little above the orbits, called fupra orbitalia; they give passage to the opthalmic nerve of Willis; there are alfo four others, two on each fide the orbit, which transmit little nerves and vessels to the finus of the os ethmoides; in the parietal bone one, though sometimes none; in each of the offa temporum, three common and three

proper;

proper: the Ist is the foramen jugale for the passage of the temporal muscle; in the 2d is the finus of the jugular vein; and the 3d is the ductus Eustachii, situated between the petrofum and the sphenoides, and leading from the mouth into the internal ear: besides these, there are also three proper foramina in the ossa temporum; ist the meatus auditorius; 2d the aquæduct of Fallopius, situated between the mastoide and styloide processes, and transmitting the hard portion of the auditory nerve; 3d a foramen behind the mastoide process. In the occipital bone are generally two foramina behind the condyloide apophyses for the passage of the vertebral veins into the lateral finuses of the dura mater. In the sphenoidal bone, befides those already mentioned, are the apertures of the finuses into the nostrils, common to them with the bones of the palate, and which are the apertures of the nares and fauces. In the upper part of the pterygoide processes, is a passage for the novum emissarium of the dura mater. In the os ethmoides, besides those common to this bone with the os frontis (already mentioned) there are the apertures of the ethmoidal finuses into the nostrils.

Q. What are the bones of the face?

A. The bones of the face are the offa nafi, malarum, ungues, plana, maxilla superior, palati, os vomer, spongiosum, maxilla inferior; to which may be added, the bones of the ear and tongue.

Q. Which are the offa nasi?

A. The offa nasi makes the upper part, or ridge of the nose.

Q. What are the offa malarum?

A. Two bones situated in the lateral and middle parts of the face, called the cheek bones; the short processes of which, together with the processus jugalis, form arches, called the offa jugalia, vel zygomaticus.

Q. Which are the offa ungues, vel lachry-

malia?

A. The offa ungues, vel lachrymalia, are two of the least bones of the face, situated in the orbits of the eyes towards the nose, very thin and transparent; between each of them and the upper jaw, is a foramen as large as a goose quill, into which the puncta lachrymalia lead, to carry off any superfluous moisture from the eyes into the nose.

Q. Which are the offa plana?

A. Situated also in the orbits beyond the offa ungues, and are near thrice as big.

Q. Which is the maxilla superior?

A. The maxilla fuperior, or upper jaw, though generally described single, consists of two bones, beingmanisestly divided by a suture, which is scarce ever obliterated. Its two processes make part of the nose. Its upper and outward parts make the lower parts of the orbits of the eyes; its lower side, all that part of the face under the cheeks, eyes, and nose, to the mouth, and two-thirds of the roof of the mouth. A little below the orbits of the eyes there are two holes in this bone, and behind

the dentes incifores one more, which divides into two, as it opens into the nofe. Between the posterior grinding-teeth and the orbits of the eyes, are two great sinuses, called antramaxilla superioris, which open into the upper part of the nose; and in the lower edge of this jaw are the alveoli, or sockets for the teeth. Part of the sides of these sinuses that lie next the nose, are only membranes, which make the cavities like drums; perhaps to give a grave sound to the voice when we let part of it through the nose.

Imposthumations sometimes happen in these sinuses; the signs of which are, great pain about the part, matter in the nose on the side diseased, stinking breath, and rotten teeth: the cure is performed by drawing out the last tooth but one, and drawing two or more, if rotten, and through their sockets making a perforation into the antrum.

Q. Which are the offa palati?

A. The offa palati are two small bones that make the back-part of the arch of the palate, or roof of the mouth: near this bone are two small foramina, through which arteries and nerves pass to the palate.

Q. Which is the os vomer?

A. The os vomer is situated between the bones of the palate and the sphenoidal bone. This bone, together with the two nasal fosse and its cartilage, are the septum nasi.

Q. Which is the os spongiosum?

A. The os spongiosum, though by some

treated as a distinct bone, is only the spongy laminæ in the nose of the os æthmoides and ossa plana.

Q. Which is the maxilla inferior?

A. The maxilla inferior, or lower jaw, is but one bone in adults, but in children two, which unite by a cartilage in the middle of the chin, and, as the child grows, hardens into bone: in children it also consists of two tables and a diploe between. This bone forms two condyloid processes; and near these are the two coronal processes; and at the inside of the chin a small rough processus innominatus. In the inside of this bone, under each processus coronalis, is a large foramen, which runs under the teeth, and passes out near the chin: in this foramen the vessels pass that belong to the teeth; and in the upper edge of the jaw are the sockets for the teeth.

Q. What number are the teeth, and how

are they distinguished?

A. The number in adults, if perfect, is generally thirty-two (fixteen in each jaw); the four fore teeth in each jaw are called incifores; the next (one on each fide the incifores) canini, called by fome the eye-teeth; the rest are the molares, or grinders; the last of which (on each side) above and below, are termed dentes sapientiæ, because they do not appear till men arrive at the years of discretion. The incifores and canini have only one single root; the two first of the molares on each side, have two roots; and the rest, some three, some four, espe-

especially in the upper jaw, where also they are spread wider, because that jaw being more spongy than the other, the teeth need more space to fix them. The cavities of the teeth are covered or lined with a fine vasculo-nervous membrane; and each branch of the roots have a foramen, or hole at their bottom, through which pass an artery, vein, and nerve, to assorbed them nutrition and sensation: these vessels and membrane are the seat of the tooth-ach.

Q. Which is the os hyoides?

A. The os hyoides, or bone of the tongue, is fituated in the middle space between the angles of the lower jaw, and adheres to the base of the tongue. The os hyoides is generally composed of five small bones; and the use of it is to give a firm basis to the tongue; and therefore several muscles of the tongue and larrynx, serving to the motions of both, are inserted in it.

Q. Which are the bones of the internal ear?

A. The bony cavity of the tympanum contains feveral little bones, called the bones of the ear; which take their names from the things they are thought to refemble, viz. malleus, incus, stapes, and orbiculare; but this last, in dried bones, passes sometimes unobferved.

Q. Which is the malleus?

A. The malleus is the hammer-like bone adhering to the membrane of the tympanum, and articulated with the incus.

Q. Which is the incus?

A. The incus, or anvil-like bone, is articulated both to the malleus and stapes.

Q. Which is the stapes?

A. The stapes, or stirrup, is a little bone, situated in the cavity of the senestra ovalis; it is joined to the incus, and its basis adheres (by means of a membrane) to the senestra ovalis of the labyrinth of the ear.

Q. Which is the os orbiculare?

A. The orbicular, or lenticular bone, is the smallest bone in the body; it is articulated with both the stapes and incus.

Q. What are the muscles, cartilages, and ligaments of the head? I mean those belong-

ing to the cranium.

A. The muscles, cartilages, and ligaments of the cranium, are thefe, viz. 1. occipitofrontalis, orbicularis palpebrarum, corrugator Supercilii, ciliaris (tarsi) cartilages and ligaments, and elevator palpebræ fuperioris; 2. the muscles of the globe of the eye, viz. elevator occuli, depreffor occuli, adductor occuli, abductor oculi, obliquus inferior, obliquus fuperior vel trochlearis; 3. the muscles, cartilages, and ligaments of the external ear, viz. cartilage and ligament, elevator auriculæ, anterior auriculæ, retractor auriculæ, helix major, helix minor, tragicus, antitragicus, tranfversus auriculæ; 4. the muscles of the internal ear, viz. laxator tympani, externus mallei, internus mallei, tenfor tympani, stapedius; 5. muscles of the nose, viz. compressor narium, elevator alæ nafi, depreffor alæ nafi, cartilages

of

of the nose; 6. muscles of the mouth, lips, chin, and cheeks, viz. sphincter oris, elevator labii superioris, depressor labii superioris vel nafalis, elevator labii inferioris, depressor labii inferioris, elevator anguli oris, depressor anguli oris, elevator menti, zygomaticus, buccinator, platysma myoides, biventer, temporalis, maffeter, pterygoideus internus, pterygoideus externus; 7. muscles of the palate, viz. pterygostaphylinus externus, thyrostaphylinus, glossoftaphylinus; 8. muscles of the os hyoides, viz. genio hyoideus, sterno hyoideus, mylohyoideus; 9. the muscles of the tongue, viz. genio-gloffus, bafio-gloffus, cerato-gloffus, ftylo-gloffus, chondro-gloffus, lingualis, the tongue itself; 10. muscles which move the head on the trunk, viz. rectus internus major, rectus internus minor, rectus major posticus, rectus minor posticus, rectus lateralis, obliquus inferior, obliquus superior.

Q. Which is the occipito frontalis?

A. The occipito-frontalis is a muscle with four sleshy bellies; it arises behind each ear from the os occipitis; and soon becoming tendinous, passes under the hairy scalp to the forehead, where it becomes broad and sleshy, adhering to the skin, and is inserted into the upper part of the orbicularis palpebrarum, into the os frontis near the nose, and by two processes into the bones of the nose. When this muscle acts from the back part, it pulls the skin of the forehead upward, wrinkling transversely, and, in some persons, the hairy scalp back-

backwards; but when the fore-part of it acts, it draws the skin with the eye-brows downward, and towards the nose when we frown.

Q. Which is the orbicularis palpebrarum?

A. The orbicularis palpebrarum furrounds the eye-lids; it arises from the upper apophysis of the maxillary bone; it shuts the eye-lids, serves to depress and draw forward the eye-brow at the same time, and elevates the lower eye-lids.

Q. Which is the corrugator supercilii?

A. The corrugator supercilii is so conjoined with the orbicularis palpebrarum, that it may be reckoned a part of it.

Q. Which is the ciliaris?

A. The ciliaris is also a very small portion of the orbicularis, next the ciliary cartilages of the eye-lids.

Q. What are the tarfi?

A. The tarsi are thin cartilages, forming the principal part of the edge of each eye-lid; their ends connected by a kind of small ligaments.

Q. What are the ligamenta tarforum lata?

A. The broad ligaments of the tarsi are situated along both edges of each orbit, the superior broader than the inferior, fixed to the edges of the cartilages, so that these ligaments and the tarsi alone, represent the eye-lids.

Q. Which is the elevator palpebræ?

A. The elevator palpebræ superioris is a very thin muscle, which arises from the periosteum at the bottom of the orbit of the eye, where the optic nerve goes through the cranium, and is inferted into the whole ciliary cartilage of the upper eye-lid by a very thin, broad tendon.

Q. What are the muscles of the globe of

the eye?

A. Four strait, viz. the elevator, depressor, adductor, and abductor; and two oblique, the superior and inferior.

Q. Which is the elevator oculi?

A. The rectus attollens, vel elevator oculi, lies on the upper part of the globe, pulling up the ball of the eye when we look up: it arises from the bottom of the orbit, between the optic nerve and the elevator palpebræ superioris, and is inserted in the upper part of the tunica sclerotis, near the cornea.

Q. Which is the depressor oculi?

A. The rectus deprimens, vel depressor oculi, pulls down the eye: it arises, and is inferted directly opposite to the elevator oculi.

Q. Which is the adductor oculi?

A. The rectus adducens, vel adductor occuli, draws the eye towards the nose: it arises from the bottom of the orbit, near the optic nerve internally, and is inserted into the tunica sclerotis on the side next the nose.

Q Which is the abductor oculi?

A. The rectus abducens, vel abductor oculi, draws the eye towards the little canthus, and hath both its origin and infertion directly opposite to the adductor. These four strait muscles terminate about the cornea by four short, thin, slat tendons: when they all act to-

F 3

gether,

gether, they draw the eye towards the bottom

Q. Which is the obliquus inferior?

A. The obliquus inferior arises from the lower side of the orbit, near its external edge, where the bones of the os maxilla superioris join together: it ascends obliquely over the depressor, and is inserted behind the tendon of the abductor: it draws the globe of the eye forward, and turns it upward.

Q. Which is the obliquus superior?

A. The obliquus superior, vel trochlearis, arises from the bottom of the orbit, between the elevator and adductor oculi, and runs obliquely towards the great canthus. In the upper and inner parts of the orbit, near its edge, there is a cartilaginous ring or pulley, through which this muscle passes in a round tendon, and is inserted near the bottom of the globe of the eye, which it pulls upward and inward, and thereby directs the pupil outward and downward.

Q. Which are the muscles of the external

ear?

A. Besides the cartilage and ligament of the ear, there is the elevator, anterior, and retractor auriculæ, the helix major and minor, the tragicus anticragicus, and the transversus auriculæ: but all these muscles are extremely small, and often scarce discernible without the help of a microscope.

Q. What are the cartilage and ligament of

the external ear?

A. The

A. The cartilage makes the greatest part of the external ear, it being the basis of all the other parts, of which this portion of the ear is made up. It is fixed to the cranium, not only by the cartilaginous portion of the meatus auditorius (which I shall describe hereafter) but also by two ligaments, one anterior, and the other posterior, opposite to one another.

Q. Which is the elevator auriculæ?

A. The attollens vel elevator auriculæ, has a very thin, tendinous origin from the tendon of the occipito frontalis, and is inferted into the back of the cartilage of the external ear.

Q. Which is the anterior auriculæ?

A. The anterior auriculæ is inferted into the back of that part of the helix which divides the concha.

Q. Which is the retractor vel retrahens au-

riculæ?

A. The retractor auriculæ arises by one, two, or three small portions from the temporal bone above the maxillary process: the upper and middle portion is inserted into the lower part of the back of the upper cavity of the external ear; and the lower portion is inserted into the back of the lower cavity. This muscle pulls the ear backward.

O Which is the helix major?

A. The helix major arises from the upper part of the outer side of the acute process of the helix; and is either inserted into the helix, or else runs along the surface of the elevator.

Q. Which is the helix minor?

A. The helix minor ascends near the helix on the fore-part of the outer ear; one end is fixed below the notch of the concha, the other above it.

Q. Which is the tragicus?

A. The tragicus arifes from the cartilage of the concha near the tragus, and terminates in the upper part of the tragus, and adjacent part of the concha.

Q. Which is the antitragicus?

A. The antitragicus arises from the outer part of the cartilage of the antitragus, and is inserted into the edge of the concha behind the antitragus, at the bottom of the helix.

Q. Which is the transversus auriculæ?

A. The transversus auriculæ is divided into two parts: that belonging to the antihelix is inferted into the back of the antihelix, and a small part of it into the back of the scapha; that belonging to the scapha is divided into several parts for a considerable length: it arises from the back of the superior cavity of the concha, and is inserted into the back of the scapha.

Q. Which are the muscles of the internal

ear?

A. The muscles of the internal ear are, the laxator tympani, externus mallei, internus mallei, tensor tympani, and stapedius.

Q. Which is the laxator tympani?

A. The laxator tympani arises from the edge of the tympanum, where the membrane of the tympanum adheres, and is inserted into the malleus.

Q. Which

Q. Which is the externus mallei?

A. The externus mallei arises from a process of the sphenoidal bone, between the os squamosum and petrosum, and is inserted into the whole length of the bony channel, which contains the auditory passage.

Q. Which is the internus mallei?

A. The internus mallei lies along the infide of the Eustachian tube, fixed in the apophysis petrofa.

Q. Which is the tenfor tympani?

A. The tenfor tympani arises from the upper part of the Eustachian tube, where it looks towards the basis of the skull, and is of a cartilaginous nature: its tendon is inserted into the handle of the malleus.

Q. Which is the stapedius?

A. The stapedius is situated within the small bony pyramid at the bottom of the tympanum; its tendon goes through a small hole in the apex of the pyramid, and is inserted into the back part of the head of the stapes.

Q. What are the muscles of the nose?

A. Two compressors, two elevators, and two depressors, besides the cartilages.

Q. Which are the compressor narium?

A. The compressor narium arises on each side from the outer part of the root of the wings of the nose, goes over the back of the anterior part of the nose, and are inserted in the moveable cartilage which forms the ala of the nares.

Q. Which are the elevator alæ nasi?

A. The

A. The elevator alæ nasi et elevator labii superioris, has its origin from the nasal process of the upper jaw bone on each side near the greater canthus of the eye: the extremities which run through the upper lip, and are inserted in the moveable cartilage, insert sibres as they pass into the upper part of the wings of the nose near the cheek.

Q. Which are the depressor alæ nasi?

A. The depressor alæ nasi arises on each side from the upper jaw bone, where the gums cover the sockets of the dentes incisores and canini: it is inserted round the root of each wing of the nose, and under, or within each nostril from the septum nasi, where that joins with the lip to the wing of the nose.

Q. What are the cartilages of the nose?

A. The inferior portion of the external nose is composed of several cartilages, which are commonly sive, the largest; the rest are uncertain. The middle cartilage is the most considerable, and supports the rest, being connected immediately to the bony parts; the other sour are connected to the middle cartilage, and to each other by means of ligaments.

Q. What are the muscles of the mouth, lips,

chin, and cheeks?

A. These are, the sphincter oris, elevator labii superioris, depressor labii superioris vel nasalis, elevator labii inferioris, depressor labii inferioris, elevator anguli oris, depressor anguli oris, elevator menti, zygomaticus, bucci-

nator, platysma myoides, biventer, temporalis, masseter, pterygoideus internus and externus.

Q. Which is the sphincter oris?

A. Orbicularis, but more properly sphincter vel constrictor oris, surrounds the whole mouth about three-fourths of an inch broad. This muscle is very much intermixed with all the muscles that are inserted into it.

Q. Which is the elevator labii superioris

proprius?

A. The elevator labii superioris proprius, arises from the bones of the upper jaw under the anterior and inferior part of the orbicularis palpebrarum, and also from the os malæ, and is inserted into the upper part of the sphincter oris. This raises the upper lip, and helps to dilate the nostrils.

Q Which is the depreffor labii superioris

proprius?

A. The depressor labii superioris propirus, is a small muscle arising from the upper jaw, near the dentes incisores, and is inserted into the upper part of the lip, and root of the cartilages of the nose; hence it is also a depressor of the nose, which action constricts the nostrils.

Q. Which is the elevator labii inferioris pro-

prius?

A. The elevator labii inferioris proprius, arises from the lower jaw, near the dentes incifores, and is inserted into the lower part of the under lip.

Q. Which is the depressor labii inferioris

proprius?

A. The depressor labii inferioris proprius, arises broad from the lower jaw at the chin, and is foon inferted into the sphincter oris.

Q. Which is the elevator anguli oris?

A. The elevator anguli oris vel labiorum communis, arifes from the fuperior maxilla under the middle of the orbit, and is inferted into the sphincter muscle near the corner of the mouth.

Q. Which is the depressor anguli oris?

A. The depressor anguli oris vel labiorum communis, arises laterally from the lower jaw near the chin, and is inferted into the sphincter opposite to the elevator anguli oris.

Q. Which is the elevator menti?

A. The elevator menti arises from the forepart of the lower jaw, from the focket of the lateral incifive tooth, extending to the focket of the next tooth on each fide.

Q. Which is the zygomaticus?

A. The zygomaticus is a muscle arising from the os zygoma, or malæ, and is inferted into the sphincter at the corner of the mouth, which it draws outward and upward.

Q. Which is the buccinator?

A. The buccinator ariseth broad and fleshy from the fore-part of the processus coronæ of the lower jaw-bone, and adhering to the gums of both jaws, is inferted into the fphincter at the angle of the lips; it brings the food between the teeth in mastication, forces the breath out of the mouth in blowing, &c. and pulls the lips or corner of the mouth outward. Steno's

Steno's falival duct perforates this muscle in the middle.

Q. Which is the platysma myoides?

A. The platysma myoides, vel latissimus colli, is a broad, membranous muscle, exceeding thin, and lies immediately under the skin; it arises from the pectoral muscle below the clavicle, and from part of the deltoide muscle; it runs obliquely forward, covering all the neck, rendering the unequal surface of the muscles even, and is inserted into the chin and depressor muscles of the lip. When it acts, it pulls down the corner of the mouth and the lower jaw: a convulsion herein is called the cynic spasm.

Q. Which is the biventer?

A. The biventer, or digastricus, arises with a fleshy belly from the upper part of the mastoid process; and descending, it contracts into a round tendon, which paffes through the stylo-hyoideus, and an annular ligament, which is fastened to the os hyoides; then growing fleshy again, it ascends towards the middle of the edge of the anterior part of the lower jaw, where it is inserted internally: when it acts, it pulls the lower jaw down by means of this trochlea, or pulley, and ferves to draw up the os hyoides, and parts annexed to it, in deglutition, as well as to prevent the action of feveral muscles which are concerned in swallowing; for which reason we cannot swallow at the fame time that we open our mouth.

Q. Which is the temporalis?

A. The temporalis arises from part of the os frontis, parietale, sphenoides, make and temporale; from whence going under the two processes of the zygoma, it is inserted externally into the processus coronalis of the lower jaw, which it pulls upward: this muscle is covered with a strong tendinous fascia.

Q. Which is the maffeter?

A. The masseter is a thick, sleshy muscle, situated at the back part of the cheek; it arises from the interior part of the os malæ and the zygomatic process, which joins this from the temporal bones: it is inserted into the angle of the lower jaw, which it pulls upward and forward: over this muscle passes Steno's salival duct.

Q. Which is the pterygoideus internus?

A. The pterygoideus internus arises from the pterygoide processes, and is inserted internally into the angle of the lower jaw, which it pulls upward.

Q. Which is the pterygoideus externus?

A. The pterygoideus externus arifes from the os maxillare and os sphenoides, and is inserted internally into the condyloide process of the lower jaw, which it pulls to one side, and forwards, or, acting with its partner, pulls the jaw directly forward.

Q. Which are the muscles of the palate?

A. The muscles of the palate are the pterygostaphylinus internus and externus, thyrostaphylinus, and glossostaphylinus.

Q. Which Q. Which is the pterygostaphylinus inter-

nus?

A. The pterygostaphylinus internus arises from the os sphenoides, near the tuba Eustachiana, and is inserted into the uvula, which it pulls up while we breathe through the mouth, or swallow.

Q. Which is the pterygostaphylinus exter-

nus?

A. The pterygostaphylinus externus, arises by the side of the last mentioned muscle, and is also inserted near it; but becomes its antagonist by being reslected on a pulley over a process at the lower part of the pterygoidal processes of the sphenoidal bone.

Q. Which is the thyroftaphylinus?

A. The thyrostaphylinus arises from the lateral part of the thyroide cartilage; and ascending towards the uvula, is inserted in the manner of an arch in the side of the velum palatinum.

Q. Which is the gloffo-staphylinus?

A. The gloffo-staphylinus passes from the tongue to the palate, which it pulls down when we breathe through the nose. The palate itself is a sort of double muscle, whose action seems only to support itself, and affist those muscles which pull it upwards.

Q. What are the muscles of the os hyoides?

A. The os hyoides is moved by five pair of muscles, viz. the geniohyoideus, sternohyoideus, mylohyoideus, coracohyoideus, and stylohyoideus.

Q. What is the geniohyoideus?

A. The geniohyoideus arises from the forepart of the lower jaw internally, and is inserted into the basis of the os hyoides, which it pulls upward and forward.

Q. Which is the sternohyoideus?

A. It is an antagonist to the last-mentioned muscle, and arises from the inside of the under part of the clavicle near the sternum; and ascending above the geniohyoideus, is inserted into the basis of the os hyoides, which it pulls downward.

Q. Which is the mylohyoideus?

A. The mylohyoideus arises from the inside of the bottom of the lower jaw, under the dentes molares, and is inserted into the basis of the os hyoides. Its common use is to move the os hyoides, tongue, and larynx, both upwards, inwards, and sideways; and when it is at rest, it has a farther use, viz. to compress the glands under the tongue, and by this means promote the discharge of the saliva into the mouth from the lower salival ducts; whence it is we use this muscle when we want saliva in the mouth.

Q. Which is the coracohyoideus?

A. The coracohyoideus is an antagonist to the last-mentioned muscle, and arises from the upper edge of the scapula; and passing obliquely under the mastoideus, is inserted into the basis of the os hyoides, and draws it obliquely downwards.

Q. Which is the ftylohyoideus?

A. The stylohyoideus arises from the styloide process,

process, and is inserted in the horn and base of the os hyoides, which it pulls upward and backward. This muscle is often perforated about the middle, by the tendon of the digastric muscle of the lower jaw.

Q. What are the muscles of the tongue, befides those of the os hyoides, already men-

tioned?

A. The muscles of the tongue (besides those of the os hyoides already mentioned) are, genioglossus, basioglossus, ceratoglossus, styloglossus, chondroglossus, lingualis, and the tongue itself.

Q. Which is the geniogloffus?

A. The genioglossus arises from the forepart of the lower jaw internally, and is inserted broad into the under part of the root of the tongue. When this muscle and its fellow act, they pull the tongue forwards, and thrust it out of the mouth.

Q. Which is the basioglossus?

A. The basing loss arises from the basis of the os hyoides, and is inserted into the tongue near its apex or tip. Its use (with the affistance of the ceratoglossus) is to draw the tongue backward, and make it shorter.

Q. Which is the ceratogloffus?

A. The ceratogloffus arises from the horn of the os hyoides; whence it has its name. It is inserted laterally into the tongue, near its root, and draws the tongue obliquely on one side; but if both act at once, the tongue is pulled directly backwards into the mouth.

G Q. Which

Q. Which is the styloglossus?

A. The styloglossus arises from the apex of the styloide process, and is inserted into the root of the tongue, which it moves upward and backward.

Q. Which is the chondrogloffus?

A. The chondroglossus arises from the cartilaginous process of the os hyoides, meeting in the basis of the tongue, where they are inserted; but this pair is not found in all subjects.

Q. Which is the lingualis?

A. The lingualis arifes pretty large and fleshy from the basis of the tongue laterally, and runs strait forwards to its apex or tip. Its use is to contract or narrow the substance of the tongue, and at the fame time to bring it backwards and downwards.

Q. What is the tongue itself?

A. The tongue itself is a muscle made up of fibres longitudinal, circular, and transverse. This is the interior part of the tongue; those I have before mentioned form the exterior part of the tongue, being inferted in it, and forming one body.

Q. What are the muscles which move the

head on the trunk?

A. The muscles which move the head on the trunk are the rectus internus major and minor, the rectus posticus major and minor, rectus lateralis, and the obliquus inferior and superior.

Q. Which is the rectus internus major?

A. The rectus capites internus major anticus, arises from the anterior part of the trans-

verte

verse apophyses of the five lower vertebræ of the neck, and passing over the two superior vertebræ is inserted into the os occipitis. This muscle bends the head forward on the first and second vertebræ of the neck.

Q. Which is the rectus internus minor?

A. The rectus capitis internus minor anticus, arises from the anterior surface of the atlas or first vertebra of the neck; it is inserted into the os occipitis. This muscle also moves the head forward on the first vertebra.

Q. Which is the rectus major posticus?

A. The rectus capitis major posticus, is one of the extensors of the head, which arises from the spinal apophysis of the second vertebra of the neck, and is inserted into the lower part of the os occipitis. It pulls the head back on the first and second vertebra.

Q. Which is the rectus minor posticus?

A. The rectus capitis minor posticus is also an extensor of the head, having its rise from the posterior part of the atlas, and is also inferted in the os occipitis, to pull the head back on the atlas or first vertebra.

Q. Which is the recturs lateralis?

A. The rectas lateralis arises from the anterior part of the transverse process of the atlas; and is inserted partly into the os occipitis, and partly into the os temporis. This turns the head on one side.

Q. Which is the rectus capitis obliquus in-

ferior?

A. The obliquius capitis inferior five major, arifes from the spinal apophysis of the second vertebra of the neck, and is inserted into the transverse apophysis of the first vertebra. When this acts, it turns the head with the atlas in a rotatory manner on the second vertebra.

Q. Which is the rectus capitis obliquus fu-

perior ?

A. The obliquus capitis superior sive minor, arises from the end of the transverse apophysis of the atlas, and is inserted into the os occipitis, and back part of the os temporis. When but one of these acts, it assists the rectus lateralis on the same side; but when they act both together, they pull the head back.

Q. What is contained in the head, or what is

the viscera of the head?

A. The parts contained in the cranium are the brain.

Q. What is the brain?

A. The brain is a foft white mass, in which all the organs of sense terminate, and the soul is supposed to reside. The brain is surrounded by two membranes called meninges and matres: these are the dura mater, and the pia mater; some make the external lamina of the latter to be a distinct membrane called arachnoides. The general mass is divided into three particular portions, viz. the cerebrum, or brain, properly so called; the cerebellum; and the medulla oblongata. And some add a fourth, viz. the medulla spinalis; which is a continuation of the

the medulla oblongata, and fills the great canal of the spina dorsi.

Q What is the cerebrum?

A. The cerebrum or brain, properly fo called, is a kind of medullary mass, of a greyish colour on its outer furface, filling all the cavity of the upper and fore part of the cranium, which lies above the transverse septum. The upper part is of an oval figure, like half an egg cut lenthways, or rather like two quarters of an egg, and parted a little from each other: it is flatter on the lower part; each lateral half of which is divided into three eminences called lobes; one anterior, one middle, and one pofterior. The human brain is three times as much in quantity as the brain of an ox; it being in general about four pounds weight. The substance of the cerebrum is of two kinds. The exterior or cortical part, and the interior or medullary part. The former is about a fixth of an inch in thickness, softish, and of a greyish or ash colour; the latter is more folid, and very white, and terminates in the beginning of the nerves. Besides the lobes I have already mentioned, the upper fide of the cerebrum is divided into two hemispheres by the processus falciformis of the dura mater; and its lower fide into four lobes; two anterior and two posterior: the latter much the largest. At the meeting of the four lobes appears the infundibulum, which is a kind of lymphatic running from the ventricles of the brain, piercing the dura mater upon the basis of the skull, and finks finks into the substance of the glandula pituitaria; which gland is feated in the fella Turcica. Between the two hemispheres is the corpus callosum, a hard white substance. The ventricles of the brain are cavities lined with a fine smooth membrane, from the pia mater. They are four in number, and all communicate with one another. The two called anterior, lateral, or fuperior are very extensive: the other two, called the third and fourth, are very small in comparifon with these. The lateral ventricles are divided by an extreme thin membrane or partition, called feptum lucidum under which is the fornix the hinder part of which divides into two parts called the crura fornicis or pedes hypocampi. In the basis of the lateral ventricles are eight prominences; two anterior, called corpora striata, from their ash colour lines; and the two thalami nervorum opticorum, fo named because these nerves arise chiefly from them. The other four are small; two anterior called nates cerebri, and two posterior called testes cerebri. Upon the beginning of the thalami nervorum opticorum, are fituated a number of blood veffels, glands, and lymphæducts, called plexus choroides. Under the beginning of the tornix is a foramen called iter ad infundibulum; and under its middle, one called foramen pofterius, which is covered with a valve named membrana vel valvula magna cerebri.

Q What is the cerebellum?

A. The cerebellum lies under the posterior lobes of the cerebrum or brain, in the lower

part of the cavity of the cranium; its figure is nearly globular, its fuperficies is not so circumvoluted and winding as that of the cerebrum, but furrowed, depressed, and largest in the middle, terminating in the vermiform process. The substance of the cerebellum is much the same as that of the cerebrum, only its cortical part is more in quantity than the medullary; the latter is elegantly branched out like shrubs or little trees, the trunks of which are the basis of the cerebellum, and are what are called its peduncles; the lobules of the cerebellum adhere in clusters to the arbusculi medullares, are surrounded by the pia mater, and compose the far greater part of the cerebellum.

Q. What is the medulla oblongata?

A. The medulla oblongata is a medullary continuation of the medullary part of the cerebrum and cerebellum, formed into a kind of tail, and extended to the great foramen in the os occipitis, where it gives origin to the medulla spinalis, and to the nerves of the brain, where it divides, or appears like two bodies, is called crura medullæ oblongatæ; its union, isthmus, and the eminence beyond it processus annularis. The medulla oblongata is a third general part of the encephalon or whole mass of the brain, cerebrum, and cerebellum.

Q. What is the medulla spinalis?

A. The medulla spinalis is a continuation of the medulla oblongata, through the great foramen of the skull, and through the bony canal of the spina dorsi, to the extremity of the os facrum;

facrum; its thickness in general is nearly equal to that of a finger; but it is not uniformly of the same size throughout. The lower part in the os facrum is called cauda equina from its resemblance. Its substance is nearly the same as that of the medulla oblongata, but fomewhat tougher and more firm; the medullary fubstance is here outwardly, that the nerves may eafily make their way out, and the cortical like part inwardly. To cover the medulla spinalis, next the bony canal of the spine internally, is a very strong tunica, which connects the vertebræ within; then the cellular or adipose coat, (which containing more or less fat, seems destined by nature to foften the former) and the dura and pia mater (which I shall by and by describe.) The dura mater, in its anterior part, is firmly connected with the vertebræ, but its posterior part is loose and fluctuating; the pia mater furrounds every part of the medulla spinalis, and all the nerves that arise from it, and enters also its longitudinal division. The arteries and veins of the medulla spinalis enter at the apertures of the vertebræ, which give passage out to the nerves, and come from the vertebrals of the neck, intercostals, and the lumbar. The nerves of the spine are thirty-two pairs, arising from the medulla, connected by and covered with membranes: the use of the spinal marrow is to give origin to these nerves, which are principally distributed to the limbs and external parts, and to fecrete and prepare a nervous Q. What

Q. What are the coverings of the brain, viz. the cerebrum, cerebellum, and medulla oblon-

gata?

A. Two membranes named dura mater and pia mater, the latter is the innermost; the former is the external membrane, which covers the whole.

Q. What is the pia mater?

A. The pia mater is a thin and exceeding fine double membrane, which immediately, and firmly, involves the brain closely, finks into all its cavities and furrows; its outer membrane is by fome made a diffinct coat, and called arach-The pia mater covers also the medulla spinalis, and its membranes adhere very closely and firmly to one another in the upper part of the head, but much less so with the dura mater. The use of this membrane is to contain the brain, and support the blood veffels, which run here in great number, with a multiplicity of turnings and windings, that the blood may not enter the brain too impetuoufly, and for the veins to unite on, that they may enter the finuses in fewer and larger branches. In short, it seems in a manner wholly composed of blood veffels; whose distribution through all its furrows and anfractuofities ferves also to fecrete proper fluids in the brain, and to form the animal spirits. The arteries are from the internal carotids and vertebrals: some of the veins discharge themselves into the sinuses of the dura mater, and others immediately into the jugular and vertebral veins.

Q. What is the dura mater?

A. The dura mater is a very compact, ftrong, thick membrane, covering the pia mater; it lines the infide of the cranium, and supplies the place of an internal periofteum, firmly adhering to its basis, and but lightly at the upper part, except at the futures; it is spread in all the holes and depressions, and covers all the eminences of the skull, to prevent their being hurtful to the brain. It has three processes; the first, named falx, begins at the crista galli, and runs backwards under the futura fagittalis to the cerebellum, dividing the cerebrum into two hemispheres: its use seems to be to divide the brain so as to render it less liable to be moved in the skull, by any violent motions of the head; and the under fide of the brain is kept steady by the inequalities of the basis of the skull which the brain is exactly fitted to. The fecond process runs from the former to the os petrofum, and prevents the cerebrum from pressing on the cerebellum: from this runs a third process, and both serve also to keep the brain steady. The dura mater receives arteries from the carotids, beautifully ramified like shrubs. Its veins are of two kinds; fome as in other parts of the body, and others of a triangular figure, called finuses, the latter give warmth to the brain. It has nerves for fensation from the fifth and eighth pair of the brain. The dura mater has a motion, faid to be peculiar to itself, and of a muscular . kind; but it is much more natural to suppose

it

it owing to the pulsations of the arteries of the brain. There are also a number of small glands in the sinuses and sides of the dura mater, and between it and the pia mater, described by Pachonius, and seem destined for the secreting of a sluid to moisten the dura mater.

Q. What are the arteries of the brain?

A. The arteries that fupply the cerebrum, cerebellum, and medulla oblongata come partly from the carotids, which enter the cranium through the canals in the apophysis petrosæ, and partly from the vertebrals which enter by the occipital foramen. I shall be more particular when I speak of the arteries of the head.

Q. What are the veins of the brain?

A. The veins of the brain are branches of the sinuses of the sinuses of the dura mater, already mentioned: their principal ramifications accompany all the cortical circumvolutions of the cerebrum, and directions of the strata of the cerebellum, running always in the duplicature of the pia mater: the veins of the plexus choroides in general, are of the number of these abovementioned, of which I shall be more particular hereafter

Q. What are the nerves of the brain?

A. In the lower part of the medulla oblongata, are distinguished the nerves of the brain, which are commonly said to be ten pair, though in reality only nine, as follows in Latin verses:

Olfaciens, cernens, oculosque movens, patiensque, Gustars, abducens, audiensque, vogansque, loquensque.

I shall also be more particular when I come to mention the nerves of the head hereafter.

Q. What is the glandula pinealis?

A. The glandula pinealis is a small gland, situated behind the thalami nervorum opticorum, in the third ventricle of the brain, adhering very closely to the plexus choroides, by which it is covered, called pinealis, from its resembling a pine apple.

Q. What is the glandula pituitaria?

A. The glandula pituitaria is a gland of the brain, about the bigness of a very large pea, partly greyish, partly reddish, and white within; it is seated in the sella of the os sphenoides, between the sphenoidal folds of the dura mater, and is covered by the pia mater as by a bag, the opening of which is the extremity of the infundibulum, from which it receives a lymph or juice, which the infundibulum derives from the plexus choroides and pineal gland; from this lymph, the gland takes its name; it also filtrates a juice itself, separating from the blood a white liquor, very subtile, and apparently very spirituous.

Q. What are the arteries of the head?

A. The arteries of the head, both external and internal, proceed from the carotids, cervicals, and vertebrals; and their branches are called by the names of the parts they are bestowed upon, as linguales, temporales, occipitales, &c. The origin of the arteries of the head I shall speak of hereafter, in describing those of the neck and trunk.

I. Arteria

out

I. Arteria carotis externa, or external carotid artery is anterior, and fmaller than the internal carotid; its trunk runs up between the external angle of the lower jaw, and the parotid gland, which it supplies as it passes; afterwards it ascends on the foreside of the ear, and ends in the temple. In this course it sends off five principal internal branches, and three principal external branches. The internal branches are, 1. Sublingualis vel ranina, to the muscles of the os hyoides, the tongue and glandulæ fublinguales. 2. Maxillaris inferior, to the maxillary, parotid, and fublingual glands, styloide, and mastoide muscles, muscles of the pharynx, and to the small flexors of the head. 3. Maxillaris externa, to the masseter, middle of the lower jaw, angles of the mouth, buccinator, and elevator menti, and a particular one to the sphincter oris, which forms a kind of coronaria labiorum, and from thence it goes to all parts of the nose, and, lastly, to the great angle of the eye, where it is ramified and loft on the musculus orbicularis palpebrarum, superciliaris. and frontalis, and is named arteria angularis. 4. Maxillaris interna, to the muscles of the palate, glandular membrane of the posterior nares, and to all the parts contained within the orbits of the eyes. A fmall branch then enters. the cranium through the sphenoidal fiffure, and is spent upon the dura mater; another small branch goes to the maxillary finus and teeth; a principal branch runs through the canal of the lower jaw to the alveoli and teeth, and goes

out at the hole near the chin to the neighbouring muscles; a third principal branch goes up between the external and internal carotid to the dura mater. 5. A principal branch goes to the maffeter muscle. The external branches are, 1. Occipitalis, to the muscles and integuments of the os occipitis, and a branch to the maftoide foramen. 2. A principal branch supplies both the external and internal ear. 3. The trunk of the external carotid runs up (as I have before observed) between the external angle of the lower jaw and parotid gland, which having supplied it, forms the temporal artery, which divides into an interior, middle, and posterior branch; the anterior to the musculus frontalis, and fometimes to the internal apophysis of the os malæ all the way to the orbits; the middle one to the musculus frontalis and occipitalis; and the posterior to the occiput. All these branches likewise furnish the integuments?

II. Arteria carotis interna, the internal carotid artery having passed the great canal of the apophysis petrosa of the os temporis, sends off a branch through the sphenoidal fissure to the orbit and eye, and foon after another through the foramen opticum, by which it communicates with the external carotid. It then runs under the basis of the brain to the side of the infundibulum, where it commonly divides into two branches; the anterior branch runs forward under the brain, and after fending off small arteries to the olfactory nerves, it divides

into

off

into two or three branches; the first of these branches goes to the anterior lobe of the brain; the fecond to the corpus callofum, falx of the dura mater, and middle lobe of the brain; the third to the posterior lobe of the brain: the posterior branch, after communicating with the vertebral artery, is ramified on, and between the superficial circumvolutions of the brain all the way to the bottom of the fulci. All these ramifications are covered by the pia mater, in the duplicature of which they are distributed, and form capillary reticular textures in great numbers; and afterwards they are lost in the inner substance of the brain. From these minute divisions of the arteries in the pia mater before they enter the brain, it would feem as if the pulse of larger arteries would make too violent an impression on so tender and delicate a part: and, perhaps, it may be from an increase of the impulse of the arteries in the brain, which strong liquors produce, that the nerves are fo much interrupted in their functions throughout the whole body, when a man is intoxicated with drinking; and may it not also be from a like cause that men are delirious in fevers.

III. Arteriæ cervicales, arise from the subclavian arteries (I shall hereafter describe) and ascend to the head through the foramina, in the transverse processes of the cervical vertebræ, and into the skull through the tenth or great foramen, and pierce the dura mater; these two arteries uniting soon after their entrance, give off branches to the cerebellum, and then paffing forward, divide and communicate with the carotids; and the carotid arteries communicating with each other, there is an entire communication between them all. These cervical arteries Winslow calls arteriæ vertebrales.

Q. What are the veins of the head?

A. The veins of the head are the jugular and the cervical or vertebral veins. Their origin I shall mention hereafter. 1. Venæ jugulares externæ, or the external jugular veins, are sometimes double from their origins; and when they are fingle, each of them divides afterwards into two, one anterior, and the other posterior, or rather superior. The anterior branch is often a branch of the internal jugular, and fometimes arises from the communication of the two jugulars, and fometimes, but very rarely, from the vena axillaris. It runs upward to the lateral parts of the lower jaw, between the angle and the chin, and fends feveral branches forwards, backwards, and inwards; forwards to the maxillary glands, digastic muscle, muscles and integuments of the chin and under lip; backwards, it fends a finall branch a little below the lower jaw, which communicates with the jugularis externa posterior; inwards to the glandulæ fublinguales, to the tongue, called venæ raninæ, to the muscles of the angles of the lips and neighbouring parts; also to the muscles of the palate, septum palati, amygdalæ, uvula, and to the membrane which lines the arch of the palate. The trunk of the

the anterior external jugular vein goes from the angle of the lower jaw to the internal angle of the orbit, fending branches on each fide to the muscles and integuments of the face, and takes the name of vena angularis. This trunk fends off branches to the upper lip, cartilages, and part of the nofe, eye-lids, and forehead, which last is called frontalis; a branch also communicates with the finuses of the dura mater, entering the orbit by the orbitary finus of the eye. The posterior branch or superior external jugular vein, runs up toward the parotid gland, and lower anterior part of the eye, giving out feveral confiderable branches toward each fide: at its origin a principal branch is fent out posteriorly, called vena muscularis, and a little higher up the vena cervicalis, (which I shall speak off hereafter) backward, it detaches the vena occipitalis to the occiput, and fends a small vein through the posterior mastoide hole, which terminates in one of the lateral finuses of the dura mater. It then communicates with the anterior external jugular, under the angle of the lower jaw, and passes through the parotid gland, sometimes giving off feveral branches, which very foon unite together, and form areolæ or meshes, through which the nerves pass. Afterwards it passes before the ear, taking the name of vena temporalis, which is distributed to the temples and lateral parts of the head, towards the occiput and forehead, sending branches also to the temporal muscle, to the neighbouring parts of the upper jaw, and to the infide of the lower

jaw. The branches of the external jugular all communicate with one another, and with the jugularis interna. 2. Vena jugularis interna, the internal jugular vein, is the largest of all those that go to the head (its origin I shall mention hereafter). This vein detaches a branch up toward the parotid gland, and angle of the lower jaw, where it fends off branches to the muscles of the os hyoides, and sometimes a branch called vena maxillaris interna. Another branch is fent backward to the occiput, communicating with a branch of the vena vertebralis, and with the lateral finus of the dura mater. Most of these branches communicate with the external jugulars. 2. Vena vertebralis, the vertebral vein (whose origin I shall mention hereafter) proceeds to the foramen occipitale, and communicates with the occipital veins and occipital finuses of the dura mater. This vein fends branches to the small interior muscles of the head, and sometimes, though not always, a branch communicates with the lateral finus of the dura mater.

Q. What are the nerves of the head?

A. The nerves of the head are ten pair proceeding from the encephalon as I have before observed, page 36.

Q. What are their names, from whence their

origin, and how distributed?

A. 1. The first pair are the nervi olfactorii, or olfactory nerves, which arise from the corpora striata of the brain, between the anterior and middle lobes; they go out through the foramina

foramina of the os cribriforme, and are immediately spread on the membrane which covers the os spongiosum, and lines all the internal parts of the nose; they communicate with the nervi opthalmici and maxillaris superior. These nerves are the immediate instruments of smelling. 2. The fecond pair are the nervi optici or optic nerves, which arise from the eminences of the cerebrum, called thalami nervorum opticorum; they pass out through their proper hole in the fphenoide bone, and enter the globe of the eye, to be expanded upon the membrana retini. The blood veffels running through the middle of these nerves, and the ramifications of the retina, are very ferviceable, whence we may deduce the reason of Picard's experiments of fuch objects as fall on the entry of the optic nerve being loft to us; and hence also an account may be given of an amaurofis, or gutta ferena. 3. The third pair are the nervi motores oculorum, which arife at the anterior part of the processus annularis, and going out at the foramen lacerum, are distributed to the globe of the eye and its muscles. 4. The fourth pair, are the nervi pathetici, and are the smallest of any; they arise from the anterior lateral part of the processus annularis of the medulla oblongata, go out at the foramina lacera, and are entirely fpent on the musculi obliqui superiores oculorum vel trochleares; and as those muscles act in ogling, staring, &c. their nerves are named pathetici. 5. The fifth pair, are the nervi gustatorii vel trigemini, and are the biggeit

gest of the brain; they arise from the sides of the annular process, giving nerves to the dura mater, then each divides into three branches; the first branches helps to form the intercostal, and then goes to the orbit, (by the name of orbitarius vel opthalmicus) lachrymal gland, fat, membranes, and palpebræ of the eye, membrana narium, the muscles and integuments of the forehead. Hence we eafily discover what parts is affected in the megrim (which is a disease causing great pain in the temple and fore part of the head) when the eye-hall and forehead are racked, and a heat is felt within the nose. Hence also we may learn how the muscles of respiration come to be so much affected on the application of any acrid irritating fubstance to the membrana narium, as to produce that violent convulfive motion fneezing; the fecond branch, or maxillaris superior, passes out through the round foramen of the sphenoide bone, and immediately gives nerves to the fat under the temporal muscle, palate, sinus sphenoidalis, and nostrils; the remaining trunk supplies the antrum Highmorianum (which is a cavity in the maxillary, or jaw bone) and teeth of the upper jaw, then comes out at the orbiter externus hole, and is spent on the musculus orbicularis palpebrarum, nose, and upper lip. The third branch, or maxillaris inferior, goes out at the fourth hole of the sphenoidal bone, and soon splitting into a great many branches, is distributed to the musculus temporalis, masseter, pterygoides, digastricus, buccinator, mylohyoideus.

hyoideus, geno-hyoideus, genioglossus, and basioglossus, glandula sublingualis, maxillaris inferior, and parotis, to the external ear, where it feems to join the portio dura to the substance of the tongue, in which it is pretty much confounded with the ninth pair; from the root of this last branch, the chorda tympani is reslected. The last ramification of this branch, which I shall take notice of, is that which furnishes the teeth of the lower jaw, and comes out at the chin, and is distributed on that and the lower lip, and again united to the seventh pair. From this short account of the large fifth pair of nerves, and by observing several phænomena which happen to those parts to which they are distributed, we might have a much farther confirmation of the general doctrine of nerves delivered, and fee, at least, the way pathed to a rational account of the phænomena, for reasoning on which we should not otherwise have the least ground. We can, for example, from the chorda tympani and the nerves of the teeth being derived from the fame common trunk, understand how the found of any vibrating body held between our teeth is fenfible to us, when another person cannot possibly hear it; by the like rule we know why, in a violent tooth ach, the muscles of the face are sometimes convulled; nor shall be furprized to hear one plagued with the ach in his upper teeth, complain of a gnawing pain deep feated in the bones of his face, or to fee his eyelids much fwelled, or the tears trickling down in great abundance; whereas the lower teeth ach-H 3

ing, the ear is pained, and the faliva flows in great quantity. We may have a distant view of some foundation in reason for the cure of the tooth-ach by strong compression of the chin, or by applying blifters behind the ears, or by burning behind or in the ear. Among a great many instances of the good effect of the actual cautery in fuch a case, I shall give one which feems to me remarkable: A man was seized with the tooth ach, and a convulsion of that whole side of his face followed whenever the pain became acute, or he attempted to speak; after he had undergone bleeding, purging, falivation, fetons, &c. without any benefit, he was cured by applying a small cauterising iron to the anti-helix. 6. The fixth pair of nerves are the nervi abducentes, which arise from the fore part of the corpora pyramidalia (which are two protuberances of the under part of the cerebellum. so called from their resemblance of a pyramid) and after piercing through the dura mater, they give off a branch, which joining with the reflected twig of the opthalmic branch, forms the original of the intercostal, and passes through the foramen lacerum, to be spent entirely on the mufculus adductor oculi. Suppofing this nerve to supply ever so little less than a due proportion of liquidum nervolum, an involuntary itrabismus, or squinting, will be occasioned. Though the fifth and fixth pair of nerves form entirely the beginning of the intercostal before it goes out of the skull; yet as several other nerves contribute towards the formation of its trunk. trunk, before it fends off any branches, I shall postpone the description of it till I have spoke of the original nerves. 7. The seventh pair are the nervi auditorii, which arise from the side of the root of the processus annularis, and entering the meatus auditorius internus, and immediately dividing, one part is expanded on the inmost camera of the ear, the other goes through the aquæductus Fallopii, and comes out of the skull between the styloide and mastoide processes; whence the reason of the one being named portio mollis, and the other dura. This last, after its exit; supplies the musculi obliqui capitis, stylohyoidei, styloglossi, stylopharyngæi, and platysma myoides, on which and to the skin of the neck, a great number of its small filaments run, which are sometimes cut in opening the jugular vein, whence follows pain at first, and a little numbness afterwards: the superior branches of this nerve supply the parotid gland, external ear, and the whole fide of the face as far forward as the chin. It is faid to communicate thrice with the fifth pair, and twice with the second vertebra. May not we hence fee some reason why the head is fo foon moved by the impression of found on our ear? 8. The eighth pair are the nervi sympathetici medii vel par vagum, which arise from the fide of the basis of the corpora olivaria, (which are two protuberances, of the medulla oblongata, fo called from their representing an olive in shape) runs to the hole common to the offa temporum and occipitis, and are there joined. H 4

joined by the accessorius Willisii, (which has its beginning from the two or three superior nerves of the medulla spinalis) mounts up and passes out with the par vagum at that common foramen just now mentioned, then separating the accessorius, goes through the musculus mastoideus, and is lost in the trapezius, and rhomboides scapulæ; while the larger trunk, (from the great number of branches it fends off, obtains the name of vagus,) runs strait down the neck near the carotid artery, and in its courfe gives feveral branches to the larynx; when entered the thorax, it splits into two, the anterior branch goes to the pericardium, and with those of the intercostal to the heart, then on the right fide turns round the fubclavian, and on the left round the ductus arteriosus, and goes up again at the fide of the cefophagus to be loft in the larynx. This recurrent branch it is, that we are earnestly cautioned to avoid in bronchotomy, though by its deep situation we are in no hazard of hurting it: if both these nerves were cut, it is probable the voice would not be entirely loft, as long as the superior branches still supply the The posterior branch goes along with the cefophagus, and fupplies the lungs, gula, and stomach very plentifully: and as all the nerves bestowed on the stomach enter at the fuperior orifice of it, the fensations here must be very acute, what remains of the par vagum is joined to the intercostal immediately below the diaphragm. 9. Nervi hypoglossi externi vel par linguale, which arise from between the corpora

pora pyramidalia and the corpora olivaria, passing out of the skull through their proper holes of the os occipitis, and after supplying the glandula thyroidea, and musculi sternohyoidei, and sternothyroidei, are lost in the substance of the tongue. Authors have disputed whether this ninth or the fifth is the gustatory nerve; the old opinion in favour of the ninth is to me most probable, because the fifth is no where else employed as an organ of sensation, and because the ninth feems to penetrate the fubstance of the tongue more, while the fifth is spent on the muscles. 10. The tenth pair are the nervi suboccipitales, which arise from the beginning of the medulla spinalis, betwixt the os occipitis and first vertebra colli; and are all, except what goes to the ganglion of the intercostal, spent on the musculi obliqui and extensores capitis. But these, as before observed, page 30, are more properly the first pair of nerves of the neck.

The only nerves of the encephalon remaining now to be described are the reslected branches of the fifth and sixth pair, which are not easily traced, being so small and pappy, and hid by the carotid artery as they go out of it; but whenever they have escaped from the os petrosum, they are joined by branches from the eighth, ninth, and tenth pair, and the first and second spinal, whence the largest ganglion of the body is formed, from which the nerve named now intercostal goes out to descend down the neck with the carotid, supplying in its course the musculi slexores of the head and neck, and

Q. What are the glands of the head?

A. The glands of the head remaining now to be described, are in the exterior part that is out of the cavity of the skull; those of the interior part, viz. the glands of the brain and its membranes, have been already spoken of. Those now to mention, are the parotides, maxillares, sublinguales, tonsillæ, linguales, labiales, buccales, fauciales palatinæ, gingivarum, and uvulares; which take their names from their respective situations, being situated in and about the mouth, palate, and tongue, to afford saliva in all parts of the mouth to keep it moist.

Those

Those more remote are chiefly concerned in mastication. In the orbit also there is the lachrymal glands; under the eye-lids are the ceraceous or sebaceous glands, the mucose glands of the pituitary membrane of the nostrils, and the ceruminose glands of the ears. The largest and most remarkable are the salival glands; the others are so small as to render a particular description unnecessary and of no significance.

Parotis, vel maxillaris superior, is the largest of the falivary glands; it is fituate behind the lower jaw, under the ear, on each fide; from this gland, there runs a very large duct, about three fingers breadth long, and of the thickness of a wheat straw, having a great number of roots: this duct, from Steno the discoverer, is called after his name, but by others ductus falivalis superior. It passes over the tendinous part of the masseter muscle, (to prevent its being compressed by that muscle, which would obstruct the saliva) through the middle of the cheek, and these perforate the buccinator muscle and the membrane of the mouth, near the second or third of the dentes molares, and at this perforation it discharges a very large quantity of its proper fluid into the mouth. When this duct is divided by an external wound, the faliva will flow out on the cheek, unless a convenient perforation be made into the mouth, and then the external wound may be healed. This gland is one of those that ferve for the fecretion of the faliva; it has the discharge of its saliva promoted by the motions of of the lower jaw: when this gland is ulcerated, there is a constant effusion of saliva; to cure which Hildanus applied the actual cautery, but if you consume the greatest part of the gland with merc. præcip. rubr. it will heal with little trouble.

Maxillaris inferior is situate between the lower jaw and the tendon of the digastric muscle; its duct passes under the musculus mylohyoideus, and enters the mouth under the tongue, near the dentes incisorii.

Sublingualis is a small gland situated under the tongue, one on each side, between the jaw

and the ceratogloffus muscle.

Tonfilla is a globular gland, about the bigness of a hazel-nut, situate upon the pterygoideus internus muscle, between the root of the tongue and the uvula, on each fide of the mouth; they are commonly called almonds of the ears, from their resembling almonds in figure. The tonfilla has no duct continued from it, but empties all its small ducts into a finus of its own; which finus, when the gland is inflamed, may eafily be mistaken for an ulcer. This gland, with its fellow, directs the masticated aliment into the pharynx, and also serves for the uvula to shut down upon when we breathe through the nofe. They are compressed by the tongue and the aliment, when the former raifes the latter over its root, and thereby opportunely emits their faliva to lubricate the food for its easier descent through the pharynx. A schirrous tumour of either of these glands is a common

fecretion

common disease, and it admits of no remedy but extirpation; the best way of doing which is by ligature. Pressure upon the surface of a gland very much promoting the fecretion that is made in it, these glands are so seated as to be pressed by the lower jaw, and its mufcles, which will be chiefly at the time when their fluid is wanted; and the force with which the jaw must be moved, being as the driness and hardness of the food, which is necessary; for all food, being to be reduced to a pulp, by being broke and mixed with faliva before it can be swallowed fit for digestion, the drier and harder foods needing more of this matter, will from this mechanism be supplied with more than moister foods, in about that proportion in which they are drier and harder; and the drier foods needing more faliva than the moister, is the reason why we can eat less, and digest less of these than those. What quantity of faliva these glands can separate from the blood in a given time will be hard to determine; but in eating of dry bread, it cannot be less than the weight of the bread; and many men, in a little time, can eat more dry bread than twice the fize of thefe glands; and some that are not used to smoaking, can fpit half a pint in smoaking one pipe of tobacco. Some men in a fallivation have spit for days, or weeks together, a gallon in four and twenty hours; and yet all thefe glands put together do not weigh more than four ounces.

The glandulæ sebaceæ are situated in the interior surface of the eye-lids; they serve for the

110 ANATOMICAL

fecretion of an oleaginous fluid, which prevents the attrition of the eye-lids from their continual motion.

The glandula lachrymalis is fituated in the orbit, above the smaller angle, with its excre-

tory ducts under the upper eye-lids.

The glandulæ ceruminosæ are small glands of a yellow colour, situate in the convex part of the membrane of the meatus auditorius of the ear, about the middle of the passage; they serve to secrete the cerumen, which they deposit for various purpose in the passage.

Glandulæ mucosæ are situated in the pituitary membrane of the nostrils, and separate

that matter which we call mucus.

DIALOGUE III.

Of the NECK, and its PARTS.

A. The fore part of the neck?

A. The fore part of the neck is the throat and the back part the nape. The neck confifts of feven bones or vertebræ; a number of muscles which serve to move the head, neck, larynx, pharynx, and the os hyoides; a number of very large arteries, as the internal and external carotids, and the vertebral ones; large veins, as the internal and external jugulars, and the vertebral veins; large nerves of the par vagum, intercostals, recurrents, diaphrag-

phragmatics, and the vertebral; a part of the spinal marrow; the aspera arteria or trachea, particularly the larynx; the pharynx, with part of the cesophagus; and the thyroide, with the other smaller glands. The bones of the neck are named vertebræ.

Q. What are the vertebræ?

A. The vertebræ or joints of the neck and back from the os occipitis to the os facrum, are twenty-four in number, those of the neck belong also to the spine forming one bony column for the reception of the spinal marrow; each vertebra is diffinguished into two parts, viz. the body and its processes. The body of each vertebræ, except the first, is softer and more spongeous than the processes, which are harder and more folid; the fore part of the body is round and convex; the hinder part fomewhat concave, its upper and lower parts are covered with a cartilage pretty thick forwards, but thin backwards, by which means we bend our body forwards; for the cartilages yield to the pressure of the bodies of the vertebræ, which in that motion come closer to one another. This could not be effected, if the harder bodies of the vertebræ were close to one another. Each vertebra has three forts of proceffes, towards its hinder part, except the first: from the hind part of each stands a process named spinalis, and from every one a process on each fide called transversalis, with one short one above it, and so below it named obliqui superiores et inferiores. By these short oblique processes

processes the vertebræ are articulated; and in each of the transverse, there is a tendon of the vertebral muscles inserted. These processes, with the hinder or concave part of the body of the vertebræ, form a large hole in each vertebra, and all the holes answering one another (from the head to the os coccygis) make a channel for the descent of the spinal marrow, which fends out its nerves to the feveral parts of the body by pairs through two small lateral holes in each space, between the vertebræ formed by the joining of four notches in the fide of each superior and inferior vertebræ. The vertebræ are articulated to one another by a ginglymus; for the two inferior oblique proceffes of all the vertebræ of the neck and back, have a little cavity-like in their extremities, wherein they receive the extremities of the two superior oblique processes of the inferior vertebræ next to it; fo that the two fuperior processes of each vertebra of the neck and back are received, and the two inferior do receive, (except the first of the neck and last of the back;) but the superior processes of each vertebra of the loins receive, and the two inferior are received, contrary to those of the neck and back. The vertebræ are all tied together by a hard membrane, made of strong and large fibres; it covers the body of all the vertebræ forwards, reaching from the first of the neck to the os facrum. There is another membrane which lines the canal, made by the large hole of each vertebra, which also ties them all togetogether; besides the bodies of the vertebræ are tied to one another by the intervening cartilages, and the tendons of the vertebral muscles which are inferted in their processes, as before mentioned, tie them together behind. From this account of the articulations of the vertebræ, it is evident their center of motion is altered in different positions of the trunk. For when we bow forwards, the fuperior moved part, bears entirely on the bodies of the vertebræ; if we bend back, the oblique processes support the weight; if we recline to one fide, we rest upon the oblique processes of that side, and part of the bodies of the vertebræ; if we stand erect, all the bodies and oblique processes have their share in our support. From this structure, the extensors have about twice the lever to act with, and consequently, twice the power to raise the trunk into an erect posture that they have to carry it beyond that posture; for then the oblique processes begin to be the center of motion, and give a like advantage to the benders; without this contrivance it would be more difficult to keep the body erect, or to recover an erect posture with considerable strength after a bend of the body. If the spine had been composed all of one bone, we could have had no motion in our backs; or had it been of two or three bones, or fewer bones or joints than it really is, they must have either not been capable of bending so much as they do, (and been so pliable for the several postures we have occasion to put ourselves in) or have bent more in each joint, which would have have pressed or bruised the spinal marrow; the ill consequences of which are sufficiently seen in persons grown crooked, or who have had distortions from external accidents. Again, if the spine had been made of several bones, without intervening cartilages, we should have had no more use of it, than if it had been but one bone; if each vertebra had had its own distinct cartilage, it might have been easily dislocated; and, lastly, the oblique processes of each superior and inferior vertebra keep the middle one, that it can neither be thrust backwards nor forwards, to compress the medulla spinalis.

Q. What are the bones of the neck?

A. The bones of the neck are the feven fuperior vertebræ of the spine, (I have already mentioned); these vertebræ are smaller than those of the back, but they are of a firmer, harder confistence: their body is more compressed than in the others, and is situated on the other part, and convex below. The breadth on the fore part increases gradually as they defcend; fo that the vertebræ of the neck taken all together represent a fort of pyramid. transverse processes of these vertebræ are perforated for the passage of the vertebral vessels to the head: and the acute or spinal processes are forked and strait; but besides this, the first and fecond vertebræ have fomething peculiar to themselves. The first, or upper vertebra is called Atlas (because it supports the head, as king Atlas did the globe of the universe, according to antient fable;) it has neither body

nor spinal process, for which reason its substance is more folid than that of any other; the foramen or hole in this, is greater than in any other vertebra, and it looks like an irregular bony ring. The upper fide of this vertebra has two cavities, into which the apophyses of the os occipitis are received; but thele two cavities together, unlike all other joints, are laterally portions of concentric circles, by which means they are but as one joint, and fo fuffer the head to move eafily fideways, which otherwise it could no more do than the knee, which also has two heads and two cavities. The under fide of this bone has a very flat articulation with the next, which fits it for a rotatory motion. In the fore part of its great hole it has a pretty large finus, in which lies the tooth like procefs of the fecond vertebra, being fastened by a ligament that rifes from each fide of the finus, that it compress not the medulla spinalis; it has two fmall finuses in its upper part, in which the tenth pair of nerves and the vertebral arteries lie. The fecond vertebra is called epiftrophæus, dentata, or axis, from a process which passes through the former bone, and is the axis upon which it turns; nevertheless, all the vertebræ of the neck contribute fomething to the rotatory motion of the head. The processus dentatus, which is long and round like a tooth, from whence its name, is fituate in the middle, between the two oblique superior processes; it is received into the aforesaid sinus of the atlas, and is strongly tied to the os occipitis, and to the the atlas by ligaments, to prevent its hurting the spinal marrow. All the rest of the vertebræ of the neck are alike. See the account of the vertebræ in general, page 111.

Q. What are the cartilages of the neck?

A. The cartilages of the bones of the neck, and of all the vertebræ in general, are of two kinds, one proper to each vertebra, the other common to the two vertebræ that lie next each other; the first I term cartilages of articulation, the other cartilages of fymphysis. The proper articular cartilages of each vertebra of the whole fpine are those four which cover the furfaces of the four small oblique or articular apophyses. The cartilages of symphysis lie between the bodies of the vertebræ, one closely joined to each bone; their height and thickness is different in each class of the vertebræ; those of the neck are not fo thick as those of the loins, nor fo thin as those of the vertebræ of the back; nor are the cartilages of an equal thickness in all their parts; those of the neck and loins appear to be the thickest on the fore side, and those of the back rather thickest on the backside. These cartilages are different from all others in the body, being made up of horizontal, concentrical rings, closest, and thinnest near the center, refembling the other cartilages of the body in nothing but their whiteness and elasticity; the interstices between the rings are filled with a mucilaginous substance, less fluid than that of the joints. All these cartilages yield to compression, (and in the inflexions of the spine, the external

external furface of the cartilage jets out on that fide toward which the inflexion is made); they restore themselves afterwards, by being freed from compression; so that a man is really taller, after lying some time, than after he has walked or carried a burthen for a great while; owing to the different state of the intervertebral cartilages. The cartilages of the larynx, &c. will be described with the muscles.

Q. What are the ligaments of the neck?

A. All the vertebræ of the spine in general are strongly connected to each other by ligaments, which are very short and strong; they cross each other obliquely, and are fixed round the edges of the body of each vertebra, covering the intervertebral cartilages, and adhering closely to them; but the first and second vertebra have both ligaments of a peculiar kind from the rest. All the vertebræ are likewise strongly connected by a ligamentary tube, which lines the inner furface of the medullary canal, representing a long flexible funnel, it's cavity at the upper part being equal to that of the occipital foramen, and ending in a small point at the os facrum: the articular ligaments of the spine are those which tie the glenoide cavities of the atlas to the condyles of the os occipitis; those that join the cartilaginous surface of the apophysis dentiformis to the anterior cavity of the first vertebra, and those by which all the oblique or articular apophyses are connected together: these are all small, short, strong ligaments, fixed by both extremities round the cartilaginous tilaginous furfaces of the apophyses, surrounding very closely all the capsular ligaments of these articulations before mentioned. The membranous ligaments of any signification will be described with the muscles.

Q. What are the mucilaginous glands of the

neck?

A. The mucilaginous glands of all the articulations of the vertebræ of the neck are very small, but are accompanied by many fatty moleculæ lying round each joint; the inner surface of the ligamentary tube just now mentioned, is lubricated by an oily or adipose substance, which I have mentioned already. See page 25.

Q. What are the muscles of the neck?

A. Besides those only peculiar to the neck, there are first those of the head and neck conjointly; then those peculiar to the neck, larynx, epiglottis, and pharynx: those of the head and neck conjointly are, biventer cervicalis, complexus, mastoideus, trachelomastoideus, splenius capitis, and rectus capitis internus major. Those only peculiar to the neck are, interspinales cervicis, intertransversalis cervicis, spinalis cervicis, transversalis cervicis, longus colli, splenius colli; the rest I must omit till I come to describe those of the thorax and abdomen. The mufcles of the larynx are, sternothyroideus, hyothyroideus vel cerato-thyroideus, cricothyroideus, crico-arytænoideus lateralis et posticus, thyro-arytænoideus, arytænoideus, and with thefe I shall first mention the cartilages and membrane

membrane of the larynx. The muscles of the epiglottis, viz. arytæno epiglotticus, and the hyo-epiglotticus, to these I shall add the œso-phagus. The muscles of the pharynx are, stylo-pharingæus, pterygopharingæus, constrictor pharyngei vel æsophagus, palato pharingæus.

Q. What is the biventer cervicalis?

A. The biventer cervicalis arises from the transverse processes of the seven superior dorsal vertebræ, and is inserted into the back part of the os occipitis.

Q. Which is the complexus?

A. The complexus is a broad and pretty long muscle situate along the back part and side of the neck; it arises from the three superior dorsal vertebræ, and the six inferior vertebræ of the neck, and is inserted with the biventer into the os occipitis and back part of the os temporis. It pulls the head and neck back.

Q. Which is the mastoideus?

A. The mastoideus is situate obliquely between the back part of the ear and lower part of the throat; it is in a manner composed of two muscles, (which Albinus terms sternomastoideus, and cleido-mastoideus) though in fact but one: it arises from the sternum and clavicula in two portions, but soon unites in one, and is inserted into the outer part of the processus mastoideus of the os temporis; over this process it sends off a very broad aponeurosis, which covers the splenius, and is inserted in the os occipitis: it pulls that side of the head it is inserted into towards the sternum, and turns the

face towards the contrary shoulder; this muscle, with its fellow, pulls the head and neck toward the breast; but acts with more force on the joints of the neck, than upon the head.

Q. Which is the trachelo mastoideus?

A. The trachelo mastoideus arises from the transverse process of the first and second vertebræ of the back, runs up under the splenius, and is inserted into the middle of the backside of the processus mastoideus.

Q. Which is the splenius capitis?

A. The splenius capitis arises from the spinal processes of the five lower vertebræ of the neck, and the five upper ones of the back, and also the linea alba colli. It is inserted into the os occipitis a little above the transverse processes of the three superior vertebræ of the neck. This muscle pulls the head and neck backward, and to the contrary side; but both of them acting together, pull them directly backward.

Q. Which is the rectus internus major?

A. The rectus capitis internus major I have already described with the muscles of the head.

Q. Which are the interspinales cervicis?

A. They are muscles arising from the superior parts of each double spinal process of the cervical vertebræ, and inserted into the inserior parts of the same. When these muscles act, they bend the neck backward, drawing the spines of the vertebræ nearer each other.

Q. Which are the intertransversales cervicis?

A. They are situate between the transverse process of the vertebræ like the interspinales;

they arise from the lower vertebra of the neck, and are inserted into that next above: these Douglas calls intervertebrales.

Q. Which is the spinalis cervicis?

A. The spinalis cervicis arises from the transverse processes of the five superior vertebræ of the back; and is inserted into the spinal processes of the second, third, fourth and sifth vertebræ of the neck. This pulls the neck backward.

Q. Which is the transversalis cervicis?

A. The transversalis vel semi-spinalis cervicis arises from the oblique processes of the four inferior vertebræ of the neck, and is inserted into the spinal process of the second vertebra of the neck. This is only a continuance of the transversalis dorsi, and moves the neck obliquely backwards, as when we look over the shoulder.

Q. Which is the longus colli?

A. The longus colli arises laterally from the bodies of the four superior vertebræ of the back, and from the anterior part of the transverse processes of the five inferior vertebræ of the neck; and is inserted into the fore part of the first and second vertebræ of the neck, which it bends forward.

Q. Which is the splenius colli?

A. The splenius colli arises from the spinal processes of the ninth and tenth vertebræ of the back, and is inserted into the transverse processes of the fifth, sixth and seventh vertebra of the neck.

122 ANATOMICAL

The rest of the muscles of the neck I shall defer mentioning till I come to speak of those of the thorax and abdomen.

The muscles of the head and neck are most of them obliquely directed; therefore they perform the oblique motions, as well as extension and flexion.

The afpera vel trachea arteria or wind pipe, I shall describe hereafter, with the contents of the thorax.

Q. What is the larynx?

A. The larynx is the thick upper part of the afpera arteria or wind-pipe, principally composed of five cartilages.

Q. What are the cartilages of the larynx?

A. They are five, viz. 1. The thyroide, or fcutiform cartilage, which stands in the anterior part, and is the largest of the five; in the fore part of this cartilage is the protuberance called pomum adami. 2. The cricoide or annular cartilage. 3, and 4. The arytænoide cartilages, which with the cricoide, make the glottis, (that is the mouth of the larynx) more easily open and contract. Between the arytænoides and fides of the thyroides, there are two finall cavities on each side; in which, if a little drink or bread fall, (as fometimes happens, when we laugh or speak when eating or drinking) it causes a violent cough, and a great tickling. 5. The fifth and last cartilage, which is fofter than the rest, is the epiglottis; its use is to cover the glottis, in eating and drinking; for the aliments, by their own weight, press it closer

down upon the glottis, and they pass over without entering the larynx, into the cesophagus or gullet: but when the aliments are passed, the epiglottis by its natural action, (which is common to all cartilages) lifts up again, and gives way to the air in breathing. While we speak, or laugh, the glottis must necessarilly be opened for the passage of the air in breathing; therefore it is not convenient to speak while we swallow.

The membrane which invests the larynx is very sensible, and is furnished with a number of exceeding small glands and oscula or openings, which discharge a lubricating sluid. The ventricles of the larynx are certain hollows, some of them smaller, some larger; they are on the inside of it, under the glottis, and serve to modulate the voice. These, with the dilatation and straitening of the muscles and cartilages of the glottis, give that wonderful variety of notes, the voice is capable of in singing, &c.

Q. Which is the sterno-thyroideus?

A. The sternothyroideus muscle arises from the sternum, and is inserted into the scutiform cartilage which it pulls downward.

Q. Which is the hyothyroideus?

A. The hyothyroideus vel cerato-thyroideus muscle arises from the os hyoides, and is inferted into the scutiform cartilage which it pulls upward.

Q. Which is the cricothyroideus?

A. The cricothyroideus arises from the fore part of the cricoide cartilage, runs under the thyroide

thyroide cartilage, and is inferted into its infide, which it pulls towards the cricoides, and ferves occasionally either to dilate or constringe the glottis.

Q. Which is the crico-arytænoideus?

A. The crico-arytænoideus lateralis arises from the lateral part of the cricoide cartilage; and is inserted into the lateral part of the arytænoides; this with its fellow serves to dilate the remula or glottis.

Q. Which is the crico-arytænoideus pof-

ticus?

A. The crico-arytænoideus posticus arises from the back part of the cricoid cartilage, and is inserted into the arytænoides to pull it backward and dilate the glottis.

Q. Which is the thyro-arytænoideus?

A. The thyro-arytænoideus arises from the scutiform, and is inserted into the fore part of the arytænoide cartilage; it serves together with the following muscle, to constringe the rimula or glottis.

Q. Which is the arytænoideus?

A. The arytænoideus is one fingle muscle, though Winslow and Douglas divide it into two or three. It arises from one arytænoidal cartilage, and is inserted into the other, which forms a sphincter for contracting the rimula, and shutting the glottis.

Q. Which is the epiglottis?

A. The epiglottis has three exceedingly small muscles, (viz. the thyroepiglotticus, arytæno-epiglotticus, and the hyoepiglotticus) which

ferve to cover the glottis, in the act of swallowing, and prevent any thing getting into it; but they are so very minute as to render any fur-

ther description unnecessary.

The œsophagus, gula or gullet, I shall describe hereafter with the contents of the thorax. The pharynx is the upper part of the œsophagus next the mouth, in which are a number of glands situated, and excretory oscula or openings are frequently discovered with them: the muscles of the pharynx serve to open and shut the œsophagus.

Q. Which is the ftylopharingæus?

A. The stylopharingæus muscle arises from the styloide process, and is inserted on both sides into the styloide and thyroide processes: this muscle with its fellow serve to elevate as well as dilate the pharynx to receive the aliments.

Q. Which is the pterygopharingæus?

A. The pterygopharingæus spheno vel salpingopharyngæus, arises from the inner wing of the os sphenoides, and is inserted partly into the pterygoide apophysis, and partly into the cartilaginous portion of the Eustachian tube: its use is to dilate the pharynx, and draw the middle part of it upwards.

Q. Which is the constrictor pharyngei?

A. The constrictor pharyngei, vel cesophagus, arises like a wing from several parts of the skull, tongue, os hyoides, cricoide and thyroide cartilages, and is inserted into the back part of the pharynx, which it draws to the fore part,

and not only constringes the pharynx for pressing down the aliment, but also compresses the tonfillæ, which fend out their liquor to lubricate the aliment, whereby it glides the more eafily down into the stomach.

Q. Which is the palatopharingæus?

A. The palatopharingæus arises from the aponeurosis of the circumflexus palati, and is inferted into the ftylopharingæus. This mufcle more properly belongs to the palate.

Winflow and fome others make many fubdivisions of the muscles of the pharynx, and give them peculiar names, but they are quite

unnecessary.

Q. What are the arteries of the neck?

A. The arteries of the neck are the vertebral arteries, and the internal and external carotids; and their branches are called by the names of the parts they are, bestowed upon, as laryngeæ, pharingeæ, &c.

Q. What are the carotid arteries?

A. The carotid arteries are two in number, as I have before observed (in describing the arteries of the head,) one called the right carotid, the other the left. They arise near each other, from the curvature, or arch of the aorta. The left immediately, but the right most commonly from the trunk of the fubclavia, as I have already observed. They run up on each side of the trachea arteria, between it and the internal jugular vein, as high as the larynx, without any ramification. During this course, they may be called carotid trunks, as each of them are afterwards wards ramified. The trunk having reached as high as the larynx, divides into two large branches, one named the external carotid artery, and the other the internal.

Q. Which is the external carotid artery, and

how distributed?

A. The external carotid artery is anterior, the internal carotid posterior; and the former lies more inward, and nearer the larynx than the latter; but the names of the blood vessels are taken from the parts they are diffributed to, and not from their situation. The external carotid is the fmallest, and yet appears by its direction to be a continuation of the common trunk; its branches may be divided into anterior or internal, and posterior or external. The first anterior or internal branch goes out from the very origin of the carotid on the infide, and fends off branches to the jugular glands, fat and skin, glandulæ thyroideæ, muscles, and other parts of the larynx, therefore termed laringeæ; it likewise sends some twigs to the pharynx. The third anterior branch, or arteria maxillaris inferior, goes to the mastoide muscle, the small flexors of the head, and muscles of the pharynx, and also to the parts just mentioned. The other branches fupplying the head and its parts I have already described. See the arteries of the head.

Q. How is the internal carotid artery distri-

buted?

A. The internal carotid artery leaving the general trunk, passes behind the external carotid,

a little more backward, and generally runs up without any ramification as high as the lower orifice of the great canal of the apophysis petrosus of the os temporis; therefore sends no branches to the neck.

Q. What is the vertebral artery?

A. The vertebral artery arises from the posterior and upper side of the subclavian; it runs up through all the holes in the transverse apophyses of the vertebræ of the neck, and in its passage sends off branches to the aspera arteria, cesophagus, muscles of the pharynx, larynx, jugular glands, and all the muscles and integuments of the neck, which take their names (as I have before observed,) from the several parts they are bestowed on. See arteriæ cervicales.

Q. What are the veins of the neck?

A. The veins of the neck are the internal and external jugulars, and the vertebral veins, which all arife from the subclavians.

Q. What are the external jugular veins?

A. The external jugular veins are sometimes double from their origin; and when they are single each of them divides afterwards into two; one anterior, and the other posterior or rather superior. The anterior in its passage sends branches to the muscles of the larynx, and to part of the mastoideus; besides those sent to the head and its parts. The posterior, a little higher up than its origin, gives off the vena cervicalis to the vertebral muscles of the neck; near the cervical vein sometimes arises the small vena cephalica which running down between the pectoral

pectoral and deltoide muscles unites with the cephalic vein of the arm, which I shall describe hereafter: both the anterior and posterior external jugular run up the neck, between the integuments and the musculus mastoidæus.

Q. How is the internal jugular vein distri-

buted ?

A. The internal jugular vein is the largest of all those that go to the neck: it runs up behind the mastoideus and coracohyoideus, along the sides of the vertebræ of the neck to the fossula of the foramen lacerum of the basis cranii; in its passage it sends off small branches to the thyroide glands, then the vena gutturalis to the thyroide gland larynx, and neighbouring muscles.

Q. What is the vertebral vein?

A. The vertebral vein arises posteriorly from the fubclavian or axillaris, fometimes by two stems, fometimes by one, which foon afterwards divides into two; the first and principal stem gives out the vena cervicalis to the neighbouring muscles, and then runs up through the holes of the transverse apophyses of the vertebræ colli. The other stem runs up on the side of the vertebræ, and communicates with the first, running in between the transverse apophyses of the fourth and fifth vertebra. Thus the vertebral vein accompanies the vertebral artery fometimes in one trunk, fometimes in feveral ftems, through all the holes of the transverse apophyses of the vertebræ, all the way to the great foramen occipitale, communicating with

the occipital veins, and small occipital sinuses of the dura mater: these veins as they pass supply the muscles of the neck, and great canal of the spinal marrow, where they form sinuses, which communicate with those on the other side; these sinuses are pretty numerous, placed one above another all the way to the occiput, communicating with one another, and at last with the occipital sinuses of the dura mater.

Q. What are the nerves of the neck?

A. The nerves of the neck proceed from the medulla spinalis, and are seven pair; though by some reckoned eight, as before observed. The first pair goes out between the first and fecond vertebræ, and after communicating with the tenth and fecond vertebral, is fpent on the musculus flexus colli, splenius, complexus, and teguments of the occipitis. The fecond pair communicates with the ninth, and with the first and third of the neck, and then is diffributed to the teguments of the neck, and fide of the head, and to the glandula parotis and external ear, where it joins with the portio dura. The third pair of cervical nerves passes out between the third and fourth vertebra, foon communicates with the fecond, and fending down a large branch, which being joined by another from the fourth pair, forms the phrenic nerve that runs along the pericardium to be lost in the diaphragm, making a small turn round that part of the pericardium which covers the apex of the heart; hence it is that fuch as have ftrong palpitations of the heart, feel a pungent acute pain

and

pain immediately above the right orifice of the stomach: the other branches of the third cervical are distributed to the musculus trapezius, and deltoides, and to the teguments on the top of the shoulder; which with the description of the eighth pair, leads us evidently to the reason that an inflammation of the liver is generally attended with a hiccup, and a suppuration of that viscus with a violent pain on the top of the shoulder; however, it is not always a certain fign of the liver being suppurated, for any other cause stimulating or stretching the nerves, such as inflammation, wounds, schirrous or steatomatous tumours, &c. may produce the same effect. The fourth cervical pair, after fending off that branch which joins with the third to form the phrenic, runs strait to the axilla, where it meets with the fifth, fixth, and feventh cervicals, and first dorsal that escape in the interstices of the musculi scaleni; these give off nerves to the muscles of the neck, scapula, arm, and thorax, and to the teguments; and the confiderable branches into which they are divided are fix; but as they properly belong to the hand and arm, I shall describe them with those parts.

Q. What are the glands of the neck?

A. The principal gland of the neck is the thyroides; besides which there are a great number of lesser ones by the sides of the carotid arteries and jugular veins, and distributed here and there among the muscles and fat. Those in the anterior part of the neck are called jugulars; and those in the hinder part occipitales

edour

and cervicales; and these are all called lymphatic glands.

Q. What are the glandulæ thyroidæ?

A. The glandulæ thyroidæ are two lymphatic glands, in figure resembling the new moon; they adhere to the larynx and œsophagus, and are of a red colour; and they have arteries, veins, and nerves, as the larynx. These glands secrete a lubricating sluid, which moistens the cartilages and muscles of the larynx.

The cesophagus, especially towards its upper

part, has a great number of glands.

DIALOGUE IV.

Of the THORAX, and its PARTS.

A. The thorax is that large part of the body fituated between the abdomen and the neck, and answers to the extent of the sternum, ribs, and vertebræ of the back, both outwardly and inwardly. The anterior part is commonly called the breast; the posterior part, the back; and the lateral parts, the right and left sides. This cavity is lined by a membrane named plura, and divided into lateral cavities by a membranous septum named mediastinum, which is a production or duplicature of the plura. The parts contained in the thorax are the heart, pericardium, trunk of the aorta, trunks.

trunks of the carotid arteries, fubclavian arteries, trunks of the vertebral and axillary arteries, the fuperior portion of the descending aorta, the intercostal arteries, the vena cava superior, vena azygos, subclavian veins, trunks of the jugular, vertebral and axillary veins, a portion of the aspera arteria, the greater part of the cesophagus, the ductus lacteus, or thoracicus, the lungs, pulmonary artery, pulmonary veins, and nerves of the several parts. The cavity of the thorax is terminated downwards by the diaphragm, which separates it from the abdomen. The heart and lungs are properly the viscera of the thorax, I shall therefore speak of them separately.

Q. What are the bones of the thorax?

A. The bones of the thorax are the twelve superior vertebræ of the back, (from the last vertebra of the neck downward,) the ribs, and the sternum. The claviculæ and scapulæ belonging properly to the upper extremities, shall speak of them hereafter. The whole spine, (as I have before observed) consists of twenty-four vertebræ; those of the neck are already spoken of; the next twelve of the spine belong to the thorax, to these the ribs are articulated. See a general description of the vertebræ, page III.

The twelve vertebræ of the thorax, or back, differ from the rest in this, that they are larger than those of the neck, and smaller than those of the loins; their acute processes slope downwards upon one another; they have in each side of their bodies a small depression, wherein

K 3

they

they receive the round extremities of the ribs, and another in their transverse processes, which receive the little tubercle near the extremity of the ribs. The articulation of the twelfth vertebra of the thorax, with the first of the loins, is by arthrodia; for both its afcending and defcending oblique processes are received. The twelve vertebræ of the back have the least motion of any, because their cartilages are thin, their acute processes are long, and very near to one another; and they are fixed to the ribs, which neither move backwards nor forwards. They are bent backwards, behind the center of motion, to make room for the parts contained in the thorax; and that they might not be made too weak by this structure, they are formed for less motion than other vertebræ; and those in particular which are bent farthest from the center of gravity have the least motion. The fore-part of the two upper vertebræ of the thorax are flat forwards, as those of the neck, to make room for the aspera arteria and gula: the third and fourth vertebræ are acute, to give way to the veffels of the lungs and heart, and bent to the right side, for the better situation of the heart, which makes that fide of the breast more convex than the other, and therefore stronger; which seems advantageous to the right arm, its motions depending upon the fupport it receives from the breaft, hence the right arm is capable of more perfect actions than the other. The vertebræ of the back are hindered from diflocating forwards by the fame provilion

vision with those of the neck; and from luxating backwards by the ribs, which are fastened to the transverse processes of the inferior vertebræ, and against the back part of the body of the next superior; they also hinder them from dislocating to either side; but the last ribs are not fixed to the transverse processes of the vertebræ of the thorax; and therefore it is that luxations are most frequently seen in this part: but the vertebræ of the loins are received into deep cavities, and are tied with much stronger

ligaments for their fecurity.

Those that have the vertebræ of the back flick out are faid to be bunch-backed; and in fuch the cartilages which are between the vertebræ are very thin and hard-forwards, but confiderably thick backwards, where the oblique processes of the superior and inferior vertebræ are at a confiderable distance from one another, which distance is filled up with a vifcous subflance. This inequality of the thickness of the cartilages happens either by a relaxation or weakness of the ligaments and muscles, which are fastened to the back side of the vertebræ, in which case their antagonists finding no oppofition, remain in a continual contraction, and confequently there can be no motion in thefe vertebræ. If this deformity has been from the womb, the bones being at that time foft and tender, the bodies of the vertebræ partake of the fame inequality as the cartilages. If the bunch be towards one shoulder, for example towards the right, then the cartilages on that fide K 4

fide are very thick, but thin and dry on the other fide; on the left fide the oblique apophyses come close together; but on the right there is a confiderable diftance betwixt them, and the ligaments and muscles are greatly extended on the right side, but those on the left are much contracted. If the vertebræ are diftorted inwards, all things have a different face; the cartilages, and fometimes the vertebræ, are very thick forwards, but very thin and hard backwards; the acute and oblique processes are very close to one another; and the ligaments upon the bodies of the vertebræ are greatly relaxed, but the muscles and ligaments which tie the processes together are very much contracted. These distortions seldom happen in the vertebræ of the loins; but fuch as are fo miserable, have little or no motion of their back.

Q. How is the sternum or breast-bone com-

posed?

A. The sternum or breast-bone is generally composed of three spongy bones, sometimes more; in the upper part of this bone, on each fide, the clavicles are articulated; the middle part is narrow, the lower part broad; to the end of this adheres a cartilage, called from its figure cartilago ensiformis vel xiphoides, which is usually fingle; sometimes it is double, and bifurcated, and not unfrequently bony throughout. There is also frequently found a variety in the form of the cartilages which join the ribs and sternum, sometimes one cartilage serving

two ribs, and fometimes a cartilage not joined to any rib; frequently in old people we find parts of them offified. On each fide of the sternum, there are seven cavities, for the articulation of the seven true ribs, to which their cartilages are fixed. The sternum serves to form the anterior part of the breast, to support the ribs and clavicles, and to defend the parts contained in the cavity of the breast: it also serves for the insertion of the mediastinum, and for the sustaining the heart itself and several muscles.

Q. How many ribs are there, and how are

they composed?

A. The costa or ribs are twenty-four in number, twelve on each fide; the middle ribs are the largest: all the ribs are harder, rounder, and more incurvated towards their articulations with the vertebræ, than at the other extremity towards the sternum, which is thinner, broader, and more fpongy. The ribs are diftinguished into true and spurious: the true ribs are the feven upper pair, so called, because their cartilages reach the sternum; the spurious, or bastard-ribs, are the five lowest pair, so called, as being smaller, shorter, and more cartilaginous than the rest, and not reaching so far as the sternum, which makes their articulations very lax, as they terminate in long, foft cartilages, which bending upwards, are joined to the upper ribs. On the infide of the middle ribs runs a pretty deep finus, or channel, for the passage of the intercostal vessels, reaching from the end next the spine, almost to its juncture with the carti-

lage. In the anterior extremities, the cartilages of the feven true ribs are all joined to the sternum; the eighth, ninth, and sometimes the tenth, cohere either with the sternum, or mutually adhere to one another, by means of their transverse cartilages. The anterior extremities of all the others, are loofe and free, between the muscles of the abdomen and the diaphragm. In the posterior extremities there are in most of them two capitulæ, or heads, which are firmly joined to the vertebræ of the back, yet so as to form movable articulations; for when they are drawn upwards, the cavity of the breast is enlarged for inspiration, and so the contrary. The ribs of infants may be broken inwards like a green stick, without the broken ends being separated; which is often done, by the idle custom of careless nurses, taking hold of their breafts, and hoifting them up on one hand frequently: the shape of children's breafts are many times quite spoiled by fuch tricks, which occasions weakness of body, crookedness, and other diseases.

Q. What are the cartilages of the thorax?

A. The cartilages of the bones of the thorax are those of the vertebræ, sternum, and ribs: but the cartilages of the vertebræ are already spoken of.

Q. What are the cartilages of the sternum

and ribs?

A. The sternum of an adult has commonly fixteen cartilages, (eight on each fide) fourteen are articular, the other two fymphyses. Of the articular

articular cartilages, two belong to the articulations of the clavicula, and twelve to those of the true ribs, from the fecond to the feventh inclusively. The two symphyses are those between the sternum, and the first rib on each fide. There is likewise another symphysis by which the upper portion of the sternum is connected to the lower, the cartilage of which is often obliterated in an advanced age. The apophysis ensiformis is often bony towards the sternum; and more or less cartilaginous at the other end. In very aged persons it has been found entirely offified, and fometimes wholly cartilaginous, even in adults. All the ribs have cartilaginous portions (as before mentioned) which differ from each other in length, breadth, incurvation, adhesions, and in their extremities are whiter, more polished, broader and thicker in the natural state than when they are dried. The cartilages of the bastard, or false ribs, are naturally more slender and pliable than those of the true ribs, the middle or inner fubstance of which acquires the confistence of bones in old age, and their extremities fometimes offify, and are immovably fixed to the sternum.

Q. What are the ligaments of the sternum

and ribs?

A. The sternum has several ligaments by which it is connected with the clavicles and ribs. It is joined to the clavicles by strong short ligaments fixed by one extremity round the edges of its superior notches; by the other, in the extremity of each clavicle; and by the middle

middle to the inter-articular cartilages already explained. All the ribs are connected to the bodies of the vertebræ, by strong short ligamentary fasciculi, fixed by one end round the fosfulæ in the vertebræ, and by the other round the head of each rib. The ribs are likewise tied to each other by thin ligaments which go obliquely from the cartilage of each rib to that of the next. The ten upper most ribs on each fide are connected to the transverse apophyses of the vertebræ of the back, by strong short articular ligaments fixed much in the fame manner as those between the heads of the ribs and bodies of the vertebræ. Both these articulations are provided with capfular ligaments. The eleventh rib on each fide having no articulation with the transverse apophyses is connected to them by auxiliary strong short ligaments fixed in its neck. The last rib is only joined by its head to the body of the twelfth vertebra of the back; but it is connected in a particular manner to the transverse apophysis of the first vertebra of the loins by a broad ligament The first true rib has no ligamentary connexion with the sternum, the cartilaginous symphysis being sufficient. The rest are closely joined to that bone, in the fame manner as the clavicles abovementioned. The cartilage of the first false rib is joined to that of the last true rib, by several short ligamentary filaments; and the other bastard ribs are connected together much in the same man-

Q. What are the muscles of the thorax.

A. The

ferted

A. The muscles of the thorax are, extenfor dorsi et lumborum vel multifidus spinæ, scalenus, serratus, superior et inferior posticus, intercostales externi et interni, triangularis sterni, and diaphragma; besides those whose actions tend chiesly to the upper extremities, which I shall describe hereafter.

Q. Which is the extensor dorsi et l'umborum, muscle?

A. The longissimus dors, multifidus spina, femispinalis, facrolumbalis, &c. are all that portion of flesh betwixt the os sacrum and the neck, which feeing there is no membrane to diffinguish it into several muscles, is all employed in the fame actions, I give it the name of extensor dorsiet lumborum vel multifidus spinæ, which arises from the upper part of the os facrum, the spine of the os ilium, the back parts of the lowermost vertebræ of the loins, and remarkably from those strong tendons which appear on their outfides. That part of this muscle which is known by the name of sacrolumbalis is inferted into all the ribs near their articulations with the transverse processes of the vertebræ, and into the transverse process of the last vertebra of the neck; besides, as this passes over the ribs, it receives an origin from every rib. The portions of this muscle, which arise from the ribs, and are inferted into other ribs above, will necessarily draw the back part of the ribs nearer together, which must always be done as the back extends. The next portion of this muscle, called longissimus dorsi, is in.

ferted into all the transverse processes of the vertebræ of the back (belonging to the thorax) and partly into the ribs, and the uppermost transverse processes of the vertebræ of the loins; the upper end of it is neither very distinct from the complexus of the head, nor spinalis of the neck. The rest of this muscle, known by the names of femi-spinalis, facer, &c. arises also from all the transverse and oblique processes of the loins and back; every portion, except the lowermost, passing over five joints, is inserted into the spinal process of the fixth vertebra above its origin, all the way up the back, and at the neck commences transversalis colli. Befides the uses of the extensor dorsi et lumborum, which its name implies, it and its fellow alternately raise the hips in walking, which any one may feel by laying his hand upon his back.

Q. Which is the scalenus muscle?

A. The scalenus arises from the transverse processes of the second, third, fourth, fifth, and sixth cervical vertebræ. It is inserted in three parts, into the first, second and sometimes the third rib, being thus divided for the transmission of the subclavian vessels; hence some anatomical writers have made three muscles of it, under the names of first, second, and third scalenus, or prior, medius, and posticus. This muscle may bend the neck; but its chief use is to support the upper ribs, and partly elevate the thorax.

Q. Which is the ferratus superior posticus?

A. The ferratus superior posticus lies immediately under the rhomboides, and arises from the spinal processes of the two inferior vertebræ of the neck, and the three superior of the thorax; but it is inferted at the bending of the fecond, third, and fourth ribs: this, with the scalenus, sustains the upper ribs, that they might not be pulled downward by the depressors of the ribs in expiration, as the lower ribs are upward in inspiration.

Q. Which is the ferratus inferior posticus?

A. The ferratus inferior posticus arises from the spinal processes of the three superior vertebræ of the loins, and two inferior of the thorax; and is inferted at the bending of the ninth, tenth, and eleventh ribs. It pulls down the ribs in expiration.

Q. Which are the intercostales externi et in-

terni !

A. The intercostales externi et interni, are eleven pair on each fide, in the interstices of the ribs; from their fituations diftinguished into external and internal; they all arise from the under edge of each rib, and are inferted into the upper edge of the rib below. The external are largest backward, having their first beginnings from the transverse processes of the vertebræ like diftinct muscles, which some call levatores costarum. The internal run all from above obliquely backward, being thickest forward and thinnest toward the spine; these are also continued betwixt the cartilages of the sternum, which fibres are perpendicular to the cartilages ;

cartilages; and between the cartilages of the lowest ribs, they are inseparable from the obliquus ascendens abdominis. These muscles, by drawing the ribs nearer to each other, pull them all upward, and dilate the thorax, they being sustained at the top by the scalenus and serratus superior posticus.

Q. Which is the triangularis sterni muscle?

A. The triangularis sterni arises from the lower and interior part of the sternum, and internally from the cartilage ensiformis; it is inserted on each side into the cartilages of the fourth, sixth, and seventh true ribs; and it is one of the constrictor or depressor muscles of the breast, which pulls the ribs to the bone of the sternum, and thereby bends its cartilages in expiration.

Q. Which is the diaphragma?

A. The diaphragma, is the transverse muscle which separates the thorax from the abdomen. This is a very broad and thin muscle, situated at the basis of the thorax, and serving as a transverse partition to separate that cavity from the abdomen: for this reason the Greeks termed it diaphragma, and the Latins, septum transverfum. Its upper fuperficies is convex, and its It is connected with the sterlower concave. num, the spurious ribs, the pericardium, the mediastinum, and the vertebræ of the loins. There are in the diaphragm two large foramina; the first is in the left side of it, and gives pasfage to the gula, and the par-vagum; the fecond is in the right fide, and the lower trunk of

the

the vena cava passes through it; there is also an interflice between the two heads of the lower part; through which pass the aorta, the vena azygos, and the ductus thoracicus. The diaphragm is covered with a membrane on the upper part from the pleura; on the lower from the peritonæum. This muscle arises on the right fide by a process from three lumbal vertebræ, and one of the thorax; and on the left, from one superior of the loins, and inferior of the thorax; and is inferted into the lower part of the sternum, and the five inferior ribs. The middle of this muscle is a flat tendon, from whence the fleshy fibres begin, and are distributed, like radii from a centre to a circumference. The uses of the diaphragm are, first, to affift in respiration; for in taking in the breath, it is preffed downwards, and in expiration, it rifes upward, into the cavity of the thorax: fecondly, to affift the necessary motions of the contents of the abdomen, viz. of the stomach, intestines, liver, and spleen; and in the promoting the fecretions of the chyle, bile, &c. and, lastly, for assisting the expulsion of the fæces, the urine, the fœtus in parturition, and of the secundines. When the diaphragm acts alone, it constricts the thorax, pulling the ribs downward, which action is generally performed to promote the ejection of the fæces.

Q. What are the vifcera of the thorax?

A. The viscera of the thorax are properly only the heart and lungs; but that cavity also contains the plura, mediastinum, pericardium.

a portion of the trachea arteria, and the greater part of the œsophagus, besides the thoracic duct, blood vesses, nerves, and glands.

Q. What is the pleura?

A. The pleura is a fine smooth, robust, and tense membrane, adhering very closely to the inner furface of the ribs, sternum, and intercoftal muscles, and lining the whole cavity of the thorax, except the diaphragm, which is covered with no other than its own proper membrane. Its structure resembles two sacks or bags (one on each fide the thorax,) which contains the two lobes of the lungs: from the conjunction of these two sacculi of the pleura in the middle of the thorax, is formed the mediaftinum, which I shall speak of by and by. The pleura is composed of a double membrane of a very firm texture, and plentifully stored with blood-veffels and nerves, in all which it refembles the peritonæum, (which I shall describe hereafter) and likewife, in that it is made up of an inner true membranous lamina, and a cellular fubstance on the outside, which is a production or continuation of the lamina. Its veffels are arteries, veins, nerves, and lymphatics. The arteries arise from the intercostals, the diaphragmatic, and the mammary ones, and are very numerous; the veins from the veins of the fame name with those arteries; but all of them discharge themselves into the trunk of the vena azygos, and the upper trunk of the The nerves are from the vertebræ of the thorax, and the diaphragmatic ones. The lymphatics

lymphatics all run to the ductus thoracicus. The use of the pieura is to make the inside of the thorax smooth and equal, and to lubricate and strengthen the whole cavity.

Q. What is the mediastinum?

A. The mediaftinum, is a double membrane continuous to the sternum, situated under it, and adhering firmly to it; it is formed by the continuation of the pleura, which comes from the sternum, and goes through the middle of the thorax to the vertebræ, dividing the cavity of the thorax longitudinally into two parts; but as it is not exactly under the middle of the sternum, but somewhat to the left side, the right part of the thorax is larger than the left: hence may be judged the uncertainty of trepaning the sternum, recommended by the ancients in some This double septum, or partition, contains in its duplicature, the heart in its pericardium, the vena cava, the cefophagus, and stomachic nerves. The mediastinum receives veins and arteries from the mammary and diaphragmatic vessels, and sometimes has proper and particular ones of its own from the aorta and cava: these are then called the mediastinal vessels. Its nerves, which are fmall, are from the diaphragmatics, and the par vagum. It has a member of lymphatics, which run to the ductus thoracicus. The ufes of the mediastinum are two: the first is to divide the breast longitudinally into two parts, by which feveral great purposes are answered; as, 1. That on one of the lobes being ulcerated, the other might

might not be immediately affected. 2. That's water, matter, or any thing elfe contained in one part of the thorax, might not, at the fame time, affect both parts of the lungs. 3. That in case of a wound in one side of the thorax, respiration might be continued in the other, and the person not be immediately suffocated. The fecond general use of the mediastinum is to fupport the heart in its pendulous state, for the benefit of its free motion, especially when we lie on our backs.

Q. What are the lungs?

A. The lungs are the largest viscus of the thorax: they are situated in the two sides of it, with the heart as it were between them; and are connected by means of the mediastinum with the fternum and vertebræ; with the heart by means of the pulmonary vessels, and immediately with the aspera arteria. The colour of the lungs in infants is a fine florid red; in adults it is darker; and in old people livid, or variegated with black and white. When inflated they have fome resemblance to the hoof of an ox; and are convex on the upper fide next the ribs, and concave underneath next the diaphragm. They are divided into two large lobes, one feated on each fide of the mediastinum; the left lobe, which is the smallest, is subdivided into two lobules, and the right, which is larger, is subdivided into three lobules. The double membrane, with which the lungs are furrounded, is continuous externally with the pleura, and internally not only immediately covering the fubstance of the lungs, but also all its inner lamina, and

and fills up the interstices which are between the bunches of the small lobes or lobules with little veficular cells. The fine capillary blood vessels are so thick upon this membrane, that it feems to be nothing but a net-work of veins and arteries. The substance of the lungs is spongeous, or vesiculous, and they seem, indeed, entirely composed of an infinite number of little lobes, or lobules, of various figures and magnitudes; but their furfaces are fo adapted to one another, as to leave but very few and small interstices. These lobes are disposed like fo many bunches of grapes, upon the fides of the bronchia, or ramifications of the trachea (described hereafter); each little lobe, or lobule, contains within its own proper membrane an infinite number of small orbicular vessels, which leave finall interstices between them, full of little membranes, like those which tie the lobes together. The bronchia, in their origin, are formed of imperfect annuli, and in their progress of cartilaginous and membranous segments, very curiously connected and joined together; these have their origin from the trachea; and after being subdivided into innumerable ramifications, finally terminate in those small veficles which form the greater part of the fubstance of the lungs, as before mentioned. These extremities (of the branches of the bronchia) open into the cavity of the veffels, which are properly formed by its membranes; but the capillary blood veffels are only spread upon the veficles like a net, with frequent and large in-L 3 olculation. osculation. The vessels which enter the lungs are the trachea, or afpera arteria, by which we draw in the air, (hereafter to be described.) The arteria pulmonalis, which comes from the right ventricle; see page 22, and the vena pulmonalis, whose trunk opens into the left ventricle of the heart, see page 34; each of these divides into two branches, for the two great lobes of the lungs, where they are fubdivided into as many branches as there are little lobes or veficles in the lungs. Wherever there is a branch of the trachea, there is also a branch of the vein and artery; and the trachea is always in the middle. Upon the branches of the trachea arteria (which I observed before are termed bronchia) runs a small artery called arteria bronchialis, and a small vein called vena pneumonica; the artery comes from the aorta, and the veins open into the subclavian. Upon the bronchi, even to the minutest ramifications, run likewise the fine threads of the eighth pair of nerves. Besides these, the lungs have lymphatics, which discharge themselves into the thoracic duct; but they are smaller, and make more frequent inosculations than are observed any where elfe. The lungs of animals before they have been dilated with air, are specifically heavier than water; but upon inflation, they become specifically lighter, and swim in water; which experiment may be made to discover whether a dead child was still-born, or not: but if the child has breathed but a little, and the experiment is made long after, the lungs may

may be collapsed, and grow heavier than water, as I have found by experiment, which may sometimes lead a man to give a wrong judgment in a court of judicature, but then it will be on the charitable side of the question. Adhesions of the lungs to the pleura are in men so common, I know not how to call it a disease; they being found so, more or less, in most adult persons, and without any inconvenience, if the lungs are not rotten.

Q. What is the pericardium?

A. The pericardium, or heart-purse, is an exceeding strong membranaceous bag, loosely including the heart; the figure of the pericardium is conic, like that of the heart itself; its fize is fuch as can conveniently contain the heart, without pressing upon it; it is connected with the mediastinum, with a great part of the diaphragm, and with the large veffels of the heart, which, together with this covering, fuftains also the heart itself in its place. The pericardium is composed of a double membrane; the exterior one is common with the pleura and mediastinum; the interior is proper, lubricous, and is continuous with the coats of the larger veffels. This membrane, when expanded upon the finger, frequently discovers a great number of foraminula, or little apertures. The arteries and veins of the pericardium are from those of the mediastinum and diaphragm, and its nerves are also from the diaphragmatics. Its lymphatics all run to the thoracic duct. The uses of the pericardium are, to support the heart in in a pendulous state, especially when we lie down; to defend the heart from the cold air taken in at the lungs, to preserve it from being injured by water, by matter, or any other extraneous stuid in the cavity of the thorax; and to contain a liquor to lubricate the surface of the heart, facilitate its motions, and abate its friction against the pericardium. The pericardium is found sometimes in persons who have been long hectic, to cohere with the substance of the heart; and there have been instances of its having been wholly wanting.

Q. What is the heart?

A. The heart is a muscular body, included in the pericardium (just now described) and fituated in the cavity of the thorax, nearly in the middle of the breaft, on the anterior part of the diaphragm, between the lobes of the lungs and the two laminæ of the mediastinum. The heart is the primary organ of the circulation of the blood, and consequently of life. It is in some measure of a conical figure, flatted on the sides, round at the top or apex, and oval at the basis; its lower part is plane, and the upper part convex. Its fituation is nearly transverie, or horizontal; fo that its base is in the right side, upon the fourth and fifth vertebræ of the thorax; and its apex or point is inclined downwards, lying with the greatest part of its bulk in the left fide of the thorax, and confequently it is there that the pulsation is felt. Its base is fixed by the veffels going to and from it; but its apex is free, and is received in a kind of cavity of the

the left lobe of the lungs, as may be observed, the lungs being extended with air. This incumbrance on the left lobe of the lungs feems to be the cause of that sides being most subject to the pleuritic pains, which have generally been found upon diffection to proceed from inflammations in the lungs. The length of the human heart is about fix fingers breadth, at the base about five fingers, and its circumference about thirteen.

At the basis of the heart, on each side, are fituated two appendices called auricles, to receive the blood; the right from the two venæ cavæ, and the left from the pulmonary veins. (See vena cava and pulmonaris.) In the right auricle, at the meeting of the cavæ, is an eminence called tuberculum Loweri, which directs the blood into the auricle; immediately below this tubercle, in the ending of the cava ascendens is the veftige of the foramen ovale, and near this in the auricle, is the mouth of the coronary veins. The left auricle is much lefs than the right; but the difference is supplied by a large mufcular cavity, which the veins from the lungs afford in that place. The fides of this mufcular cavity are thicker than the fides of the right auricle, in about that proportion in which the left ventricle of the heart is stronger than the right; their uses being to receive blood from the veins that lead to the heart, and press it into the ventricles, as a ftrength in each auricle proportionable to the strength of the ventricle that it is to fill with blood,

blood, feems necessary: and this different thickness of the coats of the auricles makes the blood in the left, which is thickest, appear through it of a paler red; but when it is let out of the auricles, it appears alike from both; which they would do well to examine, who assirm the blood returns from the lungs of a more florid colour than it went in, and offer it as an argument of the blood's being mixed with air in

lungs.

There are also two cavities in the heart called its ventricles, which receive the blood, and are hollow muscles, or two cavities in one muscle. Both these cavities receiving the same quantities of blood in the fame times, and always acting together, must be equal in fize, as it is generally supposed they equally discharge what they contain at every fystole; nevertheless, the left appears less than the right, it being found empty in dead bodies, and the right usually full of blood. Each ventricle opens at the base by two orifices, one of which answers to the auricles, the other to the mouth of a large artery; the former receives the blood through the veins, the latter delivers the blood through the arteries. The right ventricle is fituate anteriorly, and is thinner and weaker in its circumference, but usually much more capacious than the left: it receives the blood from the vena cava and the right auricle, and delivers it into the pulmonary artery, to be carried to the lungs. The left ventricle is much stronger and thicker in its sides, but is narrower and smaller than the right;

right; it receives the blood from the pulmonary vein and the left auricle, and drives it very forcibly into the aorta, and so through the

whole body.

Over the orifices of the veins at the entrance of the auricles in each ventricle, are placed valves, to hinder a return of blood while the heart contracts; those in the right ventricle are named tricuspides, those in the left mitrales: in the beginning of each artery from the heart, are placed three valves, which hinder a return of blood into the ventricles The first fort of valves open inward toward the ventricles, allowing the blood to enter the heart, but hindering it from returning the same way; the other kind open outward from the heart toward the great veffels, fuffering the blood to go out of the heart, but hindering it from returning; these last are termed semi lunar valves; but those in the pulmonary artery are more properly named figmoidales, and those in the aorta semilunares.

The inner surface of the ventricles is very uneven, many eminences and cavities being observable therein; the ventricles are divided by a septum running between the edges; and their most considerable eminencies are thick sleshy productions called columnæ. To the extremities of these pillars are fastened several tendinous cords, the other ends of which are joined to the valvulæ tricuspides. The cavities of the inner surface of the ventricles are small deep fossulæ or lacunæ placed very near each other,

with

with small prominent interstices between them. The greatest part of these lacunæ are orifices of the venous ducts.

The blood veffels of the heart are of two kinds, common and proper; its proper or peculiar veffels being the coronary arteries and veins. The common veffels of the heart are two veins, called the vena cava and the vena pulmonalis; and two arteries, the pulmonary one, and aorta. The nerves of the heart are small, and arise from the par vagum and intercostals. See Dialogue I. arteries, veins and nerves.

Q. Which are the coronary arteries?

A. The coronary arteries are two in number, and go out from the beginning of the aorta, and afterwards spread themselves round the basis of the heart, to the substance of which they send numerous ramifications One turns to the right hand, the other to the left : the right coronary artery runs in between the basis and right auricle all the way to the flat fide of the heart, and fo goes half-way round: the left artery has a like course between the basis and left auricle, and before it turns on the basis, it sends off a capital branch, which runs between the two ventricles. Another principal branch goes off from the union of the two arteries on the flat fide of the heart, which running to the apex, there joins the other branch.

Q. Which are the coronary veins?

A. The coronary veins are distributed exteriorly much in the same manner as the arteries,

but they end partly in the right auricle, and partly in the right ventricle; their trunk principally in the former, by a particular orifice, furnished with a semi-lunar valve. They likewise terminate in the left ventricle, but in smaller numbers. All the coronary veins and their ramifications communicate with each other.

Q. What is the trachea or aspera arteria?

A. The trachea or aspera arteria, vulgarly called the wind-pipe, is a large canal, partly cartilaginous, and partly membranous, extended from the mouth to the lungs. It is situated in the middle and anterior part of the neck, and is connected with the fauces, lungs and cefopha. gus. Its mouth or entrance is, by anatomists, called the larynx, the rest aspera arteria. The aspera arteria extends from the larynx to the bronchia, or lungs, being in some measure of a conic figure. Its beginning is cylindrical, and capable of admitting a finger; and its other end is somewhat narrower. It runs down into the thorax, under the sternum, between the two pleuræ, through the upper space left between the duplicature of the mediastinum, behind the thymus gland. Having reached aslow as the fourth vertebra of the back and curvature of the aorta, it divides into two lateral parts or branches, one towards the right hand, the other towards the left, which enter the lungs, and are distributed through them in the manner there described: these two branches are called bronchia, and that on the right fide is shorter than that on the left; whereas the right pulmonary

pulmonary artery is the strongest. The trachea is made up of segments of circles, or cartilaginous rings, disposed in such a manner as to form a canal open on the back part; the cartilages not going quite round; but this opening is filled by a soft glandular membrane, which completes the circumference of the canal. These circles or hoops, are in number from sixteen to twenty, and are all connected by a very strong, elastic, membranous ligament, fixed to their

edges.

The canal of the aspera arteria is lined on the infide by a particular membrane, which appears to be partly fleshy, or muscular, and partly ligamentary, perforated by an infinite number of small holes, more or less imperceptible, through which a mucilaginous fluid continually passes to defend the inner surface of the trachea against the acrimony of the air which we breathe. At the angle of the first ramification of the trachea arteria, we find on both the fore and backfides certain foft, roundish, glandular bodies, of a livid colour, and of a texture partly like that of the thymus, and partly like that of the glandulæ thyroides. There are other glands of the fame kind at the origin of each ramification of the bronchia, but they decrease proportionably, in number and fize: they are fixed immediately to the bronchia, and covered by the interlobular fubiliance; and they feem to communicate by small openings, with the cavity of the bronchia, as before observed.

The uses of the trachea are to assist in deglutition, and to be assistant to the lungs. On the exterior part of the trachea are the thyroide and bronchial glands, (which I shall hereafter mention) which secrete a humour to moisten it: its arteries are from the external carotids, the veins from the jugulars, and the nerves from the recurrent ones of the plexus cervicalis.

Q. What is the cesophagus?

A. The cefophagus, or gullet, is that membranous canal which conveys the aliment from the mouth to the stomach. It is partly muscular, and partly membranous, fituated behind the trachea arteria, and before the vertebræ of the back, from near the middle of the neck. down to the lower part of the thorax, from whence it passes into the abdomen, through a particular hole of the small or inferior muscle of the diaphragm, and ends at the upper orifice of the stomach. Its upper part is wide and open, spread behind the tongue to receive the masticated aliment, and is termed pharynx, the muscles of which I have already spoken of. The fubstance of the cofophagus is composed of feveral coats, almost in the same manner as the stomach, of which it is the continuation. The first coat, while in the thorax, is formed only by a duplicature of the posterior part of the mediaftinum, and is wanting above the thorax, and in the neck, where the outer coat of the cesophagus is only a continuation of the cellular fubstance belonging to the neighbouring parts. The fecond coat is mufcular, made

The œsophagus, from its very beginning, turns a little to the left hand, and naturally runs along the left extremities of the cartilages of the aspera arteria. The arteries of the œsophagus are from the carotids, the aorta, the intercostal, and the cæliac. The veins are from the jugulars, the azygos, and the coronary veins of the stomach; and the nerves from the par vagum. There are also certain excretory ducts, called ductus excretorii novæ vercelloni, which

arise

arise from the glands, and convey a saltish liquor into the cesophagus and stomach: see the glands of the thorax. The glands, from which these ducts have their origin, are of three kinds, viz. the gastric glands, which are conglomerate, and are situated near the left orifice of the stomach, the dorsal ones, which are situated near the fifth vertebra of the thorax; and the bronchial, thyroide, and tracheal, described below. The use of the cesophagus is to carry the meat from the mouth into the stomach, by means of the muscles of the pharynx and sleshy sibres of the gula, which perform its peristaltic motion, and also for a commixtion of the liquid serving for digestion.

Q. What are the arteries of the thorax?

A. The arteries of the thorax (besides those of the viscera I have already mentioned) are various, and proceed chiefly from the aorta already spoken of. The branches of the ascending aorta in the thorax take their names from the parts they are bestowed on; the principal branches are arteriæ coronariæ, carotides, subclaviæ, from the latter arise the thymica, diaphragmaticæ, pericardia, mediastina, trachealis, and mammaria interna: these from the aorta ascendens. From the superior portion of the aorta descendens arise arteriæ bronchiales, æsophageæ, intercostales, to their parts of the thorax. The coronary and carotid arteries are already described.

Q. Which are the fubclavian arteries?

A. The subclavian arteries are so named from their situation near the clavicles, in the transverse direction of which they run; they are two in number; one right, the other left. These arteries arise from the arch of the aorta, on each side of the left carotid, which commonly lies in the middle between them; but when both carotids go out separately, they both lie between the fubclaviæ. These arteries terminate, or rather change their name, above the middle of the two first ribs, between the anterior insertions of the musculi scaleni. They give off small arteries to the mediastinum, pericardium, and aspera arteria; afterwards they fend off the mammaria interna, cervicalis, and vertebralis, and fometimes an intercostal artery to the first ribs. The right fubclavian, at about a finger's breadth from its origin, also often produces the common carotid of the fame fide.

Q. Which is the arteria thymica?

A. The arteria thymica arises from the subclavian, communicating with and receives some branches from the mammaria interna and has some from the intercostalis superior. The mediastina and pericardia the same.

Q. Which is the arteria pericardia?

A. The arteria pericardia also arises from the anterior middle part of the subclavian, and runs down upon the pericardium all the way to the diaphragm to which it sends some small ramifications.

Q. Which is the arteria mediastina?

A. The mediastina arises sometimes immediately after the thymica, and is distributed principally to the mediastinum.

Q. Which

Q. Which is the arteria trachealis?

A. The trachealis runs up from the subclavia, in a winding course, along the aspera arteria, to the glandulæ thyroideæ and larynx, detaching small arteries to both sides, one of which runs to the upper part of the scapula.

Q. Which is the arteria mammaria interna? A. The internal mammary artery comes from the anterior and lower fide of the subclavian near the middle of the clavicle, and runs down for about one finger's breadth behind the cartilages of the true ribs, an inch distant from the sternum. In its passage, it sends branches to the thymus, mediastinum, pericardium, pleura, and intercostal muscles; and through these muscles, and between the cartilages of the ribs, to the pectoralis major, mammæ, membrana adipofa, and skin. Several of these branches communicate by anaftomofes with the mammaria interna, and other arteries of the thorax. Afterwards it goes out of the thorax, on one fide of the appendix enfiformis, and is loft in the musculus abdominis rectus, a little below its upper part; communicating at this place with the epigaftric artery, and in its course it gives off branches to the peritonæum, and the anterior parts of the abdominal muscles.

Q. Which is the arteria intercostalis superior?

A. The superior intercostal artery, when it it does not go out from the trunk of the aorta descendens, commonly arises from the lower side of the subclavian, and runs down on the inside of the two, three, or four uppermost M. 2.

true ribs, near their heads, sending off under each rib a branch, which runs along the lower edge, and supplies the intercostal muscles, and neighbouring parts of the pleura. These branches all communicate with one another, up and down the intercostal muscles. They also give branches to the musculi sterno-hyoidei, subclavius, vertebrales, and bodies of the vertebræ, pectoralis, and through the notches of the first four vertebræ to the medulla spinalis.

Q. Which are the arteriæ bronchiales?

A. The bronchial arteries go either from the foreside of the superior part of the descending aorta, first intercostal, or arteriæ æsophagea. They rise sometimes single sometimes double, sometimes triple, and adhere every where sirmly to the bronchia; their branches, or ramifications, communicate sometimes with those of the pulmonary vein, arteria æsophagea, coronaria cordis, and vena azygos.

Q. Which are the arteriæ œsophageæ?

A. The œsophageæ are generally two or three in number, though sometimes but one. They arise anteriorly from the aorta descendens, and are distributed to the œsophagus, &c. Sometimes the uppermost œsophagus produces a bronchial artery.

Q. Which are the arteriæ intercostales in-

feriores?

A. The inferior intercostals are commonly seven or eight on each side, and sometimes ten, when the superior intercostals arise likewise from the aorta descendens; in which case, these

run obliquely up upon the ribs. They arise along the backfide of the descending aorta generally in pairs, all the way to the diaphragm, and run transversely toward each side, on the bodies of the vertebræ. Those on the right fide pass behind the vena azygos; and afterwards, they run to the intercostal muscles, along the lower edge of the ribs, all the way to the sternum or near it. They send branches to the pleura, vertebral muscles, muscles of the thorax, and to the upper portions of the muscles of the abdomen, and they communicate with the arteriæ epigastricæ and lumbares. Before they take this course along the ribs, &c. they detach branches to the great canal of the spina dorsi, which communicating with the like arteries from the other side of the spine, form a kind of arterial rings, as do the arteriæ lumbares. After the above course, each intercostal about the middle of the rib fends a principal branch internally, and another externally; those that run upon the false ribs are afterwards distributed to the abdominal and other neighbouring muscles, particularly to those of the diaphragm. They also communicate with the lumbares, and fometimes with branches of the hypogastricæ

Q. Which is the arteria mammaria externa

vel thoracica superior?

A. The external mammary or superior thoracic artery, runs down in a winding course on the lateral parts of the thorax, and crosses the ribs. It gives branches to the two pectoral M 3 muscles,

muscles, to the mamma, musculus subsclavius, ferratus major, latissimus dorsi, and to the upper portions of the coraco-brachialis and biceps.

Q. Which is the arteria thoracica inferior? A. The inferior thoracic artery runs along the inferior costs of the scapula to the musculus subscapularis, teres major, and minor, infra spinatus, serratus major, latissimus dorsi, and the neighbouring intercostal muscles, communicating with the arteriæ scapulares.

Q. What are the veins of the thorax?

A. The veins of the thorax are also various, and proceed chiefly from the vena cava superior already spoken of. The principal branches are, venæ coronariæ cordis, vena azygos vel sine pari, venæ intercostales, pectoralis internæ, (from the latter arise the diaphragmaticæ superiores, mediastinæ, mammariæ internæ, thymicæ, pericardiæ, and gutturales, or tracheales) subclaviæ, jugulares, vertebrales. The coronary veins of the heart, the jugulars and vertebrals, I have already spoken of.

In describing the veins, I shall begin with the great trunks, and end with the ramifications and capillary extremities, according to their several divisions and subdivisions, as I have

before observed.

Q. Which is the vena azygos, or fine pari? A. The vena azygos, or fine pari, is a vein arifing within the thorax on the right fide, having no fellow on the left; whence it is called azygos, or vena fine pari. This vein is very confiderable, and arifes posteriorly from the

vena cava superior, a little above the pericardium, and is immediately afterwards bent backward over the origin of the right lobes of the lungs; forming an arch which furrounds the great pulmonary veffels on that fide, as the arch of the aorta does those of the left fide, with this difference only, that the curvature of the azygos is almost directly backward, whereas that of the aorta is oblique. From thence it runs down on the right fide the vertebræ dorfi, on one fide the aorta, and before the intercostal arteries I have feen this vein extremely large like the trunk of the inferior cava, from the diaphragm to the original of the renales; the true cava being through all this space very narrow, or of the fize of the ordinary azygos. The vena azygos fends out first of all two or three small veins from the top of the arch, one of which goes to the aspera arteria, and the others partly to that and partly to the bronchia, by the name of venæ bronchiales, accompanying the ramifications of the bronchial artery. And from the extremity of the arch proceeds a fmall trunk common to two or three small veins, called intercostales superiores dextræ, which bring back the blood from the first three series of intercostal muscles, and from the neighbouring parts of the pleura. These intercostal veins send branches through the intercostal muscles to the ferratus major posticus, &c. and afterwards they run along the interstices between the ribs, communicating with the venæ mammariæ. They likewile

wife fend small branches backward to the vertebral muscles, and canal of the spine, where they communicate with the venal circles or finuses, which bring back the blood from the medulla spinalis. As the azygos runs down it fends off the inferior intercostal veins on the right side, one going to each series of intercostal muscles: these veins runs along the lower edges of the ribs, and perforate the muscles which go to the posterior and external parts of the thorax. The azygos fends off likewife the left intercostal veins, but seldom the whole number; also fix or seven, more or less, inferior intercostals, which run between the aorta and vertebræ; and the same ramifications on the right fide, and likewife fome ramifications to the œfophagus; but fometimes these distributions vary, and are not all subjects. alike. Just below the last rib, the azygos fends off a large branch to the muscles of the abdomen, communicating with the last two intercostal veins: sometimes this extremity communicates either mediately or immediately with the vena adipofa, and even with the vena spermatica.

The diaphragmaticæ superiores, mediastinæ, mammariæ internæ, thymicæ, pericardiæ, and tracheales, are small veins disposed in pairs to the right and left, behind the sternum and parts near it, and some called venæ pectorales internæ.

Q. Which is the vena mediastina?

A. The right vena mediastina goes out anteriorly from the trunk of the superior cava,

a little above the origin of the azygos; the left comes from the subclavia.

Q. Which is the vena diaphragmatica fu-

perior?

A. The right superior diaphragmatica arises near the mediastina, and its branches are distributed to the pericardium. The left comes from the left subclavian, a little below the origin of the mammaria.

Q. Which is the vena mammaria interna?

A. The right internal mammaria arises from the vena cava, a little below the angle of the bifurcation. It runs along the posterior edge of the sternum, and on the cartilaginous extremities of the right ribs, together with the artery of the same name; it also sends the diaphragm a branch, and communicates with those veins. Afterwards it gives small branches to the mediastinum and others between the ribs to the integuments. The left internal mammaria arises anteriorly from the left subclavian, opposite to the cartilage or anterior extremity of the first true rib.

Q. Which is the vena thymica?

A. The right vena thymica, when it arises separately, goes out from the bisurcation; this vein often reaches no lower than the inferior part of the thymus. The left vein of the same name comes from the left subclavian, almost opposite to the sternum.

Q. Which is the vena pericardia?

A. The vena pericardia seems to go out rather from the origin of the right subclavian, but but in this there are many varieties. It goes to the upper fide of the pericardium, and other neighbouring parts; the left pericardia comes fometimes from the left fubclavian, and sometimes from the mammaria on the same fide.

Q. Which is the vena trachealis?

A. The right trachealis goes out from the upper part of the bifurcation above the mammaria of the fame fide, sometimes more backward, and sometimes from the subclavia. It is distributed to the glandulæ thyroideæ, thymus, and bronchialis, trachea arteria, and musculi sterno-hyoidei: it communicates with the internal jugular vein. The lest trachealis comes from the upper, or posterior part of the lest subsclavian near its origin. Of all these small veins, the mammaria interna, is the most considerable.

Q. Which is the vena fubelavia?

A. The right subclavian vein (as has been already said) is very short, and its course very oblique, so that it appears to rise higher than the left vein. It sends off, first of all, sour large branches, already mentioned, viz. the vertebralis, which is the first and most posterior; the jugularis et axillaris. It sends off sour large branches besides the small pectoral veins, and receives the ductus thoracicus. Also a small trunk for the left superior intercostals, which are sometimes six in number, and communicate with the inferior intercostals, and with a branch of the vena azygos. This small common intercostal trunk furnishes like-wise

wife the left bronchialis. Each subsclavian vein, near the middle of the clavicle, sends off a branch called cephalica, which descends near the surface of the body, between the deltoides and pectoralis major, and reaches the arm in the manner which shall be described hereafter.

Q. Which is the vena cava inferior?

A. The two inferior vena cava having run down about a quarter of an inch from the right auricle of the heart, within the pericardium as has been already faid,) pierces that membrane, and the tendinous portion of the diaphragm which adhere very closely to each other: at this place it gives off the venæ diaphragmaticæ inferiores vel phrenicæ, the other branches, &c. of the inferior cava I shall describe hereafter.

Q. Which are the venæ phrenicæ?

A. The venæ phrenicæ are sent off from the vena cava inferior, as abovementioned, and are distributed to the diaphragm, and appear chiefly on its lower side, one to the right hand, the other to the left. The right vein is more backward and lower than the left. The left is distributed partly to the pericardium, and partly to the diaphragm, and sometimes branches to the capsulæ renales.

Q What are the nerves of the thorax?

A. The twelve pair of dorsal or costal nerves, which all communicate with one another, as soon as they make their way out betwixt the vertebræ; each of them gives a posterior branch to the musculi erectores trunci corporis; the first after having sent off the brachial

brachial nerve, already described, is, after the same manner with the succeeding eight, bestowed on the pleura and intercostal muscles; the tenth and eleventh are most of them sent to the abdominal muscles; the twelsth communicates with the first lumbar, and is bestowed on the musculus quadratus lumbalis and iliacus internus.

Q. What are the glans of the thorax?

A. The glands of the thorax are the glandulæ bronchiales, the thymus, and dorfalis. Besides these, the cosophagus has a great number of glands, (especially towards its upper part,) as before observed.

Q. Which are the glandulæ bronchiales.

A. The glandulæ bronchiales are very obfervable glands, fituated externally in the larger divisions of the trachea and bronchia. They are of a blackish colour, and their use, like that of many others of the glands of this part of the body, is yet very little known.

Q. Which is the glandula dorfalis?

A. The glandula dorfalis is a remarkable gland fometimes found in the thorax, about the fifth vertebra of the back, adhering to the posterior part of the cesophagus. It is in disferent subjects of various sizes; being often of the size of a kidney bean; sometimes of that of an almond, and sometimes considerably larger; in others, it is much less than the smaller, and sometimes it is wholly wanting, or at least so extremely minute, and inconsiderable, that the best dissectors are not able to find

dif-

it. Sometimes also two glands are found in this part in the place of one.

Q. Which is the glandula thymus?

A. The thymus is a gland which in infants is very remarkable: it is situated in the upper part of the thorax, immediately under the fternum, and lies upon the pericardium, and on the trunk of the aorta, and of the vena cava. It extends itself from the pericardium along. the trunk of the aorta, to the beginning of the carotids, fometimes fo far as to the thyroide gland; its figure is irregular and uncertain; its colour in infants is a pale red, in adults, it is of a duskier hue : it is much larger in infants newly born, than in subjects at any more advanced period. Its length in the former is no less than three fingers breadth, and its diameter two; its thickness is about half a finger: it gradually decreases from this fize as the child grows up; in adults it is very small, and in old people it entirely disappears. Its substance is glandular and conglomerate; and it is furrounded by a membrane; it has blood veffels of the same name; its lymphatics sometimes run to the thoracic duct, fometimes to the fubclavian veins, and they have in general no valves. The nerves of the thymus are from the par vagum, or from the intercostals; there is fometimes a milky juice found in this gland in new born subjects. It has no excretory duct hitherto discovered, and its use is therefore not certainly known: possibly, according to Heister, it serves to secrete lymph, which it discharges into the thoracic duct, for the dilution of the blood and of the chyle, as the glands of the mesentery and of the pancreas do, in regard to the chyle. On this supposition, its use is much greater in the sœtus, than at any time after the birth, because the want of respiration in that state may well be supposed to subject the blood to be thicker, and to need more dilution than afterwards; nothing tending to attenuate the blood so much as respiration.

Q. What are the mammæ or breafts?

A. The mammæ or breafts, are two glandular bodies, of a roundish oval figure, situated on the anterior, and a little towards the lateral parts of the thorax: these are most remarkable in women. They differ in fize according to age and fex; in very young and very old people they are always small, also in virgins and males of all ages: but in women with child, or those who give fuck, they are generally large, often very enormous. In children of both fexes, and in males of all ages, they are commonly no more than curaneous tubercles or foft veruceæ, of a reddish colour, called papillæ, or nipples; each of them being furrounded by a fmall, thin, and pretty broad circle, or disk, more or less of a brownish colour, and an uneven surface, termed areola. The time of the breafts growing full in women, is about the age of fourteen, or that of puberty, which is fometimes fooner, fometimes later; and the most natural time of their de-

decreasing is about the forty-feventh, or fiftieth year, when their menses totally cease, and the breafts become flabby, lofing their natural confistence and solidity. The breasts, besides the common integuments of the body, (already mentioned in the beginning of this work,) are composed of a glandular subtrance, and a multitude of lactiferous ducts, or small tubes, which unite by frequent anaftomofes, and, as they approach the nipples, join and unite together, till at last they form seven, eight, or more small pipes, called tubuli lactiferi, which have feveral cross canals, by which they communicate with each other; fo that if any of them be obstructed, the milk which was brought to it might not stagnate, but pass through by the other pipes, which all terminate in the extremity of the nipple. The substance of the nipple is cavernous, almost like that of the human penis. The excretory ducts or tubes are larger in women who give fuck, and are dilated into finuses in many places forming a kind of cells, which hold the fecreted milk, and communicate with the veins and arteries. All these parts are to be seen much more distinctly in breaks that are large and full of milk than in others; in young women, indeed, they they are scarce to be distinguished at all; as alfo in fuch as have little breatts, in fuch as are emaciated, and in those of very old people.

This glandular substance of the breatts is feated in, and furrounded by, a great quantity of fat; which makes by much the greater part

of the breasts. The arteries and veins of the breasts are called mammary vessels, and are fent from the subclavian and axillary vessels. The nerves are from the dorsal ones of the spinal marrow. A sulness of the breasts, especially if there be milk sound in them, is geneally judged a mark of the loss of virginity, and a proof that a woman has been with child; but

this is not an infallible fign.

The fwelling of the breafts during the time of gestation, is owing to the consent between the breasts and the uterus; there being so near a communication between the mammary veffels and the hypogastric vessels of the womb, that a dilatation of the latter is attended with a fimilar one in the former. For the tubes which compose the glandular substance of the breasts in maids, like a sphincter muscle, contract so closely, that no part of the blood can enter them; but when the womb grows big with a fœtus, and compresses the descending trunk of the aorta, the blood flows in a greater quantity, and with a greater force, through the arteries of the breafts, and forces a paffage into their glands, which being at first narrow, admit only of a thin water; but growing wider by degrees, as the womb grows bigger, the glands receive a thick ferum; and after birth, they run with a thick milk, because the blood, which before flowed to the fœtus, and for three or four days afterwards by the uterus, beginning then to stop, dilates the mammillary glands, and confequently swells the breaft.

DIALOGUE V.

Of the Abdomen, and its Parts.

A. The abdomen begins immediately under the thorax, and terminates at the bottom of the pelvis of the offa innominata. Its circumference or outer furface is divided into regions, of which there are three anterior, viz. the epigaftric, or superior region; the umbilical, or middle region: and the hypogastric, or lower region. There is but one posterior region, named regio lumbaris.

Q. Where is the epigastric region situated?

A. The epigastric region begins immediately under the appendix ensiformis, at a small superficial depression, called the pit of the stomach, and in adult subjects ends above the navel in a transverse line from the last false ribs on each side. This region is subdivided in three parts, one middle, named epigastrium, and two lateral, termed hypochondria. The epigastrium takes in all that space which lies between the salse ribs of both sides, and the hypochondria are the places covered by the false ribs.

Q. Where is the umbilical region fituated?

A. The umbilical region being in adults, above the navel from the transverse line, where the epigastric ends, and ends below the navel at another transverse line, supposed to be drawn parallel to the former, between the two cristæ

of the offa ilium. This region is likewife divided into three parts, one middle, which is properly the regio umbilicalis, and two lateral, called ilia, or the flanks; and they comprehend the space between the false ribs and upper part of the os ilium on each fide.

Q. Where is the hypogastric region?

A. The hypogastric region is extended downward from the inferior limit of the umbilical region, and is also divided into three parts, one middle called pubis; and two lateral, called inguina, or the groins.

Q. Where is the lumbar region fituated?

A. The lumbar region is the posterior part of the abdomen, and comprehends all that space which reaches from the lowest ribs on each fide, and the last vertebra of the back, to the os facrum, and neighbouring parts of the offa ilium. The lateral parts of this region are termed the loins, but the middle part has no

proper name in men.

The bottom of the abdomen, which answers to the pelvis of the skeleton, is terminated anteriorly by the pudenda, or parts of generation, and posteriorly by the buttocks and anus. The buttocks are separated by fossa, which lead to the anus, and each buttock is terminated downward by a large fold, which d'stinguishes it from the rest of the thigh. This lumbar region takes in likewise the musculus quadratus lumborum on each fide, the lower portions of the facro lumbares of the longissimi and latissimi dorsi, facer, &c. The space between

between the anus and the parts of generation is called perinæum, and is divided into two equal lateral parts by a very distinct line, which is longer in males than in females, (as I shall make appear hereafter.) The cavity of the abdomen, formed by the parts already mentioned, is lined on the infide by a particular membrane called peritonæum. The cavity of the abdomen is separated from the cavity of the thorax by the diaphragm, and terminated below by the musculi levatores ani. This cavity of the abdomen contains the stomach and the intestines, which are commonly divided into three small parts, named duodenum, jejunum, and ilium; and three large, called cæcum, colon, and rectum. It contains likewise the mesentery, mesocolon, omentum, liver, gall, bladder, spleen, pancreas, glands of the mesentery, vasa lactea, receptaculum chyli, kidneys, renal glands, ureters, bladder, and the internal parts of generation in both fexes. Also the following bones pertaining to this lower venter, viz. the five vertebræ of the loins, os facrum, os coccygis, and offa inominata which contain the ilium, ischium, and pubis. These bones, below the vertebra, form the pelvis or bason, which is much larger in women than in men, to give room for the growth, &c. of the fœtus. The external parts of generation also belong to the abdomen, but I shall describe them hereafter.

Q. What are the bones pertaining to the

abdomen?

A. The bones belonging to the abdomen are the vertebræ of the loins, which are the five lowermost vertebræ of the spine or backbone, and complete the whole twenty-four: (see those of the neck and thorax before described.) The os sacrum, os occygis, ossainnominata or pubes, which consists of the ilium, ischium, and pubis.

Q. Does the five vertebræ of the loins differ from those of the neck and thorax which you have already described, page III

and 135.

A. Yes! The five vertebræ of the loins differ from the rest in this, that they are the broadest, and the last of them is the largest of the vertebræ. Their acute processes are broader, shorter, and wider from one another, their transverse longer to support the bowels and muscles of the back; they are not perforated as those of the neck, nor have they a dimple or finus as those of the back. The cartilages which are betwixt their bodies are thicker than any of the rest. The greatest motion of the back is performed by the vertebræ of the loins, because their cartilages are thicker, and their acute processes are at a greater distance from one another; for the thicker the cartilages are, the more we may bend our body forwards; and the greater distance there is betwixt the acute processes, the more we may bend ourselves backwards.

Q. Which is the os facrum?

A. The os facrum is the lower extremity of the spina dors, the vertebræ of which grow so close together in adults, as that they make but one large and solid bone of a triangular figure, whose basis is tied to the last vertebræ of the loins, and the upper part of its fides to the ilia, and its point to the os coccygis. This bone in infants is almost entirely cartilaginous; and in children more grown up, it always confifts of several pieces, the junctures of four or five of which may be seen even in adults; although it be one continued bone. It has four or five holes on each fide which give passage to the nerves of the intestinum rectum, bladder, and parts of generation, and to the large crural and ischiatic ones. The uses of this bone, are to ferve as a basis for the spine; to form the pelvis along with the offa innominata, and to defend the parts contained in it; to serve as a place of origin to many of the muscles; to contain in its canal or finus, the lower part or end of the spinal marrow, called cauda equina; and likewise to give passage to the nerves abovementioned.

Q. Which is the os coccygis?

A. The os coccygis is joined to the extremity of the os facrum, and is composed of three or four bones; (but sometimes all the pieces are entirely cemented together,) the lower piece is still less than the upper, till the last ends in a small cartilage, which resembles a little tail turned inward. Its use is to sustain the rectum intestinum; it yields to the pressure of

of the fœtus in women in travail, and midwives sometimes thrust it rudely and violently backwards, which occasions great pain and several bad effects.

Q. Which are the offa innominata?

A. The offa innominata are composed of three bones in young subjects, each of which has its peculiar name; the upper one is called the ilium, (the intestinum ilium lying between it and its fellow,) the anterior one the os pubis, and the lower and posterior one the os ischium. The offa innominata are joined on each fide, in the hinder parts to the os facrum, forming a very firm and strong, though somewhat moveable, articulation with it; and with this bone they also form the cavity called the pelvis. The use of these bones is to support and suftain the spina dorsi, and indeed all the parts above them; to make a firm and proper juncture of the other parts of the body with the thighs; to serve for the place of origin to feveral muscles; to form the cavity of the pelvis, and to defend its contents from external injuries.

In the center of these bones is the acetabulum coxendicis, or socket for the thigh-bone; in the bottom of which sockets is another cavity, in which lies the mucilaginous gland of this joint. When imposshumations happen in this joint they usually cause a great swelling and lameness in the hip, which in time makes a collection of matter in the external part of the hip, &c. Q. What are the cartilages pertaining to the bones of the abdomen besides those of the

vertebræ already mentioned?

A. The true cartilages of the offa innominata in adult subjects, are five in number, three common and two proper. The first and principal common cartilage is that which makes the symphisis of the offa pubis; the two other common cartilages join the offa ilium to the os facrum, but are thinner than that of the offa pubis. The proper cartilages are those that line the cotyloide cavities, and except the notches and depressions in these cavities, all the rest of the surface of the acetabulum coxendicis is covered with a very white shining smooth cartilage.

Q. What are the ligaments of the bones pertaining to the abdomen, besides those of

the vertebræ already mentioned?

A. The ligaments of the offa innominata are also of two kinds, common and proper. The common are those which go between these and the neighbouring bones, of which there is a considerable number. To these must be added the ligaments by which the os semoris is joined to the os innominatum, which I shall describe hereafter among the other ligaments of the thigh. The principal proper ligaments are four in number, two called sacro-sciatic, one broad and external, the other small and internal, one obturator, and one inguinal. The broad sacro-sciatic or internal sciatic ligament proceeds from the inside of the spine of the crista of the

os ilium to the false transverse apophyses of the os facrum and the spine of the ischium. The internal sciatic ligament adheres closely to the infide of the posterior portion of the former, going from the fourth false transverse apophyfis of the os facrum, all the way to the upper part of the os coccygis. The obturator ligament fills up all the great foramen ovale, except the oblique notch at its proper part, from which there is a transverse ligament to the infide of the upper and anterior part of the os pubis. The inguinal ligament, called from the discoverer ligamentum Fallopii, is an aponeurotic or ligamentary band, fastened by one end to the anterior and superior spine of the os ilium, and by the other to the spine of the os pubis. Another ligament runs transversely between the two angles of the cotyloide notch.

Q. Where is the mucilaginous gland belonging to the os innominatum fituated?

A. The rough unequal depression at the bottom of the acetabulum, or focket for the thigh bone, is filled by a broad flat mucilaginous gland, bordered with a fatty fubstance, and covered by a fine membrane, through which a mucilaginous liquor passes to moisten the joint, and facilitate its motions. This membrane rifes above the gland, and gives a fort of covering or coat to the ligament contained in the joint; the blood veffets of the gland pass between the bottom of the cotyloide notch, and the transverse ligament thereof.

Q. What

Q. What are the mucilaginous glands of

the joints?

A. The mucilaginous glands of the joints, where the bones are furnished with a cartilage are small glands which furnish every joint for a fliding motion, with a mucilaginous matter, for jubricating the ends of the bones, that they may move eafily upon one another; and that there may be no waste of this necesfary fluid, it is contained in the investing ligaments; which for this very reason are no where divided, except to communicate with the ligaments of the tendons. These glands are generally seated in a little fat, near the infertion of the ligaments, that they may be compressed by them when the joints are in motion, which is a proper time to have their fluid pressed out. The most considerable parcel of these glands with their fat, are seen in the joint of the knee, and the largest gland of this fort, is found in the finus, at the bottom of the acetabulum of the os innominatum, and is compressed by the ligamentum teres.

The diseases of the joints either happen from ulcers in the mucilaginous glands, when pouring out matter that cannot be discharged, and foul the ends of the bones, or elfe from fwellings in the ends of the respective ones. Either of these in time create excessive pain, which appears to be chiefly in the ligaments of the joints, notwithstanding what has been said of the infensibility of these parts. When a joint is much swelled and painful, without external inflammation, it is yulgarly called a white

fwelling, and more properly than a spina ventofa. It is fometimes in the beginning cured by evacuations, but when the limb wastes below the swelling, and the fingers or toes of the limb grow thinner at their joints, and lose their shape, the case is absolutely irrecoverable. Sometimes the ends of the bones erode, then join together, and form an anchylofis, which, though a fevere difease of itself, yet proves often the remedy of a much worfe. In like manner, the bones of the hands and feet, when they are ulcerated, fometimes unite, and are thus preserved from total ruin. But there is one case of a white swelling that is amazing, when the pain is fo great that we are forced to take off the limb, and yet neither find upon diffection, the ligaments or glands difeafed, matter in the joints, the bones carious, nor any difeased appearance, except that the ends of the bones are a little larger and fofter.

Q. What are the muscles of the abdomen?

A. The muscles peculiar to the abdomen only are five pair, (exclusive of the diaphragma, and triangularis sterni, already spoken of with the muscles of the thorax) viz. obliquus ascendens vel externus, obliquus descendens vel internus, pyramidalis, rectus abdominis, and transversalis abdominis. To these we may add those of the genital parts, anus and perinæum, viz. cremaster testis, erector penis, accelerator urinæ, transversalis penis, sphincter vestcæ urinariæ, detrusor urinæ, erector clitoridis, sphincter vaginæ, sphincter ani, elevator

ani, intertransversales lumborum, psoas parvus, quadratus lumborum, and coccygei.

Q. Which is the obliquus ascendens mus-

cle ?

A. The obliquus ascendens vel externus arises from the eight inferior ribs, the upper part of its origin being indented with the serratus major anticus, and the lower lying under a small portion of the latissimus dorsi: it is inserted into the spine of the ilium, into the os pubis, and linea alba, which is a strong tendinous line extended from the os pubis to the sternum, between the musculi recti.

Q. Which is the obliquus descendens?

A. The obliquus descendens vel internus arises, under the former muscle of the spine of the ilium, os sacrum, and spines of the loins, and is inserted partly into the lower side of the spurious ribs, and by a flat tendon into the sternum and linea alba, together with the tendon of the acendens.

Q. Which is the pyramidalis?

A. The pyramidalis is a small muscle lying in the lower part of the rectus. It has the name from its sigure, and its origin from the margin of the os pubis, with a broad sleshy head, but ends in a small round tendon in the linea alba, about three or four inches below the naval. This muscle is sometimes double and sometimes single, and sometimes they are both wanting.

Q Which is the rectus abdominis?

A. The rectus abdominis, arises from the sternum

stemum near the cartilago ensiformis, and the extremity of the two last ribs. It goes strait down to the forepart of the abdomen, and is inferted into the os pubis.

Q. Which is the transversalis abdominis?

A. The transversalis abdominis arises from the cartilago ensisormis, the extremities of the false ribs, and the transverse apophysis of the vertebræ of the loins: it is fixed to the inner side of the spine of the ilium, and then becoming a slat tendon, it passes under the rectus, and is inserted into the os pubis and linea alba. Between this tendon and the peritonæum, sometimes water is found in great quantities, constituting a dropsy in the duplicature of the peritonæum, which shews this membrane has been mistaken for part of the peritonæum.

The uses of these abdominal muscle are to fustain the viscera of the abdomen, and to compress the parts contained therein, in order to clear it of what ought to pass off by the natural outlets, to relieve the stomach by vomiting, from whatever might be hurtful to it; and laftly, to drive out by a violent expiration whatever may incommode the organs contained in the thorax. The obliquus descendens on the right fide, and afcendens on the left, acting together turn the upper part of the trunk of the body towards the left, and vicê versa; but the trunk is chiefly turned upon the thighs; the recti bend the body forward, and pull the sternum downward in expiration; the two obl que muscles and the transverse on each side near the groins, are perforated to let through the processus vaginalis with the spermatic vesfels. These perforations are distant from each other, so as to suffer the vessels to descend conveniently into the scrotum: this way the intestines or the omentum descend in ruptures.

Q. Which is the cremaster testis?

A. The cremaster testis muscle is so named because (with its fellow) it suspends the testicles, and draws them up in the act of generation: it arises from the os ilium, and upper part of the ligamentum pubis, and almost encompassing the process of the peritonæum (which furrounds the spermatic vessels as they come out of the abdomen) descends with it, and is inferted into the tunica vaginalis, upon which it is spread in several distinct portions.

Q. Which is the erector penis?

A. The erector penis and its fellow ferve for the erection of the penis. These arise on each fide from the offa ischii, and each of them is inferted into the corpus cavernofum of the same side. These muscles when they act together, press the veins of the back of the penis against the os pubis, by which they prevent the reflux of blood from the penis; and confequently when at the same time the blood flows impetuously into the part by the arteries, and cannot get back this way, the penis becomes extended and erect.

Q. Which is the accelerator urinæ?

A. The accelerator urinæ arifes tendinous from the offa ischia, and fleshy from the sphincter ani, and being expended over the bulb of the urethra, afterwards divides, and is inferted into the penis. The use of this muscle is not to accelerate the urine, for that is propelled by the detrufor urinæ, or mufcular coat of the bladder, but to protrude the femen, which is done only by this. They likewife affift the erectores in the erection of the penis, by driving the blood contained in the cavernous body of the urethra towards the glands, which is thereby diftended; the tumetaction of these muscles at the same time compressing the veins that carry off the refluent blood from the corpus cavernosum.

Q. Which is the transversalis penis?

A. The transversalis penis vel perinæi, is one of the dilators of the urethra, arising from the tubercle of the os ischium on each side, and inserted into the posterior part of the bulb of the urethra. These muscles, however, are not quite determinate and certain in their origin or insertion, and sometimes they are wholly wanting. When they act, they dilate the urethra in its posterior parts.

Q. Which is the sphincter vesicæ urinariæ?

A. The sphincter vesicæ urinariæ is composed of tranverse sibres, which form a circle round the neck of the bladder serving to close it, to prevent the involuntary discharge of the urine. In men this muscle is connected to the

fibres

fibres of the intestinum restum, and in women to those of the vagina.

Q. Which is the detrusor urinæ?

A. The detrusor urinæ is the muscular coat of the bladder, its fibres terminate in the fphincter veficæ, whereby it not only preffes the urine forward, but, when the bladder is full, becomes an antagonist to the sphincter, acting almost at right angles.

Q. Which is the erector clitoridis?

A. The erector clitoridis arises from the ischium, and is inserted into the corpora cavernosa of the clitoris, like the erector penis in men, and is faid to cause erection in the same manner.

Q. Which is the sphincter vaginæ?

A. The sphincter vaginæ arises from the sphincter ani, and surrounds the orifice of the vagina; after which it is inferted under the crura of the clitoris. Its use is to constringe the orifice of the vagina, to press out a liquor from the glands of the vagina, and embrace the penis in coition.

Q. Which is the sphincter ani?

A. The sphincter ani is a muscle near two inches broad, composed of circular fibres, which closes the extremity of the intestinum rectum, and forms the anus. It is connected forward with the accelerator urinæ in men, and with the neck of the uterus in women, and backwards with the os coccygis. This muscle furrounds the anus to close it, and to prevent involuntary falling out of the fæces. Q. Which

Q. Which is the elevator ani?

A. The elevator ani arises from the offa ischii, pubis, and sacrum, within the pelvis, and is inserted round the lower end of the intestinum rectum. It surrounds also the collum vession, glandulæ prostatæ, and vesiculæ seminales in men, and the vagina in women. The use of this muscle, is to sustain and elevate the anus, lest the sæces should be burthensome to the sphincter; and to press the prostatæ and vesiculæ seminales, in order to promote the emission of the seminal juices in coition.

Fistulæ in ano, that are within this muscle, generally run in the direction of the gut, and may be laid open into the gut with great safety; but those fiftulæ or rather abscesses that are frequently formed on the outfide of the fphincter, and usually furround it, all but where this muscle is connected to the penis, cannot be opened far into the gut, without totally dividing the sphineter, which authors say renders the sphincter ever after incapable of restraining the excrement: but this does not always hold true; for there have been many instances of the sphinster being divided, which made the patients unable to hold their excrements during the cure, but the wounds being healed, they have retained them as well as ever.

Q. Which are the intertransversales lumbo-

A. The intertransversales lumborum are small muscles seated between all the transverse processes

to

processes of the vertebræ lumborum, to bring them nearer together.

Q. Which is the ploas parvus?

A. The ploas parvus is one of the flexor muscles of the loins, which arises laterally from the body of the upper vertebra of the loins, and the last of the back, and is inserted into the os pubis, where it is joined to the ilium.

Q. Which is the quadratus lumborum?

A. The quadratus lumborum has its origin in the anterior and superior part of the posterior process of the ilium, and is inserted into all the transverse processes of the vertebræ of the loins, the last vertebra of the thorax, and the last rib. This with its fellow, acting alternately, assist the extensor dorsi et lumborum, in raising the ossa innominata in progression: or each acting singly, while the lower limbs are not moved, inclines the body to one side.

Q. Which are the coccygæi?

A. The coccygaei arise from the acute processes of the ossa is is is and are inserted into the os coccygis, which they pull forward.

Q. What is the peritonæum?

A. The peritonæum is a thin, foft, smooth, and lubricous membrane, which lines the whole cavity of the abdomen, containing most of the viscera of that part, as it were in a bag. It has a number of small glands that separate a liquor, which supplies the intestines, and facilitates their motion: the external superficies of this membrance is unequal where it adheres

The uses of the peritonæum are, to enclose the contents or viscera of the abdomen; for when it is dilated, wounded, or broken, they fall out of their proper places, and ruptures are formed. It also gives an external covering to almost all the parts contained in the abdomen, and forms the process of the peritonæum, and the tunica vaginalis of the testes.

The dropfy of the peritonæum may be diflinguished by being least prominent about the navel, for there the tendons of the muscles and the peritonæum will not separate; and the water, in some that have been diffected, has been found to have made the parts where it was contained as foul as any ulcer; therefore none of them could have been cured by operation.

Q. What are the viscera of the abdomen?

A. The viscera of the abdomen, are the stomach, omentum, duodenum, jejunum, ilium, cæcum, colon, rectum, mesentery, mesocolon, liver, gall bladder, porus biliarius, pancreas, spleen, kidneys, renal glands, ureters, and bladder; besides the vasa lactea, receptaculum chyli, and ductus thoracicus already spoken of, and the urethra and internal and external parts of generation, which I shall hereafter describe.

Q. What is the stomach?

A. The ventriculus or stomach, is a hollow membranous part, placed mostly in the left hypochondrium, immediately under the left side of the diaphragm, its left side touching the spleen, and its right covered by the thin edge of the liver. Its figure nearly resembles a pouch of a bag-pipe, but its upper side is concave, and the lower convex; it has two orifices, both on its upper parts: the left called

led cardia, is placed much higher than the right, continuous to the gula, through which the aliment passes into the stomach; its right orifice is called the pylorus, through which the aliment is conveyed out of the stomach into the duodenum: in this part, there is a circular valve which closes the stomach, and hinders a return of aliment out of the gut, but does not at all times hinder the gall from flowing into the stomach. The pylorus is connected to the upper part of the stomach by a ligament. The fize of the stomach in human subjects is various; in people addicted to gluttony it is usually very large; and in men it is generally larger than in women. Its veffels are arteries, veins, nerves, and lymphatics. The gastric arteries it receives from the cæliac; and the gastric veins all run to the vena portæ: among these are observable the vala brevia, which go off to the splenic branch, and the vena coronaria, which furrounds the stomach. Its nerves principally enter at the left orifice; they come from the par vagum, and are very large, whence it is that the stomach is so sensible: the lymphatics go to the receptaculum chyli. The substance of the stomach is membranaceous, and is composed of four coats, as the oefophagus. The first, or external coat, is membranous, the fibres of which run transversely; the fecond coat is muscular, whose fibres are chiefly longitudinal and circular; the third is nervous, this forms a multitude of wrinkles, and is furnished with a number of sanguiferous vessels, and small glands, which secrete a mucus; the fourth coat is thin, villofe, and porous, and adheres very firmly to the former. As the muscular coat contracts, the inner coat falls into folds, which increase as the stomach lessens, and consequently retard the aliment most when the stomach is nearest being empty. The use of the stomach is for the digestion of our food, that is to receive, contain, dissolve, and change what is swallowed by the mouth; and after a sufficient concoction, to expel it through the pylorus into the intestines; possibly it also absorbs, and retains the most Subtle parts of what it has thus prepared for nutrition: but I shall speak more of this hereafter.

Q. What is the duodenum?

A. The intestinum duodenum is the first of the three small guts, so called from its being about twelve fingers breadth in length. It begins from the pylorus, from which turning downwards, it first passes by the gall bladder, and then under the jejunum and mesentery, and coming in fight again in the left hypochondrium, it there commences jejunum, which is the fecond of the small guts; but the place where this ends and the other begins, is not exactly determined. About four fingers breadth from the pylorus it receives a duct from the liver and gall bladder, called ductus communis choledochus, and another from the pancreas, called pancreations, at a little distance from each other, which discharge their respective liquors liquors into it. The coats of the duodenum are thicker than those of any other of the small guts, its cavity is also greater, and its passage in a straiter direction than any of them. Near its origin it has no valves, nor rugæ or wrinkles; but in its continuation it has very numerous and remarkable ones, called by authors juga. It has also the glands of Brunnerus in great number, which serve for the secreting of a thin aqueous shuid, and it receives an artery from the cæliac and a vein from the porta.

Q. What is the jejunum?

A. The jejunum is so called from its being usually found empty, which is owing to the fluidity of the chyle, the greater stimulus of the bile in it, and the abundance of the lacteal vessels with which it is furnished. It is situated in the regio umbilicalis, and being the fecond of the small guts, it begins where the duodenum ends, and terminates where the valves are obliterated. Its length is different in various subjects; but is usually between thirteen and fixteen hands breadth long, making fomewhat more than a third part of the small guts. It is distinguished from the ilium by its coats, which are a small matter thinner, and less pale; it receives arteries from the mesentericæ, and veins from the mesaraicæ, and has also a great many connivent glands and valves.

Q What is the ilium?

A. The ilium is the continuation of the jejunum, and is the third and last of the small guts. It is situated in the hypogastrium, and very

very often some part of it in the pelvis, upon the bladder, especially in women; filling all the space between the ilia, whence its name. Its length is various, fometimes not more than fifteen, sometimes twenty-one hands breadth, or more. Its beginning is where the valves of the jejunum cease to be conspicuous, and its end is where the larger intestines begin; in which place, it is in a very fingular manner inferted into the right fide of the colon, near the upper edge of the os ilium. It has no other valves except that great one at the end which is called by many valvula coli Bauhini : its glands are, in general, more numerous towards the end than in any other part. The great length of the small guts is evidently for the convenience of a greater number of lacteals, that the chyle which misses the orifices in one place may not escape them in another. This intestine, because of its situation, falls easily down into the scrotum, by the production of the peritonæum.

Q. What are the thick and great guts?

A. The thick and great guts are the cæcum, colon, and rectum; but the cæcum here meant, is the head of the colon.

Q. What is the cæcum?

A. The cæcum, or apendicula vermiformis, is the only true cæcum, though the antients describe it otherwise. It is situated on the beginning of the colon, and is properly an appendage of that gut. It is about four singers breadth long, and the bigness of an earth-worm

or goose-quill, with a small orifice opening into the colon: this gut has seldom any thing in it. It is called cæcum, or blind, because it is open only at one end, by which it is sixed to the beginning of the colon; its other end, which is shut, is not tied to the mesentery, but to the right kidney, by means of the peritonæum. In human subjects, the cæcum is called one of the large guts, though it is the smallest by far; this mistake is owing to the ancients taking their description of the viscera from dogs and some other animals in whom the cæcum is very large, as Chelseden and others observe.

Q. What is the colon?

A. The colon may be reckoned the first of the great guts, as I have before observed. It is the greatest and widest of all the intestines, and about eight or nine hands breadth long. It begins where the ilium ends, in the cavity of the os ilium, on the right fide; from thence, ascending by the kidney on the same side, it passes under the concave side of the liver, to which it is fometimes tied, as likewife to the gail bladder, which tinges it yellow in that place; then it runs under the bottom of the stomach to the spleen, in the left side, to which it is also knit; from thence it turns down to the left kidney; and thence, passing in form of an S, it terminates in the upper part of the os facrum, in the rectum. At the beginning of this gut there is a valve formed by the production of the inmost coat of the intestines : in this

this place it hinders the excrements, which are once fallen into the colon, from returning again to the ilium. It has a strong ligament, which, running along its upper fide from the ilium to the rectum, strengthens it against the weight of the excrements, and draws it together into the cells, which, with the valvulæ conniventes, retards the paffage of the excrements, that we may not be obliged continually to go to stool. The fleshy fibres of its fecond coat are greater and stronger than those of the other intestine, because greater strength is requisite to cause the excrements to ascend. The connivent valves are larger in this than in any other of the guts, as well as the coats stronger.

Q. What is the rectum?

A. The rectum is the last of all the inteftines. It is about two hands breadth long, and three fingers breadth broad; it begins at the upper part of the os facrum, where the colon ends, and going strait down (whence its name) it is tied to the extremities of the coccyx, by the peritonæum behind, and to the neck of the bladder in men, but in women to the vagina uteri before, from whence comes the fympathy between those parts. The coats of the rectum are more thick and fleshy than those of any other of the intestines: it has in general no valves, but several rugæ: the absence of valves here, is to prevent the expulsion of the fæces from being retarded. The extremity of this gut forms the anus. The lower end of this gut is the feat of the true fiftula

in ano, which usually runs betwixt the muscular coat and the inner coat; it is cured by opening it the whole length into the cavity of the gut; it is yet better, if it can be done, to extirpate all that is fiftulous and fchirrous. The other kind of fiftula, improperly fo called, is an abscess running round the outside of the sphincter, in the shape of a horse-shoe, being a circle all but where this muscle unites with those of the penis; this is best cured by opening and removing part of the outer skin. The former of these cases happens oftenest in full habits, proceeding frequently from the piles; the latter is generally a critical discharge, and one of nature's last efforts in consumptive and fcorbutic habits of body. The inversion and sliding down of this gut is called prolapfus ani, a difease common in children, especially those who are afflicted with the stone, and not of much consequence; in men it is more rare and dangerous, being generally attended with a flux of humours. I have feen cases related of a prolapfus ani being cured by taking away a piece of the prolapsed gut with a caustic, lengthways of the gut; when the wound discharged the flux of the humours, the gut was easily reduced, and cicatrizing in that state, it never more fell down. Another, where a bold unthinking furgeon having cut off the prolapsed part, the cicatrix was so hard and contracted, that the patient could never after go to stool without a glyster, and then not without great misery. Oftentimes the piles occalion

occasion large tumours at the lower end of this gut: these are always best extirpated by ligature; for if they are cut, they will fometimes bleed excessively, and it is no easy matter to apply any thing to stop a flux of blood in that part. The guts have the same coats as the stomach, and the great guts have three membranes or ligaments, running on the outfide their whole length, and supporting the facculi into which those guts are divided. The lesser guts have, at very small distances, semi lunar valves, placed opposite to the interstices of each other, to prevent the aliment from paffing too speedily through the guts; and the better to answer that end, they are larger and more numerous near the stomach, where the food is thinner, than they are towards the colon, where the food is continually made thicker in its progrefs by a discharge of part of the chyle. This contrivance, so necessary to men because of their erect posture, when they are obliged by sickness or accidents, to lie along, becomes a great inconvenience, and calls for the help of glysters and purges. But brutes have not these valves, because they are not convenient in an horizontal posture. At the entrance of the ilium into the colon, are two very large valves, which effectually hinder the regress of the fæces into the ilium. Clysters, indeed, have been frequently known to pass them, and be vomited up; but the excrement that is fometimes vomited up I am apt to think, is fuch as had not passed into the great guts. The other valves in the colon are placed opposite, but not 111

in the same place, to each other, and make with their anterior edges an equilateral triangle; but as the gut approaches the anus, they become less remarkable, and fewer in number, as before observed. All the intestines have in their inner membrane an almost infinite number of very small glands; some of them in the large guts will appear to the naked eye when they are deceased. Their office is to difcharge into the intestines; a liquor which ferves for the attenuation of the chyle, for lubricating the intestines and in the large guts, to loften the fæces, that they may be evacuated without pain. The use of the smaller guts is to promote the formation of the chyle, to perfect its fecretions, and to propel the remaining fæces to the larger. The office of the larger guts is to receive and collect the matter of the fæces, and at a proper time to expel it. The length of the guts to that of the body is as five to one in a middle fized man; in taller men the proportion is usually less, and in fhort men greater. The intestines have veffels in great abundance, running over every part of their substance. Their arteries are from the meseraic, serving for the smaller intestines. the lower for the larger; and these make a multitude of very fingular and furprizing anaftomoses. The veins are meseraics, and go off to the vena portæ and the liver. The nerves are fent from the intercostals, and the par vagum; besides these, we are to observe the lacteal veffels already described. The rectum receives blood veffels also from the hypoga-Arics.

Q. What is the omentum?

A. The omentum, epiploon, or caul, is a fine membrane larded with fat, somewhat like net-work. It is fituate under the peritonæum, and immediately above the intestines on the furface of the fmall guts, and refembles an apron tucked up. Its outer or upper part, named ala superior, is connected to the bottom of the stomach, the spleen, and part of the intestinum duodenum; and thence defcending a little lower than the navel, is reflected and tied to the intestinum colon, the spleen, and part of the duodenum; this last part is called ala inferior, and the space between the alæ is named bursa. Sometimes both alæ are tied to the liver, and in diseased bodies adheres to the peritonæum. The uses of the omentum are, to lubricate the intestines, that they may the easier perform their peristaltic motion, and to cherish and defend them from cold; to assist in the formation of the bile, the fatty part of which is wholly owing to the veffels of the omentum; every thing (according to Malpighi) that returns from this part going to the liver through adipose ducts to the vena portæ. It ferves also to temperate the acrimony of the humours, and probably to give nourishment to the body, as all the other fat is supposed to do, when it is incapable of being nourished any other way. The arteries of the omentum come from the cæliac, and are very numerous: its veins arise chiefly from the splenic branch of the vena portæ, and its nerves are from the intercostals and par vagum.

Q. What is the melentery?

A. The mesentery is a thick fat membrane, placed in the middle of the abdomen, or midst of the intestines, particularly of the smaller ones, whence it has the name. It is almost of a circular figure, with a narrow production, to which the end of the colon and beginning of the rectum are fastened. It is about four fingers in breadth, and half a hand's breadth in diameter; its circumference is full of plaits and foldings, and about three ells in length; the intestines are tied like a border on this circumference, fo that to every inch of the mesentery, there are about three inches of the intestines fastened. Its substance is composed of membranes, fat, veffels of all kinds, and a number of glands. Its coats or membranes are two, and between these there is a cellular fubstance, which contains the fat: the meseraic veffels and glands are also placed there, which many reckon a third coat of the melentery, and that not improperly; this they call the tunica cellulofa. In the upper part it is strongly fastened to the three superior vertebræ of the loins; and in the lower, with the intestines, and particularly with the jejunum and ilium. The veffels of the mesentery are blood-veffels, nerves, lacteals, and lymphatics. The blood-veffels are the same with those of the intestines, and these make a multitude of strange meanders, and have very frequent anastomoses.

anastomoses. The nerves also come from the par vagum, and the intercostals. There are a number of glands dispersed throughout the whole mesentery, from which they take their name: these vary greatly in their size, sigure, and situation in different subjects, and in old people they frequently almost disappear. The uses of the mesentery are to preserve the jejunum and ilium from twisting in their peristaltic or vermicular motion, and to confine the rest to their places, to sustain the sanguiferous and lacteal vessels of the intestines, and to make the way for the lacteals to the receptacle the shorter.

Q. What is the mefocolon?

A. The mesocolon is that part of the mesentery connected with the great guts, and especially the colon. The mesocolon meets the middle of the colon, to which it is joined. Its lower part sticks to a part of the rectum.

Q. What is the liver?

A. The liver is the largest gland in the body, of a dusky red colour, situated immediately under the diaphragm in the right hypochondrium. Its sigure is almost round, the upper surface convex, smooth, and equal; the lower, hollow and unequal; backward, towards the ribs it is thick, and thin on its fore part, where it covers the upper side of the stomach, and some of the guts. In its middle and fore part, it is divided into two by a sissure, where the umbilical vessels enter. It is fastened in the body by two ligaments; the first, which is large and strong.

strong, comes from the peritonæum that covers the diaphragm, to which the upper side of the liver adheres, and is thus tied to it and the sternum, being named latum or suspensorium: the second is the umbilical vein, which after the birth degenerates into a ligament called teres or umbilicale; it is inserted at a small siffure in the lower edge of the liver. The ligamentum latum vel suspensorium, sustains the liver in an erect posture, or rather sixes it in its situation, while it is supported by the other viscera, they being compressed by the ab-

dominal mufcles.

In lying down, the teres prevents it from preffing on the diaphragm; and in lying on the back, they both together fuspend it, that it may not compress and obstruct the ascending vena cava. The veffels of the liver are very numerous, receiving arteries from the cæliac and mesenterica superior, called arteriæ hepaticæ; veins from the vena cava, and vena portæ; and nerves, from the plexus hepaticus of the intercostals. The biliary vessels are the ductus choledochus communis, which opens obliquely into the duodenum; the ductus cyfticus, which runs from the gall bladder to the common duct; the ductus hepaticus, which runs from the liver to the common duct; and the branches of this distributed through the liver, make what are called pori biliarii. The liver has also a great number of lymphatics, most of which open near the vena portæ, or the concave fide of the liver; from thence the

the lympha is carried by other lymphatics to the receptaculum chyli. To these vessels we may add the canalis venofus, and the great finus of the vena portæ in the fœtus. The vena cava, and the vena portæ, are accompanied in the liver with many small branches of the cœliac and meseraic arteries already mentioned. The vena portæ supplies the place of an artery, and brings the blood full of bile for fecretion, which being strained off, the vena cava returns the blood which remains. The vena portæ and the cava, enter the liver by its concave fide, and are equally distributed through all its substance; wherever there is a branch of the one, there is also a branch of the other.

Q. What are the excretory ducts of the liver?

A. The excretory ducts of the liver are the vesicula fellis, and porus biliarius.

Q. What is the vesicula fellis?

A. The vesicula fellis, or gall bladder, is a receptacle of bile, fastened to the concave part, or under side of the liver; its sigure is like that of a pear, and, in general, of the size of a small hen's egg, though it differs in bigness in almost every person. When the liver is in its natural situation, the bottom, or largest part of the gall bladder, is downwards, and the neck, or narrow part upwards; and then is touches the stomach as well as the colon, where it frequently dyes them yellow. This bladder is composed of three membranes, or coats;

strong, comes from the peritonæum that covers the diaphragm, to which the upper side of the liver adheres, and is thus tied to it and the sternum, being named latum or suspensorium: the second is the umbilical vein, which after the birth degenerates into a ligament called teres or umbilicale; it is inserted at a small fissure in the lower edge of the liver. The ligamentum latum vel suspensorium, sustains the liver in an erest posture, or rather sixes it in its situation, while it is supported by the other viscera, they being compressed by the ab-

dominal muscles.

In lying down, the teres prevents it from preffing on the diaphragm; and in lying on the back, they both together fuspend it, that it may not compress and obstruct the ascending vena cava. The veffels of the liver are very numerous, receiving arteries from the cæliac and mesenterica superior, called arteriæ hepaticæ; veins from the vena cava, and vena portæ; and nerves, from the plexus hepaticus of the intercostals. The biliary vessels are the ductus choledochus communis, which opens obliquely into the duodenum; the ductus cysticus, which runs from the gall bladder to the common duct; the ductus hepaticus, which runs from the liver to the common duct; and the branches of this distributed through the liver, make what are called pori biliarii. The liver has also a great number of lymphatics, most of which open near the vena portæ, or the concave fide of the liver; from thence the it together a certain time, and then to expel it.

Q. What is the porus biliarius?

A. The porus biliarius is another excretory veffel of the liver; and, according, to some, is the same with the ductus hepaticus: but I, as well as many others, make a distinction between them, and have already observed that the hepatic duct runs from the liver to the ductus choledechus. The porus biliarius has as many branches as the vena portæ, which it accompanies through every part of the liver. Wherever there is a branch of the one, there is a branch of the other; and these two are inclosed in one common capsule, as in a sheath. The use of this capsule is to facilitate the motion of the blood and bile, by the contraction of its fibres. All these branches unite, and make one trunk, of the bigness of a small quill, which joins the end of the cyftic duct, carrying the bile from the liver to the intestines by the common duct, as was said The infertion of the porus biliarius into the cyftic duct is oblique, with its mouth looking towards the ductus communis, by which means it is impossible that the bile, which comes from the cyftis, can enter the porus biliarius, unless the common duct is stopped.

As the liver, from its situation in the same cavity with the stomach, will be most pressed, and consequently separate most gall when the stomach is fullest, which is the time when it is most wanted; so the gall bladder, being seat-

P 2

ed against the duodenum, will have its fluid pressed out by the aliment passing thro' that gut, and consequently at a right time, and in due proportion; because the greater that quantity of aliment is, the greater will be the com-

pression: and so the contrary.

There is no way of computing with any exactness the quantity of bile that is usually fecreted by the liver in a given time: but if it is four times as much as all the falivary glands fecrete, it may be twenty-four ounces for every meal; to which being added fix ounces of faliva, which appears to be a moderate computation, supposing the pancreas in the fame time fecretes three ounces, there will then be thirty-three ounces of fluids separated for the digestion of one meal; and that these neceffary fluids may not be wasted in such quantities, they pass into the blood with the chyle, and may be foon separated again for the same use, and very likely, some of the same bile may be employed more than once, for digefting part of the fame meal. As the liver exceeds all the glands in the body in magnitude, and its excretory ducts ending in the duodenum, it feems to me to be much more capable of making those large separations from the blood, which are procured by cathartics, than the scarce visible glands of the guts. The liver ordinarily weighs, in a middle fized man, about three pounds twelve ounces: the pancreas, three ounces; and the spleen, fourteen nunces.

Q. What is the pancreas?

A. The pancreas, commonly called the Iweet-bread, is a large gland of the falivary kind, lying across the upper and back part of the abdomen, near the duodenum, behind the stomach, and between the liver and spleen. Its length is eight or nine inches, its breadth about two fingers, or two and a half, its thickness about one finger, and its weight about three ounces, as before observed. The shape of the pancreas very much resembles the tongue of a dog; it is broadest towards the duodenum, and gradually narrower towards the spleen. It is surrounded with a membrane, which is continuous with the peritonæum; it has arteries from the cæliac and splenic branches, and veins also from the splenic vein; its nerves are from the par vagum and the intercostals; and finally it has a short excretory duct, which is fituated in the middle of the pancreas, where it resembles an empty vein, and is about the thickness of a very small straw. This duct terminates in the duodenum, which it enters obliquely, four or five fingers breadth below the pylorus usually at the same orifice with the ductus choledochus, but sometimes it has a double aperture. The use of the pancreas is to secrete a peculiar liquor, called the pancreatic juice, which is of a falivose nature, and is carried by the pancreatic duct into the duodenum, where it ferves to dilute the chyle, or render it more fluid and fit to enter the mouths of the lacteals; and, perhaps, to temper and dilute the the bile, to change its viscidity, bitterness, colour, &c and make it mix with the chyle, in order to assimilate the several tastes, odours, and properties of the several foods, into one ho-

mogeneous one.

In those who die of the jaundice, for the most part, are found in the gall-bladder, and the biliary ducts, concretions of bile so light as to swim in water, yet are called gall stones: these cause the jaundice, by obstructing the ducts; many persons who have been cured of this disease, have had great numbers of these stound in their excrements.

Q. What is the spleen?

A. The spleen is a viscus of a deep blackish red colour, situated on the left side of the stomach, immediately under the diaphragm, near the ribs and above the left kidney. figure of it is a fort of depressed oval, near twice as long as broad, and almost twice as Its usual fize is five or fix broad as thick. inches long, three broad, and one inch thick; but this varies in different subjects. It is supported by the contained parts, and fixed to its place by an adhesion to the peritonæum and diaphragm; it is also connected with the stomach, by the vafa brevia, and with the pancreas, omentum, and left kidney, by mem-The substance of the spleen is vascubranes. There are two lymphatic lar and fibrofe. glands of about the bigness of a bean, situated without it, near where the vessels enter. The veffels of the spleen, considering its size, are remarkably

fremarkably larger. Its artery is from the cæliac, and is called the splenic artery; the splenic vein, is like those of the other viscera, very much ramified, and its branches are carried throughout the whole spleen; the nerves of the spleen are from the plexus splenicus; the spleen has no excretory duct; but there are in it lymphatic vessels running to the receptaculum chyli. The use of the spleen has been much controverted by authors; but the most probable opinion seems to be, that it serves to render the blood more sluid, out of which the bile is to be afterwards secreted.

What are the kidneys?

A. The kidneys are two dark-red viscera of an oblong figure, refembling a large bean; situated one on each fide the vertebræ of the loins, in the posterior part of the abdomen, upon the two last ribs. The right kidney lies under the great lobe of the liver, and is confequently lower than the left, which lies under the spleen; the concave side of the kidneys is turned inward to the vertebra, and the convex fide outward. The kidneys are connected with the loins, the lower ribs, the colon, glandulæ renales, the renal veffels, and the ureters. They have two membranes, the one common from the peritonæum, called the adipose membrane, from its being generally covered with much fat; this furrounds them but loosely, and is furnished with its own proper vessels. The other membrane is proper, is very thin, and every where applied closely to the substance of

the kidneys; the substance of the kidneys is firm and hard, and is of two kinds. The exterior, or cortical part, according to Malpighi, is glandular, but according to the discoveries of Ruysch, is throughout elegantly vascular. The interior is tubulous, and expressed by the name of tubuli urinarii Bellini; these terminate in ten or twelve papillæ, which open by a multitude of apertures into the pelvis; but these papillæ are not found in all subjects. This length answers to the distance between the two last false ribs and the os ilium, which is generally about five or fix fingers breadth; they are about half as broad as long, and half as thick as broad. The veffels of the kidneys are like those of the liver, included in a membrane, from the peritonæum. The arteries and veins are large, and called emulgents, and renal veffels; thefe are produced from the aorta and vena cava: the nerves are from the plexus renalis, proceeding from the intercostals. There is also a number of lymphatics paffing to the receptaculum chyli; and also an excretory duct, called the ureter, which I shall by and by describe. The use of the kidneys is to secrete the urine from the blood, which is brought there for that purpose by the emulgent arteries; and what remains from the fecretion, is returned by the emulgent veins, while the urine fecreted is carried off through the uteters to the bladder.

Q. What are the glandulæ renales?

A. The

A. The glandulæ renales, called also capfulæ atrabiliariæ, and renes succenturiati, are
two yellowish glands of a compressed figure,
lying on each side of the upper part of the kidneys, a little above the emulgent vessels. They
have a very narrow cavity imbued with a brownish siquor of a sweetish taste, and are about the
bigness of a large nux vomica generally in adults; in the sœtus they are larger, and often
exceed the kidneys themselves in size. The
very thin membrane that surrounds them closely involves their whole substance, and connects
them with the kidneys. They have blood vessels and nerves, and their lymphatic vessels are
numerous.

Q. What are the ureters?

A. The ureters are tubes about the bigness of goofe-quills, and about a foot long; they arise from the hollow side of the kidneys, and terminate in the urinary bladder near the neck. At their origin in the kidneys they are expanded into the form of a funnel, and this expanfion makes the pelvis of the kidneys: at their termination in the bladder, they pass obliquely for the space of an inch between its coats, which manner of entering is to them as valves, for their orifices being narrow, will admit of nothing into them from the bladder. The ureters are not strait, but somewhat bent, so as to resemble the letter S. Their substance is membranous, and they are composed of three coats; the first is from the peritonæum; the fecond is a thin muscular one; and the third, a nervous a nervous one, covered with a slimy liquor to defend it against the acrimony of the urine; and in this, there are sometimes discovered glands; the blood vessels and nerves come from the adjacent parts. Such as are subject to the gravel, and given to excess of drinking, have them sometimes so much dilated, that you may put the end of the little singer into them. Their use is to carry the urine from the kidneys to the bladder. Their obstruction causes a suppression of urine.

Q. Where is the bladder fituated, and how

composed?

A. The urinary bladder is feated in a duplicature of the peritonæum in the lower part of the pelvis; its shape is orbicular, and its coats are the same with those of the guts already defcribed, viz. an external, membranous, a middle muscular (which is the detrusor urinæ,) and an inner membranous or nervous coats which is covered with a peculiar fluid of a mucous nature, fecreted in glands fituated in this coat, and principally in that part which is near the neck of the bladder. The coats of the bladder are much thinner in the body and the fundus (which is the bottom, fituated uppermost) than they are at the neck and lowest part. The sphineter, in the neck of the bladder, closes it, to prevent an involuntary discharge of urine. The bladder in adults will conveniently hold about a pint; but it is capable of diffention to as to hold much more. It is connected in the human body, in a fingular

gular manner, by the peritonæum to the os pubis, different from other animals: it is also connected with the parts of generation by the urethra, with the navel by the urachus and umbilical arteries, and finally in men with the intestinum rectum, and in women with the vagina. The blood vessels of the bladder come from the hypogastric, the umbilical, and the hæmorrhoidal vessels in men; and in women from the spermatics also. Its nerves are from the intercostals, and principally from those of the os facrum. The bladder has three foramina; two where the ureters enter in, at which the urine passes into the bladder; and one, much larger than thefe, in the neck, for the discharge of the urine into the urethra. The nervous or inner coat of the bladder is most exceeding fensible, as is miserably experienced in the stone and gravel.

Q. What are the arteries of the abdomen?

A. The arteries of the abdomen are various, and proceed from the aorta, already defcribed. They take their names from the parts they are bestowed on, viz. arteriæ diaphragmaticæ vel phrenicæ, arteria cæliaca, ventriculi coronaria, hepatica, pylorica, gastrica, intestinalis, cysticæ et bilaria, splenica, pancreaticæ, mesentericæ, hæmorrhoidalis, renales, capsulares, spermaticæ, lumbares, sacræ, iliacæ, hypogastrica, umbilicalis, glutæa, sciatica, pudicæ, and obturatrix.

Q. Which are the arteriæ diaphragmaricæ

vel phrenicæ?

A. The

A. The arteriæ diaphragmaticæ, called also phrenicæ, arise from the aorta in two branches, as it passes under the diaphragm, which are ramified on the diaphragm, sometimes it arises from the cæliaca, and fometimes a trunk from each. They give small branches to the glandulæ renales, and membrana adipola of the kidneys, the latter being called arteriæ adiposæ. Smaller diaphragmatic arteries come from the intercostals, mammariæ internæ, mediastinæ, pericardiæ, and cæliaca.

Q. Which is the arteria cæliaca?

A. The arteria cæliaca arises from the aorta descendens, immediately below the diaphragm, and is foon after branched out to the liver, pancreas, spleen, stomach, omentum, and duodenum, which branches are named from the parts they are bestowed on, except the ventriculi coronaria bestowed upon the stomach, and the branch upon the duodenum named inteftinalis.

O. Which is the arteria ventriculi coronaria?

A. The arteria ventriculi coronaria, arises from the celiaca, and is ramified on every part of the stomach, and sends off small branches to part of the liver, neighbouring parts of the diaphragm, and omentum.

Q. Which is the arteria hepatica?

A. The hepatica, arises also from the cæliaca, and goes to the pylorus with the vena portæ, fending off the pylorica and gastrica dextra, and is then ramified with the vena portæ through all the liver.

Q. Which is the arteria pylorica?

A. The pylorica arises from the hepatica, and is ramified on the pylorus and neighbouring parts of the stomach.

Q. Which is the arteria gastrica dextra?

A. The gastrica dextra, arises also from the hepatica, and passing behind and beyond the pylorus, sends off the duodenalis vel intestination, when it does not come from the hepatica; after this, it runs along the right side of the great curvature of the stomach, and sends off branches to all the neighbouring parts on both sides, which communicate with those of the pylorica, coronaria ventriculi, and gastro-epiploicæ dextra, the latter furnishing the nearest parts of the omentum, and communicating with the mesenterica superior, at last the right gastric ends in the left, which is a branch of the splenica.

Q. Which is the duodenalis vel intestinalis?

A. The duodenal or intestinal artery runs along the duodenum on the side next the pancreas; to both which it sends branches, and also to the neighbouring part of the stomach.

Q. Which is the cysticæ and biliaria?

A. The hepatic artery having sent out the pylorica and gastrica dextra, gives two branches to the vesicula fellis, called arteria cysticæ, and another named biliaria, which is lost in the great lobe of the liver. Afterwards, the hepatic artery is ramified with the vena portæ through all the substance of the liver.

Q. Which is the arteria splenica?

222 ANATOMICAL

A. The splenica arises close by the hepatica, the hepatic branch going to the right, and the splenic to the left; immediately after its origin from the cæliaca it runs toward the left under the stomach and pancreas to the spleen.

Q. Which are the pancreaticæ?

A. The arteriæ pancreaticæ are branches of the splenica ramsied on the pancreas.

Q. Which is the arteria gastrica sinistra?

A. Under the left portion of the stomach it gives off a branch called gastrica sinistra, to the left fide of the stomach, which communicates with those of the coronaria ventriculi. This gastric artery also supplies the omentum with branches, termed gastro-epiploicæ siniftræ, which communicate with the right gaftric. By this account it appears that the coronaria ventriculi, pylorica, intestinalis, gastrica, gastro epiploicæ and consequently the hepatica, splenica, and mesenterica, communicate all together. After this, before the splenica arrives at the spleen, it gives two or three branches to the large extremity of the stomach, commonly called vafa brevia; and one to the omentum, named epiploica.

Q. Which is the mesenterica superior?

A. The arteria mesenterica superior, arises anteriorly just below the cæliaca, and gives off branches to part of the pancreas and duodenum communicating with the intestinalis; then going between the laminæ of the mesentery, and the duodenum and meseraic vein, it distributes a number of branches to the small intestines,

intestines, from the lower third part of the duodenum, to the cæcum and colon. As they approach the intestines, all these branches communicate first by reciprocal arches; then by areolæ and the meshes of all kinds of figures, from which is detached an infinite number of small ramifications, which surround the intestinal canal like an annular piece of net-work.

Q. Which is the mesenterica inferior?

A. The mesenterica inferior arises anteriorly from the aorta descendens inferior, about an inch above the bifurcation, and below the spermatic arteries; soon after it sends branches to the colon and rectum, communicating with one another; it also sends off a branch called hæmorhoidalis interna.

Q. How is the hæmorrhoidalis interna dif-

tributed?

A. The hæmorrhoidalis interna runs down behind the intestinum rectum, to which it is distributed by several ramifications, and communicates with the arteriæ hypogastricæ.

Q. Which are the arteriæ renales?

A. The renal arteries, commonly called emulgents, are commonly two in number, and go out laterally from the inferior descending aorta, immediately under the mesenterica superior, one to the right hand, the other to the lest. They exter into the depressions of the kidneys by several branches, and are ramified through all the substance of the kidneys. Sometimes the renal arteries and branches to the

the glandulæ renales, membrana adiposa of the kidneys, and even to the diaphragm.

Q. What are the arteriæ capfulares?

A. The arteriæ capsulares are the arteries of the renal glands, which arise sometimes from the arteria formation of the cæliaca; but generally the right comes from the arteria renalis of the same side, near the origin; the lest, from the arter above the renalis.

Q. Which are the arteriæ spermaticæ?

A. The arteriæ spermaticæ are commonly two in number, fometimes more; they are very fmall, and go out anteriorly from the aorta descendens inferior, near each other, about an inch below the renales, between the two mefentericæ, or between the renales and mefentericæ inferiores. They fend off small branches to the common membrane of the kidneys, named adipofæ; and afterwards theyrun down upon the ploas muscles, on the ferefide of the ureters, between the two laminæ of the peritonæum. They give several considerable branches to the peritonæum, chiefly to those parts of it which are next the mefentery, and they communicate both with the mesentericæ and adipofæ. They likewise send small arteries to the ureters. Afterwards, they pass in men through the tendinous openings of the rings of the abdominal muscles in the vagina of the peritonæum, (termed processus vaginalis) and are distributed to the testicles and epididymes, where they communicate

tum,

municate with a branch of the iliaca externa. In women they do not go out of the abdomen, but are distributed to the ovaria and uterus, and communicate with branches of the hypogastrica, at the jagged extremities of the tubæ Fallopianæ.

Q. Which are the lumbar arteries?

A. The arteriæ lumbares, or lumbar arteries, go out posteriorly from the inferior descending aorta, in five or fix pairs, or more, much in the same manner with the intercostals. The fuperior fend small branches to the neighbouring parts of the diaphragm and intercostal muscles. They are distributed on each side to the pfoas muscle, quadrati lumborum, and to the oblique and transverse muscles of the abdomen; and by perforating the oblique muscles, they become external hypogastric arteries. They go likewise to the vertebral muscles and bodies of the vertebræ; and enter the spinal canal through the lateral notches, forming rings much in the same manner with the intercostals; and they likewise give small twigs to the nerves.

Q. Which are the arteriæ facræ?

A. The arteriæ sacræ are generally two, three, or sour in number, and sometimes but one. They go out commonly from the back part of the inferior descending aorta at the biturcation, but sometimes from the lumbares, and sometimes from the iliacæ. These arteries are ramified on the os sacrum, and neighbouring parts of the peritonæum, intestinum rec-

tum, fat, &c. They likewise send small arteries to the nerves which go through the holes of the os sacrum, and they penetrate the inner substance of that bone.

Q. Which are the arteriæ iliacæ?

A. The arteriæ iliacæ are those two large lateral branches arising at the bifurcation, where the aorta descendens inserior ends at the last vertebra of the loins, and sometimes higher. They run obliquely toward the anterior and lower part of the ossailium, sending off a few very small arteries to the ossaicrum, neighbouring muscles, &c. likewise small arteries to the peritonæum, coats of the veins, fat and ureters, behind which the iliac trunks pass about three singers breadth from their origin, each iliac trunk is divided into two secondary arteries, one external, the other internal; the former has no particular name, but the latter is termed hypogastrica.

Q. How is the external iliae distributed?

A. The external iliac on each fide runs down on the iliac muscle to the ligamentum Fallopii, under which it goes out of the abdomen, and as it passes under this ligament, it detaches two considerable branches one internal the other external. The internal branch is named epigastrica, and goes out anteriorly from the external iliac; from thence it runs obliquely upward on the tendon of the transvese muscles, and to the posterior and inner part of the rectus muscle, sending ramifications

to the tendons of the neighbouring muscles, &c. and afterwards cummunicating with the mammaria interna and intercostalis inferior. It fometimes gives out a branch accompanied by a nerve to the triceps muscles, &c. through the foramen ovale. And another branch which runs down to the testicles along with the spermatic artery, and there communicates with it. The outfide branch of the external iliac divides in two at the internal labium of the os ilium, where it divides into two, and is ramified on the oblique and transverse muscles of the abdomen, communicating with the arteria lumbares. Besides these, a small twig goes off internally to the vagina of the spermatic rope, and fometimes another from the outfide to the os ilium.

Q. How is the hypogastric or internal

'iliac artery distributed?

A. The hypogastric or internal iliac, having run about an inch and a half inward and backward, bends by degrees, and ends in the umbilical artery, which ought to be looked upon as a true continuation of the trunk of the hypogastric.

Q. How is the arteria umbilicalis distri-

buted?

A. The umbilical artery (as above) ascends on the side of the bladder, and being ramissied on that viscus, and the neighbouring parts of the peritonæum, &c. it contracts, and in adults is quite closed up above the middle of the bladder: it likewise gives branches to the

Q 2

uterus, and to the neighbouring parts in both fexes. Afterwards it afcends in form of a ligament to the umbilicus, its name being taken from its use in the fœtus.

Q. Which is the arteria iliaca minor?

A. The iliaca minor arises from the convex fide of the curvature of the hypogastrica, and is the most posterior branch: it passes between the two lumbar nerves, and divides into two; one branch goes to the canal of the os facrum, the other to the iliac muscle, and inside of the os ilium, giving twigs as it passes to the ploas muscle.

Q. Which is the arteria glutæa?

A. The glutæa arises also from the convex fide of the curvature of the hypogastric, is commonly very confiderable, and fometimes the largest of all the hypogastric branches: this artery goes out of the pelvis in company with the sciatic nerve, through the upper part of the great finus of the os innominatum, and is distributed to the glutæus maximus and medius. In it passage it gives branches to the os facrum, os coccygis, mufculus pyramidalis, muscles of the anus, and to the neighbouring parts of the intestinum rectum, (forming a particular hæmorrhoidalis interna) and to the bladder and parts near it, and a pretty long branch runs down with the sciatic nerve.

Q. Which is the arteria fciatica?

A. The arteria sciatica arises near the glutæa, and gives branches to the musculus pyramidalis, quadragemini, os facrum, sciatic nerve, os ilium, os ischium, musculi glutæi, and and a branch runs under the quadratus to the articulation of the os femoris.

Q. Which is the arteria pudica?

A. The pudica arises also from the convex side of the curvature of the hypogastric; and the sirst principal branch called commonly pudica interna, gives out two branches, the sirst goes through the great sinus of the os ilium with the glutæa and sciatica, and then divides into two, one branch goes behind the spine of the ischium all the way to the origin of the

corpus cavernofum penis.

There it divides into several arteries; one goes to the fphinctor ani, named hæmorrhoidalis externa; the rest are distributed to the neighbouring integuments, to the bulb of the urethra, and to the corpus cavernosum penis. The fecond principal branch, commonly called pudica externa, runs between the bladder and intestinum rectum, and in men is distributed to the vesiculæ seminales, neck of the bladder, proftatæ, and neighbouring part of the rectum; it then runs under the os pubis on the fide of a confiderable vein; and along the penis between this vein and a nerve, being diftributed in its paffage to the corpus cavernofum. Sometimes the pudica externa goes off separately from the hypogastrica, especially in women, being distributed to the uterus, (where it communicates with the spermatic artery) and to the neighbouring parts of the vagina, &c.

Q. Which is the arteria obturatrix?

Q 3

A. The

A. The arteria obturatrix perforates the obturator muscles, from whence its name; it goes out of the pelvis at the upper part of the ligament of the foramen ovale, having first sent a small branch to the inguinal glands and integuments. In its passage it supplies the pectineus and triceps muscles, communicates with the sciatica, which goes to the articulation of the os semoris, and gives small arteries to the holes in the neck of that bone.

Q. What are the veins of the abdomen?

A. The veins of the abdomen are numerous, and all proceed from the vena cava (the vena portæ excepted) and chiefly from the vena cava inferior, which trunk, having run down about a quarter of an inch from the right auricle of the heart, within the pericardium (as I have already observed) pierces that membrane, and the tendinous portion of the diaphragm, which adheres very closely to each other; at this place it gives off the venæ diaphragmaticæ or phrenicæ. In describing the veins, I shall begin by the great trunks, and end by the ramifications in the method I have before observed. The veins of the abdomen are, venæ diaphragmaticæ or phrenicæ, hepaticæ, cysticæ gemellæ, pylorica, gastrica dextra, coronaria ventriculi, duodenalis vel inteftinalis, mesaraica, epiploica dextra, cæcalis, hæmorrhoidalis interna, splenica, pancreaticæ, gastrica sinistra, gastro-epiploica and epiploica sinistra, vasa brevia; venæ renales, capsulares et adiposæ, spermaticæ lumbares, biturcatio

ex vena cava, vena facra, venæ iliacæ, vena iliaca externa, epigastrica, hypogastrica vel iliaca interna, obturatrix, venæ hæmorrhoidales externæ.

Q. Which are the venæ diaphragmaticæ vel

phrenicæ?

A. The venæ diaphragmaticæ, or phrenicæ arise from the vena cava inferior, where they pierce the midriff, are distributed to the diaphragm, and partly to the pericardium; and sometimes they send branches to the capsulæ renales, much in the same manner as the arteriæ phrenicæ.

Q. Which are the veneæ hepaticæ?

A. The vena portæ is a large vein, the trunk is fituated chiefly between the eminences on the lower or concave fide of the liver. It may be confidered as made up of two large veins joined almost endways by their trunks, from each of which the branches go to the right and left; but first of all the trunk sends off the venæ cysticæ gemellæ, gastrica dextra, pylorica, ventriculi coronaria, duodenalis or intestinalis. The first portion of this trunk known by the name of vena portæ I have already spoken of. The other portion is the hepaticæ.

Q. Which are the venæ cysticæ gemellæ?

A. The cysticæ gemellæ run along the vesicula fellis from its neck to the bottom; they arise from the right side of the great trunk near its beginning.

Q. Which is the vena pylorica?

A. The pylorica arises from the great trunk, almost opposite to the origin of the cysticæ;

but sometimes it is only a branch of the right gastrica: it supplies the stomach and omentum, and communicates with the gastrica, coronaria ventriculi, &c.

Q. Which is the vena coronaria ventri-

culi?

A. The coronaria ventriculi, so called because it surrounds more or less the upper orifice of the stomach; and in it passage forms numerous areolæ on the sides of the stomach, it communicates with the pylorica and the veins of the great arch.

Q. Which is the vena duodenalis?

A. The duodenalis vel intestinalis, goes out from the great trunk near the cysticæ gemellæ. It is distributed chiefly to the intestinum duodenum, but sends some branches to the pancreas.

Q. Which is the vena mesaraica?

A. The mesaraica arises from the trunk of the portæ near the pancreas, and turns to the right hand; it accompanies the superior mesenteric artery and mesocolon which belong to the small intestines, the cæcum, and right portion of the colon. One branch of the mesaraica runs directly to the middle of the colon, where it is arched to the right and left; another branch divides in two, and supplies the head of the pancreas and the neighbouring parts, and is arched on the right portion and upper part of the colon. See vena epiploica.

Q Which is the vena epiploica dextra?

A. The

A. The trunk of the mesaraic sends out fometimes a particular branch to the omentum called epiploica dextra: but the mesaraiac almost immediately before it ascends over the mesenteric artery, send two large branches to the jejunum and part of the ilium, which form arches and areolæ on those intestines, like those of the artery, and sends out branches almost in the same manner.

Q. Which is the vena cæcalis?

A. This branch is the mesaraic vein called by Riolan cæcalis, arifes a little below the origin of the fecond branch from the convex side, and supplies the cæcum, appendicula vermiformis, and beginning of the colon.

Q. Which is the vena hæmorrhoidalis interna?

A. The hæmorrhoidalis interna arises from the beginning of the mesaraica, sometimes from the splenica, and sometimes between both at the bifurcation; and is thus named from the hæmorrhoidal tumors found often at its extremity next the anus. It goes to the duodenum, colon, and on the rectum, all the way to the anus; and in its course supplies the mesocolon, and forms arches which fend out numerous small ramifications round these intestines. It feems likewife to communicate, by fome capillary twigs, with the left spermatic vein.

Q. Which is the vena splenica?

A. The splenic vein is one of the great branches of the portæ, and turns to the left hand, where the meseraiac does to the right;

it runs transversely from right to left, just under the duodenum, and along the lower side of the pancreas, and in its course gives off the vena pancreaticæ, gastrica sinistra, or gastroepiploica sinistra and epiplocia sinistra. It terminates afterwards by a winding course, being divided into several branches that go to the spleen, one of which produces the small veins called by the ancients vasa brevia; and at last the vena splenica reaches the sissue of the spleen, which it enters, and supplies almost in the same manner as the splenic artery.

Q. Which are the venæ pancreaticæ?

A. The venæ pancreaticæ are several small branches sent by the splenica to the pancreas; but there are other small pancreatic veins, which do not arise from the splenica, which I have already spoken of.

Q. Which is the gastrica sinistra, or gastro-

epiploica finistra?

A. This is also a branch of the splenic vain arising at the left extremity of the pancreas, from whence it runs to the great extremity of the stomach, to which it gives several branches, and communicates with the gastrica dextra, and coronaria ventriculi: at a small distance from its origin this gastric vein sends out a branch which is distributed to the omentum, from whence the name of gastro-epiploica?

Q. Which is the vena gastro-epiploica, and

epiploica finistra?

A. The vena epiploica finistra is also a branch of the splenic, and arises at the small extremity of the pancreas, and is ramified on the omentum all the way to the colon, where it communicates with the hæmorrhoidalis interna.

Q. Which are the vafa brevia?

A. From the posterior of the branches of the splenic vein, the veins are sent off to the great extremity of the stomach, formerly known by the name of vasa brevia; which communicate with the coronaria ventriculi, and gastrica sinistra.

Q. Which are the venæ renales?

A. The vena cava inferior having got as low as the arteriæ renales, gives off the veins of the same name, (termed formerly venæ emulgentes) and which are the largest of all the veins that go from the cava inferior, from the liver to the bifurcation. They accompany the renal arteries, and are ramified in the substance of the kidneys.

Q. Which are the venæ capfulares?

A. The venæ capsulares et adiposæ, arise from the venæ renales; the former goes upwards to the glandulæ renales, the latter downwards to the fatty covering of the kidneys.

Q. Which are the venæ spermaticæ?

A. The right vena spermatica comes from the trunk of the cava inferior, a little below the venæ renales. The left comes commonly, though not always, from the left renalis; both veins accompany the spermatic arteries. In their passage small branches are sent off to the peritonæum and mesentery, and sometimes a branch to the iliac muscle and membrana adiposa of the kidneys.

Q. Which are the venæ lumbares?

A. The venæ lumbares arise posteriorly from the vena cava inferior most commonly in pairs, in the same manner as the arteriæ lumbares, but their origin often vary. They all communicate with one another, and sometimes with the intercostals. They supply the substance of the bodies of the vertebræ, muscles of the abdomen, quadratus lumborum, psoas, iliacus, &c. they send branches also backwards to the neighbouring vertebral muscles, and to the canal of the spine, and communicate with the venal sinuses in the same manner as the intercostals.

Q. Where is the bifurcation of the vena cava fituated?

A. The venæ cava inferior, having reached as low as the last vertebra of the loins, and near the bifurcation of the aorta, runs in behind the iliac artery, and is there divided into two subaltern trunks, called the right and lest iliac veins, which lie on the inside of the iliac arteries which form the bifurcation of the vena cava.

Q. Which is the vena facra?

A. The vena facra goes out from the bifurcation of the vena cava, and often from the origin of the left iliaca, and accompanies the arteria facra to the os facrum, to the nerves which

lie there, and to the membranes which cover both sides of that bone.

Q. Which are the venæ iliacæ?

A. The iliac veins arise from the bifurcation of the cava, as before mentioned, each original iliac vein is divided on the fide of the os facrum, much in the fame manner as the arteries, into two large trunks or fecondary iliac This fecond bifurcation is about a finger's breadth below that of the iliac arteries. One of these trunks is named vena iliaca externa, or anterior, the other interna, or posterior. The external vein is likewise named fimply iliaca, and the interal, hypogastrica. These veins follow nearly the course and distribution of the iliac arteries, only the hypogastric does not send off the vena umbilicalis. Near this fecond bifurcation, a particular branch goes out, to the musculus psoas, iliacus, and quadratus lumborum, and a branch of it to the transverse apophysis of the loins.

Q. Which is the vena iliaca externa?

A. The iliaca externa, a little before it leaves the abdomen, near the ligamentum Fallopii, lying on the pfoas and iliac muscles, gives off the same branches as the external iliac artery, and follows the same course. See vena epigastrica.

Q. Which is the vena epigastrica?

A. The epigastrica arises from the inside of the iliaca externa, before it leaves the abdomen, and is ramissed on the neighbouring conglobate glands and abdominal muscles, communicating with the ramifications of the mammaria which accompanies the epigastric artery. From the inside of this vein a branch is sometimes detached to the musculus obturator internus, where it joins another branch named vena obturatrix.

Q. Which is the vena hypogastrica?

A. The vena hypogastrica, or internal iliac vein, runs behind the artery of the same name, making the same kind of arch, from which branches go out to the cavity of the os sacrum, through the first and second great hole of the os sacrum to the neighbouring muscles. A large branch runs behind the great sciatic sinus to the musculi glutæi pyramidalis and gemelli. Another branch reaching the foramen ovale of the os inominatum, perforates the obturator muscles, from whence it is called vena obturatrix.

Q. How is the vena obturatrix distributed?
A. The vena obturatrix is distributed to the musculus pectinæus, triceps, and neighbouring parts, communicating with the cruralis. One branch of the obturatrix, before it perforates the muscles, runs towards the sciatic sinus to the iliac muscle os ilium and part of the obturator internus, another branch is distributed to the ureters, bladder, and internal parts of generation in both sexes. It communicates with the spermatic veins, and is more considerable in women than in men. Lastly, the hypogastric vein runs backward, and as it goes out of the pelvis it is ramisfied both

both upward and downward: it fends a large branch upward to the lower part of the os facrum, and two more downward, which going out of the pelvis are distributed to the buttocks, anus, part of the musculus pectineus, and to the external parts of generation, nearly in the same manner with the artery which accompanies them.

Q. Which are the venæ hæmorrhoidales

externæ?

A. The veins that go to the anus are termed hæmorrhoidales externæ, and they that go to the parts of generation, pudicæ intemæ. The external hæmorrhoidales communicate with the internal veins of the same name.

Q. How are the nerves of the abdomen

distributed?

A. The five pair of lumbar nerves also communicate, and give posterior branches; the first pair sends several branches to the abdominal muscles, psoas, and iliacus, while others go from it to the teguments and muscles on the superior and anterior part of the thigh, and the main trunk of it is lost in the crural. The second pair passes through the pleas muscle, and is distributed much as the former. The third is loft in the musculus pectineus. Branches proceeding from these three pair make up one trunk, which supplies the anterior part of the pelvis, and going through the foramen magnum offis ischii, is spent on the triceps muscle. This nerve is commonly known known by the name of obturator, or posterior crural nerve. By the union of branches, from the first, second, third, and fourth lumbar nerves, the anterior crural nerve is formed. The remainder of the fourth and fifth lumbar nerves join with the first, second, and third, that proceed from the os facrum (which are five or six pair). These two last of the lumbar and three first of the facrum, when united, constitute the largest nerve in the body, so well known by the name of the sciatic or ischiatic nerve. The other nerves that come out of the os facrum, are sent to the organs of generation and adjacent parts.

Q. What glands are there in the abdomen?

A. In the abdomen there are a very confiderable number of glands: the largest of them is the pancreas, after this, in fize come the glandulæ renales; then the mesaraics; and the intestinals (of Brunner and Peyer) in the intestines; all which I have already spoken of: the glands of the stomach are so small as not easily to be found. Near the receptaculum chyli, and above the os facrum, and the divisions of the iliac veffels, are many glands of various fizes and figures, called lumbares, facræ, iliacæ, &c. and they have numerous lymphatics entering into them, and discharging their contents into the receptaculum chyli. The lumbar glands have been fometimes found fwelled to the bigness of a man's fift. In the concave part of the liver, about the ingress of the vena portæ and the neck of the gall bladder,

bladder, as also about the spleen, near the ingress of the vessels, there are frequently found conglobate glands, of about the bigness of a kidney bean: these are called by authors, hepatic glands, cyftic glands, or have other names derived from the adjacent parts; and they feem to ferve the lymphatic vessels. About the left orifice of the stomach, there fometimes also is found, according to Vercellonius, a gland which he fays is equal to a kidney-bean in fize; he also says that it has ducts opening into the cavity of the stomach; in the gall-bladder there are fometimes found a number of small glands of a yellow colour, not unlike the ceruminose glands in the auditory passage. The bladder and the ureters have also sometimes a number of small glands, but they are very indeterminate in number and fize, and are not always indeed found in the fame place, especially about the ureters. Those about the bladder are usually situated towards the neck of it, and are sometimes tolerably conspicuous. The glands in the parts of generation of both fexes I shall describe hereafter with those parts.

DIALOGUE VI.

Of the ARM, and its PARTS.

Q. THAT are the parts of the arm? A. The arm comprehends the whole limb or member from the top of the shoulder to the finger's ends; to which you may add the clavicle and scapula, with the mufcles thereto belonging. The arm includes the whole member, but the fore-arm is from the elbow to the wrift only, the upper part of the arm, which joins with the scapula or shoulder-blade and clavicle, is termed humerus, which extends to the bend of the arm, the backfide of which is called the elbow or olecranon, and the forefide or flexure, ancon; from thence toward the hand is called the forearm or cubitus; the end of which towards the hand is the wrift or carpus; between the carpus and the fingers is the metacarpus, to which is articulated, pollex, the thumb; index, the forefinger; digitus medius, the middle finger; digitus annularis, the ring finger; and digitus minimus, the little finger.

Q. What are the bones of the arm?

A. The bones of the arm, including the fcapula, clavicle, hand and fingers, are thirty-two, (besides the sesamoidal bones, whose number is uncertain,) viz. scapula, clavicula, humerus, ulna, radius, each one; carpus, eight; metacarpus

metacarpus, four; phalanges of the fingers and thumb, fifteen; as may be feen at one view in the table of ofteology.

Q. Which is the clavicula?

A. The clavicula is fituated transversely on each fide, between the scapula and sternum, its figure is somewhat like that of the letter S, and it is of a fpongy and brittle substance. One end is connected to the sternum with a loofe cartilage, and the other to the processus acromion of the scapula. The uses of the clavicles are, to keep the scapulæ at a sufficient distance from the breast, by which means the shoulders are hindered from coming near together, as they do in those quadrupeds which use their four-limbs only to walk on, and not as men do their hands; to facilitate several of the motions of the arm; to serve for the place of origin for feveral muscles; and to defend the great subclavian vessels which run under them.

Q. Which is the scapula?

A. The scapula, or shoulder-blade, is a triangular bone, situated on the back and outside of the ribs, and commonly extended from the second to the seventh. It is fixed to the sternum by the clavicula, to the ribs and spine by the muscles of those parts; its outside is a little convex, and its inside somewhat concave, to fit it to the outer surface of the ribs on which it moves, and partly to give room for the subscapularis muscle. The scapula has three processes, the first runs all along the R 2 middle

middle of their outside, and is called their spine; that end of the spine which receives the extremity of the clavicula is called the proceffus acromion. The fecond process is a little lower than the acromion; it is short and sharp like a crow's bill, whence it is called coracoides; these two processes are tied to each other by a strong flat ligament, which serves to keep the head of the humerus in the cavity of the third process, and prevents its being dislocated upwards. It has a round finus tipt about its brim with a cartilage, which receives the head of the humerus; at the fore-part of this edge, close to the coracoid process, is a femicircular nitch for the passage of blood veffels, which nitch is joined at the top with a ligament, and fometimes with a bone. The use of the scapula is to receive the extremities of the clavicula and humerus, for the easier motion of the arm, and to give rife to the muscles which move the arm.

Q. Which is the os humeri?

A. The os humeri is the large bone of the arm, fituated between the shoulder and the cubit, articulated at one end with the scapula, and at the other with the ulna and radius. Its upper end or head, where it is joined to the scapula, is somewhat flat, and much larger than the socket which receives it. At the upper part are two processes for the insertions of the muscles of the arms: between these processes is a long channel or groove, in which lies a tendon of the biceps cubiti. At the

lower end are two confiderable processes, both formed to give origins to muscles of the wrist and fingers; between these processes is the joint. That part to which the upper end of the radius is fixed, is fitted not only for the motion of the elbow, but also for the rotatory motion of the radius; the rest of this joint is made up of portions of uneqal, but concentric circles, like the shanks of quadrupeds; which inequality prevents the ulna from diflocating fideways, which fo small a joint with fo much motion would be very fubject to. Of a like use is the little sinus on the forepart of the humerus, and the large one behind; the first of which receives a process of the ulna when the arm is bent, and the other the olecranon, when the arm is extended. The os humeri has evidently the most free and extensive motion of any bone in the human-body.

Q. Which is the ulna?

A. The ulna is one of the bones of the fore arm or cubit; reaching from the elbow to the wrist; it is big at its upper extremity, and grows gradually smaller towards its lower end. This bone is longer than the radius, and has a motion of flexion and extension: at its upper extremity it articulates with the os humeri and the crista of the radius, and its lower extremity articulates with the carpus, as also with the radius by means of a crista. At the upper end it has one large process or apophysis called olecranon, and a small process

on the fore part, and on one side between there is also a small cavity, which receives the upper end of the radius for its rotatory motion; and down the side of this bone, next the radius, is a sharpe edge, from which the ligament arises which connects these bones together. At the lower end is a process called styliformis, and a round head, which is received into the radius for the rotatory motion of the cubit.

Q. Which is the radius?

A. The radius is the exterior bone of the cubitus or fore arm, accompanying the ulna from the elbow to the wrist. In its upper extremity there is a glenoide cavity for its articulation with the humerus, which chiefly fits it for its rotatory motion, being also received into the ulna; for the strength of the elbow-joint receives but little advantage from theunion of these two bones. A little belowthis head is a large tubercle, into which the biceps muscle is inserted, which by the advantage of this infertion turns the cubit supine, as well as bends it. At the lower end, which is thicker, is a focket to receive the carpus, and at the fide next the ulna a small one to receive that bone, and a thin edge, into which the transverse ligament, which arises from the ulna is inserted: this ligament ties these bones conveniently and firmly together; for the ulna being chiefly articulated to the os humeri, and the radius to the carpus, a weight at the hand, bas monapole bell without

without this ligament, would be liable to pull these bones asunder.

Q What is the carpus?

A. The carpus or wrift, confifts of eight small, unequal, and irregular bones, divided into two rows, each row confifts of four bones; in the first, are the os scaphoides or naviculare, lunare, cuneiforme, and orbiculare; in the second are the os trapezium, trapezoides, magnum and unciforme. They have all obfcure motions one with another, and with those of the fecond is more confiderable, and they are moved by the same muscles which move the carpus on the radius.

Q. What is the metacarpus?

A. The metacarpus is that part of the hand between the wrift and the fingers; the inner part is called the palm, and the other the back of the hand. The metacarpus confilts of four bones, which answer to the four fingers, whereof that which fultains the fore finger is the biggest and longest, but has the least motion, and that of the little finger the most. They are all round and long, a little convex towards the back of the hand, and concave and plain towards the palm. They are hollow in the middle, and full of marrow; they touch one another only at their extremities, having spaces in the middle in which lie the musculi interossei. In the upper end of these bones there is a finus which receives the bones of the wrist; the other ends have round heads, which R 4

which are received into the finus of the first bones of the fingers.

Q. How are the fingers distinguished and

composed?

A. The fingers are four in number on each hand, exclusive of the thumb, viz. 1. index, the fore-finger; 2. medius, the middle finger; 3. annularis, the ring-finger; 4. minimus vel auricularis, the little finger. In the thumb and fingers are three bones, which make three phalanges, the upper of which (next the metacarpus) are much larger than the lower, next the extremities. These bones, on the infide, are flat and a little hollow or concave, which is necessary to make room for the flexors of the fingers, and to render their shape proper for grasping any thing. The thumb is shorter and stronger than any of the fingers, because in its actions it is to resist them all. The first phalanx, in the part where they are articulated with the bones of the metacarpus at their heads, have a glenoide cavity, by means of which articulation, they have a free motion every way. In the other extremity there are two heads with two cavities joined to the fecond phalanx, where the motions of flexion and extension are all that are possible; and the same is the case between the second phal nx and the third.

Q. What are the cartilages of the arm?

A. The cartilages of the arm, are those of the bones of the shoulder, os humeri, fore-arm, and hand. Q. What are the cartilages of the bones

of the shoulder?

A. The glenoide cavity of the scapula, which receives the superior head of the os humeri, is covered with a cartilage, which is a little raised above the edge of the bone. It is of a pliable slippery substance, yet something different from that of a cartilage, refembling in some measure the border of the acetabulum coxendix. The acromion has also a small cartilaginous surface; and the external extremity of the clavicle is crusted over with a cartilage, which is a little convex, and covers its whole triangular furface; besides which it has another moveable common cartilage already spoken of. The small cartilaginous surface of the humeral extremity of the clavicle, answering to that of the acromion, is, like that of the acromion, a little convex. Between these two cartilages of the clavicle and acromion, there is in some subjects a thin inter-articular cartilage, very smooth on both sides.

O. What are the cartilages of the os hu-

meri?

A. The cartilage by which the hemisphere of the head of the os humeri is covered, is gradually thicker towards the middle than toward the edges. The groove in the os humeri is covered by a crust, which is partly ligamentary and partly tendinous rather than cartilaginous. The lower head of the os humeri is covered by a common cartilage, and its foffulæ

fulæ near the head with a thin cartilaginous or ligamentary varnish.

Q. What are the cartilages of the fore-

arm?

A. The two figmoide cavities in the upper extremity of the ulna are covered by a cartilage common to both. The lower head of the ulna is crusted over by a cartilage; the head of the radius is covered in the same manner. All the concave fide of the basis of the radius is cartilaginous, and often divided by a finall cartilaginous prominent line. At the basis of the radius there is likewife a particular additional cartilage, or triangular production longer than it is broad, very thin, and rather flat than concave on both its smooth sides. This cartilage may be termed the inter-articular cartilage of the joint of the wrift. It is tied to the radius by very short ligaments, and sliding on the fmall head of the ulna.

Q. What are the cartilages of the hand?

A. All the bones of the carpus, metacarpus, and fingers, are crusted over with cartilages at their ends for their mutual articulations, which may be termed cartilaginous surfaces, in which the impressions and notches where the mucilaginous glands are lodged are very perceptible.

Q. What are the ligaments of the arm?

A. The ligaments of the arm are those of the bones of the shoulder, os humeri, forearm, and hand.

Q. What

Q. What are the ligaments of the shoulder?

A. The articulation of the acromion with the extremity of the clavicle, is strengthened quite round by feveral fmall strong ligaments, which go from one bone to the other. These ligaments lie very near each other, and are withal fo tightly braced over the joint as to hide it altogether, and they appear more like a cartilaginous covering than a ligamentary texture. The internal furface of these ligaments is lined with the capfula of the joint. When the fmall inter-articular cartilage is found, its whole circumference is connected to these ligaments. The articulation of the clavicle with the sternum is sustained by several ligaments fixed by one end round the pectoral extremity of the clavicle, and by the other to the sternum, as before observed. There is a long, narrow, strong ligament, which goes from one clavicle to the other, behind the furca of the sternum, and may be called the inter-clavicular ligament. The neck of the scapula, at a small distance from the edge of the glenoide cavity, gives infertion to the capfular ligament or mucilaginous bag, and to the articular ligaments of the joint of the scapula and os humeri. Besides the articular ligaments of the scapula, there are three ligamentary cords fixed to the tuberofity of the coracoide apophysis, two are inserted by the other end in the extremity of the clavicle, and the third under the acromion. A thin broad ligament alfo also reaches between the crista of the spine of the scapula and the edge of its inferior costa.

Q. What are the ligaments of the os hu-

meri?

A. The capfular or mucilaginous ligament loofely furrounds the whole articulation of the scapula, with the head of the os humeri; from its infertion round the edge of the glenoide cavity, already mentioned, it is continued over the hemisphere of the head of the os humeri. and fixed near its edges towards the muscular furfaces of the great and small tuberosities or processes. Afterwards it runs down gradually on the neck of the bone below the lowest part of the cartilaginous hemisphere, and is closely fixed in the bone, except in the groove or channel already mentioned, in which lies the tendon of the biceps cubiti. The true ligament of this joint seems to be made up of two forts of ligaments closely united together, viz. a capfular ligament which furrounds the whole articulation; and feveral true ligaments which run over, and closely adhere to the former at different distances. On the body of the os humeri there are two long, flat, thin, strong and narrow ligaments, fixed by one edge along the two lower thirds of the bone reaching to both condyles. The lower extremity of the os humeri is joined to the bones of the fore-arm, by two falciculi of ligamentary fibres, one fixed to the inner condyle, the other to the external. The capfular ligament is fixed to the condyles, and these cover them; and

and afterwards it is fixed round both sides of this lower extremity above the fossulæ, which fossulæ are only slightly varnished over with a cartilaginous substance.

Q. What are the ligaments of the fore-

arm?

A. Some of the ligaments of the bones of the fore-arm are common to them with the os humeri, some common to them with the bones of the hand, and some are proper. The two proper are the interoffeous ligament of the forearm, and one which may be termed the coronary ligament of the radius. To these may be added the annular ligaments, which only ferve for the passage of tendons; and other ligamentary expansions, which may be called muscular ligaments, and shall refer a description of them till I come to speak of the muscles. The interoffeous ligament of the forearm is very like that of the leg; it ties the ulna and radius closely together, and serves for the infertion of feveral muscles. In it are holes for the passage of the blood-vessels. The capfular ligament of the joint of the elbow runs down from its infertion in the os humeri (already mentioned) and is fixed in the olecranon, round the edge of the great figmoide cavity; it likewise runs over the head of the radius, and is fixed to the coronary ligament quite round. Thus it completely furrounds the articulation of these three bones, and serves to contain the mucilaginous liquor furn shed by the glands and fatty substance, both which

Q. What are the ligaments of the hand?

the external transverse ligament of the carpus; and may likewise be named the great oblique

A. The ligaments of the carpus are very numerous. Some of them tie each bone to one or two neighbouring bones in the same rank, but are so very short as to allow these bones only a small degree of motion. Some of them tie the bones of one row to those of the other, but they are not so short as the former, and therefore they allow these bones a more manifest motion, as we see in bending the wrist. Lastly, there are other ligaments

of the carpus, by which the three first bones of the first row are connected to the bones of the fore-arm; and to these may be added the ligaments by which the bones of the fecond row are joined to those of the metacarpus and first phalanx of the thumb. The principal ligaments belonging to the articulation of the carpus with the bones of the fore-arm, I have already mentioned, except their infertion in the carpus, which is thus: the styloide ligament of the radius is fixed to the os scaphoides vel naviculare; and the styloide ligament of the ulna is fixed first in the os cuneiforme, and then in the os unciforme, from whence it is a little stretched over the fourth bone of the metacarpus. The ligaments which lie between the two former, go from the radius and ulna to the three first carpal bones, and the mucilaginous capfula, by which these ligaments are lined are also fixed to these bones. There is likewise a considerable ligament, called the inner transverse ligament, formerly called an annular ligament.

The bones of the metacarpus, besides the short ligaments by which they are tied to the second row of the bones of the carpus, have several others, by which both their bases and heads are connected together. The bases of the third and sourth bones are not so closely tied as the rest, and therefore they have a very sensible motion, which, however, is greater in the sourth than in the third. The heads of these bones are firmly tied to each other by

a strong

a strong transverse ligament, situated in the palm of the hand, through which the tendons of the flexor muscles of the fingers have a free passage. The first phalanx of the thumb and fingers is fixed by short ligaments that pass obliquely over the articulation with the metacarpal bones. The fecond phalanx of the thumb is joined to the first by ligaments of the same kind. The other phalanges are joined by lateral ligaments almost in the same manner as the bones of the fore-arm to the os humeri. The two first phalanges of each finger have very strong ligamentary vaginæ inferted in the rough ridges on their flat sides. These vaginæ are lined with a mucilaginous membrane, which runs like a tube from one phalanx to the other, over the articulation.

Q. What are the muscles of the arm?

A. The muscles of the arm and its parts include those of the clavicle, scapula, and breast, which belong thereto, tho' situated on the superior part of the trunk; for this reason I omitted their description in describing those of the thorax. They are as follow, viz. fubclavius, trapezius, rhomboides, elevator scapulæ, serratus major et minor anticus, pectoralis, deltoida, supraspinatus, infraspinatus, teres minor et major, latissimus dorsi, subscapularis, coraco-brachialis, biceps cubiti flexor, brachiæus internus, supinator radii longus et brevis; triceps extenfor cubiti, anconæus, palmaris longus, et brevis; flexor carpi radialiset ulnaris, extensores carpi radiales, extensorulnaris,

olnaris, pronator quadratus, et teres, perforatus, perforans, lumbricales, extensor digitorum communis, minimi digiti, extensor indicis, abductor primi digiti interossai, minimi digiti ossis, metacarpi minimi digiti, extensor primi secundi et tertii internodii pollicis, slexor primi et secundi ossis pollicis, slexor tertii internodii pollicis, abductor pollicis.

Q. Which is the musculus subclavius?

A. The musculus subclavius is a small oblong muscle lying between the clavicle and first rib; it arises from the superior part of the first rib, and is inserted into almost all the middle portion of the under side of the clavicula. Its use is to draw the clavicula toward the sternum, that it may not be severed in the motions of the scapula.

Q. Which is the trapezius?

A. The trapezius, vel cucullaris, is a muscle of the scapula, which arises from the os occipitis, and from a linea alba vel ligamentum colli, from the spinal process of the last vertebra of the neck, and the ten uppermost of the back, and from a linea alba between all these processes; it is inserted into one third of the clavicle next the scapula, almost all the back part of the spine of the scapula, and as much of the processus acromion as lies between the spine of the scapula and the clavicle. This muscle draws the scapula directly backward.

Q. Which is the rhomboides?

A. The rhomboides is a thin, broad, and obliquely square, sleshy muscle, situated between

tween the basis of the scapula and the spinal dorsi. It ariseth from the spinal process of the two inferior vertebræ of the neck, and the four superior of the back; and is inserted into the whole basis of the scapula, which it draws up and backward.

Q. Which is the elevator fcapulæ?

A. The elevator scapulæ is a long and pretty thick muscle, about two singers breadth, it arises from the transverse apophyses of the four first vertebræ of the neck, and running a little obliquely, is inserted into the upper angle of the scapula for its elevation.

Q. Which is the ferratus major anticus?

A. The ferratus major anticus is a broad, fleshy, and pretty thick muscle, situated on the lateral part of the thorax, between the ribs and the scapula, by which it is covered. It arises by dentated origins (resembling a saw, from whence the name of serratus) from the anterior part of the eight superior ribs, and is inserted into the basis of the scapula, which it draws forward and upward. This muscle is an elevator of the ribs.

Q. Which is the ferratus minor anticus?

A. The ferratus minor anticus, vel pectoralis minor, arifes from the third, fourth, and fifth ribs, continues its course under the pectoralis major, and is inserted into the processus coracoides scapulæ, which it pulls forward and downward. This is also an elevator of the ribs. All the muscles inserted into the basis of the scapula are also inserted into one another.

Q. What is the musculus pectoralis?

A. The pectoralis, (by some called pectoralis major) possesses almost the whole breast, being a large, thick, sleshy muscle, covering the fore part of the breast, from the sternum, where it is very broad, to the axilla, where it contracts in its passage to the arm. It arises from near two thirds of the clavicula, next the sternum, and all the length of the os pectoris, and from the cartilages of the ribs; and is inserted into the os humeri about four singers breadth below its head, between the biceps and the insertion of the deltoides. The use of it is to draw the arm forward.

A small portion of the lower part of this muscle is often confounded with the obliques descendens abdominis; in some bodies neither the upper part nor its tendon can be easily separated from the deltoides; and in others, even that part of it that arises from the clavicula is

a distinct portion.

Q. Which is the deltoides?

A. The deltoides is a very thick, triangular muscle, covering the upper part of the arm, and forming what is called the stump of the shoulder. It is one of the elevators of the arm. It arises from the clavicula, acromion, and spine of the scapula, and is inserted near the middle of the os humeri, which bone it lifts directly upward.

O. Which

Q. Which is the fupra spinatus?

A. The supra spinatus, vel superscapularis superior, arises from the dorsum scapulæ, above the spine, and passes between the acromion and neck of the scapula, and under the ligament between the acromion and processus coracoides. It is inserted into the upper part of the os humeri, near its groove, or bony channel. It helps to lift the arm upwards, until it becomes parallel with the spina scapulæ.

The supra spinatus deltoides and coracobrachialis, assist in all the motions of the humerus, except depression; it being necessary that the arm should be raised and sustained, in

order to move it to any fide.

Q. Which is the infra spinatus?

A. The infra spinatus, vel superscapularis inferior, arises from the dorsum scapulæ below the spine, and is inserted, wrapping over part of it, at the side of the head of the os humeri; it turns the arm supine and backward.

Q. Which is the teres minor?

A. The teres minor is an abductor muscle, as well as the infra spinatus, below which it has its origin from the inferior costa of the scapula, and is inserted together with the infra spinatus, which it assists in turning the arm supine, but pulls it more downwards.

Q. Which is the teres major?

A. The teres major is a depressor muscle, which has its origin at the lower angle of the scapula, and is inserted at the under part of the os humeri about three singers breadth from the head. This muscle draws the os humeri to-

ward

ward the lower angle of the scapula, and turns the arm prone and backward.

Q. Which is the latissimus dorsi?

A. The latissimus dorsi is a very broad muscle, covering almost the whole back; it has a thin broad tendinous beginning, which arises from the spinal processes of the ilium, os sacrum, vertebræ of the loins, and from the seven inferior vertebræ of the thorax. inserted into the os humeri with the teres major, by a strong and broad tendon, with which it pulls the arm downward and backward.

Q. Which is the subscapularis?

A. The subscapularis is a muscle of the fame breadth and length with the scapula, filling up the hollow or under fide of it (whence its name) from which it arises, and is inserted into the neck of the os humeri, wrapping somewhat over it. This muscle pulls the arm to the fide and prone.

Q. Which is the coraco-brachialis?

A. The coraco-brachialis arises from the processus coracoides scapulæ, and passing over the articulation, is inferted into the middle and internal part of the humerus; this, with the deltoides and supraspinatus, lifts the arm upwards, and turns it somewhat outward.

Q. Which is the biceps cubiti flexor?

A. The biceps cubiti flexor arises with two heads, that the fibres of this muscle should not compress one another. One head arises with a long, round tendon, from the upper edge of the acetabulum scapulæ, running under the ligament ligament of the articulation, in a channel or groove on the head of the shoulder-bone, (before-mentioned,) wherein it is inclosed by a proper ligament, afterwards it becomes fleshy, and joins the other head, which arises from the processus coracoides scapulæ, in common with the coraco-brachialis muscle, and is inserted with the first head into the tubercle, at the upper head of the radius. Sometimes this muscle has a third head, which arises from the middle of the os humeri. This muscle lifts up the humerus, bends the cubit, and has as great a share as any one muscle in turning the cubit supine; the humerus being fixed by other muscles, the whole force of this muscle will be exerted upon the cubit, or the cubit being fixed by an extensor, the whole force of it will be spent in raising the arm, and therefore ought to be always reckoned among those that raise a weight at arm's length.

The rendinous expansion of this muscle, in the slexure of the cubits, is sometimes pricked in bleeding (by injudicious operators,) which generally occasions a most violent pain and inflammation, and has been attended with fatal consequences. The great mischief arising from pricking this tendinous fascia, seems owing to its lying so much upon the stretch, which may be wholly avoided by bending the elbow and

turning the cubit.

Q. Which is the brachiæus internus?

A. The brachiæus internus lies partly under the biceps, and arises from the internal part and and below the middle of the os humeri, near the infertion of the deltoides, and coraco-brachialis muscles; it is inserted in the tubercle of the ulna, a little below its upper head. This muscle helps also to bend the arm.

Q. Which is the fupinator radii longus?

A. The supinator radii longus has its origin from the exterior part of the humerus, and is inserted into the upper side of the radius near the carpus. This muscle, some say, is not a supinator, but a bender of the cubit.

Q. Which is the supinator radii brevis?

A. The supinator radii brevis, arises from the lower part of the external condyle of the os humeri, and from the upper part of the ulna, and is inserted near the tubercle of the radius. This muscle serves also to turn the palm of the hand upward, and to bend the cubit.

Q. Which is the triceps extensor cubiti vel

brachiæus externus?

A. The triceps extensor cubiti is divided into three heads. The first arises from the inferior costa of the scapula near the acetabulum; the second, from the outer and back part of the os humeri; the third, lower and more internal; and are inserted into the processus olecranon of the ulna. The first of these heads draws the arm backward.

Q. Which is the anconæus?

A. The anconæus arises from the back part of the extremities of the humerus, passes over the elbow, and is inserted into the upper part

of the ulna. This muscle is also an extensor of the cubit.

Q. Which is the palmaris longus?

A. The palmaris longus arises from the inner condyle of the os humeri; it soon after becomes a tendon, joins the ligamentum transversale carpi, and is expanded in the palm of the hand. This muscle is often wanting, but the expansion in the hand never; yet being connected to the ligament of the carpus, it must bend the carpus, and cannot constrict the palm of the hand.

Q. Which is the palmaris brevis?

A. The palmaris brevis, or caro quadrata, is in form of a small mass of slesh, very different in size in different bodies. It arises obscurely from the ligamentum transversale carpi, and seems to be inserted into the eighth bone of the carpus, and the metacarpal bone of the little singer. This helps to constrict the palm of the hand.

Q. Which is the flexor carpi radialis?

A. The flexor carpi radialis arises from the inner extuberance of the os humeri, and soon becoming a strong tendon, passes through a channel of the fifth bone of the carpus, and is inserted into the metacarpal bone of the fore finger. This not only bends the carpus upon the radius, but also the bones of the second order upon those of the first.

Q. Which is the flexor carpi ulnaris?

A. The flexor carpi ulnaris arises from the same tubercle of the humerus as the former,

and a fascia betwixt this muscle and the tensor ulnaris, contiguous to the ulna, and is inserted by a short tendon into the fourth bone of the carpus.

Q. Which are the extensores carpi radiales?

A. The first of the extensores carpi radiales, arises from the os humeri immediately below the supinator radii longus, and is inserted into the metacarpal bone of the middle singer. The first of these muscles is a bender of the cubit, as well as an extensor of the carpus.

Q. Which is the extensor ulnaris?

A. The extensor ulnaris arises from the same extuberance with the sormer, and half the ulna below the anconæus muscle; then, becoming a tendon, it runs in a small sinus at the bottom of the ulna, and is inserted into the metacarpal bone of the little singer. The extensors of the carpus being inserted into the metacarpus, at once perform the motion between the bones of the carpus, and that between the carpus and radius. The slexor and tensor ulnaris acting together turn the hand downward; the tensor and flexor radialis upward.

Q. Which is the pronator quadratus?

A. The pronator quadratus vel transversus, lies transversely on the inside of the lower extremity of the fore-arm. It arises from the lower part of the ulna near the carpus, and passing under the slexors of the singers, is inferted into the lower extremity of the radius.

Q. Which is the pronator teres?

A. The pronator teres, vel obliquus, is signated in the upper part of the ulna, opposite to the supinator brevis. It arises from the internal condyle of the os humeri, and upper and fore part of the ulna, and is inserted into the radius below the supinator brevis.

Q. Which is the perforatus, &c?

A. The perforatus, or flexor fecundi internodii digitorum, called also sublimis, arises
from the inner tubercle of the os humeri, and
from the upper part of the ulna, and the middle of the radius; then becoming four strong
tendons, it passes under the ligamentum transversale carpi, and is inserted into the beginning of the second bone of each singer.

Q. Which is the perforans?

A. The perforans, or flexor tertii internodii digitorum, arifes from half the ulna, and a great part of the ligament between the ulna and radius, then becoming four tendons, paffes under the ligamentum transversale carpi, and through the tendons of the former muscle to their infertion into the third bone of each finger. The tendons of both these muscles are tied down to the fingers by a strong ligament.

Q. Which are the lumbricales?

A. The lumbricales, or flexores primi internodii digitorum, arife from the tendons of the last mentioned muscle, and are inserted laterally toward the thumb into the beginning of the first bone of each finger.

Q. Which

Q. Which is the extensor digitorum communis !

A. The extensor digitorum communis, arifes from the external protuberance of the humerus, and at the wrift it passes under a ligament, and divides into four tendons, which are afterwards inferted into the beginning of the fecond bone of each finger. These tendons communicate upon the first joint, which keeps them from fliding off the joints of the fingers, being a little connected to the first bones.

Q. Which is the extensor minimi digiti?

A. The extensor minimi digiti vel auricularis, is a portion of the last muscle, passing under the ligament in a distinct channel.

Q. Which is the extensor indicis?

A. The extensor indicis comes from the middle and external part of the ulna, and paffing under the ligaments of the carpus, is inferted with the extensor communis into the fore finger. This muscle extends the forefinger fingly.

Q. Which are the abductor primi digiti in-

terossei, and abductor minimi digiti?

A. The abductor primi digiti interoffei, and abductor minimi digiti, are eight mufcles, one for each fide of each finger. The abductor primi digiti arises from the first bone of the thumb, and the fide of the metacarpal bone of the fore-finger. The interoffei are three pair, fitly divided into external and internal; the external arise from the metacarpal bones, whose spaces they fill up next the back

back of the hand; the internal arise from the fame bones in the infide of the hand.

Q. Which are the abductor minimi digiti? A. The abductor minimi digiti arises from the transverse ligament, and fourth bone of the carpus; these muscles are inserted, two into the first joint of each finger, and then passing obliquely over the tops of the fingers, are inferted into their last bones; they bend the first joint, and extend the two last, as in holding a pin, and in playing upon some mufical instrument. The abductors of the fore and little fingers, with the fecond and fifth interoffei mufcles acting, the fingers are divaricated, and the other four acting bring them together; these muscles which divaricate the fingers, being extenders of the fecond and third joints, we never can divaricate them without extending them a little.

Q. Which is the abductor offis metacarpi

minimi digiti?

A. The abductor offis metacarpi minimi digitiarises from the eighth bone and transverse ligament of the carpus, and is inferted into the metacarpal bone of the little finger, which it pulls toward the thumb, to constrict the palm of the hand.

Q. Which is the extensor primi internodii

pollicis? I fresh bodies as heve them. S sipilities

A. The extensor primi internodii pollicis arises, from the ulna, below the anconæus muscle and the ligament between the ulna and radius; then becoming two, three, or four tendons. tendons, is inserted into the fifth bone of the carpus, and first of the thumb. The first of these insertions can only assist the bending of the wrist upward, and in turning the arm supine.

Q. Which is the extensor secundi, &c?

A. The extensor secundis internodii pollic is arises immediately below the former, from the radius and transverse ligament, and is inserted into the second and third bone of the thumb.

Q. Which is the extensor tertii internodii

pollicis?

A. The extensor tertii internodii policis arises immediately below the last described, from the ulna and ligament, and passes over the radius nearer the ulna, to be inserted into the third bone of the thumb. This extends the thumb more towards the ulna than the former muscle, and is very much a supinator.

Q. Which is the flexor primi et secundi

offis pollicis?

A. The flexor primi et secundi ossis pollicis arises from the sitth bone and transverse ligament of the carpus, and from the beginnings of the two sirst metacarpal bones, and is inserted into the whole length of the sirst bone of the thumb, and tendinous into the beginning of the second; the sesamoid bones of the thumb in such bodies as have them, lie in this tendon, where it passes over the joint.

Q. Which is the flexor tertii internodii

pollicis?

A. The flexor tertii internodii pollicis arifes large from almost all the upper part of the radius, and becoming a round tendon, passes under the ligamentum transversale carpi to be inferted into the third bone of the thumb. This muscle singly acting draws the thumb towards the metacarpal bone of the little finger; but the last mentioned muscle acting with it, turns it toward the fore finger.

Q. Which is the adductor pollicis?

A. The adductor pollicis arises from the carpus, and almost the whole length of the metacarpal bone of the middle finger, and is inferted into the beginning of the fecond bone of the thumb. This muscle naturally enough divides into two, and might better be called a flexor than an adductor.

Q. Which is the abductor pollicis?

A. The abductor pollicis arises from the fifth bone and ligamentum transversale of the carpus, and is inferted laterally into the beginning of the second bone of the thumb to draw it towards the radius.

The muscles which bend the thumb are much less than those which bend the fingers, nevertheless, the thumb is able to refist all the fingers merely from the advantages that arife from the thickness and shortness of the bones of the thumb, compared with those of the fingers; but then the quickness of motion in the fingers will exceed that of the thumb, as much as the fingers exceed the thumb in length, and their muscles those of the thumb in largeness. Q. What

Q. What are the arteries of the arm?

A. The arteries of the arm proceed all from the subclavian: the various branches it gives off before it leaves the thorax have been already mentioned. From the arteria thoracica a branch in particular runs down between the deltoides and pectoralis, together with the vena cephalica, to which it adheres very closely, as if there were an anastomosis between them. Another branch fometimes runs between the musculus brachiæus, and anconæus internus, which communicates with a branch of the radial artery. The fuclavian artery having left the thorax immediately above the first rib, in the interstice left between the portions of the scalenus muscle, gives off the arteries above mentioned, and the external fcapulary, and then receives the name of axillaris, (because it passes under the axilla) from whence proceed the arteria humeralis, brachialis, cubitalis, and radialis.

Q. Which are the arteriæ scapulares?

A. The external scapulary artery passes thro' the notch in the superior costs of the scapula, to the musculus supraspinatus and infraspinatus, teres major et minor, and to the articulation of the scapula with the os humeri. The internal scapularis arises from the axillary artery near the axilla, and runs backward, to be distributed to the subscapularis, giving branches and ramifications to the serratus major, axillary glands, teres major, infraspinatus, and upper portion of the anconei.

Q. Which is the arteria humeralis?

A. The arteria humeralis arises from the lower and fore side of the axillaris, and runs backward between the head of the os humeri and teres major, furrounding the articulation till it reaches the posterior part of the deltoides, to which it is distributed; in its course it supplies the fuperior portions of the anconei, the capfular ligament of the joint of the shoulder, and the os humeri itself, through several holes mmediately below the great tuberofity of the iead of that bone, communicating with the scapular artery. Opposite to the origin of the humeralis the axillaris fends off a small branch in a contrary direction between the head of the os humeri and the upper part of the biceps and coraco-brachialis, which goes to the vagina and channel of the biceps, and to the periofteum, and afterwards joins the humeralis.

Q. Which is the arteria brachialis?

A. The axillary artery having given off the humeralis and its branches, as I have mentioned, passes immediately behind, the tendon of the pectoralis, where it changes its name to brachialis, which runs down on the infide of the arm over the musculus coraco-brachialis, and triceps extensor cubiti, and along the inner edge of the biceps behind the vena basilica, giving small branches on both sides to the neighbouring muscles, to the periosteum, and to the bone. Between the axilla and middle of the arm, it is covered only by skin and fat; but afterwards it is hid under the biceps, and runs

other

obliquely forward as it descends; but it does not reach the middle of the fold of the arm. In its paffage to this place it fends off many branches to the infraspinatus, teres major and minor, subscapularis, latissimus dorsi, serratus major, and other neighbouring muscles, to the common integuments, and even to the nerves. Below the fold of the arm it divides into two principal branches, which I shall mention by and by, one called cubitalis, the other radialis. From the upper and inner part the brachialis it fends off a particular branch, which runs obliquely downward and backward over the anconæus, and then turns forward again near the external condyle, where it communicates with a branch of the radialis.

Immediately below the infertion of the teres major it gives off another branch, which runs round the os humeri, and descends obliquely forward between the musculus brachizeus and triceps extensor cubiti, to both which it is diftributed in its passage. About the breadth of a finger below the fecond branch, the brachial artery fends off a third branch toward the internal condyle as the second is to the external condyle, both which communicate with the arteries of the cubit. Near the middle of the arm the brachialis fends off a branch between the brachieus and the triceps to the periosteum which penetrates the bone. About an inch lower it gives off a branch, which fends ramifications to the triceps, and runs over the inner condyle. Below the middle of the arm another branch runs behind the inner condyle in company with a confiderable nerve. A little lower another branch is sometimes detached on the foreside of the inner condyle; all these three branches communicate with branches of the cubital artery, and are termed collateral arteries. I he common trunk of the brachial artery having reached the fold of the arm, runs together with a vein and a nerve, immediately under the aponeurosis of the biceps, and passes under the vena mediana, detaching branches on each side to the neighbouring muscles.

About an inch beyond the fold of the arm, the brachial artery divides into two principal branches. The inner or posterior named cubitalis; and the outer or anterior named radialis, as has been already said. From this bisurcation the brachial artery sends branches on each side to the supinator and pronator muscles, teres, fat

and fkin.

Q. How is the arteria cubitalis distributed? A. The cubital artery, proceeding from the brachialis, as I have before observed, sinks in between the ulna and the upper parts of the pronator teres, perforatus, palmaris longus, and slexor carpi radialis: then leaving the bone, it runs down between the perforatus and palmaris longus, all the way to the carpus and great transverse ligament. In this course it winds and turns several ways, and sends out several branches; very often there is a branch of communication between the brachial and cubital arteries. This communicant branch is some

fometimes very large, and liable to be pricked by careless or injudicious blood-letters in bleeding in the basilic vein, immediately under which this branch generally lies. When the operation for an aneurism is made upon the communicant branch, it must be tied on both sides of the orifice, because the blood is liable

to flow freely into it either way.

The first branch of the cubital is a small artery which runs to the inner condyle, then turns up, and communicates with the collateral arteries before mentioned. Another small branch almost surrounds the articulation. The cubital artery running then between the heads of the ulna and radius, and having reached the interoffeous ligament, fends off two principal branches which may be called interoffeous arteries of the fore arm, one internal, the other external. The external pierces the ligament about three fingers breadth below the articulation, from which runs up a branch to the external condyle of the os humeri under the extensor ulnaris and anconæus, to which it is distributed, as also to the supinator brevis, communicating with the collateral arteries on the same fide Afterwards this external interoffeous runs down on the outfide of the figament, and is distributed to the extensor ulnaris, extensor digitorum communis, and to the extensores pollicis indicis, and minimi digitis; communicating with branches of the internal interoffeous. At the lower extremity of the ulna it unites with a branch of the internal interoiteroffeous, and is distributed, together with it, on the back of the hand, communicating with the radialis and a branch of the cubitalis.

By these communications this artery forms a fort of irregular arch, from whence branches are detached to the external interoffeous muscles, and to the external lateral parts of the The internal interoffeous artery runs down very close to the ligament, which it perforates between the pronator teres, and pronator quadratus, and goes to the back of the hand, where it communicates with the external interoffeous radialis, and the internal branch of the cubitalis. From the origin of the two arteriæ interossæ the cubitalis runs down between the perforatus, perforans, and flexor carpi ulnaris, along the ulna, fending branches to the neighbouring parts. Afterwards the cubital artery passes over the internal transverse ligament of the carpus by the side of the os pisciforme vel orbiculare, and having furnished the skin, palmaris brevis, and metacarpus, it flips under the aponeurofis palmaris, giving off a branch to the abductor minimi digiti, and another towards the thumb between the tendons of the flexors of the fingers and the bases of the metacarpal bones. A branch runs also between the third and fourth bones of the metacarpus to the back of the hand, where it communicates with the external interoffeous artery. Afterwards, having supplied the interoffeous muscles, it communicates with the radialis, and they both form an arterial arch

in the hollow of the hand in the following manner: about two fingers breadth beyond the internal annular ligament of the carpus, the cubitalis forms an arch, the convex side of which is turned to the fingers, and commonly fends off three or four branches; the first, to the inner and back part of the little finger; and the other three run in the interffices of the four metacarpal bones, near the heads of which, each of them is divided into two branches, which run on each fide the fingers internally, and at the ends of the fingers these digital arteries communicate and unite with each other. From the concave fide of this arch towards the fecond phalanx of the thumb, a branch goes to the internal lateral part thereof, and then ends near the head of the first metacarpal bone, by a communication with the radialis; having first given a branch to the forefide of the index, and another to the fide of the thumb; these communicate as the other digital arteries. This arch fends likewise small twigs to the interosseous muscles, lumbricales, palmaris, and other neighbouring parts; and lastly to the integuments.

Q. Which is the arteria radialis?

A. The radial artery proceeds from the brachialis, as before observed, and begins by detaching a small branch upward toward the fold of the arm, which also turns backward round the external condyle, communicating with the neighbouring branches of the brachial artery. The radial artery runs down on the inside of the

the radius, between the supinator longus, pronator teres, and the integuments, giving branches to these muscles, and likewise to the perforatus, perforans, and supinator brevis. From thence it runs in a winding course toward the extremity of the radius, supplying the flexor of the thumb and pronator quadratus. Having reached the extremity of the radius, it runs nearer the skin, especially toward the anterior edge of the bone, being the artery which we there feel when we examine the pulse. At the end of the radius it gives off a branch to the abductor pollicis, which detaches one to the whole internal fide of the thumb; afterwards. it runs between the first phalanx and tendons of the thumb, to the interstice between it and the first metacarpal bone, where it turns toward the hollow of the hand, and fends off a branch to the external fide of the thumb, Other branches of the radialis run transversely outward between the first two bones of the metacarpus and the two tendons of the extenfores carpi radiales; they communicate with the cubitalis, and together furnish the external interoffeous muscles and integuments of the back of the hand. Lastly, the radialis terminates in its passage over the semi interosseous muscle of the index, near the basis of the first metacarpal bone, and runs under the tendons of the flexor muscles of the fingers, where it is joined to the arch of the cubitalis. It fends off another branch along the fore part of the first bone of the metacarpus to the back of the index, where

it is lost in the integuments; it gives also a branch to the internal fide of the index, and at the end of the finger joins an opposite branch which comes from the arch of the cubitalis. Another small branch crosses the internal interoffeous muscles, and communicates with the great arch. Where the arch of the cubitalis ends at the middle finger, the radialis runs along the inner part of the first metacarpal bone, at the head of which it terminates by two branches; one runs along the inner fide of the index, and the other passes between the flexor tendons of this finger and the metacarpal bone, and having communicated with the cubital branch of the middle finger, it advances on the posterior lateral part of the index to the end of the finger, where it unites again with the first branch.

Q. What are the veins of the arm?

A. The veins of the arm, &c. likewise proceed from the subclavian, and the various branches are distinguished by the names of the parts they are bestowed on, in the same manner as the arteries already mentioned, viz. vena axillaris, cephalica, basilica.

Q. Which is the vena axillaris?

A. The subclavian vein, having sent off the branches already described, goes out of the thorax, and passes before the anterior portion of the musculus scalenus, and between the first rib and the clavicle to the axilla. Through this course it takes the name of vena axillaris, and gives off several branches, the chief

of which are the venæ musculares, thoracicæ and vena cephalica, which is sometimes double. The first veins which it sends off are the musculares, distributed to the middle portion of the musculus trapezius, to the angularis, infraspinatus and subscapularis: a little before the axillaris reaches the axilla, it sends out the venæ thoracicæ, one superior called also mammaria externa, and the other inferior. It likewise sends branches to the musculus subscapularis, teres major, teres minor, supraspinatus, latissimus dorsi, serratus major, pectoralis, and to the glands of the axilla.

Q. Which is the vena cephalica?

A. The axillaris, having reached the fide of the head of the os humeri, produces a very considerable branch named vena cephalica, and afterwards runs along the arm by the name of vena bafilica; however, the bafilica fometimes appears to be rather a branch than a continuation of the axillary trunk; in which case the cephalica and basilica might be looked upon as two principal branches of the axillaris. The cephalic vein, which is a branch of the axillaris, at a small distance from its origin, joins the small cephalica from the fubclavia or jugularis externa; having till then run near the furface of the body between the deltoides and pectoralis. The great cephalica runs down between the tendons of the last mentioned muscles and along the external edge of the biceps, to the neighbouring muscles, fat and skin communicating with the

the bafilica: a little below the external condyle of the os humeri, it fends off a branch backward. Near the fold of the arm it divides into two principal branches, one long, the other short. The long branch is named radialis externa, and the short one may be called mediana cephalica, to distinguish it from another mediana, which is a short branch of the basilica, and therefore ought to be called vena mediana basilica. The external radial vein runs along the radius between the muscles and integuments, and gives off branches to both fides, communicating and forming areolæ much like the faphena.

The mediana cephalica runs down toward the middle of the fold of the arm, under the integuments, and over the tendon of the biceps, where it joins the mediana basilica before mentioned. These two medianæ unite in an angle, the apex of which is turned downward. From this angle a confiderable branch goes down the fore-arm, uniting on one fide with the vena cephalica, and communicating on the other with the basilica, by

feveral irregular areolæ.

The name of mediana is given to this large branch as well as to the two short ones, by the union of which it is formed; this large branch is the true mediana, and that name only fufficiently distinguishes it, as the others are also termed cephalica and basilica. From this union of the two lateral medianæ, a branch goes down the infide of the fore-arm called vena wena cubiti profunda. The mediana cephalica fometimes fends down a long branch,
cailed radialis interna, almost parallel to the
externa; afterwards, having reached the extremity of the radius, it is distributed by a
numerous areolæ almost in the same course
with the radial artery; it gives off a branch
which runs superficially between the thumb
and metacarpus by the name of cephalica pollicis. The areolæ furnish the interosseous muscles and integuments, and communicate with
a small branch from the basilica called by the
ancients salvatella.

O. Which is the vena bafilica?

A. The basilic vein proceeds from the subclavian, as before mentioned, and has fometimes a double origin by a branch of communication with the trunk of the axillaris. First of all it sends off, under the head of the os humeri, a pretty large branch, which passes almost transversely round the neck of that bone, and running upon the scapula it is ramified in the deltoides; this branch may be named fub-humeralis, or articularis, as the artery; it fends down two principal branches, one on the infide of the bone, to which and the periofteum it gives fmall veins; the other turns forward, toward the middle of the arm between the bone and the biceps. Below the neck of the os humeri, near the hollow of the axilla, and behind the tendon of the pectoralis, the bafilica fends out a confiderable branch down the fide of the brachial artery, which

which supplies the neighbouring muscles on both sides. Immediately afterwards the basilica detaches two or three small veins, very closely joined to the brachial artery, and divides at the fold of the arm like the artery, having the same divisions along the whole fore arm afterwards, the basilica continues its course along the inside of the os humeri, between the muscles and integuments which it supplies. Having reached the inner condyle, and fent off obliquely in the fold of the arm, the mediana basilica, it runs along the ulna between the integuments and mufcles, a little toward the outfide by the name of cubitalis externa, and another branch, which the mediana basilica sends down the inside of the fore-arm near the ulna, may be named cubitalis interna. The bafilica, having reached the extremity of the ulna, fends feveral branches to the back-fide of the carpus, one of which named falvatella, goes to the fide of the little finger next the ring-finger, having first conmunicated with the cephalica, by means of the venal areolæ conspicuous on the back of the hand. In the other fingers the vein follows nearly the same course with the artery.

Q. What are the nerves of the arm?

A. The nerves of the arm, &c. come originally from the medulla fpinalis, and proceed immediately from the cervical nerves. The confiderable branches into which the nerves of the arm are divided are fix, which Mr. Monro names as follows.

arm, and serves the reguments as far as the

palm of the hand and fingers.

2. Musculo-cutaneus, passes through the musculus coraco brachialis, and after supplying the biceps and brachiæus internus, is spent on the teguments of the back of the cubitus and hand.

3. Muscularis, runs down the forepart of the arm to be lost in the musculi slexores car-

pi, digitorum, &c.

-UJ - 3

- 4. Ulnaris, which supplies the extensores cubiti, and teguments of the elbow, then passing through the sinuosity at the back of the external condyle of the humerus, runs along the ulna, where it gives twigs to the teguments, and neighbouring muscles; at length it is lost in the back of the hand, musculi interossei, and lumbricales, in the little singer, and side of the ring singer next to this. The course of this nerve is sufficiently felt when we lean on our elbow, by the sensibility and pricking pain in the parts to which it is distributed.
- 5. Radialis, goes down the forepart of the arm, near the radius, bestowing branches in its progress on the circumjacent muscles, and splitting at the ligamentum annulare carpi, it is sent to the thumb, fore-singer, middle-singer, and half of the ring-singer, and to the back of the hand.
- 6. Articularis, runs almost round the top of the os humeri, and ferves the musculi ex-

tensores cubiti, retractores, and elevatores humeri. By a strong and continued pressure on these nerves, by crutches, or any such hard fubstance, a palfy and atrophy of the arm may be occasioned.

Q. What are the glands of the arm?

A. The axillary glands are fituated under the arm-pits, enveloped in fat, and lie close by the axillary veffels about the scapula, and the flexure of the elbow; there are also found here and there some small glands; as also in fome places between and among the muscles. Subcutaneous glands, which fome mention, have no existence.

DIALOGUE VII.

Of the LEG, and its PARTS.

Q. WHAT are the parts of the leg?
A. Under the general denomination of the leg and its parts, I shall describe all the lower extremities, and divide them as in the table of ofteology and in the skeleton, into the thigh, leg, and foot. The crus extends from the nates or buttocks to the ends of the toes; and is divided into femur the thigh, from the ischium or hip to the knee. The fold between the belly and thigh is termed inguen, the groin; the top or forepart of the knee is termed patella, or rotula, the pan of the knee; the hinder part poples, the ham. From the knee to the instep, is properly called the tibia or leg, the forepart of which is crea, the shin; the hinder part sura the call; the outer protuberance at the love and is termed malleolus externus, the outer protuberance at the outer protuberance at the divided into three parts, viz. the tarsus, metalenus, and toes; behind is calcaneus the heel, before is tarsus the instep; from thence to the toes is the metatarsus; the top of the foot is termed dorsum pedis; the under part to tole planta pedis; and the toes digiti pedis.

Q. What are the bones of the leg?

A. As under this denomination I comprehend all the lower limb, as before observed; the bones are, the os femoris, patella, tibia; tibula, tarsus, metatarsus and digiti pedis.

Q. Which is the os femoris?

A. The os femoris is the thigh bone, and the longest and strongest bone of the whole human frame. In its upper extremity is to be observed a very large round head, and in this head a cavity is destined for the ligamentum rotundum, by means of which it is fixed in the acetabulum or socket of the os innominatum, and its luxation upwards prevented. There are two processes or apophyses near the head, called the greater and lesser trochanters, which are evidently formed for the insertion of muscles. The neck of this bone lies between

between the trochanters and the head, to which is affixed a robust annular ligament, which contains the head and neck of the bone, as it were in a case. The neck is not frait but oblique, nearly horizontal, and turning fomewhat outward, fo formed for keeping the thighs afunder, by which means we tread the firmer; the spongy cavernous aructure of this extremity of the bone, renders it less liable to fractures in this part: the apertures for the ingress of several vessels are also observable, as is the large finus between the great trochanter and the neck, into which muscles. are inferted. The middle of the thigh bone, for the conveniency of the mufcles, is a little convex forwards, and fomewhat concave backwards, which would make it subject to break backwards, if there was not a strong ridge on the backfide (termed linea afpera) which strengthens it sufficiently, and serves also for advantageous infertions for feveral muscles. The great cavity of this bone is also remarkable for containing the marrow. At the lower end of this bone are two large heads, with a cavity between them, for the articulation with the tibia; also a posterior cavity which gives passage for the large vessels to descend securely to the leg. There is besides an interior cavity for the placing of the patella: and lastly, two condyles, or tubercles, placed near the heads, for the origin of the muscles which move the foot. The strength and sirmness of this bone are furprizingly great; hence the use of the thigh.

thigh bone is to support and sustain the weight of the whole body; and its moveable articulation at the head gives way to the easy motion of the body, while the feet are unmoved.

- Q. Which is the patella?

A. The patella is a bone which covers the fore part of the joint of the knee, called also rotula, and vulgarly the knee-pan or pan-bone of the knee. It is a convex on the outside, and on the inside unequal, having an eminence and two depressions. Its substance is spongeous, and consequently it is brittle. It is connected by tendons and ligaments to the tibia and os semoris, which is the ligament by which it is connected to the thigh, and has a motion of ascent and descent in the slexion of the tibia. Its use is to secure the extensors of the tibia, lest passing over the joint, they might be too much exposed to external injuries.

The part situated from the knee to the ancle is properly called the leg, and consists of two bones besides the patella above-men-

tioned, viz. the tibia and the fibula.

Q. Which is the tibia?

A. The tibia is the bigger bone of the leg, (so called from its resemblance to an old musical pipe or slute) situated, at its anterior internal part, and continued in near a strait line, from the thigh-bone to the inner ancle; it is hard and firm with a cavity in its middle; it is almost triangular; its fore and sharp edge is called the skin. This bone is large at

its upper end, and has two finuses or sockets, which receive the two protuberances, or lower heads of the thigh-bone; and the production or process, which is between the sinuses of the tibia, is received in the finus, or cavity, which divides the two protuberances of the femur before-mentioned, and to this rough process of the tibia the cross ligaments of this joint are connected. One side of the upper end has a small process, which is received into a small sinus of the fibula, and on the fore part, a little below the patella, is another process, into which the ligament, or tendon of the patella, is inferted, and the tendons of the extenfors of the leg. Its lower extremity, which is much smaller than its upper, has a remarkable process, which forms the inner ancle, and fecures this bone from diflocating outwards: it has also a pretty large finus, which receives the convex head of the aftragalus, and the protuberance is received into the finus, in the convex head of the same bone. It has another shallow sinus in the side of its lower end, which receives the fibula. per end of this bone is triangular, as before observed, and even concave on the fide next the muscles, to make room for them; but lower, as the muscles grow less and tendinous, the bone grows rounder.

Q. Which is the fibula?

A. The fibula is the outer and fmaller bone of the leg, called also perone. It lies on the outside of the tibia, and its upper end does does not reach to the knee, but is only joined to the external fide of the tibia, receiving the small lateral process of the upper end of that bone into a small sinus, which the fibula has in its inner fide. The lower end of this bone is received into the small finus of the tibia, and then it extends into a large process, which forms the outward ancle, embracing the external fide of the astragalus. Thus the inferior processes of the tibia and fibula concur in the articulation of the tarfus, which ferves to strengthen the ancle joint, and render a luxation less easy. The tibia and fibula do not touch one another but at their ends; the space which they leave in the middle is filled up by a ftrong membranous ligament, and fome muscles, which extend the feet and toes. The fibula has no particular motion of its own, but wholly follows that of the tibia; and it feems doubtful, whether or not this bone contributes to the support of the body: its great use is for the origin of the muscles, and even its shape is suited to theirs.

The bones of the feet are those of the tar-

fus, metatarfus, and toes.

Q. Which is the tarfus, and of what bones

does it confist?

A. The tarfus is the space between the bones of the leg and the metatarfus, consisting of seven bones, viz. the astragalus or talus; calcaneum or os calcis; os naviculare; os cuboides vel cubiforme; and the three offa cuneiformia. The first of these bones, which

is the astragalus, supports the tibia, and is supported by the os calcis, which being projected backwards, makes a long lever for the muscles to act with, that extend the ancle and raife the body upon the toes. These two bones have a confiderable motion between themselves, and the astragalus also with the os naviculare, and all the rest an obscure motion one with another, and with the bones of the metatarfus; the greatest part of these motions being towards the great toe, where is the greatest stress of action. These bones thus giving way are less liable to be broke, and as a fpring under the leg, make the motions of the body, in walking, more easy and graceful, and the bones which are supported by them less subject to be fractured in violent actions.

Q. Which is the metatarfus, and of how

many bones is it composed?

A. The metatarfus is the space between the tarsus and toes, consisting of five bones, articulated to the tarsus at one end, and to the toes at the other; the metatarsal bone which supports the great toe is much the largest, the greatest stress in walking falling upon that part; under the end of this lie the two sesamoid bones, which are of the same use as the patella.

Q. What are the digiti pedis, and how are

they composed?

A. The digiti pedis is a term given by anatomists to the toes. Each of the toes, except

cept the great one, consists of three phalanges; the great toe has but two, and the two last of the little toe frequently grow together. The toes are less than the fingers, and have much less free motion.

Q. What are the cartilages of the leg?

A. The cartilages of the lower extremity are those of the os femoris, patella, tibia, fibula, and bones of the foot.

Q. What are the cartilages of the os fe-

moris?

A. No part of the os femoris is covered with a cartilage, except the uniform convexity of the head, and the articular portion of the lower extremity: the trochanters in adults have no true cartilage.

Q. What cartilage has the patella?

A. The patella has a pretty thick cartilage on its posterior or articular side.

Q. What are the cartilages of the tibia?

A. The tibia has four or five proper cartilages and two additional ones; the former are those of the two superior surfaces of the head of the tibia, that which covers the small furface on the lower part of the external condyle; also that which covers the lower surface of the basis of the tibia, continued over the outfide of the inner ancle: and there are likewife superficial cartilaginous incrustations on the back part of both ancles for the paffage of tendons. The additional cartilages of the tibia are two in number, called femi-lunar or inter-articular, in the shape of a C. they lie

on the two upper surfaces of the head of the tibia.

Q. What cartilages has the fibula?

A. Two; one lying on the upper extremity of that bone, the other covers the infide of the inferior extremity, or of the outer ancle.

Q. What are the cartilages of the bones of

the foot?

A. The aftragalus is covered by three cartilages; the first of these is for the articulation of this bone with the tibia and fibula; the fecond for the os calcis; and the third for the os scaphoides vel naviculare. The os calcis has four cartilages, of which three are superior, one large, and two small, for its triple articulation with the astragalus; the fourth is anterior, for the os cuboides. The os naviculare has two cartilages, one posterior for its articulation with the aftragalus; and one anterior, divided into three parts for the three ossa cuneiformia. The os cuboides has two remarkable cartilages, one posterior for its articulation with the os calcis, and one anterior for its articulation with the two last metacarpal bones: it has likewife a cartilage on the infide for the os cuneiforme, which is next to it, and one on the lower fide. The three offa cuneiformia have each of them a posterior cartilage for their articulations with the os naviculare; and one anterior, for the three first metacarpal bones; they have likewife small cartilaginous furfaces on their fides, for their articulations with each other; and besides, the first and third bones are joined thereby to the lateral parts of the basis of the second metatarfal bone, and the third to the os cuboides. The bases and heads of the metatarfal bones are covered with cartilages. The phalanges have cartilages in the same manner at their bases and heads, except at the heads or extremities of the last. We should beware of confounding the remains of tendons, ligaments, and aponeuroses with true cartilages, as for instance, at the posterior part of the os calcis, &c.

Q. What are the ligaments of the leg?

A. The ligaments of the lower extremity are those of the os femoris, patella, tibia, fibula, and bones of the foot.

Q. What are the ligaments of the os femoris?

A. The os femoris is connected by its upper extremity to the os innominatum, and by the lower to the bones of the leg, by means of feveral ligaments. The ligaments of the upper extremity are two in number; one furrounds the whole articulation, with the cotyloide cavity, or acetabulum, and one is contained in the articulation. To these we may, though very improperly, add a third, which is of the nature of a capfular ligament. The first is termed the orbicular ligament of the head of the os femoris, and is the most considerable, largeft, and strongest of all the articular ligaments of the human body; it is fixed quite round the border of the acetabulum, or cotyloide cavity, as before observed; and from thence largely furrounds the whole head and superior portion of the neck of the os femoris, in the lower part of which neck it is closely inserted. The other ligament of the head of the os femoris, which lies in the joint, I call internal, and inter-articular. It resembles a flat cord, broad at one end, and narrow at the other; by its narrow end, it is inferted at the two angles of the notch of the cotyloide cavity, and by the other in the os femoris, these being a fort of depression in the head of the bone for the passage of the ligament. The ligaments of the lower extremity of the os femoris, by which this bone is connected with those of the leg, are fix in number, viz. one posterior, two lateral, two middle or crucial, and one capfular. The crucial ligaments lie within the joint, and are fixed by one end to the back part of the notch or opening, which parts the two condyles; they are furrounded by the capfular ligament, but all the rest lie on the outside thereof, being closely joined to it. The two lateral ligaments are fixed one to each tuberofity of the condyles. The posterior ligament is fixed a little above the convexity of the external condyle, from whence it descends obliquely behind the great notch and internal condyle. The capfular ligament glued, as it were, to the three former, is fixed quite round the inferior extremity of the os femoris, just above the cartilage, and the posterior part of the great notch; and from the cartilage and notch, through the small space upward, already mentioned, it covers the bone; and afterwards is inverted downward, to form the capfula for the the mucilaginous liquor of the joint. What remains to be described of all these ligaments, I shall refer to the description of those of the leg.

Q. What are the ligaments of the patella?

A. The patella is fastened to the tuberosity or spine of the tibia by a broad and very strong ligament, which runs down directly from the apex of the patella; it has likewise fmall lateral ligaments, fixed in the lower part of its edge on each fide, which are inferted anteriorly, and a little laterally in the edge of the head of the tibia. The capfular ligament of the joint of the knee, of which I described one part in speaking of the lower extremity of the os femoris, is fixed round the edge of the head of the tibia, and in the edge of the patella, so that the patella itself forms a portion of the mucilaginous capfula of the joint of the knee. The crucial ligaments and those of the semilunar cartilages, are included within this capfula; but the lateral and posterior ligaments, and those of the patella, lie without it, being closely joined to its outer surface; this capfula is likewise joined to a considerable portion of the circumference of the femilunar cartilages; and it not only contains and furrounds the ligaments already named, but likewise furnishes them with a very fine vagina. There is likewife a very thin ligament fixed by one end to the lower part of the cartilaginous fide of the patella, and by the other to the anterior part of the great notch, between

between the condyles of the os femoris, the use of which seems to be to hinder the articular fat from being compressed in the motions of the knee.

Q. What are the ligaments of the tibia?

A. I have already observed, that the tibia is connected with the os femoris by feveral ligaments, two lateral, one posterior, two middle, and one capfular; and I have shewn in what manner they are fixed in the lower extremity of the os femoris and patella. Their infertion in the bones of the leg are as follows. The innermost of the two lateral ligaments is fixed pretty low down, on the inner fide of the superior part of the tibia. The external lateral ligament is fixed in the upper extremity of both tibia and fibula; both are joined to the edge of the semilunar cartilages; and lie a little behind the middle of the articulation. The posterior ligament is fixed by feveral expansions in the posterior part of the head of the tibia. One of the crucial ligaments is fixed by one end to the internal fuperficial impression in the notch of the os femoris, and, by the other, to the notch in the head behind the cartilaginous tubercle, which lies between the two superior surfaces. The other crucial ligament is fixed by one end to the external impression in the notch of the os femoris, and, by the other, between the anterior portion of the furfaces just mentioned. There are feveral other small ligaments, but these are the principal ones. Q. What Q. What are the ligaments of the fibula?

A. The fibula is joined to the tibia by nine ligaments, four at each end, and one in the middle, called the interoffeous ligament. The ligaments at the upper extremity of the fibula are short, very strong, more or less oblique and compound; two of them are anterior, two posterior, and they lie on each other; the superior ligaments surrounding the articulation more closely than the inferior. They are all glued to the capsular ligament, which runs in between them and the articulation, and they are inferted round the edges of the cartilaginous surfaces in each bone. The ligaments of the lower extremity of the fibula are disposed

which lines each fide of the bones, down to the external ancle, with the inferior edge of the basis of the tibia. The middle or interoseous ligament of the two bones of the leg,

much after the same manner, that is, two before, and two behind; which run down on the lower end of the sibula forming the outer ancle. The middle space between the tibia and

so called because it fills up all the space left between them, being stretched from one to the other. It is perforated both above and below, and sometimes in several places besides, for the passage of the blood vessels and

nerves. It is not a ligament designed to tie these bones together, but rather a ligamentary septum for the insertion of muscles, in which

respect it supplies the place of hones; and

feems partly to be a continuation of the periofteum of the tibia and fibula. At the lower part of each ancle there are commonly three strong ligaments for the connexion of the bones of the tarsus with these of the leg, one that runs forwards, one backwards, and one more or less directly downward. The other ligamentary expansions and annular ligaments of this part, as they do not belong to the bones, will be mentioned in their proper place.

Q. What are the ligaments of the foot?

A. The foot being made up of many bones, must, besides those ligaments by which it is tied to the bones of the leg, have several others to connect not only the three parts of which it is composed, but also the particular bones belonging to each part. I have already mentioned the infertion of three ligaments in each ancle for the articulation of the ancles with the foot. The ligaments of the inner ancle are all fixed in the infide of the aftragalus. The anterior and middle ligaments of the outer ancle are fixed in the outfide of the astragalus; the posterior is chiefly fixed in the outside of the os calcis. All these ligaments lie on the outlide of the capfula, which furrounds the articulation of the aftragalus with the bones of the leg. The ligaments by which the bones of the tarfus are connected with each other, are short, flat, of different breadths, and run from one bone to another in various directions. The capfular ligaments of these bones go very little further than the edges of the

the articulations of one bone with another; they adhere very closely to the true ligaments, and are covered and tied by them. The os cuboides and the three offa cuneiformia, are also connected to the metatarsal bones, by several particular ligaments. The bones of the metatarfus are connected together by their bases and heads. The first phalanges of the toes are tied to the heads of the metatarfal bones, by a fort of orbicular ligament, fet round the edges of the cartilaginous portions of the head, and those of the bases of the phalanges. The fecond and third phalanges of all the toes, being articulated by ginglymi, have lateral ligaments, which go between the sides of the bases to the sides of the heads. The capsular ligaments of all these articulations are disposed like those of the tarsus already spoken of. See the muscles of the foot.

Q. What are the mucilaginous glands of

the leg?

A. The mucilaginous glands of the lower extremity lie in the finall spaces, depressions, and superficial notches near the edges of the cartilages of each joint; they are covered by the capsular ligaments, and more or less mixed with a satty substance. The glands of the knee, which lie near the edges of the patella, are the most considerable, being disposed of in form of fringes, and supported by a great quantity of satty matter, which makes in some measure one mass with them, and is contained within the capsular ligaments. There are

other mucilaginous glands, both above and below the edges of the femilunar cartilages; and likewise in the ham, some whereof serve for the joint, the rest for the crucial ligaments. These last lie in folds formed by the internal membrane of the capfular ligament, which give particular coverings to the crucial ligaments. The mucilaginous glands of the foot answer in number and figure to the depressions between the cartilaginous edges and ligaments.

Q. What are the muscles of the leg?

A. The muscles of the lower extremities are the ploas magnus, iliacus internus, pectinæus, triceps femoris, glutæus maximus, glutæus medius, glutæus minimus, pyriformis, quadratus femoris, obturator internus et externus, fascia lata, gracilis, sartorius, semitendinosus, semimembranosus, biceps tibiæ, popliteous, rectus tibiæ, vastus externus et internus, cruræus, gasterocnemius, plantaris, solæus, tibialis anticus, et posticus, peroneus longus et brevis, extensor pollicis pedis longus et brevis, flexor pollicis pedis longus et brevis, abductor poliicis pedis, do. transversalis, extenfor digitorum pedis longus et brevis, flexor digitorum pedis brevis vel perforatus, flexor digitorum pedis longus vel perforans lumbricales pedis, abductor minimi digiti pedis, abductor secundus minimi digiti pedis et interossei pedis.

Q. Which is the ploas magnus?

A. The ploas magnus is a long thick mufcle, fituated in the abdomen, on the lumbar region, adhering to the vertebræ of the loins, from the posterior part of the os ilium to the anterior part of the thigh. It arises laterally from the bodies and transverse processes of the four superior vertebræ of the loins, and the last of the back, and is inserted with the following muscle into the lesser trochanter. This is one of the flexor muscles of the thigh, and when the ploas parvus is wanting, this is larger.

Q. Which is the iliacus internus?

A. The iliacus internus arises from the internal concave part of the os ilium, and from its lower edge; it is inserted with the psoas magnus, and employed in the same action. These move the thigh forward in walking.

Q. Which is the pectinæus?

A. The pectinæus is a small, flat and pretty long muscle, situated obliquely between the os pubis and the upper part of the os femoris. It arises from the os pubis or pectinis, near the joining of that bone with its fellow, and is inserted into the linea aspera of the thigh-bone, four singers breadth below the lesser trochanter. This bends the thigh and turns the toes outward.

Q. Which is the triceps femoris?

A. The triceps femoris is the adductor muscle of the thigh, having three heads and as many insertions; two arise under the pectineus, and a third from the os pubis and ischium; and are inferted into the whole linea afpera and the inner apophysis of the os femoris. This also bends the thigh and turns the toes outward.

Q. Which is the glutæus maximus?

A. The glutæus maximus arises from the os coccygis, the spine of the sacrum and adjoining parts; and is inserted by a strong tendon into the upper part of the linea aspera of the os semoris, sour singers breadth below the great trochanter. This extends the thigh, and both these together being contracted, occasionally assist the levatores and in supporting the anus.

Q. Which is the glutæus medius?

A. The glutæus medius ariseth under the former from the spine of the ilium, and is inserted into the superior and external parts of the great trochanter of the os semoris. This extends the thigh outward.

Q. Which is the glutæus minimus?

A. The glutæus minimus arises from the ilium under the former, and is inserted into the superior and anterior part of the great trochanter, and neck of the thigh-bone, to extend the thigh. These three muscles form the buttocks.

Q. Which is the pyriformis?

A. The pyriformis, vel iliacus externus, arises from the inside of the lower part of the os facrum, and from thence it runs transversely towards the joint of the hip, and is inserted into the upper part of the sinus, at the root of the

the great trochanter.—This affifts fomewhat in extending the thigh, but more in turning it outward.

Q. Which is the quadratus femoris?

A. The quadratus femoris is fituated transversely between the tuberosity of the ischium, and the great trochanter. It arises from the obtuse process of the ischium, and is inserted into the upper part of the linea aspera of the os femoris, between the two trochanters. This draws the thigh inward, and directs the toes outward.

Q. Which is the obturator internus?

A. The obturator internus, vel marsupialis, arises from the internal circumference of the hole that is between the ischium and the os pubis, and is inferted into the finus of the great trochanter. This turns the thigh outward.

·Q. Which is the obturator externus?

A. The obturator externus arises opposite to the former, from the external circumference of the fame hole; and is also inferted into the finus of the great trochanter. This also turns the thigh outward.

These four last-mentioned muscles acting with the extensors, prevent their turning the toes inward, and in stepping forwards are con-

tinually acting to turn the toes outward.

Q. Which is the fascia lata?

A. The fascia lata vel abductor fascialis femoris, called also musculus membranosus, arises from the fore part of the spine of the ilium. Soon after its origin it becomes entirely membranous,

branous, and closely furrounds the muscles of the thigh; after which it is inserted in the upper part of the tibia, near the head of the fibula, and from thence sends out an aponeurosis, almost over the whole external muscles of the tibia, as those of the thigh bone. About the middle of the leg it grows loose, and is so continued to the top of the foot, being connected there, and at the lower part of the leg, to the annular ligaments which tie down the tendons. When this muscle acts, the leg and thigh are drawn outwards; it also helps to extend the joint of the knee, and to elevate both the thigh and leg.

All these muscles I have now mentioned serve to move the os femoris on the pelvis.

Q. Which is the gracilis?

A. The gracilis arises from the os pubis, close to the penis, and descending by the inside of the thigh, is inserted into the inside of the tibia, near the sartorius, four or five fingers breadth below the joint of the knee. This muscle draws the leg and thigh inwards, and helps to bend the knee.

Q. Which is the fartorius?

A. The fartorius is both an abductor and elevator, ferving to move the legs upwards and forwards, and to cross each other as taylors sit with them, whence the name; it arises from the ilium, and descending obliquely is inserted into the upper and inner part of the tibia, four or sive singers breadth below the point of the knee. This is the longest muscle in the human body.

X

Q. Which

A. The semi-tendinosus vel semi-nervosus arises from the ischium, and is inserted into the upper part of the tibia, near the gracilis and fartorius. It helps to bend the leg and extend the thigh.

Q. Which is the semimembranosus?

A. The semimembranosus is a long thin muscle, partly tendinous, whence its name, situated on the backside of the thigh, a little towards the inside, being one of the five slexors of the tibia. It arises from the ischium a little above the seminervosus, and is inserted into the upper part of the tibia, but nearer the joint than the former muscle for the same use. Before it is inserted, it sends off sometimes an aponeurosis like that of the biceps. The semitendinosus and semimembranosus make the internal ham string.

Q. Which is the biceps tibiæ?

A. The biceps tibiæ, vel femoris, is a muscle with two heads, the superior from the ischium, the other from the linea aspera of the os femoris: both which join together, and are inserted by one tendon into the superior and external part of the sibula, to bend the leg, and the first head also extends the thigh. The tendon of this muscle forms the external ham string, when the knee is bent; besides the office commonly assigned to this muscle, in bending the tibia together with the fartorius and membranosus, it is likewise employed in turning the leg, together with the foot and toes.

toes outwards, when we fit down with the knees bended.

Q. Which is the popliteus?

A. The popliteus is a small muscle obliquely pyramidal, fituated under the ham, from whence its name. It arises from the outer apophysis of the os femoris, and thence running obliquely inward, is inferted into the backfide of the head of the tibia. It affifts the flexors, and draws the tibia towards the outer apophysis of the thigh bone.

Q. Which is the rectus tibiæ?

A. The rectus tibiæ, vel cruris, arises from the upper part of the acetabulum of the os inominatum, and from the processus inominatus of the ilium, and is inferted together with the three following muscles into the patella. bends the thigh and extends the tibia.

O. Which is the vaftus externus?

A. The vastus externus is a very large fleshy muscle, almost as long as the os femoris, lying on the outfide of the thigh. It arises from the fore part of the great trochanter, and fuperior part of the linea aspera of the os femoris, and is inferted into the upper and external part of the patella. This muscle extends the tibia.

O. Which is the vastus internus?

A. The vastus internus is very like the former, and fituated in the same manner, on the infide of the os femoris. It arifes from the linea aspera, and is inserted into the patella, to extend the tibia; and the fibres of this muscle being oblique, it keeps the patella in its place. Q. Which X 2

Q. Which is the cruræus?

A. The cruræus arises from the fore part of the thigh bone between the two trochanters, and lying close upon the bone, it joins its tendon with the three former muscles, which are inserted into the patella; the patella being tied down by a strong ligament to the tibia.

These three last muscles extend the tibia only, and might very properly be called exten-

for tibiæ triceps.

These ten muscles not only move the leg upon the thigh, but also the thigh upon the leg, the popliteus excepted.

Q. Which is the gasterocnemius?

A. The gasterocnemius is a pretty thick, broad and oblong muscle, which forms a great part of the calf of the leg. It arises from the posterior part of the os semoris, which soon becoming large bellies, unite, and then form a statement of the os calcis. Its use is to extend the tarsus and bend the knee.

Q. Which is the plantaris?

A. The plantaris arises from the external condyle of the os semoris, under the outer beginning of the gasterocnemius in the ham, and soon forming a small tendon is so continued betwixt the foregoing and subsequent muscles, and is inserted with them. It bends the knee and extends the tarsus.

Q. Which is the folæus?

A. The folæus vel gasterocnemius internus, arises from the upper part of the tibia, and one third

third of the fibula below the popliteus, and is inserted with the two foregoing muscles by a strong tendon into the upper and back part of the os calcis. This muscle only extends the tarsus.

Q. Which is the tibialis anticus?

A. The tibialis anticus is fituated on the fore fide of the leg, and is one of the flexor muscles; it arises from the tibia, and is inserted laterally into the internal os cuneiforme and the internal metatarsal bone. This bends and turns the tarsus inward.

Q. Which is the tibialis posticus?

A. The tibialis posticus, or abductor muscle of the foot, arises in the upper part of the tibia between that bone and the fibula, and going between the bones through a perforation in the interosseous ligament, it passes under the inner ancle, and is inserted into the os naviculare. It extends and turns inward the tarsus.

Q. Which is the peroneus longus?

A. The peroneus longus arises from the external and superior part of the sibula, and its tendon passing under the outer ancle and the muscles situated on the bottom of the foot, is inserted into the beginning of the metatarsal bone of the great toe, and the os cuneiforme next that bone. This turns the tarsus outward, and directs the force of the other extensors of the tarsus toward the ball of the great toe.

Q. Which is the peroneus brevis?

A. The

A. The peroneus brevis arises from the middle of the fibula, under a part of the former, passes under the outerancle, and is inserted into the os metatarsi of the little toe. Its use is to

extend the tarfus and turn it outward.

These two last muscles, riding over the lower end of the fibula, are often the cause of a sprain in the outward ancle, when they are vehemently exerted to fave a fall. These seven muscles (which Winflow divides into nine) ferve to move the tarfus and foot on the leg.

Q. Which is the extensor pollicis pedis

longus?

A. The extensor pollicis pedis longus arises from the upper and fore part of the fibula, and the interoffeous ligaments, and is inferted into the last bone of the great toe. This also bends the tarfus with a much longer lever than it extends the toe.

Q. Which is the extensor pollicis pedis

brevis?

A. The extensor pollicis pedis brevis, arises from the fore part of the os calcis, and foon becoming a long slender tendon, it passes obliquely over the upper part of the foot, and is inferted into the same place with the former.

Q. Which is the flexor pollicis pedis longus? A. The flexor pollicis pedis longus, is an antagonist to the extensor longus, arising opposite to it from the back part of the fibula, and its tendon passing under the inner ancle, is inserted into the last bone of the great toe. This extends the tarfus, as well as bends the toe. Q. Which

A. The flexor pollicis pedis brevis? A. The flexor pollicis pedis brevis arises from the two lesser offa cuneiformia the os cuboides and os calcis. This muscle is inserted into the offa sesamoidea, which are tied by a ligament to the first bone of the great toe, reckoning only two bones to the great toe. These muscles bend the great toe.

Q. Which is the abductor pollicis pedis?

A. The abductor pollicis pedis arises from the os calcis and os naviculare, and passing by the os cuneiforme majus, and the external sefamoid bone of the great toe, it is inserted into the first bone of the great toe. This muscle is less an abductor than a slexor; it also very much helps to constrict the foot length ways.

Q. Which is the abductor pollicis pedis

transversalis?

A. The abductor pollicis pedis transversalis arises from the lower end of the metatarsal bone of the toe next the least, and is inserted into the internal sesamoid bone. This truly is an abductor of the great toe, and helps to keep the constrictor of the bottom of the foot.

Q. Which is the extensor digitorum pedis

longus ?

A. The extensor digitorum pedis longus arises from the upper part of the tibia, and from the sibula and interosseous ligaments, then dividing into five tendons: four of them are inserted into the second bone of each lesser toe, and the fifth into the beginning of the metatar-sal bone of the least toe. The four first tendons X 4 only

only of this muscle extend the toes, but the whole five bend the tarsus.

Q. Which is the extensor digitorum brevis?

A. The extensor digitorum brevis arises, together with the extensor pollicis brevis, from the os calcis, and dividing into three small tendons, is inserted into the second joint of the three toes next the great one.

The long extensors of the toes serve not only to extend them, but also contribute to the bending of the ancle; but the short extensors arising below the ancle, extend the toes only.

Q. Which is the flexor digitorum pedis bre-

vis, vel perforatus pedis?

A. The flexor digitorum pedis brevis, vel perforatus pedis, is the flexor of the fecond phalanx, and is the inmost of all the common muscles of the toes. It arises from the under and back part of the os calcis, thence passing towards the four lesser toes, divided into four tendons, which are inserted into the beginning of the second bone or phalanx of each of the lesser toes. These tendons are divided or perforated to let through the tendons of the following muscles.

Q. Which is the flexor digitorum pedis

longus, vel perforans?

A. The flexor digitorum pedis longus, vel perforans, is the flexor of the third phalanx. It arises from the back part of the tibia, above the infertion of the popliteus, and part of the fibula; thence descending under the os calcis to the bottom of the foot, it there becomes tendinous, often crosses, croffes, and in most bodies communicates with the flexor longus pollicis pedis; then it divides into four tendons, which pass through those of the flexor brevis, and are inserted into the third bone of the four lesser toes. This muscle also extends the tarsus.

The fecond beginning of this muscle arises from the os calcis, and joins the tendons where they divide. This portion only bends the toes.

Q. Which is the lumbricales pedis?

A. The lumbricales pedis are the flexors of the first phalanx, and arise from the tendons of the perforans; they are inserted into the first bone of each of the lesser toes, which they bend.

Q. Which is the abductor minimi digiti

pedis?

A. The abductor minimi digiti pedis arises by the perforatus from the os calcis, and being part of it inserted into the metatarsal bone of the least toe, it receives another beginning from the os cuboides, and is inserted into the first bone of the least toe, which it bends and pulls outward, and very much helps to constrict the bottom of the foot.

Q. Which is the abductor fecundus minimi

digiti pedis?

A. The abductor fecundus minimi digiti pedis, arifes under the former muscle from the metatarsal bone, and is inserted into the little toe.

Q. What are the interoffei pedis?

A. The interossei pedis are seven muscles like those of the hands, arising like them from

the metatarfal bones, and are inferted into the last joints of the four leffer toes; being in their progress attached to the tendons, which extend the fecond joints of the toes, they will extend both these joints. These muscles may be fitly divided into external and internal; the internal also bend the first joints, as do all the interossei in the hand; but here the outer ones extend the first joints. The muscles that move the fingers and leffer toes fideways, are alike in number, though this motion of the toes is in a manner lost from the use of shoes. The muscles that bend or extend the last joints of the toes, will also move the second and first, and those that move the second will also move the first, as they do in the fingers.

Q. What are the arteries of the leg?

A. The arteries of the thigh, leg, and foot, are, the arteria cruralis, pudica externa, poplica, tibialis, anterior et posterior, vel suralis, plantaris externa et interna, and peronæa.

Q. Which is the arteria cruralis?

A. The iliac artery (which see) goes out of the abdomen, between the ligamentum Fallopii and tendon of the psoas, at the union of the os ilium and os pubis, and there it takes the name of arteria cruralis. It sends off, first of all, three small branches; one of which, called pudica externa, goes over the crural vein to the skin and ligament of the penis, and to the inguinal glands, communicating with the pudica interna. The second branch goes to the musculus pectineus: and the third, to the upper part of the sartorius. All these branches surnish

nish likewise the neighbouring anterior integuments. Afterwards the crural artery runs down on the head of the os femoris, and gets on the inside of the crural vein, about three fingers breadth from where it goes out of the abdomen. In this progress it is covered only by the Ikin and fat, and lies on the pectineus and triceps femoris. In changing its situation, it fends out the three following confiderable branches: the external branch runs on the upper fide of the thigh to the crureus vastus externus, rectus tibiæ, fascia lata, and glutæus medius. The middle branch runs down on the infide of the thigh, between the heads of the triceps, to which muscle it is distributed, a ramification of which perforates the triceps, and is distributed to the glutæus maximus, semitendinofus, femimembranofus, biceps, and to the neighbouring integuments. The internal branch runs backward on the quadrigemini, towards the great trochanter; and having fent a branch into the joint of the os femoris, it then runs downward, and is ramified on all the muscles that lie on the backside of that bone. one of which enters the bone itself on one fide of the linea aspera. - The arteria cruralis having detached all these branches, runs down between the fartorius, vastus internus, and triceps, giving branches to all the parts near it. It is covered by the fartorius all the way to the lower part of the thigh, where it is inflected backward over the triceps a little above the internal condyle of the os femoris. Afterward continuing

continuing its course through the hollow of the ham, it is called poplitea.

Q. How is the arteria poplitea distributed?

A. The arteria poplitea, while in the ham, is covered only by the integument, sending off branches toward each fide, some to the joint of the knee, one of which, at least, passes between the crucial ligaments. As it runs down it fends branches to the gastrocnemii and popliteus; and at the backfide of the head of the tibia, fends off a branch to each fide. Before the poplitea ends, it fends a fmall artery down on the backfide of the interoffeous ligament, very near the tibia, into which it enters by a particular hole a little above the middle portion of the bone. As the poplitea ends, it divides into two principal branches, one of which runs between the heads of the tibia and fibula, paffing from behind forwards on the interoffeous ligament, where it takes the name of tibialis anterior. The fecond branch divides into two others, the internal and largest called tibialis posterior; the other named peronæa posterior.

Q. How is the arteria tibialis anterior dif-

tributed?

A. The arteria tibialis anterior, having passed between the heads of the tibia and fibula, sends simall branches upward and laterally. Afterwards this tibial artery runs down on the fore side of the interosseous ligament, toward the outside of the tibia, between the musculus tibialis anticus and extensor policis, for about two-thirds of the length of the tibia, when it passes

passes on the foreside, under the common annular ligament, and extensor pollicis pedis, to the articulation of the foot; which giving off feveral branches to the right and left, and communicating with the tibialis posterior and peronæa posterior, so that these two are in a manner surrounded by arteries. At the joint of the foot it fends out branches, which run between the aftragalus and os calcis, being diftributed to the articulation and to the bones of the tarfus: the communications are here very numerous on all fides. Having paffed the fold of the foot, it fends off other branches on both fides, which communicating with others, make a kind of circle round the tarfus. Afterwards the tibialis anterior advances on the convex fide of the foot, as far as the interstice between the first and second metatarsal bones; between the heads of which, a large branch of it perforates the superior interosseous muscles, and forms an arch on the fide of the foot. It likewife fends two or three confiderable branches over the other metatarfal bones. Laftly, this artery terminates by two principal branches, one of which goes to the thenar, and infide of the great toe; the other is spent upon the outfide of the great toe, and the middle of the fecond toe.

Q. How is the tibialis posterior, vel suralis,

distributed?

A. The tibialis posterior, vel suralis, proceeding from the poplitea, as before observed, runs down between the solæi, tibialis posticus,

flexor digitorum communis and flexor pollicis, giving branches to these muscles, to the tibia, and to the marrow of that bone, through a particular canal in its posterior and upper part. Afterwards it runs behind the inner ancle, communicating with the tibialis anterior, then passes to the sole of the soot, and divides into two branches, one large or external, and the other small or internal, called plantaris exter-

na, et plantaris interna.

The great branch, or plantaris externa, passes on the concave side of the os calcis, obliquely under the sole of the foot, to the basis of the sisth metatarsal bone, and from thence runs in a kind of arch toward the great toe: the convex side of this arch supplies both sides of the last three toes, and the outside of the second toe, forming small communicating arches at the end, and sometimes at the middle of each toe, as in the hand. The concave side of the arch furnishes the neighbouring parts.

The small branch, or arteria plantaris interna, having reached beyond the middle of the sole of the fole of the foot, is divided into two; one goes to the great toe, the other to the first phalanges of the other toes, communicating with the ramifications from the arch already mentioned.

Q. How is the peronæa distributed?

A. The arteria peronæa, proceeding from the poplitea as before mentioned, runs down on the backfide of the fibula, between the folæus and flexor pollicis, to which, and to the neighbouring parts, it gives branches in its paf-

fage

fage all the way down to the os calcis, where it forms an arch with the tibialis posterior, between the astragalus and the tendo achillis; from thence it runs outward, and a little above the outer ancle communicates with the tibialis anterior by an arch, which sends several ramifications to the neighbouring parts.

In the description of the arteries throughout this work, I have said nothing of the cutaneous anastomoses, which are exceedingly beautiful in the fætus; nor of the frequent and considerable communications of small arteries upon the periosteum, which form a delicate kind of net-

work.

Q. What are the veins of the leg?

A. The veins of the thigh, leg, and foot, are the vena cruralis, faphena, sciatica, faphena minor, poplitea, tibialis anterior et posterior, vel suralis, plantares, and peronea.

Q. Which is the crural vein?

A. The vena cruralis goes out under the ligamentum Fallopii, on the infide of the crural artery, and immediately gives small branches to the inguinal glands, musculus pectineus, and parts of generation, which last are termed pudicæ externæ. About an inch below where it leaves the abdomen, it produces a large branch, which runs down anteriorly between the integuments and the sartorius almost all the way to the inside of the thigh, and from the condyles of the os semoris, it runs between the integuments and inner angle of the tibia to the fore part of the inner ancle, and is distributed

buted to the foot under the name of vena faphena, or faphena major. After giving out the faphena, the cruralis finks in between the muscles, and is distributed to all the inner or deep parts of the lower extremity, accompanying the crural artery to the very extremity of the foot, being all along more considerable than the artery.

Q. How is the vena faphena, or faphena

major, distributed?

A. The faphena major, arifing from the cruralis, as before-mentioned, in its paffage from the inguen to the foot, is covered only by the skin and fat; and after supplying the inferior inguinal glands and integuments, it runs down on the thigh as low as the middle of the fartorius, where it sends off several branches; and a little below the ham it runs in among the muscles situated there, and communicates with another branch, which may be termed faphena minor. Afterwards the trunk of the great faphena runs down on the infide of the tibia, lying always near the skin; and at the upper part of that bone, it fends branches forward, outward, and backward. The anterior branches go to the integuments on the upper part of the leg; the posterior, to those which cover the gasterocnemii, and the external branches also to the fat and integuments. About the middle of the tibia, it gives out a branch anteriorly, which runs along the integuments of the tibia all the way to the outer ancle : and as the faphena runs down on the infide of the tibia, it fends off branches to the neighbouring parts, and to the periosteum of the bone. At the lower part of the tibia, the faphena fends out a confiderable branch ob liquely forward over the joint of the tarfus, toward the outer ancle, fending off feveral branches. Lastly, the extremity of this trunk paffes on the forefide of the inner ancle, and runs under the skin along the interstice, between the first two metatarfal bones toward the great toe, where the vein terminates; but before it terminates at the great toe, it forms a kind of arch over the metatarfus, which communicates by feveral branches with the arch on the joint of the tarfus, and fends others to the toes.

Q. Which is the vena sciatica?

A. The sciatic vein arises from the crural, about the upper extremity of the musculus vastus internus, and runs down on the side of the crural trunk, covering the crural artery, almost as low as the ham, where it is again united to the trunk by an anastomosis. It has the name of sciatica from the sciatic nerve which it accompanies.

Q. Which is the vena faphena minor, and

how distributed?

A. The vena saphenaminor, velexterna, arises on the outside of the anastomosis of the sciatic with the crural before mentioned, and runs backward between the biceps and neighbouring muscles, and so down the backside of the leg, very near the skin, all the way to the outer ancle. In

its course downward it sends off a branch about the middle of the backside of the thigh; and immediately above and below the ham, this vein sends out other branches, which all communicate with the saphena major. About the beginning of the tendon achillis the little saphena runs outward in the integuments, toward the outer ancles, where it terminates in cutaneous ramifications sent to every side.

Q. Which is the vena poplitea?

A. The crural vein, a little above the ham, takes the name of poplitea, and as it runs down between the two condyles, it gives branches to the flexor muscles of the leg, and is ramified like the artery: afterwards it loses its name, being divided into three considerable branches, called tibialis anterior, tibialis posterior, and peronæa; of which the tibialis posterior is most frequently a continuation of the trunk, and the other two like branches.

Q. How is the tibialis anterior distributed?

A. The tibialis anterior, having sent some small branches to the muscles behind the heads of the two bones of the leg, perforates the interosseous ligament, and runs between the superior portions of the musculus tibialis anticus, and extensor digitorum pedis, sending off small superficial branches to the head of the tibia and sibula, and to the joint of the knee, which communicate with the lateral branches of the poplitea; afterwards it divides into two or three branches, which, with the anterior tibial artery, surrounds it by small communicating circles.

These

These branches, having reached the lower extremity of the leg, unite in one, which are afterwards ramified in the foot.

Q. How is the vena tibialis posterior, vel

furalis, distributed?

A. The posterior tibial vein sends off a branch from its origin toward the infide, to the gasterocnemii and solæus muscles, named suralis. Afterward the posterior tibialis runs down between the foleus and tibialis posticus, giving branches to each of them. It is divided in the fame manner as the tibialis anterior, and accompanies the artery as low as the outer ancle, furnishing the musculus tibialis posticus, and the long flexors of the toes. Lastly, it passes on the infide of the os calcis, under the fole of the foot, where it forms the vena plantares, fending ramifications to the toes nearly in the fame manner as the arteriæ plantares.

Q. How is the vena peronæa distributed?

A. The vena peronæa is likewise double, and fometimes triple, and proceeds from the poplitea, as before observed: it runs down on the infide of the fibula, in the fame manner as the arteria peronæa, which it likewise furrounds, by communicating branches, after the manner of the tibiales. It runs down as low as the outer ancle, fending ramifications to the musculi peronæi, and long flexors of the toes.

Q. What are the nerves of the leg?

A. The principal nerves are the crural and sciatic. The acterior crural nerve is formed by the union of branches, from the first, se-Y 2 cond.

cond, third, and fourth lumbar nerves, which running along the mulculus ploas, escapes with the large blood vessels out of the abdomen, and is distributed to the muscles and teguments on the fore part of the thigh: one branch of this crural nerve accompanies the vena faphena as far as the ancle. The fituation of the kidney upon, and the course of the ureter over these nerves, is the reason that in a nephritis the trunk of the body cannot be raised erect without great pain, that the thigh loses its fenfibility, and that it is drawn forwards. The remainder of the fourth and the fifth lumbar nerves join with the first, second, and third that proceed from the os facrum: these five, when united, constitute the largest nerve of the body, fo well known by the name of sciatic, or ischiatic nerve, which seems to be bigger in proportion than the nerves of any other part. When this nerve is obstructed, we see how unable we are to support ourselves, or to walk.

The sciatic nerve then goes out at the large hollow, behind the great tubercle of the os ischium, and passing over the quadrigemini muscles, runs down the posterior part of the thigh, giving off every where as it goes nerves to the teguments and muscles of the thigh and leg. At the ham it splits into two, the smaller mounting over the fibula, and serving the peronæi, slexores pedis, and extensores digitorum, is continued to the toes; the larger trunk sinks under the gasterocnemii, and then divides; one is spent on the muscles of the back.

of the leg and teguments, while the other is continued by the inner ancle to the foot, and then subdivides; one branch is distributed after the same manner as the ulnaris, and the other as the radialis in the hand.

Q. What are the glands of the leg?

A. The principal are the inguinal glands, fituated in each fide in the groin, near the crural vessels, and are in various diseases apt to grow tumid and inflamed. Abscesses are often formed in them; but their use in the body is not easily understood.

About the flexure of the knee and foot, there are also found here and there some small glands; as also in some places between and among the muscles; but these are here omitted, for the

reason I have before mentioned.

DIALOGUE VIII.

Of the Parts of Generation in both Sexes.

Q. WHAT are the parts of generation in men?

A. The parts of generation in men, are the testes, vesiculæ seminales, prostatæ, and penis; to which may be added the urethra, common to both men and women.

Q. What are the testes?

Y 3

A. The

A. The testes, or testicles, separate the seed from the blood, and are two oval bodies feated in the scrotum, which is the external coat or covering, made up of the epidermis and cutis, and immediately under the latter is a thick cellular texture closely adhering to it. The loose membrane immediately underneath the fcrotum is called dartos, which invelopes each tefficle fingly, and forms a septum or partition between the two: it is also wholly cellular, without muscular fibres, and without fat. Under the dartos is the cremaster muscle, (see page 189,) one to each tefficle: it is probably owing to the action of this muscle, that the scrotum is gathered up into rugæ by cold; as neither the ferotum nor dartos are furnished with muscular fibres, as before observed. Under this muscle is the first of the proper integuments, or coats, called proceffus vaginalis, tunica vaginalis, and elythroides; which is formed by the dilatation of the productions of the external membranes of the peritonæum through the abdominal rings: this coat is continued from the peritonæum to the testicle, which it embraces loosely, the whole body of the tefticle adhering to one end of the epididymis hereafter mentioned: it encloses the spermatic veins and arteries, termed vala preparantia; and likewife those vessels which convey the femen from the tefficles to the veficulæ seminales, called vasa deferentia; but it is divided by a feptum, or an adhesion immediately above the tefficie, so that no liquor can pass out of that part of this membrane (which encloses the spermatic vessels) into that which encloses the testicle. Large quantities of water (which is termed the hydrops testis) are sometimes found in either or both of these cavities, which disease is easily remedied by a puncture with a lancet, but rarely cured without opening the cavity where the water is contained, as in finuous ulcers: the true hernia aquosa is from the abdomen, which either extends the peritonæum into the scrotum, or breaks it, and then forms a new membrane, which thickens as it extends, as in the aneurifmal and atheromatous tumors. The dropfy in this cyst, for such it properly is, rarely admits of more than a palliative cure, by puncture or tapping, like the dropfy of the abdomen, and this with some difficulty, because the omentum usually, and sometimes the gut, descends with it. The other proper and last coat, which immediately encloses the tefticles, is called albuginea, from its white colour; it is ftrong and thick, very smooth and equal: and the branches of the vafa preparantia are finely weaved upon it. The lubitance, or kernelly part of the testicles, is of a white colour, and of a pretty firm texture, univerfally allowed to be a continuation of the evanescent branches of the spermatic artery, rolled up together. It is divided into more than twenty portions or clusters, separated from one another by as many partitions, which are productions of the albuginea; each cluster between two partitions terminates in one duct, which

which ducts (above twenty in number) meeting together, form a kind of net-work adhering to the albuginea: from this net-work arise ten or twelve other distinct ducts, which uniting, constitute the head or beginning of the epididymis; and quickly by their conflex form one larger duct; this single duct makes a roundish hard body on the upper and posterior part of the testicle, called epididymis, as it were an additional testicle; and at length terminates in the vas deferens, from whence it is evident the epididymi are the beginning of the vafa deferentia, which are excretory ducts to carry the elaborated feed into the veficulæ feminales. They pass from the epididymi of the testicles, together with the blood veffels, through the abdominal rings, all enveloped in one common sheath (the tunica vaginalis) called the spermatic cord or rope. When they have entered the abdomen, they separate from the spermatic veffels, and pass directly through the pelvis, close to the bladder, to the vesiculæ seminales. The spermatic veins and arteries are termed vafa preparantia, as before observed: the spermatic artery (one on each fide) arises from the aorta descendens inferior, as mentioned in the description of that artery: it is the smallest artery in the whole body for the length it runs: the spermatic veins accompany the arteries through the abdominal rings to the testis: and as there is no red blood found in the substance of the testis within the albuginea, most certainly fecretion begins immediately upon the entry

entry of the arterial twigs into that substance: these arteries, unlike all others, arise small, and dilate in their progress, that the velocity of the blood may be fufficiently abated for the fecretion of fo viscid a fluid as the feed. The right testicle returns its vein into the cava, and the left into the emulgent vein on the same side as before mentioned; both because it is the readiest course, and because this spermatic vein may not be obliged to cross the aorta, whose pulse would be subject to stop the blood, which returns from the testicles very slowly, by reafon of the narrow orifices of the spermatic arteries, and the largeness of the veins. testicles have many lymphæducts, which difcharge themselves into the inguinal glands. Their nerves come from the intercostal, and twenty-first of the spine. The spermatic arteries carry the blood from the aorta to the testicles, which separate that part of it which is fit for feed. The veins carry back to the cava what blood remains after the fecretion of the feed. The feed is farther purified in the epididymes, and in coition is carried by the vafa deferentia into the urethra.

Q. What are the veficulæ feminales?

A. The vesiculæ seminales are two membranous bags or receptacles, one on each side, situated at the lower and posterior part of the neck of the bladder, into which the semen is received from the vas deferens, and there deposited and accumulated, till it is thrown out by the urethra in the venereal act. They are

each of them one uninterrupted cavity, shut at one end, and sending out a duct at the other, into which duct the vas deferens opens. The external membrane is made up of muscular fibres, which in time of coition contract and press the vesiculæ, ejecting the seed through the prostate glands into the urethra.

The feed passes in time of coition from the vasa deferentia, as well as from these receptacles; for when the ducts into the urethra are distended, that is the direct course from the vasa deferentia, as well as from the vesiculæ seminales.

Q. What are the proflatæ?

A. The proftatæ are two glands, or rather one, about the fize of a walnut. These glands lie between the vesiculæ seminales and penis, under the offa pubis, almost within the pelvis, furrounding and closely embracing the beginning of the urethra. They separate a limpid glutinous humour, which is carried into the urethra by feveral ducts, which enter near those of the proftatæ. This liquor is thrown into the urethra in the act of generation, along with the femen, and is mixed with it, to make it flow more eafily. If the venereal infection reaches the prostate glands, it will sometimes make large abscesses, which are apt to form sinuses, and even make a passage into the bladder. It is often cured by opening the finuses, and confuming the difeafed parts by efcharotics: but a much better and easier way is to cut out all the fiftulous and diseased parts at once,

Q. What is the penis?

A. The shape, situation, and use of the penis are fo well known, as to need no descrip-The fubftance of the penis is composed of two spongious bodies, called corpora cavernosa, part of the urethra, the glans or nut at its extremity, and its integuments. The two corpora cavernofa, vel spongiofa, (which during erection make the chief bulk of the penis) arise from the os pubis, on each side, and are continued to the root of the glans: they arise distinct, but are soon after united, divided only by a feptum its whole length; it is strongly connected to the os pubis by a small ligament, which arises from its back. They take their name from being porous like sponge, and capable of being enlarged by the blood penetrating their substance, which forms a fort of loofe net-work internally, upon which the branches of the blood veffels are curioufly fpread. When the blood is stopped in the great veins of the penis, that which comes by the arteries runs through feveral small holes in the fide of their capillary branches into the cavities of the net-work, by which means the corpora cavernosa become distended, and the penis erected.

The glands of the penis, though in appearance a part of its body, is in reality a continuation of the spongy substance of the urethra, resected over its extremity, and expanded in the form we see. It is covered over with a thin epidermis, under which there are numerous nervous papillæ, which render it extreme-

Q. What is the urethra?

A. The urethra is a canal which runs along the under-side of the corpora cavernosa, through which both the urine and seed pass; it begins at the neck of the bladder, and runs in a furrow between the two corpora spongiosa penis to the extremity of the glans. Its whole length, without the erection of the penis, is about twelve or thirteen inches: its thickness about that of a goose quill. It consists of two thick strong membranes, with a spongy texture between them: its beginning at the neck of the bladder, is surrounded and covered by the prostate, as hath been said; at its emersion from

from which gland it becomes thicker and wider for the length of an inch, which thick part is called its bulb, from the refemblance it bears to a bulbous root. Its inner membrane is pierced with many holes here and there, thro' which, from a glandular apparatus in the spongy substance of the urethra, a mucilaginous liquor is furnished, serving to defend it against the acrimony of the urine. Besides these orifices, which some call lacunæ, Cowper, in a particular treatife, hath described and beautifully delineated three glands, two near the bulb of the urethra, one on each fide, about the bigness of a pea, and a third, less than the other two, at the bending of the urethra, under the os pubis. The first two are often found; but sometimes wanting or very small: the third is but feldom met with by the best anatomists. Where they exist, they probably ferve for the same use as the other lacunæ. Women, though their urethra is shorter and wider, are not without the same kind of lacunæ, ferving for the same use, and liable to the fame infection I shall presently mention. All these glands have excretory ducts into the urethra, and from them are fecreted all the matter which flows from the urethra in a gonorrhæa, whether venereal or not. In the venereal infection, the urethra and the glands are first inflamed by the contagious matter, that causes a heat of urine, which abates as soon as the glands begin to discharge freely; but if by chance this ditease continues till any part of the urethra urethra is ulcerated, the ulcer never heals without a cicatrix, which constricts the urethra, and occasions what is vulgarly called a caruncle.

Q. What are the parts of generation proper

to women?

A. The parts of generation proper to women are generally divided into external and internal. The external parts of generation in women are the mons veneris, which is that rifing of fat covered with hair, above the vulva or rima magna, upon the os pubis; the great doubling of the skin on each side the rima is called labia pudendi, which is only the skin fwelled by the fat underneath. These lips being a little separated, there appear the nymphæ, one on each fide the rima. In the angle which the nymphæ form in the great chink next the os pubis, is the extremity of the clitoris, strait under which appears the meatus urinarius, or orifice of the urethra. On the opposite side next the anus are the glandulæ myrtiformes, fituated in the fossa magna, or interior cavity of the pudendum muliebre; and in this angle of the chink, there is a ligament called the fork, which is torn in the first birth.

Q. What are the nymphæ?

A. The nymphæ are two membranous parts, fituated interiorly on each fide the rima; these two small bodies are red, and somewhat resemble the membranes, or wattles under a cock's throat; they are sometimes smaller, sometimes larger, and are continuous to the præputium of the clitoris. The nymphæ are spongious in their

their internal substance, and full of blood vesfels, and therefore they swell in coition; they receive vessels and nerves as the clitoris. nymphæ are full of nervous papillæ, whence their quick fense: they have also small glands that secrete a fatty matter. Their use seems to be to increase the pleasure in coition; to direct the course of the urine; and to defend the internal parts from external injuries. They are bigger in married women than in maids. nymphæ are fometimes fo large, as not only to hang without the labia pudendi, but also to prove very troublesome, so as to require extirpation. Though this operation is rarely found necessary in our parts of the world, but is frequently practifed in the East; being properly the circumcifion of women.

Q. What is the clitoris?

A. The clitoris, (which some call mentula muliebris) is fituated in the fore part of the vulva, at the angle which the nymphæ form with each other. It is generally about the bigness of the uvula, and its shape much refembles the shape of that part, yet it is sometimes found as large as the penis: but even in this case it has no urethra; for though it has a glars like the penis, it is not perforated. The prepuce, covering the glans of the clitoris, is formed of the cutis of the pudendum, and furnished with nervous papillæ: hence it is of exquisite sensibility to the touch. The clitoris, in its common natural state, lies entirely buried under the skin or prepuce; no part of it appearing

pearing but its extremity covered with the præputium. The substance of the clitoris is composed of two spongious bodies, such as those of the penis; they arise distinctly from the lower part of the os pubis, and foon after unite and form the body of the clitoris and its glans, which is of exquisite sense, as already mentioned. Before these spongious bodies unite, they are called crura, as those of the penis: the two muscles which are said to erect the clitoris in coition, in the same manner as the muscles of the penis erect that part, are already spoken of. The glans of the clitoris is usually covered with a fœtid matter, like that of the glans of the penis. It is also connected to the os pubis, by a ligament, in the same manner as the penis is in men. The clitoris receives veins and arteries from the hæmorrhoidal vessels and the pudenda; and nerves from the intercostals, which are likewise diftributed through all the parts of the vulva. The use of the clitoris is to produce a titillation in coition, and it is faid to be the chief feat of pleasure to women in that act, as the glans is in men.

Q. What are the internal parts of generation

in women?

A. The internal parts of generation in women are the vagina, uterus or womb, ovaries, and Fallopian tubes. A little lower than the clitoris, between it and the vagina, or rather just within the vagina, is the orifice of the urethra, termed meatus urinarius. The urethra in women

The

men is the same as in men, only shorter, wider, and more strait; it is naturally as big as a goose-quill.

Q. What is the vagina?

A. The vagina, or neck of the womb, is a large canal which reaches from the external orifice, or os pudendi, to the internal mouth of the uterus. It is usually about five or fix fingers breadth long in maids, and one and a half broad; but it varies in different subjects, and in married women who have borne children; its length and bigness cannot be determined, because it lengthens in the time a woman is with child, and it dilates in the time of birth. The texture of the vagina is membranous, being composed of two membranes, of which the inner, which lines its cavity, is nervous, and full of wrinkles and fulci, especially in its fore part. It has three or four small glands on that fide next the rectum, which pour into it a vifcous humour in time of copulation. The rugæ or wrinkles of this membrane, are for the friction of the balanus or glans penis, to increase the pleasure in coition, both to the man and woman; and to render the part capable of the necessary dilatation in parturition. These rugæ are largest in maids; in married women they are much fainter, and feem as if worn down; and in women who have borne children, they are almost entirely obliterated. The external membrane of the vagina is made of muscular fibres, which, as occasion requires, dilate and contract, become long or short, for adjusting its cavity to the length and bigness of the penis.

The lower part, or orifice of the vagina, is contracted with a sphincter that it may grasp the penis closely: (See muscul. sphincter, vaginæ) and to the abundance of nervous papillæ in the inner membrane, is owing its quick sensation. The oscular orifices of the excretory ducts of the glands, situated under the rugæ of the vagina before mentioned, are called lacunæ: these glands are the seat of a gonorrhæa in women, as the glands in the urethra are in the male. The vagina receives veins and arteries from the hypogastric and hæmorrhoidal vessels. It has nerves from the os sacrum.

Near the beginning of the vagina, immediately behind the orifice of the meatus urinarius, is situated a valve, called Hymen, (from the god of marriage, in the heathen mythology) which commonly passes among us for a test of virginity. The following is Mr. Cheselden's description of it: the hymen is a valve constantly found in the vagina of children, which, looking towards the orifice of the vagina, closes it, but as children grow up, and the sphincter vaginæ grows strong enough to contract and close the orifice of the vagina, this valve becoming useless ceases to encrease, and is there known by the name of glandulæ or carunculæ myrtiformes. There have been a few instances in which the edges of this growing together, it continued unperforate, until it has been necessary to make an incision to let out the menses.

Q. What is the uterus?

A. The uterus or womb, is feated at the end of the vagina, and lies also between the bladder and rectum. The os pubis is a fence to it before, the os facrum behind, and the ilium on each side, these forming as it were a bason for it, termed the pelvis; but because it must swell whilst women are with child, there is a greater space between these bones than in men; and for this reason it is, that women are bigger in the haunches than men. The figure of the uterus is like a pear, from its internal orifice to its bottom; in women not with child, the length of the uterus is about three inches; its breadth in the upper part two, and in the lower part one inch; its thickness is about an inch and a half; in maids, indeed, the uterus is much smaller than this, (the cavity being about the fize of a small almond:) but in women with child it is larger, according to the time of gestation. It presses the bowels, and reaches to the navel towards the time of delivery; whilst at other times it does not rise above the os facrum. The upper and broader part of the uterus is called the fundus; and the lower part, into which the vagina opens, is termed the cervix: its orifice into the vagina is called os uteri, and by fome os tincæ, from the refemblance it bears to a tench's mouth; it may be also compared to the mouth of a young puppy dog, and the glans penis. This internal mouth of the womb is very small in virgins, but in women who have had children, or who are with child, it is larger; and in the last it is always 2 2

always closed up with a glutinous humour. In the time of delivery, it in a wonderful manner expands itself, so as to give passage to the child. The substance of the womb is vascular, being composed of a various plexus of fleshy fibres, with a great number of vessels between: in women not with child it is compact and firm; in those with child it is spongy and finuous, and is capable of wonderful dilatation, without any diminution of its thickness. Its bottom, or fundus, grows thick as it dilates, so that in the last months of gestation it is at least an inch thick, where the placenta adheres, because its roots run into the substance of the womb. The uterus is covered externally with a membrane from the peritonæum, and internally its cavity is lined with a porous and nervous membrane; but in women with child, this inner membrane almost entirely difappears. The blood vessels of the uterus are tortuous, and make a vast number of anastomofes with one another: they open by a number of little mouths into the uterus and vagina, and are the fources of the menstrual discharge. The arteries and veins of the uterus are branches of the hypogastric and spermatic vessels, whose large ramifications inofculate with each other. as also the branches on one side of the womb with those of the other. The veins are greatly larger than the arteries, especially in women with child.

About the age of puberty, when the blood which used to be spent in the increase of the body

body, being accumulated, distends the vessels, it breaks forth once a month at those of the uterus; because of all the veins of the body, which stand perpendicular to the horizon, these only are without valves. This evacuation is called menstrua, menses, and catamenia; and

vulgarly flowers, courses, &c.

The nerves of the womb come from the intercostals, and from those which come from the os facrum. There are also several lymphatics upon its outfide, which unite by little and little into greater branches, and discharge themselves into the refervatory of the chyle. The uterus is tied by two round ligaments; and by what is called the ligamenta lata, being like two broad ligaments, though it is only a production or continuation of the peritonæum, from the fides of the womb: the ovaria are fastened to one edge of the ligamenta lata, and the tubæ Fallopianæ run along the other; thefe productions, or continuations of the peritonæum, are commonly compared to the wings of a bat, and are therefore called vespertilionis alæ. The two round ligaments, or ligamenta rotunda, arise from the sides of the uterus, and pass through the oblique and transverse muscles of the abdomen to the groin, in the same manner as the feminal vessels do in men. This way the gut passes in a hernia intestinalis in women. (See muscles of the abdomen.) These ligaments are partly continued or joined to the musculus fascia lata, on the upper part of the inside of the thigh; and from thence comes the

pain that women big with child feel in this place. The substance of these ligaments is hard, but covered with a great number of blood vessels; they are pretty big at the bottom of the womb, but they grow smaller and slatter as they approach the os pubis.

Q. What are the ovaria?

A. The ovaria, termed by the ancients testes muliebres, are two bodies of a depressed oval figure, fituated one on each fide the fundus uteri; they are commonly about half the fize of men's testicles, but differ according to the age and temperament of the subject. They are white, smooth, and largest in persons in the vigour of their age, and in women who are most mature; in such subjects they are found of two drachms weight, and furnished with a number of prominent vesicles. In old people they scarce weigh half a drachm, and are dry, corrugated, and deformed with cicatrices. The ovaria are furrounded by a ftrong white membrane from the peritonæum, and are of membranous substance, fibrous, reticulated, and full of veffels varioufly interwoven; they contain from ten to twenty, or more, pellucid eggs, which are supposed to contain the first rudiments of the fœtus; the largest of them is not so big as a pea. There are two arteries, and two veins, which pals to and from the ovaries, or testes, in the same manner that they do to and from the testicles in men; but make more windings, and the arteries dilate more fuddenly, in proportion as they are shorter. These arteries

teries and veins detach branches into the uterus and Fallopian tubes, and not only make communications betwixt the artery and vein on one fide, and those of the other, but also with the proper vessels of the uterus, which are detached from the internal iliac arteries and veins. Besides the spermatic vessels, the ovaria have nerves from the intercostals and lymphatics, which discharge themselves into the common receptacle. The ovaria are connected to the fundus uteri, by means of the ligamenta rotunda; to the Fallopian tubes, and the sides of the pelvis, by the ligamenta lata, or vespertilionis alæ; and to many other parts by means of the spermatic vessels.

Q. What are the tubæ Fallopianæ?

A. The Fallopian tubes are two canals of a tortuous figure, but approaching to a conic form, are joined to the fundus uteri one on each fide; one end of the tube is connected to the uterus, and opens into its cavity; its orifice is finall, being about the fize of a large hog's briftle. The other end is much larger, and is fimbriated, or fringed round the edges, therefore called morfus diaboli; it is free, and fluctuates about the abdomen, and when there is occasion, this extremity applies itself to the ovary, embracing it with these muscular segments or fringes. Though the extremities of these tubes are small, the middle, or widet part is about the fize of one's little finger. Their length is about fix, seven, or eight fingers breadth, different in different subjects; they

they are connected to the ovaria by the alæ vespertilionis. Their substance is membranous and cavernous: they are composed of a double membrane; the exterior one seems to be continuous with the peritonæum, and the interior, with the interior membrane of the ute-They are wrinkled on the inner furface, and are imbued with a lubricous humour; but they are not cellular in the human body, as in other animals. They are furnished with a great number of veffels having the fame veins, arteries, nerves, and lymphatics, as the ovaria: and have a cavernous substance between their membranes, by means of which they are rendered rigid in applying their mouths to the ovaria; they are also moistened on their inner surface by these vessels. Their use in generation is very great; they become erect in the time of coition, from the influx of the blood and spirits, and at that time, by a natural motion, they apply their loofe fringed extremity, to the ovaries, which are furrounded and embraced by them. In this state they convey to them the prolific matter of the male femen injected into the womb; and after one of the ovula is impregnated, they receive and convey it to the womb. The Fallopian tubes unperforated, upon the foregoing hypothesis, must cause barrenness, and feed lodged in these tubes may have the same effect; which probably is often the case of common whores, and women that use coition too frequently; and perhaps the fat in the membrane, that connects the ovaria to the tubes.

tubes, may, in very fat women, so keep these tubes from the ovaria as to interrupt impregnation; besides these cases, too much or too little of the menses may destroy or interrupt conception; but the latter case, especially in young women, is very rare. From fuch causes as these, and not from imbecility, it is most probable that barenness oftener proceeds from women than men; and though women do not propagate to fo great an age as men, it feems not to be owing to an incapacity of being impregnated, but from their menses ceasing, and those vessels being closed, which should nourish the fœtus after the impregnation, as if on purpose to prevent the propagation of a feeble and infirm species. From this consideration, it is reasonable to suppose, that the perfection of the fœtus, notwithstanding it is first formed in the male feed, depends more upon the female than the male, or elfe that nature would, for the sake of the species, have been careful to hinder men as well as women from propagating in a declining age.

The feed of all animals, and particularly of mankind, is a whitish sluid secreted from the blood in the testes as above; in which Lewen-hoeck, by the help of microscopes, discovered an infinite number of animals like tadpoles, which he and others suppose to be men in miniature, and that one of these being entered into an egg in one of the ovaria, con-

ception is performed.

DIALOGUE IX.

Of the EMBRYO, and the FOETUS in UTERO.

Q. TAT HAT is the embryo? A. The embryo, is a name given to what a woman has conceived with, from the time of conception, till the parts are perfectly formed. From conception, until the egg hath been some five or fix days in the ute. rus, it is almost impossible to describe the changes it undergoes, upon the account of its minuteness, and for want of observations; after that time we may speak of the matter with more certainty. The egg is then a round pellucid globule, of only one membrane, containing the embryo, which hath a very large head, a small body, and no legs or arms, fwimming in a large quantity of a pellucid liquor; it hath then a large flat umbilicus, by which it adheres to the obtuse extremity of the egg. The proper membrane of the egg, called amnion, by degrees fends off all around its furface small flocculent vessels whereby it grows to those vessels of the utterus from whence the menses were wont to flow before pregnancy: by degrees these flocculent vessels, round the greatest and inferior part of the egg, cease to grow, degenerating into a fecond membrane, called chorion; while those vessels at the upper and blunt end, keep increasing in their diameters, and at length form the placenta to be

be described hereafter. In the first two months, or at the end of that time, there appears not any thing bony, the several parts or members not being distinctly formed till the third month. The head is first perfected; the thorax or breast next; then the abdomen; and last of all the extremities.

Q. When does the embryo take the name

of fœtus in utero?

A. When all the parts are distinctly formed. The sectus is involved in the same coats as the embryo, viz. the chorion externally, and the amnion internally, which immediately inclosed the sectus. They contain a quantity of liquor, which is a proper medium for so tender a being as the sectus to rest in, and partly secures it from external injuries, as the aqueous humour does the crystalline in the eye; and when the membranes burst in the time of birth, this humour lubricates the vagina, to render the birth less difficult.

Q. How is the fœtus in utero nourished?

A. This is a matter of much controversy, but the most reasonable opinion is, that the fœtus is nourished both by the mouth, and by the umbilical vessels.

Q. What is the placenta?

A. The placenta is a foft, roundish mass, found in the uterus of pregnant women, which from its resemblance to a cake, derives its name; and being also like the liver, was called by the ancients hepar uterinum, or wombliver. The number of placentæ in human subjects.

subjects, answers to that of the feetuses; and as these are usually single, the placenta is usually fo too: but when there are two or more fœtuses, there are always as many placentæ; yet in this case they often cohere together, so as to feem but one; but their veffels rarely communicate from one to the other. placenta is usually about fix inches in diameter, and one inch thick in the middle, growing a little thinner towards the circumference; it is furrounded with a fmooth membrane from the chorion and amnion, and its concave part is turned towards the fœtus; it has no particular part of the uterus to adhere to, but is usually fixed to the fundus uteri. Doctor Thomas Simfon, of St. Andrew's, however, thinks that it has no place in the ovarium, nor in the uterus till once the ovum becomes contiguous to the fundus, and then every contiguous part becomes really a placenta. The placenta is composed of the extreme branches of the umbilical vein and arteries, which are for the composition of this part divided into an infinite number of exceeding small branches to join a like number of the menstrual vessels of the uterus, which vessels of the uterus are made numerous rather than large, that the separation of the placenta from them may not be attended with a flux of blood fatal to the mother; for the fides of little veffels foon collapse and close, and they are more easily stopped, being compressed by the uterus itself as it shrinks, which it begins to do from the time

time of the birth; but when the placenta is feparated before the delivery, whether untimely or not, these vessels bleed until the uterus is discharged of the sœtus. The arteries and veins of the uterus, by which the menstrual purgations are made, are joined to the umbilical arteries and veins in the placenta, the arteries of the uterus to the veins in the placenta, and the veins in the uterus to the arteries of the placenta: by these vessels a large quantity of blood is continually slowing from the mother to the sœtus and back again.

Q. What is the navel-string?

A. The navel-ftring, or umbilical bloodvessels, termed funiculus umbilicalis, passes out at the navel of the fœtus to the placenta. It is commonly about two foot long, that the fœtus may have room to move without tearing the placenta from the uterus, which being done too foon, from whatever cause, occasions a miscarriage. This umbilical cord is composed of a spongy substance enclosing the blood-veffels, viz. two arteries and one vein, twifting about each other, particularly the arteries about the vein. The arteries arise from the internal iliacs of the child, and running up on each fide of the bladder, perforate the belly where the umbilical vein entered. With the arteries and vein above-mentioned, there also passes a vessel called urachus, which arises from the top of the bladder. The umbilical vessels, with the urachus, being shrunk up after the birth, lose much of their appearance, especially near the

the navel, where they are fometimes not to be distinguished. The umbilical vein goes from the navel directly to the liver, and there enters the great trunk of the vena portæ; it has its origin from small venous tubuli, arising from the chorion, where the evanescent twigs of the arteries are lost; and at length forming one large trunk, enters the umbilious of the embryo, goes to the liver as before observed, and opens into the finus of the vena portarum. Near this entrance there goes out the ductus venosus to the great trunk of the cava, which carries part of the blood that is brought by the umbilical vein that way into the cava, while the rest circulates with the blood in the porta; the whole of it not passing through the ductus venosus, as is generally believed, but a great part of it into branches of the porta, in the liver; otherwise there need be no communication between the umbilical vein and the porta. After the birth, when the umbilical vein is stopped, it becomes a ligament, and the ductus venofus foon shrinks and almost disappears, having no longer any blood flowing through it. The blood which flows from the mother to the fœtus by the umbilical vein is returned, all but a fmall quantity, which is referved for nutrition by the two umbilical arteries.

Q. How does the blood circulate from the mother to the fortus and from the fœtus to the mother?

A. The manner in which the blood flows from the mother to the fœtus, and from the fœtus to the mother, is as follows, viz. the uterine arteries in the mother bring arterial blood to the organ of the placenta in the egg; which being taken up by the beginning of the veins of the placenta, is transmitted to the fœtus, through the umbilical vein, into the vena portarum; and from thence into the cava and heart, and the two umbilical arteries in the fœtus being continued from the aorta, carry their blood to the placenta; which blood, at the adhesion of the placenta to the uterus, is taken up by the beginning of the uterine veins, and transmitted to the cava and heart of the mother; what was arterial in the one becoming venous in the other interchangeably; but the fœtus hath some peculiarities in its own proper circulation besides.

Q. What is the magnitude and weight of

the embryo and fœtus in utero?

A. When the ovum descends into the uterus, it is supposed to be about the size of a poppy seed, and in the third month augmented to the bigness of a goose egg. Ten days after conception, the child (according to some authors) weighs half a grain; at thirty days is increased to the weight of twenty-two grains; at three months, weighs betwixt two and three ounces; and at nine months, from ten to twelve, and sometimes sixteen pounds: by which calculation it would appear, that the progress of the sectus is quickest in the beginning

ning of the growth: for, from its tenth to the thirtieth day (according to this supposition) it encreases to three and forty times its weight. But all these calculations are uncertain.

Q. How is the uterus distended?

A. As the fœtus, &c. grow in bulk, the uterus is distended, but rerains the same, or nearly the same absolute thickness; the blood-vessels, which in its natural contracted state lay folded and rolled together like threads in a clue, being straitened out, now receive blood more freely than before, and become of greater diameters. So that what the uterus loses of thickness by distension, in its membranous or cellular parts, it gains by the enlargement of its vessels. As it is gradually more and more distended, its fundus rises more and more out of the pelvis up into the abdomen towards the colon and stomach, and being stretched in every dimension, creates the bulk we see in pregnancy.

Q. What is the natural fituation of the fœ-

tus in utero?

A. The natural situation of the setus in utero for the last months of gestation (for before that time it is uncertain and various) is, as it were sitting with its head and neck bent downwards and forwards; its knees drawing up to its cheeks; its heels lying back toward its hips; and its hands hanging down, and as embracing its knees. But there is a great variety of postures of the setus, both before

and at the time of delivery, as is shewn in midwifery.

Q. What are the differences, between a

fœtus in utero and an adult?

A. The more effential differences between the human fœtus ripe for birth, and an adult appear to be as follows. In the abdomen, the umbilical vein and arteries of the navel, and the canalis venofus in the liver, are in the foetus open and pervious; in adults they are contracted and folid. The liver is very large, the stomach is filled with a glutinous fluid, and the larger intestines, and often the ilium also, with the fæces called meconium. The renes fuccenturiati are larger in the fœtus than in The kidneys themselves are not fmooth and even on the furface, as in adults, but unequal, and in some measure resemble those of a calf. The urinary bladder is of a longer shape, and extends almost to the navel. The hymen in a female fœtus is very plain and obvious. In the thorax besides a peculiar fluid, found as well in this cavity as in the abdomen, the gland thymus is larger than it is in adults. The lungs, as they have never yet been inflated by breathing, are collapsed, and of a blackish colour; and if thrown into water they fink in it contrary to what is the case in adults, or those that have breathed. In the heart, the foramen ovale between the left and right auricle, and the canalis arteriofus, between the pulmonary artery and the aorta, are open, to ferve for a peculiar circulation in the fœtus, which Aa

which has not yet breathed; and there is in the inferior trunk of the vena cava, near the heart, a remarkable valve, called by Chefelden, valvula nobilis. In the head, besides its great fize in proportion to the body, we are to observe, that the offa cranii are in several places distant from one another, especially at the fontanella; and that the futures are wanting. The brain also is softer than in adults. The teeth are imperfect, and not rooted in the gums; they lie hid or buried under the gums, to appear at a more advanced period. The meatus auditorius is not yet perfect in the fcetus, and is entirely closed up by a peculiar membrane, which is continuous with the epidermis, and naturally disappears after delivery. The bones of the whole body, excepting a very few, are either foft or absolutely imperfect : some of them are merely cartilaginous, and the articulations are not at that time perfected.

DIALOGUE

Of the EYE.

Q. TIT HAT are the parts of the eye not yet described?

A. The bony focket, muscles, cartilages and ligaments of the external parts of the eye eye are already spoken of in Dialogue the sirst. The internal parts not yet mentioned are the glandulæ sebaceæ, caruncula lachrymalis, glandulæ lachrymalis, puncta lachrymalia, orbit; the coats or tunics, viz. tunica albuginea, adnata, or conjunctiva; tunica sclerotica; tunica cornea; tunica choroides; tunica uvea, (which contains the iris and pupilla) to which may be added the retina. The humours of the eye, viz. the aqueous, vitreous, and crystalline, to which may be added the extreme thin and sine vascular membrane called tunica arachnoides, and the vessels and nerves of the eye.

Q. What are the glandulæ sebaceæ?

A. The glandulæ sebaceæ are situated in the interior surface of the eye-lids: they serve for the secretion of an oleaginous sluid, which is of great use in preventing the attrition of the eye-lids, from their continual motion.

Q. What is the caruncula lachrymalis?

A. The caruncula lachrymalis is a little eminence situated in the larger angle or canthus major of the eye, serving to direct the tears to the puncta lachrymalia, and, according to some anatomists, they help to keep them open when the eyes are shut.

Q. What is the glandula lachrymalis?

A. The glandula lachrymalis is feated in the upper and outer part of the orbits, with its excretory ducts under the upper eye-lid. This gland separates the matter of the tears, which, by the continual motion of this lid, A a 2 furnishes

furnishes at all times water enough to wash off dirt, and to keep the external surface of the eye moist, without which the cornea would dry and wrinkle by the continual action of the external air. As the tears fall off the cornea, they are stopped by the edge of the under eyelid, along which they run till they fall into the puncta lachrymalia.

Q. What are the puncta lachrymalia?

A. The puncta lachrymalia are two small holes in the inner corner or great canthus of the eye, one in each eye-lid; they are fituated at the extremities of the tarfi or cartilages, and lead to a small membranous bag or lachrymal fac, which is feated in this corner upon the os lachrymale; from the bottom of which there goes a fmall pipe or nafal canal, which pierces this bone in the nofe opening under the upper lamina of the os spongiosum. It moistens the inner membrane of the nostrils, by the superfluous humour of the lachrymal gland. Sometimes the acrimony of this humour causes fneezing, which we may hinder by prefling the angle of the eye, and fo stop its running. Between these two puncta there is a caruncle (as above-mentioned) that ferves to keep the holes open when the eyes are shut.

Q. What is the orbit of the eye?

A. The orbit of the eye is that cavity in which the eye is contained, and is in all the vacant places filled with loofe fat, which is a proper medium for the eye to rest in, and serves as a socket for its motion. The proper

ball, or bulb, are its coats or tunics, the humours, and the vessels.

Q. What is the tunica albuginea?

A. The tunica albuginea, adnata, or conjunctiva is the first membrane or coat of the eyeball; it is a smooth membrane which covers so much of the eye, as is called the white, and being resected all round, lines the two eye-lids. Being thus returned from the eye to the inside of the eye-lids, it essectually hinders any extraneous bodies from getting behind the eye into the orbit, and smooths the parts it covers, which makes the friction less between the eye and the eye-lids. It is full of small veins and arteries, which appear big in an ophthalmia or instammation of the eyes.

Q. What is the tunica sclerotica?

A. The tunica sclerotica is a thick hard and smooth coat, extended from the cornea to the optic nerve; it is opake behind, but transparent before, where it makes the third coat called cornea. Both together make one firm case of a proper form for the use of the other coats and humours.

Q. What is the cornea?

A. The cornea, so called from its substance resembling the horn of a lanthorn, is convex, transparent, and composed of various laminæ, which are nourished by many blood-vessels, so fine as not even to hinder the smallest rays of light from entering the eye. The cornea is situated in the fore-part of the eye, surrounded by the sclerotica and albuginea; it has a most exquisite.

exquisite sense, to the end that the tears, upon the least pain, may be squeezed out of the lachrymal gland, to wash off any filth, which, by sticking to the cornea, might render it opake or dim.

Q. What is the tunica choroides?

A. The tunica choroides is the fourth coat of the eye, and is so named, on account of the multitude of blood-veffels refembling the chorion; it lies immediately under the sclerotica, and is much thinner than it, being a membrane of little firmness. It is blackish, or of a dusky brown colour, more or less inclining to red. This membrane, or coat, has a great number of blood-veffels which come from the sclerotiea. It is open, or has a hole before, for the passage of the rays of light, called pupilla; the part of this coat, which makes the circumference of the hole, and lies upon the side of the crystalline humour, is the uvea.

Q. What is the uvea, you mention?

A. The uvea is the fifth coat, and is only a white circle round the back fide of the choroides near the cornea, as has been faid. In this coat we observe, first the iris, which is a circular variously coloured part, being the anterior furface of the uvea, which furrounds the pupil; it is called the iris, because in different persons it is of different colours; hence the denomination of grey, blue, brown, hafel, black eyes, &c. The iris is entirely vascular, from which arises the variety of colours in the human eyes. Secondly, the pupil, or foramen, which is round in the human eye, nearly in the middle of the iris, and is capable of dilatation and contraction. Through this aperture, the rays of light pass to the crystalline, in order to be painted on the retina, and cause vision. Thirdly, its posterior surface, which is black, and in which, when this blackness is cleared away, there appears the sphincter of the pupil, formed of circular fibres for contraction, the ciliary fibres or processes, for the dilatation of the pupil; the ciliary ligament for the motion of the vitreous and crystalline humours; the arterial and venal circles, from the veffels, are in a wonderful manner distributed over the uvea; the choroides; the ligamentum ciliare; and the vitreous and crystalline humours; the ductus nigri, so called from their black colour, placed between the processes and the ligamentum ciliare; the space between the uvea and the cornea, called the anterior camera of the eye; and that between the uvea and crystalline, called its posterior camera, which is either much smaller, or entirely wanting.

Q. What is the retina?

A. The retina is a membrane which may be called the fixth tunic or coat; it lies immediately under the tunica choroides, and is a very delicate, tender, and as it were, mucous coat of the eye, or more properly, it is only an expansion of the optic nerve at the bottom of the eye. It is the great organ of vision, and A a 4 called

net: rays of light striking upon this membrane, the sensation is conveyed by the optic nerves to the common sensorium, the brain.

Q. What is the aqueous humour of the

eye?

A. The aqueous humour lies in the fore-part of the globe, immediately under the cornea: this humour is thin and liquid, of a spirituous nature, for it will not freeze in the greatest frost. This evinces the necessity of a continual supply of this humour; which is manifest it hath, because if the cornea be pricked, and this humour squeezed out, it will be again restored in ten or twelve hours: this aqueous humour lying foremost, seems chiefly of use to prevent the crystalline from being easily bruised by rubbing, or a blow; and perhaps it serves for the crystalline humour to move forward in, while we view near objects, and backward for remoter objects.

Q. What is the crystalline humour?

A. The crystalline humour is the second, and distinctly contained in a very fine coat or membrane called aranea or arachnoides, and is sufpended by means of the ciliary ligament, between the aqueous and vitreous humour, immediately behind the pupil; in this place it hangs free, and is moveable by means of the ligament just mentioned. It is composed of a multitude of lamellæ like the coats of an onion; and therefore also pellucid and vascu-

lar. There is also a small quantity of the aqueous humour contained within or under its coat. The crystalline being a thick, compact humour, in form of a flattish convex lens situated in the middle of the eye, serves to make that refraction of the rays of light necessary to make them meet in the retina, and form an image thereon, whereby vision may be performed.

Q. What is the vitreous humour?

A. The vitreous or glassy humour is the third humour of the eye, so called from its resemblance to glass in sustion, being like a fine clear jelly in appearance; it is thicker than the aqueous, but thinner than the crystalline; and is in greater abundance than the other two. It lies behind the crystalline, and fills up the greatest part of the eye: its foreside is concave for the crystalline humour to lodge in, and its back side being convex, the tunica retina is spread over it; it serves as a medium to keep the crystalline humour and the retina at a due distance.

Q. What are the blood-yessels and nerves

of the eye?

A. The eye is furnished in a most wonderful manner with nerves and blood-vessels in all its parts. The blood vessels of the eye are branches of the carotids and jugulars (which see) distributed to all parts of the eye in an amazing manner. The extreme minute ones convey only a fine and subtile lymph thither, by which means the tunics and humours of the eye are nourished; the veins partly carry the blood back to the finuses of the dura mater, and partly to the jugulars. The nerves of the eye are very numerous; besides, the optic nerves (described in pages 68, 97.) pierce the globe of the eye from the fide of the nofe, a little on the infide of the optic axis or center; their external coat, which is a production of the dura mater, is continued to the sclerotis, as their internal is from the pia mater to the choroides: and the medullary fibres paffing through all, are expanded on the retina, upon which the images of objects are painted. The centre of this expansion is infensible, and all rays which fall upon it are loft; confequently, that point of the object from which the rays come, is invisible to the eye; the reason of this proceeds, probably, from the blood-veffels, which enter the globe of the eye with the optic nerve, and cover this part of the retina. But whatfoever the cause be, there is a manifest advantage in the optic nerves being inserted on the infide of the optic axis. For if they had pierced the eye in the axis, the middle point of every object had been invisible, and where all things conduce to make us fee best, there we had not feen at all.

DIALOGUE XI.

Of the EAR.

Q. INTHAT are the parts of the ear not

yet described?

A. The bones, muscles and cartilages of the ear are already spoken of in Dialogue the fecond Its inner substance is cartilaginous, as hath been faid, which preferves its form, without being liable to break: its use is to collect founds, and direct them into the meatus auditorius, which is the paffage that leads to the drum. Hence those who have lost their ears, hear not so acutely as others, and endeavour to supply that want by applying the palm of their hand made hollow to their ears: hence the usefulness of hearing trumpets. The internal parts of the ear not yet described are as follows, viz. meatus auditorious, cerumen, membrana tympani, tympanum, tuba eustachiana, foramina vestibulum and labyrinth, and vessels of the ear.

Q. What is the meatus auditorius?

A. The meatus auditorius is the passage of the ear which leads to and terminates in the membrana tympani, or drum of the ear. is near an inch long, and about the fourth part of an inch wide, and its paffage is not strait but crooked, paffing first upwards and then downdownwards, when it has a small tendency up wards again, and the lower part of its extremity bends a little down to the obliquity of the membrana tympani. The beginning of this passage is cartilaginous, being a continuation of the concha contracted, the end of it is bony, which makes the greatest part of the upper and back part of the meatus, as the cartilage does of the lower and fore part. The whole internal cavity of the meatus is lined with a membrane which seems to be a continuation of the skin that covers the auricula, and which grows thinner and thinner as it approaches the tympanum.

Q. What is the cerumen?

A. The cirumen or wax of the ear is secreted from a number of little glands on the back fide of the membrane whose excretory ducts bring it into the meatus. This wax by its bitterness and viscidity, together with the hairs in the membrane above-mentioned, hinders infects from approaching the membrana tympani, which it likewife preferves against the injuries of the air. When this wax is accumulated in too great quantity, it obstructs the meatus, by filling up the passage, and creates the most common kind of deafness; the remedy of which is picking the ears, or fyringing them. Those great discharges of matter from the meatus auditorius, which are commonly called imposthumes in the ear, feem to be nothing else than ulcerations or great secretions from these glands. Q. What

Q. What is the membrana tympani?

A. The membrana tympani lies at the inner extremity of the meatus; it is extended upon a bony ridge or circle of the temporal bone. This is an oval membrane; thin, dry, firm, and transparent: though it seems stretched pretty tight, yet it is not plain, but concave outwardly toward the meatus, and inwardly convex; being pulled inward by the handle of the malleus, which is tied to it. This membrane does not entirely close the passage, but has on one side a small aperture covered with a valve, letting the smoke of tobacco taken in at the mouth find a passage through it out at the ears.

Q. What is the tympanum?

A. The tympanum, or drum of the ear, is a pretty large cavity, fituated behind the membrana tympani; it is about the fourth part of an inch deep, as much wide, and about half as high. In very young children it is always found full of mucus or a purulent matter, which feems necessary to prevent founds from affecting them too much, there being no provision to shut the ears, as there is for the eyes, Cheselden gives an account of a gentleman who had four children born deaf, on which he was advised to lay blifters upon the heads of the next children he might have, which he did to three who were born afterwards, and every one of them heard well. It feems not unreasonable to suppose that too great a quan-

tity of this mucus upon the drum might be the cause of deafness in the four children, and that the discharge made by the blisters in the latter case was the cause of their escaping the same misfortune. In this cavity are to be observed the periosteum, which lines it, a very thin membrane, and furnished with a great number of blood-veffels; and the chorda tympani being a little nerve composed of a combination of little branches of the fifth and feventh pairs, this is extended in the manner of a cord under the membrana tympani. In this cavity also there are four small bones already described in Dialogue the second, of which the first is the malleolus or hammer, so called because of its shape; its manubrium or handle, which is pretty long and small; it is fastened to the membrane of the tympanum, as hath been faid, and its head is articulated by ginglymus with the body of the incus; which bone is articulated by arthrodia with the stapes, and the arbiculare lies between. The malleus is moved inward by the musculus obliquus internus, or trochliaris, or internus mallei; it extends the membrana tympani, that it may be the more affected by impulse-of sounds when they are too weak. This muscle arises from the cartilaginous part of the Eustachian tube, and paffing from thence in a proper groove, it is reflected under a small process, and thence passes on perpendicular to the membrana tympani, to be inferted into the handle of the malleus,

malleus: fometimes with a double tendon parallel to this muscle lies another extensor of the membrana tympani, called obliquus externus or externus mallei; but this is not fo obvioufly an extensor as to be known to be so without an experiment. The muscle which relaxes the membrana tympani is called externus or laxator tympani. The relaxation of the membrane is made by this muscle, without our knowledge, when founds are too ftrong; and as the pupil of the eye is contracted when we have too much light, and dilated when there is too little, from what cause soever, so when sounds are too low, or the fense of hearing imperfect, from whatever cause the extensors of the tympanum stretch it to make the impulse of sounds more effectual upon it, just as in the case of the common drum, and the cords of any mufical instrument.

Q. What is the tubæ Eustachianæ?

A. The Eustachian tube or iter ad palatum, goes from the tympanum, (which cavity is called also the barrel of the ear); the beginning of this passage is very narrow and bony, the middle is cartilaginous, and its extremity, which opens into the back-part of the nostrils, just above the uvula, is about the third part of an inch wide, membranous, and dilated by some muscular fibres; they dilate the extremity of this passage, either when we open our mouth to hear more distinctly, or when it is necessary there should be a free communication between

between the external air, and that in the cavity of the tympanum. This passage, therefore, feems to be exactly of the same use with the hole in the fide of the common drum, that is, to let the air pass in and out from the barrel of the ear to make the membrane vibrate the better, and perhaps in the ear, which is closer than a common drum, to let air in or out as it alters in denfity, and if any fluid should be seperated in the barrel of the ear, to give it a paffage out. This paffage being obstructed, as it is fometimes, by a large polypus behind the uvula, causes great difficulty of hearing, and fometimes when the meatus auditorius is obstructed, a man opening his mouth wide, will hear pretty well through this passage, which is often fo open, as that fyringing water through the nose, it shall pass through into the barrel of the ear, and cause deafness for some time. To the stapes there is one muscle called musculus stapedius; it serves to pull the stapes from off the fenestra ovalis, which otherwise it covers.

Q. What is the foramina vestibulum and

labyrinth of the ear?

A. The foramina vestibulum are three holes in the cavity of the tympanum leading to the cavity called vestibulum; which forms the middle part of the labyrinth, or innermost cavity of the ear, termed a labyrinth, from its sinuosities and windings. The first of these holes is called fenestra, ovalis; to which, as hath

been said, the base of the stapes is fixed, and completely stops it. The next hole is the fenestra rotunda, and the other leads into the cavity of the mastoide process. After these may be observed three semicircular canals, which open into the vestibulum by five orifices.

Q. What is the cochlea of the ear?

A. The cochlea of the ear is placed oppofite to the semi circular canals before-mentioned, and is in the manner of a snail shell, forming two turns and a half in a spiral form. In this are to be remarked the nucleus and the canal, which is divided into two, by a spiral lamina: the upper of these opens into the vestibulum and is called the scala vestibuli, and the lower terminates in the hollow of the tympanum through the senestra rotunda, and is call-

ed the scala tympani.

A very delicate and fine membrane carried along through the cavities of the labyrinth, is formed of an expansion of the auditory nerve, and is the primary part of the organ of hearing, just as the retina is formed of the expansion of the optic nerve, and is the primary organ of seeing. Next may be observed the auditory canal, which is distinguished into the common and proper; the common is large, and has foraminula in it, passing into the labyrinth; the upper one is narrow and large, terminating partly in the cavity of the cranium, and partly between the styloide and mastoide processes.

Q. What

Q. What are the blood-vessels and nerves of the ear?

A. The arteries of the ear are from the carotids both external and internal; the veins run partly to the jugulars, and partly to the finuses of the dura mater; the nervus auditorius enters by the hole in the internal process of the os petrofum. It confifts of two bundles, of which one is hard, the other foft. Five branches of the portio mollis enter the vestibulum, and form a delicate web, which fends flips which run through the semicircular canals; the rest of the portio mollis enters the cochlea at the center of its base, and turns with the fpiral line, of which it probably makes the membranous part. The portio dura passes through its proper passage, to be distributed among the external parts about the ear.

There have been instances of the tympanum being destroyed by an ulcer, and the auditory bones cast, without destroying the hearing. From which, and other like cases, it may be concluded that the membrana tympani, though useful in hearing is not the seat of that sense, and if any disease in that membrane should obstruct the passage of sounds, to the internal parts of the ear, which are the seat of that sense, an artificial passage through that membrane might recover hearing, as the removing the crystalline humour, when that obstructs the

light recovers fight.

In all complaints of deafness, the practitioner ought carefully to examine the state of the ear,

and learn whether the patient has or hath had any abfcefs in the external meatus of the ear affected, that might have eroded the membrane of the tympanum; and he should try whether it is found or not. This is done by holding a lighted candle to the affected ear, and if upon a strong effort of exspiration determined towards the ears, by stopping the mouth and nostrils, the flame of the candle is fenfibly acted upon, we may conclude the membrane of the tympanum to be broken, and the hearing by that ear irrecoverably loft; for the cold air gets directly through the breach into the cavity of the tympanum, and by chilling and drying the delicate periosteum of the little bones, the membranes of the cavity, and the small muscles, renders the parts unfit for performing their functions. If the wax is hardened or even stony, drop in some oil of almonds over night, quickened with a very small proportion of chemical oil of anifeeds or amber; and the next morning inject with sope suds and a little tincture of myrrh or traumatic balsam, made milk-warm or somewhat warmer. Injections are to be pushed in gently, not to hurt the membrana tympani. If there is an abscess in the external meatus it ought to be healed as foon as poffible, that the pus may not at length affect the tympanum,

Part Tip Courses et the maps with a ted meshood any whether chriton but done it ed pateriol tod nothing idesconsare blast totte, for the cold air mers directlicate privote um or the light bones, the membreses of the cavity, and the finall mulcles. towers of the year is bardened or even thony, on charitees of a poet, and (se next morning inget with tope that and a diese thete of of lonewhat warmens dejections are to be fifthe, that the presence of the ships offer the the secondary but only the paramagings MAL WALL TOWNS

A.

BDOMEN described 177, 2 Arteries of 219 Bones pertaining thereto 180 Cartilages of 183 Glands of 184, 240 Ligaments of 183 Muscles of 186 Nerves of 239 Regions of 177 Veins of 230 Viscera of 195. Abortion, a cause of, 348. Abscesses in Ano, 192. Acetabulum coxendix 182 Imposthumations in, ibid. Adami Pomum 122. Adbesion of the Lungs 150. Acromion. [See Proceffus Acromion] 244 Adipose Membrane. [See Membrana Adiposa] 5. Adnata Oculi. [See Tunica Adnata] 357. After Birth. [See Placenta] 347. Air enters the Blood by the Lungs 150. Albuginea Oculi. [See Tunica Albuginea] 357 Testes. [Sec Teftes | 325. Alæ Verspertilionis 343. Almonds of the Ears. [See Tonfils] 108. Amnion of the Fætus 347. Anafarca, Situation of, 6 Ob ervations of, ibid. Anatomy, what so called, I Use of, ilid Should be well understood by all Physicians and Surgeons, ibid. Anchylofis, how formed 16. Ancle, Sprained, Cause of, 310. Animal Spirits, what so called 35. Animals that use no Exercise in Winter, how sublisted 24. Amaurofis, Caufe of, 99. Aneurism, Operation for, and its Success above the Cubit accounted for 275. Antra Maxillæ Superioris 62 Imposthumations in, Signs of and Cure 63. Antrum Highmorianum 100. Aorta, or great Artery 19 Afcendens, ibid. Descendens 20. Aponeurofis of Muscles 16. Apsphysis of Bones 8 Dentiformis 107, Appendicula Vermiformis 199. Aquæduct of Fallopius 103. Aqueous Humour of the Eye 360. Arachnoides, Tunica 89. Arbujculi Medullares 87. Areola of the Breaft 174. A-1150 B b . 3

Arm, including all the upper Limb 242 Arteries of 271 Bones of 242 Cartilages of 248 Glands of 285 Ligaments of 250 Mucilaginous Glands of 285 Muscles of 256 Nerves of 283 Veins of 279 Why the Right has more perfect Motions than the Left 134.

Arteries, what fo called 18 Found almost empty in dead Bodies

23 Quantity of Blood in, ibid. In general 19.

Arteries of the Abdomen 219 Arm 271 Brain 91 Ears 93, 370 Eyes 94 Head 92 Leg 314 Neck 126 Nose 93 Thorax 161 Tongue 93 Temples, ibid. Teeth, ibid. Parotid Gland ibid.

Arteries, why supposed to be so minutely divided before they enter

the Brain 95.

Arteria Axillaris 271 Brachialis 272 Bronchialis 169 Bilaria 221 Capfularis 224 Carotides 93 Carotis Externa 93, 127 Interna 94, 127 Cervicalis 95 Cæliaca 220 Coronariæ Cordis 94 Ventriculi 220 Cruralis 314 Cubitalis 274 Cyflicæ 221 Diaphragmaticæ 220 Duodenalis 221 Epiploica 222 Gastrica 221 Glutæa 228 Hæmorrhoidalis 223 Hepaticæ 220 Humeralis 272 Hypogastrica 227 Iliacæ 226 Externa, ibid. Interna 227 Minor 228 Intercostalis Inferior 164 Superior 163 Intestinalis 232 Lumbaris 225 Magna vel Aortæ 19 Mammaria Externa 165 Interna 163 Mediastina 162 Mesenterica inferior 223 Superior 222 Obturatriæ 230 Oesophageæ 164 Pancreatica 222 Pericardia 94 Peronæa 318 Phrenicæ 231 Plantaris Externa 318 Interna, ibid. Poplitæa 316 Pudica interna 229 Externa, ibid. Pulmonaris 150 Pylorica 221 Radialis 277 Renalis 223 Sacra 225 Scapulares 271 Sciatica 228 Spermatica 224 Splenica 222 Subclavia 161 Suralis 317 Thoraciea 161, 165 Thymica 162 Tibialis Anterior 316 Posterior, vel Suralis 317 Trachealis 163 Vasa Brevia 222 Vertebralis 128 Umbilicalis 227.

Articulations, how moistened 15.

Arytanoide Cartilage 122.

Aspera Arteria 157.
Atlas, Vertebræ, why so called 114 Different from the rest, ibid-

Auricles of the Heart 153.

Axillary Glands 285.

Axis Vertebræ 115.

B.

Back Bone, its Structure and Advantages of TI, II3 Crooked, described I35 Luxation of, how hindered, ibid. Little or no Motion in when distorted I36.

Barrel of the Ear. [See Tympanum] 365.

Barrenness of Women, a Cause of 344.

Befilic Vein 31.

Bason. [See Pelvis] 179.

Bile, Ducts of 208 How conveyed 210 Quantity secreted 212.

Biliary Vessels 208.

Belching in Obstructions of the Menses, Cause of 106.

Bladder, Urinary, Size of 218 Vessels and Nerves of 219 Its inner Coat, why exceeding sensible 225.

Bleeding, Accidents in 262 In the Basilic Vein, why great Care

Blood, Quantity of, in the human Body 23 Why it flows per Saltum, on an Artery being opened 25 How carried from, and returned to the Heart 27.

Blood Vessels, none in the Cuticle 3 Of the Marrow 10 Of a Muscle when cut or tied destroys its Action 18 Of the Brain, why they have a Multiplicity of turnings and windings 89.

Blows, where of bad consequence 54.

Body, its Parts how distinguished 2 Parts of 7 How bent forwards 113.

Boils, Seat of 6.

Bones described 8 Substance of, how formed 11 Use of, ibid.

The exact Knowledge of those of the human Body, the Foundation of all Anatomy 12.

Bones, their Number in the human Body 12.

Bones of the Abdomen 180 Arm 242 Ear, internal, 65 Eye,

Orbit of 53 Face 61 Head 52 Leg 296 Neck 114 Note

62 Pelvis 181 Thorax 133.

Brain descr bed 84 Meninges of 85 Weight of, ibid. Ventricles of 85 Prominences of 86 Coverings of 89 Arteries of 92 Glands of 92 Nerves of 36 Veins of, ibid.

Breast Bone. [See Sternum] 136.

Breasts, Glandulous 174. Grow full at the Age of Puberty, ibid.

Decrease when the Menses cease 175. Lactiferous Ducts of, ib.

Surrounded with Fat, ibid. Why they swell in Time of Pregnancy 176. Blood Vessels of 175. Nerves of, ibid.

Bronchia, or Ramifications of the Trachea 157.

Bronchial Glands 172.

Butcher's Meat, how blowed 6.

C.

Cæcum Intestinum 199.

Canals, Semicircular, of the internal Ear 368.

Canal of the Spine, how lined 87.

Capfulæ Atrabiliariæ 217.

Capfulæ Ligaments 15.

Carbuncles, Seat of 6.

Carpus, the Wrist 247. Bones of, ibid. Cartilages of 250 Ligaments of 254 Muscles of 264 Nerves of 283 Veins of 279

Arteries of 271.

Cartilage, what so called 15 Use of, ibid. Eroded, occasions an Anchylosis, ibid.

Eb 4

Cartilages

Cartilages subject to Offisication 15 Of the Abdomen 183 Arm 248 Ears 70 Foot 392 Eye-lids 68 Hand 250 Head 66 Larynx 122 Leg 292 Neck 116 Nofe 74 Thorax 138 Vertebræ 135.

Cartilago Annularis 122 Arytenoides, ibid. Cricoides, ibid. Ensiformis 136 Epiglottis 122 Scutiformis, ibid. Thyroides, ib.

Caruncles of the Urethra 333.

Caruncula Lachrymalis 355 Myrtiformes 334.

Carotid Arteries, where they pais through the Skull 54.

Catamenia. [See Menfes] 341.

Cautery, actual, on the Antihelix, its good effects in the Tooth-ach

Cawl. [See Omentum] 205.

Cellular Substance 5.

Cephalic Vein 34.

Cephalic Vein 34.

Cerebel'um, what 86 Figure of, 87. Lobules of 16 Peduncles of, ibid. Substance of, ibid.

Cerebrum, what 85 Described 86.

Cerebellum, Wounds of, mortal 54.

Cerumen of the Ear 364 How fecreted, ibid.

Cheft. [See Thorax] 132.

Child, Dead-born, how to discover 150.

Chorion of the Fœtus 346.

Choroides Oculi. [See Tunica] 358.

Choroides Other Ear 107, 366.

Chrystalline Humour of the Eye 360.

Ciliary Ligament and Cartilages 68.

Circulation between the Foetus and Market

Circulation between the Fætus and Mother 351 How performed in a Limb after the Part has been amputated 23. .

Circumcifion of Women 335. Circumcifion of Women 335.

Clavicula. [See Os] 243.

Clavicle. [See Collar-bone] ibid.

Clinoid, Processes 55.

Clitoris, Glans of 336 Præputium of, ibid.

Coats of the Eye. [See Tunica] 357.

Cochlea of the Ear 369.

Coition 329, 336.

Colon, Intestinum 200.

Collar Bone. [See Clavicula] 243,

Colour of the Eye. [See Iris] Of the Skin, to what owing 40

Conception, manner of 344.

Colics, in Obstructions of the Menses, the Cause 106.

Conglobate. [See Glands] 43.

Conglomerate [See Glands] ibid.

Conjunctiva Occuli. [See Tunica] 357.

Conflituent Parts of the Body 7. Corenary Artery of the Heart 156 Vein of the Heart, ibid. Artery of the Stomach 219 Vein of the Stomach 231.

Coracoides. [See Processus] 244.

Common Proffitutes, [See Wheres] 344.

Cornea Oculi. [See Tunica] 357. Corpora Cavernofa. [See Penis] 331 Olivaria of the Medallis Oblongata 38 Pyramidalia of the Cerebellum 39 Striata. [See Cerebrum | 85.

Corpus Callosum. [See Cerebrum] ibid. Reticulare, Colour of, and why white in Europeans and Black in Negroes 3 Apertures of, ibid. Use of, ibid.

Cortical Part of the Brain 85.

Costa. [See Ribs] 137. Courfes. [See Menses] 341.

Comper's Glands 333. Cranium. [See Skull] 56.

Crista Galli Process of the Cranium 55. Crooked Spine, the ill Consequences of 113. Crura Clitoridis 336 Fornicis of the Brain 88.

Crura Medullæ Oblongatæ 87.

Crural Artery 314 Nerve 323 Vein 319.

Cubitus, Arteries of 274 Bones of 245 Cartilages of 249 Ligaments of 252 Muscles of 256 Nerves of 284 Veins of 280. Cutaneous Glands 285.

Cuticle. | See Cuticula | 2.

Cuticula, or Scarf Skin 2 Colour of, ibid. Pores of, and microfcopical Observations on 3 Uses of, ib'd. Why White in Europeans and Black in Negroes, ibid. Is what is raifed by Blifters and Scalds 2.

Cutis vera, or true Skin 3 Blood Vessels of 4 Foramina of, ibid. Furrows of, ibid. Glands of, ibid. Pores of, ibid. Pyramidal Papillæ of, ibid. Thickness and Use of, ibid.

Cynic Spalm, Seat of 322.

Dartos of the Testicles 326.

Dead Born Child, how to discover 150.

Deafness, Treatment of 371 Perhaps might be cured in forme Cases by perforating the Membrani Tympani, ibid. Caused fometimes by a Polypus of the Nose compressing the Eustachian Tube 368 Caufe and Cure of, when Born fo 365.

Delirium, Cause of 371. Detrufor Urinæ 191.

Dentata. [See Bones of the Neck] 114.

Dentes. See Teeth 64. Diaftole of the Heart 191.

Digestion, Quantity of Fluids separated for it at every Meal 2x2.

Digital Arteries 277.

Digiti. [See Fingers] 275 Pedis. [See Toes] 216.

Diploe of the Cranium. [See Skull] 56.

Difeases, Symptoms of, the Cause, how known 36 Of the Ear' 370 Of the Joints, Caufe of 16.

Differted Back. [See crooked Spine] 113.

Differtions

INDE X

Diftertions but feldom in the Loins 136.

Dropfy, Water in, how formed 6 How evacuated, ibid.

Drum of the Ear. [See Tympanum] 365.

Dry Food, why we can eat less and digest less of it than of moist

Duet Salival of Steno 107 Ulcers and Wounds, of the Cure, ibid. Duelus Arteriofus 353 Choledochus 210 Cyfticus, ibid. Hepaticus, ibid. Nigri 359 Thoracicus 46 Venofus 353.

Duodenum Intestinum 197.

Dura Mater 90 Arteries of gr G'ands of, ibid. Motion of 91 Nerves of, ibid. Processes of, ibid. Sinuses of, ibid. Use of 90 Veins of 91. Drunkenness, how effected 95.

E.

Ear, Bones of 6; Canals of 368 Cartilages of 70 Cochlea of 369 Description of 363 Foramina of 369 Glands Ceruminose of 364 Labyrinth of 369 Ligaments of 70 Muscles of, ibid. Nerves of 103 Veffels of 93, 370 Veftibulum of 369 Wax of 364.

Eggs of Women. [See Ovaria] 342.

Emphysema, Cause of 6.

Embry, Magnitude of 351 Weight of, ibid. How formed 346.

Encephalon. [See Brain] 84.

Ear, why fometimes pained in the Tooth-ach for Its Labyrinth, how supplied with Nerves 38.

Enfiform Cartilage 136.

Epidermis, or Scarf Skin. [See Cuticula] 2.

Epid dimis of the Tefficles 328,

Epigastrium 177.

Epigastric Region 175.

Epiglottis 124 Glands of 131 Muscles of 124

Epiphysis of Bones. [See Bones] 8.

Epiphon. [Se Omentum] 205. Epistrophæus. [See Dentata] 114.

Euftachian Tube 367.

Excretory Veffels, what fo called 44, 209.

External Parts and Integuments 2.

Extremities. [See Limb] 23.
Fro, Ball of, seated in Fat 356 Coats of. [See Tunica] 357 Description of 354 Humours of 360 Muscles of 68 Parts of 363 Veffels and Nerves of 361.

r.

Face, Bones of 6r Nerves of, where they pass 56 Why sometimes convulfed in the Tooth ach 101. Falx of the Dura Mater 96.

Fallopian Tubes 343.

Fat, what so called 6 Use of 7 How consumed, ibid. Affords Notwishment when not received in the usual way, ibid.

Femoris. [See Os] 286.

Fenestra Ovalis 66, 268 Rotunda of the Ear, ibid.

Fibres, what to called 49. Fibula. [See Os] 289.

Fiftula in Ano 192 Seat of 202 Operation for, and Remarks on, ibid.

Fingers, Arteries of 275 Bones of 255 Cartilages of 250 Ligaments of 254 Mucilaginous Glands of, ibid. Mutcles of 265. Nerves of 283 Veins of 279.

Flanks. [See Ilia] 78.

Flowers. [See Menfes] 341.

Fætus in Utero, how formed 346 Circulation of its Blood 35% Magnitude of, ibid. Nutrition of 346 Situation of 352 Time of Birth 351 Weight of, ibid. Difference of from an Adult 353.

Foot Arteries of 316 Bones of 392 Cartilages of, ibid. Glands of 325 Mucilaginous Glands of 300 Ligaments of 299 Muscles of 310 Nerves of 324 Veins of 310.

Foramen Ovale 153.

Foramina Cerebri. [See Cerebrum] 85 Of the Skull 59 Their Use, ibid. Their Number internally 59 Vestibulum of the Ear 358.

Fore Atm. [See Cubitus] 274.
Fore Skin. [See Prepeutium] 332. Fork Ligament in the Folla Magna 334. Fornix. [See Cerebrum] 85.

Forebead, why dangerous to trepan \$3.
Food required according to exercise 24 The dryer it is, the more Saliva it requires in eating 109 Dry, why we can eat less of it than of moilt Food, ibid.

Fossa Magna. [See Vulva] 334. Fluids exceed the Solids 23. Funiculus Umbilicalis 349. Fundus Uteri. [See Uterus] 339.

G.

Gall Bladdet 309. Gang'ion of Nerves, the largest of the Body ros. Generation, Parts of, in both Sexes 325. Genital Parts of Men, ibid. Of Women 334. Gastric Glands 161.

Glands described 42 Their Use 43 Conglobate, described, ibid. Conglomerate, described, ibid.

Glands, Axillary 385 Bronchial 172 Cervical 131 Ceruminofe 210 Cutaneous, are not to be found 285 Dorfal, not always to be found 172 Jugular 131 Inguinal 325 Lachrymal 110

Lacunar 333 Miliary 4 Mucilaginous 16 Mucose 110 Occipital 131 Parotid 107 Pineal 92 Pituitary, ibid. Prostate
330 Renal 217 Salival, Secretion of, how performed according to the Food 109 Weight of, ibid. Sebaceous 109 Sublingual 108 Thymus, largest in new-born infants 173 Not 10 be found in old People, ibid. Thyroide 132.

Glands of the Abdomen 240 Arm 285 Brain 92 Ears 364 Epiglottis 131 Eyes 109, 355 Head 106 Joints 16 Leg 285 Neck 131 Thorax 172 Larynx 123 Oefophagus 160 Ure-thra 332 Pharynx 125.

Glands Mucilaginous, of the Joints 16 Of the Abdomen 240 Arm 285 Head 66 Leg 300 Neck 118 Thorax, ibid.

Glands of the human Body all put together do not weigh 4 Ounces

Glans, Penis 331 Clitoris 335.

Glottis 122. Globe of the Eye, Muscles of 69 Coats of. [See Tunica] 35%

Gonorrhaa, Symptoms of 333. Great Guts. [See Guts] 199. Groins. [See Inguina] 178. Guts. [See Intestines] 199. Gustato y Nerves 99. Gutta Serena, Caufe of 98.

Hair, Description of 50. Hearing, Organ of. [See Far] 101.

Heads of Infants, why open when Born 59:

Head 50 Bones of 52 Cartilages of 68 Ligaments of 66 Mucilaginous Glands of, ibid. Muscles of, ibid. Nerves of 98

Arteries of 93 Glands of 106 Veins of 96. Head, Muscles, that move it on the Trunk 82 And its Parts deferibed 50 In many Postures of, how Suffocation is prevented 55 Contents of 83 Viscera of, ibid. Why so soon moved by the Impression of Sounds on our Ear 103.

Hefar Uterinum. [See Placenta] 347.

Hernia Aquofa. (See Hydrops Teftis] 327. Heart 152 Auricles of 153 Cavities of, ibid. Nerves of 104 Pa'pitation of 130 Valves of 155 Ventricles of 154 Veffels

of 156 Situation of 152. Heat of Urine in a Gonorrheea, Cause of 333.

Hemispheres of the Brain \$5. Hamorrhoides, beneficial to confumptive and fcorbutic Habits 2020.

Liceup, a Symptom of an inflamed Liver 132.

Hernia Aquofa, from whence 327.

Holes. [Sec Foramina] 85. Fame us, Arteries of 272 Bone of 242 Cartilages of 248 Glands of 285 Ligaments of 250 Mucilaginous Glands of 285 Mufcles of 256 Nerves of 283 Veins of 279. Rumerus

Hamerus, Processes of, not to be luxated upwards 242 Groove of, in which lies the Tendon of the Biceps Muscle 244 Has greater Motion than any Bone in the Body 245.

Humours of the Eye 360. Hump Back. [See crooked Spine] 118.

Hydrocele. [See Hydrops Testis] 327.

Hydrops Testes, Cure of, ibid.

Hymen 338.

Hypochondria 2, 177.

Hypogastric Region 2, 178.

Faundice, Remarks on those who die of it 214. Yaw, Parts of 51 Upper Sinusses of. [See Antra]

Jejunum Intestinum 198.

Ifia, or Flanks 178. Ilium, Intestinum 108.

Impostbumations, Seat of 6 In the Ear, rather Ulcerations 364.

Incus. [See Os] 65.

Infants, Still-born, how discovered 150 Deformities, how sometimes occasioned 138 Heads, why open when born 59 Ribs often depressed or broke by careless Nurses 138.

Infundibulum 85.

Inflamed Liver, a Symptom of 131.

Inguina, or Groins 178. Inguinal Glands 325.

Injections of the Ear should be very gentle 371.

Intercostal Nerve 100, 102, 104, 106.

Internal Ear. [See Ear] 101 Jugular Vein the largest Vein that goes to the Head 98 Carotid Artery fends no Branches to the Neck 128.

Intestines 199 General Description of 202.

Intestines, Arteries of 203 Coats of 202 Glands of 203 Length of, ibid. Nerves of 204 Valves of 202 Veins of 203 Use of, ibid.

Intestinum Cæcum 199 Colon 200 Duodenum 197 Jejunum 198 Ilium, ibid. Rectum 201.

Integuments, external 2, Intexication, Cause of 95.

Joints, Liquor of 15 Diseases of, the Cause, ibid. Mucilaginous Glands of, ibid. Of the Thigh, imposshumated 182.

Ischiatic Nerve, the largest in the Body 40. Hibmis of the Medulla Oblongata 87.

Iter ad Infundibulum 86.

Jugular Veins 96.

K.

Kidneys, Description of 215 Vessels of 188 Use of, ibid. Labia

Lavia Pudendi 334: Labyrineh of the Ear 368. Lachrymal Glands 110, 355. Lactea Vala 44. Lacteal Veffels, ibid. Lacunar Glands 333, 338.

Larynn, Description of 122 Cartilages of, ibid. Membrane of 123 Muscles of, ibid. Ventricles of, ibid. Glands of, ibid.

Lef. Side, why most subject to pleuritic Pains 153.

Log, including all the lower Limb 285 Arteries of 314 Bones of 286 Cartilages of 292 Glands of 325 Ligaments of 287, 294 Mucilaginous Glands of 300 Muscles of 301 Nerves of 324 Veins of 279.

Ligament, what so called 15 Description of, ibid. Kinds of, ibid.

Senfible of very acute Pains, ibid. Use of 16.

Ligaments of the Abdomen 183 Arm, &c. 250 Carpus 254 Clavicles 251 Cubitus 252 Ears 70 Eyelids 68 Femoris 294
Fibula 298 Fingers 254 Head 66 Humerus 252 Leg 287,
294 Metacarpus 255 Metatarfus 299 Neck 117 Nofe 73 Patella 296 Pelvis 183 Ribs 139 Scapula 251 Sternum 139 Farfus 299.

Ligaments of the Thorax 139 Tibia 297 Toes 299 Vertebræ

Ligamenta lata Uteri 341 Rotunda Uteri, ibid.

Ligamentum Ciliare 68 La um, ibid. Sufpensorium 208 Teres, ibid. Fallopii 184.

Lunb, Circulation of the Blood in, after a Part is amputated 23 Lower, its Parts 285 Upper, its Parts 242.

Linea Alba Colli 120.

Lips, Arteries of 93 Muscles of 74 Nerves of 100 Veins of 96: Lieux, Description of 207 Arteries of 208 Nerves of, ibid, Veins of 209 Weight of 212 Excretory Ducks of 209.

Lobes of the Brain 85.

Lains, Vertebræ of 180 Muscles of 192. Longitudinal Sinus of the Dura Mater 52.

Lumbar Region 178.

Lungs 148 Achefions of common 151 Description of 140 Veffels of 100 Nerves of 104.

Luxition of the Neck, commonly called breaking the Neck 115.

Lymphatic Veins deferibed 41. Lympbædu 7s, what fo called, ibid. Their Ufe 42 Of the Teflie

tles 3290

M.

Malleas of the internal Ear. [See Os] 65. Mumma. [See Breafts] 174.

INDEX.

Mammary Vessels 176. Mammillary Process 53.

Man, why taller after lying some Time horizontally 117 Why shorter after carrying a heavy Burden on the Head or Shoulders,

Mirrow, what so called 9 Its Use 10 Blood Vessels of 9 Nerves of 10 Bloody 9 Spinal. [See Medulla Spinalis] 87.
Medulla. [See Marrow] 9.

Medulla Oblongata 87. Medulla Spinalis, ibid. Arteries of 88 Integuments of, ibid. Nerves of 39 Substance of 88 Thickness of, ibid. Use of 89 Veins of 88 Wounds of, mortal, ibid.

Mastication 76.

Maxillary Sinus 63 Imposthumations of, their Signs and Cure, ibid. Arteries of 93 Nerves of 37.

Meatus Auditorius 363 Obstruction of, the Cure 364 Urinarius 336.

Median Vein 31.

Mediastinum 147 Uses of 148. Meditulium. [See Diploe] 56. Megrim, Disease, Seat of 100.

Membrana Tympani, by fome called the Drum of the Ear 365 How to know if found 371.

Membrana Adiposa 5. Difeases of, ibid. Substance of, ibid. Use

Membrane, what fo called 48 Description of, ibid. Names of, ibid. Use of 49.

Membranes in General 48 Abdomen. [See Peritonzum] 192 Bones. [See Periosteum] 11, 256 Brain. [See Dura and Pia Mater 89, 90 Ears, Labyrinth of 368 Eye. [See Tunica] 357 Larynx 122 Oefophagus 159 Nofe, ibid. Thorax. [See Pleura] 146.

Meninges of the Brain 84.

Menses, Description of, how produced, and why periodical 341 Obstructions of, Symptoms and Causes 106 When they commence 341, 175 When they generally cease, ibid.

Mentula Muliebris 335

Menstrua. [See Menses] 341.

Mesentery, Description of 206 Use of, ibid.

Mesocolon 207.

Metacarpus 247 Arteries of 274 Bones of 255 Cartilages of 250 Ligaments of 254 Mucilaginous Glands of 256 Muscles of 264 Nerves of 267 Veins of 279.

Metata fus 291 Arteries of 316 Bones of 291 Cartilages of 292 Glands of 325 Ligaments of 299 Mucilaginous Glands of 300 Muscles of 308 Nerves of 323 Veins of 320.

Miliary Glands 5.

Milk in Womens Breafts, Caufe of 176. Miscarriage. [See Abortion] 348.

Mons Veneris 334. Morfus Diaboli 343.

Movien.

Mation, how defiroyed 18.

edini d

Mash, Parts of 51. Maciloginous Glands of the Joints 185 Of the Neck 108 Thorax, ibid. Abdomen 184 Arm 285 Of the Acetabulum Cox-

endicis 182 Leg 300. Muscle, what so called 16 Use of 17 Action of, ibid. Aponeu-sons of 16 Description of, ibid. Dimensions of, ibid. Fascia Tendinosa of 17 Figure of 16 Insertion of, ibid. Membrane of 17 Motion of, ib d. Name of 16 Origin, ibid. Penniform or Compound, ibid. Reclilineal or Simple, ibid. Sorts of ibid. Substance of 17 Tendon of 16 Use of 17 The longest in the human Body 305 Tail of 17.

Bufeles of the Abdomen 186 Arm 256 Ear 70 Epiglottis 124 Eye 68 Foo. 310 Fore-arm 256 Fingers 266 Glottis 122

Hand 256 Head 66 Larynx 122 Leg 301 Lips 74 Loins
192 Nofe 73 Neck 118 Palate 78 Pharynx 135 Private

Tarts 189 Shoulder 257 Thigh 301 Thorax 140 Toes 310

Tongue 31 Which move it on the Trunk 82. Mufewlus Abductor Oeuli 69 Abductor Digiti Interoffei 267 Digiti Me acarpi Offis Minimi 267 Digiti Minimi 268 Primi, ibid. Pollicis 270 Pedis 311 Pedis Digiti Minimi 313 Secundus, ibid. Pedis Interoffei, ibid. Transversalis 311 Acceterator Urinæ 190 Adductor Occuli 69 Anconæus 263 Anterior Auricu æ 71 Ancitragicus 72 Arytænoideus 124 Ba-Gogloffus Sr Biceps Cubiti Flexor 261 Tibiæ 306 Biventer 75 Biventer Cervicalis 119 Brachiæus externus 263 Internus 262 Buccinator 74 Caratoglossus 81 Ceratothyroideus 123 Chondroglossus 81 Ciliari 68 Cleidomastoideus 118 Coceygei 181 Compfexus 119 Conftrictor Pharingei 125 Coracobrachialis :61 Coracohyoideus 80 Complexus 1 9 Compreffor Narium 73 Corrugator Supercilii 68 Cremafter Tefbis 189 Cricoarytenoidens la cralis 124 Posticus, ibid. Crieothyroideus 123 Cruræus 308 Cucullaris 257 Depreffor Anguli Oris 74 Alæ Nafi, ibid. Labii Inferioris, ibid. Ocerli og Deftoides 259 Detrufor Urinæ 191 Diaphragma 144 Elevaror Anguli Oris 74 Alæ Nasi 73 Auriculæ 71 Ani Costarum 143 Labii inferioris 74 Superioris, ibid. Meni 74 Occuli 69 Palpebiæ 68 Scapulæ 258 Erector Chitosidis 191 Penis 189 Extenfor Auricularis, vel Digiti Mizimi 267 Digitorum communis, ibid. Digiti primi internodii Pollicis 268 Secundi 269 Tertii, ibid. Pollicis pedis brevis Trie ps Cubiti 263 Dorn et Lumberum 141 Ulnaris 265 Carpi radia is, ibid. Fascia Lata 304 Flexor Carpi radialis 263 Ulnaris, ibid. Digi orum internodii primi 266 Secundi, Ind. Tertii, ibid. Fl. xor primi et secundi Osis Policis 269 Termintermodii Pollicis 270 Pollicis pedis brevis 310 Longus, ibid. Digitorum pedis brevis, vel Perforatus 312 Longus, vel Perforans, ibid. Frontalis. [See Occipito] Gafterochemius 308 Gracilis 305 Genioglossus 81 Geniohyoideus 89 Glossotta-phylinus 79 Glutaus maximus 303 Medius, ibid. Minimus, ib d.

ibid. Hyothyroideus 123 Infraipmacus 200 Iliacus Internus 302 Intercostales externi, interni 143 Interspinales Cervicis 120 Dorsi, Lumborum 141 Intertransversales Cervicis 120 Dorsi, Lumborum 192 Latissimus Dorsi 261 Laxator Tympani 72 Lingualis 81 Longissimus Dorsi 261 Longus Colli 121 Lumbricales 266 Pedis 313 Mallei externus 73 Internus, ibid. Masseter 78 Mastoideus 119 Multifidus Spinæ 141 Mylohyoideus 80 Nafalis Labii, Superior 74 Occipito Frontalis 67 Obliquus externus, Abdominis 187 Inferior Oculi 70 Inferior Capitis 84 Internus Abdominis 137 Superior Occuli 70 Superior Capicis, ibid. Obturator externus 304 Internus, ibid. Oesophagus 159 Orbicularis Oris 191 Palpebrarum 68 Palato pharingeus 126 Palmaris brevis 264 Longus, ibid. Pectineus 302 Pectoralis 259 Perforans 266 Perforatus, ibid. Peroneus brevis, Longus 309 Plantaris 308 Platysma-myoides 75 Popliteus 307 Pronator quadra'us 265 Teres 266 Pfoas Magnus 302 Parvus 193 Pterygoideus externus 75 Internus, ibid. Pterygopharingeus 125 Pterygostaphylinus externus 79 Internus, ibid. Pyramidalis 187 Pyriformis 303 Quadratus Femoris 304 Lumborum 193 Restus Abdominis 187 Capitis internus Major 82, 120 Minor 83 Capitis Major Posticus, ibid. Minor Posticus, ibid. Rectus Capicus Obliquus inferior 83 Superior 84 Reclus Lateral.s 83 Tibiæ 306 Retractor Auriculæ 71 Rhomboides 257 Sacer 141 Sacro lumbalis, ibid. Sartorius 305 Scalenus 142 Semimembranofus 306 Semifpinalis Cervicis 121 Dorfi 401 Semitendinofus 306 . Serratus inferior politicus 143 Superior 143 Major Anticus 258 Minor, ibid. Sphincter Ani 191 Oris [See Orbicularis] 74 Vaginæ 191 Vesica Urinariæ 190 Solæus 308 Splcnius Capitis 120 Colli 121 Spinales Cervicis 121 Dorsi 141 Sternohyoideus 80 Sternomaffoideus 119 Sternothyroideis 123 Stylogloffus 81 Stylohyoideus 80 Stylopharingeus 125 Subclavius 257 Sublimis 266 Subfcapularis 261 Subfcapularis fuperior, ibid. Inferior, ibid. Supinator radii brevis 263 Longus, ibid. Supraspinatus 260 Temporalis 75 Tensor Tympani 73 Teres Major 260 Minor, ib. Thyroaryt znoideus 124 Thyrostaphylinus 79 Tib alis Anticus 309 Posticus, ibid. Trachelomasto deus 120 Tragicus 72 Trapezius 257 Trans-versalis Abdominis 188 Cervicis 121 Dorsi 141 Penis 199 Transversus 72 Triangularis Sterni 144 Triceps Extensor Cubiti 263 Femoris 302 Vaftus Externus 307 Internus, ib. Zygomaticus 74. Myrtiformes. [See Carunculæ] 334.

N.

Nails, what 50.

Nates Cerebri 87.

Navel String. [See Funiculus] 349.

Neck, Arteries of 110 Bones of 114 Cartilages of 116 Glands of 131 Ligaments of 117 Mucilaginous Glands of 118.

Neck, Muscles of 118 Nerves of 130 Veins of 128 Luxations of, commonly called breaking the Neck 117.

C c Nectritis.

INDEX.

Nephritis, a Symptom of 324.

Nerves, Consent of 34 Description of, ibid. Division of 36 Experiments of, prove the Existence of a nervous Fluid 35 Harmony and Sympa hy of, ibid. Origin of 36 Whether vibrating Cords or traductory Tubes 35.

Nerves of the Brain 91 ist Pair 36 2d Pair 37 3d Pair, ibid. 4th Pair, ibid. 5th Pair, ibid. 6th Pair, ibid. 7th Pair 38

8th Pair, ibid. 9th Pair, 39 10th Pair, ibid.

Nerves of the Medulla Spinalis 39 1st Cervical, ibid. 2d Cervical, ibid. 3d Cervical, ibid. 4th Cervical, ibid. 5th Cervical, ibid. 6th Cervical 40 7th Cervical, ibid. 12 Twelve Dorfal, ibid. 5 Five Lumbar, ibid.

Nerves of the Abdomen 239 Arm 284 Leg 323, Neck 130

Thorax 171.

Nerves of the Bowels 204 Eye-ball 361 Stomach 196 Teeth

Nerves, Au litory 38 Abducent 370 Crural 323 Gustatory 37 Intercostal 38 Lingual 39 Maxillary 37 Motary, ibid. Occipital 39 Olfactory 36 Opthalmic 37 Optic, ibid. Pathetic, ibid. Phrenic 130 Sacral 240 Sciatic 240, 372 Sympathetic 38 Of the oblique and extensor Muscles of the Head, ib.

Nose, 93 Arteries of 83, 94 Bones of 62 Cartilages of 74
Glands of 110 Ligaments of 74 Mucilaginous Glands of 66
Muscles of 73 Nerves of 37 Veins of 128.

Noffrils, Sinufes of 61.

Nutrition of the Foctus in Utero, Manner of 347.
Nympbæ, Deferipion of 334.

0.

Occiput, Arteries of 94. Oblique Muscles of the Eye 70.

Occipital Bone, why strongly made and defended 67.

Oesephagus or Gullet 159 Arteries of 160 Coats of 159 Glands of 161 Nerves of, ibid Substance of 159 Veins of 161 Pierces the Diaphragm 159.

Omentum or Cawl 205.

Optic Nerves, where they pass 56, 68.

Orbits of the Eyes, Veffels of, where they pass 56 Arteries of 95

Contents of 356.

Os Aftragalus 291 Calcis, ibid. Canini Dentes 64 Claviculæ
213 Coccygis 181, 193 Coffæ 137 Cribiforme 55 Cuhoides
290 Cunciforme 247 Cunciformia 290 Ethmoides 55 Femoris 286 Fibula 289 Frontis 52 Humeri 244 Hyoides 65
Incifores Dentes 64 Incus 65 Innominata 182 Ilium, ibid.
Ifchium 182 Lunare 247 dalæ 62 Malleus 65 Magnum
247 Maxilla, inferior 64 Superior 62 Metacarpi 247 Metatarfi 291 Molares Dentes 64 Nafi 61 Naviculare 291 Occipitis 54 Orbiculare Auriculæ 66 Carpi 247 Palati 63 Parictale 53 Patella 238 Petrofa 54 Plana 62 Phalanges Digiti Manus 248 Digiti Pedis 291 Pubis 182 Radius 246
Rotula 288 Sacrum 181 Scaphoides 241 Scapulæ 243
Sphe-

N D E X.

Sphenoides 55 Spongiofum 63 Stapes 66 Sta Alle 27
Temporum 53 Tibia 288 Trapezium 247 Trapezoides, Ma.
Ulna 245 Unciforme 247 Ungues 62 Vertebræ 110 Vomer 63.

Officula Auditus 85. Offification, how performed 8. Offeology, Foundation of all Anatomy 14 Table of, ibid.

Ovaria of Women, Description of 342.

P.

Pain of the Thigh in Pregnancy, from whence 341. Palate, Bones of 63 Muscles of 78.

Palpitation of the Heart, a Symptom of 130.

Pachomius's Glands 91. Pancreas, Description of 213 Use of, ibid. Weight of, ibid. Pancreatic Juice, Quantity fecreted 212.

Par Linguale 39 Vagum 38.

Parietal Bones the largest in the Skull 53 What makes the Treelike Appearance on their Infide 60.

Parotid Gland 107 Arteries of 93.

Parts of the Body 7. Patella, or Knee-pan. [See Os] 288 Cartilages of 292 Liga. ments of 296 Mucilaginous Glands of 300.

Pedis Hypocampi. [See Fornix] 86.

Pelvis, Bones of 179. Penis, Description of 33x Corpora Cavernosa of, ibid. Erection of, the Cause, ibid. Glands of, ibid. Præputium of, ibid.

Vessels of, ibid. Penis of Women. [See Clitoris] 335.

Pericardium, Description of 151 Adhesion of, ibid.

Pericranium, Description of 56. Peritonaum, Description of 193.

Perinæum 178.

Peritonæum, Dropfy of, how to discover 195. Perspiration insensible, Quantity of, secreted in a Day 5.

Pharynx 93, 159 Muscle of 99 Glands of, ibid.

Phrenic Nerve 130.

Pituitary Gland. [See Gland] 92.

Periofteum, what so called II Full of Blood Vessels and Nerves, ibid. Constitutes the first Rudiments in the Bones of the Fætus, ibid. Use of, ibid.

Pia Mater, Description of 89 Arteries of, ibid. Use of, ibid. Veins of, ibid. Almost wholly composed of Blood Vessels, and from whence, ibid.

Piles, how best extirpated 203.

Pleura, Description of 146 Uses of 147.

Pleuritic Pains, from whence they proceed 153.

Plexus Choroides 86.

Pinguedo. [See Fat] 6. Polypus behind the Uvula occasions Deafness 368.

Pomum Adami 122.

Pores

I N D E X.

Pores of the Cuticle 3. Portio Mollis et dura 103.

Pracessus Acromion 244 Ciliares 359 Coracoides 244 Dentatus 115 Vaginalis. [See Tunica] 326.

Processes of the Vertebræ 111.

Proffate Glands 330. Proftatæ, ibid.

Prolapsus Ani 202 Ca'es of, ibid.

Placenta, Description of 347 Separation of, occasions a Miscarriage 349.

Præputium 332.

Pudenda. [See Labia] 3.34.

Pulmonary Artery 22 Vein 27. Pulse, the Artery we feel when we feel the Pulse 278 Cause of 24. Pterygoid Processes 55.

Puneta Lachrymalia 62, 356. Pupil of the Eye 358.

Ramifications of the Arteries and Veins branched like the Stamina of the Leaves of Plants 22.

Receptaculum Chyli 45 Will contain about one Ounce of Water 46.

Restum Intestinum 261.

Recurrent Nerves cautionfly to be avoided in Bronchotomy 104 If both cut, the Voice would not be enti ely lost, as some affirm, ib. Renal Glands 216.

Renes Succenturiati 217. Regio Umbilicalis 2.

Reticulum Mucofun 2

Reticulum Mucofum 3. Retina Occuli. [See l'unica] 359.

Respiration, why affected in sneezing too.

Ribs, their Veffels and Nerves pass in a Furrow in the under Edge of each Rib 137 Cartilages of 158 Ligaments of 139 Of Infants, often depressed or broke by careles Nurses 138.

Rima Magna. [See Vulva] 334. Rotula. [See Patella] 288.

Saliva, how promoted 80 Secretion of, how performed 107 Quantity fecreted always according to the Drynefs of the Food 109 Quantity of 212.

Salival Ducts of Steno 107 Ulcer and Wounds of, the Cure, ibid. Glands of 108.

Salivation 109 These that are down in it sometimes spit a Gallon in twenty-four Hours, ibid.

a D

Solvatella Vein 31.

Scala Tympani 309 Vestibuli, ibid.

Scopela, Ribs of 243 Sinus of 244 Spine of, ibid.

Schirrous

I N D E X,

Schirrous Tonfils, Extirpation of 108. Sciatic Ligament 183 Nerve 240, 324 The largest in the Body, ibid.

Sclerotica Oculi. [See Tunica] 357.

Scrobiculus Cordis 2.

Scarf Skin. [See Cuticula] 2.

Scutiform Cartilage 122. Sebaceous Glands 355.

Seed, Secretion of 328 How paffed and ejected in Time of Coition

329 Description of 345. Sella Turcica, vel Equina 55.

Semen Masculinum. [See Seed] 345.

Seminal Vessels. [See Vafa Deferentia] 326.

Septum Lucidum Cerebri 85.

Skoulder. [See Humerus] 242 Top Pain of, in Suppurations of the Liver, &c. how occasioned 131.

Shoulder-blade. [See Scapula] 243.

Shoulder not to be diflocated upwards 244 The greatest Motion of any Joint in the Body 245 Cartilages of 249 Ligaments of 25I.

Sinus Frontalis 53.

Sinus Maxilla Superioris 63.

Sinus Cervicalis 130 Sphenoidalis 56 Vena Portæ 208.

Singing, Remarks on 123.

Skin, from whence its Colour 3.

Smelling, immediate Instrument of 99.

Skull, Diploe of 56 Foramina of 59 Imperfection of, in newborn Children 57 Lining of 90 Shape of 57 Sutures of, ibid. Tables of 56.

Seap Suds, a proper Injection for the Ear 371.

Spermatic Cord 328 Artery the smallest in the whole Body for its Length, ibid.

Spine, Bones of 112 Crooked 114, 135 Joints of, why fo many, 113 Luxations of, ibid.

Spirits. [See Animal Spirits] 35.

Spine Canal, how lined 38 Lined with a ligamentary Tube 117 Crooked, how formed 135.

Spiral Marrow 88 Its use, ibid. Processus 111.

Spleen, Description of 214 Weight of 215 Use of, ibid. Vessels of, ibid. Size of 214.

Sprain of the Ancle, a Caufe of 310.

Steno's Salival Duct 107.

Stomach, Description of 195 Coats of 196 Nerves of, ibid. Vef-

fels of, ibid. Stomach, very large in Gluttons, ibid. Senfation of, why very acute 104 Pain of, in Palpitations of the Heart, how occasioned

Stomach Ach in Obstructions of the Menses, the Cause 106.

Sternum. [See Os Sternum] 136 The Uncertainty of Trepanning

Squinting, involuntary, how occasioned 102.

Stra-

INDEX.

Strabifmut. [See Squinting] ibid.

Stapes. [See Os Stapes] 66.

Styliform Process 54. Sublingual Glands 108.

Suppuration, how performed 23 Of the Liver, and Symptoms of

Still Born Infants, how to discover 150.

Styloide Process 125.

Sutures of the Skull 57 How formed, ibid. Why open in newsborn Infants 59 Use of 58 True 57 Spurious, ibid.

Sutura Coronalis 58 Lambdoidalis, ibid. Sagittalis, ibid. Squamofa, ibid. Transversalis, ibid.

Swallowing, Action of, 122 Why we cannot when our Mouth is

Swelling. [See White Swelling] 15.

Sweat, what 5.

Sympathy of Nerves, the Knowledge of, of great Use in Physic 35 Of the Parts from whence 106.

Sympathetic Nerves. [See Par Vagum] 38. Synovia Liquor of the Joints 15.

Systele of the Heart 18.

T.

Table, Ofteology of 14.

Tables of the Skull 56.

Tarfus, Arteries of 290 Bones of, ibid. Cartilages of 68 Glands of 294, 325 Ligaments of 68 Mucilaginous Glands of 300 Muscles of 308 Nerves of 323 Veins of 322.

Ta fi Cartilage of the Eye lids 68.

Teylor's Muscle 305. Tears, Matter of 355.

Teeth 64 Names of, ibid. Nerves of 65 Number of 64 Veffels of, where they pass, ibid. Of the upper Jaw, why their Roots spread more than those of the under Jaw 65 Nutrition of, ibid. Sensation of, ibid. Arteries of 93.

Tendons, what fo called 18.

Testes Cerebri 86 Or Testicles 325. Thatami Nervorum Opticorum 86.

Thigh, Arteries of 314 Bone of 286 Cartilages of 292 Glands of 325 Ligaments of 294 Mucilagineus Glands of 300 Muscles of 301 Nerves of 323 Veins of 320.

Thigh, Numbuefs of, in a Nephritis, the Cause 324 Pain of in Pregnancy, the Cause 341.

Tho acic Duct 46.

Thorax 132 Arteries of 161 Bones of 133 Cartilages of 138
Glands of 172 Ligaments of 139 Muc laginous Glands of 138
Mu'cles of 141 Neives of 171 Veins of 166 Viscera of 145.

Thumb. [see Pollex] 248.

Thymus Gland 173 Largest in new-born Infants, ibid. Not to be found in old People, ibid.

Thyroide Glands 132.

N D E X.

Tobacco, Smoaking of, promotes spitting in those not accustomed

Tous, Arteries of 316 Bones of 291 Cartilages of ibid. Glands to it 109. of 300 Ligaments of 298 Mucilaginous Glands of 300 Mus-

cles of 310 Nerves of 324 Veins of 320. Trachea, Description of 157 Cartilaginous Rings of, ibid. Glands

of, ibid. Membrane of 158 Use of, ibid. Tongue, Bones of 65 Muscles of 81 Nerves of 105 Vessels of

93, 98. Tonfils Schirrous, Extirpation of 108.

Tooth Ach, by the Sympathy of the Nerves, may affect the Eye and Ear 37 A remarkable Cure of 102.

Trepan, or Trephine, not to be applied to the Forehead 53.

True Skin. [See Skin] 3. Tuba Eustachiana 367 Passes from the Mouth to the Ear 368 Obstruction of, causes Desfness, ibid.

Tubæ Fallopiana 343 Eustachianæ 367.

Tuberculum Loweri 153. Tunica, Albuginea Occuli 357 Adnata Occuli, ibid. Choroides Occuli 358 Conjunctiva Occuli 357 Cornea Occuli, ibid. Retina Occuli 359 Sclerotica Occuli 357 Uvea Occuli 358.

Tunica Vaginalis 326. Tunics of the Eye 355.

Tympanum, or Drum of the Ear 365 Not really the Organ of Hearing 370 Destroyed, Instances of, without loss of Hearing, ibid. Always full of Mucus in new-born Children 365 Membrane of, how to know when found 371.

V.

Vagina, Description of 337. Valves of Veins 27. Lacteal Veffels 44 Lymphatic Veffels 41.

Valvula Magna Cerebri 86. Vas, Brevia Arteria 235 Vena 239 Deferentia 326 Laclea 45 Præparantia 326.

Vein, what fo called 25.

Vein', Quantity of Blood in 23 Described 25 Their Ufe, ibid. Double the Number of the Arteries 26 Why fullest when warm, ibid. Subcutaneous, not in two Perfons alike 22 Accompany the Arteries to return the Blood to the Heart 23 How dilated 24 Valves of, ibid.

Veins of the Abdomen 230 Arm 279 Brain 91 Ears 93 Eyes 361 Head 96 Leg 319 Neck 128 Nofe 129 Thorax 166. Vein, Basi ic 282 Cephalic 280 Jugular 128 Median 281 Ver-

Vena Cava 28 Afcendens 31 Descendens 28 Portæ 33 Pul-monaris 34 Adiposæ 235 Axillaris 279 Azygos 166 Basilica 282 Bronchiales 164 Capfulares 235 Cephalica 280 Cæcalis 233 Coronariæ Cordis 156 Ventriculi 232 Cruralis 319 Cysticæ Gemellæ 231 Diaphragmaticæ inferiores, ibid. Superiores 169 Duodenalis 232 Epigastrica 237 Epiploica dextra 232 Sinistia 235 Galtrica dextra 221 Epiploica 235 Sinistra CC4

Sinistra 234 Hæmorrhoidales externæ 239 Interna 233 He-paticæ 231 Hypoga Irica 238 Iliacæ 237 Iliaca externa, ib. Interna, ibid. Intercostales 167 Intestinalis 232 Jugularis externa 128 Anterior 97 Posterior, ibid. Interna 129 Lum-bares 236 Mammaria externa, ibid. Interna 169 Mediana 281 Mediastinæ 168 Mesaraica 232 Obturarrix 238 Pancreaticæ 124 Pericardiæ 169 Peronea 323 Phrenicæ 171 Plantaris 323 Pylorica 231 Poplitea 322 Renales 235 Sacra 236 Saphena 320 Minor 321 Sciatica, ibid. Spermaticæ 235 Splenica 233 Subclaviæ 170 Suralis 323 Thoracicæ 166 Thymicæ 169 Tibialis anterior 322 Posterior 323 Trachealis 170 Vafa Brevia 235 Vertebralis 129.

Ventricles of the Heart, how filled and emptied 23 Of the Brain 850

Ventriculus. [See Stomach] 195.

Vertebræ of the Loins III Neck, ibid. Thorax 133.

Vertebral Arteries 129 Nerves 130 Veins 129.

Vesica Fellis. [See Gall Bladder] 209 Urinaria. [See Bladder] 218.

Veficulæ Seminales 329. Vestibulum of the Ear 368.

Virginity, Loss of, how to know, tho' not infallible 176.

Vomiting in a Nephritis, the Caufe 106.

Viscera of the Abdomen 195 Head 83 Thorax 145.

Vitreous Humour of the Eye 361.

Voice, loft, or greatly impaired by cutting the recurrent Nerves 104 Harmony of 123.

Vulva. [See Generation] 334.

Umbilical Region 177 Cord. [See Funiculus] 349.
Ulna. [See Os Ulna] 245.

Urachus. [See Arteria Umbilicalis] 227.

Uterus, Description of 217 Large in Nephritic Patients and Drunkards 218.

Ureibra, Description of 332 Women's shorter than Men's 333 Caruncles of, ibid. Glands of, ibid. Ulcers of 334.

Urine, Suppression of, the Cause 218.

Uterus, Description of 339 Cervix of, ibid. Distension of, ibid. Fundus of, ibid. Vessels of 340. Uvea Oculi, [See Tunica] 357.

W.

White Swelling, Caufe and Cure of 15. Whores that are common, why they do not conceive 344. Windpipe. [See Trachea] 157. Womb. [See Uterus] 339. Wemen, Circumcifion of 335 Genital Parts of 334 Fat, why barren 345.

Xir heide Carrilage. [See Ensiform Cartilage] 136.

EXPLANATION of the FIGURES in the following TABLES.

TABLEI

S Frontis

Sutura Coronalis

3 Os Verticis

4 Sutura Squamofa

5 Os Temporis

6 Processus Mammillaris

7 Os Mala 8 Osfa Nasi

9 Ossa Maxillaris superiora 10 Os Maxillæ inferioris

11 Vertebræ Colli

12 Vertebræ Lumborum

13 Os Sacrum

14 Sternum

14 Sternum 15 Scapula 16 Costæ Veræ

17 Costæ Nothæ 18 Clavicula

19 Processus Coracoideus

20 Os Humeri

21 Ulna

22 Radius

23 Os Ilium

25 Ifchium
26 Os Pubis
27 Foramen Magnum
28 Os Femoris
29 Trochanter Mai

28 Os Femoris 29 Trochanter Major 30 Trochanter Minor

31 Patella

32 Tibia

33 Fibula

34 Talus

35 Os Calcaneum

36 Offa Tarfi

TABLE

r Os Parietalia

2 Sutura Sagittalis

3 Sutura Lambdoidalis

4 Os Occipitis

5 Sutura Squamofa 6 Maxilla Inferior

7 Vertebræ Golli

8 - Dorsi

9 --- Lumborum

30 Os Sacrum

TI Os Coccygis

+ Costæ Veræ, No. 7

Coffæ Nothæ, No. 5 H

12 Clavicula

17 Scapula

14 Spina Scapulæ

x 5 Acromion

16 Os Humeri

17 Ulna

18 Radius

19 Offa Carpi

20 Offa Metacarpi 21 Offa Digitorum

22 Ilium

23 Ischium

23 Ischium
24 Os Femoris
25 Collum Ossis Femoris
26 Trochanter Major
27 Trochanter Minor
28 Condylus exterior Ossis Femoris
29 Condylus interior Ossis Femoris
30 Tibia
31 Fibula

31 Fibula

32 Os Calcaneum

33 Offæ Tarfi

34 Offæ Metatarfi

A B L E III. T

I Frontales

2 Orbicularis Palpebræ

3 Zygomaticus Major

4 Nafales Labii fuperior

5 Depreffor Labi inferior

6 Depreffor anguli oris

7 Platifma myoides

8 Pectoralis

9 Latiffimus dorfi

To Serratus magnus

Externus obliquus abdominis

12 Recti abdom nis

13 Pyramidales
14 Linea alba

15 Gracilis

36 Adductor longus tricipitis femoris

17 Pectineus

18 Pfoas magnus

19 lliacus internus

20 Sartorius

21 Glutæus medius

22 Fascialis

20 12

23 Vaftus extermus

TABLE III.

- 24 Rectus femoris
- 25 Vastus internus
- 26 Pars bicipitis
- 27 Pars gastrocnemii
- 28 Soleus
- 30 Extensor longus digitorum pedis
 31 Tibialis anticus
 32 Deltoides
 33 Triceps
 34 Biceps

- 34 Biceps
 35 Brachiæus externus
- 36 Supinator longus
- 37 Pronator rotundi radii
- 38 Radialis internus
- 39 Palmaris longus
- 40 Sublimis
- 41 Ulnaris internus
- 40 Sublimis
 41 Ulnaris internus
 42 Abductor longus pollicis
 43 Radialis externus longus

TABLE IV.

- 1 Temporalis
- 2 Mastoidæus
- 3 Trapezius
- 4 Deltoides
- 5 Brachiæus
- 6 Gemellus
- 7 Palmaris longus
- 8 Sublimis
- 9 Ulnaris internus
- 10 Radialis externus longior
 11 Extenfor communis digitorum
 12 Infra spinatus
 13 Latissimus dorsi

- 74 Obliquus externus abdominis 15 Glutæus medius 16 Glutæus major
- 16 Glutæus major
- 17 Gracilis
- 18 Adductor magnus femoris
 19 Semtiendinofus
 20 Biceps Cruris

- 21 Vastus externus
- 22 Gastrocnemius
- 23 Soleus
- 24 Tendo Achillis

ABLE

- 1 Occipitalis
- 2 Attollens auriculam
- 3 Orbiculares palpebrarum
- 4 Latissimus colli

TABLE V.

Mastoidaus 6 Trapezius 7 Deltoides 8 Biceps 9 Brachialis internus To Triceps II Surmator longus 12 Ralialis internus 23 Radialis externus longior 14 Radialis externus brevior 15 Ilnaris externus Abductor pollicis longus manus 27 Infraspinatus 18 Teres minor 19 Teres major 20 Latissimus dorsi 21 Pectoralis 22 Serratus magnus 23 Obliquus externus abdominis
24 Tenfor vaginæ femoris 25 Glutæus medius 26 Glutæus magnus 27 Semitendinofus 28 Biceps cruris 20 Vaftus externus 30 Rectus Cruris 31 Gastroenemius 32 Soleus 33 Tendo Achillis 34 Peroneus longus 35 Peroneus brevis 36 Extensor longus digitorum pedis 37 Tibialis anticus 38 Ligamentum à patella ad tibiam pertinens 39 Vastus Internus Bush then Blico I se 40 Sartorius 42 Triceps parfquæ longus vocatur 43 Triceps pars quæ brachialis externus vocatur 44 Brachialis internus

AS Biceps brachii 46 Pronator teres 47 Palma is longus 48 Sublimis 49 Ulnaris internus 50 Ulnaris externus

TABLE VI. FIGURE I.

- I The larynx
- 2 The internal jugular vein
- 7 The fubclavian vein
- 4 The vena cava defeenders
- The right auricle of the heart

TABLE VI. FIGURE I.

- 6 The right ventricle
- 7 Part of the left ventricle
- 8 The aorta afcendens
- 9 The arteria pulmonalis
- The right lobe of the lungs, part of which is cut off to shew the great blood veffels II The left lobe of the lungs
- 12 The diaphragm
- 13 The liver
- 14 The ligamentum rotundum
- The gall bladder
 The stomach pressed by the liver toward the left side
- 17 The fmall guts
- 18 The spleen

FIGURE II.

- 2 The right ventricle of a fœtus distended with wax
- 2 The right auricle
- 3 The left auricle
- 4 Branches of the pulmonary veins of the right lobe of the lungs
- 5 The arteries of the left lobe of the lungs
 6 The vena gava descendens
- 7 The aorta ascendens
- 8 The arteria pulmonalis
- 9 The ductus arteriosus

TABLE VII. FIGURE I.

- A. The upper orifice of the stomach B. The stomach
 C. The pylorus

- D D. Arteries
- E F. Nerves which accompany the arteries
- G. The duodenum

- H. H. I. I. The fmall guts
 K. The valve in the colon
 L. The appendicle of the cocum
- M. N. The colon
- O. The rectum
- P. The constrictor of the anus
- Q. Q. The lifters up of the anus R. The anus.

FIGURE II.

- A. A kidney divested of its external coat

 B. A kidney in its natural state
- C. The vena cava
- D. The aorta
- E. E. The renal glands with their vessels, &co, F. The emulgent vessels
- G. G. The ureters
- H. The urinary bladder
- 1. The neck of the bladder

TABLE VII. FIGURE II.

K. L. The testicles

M. The process of the peritoneum (in which the spermatic vessels go) cut off alisamenta pulmonsiis

N. The cremaster muscle cut off

O. O. The spermatic vessels

Q. The Epididymis

R. R. The vafa deferrentia S. The corpus glandofum

T. T. The two bodies which compose the penis, and appear when the skin t. t. is drawn aside

U. U. The prepuce

V. The glans penis W. The extraordinary infertion of the spermatic vein into the emulgent X. X. Vesiculæ seminales

Y. Y. The infertion of the ureters Z. The beginning of the urethra

r. r. Veins which run and unite on the back of the yard

s. s. Arteries with nerves on each fide v. Veins opened to shew their valves

FIGURE III.

A. A. A. The parenchymous substance of the pancreas laid

B. The pancreatic duct with its branches C. C. C. C. C. C. C.

D. The bile duct joining the pancreatic duct

E. The duodenum opened

F. The orifice of the bile and the pancreatic ducts

A B L E VIII.

I Aorta

A. Valvulæ femilunares

2 Arteria coronaria mag.

3 Ligamentum arteriofum

4 Arteriæ fubclaviæ

5 --- carotides

___ vertebrales 7 --- temporales

8 ____ occipitales

o Contorsiones carotides

C. Glandula pitu taria

D. Arteriæ opthalmicæ

10 Contorsiones vertebrales

II Ramificationes arter &

12 Arteriæ mammariæ 13 —— cubitales

14 Arteria aorta descendens

15 --- bronchia'is

16 Arteriæ intercostales

17 Arteria coeliaca

18 Arteria hæpaticæ

TABLE VIII.

	The second secon
19 Arteria cystica	allahmilana mus 2 3
20 coronaria inferiora stomachi	sinor so a
21 — pylorica	aissagad g
22 — epiploica	kingspering
23 coronaria superiora stomachi	erroumso
24 Arteriæ phrenicæ	so Procedus madoldens
25 Arteria splenica	Taurenthus susself ex
26 mesenterica superior	Procedur Historias
27 — inferior	sitabila Ot the
28 Arteriæ emulgentes	zabionspile O at
29 vertebrales lumborum	The Tag - Et
30 fpermeticæ	1 71381 01
31 Arteria facra	17 - en nis
32 Arteriæ iliacæ	munity - 8s
33 ——— externæ	rotten be el Bull or
34 internæ	
35 — umbilicales	
36 —— epigastricæ	2 2' Mexille inferior
37 ——— penis	2 Procedus canadis
38 ——— crurales	ag ge
30	esm to 7 a c
TABLE	Ze Denier berleit.XI
1 The brain	inium
2 The cerebellum	ancion as
3 The corpus pyramidalis	zo Chariquerum
4 The annular protuberance	oo Foranen
5 Processus mammillaris	
6 Optic nerves	
7 Motores oculorum	
8 The fourth pair of nerves	
9 The fifth pair spreading into three branches	
The fixth pair	
11 The seventh pair	
12 The eighth pair	
13 The recurrent nerves joined with the eighth pair	
14 The recurrent nerves after leaving the eighth pair	
15 The trunks of the eighth pair	
16 Intercostal nerves	
17 Phrenic nerves	
18 Branches of nerves going to the spermatic vessels, testicles,	
uterus, &c.	
19 Branches of the 9th pair	
20 The fciatic and caural nerves	
21 The brachial nerves	
+ Communications between the dorfal	and the intercoffal nerves
+ Communications between the contai	and the intercontar nerves
TABLE X.	
	The state of the s
r Sutura coronalis	
2 — fagittalis	
3 —— lambdoidalis	
4 —— fquamofa	5 Sutura
	J Ductura

TABLE X

softwa simula or

Sutura transversalis 6 Os frontis 7 - bregmatis 8 — occipitis g - temporis to Processus mastoidæus zz Meatus auditorius 12 Processus styliformis 13 --- jugalis 14 Os sphænoides 15 - mali 16 - nasi 17 - unguis 18 - plenum 19 Ductus ad nasum 20 Maxilla fuperior
21 Foramen maxillæ fuperioris
22 Maxilla inferior
31 Proceffus coronalis 21 Processus coronalis 24 - condyloides 25 Foramen 26 Dentes inciforii 27 ---- canini 28 --- molares 29 Os triquetrum 30 Foramen

the Clare to the follower to the formette vellely

alegampl ---- 6

