

**Observations on the epidemical diseases in Minorca, from the year 1744 to 1749 : to which is prefixed a short account of the climate, productions, inhabitants, and endemial distempers of that island / by George Cleghorn.**

**Contributors**

Cleghorn, George, 1716-1789.

**Publication/Creation**

London : printed for T. Cadell and G. Robinson, 1779.

**Persistent URL**

<https://wellcomecollection.org/works/gveschbj>

**License and attribution**

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>





Royal Ordnance Medical Library,

JANUARY 10, 1815.

*The Time allowed for keeping each Book.*

A Folio - - - - -	3	} Weeks.
A Quarto or Octavo - - - - -	2	
Smaller Books - - - - -	1	

N. B. If the Volume is not wanted by any other Subscriber, the Period may be renewed.

*The undermentioned Fines are imposed for keeping Books beyond the Time specified.*


A Folio - - - - -	3d.	} each Day.
A Quarto or Octavo - - - - -	2d.	
Smaller Books - - - - -	1d.	

TRO  
Strongroom  
RAMC  
GH.  
/CLE

22101945954



B.  
45 a 34 - 8




OBSERVATIONS

ON THE

EPIDEMICAL DISEASES

IN

MINORCA.







Faint, illegible text or markings below the stamp.

OBSERVATIONS

ON THE

EPIDEMICAL DISEASES

IN

MINORCA

Faint, illegible text at the bottom of the page.

OBSERVATIONS  
ON THE  
EPIDEMICAL DISEASES  
IN  
MINORCA.



From the Year 1744, to 1749.

TO WHICH IS PREFIXED,

A Short Account of the CLIMATE, PRODUCTIONS, INHABITANTS, and ENDEMIAL DISTEMPERS, of that Island.



---

BY GEORGE CLEGHORN,  
Lecturer of Anatomy in the University of DUBLIN,  
formerly Surgeon to the Twenty-second Regiment  
of Foot.

---

FOURTH EDITION.

---



---

L O N D O N:  
Printed for T. CADELL, in the Strand; and  
G. ROBINSON, in Pater-noster Row.

MDCCLXXIX.



OBSERVATIONS

ON THE

EPIDEMICAL DISEASES

IN

MINORCA

From the Year 1744 to 1749.

to which is added

A Short Account of the Climate, Population, Inhabitants, and Principal Diseases of the Island.

By GEORGE CLEBORNE,

Professor of Anatomy in the University of Durham, formerly Surgeon to the Tercio of the Kingdom of Castile.

FOURTH EDITION.

Printed by W. Johnston, at the Press of the University of Durham, in the Year 1754.

LONDON,

Printed by T. Cadell, in the Strand, and C. Richardson, in Pall-mall.

1754.





TO THE  
Society of SURGEONS  
OF HIS  
MAJESTY'S Royal Navy.

GENTLEMEN,

AS many of you must be sensible, how little the best Information we can acquire in this temperate Climate, qualifies us for treating the Diseases which are frequent in warmer Latitudes,

A 3 with



with the desired Success, I take the Liberty to address the following Sheets to your Society.

They contain, 'tis true, an Account of the Diseases only of a small, remote Part of the *British* Dominions; but of a Part in which Numbers of his Majesty's Subjects, beside the Natives and those employed in the Protection of the Place, are often brought together, both in Time of Peace and War: And as the Qualities of the Air, and the Course of the Seasons in *Minorca* correspond nearly with those in several other Parts of the World, to which our  
Fleets

## DEDICATION.

vii

Fleets frequently repair, it is probable the Diseases may likewise be similar.

Would all who practise Physick in our Factories and Colonies abroad embrace the Opportunity their Situation affords, to make proper Observations on the Sick, and communicate them to the Publick, we should soon have a more exact and ample History of Diseases, than we are yet possessed of; and future Practitioners would be enabled to shun the Dangers into which many have fallen, and to conduct those committed to their Care, through

A 4

the



the Disorders they are exposed to, with Satisfaction and Honour to themselves, and no small Benefit to their Country. It is therefore with great Pleasure, that I see this is particularly recommended, in the Plan for publishing Medical Observations, which you have lately established, and will, it is hoped, prosecute with the Vigour so useful an Institution deserves.

For my own Part, I must confess, I had not been long in *Minorca* 'ere I had great Reason to wish that some of the Practitioners who had been there before me, and who must have seen how widely the predominant Distempers in  
this



this Island differ from those in *England*, had been at the Pains to furnish their Successors with some Hints, some Observations, by which the fatal Consequences frequently attending these Diseases, might have been timely foreseen, or happily prevented.

Being therefore fully convinced, that some Remarks of this Nature might be useful to those who should afterwards practise in this Island, I determined to observe and record, with the utmost Care and Impartiality, whatever should appear conducive to a thorough Knowledge of its Diseases and their Cure; imagining that,  
next



next to the immediate Care of the Sick, this would be the most essential Service, which one in my Station could render to the Public.

With this View, in the Year 1743, I began to keep a Diary of the Weather, to remark the Course of the Seasons, to describe the Diseases they produced; and that commonly in the Chambers of the Sick: Which Diary I continued, with no small Labour and Assiduity, in the midst of an extensive Practice, both among the *English* and *Natives*, till the Year 1749; when the Removal of the Regiment, in which I have the  
Honour

Honour to serve, obliged me to leave the Island, and furnished me with Leisure for revising my Observations, and collecting, from a vast Multitude of Cases, such general Remarks as appeared worthy to be communicated to the Public.

You will observe that amongst the Epidemicks of *Minorca*, Tertian Fevers make the most considerable Part. The Diversity of their Types, the Violence of their Symptoms, their fallacious Intermissions, their sudden, and too frequently pernicious Events, rendered it necessary to give an explicit



explicit Account of them ; and the more so, as they seldom appear in this Manner in the northern Parts of *Europe* ; though in *Greece*, in *Italy*, and the adjacent Countries, it is evident from the Remains of Antiquity\*, and the Writings of the most judicious Moderns †, that they ever have been, and still are, very frequent, and that with a surprizing Constancy and Uniformity of Symptoms; though to one who

\* Hippocrat. Aphor. § iii. No. 21, and De Morb. Vulg. Lib. vii. Asclepiad. apud Cæl. Aurel. de Morb. acut. Lib. ii. Cap. x. Galen de Morb. temp. sub finem.

† River. Lib. xvii. §. iii. Cap. i. Lancif. Epid. Torti Therapeut. Special. Bianch. Hist. hepat. p. 3, &c.

never



never had an Opportunity of observing them in all Shapes, nor seen them amply described, they have the Appearance of great Irregularity and Confusion.

And it is more than propable, from the Accounts of several Physicians\* and Travellers, that epidemical Tertians are not wholly confined to the Coasts and Islands of the *Mediterranean*; but that they are equally frequent and destructive in many other Parts of the Globe, and perhaps may be deemed the anniversary autum-

\* Spigel. de Semitertian. Lib. iii. Cap. i. Tenent on the Diseases of Virginia, p. 12. Warren on the Fever of Barbadoes, p. 70.



nal Distempers of most hot Countries in the World.

A Tertian, it is true, when once discovered, may for the most part be readily cured, as we have so safe and efficacious a Remedy as the *Bark* in our Hands: But in warm Climates, such is the rapid Progress of this Distemper, that it is requisite we should know it in the Beginning, that no Opportunity may be lost of giving the Remedy in a sufficient Quantity to avert those Dangers into which the Sick are otherwise very soon precipitated: Yet the Variableness of its Aspect, and its frequently personating other acute Diseases, render



render this often difficult to the most experienced; and much more so to those who have seldom or never seen such Tertians, as they are only now and then to be met with in *England*.

These Considerations induce me to believe, that the Account of them contained in the following Pages, will neither be unacceptable nor useless to many of your Society; and particularly to those, who, by their Station in his Majesty's Service, are often obliged to take Care of Numbers of their Fellow Subjects, in Climates exposed to such Disorders; whilst at the same Time their  
quick




quick Transition from one Place to another, prevents their acquiring a competent Knowledge of the various Epidemicks from their own Observation.

To you, therefore, Gentlemen, I address these Remarks with the utmost Deference and Esteem, being in Hopes, that the Motives which prevailed with me to offer them to you and the Public, will sufficiently plead my Apology to both, for any Imperfections that may appear in this Performance, with regard to Method or Expression. I am,

*Your most humble Servant,*

LONDON, May 1.  
1751.

G. CLEGHORN.



# C O N T E N T S.

## INTRODUCTION.

<b>T</b> HE <i>Air in Minorca</i>	Page 2
<i>The Seasons of the Year</i>	ibid.
<i>Heat and Cold</i> - - -	3
<i>Winds</i> - - -	4
<i>Rain, and other Meteors</i> -	7
<i>The Soil and its Productions</i> -	12
<i>Corn</i> - - -	13
<i>Wine</i> - - -	14
<i>Pulse</i> - - -	16
<i>Roots, Sallads, and Pot-herbs</i> -	17
<i>Fruits and Trees cultivated</i> -	20
<hr/> <i>Indigenous</i> -	36
a	<i>Minerals</i>



<i>Minerals</i>	-	-	-	Page 46
<i>Fish</i>	-	-	-	47
<i>Cattle</i>	-	-	-	49
<i>Wild Fowl</i>	-	-	-	54
<i>Water Fowl</i>	-	-	-	55
<i>Birds of Passage</i>	-	-	-	56
<i>Eatable Snails</i>	-	-	-	58
<i>The natural temper of the Minorquines</i>				59
<i>Their Diet</i>	-	-	-	61
<i>Their Festivals and Publick Diversions</i>				64
<i>Their ancient Usages</i>	-	-	-	68
<i>Endemial Diseases</i>	-	-	-	74
<i>Obstructions of the Abdominal Viscera</i>				75
<i>Hæmorrhoids</i>	-	-	-	78
<i>Ulcers of the Legs</i>	-	-	-	ibid.
<i>Ruptures</i>	-	-	-	80
<i>Sore Eyes</i>	-	-	-	ibid.
<i>Nepbritick Pains</i>	-	-	-	81
<i>Convulsion of the Jaw in Children</i>				ibid.
<i>Accidents from endemial Causes</i>				83
<i>Their Notions of Witchcraft and evil Spirits</i>	-	-	-	84

## C H A P. I.

<i>The Weather described from the Year 1744</i>	
<i>to 1749</i> - - -	Page 87
<i>A Thermometrical Table</i> -	130
<i>A short Account of the epidemical Diseases</i>	
<i>from 1744 to 1749</i> -	131

## C H A P. II.

<i>Tertian Fevers, why so called</i>	147
<i>The Diversity of Tertians occasioned by the</i>	
<i>different Types of their Periods, viz.</i>	
<i>Simple Tertians true and spurious</i>	152
<i>Double Tertians</i> - - -	ibid.
<i>True, Spurious, and Subintrant Double</i>	
<i>Tertians</i> - - -	154
<i>Triple Tertians</i> - - -	ibid.
<i>Semitertians</i> - - -	155
<i>Intermittent, Remittent, and Continual</i>	
<i>Tertians</i> - - -	156
<i>Anticipating and postponing Tertians</i>	157



<i>The Diversity of Tertians occasioned by the different Symptoms of their Paroxysms,</i>	Page 158
<i>Lipyrria, Febris Syncopalis, Affodes, E- lodes, &amp;c.</i>	160
<i>The Constancy of Nature in the Course of Tertians - - -</i>	162
<i>Their most frequent Form in their Be- ginning - - -</i>	163
<i>In their Progress - -</i>	164
<i>In their State and Declension</i>	166
<i>Signs of Security in Tertians -</i>	170
<i>of Danger - -</i>	171
<i>of Death, and when to be expected</i>	175
<i>Appearances on opening the Bodies of the Dead</i>	180
<i>What Species of Tertians are most common in Minorca - - -</i>	184
<i>Management of the Non-naturals in the Cure of Tertians - -</i>	186
<i>Of Bleeding - -</i>	196
<i>Of Vomiting and Purging -</i>	201
<i>When Tertians may be left to Nature</i>	205
<i>When</i>	



C O N T E N T S.

xxi

<i>When they require the Bark</i>	Page 206
<i>When Purgatives should be joined with the Bark</i>	- - - 211
<i>When Cordials</i>	- - - 212
<i>Prejudices against the Bark considered</i>	215
<i>Its Uses recommended in several Cases, where forbid by Authors</i>	- 221
<i>The Cure of the Symptoms in Tertians</i>	226
<i>Acute Pains in the Back and Limbs</i>	ibid.
<i>Vomiting and Purging</i>	- 227
<i>Restlessness and Head-achs</i>	- ibid.
<i>Hæmorrhages from the Nose</i>	228
<i>Pain and Inflammation in the lower Belly</i>	ibid.
<i>Coma, and Hiccup</i>	229
<i>Abscesses near the Hip</i>	- ibid.
<i>Parotids</i>	- - - 230
<i>Of Relapses of Tertians, their Prevention and Cure</i>	231
<i>Of Sporadick Tertians</i>	- - 234
<i>The frequency of epidemical Tertians in many Parts of the Globe</i>	235
	<i>The</i>



*The Bark recommended by the best Practitioners of different Nations* Page 236

## C H A P. IV.

*Of the Sudamina, or Rash* - 238  
*Of the Effere, called Favas in Minorca* 240  
*Of the Cholera Morbus* - 243

## C H A P. V.

*In what Manner Dysenteries begin* 244  
*How they terminate, and the Appearances in such as die of them* - 246  
*Ipecacuanha & Vitrum Antimonii recommended in some Dysenteries* - 249  
*The antiphlogistick Method in others* 251  
*Hardened Excrements commonly collected in the Guts, when the Ilium, Colon, or Rectum are inflamed* - 252  
*Calomel with Opium, the best Remedy to dislodge them* - - - ibid.  
*Of old inveterate Dysenteries* 254  
*Opium of great Service in them* ibid.

*In*



*In what Sort of Dysenteries the Bark is  
successful* - - Page 256

## C H A P. VI.


<i>General Remarks on the Summer and Win- ter Fevers</i> - - -	257
<i>Of the Pleurisy in the Year 1746</i>	261
<i>Its treacherous Remissions</i> -	266
<i>On what Days the Patients died, and the Appearances in the dead Bodies</i>	267
<i>A Case of an Abscess in the Liver</i>	269
<i>The natural Evacuations which terminate Pleurisies</i> - - -	270
<i>The capital Remedies in the Cure of Pleurisies</i> - - -	276
<i>Other collateral Helps</i> - - -	282
<i>Of the inflammatory Diseases, epidemical at the same Time with the Pleurisy</i>	285
<i>Extracts from Letters concerning the Pleurisy</i> - - -	286



## C H A P. VII.

<i>Of the Small-Pox in the Year 1742</i>	292
<i>Of the Small-Pox in 1746</i>	- 296
<i>Their pestilential Disposition</i>	- 297
<i>Two Cases cured by the early Use of the antiphlogistick Method</i>	- 305
<i>Three, wherein this Method failed</i>	306
<i>The Danger of the natural Small-Pox, and the Safety of Inoculation</i>	- 310





[ 1 ]

---

---

T H E

## INTRODUCTION.

WHEN I determined to write upon the Fevers, which have, of late Years, been epidemical in *Minorca*, I thought it would not be improper, in the first Place, to give a short Account of the Nature of the Climate; the Qualities of the Soil, and its Productions; the Manners of the Inhabitants; their Diet, and Way of Life; and to mention some other Diseases to which they are peculiarly subject.

The following Introduction is intended for this Purpose; and the Reader will  
B easily



easily discover by the Style, that it was composed during my Residence in that Island.

The Air is much more clear, and pure, than in *Britain*; being seldom darkened with thick Fogs: Yet the low Vallies are not free from Mists and unwholesome Vapours; and in windy Weather the Spray of the Sea is driven over the whole Island; as is evident, from the briny Dew found on the Leaves of Vegetables growing in the most inland Parts of it. Hence it is, that Utensils of Brass or Iron are extremely susceptible of Rust, in spite of all Endeavours to preserve them; and household Furniture, if not frequently exposed to Fire or the Sun, contracts a Mouldiness, and is destroyed by the Moisture.

The Summers are dry, clear, calm, and excessive hot; the Autumns moist, warm, and unequal; at one Time perfectly serene, at another cloudy and tempestuous:

## INTRODUCTION.

3

Nor are the Summer and Autumnal Seasons of one Year much different from those of another. During the Winter, Storms of any Kind, are neither frequent, nor of long Continuance (though sometimes very violent) and whenever they cease, the Weather returns to its usual Serenity. The Spring is always variable, but bears a stronger Resemblance of the Winter than the Summer Season.

The Changes of Heat and Cold, are neither so sudden nor so great in this Climate as in many others. In the Compass of a Year, the Mercury in FAHRENHEIT'S Thermometer, seldom rises much above the 80th, nor falls below the 48th Degree; and though it has been observed, in extraordinary Seasons, to reach the 87th, and descend to the 41st Degree; yet it attains not usually to either of these Extremities, nor continues at them long. In Summer there is scarcely ever four or five Degrees Difference between the Heat



## INTRODUCTION.

of the Air at Noon, and at Night, and in Winter the Variation is still less considerable.

What I have said with Regard to the Thermometer, is to be understood of one kept within Doors, and so situated, as neither to be affected by the solar Rays, nor the Influence of a Fire. For if it be exposed to the Summer's Sun, it will rise 12, 14, or 16 Degrees higher than in the House; and in other Seasons, the Difference between the Air in the Shade, and that heated by the direct Rays of the Sun, is frequently found much greater. Yet even in the Dog Days, the Heat of the Atmosphere, at least in open Places, where there is a free Circulation of Air, seldom surpasses that of the Blood of a Man in Health.

The Winds about the vernal and autumnal Equinoxes, and sometimes during the Winter, are extremely boisterous. In the other Seasons, they are usually moderate;

derate; and according to the Seamen's Observations, they rarely blow in the same Direction near the Islands adjacent to the *Gulph of Lyons*, as in the open Sea.

During the Summer, in the Mornings and Evenings there is commonly a perfect Calm: But the Middle of the Day is cooled by refreshing breezes, which rise in the East, and following the Course of the Sun, increase gradually till two or three o'Clock in the Afternoon; and die away insensibly as Night approaches. This renders the intense Heat of the Sun less dangerous and inconvenient: But if these Breezes intermit for a Day, the Natives themselves grow languid and inactive from the Sultriness of the Season.

The Northerly Winds in general are cold, dry, and healthy; they dispel the Mists, and make a clear blue Sky; whilst those which blow from the opposite Points render the Air warm, moist, and unhealthy. That the Influence of the



North Wind is superior to the rest, is evident from hence, that the Tops of all the Trees incline to the South, and the Branches on the North Side are bare and blasted. The next to it in Force is the North-West. Both these Winds are frequent towards the Close of Winter, and in Spring; and as they are dry and cold, they shrivel up the Leaves of the Vegetables, destroy their tender Shoots, and are often excessively detrimental to the Vineyards and rising Corn. The piercing Blasts, which blow at the same Season from the North-East, as they are moister, and more frequently attended with Rain, are not so prejudicial. The South and South-East Winds are by much the most unhealthy. In whatever Seasons they blow, the Air is foggy, and affects the Breathing; but in the Summer they are sultry and suffocating. An excessive Dejection of Spirits is then a universal Complaint; and, on exposing the Thermometer to the Rays of the Sun, the Mercury has frequently risen above  
the

## INTRODUCTION.

7

the 100th Degree. The West Wind is usually drier than the South: The East is cold and blustering in the Spring, and sultry in the Summer.

The Weather in this Climate is generally fair and dry: But when it rains, the Showers are heavy, though of short Continuance; and they fall most commonly in the Night.

The Sky in Summer is clear, and of a beautiful Azure, without Clouds or Rain; but moderate Dews descend regularly after Sun-set.

As the Autumn draws near, the Weather becomes less serene; Whirlwinds and Thunder are frequent. In the Nights, Lightning, and those Meteors called falling Stars, are very common.

At that Season, Water-spouts are often seen in the Neighbourhood of the Island; and sometimes they break upon the Shore.



LUCRETIVS\* has described them elegantly, and the various Shapes, in which they appear, are well represented by Dr. STUART in the *Philosophical Transactions* †.

About the Autumnal Equinox, a sudden Alteration in the Weather ensues; the Skies are darkened with Clouds, and the Rain falls in such Quantities, that the Torrents thereby occasioned, pouring from the Hills, tear up Trees by the Root, carry away Cattle, break down Fences, and do considerable Mischief to the Gardens and Vineyards. But then these anniversary Rains are much more violent than lasting, always falling in sudden heavy

\* Nam fit ut interdum tanquam demissa columna,  
In mare de cœlo descendat; quam freta circum  
Ferviscunt, graviter spirantibus incita flabris,  
Et quæcunque in eo, tum sunt deprensa tumultu,  
Navigia, in summum veniunt vexata periculum.

*De Rer. Nat. l. vi.*

† Plate I. Vol. 4. Part II.

Showers,

## INTRODUCTION.

9

Showers, with Intervals of fair Weather. As they come after a long scorching Summer, they are very acceptable and beneficial; for they mitigate the excessive Heat of the Air, give a Check to the epidemical Diseases, and by softening the Sun-burnt Earth, render it capable of being cultivated. They are commonly accompanied with Thunder, Lightning, and Squalls of Wind for the most Part from the North. Such Storms as these are seldom seen in cold Countries; but they are frequent enough in warmer Climates; and the Descriptions given of them by VIRGIL \*, are equally just and poetical.

Hail

\* Sæpe etiam, immensum cœlo venit agmen aquarum,  
 Et scdam glomerant tempestatem imbribus atris  
 Collectæ ex alto nubes: ruit arduas Æther,  
 Et pluvîâ ingenti fata læta boumque labores  
 Diluit: implentur fossæ, et cava flumina crescunt  
 Cum sonitu, fervetque fretis spirantibus Æquor.  
 Ipse Pater, mediâ nimborum in nocte, corusca,  
 Fulmina molitur dextrâ; quo maximo motu

Terra



## INTRODUCTION.

Hail and Snow are often intermixed with the Rains which fall in Winter and in Spring; but the Snow, for the most Part, dissolves immediately; and Ice is here an uncommon Appearance.

It is not my Design to give a geographical Account of the Island, its Situation, Harbours, Cities, and other Particulars, of this Nature: these may be met with elsewhere. I shall only observe that the whole of it is what the Sailors term low Land; excepting a few Hills near the Center, of which the most con-

*Terra tremit: fugere feræ, et mortalia corda  
Per gentes humiles stravit pavor; ille flagranti  
Aut Atho, aut Rhodopen, aut alta Ceraunia telo  
Dejicit: ingeminant Austri, et densissimus imber,  
Nunc nemora ingenti vento, nunc litora plangunt.*

*VIRG. Georg. L. i. ver. 322.*

*Effusis imbribus atris  
Tempestas sine morâ furit, tonitruque tremiscunt  
Ardua terrarum, et campi: ruit æthere toto  
Turbidus imber aquâ, densisque nigerrimus Austris.*

*Æneid. L. v. ver. 693.*

siderable,

## INTRODUCTION.

11

siderable, named *Toro* by the Inhabitants, may be discovered from Sea, in clear Weather, at twelve or fourteen Leagues distance.

The Surface of the Island is rough and unequal; and in many Places divided, by long narrow Vales of a considerable Depth, which are called *Barranco's* by the Natives. They begin towards the Middle of the Island, and after several Windings, terminate at the Sea. The South-West Side is more plain, and regular, than towards the North-East; where the Hills are higher, with low marshy Vallies between them, the Soil less fruitful, and the whole Tract unhealthy to Man and Beast. Near the Towns and Villages the Fields are well cultivated, and enclosed by Stone Walls: But the rest, for the most Part, is rocky, or covered with Woods and Thickets. There are some Pools of standing Water, and but very few Rivulets; one cannot therefore easily  
account



account for Cardinal DE RETZ's\* Mistake, when he describes a Number of Streams pouring into *Mabon* Harbour, unless we suppose him to have seen it in the rainy Season.

The Soil is light, thin, and very stony, with a good deal of Sea Salt, and some calcareous Nitre intermixed. In most places there is so little Earth, that the Island appears to be but one large irregular Rock, covered here and there with Mould, and an infinite Variety of Stones. Notwithstanding this, it is not only extremely proper for Vineyards, but produces more Wheat and Barley, than could at first Sight be imagined. And, if the Peasants may be credited, it would always yield a Quantity of Corn, and Wine, sufficient for the Natives; did not the Violence of the Winds, and the excessive Drought of the Weather, in dif-

\* See his Memoirs.

ferent Seasons frequently injure their Crops.

The Fields commonly lie fallow for two Years, and are sown the third. About the latter End of Winter, or the Beginning of Spring, they are first broke up: and next Autumn, as soon as the Rains fall, they are again ploughed, and prepared for receiving the proper Seeds. The Tillage is neither a laborious, nor an expensive Work; for a Plough so light as to be transported from Place to Place, on the Ploughman's Shoulder, and to be drawn by a Heifer, or an Ass sometimes assisted by a Hog, is sufficient for opening so thin a soil. The later the Harvest happens, the more plentiful it proves. The Barley is usually cut down about the 20th of *May* New-Style, and the Wheat is reaped in *June*; so that the whole Harvest is commonly got in by Midsummer Day. The Grain is not threshed with Flails as in *England*, but trodden out on a smooth Piece of Rock by  
Oxen



Oxen and Asses, according to the Practice of the eastern Nations.

In the planting of Vines, it is customary to lay a large Stone on every Slip; this is a Protection, as VIRGIL observes\*, against the excessive Heat of the Sun, which would otherwise, in so shallow a Soil, deprive it of all Moisture; and at the same Time prevents the Mould being washed away by the immoderate Rains. *September* is the season for the vintage. After the Grapes are trampled, but before they are pressed, they are sprinkled over with Powder of Alabaster (1), in order to give the Wine a brighter red Colour. Such Wines as are made with Care, from the Fruit of old Vineyards, still deserve the Commendations for-

\* Hoc, effusus munimen ad imbres :  
Hoc ubi hiulca siti findit canis æstifer arva.

*Georg. 2.*

(1) Called by the natives *Parell*; it is got out of the same Pits with the Plaister of Paris.

merly

merly given them by PLINY\*. And they have one Property, seldom to be met with in Wines of this Complexion. I mean that of keeping the Body open, which renders them less heating, and therefore less prejudicial. But since the Commencement of the War, as *Mabon* Harbour has become the Rendezvous for Ships of different Nations, the Quantity of the Wine has been more regarded by the Proprietors, than its Quality. Hence the greatest Part of it grows sour in the Beginning of the Summer: And this is probably the principal Reason, why Dy-fenteries have of late Years, been both more frequent, and more fatal, than usual.

The Natives hang up Clusters of ripe Grapes (1) to the Ceilings of their Chambers, in order to dry for Winter Store.

\* Lib. 24. C. 6. Vina Balearica conferuntur Italiae primis.

(1) Uvæ Penfiles, *Panjois*.



The *Sapa Vini* (2), together with various Roots and Fruits preserved by being boiled in it, amongst other Uses, makes a considerable Part of their Bill of Fare on Fast Days.

In some Places there are Fields of Hemp (3), Flax (4), and Tobacco (5): They likewise sow Beans (6), Chichlings (7), Chich Pease (8), two Species of the Kidney Bean (1), and Lentils (2): These being a considerable Part of their Diet at such Times as they are prohibited from eating Meat, by their Religion.

(2) *Arrop.*

(3) *Cannabis, Canem.*

(4) *Linum, Lli.*

(5) *Nicotiana, Tabach.*

(6) *Faba, Favas.*

(7) *Lathyrus, Guixes.*

(8) *Cicer, Ciurous, Garravanfos.*

(1) *Phaseolus*, the common white large Kidney-Bean is called *Mongeta*; a smaller Sort with a black Spot in its middle *Fesol*, or *Guixon*.

(2) *Lens, Llentias.*

There

There are a few Pease (3) in the Vineyards and Gardens; but these are commonly reserved for the Tables of the Wealthy.

In the moist fenny Grounds, they plant quantities of Canes (4), which they make use of instead of Laths, to support the Tiles, in the Roofs of their Buildings. They have also in some few Places *Indian* Wheat (5), Job's-Tears, (6), and *Indian* Reed (7): The hard, stony Seeds of the two last being perforated and strung, serve them as beads for their Rosaries.

Their Gardens are more for Use than Show, and furnish most Kinds of Pot-Herbs, Roots, and Sallading, in great

(3) Pisum, *Peas*.

(4) Arundo donax, *Canya*,

(5) Mayz, *Blad de las indias*.

(6) Lachryma Job, *Lagrimas de viu*.

(7) Cannacorus, *Mariettas*.



Plenty: And the Herb Market of *Mahon*, has been as useful to the *British* Fleet in restoring the Health of the Sailors, as the Harbour in refitting and securing the Ships. There are at all Seasons, Cabbages (1), Coleworts (2), Lettice (3), Spinage (4), Endive (5), Beets (6), Parsley (7), Cresses (8), Leeks (9), Onions (10), Garlick (11), Sellery (12), Radishes (13), Horse Radish (14),

- (1) *Brassica capitata, Cols capdelladas.*
- (2) *Brassica aperta, Cols abertas.*
- (3) *Lactuca, Llatugas.*
- (4) *Spinachia, Espinachs.*
- (5) *Cichorium five Endivia, Escarolas.*
- (6) *Beta, Bledas.*
- (7) *Apium five Petro selium, Jullivert.*
- (8) *Nasturtium, Murrija.*
- (9) *Porrum, Porras.*
- (10) *Cæpe, Cebas.*
- (11) *Allium, Alls, Ai'as.*
- (12) *Apium dulce, Apits.*
- (13) *Raphanus, Ravas.*
- (14) *Raphan. rust. Cochlear. Spec. Ravas de cavall.*



Sage (1), Mint (2); Sweet Marjoram (3), Wild Marjoram (4), Thyme (5), &c. You have besides these in Winter, Carrots (6), Parsnips (7), Turnips (8), Artichoaks (9), Asparagus (10), and Colly-flower (11): In Summer, Love-Apples (12), Mad-Apples (13), Guinea Pepper (14), together with various Kinds of Cucumbers (15), Pompions (16), Musk-melons (17), and Water-melons (18), in great

- (1) *Salvia, Sauvia.*
- (2) *Mentha, Herba sana.*
- (3) *Majorana, Morredux.*
- (4) *Origanum, Orenge.*
- (5) *Thymus, Them.*
- (6) *Daucus, Bastenagues.*
- (7) *Pastinaca, Xaroviás.*
- (8) *Rapum, Naps.*
- (9) *Cynara, Carxofas.*
- (10) *Asparagus, Esparachs.*
- (11) *Brassica cauliflora, Cols flos.*
- (12) *Lycopersicon, mala Aurea, Tomatils, Tomatigues.*
- (13) *Melongena, mala infana, Auberginias.*
- (14) *Capficum, Pebres.*
- (15) *Cucumis, Cubombros.*
- (16) *Pepo, Carabusas.*
- (17) *Melo, Melons.*
- (18) *Anguria, Sindrias.*



## INTRODUCTION.

Plenty and Perfection. But as the Climate is liable to severe Droughts, every Garden is supplied with a deep Well; from whence, by means of the *Persian Wheel* \*, the Gardener fills his Reservoir, and from it conveys the Water by Stone Canals, to the different Beds as Occasion requires.

Beside the Fruits common in *England*; such as Cherries (1), Apples (2), Pears (3), Apricocks (4), Plumbs (5), Peaches (6), Medlars (7), Mulberries (8), Quinces (9), and Walnuts (10); there are several others ripen here, which in colder

\* See a Figure of the *Persian Wheel* in *Shaw's Travels*.

- (1) *Cerasus, Cireras.*
- (2) *Malus, Pomas.*
- (3) *Pyrus, Pyras.*
- (4) *Armeniaca, Aubercoes.*
- (5) *Prunus, Prunes.*
- (6) *Persica, Presechs.*
- (7) *Mespilus, Nesples.*
- (8) *Morus, Moros.*
- (9) *Cydonia, Codens.*
- (10) *Nux. Juglans, Nous.*

Countries,



Countries, seldom or never come to Perfection, even with the Help of a Hot-bed; such as the large juicy Pomegranates (1), of which there is great Abundance in every Garden; Lemons (2), Citrons (3), and Oranges (4), which are all, of late years, become very common in this Island; the Almonds (5), which thrive perfectly well in this Soil; and the *Indian Figs* (6), which are the principal Sustenance of whole Families in *September*: the prickly Shrub that bears them grows wild among the Rocks, and is often made use of as a Fence to their Gardens. To these may be added some Fruits of smaller Esteem, as the Jujubes (7), the true Services (8), *Ne-*

(1) *Punica, Magranas.*

(2) *Limon, Llimons.*

(3) *Citreum, Cidros.*

(4) *Aurantium, Teronges.*

(5) *Amygdalus, Merlas.*

(6) *Opuntia vulgo Herbariorum, Figas Moriscas.*

(7) *Zisiphus, Gingols.*

(8) *Sarbus legitima, Ser-ves.*



## INTRODUCTION.

*apolitan* Medlars (9), and the Berries of the Nettle-Tree (10).

In enumerating the Trees which adorn their Gardens, I must not omit the Cypress (1), Laurel (2), Poplar (3), *Egyptian* Thorn (4), Bead-Tree (5), and a beautiful Species of Dog's-Bane (6). Neither can I forbear to mention the Fig-Tree (7), which not only produces large Quantities of excellent Fruit, (some Kinds of it, two Crops in a Year) but affords a convenient Shade, under which the Peasants usually regale themselves. Nor must the Palm-Tree (8) be past over in silence; for although the Dates in

(9) *Mespilus folio Apii lacin. C. B. Azarolas.*

(10) *Celtis, Lotus, Lladons.*

(1) *Cupressus, Cipré.*

(2) *Laurus, Lloré.*

(3) *Populus, Poll.*

(4) *Acacia, Aroma.*

(5) *Azederach, Suclemoro.*

(6) *Apocynum, Sedé.*

(7) *Ficus, Figuera.*

(8) *Palma, Fascé.*

this



this Country never come to Perfection, yet its inner Branches, when blanched, serve as Ornaments for their Proceffions on *Easter Sunday*; and the others are used by the Children in *Passion Week*, for striking the Earth, which is superstitiously termed by the Inhabitants, beating of *Judas*. The Minds of the People are at that Time so inflamed by the Sermons of their Priests, that it is dangerous for the Jews to be seen abroad.

The Vegetables hitherto mentioned, are produced by Culture. I come next to speak of the indigenous, which grow spontaneously in the Island; and as there is a vast Variety of these, I shall not pretend to treat them so fully as the Subject deserves; (nor am I equal to the Task, Botany, tho' sometimes my Amusement, never having been my Study) all I shall aim at, is, briefly to point out the most remarkable amongst them.

In the first Place, may be mentioned such Plants, as serve the Natives for Sal-



## INTRODUCTION.

lading and Pot-Herbs; viz. Succory (9), Prickly Rock Asparagus (10), Alexanders (1), Buckshorn Plantain (2), Goat's-Beard (3), Fennel (4), Hawkweed (5), Purflane (6), Sow-Thistle (7), Sorrel (8), Water-Cresses (9), Capers (10), and Samphire (1)

To the same Class belong Borrage (2), Blites (3), Beets (4), Orrache (5), Dan-

- (9) Cichorium, *Camarotjes*.
- (10) Asparagus, *Corruda*, *Esparachs*.
- (1) Smyrnum, *Cugulls*.
- (2) Coronopus, *Cornicellis*.
- (3) Scorzonera, *Tragopogon*, *Cuxa de Dona*.
- (4) Fœniculum, *Fenoi*.
- (5) Hieracium, *Cascunias*.
- (6) Portulaca, *Verdulagas*.
- (7) Sonchus, *Llecsons*.
- (8) Acetosa, *Vinagrellas*.
- (9) Syfimbrium, *Crexechs*.
- (10) Capparis, *Taparas*.
- (1) Crithmus, *Fenoi mari*.
- (2) Borrage, *Borrajtes*.
- (3) Blitum, *Blets*.
- (4) Beta, *Bledas*.
- (5) Attriplex.

delion

delion (6), and Spatling Poppy (7). The Luxury of the present Age, seldom or never, indeed, allows these a Place at the Table, but in Times of Scarcity they have served as common Food; particularly in the Year 1635, when a swarm of Locusts had destroyed the Harvest.

Of medicinal Plants, there is such Variety, that barely to recount them, will (I am afraid) be tedious. Common Wormwood (8), Sea-Wormwood (9), Brank-Ursine (10), True Maiden-Hair (1), Agrimony (2), Winter Cherry (3), Chickweed (4), Male Pimpernel (5),

(6) Dens Leonis, *Caxal de veia*.

(7) *Lychnis species, five spumeum papaver, Cou-*  
*liffe.*

(8) Absynthium Vulgare, *Donzell.*

(9) Absynth. Maritimum, *Donzell mari.*

(10) Acanthus, *Carnera.*

(1) Adiantum verum, *Falzia.*

(2) Agrimonia.

(3) Alkekengi, *Orval fi.*

(4) Alfine, *Tina.*

(5) Anagallis flore phæniceo.

Female



Female Pimpernel (6), Common Rest Harrow (7), Yellow Rest Harrow (8), Snap-Dragon (9), Goose Grass (10), Friar's Cowl (1), Cuckow-Pint (2), Spleen-Wort (3), Rough Spleen-Wort (4), Brook-Lime (5), Black Briony (6), Bugloss (7), Oxe-Eye (8), Shepherd's Purse (9), Mary-Gold (10), Small red Centaury (1), Small yellow Centaury (2),

(6) *Anagallis flore cæruleo.*

(7) *Anonis Vulgaris, Resta bovis.*

(8) *Anonis lutea viscosa, spinis carens, C. B. Motxes.*

(9) *Antirrhinum, Cullons de gat.*

(10) *Aparine, Amor de Hortola, & Rabosa.*

(1) *Arifaram, Frare Cugat.*

(2) *Arum, Rapa.*

(3) *Asplenium, Dauradella.*

(4) *Lonchitis.*

(5) *Veronica species, Becabunga.*

(6) *Tamnus, Bryonia nigra.*

(7) *Buglossum, Llengua bovina.*

(8) *Bupthalmum, Bulichs.*

(9) *Bursa Pastoris, Bosa de Pastor.*

(10) *Calendula, Lleua mal.*

(1) *Centaur. minus vulgar. Centauré.*

(2) *Centaur. min. flav.*

Germander (3), Ground-Pine (4), great Celandine (5), Hemlock (6), Male-Cistus (7), Female Cistus (8), Golden flowered Samphire (9), Sow-bread (10), Hound's-Tongue (1), Long-rooted sweet Cyperus (2), Wild Carrots (3), Great Dragons (4), Viper's Buglos (5), Dwarf-Elder (6), Golden Cassidony (7), Horse Tail (8), Wild Rocket (9), Base Wild

- (3) Chamædrys, *Ufinetta*.  
 (4) Chamæpitys, *Iva*.  
 (5) Chelidonium majus, *Celedonia*.  
 (6) Cicuta.  
 (7) Cistus flore rubro, *Stipa*.  
 (8) Cistus flore albo.  
 (9) Asteris species, Crithmum chrysanthemum, *Salfons*.  
 (10) Cyclamen, *Pa de Porc*.  
 (1) Cynoglossum, *Llepafera*.  
 (2) Cyperus longus.  
 (3) Daucus Sylvester, *Bastenagues salvages*.  
 (4) Dracunculus major, *Rapa mascle*.  
 (5) Echium.  
 (6) Ebulus, *Ebul*.  
 (7) Elichrysum, *Mansmillas*.  
 (8) Equisetum, *Coua de Cavall*.  
 (9) Eruca, *Ruca*.

Rocket



## INTRODUCTION.

Rocket (10), Sea-Holly (1), Bank-Cresses (2), Giant Fennel (3), Fern (4), Fumitory (5), Dog's-Grass (6), Greater Turn-sol (7), Mules Fern (8), St. John's-Wort (9), Hypocistis (10), Glass-Wort (1), Wild Cucumber (2), Sharp-pointed Dock (3), Hart's-Tongue (4), Wood Sorrel (5), Mallows (6), White Hoar Hound (7), Stinking Hoar Hound (8), Base Hoar

- (10) Refeda.  
 (1) Eryngium, *Card panical*.  
 (2) Erysimum.  
 (3) Ferula, *Canya fillera*.  
 (4) Felix, *Falguera*.  
 (5) Fumaria, *Fumisterra*.  
 (6) Gramen caninum, *Gram*.  
 (7) Heliotropium majus, *Girafol*.  
 (8) Hemionatis.  
 (9) Hypericum, *Transflorina*, *Herba de san Juan*.  
 (10) Hipocistis, *Margalidetas*.  
 (1) Kali, *Sofa*.  
 (2) Cucumis afininus.  
 (3) Lapathum acutum, *Paredellas*.  
 (4) Lingua Cervina, *Llengua de Cero*.  
 (5) Oxys, *Rujula*.  
 (6) Malva, *Mauves*.  
 (7) Marhubium Album, *Malrubins*.  
 (8) Ballote, *Malrubins bords*.

Hound



## INTRODUCTION.

29

Hound (9), Dog's Mercury (10), Sciatica  
 Cresses (1), Roman Fennel-Flower (2),  
 Broom Rape (3), Peony (4), Corn Poppy  
 (5), Yellow flowered Horn Poppy (6),  
 Pellitory of the Wall (7), Thorough Wax  
 (8), Periwinkle (9), Ivy (10), Honey-  
 suckle (1), Bind-Weed (2), Burnet (3),  
 Common Plantain (4), Water Plantain (5),  
 Knot-Grass (6), Polypody of the Oak (7),

- (9) Stachys.
- (10) Mercurialis, *Malcorages*.
- (1) Nasturtium sylvest. Iberis, *Murrissa bord.*
- (2) Nigella, *Niella*.
- (3) Orobanche.
- (4) Pæonia, *Pampelonia*.
- (5) Papaver. rubr. *Rofellas*.
- (6) Papaver corniculatum.
- (7) Parietaria, *Maias*.
- (8) Perfoliata.
- (9) Pervinca, *Proenga*.
- (10) Hædera, *Eura*.
- (1) Caprifolium, *Madrefelva*.
- (2) Convolvulus, *Corregiola*.
- (3) Pimpinella fanguis orb. *Pantinella*.
- (4) Plantago *Plantage*.
- (5) Plantago Aquatica.
- (6) Polygonum, *Ceutnous*.
- (7) Polypodium, *Polipodi*.

Flea-



## INTRODUCTION.

Flea-wort (8), Cinquefoil (9), Small Mad-  
 der (10), Butcher's Broom (1), Willow-  
 wort (2), Round-leaved Water Pimper-  
 nel (3), Scabious (4), Shepherd's Needle  
 (5), Sea Onion (6), Figwort (7), Greater  
 House Leek (8), Smaller House Leek (9),  
 Water-Parfnip (10), Night-shade (1),  
 Loufe-wort (2), Thorn Apple (3), White  
 Mullein (4), Vervain (5), Swallow-

- (8) Pfyllium, *Herba Pucera.*  
 (9) Quinque folium, *Pau de Christ.*  
 (10) Rubia Minor.  
 (1) Ruscus, *Brufe.*  
 (2) Salicaria.  
 (3) Samolus.  
 (4) Scabiosa, *Scabiosa.*  
 (5) Scandix, *peſten Vener.*  
 (6) Scilla, ornithogall. *Ceba marina.*  
 (7) Scrophularia, *Herba pudent.*  
 (8) Sedum majus, *Conſolva.*  
 (9) Sedum minus.  
 (10) Sium, *Apits de Sichia.*  
 (1) Solanum, *Morella.*  
 (2) Delphinium, Staphisagria, *Mata poi.*  
 (3) Stramonium, *Orval.*  
 (4) Verbascum, *Trapo.*  
 (5) Verbena, *Herba Verbena.*

wort



wort (6), Navel-wort (7), Common Nettles (8), and Roman Nettles (9).

To these we may add such others, as have been transplanted formerly from foreign Countries, but are now so far naturalized here, as to grow wild in several Places; such as the *American* Night-shade (10), the *Mexican* jasmine (1), the Passion Flower (2), the great Spurge (3), and the Aloes (4): Which last seems to have been industriously cultivated near the Farm Houses, as a Remedy for the Accidents to which Country Labourers are much exposed; it having been formerly in great

(6) *Asclepias*.

(7) *Cotyledon*, Umbilic. Vener. *Cocas de Parets*.

(8) *Urtica*. Vulg. *Ortigas*.

(9) *Urtica* Romana.

(10) *Phytolacca*, *Rems de moro*.

(1) *Jallapa flore purpureo*, *Juan de Notxe & Pedro de Notxe*.

(2) *Granadilla*, *Passionera*.

(3) *Ricinus*, *Cagamutxe*.

(4) *Aloes*, *Azibare*.

Esteem



Esteem (as we learn from DIOSCORIDES\*) for healing recent Wounds.

Beside these there is a Number of Aromatick Plants, which thrive luxuriantly in this Soil, and by imbalming the Air with fragrant Exhalations, contribute to preserve as well as restore the Health of the Inhabitants: The principal of these are Lavender Cotton (5), Starwort (6), Mountain-Mint (7), Horse-Mint (8), Penny-royal (9), Flea-bane of various Kinds (10), Clary (1), Poley-Moun-

\* Lib. ii. cap. 23. Aloe nascitur in Arabia, Asia, et aliis locis Maritimis, inutilis quidem succo extrahendo, sed conveniens recentibus vulneribus, si confusa Emplastri formâ, applicetur.

(5) Santolina, Abrot. fœm. *Camomela.*

(6) Asteriscus foliis ad florem mollibus. An Asterisc. annus aquat. patul. Tourn?

(7) Calamintha.

(8) Mentha, Mentastrum, *Mendafira.*

(9) Mentha, Pelugium, *Puriol.*

(10) Asteris species, *Conyzæ, Olivarda.*

(1) Sclarea, Horminum, *Tarach.*



tain (2), Syrian Herb Mastick (3), Rue (4), *French* Lavendar (5), Water-Germander (6), *St. Peter's* Wort (7), and strong smelling Clover (8).

The Garlick makes so considerable a Part, both of their Food and Physick, that it deserves to be more particularly mentioned. Several Kinds of it grow here in such Plenty that the Milk of the Cattle, and even the Meat itself, frequently participate of its Taste. The only Sort which the Natives use, is a *Scorodo-Prasum*, (9) milder than the Garden Garlick, and sweeter than the Leek. What is commonly eaten by the Soldiers and Sailors, is

(2) *Polium montanum*, *Polio*.

(3) *Chamædrys maritima*, *incana fructif. fol. lanceol. Frigola*.

(4) *Ruta*, *Ruda*.

(5) *Stæchas*, *Tumani*.

(6) *Chamædrys pallust. Scordium*, *Scordi*.

(7) *Ascyrum exiguo folio, flore magno, Stepara grega*.

(8) *Trifol. bituminosum*.

(9) *Allium Capite sphærico, folio latiore, Porradallt*.

D

another



## INTRODUCTION.

another more acrid Species, with a triangular Stalk (10).

Of the Thistle Tribe, the following deserve our Notice: The milky (1), and golden Thistle (2), are both Esculent; the Flowers of the prickly wild Artichock (3) serve, instead of Rennet, to curdle Milk; in *May*, the Bees feed chiefly on the white flowered Star Thistle (4), which affords the finest Honey; and the broad yellow flowered Fish Thistle (5) yields a coarser Sort, about a Month later.

The Bounds to which I propose to confine myself, will not permit me to enumerate the different Spurges (6), whereof

(10) *Allium caule triangulo.*

(1) An *Carduus lact. peregrin. maj. Sem. fusco?*  
*Card. Gallofe.*

(2) *Scolymus chrysanthem, ann. Caderlinas.*

(3) *Cynara Sylvestris.*

(4) *Carduus stellat. five Calcitrapa, flore Albo.*  
*Card. Blanc.*

(5) *Carlina. Arcana flore luteo patulo, Card.*  
*Segrelle.*

(6) *Tithymalus, Lletrera, Baladre.*

the



the Arboreſcent are the moſt beautiful; nor the Orchis's, among which thoſe that reſemble the Bee (7), and Butterfly (8), have the Preference; much leſs will they allow me to deſcribe the various Species of Linarias (9), Sea-Lavender (10), Rattle Graſs (1), and the vaſt Profuſion of Ranunculus's (2), and Geraniums(3), which enamel the Meadows: Yet, I cannot omit a few Plants, with bulbous Roots, and liliaceous Flowers, which are a peculiar Ornament to the Fields; *viz.* A white Star of *Bethlem* (4), a blue Muſcari (5), a Fleſh-coloured Corn Flag (6), Vernal

- (7) Orchis fucum referens.  
 (8) Orchis Papilionem referens.  
 (9) Linaria, one kind is called *Gall ſave*.  
 (10) Limonium.  
 (1) Pedicularis.  
 (2) Ranunculus.  
 (3) Geranium.  
 (4) Ornithogalli ſpecies, an Lilium Alexandrinum? *Liri Bord.*  
 (5) Muſcari, *Aiaſas*.  
 (6) Gladiolus, *Coltells*.



## INTRODUCTION.

Daffodill (7), with a yellow Cup, an Autumnal Daffodill with a large white Flower (8), Snow-Drop (9), Saffron (10), Meadow Saffron (1) *Spanish* Nut (2), and two Species of Asphodell (3,) both of which are luxuriant in the richest Soils, and from their Flowers, the Bees extract a considerable Share of the finest Honey.

Besides, in all the uncultivated Parts of the Island, thick, evergreen Bushes, of Mastick (4), Mock-Privet (5), and dwarf Olive (6), sprout up in such Abundance, that the broken surface of the Ground is

- (7) *Narcissus medio luteus.*  
 (8) *Narcissus Maritimus* C. B. *Azucena.*  
 (9) *Narcisso-leucoium, Aiasas.*  
 (10) *Crocus, Safrá.*  
 (1) *Colchicum.*  
 (2) *Sifyrinchium.*  
 (3) 1. *Asphodelus albus non ramos* C. B. 2. *Asphodelus minor foliis fistulosis, 1. Porraces, Albuens, 2. Cebollas.*  
 (4) *Lentiscus, Mata.*  
 (5) *Phillyrea, Ledern.*  
 (6) *Oleastellus, Ullastre.*



at all Seasons covered with an agreeable Verdure.

Intermixed with these, is an immense Quantity of Myrtle (7), sweet Gum Cistus (8), and Rosemary (9;) which jointly perfume the Air with an exquisite fragrance; and from the Flowers of the last, a delicious Kind of Honey is collected early in the Spring.

There is also a great deal of the Strawberry Shrub (10), Heath of various Kinds (1), and a long coarse Grass (2), on which the Goats, and black Cattle feed, when they are deprived of better Sustenance, by the Severity of the Winter.

(7) *Myrtus, Murta.*

(8) *Cistus labdanifera, Stepara stepa.*

(9) *Rosmarinus, Romani*

(10) *Arbutus Comarus Theoph. Arbesé.*

(1) *Erica, Bruc mascle, Bruc femelle & spel.*

(2) *Graminis Species, Carex.*



Shrub-Trefoil (3,) Stinking Bean-Trefoil (4), Sea-Grape (5), Shrub-Mullien (6), Sloe-Tree (7), Dwarf-Palm (8), Widow-wail (9), and another Kind of Spurge-Olive (10), beside that which affords the *Grana Cnidia* (1), likewise grow in several Places: But Brambles (2), Wild Roses (3), and some other thorny Bushes (4), are every where so common, that it is requisite for those who would pass through the Thickets, to dress like the Peasants, in short Jackets and leathern Spatterdashies.

(3) *Dorycenium Monspeliens.*

(4) *Anagyris foetida, Garroue bord.*

(5) *Ephedra Anabasis Bellon. Trompera.*

(6) *Phlomis fruticosa.*

(7) *Prunus Sylvester, Prunoné, Arrinoné.*

(8) *Palma humilis folio flabelliformi, Garbaions.*

(9) *Chamelæa tricoccus.*

(10) *Thymelæa lanuginosa foliis sedi minor. C. B.*

*P. Pala marina.*

(1) *Thymelæa fol. Lini, Matapoll.*

(2) *Rubus, Romaguera.*

(3) *Rosa Sylvestris, Gaverera.*

(4) 1 *Genistæ spinosæ Species.* 2. *Vel. Hierac. vel. Dent. Leon. species, 1. Argelaguera, 2 Saccorell.*

Hitherto



Hitherto I have treated of the lower Vegetables: I proceed next to the Trees. And here, I must again mention the Myrtle, Mock Privet, Lentiscus, and Arbutus, as they often rise to the Height of Trees; and mixing with Pines (5), Wild Olives (6), and great Holm-Oakes (7), which never lose their Verdure, supply the Cattle with Shelter during the excessive hot or cold Weather; and with Leaves to feed on, when the Summer's Sun, or the Blasts of the Winter, have destroyed the Grass of their Pastures.

But it is not the Cattle alone, which have been furnished with Food, from these Woods and Thickets; for in Times of Famine, the Inhabitants of this Island have been obliged to have Recourse to Acorns, wild Dates (8), the Berries of the

(5) *Pinus Pi.*

(6) Oleaster *Ullaftre.*

(7) *Ilex, Ufnas.*

(8) This is the Fruit of the *Palma humilis folio fabelli formi.*



Myrtles the Bramble, the Arbutus, and the like; which, if we may believe the Poets\*, was the Nourishment of the primitive Race of Mankind.

From the Qualities of the Soil, and the Abundance of uncultivated Fruit it produces, one might justly expect, according to VIRGIL'S Remark †, that the Olive would thrive well in this Island; yet the Natives take so little Care to propagate it, that they are obliged to import almost all their Oil from *France, Spain, and Majorca*; and as they purchase it with ready Money, it is a considerable Detriment to the Country. I know it is commonly alledged that they cannot have Olives of their own Growth in Plenty and Perfection, because there are but few Hills high

\* LUCRET. lib. v. VIRGIL'S *Georg.* lib. ii. OVID'S *Metam.* lib. i.

† Difficiles primum terræ, collesque maligni  
Tenius ubi Argilla, et dumosis calculus Arvis,  
Palladiâ gaudent sylva vivacis Olivæ,  
Indicio est, tractu surgens Oleaster eodem  
Plurimus, et strati baccis sylvestribus Agri.

enough



enough to protect them from the cold northerly Winds: But I have been well assured by several very good Judges in such Matters, that there are many Places, particularly in the *Termino* of *Ferarias*, where this tender Fruit might be sufficiently sheltered; and that the Want of this most useful Commodity, is rather to be ascribed to the Inattention and Neglect of the People, than to any other Cause. I have likewise been assured, that a *Spanish* Governor of this Island, made use of his Authority in obliging the Inhabitants to cultivate the Olive; and that his Endeavours would have had the desired Effect, had they been duly seconded in succeeding Times. Wherefore it were to be wished that some of his *British* Successors would follow so laudable an Example; for without their Assistance, it is to be feared, that a people so tenacious of the Customs of their Ancestors will scarcely come into a new Project, however advantageous it might prove to Posterity.

From



From the Berries of the Mastick they extract an Oil (as DIODORUS SICULUS tells us they did in his Time) which is commonly used for Lamps; and sometimes by poor People for frying Fish; in which Case they previously correct its Astringency by dipping in it a Piece of soft Bread. The Leaves of the Myrtle are very serviceable in tanning their Leather, and in dying their Cloaths black, which is a favourite Colour among the *Spaniards*; and its tough flexible Boughs being properly twisted, make the best and most durable Rope, for the *Persian* Wheel above-mentioned.

The Berry-bearing Savine (9) grows to a considerable Height in several Places near the Sea Coast; and the marshy Valleys afford plenty of Oziers (10), and Tamarisk (1), of which the Islanders make

(9) *Sabina folio Cupressi*, C. B. *Savina*.

(10) *Salix viminea*, *Vime*.

(1) *Tamariscus*, *Tamarell*.

Hoops



Hoops for their Casks. The Carob-Tree (2), the Hawthorn (3), and Evergreen, Privet (4), are likewise to be met with in the Woods; as also few Specimens of the Agnus Castus (5). \*

And now having taken a View of the Vegetable Kingdom, it would be no difficult Task for me to shew, that the indigenous Plants are well adapted to the Nature of the Climate, and wisely calculated for preserving or restoring the Health of the Inhabitants: But as it would be tedious to descend too far into Particulars, I shall only observe, that even the Shape and Figure of the Trees seem to be somewhat more than merely the Effect of Chance; none of them grow tall or lofty;

(2) *Ceratia*, *Garrove*.

(3) *Mespilus apii* fol. sylv. spin. *Spinal*.

(4) *Alaternus*, *Llampugal*.

(5) *Vitex*, *Alloch*.

\* The Reader may be pleased to take Notice, that the *Latin* Names are those by which they are described by *Tournefort*; the others in *Italicks*, are those by which the Natives distinguish them.



they never lose their Leaves; and the greatest Part of them have their Trunks so bent by the North Wind, that their Tops, which are broad and bushy, stretch out almost horizontally towards the South, and furnish a commodious Shelter to Man and Beast, against all Inclemency of the Weather. What still makes them fitter for this Purpose is the numerous Climbers that are commonly joined with them; particularly, the Traveller's Joy (6), and prickly Bind-weed (7), which are often so interwoven with the wide-spreading Branches of the Lentisc and wild Olive, as to render the Retreat which they afford almost impenetrable by Rain or Sun. Had not Nature provided such a necessary Cover, few or no Animals could live on these Rocks in the tedious scorching Summers; nor could they well endure the heavy Rains and chilling Blasts to which the Climate is liable.

(6) Clematis, *Vid. Auba.*

(7) Smilax Aspera, *Arritges.*

From



From what has been said, it sufficiently appears, that these evergreen Woods and Thickets, which Nature has surprizingly raised upon a Rock, are not only a great Ornament to this Island, but of infinite Advantage to the Inhabitants; they furnish them with Fuel, and their Cattle with Food and Shelter; and as the finer Soil is washed away by the anniversary Rains, their Fields would soon become barren, were they not constantly supplied with fresh Manure from the Leaves of the Vegetables, intermixed with the Dung of the Animals which feed in the Woods. The Natives, therefore, are much to be blamed in felling so many of their Trees, and stubbing up the Roots so rashly, as they have done of late Years, for immediate Profit, since the Damage will soon be sensibly felt, and not easily retrieved, by their Posterity.

The



The mineral Kingdom affords less Variety here than any other Part of Nature's Works. The Soil is of two Kinds; whereof one is light, blackish, and very fruitful; the other, called *Terra Agra* by the Natives, is a heavy, barren, red-coloured Earth. There is a Potter's Clay, of which they make Tiles, Bricks, and a coarse Kind of earthen Ware. Limestone and Plaister of Paris are found here in great Abundance, and supply the Inhabitants with different Sorts of Cement for their Buildings. The Stone commonly used in building is a white soft Grit, which is easily cut into any Form, and quickly cemented by means of the Plaister of Paris; so that they raise the Walls, and turn Arches with great Expedition: But in many Places of the Island, especially towards the North-East Side, the only Stone to be met with, is found in Shivers like Slate. In Caves not far from *Ciudadella*, are an infinite Number of beautiful *Stalactites*, some of which are hard enough to admit of a  
Polish.



Polish. There are some Fossil Shells; but no Kind of Metal, as far as I know, occurs in *Minorca*; though there is a good deal of metallic Ore in a small Island adjoining to it\*.

With respect to the Animals, I shall only mention such as are used for Food by the Inhabitants; beginning with the Fish, of which there is both Plenty and Variety. Some of them are to be caught at all Times either in the Bays or Harbours, or in the deeper Water at Sea; whilst others come regularly in Shoals at certain Seasons of the Year: But as many of them are unknown in *Britain*, or at least have no Name in our Language, I must refer the Reader to the Bottom of the Page †, where he will find inserted the

\* *Columba* Island.

† Pisces littorales, *Peix litoral.*

(1) *Polypi prima species Rondeletii, Pop Juen.*

(1) *Polypi secunda species Rond. Pop. ver.*

(3) *Loligo magna Rond. Eluja.*

(4) *Loligo parva Rond. Calamar.*

Names



Names given them in *Minorca*, and those bestowed on them by the best Authors who have treated on Fishes.

- Pisces littorales, *Peix litoral.*
- (5) Sepia Rond. *Sipia.*
- (6) Urtica, *Ortiga.*
- (7) Anguilla Salv. *Anguila.*
- (8) Conger Rond. *Congre.*
- (9) Muræna omnium Autorum, *Morena mascle,*  
y *femele.*
- (10) Salpa Rond. *Saupá.*
- (11) Sargus Rond. *Sarch.*
- (12) *Morruda.*
- (13) Scaras Onias Rond. *Variada.*
- (14) Sparns Rond. *Esperai.*
- (15) Aurata Rond. *Orada.*
- (16) Melanurus Rond. *Oblada.*
- (17) Mormyrus Rond. *Mabre.*
- (18) Erythrinus Rond. Rubellio Aldr. *Pagell.*
- (19) Pagrus Rond. *Pagre.*
- (20) Dentex sive Synodon Aldr. *Dental.*
- (21) Coracinus Rond. *Curbai.*
- (22) Buglossus, seu solea Rond. *Llenguada.*
- (23) Passer Bellonii, *Pedus.*
- (24) Cantharus *Cantara.*
- (25) Lupus Rond, *Llep.*
- (26) An Acarnan Rond? *Besuc.*
- (27) Scorpius major Rond. Gefn. *Cap rot x Rotje.*
- (28) Scorpius

The Island abounds with Cattle, Sheep,  
and Goats, which furnish the Inhabitants  
with

Pisces littorales, *Peix litoral.*

- (28) Scorpius minor five Scorpæna Rond. Gefn.  
*Rafelé Scorpera.*  
(29) An Anthiæ secunda species Rond. ? an Phycis  
Rond. ? *Mollera, Molle.*  
(30) Mullus Barbatu Rond. *Moll.*  
(31) Channa Rond. *Serra.*  
(32) Merula Salv. & Rond. *Mero, Enfos.*  
(33) Turdus varia specie, *Tort musot, flavafado, Bo-  
vos, pintado, Griwia.*  
(34) An Cestreu Rond. *Llifa.*  
(35) An mugil Cephal. Rond. *Cap pla.*  
(36) Julis Rond. *Donzella.*

Pisces testacei & crustacei, *Peix de Clofca.*

- (1) Astacus Rond. *Grumant.*  
(2) Locusta Rond. five Carabus, *Llangofa.*  
(3) Squilla lata Rond. *Sigala.*  
(4) Pagurus, *Cabre.*  
(5) Cancræ varii, *Cranchs peluts, reals, Jueus.*  
(6) Squilla parva, *Gambe.*  
(7) Echinus varia specie, *Voga mari.*  
(8) Pinna magna, *Nacre.*  
(9) Pholas, *Datil.*  
(10) Testudo marina, *Tortuga.*  
(11) Musculus, *Muscle.*  
(12) Tellinæ, *Cluifas.*
- E
- (13) Ofreum



## INTRODUCTION.

with Cheefe and Wool, both for their  
own Use, and for Exportation. As Lard  
or

Pisces littorales,      *Peix litoral.*

- (13) Ostreum, *Ostia.*
- (14) Lepas, Patella, *Pagellida.*
- (15) Peecten, *Cupina gravada.*
- (16) Cancellus, *Hermitan.*
- (17) Cochlea varia specie, *Cornes & Caragols.*
- (18) Murex, *Corns de fell.*
- (19)                    *Pau de Cabrit.*

Pisces Pelagii,      *Peix de alt al mar.*

- \* (1) Phocæna Rond. *Delfi.*
- \* (2) Centrine Rond. *Peix pore.*
- (3) Squatina Rond. *Escat, Escat vexigal.*
- \* (4) Zygæna Rond. *Llunada.*
- (5) Catulus maximus forte *Canicula faxatalis*  
Rond. *Gats.*
- (6) Catulus major vulg. *Canicula Arist. Rond.*  
*Aldr. Gatons, Pintarotge.*
- (7) Mustelus lævis primus Salv. *Musola.*
- (8) Galeus Acanthias, Spinax Aldr. *Caso.*
- (9) Xiphias Piscis, *Peix de Espasa.*
- (10) Pastinaca Aspera Bellon. *Romaguera.*
- (11) Pastinaca mar. læv. Bellon. *Ferrasa.*
- (12) Aquila Bell. & Salv. *Mila.*
- (13) Rana Piscatrix, *Buldroy.*
- \* (14) Torpedo, *Tremulo.*
- (15) Raia varia specie, *Ratjada, Caputxi, Clavell,*  
*Clavell borrell, Cardayre.*

(16) Faber



or Oil is commonly used in dressing their Victuals, they make but little Butter; and this by a very singular Process: They boil the Whey, which is pressed from the Curd in making Cheese, and skim off the Part which rises to the Top; when they have collected a proper Quantity of this, they

*Pisces Pelagii. Peix de alt al mar.*

- (16) Faber five Gallus mar. Rond. *Gall, Peix de san Pedro.*  
 (17) Milvus Salv. Hirundo Rond. *Xurigué.*  
 (18) Mugil alat. Rond. Hirundo Plin. *Uraola.*  
 (19) Cuculus Aldr, *Gallina, Gallinetta.*  
 (20) Lyra prior Rond. *Grenau, Peix de san Rafael.*  
 (21) Mullus imberbis Rond. potius Cuculi species, *Cabot de la mar.*  
 (22) Gurnardus griseus, *Uliora, Baluerna.*  
 (23) Draco five Araneus Plinii, *Arana.*  
 (24) An Ophidion Rond. ? an Acus lumbriciformis Willough. ? *Drago, Saltan cono.*  
 (25) Uranoscopus, Callyonimus, *Rata.*  
 (26) An Perca marina Rond ? *Serran imperial.*  
 (27) An Stromateus Rond ? *Llampuga.*  
 (28) Pompilus, *Pampul.*

*Pisces gregales autumnales.*

- (1) Acus vulgaris oppian. *Aguia.*  
 (2) Sardinia Rond. *Sardina.*  
 (3) Thrissa Rond. *Alatx.*



they work it a considerable Time with their bare Feet or Arms, which is the only Method of churning they are acquainted with; then by the Addition of cold Water, the Butter which floats upon the Surface is separated; and after being washed, is boiled, till the watery Particles are evaporated; by this Process it acquires, when cold, the Taste and Consistency of a thick, sweet, Oil.

*Pisces gregales hyemales.*

- (1) *Pelamys* Bellon. *Amia* Rond. *Bonitol.*

*Pisces gregales vernaes.*

- (1) An *Thynni* species? *Siruvia*, *Sirviola.*  
 (2) *Sphyræna*, five *Sudis* Aldr. *Espet.*

*Pisces gregales Æstivi.*

- (1) *Scomber*, *Veirat*, *Cavallar.*  
 (2) *Trachurus* Ald. *Saurell.*  
 (3) *Mænæ duplex* species, *Mora* & *Xucla.*  
 (4) *Smaris*, *Gerretts.*  
 (5) *Boops* Rond. *primus*, *Vega*  
 (6) *Encrasicholus* Aldr. *Anxove*, *Aledroc.*

*N. B.* In the above Catalogue of Fish, it is to be observed that *Willoughby's* Names are commonly made use of; and such as are marked with an Asterisk are seldom or never brought to Table.

Beef



Beef and Mutton, though commonly lean, are eatable throughout the Year: But in the Spring, while the Grass is tender, the latter is in its greatest Perfection; and the former in Summer, when the Cattle feed on the Stubble and Leaves of the Evergreens.

The Goats are fatest in Autumn, and are slaughtered from *September* to *January*, chiefly for the Use of the common People.

But of all the Kinds of Meat none is here in so great Plenty and Perfection as Pork; nor is any other so much esteemed by the Natives. It is in Season from *September* to *Lent*; Bacon is to be had at all Times; and being fried or boiled, is commonly eat with Bread for Breakfast; they enrich their Broth with Hog's Lard; and from the same Animal they make a great Variety of Puddings, particularly Sausages\*, scarce inferior to those from *Bologna*.

\* Sobreassados.



## INTRODUCTION.

They have likewise plenty of Rabbits ; as also Hedge-Hogs, and Land Turtles, which are sometimes eat by the Poor.

Their domestick Fowls are Turkeys (1), Geese (2), Ducks (3), Cocks and Hens (4), in great Numbers.

In the Woods and Fields (besides various Kinds of Owls and Birds of Prey, which I pass over, as they never make a Part of Diet) we have Ring Doves (5), Red Legg'd Partridges (6), Stone Curlews (7), Quails (8), Blackbirds (9),

(1) Gallo pavo five meleagris, *Galls de India.*

(2) Anser domesticus, *Ojas.*

(3) Anas domestica, *Anades.*

(4) Gallus gallinac & Gallina, *Galls & Gallinas.*

(5) Palumbus torquatus Aldr. *Tudons.*

(6) Perdix ruffa, *Perdius.*

(7) Oedicnemus Bellon. Charadrius Gefner. *Sabbellins.*

(8) Coturnix, *Gualleras.*

(9) Merula vulgaris, *Torts.*

Solitary

Solitary Sparrows (1), Nightingales (2), Gold-finches (3), and an infinite Number of other small Birds.

Wild Ducks of different Kinds (4), Wigeons (5), Teal (6), Coots (7), and several Sorts of Water Hens (8), are common about the Ponds and Marshes, Kings-Fishers (9), are frequent about the Shore; and Flocks of Rock-Pigeons (10) breed in the Caves and Hollows, formed by the dashing of the Waves round the Coasts.

(1) *Passer solitarius*, *Melleres*.

(2) *Luscinia* seu *Philomela*, *Roffinols*.

(3) *Carduelis Cardeneras*.

(4) *Anas fera*, *varia specie*, *Anades rosas de coll blau*, *sayardes*, *soteras*.

(5) *Penelope*.

(6) *Querquedula*, *Anadons*.

(7) *Fulica Fotges*.

(8) *Gallinula varia specie*, *Pollo's de Riu Gallets de Riu Titetas*.

(9) *Ispida*, an *Veterum Alcyon*?

(10) *Columba rupicola*, *Coloms*.



## INTRODUCTION.

Besides all these, Swifts (1), Swallows (2), Sand Martins (3), Turtles (4), Bee-Eaters (5), Hoopo's (6), and Sky-Larks (7), arrive here in the Spring, and, after bringing forth their Young, leave the Island in Autumn.

Wood-cocks (8), Snipes (9), a small Sort of Pigeon (10), green and gray Plovers (11), Red-wings (12), Fieldfares (13), Oxe-Eyes (14), Chaffinches (15),

- (1) *Hirundo Apus, Vinjolas.*
- (2) *Hirundo domestica, Uranellas.*
- (3) *Hirundo riparia, Culs blancs.*
- (4) *Turtur, Tortora.*
- (5) *Merops sive Apiafter, Abeyrols.*
- (6) *Upupa Aldrov. Puputs.*
- (7) *Alauda, Turrolas.*
- (8) *Sclopax, Segues.*
- (9) *Gallinago minor Aldr. Bogafines.*
- (10) *An Columba livia Gefner.? Xexels.*
- (11) *Pluvialis viridis & cinericea. Kilots & Jages.*
- (12) *Turdus iliacus, Torts borrell.*
- (13) *Turdus pilaris.*
- (14) *Fringillago, Ulls de bou.*
- (15) *Fringilla, Pinsans.*



Starlings (1), Foreign Quails (2), and Daker-Hens (3), make their Appearance about the End of *October*, and remain during the Winter. Cranes (4), Wild Geese (5), and Curlews (6), sometimes stop here, as if it were to recruit themselves for a further Flight; and now and then we meet with a Flamenco (7).

It is observed, that the Flesh of such Fowls as feed on Land, frequently tastes of Mastick or Garlick; and the Water-Fowl are best in bad Weather, when the Storms prevent their going to Sea, and living upon Fish.

But as several of the Animals above-mentioned, are only to be met with at

- (1) *Sturnus, Estornells.*
- (2) *Coturnix, Gualleras babarescas.*
- (3) *Ortygometra, an Rallus terrestris?*
- (4) *Grus, Gruas.*
- (5) *Anser ferus, Ojas salvages.*
- (6) *Numenius five Aquata.*
- (7) *Phænicopterus, Flamencos.*

the



the Tables of the Opulent, the plentiful Provision of Snails (1), with which Nature has furnished this Island, are of infinite Service for the Maintenance of the poorer Families, who eat them boiled, after having been kept within Doors long enough to lose their earthy Taste. In dry Weather, when they are in their Prime, they lurk in the Chinks of the Earth and Crannies of the Rocks, and commonly stick together in large Clusters like Grapes; which probably induced the *Romans* to give them the Name of *Cochleæ Cavaticæ*: But in wet moist Weather, they leave these Places of Retirement in quest of Food, and are frequently to be met with on the Stalks of the *Asphodells*, the Shoots of the Vines, and other Vegetables; for what *PLINY*\* says of their never coming out of their Caves, or feeding on Greens, is intirely fabulous.

(1) *Caragols, Boves, Mongetas, Caragolins.*

\* *Lib. viii. Cap. xxxix.*



Having given these short Hints relating to the natural History of *Minorca*, I shall now describe the Temper and Manners of the People, so far as may be necessary to introduce an Account of their Diseases.

The Natives of this Island, are commonly thin, lean, and well built, strong and active, of a middle Stature, and an Olive Complexion. Their Hair, for the most part, is black and curled; in many Chestnut coloured; in some red. In a Word, the young People are either of a sanguine or choleric Constitution; while those of more advanced Years, become dry, meagre, and, what the Ancients called atrabilious. Such is the natural Impetuosity of their Temper, that the slightest Cause provokes them to Anger; and they are equally incapable of forgiving and forgetting an Injury. Hence it is, that Quarrels about the merest Trifles daily break out even among Neighbours  
and



and Relations; and Family Disputes are hereditarily transmitted from Father to Son. Thus although Lawyers and Pettifoggers are very numerous in this Country, yet there is still too few for the Clients.

They don't commonly live to so great an Age as the Inhabitants of more Northernly Countries; though, perhaps, they are equal in this Respect with their nearest Neighbours on the Continent. Girls soon arrive at Maturity, and soon grow old. The *Menses*, for the most part, appear before Fourteen, and frequently at eleven Years of Age; in some they return twice a Month; in others every three Weeks, and continue from three to seven Days. Both Sexes are, by Constitution, extremely amorous: They are often betrothed to each other, while Children, and marry at Fourteen. The Women have easy Labours, and commonly return in a few Days to their usual domestic Business. But lest the Family should become



come too numerous for their Income, it is a Practice among the poorer Sort, to keep their Children at the Breast for two or three Years, that, by this Means, the Mothers may be hindered from breeding.

Bread of the finest Wheat Flour, well fermented and well baked, is more than half the Diet of People of all Ranks. Rice, Pulse, Cuscawowe, Vermicelli, Herbs and Roots from the Fields and Gardens, Summer Fruits, pickled Olives, and Pods of the *Guinea* Pepper, make up almost the other Half; so that scarce a Fifth of their whole Food is furnished from the Animal Kingdom; and of this, Fish makes by much the most considerable Portion. On *Fridays*, and other Fast-Days, they abstain entirely from Flesh; and during Lent, they live altogether on Vegetables and Fish, excepting *Sundays*, when they are permitted the Use of Eggs, Cheese and Milk. The most of their Dishes are high seasoned with Pepper,



per, Cloves, Cinnamon, and other Spices; many of them are tinged with Saffron, many sweetened with Honey or Sugar; and Garlick, Onions, or Leeks, are almost constant Ingredients. They eat a great deal of Oil, and that none of the sweetest or best flavoured; using it not only with Sallads, but also with boiled and fried Fish, Greens, Pulse, &c. instead of Butter. A Slice of Bread soaked in boiled Water, with a little Oil and Salt, is the common Breakfast of the Peasants, well known by the Name of *Oleagua*. Their ordinary Meals are very frugal, and consist of little Variety: But on Festivals, and other solemn Occasions, their Entertainments are to the last Degree profuse and extravagant. The Bill of Fare of a Country Farmer's Wedding Dinner, would scarce be credited. This Custom, with many others, they seem to have borrowed from the Eastern Nations.

People of all Conditions use Wine at their Meals; and though Drinking to  
Excess



Excess is not common, yet the Vulgar are not exempted from private Debauches, Drams of Anniseed Water being too much in Vogue. In Summer the excessive Heat obliges them to have frequent Recourse to large Draughts of cold Water, the greatest Part of which is but very indifferent; for the Cisterns are seldom clean, and the Water which the Springs and Rivulets afford, is often brackish, and always hard; so that it is neither proper for Washing, nor for boiling Pulse, and leaves a stony Crust on the Sides of the Tea-Kettles, and other Vessels in which it has been frequently boiled.

Though the Natives make three or four plentiful Meals a Day, they are generally costive; and many, in perfect Health, have no occasion to ease themselves oftener than twice a Week.

They are so much addicted to the Use of Tobacco, as never to be without a  
Pipe,



Pipe, either in their Mouths or their Pockets. In Summer almost every one sleeps an Hour or two after Dinner, and some follow this Practice throughout the Year.

A Fourth of their Time is made up of Holidays, on which, tho' Work is prohibited, Sports and Pastimes are allowed. A considerable Part of those Days is spent in the Churches or in Processions: At Night the more Sedate divert themselves in their Houses with Musick and Cards; whilst the young Men serenade their Mistresses, in the Streets, with the jarring Musick of their Guitars, and extemporary Love Songs of their own composing.

In the Interval between the Harvest and the Vintage, there are a Number of publick Diversions in different Places of the Island. Whether because the People have then most Leisure, or are in Reality more chearful, from the Serenity of the Weather,



ther, as one of their Proverbs \* seems to hint, I shall not determine. To their Horse and Foot Races †, notwithstanding the immoderate Heat of the Season, Men, Women, and Children, flock from all Quarters, and expose themselves to the Sun in the middle of the Day, dancing in the open Air on the scorching Rocks, rattling their Castanets in Concert with the Musick of the Guittar. Nor does their Mirth finish with the Day; as soon as it grows dark, some Pieces of Pine Tree are lighted as a Torch in the middle of the Street, where the Croud assembles, and continues dancing till Morning.

Every Year, about Midsummer, they have likewise a Diverfion in *Mabon Har-*

\* *En lo Estiu tout Hom. Viu.* In Summer every Body is alive.

† On St. JOHN'S Day, *June* the 24th, and the *Sunday* following. On St. PETER'S Day, *June* 29. On St. JAMES'S Day, *July* 25. On St. LAURENCE'S Day, *August* 10. On St. BARTHOLOMEW'S Day, *August* 24, and the 29th of the same Month. On St. GRACIA'S Day, *September* 8.

F

bour,



bour \*, which may properly be called a Boat-Race; and one cannot observe the Eagerness of the Boat-Men, and the Solicitude of their Friends on Shore, without calling to Mind VIRGIL'S Description of a similar Contest on the Anniversary of *Anchises's* Funeral †.

In the Carnival too, this People, however grave and serious at other Times, freely indulge themselves in all Manner of

\* *Viz.* on St. PETER'S Day, June 29.

† Viridem Æneas frondenti ex Ilice metam  
 Constituit signum nautis, pater; unde reverti  
 Scirent, et longos ubi circumflectere cursus.  
 Confidunt transtris intentaque brachia remis  
 Intenti expectant; exsultantiaque haurit  
 Corda pavor pulsans laudumque arrecta cupido  
 Inde ubi clara dedit sonitum tuba, finibus omnes,  
 Haud mora, profluere suis, ferit Æthera clamor  
 Nauticus adductis spumant freta versa lacertis  
 Infindunt pariter sulcos, totumque dehiscit  
 Convulsum remis, rostrisque stridentibus æquor,  
 Tum plausu, sonituque virum, strepituque frementum  
 Consonat omne nemus, vocemque inclusa volutant  
 Littora, pulsati colles clamore resultant.

*Æn. v.*

ludicrous



ludicrous Sports and Amusements; especially during the last Week of it, when their Revelling and extravagant Mirth resembles more the ancient *Bacchanalia*, than the Diversions of any modern civilized Nation: For Night and Day, the Streets are filled with People in Masks, and dressed in the most ridiculous Habits; while the jarring Sound of Castanets, Pipes and Tabors, Violins, Guittars, and their more discordant vocal Musick, heightened with Screaming, Shouting, and every wild Demonstration of intemperate Joy, make almost one continued Uproar.

After this Excess of Mirth, the Fasts enjoined by their Religion succeed, and continue till the Expiration of *Lent*; at which Time a Sheep or a Lamb is slaughtered by each Family; and on the joyful Night which puts an End to this mortifying Season, they endeavour, as it were, by one voracious Meal to make up for the Abstinence they have been compelled to observe. This proves fatal to some, and



would undoubtedly be so to many more, did not Nature prevent the Effects of their Intemperance, by a *Cholera Morbus*, or some such Relief.

The last Thing to be taken Notice of with regard to the Manners of these Islanders, is, that their great Veneration for Antiquity, and the little Intercourse they formerly had with other Nations, hath occasioned a Number of old Customs to be still kept here at this Day. Thus poetical Disputes are much in Vogue amongst the Peasants \*. One of them sings some extemporary Verses on whatever Subject he pleases, and accompanies them with the Musick of his Guittar: he is immediately answered in the same Number of unpremeditated Lines, by another, who endeavours to excel or ridicule him; and this alternate Contest lasts, to the no small Amusement of their attentive Companions,

\* These Verses are called *Glosses*, and those who excel in making them, *Glossodors*.

until



until the Wit of the rival Poets be exhausted. These are the *Carmina Amœbœa* of the ancient *Greeks*, in Imitation of which some of the Pastorals of THEOCRITUS, and VIRGIL, were written. In Imitation of the Ancients too, it is customary for Lovers to pelt their Mistresses with Oranges, as a Mark of their Regard\*; though this is a Diversion reserved for the Carnival. The Practice of throwing Nuts and Almonds at Weddings, which VIRGIL † mentions, is likewise retained. Soon after a Person is deceased, his Friends

\* Malo me Galatea petit lasciva puella, &c.

VIRG. *Eclog.* iii.

Malo ego te ferio, tu si me diligis, illud  
 Suscipe, me imperti et virginitate tuâ,  
 Hoc fieri, si posse negas, Hoc suscipe malum et  
 Quam pereat parvo tempore, forma vide.

PLATO'S Epigram paraphrased by BENTINUS.

Which is thus translated.

Fair, if you love me, take the Fruit I throw,  
 And, in return, your Virgin Sweets bestow;  
 Or take the Orange, tho' you scorn my Suit,  
 See Beauty's Emblem in the fading Fruit.

† Sparge marite nuces. *Eclog.* viii.



and Relations assemble in the House to bewail their Loss, and commemorate his Virtues, Weeping and Howling with all the seeming Agonies of Distress; and that this was an ancient Custom, appears by the elegant and pathetick Funeral Dirge, which we find in LUCRETIVS \*. And in this Island, as in the Southern Provinces of *France*, and in *Italy*, the dead Bodies are not nailed up in Coffins, but carried in an open Litter to the Grave, which, we learn from Authors, was also practised, on some Occasions, among the old *Romans* †. Another Instance of their inviolable Attachment to ancient Usages, is the Manner in which the Women wear

\* At jam non Domus accipiet te læta neque Uxor  
Optima; nec dulces occurrent Oscula Nati,  
Præripere et tacitâ pectus dulcedine tangent,  
Non poteris factis, tibi fortibus esse tuisque  
Præsidio: Miser, O Miser, aiunt, omnia ademit  
Una Dies infesta tibi tot Præmia Vitæ.      Lib. vi.

† See CORNELIVS NEPOS on the Burial of T. POMPON. ATTICUS and LISPIUS's Notes on this Passage.

their



their Hair ; for, contrary to the Custom of all the neighbouring Countries, excepting *Majorca*, they force it with Fillets to the back Part of the Head, and bind it in a *Queue* of a considerable Length. When the natural Locks are not long enough for this Purpose, false ones are added to supply the Deficiency ; for nothing is reckoned more unbecoming than to be destitute of a Tail, or to appear with one too short. But not to be tedious with too many Instances, I shall only add, that in the Use of the Sling the present Inhabitants of *Minorca* are no less dextrous than their Fore-fathers, who are said to have driven the *Romans* off their Coasts by Vollies of Stones \*. The Shepherds, or those that tend the Cattle, but seldom miss such of them as fall under their Displeasure ; and by this Means they have their Flocks and Herds so much at command, that even the cracking of the empty Sling is sufficient to intimidate them, and bring

FLOR. *Epitom.* Lib. iii.

F 4

them



them together in such Parts of the Pasture as their Keeper pleases: And as the Cattle are often maimed and hurt by the severe Chastisement inflicted by Means of this Instrument, the Farmers find it expedient to forbid the Use of it to such of their Servants as are of a cruel and mischievous Disposition.

The mentioning of their Cattle puts me in Mind of two other singular Practices which prevail here; *viz.* that of castrating Animals by bruising their Testicles, which, we learn from ALBUCA-SIS\*, was customary among the *Arabians*; and the Way of slaughtering Oxen, by thrusting a Knife into the *Medulla Spinalis*, immediately behind the *Occiput*, which is so much preferable to the Method of knocking them on the Head, that it is surprizing other Nations do not fall into it.

\* Chirurg. p. 2. cap. lxi.



Having gone through what I intended to remark, in Relation to the Natives, I should next, according to the Plan of this Introduction, give a circumstantial Account of the Diet, and common Way of Life, of the *British* Soldiers in this Island : But as this would be a disagreeable Task, I shall only observe, that the Excess of Drinking, is, among them, an universal Vice, confirmed into a constant Habit. *Pudet hæc opprobria nobis, &c.*

But however different the *Spaniards* be from the *English*, in their Meat, Drink, Exercise, Affections of the Mind, and Habit of Body ; yet the Health of those of both Nations is equally influenced by the Seasons. An epidemical Distemper seldom or never attacks the one Class of Inhabitants, while the other remains unhurt ; and surprizing as it may appear, it is nevertheless true, that the Peasants remarkable for Temperance and Regularity, and the Soldiers, who, without Meat and Cloaths, frequently lie abroad drunk, exposed



exposed to all Weathers, have Diseases almost similar, both as to their Violence and Duration. Hence it is evident how far the Power of the Air is superior to that of the other Non-naturals in producing Disorders of the Animal Oeconomy.

The Diseases which, from their Frequency in this Island, I reckon *Endemial*, may be divided into two Classes; the *Epidemick*, or such as affect Numbers together at particular Seasons; and the *Sporadick*, which are equally common at all Times of the Year. To the first belong the *Rash*, *Effere*, *Cholera Morbus*, *Tertian Fevers*, *Fluxes*, *Pleurisies*, *Peripneumonies*, *Erysipelatous Fevers*, and those that are accompanied with *Catarrhs*: To the second, *Obstructions of the Abdominal Viscera*, the *Hæmorrhoids*, *Ulcers of the Legs*, *Ruptures*, *Inflammations of the Eyes*, and *Nephritick Pains*.

With



With respect to the Epidemicks, it may not be improper to observe in general, that acute Fevers are more frequent here than in *England*; that they are much more violent, but of shorter Duration; that they oftner terminate completely by a manifest *Crisis*; and in all respects agree much more exactly with what the Ancients have said concerning such Evacu-ations, and the Periods in which they usually happen.

The most common Diseases of the Sporadick Kind, are Obstructions, Indurations, and Swellings of the glandular Viscera in the lower Belly; together with Wind in the first Passages, and bad Digestion. The Causes of these Obstructions seem to be, in the first Place, a Scarcity of good Water; for, as HIPPOCRATES has well observed \*, *Where there are no Rivers, and the Inhabitants are obliged to drink stagnating, ill-scented Well-Water, such must*

\* De Aere Aquis & locis sub finem.

*needs*



*needs hurt both the Belly and the Spleen.* And it is remarkable, that large Splens, like those described by *Trallianus* \*, and hard tumified Livers, are not only common to the human Species here, but also to Brutes; particularly the Sheep that feed on the North-East Side of the Island, where the Waters are very brackish; though the Butchers, to whom this Fact is well known, generally ascribe it to their eating the Horse-Mint and Penny-Royal, which abound in the Pastures.

*Secondly,* The intense and long continued Summer Heats, by dissipating the finest Particles of the animal Juices, necessarily leave the rest of a grosser and more earthy Nature. And thus, is a large Proportion of that Kind of Matter generated in the Blood, which the Ancients called *Atrabilious*; and this being deposited in the Viscera, occasions the above-mentioned Obstructions †.

\* Lib. viii. Cap. xii.

† See Boerhaave's Aphorisms de Melancholia.

*Thirdly,*



*Thirdly*, Another Cause of these Obstructions, is the Frequency of acute Diseases; and more especially of Tertian Fevers; which, as they frequently relapse, and go off with imperfect *Crisis*, weaken the Tone of the *Chylopoetick Viscera*, and at last terminate in hard scirrous Tumours of the Liver and Spleen.

It is likewise probable, that their living so much upon Pulse and crude Vegetables; the Abuse of Spirituous Liquors; their high seasoned Diet; their passionate Temper, and immoderate Use of Venery, co-operate with the Causes just now mentioned, in producing the same Effects.

But such is the Goodness of Providence, that every Climate seems to yield domestick Antidotes for the Endemial Diseases\*. Accordingly we find, that this Island

\* See the Quotations from RAY and BENORVINUS in DERH. *Physico Theolog.* Book x.

abounds



abounds with Whey, Honey, Summer-Fruits, gentle Catharticks, *Hieracea*, *Cicchoracea*, and all that Class of Plants and Roots with aperient saponaceous Juices, which Antiquity has recommended as Specificks or approved Remedies in obstructed Bowels.

But in these Distempers, it is universally allowed, that nothing is of so much Advantage, as the *Hæmorrhoidal Flux*; and therefore, however frequent and troublesome the Piles may be in this Climate, they ought to be considered rather as a Benefit of Nature, and a Remedy, than as a Misfortune, or a Disease; more especially as they prevent Pleurifies and Peripneumonies, according to the Doctrine of HIPPOCRATES\*.

BAGLIVI † tells us, that at *Rome*, Ulcers of the Legs are almost incurable, and

\* De Humor. & Epid. Lib. vi,

† Prax. Med. Lib. i. P. 102.



Wounds in them difficult to heal; while the like Accidents on the Head, are quickly cured without any Trouble. The same Thing happens here, insomuch that it is a Proverb among the Natives, "*Minorca is good for the Head, but bad for the Shins* \*." This, perhaps, is owing partly to the Redundancy of *Atrabilious* Particles in the Blood, which naturally flowing into the inferior Branches of the *Aorta*, constantly keep open any Outlet through which they have once found the Way; partly to the large obstructed *Viscera*, compressing the *Vena Cava*, and hindering the free Return of the Fluids from the inferior Extremities. Hence we find, that Ulcers of the Legs, with black *Cicatrices* (such as may daily be seen among the Soldiers and *Spaniards*) are mentioned both by HIPPOCRATES † and CELSUS ‡, as the Effect of overgrown Spleens.

\* *Minorca es bo de Cap y mal de Camas.*

† De morb. intem.

‡ Lib. ii. Cap. vii.

And



And now it plainly appears, why Ruptures are so common in this Place; for the other Bowels being swelled beyond their natural Size, the Intestines are too much confined; and from the Nature of the Aliment, being frequently distended with Wind, it is not to be wondered at, that they often push through the Rings of the Abdominal Muscles.

In so hot and dry a Country as this, all the Parts of the Body, are very subject to topical Inflammations: But the Eyes are most particularly affected in this Way; which seems to be occasioned chiefly by the strong dazzling Light reflected, during the Summer Season, from the white Rocks and Sand; to which we may perhaps add, that the floating Particles of Salt, Dust, and minute Insects, wherewith the Air often abounds, do likewise injure the tender Texture of this Organ, and give rise to *Ophthalmia's*, or increase them.

Whatever



## INTRODUCTION.

Whatever dissipates the finer Parts of our Fluids, and increases the Proportion of Earth, and fixed Matter (and several such Causes I have already had Occasion to take Notice of) will be apt to produce Sand and Gravel in the urinary Passages; though it is probable, that the calculous Concretions and nephritick Pains are chiefly owing to the Waters; which, as I formerly observed, are mostly hard and brackish, and let fall a large Quantity of stony Sediment after boiling.

The Convulsion of the lower Jaw in Children, ought likewise to be reckoned among the sporadick Distempers. And as it is both very frequent and fatal in this Island, and some of the neighbouring Countries, I shall give a Description of it from HYACINTHUS ANDREAS, a *Spaniard*, who, about the End of the last century, published an Abridgment of RIVIERUS, under the Name of *Praxis Medica Gotholanorum*, with very little Addition of his own, excepting an Account of  
G this



## INTRODUCTION.

this Disease, which most other Authors have omitted. *In hac Urbe nostrâ Barchinonensi, affliçantur plurimi infantes, adeo feroci convulsione mandibulæ inferioris, ut eâ apprehensi, nullo possint motu illam movere, et abhinc suctus lactis impeditur omnino. Emergit hoc malum, ex causa humiditatis regionis, et potissimum si matres prægnationis tempore, minus sobrie vixerint; et usæ fuerint alimentis humidis, et potibus gelidarum eximiis: et quanquam istas duas inveniamus causas, adeo manifestas, existimo tamen potius hanc cladem insolescere, ex peculiari cæli vel astrorum influxu, quam ex illis duabus: Nam in plurimis aliis humidis regionibus, laute bibunt mulieres, et tamen non affliçantur infantes (ita attestantur medici) morbo isto diro, quemadmodum in hac nostra civitate, in quâ tot interficit mala ista convulsio, ac variolæ aut morbilli. Unde si in toto orbe preman- tur infantes unico tyranno, nempe variolis, in hac quidem civitate, duplici consliçantur; sciz. Variolis et convulsione mandibularum, quæ a nostris mulierculis et obstetricibus*



*cibus vocantur barrettas, in quarum periculum incurrunt recenter nati, usque ad nonum sui nativitatis diem, eoque transacto, omne discrimen cessare docuit semper experientia.* It is needless to add the Remedies prescribed by our Author, who ingenuously confesses, the Disease is so seldom curable, that in twenty Years Practice, he had scarce known six recover.

To these we may add the following Hurts arising from endemial Causes. The Children and Peasants are often ulcerated with the caustick milky Juice of the Fig-Trees and Spurges, which are common in the Fields: In drinking corrupted Waters, sometimes Leeches are swallowed, which I have known to occasion extraordinary Symptoms, such as Coughs, Nausea, Spitting of Blood, &c. to the great Surprize of both the Physician and Patient, who were entirely ignorant of the Cause of these Complaints. The *Pastinacæ Marinae*, and *Aquila*, wound dangerously



roufly with the Stings in their Tails; and the *Scorpius*, *Scorpæna*, and *Draco*, with the Prickles of their Back; (for which Reason the Fishermen are obliged by Law to cut off these Weapons before they bring the Fish to Market). In the hot Weather, the Viper, the Land Scorpion, and the small black Field Spider, are reckoned poisonous. In the Spring the Hedge-Hogs, *Flagrantes æstu venereo*, are said to pollute the Waters, to which they have Access, and thereby occasion a Strangury and Priapism to those who drink of them: and at this Season the Flesh of these Animals has the same Effect, tho' it is wholesome and innocent enough at other Times of the Year. But as Accidents of this Class rarely occur, it is sufficient barely to have mentioned them.

In the opinion of the Natives, no Diseases are more frequent here than Witchcraft, Charms, and evil Spirits: Those nevertheless, I shall entirely omit, having neither Leisure nor Inclination to enlarge upon  
upon



upon the Craft of the Clergy, and the Credulity of their Flocks: But whoever desires to see this Subject copiously handled, may consult the learned Father FEIJOO, who, in his useful and elaborate Volumes on vulgar Errors, has taken Occasion to expose the Tricks of those pious Jugglers, who pretend to exorcise evil Spirits, together with several other Impositions of the same Nature.

These particulars I thought it necessary to premise, before we enter on the History of the epidemical Diseases. In Excuse for whatever Errors I may have committed in this mixt Essay, I hope I may be allowed to plead, my having wrote it in a remote Corner of the World, which afforded me very little Assistance either from Men or Books.

*Qualemcunque igitur veniam dignare libellum  
Sortis et excusa conditione mee.*

OVID.

*Minorca, An. Dom. MDDCXLVII.*



INSTRUCTIONS

The first part of the Instructions is a general  
 statement of the principles which should govern  
 the conduct of the Court in the exercise of  
 its jurisdiction. It is in these principles that  
 the Court is to be guided, and it is to these  
 principles that the Court is to refer in all  
 cases where the law is not clear.

The second part of the Instructions is a  
 statement of the powers of the Court. It is  
 in these powers that the Court is to be  
 guided, and it is to these powers that the  
 Court is to refer in all cases where the  
 law is not clear.

The third part of the Instructions is a  
 statement of the duties of the Court. It is  
 in these duties that the Court is to be  
 guided, and it is to these duties that the  
 Court is to refer in all cases where the  
 law is not clear.

Approved by the Board of Directors



---

A N  
E S S A Y  
O N T H E  
E P I D E M I C A L D I S E A S E S  
I N  
M I N O R C A.


---

C H A P. I.

*Of the Weather from the Year 1744 to  
1749.*

I AM sorry it was not in my Power, for  
want of Conveniency and proper  
Instruments, to determine exactly the  
Weight of the Air, the Quantity of Rain,  
and Force of the Winds: notwithstanding  
which Omissions, I flatter myself, the

CHAP.  
I.



G 4 following



CHAP. following short Account of the principal  
 I. Variations of the Weather, taken from a  
 ~~~~~ Diary regularly kept, with very few In-  
 terruptions, will neither be entirely useless  
 nor unacceptable: And, in order to make  
 it better understood, I shall premise a few  
 explanatory Observations.

In the first Place, in speaking of Days  
 and Months, I use the old Stile, as is cus-  
 tomary among the *English* in *Minorca*;  
 though the Natives reckon by the New:  
 And where it appeared necessary to dis-  
 tinguish the Forenoon from the After-  
 noon, I have annexed a. m. or p. m. to  
 the particular Days.


Secondly, Wherever a rainy Day occurs,  
 without any Dots after it, the Reader will  
 be pleased to take Notice, that only driz-  
 zling Rain, or a slight Shower, happened  
 on that Day: But if two Dots be placed  
 after it, in this Manner . . it signifies one  
 or more smart Showers to have fallen: If  
 three Dots . . . great Rains: If four . . .  
 extreme



extreme heavy Rains, or rather Floods of  
Water.

CHAP.

I.



Thirdly, Wherever the Thermometer is mentioned, it is to be understood of a large mercurial one, graduated according to FAHRENHEIT'S Scale, and kept in a proper Place within Doors; except when I describe the Heat of the Sun's Rays; which was measured by a smaller Instrument of the same Kind, hung out at an upper Window, in the open Street, at a considerable Distance from the Walls of Houses. Both these Thermometers, being dipt in Snow, fell to the 32d Degree. Both were raised to the 96th, 97th, or 98th Degrees, by the natural Warmth of Persons in Health. Nor did I observe any other Difference between them, excepting that the Mercury in the smallest was soonest affected by Heat or Cold, and consequently a little quicker in its Motions; which occasioned its being found sometimes a Degree higher in Summer,  
and



CHAP. and as much lower in Winter, than the  
 I. Quicksilver in the large Thermometer.

Fourthly, The Height of the Thermometer, at a Medium, during each Month, is calculated from Observations, made about three o'Clock in the Afternoon, when the Mercury is commonly higher by one or two Degrees in Winter, and two or three in Summer, than in the Mornings and Evenings.

Fifthly, The Difference between the Heat of Air warmed by the direct Rays of the Sun, and that in the Shade, is at a Medium in Summer, about 13 Degrees. Whenever it was remarkably greater, Notice will be taken of it; as likewise of the most extraordinary Heights, to which the Mercury rose, when the Thermometer was exposed to the Sun, at other Times of the Year.

Sixthly, The Mercury seldom or never sinks below the 48th Degree in *Minorca*,  
 except



except when sharp Winds blow from the North, which cause the Weather to seem as intensely Cold as it is in *England*, when the Thermometer has been ten Degrees lower. And upon the Descent of heavy Rains, particularly in the Autumn, the Cold affects us much more sensibly than one would imagine, from the Alteration of the Thermometer.

CHAP.  
I.  
~

These Things being premised, I proceed to describe the most remarkable Changes of the Weather, in each Month, during the Period of Time that produced the epidemical Diseases, which are the Subject of this Treatise.

A. D. MDCCXLIV.

*January* was mild and temperate in the Beginning: But after the Rains, about the Middle of the Month, it continued, for the most Part, cold and cloudy.

A. D.  
1774. W

Rainy



CHAP. Rainy days, 1, 6, 9, .. 10; .. 11, 14,  
 I. 15, .. 19, .. with Hail, 20 with Hail,  
 24, 27, 30.  
 1744.

The first Days of *February* were cloudy, cold, rainy, and stormy: From the 4th to the 19th, the Weather was moderate, and seasonable, without excessive Rains or immoderate Cold: From which, to the 26th, it was fair and warm: But afterwards to the End of the Month, rough and stormy as in the Beginning.

*March*, during the first Week, was sometimes warm and sometimes cold: But in the three last Weeks, piercing Winds blew constantly from the North, with frequent Rains, and sometimes Hail.

The Weather continued colder, and more disagreeable than usual, till much Rain had fallen about the Middle of *April*; the Remainder of which Month was mostly fair and temperate.

The



The first Fortnight of *May* was likewise fair and temperate; the latter Part of it dry and warm, without any Rain.

CHAP.

I:

1744.

*June* was clear, calm, dry, and sultry, as this and the two following Months, are constantly observed to be; the Weather varying much less in these than in the other Months.

Rainy Day, 9.

In *July* the Northerly Winds were higher and more frequent than ordinary, infomuch, that on some Days the Cold was more troublesome than the Heat; nor do I ever remember to have felt so little of it in this Month, in *Minorca*,

Rainy Days, 6, at Night . . . . with Thunder and Lightning, 17 p. m.

Coldest Day, 8. Therm. 70. } Height at a  
Hottest Days, 21, 22, 31. 80. } Med. 76.<sup>12</sup>/<sub>32</sub>

During



CHAP. During the first Days of *August*, the  
 I. Northerly Winds continued : But the  
 1744. rest of the Month was either calm, or the  
 Breezes came from the South, the Air  
 being very hot and sultry.

Rainy Days, 2. a. m. 14. a. m.

Coldest Days, 3, 4. Therm. 73. } Height at a  
 Hottest Days, 27, 28, 29, 30. 80. } Med. 77 $\frac{2}{3}$

In *September* the Weather is always very unequal; fair Intervals, and short violent Storms reciprocally succeeding each other: But this Year the anniversary Rains were not so constantly attended with Northerly Winds, as they commonly are.

Rainy Days, 7 a. m. . . . . with Thunder and Lightning, and at Night . . . . 11,  
 at Night, . . . . 15, at Night, . . . . 18,  
 at Night, . . . . 19, at Night, . . . . 22,  
 at Night, . . . . with Lightning, 23, at  
 Night, . . . . with Lightning.

Coldest



|                        |            |               |       |
|------------------------|------------|---------------|-------|
| Coldest Days, 17, 26.  | Therm. 71. | } Height at a | CHAP. |
| Hottest Days, 1, 2, 3. | 76.        |               |       |

1744.

In the first Part of *October*, the Winds were mostly from the North or North West, the Weather variable and unsettled: But from the 14th, to the end of the Month, it was calm, warm, and fair.

Rainy Days, 2, at Night, 4, at Night, . . 5, at Night, . . 13, a. m. and p. m. . . . .

|                     |            |               |
|---------------------|------------|---------------|
| Coldest day, 14.    | Therm. 65. | } Height at a |
| Hottest Days, 1, 2. | 71.        |               |

*November* was remarkable for bad Weather. All the first Day, and the second before Noon, it blew violently from the North, with heavy Rain: From the 7th to the 22d the Weather was constantly cold, cloudy, and stormy, with Hail, Rain, and high Winds, from the North. After a few fair Days, this Month ended stormy, as it began.

Coldest



CHAP.  
I.

Coldest Day,  
Hottest Day, 1.

Th. 50.  
67. }

Height at a Med. 56.

1744.

From the First of *December*, to the 13th, the Air was cold, but for the most Part clear; the Wind North or North-West. But at that Time, shifting suddenly to North-East, a violent Storm began, and continued with great Fury to the 17th, more especially in the Nights, with Hail, Rain, and Flakes of Snow. On the 18th the Weather again cleared up; but on the 21st it once more became cold, stormy, and rainy, and, excepting one fair Day or two, it continued so to the End of the Month.

Coldest Days, 14, 15.  
Hottest Days, 26, 31.

Therm. 44.  
57. }

Height at a  
Med. 51  $\frac{1}{2}$ .

A. D. MDCCXLV.

This Year began with strong piercing northerly Winds; nor did the Coldness of the Air decrease much, before the latter End of *January*, though the Weather



ther was mostly calm and clear with Sunshine.

CHAP.  
I.

1745.

Rainy Days, 6, at Night, . . with Hail, 13, at Night, . . 15, at Night, . . 26, at Night. . . .

Coldest Day, 6. Therm. 43. } Height at a  
Hottest Days 26 to 31. 57. } Med. 52.

The first Week of *February* was pleasant and serene; the Remainder was mostly cold, cloudy, and overcast.

Rainy Days, 8, p, m. . . . 11, at Night, . . . 15, . . 21, . . . 24, a. m, . . . 25, 28, . .

Coldest Day, 26 Therm. 46. } Height at a  
Hottest Days, 7, 8. 61. } Med. 55.

*March*, this Year, excepting the first Day, which was windy, contrary to Custom, was warm, calm, and dry, without any Storms, the winds being mostly from the South or West.

Rainy Days, 22, a. m. 23, . .

H

Coldest



CHAP. Coldest Day, 1. Therm. 51. } Height at a  
 I. Hottest Days, 12. 63. } Med. 59.



1745. *April* was likewise warm and temperate, but somewhat more windy and rainy than the preceding Month.

Rainy Days, 3, 8, 23, . . . 27, 28.

Coldest Day, 17. Therm 58. } Height at a  
 Hottest Days, 25, 26. 65. } Med. 61.  $\frac{2}{3}$ .

On the 19th, Therm, in the Shade 62, exposed to the Sun 80.

The Heat of the Weather increased greatly in *May*, notwithstanding some unseasonable Rains, and northerly Winds, about the End of the Month.

Rainy Days, 1, 7, 8, . . 16, . . 23, at Night, . . . . 24, p. m. . . . . 25.

Coldest Days, 1, 2. Therm 62. } Height at a  
 Hottest Day, 17. 74. } Med. 68  $\frac{8}{11}$ .

On



On the 4th, Therm, in the Shade 65, CHAP.  
exposed to the Sun 88. I.

1745.

The Beginning of *June* was likewise unseasonable, with Rains and northerly Winds. The rest of the Month was calm, dry, and hot, as usual.

Rainy Days, 4, a. m. . . . and at Noon, . . . 7, at Night . . . .

|                       |            |               |
|-----------------------|------------|---------------|
| Coldest Days, 4 to 7. | Therm. 69. | } Height at a |
| Hottest Day, 28.      | 82.        |               |

On the 28th, Therm. in Shade 82, exposed to the Sun 98.

*July* was a little more temperate than usual, the Heat of the Air being frequently allayed by Breezes or Showers.

Rainy Days, 16, a. m. . . . 17, a. m. . . . and p. m. . . . and at Night, . . . . with Thunder, 27, p. m. . . 29, at Night, . . . . with Lightning.



CHAP. Coldest Days, 21. 22. Therm. 75. } Height at a  
 I. Hottest Days, 6, 25, 26. 80. } Med.  $77\frac{1}{2}$ .

1745.

On the 7th, Therm. in the Shade 79, exposed to the Sun 100. 25th Therm. in the Shade 80, exposed to the Sun 96.

*August*, excepting the three first Days, was excessive hot and sultry, till the latter End, when the Air was refreshed by brisk northerly Breezes.

Rainy Days, none.

Coldest Days, 1, 23 to 27. Th. 74. } Height at a  
 Hottest Day, 9. 82. } Med.  $77\frac{1}{2}$ .

On the 9th. Th. in the Shade 82, exposed to the Sun 95.

The Beginning of *September* was very sultry, though not always serene. But from the 12th to the End, it was constantly either cloudy, rainy, or stormy with violent Squalls from the North.

Rainy



Rainy Days, 12, at Night, . . . . 13, at  
 Night, . . . 14, at Night, with Thunder,  
 16, p. m. . . . . and at Night, . . . . with  
 Thunder, 17, p. m. . . . . 18, a. m. . . .  
 and at Night, . . . . with Thunder and  
 Lightning, 19, a. m. . . . . and at Night, . . . .  
 20, . . 27, . . . . at Night, with Thunder  
 and Lightning, 28, . . 30, at Night, with  
 Lightning.

CHAP.  
 I.  
 ~~~~~  
 1745.

Much Lightning in the Nights of the  
 20th, 21st, and 22d.

Coldest Day, 19. Th. 69. } Height at a Med.  $75\frac{1}{2}$ .  
 Hottest Day, 9. 80. }

On the 24th, Th. in the Shade 73,  
 exposed to the Sun 89.

The Weather, during *October*, was  
 fair, pleasant, and serene, excepting the  
 rainy Days, and some few others, the  
 Winds being mostly moderate and nor-  
 therly.



CHAP. Rainy Days, 1. a. m. 3, a. m. . . . 7,  
 I. at Night, . . . with Thunder and Light-  
 ning, 8, a. m. . . . 17, at Night, with  
 1745. Lightning, 25, 26, at Night, . . . 27. a.  
 m. . . .

Coldest Days, 29, 30. Therm. 61. } Height at a  
 Hottest Days, 1, 2. 69. } Med. 65  $\frac{20}{31}$ .

Almost all *November* was either cloudy or wet: From the Beginning to the 24th, the Wind was mostly from the South, or South-West. Afterwards it changed to the North; and the Air continued cold and damp to the End of the Month.

Rainy Days, 2, 5, 7, at Night, . . . 10,  
 a. m. . . . and p. m. . . and at Night, . . .  
 11, at Night, . . . 14, 15, at Night, . . .  
 with Hail, 21, a. m. . . . 26, . . . 27, . . .  
 30. . . .

Coldest Day, 27. Th. 50. } Height at a Med. 58  $\frac{6}{70}$ .  
 Hottest Day, 8. 65. }



The first Week of *December*, was wet and cold, with high easterly and northerly Winds. The Remainder was somewhat more temperate and calm; the Winds being, for the most part, South-West or southerly.

CHAP.  
I.  
1745.

Rainy Days, 1, . . . 2, . . . 5, at Night,  
. . . 7, p. m. . . 12, . . . 18, . . . 20, p. m.  
. . . and at Night, . . . 21. . .

Coldest Days, 5, 6, 9. Therm. 48. } Height at a  
Hottest Day, 27. 60. } Med. 53  $\frac{1}{3}$ .

## A. D. MDCCXLVI.

The greatest Part of *January* was fair and clear, with cold easterly Winds.

Rainy Days 6, . . . 10, . . . 11, a. m. . .  
23, 31, . . . with Hail.

Coldest Days, 7, 12. Therm. 48. } Height at a  
Hottest Day, 28. 57. } Med. 52  $\frac{2}{3}$ .

H 4

During



CHAP. I. During *February*, the Weather resembled that of the preceding Month, though somewhat colder, and more windy.

1746.

Rainy Days, 3, . . 6, a. m. . . . 7. a. m. . . 23, a. m. . . 28. . .

Coldest Days, 2, 3, 15. Therm. 45. } Height at a  
Hottest Days, 20. . . . . 57. } Med. 51  $\frac{1}{2}$ .

On the first of *March*, there was a remarkable Storm from the North, with Flakes of Snow in the Evening. Next Morning, the Fields were covered with Snow a Foot deep, which melted away soon after Sun-rising. But some more fell the following Night, and continued upon the Ground for three Days ere it melted: Which is so extraordinary a Phænomenon in that Part of the World, that it had never before happened above once or twice in the Memory of Man. From the 5th Day, the Cold gradually decreased to the 15th, and afterwards, to the



the End of the Month, we had fine moderate Weather. CHAP. I.

1746.

Rainy Days, 13, 14, .. 28, a. m. ...  
30, a. m. .. 31. ..

Coldest Days, 3, 4.	Therm. 42.	} Height at a Med. 54 $\frac{2}{3}$
Hottest Days, 24, 27, 30, 31.	60.	

In *April* the Weather was exceeding changeable, and often rainy or overcast.

Rainy Days, 3, at Night, ..... 4, a.  
m. .... 20, p. m. .... 11, ... with  
Hail, 13, ..... with a Storm of Wind,  
14, 17, at Night, ..... 18, a. m. ..  
23. ..

Coldest Day, 13.	Therm. 54.	} Height at a Med. 59 $\frac{2}{3}$
Hottest Day, 30.	68.	

On the 3d, Th. in the Shade 57, exposed to the Sun 83.  
29th, 65, 88.

*May* was calm, serene, and seasonable, seldom obscured with Clouds or Rain, until it drew near an End.

Rainy



CHAP.

I.

1746.

Rainy Days, 9, 21, 24, a. m. .. 29,  
30, p. m. .... with Thunder and Light-  
ning, and at Night, .... with Thunder  
and Lightning.

Coldest Day, 11.	Therm. 64.	} Height at a Med. 68 $\frac{2}{3}$ .
Hottest Days, 4, 5, 18, 20.	70.	

*June*, as usual, was dry, sultry, and  
serene.

Rainy Day, 15th.

Coldest Day, 11.	Therm. 69.	} Height at a Med. 73 $\frac{1}{2}$ .
Hottest Days, 26, 27.	79.	

The Heat gradually increased, from  
the End of *June* to the 20th of *July*,  
being very troublesome and uneasy, even  
to those, who never exposed themselves  
to the Sun: But during the latter Part  
of the Month, it was mitigated by daily  
Breezes.

Rainy Days, 28, a. m. 29, a. m. and  
at



at Night, . . . . 30, a. m. . . . . with Thunder and Lightning. CHAP. I.

Coldest Days, 8, 29, 30. Th. 77. } Height at a  
Hottest Day, 19. 87. } Med. 80  $\frac{2}{3}$ .

1746.

*August* was more temperate than the preceding Month, the Heat being somewhat allayed by the northerly Winds.

Rainy Days, 1, a. m. 19, a. m. 27, p. m. . . . . with Thunder and Lightning, 31, at Night, . . . . with Thunder and Lightning.

Coldest Day, 29. Therm. 70. } Height at a  
Hottest Day, 15. 81. } Med. 76  $\frac{1}{3}$ .

*September* began with Storms of Wind and Rain; from the 4th to the 14th it was clear and warm; from which to the End of the Month the Sky was constantly either darkened with heavy Rains, or was tempestuous with strong northerly Winds; the Weather being much colder than usual at that Season of the Year.

Rainy



CHAP. I. Rainy Days, 2, p. m. . . . . with  
 1746. Thunder and Lightning, 3, a. m. . . . .  
 with Thunder and Lightning, and Hail-  
 stones of an Inch Diameter, 15, a. m. . . . .  
 16, a. m. . . and at Night, . . . . 17, . . . .  
 with Thunder and Lightning, 18, . .  
 19, . . 20, at Noon, . . . and at Night, . . . .  
 22, p. m. . . 23, at Night, . . . . 24, . .  
 and at Night, . . . 26, and at Night, . . . .  
 28, at Noon. . . . .

Coldest Days, 22, 26, Therm, 58. } Height at a  
 Hottest Day, 13. 74. } Med. 67  $\frac{2}{3}$

The first and last Week of *October*,  
 were mostly pleasant and serene: The  
 two intermediate Weeks were rainy and  
 over-cast; and the whole Month was  
 uncommonly cold, with northerly Winds.

Rainy Days, 8, p. m. and at Night, . . . .  
 9, at Night, . . . 10, . . . at Night, . . . .  
 14, p. m. . . 18, 20, p. m. . . . . 21, . . . .  
 with Thunder and Lightning, 22,  
 P.



p. m. . . . . and at Night, . . . . 30, CHAP.  
a. m. . . . . with Hail. I.

1740.

Coldest Days, 8, 9. Therm. 54. } Height at a  
Hottest Day, 1. 68. } Med. 58  $\frac{2}{3}$ .

During *November* the Weather was calm and seasonable, with much Sunshine on the Days free from Rain.

Rainy Days, 1, at Night, . . 2, . . .  
and at Night, . . . . 3, at Night, . . . .  
4, . . . 5, . . 12, at Night, . . . 14, . .  
17, a. m. . . 19, a. m. . . 26, p. m. . . .

Coldest Day, 20. Therm. 54. } Height at a  
Hottest Days, 28, 29. 62. } Med. 57  $\frac{1}{3}$ .

In *December* there was no extraordinary cold Weather, and very seldom high Winds; nor was the Serenity of the Sky much interrupted by Clouds of Rain.

Rainy Days, 14, . . . 15, . . 29, at  
Night, . . . 30, . . 31, . . . and at  
Night. . . . .

Coldest Days, 11 to 15. Th. 54. } Height at a  
Hottest Day, 59. } Med. 56.

A. D.



CHAP.  
I.

A. D. MDCCXLVII.

1747.

The Weather of *January*, was like that of the preceding Month, but somewhat more wet and cold in the Beginning.

Rainy Days, 1, a. m. . . . 2,  
p. m. . . and at Night . . . 4, at  
Night, . . 12, . . 14. . .

Coldest Day, 5.  
Hottest Days, 20, 28.

Th. 50. } Height at a  
59. } Med. 55  $\frac{4}{7}$ .

During the first Fortnight of *February*, we had warm, fair, Summer-like Weather: The third Week was rainy and tempestuous, with some Hail and Snow; the fourth very changeable, the Sun-shine being often interrupted with Showers.

Rainy Days, 16, p. m. . . . and at  
Night, . . . . 17, a. m. . . and p. m. . .  
with Hail and Snow, 19, at Night, . . .  
21, . . 24 at Night, . . 25, 27, 28.

Coldest



Of the WEATHER.

III

CHAP.

I.

1747.

Coldest Day, 18.  
Hottest Days, 5, 10.

Therm. 50. } Height at a  
62. } Med. 55 $\frac{20}{28}$

On the 4th, Therm. in the Shade 61,  
exposed to the Sun 78.

From the Beginning of *March*, to the  
21st, the Weather was cold, and often  
wet; the Winds mostly from the North,  
and sometimes very boisterous: The  
Remainder of the Month was fair and  
moderate, with westerly or South-West  
Winds.

Rainy Days, 1, 2, . . . 3, . . . and at  
Night, . . . 9, p. m. 14, a. m. . . 17, at  
Night, . . . 19, a. m. . . 20, a. m. . .  
with Hail.

Coldest Days, 6, 13 to 17.  
Hottest Day, 30.

Th. 50. } Height at a  
63. } Med. 54 $\frac{17}{27}$ .

*April* was fair and dry, but sometimes  
very windy; in the fore Part of the  
Month especially; and the Wind being  
then at North, did great Damage to the  
Fields



CHAP. Fields and Vineyards. About the Middle  
 I. and to the End, the South-East pre-  
 vailed.  
 1747.

Rainy Days, 9, at Night . . . 10, at  
 Night. . .

Coldest Days, 2, 3. Therm. 56. } Height at a  
 Hottest Days, 27 to 30. 68. } Med. 61 $\frac{2}{3}$ .

Betwixt the 1st and the 9th of *May*,  
 the Weather was often disagreeable, with  
 Rain, Clouds, or much Wind. But al-  
 most all the rest of the Month was clear,  
 calm, and warm.

Rainy Days, 1, 6, p. m. . . . 7,  
 p. m. . . . with Thunder and Light-  
 ning, 8, at Noon, . . . 28, p. m. with  
 Thunder and Lightning, 29, a. m. . .

Coldest Days, 1, 7 to 10. Th. 67. } Height at a  
 Hottest Days, 31. 75. } Med. 71.

On the 4th, Therm. in the Shade 70,  
 exposed to the Sun 88.

*June*



*June* was hot and dry as usual. On the 27th of the Month, dark Clouds suddenly rose from the North, a little before Sun-set, and a Storm of Wind succeeded, which lasted all that Night.

Rainy Day, 23, a. m.

Coldest Day, 23. Th. 71. } Height at a Med. 76.  
Hottest Days, 26, 27. 80. }

On the 24th, Th. in the Shade 71, exposed to the Sun 94.

During *July*, hot suffocating Winds with unwholesome Vapours, from the South and East, were predominant.

Rainy Days, 6, a. m. . . . 7, a. m. . . .  
18, at Night, . . . . 26, p. m. . . .

Coldest Day, 8. Th. 73. } Height at a Med. 79½.  
Hottest Day, 21. 84. }

On the 12th, Therm. in Shade 80, exposed to the Sun 99. On the 21st,  
I Therm.



CHAP. Therm. in the Shade 84, exposed to the  
I. Sun 100.

1747.

The Beginning of *August* was intolerably hot; nor was the rest much more temperate, though it rained plentifully about the Middle and End of it, the Winds being most commonly from the East and South-East.

Rainy Days, 9, p. m. 12, at Night, . . . .  
with Thunder and Lightning, 15, at  
Night, . . . . with Thunder and Light-  
ning 16, p. m. . . . . with Thunder and  
Lightning, 28, at Night, . . . . 29, a. m.  
with Thunder and Lightning, 30, a. m.  
. . . . and much Lightning at Night.

Coldest Day, 30. Therm. 74. } Height at a  
Hottest Days, 8, 10, 11. 84. } Med. 80  $\frac{4}{11}$ .

On the 8th, Therm, in the Shade 84,  
exposed to the Sun 100. On the 15th,  
Therm. in the Shade 79, exposed to the  
Sun 98.

The



The slight Showers, which fell first in *September*, rather increased than allayed the Warmth of the Air: But the Storms of Wind and Rain about the Middle of the Month put an End to the hot Weather for that Season.

Rainy Days, 8, 11, a. m. 13, p. m. and at Night, . . . . 14, . . . . and at Night, . . . . 18, at Night, . . . with Thunder and Lightning, 19, . . . and at Night, . . . . 20, . . . 21, at Noon, . . . with Hail, 23, p. m. and at Night, . . . 24, a. m. . . . and at Night, . . . . 25, . . 28, p. m. . . . and at Night, . . . . with Thunder and Lightning, 29, . . . and at Night, . . . . 30, a. m. . . . .

Coldest Day, 21.	Th. 63.	} Height at a Med. 72 $\frac{22}{30}$ .
Hottest Day, 10.	81.	

On the 1st, Th. in the Shade	79,	exposed to the Sun	98.
2d,	80,		100.
10th,	81,		98.

Excepting a Shower or two, and a Storm of Wind from the North-East on



CHAP. the 21st, *October*, was entirely calm and  
I. serene.

1747. Rainy Days, 4, p. m. . . and at Night, . .  
17, p. m. . . 19, p. m. . . 21.

Coldest Days, 19, 25, 27, 28.	Th. 62.	} Height at a Med. $65\frac{1}{2}$ .
Hottest Days, 10, 11, 12, 14.	70.	

On the 2d, Th. in the Shade 67,	exposed to the Sun 90.
9th,	69, 86.
23d,	64, 83.
26th,	63, 88.

Nor was *November* less serene, though  
sometimes more windy, especially at the  
Close of the Month; when it blew vio-  
lently from the North.

Rainy Day, a. m. . . . with Hail.

Coldest Day, 27.	Therm. 51.	} Height at a Med. $60\frac{2}{3}$ .
Hottest Day, 2.	67.	

On the 16th, Th. in the Shade 59,	exposed to the Sun 84.
22d,	62, 82.
27th,	51, 74.
28th,	54, 78.

From the 1st to the 6th of *December*  
the Wind blew strong from the South-  
West;



West; as it did from the South-East, from the 20th to the 24th: The other Days of this Month were mostly fair, calm, and agreeable; the whole of the Month was uncommonly warm for the Season, excepting the last Day, when the Wind changed to the North.

CHAP.  
I.  
1747.

Rainy Days, 8, a. m. . . . and at Night, . . 9, 12, 14, . . with Thunder and Lightning, and at Night, . . 17, . . 24, a. m. . . . 25, p. m. . . . 28, p. m. . .

Coldest Day, 31.  
Hottest Days, 3, 5.

Therm. 53. } Height at a  
64. } Med. 58  $\frac{1}{3}$ .

On the 5th, Th. in the Shade 64, exposed to the Sun 88.  
6th, 63, 84.

A. D. MDCCXLVIII.

The sharp northerly Wind, which began on the last of *December*, was attended with some Hail and Snow on the 4th, and continued to the 6th of *January*: It then shifted to the North-West, and

I 3                      though



CHAP. though the Weather was mostly fair and  
 I. dry, yet the Air continued cold till the  
 1748. Rains fell, near the latter End of the  
 Month.

Rainy Days, 2, a. m. . . and p.  
 a. m. . . 5, a. m. . . . 6, at Night, . . 15,  
 at Noon, . . 18, at Night, . . 22, at  
 Night, . . . 23, and at Night, . . . 25,  
 p. m. . .

Coldest Day, 4.  
 Hottest Day, 22.

Therm. 43. } Height at a  
 57. } Med.  $51\frac{1}{2}$ .

On the 8th, Th, in the Shade 44, exposed to the Sun 64.  
 18th, 50, 68.

The Weather continued mild and tem-  
 perate to the 17th of *February*, tho' the  
 Rains that fell in the second Week were  
 accompanied with high southerly, and  
 South East Winds: Afterwards, to the  
 End of the Month, we had dry, cold,  
 North, and North-West Winds, which  
 did much Mischief to the Corn.

Rainy Days, 6, at Night, 8, 9, . . 10,  
 a. m.







CHAP. p. m. . . . and at Night, . . . . 20, p. m.  
 J. 21, 30, p. m.

1748.

Coldest Days, 19, 21. Therm 50. } Height at a  
 Hottest Days, 27, 29, 30. 63. } Med. 55  $\frac{2}{3}$ .

On the 14th, Th. in the Shade 51, exposed to the Sun, 76.  
 27th, 63, 82.

*April* was cold and stormy from the 14th to the 20th; all the rest of the Month was calm, clear, and warm, excepting the 25th, when it rained, and blew fresh from the North-West.

Rainy Days, 6, at Night, 14, p. m. . . . with Thunder, 15, at Night, . . . . 16, p. m. . . . with Thunder and Lightning, and at Night, . . . with Thunder and Lightning, 19, 24, at Night, . . . 25, . . . 30.

Coldest Day, 16. Th. 55. } Height at a  
 Hottest Days, 24, 28, 29, 30. 65. } Med. 61  $\frac{1}{2}$ .

*May* was clear, calm, and warm, excepting  
 ing



ing the 15th and 16th Days, when it blew violently from the North.

CHAP.  
I.

1748.

Rainy Days, 20, p. m. 21, p. m.

Coldest Day, 16. Therm. 61. } Height at a  
Hottest Days, 25, 30, 31. 72. } Med. 68  $\frac{28}{31}$ .

About the Middle of *June*, the North-East Winds blowing fresh for some Days, allayed the Heat of the Air, but did great Damage to the Fruit Trees.

Rainy Day, 26, at Noon. . . . .

Coldest Day, 18. Therm. 71. } Height at a  
Hottest Days, 12, 13. 80. } Med. 75  $\frac{28}{30}$ .

The first Weeks of *July* were extremely hot and sultry, especially from the 6th to the 16th; the Mercury in the Thermometer rising daily above the 80th Degree. Nor was the latter Part of it much more temperate, tho' it blew from the North pretty briskly on the 17th and 18th, and moderately about the End of the Month.

Rainy



CHAP.  
I.

1748. Rainy Days, 2, a. m. 16, at Night,  
17th, at Night, 18, at Noon, . . . . 31, at  
Night, . . . with Thunder and Lightning.

Coldest Days, 18.	Therm. 74.	} Height at a Med. 71 $\frac{5}{11}$ .
Hottest Days, 14, 15, 16.	84.	

On the 5th, Th. in the Shade 79,	exposed to the Sun 96.
7th,	81, 100.
23d,	81, 98.

The excessive Heat of the Season continued to the 19th of *August*; then it intermitted for some Days, the Clouds threatening Rain, with boisterous Winds from the North, especially in the Night-time. On the 24th it again became sultry, and continued so to the End of the Month.

Rainy Days, 3, a. m. 7, at Night, 19, a. m. with Thunder and Lightning, and at Night. . . . .

Coldest Days,	Therm. 74.	} Height at a Med. 79 $\frac{5}{11}$ .
Hottest Days, 15.	85.	

On



On the 9th, Th. in the Shade 77, exposed to the Sun 95.  
 10th, 79, 96.  
 14th, 84, 101.  
 15th, 85, 100.

CHAP.  
 I.  
 1748.

In *September*, though the Sky was often over-cast, with Signs of Rain, yet the Showers were neither so heavy nor so frequent as they commonly are about this Time of the Year: But large Dews falling in the Nights, and constant Breezes from the North, rendered the Air temperate, as it commonly is in this Month.

Rainy Days, 8, at Night, 9, at Night, with Thunder and Lightning, 17, at Night, 18, at Night, 23, at Noon, . . . . and at Night, . . . . 24, a. m. . .

Coldest Day, 25. Th. 66. } Height at a  
 Hottest Day, 7, 8. 80. } Med. 73  $\frac{1}{3}$ .

On the 22d, Th. in the Shade 73, exposed to the Sun 96.

The first Ten Days in *October*, were mostly



CHAP. mostly fair, clear, and calm, and also the  
 I. last Week of the Month: All the inter-  
 1748. mediate Days were either wet, or windy  
 and overcast.

Rainy Days, 1, at Noon, . . . 9, p. m.  
 10, p. m. . . and at Night, . . . . 11, . . . .  
 12, at Night, . . . . 14, . . . . 15, . . . 19,  
 20, at Night, . . . . 21, at Night, . . . 22,  
 23, at Noon, . . . . 27, at Noon. . . .

Coldest Days, 12, 13. Therm. 58. } Height at a  
 Hottest Day, 7. 72. } Med.  $65\frac{1}{2}$ .

Towards the Beginning and End of  
*November* the Weather was temperate  
 and serene. During the middle Part of  
 the Month it was frequently cloudy and  
 cold with dry North Winds.

Rainy Days, 8, at Noon, . . . . with  
 Thunder and Lightning, and at Night,  
 . . . . 11, a. m. 20, . . . . 22, 26, . . . .

Coldest Day, 23. Therm. 53. } Height at a  
 Hottest Days, 1, 2, 3. 66. } Med.  $58\frac{3}{8}$

In



CHAP.

I.

1748.

In *December* the Wind was either south-erly or westerly : Hence the Weather was never cold ; and though some Days were foggy, yet by much the greatest Part of this Month was dry, clear, and very pleasant.

Rainy Days, 14, . . . at Night, 21.

Coldest Day, 9 to 17.	Therm. 56.	} Height at a
Hottest Days, 27, to 31.	60.	

A. D. MDCCXLIX.

So mild was the Weather in *January*, that it scarce seemed to be Winter. And the preceding Autumn having been uncommonly dry, and the Showers which fell during the three first Months of this Year, being neither frequent nor plentiful, the Corn in many Places perished for want of Rain.

Rainy Days, 4, a. m. 7, at Noon, and  
at



CHAP. at Night, . . . 18, p. m. 19, at Night,  
I. 20, at Noon.

1749.

Coldest Days, 6, 7.	Therm. 53.	} Height at a Med. 55 $\frac{3}{4}$ .
Hottest Days, 1, 2, 16, 17.	58.	


*February* was likewise mild and temperate; the Mercury in the Thermometer fluctuating between the 54th and 57th Degree: A little Rain fell on the first Days, a heavy Shower on the 12th, slighter Showers on the 22d and 23d; during the rest of this Month the Air was clear and dry.

*March* was fair, warm, and without Rain, both towards the Beginning and latter End; the Thermometer standing above the 60th Degree. Between the 9th and 22d Day, frequent Storms of Rain, with some Hail intervened, which sunk the Mercury to the fifty second.

The Weather, during almost all *April*, was mild, clear, and agreeable.

Rainy





Rainy Days, 7, a. m. . . . and p. m. . . .  
10, p. m. . . . and at Night. . . .

Coldest Days, 8, 9. Therm. 56. } Height at a  
Hottest Day, 30. 69. } Med. 62  $\frac{2}{3}$ .

In *May*, sudden heavy Rains sometimes interrupted the Serenity of the Weather; and at the Close of the Month, a strong westerly Wind did much Damage to the Vineyards. The Harvest this Year was so poor and scanty, especially in the Northern Parts of the Island, where the Droughts were most excessive, that they scarcely reaped as much Corn as they had sown.

Rainy Days, 3, p. m. . . . 11, at Noon, . . . 12, at Noon, . . . 16, 19, 25, at Night, . . . 31, at Night, . . . with Thunder and Lightning.

Coldest Day, 4. Therm. 64. } Height at a  
Hottest Days, 24, 25. 73. } Med. 68  $\frac{1}{2}$ .

During



## CHAP.

## I.

1749.

During the Month of *June* the Air was dry and excessive hot, though somewhat more temperate between the 14th and the 25th; frequent Gales coming from the North.

Rainy Day, 15, p. m.

Coldest Day, 1.	Therm. 69.	} Height at a Med. 74 $\frac{3}{8}$ .
Hottest Day, 30.	79.	

On the 29th, Th. in the Shade 76, exposed to the Sun 96.

The Weather has seldom been felt so extremely hot and sultry as this Year in *July*; the Quicksilver rising daily above the 80th Degree of the Thermometer, and never falling below the 79th, even in the Night-time, till the Rains which fell in the latter Part of the Month had cooled the Air.

Rainy Days, 25, at Noon, . . . 26, p. m. . . . .

Coldest



Coldest Days, 26, 27. Therm. 77. } Height at a  
 Hottest Day, 20. 86. } Med.  $82\frac{6}{31}$ .

1749.

On the 3d, Th. in the Shade	82,	exposed to the Sun	102.
4th,	84,		100.
13th,	84,		104.
14th,	85,		104.

In the Beginning of *August*, a general Relief of all his Majesty's Troops in the Island, obliged me to embark for *Ireland*, and put an End to these Observations.





A Table shewing the greatest, least, and mean Height of the Mercury at Three o'Clock in the Afternoon, in each Month, from the Year 1744 to 1749.

		1744.	1745.	1746.	1747.	1748.	1749.
January.	G		57	57	59	57	58
	L		43	48	50	43	53
	M		52	52 $\frac{23}{31}$	55 $\frac{4}{31}$	51 $\frac{15}{31}$	55 $\frac{3}{31}$
February.	G		61	57	62	60	57
	L		46	45	45	48	54
	M		55	51 $\frac{16}{28}$	55 $\frac{20}{28}$	55	
March.	G		63	60	63	63	64
	L		51	42	50	50	52
	M		59	54 $\frac{5}{31}$	54 $\frac{17}{31}$	55 $\frac{9}{31}$	
April.	G		65	68	68	65	69
	L		58	54	56	55	56
	M		61 $\frac{24}{30}$	59 $\frac{22}{30}$	61 $\frac{22}{30}$	61 $\frac{15}{30}$	62 $\frac{23}{30}$
May.	G		74	70	75	72	73
	L		62	64	67	61	64
	M		68 $\frac{8}{31}$	68 $\frac{3}{31}$	71	68 $\frac{7}{31}$	68 $\frac{1}{2}$
June.	G		82	79	80	80	79
	L		69	69	71	71	69
	M		73 $\frac{1}{2}$	73 $\frac{12}{30}$	76	75 $\frac{28}{30}$	74 $\frac{6}{30}$
July.	G	80	80	87	84	84	86
	L	70	75	77	73	74	77
	M	76 $\frac{19}{31}$	77 $\frac{12}{31}$	80 $\frac{25}{31}$	79 $\frac{7}{31}$	79 $\frac{5}{30}$	82 $\frac{6}{31}$
August.	G	80	82	81	84	85	
	L	73	74	70	74	74	
	M	77 $\frac{2}{21}$	77 $\frac{25}{31}$	76 $\frac{15}{31}$	80 $\frac{4}{31}$	79 $\frac{4}{31}$	
Septemb.	G	76	80	74	81	80	
	L	71	69	58	63	66	
	M	73 $\frac{2}{30}$	74 $\frac{12}{30}$	67 $\frac{7}{30}$	72 $\frac{22}{30}$	73 $\frac{21}{30}$	
October.	G	71	69	68	70	72	
	L	65	61	54	62	58	
	M	68 $\frac{5}{24}$	65 $\frac{20}{31}$	58 $\frac{21}{31}$	65 $\frac{21}{31}$	65 $\frac{21}{31}$	
Novemb.	G	67	65	62	67	66	
	L	50	50	54	51	53	
	M	56	58 $\frac{6}{30}$	57 $\frac{17}{30}$	60 $\frac{12}{30}$	58 $\frac{3}{30}$	
Decemb.	G	57	60	59	64	60	
	L	44	48	54	53	56	
	M	51 $\frac{1}{2}$	53 $\frac{10}{31}$	56	68 $\frac{10}{31}$	57 $\frac{13}{31}$	



## C H A P. II.

*Of the Rise, Progress, Declension, and Succession of the epidemical Diseases, from the Year 1744 to 1749.*

A. D. MDCCXLIV.

I N the Beginning of the Year, few Diseases appeared, excepting some Pleurifies and Tertian Fevers; both which are common in *Minorca*, and are frequently met with in all Seasons; the first being generally the chief amongst Vernal Epidemics, as the other is constantly amongst the Autumnal.

CHAP.  
II.

A. D.  
1744.

During the cold Weather in *March*, many Adults were affected with Catarrhs; and almost all the Children were seized with a smart Fever, attended by a Cough:

K 2

Which



CHAP. Which Disorders did not cease, till the  
 II. Beginning of Summer.

1744.

As the Summer and Autumnal Weather of one Year, never varies much from that of another, so the same Tribe of Distempers return regularly with the Seasons, and succeed each other in the following Order.

Towards the End of *June*, the young Children (who constantly suffer first by excessive Heat or Cold) are attacked with a Vomiting, Purging, and periodical Fever, often of the erratic Kind, without any fixed Type.

In the following Month, Tertian Fevers of various Forms appear among People of all Ages, and spreading from one to another, by Contagion \*, they conti-

\* These tertian Fevers have as good a Right to be called contagious, as the Measles, Small Pox, or any other Disease; for although, in that Season, there certainly is a peculiar Disposition in the Air to affect Numbers in the same Way, yet those who are much conversant among the Sick are most liable to catch the Distemper.



nue to increase daily till about the time of the Autumnal Equinox; when they rage with the utmost Fury amongst Persons of all Ranks and Constitutions, whether Natives or Foreigners. Afterwards they gradually decline; and as soon as the Winter begins, (their Contagion being rendered inactive by the Cold) few of those who have hitherto escaped are infected: yet some primary Fevers continue to *January*, and Relapses are exceeding frequent in the latter Months of the Year.

CHAP.

II.

1744.

About the Time when the Tertians begin, the *Cholera Morbus*, Rash, and *Effere*, become frequent and epidemical in a lesser Degree: But are seldom met with after *September*, whilst the Tertians continue till Winter.

Diarrhœa's, Dysenteries, and Tenesmus, likewise make their Appearance as Epidemicks, in Summer and Autumn: But some Years they occur so seldom, as scarce



CHAP. to deserve that Name; whilst in others  
 II. they are almost as numerous as the Ter-  
 1744. tians themselves.

There seems likewise to be a near Alliance amongst all the Diseases above-mentioned. Those who have the Rash, or *Effere*, to a great Degree, are very liable to Tertian Fevers; on the other hand, in the Paroxysms of Tertians, these cutaneous Eruptions are apt to break out. The *Cholera Morbus* sometimes hath its regular Periods, like a Tertian, as the Paroxysms of Tertians are frequently attended with a *Cholera*. Sometimes a Tertian is changed into a Dysentery; or a Dysentery becomes a Tertian; and when one of these Diseases is suppressed, the other often ensues; nor is it uncommon for Dysenterick Fevers to put on the Form of Tertians; and for the Fits of Tertians, to be regularly accompanied by Gripes and Stools.

It is remarkable, that both Dysenteries and Tertians, without any manifest Cause,  
 are



are sometimes more universal and severe in one Part of the Island in one Year, than another ; and often seem, as it were, to attack particular Families with uncommon Severity ; whilst others in the same Place, the same Circumstances and Way of Life, escape. Those, however, who live in low Vallies, or near stagnating corrupted Waters, are commonly the greatest Sufferers.

CHAP.

II.

1744.

To me it appears probable, that all the Summer and Autumnal Diseases are the Consequence of Nature's attempting to free the Body from noxious Humours, either by throwing them upon the Skin, or by conveying them through the Liver and other Organs of Secretion which open into the Intestines. And if, with the Ancients, we allow distempered Bile to be the Cause of Tertian Fevers, it will perhaps be easier to account for their more obvious Appearances, than by supposing them to arise from a Lentor in the smaller

K 4

Arteries,



CHAP. Arteries, according to the modern Theory :  
 II. But this is transiently offered as a Con-  
 ~~~~~  
 1744. jecture ; to record Facts without framing  
 any Hypothesis, being my principal  
 Design.

In *July* 1744, Tertian Fevers were numerous ; but during the two ensuing Months, they were neither so universal nor so malignant, as they commonly are at this Time of the Year : Yet obstinate Relapses were frequently met with till *January*.

A little before the Equinox, Dysenteries of a bad Kind began to appear ; and being propagated by Contagion, like the Tertian Fevers, they soon encreased to such a Degree, that it was doubtful which of the two Distempers were most universal. This Year, in the Space of three Months, I had more Patients with Bloody-Fluxes, than in all the former Part of my Life ; and as no manifest Quality in the Air, or Variation in the Weather, seemed capable  
 of



of producing them, it is probable that their uncommon Frequency and singular Severity was chiefly owing to the four damaged Wine, which, for the Sake of Gain, was at that Time sold in the Island by those whose Duty it was to have prevented such Abuses. But be this as it will, it is certain that incredible Havock was made by these Distempers among the lower Sort of the Natives, as well as among the Soldiers and Sailors of his Majesty's Fleet, which was then in the Harbour; the most Part of those who were attacked died about the Winter Solstice or sooner; the rest continued weak, lean, and pale, like so many shadows, during the Winter and Spring; nor did any Method we could suggest, entirely protect them from the Torments of those painful Distempers, until the Heat of the Summer, by increasing Perspiration, had diminished the Tendency of the Humours to the Intestines.

CHAP.

II.

1744.

In *December* some were cut off by the  
Pleurisy;



CHAP. Pleurisy; and many worn out with Age,  
 II. Intemperance, or frequent Sickness, died  
 1744. about the End of this Month and the  
 Beginning of *January*.

A. D. MDCCXLV.

About the latter End of *January*, a manifest Alteration to the better was perceptible among the Sick. Those who had been reduced to the last Extremity, by repeated Relapses of Tertians, were now restored to their former Health: And such as were ill of Fluxes, and had been despaired of, began to shew Signs of Recovery: Nor did any new Distemper, excepting a few Pleurisies, appear during the Spring.

Towards the End of *May*, and Beginning of *June*, heavy Rains having occasioned a sudden Change in the Air from Heat to Cold, some were attacked with a Looseness, Gripes, and Cholick Pains: And others with an *Angina*.

*June*



*June* was scarce ended, when the Tertian Fevers and *Cholera Morbus* began; and increasing daily according to Custom, came to their greatest Degree of Frequency in *September*; from which Time they gradually declined, and seldom appeared after the Winter Solstice. These Distempers, it was observed, were this Year often attended with fixed Pains of the Side, and sometimes with a Spitting of Blood.

In *July* and *August* a slight Jaundice, without a Fever, which soon yielded to Purgatives, and saponaceous Medicines, was a common Distemper.

In *September*, Dysenteries broke out, and continued to the Winter: But they were neither so numerous nor so malignant as in the preceding Year.

As the Dysenteries and Tertian Fevers disappeared, the Pleurisy, which had been fatal



CHAP. fatal to a few in the Autumn, became  
 II. more frequent; and towards the Close of  
 the Year, it raged more violently than I  
 had ever known, at least among the *En-*  
 1746. *glish* Inhabitants of the Island.

A. D. MDCCXLVI.

I come now to a Year remarkable for dreadful Distempers and much Mortality. For the Pleurisy above-mentioned continued to make great Destruction till *April*: At which Time it began to abate; and vanished entirely about the Summer Solstice. Together with the Pleurisy, two other Diseases equally fatal, the *Phrenitis* and *Paraphrenitis*, were also epidemical: And so likewise was an Erysipelatous Fever. A few had large Tumours of the Parotids; and others had Inflammations of the Throat. But these Diseases were peculiar to Adults. Amongst the Children, obstinate, and often fatal Chin-coughs, became frequent in *March*, and continued till Summer: Some were suddenly suffocated by an *Angina*, without any appa-  
 rent



rent Swelling: And to compleat the Measure of Evils, the Island was this Year visited with the Small Pox, and of the most pestilential Kind.

CHAP.  
II.  
1746.

The Chin-cough had no sooner disappeared than a periodical Fever, accompanied with vomiting and purging, took its Place, which proved alike fatal to many Children during the Summer.

In *July* the Tertain Fevers broke out as usual, and their Violence being augmented by the excessive Heat of the Season, many Persons died suddenly about the seventh Day of the Distemper: But the cold Weather of *September* prevented their becoming so general and continuing so long as they commonly do; for about the End of *October* they gave way to Catarrhs and Quartan Agues.

A. D. MDCCXLVII.

The first Part of this Year, though not very unhealthy, produced some few Intermittents,



CHAP.  
II.  
1747.

mittents, together with inflammatory and catarrhal Fevers. And the extraordinary Heat of *May* ushered in the Summer Diseases somewhat sooner than usual. For about the End of that Month the *Cholera Morbus* carried off many Children, and in *June* the Tertians became universal. Besides, towards the End of *June*, the Diarrhœa, Dysentery, and Tenesmus broke out and raged violently for some Weeks: But contrary to Expectation, they continued only a short Time, and after the Beginning of *September*, could scarcely be reckoned epidemical.

In the mean while, as the Dysenteries decreased, the Number of Tertians were multiplied, and being of a malignant Kind, great Mortality ensued; especially in the Southern Parts of the Island. General WYNYARD's Regiment, which was quartered in *Mabon*, was so severely handled by these Fevers, that between *June* and *November* One and Forty Men died; and the most Part of those who survived,  
continued



continued weak, sickly, and ill-coloured till the Spring.

CHAP.  
II.

1747.

In *October* a few complained of Coughs, and sore Throats: And the latter End of the Year produced Pleurifies; which (as it commonly happens) did not affect the English so much as the *Spaniards*.

A. D. MDCCXLVIII.

Pleurifies were exceedingly mortal in the Beginning of the Year, and carried off divers in the Spring.

In *March* many erysipelatous Phlegmons were observed to break out on the Extremities, which generally terminated in Abscesses.

Towards the Beginning of *April*, some complained of sore Throats and rheumatic Pains. And about the 10th Day of the Month, there appeared suddenly a  
catarrhal



CHAP.  
II.  
~  
1748.

catarrhal Fever, which for three Weeks raged so universally, that almost every body in the Island was seized with it. This Disease exactly resembled that which was so epidemical in the Year 1733\*. For in most Part of the Sick, the Feverish Symptoms went off with a plentiful Sweat in two or three Days; while the Cough and Expectoration continued some Time longer. In a few athletick Persons, who were not blooded in Time, it terminated in a fatal Pleurisy or Phrensy. In one or two, the Vessels on the Coats of the Intestines burst, and the Patients expired after voiding an incredible Quantity of Blood by the *Anus*.

The ensuing Summer proved very unhealthy to the Children; many of them dying of the *Cholera Morbus*, and periodical Fevers; whilst others were much affected with cutaneous Eruptions of different Sorts.

The Tertians began in *July*, and finished

\* Med. Essays, Vol. II. Art. 2.



nished their annual Career about the End of *November*; many Persons died suddenly during the Dog Days, and about the Equinox.

CHAP.  
II.  
1748.

In the mean Time Dysenteries prevailed in *Ciudadella* among the lower Sort of the Natives, whose Bread was made of damaged Wheat, without affecting the Soldiers and others in the same town, who lived on better provisions.

From the End of *November* to the Middle of *January*, Pleurifies were predominant in all Parts of the Island; and large critical Abscesses, together with various Kinds of cutaneous Eruptions, were more frequent both among the *English* and *Spaniards* than had ever been observed.

A. D. MDCCXLIX.

That Part of this Year, during which I resided in *Minorca*, was remarkable for

L

dry



CHAP. dry Weather and few Distempers. How-  
 II. ever, in *March* Pleurifies and catarrhal  
 Fevers now and then appeared; in *April*  
 1749. some had *Anginas* with *Aphtæ* in the  
 Mouth, and spontaneous Salivations;  
 others complained of Pains in the Ears,  
 and Eruptions on the Face. In *June* and  
*July* there were some Specimens of the  
 Summer's Diseases, but so few that they  
 scarcely deserve to be called epidemical.





## C H A P. III.

## Of TERTIAN FEVERS.

HAVING described the State of  
 the Weather, and the Succession of  
 the epidemical diseases, I come now to  
 treat particularly of each, beginning with  
*Tertian Fevers*, which are, by much, the  
 most frequent of any.


CHAP.  
 III.

These Fevers never proceed uniformly,  
 with constant uninterrupted Violence  
 from their first beginning to their final  
 Termination; but, sooner or later, have  
 periodical Intervals, more or less evident,  
 so that the Patient is by Turns one Day  
 better and another worse. They are  
 called *Tertians*, or Fevers of the Tertian  
 Kind, because each particular period or  
 Revolution of the Disease (which the  
*Romans* expressed by the Word *Circuitus*)

L 2

is



CHAP.  
III.

is compleated in the Space of about Forty-eight Hours; and a new Attack is begun on the alternate Days, or every third Day, if we include the Day on which the genuine Paroxysms happen, together with the intermediate one, according to the common Method of Reckoning, used by Physicians,


The Fevers belonging to this Tribe assume so many, and such various shapes, that though they are essentially the same, yet their Appearance is often very different; insomuch that when I reflect on the several Species I have seen, I almost despair of being able to give the Reader any tolerable Idea of them. Nevertheless, as it is of such Importance to be acquainted with the specifick Properties of each, that without an exact Knowledge of them, we can neither foretell the Return of the Fits or Intervals, nor consequently administer Food or Medicines at the most convenient Hours; I flatter myself it will not be thought unnecessary to dwell a little on  
this



this Subject, and endeavour to describe the particular Types and Forms, in which the various Kinds of Tertians appear, especially as no Author who has fallen into my Hands, is sufficiently clear and accurate on this Head. For these Distempers are slightly past over by most of the Moderns; and although the *Greeks* and *Arabians* had treated of them at great Length, yet in their voluminous Writings we do not find them described as they really are, but as they would be, if GALEN'S Theory of the Four Humors was well grounded; their Distinctions being deduced from different Mixtures of Bile and *Pituita*, which they assign as the Cause of Tertians; and having once laid it down as an Axiom, that each Species arises from a particular Humor, which must produce such and such Effects, they ascribe Symptoms to the Fever, from a preconceived Hypothesis, which seldom or never accompany it in reality\*.


CHAP.

III.



\* Vid. Simon Simon. apud Seannert. tom. ii. lib. ii. c. xvii.



CHAP.  
III.

Wherefore, in order to avoid the like Mistakes, I shall leave the Reader to form what Idea of the Causes of these Fevers he thinks fit, and endeavour to shew as distinctly as I can, First, the Differences which arise amongst them from the different Types or Forms of their Periods. Secondly, the Distinctions occasioned by the most obvious Symptoms of their Paroxysms; and, Thirdly, I shall describe the various Appearance of the more frequent epidemical Tertians in their Rise, Progress, and Termination, with as much Exactness, as their own irregular Anomalous Disposition will permit.

But it may not be improper, in the first Place, to apprize the Reader, that I have used the Appellations commonly given to the various Fevers of this Class, in a Manner somewhat different from several who have wrote upon this Subject, tho' not without explaining as clearly as I could in what Sense I would be understood:




stood: And indeed whoever consults these Writers, will find that some of them so limited their Definitions as to make it doubtful if ever any Disease existed to which the Name was applicable; whilst others apply their Terms in so loose and vague a Signification, as to include several Species, which ought to be distinguished, in order to understand their History and Cure\*. It may likewise be necessary to inform those, who are not much conversant with the Hippocratical Writings, that by the *odd Days* are meant the 1, 3, 5, 7, &c. by the even Days the 2, 4, 6, 8, &c. and that however singular such a Distinction may appear in a

\* For Instance, if we look into several of the Galenical Writers, we shall find, that in order to constitute the *Tertiana pura* or *exquisita*, and some Kinds of the *Semitertiana*, a Number of different Circumstances are required, which seldom or never occur in the same Patient. On the other hand, SPIGELIUS, who wrote a particular Treatise *De Semitertiana*, has extended the Meaning of that Denomination so far, that it includes almost every Species of the Tertian Tribe.



CHAP.  
III.



Country where no such uniform Diversity is observable, yet it will be found next to impossible to give a tolerable Idea of the Diseases in that Climate without the Assistance of this Distinction.

In this Treatise, Tertian Fevers are called *simple*, when there is but one Fit and one Interval during each Period of the Distemper. When the Paroxysms do not exceed twelve hours, they are named *true Tertians*; when they continue longer, they are called *Spurious*. In the *true simple* Tertian, the Fit, for the most Part, comes on in the middle of the Day, and goes off the same Evening. In the *spurious simple* Tertian it comes on much earlier, and often lasts above eighteen Hours.

They are called *double* Tertians, when there are two Fits and two Intervals within the Time of each Period. But commonly there is some Difference between the two Fits, either in respect to  
the




the Hour they come at, the Time of their Duration, or the Nature and Violence of their concomitant Symptoms; whilst at the same Time the third Paroxysm of the Disease resembles the First, the Fourth agrees with the Second, the Fifth with the Third, and so on.

CHAP.  
III.

Some double Tertians begin in this Manner: On the Evening of *Monday*, for Example, a slight Fit comes on, and goes off early next Morning: But on *Tuesday* towards the Middle of the Day, a more severe Paroxysm begins and continues till Night. Then there is an Interval to *Wednesday* Evening; when a slight Fit commences a new Period of the Fever, which proceeds in the same Manner as the first; so that (according to the Way Physicians calculate the Days of Diseases by beginning to reckon from the first Hour of their Invasion) both Paroxysms happen on the odd Days, while the greatest Part of the even Days is calm and undisturbed.

But



CHAP.  
III.

But in most double Tertians, the Patient has a Fit every Day of the Disease; the severe one commonly appearing at Noon on the odd Days; the slight one towards Evening on the even Days; tho' sometimes the worst of the two Fits happen on the even Days.

Double Tertians, if their Fits do not exceed twelve Hours, are named *True*; if they are somewhat more protracted, they are called *Spurious*; and if the Fits are so far protracted, that one is scarce ended when another begins, they are called *Subintrant*.

There is a Tertian Fever sometimes to be met with, during each Period of which, there are three different Fits, and as many Intervals. For Example, towards *Monday* Noon the Patient is seized with a Paroxysm, which declines about five or six o'Clock the same Evening: A few Hours after, another Fit begins and  
continues



continues until Morning; from which Time there is an Interval to *Tuesday* Evening, when a third Fit comes on, and lasts most Part of the Night. On *Wednesday* there are again two Paroxyfms, as on *Monday*, and one on *Thursday* like that of *Tuesday*; and thus the Fever goes on, with a double Fit on each of the odd Days, and a fingle Fit on the even Days.

CHAP.  
III.  
~

This uncommon Kind of Tertian is the true Semitertian of HOFFMAN \* and some others: But I fhall call it a *Triple Tertian*, in order to diftinguifh it from another very common Fever, which proceeds in the following Manner.

A Fit begins on *Monday* Noon, for Example, and goes off the fame Night. On *Tuesday* Afternoon a fecond Fit comes on, and gradually increafes till *Wednesday* Night, when it terminates. On *Thursday*

\* Med. Rat. tom. iv. § i. c. v.

Morning



CHAP. Morning there is such another Interval as  
 III. happened on *Tuesday* Morning: But on  
*Thursday* Afternoon another long Fit,  
 like the preceding, commences, and re-  
 turning regularly every other Day, leaves  
 only a short Interval of ten or twelve  
 Hours during the Eight and Forty. This  
 Fever I shall call, with CELSUS\* and  
 AGATHINUS †, a *Semitertian*,

But the various Kinds of Tertians  
 hitherto mentioned, differ among them-  
 selves, according as their periodical Inter-  
 vals are more or less tranquil and free  
 from Disturbance. When there is a com-  
 plet Apyrexia or entire Vacation of Fever  
 between the Fits, they are called Inter-  
 mittents; when the Apyrexies are more  
 imperfect and obscure, Remittents; and  
 continual Tertians, when the Paroxysms  
 and their Intervals are less perceptible,  
 though the Violence of the Symptoms is

\* CEL. lib. ii. c. iii.

† Apud Galen de Febr. Different. lib. ii. c. ii.


some-



somewhat abated on the alternate Days. In double Tertians that Interval is the most considerable which follows the severe Fit; for the slight Fit oftener ends in a Remission, than Intermission, and frequently lingers till the other approaches: Hence it is that the Night preceding the vehement Fit is much more restless, than that which comes after it, as has been observed by HIPPOCRATES\*.

CHAP.

II.



These are the Differences of Tertian Fevers, which proceed from the Types of their Periods. But in some, each Period is compleated in a few Hours less than Eight and Forty; in others, in a few Hours more: The former are therefore called anticipating, the latter postponing Tertians. In double Tertians the vehement Fit often comes on a little earlier in each Period, while the slight Fit returns at the same Hour, or perhaps later and later every other Day: So that the

\* Morb. Vulg. l. vi. § 2.



CHAP. Motions of one have no Influence on those  
 III. of the other; from whence it appears,  
 that each of these Fits hath its own proper independent Cause.

With Regard to the Types of the Periods, and the Time of their Duration, this may be sufficient. I come now to the Differences of Tertian Fevers formed by the Symptoms attending their Paroxysms.

Many Authors, (and HOFFMAN in particular\*) have carefully enumerated the *Phænomena* of the Fits in regular distinct Tertians, according to the Order in which they succeed: But so far are the Paroxysms of the epidemical Fevers, which are the Subject of our Discourse, from having constantly one and the same Appearance, that no two Distempers can differ more than they often seem to do from one another. In both simple and double Ter-

\* Med. Rat. tom. iv. § i. c. ii.

tians,



tians, as well as Semitertians, they often begin with a slight Shivering; sometimes without any Sense of Cold; and frequently while the Patient complains of Cold his Skin is actually warmer than in a natural State. In some a most intense Degree of Heat is brought on; in others it is moderate. In a few they terminate by Stool or Urine rather than by Sweat: Sometimes they are so complicated with fixed Pains of the Head, Breast, Belly, Back, or Limbs, as to personate with great Exactness, a Phrensy, Pleurisy, *Hepatitis* *Lumbago*, or Rheumatism; especially if the Apyrexies are obscure or imperfect. Sometimes one or two Symptoms of the Fit predominate with such violence, that the rest are obscured or altogether eclipsed. Hence we so frequently meet with *Hemicrania's*, *Choleras*, *Dysenteries*, and *Chin-coughs*, returning regularly at stated Periods: And several Fevers of this Class, upon Account of some predominant Symptom, have had particular Names bestowed upon them: For Example, when the  
Cold,

CHAP.

III.





CHAP.  
III.

Cold, which ushers in the Fits, continues longer than ordinary, and is so very intense that the Surface of the Body is chilled, whilst a burning Heat is felt in the Bowels, the Fever is called *Lipyria*. When the Anxiety and Dejection of Mind, which commonly happens in the first Stage of the Paroxysm, is so far augmented, as to degenerate into an absolute Fainting, the Disease is termed *Febris Syncopalis*. When exquisite Pain and burning Heat in the Bowels, renders the Patient restless, and incapable of continuing many Moments in the same Posture, *Affodes* is the Name appropriated to the Fever. And it is called *Elodes*, when the Skin is constantly covered with Sweat; either from the *Viscera* being inflamed, or from a general Dissolution of the Blood \*. For although in the Course of these malignant Fevers it often happens that you

\* Atque hoc in totum de sudoribus animadvertere oportet, quod nonnulli quidem ex corporis dissolutione, quidem ex inflammationis vehementia contingant.

HIPPOCRAT. *Prænot.* lib. 5



can neither distinguish Paroxysms nor Intervals, yet it is evident that they belong to the Tertian Tribe, since, for the most Part, in their Beginning the Periods are sufficiently distinct; and as soon as that Violence of the Symptoms which occasioned the Confusion in the Height of the Distemper abates, they again become more regular, and assume either their former or analogous Types\*.

Whoever rightly comprehends what I have already said on Tertian Fevers, will easily see that it would be endless to reckon up the different Species into which they might be divided, and to ascribe particular Names to each. Yet there is another Circumstance which renders this subject still more perplexing. For such is the variable Disposition of

\* *Febris Syncopalis minuta subtilis, est febris acuta faciens cadere pulsum & virtutem in paroxysmo uno aut duobus paroxysmis, cum additione dissolutionis accidente in corpore cum velocitate.—Plures paroxysmi hujus febris, sunt paroxysmi tertianæ.*

AVICEN. *de Feb. c. liii.*

M

these




CHAP. these Diseases, that they often change  
III. from one Appearance to another, and  
feldom retain the same Form from their  
Beginning to their Termination; each  
Period sometimes assuming a new Type;  
and every Paroxysm being attended with  
different Symptoms. This led me at first  
to apprehend that they were confused,  
anomalous, and altogether without Order:  
But after being for some Time acquainted  
with them, I began to discover their Re-  
gularity; and the longer I was conversant  
among the Sick, the more I was sur-  
prised at the Constancy of Nature in the  
Production and Progress of Tertian Fe-  
vers; their Periods being perfectly similar  
in the *Spaniards* and in the *English*; and  
sometimes not very different in him who  
lies upon the bare Ground, destitute of  
Assistance, and those who are treated in  
the most judicious Methods, under every  
Advantage of Fortune: and frequently  
neither the Patients Intemperance, nor  
unskilful Management, can alter their  
stated Course, and prevent their termi-  
nating



nating in Recovery. So much are those mistaken, who imagine that the Bent of Nature in acute Diseases can be altered or controuled by every trifling Accident or insignificant prescription.

CHAP.


III.



The most Part of these Fevers make their first Appearance in the Shape of a true simple, or double intermittent Tertian. The cold Fit seldom lasts above an Hour or two; and as it goes off, some bilious Matter is commonly discharged by Vomit or Stool. Then follows an intense Heat over the whole Body, which raises the Mercury in the Thermometer to the 103d or 104th Degree: And lastly, a profuse Sweat puts an End to the Paroxysm. The Apyrexia is tolerably compleat, though for the most Part the Patient complains of a disagreeable Taste, Loss of Appetite, Head-ach, and Pain in the small of his Back, and Pit of his Stomach upon a full Inspiration. The Pulse during the Intermision is almost natural: in the Fits it varies according to their



CHAP.  
III.



predominant Symptoms: When they are attended by acute Pains in the *Præcordia*, it becomes small and obscure, so as to indicate greater Weakness than there really is: on the other Hand, when drowsy lethargick Symptoms come on, it often resembles that of a Person in full Health, Though the Sick is in the utmost Danger. The Urine, whether made in the Time of the Paroxysm or Interval, is always clear, frothy, and of a deep red Colour, without any Separation. Blood drawn from a Vein is most commonly florid like scarlet, without any sizy Crust: the *Serum* is sometimes tinged with yellow, but oftner red like the *Lotura Carnium*, and in great Quantity. Sometimes the *Serum* and *Crassamentum* remain united in a lax gelatinous Mass.


As the Fever advances to its Height, the Coldness and Shivering which usher in the Paroxysms become less, or intirely imperceptible: in which case a *Cholera Morbus*, or acute Pains in the Back or  
Limbs,



Limbs, often supply their Place: frequently the Shiverings are intermixed with Flushings of Heat. In the mean Time, the Paroxysms themselves become longer, and bring on more formidable Symptoms; such as Head-achs, Raving, *Sopors*, Apoplectick Fits, Bleeding at the Nose, Cough, Difficulty of Breathing, Palpitation of the Heart, Irregularity of the Pulse, Sickness and Anxiety, Pain about the upper Orifice of the Stomach, Vomiting and Purgings; Heat, Tension, Pain and Pulsation in the abdominal *Viscera*: *Subsultus Tendinum*, and an infinite Variety of other Complaints, which do not entirely cease with the Sweat, that carries off the Paroxysm; so that the Apyrexia is not only shortened, but rendered more obscure.

Besides, it often happens, during the second, third, fourth, or fifth Period, that the Tertian becomes double, though at first it was simple: Or if it was double from the Beginning, the weaker Fit continues,



CHAP.  
III.

tinues, without any Intermission, till the stronger comes on, and both being blended together, the Disease puts on the Appearance of a Semitertian, having one very long Fit, with a short Interval every Forty-eight Hours. Sometimes a double Tertian degenerates into a triple Tertian, two Fits instead of one happening on the odd Days.

It must likewise be observed, that, in the Progress of the Fever, the regular Order of the Periods is frequently disturbed, by the Paroxysms changing their Hour of Invasion, and attacking unawares without any previous Cold. Nor are anticipating Fits always a bad Sign, or those which postpone a good one, as some Authors insinuate; on the contrary, the first frequently shew the Strength of Nature, as the others do her Weakness.

After this Manner these proteiform Distempers continue to vary their Shape in every Period, and to produce longer,  
more



more severe, or more frequent Paroxysms, till they arrive at their Height; about which Time the Fits and Intervals are often so confused, that they are scarcely to be distinguished: Nevertheless, if Death be not speedily the Consequence of this Confusion, they commonly again put on a more simple or regular Form, and, after one or more slight Paroxysms, go away of their own Accord.

CHAP.  
III.  
~

Those Fevers which come to their Height in the third Period, terminate in the fourth or fifth Period. Those which come to their Height in the fourth Period, terminate in the fifth or sixth. And those which come to their Height in the fifth, terminate in the sixth or seventh. When the most vehement Paroxysms happen on the odd Days, the *Crisis* will be on the odd Days: When they happen on the even Days, the great Changes of the Distemper will likewise be on the even Days.

If the Fever increases to the seventh  
M 4 Period,



CHAP.  
III.

Period, it probably will not cease before the Ninth: But it rarely happens that intermitting or remitting Tertians run out to so great a Length, Yet I have seen every Year a few of the continual Kind, which began with great Mildness, and increasing by slow Degrees, broke out violently in the third or fourth Week; and soon after ended in Intermitments; though some of them have continued without any considerable Interval for six or seven Weeks. HIPPOCRATES \* has accurately

\* The other Fevers were altogether of the continual Kind without any Intermissions; and the Paroxysms in all like the Semitertians, one Day better, another worse. And of all the Fevers that then reigned, these were the most vehement, tedious, and painful, beginning very mildly, but increasing always, growing worse and worse on the critical Days. After a little Abatement they soon grew bad again, had stronger Fits on the critical Days, and for the most Part worse: Shiverings were universally irregular and uncertain, seldom and very little in these, but more in other Fevers: Sweats were common, but here least of all; and so far from easing the Patient,



accurately described such continual Tertians, and tells us they are apt to terminate in Dysenteries, Lienteries, and Tenesmus, which we find in fact to be too often the Case.

CHAP.  
III.  
~


But it is much more common to meet with Tertians, which set out furiously, with severe subintrant double Paroxysms; so that for some Days, they have little or no Interval. On the third or fifth Day a profuse Sweat commonly brings on an Intermiffion; and afterwards the Disease assumes the Type of a double intermitting Tertian, or of a Semitertian. Such Fevers I have frequently observed to terminate spontaneously on the seventh, ninth, and eleventh Days, and for the most Part, they are less to be feared than those which begin deceitfully in the Shape of a slight double or simple Tertian.

Patient, that, on the contrary, they did him harm.—  
The Belly in all was disturbed, and in a bad Manner,  
but worst of all by much in these, &c. &c.

CLIFTON'S HIPPOCRAT. p. 57, 58.

For



CHAP.  
III.

For however mild and insignificant these Intermittents may at first seem to be, we are never to trust Appearances, till they have performed three or four Revolutions. Then indeed, if the Paroxysms are not attended with acute Pains in the *Viscera*, and do not last above twelve Hours; if they decline with plentiful warm Sweats, and leave the Intervals tolerably free; if the Patient bears the Distemper well, and begins to have an Appetite for Victuals; if small Pustles break out in the Inside of the Mouth, or Scabs about the Lips\*; if the Urine has recovered its natural Complexion, or is cloudy and turbid, or lets fall a white or a pale red Sediment; I say, if all these Signs concur, about the third or fourth


\* It is to be observed, that these are only to be looked upon as a good Sign, when they come in the Declension of the Disease, attended with other Signs of Concoction; for if they break out in the Beginning, they prognosticate a dangerous or a tedious Illness.

Period,



Period, we may safely prognosticate a speedy Recovery.

CHAP.  
III.



On the other Hand, it denounces Danger, when, about this Time of the Disease, the Paroxysms are long and protracted; or are accompanied with an obstinate *Delirium*, an intense *Coma*, great Anxiety, and Pain in the Loins, or about the upper Orifice of the Stomach; when the Patient has an utter Aversion to Food, and even in the Intervals is so feeble, and attended with such a swimming in the Head, that he can scarcely walk about; when the *Hypocondria* and Epigastrick Region, are swelled, hard, and painful to the Touch; when numerous Blotches, like the stinging of Nettles, frequently break out on the Skin; when the Urine continues thin, clear, high-coloured, or covered with an Ash-coloured Membrane like a Cobweb; and lastly, it denounces Danger, when larger Evacuations come on than the Strength can well bear, such as Vomiting, Purgings,  
Bleed-



CHAP.  
III.

Bleeding of the Nose, Colliquative Sweats, or the like. For Fevers with these Appearances, sometimes are immediately changed into mortal Dysenteries; sometimes they become continual Tertians, and run out to a great Length; but, for the most Part, they preserve the Form of remitting or intermitting \* Fevers, and daily

\* In this and some other Parts of this Treatise, it is inculcated that the Danger in Tertian Fevers, is rather to be estimated from the Symptoms of the Paroxysms, than the Length and Serenity of the Intermiſſions: On which Subject, as FRANCISCUS TORTI has expressed himself with great Clearness, I shall here add a few Remarks, from his *Therapeutice Specialis*, a valuable Work, which I never had the good Fortune to meet with till very lately. He tells us, that Periodical Fevers, and principally double or simple Tertians, become malignant, either when they degenerate into continual acute Fevers, or when they still retain their Intermiſſions, but are accompanied with one or other of the following Symptoms; which commonly proves fatal in the second or third Fit, after its appearing in the formidable Manner he describes: 1. A Vomiting or Looseness, like a *Cholera Morbus* or Dysentery. 2. A Looseness which often resembles the *Fluxus Hepaticus*, and sometimes a purging of *Atrabilis*.




daily growing stronger, prove very dangerous about the sixth or seventh Period; and though the Patient may escape, after a great

CHAP.

III.

3. A *Cardialgia*. 4. Cold Sweats. 5. A *Syncope*. 6. A Constant Coldness, neither succeeded by Heat, nor Sweat. 7. A lethargic Disposition, little different from an Apoplexy. Hence he makes so many different Species of malignant Intermitents, and names them from the predominant Symptoms of the Fit. 1. *Febris Cholericæ seu Dysentericæ*. 2. *Subcruenta seu Atabilaris*. 3. *Cardiaca*. 4. *Diaphoretica*. 5. *Syncopealis*. 6. *Algida*. 7. *Lethargica*. He describes, with great Accuracy, the Manner in which each Species occasions Death; and takes Notice that in the six Species first mentioned, the Pulse is constantly small, feeble, and greatly depressed; whereas in the Seventh it is rather full, strong, and slow (as in an Apoplexy), than weak and quick. This, which he calls the *Lethargica*, is extremely common in *Minorca*; the *Cardiaca* and *Cholericæ* are likewise frequent; and all the rest are now and then to be met with, excepting perhaps the *Subcruenta*, which I never took Notice of. Our author likewise remarks, that it is not unusual for some of the abovementioned Symptoms to attend continual periodical Fevers, though seldom in so great a Degree of Violence as in the Intermitents. And he observes, that intermittent Quartans rarely kill in the Fit, as the Tertians do, but often prove fatal by becoming



CHAP.  
III.

a great Struggle, by means of some critical Discharge, such as Purging, Sweating, Parotids, or Abscesses near the Hip; yet his Constitution is commonly so shattered, that he long continues exposed to irregular Paroxysms, Night Sweats, Fluxes, Obstructions of the *Chylopoetick Viscera*, and every kind of chronick Disorder.

Those Fevers are most to be dreaded, whose Violence is greatest on the even Days; and if the Paroxysm stops on the third, fifth, or seventh Day, but continues on the fourth, sixth, or eighth Day, we must be upon our Guard, lest a sudden Storm should succeed this treacherous Intermission\*.

But


becoming continual Fevers. The whole Work deserves to be carefully perused, but particularly the first Chapter of the third Book, from which the above Remarks are collected, with a view to raise, rather than gratify, the Reader's Curiosity.

\* I inserted this caution, having seen a few double Tertians change into simple Tertians, by the Fits intermitting



But the utmost Danger is to be apprehended, if a few Drops of Blood fall from the Nose: if black Matter like the Grounds of Coffee, is discharged upwards or downwards: if the Urine is of a dark Hue and a strong offensive smell: if the

CHAP.  
III.




terminating on the odd Day, and nevertheless prove fatal soon afterwards; which is a case I had not found so much as hinted at in any Author except HIPPOCRATES, who tells us in the *Prænot. Coac. Quibus tertio die subsistit accessio et quarto ingravescit malum.* Yet from the following passage in Torti, I imagine that experienced Physician must have met with Accidents of the same Kind. *Suspecta itaque ab exordio erit, ne in continuam degeneret, intermittens, quæ cum paucis aut nullo rigore solet invadere, sed potius cum sensu caloris. Item quæ primo die leviusculam (dum modo tamen vere febrilem) infert accessionem; altero vero die (non altera periodo) fortiozem, et sic progrediendo modum servat graviozem per dies pares, loquendo per modum exempli, de tertiana duplici ab ortu. Quod si eadem sic orta in simplicem statim mutetur, etsi hoc laudabile sit, tamen non desinit esse suspectum, si primam accessionem validam, debilis, ut supra, immediate præcessit: potest enim facile ille typus mutari qui ordine inverso potuit incipere.*

Therap. Special. l. iii. c. .

whole



CHAP.  
III.

whole Skin is tinged with a deep yellow, or any where discoloured with livid Spots or Suffusions \*: if a cadaverous Smell is perceptible about the Patient's Bed : if in the Time of the Fit he continues cold and chilly, without being able to recover Heat ; or if he becomes extremely hot, speechless and stupid ; has frequent Sighs, Groans, or Hickups ; and lies constantly on his Back, with a ghastly Countenance, his Eyes half shut, his Mouth open, his Belly swelled to an enormous Size, with an obstinate Costiveness, or an involuntary Discharge of the Excrements: which formidable Symptoms, as they seldom appear before the third Revolution of the Disease, so they frequently come on, both in double and simple Intermittents, during the fourth, fifth, or sixth Period, even where the smallest Danger was not foreseen : But at whatever Time the greatest Part of them concur, they afford

\* The *English* in *Minorca* are more liable than the Natives to become yellow in these Fevers.

a melan-



a melancholy Prognostick: For notwithstanding they sometimes go entirely off with the Paroxysm, and the Patient seems to be left in a fair Way of Recovery, yet most commonly they return in the next Period with double Violence, and terminate in sudden Death. Agreeable to this HIPPOCRATES tells us, "In the Summer  
 " Intermittent Fevers and the *Cholera*  
 " *Morbis* prevail; and as they sometimes  
 " degenerate into malignant acute Diseases, we should be upon our Guard;  
 " the fifth, seventh, and ninth Days  
 " point out the Danger; but we must be  
 " cautious to the fourteenth\*." For the fifth Day, if the bad Symptoms just now mentioned, appear on it, will indicate Death on the seventh. In the same Manner the seventh will indicate the ninth, and the ninth the eleventh to be fatal, provided the Paroxysms are aggravated on the odd Days; for if they are most vehement on the even Days, one of

\* De Morb. Popular. l. vii.



CHAP. these Days will prove indicatory as well  
 III. as critical. Hence we find GALEN\*  
 lays down as a general Rule, that those  
 who grow remarkably worse on the fourth  
 Day, die on the sixth; and with equal  
 Justice he might have said the same Thing  
 of the sixth Day, and with regard to the  
 eighth, and of the eighth, with respect to  
 the tenth.

Nor is there only a Possibility, in many  
 Cases, of foretelling the Day, but like-  
 wise the Hour on which the Patient will  
 expire; for that Stage of the Paroxysm,  
 which he usually got over with most  
 Difficulty, will most probably in the End  
 prove fatal, as GALEN † has also re-  
 marked. I have seen some expire in  
 what may be called the first Stage of the  
 Paroxysm; the Skin being chilled and wet

\* Etenim qui in quarto ad pejorem statum recidunt,  
 plerumque sexto moriuntur.

*De Dieb. decret. l. i. cap. iv.*

† *De Crisib. l. iii. c. x.*







CHAP.  
III.

hundred Persons, who perished in these Fevers, and constantly found one or other of the adipose Parts in the lower Belly, (the Cawl, Mesentery, Colon, &c.) of a dark black Complexion, or totally corrupted; the *Vesica Fellea* full and turgid, and the Stomach and Intestines overflowing with bilious Matter; the Spleen large, sometimes weighing four or five Pounds, and so excessively soft and rotten, that it had more the Appearance of congealed Blood wrapt up in a Membrane, than of an organical Part. In the Cavity of the Head and Breast, nothing extraordinary was met with, excepting yellow *Serum*, when the Skin was tinged with the same Colour.

Some Observations relating to malignant Tertians are handed down to us by COELIUS AURELIANUS, which, for the Reader's Information, I shall transcribe at the Bottom of the Page\*; and whoever

\* Hæc passio (nempe apprehensio sive oppressio) lethargiæ similis est, HIPPOCRATES et DIOCLES αφωριασιν appellavit,



ever is acquainted with this Author's CHAP.  
 Style, will easily discover how nearly the } III.  
 preceding Remarks agree with those of  
 the

appellavit, PRAXAGORAS *Κυματωδον* ANTIGENES *ανωδιαν*, ASCLEPIADES Catalepsin—DIOCLES ait (a) (a) Inter-  
*defectivas febres* tutas et innoxias esse frequentius quam mittentes,  
 sunt continuæ, quamquam et in his periclitentur, qui  
 in accessionibus apprehensi conticescant, vel raptu  
 quodam alterno per membra tentantur, cum supra  
 dictis: quod sæpe, inquit, est accidens pueris. Item  
 PRAXAGORAS ait esse quasdam febres ex anno duo-  
 decimo usque ad annum XVI aut XVII, quæ, quadem  
 privata pernicie, mortis habent effectum. Atque id  
 in servis magis quam liberis evenire, sed (b) *excesso die-* (b) Impari-  
*rum numero* passionis fiunt, catochæ vehementes, ut bus diebus.  
 etiam voce capiantur ægrotantes: Horum aliquos  
 etiam lethargicos fieri. Denique, inquit, quidem  
 liberati, et deinde (c) *sanitatis creduli*, plurimum quic- (c) Sani sibi  
 quam sumentes, repente in mortem venerunt. — vivi.  
 Item ARCHIGENES ait difficiles esse periodicos typos  
 horum difficiliiores esse Tertianas, in quibus quo-  
 tidianæ accessiones fiunt, sed alternâ diei interposi-  
 tione, (d) *suæ similitudini respondeant*, quando in accef- (d) Sibi si-  
 sione vehemens occurrerit oppressio, et appellavit PITOS. miles sint,  
 Difficilis, inquit etiamsi quotidianis diebus accessiones  
 sibi similes fiunt, et in accessione ægrotantes supra dicta  
 patiantur. Dehinc progrediens paululum idem dixit  
 periodicos Typos non esse perniciosos, sed horum esse  
 N 3 molestos,



CHAP. the old *Greek* and *Roman* Physicians. In  
 III. some Circumstances however there seems  
 ~~~~~ to be a Difference; for according to  
 them,

(e) Profundus somnus.

(f) Somno turbulento.

(g) Tertianas duplices.

(b) Non vera sed in specie tantum.

molestos, quibus accessionis tempore (e) *Pressure vehementes* eveniunt, et quodam (f) *Nubilo* corpus demergitur, quod item Rhigos vocavit; sed hoc, inquit, est accidens magis tertianis, aliquando etiam quotidianis, quæ similibus respondeat accessionibus. — Apud Roman vero inquit ASCLEPIADES frequentare advertimus (g) *has febres*, cum corporis atque mentis oppressione, in similitudine lethargiæ, quæ secundo vel tertio in statu accessionis constitutæ, statim recalescitur corpore, vel cessante vehementia, in resumptionem et resurrectionem mediocrem revocant ægrotantes. At si (b) *levi figmento* cessaverint, semel apprehensio ægro nullam resurrectionem dabunt, sed in sudores, et respirationem celerum, in pulsum febricitantem desinunt et occidunt. — Autumni tempore hæc passio magis irruit corporibus atque puerilibus frequenter ætatibus; item mulieribus humorosis, et vacuis corporibus, et edacibus hominibus. — Præterea omni febriculæ hæc passio irruere potest sive continuis sive demissionibus intercapedinatis, hæmitritaicis etiam febribus, vel quotidianis, vel tertianis, vel quartanis, frequentius tamen quotidianis accessionibus, vel tertianis, ægrotantes ista passione afficiuntur, et propterea, diuturnis accessionibus admoniti, tertianis similitudinem servant, ad typum quotidianum: sed omnium earum febrium



them, Children are most liable to malignant Tertians; whereas by what I have observed, they were more frequent among Adults and those of an advanced Age. They likewise talk, as if the Fits were more to be dreaded when they come on with a Numbness of the Joints and

CHAP.  
III.

februm gravius, quoties cum articularum frigido torpore fuerit (i) *Qualitas*: levius, quoties tremore ægro-  
tantes afficiuntur: Item magis et magis levius, quoties sine his quæ supra diximus, solo fervore febres initium accipiunt. *Afterwards, having accurately enumerated the Signs of the Distemper, our Author proceeds to describe it in the last Stage.* Cum sudore sæpius plurimo atque ferventi et in demissione sinceritati propinquantes rursus admonentur. At si ad pejora passio fuerit devoluta, fervor plurimus corporis in superficie, magis sentitur respiratio, oculorum conversio, menti quoque fixa conductio, manuum contractus, et musculorum, qui buccas colligunt, tanquam ridentium, sudor igneus, et quibusdam in vultu et thorace emergentes discolors vel stantes in rotunditate (k) *Maculæ*, in similitudinem (l) *Scatebrarum corporis*, quas Græci *λοιδες* vocant, et e magnitudine repentinus virium casus, gutturis stridor, quem ronchum vocant, torpor frigidus, albidus vultus, et in multimo (m) *Effatio*, atque vitæ periculum.

(i) Mos febris.

(k) Exanthemata.  
(l) Sodoris guttularum.

(m) Suffocatio.

Lib. ii. cap. x. *De morb. acut.*




CHAP.  
III.

Tremblings; but the most formidable Paroxysms, which I have seen, broke out into a burning Heat at the Beginning, without any previous Cold.

From what has been said, I flatter myself that young Practitioners will be enabled, not only to distinguish Tertians from all other Fevers, but likewise to foresee the Times of their Paroxysms, and Intervals, and to make a rational Conjecture, concerning their Event. Before I proceed to the Cure, I shall only remark, that although we find them appearing every Year in the several Forms and Types described in the Beginning of this Chapter, yet the true simple and double Tertian, and the Semitertian, are by much the most frequent. In *July*, when they first break out, their Type is commonly simple and regular; their Paroxysms are of short Duration; and after three, four, or five Periods, they vanish of their own Accord. So just is HIPPOCRATES'S Observation, that true Tertians





tians generally end within five, seven, or at most, nine Revolutions \*, notwithstanding what some of the Moderns have alledged to the contrary. As the Season advances, the Tertians become more dangerous and difficult, often terminating in those malignant Fevers, called *Syncopalis*, *Lipyrria*, *Assodes*, &c. especially if much Rain, without Wind, fall during the Dog - Days. About the Time of the Equinox, they assume a surprizing Variety of Forms, and very often counterfeit continual Fevers, having long redoubled Paroxysms. But as the Winter draws near, their Type becomes more simple, and though they prove tedious and obstinate in cold Weather, yet they are more regular, and less dangerous than in the Summer.

In the Cure of Tertians, the Symptoms of the Paroxysms are more to be regarded than the Types of the Pe-

\* Prænot. Coac.

riods;



CHAP. riods; for *cæteris paribus*, simple, double,  
 III. triple, intermittent, and remittent Ter-  
 tians, together with the Semitertians, all  
 require the same Sort of Treatment.

As soon as the Fit comes on, the Patient should be put to Bed, and covered with Cloaths in such a Manner, that he may neither be suffocated with immoderate Heat, nor the critical Sweats be suppressed by the Admission of too much Cold. If it can conveniently be done, his Bed should be placed in a large Room, where he may breathe freely, in pure temperate Air. In the Summer it will be necessary to keep out the Sun, and to refresh the Air frequently by sprinkling the Floor with Vinegar and Water, and opening the Windows to the North; during Winter, or the autumnal Rains, the Cold and Moisture of the Weather must be corrected by a Fire.

While the cold Fit continues, the Patient should abstain from drinking. For  
 at



at that Time the *Vena Cava* and sub-clavian Vein are so full and turgid, by the Blood's being driven from the Surface of the Body to the internal Parts\*, that the Discharge of the thoracick Duct is prevented: Hence the Liquor swallowed down, remains in the alimentary Tube, loads the Bowels, and creates Anxiety. And therefore if the Patient is thirsty, he must endeavour to allay it in the best manner he can, by washing his Mouth often, and eating Slices of Lemons sprinkled with Sugar. In the mean Time, if a *Nausea* or Inclination to vomit supervene, these salutary Attempts of Nature should be promoted by large Draughts of warm Water or weak Broth, which commonly bring up a Quantity of bilious Matter, to the immediate Relief of the Sick.

CHAP.  
III.


\* Sanguis presentem horrorem metuens, ad partes maxime calidas concurrat.

HIP. lib. de Flat.

When



CHAP.  
III.



When the first Stage of the Paroxysm is over, or (to use HIPPOCRATES'S Phrase) *When the Heat descends to the Feet* \*, Liquids may then be given; but with such Moderation, that the Patient be neither allowed to take as much as he pleases, nor suffered to be tormented with Thirst. As soon as the Sweat breaks out, he may then be permitted to drink at Pleasure.

The *Spanish* Physicians deny them Liquors altogether till the Fit be at its Height: They then give them in small Quantity; always encouraging the Sick, with CELSUS'S Remark †, “*That the Thirst will cease with the Fit; which will continue longer if they drink.*” On

\* Per totum morbum istud præcipue observari debet; ut cum frigidi pedes fuerint, tum a sorbitione exhibenda, tum maxime a potu abstinemus. Cum vero calor ad pedes descenderit, tunc dare convenit.

HIP. de Vict. Rat. in Morb. acut.


† Lib. ii. cap. vi.



the other hand, our People run into the opposite Extreme, by permitting the free use of Liquids in every Stage of the Paroxysm. The first of these Errors, especially in Tertians which have long Paroxysms, besides torturing the Patient, augments the feverish Heat, occasions a Putrefaction of the animal Juices, destroys the Solids, and throws Life itself into the utmost Danger. By the contrary Error, the Stomach is unloaded, Nature is diverted from expelling the critical Sweat, and consequently the Paroxysm is prolonged.

The proper Drinks on this Occasion, are those which allay Heat, prevent Putrefaction, dissolve the acrimonious Particles of the Blood, and convey them through the Kidneys and Pores of the Skin, such as boiled Water, sweetened with Sugar, and rendered agreeably acid with Lemon Juice, Spirit of Vitriol, or sliced Apples infused in it; Barley-Water mixed with simple Oxymell; Almond-Milk,

CHAP.  
III.





CHAP. Milk, and other Emulfions; to all which  
 III. Nitre must be occasionally added.


It is a noted Question among Physicians, whether during the Fit the Drink should be warm or cold? The *Spaniards* generally give crude Water, cold from the Cistern; and we find by Experience that this, if it be not hastily swallowed down in great Quantities, is not only safe and innocent in Summer Fevers, but much preferable to warmer Liquors, as it quenches Thirst more effectually, strengthens at the same Time the Tone of the Vessels relaxed and enervated by Heat, and preventing the Tendency of the Blood to a putredinous Thinness. Hence after each Draught the Body seems to acquire fresh Vigour, whereby it is enabled to perform the Concoction of the febrile Matter, and discharge it by the proper Emunctories\*. And therefore,

In hac curatione observandum est, a spirit. sulph.  
 gutt. xx. cum libra aquæ frigidæ exhibitis sudores copiosos



fore, they are greatly to be blamed, who refuse their Patients so powerful and agreeable a Remedy, in spite of the earnest Call of Nature, contrary to the Advice of the best Practitioners\*. Nevertheless, as there is a manifest Hazard of the Blood's being coagulated by the sudden Application of intense Cold, we must beware of giving Ice Water, as the *Italians* and *Sicilians* do, unless the Patient has been accustomed to it when in Health. And if the Bowels are inflamed, the safest Way is to give the Drink luke-warm, or a very little colder.

CHAP.  
III.



During the Time of the critical Sweats, the Patient's Shirt and Sheets should fre-

quenter piofos fuisse provocatos, quod multis experimentis nobis innotescit, qui plures tertianas febres curatas vidimus, eodem remedio, in summo accessionis æstu, et urgente siti exhibito, unde copiosi sudores provocabantur, a quibus non solum Paroxysmus, sed etiam totus morbus integre solvebatur.

River. Obs. xix. Cent. i.

\* Vid. HOFFM. t. iv. § i. c. i. & t. iii. § ii. c. xi.

quently



CHAP.  
III.

quently be changed; for when once they are thoroughly wet, they do not readily absorb the Sweat; and besides this, there is a Chance lest the morbid Exhalations should be absorbed from the wet Linen by the cutaneous Veins, and again conveyed into the Mass of Blood.

When the Fit is over, and the Patient has got a little rest after his Fatigue, it is needless, or rather hurtful, to confine him constantly to Bed, as is the *Spanish* Custom; yet he ought to keep within Doors, or at least avoid being exposed to the Sun, and all violent Exercise; which is apt to bring on Inflammations in the Bowels, and turn an Intermittent into a continual Fever.

In the Paroxysms, no Kind of Food should be given, unless the Fits are of an uncommon Length, and the Patient very weak. But every three or four Hours during the Intervals, a few Spoonfuls of thin light Nourishment will be necessary



cessary to support Nature. The *French* and *Spaniards* use weak Mutton or Chicken-Broth in all Fevers; which Practice is justly condemned by RIVERIUS\*; as Panada and other vegetable Dishes are not only more agreeable to the Generality of the Sick, but likewise better adapted to prevent the spontaneous Tendency of the Humours to an alcalescent State. Nor can I think the Summer Fruits are improper upon this Occasion, tho' they are forbid by most Authors, who have in this respect blindly copied after GALEN. ALEXANDER TRALLIAN † prescribes

CHAP.  
III.

\* Prax. Med. lib. xvii. c. i.

† Pomaceos fructus, exquisitis tertianis liberaliter exhibere convenit, nempe uvam dulcem, Persica cocta, & non cocta, peponumque medullam; præsertim autem si etiam ægri siti vexentur. Ego sane novi me frequenter ægros, ne amplius accessione febris infestarentur, impedivisse, cum peponas prabe refrigeratos, horâ ante incursum morbi, assumere jussissem, et rursus aquam temperatam copiosam, et quantam potuerint peponi superbibendam præcepissem. Secutus itaque est, non multo post aquam epotam, aliquibus sane sudor, aliis copiosa bilis per alvum. Offendi autem

ego



CHAP.  
III.

prescribes the free Use of Grapes and Peaches in Tertians, and boasts of the many Cures he had performed to the great Surprize of his Cotemporaries, by giving Water Melons, and large Draughts of cool Water an Hour before the Fit. AVICENNA recommends Pomegranates and ripe Plumbs, but above all the Water Melons\* And GALEN himself, though he was unreasonably prejudiced against the

ego plerosque Romæ medicos, qui ne nomen quidem peponum tanquam bilem procreantium proferre audebant. Quum itaque ego cuidam aliquando; et sitienti vehementer et æstu flagranti defatigatque injunxissem, ut peponem assumeret, quidam præsens medicus exclamavit homo cur ægrum magis vis occidere: Annon didicisti quod pepon bilem producat? Lege Galenum de alimentis, ubi dicat manifeste peponem comestum cholericos efficere. Laboravi igitur ego, non parum ut iis persuaderem, qui intellectu assequi poterant, Galenum non dicere hic ipsos pepones bilem creare, sed choleram efficere. Proinde tertianâ febre detentis, cum fiducia dare debent.—Quomodo enim ea, quæ refrigerant et humectant, bilem creare possint, non video.

Lib. xii. c. vi.

\* In Opere venet de Febr. c. xxxviii. p. 31.

Summer



Summer Fruits (a Surfeit of which threw him into a Fit of Sickness when a Boy) yet in the present Case he allows the Use of such of them as are not difficult to be digested †. And indeed in several other Places of his Works, he is obliged to contradict his own Opinion of their being unwholesome Nourishment †; particularly where he tells us §, “that those who are set to watch the Vineyards, and live for two Months on Grapes, Figs, and Bread, become fat and lusty.” Which Observation is annually confirmed in *Minorca*; it being remarkable that the Persons appointed for the same Purpose there commonly continue in good Health, though in that Season Tertians usually rage with the greatest Violence.

Before I have done with the Diet of the Sick, it may be noted, that altho' in the Beginning of these Fevers they commonly

† Ad Glauco. l. i. c. ix.

† De Alim. Facult. l. ii. c. ii.

§ Ibid. c. ix.



loath every thing, except thin, cooling, acescent Liquors; yet after some Days Wine becomes less disagreeable, and towards the Decline of the Disease, a prudent Use of it, mixed with Water, or made into Whey, is often absolutely necessary to keep up the failing Strength, and prevent Weakness, the worst of all Distempers.

With regard to bleeding it has been warmly debated among both antient and modern Authors where it ought or ought not to be used in Tertian Fevers. Much has been wrote on both Sides; but the Question appears to me too general to admit of a positive Answer. CELSUS\* has justly observed that “Medicines differ, “according to the Nature of the Climate; “one Kind being necessary in *Rome*, “another in *Egypt*, and a third in “*France*.” And since we daily meet with a remarkable Diversity of Symptoms

\* De Med. Præfat.



among Fevers of the Tertian Tribe even in the same Climate at the same Season of the Year, it cannot surely be surprizing that any one Remedy is not equally beneficial in all Cases, and at all Times.

CHAP.  
III.

For my own Part, when I was called early enough, in the Beginning of these Fevers I used to take away some Blood (unless there was a strong Contraindication) from People of all Ages; namely, from robust Adults ten or twelve Ounces; from others a smaller Quantity in Proportion to their Strength and Years. And farther, if a violent Head-ach, and obstinate *Delirium*, and great Heat or Pains of the Bowels were urgent, within a Day or two, I repeated the Bleeding. By which seasonable Evacuation the Vehemency of the Paroxysms is somewhat diminished; the Apyrexies become more compleat; the Operation of Emeticks and Catharticks is rendered safer and more successful; and the terrible Symptoms, which often make their Appearance about



CHAP.  
III.

the Height of the Distemper, such as raving *Sopor*, Difficulty of Breathing, Inflammations of the *Abdominal Viscera*, &c. are either prevented or mitigated.

But if before I was called, the Fever had already continued some Time, and the Mass of Blood appeared to be too much melted down, or inclinable to a putrid Dissolution, which is often the Case, during the extreme hot Weather, about the fourth Period of the Distemper, and is readily known by the great Alteration of the Patient's Looks and his sudden Loss of Strength; or if the first Paroxysms of the Disease were attended with profuse Evacuations, whether by Vomiting, Purgings, Sweating, or a Hæmorrhage from the Nose: In all these Circumstances I either omitted the Bleeding entirely, or took away a very small Quantity, though some importunate Symptoms might seem to require a much larger Evacuation,

But



But when Bleeding is allowed to be necessary, it is asked at what particular Time of the periodical Revolution it ought to be performed? The Ancients\* believed that “to open a Vein in the “Height of the Fever, was to destroy “the Patient; and therefore they order you to wait for the Intermiffion, or the Hour on which the Symptoms are generally most moderate. But the Experience of the present Age has taught us that this Operation is safe enough, at any Time of the Period, unless while the cold Fit lasts, or is soon expected; or while the Skin is covered with critical Sweats. Of late Years, encouraged by the Example of some Practitioners of Reputation †, I commonly opened a Vein in the Beginning of the hot Fit; by which Means the Sick were immediately relieved; the im-

\* Si vehemens febris urget, in ipso impetu ejus, sanguinem mittere hominem jugulare est, expectanda ergo intermissio, &c. CÆL. I. ii. c. x.

† Astruc on Fevers, p. 71. Gourraigne de Febribus.



CHAP.  
III.


moderate Heat of the Body (which is often productive of fatal Effects) was diminished; and the critical Sweats were brought on sooner, and in greater Abundance. But when that Time of the Paroxysm was passed before my being called, I bled in the Evening, when it abated or went off, that I might be at Liberty next Day to make use of the Remission or Intermiffion, which commonly happens in the Morning, to evacuate the first Passages.

For, the disagreeable Taste in the Mouth, loathing of Food, Giddiness, Pain in the Forehead and Loins, and other constant Attendants of Tertian Fevers, make it evident, that the Stomach and Intestines are overloaded with noxious Humours, and particularly with corrupted Bile; from which if they be not early discharged, very threatening Symptoms will be apt to ensue about the State of the Disease; such as violent Vomiting, Redoubling or Continuation of the Paroxysms,



roxyfms, Ravings, Restleffness, Pain, Inflammation, Gangrene of the abdominal Bowels, and laftly, fudden Death. Wherefore it is not only neceffary to wash the Stomach with warm Water or weak Broth in the beginning of the Fits, when indicated by an Inclination to vomit, and to keep the Belly constantly open with Glyfters; but likewise to empty the firft Paffages by more powerful Means at the firft convenient Interval,

CHAP.  
III.



It is a controverted Point whether it is beft to difcharge thefe noxious Humours by Vomit or Stool. At firft View Vomits feem to be moft eligible, as they quickly empty the fuperior Part of the alimentary Tube, which appears to be the principal Seat of the morbid Matter. But it muft be confidered that whatever irritates much, and produces violent Comotions, ought to be avoided in the prefent Cafe, *Cave ne inducas Effervescen-  
tiam Bilioforum*, is a Caution given by AVICENNA; and the Spaniards no more than



CHAP.  
III.

than the *Italians* \*, if their Physicians may be credited, cannot well bear rough Medicines of any Kind. Besides, the Inflammations of the Bowels, too frequently accompanying Tertians, are exasperated beyond Expression by the strong Contraction of the Diaphragm and abdominal Muscles in this Operation: And if the Spleen or Liver are disposed to become putrid (which is no uncommon Case in these Fevers) it is needless to point out the dangerous Consequences, that may result from the repeated Efforts of vomiting. For which Reasons, mild Purgatives, though less powerful Remedies, are the safest, and therefore to be preferred in the Generality of Cases. Those which I have found most beneficial are *Senna*, *Manna*, *Cremor Tartari*; but above all the *Sal Catharticum Amarum*, which neither gripes nor disturbs the Body, and seldom fails of having the desired Effect in a few Hours, a Circumstance of great Moment where the Intervals are short.

\* Vid. Bagliv. lib. i. c. xv. §v.

But



But if Vomits are to be used, they should be given in the Beginning of the Disease, before repeated Paroxysms have brought on Inflammations, or too much dissolved the Texture of the Blood; taking Care that the Operation does not interfere with the Fit, lest some sudden Mischief should arise from the united Shock of the Remedy and the Disease.

CHAP.

III.

When I first became acquainted with these Diseases, the uncommon Violence of their Symptoms induced me to lay the principal Stress of the Cure on Evacuations; and to have recourse to frequent Bleeding upon account of the Inflammations of the *Viscera*: endeavouring at the same time by repeated Cathartics to discharge the corrupted Humours from the Intestines: But when Experience had convinced me that the Bark was both a safe and effectual Remedy in these Circumstances, I then plainly perceived that such profuse Evacuations were unnecessary if not prejudicial; and of late Years, as

I fel-



CHAP. I seldom omitted to bleed and purge once  
 III. or twice, I rarely repeated either Operation  
 oftener.


In Semitertians and Remittents, which approach to the Nature of continual Fevers, I give a Cathartick early in the Morning of that Day on which the Symptoms are most moderate; hastening the Operation with Glysters (if occasion requires) so that it may be finished before the Middle of the Day, about which Time the Patient commonly grows worse. In true simple and double Tertians there is generally an Interval every Morning, in which the Purgative may be administered, but that which succeeds the worst Fit, is the most proper, as it is more calm, and continues longer than the other.

Another Inducement to purge in the Beginning of Tertians, is, that these Fevers are sometimes accompanied with Worms in the first Passages.

Proper



Proper Evacuations being premised, if possible, within the first four or five Days of the Distemper, I carefully examine the Condition of the Patient, during the third Revolution, and determine accordingly in what Manner it will be necessary to proceed. If the Paroxysms of that Revolution be neither longer, nor attended with more threatening Symptoms, than those of the second: if the Patient preserves his Strength, bears his Illness easily, and Signs of Concoction appear in the Urine; I frequently trust the whole Business to Nature, which commonly terminates the Fever about the fourth or fifth Revolution, and, for the most Part, with an Increase of some of the natural Evacuations; so that Sweats, cloudy or thick Urine, and bilious Stools often supervene, and sometimes a spontaneous Efflux of Spittle, or a copious Expectoration of pituitous Matter.

CHAP.  
III.

But



CHAP.  
III.

But if the Paroxysm on the fifth Day be evidently the longest and most severe that has happened; if it be attended with any doubtful or dangerous Symptom; if the Sick become giddy, feeble, and languid; in these Cases, without Delay, I have recourse to the Bark; and the same Evening, as soon as the Sweats have procured a Remission, I order two Scruples or a Drachm of it in Powder to be given every two or three Hours, or every Hour and a Half, so that five or six Drachms may be taken before next Day at Noon, with as little Interruption to their Sleep as may be; and the Assistants are strictly enjoined to comply punctually with these Directions, lest if this Interval escape, we should not afterwards have a favourable Opportunity of giving a sufficient Quantity of the Medicine, as the Fits about this Period of the Disease are wont to become double, subintrant, or continual. Yet it is not always in our Power to put an immediate Stop to the  
Fever



Fever by this Means: On the contrary, do what we can, it will often proceed in its Career, and, in spite of all our Attempts, run obstinately on to the seventh or ninth Day: But the great Advantage which accrues from the early Use of the Bark, is, that it invigorates the Powers of the Body, it prevents or removes the dangerous Symptoms, and in Tertians, which of their own Accord would continue to the End of the second Week or longer, it brings on a Crisis sooner, and with much less Disturbance. In short, to use the Expression of one of the greatest Promoters of medical Knowledge in this Age\*, “it proves an excellent Assistant to Nature, in what the Antients called the Concoction and Maturation of the morbid Matter;” and (I must add) in the Expulsion of it likewise, sensibly or insensibly, by the most convenient Outlets. For so far is it from suppressing any beneficial Discharge, as

\* MONRO on the Use of the Bark in Small Pox and Gargrenes, Med. Essays, vol. v. art. x.

some



CHAP.  
III.

some have asserted, that we daily observe a laudable Separation in the Urine, warm, profuse, universal Sweats, plentiful bilious Stools, and sometimes the Hæmorrhoids and Menses coming on after it has been used; though it effectually restrains the colliquative Night Sweats, to which Persons weakened by tedious Intermittents are incident.

Having given the Bark in the Manner directed, on the fifth Day of the Fever, if a Paroxysm comes on on the sixth, and declines the same Evening, I order a few more Doses to be taken, with a View to prevent, if possible, or at least to mitigate the Fit expected on the Seventh. Yet it sometimes happens that the Fit of the sixth Day unites with that of the seventh, no Remission intervening, so that the Heat, Restlessness, Raving, and other Complaints being greatly augmented, the Case seems to be much more desperate than ever. But these Commotions which follow upon the Use of the Bark in this Stage of the Fever, are more dangerous

in



in Appearance than Reality; and so far from being alarmed at the Sight of them, I commonly give Expectation that a Remission with profuse Evacuations will happen the next Evening; at the same Time giving positive Assurances that if the Patient takes as much Bark in that Interval, as he did in the former, he will either have no more Fits, or moderate ones, which will quickly yield to the same Sort of Management.

CHAP.  
III.

By this Method, when Assistance is timely called, the most formidable Kinds of intermitting and remitting Tertians, whether appearing in their own proper Colours, or personating other Distempers, may be certainly and speedily brought to a happy Conclusion about the End of the first Week or beginning of the Second.

But if the Fever has been neglected in the Beginning, and you are not called till about the third or fourth Period; when by the Use of spirituous Liquors, strong Vomits, or violent Exercise, the Bowels

P

are



CHAP.  
III.

are inflamed; or for want of seasonable Evacuations, the first Passages are overcharged with corrupted Gall, and other putrid Matter; the Fits tedious and subintrant; or perhaps attended with a *Stupor*, *Syncope*, *Cholera Morbus*, cold Sweats, and great Weakness. These, indeed, are terrible Cases, yet such as too frequently happen, and involve the Physician in the greatest Perplexity. So many Indications and Contraindications present themselves at the same time, that by obviating one Symptom you hazard the Increase of another\*: Nor, under such Circumstances, is it easy to lay down Rules for managing the Sick: All that can be with Safety advised, is, to palliate

\* Sæpissime ad ægrotos vocatus, tantam tamque confusam, mirabar symptomatum turbam, ut purgatione, an venæsectio vel neutra, imprimis foret eligenda decernere anceps et summi ponderis negotium esset. Nec doctorum, ut gravissimorum, consilia poterant in illis casibus suffragari, quin in Alterutra operatione tentanda adhuc extaret ambiguitas.

Guidet. de tertian. autumn. apud Bianch. hist.


Hepat. par. iii. p. 287.

the



the most pressing Complaints in the Manner hereafter proposed; in the mean Time carefully watching the Evening, the Night, and the early Part of the Morning; for a Remission; and the Moment it offers, to fly to the Bark as to the only Remedy which can avert the impending Danger.

CHAP.  
III.



If the Patient seems strong enough to bear Purging, I divide an Ounce or six Drachms of *Sal Catharticum amarum*, and half an Ounce of Bark, into four equal Parts, and order one to be taken every two Hours: The effect of this is, that the next Fit is mitigated, and an Intermision commonly ensues, in which the Bark without the Purgative must be repeated to finish the Cure\*.

\* Si tamen vacuatione opus sit, et ab urgente febre, tempus ad exhibendum catharticum denegetur, post V. S. cortex Peruvianus, cum purgante medicamento idoneo conjunctus, statim exhibeatur.

GEOFFR. Mat. Med. vol. ii. p. 188.


Tuto igitur in perniciosis his febribus, febrifugum quocunque tempore potissimum purgantibus immixtum propinetur

BIANCH. Hist. hep. par. iii. p. 287.

P 2

But



CHAP.  
III.

But if the Patient is so excessively feeble, that there is a manifest Risk of his dying in the next Period of the Fever, instead of the *Sal Catharticum*, I give the Bark with Cordials (among which Wine is by much the best) and endeavour to have six or seven Drachms of it taken in the Space of ten or twelve Hours; having found by Experience that the Paroxysms, if a smaller Quantity is given, too frequently come on earlier than usual, and make all Attempts to preserve Life unsuccessful\*.

When the Sick are out of Danger, and have recovered a sufficient Degree of Strength, Bleeding and Purging may safely be directed, notwithstanding the Use of the Bark, if these Evacuations are

\* For which Reason, in such deplorable Cases, TORTI orders half an Ounce or six Drachms of Bark to be swallowed at once; which he affirms is much more powerful than the same Quantity divided into several Doses.


Vid. Therap. Spec. l. iii. c. iii.

indicated;



indicated; and so far is opening the Belly occasionally from bringing on a Relapse, that those who have had the best Opportunities of making the Trial\*, have found it to be the most probable Means of preventing a Return of the Fever.

CHAP.  
III.



This is the Method of using the *Cortex*, into which I at length fell, after trying a

\* Qui distitant febrem per corticem Peruvianum deletam, si postea Cathartica propinentur, revocari, hosce toto quidem cœlo hallucinari, assidua nos docuit experientia, qua ægros post exhibitum corticem a febre liberos, per subrogata purgantia tutius a relapsu præcaveri recognovimus.

BLANCH. Hist. hep. par. iii. p. 283.

Mixtionem Rhabarbari cum Chinâ ipse ego apud Italos, xx circiter abhinc annis, primus faustam prosperamque hisce in casibus expertus fui, felicemque exitum, cum aliis, per epistolas communicavi.

Lancif. lib. ii. epid. iv. c. vi.

Cortex cum rhabarbaro anno 1710, non ea felicitate stimulat alvum, qua anno 1708 & 1709, et propterea opus erat illius remedii usum grandioribus catharticiis interrumpere.

Lanc. epid. iv. c. viii.



CHAP.  
III.

great Variety of others. Sometimes, indeed, extreme Weakness, or some formidable Symptom, obliged me to have Recourse to it in the second Period of the Fever; but I never chose to give it before the third, nor to delay it after the fourth, in all Cases of any Consequence, provided there was a proper Interval for its Administration; and with its Assistance, if the Patients were not altogether exhausted, I had the Pleasure of seeing them happily rescued from Tertians of the most malignant Kind, such as the *Lipyrria*, *Assodes*, *Febris Syncopalis*, &c. What the Poet says on a different Occasion\* being literally true in the present Case.

*Hi motus, et hæc certamina tanta  
Pulveris exigui jactu compressa quiescunt.*

And the more Experience I had of the Bark, the more I was convinced of both its Innocency and Efficacy; so that I heartily wish I had always given it with

VIRGIL, Georg. iv.



as much Freedom as I did during the last seven Years of my Stay in *Minorca*: But the Prejudices against this Medicine, which I had early imbibed from some of the most approved Authors, made me for a long time use it with too much Diffidence\*.

CHAP.

III.

In the first Place, I suspected that the Relapses, so frequent from *July* to *January*, were in some Measure owing to the general Use of the Bark: And as I observed,

\* Major medentium pars apud nostrates, ut vulgi calumniis, et assiduis ægrorum querimoniis se subducerent, in hac tempestate ad Peruvianum corticem confugere; At parum prospere; nam in perniciosos scopulos ægrotantes suos sæpe deduxere. Ufu febrifugi, per aliquot dies, equidem latebat sub cinere doloso ignis, verum postliminio violenter recrudescibat.

Ramaz. Conf. epid. 1690.

Adverte quæso mi Nepos, et diligenter observa febres intermittentes post epotam chinam nunquam ad veram et perfectam apyrexiam pertingere, qualis contingit, quando natura sponte per sudorem aut alias vias accessionem discutit, &c. &c. &c.

Ramaz. de usu & abusu chinæ.

P 4

that



CHAP.  
III.


that the greatest Number of Tertians went away of their own Accord in a Fort-night's Time, I thought it would be more advantageous to the Patient to suffer a few Paroxysms, and, when no immediate Danger appeared, to wait the spontaneous Termination of the Fever, than to hazard a Return by having it prematurely suppressed: but afterwards, by comparing a Number of Cases which had terminated of their own Accord, with others wherein the Bark had been given, I evidently saw that those were most liable to a Relapse whose Strength had been most impaired by the primary Fever, whether they had been cured by Art or Nature; so that a Delay in giving the Bark, seems frequently to have produced the Effects ascribed to its having been used too early.

Besides, while I was waiting for the spontaneous *Crisis*, even in Cases where the Mildness of the Fits, and the Length of the Intermissions, afforded the most flattering Hopes of a favourable Issue, I

now



now and then had the Mortification to find my Patient unexpectedly seized with a violent malignant Paroxysm, attended by a *Stupor*, Speechlessness, and apoplectick Symptoms, which seldom indeed proved fatal immediately, but were often succeeded by such insuperable Weakness, that the Bark could not be administered, or was given unsuccessfully, so that he died in the next Period of the Fever. The first two or three Accidents of this Kind I was willing to impute to some concealed Irregularity in the Use of the Non-naturals; but I have since been taught by too many Instances, that during the Months of *July*, *August*, and *September*, it is very common for these fallacious Fevers about the End of their second Week, suddenly to change from the mildest to the most formidable Aspect; and consequently that it is dangerous to permit their continuing so long. The Repetition of such Accidents first induced me to think of writing on this Subject; considering it as an indispensable Duty to point

CHAP.  
III.

out  
of



CHAP. out the Danger of such Omissions, to  
 III. Practitioners of less Experience, in order  
 to prevent them from being misled by  
 the plausible Theory of some Authors,  
 and the positive Assertions of others,  
 though delivered in so dogmatical a Man-  
 ner, as if they were wholly the Result of  
 careful Observation. So that I must in-  
 culcate it as a Rule of the utmost impor-  
 tance in the Cure of these epidemical  
 Tertians, never to wait longer than the  
 End of the first Week, or the Beginning  
 of the second, for their spontaneous Ter-  
 minations, but without farther Delay to  
 apply to the Bark.

Secondly, Before I learned, by observ-  
 ing the Course of these Fevers when left  
 to themselves, that it is customary for  
 them, in their Progress, to vary their Type,  
 and increase in Violence to the third,  
 fourth, or fifth Periods, I suspected that  
 the hasty Administration of the Bark not  
 only produced the Inflammation of the  
 Bowels, *Delirium*, and other bad Symp-  
 toms, which supervene about the Height  
 of



of the Disease, but likewise occasioned the Paroxysm to double, or perhaps continue without Remission\* ; whereas the

CHAP.  
III.

Fact


\* *Chinam chinæ dare impuro corpore, id est in principiis morborum, nullis precedentibus signis coctionis, et corpore non purgato, piaculum est in Aere romano; methodus damnabilis ac perniciosæ. Dicunt multi dandum esse, ut impetus symptomatum tunc furentium coerceatur ac refrænetur; ut inde æger, symptomatum sopita vi, diutius possit morbo resistere. At si tu loco frænandorum symptomatum, parvos humores per chinam chinæ in aliquo viscere figas, ac concludas, & ita internam parias inflammationem, ut frequentissime observari, nonne tu culpandus eris? Nonne tu reus necis lege aquilia puniendus?—Febricitantes meos curo per leges coctionis & crisiûm—et raro cum recidivâ: quam recidivam singulis momentis ab usu chinæ chinæ expectato.*

BAGL. de Fib. Mot. Spec. c. xiii.

Pariter rubris existentibus urinis, et supra modum tincti, cave cane pejus et angue, ne chinam chinæ præscribas, sive sint acutæ sive sint intermittentes febres; nam acutæ, facta internâ inflammatione statim præcipitabunt in deterius; intermittentes vero statim fi-ent continuæ, graves, periculosæ; quare si in aliis morbis, certe quando isti conjunctam habent urinam nimis rubram, patiens, longa, prudensque humorum coctio,

semper



CHAP.  
III.

Fact is, that many Tertians which intermit during the first Week, are very apt of their own Accord, to become continual Fevers in the second, and extend to the seventeenth or one-and-twentieth Day, or farther, before an Intermission is again perceptible; but if any interval, however obscure, appears about the fifth or seventh Days, and you can give them five or six Drachms of Bark as above directed, you will probably obtain a more perfect Remission in the next Period; during which the like Quantity must be given; and thus by repeating the Remedy as Opportunity offers, the Disease may be brought to an End, about the ninth, eleventh, or thirteenth Day. Nor are we to hesitate in giving the Bark, upon Account of the

*semper expectanda; si secus feceris, vel mortem vel longos et incurabiles morbos expectato. Romæ scribimus in Aere Romano—sancte fateor fere centies hujusmodi veritatem expertus sum, Romæ in ægrotantibus et sæpe cum magno animi mærore; quando medicos in contrariam ire sententiam observabam.*

BAGL. Prax. Med. p. 71, &c.

Crudity



Crudity and Redness of the Urine, which I have frequently observed to become paler, turn cloudy, or let fall a Sediment, by the Use of this Medicine.

CHAP.  
III.

Thirdly, I was a long time in doubt, whether the Bark might be given without Prejudice, while the first Passages were full of vitious Humours, and the Bowels were inflamed, or affected with inveterate Obstructions\*; but I have now good Reason

\* Vid. BOERHAAV. Aphor. 767.

Nonnulli in his casibus (nempe tritaephyis, hemi-tritaëis et aliis malignis tertianis) solent more solito chinam chinæ præscribere; quo autem cum successu, pluribus in locis hujus operis animadverti. Nam hoc remedium impuro corpori dare, sæpe in ægroti perniciem vertitur; potissimum in maximo apparatu humorum in mesenterio.

BAGL. Prax. Med. p. 58.

Romæ scribo et in Aere romano: Et ideo garriant quicquid velint chinæ chinæ fautores: aliis forsan in urbibus egregium est remedium, hic noxium experior.

BAGL. Ibid.

Si chinam dederis (ut fataliter plurimi faciunt) ventre adhuc humoribus onusto, tria expectato, aut inflammationem,



CHAP. Reason for asserting, that in these very  
 III. Cases this Medicine is of the greatest Use;  
 as it averts sudden Death, and gains us  
 Time to join with it other Means towards  
 compleating a Cure. For the Quantity of  
 acrimonious Contents in the *Primæ Viæ* is  
 the Effect of the Alteration produced in  
 the circulating Fluids by the Fever; and  
 the longer this continues the more Impu-  
 rities will be accumulated, till at last  
 they bring on a violent *Cholera Morbus*;  
 or perhaps make their Way through the  
 Lacteals into the Habit, and there occasion  
 very fatal Effects; all which might have  
 been prevented by the Use of the Bark,  
 which removes the Cause of these Impu-  
 rities by putting a Stop to the Fever; and

tionem, aut lentam ac diuturnam febrem, aut mortem.  
 Observa bene, et si falsa dixero, me redarguas. Ro-  
 mæ scribo et in Aere romano.

BAGL. de Fib. Motr. c. xiii.

In semitertiana (inquit HOFFMANNUS) omittantur  
 pulveres adstringentes nec non cortex chinæ, &c.

Med. Rat. tom. 4. § i. c. v.

by



by corroborating the Solids, enables them to throw off the excrementitious Fluids by the proper Emunctories.

CHAP.  
III.  
~

Inflammations of the *Abdominal Viscera* are likewise natural Effects of Tertian Fevers: For we find that they often come on by little and little, and increase with every Paroxysm, till at last they end in a Gangrene: Whereas the *Cortex*, by bringing the Fever to a speedy Conclusion, impedes the farther Progress of the Inflammation; so that it afterwards goes off gradually of its own Accord, as I have had Occasion to observe in a Multitude of Instances where acute fixed Pains, Tension, and other Symptoms, made the Nature of the Disease too plain to be doubted of.


When the Bowels are obstructed, we are told\* that the Fever should be permitted to continue, in order to remove the Obstruction; and this in many Cases

\* VANSWIET. in Aph. BOERHAAV. 767.

may



CHAP.  
III.



may be adviseable: But at the same Time it is well known, that if the obstructing Matter be suddenly dissolved and pushed into the Blood, it is capable of occasioning the most terrible Effects\*. And therefore I have commonly found it expedient, in Persons troubled with hard overgrown Livers and Spleens, to prevent the Repetition of long, severe, burning Paroxysms, lest worse Consequences should ensue. After the Sick recovered their Strength, I endeavoured to reduce the Swellings of the Belly, by the Use of saponaceous Gum Pills, washed down with an Infusion of Juniper Berries.

When there is an icteritious Colour in the Eyes, we are likewise told †, that the *Cortex* should not be administered; tho' in my Opinion, it is for the most Part dangerous to delay it, after the first Appearance of that Symptom; which is

\* Aph. BOERHAAV. 1104.

† VANSWIET. ubi supr. & HUXHAM on Fevers.



often succeeded by a Yellowness of the whole Body, arising in this as well as in other malignant \* Fevers from a total Corruption, or gangrenous Disposition of the Mass of Blood, and is too frequently the Harbinger of Death.

CHAP.  
III.

Upon the Whole, I am convinced that the unhappy *Metastases*, which some have observed to follow the Use of the Bark †, are exceeding rare, and ought rather to be ascribed to other Causes than to this Medicine. And I will venture to affirm, that more bad Consequences ensue from giving it too late than too soon; Prostration of Strength, sudden Death, or the most obstinate chronick Diseases, if the Sick recover, being the usual Effects of Delay: Whereas the worst that commonly happens from the too early use of it is, that it does not at once restrain the Paroxysms, like a Charm, without any sen-

\* See WARREN on the malignant Fever of *Barbadoes*, p. xii.

† *Med. Essays*, vol. iv. art: xxiv.

Q

fible



CHAP.  
III.

fible Evacuation, as it frequently does, when given after the Fever has arrived naturally to its Height, and begins to decline of its own Accord.

Having delivered the general Method of treating Tertians, it is needless to say much about their Symptoms; as they commonly disappear with the Fever itself, and seldom require a separate Cure. Nevertheless, it will not be amiss to take Notice of a few, which I have found the most troublesome.

In the first Stage of the Paroxysms, acute Pains in the Back and Limbs, with and without *Rigors* or Chilness, frequently happen; and I have sometimes known them so intolerable, and accompanied with such inexpressible Anxiety, that Persons of the soundest Judgment and Morality have been in Hazard of destroying themselves to get rid of it. Before I was aware that these Pains were the Fore-runners of Tertian Paroxysms, I commonly had  
recourse



recourse to Bleeding, and without observing it was attended with any ill Effect: But of late Years they never alarmed me, being well assured they would go away of themselves, as the hot Fit advanced.

CHAP.  
III.  
~

The Vomiting and *Nausea*, after washing the Stomach two or three Times with warm Water, is commonly allayed by Draughts of Salt of Wormwood, Lemon-Juice and Mint-Water. And if a constant Vomiting or Purgings hinders the Bark from being retained, small Doses of *Laudanum* should be joined with it: But a moderate bilious Discharge is commonly beneficial, and therefore should never be suppressed.

Restlessness and Head-achs, are the inseparable Companions of the hot Fit; so that the Sick must either be encouraged to bear them with Patience, or must be amused with some innocent Prescription till the Sweats carry them off. When the Head-achs are constant and troublesome in the Apyrexies, I order the Limbs

Q 2

to



CHAP. to be frequently bathed in warm Water,  
 III. and Cataplasms of Horse-radish and Leaven  
 } to be applied to the Soles of the Feet.

Hæmorrhages of the Nose are often of signal Service, in removing the obstinate Head-achs, and Pains in the abdominal *Viscera*, whether they are *e directo* or not; though for the most part they happen from the Nostril of that Side, in which the Pains are fixed; for which Reason they ought not to be hastily stopt, unless they continue too long, or come at Time of great Weakness.

When the Belly is sore and painful to the Touch, Glysters, and warm Fomentations, give great Relief. When the Pains are extremely violent, Bleeding must be repeated: but during the hot Weather, we must not be too free with the Lancet after the fourth Period of the Distemper; as the Weakness commonly is then too great to admit of any such Evacuation with Safety.


If



If the Patient continues comatous and stupified longer than ordinary, we must endeavour to rouse him from that lethargick State by scarifying, cupping, and blistering the Neck and Back.

CHAP.

III.



Troublesome Hickups often cease, upon applying Cupping-Glasses without Scarification to the Pit of the Stomach, and giving small Doses of *Laudanum* with *Tincture of Castor*.

If acute Pain and Pulsation in the Hip, point out that a critical Abscess is about to be formed there (a Case that hath occurred to me five or six Times) it must be brought forward by Cupping-Glasses, and Cataplasms applied to the Place; and as soon as a Fluctuation can be perceived, it must be opened by a deep Incision; otherwise, the Matter being lodged under the posterior double Edge of the *Glutæus externus*, instead of pointing outwards, may insinuate itself betwixt the Muscles of the Thigh; or may perhaps work its

Q 3

Way



CHAP. Way into the Cavity of the *Pelvis*, by the  
 III. Hole through which the *Musculus Pyri-*  
 ~~~~~ *formis*, and the sciatick Nerve pass out;  
 which seem'd to be the Case in one Man,  
 who had first an Abscess in his right Hip,  
 and some Time afterwards another in the  
 left, and died consumptive from the im-  
 moderate Discharge.

Parotids must also be brought to suppu-  
 rate as soon as possible: But these are not  
 very common in *Minorca*; nor are they  
 much to be wished for, being oftener symp-  
 tomatical than critical.

To conclude, as Tertians personate al-  
 most all Distempers, so there are but few  
 Disorders to which the human Body is  
 incident, that will not sometimes appear in  
 the Course of these Fevers; and these,  
 when importunate, must be mitigated, as  
 in other acute Diseases.

It has been already remarked, that  
 whether the Fever is cured by Art or Na-  
 ture,



ture, there is a Hazard of its returning within a Fortnight or three Weeks; nor could I ever fall upon any certain Method of preventing one Relapse after another, till the cold Weather, which commonly sets in about *Christmas*, had braced the Solids, and given them sufficient Strength to throw off the excrementitious Humours by the proper Out-lets; for which Reason I always advised those who were attacked in the Beginning of the Season, to leave the Island, if their Circumstances would permit, and not to return until the Spring; And there are many Instances of Persons being greatly recovered by the Change of Air, even in the first two or three Days of their being at Sea. But if they were obliged to remain in the Island, the best Chance they had of escaping a Relapse was to take a Dose of the *Cortex* every Morning and Evening for several Weeks; and now and then a gentle Purgative, if a Bitterness in the Mouth, Loss of Appetite, Swimming in the Head, or Sickness at the Stomach, the common Forerunners of it, should be perceived.

Q 4

But



## Of TERTIAN FEVERS.

But if, notwithstanding all Precautions the Fever returns, the Patient must be treated as in the first Attack; with this Difference, that as Relapses are attended with less Heat and Inflammation, Bleeding, especially in the Fit, must be used with more Caution, or altogether omitted, and Vomits may be given with greater Freedom. And, as they happen mostly in a cooler Season, they do not threaten such immediate Destruction, and consequently may be longer trusted without the Bark, if you are disposed to attempt a Cure by other Methods; though that Remedy must never be delayed when the Fits are violent or protracted. In this Manner the Patient must be content to go on till the Turn of the Year alters the Constitution of his Body, and restores him to his former Health.

It has been insinuated by some, that a long continued Use of the Bark, was liable to bring on nervous Complaints and Lowspiritedness: but in the Multitude of Cases, wherein I have been obliged



to make Use of it, I have not been able to discover that it had these or any other ill Effects, when given in the Manner above-mentioned. And as to its occasioning Dropsies, I do not remember to have met with above one Instance these ten Years, where there was a Possibility of alledging, it gave Rise to that Distemper.

CHAP.  
III.



It is not uncommon for a Looseness to come on in the room of a Relapse. Sometimes instead of a Tertian, a Quartan supervenes; which nevertheless may be speedily taken off by the Bark; but if left to Nature alone it will probably continue to the Spring; or perhaps first turn to a double Quartan, and afterward to a triple Quartan; which, as CELSUS observes, is a dangerous Distemper\*.

With respect to the Tertians, which now and then appear at other Times of the Year, they are more of a chronick

\* Lib iii. c. xv.

Disposition



CHAP.  
III.

Disposition than those of the Summer and Autumn; the cold Fits being longer, the hot Fits less severe, and the Intermittions, more regular and perfect. Add to this, that they are not so commonly attended with critical Eruptions about the Mouth; nor do the Sick discharge such Quantities of Gall either upwards or downwards; and the Urine oftener deposites a lateritious Sediment: But so far is this Sediment from being the *Criterion* of Intermittents, that I have frequently seen it in Pleurifies, and other inflammatory Fevers; while in both Tertians and Quartans, I have, for the most Part, found the Urine clear without any Separation; and in Tertians the Sediment was oftener more like Chalk than Brick-dust.


Thus have I given the History of these Fevers, collected from an almost infinite Number of Cases, carefully minuted in the Chambers of the Sick, without trusting to Memory, or regarding what others had said on the Subject before me, or advancing



vancing any Thing, but from reiterated Examination. And the Observations relating to the Cure, which are by much the most material, have been confirmed again and again by the Experience of all who have practised Physic with Attention, among either the *English* or *Spanish* Inhabitants of *Minorca*; so that I am in hopes these Remarks will not be altogether useless to the Publick. It is well known, that contagious intermittent and remittent Fevers of the Tertian Kind, are anniversary Distempers in several of the warmer Climates both in *Europe*, *Africa*, and *America*; and even in the Northern Parts of *Europe*, they are often epidemical after extraordinary hot dry Summers\*. By all I have been able to learn of them from Authors, there is a great Analogy among them every where: And though in some

CHAP.

III.



\* HOFFM Med. Rat. tom. iv. cap. iv. sect. i.  
SHORT'S History of the Weather, &c. Anno Domini 1237. 1540. 1558. 1574. 1652. 1657. 1669.  
WINTRINGHAM, Anno 1719, and VANSWIET.  
Comment. in Aph. BOERHAAV. 767.

Places



CHAP.  
III.

Places they may require more or less copious Evacuations, than we find beneficial in *Minorca*, yet I am persuaded it may safely be laid down, as a general Rule in all Cases of Danger, to give the Bark liberally and without Hesitation about the third or fourth Period of the Disease, whether Evacuations have been previously used or not. Which Practice is agreeable to the Observations of the ablest and most experienced Physicians of different Nations; such as MORTON\* and SYDENHAM† in *England*, BARTHOLIN‡ in *Denmark*, HOFFMAN || in *Germany*, GEOFFROY § in *France*, RODRIGUEZ\*\*, and many others, in *Spain*, GUIDETTUS and BIANCHI in *Piedmont* ††, TORTI in *Modena* ‡‡, MUSITANUS ||| in *Naples*,

\* Exercit de morb. acut.

† Epist. respon. I.

‡ Ephemerid. German.

|| Med. Rat. tom. iv. § i- c. i. obs. 5.

§ Mater. Med. tom. ii.

\*\* Palæstr. Med. tom. ii. Disc. 12.

†† Hist. Hep. part. iii.

‡‡ Therapeut. special. Sparsim.

||| Pyretolog. c. xxiii.

and



and TRAVERSARIUS § in *Pesaro*; nay even in *Aere Romano*, where BAGLIVI declaimed against the Use of the Bark with so much Passion, LANCISI\* has of late Years, as well as the Cardinal de LUGO † and Father FABRI ‡ formerly, had incontestible Proofs of its being not only an innocent, but a necessary, powerful Remedy in the Cure of Tertian Fevers.

CHAP.  
III.

§ Apud Lancif. l. ii. epid. iv. c. viii.


\* Epid. Sparfim.

† Antym. Conyg. pulv. peruv. vind.

‡ Id. Ibid.







## C H A P. IV.

*Of the RASH, ESSERE, and CHOLERA  
MORBUS.*

CHAP.  
IV.

**H**AVING given an Account of Tertian Fevers, which was the chief Design of this Treatise, I shall endeavour to bring my Remarks on the other epidemical Diseases, into as narrow Bounds as possible, by passing over such Circumstances as are taken Notice of by the Generality of Authors.

The cutaneous Eruption which we call the Rash, or prickly Heat, is the *Sudamina* or *Papulæ Sudoris* of the Romans, and the ἰδρωα of HIPPOCRATES\*, who justly places it among the Summer Diseases; it being so frequent in warm Countries,

\* Aphor. § iii. No. 21.

that



that few People escape having more or less of it during the hot Weather; though Children are much more affected by it than others. It consists of numerous minute Pimples, or rather small, round, red Spots, just perceivable by the Touch, as a Kind of Roughness on the Skin, which break out on different Parts of the Body, especially after Exercise or drinking cold Water,

This Eruption is commonly looked upon as a Sign of Health; and indeed while it continues fresh on the Skin, no Inconveniency arises from it except a frequent Itching: But if accidentally the Pimples are driven in, by catching Cold, bathing in the Sea, or any other Error in the Non-naturals, the Case too often becomes dangerous. And I have constantly observed, that those who had a great deal of this Eruption in the Summer, were subject either to Fluxes, Hæmorrhages from the Nose, or Fevers, upon the Alteration of Weather about the autumnal Equinox.

Where-



CHAP.  
IV.

Wherefore, I commonly advised them, as soon as the slightest Signs of a Retrocession appeared (such as Sickness, Headach, and preternatural Heat) to lose some Blood immediately, and make use of mild Catharticks, vegetable Food, and cooling acefcent Liquors: By which Means, the bad Consequences abovementioned, if I am not greatly deceived, have been frequently prevented, notwithstanding the vulgar Prejudices against Evacuations in this and other similar Cases.

The *Essere*, so called by the *Arabians* who first described them, though not uncommon in *Minorca*, appear much more seldom than the *Rash*. They are hard, flat Tubercles, of a pale Colour and different Forms, and break out chiefly while one is warm in Bed, sometimes on one Part, sometimes on another, occasioning such an intolerable Itching, that it is impossible to refrain from scratching, and the more they are rubbed the larger they become. They seldom continue out above

an




an Hour or two at a Time, but suddenly vanish of themselves, the Skin recovering its natural Colour, and come back again as unexpectedly as they went off; the Patient having more or less Anxiety about the Pit of the Stomach, during the Time of their disappearing. The *Spaniards* call them *Ronchados*, from *Roncho* a Nettle in the *Castilian* Language, they having exactly the Appearance of Swellings occasioned by the Sting of Nettles: In *Minorca* they go by the Name of *Favas*, probably from their frequently resembling Beans in Shape and Magnitude.

It is commonly allowed by Authors, and not without Reason, that those who have this Sort of Eruption, should make use of Bleeding, Purging, and proper Alteratives; otherwise a Fever is liable to ensue\*.

The *Essere*, as has been already observed, often accompany Tertian Fevers, and appear most commonly in the hot Fit.

\* Vid. Sennert. lib. v. part i. cap. xxvi.



CHAP. Sometimes I have seen them so numerous  
IV. that the whole Body was disfigured by  
 them, and painted in many Places with  
 all the Colours of the Rain-bow. In a  
 few such Instances, which contrary to Ex-  
 pectation proved suddenly mortal, the Pa-  
 tient's Death seemed to be occasioned by  
 these fiery Pustules being accidentally  
 thrown upon the Brain, instead of the Skin.  
 For which Reason, wherever I met with  
 them in great Abundance, I took Care to  
 make as liberal Evacuations as the Strength  
 and Age of the Patient would permit, and  
 proceeded speedily to give the Bark; which  
 effectually cures both them and the Fever.

As to the *Cholera Morbus*, it must be  
 managed according to the Method first  
 hinted by HIPPOCRATES\*, and of late  
 described more fully by SYDENHAM †.  
 To which I must add, that if feverish

\* Cholerae morbo conferunt, si quidem dolor adsit,  
 anodyna; venter autem superior, tum inferior poti-  
 onibus humectantibus curandus.

HIP. de Affect.

† De Morb. Acut § iv. c. ii.



Complaints, or fixed Pains in the Belly or Sides come on, after the Evacuation is stopt, (as is frequently the Case) it will be necessary to bleed and keep the Belly open.

CHAP.  
IV.  
~


The *Spanish* Physicians have often assured me, that they found nothing more beneficial in violent deplorable *Choleras* than drinking of cold Water. Which Practice is recommended by many of the Ancients\*.

When the *Cholera Morbus* returns periodically, like a Tertian, it must be cured as these Fevers are; but it denounces much more Danger when it attends the Fits about the third or fourth Period, than it does in the Beginning of the Disease.

\* Sin autem omnia antiqua stercora dejecta fuerint, et biliosi humores transierint, biliosusque, vomitus et distentio adsit, fastidium, anxietas, virium labefactatio, tunc frigida aqua Cyathi duo aut tres propinandi sunt ad ventris astrictionem, ut retrogradus humorum cursus cohibeatur, atque stomachus ardens refrigeretur. Assidue vero id, quum potam aquam vomuerit, facito.

Aret. Cappad. de curat. M. A. l. ii. c. iv:  
See likewise Cæl. Aurel. de morb. acut. l. iii. c. xxi.





## C H A P. V.

*Of the D Y S E N T E R Y.*

CHAP. V. **B**Y perusing the Accounts which I kept of the Sick, while Dysenteries were epidemical in *Minorca*, I find that they began three different Ways.

*First*, Sharp indigested Humours flowing from the Stomach, or (what seems to be more frequently the Case) an increased Secretion of the Bile and other Liquids necessary for Chylification, produce a simple Looseness; which gradually washes off the *Mucus* from the Intestines, erodes their Membranes, and at last brings on severe Gripes with bloody Stools.


*Secondly*,



*Secondly*, Others are seized with *Horrors*, *Rigors*, and all the Train of feverish Symptoms, which commonly attend the first Attack of *acute* Diseases; and in a short Time afterwards, a painful frequent Evacuation of Slime streaked with Blood, ensues.

CHAP

V.



*Thirdly*, Others have no proceeding Fever, but are seized at first with a Twisting of the Guts, which (as they express it) draws up their Bowels into Knots; and many, instead of griping Pains, which shift from Place to Place, and come at Intervals, have acute fixed ones in some particular Part of the Belly, which occasion Complaints as various as their Seat; some being tormented with Stitches about the bastard Ribs, interrupting their breathing freely, as in the Pleurisy; others with a Pain reaching from the one *Hypochondrium* to the other, cutting them, as it were, in two; while others complain only of a Pain about the *Pelvis*, with a constant, fruitless, Straining to Stool; though the Body is, for



CHAP.

V.

the most Part, costive, or discharges nothing but bloody Slime.

But in which soever of these Ways Dysenteries begin, in Process of Time the Case comes to be pretty much the same; the Intestines are irritated, inflamed, and ulcerated; a Fever, for the most Part of the periodical Kind, comes on; the constant *Stimulus* in the Bowels diminishes the cutaneous Discharges; and, of Course, a greater Proportion of Fluids is thrown upon the Guts; the Flux increases, and what is discharged becomes more ichorous, and offensive to the Smell; the Strength decays; and Death, or a tedious Recovery, is commonly the Consequence.


Upon opening the Bodies of the Dead, I have constantly found the great Guts either entirely mortified, or partly inflamed, partly mortified, the *Rectum* being generally most affected: in many I have seen scirrhous Tubercles straitening the Cavity of the *Colon* in several Places; in



a few, there were small Abscesses in the cellular Membrane of the *Peritoneum* contiguous to the *Colon* and *Rectum*; sometimes the small Guts were perfectly sound in Appearance: but more frequently their lower Part was inflamed, the Convolution being often preternaturally connected to each other by Membranes, as the Lungs sometimes are to the *Pleura*. In two People the *Omentum* was almost entirely wasted (the small Remains of it being quite black) while purulent Water was found in the Cavity of the *Abdomen*; in several it was inflamed and adhered both to the Guts and *Peritoneum*; for the most Part the Gall-Bladder was full of dark Bile, and the Spleen, more or less, in a putrid Condition.

CHAP.

V.




Almost all the Dysenteries, which fell under my Observation, unless they were speedily cured in the Beginning, at best proved obstinate, and too frequently fatal, in spite of the many boasted Specificks for this Distemper; for which Reason, when-

R 4

ever



CHAP.  
V.

ever they are epidemical, the utmost Diligence should be used in applying the proper Remedies before the Strength of the Patient be exhausted, and the Coats of the Intestines too much injured. And even let the utmost Care be taken of them never so early, the Event is often dubious or unsuccessful; tho' among the common Soldiers this seems frequently to be owing to a want of means for keeping them clean, and Conveniences for easing themselves, without being obliged to get often out of Bed, and expose their Bodies to the open Air. Which Conveniences it were to be wished, that those who have the Direction of our Fleets and Armies, would order to be provided both in the Ships and Hospitals.

I shall not enter upon a minute Description of all the Rules necessary to be observed in the Cure of Dysenteries; but only to point out such Methods of treating the Sick, as Experience taught me to be the most beneficial.

When




When they begin like a simple *Diar-*  
*rhœa*, without Fever or fixed Pains in the  
Belly, the first Thing to be done, is to  
empty the Intestines of their acrimonious  
Contents as soon as possible: And the most  
efficacious Remedies for this Purpose, as  
far as I know, are the *Radix Ipecacuanbæ*,  
and the *Vitrum Antimonii Ceratum*. The  
latter, I used to give from five to ten  
Grains early in the Morning: Of the for-  
mer, I directed ten or fifteen Grains in  
Powder, to be divided into three Doses,  
and to be taken in the Forenoon at the  
Interval of two Hours, or an Hour and  
a half between each Dose. The most  
common Effect of both was to procure a  
thorough Evacuation upwards and down-  
wards during the Day; and they often  
threw the Patient into a Sweat the ensuing  
Night.

The same Remedies I have likewise  
found to be serviceable in obstinate Relap-  
ses of intermittent Fevers: But I prefer  
the *Ipecacuanbæ*, as being certain in its  
Operation;

CHAP.  
V.



CHAP.  
V.

Operation; whereas the other sometimes did not produce the intended Discharge, at other Times occasioned greater Comotions than were expected. Nevertheless, I must acknowledge that now and then, in desperate Bloody Fluxes, I have known the antimonial Medicine to be successful, after every Thing else had been tried to no Purpose. But I return to the Method of curing them in the Beginning.

At first I repeat the abovementioned Evacuations every other Day for three or four Times, and afterwards at greater Intervals, if the Disease continues; endeavouring, in the mean while, by proper Drinks, to blunt the Acrimony in the first Passages, and defend their sensible Membranes from being eroded; and every Night at Bed-time, I give a small Dose of Opium, sufficient to mitigate Pain, procure Rest, and promote Perspiration, without keeping the Sick in a constant State of Stupidity, or preventing a due Discharge by Stool; as I have seen some


do



do very injudiciously by the too liberal Use  
of Anodynes,

CHAP.

V.



But when the Dysentery began in the second or third Way above described, which was commonly the Case in the Year 1747, I confided principally in the early Use of the antiphlogistick Method, with a View to hinder the Guts from being farther inflamed and ulcerated. Wherefore, when I was called in Time to young People, I took away ten or twelve Ounces of Blood immediately, and afterwards repeated the Operation once or twice at proper Intervals, according as the Violence of the Pain and Degree of Fever indicated: At the same Time I ordered emollient Glysters and Fomentations to be frequently made use of; and plenty of warm, soft, diluent Liquors, for common Drink; shunning *Opium* as much as the intolerable Torture of the Distemper would permit.

As soon as the Fever was somewhat abated, and the Violence of the Pains asswaged,



CHAP.  
V.


swaged, I next endeavoured to procure a sufficient Discharge by Stool. For it is to be observed, that wherever the *Ilium*, *Colon*, or *Rectum* are inflamed, hardened Excrements are, for the most Part, either the Cause or the Consequence of the Disease; nor can we hope for a compleat Recovery while such offensive Matter remains in the Bowels; wherefore Purgatives of the mildest Kind may first be made use of, such as Whey, weak Broth, sweet Oil, Solution of *Manna*, *Cremor Tartari*, &c. proceeding by Degrees to the more active, till the End proposed be obtained; after the same Manner as we are directed to do, in SYDENHAM's bilious Cholick; which indeed is a Disease nearly allied to the Dysenteries I am now treating of, and is always frequent at the Time they are epidemical. In both Cases, when other Means have failed, six or seven Grains of *Calomel* with a Grain of *Opium* at Night, after the Use of the *Semicupium*, and a purging Apozem made of *Senna*, *Manna*, and *Sal Catharticum* the next Day, have proved



proved successful, and brought off a prodigious Quantity of round, hard, foetid Lumps, to the great Relief of the Patient; nor is it easy to conceive how so much had been collected, or where it had lodged so long, as in some Cases I have observed: the Patient having eat nothing for two or three Weeks, that could furnish much Excrement, and during that Time had taken several Glysters and Common Catharticks, which brought away liquid Stools.

CHAP.

V.




Having by such Methods removed the Inflammation of the Intestines, and dislodged their irritating Contents, the Tenderness of the Bowels must be mitigated by Anodynes, and the Use of such Medicines, as supply the Want of the intestinal *Mucus*.

By these Means, Dysenteries, if taken in Time, have been speedily carried off; as well as the *Tenesmus*, which is a Disease very nearly allied to the former, and  
equally



CHAP.  
V.



equally fatal, whatever CELSUS may al-  
ledge\* to the contrary. But if they were  
neglected in the Beginning, or did not  
soon yield to the Remedies, above men-  
tioned, the Case commonly became in-  
veterate and perplexing; the same Medi-  
cine which gave Relief to one, often doing  
Harm to another, as HOFFMAN justly  
observes †, In general, it answered best  
to avoid such Diet as would afford a large  
Quantity of putrid, irritating Excrement;  
to drink plentifully of mild gentle deter-  
sive Balsamicks; to give Liquids of the  
same Kind in Glysters; in short, to pursue  
with Diligence the Directions given us by  
the great BOERHAAVE in his Aphorisms ‡,  
with this Addition, that it was absolutely  
necessary to give *Opium* twice a Day, in  
order to obtain some Respite from perpe-  
tual Torment, and gradually to increase  
the Dose, from half a Grain to five or six,  
as Use made it familiar; and once or twice

\* De Med. lib. iv. c. xxi.

† Med. Rat. t. vi. § ii. c. vii.

‡ No. 966, 976.




a Week or oftener, as the Strength would allow, and the Symptoms indicated, to hinder the acrid Matter from being accumulated in the Intestines by giving Glysters, Catharticks, or small Doses of *Ipecacuanbæ*.

If by these Means the Patient can be kept alive during the first severe Winter Weather, he stands a good Chance of holding out to the Summer, which commonly restores him to his former Health, when he must be weaned by Degrees from the Use of *Opium*; from the Continuance of which Medicine, in such Cases, I have never found any ill Effects ensue; on the contrary, I could name many People who have been obliged to take it in this Manner from *September* to the following *June*; and as far as I am able to judge, it is to the *Opium* they are chiefly indebted for their Lives, and the perfect Health they at present enjoy.

The great Similitude there is in many Respects between Tertian Fevers and Dysenteries,




CHAP.  
IV.

Dysenteries, induced me frequently to make Use of the Bark in the last named Disease. When the Fever and Gripes were regularly exasperated, either every Day or every other Day at stated Periods, it has often effectually put a Stop to both; especially if the Exacerbation began with Chillness, and terminated in Sweats: At other Times it removed the Fever, the Flux continuing without much Alteration. In some Cases I have given it merely with a View to prevent the Mortification of the Intestines in the last Stage of the Distemper: But I am sorry to say it was seldom so successful as I could have wished.







## C H A P. VI.


*Of the PLEURISY and other INFLAM-  
MATORY FEVERS, frequent in the Win-  
ter and Spring.*

**T**HE anniversary epidemical Fevers CHAP.  
VI.  
in *Minorca* may be divided into two  
Classes, which at present, for Distinction's  
Sake, we shall call the Summer and the  
Winter Fevers. The former break out in  
*June* or *July*, and cease about *January*,  
or somewhat sooner. The latter seldom  
appear before *November*, and are rarely  
seen after the Summer Solstice; so that  
the one seems to be the Offspring of ex-  
cessive Heat; the other of the sudden Cold,  
which the North Winds frequently occa-  
sion. And as the same Sort of Weather

S

recurs



CHAP.  
VI.

recurs regularly every Summer and Autumn; so likewise do the Diseases: Whereas those of the Winter and Spring, agreeable to the variable Disposition of these Seasons, are neither so uniform nor so certain; being some Years more frequent in one Month, others in another. The Summer Fevers are by much the most universal, making up two Thirds, or sometimes three Fourths of the whole annual Diseases, and attacking the Inhabitants of every Rank, whether Natives or Foreigners, without Distinction: Whereas the others are less injurious to the *English* than to the *Spaniards*; and especially to the Peasants, whose Houses are commonly built upon rising Grounds, and not so well adapted for keeping out the piercing Cold of the Winter, as sheltering them from the Summer's Heat; so that I have known particular Corners of the Country almost depopulated, while the Towns and Villages escaped any remarkable Mortality.

Both



Both these Classes of Fevers, and indeed almost all others which happen in that Climate, whether primary or symptomatical Diseases, may be termed periodical, having Remissions at Intervals, more or less considerable: But those of the Summer, as has been already observed, generally assume some one or other of the Tertian Types, being worse one Day and better the next alternately: Whereas the Winter Fevers, though they often counterfeit Tertians, especially in their Beginning, yet for the most part have Exacerbations equally strong every Day, coming on about Noon with or without cold Shiverings, and terminating towards Morning, sometimes with gentle Sweats, sometimes without any sensible Evacuation.

It may likewise be remarked, that as the Summer Fevers are generally complicated with Fluxes and painful Obstructions in the *Chylopoetick Viscera*; so are those of the Winter, with Coughs, Catarrhs, and topical Inflammations of the



vital Organs, the Brain, the Lungs, the Heart itself; and hence it is that the latter are more destructive in proportion to their Numbers. Notwithstanding which, as the former are much more constant and universal, if we calculate one Year with another, we shall find that “ the Autumn produces “ the most acute and most fatal Diseases “ of any Season; and the Paroxysms in the “ Evenings bear some Resemblance to it. “ For as the Day of any particular Disease, “ is to the Year that contains the Period or “ Circle of Diseases; so is the Paroxysm “ in the Evening to the Autumn\*.” And indeed the Conformity between such of these Diseases as are described in the Hippocratical Writings and the Appearance of them at this Day in Climates near the Latitude of *Greece*, will be abundantly evident to every one who considers the preceding Account with any Degree of Attention.


\* HIP. de Morb. Vulg. l. ii.

Having



Having premised these general Reflections, in order to give the clearest Idea I can of the Winter Fevers, I shall describe them as they appeared during the last Month of the Year 1745, and the first Part of 1746, when the uncommon Destruction they made among the *English* rendered them more immediately the Object of my Attention; and as the Generality of them is called *Mal de Costat* by the Vulgar, and the *Pleurisy* by Physicians, from their being accompanied with Pains in the Side, I have in Compliance to prevailing Custom, retained that Name in the Title of this Chapter, tho' it will appear in the Sequel, that they ought rather to be termed Peripneumonies, agreeable to the Observations of ZECHIUS\*, HOFFMAN†, and others,


CHAP.  
III.



These Pleurifies began commonly like an Ague Fit, with shivering and shaking,

\* Apud Bonet. Sepulchr. Anat. lib. ii. § iv.  
† Med. Rat. tom. iv. § 2. cap iv.



CHAP.  
VI.

flying Pains, all over the Body, bilious Vomitings and Purgings, which were soon succeeded by quick Breathing, immoderate Thirst, inward Heat, Head-ach, and other feverish Symptoms. In a few Hours the Respiration became more difficult and laborious; the most Part of the Sick being seized with Stitches in their Sides, striking upwards to the Clavicle, and Shoulder Blade; obliquely downwards along the Cartilages of the bastard Ribs; or else darting across from the Breast-Bone to the *Vertebrae* of the Back; so that they could neither cough, nor make a full Inspiration without great Pain. Many complained chiefly of a Load and Oppression in their Breast, as if a Mill-stone had been laid upon it; some of a Heaviness and fluttering about the Heart, which at one Time seemed to glow with extraordinary Heat, at another to be chilled with Cold, as if it had been dipt in Ice-Water. In a few of the Sick, these Complaints preceded the Fever, in others they did not come on till the Day after,


In



In the Progress of the Disease, it was not uncommon for the Pains to move about in the Thorax from one Place to another. Sometimes they would shift from the Breast to the Limbs, and of a sudden return to the Bowels; and I have seen Cases wherein, after leaving one Side, they have attacked the other unexpectedly, and proved fatal in a very short Time. the left Side of the *Thorax* was not near so liable to be affected as the other; forty-two out of sixty Patients who were seized about the same Time, having had the Disease in the right. But whichsoever side was affected, the Sick lay easiest on the opposite; though the Generality were obliged to lie upon their Backs, or to sit up in Bed with their Heads erect. Many were drowzy and inclinable to Sleep, but they raved at Intervals, or were much disturbed with extravagant Dreams. Some laughed in their Sleep; others would awake in a Fright and start out of Bed, imagining that the House was in Flames; that those about them were endeavouring to push them over

CHAP.

VI.





CHAP.  
VI.

a Precipice; to pierce their Sides with Dag-  
gers; to bind them down with Cords, or  
iron Hoops, and Things of the like  
Nature.

In the mean Time, the external Heat  
of the Body was in several very moderate;  
in some less than natural; but for the most  
part it was so intense as to raise the Mer-  
cury in FAHRENHEIT'S Thermometer to  
the 102d Degree, and often in the After-  
noon to the 104th. The Pulse was like-  
wise very variable, not only in different  
Persons, but in the same at different  
Times, and, in respect to its Strength, in  
different Arms, that of the pained Side  
being most obscure; and I have frequently  
found it like that of a Man in Health, or  
even slower than natural\*, while the

\* Pulsus in Pleuritide minus celorem, aut fortem  
(febre tamen acutâ in summo vigore nihilo minus sub-  
sistente) sæpius notavi; Pulsus igitur celeritas & mag-  
nitude non semper cum febra inflammatoriâ sociantur.  
—Qui in pleuræ aut pulmonum inflammationibus,  
pulsui nimium fidunt, decipiuntur, &c. &c.

O'Connel de Morb. Acut. p. 235.

Patient



Patient was in the greatest Danger; so that it could neither be depended upon as a prognostick Sign, nor as an Indication of Cure. Nor was the Colour or Consistence of the Blood more to be trusted; in many it had a white or pale yellow Crust, the *Serum* being of the same Complexion; but for the most part it was red and florid. It frequently changed its Appearance, in the Space of a few Hours, in the same Person; what was drawn in the Morning having a Crust; that in the Afternoon none, *et vice versa*. And I could never positively determine which Sort afforded the best or the worst Prognostick. The Signs from which one could pronounce the Patient's Recovery, with the most Certainty, were his being able to sleep sound in the natural Posture, and to make a full Inspiration without Difficulty, while his Thirst and inward Heat were moderate.

Beside some Abatement of the Fever, which commonly happened every Morning, it was remarkable, that upon the third

Day

CHAP.  
VI.




CHAP. Day or beginning of the fourth, there was  
 VI. frequently a great Remission, sometimes a  
 ~~~~~ total Cessation of every violent Symptom;  
 so that the Sick were thought to be out of  
 Danger: but on the fourth or fifth a De-  
 lirium suddenly came on, or the Breathing  
 became more difficult than ever, and one  
 or both of these Symptoms encreasing  
 hourly, the Patient expired in a Day or  
 two, either suffocated or raving mad; un-  
 less Nature or Art assisting, he had the  
 good Fortune to escape by Means of some  
 of the Evacuations to be hereafter menti-  
 oned. It sometimes happened that the re-  
 markable Remission fell out on one of the  
 Days between the fourth and the seventh;  
 in which Case the Exacerbation succeeded  
 on the following Day.

Out of twenty one Patients, whom I  
 lost by this Distemper, four died upon the  
 fourth Day; three the fifth Day; three the  
 sixth; three the seventh; three the eighth;  
 two the eleventh; one the fourteenth; and  
 the remaining two, though the Day cannot  
 positively



positively be determined, yet from some Circumstances, it is evident, it must have been the fourth or fifth. And indeed such was the rapid Progress of these mortal Pleurisy, that if any of them survived the seventh Day, it seemed to be entirely owing to Bleeding.

CHAP.  
VI.

I examined fourteen of the Bodies, in all which the Lungs were principally affected; whilst in several the *Pleura* was perfectly sound, or only slightly attached to the Lungs, which is a common Case in Adults, as every one knows, who is accustomed to Dissections. In many, the Lungs were converted into a hard Liver-like Substance, and sunk in Water; in some the Diaphragm was inflamed; in others, large firm *Polypi* were taken out of the Ventricles of the Heart and large Vessels adjoining. Abscesses, or rather half-formed Abscesses, with a sanious *Ichor*, and a rotten gelatinous substance, instead of concocted Matter, were frequently found, even in those who died so early as the fourth Day, either in the Lungs, or between the

Lungs



CHAP.  
VI.


Lungs and *Pleura*, where they adhered, or between the Membranes of the *Mediastinum*, near the Diaphragm. And these Abscesses had sometimes emptied themselves into the Cavity of the *Thorax*, so that the Lungs floated in purulent *Serum*; their external Membrane, and likewise the *Pleura*, being greatly thickened and converted, as it were, into a white Crust\*, like melted Tallow grown cold, part of them being

\* I am now doubtful if this Crust was the *Pleura* and external Coat of the Lungs, changed from a natural State by soaking in a purulent Fluid, or if it was not altogether a preternatural Substance, formed by Fluids deposited on these Membranes, and compacted together by the Motion of the Lungs; for I have lately been informed by Mr. HUNTER, that, in those who die of internal Inflammations, he generally finds the Surface of the Cavities and the Bowels furred over, as it were, with a thick Slough of an Ash Colour, somewhat inclining to yellow, which he had an Opportunity of observing in all the different Degrees of Consistence, between a soft *Mucus* slightly adhering to the Parts, and a solid fibrous *Lamella*, so closely attached to them, that at first View they appeared to be inseparably united: Yet after Maceration in Water, he could peel off the whole of this Slough from the smooth, sound, natural Membrane which it covered. This Observation confirms HAL-



being grown cold, part of them being eroded and detached from the rest. In some dies the *Pericardium* was full of purulent *Serum*, its internal Membrane and the outer Surface of the Heart being affected in the same Manner as I have just now described those of the Lungs and *Pleura*. In two People whose Heads were examined, the *Sinuses* of the *Dura Mater* were stuffed and distended with Blood, the Membrane itself being found, and the *Pia Mater*, together with the *Plexus Choroides* was inflamed and much thicker than in a natural State.

CHAP.  
VI.




In one Person, whom I imagined to have died of a Pleurisy, the Lungs and *Pleura* were found; but there were *Polypi* in the Heart, the Diaphragm was inflamed,

LER's Opinion, that the membranous Adhesions, so frequently met with between the Lungs and *Pleura*, are generally formed by the Coagulation of the watery Fluid which oozes from the exhalent Vessels of these Parts. See Lin. Prim. Physiolog. No. 262.

and



CHAP.  
VI.



and a large Abscess was found in the right Lobe of the Liver, which had discharged itself into the *Abdomen*, where much purulent stinking Matter was collected; Part of the *Colon* and Teguments near the Liver being sphacelated, the rest of the Guts inflamed, adhering to each other, and beginning to mortify. This Man, for the first four Days of his Distemper, had no great Pain; on the fifth and sixth Day it became violent; after this he expectorated freely, which gave Hopes of Recovery to the twelfth, when he died, contrary to Expectation, putting me in mind of a Prognostick in \* HIPPOCRATES, which seemed to be applicable to his Case.

Among the natural Evacuations which terminated these Pleurifies, the most frequent was a plentiful Expectoration without hard Coughing. When this Discharge began early, and continued freely, it kept

\* Quibus autem pleuriticis, dolores initio mites sunt, ingravescent autem quinto aut sexto facile ad duodecimum usque perveniunt; ac raro illi servantur.


Prænot. coac.

off



off or mitigated the dangerous Symptoms so liable to come on about the fourth or fifth Day, and the Fever disappeared about the seventh: But if the Spitting did not begin before the Exacerbation of the fourth or fifth Day, it often proved insufficient to save the Patient. If he did recover, the Fever seldom left him before the fourteenth Day, and frequently continued much longer.

CHAP.  
VI.



HIPPOCRATES\* describes the Colour and Consistence of the expectorated Matter, which generally prognosticates Death or Recovery: But at the same Time he takes Care to inform us, that every Sort of Spitting which does not remove the Pain, is bad; and every Sort that does remove it is good: The latter Part of which Remark I have seen verified in some who owed their Lives to that Evacuation, though the Matter was always thin, crude, and ill-coloured.

The next critical Discharge to be mentioned, is a copious Efflux of Urine,

\* Prænot coac.

which



CHAP. VI. which soon after being made turns thick, and is either of a pale red, letting fall a lateritious Sediment, or milky, as if mixed with laudable *Pus*, and deposited an equal smooth white one. Such Urine alone terminated the Disease in some; and in many it proved an Assistant to the Expectoration.


Sweats were common in these Pleurifies: in the Beginning indeed, they were oftener symptomatical than critical: But after the Obstruction of the Head and Breast were removed by the Evacuations abovementioned, they seldom failed of coming on to abate the Fever, and compleat the Cure. And tho' they did not always fall out on the critical Days, yet it is remarkable, that those which were most profuse, and brought about the greatest Changes, which really happened on the 4th, 7th, 9th, 11th, 14th, 17th, and 21st Day, much more frequently than on others. Which is a Circumstance I had not always attended to, though I find from my Notes it is a true one.

It



It was common for these Diseases to begin with a Vomiting and Purging of green or yellow bilious Matter: But there were few Instances where either of these Evacu- ations could be reckoned critical, except in one Man who laboured under a dry Pleu- rify with crude urine, and had got to the eleventh Day, with the utmost Difficulty, when a Purging of porraceous Bile came on, and gave a Turn to the Distemper, which afterwards went off gradually by Sweat and Expectoration.

CHAP.  
VI.



One Person who had a violent Pleurisy in the right Breast, and had been four Times bled, was seized with a Hæmor- rhage from the right Nostril on the seventh Day; and after losing four or five Ounces, a most profuse universal Sweat came on, and put him out of Danger.

Another Way which Nature took to re- lieve herself, was, by translating the mor- bifick Matter from the Bowels to the Sur-  
T face



CHAP.  
VI.

face of the Body. In three People the Disease was, immediately after its Invasion, changed into the erysipelatous Fever, described by SYDENHAM, and was cured as he directs. In seven or eight, the second or third Day the Fever and all other Complaints vanished, an *Erysipelas* appearing on the inferior Extremities; which in some fell lower and lower every Hour, and soon made its Exit by the Toes; in others, being more of the Nature of a *Phlegmon*, it settled in one of the Legs, and formed an Abscess; which degenerated into Fistulous Ulcers, and proved very difficult to be cured.

By reflecting upon these Cases, and considering that an *Erysipelas* of the Bowels would spread from Place to Place inwardly, as it does on the Skin; I could account for several Appearances in the predominant Distempers, which at first seemed somewhat extraordinary; such as the Inconstancy and Mutability of the Pains, and

the




the frequent Alteration of the Pulse and Breathing, according as the morbidick Matter shifted its Situation, and fixed upon different Organs, the Heart, the Lungs, the Midriff, or containing Membranes of the Thorax. And I imagined that the treacherous Remission, so frequent on the third or fourth Day, happened whilst the morbidick Matter was shifting from the Breast; which after being re-assumed into the Mass of Blood, and falling upon the Head, or returning again to the Lungs, often occasioned tragical Effects.

CHAP.  
VI.

When these Pleurifies first became epidemical, their quick Progress, and uncommon Mortality, surprized me greatly. I attempted to cure them by bleeding once or twice a Day, if the complaints were violent, as I had always used to do in inflammatory Fevers: But the Remissions in the Mornings sometimes induced me to omit the Operation; and the Cessation of the



CHAP.  
VI.

Symptoms, which generally happened about the third Day, made me imagine the Danger was over; so that before the Patients were blooded above twice or three Times, the Exacerbation came on upon the fourth or fifth Days, and defeated all Attempts by Bleeding, Blistering, or otherwise to relieve them.

These unforeseen Events startled me greatly, and led me again to review the whole Progress of the Disease, its Symptoms, and Issue. I had observed that some escaped by Means of Expectoration and purulent Urine, without much Assistance from *Plebotomy*: and considering the periodical Revolutions of the Fever, the quick Transition of the Stitches from one Part to another, together with the prevailing Colour of the Blood, as well as that of the Spitting, and other Excretions, I was apprehensive that these were what Authors call bilious Pleurifies, which they alledge are exasperated by large Evacuations



cuations\*; particularly DURETUS†, who exclaims with great Vehemence against those Physicians who trust principally to Bleeding in the Cure of these Diseases, without waiting for the natural Evacuations. These Motives induced me to use the Lancet with more Caution; and to rely chiefly on the speedy Application of Blisters for restraining the Symptoms. But this Management proved less successful than the former; and I was convinced in a short Time, that instead of too much, too little Blood had been taken away in the Beginning; having been sometimes misled by the insidious Intervals of the Disease: at others having trusted too much to the faint Attempts which Nature

CHAP.  
VI.  
~

\* Ballon. Epid. Sparsim. BIANCH. Hist. Hep. p. iii. § viii. &c. BAGL. Prax. Med. l. i. c. ix. LANCIS. Epid. Rom. c. vi.

† O homines reipublicæ calamitosos atque funestos! ipsam pleuritidem, quæ sua sponte nullius operis indigens cum tali spūto quiesceret, ex eventu reddunt mortiferam.

Duret. in Prænot. Coac.



CHAP.  
VI.

made to relieve herself by Expectoration and Urine; the latter often becoming crude on the fourth Day, as the Delirium advanced, though it had promised fair on the second or third; the former frequently being checked about that Period of the Disease, by the immoderate Heat of the Lungs rendering the Matter viscid, globular, and not to be discharged, but with the utmost Difficulty.

I then began to bleed more plentifully, and repeated it so as to take away thirty or forty Ounces within the three first Days of the Distemper; and endeavoured, by bathing the Legs and blistering of them on the third Day, to prevent the fatal Symptoms from coming on about the fourth or fifth; giving Nitre at the same Time liberally, and Camphire in small Doses, to promote the thinner Secretions. This Method succeeded well in several Cases; Expectoration and Urine being thereby increased: but if they were not

increased,



increased, the Commotions which arose on the fourth, fifth, or sixth Day, made it always necessary to have Recourse to Bleeding again, and more Blisters, in order to relieve the Oppression of the Head and Breast; and though for the most Part the Patients did escape, yet they recovered with Difficulty, and continued coughing, spitting, and sweating in the Nights, for several Weeks.

CHAP.  
VI.  
~

At last, about the middle of *March*, when the Disease raged with the utmost Fury, having found there was an absolute Necessity for bleeding largely without Delay, in order to preserve Life, I began to put in Practice the following Method of Cure, which seldom or never failed, not only in young robust People, but even in those of a more advanced Age, provided I saw the Sick before the End of the third Day.

If I was called, for Example, in the Morning, - the Patient was immediately

T 4

laid



CHAP.  
VI.

laid in a horizontal Position, and bled at the Arm until his Pains abated, or he began to faint; neither of which commonly happened before  $\bar{z}$ xvi,  $\bar{z}$ xx, or  $\bar{z}$ xxiv. were taken away. If the Symptoms continued, I ordered about the same Quantity to be taken from the other Arm in the Afternoon, without regarding the Urine, Expectoration, or Appearance of the Blood. Next Morning, though there might be a great Alteration for the better; yet, if there was the least room to suspect, that any Obstruction remained in the Head or Breast, the Bleeding was repeated: And by carefully weighing the Blood\*, I found, that between forty-eight and fifty-four Ounces were frequently taken away during the first four-and-twenty Hours of my Attendance. This sudden copious Evacuation, commonly procured a Cessation of all violent Symptoms, and afforded

\* In weighing the Blood, I used the Island Weights, fourteen Ounces of which answer nearly to sixteen Ounces Averdupois.



an Opportunity to give an Antiphlogistick Purge the next Day. But if the Symptoms did not cease; or if the Pains and Difficulty of Breathing returned the Day after the Purge had been given; or if there was Reason to suspect from the Head-ach, Giddiness, Tingling of the Ears, and disturbed Rest, that the Brain was in Danger of being affected; I had again immediate Recourse to Bleeding, taking away at different Times to the Amount of  $\bar{z}$  xii,  $\bar{z}$  xviii, or  $\bar{z}$  xxiv, in the Space of a Day, either by the Lancet or cupping Glasses, or both, as occasion required; by which Means the impending Storm was happily averted; and as soon as the Commotions were quelled, the Purgative was repeated every other Day for three Times, unless some of the critical Evacuations appeared with such visible good Effects, as rendered it unnecessary.

CHAP.


VI.



In this Manner, I found with SYDENHAM, that Pleurisies of the most fatal

Ten-



CHAP.  
VI.

Tendency, might be happily cured in the Space of a few Days; and with as much Certainty as any Distemper whatever. And it was no less remarkable to observe, how quickly the Sick recovered their usual Health and Strength, notwithstanding the great Loss of Blood they had sustained; while many who had been bled more sparingly, continued in a languid, infirm State for Months, without being able to get rid of the Cough and Pains in the Breast.

Hitherto I have only mentioned the capital Remedies made use of in the Cure of these Distempers. But it may not be improper to take Notice of some other collateral Helps, which were generally used in the several Ways of treating them above described.


In the first Place, the Sick had warm Barley-Water with Oxymel for common Drink; and nitrous antiphlogistick Medicines



cines in the first Days of the Distemper; their Bellies being kept open with Clysters.

CHAP.

VI.




Oily Linctuses were exceeding useful in allaying the Cough; and small Doses of Anodynes were often necessary, both for that Purpose, and to procure Sleep when the Height of the Distemper was over.

If the Pleurisy began with a bilious Vomiting, it was expedient to promote that Evacuation by large Draughts of warm Water, in order to carry it off the sooner.

To ease the Pains in the Breast, the large leaves of the *Opuntia*, toasted in an Oven, and split through the Middle, were frequently applied: These being thick and succulent, retain the Heat a long Time, and produce all the good Effects that attend anodyne emollient Cataplasms

and



CHAP.  
VI.

and Fomentations; as I have frequently experienced in Tertian Fevers, Dysenteries, and other Diseases with inflamed Bowels, as well as in this Disease, since I first learned the Virtues of these Leaves from the Natives of *Minorca*.

After Bleeding twice or thrice, Blisters laid upon the Part affected were often successful in removing obstinate Stitches: But nothing affords such certain and immediate Relief, as the Scarificator and Cupping Glass; insomuch that among the many Trials I have had of them, I do not remember an Instance where they did not either carry off the Complaint entirely, or mitigate it considerably: But my Glasses were both wider and deeper than those commonly used in *England*.

The Phrensies and Anginas, which now and then appeared during this Constitution, required the same Method of Cure: And as to the Chin-cough, which  
was



was so fatal to Children, the chief Difference between it and the Pleurisy, seems to have been, that in the one the morbifick Matter was thrown into the Vesicles of the *Trachæa*; in the other, it stagnated in the Extremities of the pulmonary Arteries.

CHAP.  
IV.

During the same Period of Time, catarrhal Fevers appeared among the Adults, attended with Coughs, Pains in the Head and Bones, and frequently a *Delirium*; some of which suddenly terminated on the seventh Day by profuse Sweats; but most of them had partial *Crises* sooner, and went off by Degrees. When large Bleeding was neglected in the Beginning, they were liable to degenerate into Pleurisy. This was likewise the Case with the Tertian Fevers; and indeed it was remarkable, that not only all acute Diseases of this Season, but even accidental Hurts and Bruises required more plentiful Evacuations than ordinary; so inclinable was this

Con-



CHAP.  
VI.

Constitution of the Air to promote Inflammations.

I shall conclude this Chapter with a Paragraph or two from some Letters, occasioned by the Pleurisies of which we have been speaking, in order to confirm what has been said on them by the Testimony of my Friend Dr. FONT, an eminent Physician of *Ciudadella* in *Minorca*, who has deservedly enjoyed a most extensive Practice for upwards of thirty Years.

*Didaco FONT, M. D. G. C. S.*

—Novus annus funera densa produxit,  
grassante febre inflammatoriâ, quæ caput  
et organa respirationis potissimum afficit,  
modo pleuritis; modo peripneumonia, in-  
terdum phrenitis, interdum paraphrenitis  
adpellanda. Hic morbus, ut vehemens,  
largis et repetitis venæsectionibus, cum  
interpositâ catharsi, in herba felicissime  
jugulatur.

Complures liberale sputum,  
fine



fine multa sanguinis jacturâ, periculo eripuit. Nec defunt, quibus urinæ purulentæ Fluxus diu perseverans, salutis fuit. At, nisi istius modi auxilia mature accedant, væ ægris! nam postquam morbus per triduum impune sævit, altasque egit radices, sero plerumque medicina paratur; et neque venæsectiones, neque epispastica, neque cucurbitulæ, neque pectoralia tantopere decantata, impedire valent, quo minus, juxta Hippocratis effatum, "*Septimo die vel celerius succumbant, aut mente læsi, aut Orthopnæâ suffocati.*"

CHAP.  
VI.  
~

A te itaque peto, vir experientissime, ut dato otio, his quæsitis responsum præbeas.

Annon pleuritis morbus anniversarius in hâc insulâ et quibus mensibus?

An semper febrem continuam, cum celeri pulsu, acri calore, &c. comitem sibi adsciscat? An potius febri peridicâ remittente conjungatur.

Nonne



CHAP.  
VI.

Nonne tutius est, morbi resolutionem per venæsectiones, quam primum tentare, quam Coctionem et crisin naturæ præstolari?

Ubi resolutio tentanda est, quâ mensurâ et quibus intervalis, cruorem detrahere conveniat?

Datum Magone tertio die Maii, S. V.

Anno M,DCC,XLVI.

G. C. Diadacus FONT, S.

—Anniversarius hac in parte insulæ morbus est pleuritis, hiemeque viget, quantum ex usu observavi; et ubi rigida hyems plus justo prolongatur ad medium aut finem veris excurrere solet; immo anno 1730, quamvis solito modo procederent tempestates, memini hunc morbum, cum aliis inflammationibus internis, magnam stragem fecisse; ægris vel mente læsis, vel orthopnæâ suffocatis.

Pleuritis



Pleuritis non semper febrem acutam continuam habet adjunctam cum siti calore et celeri pulsu: Imo potius in ipsius principio, febris est fere semper mitis; pulsus quoad celeritatem, parum distat a naturali, calorque non est nimius, nec mordax; et in multis propensionem ad somnum observavi. Nequaquam vero, licet ita ingrediatur, leniter tractanda est; sed eodem modo ac si cum vehementi febre invasisset.

Præterea, pleuritis sæpe sociatur febre quæ periodice remittit, et intenditur, modo quotidie, modo de tertio in tertium\*. Et aliquando sola febris tertiana intermittens adest; sicuti observavi in quatuor ægris, quorum unus in initio septimi paroxysmi e vita decessit.

Quoad curationem; ubi tempestive vocatus sum, primo scil. die, vel secundi initio,

Vide GALEN. Avicen. & Mercurial apud BIANCH. p. iii. Schol. 25 & Spigel. de semitertian, cap. iii.



CHAP.  
VI.

depositâ morâ, sanguinis circiter quatuor libras, partitis vicibus, nuchthemer ispatio, si æger robustus sit, detrahere, jubeo; quâ subitâ, et copiosâ evacuatione, morbus quandoque in herbâ resolvitur. Sin vero persistat, ante diem quartum, totidem fere sanguinis libras, partitis vicibus noviter extraho: Et similiter prosequor, licet ante diem quartum, magna et notabilis omnium symptomatum remissio contingat. Quæ quidem remissio, talis et tanta esse solet, ut æger et reliqui rem jam in tuto putent; et ipse olim ita existimavi, donec infausti eventus aliter docuissent. Quippe veniente die quarto, aut quinto, supra modum exacerbantur symptomata et furente novâ procella, æger qui mox convalesceret sperabatur, orco traditur: Quod nullâ aliâ methodo quam supra præscriptâ præcavere potui; diluentibus interim, expectorantibus, clysteribus, cucurbitalis et epispasticis, pro ratione symptomatum, diligenter adhibitis; ut et purgantibus, post septimum diem. En methodus medendi, quæ meis

et



et ægrorum votis respondet! Ex centum qui corripuntur, nonaginta, aut in initio resolutione, aut postea manifestâ crisi servantur. Quæ crisis, aut per sudorum, aut per expectorationem, aut per urinæ profluvium, modo sanguinolentæ, modo puriformis, aut per diarrhœam perficitur; adjunctis vitæ viribus, sanguinis evacuatione, et supra memoratis remediis legitime exhibitis. Quibus inter initia spretis, Mors ut plurimum, succedit.


CHAP.  
VI.



Datum Civitellæ Maii 26. S. N.

ANNO MDCCLXVI.





## C H A P. VII.

*Of the S M A L L - P O X.*

CHAP.  
VII.


**T**HE Small-Pox were twice epidemical in *Minorca*, while I resided there, *viz.* in the Year 1742, and 1746. Without entering into a minute Description of the Disease, I shall only endeavour to give a general View of its Nature and Effects, in each of these Years; from which it will appear how widely the Infection of the one differed from that of the other.

About the middle of *March* 1742, the Small-Pox broke out in *Mabon*, to the  
great



great Consternation of the Natives, who had not seen them since the Year 1725, but well remembered the Destruction which they then occasioned. The Contagion quickly spread about, and was soon conveyed to the other Towns and Villages; in-somuch that before the End of *April* it prevailed in every part of the Island. As it attacked almost all who were under Seventeen, and many of a more advanced Age, the Sick were so numerous during *May* and *June*, that every House might be called an Hospital. About the End of *July* the Disease suddenly disappeared, most of those who were susceptible of it, having by that Time undergone it.

CHAP.  
VII.



During the first six or eight Weeks, the Distemper was favourable, and seldom proved fatal: But its Virulence increased with the Heat of the Weather; so that in *June* and *July* it was not uncommon, both at *Mabon* and *Ciudadella*, to bury ten or twelve of a Day. Nevertheless,



CHAP. in Proportion to the Numbers, not many  
VII. died; and what Mortality there was, hap-  
pened chiefly among Children at the Breast  
and the common Soldiers.

Among such Multitudes of Sick, all the different Species of Small-Pox, described by Authors, were sometimes met with; But both the distinct and confluent had a nearer resemblance to what SYDENHAM calls the anomalous, than to the regular Small-Pox. For though the fluxed Kind commonly appeared on the second or third Day, it was often the fourteenth or seventeenth before the Roughness to the Touch shewed, that those on the Face began to turn; and on the Legs and Arms they frequently continued fresh till near the thirtieth. In these Kinds of Small-Pox, the secondary Fever raged violently between the fourteenth and twenty-fourth Day; and almost all who perished by the Disease died on one or other of the intermediate Days.


The




The Constancy of Nature in promoting one or other of the common Evacuations, whereby Part of the morbid Matter was discharged, contributed to save great Numbers; for a plentiful Salivation commonly appeared with the Eruption, and continued till the Suppuration was compleated, not only in Adults, but in those of all Ages; even in Children at the Breast, whose want of Sense to spit freely, was probably the chief Cause of their suffering more than others. About the seventh Day or sooner, the Swelling of the Face came on; and it was regularly succeeded by a Tumefaction of the Hands, and sometimes of the Feet. But it was most remarkable in this epidemical Small-Pox, that a moderate Looseness commonly appeared, as the Pustules began to dry, and continued till the Skin was in some Measure cleared of the thick black Crust that covered it; by which Means, the Defect of Perspiration was supplied, the Symptoms

CHAP.

VII.





CHAP.  
VII.

of the secondary Fever were restrained, and many People were happily rescued from the Jaws of Death: From whence we may learn, how reasonable it is to give Purgatives in this Stage of the Disease, according to the Rules laid down for that Purpose, by Drs. FRIEND and MEAD.


In *December*, 1745, the Small-Pox were brought from *Constantinople* by one of his Majesty's Ships; and the following Year made a slow but fatal Progress over the whole Island. During the Spring 1746, they confined themselves to *St. Philip's Castle*, without reaching *Mabon*, though they are but two small Miles asunder. In the Summer and Autumn they were frequent in *Mabon* and the neighbouring Parts. Then they travelled Northward to *Ciudadella*, and disappeared in the Spring; having carried off almost all the Children who survived the Chin-cough and Summer Fevers of the Year 1746. It

was



was, however, very remarkable, that the longer the Infection continued in the Island, it became the milder; so that there was much less Mortality in the Northern Parts than in the Southern, where it first broke out.

CHAP.  
VII.



Whilst the Distemper raged at *St. Philip's Castle*, as I lived at *Mabon*, I saw but few of the Sick; but I was informed by the Physicians who attended them, that the Pustules were commonly of the confluent Kind, and often intermixed with Purples; that they seldom rose or filled well, but continued either hard like Tubercles, or quite empty, or had a small Quantity of *Ichor*, with a black Spot in their Middle, and frequently seemed to wither before they were ripe; that the Fever, instead of going off with the Eruption, increased as the Disease advanced, being generally accompanied with a *Coma*, *Delirium*, Difficulty of Breathing, incessant Vomiting and Loathing of Food; that



CHAP.  
VII.


that the Face seldom or never swelled, but about the Time that this ought to have happened, a Soreness of the Mouth or Throat gave great Uneasiness, the Skin separated from the Inside of the Lips, and the Breath became very fetid; that three fourths of the Infected, in spite of all that could be contrived for their Preservation, perished between the sixth and the fourteenth Days of the Fever. And the most of those who survived, remained blind, consumptive, or lame with *Caries* of the Bones, fordid Ulcers, &c. so that this Disease approached the nearest to the Plague of any that had been known in the Island.

From these Accounts (which were too well confirmed by my own Experience afterwards) I conjectured, that the extraordinary Mortality of the Disease was owing, partly to the variolous Matter abounding so much in the Blood, that the Skin was not capable of receiving the  
whole



whole Quantity; partly to the peculiar Disposition of the Air at that Time, which, as we have seen in the former Chapter, rendered the Head and Breast extremely liable to Inflammations. Hence, I imagined, that many of the Sick died of a *Phrenitis* or *Peripneumonia*, on the sixth, seventh, or eighth Days, before the Pustules had time to ripen; while in others, the Pustules having terminated in a Gangrene, the corrosive *Ichor* received into the Blood, proved fatal about the latter End of the second Week. It appeared to me that the most probable Way of averting these Disasters would be, to make large Evacuations in the first Days of the Distemper, and to replenish the Vessels with mild antiputrescent Liquors; by which Means, either the Eruption would be prevented, or the Pustules would be fewer in Number; or at least disposed to suppurate rather than mortify. This Method of treating the Small-Pox is countenanced by BALLONIUS\*, and


CHAP.  
VII.



\* Ephem, & Epid. l. i.

strongly



CHAP.  
VII.

strongly recommended by BOERHAAVE\*, who probably took the first Hint from RHAZIS†; and if ever so bold a Practice could be justified, it was in such a pestilential Kind as this, which destroyed almost all who were left to Nature, or managed in the common Way. These Considerations, and the visible good Effects of Bleeding and Purging largely, in the predominant Pleurifies, gave just Cause to expect some Benefit from a similar Method in the Cure of the Small-Pox; which Apprehension was confirmed by the following Accident.

A young Man, about Twenty-six Years of Age, was seized on *Wednesday, May 21,*

\* Aph. 1393.

† Si antequam apperere incipiant variolæ, ægrum medicus inveniat, minuatur sanguinis multitudo— Venter autem si strictus fuerit, infusiones quotidie in potu fumantur, ex hoc enim aut omnino prohibebitur pustularum egressio, aut si quid egressum fuerit, parvum erit.


Vid. Op. Venet. de Febr. cap, xviii. p. 105.

between



between seven and eight o'Clock in the Morning, with a Coldness and Shivering, which was soon succeeded by common feverish Complaints, and a Pain under his left Breast, straitening Respiration. On *Thursday* Morning, when I first visited him, imagining the Case to be a Pleurisy, I ordered him to be treated accordingly. In Consequence of which, seventeen Ounces of Blood were taken away immediately; twenty Ounces more in the Afternoon, and fifteen on *Friday* Morning; his Belly was kept open by Glysters; his Drink was Barley-Water, to which Oxymell and Nitre were added; and sometimes a Decoction of Tamarinds; and Leaves of the *Opuntia* were frequently applied to his Side. On *Friday* Forenoon his Pains were easier: but he complained greatly of an Inclination to vomit, and after drinking warm Water, threw up a considerable Quantity of bilious Matter. In the Afternoon an Eruption was discovered all over his Body, which from  
the



CHAP.  
VII.  


the Appearance it had on his Face, and its being attended with a Ptyalism, I suspected it to be the confluent Small Pox; and Dr. SEGUI, an eminent Physician of *Mabon*, who was consulted on this Emergency, confirmed my Opinion. Our Patient was bled on Friday Evening thirteen Ounces; and afterwards drank plenty of warm Water, in order to carry off his Vomiting; a Glyster was likewise injected, his Head shaved, his whole Skin cleaned and washed. On *Saturday* Morning the Fever being moderate, and the Eruption thick all over him, he took a mild Purge, which procured six or seven Stools, and a Grain of Opium was given at Bed-time. On *Sunday* Morning, after a restless disturbed Night, his Spitting having ceased, he complained of his Throat being so sore that he could scarce swallow, and that the Stitch had moved from his left Breast, and fixed about the lower Ribs in the right Side. These Symptoms induced us to bleed him a fifth

Time,



CHAP.  
VII.

Time, to the Quantity of eleven Ounces, which were of a crimson Colour, as all the rest of his Blood had been, without any fizy Crust. In the Afternoon he spit considerably, slept sound, was free from Pain; and after a good Night's Rest, on *Monday* Morning his Purge was repeated, which gave six or seven Motions, without checking the Salivation. After this, all farther Evacuations were laid aside (excepting Glysters to keep the Belly open) the Pustules, which were small and numerous all over him, began to fill with good Matter, and every thing went on according to our Wish. On *Tuesday* Morning his Face swelled; but abated on *Thursday* Afternoon, the Pustules beginning to dry; while his right Foot swelled, and became painful, the Ptyalism still continuing. *Monday* Morning, *June 2*, his Skin being dry every where, he took Physick, which was repeated on the *Friday* following.



CHAP. lowing. He soon gained Strength, and  
 VII. still enjoys a good State of Health.

The next Patient I attended in this Distemper was a *Jew's* Daughter, about five Years of Age, to whom I was called on *Saturday* Morning, *May 24*, when the Pustules were just beginning to appear, she having been feverish since *Wednesday* Morning. She was thrice bled by my Order, in the Space of four-and-twenty Hours; four Ounces being taken away each Time, she was strictly kept to the cooling Regimen; Glysters were frequently injected; and she commonly took a Paregorick at Night. The Pustules were low, small, confluent, and attended with a considerable Salivation. Her face swelled on *Tuesday*, her Hands the Day following, and her Feet on *Friday*. On *Sunday, June 1*, the Pustules began to dry on her Face. On *Monday* she was purged, and gradually grew strong. This and the former

mer



mer Patient were the two first in *Mabon*, who at that Time recovered of the Small-Pox.


CHAP.  
VII.  
~

About the End of *May*, and the Beginning of *June*, three or four Adults were attacked with violent feverish Symptoms; and as they never had the Small-Pox, it was suspected they had caught the Infection. They were all treated in the same Manner with the Man, whose Case has been already related, and in six or seven Days got perfectly well without Signs of Eruption. Nor did any of them take the Small-Pox afterwards.

Elevated with this Success, I began to flatter myself, that I had now hit upon a Method of Cure which would generally answer: But it was not long before Experience convinced me of my Mistake.

A young man, about one-and-twenty, was seized at Noon, *Monday, June 16,*  
X with



CHAP.  
VII.

with feverish Symptoms, and a violent Pain in his Stomach, Loins, and left Breast. He lost fifteen Ounces of Blood on *Tuesday* Morning, seventeen Ounces the same Afternoon, and fifteen on *Wednesday* Morning: All this Day he was sick and squeamish, though he drank much warm Water, and vomited up Bile along with it. He passed an uneasy Night, with great Pain at his Stomach and in the Small of his Back, and on *Thursday* Morning took a mild Purgative, which operated several Times upwards, and procured six or seven Stools. On *Friday* Morning the Small-Pox began to appear in his Face; his Stomach was somewhat relieved, but the Pain in his Back continued as bad as ever; his Mouth and Throat were very sore; and I was informed he had been delirious all Night. He was again blooded to twelve Ounces, which was red and florid, like all the rest which had been taken. In the Afternoon, the Pain of his Back being severe, the part

was



was cupped and scarified, and a Glyster was injected, which brought away much hard stinking Excrement. Notwithstanding this, he got no Manner of Relief; the Raving, Vomiting, and Restlessness increased, the Pustules never filled, his Mouth was ulcerated, and his Breath became offensive to the Smell; on *Sunday* Morning he lay quite stupified and senseless; black Spots appeared in the Middle of the Pustules; and he expired that Day about Noon. Upon opening the Body nothing extraordinary was found, except a monstrous large Gall-Bladder full of thick, green Bile, and slight Signs of Inflammation, in the villous Coat of the Stomach, the small Guts, and *Pia Mater*.

A young Lad of fifteen or sixteen Years, felt some Difficulty of Breathing, and a Pain in his Back on Tuesday Noon, *June* 24. He concealed his being ill to *Thursday*, when he complained much of Heat, Thirst, and Head-ach, Sicknes at his



CHAP.  
VII.  
~

Stomach, and a painful Oppression on both Sides of the *Thorax*. Eleven Ounces of Blood were taken on *Thursday* Morning, eight Ounces that Afternoon, and ten next Morning; Glysters and diluent Drinks were used, &c. None of his Blood appeared fizy; and before the last ten Ounces were taken, something like the Small-Pox were seen on his Face. But the Eruption never advanced; and on *Saturday* Morning Purples came out all over him. He then began to use the Bark every two Hours, and took in all about six or seven Drachms. In the mean time he was very delirious, vomited often, and breathed with great Difficulty. On *Monday* Morning he was seized with a Hæmorrhage from the Nose, which continued till seven at Night, when he died with black Spots all over him.


Soon after this, a robust middle-aged Man fell sick, and was treated in the same Way; on the second and third Days of the

Disease



Disease he lost forty-nine Ounces of Blood; on the third Day the Eruption began to appear; on the fourth he bled some Spoonfuls from the Nose; and purple Spots broke out all over him. The Small-Pox never rose; he raved incessantly, though he was afterwards twice blooded, and had Blisters applied. He died on the eleventh Day.

CHAP.  
VII.



These unfortunate Cases made me lay aside large Evacuations for the future, and content myself with endeavouring to alleviate the Symptoms in the common Way, by moderate Bleeding, Blistering, Anodynes, Bark, and Cordials, according as they were indicated. But in spite of all my Attempts to cure them, more perished than recovered. Nor could I find that any other Practitioner had better Success, until Time had corrected the Malignity of the Disease. In short, this epidemical Small-Pox sufficiently verified the *English* Proverb, "That there is one Sort in which the Nurse cannot kill, and another which the



CHAP. “the Physician can never cure.” And  
 VII. since it is a Matter of mere chance, whether the best or the worst Kind is got in the natural Way, it is evident what great Honour is due to the Memory of those Gentlemen who first introduced the Practice of Inoculation into this Kingdom, where the Safety and Security of it has been confirmed by the Experience of thirty Years,

This is all I have to offer to the Publick with regard to some of the most destructive Distempers incident to Mankind, whose Nature and Effects my Situation afforded many opportunities of observing. I have related the bad as well as the good Success of my Endeavours to cure them, with that Fidelity which is required in Affairs that so nearly concern the Lives of our Fellow-Creatures; and if these Pages should be found serviceable to others, by pointing out what will prove hurtful or beneficial in similar Cases, I shall not only think the


Pains



Pains they cost me well bestowed, but  
esteem it a singular Happiness that Fortune  
put it in my power to contribute so far to  
the Welfare of Society.

CHAP.

VII.



*Hoc opus, hoc studium, parvi properemus et  
ampli,*

*Si patriæ volumus, si nobis vivere cari.*

H O R.

F I N I S.



OF THE

THEY COLLECTED AND WELL KNOWN TO ALL  
THEY ARE IN THE HANDS OF THE  
THEY ARE IN THE HANDS OF THE

THEY ARE IN THE HANDS OF THE  
THEY ARE IN THE HANDS OF THE

THEY ARE IN THE HANDS OF THE  
THEY ARE IN THE HANDS OF THE  
1100.

THEY ARE IN THE HANDS OF THE  
THEY ARE IN THE HANDS OF THE

THEY ARE IN THE HANDS OF THE  
THEY ARE IN THE HANDS OF THE  
F I N I S H

THEY ARE IN THE HANDS OF THE  
THEY ARE IN THE HANDS OF THE

THEY ARE IN THE HANDS OF THE  
THEY ARE IN THE HANDS OF THE

THEY ARE IN THE HANDS OF THE  
THEY ARE IN THE HANDS OF THE

THEY ARE IN THE HANDS OF THE  
THEY ARE IN THE HANDS OF THE

THEY ARE IN THE HANDS OF THE  
THEY ARE IN THE HANDS OF THE



