

Key to the course of mathematics / composed for the use of the Royal Military Academy, by Charles Hulton.

Contributors

Dowling, Daniel.
Hulton, Charles.
Royal Military Academy.

Publication/Creation

London : M. Iley, etc., 1818.

Persistent URL

<https://wellcomecollection.org/works/jmewvgk8>

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Supp. 59016/13

~~X~~ ~~X~~




X - X
y - 4.





Digitized by the Internet Archive
in 2016 with funding from
Wellcome Library

<https://archive.org/details/b28741912>



K E Y
TO
HUTTON'S COURSE
OF
Mathematics.

Entered at Stationers' Hall.

KEY

TO THE COURSE

OF

MATHEMATICS,

COMPOSED FOR

THE USE OF THE ROYAL MILITARY ACADEMY,

BY

CHARLES HUTTON, LL. D. F. R. S.

THE LATEST EDITION,

Enlarged and Corrected.

BY

DANIEL DOWLING,

PROFESSOR OF ASTRONOMY AND THE MATHEMATICS;
LECTURER ON NATURAL PHILOSOPHY;
AND MASTER OF THE CLASSICAL, COMMERCIAL, AND MILITARY
SCHOOL,

Mansion-House, Highgate.

LONDON:

PRINTED FOR MATTHEW ILEY, BOOKSELLER, SOMERSET-STREET, PORTMAN-SQUARE; E. KERBY, STAFFORD-STREET, BOND-STREET; SCATCHERD AND LETTERMAN, AVE-MARIA-LANE; BALDWIN, CRADOCK AND JOY, AND SHERWOOD, NEELY AND JONES, PATERNOSTER-ROW; J. DEIGHTON, AND J. NICHOLSON, CAMBRIDGE; AND MAY BE HAD OF ALL BOOKSELLERS IN TOWN AND COUNTRY.

1818.

TO

THE RIGHT HONORABLE

LORD ABERCROMBIE, BARRISTON

AT THE BAR

ONE OF HIS MAJESTY'S MOST HONORABLE JUSTICES

OF THE KING'S BENCH

MEMBER OF PARLIAMENT FOR THE UNIVERSITY

OF CAMBRIDGE

AND

DEPUTY

IN THE HOUSE OF COMMONS

FOR HIS MAJESTY'S UNIVERSITY OF CAMBRIDGE

AND OF RESPECT

FOR THE BARRISTERS AND DISTINGUISHED GENTLEMEN

IN THE HOUSE OF COMMONS

AT HIS MAJESTY'S MOST GRACIOUS

AND MUCH OBLIGED SERVANT

THE AUTHOR

LIST OF SUBSCRIBERS.

- HIS GRACE THE DUKE OF NORFOLK, Lower Brook Street.
HIS GRACE THE DUKE OF BUCCLEUCH, South Audley Street.
HER GRACE THE DUCHESS DOWAGER OF BUCCLEUCH,
Montagu House, Whitehall.
HIS GRACE THE DUKE OF MONTROSE, Grosvenor Square.
HER GRACE THE DUCHESS OF MONTROSE, Grosvenor Square.
THE RT. HON. THE EARL OF COURTOWN, 11, Hill Street.
THE RT. HON. LORD VISCOUNT SYDNEY, Wansford.
THE RT. HON. LORD VISCOUNT PALMERSTON, Secretary at
War.
THE RT. HON. LADY PETRE, Grosvenor Square.
THE RT. HON. LORD MONTAGU, Whitehall.
THE RT. HON. LADY MONTAGU, Ditto.
THE RT. HON. LORD DOUGLAS, Bruton Street.
THE RT. HON. LORD GRAHAM, Mynadoc Castle.
THE RT. HON. LADY CHARLOTTE FINCH HATTON, Wans-
ford.
THE RT. HON. LADY CAROLINE GRAHAM, Grosvenor Square.
THE RT. HON. LADY LUCY GRAHAM, Ditto.
THE RT. HON. LORD CLIVE, Grafton Street.
THE HON. WILLIAM TEMPLE, Stanhope Street.
THE HON. MISS TEMPLE, Great Stanhope Street, May Fair.
THE HON. CAPT. GEORGE DOUGLAS, R. N.
PRINCE CHARLES DE BROGLIE, Paris.
LE MARQUIS DE BONNEVALLE, Mortimer Street.
LE COMTE DE GUISE, Paris.
LE COMTE DESCAR, Paris.
LE COMTE DE LOPPINOT, Trinidad.
FINCH HATTON, ESQ. Wansford.
CAPT. SCOTT, R. N. Petersham.
WILSON LOWRY, ESQ. R. A. & F. R. S. Titchfield Street.
BOND, JOHN, D. D. Hanwell Paddock.
BURNEY, WILLIAM, LL. D. Naval Academy, Gosport.
BAKER, MR. WILLIAM, Englefield Green.
BARKER, RICHARD, ESQ. Tavistock Street, Bedford Square.

LIST OF SUBSCRIBERS

- LANE, REV. DR. ...
- LANE, JOHN, ESQ. ...
- LANE, REV. DR. JAMES ...
- LANE, MRS. ...
- LANE, T. ESQ. ...
- LANE, REV. MR. ...
- LANE, GEORGE, ESQ. ...
- LANE, REV. DR. ...
- LANE, JOSEPH, ESQ. ...
- LANE, CHRISTIAN, ESQ. ...
- LANE, THOMAS, ESQ. ...
- LANE, J. ESQ. ...
- LANE, REV. DR. ...
- LANE, REV. DR. ...
- LANE, REV. ...
- LANE, WILLIAM, ESQ. ...
- LANE, WILLIAM, ESQ. ...
- LANE, G. A. ESQ. ...
- LANE, W. ESQ. ...
- LANE, REV. DR. ...
- LANE, REV. ...
- LANE, JOHN, ESQ. ...
- LANE, GEORGE, ESQ. ...
- LANE, GEORGE, ESQ. ...
- LANE, GEORGE, ESQ. ...
- LANE, J. ESQ. ...
- LANE, H. ESQ. ...
- LANE, JAMES, ESQ. ...
- LANE, REV. ...
- LANE, REV. ...
- LANE, GEORGE, ESQ. ...
- LANE, REV. DR. ...
- LANE, VINCENT, ESQ. ...
- LANE, CHARLES, ESQ. ...
- LANE, MR. W. ...
- LANE, R. H. ESQ. ...

The names of several subscribers cannot be inserted

P R E F A C E.

HAVING several years ago introduced DR. HUTTON'S COURSE OF MATHEMATICS into the number of Class Books used in my own SEMINARY, it was convenience first prompted me to put in manuscript a KEY to the more useful parts of it.

After some time, and when Dr. Hutton had enlarged his work, I indeed substituted his Three Volumes, almost exclusively, for the excellent French Authors till then generally and deservedly in the hands of the Mathematical Pupils.

But though my manuscript was thus acquiring bulk, still I had no intention of submitting it to the Public, greatly as I was convinced of its value in the school: and it was only at the instigation of some friends, highly read in the Mathematics, that I was induced to complete the KEY.

I confess I had long been desirous of contributing to the advancement of a science I had sedulously cultivated from my earliest years; and, on mature deliberation, I thought I could in no way more effectually accomplish this object than by rendering Dr. Hutton's Treatise, if possible, of more extensive and general utility; being well aware that the want of a KEY to his com-

prehensive COURSE, had prevented many from deriving such essential benefit from it, as might otherwise have been expected.

Accordingly I took the opinion of several of the most eminent Mathematicians of the present day, who all, after perusal, so warmly extolled the solutions, and commended the plan, that I could be no longer in doubt what line to pursue.

It is on these grounds then I venture to lay my labours before an impartial public, in the hope they will be found useful, not only to every description of Students in the Mathematics, but also to Officers of the Army, Navy, and the Honourable Company's service, as well as to others who may wish to *renew* their acquaintance with so many elegant parts of Mathematical science as are to be found in the COURSE.

With regard to the arrangement of the KEY, it was necessary it should be the same as that pursued by Dr. Hutton in his Three Volumes; but the facility of finding any question or proposition is rendered greater than is often to be met with.

I have in general adhered, with scrupulous exactness, to the rules laid down for the Student; yet in some few instances I have deviated a little, where deviation was warrantable: and these cases more frequently occur, perhaps, in the solution of Examples by logarithms. Many theorems I have demonstrated in two or more ways, where it appeared to me necessary; and in *Analytical Plane*

Trigonometry there is given, with very few exceptions, an easy geometrical construction, as best calculated to be more generally serviceable.

It has likewise been deemed expedient, sometimes for illustration, and sometimes for the detection of error, to subjoin occasional notes and remarks. By these it will be seen, there are some imperfections in Dr. Hutton's text and Answers.

These imperfections, however, if the extent and multifariousness of the work be taken into consideration, are comparatively few; and, in alluding to them, I by no means wish to infer, that the KEY may be expected entirely free from error.

It has nevertheless been uniformly my aim throughout, to send it forth with as few blemishes as possible; and in this, I trust, I have succeeded nearly to my wishes. Should there, however, be discovered defects of which I am not at present aware, I shall highly esteem the favour of their being pointed out to me by letter, in order to correct them. I own *much* novelty is not to be looked for in a work of the present kind; though at the same time I think the reader will be gratified in some degree by often meeting it unexpectedly.

Here it may be allowed me to state, that I have spared no pains or expense to render the work beautiful in execution; and if the price at first sight be thought a little more than the Book might have been afforded for, the difficulty of

printing such matter, and the scanty number of impressions in the edition, will argue that I could have had no mercenary object in view by fixing the price at £1 .. 4s. the copy.

The geometrical figures and *style* of printing, it is presumed, will be some enhancement to the value; while the size of the volume and the quality of the paper will enable and entitle it to be bound uniform with the COURSE to which it is the KEY.

But it is not for myself to judge how far I have succeeded; since, unless the book shall be useful to the public, my labours will in a measure be incomplete.

Mansion-House, Highgate,
November 15, 1817.

CONTENTS.

ARITHMETIC.

	Page
<i>NUMERATION</i>	1
<i>Notation</i>	2
<i>Addition</i>	<i>ib.</i>
<i>Subtraction</i>	3
<i>Multiplication</i>	<i>ib.</i>
<i>Division</i>	8
<i>Reduction</i>	14
<i>Compound Addition</i>	16
<i>Compound Subtraction</i>	17
<i>Compound Multiplication</i>	19
<i>Compound Division</i>	22
<i>Rule of Three</i>	27
<i>Compound Proportion</i>	30
<i>Reduction of Vulgar Fractions</i>	31
<i>Addition of Vulgar Fractions</i>	37
<i>Subtraction of Vulgar Fractions</i>	38
<i>Multiplication of Vulgar Fractions</i> :	39
<i>Division of Vulgar Fractions</i>	40
<i>Rule of Three in Vulgar Fractions</i>	41
<i>Addition of Decimals</i>	42
<i>Subtraction of Decimals</i>	<i>ib.</i>
<i>Multiplication of Decimals</i>	<i>ib.</i>
<i>Division of Decimals</i>	44

	Page
<i>Reduction of Decimals</i>	45
<i>Duodecimals</i>	48
<i>Involution</i>	50
<i>Evolution</i>	<i>ib.</i>
<i>Arithmetical Proportion</i>	55
<i>Geometrical Progression</i>	56
<i>Single Fellowship</i>	57
<i>Double Fellowship</i>	59
<i>Simple Interest</i>	61
<i>Compound Interest</i>	63
<i>Alligation</i>	66
<i>Single Position</i>	68
<i>Double Position</i>	69
PRACTICAL QUESTIONS IN ARITHMETIC	72
<i>Logarithms</i>	81

ALGEBRA.

<i>Addition in Algebra</i>	82
<i>Subtraction in Algebra</i>	83
<i>Multiplication in Algebra</i>	84
<i>Division in Algebra</i>	85
<i>Algebraic Fractions</i>	88
<i>Involution</i>	94
<i>Evolution</i>	95
<i>Reduction of Surds</i>	98
<i>Addition of Surds</i>	101
<i>Subtraction of Surds</i>	102
<i>Multiplication of Surds</i>	103
<i>Division of Surds</i>	105
<i>Involution of Surds</i>	106
<i>Evolution of Surds</i>	107
<i>Infinite Series</i>	108
<i>Arithmetical Progression</i>	115
<i>Application of Arithmetical Progression</i>	<i>ib.</i>

	Page
<i>Geometrical Proportion</i>	118
<i>Simple Equations</i>	119
<i>Quadratic Equations</i>	137
<i>Cubic and Higher Equations</i>	148
<i>Application of Cardan's Rule</i>	152
<i>Algebra applied to Geometry</i>	153

PLANE TRIGONOMETRY.

<i>Theorems</i>	169
<i>Heights and Distances</i>	177

MENSURATION.

<i>Mensuration of Planes</i>	193
<i>Mensuration of Solids</i>	201
<i>Artificers' Work</i>	206

NATURAL PHILOSOPHY.

<i>Laws of Gravity</i>	216
<i>Practical Gunnery</i>	<i>ib.</i>
<i>Hydrostatics</i>	222
<i>Pneumatics</i>	224

PRACTICAL QUESTIONS IN MENSURATION	227
. IN SPECIFIC GRAVITY	250
BALLS AND SHELLS	252
DISTANCE ESTIMATED BY SOUND	257
PRACTICAL QUESTIONS IN PHILOSOPHY	258

FLUXIONS.

<i>The Direct Method of Fluxions</i>	275
<i>The Inverse Method of Fluxions</i>	282

	Page
<i>Tabular Fluents</i>	309
<i>Maxima et Minima</i>	296
<i>Tangents</i>	302
<i>Length of Curve Lines</i>	303
<i>Quadratures</i>	304
<i>The Surfaces of Solids</i>	307
<i>The Solidity of Bodies</i>	309
<i>Logarithms</i>	310
<i>Points of Inflexion</i>	312
<i>Radius of Curvature</i>	<i>ib.</i>
<i>Involute and Evolute Curves</i>	314
<hr/>	
<i>Isoperimetry</i>	317
<i>Analytical Plane Trigonometry</i>	321
<i>Resolution of Spherical Triangles</i>	328
<i>Geodesic Operations</i>	336
<i>Polygonometry</i>	345
<i>Division of Surfaces</i>	353
<i>Solution of Equations</i>	362
<i>Equations to Curves</i>	370
<i>Loci of Equations</i>	371
<i>Fluxions and Fluents</i>	375
<i>Lateral Pressure of Fluids</i>	<i>ib.</i>
<i>DOWLING'S Rule for the Pressure on Dam-gates</i>	376

ERRATA.

- Page 165, last line, AC should be BC.
 Page 172, last line, -64° should be $=64^\circ$.
 Page 190, 21st line, $71^\circ .. 16'$ should be $37^\circ .. 28'$.
 Page 261, 25th line, "solid" should be "the solid."
 Page 271, 2nd line from the bottom, $\overline{32\frac{1}{2}}$ should be $\overline{32\frac{1}{4}}$.
-

K E Y

TO

HUTTON'S COURSE OF MATHEMATICS.

NUMERATION.

(Key to Vol. I. page 6.)

THIRTY-FOUR.

Ninety-six.

Three hundred and eighty.

Seven hundred and four.

Six thousand, one hundred and thirty-four.

Nine thousand and twenty-eight.

Fifteen thousand and eighty.

Seventy-two thousand and three.

One hundred and nine thousand, and twenty-six.

Four hundred and eighty-three thousand, five hundred.

Two millions, five hundred thousand, six hundred, and thirty-nine.

Seven millions, and five hundred and twenty-three thousand.

Thirteen mill. four hundred and five thous. six hundred and seventy.

Forty-seven mill. fifty thous. and twenty-three.

Three hundred and nine mill. twenty-five thous. six hundred.

Four thous. seven hund. and twenty-three mill. five hund. and seven thous. six hund. and eighty-nine.

Two hund. seventy-four thous. eight hund. and fifty-six mill. three hundred and ninety thousand.

Six billions, five hund. seventy-eight thous. six hund. millions, three hund. and seven thous. and twenty-four.

(Key to Vol. I. page 6.)

NOTATION.

57
 286
 9210
 27594
 640481
 3260106
 408255192
 27008096204
 200540110016
 21000810064150

(Page 11.)

ADDITION.

Ex. 5. 3426
 9024
 5106
 8890
 1204
 Ans. 27650

Ex. 6. 509267
 235809
 72920
 8392
 420
 21
 9
 Ans. 826838

Ex. 7. 2
 19
 817
 4298
 50916
 730205
 9180634
 Ans. 9966891

Qu. 8. Jan. 31	Quest. 9. Apr. 16	Quest. 10. 52714	Inf.
Feb. 28	May 31	5110	Cav.
Mar. 31	Jun. 30	6250	Drag.
Apr. 30	July 31	3927	Lt. H.
May 31	Aug. 31	928	Art.
Jun. 30	Sep. 30	1410	Pion.
July 31	Oct. 31	250	Sapp.
Aug. 31	Nov. 24	406	Min.
Sep. 30			
Oct. 31	Ans. 224 Days.	Ans. 70995	Total.
Nov. 30			
Dec. 31			

Ans. 365 Days.

(Key to Vol. I. page 12.)

SUBTRACTION.

Ex. 4. From 5331806
Take 5073918

Rem. 257888

Proof 5331806

Ex. 5. From 7020974
Take 2766809

Rem. 4254165

Proof 7020974

Ex. 6. From 8503402
Take 574271

Rem. 7929131

Proof 8503402

Ques. 7. 1727
1642

Ans. 85 Years.

Ques. 8. 2548
1815

Ans. 733 Years.

Ques. 9. 4000
1656

Ans. 2344 Years.

Ques. 10. 1815
1150

Ans. 665 Years.

Ques. 11. 1441
1330

Ans. 111 Years.

Ques. 12. 1492
1302

Ans. 190 Years.

(Page 16.)

MULTIPLICATION.

Ex. 1. Mul. 123456789
by 3

370370367 Prod.

Ex. 2. Mul. 123456789
by 4

493827156 Prod.

Ex. 3. Mul. 123456789
by 5

617283945 Prod.

Ex. 4. Mul. 123456789
by 6

740740734 Prod.

(Key to Vol. I. page 16.)

<p>Ex. 5. Mul. 123456789 by 7</p> <hr style="width: 100%;"/> <p>864197523 Prod.</p> <hr style="width: 100%;"/>	<p>Ex. 6. Mul. 123456789 by 8</p> <hr style="width: 100%;"/> <p>987654312 Prod.</p> <hr style="width: 100%;"/>
--	--

<p>Ex. 7. Mul. 123456789 by 9</p> <hr style="width: 100%;"/> <p>1111111101 Prod.</p> <hr style="width: 100%;"/>	<p>Ex. 8. Mul. 123456789 by 11</p> <hr style="width: 100%;"/> <p>1358024679 Prod.</p> <hr style="width: 100%;"/>
---	--

Ex. 9. Mul. 123456789
by 12

1481481468 Product.

<p>Ex. 10. Mul. 302914603 by 16</p> <hr style="width: 100%;"/> <p>1817487618 302914603</p> <hr style="width: 100%;"/> <p>4846633648 Prod.</p> <hr style="width: 100%;"/>	<p>Ex. 11. Mul. 273580961 by 23</p> <hr style="width: 100%;"/> <p>820742883 547161922</p> <hr style="width: 100%;"/> <p>6292362103 Prod.</p> <hr style="width: 100%;"/>
--	---

<p>Ex. 12. Mul. 402097316 by 195</p> <hr style="width: 100%;"/> <p>2010486580 3618875844 402097316</p> <hr style="width: 100%;"/> <p>78408976620 Prod.</p> <hr style="width: 100%;"/>	<p>Ex. 13. Mul. 82164973 by 3027</p> <hr style="width: 100%;"/> <p>575154811 164329946 2464949190</p> <hr style="width: 100%;"/> <p>248713373271 Prod.</p> <hr style="width: 100%;"/>
---	---

Ex. 14. Mul. 7564900
by 579

68084100
52954300
37824500

4380077100 Prod.

(Key to Vol. I. page 16.)

Ex. 15. Mul. 8496427
 by 874359

76467843
 42482135
 25489281
 33985708
 59474989
 67971416

7428927415293 Prod.

Ex. 16. Mul. 2760325
 by 37072

5520650
 19322275
 193222750
 8280975

102330768400 Prod.

(Page 17.)

When there are Ciphers in the Factors.

Ex. 3. Multiply 81503600
 by 7030

2445108000 Product by 30
 5705252000 - - Product by 7000

572970308000 PRODUCT by 7030

Ex. 4. Multiply 9030100
 by 2100

903010000 Product by 100
 18060200 - - Product by 2000

18963210000 PRODUCT by 2100

Ex. 5. Multiply 8057069
 by 70050

402853450 Product by 50
 5639948300 - - Product by 70000

564397683450 PRODUCT by 70050

(Key to Vol. I. page 18.)

When the Multiplier is the Product of two or more Numbers in the Multiplication Table.

Ex. 2. Here the multiplier 36 is the product of 6 by 6; or, of 4 by 9; therefore (always preferring the product of two numbers to that of three or more) it is

$\begin{array}{r} 31704592 \text{ Multiplicand.} \\ 6 \text{ First multiplier.} \\ \hline 190227552 \text{ Prod. by 6.} \\ 6 \text{ Second multiplier.} \\ \hline 1141365312 \text{ Prod. by } 6 \times 6 = 36. \end{array}$	or,	$\begin{array}{r} 31704592 \text{ Multiplicand.} \\ 9 \text{ First multiplier.} \\ \hline 285341328 \text{ Prod. by 9.} \\ 4 \text{ Second multipl.} \\ \hline 1141365312 \text{ Prod. by } 4 \times 9 = 36. \end{array}$
--	-----	---

Ex. 3. Here the multiplier 72 is the product of 12 by 6; or, of 8 by 9; therefore

$\begin{array}{r} 29753804 \text{ Multiplicand.} \\ 12 \text{ First multiplier.} \\ \hline 357045648 \text{ Prod. by 12.} \\ 6 \text{ Second multipl.} \\ \hline 2142273888 \text{ Prod. by } 6 \times 12 = 72. \end{array}$	or,	$\begin{array}{r} 29753804 \text{ Multiplicand.} \\ 8 \text{ First multiplier.} \\ \hline 238030432 \text{ Prod. by 8.} \\ 9 \text{ Second multipl.} \\ \hline 2142273888 \text{ Prod. by } 9 \times 8 = 72. \end{array}$
--	-----	---

Ex. 4. The multiplier 96 being the product of 12 by 8, it follows that

$\begin{array}{r} 7128368 \text{ Multiplicand.} \\ \text{by } 12 \\ \hline 85540416 \text{ Prod. by 12.} \\ \text{and by } 8 \\ \hline 684323328 \text{ Prod. by } 8 \times 12 = 96, \text{ which was required.} \end{array}$

Ex. 5. In this example the multiplier 108 is the product of 12 by 9, therefore

$\begin{array}{r} 160430800 \text{ Multiplicand.} \\ 12 \text{ First multiplier.} \\ \hline 1925169600 \text{ Prod. by 12.} \\ 9 \text{ Second multiplier.} \\ \hline 17326526400 \text{ Prod. by } 9 \times 12 = 108. \end{array}$

(Key to Vol. I. page 18.)

Ex. 6. Here the multiplier 1320 is the continued product of 10, 11, and 12; therefore,

$$\begin{array}{r}
 61835720 \text{ Multiplicand.} \\
 11 \text{ First multiplier.} \\
 \hline
 680192920 \text{ Prod. by 11.} \\
 10 \text{ Second multiplier.} \\
 \hline
 6801929200 \text{ Prod. by } 10 \times 11 = 110. \\
 12 \text{ Third multiplier.} \\
 \hline
 81623150400 \text{ Prod. by } 12 \times 10 \times 11 = 1320. \\
 \hline
 \end{array}$$

Quest. 7. In this question the multiplier 500 is the continued product of 10 by 10 by 5, wherefore, $104 \times 10 \times 10 \times 5 = 104 \times 500$.

$$\begin{array}{r}
 104 \text{ battalions.} \\
 10 \\
 \hline
 1040 \text{ Prod. by 10.} \\
 10 \\
 \hline
 10400 \text{ Prod. by } 10 \times 10 = 100. \\
 5 \\
 \hline
 \end{array}$$

Ans. 52000 men. *Prod. by* $5 \times 10 \times 10 = 500$.

Or, more concisely, annex two ciphers to the multiplicand, and multiply by 5. Thus,

104, with two ciphers annexed, is 10400 *Prod. by* 100.
 which \times^{ed} by 5

gives the *Product by* $500 = 52000$ men. Ans.

Quest. 8. Here, either 250 or 320 may be the multiplier; the former being the continued product of 5 by 5 by 10; and the latter of 4, 8, and 10. Therefore,

$$\begin{array}{r}
 320 \text{ waggons.} \\
 5 \\
 \hline
 1600 \text{ Prod. by 5.} \\
 5 \\
 \hline
 8000 \text{ Prod. by 25.} \\
 10 \\
 \hline
 \end{array}
 \qquad
 \text{or,}
 \qquad
 \begin{array}{r}
 250 \text{ loaves.} \\
 4 \\
 \hline
 1000 \text{ Prod. by 4.} \\
 8 \\
 \hline
 8000 \text{ Prod. by 32.} \\
 10 \\
 \hline
 \end{array}$$

Ans. 80000 loaves. *Prod. by* 250. Ans. 80000 loaves. *Prod. by* 320.

(Key to Vol. I. page 20.)

DIVISION.

Ex. 3.

4) 73146085 (18286521 $\frac{1}{4}$ Quotient.

$$\begin{array}{r}
 4 \overline{) 73146085} \\
 \underline{32} \\
 41 \\
 \underline{32} \\
 11 \\
 \underline{8} \\
 34 \\
 \underline{32} \\
 26 \\
 \underline{24} \\
 20 \\
 \underline{20} \\
 8 \\
 \underline{ 8} \\
 5 \\
 \underline{ 4} \\
 1 \text{ remainder.}
 \end{array}$$

Ex. 4.

7) 5317986027 (759712289 $\frac{4}{7}$ Quot.

$$\begin{array}{r}
 7 \overline{) 5317986027} \\
 \underline{49} \\
 41 \\
 \underline{35} \\
 67 \\
 \underline{63} \\
 49 \\
 \underline{49} \\
 8 \\
 \underline{ 7} \\
 16 \\
 \underline{14} \\
 20 \\
 \underline{14} \\
 62 \\
 \underline{56} \\
 67 \\
 \underline{63} \\
 4 \text{ remainder.}
 \end{array}$$

(Key to Vol. I. page 20.)

Ex. 7. $97 \overline{) 137896254} (1421610\frac{34}{97} \text{ Quotient.}$

97							
408	-						
388							
209	--						
194							
156	----						
97							
592	----						
582							
105	----						
97							
84 remainder.							
97 divisor.							

Ex. 8.

$$764 \overline{) 35821649} (46886\frac{745}{764} \text{ Quot.}$$

3056

5261

4584

6776

6112

6644

6612

5329

4584

745 remainder.

764 divisor.

Ex. 9.

$$5201 \overline{) 72091365} (13861\frac{304}{5201} \text{ Quo.}$$

5201

20081

15603

44783

41608

31756

31206

5505

5201

304 remainder.

5201 divisor.

(Key to Vol. I. page 20.)

Ex. 10. $57606 \overline{)4637064283(80496\frac{11707}{57606} \text{ Quotient.}}$
 $\underline{460848}$
 $\cdot 285842$
 $\underline{230424}$
 554188
 $\underline{518454}$
 357343
 $\underline{345636}$
 $\cdot 11707 \text{ remainder.}$
 $\underline{57606 \text{ divisor.}}$

Quest. 11.
 $3 \overline{)471(157 \text{ men. Ans.}}$
 $\underline{3}$
 17
 $\underline{15}$
 21
 $\underline{21}$
 $\cdot \cdot$

Quest. 12.
 $18 \overline{)378(21 \text{ miles. Ans.}}$
 $\underline{36}$
 18
 $\underline{18}$
 $\cdot \cdot$

Quest. 13. $365 \overline{)£38330(£105\frac{5}{365} \text{ Answer.}}$
 $\underline{365}$
 1830
 $\underline{1825}$
 $\cdot \cdot \cdot 5 \text{ remainder.}$
 $\underline{365 \text{ divisor.}}$

* This answer differs from that given in the Course.

(Key to Vol. I. page 21.)

CONTRACTIONS IN DIVISION.

Ex. 2.	4)52619675	Ex. 3.	5)1379192
	<u>13154918$\frac{3}{4}$ Quotient.</u>		<u>275838$\frac{2}{5}$ Quotient.</u>
Ex. 4.	6)38672940 Dividend.	Ex. 5.	7)81396627 Dividend.
	<u>6445490 Quotient.</u>		<u>11628089$\frac{4}{7}$ Quotient.</u>
Ex. 6.	8)23718920 Dividend.	Ex. 7.	9)43981962 Dividend.
	<u>2964865 Quotient.</u>		<u>4886884$\frac{6}{9}$ Quotient.</u>
Ex. 8.	11)57614230 Dividend.	Ex. 9.	12)27980373 Dividend.
	<u>5237657$\frac{3}{11}$ Quotient.</u>		<u>2331697$\frac{9}{12}$ Quotient.</u>

(Page 22.)

When Ciphers are annexed to the Divisor.

Ex. 3. 23000)7380964(320 $\frac{2}{3}$ $\frac{0}{3}$ $\frac{0}{8}$ $\frac{6}{8}$ $\frac{4}{8}$ Quotient.*

 ••• 69 •••

 —

 48

 —

 46

 —

 • 20964 remainder.

 —

 23000 divisor.

 —

* *A small typographical error in the Course.*

Ex. 4. 5800)2304109(397 $\frac{1}{5}$ $\frac{5}{8}$ $\frac{0}{8}$ $\frac{0}{8}$ Quotient.

 •• 174 ••

 —

 564

 —

 522

 —

 421

 —

 406

 —

 • 1509 remainder.

 —

 5800 divisor.

 —

(Key to Vol. I. page 23.)

When the Divisor is the Product of two or more Numbers in the Multiplication Table.

Ex. 2. Here the divisor 72 is the product of 8 by 9; or, of 12 by 6; therefore,

8)7014596 Dividend.

$$\begin{array}{r} \underline{\quad\quad\quad} \\ 9) 876824 \dots\dots\dots 4 \text{ first remainder} \\ \underline{\quad\quad\quad} \\ 97424 \dots\dots\dots 8 \text{ second remainder} \end{array} \left. \vphantom{\begin{array}{r} 9) 876824 \\ \underline{\quad\quad\quad} \\ 97424 \end{array}} \right\} = \frac{68}{72}$$

Hence $97424 \frac{68}{72}$ Quotient.

or,

12)7014596 Dividend.

$$\begin{array}{r} \underline{\quad\quad\quad} \\ 6) 584549 \dots\dots\dots 8 \text{ first remainder} \\ \underline{\quad\quad\quad} \\ 97424 \dots\dots\dots 5 \text{ second remainder} \end{array} \left. \vphantom{\begin{array}{r} 6) 584549 \\ \underline{\quad\quad\quad} \\ 97424 \end{array}} \right\} = \frac{68}{72}$$

Hence, as before, $97424 \frac{68}{72}$ Quotient.

Ex. 3. Here the divisor 132 is the product of 11 by 12; wherefore,

11)5130652 Dividend.

12) 466422.....10 first remainder.

38868.....6 second remainder.

Hence $38868 \frac{76}{132}$ Quotient.

Ex. 4. In this example, 240, the divisor, is the continued product of 10, 2, and 12, that is, of 20 and 12; for which reason,

20)83016572 Dividend.

12) 4150828.....12 first remainder.

345902.....4 second remainder.

Whence it is evident that $345902 \frac{92}{240}$ is the quotient required.

(Key to Vol. I. page 23.)

COMMON DIVISION, PERFORMED CONCISELY.

Ex. 2. $238)79165238(332627\frac{12}{238}$ Quotient.

$$\begin{array}{r}
 \cdot 776 \quad || \\
 \cdot 625 \quad || \\
 1492 \quad || \\
 \cdot \cdot 643 \quad | \\
 1678 \quad | \\
 \cdot \cdot 12 \text{ remainder.} \\
 \hline
 \end{array}$$

Ex. 3. $5317)29137062(5479\frac{5212}{5317}$ Quotient.

$$\begin{array}{r}
 \cdot 25520 \quad || \\
 \cdot 42526 \quad | \\
 \cdot 53072 \quad | \\
 \cdot 5219 \text{ remainder.} \\
 \hline
 \end{array}$$

Ex. 4. $7803)62015735(7947\frac{5294}{7803}$ Quotient.

$$\begin{array}{r}
 \cdot 73947 \quad || \\
 \cdot 37203 \quad | \\
 \cdot 59915 \quad | \\
 \cdot 5294 \text{ remainder.} \\
 \hline
 \end{array}$$

(Page 31.)

REDUCTION.

Ex. 3. £ 24
20

480 shillings.

12

5760 pence.

4

Ans. 23040 farthings.

Ex. 4. $4)337587$ farthings. $12)84396\frac{3}{4}$ pence. $20)7033$ shillings.

s. d.

Ans. £ 351 .. 13 .. $0\frac{3}{4}$

(Key to Vol. I. page 31.)

Ex. 5. 36 guineas.
 21

 756 shillings.
 12

 9072 pence.
 4

 Ans. 36288 farthings.

Ex. 6. 4) 36288 farthings.

 12) 9072 pence.

 21) 756 shillings.

 Ans. 36 guineas.

Ex. 7. lb. dwt. gr.
 59 .. 13 .. 5
 12

 708 oz.
 20

 14173 dwt.
 24

 Ans. 340157 grains.

Ex. 8. 24) 8012131 grains.

 20) dwt. 333838 19 gr.

 12) oz. 16691 18 dwt.

 lb. 1390 11 oz.

 lb. oz. dwt. gr.
 Ans. 1390 .. 11 .. 18 .. 19

Ex. 9. tons. cwt. qr. lb. oz. dr.
 35 .. 17 .. 1 .. 23 .. 7 .. 13
 20

 717 cwt.
 4

 2869 qrs.
 28

 80355 lb.
 16

 1285687 oz.
 16

 Ans. 20571005 drams.

Qu. 10. 25000 miles.
 8

 200000 furlongs.
 40

 8000000 poles.
 5½

 44000000 yds.
 3

 132000000 feet.
 12

 1584000000 inches.
 3

 Ans. 4752000000 b. corns.

(Key to Vol. I. page 31.)

days. hrs. min. sec.	days. hrs. min. sec.
Quest. 11. 365 .. 5 .. 48 .. 45 $\frac{1}{2}$	Ques. 12. 29 .. 12 .. 44 .. 3
24	24
-----	-----
8765 hours.	708 hours.
60	60
-----	-----
525948 minutes.	42524 minutes.
60	60
-----	-----
Ans. 31556925 $\frac{1}{2}$ seconds.	Ans. 2551443 seconds.
-----	-----

(Page 32.)

COMPOUND ADDITION.

Ex. 2. Ans. £ 59 18 5 $\frac{1}{4}$	Ex. 3. Ans. £ 79 7 1
Ex. 4. Ans. £ 174 2 4 $\frac{3}{4}$	Ex. 5. Ans. £ 210 5 11 $\frac{1}{4}$
Ex. 6. Ans. £ 170 3 1	Ex. 7. Ans. £ 141 11 3 $\frac{1}{4}$
Ex. 8. Ans. £ 930 4 11 $\frac{1}{4}$	

		s.	d.	
Quest. 9,	£ 197	13	7 $\frac{1}{2}$	Butcher's bill.
	59	5	2 $\frac{3}{4}$	Baker's bill.
	85	0	0	Brewer's bill.
	103	13	0	Wine Merchant's bill.
	75	0	3	Corn-chandler's bill.
	27	15	11 $\frac{1}{4}$	Cheese-monger's bill.
	55	3	5 $\frac{3}{4}$	Tailor's bill.
	127	3	0	Sundry charges.
	100	0	0	Pocket-money.
	£ 830	14	6 $\frac{1}{4}$	Sum Total.

(Page 35.)

Ex. 1. Ans. 235 .. 6 .. 15	Ex. 2. Ans. 81 .. 3 .. 21
lb 3 3 9	oz. dwt. gr.
Ex. 3. Ans. 79 .. 9 .. 6 .. 1	Ex. 4. Ans. 76 .. 2 .. 2 .. 15
	3 3 9 gr.

(Key to vol. I. page 35.)

Ex. 5.	Ans.	lb.	oz.	dr.	Ex. 6.	Ans.	cwt.	qr.	lb.
		70	.. 7	.. 7			54	.. 3	.. 22
Ex. 7.	Ans.	mil.	fur.	pol.	Ex. 8.	Ans.	yds.	ft.	in.
		71	.. 1	.. 36			234	.. 0	.. 11
Ex. 9.	Ans.	yds.	qr.	nl.	Ex. 10.	Ans.	En.ells.	qr.	nl.
		331	.. 3	.. 1			362	.. 1	.. 0
Ex. 11.	Ans.	A.	R.	P.	Ex. 12.	Ans.	A.	R.	P.
		303	.. 3	.. 34			402	.. 3	.. 11
Ex. 13.	Ans.	tuns.	hhd.	gall.	Ex. 14.	Ans.	hhds.	gall.	pts.
		137	.. 2	.. 51			86	.. 35	.. 0
Ex. 15.	Ans.	hhds.	gall.	pts.	Ex. 16.	Ans.	hhds.	gall.	pts.
		56	.. 13	.. 6			124	.. 53	.. 7

(Page 36.)

COMPOUND SUBTRACTION.

Ex. 3.	Ans.	£ 51	17	7 $\frac{3}{4}$	Ex. 4.	Ans.	£ 217	2	7 $\frac{1}{2}$
Quest. 5.		£ 73	0	5 $\frac{1}{4}$	minuend.				
		19	13	10	subtrahend.				
	Ans.	£ 53	6	7 $\frac{1}{4}$	remainder.				

(Page 37.)

Quest. 6.		£ 100	0	0	money lent by A.					
		73	12	4 $\frac{3}{4}$	in value repaid by B.					
	Ans.	£ 26	7	7 $\frac{1}{4}$	balance in favour of A.					
Quest. 7.	Land-tax	£ 0	14	6	}	£ 20	12	0	half-year's rent.	
	Repairs	1	3	3 $\frac{1}{4}$			1	17	9 $\frac{1}{4}$	disbursed.
						Ans.	£ 18	14	2 $\frac{3}{4}$	due.

(Key to Vol. I. page 37.)

Quest. 8.	Debts.	Effects.
A.	£ 35 7 6	£23 7 5 Cash.
B.	91 13 0 $\frac{1}{2}$	53 11 10 $\frac{1}{4}$ Wares.
C.	53 0 7 $\frac{1}{4}$	63 17 7 $\frac{3}{4}$ Household furn.
D.	87 5 0	25 7 5 Recov ^{ble} debts.
E.	111 3 5 $\frac{3}{4}$	
	<hr/>	
	£378 9 7 $\frac{1}{2}$	£166 4 4
	166 4 4	amount of debts.
	<hr/>	value of effects.
Ans.	£ 212 5 3 $\frac{1}{2}$	deficient.

EXAMPLES OF WEIGHTS AND MEASURES.

Ex. 1.	Ans. 3 lb. 10 oz. 5 dwt. 17 gr.	Ex. 2.	Ans. 4 lb. 2 oz. 8 dwt. 5 gr.
Ex. 3.	Ans. 43 lb. 3 oz. 3 dwt. 9 gr. 15	Ex. 4.	Ans. 2 cwt. 1 qr. 7 lb.
Ex. 5.	Ans. 53 lb. 11 oz. 7 dr.	Ex. 6.	Ans. 6 mil. 5 fur. 6 pol.
Ex. 7.	Ans. 23 yds. 0 ft. 7 in.	Ex. 8.	Ans. 8 yds. 1 qr. 3 nls.
Ex. 9.	Ans. 1 yd. 2 qr. 1 nl.	Ex. 10.	Ans. 0 A. 3 R. 6 P.
Ex. 11.	Ans. 34 A. 1 R. 27 P.		

(Page 38.)

Ex. 12.	Ans. 8 tuns. 0 hhds. 50 gal.	Ex. 13.	Ans. 2 hhds. 50 gal. 6 pts.
Ex. 14.	Ans. 4 hhds. 47 gal. 4 pts.	Ex. 15.	Ans. 52 hhds. 9 gal. 4 pts.
Ex. 16.	Ans. 3 lasts. 1 qr. 2 bu.	Ex. 17.	Ans. 4 B. 4 G. 2 P.
Ex. 18.	Ans. 54 mo. 0 w. 6 da.	Ex. 19.	Ans. 42 da. 6 hrs. 49 min.
Quest. 20.	236 yds. 0 ft. 0 in. line of defence.		
	146 yds. 1 ft. 4 in. part terminated by the shoulder and curtain.		
Ans.	<hr/> 89 yds. 1 ft. 8 in. length of the face required.		

(Key to Vol. I, page 39.)

COMPOUND MULTIPLICATION.

$$\begin{array}{r} \text{Ex. 2.} \quad \text{£ } 0 \quad 7 \quad 8 \\ \qquad \qquad \qquad 4 \text{ lb.} \\ \hline \text{£ } 1 \quad 10 \quad 8 \text{ Ans.} \\ \hline \end{array}$$

$$\begin{array}{r} \text{Ex. 3.} \quad \text{£ } 0 \quad 0 \quad 9\frac{1}{2} \\ \qquad \qquad \qquad 6 \text{ lb.} \\ \hline \text{£ } 0 \quad 4 \quad 9 \text{ Ans.} \\ \hline \end{array}$$

$$\begin{array}{r} \text{Ex. 4.} \quad \text{£ } 0 \quad 1 \quad 8\frac{1}{2} \\ \qquad \qquad \qquad 7 \text{ lb.} \\ \hline \text{£ } 0 \quad 11 \quad 11\frac{1}{2} \text{ Ans.} \\ \hline \end{array}$$

$$\begin{array}{r} \text{Ex. 5.} \quad \text{£ } 0 \quad 2 \quad 7\frac{1}{2} \\ \qquad \qquad \qquad 9 \text{ st.} \\ \hline \text{£ } 1 \quad 3 \quad 7\frac{1}{2} \text{ Ans.*} \\ \hline \end{array}$$

* Differing by the price of 1 stone of beef from the given Ans., £1. 1s. being the price of 8 stone.

$$\begin{array}{r} \text{Ex. 6.} \quad \text{£ } 2 \quad 17 \quad 10 \\ \qquad \qquad \qquad 10 \text{ cwt.} \\ \hline \text{£ } 28 \quad 18 \quad 4 \text{ Ans.} \\ \hline \end{array}$$

$$\begin{array}{r} \text{Ex. 7.} \quad \text{£ } 3 \quad 7 \quad 4 \\ \qquad \qquad \qquad 12 \text{ cwt.} \\ \hline \text{£ } 40 \quad 8 \quad 0 \text{ Ans.} \\ \hline \end{array}$$

CONTRACTIONS.

When the Multiplier exceeds 12.

$$\begin{array}{r} \text{Ex. 2.} \quad \text{£ } 4 \quad 7 \quad 2 \\ \qquad \qquad \qquad 10 \text{ cwt.} \\ \hline \text{£ } 43 \quad 11 \quad 8 \text{ pr. of 10 cwt.} \\ \qquad \qquad \qquad 2 \text{ cwt.} \\ \hline \text{Ans. } \text{£ } 87 \quad 3 \quad 4 \text{ pr. of 20 cwt.} \\ \hline \end{array}$$

$$\begin{array}{r} \text{Ex. 3.} \quad \text{£ } 3 \quad 7 \quad 6 \\ \qquad \qquad \qquad 6 \text{ tons.} \\ \hline \text{£ } 20 \quad 5 \quad 0 \text{ pr. of 6 tons.} \\ \qquad \qquad \qquad 4 \text{ tons.} \\ \hline \text{Ans. } \text{£ } 81 \quad 0 \quad 0 \text{ pr. of 24 tons.} \\ \hline \end{array}$$

$$\begin{array}{r} \text{Ex. 4.} \quad \text{£ } 0 \quad 1 \quad 6 \\ \qquad \qquad \qquad 9 \text{ ells.} \\ \hline \text{£ } 0 \quad 13 \quad 6 \text{ price of 9 ells.} \\ \qquad \qquad \qquad 5 \text{ ells.} \\ \hline \text{Ans. } \text{£ } 3 \quad 7 \quad 6 \text{ price of 45 ells.} \\ \hline \end{array}$$

(Key to Vol. I. page 40.)

<p>Ex. 5. $\begin{array}{r} \text{£}0\ 2\ 3 \\ \text{9 gallons.} \\ \hline \text{£}1\ 0\ 3 \text{ price of 9 gals.} \\ \text{7 gallons.} \\ \hline \text{Ans. } \text{£}7\ 1\ 9 \text{ pr. of 63 gals.} \end{array}$</p>	<p>Ex. 6. $\begin{array}{r} \text{£}1\ 4 \\ \text{10 barrels.} \\ \hline \text{£}12\ 0 \text{ pr. of 12 barrels.} \\ \text{7 barrels.} \\ \hline \text{Ans. } \text{£}84\ 0 \text{ pr. of 70 barrels.} \end{array}$</p>
---	--

<p>Ex. 7. $\begin{array}{r} \text{£}1\ 12\ 8 \\ \text{12 quarters.} \\ \hline \text{£}19\ 12\ 0 \text{ price of 12 qrs.} \\ \text{7 quarters.} \\ \hline \text{Ans. } \text{£}137\ 4\ 0 \text{ pr. of 84 qrs.} \end{array}$</p>	<p>Ex. 8. $\begin{array}{r} \text{£}1\ 3\ 4 \\ \text{12 quarters.} \\ \hline \text{£}14\ 0\ 0 \text{ price of 12 qr.} \\ \text{8 quarters.} \\ \hline \text{Ans. } \text{£}112\ 0\ 0 \text{ price of 96 qr.} \end{array}$</p>
--	--

<p>Ex. 9. $\begin{array}{r} \text{£}0\ 5\ 9 \\ \text{12 days.} \\ \hline \text{£}3\ 9\ 0 \text{ wages for 12 d}^s. \\ \text{10 days.} \\ \hline \text{Ans. } \text{£}34\ 10\ 0 \text{ wages for 120 d}^s. \end{array}$</p>	<p>Ex. 10. $\begin{array}{r} \text{£}0\ 13\ 4 \\ \text{12 reams.} \\ \hline \text{£}8\ 0\ 0 \text{ for 12 reams.} \\ \text{12 reams.} \\ \hline \text{Ans. } \text{£}96\ 0\ 0 \text{ for 144 reams.} \end{array}$</p>
---	--

When the Multiplier is not produced by the Multiplication of simple Numbers.

<p>Ex. 2. $\begin{array}{r} \text{£}2\ 5\ 3\frac{1}{4} \\ \text{4 quarters.} \\ \hline \text{£}9\ 1\ 1 \text{ for 4 qrs.} \\ \text{7 quarters.} \\ \hline \text{£}63\ 7\ 7 \text{ for 28 qrs.} \\ \text{Add } \text{£}2\ 5\ 3\frac{1}{4} \text{ for 1 qr.} \\ \hline \text{Ans. } \text{£}65\ 12\ 10\frac{1}{4} \text{ for 29 qrs.} \end{array}$</p>	<p>Ex. 3. $\begin{array}{r} \text{£}3\ 15\ 2 \\ \text{6 loads.} \\ \hline \text{£}22\ 11\ 0 \text{ for 6 lds.} \\ \text{9 loads.} \\ \hline \text{£}202\ 19\ 0 \text{ for 54 lds.} \\ \text{Subtract } \text{£}3\ 15\ 2 \text{ for 1 ld.} \\ \hline \text{Ans. } \text{£}199\ 3\ 10 \text{ for 53 lds.} \end{array}$</p>
---	---

(Key to Vol. I, page 42.)

COMPOUND DIVISION.

Ex. 2. $3)£432 \ 12 \ 1\frac{3}{4}$ Divd. Ex. 3. $4)£507 \ 3 \ 5$ Divd.

$$\begin{array}{r} \underline{\underline{£144 \ 4 \ 0\frac{1}{2}}} \text{ Quot.} \end{array}$$

$$\begin{array}{r} \underline{\underline{£126 \ 15 \ 10\frac{1}{4}}} \text{ Quot.} \end{array}$$

Ex. 4. $5)£632 \ 7 \ 6\frac{1}{2}$ Divd. Ex. 5. $6)£690 \ 14 \ 3\frac{1}{4}$ Divd.

$$\begin{array}{r} \underline{\underline{£126 \ 9 \ 6}} \text{ Quot.} \end{array}$$

$$\begin{array}{r} \underline{\underline{£115 \ 2 \ 4\frac{1}{2}}} \text{ Quot.} \end{array}$$

Ex. 6. $7)£705 \ 10 \ 2$ Divd. Ex. 7. $8)£760 \ 5 \ 6$ Divd.

$$\begin{array}{r} \underline{\underline{£100 \ 15 \ 8\frac{3}{4}}} \text{ Quot.} \end{array}$$

$$\begin{array}{r} \underline{\underline{£ \ 95 \ 0 \ 8\frac{1}{4}}} \text{ Quot.} \end{array}$$

Ex. 8. $9)£761 \ 5 \ 7\frac{3}{4}$ Divd. Ex. 9. $10)£829 \ 17 \ 10$ Divd.

$$\begin{array}{r} \underline{\underline{£ \ 84 \ 11 \ 8\frac{3}{4}}} \text{ Quot.} \end{array}$$

$$\begin{array}{r} \underline{\underline{£ \ 82 \ 19 \ 9\frac{1}{4}}} \text{ Quot.} \end{array}$$

Ex. 10. $11)£937 \ 8 \ 8\frac{3}{4}$ Dividend.

$$\begin{array}{r} \underline{\underline{£ \ 85 \ 4 \ 5 \ 0\frac{7}{11}}} \text{ Quotient.} \end{array}$$

Ex. 11. $12)£1145 \ 11 \ 4\frac{1}{4}$ Dividend.

$$\begin{array}{r} \underline{\underline{£ \ 95 \ 9 \ 3 \ 1\frac{5}{12}}} \text{ Quotient.} \end{array}$$

When the Divisor exceeds 12, and is the Product of two or more Numbers in the Multiplication Table.

Ex. 2.

$$2)£150 \ 6 \ 8 \text{ for 20 cwt.}$$

$$10)£ \ 75 \ 3 \ 4 \text{ for 10 cwt.}$$

$$\text{Ans. } \underline{\underline{£ \ 7 \ 10 \ 4}} \text{ for 1 cwt.}$$

Ex. 3.

$$6)£ \ 98 \ 8 \text{ Dividend.}$$

$$6)£ \ 16 \ 8 \text{ Quotient by 6.}$$

$$\text{Ans. } \underline{\underline{£ \ 2 \ 14 \ 8}} \text{ Quot. by 36.}$$

(Key to Vol. I. page 42.)

Ex. 4.

8) £71 13 10 Dividend.

7) £ 8 19 2 $\frac{3}{4}$ Quot. by 8.

Ans. £ 1 5 7 $\frac{1}{4}$ Quot. by 56.

Ex. 5.

8) £44 4 0 Dividend.

12) £ 5 10 6 Quot. by 8.

Ans. £ 0 9 2 $\frac{1}{2}$ Quot. by 96.

Quest. 6. 4) £31 10 0 per cwt.

7) £ 7 17 6 per qr.

4) £ 1 2 6 per 4 lb.

Ans. £ 0 5 7 $\frac{1}{2}$ per lb.

(Page 43.)

When the Divisor is not the Product of two or more Numbers in the Multiplication Table.

Ex. 2.

57) £39 14 5 $\frac{1}{4}$ (£0 13 11 $\frac{1}{4}$ Quotient.

20		
794		
224		
53		
12		
641		
71		
14		
4		
57		
..		

(Key to Vol. I. page 48.)

Ex. 3. $43)£125 \ 4 \ 9 \ (\text{£}2 \ 18 \ 3 \ \text{Quotient.}$

$$\begin{array}{r}
 \cdot 39 \\
 20 \\
 \hline
)784 \text{--} \\
 354 \\
 \cdot 10 \\
 12 \\
 \hline
)129 \text{----} \\
 \dots \\
 \hline
 \end{array}$$

Ex. 4. $97)£542 \ 7 \ 10 \ (\text{£}5 \ 11 \ 10 \ \text{Quotient.}$

$$\begin{array}{r}
 \cdot 57 \\
 20 \\
 \hline
)1147 \text{--} \\
 \cdot 177 \\
 \cdot 80 \\
 12 \\
 \hline
)970 \text{----} \\
 \dots \\
 \hline
 \end{array}$$

Ex. 5. $127)£123 \ 11 \ 2\frac{1}{2} \ (\text{£}0 \ 19 \ 5\frac{1}{2} \ \text{Quotient.}$

$$\begin{array}{r}
 20 \\
 \hline
)2471 \text{--} \\
 1201 \\
 \cdot \cdot 58 \\
 12 \\
 \hline
)698 \text{----} \\
 \cdot 63 \\
 4 \\
 \hline
)254 \text{----} \\
 \dots \\
 \hline
 \end{array}$$

EXAMPLES OF WEIGHTS AND MEASURES.

<p>Ex. 1. $7) \begin{array}{l} \text{lb. oz. dwt. gr.} \\ 17 \dots 9 \dots 0 \dots 2 \end{array} \text{ Divd.}$</p> <hr style="width: 50%; margin-left: 0;"/> <p style="text-align: center;">$2 \dots 6 \dots 8 \dots 14 \ \text{Quot.}$</p>	<p>Ex. 2. $12) \begin{array}{l} \text{lb} \ \frac{3}{4} \ \text{oz} \ \text{gr.} \\ 17 \dots 5 \dots 2 \dots 1 \dots 4 \end{array} \text{ Div.}$</p> <hr style="width: 50%; margin-left: 0;"/> <p style="text-align: center;">$1 \dots 5 \dots 3 \dots 1 \dots 12 \ \text{Quo.}$</p>
--	--

(Key to Vol. I. page 43.)

Ex. 3. cwt. qr. lb. cwt. qr. lb. Quotient.
 53)178 .. 3 .. 14 (3 .. 1 .. 14
 ·19 | |
 4 | |
 — | |
)79 -- | |
 26 | |
 28 | |
 — | |
)742 ----- | |
 212 | |
 ... | |
 — | |

Ex. 4. mil. fur. po. yd. ft. mil. fur. po. yds. ft. in. Quo.*
 39)144 .. 4 .. 2 .. 1 .. 2 (3 .. 5 .. 25 .. 3 .. 2 .. 6²⁴/₃₉
 27 | | | |
 8 | | | |
 — | | | |
)220 -- | | | |
 ·25 | | | |
 40 | | | |
 — | | | |
)1002 ----- | | | |
 ·222 | | | |
 ·27 | | | |
 11 | | | |
 — | | | |
 2)299 half yds.-- | | | |
 yds. ft. in. | | | |
 149 .. 1 .. 6 | | | |
 2 .. 0 add -- | | | |
 — | | | |
 -)150 .. 0 .. 6 (----- | | | |
 33 | | | |
 ·3 | | | |
 — | | | |
)99 -- | | | |
 21 | | | |
 12 | | | |
 — | | | |
)258 ----- | | | |
 ·24 remainder. | | | |
 — | | | |
 39 divisor. | | | |

* As this Quotient and Dr. Hutton's disagree, we subjoin the proof.

mil. fur. po. yds. ft. in.
 3 .. 5 .. 25 .. 3 .. 2 .. 6²⁴/₃₉
 39
 —————
 Proof 144 .. 4 .. 2 .. 1½ .. 0 .. 6 evidently = 144 .. 4 .. 2 .. 1 .. 2 .. 0

(Key to Vol. I. page 45.)

Ex. 5. yds. qrs. nls. yds. qr. nls.
 47)534 .. 2 .. 2 (11 .. 1 .. 2 *Quotient.*
 ·64
 17
 4
 ———
)70 --
 23
 4
 ———
)94 -----
 ..
 ———

Ex. 6. A. R. P. A. R. P.
 51)71 .. 1 .. 33 (1 .. 1 .. 24 $\frac{2}{3}$ *Quotient. **
 20
 4
 ———
)81 --
 30
 40
 ———
)1233 -----
 ·213
 ··9 remainder.
 ———
 51 divisor.
 ———

* If 77 acres be written in the Dividend instead of 71, the Quotient is 1 A. 2 R. 3 P., the Answer given in the WORK.

Ex. 7. tuns. hhd. gal. pts. tu. hhd. gal. pts.
 65)7 .. 0 .. 47 .. 7 (0 .. 0 .. 27 .. 7 *Quotient.*
 4
 ———
 28 ---
 63
 ———
)1811 -----
 ·511
 ·56
 8
 ———
)455 -----
 ...
 ———

(Key to Vol. I. page 46.)

Quest. 8. cwt. gr. cwt. gr. £. s. d. £. s.
As 7 .. 1 : 43 .. 2 ∴ 1 : 6, ∴ 26 .. 10 .. 4 : 159 .. 2. Ans.

Quest. 9. men. men. £. s. £. s.
As 750 : 3500 ∴ 3 : 14, ∴ 2381 .. 5 : 13212 .. 10. Ans.

Quest. 10. ft. ft. ft. ft. yd. yds.
As $3 \times 3 : 27 \times 20$, that is, As 1 : 60, ∴ 1 : 60. Ans.

Quest. 11. bu. bu. s. d. 34 .. 6 s. d.
As 36 : 6, that is, As 6 : 1, ∴ 34 .. 6 : $\frac{34 \cdot 6}{6} = 5 \cdot 9$. Ans.

Quest. 12. ft. ft. stones. stones.
As 2 : 3 ∴ 6352 : 9528. Ans.

Quest. 13. oz. lb. oz. dwt. s. d. s. d.
As 1 : 73 .. 5 .. 15 ∴ 5 .. 9 : £253 .. 10 .. 0 $\frac{3}{4}$. Ans.

Quest. 14. men. men. days. days.
As 1124 : 536 ∴ 281 : 134, ∴ 365 : 174 $\frac{16}{81}$. Ans.

Quest. 15. s. d. s. s. d.
As £1 : 3 .. 6 ∴ £763 .. 15 : £133 .. 13 .. 1 $\frac{1}{2}$. Ans.

(Page 47.)

Quest. 16. hrs. hrs. days. days.
As 6 : 4, that is, As 3 : 2, ∴ 12 : 8. Ans.

Quest. 17. sh. gs. sh. bu. bu. qr. bu.
As 6 : 90 = 1890 ∴ 1 : 315, ∴ 1 : 315 = 39 .. 3. Ans.

Quest. 18. s. d. s. d.
As £977 : £420 .. 6 .. 3 $\frac{1}{4}$ ∴ £1 : £0 .. 8 .. 7 $\frac{1}{4}$. Ans.

Quest. 19. horse. horse. days. days.
As 2000 : 3000, that is, As 2 : 3, ∴ 18 : 27. Ans.

Quest. 20. day. days. s. d. s. d.
As 1 : 365 ∴ £1 .. 5 .. 6 : £465 .. 7 .. 6.

But £630 - £465 .. 7 .. 6 = £164 .. 12 .. 6. Ans.

Quest. 21. cwt. cwt. qr. lb. s. d. s. d.
As 1 : 33 .. 0 .. 24 ∴ £0 .. 16 .. 4 : £27 .. 2 .. 6. Ans.

Quest. 22. da. da. lb. lb.
As 80 : 54 ∴ 1 $\frac{1}{2}$: 1 $\frac{1}{80}$. Ans.

(Key to Vol. I. page 47.)

- Quest. 23. As $10 \text{ s. } 6 \text{ d.} : £20 :: 1 : 38 \frac{12}{20}$. Ans.
- Quest. 24. As $1 \text{ ch. } 75 \text{ bu. } 7 :: £1 \text{ } 13 \text{ s. } 6 \text{ d.} : £125 \text{ } 19 \text{ s. } 0 \frac{1}{6}$. Ans.
- Quest. 25. As $8 \text{ s. } 4 \text{ d.} : 7 \text{ s. } 3 \text{ d.} :: 8 \text{ oz. } 6 \text{ dr.} : 15 \frac{36}{100}$. Ans.
- Quest. 26. As $1 \text{ A. } 173 \text{ R. } 2 \text{ P. } 14 :: £1 \text{ } 7 \text{ s. } 8 \text{ d.} : £240 \text{ } 2 \text{ s. } 7 \frac{1}{20}$. Ans.
- Qu. 27. As $19 \frac{1}{2} \text{ cwt. } 132 \text{ qr. } 1 \frac{1}{4} :: £10 \text{ } 4 \text{ s.} : £69 \text{ } 4 \text{ s. } 2 \text{ qr. } 1 \frac{3}{8}$. Ans.
- Quest. 28. As $3 \text{ qrs. } 3 \frac{1}{2} \text{ yds. } = 14 :: 7 = 1 \frac{3}{4} : 8 \text{ } 0 \text{ } 2 \frac{2}{3}$. Ans.
- Quest. 29. As $5 \text{ yds. } 191 \text{ qr. } 1 :: 14 \text{ s. } 2 \text{ d.} : £27 \text{ } 1 \text{ s. } 10 \frac{1}{2}$. Ans.
- Quest. 30. As $365 \text{ days. } 1 :: £2107 \text{ } 12 - £500 : £4 \text{ } 8 \text{ s. } 1 \frac{19}{365}$. Ans.

(Page 48.)

- Quest. 31. As $13 \frac{1}{2} \text{ po. } 40 :: 4 : 11 \text{ } 4 \text{ yds. } 2 \text{ ft. } 0 \frac{18}{27}$. Ans.
- Quest. 32. As $4 \text{ qrs. } 266 :: 2 : 133, :: 7 \text{ s. } 9 \frac{1}{2} \text{ d.} : £25 \text{ } 18 \text{ s. } 1 \frac{3}{4}$. Ans.
- Quest. 33. As $3 \text{ cwt. } 1 \text{ qr. } 5 \text{ lb. } 0 \text{ mil. } 14 :: 96 : 151 \text{ } 3 \text{ fur. } 3 \frac{1}{3}$. Ans.
- Quest. 34. As $1 \text{ oz. } 1 \text{ lb. } 7 \text{ dwt. } 14 :: 6 \text{ s. } 4 \text{ d.} : £6 \text{ } 4 \text{ s. } 9 \frac{1}{5}$. Ans.
- Quest. 35. As $1 \text{ A. } 547 :: \frac{15 \text{ s. } 9 \text{ d.}}{2} = 7 \text{ s. } 10 \frac{1}{2} \text{ d.} : £211 \text{ } 19 \text{ s. } 3$. Ans.
- Quest. 36.* As $\frac{9 \text{ m.}}{6 \text{ da.}} : \frac{27 \text{ ft.}}{4 \text{ da.}} :: 6 : 27 :: 2 : 9, :: 16 : 72$. Ans.

* This is a question in Compound Proportion. See Hutton's Practical Arithmetic, Quest. 5, page 38. Lond. 1815.

(Key to Vol. I. page 48.)

Quest. 37. $\begin{array}{c} \text{day} \text{ days.} \quad d. \\ \text{As } 1 : 365 : : 14\frac{1}{2} \times 20 = \end{array} \begin{array}{c} s. \quad d. \\ \text{£}1 \text{ .. } 4 \text{ .. } 2 : \end{array} \begin{array}{c} s. \quad d. \\ \text{£}441 \text{ .. } 0 \text{ .. } 10. \end{array} \text{ Ans.}$

Quest. 38. $\begin{array}{c} \text{ells. qrs.} \quad \text{ells. qrs.} \\ \text{As } 18 \times 3 : 50 \times 4 : : \end{array} \begin{array}{c} s. \quad d. \\ \text{£}1 \text{ .. } 19 \text{ .. } 6 : \end{array} \begin{array}{c} s. \quad d. \\ \text{£}7 \text{ .. } 6 \text{ .. } 3\frac{1}{2}\frac{5}{7}. \end{array} \text{ Ans.}$

Quest. 39. $\begin{array}{c} \text{yd. ft.} \quad \text{yds. ft.} \\ \text{As } 1 \times 2\frac{1}{2} : 20 \times 9 = 180 : : \end{array} \begin{array}{c} \text{yd. yds.} \\ 1 : 72. \end{array} \text{ Ans.}$

Quest. 40. $\begin{array}{c} s. \\ \text{As } \text{£}1 : \text{£}384 \text{ .. } 16 : : \end{array} \begin{array}{c} s. \quad d. \\ \text{£}0 \text{ .. } 2 \text{ .. } 9\frac{1}{2} : \end{array} \begin{array}{c} s. \quad d. \\ \text{£}53 \text{ .. } 14 \text{ .. } 2\frac{1}{2}. \end{array}$
 But $\begin{array}{c} s. \\ \text{£}384 \text{ .. } 16 - \end{array} \begin{array}{c} s. \quad d. \\ \text{£}53 \text{ .. } 14 \text{ .. } 2\frac{1}{2} = \end{array} \begin{array}{c} s. \quad d. \\ \text{£}331 \text{ .. } 1 \text{ .. } 9\frac{1}{2}. \end{array} \text{ Ans.}$

Quest. 41. $\begin{array}{c} \text{hours. min.} \quad \text{hour. miles.} \quad \text{miles.} \\ \text{As } 23 \text{ .. } 56 : 1 : : 25000 : 1044\frac{816}{1436}. \end{array} \text{ Ans.}$

Quest. 42. $\begin{array}{c} s. \quad s. \\ \text{As } 10 : 8, \text{ that is, } \end{array} \begin{array}{c} \text{bot. bottles.} \\ \text{As } 5 : 4, : : 20 : 16. \end{array} \text{ Ans.}$

Quest. 43. $\begin{array}{c} \text{qr. qr. bu.} \\ \text{As } 1 : 43 \text{ .. } 5 : : \end{array} \begin{array}{c} s. \quad d. \\ \text{£}1 \text{ .. } 8 \text{ .. } 6 : \end{array} \begin{array}{c} s. \quad d. \\ \text{£}62 \text{ .. } 3 \text{ .. } 3\frac{1}{3}. \end{array} \text{ Ans.}$

Quest. 44. $\begin{array}{c} \text{qrs. yd. yd. qrs. yds.} \\ \text{As } 5 \times 1 : 1 : : 3 \times 50 = 150 : 30. \end{array} \text{ Ans.}$

Quest. 45. $\begin{array}{c} \text{oz. gr.} \\ \text{As } 1 : 1, \text{ that is, } \end{array} \begin{array}{c} s. \\ \text{As } 480 : 1, : : \end{array} \begin{array}{c} s. \quad d. \\ \text{£}4 \text{ .. } 4 : \end{array} \begin{array}{c} s. \quad d. \\ \text{£}0 \text{ .. } 0 \text{ .. } 2\frac{1}{10}. \end{array} \text{ Ans.}$

Quest. 46. $\begin{array}{c} \text{cwt. lb.} \quad \text{£. s.} \quad \text{£. £. s.} \\ \text{As } 3 : 1 : : 40 \text{ .. } 12 + 10 = 50 \text{ .. } 12 : \end{array} \begin{array}{c} s. \\ \text{£}0 \text{ .. } 3\frac{4}{36}. \end{array} \text{ Ans.*}$

* There is a minute error in the Denominator of the Fraction given in the last Edition of the Course, although in the Sixth Edition the Answer is right.

COMPOUND PROPORTION.

(Page 50.)

Quest. 2. $\begin{array}{c} \text{yd.} \quad \text{yds.} \\ \text{As } \text{£}100 \times 1 : \text{£}750 \times 7 : : 2 : 105 : : \end{array} \begin{array}{c} s. \\ \text{£}5 : \text{£}262 \text{ .. } 10. \end{array} \text{ Ans.}$

Quest. 3. $\begin{array}{c} \text{pers. mo.} \quad \text{pers. mo.} \\ \text{As } 8 \times 9 : 18 \times 12 : : 1 : 3, : : \end{array} \text{£}200 : \text{£}600. \dagger \text{ Ans.}$

† Dr. Hutton's Answer to this Question is half the true Answer.

(Key to Vol. I. page 56.)

Ex. 5. $1362 \div 25 = 54\frac{12}{25}$. equivalent mixed number.

Ex. 6. $2918 \div 17 = 171\frac{1}{17}$. equivalent mixed number.

(Page 57.)*To reduce a whole Number to an equivalent Fraction with a given Denominator.*

Ex. 2. $\frac{12 \times 13}{13} = \frac{156}{13}$ fraction required.

Ex. 3. $\frac{27 \times 11}{11} = \frac{297}{11}$ fraction required.

(Page 58.)*To reduce a Compound Fraction to an equivalent Simple Fraction.*

Ex. 3. $3 \times 4 = 12$ numerator,
and $7 \times 5 = 35$ denominator.

Therefore $\frac{12}{35}$ fraction required.

Ex. 4. $2 \times 3 \times 5 = 30$ numerator,
and $3 \times 5 \times 9 = 135$ denominator.

Hence $\frac{30}{135} = \frac{2}{9}$ Ans.

Or,

Cancelling 3 and 5 in the Numerator and also in the Denominator of the compound fraction, the simple fraction is $\frac{2}{9}$ as before.

Ex. 5. $3\frac{1}{2} = \frac{7}{2}$, And $\frac{2 \times 8 \times 7}{8 \times 8 \times 2} = \frac{7}{8}$ fraction required.

Ex. 6. $\frac{2 \times 5 \times 7 \times 4}{7 \times 8 \times 2 \times 1} = \frac{5}{2}$ fraction required.

Ex. 7. $\frac{2 \times 5}{5 \times 6} = \frac{1}{3}$, and $2 \frac{1}{3} = \frac{7}{3}$ fraction required.

(Key to Vol. I. page 56.)

To reduce Fractions of different Denominators to equivalent Fractions having a common Denominator.

Ex. 2. $2 \times 9 = 18$ First numr.
 $5 \times 7 = 35$ Sec. numr. Therefore $\frac{18}{63}$ and $\frac{35}{63}$ frac'. required.
 And $7 \times 9 = 63$ Com. denr.

(Page 59.)

Ex. 3. $2 \times 5 \times 4 = 40$ First numr.
 $3 \times 3 \times 4 = 36$ Sec. numr. Hence $\frac{40}{60}$, $\frac{36}{60}$, and $\frac{45}{60}$ Ans.
 $3 \times 3 \times 5 = 45$ Third numr.
 And $3 \times 5 \times 4 = 60$ Com. denr.

Ex. 4. $2\frac{3}{5} = \frac{13}{5}$, and $4 = \frac{4}{1}$.

Then $5 \times 5 \times 1 = 25$ First numerator.

$13 \times 6 \times 1 = 78$ Second numerator.

$4 \times 5 \times 6 = 120$ Third numerator.

And $6 \times 5 \times 1 = 30$ Common denominator.

Wherefore $\frac{25}{30}$, $\frac{78}{30}$, and $\frac{120}{30}$ are the fractions required.

(Page 60.)

To find the Value of a Fraction in parts of the Integer.

Ex. 3. 3 numerator.
 20 next inferior denomination.
 — s. d.
 denr. 8) 60 (7 .. 6 value sought.
 ..

Ex. 4. 2 numerator.
 21 next inferior denomination.
 —
 denr. 9) 42 Product.
 —
 4s. 8d. Ans.

(Key to Vol. I. page 60.)

Quest. 5. $\begin{array}{r} s. \quad d. \\ 2 \dots 6 \\ \hline 3 \end{array}$ numerator of the given fraction.

denr. $\overline{4) 7 \dots 6}$ Product.

$\begin{array}{r} s. \quad d. \\ 1 \dots 10\frac{1}{2} \end{array}$ Ans.

Quest. 6. $\begin{array}{r} s. \quad d. \\ 4 \dots 10 \\ \hline 2 \end{array}$ numerator of the given fraction.

denr. $\overline{5) 9 \dots 8}$ Product.

$\begin{array}{r} s. \quad d. \\ 1 \dots 11\frac{1}{5} \end{array}$ Ans.

Quest. 7. 4 lb. Troy. numr. of the given fraction,
12 next inferior denomination.

denr. $\overline{5) 48}$ ounces.

$\begin{array}{r} \hline \text{dwt.} \\ \text{oz. } 9 \dots 12. \end{array}$ Ans.

Quest. 8. 5 numerator of the given fraction.
4 next inferior denomination.

denr. $\overline{16) 20}$ quarters.

$\begin{array}{r} \text{qr. lb.} \\ 1 \dots 7 \end{array}$ Ans.

Quest. 9. 7 numerator of the given fraction.
4 next inferior denomination.

denr. $\overline{8) 28}$ roods.

$\begin{array}{r} \text{ro. poles.} \\ 3 \dots 20. \end{array}$ Ans.

(Key to Vol. I. page 60.)

Quest. 10. 3 numerator of the given fraction.
 24 next inferior denomination.

denr. 10) 72 hours.

hrs. min.
 7 .. 12 Ans.

(Page 61.)

To reduce a Fraction from one Denomination to another.

Ex. 3. $\mathcal{L} \frac{2 \times 20 \times 12}{15 \times 1 \times 1} = \frac{480}{15} = \frac{32}{1} d.$ value sought.

Ex. 4. q. $\frac{2}{5} \times \frac{1}{4} \times \frac{1}{12} \times \frac{1}{20} = \mathcal{L} \frac{1}{2400}$ fraction required.

Ex. 5. Cwt. $\frac{2 \times 4 \times 28}{7 \times 1 \times 1} = \text{lb. } \frac{32}{1}$ fraction required.

Ex. 6. Dwt. $\frac{3}{5} \times \frac{1}{20} \times \frac{1}{12} = \frac{1}{400}$ lb. Troy.

Ex. 7. Cr. $\frac{3}{8} \times \frac{5}{1} \times \frac{1}{21} = \frac{5}{56}$ of a guinea.

Ex. 8. H. cr. $\frac{5}{6} \times \frac{5}{2} = \frac{25}{12}$ of a shilling.

Ex. 9. $\frac{s.}{2} \dots \frac{d.}{6} = 30$, and $\mathcal{L}1 = 240$, therefore
 $\mathcal{L} \frac{30}{240} = \mathcal{L} \frac{1}{8}$ is the fraction required.

(Key to Vol. I. page 61.)

Ex. 10.	$\begin{array}{r} \text{s.} \quad \text{d.} \quad \text{gr.} \\ 17 \dots 7 \dots 3\frac{3}{5} \text{ numerator.} \\ \hline 211 \text{ pence.} \\ \hline 847 \text{ farthings.} \\ \hline \end{array}$	$\begin{array}{r} \text{£ 1 denominator.} \\ 20 \\ \hline 20 \text{ shillings.} \\ 12 \\ \hline 240 \text{ pence.} \\ 4 \\ \hline 960 \text{ farthings.} \\ 5 \\ \hline \end{array}$
	Numerator 4238 fifth-parts of a farthing.	Denominator 4800 fifth-parts of a farthing.

Wherefore $\text{£} \frac{4238}{4800} = \text{£} \frac{2119}{2400}$ is the fraction required.

ADDITION OF VULGAR FRACTIONS.

(Page 62.)

Ex. 4. $\frac{3+6}{7} = \frac{9}{7} = 1\frac{2}{7}$. Ans.

Ex. 5. $\frac{3}{4} = \frac{27}{36}$, and $\frac{5}{9} = \frac{20}{36}$. But $\frac{27+20}{36} = \frac{47}{36} = 1\frac{11}{36}$ sum reqd.

Ex. 6. $\frac{2}{7} = \frac{4}{14}$, and $\frac{4+5}{14} = \frac{9}{14}$ sum of the given fractions.

Ex. 7. $2 \times 5 \times 7 = 70$ First numerator.
 $3 \times 3 \times 7 = 63$ Second numerator.
 $5 \times 3 \times 5 = 75$ Third numerator.
 And $3 \times 5 \times 7 = 105$ Common denominator.
 But $\frac{70 + 63 + 75}{105} = 1\frac{103}{105}$. Ans.

(Key to Vol. I. page 62.)

- Ex. 8. $5 \times 5 \times 6 = 150$ First numerator.
 $3 \times 9 \times 6 = 162$ Second numerator.
 $1 \times 9 \times 5 = 45$ Third numerator.
 And $9 \times 5 \times 6 = 270$ Common denominator.
 But $2 + \frac{150+162+45}{270} = 3\frac{30}{270}$ the sum required.
- Ex. 9. $\frac{3}{5} + \frac{4}{5}$ of $\frac{1}{3} = \frac{13}{15}$. Reducing this fraction and $\frac{3}{20}$ to
 a common denominator, the new fractions are $\frac{52}{60}$ and $\frac{9}{60}$.
 But $9 + \frac{52+9}{60} = 10\frac{1}{60}$ sum required.

- Ex. 10. $\mathcal{L}\frac{2}{3} \times \frac{20}{1} = \frac{40}{3} = \frac{120}{9} sh.$
 And $\frac{120+5}{9} = \frac{125}{9} sh. = 13..10..2\frac{2}{3}$ Ans.

- Ex. 11. $\frac{3}{5} \times \frac{12}{1} = \frac{36}{5} = \frac{108}{15}$. And $\frac{108+4}{15} = \frac{112}{15} = 7..1\frac{1}{3}$. Ans.

- Ex. 12. $\mathcal{L}\frac{1}{7} \times \frac{20}{1} = \frac{20}{7}$. And $\frac{5}{12}d. \times \frac{1}{12}s. = \frac{5}{144}sh.$ Then

$$\frac{20 \times 9 \times 144 + 2 \times 7 \times 144 + 5 \times 7 \times 9}{7 \times 9 \times 144} = \frac{3139}{1008} = 3..1..1\frac{1}{27}$$
 Ans.

SUBTRACTION OF VULGAR FRACTIONS.

(Page 63.)

Quest. 3. $\frac{7-5}{12} = \frac{2}{12} = \frac{1}{6}$ Ans.

Quest. 4. $\frac{3}{13} = \frac{9}{39}$, and $\frac{9-4}{39} = \frac{5}{39}$ Ans.

(Key to Vol. I. page 63.)

Quest. 5. $\frac{5}{12} = \frac{65}{156}$, and $\frac{7}{13} = \frac{84}{156}$. But $\frac{84-65}{156} = \frac{19}{156}$. Ans.

Quest. 6. $\frac{2}{7}$ of $4\frac{1}{6} = 1\frac{4}{21} = 1\frac{32}{168}$, and $\frac{3}{8} = \frac{63}{168}$.

But $5\frac{63}{168} - 1\frac{32}{168} = 4\frac{31}{168}$ Ans.

Quest. 7. $\frac{2}{3} \times \frac{3}{4} = \frac{1}{2} sh. = \frac{9}{18} sh.$ and $\mathcal{L} \frac{5}{9} \times \frac{20}{1} = \frac{100}{9} = \frac{200}{18}$.

But $\frac{200-9}{18} = \frac{191}{18} sh. = 10..7..1\frac{1}{3}$. Ans.

Quest. 8. $\frac{2}{7}$ of $\mathcal{L} 5\frac{1}{6} = \mathcal{L} 1\frac{20}{21}$, and $\frac{3}{5} \times \mathcal{L} \frac{1}{20} = \mathcal{L} \frac{3}{100}$.

But $\mathcal{L} 1\frac{20}{21} - \mathcal{L} \frac{3}{100} = \mathcal{L} 1\frac{937}{2100} = \mathcal{L} 1..8..11\frac{2}{3}$. Ans.

MULTIPLICATION OF VULGAR FRACTIONS.

Ex. 3. $\frac{2}{7} \times \frac{5}{8} = \frac{1}{7} \times \frac{5}{4} = \frac{5}{28}$ Product required.

Ex. 4. $\frac{4}{15} \times \frac{5}{24} = \frac{1}{3} \times \frac{1}{6} = \frac{1}{18}$ Product required.

Ex. 5. $\frac{3}{7} \times \frac{4}{9} \times \frac{14}{15} = \frac{1}{1} \times \frac{4}{9} \times \frac{2}{5} = \frac{8}{45}$ Ans.

(Page 64.)

Ex. 6. $\frac{1}{2} \times \frac{2}{3} \times \frac{3}{1} = \frac{1}{1} \times \frac{1}{1} \times \frac{1}{1} = \frac{1}{1} = 1$. Ans.

Ex. 7. $\frac{7}{9} \times \frac{3}{5} \times \frac{61}{14} = \frac{1}{3} \times \frac{1}{5} \times \frac{61}{2} = \frac{61}{30} = 2\frac{1}{30}$. Prod. reqd.

(Key to Vol. I. page 64.)

$$\text{Ex. 8. } \frac{5}{6} \times \frac{2}{3} \times \frac{6}{7} = \frac{5}{1} \times \frac{2}{3} \times \frac{1}{7} = \frac{10}{21} \text{ Ans.}$$

$$\text{Ex. 9. } \frac{6}{1} \times \frac{2}{3} \times \frac{5}{1} = \frac{2}{1} \times \frac{2}{1} \times \frac{5}{1} = \frac{20}{1} = 20 \text{ Ans.}$$

$$\text{Ex. 10. } \frac{2}{9} \times \frac{3}{5} \times \frac{5}{8} \times \frac{23}{7} = \frac{1}{3} \times \frac{1}{1} \times \frac{1}{4} \times \frac{23}{7} = \frac{23}{84} \text{ Product.}$$

$$\text{Ex. 11. } \frac{23}{7} \times \frac{146}{33} = \frac{3358}{231} = 14\frac{124}{231} \text{ Product required.}$$

$$\text{Ex. 12. } \frac{5}{1} \times \frac{2}{3} \times \frac{2}{7} \times \frac{3}{5} \times \frac{25}{6} = \frac{1}{1} \times \frac{2}{3} \times \frac{1}{7} \times \frac{1}{1} \times \frac{25}{1} = \frac{50}{21} = 2\frac{8}{21} \text{ Ans.}$$

DIVISION OF VULGAR FRACTIONS.

$$\text{Ex. 3. } \begin{array}{r} 4) 16(4 \\ 5) 25(5 \end{array} \text{ Quotient.}$$

$$\text{Ex. 4. } \begin{array}{r} 3) 7(28 \\ 4) 16(48 \end{array} = \frac{7}{12} \text{ Quotient.}$$

$$\text{Ex. 5. } \begin{array}{r} 7) 14(84 \\ 6) 9(63 \end{array} = 1\frac{1}{3} \text{ Quotient.}$$

$$\text{Ex. 6. } \begin{array}{r} 15) 5(35 \\ 7) 6(90 \end{array} = \frac{7}{18} \text{ Quotient.}$$

$$\text{Ex. 7. } \begin{array}{r} 3) 12(4 \\ 5) 35(7 \end{array} \text{ Quotient.}$$

$$\text{Ex. 8. } \begin{array}{r} 3) 2(10 \\ 5) 7(21 \end{array} \text{ Quotient.}$$

(Page 65.)

$$\text{Ex. 9. } \begin{array}{r} 3) 9(3 \\ 1) 16(16 \end{array} \text{ Quotient.}$$

(Key to Vol. I. page 65.)

Ex. 10. $\frac{2}{1} \Big) \frac{3}{5} \left(\frac{3}{10} \right.$ *Quotient.*

Ex. 11. $\frac{86}{9} \Big) \frac{22}{3} \left(\frac{198}{258} = \frac{33}{43} \right.$ *Quotient.*

Ex. 12. $\frac{38}{7} \Big) \frac{2}{9} \left(\frac{14}{342} = \frac{7}{171} \right.$ *Quotient.*

RULE OF THREE IN VULGAR FRACTIONS.

Quest. 2. As $\frac{1}{1}$ oz. : $\frac{27}{8}$ oz. :: £0 .. 6 .. 8 = £ $\frac{1}{3}$: £ $\frac{27}{4}$ = £1 .. 1 .. 4 $\frac{1}{2}$. Ans.

Quest. 3. As $\frac{6}{32}$: $\frac{5}{32}$, that is, As 6 : 5, £273 .. 2 .. 6 : £227 .. 12 .. 1. Ans.

Quest. 4. As £ $\frac{100}{1}$: £108 $\frac{1}{2}$:: £1230 : £1336 .. 1 .. 9. Ans.

Quest. 5. As £ $\frac{100}{1}$: £3 $\frac{1}{4}$:: £273 $\frac{3}{4}$: £8 .. 17 .. 11 $\frac{1}{4}$. Ans.

Quest. 6. As £73 $\frac{1}{10}$: £250 $\frac{1}{2}$:: $\frac{1}{8}$: $\frac{3}{7}$. Ans.

Quest. 7. As 7 $\frac{3}{4}$ inches. : 12 in. :: 12 in. : 18 $\frac{18}{31}$. Ans.*

Quest. 8. As $\frac{3}{4}$ yd. : 2 $\frac{1}{2}$ yds., that is, As 3 : 10, :: 9 $\frac{1}{2}$ yds. : 31 $\frac{2}{3}$. Ans.

* In some Editions of the *Course* (and perhaps in the last) there is an Error in the Answer.

(Page 66.)

Quest. 9. As 8 $\frac{1}{2}$ sh. : 5 oz. :: 6 $\frac{9}{10}$ sh. : 4 $\frac{1}{17}$ oz. Ans.

Quest. 10. As 11 $\frac{11}{12}$ po. : 40 po. :: 4 po. : 13 $\frac{61}{43}$. Ans.

Quest. 11. As 11 $\frac{9}{10}$ hrs. : 13 $\frac{5}{8}$ hrs. :: 35 $\frac{1}{2}$ days. : 40 $\frac{615}{952}$. Ans.

Quest. 12. $\frac{7}{8}$ yd. : $\frac{13}{8}$ yds., that is, As 7 : 13, :: 2 $\frac{1}{2}$ yds. : $\frac{65}{14}$ = 4 $\frac{9}{14}$ yards.

But 4 $\frac{9}{14}$ × 976 = 4531 $\frac{3}{7}$ = 4531 .. 1 .. 2 $\frac{6}{7}$. Ans.

(Key to Vol. I. page 67.)

DECIMAL FRACTIONS.

ADDITION OF DECIMALS.

<p>Ex. 2.</p> <table style="margin-left: 20px;"> <tr><td>276,</td></tr> <tr><td>39,213</td></tr> <tr><td>72014,9</td></tr> <tr><td>417,</td></tr> <tr><td>5032,</td></tr> <tr><td style="border-top: 1px solid black;">Sum 77779,113</td></tr> </table>	276,	39,213	72014,9	417,	5032,	Sum 77779,113	<p>Ex. 3.</p> <table style="margin-left: 20px;"> <tr><td>7530,</td></tr> <tr><td>16,201</td></tr> <tr><td>3,0142</td></tr> <tr><td>957,13</td></tr> <tr><td>6,72119</td></tr> <tr><td>,03014</td></tr> <tr><td style="border-top: 1px solid black;">Sum 8513,09653</td></tr> </table>	7530,	16,201	3,0142	957,13	6,72119	,03014	Sum 8513,09653
276,														
39,213														
72014,9														
417,														
5032,														
Sum 77779,113														
7530,														
16,201														
3,0142														
957,13														
6,72119														
,03014														
Sum 8513,09653														

Ex. 4. Ans. 17500,9718.

SUBTRACTION OF DECIMALS.

(Page 68.)

<p>Ex. 2.</p> <table style="margin-left: 20px;"> <tr><td>From 2,73</td></tr> <tr><td>Take 1,9185</td></tr> <tr><td style="border-top: 1px solid black;">0,8115 Rem.</td></tr> </table>	From 2,73	Take 1,9185	0,8115 Rem.	<p>Ex. 3.</p> <table style="margin-left: 20px;"> <tr><td>From 214,81</td></tr> <tr><td>Take 4,90142</td></tr> <tr><td style="border-top: 1px solid black;">Rem. 209,90858</td></tr> </table>	From 214,81	Take 4,90142	Rem. 209,90858
From 2,73							
Take 1,9185							
0,8115 Rem.							
From 214,81							
Take 4,90142							
Rem. 209,90858							
<p>Ex. 4.</p> <table style="margin-left: 20px;"> <tr><td>From 2714,</td></tr> <tr><td>Take ,916</td></tr> <tr><td style="border-top: 1px solid black;">Rem. 2713,084</td></tr> </table>	From 2714,	Take ,916	Rem. 2713,084				
From 2714,							
Take ,916							
Rem. 2713,084							

MULTIPLICATION OF DECIMALS.

(Page 69.)

<p>Ex. 2.</p> <table style="margin-left: 20px;"> <tr><td>Multiply 79,347</td></tr> <tr><td>by 23,15</td></tr> <tr><td style="border-top: 1px solid black;">Product 1836,88305</td></tr> </table>	Multiply 79,347	by 23,15	Product 1836,88305	<p>Ex. 3.</p> <table style="margin-left: 20px;"> <tr><td>Multiply ,63478</td></tr> <tr><td>by ,8204</td></tr> <tr><td style="border-top: 1px solid black;">Product ,520773512</td></tr> </table>	Multiply ,63478	by ,8204	Product ,520773512
Multiply 79,347							
by 23,15							
Product 1836,88305							
Multiply ,63478							
by ,8204							
Product ,520773512							

(Key to Vol. I, page 69.)

Ex. 4. Multiply ,385746
 by ,00464

,00178986144 Product.

CONTRACTION I.

- Ex. 2. $2.714 \times 100 = 271.4$ Product required.
 Ex. 3. $.916 \times 1000 = 916$ Product required.
 Ex. 4. $21.31 \times 10000 = 213100$ Product required.

CONTRACTION II.

(Page 70.)

Ex. 2. 480,14936 Multiplicand.
 61427.2 Multiplier reversed.

9602987
 3361045
 96030
 19206
 480
 288

1308.0036 Product required.

Ex. 3. 2490,30480 Multiplicand.
 682375.0 Multiplier reversed.

124515240
 17432134
 747091
 49806
 19922
 1494

1427.65687 Product required.

(Key to Vol. I. page 70.)

Ex. 4. Ans. 235·104|0907965204.

 DIVISION OF DECIMALS.

(Page 71.)

- Ex. 3. 54·25)123·70536(2·2802 Quotient.
 Ex. 4. ·7854)12·0000000(15·278 Quotient.
 Ex. 5. 100)4195·68(41·9568 Quotient.
 Ex. 6. ·153)·8297592(5·4232 Quotient.

 CONTRACTION I.

(Page 72.)

- Ex. 2. Quotient 1·281875
 Ex. 3. Quotient ·04412037037 &c.
 Ex. 4. Quotient ·0007721519 *nearly*.

 CONTRACTION II.

- Ex. 2. 100)5·16(·0516 Quotient.
 Ex. 3. 10)419(41·9 Quotient.
 Ex. 4. 1000)·21(·00021 Quotient.

 CONTRACTION III.

(Page 73.)

- Ex. 2. 230·409)41092351(17·8345 Quotient.
 180514
 19228
 796
 105
 13
 2
 —

(Key to Vol. I. page 75.)

Ex. 5. $\overset{\text{lb.}}{2 \cdot 15} \div (28 \times 4) = \overset{\text{cwt.}}{.019196} \text{ \&c. the decimal required.}$

Ex. 6. $\overset{\text{yds.}}{24} \div 1760 = \overset{\text{miles.}}{.013636} \text{ \&c. Ans.}$

Ex. 7. $\overset{\text{poles.}}{.056} \div (40 \times 4) = \overset{\text{acres.}}{.00035} \text{ the decimal required.}$

Ex. 8. $\overset{\text{pt.}}{1 \cdot 2} \div (8 \times 63) = \overset{\text{hhd.}}{.00238} \text{ \&c. Ans.}$

Ex. 9. $\overset{\text{min.}}{14} \div (60 \times 24) = \overset{\text{day}}{.009722} \text{ \&c. decimal required.}$

Ex. 10. $\overset{\text{pt.}}{.21} \div 16 = \overset{\text{peck}}{.013125} \text{ Ans.}$

Ex. 11.* $28'' \text{ .. } 12''' = \frac{28}{60} + \frac{12}{60} \text{ of } \frac{1}{60} \text{ minute.}$

Also $\overset{\text{min.}}{28} \div 60 = .466\dot{6} \text{ minute.}$

And $\overset{\text{min.}}{12} \div 3600 = 003\dot{3} \text{ minute.}$

Sum of both decimals $.47'$ the decimal required.

Otherwise,

$$28'' \text{ .. } 12''' = 1692''', = \frac{1692}{3600} \text{ minute} = .47' \text{ as before.}$$

* See this question solved the 6th in the next article.

When there are several Numbers to be reduced Collectively to the Decimal of the highest Denomination.

(Page 76.)

Ex. 2. $\begin{array}{r|l} \text{qr. } 4 & 1 \cdot \\ \text{d. } 12 & 3 \cdot 25 \\ \text{s. } 20 & 17 \cdot 27083\dot{3} \\ \hline \text{£ } & 19 \cdot 863541\dot{6} \end{array} \text{ decimal required.}$

Ex. 3. $\begin{array}{r|l} \text{d. } 12 & 6 \cdot \\ \text{s. } 20 & 15 \cdot 5 \\ \hline \text{£ } & 0 \cdot 775 \end{array} \text{ decimal required.}$

(Key to Vol. I. page 76.)

$$\begin{array}{r|l} \text{Ex. 4.} & \text{qr. 4} \quad | \quad 2 \cdot \\ & \text{d. 12} \quad | \quad 7 \cdot 5 \\ & \text{sh. 0.625 decimal required.} \end{array}$$

$$\begin{array}{r|l} \text{Ex. 5.} & \text{gr. 24} \quad | \quad 16 \cdot \\ & \text{dwt. 20} \quad | \quad 12 \cdot 666 \\ & \text{oz. 12} \quad | \quad 5 \cdot 6333 \\ & \text{lb. 0.46944. Ans.} \end{array}$$

$$\begin{array}{r|l} \text{Ex. 6.} & \text{(Referred to, above.)} \\ & \text{thirds 60} \quad | \quad 12'' \\ & \text{seconds 60} \quad | \quad 28 \cdot 2'' \\ & \text{Minute 0.47'. Ans. [See Ex. 11. last article.]} \end{array}$$

DUODECIMALS.

(Page 77.)

$$\begin{array}{r} \text{Ex. 3.} \quad \text{ft. in.} \\ 4 \quad \cdot \quad 7 \\ 9 \quad \cdot \quad 6 \\ \hline 41 \quad \cdot \quad 3 \\ 2 \quad \cdot \quad 3 \quad \cdot \quad 6 \end{array}$$

Sq. feet. $43 \quad \cdot \quad 6 \quad \cdot \quad 6$ Prod.

$$\begin{array}{r} \text{Ex. 4.} \quad \text{ft. in.} \\ 12 \quad \cdot \quad 5 \\ 6 \quad \cdot \quad 8 \\ \hline 74 \quad \cdot \quad 6 \\ 8 \quad \cdot \quad 3 \quad \cdot \quad 4 \end{array}$$

Sq. ft. $82 \quad \cdot \quad 9 \quad \cdot \quad 4$ Prod.

$$\begin{array}{r} \text{Ex. 5.} \quad \text{ft. in. } \frac{1}{2}\text{ths.} \\ 35 \quad \cdot \quad 4 \quad \cdot \quad 6 \\ 12 \quad \cdot \quad 3 \end{array}$$

$$\begin{array}{r} 424 \quad \cdot \quad 6 \quad \cdot \quad 0 \\ 8 \quad \cdot \quad 10 \quad \cdot \quad 1 \quad \cdot \quad 6 \\ \hline \text{Sq. feet } 433 \quad \cdot \quad 4 \quad \cdot \quad 1 \quad \cdot \quad 6 \text{ Prod.} \end{array}$$

$$\begin{array}{r} \text{Ex. 6.} \quad \text{ft. in. } \frac{1}{2}\text{ths.} \\ 64 \quad \cdot \quad 6 \quad \cdot \quad 0 \\ 8 \quad \cdot \quad 9 \quad \cdot \quad 3 \end{array}$$

$$\begin{array}{r} 516 \quad \cdot \quad 0 \quad \cdot \quad 0 \\ 48 \quad \cdot \quad 4 \quad \cdot \quad 6 \quad \cdot \quad 0 \\ 1 \quad \cdot \quad 4 \quad \cdot \quad 1 \quad \cdot \quad 6 \quad \cdot \quad 0 \\ \hline \text{Sq. ft. } 565 \quad \cdot \quad 8 \quad \cdot \quad 7 \quad \cdot \quad 6 \quad \cdot \quad 0 \text{ Pro.} \end{array}$$

* * In this Rule the second term of the Product, to adapt it to the capacity of workmen, is generally denominated inches; but it is requisite the student should clearly understand that inches con-

(Key to Vol. I. page 77.)

stitute the *third* term in Square Measure, and the *fourth* in Solid; of which last species, however, the author of the *COURSE* has given no example.

Now it is only necessary to remember that, in Decimals, Duodecimals, Time, and Degrees, the value decreases *gradatim* in Geometrical Progression from the left hand to the right; and that the common ratio in Decimals is 10, in Duodecimals 12, and in Time and Degrees 60. Hence the three series decreasing from unity are,

	0.	1.	2.	3.	4.	5.	Terms.
Decimals.	$\frac{1}{1}$,	$\frac{1}{10}$,	$\frac{1}{100}$,	$\frac{1}{1000}$,	$\frac{1}{10000}$,	$\frac{1}{100000}$,	&c.
Duodecimals.	$\frac{1}{1}$,	$\frac{1}{12}$,	$\frac{1}{144}$,	$\frac{1}{1728}$,	&c.		
Time and Degrees.	$\frac{1}{1}$,	$\frac{1}{60}$,	$\frac{1}{3600}$,	$\frac{1}{216000}$,	&c. called also <i>minutes, seconds, thirds, fourths, fifths, &c.</i>		

The object of Duodecimals is to ascertain, by an easy method, the number of square, or of solid feet, in a piece of work or other body to which the Rule is applicable; and it is evident, that in multiplying length by breadth, feet into feet, give *square feet*; and inches into inches, *square inches*: feet, therefore, into inches must give *12th parts of square feet for the second term of the Product*. In the subsequent pages of the *KEY*, when admeasurement of this kind occurs, we shall place *pts.* (meaning *12th parts of sq. feet*) before inches; as 193 sq. ft. 10 pts. 7 inches, and in the meantime it is thought a further elucidation to subjoin an example of Solid Measure, as very proper to come under the present Rule.

$\begin{matrix} \text{ft.} & \text{in.} & & \text{ft.} & \text{in.} \\ \text{Required the solidity of a wall} & 130 \text{ .. } 8 \text{ long,} & & 10 \text{ .. } 4 \text{ high, and} \\ \text{ft. in.} & & & & & \\ 2 \text{ .. } 3 \text{ in thickness throughout.} & & & & & \end{matrix}$

$\begin{matrix} \text{ft.} & \text{in.} \\ 130 \text{ .. } 8 \text{ long.} \\ \text{Multiplier } 10 \text{ .. } 4 \text{ high.} \end{matrix}$

$\begin{matrix} \text{Sq}^{\text{r}}. \text{ feet} & 1306 \text{ .. } 8 & \text{Prod. of the length by 10 feet high.} \\ & 43 \text{ .. } 6 \text{ .. } 8 & \text{Prod. of the length by 4 inches high.} \end{matrix}$

$\begin{matrix} \text{Sq}^{\text{r}}. \text{ feet} & 1350 \text{ .. } 2 \text{ .. } 8 & \text{area of either side of the wall.} \\ \text{Multiplier} & 2 \text{ .. } 3 \text{ .. } 0 & \text{thickness of the wall.} \end{matrix}$

$\begin{matrix} \text{Solid feet} & 2700 \text{ .. } 5 \text{ .. } 4 & \text{Prod. of the area by 2 feet thick.} \\ & 337 \text{ .. } 6 \text{ .. } 8 \text{ .. } 0 & \text{Prod. of the area by 3 inches thick.} \end{matrix}$

$\begin{matrix} \text{Cubic feet} & 3038 \text{ .. } 0 \text{ .. } 0 \text{ .. } 0 & \text{the content required.} \end{matrix}$

(Key to Vol. I. page 79.)

INVOLUTION.

Quest. 1. $45 \times 45 = 2025.$ Ans.

Quest. 2. $4.16 \times 4.16 = 17.3056.$ Ans.

Quest. 3. $3.5 \times 3.5 \times 3.5 = 42.875.$ Ans.

Quest. 4. $.029 \times .029 \times .029 \times .029 \times .029 = .000000020511149.$
Ans.

Quest. 5. $\frac{2}{3} \times \frac{2}{3} = \frac{4}{9}$ Ans.

Quest. 6. $\frac{5}{9} \times \frac{5}{9} \times \frac{5}{9} = \frac{125}{729}$ Ans.

Quest. 7. $\frac{3}{4} \times \frac{3}{4} \times \frac{3}{4} \times \frac{3}{4} = \frac{81}{256}$ Ans.

EVOLUTION.

(Page 83.)

Quest. 3. $\begin{array}{r} \dot{2}0\dot{2}5(45 \text{ Ans.} \\ 16 \\ 85) \overline{425} \\ \underline{425} \\ \dots \end{array}$

Quest. 4. $\begin{array}{r} \dot{1}7.\dot{3}0\dot{5}6(4.16 \text{ Root required.} \\ 16 \\ 81) \overline{130} \\ \underline{81} \\ 826) \overline{4956} \\ \underline{4956} \\ \dots \end{array}$

Quest. 5. $\begin{array}{r} \dot{0}00\dot{7}2\dot{9}(.027 \text{ Root required.} \\ 4 \\ 47) \overline{329} \\ \underline{329} \\ \dots \end{array}$

(Key to Vol. I. page 84.)

Quest. 3. $\sqrt{\frac{9}{12}} = \sqrt{\cdot 75} = \cdot 8660254 \text{ \&c. } \text{Ans.}$

Quest. 4. $\sqrt{\frac{5}{12}} = \sqrt{\cdot 4166\dot{6}} = \cdot 645497 \text{ \&c. } \text{Root reqd.}$

Quest. 5. $\sqrt{17\frac{3}{8}} = \sqrt{17\cdot 3750} = 4\cdot 168333. \text{ Ans.}$

The 4th, 8th, 16th, 32nd, &c. Root, being an Integer Power of 2.

Quest. 2. $\sqrt[4]{97\cdot 41} = \sqrt{\sqrt{97\cdot 41}} = \sqrt{9\cdot 86965} \text{ \&c. } = 3\cdot 1416 \text{ very nearly. } \text{Ans.}$

CUBE ROOT.

(Page 86.)

Ex. 2.
$$\begin{array}{r|l} 3 \times 8^2 = 192 & 571482\cdot 190 \text{ (82\cdot 9 \&c. Root reqd.)} \\ 3 \times 8 = 24 & 512 \end{array}$$

1st Divisor 1944) 59482

$$\left. \begin{array}{l} 3 \times 8^2 \times 2 = \text{-----} 384 \\ 3 \times 8 \times 2^2 = \text{-----} 96 \\ 2^3 = \text{-----} 8 \end{array} \right\} \text{add}$$

$$\begin{array}{r|l} 3 \times 82^2 = 20172 & \\ 3 \times 82 = 246 & \end{array} \quad \begin{array}{l} \\ \\ \hline 39368 \text{ Subtrahend.} \end{array}$$

2d Div^r. 201966 - - - - -) 20114190 Resolvend.

$$\left. \begin{array}{l} 3 \times 82^2 \times 9 = \text{----} 181548 \\ 3 \times 82 \times 9^2 = \text{----} 19826 \\ 9^3 = \text{-----} 729 \end{array} \right\} \text{add}$$

18353789 Subtrahend.

1760401 Remainder.

Quest. 3. Ans. 11\cdot 76 and 1778424 Remainder.

Quest. 4. Ans. 11\cdot 0027 and 19659410317 Remaining.

(Key to Vol. I. page 89.)

$$\begin{array}{l} \text{And } 6.912 = n + 1 \text{ times } A. \quad \text{Also } 8 = n + 1 \text{ times } P. \\ 4 = n - 1 \text{ times } P. \quad 3.456 = n - 1 \text{ times } A. \end{array}$$

10.912 sum.

11.456 sum.

But, As $10.912 : 11.456 :: 1.2 (=r) : 1.25982$. First approximation to the root required.

Proceeding now with 1.25982 as r , the result is,
 1.259921 the Root, *nearly*.

Quest. 2. Here $P = 3214$
 $n = 3$
 $r = 14.7$
 $A = 3176.523$, whence 14.75758 . Ans.

Quest. 3. $\sqrt[4]{2} = \sqrt[2]{\sqrt[2]{2}} = \sqrt{1.4142136} = 1.189207$ &c. Ans.

Quest. 4. Here $P = 97.41$
 $n = 4$
 $r = 3.1416$
 $R = 3.1415999$, which was required.*

* See the last Quest. page 84, vol. i. of the *Course*, and its Answer in the **KEY**.

Quest. 5. $P = 2$
 $n = 5$
 $r = 1.14$
 $A = 1.9254145824$.

And, As $19.5524874944 : 19.7016583296 :: 1.14 (=r) : 1.148697$
 first approximation to the 5th Root of 2. Proceeding next with 1.148697 as r , the result is 1.148699 &c. Root, *nearly*.

Quest. 6. $\sqrt[6]{21035.8} = \sqrt[3]{\sqrt[2]{21035.8}}$ or
 $\sqrt[2]{\sqrt[3]{21035.8}} = \sqrt{27.60491056} = 5.254037$. Ans.

Quest. 7. $\sqrt[6]{2} = \sqrt[2]{\sqrt[3]{2}} = \sqrt{1.259921} = 1.122462$. Ans.

Or, by the Rule for GENERAL ROOTS,

As $(n+1)^{\frac{1}{2}} A + (n-1)^{\frac{1}{2}} P : P \oslash A :: r : R \oslash r$. That is,
 As $11.5464509325635 : .129585447839 :: 1.11 : .012457 = R \oslash r$.

Add, Root of the assumed Power, 1.11

Root (*nearly*) 1.122457

And 1.122457 (*the Root last found*) is, by another approximation, discovered to be $.000005$ too little, consequently 1.122462 Root, *as before*. Ans.

(Key to Vol. I. page 89.)

Quest. 8. $P = 21035 \cdot 8$

$$n = 7$$

$$r = 4$$

$$A = 16384, \text{ whence } 4 \cdot 1 \text{ is the 1st approximation.}$$

But $4 \cdot 1^7 = 19475 \cdot 4273881$, whereby is obtained $4 \cdot 145392$ the Root, *nearly*.

Quest. 9. $P = 2$

$$n = 7$$

$$r = 1 \cdot 1$$

$$A = 1 \cdot 7777771$$

From which, by two approximations, is found $1 \cdot 104089$. Ans.

Quest. 10. $\sqrt[3]{21035 \cdot 8} = \sqrt[2]{\sqrt[2]{21035 \cdot 8}} = 3 \cdot 470323$. Root reqd.

Quest. 11. $\sqrt[3]{2} = \sqrt[2]{\sqrt[2]{2}} = \sqrt[4]{2} = \sqrt{1 \cdot 189207} = 1 \cdot 090508$. Ans.

Quest. 12. $\sqrt[3]{21035 \cdot 8} = \sqrt[3]{27 \cdot 60491056} = 3 \cdot 022239$.
Ans.

Quest. 13. $\sqrt[3]{2} = \sqrt[3]{1 \cdot 259921} = 1 \cdot 080059$. Ans.

ARITHMETICAL PROPORTION.

(Page 112.)

PROB. I.

Quest. 2. $\frac{(12 + 1) \times 12}{2} = 78$. Ans.

(Page 113.)

Quest. 3. $\frac{(24 + 1) \times 24}{2} = 300$. Ans.

Quest. 4. $\frac{(103 + 1) \times 52}{2} = 2704s. = \text{£}135 \cdot 4$. Ans.

PROB. II.

Quest. 2. $\frac{70 - 10}{20} = 3$ the common difference.

And $\frac{(70 + 10) \times 21}{2} = 840$ the sum of the series.

(Key to Vol. I. page 113.)

$$\text{Quest. 3. } \frac{103 - 1}{52 - 1} = \frac{102}{51} = 2. \text{ Ans.}$$

(Page 114.)

PROB. III.

Quest. 2. $3 \times 20 = 60$ the difference of the extremes,
 And $70 - 60 = 10$ the less extreme.

$$\text{But } \frac{(70 + 10) \times 21}{2} = 840 \text{ the sum of the series.}$$

Therefore 10, and 840. Ans.

Quest. 3. $2 \times 51 = 102$ the difference of the extremes.

And $102 + 1 = 103$ the greater extreme.

$$\text{Also } \frac{(103 + 1) \times 52}{2} = 2704 = \mathcal{L}135 \text{ .. } 4 \text{ the sum of the series.}$$

Therefore $\mathcal{L}135 \text{ .. } 4$ the debt, and $\mathcal{L}5 \text{ .. } 3$ the last payment. Ans.

GEOMETRICAL PROGRESSION.

(Page 117.)

Quest. 1. $1 \times 512 = 512$ the greater extreme,

$$\text{And } \frac{511 + 512}{2 - 1} = 1023 \text{ the sum of the series.}$$

Therefore 512, and 1023. Ans.

Quest. 2. $\mathcal{L}1 \times \mathcal{L}2048 = \mathcal{L}2048$ the greater extreme,

$$\text{and } \frac{\mathcal{L}2048 + \mathcal{L}2047}{2 - 1} = \mathcal{L}4095 \text{ sum of the series.}$$

Therefore $\mathcal{L}4095$ the debt, and $\mathcal{L}2048$ the last payment. Ans.

(Key to Vol. I. page 121.)

SINGLE FELLOWSHIP.

Quest. 3. As £120 : £30 :: £75 : £18 .. 15^{s.} = C's share,
and consequently £11 .. 5 = D's share.

Quest. 4.
$$\begin{array}{r} \text{As } £700 : £125 .. 10 :: £123 : £22 .. 1 .. 0 .. 2\frac{2}{5} = \text{E's share,} \\ 700 : 125 .. 10 :: 358 : 64 .. 3 .. 8 .. 0\frac{3}{5} = \text{F's share.} \\ 700 : 125 .. 10 :: 219 : 39 .. 5 .. 3 .. 1\frac{1}{5} = \text{G's share.} \end{array}$$

£125..10

Quest 5.
$$\begin{array}{r} \begin{array}{l} 250 \\ 350 \\ 400 \\ 500 \end{array} \left. \vphantom{\begin{array}{l} 250 \\ 350 \\ 400 \\ 500 \end{array}} \right\} \text{are as } \left\{ \begin{array}{l} 5 \\ 7 \\ 8 \\ 10 \end{array} \right. \left\| \begin{array}{l} \text{As } 30 : 5 :: 700 : 116 .. 13 .. 4 \dots \text{1st village.} \\ \text{As } 30 : 7 :: 700 : 163 .. 6 .. 8 \dots \text{2d village.} \\ \text{As } 30 : 8 :: 700 : 186 .. 13 .. 4 \dots \text{3d village.} \\ \text{As } 30 : 10 :: 700 : 233 .. 6 .. 8 \dots \text{4th village.} \end{array} \right. \\ \text{Sum } 30 \qquad \qquad \qquad \underline{\underline{£700}}$$

Quest. 6.
$$\begin{array}{r} £500 \\ 320 \\ 75 \end{array} \left. \vphantom{\begin{array}{r} £500 \\ 320 \\ 75 \end{array}} \right\} \text{are as } \left\{ \begin{array}{r} £100 \\ 64 \\ 25 \end{array} \right. \\ \text{Sum } \underline{\underline{£189}}$$

And
$$\begin{array}{r} \text{As } £189 : £100 :: 37 .. 2 .. 14 : 20 .. 3 .. 39\frac{139}{179} \dots \text{L's share.} \\ 189 : 64 :: 37 .. 2 .. 14 : 13 .. 1 .. 30\frac{46}{179} \dots \text{M's share.} \\ 189 : 25 :: 37 .. 2 .. 14 : 3 .. 0 .. 23\frac{173}{179} \dots \text{N's share.} \end{array}$$

37..2..14

Quest. 7.
$$\begin{array}{r} £ 57 .. 15 .. 0 \\ 108 .. 3 .. 8 \\ 22 .. 0 .. 10 \\ 73 .. 0 .. 0 \\ \hline £260 .. 19 .. 6 \text{ Sum. And,} \end{array}$$

(Key to Vol. I, page 121.)

$$\begin{array}{r}
 \text{As } \begin{array}{cccccc} \mathcal{L}. & s. & d. & \mathcal{L}. & s. & d. \end{array} : \begin{array}{cccccc} \mathcal{L}. & s. & d. & \mathcal{L}. & s. & d. \end{array} \text{ qrs.} \\
 260 \dots 19 \dots 6 : 170 \dots 14 \dots : 57 \dots 15 \dots 0 : 37 \dots 15 \dots 5 \dots 2 \frac{5302}{10439} & O. \\
 260 \dots 19 \dots 6 : 170 \dots 14 \dots : 108 \dots 3 \dots 8 : 70 \dots 15 \dots 2 \dots 2 \frac{7498}{10439} & P. \\
 260 \dots 19 \dots 6 : 170 \dots 14 \dots : 22 \dots 0 \dots 10 : 14 \dots 8 \dots 4 \dots 0 \frac{4720}{10439} & Q. \\
 260 \dots 19 \dots 6 : 170 \dots 14 \dots : 73 \dots 0 \dots 0 : 47 \dots 14 \dots 11 \dots 2 \frac{3358}{10439} & R. \\
 \hline
 & \mathcal{L} \ 170 \dots 14 \dots 0 \dots 0
 \end{array}$$

(Page 122.)

Quest. 8. $\mathcal{L}900 - \mathcal{L}540 = \mathcal{L}360$ the loss sustained by S, T, and V.

$$\text{And } \left\{ \begin{array}{l} \mathcal{L}360 \times \frac{1}{3} = \mathcal{L} 45 = \text{S's loss.} \\ 360 \times \frac{1}{4} = 90 = \text{T's loss.} \\ 360 \times \frac{1}{8} = 225 = \text{V's loss.} \end{array} \right\} \text{ Ans.}$$

$\mathcal{L}360$

$$\begin{array}{r}
 \text{Quest. 9.} \\
 \frac{1}{2} \text{ of } 25 = 12 \dots 6 \\
 \frac{1}{3} \text{ of } 25 = 8 \dots 4 \\
 \frac{1}{4} \text{ of } 25 = 6 \dots 3 \\
 \frac{1}{5} \text{ of } 25 = 5 \dots 0 \\
 \hline
 \text{Sum } 32 \dots 1
 \end{array}$$

$$\text{And } \left\{ \begin{array}{l} \text{As } \begin{array}{cccccc} s. & d. & s. & s. & d. & s. & d. & \text{qr.} \end{array} \\
 32 \dots 1 : 25 \dots : 12 \dots 6 : 9 \dots 8 \dots 3 \frac{4}{77} \text{ ---- W.} \\
 32 \dots 1 : 25 \dots : 8 \dots 4 : 6 \dots 5 \dots 3 \frac{5}{77} \text{ ---- X.} \\
 32 \dots 1 : 25 \dots : 6 \dots 3 : 4 \dots 10 \dots 1 \frac{5}{77} \text{ ---- Y.} \\
 32 \dots 1 : 25 \dots : 5 \dots 0 : 3 \dots 10 \dots 3 \frac{1}{77} \text{ ---- Z.} \end{array} \right\} \text{ Ans.}$$

25 shillings.

Otherwise.

The fractions $\frac{1}{2}$, $\frac{1}{3}$, $\frac{1}{4}$, $\frac{1}{5}$, brought to a common denominator, are, $\frac{30}{60}$, $\frac{20}{60}$, $\frac{15}{60}$, and $\frac{12}{60}$; but the shares will be as the numerators of these new fractions, the sum of which is 77. That is,

$$\text{As } \begin{array}{cccccc} s. & s. & s. & s. & d. & \text{qr.} \\
 77 : 25 \dots : 30 : 9 \dots 8 \dots 3 \frac{4}{77} \text{ ---- W.} \\
 77 : 25 \dots : 20 : 6 \dots 5 \dots 3 \frac{5}{77} \text{ ---- X.} \\
 77 : 25 \dots : 15 : 4 \dots 10 \dots 1 \frac{5}{77} \text{ ---- Y.} \\
 77 : 25 \dots : 12 : 3 \dots 10 \dots 3 \frac{1}{77} \text{ ---- Z.} \end{array} \left. \vphantom{\begin{array}{l} 77 : 25 \dots : 30 : 9 \dots 8 \dots 3 \frac{4}{77} \text{ ---- W.} \\ 77 : 25 \dots : 20 : 6 \dots 5 \dots 3 \frac{5}{77} \text{ ---- X.} \\ 77 : 25 \dots : 15 : 4 \dots 10 \dots 1 \frac{5}{77} \text{ ---- Y.} \\ 77 : 25 \dots : 12 : 3 \dots 10 \dots 3 \frac{1}{77} \text{ ---- Z.} \end{array}} \right\} \text{ Ans. as before.}$$

(Key to Vol. I. page 123.)

Quest. 4. $\begin{matrix} s. & mo. & off. \\ 40 \times 6 \times 4 = & 960 & \text{for the officers.} \end{matrix}$

$\begin{matrix} s. & mo. & pet. off. \\ 30 \times 6 \times 12 = & 2160 & \text{for the midshipmen.} \end{matrix}$

$\begin{matrix} s. & mo. & seam. \\ 22 \times 3 \times 110 = & 7260 & \text{for the sailors.} \end{matrix}$

And $\begin{matrix} 960 \\ 2160 \\ 7260 \end{matrix} \left. \vphantom{\begin{matrix} 960 \\ 2160 \\ 7260 \end{matrix}} \right\} \text{are as } \begin{matrix} 16 \\ 36 \\ 121 \end{matrix}$

Sum 173

But

	\pounds	$s.$	$d.$	$qr.$	
As 173 : 1000 :: 16 :	£92	9	8	$2\frac{132}{1038}$	--- Officers.
173 : 1000 :: 36 :	208	1	10	$0\frac{816}{1038}$	--- Midshipmen.
173 : 1000 :: 121 :	699	8	5	$1\frac{90}{1038}$	--- Seamen.
	<u>£1000</u>	<u>0</u>	<u>0</u>	<u>0</u>	

And,

	\pounds	$s.$	$d.$	$qr.$	
4) 92..9.. 8..	$23\frac{2}{173}$	2	5	$0\frac{92}{173}$	for each Offr. } Ans.
12) 208..1..10..	$17\frac{6}{173}$	6	9	$3\frac{60}{173}$	---- Midsh. } Ans.
110) 699..8.. 5..	$6\frac{7}{173}$	7	2	$0\frac{8}{173}$	---- Seam. } Ans.

(Page 124.)

Quest. 5. $\begin{matrix} mo. \\ \pounds 1000 \times 12 = & \pounds 12000 & \text{---- H.} \\ 1500 \times 10 = & 15000 & \text{---- I.} \\ 2800 \times 7 = & 19600 & \text{---- K.} \end{matrix}$

Now $\begin{matrix} \pounds 12000 \\ 15000 \\ 19600 \end{matrix} \left. \vphantom{\begin{matrix} \pounds 12000 \\ 15000 \\ 19600 \end{matrix}} \right\} \text{are as } \begin{matrix} \pounds 60 \\ 75 \\ 98 \end{matrix}$

Sum £233

(Key to Vol. I. page 124.)

And
 As £233 : £1776 .. 10 : : £60 : £457 .. 9 .. 4 $\frac{1}{4}$ --H. }
 233 : 1776 .. 10 : : 75 : 571 .. 16 .. 8 $\frac{1}{4}$ --I. } Ans.*
 233 : 1776 .. 10 : : 98 : 747 .. 3 .. 11 $\frac{1}{4}$ --K. }

£1776 .. 9 .. 11 $\frac{3}{4}$

Of the farthing deficient, $\frac{23}{233}$ be- }
 long to H, $\frac{87}{233}$ to I, and $\frac{123}{233}$ to }
 K, amounting to - - - - - } $\frac{1}{4}$

£1776 .. 10 Proof.

* This is the Answer given by Dr. Hutton, but is not *strictly* correct.

Quest. 6. $(£20 \times 12) + (£20 \times 8) = \dots \dots \dots £400 \dots X.$
 $(£30 \times 12) + (£20 \times 9) + (£40 \times 7) = £820 \dots Y.$
 $(£60 \times 12) + (£10 \times 7) - (£30 \times 6) = £610 \dots Z.$

And $\left. \begin{matrix} £400 \\ 820 \\ 610 \end{matrix} \right\}$ are as $\left\{ \begin{matrix} £40 \\ 82 \\ 61 \end{matrix} \right.$

Sum £ 183

But || As £183 : £50 : : £40 : £10 .. 18 .. 6 .. 3 $\frac{4}{61}$ --- X.
 183 : 50 : : 82 : 22 .. 8 .. 1 .. 0 $\frac{12}{61}$ --- Y.
 183 : 50 : : 61 : 16 .. 13 .. 4 .. 0 --- Z.

£ 50 .. 0 .. 0

SIMPLE INTEREST.

(Page 126.)

Quest. 4. $\frac{£450 \times 5}{100} = \frac{£45}{2} = £22 .. 10. \text{ Ans.}$

(Key to Vol. I. page 126.)

$$\text{Quest. 5.} \quad (\text{£} \overset{s.}{715} \dots \overset{d.}{12} \dots 6) \times 4\frac{1}{2} = \text{£} \overset{s.}{32} \dots \overset{d.}{4} \dots 0\frac{3}{4} \text{ Ans.}$$

$$\frac{\text{£} 715 \dots 12 \dots 6}{100} \times 4\frac{1}{2} = \text{£} 32 \dots 4 \dots 0\frac{3}{4} \text{ Ans.}$$

$$\text{Quest. 6.} \quad \frac{\text{£} 720 \times 3}{20} = \text{£} 108. \text{ Ans.}$$

$$\text{Quest. 7.} \quad (\text{£} \overset{s.}{355} \dots \overset{d.}{15}) \times 4 = \text{£} \overset{s.}{56} \dots \overset{d.}{18} \dots 4\frac{3}{4}. \text{ Ans.}$$

$$\frac{(\text{£} 355 \dots 15)}{25} \times 4 = \text{£} 56 \dots 18 \dots 4\frac{3}{4}. \text{ Ans.}$$

(Page 127.)

$$\text{Quest. 8.} \quad (\text{£} \overset{s.}{32} \dots \overset{d.}{5} \dots 8) \times 4\frac{1}{4} \times 7 = \text{£} \overset{s.}{9} \dots \overset{d.}{12} \dots 1. \text{ Ans.}$$

$$\frac{(\text{£} 32 \dots 5 \dots 8)}{100} \times 4\frac{1}{4} \times 7 = \text{£} 9 \dots 12 \dots 1. \text{ Ans.}$$

$$\text{Quest. 9.} \quad \frac{\text{£} 170 \times 1\frac{1}{2}}{20} = \text{£} 12 \dots 15. \text{ Ans.}$$

$$\text{Quest. 10.} \quad \frac{\text{£} \overset{s.}{205} \dots \overset{d.}{15}}{100} = \text{£} \overset{s.}{2} \dots \overset{d.}{1} \dots 1\frac{3}{4}. \text{ Ans.}$$

$$\text{Quest. 11.} \quad (\text{£} \overset{s.}{319} \dots \overset{d.}{0} \dots 6) \times 5\frac{3}{4} \times 3\frac{3}{4} = \text{£} \overset{s.}{68} \dots \overset{d.}{15} \dots 9\frac{1}{2}. \text{ Ans.}$$

$$\frac{(\text{£} 319 \dots 0 \dots 6)}{100} \times 5\frac{3}{4} \times 3\frac{3}{4} = \text{£} 68 \dots 15 \dots 9\frac{1}{2}. \text{ Ans.}$$

$$\text{Quest. 12.} \quad \frac{\text{£} 107 \times 4\frac{3}{4} \times 117}{365 \times 100} = \text{£} \overset{s.}{1} \dots \overset{d.}{12} \dots 7. \text{ Ans.}$$

$$\text{Quest. 13.} \quad \frac{\text{£} (\overset{s.}{17} \dots \overset{d.}{5}) \times 4\frac{3}{4} \times 117}{365 \times 100} = \text{£} \overset{s.}{0} \dots \overset{d.}{5} \dots 3. \text{ Ans.}$$

(Key to Vol. I. page 127.)

Quest. 14. $7\frac{1}{2}$ per cent. per ann. being at the rate of 5 per cent. for 8 months, *the time given*, it will be

$$\frac{\text{£}712 \text{ .. } 6) \times 5}{100}, \text{ or, } \frac{\text{£}712 \text{ .. } 6}{20} = \text{£}35 \text{ .. } 12 \text{ .. } 3\frac{1}{2}. \text{ Ans.}$$

COMPOUND INTEREST.

(Page 128.)

Ex. 2.	20)	£50 ..	^{s.} 0 ..	^{d.} 0 ..	^{qr.} 0	PRINCIPAL.
		2 ..	10 ..	0 ..	0	First year's interest.
	20)	£52 ..	^{s.} 10 ..	^{d.} 0 ..	^{qr.} 0	Amount in one year.
		2 ..	12 ..	6 ..	0	Second year's interest.
	20)	£55 ..	^{s.} 2 ..	^{d.} 6 ..	^{qr.} 0	Amount in two years.
		2 ..	15 ..	1 ..	2	Third year's interest.
	20)	£57 ..	^{s.} 17 ..	^{d.} 7 ..	^{qr.} 2	Amount in three years.
		2 ..	17 ..	10 ..	$2\frac{3}{10}$	Fourth year's interest.
	20)	£60 ..	^{s.} 15 ..	^{d.} 6 ..	^{qr.} $0\frac{3}{10}$	Amount in four years.
		3 ..	0 ..	9 ..	$1\frac{43}{200}$	Fifth year's interest.
	Ans.	£63 ..	16 ..	3 ..	$1\frac{103}{200}$	AMOUNT TOTAL.

Ex. 3. $\text{£}1 + \frac{\text{£}1 \times 1}{40}^{\text{yr.}} = \text{£}1.025$ the amount of £1 in $\frac{1}{2}$ year.

And $(\text{£}1.025)^{10} = 1.28009$ Amount of £1 in 10 half-years.
by £50 PRINCIPAL.

Prod. $\text{£}64 \text{ .. } 0 \text{ .. } 1$ Amount required.

(Key to Vol. I. page 128.)

Ex. 4. $\text{£}1 + \frac{\text{£}1 \times 1^{\text{yr.}}}{80} = \text{£}1.0125$ the amount of $\text{£}1$ in $\frac{1}{4}$ year.

And $(\text{£}1.0125)^{20} = \text{£}1.28202$ nearly, for 20 quarters.
by $\text{£}50$ PRINCIPAL.

Prod. $\text{£}64..2..0..0.96$ Amount required.

Ex. 5. $\frac{\text{£}1 \times 4}{100} + \text{£}1 = \text{£}1.04$ the amount of $\text{£}1$ at 4 per cent. per ann. for 1 year.

And $(\text{£}1.04)^6 = \text{£}1.265319018496$ Amount of $\text{£}1$ in 6 years.
by $\text{£}370$ PRINCIPAL.

Prod. $\text{£}468.16803684252$ Amount Total.
Subtract 370 PRINCIPAL.

Rem. $\text{£}98..3..4..1.3153688192$ Int. reqd.

Ex. 6. $\text{£}1 + \frac{\text{£}1 \times 4\frac{1}{2}}{100} = \text{£}1.0225$ the amount of $\text{£}1$ at $4\frac{1}{2}$ per cent. per ann. for $\frac{1}{2}$ year.

And $(\text{£}1.0225)^5 = \text{£}1.117676$ Amount of $\text{£}1$ in $2\frac{1}{2}$ years.
by $\text{£}410$ PRINCIPAL.

Prod. $\text{£}458.2476$ Amount Total.
Subtract 410 PRINCIPAL.

Rem. $\text{£}48..4..11\frac{1}{4}$ The interest required.

☞ The two last Examples may be more concisely performed without Subtraction, if the *decimal part only*, in the amount of $\text{£}1$ for the time given, be *multiplied* by the PRINCIPAL.

(Key to Vol. I. page 128.)

Ex. 7. 5 per cent. per ann. payable quarterly, being $1\frac{1}{4}$ per cent. per quarter,

	s.	d.	qr.	
80) £217	0	0	0	PRINCIPAL.
1st qr's int.	2	14	3	
<hr/>				
80) £219	14	3	0	
2d qr's int.	2	14	11 0.55	
<hr/>				
80) £222	9	2	0.55	
3d qr's int.	2	15	7 1.506875	
<hr/>				
80) £225	4	9	2.056875	
4th qr's int.	2	16	3 2.8757109375	
<hr/>				
80) £228	1	1	0.9325859375	
5th qr's int.	2	17	0 0.66165732421875	
<hr/>				
80) £230	18	1	1.59424326171875	
6th qr's int.	2	17	8 2.869928040771484375	
<hr/>				
80) £233	15	10	0.464171302490234375	
7th qr's int.	2	18	5 1.5058021412811279296875	
<hr/>				
80) £236	14	3	1.9699734437713623046875	
8th qr's int.	2	19	2 0.57462466054714202880859375	
<hr/>				
80) £239	13	5	2.54459810431850433349609375	
9th qr's int.	2	19	11 0.081807476303981304168701171875	
<hr/>				
Amount } Total. }	£242	13	4 2.626405580622485637664794921875.	
<hr/>				

Otherwise,

The amount of £1 for a quarter of a year, at the rate given, is £1.0125. And

$(£1.0125)^9 = £1.118292178510262147076129913330078125$
multiply by the Principal - - - - - £217

Amt. Total £242 13 4 2.626405580622485637664794921875.

(Key to Vol. I. page 130.)

ALLIGATION.

ALLIGATION MEDIAL.

Quest. 2.

lb.	s.	s.	d.		s.
5	×	7	=	35	.. 0 value of Tea at 7.
9	×	8 $\frac{1}{2}$	=	76	.. 6 - - - - - 8. d.
14 $\frac{1}{2}$	×	5 $\frac{5}{6}$	=	84	.. 7 - - - - - 5 .. 10.
					s. d.
Divide by				28 $\frac{1}{2}$ - - -)	196 .. 1 (6 .. 10 $\frac{1}{2}$ nearly, per lb.

Quest. 3.

gal.	s.	d.	s.	d.	s.	d.
4	×	(4 .. 10)	=	19	.. 4 value of Wine at 4 .. 10.	
7	×	(5 .. 3)	=	36	.. 9 - - - - - 5 .. 3.	
9 $\frac{3}{4}$	×	(5 .. 8)	=	55	.. 3 - - - - - 5 .. 8.	
					s. d.	
Divide by				20 $\frac{3}{4}$ - - - - -)	111 .. 4 (5 .. 4 $\frac{1}{2}$ Ans.	

Quest. 4.

s.	d.	bu.	s.	d.	s.	d.
(3 .. 5)	×	3	=	10	.. 3 value of Flour at 3 .. 5.	
(5 .. 6)	×	4	=	22	.. 0 - - - - - 5 .. 6.	
(4 .. 8)	×	5	=	23	.. 4 - - - - - 4 .. 8.	
					s. d. qr.	
Divide by				12) 55 .. 7 (4 .. 7 .. 2 $\frac{1}{3}$ Ans.		

Quest. 5.

s.	bu.	s.	bu.	s.	bu.	s.			
5	×	10	+ 3	×	18	+ 2	×	20	= 3. Ans.
48 bu.									

Quest. 6.

car.	oz.	car.	oz.	car.	oz.				
22	×	7	+ 21	×	12 $\frac{1}{2}$	+ 19	×	17	= 20 $\frac{19}{73}$ carats fine. Ans.
36 $\frac{1}{2}$ oz.									

Quest. 7.

lb.	oz.	lb.	oz.	oz.		
3	×	9	+ (5 .. 8)	×	10	= 7 $\frac{61}{63}$ ounces fine. Ans.
lb.	lb.	oz.	lb.	oz.		
3	+ 5	.. 8	+ 1	.. 10		

(Key to Vol. I. page 132.)

ALLIGATION ALTERNATE.

RULE I.

Quest. 2. $5 \left\{ \begin{array}{l} 6 \\ 4 \end{array} \right\}$ equal quantities. Ans.

Quest. 3. $d. \left\{ \begin{array}{l} 4 \\ 6 \\ 11 \end{array} \right\} \begin{array}{l} - - 4 \\ - - 4 \\ - - 3+1=4 \end{array} \begin{array}{l} lb. \\ lb. \\ \end{array} \right\}$ Therefore equal quantities.

Quest. 4. $46 \left\{ \begin{array}{l} d. \left(\begin{array}{l} 30 \\ 44 \\ 48 \\ 56 \end{array} \right) \begin{array}{l} - - 2 \text{ at } 2 \dots 6 \\ - - 10 \text{ at } 3 \dots 8 \\ - - 16 \text{ at } 4 \dots 0 \\ - - 2 \text{ at } 4 \dots 8 \end{array} \end{array} \right\}$ or, $46 \left\{ \begin{array}{l} d. \left(\begin{array}{l} 30 \\ 44 \\ 48 \\ 56 \end{array} \right) \begin{array}{l} - 10 \text{ at } 2 \dots 6 \\ - 2 \text{ at } 3 \dots 8 \\ - 2 \text{ at } 4 \dots 0 \\ - 16 \text{ at } 4 \dots 8 \end{array} \end{array} \right\}$

Or,

$46 \left\{ \begin{array}{l} d. \left(\begin{array}{l} 30 \\ 44 \\ 48 \\ 56 \end{array} \right) \begin{array}{l} - - 2+10=12 \\ - - 10+ 2=12 \\ - - 16+ 2=18 \\ - - 2+16=18 \end{array} \end{array} \right\}$ which are as $\left\{ \begin{array}{l} bu. \quad s. \quad d. \\ \left(\begin{array}{l} 2 \text{ at } 2 \dots 6 \\ 2 \text{ at } 3 \dots 8 \\ 3 \text{ at } 4 \dots 0 \\ 3 \text{ at } 4 \dots 8 \end{array} \right) \end{array} \right\}$

Quest. 5. $c. \text{ fine} \left\{ \begin{array}{l} 16 \\ 18 \\ 23 \\ 24 \end{array} \right\} \begin{array}{l} - - 3 \text{ of } 16 \\ - - 2 \text{ of } 18 \\ - - 3 \text{ of } 23 \\ - - 5 \text{ of } 24 \text{ or pure.} \end{array} \right\}$ c. fine.

Quest. 6. $s. \left\{ \begin{array}{l} 12 \\ 10 \\ 1 \\ 0 \end{array} \right\} \begin{array}{l} - - - 8 - - - - \text{Brandy} \\ - - - 7 - - - - \text{Wine} \\ - - - 2 - - - - \text{Cider} \\ - - - 4 - - - - \text{Water} \end{array} \right\}$ gallons.

(Key to Vol. I. page 133.)

RULE II.

Quest. 2.

	d.	gal.	gal.	gal.	gal.	gal.	gal.	s.	d.													
64	}	d.	48	--	8	+	2	=	10		As	60	:	10	::	18	:	3	at	4	..	0
		60	--	2	+	8	=	10		60	:	10	::	18	:	3	at	5	..	0		
		66	--	4	+	16	=	20		60	:	20	::	18	:	6	at	5	..	6		
		72	--	16	+	4	=	20		60	:	20	::	18	:	6	at	6	..	0		
					Sum				60									18				

} Ans.

RULE III.

(Page 134.)

Quest. 2.

	s.	lb.	lb.	lb.	lb.	s.										
8	}	s.	12	--	4		As	4	:	20	::	4	:	20	at	12
		10	--	2		4	:	20	::	2	:	10	at	10		
		6	--	2		4	:	20	::	2	:	10	at	6		
		4	--	4		4	:	20	::	4	:	20	at	4		

} Ans.

Quest. 3.

	c. fine	oz.	oz.	oz.	oz.	oz.	c. fine.												
20	}	c. fine	15	-----	2		As	2	:	5	::	2	:	5	of	15			
		17	-----	2		2	:	5	::	2	:	5	of	17					
		22	5	+	3	+	2	=	10		2	:	5	::	10	:	25	of	22
		18	-----	2		2	:	5	::	2	:	5	of	18					

} Ans.

SINGLE POSITION.

(Page 136.)

Quest. 2. Suppose 12.

$$\frac{12 \times 7}{6} = 14. \quad \text{And, As } 14 : 12 :: 21 : 18. \quad \text{Ans.}$$

Quest. 3. Suppose 24.

$$24 + 12 + 8 + 6 = 50. \quad \text{And, As } 50 : 24 :: 75 : 36. \quad \text{Ans.}$$

(Key to Vol. I. page 139.)

21 First supposition.
18 Second supposition.

3 *diff.* of the assumed numbers.
Multiply by 6 greater error.

Divide by the *diff.* of the errors. $3)18(6$ to be taken from the greater
supposition.

Wherefore $\overset{\text{yrs.}}{15}$ and $\overset{\text{yrs.}}{45}$ Ans.

	work da. id. da.	work da. id. da.
Quest. 3. Ist. Suppose	15 and 5 <u>3</u> <u>1</u>	2dly. Suppose 18 and 2 <u>3</u> <u>1</u>
	$45s. - 5s. = 40s.$	$54s. - 2s. = 52$ <i>s.</i>

With $\overset{s.}{£2}..4$ given. Compare $2..0$ First result.	With $\overset{s.}{£2}..4$ given. Compare $2..12$ Second result.
<u>4 too little.</u>	<u>8 too much.</u>

Now $18 - 15 = 3$ *diff.* of the assumed numbers; but

Sum of the errors $\frac{3 \times 8}{12} = 2$ ^{da.} to be taken from 18. ^{da.}

Consequently, 16 the number of working days }
and, 4 the number of idle days. } Ans.

Quest. 4. Ist. Suppose A.	$\overset{gs.}{70}$ and B. $\overset{gs.}{70}$ won 20 lost 20 <u> </u> <u> </u>
End of the 1st Game	90 50
$\frac{2}{3}$ of 90 lost.	60 won 60 <u> </u> <u> </u>
End of their Play	30 110 <u> </u> <u> </u>

(Key to Vol. I. page 139.)

	2dly. Suppose A. ^{gs.} 130	and B. ^{gs.} 130
	won 20	lost 20
	End of the 1st Game. 150	110
$\frac{2}{3}$ of 150 lost.	100	won 100
	End of their Play. 50	210

^{gs.} With $30 \times 4 = 120$	^{gs.} With $50 \times 4 = 200$
Compare 110 First result.	Compare 210 Second result.
10 too few.	10 too many.

From which it is evident, that *half the Sum* of the two Suppositions, or, $\frac{70 + 130}{2} = 100$. Ans.

Quest. 5. 1st. Suppose £100 the income.
1st error is £ 20 too much.

2dly. Suppose £120 the income.
2nd error is £4 too much.

Difference of the errors is 16.

£ 120 second supposition.	£100 first supposition.
20 first error.	4 second error.
£2400 Prod.	£400 Prod.

And, $\frac{£2400 + 400}{\text{Diff. of the err. } 16} = £125$, the INCOME required.

Lastly, $£125 - \frac{£125}{5} = £100$ per ann. expended by A. and

$£125 + \frac{£125}{5} = £150$ per ann. expended by B.

(Key to Vol. I. page 140.)

PRACTICAL QUESTIONS IN ARITHMETIC.

Quest. 1. $\frac{\text{miles} \quad \text{feet}}{100000000 \times 5280} = 264000000''$ the time of flight.

And $365 \text{ days} \dots 6 \text{ hrs.} = 31557600''$, wherefore

$$\frac{264000000''}{31557600} = 8\frac{4808}{13149} \text{ years. Ans.}$$

Quest. 2. $\frac{\text{miles} \quad \text{feet}}{100000000 \times 5280} = 5866666\dot{6}$ time of the ball.

Hence $782222\frac{2}{9}$ ($= \frac{5866666\dot{6}}{7\frac{1}{2}}$) to 1, ratio required.

Quest. 3. $\frac{\text{pa.} \quad \text{in.}}{70 \times 28 \times 60'} = 1\frac{1}{13\frac{1}{2}}$ mile. Ans.

Quest. 4. $\frac{\text{pa.} \quad \text{in.}}{120 \times 28 \times 60'} = 3\frac{2}{11}$ miles per hour. Ans.

And $20 \div 3\frac{1}{2} = 6\frac{2}{7}$ the hours necessary for march.

Also $6\frac{2}{7} + 1 = 7 \dots 17\frac{1}{7}$ Ans.

Quest. 5. $700 \text{ yds.} - 220 \text{ yds.} = 480$. And $29 \text{ da.} - 11 \text{ da.} = 18$.

Now, As $220 : 480 :: 12 \times 11 = 132 : 288$, men involved in days.

But $\frac{288}{18} = 16$ men, the numb. that must work.

Lastly, $16 - 12 = 4$ additional required.

Quest. 6. $\frac{\text{inches.} \quad \text{miles.} \quad \text{yds.} \quad \text{ft.} \quad \text{in.}}{12 \times 3 \times 5\frac{1}{2} \times 40 \times 8} = 7891 \dots 728 \dots 2 \dots 8$. Ans.

(Key to Vol. I. page 140.)

Quest. 7. $\begin{matrix} \text{yds.} & \text{yds.} \\ 11 \times 3' = 33 & \text{travelled by A in } 3', \text{ that is, every } 34 \\ \text{yards of B gain 1 yard on A; now half the circuit of the} \\ \text{wood} = 268 \text{ yards, therefore} \end{matrix}$

$\begin{matrix} \text{yd.} & \text{yds.} & \text{yds.} & \text{yds.} \\ \text{As } 1 : 34 :: 268 : 9112 & \text{travelled by B when he overtakes} \end{matrix}$

A. But $\frac{9112}{2 \times 34} = 17$ revolutions made by B, and consequently $16\frac{1}{2}$ by A.

(Page 141.)

Quest. 8. It is manifest that $\frac{1}{12}$ of the work = A's daily labour; and that $\frac{1}{14}$ of it = the daily labour of B.

But $\frac{1}{12} + \frac{1}{14} = \frac{26}{168}$ parts of the work performed in a day when both work.

Again, $\frac{26}{168} \times 1 = 6\frac{2}{3}$ days. Ans.

Or, more clearly, As $\frac{26}{168}$, that is, $\frac{13}{84}$: whole work = Unity or 1 :: 1 day : $6\frac{2}{3}$ days.

Quest 9. $\frac{3}{4}$ of $\frac{3}{5} = \frac{9}{20}$.

And, $\frac{9}{20} \times 1800 = \pounds 4000$. Ans.

Quest. 10.

1st. Suppose $\pounds 100$
 $\pounds 25 + \pounds 20 = 45$

—
 $\pounds 55$

80

1st error $\pounds 25$ too little.

2dly. Suppose $\pounds 400$

$\pounds 100 + \pounds 20 = 120$

—
 $\pounds 280$

230

should be

2d error $\pounds 50$ too much.

Wherefore, $\frac{1 \text{ sup. } 2 \text{d err. } 2 \text{d sup. } 1 \text{st error.}}{\text{Sum of the errors, } 25 + 50} = \pounds 200$. Ans.

Quest. 11. Because the hands appear in conjunction 11 times in 12 hours, and that their respective velocities are always in the same ratio, it follows that,

$\begin{matrix} \text{hr.} & \text{hr.} & \text{hr.} & \text{min.} & \text{sec.} \\ \text{As } 11 : 12 :: 1 : 1\frac{1}{11} = 1 \dots 5 \dots 27\frac{3}{11} \end{matrix}$ Ans.

(Key to Vol. I. page 141.)

Quest. 12. $\overset{\text{we.}}{\pounds 21} \times 52 = \pounds 1092.$
 And $\pounds 1500 - \pounds 1092 = \pounds 408$, *saved.* Ans.

Quest. 13. $\overset{\text{oranges.}}{180} \div 2 = 90$
 $\overset{d.}{180} \div 3 = 60$

 5) $\overset{360}{\quad}$ 150 prime cost.
 $72 \times 2 = 144$ sold for.

 6 pence *loss.* Ans.

Quest. 14. $\overset{\text{we.}}{\text{As } 20} \times \overset{\text{oz.}}{8} : \overset{\text{we.}}{12} \times \overset{\text{oz.}}{20} :: \overset{\text{men.}}{1500} : \overset{\text{men.}}{2250}$. Ans.

Quest. 15. $\frac{15550 \text{ miles}}{23 \text{ hrs. } 56'} = 649 \frac{259}{359}$ miles per hour. Ans.

Quest. 16. $\frac{1}{4} + \frac{3}{7}$ of $\frac{3}{4} = \frac{4}{7}$. And $1 - \frac{4}{7} = \frac{3}{7}$, *remaining.*
 But $\frac{3}{7}$ $\overset{s.}{\pounds} 820$ ($\overset{d.}{\pounds} 1913 \text{ .. } 6 \text{ .. } 8$). Ans.

Quest. 17. $\overset{\text{men}}{\text{As } 1500} \times \overset{\text{we.}}{8} : \overset{\text{men}}{1000} \times \overset{\text{we.}}{5} :: \overset{\text{oz.}}{16} : \overset{\text{oz.}}{6\frac{2}{3}}$. Ans.

Quest. 18. $\text{As } 7 : 9 :: \pounds 8400 : \pounds 10800$ *elder brother's FORTUNE.*
 And $\pounds 10800 + \pounds 8400 = \pounds 19200$. Ans.

(Page 142.)

Quest. 19. $\overset{\text{hrs.}}{\text{As } 11} : \overset{\text{hrs.}}{12} :: \overset{\text{hrs.}}{5} : \overset{\text{hrs.}}{5\frac{5}{11}} = 5 \text{ .. } 27 \text{ .. } 16\frac{4}{11}$. Ans.

Quest. 20. $3 \times \overset{\text{men}}{5} \times \overset{\text{men}}{20} = 300$. Ans.

For it is evident that, if the work be three times as great, and the time but one fifth part the former, fifteen times as many men will be required.

(Key to Vol. I. page 142.)

Quest. 21. $1 - \frac{7}{18} = \frac{11}{18}$ the residue after the elder son's fortune.

And $\frac{7}{18}$ of $\frac{11}{18} = \frac{77}{324}$ the younger son's patrimony.

Now $\frac{11}{18} = \frac{126}{324}$, and $\frac{126 - 77}{324} = \frac{49}{324}$ the difference of the son's portions. = £514 $\frac{1}{3}$ by the question.

Also $\frac{126 + 77}{324} = \frac{203}{324}$ the sum of both sons' property; there-

fore $1 - \frac{203}{324} = \frac{121}{324}$ the provision made for the RELICT.

shares shares s. d.
And As 49 : 121 :: £514 $\frac{1}{3}$: £1270 .. 1 .. 9 $\frac{1}{3}$. Ans.

Quest. 22. $1 - \frac{13}{20} = \frac{7}{20}$ of the estate, for the younger child.

But $\frac{13 - 7}{20} = \frac{6}{20} = \frac{3}{10}$, therefore,

As 3 : 10 :: £1200 : £4000. Ans.

Quest. 23. $1\frac{1}{2} \times 7 = 10\frac{1}{2}$ A. rode more than B. And

$\frac{10\frac{1}{2}}{2} = 5\frac{1}{4}$. Also $\frac{100}{2} = 50$.

But $50 + 5\frac{1}{4} = 55\frac{1}{4}$ for A. And
 $50 - 5\frac{1}{4} = 44\frac{3}{4}$ for B.

Lastly $\frac{55\frac{1}{4}}{7} = 7\frac{5}{8} =$ A's rate per hour.

And $\frac{44\frac{3}{4}}{7} = 6\frac{1}{2}$ B's rate per hour.

} Ans.

(Key to Vol. I. page 142.)

Quest. 24. $\begin{matrix} \text{h.} & \text{h.} \\ \text{From 8. A. M. to 4. P. M. there are 8 hours, which} \\ \times^{\text{ed}} \text{ by 3, give 24 miles walked by A, when B sets out from} \\ \text{London.} \end{matrix}$

$\begin{matrix} \text{miles} & \text{miles} & \text{miles} & \text{miles} & \text{miles} & \text{miles} \\ \text{Also } 130 - 24 = 106; \text{ And } 4 + 3 = 7 \text{ per hour.} \end{matrix}$

$\begin{matrix} \text{miles} \\ \text{But } \frac{106}{7} = 15\frac{1}{7} \text{ hours, the time of B. on the road.} \end{matrix}$

$\begin{matrix} \text{hrs.} & \text{miles} & \text{miles} & \text{miles} \\ \text{Again } (15\frac{1}{7} \times 3) + 24 = 69\frac{3}{7}, \text{ A's distance from Exeter when he} \\ \text{meets B, consequently } 60\frac{4}{7} \text{ miles from London.} \end{matrix}$

Quest. 25. $\begin{matrix} \text{yds.} & \text{yd.} & & \text{yds.} & \text{miles} & \text{yds.} \\ (100 - 1) \times 50 \times 2 = 10100 = 5 \text{ .. } 1300. \text{ Ans.} \end{matrix}$

Quest. 26. $\begin{matrix} \text{str.} & \text{str.} & \text{hrs.} & \text{str.} \\ (24 + 1) \times 12 = 300. \text{ Ans.}^* \end{matrix}$

* See this Question, page 113, line 1, vol. i. of Hutton's Course.

Quest. 27. $\begin{matrix} (1 \times 2^{63} \times 2) - 1 & \text{quarters} & \text{bu.} & \text{p.} & \text{gal.} \\ \hline = 4691249611844 \text{ .. } 2 \text{ .. } 0 \text{ .. } 1 \\ (2 - 1) \times 7680 \times 512 \text{ pints in 1 qr.} \end{matrix}$

$\begin{matrix} \text{qrs.} & \text{bu.} & \text{p.} & \text{gal.} & \text{s.} & \text{d.} \\ \text{And } 4691249611844 \text{ .. } 2 \text{ .. } 0 \text{ .. } 1 \times 27 \text{ .. } 6 = \end{matrix}$

$\begin{matrix} \text{s.} & \text{d.} & \text{qr.} \\ \text{£}6450468216285 \text{ .. } 17 \text{ .. } 3 \text{ .. } 3\frac{2757}{2768}. \text{ Ans.} \end{matrix}$

(Page 143.)

Quest. 28.

1st. Suppose £400	2dly. Suppose £464
End of 1st year it is 600	End of 1st year it is 680
2d year - - - 850	2d year - - - 950
3d year - - 1162·5	3d year - - 1287·5
4th year - 1553·125	4th year - - 1709·375
10342·1875	should be 10342·1875

1st error £ 8789·0625 *too little.* 2d err. £8632·8125 *too little.*

The *difference* of the errors is £156·25

The *difference* of the products arising is £625000.

But $\frac{£625000}{£156\cdot25} = £4000. \text{ Ans.}$

(Key to Vol. I. page 143.)

Quest. 29. ($\overset{s.}{£1012} .. 10$) - $\overset{s.}{£750} = \overset{s.}{£262} .. 10$ the amount of *int.*

And $\frac{\overset{s.}{£262} .. 10}{7 \text{ yrs.}} = \overset{s.}{£37} .. 10$ the *yearly* interest.

Lastly $\frac{(\overset{s.}{£37} .. 10) \times \overset{£}{100}}{\overset{£}{750}} = \overset{£}{5}$ per cent. Ans.

Quest. 30.

1st. Suppose A. $\overset{£}{400}$ 2dly. Suppose A. $\overset{£}{600}$

P. Quest. $\left\{ \begin{array}{l} \text{B. } 185 \\ \text{C. } 280 \end{array} \right.$ P. Quest. $\left\{ \begin{array}{l} \text{B. } 385 \\ \text{C. } 480 \end{array} \right.$

$\overset{£}{865}$ $\overset{£}{1465}$
 $\overset{£}{1000}$ should be $\overset{£}{1000}$

1st error $\overset{£}{135}$ *too little.* 2d error $\overset{£}{465}$ *too much.*

And $\frac{\overset{£}{135} \times \overset{£}{600} + \overset{£}{465} \times \overset{£}{400}}{\overset{£}{135} + \overset{£}{465}} = \overset{£}{445} = \text{A's dividend.}$

Wherefore $\overset{£}{445}$ for A. $\overset{£}{230}$ for B. and $\overset{£}{325}$ for C. Ans.

Quest. 31. $4 + 5 = 9$, and $\overset{\text{hrs.}}{12} \overset{\text{h.}}{9} = 1\frac{1}{3}$ one of the fifths of the time till midnight.

But $\overset{\text{hr.}}{1\frac{1}{3}} \times 4 = \overset{\text{hrs.}}{5\frac{1}{3}} = 5 .. 20$, consequently 20 *min. past 5.* Ans.

Quest. 32. $\frac{2}{5}$ of $\frac{4}{9}$ of $\frac{3}{16} = \frac{1}{30} = \frac{8}{240}$, and $\frac{3}{16} = \frac{45}{240}$.

But $\frac{45 - 8}{240} = \frac{37}{240}$ left unsold.

Also $\overset{£}{1200} \times \frac{37}{240} = \overset{£}{185}$ the worth of $\frac{37}{240}$ of the ship. Ans.

(Key to Vol. I. page 143.)

<p>Quest. 33.</p> <table style="margin-left: 20px;"> <tr> <td style="text-align: right;">ac.</td> <td></td> </tr> <tr> <td style="text-align: right;">Ist. Suppose A.</td> <td style="text-align: right;">200</td> </tr> <tr> <td style="text-align: right;">Per Quest. } B.</td> <td style="text-align: right;">300</td> </tr> <tr> <td style="text-align: right;"> } C.</td> <td style="text-align: right;">364</td> </tr> <tr> <td></td> <td style="text-align: right;"><hr style="width: 50px; margin-left: 0;"/></td> </tr> <tr> <td></td> <td style="text-align: right;">864</td> </tr> <tr> <td></td> <td style="text-align: right;">1200</td> </tr> <tr> <td></td> <td style="text-align: right;"><hr style="width: 50px; margin-left: 0;"/></td> </tr> <tr> <td style="text-align: right;">Ist error</td> <td style="text-align: right;">336 <i>too little.</i></td> </tr> </table>	ac.		Ist. Suppose A.	200	Per Quest. } B.	300	} C.	364		<hr style="width: 50px; margin-left: 0;"/>		864		1200		<hr style="width: 50px; margin-left: 0;"/>	Ist error	336 <i>too little.</i>	<table style="margin-left: 20px;"> <tr> <td style="text-align: right;">ac.</td> <td></td> </tr> <tr> <td style="text-align: right;">2dly. Suppose A.</td> <td style="text-align: right;">400</td> </tr> <tr> <td style="text-align: right;">Per Quest. } B.</td> <td style="text-align: right;">500</td> </tr> <tr> <td style="text-align: right;"> } C.</td> <td style="text-align: right;">564</td> </tr> <tr> <td></td> <td style="text-align: right;"><hr style="width: 50px; margin-left: 0;"/></td> </tr> <tr> <td></td> <td style="text-align: right;">1464</td> </tr> <tr> <td></td> <td style="text-align: right;">1200</td> </tr> <tr> <td></td> <td style="text-align: right;"><hr style="width: 50px; margin-left: 0;"/></td> </tr> <tr> <td style="text-align: right;">2d error</td> <td style="text-align: right;">264 <i>too much.</i></td> </tr> </table>	ac.		2dly. Suppose A.	400	Per Quest. } B.	500	} C.	564		<hr style="width: 50px; margin-left: 0;"/>		1464		1200		<hr style="width: 50px; margin-left: 0;"/>	2d error	264 <i>too much.</i>
ac.																																					
Ist. Suppose A.	200																																				
Per Quest. } B.	300																																				
} C.	364																																				
	<hr style="width: 50px; margin-left: 0;"/>																																				
	864																																				
	1200																																				
	<hr style="width: 50px; margin-left: 0;"/>																																				
Ist error	336 <i>too little.</i>																																				
ac.																																					
2dly. Suppose A.	400																																				
Per Quest. } B.	500																																				
} C.	564																																				
	<hr style="width: 50px; margin-left: 0;"/>																																				
	1464																																				
	1200																																				
	<hr style="width: 50px; margin-left: 0;"/>																																				
2d error	264 <i>too much.</i>																																				

$$\text{And } \frac{\begin{array}{c} \text{ac.} \\ 336 \end{array} \times \begin{array}{c} \text{ac.} \\ 400 \end{array} + \begin{array}{c} \text{ac.} \\ 264 \end{array} \times \begin{array}{c} \text{ac.} \\ 200 \end{array}}{\begin{array}{c} \text{ac.} \\ 264 \end{array} + \begin{array}{c} \text{ac.} \\ 336 \end{array}} = 312. \text{ A's share.}$$

Wherefore 312 acres for A. 412 acres for B. and 476 acres for C. Ans.

Quest. 34. $\frac{2}{7}$ of $\frac{3}{8} = \frac{9}{84}$. And $\frac{8}{15}$ of $\frac{5}{16} = \frac{14}{84}$.

But $10\frac{9}{84} - \frac{14}{84} = 9\frac{79}{84}$. Ans.

Quest. 35. $\frac{2}{3}$ of $\frac{4}{5}$ of $1\frac{1}{2} = \frac{4}{5}$. Also $1 \div \frac{4}{5} = \frac{5}{4}$.

And $\frac{5}{4} \times \frac{5}{4} = \frac{25}{16} = 1\frac{9}{16}$. Ans.

Quest. 36. As $8\frac{1}{2} : 12 :: 12 : 16\frac{1}{2}$. Ans.

Quest. 37. As $12 : 15 :: \text{£}5 : \text{£}6\frac{1}{4}$.

And, As $\text{£}100 + \text{£}6\frac{1}{4} : \text{£}100 :: \text{£}138\frac{1}{8} : \text{£}130$. Ans.

Quest. 38. A. £4. B. £3 } Per Quest.
A. 8. B. 6 C. £5 }
But $\text{£}8 + \text{£}6 + \text{£}5 = \text{£}19$, therefore
 $\frac{8}{19}$, $\frac{6}{19}$, and $\frac{5}{19}$, of the fortune, are the three shares.

Now $\frac{8}{19}$) £4000 A's share (£9500. Ans.

Or, more clearly, As $8 : 19 :: \text{£}4000 : \text{£}9500$ the whole FORTUNE.

(Key to Vol. I. page 144.)

Quest. 39. As $60' : \frac{2'}{3} :: 1760 \times 10 : \frac{1}{9}$ of $1760 = 195\frac{5}{9}$ the hare had increased her distance from the dog before she was observed.

And $18 - 10 = 8$ miles per hour, the excess of the dog's speed above that of the hare. Wherefore,

As $1760 \times 8 : 40 + 195\frac{5}{9} :: 60' : 1' .. 0''\frac{5}{22}$ time of the course.

Also As $60' : 1' .. 0''\frac{5}{22} = 1'\frac{1}{264} :: 1760 : 530$ run by the Dog.

Quest. 40. Prudent Officer's annual state.

	£.	s.	d.	qr.	years.				
Divide by 20 and add, yearly.	10	0	0		1st.				
	20	10	0		2d.				
	31	10	6		3d.				
	43	2	0	1,2	4th.				
	55	5	1	2,06	5th.				
	68	0	4	2,36	6th.				
	81	8	4	3,28	7th.				
	95	9	9	3,44	8th.				
	110	5	3	3,01	9th.				
	125	15	6	3,76	10th.				
	142	1	4	1,15	11th.				
	159	3	5	0,41	12th.				
	177	2	7	0,63	13th.				
	195	19	8	2,86	14th.				
	215	15	8	2,21	15th.				
	236	11	5	3,92	16th.				
	258	8	0	3,51	17th.				
	281	6	5	2,89	18th.				
	305	7	9	0,43	19th.				
	330	13	1	3,05	20th.				
	357	3	9	2,61	21st.				
	385	0	11	3,74	22d.				
	414	6	0	2,13	23d.				
	445	0	4	0,43	24th.				
	477	5	4	1,25	25th.				
	511	2	7	2,12	26th.				
	546	13	9	0,42	27th.				
	584	0	5	1,44	28th.				
	623	4	6	2,52	29th.				
	664	7	9	1,44	30th.				
£400	0	0	0	Annuity	707	12	2	0,31	31st.
37	12	11	3,47 = int. of	752	19	9	1,53	32d.	

Sum £437 12 11 3,47 to spend per ann. Ans.

(Key to Vol. I. page 144.)

Prodigal Officer's annual state.

	£.	s.	d.	qr.	years.
$\text{£}10 + \text{£}1..10 + \text{£}0..13..9..2,4 \text{ instr.}$	= 12	3	9	2,4	1st.
$\text{£}0..12..2..1,12 + \text{£}11..10 + \text{ins.}$	= 25	15	1	3,17	2d.
$\text{£}1..9..1..3,65$					
(In which manner proceed <i>annually</i> .)	40	17	1	3,74	3d.
	57	13	3	2,41	4th.
	76	7	5	0,28	5th.
	97	3	9	3,42	6th.
	120	7	3	1,11	7th.
	146	3	1	0,56	8th.
	174	17	2	1,82	9th.
	206	16	2	1,06	10th.
	242	7	4	2,15	11th.
	281	18	11	0,73	12th.
	325	19	11	0,69	13th.
	375	0	5	3,20	14th.
	429	11	10	0,22	15th.
	490	6	6	0,75	16th.
	557	18	5	1,35	17th.
	633	3	1	3,11	18th.
	716	17	10	1,90	19th.
	810	1	10	0,92	20th.
	913	16	5	2,17	21st.
	1029	5	10	3,30	22d.
	1157	15	10	3,57	23d.
	1300	16	3	3,61	24th.
	1459	19	11	2,56	25th.
	1637	3	4	1,60	26th.
	1834	7	2	0,02	27th.
	2053	16	7	1,60	28th.
	2168	0	4	3,88	29th.
	2424	10	1	2,78	30th.
$\text{£}400..0..0..0 \text{ Annuity}$	2709	19	7	1,09	31st.
$333..0..11..1,53 = 11 \text{ per ct. on}$	3027	14	0	3,76	32d.

Diff. £ 66..19..0..2,46 to spend per ann. Ans.

END OF THE ARITHMETIC.

LOGARITHMS.

(Key to Vol. I. page 160.)

	Numbers.	Logarithms.
Ex. 7.	3·1416 - - - - -	0·497151
	82· - - - - -	1·913814
	{ 73 its log. ·863323 } diff. - - -	0·250539
	{ 41 its log. ·612784 }	
	<u>Prod. 458·6736 - - - - -</u>	<u>2·661504</u>

	Numbers.	Logarithms.
Ex. 8.	·02196 - - - - -	-2·4647875
	751·3 - - - - -	2·8758134
	{ 6 - - - - -	0·7781513 }
	{ 941 log. arithmetical comp. -	-3·0264104 }
	<u>Prod. ·1396891 - - - - -</u>	<u>-1·1451626</u>

	Numbers.	Logarithms.
Ex. 9.	As 7241 log. arithmetical comp. -	4·1402015
	: 3·58 - - - - -	0·5538830
	:: 20·46 - - - - -	1·3109056
	<u>: ·01011556 - - - - -</u>	<u>-2·0049901</u>

	Numbers.	Logarithms.
Ex. 10	As $\sqrt{724}$. log. arithmetical comp. -	2·5701307
	: $\sqrt{\frac{58}{13}}$. - - - - -	0·3247423
	:: 6·927. - - - - -	0·8405452
	<u>: ·54377375 - - - - -</u>	<u>-1·7354182</u>

ALGEBRA.

ADDITION OF ALGEBRA.

(Key to Vol. I. page 167.)

CASE I.

- | | |
|------------------------------------|---|
| Ex. 7. Ans. $69xy$. | Ex. 8. Ans. $-29y^2$. |
| Ex. 9. Ans. $28a - 20b$. | Ex. 10. Ans. $93 - 73\sqrt{x} - 20xy$. |
| Ex. 11. Ans. $21xy - 15x + 20ab$. | |
-

(Page 168.)

CASE II.

- | | |
|----------------------------|--|
| Ex. 4. Ans. $6a^2$. | Ex. 5. Ans. $-19b^2y^3$. |
| Ex. 6. Ans. $3ab - 5$. | Ex. 7. Ans. $3\sqrt{a^2x} = 3ax^{\frac{1}{2}}$. |
| Ex. 8. Ans. $-\sqrt{ax}$. | Ex. 9. Ans. $4y + 7ax^{\frac{1}{2}}$. |
-

(Page 169.)

CASE III.

- | | |
|--|--|
| Ex. 4. Ans. $9x^2y^2 - 11x^2y + 3axy$. | |
| Ex. 5. Ans. $25ax + x^2 + 3xy + 8y^2 + 26$. | |
| Ex. 6. Ans. $9\sqrt{ax} + 2x + 7\sqrt{xy} + 5y + 19$. | |
| Ex. 7. Ans. $-3x^2y - 3xy^2$. | |

(Key to Vol. I. page 169.)

- Ex. 8. Ans. $5\sqrt{xy} + 17x + 4\sqrt{x-y} - 9$.
- Ex. 9. Ans. $3a^2 + 3a + x^2 - 3x + \sqrt{x} - 2y - 4$.
- Ex. 10. Ans. $4a - 4b$.
- Ex. 11. Ans. $8a - 12x$.
- Ex. 12. Ans. $2x - 4a - b + 5$.
- Ex. 13. Ans. $10b - 3a + c$.
- Ex. 14. Ans. $2a$.
- Ex. 15. Ans. $-2b - 3c - d - 17$.
- Ex. 16. Ans. $b^2 + 2ab + bc - b - c$.
- Ex. 17. Ans. $a^3 + ab^2 - abc + b^2c$.
- Ex. 18. Ans. $12x + 6a - 4b - 12c + 40$.

SUBTRACTION OF ALGEBRA.

(Page 170.)

- Ex. 7. Ans. $10x^2y + 4$.
- Ex. 8. Ans. $-2\sqrt{xy} + 2x\sqrt{xy} + 2xy - 3$.
- Ex. 9. Ans. $-2x^2 + \sqrt{x} - 2b - 6$.

(Page 171.)

- Ex. 10. Ans. $-2xy + 20$.
- Ex. 11. Ans. $7x^3 - 2x^2 + 2a + 2b$.
- Ex. 12. Ans. $3xy^3 - 4x^2y^2 + 8a(xy + 10)^{\frac{1}{2}}$.
- Ex. 13. Ans. $2b$.
- Ex. 14. Ans. $3a + 3b$.
- Ex. 15. Ans. $a - 9b$.

(Key to Vol. I. page 171.)

- Ex. 16. Ans. $4a - 9x$.
 Ex. 17. Ans. $-4x - 5a + 3b - 3$.
 Ex. 18. Ans. $a + b - 10$.
 Ex. 19. Ans. $c - d$.
 Ex. 20. Ans. $-3a^2 + 2ab - 3c + bc - b$.
 Ex. 21. Ans. $a^3 + 3b^2c - b^2$.
 Ex. 22. Ans. $8x + 9a - 8b - 6d + 50$.
 Ex. 23. Ans. $x - 12a - 2b + 12c - 10$.
 Ex. 24. Ans. $10x + 14a - 8b - 7c$.

MULTIPLICATION OF ALGEBRA.

(Page 172.)

When both Factors are Simple Quantities.

- Ex. 9. Ans. $-12ax^2$. Ex. 10. Ans. $6acx$.
 Ex. 11. Ans. $-12xy$. Ex. 12. Ans. $25ax^2yz$.

 (Page 173.)
When one of the Factors is a Compound Quantity.

- Ex. 4. Ans. $48ax - 8a^2c$.
 Ex. 5. Ans. $-50ac + 14ab$.
 Ex. 6. Ans. $8abx - 2ab^2 + 6a^2b^2$.
 Ex. 7. Ans. $12c^2xy + 4x^2y$.
 Ex. 8. Ans. $-40x^4 + 12x^2y^2$.
 Ex. 9. Ans. $6a^3x^2 - 4ax^4 - 12abx^2$.

(Key to Vol. I. page 174.)

When both Factors are Compound Quantities.

$$\begin{array}{r} \text{Ex. 1. Mul. } 10ac \\ \text{by } 2a \\ \hline \text{Prod. } 20a^2c \\ \hline \end{array}$$

$$\begin{array}{r} \text{Ex. 2. Mul. } 3a^2 - 2b \\ \text{by } 3b \\ \hline \text{Prod. } 9a^2b - 6b^2 \\ \hline \end{array}$$

$$\begin{array}{r} \text{Ex. 3. Mul. } 3a + 2b \\ \text{by } 3a - 2b \\ \hline \text{Prod. } 9a^2 - 4b^2 \\ \hline \end{array}$$

$$\begin{array}{r} \text{Ex. 4. Mul. } x^2 - xy + y^2 \\ \text{by } x + y \\ \hline \text{Prod. } x^3 + y^3 \\ \hline \end{array}$$

$$\begin{array}{r} \text{Ex. 5. Mul. } a^3 + a^2b + ab^2 + b^3 \\ \text{by } a - b \\ \hline a^4 + a^3b + a^2b^2 + ab^3 \text{ --- Prod. by } a. \\ - a^3b - a^2b^2 - ab^3 - b^4 \text{ --- Prod. by } -b. \\ \hline a^4 \quad * \quad * \quad * \quad - b^4 \text{ Product by } a - b. \\ \hline \end{array}$$

Ex. 6. Ans. $a^4 + a^2b^2 + b^4$.

Ex. 7. Ans. $3x^4 + 4x^3y - 13x^2 - 4x^2y^2 + 24xy - 30$.

Ex. 8. Ans. $9a^4 - 18a^3x + 2a^2x^2 - 6ax^3 - 35x^4$

Ex. 9. Ans. $6x^6 - 5x^5y^2 + 15x^3y^3 - 6x^4y^4 - 3x^2y^5 + 9y^6$.

Ex. 10. Ans. $a^3 - a^2b - ab^2 - 2b^3$.

DIVISION OF ALGEBRA.

(Page 175.)

When the Divisor and Dividend are both Simple Quantities.

Ex. 3. $8x)16x^2(2x$. Quotient.

Ex. 4. $-3a^2x)12a^2x^2(-4x$. Quotient.

Ex. 5. $3ay)-15ay^2(-5y$. Quotient.

Ex. 6. $-8axz)-18ax^2y(\frac{9xy}{4z}$. Quotient.

(Key to Vol. I. page 176.)

When the Divisor is Simple, and the Dividend a Compound Quantity.

$$\text{Ex. 4. Ans. } 3b - 4x + \frac{1}{2}. \quad \text{Ex. 5. Ans. } x + 2 + \frac{2a - 5}{x}.$$

$$\text{Ex. 6. Ans. } 4x - 3a + 2c. \quad \text{Ex. 7. Ans. } 2a^2 - 3x - 5.$$

$$\text{Ex. 8. Ans. } 3x^2 - 3ab - \frac{d^2}{c} \quad \text{Ex. 9. Ans. } \frac{5+y}{7} - \frac{6y}{7a}.$$

$$\text{Ex. 10. Ans. } \frac{3\frac{1}{3}d^2b}{a} - 10b^2. \quad \text{or, } \frac{10d^2b}{3a} - 10b^2.$$

(Page 178.)

When the Divisor and Dividend are both Compound Quantities.

$$\text{Ex. 1. } \begin{array}{r} a+2x \overline{) a^2+4ax+4x^2} \\ \underline{a^2+2ax} \\ 2ax+4x^2 \\ \underline{2ax+4x^2} \\ 0 \end{array} \quad \text{Quotient.}$$

$$\text{Ex. 2. } \begin{array}{r} a-z \overline{) a^3-3a^2z+3az^2-z^3} \\ \underline{a^3-a^2z} \\ -2a^2z+3az^2-z^3 \\ \underline{-2a^2z+2az^2} \\ az^2-z^3 \\ \underline{az^2-z^3} \\ 0 \end{array} \quad \text{Quotient.}$$

$$\text{Ex. 3. } \begin{array}{r} 1+a \overline{) 1-a+a^2-a^3+a^4-a^5+a^6-\&c.} \\ \underline{1+a} \\ -a \text{ rem.} \\ \underline{-a-a^2} \\ +a^2 \text{ rem.} \\ \underline{+a^2+a^3} \\ -a^3 \text{ rem. } \&c. \end{array} \quad \text{Quot.}$$

(Key to Vol. I. page 178.)

Ex. 4.

$$\begin{array}{r}
 3x-6)12x^4 \text{ -----} 192(4x^3+8x^2+16x+32. \text{ Quo.} \\
 \underline{12x^4-24x^3} \\
 24x^3 \\
 \underline{24x^3-48x^2} \\
 48x^2 \\
 \underline{48x^2-96x} \\
 96x-192 \\
 \underline{96x-192} \\
 \cdot \\

 \end{array}$$

Ex. 5. Divisor $a^2-2ab+b^2$

Quotient $(a^3-3a^2b+3ab^2-b^3)$.

$$\begin{array}{r}
 \text{Dividend } a^5-5a^4b+10a^3b^2-10a^2b^3+5ab^4-b^5 \\
 \underline{a^5-2a^4b+a^3b^2} \\
 -3a^4b+9a^3b^2-10a^2b^3 \\
 \underline{-3a^4b+6a^3b^2-3a^2b^3} \\
 3a^3b^2-7a^2b^3+5ab^4 \\
 \underline{3a^3b^2-6a^2b^3+3ab^4} \\
 -a^2b^3+2ab^4-b^5 \\
 \underline{-a^2b^3+2ab^4-b^5} \\
 \\

 \end{array}$$

Ex. 6.

$$\begin{array}{r}
 2z-3a)48z^3-96az^2-64a^2z+150a^3(24z^2-12az-50a^2. \text{ Quot.} \\
 * \underline{-24az^2-64a^2z} \\
 * \underline{-100a^2z+150a^3} \\
 * \\
 *
 \end{array}$$

Ex. 7. Divisor $b^3-3b^2x+3bx^2-x^3$

Quotient $(b^3+3b^2x+3bx^2+x^3)$

$$\begin{array}{r}
 \text{Dividend })b^6 * -3b^4x^2 * +3b^2x^4 * -x^6 \\
 * \underline{3b^5x-6b^4x^2+b^3x^3+3b^2x^4} \\
 * \underline{3b^4x^2-8b^3x^3+6b^2x^4} \\
 * \underline{b^3x^3-3b^2x^4+3bx^5-x^6} \\
 * \\
 *
 \end{array}$$

(Key to Vol. I. page 180.)

*To reduce an Improper Fraction to a Whole or Mixed Quantity.*Ex. 3. $5)33(6\frac{3}{5}$. And

$$a)2ax - 3x^2(2x - \frac{3}{a}x^2.$$

Therefore $6\frac{3}{5}$, and $2x - \frac{3x^2}{a}$ are mixed quantities equivalent to the given fractions.

Ex. 4. Ans. $2ax$, and $2a + 2b + \frac{4b^2}{a-b}$.Ex. 5. Ans. $3x - 3y$, and $2x^2 + 2xy + 2y^2$.Ex. 6. Ans. $2a - \frac{4}{5} + \frac{6}{5a}$. Or, $2a - \frac{4a-6^*}{5a} = 2a + \frac{6-4a}{5a}$.Ex. 7. Ans. $5 - \frac{5a^2 + 10a + 20}{3a^3 + 2a^2 - 2a - 4}$.

* Here, if the fractional part $\frac{4}{5} + \frac{6}{5a}$ be reduced to a common denominator and expressed as one quantity, the sign + of $\frac{6}{5a}$ must be changed to minus, since the leading sign denotes that the whole compound quantity is to be subtracted, affecting $\frac{6}{5a}$ in a manner contrary to that intended by $-\frac{4}{5} + \frac{6}{5a}$. [See an Example, the last line, page 88 of the KEY.] Now, because changing plus to minus, and minus to plus, is in reality changing Addition to Subtraction, and Subtraction to Addition, it follows that, in all compound quantities under the same VINCULUM, of which the leading sign is minus, all the other signs must invariably be made contrary when the VINCULUM is broken.  The Course offers no Explanation on this subject!

(Page 181.)

*To reduce Fractions to a Common Denominator.*Ex. 3. $2a \times 2c = 4ac$ First Numerator. $3b \times x = 3bx$ Second Numerator.And $x \times 2c = 2cx$ Common Denominator.Therefore $\frac{4ac}{2cx}$, and $\frac{3bx}{2cx}$. Ans.

(Key to Vol. I. page 181.)

Ex. 4. $2a \times 2c = 4ac$ First Numerator.
 $(3a + 2b) \times b = 3ab + 2b^2$ Second Numerator.
 And $b \times 2c = 2bc$ Common Denominator.

Wherefore $\frac{4ac}{2bc}$, and $\frac{3ab + 2b^2}{2bc}$ are the fractions required.

Ex. 5. $5a \times 2c \times 1 = 10ac$ First Numerator.
 $3b \times 3x \times 1 = 9bx$ Second Numerator.
 $4d \times 3x \times 2c = 24cdx$ Third Numerator.
 And $3x \times 2c \times 1 = 6cx$ Common Denominator.

Consequently the new fractions are $\frac{10ac}{6cx}$, $\frac{9bx}{6cx}$, and $\frac{24cdx}{6cx}$.

Ex. 6. $5 \times 4 \times b = 20b$ First Numerator.
 $3a \times 6 \times b = 18ab$ Second Numerator.
 $(2b^2 + 3a) \times 6 \times 4 = 48b^2 + 72a$ Third Numerator.
 And $6 \times 4 \times b = 24b$ Common Denominator.

Wherefore $\frac{20b}{24b}$, $\frac{18ab}{24b}$, and $\frac{48b^2 + 72a}{24b}$ are the fractions required, admitting however of reduction to *lower terms*.

Ex. 7. Ans. $\frac{4a + 4b}{12a + 12b}$, $\frac{6a^3 + 6a^2b}{12a + 12b}$, and $\frac{24a^2 + 12b^2}{12a + 12b}$; or, in
lowest terms, $\frac{2a + 2b}{6a + 6b}$, $\frac{3a^3 + 3a^2b}{6a + 6b}$, and $\frac{12a^2 + 6b^2}{6a + 6b}$.

Ex. 8. Ans. $\frac{9b}{12a^2}$, $\frac{8ac}{12a^2}$, and $\frac{6ad}{12a^2}$.

(Page 182.)

To find the Greatest Common Measure of Fractions.

Ex. 3. Dividing the given divisor by b , it becomes $a + 2$.
 And $a + 2 \overline{) a^2 + 2a}$ * $-4(a - 2)$ Quotient.

$$\begin{array}{r} a^2 + 2a \\ \underline{a^2 + 2a} \\ -2a - 4 \\ \underline{-2a - 4} \end{array}$$

* without remainder.

$a + 2$, therefore, is the greatest common measure.

(Key to Vol. I. page 183.)

Ex. 4.
$$\frac{(a^4 - b^4)a^5}{a^5 - ab^4} * -a^3b^2(a$$

Divide by ab^2 - - - -) $ab^4 - a^3b^2$ remainder.

Quot.
$$\frac{b^2 - a^2}{b^2 - a^2} = -a^2 + b^2) a^4 * -b^4(-a^2 - b^2$$

$$\frac{a^4 - a^2b^2}{a^4 - a^2b^2}$$

$$\frac{a^2b^2 - b^4}{a^2b^2 - b^4}$$

Here, $-a + b$ is the last divisor and greatest common measure ; but it is evident that $a - b$ will likewise divide both terms of the given fraction without a remainder.

Ex. 5. The greatest common measure is 1 ; that is, the fraction is incommensurate. But, if instead of $\frac{a^3x + 2a^2x^2 + 2ax^3 + x^4}{5a^5 + 10a^4x + 5a^2x^2}$ it was intended to submit $\frac{a^3x + 2a^2x^2 + 2ax^3 + x^4}{5a^5 + 10a^4x + 5a^3x^2}$, the greatest common measure is $a + x$.

(Page 184.)

To reduce a Fraction to its Lowest Terms.

Ex. 3. The greatest common measure, by CASE IV. is $c - b$.

Hence $c - b) \frac{c^3 - b^3}{c^4 - b^2c^2} (\frac{c^2 + bc + b^2}{c^3 + bc^2}$ lowest terms.

Ex. 4. The greatest common measure being $a^2 - b^2$,

It is $a^2 - b^2) \frac{a^2 - b^2}{a^4 - b^4} (\frac{1}{a^2 + b^2}$ lowest terms.

Ex. 5. The greatest common measure is $a - b$. Hence

$a - b) \frac{a^4 - b^4}{a^3 - 3a^2b + 3ab^2 - b^3} (\frac{a^3 + a^2b + ab^2 + b^3}{a^2 - 2ab + b^2}$ lowest terms.

Ex. 6. The greatest common measure being $a^2 + 2ac + c^2$, It is

$a^2 + 2ac + c^2) \frac{3a^5 + 6a^4c + 3a^3c^2}{a^3c + 3a^2c^2 + 3ac^3 + c^4} (\frac{3a^3}{ac + c^2}$ lowest terms.

(Key to Vol. I. page 184.)

Ex. 7. The greatest common measure is $a+b$.

$$\text{Wherefore } a+b) \frac{a^3 - ab^2}{a^2 + 2ab + b^2} \left(\frac{a^2 - ab}{a+b} \right) \text{ lowest terms.}$$

ADDITION OF ALGEBRAIC FRACTIONS.

(Page 185.)

$$\text{Ex. 4. } \frac{(4x \times 5b) + (2x \times 3a)}{3a \times 5b} = \frac{20bx + 6ax}{15ab} \quad \text{Ans.}$$

$$\text{Ex. 5. } \frac{(a \times 4 \times 5) + (a \times 3 \times 5) + (a \times 3 \times 4)}{3 \times 4 \times 5} = \frac{47}{60}a. \quad \text{Ans.}$$

$$\text{Ex. 6. } \frac{(2a-3) \times 8 + (5a \times 4)}{4 \times 8} = \frac{9a-6}{8}. \quad \text{Ans.}$$

$$\text{Ex. 7. } \frac{(11a+3) \times 4 + (18a-5) \times 5}{5 \times 4} = 6\frac{7}{10}a - \frac{13}{20} \quad \text{or}$$

$$6a + \frac{14a-13}{20}. \quad \text{Ans.}$$

$$\text{Ex. 8. } \frac{(3a^2 \times 3b) + 4b(a+b) + 6a}{4b \times 3b} = 6a + \frac{9a^2 + 4a + 4b}{12b}. \quad \text{Ans.}$$

$$\text{Ex. 9. } \frac{(5a \times 5 \times 7) + (6a \times 4 \times 7) + (3a+2) \times 4 \times 5}{4 \times 5 \times 7} = 2a + \frac{123a+40}{140}. \quad \text{Ans.}$$

$$\text{Ex. 10. } \text{Ans. } 2\frac{13}{24}a + 3.$$

$$\text{Ex. 11. } \text{Ans. } 10\frac{1}{8}a.$$

SUBTRACTION OF ALGEBRAIC FRACTIONS.

(Page 186.)

$$\text{Ex. 3. } \frac{10a}{9} - \frac{4a}{7} = \frac{70a-36a}{63} = \frac{34}{63}a. \quad \text{Ans.}$$

(Key to Vol. I. page 186.)

$$\text{Ex. 4. } \frac{6a}{1} - \frac{3a}{4} = \frac{24a-3a}{4} = \frac{21}{4}a = 5\frac{1}{4}a. \quad \text{Ans.}$$

$$\text{Ex. 6. } \frac{5a}{4} - \frac{2a}{3} = \frac{15a-8a}{12} = \frac{7}{12}a. \quad \text{Ans.}$$

$$\text{Ex. 7. } \frac{2a+6}{9} = \frac{10a+30}{45}, \text{ and } \frac{4a+8}{5} = \frac{36a+72}{45}.$$

$$\text{But } \frac{36a+72-10a-30}{45} = \frac{26a+42}{45}. \quad \text{Ans.}$$

$$\text{Ex. 8. The minuend reduced is } \left\{ \frac{4ac+2a}{c} \right.$$

$$\text{The subtrahend reduced is } \left\{ \frac{2ac-a+3b}{c} \right.$$

$$\text{Hence, The difference required is } \left\{ \frac{2ac+3a-3b}{c} \right.$$

MULTIPLICATION OF ALGEBRAIC FRACTIONS.

(Page 187.)

$$\text{Ex. 4. } \left. \begin{array}{l} 4a \times 6a = 24a \quad \text{Numerator.} \\ 3 \times 5c = 15c \quad \text{Denominator.} \end{array} \right\} = \frac{8a}{5c} \quad \text{Ans.}$$

$$\text{Ex. 5. } \frac{3a}{4} \times \frac{4b^2}{3a} = b^2. \quad \text{Ans.}$$

$$\text{Ex. 6. } \frac{3a}{b} \times \frac{8ac}{b} \times \frac{4ab}{3c} = \frac{32a^3}{b}. \quad \text{Ans.}$$

$$\text{Ex. 7. } \text{Ans. } \frac{12a^3c+3a^3b}{2bc}.$$

$$\text{Ex. 8. } \text{Ans. } \frac{8a^4-4a^2b^2-4b^4}{3abc+3b^2c}.$$

$$\text{Ex. 9. } \text{Ans. } \frac{12a^2-3}{2a+b}.$$

(Key to Vol. I. page 187.)

Ex. 10. Reducing the fractions to a common denominator, and collecting like terms, it will be $\frac{4a^3 + 2ax - x^2}{4a^2} \times \frac{4x^3 - 2ax + a^2}{4x^3} = \frac{4a^5 - 8a^4x + 16a^3x^3 - 5a^2x^2 + 2a^3x + 2ax^3 + 8ax^4 - 4x^5}{16a^2x^2}$, the product required.

DIVISION OF ALGEBRAIC FRACTIONS.

(Page 188.)

Ex. 5. $\frac{11}{12} \Big) \frac{3x}{4} \left(\frac{36x}{44} = \frac{9}{11}x. \text{ Quotient.}$

Ex. 6. $\frac{3x}{1} \Big) \frac{6x^2}{5} \left(\frac{6x^2}{15x} = \frac{2}{5}x. \text{ Quotient.}$

Ex. 7. $\frac{4x}{3} \Big) \frac{3x+1}{9} \left(\frac{9x+3}{36} = \frac{1}{12}x + \frac{1}{4}. \text{ Quotient.}$

Ex. 8. $\frac{x}{3} \Big) \frac{4x}{2x-1} \left(\frac{12x}{2x^2-x} = \frac{12}{2x-1}. \text{ Quotient.}$

Ex. 9. $\frac{3a}{5b} \Big) \frac{4x}{5} \left(\frac{4bx}{3a} = \frac{1\frac{1}{3}bx}{a}. \text{ Quotient.}$

Ex. 10. $\frac{5ac}{6d} \Big) \frac{2a-b}{4cd} \left(\frac{12a-6b}{20ac^2} = \frac{6a-3b}{10ac^2}. \text{ Quotient.}$

Ex. 11. Ans. $\frac{10a^5 - 10ab^4 - 10a^4b + 10b^5}{6a^4 - 7a^3b - 4a^2b^2 + 5ab^3}.$

INVOLUTION.

(Page 190.)

Ex. 1. $3a^2 \times 3a^2 \times 3a^2 = 27a^6.$ The Cube required.

Ex. 2. $2a^2b \times 2a^2b \times 2a^2b \times 2a^2b = 16a^8b^4.$ The Power required.

(Key to Vol. I. page 190.)

Ex. 3. Ans. $-64a^6b^9$. Ex. 4. Ans. $\frac{a^8x^4}{16b^8}$.

Ex. 5. Ans. $a^5 - 10a^4x + 40a^3x^2 - 80a^2x^3 + 80ax^4 - 32x^5$.

<p>Ex. 6. <i>Root</i> $2\sqrt{a}$ Mul. by $2\sqrt{a}$ <hr style="width: 50%; margin-left: 0;"/> <i>Square</i> $4a$ Mul. by $2\sqrt{a}$ <hr style="width: 50%; margin-left: 0;"/> <i>Cube</i> $8a\sqrt{a}$ Mul. by $2\sqrt{a}$ <hr style="width: 50%; margin-left: 0;"/> <i>4th Power</i> $16aa = 16a^2$ Mul. by $2\sqrt{a}$ <hr style="width: 50%; margin-left: 0;"/> <i>5th Power</i> $32a^2\sqrt{a}$ Mul. by $2\sqrt{a}$ <hr style="width: 50%; margin-left: 0;"/> <i>6th Power</i> $64a^2a = 64a^3$. Ans.</p>	or,	<p>$2a^{\frac{1}{2}}$ <i>Root.</i> <hr style="width: 50%; margin-left: 0;"/> $4a^{\frac{1}{1}}$ <i>Square.</i> <hr style="width: 50%; margin-left: 0;"/> $8a^{\frac{3}{2}}$ <i>Cube.</i> <hr style="width: 50%; margin-left: 0;"/> $16a^{\frac{2}{1}}$ <i>Biquadrate.</i> <hr style="width: 50%; margin-left: 0;"/> $32a^{\frac{5}{2}}$ <i>Sursolid.</i> <hr style="width: 50%; margin-left: 0;"/> $64a^{\frac{3}{1}}$ <i>Sixth Power.</i></p>
---	-----	---

EVOLUTION.

(Page 193.)

To determine the Roots of Simple Quantities.

Ex. 5. $\sqrt{2a^2b^4} = \sqrt{2} \times \sqrt{a^2} \times \sqrt{b^4} = ab^2\sqrt{2}$. Root required.

Ex. 6. $\sqrt[3]{-64a^3b^6} = \sqrt[3]{-64} \times \sqrt[3]{a^3} \times \sqrt[3]{b^6} = -4ab^2$. Root reqd.

Ex. 7. $\sqrt{\frac{8a^2b^2}{3c^3}} = \frac{\sqrt{4}\sqrt{2}\sqrt{a^2}\sqrt{b^2}}{\sqrt{c^2}\sqrt{3c}} = \frac{2ab}{c}\sqrt{\frac{2}{3c}}$. Root required.

Ex. 8. $\sqrt[4]{81a^4b^6} = \sqrt[4]{81}\sqrt[4]{a^4} \times \sqrt[4]{b^4} \times \sqrt[4]{b^2} = 3ab\sqrt{b}$. Root reqd.

Ex. 9. $\sqrt[5]{-32a^5b^0} = \sqrt[5]{-32}\sqrt[5]{a^5} \times \sqrt[5]{b^5} \times \sqrt[5]{b} = -2ab\sqrt[5]{b}$. Ans.

(Key to Vol. I. page 195.)

To extract the Square Root of a Compound Quantity.

Ex. 3.
$$\begin{array}{r} a^4 + 4a^3 + 6a^2 + 4a + 1 \text{ the Root.} \\ a^4 \end{array}$$

1st Divisor $2a^2 + 2a$) $4a^3 + 6a^2$ *Dividend.*
 $4a^3 + 4a^2$

2d Divisor $2a^2 + 4a + 1$) $2a^2 + 4a + 1$ *Dividend.*
 $2a^2 + 4a + 1$
 . . .

Ex. 4.
$$\begin{array}{r} a^4 - 2a^3 + 2a^2 - a + \frac{1}{4} \text{ the Root.} \\ a^4 \end{array}$$

1st Divisor $2a^2 - a$) $-2a^3 + 2a^2$ *Dividend.*
 $-2a^3 + a^2$

2d Divisor $2a^2 - 2a + \frac{1}{2}$) $a^2 - a + \frac{1}{4}$ *Dividend.*
 $a^2 - a + \frac{1}{4}$
 . . .

Ex. 5.
$$\begin{array}{r} a^2 - ab \left(a - \frac{b}{2} - \frac{b^2}{8a} - \frac{b^3}{16a^2} - \&c. \text{ Root reqd.} \right. \\ a^2 \end{array}$$

1st Divisor $2a - \frac{b}{2}$) $-ab$ *Dividend.*
 $-ab + \frac{b^2}{4}$

2d Divisor $2a - b - \frac{b^2}{8a}$) $-\frac{b^2}{4}$ *Dividend.*
 $-\frac{b^2}{4} + \frac{b^3}{8a} + \frac{b^4}{64a^2}$

3d Divisor $2a - b - \frac{b^2}{4a} - \frac{b^3}{16a^2}$) $-\frac{b^3}{8a} - \frac{b^4}{64a^2}$ *Dividend, &c.*

(Key to Vol. I. page 195.)

To determine the Roots of Powers in General.

Ex. 3.
$$\frac{a^2 - 2ab + 2ax + b^2 - 2bx + x^2}{a^2} \quad (a - b + x. \text{ Ans.}$$

Divisor $2a) \cdot -2ab$ *Dividend.*

$$a^2 - 2ab \quad + b^2$$

Divisor $2a) \cdot \cdot +2ax \cdot$ *Dividend.*

$$a^2 - 2ab + 2ax + b^2 - 2bx + x^2$$

.

Ex. 4.
$$\frac{a^6 - 3a^5 + 9a^4 - 13a^3 + 18a^2 - 12a + 8}{a^6} \quad (a^2 - a + 2. \text{ Ans.}$$

Divis. $3a^4) \cdot -3a^5$ - - - - - *Dividend.*

$$a^6 - 3a^5 + 3a^4 - a^3$$

Divis. $3a^4) \cdot \cdot +6a^4 - 12a^3$ - - *Remainder.*

$$a^6 - 3a^5 + 9a^4 - 13a^3 + 18a^2 - 12a + 8$$

.

Ex. 5.
$$\frac{81a^4 - 216a^3b + 216a^2b^2 - 96ab^3 + 16b^4}{81a^4} \quad (3a - 2b. \text{ Ans.}$$

Div. $108a^3) \cdot -216a^3b$ *Dividend.*

$$81a^4 - 216a^3b + 216a^2b^2 - 96ab^3 + 16b^4$$

.

Ex. 6.
$$\frac{a^5 - 10a^4 + 40a^3 - 80a^2 + 80a - 32}{a^5} \quad (a - 2 \text{ Root sought.}$$

Divis. $5a^4) \cdot -10a^4$ *Dividend.*

$$a^5 - 10a^4 + 40a^3 - 80a^2 + 80a - 32$$

.

(Key to Vol. I. page 195.)

Ex. 7. $1 - x^2 \left(1 - \frac{x^2}{2} - \frac{x^4}{8} - \frac{x^6}{16} - \&c. \right)$ Root required.

$$\begin{array}{r}
 1 \\
 \hline
 \text{Divisor } 2) \cdot -x^2 \quad \text{Dividend.} \\
 \hline
 1 - x^2 + \frac{x^4}{4} \quad \text{Subtrahend.} \\
 \hline
 \text{Divisor } 2) \cdot \cdot -\frac{x^4}{4} \quad \text{Rem. and Dividend.} \\
 \hline
 1 - x^2 \quad * \quad +\frac{x^6}{8} + \frac{x^8}{64} \quad \text{Subtrahend.} \\
 \hline
 \text{Divisor } 2) \cdot \cdot \cdot -\frac{x^6}{8} - \frac{x^8}{64} \quad \text{Remainder.} \\
 \hline
 1 - x^2 \quad \&c. + \&c. \quad \text{Subtrahend.} \\
 \hline
 \hline
 \end{array}$$

Ex. 8. $1 - x^3 \left(1 - \frac{x^3}{3} - \frac{x^6}{9} - \&c. \text{ AD INFINITUM. } \right)$ Cube root reqd.

$$\begin{array}{r}
 1 \\
 \hline
 \text{Div. } 3) \cdot -x^3 \quad \text{Dividend.} \\
 \hline
 1 - x^3 + \frac{x^6}{3} + \frac{x^9}{27} \quad \text{Subtrahend.} \\
 \hline
 \text{Div. } 3) \cdot \cdot -\frac{x^6}{3} - \frac{x^9}{27} \quad \text{Remainder.} \\
 \hline
 1 - x^3 \quad \&c. + \&c. \quad \text{Subtrahend.} \\
 \hline
 \hline
 \end{array}$$

SURDS.

(Page 196.)

To reduce a Rational Quantity to the Form of a Surd.

Ex. 3. $6 \times 6 \times 6 = 216$, wherefore $\sqrt[3]{216}$, or, $(216)^{\frac{1}{3}}$. form reqd.

(Key to Vol. I. page 196.)

Ex. 4. $\frac{1}{3}ab \times \frac{1}{3}ab = \frac{1}{9}a^2b^2$, wherefore $\sqrt{\frac{1}{9}a^2b^2}$, or, $(\frac{1}{9}a^2b^2)^{\frac{1}{2}}$. Ans.

Ex. 5. $2 \times 2 \times 2 \times 2 = 16$, wherefore $\sqrt[4]{16}$, or, $(16)^{\frac{1}{4}}$. Ans.

Ex. 6. $a^{\frac{1}{3}} \times a^{\frac{1}{3}} \times a^{\frac{1}{3}} \times a^{\frac{1}{3}} \times a^{\frac{1}{3}} = a^{\frac{5}{3}} = \sqrt[3]{a^5}$. form required.

Ex. 7.

$(a+x) \times (a+x) = a^2 + 2ax + x^2$, hence $\sqrt{a^2 + 2ax + x^2}$. Ans.

Ex. 8. $\left\{ (a-x) \times (a-x) \times (a-x) \right\}^{\frac{1}{3}} = \sqrt[3]{a^3 - 3a^2x + 3ax^2 - x^3}$. the form required.

(Page 197.)

To reduce Quantities to a Common Index.

Ex. 3. $\frac{1}{4} \frac{1}{3} (\frac{4}{3})$ the form for 4.

And $\frac{1}{4} \frac{1}{2} (\frac{4}{2}) = \frac{2}{1}$ the form for 5.

But $4^4 = 256$, and $5^2 = 25$,

Hence $(256^{\frac{1}{3}})^{\frac{1}{4}}$, and $25^{\frac{1}{2}}$. Ans.

Ex. 4. $\frac{1}{6} \frac{1}{3} (\frac{6}{3}) = \frac{2}{1}$ the form for a .

And $\frac{1}{6} \frac{1}{4} (\frac{6}{4}) = \frac{3}{2}$ the form for x .

Hence $(a^2)^{\frac{1}{6}}$, and $(x^{\frac{3}{2}})^{\frac{1}{6}}$. Ans.

Ex. 5. For the form of the Square Root,

And $\left. \begin{array}{l} \frac{1}{2} \frac{2}{1} (\frac{4}{1}) \text{ for the index of } a, \\ \frac{1}{2} \frac{3}{1} (\frac{6}{1}) \text{ for the index of } x, \end{array} \right\} \text{giving } \sqrt{a^4}, \text{ and } \sqrt{x^6}. \text{ Ans.}$

For the form of the Cube Root,

And $\left. \begin{array}{l} \frac{1}{3} \frac{2}{1} (\frac{6}{1}) \text{ for the index of } a, \\ \frac{1}{3} \frac{3}{1} (\frac{9}{1}) \text{ for the expon. of } x, \end{array} \right\} \text{hence } \sqrt[3]{a^6}, \text{ and } \sqrt[3]{x^9}. \text{ Ans.}$

(Key to Vol. I. page 197.)

For the form of the n th Root,

And $\left. \begin{array}{l} \frac{1}{n} \times 2 \left(\frac{2n}{1} \text{ for the Index of } a, \right) \\ \frac{1}{n} \times 3 \left(\frac{3n}{1} \text{ for the Index of } x, \right) \end{array} \right\} \text{wherefore } \sqrt[n]{a^{2n}}, \text{ and } \sqrt[n]{x^{3n}}.$

But it is evident that $a^{\frac{4}{2}} = a^2$, and that $x^{\frac{6}{2}} = x^3$, consequently the given quantities have the same radical sign, namely 1, without any operation.

Ex. 6. $\frac{1 \times 2}{3 \times 2} = \frac{2}{6}$ for the form of $(a+x)$, And

$\frac{1 \times 3}{3 \times 2} = \frac{3}{6}$ for the form of $(a-x)$

Consequently $\left. \begin{array}{l} (a^2 + 2ax + x^2)^{\frac{1}{6}} \\ \text{And } (a^3 - 3a^2x + 3ax^2 - x^3)^{\frac{1}{6}} \end{array} \right\} \text{Ans.}$

Ex. 7. $\frac{1 \times 4}{2 \times 4} = \frac{4}{8}$, and $\frac{1 \times 2}{2 \times 4} = \frac{2}{8}$

Therefore $\left. \begin{array}{l} (a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4)^{\frac{1}{8}} \\ \text{And } (a^2 - 2ab + b^2)^{\frac{1}{8}} \end{array} \right\} \text{Ans.}$

(Page 198.)

To reduce Complicated Surds to Simpler Terms.*

Ex. 3. $\sqrt{75} = \sqrt{25} \times \sqrt{3} = 5\sqrt{3}$. Ans.

Ex. 4. $\sqrt{\frac{44}{75}} = \frac{\sqrt{4}\sqrt{11}}{\sqrt{25}\sqrt{3}} = \frac{2}{5} \times \frac{\sqrt{11}}{\sqrt{3}} = \frac{2}{5} \times \frac{\sqrt{33}}{\sqrt{9}} = \frac{2}{15}\sqrt{33}$. Ans.

Ex. 5. $\sqrt[3]{189} = \sqrt[3]{27} \sqrt[3]{7} = 3\sqrt[3]{7}$. Ans.

Ex. 6. $\sqrt[3]{\frac{135}{32}} = \frac{\sqrt[3]{27}\sqrt[3]{5}}{\sqrt[3]{8}\sqrt[3]{4}} = \frac{3\sqrt[3]{5}}{2\sqrt[3]{4}} = \frac{3}{2} \times \frac{\sqrt[3]{10}}{\sqrt[3]{8}} = \frac{3}{4}\sqrt[3]{10}$. Ans.

* Quantities consisting of a *rational* and an *irrational* part involved together are termed *Complicated Surds*.

(Key to Vol. I. page 198.)

Ex. 7. $\sqrt{75a^2b} = \sqrt{25} \sqrt{3} \sqrt{a^2} \sqrt{b} = 5a \sqrt{3b}$. Ans.

ADDITION OF SURDS.

(Page 199.)

Ex. 3. $\sqrt{27} = \sqrt{9} \sqrt{3} = 3\sqrt{3}$
 And $\sqrt{48} = \sqrt{16} \sqrt{3} = 4\sqrt{3}$
 $\underline{\hspace{1.5cm}}$
 $7\sqrt{3}$ Sum required.

Ex. 4. $\sqrt{50} = \sqrt{25} \sqrt{2} = 5\sqrt{2}$
 And $\sqrt{72} = \sqrt{36} \sqrt{2} = 6\sqrt{2}$
 $\underline{\hspace{1.5cm}}$
 $11\sqrt{2}$ Sum required.

Ex. 5. $\sqrt{\frac{3}{5}} = \sqrt{\frac{45}{75}} = \frac{\sqrt{9} \sqrt{5}}{\sqrt{25} \sqrt{3}} = \frac{3}{5} \times \frac{\sqrt{5}}{\sqrt{3}} = \frac{3}{5} \sqrt{\frac{15}{9}} = \frac{3}{15} \sqrt{15}$.

And $\sqrt{\frac{1}{15}} = \sqrt{\frac{5}{75}} = \frac{\sqrt{1} \sqrt{5}}{\sqrt{25} \sqrt{3}} = \frac{1}{5} \times \frac{\sqrt{5}}{\sqrt{3}} = \frac{1}{5} \sqrt{\frac{15}{9}} = \frac{1}{15} \sqrt{15}$.

$\underline{\hspace{1.5cm}}$
 Sum and Answer $\frac{4}{15} \sqrt{15}$.

But $\frac{4}{15} \sqrt{15} = \frac{4}{1} \times \frac{\sqrt{15}}{\sqrt{225}} = 4 \sqrt{\frac{1}{15}}$ the other Ans. given in the COURSE.

Ex. 6. $\sqrt[3]{56} = \sqrt[3]{8} \sqrt[3]{7} = 2\sqrt[3]{7}$
 And $\sqrt[3]{189} = \sqrt[3]{27} \sqrt[3]{7} = 3\sqrt[3]{7}$
 $\underline{\hspace{1.5cm}}$
 $5\sqrt[3]{7}$ Sum required.

(Key to Vol. I. page 199.)

$$\text{Ex. 7. } \sqrt[3]{\frac{1}{4}} = \frac{\sqrt[3]{128}}{\sqrt[3]{512}} = \frac{\sqrt[3]{8} \sqrt[3]{16}}{\sqrt[3]{64} \sqrt[3]{8}} = \frac{1}{4} \sqrt[3]{16} = \frac{1}{4} \sqrt[3]{8} \sqrt[3]{2} = \frac{1}{2} \sqrt[3]{2}.$$

$$\text{And } \sqrt[3]{\frac{1}{32}} = \frac{\sqrt[3]{16}}{\sqrt[3]{512}} = \frac{\sqrt[3]{1} \sqrt[3]{16}}{\sqrt[3]{64} \sqrt[3]{8}} = \frac{1}{8} \sqrt[3]{16} = \frac{1}{8} \sqrt[3]{8} \sqrt[3]{2} = \frac{1}{4} \sqrt[3]{2}.$$

Sum and Answer $\frac{3}{4} \sqrt[3]{2}.$

$$\text{Ex. 8. } 3\sqrt{a^2b} = 3\sqrt{a^2} \sqrt{b} = 3a \sqrt{b}$$

$$\text{And } 5\sqrt{16a^4b} = 5\sqrt{16} \sqrt{a^4} \sqrt{b} = 20a^2 \sqrt{b}$$

Sum $\underline{\underline{(20a^2 + 3a)\sqrt{b}}}$. Ans.

SUBTRACTION OF SURDS.

$$\text{Ex. 3. } \sqrt{75} = \sqrt{25} \sqrt{3} = 5\sqrt{3}$$

$$\text{And } \sqrt{48} = \sqrt{16} \sqrt{3} = 4\sqrt{3}$$

Difference $\underline{\underline{\sqrt{3}}}$. Ans.

$$\text{Ex. 4. } \sqrt[3]{256} = \sqrt[3]{64} \sqrt[3]{4} = 4\sqrt[3]{4}$$

$$\text{And } \sqrt[3]{32} = \sqrt[3]{8} \sqrt[3]{4} = 2\sqrt[3]{4}$$

Difference $\underline{\underline{2\sqrt[3]{4}}}$. Ans.

$$\text{Ex. 5. } \sqrt{\frac{3}{4}} = \sqrt{\frac{9}{12}} = \frac{\sqrt{9} \sqrt{1}}{\sqrt{4} \sqrt{3}} = \frac{3}{2} \sqrt{\frac{1}{3}} = \frac{3}{2} \sqrt{\frac{3}{9}} = \frac{1}{2} \sqrt{3}$$

$$\text{And } \sqrt{\frac{1}{3}} = \sqrt{\frac{4}{12}} = \frac{\sqrt{4} \sqrt{1}}{\sqrt{4} \sqrt{3}} = \sqrt{\frac{1}{3}} = \sqrt{\frac{3}{9}} = \frac{1}{3} \sqrt{3}$$

Difference $\underline{\underline{\frac{1}{6} \sqrt{3}}}$. Ans.

(Key to Vol. I. page 199.)

$$\text{Ex. 6.} \quad \sqrt{\frac{2}{3}} = \sqrt{\frac{16}{24}} = \frac{\sqrt{16}\sqrt{1}}{\sqrt{4}\sqrt{6}} = 2\sqrt{\frac{1}{6}} = 2\sqrt{\frac{6}{36}} = \frac{4}{12}\sqrt{6}$$

$$\text{And} \quad \sqrt{\frac{3}{8}} = \sqrt{\frac{9}{24}} = \frac{\sqrt{9}\sqrt{1}}{\sqrt{4}\sqrt{6}} = \frac{3}{2}\sqrt{\frac{1}{6}} = \frac{3}{2}\sqrt{\frac{6}{36}} = \frac{3}{12}\sqrt{6}$$

$$\text{Difference} \quad \frac{1}{12}\sqrt{6}. \text{ Ans.}$$

$$\text{Ex. 7.} \quad \sqrt[3]{2\frac{7}{9}} = \sqrt[3]{\frac{25}{9}} = \frac{\sqrt[3]{125}}{\sqrt[3]{45}} = \frac{5}{\sqrt[3]{45}}$$

$$\text{And} \quad \sqrt[3]{\frac{3}{5}} = \frac{\sqrt[3]{27}}{\sqrt[3]{45}} = \frac{3}{\sqrt[3]{45}}$$

$$\text{Difference} \quad \frac{2}{\sqrt[3]{45}}$$

$$\text{But} \quad \frac{2}{\sqrt[3]{45}} = \frac{\sqrt[3]{8}}{\sqrt[3]{45}} = \frac{\sqrt[3]{8^3/2025}}{45} = \frac{2}{45}\sqrt[3]{27^3/75} = \frac{2}{15}\sqrt[3]{75}. \text{ Ans.}$$

$$\text{Ex. 8.} \quad \sqrt{24a^2b^2} = \sqrt{4}\sqrt{6}\sqrt{a^2}\sqrt{b^2} = 2ab\sqrt{6}$$

$$\text{And} \quad \sqrt{54b^4} = \sqrt{9}\sqrt{6}\sqrt{b^4} = \dots = 3b^2\sqrt{6}$$

$$\text{Therefore} \quad \left\{ \begin{array}{l} \text{*when } 2ab > 3b^2 \text{ - - } (2ab - 3b^2)\sqrt{6} \\ \text{when } 2ab < 3b^2 \text{ - - } (3b^2 - 2ab)\sqrt{6} \end{array} \right\} \text{ Ans.}$$

* It may be necessary to mention that $>$ signifies *greater than*, and $<$ *less than*.

MULTIPLICATION OF SURDS.

(Page 200.)

Ex. 3. $3 \times 2 = 6$ the Product of the rational Quantities.

And $\sqrt{2} \times \sqrt{8} = \sqrt{16} = 4$ the Product of the Surd Quantities.

Consequently 24 *Product* required.

(Key to Vol. I. page 200.)

Ex. 4. $\frac{1}{3} \times \frac{3}{4} = \frac{1}{4}$ the Product of the rational Quantities.

And $\sqrt[3]{4} \times \sqrt[3]{12} = \sqrt[3]{48} = \sqrt[3]{8} \sqrt[3]{6} = 2\sqrt[3]{6}$ Product of the Surds.

Wherefore $\frac{1}{4} \times 2\sqrt[3]{6} = \frac{1}{2}\sqrt[3]{6}$. Ans.

Ex. 5. $\frac{5}{3} \times \frac{9}{10} = \frac{3}{2}$ the Product of the rational Quantities.

And $\sqrt{\frac{3}{8}} \times \sqrt{\frac{2}{5}} = \sqrt{\frac{6}{40}} = \frac{\sqrt{1}\sqrt{6}}{\sqrt{4}\sqrt{10}} = \frac{1}{2}\sqrt{\frac{6}{10}} = \frac{1}{2}\sqrt{\frac{60}{100}} = \frac{1}{20}\sqrt{60}$

But $\frac{1}{20}\sqrt{60} = \frac{1}{10}\sqrt{15}$ the Product of the Surd Quantities.

Therefore $\frac{3}{2} \times \frac{1}{10}\sqrt{15} = \frac{3}{20}\sqrt{15}$. Ans.

Ex. 6. $2 \times 3 = 6$ the Product of the rational Parts.

And $\sqrt[3]{14} \times \sqrt[3]{4} = \sqrt[3]{56} = \sqrt[3]{8} \sqrt[3]{7} = 2\sqrt[3]{7}$ Prod. of the Surds.

Wherefore $12\sqrt[3]{7}$. Ans.

Ex. 7. $2 \times 1 = 2$ the Product of the Co-efficients.

And $a^{\frac{2}{3}} \times a^{\frac{4}{3}} = a^{\frac{6}{3}} = a^2$ the Product of the Surds.

Consequently $2a^2$. Ans.

Ex. 8. $\frac{1}{3} = \frac{4}{12}$, and $\frac{3}{4} = \frac{9}{12}$, the new indices with a com. radical.

But $\sqrt[12]{a+b^4} \times \sqrt[12]{a+b^9} = (a+b)^{\frac{13}{12}}$. That is,

$(a^{13} + 13a^{12}b + 78a^{11}b^2 + 286a^{10}b^3 + 715a^9b^4 + \&c. \&c.)^{\frac{1}{12}}$. Ans.

Ex. 9. $2x + \sqrt{b}$ *Multiplicand.*
 $2x - \sqrt{b}$ *Multiplier.*

$4x^2 + 2x\sqrt{b}$ Prod. by $2x$.
 $-2x\sqrt{b} - b$ Prod. by $-\sqrt{b}$.

$4x^2$ * $-b$ PROD. by $(2x - \sqrt{b})$ which was reqd.

(Key to Vol. I. page 200.)

Ex. 10. $(a + 2\sqrt{b})^{\frac{1}{2}}$ *Multiplicand.*
 $(a - 2\sqrt{b})^{\frac{1}{2}}$ *Multiplier.*

$\sqrt{(a^2 - 4b)}$ PRODUCT required.

Ex. 11. $2x^{\frac{1}{n}} \times 3x^{\frac{1}{n}} = 6(x^2)^{\frac{1}{n}}$ or $6\sqrt[n]{x^2}$. Ans.

Or if, as in the Sixth Edition,* the question be $2x^{\frac{1}{n}}$ and $3x^{\frac{1}{m}}$,
First reducing the Surds to a common radical, it is

$$2x^{\frac{m}{mn}} \times 3x^{\frac{n}{mn}}$$

Next, $2x^m \times 3x^n = 6x^{m+n}$, therefore, subscribing the com-

mon index, $6x^{\frac{m+n}{mn}} = 6\sqrt[mn]{x^{m+n}}$. Ans.

* There happens to be two SIXTH EDITIONS of Hutton's Course (if both can be well termed Editions,) the one printed in 1811, and the other (whereof a part, at least, was certainly printed subsequently to the corresponding part of the former) purporting to have been printed in 1815. Here we allude to the latter Edition, whereas in page 30 of the Key the allusion applies to the former.

Ex. 12. $4x^{\frac{1}{n}} \times 2y^{\frac{1}{n}} = 8(xy)^{\frac{1}{n}}$, or $8\sqrt[n]{xy}$. Ans.

DIVISION OF SURDS.

(Page 201.)

Ex. 3. $4 \div 2 = 2$ the Quotient of the Co-efficients.

And $\sqrt{50} \div \sqrt{5} = \sqrt{10}$. Therefore $2\sqrt{10}$. Ans.

Ex. 4. $6 \div 3 = 2$ the Quotient of the Co-efficients.

And $\sqrt[3]{100} \div \sqrt[3]{5} = \sqrt[3]{20}$. Therefore $2\sqrt[3]{20}$. Ans.

Ex. 5. $\frac{5}{6} \div \frac{3}{4} = \frac{10}{9}$ the Quotient of the Co-efficients.

And $\sqrt{\frac{1}{50}} \div \sqrt{\frac{2}{5}} = \sqrt{\frac{5}{100}} = \frac{1}{10}\sqrt{5}$. Quotient of the Surds.

But $\frac{10}{9} \times \frac{1}{10}\sqrt{5} = \frac{1}{9}\sqrt{5}$. Ans.

(Key to Vol. I. page 201.)

Ex. 6. $\frac{3}{4} \div \frac{3}{5} = \frac{5}{4}$ the Quotient of the Co-efficients.

And $\sqrt[3]{\frac{3}{16}} \div \sqrt[3]{\frac{2}{5}} = \sqrt[3]{\frac{15}{32}} = \sqrt[3]{\frac{30}{64}} = \frac{1}{4} \sqrt[3]{30}$. Quot. of the Surds.

But $\frac{5}{4} \times \frac{1}{4} \sqrt[3]{30} = \frac{5}{16} \sqrt[3]{30}$. Ans.

Ex. 7. $\frac{4}{5} \div \frac{2}{3} = \frac{6}{5}$ the Quotient of the Co-efficients.

And $\frac{1}{2} - \frac{1}{3} = \frac{1}{6}$ for the exponent of the Quotient Surd.

Therefore $\frac{6}{5} a^{\frac{1}{6}}$. Ans.

Ex. 8. $\frac{4}{3} - \frac{2}{3} = \frac{2}{3}$ for the exponent of a in the Quotient.

Therefore $a^{\frac{2}{3}}$, or $\sqrt[3]{a^2}$. Ans.

Ex. 9. $3 \div 4 = \frac{3}{4}$ the Quotient of the Co-efficients.

And $\frac{x}{n}, \frac{x}{m}$, reduced to a common radical are $\frac{m}{mn}$, and $\frac{n}{mn}$.

Therefore $\frac{3}{4} a^{\frac{m-n}{mn}}$. Ans.

INVOLUTION OF SURDS.

Ex. 4. $2\sqrt[3]{2} \times 2\sqrt[3]{2} = 4\sqrt[3]{4}$. Ans.

Ex. 5. $\sqrt{3} \times \sqrt{3} \times \sqrt{3} = \sqrt{3^3} = \sqrt{27} = \sqrt{9} \sqrt{3} = 3\sqrt{3}$. Ans.

Ex. 6. $\frac{1}{3} \times \frac{1}{3} \times \frac{1}{3} = \frac{1}{27}$ the power required of the Co-efficient.

And, $\sqrt{3} \times \sqrt{3} \times \sqrt{3} = \sqrt{27} = 3\sqrt{3}$ the Cube of the Surd.

But $\frac{1}{27} \times 3\sqrt{3} = \frac{1}{9} \sqrt{3}$. Ans.

(Key to Vol. I. page 201.)

Ex. 7. $2^{\frac{1}{2}} \times 2^{\frac{1}{2}} \times 2^{\frac{1}{2}} \times 2^{\frac{1}{2}} = 2^2$ or $16^{\frac{1}{2}} = 4$ the power of the Surd.

And $\frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} = \frac{1}{16}$ the power of the Co-efficient.

But $\frac{1}{16} \times 4 = \frac{1}{4}$ Ans.

(Page 202.)

Ex. 8. Ans. $\sqrt[n]{a^m}$, or $a^{\frac{m}{n}}$.

Ex. 9. $2 + \sqrt{3}$ - - - - - Given Root.
 $2 + \sqrt{3}$ - - - - - Multiplier.

4 + 2√3 - - - - - Prod. by 2.
 + 2√3 + 3 - - - - - Prod. by √3.

Sum 4 + 4√3 + 3 = 7 + 4√3, the Square required.

EVOLUTION OF SURDS.

Ex. 3. $6^3 = 216$.

Now $\sqrt{216} = \sqrt{36} \sqrt{6} = 6\sqrt{6}$. Ans.

Ex. 4. $\sqrt[3]{\frac{1}{8}a^3b} = \sqrt[3]{\frac{1}{8}} \sqrt[3]{a^3} \sqrt[3]{b} = \frac{1}{2}a \sqrt[3]{b}$. Ans.

Ex. 5. $\sqrt[4]{16a^2} = \sqrt[4]{16} \sqrt[4]{a^2} = 2\sqrt{a}$. Ans.

Ex. 6. $\frac{x}{n} \div \frac{m}{1} = \frac{x}{mn}$, therefore $x^{\frac{1}{mn}}$. Ans.

Ex. 7. $a^2 - 6a\sqrt{b} + 9b$ ($a - 3\sqrt{b}$ Root required.)
 a^2

Divisor $2a) \cdot - 6a\sqrt{b}$ Dividend.

$a^2 - 6a\sqrt{b} + 9b$

(Key to Vol. I. page 204.)

INFINITE SERIES.

Fractional Quantities reduced to Infinite Series by Division.

Ex. 3.

$$a+c)b \dots - \left(\frac{b}{a} - \frac{bc}{a^2} + \frac{bc^2}{a^3} - \&c. \right) = \frac{b}{a} \times \left(1 - \frac{c}{a} + \frac{c^2}{a^2} - \&c. \right) \text{ Ans.}$$

$$\begin{array}{r}
 b + \frac{bc}{a} \\
 \hline
 \cdot - \frac{bc}{a} \\
 \frac{bc}{a} - \frac{bc^2}{a^2} \\
 \hline
 \cdot + \frac{bc^2}{a^2} \\
 + \frac{bc^2}{a^2} + \frac{bc^3}{a^3} \\
 \hline
 \cdot - \frac{bc^3}{a^3} \text{ remainder.} \\
 \hline
 \hline
 \end{array}$$

$$\text{Ex. 4. } a-b)a \dots - - - - \left(1 + \frac{b}{a} + \frac{b^2}{a^2} + \frac{b^3}{a^3} + \frac{b^4}{a^4} + \&c. \right) \text{ Ans.}$$

$$\begin{array}{r}
 a-b \\
 \hline
 \cdot + b \\
 + b - \frac{b^2}{a} \\
 \hline
 \cdot + \frac{b^2}{a} \\
 + \frac{b^2}{a} - \frac{b^3}{a^2} \\
 \hline
 \cdot + \frac{b^3}{a^2} \&c. \\
 \hline
 \hline
 \end{array}$$

(Key to Vol. I. page 204.)

Ex. 5. $\frac{1+x}{1+x} (1-2x+2x^2-2x^3+2x^4-\&c. \text{ series reqd.})$

$$\begin{array}{r}
 \frac{1+x}{1+x} \\
 \cdot -2x \\
 \hline
 -2x-2x^2 \\
 \hline
 \cdot +2x^2 \\
 +2x^2+2x^3 \\
 \hline
 \cdot -2x^3 \\
 -2x^3-2x^4 \\
 \hline
 \cdot +2x^4. \&c. \\
 \hline
 \end{array}$$

Ex. 6. $a^2+2ab+b^2) a^2 - \dots - (1 - \frac{2b}{a} + \frac{3b^2}{a^2} - \frac{4b^3}{a^3} + \&c. \text{ Ans.})$

$$\begin{array}{r}
 \frac{a^2+2ab+b^2}{a^2+2ab+b^2} \\
 \cdot -2ab-b^2 \\
 \hline
 -2ab-4b^2-\frac{2b^3}{a} \\
 \hline
 \cdot +3b^2+\frac{2b^3}{a}. \&c. \\
 \hline
 \end{array}$$

Ex. 7. $\frac{1+1}{1+1} (1-1+1-1+\&c. \text{ the series required.})$

$$\begin{array}{r}
 \frac{1+1}{1+1} \\
 \cdot -1 \\
 \hline
 -1-1 \\
 \hline
 \cdot +1 \\
 +1+1 \\
 \hline
 \cdot -1 \\
 -1-1 \\
 \hline
 \cdot +1 \text{ remainder.} \\
 \hline
 \end{array}$$

(Key to Vol. I. page 205.)

Compound Surds reduced in Infinite Series.

Ex. 2. $\frac{1}{1} + 1(1 + \frac{1}{2} - \frac{1}{8} + \frac{1}{16} - \frac{1}{128} + \&c. \text{ Ans.}$

$$\frac{2 + \frac{1}{2}) \cdot +1}{+1 + \frac{1}{4}}$$

$$\frac{2 + 1 - \frac{1}{8}) \cdot -\frac{1}{4}}{-\frac{1}{4} - \frac{1}{8} + \frac{1}{64}}$$

$$\cdot \frac{+\frac{1}{8} - \frac{1}{64} \cdot \&c.}{\underline{\hspace{2cm}}}$$

Ex. 3. $\frac{1}{1} - 1(1 - \frac{1}{2} - \frac{1}{8} - \frac{1}{16} - \frac{1}{128} - \&c. \text{ Root required.}$

$$\frac{2 - \frac{1}{2}) \cdot -1}{-1 + \frac{1}{4}}$$

$$\frac{2 - 1 - \frac{1}{8}) \cdot -\frac{1}{4}}{-\frac{1}{4} + \frac{1}{8} + \frac{1}{64}}$$

$$\cdot \frac{-\frac{1}{8} - \frac{1}{64} \cdot \&c.}{\underline{\hspace{2cm}}}$$

Ex. 4. $a^2 + x(a + \frac{x}{2a} - \frac{x^2}{8a^3} + \frac{x^3}{16a^5} - \&c. \text{ Root in the series.}$

$$\frac{a^2}{2a + \frac{x}{2a}) \cdot +x}{+x + \frac{x^2}{4a^2}}$$

$$\frac{2a + \frac{x}{a} - \frac{x^2}{8a^3}) \cdot -\frac{x^2}{4a^2}}{-\frac{x^2}{4a^2} - \frac{x^3}{8a^4} + \frac{x^4}{64a^6}}$$

$$\cdot \frac{+\frac{x^3}{8a^4} - \frac{x^4}{64a^6} \cdot \&c.}{\underline{\hspace{2cm}}}$$

(Key to Vol. I. page 205.)

Ex. 5.
$$a^2 - 2bx - x^2 \left(a - \frac{2bx + x^2}{2a} - \frac{4b^2x^2 + 4bx^3 + x^4}{8a^3} - \&c. \right)$$

$$\frac{a^2}{2a - \frac{2bx + x^2}{2a}} = \frac{2bx - x^2}{-2bx - x^2 + \frac{4b^2x^2 + 4bx^3 + x^4}{4a^2}}$$

$$\frac{4b^2x^2 + 4bx^3 + x^4}{4a^2} \cdot \&c.$$

Wherefore the series is $a - \frac{2bx + x^2}{2a} - \frac{4b^2x^2 + 4bx^3 + x^4}{8a^3} \&c.$ IN INFINITUM.

(Page 207.)

The Extraction of Roots of Binomial Quantities, and the Conversion of Binomial Surds to Infinite Series.

Ex. 3. Here $\frac{a^2}{a-x} = a^2(a-x)^{-1}$, and

$$P = a,$$

$$Q = \frac{-x}{a},$$

$$\frac{m}{n} = \frac{-1}{1}; \text{ therefore}$$

$$P \frac{m}{n} = (a)^{-1} = a^{-1} = \frac{1}{a} = A, \text{ the 1st term of a series.}$$

$$\frac{m}{n} A Q = \frac{-1}{1} \times \frac{1}{a} \times \frac{-x}{a} = \frac{x}{a^2} = B, \text{ the 2d term.}$$

$$\frac{m-n}{2n} B Q = \frac{-2}{2} \times \frac{x}{a^2} \times \frac{-x}{a} = \frac{2x^2}{2a^3} = \frac{x^2}{a^3} = C, \text{ 3d term.}$$

$$\frac{m-2n}{3n} C Q = \&c. \&c. \quad \frac{x^3}{a^4} = D, \&c.$$

(Key to Vol. I. page 207.)

$$\text{But } a^2 \times \left(\frac{1}{a} + \frac{x}{a^2} + \frac{x^2}{a^3} + \frac{x^3}{a^4} + \frac{x^4}{a^5} + \frac{x^5}{a^6} + \&c. \right) = a + x + \frac{x^2}{a} + \frac{x^3}{a^2} + \frac{x^4}{a^3} + \frac{x^5}{a^4} + \&c. \text{ Ans.}$$

$$\text{Ex. 4. } \left(\frac{1}{a^2+x^2} \right)^{\frac{1}{2}} = 1 \times (a^2+x^2)^{-\frac{1}{2}} = (a^2+x^2)^{-\frac{1}{2}}.$$

$$\text{Also } P = a^2,$$

$$Q = \frac{x^2}{a^2},$$

$$\frac{m}{n} = \frac{-1}{2}; \text{ therefore}$$

$$P \frac{m}{n} = (a^2)^{-\frac{1}{2}} = \frac{1}{a} = A, \text{ the 1st term of the series sought.}$$

$$\frac{m}{n} \text{AQ} = \frac{-1}{2} \times \frac{1}{a} \times \frac{x^2}{a^2} = \frac{-x^2}{2a^3} = B, \text{ the 2d term.}$$

$$\frac{m-n}{2n} \text{BQ} = \frac{-3}{4} \times \frac{-x^2}{2a^3} \times \frac{x^2}{a^2} = \frac{3x^4}{8a^5} = C, \text{ the 3d term, \&c.}$$

$$\text{Hence the series is } \frac{1}{a} - \frac{x^2}{2a^3} + \frac{3x^4}{8a^5} - \frac{5x^6}{16a^7} + \&c. \text{ IN INFINITUM.}$$

$$\text{Ex. 5. Here } \frac{a^2}{(a-b)^2} = a^2 \times (a-b)^{-2}. \text{ and}$$

$$P = a,$$

$$Q = \frac{-b}{a},$$

$$\frac{m}{n} = \frac{-2}{1}; \text{ therefore}$$

$$P \frac{m}{n} = (a)^{-2} = \frac{1}{a^2} = A, \text{ the 1st term of a series to be } \times^{\text{ed}} \text{ by } a^2.$$

$$\frac{m}{n} \text{AQ} = \frac{-2}{1} \times \frac{1}{a^2} \times \frac{-b}{a} = \frac{2b}{a^3} = B, \text{ the 2d term.}$$

$$\frac{m-n}{2n} \text{BQ} = \frac{-3}{2} \times \frac{2b}{a^3} \times \frac{-b}{a} = \frac{3b^2}{a^4} = C, \text{ the 3d term, \&c.}$$

$$\text{And } a^2 \times \left(\frac{1}{a^2} + \frac{2b}{a^3} + \frac{3b^2}{a^4} + \&c. \right) = 1 + \frac{2b}{a} + \frac{3b^2}{a^2} + \frac{4b^3}{a^3} + \frac{5b^4}{a^4} + \&c. \text{ Ans}$$

(Key to Vol. I. page 207.)

Ex. 6. Here, $P=a^2$,

$$Q = \frac{-x^2}{a^2}, \text{ and}$$

$$\frac{m}{n} = \frac{1}{2}; \text{ therefore}$$

$$P^{\frac{m}{n}} = (a^2)^{\frac{1}{2}} = a = A.$$

$$\frac{m}{n}AQ = \frac{1}{2} \times \frac{a}{1} \times \frac{-x^2}{a^2} = \frac{-x^2}{2a} = B. \text{ \&c.}$$

Hence $a - \frac{x^2}{2a} - \frac{x^4}{8a^3} - \frac{x^6}{16a^5} - \frac{5x^8}{128a^7} - \text{\&c.}$ the Root reqd.

Ex. 7. Here, $P=a^3$,

$$Q = \frac{-b^3}{a^3}, \text{ and}$$

$$\frac{m}{n} = \frac{1}{3}; \text{ therefore}$$

$$P^{\frac{m}{n}} = (a^3)^{\frac{1}{3}} = a = A, \text{ the 1st term of the series.}$$

$$\frac{m}{n}AQ = \frac{1}{3} \times \frac{a}{1} \times \frac{-b^3}{a^3} = \frac{-b^3}{3a^2} = B, \text{ the 2d term, \&c.}$$

Hence $a - \frac{b^3}{3a^2} - \frac{b^6}{9a^5} - \frac{5b^9}{81a^8} - \text{\&c.}$ the Cube Root required.

Ex. 8. In this Example, $P=a^5$, $Q=\frac{x^5}{a^5}$, and $\frac{m}{n}=\frac{1}{5}$; therefore

$$P^{\frac{m}{n}} = (a^5)^{\frac{1}{5}} = a = A, \text{ the 1st term of the series.}$$

$$\frac{m}{n}AQ = \frac{1}{5} \times \frac{a}{1} \times \frac{x^5}{a^5} = \frac{x^5}{5a^4} = B, \text{ the 2d term.}$$

$$\frac{m-n}{2n}BQ = \frac{-4}{10} \times \frac{x^5}{5a^4} \times \frac{x^5}{a^5} = \frac{-4x^{10}}{50a^9} = \frac{-2x^{10}}{25a^9} = C, \text{ the 3d term.}$$

Wherefore $a + \frac{x^5}{5a^4} - \frac{2x^{10}}{25a^9} + \frac{6x^{15}}{125a^{14}} \text{\&c.}$ Ans.*

* The sign of the 3d Term of the Answer given in the Course is different from the sign here given.

(Key to Vol. I. page 208.)

$$\text{Ex. 9. } \sqrt{\frac{a^2 - b^2}{a^2 + b^2}} = (a^2 - b^2)^{\frac{1}{2}} \times (a^2 + b^2)^{-\frac{1}{2}}.$$

$$\text{Now } (a^2 - b^2)^{\frac{1}{2}} = a - \frac{b^2}{2a} - \frac{b^4}{8a^3} - \&c.$$

$$\text{And } (a^2 + b^2)^{-\frac{1}{2}} = \frac{1}{a} - \frac{b^2}{2a^3} + \frac{b^4}{8a^5} \&c. \text{ Whence}$$

$$1 - \frac{b^2}{a^2} + \frac{b^4}{2a^4} - \frac{b^6}{2a^6} \&c. \text{ the Square Root required.}$$

But, as this method is tedious and complex, it is better to extract the Square Root of the Quotient in an Infinite Series. Thus,

$$a^2 + b^2) a^2 - b^2 (1 - \frac{2b^2}{a^2} + \frac{2b^4}{a^4} - \frac{2b^6}{a^6} + \frac{2b^8}{a^8} - \frac{2b^{10}}{a^{10}} + \frac{2b^{12}}{a^{12}} - \&c. \text{ Quot.}$$

$$\text{And } \sqrt{1 - \frac{2b^2}{a^2} + \frac{2b^4}{a^4} - \&c.} = 1 - \frac{b^2}{a^2} + \frac{b^4}{2a^4} - \frac{b^6}{2a^6} + \&c. \text{ Ans.*}$$

as before.

* This Answer, however, differs from that given by Dr. Hutton.

$$\text{Ex. 10. } \sqrt[3]{\frac{a^3}{a^3 + b^3}} = \sqrt[3]{a^3} \times (a^3 + b^3)^{-\frac{1}{3}} = a(a^3 + b^3)^{-\frac{1}{3}}.$$

$$P = a^3 \left\{ \begin{array}{l} \frac{m}{n} = (a^3)^{-\frac{1}{3}} = \frac{1}{\sqrt[3]{a^3}} = \frac{1}{a} = A, \text{ the 1st term of a series.} \\ \frac{m}{n} \wedge Q = \frac{-1}{3} \times \frac{1}{a} \times \frac{b^3}{a^3} = \frac{-b^3}{3a^4} = B, \text{ the 2d term.} \\ \frac{m}{n} = \frac{-1}{3} \left\{ \begin{array}{l} \frac{m-n}{2n} BQ = \frac{-4}{6} \times \frac{-b^3}{3a^4} \times \frac{b^3}{a^3} = \frac{2b^6}{9a^7} = C, \text{ the 3d term.} \\ \frac{m-2n}{3n} CQ = \frac{-7}{9} \times \frac{2b^6}{9a^7} \times \frac{b^3}{a^3} = \frac{-14b^9}{81a^{10}} = D, \text{ the 4th term, \&c.} \end{array} \right. \end{array} \right.$$

$$Q = \frac{b^3}{a^3} \left\{ \begin{array}{l} \frac{m}{n} \wedge Q = \frac{-1}{3} \times \frac{1}{a} \times \frac{b^3}{a^3} = \frac{-b^3}{3a^4} = B, \text{ the 2d term.} \\ \frac{m}{n} = \frac{-1}{3} \left\{ \begin{array}{l} \frac{m-n}{2n} BQ = \frac{-4}{6} \times \frac{-b^3}{3a^4} \times \frac{b^3}{a^3} = \frac{2b^6}{9a^7} = C, \text{ the 3d term.} \\ \frac{m-2n}{3n} CQ = \frac{-7}{9} \times \frac{2b^6}{9a^7} \times \frac{b^3}{a^3} = \frac{-14b^9}{81a^{10}} = D, \text{ the 4th term, \&c.} \end{array} \right. \end{array} \right.$$

$$\frac{m}{n} = \frac{-1}{3} \left\{ \begin{array}{l} \frac{m-n}{2n} BQ = \frac{-4}{6} \times \frac{-b^3}{3a^4} \times \frac{b^3}{a^3} = \frac{2b^6}{9a^7} = C, \text{ the 3d term.} \\ \frac{m-2n}{3n} CQ = \frac{-7}{9} \times \frac{2b^6}{9a^7} \times \frac{b^3}{a^3} = \frac{-14b^9}{81a^{10}} = D, \text{ the 4th term, \&c.} \end{array} \right.$$

$$\frac{m-2n}{3n} CQ = \frac{-7}{9} \times \frac{2b^6}{9a^7} \times \frac{b^3}{a^3} = \frac{-14b^9}{81a^{10}} = D, \text{ the 4th term, \&c.}$$

$$\text{But } a \times \left(\frac{1}{a} - \frac{b^3}{3a^4} + \frac{2b^6}{9a^7} - \&c. \right) = 1 - \frac{b^3}{3a^3} + \frac{2b^6}{9a^6} - \frac{14b^9}{81a^9} + \&c. \text{ Ans.*}$$

* By which it may be perceived that there is an Error in the Numerator of the 3d Term of Hutton's Answer.

(Key to Vol. I. page 209.)

ARITHMETICAL PROPORTION.

Ex. 3. The 1st term is 1, the last term 100, also the number of terms 100.

But $(1+100) \times 50 = 5050$ *Sum required.*

(Page 210.)

Ex. 4. $(98 \times 2) + 1 = 197$ the last term of the series.

And $\frac{1+197}{2} \times 99 = 9801$ *the Sum required.*

Ex. 5. $10 - (\frac{1}{3} \times 20) = 3\frac{1}{3}$ the 1st term of the series.

And $\frac{10+3\frac{1}{3}}{2} \times 21 = 140$ *the Sum required.*

Ex. 6. Here the 1st term is 4 *yds.*, the last term 400 *yds.*, and the number of terms 100.

But $\begin{matrix} \text{yds.} & \text{yds.} & \text{--- yds.} & \text{miles} & \text{yds.} \\ (4+400) \times 50 = 20200 = 11 \text{ .. } 840. & \text{Ans.} \end{matrix}$

APPLICATION OF ARITHMETICAL PROGRESSION TO MILITARY AFFAIRS.

QUESTION I.

The 1st term is 1, the com. diff. 2, and the number of terms 30.

But $\begin{matrix} \text{men} & \text{men} & \text{man} & \text{men} \\ (2 \times 29) + 1 = 59 \end{matrix}$ the last term, or *number of men in the last rank.*

And $\frac{1+59}{2} \times 30 = 900$ men, STRENGTH of the BATTALION. Ans.

(Key to Vol. I. page 211.)

QUESTION II.

The 1st term is 2 , the com. diff. $1\frac{1}{2}$, and the number of terms 12 .

But $(1\frac{1}{2} \times 11) + 2 = 18\frac{1}{2}$ leagues, *last day's march*.

And $(2 + 18\frac{1}{2}) \times 6 (= \text{half } n^{\circ} \text{ terms}) = 123$ leagues, LENGTH of the MARCH. Ans.

QUESTION III.

The 1st term is 15 , the last term 3 , and the com. diff. 2 .

But $\frac{15 - 3 + 2}{2} = 7$ the number of terms, or *nights employed*.

And $\frac{3 + 15}{2} \times 7 = 63$ yards, LENGTH of the WHOLE SAP. Ans.

(Page 212.)

QUESTION IV.

The 1st rank is 4 , the last rank 9 , and the number of ranks 6 .

But $\frac{9 - 4}{6 - 1} = 1$ the common difference.

And $(4 + 9) \times 3 = 39$ the NUMBER of GABIONS. Ans.

QUESTION V.

If the straight line joining the two Camps pass through the Post in question, it is evident that $18\frac{1}{2}$ leagues is the Sum of the series of either detachment.

(Key to Vol. I. page 212.)

Wherefore, for the 1st Detachment,

$$\left. \begin{array}{l} \text{leag.} \\ (18\frac{1}{2} \times 2) \div 5 (= \text{No. of terms}) = 7\frac{2}{5} \text{ the sum of the Extremes.} \\ (1\frac{1}{2} \times 5) - 1\frac{1}{2} = 6 \text{ the difference of the Extremes.} \end{array} \right\}$$

Hence (by $\frac{1}{2}$ sum and $\frac{1}{2}$ diff. &c.) $\frac{7}{10}$, $2\frac{2}{10}$, $3\frac{7}{10}$, $5\frac{2}{10}$, $6\frac{7}{10}$ the SERIES.

For the 2d Detachment,

$$\left. \begin{array}{l} \text{leag.} \\ (18\frac{1}{2} \times 2) \div 4 (= \text{No. of terms}) = 9\frac{1}{4} \text{ the sum of the Extremes.} \\ (2 \times 4) - 2 = 6 \text{ the difference of the Extremes.} \end{array} \right\}$$

Hence (by half sum and half diff. &c.) $1\frac{5}{8}$, $3\frac{5}{8}$, $5\frac{5}{8}$, $7\frac{5}{8}$ the SERIES.

* * * But as this Post may be any where in a line passing at right angles through the middle of the right line joining the two Camps, the Question admits of an Infinity of Answers.

QUESTION VI.

It is manifest that the sum of the series of the detachment must be some multiple of 8, and at the same time equal to 8 times the number of terms; therefore, constructing the Series, it will be,

	<i>Days</i>	1st.	2d.	3d.	4th.	5th.	6th.	7th.	8th.	&c.
Series in Leagues,		2,	5,	8,	11,	14,	17,	20,	23,	&c.

	<i>Daily</i>									
<i>Sum of the Series,</i>		2,	7,	15,	26,	40,	57,	77,	100,	&c.

	<i>Daily</i>									
<i>Progress of the Deserter,</i>		8,	16,	24,	32,	40,	48,	56,	64,	&c.

By which it appears, that no term except the 5th can answer the conditions given.

Consequently 5 days, and 40 leagues. Ans.

ALGEBRAICALLY.

The 1st term is 2, the com. diff. 3, and if x be put for the number of terms, it will be, $3(x-1)+2=3x-1$ the last term of the SERIES.

And, $\frac{2+3x-1}{2} \times x = \frac{3x+1}{2} \times x = \frac{3x^2+x}{2}$ the Sum of the SERIES.

(Key to Vol. I, page 212.)

But (by the Quest.) $\frac{3x^2+x}{2} = 8x$, that is, $3x^2+x=16x$, or,
(dividing by x) $3x+1=16$, hence $x=5$ days.

Consequently, as before, 5 days, and 40 leagues. Ans.

(Page 213.)

QUESTION VII.

The 1st term is $\frac{1}{2}$, the last $9\frac{1}{2}$, and the number of terms 7.
leag. leag. leag.

Also $\frac{9\frac{1}{2}-\frac{1}{2}}{7-1} = 1\frac{1}{2}$ the common difference.

Consequently,

Series in Leagues.	The Convoy had advanced in Leagues.	Days.	The Escort had advanced in Leagues.	Dist. of the Escort from the Convoy, at the end of each March. Leagues.	Nights.
$\frac{1}{2}$	5	1	$\frac{1}{2}$	$4\frac{1}{2}$	1st
2	10	2	$2\frac{1}{2}$	$7\frac{1}{2}$	2d
$3\frac{1}{2}$	15	3	6	9	3d
5	20	4	11	9	4th
$6\frac{1}{2}$	25	5	$17\frac{1}{2}$	$7\frac{1}{2}$	5th
8	30	6	$25\frac{1}{2}$	$4\frac{1}{2}$	6th
$9\frac{1}{2}$	35	7	35	0	7th

Whereby the Answer is obvious.

GEOMETRICAL PROPORTION.

(Page 220.)

Ex. 3. $3^{11} = 177147$

And $\frac{(3 \times 177147) - 1}{3 - 1} = 265720$. Ans.

Ex. 4. $\left(\frac{1}{3}\right)^{11} = \frac{1}{177147}$.

And $1 - \left(\frac{1}{3} \times \frac{1}{177147}\right) = \frac{531440}{531441}$.

Lastly $\frac{531440}{531441} \div \frac{2}{3} = \frac{1594320}{1062882} = \frac{265720}{177147}$. Ans.

(Key to Vol. I. page 226.)

Ex. 2.	Given.	$6x - 15 = x + 6$
	From both sides take x .	$5x - 15 = 6$
	Transp. -15 .	$5x = 6 + 15 = 21$
	Divide by 5.	$x = \frac{21}{5} = 4\frac{1}{5}$ Ans.

Ex. 3.	Given.	$8 - 3x + 12 = 30 - 5x + 4$
	From both sides take $-3x + 4$.	$8 + 8 = 30 - 2x$
	Collect and transp. $-2x$.	$2x + 16 = 30$
	Transp. 16.	$2x = 30 - 16 = 14$
	Divide by 2.	$x = 7$ Ans.

Ex. 4.	Given.	$x + \frac{x}{3} - \frac{x}{4} = 13$
	\times^{ly} by 3.	$3x + x - \frac{3x}{4} = 39$
	\times^{ly} by 4.	$12x + 4x - 3x = 156$
	Collect.	$13x = 156$
	Divide by 13.	$x = 12$ Ans.

Ex. 5.	Given.	$3x + \frac{x}{2} + 2 = 5x - 4$
	Take $3x$ from both sides.	$\frac{x}{2} + 2 = 2x - 4$
	Transp. -4 .	$\frac{x}{2} + 6 = 2x$
	\times^{ply} by 2.	$x + 12 = 4x$
	From both sides take x .	$12 = 3x$
	Divide by 3.	$4 = x$, or $x = 4$ Ans.

Ex. 6.	Given.	$4ax + \frac{a}{3} - 2 = ax - bx$
	\times^{ly} by 3.	$12ax + a - 6 = 3ax - 3bx$
	From both sides take $3ax$.	$9ax + a - 6 = -3bx$
	Transp. $-3bx$, and $a - 6$.	$9ax + 3bx = 6 - a$
	Divide by $9a + 3b$.	$x = \frac{6 - a}{9a + 3b}$ Ans.

(Key to Vol. I. page 226.)

Ex. 7.

Given.	$\frac{x}{3} - \frac{x}{4} + \frac{x}{5} = \frac{1}{2}$	
\times^{ly} by 3.	$x - \frac{3x}{4} + \frac{3x}{5} = \frac{3}{2}$	
\times^{ly} by 4.	$4x - 3x + \frac{12x}{5} = 6$	
Collect, and \times^{ly} by 5.	$5x + 12x = 30$	
Collect and divide by 17.	$x = \frac{30}{17} = 1\frac{13}{17}$	Ans.

Ex. 8.

Given.	$\sqrt{4+x} = 4 - \sqrt{x}$	
Square both sides.	$4+x = 16 - 8\sqrt{x} + x$	
From both sides take x .	$4 = 16 - 8\sqrt{x}$	
Transp. $- 8\sqrt{x}$, and 4.	$8\sqrt{x} = 16 - 4 = 12$	
Square both sides.	$64x = 144$	
Divide by 64.	$x = 2\frac{1}{4}$	Ans.

Ex. 9.

Given.	$4a+x = \frac{x^2}{4a+x}$	
\times^{ly} by $4a+x$.	$16a^2 + 8ax + x^2 = x^2$	
Take x^2 from both sides, } and transp. $16a^2$. }	$8ax = -16a^2$	
Divide by $8a$.	$x = -2a$	Ans.

Ex. 10.

Given.	$\sqrt{4a^2+x^2} = \sqrt[4]{4b^4+x^4}$	
Square both sides.	$4a^2 + x^2 = \sqrt[2]{4b^4+x^4}$	
Square both sides.	$16a^4 + 8a^2x^2 + x^4 = 4b^4 + x^4$	
From both sides take x^4 .	$16a^4 + 8a^2x^2 = 4b^4$	
Transp. $16a^4$.	$8a^2x^2 = 4b^4 - 16a^4$	
Divide by 4.	$2a^2x^2 = b^4 - 4a^4$	
Divide by $2a^2$.	$x^2 = \frac{b^4 - 4a^4}{2a^2}$	
Extract the Sq. Root } on both sides. }	$x = \sqrt{\frac{b^4 - 4a^4}{2a^2}}$	Ans.

(Key to Vol. I. page 226.)

Ex. 11.	Given.	$\sqrt{x + \sqrt{2a+x}}$	$= \frac{4a}{\sqrt{2a+x}}$
	\times by $\sqrt{2a+x}$.	$\sqrt{2ax+x^2} + 2a+x$	$= 4a$
	Transp. $2a+x$.	$\sqrt{2ax+x^2}$	$= 4a - 2a - x = 2a - x$
	Square both sides.	$2ax+x^2$	$= 4a^2 - 4ax + x^2$
	Cancel x^2 .	$2ax$	$= 4a^2 - 4ax$
	Transp. $-4ax$.	$6ax$	$= 4a^2$
	Divide by $6a$.	x	$= \frac{4a^2}{6a} = \frac{2}{3}a$. Ans.

Ex. 12.	Given.	$\frac{a}{1+2x} + \frac{a}{1-2x}$	$= 2b$
	Reduce the fractions.	$\frac{2a}{1-4x^2}$	$= 2b$
	\times by $1-4x^2$.	$2a$	$= 2b - 8bx^2$
	Transpose $-8bx^2$, and $2a$.	$8bx^2$	$= 2b - 2a$
	Divide by $8b$.	x^2	$= \frac{b-a}{4b} = \frac{1}{4} \left(\frac{b-a}{b} \right)$
	Extr. the Square Root on } both sides. }	x	$= \frac{1}{2} \sqrt{\frac{b-a}{b}}$. Ans.

Ex. 13.	Given.	$a+x$	$= \sqrt{a^2+x} \sqrt{4b^2+x^2}$
	Square both sides.	$a^2+2ax+x^2$	$= a^2+x \sqrt{4b^2+x^2}$
	Cancel a^2 , and divide by x .	$2a+x$	$= \sqrt{4b^2+x^2}$
	Square both sides.	$4a^2+4ax+x^2$	$= 4b^2+x^2$
	Cancel x^2 .	$4a^2+4ax$	$= 4b^2$
	Transp. $4a^2$, and div. by 4 .	ax	$= b^2 - a^2$
	Divide by a .	x	$= \frac{b^2}{a} - a$. Ans.

(Page 228.)

To exterminate Two Unknown Quantities.

RULE. I.

Ex. 2.	Given*	$\left\{ \begin{array}{l} \frac{x}{2} + 2y = a \\ \frac{x}{2} - 2y = b \end{array} \right\}$
--------	--------	--

* This Equation belongs more properly to Rule III, admitting there of a much Shorter and more Elegant Solution.

(Key to Vol. I. page 228.)

In the 1st equat. transp. $2y$, and \times^{ly} by 2.	$x = 2a - 4y$
In the 2d equat. transp. $-2y$, and \times^{ly} by 2.	$x = 2b + 4y$
Put these values of x equal to one another.	$2a - 4y = 2b + 4y$
Transpose $-4y$ and $2b$.	$2a - 2b = 8y$
Divide by 8.	$\frac{a-b}{4} = y$

But $x = 2a - 4y = 2a - (a - b) = a + b$, and the Answer *obvious*.

Ex. 3. Given. * $\left\{ \begin{array}{l} 3x + y = 22 \\ x + 3y = 18 \end{array} \right\}$

In the first equat. transpose $3x$.	$y = 22 - 3x$
In the 2d equat. tr. x , and divide by 3.	$y = \frac{18 - x}{3}$
Put these values of y equal to one another.	$22 - 3x = \frac{18 - x}{3}$
Multiply by 3.	$66 - 9x = 18 - x$
Cancel $-x$, and transpose.	$66 - 18 = 8x$
Collect and divide by 8.	$6 = x$

But $y = 22 - 3x = 22 - 18 = 4$.

Wherefore $x = 6$, and $y = 4$. Ans.

* The Observation made on the last Equation is equally applicable to the present.

Ex. 4.† Given $\left\{ \begin{array}{l} \frac{1}{2}x + \frac{1}{3}y = 4 \\ \frac{1}{3}x + \frac{1}{2}y = 3\frac{1}{2} \end{array} \right\}$

In the 1st equat. tr. $\frac{1}{2}x$, and \times^{ly} by 3.	$y = 12 - \frac{3x}{2}$
In the 2d equat. tr. $\frac{1}{3}x$, and \times^{ly} by 2.	$y = 7 - \frac{2x}{3}$
Put these values of y equal to one another.	$12 - \frac{3x}{2} = 7 - \frac{2x}{3}$
Take 7 from both sides.	$5 - \frac{3x}{2} = -\frac{2}{3}x$
Multiply by 2, and by 3.	$30 - 9x = -4x$
Cancel $-4x$ on both sides, and tr.	$30 = 5x$
Divide by 5.	$6 = x$

† This Example and the next are likewise Equations in the Third Rule.

(Key to Vol. I. page 228.)

But $y = 12 - \frac{3x}{2} = 12 - \frac{18}{2} = 12 - 9 = 3$, and the Ans. *obvious*.

Ex. 5.

Given $\left\{ \begin{array}{l} \frac{2x}{3} + \frac{3y}{5} = \frac{22}{5} \\ \frac{3x}{5} + \frac{2y}{3} = \frac{67}{15} \end{array} \right\}$

In the 1st equat. tr. $\frac{2x}{3}$, and \times by 5. $3y = 22 - \frac{10x}{3}$

Divide by 3. $y = \frac{22}{3} - \frac{10x}{9}$

In the 2d equat. tr. $\frac{3x}{5}$, and div. by $\frac{2}{3}$. $y = \frac{67}{10} - \frac{9}{10}x$

Put these values of y equal to one another. $\frac{67}{10} - \frac{9}{10}x = \frac{22}{3} - \frac{10x}{9}$

Reduce. $603 - 81x = 660 - 100x$

Cancel $81x$ on both sides. $603 = 660 - 19x$

Take 603 from both sides. $0 = 57 - 19x$

Transpose, and divide by 19. $x = 3$

But $y = \frac{22}{3} - \frac{10x}{9} = \frac{22}{3} - \frac{30}{9} = \frac{22}{3} - \frac{10}{3} = \frac{12}{3} = 4$, and the Ans. *obvious*.

Ex. 6.

Given $\left\{ \begin{array}{l} x + 2y = s \\ x^2 - 4y^2 = d^2 \end{array} \right\}$

In the 1st equat. tr. $2y$. $x = s - 2y$

In the 2d equat. tr. $-4y^2$, and extr. $\sqrt{\quad}$. $x = \sqrt{d^2 + 4y^2}$

Put these values of x equal to one another, and Square. $s^2 - 4sy + 4y^2 = d^2 + 4y^2$

Cancel $4y^2$. $s^2 - 4sy = d^2$

Transp. $-4sy$, and d^2 . $s^2 - d^2 = 4sy$

Divide by $4s$. $\frac{s^2 - d^2}{4s} = y$

But $x + 2y = s$, that is, $x + \frac{s^2 - d^2}{2s} = s$

Transpose $\frac{s^2 - d^2}{2s}$ $x = s - \frac{s^2 - d^2}{2s} = \frac{2s^2 - s^2 + d^2}{2s} = \frac{s^2 + d^2}{2s}$

Therefore $x = \frac{s^2 + d^2}{2s}$, and $y = \frac{s^2 - d^2}{4s}$. Ans.

(Key to Vol. I. page 228.)

Ex. 7.

$$\text{Given } \left\{ \begin{array}{l} x - 2y = d \\ x : y :: a : b \end{array} \right\}$$

$$\text{By the 1st equat. } \left| \begin{array}{l} x = d + 2y \end{array} \right.$$

$$\text{By the ratio given. } \left| \begin{array}{l} x = \frac{ay}{b} \end{array} \right.$$

$$\text{Therefore (Eucl. i. Ax. 1.) } \left| \begin{array}{l} \frac{ay}{b} = d + 2y \end{array} \right.$$

$$\times^{\text{ly}} \text{ by } b. \left| \begin{array}{l} ay = bd + 2by \end{array} \right.$$

$$\text{Transp. } 2by. \left| \begin{array}{l} ay - 2by = bd \end{array} \right.$$

$$\text{Divide by } a - 2b. \left| \begin{array}{l} y = \frac{bd}{a - 2b} \end{array} \right.$$

$$\text{But } x = d + 2y = d + \frac{2bd}{a - 2b} = \frac{ad - 2bd + 2bd}{a - 2b} = \frac{ad}{a - 2b}$$

Wherefore the values of x and y are known.

(Page 229.)

RULE II.

Ex. 2.

$$\text{Given } \left\{ \begin{array}{l} 2x + 3y = 29 \\ 3x - 2y = 11 \end{array} \right\}$$

$$\text{In the 1st equat. } \left| \begin{array}{l} x = \frac{29 - 3y}{2} \end{array} \right.$$

$$\text{Subst. this val. of } x \text{ for } x \text{ in the 2d equat. } \left| \begin{array}{l} \frac{87 - 9y}{2} - 2y = 11 \end{array} \right.$$

$$\times^{\text{ly}} \text{ by } 2, \text{ collect, tr. and div. by } 13.$$

Wherefore, by step 1st.

$$\left. \begin{array}{l} 5 = y \\ \frac{29 - 15}{2} = 7 = x \end{array} \right\} \text{Ans.}$$

Ex 3.

$$\text{Given* } \left\{ \begin{array}{l} x + y = 14 \\ x - y = 2 \end{array} \right\}$$

$$\text{In the 1st equation. } \left| \begin{array}{l} x = 14 - y \end{array} \right.$$

$$\text{Subst. this val. of } x \text{ for } x \text{ in the 2d equat. } \left| \begin{array}{l} 14 - y - y = 2 \end{array} \right.$$

$$\text{Tr. collect, and divide by } 2. \left| \begin{array}{l} 6 = y \end{array} \right\} \text{Ans.}$$

$$\text{And, by step 1st. } \left| \begin{array}{l} 8 = x \end{array} \right\}$$

* This Equation solved by the Rule is neither a short nor elegant Operation.

For

Add the given Equations, and divide by 2.

Subtract the 2d Equat. from the 1st, and div. by 2. $\left. \begin{array}{l} x = 8 \\ y = 6 \end{array} \right\} \text{Ans.}$

(Key to Vol. I. page 230.)

Ex. 4.

$$\text{Given } \left\{ \begin{array}{l} x : y :: 3 : 2 \\ x^2 - y^2 = 20 \end{array} \right\}$$

$$\text{By the given ratio. } \left| \begin{array}{l} x = \frac{3y}{2} \end{array} \right.$$

$$\text{Square both sides. } \left| \begin{array}{l} x^2 = \frac{9y^2}{4} \end{array} \right.$$

$$\text{Subst. this val. of } x^2 \text{ for } x^2 \text{ in the given equat. } \left| \begin{array}{l} \frac{9y^2}{4} - y^2 = 20 \end{array} \right.$$

$$\times^{\text{ly}} \text{ by 4, and collect. } \left| \begin{array}{l} 5y^2 = 80 \end{array} \right.$$

$$\text{Divide by 5, and extr. } \sqrt{} \left| \begin{array}{l} y = 4 \end{array} \right.$$

$$\text{But, by the Quest. } \left| \begin{array}{l} x : y, \text{ or } 4 :: 3 : 2 \end{array} \right.$$

$$\text{Wherefore, } \left| \begin{array}{l} x = 6 \end{array} \right.$$

Ex. 5.

$$\text{Given } \left\{ \begin{array}{l} \frac{x}{3} + 3y = 21 \\ \frac{y}{3} + 3x = 29 \end{array} \right\}$$

$$\text{In the 1st equat. tr. } 3y, \text{ and } \times^{\text{ly}} \text{ by 3. } \left| \begin{array}{l} x = 63 - 9y \end{array} \right.$$

$$\text{Subst. this val. of } x \text{ for } x \text{ in the 2d equat. } \left| \begin{array}{l} \frac{y}{3} + 189 - 27y = 29 \end{array} \right.$$

$$\times^{\text{ly}} \text{ by 3, tr. and collect. } \left| \begin{array}{l} 480 = 80y \end{array} \right.$$

$$\text{Divide by 80. } \left| \begin{array}{l} 6 = y \end{array} \right.$$

$$\text{But, by the 2d equat. } \left. \begin{array}{l} \frac{6}{3} + 3x = 29, \text{ or} \\ 9 = x \end{array} \right\} \text{Ans.}$$

Ex. 6.

$$\text{Given } \left\{ \begin{array}{l} 10 - \frac{x}{2} = \frac{y}{3} + 4 \\ \frac{x-y}{2} + \frac{x}{4} - 2 = \frac{3y-x}{5} - 1 \end{array} \right\}$$

$$\text{In the 1st equat. cancel 4. } \left| \begin{array}{l} 6 - \frac{x}{2} = \frac{y}{3} \end{array} \right.$$

$$\text{Transp. and } \times^{\text{ly}} \text{ by 2. } \left| \begin{array}{l} 12 - \frac{2y}{3} = x \end{array} \right.$$

$$\text{Reduce the 2d equat. } \left| \begin{array}{l} 19x = 22y + 20 \end{array} \right.$$

$$\text{For } x \text{ subst. } 12 - \frac{2}{3}y. \left| \begin{array}{l} 228 - \frac{38y}{3} = 22y + 20 \end{array} \right.$$

$$\times^{\text{ly}} \text{ by 3. } \left| \begin{array}{l} 684 - 38y = 66y + 60 \end{array} \right.$$

$$\text{Transp. and div. by 104. } \left| \begin{array}{l} 6 = y \end{array} \right.$$

$$\text{But (by step. 2d). } \left. \begin{array}{l} 12 - \frac{2}{3}y, \text{ that is, } 12 - 4 = 8 = x \end{array} \right\} \text{Ans.}$$

(Key to Vol. I. page 230.)

Ex. 7.

$$\text{Given } \left\{ \begin{array}{l} x : y :: 4 : 3 \\ x^3 - y^3 = 37 \end{array} \right\}$$

By the given ratio.	$x = \frac{4y}{3}$	
Cube both sides.	$x^3 = \frac{64y^3}{27}$	
Subst. this val. of x^3 for x^3 in the given equat. } \times^{ly} by 27.	$\frac{64y^3}{27} - y^3 = 37$	
Collect and divide by 37.	$64y^3 - 27y^3 = 999$	
Extr. $\sqrt[3]{}$ on both sides.	$y^3 = 27$	
But, by step 1st.	$y = 3$	} Ans.
	$x = \frac{4y}{3} = \frac{12}{3} = 4$	

(Page 231)

RULE III.

Ex. 1.

$$\text{Given } \left\{ \begin{array}{l} \frac{x+8}{4} + 6y = 21 \\ 5x + \frac{y+6}{3} = 23 \end{array} \right\}$$

Reduce the 1st equat.	$x + 24y = 76$	
Reduce the 2d equat.	$15x + y = 63$	
\times^{ly} the 1st equat. red. by 15.	$15x + 360y = 1140$	
From this equat. subtr. the 2d } reduced. }	$359y = 1077$	
Divide by 359.	$y = 3$	} Ans.
But, by step. 1st (subst. 3 for y .)	$x + 72 = 76, \text{ or } x = 4$	

Ex. 2.

$$\text{Given } \left\{ \begin{array}{l} \frac{3x-y}{4} + 10 = 13 \\ \frac{3y+x}{2} + 5 = 12 \end{array} \right\}$$

Reduce the 1st equation	$3x - y = 12$
Reduce the 2d equation	$x + 3y = 14$

(Key to Vol. I. page 231.)

$$\begin{array}{l}
 \times^{\text{ly}} \text{ the 2d equat. reduced, by 3. } 3x+9y=42 \\
 \text{From this equat. take the 1st reduced. } 10y=30 \\
 \text{Divide by 10. } y=3 \\
 \text{But, by step. 2d, subst. 3 for } y. x+9=14, \text{ or, } x=5 \left. \vphantom{\begin{array}{l} 3x+9y=42 \\ 10y=30 \\ y=3 \end{array}} \right\} \text{Ans.}
 \end{array}$$

$$\text{Ex. 3.} \quad \text{Given } \left\{ \begin{array}{l} \frac{3x+4y}{5} + \frac{x}{4} = 10 \\ \frac{6x-2y}{3} + \frac{y}{6} = 14 \end{array} \right\}$$

$$\begin{array}{l}
 \text{Reduce the 1st equat. } 17x+16y=200 \\
 \text{Reduce the 2d equat. } 12x-3y=84 \\
 \times^{\text{ly}} \text{ the 1st equat. red. by 3. } 51x+48y=600 \\
 \times^{\text{ly}} \text{ the 2d equat. red. by 16. } 192x-48y=1344 \\
 \text{Add together the 2 last steps. } 243x=1944 \\
 \text{Divide by 243. } x=8 \\
 \text{But (by the 2d equat. reduced) } 96-3y=84, \text{ or } y=4 \left. \vphantom{\begin{array}{l} 17x+16y=200 \\ 12x-3y=84 \\ 51x+48y=600 \\ 192x-48y=1344 \\ 243x=1944 \\ x=8 \end{array}} \right\} \text{Ans.} \\
 \text{substituting 8 for } x.
 \end{array}$$

$$\text{Ex. 4.} \quad \text{Given } \left\{ \begin{array}{l} 3x+4y=38 \\ 4x-3y=9 \end{array} \right\}$$

$$\begin{array}{l}
 \times^{\text{ly}} \text{ the 1st equat. by 4. } 12x+16y=152 \\
 \times^{\text{ly}} \text{ the 2d equat. by 3. } 12x-9y=27 \\
 \text{Subtr. the 2d step from the 1st. } 25y=125 \\
 \text{Divide by 25. } y=5 \\
 \text{By the question, subst. 5 for } y. 3x+20=38, \text{ or } x=6 \left. \vphantom{\begin{array}{l} 12x+16y=152 \\ 12x-9y=27 \\ 25y=125 \\ y=5 \end{array}} \right\} \text{Ans.}
 \end{array}$$

(Page 233.)

To exterminate Three or more Unknown Quantities.

$$\text{Ex. 2.} \quad \text{Given } \left\{ \begin{array}{l} x+y+z=18 \\ x+3y+2z=38 \\ x+\frac{1}{3}y+\frac{1}{2}z=10 \end{array} \right\} \text{ or } \left\{ \begin{array}{l} x=18-y-z \\ x=38-3y-2z \\ x=10-\frac{1}{3}y-\frac{1}{2}z \end{array} \right\}$$

$$\begin{array}{l}
 \text{Putting the 1st equat. equal to } \left. \vphantom{\begin{array}{l} 2y+z=20 \\ 4y+2z=40 \\ 4y+3z=48 \end{array}} \right\} \\
 \text{the 2d, and reducing. } 2y+z=20 \\
 \times^{\text{ly}} \text{ by 2. } 4y+2z=40 \\
 \text{Putting the 1st equat. equal to } \left. \vphantom{\begin{array}{l} 4y+2z=40 \\ 4y+3z=48 \end{array}} \right\} \\
 \text{the 3d, and reducing. } 4y+3z=48 \\
 \text{By Subtraction. } z=8 \\
 \text{But (by step 2d) substituting 8 for } z, 2y+8=20, \text{ or } y=6 \left. \vphantom{\begin{array}{l} 2y+z=20 \\ 4y+2z=40 \\ 4y+3z=48 \end{array}} \right\} \text{Ans.} \\
 \text{And, subst. 6 for } y \text{ and 8 for } z, x+14=18, \text{ or } x=4
 \end{array}$$

(Key to Vol. I. page 233.)

Ex. 3. Given $\left\{ \begin{array}{l} x + \frac{1}{2}y + \frac{1}{3}z = 27 \\ x + \frac{1}{3}y + \frac{1}{4}z = 20 \\ x + \frac{1}{4}y + \frac{1}{5}z = 16 \end{array} \right\}$ or $\left\{ \begin{array}{l} x = 27 - \frac{1}{2}y - \frac{1}{3}z \\ x = 20 - \frac{1}{3}y - \frac{1}{4}z \\ x = 16 - \frac{1}{4}y - \frac{1}{5}z \end{array} \right\}$

Putting the 1st equat. equal to the 2d. $2y + z = 84$
 Putting the 1st equat. equal to the 3d. $15y + 8z = 660$
 \times ^{ly} the 1st step by 8. $16y + 8z = 672$
 From this equat. subtr. the 2d step. $y = 12$
 But, by step 1st $2y + z = 84$, hence $z = 60$
 And, subtr. 12 for y , and 60 for z &c. $x = 1$ } Ans.

Ex. 4. Given $\left\{ \begin{array}{l} x - y = 2 \\ x - z = 3 \\ y + z = 9 \end{array} \right\}$

Subtr. the 1st equat. from the 2d. $y - z = 1$
 Add this equat. to the 3d. $2y = 10$, or $y = 5$
 Subtr. the 1st step from the 3d } $2z = 8$, or $z = 4$ } Ans.
 equat. }
 But, (by the 2d equat.) $x - 4 = 3$, or $x = 7$

Ex. 5. Given $\left\{ \begin{array}{l} 2x + 3y + 4z = 34 \\ 3x + 4y + 5z = 46 \\ 4x + 6y + 8z = 58 \end{array} \right\}$ or $\left\{ \begin{array}{l} x = \frac{34 - 3y - 4z}{2} \\ x = \frac{46 - 4y - 5z}{3} \\ x = \frac{58 - 6y - 8z}{4} \end{array} \right\}$

Putting the 1st equat. equal to the 2d. $y + 2z = 10$.

But, on comparing the 1st and 3d equations, an absurdity appears, and therefore no three numbers can answer the conditions.

If, however, in the last of the given Equations $5y$ be intended, and if 58 be a mistake for 66 , then, $x = 6$, $y = 2$, and $z = 4$.

NOTE.—In some Editions of the Course the last Equation is read $4x + 5y + 6z = 58$.

(Page 236.)

QUESTIONS FOR PRACTICE.

Quest. I. Let x represent the greater number, and y the less.

By the Quest. $\left\{ \begin{array}{l} x - y = 4 \\ x^2 - y^2 = 64 \end{array} \right\}$ or $\left\{ \begin{array}{l} x = 4 + y \\ x^2 = 64 + y^2 \end{array} \right\}$

(Key to Vol. I. page 236.)

$$\begin{array}{l}
 \text{Square both sides of the 1st equation transposed.} \\
 \text{Compare the two values of } x^2. \\
 \text{Cancel } y^2. \\
 \text{Transpose 16, and divide by 8.} \\
 \text{But } x-y=4, \text{ that is,}
 \end{array}
 \left. \begin{array}{l}
 x^2 = y^2 + 8y + 16 \\
 y^2 + 8y + 16 = 64 + y^2 \\
 8y + 16 = 64 \\
 y = 6 \\
 x = 10
 \end{array} \right\}$$

Consequently 10 and 6 are the numbers sought.

Quest. 2. If x be put for the 1st number, and y for the 2d, it is,

$$\text{By the Quest. } \left\{ \begin{array}{l} \frac{x}{2} + \frac{y}{3} = 9 \\ \frac{x}{4} + \frac{y}{5} = 5 \end{array} \right\}$$

$$\begin{array}{l}
 \text{Reduce the 1st. equat.} \\
 \text{Reduce the 2d equat.} \\
 \times^{\text{ly}} \text{ the 1st equat. reduced, by 2.} \\
 \text{From this equat. take the 2d} \\
 \text{equat. reduced.} \\
 \text{And, because by the 1st equat.} \\
 \text{reduced, } (24 + 2y = 54.)
 \end{array}
 \left. \begin{array}{l}
 3x + 2y = 54 \\
 5x + 4y = 100 \\
 6x + 4y = 108 \\
 x = 8 \\
 y = 15
 \end{array} \right\} \text{The two numbers.}$$

Quest. 3. Put x for one of the parts, and y for the other.

$$\text{By the Quest. } \left\{ \begin{array}{l} x + y = 20 \\ \frac{x}{3} + \frac{y}{5} = 6 \end{array} \right\}$$

$$\begin{array}{l}
 \text{Reduce the 2d equat.} \\
 \times^{\text{ly}} \text{ the 1st equation by 5.} \\
 \text{From this equat. take the} \\
 \text{2d equat. reduced.} \\
 \text{And, by the 1st equation,}
 \end{array}
 \left. \begin{array}{l}
 5x + 3y = 90 \\
 5x + 5y = 100 \\
 2y = 10, \text{ or } y = 5 \\
 20 - y (= 20 - 5) = x = 15
 \end{array} \right\}$$

Wherefore 15 and 5 are the parts required.

Quest. 4. Put x for the 1st number, y for the 2d, and z for the 3d.

$$\text{By the Quest. } \left\{ \begin{array}{l} x + y = 7 \\ x + z = 8 \\ y + z = 9 \end{array} \right\}$$

$$\begin{array}{l}
 \text{Subtr. the 1st equat. from the 2d.} \\
 \text{Add this rem. to the 3d equation.} \\
 \text{Subtr. the 1st step from the 3d equat.} \\
 \text{But } x+4=7, \text{ hence}
 \end{array}
 \left. \begin{array}{l}
 z - y = 1 \\
 2z = 10, \text{ or, } z = 5 \\
 2y = 8, \text{ or, } y = 4 \\
 x = 3
 \end{array} \right\}$$

Ans.

(Key to Vol. I. page 236.)

Quest. 5.* Let x represent the son's portion, and y that of the daughter.

$$\text{By the Quest. } \left\{ \begin{array}{l} x+y=\text{£}2800 \\ x:y::5:2 \end{array} \right\} \text{ or } \left\{ \begin{array}{l} x=\text{£}2800-y \\ 2x=5y \end{array} \right\}$$

$$\begin{array}{l} \text{Mul. the 1st equat. transposed, by 2.} \\ \text{From this equat. subtr. the 2d equat.} \\ \text{Transp. } 7y, \text{ and divide by 7.} \\ \text{Therefore, (by the 1st equat.)} \end{array} \left. \begin{array}{l} 2x=\text{£}5600-2y \\ 0=5600-7y \\ y=800 \\ x=2000 \end{array} \right\} \text{Ans.}$$

* In common Arithmetic this Question admits of the easiest solution, for
 $2+5) \text{£}2800$ ($\text{£}400$ which $\times \left\{ \begin{array}{l} 2=\text{£}800 \text{ the daughter's portion.} \\ 5=2000 \text{ the son's portion.} \end{array} \right.$

$$\begin{array}{r} \text{£}2800 \\ \hline \end{array}$$

Quest. 6.

$$\text{Given } \left\{ \begin{array}{l} A+B+C=\text{£}400 \\ 2A+\text{£}20=B \\ A+B=C \end{array} \right\}$$

$$\begin{array}{l} \text{Substitute B's and C's} \\ \text{contr. in terms of A's.} \\ \text{Collect.} \\ \text{Tr. and divide by 6.} \\ \text{Therefore} \\ \text{And} \end{array} \left. \begin{array}{l} A+(2A+\text{£}20)+(A+2A+\text{£}20)=\text{£}400 \\ 6A+\text{£}40=\text{£}400 \\ A=60 \\ B=140 \\ C=200 \end{array} \right\} \text{Ans.}$$

Otherwise.

It is manifest, by the 3d equat. given, that $c=\text{£}200$
 And (by comparing the 1st and 2d equat^s.) that $3A=150$
 Whence it follows that $A=\text{£}60$, $B=\text{£}140$, and $c=\text{£}200$ as before.

(Page 237.)

Quest. 7. Put x for the number of half guineas, and y for the number of crowns.

$$\begin{array}{l} \text{By the Quest.} \\ \text{And since } (\text{£}100=4000 \text{ sixpences}) \\ \times^{\text{ly}} \text{ the 1st equat. by 21.} \\ \text{From this equat. take the 2d equation.} \\ \text{Divide by 11.} \\ \text{From 202 take 22.} \end{array} \left. \begin{array}{l} x+y=202 \\ 21x+10y=4000 \\ 21x+21y=4242 \\ 11y=242 \\ y=22 \\ x=180 \end{array} \right\}$$

Consequently 22 crowns, and 180 half guineas. Ans.

(Key to Vol. I. page 237.)

Quest. 8. By the Quest. $\left\{ \begin{array}{l} 2B - 20 = A + 10 \\ B + 10 = 3A - 30 \end{array} \right\}$

\times the 1st equat. by 3. $\left. \begin{array}{l} 6B - 60 = 3A + 30 \\ 5B - 70 = 60 \end{array} \right\}$
 From this equat. take the 2d equat. $\left. \begin{array}{l} 6B - 60 = 3A + 30 \\ 5B - 70 = 60 \end{array} \right\}$
 Transp. -70 gs. and divide by 5. $\left. \begin{array}{l} B = 26 \\ A = 22 \end{array} \right\}$ Ans.
 Therefore, by the 1st equat.

Quest. 9. Put x for the money the tippler had at first.

	<i>s.</i>	<i>s.</i>
2 <i>s.</i> spent at the 1st house left	$x - 2$	2
Add $x - 2$, borrowed; had	$2x - 4$	
2 <i>s.</i> spent at the 2d house left	$2x - 6$	6
Add $2x - 6$, borrowed; had	$4x - 12$	
2 <i>s.</i> spent at the 3d house left	$4x - 14$	14
Add $4x - 14$, borrowed; had	$8x - 28$	
2 <i>s.</i> spent at the 4th house left	$8x - 30 = 0$	

Therefore $8x = 30$, that is, $x = 3 \frac{3}{8}$. Ans.

Quest. 10. Put x for the charge made for the man, and y that made for his wife.

By the Quest. $\left\{ \begin{array}{l} 1 = \text{the child's reckoning.} \\ 1 + \frac{x}{4} = y. \text{ the woman's reckoning.} \\ 2 + \frac{x}{4} = x. \text{ the man's reckoning.} \end{array} \right.$

\times the last equat. by 4. $\left. \begin{array}{l} 8 + x = 4x \\ \frac{8}{3} = x = 2 \dots 8 = 32d. \end{array} \right\}$
 Transp. x , and divide by 3. $\left. \begin{array}{l} 8 + x = 4x \\ \frac{8}{3} = x = 2 \dots 8 = 32d. \end{array} \right\}$ Ans.
 And, by the 2d equat. $\left. \begin{array}{l} 8 + x = 4x \\ y = 1 \dots 8 = 20d. \end{array} \right\}$

Quest. 11. Let x, y, z respectively be the quantities of brandy, wine, and cider, in gallons.

By the Quest. $\left\{ \begin{array}{l} x + y + z = 60 \\ z - 6 = x \\ z + \frac{1}{3}x = y \end{array} \right\}$ or $\left\{ \begin{array}{l} x = 60 - y - z \\ x = z - 6 \\ x = 5y - 5z \end{array} \right\}$

(Key to Vol. I. page 237.)

$$\begin{array}{l}
 \text{Putting the 1st equat. equal to} \\
 \text{the 2d.} \\
 \text{Putting the 1st equat. equal to} \\
 \text{the 3d.} \\
 \times^{\text{ing}} \text{ the 1st result by 2.} \\
 \text{To this adding the 2d result,} \\
 \text{and dividing by 8.} \\
 \text{By the 1st result.} \\
 \text{And by the 1st equation.}
 \end{array}
 \left\{ \begin{array}{l}
 y + 2z = 66 \\
 6y - 4z = 60 \\
 2y + 4z = 132 \\
 y = 24 \\
 z = 21 \\
 x = 15
 \end{array} \right.$$

Wherefore 15 galls. Brandy, 24 galls. Wine, and 21 galls. Cider. Ans.

Quest. 12. Put x for the number of troops in the whole army, z for the side of the 1st square, and $z+1$ for the side of the 2d square.

$$\text{Then, by the Question. } \left\{ \begin{array}{l}
 z^2 + 284 = x. \\
 z^2 + 2z + 1 - 25 = x.
 \end{array} \right.$$

$$\text{Therefore (Eucl. i. Ax. 1.) } z^2 + 284 = z^2 + 2z - 24.$$

$$\text{Cancel } z^2, \text{ and reduce. } \quad 154 = 2z$$

$$\text{But } 154^2 + 284 = x = 24000 \text{ men. Ans.}$$

Quest. 13. Let x represent the number sought.

$$\begin{array}{l}
 \text{Then, by the Question.} \\
 \times^{\text{ing}} \text{ extremes and means.} \\
 \text{Cancel } x^2 + 10x + 24.
 \end{array}
 \left\{ \begin{array}{l}
 x + 3 : x + 5 :: x + 5 : x + 8 \\
 x^2 + 11x + 24 = x^2 + 10x + 25 \\
 x = 1 \text{ the number sought.}
 \end{array} \right.$$

Quest. 14. For the three traders put A , B , and C .

$$\text{By the Quest. } \left\{ \begin{array}{l}
 A + B + C = \text{£}760. \\
 A + B - \text{£}240 = C. \\
 B + C - \text{£}360 = A.
 \end{array} \right\} \text{ or, } \left\{ \begin{array}{l}
 A = \text{£}760 - B - C. \\
 A = \text{£}240 - B + C. \\
 A = B + C - \text{£}360.
 \end{array} \right.$$

$$\text{Comparing the 1st and 2d equations. } \left\{ \begin{array}{l}
 \text{£}520 = 2C, \text{ that is, } C = \text{£}260.
 \end{array} \right.$$

$$\text{Comparing the 1st and 3d equations. } \left\{ \begin{array}{l}
 \text{£}1120 = 2C + 2B, \text{ that is, } B = \text{£}300.
 \end{array} \right.$$

$$\text{By the 1st equat. therefore, } \quad \text{£}200 = A.$$

Consequently, the 1st £200, the 2d £300, and the 3d £260. Ans.

(Page 238.)

Quest. 15. Let x and y represent the two numbers sought.

$$\text{By the Question } x : y :: 3 : 4, \text{ hence } 4x = 3y, \text{ or, } 12x = 9y.$$

$$\text{Also } xy = 12x + 12y.$$

(Key to Vol. I. page 238.)

$$\begin{array}{l|l} \text{For } 12x \text{ substitute } 9y, & xy = 9y + 12y = 21y. \\ \text{Divide by } y, & x = 21. \\ \text{But} & x : y :: 3 : 4 :: 21 : 28. \end{array}$$

Therefore 21 and 28 are the numbers required.

Quest 16. Put x for the reckoning, and y for the number of persons.

$$\text{It is by the Question. } \left\{ \begin{array}{l} \frac{x}{y+4} + 1 = \frac{x}{y-3} - 1 = \frac{x}{y} \end{array} \right\}$$

$$\text{Comparing the 1st expression with the 3d, } \left. \begin{array}{l} \text{and reducing.} \\ \end{array} \right\} x = \frac{1}{4}(y^2 + 4y)$$

$$\text{Comparing the 2d expression with the 3d, } \left. \begin{array}{l} \text{and reducing.} \\ \end{array} \right\} x = \frac{1}{3}(y^2 - 3y)$$

$$\begin{array}{l|l} \text{Therefore (Eucl. i. Ax. 1.)} & 3y^2 + 12y = 4y^2 - 12y. \\ \text{Reduce.} & 24y = y^2. \\ \text{Divide by } y. & 24 = y = \text{the numb. of persons.} \end{array}$$

$$\text{But } \frac{x}{28} + 1 (\text{substituting } 24 \text{ for } y) = \frac{x}{24}, \text{ that is, } 24x + 672 = 28x,$$

or $x = 168$ shillings.

$$\text{Now } 168s. = 8 \text{ guineas, and } \frac{168s.}{24} = 7 \text{ shillings for each person.}$$

Therefore

24 persons, 7s. paid by each, and 8 guineas the whole reckoning. Ans.

Quest. 17. Let x represent the 1st horse, and y the 2d.

$$\text{Then, by the Quest. } \left\{ \begin{array}{l} x + 18 = 2y + 6, \text{ that is, } x = 2y - 12. \\ 3x + 9 = y + 18. \end{array} \right.$$

$$\text{Substitute } 2y - 12 \text{ for } x. \left| \begin{array}{l} 6y - 27 = y + 18. \\ \end{array} \right.$$

$$\text{Reduce. } \left| \begin{array}{l} 5y = 45, \text{ or, } y = 9. \\ \end{array} \right.$$

$$\text{But } \left| \begin{array}{l} x = 2y - 12 = 18 - 12 = 6. \\ \end{array} \right.$$

Therefore the 1st horse was worth £6, and the 2d horse £9. Ans.

Quest. 18. Let x and y represent the two numbers ;

$$\text{By the Question. } \left\{ \begin{array}{l} x : y :: 2 : 3, \text{ that is, } 3x = 2y. \\ x + 6 : y + 6 :: 4 : 5, \text{ that is, } 5x + 6 = 4y. \end{array} \right.$$

$$\text{From double the 1st equat. take the 2d. } \left| \begin{array}{l} x - 6 = 0, \text{ or, } x = 6. \\ \end{array} \right.$$

$$\text{But, by the 1st equation } \left| \begin{array}{l} 3x = 2y = 18, \text{ hence } y = 9. \\ \end{array} \right.$$

The two numbers therefore are 6 and 9.

(Key to Vol. I. page 238.)

Quest. 19. Put x and y for the two numbers sought.

$$\text{By the Question } \left\{ \begin{array}{l} x:y::x+y:20, \\ x:y::x-y:10, \end{array} \right\} \text{ or, } \left\{ \begin{array}{l} 20x=xy+y^2 \\ 10x=xy-y^2 \end{array} \right\}$$

$$\begin{array}{l} \text{Add the 1st equat. to the 2d, and div. by } x. \\ \text{And by the 1st equat. substituting 15 for } y. \\ \text{Reduce.} \end{array} \left| \begin{array}{l} 30=2y, \text{ or, } y=15 \\ 20x=15x+225 \\ 5x=225, \text{ and } x=45. \end{array} \right.$$

Therefore 15 and 45 are the numbers required.

Quest. 20. If x be put for the greater number, and y for the less.

$$\text{By the Question. } \left\{ \begin{array}{l} x-y:x+y::2:3 \\ x+y:xy::3:5 \end{array} \right\} \text{ or, } \left\{ \begin{array}{l} 3x-3y=2x+2y \\ 5x+5y=3xy \end{array} \right\}$$

$$\begin{array}{l} \text{Reduce the 1st equation.} \\ \text{Add the 2d equat. to the 3d.} \\ \text{Divide by } x. \\ \text{But} \end{array} \left| \begin{array}{l} x=5y, \text{ that is, } x-5y=0. \\ 6x=3xy \\ 6=3y, \text{ hence } y=2. \\ x=5y=10. \end{array} \right.$$

Therefore 2 and 10 are the Answer.

Quest. 21. Let x , y , and z represent the three numbers.

$$\text{By the Question. } \left\{ \begin{array}{l} x:z::5:9 \\ x, y, z \text{ have equal diff.} \\ x+y+z=63 \end{array} \right\} \text{ or, } \left\{ \begin{array}{l} x=\frac{5}{9}z. \\ x=2y-z. \\ x=63-y-z. \end{array} \right\}$$

$$\begin{array}{l} \text{Comparing the 1st and 2d equa-} \\ \text{tions, and reducing.} \\ \text{Comparing the 1st and 3d equa-} \\ \text{tions, and reducing.} \\ \text{Therefore} \end{array} \left\{ \begin{array}{l} 14z=18y. \\ 14z=567-9y. \\ 18y=567-9y, \text{ that is, } y=21. \end{array} \right.$$

But $14z=18y$, consequently $z=27$, and $x=\frac{5}{9}z=15$.

Hence 15, 21, and 27 are the three numbers required.

Quest. 22. Put x for the greater part, and y for the less;

$$\text{By the Question. } \left\{ \begin{array}{l} x+y=24 \\ \frac{x}{y}:\frac{y}{x}::4:1 \end{array} \right\} \text{ or, } \left\{ \begin{array}{l} x=24-y. \\ \frac{x}{y}=\frac{4y}{x} \end{array} \right\}$$

(Key to Vol. I. page 238.)

$$\begin{array}{l} \text{Substituting } 24-y, \text{ for } x. \\ \text{Reduce.} \\ \text{Extract } \sqrt{\quad} \text{ on both sides.} \\ \text{But } x=24-y. \end{array} \left| \begin{array}{l} \frac{24-y}{y} = \frac{4y}{24-y} \\ 576-48y+y^2=4y^2 \\ 24-y = 2y, \text{ or } y=8 \\ \text{that is, } x=16. \end{array} \right.$$

8 and 16, therefore, are the two parts.

Quest. 23. Let x be the age of the elder, and y the age of the younger son.

$$\text{By the Question. } \left\{ \begin{array}{l} x+y+18=2x \\ x-y-6=y \end{array} \right\} \text{ or, } \left\{ \begin{array}{l} x=y+18 \\ x=2y+6. \end{array} \right.$$

Comparing the 1st and 2d equations, and reducing, $y=12$.

$$\text{But } 2y+6=x=30.$$

Hence 12 and 30 are the Answer.

(Page 239.)

Quest. 24. Assume $x, y, z,$ and $w,$ for the four numbers.

$$\text{By the Question. } \left\{ \begin{array}{l} x+y+z=13 \\ x+y+w=15 \\ x+z+w=18 \\ y+z+w=20 \end{array} \right.$$

$$\begin{array}{l} \text{Take the 1st equat. from the 2d.} \\ \text{Subst. } 2+z \text{ for } w \text{ in the 3d equat.} \\ \text{From this equat. take the 1st.} \\ \text{Subst. known values in the 4th.} \\ \text{Collect.} \\ \text{But} \\ \text{And} \\ \text{Lastly} \end{array} \left| \begin{array}{l} w=2+z \\ x+2z=16 \\ z-y=3, \text{ or } z=y+3 \\ y, +y+3, +2+y+3=20 \\ 3y+8=20, \text{ or } y=4 \\ z=y+3=4+3=7 \\ x=16-2z=16-14=2 \\ w=2+z=2+7=9. \end{array} \right.$$

Therefore the numbers sought are, 2, 4, 7, and 9.

Quest. 25. Let the four parts be severally represented by $x, y,$
 $w,$ and $z.$

$$\text{By the Question. } \left\{ \begin{array}{l} x+3=y-3 \\ x+3=3w \\ x+3=\frac{1}{3}z \\ x+y+z+w=48 \end{array} \right\} \text{ or, } \left\{ \begin{array}{l} x=y-6 \\ x=3w-3 \\ x=\frac{1}{3}(z-9) \\ \text{-----} \end{array} \right\}$$

(Key to Vol. I. page 239.)

$$\begin{array}{l} \text{Comparing the 1st and 2d values} \\ \text{of } x. \end{array} \left. \vphantom{\begin{array}{l} \text{Comparing the 1st and 2d values} \\ \text{of } x. \end{array}} \right\} \frac{y}{3} - 1 = w$$

$$\begin{array}{l} \text{Comparing the 1st and 3d values} \\ \text{of } x. \end{array} \left. \vphantom{\begin{array}{l} \text{Comparing the 1st and 3d values} \\ \text{of } x. \end{array}} \right\} 3y - 9 = z$$

$$\begin{array}{l} \text{In the 4th equat. given, express} \\ x, w, z, \text{ in terms of } y. \end{array} \left. \vphantom{\begin{array}{l} \text{In the 4th equat. given, express} \\ x, w, z, \text{ in terms of } y. \end{array}} \right\} y - 6 + y + \frac{y}{3} - 1 + 3y - 9 = 48.$$

$$\text{Collect and reduce. } y = 12.$$

$$\begin{array}{l} \text{But} \\ \text{And} \\ \text{Also} \\ \text{Therefore} \end{array} \left| \begin{array}{l} y - 6 = x = 12 - 6 = 6. \\ \frac{1}{3}y - 1 = w = 4 - 1 = 3. \\ 3y - 9 = z = 36 - 9 = 27. \\ 6, 12, 3, \text{ and } 27 \text{ are the parts sought.} \end{array} \right.$$

QUADRATIC EQUATIONS.

(Page 243.)

$$\begin{array}{l} \text{Ex. 1.} \\ \text{Complete the square.} \\ \text{Extract the square root.} \\ \text{Transpose.} \end{array} \left| \begin{array}{l} \text{Given. } x^2 - 6x = 40 \\ x^2 - 6x + 9 = 49 \\ x - 3 = \pm 7 \\ x = 3 \pm 7 = 10, \text{ or, } -4. \text{ Ans.} \end{array} \right.$$

$$\begin{array}{l} \text{Ex. 2.} \\ \text{Complete the square.} \\ \text{Extract the square root.} \\ \text{Transpose.} \end{array} \left| \begin{array}{l} \text{Given. } x^2 - 5x = 24 \\ x^2 - 5x + 6.25 = 30.25 \\ x - 2.5 = \pm 5.5 \\ x = 2.5 \pm 5.5 = 8, \text{ or, } -3. \text{ Ans.} \end{array} \right.$$

$$\begin{array}{l} \text{Ex. 3.} \\ \text{Complete the square.} \\ \text{Extract the square root.} \\ \text{Transpose.} \end{array} \left| \begin{array}{l} \text{Given (when reduced). } x^2 + \frac{4}{5}x = \frac{204}{5} \\ x^2 + \frac{4x}{5} + \frac{4}{25} = \frac{1024}{25} \\ x + \frac{2}{5} = \pm \frac{32}{5} \\ x = \pm \frac{32-2}{5} = 6, \text{ or } -6\frac{4}{5}. \text{ Ans.*} \end{array} \right.$$

* Hutton's Answers in general to the Quadratic Equations are imperfect, in as much as he gives only ONE of the Roots of each Equation.

$$\text{Ex. 4.} \quad \text{Given. } \left| x^2 - \frac{1}{2}x = 14. \text{ (By doubling and transposition.)} \right.$$

(Key to Vol. I. page 243.)

$$\begin{array}{l|l} \text{Complete the square.} & x^2 - \frac{1}{2}x + \frac{1}{16} = 14\frac{1}{16} = 14.0625 \\ \text{Extract the square root.} & x - \frac{1}{4} = \pm 3\frac{3}{4} \\ \text{Transpose.} & x = \frac{1}{4} \pm 3\frac{3}{4} = 4, \text{ or } -3\frac{1}{2}. \text{ Ans.} \end{array}$$

$$\begin{array}{l|l} \text{Ex. 5.} & \text{Given. } x^4 - \frac{2}{3}x^2 = \frac{40}{3} \\ \text{Complete the square.} & x^4 - \frac{2}{3}x^2 + \frac{1}{9} = \frac{121}{9} \\ \text{Extr. the square root.} & x^2 - \frac{1}{3} = \pm 3\frac{2}{3} \\ \text{Transpose.} & x^2 = \frac{1}{3} \pm 3\frac{2}{3} = 4, \text{ or } -3\frac{1}{3} \text{ an irrational Surd.} \\ \text{Extr. the square root.} & x = 2, \text{ or } -2. \text{ Ans.} \end{array}$$

$$\begin{array}{l|l} \text{Ex. 6.} & \text{Given } (\times \text{ing by } 3). \quad x - \frac{3}{2}\sqrt{x} = 4\frac{1}{2} = \frac{9}{2} \\ \text{Complete the square.} & x - \frac{3}{2}\sqrt{x} + \frac{9}{16} = 5\frac{1}{16} \\ \text{Extract the square root.} & \sqrt{x} - \frac{3}{4} = \pm 2\frac{1}{4} \\ \text{Transpose.} & \sqrt{x} = \frac{3}{4} \pm 2\frac{1}{4} = 3, \text{ or } -1\frac{1}{2} \\ \text{Square both sides.} & x = 9, \text{ or } 2\frac{1}{4}. * \text{ Ans.} \end{array}$$

* When $x=2\frac{1}{4}$ its negative root must be taken to answer the conditions.

$$\begin{array}{l|l} \text{Ex. 7.} & \text{Given. } x^2 + \frac{4}{3}x = 1\frac{1}{2} \\ \text{Complete the square.} & x^2 + \frac{4}{3}x + \frac{4}{9} = \frac{70}{36} = 1.944444 \\ \text{Extract the square root.} & x + \frac{2}{3} = \pm 1.39444. \\ \text{Transpose.} & x = .727766, \text{ or } -2.06111. \text{ Ans.} \end{array}$$

$$\begin{array}{l|l} \text{Ex. 8.} & \text{Given. } x^6 + 4x^3 = 12 \\ \text{Complete the square.} & x^6 + 4x^3 + 4 = 16 \\ \text{Extract the square root.} & x^3 + 2 = \pm 4 \\ \text{Transpose.} & x^3 = \pm 4 - 2 = 2, \text{ or } -6. \end{array}$$

(Key to Vol. I. page 243.)

Extr. the Cube root. | $x = \sqrt[3]{2} = 1.259921$, or $\sqrt[3]{-6} = -1.81712$.

Ex. 9.	Given.	$x^2 + 4x = a^2 + 2$
	Complete the square.	$x^2 + 4x + 4 = a^2 + 6$
	Extract the square root.	$x + 2 = \pm \sqrt{a^2 + 6}$
	Transpose.	$x = \pm \sqrt{a^2 + 6} - 2$.

QUESTIONS FOR PRACTICE.

(Page 245.)

Quest. 1.	Put	x for the number sought.
	By the Question.	$x^2 + x = 42$
	Complete the square.	$x^2 + x + \frac{1}{4} = 42\frac{1}{4}$
	Extract the square root.	$x + \frac{1}{2} = \pm 6\frac{1}{2}$
	Transpose.	$x = 6$, or -7 .

Quest. 2. Let x represent the greater number, and y the less.

By the Question $\left\{ \begin{array}{l} y : x :: x : 12 \\ x^2 + y^2 = 45 \end{array} \right\}$ or, $\left\{ \begin{array}{l} x^2 = 12y \\ x^2 = 45 - y^2 \end{array} \right\}$

Therefore	$y^2 + 12y = 45$
Complete the sq ^r .	$y^2 + 12y + 36 = 81$
Extract the sq ^r . root.	$y + 6 = \pm 9$
Transpose.	$y = 3$, or, -15
But	$12y = x^2 = 36$, or, -180
Therefore	$x = 6$, or $\sqrt{-180}$, an irrational Surd.

(Page 246.)

Quest. 3. Assume x as the greater, and y as the less number.

By the Question $\left\{ \begin{array}{l} x = y + 2 \\ x^3 - y^3 = 98 \end{array} \right\}$ or, $\left\{ \begin{array}{l} x^3 = y^3 + 6y^2 + 12y + 8 \\ x^3 = y^3 + 98 \end{array} \right\}$

Therefore	$6y^2 + 12y = 90$
Divide by 6.	$y^2 + 2y = 15$
Complete the square.	$y^2 + 2y + 1 = 16$
Extract the square root.	$y + 1 = \pm 4$
Transpose.	$y = 3$, or, -5
But	$x = y + 2 = 5$, or -3 . } Ans.

(Key to Vol. I. page 246.)

Quest. 4. Put x and y for the two numbers sought.

$$\text{By the Quest. } \left\{ \begin{array}{l} x + y = 6 \\ x^3 + y^3 = 72 \end{array} \right\} \text{ or, } \left\{ \begin{array}{l} x^3 = 216 - 108y + 18y^2 - y^3 \\ x^3 = 72 \end{array} \right\}$$

$$\begin{array}{l|l} \text{Therefore} & 18y^2 - 108y = -144 \\ \text{Divide by 18.} & y^2 - 6y = -8 \\ \text{Complete the square.} & y^2 - 6y + 9 = 1 \\ \text{Extract the square root.} & y - 3 = \pm 1 \\ \text{Transpose.} & y = 4, \text{ or } 2. \\ \text{But} & x = 6 - y = 2, \text{ or } 4. \end{array}$$

That is, no numbers except 4 and 2 will answer the conditions of the Question.

Quest. 5. Let x and y represent the two numbers proposed.

$$\text{By the Question } \left\{ \begin{array}{l} xy = 20 \\ x^3 - y^3 = 61 \end{array} \right\} \text{ or, } \left\{ \begin{array}{l} x = \frac{20}{y}, \text{ and } x^3 = \frac{8000}{y^3} \\ \dots\dots\dots x^3 = y^3 + 61. \end{array} \right.$$

$$\begin{array}{l|l} \text{Therefore} & y^3 + 61 = \frac{8000}{y^3} \\ \text{Reduce.} & y^6 + 61y^3 = 8000 \end{array}$$

$$\begin{array}{l|l} \text{Complete the square.} & y^6 + 61y^3 + 930\frac{1}{4} = 8930\cdot 25 \\ \text{Extract the square root.} & y^3 + 30\frac{1}{2} = \pm 94\cdot 5 \\ \text{Transpose.} & y^3 = 64. \text{ or } -125 \\ \text{Extract the cube root.} & y = 4. \text{ or } -5 \\ \text{But} & xy = 20 = 4x, \text{ or, } -5x \\ \text{Consequently} & x = 5. \text{ or } -4. \end{array}$$

Quest. 6.

$$\text{Given } \left\{ \begin{array}{l} x + y = 11 \\ x^2 y^2 = 784 \end{array} \right\} \text{ or, } \left\{ \begin{array}{l} x = 11 - y \\ x = \frac{28}{y} \end{array} \right\}$$

$$\begin{array}{l|l} \text{Therefore} & 11 - y = \frac{28}{y} \\ \text{Reduce.} & y^2 - 11y = -28 \\ \text{Complete the square.} & y^2 - 11y + 30\cdot 25 = 2\cdot 25 \\ \text{Extract the square root.} & y - 5\cdot 5 = \pm 1\cdot 5 \\ \text{Transpose.} & y = 7, \text{ or } 4. \\ \text{But} & x = \frac{28}{y} = 4. \text{ or } 7. \end{array}$$

(Key to Vol. I. page 246.)

Quest. 7.

$$\text{Given } \left\{ \begin{array}{l} x+y=5 \\ \frac{x}{y}+\frac{y}{x}=4\frac{1}{4} \end{array} \right\} \text{ or, } \left\{ \begin{array}{l} x=5-y \\ \frac{x^2+y^2}{xy}=4\frac{1}{4} \end{array} \right\}$$

Substitute $5-y$ for x .	$\frac{25-10y+2y^2}{5y-y^2}=4\frac{1}{4}$
Multiply by $5y-y^2$.	$2y^2-10y+25=21\frac{1}{4}y-4\frac{1}{4}y^2$
Transpose and collect.	$6\frac{1}{4}y^2-31\frac{1}{4}y=-25$
Divide by $6\frac{1}{4}$.	$y^2-5y=-4$
Complete the sq ^r and extract.	$y-2\frac{1}{2}=\pm 1\frac{1}{2}$
Transpose.	$y=4, \quad \text{or } 1.$
Consequently	$x=1, \quad \text{or } 4. \quad \left. \vphantom{\frac{25-10y+2y^2}{5y-y^2}} \right\} \text{Ans.}$

Quest. 8.

$$\text{Given } \left\{ \begin{array}{l} x+y=12 \\ xy=8x-8y \end{array} \right\} \text{ or, } \left\{ \begin{array}{l} x=12-y \\ \text{-----} \end{array} \right\}$$

Substitute $12-y$ for x in the } 2d equat. }	$12y - y^2 = 96 - 16y$
Change the signs, and reduce.	$y^2 - 28y = -96$
Complete the square.	$y^2 - 28y + 196 = 100$
Extract the square root.	$y - 14 = \pm 10$
Transpose.	$y = 24. \text{ (not applicable), or } 4.$
Now	$x = 12 - y = 12 - 4 = 8.$

Quest. 9. Let x represent the greater part, y the less.

$$\text{By the } \left\{ \begin{array}{l} x+y=10 \\ 16y^2-112=4x^2 \end{array} \right\} \text{ hence } \left\{ \begin{array}{l} 4x^2=400-80y+4y^2 \\ 4x^2=16y^2-112 \end{array} \right\}$$

Therefore, and by reduction,	$y^2 + \frac{20}{3}y = 42\frac{2}{3}$
Complete the square, extract, } and transpose. }	$y=4. \text{ or } -10\frac{2}{3} \text{ inapplicable.}$
Consequently	$x=6.$

Quest. 10. Put x and y for the numbers required, x being the greater.

$$\text{By the } \left\{ \begin{array}{l} x^2+xy=104 \\ x^2+y^2=89 \end{array} \right\} \text{ or, } \left\{ \begin{array}{l} x^2=104-xy \\ x^2=89-y^2, \text{ and } x=\sqrt{89-y^2} \end{array} \right\}$$

(Key to Vol. I. page 246.)

$$\begin{array}{l|l}
 \text{Therefore} & y^2 + 15 = xy = y\sqrt{89 - y^2} \\
 \text{Square both sides.} & y^4 + 30y^2 + 225 = 89y^2 - y^4 \\
 \text{Transpose.} & 2y^4 - 59y^2 = -225 \\
 \text{Divide by 2, and complete.} & y^4 - \frac{59}{2}y^2 + \frac{3481}{16} = \frac{1681}{16} \\
 \text{Extract the square root.} & y^2 - \frac{59}{4} = \pm \frac{41}{4} \\
 \text{Transpose.} & y^2 = \frac{59}{4} \pm \frac{41}{4} = 25, \text{ or } 4\frac{1}{2}.
 \end{array}$$

$$\begin{array}{l|l}
 \text{Extract the square root.} & y = 5. \text{ or } 2.12131 \text{ \&c.} \\
 \text{But} & x = \sqrt{89 - y^2} = 8. \text{ or } 9.19238 \text{ \&c.}
 \end{array}$$

Quest. 11. Let x be the 1st digit, y the 2d, and z the number sought.

$$\text{By the Question} \left\{ \begin{array}{l} 10x + y = z \\ \frac{10x + y}{xy} = 5\frac{1}{3} \end{array} \right\} \text{ also } \left\{ \begin{array}{l} 10x + y - 9 = 10y + x \end{array} \right.$$

$$\begin{array}{l|l}
 \text{Collect the last equat. and divide by 9.} & x = y + 1 \\
 \text{Substitute } y + 1 \text{ for } x \text{ in the 2d equat.} & \frac{11y + 10}{y^2 + y} = 5\frac{1}{3} \\
 \text{Multiply by } y^2 + y. & 11y + 10 = 5\frac{1}{3}y^2 + 5\frac{1}{3}y \\
 \text{Reduce, and divide by } 5\frac{1}{3}. & y^2 - \frac{17}{16}y = \frac{30}{16} \\
 \text{Complete the square, extract the sq. } \left. \begin{array}{l} \text{root, and transpose.} \\ \text{But} \\ \text{And} \end{array} \right\} & \left. \begin{array}{l} y = 2 \\ x = y + 1 = 3 \\ z = 10x + y = 30 + 2 = 32. \end{array} \right.
 \end{array}$$

Quest. 12. Let the three parts be severally represented by x , y , and z .

$$\text{By the Question} \left\{ \begin{array}{l} x + y + z = 20 \\ xyz = 270 \\ x - (y + 2) = y - z \end{array} \right\} \text{ or, } \left\{ \begin{array}{l} x = 20 - y - z \\ x = \frac{270}{yz} \\ x = 2y - z + 2. \end{array} \right.$$

(Key to Vol. I. page 246.)

Compare the 2d and 3d equat ^s .	$3y^2 = 75$, or $y = 5$
Compare the 1st and 2d } equat ^s . substituting 5 for y .	$2z^2 - 16z = -14$.
Divide by 2, and complete } the square.	$z^2 - 8z + 16 = 9$
Extract the sq ^r root, and transp.	$z = 7$ or 1
But	$x = 13 - y - z = 8 - z = 1$ or 7 .
Therefore	$1, 5$, and 7 are the parts sought.

Quest. 15. Put x, y , and z for the three numbers proposed.

By the Question $\left\{ \begin{array}{l} x + y + z = 12 \\ x - y = y - z \\ x^4 + y^4 + z^4 = 962 \end{array} \right\}$ or, $\left\{ \begin{array}{l} x = 12 - y - z \\ x = 2y - z \\ \dots \dots \dots \end{array} \right\}$

Compare these two va- } lues of x .	$12 = 3y$, or $y = 4$
Substitute 4 for y in the } 2d and 3d equations	$x + z = 8$, and $x^4 + z^4 = 706$
Put $a = \frac{x+z}{2}$, and $n = \frac{x-z}{2}$	$x = a + n$, and $z = a - n$.
Therefore	$\left\{ \begin{array}{l} x^4 = \overline{a+n}^4 = a^4 + 4a^3n + 6a^2n^2 + 4an^3 + n^4 \\ z^4 = \overline{a-n}^4 = a^4 - 4a^3n + 6a^2n^2 - 4an^3 + n^4 \end{array} \right.$
Add, and divide by 2.	$353 = a^4 + 6a^2n^2 + n^4$
Substitute 4 (its known } value) for a .	$353 = 256 + 96n^2 + n^4$
Transpose, and com- } plete the square.	$n^4 + 96n^2 + 2304 = 2401$
Extract the square root, } and transpose.	$n^2 = \pm 49 - 48 = 1$, or -97 an irrat. Surd.
Extract the square root.	$n = \pm 1$.

But $x = a + n = 4 \pm 1 = 5$ or 3 , and $z = a - n = 4 \mp 1 = 3$, or 5 .
Consequently 3, 4, and 5, are the numbers sought.

(Page 247.)

Quest. 16. For the proposed three numbers, substitute x, y , and z .

By the Question $\left\{ \begin{array}{l} 2y = x + z \\ z^2 + xy = 28 \\ x^2 + yz = 44 \end{array} \right\}$ or, $\left\{ \begin{array}{l} y = \frac{1}{2}(x + z) \\ \dots \dots \dots \\ \dots \dots \dots \end{array} \right\}$

(Key to Vol. I. page 247.)

$$\text{Sub. } \frac{x+z}{2} \text{ for } y \text{ in the 2d \& 3d equat}^{\text{s}} \left\{ \begin{array}{l} z^2 + \frac{x^2+xz}{2} = 28, \text{ or, } 2z^2 + x^2 + xz = 56 \\ x^2 + \frac{z^2+xz}{2} = 44, \text{ or, } 2x^2 + z^2 + xz = 88 \end{array} \right.$$

By subtraction. $x^2 = z^2 + 32$
 Put $r+s=x$. $x^2 = r^2 + 2rs + s^2$
 and $r-s=z$. $z^2 + 32 = r^2 - 2rs + s^2 + 32$

Therefore (Eucl. i. Ax. 1.) $4rs = 32$, and $r = \frac{8}{s}$.

But

$$x = r + s = \frac{8}{s} + s = \frac{8 + s^2}{s}$$

And

$$z = r - s = \frac{8}{s} - s = \frac{8 - s^2}{s}$$

Also

$$x^2 + \frac{xz + z^2}{2} = 44 = \frac{64 + 16s^2 + s^4}{s^2} + \frac{64 - s^4 + 16s^2 + 64 + s^4}{2s^2}$$

Reduce.

$$44 = \frac{s^4 + 8s^2 + 128}{s^2}, \text{ or } 44s^2 = s^4 + 8s^2 + 128.$$

By further reduction.

$$s^4 - 36s^2 = -128.$$

Complete the square.

$$s^4 - 36s^2 + 324 = 196.$$

Extract the square root and transpose.

$$s^2 = 32, \text{ or } 4.$$

Extract the square root.

$$s = 5.6568542 \text{ \&c. or } 2.$$

Wherefore,

$$r = \frac{8}{s} = \frac{8}{\sqrt{32}}, \text{ or } \frac{8}{2} = 4.$$

And

$$x = r + s = \frac{8}{\sqrt{32}} + \sqrt{32}, \text{ or } 4 + 2 = 6. \text{ Likewise}$$

$$z = r - s = \frac{8}{\sqrt{32}} - \sqrt{32}, \text{ or } 4 - 2 = 2.$$

(Key to Vol. I. page 247.)

Rejecting the Surd values of x and z as irrelevant to the question,

$$y = \frac{x+z}{2} = \frac{6+2}{2} = 4. \text{ Consequently } 2, 4, 6 \text{ are the numbers required.}$$

Quest. 17.

$$\text{Given } \begin{cases} A + B + C = \mathcal{L}1444 \\ B + \sqrt{A} = 920 \\ B + \sqrt{C} = 912 \end{cases}$$

$$\text{Put } \begin{cases} x^2 + 2xy + y^2 = A \\ x^2 - 2xy + y^2 = C \end{cases}$$

Then by comparison and substitution,

$$\mathcal{L}1444 - x^2 - 2xy - y^2 - x^2 + 2xy - y^2 + x + y = \mathcal{L}920.$$

$$\mathcal{L}1444 - x^2 - 2xy - y^2 - x^2 + 2xy - y^2 + x - y = 912.$$

$$\begin{array}{r} \text{By subtraction} \quad \text{-----} \quad 2y = \quad 8. \text{ And} \\ \mathcal{L}2888 - 4x^2 \qquad \qquad -4y^2 \quad +2x = 1832. \text{ by} \\ \text{addition.} \end{array}$$

Transpose, and substitute $\mathcal{L}8$ for $2y$.

$$\mathcal{L}992 = 4x^2 - 2x, \text{ hence } \mathcal{L}248 = x^2 - \frac{1}{2}x.$$

Complete the sq^r, extract the square root, and transpose.

$$x = \mathcal{L}16.$$

But $y = \mathcal{L}4$. Therefore

$$x + y = \mathcal{L}20 = \sqrt{A}.$$

And

$$x - y = \mathcal{L}12 = \sqrt{C}.$$

Consequently $A = \mathcal{L}400$, $B = \mathcal{L}900$, and $C = \mathcal{L}144$.Quest. 18. Let x , y , and z , represent the three numbers sought.

$$\text{By the Question. } \begin{cases} x + z = 2y \\ x^2 + y^2 + z^2 = 93 \\ 3x + 4y + 5z = 66 \end{cases} \text{ or } \begin{cases} x = 2y - z \\ \text{-----} \\ x = \frac{66 - 4y - 5z}{3} \end{cases}$$

Compare these two values of x .

$$10y + 2z = 66, \text{ or } y = \frac{66 - 2z}{10}.$$

Substitute in the 1st equation $\frac{66 - 2z}{10}$ for y .

$$x = \frac{66 - 2z}{5} - z = \frac{66 - 7z}{5}.$$

(Key to Vol. I. page 247.)

x and y being now known in terms of z , it is

$$x^2 + y^2 + z^2 = \frac{4356 - 924z + 49z^2}{25} + \frac{4356 - 264z + 4z^2}{100} + z^2 = 93.$$

Reduce, complete the square, extract the square root, and transpose, $z=8$.

But

$$y = \frac{66 - 2z}{10} = \frac{66 - 16}{10} = \frac{50}{10} = 5.$$

And

$$x = 2y - z = 10 - 8 = 2.$$

Therefore
2, 5, and 8, are the numbers required.

Quest. 19. For the two numbers proposed, put x and y .

By the Question. $\left\{ \begin{array}{l} xy + x + y = 47 \\ x^2 + y^2 - x - y = 62 \end{array} \right\}$

Assume $m = \frac{1}{2}(x+y)$ and $n = \frac{1}{2}(x-y)$.

By substitution.	$xy + x + y = m^2 - n^2 + 2m = 47$
And	$x^2 + y^2 - (x+y) = 2m^2 + 2n^2 - 2m = 62$
Subtr. half the last equat. } from the equat. above it. }	$3m - 2n^2 = 16$
Add the last subtrahend to } its minuend. }	$2m^2 + m = 78$
Divide by 2, and complete } the square. }	$m^2 + \frac{1}{2}m + \frac{1}{16} = 39.0625$
Extract the square root and } transpose. }	$m = 6$
But	$3m - 2n^2 (= 18 - 2n^2) = 16$
Therefore	$2n^2 = 2, \text{ or } n = 1.$

Now $m+n=x=7$, and $m-n=y=5$. Which were to be determined.

(Key to Vol. I. page 253.)

CUBIC AND HIGHER EQUATIONS.

Ex. 5. Assume 4 and 5 as values of x .

$$x \text{ being } 4 \left\{ \begin{array}{rcl} 20 & = & 5x = 25 \\ 160 & = & 10x^2 = 250 \\ 64 & = & x^3 = 125 \\ \hline 244 & \text{Sum} & 400 \\ 260 & \text{should be} & 260 \\ \hline \end{array} \right\} x \text{ being } 5.$$

Too little 16 Errors 140 *Too much.*

The sum of the errors is 156. And

As $156 : 1. :: 16 : \cdot 1$. Hence $x=4\cdot 1$ nearly.Proceeding now with $4\cdot 1$ and $4\cdot 2$, x is found $4\cdot 1179$ &c. Ans.Ex. 6. Assume $3\cdot 5$ and $3\cdot 9$ as values of x .

$$x \text{ being } 3\cdot 5 \left\{ \begin{array}{rcl} 42\cdot 875 & = & x^3 = 59\cdot 319 \\ -7 & = & -2x = -7\cdot 8 \\ \hline 35\cdot 875 & \text{Sum} & 51\cdot 519 \\ 50 & \text{should be} & 50\cdot \\ \hline \end{array} \right\} x \text{ being } 3\cdot 9$$

Too little $14\cdot 125$ Errors $1\cdot 519$ *Too much.*

The sum of the errors is $15\cdot 644$. AndAs $15\cdot 644 : \cdot 4 :: 1\cdot 519 : \cdot 038839$ the excess of $3\cdot 9$.But $3\cdot 9 - \cdot 038839 = 3\cdot 8611 = x$ nearly.Proceeding, therefore, with $3\cdot 861$ and $3\cdot 862$, x is found *more nearly* $3\cdot 8648$. Ans.

(Page 254.)

Ex. 7. By a few trials x is discovered to be between $5\cdot 1$ and $5\cdot 2$. Wherefore assuming these,

$$x \text{ being } 5\cdot 1 \left\{ \begin{array}{rcl} -117\cdot 3 & = & -23x = -119\cdot 6 \\ 52\cdot 02 & = & 2x^2 = 54\cdot 08 \\ 132\cdot 625 & = & x^3 = 140\cdot 608 \\ \hline 67\cdot 345 & \text{Sum} & 75\cdot 088 \\ 70\cdot & \text{should be} & 70\cdot \\ \hline \end{array} \right\} x \text{ being } 5\cdot 2.$$

Too little $2\cdot 655$ Errors $5\cdot 088$ *Too much.*

(Key to Vol. I. page 254.)

The sum of the errors is 7.743. And

As $7.743 : .1 :: 2.655 : .03428$ the deficiency of 5.1.But $5.1 + .03428 = 5.13428 = x$ nearly.By another approximation x is found (more nearly) 5.13457.Ex. 8. Assuming 14.9 and 15 as values of x .

$$x \text{ being } 14.9 \left\{ \begin{array}{l} 3307.949 = x^3 = 3375 \\ -3774.17 = -17x^2 = -3825 \\ 804.6 = 54x = 810 \\ \hline 338.379 \quad \text{Sum} \quad 360 \\ 350. \quad \text{should be} \quad 350 \end{array} \right\} x \text{ being } 15.$$

Too little 11.621 Errors 10 Too much.

The sum of the errors is 21.621. And

As $21.621 : .1 :: 10 : .04625$ the excess of 15.But $15 - .04625 = 14.9537 = x$ nearly.By another approximation x is found (more nearly) 14.95407.Ex. 9. A few trials determine x between 10.2 and 10.3,Let these, therefore, be assumed as x . It is

$$x \text{ being } 10.2 \left\{ \begin{array}{l} 10824.3226 = x^4 = 11255.0881 \\ -312.12 = -3x^2 = -318.27 \\ -765 = -75x = -772.5 \\ \hline 9747.2026 \quad \text{Sum} \quad 10164.3181 \\ 10000. \quad \text{should be} \quad 10000. \end{array} \right\} x \text{ being } 10.3.$$

Too little 252.7974 Errors 164.3181 Too much.

The sum of the errors is 417.1155. And

As $417.1155 : .1 :: 164.3181 : .03939$ the excess of 10.3.But $10.3 - .03939 = 10.2606 = x$ nearly.Yet another approximation determines x (more nearly) 10.2609.Ex. 10. After a few trials assuming 1.284 and 1.285 as values of x , it is

$$x \text{ being } \left. \begin{array}{l} 1.284 \\ 1.284 \end{array} \right\} \left\{ \begin{array}{l} 5.436153 = 2x^4 = 5.453085 \\ -33.869988 = -16x^3 = -33.949186 \\ 65.94624 = 40x^2 = 66.049 \\ -38.52 = -30x = -38.55 \\ \hline -1.007595 \quad \text{Sum} \quad -0.997101 \\ -1. \quad \text{should be} \quad -1. \end{array} \right\} x \text{ being } 1.285.$$

Too much .007595 Errors .002899 Too little.

(Key to Vol. I. page 254.)

The sum of the errors is $\cdot 001371$. AlsoAs $\cdot 001371 : \cdot 001 :: \cdot 0000367 : \cdot 0000268$ the deficiency $8\cdot 64$.But $8\cdot 64 + \cdot 0000268 = 8\cdot 6400268 = x$. Ans.Ex. 13. After a few trials assuming $1\cdot 84$ and $1\cdot 85$, as values of x ,

$$x \text{ being } \left. \begin{array}{l} 1\cdot 84 \\ 1\cdot 84 \end{array} \right\} \begin{array}{l} 22\cdot 92457472 = 2x^4 = 23\cdot 4270125 \\ -43\cdot 606528 = -7x^3 = -44\cdot 321375 \\ 37\cdot 2416 = 11x^2 = 37\cdot 6475 \\ -5\cdot 52 = -3x = -5\cdot 55 \end{array} \left. \begin{array}{l} \\ \\ \\ \end{array} \right\} x \text{ being } 1\cdot 85$$

$$\begin{array}{r} 11\cdot 03964672 \quad \text{Sum} \quad 11\cdot 2031375 \\ 11\cdot \quad \quad \quad \text{should be} \quad 11\cdot \end{array}$$

Too much $\cdot 03964672$ Errors $\cdot 2031375$ Too much.The difference of the errors is $\cdot 1634907$. NowAs $\cdot 1634907 : \cdot 01 :: \cdot 03964672 : \cdot 00242$ the excess of $1\cdot 84$.But $1\cdot 84 - \cdot 00242 = 1\cdot 83758 = x$ nearly.Ex. 14. Let 18 and 19 be severally assumed and substituted for x . x being 18. x being 19.

$$\left\{ \begin{array}{l} \sqrt[5]{964\cdot 5147186}^3 = (3x^2 - 2\sqrt{x} + 1)^{\frac{3}{5}} = \sqrt[5]{1075\cdot 2822022}^3 \\ -\sqrt[5]{31\cdot 2577917}^5 = -(x^2 - 4x\sqrt{x} + 3\sqrt{x})^{\frac{5}{9}} = -\sqrt[5]{42\cdot 8003803}^5 \end{array} \right\}$$

But,

Log. of $964\cdot 5147186$ is $2\cdot 9843088$. which multiplied by 3, divided by 5, becomes(Log. of $61\cdot 7426571$. or) $1\cdot 7905853$ the log. of the 5th root of the cube of $964\cdot 5147186$. AndLog. of $31\cdot 2577917$ is $1\cdot 4949583$. which multiplied by 5, divided by 9, becomes(Log. of $6\cdot 7691234$. or) $0\cdot 8305324$ the log. of the 9th root of the 5th power of $31\cdot 2577917$. AlsoLog. of $1075\cdot 2822022$ is $3\cdot 0315234$ which multiplied by 3, divided by 5, becomes(Log. of $65\cdot 904338$. or) $1\cdot 8189140$ the log. of the 5th root of the cube $1075\cdot 2822022$. Again

(Key to Vol. I. page 254.)

Log. of 42·8003803 is 1·6314476 which multiplied by 5,
divided by 9, becomes
(Log. of 8·060459. or) 0·9063598 the log. of the 9th root of
the 5th power of 42·8003803.

Substituting these four natural numbers found, it is,

$$\begin{array}{r}
 61\cdot7426571 \\
 - 6\cdot7691234 \\
 \hline
 54\cdot9735337 \\
 56\cdot
 \end{array}
 \quad
 \begin{array}{r}
 65\cdot904338 \\
 - 8\cdot060459 \\
 \hline
 57\cdot843879 \\
 \text{Sum} \\
 \text{should be } 56\cdot
 \end{array}$$

Too little 1·0264663 *Errors* 1·843879 *Too much.*

The sum of the errors is 2·8703453. And

As 2·8703453 : 1 : : 1·0264663 : ·357 *the deficiency of 18.*

But $18 + \cdot357 = 18\cdot357 = x$ *nearly.*

Proceeding with 18·357 and 18·358, x is found (*more nearly*)
18·36087 &c. Ans.

CARDAN'S RULE FOR CUBIC EQUATIONS.

(Page 256.)

Ex. 2.

Substituting $z+3$ for x

$$\begin{array}{r}
 x^3 = z^3 + 9z^2 + 27z + 27 \\
 - 9x^2 = -9z^2 - 54z - 81 \\
 + 28x = +28z + 84 \\
 \hline
 \end{array}$$

Therefore $z^3 + z + 30 = 30$

That is, $z^3 + z = 0$, or $z = 0$, consequently $x = 3$ in one of
the roots of the given equation.

Then,

$x - 3) x^3 - 9x^2 + 28x - 30 (= x^2 - 6x + 10, \text{ the equation de-}$
pressed one degree.

$$\begin{array}{l}
 \text{By transposition.} \\
 \text{Complete the square.} \\
 \text{Extract the square root.} \\
 \text{Transpose.}
 \end{array}
 \left|
 \begin{array}{r}
 x^2 - 6x = -10 \\
 x^2 - 6x + 9 = -1 \\
 x - 3 = \pm\sqrt{-1} \\
 x = 3 \pm\sqrt{-1}
 \end{array}
 \right.$$

Therefore 3, $3 + \sqrt{-1}$, and $3 - \sqrt{-1}$ are the three values of x .

(Key to Vol. I. page 256.)

Ex 3. Substitute $z + \frac{7}{3}$ for x ;

$$\begin{aligned} x^3 &= z^3 + 7z^2 + \frac{147}{9}z + \frac{343}{27} \\ - 7x^2 &= -7z^2 - \frac{294}{9}z - \frac{1029}{27} \\ + 14x &= + \frac{126}{9}z + \frac{882}{27} \end{aligned}$$

Therefore $z^3 - \frac{7}{3}z + \frac{196}{27} = 20$, or, $z^3 - \frac{7}{3}z = 12\frac{29}{27}$.Here $a = -\frac{7}{3}$, $b = 12\frac{29}{27}$, $c = -\frac{7}{9}$, and $d = 6\frac{19}{27}$. Whence

$$\begin{aligned} \sqrt[3]{d + \sqrt{d^2 + c^3}} + \sqrt[3]{d - \sqrt{d^2 + c^3}} &= \sqrt[3]{\frac{172}{27} + \sqrt{\frac{29584 - 343}{729}}} + \\ \sqrt[3]{\frac{172}{27} - \sqrt{\frac{29584 - 343}{729}}} &= \sqrt[3]{\frac{172 + 171}{27}} + \sqrt[3]{\frac{172 - 171}{27}} = \sqrt[3]{\frac{343}{27}} + \\ \sqrt[3]{\frac{1}{27}} &= \frac{7}{3} + \frac{1}{3} = \frac{8}{3} = 2\frac{2}{3} = z. \text{ But } z + \frac{7}{3} = x = 2\frac{2}{3} + 2\frac{1}{3} = 5 \text{ the value} \\ &\text{of the unknown quantity in one of the roots of the given equation.} \end{aligned}$$

Then,

$(x - 5)x^3 - 7x^2 + 14x - 20$ ($= x^2 - 2x + 4$ the equation depressed one degree.

By transposition.	$x^2 - 2x = -4$
Complete the square.	$x^2 - 2x + 1 = -3$
Extract the square root.	$x - 1 = \pm \sqrt{-3}$
Transpose.	$x = 1 \pm \sqrt{-3}$.

Therefore 5, $1 + \sqrt{-3}$, and $1 - \sqrt{-3}$ are the three values of x .

APPLICATION OF ALGEBRA TO GEOMETRY.

(Page 363.)

PROBLEM VI.

Put x for the hypotenuse,
 b for the base (3) of the triangle; and
 d for the difference (1) of the hypotenuse and
perpendicular.

(Key to Vol. I. page 363.)

Then (because Eucl. i. 18. the hypotenuse is greater than the perpendicular) $x - d =$ the perpendicular.

But (Eucl. i. 47.) $x^2 = (x - d)^2 + b^2 = x^2 - 2dx + d^2 + b^2$, that is, $2dx = d^2 + b^2$ and $x = \frac{d^2 + b^2}{2d} =$ (substituting known values) $\frac{1^2 + 3^2}{2 \times 1} = \frac{10}{2} = 5$.

Wherefore the triangle is determined.

PROBLEM VII.

Put x for the less side about the right angle,
 d for the difference (1) of the base and perpendicular,
 also a for the hypotenuse (5).

Then (Eucl. i. 47.) $a^2 = x^2 + (x + d)^2 = x^2 + x^2 + 2dx + d^2 = 2x^2 + 2dx + d^2$, that is, $x^2 + dx = \frac{a^2 - d^2}{2}$. Completing the square, extract-

ing the square root, and transposing, $x = -\frac{1}{2}d \pm \sqrt{\frac{a^2 - d^2}{2} + \frac{1}{4}d^2}$.

$=$ (substituting known values) $-\frac{1}{2} \pm \sqrt{\frac{25 - 1}{2} + \frac{1}{4}} = -\frac{1}{2} \pm 3\frac{1}{2} = 3$

or -4 , of which values the last (being irrelevant) is to be rejected.

Wherefore, x being known, the triangle is determined.

PROBLEM VIII.

If a be put for the perpendicular, and
 b for the base of the triangle;
 c for the area of the rectangle; and
 x for the length of the rectangle parallel to the base of
 the triangle, it will be

As $b : a :: x : \frac{ax}{b}$ the difference of the altitudes of the rectangle
 and triangle.

Therefore the altitude of the rectangle is $a - \frac{a}{b}x$.

But $c = x (a - \frac{a}{b}x) = ax - \frac{a}{b}x^2$, that is, $x = \frac{1}{2}b \pm \sqrt{\frac{b^2}{4} - \frac{b}{a}c}$ the

(Key to Vol. I. page 363.)

length of the rectangle; consequently $\frac{c}{\frac{1}{2}b \pm \sqrt{\frac{b^2}{4} - \frac{b}{a}c}}$ = the breadth;
 which were to be determined.

PROBLEM IX.

Put a and b for the two segments of the base,
 x for the side adjacent to a ,
 y for the other side of the triangle; and
 $m : n$, the ratio x has to y .

Then, (since $m : n :: x : y$) it follows that $y = \frac{n}{m} x$.

But $x^2 - a^2 = y^2 - b^2 = \frac{n^2 x^2}{m^2} - b^2$.

That is, $m^2 x^2 - n^2 x^2 = m^2 a^2 - m^2 b^2$.

Hence $x = \sqrt{\frac{m^2 a^2 - m^2 b^2}{m^2 - n^2}}$
 And $y = \frac{n}{m} \sqrt{\frac{m^2 a^2 - m^2 b^2}{m^2 - n^2}}$ } which were to be determined.

PROBLEM X.

Put b for half the base,
 c for the line from the vertex,
 a for the sum of the two sides; and
 x for one of the sides.

Then $a - x$ is the other side of the triangle.

But (Theor. 38) $2b^2 + 2c^2 = a^2 - 2ax + 2x^2$. Wherefore
 $x = \frac{1}{2}a \pm \sqrt{b^2 + c^2 - \frac{1}{4}a^2}$, and $a - x = \frac{1}{2}a \mp \sqrt{b^2 + c^2 - \frac{1}{4}a^2}$.

Consequently the triangle is determined.

(Page 364.)

PROBLEM XI.

Let a represent one of the sides of the triangle,
 b the other side,
 c the line bisecting the vertical angle, and
 x the base;

(Key to Vol. I. page 364.)

it will be, As $(a+b) : x :: a : \frac{ax}{a+b}$ the segment of the base, adjacent to a .

And, As $(a+b) : x :: b : \frac{bx}{a+b}$ the seg. of the base, adjacent to b .

Wherefore (*Because in any triangle having the vertical angle bisected, the rectangle of the two sides less the rectangle of the segments of the base is equal to the square of the line bisecting the vertical angle*) [Theor. lxiv.] $ab - \left(\frac{ax \times bx}{a+b}\right) = c^2$.

Hence $x^2 = \frac{a^3b + 2a^2b^2 + ab^3 - a^2c^2 - 2a^2cb - b^2c^2}{ab}$; which equation

resolved, $x = (a+b)\sqrt{\frac{ab-c^2}{ab}}$ = the base of the triangle Q. E. I.

PROBLEM XII.

If a be put for the line joining the acute angle at the base and the middle of the perpendicular,

b for the line from the middle of the base to the vertical angle,

x for half the base, and

y for half the perpendicular;

Then $(2y)^2 = b^2 - x^2$; or $y^2 = \frac{b^2 - x^2}{4}$.

Also $a^2 = \frac{b^2 - x^2}{4} + 4x^2$, or $4a^2 - b^2 = 15x^2$, that is,

$x = \sqrt{\frac{4a^2 - b^2}{15}}$, hence the base of the triangle = $2\sqrt{\frac{4a^2 - b^2}{15}}$

And $y^2 = \frac{b^2 - x^2}{4} = \frac{16b^2 - 4a^2}{4 \times 15}$, or $y = \sqrt{\frac{4b^2 - a^2}{15}}$. Wherefore

$2y = 2\sqrt{\frac{4b^2 - a^2}{15}}$ = the perpendicular. Lastly,

$\sqrt{\frac{12a^2 + 12b^2}{15}}$ = the hypotenuse. Q. E. I.

PROBLEM XIII.

Put a for the radius of the inscribed circle,

p for the perimeter of the triangle,

x for half the sum of the base and perpendicular, and

y for half their difference:

(Key to Vol. I. page 364.)

Then $p - 2x =$ the hypotenuse, and $\frac{ap}{2} =$ the area of the triangle.

But $x^2 - y^2 = ap$, and $(x+y)^2 + (x-y)^2$ being equal to $(p-2x)^2$ or $p^2 - 4px + 4x^2$, it follows that, $p^2 - 4px + 4x^2 = 2x^2 + 2y^2$.

And $2 \times (x^2 - y^2) = 2 \times ap$. Therefore by addition,

$$\left\{ \begin{array}{l} 2x^2 - 2y^2 = 2ap \\ 2x^2 + 2y^2 = p^2 - 4px + 4x^2 \end{array} \right\}$$

$$4x^2 = p^2 + 2ap - 4px + 4x^2, \text{ or}$$

$$p^2 + 2ap - 4px = 0. \text{ Hence } x = \frac{p^2 + 2ap}{4p} = \frac{1}{2}(\frac{1}{2}p + a)$$

And $y = \sqrt{x^2 - ap} = \sqrt{\frac{1}{4}(\frac{1}{2}p + a)^2 - ap}$. Therefore


$$\left. \begin{array}{l} \frac{1}{2}(\frac{1}{2}p + a) + \sqrt{\frac{1}{4}(\frac{1}{2}p + a)^2 - ap} = \text{the base or perpendicular} \\ \frac{1}{2}(\frac{1}{2}p + a) - \sqrt{\frac{1}{4}(\frac{1}{2}p + a)^2 - ap} = \text{the perpendicular or base} \end{array} \right\} \text{Q. E. I.}$$

Otherwise.

Let ABC be the given triangle, DEF the inscribed circle, whereof OD, OE, OF are three radii at right angles to the sides of the triangle, AF is equal to AD , and BF to BE : also EC and CD are equal.

Put p for the perimeter,
 r for the given radius.
 x for EB , and
 y for AD .

Then $AC = r + y$; $BC = r + x$; and
 $AB = x + y$.



But $x + y + (r + y) + (r + x) = p$, or $x + y = \frac{p - 2r}{2} = \frac{1}{2}p - r$, and

$$(x + y)^2 = (r + y)^2 + (r + x)^2; \text{ or } r(x + y) = xy - r^2.$$

In the 1st equation $y = \frac{1}{2}p - r - x$; substitute, therefore, this value for y in the 2d equation, and reduce.

$$x^2 - (\frac{1}{2}p - r)x = -\frac{1}{2}rp. \text{ Wherefore}$$

$$x = \frac{1}{2}(\frac{1}{2}p - r) \pm \sqrt{\frac{1}{4}(\frac{1}{2}p - r)^2 - \frac{1}{2}rp}. \text{ And}$$

$$y = \frac{1}{2}(\frac{1}{2}p - r) \mp \sqrt{\frac{1}{4}(\frac{1}{2}p - r)^2 - \frac{1}{2}rp}. \text{ Hence, by adding } r \text{ to each,}$$

$$BC = \frac{1}{2}(\frac{1}{2}p + r) \pm \sqrt{\frac{1}{4}(\frac{1}{2}p - r)^2 - \frac{1}{2}rp}$$

$$AC = \frac{1}{2}(\frac{1}{2}p + r) \mp \sqrt{\frac{1}{4}(\frac{1}{2}p - r)^2 - \frac{1}{2}rp} \left. \right\} \text{Q. E. I.}$$

(Key to Vol. I. page 364.)

PROBLEM XIV.

Let a represent the base,
 x one of the segments of the base by the perpendicular,
 r the side of the triangle adjacent to x ,
 s the other side of the triangle,
 b the perpendicular; and
 $m : n$ the ratio of $r : s$.

Then $a - x$ is the other segment of the base. Also
 $x^2 + b^2 = r^2$, and $x^2 + b^2 + a^2 - 2ax = s^2$.

But $m^2 : n^2 :: (x^2 + b^2) : (x^2 + b^2 + a^2 - 2ax)$ Wherefore
 $(m^2 - n^2)x^2 - 2am^2x = (n^2 - m^2)b^2 - a^2m^2$. That is,

$$x^2 - \frac{2am^2}{m^2 - n^2}x = -b^2 - \frac{a^2m^2}{m^2 - n^2}, \text{ which equation resolved,}$$

$$x = \frac{am^2}{m^2 - n^2} \pm \sqrt{\frac{a^2m^4}{(m^2 - n^2)^2} - \frac{a^2m^2}{m^2 - n^2} - b^2}.$$

$$r = \sqrt{b^2 + \left(\frac{am^2}{m^2 - n^2} \pm \sqrt{\frac{a^2m^4}{(m^2 - n^2)^2} - \frac{a^2m^2}{m^2 - n^2} - b^2}\right)^2}. \text{ And}$$

$$s = \frac{n}{m} \sqrt{b^2 + \left(\frac{am^2}{m^2 - n^2} \pm \sqrt{\frac{a^2m^4}{(m^2 - n^2)^2} - \frac{a^2m^2}{m^2 - n^2} - b^2}\right)^2}. \text{ Conse-}$$

quently the triangle is determined, which was to be done.

PROBLEM XV.

Put a for the hypotenuse,
 c for the side of the given square,
 x for the perpendicular, and
 y for the base of the triangle.

Then As $x : y :: (x - c) : c$, or $y(x - c) = cx$, that is $xy = c(x + y)$

But $a^2 = x^2 + y^2$. Add $2xy$ to both sides of the equation,
 $a^2 + 2xy = x^2 + 2xy + y^2$.

Substitute $2c(x + y)$ for $2xy$ on the 1st side,
 $a^2 + 2c(x + y) = x^2 + 2xy + y^2$, or
 $a^2 = (x + y)^2 - 2c(x + y)$. Therefore
 $x + y = c \pm \sqrt{a^2 + c^2}$, and
 $y = c - x \pm \sqrt{a^2 + c^2}$.

(Key to Vol. I. page 364.)

Substitute this value of y in the equation $xy=c(x+y)$.

$$x(c-x \pm \sqrt{a^2+c^2})=c(c \pm \sqrt{a^2+c^2}).$$

That is, $x^2 - (c \pm \sqrt{a^2+c^2})x = -c(c \pm \sqrt{a^2+c^2})$.*

Which equation resolved,

$$x = \frac{1}{2}(c \pm \sqrt{a^2+c^2}) \pm \sqrt{\frac{1}{4}a^2 - \frac{1}{2}c^2 \mp \frac{1}{2}c\sqrt{a^2+c^2}} = \text{the perpend.}$$

And


$$y = \frac{1}{2}(c \pm \sqrt{a^2+c^2}) \mp \sqrt{\frac{1}{4}a^2 - \frac{1}{2}c^2 \mp \frac{1}{2}c\sqrt{a^2+c^2}} = \text{the base.}$$

But a plane right angled triangle is determined when the base and perpendicular are known; therefore, &c.

* Because (line 11 of the Solution) the value of $x+y$ is $c \pm \sqrt{a^2+c^2}$.

PROBLEM XVI.

Let ABC be the given circle, whereof o is the center. Inscribe in it the equilateral triangle pqr (Eucl. iv. 15); join oq , or , op ; and produce op till ps equal the half of pr , or of pq . Draw sr , and parallel to sr through the point p draw pm meeting or in m ; and through m draw ml parallel to rp ; through L , LN parallel to pq ; and join mn .



L, M, N are the centers of three circles that shall touch one another, and the circumference of the given circle ABC .

For bisect pq in H , and join SH ; and parallel to SH draw PF .

(Key to Vol. I. page 364.)

Because POQ is an isocetes triangle, and that, LN is parallel to the base, PL is equal to QN : And because SH, PF are parallel, (PH, LF being also parallel,) and that, SP, PH are equal, PL, LF are equal. In the same manner it may be proved that QN is equal to NF , to NE , to EM , to MR , &c. Moreover it is evident that, NF, FL are in the same straight line.

Putting, therefore, $a = \text{radius of the given circle, and}$
 $x = \text{radius of one of the inscribed circles,}$

It is, (because $NF = \frac{1}{2}MN$, and, by similar triangles $OE = \frac{1}{2}OM$)

$$(a-x)^2 - \left(\frac{a-x}{2}\right)^2 = x^2. \quad \text{Whence}$$

$$x = -3a \pm \sqrt{12a^2} = 2a\sqrt{3} - 3a. \quad \text{Which was required.}$$

PROBLEM XVII.

Put p for the perimeter of the triangle.
 a for the perpendicular falling on the hypotenuse,
 x for the greater, } if unequal.
 y for the less side, }

Also z for the hypotenuse.

$$\text{Then } z = p - x - y. \quad \text{And}$$

$$z^2 = x^2 + y^2 = p^2 - 2p(x+y) + x^2 + 2xy + y^2.$$

$$\text{That is, } p(x+y) - \frac{1}{2}p^2 = xy; \quad \text{and by comparison,}$$

$$(a+p) \times (x+y) = ap + \frac{1}{2}p^2. \quad \text{That is,}$$

$$x+y = p \left(\frac{a + \frac{1}{2}p}{a+p}\right), \quad \text{and, } y = p \left(\frac{a + \frac{1}{2}p}{a+p}\right) - x.$$

By substituting these values for $(x+y)$, and for y , in the foregoing equations, the following equation is obtained:

$$(a+p)x - p\left(a + \frac{1}{2}p\right)x = \frac{1}{2}ap^2. \quad \text{Hence}$$

$$x = p \left(\frac{a + \frac{1}{2}p}{2a + 2p}\right) \pm \frac{p}{2(a+p)} \sqrt{\left(a - \frac{1}{2}p\right)^2 - 2a^2}. \quad \text{And}$$


$$y = \frac{p\left(a + \frac{1}{2}p\right)}{2a + 2p} \mp \frac{p}{2a + 2p} \sqrt{\left(a - \frac{1}{2}p\right)^2 - 2a^2}. \quad \text{Also}$$

$$z = p - x - y = \frac{p^2}{2a + 2p}. \quad \text{Wherefore the triangle is determined.}$$

(Key to Vol. I. page 364.)

PROBLEM XVIII.

In the annexed fig. let CD be the greater, and AD the less of the two lines of which the difference is given, and let DH be a production of CD, and AH perpendicular to CH, CH is equal to HD, because the angle ADH is equal to the sum of the angles ACD, CAD, together equal to half a right angle, and the angle at H a right angle.



If, therefore, a be put for AC the hypotenuse,
 x for CD,
 y for AD,
 b for the difference of x and y ,
 r for DH, and
 s for AH,

Then (Theor. 4.) $r = s = \frac{y}{\sqrt{2}}$.

But $x^2 + y^2 + 2rx = a^2 = x^2 + y^2 + xy\sqrt{2}$.

Now substituting $x - b$ for y , and c for the $\sqrt{2}$, it will be $x^2 - 2bx + b^2 + x^2 + cx^2 - cbx = a^2$, that is, $(2+c)x^2 - (2+c)bx + b^2 = a^2$. Whence

$$x = \frac{b}{2} \pm \sqrt{\frac{a^2 - b^2}{2+c} + \frac{1}{4}b^2}, \text{ and}$$

$y = \sqrt{\frac{a^2 - b^2}{2+c} + \frac{1}{4}b^2} - \frac{1}{2}b$. Consequently the radius of the inscribed circle is known, and the triangle determined,

For put $m = \frac{1}{2}b \pm \sqrt{\frac{a^2 - b^2}{2+c} + \frac{1}{4}b^2} = CD$,

$$n = \sqrt{\frac{a^2 - b^2}{2+c} + \frac{1}{4}b^2} - \frac{1}{2}b = AD.$$

And let de, df, dg , be three radii at right angles to the sides of the triangle; likewise put w for Ag , and z for cg .

$$z^2 - w^2 = m^2 - n^2$$

(Key to Vol. I. page 365.)

Also $z + w = a$; and by division

$$z - w = \frac{m^2 - n^2}{a}, \text{ that is, } z = \frac{a^2 + m^2 - n^2}{2a}; w = \frac{a^2 - m^2 + n^2}{2a}$$

And $dg = \sqrt{m^2 - \frac{(a^2 + m^2 - n^2)^2}{4a^2}} = \text{the radius of the inscribed circle.}$

PROBLEM XIX.

Put a for half the base, d for the difference of the two sides,
 b for the perpendicular,
 x for the excess of the greater segment of the base above a .
 y for the greater side of the triangle, and
 z for the less.

Then $y^2 = b^2 + (a+x)^2$, or $y = \sqrt{b^2 + (a+x)^2}$. Also
 $z^2 = b^2 + (a-x)^2$, or $z = \sqrt{b^2 + (a-x)^2}$.

But $\sqrt{b^2 + (a+x)^2} - d = \sqrt{b^2 + (a-x)^2}$. Hence
 $b^2 + (a+x)^2 - 2d\sqrt{b^2 + (a+x)^2} + d^2 = b^2 + (a-x)^2$: and by red.
 $4ax + d^2 = 2d\sqrt{b^2 + (a+x)^2}$, or $16a^2x^2 + 8ad^2x + d^4 = 4d^2(b^2 + a^2 + 2ax + x^2)$.

Consequently $x = \sqrt{\frac{4d^2(a^2 + b^2) - d^4}{16a^2 - 4d^2}}$, which value put equal c .

$y = \sqrt{(a+c)^2 + b^2}$. } which were to be determined.
 $z = \sqrt{(a-c)^2 + b^2}$. }

PROBLEM XX.

Let a represent the perpendicular,
 b half the base,
 c the product of the two sides,
 x the excess of the greater segment of the base by the
 perp. above b ;

And put y for the greater side of the triangle,
 z for the less.

Then $y = \sqrt{a^2 + (b+x)^2} = \frac{c}{z}$

$$z = \sqrt{a^2 + (b-x)^2} = \frac{c}{y} = \frac{c}{\sqrt{a^2 + (b+x)^2}}, \text{ (or squared)}$$

$$a^2 + (b-x)^2 = \frac{c^2}{a^2 + (b+x)^2}. \text{ Hence}$$

(Key to Vol. I. page 365.)

$$e^2 = (a^2 + b^2)^2 + x^4 + 2(a^2 - b^2)x^2, \text{ that is, } x^4 + 2(a^2 - b^2)x^2 = (a^2 + b^2)^2 - c^2.$$

And, this equation resolved,

$$x^2 = -(a^2 - b^2) \pm \sqrt{(a^2 + b^2)^2 - c^2 + (a^2 - b^2)^2}.$$

or $x = \sqrt{(b^2 - a^2) \pm \sqrt{(a^2 + b^2)^2 - c^2 + (a^2 - b^2)^2}}$, which value put = d .


$$\left. \begin{aligned} y &= \sqrt{a^2 + (b+d)^2} \\ z &= \sqrt{a^2 + (b-d)^2} \end{aligned} \right\} \text{which were to be determined.}$$

PROBLEM XXI.

Let ABC be the triangle, and AF, BE, and CD the three given lines.

If a represent AF, b , CD. And c , BE; also

If x be put for BC, y for AB, and z for AC;



Then (Theor. 38) $y^2 + z^2 = 2a^2 + \frac{1}{2}x^2$, that is,

$$y^2 + z^2 - \frac{1}{2}x^2 = 2a^2. \text{ For the same reason}$$

$$x^2 + z^2 - \frac{1}{2}y^2 = 2b^2. \text{ And}$$


$$x^2 + y^2 - \frac{1}{2}z^2 = 2c^2. \text{ Comparing the 1st equation with double the sum of the 2d and 3d.}$$

$4\frac{1}{2}x^2 = 2(2b^2 + 2c^2 - a^2)$ or, divided by $4\frac{1}{2}$, and the square root extracted,

$$\left. \begin{aligned} x &= \frac{2}{3}\sqrt{2b^2 + 2c^2 - a^2} \\ y &= \frac{2}{3}\sqrt{2a^2 + 2c^2 - b^2} \\ z &= \frac{2}{3}\sqrt{2a^2 + 2b^2 - c^2} \end{aligned} \right\} \text{Q. E. I.}$$

PROBLEM XXII.

Conceiving the figure constructed (as in the accompanying diagram) draw lines from the center to each of the angles, and to the points of contact. Produce either of the lines joining the angular points and center (as BO) indefinitely through the opposite side, and on it pro-



(Key to Vol. I. page 365.)

duced let fall a perpendicular (as AG) from that angular point from which the perpendicular falls without the triangle.

Put a for Ad ,
 b for dB , and
 c for cf . Also
 x for $od=oe=of$.


Then $\sqrt{b^2+x^2}$ [BO] : x [od] :: $a+b$ [AB] : $\frac{ax+bx}{\sqrt{b^2+x^2}}$ [AG].

But BO : Bd :: AB : BG. that is, $\frac{ab+b^2}{\sqrt{b^2+x^2}} - \sqrt{b^2+x^2} = \frac{ab-x^2}{\sqrt{b^2+x^2}}$.

And AG : OG :: cf : of, whence $ax+bx : ab-x^2 :: c : x$.

Therefore $x = \sqrt{\frac{abc}{a+b+c}}$. Which was required.

PROBLEM XXIII.



Let ABC be the proposed triangle, BFDE the inscribed square, OG and OP radii of the inscribed circle at right angles to AB and AC, and BD a diagonal of the inscribed square; also let BQ be perpendicular to AC from the right angle. Put

a for the side of the given square,
 b for the radius of the given circle, and
 x for the segment AQ of the base AC by the perpend. BQ,

Then $FG = a - b$ since $GB = OG$; and

$a - b : a :: b : \frac{ab}{a - b} = BQ$ (because $GF : BF :: OD : BD :: OP : BQ$).

Wherefore (since $BD = \sqrt{2a^2}$), $DQ = \sqrt{2a^2 - \frac{a^2b^2}{(a-b)^2}}$. Let this

(Key to Vol. I. page 365.)

value of dQ be recognized in c , and put d for $\frac{ab}{a-b}$. It is


As $x : d :: d : \frac{d^2}{x} = cQ$ [Theor. 87]. And $x + c : \frac{d^2}{x} - c :: x : d$.*

Hence $dx + dc = d^2 - cx$. That is $x = \frac{d^2 - dc}{d + c} = AQ$, wherefore the triangle is determined.

* For the triangles AFD , DEC are similar; also the triangle AQB is similar to AFD , and consequently to DEC . Therefore $AD : DC :: AF : DE (= DF) :: AQ : QB$.

PROBLEM XXIV.

Having constructed the figure, draw lines from the center of the circle to the points of contact, as OG , OF , OE . Also produce one of the given lines (as BO) indefinitely beyond the center, and on it produced, from either of the other angles let fall a perpendicular (as CD).



Then because the angles ABC , BCA , and CAB are together equal to two right angles, the angles OBC , OCB , and OAC are together equal to half of two right angles. But the angle OBC together with the angle OCB , is equal to the angle COD , therefore the angles COD , OAE are equal to the angles COD and OCB , each to each, and either pair to a right angle. Put $a = BO$

$$b = AO$$

$$c = CO, \text{ and}$$

put x for the radius of the inscribed circle.

Then because $AO : OE :: CO : OD$, $b : x :: c : \frac{c}{b}x = OD$.

Therefore $CD = \sqrt{c^2 - \frac{c^2x^2}{b^2}} = \frac{c}{b} \sqrt{b^2 - x^2}$. But, (Theor. 36)

$$BC^2 = BO^2 + CO^2 + 2(BO \times OD), \text{ or } AC = \sqrt{\frac{b(a^2 + c^2) + 2acx}{b}}$$


(Key to Vol. I. page 365.)

Now $BC : CD :: BO : OF$. That is, $\sqrt{\frac{b(a^2+c^2)+2acx}{b}} : \frac{c}{b} \sqrt{b^2-x^2} :: a : x$. Hence $bx^2 (b \times a^2 + c^2 + 2acx) = a^2c^2(b^2 - x^2)$, and by reduction, $x^3 + \left(\frac{a^2b^2 + a^2c^2 + b^2c^2}{2abc}\right)x^2 = \frac{1}{2}abc$, an equation in which x is determinable, and, x known, *the triangle is determined.*

The equation, however, arising from this problem, as it contains three dimensions of the unknown quantity, admits of *no Geometrical construction by means of a CIRCLE and RIGHT LINES.*

PROBLEM XXV.

Construct a right angled triangle, and inscribe in it a circle; also draw right lines from the center to the points of contact, and to the two acute angles, as in the figure ABCDEFG. Put x for the base (AB), y for the perpendicular (BC) and let a represent the given radius (DG, DE, or DF), b the hypotenuse (AC).



Then, because GB and DF are equal, $y - a$ is the expression for CG ; and $x - a = AF$. Also CE and CG are equal [Eucl. i. 26]; and AF is equal to AE .

But $AE + CE = AC$; that is, $(y - a) + (x - a) = b$, or $x + y = 2a + b$.

Now $x^2 + y^2 = b^2$; comparing, therefore, the double of this equation, with the square of the preceding, $x^2 - 2xy + y^2 = b^2 - 4ab - 4a^2$. Hence

$x - y = \sqrt{b^2 - 4ab - 4a^2}$, consequently

$x = a + \frac{1}{2}b \pm \frac{1}{2}\sqrt{b^2 - 4ab - 4a^2}$, and

$y = a + \frac{1}{2}b \mp \frac{1}{2}\sqrt{b^2 - 4ab - 4a^2}$.

Wherefore the triangle is determined.

(Key to Vol. I. page 365.)

PROBLEM XXVI.

Let ABC be the triangle, and BD the line bisecting the vertical angle.

Draw the diameter FG at right angles to the base. If BD be produced it will meet the circumference in G, because equal angles stand on equal arcs. Put

a for BD,
 b for AE or EC,
 c for FG,

And x for EG.

Then [Theor. 61]

$x(c-x) = b^2$, hence

$x = \frac{1}{2}c \pm \sqrt{\frac{1}{4}c^2 - b^2}$; for this value of x put e ; and join BF, also let y represent DG.

The angle GBF being [Theor. 52] a right angle, the triangles GDE, GFB, are [Theor. 84] similar; therefore $y : e :: c : a + y$, that is, $ay + y^2 = ec$, hence $y = \pm \sqrt{ec + \frac{1}{4}a^2} - \frac{1}{2}a$. Put f for this value of y ; and $z = DC$.

Then [Theor. 61] $af = 2bz - z^2$, that is,

$z = b \pm \sqrt{b^2 - af}$. Put this value of $z = g$, and from B draw BH at right angles to FG.

GD : GB :: DE : BH, that is,

As $f : (a + f) :: (b - g) : (b - g + \frac{ab + ag}{f}) = BH$ the distance of the perpendicular from the middle of the base, = EP. Join BP.

GE : EH (=BP) :: GD : DB, that is,


As $e : BP :: f : a$, or

$BP = \frac{ae}{f} =$ the perpendicular of the triangle ABC. But

$AB = \sqrt{AP^2 + BP^2} = \sqrt{\left\{ b + (b - g + \frac{ab + ag}{f}) \right\}^2 + p^2}$. [where p represents PB]; and

$BC = \sqrt{PC^2 + BP^2} = \sqrt{\left\{ b - (b - g + \frac{ab + ag}{f}) \right\}^2 + p^2}$.

Wherefore the triangle is determined, which was to be done.



K E Y

TO

HUTTON'S COURSE OF MATHEMATICS.

PLANE TRIGONOMETRY.


(Key to Vol. II. page 9.)

THEOREM I.

Ex. 2.

1. Geometrically.

Draw an indefinite line, and take in it AB 365 poles from any convenient scale of equal parts. At A make the angle BAC $57^{\circ} .. 12'$, and at B make the angle ABC $24^{\circ} .. 45'$, ABC is the triangle in species. But



AC measures $154\frac{1}{3}$ poles } on the scale employed for AB.
 BC - - - - - 310 poles }

And the angle ACB by a line of Chords or Goniometer* is $98^{\circ} .. 3'$.

* Any Instrument to measure Angles by Inspection.

2. Arithmetically.

$180^{\circ} - \{ 57^{\circ} .. 12' (= \angle A) + 24^{\circ} .. 45' (= \angle B) \} = 98^{\circ} .. 3'$ the angle C.

(Key to Vol. II. page 9.)

And

As Sin. 98° .. 3'	∠ C - - -	Log. ar. co. -	10·004301
: Sin. 24° .. 45'	∠ B - - - - -	Log.	9·621861
:: 365 poles	AB - - - - -	Log.	2·562293
: 154·33 poles AC			Log. 2·188455

Also,

As Sin. 98° .. 3'	∠ C - - -	Log. ar. co. -	10·004301
: Sin. 57° .. 12'	∠ A - - - - -	Log.	9·924572
:: 365 poles	AB - - - - -	Log.	2·562293
: 309·86 poles BC			Log. 2·491166

3. *Instrumentally.*


The sine of 98° .. 3' = sine 81° .. 57', and the extent from 81° .. 57' to 24° .. 45' on the line of sines, reaches from 365 to 154 $\frac{1}{3}$ on the line of numbers, for the length of the side AC in poles. Likewise the extent from 81° .. 57' to 57° .. 12' on the line of sines reaches from 365 to 310 *nearly*, on the line of numbers, for the length of BC in poles.

(Page 10.)

Ex. 3.

1. *Geometrically.*

Draw an indefinite line, and take in it AC 120 feet. At A make the angle CAB 57° .. 28', and with a radius of 112 from the scale on which AC measures 120, describe the arc *mno* cutting AB in the points B and B, either of the triangles ABC answers the conditions, shewing that the case is ambiguous. Now



AB measures $\left\{ \begin{matrix} 16\frac{1}{2} \\ 112\frac{1}{2} \end{matrix} \right\}$ feet, on the scale employed for AC.

The angle B $\left\{ \begin{matrix} 115^\circ \dots 25' \\ 64^\circ \dots 35' \end{matrix} \right\}$ by a Goniometer, or line of Chords.

And the angle c $\left\{ \begin{matrix} 7^\circ \dots 7' \\ 57^\circ \dots 57' \end{matrix} \right\}$ by subtraction, or admeasurement.

(Key to Vol. II. page 10.)

2. *Arithmetically.*

$$\begin{array}{l}
 \text{As } 112 \text{ feet } BC \text{ - - - - - } \text{Log. ar. co. - } 3.95078 \\
 : 120 \text{ feet } AC \text{ - - - - - } \text{Log. } 2.07918 \\
 :: \text{Sin. } 57^\circ .. 28' \angle A \text{ - - - - - } \text{Log. } 9.92587 \\
 \\
 : \text{Sin. } \left\{ \begin{array}{l} 64^\circ .. 35' \\ 115^\circ .. 25' \end{array} \right\} \angle B \text{ - - - - } \text{Log. } 9.95583
 \end{array}$$

Hence

$$\text{The angle } c = \left\{ \begin{array}{l} 7^\circ .. 7' \\ 57^\circ .. 57' \end{array} \right\} \text{ the supplement of } \angle (A+B)$$

And

$$\begin{array}{l}
 \text{As Sin. } 57^\circ .. 28' \angle A. \\
 : \text{Sin. } \left\{ \begin{array}{l} 7^\circ .. 7' \\ 57^\circ .. 57' \end{array} \right\} \angle C. \\
 :: 112 \text{ feet, } BC. \\
 : \left\{ \begin{array}{l} 16.47 \\ 112.6 \end{array} \right\} \text{ feet, } AB
 \end{array}$$

3. *Instrumentally.*

The extent from 112 to 120 on the line of numbers reaches from $57^\circ .. 28'$ to $64^\circ .. 35'$ (=sine of $115^\circ .. 25'$) on the line of sines, for the $\angle B$. Hence $\angle C = \left\{ \begin{array}{l} 7^\circ .. 7' \\ 57^\circ .. 57' \end{array} \right\}$

Again the extent from $57^\circ .. 28'$ to $7^\circ .. 7'$, or $57^\circ .. 57'$ on the line of sines, reaches from 112 to $16\frac{1}{2}$ or 112.6 on the line of numbers, the two values of AB in feet.

(Page 12.)

THEOREM II.

Ex. 2. [See the figure in Example 2, Theorem 1.]

1. *Geometrically.*

From a scale of equal parts take AB 365 poles. At A make the angle BAC $57^\circ .. 12'$, and from A with a radius of 154.33 on the scale on which AB measures 365, determine the point c ; lastly join BC . ABC is the triangle required.

BC measures 309.8 on the scale employed for AB .

(Key to Vol. II. page 12.)

And

By the Goniometer, or a line of Chords $\left\{ \begin{array}{l} \angle B = 24^\circ .. 45' \\ \angle C = 98^\circ .. 3' \end{array} \right\}$ 2. *Arithmetically.*

As 519.33 $AB + AC$ - - - - - Log. ar. co. - 3.2845566
 : 210.67 $AB - AC$ - - - - - Log. 2.3236027
 :: Tang. $61^\circ .. 24'$, half supp. $\angle A$ - - - - - Log. 10.2634301

: Tang. $36^\circ .. 39'$ half the diff. of the $\angle^{les} B$ and c . Log. 9.8715894

But $61^\circ .. 24' + 36^\circ .. 39' = 98^\circ .. 3'$ the angle c .
 And $61^\circ .. 24' - 36^\circ .. 39' = 24^\circ .. 45'$ the angle B .

Again,

As Sin. $98^\circ .. 3'$ $\angle C$ - - - Log. ar. co. - 10.004301
 : Sin. $57^\circ .. 12'$ $\angle A$ - - - - - Log. 9.924572
 :: 365 poles AB - - - - - Log. 2.562293

: 309.86 poles BC - - - - - Log. 2.491166

3. *Instrumentally.*

The extent from $519\frac{1}{3}$ to 210.7 on the line of numbers reaches from $61^\circ .. 24'$ beyond 45° on the line of tangents; but this excess above 45° applied backwards from 45° reaches to $36^\circ\frac{2}{3}$ half the diff. of the $\angle^{les} B$ and c . Hence $c = 98^\circ .. 3'$ and $B = 24^\circ .. 45'$. Again the extent from $81^\circ .. 57'$ to $57^\circ .. 12'$ on the line of sines reaches from 365 to 309.8 on the line of numbers, for the length of BC in poles.

Ex. 3. [See the Greater triangle ABC in the figure to Example 3, Theorem 1.]

1. *Geometrically.*

Take AC 120 yards from any convenient scale of equal parts, and at c make the angle ACB $57^\circ .. 57'$; then from c with a radius of 112 from the scale on which AC measures 120, determine B ; lastly join AB .

AB measures 112.6 on the scale employed for AC .

And

By the Goniometer, or a line of Chords $\left\{ \begin{array}{l} \angle A = 57^\circ .. 28' \\ \angle B = 64^\circ .. 35' \end{array} \right\}$

(Key to Vol. II. page 12.)

2. *Arithmetically.*

As 232 yards AC + BC - - - - - Log. ar. co. - 3·6345120
 : 8 yards AC - BC - - - - - Log. 0·9030900
 :: Tang. 61° .. 1' .. 30'' half supp. ∠ C - - - - Log. 10·2566460
 : Tang. 3° .. 34' half the diff. of the ∠^{les} A and B. Log. 8·7942480

But 61° .. 1'½ + 3° .. 34' = 64° .. 35'½ ∠ B. } *
 And 61° .. 1'½ - 3° .. 34' = 57° .. 27'½ ∠ A. }

* These angles differ half a minute from the angles given in the figure, because the calculation throughout has not been carried to seconds.

Again,

As Sin. 57° .. 27'½ ∠ A - - - - - Log. ar. co. - 10·0741722
 : Sin. 57° .. 57' ∠ C - - - - - Log. 9·9281834
 :: 112 yards BC - - - - - Log. 2·0492180
 : 112·61 yards AB - - - - - Log. 2·0515736

3. *Instrumentally.*

The extent* from 232 to 8 on the line of numbers reaches from 61° .. 1' beyond 45° on the line of tangents; but this excess above 45°, applied backwards from 45°, reaches to 3° .. 34' half the diff. of the ∠^{les} A and B.

Hence { ∠ B = 64° .. 35' }
 { ∠ C = 57° .. 27' }

Again the extent from 57° .. 27' to 57° .. 57' on the line of sines reaches from 112 to 112·6 on the line of numbers, for the length of AB in yards.

* It is necessary to employ some multiple of 232 and 8, instead of the numbers themselves, and such that the multiple of 8 may exceed 100, if a two-foot scale be employed.

(Page 14.)

THEOREM III.

Ex. 2. [See the figure in Example 2, Theorem I.]

(Key to Vol. II. page 14.)

1. Geometrically.

From any convenient scale of equal parts take AB 365 poles, and with a radius 154.33 from the same scale, and center A , describe an arc; also from B as a center with a radius 309.86 of the scale on which AB measures 365, cut the former arc in c . Join AC and BC .

$$\text{By a line of Chords, or Goniometer } \left\{ \begin{array}{l} \angle A = 57^\circ \dots 12' \\ \angle B = 24^\circ \dots 45' \\ \angle C = 98^\circ \dots 3' \end{array} \right\}$$

* A typographical error in the Course.

2. Arithmetically.

As 365 poles AB - - - - -	Log. ar. co. - 3.4377071
: 469.19 poles $AC + BC$ - - - - -	Log. 2.6666958
:: 155.53 poles $BC - AC$ - - - - -	Log. 2.1918142
<hr/>	
: 197.8 poles <i>diff. of the segments of AB by a perp. from c</i> } - - -	Log. 2.2962171
<hr/>	

Hence $Ap = 83.6$ poles, and $Bp = 281.4$, by $\frac{1}{2}$ sum and $\frac{1}{2}$ diff. of the segments of the base.

And

As 154.33 pol. AC : 83.6 pol. Ap :: $\text{Sin. } 90^\circ \angle APC$: $\text{Cos. } 57^\circ \dots 12' \angle A$.
 As 309.86 pol. BC : 281.4 pol. Bp :: $\text{Sin. } 90^\circ \angle BPC$: $\text{Cos. } 24^\circ \dots 45' \angle B$.

Wherefore, by subtraction, $\angle C = 98^\circ \dots 3'$.

3. Instrumentally.

The extent from 365 to 469.2 reaches from $155\frac{1}{2}$ to 198 nearly, on the line of numbers, for the difference of Ap and Bp .

Hence $Ap = 83\frac{1}{2}$ poles, and $Bp = 281\frac{1}{2}$ poles.

Again the extent from $154\frac{1}{2}$ to $83\frac{1}{2}$ on the line of numbers, reaches on the line of sines from 90° to $32^\circ \dots 48'$, of which the complement is $57^\circ \dots 12'$ the $\angle A$.

Also the extent from 310 to $281\frac{1}{2}$ on the line of numbers reaches on the line of sines from 90 to $65^\circ \dots 15'$, of which the complement is $24^\circ \dots 45'$ the $\angle B$; and consequently the $\angle C = 98^\circ \dots 3'$ as before.

(Key to Vol. II. page 15.)

Ex. 3. [See the Greater triangle ABC in the fig. to Ex. 3. Th. 1. but transpose B and c.]

1. Geometrically.

Take AB 120 from any convenient scale of equal parts, and with A as a center at the distance of 112.6 from the same scale describe an arc; lastly cut this arc with 112 described from the center B.

$$\text{By a line of Chords, or Goniometer } \left\{ \begin{array}{l} \angle A = 57^\circ .. 28' \\ \angle B = 57^\circ .. 57' \\ \angle C = 64^\circ .. 35' \end{array} \right.$$

2. Arithmetically.

$$\begin{array}{rcl} \text{As } 120 \text{ AB} & - & \text{Log. ar. co. } -3.9208188 \\ : 224.6 \text{ AC} + \text{BC} & - & \text{Log. } 2.3514098 \\ \therefore \quad .6 \text{ AC} - \text{BC} & - & \text{Log. } -1.7781513 \\ & & \hline : 1.123 \text{ diff. of the segments of AB} & - & \text{Log. } 0.0503799 \end{array}$$

Wherefore, by $\frac{1}{2}$ Sum and $\frac{1}{2}$ diff. $Ap = 60.566$, and $Bp = 59.434$.

And

$$\text{As } 112.6 \text{ AC} : 60.566 \text{ Ap} :: \text{Sin. } 90^\circ \angle \text{ApC} : \text{Cos. } 57^\circ .. 28' \angle A.$$

Also

$$\begin{array}{l} \text{As } 112 \text{ BC} : 59.434 \text{ Bp} :: \text{Sin. } 90^\circ \angle \text{BpC} : \text{Cos. } 57^\circ .. 57' \angle B. \\ \text{Hence, by subtraction, } \angle C = 64^\circ .. 35'. \end{array}$$

3. Instrumentally.

The extent on the line of numbers from 120 to 224.6 will reach from any multiple of .6 to the same multiple of $1\frac{1}{3}$ nearly, the diff. of Ap and Bp.

Hence $Ap = 60\frac{2}{16}$, and $Bp = 59\frac{7}{16}$.

Again the extent from 112.6 to $60\frac{2}{16}$ on the line of numbers will reach from 90° to $32^\circ .. 32'$ on the line of sines; wherefore $\angle A = 57^\circ .. 28'$ the complement of the 4th term. In like manner,

The extent from 112 to $59\frac{7}{16}$ on the line of numbers will reach on the line of sines from 90° to $32^\circ .. 3'$, of which the complement is $57^\circ .. 57' = \angle B$; and (by Eucl. i. 32.) $\angle C = 64 .. 35'$ which might be proved by a third proportion.


(Key to Vol. II. page 16.)

THEOREM IV.

1. Geometrically.

Ex. 2.

Take AB 180 from any scale of equal parts and at A make the angle BAC $62^\circ .. 40'$, lastly from B raise the perpendicular BC meeting AC in c .



$$\left. \begin{array}{l} AC=392 \\ EC=348.2 \end{array} \right\} \text{ on the scale employed for } AB.$$

2. Arithmetically.

As Radius AB - - - - -	Log. ar. co. - 10.	
: Tang. $62^\circ .. 40'$ BC - - - - -	Log. 10.2866141	
:: 180 AB - - - - -	Log. 2.2552725	
: 348.2464 BC - - - - -	Log. 2.5418866	

And

As Radius AB - - - - -	Log. ar. co. - 10.	
: Secant $62^\circ .. 40'$ AC - - - - -	Log. 10.3380298	
:: 180 AB - - - - -	Log. 2.2552725	
: 392.0146 AC - - - - -	Log. 2.5933023	

3. Instrumentally.

The extent from 45° to $62^\circ .. 40'$ on the line of tangents, reaches from 180 to $348\frac{1}{2}$ on the line of numbers, for the length of BC .

And the extent from $62^\circ .. 40'$ to 90° on the line of sines, reaches from $348\frac{1}{2}$ to 392 on the line of numbers for the length of AC .


(Key to Vol. II. page 22.)

MENSURATION OF HEIGHTS AND DISTANCES.

* * * In the subsequent Examples it has been thought unnecessary to detail the Construction, except in a few intricate cases.

Ex. 6

Let the $\angle c$, in the accompanying figure, be the angle of inclination of the ladder to the horizon, and the fort as there represented, the ditch extending or supposed to extend from the foot of the ladder to the base of the perpendicular tower. It is



As Radius - - - - - Log. ar. co. - 10
 : Tang. $62^\circ .. 40'$ $\angle c$ - - - - - Log. 10.286614
 :: 36 feet, the breadth of the ditch, - Log. 1.556303

 : 69.649 feet, the height of the wall. - Log. 1.842917

And,

As Sin. $62^\circ .. 40'$ $\angle c$ - - - Log. ar. co. - 10.0514158
 : Sin. 90° \angle subtended by the ladder, Log. 10. -----
 :: 69.649 feet, the height of the wall, Log. 1.8429169


 : 78.403 feet, the length of the ladder. Log. 1.8943327

Or, instead of the last proportion,

As Rad. : Sec. $62^\circ .. 40'$ $\angle c$: : 36, breadth of the ditch : 78.403, length of the ladder.

Ex. 7.

Let b in the annexed diagram be the foot of the shoar, and the building as there represented. It will be



(Key to Vol. II. page 22.)

As 11 feet, the extent of strut, Log. ar. co. -2.9586073
 : 23.83 feet, the height of the jamb, Log. 1.3771790
 :: Radius - - - - - Log. $10.-----$
 : *Tang.* 65° .. $13' \angle B$ - - - - - Log. 10.3357863

And,

As Sin. 65° .. $13' \angle B$ - - - Log. ar. co. -10.0419622
 : Sin. $90^\circ \angle$ subtended by the shoar, Log. $10.-----$
 :: 23.83 feet, the height of the jamb, Log. 1.3771790
 : 26.251 feet, the length of the shoar. Log. 1.4191412

This Question is easily solved by Common Arithmetic, on the Principles deduced from Eucl. i. 47.

Ex. 8. Here are given two right angled triangles, the perpendicular of the one being 33 feet, of the other 21 feet, and the hypotenuse of either, 40 feet, to find the *sum* of the bases.

Now

$\sqrt{40^2 - 33^2} = \sqrt{1600 - 1089} = 22.605$ feet, the base of the 1st triangle.

And

$\sqrt{40^2 - 21^2} = \sqrt{1600 - 441} = 34.044$ feet, the base of the 2d triangle.

But

$\overset{\text{ft.}}{22.605} + \overset{\text{ft.}}{34.044} = 56.649$ feet, the *required breadth of the street*.

This solution being much shorter than by Sines and Tangents, it would be superfluous to add the Trigonometrical proportions.

Ex. 9. Here are given the base (15 feet) and hypotenuse (39 feet) of a right angled triangle, to find the *sum* of the hypotenuse and perpendicular. Now

$\sqrt{39^2 - 15^2} = \sqrt{1521 - 225} = 36$ feet, left standing.
 Add 39 feet, the part broken off.

Sum 75 feet, the whole height.

(Key to Vol. II. page 22.)

Trigonometrically.


As $\frac{\text{ft.}}{39} : \frac{\text{ft.}}{15} :: \text{Rad.} : \text{Sin. } 22^\circ .. 37'$ the vertical \angle , and

As $\text{Tang. } 22^\circ .. 37' : \text{Rad.} :: \frac{\text{ft.}}{15} : \frac{\text{ft.}}{36}$ left standing.

Hence, as before, the whole height 75 feet.

Ex. 10.

Let A in the marginal figure be the observed angle $52^\circ .. 30'$, and the tower as there represented. It is




As Radius - - - - - Log. ar. co. - 10. _____
 : Tang. $52^\circ .. 30'$ $\angle A$ - - - - - Log. 10.1150195
 :: 170 feet, the distance of A from the tower, Log. 2.2304489
 : 221.54 feet, the altitude required. - - - - - Log. 2.3454684

Otherwise.

As Cos. $52^\circ .. 30'$ $\angle A$: Sin. $52^\circ .. 32'$ $\angle A$:: 170 feet : 221.54 feet,
 as before.

Ex. 11.

Let H in the annexed figure be horizontal with the top of the tower, and the ship* as there represented. The height of the tower subtends at the vessel an angle of 35° . [*Eucl. i. 29.*]



Therefore,


As Sin. 35° the complement of the \angle } Log. ar. co. - 10.2414087
 of depression, }
 : Cos. 35° (the same angle), - - - - - Log. 9.9133645
 :: 143 feet, the height of the tower, - - - - - Log. 2.1553360
 : 204.22 feet, the distance required. - - - - - Log. 2.3101092

* The Ship is purposely diminished in the diagram to a Model at anchor.

(Key to Vol. II. page 23.)

Ex. 14.

Let A and D be the two stations, E the top of the hill, and c the top of the tower. It will be




$$\begin{array}{l}
 \text{As Sin. } 17^{\circ} .. 15' \angle ADC \text{ --- Log. ar. co. --- } 10.5279144 \\
 : \text{ Sin. } 33^{\circ} .. 45' \angle D \text{ --- --- --- Log. } 9.7447390 \\
 :: 200 \text{ feet AD --- --- --- Log. } 2.3010300 \\
 \hline
 : 374.7 \text{ feet AC --- --- --- Log. } 2.5736834
 \end{array}$$

And

$$\begin{array}{l}
 \text{As Sin. } 130^{\circ} \angle AEC \text{ --- --- --- Log. ar. co. --- } 10.1157460 \\
 : \text{ Sin. } 11^{\circ} \angle EAC \text{ --- --- --- Log. } 9.2805988 \\
 :: 374.7 \text{ feet AC --- --- --- Log. } 2.5736834 \\
 \hline
 : 93.33148 \text{ feet, the height required. --- Log. } 1.9700282
 \end{array}$$

Ex. 15.

If A and B represent the two windows at which the observations were made, and c the top of the steeple, it will be,



$$\begin{array}{l}
 \text{As Sin. } 2^{\circ} .. 30' \angle ACB \text{ --- --- --- Log. ar. co. --- } 9.3603204 \\
 : \text{ Sin. } 127^{\circ} .. 30' \angle CBA \text{ --- --- --- Log. } 9.8994667 \\
 :: 18 \text{ feet AB --- --- --- Log. } 1.2552725 \\
 \hline
 : 327.38 \text{ feet AC --- --- --- Log. } 2.5150596
 \end{array}$$

(Key to Vol. II. page 23.)


And,

$$\begin{array}{l}
 \text{As Sin. } 90^\circ \angle CDA \text{ - - - - - Log. ar. co. - } 10 \cdot \text{-----} \\
 : \text{Sin. } 40^\circ \angle DAC \text{ - - - - - Log. } 9 \cdot 808067 \\
 :: 327 \cdot 38 \text{ feet AC - - - - - Log. } 2 \cdot 515060 \\
 \\
 : 210 \cdot 44 \text{ feet, the height required. - - - Log. } 2 \cdot 323127
 \end{array}$$

Lastly,

$$\begin{array}{l}
 \text{As Sin. } 90^\circ \angle CDA \text{ - - - - - Log. ar. co. - } 10 \cdot \text{-----} \\
 : \text{Sin. } 50^\circ \angle ACD \text{ - - - - - Log. } 9 \cdot 884254 \\
 :: 327 \cdot 38 \text{ feet AC - - - - - Log. } 2 \cdot 515060 \\
 \\
 : 250 \cdot 79 \text{ feet, the distance sought. - - Log. } 2 \cdot 399314
 \end{array}$$

Ex. 16.



Let A and B represent the two stations; and CD the object observed. The $\angle BDA$ [Eucl. i. 29.] is 27° ; the $\angle ABD$ (by comparison) 15° ; the $\angle BAD$ [Eucl. i. 32.] is 138° ; the $\angle CBD$ (by comparison) 8° ; and the $\angle DCB$ (by subtraction, and Eucl. i. 32.) is 109° .

Wherefore,

$$\begin{array}{l}
 \text{As Sin. } 27^\circ \angle BDA \text{ - - - - - Log. ar. co. - } 10 \cdot 342953 \\
 : \text{Sin. } 15^\circ \angle ABD \text{ - - - - - Log. } 9 \cdot 412996 \\
 :: 264 \text{ feet AB - - - - - Log. } 2 \cdot 421604 \\
 \\
 : 150 \cdot 5 \text{ feet AD, distance required. - - Log. } 2 \cdot 177553
 \end{array}$$


And

$$\begin{array}{l}
 \text{As Sin. } 27^\circ \angle BDA \text{ - - - - - Log. ar. co. - } 10 \cdot 342953 \\
 : \text{Sin. } 138^\circ \angle BAD \text{ - - - - - Log. } 9 \cdot 825511 \\
 :: 264 \text{ feet AB - - - - - Log. } 2 \cdot 421604 \\
 \\
 : 389 \cdot 1 \text{ feet BD - - - - - Log. } 2 \cdot 590068
 \end{array}$$

(Key to Vol. II. page 23.)

To render this more plain,

Let MN be the true horizon, KOD the boundary of the visual horizon to E any altitude whatever above F the surface of the Earth, of which the center is c; and let IH be a tangent to the circle at F, parallel to MN, and cutting the tangents ED and EK in the points G and L; $EG + GF = ED$. For draw CH through the point D, and CI through the point K; also join KF. The three triangles CFH, CDE, and CFI are in all respects equal. But the triangles EFG, HDG, are similar to the triangle EDC, consequently GF is equal to GD, and $EG + GF = ED$. Or, because CKE, CFI are both right angles, and that CK is equal to CF, the angles LKF, LFK are equal, and [Eucl. i. 6.] KL equal to LF; hence $EL + LF = EK = ED$. Therefore,



As Rad. EF	-----	Log. ar. co.	- 10. -----
: 2.5 miles, (<i>altitude given</i>)	-----	Log.	0.3979400
:: Tang. 87° .. 58' ∠ CED	-----	Log.	11.4497317

: 70.416 miles FG	-----	Log.	1.8476717

:: Secant 87° .. 58' ∠ CED	-----	Log.	11.4500052

: 70.46 miles EG	-----	Log.	1.8479452

miles miles miles
But $70.416 + 70.46 = 140.876 = ED$ as before.

Whence, also, 7936 miles, the diameter of the Earth.

(Page 24.)

Ex. 18. Here are given the three angles of a plane triangle, and one side, to find the other two sides. Now if the fort be denoted by F, and the two ships of war respectively by A and B, it is


(Key to Vol. II. page 24.)

As Sin. $11^\circ \angle AFB$ - - - - -	Log. ar. co. - 10.719401
: 440 yards AB - - - - -	Log. 2.643453
:: Sin. $83^\circ .. 45' \angle ABF$ - - - - -	Log. 9.997411
<hr/>	
: 2292.26 yards AF - - - - -	Log. 3.360265
<hr/>	
:: Sin. $85^\circ .. 15' \angle BAF$ - - - - -	Log. 9.998506
: 2298.05 yards BF - - - - -	Log. 3.361360

Therefore $\left\{ \begin{array}{l} 2292.26 \text{ yards, distance of the 1st ship} \\ 2298.05 \text{ yards, distance of the 2d ship} \end{array} \right\}$ Ans.

Ex. 19.

Let A and B be the two stations, and c the house. By the question the $\angle ABC$ is $68^\circ .. 2'$, the $\angle BAC$ $73^\circ .. 15'$, consequently the $\angle ACB$ $38^\circ .. 43'$; also AB is 400 yards. But,




As Sin. $38^\circ .. 43' \angle ACB$ - - -	Log. ar. co. - 10.203794
: 400 yards AB - - - - -	Log. 2.602060
:: Sin. $68^\circ .. 2' \angle ABC$ - - - - -	Log. 9.967268
<hr/>	
: 593.09 yards AC - - - - -	Log. 2.773122
<hr/>	
:: Sin. $73^\circ .. 15' \angle BAC$ - - - - -	Log. 9.981171
: 612.38 yards BC - - - - -	Log. 2.787025

Wherefore $\left\{ \begin{array}{l} 593.09 \text{ yards, distance of A} \\ 612.38 \text{ yards, distance of B} \end{array} \right\}$ Ans.

Ex. 20.

Let c be the tree observed, A and B the two stations of observation, and CD the breadth required. It is



(Key to Vol. II, page 24.)


As Sin. $47^\circ .. 48'$	$\angle ACB$	- - -	Log. ar. co. -	10·130296
: Sin. $79^\circ .. 12'$	$\angle BAC$	- - - - -	Log.	9·992239
:: 500 yards AB	- - - - -	- - - - -	Log.	2·698970
				<hr/>
: 662·99 yards BC	- - - - -	- - - - -	Log.	2·821505

And

As Sin. 90°	$\angle CDB$	- - - - -	Log. ar. co. -	10·————
: Sin. 53°	$\angle CBD$	- - - - -	Log.	9·902349
:: 662·99 yards BC	- - - - -	- - - - -	Log.	2·821505
				<hr/>
: 529·48 yards CD,	<i>breadth required</i>	-	Log.	2·723854

Ex. 21.

If A and B represent the two headlands, and C the point inland, there are given AC 735 yards, BC 840 yards, and the $\angle C$ $55^\circ .. 40'$; wherefore it will be



As 1575 yards AC+BC	- - - -	Log. ar. co. -	4·802719
: 105 yards BC-AC	- - - - -	Log.	2·021189
:: Tang. $62^\circ .. 10'$ half sup.	$\angle ACB$	- -	Log. 10·277379
			<hr/>
: Tang. $7^\circ .. 11'$ half diff.	\angle^{ies} A and B.	Log.	9·101287

Hence

The \angle at A = $69^\circ .. 21'$ and the \angle at B = $54^\circ .. 59'$.


Again

As Sin. $69^\circ .. 21'$	$\angle A$	- - - -	Log. ar. co. -	10·028839
: Sin. $55^\circ .. 40'$	$\angle C$	- - - - -	Log.	9·916859
:: 840 yards BC	- - - - -	- - - - -	Log.	2·924279
				<hr/>
: 741·2 yards AB,	<i>distance required</i>	-	Log.	2·869977

(Key to Vol. II. page 24.)

Ex. 22.

Let *c* be the point of land; *A* the former and *B* the latter place of observation. In the triangle *ABC* are given




the angle *A* $56^{\circ}..15'$, the angle *B* $101^{\circ}..15'$, and the side *AB* 12 miles; consequently the angle *c* is $22^{\circ}..30'$. But

As Sin. $22^{\circ}..30'$ $\angle C$ - - - -	Log. ar. co. -	10.417160
: Sin. $56^{\circ}..15'$ $\angle A$ - - - -	Log.	9.919846
:: 12 miles <i>AB</i> - - - -	Log.	1.079181
: 26.0728 miles <i>BC</i> , distance required	Log.	1.416187

(Page 25.)

Ex. 23.

If *AB* in the accompanying diagram be the base line, *C* the house, and *D* the mill, there are given the base line *AB* 600 yards, the $\angle DAB$ $95^{\circ} 20'$, the $\angle CAB$ $58^{\circ}..20'$, the $\angle ABC$ $98^{\circ}..45'$, and the $\angle DBA$ $53^{\circ}..30'$; hence the $\angle DAC$ is 37° , the $\angle ACB$ $22^{\circ}..55'$, and the $\angle ADB$ $31^{\circ}..10'$. But



As Sin. $22^{\circ}..55'$ $\angle ACB$ - - -	Log. ar. co. -	10.409613
: Sin. $98^{\circ}..45'$ $\angle ABC$ - - - -	Log.	9.994916
:: 600 yards <i>AB</i> - - - -	Log.	2.778151
: 1522.92 yards <i>AC</i> - - - -	Log.	3.182680

(Key to Vol. II. page 25.)

And

$$\begin{aligned} \text{As Sin. } 31^\circ \text{ .. } 10' \angle_{ADB} & \text{ - - Log. ar. co. - } 10.286065 \\ : \text{ Sin. } 53^\circ \text{ .. } 30' \angle_{ABD} & \text{ - - - - - Log. } 9.905178 \\ :: 600 \text{ yards AB} & \text{ - - - - - Log. } 2.778151 \\ & \text{-----} \\ : 931.96 \text{ yards AD} & \text{ - - - - - Log. } 2.969394 \end{aligned}$$

Also

$$\begin{aligned} \text{As } 2454.88 \text{ yards AC} + \text{AD} & \text{ - - Log. ar. co. - } 4.6189697 \\ : 590.96 \text{ yards AC} - \text{AD} & \text{ - - - - - Log. } 2.7715581 \\ :: \text{Tang. } 71^\circ \text{ .. } 30' \text{ half sup. } \angle_{DAC} & \text{ - Log. } 10.4754801 \\ : \text{Tang. } 36^\circ \text{ .. } 18' \text{ half the diff. of} & \text{-----} \\ \text{the } \angle_{les} \text{ ADC and ACD} & \left. \begin{array}{l} \\ \end{array} \right\} \text{ Log. } 9.8660079 \end{aligned}$$

Wherefore

The $\angle_{ADC} = 107^\circ \text{ .. } 48'$ and the $\angle_{ACD} = 35^\circ \text{ .. } 12'$.


Lastly,

$$\begin{aligned} \text{As Sin. } 107^\circ \text{ .. } 48' \angle_{ADC} & \text{ - - Log. ar. co. - } 10.0213040 \\ : \text{ Sin. } 37^\circ \text{ .. } 0' \angle_{DAC} & \text{ - - - - - Log. } 9.7794630 \\ :: 1522.92 \text{ yards AC} & \text{ - - - - - Log. } 3.1826771 \\ & \text{-----} \\ : 962.59 \text{ yards DC, distance required.} & \text{* Log. } 2.9834441 \end{aligned}$$

* Differing about 2 yards from the Answer given with the Question.

Ex. 24.

Conceiving the figure constructed as in the marginal diagram, the question gives AB 500 yards; AC and BD each 100 yards; AD 550 yards; and BC 560 yards. Hence it is,



$$\begin{array}{cccc} \text{yds.} & \text{yds.} & \text{yds.} & \text{yds.} \\ \text{As } 560 \text{ BC} : 600 \text{ AB} + \text{AC} & :: & 400 \text{ AB} - \text{AC} : 428.57 \text{ the diff. of the seg-} \\ & & & \text{ments of BC by a perpendicular from A. And, by } \frac{1}{2} \text{ sum and } \frac{1}{2} \\ & & & \text{diff. EB} = 494.28 \text{ yards, and EC} = 65.72 \text{ yards. Now} \end{array}$$

(Key to Vol. II. page 25.)

$$\text{As } \overset{\text{yds.}}{500} \text{ AB} : \overset{\text{yds.}}{494.28} \text{ EB} :: \text{Rad.} : \text{Sin. } 81^\circ .. 20' \angle \text{EAB.}$$

And

$$\text{As } \overset{\text{yds.}}{100} \text{ AC} : \overset{\text{yds.}}{65.72} \text{ EC} :: \text{Rad.} : \text{Sin. } 41^\circ .. 5' \angle \text{CAE.}$$

Again,

$$\text{As } \overset{\text{yds.}}{550} \text{ AD} : \overset{\text{yds.}}{600} \text{ AB} + \text{BD} :: \overset{\text{yds.}}{400} \text{ AB} - \text{BD} : \overset{\text{yds.}}{436.36} \text{ the diff. of the segments of AD by a perpendicular from B. Wherefore, by } \frac{1}{2} \text{ sum and } \frac{1}{2} \text{ diff. } \text{AF} = 493.18, \text{ and } \text{FD} = 56.82. \text{ And}$$

$$\text{As } \overset{\text{yds.}}{100} \text{ BD} : \overset{\text{yds.}}{56.82} \text{ FD} :: \text{Rad.} : \text{Sin. } 34^\circ .. 37' \angle \text{FBD. Also}$$

$$\text{As } \overset{\text{yds.}}{500} \text{ AB} : \overset{\text{yds.}}{493.18} \text{ AF} :: \text{Rad.} : \text{Sin. } 80^\circ .. 28' \angle \text{ABF. Consequently the } \angle \text{AOB} = 57^\circ .. 30', \text{ the } \angle \text{OAB} = 57^\circ .. 35', \text{ and the } \angle \text{OBA} = 64^\circ .. 55'.$$

But


$$\begin{aligned} \text{As Sin. } 57^\circ .. 30' \angle \text{AOB} : \overset{\text{yds.}}{500} \text{ AB.} \\ :: \text{Sin. } 57^\circ .. 35' \angle \text{OAB} : 500.47 \text{ OB one of the distances required.} \\ :: \text{Sin. } 64^\circ .. 55' \angle \text{OBA} : 536.81 \text{ nearly, AO the other distance} \\ \text{sought. Ans.*} \end{aligned}$$

* Differing in the decimal only from the Answer given with the Question.

Ex. 25.

1. Geometrically.

Construct a triangle, as ABC, having its three sides respectively equal to the three given straight lines, viz. AB $266\frac{1}{4}$ yds. AC 530 yds. and BC $327\frac{1}{2}$ yds. At the point A make the angle CAD equal to the observed angle BSC $29^\circ .. 50'$, and at the point c make the angle ACD equal to the observed $\text{ASB } 13^\circ .. 30'$. Then about the triangle ACD describe a circle, as CSAD, (Eucl. iv. 5.) and join DB. If DB be produced, it will meet the circumference in s, the station of the observer. But, on the scale employed for AB,



(Key to Vol. II, page 25.)

$$\left. \begin{array}{l} SA \\ SB \\ SC \end{array} \right\} \text{measures } \left\{ \begin{array}{l} 757.12 \text{ yds.} \\ 537 \text{ yds.} \\ 655.3 \text{ yds.} \end{array} \right\} \text{Ans.}$$

2. *Trigonometrically.*

The figure being as already constructed, from B demit the perpendicular BP upon AC. It is

$$\begin{array}{l} \text{As Sin. } 136^\circ .. 40' \angle ADC - - \text{Log. ar. co.} - 10.1635229 \\ \quad : 530 \text{ yards AC} - - - - - \text{Log.} \quad 2.7242759 \\ \therefore \text{Sin. } 13^\circ .. 30' \angle ACD - - - - - \text{Log.} \quad 9.3681853 \\ \\ \quad : 180.29 \text{ yards AD} - - - - - \text{Log.} \quad 2.2559841 \\ \\ \therefore \text{Sin. } 29^\circ .. 50' \angle DAC - - - - - \text{Log.} \quad 9.6967745 \\ \\ \quad : 384.21 \text{ yards DC} - - - - - \text{Log.} \quad 2.5845733 \end{array}$$

Likewise,

$$\begin{array}{l} \text{As } 530 \text{ yards AC} - - - - - \text{Log. ar. co.} - 3.2757241 \\ \quad : 593.75 \text{ yards BC} + AB - - - - - \text{Log.} \quad 2.7736036 \\ \therefore 61.25 \text{ yards BC} - AB - - - - - \text{Log.} \quad 1.7871061 \\ \\ \quad : 68.617 \text{ yards diff. of AP and PC} - - \text{Log.} \quad 1.8364338 \end{array}$$

Hence, by $\frac{1}{2}$ sum and $\frac{1}{2}$ diff. AP=230.692 yards, and CP=299.308 yds. Wherefore [by Theor. I. Plane Trigonometry] the $\angle ABP=60^\circ .. 2'$, the $\angle BAC=29^\circ .. 58'$, the $\angle CBP=66^\circ .. 3'$, and the $\angle BCA=23^\circ .. 57'$. Also by addition the $\angle BCD=71^\circ .. 16'$.

Again,

$$\begin{array}{l} \text{As } 711.71 \text{ yards CD} + CB - - \text{Log. ar. co.} - 3.1476969 \\ \quad : 56.71 \text{ yards CD} - CB - - - - - \text{Log.} \quad 1.7536596 \\ \therefore \text{Tang. } 71^\circ .. 16' \text{ half sup. } \angle BCD - \text{Log.} \quad 10.4696339 \\ \\ \quad : \text{Tang. } 13^\circ .. 13' \text{ half the diff. of} \\ \quad \quad \text{the } \angle^{\text{les}} \text{ CDB, CBD.} \} - \text{Log.} \quad 9.3709904 \end{array}$$

Consequently the $\angle DBC=84^\circ .. 29'$, and the $\angle CDB=58^\circ .. 3'$. But $\angle CDA 136^\circ .. 40' - \angle CDB 58^\circ .. 3' = \angle BDA 78^\circ .. 37'$.

And

$$\begin{array}{l|l} \text{As Sin. } 13^\circ .. 30' \angle ASD & \text{As Sin. } 29^\circ .. 50' \angle CSD \\ \quad : \text{Sin. } 78^\circ .. 37' \angle SDA & \quad : \text{Sin. } 58^\circ .. 3' \angle CDS \\ \therefore 180.29 \text{ yards AD} & \therefore 384.21 \text{ yards CD} \\ \quad : 757.14 \text{ yards SA.} & \quad : 655.30 \text{ yards es.} \end{array}$$

(Key to Vol. II. page 25.)

Also

$$\begin{aligned} \text{As } 266.25 \text{ yards } AB \\ : 757.14 \text{ yards } SA \\ :: \text{Sin. } 13^\circ .. 30' \angle BSA \\ : \text{Sin. } 138^\circ .. 25' \angle ABS. \end{aligned}$$


$$\begin{aligned} \text{As Sin. } 138^\circ .. 25' \angle ABS. \\ : \text{Sin. } 28^\circ .. 5' \angle BAS \\ :: 757.14 \text{ yards } SA \\ : 537.10 \text{ yards } SB. \end{aligned}$$

$$\text{Wherefore } \left\{ \begin{array}{l} SA \ 757.14 \\ SB \ 537.1 \\ SC \ 655.3 \end{array} \right\} \text{ yds. Ans.}$$

Ex. 26.

1. Geometrically.

Construct a triangle, as ABC, having its three sides respectively equal to the three given straight lines, viz. AB 600 yards, AC 800 yards, and BC 400 yards. At the point A make the angle CAE equal to the observed angle BSC $22^\circ .. 30'$, and at the point C make the angle ACE equal to the observed angle ASB $33^\circ .. 45'$. Then about the triangle AEC describe (*Eucl.* iv. 5.) a circle, as AECS, and join BE. If BE be produced, it will meet the opposite circumference in s, the station of the observer.



But, on the scale employed for AB.

$$\left. \begin{array}{l} SA \\ SB \\ SC \end{array} \right\} \text{ measures } \left\{ \begin{array}{l} 710\frac{1}{3} \text{ yds.} \\ 1042 \text{ yds.} \\ 934\frac{1}{6} \text{ yds.} \end{array} \right\} \text{ Ans.}$$

2. Trigonometrically.

The figure being as already constructed, from B demit the perpendicular BP upon AC. The angle ASC = $56^\circ .. 15'$ the sum of the observed angles : and the angle AEC = $123^\circ .. 45'$ the supplement of the $\angle ASC$ [*Theor.* 54. *Geometry*]. Hence it is

(Key to Vol. II. page 25.)

Now

$$\begin{array}{l}
 \text{As Sin. } 33^\circ 45' : \text{Sin. } 41^\circ .. 7\frac{1}{2} :: 600_{AB} : 710.3_{SA} \\
 \text{Sin. } 22^\circ 30' : \text{Sin. } 63^\circ .. 21\frac{1}{2} :: 400_{BC} : 934.14_{SC} \\
 \text{Sin. } 22^\circ .. 30' : \text{Sin. } 94^\circ .. 8\frac{1}{2} :: 400_{BC} : 1041.85_{SB}
 \end{array}
 \left. \vphantom{\begin{array}{l} \text{As Sin. } 33^\circ 45' : \text{Sin. } 41^\circ .. 7\frac{1}{2} \\ \text{Sin. } 22^\circ 30' : \text{Sin. } 63^\circ .. 21\frac{1}{2} \\ \text{Sin. } 22^\circ .. 30' : \text{Sin. } 94^\circ .. 8\frac{1}{2} \end{array}} \right\} \text{Ans.}^*$$

* By which it should appear that the Answer given with the Question is correct only in the proximity of the correctness of *Geometrical Construction* to that of *Trigonometrical Calculation*.

MENSURATION OF PLANES.

(Page 27.)

1. The Area of Parallelograms.

$$\text{Ex. 2. } \frac{\text{Chains } 35.25 \times \text{Chains } 35.25}{10} = 124.25625 = 124 \text{ Acres } \text{A. R. P. } 1 .. 1. \text{ Ans.}$$

$$\text{Ex. 3. } 12.5 \times .75 = 9.375 = 9\frac{3}{8} \text{ feet. Ans.}$$

$$\text{Ex. 4. } \frac{\text{Chains } 6.2 \times \text{Chains } 5.45}{10} = 3.379 = 3 \text{ Acres } \text{A. R. P. } 1 .. 20.64. \text{ Ans.}$$

$$\text{Ex. 5. } \frac{\text{ft. } 37 \times \text{ft. } 5\frac{1}{4}}{9} = 21\frac{7}{9} \text{ square yards. Ans.}$$

(Page 28.)

2. The Area of Triangles.

RULE 1.

$$\text{Ex. 2. } \frac{\text{ft. } 40 \times \text{ft. } 30}{2 \times 9} = 66\frac{2}{3} \text{ square yards. Ans.}$$

(Key to Vol. II, page 28.)

Ex. 3. $\frac{\text{ft. } 24 \cdot 5 \times \text{ft. } 25 \cdot 25}{9} = 68 \cdot 7361 = 68\frac{5}{7}\frac{3}{2}$ square yards. Ans.

Ex. 4. $(9 \text{ .. } 2) \times (11 \text{ .. } 10) = 108 \text{ .. } 5 \text{ .. } 8 = 108\frac{1}{3}\frac{7}{6}$ sqr. feet. Ans.*

* The Answer being 108 ft. 68 in. instead of 108 ft. 5 $\frac{1}{3}$ in. as given by Hutton, the student is referred to the 49th page of the KEY for more information.

(Page 29.)

RULE II.

Ex. 2. $\frac{\frac{1}{2} \times \text{ft. } 25 \times \text{ft. } 21 \cdot 25 \times \cdot 7071068^*}{9} = 20 \cdot 86947$ Ans.

Or,

As Rad. - - - - -	Log. ar. co. - 10. _____
: Sin. 45° the contained ∠, - - - -	Log. 9·8494850
: : 265·625 ft. half prod. of the sides,	Log. 2·4242639
: 187·82523 ft. = 20·86947 <i>sqr. yds.</i>	} Log. 2·2737489
<i>the area sought.</i>	

* Nat. Sin. 45°.

(Page 30.)

RULE III.

Ex. 2. $\frac{\text{ft. } 30 + \text{ft. } 40 + \text{ft. } 50}{2} = 60$ half Sum of the sides. And,

- ft. ft. ft.
- 60 - 30 = 30 First remainder.
- 60 - 40 = 20 Second remainder.
- 60 - 50 = 10 Third remainder.

But

$\sqrt{60 \times 30 \times 20 \times 10} = \sqrt{360000} = 600$ square feet, or 66 $\frac{2}{3}$ square yards. Ans.

(Key to Vol. II. page 30.)

Ex. 3. $\frac{\text{links } 2569 + \text{links } 4900 + \text{links } 5025}{2} = 6247$ half *Sum* of the sides. And

$\text{links } 6247 - \text{links } 2569 = 3678$ First remainder.
 $\text{links } 6247 - \text{links } 4900 = 1347$ Second remainder.
 $\text{links } 6247 - \text{links } 5025 = 1222$ Third remainder.

But

$\sqrt{\frac{6247 \times 3678 \times 1347 \times 1222}{100000}} = 61 \text{ AC. } 1 \text{ RO. } 39 \text{ PO. Ans.}$

3. *The Area of Trapezoids.*

Ex. 2. $\frac{\text{in. } 15 + \text{in. } 11}{2} \times 12.5 \text{ feet} = (1 \text{ ft. } 1 \text{ in.}) \times (12 \text{ ft. } 6 \text{ in.}) = 13 \frac{1}{4} \text{ sq. ft. Ans.}$

(Page 31.)

Ex. 3. $\frac{\text{links } 110_{AP} \times \text{links } 352_{CP}}{2} = 19360$ square links, the area of APC.

And

$\frac{\text{links } (352_{CP} + 595_{DQ}) \times \text{links } 635_{PQ}}{2} = 300672.5$ square links, the area of CPQD.

Again,

$\frac{\text{links } 365_{BQ} \times \text{links } 595_{DQ}}{2} = 108587.5$ sq. links, the area of DQB.

But

$\text{sq. links } (APC) \quad (CPQD) \quad (DQB) \quad \text{Acres}$
 $100000) 19360 + 300672.5 + 108587.5 (4.2862 \text{ whole area.}$

4

Roods 1.1448
40

Perches 5.7920


(Key to Vol. II. page 31.)

4. *The Area of Trapezia.*

$$\text{Ex. 2. } \frac{\overset{\text{ft.}}{28} + \overset{\text{ft.}}{33\frac{1}{2}}}{2} \times \overset{\text{ft.}}{65} = 222\frac{1}{2} \text{ square yards. Ans.}$$

Ex. 3. Construction.

From any convenient scale of equal parts take AC 378 yards, and by the same scale make AE 100 yards, and CF 70 yards. Then from E and F at right angles to AC draw ED, FB; and from the point A



yards on the scale employed for AC, determine D; also from the point C, with a radius of 265 yards on the same scale, determine B. Lastly, join AB and DC. ABCD is the field required.

Calculation.

$$\sqrt{\overset{\text{yds.}}{(220_{AD})^2} - \overset{\text{yds.}}{(100_{AE})^2}} = \overset{\text{yds.}}{196 \text{ nearly, DE.}}$$

And

$$\sqrt{\overset{\text{yds.}}{(265_{BC})^2} - \overset{\text{yds.}}{(75_{CF})^2}} = \overset{\text{yds.}}{255.5 \text{ BF.}}$$

$$\text{But } \frac{\overset{\text{yds.}}{196} + \overset{\text{yds.}}{255.5} \times \overset{\text{yds.}}{378_{AC}}}{2 \times 4840 \text{ sq. yds.}} = \overset{\text{A.}}{17} \dots \overset{\text{R.}}{2} \dots \overset{\text{P.}}{21}. \text{ Ans.}$$

(Page 32.)

5. *The Area of irregular Polygons.*

EXAMPLE. [See the Figure given with the Problem.]

AC	55	×	9	=	495	area of	ABC
AC	55	×	6½	=	357½	- - - -	AGC
Dg	23	×	22	=	506	- - - -	GDC
½FD	26	×	12	=	312	- - - -	FGD
½FD	26	×	8	=	208	- - - -	FDE

Sum 1878½ whole area. Ans.

(Key to Vol. II. page 32.)

Otherwise.

$$\frac{55_{AC} \times (18_{Bn} + 13_{Gm})}{2} = 852\frac{1}{2} \text{ ABCG.}$$

$$\frac{44_{GC} \times 23_{Dq}}{2} = 506 \text{ CGD.}$$

$$\frac{52_{FD} \times (12_{GO} + 8_{EP})}{2} = 520 \text{ FGDE.}$$

$$\text{Sum } \underline{1878\frac{1}{2}} \text{ Ans. as before.}$$

(Page 33.)

6. The Area of regular Polygons.

- Ex. 2. (Side)(Side) (Tab. area)
 $20 \times 20 \times 0.4330127 = 173.20508. \text{ Ans.}$
- Ex. 3. $20 \times 20 \times 2.5980762 = 1039.23048. \text{ Ans.}$
- Ex. 4. $20 \times 20 \times 4.8284271 = 1931.37084. \text{ Ans.}$
- Ex. 5. $20 \times 20 \times 7.6942088 = 3077.68352. \text{ Ans.}$

(Page 36.)

The Length of an Arc of a Circle.

- Ex. 1. (Const. dec.) (Rad.) feet
 $.01745 \times 30^\circ \times 9\text{ft.} = 4.7115. \text{ Ans.}$
- Ex. 2. (Const. dec.) (Rad.) feet
 $.01745 \times 12^\circ \frac{1}{2} \times 10\text{ft.} = 2.1231 \text{ length of the arc required.}$

(Page 37.)

The Areas of Circles.

- Ex. 2. (C.) (D.)
 $\frac{22 \times 7}{4} = 5\frac{1}{2} \times 7 = 38\frac{1}{2}. \text{ Ans.}$

Or,

- (Circular ar.) (D²)
 $.7854 \times 49 = 38.4846 \text{ area required, nearly as before.}$

(Key to Vol. II, page 37.)

$$\text{Ex. 3. } \begin{array}{l} \text{(Circul. a.)} \quad \text{(D}^2\text{.)} \quad \text{sq. feet} \quad \text{sq. yds.} \\ \cdot 7854 \times 12\frac{1}{4} = 9\cdot 62105 = 1\cdot 069. \quad \text{Ans.} \end{array}$$

$$\text{Ex. 4. } \begin{array}{l} \text{(Recip. dec.)} \quad \text{(C}^2\text{.)} \quad \text{sq. feet.} \\ \cdot 07958 \times 144 = 11\cdot 45952. \quad \text{Ans.} \end{array}$$

(Page 38.)

The Areas of Circular Rings.

$$\text{Ex. 2. } \begin{array}{l} 20 \times 20 = 400 \text{ the square of the greater diameter.} \\ 10 \times 10 = 100 \text{ the square of the less diameter.} \end{array}$$

$$\begin{array}{r} \text{300 difference of the squares.} \\ \text{Multiply by } \cdot 7854 \\ \hline \end{array}$$

$$\text{PRODUCT } 235\cdot 62 \text{ area required.}$$

Or,

$$\begin{array}{l} 20 + 10 = 30 \text{ sum of the diameters.} \\ 20 - 10 = 10 \text{ diff. of the diameters.} \end{array}$$

$$\begin{array}{r} \text{300 rectangle under the sum and diff.*} \\ \text{Multiply by } \cdot 7854 \\ \hline \end{array}$$

$$\text{PRODUCT } 235\cdot 62 \text{ area required, as before.}$$

* EUCLID, Book ii. Prop. 5.

(Page 39.)

The Areas of Sectors of Circles.

$$\text{Ex. 2. } \begin{array}{l} \text{(Rad.)} \quad \text{(Arc)} \\ \frac{10 \times 20}{2} = 100. \quad \text{Ans.} \end{array}$$

$$\text{Ex. 3. } \begin{array}{l} \text{(Rad.)} \\ 25 \times 2 = 50 \text{ the diameter of the circle.} \end{array}$$

$$\text{And } 50 \times 50 = 2500 \text{ the square of the diameter.}$$

$$\begin{array}{l} \text{(Circ. a.)} \quad \text{(D}^2\text{.)} \\ \text{Also } \cdot 7854 \times 2500 = 1963\cdot 5 \text{ area of the whole circle.} \end{array}$$

(Key to Vol. II. page 39.)

But

$$\begin{matrix} \text{(Circ.)} & \text{(Giv. arc)} & \text{(Whole area)} & \text{(Area reqd.)} \\ \text{As } 360^\circ & : 147^\circ \text{ .. } 29' & :: 1963\cdot5 & : 804\cdot3986. \end{matrix} \text{ Ans.}$$

(Page 40.)

The Areas of Segments of Circles.

Ex. 3. $\begin{matrix} \text{(D.)} & \text{(Alt.)} \\ 50 & 18 \end{matrix} \cdot 36$ Quotient.


Entering the *Table of Circular Segments* with $\cdot 36$, the corresponding area is $\cdot 25455$. Lastly,

$$\begin{matrix} \text{(Tab. ar.)} & \text{(D}^2\text{.)} \\ \cdot 25455 & \times 2500 = 636\cdot375. \end{matrix} \text{ Ans.}$$

Ex. 4.

Construction.

From any convenient scale of equal parts take EC 10, and about E at the distance EC describe the circle ABC. Place the right line AB 16 from the same scale, in the circle ABC, [*Eucl.* iv. 1.] and join AE, EB. Bisect AB in D, and through the point D draw EC. The line AB divides the circle into two segments, either of which is the segment whereof the area is required.



Calculation.

$$\begin{array}{r} \text{As } 10 \text{ AE} \text{ - - - - -} \text{ Log. ar. co. - } 1\cdot \text{ - - - - -} \\ \text{: } 8 \text{ AD} \text{ - - - - -} \text{ Log. } 0\cdot 9030900 \\ \text{: : Radius - - - - -} \text{ Log. } 10\cdot \text{ - - - - -} \\ \text{: Sin. } 53^\circ \text{ .. } 8' \angle \text{AEC, nearly - - - -} \text{ Log. } 9\cdot 9030900 \end{array}$$

Wherefore the arc ACB contains $106^\circ \text{ .. } 16'$ nearly.

Again,

$$\begin{matrix} \text{(Circ. a.)} & \text{(D}^2\text{.)} \\ \cdot 7854 & \times 400 = 314\cdot16 \end{matrix} \text{ the area of the whole circle.}$$

But

$$\begin{matrix} \text{(Circ.)} & \text{(Arc ACB)} & \text{(Whole area)} & \text{(Area ACBE)} \\ \text{As } 360^\circ & : 106^\circ \text{ .. } 16' & :: 314\cdot16 & : 92\cdot728. \end{matrix}$$

(Key to Vol. II. page 40.)

Now

$$\sqrt{100_{AE}^2 - 64_{AD}^2} = \sqrt{36} = 6 = ED. \text{ And}$$

$$\frac{16_{AB} \times 6_{ED}}{2} = \frac{96}{2} = 48 \text{ area of the triangle } AEB.$$

Hence, by subtraction, $44.728 =$ the area of the segment AEC .

Consequently,

269.432 and 44.728 are the *two* Answers.*

* Evincing a defect in the Answer accompanying the Question.

(Page 42.)

The Areas of irregular Figures.

$$\text{Ex. 2. } \frac{17.4 + 24.4}{2} = 20.9 \text{ mean breadth of the two ends.}$$

And

$$\frac{20.9 + 20.6 + 14.2 + 16.5 + 20.1}{5} \times 84 = 1550.64. \text{ Ans.}$$

The Areas of Ellipses.

(Tra. di.) (Conj. di.) (Const. dec.)

$$\text{Ex. 1. } 70 \times 50 \times .7854 = 2748.9 \text{ area required.}$$

(Conj. di.) (Tra. di.) (Const. dec.)

$$\text{Ex. 2. } 24 \times 28 \times .7854 = 339.2928. \text{ Ans.}$$

(Page 43.)

The Areas of Elliptic Segments.

(Alt.) (Gr. ax.)

$$\text{Ex. 2. } 10 \div 35 = 28\frac{2}{7} \text{ the versed sine; to which in the Table of Circular Segments corresponds } .18518$$

Multiply by $875 = 35 \times 25$, the axes.

$$\text{PRODUCT } \underline{\underline{162.0325}} \text{ area required.}$$

(Key to Vol. II. page 48.)

(Alt.) (Less ax.)
 Ex. 3. $5 \div 25 = .2$ the versed sine; with which, entering the
Table of Circular Segments, the corresponding area is
 found $.11182$
 Multiply by $875 = 35 \times 25$, the two axes.
 PRODUCT 97.8425 area sought.

The Areas of Parabolæ and Parabolic Segments.

(Base) (Alt.)
 Ex. 2. $\frac{2}{3} \times 16 \times 10 = 106\frac{2}{3}$. Ans.

MENSURATION OF SOLIDS.

(Page 44.)

The Superficies of Prisms and Cylinders.

(Side) (Side) (No.)
 Ex. 1. $20 \times 20 \times 6 = 2400$ the surface of the cube.

(Length) (Breadth) (Sides) ft.
 Ex. 2. $20 \text{ ft.} \times 1.5 \text{ ft.} \times 3 = 90$ area of the three sides.

And $\frac{1.5 + 1.5 + 1.5}{2} = 2.25$ half sum of the three sides of either
 triangle forming an end of the given prism.

(Half sum) (Side)
 Also $2.25 - 1.5 = .75$ one of the three equal remainders. But
 $\sqrt{.75 \times .75 \times .75 \times 2.25} = .97432$ area of either end of
 the prism.

Multiply by 2

PRODUCT 1.94864 area of both ends.
 Add 90 area of the sides.

Sum 91.94864 whole superficies.

(Di.) (Circumf.) (Di.) (Circumf.)
 Ex. 3. As $1 : 3.1416 :: 2 \text{ ft.} : 6.2832 \text{ ft.}$ of the given cylinder.

(Key to Vol. II. page 44.)

(Circumf.) (Length) sq. feet.
And $6\cdot2832 \text{ ft.} \times 20 \text{ ft.} = 125\cdot664$ convex surface required.

ft. ft. ft. ft. sq. feet
Ex. 4. $(2\cdot5 + 3\cdot166 + 2\cdot5) \times 2\cdot66 = 21\cdot77$ the area of the bottom
and two ends of the cistern. Also

ft. ft. sq. feet
 $(2 \times 3\cdot166) \times 2\cdot5 = 15\cdot8333$ area of the two sides.

Hence the whole internal area is $37\frac{1}{8}$ square feet.

sq. feet lb. lb. d.
But $37\frac{1}{8} \times 7 = 263\frac{5}{8}$, which at 3 per lb. = £3 .. 5 .. $9\frac{3}{4}$ nearly.

(Page 45.)

The Superficies of Pyramids and Cones.

Perim of the ba. (Sl. ht.)*
Ex. 1. $\frac{(3 \text{ ft} + 3 \text{ ft.} + 3 \text{ ft.}) \times 20 \text{ ft.}}{2} = 90$ square feet. Ans.

* The slant height of Pyramids and Pyramidal frusta is an ambiguous expression. Dr. Hutton means the line that in the plane of the perpendicular bisects a side.

(Di.) (Circumf.) (Di.) (Circumf.)
Ex. 2. As 1 : $3\cdot1416$:: 8·5 ft. : $26\cdot7036$ ft. round the base.

(Circumf.) (Sl. ht.)
And $\frac{26\cdot7036 \text{ ft.} \times 50 \text{ ft.}}{2} = 667\cdot59$ square feet. Ans.

The Surface of the Frusta of Pyramids and Cones by Sections parallel to the Base.

ft. in. ft. in. (Sides) (Sl. ht.)
Ex. 1. $\frac{(3 \text{ .. } 4) + (2 \text{ .. } 2)}{2} \times 4 \times 10 \text{ ft.} = 110$ sq. feet. Ans.

ft. ft. (Sl. ht.)
Ex. 2. $\frac{6 + 8\cdot4}{2} \times 12\cdot5 \text{ ft.} = 90$ square feet. Ans.

(Page 46.)

The Solidity of Prisms and Cylinders.

in. in. in. cubic in.
Ex. 1. $24 \times 24 \times 24 = 13824$. Ans.

(Key to Vol. II, page 46.)

$$\text{Ex. 2. } \begin{array}{ccc} \text{(Length)} & \text{(Breadth)} & \text{(Thickness)} \\ (3 \text{ ft. } 2 \text{ in.}) & \times (2 \text{ ft. } 8 \text{ in.}) & \times (2 \text{ ft. } 6 \text{ in.}) = 21\frac{1}{5} \text{ solid ft. } \text{Ans.} \end{array}$$

$$\text{Ex. 3. } \begin{array}{cc} \text{c. in.} & \text{cub. in.} \\ 282) 21\frac{1}{5} \times 1728 & (129\frac{17}{47} \text{ gallons of water. } \text{Ans.} \end{array}$$

Ex. 4. Because $3^2 + 4^2 = 5^2$ it follows [Eucl. i. 48] that the angle contained by the two sides, 3 and 4 respectively, is a right angle. Therefore

$$\frac{3 \times 4}{2} \times 10 \text{ feet} = 60 \text{ cubic feet, content required.}$$

$$\text{Ex. 5. } \begin{array}{cccc} \text{ft.} & \text{ft.} & \text{(Rec. dec.)} & \text{(Length)} & \text{sol. feet.} \\ 5.5 \times 5.5 \times .07958 \times 20 \text{ feet} & = & 48.1459. & \text{Ans.} \end{array}$$

(Page 47.)

The Solidity of Pyramids and Cones.

$$\text{Ex. 1. } \frac{\begin{array}{c} \text{(Base)} \\ 30 \times 30 \end{array}}{3} \times \begin{array}{c} \text{(Alt.)} \\ 25 \end{array} = 7500 \text{ content required.}$$

$$\text{Ex. 2. } \frac{\begin{array}{c} \text{(Sides)} \\ 3+3+3 \end{array}}{2} = \frac{9}{2} = 4.5 \text{ half sum of the sides.}$$

And $4.5 - 3 = 1.5$ one of the three equal remainders.

$$\text{But } \sqrt{\begin{array}{c} \text{(Alt.)} \\ 4.5 \times 1.5 \times 1.5 \times 1.5 \times 30 \div 3 \end{array}} = 3.897117 \times 10, \text{ or } 38.97117 \text{ the solidity required.}$$

$$\text{Ex. 3. } \frac{\begin{array}{c} \text{ft.} \quad \text{ft.} \quad \text{ft.} \\ 5+6+7 \end{array}}{2} = 9 \text{ ft. half sum of the three sides.}$$

And the three remainders are $\begin{array}{ccc} \text{ft.} & \text{ft.} & \text{ft.} \\ 2, & 3, & 4. \end{array}$

$$\text{But } \sqrt{9 \times 2 \times 3 \times 4 \text{ ft.}} = \sqrt{216 \text{ ft.}} = 14.6969385 \text{ ft. area of the base.}$$

$$\text{Lastly } \frac{\begin{array}{c} \text{ft.} \\ 14.6969385 \times 14.5 \text{ ft.} \end{array}}{3} \begin{array}{c} \text{(Alt.)} \\ \text{solid feet} \end{array} = 71.0352. \text{ Ans.}$$

(Key to Vol. II. page 47.)

Ex. 4. $\begin{matrix} \text{(Tab. area)} & \text{(Side sqd.)} & \text{sq. ft.} \\ 1.7204774 \times 4 \text{ ft.} & = & 6.8819096 \text{ area of the base.} \end{matrix}$
 Multiply by $\underline{4 \text{ ft.} = \frac{1}{3} \text{ of } 12 \text{ ft. altitude.}}$
 PRODUCT $\underline{27.5276384}$ solid feet. Ans.

Ex. 5. $\begin{matrix} \text{(Tab. area)} & \text{(Side sqd.)} & \text{square feet} \\ 2.5980762 \times .25 \text{ feet} & = & .64951905 \text{ area of the base.} \end{matrix}$
 And $\frac{\begin{matrix} \text{sq. feet} & \text{(Alt.)} \\ .64951905 \times 6.4 \text{ ft.} & \text{cubic feet} \end{matrix}}{3} = 1.38564064.$ Ans.

Ex. 6. $\begin{matrix} \text{(Recip. dec.)} & \text{(c}^2\text{)} & \text{sq. feet} \\ .07958 \times 81 & = & 6.44598 \text{ area of the base.} \end{matrix}$
 Multiply by $\underline{3.5 \text{ ft.} = \frac{1}{3} \text{ of } 10\frac{1}{2} \text{ ft. altitude.}}$
 PRODUCT $\underline{22.56093}$ solid feet. Ans.

(Page 48.)

The Solidity of the Frusta of Cones and Pyramids.

Ex. 1. $\begin{matrix} \text{ft.} & \text{ft.} & \text{sq. ft.} \\ .5 \times .5 & = & .25 \text{ area of the less end.} \end{matrix}$ And
 $\begin{matrix} 1.25 \times 1.25 & = & 1.5625 \text{ - - - - - greater end.} \\ & & \text{sq. ft.} \end{matrix}$

But $\sqrt{.25 \times 1.5625 \text{ ft.}} = .625$ mean proportional.

Again
 $\frac{\begin{matrix} \text{sq. ft.} & \text{sq. ft.} & \text{sq. ft.} & \text{sq. feet} \\ .25 + .625 + 1.5625 & = & 2.4375 & \text{sq. ft.} \end{matrix}}{3} = \frac{2.4375}{3} = .8125$ mean area.

Multiply by $\underline{24 \text{ altitude.}}$

PRODUCT $\underline{19.5}$ solid feet. Ans.

Ex. 2. $\begin{matrix} \text{ft.} & \text{ft.} & \text{ft.} & \text{ft.} & \text{sq. ft.} \\ (1.5)^2 + (1.5)^2 + (1.5 \times .5) & = & 2.5625 & \text{sum of the products.} \end{matrix}$

But $\frac{\begin{matrix} \text{(Tab. ar.)} & \text{(Sum of prods.)} & \text{(Alt.)} \\ 1.7204774 \text{ ft.} \times 2.5625 \text{ ft.} \times 5 \text{ ft.} & \text{cubic feet} \end{matrix}}{3} = 9.31925.$ Ans.

(Key to Vol. II. page 48.)

$$\text{Ex. 3. } \begin{matrix} (Gr. di^2.) & (L. di^2.) & (Gr. di) & (L. di.) \\ 64 + 16 + (8 \times 4) = 112 \end{matrix} \text{ sum of the products.}$$

$$\text{And } \begin{matrix} (Tab. ar.) & (Sum of pro.) & (Alt.) \\ .7854 \times 112 \times 18 \\ 3 \end{matrix} = 527.7888. \text{ Ans.}$$

$$\text{Ex. 4. } \begin{matrix} (Gr. di^2.) & (L. di^2.) & (Gr. di) & (L. di.) \\ 400 + 100 + (20 \times 10) = 700 \end{matrix} \text{ sum of the prods.}$$

$$\text{And } \begin{matrix} (Tab. ar.) & (Sum of pro.) & (Alt.) \\ .07958 \times 700 \times 25 \\ 3 \end{matrix} = 464.216 \text{ solidity required.}$$

$$\text{Ex. 5. } \begin{matrix} (B. di^2.) & (H. di^2.) & (B. di) & (H. di.) & \text{sq. in.} \\ 784 \text{ in.} + 400 \text{ in.} + (28 \text{ in.} \times 20 \text{ in.}) = 1744 \end{matrix} \text{ sum of the prods.}$$

$$\text{And } \begin{matrix} (Tab. ar.) & (Sum of prods.) & (Alt.) & \text{cub. inches} \\ .7854 \text{ in.} \times 1744 \text{ in.} \times 40 \\ 3 \end{matrix} = 18263.168 \text{ the content of}$$

the whole Cask. But

c. in. cub. inches

231) 18263.168 (79.06133 gallons of wine. Ans.

(Page 50.)

The Superficies of Spheres and Spherical Segments.

$$\text{Ex. 1. } \begin{matrix} (c.) & (D.) \\ 22 \times 7 = 154 \end{matrix} \text{ the superficies required.}$$

$$\text{Ex. 2. } \begin{matrix} (D^2.) & (Const. dec.) & \text{sq. inches.} \\ 24 \text{ in.} \times 24 \text{ in.} \times 3.1416 = 1809.5616 \end{matrix} \text{ superficies required.}$$

$$\text{Ex. 3. } \begin{matrix} (Di.) & (Cir.) & \text{square miles} \\ 7957.75 \text{ mls.} \times 25000 \text{ mls.} = 198943750. \end{matrix} \text{ Ans.}$$

$$\text{Ex. 4. } \begin{matrix} (Di.) & (Cir.) & (Di.) & (Cir.) \\ \text{As } 1 : 3.1416 :: 42 \text{ in.} & 131.9472 \text{ in.} \end{matrix} \text{ the circumference of the sphere of which the segment proposed is a part.}$$

$$\text{But } \begin{matrix} (Whole cir.) & (Alt.) & \text{square inches.} \\ 131.9472 \text{ in.} \times 9 \text{ in.} = 1187.5248. \end{matrix} \text{ Ans.}$$

$$\text{Ex. 5. } \begin{matrix} (Di.) & (Cir.) & (Di.) & (Cir.) \\ \text{As } 1 : 3.1416 :: 12.5 \text{ ft.} & 39.27 \text{ ft.} \end{matrix} \text{ the circumference of the sphere of which the zone proposed is a part.}$$

(Key to Vol. II. page 50.)

(Whole cir.) (Alt.) sq. ft.
 But $39.27 \text{ ft.} \times 2 \text{ ft.} = 78.54$ convex surface required.

(Page 51.)

The Solid Content of Spheres.

Ex. 1. $(Di^3) (Const. dec.)$
 $(12 \times 12 \times 12) \times .5236 = 904.7808$. Ans.

Ex. 2. $(Cir.^3) (Const. dec.)$ square miles
 $25000^3 \text{ mls.} \times .01688 \text{ mls.} = 263750000000$. Ans.

(Page 52.)

The Solidity of Spherical Segments.

Ex. 1. $(Di.) (Dbl. alt.) (Alt.)^2$ sq. ft.
 $(8 \text{ ft.} \times 3) - 4 \text{ ft.} = 20 \text{ ft.}$ which \times^{ed} by 4 ft. = 80
 Multiply by the constant decimal .5236 feet.
 PRODUCT cubic feet 41.888. Ans.

Ex. 2. Here 3 times the square of the radius of the base = 300
 The square of the height is 81

Sum	381
Multiply by the height	9
<hr/>	
PRODUCT	3429
Multiply by the const. decimal	.5236
<hr/>	
Ans.	1795.4244

ARTIFICERS-WORK.

(Page 83.)

BRICKLAYING.

Ex. 1. $\frac{\text{ft. in. ft. in. h. bricks}}{(57 \text{ .. } 3) \times (24 \text{ .. } 6) \times 5} = \frac{\text{sq. ft. pts. in. rods. yds.}}{2337 \text{ .. } 8 \text{ .. } 6} = 8 \text{ .. } 17\frac{2}{3}$. Ans.
 3 half bricks.

(Key to Vol. II. page 83.)

$$\text{Ex. 2. } \frac{\text{ft. in.} \quad \text{ft. in.} \quad \text{h. bricks}}{(62 \text{ .. } 6) \times (14 \text{ .. } 8) \times 5}{9 \times 3 \text{ half bricks}} = 169 \cdot 753 \text{ sq. yds. Ans.}$$

$$\text{Ex. 3. } \frac{\text{(Length)} \quad \text{(Height)} \quad \text{h. bricks}}{24 \cdot 75 \text{ ft.} \times 17 \cdot 5 \text{ ft.} \times 4}{2 \times 3 \text{ half bricks}} \quad \begin{array}{l} \text{sq. ft.} \quad \text{sq. yards} \\ \div 9 = 32 \cdot 083\bar{3}. \text{ Ans.} \end{array}$$

$$\text{Ex. 4. } \frac{\text{ft. in.} \quad \text{ft. h. br.}}{(28 \text{ .. } 10) \times 20 \times 5}{3 \text{ half bricks}} \quad \begin{array}{l} \text{sq. ft. 12ths} \\ = 961 \text{ .. } 1\frac{1}{3} \text{ the first 20 feet up.} \end{array}$$

$$\frac{\text{ft. in.} \quad \text{ft. h. bricks}}{(28 \text{ .. } 10) \times 20 \times 4}{3 \text{ half bricks}} = 768 \text{ .. } 10\frac{2}{3} \text{ the next 20 feet.}$$

$$\frac{\text{ft. in.} \quad \text{ft. in.}}{(28 \text{ .. } 10) \times (15 \text{ .. } 8)} = 451 \text{ .. } 8\frac{2}{3} \text{ brick and half thick.}$$

$$\frac{\text{ft. in.} \quad \text{ft. in.} \quad \text{h. bricks.}}{(28 \text{ .. } 10) \times (10 \text{ .. } 6)^* \times 2}{2 \times 3 \text{ half bricks}} = 100 \text{ .. } 11 \text{ in the gable.}$$

$$\begin{array}{r} \text{sq. ft.} \quad \text{sq. yards} \\ \text{Divide by 9) } 2282 \text{ .. } 7\frac{2}{3} \text{ (} 253 \cdot 626 \text{. Ans.} \end{array}$$

* If four courses of bricks make a foot, 42 courses make $10\frac{1}{2}$ feet.

(Page 84.)

MASON-WORK.

$$\text{Ex. 1. } \frac{\text{ft. in.} \quad \text{ft. in.} \quad \text{ft. cub. feet.}}{(53 \text{ .. } 6) \times (12 \text{ .. } 3) \times 2} = 1310\frac{3}{4}. \text{ Ans.}$$

$$\text{Ex. 2. } \frac{\text{ft. in.} \quad \text{ft. in.} \quad \text{ft. ft.} \quad \text{sq. ft.} \quad \text{cub. feet.}}{(24 \text{ .. } 3) \times (10 \text{ .. } 9) \times 2} = 24 \cdot 25 \times 21 \cdot 5 = 521 \cdot 375. \text{ Ans.}$$

(Key to Vol. II. page 84.)

$$\text{Ex. 3. } \begin{array}{cccccc} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} & \text{sq. ft.} & \text{pts. in.} & \text{sq. ft.} \\ (5 \text{ .. } 7) \times (1 \text{ .. } 10) & = & 10 \text{ .. } 2 \text{ .. } 10 & = & 10 \cdot 2361 \end{array}$$

Multiply by 8s.

$$\text{PRODUCT } \underline{\underline{\text{£}4 \text{ .. } 1 \text{ .. } 10 \cdot 608. \text{ Ans.}}}$$

$$\text{Ex. 4. } \begin{array}{cccccc} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} & \text{sq. ft.} & \text{pts.} \\ (4 \text{ .. } 6) \times (3 \text{ .. } 2) & = & 14 \text{ .. } 3 \end{array}$$

$$\text{And } \begin{array}{cccccc} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} & & \\ (4 \text{ .. } 4) \times (1 \text{ .. } 9) & = & 7 \text{ .. } 7 \end{array}$$

$$\text{Sum } \underline{\underline{21 \text{ .. } 10 = 21\frac{1}{2} \text{ sq. feet. Ans.}}}$$

(Page 85.)

CARPENTERS' WORK.

$$\text{Ex. 1. } \frac{\begin{array}{cccc} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} \\ (48 \text{ .. } 6) \times (24 \text{ .. } 3) \end{array}}{100 \text{ sq. feet.}} = \begin{array}{cc} \text{sqs.} & \text{sq. feet} \\ 11 \text{ .. } 76\frac{1}{8} \end{array} \text{ Ans.}$$

$$\text{Ex. 2. } \frac{\begin{array}{cccc} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} \\ (36 \text{ .. } 3) \times (16 \text{ .. } 6) \end{array}}{100 \text{ sq. feet}} = \begin{array}{cc} \text{sqs.} & \text{sq. feet} \\ 5 \text{ .. } 98\frac{1}{3} \end{array} \text{ Ans.}$$

$$\text{Ex. 3. } \frac{\begin{array}{cccc} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} \\ (173 \text{ .. } 10) \times (10 \text{ .. } 7) \end{array}}{100 \text{ sq. feet.}} = \begin{array}{c} \text{squares} \\ 18 \cdot 3973 \end{array} \text{ Ans.}$$

$$\text{Ex. 4. } \frac{\begin{array}{cccc} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} \\ (52 \text{ .. } 8) \times (30 \text{ .. } 6) \times 3 \end{array}}{2 \times 100 \text{ sq. feet}} = \begin{array}{c} \text{squares} \\ 24 \cdot 095 \end{array}$$

Multiply by 10·5 shillings.

$$\text{PRODUCT } \underline{\underline{\text{£}12 \text{ .. } 12 \text{ .. } 11 \text{ .. } 3 \cdot 4 \text{ Ans.}}}$$

(Key to Vol. II. page 86.)

Ex. 5. $\begin{matrix} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} & \text{sq. ft.} & \text{pts.} \\ (83 \dots 8) \times (12 \dots 6) & = & 1045 \dots 10 & \text{whole area.} & \text{Also} \\ \text{ft.} & \text{in.} & \text{ft.} & \text{in.} & \text{sq. ft.} & \text{pts.} \\ (7 \dots 8) \times (3 \dots 6) \times 3 & = & 80 \dots 3 & \text{---} & \text{the three windows.} \\ \text{ft.} & \text{ft.} & \text{in.} & & & \\ \text{And } 7 \times (3 \dots 6) & = & 24 \dots 6 & \text{---} & \text{the door.} \end{matrix}$

Sum 104 .. 9

Half sum - - - - 52 .. 4½ square yards

Total 1098·2083 = 122·02314

Multiply by 6sh.

PRODUCT $\begin{matrix} \text{s.} & \text{d.} & \text{qrs.} \\ \text{£}36 \dots 12 \dots 1 \dots 2 \cdot 664 & \text{Ans.}^* \end{matrix}$

* *Dr. Hutton, in his Answer, gives more than three farthings too much.*

SLATING AND TILING.

Ex. 1. $\begin{matrix} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} & \text{sq. ft.} & \text{pts.} & \text{in.} & \text{sq. yds.} \\ (45 \dots 9) \times (34 \dots 3) & = & 1566 \dots 11 \dots 3 & = & 174 \frac{5}{8} & \text{Ans.} \end{matrix}$

Ex. 2. $\begin{matrix} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} & \text{ft.} & \text{in.} \\ (27 \dots 5) + 2(1 \dots 4) & = & 30 \dots 1 & \text{between the ridges at bottom.} \end{matrix}$

But $\begin{matrix} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} & \text{sq. ft.} & \text{pts.} & \text{in.} \\ (43 \dots 10) \times (30 \dots 1) & = & 1318 \dots 7 \dots 10 \\ \text{Of which the half is} & & 659 \dots 3 \dots 11 \end{matrix}$

Sum 1977 .. 11 .. 9 flat and half.

Or, divided by 100 - - - 19·7798 squares, *nearly.*

Multiply by 25·5sh.

PRODUCT £25 .. 4 .. 4½. Ans.*

* *Differing $\begin{matrix} \text{s.} & \text{d.} \\ \text{£}0 \dots 14 \dots 10 \frac{3}{4} \end{matrix}$ from Dr. Hutton's Answer.*

OTHERWISE, *and more correctly.*

$\begin{matrix} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} \\ (30 \dots 1) \times (30 \dots 1) & = & 905 \cdot 00674 & \text{square of the hypotenuse.} \end{matrix}$

(Key to Vol. II. page 86.)

$$\text{And } 2\sqrt{\frac{905.00674 \text{ ft.}}{2}} = 42.544 \text{ nearly, girt of the ridges.}$$

$$\begin{array}{r} \text{ft.} \quad \text{ft.} \quad \text{sq. feet} \quad \text{squares} \\ \text{But } 42.544 \times 43.83 = 1864.70453 = 18.6470453 \\ \text{Multiply by} \quad \quad \quad 25.5sh. \end{array}$$

$$\text{PRODUCT } \underline{\underline{\pounds 23 \text{ .. } 15 \text{ .. } 6. \text{ Ans.}^*}}$$

* Differing from Hutton's Answer $\pounds 0 \text{ .. } 13 \text{ .. } 11\frac{1}{2}$ in a contrary direction. The mean amount of this and the former result, however, is $\pounds 24 \text{ .. } 9 \text{ .. } 11\frac{1}{2}$, being in excess only $5\frac{1}{2}d.$ above the Answer given.

PLASTERING.

(Page 87.)

$$\text{Ex. 1. } \frac{\begin{array}{cc} \text{ft.} & \text{in.} \\ (43 & .. & 3) \end{array} \times \begin{array}{cc} \text{ft.} & \text{in.} \\ (25 & .. & 6) \end{array}}{9 \text{ sq. feet}} = 122\frac{1}{2} \text{ square yards. Ans.}$$

$$\text{Ex. 2. } \frac{\begin{array}{cc} \text{ft.} & \text{in.} \\ (21 & .. & 8) \end{array} \times \begin{array}{cc} \text{ft.} & \text{in.} \\ (14 & .. & 10) \end{array}}{9 \text{ sq. feet}} \times 10 = \pounds 1 \text{ .. } 9 \text{ .. } 8\frac{3}{4} \text{ Ans.}$$

$$\text{Ex. 3. } \begin{array}{cc} \text{ft.} & \text{in.} \\ (18 & .. & 6) \end{array} \times \begin{array}{cc} \text{ft.} & \text{in.} \\ (12 & .. & 3) \end{array} = 226 \text{ .. } 7 \text{ .. } 6 = 25.18 \text{ sq. yds. of ceiling.}$$

$$\text{And } 2 \begin{array}{cc} \text{ft.} & \text{in.} \\ (18 & .. & 6) \end{array} + \begin{array}{cc} \text{ft.} & \text{in.} \\ (12 & .. & 3) \end{array} \times \begin{array}{cc} \text{ft.} & \text{in.} \\ (10 & .. & 6) \end{array} = 645 \text{ .. } 9 \text{ whole area of wall.}$$

$$\text{Subtr. } \left\{ \begin{array}{l} \begin{array}{cc} \text{ft.} & \text{in.} \\ (3 & .. & 8) \end{array} \times 7 = 25 \text{ .. } 8 \text{ door.} \\ \begin{array}{cc} \text{ft.} & \text{ft.} \\ 5 \times 5 = 25 \text{ fire-place.} \end{array} \end{array} \right\} 50 \text{ .. } 8 \text{ deduction.}$$

$$\text{Remainder } \underline{\underline{595 \text{ .. } 1}} = 66.12 \text{ rendering.}$$

$$\text{But } 66.12 \times 3 + 25.18 \times 8 = \pounds 1 \text{ .. } 13 \text{ .. } 3\frac{3}{4} \text{ Ans.}^*$$

* Being a halfpenny more than the estimate of Dr. Hutton.

(Key to Vol. II. page 87.)

Ex. 4. $\frac{2(14..5 + 13..2) \times (9..3) - (7 \times 4)}{9 \text{ sq. feet}}$ $\overset{\text{ft. in. ft. in. ft. in. ft. ft.}}{\text{sq.yds. ft. pts. in.}} = 53..5..3..6$ rendering.

$(55..2) \times (8..6) = 39..0..11 = 4..3..0..11$ of cornice.

And $\frac{(13..7) \times (12..4)}{9 \text{ sq. feet}} = \dots\dots\dots 18..5..6..4$ of ceiling.

Whereby the Answer is obvious.

PAINING.

Ex. 1. $\frac{(65..6) \times (12..4)}{9 \text{ sq. feet}}$ $\overset{\text{ft. in. ft. in.}}{\text{sq. ft. pts. in.}} = 89\frac{1}{3}$ square yards. Ans.

Ex. 2. $2(20 + 14..6) \times (10..4) = 713$ sq. ft. whole area of wall.

Subtr. $\left\{ \begin{array}{l} \text{ft. in. sq. ft. pts.} \\ 4(4..4) = 17..4 \text{ fire-place.} \\ \text{ft. ft. in. sq. ft.} \\ 2 \times 6 \times (3..2) = 38 \text{ windows.} \end{array} \right\} \overset{\text{pts.}}{55..4} \text{ deduction.}$

Remainder $\overset{\text{sq. yds.}}{657\frac{2}{3}} = 73\frac{2}{27}$ Ans.

(Page 88.)

Ex. 3. $2(24..6 + 16..3) \times (12..9) = 1039..1..6$ inside area.

Add

{	Outside of the door	$7 \times (3..6) =$	$24..6..0$
	Inside of the shutters	$2(7..9) \times (3..6) =$	$54..3..0$
{	Window-breaks, cills, and	$\times (1..3) =$	$60..0..0$
	soffits $4(8\text{ft. } 6\text{in.} + 3\text{ft. } 6\text{in.})$		

Sum $1170..10..6$

Deduct fire-place $5 \times (5..6) = 27..6..0$

Remainder $\overset{\text{sq.yds. ft. 12ths}}{1150..4..6} = 127..7..4\frac{1}{2}$

Multiply by $6d.$

Product and Answer $\underline{\underline{\pounds 3..3..10\frac{3}{4}}}$

(Key to Vol. II. page 88.)

GLAZING.

$$\text{Ex. 1. } \begin{array}{ccc} \text{ft.} & \text{ft.} & \text{sq. feet} \\ 4.25 \times 2.75 = 11.6875. & \text{Ans.} & \end{array}$$

$$\text{Ex. 2. } \frac{\begin{array}{cc} \text{ft.} & \text{in.} \\ 12 & 6 \end{array} \times \begin{array}{cc} \text{ft.} & \text{in.} \\ 6 & 9 \end{array}}{2} \times \begin{array}{c} d. \\ 10 \end{array} = \begin{array}{ccc} \text{s.} & \text{d.} \\ \mathcal{L}1 & 15 & 1\frac{3}{4}. \end{array} \text{ Ans.}$$

$$\text{Ex. 3. } 3 \begin{array}{ccc} \text{ft.} & \text{in.} & \text{ft. in.} \\ 7 & 10 & 6 & 8 & 5 & 4 \end{array} \times \begin{array}{cc} \text{ft.} & \text{in.} \\ 3 & 11 \end{array} \times \begin{array}{c} d. \\ 14 \end{array} = \begin{array}{ccc} \text{s.} & \text{d.} \\ \mathcal{L}13 & 11 & 10\frac{1}{2}. \end{array} \text{ Ans.}$$

(Page 89.)

$$\text{Ex. 4. } \begin{array}{ccc} \text{ft.} & \text{in.} & \text{ft. in.} \\ 7 & 9 & 6 & 6 & 5 & 3\frac{1}{4} \end{array} \times \begin{array}{cc} \text{ft.} & \text{in.} \\ 3 & 9 \end{array} = \begin{array}{c} \text{sq. ft. pts. in.} \\ 219.7.3\frac{3}{4} \end{array} \text{ rectang. frames.}$$

$$\text{Add } \begin{array}{cc} \text{ft.} & \text{in.} \\ 1 & 10\frac{1}{2} \end{array} \times \begin{array}{cc} \text{ft.} & \text{in.} \\ 3 & 9 \end{array} = \begin{array}{c} 7.0.4\frac{1}{2} \end{array} \text{ oval window.*}$$

$$\begin{array}{r} \text{Sum } 226.7.8\frac{1}{4} \text{ Total.} \\ \text{Multiply by } 13d. \end{array}$$

$$\text{PRODUCT } \mathcal{L}12.5.6\frac{1}{4}. \text{ Ans.}$$

* The oval window is computed as rectangular; else the expense is only $\mathcal{L}12.3.9\frac{1}{2}$, the area of the oval being 5.52234375 square feet.

PAVING.

$$\text{Ex. 1. } \frac{\begin{array}{cc} \text{ft.} & \text{in.} \\ 35 & 4 \end{array} \times \begin{array}{cc} \text{ft.} & \text{in.} \\ 8 & 3 \end{array}}{9 \text{ sq. feet}} \times \begin{array}{c} \text{s.} & \text{d.} \\ 3 & 4 \end{array} = \begin{array}{ccc} \text{s.} & \text{d.} \\ \mathcal{L}5 & 7 & 11\frac{1}{2}. \end{array} \text{ Ans.}$$

$$\text{Ex. 2. } \frac{\begin{array}{cc} \text{ft.} & \text{in.} \\ 27 & 10 \end{array} \times \begin{array}{cc} \text{ft.} & \text{in.} \\ 14 & 9 \end{array}}{9 \text{ sq. feet}} \times \begin{array}{c} \text{s.} & \text{d.} \\ 3 & 2 \end{array} = \begin{array}{ccc} \text{s.} & \text{d.} \\ \mathcal{L}7 & 4 & 5\frac{1}{4}. \end{array} \text{ Ans.}$$

$$\text{Ex. 3. } 45 - \begin{array}{cc} \text{ft.} & \text{in.} \\ 5 & 3 \end{array} = \begin{array}{cc} \text{ft.} & \text{in.} \\ 39 & 9 \end{array} \text{ breadth laid with pebbles.}$$

(Key to Vol. II. page 89.)

$$\text{And } \frac{\text{ft. ft. in.}}{63 \times (39 \dots 9)} \times \frac{\text{s. d.}}{(2 \dots 6)} = \text{£}34 \dots 15 \dots 7\frac{1}{2} \text{ pebbled.}$$

9 sq. feet

$$\text{Also } \frac{\text{ft. ft. in.}}{63 \times (5 \dots 3)} \times \frac{\text{s.}}{3} = 5 \dots 10 \dots 3 \text{ foot path.}$$

9 sq. feet

$$\text{Sum } \text{£}40 \dots 5 \dots 10\frac{1}{2} \text{ Ans.}$$

PLUMBERS' WORK.

$$\text{Ex. 1. } \frac{\text{ft. in. ft. in. lb. lb. oz.}}{(30 \dots 6) \times (3 \dots 3) \times 8\frac{1}{2}} = 1091 \dots 3 \text{ Ans.}$$

(Page 90.)

$$\text{Ex. 2. } \frac{\text{ft. ft. lb. lb.}}{43 \times 32 \times 9 \cdot 831} = 13527 \cdot 456 \text{ roofing.}$$

$$\text{And } \frac{\text{ft. ft. lb. lb.}}{57 \times 2 \times 7 \cdot 373} = 840 \cdot 522 \text{ guttering.}$$

$$\text{Sum } 14367 \cdot 978 = 128 \cdot 286 \text{ of lead.}$$

Multiply by 18s. per cwt.

$$\text{PRODUCT } \text{£}115 \dots 9 \dots 1\frac{3}{4} \text{ Ans.}$$

TIMBER MEASURING.

The Superficies of Boards.

$$\text{Ex. 1. } \frac{\text{ft. in. ft. in. d. s. d.}}{(12 \dots 6) \times (0 \dots 11) \times 1\frac{1}{2}} = \text{£}0 \dots 1 \dots 5 \cdot 17 \text{ Ans.}$$

By the Sliding Rule.*

Set 12 on B to 11 inches on A; then opposite 12½ feet on B are 11½ feet on A, which multiplied by 1½d., the result is 17½d. nearly. Ans.

$$\text{Ex. 2. } \frac{\text{ft. in. ft. in. sq. ft. pts. in.}}{(11 \dots 2) \times (1 \dots 10)} = 20 \dots 5 \dots 8 \text{ Ans.}$$

* The Directions for the Sliding Rule being sufficiently intelligible in Hutton's text, it would be superfluous to give more Examples solved in that way.

(Key to Vol. II. page 90.)

$$\text{Ex. 3. } \begin{array}{cccccc} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} & d. & s. & d. \\ (12 \dots 9) \times (1 \dots 3) \times 2\frac{1}{2} = \mathcal{L}0 \dots 3 \dots 3\frac{3}{4}. & \text{Ans.} \end{array}$$

$$\text{Ex. 4. } \begin{array}{cccccc} \text{in.} & & \text{in.} & \text{in.} & \text{in.} & \text{ft.} & \text{in.} \\ (13\frac{1}{2} \times 2) + 14\frac{1}{2} + 18 + 11\frac{1}{4} = 70\frac{3}{4} = 5 \dots 10\frac{3}{4}. & \text{mean breadth.} \end{array}$$

$$\text{But } \begin{array}{cccccc} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} & d. & s. & d. \\ (5 \dots 10\frac{3}{4}) \times (17 \dots 6) \times 3 = \mathcal{L}1 \dots 5 \dots 9\frac{1}{2}. & \text{Ans.} \end{array}$$

(Page 91.)

The Solidity of Squared Timber.

$$\text{Ex. 1. } \begin{array}{cccccc} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} & \text{ft.} & \text{in.} & \text{ft.} & \text{c.ft.} & \text{12s.} & \text{144s.} & \text{in.} \\ (18 \dots 6) \times \frac{1 \dots 6 + 1 \dots 3}{2} \times \frac{1 \dots 3 + 1}{2} = 28 \dots 7 \dots 4 \dots 10\frac{1}{2}. & \text{Ans.*} \end{array}$$

* Dr. Hutton writes in his Answer 7 inches, meaning (it may be supposed) 1008 inches for 1066½ inches, the exact Answer being 28 ft. 1066½ in. See the 49th page of the KEY.

$$\text{Ex. 2. } \begin{array}{cccc} \text{ft.} & \text{ft.} & \text{ft.} & \text{solid feet} \\ 24 \cdot 5 \times 1 \cdot 04 \times 1 \cdot 04 = 26 \cdot 4992 \text{ or } 26\frac{1}{2} \text{ cubic ft. } & \text{nearly.} & \text{Ans.} \end{array}$$

$$\text{Ex. 3. } \begin{array}{ccc} \text{in.} & \text{in.} & \text{in.} \\ \frac{19\frac{1}{8} + 9\frac{7}{8}}{2} = 14\frac{1}{2} & \text{mean breadth and thickness.} \end{array}$$

$$\text{But } \begin{array}{ccc} \text{ft.} & \text{in.} & \text{in.} \\ 20 \cdot 38 \times \frac{14 \cdot 5 \times 14 \cdot 5}{144 \text{ sq. in.}} = 29 \cdot 7562 \text{ solid feet.} & \text{Ans.} \end{array}$$

(Page 92.)

$$\text{Ex. 4. } \begin{array}{cccccc} \text{ft.} & \text{ft.} & \text{ft.} & \text{ft.} & \text{ft.} & \text{solid feet.} \\ \frac{178 + 1 \cdot 04}{2} \times \frac{1 \cdot 23 + 0 \cdot 91}{2} \times 27 \cdot 36 = 41 \cdot 278. & \text{Ans.} \end{array}$$

The Solidity of Round Timber.

$$\text{Ex. 1. } \begin{array}{ccc} \text{ft.} & \text{ft.} & \text{ft.} \\ 3\frac{1}{2} \times 3\frac{1}{2} = 12\frac{1}{4} & \text{the square of the } & \text{quarter-girt.} \\ \text{Multiply by } 9\frac{1}{2} & \text{ft. the length.} \end{array}$$

$$\text{PRODUCT } 116\frac{3}{8} \text{ solid feet. Ans.*}$$

* Here Dr. Hutton is in error by the Rule (which is fundamentally wrong) 72 cubic inches, or the twenty-fourth part of a solid foot.

Nota Bene. This and the three following Questions being solved in the manner directed, the Answers must not be considered the true content.

(Key to Vol. II. page 92.)

Ex. 2. $\frac{\text{ft. ft.}}{2} \frac{14+2}{2} = \text{ft. } 8$ mean girt; whereof $\frac{1}{4}$ is 2 feet, *quarter-girt*.

But $2 \times 2 = 4$ the square of the *quarter-girt*.

Multiply by 24 ft. the length.

PRODUCT 96 solid feet. Ans.

Ex. 3. $\frac{\text{ft.}}{4} \frac{3.5}{4} = \text{ft. } 0.7875$ the *quarter-girt*. And

$\frac{\text{feet}}{0.7875} \times \frac{\text{feet}}{0.7875} \times \frac{\text{feet}}{14.5} = \frac{\text{solid feet}}{8.992265625}$. Ans.

Ex. 4. $\frac{\text{ft. ft. ft. ft. ft.}}{5} \frac{9.43+7.92+6.15+4.74+3.16}{5} = \text{ft. } 6.28$ mean girt.

Now $\frac{\text{ft.}}{4} \frac{6.28}{4} = \text{ft. } 1.57$ the *quarter-girt*.

And $1.57 \times 1.57 = 2.4649$ square of the *quarter-girt*.
Multiply by 17.25 ft. the length.

PRODUCT 42.519525 solid feet. Ans.

END OF THE MENSURATION.

NATURAL PHILOSOPHY.

(Key to Vol. II, page 151.)

The Laws of Gravity.

- Ex. 1. As $1''^2 : 7''^2 :: 16\frac{1}{12}$ ft. : $788\frac{1}{12}$ feet, whole descent.
 And $1'' : 7'' :: 32\frac{1}{6}$ ft. : $225\frac{1}{6}$ feet, velocity acquired.
- Ex. 2. As $32\frac{1}{6}$ ft. : 100 ft. :: $1'' : 3''\frac{21}{193}$ time of the fall.
 And $1''^2 : 3''\frac{21}{193}]^2 :: 16\frac{1}{12}$ ft. : $155\frac{85}{193}$ ft. passed through.
- Ex. 3. As $16\frac{1}{12}$ ft. : 400 ft. :: $1''^2 : 4''\frac{76}{77}]^2$ time in motion.
 And $1'' : 4''\frac{76}{77} :: 32\frac{1}{6}$ ft. : $160\frac{32}{77}$ ft. velocity generated.

QUESTIONS IN PRACTICAL GUNNERY.

(Page 162.)

Ex. 1.

	lb.	lb.	lb.	lb.	ft.	ft.	in.	
As	$\sqrt{196}$	$\sqrt{18}$	$\cdot 14$	$4\cdot2426407$::	1600	: 485 - 13	shell.
	$\sqrt{90}$	$\sqrt{8}$	$\cdot 9\cdot486833$	$2\cdot828427$,				} 10 shell.
					::	1600	: 477	
	$\sqrt{48}$	$\sqrt{4}$	$\cdot 6\cdot9282032$	2	::	1600	: 462 - 8	shell.
	$\sqrt{16}$	$\sqrt{2}$	$\cdot 4$	$1\cdot4142136$::	1600	: 566 - $5\frac{1}{2}$	shell.
	$\sqrt{8}$	$\sqrt{1}$	$\cdot 2\cdot8284271$	1	::	1600	: 566 - $4\frac{2}{3}$	shell.

ft. ft. ft. ft. ft.
 Wherefore 485, 477, 462, 566, and 566, respectively. Ans.

(Key to Vol. II. page 162.)

Ex. 2. As Sin. $2 \times (45^\circ \angle \text{gr. range})$ Log. ar. co. — 10.
 : Sin. $2 \times (30^\circ \text{ .. } 16' \text{ elev. prop.})$, Log. 9.9398396
 :: 1000 yds. given range, - - - - Log. 3. - - - -
 : 870.64 yds, range sought. - - - Log. 2.9398396

(Page 163.)

Ex. 3. As 3750 ft. range found, Log. ar. co. — 4.4259687
 : 2810 ft. range proposed, - - - Log. 3.4487063
 :: Sin. $2 \times (45^\circ \text{ trial elevation})$, - - Log. 10. - - - -
 : Sin. $2 \times \left\{ \begin{array}{l} 24^\circ \text{ .. } 16' \\ 65^\circ \text{ .. } 44' \end{array} \right\} \text{elevat. reqd.}$ Log. 9.8746750

Ex. 4. As Sin. $2(32^\circ \text{ .. } 12' \text{ giv. elev.})$ Log. ar. co. — 10.0448741
 : Sin. $2 \times (45^\circ \angle \text{ of great. range})$, Log. 10. ———
 :: 3250 ft. proposed range, - - - Log. 3.5118834
 : 3603.7 ft. greatest range. - - - Log. 3.5567575

But $3603.7 \text{ ft.} \div 2 = 1802 \text{ feet}$ nearly, the impetus required.

Again,

$2\sqrt{1802 \times 16 \frac{1}{2}} \text{ ft.} = 2\sqrt{28982 \frac{1}{2}} \text{ ft.} = 340.4 \text{ ft.}$ the velocity demanded.

And

As 485 ft. tab. velocity, - - - Log. ar. co. — 3.3142583
 : 340.4 ft. velocity necessary, - - - Log. 2.5319896
 :: $\sqrt{9 \text{ lb. tab. charge}}$, - - - - - Log. 0.4771213
 : $\sqrt{4.4334 \text{ lb.}} = 4 \text{ lb. } 6 \frac{47}{50} \text{ oz. charge req.}^* \text{ Log. } .3233692$

* In the Impetus and Velocity we agree nearly with Hutton, but differ more than $\frac{1}{2}$ lb. in the Charge of Powder. Indeed if the Velocity be taken at 340 ft. as he states it, we differ $\frac{3}{4}$ lb. in the Charge.

Ex. 5. As Sin. $2 \times (25^\circ \text{ .. } 12' \text{ actual elevation})$, } Log. ar. co. — 10.1132199
 : Sin. $2 \times (36^\circ \text{ .. } 15' \text{ prop. elevation})$, Log. 9.9794195
 :: 3500 ft. range measured, - - - - - Log. 3.5440680
 : 4332.1 ft. range sought. - - - - - Log. 3.6367074

(Key to Vol. II. page 163.)

Ex. 6. As $\begin{matrix} \text{ft.} & \text{ft.} & \text{ft.} & \text{ft.} & \text{lb.} & \text{lb.} \\ 4000 & : & 3000 & \cdot\cdot & 4 & : & 3 & :: & 9 & : & 6\frac{3}{4}. \end{matrix}$ Ans.

Ex. 7. Because [Theor. iii. page 156, vol. ii.] $T'' = 2\sqrt{\frac{H}{g}}$ ft. and that $g = 16\frac{1}{2}$ feet; if, rejecting the fraction, as in comparison small, g be taken = 16 feet, it is

$$T'' = 2\sqrt{\frac{H}{16}} \text{ft.} = 2\sqrt{\frac{\frac{1}{4}R}{16}} \text{ft.} = 2 \times \frac{\sqrt{\frac{1}{4}R}}{\sqrt{16}} \text{ft.} = \frac{1}{2}\sqrt{\frac{1}{4}R} \text{ft.}$$

$$\text{But } \frac{1}{2}\sqrt{\frac{1}{4}R} \text{ft.} = \frac{1}{2} \times \sqrt{\frac{1}{4}R} \text{ft.} = \frac{1}{2} \times \frac{1}{2}\sqrt{R} \text{ft.} = \frac{1}{4}\sqrt{R} \text{ft.}$$

Wherefore the time in seconds = $\frac{1}{4}\sqrt{\text{range in feet.}}$ Ans.

Ex. 8. As Tang. $45^\circ \angle$ gr. range, Log. ar. co. - 10. _____
 : Tang. 32° given \angle of elevation, Log. 9.7957892
 :: 812.5 ft. $\frac{1}{4}$ proposed range, - - Log. 2.9098234
 : 507.72 ft. greatest altitude. - - - Log. 2.7056126

Now $2\sqrt{\frac{507.72}{16\frac{1}{2}}}$ ft. = $2 \times 5''.6 = 11''.2 = 11\frac{1}{4}$ seconds nearly. Ans.

Ex. 9. $90^\circ - (32^\circ .. 30') = 57^\circ .. 30'$. And
 $(32^\circ .. 30') - (8^\circ .. 15') = 24^\circ .. 15'$. But

The log.-sin. of $57^\circ .. 30'$ - - - - is 9.9260292

- - - - of $24^\circ .. 15'$ - - - - is 9.6135446

The ar. co. of twice the log. cos. $8^\circ .. 15'$ is - 20.0090356

The log. of 4(3000 ft. impetus) - - - - is 4.0791812

The sum of these 4 logarithms - - - - is 3.6277906 which
 is the log. of 4244 feet range on the ASCENT.

Again

$$\frac{90^\circ - (8^\circ .. 15')}{2} = 40^\circ .. 52'\frac{1}{2}. \text{ And}$$

The log.-sin. of $57^\circ .. 30'$ - - - - is 9.9260292

- - - - of $40^\circ .. 52'\frac{1}{2}$ - - - - is 9.8158506

The ar. co. of twice the log.-cos. of $8^\circ .. 15'$ is - 20.0090356

The log. of 4(3000 ft. impetus) - - - - is 4.0791812

The sum of these 4 logarithms - - - - is 3.8300966 which
 is the log. of 6762.3 ft. range on the DESCENT.*

* By which it should appear that Dr. Hutton had found 17.3 feet more than the true range on the descending plane.

(Key to Vol. II. page 163.)

Ex. 10. Here the impetus (by Art. 98) is 3000 ft. *nearly*. And

$$2\sqrt{3000 \times 16\frac{1}{2}} \text{ ft.} = 2\sqrt{48250} \text{ ft.} = 439.363 \text{ ft. the VELOCITY.}$$

But

$$\text{As } 485^* : 439.363 :: \sqrt{9 \text{ lb.}} : \sqrt{7.38209 \text{ lb.}} \text{ nearly, the CHARGE.}$$

Ans. †

* See Art. 107, page 162, vol. ii.

† This Answer and the Answer given by Dr. Hutton differ in the decimal only.

Ex. 11. Finding the impetus, as in the foregoing Example, to

$$\text{be nearly 3000 feet; (for } \sqrt{\frac{2(7.375) \text{ lb.}}{196}} \times 1600 = 438.4 \text{ feet}$$

$$\text{the velocity: and } \frac{438.4 \text{ ft.}^2}{2\sqrt{16}} = 3003.04 \text{ ft. the impetus) it is,}$$

$$\text{As Rad.: Tang. } 8^\circ .. 15' :: \frac{3000 \text{ ft.}}{2} : 217.49 \text{ ft.}$$

And

$$\begin{array}{l} \text{As Radius, - - - - - Log. ar. co. - 10.} \\ : \text{Cos. } 8^\circ .. 15' \angle \text{ of descent, - - - Log. } 9.9954822 \\ :: 6745 \text{ ft. given range, - - - - - Log. } 3.8289820 \\ \hline : 6675.2 \text{ ft. horizontal range. - - - - - Log. } 3.8244642 \end{array}$$

But

$$\frac{6675.2}{4} \text{ ft.} - 217.49 \text{ ft.} = 1451.3 \text{ ft.}$$

And

$$\begin{array}{l} \text{As } 1500 \text{ ft. } \frac{1}{2} \text{ the impetus, - Log. ar. co. - 4.8239087} \\ : 1451.3 \text{ ft. - - - - - Log. } 3.1617572 \\ :: \text{Cos. } 8^\circ .. 15' \angle \text{ of descent, - - - Log. } 9.9954822 \\ \hline : \text{Cos. } 16^\circ .. 46'. - - - - - \text{Log. } 9.9811481 \end{array}$$

Now

$$\left. \begin{array}{l} 45^\circ - \frac{(8^\circ .. 15') + (16^\circ .. 46')}{2} = 32^\circ .. 29'\frac{1}{2} \\ \text{Also } 90^\circ - (32^\circ .. 29'\frac{1}{2} + 8^\circ .. 15') = 49^\circ .. 15'\frac{1}{2} \end{array} \right\} \text{Ans.}$$

Otherwise.

Because (*Theor. i. page 159.*) $R = \frac{cs}{c^2} \times 4a$, or $\frac{c^2 R}{4a} = cs$; it is,
 (having first found the impetus to charge $7\frac{3}{8}$ lb. to be 3000 feet
 nearly, as before) $cs = \frac{\text{cos.}^2 \text{ of } (8^\circ .. 15') \times 6745 \text{ ft.}}{12000 \text{ ft.}}$

(Key to Vol. II. page 163.)

IN LOGARITHMS.

Twice the log-cosine of $8^{\circ} .. 15'$ = - - 19.9909644

The logarithm of 6745 ft. range = - - 3.8289820

Log. ar. co. of 4(3000 ft. impetus) = - - 5.9208188

Log. Sum	19.7407652	which
----------	------------	-------

is the logarithm of the cos. of direction above the horizon + the log-sine of direction above the plane. But the angle of direction above the plane is evidently $8^{\circ} .. 15'$ greater than the angle of direction above the horizon. Entering, therefore, the table of log-sines, find the angle of which the log-cosine added to the log-sine of the angle $8^{\circ} .. 15'$ greater, shall approach the nearest to 19.7407652.

Now the log.-cosine of $32^{\circ} .. 30'$ is 9.9260292Log.-sine of $40^{\circ} .. 45'$ is 9.8147534


Sum	19.7407826
-----	------------

Also log.-cosine of $49^{\circ} .. 15'$ is 9.8147534Log.-sine of $57^{\circ} .. 30'$ is 9.9260292

Sum	19.7407826
-----	------------

Consequently $32^{\circ} .. 30'$ and $49^{\circ} .. 15'$ are the elevations required.

GEOMETRICALLY.



Draw any right line AH representing the horizon, and at A erect the perpendicular AP = 3000 ft. on any convenient scale of equal parts. Bisect AP in B, and draw BD parallel to AH. Also

(Key to Vol. II. page 163.)

make the angles PAQ, HAI, each, equal to 8° .. 15'. And from c (the point of intersection of AQ and BD) at the distance CA or CP, describe the arc ADQP. By the scale employed for AP make AI = 6745 ft. and take Am one fourth-part of AI. If through the point m there be drawn the line oonm parallel to AP, the two points oo in the circumference ADQP will indicate the elevations required. From A therefore, draw straight lines through the points o and o; the angle oAn measured by a line of chords, or goniometer, is $\left\{ \begin{array}{l} 49^\circ \text{ .. } 15' \\ 32^\circ \text{ .. } 30' \end{array} \right\}$ nearly, the two elevations sought.*

* Again the KEY and COURSE are not agreed in the Answer.

Ex. 12. $45^\circ - (8^\circ \text{ .. } 30') = 36^\circ \text{ .. } 30'$ the angle of direction above the plane.

And $\sqrt{\frac{2(2304)}{16}}$ ft. † = 16''·97 time of greatest horizontal range.

Lastly, As Sin. 45°	- - - - -	Log. ar. co. -	10·1505150
: Sin. 36° .. 30'	- - - - -	Log.	9·7743876
:: 16''·97 time of greatest hor. range,		Log.	1·2296818
: 14''·275 the time sought.	- - - - -	Log.	1·1545844

Otherwise, and more correctly.

Because, [Theor. iv. page 159, vol. ii.] $2T = \frac{s}{c} \sqrt{a}$ (if 16 feet, instead of $16\frac{1}{2}$ feet, be substituted for g) it is,

Half the log. of 2304 ft. - - -	is	1·6812412 = log. \sqrt{a} .
The log. sine of 36° .. 30' - -	is	9·7743876 = log. of s.
The ar. co. log-cos. 8° .. 30' -	is	- 10·0047967 = -log. of c.

The log. of 28''·868 is the Sum = 1·4604255 = log. of 2T''.

Wherefore 14''·434 is the time required.

† Because $T = \sqrt{\frac{tR}{g}}$; and that here $t = 1$, $R = 2a$, and $g = 16$ nearly.

(Key to Vol. II. page 229.)

HYDROSTATICS.

To find the Specific Gravity of a Solid Body heavier than Water.

EXAMPLE. lb. lb. lb.
 $10 - 6\frac{3}{4} = 3\frac{1}{4}$ weight lost in water.

Therefore,

As $3\frac{1}{4}$ lb. loss in water,
 : 10 lb. weight in air,
 :: 1000 lb. specific gravity of water,
 : 3077 lb. *specific gravity required.*

To find the Specific Gravity of a Solid Body lighter than Water.

EXAMPLE. Here the aggregate body weighs 33lb. in air, and 6lb. in water. Also, the denser body weighs 18lb. in air, and 16lb. in water. But $33\text{lb.} - 6\text{lb.} = 27\text{lb.}$ lost by the *aggregate* body in water; and $18\text{lb.} - 16\text{lb.} = 2\text{lb.}$ lost by the *denser* body in water. Therefore,

As $27\text{lb.} - 2\text{lb.} = 25\text{lb.}$ difference of loss,
 : 15lb. weight of the lighter body in air,
 :: 1000lb. specific gravity of water,
 : 600lb. *specific gravity required.*

(Page 230.)

To find the Specific Gravity of a Fluid.

EXAMPLE. oz. oz. oz.
 $40 - 35.61 = 4.39$ loss in the fluid. Therefore,

As 40 oz. absolute weight in air,
 : 4.39 oz. loss of weight,
 :: 7425 oz. specific gravity of cast iron,
 : 814.9 oz. *the specific gravity required.**

* A cubic foot of the fluid in question weighs 185.1 ounces less than Dr. Hutton imagined.

(Key to Vol. II. page 230.)

Specifically to find the Quantity of each of two known Ingredients in a binary Compound.*

EXAMPLE. $8784 - 7320 = 1464$ First difference.
 $9000 - 8784 = 216$ Second difference.
 $9000 - 7320 = 1680$ Third difference.

Also $8784 \times 1680 = 14757120$
 $9000 \times 1464 = 13176000$
 $7320 \times 216 = 1581120$

But, As $14757120 : 13176000 :: 112\text{lb.} : 100\text{lb. of copper.}$ } Ans.
 And $14757120 : 1581120 :: 112\text{lb.} : 12\text{lb. of tin.}$ }

* The Ingredients may be themselves Compound.

(Page 231.)

To determine the Magnitude of Bodies by their Weight.

Ex. 1. As 2520 oz. specific gravity of common stone,
 : 1792 oz. absolute weight in ounces,
 :: 1728 cubic inches, in a cubic foot,
 : $1228\frac{2016}{525}$ cubic inches, content required.

Ex. 2. As 937 oz. specific gravity of gunpowder,
 : 16 oz. absolute weight in ounces,
 :: 1728 cubic inches, in a cubic foot,
 : $29\frac{1}{2}$ cubic inches, content required.

(Page 232.)

Ex. 3. As 925 oz. specific gravity of oak,
 : 35840 oz. absolute weight in ounces,
 :: 1 cubic foot,
 : $38\frac{138}{185}$ cubic feet, content required.

To determine the Weight of Bodies by their Magnitude.

Ex. 1. ft. ft. ft.
 $63 \times 12 \times 12 = 9072$ solid feet in the block.
 And, As 1 solid foot,
 : 9072 solid feet, content of the block,
 :: 2700 oz. specific gravity of marble,
 : $24494400 \text{ oz.} = 683.4 \text{ tons, weight required.}$

(Key to Vol. II. page 232.)

Ex. 2. As 1728 cubic inches, in a cubic foot,
 : 35.25 cubic inches, in 1 pint ale measure,
 :: 937 oz. specific gravity of gunpowder,
 : 19.1 oz. *the weight required.*

Ex. 3. $10 \times 3 \times 2\frac{1}{2} = 75$ cubic feet in the block.

And, As 1 cubic foot,
 : 75 cubic feet, content of the block,
 :: 925 oz. specific gravity of dry oak,
 : 69375 oz. = 4335 $\frac{1}{6}$ lb. *weight required.*

PNEUMATICS.

(Page 246.)

Ex. 1. $\begin{array}{l} \text{The log. of } 29.68 \text{ inches is } 1.4724639 \\ \text{The log. of } 25.28 \text{ inches is } 1.4027771 \end{array}$

$\begin{array}{r} \text{Difference } \cdot 0696868 \\ \text{Multiply by } 10592 \end{array}$

PRODUCT 738.1225856 fathoms.

Again $55^\circ - 50^\circ = 5^\circ$. *difference of temperature.*

But $\frac{5}{435} \times 738.1 = 8.5$ *excess of 738.1 fathoms.*

And $738.1 - 8.5 = 729.6$ fathoms, the height *nearly.*

Ex. 2. $\begin{array}{l} \text{The log. of } 29.45 \text{ inches is } 1.4690853 \\ \text{The log. of } 26.82 \text{ inches is } 1.4284588 \end{array}$

$\begin{array}{r} \text{Difference } \cdot 0406265 \\ \text{Multiply by } 10000 \end{array}$

PRODUCT 406.265 fathoms.

(Key to Vol. II. page 246.)

Again $33^\circ - 31^\circ = 2^\circ$. *difference of temperature.*

And $\frac{2}{435} \times 406.265 = 1.868$ *deficiency of 406.2 &c.*

But $406.265 + 1.868 = 408.133$ fathoms. Ans.

Or, by the other formula;

The *difference of the logarithms, as before*, being .0406265
 Multiply by 10592
 PRODUCT 430.315888

Again $55^\circ - 33^\circ = 22^\circ$ *difference of temperature.*

And $\frac{22}{435} \times 430.315 = 21.763$ *excess of 430.3 &c.*

But $430.315 - 21.763 = 408.552$ fathoms. Ans.*

* In this Example the Answer by either formula differs about 11 fathoms from HUTTON'S Answer.

(Page 247.)

Ex 3. The log. of 1. is - - - .0000000
 The log. of .25 - - - 1.3979400
 Difference .6020600
 Multiply by 10000
 PRODUCT 6020.6 fathoms. Ans.

(Page 255.)

Ex. 1. $57^\circ - 43^\circ = 14^\circ$ *diff. of temperature in the quicksilver.*
 And $\frac{14}{9600} \times 25.28 = .0368$ *correction for 14° temperature.*
 Add 25.28 *observed alt. of the quicksilver.*
 Sum 25.3168 *Barometer corrected for 57° T.*

(Key to Vol. II. page 255.)

Again,

inches

The log. of 29·68	is	1·4724639
The log. of 25·316	is	1·4033951

<i>Difference</i>	·0690688
Multiply by	10000

PRODUCT	690·688 fathoms.
---------	------------------

And

$$\frac{57^\circ + 42^\circ}{2} = 49^\circ \frac{1}{2} \text{ mean temperature of the air.}$$
But $49^\circ \frac{1}{2} - 31^\circ = 18^\circ \frac{1}{2}$.

Also

$\frac{18 \cdot 5}{435}$	fathoms	fathoms	fathoms
$\times 690 \cdot 688 = 29 \cdot 37$ the deficiency of 690·6 &c.			

fathoms fathoms fathoms

Therefore $690 \cdot 688 + 29 \cdot 37 = 720 \cdot 058$ altitude required.*

* The Altitude here found is about 4 feet more than Dr. Hutton gives.

(Page 256.)

Ex. 2. $41^\circ - 38^\circ = 3^\circ$ difference of temperature by the attached thermometer.

And $\frac{3}{9600} \times 29 \cdot 45 = \cdot 0092$ correctional altitude for 3° temper.

Add 29·45 observed height of the quicksilver.

Sum	29·4592	Base-alt. corrected for 41° temperature of the quicksilver.
-----	---------	--

Now the log. of 29·459 is 1·4692180

The log. of 26·82 is 1·4284588

<i>Difference</i>	·0407592
Multiply by	10000

PRODUCT	407·592 fathoms.
---------	------------------

(Key to Vol. II. page 256.)

Again,

$$\frac{31^{\circ} + 35^{\circ}}{2} = 33^{\circ} \text{ mean temperature by the detached therm.}$$

And

$$33^{\circ} - 31^{\circ} = 2^{\circ} \text{ above the temperature reduced to the formula.}$$

$$\text{But } \frac{2}{435} \times 407.592 = 1.87 \text{ the deficiency of } 407.5 \text{ \&c.}$$

Therefore the altitude required is 409.462 fathoms.*

* Which is nearly but not exactly the Answer given with the Question.

PRACTICAL QUESTIONS IN MENSURATION.

(Page 259.)

Quest. 1. ft. ft. sq. ft.
 $28 \times 20 = 560$ area of the 1st floor. And
 $2(14 \times 10) = 280$ area of the two less floors.

Difference 280 square feet.

Sum 840 square feet = 8.4 SQUARES.

Multiply by 45s.

PRODUCT £18.. 18

Therefore the *Diff.* is 280 sq. feet; and the *Amount* 18 guineas.

Quest. 2. sq. yd. sq. in. ft. in.
 $1 = 1296$, and $14 \text{ .. } 3 = 171$ inches in length.

But $\frac{1296}{171} = 7\frac{11}{19}$ inches in width. Ans.

Quest. 3. sq. yds.
 $114\frac{2}{3} = 1032$ square feet.

Now $\frac{1032}{28 \text{ ft. } b.} = 36\frac{6}{7}$ feet in length. Ans.

(Key to Vol. II. page 259.)

Quest. 4. $2 \begin{matrix} \text{in.} & \text{in.} \\ (7 \times 2\frac{1}{2}) \end{matrix} = 35$ square inches.

But $\frac{\begin{matrix} \text{sq. in.} \\ 35 \end{matrix}}{3 \text{ in. } l.} = 11\frac{2}{3}$ inches deep. Ans.

Quest. 5. $\begin{matrix} \text{sq. yd.} & \text{sq. in.} \\ 1 \end{matrix} = 1296$, and

As $\begin{matrix} d. & d. & \text{sq. in.} & \text{sq. in.} \\ 6 & ; & 38 & :: & 1296 & : & 8208 \end{matrix}$ internal area of the cistern.

Now $2 \begin{matrix} \text{in.} & \text{in.} & \text{sq. in.} \\ (102 \times 21) \end{matrix} = 4284$ internal area of both sides.

And $8208 - 4284 = 3924$ area of the bottom and two ends.

Also $\begin{matrix} \text{in.} & \text{in.} & \text{in.} \\ (2 \times 21) + 102 \end{matrix} = 144$ length of the bottom, and depth of both ends.

But $\frac{\begin{matrix} \text{sq. in.} \\ 3924 \end{matrix}}{144 \text{ in. } l.} = 27\frac{1}{4}$ inches wide. Ans.

Quest. 6. $\begin{matrix} \text{ft.} & \text{in.} & \text{ft.} & \text{in.} & \text{sq. ft.} \\ (47 \text{ .. } 9) \times (47 \text{ .. } 9) \end{matrix} = 2279\frac{1}{16}$ whole area.
Subtract $(47 \text{ ft. } 9 \text{ in.}) \times 4 \text{ ft.} = 191$ purbeck stone.

Divide by 9 sq. ft. Rem. $2088\frac{1}{16}$ paved with flint.

Quotient $232\frac{1}{44}$ square yards.
Multiply by $6d.$ per yard.

PRODUCT $\underline{\underline{\pounds 5 \text{ .. } 16 \text{ .. } 0\frac{1}{2}}}$. Ans.

Quest. 7. $\sqrt{\begin{matrix} \text{ft.} & \text{ft.} & \text{ft.} & \text{ft.} \\ (26\frac{2}{3} \times 26\frac{2}{3}) - (22 \times 22) \end{matrix}} = 15.07$ feet. And
 $\sqrt{\begin{matrix} \text{ft.} & \text{ft.} & \text{ft.} & \text{ft.} \\ (26\frac{2}{3} \times 26\frac{2}{3}) - (14 \times 14) \end{matrix}} = 22.69$ feet.

Sum $\underline{\underline{37.76}}$ feet nearly. Ans.

Quest. 8. $18d. = \pounds 0.075$; and,

As $\pounds 0.075 : \pounds 100 :: 1 \text{ sq. ft.} : 1333\frac{1}{3} \text{ sq. ft.}$ area of the court.

Also $\frac{1}{2} \times 88 \text{ ft.} = 44$ feet, half the base.

(Key to Vol. II. page 259.)

But $\frac{\overset{\text{sq. feet}}{1333\frac{1}{3}}}{44 \text{ ft.}} = 30.303$ feet, the perpendicular.

And $\sqrt{44^2 + 30.303^2} \text{ ft.} = \sqrt{1936 + 918.27} \text{ ft.} = 53.425 \text{ ft. } \frac{1}{2} \text{ sum.}$
 Multiply by $\frac{2}{2}$

PRODUCT and Answer 106.85 ft.

(Page 260.)

Quest. 9. $\sqrt{86^2 - 76^2} \text{ ft.} = 40.24922$ feet, height of the *lower* column above the STATUE. But

$\overset{\text{ft.}}{50} - \overset{\text{feet}}{40.24922} = 9.75078$ feet, height of the STATUE.

Hence the excess of the height of the *higher column* above that of the Statue is 54.24922 feet.

Now $\sqrt{97^2 - 54.24922^2} \text{ ft.} = 80.41157$ feet, horizontal distance of the higher column from the center of the figure's base.

And

$\overset{\text{feet}}{76} + \overset{\text{feet}}{80.41157} = 156.41157$ feet, distance between the COLUMNS.

Again,

$\overset{\text{ft.}}{64} - \overset{\text{ft.}}{50} = 14$ difference of altitude of the Columns.

Wherefore

$\sqrt{156.41157^2 + 14^2} \text{ ft.} = 157.03$ feet. Ans.

Quest. 10. $\frac{\overset{\text{ft.}}{16.5}}{2} = 8.25$ feet, circumference of the wheel.

And, As $\overset{(\text{c.})}{3.1416} : \overset{(\text{D.})}{1} :: \overset{(\text{Circ.})}{8.25 \text{ ft.}} : \overset{(\text{Di.})}{2.626 \text{ ft.}}$ Ans.

(Key to Vol. II. page 260.)

Quest. 11.* Because the circumferences of circles are to one another as their diameters, and *vice versâ* the diameters in the ratio of the circumferences; it follows (the circumference of the greater circle described being twice the circumference of the less) that the radius of the outer ring is double the radius of the inner. But the distance between the rings is 5 feet, wherefore the diameter of the outer ring is 20 feet.

(D.) (Circ.) (Di.) (Circ.)
And, As 1 : 3·14·16 :: 20 ft. : 62·832 ft. Ans.

* This Question, like many others in Dr. Hutton's Book, was first taken from a celebrated old Work, but in which the conditions are frequently expressed in rather clumsy, sometimes ungrammatical terms. Although the observation may not exactly apply to the present Example, yet the mention of the height of the wheels was superfluous, inasmuch as (unless it had been meant to be said that the one wheel revolved only once and the other twice, a thing impossible) the size of the wheels, if it did not exceed 10 feet, had no influence on the Answer; and then only because the revolutions or turns in the Question could not have been performed. It cannot be here objected that the wheels might have been of different heights, since two-wheeled vehicles of this description have never been in use.

Quest. 12. Because a guinea for a yard in length is 7s. per foot, and that 8d. = $\frac{2}{3}$ of a shilling, it is manifest that

As $\frac{2}{3}$ s. : 7s. that is, As 2s. : 21s.
:: the number of feet *in length* palisaded,
: the number of *square* feet in the area.

Hence, As the number of feet *in length* constituting a side of the Δ ,
: the number of *square* feet in the area paved,
:: $\frac{2}{3}$: 21 :: 2 : 63.

But in the table of regular polygons the area of an equilateral triangle is ·4330127 when the side is 1.

Now if x be substituted for the side of the triangle sought, whilst a represents the tabular area ·4330127, it will be

As $(a \times x^2) : x : 63 : 2$.
That is, $2ax^2 = 63x$, or $x = \frac{63}{2a}$.

Therefore

·8660254) 63· - - - - - (72·746 feet. Ans.

(Key to Vol. II. page 260.)

Quest. 13. $\sqrt{13^2(=AB^2)+31\cdot2^2(=BC^2)}=33\cdot8=AC.$

Wherefore in the triangle ACD are given the three sides to find the area; which (*page 29. vol. ii.*) is 207·322
 Add the area of ABC = $\frac{1}{2} \times 13 \times 31\cdot2 = 202\cdot8$

Sum and Answer 410·122

Quest. 14. $(24 \text{ .. } 8) \times (14 \text{ .. } 6) = 357\frac{2}{3}$ square feet.
 Multiply by 8 lb.

PRODUCT 2861 $\frac{1}{3}$ lb. of lead.


And As lb. lb. $112 : 2861\frac{1}{3} ::$ £·9 : £22 .. 19 .. 10 $\frac{1}{4}$. Ans.

Quest. 15.

sq. in.	in.	in.	breadth of the	1st cut.	inches left.
144 ÷ 27	=	5·3	breadth of the	1st cut.	52·6 length.
144 ÷ 52·6	=	2·734	- - - - -	2d cut.	24·266 breadth.
144 ÷ 24·266	=	5·934	- - - - -	3d cut.	46·732 length.
144 ÷ 46·732	=	3·0814	- - - - -	4th cut.	21·1846 breadth.
144 ÷ 21·1864	=	6·7974	- - - - -	5th cut.	39·9346 length.
144 ÷ 39·9346	=	3·6059	- - - - -	6th cut.	17·5787 breadth.
144 ÷ 17·5787	=	8·1917	- - - - -	7th cut.	31·7429 length.
144 ÷ 31·7429	=	4·5364	- - - - -	8th cut.	13·0423 breadth.
144 ÷ 13·0423	=	11·0409	- - - - -	9th cut.	20·702 length.
144 ÷ 20·702	=	6·956	- - - - -	10th cut.	6·0863 breadth.

There being now less than a square foot remaining, the dimensions are evidently 20·702 inches in *length*, and 6·0863 inches in *breadth*. Ans.

Quest. 16.



Construct any triangle ABC, obtuse angled at c; and let AB be supposed 40 poles; and AC 20 poles. It will be,

- As 400 poles half the product of AB X AC,
- : 160 poles or 1 acre,
- :: Radius,
- : Sin. 156° .. 25' the angle ACB.

(Key to Vol. II. page 260.)

And

As 60 poles, the sum of AB and AC,
 : 20 poles, AB - AC,
 :: Tang. $11^{\circ}..47'\frac{1}{2}$ half the supplement of the $\angle ACB$,
 : Tang. $3^{\circ}..59'$ the difference of the \angle les A and B.

Hence the angle opposite to the side AC* is $7^{\circ}..48'\frac{1}{2}$, and the angle BAC is $15^{\circ}..46'\frac{1}{2}$.

But As Radius AC,
 : Sin. $15^{\circ}..46'\frac{1}{2}$ the $\angle BAC$,
 :: 20 poles, the side AC,
 : 5.4344 poles, the perpendicular cp.


Now $\frac{5.4344 \text{ poles}}{2} = 2.7172$ half the perpendicular cp.

And $\frac{160 \text{ sq. p.}}{2.7172 p.} = 58.884$ &c. poles AB the third side sought.

* The side BC being by the construction greater than AC.

Or,

If 40 poles be the side opposite to the obtuse angle, then as in the following diagram, AC being 20 poles, it will be $\frac{160 \text{ sq. p.}}{20 p.} \times 2 = 16$



poles BE a perpendicular from B meeting AC produced. Wherefore [Eucl. i. 47.] $AE = 36.66$ poles. And by subtraction, $CE = 16.66$ poles; also, by the last-mentioned Theorem, $CB = 23.099$ poles, the third side of the triangle.

Otherwise, without producing AC.

(1 acre) (AB)
 $(160 \text{ sq. p.} \div 40 \text{ poles}) \times 2 = 8 = CD$ a perpendicular from c upon AB.

And

$AC^2 - CD^2 = AD^2$ Hence $AD = \sqrt{20^2 - 8^2} = \sqrt{336} = 18.3303$ poles.

But 40 poles AB, - 18.3303 poles AD, = 21.6696 poles DB.

(Key to Vol. II. page 260.)

But $DB^2 + DC^2 = BC^2$. Or in numbers,

$$BC = \sqrt{21.6696^2 + 8^2} = 23.09 \text{ poles, as before.}$$

Consequently 58.884 poles, and 23.09 poles. Ans.

Courses (br.)

Quest. 17. $38 \div 4 = 9\frac{1}{2}$ feet, the height of the triangular gable.

Also $\frac{1}{3} \times 40 \text{ ft.} = 13\frac{1}{3}$ feet, the height of each of the three portions of wall to the eaves.

It is, therefore,

ft.	in.	ft.	in.	sq. ft.	pts.	sq. ft.	pts.
(24 .. 6)	×	(13 .. 4)	=	326 .. 8	two bricks thick - -	=	1306 .. 8 half brick.
(24 .. 6)	×	(13 .. 4)	=	326 .. 8	brick and half thick	=	980 .. 0 - - - - -
(24 .. 6)	×	(13 .. 4)	=	326 .. 8	brick thick - - - - -	=	653 .. 4 - - - - -
(24 .. 6)	×	(4 .. 9)	=	116 .. 4 $\frac{1}{2}$	half brick thick - -	=	116 .. 4 $\frac{1}{2}$ - - - - -

Divide by 3) 3056 .. 4 $\frac{1}{2}$ - - - - -

Standard thickness 1018.792 of work.

But $1018.792 \div 272.25^* = 3.742$ square rods.

Multiply by £5.5 given price per rod.

PRODUCT £20 .. 11 .. 7.44. Ans.

* See page 82, vol. ii. of the Course of Mathematics.

(Page 261.)

Quest. 18. 500 feet = 6000 inches, which \div by 10 = 600 the number of bricks, end to end, in a half-brick course.

ft. (bricks)
But $10 \times 4 \times 600 = 24000$ bricks in the wall, $\frac{1}{2}$ brick thick.
Multiply by 3

PRODUCT 72000 bricks. Ans.

Quest. 19. in. in. in.
 $10 \times 5 \times 3 = 150$ cubic inches in a brick.

(Key to Vol. II. page 261.)

$$\text{And } \frac{\text{ft.} \quad \text{ft.} \quad \text{ft.}}{3} 100 \times 100 \times 100 \text{ (cub. in. in 1 ft.)} \times 1728 = 576000000 \text{ solid in. of PYRAMID.}$$

$$\text{But } \frac{\text{cubic inches} \quad \text{cub. in.}}{150} 576000000 \div 150 = 3840000 \text{ bricks. Ans.}$$

$$\text{Quest. 20.* } \frac{\text{ft.} \quad \text{ft.}}{2} \times 12 \times 16 = 96 \text{ sq. feet, area of the triangle.}$$

$$\text{And, As } \frac{\text{sq. ft.} \quad \text{sq. ft.} \quad \text{ft.} \quad \text{ft.}}{96 : 24 :: 12^2 : 6^2}.$$

Wherefore the base of the less triangle is 6 feet.†

$$\text{Again, As } \frac{\text{sq. ft.} \quad \text{sq. ft.} \quad \text{ft.} \quad \text{ft.}}{96 : 24 \therefore 4 : 1 :: 16^2 : 8^2}.$$

Consequently the perpendicular sought is 8 feet.

$$\text{Lastly, } \sqrt{6^2 + 8^2} \text{ ft.} = \sqrt{100} \text{ ft.} = 10 \text{ feet the hypotenuse.}$$

* This Question is very *curiously* worded!

† The base being 6 feet or half the base of the greater triangle, it is evident that the perpendicular must be 8 feet or half the perpendicular given; and the hypotenuse of the less triangle half the hypotenuse of the greater.

$$\text{Quest. 21. } \frac{\text{in.} \quad \text{in.}}{14 \times 2} = 28 = 2\frac{1}{3} \text{ feet. And}$$

$$840 \text{ links} = 554.4 \text{ feet. Also}$$

$$612 \text{ links} = 403.92 \text{ feet. Therefore,}$$

556.73 feet and 406.253 feet are the diameters of the outer ellipse.

$$\text{But } \frac{\text{ft.} \quad \text{ft.}}{556.73 \times 406.253 \times .7854} = 177637.66 \text{ square feet area.}$$

$$\text{And } \frac{\text{ft.} \quad \text{ft.}}{554.4 \times 403.92 \times .7854} = 175877.17 \text{ square feet area.}$$

Difference 1760.49 built upon.

$$\text{Again } \frac{\text{sq. ft.} \quad \text{sq. ft.} \quad \text{acres} \quad \text{A.} \quad \text{R.} \quad \text{P.}}{175877.17 \div 43560^*} = 4.0375 = 4 \text{ .. } 0 \text{ .. } 6 \text{ inclosed by the walls.}$$

* The number of square feet in an acre.

$$\text{Quest. 22. } \frac{\text{in.} \quad \text{in.} \quad \text{in.}}{7 \times 7} = 49 \text{ the sqr. of the diameter of the 1st pillar.}$$

$$\text{And, As } \frac{\text{cub. ft.} \quad \text{cub. ft. (1st D}^2\text{.)} \quad \text{(2d D}^2\text{.)}}{4 : 40 :: 49 \text{ in.} : 490 \text{ inches.}}$$

$$\text{But } \frac{\text{sq. in.}}{\sqrt{490}} = 22.136 \text{ inches. Ans.}$$

(Key to Vol. II. page 261.)

Quest. 23. $\frac{\text{sq. yds.}}{4840} \div 2 = 2420 = \frac{1}{2} \text{ acre.}$

And $\frac{\text{sq. yds.}}{2420} \div .7854 = 3081.23$, square of the diam. of the POND.

Lastly, $\frac{\text{sq. yds.}}{\frac{1}{2}\sqrt{3081.23}} = 27\frac{3}{4}$ yards, *the radius required.*

Qu. 24. $\frac{\text{feet}}{22.75} + (\frac{1}{2} \times 22.75) + (1.66 \text{ for the eave boards}) = 35.791.$

(*Breadth and half*) (*Length*) $\frac{\text{sq. ft.}}{\text{And } 35.791 \text{ ft.} \times 32.75 \text{ ft.} \div 100 = 11.7217708 \text{ SQUARES.}}$
 Multiply by 15 shillings.

PRODUCT $\underline{\underline{\text{£}8..15..9\frac{1}{2} \text{ nearly. Ans.}}}$

Quest. 25. $\frac{\text{in. in.}}{9 \times 9} = 81$ square of the circumf. of the 1st cable.

Also $12 \times 12 = 144$ square of the circumf. of the 2d cable.

Now, As $\frac{\text{sq. in.}}{81} : \frac{\text{sq. in.}}{144} :: \frac{\text{lb.}}{22} : 39\frac{1}{5}$ weight per yard.
 Multiply by 2 yards in a fathom.

PRODUCT $\underline{\underline{78\frac{2}{5} \text{ lb. Ans.}}}$

Quest. 26.* $\frac{\text{in. in. in.}}{(40 \times 2) + 26} = 106$ breadth of the bottom and 2 sides spread out. Again,

$\frac{\text{in. in. sq. in.}}{106 \times 74} = 7844$ area of the bottom and sides.

Also $\frac{\text{in. in.}}{40 \times (26 \times 2)} = 2080$ area of both ends.

So $26 \times (16 \times 3) = 1248$ area of the three stays.

Divide the Sum by 144) $11172 (77.58\dot{3}$ square feet in the WHOLE.

Multiply by $.25 \text{ cwt. per sq. foot.}$

PRODUCT $19.39583 \text{ cwt. of lead.}$

Multiply by $\text{£}1.1 \text{ per cwt. for work, \&c.}$

PRODUCT $\text{£}21..6..8\frac{1}{2}$

Subtract $3..6$ balance per Question.

Remainder $\underline{\underline{\text{£}21..3..2\frac{1}{2} \text{ amount of paving.}}}$

* For what purpose is the Question encumbered with the mention of sheet lead $\frac{1}{10}$ of an inch in thickness weighing 5.899 lb. "the square foot?"

(Key to Vol. II. page 261.)


Again,
 $\text{£}21 \text{ .. } 3 \text{ .. } 2\frac{1}{2} \div 7 = \text{£} \frac{21 \cdot 16041}{\cdot 02916} = 725 \cdot 5$ a number equal to the
 number of square feet in the SHOP.

sq. ft. ft. ft. in.
 But, $725 \cdot 5 \div 22 \cdot 83 = 32 \text{ .. } 0\frac{3}{4}$ the length required.

(Page 262.)

Construction.

Quest. 27. Draw any indefinite right line, and take in it KH corresponding to 30 on some convenient scale of equal parts; then by the same scale make KC and HI, each equal to 25. And about K at the distance KC, and H at the distance HI, describe the circles DAB, BAE, intersecting one another in the points A and B. If the straight line AB be drawn cutting KH in o, it will be, (because KI=CH=5, and IO=OC=10)



(Alt.) (Di.) (Quot.)

$10 \div 50 = \cdot 2$ The number wherewith to enter the Table.

But in the Table of Circular segments the number answering to $\cdot 2$ is $\cdot 111823$. Therefore

(Tab. area) (Di.) (Di.)
 $(\cdot 111823 \times 50 \times 50) \times 2 = 559 \cdot 119$. Ans.


Quest. 28. in. in. (ft.) 12th pts. of a cub. ft.
 $1\frac{1}{4} \times 1\frac{1}{4} \times 20 = 31 \cdot 25$
 $\frac{7}{8} \times \frac{7}{8} \times 50 = 38 \cdot 28125$

Now, As 31·25 :: 38·28125 :: 1120 lb. : 1372 wt. of the 50 proposed.
 Multiply by $3\frac{1}{2}d.$ the price per lb.

PRODUCT $\text{£}20 \text{ .. } 0 \text{ .. } 2$. Ans.

(Key to Vol. II. page 262.)

Quest 29. Describe an equilateral triangle as DCE, and bisect the base in F. Also produce DE both ways to A and B; and about F at the distance FC describe the semicircle ACB.



Supposing now the area of DCE to be 100, the side of the triangle is $\sqrt{230.94} = 15.17$ &c. because

$$\begin{array}{cccc} \text{(Tab. area)} & \text{(Giv. area)} & \text{(Side}^2) & \text{(Side}^2) \\ \text{As } .4330127 & : 100 & :: 1 & : 230.94. \end{array}$$

But $CF = \text{Radius} = \sqrt{230.94 - \frac{1}{4}(230.94)}$ because $DF = FE = \frac{1}{2}DC$.

Therefore $2\sqrt{230.94}$ &c. = 57.73 &c. = 26.32148 &c. Ans.

Quest. 30. 14 lb. = 224 oz. And

$\begin{array}{cccc} \text{oz.} & \text{oz.} & \text{cub. in.} & \text{cub. in.} \\ \text{As } 11325 & : 224 & :: 1728 & : 34.17854 \text{ solidity of 1 yard of pipe.} \end{array}$

Again, $\begin{array}{cccc} \text{in.} & \text{in.} & \text{(Const. dec.) in.} & \\ \text{As } 1\frac{1}{4} \times 1\frac{1}{4} \times .7854 \times 36 & = & 44.17875 & \text{cavity in 1 yd. of pipe.} \end{array}$

Now, $\begin{array}{cccc} \text{cub. in.} & \text{cubic inches} & \text{inches} & \text{sq. inches} \\ \text{As } 44.17875 : (44.17875 + 34.17854) & :: & (1\frac{1}{4} \times 1\frac{1}{4}) & : 2.771315 \end{array}$
the square of the diameter of the external circumference of the pipe.

$$\text{But } \frac{\sqrt{2.771315} - 1.25}{2} = .20736 \text{ \&c. thickness required. Ans.}$$

Quest. 31. $\begin{array}{cc} \text{s.} & \text{d.} \\ 2 \text{ .. } 4 & = \mathcal{L}.116. \text{ And} \end{array}$

$\begin{array}{cccc} & \text{sq. ft.} & \text{sq. ft.} & \\ \text{As } \mathcal{L}.116 : \mathcal{L}10 & :: 1 & : 85.7143 \text{ the area of the semi-circle.} \\ & \text{Multiply by} & 2 & \end{array}$

PRODUCT 171.4286 area of the whole circle.

$\begin{array}{cccc} & \text{sq. feet} & \text{(Const. dec.)} & \text{sq. feet.} \\ \text{But } 171.4286 \div .7854 & = & 218.269, & \text{square of the diameter.} \end{array}$

Lastly, $\begin{array}{cc} \text{sq. feet} & \\ \sqrt{218.269} & = 14.7739 \text{ feet. Ans.} \end{array}$

(Key to Vol. II. page 262.)

Quest. 32. $(Di.) (Area) (Di.) (Area)$
As 1 : .78539816 :: $\frac{1}{3}$: .26179939.*

But the versed sine of .26179939 is .36753395.

Now the semichord is a mean proportional between the segments of the diameter ; [*Euclid*, iii. 35.]. Therefore

$$2\sqrt{(1 - .36753395) \times .36753395} = .96426162 \text{ chord to diameter 1.}$$

And

$(Di.) (Di.) (Chord) (Chord)$
As 1 : 289 :: .96426162 : 278.6716 *the chord required.*

* More briefly $.7854 \div 3 = .2618$ the area, *nearly*, of one third part of a circle whose diameter is Unity.

Quest. 33. $\begin{matrix} \text{sq. ft.} & \text{sq. inches} & & \text{in.} \\ 64.3 = 9259.2, & \text{which} \times^{\text{ed}} & \text{by} & \frac{1}{8} = 1157.4 \text{ cubic inches.} \end{matrix}$

$\begin{matrix} \text{cub. in.} & \text{oz.} & \text{ounces} & \text{cwt.} \\ \text{And } 1157.4 \times 6\frac{1}{2} = 7586.757 = 4.233 \text{ of lead in the cistern.} \end{matrix}$

But, As $\begin{matrix} \text{cwt.} & \text{cwt.} & & \text{s.} & \text{d.} \\ 19\frac{1}{2} : 4.233 :: \text{£}21 : \text{£}4 \text{ .. } 11 \text{ .. } 2\frac{1}{2} \text{ nearly.} \end{matrix}$ Ans.

Quest. 34. Because in any sphere the square of the diameter \times^{ed} by 3.1416 = the SUPERFICIES, and the cube of the diameter \times^{ed} by .5236 = the SOLIDITY, it follows that the Answer is 6, or $3.1416 \div .5236$.

For if x be put for the diameter of the globe, then

$$\begin{array}{l|l} \text{By the Question.} & x^3 \times .5236 = x^2 \times 3.1416 \\ \text{Divide by } x^2. & x \times .5236 = 1 \times 3.1416 \\ \text{Divide by } .5236. & x = \frac{3.1416}{.5236} = 6. \text{ Ans.} \end{array}$$

Quest. 35. $\begin{matrix} \text{in.} & \text{in.} & \text{sq. in.} \\ 22\frac{1}{2} \times 22\frac{1}{2} = 506.25. \end{matrix}$ And

$$\begin{matrix} \text{in.} & \text{in.} & \text{sq. in.} \\ (22\frac{1}{2} \times 2) \times (22\frac{1}{2} \times 2) = 2025. \end{matrix}$$

But, As $\begin{matrix} \text{sq. in.} & \text{sq. in.} & \text{bu.} & \text{bu.} \\ 506.25 : 2025 :: 3 : 12. \end{matrix}$ Ans.

(Page 263.)

Quest. 36. The diameter of the inner circle being 42 inches, and the diameter of the outer circle 56.5 inches, it is

$$\begin{matrix} \text{in.} & \text{in.} & (\text{Const. dec.}) & \text{sq. in.} \\ (56.5^2 - 42^2) \times .7854 = 1121.7475 \text{ area of the curb.} \end{matrix}$$


(Key to Vol. II. page 263.)

sq. in.
 Now $1121.7475 = 7.789$ square feet.
 Multiply by $8d.$ per square foot.

PRODUCT $\underline{\underline{\pounds 0 .. 5 .. 2\frac{1}{4}}}$ Ans.

Quest. 37. $\begin{matrix} \text{ft.} & \text{ft.} & \text{sq. ft.} \\ 100 \times 80 \div 2 = 4000 \end{matrix}$ area of
 the walk proposed.

And if x be put for the breadth of
 the walk, it will be



By the Question. $180x - x^2 = 4000$
 Change the signs, complete, &c. $x = 90 \pm 64.0312$
 Wherefore 154.0312 ft. and 25.9688 ft. are
 the 2 values of x in the equation; but the first being inappli-
 cable, the second or 25.9688 ft. is the Answer.

Quest. 38. $\begin{matrix} \text{ft.} & \text{ft.} & \text{sq. ft.} & \text{sq. ft.} & \text{sq. ft.} & \text{ft.} \\ \sqrt{(26^2 - 10^2)} = \sqrt{(676 - 100)} = \sqrt{576} = 24 \end{matrix}$ the part
 left standing.

And $24 + 26 = 50$ the height required.

Quest. 39. $\begin{matrix} \text{in.} & \text{in.} & \text{sq. in.} \\ 60 \times 60 \times .7854 = 2827.44 \end{matrix}$ area of either circular side
 of the stone; which, divided by $7 = 403.92$ each man's share
 of the side.

Now $403.92 \div .7854 = 514.2856$ square of the diameter for
 the last man.

And

inches
 $\sqrt{3600 - (514.2856 \times 1)} = 55.5492$ diameter left by the 1st man.
 $\sqrt{3600 - (514.2856 \times 2)} = 50.7092$ - - - - - 2d man.
 $\sqrt{3600 - (514.2856 \times 3)} = 45.3557$ - - - - - 3d man:
 $\sqrt{3600 - (514.2856 \times 4)} = 39.2792$ - - - - - 4th man.
 $\sqrt{3600 - (514.2856 \times 5)} = 32.0715$ - - - - - 5th man.
 $\sqrt{3600 - (514.2856 \times 6)} = 22.6778$ - - - - - 6th man.

(Key to Vol. II. page 263.)

Wherefore

$$\left. \begin{array}{l} \text{inches} \quad \text{inches} \quad \text{inches} \quad \text{inches} \\ 1st. 4.4508 - 2nd. 4.84 - - 3rd. 5.3535 - 4th 6.0765 \\ 5th. 7.2079 - 6th. 9.3935 - 7th. 22.6778 - - - - - \end{array} \right\} \text{Ans.}$$

$$\text{Quest. 40. } 16\frac{1}{2} \text{ ft.} \times 16\frac{1}{2} \text{ ft.} = 272\frac{1}{4} \text{ sq. ft. area of the old KILN.}$$

$$\text{And } \sqrt{3 \times 272\frac{1}{4}} \text{ sq. ft.} = \sqrt{816\frac{3}{4}} \text{ sq. ft.} = 28 \text{ ft. } 7 \text{ in. Ans.}$$

$$\text{Quest. 41. } 3 \text{ in.} \times 3 \text{ in.} \times 3 \text{ in.} = 27 \text{ cubic inches in a 3-inch cube.}$$

$$\text{Also } 12 \times 12 \times 12 = 1728 \text{ cubic inches in a 12-inch cube.}$$

$$\text{But } 1728 \div 27 = 64 \text{ cubes. Ans.}$$

$$\text{Quest. 42. } 1 \text{ acre} = 4840 \text{ square yards. And}$$

$$\text{As } 355 : 452 :: 4840 : 6162.4788 \text{ square of the diam. sought.}$$

$$\text{But } \sqrt{\frac{6162.4788 \text{ sq. yds.}}{2}} = 39\frac{1}{4} \text{ yards. Ans.}$$

$$\text{Quest. 43. As } 22 : 7 :: 64 \text{ ft.} : 20.36 \text{ ft. diameter of the base of the CONE.}$$

Hence the radius of the base is 10.18 ft. *nearly*.

$$\text{And } \sqrt{118^2 + 10.18^2} \text{ ft.} = 118.43 \text{ ft. the slant side of the cone.}$$

$$\text{Again } \frac{118.43 \text{ ft.} \times 64 \text{ ft.}}{2} = 3789.76 = 421.08 \text{ square yards.}$$

Multiply by 8d. per square yard.

$$\text{PRODUCT } \underline{\underline{\pounds 14 \text{ .. } 0 \text{ .. } 8.64. \text{ Ans.}}}$$

$$\text{Quest. 44. } 18\frac{1}{2} \text{ in.} \times 18\frac{1}{2} \text{ in.} \times 8 \text{ in. (Const. dec.)} = 2150.4252 \text{ in the BUSHEL.}$$

$$\text{And } \sqrt{2150.425 \div 7.5 \div 7854} = 19.1067 \text{ inches. Ans.}$$

Otherwise.

$$\sqrt{\left(\frac{18.5 \text{ in.} \times 18.5 \text{ in.} \times 8 \text{ in.}}{7.5 \text{ in.}} \right)} = 19.1067 \text{ inches as before. Ans.}$$

(Key to Vol. II. page 263.)

Quest. 45. $\begin{matrix} \text{in.} & \text{in.} & (\text{Const. dec.}) & \text{sq. inches} \\ 72 \times 72 \times 3 \cdot 1416 = 16286 \cdot 0544 & \text{surface of the BALL.} \\ \text{Multiply by} & & 3\frac{1}{2}d. & \text{per sq. inch.} \\ \hline \text{PRODUCT } \mathcal{L}237 \cdot 10 \cdot 1. & \text{Ans.} \end{matrix}$

Quest. 46. $\frac{\begin{matrix} \text{ft.} & \text{ft.} \\ 4 & - & 1 \cdot 5 \end{matrix}}{2} = 1 \cdot 25. \text{ And}$

$\begin{matrix} \text{ft.} & \text{ft.} & \text{ft.} & \text{ft.} & \text{sq. feet.} \\ (8 \times 8) - (1 \cdot 25 \times 1 \cdot 25) = 62 \cdot 4375 & \text{sqr. of the altitude.} \end{matrix}$

Now $\sqrt{\begin{matrix} \text{sq. feet} & \text{ft.} \\ 62 \cdot 4375 \end{matrix}} = 7 \cdot 9$ the perpendicular height.

Again,

$\begin{matrix} \text{ft.} & \text{ft.} & \text{ft.} & \text{ft.} & \text{ft.} & \text{ft.} & \text{sq. ft.} \\ (4 \times 4) + (1 \cdot 5 \times 1 \cdot 5) + (4 \times 1 \cdot 5) = 24 \cdot 25 & \text{sum of the 3 products.} \\ \text{Multiply by} & & & & & & \cdot 7854 \text{ constant decimal.} \end{matrix}$

PRODUCT 19·04595 of which one-third is

$\begin{matrix} \text{sq. feet} & \text{ft.} \\ 6 \cdot 34865; \text{ this } \times^{\text{cd}} \text{ by } 7 \cdot 9 = 50 \cdot 154335 & \text{solid feet of MARBLE.} \\ \text{Multiply by} & & 12s. & \text{per solid foot.} \end{matrix}$

PRODUCT $\mathcal{L}30 \cdot 1 \cdot 10\frac{1}{4}$. Ans.

(Page 264.)

Quest. 47.

As $\begin{matrix} \text{in.} & \text{in.} & \text{in.} \\ \sqrt[3]{3} : \sqrt[3]{1} :: 20 : 20\sqrt[3]{\frac{1}{3}} = 13 \cdot 867226 & \text{alt. of the uppermost part.} \\ \sqrt[3]{3} : \sqrt[3]{2} :: 20 : 20\sqrt[3]{\frac{2}{3}} = 17 \cdot 471606 & \text{do. and middle part.} \end{matrix}$

$\text{Difference } 3 \cdot 604380 \text{ - - - middle part.}$

Lastly $\begin{matrix} \text{in.} & \text{inches} \\ 20 - 17 \cdot 471606 = 2 \cdot 528394 & \text{- - - lowest part.} \end{matrix}$

Consequently $\begin{matrix} \text{in.} & \text{in.} & \text{in.} \\ 2 \cdot 528394, 3 \cdot 60438, * & \text{and } 13 \cdot 867226 & \text{respectively.} \\ & & \text{Ans.} \end{matrix}$

* Differing in the 3d place of decimals from Dr. Hutton.

(Key to Vol. II, page 264.)

Quest. 48. $300 \times 200 \times 1 = 60000$ cubic feet of earth.


And $2 \left\{ \begin{array}{l} \text{ft.} \quad \text{ft.} \quad \text{ft.} \quad \text{ft.} \\ (300 \times 8) + (200 \times 8) \end{array} \right\} + 4(8 \times 8) = 8256$ area of the surface of the ditch.

But $60000 \div 8256 = 7 \frac{23}{80}$ feet. Ans.

* The reason will be apparent on constructing the figure.

Quest. 49.

Let HBIG (in the marginal diagram) represent the earth; and let HI be the true horizon of the observer at A; also F, D two points in the visual horizon, and c the earth's center.



Then, because spherical curve surfaces are to one another as their altitudes, † if BE be taken $= \frac{1}{3}$ of BG: it will be

As CE : CB :: CB : CA, that is, As $\frac{2}{3}$ CB : CB :: CB : $\frac{3}{2}$ CB.

Consequently AB = BG, or in other words, the altitude of the observer above the earth's surface is equal to the earth's diameter. Ans.

NOTE.—Should the Student find any difficulty in tracing the first proportion, he has only to remember that CB is supposed equal to CD, and that if D be connected to c and E by right lines, the triangles ACD and DCE are equiangular; hence, As CE : CD = CB :: CD = CB : CA.

† Cor. ii. page 50. vol. ii.

Quest. 50. $\frac{1}{40}$ in. \times $\frac{1}{40}$ in. = $\frac{1}{1600}$ sq. inch. square of the diameter of the wire.

And $1728 \div \frac{1}{1600}$ sq. in. \times .785398 = 3520252.69329 inches.

Lastly, $3520252.69 = 97784.797 = 55 \text{ .. } 984.797$. Ans.

(Key to Vol. II. page 264.)

Quest. 51. The weight of a CUBIC INCH of cast iron is ^{oz.} 4.0509.

And As ^{oz.} 4.0509 : (^{oz.} 16 × 24) :: 1 : $\frac{128}{1.3503}$ the solidity of the ball.

Hence the diameter of the ball is 5.547 inches.

Add for windage .1 inch.

Sum 5.647 inches. Ans.

Otherwise.

The diameter of a 24lb. shot, in the TABLE, is 5.5469 inches.

To which add $\frac{1}{10}$ inch for windage .1

Sum and Answer 5.6469 in. as before.

	lb.	lb.	in.	inches	inches
Quest. 52. As	9	: 1	:: 4 ³	: 1.9230 ³ diameter.	1.9622 caliber.
	9	: 2	:: 4 ³	: 2.4228 ³ - - - - -	2.4723 - - - - -
	9	: 3	:: 4 ³	: 2.7734 ³ - - - - -	2.8301 - - - - -
	9	: 4	:: 4 ³	: 3.0526 ³ - - - - -	3.1149 - - - - -
	9	: 6	:: 4 ³	: 3.4943 ³ - - - - -	3.5656 - - - - -
	9	: 9	:: 4 ³	: 4.0000 ³ - - - - -	4.0816 - - - - -
	9	: 12	:: 4 ³	: 4.4026 ³ - - - - -	4.4924 - - - - -
	9	: 18	:: 4 ³	: 5.0397 ³ - - - - -	5.1425 - - - - -
	9	: 24	:: 4 ³	: 5.5469 ³ - - - - -	5.6601 - - - - -
	9	: 32	:: 4 ³	: 6.1051 ³ - - - - -	6.2297 - - - - -
	9	: 36	:: 4 ³	: 6.3496 ³ - - - - -	6.4792 - - - - -
	9	: 42	:: 4 ³	: 6.6844 ³ - - - - -	6.8208 - - - - -

Whereby the Answer is obvious.

(Page 265.)

Quest. 53. For a ten-inch Mortar (the process for the other calibers being the same.)

Mortar 10 inches, { windage $\frac{1}{60}$ of 10 inches = $\frac{1}{6}$ inch.
 hollow of the shell $\frac{7}{10}$ of 10 in. = 7 inches.

And 10 in. - $\frac{1}{6}$ inch = $9\frac{5}{6}$ inches, external diameter of the Shell.

(Key to Vol. II. page 265.)

Now $\left\{ \begin{array}{l} \text{cube of the external diameter is } 950.828 \text{ inches.} \\ \text{cube of the internal diameter is } 343.000 \text{ inches.} \end{array} \right.$

Difference 607.828 inches.

But $\frac{9^*}{64} \times 607.828 \text{ inches} = 85.476 \text{ lb. weight of the shell.}$

(*Int. di³.*) cub. in.†

And $343 \div 57.3 = 5.986 \text{ lb. of gunpowder requisite.}$


Proceeding in the same manner with the other given calibers, the following results are obtained.

Caliber	Diameter	Weight	Powder
in. 4.6 - - -	in. 4.523 - - -	lb. 8.320 - - -	lb. 0.583
5.8 - - -	5.703 - - -	16.677 - - -	1.168
8. - - -	7.867 - - -	43.764 - - -	3.065
13. - - -	12.783 - - -	187.791 - - -	13.151.

* See page 270, vol. ii. of the COURSE.

† Page 271 same volume.

Quest. 54. Let ABC represent the conical glass, F the center of the sphere, and FD a radius at right angles to BC the slant side of the cone; if a circle be described about F at the distance FD it will represent the globe *in plano*: H, therefore, is the lowest point of the sphere, and CHF are in the same straight line, which, if produced to G, will bisect AB.



By the Question $AG = 2.5$ inches, and $GC = 6$ inches.

But $\sqrt{AG^2 + GC^2} = AC$; that is, $AC = \sqrt{42.25} = 6.5$ inches.

Now, (*because of similar triangles*) it is,

in. in. in. in.
As $2.5 [AG] : 6.5 [AC] :: 2 [FD] : 5.2 [FC]$.


Consequently $FG = .8$ inch, and $GH = 2.8$ inches.

Again, (*page 51. vol. ii. Prob. ix.*) the solidity of the segment immersed is $\left\{ \begin{array}{l} \text{in.} \quad \text{in.} \\ (4 \times 3) - (2.8 \times 2) \end{array} \right\} \times 2.8 \times 2.8 \times .5236 = 26.272$ cubic inches.

cub. in. cub. in. pts.
Lastly, As $282 : 26.272 :: 8 : \frac{3}{4}$ of a pint *nearly*. Ans.

(Key to Vol. II. page 265.)

Quest. 55. Let ro be the surface of the water round the ball, the cone being perfectly upright; and let HKI be a horizontal tangent at K to the sphere immersed; FAG a line parallel to HI from G a point of contact of the globe and cone, D the center of the sphere, and DG a radius at right angles to BC the slant side of the glass. Join AB , aH , ai ; and from c draw CE perpendicular to AB ; CE passes through the points K , D , and bisects AB in E .



By the Question $AB=5$ inches, $EC=6$ inches, and DG or $DK=2$ inches; also, by the last example, the slant side $BC=6.5$ inches, and $DC=5.2$ inches. But,

$$\text{As } 6.5 \text{ [BC]} : 6 \text{ [EC]} :: 5.2 \text{ [CD]} : 4.8 \text{ [GC]}.$$

Likewise

$$\text{As } 5.2 \text{ [CD]} : 4.8 \text{ [GC]} :: 4.8 \text{ [GC]} : 4.43077 \text{ [ac]}.$$

Consequently,

By subtraction $CK=3.2$ inches, and $aK=1.23077$ inches.

And

$$\text{As } 6 \text{ [CE]} : 3.2 \text{ [CK]} :: 5 \text{ [AB]} : 2.666 \text{ [HI]}.$$

Now the unguia noG above the level ro is equal to the cone abc above the same level; wherefore, if Q be put for the DIAMETER in inches, and R for the ALTITUDE in inches of a cone equal to $\frac{1}{3}$ the given cone; also s for the solidity in inches, of the cone abc ; it will be

$$\frac{HI^2 \times .7854 \times ac}{3} - \frac{Q^2 \times .7854 \times R}{3} = s.$$

That is, substituting numeral values,

$$\frac{1}{3}(2.666^2 \times .7854 \times 4.43077) - \frac{1}{3}(2.924^2 \times .7854 \times 3.509), * \text{ or}$$

cub. in. cub. in. cub. in.

$8.2487 - 7.8543 = .3944$ the content of the cone abc , or of the unguia gno . But,

$$\text{As Cone } aHI : \text{Cone } abc :: aK^3 : am^3.$$


Hence $am = .684$ of an inch.

$$\text{Lastly, } 1.23077 \text{ in.} - .684 \text{ in.} = .54677 \text{ in.} \text{ Ans.}$$

* More correctly 3.508821 inches.

(Key to Vol. II. page 265.)

Quest. 56. Let ABC (as in the two preceding Examples) represent the conical glass, rao the level of the water before the sphere is immersed, and FKG its surface after the ball has sunk; also let D be the center of the sphere, Dl a radius at right angles to BC , and HbI a line parallel to FG or ro from the point of contact I of the sphere and glass.



Then $ac=3.508821$ inches. [See the last Question.]
 $ro=2.924$ and $ao=1.462$ inches.

Also the solidity of the cone cro is 7.854 cubic inches.*

Now, As $ca : ao :: ib : bD$. Or, in numbers,

As $3.508821 : 1.462 :: 1 : .41666$. That is, bI is to bD in the ratio of unity to $.41666$. Again,

As $db : bI : ib : bc$. Consequently,

$bc=2.4$ inches, and by addition, $DC=2.81666$ inches.

Hence, (still considering bI as unity,) $Dl=1.08333$ inches, $CK=3.9$ inches; and $KG=1.625$ inches, which doubled $=3.25$ inches $=FG$. And

$\frac{(FG) (Const. dec.) (CK)}{3} = \frac{3.25^2 \times .7854 \times 3.9}{3} = 10.78454$ the solidity of the cone CFG .

Likewise $(2Dl)^3 \times .5236 =$ the solidity of the immersed globe $= (2\frac{1}{8} \text{ in.})^3 \times .5236 = 5.325682$ cubic inches. Now, by the Question, the solidity of the sphere immersed is equal to the solid content of the conical frustum $FROG$. Wherefore, by subtracting this frustum from the cone CFG (above found to be 10.78454 cubic inches when Dl is 1 inch) there remain 5.45886 cubic inches for the cone cro ; but it is known to contain 7.854 cubic inches.

By proportion, then,

As $5.45886 : 7.854 :: 10.78454 : 15.5163$ cubic inches, the true content of the cone CFG .

Subtract 7.854 cub. in. (the cone cro).

Remainder 7.6623 solidity of the ball.

* Because equal to $\frac{1}{3}$ the whole cone ABC ; that is, equal to $\frac{1}{3} (5^2 \times .7854 \times 2 \text{ in.}) = 7.854$ cubic inches.

(Key to Vol. II. page 265.)

Lastly $\sqrt[3]{\frac{7.6623}{.5236}}$ inches = 2.445981 inches. Ans.

Quest. 57. As $\begin{matrix} (Cir.) & (mil. Cir.) & (D) & (mil. Di.) \\ 3.1416 & : & 25000 & :: 1 : 7957.74 \end{matrix}$ the diameter of the earth. And


As $\begin{matrix} \text{miles} & \text{miles} & (Cir.) & (Arc.) \\ 25000 & : & 49.5933 & :: 360^\circ : 0^\circ .. 42'.85 \end{matrix}$ the angle of Oxford and London, at the center of the Earth.

Wherefore,

As Radius - - - - - Log. ar. co. - 10. _____
 : Sec. $0^\circ .. 42' .. 51''$ - - - - - Log. 10.0000337
 :: 3978.87 miles (*the Earth's radius*) - - - Log. 3.5997597
 : 3979.178 miles, *balloon's dist. from the* }
 Earth's center - - - - - } - Log. 3.5997934

But $3979.178 - 3978.87 = .308$ or 542.08 yards. Ans.

Quest. 58.* Construct the triangle ABC having its three sides respectively equal to the sides given: namely, AB 213 yards, AC 424 yards, and BC 262 yards. Then at A make the $\angle CAE$ equal to the observed $\angle CSB 29^\circ .. 50'$; and at c make the $\angle ACE$ equal to the observed $\angle ASB 13^\circ .. 20'$.



Through the three points AEC, (E being the intersection of AE and EC) describe a circle; and join EB. If EB be produced, it will cut the opposite circumference in s, the station of the observer.

Draw AC, and cs; the positions are Geometrically determined.

Trigonometrically.

$180^\circ - 43^\circ .. 20'$ (the sum of the \angle^{ics} at s) = $136^\circ .. 40'$ $\angle AEC$; whence AE, EC, ap, pc, (p being the point in which a perp. from

* This Question produces a figure the same in species as the figure arising from the 25th Example of HEIGHTS and DISTANCES, vol. ii. page 25. And the sides here are respectively to the sides there, as 212 to 265, which is evident by comparing AB given, in the one, with AB, in the other.


(Key to Vol. II. page 265.)

B cuts AC;) also the \angle^{les} BCA, BAC, &c. &c. are all known. [See Ex. 25. page 25. vol. ii.]

$$\text{And, by resolution, } \left. \begin{array}{l} AS=605\cdot71 \text{ yards.} \\ BS=429\cdot68 \text{ yards.} \\ CS=524\cdot24 \text{ yards.} \end{array} \right\} \text{Ans.}$$

(Page 266.)

Quest. 59. Construct a triangle ABC, having its three sides AB, AC, and BC, equal to the given sides, each to each: and at the point A make the $\angle CAE$ equal to the observed $\angle BSC$ $22^\circ .. 30'$: also at C make the $\angle ACE$ equal to the observed $\angle ASB$ $33^\circ .. 45'$; and from B draw a straight line to E, the point of intersection of AE and CE.



Then through the three points AEC, which are not in the same right line, describe a circle; and produce BE until it meet the opposite circumference at s, the station of the observer. If the lines SA and SC be drawn, the figure is Geometrically determined, and the distances nearly as in the Answer given.

Trigonometrically.

The $\angle AEC$ is the supplement of the sum of the \angle^{les} at s, hence equal to $123^\circ .. 45'$. And,

$$\begin{aligned} \text{As Sin. } 123^\circ .. 45' \angle AEC : 12 \text{ furl. AC} &:: \text{Sin. } 22^\circ .. 30' \angle EAC : \\ &5\cdot523 \text{ furlongs EC.} \\ :: \text{Sin. } 33^\circ .. 45' \angle ACE : 8\cdot0181 \text{ \&c. furlongs AE.} \end{aligned}$$

Demitting a perpendicular from B upon AC, and proceeding in all respects as in Example 26,* page 25 of the second volume of the COURSE, the Answer is found as below:

$$\left. \begin{array}{l} AS 10\cdot6545 \text{ furlongs} \\ BS 15\cdot6277 \text{ furlongs} \\ CS 14\cdot0121 \text{ furlongs} \end{array} \right\} \text{Ans.}$$

* The present Example is in fact in all the angles the same as the Example to which we allude, and consequently the sides *here* will be to the analogous sides *there* in the ratio of 1320 yards, or 6 furlongs, BC *here*, to 400 yards BC *there* $\therefore 3\cdot3 : 1$, either in yards or in furlongs.

(Key to Vol. II. page 266.)

Quest. 60. $5 \text{ dwt. gr.} \dots 9\frac{1}{2} = 129\frac{1}{2}$ grains of standard gold in a guinea.

Now, As $21 : 20 :: 129\frac{1}{2} : 123\frac{1}{3}$ the value of £1 in standard gold.

But, $\frac{\text{gr. } 123\cdot3 \times \text{£}480000000}{437\cdot5 \text{ gr. in an oz. avoird.}^*} = 135314285\cdot7142857$ the weight of the cube in ounces Avoirdupois.

And, As $17724\ddagger : 135314285\cdot7 \text{ \&c.} :: 1 : 7634\cdot523$ the solidity of the cube in feet.

Lastly, $\sqrt[3]{7634\cdot523 \text{ ft.}} = 19\cdot69069$ feet. Ans. †

* See the First Volume of the Mathematics, page 26.

† By reference to the table of Specific Gravity, given in the Second Volume of the Course, page 231.

‡ Here we differ from Dr. Charles Hutton about 1 foot in the side of the cube, and think the gold would occupy at least *one thousand one hundred and five cubic feet* more than the space allotted it by the Doctor.

Quest. 61. $\frac{20\text{ ft.} + 22\text{ ft.}}{2} = 21$ feet, mean breadth of the ditch.

And $21 \text{ ft.} \times 9 \text{ ft.} \times 1000 \times 1728 = 326592000$ cubic inches, content of the ditch.

But $\frac{\text{cub. inches } 326592000 \div 282}{\text{cub. in.}} = 1158127\frac{3}{7}$ gallons. Ans.

Quest. 62. As $7930^3 : 2160^3 :: \text{Earth's solidity} : \text{Moon's solid}^y$;

And

As $7930^2 : 2160^2 :: \text{Earth's surface} : \text{Moon's surface}$.

But

$7930^3 = 49867725700.$ $2160^3 = 1007769600.$
 $7930^2 = 62884900.$ $2160^2 = 4665600.$

Hence

Solidity of the Earth : Solidity of the Moon :: $49\frac{1}{2} : 1$ }
 Surface of the Earth : Surface of the Moon :: $13\frac{1}{2} : 1$ } Ans.

(Key to Vol. II. page 267.)

PRACTICAL QUESTIONS IN SPECIFIC GRAVITY.

To determine the Magnitude of a Body by its Weight.

Ex. 1. As 2520 oz. tabular specific gravity of COMMON STONE,
 : 1792 oz. weight of the given block,
 :: 1728 cubic inches in 1 cubic foot,
 : $1228\frac{4}{5}$ cubic inches in the block. Ans.

Ex. 2. As 937 oz. tabular specific gravity of GUNPOWDER,
 : 16 oz. weight of the quantity given,
 :: 1728 cubic inches in 1 cubic foot,
 : $29\frac{1}{2}$ cubic inches nearly. Ans.

Ex. 3. As 925 oz. tabular specific gravity of DRY OAK,
 : 35840 oz. in one ton,
 :: 1 cubic foot,
 : $38\frac{138}{185}$ cubic feet. Ans.

To determine the Weight of a Body by its Magnitude.

Ex. 1. $\begin{matrix} \text{ft.} & \text{ft.} & \text{ft.} \\ 63 \times 12 \times 12 = & 9072 \text{ cubic feet in the block.} \end{matrix}$

And

As 1 cubic foot,
 : 9072 cubic feet,
 :: 2700 oz. tabular specific gravity of MARBLE,
 : 24494400 oz. in the block, = 683.4 tons. Ans.

Ex. 2. As 1728 cubic inches in 1 cubic foot,
 : $35\frac{1}{4}$ cubic inches in an ale pint,
 :: 937 oz. tabular specific gravity of gunpowder,
 19.1 oz. the weight required.

Ex. 3. $\begin{matrix} \text{ft.} & \text{ft.} & \text{ft.} \\ 10 \times 3 \times 2\frac{1}{2} = & 75 \text{ solid feet of oak in the block.} \end{matrix}$

(Key to Vol. II. page 267.)

And,

As 1 cubic foot,
 : 75 cubic feet, content of the block,
 :: 925 oz. tabular specific gravity of OAK,
 : 69375 oz. = $4335\frac{15}{16}$ lb. the weight required.

** The last six Questions are found at the 231st and following page of the second volume of the COURSE; and the three following, with only a slight variation in the first, in pages 229 and 230.

(Page 268.)

To ascertain the Specific Gravity of a Body.

lb. lb. lb.
 Ex. 1. $10 - 6\frac{1}{6} = 3\frac{5}{6}$ difference of weight in water and air.

And

As $3\frac{5}{6}$ lb. weight lost in water,
 : 10 lb. weight in air,
 :: 1000 oz. specific gravity of WATER,
 2609 oz. specific gravity required.

lb. lb. lb.
 Ex. 2. $18 + 15 = 33$ weight of the aggregate body in air.
 $18 - 16 = 2$ lost by the copper in water.
 Also $33 - 6 = 27$ lost by the aggregate mass in water.
 And $27 - 2 = 25$ difference of loss.

But, As 25 lb. : 15 lb. :: 1000 oz. : 600 oz. Ans.

(Page 269.)

To determine the Quantity of each Ingredient in a Body compounded of Two known Ingredients.

oz. oz. oz.
 Ex. 1. $9000 - 7320 = 1680$ first difference.
 $8784 - 7320 = 1464$ second difference.
 $9000 - 8784 = 216$ third difference.

Also,

$8784 \times 1680 = 14757120.$
 $9000 \times 1464 = 13176000.$
 $7320 \times 216 = 1581120.$

(Key to Vol. II. page 269.)

$$\begin{array}{r}
 \text{But,} \\
 \text{As } 14757120 : 112 : : 13176000 : 100 \text{ copper } \left. \vphantom{\begin{array}{l} 14757120 \\ 13176000 \end{array}} \right\} \text{Ans.} \\
 \quad \quad \quad \text{lb.} \quad \quad \quad \text{lb.} \\
 \quad \quad \quad 14757120 : 112 : : 1581120 : 12 \text{ tin} \\
 \text{Sum and Proof } \quad \quad \quad \underline{\quad} \quad 112 \text{ lb.}
 \end{array}$$

BALLS AND SHELLS.

To determine the Weight of an Iron Ball of known Diameter.

$$\begin{array}{l}
 \text{Ex. 1.} \quad \begin{array}{cccc} \text{in.} & \text{in.} & \text{in.} & \text{cub. inches} \end{array} \\
 6.7 \times 6.7 \times 6.7 = 300.763 \text{ cube of the diameter.} \\
 \text{And,} \\
 \text{lb. } \frac{9}{84} \times 300.763 = 42.2948 \text{ lb. Ans.}
 \end{array}$$

$$\begin{array}{l}
 \text{Ex. 2.} \quad \begin{array}{cccc} \text{in.} & \text{in.} & \text{in.} & \text{cubic inches} \end{array} \\
 5.54 \times 5.54 \times 5.54 = 170.031464 \text{ cube of the diameter.} \\
 \text{Now,} \\
 \text{lb. } \frac{9}{84} \times 170.031464 = 23.99 \text{ lb. or } 24 \text{ lb. nearly. Ans.}
 \end{array}$$

(Page 270.)

To determine the Weight of a Leaden Ball of known Diameter.

$$\begin{array}{l}
 \text{Ex. 1.} \quad \begin{array}{cccc} \text{in.} & \text{in.} & \text{in.} & \text{cubic inches.} \end{array} \\
 6.6 \times 6.6 \times 6.6 = 287.496 \text{ cube of the diameter.} \\
 \text{But,} \\
 \text{lb. } \frac{3}{14} \times 287.496 = 61.606 \text{ lb. Ans.}
 \end{array}$$

$$\begin{array}{l}
 \text{Ex. 2.} \quad \begin{array}{cccc} \text{in.} & \text{in.} & \text{in.} & \text{cub. inches} \end{array} \\
 5.3 \times 5.3 \times 5.3 = 148.877 \text{ cube of the diameter.} \\
 \text{And,} \\
 \text{lb. } \frac{3}{14} \times 148.877 = 32 \text{ lb. nearly. Ans.}
 \end{array}$$

To ascertain the Diameter of an Iron Ball of known Weight.

$$\begin{array}{l}
 \text{Ex. 1.} \quad \begin{array}{cccc} \text{lb.} & \text{lb.} & \text{cub. in.} & \text{cub. in.} \end{array} \\
 \text{As } 9 : 42 : : 64 : 298.66 \text{ cube of the diameter.}
 \end{array}$$

(Key to Vol. II. page 270.)

But $\sqrt[3]{298\cdot66 \text{ in.}} = 6\cdot685 \text{ inches. Ans.}$

Or, by the Rule,

$$\sqrt[3]{42 \times \frac{64}{9}} = 6\cdot685 \text{ the diameter in inches, as before.}$$

lb. lb. cub. in. cub. in.

Ex. 2. As 9 : 24 :: 64 : 170·66 cube of the diameter.

And $\sqrt[3]{170\cdot66 \text{ in.}} = 5\cdot54 \text{ inches. Ans.}$

That is,

$$\sqrt[3]{24 \times 7\frac{1}{3} \text{ lb.}} = 5\cdot54 \text{ the diameter in inches, as before.}$$

To ascertain the Diameter of a Leaden Ball of known Weight.

lb. lb. cub. in. cub. in.

Ex. 1. As $\frac{3}{4}$: 64 :: 1 : 298·66 cube of the diameter.

Now,

$$\sqrt[3]{298\frac{2}{3} \text{ in.}} = 6\cdot685 \text{ inches. Ans.}$$

lb. lb. cub. in. cub. in.

Ex. 2. As $\frac{3}{4}$: 8 :: 1 : 37 $\frac{1}{3}$ cube of the diameter.

Hence, by extraction of the cube root, 3·343 inches. Ans.

(Page 271.)

To determine the Weight of an Iron Shell of known diameter and thickness.

Ex. 1. $\begin{array}{cccc} \text{in.} & \text{in.} & \text{in.} & \text{cub. inches} \\ 12\cdot8 \times 12\cdot8 \times 12\cdot8 & = & 2097\cdot152 & \text{the outer diameter cubed.} \end{array}$

And $9\cdot1 \times 9\cdot1 \times 9\cdot1 = 753\cdot571$ the internal diam^r. cubed.

$$\begin{array}{r} \text{Multiply by } \underline{1343\cdot581} \text{ difference of the cubes.} \\ \frac{2}{64} \end{array}$$

$$\text{PRODUCT } \underline{188\cdot941 \text{ lb. Ans.}}$$

(Key to Vol. II. page 271.)

$$\begin{array}{l} \text{Ex. 2.} \quad \begin{array}{cccc} \text{in.} & \text{in.} & \text{in.} & \text{cub. inches} \\ 9.8 \times 9.8 \times 9.8 & = & 941.192 & \text{the outer diameter cubed.} \\ \text{And } 7 \times 7 \times 7 & = & 343 & \text{the inner diameter cubed.} \end{array} \end{array}$$

$$\begin{array}{r} \text{Multiply by} \quad \frac{598.192 \text{ difference of the cubes.}}{64} \\ \hline \text{PRODUCT } 84.12 \text{ lb. Ans.} \end{array}$$

To determine the Quantity of Gunpowder necessary to fill a Shell of known Capacity.

$$\begin{array}{l} \text{Ex. 1.} \quad \begin{array}{cccc} \text{in.} & \text{in.} & \text{in.} & \text{cub. inches} \\ 9.1 \times 9.1 \times 9.1 & = & 753.571 & \text{cube of the internal diameter.} \\ \text{Divide by } 57.3 \text{ cub. inches} & \text{---} & \text{the Quotient is } 13.15 \text{ lb. Ans.} \end{array} \end{array}$$

$$\text{Ex. 2.} \quad \frac{\begin{array}{ccc} \text{in.} & \text{in.} & \text{in.} \\ 7 \times 7 \times 7 \end{array}}{57.3} = 6 \text{ lb. nearly. Ans.}$$

To determine the Quantity of Powder necessary to fill a Rectangular Box of known Dimensions.

$$\text{Ex. 1.} \quad \frac{\begin{array}{ccc} \text{in.} & \text{in.} & \text{in.} \\ 15 \times 12 \times 10 \end{array}}{30} = 60 \text{ lb. Ans.}$$

$$\text{Ex. 2.} \quad \frac{\begin{array}{ccc} \text{in.} & \text{in.} & \text{in.} \\ 12 \times 12 \times 12 \end{array}}{30} = 57.6 \text{ lb. Ans.}$$

To ascertain the Weight of Powder sufficient to fill a hollow Cylinder of known Length and Diameter.

$$\text{Ex. 1.} \quad \frac{\begin{array}{ccc} \text{in.} & \text{in.} & \text{in.} \\ (10 \times 10) \times 20 \end{array}}{38.2} = 52.35 \text{ lb. Ans.}$$

(Page 272.)

$$\text{Ex. 2.} \quad \frac{\begin{array}{ccc} \text{in.} & \text{in.} & \text{in.} \\ (4 \times 4) \times 12 \end{array}}{38.2} = 5\frac{5}{191} \text{ lb. Ans.}$$

(Key to Vol. II. page 272.)

To determine the Size of a Shell that will just contain a given Weight of Powder.

Ex. 1. $\sqrt[3]{13\frac{1}{6} \times 57.3} = 9.1$ nearly, the diameter in inches. Ans.

Ex. 2. $\sqrt[3]{6 \times 57.3} = 7$ nearly, the diameter in inches. Ans.

To determine the Side of a Cubical Box that will just hold a specified Portion of Gunpowder by Weight.

Ex. 1. $\sqrt[3]{50 \times 30} = 11.44$, the side required, in inches.

Ex. 2. $\sqrt[3]{400 \times 30} = 22.89$ &c. the side required, in inches.

To determine the Length a given Weight of Gunpowder will occupy of a Cylinder of known Diameter.

Ex. 1. $\frac{\text{lb. cub. in.} \quad \text{cub. inches.}}{12 \times 38.2 \quad 458.4 \times 9}$
 $\frac{\quad \quad \quad \text{in.}}{(6\frac{2}{3} \times 6\frac{2}{3}) \text{ in.}} = \frac{\quad \quad \quad}{400 \text{ sq. in.}} = 10.314$ in. the length reqd.

Ex. 2. $\frac{\text{lb. cub. in.} \quad \text{cub. inches}}{20 \times 38.2 \quad 764}$
 $\frac{\quad \quad \quad \text{in.}}{(8 \times 8) \text{ in.}} = \frac{\quad \quad \quad}{64 \text{ sq. in.}} = 11\frac{1}{6}$ inches. Ans.

(Page 273.)

To ascertain the Number of Balls in a Triangular Pile.

Ex. 1. $\frac{30 \times 31 \times 32}{6} = 4960$ balls. Ans.

Ex. 2. $\frac{20 \times 21 \times 22}{6} = 1540$ balls. Ans.

(Page 274.)

To ascertain the Number of Balls in a Square Pile.

Ex. 1. $\frac{30 \times 31 \times 61}{6} = 9455$ balls. Ans.

(Key to Vol. II, page 274.)

$$\text{Ex. 2. } \frac{20 \times 21 \times 41}{6} = 2870 \text{ balls. Ans.}$$

To ascertain the Number of Balls in a Rectangular Pile.

$$\text{Ex. 1. } \begin{array}{l} \text{balls} \quad \text{balls} \\ 46 \times 3 = 138 \text{ thrice the number of shot in the length of} \\ \text{the base of the pile. And} \\ 15 - 1 = 14 \text{ one ball less than the number of shot in the} \\ \text{breadth of the base of the pile.} \end{array}$$

$$\text{But } \frac{\text{balls} \quad \text{balls} \quad (138 - 14) \times 15 \times 16}{6} = 4960 \text{ balls. Ans.}$$

$$\text{Ex. 2. } \begin{array}{l} \text{shot} \quad \text{shot} \\ 59 \times 3 = 177 \text{ thrice the number of balls in the length of} \\ \text{the base of the pile. And} \\ 20 - 1 = 19 \text{ one shot less than the number of balls in} \\ \text{the breadth of the pile.} \end{array}$$

$$\text{But } \frac{\text{shot} \quad \text{shot} \quad (177 - 19) \times 20 \times 21}{6} = 11060 \text{ shot. Ans.}$$

To ascertain the Number of Balls in an Incomplete Pile.

$$\text{Ex. 1. } \begin{array}{l} \frac{1}{6} \times 40 \times 41 \times 42 = 11480 \text{ shot, for the complete Pile.} \\ \text{And } \frac{1}{6} \times 19 \times 20 \times 21 = 1330 \text{ shot wanting.} \end{array}$$

$$\text{Difference } \underline{\underline{10150 \text{ shot. Ans.}}}$$

(Page 275.)

$$\text{Ex. 2. } \begin{array}{l} \frac{1}{6} \times 24 \times 25 \times 26 = 2600 \text{ shot, for the complete pile.} \\ \text{And } \frac{1}{6} \times 7 \times 8 \times 9 = 84 \text{ shot wanting.} \end{array}$$

$$\text{Difference } \underline{\underline{2516 \text{ shot. Ans.}}}$$

$$\text{Ex. 3. } \begin{array}{l} \frac{1}{6} \times 24 \times 25 \times 49 = 4900 \text{ balls, for the complete pile.} \\ \text{And } \frac{1}{6} \times 7 \times 8 \times 15 = 140 \text{ balls wanting.} \end{array}$$

$$\text{Difference } \underline{\underline{4760 \text{ balls. Ans.}}}$$

(Key to Vol. II, page 275.)

shot shot shot
Ex. 4. $40 - 11 = 29$ in the top row of either side of the pile.

And

$20 - 11 = 9$ in the top row of either end of the pile.

Therefore the length of the base of DEFECT is 28 shot, and the breadth 8 shot.

But, $(40 \times 3 - 19) \times 20 \times 21 \times \frac{1}{6} = 7070$ shot, for the whole pile.

And $(28 \times 3 - 7) \times 8 \times 9 \times \frac{1}{6} = 924$ shot deficient.

Difference 6146 shot. Ans.

DISTANCE ESTIMATED BY THE VELOCITY OF SOUND.

Ex. 1. $\frac{3}{4}$ mls. $\times 12 = \frac{36}{4} = 2\frac{1}{2}$ miles. Ans.

Or, more correctly,

As $1'' : 12'' :: 1142 \text{ feet} : 13704 \text{ feet} = 2 \text{ miles} \text{ .. } 1048 \text{ yds.}$ Ans.

Ex. 2. $4''\frac{2}{3} \times 8 = 37\frac{1}{3}$ seconds. Ans.

Ex. 3. As $4''\frac{2}{3} : 7'' :: 1 \text{ mile} : 1\frac{1}{2} \text{ mile}$. Ans.

Ex. 4. As $70 \text{ pulsat.} : 6 \text{ pulsat.} :: 60'' : 5''\frac{1}{7}$ time of the sound in seconds.

And

As $4''\frac{2}{3} : 5''\frac{1}{7} :: 1 \text{ mile} : 1\frac{5}{9} \text{ mile} = 1 \text{ .. } 179\cdot6 \text{ yds.}$ Ans.*

More correctly,

The time being, as already found, $5\frac{1}{7}$ seconds, it is

* Differing 18·4 yards from Dr. Hutton's Answer, which differs 395½ yards from the truth by Rule, allowing sound to travel regularly 1142 feet per second. But the Student will bear in mind that, owing to currents of air and other circumstances, the velocity of sound cannot be exactly depended on.

(Key to Vol. II. page 275.)

$$\text{As } 1'' : 5''\frac{1}{7} :: 1142 : 5873\frac{1}{7} = 1 : 593\frac{1}{7}. \quad \text{Ans.}$$

(Page 276.)

$$\text{Ex. 5. } \begin{array}{c} \text{pulsat.} \\ \text{As } 75 : 5 : 60'' : 4'' \end{array} \text{ time of the sound in seconds.}$$

But,

$$1142 \times 4 = 4568 = 1522\frac{2}{3} \text{ yards.} \quad \text{Ans.}$$

$$\text{Ex. 6. } \begin{array}{c} \text{mile} \\ \text{As } 4''\frac{2}{3} : 33'' :: 1 : 7\frac{1}{4}. \end{array} \text{ Ans.}$$

PRACTICAL QUESTIONS IN PHILOSOPHY.

$$\text{Quest. 1. } \begin{array}{c} \text{in.} \\ 3 \times 3 \times 3 \times \cdot 5236 = 14 \cdot 1372 \text{ cub. in. cast iron in the ball.} \\ \text{in.} \\ \text{Multiply by } \cdot 258 \text{ lb. Avoirdupois.} \end{array}$$

$$\text{PRODUCT } \underline{\underline{3 \cdot 6473976 \text{ lb.}}} \quad \text{Ans.}$$

$$\text{Quest. 2. } \begin{array}{c} \text{in.} \\ 5^3 - 3^3 = 98 \text{ diff. of the cubes of the diameters.} \\ \text{in.} \\ \text{cub. in.} \end{array}$$

Now

$$\text{lb. } \frac{9}{84} \times 98 = 13 \cdot 78125 \text{ lb.} \quad \text{Ans.}^*$$

* *By which it appears that the weight of the Shell is more than $\frac{1}{2}$ lb. above Dr. Hutton's estimate.*

$$\text{Quest. 3. } \begin{array}{c} \text{mile} \\ \text{As } 4''\frac{2}{3} : 17'' :: 1 : 3\frac{9}{4}. \end{array} \text{ Ans.}$$

$$\text{Quest. 4. } \text{By Question 62, page 266 of the second volume of the COURSE, the bulk of the Earth is to that of the Moon as } 49\frac{1}{2} \text{ to 1.}$$

(Key to Vol. II. page 276.)

Therefore $\frac{49.5 \times 10}{7} = 70.7$, to 1. Ans.

Quest. 5. Referring to the Table of Specific Gravity in vol. ii. page 231,

A cubic foot of brass weighs 8000 oz. avoirdupois.
 A cubic foot of marble - - - 2700 oz.

	—		
Difference	5300 oz.		
Multiply by	$1\frac{1}{2}$		
	—	lb.	oz.
PRODUCT	7950 oz.	= 496 ..	14. Ans.
	—		

Quest. 6.* $63 \times 12 \times 12 = 9072$ cubic feet. And

As 1 cubic foot,
 : 9072 cubic feet,
 :: 2700 oz. tabular specific gravity of MARBLE,
 : 24494400 oz. = 683.4 tons. Ans.

* The present Question was surely a favourite, for this is the third time of its appearance in the second volume. See pages 232, 267, and 276.

Quest. 7. It is evident, since the *momenta* must be equal, that the velocities will be inversely as the weights. Hence

lb. lb. ft. ft.
 As 32 : 10000 :: 20 : 6250 per second. Ans.

(Page 277.)

Quest. 8. Multiplying the velocities respectively by the weights, it is

(v) (w)
 $1000 \times 1 = 1000$ the momentum of the less body.
 And $1 \times 25 = 25$ the momentum of the greater body.
 But As 25 : 1000 :: 1 : 40.

Wherefore the momentum of the *less* is to that of the *greater* body as 1 to 40. Ans.

(Key to Vol. II. page 277.)

Quest. 9. $\begin{matrix} \text{ft.} & \text{lb.} \\ 100 \times 20 = 2000 \end{matrix}$ momentum of the 1st body.

$\begin{matrix} (\text{m}) \\ \text{And } 2000 \div 8 \text{ lb.} = 250 \end{matrix}$ feet velocity of the less body per second. Ans.

Or, the velocities being in the inverse ratio of the weights, it is

$\begin{matrix} \text{lb.} & \text{lb.} & \text{ft.} & \text{ft.} \\ \text{As } 8 : 20 :: 100 : 250 \end{matrix}$ per second, *as before.*

Quest. 10. $\begin{matrix} \text{lb.} & \text{lb.} & (\text{v.}) & (\text{v.}) \\ \text{As } 60 : 100 :: 8 : 13\frac{1}{3} \end{matrix}$ the ratio to 1, the velocity of the less body has to that of the greater. But $13\frac{1}{3}$ are to 1 as 40 to 3. Ans.

$\begin{matrix} (\text{Force}) \text{ lb. \&c.} \\ \text{Quest. 11. } 48 \div 8 = 6 \end{matrix}$ velocity of the greater body.
 And $1 \div 1 = 1$ velocity of the less body.
 Hence the velocity of the greater is to the velocity of the less, as 6 to 1. Ans.

Quest. 12. $\begin{matrix} \text{min.} & (\text{v.}) & \text{min.} & (\text{v.}) \\ \text{As } 1 \times 40 : 120 \times 1 :: 1 : 3 \end{matrix}$ the ratio of the space described by the swifter, to that described by the slower body. Ans.

Quest. 13. $\begin{matrix} \text{min.} & (\text{v.}) & \text{min.} & (\text{v.}) & \text{ft.} & \text{ft.} \\ \text{As } 1 \times 1 : 12 \times 30 :: 5 : 1800 \end{matrix}$ by the swifter body.
 Subtract 5 by the slower body.

Difference 1795 feet. Ans.

Quest. 14. It is manifest that if the bodies had moved with equal velocities, the slower body would have passed over $5 \times 5 = 25$ miles.

$\begin{matrix} \text{miles} & \text{miles} \\ \text{Now, As } 25 : 50 :: 1 : 2 \end{matrix}$ the ratio of the time of the slower body to that of the faster. Ans.

Quest. 15. It will be,

$\begin{matrix} \text{in.} & \text{ft. high} & \text{in.} & \text{ft. high} & \text{hrs.} & \text{hrs.} & \text{min.} \\ \text{As } 2(18 \times 104) : 40 \times 73 :: 13 : 10 \text{ .. } 8\frac{1}{2} \end{matrix}$ Ans.

(Key to Vol. II. page 278.)

Quest. 16. $100 - 7\frac{1}{2} = 92\frac{1}{2}$ dist. of the power from the fulcrum.

And

As $7\frac{1}{2} : 92\frac{1}{2} :: 168 (= 1\frac{1}{2} \text{ cwt.}) : 2072$. Ans.

Quest. 17. $1\frac{1}{2} \times 28 = 42$ for the weight in the palm.

And $1\frac{1}{2} \times 12 = 18$ for the weight at the elbow.

Difference 24 lb. Ans.


Quest. 18. $70 \div 2 = 35$, that is, the ratio of the distance of the power to that of the weight, from the fulcrum, is 35 to 1.

Now $9\frac{1}{2} \text{ cwt.} = 1064 \text{ lb.}$

And $1064 \text{ lb.} \div 35 = 30.4 \text{ lb.}$ Ans.

Quest. 19.

As $36 : 2 :: 10 : \frac{5}{9}$,
which \times^{ed} by 2 =
 $1\frac{1}{9} \text{ lb.}$ balanced at the
fulcrum: hence $8\frac{8}{9} \text{ lb.}$
of the beam remain
unbalanced.



But $8\frac{8}{9} \text{ lb.} \times 18 \times \frac{1}{2} = 80 \text{ lb.}$ required at the short end to keep
the beam in equilibrio. And

$95 - 80 = 15$ the real weight to be balanced.

Now, As $1 : 15 :: 2 : 30$ the dist. of 1 from the fulcrum.

$2 : 15 :: 2 : 15$ - - - - - 2

$3 : 15 :: 2 : 10$ - - - - - 3

$4 : 15 :: 2 : 7\frac{1}{2}$ - - - - - 4

$5 : 15 :: 2 : 6$ - - - - - 5

&c. &c. &c.

Hence 30, 15, 10, $7\frac{1}{2}$, 6, $4\frac{2}{7}$, $3\frac{3}{4}$, $3\frac{1}{3}$, 3, &c. Ans.

Quest. 20. As $4 : 2.5 :: 200 : 125$ for the shorter end. }
 $4 : 1.5 :: 200 : 75$ for the longer end. } Ans.

(Key to Vol. II. page 278.)

Quest. 21. $\sqrt{\text{lb. lb.}}$
 $\sqrt{40 \times 90} = 60 \text{ lb.}$ *the true weight of the body.*

But, As lb. lb.
 60 : 40 :: 3 : 2 *the proportion required.*

Quest. 22. ft. ft. ft.
 $13 - 1 = 12$ dist. of the power from the fulcrum, 1
 foot occasioning by the Question a *diff.* of 210 lb. in the
 power required.

Now, As ft. ft. lb. lb.
 1 : 12 :: 210 : 2520. Ans.

(Page 279.)

Quest. 23. Without discussing the merits of this Example, it
 will be,

As 1 inch (EF) advanced towards the fulcrum by the weight,
 : 12 inches (AG) advanced towards the same by the power,
 :: 2 lb. the power,
 : 24 lb. *the weight.*

Which is evident, since there is said to be an equilibrium in both
 situations.

Quest. 24. As in. in. in. in.
 50 : 30 :: 40 : 24 = BD.

And

As 30 in. : 40 in. :: 3 : 4 :: P : Q.

Therefore the ratio of P to Q is that of 3 to 4; and the altitude
 BD, 24 inches. Ans.

Quest. 25. ft. ft. ft.
 $6 + 6 = 12 = 144$ inches the diameter of the circle.

(Di.) (Circ.) (Di.) (Circumf.)
 And, As 1 : 3.1416 :: 144 in. : 452.3904 in. orb. of the power.
 Multiply by 50 lb. the power.

PRODUCT and Answer 22620 lb. nearly.


Quest. 26. As ft. ft. lb. lb.
 20 : 30 :: 150 : 225. Ans.

(Key to Vol. II. page 279.)

Quest. 27. As $\frac{2 \text{ in.}}{12 \text{ in.}} :: \frac{150 \text{ lb.}}{900 \text{ lb.}}$ Ans.

Quest. 28.

Let bc represent the pillar and ap a perpendicular from the lowest point of the base; the $\angle apc$, by the question, is 15° , and ac 30 feet: wherefore it is,



As $\text{Sin. } 15^\circ \angle apc$, - - - - - $\text{Log. ar. co.} - 10.5870038$
 : $\text{Sin. } 75^\circ \angle pac$, - - - - - $\text{Log. } 9.9849438$
 :: 30 feet ac , - - - - - $\text{Log. } 1.4771213$

 ; 111.96 feet pc , the length required - $\text{Log. } 2.0490689$

Quest. 29. $\frac{2}{5} \times 12\sqrt{\frac{4}{5}} \text{ ft.} = \frac{48}{5}\sqrt{\frac{1}{5}} \text{ ft.} = \frac{48}{25}\sqrt{5} \text{ ft.} = 4.29325056$
 feet. Ans. See page 198, &c. vol. ii. of the MATHEMATICS.

Quest. 30. $\frac{1}{2} \times 12\sqrt{4 \div 5} \text{ ft.} = 12\sqrt{\frac{1}{5}} \text{ ft.} = \frac{12}{5}\sqrt{5} \text{ ft.} = 5.3665632$
 feet. Ans. See the second vol. of the MATHEMATICS, p. 199.

(Page 280.)

Quest. 31.

For the rectangular wall.

Because $FE^* = AQ\sqrt{\frac{m}{3n}}$ [*Prop. xlv. Statics*], and that

As $\text{Rad.} : \text{Sin. } 60^\circ :: 12 \text{ feet} : 10.391 \text{ feet } AQ$, it is

$$FE = 10.391 \sqrt{\frac{4}{15}} \text{ ft.} = \frac{20.782}{15} \times \sqrt{15} \text{ ft.} = 5.365889 \text{ feet.}$$

* The thickness of the wall.

(Key to Vol. II. page 280.)


And for the triangular wall.

Because $FE = AQ \sqrt{\frac{m}{2n}}$ [*Prop. xlv. Statics*], and that

$AQ = 10.391$ feet, it is $FE = 10.391 \sqrt{.4} \text{ ft.} = 6.571845$ feet.*

* Dr. Hutton's Answer to this part of the Question is evidently wrong, in as much as $12 \sqrt{\frac{3}{10}} = 6.57267$ and not 6.53667 .

Quest. 32. Let HA be the surface of the water and AC its depth, which by the Question is 10 feet. Also let $FEDC$ be a perpendicular section of the wall, of which the height is 12 feet and thickness sought. Take CB equal to CA 10 feet, and draw the diagonal AB ; the triangle ABC represents the pressure of the water against AC . [*Corol. 3, prop. lxi. Hydrostatics.*] If in CA , therefore, ch be assumed equal to $\frac{1}{3}$ of CA , the whole pressure of the triangle ABC may be resolved to the point h . But the center of gravity of the wall is the intersection of the diagonals FD , EC , which being referred to a point g in the base, $dg = \frac{1}{2}CD$. Consequently putting x for CD the thickness sought, and considering D as the fulcrum, the weight at g , and the power at h , it is



$$(A) \dagger \quad (n) \quad (Dg) \quad (AC) \quad (CB) \quad (m) \quad (Ch)$$

$$12x \times 11 \times \frac{1}{2}x = \frac{1}{2} (10 \text{ ft.} \times 10 \text{ ft.}) \times 7 \times 3\frac{1}{3} \text{ ft.} = 1166\frac{2}{3} \text{ feet.}$$

That is,

$$66x^2 = 1166\frac{2}{3}, \text{ or } x^2 = 17.67\ddot{6}\ddot{7}.$$


Hence by extracting the square root on both sides,

$$x = 4.204374825 \text{ \&c. feet. Ans.}$$

† Where A represents the area of the wall, n its specific gravity, and m the specific gravity of water.

Quest. 33.

If FDC represent a perpendicular section of the wall, FC by the Question is 12 feet: also the water and the point h are as in the last example; but in this instance $dg = \frac{2}{3}DC$ [*Art. 236, page 199, vol. ii.*] Wherefore, putting $x = DC$ the thickness required, it is



(Key to Vol. II. page 280.)

$$(A) (n) (Dg) \quad (AC) \quad (CB) \quad (m) (Ch)$$

$$6x \times 11 \times \frac{2}{3}x = \frac{1}{2}(10 \text{ ft.} \times 10 \text{ ft.}) \times 7 \times 3\frac{1}{3} \text{ ft.} = 1166\frac{2}{3} \text{ feet.}$$

That is,

$$44x^2 = 1166\frac{2}{3} \text{ feet, and } x^2 = 26\cdot51\dot{5}\dot{1} \text{ feet.}$$

Hence, by extracting the square root on both sides,

$$x = 5\cdot1492865054 \text{ feet. Ans.*}$$

☞ It is evident by the proposition quoted, that the thickness of a triangular wall must always be to that of a rectangular wall, *cæteris paribus*, as $\sqrt{\frac{3}{10}} : \sqrt{\frac{1}{5}} \therefore \sqrt{3} : \sqrt{2} \therefore \sqrt{3} : \sqrt{2} \therefore 17320508 : 14142136$.

* The reader is requested to examine attentively the 15th Problem (as it is termed) of the Promiscuous Exercises in the third volume of the COURSE, where (after noticing a Question proposed by Mr. Burton, of Salton, in 1795, and answered in the Ladies' Diary the following year by Gentlemen in different parts of the kingdom, one of whom was Dr. O. G. Gregory,) the Author gives a theorem for the thickness of a triangular wall to support a body of water of equal height; the accuracy of which theorem we dispute.

$$\text{Quest. 34. As } \begin{matrix} \text{(Times)} & \text{(Time)} & & \text{miles} & & \text{miles} \\ \sqrt{4} : \sqrt{1} \therefore 2 : 1 :: 95000000 : 47500000. \text{ Ans.} \end{matrix}$$

$$\text{Quest. 35. As } 1^2 : 106^2 :: 1 : 11236. \begin{matrix} \text{(Time)} & \text{(Times)} \\ \text{Therefore } 11236 \text{ times as hot. Ans.} \end{matrix}$$

$$\text{Quest. 36. } 95^2 \div 495^2 = \frac{9025}{245025} = \frac{1}{27} \text{ nearly. Ans.}$$

$$\text{Quest. 37. As } \begin{matrix} \text{lb.} & \text{lb.} & \text{S. di.} & \text{S. di.} \\ 10 : 112 :: 1 : 11\cdot2. \end{matrix}$$

S. di.
But, $\sqrt{11\cdot2} = 3\cdot3466401$ semi-diameters of the earth from the earth's surface.

Or,

$$\text{As } \begin{matrix} \text{lb.} & \text{lb.} & \text{lb.} & \text{lb.} & \text{S. di.} \\ 112 : 10 \therefore 56 : 5 :: 1 : \frac{5}{56} \end{matrix} \text{ of a semidiameter of the earth from the earth's center.}$$

$$\text{Quest. 38. } \begin{matrix} \text{miles} & \text{miles} \\ 7930 \div 2 = 3965 \end{matrix} \text{ the semidiameter of the earth.}$$

$\begin{matrix} \text{miles} & \text{miles} & \text{miles} \\ 3965 + 50 = 4015 \end{matrix}$ distance of the body from the earth's center. It will therefore be,

$$\text{As } \frac{1}{3965^2} : \frac{1}{4015^2} \therefore \frac{1}{15721225} : \frac{1}{16120225} :: \begin{matrix} \text{oz.} & \text{oz.} & \text{dr.} \\ 16 : 15 \cdot 9\frac{5}{8} \end{matrix} \text{ nearly. Ans.}$$

(Key to Vol. II. page 280.)

Quest. 39. As $\overset{\text{miles}}{2160^3} : \overset{\text{miles}}{7930^3} :: 1 : 49.5$ the ratio of bulk.

Now $49.5 \times 10 = 495$ quantity of matter in the earth.

And $1 \times 7 = 7$ - - - - - in the moon.

But, As $495 \div 7 : 495 :: 30 : 29\frac{2}{3}\frac{2}{3}$.

And $30 - 29\frac{2}{3}\frac{2}{3} = \frac{2}{3}\frac{1}{3}\frac{2}{3} = \frac{105}{51}$ diameters of the earth from the earth's center.

Also $\frac{\text{Di. } 251^*}{502} - \frac{\text{Di. } 210^\dagger}{502} = \frac{\text{Di. } 41}{502} = 647.669$ miles below the surface of the earth. Ans.

* That is, one semidiameter of the earth.

† A fraction equal to $\frac{105}{251}$ diameters, as above.

(Page 281.)

Quest. 40. Put x for the distance of the point sought, in diameters of the earth from the earth; a for the quantity of matter in our own globe; and b for that in the moon; also let d be the mutual distance of the orbs in diameters of the earth;

Then, As $10a : 7b :: x^2 : \overline{d-x}^2$.

Whence $x = 26\frac{9}{11}$. Therefore the point required is $26\frac{9}{11}$ of the earth's diameters from the earth's center, and $3\frac{2}{11}$ times the diameter of the earth from the center of the moon.

Quest. 41. As $1''^2 : 11''^2$, that is, As $1 : 121 :: 16\frac{1}{2}$ feet : $1946\frac{1}{2}$ feet. Ans.

Quest. 42. As $1''^2 : 6''^2$, that is, As $1 : 36 :: 16\frac{1}{2}$ ft. : 579 ft. the depth of the 6'' well.

And As $1''^2 : 10''^2$, that is, As $1 : 100 :: 16\frac{1}{2}$ ft. : $1608\frac{1}{2}$ feet, the depth of the 10'' well.

But $1608\frac{1}{2}$ ft. - 579 ft. = $1029\frac{1}{2}$ feet *difference*. Ans.

Quest. 43. As $1''^2 : 19\frac{1}{2}''^2$, that is, As $1 : 380.25 :: 16\frac{1}{2}$ feet : $6115\frac{1}{6}$ feet. Ans.

(Key to Vol. II. page 281.)

Quest. 44. As $16\frac{1}{2}$ ft. : 400 ft. :: 1" : 25", hence 5" nearly. Ans.

Quest. 45. $1''$ $2''$ $3''$ &c. series in seconds.
 $16\frac{1}{2}$ ft. $64\frac{1}{3}$ ft. $144\frac{2}{3}$ ft. spaces passed through at the
 end of each second.

Also, $1''$ $2''$ $3''$ &c. series in seconds.
 $16\frac{1}{2}$ ft. $48\frac{1}{4}$ ft. $80\frac{5}{8}$ ft. spaces for each second.

Likewise, 100 ft. - $80\frac{5}{8}$ ft. = $19\frac{7}{8}$ ft.

But, As $80\frac{5}{8}$ ft. : $19\frac{7}{8}$ ft. :: $3''$: $\cdot 6088''$.

Again, $3'' + \cdot 6088 = 3\cdot 6088''$ the whole time of falling. } Ans.
 Lastly, As $1''^2$: $3\cdot 6088''^2$:: $16\cdot 083$ ft. : $209\cdot 45594434752$ ft. }

Quest. 46. If x be put for the number of seconds the stone is in motion, it will be by the question

$$16\frac{1}{2}x^2 + 1142x = 1142 \text{ ft.} \times 10'' = 11420. \text{ That is,}$$

$$x^2 + \frac{1142x}{16\cdot 083} = \frac{11420}{16\cdot 083}; \text{ or, reduced, } x^2 + 71x = 710. \text{ nearly.}$$

$$\text{Hence } x = 8''\cdot 8876.$$

But, As $1''^2$: $8''\cdot 8876^2$:: $16\frac{1}{2}$ ft. : $1270\cdot 3865$ ft. Ans.

Quest. 47. As $3\cdot 14159$: 1 :: 193 : $19\cdot 6$.
 (Cir.) (Di) in. in.

And $19\cdot 6$ in. $\times 2 = 39\cdot 2$ in. or more correctly $39\frac{1}{4}$ in. Ans.

(Page 282.)

Quest. 48.

As $1''^2$: $2''^2$:: $39\frac{1}{8}$ in. : $156\frac{1}{2}$ in. length of a pendl. of $2''$ }
 $1''^2$: $\frac{1''^2}{2}$:: $39\frac{1}{8}$ in. : $9\frac{2}{3}\frac{5}{8}$ inches - - - - - $\frac{1''}{2}$ } Ans.
 $1''^2$: $\frac{1''^2}{4}$:: $39\frac{1}{8}$ in. : $2\frac{5}{8}\frac{7}{8}$ inches - - - - - $\frac{1''}{4}$ }

Quest. 49. As 12 : $39\frac{1}{8}$:: 3600 : $11737\cdot 5$ sqr. of vibr. per min.
 in. in. (Sq. vib. per m.)

And, As 6 : $39\frac{1}{8}$:: 3600 : 23475 sqr. of vibr. per min.

Now $\sqrt{11737\cdot 5} = 108\cdot 33$ vibr. per min. by a pendl. of 1 foot.

And $\sqrt{23475\cdot 0} = 153\cdot 21$ vibr. per min. by a pendl. of $\frac{1}{2}$ foot.

(Key to Vol. II. page 282.)

vibr.
Again, $153.21 \times 60' = 9192.6$ vibrations in an hour.
And, $108.33 \times 60' = 6499.8$ vibrations in an hour.

Difference $\underline{\underline{2692.8}}$ Ans.

Quest. 50. $\begin{matrix} \text{in.} & \text{in.} & (\text{Sq. vib. per m.}) \\ \text{As } 18 : 39\frac{1}{8} :: 3600 : 7825 \end{matrix}$ square of vibr. per min.

But, $\sqrt{7825} = 88.4$ vibrations per minute.

And, $88.4 \div 60 = 1.473$ vibrations per second.

Also, $8 \text{ vib.} \div 1.473 = 5''.4$ time marked by the pendulum.

Now putting x for the time of the fall in seconds, there arises the following equation:

$$16\frac{1}{2}x^2 + 1142x = 1142 \text{ ft.} \times 5''.4 = 6167. \quad \text{That is,}$$

$$x^2 + 71x = 383.44. \quad \text{Whence } x = 5''.04.$$

But $16\frac{1}{2} \text{ ft.} \times 5''.04 \times 5''.04 = 408.54$ feet; or (by keeping several more decimal places from the beginning of the operation) 412.61 feet. Ans.

Quest. 51. $\begin{matrix} \text{As Rad.} & - & - & - & \text{Log. ar. co.} & - & 10.0 & \text{————} \\ & : & \text{Co. Sin. } 10^\circ & \text{giv. } \angle, & - & \text{Log. } 9.9933515 \\ & :: & 20 \text{ feet,} & - & - & - & - & \text{Log. } 1.3010300 \\ & & & & & & & \underline{\underline{19.696 \text{ feet}}} & - & - & - & \text{Log. } 1.2943815 \end{matrix}$

Now $20 \text{ ft.} - 19.696 \text{ ft.} = .304 \text{ ft.}$ versed Sine of 10° (to 20 ft. Rad.)

But, As $\sqrt{16\frac{1}{2} \text{ ft.}} : \sqrt{.304 \text{ ft.}} :: 32\frac{1}{6} \text{ ft.} : 4.4213 \text{ ft.}$ Ans.

Quest. 52. As $16\frac{1}{2} \text{ ft.} : 10 :: 1''^2 : .62176$ square of the time in seconds for the perpendicular altitude.

Now, $\sqrt{.62176''} = .78852''$ the time for 10 feet.

And, As $10 \text{ ft.} : 100 \text{ ft.} :: .78852'' : 7''.8852$ the time reqd. }
Also $\frac{100 \times 2}{7''.8852} = 25.364$ feet per second velocity acqd. } Ans.

Quest. 53. $\frac{60^2 \times 39\frac{1}{8} \text{ in.}}{(40 \times 40) \text{ vib.}} = 88\frac{1}{2}$ inches, the distance of the center of oscillation from the axis of motion.

(Key to Vol. II. page 282.)

And, As $\overset{\text{in.}}{118} : \overset{\text{in.}}{88^*} :: \overset{\text{in.}}{30} : \overset{\text{in.}}{22.36}$ the length of the chord to radius 88 inches. Referring now to Prop. lviii. of Statics, [See pp. 219, 220, and 221, vol. ii.]

$$\left. \begin{array}{l} a = 16\frac{1}{2} \text{ feet.} \\ b = 1 \text{ lb.} \\ c = 22.36 \text{ in.} \end{array} \right\} \begin{array}{l} g \\ i \\ o \end{array} = 88 \text{ inches.}$$

and $p = 500 \text{ lb.}$

Therefore,

$$\frac{bii + gop}{bio} \times c \times \sqrt{\frac{2a}{o}} = 501 \times \frac{22.36 \text{ in.}}{12 \text{ in.}} \times 2.094 = 933.53 \times 2.094 =$$

$1954.81182 \text{ feet per second, the velocity of the BALL.}$ And $\frac{ci}{o} \sqrt{\frac{2a}{o}} = 1.863 \text{ ft.} \times 2.094 = 3.9018199 \text{ feet, the velocity with which the center of oscillation passes the perpendicular.}$

* The fraction is omitted as of comparatively small value, but this omission occasions a little deviation in the Answer. The data were not sufficiently clear to determine correctly the center of gravity which might have been higher than the center of oscillation, and hence the center of gyration not precisely 88 inches.

Quest. 54. As $1000 : 925 :: 12 : 11\frac{1}{10}$ inches. Ans.

Quest. 55. Because the area of a sphere of which the diameter is 1 foot = 18.8496 inches, [Prob. vii. page 49. vol. ii.] it is $18.8496 \times \frac{1}{2} = 113.0976$ inches.

And $\sqrt{113.0976} = 10.6347$ inches. Now,

As 1000 oz. : 925 oz. :: 10.6347 inches : 9.9567 inches. Ans.

(Page 283.)

Quest. 56. Let x^3 be solid content of the wood in inches.

Then, As $x^3 : (x^3 - 3x^2) :: 1000 \text{ oz.} : (1000 - \frac{3000}{x}) \text{ oz.}$ the specific gravity of the wood.

Also, As $x^3 : (x^3 - 4\frac{8}{103}x^2) :: 1030 \text{ oz.} : (1030 - \frac{4200}{x}) \text{ oz.}$ the specific gravity of the same wood. Therefore $30x = 1200$,

Or, $x = 40$. But $(1000 - \frac{3000}{40}) \text{ oz.} = 925 \text{ oz.}$

(Key to Vol. II. page 283.)

Now in the table of specific gravity 925 oz. correspond to OAK.
And it is evident that each side of the cube is 40 inches. Q. E. I.

$$\begin{aligned} \text{Quest. 57. } & 12 \overset{\text{oz.}}{\div} (12 - 7) = 2\frac{2}{5} \\ & \text{And } 14\frac{1}{2} \overset{\text{oz.}}{\div} (14\frac{1}{2} - 9) = 2\frac{7}{11} \\ & \text{But, As } 2\frac{2}{5} : 2\frac{7}{11} :: 145 : 132. \text{ Ans.} \end{aligned}$$

$$\begin{aligned} \text{Quest. 58. } & 171 \overset{\text{gr.}}{\div} (171 - 120) = 3\frac{2}{9} \text{ glass.} \\ & \text{And } 102 \overset{\text{gr.}}{\div} (102 - 79) = 4\frac{10}{3} \text{ magnet.} \\ & \text{But, As } 3\frac{2}{9} : 4\frac{10}{3} :: 10 : 13 \text{ nearly. Ans.} \end{aligned}$$

$$\begin{aligned} \text{Quest. 59. } & 63 \overset{\text{oz.}}{\div} 10.36 = 6.08108 \text{ had the crown been gold.} \\ & 63 \overset{\text{oz.}}{\div} 5.85 = 10.76923 \text{ - - - - - silver.} \end{aligned}$$

$$\begin{array}{r} \text{By alligation, } 8.2245 \left\{ \begin{array}{l} 6.08108 \text{ } 2.54473 \text{ cubic inches.} \\ 10.76923 \text{ } 2.14342 \text{ cubic inches.} \end{array} \right. \end{array}$$

4.68815 Sum

$$\begin{array}{r} \text{By division, } 4.68815 \left\{ \begin{array}{l} 2.54473 \text{ } .5428 \text{ gold.} \\ 2.14342 \text{ } .4572 \text{ silver.} \end{array} \right. \end{array}$$

$$\begin{aligned} \text{But } .5428 \times 63 \text{ oz.} &= 34.1884 \text{ oz. gold} = \text{lb. } 2 \text{ .. } 10 \text{ .. } 3 \text{ .. } 22\frac{1}{4} \text{ gold.} \\ \text{And } .4572 \times 63 \text{ oz.} &= 28.8036 \text{ oz. silver} = \text{lb. } 2 \text{ .. } 4 \text{ .. } 16 \text{ .. } 1\frac{3}{4} \text{ silver.} \end{aligned}$$

NOTE.—Although this solution is generally considered correct, yet in point of fact it is erroneous; for an alloy of gold and silver has less bulk than the sum of the bulks of the two metals before the alloy: and consequently Archimedes was deceived when he thought he had discovered the true method of examining the crown. Independent, indeed, of the above circumstance, much depends on the temperature of the atmosphere and water at the time of observation.

$$\text{Quest. 60. } 3.84 \text{ lb.} \times 12 \div 1.4921 \text{ oz.} = 30.816 \text{ cubic inches of glass in the bottle.}$$

And 231 cubic inches = 1 gallon, the given quant. of brandy.

Also 124.0008 oz. = weight of the brandy.

The bottle and brandy therefore weigh 170.0808 oz.

- - - - - contain 261.816 cub. in.

(Key to Vol. II. page 283.)

But $261.816 \times .59542 \text{ oz.} = 155.89048272 \text{ oz.}$ the weight of an equal bulk of salt water.

Lastly, $170.0808 \text{ oz.} - 155.89048272 \text{ oz.} = 14.1903728 \text{ oz.}$ Ans.

Quest. 61. ^{cub. in.} $231 \times 5 = 1155$ cubic inches of brandy.

And $1728 \div 8 = 216$ cubic inches of oak in the cask.

Also $925 \text{ oz.} \div 8 = 115.625 \text{ oz.}$ weight of the wood.

But 1371 cubic inches = the content of the $\frac{1}{2}$ anker and brandy.

And $1371 \times .59542 \text{ oz.} = 816.32082 \text{ oz.}$ the gravity of an equal bulk of salt water.

Now the brandy weighs 620.04 oz.; hence 80.691 *absolute* oz. are necessary to sink the cask.

Again,

As the wt. of lead : the wt. of salt water :: 11325 oz. : 1030 oz.

Consequently $\frac{1}{10.99514}$ is the proportion of loss of weight lead sustains in salt water.

But,

^{oz.} $80.691 - \frac{1}{10.99514}$ of ^{oz.} 80.691 : ^{oz.} 80.691 :: ^{oz.} 80.691 : ^{oz.} 89.743. Ans.

Quest. 62. ^{cub. ft.} As 1 : ^{cub. ft.} 50000 :: 1000 oz. : $1395\frac{5}{6}$ tons. Ans.

(Page 284.)

Quest. 63. ^{in.} $30 \times 14 \div 12 = 35$ feet, height of water.

And $1000 \text{ oz.} \div 1.222$ (specif. gr. of air) = 818.18 the ratio to 1, that the weight of water bears to that of air; hence

$35 \text{ ft.} \times 818.18 = 28818.1818$ feet = 5.458 miles *nearly*, the height of air. Ans.*

(See art. 368, page 243, vol. ii.)

* Differing from Dr. Hutton more than 348 feet in the height of the atmosphere, though agreeing with him in the water-barometer.

Quest. 64. ^{ft.} As $16\frac{1}{2}$: a :: $\frac{32\frac{1}{6}}{6}$: $v.^2$ of mercury per second
^{ft.} As $16\frac{1}{2}$: b :: $\frac{32\frac{1}{6}}{6}$: $v.^2$ of water per second
^{ft.} As $16\frac{1}{2}$: c :: $\frac{32\frac{1}{6}}{6}$: $v.^2$ of air per second
 where $a = 30$ inches, $b = 35$ feet, and $c = 5.5240$ miles.

(Key to Vol. II. page 284.)

Therefore

The velocity of quicksilver is 12·681 ft. per second.

The velocity of water - - - 47·447 ft. per second.

The velocity of air - - - 1369·8 ft. per second.

(See pages 233, 234, and 235, vol. ii.)

	ft.	ft.	ft.				
Quest. 65.	10-1=9	and	1 × 9 × 4 =	36	but	√ 36 =	6·
	10-2=8	- -	2 × 8 × 4 =	64	- -	√ 64 =	8·
	10-3=7	- -	3 × 7 × 4 =	84	- -	√ 84 =	9·16515
	10-4=6	- -	4 × 6 × 4 =	96	- -	√ 96 =	9·79796
	10-5=5	- -	5 × 5 × 4 =	100	- -	√ 100 =	10·
	10-6=4	- -	6 × 4 × 4 =	96	- -	√ 96 =	9·79796
	10-7=3	- -	7 × 3 × 4 =	84	- -	√ 84 =	9·16515
	10-8=2	- -	8 × 2 × 4 =	64	- -	√ 64 =	8·
	10-9=1	- -	9 × 1 × 4 =	36	- -	√ 36 =	6·

Put b for the height of the fluid above the orifice, in inches ; $a = .031416$ of a square inch the area of the orifice ; and let $g = 193$ inches the descent in the first second, of a heavy body falling freely at the surface of the earth. Then

$a \times 2\sqrt{gb} = 2a\sqrt{gb}$; which taken for the several heights, the result is,

cubic inches	per second discharged through the hole	height.
3·0238		9 feet high.
4·2763	- - - - -	8 feet high.
5·2374	- - - - -	7 feet high.
6·1884	- - - - -	6 feet high.
6·7613	- - - - -	5 feet high.
7·4067	- - - - -	4 feet high.
8·0002	- - - - -	3 feet high.
8·5525	- - - - -	2 feet high.
9·0713	- - - - -	1 foot high.

Sum $58·5179 \times 60'' \times 10' \div 282$ cub. in. = 124·5 gallons. Ans.*

* This Answer differs about $\frac{1}{3}$ of a gallon from the Answer given with the Question.

(Page 285.)

Quest. 66. As 1 : 3·1416 :: 100 ft. : 314·16 feet the circumference of the globe.

(Key to Vol. II. page 285.)

And $314.16 \text{ ft.} \times 100 \text{ ft.} = 31416 \text{ sq. feet}$ the surface of the globe.

This \times^{cd} by $144 = 4523904$ square inches of surface.

Again, $100 \times 100 \times 100 \times .5236 = 523600$ cubic feet contained.

Now, As $1 : 523600 :: 1\frac{2}{9} : 639956$ of air displaced.

And, As $9000 \text{ oz.} : 639956 \text{ oz.} :: 1 \text{ cub. ft.} : 71.1062$ cubic feet of copper in the globe.

But 71.1062 cubic feet $= 71.1062$ square feet 12 inches thick.

Therefore (reciprocally) $31416 : 12 :: 71.1062 : .0271$ of an inch. Ans.*

* Which is nearly, but not exactly, Dr. Hutton's Answer.

Quest. 67. As $1 : 3.1416 :: 100 \text{ ft.} : 314.16 \text{ ft.}$ circumf. of the balloon.

Also $314.16 \text{ ft.} \times 100 \text{ ft.} = 31416$ square feet of surface.

But $\frac{31416}{100 \times 12} = 26.18$ cubic feet of copper in the balloon.

Now, As $1 : 26.18 :: 9000 : 235620$ the weight of copper in the balloon.

And $100 \times 100 \times 100 \times .5236 = 523600$ cubic feet in the sphere.

Also, As $1 : (\frac{1}{10} \text{ of } 1\frac{2}{9}) \text{ oz.} :: 523600 : 63995.6 \text{ oz.}$ of inflam. air.

And, As $1 : 1\frac{2}{9} \text{ oz.} :: 523600 : 639956 \text{ oz.}$ of atmospheric air displaced.

But 235620 oz. (wt. of the copp.) $+ 63995.6 \text{ oz.}$ (wt. of the gas) $= 299615.6 \text{ oz.}$ weight of the inflated balloon.

Therefore

$639956 \text{ oz.} - 299615.6 \text{ oz.} = 340340.4 \text{ oz.} = 21271.28 \text{ lb.}$ Ans.*

* Being about $1\frac{1}{2} \text{ lb.}$ less than given in the COURSE.

Quest. 68. $29\frac{1}{2} \text{ inches} \times 14 = 34.416 \text{ feet} = 413 \text{ inches}$, the height to which water ought to rise in a Toricellian tube.

And $413 \text{ in.} + 30 \text{ in.} = 443 \text{ inches.}$

(Key to Vol. II. page 285.)

Now if x represent the height sought in inches, it will be
As $443 - x : 413 :: 36 : 36 - x$.

Therefore $x^2 - 479x = -1080$ inches.

Hence $x = 239.5 \mp 237.2345885 = 2.2654115$. Ans.

ft. ft. (*Con. de.*) ($\frac{1}{3}$ *Alt.*) cub. ft.
Quest. 69. $8 \times 8 \times .7854 \times 4 = 201.0624$ cont. of the Paraboloid.

Also 30.9 inches $\times 14 = 432.6$ inches $= 36$ feet *nearly* for the altitude of water in vacuo.

Likewise 5 fathoms $= 30$ ft.; 10 fath^s. $= 60$ ft.; 15 fath^s. $= 90$ ft.
and 20 fath^s. $= 120$ ft.

	ft.	ft.	ft.	cub. feet	cub. feet	
But, As	$36 +$	$30 :$	$36 ::$	$201.0624 :$	109.6704	at 5 fathoms.
	$36 +$	$60 :$	$36 ::$	$201.0624 :$	75.3984	at 10 fathoms.
	$36 +$	$90 :$	$36 ::$	$201.0624 :$	54.8352	at 15 fathoms.
	$36 +$	$120 :$	$36 ::$	$201.0624 :$	46.4	at 20 fathoms.

cub. ft. (*Const. dec.*)
Again, $109.6704 \div .7854 = 139.6363$ - - 5 fathoms.
 $75.3984 \div .7854 = 96.0000$ - - 10 fathoms.
 $54.8352 \div .7854 = 69.8181$ - - 15 fathoms.
 $46.4000 \div .7854 = 59.0769$ - - 20 fathoms.

And $\frac{1}{2}\sqrt{139.6363} = 5.90845$ ft. depth of air within at 5 fath.
 $\frac{1}{2}\sqrt{96.0000} = 4.89898$ ft. - - - - - 10 fath.
 $\frac{1}{2}\sqrt{69.8181} = 4.17786$ ft. - - - - - 15 fath.
 $\frac{1}{2}\sqrt{59.0769} = 4.15692$ ft. - - - - - 20 fath.

	ft.	feet	feet	
But	$8 -$	$5.90845 =$	2.09154	height of water within, at 5 fathoms.
	$8 -$	$4.89898 =$	3.10101	- - - - - 10 fathoms.
	$8 -$	$4.17786 =$	3.82213	- - - - - 15 fathoms.
	$8 -$	$4.15692 =$	4.15692	- - - - - 20 fathoms.

Ans.*

* Differing the tenth part of an inch at the third depth, and considerably less at the other depths, from the altitude found by Dr. HUTTON.

END OF THE PRACTICAL EXERCISES IN PHILOSOPHY.

FLUXIONS.

(Key to Vol. II. page 299.)

PRACTICAL EXAMPLES IN FINDING FLUXIONS.

Ex. 1. The fluxion of a being 0 (page 289), it is
 $a\dot{x}y + ax\dot{y}$, or $a(\dot{x}y + x\dot{y})$. Ans.

Ex. 2. The fluxion of xyz being $\dot{x}yz + x\dot{y}z + xy\dot{z}$, it is
 $b\dot{x}yz + bx\dot{y}z + bxy\dot{z}$, or $b(\dot{x}yz + x\dot{y}z + xy\dot{z})$. Ans.

Ex. 3. $cx \times (ax - cy) = acx^2 - c^2xy$; but the fluxion of acx^2 is
 $2acx\dot{x}$, and the fluxion of c^2xy is $c^2\dot{x}y + c^2x\dot{y}$.

Therefore $2acx\dot{x} - (c^2\dot{x}y + c^2x\dot{y}) = c(2ax\dot{x} - c\dot{x}y - cx\dot{y})$. Ans.

Ex. 4. Ans. $mx^{m-1}y^n\dot{x} + ny^{n-1}x^m\dot{y} = (my\dot{x} + nx\dot{y})x^{m-1}y^{n-1}$.

Ex. 5. Ans. $mx^{m-1}y^n z^r\dot{x} + ny^{n-1}x^m z^r\dot{y} + rz^{r-1}x^m y^n\dot{z}$.

Ex. 6. $(x+y) \times (x-y) = x^2 - y^2$.

Therefore $2x\dot{x} - 2y\dot{y}$. Ans.

Ex. 7. Here the fluxion of $x^2 = 2x\dot{x}$
Multiply by $2a$

PRODUCT $4ax\dot{x}$. Ans.

Ex. 8. The fluxion of x^3 is $3x^2\dot{x}$
Multiply by 2

PRODUCT $6x^2\dot{x}$. Ans.

(Key to Vol. II. page 299.)

Ex. 9. The fluxion of x^4y is $4x^3y\dot{x} + x^4\dot{y}$
 Multiply by 3

$$\text{PRODUCT } \underline{12x^3y\dot{x} + 3x^4\dot{y}. \text{ Ans.}}$$

Ex. 10. The fluxion of $x^{\frac{2}{3}}y^4$ is $\frac{2}{3}x^{-\frac{1}{3}}y^4\dot{x} + 4x^{\frac{2}{3}}y^3\dot{y}$
 Multiply by 4

$$\text{PRODUCT } \underline{2\frac{2}{3}x^{-\frac{1}{3}}y^4\dot{x} + 16x^{\frac{2}{3}}y^3\dot{y}. \text{ Ans.}}$$

Ex. 11. The fluxion of ax^2y is $2axy\dot{x} + ax^2\dot{y}$. And
 the fluxion of $x^{\frac{1}{2}}y^3$ is $\frac{1}{2}x^{-\frac{1}{2}}y^3\dot{x} + 3x^{\frac{1}{2}}y^2\dot{y}$. Therefore

$$2axy\dot{x} + ax^2\dot{y} - \frac{1}{2}x^{-\frac{1}{2}}y^3\dot{x} - 3x^{\frac{1}{2}}y^2\dot{y}, \text{ or}$$

$$\left(2axy - \frac{\frac{1}{2}y^3}{\sqrt{x}}\right)\dot{x} + (ax^2 - 3\sqrt{xy^2})\dot{y}. \text{ Ans.}$$

Ex. 12. The fluxion of $4x^4$ is $16x^3\dot{x}$.
 - - - - - of x^2y is $2xy\dot{x} + x^2\dot{y}$. And
 - - - - - of $3byz$ is $3by\dot{z} + 3byz$. Therefore
 $(16x^3 - 2xy)\dot{x} + (3bz - x^2)\dot{y} + 3by\dot{z}$. Ans.

$$\text{Ex. 13. Ans. } \frac{1}{n} x^{\frac{1-n}{n}} \dot{x}.$$

$$\text{Ex. 14. Ans. } \frac{m}{n} x^{\frac{m-n}{n}} \dot{x}.$$

$$\text{Ex. 15. Ans. } -\frac{m}{n} x^{\frac{-m+n}{n}} \dot{x} = \frac{-\frac{m}{n}\dot{x}}{x^{\frac{m+n}{n}}}.$$

$$\text{Ex. 16. Ans. } \frac{1}{2}x^{-\frac{1}{2}}\dot{x} = \frac{\frac{1}{2}\dot{x}}{\sqrt{x}} = \frac{\dot{x}}{2\sqrt{x}}.$$

$$\text{Ex. 17. Ans. } \frac{1}{3}x^{-\frac{2}{3}}\dot{x} = \frac{\frac{1}{3}\dot{x}}{\sqrt[3]{x^2}} = \frac{\dot{x}}{3x^{\frac{2}{3}}}.$$

(Key to Vol. II. page 299.)

Ex. 18. Ans. $\frac{2}{3}x^{-\frac{1}{3}}\dot{x} = \frac{\frac{2}{3}\dot{x}}{\sqrt[3]{x}} = \frac{\dot{x}}{1\frac{1}{2}x^{\frac{1}{2}}}$.

Ex. 19. Ans. $\frac{3}{2}x^{\frac{1}{2}}\dot{x} = 1\frac{1}{2}\sqrt{xx}$.

Ex. 20. Ans. $\frac{3}{4}x^{-\frac{1}{4}}\dot{x} = \frac{\frac{3}{4}\dot{x}}{\sqrt[4]{x}} = \frac{\dot{x}}{1\frac{1}{3}\sqrt[4]{x}}$.

Ex. 21. Ans. $1\frac{1}{3}x^{\frac{1}{3}}\dot{x} = \frac{4}{3}\sqrt[3]{xx}$.

Ex. 22. Ans. $\frac{1}{2}(a^2+x^2)^{-\frac{1}{2}} \times 2x\dot{x} = \frac{x\dot{x}}{\sqrt{a^2+x^2}}$.

Ex. 23. Ans. $\frac{1}{2}(a^2-x^2)^{-\frac{1}{2}} \times -2x\dot{x} = \frac{-x\dot{x}}{\sqrt{a^2-x^2}}$.

Ex. 24. The fluxion of $2rx-x^2$ being $2r\dot{x}-2x\dot{x}$, or $2(r-x)\dot{x}$, it is $\frac{1}{2}(2rx-x^2)^{-\frac{1}{2}} \times 2(r-x)\dot{x}$, or

$$\frac{(r-x)\dot{x}}{\sqrt{2rx-x^2}}. \text{ Ans.}$$

Ex. 25. Ans. $-\frac{1}{2}(a^2-x^2)^{-\frac{3}{2}} \times -2x\dot{x} = \frac{x\dot{x}}{\sqrt{(a^2-x^2)^3}}$.

Ex. 26. The fluxion of $ax-x^2$ being $a\dot{x}-2x\dot{x}$, or $(a-2x)\dot{x}$, it is $\frac{1}{3}(ax-x^2)^{-\frac{2}{3}} \times (a-2x)\dot{x}$, or

$$\frac{(a-2x)\dot{x}}{3\sqrt[3]{ax-x^2}^2}. \text{ Ans.}$$

(Page 300.)

Ex. 27. Ans. $\frac{2(\pm x\dot{x})\dot{x}}{\sqrt{a^2 \pm x^2}} = \frac{\pm 2x\dot{x}^2}{(a^2 \pm x^2)^{\frac{1}{2}}}$.

Ex. 28. Ans. $\frac{3}{2}(a^2-x^2)^{\frac{1}{2}} \times -2x\dot{x} = -3x(a^2-x^2)^{\frac{1}{2}}\dot{x}$.

(Key to Vol. II. page 300.)

Ex. 29. Ans. $\frac{xz+x\dot{z}}{2\sqrt{xz}}$.

Ex. 30. The fluxion of $xz - z^2$ being $xz+x\dot{z}-2z\dot{z}$, or

$(x-2z)\dot{z}+z\dot{x}$, it is $\frac{1}{2}(xz-z^2)^{-\frac{1}{2}} \times z\dot{x}+(x-2z)\dot{z}$, that is,

$$\frac{z\dot{x}+(x-2z)\dot{z}}{2\sqrt{xz-z^2}} \quad \text{Ans.}$$

Ex. 31. Ans. $\frac{1}{2a} \times x^{-\frac{3}{2}} \times -\dot{x} = \frac{-\dot{x}}{2ax^{\frac{3}{2}}}$.

Ex. 32. The fluxion of the numerator drawn into the denominator is $3ax^2\dot{x}(a+x) = 3a^2x^2\dot{x}+3ax^3\dot{x}$. And the fluxion of the denominator drawn into the numerator is $ax^3\dot{x}$.

$$\text{Hence } \frac{(3a^2x^2+2ax^3)\dot{x}}{a^2+2ax+x^2} \quad \text{Ans.}$$

Ex. 33. Here the fluxion of the numerator drawn into the denominator is $mx^{m-1}y^n\dot{x}$. And the fluxion of the denominator drawn into the numerator is $nx^m y^{n-1}\dot{y}$.

$$\text{Hence } \frac{mx^{m-1}y^n\dot{x}-nx^m y^{n-1}\dot{y}}{y^{2n}} \quad \text{Ans.}$$

Ex. 34. The fluxion of the numerator drawn into the denominator is $z(xy+x\dot{y})$. And the fluxion of the denominator drawn into the numerator is xyz .

$$\text{Therefore } \frac{z(xy+x\dot{y})-xyz}{z^2} \quad \text{Ans.}$$

Ex. 35. $\frac{c}{xx} = cx^{-2}$. Consequently $-2cx^{-3}\dot{x}$, or $\frac{-2c\dot{x}}{x^3}$. Ans.

Or, without the negative index ;

The fluxion of the numerator is 0, which drawn into the denominator is 0; and the fluxion of the denominator is $2x\dot{x}$, which drawn into the numerator is $2cx\dot{x}$: this subtracted from 0 leaves $-2cx\dot{x}$: lastly $\frac{-2cx\dot{x}}{x^4} = \frac{-2c\dot{x}}{x^3}$ as before.

(Key to Vol. II. page 300.)

Ex. 36. In this Example the rectangle under the denominator and fluxion of the numerator is $3(a-x)\dot{x} = 3a\dot{x} - 3x\dot{x}$; and the rectangle under the numerator and fluxion of the denominator is $3x\dot{x}$. Wherefore $\frac{(3a-6x)\dot{x}}{(a-x)^2}$. Ans.

Ex. 37. The rectangle under the denominator and fluxion of the numerator is $(x+z)\dot{z} = x\dot{z} + z\dot{z}$; and the rectangle under the numerator and fluxion of the denominator is $z(\dot{x} + \dot{z}) = z\dot{x} + z\dot{z}$.

Therefore $\frac{\dot{z}x - z\dot{x}}{(x+z)^2}$. Ans.

Ex. 38. Here the fluxion of the numerator drawn into the denominator is $2xz^2\dot{x}$; and the fluxion of the denominator drawn into the numerator is $2x^2z\dot{z}$. Consequently

$\frac{2(xz^2\dot{x} - x^2z\dot{z})}{z^4}$, or $\frac{xz\dot{x} - x^2\dot{z}}{\frac{1}{2}z^3}$. Ans.

Ex. 39. The fluxion of $x^{\frac{2}{3}}$ is $\frac{2}{3}x^{-\frac{1}{3}}\dot{x}$. And the fluxion of $y^{\frac{3}{2}}$ is $\frac{3}{2}\sqrt{y}\dot{y}$.

Wherefore $\frac{\frac{2}{3}y^{\frac{3}{2}}x^{-\frac{1}{3}}\dot{x} - \frac{3}{2}x^{\frac{2}{3}}y^{\frac{1}{2}}\dot{y}}{y^3}$. Ans.

Otherwise.

$\frac{x^{\frac{2}{3}}}{y^{\frac{3}{2}}} = x^{\frac{2}{3}}y^{-\frac{3}{2}}$ of which the fluxion is

$\frac{2}{3}x^{-\frac{1}{3}}y^{-\frac{3}{2}}\dot{x} - \frac{3}{2}x^{\frac{2}{3}}y^{-\frac{5}{2}}\dot{y} = \frac{\frac{2}{3}\dot{x}}{x^{\frac{1}{3}}y^{\frac{3}{2}}} - \frac{\frac{3}{2}x^{\frac{2}{3}}\dot{y}}{y^{\frac{5}{2}}} = \frac{\frac{2}{3}y^{\frac{5}{2}}\dot{x} - \frac{3}{2}xy^{\frac{3}{2}}\dot{y}}{x^{\frac{1}{3}}y^4} =$

$\frac{\frac{2}{3}y^{\frac{3}{2}}\dot{x} - \frac{3}{2}xy^{\frac{1}{2}}\dot{y}}{x^{\frac{1}{3}}y^3} = \frac{\frac{2}{3}y^{\frac{3}{2}}x^{-\frac{1}{3}}\dot{x} - \frac{3}{2}x^{\frac{2}{3}}y^{\frac{1}{2}}\dot{y}}{y^3}$ as before.

Ex. 40. The fluxion of axy^2 is $a\dot{x}y^2 + 2axy\dot{y}$; and the fluxion of z is \dot{z} .

Therefore $\frac{az(\dot{x}y^2 + 2xy\dot{y}) - axy^2\dot{z}}{z^2}$. Ans.

(Key to Vol. II. page 300.)

Ex. 41. $\frac{3}{\sqrt{(x^2-y^2)}} = 3(x^2-y^2)^{-\frac{1}{2}}$; and the fluxion of x^2-y^2 is $2x\dot{x}-2y\dot{y}$. Therefore $- \frac{3}{2}(x^2-y^2)^{-\frac{3}{2}} \times 2(x\dot{x}-y\dot{y}) = \frac{-3(x\dot{x}-y\dot{y})}{(x^2-y^2)^{\frac{3}{2}}}$. Ans.

Otherwise.

The fluxion of 3 is 0; and the fluxion of $\sqrt{x^2-y^2}$ being $\frac{1}{2}(x^2-y^2)^{-\frac{1}{2}} \times 2x\dot{x}-2y\dot{y}$, the rectangle under the numerator and fluxion of the denominator is $\frac{3(x\dot{x}-y\dot{y})}{(x^2-y^2)^{\frac{1}{2}}}$; this subtracted from 0 leaves $\frac{-3(x\dot{x}-y\dot{y})}{(x^2-y^2)^{\frac{1}{2}}}$, which, divided by the square of $\sqrt{x^2-y^2} = x^2-y^2$, becomes $\frac{-3(x\dot{x}-y\dot{y})}{(x^2-y^2)^{\frac{3}{2}}}$. as before.

Ex. 42. Here the fluxion of the quantity is $a\dot{x}$, which divided by the quantity becomes $\frac{a\dot{x}}{ax} = \frac{\dot{x}}{x}$. Ans.

Ex. 43. In this Example the fluxion of the quantity is \dot{x} , which divided by the quantity becomes $\frac{\dot{x}}{1+x}$. Ans.

Ex. 44. Ans. $\frac{\dot{x}}{1-x}$.

Ex. 45. Ans. $\frac{2x\dot{x}}{x^2} = \frac{2\dot{x}}{x} = \frac{\dot{x}}{\frac{1}{2}x}$.

Ex. 46. Ans. $\frac{\frac{1}{2}z^{-\frac{1}{2}}\dot{z}}{z^{\frac{1}{2}}} = \frac{\frac{1}{2}\dot{z}}{z} = \frac{\dot{z}}{2z}$.

Ex. 47. Ans. $\frac{mx^{m-1}\dot{x}}{x^m} = \frac{m\dot{x}}{x} = \frac{\dot{x}}{\frac{1}{m}x}$.

(Key to Vol. II. page 301.)

Ex. 48. $\frac{2}{x^2} = 2x^{-2}$, of which the fluxion is $-4x^{-3}\dot{x} = \frac{-4\dot{x}}{x^3}$,

and this fluxion divided by $\frac{2}{x^2} = \frac{-2\dot{x}}{x}$. Ans.

Ex. 49. The fluxion of $\frac{1+x}{1-x}$ is $\frac{-2x\dot{x}}{(1-x)^2}$, which divided by $\frac{1+x}{1-x}$

becomes $\frac{-2x\dot{x}}{1-x^2}$. Ans.

Ex. 50. The fluxion of $\frac{1-x}{1+x}$ is $\frac{2x\dot{x}}{(1+x)^2}$, which divided by $\frac{1-x}{1+x}$

becomes $\frac{2x\dot{x}}{1-x^2}$. Ans.

Ex. 51. Ans: $c^x \dot{x} \times \text{hyp. log. of } c$. [See Art. 25, page 295, vol. ii. of the Mathematics.]

Ex. 52. Ans. $10^x \dot{x} \times (2.3025850929940456840179914 \text{ \&c. the hyperbolic logarithm of } 10.)$

Ex. 53. Ans. $(a+c)^x \dot{x} \times \text{hyp. log. of } (a+c)$.

Ex. 54. Ans. $100^{xy} \times (xy + x\dot{y}) \times (4.605170185988091368 \text{ \&c. the hyperbolic logarithm of } 100.)$

Ex. 55. The fluxion of the given quantity considering *only* the root variable is $zx^{z-1}\dot{x}$, and the fluxion of x^z considering *only* the exponent variable is $x^z \dot{z} \times \text{hyp. log. of } x$. The sum of these two fluxions is

$$zx^{z-1}\dot{x} + x^z \dot{z} \times \text{hyp. log. of } x. \text{ Ans.}$$

Ex. 56. The fluxion of y^{10x} considering the exponent constant is $10xy^{10x-1}\dot{y}$, and the fluxion of y^{10x} regarding the root constant is $10y^{10x}\dot{x} \times \text{hyp. log. of } y$; wherefore $10xy^{10x-1}\dot{y} + 10y^{10x}\dot{x} \times \text{hyp. log. of } y$. Ans.

Ex. 57. Ans. $x^x \dot{x} + x^x \dot{x} \times \text{hyp. log. of } x$, or $(1 + \text{hyp. log. of } x) x^x \dot{x}$.

Ex. 58. The fluxion of $(xy)^{xz}$, the exponent considered constant, is $xz(xy)^{xz-1} \times (x\dot{y} + y\dot{x})$, and the fluxion of $(xy)^{xz}$, considering the root constant, is $(xy)^{xz} \times (x\dot{z} + xz) \times \text{hyp. log. of } (xy)$. Therefore $xz(xy)^{xz-1} \times (x\dot{y} + y\dot{x}) + (xy)^{xz} \times (x\dot{z} + xz) \times \text{hyp. log. of } (xy)$. Ans.

(Key to Vol. II. page 301.)

Ex. 59. Ans. $\dot{x}y \pm x\ddot{y}$.Ex. 60. Ans. $\ddot{x}y^2 \pm 2x\dot{y}^3$.For $\dot{x}y^2 = x\dot{y}y$, of which the fluxion is $\ddot{x}y + x\ddot{y} + x\dot{y}\dot{y} = \ddot{x}y^2 + 2x\dot{y}^3$.Ex. 61. The first fluxion of xy is $\dot{x}y + x\dot{y}$, of which the fluxion is $\ddot{x}y + \dot{x}\dot{y} + x\ddot{y} + x\dot{y}\dot{y} = \ddot{x}y + 2\dot{x}\dot{y} + x\ddot{y}$. Ans.Ex. 62. The first fluxion of xy being as in the last Example $\dot{x}y + x\dot{y}$, it will be $\ddot{x}y + \dot{x}\dot{y} + x\ddot{y} = 2\dot{x}\dot{y} + x\ddot{y}$. Ans.Ex. 63. The first fluxion of x^n is $nx^{n-1}\dot{x}$, and the fluxion of this fluxion is $(n^2 - n)x^{n-2}\dot{x}^2 + nx^{n-1}\ddot{x}$. Ans.Ex. 64. The first fluxion of x^n being, as in the last Example, $nx^{n-1}\dot{x}$, it will be $(n^2 - n)x^{n-2}\dot{x} \times \dot{x}$ for the second fluxion, and $(n^3 - 3n^2 + 2n)x^{n-3}\dot{x}^3$ for the third fluxion, \dot{x} being constant. Ans.Ex. 65. The first fluxion of xy is $\dot{x}y + x\dot{y}$, of which the fluxion is $\ddot{x}y + 2\dot{x}\dot{y} + x\ddot{y}$; and the fluxion of this fluxion is $\ddot{\ddot{x}}y + \ddot{x}\dot{y} + 2\ddot{x}\dot{y} + 2x\ddot{\dot{y}} + x\dot{y}\dot{y} = \ddot{\ddot{x}}y + 3\ddot{x}\dot{y} + 3x\ddot{\dot{y}} + x\dot{y}\dot{y}$. Ans.

FLUENTS.

(Page 303.)

Ex. 8. Ans. y^n .Ex. 9. Ans. $\frac{-1}{z}$.Ex. 10. $\frac{a\dot{y}}{y^n} = ay^{-n}\dot{y}$, of which the fluent is $\frac{1}{y^{n-1}}a$. Ans.Ex. 11. Ans. $\frac{1}{3}(a+x)^5$.Ex. 12. Ans. $\frac{1}{8}(a^4+y^4)^2$.Ex. 13. Ans. $\frac{1}{15}(a^3+z^3)^5$.Ex. 14. Ans. $\frac{1}{mn+n}(a^n+x^n)^{m+1}$.

(Key to Vol. II. page 303.)

Ex. 15. Ans. $\frac{1}{8}(a^2+y^2)^4$.

Ex. 16. Ans. $(a^2+z^2)^{\frac{1}{2}}$.

Ex. 17. Ans. $-2\sqrt{a-x}$.

(Page 304.)

When a Compound Root is multiplied by some power of the flowing quantity such that, of this factor the Index plus 1 is a multiple of the Index of the unknown quantity under the vinculum.

Ex. 1. Put z for $a+cx$, so shall $x = \frac{z-a}{c}$; and taking the fluxion

on both sides, $\dot{x} = \frac{\dot{z}}{c}$. Consequently $x^3\dot{x} = \left(\frac{z-a}{c}\right)^3 \times \frac{\dot{z}}{c} =$

$\frac{z^3 - 3az^2 + 3a^2z - a^3}{c^4} \dot{z}$. Hence $(a+cx)^{\frac{1}{2}}x^3\dot{x} = z^{\frac{1}{2}} \times$

$\frac{z^3 - 3az^2 + 3a^2z - a^3}{c^4} \dot{z}$, an expression equivalent to

$\frac{z^{\frac{7}{2}} - 3az^{\frac{5}{2}} + 3a^2z^{\frac{3}{2}} - a^3}{c^4} \dot{z}$, which is equal to $\frac{z^{\frac{7}{2}}\dot{z}}{c^4} - \frac{3az^{\frac{5}{2}}\dot{z}}{c^4} +$

$\frac{3a^2z^{\frac{3}{2}}\dot{z}}{c^4} - \frac{a^3\dot{z}}{c^4}$, of which the fluent is $\frac{2z^{\frac{9}{2}}}{9c^4} - \frac{6az^{\frac{7}{2}}}{7c^4} + \frac{6a^2z^{\frac{5}{2}}}{5c^4} -$

$\frac{a^3z}{c^4}$.

Now substituting $a+cx$ for z , this fluent becomes

$$\frac{\frac{2}{9}(a+cx)^{\frac{9}{2}} - \frac{6}{7}a(a+cx)^{\frac{7}{2}} + \frac{6}{5}a^2(a+cx)^{\frac{5}{2}} - a^3(a+cx)}{c^4}. \text{ Ans.}$$

Ex. 2. Assuming $z = a+cx$ as in the last Example, $x^2\dot{x} =$
 $\left(\frac{z-a}{c}\right)^2 \times \frac{\dot{z}}{c} = \frac{z^2 - 2az + a^2}{c^3} \dot{z}$. Also $(a+cx)^{\frac{3}{4}}x^2\dot{x} = z^{\frac{3}{4}} \times$

$\frac{z^2 - 2az + a^2}{c^3} \dot{z}$, an expression equivalent to $\frac{z^{\frac{11}{4}} - 2az^{\frac{7}{4}} + a^2z^{\frac{3}{4}}}{c^3} \dot{z}$,

(Key to Vol. II. page 304.)

and equal to $\frac{z^{\frac{1}{4}} \dot{z} - 2az^{\frac{7}{4}} \dot{z} + a^2 z^{\frac{9}{4}} \dot{z}}{c^3}$; now of this fluxion the
 fluent is $\frac{\frac{4}{15} z^{\frac{5}{4}} - \frac{8}{9} az^{\frac{9}{4}} + \frac{4}{5} a^2 z^{\frac{13}{4}}}{c^3}$. Substituting therefore $a +$
 cx for z , it is $\frac{\frac{4}{15} (a+cx)^{\frac{5}{4}} - \frac{8}{9} (a+cx)^{\frac{9}{4}} + \frac{4}{5} a^2 (a+cx)^{\frac{13}{4}}}{c^3}$. Ans.

Ex. 3. Let y represent $a + cx^2$, then $x^2 = \frac{y-a}{c}$. But the fluxion of
 this equation is $2x\dot{x} = \frac{\dot{y}}{c}$; or by dividing both sides by 2,
 $x\dot{x} = \frac{\dot{y}}{2c}$: therefore (multiplying the first side by x^2 , and
 the second by its equivalent,) $x^3\dot{x} = (\frac{y-a}{c^2})\dot{y}$. Conse-
 quently $d(a+cx^2)^{\frac{1}{3}} = dy^{\frac{1}{3}} \times (\frac{y-a}{c^2})\dot{y} = \frac{dy^{\frac{4}{3}}\dot{y} - ady^{\frac{1}{3}}\dot{y}}{c^2}$. Now
 of this fluxion the fluent is $\frac{\frac{3}{7} dy^{\frac{7}{3}} - \frac{3}{4} ady^{\frac{4}{3}}}{c^2}$, or, (substituting
 $a+cx^2$, for y), $\frac{\frac{3}{7} d(a+cx^2)^{\frac{7}{3}} - \frac{3}{4} ad(a+cx^2)^{\frac{4}{3}}}{c^2}$. Ans.

Ex. 4. If x be assumed for $a+z$, and their fluxions put equal,
 it will be $\dot{x} = \dot{z}$; therefore $z\dot{z} = x\dot{x}$. Consequently
 $c(a+z)^{-\frac{1}{2}}z\dot{z} = cx^{-\frac{1}{2}}x\dot{x} = cx^{\frac{1}{2}}\dot{x}$, of which the fluent is $\frac{2}{3}cx^{\frac{3}{2}}$.
 This, by substituting $a+z$ for x , becomes $\frac{2}{3}c(a+z)^{\frac{3}{2}}$. Ans.*

* The fluent arising from a given fluxion often requires (as in the present
 Example) a correction to make it cotemporaneous with the fluxion. [See
 page 313, vol. ii.]

Ex. 5. Assuming $x = a + z^n$, and taking the fluxion of the equa-
 tion, $\dot{x} = nz^{n-1}\dot{z}$; also $x - a = z^n$, and by squaring both
 sides, $x^2 - 2ax + a^2 = z^{2n}$. Consequently $(x^2 - 2ax + a^2)\dot{x}$
 $= nz^{3n-1}\dot{z}$, and $(a+z^n)^{-\frac{1}{2}}cz^{3n-1}\dot{z} = x^{-\frac{1}{2}} \times (\frac{x^2 - 2ax + a^2}{n})$
 $\times \dot{x} = \frac{cx^2\dot{x} - 2acx\dot{x} + a^2\dot{x}}{n\sqrt{x}} = \frac{cx^{\frac{3}{2}}\dot{x} - 2acx^{\frac{1}{2}}\dot{x} + a^2cx^{-\frac{1}{2}}\dot{x}}{n}$,

(Key to Vol. II. page 304.)

of which the fluent is $\frac{\frac{2}{5}cx^{\frac{5}{2}} - \frac{4}{3}acx^{\frac{3}{2}} + 2a^2cx^{\frac{1}{2}}}{n}$. Substituting therefore $a+z^n$ for x , it is $\frac{c}{n} \times \left\{ \frac{2}{5}(a+z^n)^{\frac{5}{2}} - \frac{4}{3}a(a+z^n)^{\frac{3}{2}} + 2a^2(a+z^n)^{\frac{1}{2}} \right\}$ Ans.

Ex. 6. Ans. $\frac{\sqrt{a^2+z^2}^{\frac{3}{2}} \times 2z^2 - 3a^2}{15a^4z^5}$.

Ex. 7. Ans. $\frac{\sqrt{a-x^n}^{\frac{3}{2}}}{-\frac{7}{2}nax^{\frac{7}{2}n}} \times 1 + \frac{4x^n}{5a} + \frac{8x^{2n}}{15a^2}$ which reduced is $\frac{\sqrt{a-x^n}^{\frac{3}{2}} \times 30a^2 + 24ax^n + 16x^{2n}}{105na^3x^{\frac{7}{2}n}}$.

When there are several Terms involving Two or more variable Quantities.

Ex. 3. Ans. $x^{\frac{1}{2}}y^2$.

Ex. 4. Ans. $\frac{x}{y}$.

Ex. 5. Ans. $ax^2y^{-\frac{1}{2}}$.

(Page 305.)

When the Fluxion of a Quantity is divided by the Quantity.

Ex. 4. Ans. The hyp. log. of $a+x$.

Ex. 5. Ans. The hyp. log. of $a+x^3$.

(Page 309.)

The Comparison of Fluents.

Ex. 4. The proposed fluxion is found to agree with the 1st FORM; and $x^4\dot{x} = x^{n-1}\dot{x}$; hence $n = 5$.

Therefore $\frac{ax^5}{5}$, or $\frac{a}{5}x^5$. Ans.

(Key to Vol. II. page 309.)

Ex. 5. This fluxion corresponds with the 2nd FORM,

Wherein $x^2 = x^n$, and $n = 2$.Also $m - 1 = \frac{2}{3}$, or $m = \frac{5}{3}$.Therefore $2 \times \frac{3}{10} (10 + x^2)^{\frac{5}{3}}$, or

$$\frac{3}{5} (10 + x^2)^{\frac{5}{3}}. \text{ Ans.}$$

Ex. 6. There being numerous typographical errors in the fluxional part of Hutton's Course, it is imagined the present Example was misprinted. If the fluent of $\frac{ax}{(c^2 + x^2)^{\frac{3}{2}}}$ be really required, it is $\frac{a}{c} \times \text{arc to tang. } \frac{1}{c} \sqrt{x}$, or $\frac{a}{2c} \times$ arc to cos. $\frac{c^2 - x^2}{c^2 + x^2}$; according to the 11th FORM.*

* In the 11th Form tang. $\sqrt{\frac{x}{a}}$ should be tang. $\sqrt{\frac{x^n}{a}}$, and consequently the Answer to this Example, as well as to Examples 18 and 19, and to all others by the 11th Form are in part incorrect. See KEY, page 295.

Ex. 7. In this Example the fluxion agrees with the 3d FORM, and $x^2 = x^{m \cdot n - 1}$; hence $m = 3$, and $n = 1$.

$$\text{Consequently } \frac{3}{4} \times \frac{1}{3} a \times \frac{x^3}{(a-x)^3}, \text{ or } \frac{ax^3}{(a-x)^3}. \text{ Ans.}$$

(Page 310.)

Ex. 8. Here the fluxion is of the 4th FORM, also $m = n = 2$; wherefore the fluent of $\frac{(c^2 - x^2)^{\frac{1}{2}} \dot{x}}{x^5}$ is $\frac{-1}{4c^2} \times \frac{(c^2 - x^2)^2}{x^4} =$

$$\frac{-c^2}{4x^4} + \frac{1}{2x^2} - \frac{1}{4}c^2. \text{ Ans.}$$

Ex. 9. Ans. $\frac{-1}{6x^3} - \frac{3}{4x^2}$; by the 4th FORM.Ex. 10. Ans. $x^3 y^2$; by the 5th FORM.Ex. 11. Ans. $xy^{\frac{1}{3}}$; by the 5th FORM.Ex. 12. Ans. $\frac{3}{a} \times \text{hyp. log. of } x$; by the 7th FORM.Ex. 13. Ans. $-\frac{1}{2}a \dagger \times \text{hyp. log. of } 3 - 2x$; by the 8th FORM.

† Because the fluent of $\frac{ax}{3-2x} = -\frac{1}{2}a \times \text{fluent of } \frac{-2\dot{x}}{3-2x}$.

(Key to Vol. II. page 310.)

Ex. 14.* Ans. $\frac{3}{2}$ hyp. log. of $\frac{x}{2-x}$, by the 9th FORM, which is likewise misprinted, and ought to be $\frac{x^{-1}x}{a \pm x^n}$.

* There is a typographical error in the denominator of this Example in every edition of the COURSE.

Ex. 15.† The given Example, when corrected, is equal to $\frac{2}{3} \times \frac{x^{-1}x}{\frac{1}{3}-x^2}$, a FORM agreeing with the 9th in the Table; therefore the hyp. log. of $\frac{x^2}{\frac{1}{3}-x^2}$. Ans.

† The latter part of this Example is misprinted in the Mathematics. It ought to be $\frac{2x^{-1}x}{1-3x^2}$ instead of $\frac{2x^{-1}3x^2}{1-3x^2}$.

Ex. 16. Ans. $\frac{3}{4}$ hyp. log. of $\frac{1+x^2}{1-x^2}$, by the 10th FORM, † a being 1 and $n=4$.

† Which in every copy we have seen of HUTTON is misprinted. It ought to be $\frac{x^{\frac{1}{2}n-1}}{a-x^n}$

Ex. 17. Ans. $\frac{a}{1} \times \frac{1}{5\sqrt{2}} \times$ hyp. log. of $\frac{\sqrt{2+\sqrt{x^5}}}{\sqrt{2-\sqrt{x^5}}}$, by the 10th FORM, n being 5, and a in the form = 2.

Ex. 18. Ans. $\left. \begin{array}{l} \frac{2}{1} \times \frac{2}{4\sqrt{1}} \times \text{arc to tan. } \sqrt{\frac{x^{\dagger}}{1}} \\ \text{Or, } \frac{2}{1} \times \frac{1}{4\sqrt{1}} \times \text{arc to cos. } \frac{1-x^4}{1+x^4} \end{array} \right\} \text{by the 11th FORM.}$

$\left. \begin{array}{l} \text{That is, The arc to the tangent of } x^{\frac{1}{2}}; \\ \text{or, half the arc to the cosine of } \frac{1-x^4}{1+x^4}. \end{array} \right\} \text{by reduction.}$

† See the note in page 295 of the KEY.

(Key to Vol. II. page 310.)

Ex. 19. Ans. $\frac{a}{1} \times \frac{2}{5\sqrt{2}} \times \text{arc to tang. } \sqrt{\frac{x}{2}};^*$ } *by FORM II.*
 or, $\frac{a}{1} \times \frac{1}{5\sqrt{2}} \times \text{arc to cos. } \frac{2-x^5}{2+x^5}.$ }
 And, $\frac{1}{5} a\sqrt{2} \times \text{arc to tang. } \frac{1}{2}\sqrt{2x};$ } *by reduction.*
 or, $\frac{1}{10} a\sqrt{2} \times \text{arc to cos. } \frac{2-x^5}{2+x^5}$ }

* See the note referred to in the last Example.

Ex. 20. Ans. $\frac{6}{4} \times \text{hyp. log. of } \sqrt{x^4 + \sqrt{1+x^4}},$ *by the 12th FORM.*
 and, $\frac{3}{2} \text{ hyp. log. } x^2 + \sqrt{1+x^4},$ *by reduction.*

Ex. 21. Changing the given expression to its equivalent $\frac{ax^{\circ}x}{\sqrt{-4+x^2}} = \frac{a}{1} \times \frac{x^{\circ}x}{\sqrt{-4+x^2}},$ the FORM agrees with the 12th in the Table, n being 2, and $a = -4.$ Therefore $a \times \text{hyp. log. of } x + \sqrt{x^2-4}.$ Ans.

NOTE. x° is always equivalent to unity, which may be proved thus; $x^{\circ} = x^{1-1} = x^1 \times x^{-1} = \frac{x}{1} \times \frac{1}{x} = \frac{x}{x} = 1.$

Ex. 22. Ans. $\frac{3}{2} \times \text{arc to sin. } x^2,$ or $\frac{3}{4} \times \text{arc to vers. } 2x^4,$ *by the 13th FORM* n being 4 and $a=1.$

Ex. 23. Ans. $a \times \text{arc to sin. } \frac{1}{2}x,$ or $\frac{1}{2}a \times \text{arc to vers. } \frac{1}{2}x^2,$ *by the 13th FORM,* a in the FORM being 4 and $n=2.$ See Example 21 above, or Examples 6 and 13, page 286.

Ex. 24. *By the 14th FORM,* n being 2, and $a=1,$ the fluent is the hyp. log. of $\frac{1-\sqrt{1-x^2}}{1+\sqrt{1-x^2}} = 2 \text{ hyp. log. } \frac{1-\sqrt{1-x^2}}{x}.$

Ex. 25. $\frac{ax}{\sqrt{ax^2+x^{\frac{1}{2}}}} = \frac{ax}{\sqrt{x^{\frac{1}{2}}\sqrt{ax^{\frac{3}{2}}+1}}} = \frac{ax^{-\frac{1}{4}}x}{\sqrt{ax^{\frac{3}{2}}+1}} = \frac{ax^{-\frac{1}{4}}x}{\sqrt{a\sqrt{x^{\frac{3}{2}}+a^{-1}}}}$

(Key to Vol. II. page 310.)

$$= \frac{a^{\frac{1}{2}} x^{-\frac{1}{4}} \dot{x}}{\sqrt{\frac{1}{a} + x^{\frac{3}{2}}}} = \sqrt{a} \times \frac{x^{-\frac{1}{4}} \dot{x}}{\sqrt{a^{-1} + x^{\frac{3}{2}}}}, \text{ a form agreeing with the}$$

12th in the Table, n being $\frac{3}{2}$, and a under the vinculum in the Table $= \frac{1}{a}$ in the proposed Example; therefore

$$\frac{4}{3} \sqrt{a} \times \text{hyp. log. of } \sqrt{x^{\frac{3}{2}} + \sqrt{a^{-1} + x^{\frac{3}{2}}}}. \text{ Ans.}$$

Otherwise,

By the 14th FORM. The given fluxion divided by x becomes

$$\frac{ax^{-1} \dot{x}}{\sqrt{a+x^{-\frac{3}{2}}}}, \text{ of which the fluent is } -\frac{2}{3} \sqrt{a} \times \text{hyp. log. } \frac{\sqrt{a+x^{-\frac{3}{2}}} - \sqrt{a}}{\sqrt{a+x^{-\frac{3}{2}}} + \sqrt{a}},$$

reducible to simpler terms.

Ex. 26. Correcting the typographical error in the denominator, &c. and dividing the numerator by 2, the fluxion becomes

$$\frac{x^{-1} \dot{x}}{\sqrt{x^2-1}}, \text{ agreeing with the 15th FORM. Therefore twice}$$

$$\text{the arc to secant } x, \text{ or the arc to cosine } \frac{2-x^2}{x^2}. \text{ Ans.}$$

(Page 311.)

$$\text{Ex. 27. } \frac{ax}{\sqrt{x^{\frac{1}{2}} - ax^2}} = \frac{ax}{\sqrt{x^{\frac{1}{2}} \sqrt{-ax^{\frac{3}{2}} + 1}}} \text{ which is equal to}$$

$$\sqrt{a} \times \frac{x^{-\frac{1}{4}} \dot{x}}{\sqrt{-x^{\frac{3}{2}} + a^{-1}}}, \text{ a form agreeing with the 13th of}$$

$$\text{the Table; therefore } \left. \begin{array}{l} \frac{4}{3} \sqrt{a} \times \text{arc to sin. } \sqrt{a^{-1} x^{\frac{3}{2}}}, \\ \text{or, } \frac{2}{3} \sqrt{a} \times \text{arc to vers. } 2a^{-1} x^{\frac{3}{2}} \end{array} \right\} \text{ Ans.}$$

Ex. 28. Ans. Circular segment to diameter 2 and versed sine x , by the 16th FORM.

Ex. 29. Ans. $\frac{a^x}{\text{hyp. log. of } a}$, by the 17th FORM, n being 1.

Ex. 30. Ans. $\frac{3a^{2x}}{\text{twice hyp. log. of } a}$, by the 17th FORM, n being 2.

Ex. 31. Ans. $3z^x$, by the 18th FORM.

(Key to Vol. II. page 311.)

Ex. 32. $(1+x^3)x\dot{x} = (x+x^4)\dot{x} = x\dot{x} + x^4\dot{x}$, of which the fluent by the 1st FORM is $\frac{1}{2}x^2 + \frac{1}{5}x^5$. Or if the given fluxion be $(1+x^3)x^2\dot{x}$ the fluent is $\frac{1}{3}x^3 + \frac{1}{6}x^6$. Ans.

Ex. 33. $(2+x^4)x^{\frac{3}{2}}\dot{x} = (2x^{\frac{3}{2}} + x^{\frac{11}{2}})\dot{x} = 2x^{\frac{3}{2}}\dot{x} + x^{\frac{11}{2}}\dot{x}$, of which the fluent by the 1st FORM is $\frac{4}{5}x^{\frac{5}{2}} + \frac{2}{13}x^{\frac{13}{2}}$. Ans.

Ex. 34. $x^2\dot{x}\sqrt{a^2+x^2} = \frac{x^2\dot{x}(a^2+x^2)}{\sqrt{a^2+x^2}} = \frac{a^2x^2\dot{x}}{\sqrt{a^2+x^2}} + \frac{x^4\dot{x}}{\sqrt{a^2+x^2}}$.
 Multiplying both numerator and denominator of the first part by x , and of the second by x^3 , the result is $\frac{a^2x^3\dot{x}}{\sqrt{a^2x^2+x^4}} + \frac{x^7\dot{x}}{\sqrt{a^2x^6+x^8}} = x^2\dot{x}\sqrt{a^2+x^2}$.

Thus transformed, the quantity without the vinculum is a constant part of the fluxion of the highest term under the vinculum; namely in the first fluxion, $\frac{1}{4}$ th part; and in the second, $\frac{1}{8}$ th part.

Adding therefore to the numerator in the one instance $\frac{1}{4}$ th of, and in the other $\frac{1}{8}$ th of, the fluxion of the leading term respectively under the vinculum, there arise

$$a^2 \times \frac{(\frac{1}{2}a^2x\dot{x} + x^3\dot{x})}{\sqrt{a^2x^2+x^4}} \text{ and } \frac{\frac{3}{4}a^2x^5\dot{x} + x^7\dot{x}}{\sqrt{a^2x^6+x^8}}, \text{ two perfect fluxions.}$$

Now the fluents of these by the 2nd FORM are $\frac{1}{2}a^2\sqrt{a^2x^2+x^4}$ and $\frac{1}{4}\sqrt{a^2x^6+x^8}$; the former equal to $\frac{1}{2}a^2x\sqrt{a^2+x^2}$, the latter equal to $\frac{1}{4}x^3\sqrt{a^2+x^2}$. But from the former there must be deducted the fluent of $\frac{\frac{1}{2}a^2x\dot{x}}{\sqrt{a^2x^2+x^2}}$ which by the 12th

FORM is $\frac{1}{2}a^2 \times \text{hyp. log. of } x + \sqrt{a^2+x^2}$; and from the latter, the fluent of $\frac{\frac{3}{4}a^2x^5\dot{x}}{\sqrt{a^2x^6+x^8}}$, an expression equivalent

to $\frac{\frac{3}{4}a^2x^2\dot{x}}{\sqrt{a^2+x^2}}$, and consequently its fluent (see the fluent

just found,) $\frac{3}{4}a^2 \times (\frac{1}{2}x\sqrt{a^2+x^2} - \frac{1}{2}a^2 \times \text{hyp. log. of } x + \sqrt{a^2+x^2})$. Therefore $\frac{1}{2}a^2(x\sqrt{a^2+x^2} - \text{hyp. log. of } x + \sqrt{a^2+x^2}) + \left\{ (\frac{1}{4}x^3\sqrt{a^2+x^2} - \frac{3}{4}a^2(x\sqrt{a^2+x^2} - a^2 \times$

(Key to Vol. II. page 312.)

$$\text{hyp. log. of } \sqrt{a^2 + x^2} \left. \vphantom{\sqrt{a^2 + x^2}} \right\} \text{ Or } \left(\frac{1}{4}x^3 + \frac{1}{8}a^2x \right) \times \sqrt{a^2 + x^2} - \frac{1}{8}a^4 \times \text{hyp. log. of } (x + \sqrt{a^2 + x^2}). \text{ Ans.}$$

To find Fluents by Infinite Series.

Ex. 1.

For the ascending series.

$$a-x) \quad bx \left(\frac{bx}{a} + \frac{bx^2}{a^2} + \frac{bx^3}{a^3} + \frac{bx^4}{a^4} + \frac{bx^5}{a^5} = \frac{b}{a} \times \left(x + \frac{x^2}{a} + \frac{x^3}{a^2} + \frac{x^4}{a^3} + \&c. \right) \text{ Wherefore}$$

$$\frac{bx\dot{x}}{a-x} = \frac{b}{a} \left(x\dot{x} + \frac{x^2\dot{x}}{a} + \frac{x^3\dot{x}}{a^2} + \frac{x^4\dot{x}}{a^3} + \frac{x^5\dot{x}}{a^4} + \&c. \right) \text{ of which}$$

$$\text{the fluent is } \frac{b}{a} \left(\frac{x^2}{2} + \frac{x^3}{3a} + \frac{x^4}{4a^2} + \frac{x^5}{5a^3} + \frac{x^6}{6a^4} + \&c. \right) \text{ Ans.}$$

And for the descending series.

$$-x+a) \quad bx \left(-b - \frac{ab}{x} - \frac{a^2b}{x^2} - \frac{a^3b}{x^3} - \frac{a^4b}{x^4} - \frac{a^5b}{x^5} \&c. \right)$$

$$\text{Therefore } \frac{bx\dot{x}}{a-x} = -bx - \frac{abx}{x} - \frac{a^2bx}{x^2} - \frac{a^3bx}{x^3} - \frac{a^4bx}{x^4} - \&c.$$

$$\text{of which the fluent is } -bx, -ab \times \text{hyp. log. of } x, +a^2bx^{-1} + \frac{1}{2}a^3bx^{-2} + \frac{1}{3}a^4bx^{-3} + \&c. \text{ Ans.}$$

Ex. 2.

For the ascending series.

$$a+x) \quad b \left(\frac{b}{a} - \frac{bx}{a^2} - \frac{bx^2}{a^3} - \frac{bx^3}{a^4} - \&c. = \frac{b}{a} \times \left(1 - \frac{x}{a} - \frac{x^2}{a^2} - \frac{x^3}{a^3} - \&c. \right) \right)$$

$$\text{Wherefore } \frac{bx\dot{x}}{a+x} = \frac{b}{a} \left(\dot{x} - \frac{x\dot{x}}{a} - \frac{x^2\dot{x}}{a^2} - \frac{x^3\dot{x}}{a^3} - \&c. \right) \text{ of which the fluent}$$

$$\text{is } \frac{b}{a} \times \left(x - \frac{x^2}{2a} - \frac{x^3}{3a^2} - \frac{x^4}{4a^3} - \frac{x^5}{5a^4} - \&c. \right) \text{ Ans.}$$

And for the descending series.

$$x+a) \quad b \left(\frac{b}{x} - \frac{ab}{x^2} - \frac{a^2b}{x^3} - \frac{a^3b}{x^4} - \frac{a^4b}{x^5} - \frac{a^5b}{x^6} - \&c. \right) \text{ Hence}$$

$$\frac{bx\dot{x}}{a+x} = \frac{bx}{x} - \frac{abx}{x^2} - \frac{a^2bx}{x^3} - \frac{a^3bx}{x^4} - \&c. \text{ of which the fluent is}$$

$$b \times \text{hyp. log. of } x, +abx^{-1} + \frac{1}{2}a^2bx^{-2} + \frac{1}{3}a^3bx^{-3} + \&c. \text{ Ans.}$$

(Key to Vol. II. page 312.)

Ex. 3. $a+x) -3 \left(-\frac{3}{a} + \frac{3}{a^2}x - \frac{3}{a^3}x^2 + \frac{3}{a^4}x^3 - \frac{3}{a^5}x^4 + \&c. \right)$

Therefore $-\frac{3}{a}\dot{x} + \frac{3}{a^2}x\dot{x} - \frac{3}{a^3}x^2\dot{x} + \frac{3}{a^4}x^3\dot{x} - \&c.$

And consequently $-\frac{3}{a}x + \frac{3}{2a^2}x^2 - \frac{3}{3a^3}x^3 + \frac{3}{4a^4}x^4 - \&c.$ Ans.

Ex. 4. $1+x-x^2) 1-x^2+2x^4(1-x+x^2-2x^3+5x^4-7x^5+\&c.$

Wherefore $\dot{x}-x\dot{x}+x^2\dot{x}-2x^3\dot{x}+5x^4\dot{x}-7x^5\dot{x}+\&c.$

And $x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{2} + x^5 - \frac{7x^6}{6} + \&c.$ Ans.

Ex. 5. $a^2+x^2)b \left(\frac{b}{a^2} - \frac{b}{a^4}x^2 + \frac{b}{a^6}x^4 - \frac{b}{a^8}x^6 + \frac{b}{a^{10}}x^8 - \frac{b}{a^{12}}x^{10} + \&c. \right)$

Whence $\frac{bx}{a^2} - \frac{bx^2\dot{x}}{a^4} + \frac{bx^4\dot{x}}{a^6} - \frac{bx^6\dot{x}}{a^8} + \frac{bx^8\dot{x}}{a^{10}} - \frac{bx^{10}\dot{x}}{a^{12}} + \&c.$

That is,

$$z = \frac{bx}{a^2} - \frac{bx^3}{3a^4} + \frac{bx^5}{5a^6} - \frac{bx^7}{7a^8} + \frac{bx^9}{9a^{10}} - \frac{bx^{11}}{11a^{12}} + \frac{bx^{13}}{13a^{14}} - \&c. \text{ Ans.}$$

Ex. 6. $a+x) a^2+x^2 (a-x + \frac{2x^2}{a} - \frac{2x^3}{a^2} + \frac{2x^4}{a^3} - \frac{2x^5}{a^4} + \frac{2x^6}{a^5} - \&c.$

And $a\dot{x} - x\dot{x} + \frac{2x^2\dot{x}}{a} - \frac{2x^3\dot{x}}{a^2} + \frac{2x^4\dot{x}}{a^3} - \frac{2x^5\dot{x}}{a^4} + \&c.$ whence

$$z = ax - \frac{1}{2}x^2 + \frac{2x^3}{3a} - \frac{2x^4}{4a^2} + \frac{2x^5}{5a^3} - \frac{2x^6}{6a^4} + \frac{2x^7}{7a^5} - \&c. \text{ Ans.}$$

Ex. 7. $z = \frac{3}{2}(a+x)^2.$ Ans.

Ex. 8. $\sqrt{a^2+x^2} = a + \frac{x^2}{2a} - \frac{x^4}{8a^3} + \frac{x^6}{16a^5} - \&c.$ therefore

$$2a\dot{x} + \frac{x^2\dot{x}}{a} - \frac{x^4\dot{x}}{4a^3} + \frac{x^6\dot{x}}{8a^5} - \&c. = \dot{z}, \text{ and}$$

$$2ax + \frac{3x^3}{9a} - \frac{x^5}{20a^3} + \frac{x^7}{54a^5} - \&c. = z. \text{ Ans.}$$

Ex. 9. $\sqrt{a^2-x^2} = a - \frac{x^2}{2a} - \frac{x^4}{8a^3} - \frac{x^6}{16a^5} - \frac{5x^8}{128a^7} - \&c.$ therefore

(Key to Vol. II. page 312.)

$$4ax - \frac{2x^2x}{a} - \frac{x^4x}{2a^3} - \&c. = z; \text{ and,}$$

$$4ax - \frac{2x^3}{3a} - \frac{x^5}{10a^3} - \&c. \&c. = z. \text{ Ans.}$$

Ex. 10. $\frac{5a}{\sqrt{x^2 - a^2}} = \frac{5a}{x} - \frac{10x}{a} - \frac{40x^3}{a^3} - \frac{80x^5}{a^5} - \&c.$ whence

$$5ax^{-1}x - \frac{10xx}{a} - \frac{40x^3x}{a^3} - \frac{80x^5x}{a^5} - \&c. = z. \text{ And}$$

$$5a \times \text{hyp. log. of } x - \frac{5x^2}{a} - \frac{10x^4}{a^3} - \frac{80x^6}{6a^5} - \&c. = z. \text{ Ans.}$$

Ex. 11. $\sqrt[3]{a^3 - x^3} = a - \frac{x^3}{3a^2} - \frac{x^6}{9a^5} - \frac{5x^9}{81a^8} - \&c.$ wherefore

$$2ax - \frac{2x^3x}{3a^2} - \frac{2x^6x}{9a^5} - \&c. = z; \text{ consequently}$$

$$2ax - \frac{x^4}{6a^2} - \frac{2x^7}{63a^5} - \&c. = z. \text{ Ans.}$$

Ex. 12. $\frac{3ax}{\sqrt{ax - xx}} = \frac{3x}{1} \times \frac{a}{\sqrt{ax - xx}}.$ Likewise

$$\frac{a}{\sqrt{ax - xx}} = \frac{\sqrt{a}}{\sqrt{x}} - \frac{2a^{\frac{3}{2}}}{x^{\frac{3}{2}}} - \frac{8a^{\frac{5}{2}}}{x^{\frac{5}{2}}} - \frac{16a^{\frac{7}{2}}}{x^{\frac{7}{2}}} - \&c.$$

Hence, \times lying by $3x$, it will be $3\sqrt{ax}^{-\frac{1}{2}}x - 6a^{\frac{3}{2}}x^{-\frac{3}{2}}x - 24a^{\frac{5}{2}}x^{-\frac{5}{2}}x - 48a^{\frac{7}{2}}x^{-\frac{7}{2}}x - \&c. = z.$

Therefore $6\sqrt{ax}^{\frac{1}{2}} + 12a^{\frac{3}{2}}x^{-\frac{1}{2}} + 16a^{\frac{5}{2}}x^{-\frac{3}{2}} + 19\frac{1}{3}a^{\frac{7}{2}}x^{-\frac{5}{2}} + \&c. = z. \text{ Ans.}$

Ex. 13. $\sqrt[3]{x^3 + x^4 + x^5} = x + \frac{x^2}{3} + \frac{2x^3}{9} + \&c.$ therefore

$$2xx + \frac{2x^2x}{3} + \frac{4x^3x}{9} + \&c. = z; \text{ and}$$

$$x^2 + \frac{2x^3}{9} + \frac{x^4}{9} + \&c. = z. \text{ Ans.}$$

(Key to Vol. II. page 312.)

Ex. 14. $\sqrt{ax-x^2} = a^{\frac{1}{2}}x^{\frac{1}{2}} - \frac{x^{\frac{3}{2}}}{2a^{\frac{1}{2}}} - \frac{x^{\frac{5}{2}}}{8a^{\frac{3}{2}}} - \frac{x^{\frac{7}{2}}}{16a^{\frac{5}{2}}} - \&c.$ wherefore

$$5a^{\frac{1}{2}}x^{\frac{1}{2}}\dot{x} - \frac{5x^{\frac{3}{2}}\dot{x}}{2\sqrt{a}} - \frac{5x^{\frac{5}{2}}\dot{x}}{8\sqrt{a^3}} - \frac{5x^{\frac{7}{2}}\dot{x}}{16\sqrt{a^5}} - \&c. = \dot{z};$$

and consequently $\frac{10a^{\frac{1}{2}}x^{\frac{3}{2}}}{3} - \frac{x^{\frac{5}{2}}}{\sqrt{a}} - \frac{5x^{\frac{7}{2}}}{28\sqrt{a^3}} - \&c.$ Ans.

(Page 314.)

The Correction of Fluents.

Ex. 2. Because $\dot{z} = 5x\dot{x}$, it follows that $z = \frac{5}{2}x^2$. Putting, therefore, $z=0$, and $a=x$, it is $0 = \frac{5}{2}a^2$, and by subtraction $z-0 = \frac{5x^2-5a^2}{2}$, that is, $z = \frac{5}{2}(x^2 - a^2)$. Ans.

Otherwise,

z being unconditionally equal to $\frac{5}{2}x^2$, it will be conditionally $z = \frac{5}{2}x^2 + c$, and by substituting 0 and a agreeably to the conditions, $c = -\frac{5}{2}a^2$; writing, therefore, this value of c for c , it is $z = \frac{5}{2}x^2 - \frac{5}{2}a^2$, as before.

Ex. 3. Since $\dot{z} = 3x\sqrt{a+x}$, manifestly $z = 2(a+x)^{\frac{3}{2}}$. Wherefore substituting 0 both for x and z , there arises $0 = 2(a+0)^{\frac{3}{2}} = 2\sqrt{a^3}$. Consequently the correct fluents are $z = 2(a+x)^{\frac{3}{2}} - 2\sqrt{a^3}$. Ans.

Or,

Putting c for the correction, $z = 2(a+x)^{\frac{3}{2}} + c$; that is, (writing 0 both for z and x), $0 = 2(a+0)^{\frac{3}{2}} + c$; hence $c = -2\sqrt{a^3}$, and the correct fluents are as before $z = 2(a+x)^{\frac{3}{2}} - 2\sqrt{a^3}$. Ans.

Ex. 4. Here $z = 2a \times \text{hyp. log. of } (a+x)$, and by substituting 0 both for x and z , agreeably to the conditions given, it is $z - 0 = 2a \times \text{hyp. log. of } (a+0)$, wherefore the correct fluents are $z = 2a \times (\text{hyp. log. of } (a+x) - \text{hyp. log. of } a) = 2a \times \text{hyp. log. } \frac{(a+x)}{a}$. Ans.

(Key to Vol. II. page 314.)

Ex. 5. By the 11th* Form $z = \frac{2}{a} \times \text{arc to tang. } \frac{x^{\frac{1}{2}}}{a}$, or $\frac{1}{a} \times \text{arc to cosine } \frac{a^2 - x^2}{a^2 + x^2}$. If, therefore, 0 be substituted both for x and z , there will arise in the *first* instance, $0 = \frac{2}{a} \times \text{arc to tang. of } 0^{\frac{1}{2}} a^{-1}$, that is $\frac{2}{a} \times 0$; and in the *second* instance, $0 = \frac{1}{a} \times \text{arc to cos. } \frac{a^2 - 0}{a^2 + 0} = \frac{1}{a} \times \text{arc to cos. 1}$. Consequently the correct fluents according to the Table are

$$z = \left. \begin{array}{l} \frac{2}{a} \times \text{arc to tang. } x^{\frac{1}{2}} a^{-1} \\ \frac{1}{a} \times (\text{arc to cos. } \frac{a^2 - x^2}{a^2 + x^2} - \text{arc to cos. 1.}) \end{array} \right\} \text{Ans.}$$

Or by Infinite Series.

$$\dot{z} = \frac{2\dot{x}}{a^2} \times (1 - \frac{x^2}{a^2} + \frac{x^4}{a^4} - \frac{x^6}{a^6} + \frac{x^8}{a^8} - \frac{x^{10}}{a^{10}} + \&c.) \text{ and}$$

$$z = \frac{2x}{a^2} - \frac{2x^3}{3a^4} + \frac{2x^5}{5a^6} - \frac{2x^7}{7a^8} + \frac{2x^9}{9a^{10}} - \frac{2x^{11}}{11a^{12}} + \&c.$$

But substituting 0 both for x and z , it is $0 = \frac{0}{a^2} - \frac{0}{3a^4} + \frac{0}{5a^6} - \&c.$ signifying that $z = x - 0$, or that the fluents require no correction.

* We have taken the 11th form throughout as it is printed in Hutton; suggesting that the reader will easily accommodate our answers to any change in that form. Thus, when $\sqrt{xa^{-1}}$ is changed to $\sqrt{x^na^{-1}}$, as it ought to be, will $x^{\frac{1}{2}}$ in the 1st and 7th lines of this page become x , and examples 6, 18, and 19, in the comparison of fluents be accordingly adjusted in the tangent value of each fluent. It is the business of the KEY to shew how the fluents of proposed fluxions are found by the Table, not to construct or alter the Forms.

(Key to Vol. II, page 318.)

MAXIMA ET MINIMA.

Ex. 1. Put x for one of the parts of the given line or quantity, so shall $a - x$ be the other part. Now the rectangle under x and $a - x$ is $ax - x^2$, of which the fluxion is $ax - 2x\dot{x}$, by the question a maximum, and consequently $a - 2x = 0$. Hence $x = \frac{1}{2}a$. Ans.

Ex. 2. If x and y represent the two parts, it is evident that $\dot{x} = -\dot{y}$, and *vice versa*, $\dot{y} = -\dot{x}$, according as x increases or decreases, the fluxion of both parts being simultaneous and equal. But by the question $x^m y^n$ is a maximum; therefore $mx^{m-1}y^n = ny^{n-1}x^m$; and, As $x^m : y^n :: mx^{m-1} : ny^{n-1} :: \frac{mx^m}{x} : \frac{ny^n}{y} :: \frac{m}{x} : \frac{n}{y}$, consequently $my = nx$, and, As $x : y :: m : n$. Ans.*

* Signifying that either part must be to the other as the power of the former to the power of the latter. Whereby it is evident that $x = am \div (m+n)$ and $y = an \div (m+n)$.

Ex. 3. Let $x, y,$ and z be assumed for the three parts, then xyz by the question is a maximum: and because xyz is a maximum, xy is a maximum; and for the same reason xz is a maximum. But $\dot{x} = \dot{y}$, therefore, $x = y$; and $\dot{x} = \dot{z}$, consequently, $x = z$. Hence $x = y = z$. Ans.

Ex. 4. If x, y, z be assumed, xy^2z^3 is, by the question a maximum; And taking xy^2 a maximum $xy^2 - 2xy\dot{y} = 0$, hence $y = 2x$: Again, taking xz^3 a maximum, $xz^3 - 3xz^2\dot{z} = 0$, consequently $z = 3x$. Wherefore $x = \frac{1}{6}a, y = \frac{1}{3}a,$ and $z = \frac{1}{2}a$. Ans.

Corollary. Into whatever number of parts a be divided, the divisions will be in the direct ratio of the powers.

Ex. 5. Put x for the value of the fraction sought, then by the question $x^m - x^n$ is a maximum, and for that reason $mx^{m-1} = nx^{n-1}$. Now because m must be to n in the ratio of 1 to some number whether integral or fractional, if m be assumed as unity, whilst n represents the number to which 1 has the same ratio as m to n , it will be $1 = nx^{n-1}$, and by transposition, &c. $x = \sqrt[n-1]{\frac{1}{n}}$; wherefore $\sqrt[n-1]{\frac{m}{mn}}$ is the fraction required.

Ex. 6. Because the fluxions of the two parts must be at all times equal, the one negative and the other positive, if x and y be assumed, it will be $\dot{x} = -\dot{y}$, and $-\dot{x} = \dot{y}$. But by the question $2x^2 + xy + 3y^2$ is a minimum, therefore (taking the fluxion and rejecting the fluxional letters,) $4x + y - x - 6y$, that is, $3x - 5y$, is a minimum. Now $x + y$ being equal

(Key to Vol. II. page 318.)


to 80, there may be substituted $80-x$ for y , or $80-y$ for x . Consequently $3x=400-5y$; or $240-3y=5y$. Hence $x=50$, and $y=30$. Ans.

Ex. 7. Let ABC be the triangle, and draw DE parallel to BC , and EF parallel to AB . Also put a for AB the perpendicular, b for BC the base, and x for BD the altitude of the rectangle sought. Then $AD=a-x$.

But, As $a : b :: (a-x) : \frac{(ba-bx)}{a} =$

DE ; therefore $\frac{bax-bx^2}{a} = \frac{b}{a} \times (ax-$


$x^2) =$ the area of the rectangle required. Now the fluxion of $ax-x^2 = ax-2xx=0$, by the question; hence $x=\frac{1}{2}a$, or the rectangle will have half the altitude of the triangle. Q. E. I.




Ex. 8. In the quadrant ABC , draw FD parallel to CB ; DE parallel to AC , and join CD . Also put r for the radius CD , and x for CE . $ED = \sqrt{r^2-x^2}$, and $CE \times ED =$ area of the rectangle $= x\sqrt{r^2-x^2}$; of

which the fluxion is $\frac{r^2x-2x^2x}{\sqrt{r^2-x^2}}$. Whence

$r^2 = 2x^2$, or $x = r\sqrt{\frac{1}{2}} = r \times .70710678$ &c. Ans.




Ex. 9. Ans. When the sides about the right angle are equal, and the hypotenuse a tangent to the circle at the middle point of the arc. See the annexed diagram, where ABC represents the quadrant, F the middle point of the arc, DE a tangent to the circle at F , and DBE the triangle required.



(Key to Vol. II. page 318.)

Ex. 10. Ans. *For the triangle.* When the isocles sides are tangents to the curve, and the altitude of the triangle double the length of the subtangent.

For the rectangle. When the altitude is half the altitude of the circumscribing triangle; which is evident by Ex. 7, in maxima and minima. See the figure in the margin, where ACB is the semi-ellipse; CD half the transverse diameter; ET and FT tangents to the curve at the points t, t ; ST the subtangent equal to half of TD ; AB the conjugate diameter produced both ways to meet the tangents TF and TE in the points F and E ; ETF the required triangle; and $tppt$ the inscribed rectangle.



(Page 319.)

Ex. 11. Ans. The triangle and rectangle will be precisely as in the semi-ellipse in Ex. 10.

Ex. 12. Ans. The same limitations as given in the Answer to the last two Questions.

Ex. 13.


Put a for BC , the altitude of the cone;

b for AD , the diameter of the base;

x for $fg = dh$, the diameter of the cylinder;

and m for $\frac{3.1416}{4} = .7854$ the

area of a circle whose diameter is 1.



Then,

As $1^2 : x^2 :: m : mx^2 =$ the area of the circle $fsgr$.


Again, because the triangle Adf is similar to the triangle ACB ,

It is, As $\frac{1}{2}b : a :: (\frac{1}{2}b - \frac{1}{2}x) : df = \frac{ab - ax}{b}$; which multiplied by

(Key to Vol. II. page 319.)


the area of the cylindrical base, $mx^2 = \frac{ambx^2 - amx^3}{b} =$
 the solidity of the cylinder and a maximum. Therefore
 $\frac{2abmx\dot{x} - 3amx^2\dot{x}}{b} = 0$, or $x = \frac{2}{3}b$; and $df = \frac{1}{3}a$. Hence the
 diameter of the greatest cylinder that can be inscribed in any
 cone is $\frac{2}{3}$ of the diameter of the base of the cone, and the alti-
 tude of the cylinder $\frac{1}{3}$ the altitude of the cone. *Which was re-
 quired.*

Ex. 14. By Example 1. in max-
 ima it is evident that the base of
 the cistern must be a square,
 therefore put x for the side of the
 base, op ; and y for the altitude
 of the cistern, om . Then $4x =$ the
 periphery of the base, $x^2 =$ area of



the base, and $4xy =$ internal superficies of the sides of the cistern.
 Also the solidity is x^2y , which put $= a$ the quantity of water given.
 Now since $x^2y = a$, and $4xy$ the internal superficies of the sides, it
 follows that $\frac{4}{x} \times \frac{x^2y}{1} = \frac{a}{1} = 4xy = \frac{4a}{x} =$ the internal superficies
 of the sides. Therefore $x^2 + \frac{4a}{x} =$ the whole part to be covered
 with lead and a minimum. But the fluxion of $x^2 + 4ax^{-1}$ is $2x\dot{x}$
 $- 4ax^{-2}\dot{x} = 0$. Hence $2x^3 = 4a$, or $x^3 = 2a$, and $x = \sqrt[3]{2a}$. Again,
 since $x^2y = a$, and $x^3 = 2a$, it follows that, $2x^2y = x^3$, or $2y = x$;
 consequently the internal side of the base must be double the
 altitude. Q. E. I.

Ex. 15. For one quart ale measure in
 cubic inches put a , and let x represent the
 diameter of the tankard, cd , or ab ; y the
 depth, ac , or bd . Moreover put $m = 3.1416$
 the circumference of a circle of which the




diameter is unity. Then, As $1 : m :: x : xm$, the circumference
 of the base: consequently xym is the concave superficies of the
 cylinder. Again the area of the base $= \frac{mx^2}{4}$ which drawn into
 the altitude y , gives $\frac{mx^2y}{4} =$ the solidity $= a$. Hence, (as in the
 last example.) $\frac{4a}{x} + \frac{mx^2}{4} =$ the quantity of silver a minimum.


(Key to Vol. II. page 319.)

But the fluxion of $4ax^{-1} + \frac{1}{4}mx^2$ is $\frac{mxx}{2} - \frac{4ax}{x^2} = 0$. Therefore $mx^3 = 8a$, and $x = 2\sqrt[3]{\frac{a}{m}}$. Now $\frac{mx^2y}{4}$ being equal to a , and mx^3 equal to $8a$, it follows that, $2mx^2y = mx^3$; or, dividing by mx^2 , $2y = x$; that is, half the diameter of the base is equal the altitude. But $a = 70\frac{1}{2}$ cubic inches; wherefore $y = 2.8205$ inches, and $x = 5.641$ inches. Q. E. I.

Ex. 16. Let the axis of the parabola be ns parallel to AC , and let rm be the ordinate or base. Also put a for AB ; b for AC ; and x for BN . Then, As $a : b :: x : a^{-1}bx = ns$; likewise $rn \times rn = BN \times NA = nm^2 = \overline{a-x} \times x = ax - x^2$. Therefore $rm = 2\sqrt{ax - x^2}$. But every parabola is $\frac{2}{3}$ of a parallelogram of the same base and altitude, consequently $\frac{2}{3} \times rm \times ns = \text{area of the parabola} = \frac{2}{3} \times \frac{bx}{a} \times 2\sqrt{ax - x^2} = \frac{4bx}{3a} \sqrt{ax - x^2}$; of which the fluxion is $3ax^2\dot{x} - 4x^3\dot{x} = 0$. Hence $x = \frac{3}{4}a$, $rm = AB \times \sqrt{\frac{3}{4}}$, and $ns = \frac{3}{4}$ of AC . Ans.



Ex. 17. Let ML represent the less, and BE the greater axis of the ellipse; and let ev be parallel to AC the axis of the cone, meeting BD the diameter of the base in v . Also let the diameters EF and pn be parallel to BD , and such that pn may pass through o the center of the ellipse considered as variable by the motion of E on the line AD . If a be now put for the axis of the cone, b for the radius of the cone's base, and x for cv , then will $b+x = BC + cv = Bv$. And, As $b : a :: b-x : \frac{ab-ax}{b} = ev$. Where-



fore (Euc. B. I. Prop. xlvii.) $BE = \sqrt{a^2 \times b - x^2 + b^2 \times b + x^2}$. Moreover because of the similar triangles BEF and BOP , EBD and EOH , and that $OB = OE = \frac{1}{2}BE$, $on = b$; likewise $op = x$. But

(Key to Vol. II. page 319.)

$$op \times on = ol^2 = bx; \text{ that is } ML = 2\sqrt{bx}; \text{ therefore } BE \times ML = \frac{\sqrt{4bx \times a^2 \times b-x^2 + b^2 \times b+x^2}}{b}$$

Now the area of an ellipse being in the constant ratio of the rectangle of its two diameters [Cor. 2. Theor. iii. of the Ellipse, Conic Sections.], that is, in the ratio of .785398163397448309615660845819875721049292349843776455243736148076954101 &c. to 1; and the area being at the last given expression a *maximum*, the expression itself is a *maximum*: for which reason its fluxion $b^4\dot{x} + 4b^3x\dot{x} + 3b^2x^2\dot{x} + a^2b^2\dot{x} - 4a^2bx\dot{x} + 3a^2x^2\dot{x} = 0$. Consequently x^2

$$\frac{-4bx \times (a^2 - b^2)}{3a^2 + 3b^2} = \frac{-b^2}{3}, \text{ or } x = \frac{2b \times a^2 - b^2 \pm b\sqrt{a^4 - 14a^2b^2 + b^4}}{3a^2 + 3b^2},$$

by which the ellipse is determined.

If, however, $a^4 - 14a^2b^2 + b^4$ be, as it sometimes may, negative, the solution (since the root of a negative quantity is impossible) in that case fails. To determine the limit, suppose that $a^4 - 14a^2b^2 + b^4 = 0$, or, which is the same, put $a^4 + b^4 = 14a^2b^2$; then $a^2 = b^2 \times \sqrt{48 + 7}$, and $a = b \times 2 + \sqrt{3}$; consequently, As $a : b :: (2 + \sqrt{3}) : 1$. Hence if the ratio of the altitude of the cone to half the diameter of the base, be not greater than that of $2 + \sqrt{3}$ to 1, that is, if the angle ADB be not greater than 75° , the fluxion of the ellipse cannot be equal to 0, for the area would constantly increase as E approaches to D , and be greatest when E is at D .

Again, if the ellipse do not *increase continually* from the vertex to the base of the cone, as in the case when the angle ADC is not greater than 75° , the elliptical area will increase till it arrives at a certain point determinable, afterwards decrease, and again increase till it coincides with the base of the cone. [See Simpson's Fluxions, where it is proved that, unless the angle which the slant side makes with the axis of the cone be less than $11^\circ \cdot 57'$, the greatest ellipse will be less than the base of the cone.]

Ex. 18. The fluxion of the hyp. log. of x^x (See Art. 68, page 327, and Ex. 57, page 300, vol. ii.) is $x + \dot{x}$ (hyp. log. of x .) which by the question is a maximum, and therefore equal to 0; hence the hyp. log. of $x = -1$. Now the number answering to any hyperbolic logarithm z ,* (see page 328, vol. ii.) is $1 + z + \frac{z^2}{2} +$

* Let z be denoted by $\frac{1+n}{1}$, that is by $1+n$.

(Key to Vol. II. page 319.)

$\frac{z^3}{2 \cdot 3} + \frac{z^4}{2 \cdot 3 \cdot 4} + \&c.$; consequently writing -1 for z , there arises this equation, $x = 1 - 1 + \frac{1}{2} - \frac{1}{2 \cdot 3} + \frac{1}{2 \cdot 3 \cdot 4} - \frac{1}{2 \cdot 3 \cdot 4 \cdot 5} + \&c.$ which series summed $= 0.367878 \&c.$ Ans.

THE METHOD OF TANGENTS.

(Page 320.)

Ex. 2. $ax - x^2$ being equal to y^2 , their fluxions $a\dot{x} - 2x\dot{x}$ and $2y\dot{y}$ are equal, therefore $\frac{\dot{x}}{\dot{y}} = \frac{2y}{a - 2x} = \frac{y}{\frac{1}{2}a - x}$, or $\frac{y\dot{x}}{\dot{y}} = \frac{y^2}{\frac{1}{2}a - x} =$ the subtangent. Ans.

Ex. 3. ax being equal to y^2 , their fluxions $a\dot{x}$ and $2y\dot{y}$ are equal, hence $\frac{\dot{x}}{\dot{y}} = \frac{2y}{a}$, and $\frac{y\dot{x}}{\dot{y}} = \frac{2y^2}{a} = \frac{2ax}{a} = 2x$; that is, the subtangent is double its corresponding absciss. Q. E. 1.

Ex. 4. The equation being $c^2(ax - x^2) = a^2y^2$, the fluxion of the one side is equal to the fluxion of the other; wherefore $c^2 \times \overline{a\dot{x} - 2x\dot{x}} = 2a^2y\dot{y}$, or $\frac{\dot{x}}{\dot{y}} = \frac{2a^2y}{c^2(a - 2x)}$. Consequently $\frac{y\dot{x}}{\dot{y}} = \frac{2a^2y^2}{c^2(a - 2x)} = \frac{a^2y^2}{c^2(\frac{1}{2}a - x)} = \frac{c^2(ax - x^2)}{c^2(\frac{1}{2}a - x)} = \frac{ax - x^2}{\frac{1}{2}a - x}$, which was required.

Ex. 5. The equation of the curve being $c^2(ax + x^2) = a^2y^2$, it follows that, $c^2(a\dot{x} + 2x\dot{x}) = 2a^2y\dot{y}$; hence $\frac{y\dot{x}}{\dot{y}} = \frac{a^2y^2}{c^2(\frac{1}{2}a + x)} = \frac{ax + x^2}{\frac{1}{2}a + x} =$ the subtangent. Wherefore the point through which the tangent must pass being determined, the tangent may be drawn. Q. E. F.

(Key to Vol. II. page 320.)

Ex. 6. Because $xy=a^2$, it is manifest that $x = \frac{a^2}{y} = a^2y^{-1}$. Now taking the fluxion of this equation, $\dot{x} = \frac{-a^2\dot{y}}{y^2}$ and $\frac{\dot{x}}{\dot{y}} = \frac{-a^2}{y^2}$; consequently $\frac{y\dot{x}}{\dot{y}} = \frac{-a^2}{y}$ = the value of the sub-tangent, which being a negative quantity, indicates that the tangent and vertex will be on different sides of the ordinate, but the sub-tangent equal to the absciss. Q. E. I.

THE LENGTH OF CURVE LINES.

(Page 323.)

Ex. 2. $x = \frac{y^n}{a^{n-1}}$ being a general expression for the parabola, $\dot{x} = \frac{ny^{n-1}\dot{y}}{a^{n-1}}$; and because $\dot{z} = \sqrt{\dot{y}^2 + \dot{x}^2}$, it is evident that, $\dot{z} = \sqrt{\dot{y}^2 + \frac{n^2y^{2n-2}\dot{y}^2}{a^{2n-2}}} = \dot{y}\sqrt{1 + \frac{n^2y^{2n-2}}{a^{2n-2}}}$, of which the fluent in an infinite series is $y + \frac{n^2y^{2n-1}}{2n-1 \times 2a^{2n-2}} - \frac{n^4y^{4n-3}}{4n-3 \times 8a^{4n-4}} + \frac{n^6y^{6n-5}}{6n-5 \times 16a^{6n-6}} - \&c. = z$. Ans.

Ex. 3. Because $ax^2=y^3$, it follows that $x = \frac{\sqrt{y^3}}{\sqrt{a}}$, and $\dot{x} = \frac{3y^{\frac{1}{2}}\dot{y}}{2\sqrt{a}}$. Wherefore $\dot{z} (= \sqrt{\dot{y}^2 + \dot{x}^2}$ as in the last example) = $\sqrt{\dot{y}^2 + \frac{9y\dot{y}^2}{4a}} = \frac{\dot{y}\sqrt{4a+9y}}{2\sqrt{a}}$. Now the fluent of this fluxion is $\frac{\sqrt{4a+9y}^{\frac{3}{2}}}{27a^{\frac{1}{2}}}$ which corrected by taking $y=0$, becomes $\frac{\sqrt{4a+9y}^{\frac{3}{2}}}{27a^{\frac{1}{2}}} - \frac{8a}{27} = z$. Ans.

(Key to Vol. II. page 323.)

Ex. 4. Let the semi-transverse axis of the ellipse be represented by a , and the semi-conjugate by b , then the equation of the

curve is $\frac{a^2 y^2}{b^2} = 2ax - x^2$. Hence $x = a \pm \sqrt{a^2 - \frac{a^2 y^2}{b^2}} = a \pm$

$\frac{a\sqrt{b^2 - y^2}}{b}$, and $\dot{x} = \frac{-ay\dot{y}}{b\sqrt{b^2 - y^2}}$. Again, $z (= \sqrt{x^2 + y^2}) =$

$\sqrt{y^2 - \frac{a^2 y^2 \dot{y}^2}{b^2(b^2 - y^2)}} = \dot{y} \sqrt{1 - \frac{a^2 y^2}{b^4 - b^2 y^2}}$. Now converting

$\frac{a^2 y^2}{b^4 - b^2 y^2}$ to an infinite series, the last fluxion becomes $\dot{y} \times$

$\sqrt{1 - \frac{a^2 y^2}{b^4} + \frac{a^2 y^4}{b^6} - \frac{a^2 y^6}{b^8} + \&c.} = \dot{y} - \frac{a^2}{2b^4} y^2 \dot{y} + (\frac{a^2}{2b^6} +$

$\frac{a^4}{8b^8}) y^4 \dot{y} - (\frac{a^2}{2b^8} + \frac{a^4}{4b^{10}} + \frac{a^6}{16b^{12}}) y^6 \dot{y} + \&c.$ Wherefore $z = y$

$- \frac{a^2 y^3}{6b^4} + (\frac{a^2}{b^2} + \frac{a^4}{4b^4}) \times \frac{y^5}{10b^4} - (\frac{a^2}{b^2} + \frac{a^4}{2b^4} + \frac{a^6}{8b^6}) \times$

$\frac{y^7}{14b^6} + \&c.$ Ans.

Ex. 5. If a and b respectively be assumed, as in the last Example, for the semi-transverse and semi-conjugate axes of the hyperbola, the equation of the curve will be $\frac{a^2 y^2}{b^2} = 2ax$

$+ x^2$; hence $x = \frac{a\sqrt{b^2 + y^2}}{b} - a$, and $\dot{x} = \frac{ay\dot{y}}{b\sqrt{b^2 + y^2}}$. Pro-

ceeding, therefore, in all respects as in the last Example,

$z = y + \frac{a^2 y^3}{6b^4} - \frac{y^5}{10b^4} (\frac{a^2}{b^2} + \frac{a^4}{4b^4}) + \frac{y^7}{14b^6} (\frac{a^2}{b^2} + \frac{a^4}{2b^4} + \frac{a^6}{8b^6})$

$- \&c.$ Ans.

QUADRATURES.

(Page 325.)

Ex. 3. Extracting the $(m+n)$ th root on both sides, the general equation becomes $y = a^{\frac{m}{m+n}} z^{\frac{n}{m+n}}$; and putting w for yz there

(Key to Vol. II. page 325.)

arises $w = a^{\frac{m}{m+n}} z^{\frac{n}{m+n}} z$. Wherefore w (the fluent or area) is

$$a^{\frac{m}{m+n}} \times \frac{z^{\frac{n}{m+n}+1}}{\frac{n}{m+n}+1} = \frac{a^{\frac{m}{m+n}} z^{\frac{n}{m+n}} (m+n)z}{m+2n} = \frac{m+n}{m+2n} \text{ times the}$$

rectangle under the absciss and ordinate. Ans.

COR. 1. When $m=n$, the figure is the common parabola, and the area $= \frac{2}{3}zy$. ($= \frac{2}{3}xy$.)

COR. 2. When $m=0$, the figure is a right angled isocetes triangle, and its area $= \frac{1}{2}zy$. ($= \frac{1}{2}xy$.)

Ex. 4. Let a represent the transverse axis, and c the conjugate,

so shall $y = \frac{c}{a} \sqrt{ax-x^2}$, and $\dot{w} = \frac{c}{a} \times \dot{x} (ax-x^2)^{\frac{1}{2}}$. Now the

fluxion of w is the fluxion of the area bounded by the curve, absciss, and ordinate, to any portion of the elliptic curve having the transverse diameter a for its axis, x for its absciss, and y for its ordinate. But $\dot{x} (ax-x^2)^{\frac{1}{2}}$ [see Ex. 2. page 324, vol. ii.] is the fluxion of the corresponding area of a circle of which the diameter is a . Again, the fluent of


$$\dot{x} (ax-x^2)^{\frac{1}{2}} \text{ is } x\sqrt{ax} \times \left(\frac{2}{3} - \frac{x}{5a} - \frac{x^2}{28a^2} - \frac{3x^3}{216a^3} - \frac{15x^4}{2112a^4} - \right.$$

$$\left. \&c.) \text{ And the fluent of } w \text{ is } \frac{c}{a} \times x\sqrt{ax} \times \left(\frac{2}{3} - \frac{x}{5a} - \frac{x^2}{28a^2} \right.$$

$-\&c.)$ Consequently the area of any segment of an ellipse is, to the area of the corresponding segment of the circumscribing circle, as the less axis of the ellipse is to the diameter of that circle; and the wholes in the same ratio.

Q. E. D.

Ex. 5. Let B in the annexed figure be the vertex of an hyperbola, cd a tangent to the curve at the point D, BC the semi-transverse axis, and AB one of the abscisses to the ordinate AD. Also put a for BC, b for the semi-conjugate axis



of the hyperbola, x for the subtangent AC, and y for the ordinate


AD. Then by the property of the curve, $y = \frac{b}{a} \sqrt{x^2 - a^2}$.

(Key to Vol. II. page 325.)

Wherefore \dot{w} , that is, $y\dot{x} = \frac{cx}{a} \sqrt{x^2 - a^2}$ the fluxion of the hyperbolic area, of which the correct fluent or area is $\frac{cx\sqrt{x^2 - a^2}}{2a} - \frac{1}{2}ac \times \text{hyp. log.} \frac{x + \sqrt{x^2 - a^2}}{a}$. Q. E. I.

Ex. 6. Let $AIBIC$ be an hyperbola, DE and EF the asymptotes, B the vertex, and $GBIK$ the required space between the curve and asymptote. Also put a for EK , x for CK , and y for IK . Then by the property of the curve, $y = \frac{a^2}{a+x}$.


Now if a be considered unity, the value of y is $\frac{1}{1+x}$, and $y\dot{x} = \frac{\dot{x}}{1+x}$ = the fluxion



of the area sought. But $\frac{\dot{x}}{1+x} = \dot{x} - x\dot{x} + x^2\dot{x} - x^3\dot{x} + x^4\dot{x} - \&c.$ in an infinite series, and the fluent of this fluxion is $x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \frac{1}{5}x^5 - \&c.$ the area $GBIK$. Q. E. I.

Ex. 7. Let the curve be AOC , PQ the asymptotes, QC , and the area of quadrature $CCQRO$. Then by the given equation, $y = \frac{a^{\frac{m+n}{n}}}{\sqrt[n]{x^m}} = a^{\frac{m+n}{n}} x^{-\frac{m}{n}}$, conse-

quently $y\dot{x} = a^{\frac{m+n}{n}} x^{-\frac{m}{n}} \dot{x}$. Now of this fluxion the fluent is $\frac{a^{\frac{m+n}{n}} x^{-\frac{m}{n}}}{-\frac{m}{n}} = -\frac{na^{\frac{m+n}{n}} x^{-\frac{m}{n}}}{m}$ the correct area when x is 0 and Q



n greater than m . But if n be less than m when x is 0, the area is infinite, because the index of x being negative, x (that is, 0) descends to the denominator, and $na^{\frac{m+n}{n}}$ a finite quantity is divided by 0.

In this case, however, the area $AQRO$ will be finite, and its

(Key to Vol. II. page 325.)

fluxion = $-a^{\frac{m+n}{n}} x^{\frac{-m}{n}}$; of which the fluent (reduced, and its signs changed) is $\frac{na^{\frac{m+n}{n}} x^{\frac{n-m}{n}}}{m-n}$, evidently wanting no correction because vanishing when x is infinite. Q. E. I.

☞ This example may be seen at some length, and treated in a different manner, in VINCE'S excellent treatise on Fluxions, 4th edit. p. 70.


THE SURFACES OF SOLIDS.

(Page 326.)

Ex. 2.

1. For an Oblong Spheroid.

Let FCADG be a semi-spheroid generated by the rotation of the quarter ellipse HFCA about the semi-transverse axis HA; and let CBD be a plane parallel to the base of the semi-spheroid meeting the transverse axis in any point B. If a be put for AH, b for FH, x for BH, y for BC, z for FC, and s for the superficies generated by FC, it will be, by the



property of the ellipse, $y = \frac{b}{a} \sqrt{a^2 - x^2}$. Hence $\dot{y} = \frac{-bx\dot{x}}{a\sqrt{a^2 - x^2}}$,

and $\sqrt{x^2 + y^2} = \sqrt{x^2 + \frac{b^2 x^2 x^2}{a^2(a^2 - x^2)}} = \frac{x\sqrt{a^4 - (a^2 - b^2)x^2}}{a\sqrt{a^2 - x^2}}$. Now

substituting e for $(a^2 - b^2)^{\frac{1}{2}}$ the eccentricity of the ellipse, the last fluxion becomes $\frac{x\sqrt{a^4 - e^2 x^2}}{a\sqrt{a^2 - x^2}} = \frac{ex\sqrt{a^4 e^{-2} - x^2}}{a\sqrt{a^2 - x^2}}$. Consequently

\dot{s} , that is, $2cyz$ or $2cy\sqrt{x^2 + y^2}$, = $\frac{2bcex\sqrt{a^4 e^{-2} - x^2}}{a^2}$, of which the

fluent† expressed in an infinite series is $2bcx \left(1 - \frac{e^2 x^2}{6a^4} - \frac{e^4 x^4}{40a^8} - \right.$

$\left. \frac{e^6 x^6}{112a^{12}} - \&c. \text{ in infinitum.} \right)$

* Where c represents the circumference of the circle of which the diameter is UNITY.

† Which might also be expressed in finite terms.

(Key to Vol. II. page 326.)

2. For an Oblate Spheroid.

Because AH (see the last figure) is in an oblate spheroid less than FH , $\sqrt{a^2 - b^2}$ is an imaginary value, the square root of a negative quantity being impossible. In this case, therefore, put

$e = \sqrt{b^2 - a^2}$, and take $m : a :: a : e$, so shall $m = a^2 e^{-1}$. Hence

$$\dot{s} = \frac{2bcx\sqrt{a^4 e^{-2} + x^4}}{a^2} = \frac{2bcx\sqrt{m^2 + x^2}}{m} = \frac{2bc}{m} \times x\sqrt{m^2 + x^2},$$

of which the fluent is $\frac{bcx}{m} \sqrt{m^2 + x^2} + bcm \times \text{hyp. log.} \frac{x + \sqrt{m^2 + x^2}}{m}$.

Ex. 3. If a denote the parameter of the axis, the equation of the generating curve will be $ax = y^2$; hence $x = \frac{y^2}{a}$, and $\dot{x} = \frac{2y\dot{y}}{a}$.

Consequently $\sqrt{x^2 + y^2} = \sqrt{4a^{-2}y^2\dot{y}^2 + y^2} =$

$$\frac{y\sqrt{4y^2 + a^2}}{a}, \text{ and } 2cy\dot{z} = \frac{2cy\dot{y}}{a} \times (4y^2 + a^2)^{\frac{1}{2}},$$

of which the fluent is $\frac{c\sqrt{4y^2 + a^2}^{\frac{3}{2}}}{6a}$ or, (when corrected by taking $y=0$),

$$\frac{c\sqrt{4y^2 + a^2}^{\frac{3}{2}}}{6a} - \frac{1}{6}a^2c. \text{ Ans.}$$

Ex. 4. Put a for the semi-transverse axis of the generating hyperbola, b for the semi-conjugate, and x for the distance from the center of any ordinate to the curve. Then,

by the property of the hyperbola, $y = \frac{b}{a}\sqrt{x^2 - a^2}$; hence

$$\dot{y} = \frac{bx\dot{x}}{a\sqrt{x^2 - a^2}}, \text{ and } \sqrt{x^2 + y^2} = \frac{x\sqrt{a^2 + b^2} \times (x^2 - a^4)}{a\sqrt{x^2 - a^2}}.$$

Consequently $2cy\dot{z} = \frac{2bcx\sqrt{a^2 + b^2} \times (x^2 - a^4)}{a^2}$; of which

the fluent (assuming $n^2 = \frac{a^4}{a^2 + b^2}$) is $\frac{bcx\sqrt{x^2 - n^2}}{n} - bcn \times$

$\text{hyp. log. of } x + \sqrt{x^2 - n^2}$. Lastly, correcting this fluent by


(Key to Vol. II. page 326.)

putting $x=a$, the true fluent is $(\frac{bcx}{n} \sqrt{x^2-n^2} - b^2c) - bcn$
 \times hyp. log. of $\frac{x + \sqrt{x^2-n^2}}{(a^2+bn)a^{-1}} = (\frac{bcx}{n} \sqrt{x^2 - \frac{n^2}{2x}} - b^2c) - bcn$
 \times hyp. log. of $\frac{a}{a^2+bn} (2x - \frac{n^2}{2x} - \frac{n^4}{8x^3} - \frac{n^6}{16x^5} - \frac{5n^8}{128x^7} - \&c.)$

THE SOLIDITY OF BODIES.

(Page 327.)

Ex. 2. Let AB be the transverse and CD the conjugate axis of the generating ellipse; also let FBG be a plane at right angles to, and cutting the transverse axis in any point H. Then if the greater axis of the solid be represented by a , the less by b , AH by x , and HG by y , it



will be, by the property of the ellipse, As $a^2 : b^2 :: x(a-x) : y^2$, whence $y^2 = \frac{b^2}{a^2}(ax-x^2)$, and $cy^2x = a^{-2}b^2c(axx+x^2x)$, of

which the fluent is $a^{-2}b^2c(\frac{1}{2}ax^2 - \frac{1}{3}x^3)$ the content of the segment BGF. Now when H coincides with B, x is equal to a , and $a^{-2}b^2c(\frac{1}{2}ax^2 - \frac{1}{3}x^3)$ is the fluent of the whole spheroid, which, by substituting a for x , becomes $a^{-2}b^2c(\frac{1}{2}a^3 - \frac{1}{3}a^3) = \frac{1}{2}ab^2c - \frac{1}{3}ab^2c = \frac{1}{6}ab^2c$.

COROLLARY 1. When $AB=CD$, the solid is a sphere, and its content truly expressed by $\frac{1}{6}a^3c$.

COROLLARY 2. A sphere and spheroid, whether oblong or oblate, are each $\frac{2}{3}$ of the solidity of the circumscribing cylinder, for $\frac{1}{6}ab^2c$ being the content of a cylinder of which the diameter of the base is b , and altitude a , it is evident that, As $\frac{1}{6} : \frac{1}{4} :: \frac{2}{3} : 1 ::$ content of a sphere or spheroid : the content of the circumscribing cylinder.

Ex. 3. The equation of the generating curve being $a^{m-n}x^n = y^m$, it follows that $y = a^{\frac{m-n}{m}} x^{\frac{n}{m}}$. Hence $cy^2x = ca^{\frac{2m-2n}{m}} x^{\frac{2n}{m}} x$, of

(Key to Vol. II. page 327.)

$$\text{which the fluent is } ca^{\frac{2m-2n}{m}} \times \frac{x^{\frac{2n}{m}+1}}{\frac{1}{m}(2n+m)} = ca^{\frac{2m-2n}{m}} \times \frac{mx^{\frac{2n+m}{m}}}{2n+m} = ca^{\frac{2m-2n}{m}} \times \frac{mx}{2n+m} = cy^2 \times \frac{mx}{2n+m}. \text{ Ans.}$$

COR. 1. If $m=2$, and $n=1$, the solid is the common paraboloid, and equal to $\frac{1}{2}$ its circumscribing cylinder.

COR. 2. If $m=n$, the curve is changed to a straight line, and the solid generated is a cone, equal to $\frac{1}{3}$ its circumscribing cylinder.

Ex. 4. Because the equation of the curve is $\frac{b^2}{a^2}(ax+x^2)=y^2$, it

follows that, $cy^2\dot{x} = \frac{b^2c}{a^2}(ax\dot{x}+x^2\dot{x})$ of which the fluent is

$$\frac{b^2c}{a^2}\left(\frac{1}{2}ax^2 + \frac{1}{3}x^3\right). \text{ The content required.}$$

OF LOGARITHMS.

(Page 329.)

Ex. 2. The fluxion of $\frac{a+x}{b}$ is $\frac{\dot{x}}{b}$, which divided by $\frac{a+x}{b}$ becomes

$$\begin{aligned} \frac{\dot{x}}{a+x} &= \dot{x} \times \frac{1}{a+x} = \dot{x} \times \left(\frac{1}{a} - \frac{x}{a^2} + \frac{x^2}{a^3} - \frac{x^3}{a^4} + \frac{x^4}{a^5} - \frac{x^5}{a^6} + \&c. \right) \\ &= \frac{\dot{x}}{a} - \frac{x\dot{x}}{a^2} + \frac{x^2\dot{x}}{a^3} - \frac{x^3\dot{x}}{a^4} + \frac{x^4\dot{x}}{a^5} - \&c. \text{ of which the fluent is } \frac{x}{a} \\ &\quad - \frac{x^2}{2a^2} + \frac{x^3}{3a^3} - \frac{x^4}{4a^4} + \frac{x^5}{5a^5} - \frac{x^6}{6a^6} + \&c. \text{ the logarithm reqd.} \end{aligned}$$

Ex. 3. The fluxion of $a-x$ is $-\dot{x}$, which divided by $a-x$ be-

$$\begin{aligned} \text{comes } \frac{-\dot{x}}{a-x} &= \dot{x} \times \frac{-1}{a-x} = \dot{x} \times \left(-\frac{1}{a} - \frac{x}{a^2} - \frac{x^2}{a^3} - \frac{x^3}{a^4} - \frac{x^4}{a^5} - \&c. \right) \\ &= -\frac{\dot{x}}{a} - \frac{x\dot{x}}{a^2} - \frac{x^2\dot{x}}{a^3} - \frac{x^3\dot{x}}{a^4} - \frac{x^4\dot{x}}{a^5} - \&c. \text{ of which the fluent is } - \\ &\quad \frac{x}{a} - \frac{x^2}{2a^2} - \frac{x^3}{3a^3} - \frac{x^4}{4a^4} - \frac{x^5}{5a^5} - \&c. \text{ IN INFINITUM, the logarithm} \\ &\quad \text{required.} \end{aligned}$$

(Key to Vol. II. page 329.)

Ex. 4. Taking $\frac{a+x}{a-x} = 3$, the value of a is $2x$; wherefore $\frac{x}{a} = \frac{1}{2}$, and $\frac{x^2}{a^2} = \frac{1}{4}$ a constant factor in every succeeding term of the series. Likewise $2m = 2$ for hyperbolic logarithms, and $\cdot 8685889638065036553022578$ &c. for common logarithms.

Dividing $2m$, therefore, by 2, the quotients successively by 4, and afterwards collecting the whole of the 1st quotient, $\frac{1}{3}$ of the 2nd quotient, $\frac{1}{3}$ of the 3rd quotient, $\frac{1}{7}$ of the 4th quotient, $\frac{1}{9}$ of the 5th quotient, $\frac{1}{11}$ of the 6th quotient, $\frac{1}{13}$ of the &c. there arise

In the one instance.

1.
 .083333333333
 .0125
 .002232142857
 .000434027777
 .000088778409
 .000018780048
 .000004069010
 .000000897582
 .000000200773
 .000000045413
 .000000010366
 .000000002384
 .000000000551
 .000000000128

1.098612288631 = the hyp. log. of 3, true to 10 decimal places.

In the other instance.

.434294481903
 .036191206825
 .005428681023
 .000969407325
 .000188495868
 .000038555973
 .000008156071
 .000001767148
 .000000389813
 .000000087194
 .000000019722
 .000000004501
 .000000001035
 .000000000239
 .000000000055*

.477121254700 = the com. log. of 3, true to 10 decimal places.

* Here 6 are carried from the preceding column (which is omitted for the sake of uniformity,) making the amount of this column 70.

Ex. 5. Ans. { The com. log. of 5 is 0.698970004336018804786 &c.
 { The hyp. log. of 5 is 1.6094379127 &c.

Ex. 6. Ans. { The com. log. of 11 is 1.0413926851582250407 &c.
 { The hyp. log. of 11 is 2.397895273016 &c.

(Key to Vol. II. page 331.)

POINTS OF INFLEXION.

Ex. 2. It is imagined the given equation was misprinted, $ay = a\sqrt{ax} + xx$ being intended; in which case $x = \frac{1}{4}a$, and $y = \frac{9a}{16}$. Ans.

Ex. 3. Ans.
$$\begin{cases} x = a(\sqrt{\frac{4}{3}} - 1)^{\frac{1}{2}} \\ y = a\sqrt{\sqrt{\sqrt{\frac{4}{3}} - 1} + (\sqrt{\frac{4}{3}} - 1)^{\frac{3}{2}}} \end{cases}$$

Ex. 4. Put a for PF , b for $EI=AF$, x for FD , and y for DE . Then $x^2 = (a+y)^2 \times (b^2 - y^2) \times y^{-2}$. Whence $y^3 + 3ay^2 = 2ab^2$, an equation in which y is determinable. Now if q be assumed as the value of y in this cubic equation, the value of x is $(a^2b^2q^{-2} + 2ab^2q^{\frac{1}{2}} + b^2 - a^2 - 2aq - q^2)^{\frac{1}{2}}$; and if $a=b$, then $y = a\sqrt{3} - a$, and $x = (a^4q^{-2} + 2a^3q^{\frac{1}{2}} - 2aq - q^2)^{\frac{1}{2}}$. Ans.

RADIUS OF CURVATURE.

(Page 333.)

Ex. 2. Here $y = \frac{c}{a}\sqrt{ax - x^2}$; hence $\dot{y} = \frac{c^2\dot{x}(a-2x)}{2a^2y} = \frac{(ac+2cx)\dot{x}}{2a\sqrt{ax-x^2}}$, and $-\ddot{y} = \frac{a^2\dot{y}^2 - c^2\dot{x}^2}{a^2y} = \frac{ac\dot{x}^2}{4(ax-x^2)^{\frac{3}{2}}}$.

Wherefore $\sqrt{\dot{y}^2 + \dot{x}^2} = \frac{\sqrt{c^2\dot{x}^2(a-2x)^2}}{4a^2(ax-x^2)} + \dot{x}^2 = \frac{\dot{x}}{2a} \times \frac{\sqrt{a^2c^2 + 4(ax-x^2)(a^2-c^2)}}{ax-x^2} = \dot{z}$; whence it follows that

$$\frac{\dot{z}^3}{-\dot{x}\ddot{y}} = \frac{[a^2c^2 + 4(ax-x^2)(a^2-c^2)]^{\frac{3}{2}}}{2a^4c} = \text{the radius of curvature required.}$$

(Key to Vol. II. page 333.)

Ex. 3. In this Example $y = \frac{c}{a} \sqrt{ax+x^2}$; and $\dot{y} = \frac{(a+2x)c^2 \dot{x}}{2a^2 y}$
 $= \frac{(ac+2cx)x \dot{x}}{2a \sqrt{ax+x^2}}$, also $-\ddot{y} = \frac{ac \dot{x}^2}{4(ax+x^2)^{\frac{3}{2}}}$.


Consequently $\sqrt{\dot{y}^2 + \dot{x}^2} = \sqrt{\frac{(a+2x)^2 c^2 \dot{x}^2}{4a^2(ax+x^2)} + \dot{x}^2} =$
 $\frac{\dot{x} \sqrt{c^2(a+2x) + 4a^2(ax+x^2)}}{4a^2(ax+x^2)} = \frac{\dot{x} \sqrt{a^2 c^2 + (a^2 + c^2) \times 4(ax+x^2)}}{2a \sqrt{ax+x^2}}$

$= \dot{z}$. Wherefore $\frac{\dot{z}^3}{-\dot{x}\ddot{y}} = \frac{a^2 c^2 + (a^2 + c^2) \times 4(ax+x^2)^{\frac{3}{2}}}{2a^4 c}$ the

radius of curvature required.

Ex. 4.

Let ABC be a cycloid, BEDF its generating circle, and AC the right line on which the circle rolls. Also let AD be equal to DC, and DB a diameter of the circle, at right angles to AC. Assume in the curve any point H, of which the center and radius of curvature are required; and draw the ordinate HG. If



from E the point of intersection of this ordinate and the generating circle, a right line be drawn to D, ED shall equal half the radius of curvature at the point H, and the center K shall be in a straight line parallel to ED, and passing through H; that is, HI shall be half of HK. For, put a for the radius of the generating circle, x for the absciss BG, y for the ordinate GH, and w for the arc BE. Then by the property of a circle $GE = \sqrt{2ax - x^2}$; and $y = w + \sqrt{2ax - x^2}$ by the property of a cycloid. Consequently

(Key to Vol. II. page 383.)

$$\dot{y} = \dot{w} + \frac{ax - xx}{\sqrt{2ax - x^2}}. \text{ Now } \dot{w} = \frac{ax}{\sqrt{2ax - x^2}}, \text{ wherefore } \dot{y} = \frac{2ax - xx}{\sqrt{2ax - x^2}} = \frac{\dot{x}}{x} \sqrt{2a - x}; \text{ hence } \ddot{y}, \text{ considering } \dot{x} \text{ constant, is } \frac{-a\dot{x}^2}{x\sqrt{2ax - x^2}}.$$

These values of \dot{y} and \ddot{y} being substituted in the general expression for $r = \frac{(x^2 + y^2) \times \sqrt{x^2 + y^2}}{-\dot{x}\dot{y}} = \frac{(1 - y^2)\sqrt{1 - y^2}}{-\ddot{y}}$, the value of r is $2\sqrt{4a^2 - 2ax}$.* Ans.

COR. 1. When $x = 0$, the radius of curvature is 4 times the radius of the generating circle.

COR. 2. When $x = 2a$, the radius of curvature vanishes.


Hence L is the center, and LB the radius of curvature to the point B ; and if h be assumed as any other point in the curve, (gh being the ordinate and bg the absciss,) hi is equal and parallel to FD , and $kh(=2ih) =$ the radius of curvature from the center k , to the curve at h .

* This expression is equal to $2\sqrt{d^2 - dx}$ by considering $a = \frac{1}{2}d$, and recognizing in d the diameter of the generating circle.

INVOLUTE AND EVOLUTE CURVES.

(Page 272.)

EX. 2. Let ABC be half the curve of a common cycloid, and ADE its evolute, D being the center, and DB the radius of curvature to the point B ; also let Bm and FD be parallel to AG the line on which the generating circle of the given curve rolls, whilst od and Br are at right angles to that line. By the property of the cycloid the arc $ABC = 2CG$. If, therefore, a be put for ABC , x for Av , y for Bv , and z for AB , then $BC = a - z$, $cm = \frac{1}{2}a - y$, and, As $a^2 : \frac{1}{2}a :: a - z^2 : \frac{1}{2}a - y$. Consequently



(Key to Vol. II. page 333.)

$y = \frac{2az - z^2}{2a}$, and $\dot{y} = \frac{a\dot{z} - z\dot{z}}{a}$. Now $\dot{y} = \sqrt{\dot{z}^2 - \dot{x}^2}$, or $\dot{y}^2 = \dot{z}^2 - \dot{x}^2$, hence $\dot{x}^2 = \frac{\dot{z}^2(2az - z^2)}{a^2}$, $\dot{x} = \frac{\dot{z}\sqrt{2az - z^2}}{a}$, and (considering \dot{z} constant,) $\ddot{x} = \frac{\dot{z}^2(a - z)}{a\sqrt{2az - z^2}}$; hence $\frac{\dot{y}\ddot{z}}{\ddot{x}} = \sqrt{2az - z^2} = DB = DA$.

But when a and z are equal, by the coincidence of EDB with EGC , ADE also coincides with, and is equal to CGE ; and $GE = AH = GC = \frac{1}{2}a$. Therefore, because it is plain that, As $AE^2 : AD^2 :: AH : AF$, it follows that, the evolute ADE is a semi-cycloid equal and similar to the involute ABC , but in a contrary direction. Consequently the evolute of the common Cycloid is a pair of contiguous semi-cycloids in contrary directions, the generating circles of both involute and evolute being equal, and rolling on parallel lines, of which the distance is a diameter of either circle. Q. E. D.

END OF VOL. II.

(continued from page 217)

$$\frac{2x^2 - 3x + 1}{x^2 - 2x + 1} = \frac{2x^2 - 3x + 1}{(x-1)^2} = \frac{2(x-1)^2 + 3(x-1) + 4}{(x-1)^2} = 2 + \frac{3(x-1) + 4}{(x-1)^2}$$

$$= 2 + \frac{3x - 3 + 4}{(x-1)^2} = 2 + \frac{3x + 1}{(x-1)^2}$$

$$= 2 + \frac{3(x-1) + 4}{(x-1)^2} = 2 + \frac{3(x-1)}{(x-1)^2} + \frac{4}{(x-1)^2} = 2 + \frac{3}{x-1} + \frac{4}{(x-1)^2}$$

But when a root is one, the remainder of the division is not zero, and the remainder is equal to the remainder of the division of the numerator by the denominator. In this case, the remainder is 4, and the remainder of the division of the numerator by the denominator is 4. This is because the remainder is 4, and the remainder of the division of the numerator by the denominator is 4. This is because the remainder is 4, and the remainder of the division of the numerator by the denominator is 4.

END OF VOLUME

K E Y

TO

HUTTON'S COURSE OF MATHEMATICS.

ISOPERIMETRY.

(Key to Vol. III. page 52.)

- Ex. 1. $36 \div 3 = 12$ side of the equilateral triangle.
 $36 \div 4 = 9$ side of the square of equal perimeter.
 $36 \div 6 = 6$ - - - - - hexagon.
 $36 \div 12 = 3$ - - - - - dodecagon.

Also $\sqrt{12}^2 \times 0.4330127 = 62.3538288$ area of the eq. lat. triang.

And $9^2 \times 1.0000000 = 81.0000000$ area of the square.

So $6^2 \times 2.5980762 = 93.5307432$ area of the hexagon.

And $3^2 \times 11.1961524 = 100.7653716$ area of the dodecagon.

Lastly $\sqrt{36}^2 \times .07958 = 103.1356800$ area of the circle.

Ex. 2. Because a triangle whose sides are 3, 4, 5, is right angled, it follows that,

$3 \times \frac{4}{2} = 3 \times 2 = 6.0000$ is the area of the right angled triangle.

Now $\frac{3+4+5}{3} = 4$ is the side of the equilateral triangle of equal perimeter.

But $4^2 \times 0.4330127 = 6.9282032$ the area of the equilateral triangle. Therefore $.9282032$ is the required difference.

Ex. 3. By Theor. v. Elem. of Isoperimetry (see page 36, vol. iii.) the angle contained by the sides 8 and 11 is a right angle; therefore 44 is the area required. And $\sqrt{8^2+11^2} = \sqrt{185} = 13.6014705$ the length of the third side. Q. E. I.

(Key to Vol. III. page 52.)

- Ex. 4. $\overline{12}^2 \times .07958 = 11.45952$ the area of the circle.
 But (*Theor.* xi. page 36. *vol.* iii.) it will be,
 As 12 : 15, $\therefore 4 : 5 :: 11.45952 : 14.3244$ the area of the
 polygon.
 Whereby the Answer is obvious.

(Page 53.)

- Ex. 5. By *Theor.* xxiii. page 44, *vol.* iii. the parallelopiped required is a cube; therefore $18 \div 3 = 6 =$ the length of the side of the cube. Consequently $6 \times 6 \times 6 = 216$ is an expression for both the surface and solidity. Q. E. I.
- Ex. 6. It is evident by *Theor.* xxiii. Isoperimetry, and by the last Example that, the square prism proposed is a cube; therefore $546 \div 6 = 91 =$ the surface of any one of the sides. But $\sqrt{91} = 9.539392 =$ the side of the cube: consequently $\overline{9.539392}^3 = 868.084669421158924288$ is the solidity required.
- Ex. 7. The surface of the cylinder being a *minimum*, the species (*Theor.* xxvii. *Isoperimetry*) is the Archimedean. But a cylinder circumscribing a sphere, is to the sphere as 3 : 2; therefore 113.097312 is the solidity of the sphere circumscribed, of which the diameter is 6. Consequently the surface of the cylinder, is expressed by the same figures as its solidity namely, by 169.645968 &c. Q. E. I.
- Ex. 8. Because, by the question, the solidity is a *maximum*, the slant side (*Theor.* xxxiii.) is triple the radius of the base; and consequently the whole surface of the cone quadruple the area of the base. But $201.061952 \div 4 = 50.265488$, which divided by $.78539 = 64$ the square of the diameter of the base of the cone. Hence the radius of the base is 4; the length of the slant side 12; and the altitude $\sqrt{128} = 11.3137085$, of which $\frac{1}{3}$ part is 3.77123617, which \times ^{int} 50.265488 (the area of the base) = 189.56302643830096. Ans.
- Ex. 9. Since the capacity of the pyramid, given in species, is a *maximum*, it is evident (*Theor.* xxxv.) that, the length of the slant side is triple the radius of a circle inscribed in the base; and (*Theor.* i. *Cor.* 2.) that, the base is equilateral. Now the side of an equilateral triangle circumscribing a circle whose diameter is d , is to that diameter, as $d\sqrt{3}$ to d . Consequently the areas are as 1.2990381 to .7854. But (*Theor.* xxxiv.) the

(Key to Vol. III. page 53.)

ratio of the surfaces of the bases of a right cone and its circumscribing pyramid, is the ratio of the surfaces *total*.

Therefore

As $1.2990381 : .7854 :: 43.30127 : 26.18$ the surface of the inscribed cone. Hence, as in the last example, it will be, $26.18 \div 4 = 6.545 =$ the area of the cone's base, which divided by $.78539 = 8.33343$ the square of the diameter of the base of the cone. Hence the radius is 1.443 &c. Also the slant side of the cone (and of the triangle) is 4.33 . Moreover, the altitude is $\sqrt{4.33^2 - 1.443^2} = \sqrt{18.7489 - 2.082249} = \sqrt{16.66666} = 4.08248$. Again 2.886 (the diameter of the circle inscribed in the base,) $\times 1.7320508$ (the square root of 3) $= 4.9986986 =$ the side of the equilateral triangle forming the base, and therefore the area of the base of the pyramid $= 10.782$ nearly.

Lastly, $\frac{10.782 \times 4.08248}{3} = 14.67243$ &c. *the capacity sought.*

(Di.) (Di.) (Di.) (Const. dec.)

Ex. 10. *For the sphere.* $20 \times 20 \times 20 \times .5236 = 4188.8$ the solidity of the sphere.

For the cone. As $4 : 9 :: 4188.8 : 9424.8$ the solidity of the cone.

And *For the cylinder.* As $2 : 3 :: 4188.8 : 6283.2$ the solidity of the cylinder. Q. E. I.

Ex. 11. The surface 28.274337 (of the sphere,) divided by $3.141593 = 8.1 =$ the square of the diameter of the sphere; therefore the diameter is $= 2.846$. But $28.274337 \times 2.846 \times \frac{1}{6} = 13.410518$ the solidity of the sphere. Now (*Theor. xxxvi. Isoperimetry, vol. iii.*) it will be, As $28.274337 : 35 :: 13.410518 : 16.6$ the solidity of the irregular polyedron circumscribed about the sphere.

Ex. 12. *For the sphere.* The solidity (500) of the sphere divided by $.5236$ gives 954.927 &c. for the cube of the diameter of the sphere, hence the diameter is 9.84744 . But the content of any sphere divided by $\frac{1}{6}$ of the diameter gives the surface; therefore $500 \div 1.64124 = 304.6477$ THE SURFACE OF THE SPHERE.

For the cylinder. As $3 : 2 :: 500 : 333\frac{1}{3}$ the solidity of the inscribed sphere (*Theor. xxvi. def. page 45, vol. iii.*). And $333\frac{1}{3} \div .5236 = 636.618 =$ the cube of the diameter, whence the diameter $= 8.6025$. And $74.0035^* \times .7854 \times 6 = 348.7340934$ THE SURFACE OF THE CYLINDER.


* The square of 8.6025 .

(Key to Vol. III. page 53.)


Lastly, for the cone. As $9 : 4 :: 500 : 222\frac{1}{3}$ the solidity of the inscribed sphere; and $222\frac{1}{3} \div \cdot 5236 = 424\cdot 412$ the cube of the diameter, from which the diameter is obtained $7\cdot 515$. But $222\frac{1}{3} \div 7\cdot 515 \times \frac{1}{6} = 177\cdot 423$ nearly = the surface of the sphere; consequently it will be, As $4 : 9 :: 177\cdot 423 : 399\cdot 2017$ THE SURFACE OF THE EQUILATERAL CONE. Now $7\cdot 515 \times \sqrt{3} = 13\cdot 016361762$ the diameter of the base, and length of the slant line. Which were to be determined.

Ex. 13. For the circumscribing cylinder.

Because, As $1\cdot 000 : 3\cdot 1415926535$ &c. so is the diameter of a circle, to the circumference; and that, radius $\times \frac{1}{2}$ circumference = the area = $\cdot 7853981634$ nearly; and because (it is prov'd page 49, vol. ii.) that, the superficies of the sphere is equal to the curve surface of its circumscribing cylinder = diameter \times circumference = $3\ 1415926535 = 4$ times the area of the base = 4 times the area of a great circle of the sphere, it follows that the whole area of the cylinder ABCD is equal to 4 great circles + the circle CSDH + the opposite circle ARBG; that is, equal to 6 great circles of the inscribed sphere. Therefore the cylinder's convex and whole surface are, respectively as 4 and 6, to the surface of the inscribed sphere, whose superficies is 4.



For the circumscribing equilateral cone. Because, when $MN=1$, then $IN=\sqrt{3}$, (since, As Rad. (MN) : 1 :: Tang. 60° ($\angle IMN$) : $\sqrt{3}$, and the doubles in the same ratio), it is evident that the circumference $IOLP = 3\cdot 1415926535897 \times \sqrt{3}$; and, IK being by the question equal to IL , the convex surface of the cone is truly expressed by $\frac{1}{2}\sqrt{3} \times 3\cdot 1415926535897 \times \sqrt{3} = \frac{1}{2} \times 3\cdot 1415926535 \times \sqrt{3} \times \sqrt{3} = \frac{1}{2} \times 3\cdot 1415926535 \times 3 = 1\frac{1}{2} \times 3\cdot 1415$ &c. = $4\cdot 71238895025$. But the area of the circle $IOLP$ (or, the area of the cone's base to rad. $\frac{1}{2}\sqrt{3}$) is $\sqrt{3} \times \sqrt{3} \times \cdot 7853981633$ &c. = $3 \times \cdot 78539$ &c. = $2\cdot 356194490125$; which is to the convex surface of the cone as 1 to 2; or, as 3 to 6. Lastly, the surface of a globe, whose diameter is unity, being, as above, $3\cdot 1415926535$ = the surface of the sphere inscribed in the equilateral cone IKL, it is, As $3\cdot 1415926535 : 4\cdot 71238895025 :: 4 : 6$. And, As $3\cdot 1415926535 : 4\cdot 71238895025 + 2\cdot 356194490125 :: 4 : 9$.



Which was to be shown.

(Key to Vol. III. page 53.)

Ex. 14. Since, in the last example, the surfaces, of a sphere, its circumscribing cylinder, and circumscribing equilateral cone are demonstrated to be in the ratio of the numbers 4, 6, and 9, it is plain, by Theor. xxxvi. Isoperimetry, that their solidities are in the same ratio. Which was required to be proved.

ANALYTICAL PLANE TRIGONOMETRY.

(Page 76.)

Ex. 7. Because $\sin. 18^\circ = \text{co. sin. } 72^\circ = \frac{1}{2}$ the side of a decagon inscribed in a circle $= \frac{1}{2}\sqrt{\frac{5}{4}} - \frac{1}{4}$, if the surd be reduced to simpler terms by \times lying the denominator of $\frac{1}{2}$ by 4, and the numerator of the surd by 4 in the form of the square root ($=16$.) there arises $\frac{1}{2}\sqrt{\frac{5}{4}} = \frac{1}{8}\sqrt{\frac{80}{4}} = \frac{1}{8}\sqrt{20} = \frac{1}{8}\sqrt{4}\sqrt{5} = \frac{1}{8} \times 2 \times \sqrt{5} = \frac{1}{4}\sqrt{5}$; therefore $\frac{1}{2}\sqrt{\frac{5}{4}} - \frac{1}{4} = -\frac{1}{4} + \frac{1}{4}\sqrt{5}$, which \times cd by the rad. $= \text{rad.} \times (-\frac{1}{4} + \frac{1}{4}\sqrt{5}) = \frac{1}{4} \text{ rad.} \times (-1 + \sqrt{5})$. And so for $\sin. 54^\circ = \text{co. sin. } 36^\circ$ of which the equation is $\frac{1}{2}\sqrt{\frac{5}{4}} + \frac{1}{4}$ &c. Q. E. D.

Ex. 8. If P and Q be any two arcs whatever, since $\sin. (P+Q) = \sin. (P-Q) + 2 \cos. P \times \sin. Q$, it is evident that when $P=60^\circ$, then $\cos. P = \cos. 60^\circ = \text{half the radius} = \frac{1}{2}$, and the $\sin. (60^\circ + Q) = \sin. Q + \sin. (60^\circ - Q)$. Q. E. D.


Ex. 9. *First.* Let A , and $60^\circ - A$, be the two arcs, then $\frac{1}{2}$ their diff. is $A - 30$. But, As $\sin. (A - 30^\circ) : \sin. A - \sin. (60^\circ - A) = \sin. A \times \cos. 30^\circ - \sin. 30^\circ \times \cos. A :: \sin. A - \sin. 60^\circ \times \cos. A + \sin. A \times \cos. 60^\circ = \sin. A \times \cos. 30^\circ - \sin. 30^\circ \times \cos. A : \frac{3}{2} \sin. A - \sin. 60^\circ \times \cos. A = \frac{1}{2}\sqrt{3} \times \sin. A - \frac{1}{2} \cos. A : \frac{3}{2} \sin. A - \frac{1}{2}\sqrt{3} \times \cos. A \therefore 1 : \sqrt{3}$.

Next. Let A and $90^\circ - A$ be the two arcs; then $\frac{1}{2}$ their difference is $A - 45^\circ$. But,


As $\sin. (A - 45^\circ) : \sin. A - \sin. (90 - A) = \sin. A \times \cos. 45^\circ - \sin. 45^\circ \times \cos. A :: \sin. A - \cos. A = \frac{1}{\sqrt{2}} : 1 \therefore 1 : \sqrt{2}$. Q. E. D.

(Key to Vol. III. page 76.)


Ex. 10. Let ABC be any arc whatever, bisected in the point B ; then AD is the sine of the arc ABC , and DC is its versed sine. But the angle ADC is a right angle, and the square $AEFD$, together with the square $CDGH$, equal to the square $AEKC$ (*Eucl. i. 47*) Now AC the chord of the arc ABC is bisected at L , and AL ($=LC$) is the sine of the arc $AB = BC = \frac{1}{2} ABC$, therefore &c. Q. E. D.



Ex. 11. If AF be any arc whatever, and AFD its double, then is AG the sine of the mean arc, AD twice the sine of the mean arc, AC the versed sine of double the arc AF , and AB the rad. of the circle. But, As $AB : AG :: AD : AC$, that is, As rad. : sin. mean arc. :: twice sin. mean arc : vers. sin. double arc; or, putting a for the radius, b for the sine of the arc AF , and c for the half of AC , it will be, As $a : b :: 2b : 2c$, wherefore $\frac{1}{2}a : b :: b : 2c$. Hence "the sine of an arc," &c. Q. E. D.




Otherwise. Let AB be an arc of a circle, ABC its double, and AGC the chord of the double arc; then DCA being an angle in a semicircle is a right angle, and AGC is a mean proportional between the diameter AD and the segment of the base AP . (*Eucl. vi. 8.*) But $4AG^2 (=4 \sin^2$ of the arc $AB) = AGC^2 = AD \times AP = 2AE \times AP$. That is, $AG^2 = \frac{1}{2}AE \times AP$. Wherefore AG ($=\sin$ arc AB .) is a mean proportional between half the radius and AP (the versed sine of double the arc.) Q. E. D.



(Key to Vol. III. page 76.)


Ex. 12. Let BAC be any angle whatever, AB its secant, and CB its tangent. In BA take BD equal to BC , and join CD ; then parallel to DC draw AE , and produce BC to E . Because BD and BC are equal, their opposite angles are equal: but the angle BDC is evidently equal to BAC together with half its complement; therefore the angle BEA is equal to the angle BAC together with half its complement. Whence it follows, that the angle CAE is half the complement of BAC . Now CE is the tangent of the angle CAE , and equal to AD : wherefore EB is equal to AB ; that is, the secant of BAC is equal to the tangent of the same angle, together with the tangent of half its complement.



Q. E. D.


* * This problem may likewise be demonstrated by the algebraical expressions given in the III. chap. of Hutton's 3d volume, or in WOODHOUSE's elegant *Treatise on Plane and Spherical Trigonometry*, Cambridge, 1813.

Ex. 13. Let ABC be the triangle, whereof BC is the base; and since by the hypothesis AB is unequal to AC , AD may be supposed equal to AB . Join BD , and from A draw AE at right angles to BD . Then, because ABD is isosceles, and the vertical \angle at A the same in the two triangles ABD and ABC , it is evident that the



sum of the angles at the base of the one triangle is equal to that of those at the base of the other; and that the $\angle ABD$ is half the sum of the $\angle^{les} ABC, ACB$; but the $\angle CBD$ is the excess of the $\angle ABC$ above that half sum, consequently equal to half their difference. Now, As $BC : DC$ (the difference of the sides) $:: \sin. \angle ADB (= \angle ABD) : \sin. \angle DBC$.


Again, for the second Theorem of the Example. Let ABC be the given triangle, and produce CB indefinitely; also make $BD = AB$. Join AD , and from B draw BE perpendicular to AD , and BF parallel to AC . Then it is manifest that the $\angle BDA = \angle DAB = \frac{1}{2} \angle ABC$. It is likewise evident that the $\angle DBA$ (external \angle) is equal to both



(Key to Vol. III. page 76.)

the angles at the base of the triangle ABC , therefore the $\angle ABE$ equal to half the sum of $\angle^{les} BAC, ACB$; and EBF being the excess of the greater above that half sum, is equal to half their difference. But, As $AC : CD$ (sum of the sides) $:: \text{co. sin. } \angle DBE : \text{co. sin. } EBF$. Which was to be proved.

Ex. 14. Because the three arcs constitute an entire circumference, and are to one another as 1, 2, 4, it follows that their chords (see the accompanying figure, wherein the three chords are $DB, BA,$ and AD) are 177.9, 142.76, 79.24. This question may likewise be solved by the common rules of Trigonometry, since it only requires a triangle of which the perimeter shall be 400, and the angles $25^{\circ}\frac{5}{7}, 51^{\circ}\frac{3}{7},$ and $102^{\circ}\frac{6}{7}$; for the sines of the angles being as the sides which subtend them, it is evident that $AD=79.239$ &c. $BD=177.924$ &c. and $AB=142.756$. Which was required to be shown.



Ex. 15. Because (page 57, vol. iii.) “The sides of plane triangles are proportional to the sines of their opposite angles, and vice versa,” the sides must also be as 17, 15, and 8. But 160 divided into three parts in that ratio, gives 68, 60, and 32 for the sides of the triangle. Consequently the angles are $89^{\circ}..58', 61^{\circ}..57',$ and $28^{\circ}..5'$. Which was required.

Ex. 16. The greater logarithm = 2.5378191
 The less logarithm = 2.2407293


—————
 The difference of the \log^s . is 0.2970898

Adding 10 to the index* of this difference, the result is the log. tangent of $63^{\circ}..13'$, from which when 45° have been taken, the remainder is $18^{\circ}..13'$, of which the $\log. \text{ tang.} = 9.5173353$. And the $\log. \text{ co. tang.}$ of $18^{\circ}..40'$ (half the given angle) = 10.4712979 , of which the sum (abating 10 in the index for the 10 added to the first difference) is $9.9886332 = \log. \text{ tang.}$ of $44^{\circ}..15'$, therefore $115^{\circ}..35'$ and $27^{\circ}..5'$ are the angles required. (See vol. iii. page 70.)

* It will be observed, that in many of our solutions we scrupulously follow Dr. Hutton's Rules, whereas in others we abbreviate the operation by means not mentioned in the COURSE. In this Example the addition of 10 to the index of the $\log.$ difference and the subsequent abatement of 10 in the $\log.$ sum of the tangent and co-tangent might be avoided, and less liability to error ensue, if the $\log.$ tangents, &c. to radius 1, were used. For this use see various parts of the KEY.

(Key to Vol. III. page 76.)

Ex. 17. The fractions $\frac{1}{3}$, $\frac{1}{4}$, $\frac{1}{5}$, being to one another as the numbers 20, 15, and 12, if these be assumed as the sides of the triangle, the angles by the *second* formula (page 56, vol. iii.) are, $A=36^\circ .. 43'$, since



$$\frac{400+225-144}{2 \times 15 \times 20} = .801666 = \text{natural co. sin. } 36^\circ .. 43'.$$

$$B=94^\circ .. 56', \text{ because } \frac{144+225-400}{2 \times 12 \times 15} = -.086111 = \text{nat. co. sin.}$$

$$94^\circ .. 56'. \text{ and } c=48^\circ .. 21' \text{ since } \frac{144+400-225}{2 \times 12 \times 20} = .664583 =$$


$$\text{nat. co. sin. } 48^\circ .. 21', \text{ nearly. But by formula xxxiv. } \sin. \frac{1}{2} \angle A = \sqrt{\frac{(\frac{1}{2}s-b) \cdot (\frac{1}{2}s-c)}{bc}} = \sqrt{\frac{3.5 \times 8.5}{300}} = .3149073 = \sin. 18^\circ .. 21' .. 23'',$$

therefore $A=36^\circ .. 42' .. 46''$.

$$\text{And } \sin. \frac{1}{2} \angle B = \sqrt{\frac{(\frac{1}{2}s-a) \cdot (\frac{1}{2}s-c)}{ac}} = \sqrt{\frac{11.5 \times 8.5}{180}} = .736925$$

$= \sin. 47^\circ .. 28' .. 1''$, therefore $B=94^\circ .. 56' .. 2''$, consequently $c=48^\circ .. 21' .. 12''$. Q. E. I.

Ex. 18. Let AB be the secant of 60° and CD the tangent of 45° ; and join CE. Then because DAC is an angle of 45° , and ACD a right angle, the angle CDA is likewise 45° , and $CD=AC=Rad.$ Again because EAC is an angle of 60° and $AE=AC$, the triangle AEC is equilateral, and $CE=Rad.$ But the angle at $B=30^\circ$, and the angle $BCE = \text{compl. } 60^\circ = 30^\circ$; therefore $EB=EC=Rad.=DC$. Now $AE=EB$, that is, $AB = \text{twice } DC$. Moreover it is manifest that the square of AD (that is, of the secant of 45°) $=2r^2$, and $AB (=2r = \text{secant of } 60^\circ)$ multiplied by $CD (=r = \text{tang. } 45^\circ) = 2r^2$. Wherefore AD is a mean proportional between CD and AB. Q. E. D.



This problem (like most of the others which we prove geometrically) admits of easy demonstration by means of the algebraic expressions for the secant, tangent, &c. Whoever desires to see these subjects treated in a manner as picturesque as elegant, is referred to Woodhouse's Trigonometry, 2nd edition.

(Key to Vol. III. page 77.)

Ex. 19. Because the rectangle of the sines of any two arcs is equal to the rectangle of half the radius and the difference of the co-sines of the sum and difference of the two arcs $= \frac{1}{2}$ sum of the chords $\times \frac{1}{2}$ difference of the chords of the two arcs $= \frac{1}{4}sd$, (where s is the sum and d the difference of the chords,) it follows that 4 times the rectangle of the sines of the two arcs $= sd$. But the difference of the squares of the chords is equal to a rectangle under the sum and difference of those chords $= sd$; therefore "4 times the rectangle &c." Q. E. D.

Otherwise.

Let A and B represent any two unequal arcs, whereof A is the greater; then, since $\sin. (P+Q) \times \sin. (P-Q) = \sin^2. P - \sin^2. Q$, put $P+Q$ for A , and $P-Q$ for B , so shall $P = \frac{1}{2}(A+B)$, and $Q = \frac{1}{2}(A-B)$; and $\sin. A \times \sin. B = \sin^2. \frac{1}{2}(A+B) - \sin^2. \frac{1}{2}(A-B)$.

Consequently $4(\sin. A \times \sin. B) = 4 \times \sin^2. \frac{1}{2}(A+B) - 4 \times \sin^2. \frac{1}{2}(A-B) = \text{chord}^2 \text{ of } \frac{1}{2}(A+B) - \text{chord}^2 \text{ of } \frac{1}{2}(A-B)$. Which was to be proved.

Ex. 20.

$$\sqrt{\frac{(\frac{1}{2}s-b) \cdot (\frac{1}{2}s-c)}{bc}} = \sqrt{\frac{\frac{5+6+7}{2} - 6 \times \frac{5+6+7}{2} - 7}{6 \times 7}} = \sqrt{\frac{1}{7}}$$

Also,

$$\sqrt{\frac{(\frac{1}{2}s-a) \cdot (\frac{1}{2}s-c)}{ac}} = \sqrt{\frac{\frac{5+6+7}{2} - 5 \times \frac{5+6+7}{2} - 7}{5 \times 7}} = \sqrt{\frac{3}{35}}$$

And,

$$\sqrt{\frac{(\frac{1}{2}s-a) \cdot (\frac{1}{2}s-b)}{ab}} = \sqrt{\frac{\frac{5+6+7}{2} - 5 \times \frac{5+6+7}{2} - 6}{5 \times 6}} = \sqrt{\frac{2}{5}}$$

Now* $10 + \log. \text{ of } \sqrt{\frac{1}{7}} = 9.5774504 = \log. \sin. 22^\circ .. 12' .. 27''$.

$10 + \log. \text{ of } \sqrt{\frac{3}{35}} = 9.6795106 = \log. \sin. 28^\circ .. 33' .. 42''$.

$10 + \log. \text{ of } \sqrt{\frac{2}{5}} = 9.8010299 = \log. \sin. 39^\circ .. 13' .. 51''$.


Wherefore,

$$\left. \begin{aligned} \angle A &= 2(22^\circ .. 12' .. 27'') = 44^\circ .. 24' .. 54'' \\ \angle B &= 2(28^\circ .. 33' .. 42'') = 57^\circ .. 7' .. 24'' \\ \text{And } \angle C &= 2(39^\circ .. 13' .. 51'') = 78^\circ .. 27' .. 42'' \end{aligned} \right\} \text{Ans.}$$

* See page 74, vol. iii.

(Key to Vol. III. page 77.)

Ex. 21. Let ABC be a semi-circle, BD the tangent of the arc BE, and EC its co-tangent.



Then because $\frac{\text{rad.}^2}{\text{tang.}}$ A


$= \text{co-tang.}$, if x be put for the length of the tangent (BD) and a for the radius of the circle, then $\frac{a^2}{x} = \text{the co-tangent (EC)}$. But

by the question $4a = x + \frac{a^2}{x}$; that is, $x^2 - 4ax = -a^2$. Hence x^2

$- 4ax + 4a^2 = 3a^2$; and $x = 2a \pm \sqrt{3a^2} = 2a \pm a\sqrt{3}$. Now when radius is unity this equation becomes $x = 2 \pm \sqrt{3} = 3.7320508$ or $.2679492$, of which the logarithms are $.5719475$ and $-1.4280525 = \text{the log. tangents}^*$ of 75° and 15° , either of which is the arc required.

* To radius 1. If 10 be added to the index, the log. sum will be the common log. tangent.

Ex. 22. Let ABC be a quadrant of a circle, of which the center is D; and let the radii DA and DC be produced indefinitely towards F and G. Then in DG take Dk equal to a , and in DF take Dl equal to Dk. Join kl, so shall kl, if it cut or touch the quadrant, indicate two arcs either of which will answer the conditions. Ar, therefore, is an



arc of which the sine added to the cosine is equal to a , that is, equal to Dk; which is evident, since Dq is equal to the sine, and $qr = qk$, equal to the cosine. Or, Av, is the arc, in which case $ov = di$ is the sine, and $vi = ik$, the cosine; together equal to Dk = a .

By which it is manifest, that the limits of possibility are double the sine of 45° on the one hand, and the sine of 90° on the other; for when a exceeds the former limit, and is equal to Ds, then st neither cuts nor touches the quadrant; and when a falls short of the latter limit and is equal to Dn, the line nm cannot meet the curve. Consequently DC is the least, and DE the greatest possible value of a . Q. E. I.


(Key to Vol. III. page 77.)

Ex. 23. If AB represent the arc sought, AF its tangent, CF its secant, and DE its co-tangent, then by the question $CF=DE$. But the

$$\text{co-tang.} = \frac{\text{rad.}^2}{\text{tang.}}, \text{ and sec.} =$$

$\sqrt{\text{rad.}^2 + \text{tang.}^2}$ Putting, therefore, x for AF (the tangent of the arc required) the equation is $\frac{\text{rad.}^2}{x}$

$= \sqrt{\text{rad.}^2 + x^2}$, that is, (if radius be unity,) $1 = x^2 + x^4$, whence, by completing the square of this quadratic, $x^4 + x^2 + \frac{1}{4} = \frac{5}{4}$, and $x^2 = -\frac{1}{2} \pm \sqrt{\frac{5}{4}}$, consequently $x = \sqrt{-\frac{1}{2} \pm \sqrt{\frac{5}{4}}} = .786151$ of which the log. is $.8955060 = \log. \text{ tang.}$ of $38^\circ .. 10' .. 21'' .. 42''' = \text{the arc AB. Q. E. I.}$



RESOLUTION OF SPHERICAL TRIANGLES.

(Page 108.)

Ex. 2. $\text{Tang. } c = \text{tang. } a \times \text{co. sin. } B = \log. \text{ tang. } a + \log. \text{ co. sin. } B - 10 = 10.6851149 + 9.9469372 - 10 = 10.6320521 = \log. \text{ tangent of } 76^\circ .. 52' = \text{the side } c$. And $\text{sin. } b = \text{sin. } a \times \text{sin. } B = \log. \text{ sin. } A + \log. \text{ sin. } B - 10 = 9.9909338 + 9.6680265 - 10 = 9.6589603 = \log. \text{ sin. of } 27^\circ .. 8' = \text{the side } b$. And $\text{tang. } c = \text{co. tang. } B \div \text{co. sin. } A = \log. \text{ cot. } B - \log. \text{ co. sin. } a + 10 = 10.2789107 - 9.3058189 + 10 = 10.9730918 = \log. \text{ tang. of } 83^\circ .. 56' = \text{the angle } c$. Ans.

Ex. 3. $\text{Tang. } a = \text{tang. } b \div \text{co. sin. } c = \log. \text{ tang. } 117^\circ .. 34' - \log. \text{ co. sin. of } 31^\circ .. 51' + 10 = 10.2822906 - 9.9291289 + 10 = 10.3531617 = \text{the log. tang. of } 66^\circ .. 5' \text{ or } 113^\circ .. 55'$, which last $= a = \text{the hypotenuse}$. And $\text{cos. } B = \text{co. sin. } b \times \text{sin. } c = \log. \text{ cos. } 117^\circ .. 34' + \log. \text{ sin. } 31^\circ .. 51' - 10 = 9.6653749 + 9.7223848 - 10 = 9.3877597 = \log. \text{ co. sin. of } 75^\circ .. 52'$, or $104^\circ .. 8'$, whereof the latter $= B$, the third angle of the triangle.

Again, $\text{tang. } c = \text{sin. } b \times \text{tang. } c = \log. \text{ sin. } 117^\circ .. 34' + \log. \text{ tang. } 31^\circ .. 51' - 10 = 9.9476655 + 9.7932560 - 10 = \text{tang. } 28^\circ .. 51' = c$, the third and last side sought.

(Key to Vol. III. page 108.)

Ex. 4. $\text{Co. sin. } a = \text{co. sin. } b \times \text{co. sin. } c = \log. \text{co. sin. } 27^\circ .. 6' + \log. \text{co. sin. } 76^\circ .. 52' - 10. = 9.9494938 + 9.3564426 - 10 = 9.3059364 = \log. \text{co. sin. } 78^\circ .. 20' = a = \text{hypotenuse.}$ Also, $\text{Tang. } b = \text{tang. } b \div \text{sin. } c = \log. \text{tang. } 27^\circ .. 6' - \log. \text{sin. } 76^\circ .. 52' + 10 = 9.7090374 - 9.9884894 + 10 = 9.7205480 = \log. \text{tang. } 27^\circ .. 44' = \text{the angle } b.$ And, $\text{tang. } c = \text{tang. } c \div \text{sin. } b = \log. \text{tang. } 76^\circ .. 52' - \log. \text{sin. } 27^\circ .. 6' + 10. = 10.6320468 - 9.6585312 + 10 = 10.9735156 = \log. \text{tang. } 83^\circ .. 56' = \text{the angle } c.$ Which was to be done.

Ex. 5. $\text{Sin. } a = \text{sin. } b \div \text{sin. } B$, hence, $-1.8271887 = \log. \text{sin. } b.$
 $-1.8710735 = \log. \text{sin. } B.$

$\log. \text{diff. } -1.9561152 = \log. \text{sin. of an } \angle$
of $64^\circ .. 40' \frac{1}{2}$, or of its supplement $= a = \text{the hypotenuse.}$ And

$\text{Sin. } c = \text{tang. } b \div \text{tang. } B$, hence $-1.9574850 = \log. \text{tang. } b.$
 $0.0455626 = \log. \text{tang. } B.$

$\log. \text{diff. } -1.9119224 = \log. \text{sin. of an } \angle$
of $54^\circ .. 44'$, or of its supplement $= \text{the side } c.$ Lastly,

$\text{Sin. } c = \text{co. sin. } B \div \text{co. sin. } b$, hence, $-1.8255109 = \log. \text{cos. } B.$
 $-1.8697037 = \log. \text{co. sin. } b.$

$\log. \text{diff. } -1.9558072 = \log. \text{sin. of an } \angle$
of $64^\circ .. 35'$, or of its supplement $= \text{the angle } c.$

Ex. 6. Given $B = 48^\circ$, $c = 64^\circ .. 35'$, and A a right angle. Now
 $\text{co. sin. } a = \text{cot. } B \times \text{cot. } c.$ But

$\text{Log. cot. } 48^\circ .. 0'$ is -1.9544374

$\text{Log. cot. } 64^\circ .. 35'$ is -1.6768686

$\text{Log. cos. } 64^\circ .. 40' \frac{1}{2} = a$, is -1.6313060 log. sum.

And $\text{cos. } b = \text{cos. } B \div \text{sin. } c.$ Therefore

$\text{Log. cos. } 48^\circ$ is -1.8255109

$\text{Log. cos. } 64^\circ .. 35'$ is -1.9557890

$\text{Log. cos. } 42^\circ .. 12' = b$, is -1.8697219 log. diff.

Likewise $\text{cos. } c = \text{cos. } c \div \text{sin. } B.$ Hence,

$\text{Log. cos. } 64^\circ .. 35'$ being -1.6326576

$\text{Log. sin. } 48^\circ .. 0'$ being -1.8710735

$\text{Log. cos. } 54^\circ .. 44' = c$, is -1.7615841 log. sum.

Therefore $a = 64^\circ .. 40' \frac{1}{2}$, $b = 42^\circ .. 12'$, and $c = 54^\circ .. 44'$. Ans.

(Key to Vol. III. page 109.)

Ex. 7. Taking the *quadrantal* side *a* as radius; $115^\circ .. 20'$ the supplement of the $\angle A$, as hypotenuse; and $42^\circ .. 12'$ the $\angle C$ as a side; it will be (changing *like* for *unlike* in the determinations,) by Table 1. page 102,

$$\text{Sin. } c = \frac{\text{sin. } 42^\circ .. 12'}{\text{sin. } 115^\circ .. 20'}$$

$$\text{Cos. } b = \frac{\text{tang. } 42^\circ .. 12'}{\text{tang. } 115^\circ .. 20'}$$

$$\text{And Cos. } B = \frac{\text{cos. } 115^\circ .. 20'}{\text{cos. } 42^\circ .. 12'}$$

By Logarithms.

Log. sin. $42^\circ .. 12'$	= -1.8271887
Log. ar. co. of sin. $115^\circ .. 20'$	= 0.0439114

Log. sum	- 1.8711001*
Add 10 to the index	10. -----

Sin. $c = \text{sin. } 48^\circ .. 0' .. 13''\frac{1}{2}$	Log. 9.8711001

* -1.8711001 is the log. sin. $48^\circ .. 0' .. 13''\frac{1}{2}$ to radius 1.

And for the cos. *b* it is,

Tang. $42^\circ .. 12'$	- - - - - Log. -1.9574850
Tang. $115^\circ .. 20'$	- - - Log. ar. co. -1.6752372

Cos. $b = \text{cos. } 115^\circ .. 25' .. 14''\frac{1}{2}$	Log. -1.6327222

Lastly for the cos. *A* it will be,

Cos. $115^\circ .. 20'$	- - - - - Log. -1.6313258
Cos. $42^\circ .. 12'$	- - - - - Log. ar. co. 0.1302963

Cos. $B = \text{cos. } 125^\circ .. 16' .. 53''$	Log. -1.7616221

Wherefore $\left. \begin{array}{l} B = 125^\circ .. 16' .. 53'' \\ b = 115^\circ .. 25' .. 14''\frac{1}{2} \\ c = 48^\circ .. 0' .. 13''\frac{1}{2} \end{array} \right\} \text{Which were required.}$

(Key to Vol. III. page 110.)

Ex. 9. Given $c = 114^\circ .. 30'$, $a = 56^\circ .. 40'$, and $c = 125^\circ .. 20'$.
Hence it is by logarithms, (using the common indices,)

As Sin. $114^\circ .. 30'$ (c)	- Log. ar. co.	- 10.0409771
: Sin. $125^\circ .. 20'$ (c)	- - - - -	Log. 9.9115844
:: Sin. $56^\circ .. 40'$ (a)	- - - - -	Log. 9.9219401
<hr style="width: 100%;"/>		
: Sin. $48^\circ .. 30'$ (A)	- - - - -	Log. 9.8745016
<hr style="width: 100%;"/>		

And

$$\text{Co. tang. } \frac{1}{2} \angle B = \frac{\text{Tang. } \frac{1}{2} \text{ diff. of } A \text{ and } c \times \sin. \frac{1}{2} \text{ sum of } a \text{ and } c.}{\text{Sin. } \frac{1}{2} \text{ diff. of } a \text{ and } c.}$$

But $\frac{1}{2}$ diff. of A and $c = 38^\circ .. 25'$, $\frac{1}{2}$ sum of a and $c = 85^\circ .. 35'$,
and $\frac{1}{2}$ diff. of a and $c = 28^\circ .. 55'$; wherefore

+ Log. tang. $38^\circ .. 25'$	=	9.8993082
+ Log. sin. $85^\circ .. 35'$	=	9.9987084
- Log. sin. $28^\circ .. 55'$ ar. co.	=	- 10.3255703

Log. sum - - - - - 10.2135869 = Log. co.
tang. $31^\circ .. 27' \frac{1}{2}$, which \angle doubled = $62^\circ .. 55' = \angle B$. But

As Sin. $48^\circ .. 30'$ (A)	- - Log. ar. co.	- 10.1255439
: Sin. $56^\circ .. 40'$ (a)	- - - - -	Log. 9.9219401
:: Sin. $62^\circ .. 55'$ (B)	- - - - -	Log. 9.9495585
<hr style="width: 100%;"/>		
: Sin. $83^\circ .. 19'$ (b)	- - - - -	Log. 9.9970425

Whereby the Answer is obvious.

Ex. 10. Given $A = 48^\circ .. 30'$, $c = 125^\circ .. 20'$, and c (it was intended)
to be given) = $114^\circ .. 30'$. Therefore it will be

As Sin. $125^\circ .. 20'$ (c)	- - Log. ar. co.	- 10.0884156
: Sin. $114^\circ .. 30'$ (c)	- - - - -	Log. 9.9590229
:: Sin. $48^\circ .. 30'$ (A)	- - - - -	Log. 9.8745016
<hr style="width: 100%;"/>		
: Sin. $56^\circ .. 40'$ (a)	- - - - -	Log. 9.9219401

$$\text{And Tang. } \frac{1}{2} b = \frac{\text{Tang. } \frac{1}{2} \text{ diff. of } a \text{ and } c \times \sin. \frac{1}{2} \text{ sum of } A \text{ and } c.}{\text{Sin. } \frac{1}{2} \text{ diff. } A \text{ and } c.}$$

But $\frac{1}{2}$ diff. of a and $c = 28^\circ .. 55'$, $\frac{1}{2}$ sum of A and $c = 86^\circ .. 55'$,
and $\frac{1}{2}$ diff. of A and $c = 38^\circ .. 25'$. Consequently it is

(Key to Vol. III. page 110.)

$$\begin{array}{r}
 + \text{Log. tang. } 28^\circ .. 55' = 9.7422609 \\
 + \text{Log. sin. } 86^\circ .. 55' = 9.9993640 \\
 \hline
 \text{Subtract log. sin. } 38^\circ .. 25' = 9.7933543 \\
 \hline
 \text{Log. tang. } 41^\circ .. 36' = 9.9482706 \text{ diff.} \\
 \text{Multiply by } 2 \\
 \hline
 \text{PRODUCT } 83^\circ .. 12' = \text{the side } b.
 \end{array}$$

$$\begin{array}{r}
 \text{Again, As Sin. (a) } 56^\circ .. 40' \text{ Log. ar. co. } - 10.0779768 \\
 : \text{Sin. (A) } 48^\circ .. 30' \text{ - - - Log. } 9.8744561 \\
 :: \text{Sin. (b) } 83^\circ .. 12' \text{ - - - Log. } 9.9969342 \\
 : \text{Sin. (B) } 62^\circ .. 55' \text{ - - - Log. } 9.9493671
 \end{array}$$

Ex. 11. Given $c=114^\circ .. 30'$, $a=56^\circ .. 40'$, and $B=62^\circ .. 54'$.

Now,

$$\text{Tang. } \frac{1}{2} \text{ diff. of A and c} = \frac{\text{Co. tang. } \frac{1}{2} B \times \text{sin. } \frac{1}{2} \text{ diff. of } a \text{ and } c.}{\text{Sin. } \frac{1}{2} \text{ sum of } a \text{ and } c.}$$

And

$$\text{Tang. } \frac{1}{2} \text{ sum of A and c} = \frac{\text{Co. tang. } \frac{1}{2} B \times \text{co. sin. } \frac{1}{2} \text{ diff. of } a \text{ and } c.}{\text{Co. sin. } \frac{1}{2} \text{ sum of } a \text{ and } c.}$$

But $\frac{1}{2} B=31^\circ .. 27'$, $\frac{1}{2}$ diff. of a and $c=28^\circ .. 55'$, and $\frac{1}{2}$ sum of a and $c=85^\circ .. 35'$. Therefore it is

$$\begin{array}{r}
 + \text{Log. co. tang. } 31^\circ .. 27' = 10.2132480 \} \\
 + \text{Log. sin. } 28^\circ .. 55' = 9.6844297 \} \\
 \hline
 19.8976777 \\
 - \text{Log. sin. } 85^\circ .. 35' = - - - - - 9.9987084 \\
 \hline
 \text{Log. tang. } 38^\circ .. 24' = 9.8989693 \text{ diff.}
 \end{array}$$

And,

$$\begin{array}{r}
 + \text{Log. co. tang. } \frac{1}{2} B=31^\circ .. 27' = 10.2132480 \} \\
 + \text{Log. co. sin. } 28^\circ .. 55' = 9.9421688 \} \\
 \hline
 20.1554168 \\
 - \text{Log. co. sin. } 85^\circ .. 35' = - - - - - 8.8865418 \\
 \hline
 \text{Log. tang. } 86^\circ .. 55' = 11.2688750 \text{ diff.}
 \end{array}$$

Consequently $A=48^\circ .. 30'$, and $c=125^\circ .. 20'$.

(Key to Vol. III. page 110.)

Lastly, As Sin. (A) $48^\circ .. 30'$ Log. ar. co. -10.1255439
 : Sin. (a) $56^\circ .. 40'$ - - - - - Log. 9.9219401
 :: Sin. (B) $62^\circ .. 54'$ - - - - - Log. 9.9494938
 : Sin. (b) $83^\circ .. 15'$ - - - - - Log. 9.9969778

Ex. 12. Given $c = 125^\circ .. 20'$, $b = 83^\circ .. 12'$, and $A = 48^\circ .. 30'$.

Ans. $a = 56^\circ .. 40'$
 $c = 114^\circ .. 30'$ } Which were required.
 and $B = 62^\circ .. 54'$

Ex. 13. Given $A = 48^\circ .. 31'$, $B = 62^\circ .. 56'$, and $c = 125^\circ .. 20'$.

$a = 56^\circ .. 40'$
 $b = 83^\circ .. 12'$ } Ans.
 $c = 114^\circ .. 30'$

Ex. 14. Given $A = 50^\circ .. 12'$, $c = 58^\circ .. 8'$, and $a = 62^\circ .. 42'$.

Wherefore it will be,

As Sin. (A) $50^\circ .. 12'$ - - - - - Log. ar. co. -10.1144385
 : Sin. (a) $62^\circ .. 42'$ - - - - - Log. 9.9487147
 :: Sin. (c) $58^\circ .. 8'$ - - - - - Log. 9.9290504
 : Sin. (c) $79^\circ .. 11'$, or its suppl. $100^\circ .. 49'$, Log. 9.9922036

Now, $\text{co. tang. } \frac{1}{2} B = \frac{\text{Tang. } \frac{1}{2} \text{ diff. } A \text{ and } c \times \text{sin. } \frac{1}{2} \text{ sum } a \text{ and } c}{\text{Sin. } \frac{1}{2} \text{ diff. } a \text{ and } c}$

But $\frac{1}{2}$ diff. A and $c = 3^\circ .. 58'$, $\frac{1}{2}$ sum a and $c = 70^\circ .. 56' \frac{1}{2}$,
 and $\frac{1}{2}$ diff. a and $c = 8^\circ .. 15'$.

Consequently

Add $\left\{ \begin{array}{l} \text{log. tang. } 3^\circ .. 58' = 8.8409977 \\ \text{log. sin. } 70^\circ .. 56' \frac{1}{2} = 9.9755175 \end{array} \right.$
18.8165152
 and subtract $\text{log. sin. } 8^\circ .. 15' = 9.1568296$

$\text{Log. co. tang. } 65^\circ .. 26' = 9.6596856$ diff.
 Multiply by 2

PRODUCT = $\angle B = 130^\circ .. 52'$, or, by taking the
 other value of $c (= 100^\circ .. 49')$, $\angle B = 156^\circ .. 17'$; hence, by
 common analogy,

(Key to Vol. III. page 110.)

As Sin. (A) 50° .. 12' Log. ar. co. — 10·1144385
 : Sin. (a) 62° .. 42' - - - - Log. 9·9487147
 :: Sin. (B) 130° .. 52' - - - - Log. 9·8786563
 : Sin. (b) 119° - - - - - Log. 9·9418095,

Or taking B = 156° .. 17', b is 152° .. 17'. Ans.

Ex. 15. Because (*Theor. v. Spherical Trigonometry.*)

As 360° : 1'' :: $\frac{1}{2}$ surface of the earth : area required, it will be, putting $a = 3·14159$ &c. and $d = 7957·75$ the diameter given,

As 1296000'' : 1'' :: $\frac{ad^2}{2}$: area required; that is, $\frac{ad^2}{2 \times 1296000}$
 $= \frac{ad^2}{2592000}$ = area of the triangle measured on the \odot 's surface.

Now, $d = 7957·75$, whereof the log. is 3·9007903
 Multiply by 2

Product 7·8015806 = log. d^2


And, $a = 3·141592$, whereof the log. is ·4971500

Sum 8·2987306 = log. ad^2

Lastly, the log. of 2592000 is - - - 6·4136350

Difference 1·8850956 = log. of the area = 76·75299 square miles and upwards. Ans.

Ex. 16. — *First for the plane angle between the base and each face.* Since by the properties of a hexagon each side of the base = radius, and because by the question, the altitude is double each side of the base, that is, double the radius; it will be




As Sin. 90° - - - Log. ar. co. — 10·0000000
 : 1 (=CB) - - - - - Log. 0·0000000
 :: Sin. 60° ($\angle DBC$) - - - - - Log. 9·9375306
 : CD ·8660254 - - - - - Log. — 1·9375306

(Key to Vol. III. page 110.)

$$\begin{aligned}
 \text{And, As } 2 \text{ (CE) - - - - - Log. ar. co. - } & 1.6989700 \\
 \text{: Rad. (CE) - - - - - Log. } & 10.0000000 \\
 \text{: : } .8660254 \text{ (CD) - - - - - Log. - } & 1.9375306 \\
 & \hline
 \text{: Co. tang. } 66^\circ .. 35' .. 11'' \frac{1}{2} \angle \text{CDE - } & \text{Log. } 9.6365006 \\
 & \hline
 \end{aligned}$$

Next, for the plane angle between each pair of lateral faces. Because CD is double of AD or of DB, it will be,

$$\begin{aligned}
 \text{As } 2 \text{ (CD) - Log. ar. co. - } & 1.6989700 \\
 \text{: } 1 \text{ (AD) - - - - - Log. } & 0.0000000 \\
 \text{: : Rad. (CD) - - - - - Log. } & 10.0000000 \\
 & \hline
 \text{: Co. tang. } 63^\circ .. 26' .. 5'' \frac{3}{4} \} & 9.6989700 \\
 \angle \text{CAD - - - - - Log. } & \}
 \end{aligned}$$



But if DE and DF be drawn at right angles to AC and CB, the angle FDE is the measure of the inclination of the plane CDB to the plane CDA. Now it is evident that, FDE is bisected by the straight line CD; consequently the angle CDE is half the measure of the angle required. Again the two triangles EDC and DAC are similar, and the angle CAD = \angle CDE. Wherefore the angle CDE = $63^\circ .. 26' .. 5'' \frac{3}{4}$ and the whole angle FDE = $126^\circ .. 52' .. 11'' \frac{1}{2}$.

And since the magnitude of a trilateral solid angle is measured by the excess of the sum of the three angles made by its bounding planes above 2 right angles, it follows that, As 360° : $\{ 2(66^\circ .. 35' .. 11'' \frac{1}{2}) + 126^\circ .. 52' .. 11'' \frac{1}{2} - 180^\circ \}$:: 1000 : 222.4143 either solid angle at the base of the pyramid. And for the solid angle at the vertex, since its measure is the excess of the sum of the angles of inclination of the several planes forming it, above double the number of those planes multiplying 90° , when from the product 4 right angles have been taken; it follows that,

$$\text{As } 360^\circ : \{ 6(126^\circ .. 52' .. 11'' \frac{1}{2}) - 720^\circ \} :: 1000 : 114.497685$$

the solid angle at the vertex of the pyramid.

☞ The plane angles agree with Dr. Hutton's answer, but the solid angles disagree.

(Key to Vol. III. page 146.)

GEODESIC OPERATIONS, AND THE FIGURE OF THE EARTH.

Ex. I. The arc of a great circle passing through the two distant objects will cut the horizon, and give two similar right angled spherical triangles for solution. Now the hypotenuses of these will be in the ratio of their perpendiculars, that is, in the ratio of 1" to 25' .. 47" \therefore 1 : 1547. Hence, (the sum of the hypotenuses being 66° .. 30' .. 39",) the hypotenuse of the greater triangle is 66° .. 28' .. 4" $\frac{1}{5}$, and of the less, 2' .. 34" $\frac{4}{5}$. Therefore, [*Case 1. Right angled spherical triangles.*]

$$\begin{array}{r}
 \text{As Cos. } 0^\circ \text{ .. } 25' \text{ .. } 47'' \text{ - - - - - Log. ar. co. - } 10.0000122 \\
 : \text{Cos. } 66^\circ \text{ .. } 28' \text{ .. } 4'' \frac{1}{5} \text{ - - - - - Log. } 9.6012600 \\
 :: \text{Radius or sin. } 90^\circ \text{ - - - - - Log. } 10. \text{ ---} \\
 \\
 : \text{Sin. } 66^\circ \text{ .. } 28' \text{ .. } 1'' \cdot 67 \text{ - - - - - Log. } 9.6012722
 \end{array}$$

$$\begin{array}{r}
 \text{But } \frac{1}{1547} \times (66^\circ \text{ .. } 28' \text{ .. } 1'' \cdot 67) = 0^\circ \text{ .. } 2' \text{ .. } 34'' \cdot 7 \\
 \text{Add } 66^\circ \text{ .. } 28' \text{ .. } 1'' \cdot 67
 \end{array}$$

$$\begin{array}{r}
 \text{Sum and Answer* } \quad \underline{\underline{66^\circ \text{ .. } 30' \text{ .. } 36'' \text{ } 37}}
 \end{array}$$

* Which we submit to be more than half a second nearer the true horizontal angle, than is the Answer given by Dr. Hutton.

Otherwise,†

First taking the trouble to correct the formula which in its present state is imperfect, it becomes

$$\text{Log. sin. } \frac{1}{2} c = \frac{1}{2} \left\{ 20 + \log. \sin. \frac{1}{2} (c + h - h') + \log. \sin. \frac{1}{2} (c + h' - h) - \log. \cos. h' - \log. \cos. h. \right\}$$

$$\begin{array}{r}
 \text{Then } c = 66^\circ \text{ .. } 30' \text{ .. } 39'' \\
 \quad h = \quad \quad 25' \text{ .. } 47'' \\
 \quad h' = \quad \quad - 1''
 \end{array}$$

$$\begin{array}{l}
 \text{Wherefore } \frac{1}{2} (c + h - h') = 33^\circ \text{ .. } 28' \text{ .. } 13'' \frac{3}{5}, \text{ and} \\
 \frac{1}{2} (c + h' - h) = 33^\circ \text{ .. } 2' \text{ .. } 25'' \frac{1}{5}.
 \end{array}$$

† The reader, if he desires it, will find the figure, and other particulars, in *Woodhouse's Trigonometry*, 2nd edition.

(Key to Vol. III. page 146.)

Consequently it is

Radius square* - - - - -	Log.	20	-----
Sin. 33° .. 28' .. 13'' ¹ / ₂ - - - - -	Log.	9.7415504	
Sin. 33° .. 2' .. 25'' ¹ / ₂ - - - - -	Log.	9.7365801	
Co. Sin. - - - - - 1" - - - - -	Log. ar. co.	-10	-----
Co. Sin. - - - 25' .. 47" - - - - -	Log. ar. co.	-10.0000122	-----
- - - - - Divide by 2 - - - - -		19.4781427	log. sum.

Sin. 33° .. 15' .. 18" · 19 - - - - -	Log.	9.7390713	half sum.
Multiply by 2			

PRODUCT 66° .. 30' .. 36" · 38 ∠ required, agreeing to the ¹/₁₀₀ part of a second with the angle *before found*.

* If the log. sines to Radius 1 be used, the operation will be—

Log. sin. 33° .. 28' .. 13'' ¹ / ₂ is	-1.7415504
Log. sin. 33° .. 2' .. 25'' ¹ / ₂ is	-1.7365801
Log. ar. co. cos. -1" + Log. ar. co. cos. 25' .. 47"	0.0000122

Divide by 2	-1.4781427

Sin. 33° .. 15' .. 18" · 19	Log. -1.7390713

NOTE.—This example is a partial statement of the angular distance, altitude, and depression of *Fiennes* and *Watten*, as observed at CALAIS in the Trigonometrical Survey of 1787, made for the purpose of connecting the observations of Greenwich and Paris: and here we cannot help expressing our unqualified opinion that the answer given with the question was a mere guess.

(Page 147.)

Ex. 2. Because the two measured distances D and d are equal, putting x for the height of the tower, and taking the formula given in Cor. to Prob. 4, page 129, vol. iii. it will be, $x =$

$$\sqrt{\frac{2d^2}{\cot^2 A + \cot^2 C - 2 \cot^2 B}} = \sqrt{\frac{d}{\frac{1}{2} \cot^2 A + \frac{1}{2} \cot^2 C - \cot^2 B}}$$

$$= \sqrt{\frac{84}{\frac{1}{2} \cot^2 36^\circ .. 50' + \frac{1}{2} \cot^2 14^\circ - \cot^2 21^\circ .. 24'}}$$

Therefore by the rule page 129 as above,

$$2x$$

(Key to Vol. III. page 147.)

$$\begin{array}{r} 2 \log. \cot. 36^\circ .. 50' = 0.2510324 \text{ of which the nat. num. is } 1.78251 \\ 2 \log. \cot. 14^\circ .. 0' = 1.2064578 \text{ - - - - - } 16.08635 \end{array}$$

$$\text{Sum of the nat. numbers } \underline{17.86886}$$

$$\begin{array}{r} \text{Half sum } 8.93443 \\ 2 \log. \cot. 21^\circ .. 24' = 0.8136590 \text{ of which the nat. num. is } \underline{6.51117} \end{array}$$

$$\text{And } \underline{.3844000} = \log. \text{ of the remainder } 2.42326$$

$$\begin{array}{l} \text{Subtract } .1922000 = \text{half log. of the remainder.} \\ \text{From } 1.9242793 \text{ log. of 84 feet} = \underline{D.} \end{array}$$

$$\log. \text{ of } x = 1.7320793 \text{ log. difference, of which the natural number is } 53.9609 \text{ feet. Ans.}$$

INVESTIGATION OF GENERAL ROY'S RULE.

Ex. 3. Because in any spherical triangle the excess of its three angles above two right angles, is the measure of the area of the triangle, (*Cor. 1. Theor. v. Spher. Trig. vol. iii. p. 84*), and because the three angles of a spherical triangle— 180° , are to 180° , as the surface or area of the triangle, to the area of a great circle of the sphere*; if A, B, C, be severally put for the three angles, a for the area of the triangle, and q for the area of a great circle of the sphere, ($=\frac{1}{4}$ the surface of the sphere,) it will be, $\frac{a \times 180^\circ}{q} = A + B + C - 180^\circ$; therefore,

since the square of the radius multiplying 3.1415926535897932384626433832795028841971693993751058209749445923078174062962089986280348253421170679821480865132723066470938446319 &c. = the area of a great circle of the sphere $=\frac{1}{4}$ the superficies of the sphere, it follows that,

$$\frac{a \times 180^\circ}{\text{rad.}^2 \times 3.141592 \text{ \&c.}} = A + B + C - 180^\circ. \text{ But the length}$$

of the earth's radius in feet, (*the earth considered a perfect sphere, and the length of a degree on its surface taken at*

$$365154.6 \text{ feet,}) = \frac{180^\circ \times 365154.6}{3.1415926} \text{ feet; whence } A + B + C$$

* Theorem v. page 83, vol. iii.

(Key to Vol. III. page 147.)

$$-180^\circ = \frac{a \times 180^\circ}{365154 \cdot 6^2 \times \frac{180^\circ}{3 \cdot 1415}} \times 3 \cdot 14159 = \frac{3 \cdot 14159 \times a}{180^\circ \times 365154 \cdot 6^2}$$

feet = the excess of $A + B + C$ above two right angles. This

excess expressed in seconds is $\frac{3 \cdot 14159 \times 60 \times 60 \times a}{180^\circ \times 365154 \cdot 6^2} =$

$$\frac{62 \cdot 831853 \times a}{365154 \cdot 6^2}.$$

Now in logarithms the excess will be, $\log.$

$$a + \log. 62 \cdot 831853 - 2 \log. 365154 \cdot 6 = \log. a + 1 \cdot 7981799 - 11 \cdot 1249536 = \log. a - 9 \cdot 3267737. \text{ Which was required.}$$

Ex. 4. Half the sum of the sides being 22.5, and the three remainders (*see page 29, vol. ii.*) being 12.5, 5.5, and 4.5, half the sum of the logarithms of these 4 numbers is the $\log.$ of the area of the triangle.

But the Log. of 22.5.	is	1.3521825
- - - - - 12.5.	is	1.0969100
- - - - - 5.5.	is	0.7403627
- - - - - 4.5.	is	0.6532125

$$\text{Log. sum} = 3 \cdot 8426677$$

Log. half sum = 1.9213338 which is the $\log.$ of 83.432233 square miles.

Now 2580^2 feet = 1 square mile,
and $2 \log.$ of 5280 feet = 7.4452678

$$\text{Log. sum } 9 \cdot 3666016 = \text{the } \log. \text{ of the area of the triangle in feet.}$$

$$\text{Subtract the const. log. } 9 \cdot 3267737$$

$$\left. \begin{array}{l} \text{The } \log. \text{ of the excess of the} \\ \text{3 angles above } 180^\circ \text{ is} \end{array} \right\} \cdot 0398279 = \text{L. of } 1'' \cdot 096.$$

Ex. 5. $\left. \begin{array}{l} \text{mils. mils.} \\ 15 \times 12 = 180 \text{ sq. mils. of} \\ \text{which the } \log. \text{ is} \end{array} \right\} \text{---} 2 \cdot 2552725$

And $\left. \begin{array}{l} 1760 \text{ yds.} \times 3^2 = 5280^2 \\ \text{feet of which the } \log. \text{ is} \end{array} \right\} \text{---} 7 \cdot 4452678$

$$\text{Log. sum } 9 \cdot 7005403 \text{ the } \log. \text{ of the area of the triangle in feet.}$$

$$\text{Subtract the const. log. } 9 \cdot 3267737$$

$$\text{Difference } \cdot 3737666 \text{ log. of } 2'' \cdot 3646$$

Wherefore $2'' \dots 21''' \dots 52'''' \frac{1}{2}$. Ans.

(Key to Vol. III. page 147.)

Otherwise.

24 miles in ft. = 126720, of which the log. is 5·1028452

And 15 miles in ft. = 79200, of which the log. is 4·8987252

<i>Sum of the logs.</i>	10·0015704
Deduct (see Art. 4, page 135, vol. iii.)	9·6278037

The com. log. of the spherical excess in seconds is 0·3737667 the
log. of $2'' \dots 21'' \dots 52'' \frac{1}{2}$ exactly as before.

Ex. 6.

As the sum of the sides (41) - - - - - Log. ar. co. - 2·3872161

: difference of the sides (5) - - - - - Log. 0·6989700

∴ Cot. $29^\circ \dots 12' \dots 18''$, ($\frac{1}{2}$ cont^d \angle) - - - - - Log. 10·2527696: Tang. $12^\circ \dots 18' \dots 42'' \frac{1}{2}$ (half diff. of the other 2 \angle 's) 9·3389557And the complement of half the included \angle is $60^\circ \dots 47' \dots 42''$.

Hence the three angles of the plane triangle are $58^\circ \dots 24' \dots 36''$,
 $93^\circ \dots 6' \dots 24'' \frac{1}{2}$, and $48^\circ \dots 28' \dots 59'' \frac{1}{2}$. And the third side =
26·172355 miles.

Now half the sum of the sides = 33·586 miles; and the three
remainders are 15·586, 10·586, and 7·403 (page 29, vol. ii.)

But the

Log. of 33·586 is 1·5261583

- - - - 15·586 is 1·1927347

- - - - 10·586 is 1·0247319

- - - - 7·403 is 0·8694077

<i>Log. sum</i>	4·6130326
-----------------	-----------

Half sum 2·3065163 log. of the area in miles.

And twice log. of 5280 is 7·4452678

<i>Log. sum</i>	9·7517841	log. of the area in feet.
Subtract the const. log.	9·3267737	

<i>Difference</i>	0·4250104	the log. of $2'' \cdot 6607$.*
-------------------	-----------	---------------------------------

* The same result will obtain, if the two given sides be converted into minutes and seconds, and the triangle solved spherically. (See Theor. 5, ch. iv.)

(Key to Vol. III. page 147.)

Ex. 7. As rad. $\overset{\text{ft.}}{+38}$: rad. :: given base : base required.

But the radius of the earth, at a mean calculation, is more than 21 millions of feet, therefore, as in *Prob. ix. page 136, vol. iii.* since the required base (*by multiplying means and extremes*)

$$= \frac{\text{rad.} \times \text{given base}}{\text{rad.} + 38 \text{ ft.}}, \text{ or, (by further comparison, and}$$

the division of the given base $\times 38 \text{ ft.}$, *by the rad. + 38 ft., ex-*

pressed symbolically,) $B - b = B \times \left(\frac{h}{r} - \frac{h^2}{r^2} + \frac{h^3}{r^3} - \&c. \right)$ Now

neglecting the second and all the succeeding terms of the

series as insignificant, it will be $B - b = \frac{Bh}{r}$.

By Logarithms.

The log. of 34286 ft. is - - 4.5351168

And the log. of 38 ft. - - - 1.5797836

Log. sum 6.1149004

Deduct (l. rad. of the earth in ft.) 7.3223947

Remainder - 2.7925057 the log. of .062016267

Consequently $34286 - .062016267 = 34285.937983732 \text{ ft. Ans.}$

Ex. 8. If the altitude be corrected by subtracting 44'' the refraction in altitude, and then by adding 6'' for the parallax, the true altitude of the sun is $52^\circ .. 34' .. 22''$, the co. altitude $37^\circ .. 25' .. 38''$, the co. lat. $41^\circ .. 9'$, the co. declination $71^\circ .. 30'$, and the proportional hour angle $27^\circ .. 8' .. 30''$. Wherefore,

As Sin. co. alt. $37^\circ .. 25' .. 38''$ Log. ar. co. - 10.2162728

: Sin. hour $\angle 27^\circ .. 8' .. 30''$ - - - Log. 9.6591478

:: Sin. co. dec. $71^\circ .. 30' .. 00''$ - - - Log. 9.9769566

: Sin. \angle reqd. or of its supplement - Log. 9.8523772 the log.
sin. of $45^\circ .. 23' .. 2\frac{1}{2}''$. Ans.

Ex. 9. The part of the ecliptic between the beginning of *Aries* and the center of the sun being considered as the hypothenuse of a right angled spherical triangle, and the obliquity of the ecliptic as the angle adjacent to the side sought; the Tang. of the sun's right ascension = Tang of his longitude \times^d by the cosine of the obliquity of the ecliptic. But

(Key to Vol. III. page 147.)

Log. tang. $29^{\circ}..13'..43''$ is 9.7478285
 Log. cosine $23^{\circ}..27'..40''$ is 9.9625259

Their sum (abating 10 in the index) is 9.7103544 which is the
 log. tang. of $27^{\circ}..10'..13''..50''\frac{1}{2}$ = the sun's right ascen-
 sion. Q. E. I.

Ex. 10. By Theorem 11 in the Scholium to Prob. xii. alluded
 to in the question, the cosine of the sun's longitude = co. sin.
 of his right ascension \times^d by the cosine of his declination.
 Now the

Log. co. sin. $32^{\circ}..46'..52''\frac{1}{2}$ is 9.9246636
 Log. co. sin. $13^{\circ}..13'..27''$ is 9.9883281

Their sum (abating 10 in the index) is 9.9129917 which is the
 log. cos. of $35^{\circ}..4'..12''..31'''$ = the sun's longitude. Q. E. I.

Ex. 11. The Theorem for the longitude being (*see Theorem iii.*
page 142, vol. iii.) "that its tangent = the sine of the obli-
 quity of the ecliptic \times^d by the tangent of the declination
 \times^d by the secant of the right ascension + the tangent of the
 right ascension \times cosin. of the obliquity of the ecliptic;" it
 will be by logarithms as follows, *to radius 1* :

Log. sin. $23^{\circ}..27'..40''$ is -1.6000212
 Log. tang. $62^{\circ}..50'..0''$ 0.2897176
 Log. sec. $162^{\circ}..50'..34''$ 0.0197698

Log. sum -1.9095086 the com. log. of .8119113

And,

Log. tang. $162^{\circ}..50'..34''$ is -1.4895840
 Log. co. sin. $23^{\circ}..27'..40''$ -1.9625259

Log. sum -1.4521099 the com. log. of .2832108

The sum of the two natural numbers is 1.0951221

Which is the natural tangent of $132^{\circ}..24'..2''$ = longitude of α
Ursæ Majoris.

Again. The Theorem for the latitude (*see Theorem iv. page*
142, vol. iii.) being "that its sine = sin. of the declination \times
 co. sin. of the obliquity of the ecliptic, - sin. of the right
 ascension \times co. sin. of the declination \times sin. of the eliptic's
 inclination to the equator," the solution will be by logarithms
 as follows :

(Key to Vol. III. page 147.)

Log. sin. $62^{\circ}..50'$ is -1.9492349 Log. co. sin. $23^{\circ}..27'..40''$ is -1.9625259

 Log. sum -1.9117608 the com. log. of $.8161328$

And,

Log. sin. $162^{\circ}..50'..34''$ is -1.4698143 Log. co. sin. $62^{\circ}..50'..0''$ is -1.6595173 Log. sin. $23^{\circ}..27'..40''$ is -1.6000212

 Log. sum -2.7293527 the com. log. of $.0536232$

The *difference* of the two natural numbers is $.7625096$ Which is the nat. sin. of $49^{\circ}..41'..9'' = \text{lat. of } \alpha \text{ Ursæ Majoris. Q. E. I.}$

Ex. 12. Putting m for 60833, M for 60494, also l for $49^{\circ}..3'$ and L for $12^{\circ}..32'$, the Theorem (page 114) where d = the *greater*, and c the *less* axis of the earth, is

$$\frac{d}{c} = \sqrt{\frac{(M^{\frac{1}{3}} \sin. L + m^{\frac{1}{3}} \sin. l) \cdot (M^{\frac{1}{3}} \sin. L - m^{\frac{1}{3}} \sin. l)}{(m^{\frac{1}{3}} \text{co. sin. } l + M^{\frac{1}{3}} \text{co. sin. } L) \cdot (m^{\frac{1}{3}} \text{co. sin. } l - M^{\frac{1}{3}} \text{co. sin. } L)}}$$

which may be more simply expressed, As $d : c :: \text{sq. root of the rectangle of the sum and difference of } M^{\frac{1}{3}} \sin. L \text{ and } m^{\frac{1}{3}} \sin. l : \text{sq. root of the rectangle of the sum and difference of } M^{\frac{1}{3}} \text{co. sin. } L \text{ and } m^{\frac{1}{3}} \text{co. sin. } l.$ Now

Log. $M^{\frac{1}{3}}$ is 1.5939041 Log. sin. L is -1.3364794

 Log. sum $.9303835$ com. log. of $8.5189.$

Log. $m^{\frac{1}{3}}$ is 1.5947131 Log. sin. l is -1.8781090

 Log. sum 1.4728221 com. log. of $29.7046.$

The *sum* of these two natural numbers is 38.2235 Their *difference* is 21.1857 The log. of 38.2235 is 1.5823300 The log. of 21.1857 is 1.3260285

 Log. sum 2.9083585

Half sum 1.4541792 com. log. of $28.456.$

(Key to Vol. III. page 147.)

Again,

$$\text{Log. } m^{\frac{1}{3}} \quad \text{is} \quad 1.5947131$$

$$\text{Log. co. sin. } l. \quad \text{is} \quad -1.8165066$$

$$\text{Log. sum} \quad 1.4112197 \quad \text{c. log. of } 25.776.$$

$$\text{Log. } M^{\frac{1}{3}} \quad \text{is} \quad 1.5939041$$

$$\text{Log. co. sin. } L. \quad -1.9895254$$

$$\text{Log. sum} \quad 1.5834295 \quad \text{c. log. of } 38.320.$$

The *sum* of these two natural numbers is 64.096Their *difference* is 12.544

The log. of 64.096 is 1.8068309

The log. of 12.544 is 1.0984360

The half sum of these logs. is 1.4526334 com. log. of 28.355

Therefore, As $d : c :: 28456 : 28355 ::$ ratio of the axes. Q. E. D.

Ex. 13. Let R be a degree of the earth's equator, M and m respectively the first and last degrees of latitude; then R , M , m , are severally to each other as the radii of their osculatory circles; in the same ratio are AC , $\frac{CD^2}{AC}$, and $\frac{AC^2}{CD}$. See the fig. to Prob. xiii. page 143, vol. iii.

Hence, As $R^2 : m^2 :: AC^2 : \frac{AC^4}{CD^2}$. And,

As $R : M :: AC : \frac{CD^2}{AC}$. By composition, therefore, $R^3 : Mm^2 :: 1 : 1$. Wherefore

$R = \sqrt[3]{Mm^2}$, that is, R is the first of two mean proportionals between M and m . Which was to be proved.

Ex. 14. The length of a degree of the meridian in any latitude being to the length of a degree of the meridian in any other latitude, in the direct ratio of the radii of the respective osculatory circles; if R be put for the radius of curvature at any point of the terrestrial meridian, c , d , and e being as in Proposition xiii. Trigonometrical Surveying, vol. iii. of the COURSE, it will be

(Key to Vol. III. page 147.)

$$R = \frac{d^2}{c(d^2 - e^2 \sin^2 L)^{\frac{3}{2}}} \therefore (1 - \frac{e^2}{d^2} \times \sin^2 L)^{-\frac{3}{2}} \therefore 1 + \frac{\frac{3}{2}e^2}{d^2} \times \sin^2 L + \frac{\frac{15}{4}e^4}{d^4} \times \sin^4 L + \&c. \therefore 1 + \frac{\frac{3}{2}e^2}{d^2} \times \sin^2 L \text{ very nearly.}$$

Therefore the increase of a degree of the terrestrial meridian in receding from the equator, is in the ratio of $\frac{3e^2}{2d^2} \times \sin^2 L$ nearly, that is, very nearly in the duplicate ratio of the sine of the latitude. Q. E. D.

(Page 148.)

Ex. 15.* Because p and m are given, the radii of curvature to p and m are determinable. Let these be r and $r + n$; the radius of curvature to d will be $\frac{r(r+n)}{(r+n) - n \sin^2 a}$. Now to compare the degrees of the three several arcs whereof the radii of curvature are r , $r + n$, and $\frac{r(r+n)}{(r+n) - n \sin^2 a}$, substitute the given lengths d , m , and $p (=m + v)$ for the radii, (since the degrees are in the direct ratio of the radii of curvature,) and it will be, $d = \frac{m(m+v)}{(m+v) - v \sin^2 a} = \frac{m \times p}{p - v \sin^2 a} = \frac{pm}{p + (m-p) \sin^2 a}$. Q. E. D.

* At page 45, No. 98 of the Ladies' Diary, an example proposed in 1800 by Capt. (now Col.) Mudge, and applicable to this Theorem, may be seen solved, and the Theorem investigated at more length.

POLYGONOMETRY.

(Page 160.)

Ex. 6. Because twice the area of the hexagon = $\left\{ \begin{array}{l} AB. BC. \sin. B. \\ + AB. DC. \sin. (B + C) + BC. DC. \sin. C + DE. EF. \sin. E + DE. \end{array} \right.$

(Key to Vol. III. page 160.)

$$\left. \begin{aligned} &AF. \sin. (E+F) + EP. AF. \sin. F. \} = \{ 1284 \times 1782 \times \sin. 32^\circ + \\ &1284 \times 2400 \times \sin. 80^\circ + 1782 \times 2400 \times \sin. 48^\circ + 2700 \times 2860 \\ &\times \sin. 66^\circ + 2700 \times 4621.5 \times \sin. 25^\circ .. 28' .. 22'' + 2860 \times \\ &4621.5 \times \sin. 88^\circ .. 31' .. 38''. \} \text{ it will be by logarithms,} \end{aligned}$$

$$\begin{array}{rcl} 1284 & \log. & 3.1085650 \\ 1782 & \log. & 3.2509077 \\ 32^\circ & \log. \sin. & -1.7242097 \end{array}$$

$$\text{Log. sum} \quad \underline{\underline{6.0836824}} \text{ the log. of } 1212501.95$$

$$\begin{array}{rcl} 1284 & \log. & 3.1085650 \\ 2400 & \log. & 3.3802112 \\ 80^\circ & \log. \sin. & -1.9933515 \end{array}$$

$$\text{Log. sum} \quad \underline{\underline{6.4821277}} \text{ the log. of } 3034882.63$$

$$\begin{array}{rcl} 1782 & \log. & 3.2509077 \\ 2400 & \log. & 3.3802112 \\ 48^\circ & \log. \sin. & -1.8710735 \end{array}$$

$$\text{Log. sum} \quad \underline{\underline{6.5021924}} \text{ the log. of } 3178281.75$$

$$\begin{array}{rcl} 2700 & \log. & 3.4313638 \\ 2860 & \log. & 3.4563660 \\ 66^\circ & \log. \sin. & -1.9607302 \end{array}$$

$$\text{Log. sum} \quad \underline{\underline{6.8484600}} \text{ the log. of } 7054400.$$


$$\begin{array}{rcl} 2700 & \log. & 3.4313638 \\ 4621.5 & \log. & 3.6647830 \\ 25^\circ .. 28' .. 22'' & \log. \sin. & -1.6335514 \end{array}$$

$$\text{Log. sum} \quad \underline{\underline{6.7296982}} \text{ the log. of } 5366587.65$$

(Key to Vol. III. page 160.)

2860	log.	3.4563660	
4621.5	log.	3.6647830	
88° .. 31' .. 38"	log. sin.	-1.9998565	
		Log. sum 7.1210055	the log. of 13213124.61
		Sum of the nat. numbers	33059778.59
		Half sum and AREA required	16529889.29

Which problem solved by the formula, for the surface or area? in page 155, third vol. of the Course, will be as follows :



Let ABCDEF be the hexagon given in Ex. 1. Then by plane trigonometry,

- $\angle EDF = 34^\circ \dots 4' \dots 14''$ $BF = 5136.3$
- $\angle FAE = 31^\circ \dots 20' \dots 23''$ $CE = 4585.7$
- $\angle ABF = 59^\circ \dots 36' \dots 24''$
- $\angle ECD = 27^\circ \dots 38' \dots 36''$
- $\angle CED = 24^\circ \dots 21' \dots 24''$
- $\angle AFB = 13^\circ \dots 51' \dots 58''$
- $\angle BCF = 72^\circ \dots 18' \dots 20''$
- $\angle BFC = 19^\circ \dots 18' \dots 4''$
- $\angle CFD = 26^\circ \dots 22' \dots 34''$
- $\angle FCE = 32^\circ \dots 3' \dots 4''$
- $\angle EBC = 56^\circ \dots 41' \dots 31''$
- $\angle BEC = 18^\circ \dots 57' \dots 5''$
- $\angle AFE = 91^\circ \dots 28' \dots 22'' = a'$
- $\angle BFE = 77^\circ \dots 36' \dots 24'' = b'$
- $\angle CFE = 58^\circ \dots 18' \dots 20'' = c'$
- $\angle DFE = 31^\circ \dots 55' \dots 46'' = d'$
- $\angle AEF = 57^\circ \dots 11' \dots 15'' = a''$
- $\angle BEF = 70^\circ \dots 41' \dots 31'' = b''$
- $\angle CEF = 89^\circ \dots 38' \dots 36'' = c''$
- $\angle DEF = 114^\circ \dots \text{---} = d''$

(Key to Vol. III. page 160.)

Wherefore, by Prob. 1, Polygonometry, the area ABCDEF =
 $\frac{1}{2}FE^2 \times \left\{ \frac{\text{Sin. } a'' \cdot \text{Sin. } b'' \cdot \text{Sin. } (a' - b')}{\text{Sin. } (a' + a'') \cdot \text{Sin. } (b' + b'')} + \right.$
 $\frac{\text{Sin. } b'' \cdot \text{Sin. } c'' \cdot \text{Sin. } (b' - c')}{\text{Sin. } (b' + b'') \cdot \text{Sin. } (c' + c'')} + \frac{\text{Sin. } c'' \cdot \text{Sin. } d'' \cdot \text{Sin. } (c' - d')}{\text{Sin. } (c' + c'') \cdot \text{Sin. } (d' + d'')} +$
 $\left. \frac{\text{Sin. } d' \cdot \text{Sin. } d''}{\text{Sin. } (d' + d'')} \right\}$ which admits of the following solution
 by logarithms.

$a'' = 57^\circ .. 11' .. 15''$	log. sin.	9.9245111
$b'' = 70^\circ .. 41' .. 31''$	log. sin.	9.9748590
$(a' - b') = 13^\circ .. 51' .. 58''$	log. sin.	9.3795850
$(a' + a'') = 148^\circ .. 39' .. 37''$	log. sin.	-10.2839042 ar. co.
$(b' + b'') = 148^\circ .. 17' .. 55''$	log. sin.	-10.2794337 ar. co.

Log. sum rejecting 10 in the index — 1.8422930 which is the
 logarithm of .69549.

$b'' = 70^\circ .. 41' .. 31''$	log. sin.	9.9748590
$c'' = 89^\circ .. 38' .. 36''$	log. sin.	9.9999916
$(b' - c') = 19^\circ .. 18' .. 4''$	log. sin.	9.5192144
$(b' + b'') = 148^\circ .. 17' .. 55''$	log. sin.	-10.2794337 ar. co.
$(c' + c'') = 147^\circ .. 56' .. 56''$	log. sin.	-10.2951710 ar. co.

Log. sum rejecting 10 in the index — 0.0486697 which is the
 logarithm of 1.1191.

$c'' = 89^\circ .. 38' .. 36''$	log. sin.	9.9999916
$d'' = 114^\circ$	log. sin.	9.9607302
$(c' - d') = 26^\circ .. 22' .. 34''$	log. sin.	9.6476388
$(c' + c'') = 147^\circ .. 56' .. 56''$	log. sin.	-10.2951710 ar. co.
$(d' + d'') = 145^\circ .. 55' .. 46''$	log. sin.	-10.2516465 ar. co.

Log. sum rejecting 10 in the index — 0.1351781 which is the
 logarithm of 1.3651.

$d' = 31^\circ .. 55' .. 46''$	log. sin.	9.7233527
$d'' = 114^\circ$	log. sin.	9.9607302
$(d' + d'') = 145^\circ .. 55' .. 46''$	log. sin.	-10.2516465 ar. co.

Log. sum rejecting 10 in the index — 1.9357294 which is the
 logarithm of .86244.

(Key to Vol. III. page 160.)

The sum of these 4 natural numbers is 4·04213, of

which the log. is 0·6066103
 $\frac{1}{2}FE^2 = 4089800$; its log. is 6·6117021

Log. sum 7·2183124 which is the
 logarithm of 16531505·7, the AREA REQUIRED.

☞ We could have solved this question in various other ways, and with much less trouble, but we thought it proper to find the area by *Lhuillier's* theorem as well as by reducing the expression for the surface given in Problem I. Although the results do not exactly agree, (which indeed may be owing to some trifling error which we are not at present aware of,) yet the difference is not comparatively great.

$$\begin{aligned} \text{Ex. 7. } \text{Tang. } \angle D &= \frac{BC \cdot \sin. C + AB \cdot \sin. (B + C)}{DC + CB \cdot \cos. C + AB \cdot \cos. (B + C)} = \\ &= \frac{30 \times \sin. 82^\circ .. 37' + 24 \times \sin. 170^\circ .. 19'}{34 + 30 \times \cos. 82^\circ .. 37' + 24 \times \cos. 170^\circ .. 19'} = \\ &= \frac{30 \times \sin. 82^\circ .. 37' + 24 \times \sin. 9^\circ .. 41'}{34 + 30 \times \cos. 82^\circ .. 37' - 24 \times \cos. 9^\circ .. 41'}. \end{aligned} \quad \text{Now the}$$

Log. of 30 is 1·4771213
 Log. sin. of 82° .. 37' is -1·9963841

Sum 1·4735054 log. of 29·751

Log. of 24 is 1·3802112
 Log. sin. of 9° .. 41' is -1·2258328

Sum 0·6060440 log. of 4·0369

33·7879 Numerator.

Log. of 30 is 1·4771213
 Log. cos. of 82° .. 37' is -1·1089272

Sum 0·5860485 log. of 3·8552

Log. of 24 is 1·3802112
 Log. cos. of 9° .. 41' is -1·9937679

Sum 1·3739791 log. of 23·658

(Key to Vol. III. page 160.)

But $34 + 3.8552 - 23.658 = 14.1972$ Denominator.

Again, the log. of 33.7879 is 1.5287611

log. of 14.1972 is 1.1512027

Log. difference 0.3775584Add 10 to the index 10

Sum 10.3775584 the log. tan-

gent of $67^\circ .. 14' = \angle ADC$. Wherefore [*Eucl. B. I. Prop.*
xxxii. *Cor. i.*] $\angle DAB = 103^\circ .. 5'$.Now the side $AD = AB. \cos. 103^\circ .. 5' + BC. \cos. 9^\circ .. 41' + CD. \cos.$
 $67^\circ .. 14' = -24 \times \cos. 76^\circ .. 55' + 30 \times \cos. 9^\circ .. 41' + 34 \times \cos.$
 $67^\circ .. 14'$.

In Logarithms.

The log. of -24 is 1.3802112

Log. cos. $76^\circ .. 55'$ is -1.3548150

Log. sum .7350262 the log. of -5.4328

The log. of 30 is 1.4771213

Log. cos. $9^\circ .. 41'$ is -1.9937679

Log. sum .4708892 the log. of 29.572

The log. of 34 is 1.5314789

Log. cos. $67^\circ .. 14'$ is -1.5876876

Log. sum 1.1191665 the log. of 13.157

Sum = AD = 37.2962

Now conceiving a line to be drawn from A to C, dividing the
trapezium into two triangles, the double area will be $AB. BC. \sin. B. + CD. DA. \sin. D. = 24 \times 30 \times \sin. 87^\circ .. 42' + 34$
 $\times 37.2962 \times \sin. 112^\circ .. 46'$.

In Logarithms.

The log. of 24 is 1.3802112

log. of 30 is 1.4771213

Log. sin. $87^\circ .. 42'$ is -1.9996500

Log. sum 2.8569825 the log. of 719.42

(Key to Vol. III. page 160.)

The log. of 34 is 1.5314789
 log. of 37.2962 is 1.5716646
 Log. sin. $112^\circ .. 46'$ is -1.9647726

Log. sum 3.0679161 the log. of 1169.27

Sum of the nat. numbers 1888.69

Half sum and AREA 944.34

Otherwise without a diagonal.

The double area is $AB. BC. \sin. B. + AB. DC. \sin. (B+C) + BC. DC. \sin. C = 24 \times 30 \times \sin. 87^\circ .. 42' + 24 \times 34 \times \sin. 170^\circ .. 19' + 30 \times 34 \times \sin. 82^\circ .. 37'$.

In Logarithms.

The log. of 24 is 1.3802112
 log. of 30 is 1.4771213
 Log. sin. $87^\circ .. 42'$ is -1.9996500

Log. sum 2.8569825 log. of 719.42

The log. of 24 is 1.3802112
 log. of 34 is 1.5314789
 Log. sin. $170^\circ .. 19'$ is -1.2258328

Log. sum 2.1375229 log. of 137.253

The log. of 30 is 1.4771213
 log. of 34 is 1.5314789
 Log. sin. $82^\circ .. 37'$ is -1.9963841

Log. sum 3.0049843 log. of 1011.54

Sum of the nat. numbers 1868.213

Half sum and AREA 934.106

Or, by drawing a diagonal from B to D.

Twice the area will be $BC. CD. \sin. C + DA. AB. \sin. A = (30 \times 34) \times \sin. 82^\circ .. 37' + (37.2962 \times 24) \times \sin. 76^\circ .. 55'$.

(Key to Vol. III. page 160.)

In Logarithms.

The log. of 30 is 1.4771213

log. of 34 is 1.5314789

Log. sin. $82^\circ .. 37'$ is -1.9963841

 Log. sum 3.0049843 log. of 1011.5

The log. of 37.2962 is 1.5716646

log. of 24 is 1.3802112


Log. sin. $76^\circ .. 55'$ is -1.9885776

 Log. sum 2.9404534 log. of 871.87

 Sum of the nat. numbers 1883.37

 Half sum and AREA 941.68*

* The result by these methods is by no means coincident, and proves the necessity of ascertaining the sides and angles more minutely accurate than in common cases is usual.

Ex. 8. Here $a' = 89^\circ .. 14'$ $b' = 68^\circ .. 11'$ $c' = 36^\circ .. 24'$ $d' = 19^\circ .. 57'$ $a'' = 25^\circ .. 18'$ $b'' = 69^\circ .. 24'$ $c'' = 94^\circ .. 6'$ $d'' = 121^\circ .. 18'$ $(a' - b') = 21^\circ .. 3'$ $(a' + a'') = 114^\circ .. 32'$ $(b' + b'') = 137^\circ .. 35'$ $(b' - c') = 31^\circ .. 47'$ $(c' + c'') = 130^\circ .. 30'$ $(c' - d') = 16^\circ .. 27'$ $(d' + d'') = 141^\circ .. 15'$ 

Wherefore it is,

+ log. sin. $25^\circ .. 18'$ - - - - - 1.6307917+ log. sin. $69^\circ .. 24'$ - - - - - 1.9713035+ log. sin. $21^\circ .. 3'$ - - - - - 1.5549868- log. sin. $114^\circ .. 32'$ ar. co. 0.0410923- log. sin. $137^\circ .. 35'$ ar. co. 0.1710070

 Log. sum -1.3691813 log. of .23398

(Key to Vol. III. page 160.)

+	log. sin.	69° .. 24'	- - - -	- 1.9713035
+	log. sin.	94° .. 6'	- - - -	- 1.9988871
+	log. sin.	31° .. 47'	- - - -	- 1.7215704
-	log. sin.	137° .. 35'	ar. co.	0.1710070
-	log. sin.	130° .. 30'	ar. co.	0.1189545

Log. sum - 1.9817225 log. of .95879

+	log. sin.	94° .. 6'	- - - -	- 1.9988871
+	log. sin.	121° .. 18'	- - - -	- 1.9316911
+	log. sin.	16° .. 27'	- - - -	- 1.4520603
-	log. sin.	130° .. 30'	ar. co.	0.1189545
-	log. sin.	141° .. 15'	ar. co.	0.2034788

Log. sum - 1.7050718 log. of .50707

+	log. sin.	19° .. 57'	- - - -	- 1.5330090
+	log. sin.	121° .. 18'	- - - -	- 1.9316911
-	log. sin.	141° .. 15'	ar. co.	0.2034788

Log. sum - 1.6681789 log. of .46578

Sum of the nat. numbers 2.16562

Now $\frac{1}{2} pQ^2 = 3220722$, its log. 6.5079533
 The logarithm of 2.16562 is 0.3355822

Log. sum 6.8435355 which is the logarithm of 6974859.677 the area in square links, which divided by 100000, the number of square links in an acre, the quotient is 69.74859677 acres. Ans.

DIVISION OF SURFACES.

(Page 172.)

Ex. 1. $\frac{20+18+16}{2} = 27$, and

(Key to Vol. III. page 172.)

$$27 \left\{ \begin{array}{l} -20 \\ -18 \\ -16 \end{array} \right\} = \left\{ \begin{array}{l} 7 \\ 9 \\ 11 \end{array} \right\} \text{ the three remainders.}$$

Also $\sqrt{27 \times 7 \times 9 \times 11} = 136.8$ *nearly*. Therefore the proposed triangular field contains 13.68 acres *nearly*.

Consequently 9.68 ac. and 4 ac. are the two intended divisions.

Let the side 20, be AB; 18, BC, and 16, AC. Likewise let the 4 acres be contiguous to AC; and let D be the termination of the fence in the side AB. Then

Acres	Acres	Chains	Chains	}	the two distances of the dividing line.
As 13.68	: 4	:: 20	: 5.848 = AD		
				Hence	14.152 = BD

Again,

The angle A = $58^\circ .. 45'$ because $\frac{400 + 256 - 324}{2 \times 16 \times 20} = .51875$
 = nat. co. sin. $58^\circ .. 45'$. And

As 21.848 (AC + AD)	- - - - -	Log. ar. co. -	2.6605883
: 11.848 (AC - AD)	- - - - -	Log.	1.0736450
:: Cot. $29^\circ .. 22' \frac{1}{2}$ (half the contained \angle^{les})	- -	Log.	10.2495716

: Tang. $43^\circ .. 56'$ (half the diff. of the \angle^{les} ADC, ACD)	Log.	9.9838049
--	------	-----------

Whence the angle ADC = $104^\circ .. 33' \frac{1}{2}$. But

As Sin. $104^\circ .. 33' \frac{1}{2}$	- - -	Log. ar. co. -	10.0141730
: Sin. $58^\circ .. 45'$	- - - - -	Log.	9.9319213
:: 16 chains	- - - - -	Log.	1.2041200

: 14.132 chains = CD	- - - - -	Log.	1.1502143
----------------------	-----------	------	-----------

Therefore the length of the dividing line is 14.132 chains, and its distance from the angle which is neither most nor least acute, 5.848 chains.

Ex. 2. $\frac{5 + 12 + 13}{2} = 15$. And

$$15 - \left\{ \begin{array}{l} 5 = 10 \\ 12 = 3 \\ 13 = 2 \end{array} \right\} \text{ the three remainders of the half sum and sides.}$$

Also $\sqrt{15 \times 10 \times 3 \times 2} = 30$ the area of the triangle.

Therefore $m = 20$, and $n = 10$ adjacent to the longest side, AB.

Now let the side 12 be BC, and F the point in it to be de-

(Key to Vol. III. page 172.)

terminated. $CF = CB \sqrt{\frac{m}{m+n}} = 12 \sqrt{\frac{20}{30}} = 12 \sqrt{\frac{2}{3}} = 4\sqrt{6} = 9.7979588.$


Again. Let the other extremity of the dividing line be g , and it will be, As $CF : FG :: CB : AB.$ Hence

$$\frac{9.7979588 \times 13}{12} = 10.61445536 = FG.$$

Wherefore the length of the dividing line is 10.61445536 , and the distance of its extremity in BC from the longest side is 2.2020412 ; and of its other extremity $\frac{5}{12}$ (2.2020412), or $.91751716$, which were to be determined.*

* This and the last example admit, each, of two answers.

Ex. 3. Construct a triangle having its three sides respectively equal to the three given lines, namely, $AB=24, AC=25,$ and $BC=7.$ Bisect AB in $D,$ and join $CD,$ the triangle ABC is bisected by CD [*Eucl. i. 38.*].



Also $\sqrt{AD \times AC} = \sqrt{12 \times 25} = \sqrt{300} = 17.3205081 =$ the distance of the required line (to be taken in AC and AB) from the point A the angular point of the acutest angle of the given triangle. [*Eucl. i. 19.*] If therefore in $AC,$ AF be taken equal $\sqrt{300},$ and from AB there be cut a part AE equal to $AF,$ the straight line joining the points E and F will be the shortest line possible that can divide the triangle ABC into two equal parts. (*For the demonstration, see page 163, vol. iii. of the COURSE.*)

To ascertain the length of this line, $EF,$ the angle $A = 16^\circ..15'$ because $\frac{25^2 + 24^2 - 7^2}{2(24 \times 25)} = .96 = \text{nat.co. sin. } 16^\circ..15'$ nearly. Therefore the angle $AEF = \frac{1}{2}$ the supplement of $16^\circ..15',$ or $81^\circ..52'\frac{1}{2}.$


And

As	Sin. $81^\circ..52'\frac{1}{2}$	- - -	Log. ar. co.	- 10.0043815
:	Sin. $16^\circ..15'$	- - - - -	Log.	9.4468927
::	$17.3205081 = AF$	- - - - -	Log.	1.2385607
	$4.89592 = EF$	- - - - -	Log.	.6898349

(Key to Vol. III. page 172.)

Consequently the position and length of the proposed shortest-possible line are determined. Q. E. F.

Ex. 4. Construct a triangle ABC right-angled at c, and having the side AC = 6 and the side BC = 8, the hypotenuse AB will necessarily be 10; which three numbers are the three sides given. But the area of the triangle = $\frac{8 \times 6}{2} = 24 = 16(1.5)$. Therefore 7



$\times 1.5$, and 9×1.5 are the two divisions proposed. Let m = the former, n = the latter; and assume in BC any point G through which a line parallel to AC shall give the trapezium ADGC to the triangle BDG, as m is to n . $BG = BC \sqrt{\frac{n}{m+n}}$ [Case 2. Prob. 1. chap. vii. vol. iii.] = $8 \sqrt{\frac{9}{16}} = \frac{24}{4} = 6$. Consequently $GC = 2$. Parallel, therefore, to AC, draw DG, so that GC may be equal to 2, the line DG is the line required. For,

As $10 : 8 :: \sin. 90 : \sin. 53^\circ .. 7' .. 48'' \frac{1}{3} =$ the angle BAC, of which the half is $26^\circ .. 33' .. 54'' \frac{1}{6}$; and the half of the angle ACB is 45° .

Now let AF and CF be two lines bisecting the angles A and c, F is the center of the inscribed circle to the triangle ABC. But

As Sin. $108^\circ .. 26' .. 5'' \frac{5}{6}$ ($= \angle AFC$)	Log. ar. co. - 10.0228788
: Sin. $26^\circ .. 33' .. 54'' \frac{1}{6}$ ($= \angle FAC$)	- - - - - Log. 9.6505149
:: 6 (=side AC)	- - - - - Log. 0.7781513
: $2.8284271 = \sqrt{8} = \sqrt{2^2 + 2^2} = FC$	- - - Log. 0.4515450

That is, the point F is in the line GD, which was to be proved.


Hence the length of the dividing line is $4\frac{1}{2}$.

It is evident that a line drawn through the point E parallel to BC would not divide the triangle ABC in the ratio required, but the position of a line parallel to BC to effect this is determinable by Case 2. Prob. 1. See lm in the following diagram.

(Key to Vol. III. page 172.)

Again. As Sin. $62^\circ .. 52' .. 8''$ - - Log. ar. co. - 10.0506269
 : Sin. 70° - - - - - Log. 9.9729858
 :: 17 (*side subt. $\angle 62^\circ .. 52' .. 8''$*) Log. 1.2304489
 : 17.95 *base and longest side of the Δ .* Log. 1.2540616

Constructing, then, a triangle of which the three sides are respectively equal to 14, 17, and 17.95, (which let be ABC,) it is required to divide it into three equal parts by lines parallel to AC. Assume D and E in AB, as the extremities of the lines sought.



$$\left. \begin{aligned} BD &= BA \sqrt{\frac{1}{3}} = 14 \sqrt{\frac{1}{3}} = \frac{14}{3} \sqrt{3} = 8.0829037 \\ BE &= BA \sqrt{\frac{2}{3}} = 14 \sqrt{\frac{2}{3}} = \frac{14}{3} \sqrt{6} = 11.4309519 \end{aligned} \right\}$$

Take, therefore, $BD = 8.0829037$, and
 $BE = 11.4309519$, and parallel to AC draw DF and EG. It will be,

$$\left. \begin{aligned} \text{As } BA : AC &:: BE : EG, \text{ and} \\ \text{As } BA : AC &:: BD : DF, \text{ that is,} \\ \text{As } 14 : 17.95 &:: 11.4309519 : 14.6561133 = EG \\ \text{As } 14 : 17.95 &:: 8.0829037 : 10.3634374 = DF \end{aligned} \right\} \text{Ans.}$$

Ex. 6. Suppose 56° one of the angles of the triangle, then $66^\circ .. 3'$ is the 3d angle. Also suppose 58° to be one of the angles, in which case the 3d angle is $64^\circ .. 3'$. It will be,

As Sin. $57^\circ .. 57'$ - - Log. ar. co. - 10.0718166
 : Sin. 56° - - - - - Log. 9.9185742
 :: 112.65 - - - - - Log. 2.0517312
 : 110.19 - - - - - Log. 2.0421220

And

As Sin. $57^\circ .. 57'$ - - Log. ar. co. - 10.0718166
 : Sin. $66^\circ .. 3'$ - - - - - Log. 9.9608987
 :: 112.65 - - - - - Log. 2.0517312
 : 121.46 - - - - - Log. 2.0844465

Now $121.46 - 110.19 = 11.27$, but should be 8. Again,

(Key to Vol. III. page 172.)

As Sin. $57^\circ .. 57'$ - - Log. ar. co. - 10·0718166
 : Sin. 58° - - - - - Log. 9·9284205
 :: 112·65 - - - - - Log. 2·0517312

: 112·71 - - - - - Log. 2·0519683

And

As Sin. $57^\circ .. 57'$ - - Log. ar. co. - 10·0718166
 : Sin. $64^\circ .. 3'$ - - - - - Log. 9·9538448
 :: 112·65 - - - - - Log. 2·0517312

: 119·51 - - - - - Log. 2·0773926


And $119·51 - 112·71 = 6·8$, but should be 8.

In the first instance the error is 3·27 *too much*, in the 2d it is 1·2 *too little*. The *sum* of the errors is 4·47. But,

As $4·47 : 2^\circ :: 1·2 : 0^\circ .. 32'·22$ the *excess* of 58° , Therefore the three angles are $57^\circ .. 57'$, $57^\circ .. 28'$, and $64^\circ .. 35'$. Hence the three sides of the triangle are 112·65, 112·05, and 120·05, *nearly*.

Construct, then, a triangle, as ABC, having its three sides equal to those three numbers, and let P be the determinate point within the triangle.

Then because three equal divisions are proposed, make AD equal to one third of AC, and draw DE parallel to AB. Likewise make BF equal to one third of BA, and parallel to BC draw FG, the point of intersection P of these parallels is the determinate point. To this point, therefore, from the three angles let CP, AP and BP be drawn; these will be the lines required. (For demonstration see page 167, vol. iii.)




REMARK. Had AB been bisected, a line from c to the point of bisection would have passed through P, and a line from A or B to the middle of the opposite side, by intersecting the line from c, would have determined P. It is therefore evident that PE is equal to PD, and each equal to $\frac{1}{3}$ of AB, that is, equal one third of $112·65 = 37·55$. Consequently $CD = \frac{2}{3}(120·05) = 80·03$, and $CE = \frac{2}{3}(112·05) = 74·7$. Likewise the angle CDE ($= \angle CAB$) = $57^\circ .. 28'$, and the angle CED ($= \angle CBA$) = $64^\circ .. 35'$. Moreover $EP = 37·55$, $EB = 37·35$, and the angle BEP = the supplement of the

(Key to Vol. III. page 172.)

angle $ABC = 115^\circ .. 25'$. Also $AD = 40.015$ and the angle $ADP = 122^\circ .. 32'$ the supplement of $57^\circ .. 28'$ the angle BAC or EDC . Here, then, are three separate triangles, in each of which are given two sides and the included angle to find the third side. Hence $CP = 67.693$, $BP = 63.065$, and $AP = 68.021$. Which were to be determined.


Ex. 7. $\sqrt{28^2 + 45^2} = \sqrt{2809} = 53 =$ the hypotenuse. Construct, therefore, a triangle of which the three sides are 28, 45, and 53, which let be DEF .

Bisect DF in G , DE in H , and EF in I . Join GH , GE , and GI , the lines GH , GE , and GI , are the lines required. For G is the middle of the hypotenuse, and the four triangles are evidently equal. [Eucl. i. 37.] It is likewise manifest that GI is parallel to DE , and that GH is parallel to EF . Consequently $GH = 14$, and $GI = 22\frac{1}{2}$. Lastly, $GE = \sqrt{GH^2 + GI^2} = \sqrt{14^2 + 22.5^2} = \sqrt{702.25} = 26.5$. Wherefore 14, 26.5, and 22.5 are the three lines required.




UNIVERSALLY.

Bisect DF in G , DG in L , and GF in K . Draw EG ; and parallel to EG , through the points K and L , draw KI and LH . Join GI and GH , then DHG , HGE , EGI and IGF , are the four triangles required, and in all respects as before.



(Page 173.)

Ex. 8.




Construct the given rectangle, $ABCD$, and let P be the point in

(Key to Vol. III. page 173.)

the longest side $\frac{4}{5}$ of AB from A. Join PC and PD, and through the point B draw BF parallel to PC, and through the point A draw AE parallel to PD: also produce CD both ways to E and F. Take EG equal to one fifth part of EF, and connect PG; GP is the line required. (See *Demonstration*, page 168, vol. iii.) Because EF is double of DC, EG = 6, and GD = 2. Through P, therefore, draw Pr parallel to AD or to BC, rG = 2. Consequently PG = $\sqrt{Pr^2 + rG^2} = \sqrt{9^2 + 2^2} = \sqrt{85} = 9.2195445$. Q. E. I.

Ex. 9.

Let ABCDEF be the given hexagon, and o the center of the circumscribing circle. Draw any two lines as GH, IK, to represent the lines sought; also join OF, OE, OD, and OA. It is evident (since by the hypothesis OG and OK are equal) that, FG and EK must be equal.



If, therefore, a be substituted for $AF = FE = ED = 12$ by the question, x for $FG = EK$, and am for the area of the equilateral triangle FOE, the area of the triangle GOF will be truly expressed by xm , and the area of the triangle AOG by $(a-x)m$. [Eucl. B. vi. Prop. 1.]

But the two triangles GOF, KOE are in all respects equal, and the triangles AOG, HOD, DOK, are likewise equal and similar; consequently the whole area GOKEF = $am + 2xm$, and the whole area KOHD = $2(a-x)m = 2am - 2xm$. Now by the question these two whole areas are equal; wherefore $am + 2xm = 2am - 2xm$, that is, $a + 2x = 2a - 2x$. Hence $x = \frac{1}{4}a = 3$, and the angle GOF = $13^\circ .. 53' .. 52''.8$.

Also by analogy,


As Sin. $13^\circ .. 53' .. 52''.8$ (\angle GOF)	Log. ar. co. -	10.6194376
: Sin. 60° (\angle OFG)	- - - - -	Log. 9.9375306
:: 3, (side FG)	- - - - -	Log. 0.4771213
		<hr/>
: 10.81656 (side OG)	- - - - -	Log. 1.0340895
		<hr/>

(Key to Vol. III. page 173.)

But $2 \times 10.81656 = 21.63312 = GH = IK$, which being the *only equal lines* that can possibly divide the hexagon into four equal parts, are necessarily the shortest possible. Q. E. I.

Ex. 10. Let ABC be the proposed triangle, $AB=5$, $AC=6$, and $BC=7$.

Draw mn at right angles to BC , and dividing the triangle ABC into two equal parts, by Problem I. Case 3, page 161, vol. iii.; and draw og parallel to BC , dividing the triangle ABC into two equal parts, by Prob. I. as above, Case 2. So shall mn , og be the two lines required.



The demonstration is obvious.

* * * This example admits of three answers, since it is immaterial whether mn be drawn at right angles to BC , to AB , or to AC .

SOLUTION OF EQUATIONS.

(Page 186)

Ex. 4. Because the 1st term of an equation is the unknown quantity raised to the power denoted by the number of roots, the 1st term of the equation to be produced must be x^2 .

And because the 2nd term will contain the unknown quantity raised to a power less than the former by *unity*, with a coefficient equal to the sum of the roots taken with contrary signs, it follows that $(-\frac{3}{4} - \frac{4}{5})x = -\frac{31}{20}x$ is the *second term* of the equation sought. Also, because the unknown quantity raised to a power less by 2 than that of the 1st term, with a co-efficient equal to the sum of all the products that can arise by multiplying all the roots of the equation two and two, will constitute the third term of the equation, $(\frac{3}{4} \times \frac{4}{5}) = \frac{12}{20} = \frac{3}{5}$ must be the *third and last term* of the equation required.

Therefore $x^2 - \frac{31}{20}x + \frac{3}{5} = 0$. Ans.

(Key to Vol. III. page 187.)

Ex. 5. It is evident by the last example that the leading term will be x^3 .

Now the sum of 2, 5, and -3 taken with contrary signs is -4 ; consequently the 2nd term must be $-4x^2$.

Likewise the sum of (2×5) , (2×-3) , and (5×-3) is $10 - 6 - 15 = -11$; therefore $-11x$ is the 3rd term of the equation.

And since the 4th term will consist of the unknown quantity raised to a power less by 3 than that of the 1st term, with a co-efficient equal to the sum of all the possible trinary products of the roots taken with contrary signs, it follows that $(-2 \times -5 \times +3)x^0 = 30$ is the 4th and last term of the equation proposed.

Hence, when connected, $x^3 - 4x^2 - 11x + 30 = 0$. Ans.

Ex. 6. It is manifest by examples 4th and 5th, [and Theor. i. page 176] that the powers of the unknown quantity in the several terms will be $x^4 \dots x^3 \dots x^2 \dots x^1 \dots x^0$, and that the sign and co-efficient of the leading term will be $+1$.

Now for the co-efficient of the second term (equal to the sum of all the roots taken with contrary signs) it will be, $-1 - 4 + 5 - 6 = -6$ the co-efficient of the 2nd term.

And for the co-efficient of the 3d term, (equal to the sum of all the binary products of the roots,) it will be

$(1 \times 4) + (1 \times -5) + (1 \times 6) + (4 \times -5) + (4 \times 6) + (-5 \times 6) = 4 - 5 + 6 - 20 + 24 - 30 = -21$ the co-efficient of the 3d term.

Next for the co-efficient of the 4th term (equal to the sum of all the trinary products of the roots taken with contrary signs)

it will be $(-1 \times -4 \times +5) + (-1 \times -4 \times -6) + (-1 \times +5 \times -6) + (-4 \times +5 \times -6) = 20 - 24 + 30 + 120 = +146$ the co-efficient of the 4th term.

Lastly, since the co-efficient of the 5th term will be the continued product of all the roots taken with contrary signs, it will evidently be, $-1 \times -4 \times +5 \times -6 = -120$ the co-efficient of that term.

Hence by uniting the co-efficients with the powers of the unknown quantity, and connecting the terms, there arises $x^4 - 6x^3 - 21x^2 + 146x - 120 = 0$ the equation required.

Ex. 7. Here $p = 347$, $q = 22110$ and the equation agrees with the 1st FORM.

Now taking the tang. $\Lambda = \frac{2}{p} \sqrt{q}$, the two roots of the proposed equation are $x = + \text{tang. } \frac{1}{2} \Lambda \sqrt{q}$, and $x = - \text{cot. } \frac{1}{2} \Lambda \sqrt{q}$.

(Key to Vol. III. page 187.)

Again,

$$\begin{array}{r} \text{Log. cot. } 36^\circ \text{ .. } 21' \text{ .. } 8'' \frac{1}{2}, (\frac{1}{2}A) \text{ - is } 0.1831340 \\ \text{Also log. } \sqrt{q} \text{ (as above) - - - is } 0.8668660 \end{array}$$

$$\text{The log. sum - - - - - is } 1.0000000 = \text{log. of } 10.$$

Consequently 10, and $-\frac{5}{12}$ are the *two* values of x in the proposed equation. Q. E. I.

Ex. 9. In this example $p = \frac{264}{25}$, $q = \frac{695}{25}$, and the equation is of the 4th FORM. Hence,

$$\text{If } \sin. A = \frac{2}{p} \sqrt{q} = \frac{50}{264} \sqrt{q}, \text{ } x \text{ in one root is equal to } \text{tang. } \frac{1}{2} A \sqrt{q}, \text{ and in the other root equal to the cot. } \frac{1}{2} A \sqrt{q}.$$

In Logarithms.

$$\begin{array}{r} \text{Log. of } 695 \text{ - - is } 2.8419848 \\ \text{Log. of } 25 \text{ - - is } 1.3979400 \end{array}$$

$$\begin{array}{r} \frac{1}{2} \text{ the log. diff. - is } 0.7220224 = \text{log. } \sqrt{q}. \\ \text{Log. of } 50 \text{ - - - is } 1.6989700 \\ \text{Ar. co. log. } 264 \text{ is } -3.5783961 \end{array}$$

$$\text{Sum of the 3 last logs. is } -1.9993885 = \text{log. sin. } 86^\circ \text{ .. } 57' \text{ .. } 36'' = A.$$

Wherefore

$$\begin{array}{r} \frac{1}{2} A = 43^\circ \text{ .. } 28' \text{ .. } 48'', \text{ of which the log. tang. is } -1.9769470 \\ \text{But log. } \sqrt{q} \text{ (already found) - - - - - is } 0.7220224 \end{array}$$

$$\text{The log. sum is } 0.6989694 = \text{the log. of } 5 = x \text{ in one of the roots of the equation.}$$

Again,

$$\begin{array}{r} \text{Log. cot. of } 43^\circ \text{ .. } 28' \text{ .. } 48'' \text{ is } 0.0230530 \\ \text{And log. } \sqrt{q} \text{ (as above) } 0.7220224 \end{array}$$

$$\text{The log. sum is } 0.7450754 \text{ the log. of } 5.56 = \frac{139}{25}. \text{ Therefore } 5 \text{ and } 5.56 \text{ are the two values of } x \text{ in the given equation. Q. E. I.}$$

Ex. 10. The proposed equation is of the 4th FORM, wherein $p = 24113$, and $q = 481860$.

$$\text{Taking, therefore, } \sin. A = \frac{2}{p} \sqrt{q} \text{ the two values of } x \text{ are the tangent and cotangent of } A \sqrt{q}.$$

(Key to Vol. III. page 187.)

In Logarithms.

$$\begin{aligned} \frac{1}{2} \log. 481860 &= 2.8414604 = \log. \sqrt{q} \\ \text{Log. of } 2 &= 2.3010300 \\ \text{Ar. co. log. } 24113 &= -5.6177488 \end{aligned}$$

The log. sum = -2.7602392 the log. sin. $3^\circ..18'..2''..24''' = A$.

Consequently $\frac{1}{2} A = 1^\circ..39'..1''..12'''$ of } -2.4595690
 which the log. tang. is - - - - - }

The log. \sqrt{q} (already found) is 2.8414604

The log. sum is 1.3010294 the logarithm of 20.

Again,

$$\begin{aligned} \text{Log. cot. of } 1^\circ..39'..1''..12''' &\text{ is } +1.5404310 \\ \text{And log. } \sqrt{q} \text{ (as above) is } &2.8414604 \end{aligned}$$

The log. sum is 4.3818914 the log. of 24093. Wherefore 20 and 24093 are the two values required.

Ex. 11. Here $p=3$, $q=1$, and the equation agrees with the 2nd FORM for Cubic equations; consequently,

$$\text{If } \sin. B = \frac{p}{3q} 2\sqrt{\frac{1}{3}p}, \text{ tang. } A = \sqrt[3]{\tan. \frac{1}{2}B}.$$

$$\text{Then } x = \mp \text{co. sec. } 2A \times 2\sqrt{\frac{1}{3}p}.$$

But $\frac{\frac{1}{3}p}{q} \times 2\sqrt{\frac{1}{3}p} = \frac{1}{1} \times 2\sqrt{1} = 2 = \sin. B$ and evidently greater

than *Unity*: wherefore taking co. sec. of $3A = \frac{\frac{1}{3}p}{q} \times 2\sqrt{\frac{1}{3}p} = 2 = \text{co. sec. } 30^\circ$, it will be, $A=10^\circ$. And

$$x = \pm \sin. A \times 2\sqrt{\frac{1}{3}p} = \pm \sin. 10^\circ \times 2.$$

$$x = \pm \sin. (60^\circ - A) 2\sqrt{\frac{1}{3}p} = \pm \sin. (60^\circ \mp 10^\circ) \times 2 = \pm 2 \times \begin{cases} 50^\circ \\ 70^\circ \end{cases}$$

And

$$x = \pm \sin. (60^\circ + A) 2\sqrt{\frac{1}{3}p} = \pm \sin. (60^\circ \pm 10^\circ) \times 2 = \pm 2 \times \begin{cases} 70^\circ \\ 50^\circ \end{cases}$$

Again, [*Theor. 2. page 178. vol. iii.*] it is evident by the variation and permanency of the signs of the proposed equation that one of the roots, *only*, is *positive* and two *negative*, and that the double sine of 70° is the positive root. Consequently

(Key to Vol. III. page 187.)

$$x = \left\{ \begin{array}{l} +1.8793852 = \sin. 70^\circ \\ -1.5320888 = -\sin. 50^\circ \\ - .3472964 = -\sin. 10^\circ \end{array} \right\} \text{ to Radius 2.}$$

Ex. 12. Here $p=1$, and $q=6$; wherefore $\frac{\frac{1}{3}p}{q} = \frac{1}{18}$, and $2\sqrt{\frac{1}{3}p} =$

$$\sqrt{\frac{12}{9}} = \frac{2}{3}\sqrt{3}. \text{ Hence } \frac{\frac{1}{3}p}{q} \times 2\sqrt{\frac{1}{3}p} = \frac{1}{27}\sqrt{3}.$$

In Logarithms.

$$\begin{array}{r} \frac{1}{2} \log. \text{ of } 3 = 0.2385606 \\ \log. \text{ of } 1 = 0.0\text{---} \\ \text{Ar. co. log. of } 27 = -2.5686362 \end{array}$$

$$\text{The log. sum} = -2.8071968 \text{ the log. sin. } \left\{ \begin{array}{l} 3^\circ..40'..41'' \\ 176^\circ..19'..19'' \end{array} \right\} = \mathbf{B}$$

$$\text{Or, } \frac{1}{2}B = \left\{ \begin{array}{l} 1^\circ .. 50' .. 20'' \frac{1}{2} \\ 88^\circ .. 9' .. 39'' \frac{1}{2} \end{array} \right\}$$

But log. tang. $A = \log. \sqrt[3]{\text{tang. } 88^\circ .. 9' .. 39'' \frac{1}{2}} = 0.4977964$ the
log. tang. of $72^\circ .. 22' .. 4''$, which doubled $= 2A = 144^\circ .. 44' .. 8''$.

And co. sec. $144^\circ .. 44' .. 8'' = \text{co. sec. } 35^\circ .. 15' .. 52'' = \text{sec. } 54^\circ .. 44' .. 8''$.

Hence $(\text{co. sec. } 2A \times 2\sqrt{\frac{1}{3}p})x =$

$$\sqrt[3]{3} \times \left\{ \begin{array}{l} \text{co. sec. } 144^\circ .. 44' .. 8'' \\ \text{co. sec. } 35^\circ .. 15' .. 52'' \\ \text{sec. } 54^\circ .. 44' .. 8'' \end{array} \right\} \text{ Ans. } *$$

* In this Example the Answer given in the COURSE differs 12 SECONDS from the Answer in the Key.

Ex. 13. Given $25x^3 + 75x - 46 = 0$: that is, divided by 25,

$$x^3 + 3x - \frac{46}{25} = 0.$$

Here $p = 3$, $q = \frac{46}{25}$, and the equation is of the 1st CUBIC

FORM. Wherefore

$$\text{Tang. B.} = \frac{\frac{1}{3}p}{q} \times 2\sqrt{\frac{1}{3}p} = \frac{25}{46} \times 2 = \frac{50}{46}.$$

In Logarithms.

$$\begin{array}{r} \text{Log. of } 50 \text{ is } 1.6989700 \\ \text{Ar. co. log. } 46 \text{ is } -2.3372422 \end{array}$$

$$\text{The log. sum is } 0.0362122 = \log. \text{ tan. } \mathbf{B}.$$

(Key to Vol. III. page 187.)

$$\text{That is, } B = \left\{ \begin{array}{l} 47^\circ \dots 23' \dots 9'' \frac{2}{3}, \text{ or } \\ 132^\circ \dots 36' \dots 50'' \frac{3}{5} \end{array} \right\}$$

Taking the latter value,

$$\frac{1}{2}B = 66^\circ \dots 18' \dots 25'' \cdot 3 \text{ of which the log. tang. is } 0.3576517$$

But log. tang. $A = \log. \sqrt[3]{\text{tang. } \frac{1}{2}B} = 0.1192172$ the
log. tang. of $52^\circ \dots 46' \dots 1'' \frac{3}{4}$ nearly. Consequently

$$2A = 105^\circ \dots 32' \dots 3'' \frac{1}{2} \text{ nearly, or (the supplement of that arc =)} \\ 74^\circ \dots 27' \dots 56'' \frac{1}{2}.$$

$$\text{Hence } x (= \cot. 2A \times 2\sqrt{\frac{1}{3}p}) = 2 \cot. \left\{ \begin{array}{l} 74^\circ \dots 27' \dots 56'' \frac{1}{2} \\ 105^\circ \dots 32' \dots 3'' \frac{1}{2} \end{array} \right\} \text{ Ans.}^*$$

* Differing 8.5 seconds from the Answer given with the Question.

Ex. 14. Because $x^4 - 8x^3 - 12x^2 + 84x - 63 = 0$, it follows that
 $x^4 - 8x^3 = 12x^2 - 84x + 63$. Hence, by adding $16x^2$ to both
sides of the equation,

$$x^4 - 8x^3 + 16x^2 = 28x^2 - 84x + 63 = 7(4x^2 - 12x + 9), \text{ Or}$$

$$(x^2 - 4x)^2 = 7(2x - 3)^2; \text{ that is, } x^2 - 4x = \pm \sqrt{7} \times (2x - 3).$$

Consequently $x^2 - (4 \pm 2\sqrt{7})x = \mp 3\sqrt{7}$ a quadratic equa-
tion; which solved gives $x = 2 + \sqrt{7} \pm \sqrt{11 + \sqrt{7}}$. Ans.

Ex. 15. If $548x^2$ be added to both sides of the given equation,
it becomes,

$$x^4 + 36x^3 + 148x^2 - 3168x + 7744 = 548x^2.$$

And extracting the square root on both sides,

$$x^2 + 18x - 88 = \pm x\sqrt{548}. \text{ That is,}$$

$$x^2 + (18 \mp \sqrt{548})x = 88; \text{ and solving this quadratic equa-} \\ \text{tion, } x = 12.46766 \text{ very nearly. Ans.}^*$$

* We cannot help remarking the great disparity between our Answer and that
of Dr. Hutton, in the present Example.

Ex. 16. Here, as in Ex. 15, the co-efficient of the 4th term divided
by that of the second gives the square root of the last term
for the quotient; therefore, adding $256x^2$ to both sides of
the biquadratic, there arises

$$x^4 + 24x^3 + 142x^2 - 24x + 1 = 256x^2.$$

And by extracting the square root on both sides,

$$x^2 + 12x - 1 = \pm 16x. \text{ Hence}$$

(Key to Vol. III. page 187.)

$$x^2 + \left\{ \begin{array}{l} -4 \\ +28 \end{array} \right\} x = 1. \quad \text{Wherefore}$$

$$x = \left\{ \begin{array}{l} 2 \pm \sqrt{5} \\ -14 \pm \sqrt{197} \end{array} \right\} \quad \text{Ans.}$$

Ex. 17. Adapting the given equation to Euler's Rule, it will be $x^4 - 0x^2 - 12x - 5 = 0$. Hence the roots are $1 \pm \sqrt{2}$ and $-1 \pm 2\sqrt{-1}$.

Ex. 18. Adding x^2 to both sides of the given equation, there arises $x^4 - 12x^3 + 48x^2 - 72x + 36 = x^2$, two perfect squares. And, by extraction of the square root, $x^2 - 6x + 6 = x$, or $x^2 - 7x = -6$.

But the two roots of this quadratic (*and consequently two roots of the biquadratic*) are 3.5 ± 2.5 , that is, 6 and 1.

Now the other two roots of the proposed equation admit of being found in a variety of ways; for it is evident, by the alternation of the signs, that they are positive, and by the co-efficient (-12) of the second term, that their sum is 6.

Let them be found, however, in the following manner:

Divide the given equation by $x-6$, and it is depressed to the cubic equation $x^3 - 6x^2 + 11x - 6 = 0$.

In like manner divide the given equation by $x-1$, and it is depressed to the cubic equation $x^3 - 11x^2 + 36x - 36 = 0$.

Putting these two cubic equations equal, and reducing, the result is $x^2 - 5x = -6$.

Now the two roots of this quadratic, (*and consequently the other two roots of the biquadratic*), are $2.5 \pm .5$, that is, 3 and 2.

By which it is plain, that the four roots required are 1, 2, 3, and 6. Q. E. I.

Ex. 19. If the given equation be divided $x+a$, it will be depressed to $x^4 - 6ax^3 - 74a^2x^2 + 6a^3x + a^4 = 0$, a biquadratic equation. Whence it follows that $-a$ is one of the roots of the sursolid equation.

Let $81a^2$ be now added to both sides of the new equation, and there will arise $x^4 - 6ax^3 + 7a^2x^2 + 6a^3x + a^4 = 81a^2$, two perfect squares; and by extraction of the square root, $x^2 - 3ax - a^2 = 9a$, or $x^2 - 3ax = a^2 + 9a$.

But the two roots of this quadratic (*and consequently two roots of the biquadratic, and also of the proposed equation*), are $\frac{3}{2}a + \sqrt{\frac{1}{4}a^2 + 9a}$, and $\frac{3}{2}a - \sqrt{\frac{1}{4}a^2 + 9a}$, by which the remaining two roots are known. Or the biquadratic may be transformed into an equation that shall want the 2nd term, (and to which Euler's rule will be applicable), by assuming


(Key to Vol. III. page 187.)

$m = -\frac{3}{2}a$, and substituting $z - m$ (that is, $z + \frac{3}{2}a$) for x . Or, by the rule for the solution of biquadratic equations, page 183, vol. iii. of the Course, the biquadratic above found being of the form $x^4 + 2px^3 = qx^2 + rx + s$, where $p = -3a$, $q = 74a^2$, $r = -6a^3$, and $s = -a^4$, the four roots are obtained from two quadratic equations. These roots, by whatever method determined, are surd quantities, more or less complicated, and in the simplest form are $-3a + a\sqrt{10}$, $-3a - a\sqrt{10}$, $6a + a\sqrt{37}$, and $6a - a\sqrt{37}$.

EQUATIONS TO CURVES.

(Page 205.)

Ex. 2. If in the right line BC there be taken a point a , and if the points b, c, d, e, f, g , be assumed such that ab, ac, ad, ae, af , and ag , be in arithmetical progression, then if ah, bi, ck, dl , &c. be in geometrical progression, and perpendicular ordinates at those points, the curve $Donmlk i h E$ passing through the extremities of the ordinates, is the logarithmic curve, BC an asymptote to the curve, and AB (being parallel to, and greater by the least assignable magnitude than, the greatest ordinate,) is the other asymptote.* Because the abscissæ de, df , and dg , are in arithmetical progression, and their ordinates em, fn , and go , in geometrical progression, it follows that, when $dl = 1$, and $em = a$, then $fn = a^2$, $go = a^3$, &c.



Wherefore, if x represent any part whatever of the arithmetical series ascending, measured from d , the corresponding ordinate is a^x ; that is, $y = a^x$, a being the number of which the logarithm is 1 in the system of logarithms represented by the curve.

* For the square of any term of a geometrical series is equal to the rectangle of any two terms equidistant from it; consequently the square of dl is equal to $og \times ah$: but og increases as ah decreases; yet, it is plain, ah can never be equal to 0, nor og infinite. Therefore BC continually approaches the curve, and when og is inconceivably near infinite by ah being inconceivably near nothing, the distance of og from AB is less than the least assignable magnitude, and AB is an asymptote to the curve.

(Key to Vol. III. page 205.)


Ex. 3. Let the curve be MB, and the generating semicircle ADB; also let AP be any absciss whatever, end PM the corresponding ordinate; then by the properties of the curve, As AP : PD :: AB : PM.

But the angle ADB is a right angle, [Eucl. iii. 31.] and it is As AP : PD :: PD : PB. [Eucl. vi. 8.]

If, therefore, d be put for AB, the value of PD is $\sqrt{dx-x^2}$, and

As $x : \sqrt{dx-x^2} :: d : y$. That is

$$xy = d \sqrt{dx-x^2}, \text{ and } y = \frac{d \sqrt{dx-x^2}}{x} = d \sqrt{\frac{dx-x^2}{x^2}} = d \sqrt{\frac{d-x}{x}}. \text{ Q. E. I.}$$



LOCI OF EQUATIONS.



(Page 208.)

Ex. 4. First let n be some odd number, and x positive, so shall $y = \frac{a^{n+1}}{x^n}$.

Next let n be some odd number and x negative, so shall $y = -\frac{a^{n+1}}{x^n}$.

By which it is evident that the loci of the curve will be in the vertical angles ABC, abc , which situation corresponds to the common hyperbola, ac , and ac being asymptotes to the curve.

Again let n be any even number whatever; it is plain, since x^n must be positive, that y must also be positive; and in this case the curve will fall in the adjacent angles ABC and ABD, having AB, BC, and AB, BD for asymptotes.



(Key to Vol. III. page 208.)

Whereby the principal properties of the curve are apparent.

Q. E. D.

Ex. 5. Here $y = \frac{bc + bx}{a + c + x}$, by which it appears that the ordinate can meet the curve in *one point only*, there being a single corresponding value of y for each value of x . Now when $x = 0$, $y = \frac{bc}{a + c}$, consequently the curve does not pass through the point from which the values of x are measured.

Again it is manifest, that the quantity $a + c + x$ is greater than the quantity $c + x$, therefore, although y must increase when x increases, yet y can never be equal to b , except when x becomes infinite; in which case since a and c (*finite quantities*;) bear no comparison with x (*an infinite quantity*;) y may be truly expressed by $b \times \frac{x}{x}$, that is, by b . The asymptote, therefore, will touch the curve at an infinite distance. But if x be supposed negative, then $y = b \times \frac{c - x}{a + c - x}$ and when $x = c$, $y = b \times \frac{c - c}{a + c - c} = \frac{0}{a} = 0$. Wherefore B is a point in the curve as often as $AB = c$.

On the other hand, if c be less than x , it is plain that $c - x$ is negative, for which reason the ordinate to the curve will be negative until x become equal to $a + c$; at which situation $y = b \times \frac{-a}{0}$, that is *infinity*; and, therefore, x infinite.


Consequently if any point M be assumed in the straight line on which x is measured, so that $a + c = AM$, and if through M there be drawn a right line parallel to an ordinate, that right line shall be an asymptote to the curve.

Lastly, if a and c together be less than x , (x being still negative), then $c - x$ and $a + c - x$ are both negative; but y positive, because b multiplying a negative quantity divided by a negative quantity is positive: and since, in this case, $x - c$ must be always greater than $x - a - c$, it follows that y will always exceed b . Hence the several *loci* of the curve are confined to the angular space vertical to that in which they fall in the first supposition. The ratio, however, of $x - c$ to $x - a - c$, approaches more and more to the ratio of

(Key to Vol. III. page 208.)

1 to 1 as x increases, and is the ratio of 1 to 1 when x is infinite. This signifies that the curve continually approaches to its asymptote in this angular space likewise, and must therefore be the common hyperbola. To prove which, let ab be added to both sides of the given equation, then $y(a+c+x) + ab = b(a+c+x)$, and $ab = (b-y) \times (a+c+x)$. Whereby the truth of the inference is manifest. Q. E. D.

Ex. 6. If oo be assumed equal to a the line of the abscissæ, and if sL be an ordinate at right angles to this line, it is manifest, by the equation of the curve, that $x=a$, when $y=0$; therefore, the curve must pass through the point o , and have tot a line parallel to sL , for a tangent at that point.



It likewise appears by the equation of the curve, that when x vanishes, $y = \frac{1}{2}a$; wherefore, if or be drawn from the extremity o of the line oo , equal to $\frac{1}{2}a$, and parallel to tt , r will be a point of the curve on either side of the line of the abscissæ.

Finding in like manner other values of y , the several points L, L, L, L , are determined to be loci of the equation, such that, if a circle be described on the half of a , nL is always equal to the radius of that circle. For, putting $oL = z = \sqrt{x^2 + y^2}$, and the angle $oOL = v$, it will be, As $z : \text{rad.} :: x : \cos. v$, that is, taking *Unity* for radius, $x = \cos. v$. Consequently $a^2 z^2 = 4(z^2 - \frac{1}{2}az \times \cos. v)^2$; or by extracting the square root on both sides, dividing by z , and transposing, $z = \frac{1}{2}(a + a. \cos. v)$

Now because $or = \frac{1}{2}a$, if rn be drawn at right angles to oL , the proportion is,

As $or (= \frac{1}{2}a) : \text{rad.} (1) :: on : \cos. v$, that is, $on = \frac{1}{2}a. \cos. v$.

But $z = \frac{1}{2}(a + a. \cos. v) = \frac{1}{2}a + on = oL = (\text{because } or = \frac{1}{2}a,) or + oL - nL$.


Hence $or (= ro) = nL$; and nL is every where equal to or ; FF , or LL , is equal to $2or$; $oL = or \pm on$; and n always bisects LL .

* * * Many other properties of the Cardioide may be seen in the *Philos. Trans.* for 1741.

(Key to Vol. III. page 208.)

Ex. 7. Here $y = \frac{ax^3 + bx^2 + cx + d}{x}$, which is infinite when

$x=0$. And when x is infinite y is also infinite, because equal to infinity *plus* a finite quantity *plus* nothing. Again, when $x=unity$ the value of y is $a+b+c+d$; or if x be any arbitrary quantity, and either one or more of the co-efficients $=0$, then the corresponding term or terms of the equation must vanish or become wanting.



By which it appears, that the figure has four infinite branches or legs; and that since in two of them the tangent to the curve coincides with the asymptote when the point of contact is at an infinite distance, but in the other two recedes *in infinitum* and is no where found, necessarily the former are diverging hyperbolic legs in contrary directions, as CD and EF about the asymptote AB , and the latter converging parabolic legs as GD and HE meeting the hyperbolic branches in D and E . It is likewise evident that the simplest equation to the curve is when the rectangle under the absciss and ordinate is equal to $x^3 + a^3$, for the loci of the curve are in this case easily determinable by a common parabola whose absciss is $ax^2 + bx + c$, and an hyperbola having its absciss $= d \div x$, for y will be at all times equal to the *sum* or *difference* of the correspondent ordinates of those two conic sections.

Ex. 8. FIRST FOR THE CISSOID. Because the equation of the curve is $x^3 = y^2 \times (d-x)$, and because it is plain that $d-x$ decreases as x increases, it follows that y , and consequently y^2 , must continually increase; otherwise $(d-x)y^2$ could not be equal to x^3 .

Again, it is manifest that x can never equal d , for in that event x^3 would equal 0, which is absurd; wherefore y or PM may approach within less than the least assignable magnitude of BX , but can never coincide with it. *That is, BX is an asymptote* to the curve AMQ .*

NEXT FOR THE WITCH. It is evident by the equation of the curve that the value of $d-x$, and consequently that of

* The curve AMQ bisects the curve ATB , and the cissoidal space under the diameter AB , the asymptote BX and the curve AMQ infinitely produced, is six times the area of the generating semicircle.

(Key to Vol. III. page 208.)

$d\sqrt{\frac{d-x}{x}}$, must decrease by the increase of x . Now when $x=d$, y vanishes; but when $x=0$, y is infinite, since $d\sqrt{\frac{d-0}{0}}$ is infinite. If, therefore, a tangent be drawn to the generating semicircle at the point Δ , that tangent will be an asymptote to the curve MB , inasmuch as PM may approach within less than the least assignable magnitude of the tangent, but can never coincide with it.

FLUXIONS AND FLUENTS.

(Page 229.)

Ex. 5. Ans. $-\frac{2}{3}x^2 \times (a^2 - x^2)^{\frac{3}{2}} - \frac{4}{15}a^2$.

Ex. 6. Ans. $-2x^4 \times (a^2 - x^2)^{-\frac{1}{2}} - \frac{8a^2x^2 + 16a^4}{3}$.

Ex. 7. Ans. $\frac{(a-x^n)^{\frac{3}{2}}}{-\frac{7}{2}anx^{\frac{7}{2}n}} \times \left[1 + \frac{\frac{4}{5}ax^n + \frac{8}{15}x^{2n}}{a^2} \right] =$


$$-\frac{(a+x^n)^{\frac{3}{2}} \times 30a^2 + 24ax^n + 16x^{2n}}{105na^3x^{\frac{7}{2}n}}$$

LATERAL PRESSURE OF FLUIDS.

(Page 262.)

Ex. 3. Let $ACDB$ be a perpendicular section of the canal, and let the sides AC and BD of the trapezium AD , meet, when produced, in E ; also let EH be perpendicular to AB , and parallel to CD draw CF and DG . It will be, As $AF : FC :: AH : HE :: 3 \text{ ft.} : 8 \text{ ft.} :: 10 \text{ ft.} : 26\frac{2}{3} \text{ ft.} = EH$.

Now if o be assumed as the center of gravity of the trapezium, m as the center of gravity of the triangle ABE , and n that of the triangle CDE , then Hm [*Statics, art. 229. vol. ii.*] = $8\frac{2}{3}$ ft. and $En = 12\frac{4}{9}$ ft. But, As trapez. $AD : \Delta CDE :: nm : mo :: (26\frac{2}{3})^2 - (18\frac{2}{3})^2 : (18\frac{2}{3})^2 :: 5\frac{1}{3} \text{ ft.} : 5\frac{1}{3} \text{ ft.} = mo$.



(Key to Vol. III. page 263.)

Hence the depth of o below the surface of the water is $3\frac{5}{7}\frac{1}{2}$ ft. which drawn into 136, the area of the dam-gate, the result is 511.9 *very nearly*.

But 511.9 cubic feet of water weigh 511900 oz. avoirdupois = the pressure = $31993\frac{3}{4}$ lb. = 14.28292 tons. Ans.

Otherwise and more elegantly by Mr. DOWLING's Rule, as practised at the Academic Institution, Highgate.

To $\frac{1}{3}$ the difference of the breadth at the surface and bottom in feet add the less breadth, and multiply the sum by half the square of the depth for the number of Cubic feet of water whereof the absolute weight shall be equal to the pressure required.

$\frac{1}{3}$ the diff. of the breadths is 2 feet.
Add the less breadth - - - - 14 feet.

Sum 16 feet.

Multiply by $\frac{1}{2}$ the square of the depth = 32 square feet.

PRODUCT 512 cubic feet.

But the absolute weight of 512 cubic feet of water is 14.28571 &c. tons *the true pressure required*.

F I N I S.

Prospectus

OF THE

ADVANTAGES ENJOYED BY THE PUPILS

AT THE

ACADEMIC INSTITUTION,

MANSION-HOUSE, HIGHGATE,

CONDUCTED BY

DANIEL DOWLING,

PROFESSOR OF THE MATHEMATICS; LECTURER IN NATURAL
PHILOSOPHY AND ASTRONOMY; AND AUTHOR OF
THE KEY TO HUTTON'S COURSE OF
MATHEMATICS.

Situation.

THE Mansion-House stands on Highgate Hill, on a spot the most healthy and picturesque in Middlesex, commanding a beautiful view of London and the Thames, with the hills of Surrey and Kent beyond them, the groves of Essex and Epping Forest on the left, and the village of Hampstead, almost contiguous, on the right; whilst a delightful assemblage of verdant fields, neat cottages, and shady lawns, constitute the fore ground of an enchanting prospect of more than 200 square miles.

Annexed to the Mansion-House (formerly the residence of Lady Cave) are eighteen acres of good garden, arable, and meadow land, affording Mr. Dowling a constant supply of milk, fruit, and every useful vegetable in season.

Stages and Post.

There are stage-coaches to and from Highgate every hour of the day, with three deliveries of letters and two departures of post.

Age, Accommodation, and Treatment.

Young Gentlemen, of all ages, are received and educated according to their future destination in life. The school-room and play-ground are dry, comfortable, and extensive, with arbours to retire to when it rains.

Mr. DOWLING treats his scholars with the utmost mildness, endeavours to form in them the habits and disposition of gentlemen, and cares for their health, comfort, and religious concerns with parental solicitude.

Each pupil has a separate bed, and the whole family breakfast and dine together.

Instruction.

THE ENGLISH LANGUAGE, ELEGANCE IN COMPOSITION, HISTORY, PLAIN AND ORNAMENTAL WRITING, AND ARITHMETIC, are taught indiscriminately to all the pupils, and on a plan which insures incredible success, in comparatively a short time.

The LATIN and GREEK languages, essential embellishments to a good education, are likewise indefatigably pursued, unless the age and views of the student determine otherwise.

The STUDY OF THE MATHEMATICS, not so much, indeed, for the exquisite pleasure wherewith it fills the mind

in tracing the beautiful relations of quantity and figure, as for its communication of an invaluable habit of patient research, a love of truth, and a system of accurate reasoning, forms a grand feature in the plan of education.

And whether the youth be intended for the civil or military department of government, the land or sea service of his country, the Honourable East India Company's service, or the useful walks of private life, he is shortly made master of the branches more immediately requisite, by proper books, a well directed attention, and the use of the best Instruments.

The FRENCH LANGUAGE is also unremittingly followed up, and soon acquired in perfection, because, a considerable part of the family being French, that language is spoken in its purity in the House.

The Pupils have likewise an excellent opportunity of learning SPANISH and the other mercantile tongues.

BOOK-KEEPING, with all the routine of the Counting-House, and the forms of actual business, engages a reasonable portion of the time of the future Merchant; and it has been often found, that the information obtained in this department, under Mr. DOWLING, has been superior to that received by the generality of articed clerks.

Lectures.

To Mr. DOWLING's Lectures in Natural Philosophy and Astronomy, illustrated by the most complete set of apparatus possible, and one of the largest telescopes in England, the pupils have free access, and are assisted in their experiments and observations. Or, if the study of Chemistry be deemed expedient, there is a laboratory fitted up with all the necessary utensils, tests, and re-agents.

Dancing, Drawing, Music, and Fencing.

Approved Masters in each of these polite branches of Education, attend two or three times a week, at suitable hours, appropriated expressly for that purpose, without encroaching too much either on *Business* or *Recreation*.

Terms.

Pupils, under 10 years of age, pay 40 guineas per annum.

from 10 to 15 years, 50 guineas per annum.

from 15 to 20 years, 60 guineas per annum.

Subject to an extra charge for washing, and every branch of education not generally considered plain.

Vacations.

The usual vacations are, a month at Christmas, and at Midsummer; for either of which, if a pupil continues in the House, the charge is 5 guineas.

Mansion-House, Highgate,

4th Nov. 1817.

Faint, illegible text, possibly bleed-through from the reverse side of the page. The text is too light to transcribe accurately.

