An epitome of the natural history of the insects of China. Comprising figures and descriptions of upwards of one hundred new, singular, and beautiful species; together with some that are of importance in medicine, domestic economy, &c.; The figures are accurately drawn, engraved and coloured, from specimens of the insects; the descriptions are arranged according to the system of Linnæus; with references to the writings of Fabricius, and other systematic authors / By E. Donovan.

Contributors

Donovan, E. 1768-1837.

Publication/Creation

London: Printed for the author, by T. Bensley, 1798.

Persistent URL

https://wellcomecollection.org/works/ub7qpd77

License and attribution


This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.


You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org







44collater. J. J. Digitized by the Internet Archive in 2017 with funding from Wellcome Library

https://archive.org/details/b28758262

Sonovan (E) Gritome of the Natural Austory
of the Insects of China, original edition,
4 to, 50 beautifully wlowed places, 17.98.

Elf. 14. 0

S/r/:
14. ny. 83.

AN

EPITOME

OF THE

NATURAL HISTORY

OF THE

INSECTS OF CHINA:

COMPRISING

FIGURES AND DESCRIPTIONS

OF

UPWARDS OF ONE HUNDRED NEW, SINGULAR, AND BEAUTIFUL SPECIES; TOGETHER WITH SOME THAT ARE OF IMPORTANCE IN MEDICINE, DOMESTIC ECONOMY, &c.

THE FIGURES

ARE ACCURATELY DRAWN, ENGRAVED, AND COLOURED, FROM SPECIMENS OF THE INSECTS;

THE DESCRIPTIONS

ARE ARRANGED ACCORDING TO THE SYSTEM OF LINNÆUS; WITH REFERENCES TO THE WRITINGS OF FABRICIUS, AND OTHER SYSTEMATIC AUTHORS.

By E. DONOVAN,

AUTHOR OF THE NATURAL HISTORY OF BRITISH INSECTS, &C.

LONDON:

PRINTED FOR THE AUTHOR, BY T. BENSLEY;

AND SOLD BY WHITE, FLEET-STREET; FAULDER, BOND-STREET; BELL, OXFORD-STREET, &c.

1798.

HISTORICAL MESHOAL

Wello

ADVERTISEMENT.

The History, and Present State of China, as far as relates to its government, its sciences, and its arts, has long been considered worthy of the deepest research; and much information has been collected on these subjects, with considerable labour and expence, under the auspices of several of the most powerful courts in Europe. If the natural productions of that country have attracted less attention, it is only because their value and importance are less generally known; upon better acquaintance they would at once excite our astonishment, and convince us of their utility.

General readers are sometimes disposed to consider the Insect World as too minute and frivolous to deserve investigation. Let them remember, however, that to a foreign Insect the staple commodity of this kingdom is indebted for its richest dye; that from another we derive the most costly articles of dress, and splendid ornaments of luxury; and the utility of many others is apparent in various branches of medicine, the arts, and domestic economy.

Prompted by these reflections, the Author determined to submit to the Public a Series of Engravings illustrative of the Entomology of China.—But flattering as the prospect of encouragement to his project appeared, he would not presume to publish his Work till the result of the late Embassy of Earl Macartney to that country was fully known: of that the public are now in possession; and, though, in common with every friend to the commercial advantages and scientific inquiries of this country, the Author must regret its issue, it is perhaps, on the whole, more favourable to the present Publication than if the event had been different. If indeed a more general intercourse had been established between the two nations, and the language of China had been better understood, it is impossible to calculate the advantages which Entomology, amongst other sciences, might have derived; for the Chinese, like their neighbours the Japanese, are well acquainted with the natural productions of their empire, and Zoology and Botany, in particular, are favourite studies amongst them. To what degree of excellence they have arrived in their scientific researches we are not informed, but we must not affect to despise the instruction of a people amongst whom the most useful arts, and sciences, first dawned and acquired a high degree of perfection, when "Europe had scarcely a few savages scattered over her forests."

The few, but interesting hints, which Sir George Staunton has given on the practical Entomology of China, induces us to look forward to a period when some of the Insects, as well as Plants, of that vast empire may be no less objects of curiosity, than of national utility and importance; the Chinese Cochineal Insect³, and that from which the wax of the east is procured,

² Dr. Anderson has found eight species of Cocci at Madras. One of these, he says, was found on a young citron-tree, Citrus Sinensis, just landed from China; it was more deeply intersected between the abdominal rings than any of those of the coast, and he names it therefore C. Diacopeis.—Collection of Letters from Madras, Jan. 28, 1788.—The Cactus Cochinilifer has lately been found by Mr. Kincaid, at Canton; its Chinese name is Pau wang.—This has been transmitted to the Nopalry of the Hon.


are two species that deserve particular attention. The medical precepts of the Chinese will certainly find few votaries in Europe, but as articles of medicine, amongst others, the Meloe Cichorei, which were the cantharides of the ancients, and are now used by the Chinese, may be of importance, as it possesses more virtues than the Meloe Vesicatorius used in our pharmacy: the Curculio regalis, Buprestis vittata, and many others, are also employed in articles of jewellery in the eastern parts of the world, and may vie with the richest gems in beauty and splendour.


These observations we presumed to offer on the Entomology of China, in submitting the Design of this undertaking to the public. We have solicitously endeavoured to gratify the curiosity and expectation such observations were calculated to excite, and trust not altogether unsuccessfully. On the economical purposes of the Chinese Insects we can offer little except conjectures; those may, however, assist the inquiries of future observers; and the general reader will not be disposed to regard it with less favour, if novelty and beauty supply the deficiency of useful information. It embraces, in one view, a variety of the most uncommon and brilliant species of that fertile region, portrayed in an elegant and faithful manner; and classically arranged according to the favourite system of Linnæus. Thus, whilst our Epitome of the Chinese Insects exhibits a splendid display of this beauteous race, it may insensibly lead to a comprehensive survey of the system itself; and, by conveying instruction in its most pleasing form, facilitate the study of this charming, but much neglected science.


From its commencement, the Author has been encouraged by the liberal attention of several persons, whose names would reflect honour on this undertaking, were he at liberty to mention them. He has already observed that his own collection includes several thousand specimens, collected by amateurs of the first celebrity b. To this he may add, that every other collection he was desirous of consulting, has been kindly open to his inspection; and every information communicated with a readiness that merits his warmest thanks. Amongst these, he cannot refrain noticing the valuable collection of Drawings and MSS before alluded to; the specimens of Insects collected in the journey of his Excellency Earl MACARTNEY, in the late embassy to China; and the very magnificent collections of Mr. Francillon and Mr. Drury, from both of whom he had unreserved permission to figure and describe whatever his own cabinet could not furnish. And, finally, he must own himself particularly indebted to the favours of the Right Hon. Sir J. Banks, Bart. K. B. whose invaluable cabinet and library have afforded him every assistance in completing his design, and for which he begs leave to testify his most grateful acknowledgments.

East India Company, at Madras, and promises to be of future advantage to the commercial concerns of Great Britain. We have been unable to procure any of the Chinese Cochineal insects, and purposely omit that species which Sir G. Staunton has noticed, because it has no relation to the productions of China.

b The late Duchess Downger of Portland, - Tunstall, Esq. Governor Holford, Smeethman, Ellis, Keate, Yeats, Forster, Eaily, &c. &c.















* Scarabaus Molofsus.

: Scarabaus Seniculus.

: Scarabaus Bucephalus.

London Published as the Act directs by E. Denovan May 11798.

SCARABÆUS NASICORNIS.

GENERIC CHARACTER.

Antennæ terminate in a kind of club, divided longitudinally into laminæ. Second joint of the foremost pair of legs furnished with spines or teeth.

SPECIFIC CHARACTER.

A scutellum. Thorax armed with three prominences. Horn on the head recurved. Wing-cases smooth.

SCARABÆUS NASICORNIS: scutellatus thorace prominentia triplici, capitis cornu recurvo, elytris lævibus.

Linn. Syst. Nat.

Fab. Ent. Syst. 1. p. 14. 38.

The male of this species is furnished with a long recurved horn on the head; the female has only a small rising on that part. It is found in Europe as well as China.

SCARABÆUS SENICULUS.

SPECIFIC CHARACTER.

Without fcutellum. The anterior part of the thorax furnished with two horns, posterior of the head bidentated.

Scarabæus Seniculus: exécutellatus thorace antice clypeo postice bicorni. Fab. Ent. Syst. 1. p. 43. 142.

The annexed figures exhibit the two fexes of Scarabæus Seniculus. In fome specimens the spots are very indistinct and reddish, in others the wing-cases have faint red striæ. The semale has the rudiments of horns on the thorax.

SCARABÆUS MIDAS.

SPECIFIC CHARACTER.

Without fcutellum. Thorax armed with three horns: a horn on each fide of the head.

Scarabæus Midas: exfcutellatus thorace tricorni, capitis, clypeo finuato bicorni.

Fab. Ent. Syft. T. 1. p. 45. 148.

The figure of this rare species is taken from a specimen in the collection of Mr. Drury, of London, on affurance that it was received from China. Another, in the cabinet of Sir J. Banks, Bart. described by Fabricius, is noted from America.

The horns on the head of this infect have a very uncommon appearance, and which authors have compared to a pair of ears. The specific name Midas has been aptly given from this circumstance.

SCARABÆUS BUCEPHALUS.

SPECIFIC CHARACTER.

Without fcutellum. Thorax blunted in front, armed with four teeth or horns. Shield of the head angulated, furnished with a horn.

Scarabæus Bucephalus: exfcutellatus thorace retufo quadridentato, capitis clypeo angulato: cornu emarginato. Fab. Ent. Syft. T. 1. p. 51. 166.

This fpecies has been confounded with S. Moloffus. Both infects are given on the fame plate, that the difference may be precifely observed.

SCARABÆUS MOLOSSUS.

SPECIFIC CHARACTER.

Without feutellum. Thorax blunted, armed with two teeth, or horns: impressed on each side. Front of the head lunated; with a horn. Wing-cases smooth.

Scarabæus Molossus: exfcutellatus thorace retufo bidentato utrinque impresso, elypeo lunato unicorni integro, elytris lævibus. Linn. Syst. Nat. 2. 543. 3.

Fab. Ent. Syft. 1. p. 51. 167.

S. Moloffus and S. Bucephalus are very common in China. The first seems a local species, the latter is faid to be found in other parts of the East Indies. Olivier has given three varieties of Scarabæus Moloffus. The specimen figured in the annexed plate is the var. c. of that author.

The larvæ of the larger kinds of coleopterous infects, abounding in unctuous moifture, are not less efteemed as food among some modern nations, than they were by the epicures of antiquity. In Jamaica, and other islands in the West Indies, the Macokko a larva is an article of luxurious food; and in China most infects in that state are appropriated to the same purpose. Thus also the Romans introduced the larvæ of the Lucani b and Cerambyes c in their voluptuous repasts; previously feeding them on farinaceous substances to give consistence to the animal juices.

The learned author of the last account we have of China, says, "Under the roots of the canes is found a large white grub, which being fried in oil is eaten as a dainty by the Chinese." Perhaps this is the larva of Scarabæus Molossus, which, like many other of the Scarabæi, may live sedentary in the ground, and substitution the roots of plants: the general description and abundance of this insect in China favours such opinion. The same author observes, in another part of his work, that "the aurelias of the filk worm which is cultivated in China, after the filk is wound off, furnish an article for the table." This also is a very ancient custom among the Asiatics, and even Europeans, before the fixteenth century, if we may credit Aldrovandus: it is certain the worms, if not the aurelias, were administered in medicine in early ages.

2 Prionus damicornis. Fab. Ent. Syft.—Cerambyx damicornis. Linn. Mant.
b Stag beetles.
c Capricorn, or Goat beetles.

⁶ The larvæ of the Scarabæi live in the trunks of decayed trees, in putrid and filthy animal fubflances, or in the earth. The laft are the most injurious, because they destroy the roots of plants. All the known kinds of these larvæ are of an unwieldy form, and whitish colour, the skin free from hairs, and only the head and fore seet desended with a shelly covering.

e The German foldiers fometimes fry and eat filk worms. Aldrov.

f Silk worms dried, powdered, and put on the crown of the head, help the vertige and convulfions; mundify or cleanfe the blood, &c. &c. Schroderus, Serapio, &c. &c.

SCARABÆUS SACER.

SACRED BEETLE.

SPECIFIC CHARACTER.

No fcutellum. Front of the head divided into fix dentations. Thorax unarmed, margin crenulated.

Wing-cases smooth. Shanks of the posterior legs hairy.

Scarabæus Sacer: exfcutellatus clypeo fex dentato thorace inermi crenulato, tibiis pofticis ciliatis, elytris lævibus. Linn. Syft. Nat.

Fab. Ent. Syft. 1. p. 62. 205.

Scarabæus Sacer is a native of China; it is also found in other parts of the East Indies, in Egypt, Barbary, the Cape of Good Hope, and other countries of Africa, and throughout the fouth of Europe.

A few remains of ancient monuments, and fome fragments of historic information, preferved from an early period of the world, afford certain and interesting details of this inconfiderable creature. Those remains evince indeed but the first dawning of natural and moral philosophy on the human mind, but, connected with the history of the insect before us, are too important to be passed over in filence.

The Scarabæus was held in profound veneration by the people of Egypt. ^g They regarded it as a vifible deity; but a more refined fystem of religious worship prevailed in their temples among the priests and fages. ^h They deemed it only the symbol of their god, and, ascribing both sexes to the Scarab, it became a striking emblem of a self-created and supreme first cause. ⁱ

This infect was more especially the symbol of their god Neith, k whose attribute was power supreme in governing the works of creation, and whose glory was increased, rather than diminished, by the presence of a superior being, *Phtha*, the creator. The theological definition of the two powers being independent, yet centering in one spirit, is implied by the sigurative union of two sexes in the Scarab. In the latter sense it signified therefore but one omnipotent power. The Scarab, typisying Neith, was carved or painted on a

8 The species worshipped by the Egyptians is precisely noted by Linnæus. Scarabæus facer: and also by Olivier. Scarabé faceé.
"Cet insecte étoit autresois en veneration en Egypt." Oliv.

h Eschenbach, Jablonski, Savary, Se.

- i "The father, mother, male and female art thou." Synefius. Hymn. Phtha.—" The Egyptian spirit Phtha gave chaos form, and then created all things." Jamblichus de Mysteriis, sett. 8.
- k Nith. The disposer of all things, &c. Jablonski. "Sais," of the Delta, "the capital of its district, is a considerable city, of which Amasis was king. Neith, the Minerva of the Greeks, is the titular divinity." Plato in Timeo.—"On the door of the temple of Neith was engraved in hieroglyphics, "I am what is, what was, what shall be; mortal has never raised my veil; the sun is the fruit of my womb." Proclus Commentary on the Timeus of Plato, &c.—These passages demonstrate Neith and Phtha to be two attributes of one spirit. The third attribute is Cneph, or divine goodness. Savary.

ring, and worn by the foldiers, as a token of homage to that power who disposed of the fate of battles; I and sculptured on astronomical tables, or on columns, m it expressed the divine wisdom which regulates the universe and enlightens man.

- 1 Authors quote a doubtful paffage in Herapollo Hieroglyph. lib. 1. to fupport this opinion. That fuch rings were worn by the ancient Egyptians is beyond conjecture, many remains of them, and fome very perfect, have been found in the fubterranean caverns and fepulchres in the Plain of Mummies near Saccara and Giza. Those which we have examined, are remarkable for the convexity, or full relieve of the figure sculptured on them, in some it is of the natural fize of the infect, but generally smaller; the stone, cornelian, without a rim, and turning on a fwivel ring of gold.
- Innæus fays the Scarabæus facer is feulptured on the antique Egyptian columns in Rome. "Hie in columnis antiquis Romæ exfeulptus ab Ægyptiis." Syft. Nat. Does Linnæus allude to any remains of those colossal obelisks, which Augustus transported to Rome when he subjugated Egypt, or others of more recent date? It would increase the interest of our enquiries, to learn, that the Scarabæus was among the hieroglyphics, on the two very ancient obelisks, carried from Heliopolis, the city of the Sun.

The indefinite and visionary interpretations, imposed on most Egyptian hieroglyphics through a long feries of ages, will barely support a few conjectures on their original fignification. Those which related to local incidents, history, or the arts, are veiled in prosound obscurity. The phænomena of nature, and astronomical calculations, inscribed in those characters, are scarcely better understood, though the knowledge of those sciences have been in part handed down to us from the learned Egyptians in remote ages. We are informed by ancient writers, that the Scarabæus engraved on the astronomical tables of these people, implied the divine Wisdom which governed the motion and order of the celestial bodies; that those tables were huge and massly stones, or columns of granite, with the characters and figures, large, and highly embosses; in short, such as were supposed capable of long resistance to the corroding hand of time. Among those the Scarab was probably the most conspicuous, its size gigantic, and the figure frequently repeated; for this we have observed, even on small Egyptian antiques.

Various valuable remains of tablets, with figures of the Scarabans facer, are preferred in the British Museum and other collections of antiquities in this country. Those we have examined are of various descriptions, some smaller than the insect itself, others of a monstrous size. The stones on which they are sculptured generally green rephritic or jude flowe, or a kind of bafaltes, and black marble; the figure basso relieve, on a tablet or slab, but oftener in relieve, with the prominent characters of the insect very accurately defined, particularly the fix dentations of the clypeo, and those of the tibiæ. The reverse of the embossed side is flat and smooth, and abounds in characters altogether unknown, though, from the number of religious objects of worship occasionally interspersed, we may presume they contain an ample store of the ancient sacerdotal language; the most remarkable were the scarab, the sceptre and eye, the human sigure with a dog's head, the hawk, and the Ibis, or sacred bird. On the thorax of one sine specimen we remarked sour elegant sigures. One of them is holding a cornne-pia in the left hand, and a branch in the right: this is perhaps a subordinate deity of the Nile, that river having been once sound depictured on an antique Alexandrian coin, like an aged man, holding the cornucopia, and a branch of the Papyrus; denoting its abundance and produce.

The digreffion on the mythological hiftory of this infect may be confidered by fome as a tedious deviation from the purfuit of the naturalit; with others we truft it will be more favourably received; for it proves to the unprejudiced mind how deeply the hiftory of nature, and in the prefent inflance the science of Entomology, involves a most important enquiry into the first philosophical opinions of the human race. The means, however trifling, must not be contemned, which illumine the most fublime of all human researches,—The Study of Mankind.

a Cijris, or the fun. b Anabis. c Herus, a famous deity, had three cities dedicated, called by the Greeks the cities of Apollo.
e The mammies of these birds are found in urns, in the fubterranean galleries called the well of birds near Samura, supposed from the fize to be the Tantalus
This of Linearus.

SCARABÆUS LEEL

SPECIFIC CHARACTER.

Without fcutellum. Obfcure copper colour. Under fide black. Shield of the head margined.

Scarabæus Leei: exfcutellatus obfcure cupreus fubtus niger, clypeo emarginato.

Fab. Ent. Syft. 1. 65, 215.

SCARABÆUS CINCTUS.

SPECIFIC CHARACTER.

Without fcutellum. Unarmed, black. Margin of the clytra pale. Shield of the head margined.

Scarabæus cinctus: exfcutellatus muticus niger elytrorum margine pallido; elypeo emarginato.

Fab. Ent. Syll. 1. 69. 231.

CETONIA CHINENSIS.

SPECIFIC CHARACTER.

Shining green and gold colour. Shield of the head margined, with two fpines. Posterior part of the thorax lobed. Wing-cases terminate in an acute spine.

CETONIA CHINENSIS: ænea clypeo emarginato subspinoso, thorace postice lobato, elytris acuminatis.

Fab. Ent. Syst. 1. p. 2. 126. 6.

This, and the next species, are described by Fabricius, in the two new genera, Cetonia and Melolontha. They are Scarabæi of the Linnæan system. Cetonia Chinensis is a fine and very rare insect.

MELOLONTHA VIRIDIS.

SPECIFIC CHARACTER.

Above green. Beneath golden colour, or bronze.

MELOLONTHA VIRIDIS: glabra fupra viridis fubtus aurea. Fab. Ent. Syft. 1. p. 2. 160. 23.

Oliv. Inf. tab. 21. fig. 6. tab. 3. 18.

This fpecimen nearly corresponds with one of the Melolontha viridis figured by Olivier, which came from the Cape of Good Hope: our insect is undoubtedly from China.










: Cetonia Chinensis.

. Melolontha viridis.







CURCULIO CHINENSIS.

GENERIC CHARACTER.

Antennæ club-fhaped, and inferted in a horny probofcis or fnout.

SPECIFIC CHARACTER.

Beak long. Thighs dentated. Body covered with yellowish powder, except a stripe of black on the sides of the thorax and wing cases: a spine on each side of the beak, at the apex.

Curculio Chinensis: longiroftris, femoribus dentatis 3, corpore polline flavescente obtecto, lateribus nigris, roftro utrinque spinoso.

This infect feems nearly allied to Curculio mucoreus, an Indian species, described by Linnæus, but not figured by any author: the lateral stripe of black; and the denticulations on the posterior thighs of our infect clearly removes it, however, from the Linnæan species. Among the Curculiones described since by Fabricius, we have not discovered any with which our species can be confounded; and deeming it a non-descript, we name it specifically Chinensis; a name, perhaps, too local, but expressive of the native place of our specimen; and not applied by Fabricius, in his last work, to any insect of the same genus, though that author has exhausted almost every applicable name, in the descriptions of more than four hundred of its species.

The only specimen of this curious insect we have seen, is in the possession of Mr. Drury of London, from whose extensive collection we have been liberally permitted to copy, and describe, such Insects as we considered worthy of noticing in this work.

Except the lateral black ftripes, and the roftrum, Curculio Chinenfis is totally covered with a bright brown powder, or rather, with very minute hairs which adhere but flightly, and refemble that fubfiance. We observe a fimilar farinaceous appearance on the Curculiones, Lasten, Niveus, &c. and especially on that gigantic beetle Scarabaus Elephan.

CURCULIO LONGIPES.

SPECIFIC CHARACTER.

General colour blackish brown. Wing-cases ferruginous. Beak long, margined with small irregular tubercles on each side, and at the apex. Anterior legs longest.

Curculio longipes: longiroftris nigricans elytris ferrugineis, roftro emarginato, pedibus anticis longioribus. Fab. Ent. Syft. 1. p. 2. 395. 4.

Oliv. fig. 191.

All the specimens of this Curculio that have come under our inspection, are natives of China. Fabricius says it is from the Cape of Good Hope b.

CURCULIO BARBIROSTRIS.

SPECIFIC CHARACTER.

Entirely black. Beak long, and bearded. Three spines on the anterior shank of the legs.

Curculio Barbirostris: longiroftris niger roftro barbato, tibiis anticis tridentatis. Fab. Ent. Syft. 1.

Uncommonly fcarce. Fabricius refers for this infect to the collection of Cramer. We have only feen the fpecimen from which our figure is taken.

b The inaccuracy of this remark of Fabricius will appear by comparing the descriptions and fynonyms of Curculio longipes, in his Species Insection, with the same species in his Entomologia Systematica. In the first he describes Curculio longipes, quotes fig. 3. tab. 33. vol. 2. of Mr. Drury's work, and gives its habitat, Cape of Good Hope. In his last work he gives the same description, changes the reference from Mr. Drury's work to that of Olivier, fig. 191; but continues the other references, by which the Chinese species is probably unintentionally left as a native of Africa.

CURCULIO VERRUCOSUS.

SPECIFIC CHARACTER.

Snout fhort, colour throughout black, with a faint braffy gloss. Wing-cases beset with striæ of elevated points, and terminated at the apex in a single obtuse protuberance.

Curculio verrucosus: brevirostris aeneo niger elevato punctatus elytris postice verrucosis. Linn. Syst.

Nat. 2, 618. 90.

Fab. Ent. Syst. 1. p. 2. 478. 353.

This can be no other than Curculio verrucofus, though our specimen came from China, and that noticed by Fabricius from the Cape of Good Hope. The three elevated striæ of warts on each of the wing cases, and the intermediate double row of impressed points, are decisive in our opinion of this species.—

We have another that is mottled with clay colour, which is probably found in Africa.

CURCULIO SQUAMOSUS.

SPECIFIC CHARACTER.

Probofcis fhort. Entirely covered with green fcales. Impressed lines on the thorax and probofcis.

Curculio squamosus: breviroftris viridi fquamofus roftro thoraceque canaliculatis. Fab. Ent. Syft. 1.

p. 2. 452. 244.

A fmall, but fuperb fpecies, being totally covered with minute scales of an oblong form, and resplendent green colour, interspersed with changeable sparks of gold and crimson, in various reslections of light. This scaly covering is not unlike that of Curculio argentatus sound in England; but of a brilliance scarcely inferior to the gem-like spangles on the Curculio imperialis of Brazil.—Curculio squamosus is represented in its natural size, in the annexed plate; and, in justice to an insect of such uncommon beauty, an additional plate is given, to exhibit its appearance in the opaque microscope.—It is extremely common in China.

CURCULIO PULVERULENTUS.

SPECIFIC CHARACTER.

Entirely greyish. Head flat, and marked with longitudinal lines: an impressed canal down the middle.

CURCULIO PULVERULENTUS: brevirostris cinereus capite plano canaliculato. Fab. Ent. Syst. 1. p. 2. 452. 243.

Like Curculia squamosus this insect is entirely black, when divested of its scaly covering. It has been supposed a variety of C. squamosus, but Fabricius considers them as two species, and we are of the same opinion.

CURCULIO PERLATUS.

SPECIFIC CHARACTER.

Black. Abdomen white, with elevated shining black spots. Beak short and surrowed.


Curculio perlatus: brevirostris niger abdomine albo: punctis elevatis glabris atris, rostro sulcato.


Fab. Ent. Syst. 1. p. 2. 459. 271.

Neither this nor the two preceding species have been figured in any work. We have given a figure of the under side of this insect, to illustrate the specific character given by Fabricius.



Curculio pulverulentus.







Cerambyx Rubus.* Cerambyx Reticulator::

Cerambyx Farinosus **

London Published as the Act directs by E. Donovan Apls. 1798.

CERAMBYX FARINOSUS.

GENERIC CHARACTER.

Articulations of the antennæ gradually decrease in fize towards the end. Thorax spined or gibbous.

Elytra narrow, and of equal length throughout.

SPECIFIC CHARACTER

AND

SYNONYMS.

Thorax spined. Shells spotted with a white farinaceous substance. Antennæ longer than the body.

CERAMBYX FARINOSUS. Linn. Syft. Nat. 2. 626. 24.

Fab. Spec. Inf. 1. p. 214. 25.

Stenocorus Farinofus: thorace spinosa piceus elytris punctis sparsis farinosis, antennis longis.

Fab. Ent. Syft. 1. 295. 11.

CERAMBYX CHINENSIS. Forft. N. Sp. Cent. 1. 39.

Oliv. Inf. 66. tab. 7. fig. 46.

Among fome Chinese drawings of the late Mr. Bradshaw, we observed one on which the metamorphosis of this Cerambyx seemed to be delineated. The larva was partly concealed in the hollow of a piece of decayed wood; it was of a whitish colour, with the head and tail black, as described by Fabricius.

CERAMBYX RUBUS.

GENERIC CHARACTER.

Articulations of the antennæ gradually decrease in fize towards the end. Thorax spined or gibbous.

Elytra narrow, and of equal length throughout.

SPECIFIC CHARACTER

AND

SYNONYMS.

Thorax fpined, bimaculated. Base of the shells studded with small protuberances. Apex of the shells terminate in a sharp point.

CERAMBYX RUBUS. Linn. Syst. Nat. 2. 625. 21.

LAMIA RUBUS: thorace fpinoso bimaculato, elytris basis spiceque mucronatis.

Fab. Ent. Syst. 2. 290. 89.

Oliv. Inf. 67. tab. 7. sig. 57.

This species, we find, is not uncommon in China. It is the largest of its genus that we are acquainted with from that country.

CERAMBYX RETICULATOR.

GENERIC CHARACTER.

Articulations of the antennæ gradually decrease in fize towards the end. Thorax gibbous or spined.

Elytra narrow, and of equal length throughout.

SPECIFIC CHARACTER

AND


SYNONYMS.

Thorax fpined. Thorax and fhells yellow brown; the first marked with longitudinal black streaks, the latter reticulated with the same colour.

Lamia Reticulator: thorace fpinoso nigra thorace elytrisque sulvis, thorace nigro lineato, elytris reticulatis. Fab. Ent. Syst. 1. p. 2. 278 44

Oliv. Inf. 67. tab. 12. fig. 85.

Cerambyx reticulator is a very rare species; we have never seen more than two specimens of it. It is altogether a beautiful insect; but the singular structure of the antennæ deserves particular notice: it is entirely brown except the first articulation, which is black; the second has a large verticillated tust of black hair at the summit; at the base of this articulation it has another tust, but smaller; and a similar tust, but still smaller, is situated on the third and sourth articulations.





London Published as the Act directs by E. Donoran Mary 1. 1798.

BUPRESTIS VITTATA.

GOLDEN-STRIPE BUPRESTIS.

GENERIC CHARACTER.

Antennæ fetaceous, as long as the thorax. Head half retracted, or drawn back within the thorax.

SPECIFIC CHARACTER.

Elytra impressed with points or spots: marked with four longitudinal ridges, and terminated in two teeth or spines; a ribbon-like stripe of golden yellow, down each elytron.

Buprestis Vittata: elytris bidentatis punctatis: lineis quatuor elevatis viridi æneis vittaque lata aurea.

Fab. Ent. Syft. 1. p. 2. 186. 5.

Bupreste Bande-dorée. Oliv. Inf. 32. tab. 3. fig. 17.

The Buprestides form an extensive, and most brilliant tribe of coleopterous insects. Brasil and New Holland produce some gigantic species, but none more beautiful than those of India. We need adduce no other proof of this, than Buprestis chrysis, sternicorus, attenuata, ocellata, and vittata. These wrought into various devices, and trinkets, decorate the dresses of the natives in many parts of India. The Buprestis vittata in particular, is much admired among them. It is, we believe, entirely peculiar to China, where it is found in vast abundance, and distributed from thence at a low price among the other Indians. The Chinese, who always profit by the curiosity of Europeans, collect vast quantities of this Buprestis, and other gay insects, in the interior of the country, and traffic with them.

A confiderable error feems to have arisen concerning this species, and the true Buprestis ignita of Linnæus. All authors, except Fabricius and Olivier, have confidered the Buprestis vittata, and ignita, the same. Fabricius, in his Species Insectorum, refers to Mus. Dom. Banks only, for the Buprestis vittata; and to the 14th figure, plate 6, in Sulzer's work, for a figure of Buprestis ignita. In the Entomologia Systematica we however find the same reference to the figure of Sulzer, for B. vittata; and, to increase the perplexity, precisely the same reference under that of B. ignita also.

^{*} A pair of slippers embroidered with little pieces of the wing cases of this insect is preserved in the Leverian Museum

⁷ Vol. I. part 2. p. 186. 5.

⁼ Vol. I. part 2. p. 192. 28.

The Buprestis ignita of Linnæus, we imagined, was unknown in the cabinets of the curious in this country, till we discovered an insect nearly corresponding in character with it, in the collection of Mr. Drury, and which we presume is the B. ignita, or at least a variety of it. It has not the brilliance of colours that so eminently distinguishes B. vittata, but in form and fize it agrees with it. The only figure of that species is given by Olivier, from a specimen formerly in the cabinet of Gigot d'Orcy, of Paris. We have examined the specimen in the cabinet of Sir J. Banks, referred to by Fabricius as B. vittata, and find the figure in Sulzer is of that species, as well as the specimen we have represented.

Fabricius has given as a part of the specific distinction of these insects, that B. ignita has three spines at the end of each wing case, or elytron, and B. vittata no more than two. This may form a sufficient characteristic in those species; but we must remark, that it is not so in Buprestis occilata. We have two specimens that have two spines at the end of each elytron, and another with three, as Fabricius has described it. We also find several insects nearly allied to B. vittata, the stripe of gold on each side excepted; one of these has fix teeth, another four teeth, and a third only two.

The Buprestides are supposed, for the most part, to undergo their transformations in the water, or marshy ground.

Canna Indica.—Indian flowering Reed.

This plant is common in China, and is found also in most other parts of Asia, Africa, and America. In our climate it requires to be placed in the stove, where it produces an abundant succession of flowers throughout the summer. It bears a berry which is perfectly hard and round, of a black colour, and highly polished. It is called Indian Shot.

BUPRESTIS OCELLATA.

OCELLATED BUPRESTIS.

SPECIFIC CHARACTER

AND

SYNONYMS.

Shining green. Elytra terminate in three points: a large round yellow fpot in the middle of each elytron, and two golden-red fpots, one above and the other below it.

Buprestis ocellaria viridi nitens elytris tridentatis: maculis duabus aureis ocellarique flava. Fab. Ent.

Syft. 4. p. 2. 193. 30.

De Geer Inf. 7. 633. 30. tab. 47. fig. 12.

Bupreste Oculé. Oliv. Inf. 32. tab. 2. fig. 3.


Sulz. Hift. Inf. tab. 6. fig. 15.

The Buprestis occilata is a native of China, and is very rare: specimens of it have been found in parcels of the common fort. Olivier says it is from Chandernagore in the East Indies. Mr. Drury has an extraordinary variety of this insect from China, in which the two spots unite at the suture so as to form only one large spot on the back when the wing cases are closed.

These spots on the wing cases are strikingly characteristic of this species; at least we have not obferved them in any other of the Buprestis genus. They are usually situated in the centre of each elytron; are somewhat pellucid, and in fine specimens are cream colour, surrounded with a crimson circle. These spots are sometimes brown; probably they become so after the insect dies.

13

Our figures represent this infect with expanded wings; one of those is designed to exhibit the beautiful appearance of the under furface, particularly the effulgent abdomen and purple colour of the interior part of the shells: this is seen hovering over the flower of the Canna Indica: the other represents the upper furface.







Mylabris Cichorei: : Sagra femorata.

TENEBRIO FEMORATUS.

GENERIC CHARACTER.

Antennæ like a ftring of beads, extreme articulation roundish. Thorax plano-convex, margined. Head porrected. Elytra rather stiff.

SPECIFIC CHARACTER.

Greenish copper colour. Thighs and shanks of the posterior legs dentated.

SAGRA FEMORATA: viridi ænea femoribus tibiifque posticis dentatis. Fab. Ent. Syst. T. 1. p. 51, 55. 1.

This infect may be referred to the Linnæan genus Tenebrio. It is described by Fabricius in the Species Insectorum, in the new genus Alurnus; in the Entomologia Systematica of the same author, it is removed to another new genus, Sagra, and its former specific name, semoratus, necessarily altered to semorata. As the characters of these Fabrician genera are taken from the palpi, maxilla, and labium, we prefer the more obvious characters of the Linnæan genus.

The figures quoted by Fabricius differ, in some respects, from the Chinese specimens. Sulzer reprefents it of a green colour, without a tinge of red purple, or copper colour, so predominant in every specimen we have seen. Our insects are of two kinds: one is of a glowing purple, resplendent as metal, and changeable to green or yellow; the other is purple also, but not quite so vivid, being tinged with green.

MELOE CICHOREI.

GENERIC CHARACTER.

Antennæ moniliform: last joint oblong. Thorax roundish. Elytra soft and slexible. Head inflected and gibbous.

SPECIFIC CHARACTER.

Black. Wing-cases yellow, with three transverse black bands.

Meloe cichorei: nigra elytris flavis: fasciis tribus nigris. Linn. Syst. Nat. 2, 680, 5.

Mylabris Cichorei. Fab. Ent. Syst. T. 1. p. 2, 88, 2.


This infect is very common in China, and some other parts of the East Indies. The small specimen is


rare, but is, probably, only a variety of the other fort. According to Olivier, the Meloe cichorei is used by the Chinese in their medical preparations instead of the Cantharis vesicatorius of the Europeans, and is supposed to be more efficacious in certain cases. The same author quotes a passage in Dioscoride Mat. Med. Lib. 2. to prove that it is also the Cantharides of the ancients i.

i "Les Cantharides des anciens et celle des Chinois ne font pas les mêmes que celles des Européens. Les Chinois employient le Mylabre de la Chicorée, &c. &c."—"The Cantharides of the ancients, and those of the Chinese, are not the same as
ours. The Chinese employ the Mylabre de la Cichorei, and it appears from Dioscoride Mat. Med. Lib. 2. Cap. 65, the ancient
Cantharides were the same as those now used by the Chinese." "The most efficacious sort of Cantharides," says Dioscoride,
"are of many colours, having yellow transverse bands; the body oblong, big, and fat; those of only one colour are without
strength." The description Dioscoride has given, does not agree with our species of Cantharides, as they are of a fine green
colour, but is more applicable to the Mylabre de la Cichorei, which is very common in the country where Dioscoride lived.
Olivier's Entomologie, ou Hist. Nat. des Insectes. Vol. I. Introd.

The Cantharides of the ancients, are by no means to be confounded with those of medical writers in the last century. By the term Cantharides, in an European Pharmacopeia, we understand the Meloe vesicatorius * of Linnæus, an insect whose medicinal properties are very generally known †. The Cantharides of the ancients can scarcely be ascertained; it was a term indiscriminately applied to several kinds of insects, and too often without regard to their physical virtues. Pliny speaks of the Cantharis as a small beetle that eats and consumes corn; and of another that breeds in the tops of ashes and wild olives, and shines like gold. The ancients were certainly well acquainted with our common fort, though it is consounded with others in a general appellation ‡. Hippocrates, Galen, Pliny, Matthielus, and other physical writers of antiquity, treat of the medicinal uses of Cantharides; but it is not clear that they alluded to only one species §.

- * Geoffroy calls this a Cantharis. The Linnæan Cantharis is a diftinct genus.
- + Applied externally to raife blifters. It is a violent poifon taken inwardly, except in fmall portions.
- 2 The common sort has been called Musea Hispanica by some Latin authors, and hence Spanish fly by Boyle.
- § Olivier endeavours to prove that the Mylabris Cichorei is the ancient Cantharides; the authority of his opinion is credible, and the inference natural, if not conclusive. But if it were in use, so also might the common fort, for Dioscoride, whom he quotes, mentions those of only one colour also. The ancients often consounded the term Scarabæus with Cantharis; but whether because they knew that the common kinds of Scarabæi produce the same effects as the Cantharis, is uncertain.—The Scarabæus auratus, and Meloloutha, several Coccinellæ, Cimen nigro-lineatus, Se. Se. have a place in the Materia Medica as Cantharides.





Mantis Oculata. Mantis Flabellicomis:

MANTIS OCULATA.

CORNUTED - EYE CAMEL CRICKET.

GENERIC CHARACTER.

Head flightly attached to the thorax, unfteady: the mouth armed with jaws and palpi. Antennæ fetaceous.

SPECIFIC CHARACTER

AND

SYNONYMS.

Thorax filiform, or like a thread, triangular. Eyes oblong, prominent, and terminated each in a fharp spine.

Mantis Oculata thorace triangulo filiformi, oculis oblongis porrectis acuminato fpinofis. Fab. Ent. Syft.

T. 2. 19. 26.

Mantis bicornis thorace lævi, capite bipartito fubulato. Linn. Syft. Nat. 2. 691. 11.

Two figures of the Mantis, very much refembling our species, is given in the work of Stoll on Cicadæ, &c. one kind he calls La Mante ètroitement cornue, the other La Mante Chinoise ètroites cornue. The first is from the coast of Coromandel and Tranquebar, the other, as its name implies, is a Chinese insect. We cannot discover any material difference between these figures and our specimen, and are inclined to consider them altogether as one species.

Few of the Cicadæ and Locustæ described by Fabricius, in his Species Insectorum, have any reference to figures, because a very small number of them had been figured till Stoll published his work on those genera. Stoll has occasion to mention with regret, that Fabricius has scarcely noticed any of his plates. It is a considerable disadvantage to the works of that author, as well as to the naturalist that consults them, that no scientific names, or definitions, are given to the figures of many rare insects included amongst them.

The Mantis Oculata of Fabricius is an African insect, and was described from the collection of the Right Hon. Sir J. Banks, Bart. we have compared our Chinese specimen with it, and find it is precisely the same species.

MANTIS FLABELLICORNIS.

FAN-HORNED CAMEL CRICKET.

SPECIFIC CHARACTER.

A broad membrane on each fide of the thorax, next the head. Anterior thighs terminate in a spine, the second joint beset with spines, and surnished with a lobe on one fide. Antennæ pectinated.

Mantis flabellicornis: thorace apice dilato membranaceo, femoribus anticis spina terminatis reliquis lobo, antennis pectinatis. Fab. Ent. Syst. 2. p. 16. sp. 16.

This Mantis is described by Fabricius only. Stoll has given the figure of an insect not unlike it in his publication; and we have seen a specimen similar to it, which was found by Professor Pallas near the Caspian sea. It is allied to Mantis Gongyloides, a native of Africa and Asia, but bears a closer affinity to Mantis Pauperata a from Java, Molucca, and perhaps other islands in the Indian sea.

Fabricius enumerates fifty-one species of this genus in his last system; a considerable portion of these are from Asia: had he included the lately discovered kinds in America and New Holland, his genus would have been far more comprehensive. Few naturalists have had the opportunity of observing the manners of these creatures in distant countries; nor can we always rely on the information those sew have given. Of the European species we can speak with more precision, because some indefatigable naturalists have attended minutely to them; Roesel in particular has treated at considerable length on the manners of the Mantis Religiosa of Linnæus.

Descriptions can only convey an impersect idea of the extraordinary appearance of many creatures included in the Mantis and Locusta genera. Among them are found species that bear a similitude to the usual forms of other insects; but, from these we almost imperceptibly descend to others, bearing as strong a similitude to the vegetable part of creation; seeming as if Nature designed them to unite the appearance of a vegetable with the vital functions of an animal, to preserve them from the ravages of voracious creatures, or to connect that chain of progressive and universal being, which

" The great directing MIND of ALL ordains."

Many of these creatures assume so exactly the appearance of the leaves of different trees, that they fur-

2 Figured by Stoll under the name of La Mante Gouttenfe Brune?

nish the entomologist with unerring specific distinctions; thus we have L. citrifolia, laurifolia, myrtifolia, olcifolia, graminifolia, and others, equally expressive of their resemblance in form, and colours, to the leaves of those respective plants. Travellers in countries that produce these creatures, have been struck with the phænomenon, as it must appear, of animated vegetable substances; for the manners of the Mantis, in addition to its structure, are very likely to impose on the senses of the uninformed. They often remain on the trees for hours without motion, then suddenly spring into the air, and, when they settle, again appear lifeless. These are only stratagems to deceive the more cautious insects which it feeds upon; but some travellers who have observed it, have declared they saw the leaves of those trees become living creatures, and take slight.

M. Merian informs us of a fimilar opinion among the Indians, who believed these insects grew like leaves on the trees, and when they were mature, loosened themselves and crawled, or flew, away. From the credulous, and unscientific, marvellous reports of such extraordinary creatures may be expected; but, to these we must add the authority of a naturalist, whose works are a valuable addition to our present knowledge of the history of nature: these are the works of Piso b.

"Those little animals," says that author, "change into a green and tender plant, which is of two hands breadth. The feet are fixed into the ground first; from these, when necessary humidity is attracted, roots grow out, and strike into the ground; thus they change by degrees, and in a short time become a perfect plant. Sometimes only the lower part takes the nature and form of a plant, while the upper part remains as before, living and moveable: after some time the animal is gradually converted into a plant. In this Nature seems to operate in a circle, by a continual retrograde motion."

Has the father of inventive romance outdone this account of Pifo in his well-known extravagant poetical effusions c? It may be imagined he has not; but before we difmifs his account with a hafty

b Gulielmi Pifonis, Amfterl. 1657.

Luna quater junctis implerat cornibus orbem, &c. Ovidii Metamorph. Sorores Phaethontis in Arbores.

Four times, revolving, the full moon return'd,
So long the mother and the daughters mourn'd;
When now the eldeft, Phaëthufia, ftrove
To reft her weary limbs, but could not move;
Lampetia would have help'd her, but she found
Herself withheld, and rooted to the ground:
A third in wild affliction, as she grieves,
Would rend her hair, but fills her hands with leaves:
One sees her thighs transform'd, another views
Her arms shot out, and branching into boughs.
And now their legs, and breasts, and bodies, stood
Crusted with bark, and hardening into wood;
But still above were semale heads display'd,
And mouths, that call'd the mother to their aid.

What could, alas! the weeping mother do?

From this to that with eager hafte she flew,

And kiss'd her sprouting daughters as they grew:

She tears the bark that to each body cleaves,

And from the verdant singers strips the leaves:

The blood came trickling where she tore away

The leaves and bark: the maids were heard to say,

"Forbear, mistaken parent, oh, forbear!

A wounded daughter in each tree you tear:

Farewel for ever." Here the bark increas'd,

Clos'd on their faces, and their words suppress'd.

Addison's Tranf. Phaeton's Sifters transformed into Trees.

cenfure, let us attend to the opinion of a few authors, nearly contemporary with himfelf: we find these collected in the criticisms of Roesel on that passage.

From these it appears that the works of Piso were much admired when first published, but we rely less implicitly on the information it contains, than his readers in the last century. Roesel treats his account with more than merited feverity; not because he could contradict the relation of Piso, but, because he had never observed the same circumstance attend the Wandering Leaf, or Mantis Oratoria, in Europe 4. This reasoning is neither so conclusive, or liberal, as we should expect from Roesel; and more especially as he afterwards describes even the first symptom of the transformation as related by Pifo. When he speaks of the death of the European species his words are, "As their dissolution approaches, their green eyes become brown, and they unavoidably lofe their fight: they remain a long while on the fame fpot, till at laft they fall quite exhausted and powerless, as if asleep." What is this but substantiating part of the evidence of Pifo, which he has laboured before to difcredit? As to the change after they remained long on the ground, fuch as fending forth fibres, roots, and ftems, from the body of the infect, it is only aftonishing fuch a well-informed naturalist should have deemed it matter of surprise. Could be be ignorant of the many inflances that occur, of animal fubflances producing plants ? or was he not informed that the pupa which commonly fends forth a bee, a wasp, or cicada, has sometimes become the nidus of a plant, thrown up ftems from the fore part of the head, and changed in every respect into a vegetable, though still retaining the shell and exterior appearance of the parent insect at the root f? We own at first fight with Roefel that the account of Pifo feems "an inattentive and confounded observation," but that an insect may firike root into the earth, and, from the co-operation of heat, and moisture, congenial to vegetation, produce a plant of the cryptogamia kind, cannot be disputed. We have seen species of clavaria both of the undivided and branched kinds, four times larger than the infect from which they fprang; and can we then deny that the infect mentioned by Pifo might not produce a plant of a proportionate magnitude? In fhort, are we fo well acquainted with the productions of Brafil as to contradict any of his affertions, concerning this transformation? Pifo does not fay of what kind this vegetable was; it must furely be of the fungi

We quote this part of the poem at length; for as we peruse it, every successive line bears a stronger similitude to the wonderful transformation of the Mantis as related by Piso; we might be almost tempted to condemn the description of the naturalist as a service copy of Ovid's verses, if the similar transformation of other insects above noticed, had not occurred to the knowledge of every entomologist.

- d Among the annotations on the last edition of Rocsel's Infecten Bellustigung we find one relating to this part of the works of Piso.

 44 Der feel Her geheime Rath Trew, &c. Counf. Trew affures Mr. Roesel that Piso not only very often gave out the credible observations of others, as his own, but himself believed the most incredible relations, and pretended to be an eye witness thereof." We quote this in justice to the remarks of Roesel. Note in page 10, sellion Das Wandlende Blat.
 - e Such as Mucor crustaceus, &c.

f Specimens of these vegetated animals, are frequently brought from the West Indies; we have one of the cicada from the pupa, as well as others produced from wasps and bees in the perfect or winged state. Mr. Drury has a beetle in the perfect state, from every part of which, small stalks and sibres have sprouted forth; they are very different from the tusts of hair that are observed on a sew coleopterous insects, such as the Buprestis sascicularis, of the Cape of Good Hope; they are certainly a vegetable production.

kind: reasoning then from analogy, it might be an unknown species of *clavaria* with numerous and spreading branches; and, finally, the colour of his plant, on which authors lay much stress, might be green, though a colour not so predominant in that tribe of vegetables as some others.

The largest and most interesting of the Indian species of Mantis is found in the isle of Amboyna. Stoll contradicts the account of Renard s, who says these creatures are sometimes thirteen inches in length; but we have a specimen almost of that size. It is related by Renard, and others, that the larger kinds of Mantes go in vast troops, cross hills, rivers, and other obstacles that oppose their march, when they are in quest of food. If they substituted entirely on vegetables, a troop of these voracious creatures would desolate the land in their excursions; but they prefer insects, and clear the earth of myriads that insect it: if these become fearce from their ravages, they sight and devour one another. When they attack the plants, they do great mischies. It is said of some Locusts and Mantes that the plants they bite wither, and appear as if secrebed with fire: we have not heard of this pestilential property in any of the larger species of Mantes.

Of the smaller kinds, the Mantis Oratoria is the most widely disfused, being found in Africa and Asia as well as in all the warmer parts of Europe. These creatures are esteemed sacred by the vulgar in many countries, from their devout or supplicating posture. The Africans worship them; and their trivial names in many European languages imply a superstitious respect for them h.

England produces no species of this tribe. The entomologists in this country must consequently rely on the accounts of those, who have observed them in other parts of the world. We shall select a few remarks from Roefel's extensive description of Mantis Oratoria and Gongyloides, because, if we may prefume from the analogy they bear in form to Mantis Flabellicornis, the history of one will clearly elucidate that of the other.

Rocfel fays, fome of the Mantes are local in Germany; they are found chiefly in the vintages at Moedting in Moravia, where they are called Weinhandel i. The males die in October, the females foon after k. The young brood are preferved in the egg ftate, in a kind of oblong bag, of a thick fpongy fubftance; this bag is imbricated on the outfide; it is fastened lengthwise to the branch of some plant i. As the eggs ripen they are protruded through the thick substance of the bag, and the larva, which are about half an inch in length, burst from them. Roesel, wishing to observe the gradual progress of these creatures, to the winged state, placed the bag containing the eggs in a large glass, which he closed, to prevent their escape. From the time they were first hatched they exhibited marks of a savage disposition. He put different sorts of plants into the glass, but they refused them, to prey on one another: this determined him to supply them with other infects to eat: he put ants into the glass to them, but they then betrayed as much cow-

⁸ Poissons des Molucques par M. Renard, Amsterl. 1754. h Louva Dios by the Portuguese. Presque Dieu by the French.

Probably a provincial term for a dealer in wine. k Goetz, in his Beytrage, observes, that they live sometimes ten years.


1 To that of the vine in Mantis Oratoria.

ardice, as they had barbarity before; for the inftant the Mantes faw the ants they tried to escape in every direction. By this Roesel found the ants were the greatest persecutors of the Mantes. He next gave them some of the common musca (house slies), which they seized with eagerness in their fore claws, and tore in pieces: but, though these creatures seemed very fond of the slies, they continued to destroy one another through savage wantonness. Despairing at last, from their daily decrease, of rearing any to the winged state, he separated them into small parcels in different glasses; but, here, as before, the strongest of each community destroyed the rest.

Another time, he received feveral pair of Mantes in the winged state; profiting by his former observation, he put each pair [a male and semale] into a separate glass, but they still shewed signs of an eternal enmity towards one another, which neither sex nor age could soften; for the instant they were in sight of each other, they threw up their heads, brandished their fore legs, and waited the attack: they did not remain long in this posture, for the boldest throwing open its wings, with the velocity of lightning, rushed at the other, and often tore it in pieces with the crockets and spines of the fore claws. Roefel compares the attack of these creatures to that of two hussars; for they dexterously guard and cut with the edge of the fore claws, as those soldiers do with their sabres, and sometimes at a stroke, one cleaves the other through, or severs its head from the thorax. After this the conqueror devours his vanquished antagonist.

We learn from Roefel alfo, the manner in which this creature takes its prey, in which refpect we find it agrees with what is related of the extra European species. The patience of this Mantis is remarkable, and the posture to which superstition has attributed devotion, is no other, than the means it uses to catch it. When it has fixed its eyes on an insect, it very rarely loses sight of it, though it may cost some hours to take. If it sees the insect a little beyond its reach, over its head, it slowly erects its long thorax, by means of the moveable membranes that connect it to the body at the base; then, resting on the four posterior legs, it gradually raises the anterior pair also; if this brings it near enough to the insect, it throws open the last joint, or crocket part, and snaps it between the spines, that are set in rows on the second joint. If it is unsurfacesful, it does not retract its arms, but holds them stretched out, and waits again till the insect is within its reach, when it springs up and seizes it. This is the uncommon posture before alluded to. Should the insect go far from the spot, it slies, or crawls after it, slowly on the ground like a cat, and when the insect stops, erects itself as before. They have a small black pupil or sight which moves in all directions within the parts we usually term the eyes, so that it can see its prey in any direction without having occasion to disturb it, by turning its head.

The most prevalent colour of this tribe of insects is fine green, but many of these fade or become brown after the insect dies: some are finely decorated with a variety of vivid hues; the most beautiful of these, that we have seen, are from the Moluccas.





* Gryllus vittatus.

* Gryllus masutus.

GRYLLUS NASUTUS.

GENERIC CHARACTER.

Head inflected, armed with jaws, and furnished with palpi. Antennæ either setaceous or filiform. Wings wrapped round the body, and concealed under the elytra. Feet armed with two nails or crochets.

Acridæ.

SPECIFIC CHARACTER.

Head conic. Antennæ ensiform, or fword shaped. Body green.

GRYLLUS NASUTUS: capite conico, antennis enfiformibus, corpore viridi. Linn. Syft. Nat. 2. 692. 1.

This species is found in Africa, Asia, and the south of Europe. Its varieties are numerous; and in size and colour depends on the climate they breed in. Sulzer represents it with red wings: in the Chinese specimens these are tinged with green.

GRYLLUS VITTATUS.

SPECIFIC CHARACTER.

Head prominent, testaceous. A filver stripe on each fide of the head and thorax, and along the posterior thighs.

TRUXALIS VITTATUS: capite prominulo testaceus capite thorace femoribusque posticis vitta laterali argentea. Fab. Syst. Ent. T. 2. p. 27. 3.

Gryllus vittatus has not been figured by any author. A fingle specimen of it has been recently brought from China, and is in the possession of Mr. Francillon.

GRYLLUS GRYLLOTALPA.

MOLE CRICKET.

Acheta.

SPECIFIC CHARACTER.

Wings much longer than the elytra, and when folded, appear like tails. Anterior pair of feet palmated.

GRYLLUS GRYLLOTALFA: alis caudatis elytro longioribus, pedibus anticis palmatis.

Linn. Syft. Nat. 2. 693. 10.

This differs in no respect from the European species of the same name, except in size and colour. The Mole Cricket of this country, for example, being twice as large, and more of a mouse colour. Mr. Abbot has sent us a variety similar to the Chinese sort, but of a darker colour, from North America.

GRYLLUS ACUMINATUS.

Tettigoniæ.

SPECIFIC CHARACTER.

Thorax roundish, emarginated. Vertex subulated, or awl shaped. Wings greenish, and of equal length.

GRYLLUS ACUMINATUS. Linn. Syft. Nat. 2. 696. 23.

Locusta Acuminata: thorace rotundato utrinque emarginato, vertice subulato, alis virescentibus æqualibus.

Fab. Ent. Syst. T. 2. p. 39. Sp. 22.


Inhabits China and every other part of India.






Gryllus flavicornis : Gryllus Gryllotalpa.

Soundard Problems as whether down by E.Domeson Catalogary of







LEPIDOPTERA.



Gryllus morbillosus.

and a Published as the Art direct by E. Denevan, July 2 2744.





*Locusta perspicillata. **Locusta acuminata.

GRYLLUS PERSPICILLATUS.

SPECIFIC CHARACTER.

Thorax deflexed. Wing cases concave, green: a process at the base of each that fold one over the other; the lower one transparent, or glassy: upper one coriaceous.

Locusta Perspicillata: thorace deflexo, elytris concavis viridibus: bafi ocello dorfali fenestrato.

Fab. Ent. Syst. 2. p. 36. 10.

Fabricius erroneously describes this as a native of America. It is not figured elsewhere.

GRYLLUS FLAVICORNIS.

Locustæ.

SPECIFIC CHARACTER.

Upper part of the thorax carinated or keeled, green, without spots. Base of the wings red. Posterior thighs red, with yellow teeth.

GRYLLUS FLAVICORNIS: thorace fubcarinato viridis elytris immaculatis, alis bafi rufis, tibiis posticis fanguineis flavo ferratis. Fab. Ent. Syst. T. 2. p. 52. 23.

Described by Fabricius; but not figured. It is abundant in China.

GRYLLUS MORBILLOSUS.

SPECIFIC CHARACTER.

Thorax fquare, warted, bright red. Wing cases susceptibles with white spots. Wings red.

Gryllus morbillosus: thorace quadrato verrucoso rubro, elytris susceptibles albo punctatis, alis russ.

Linn. Syst. Nat. 2. 700. 38.

The Gryllus morbillosus appears in the early addition of the Systema Naturae, and the works of Roesel, as an Indian species: Mr. Drury affures us he has received it several times from China. Another sort is

also found at the Cape of Good Hope, which is rather larger and deeper in colour than the Chinese variety.

When this infect is at reft, the wings are folded, and much of its beauty is concealed: but when these are expanded, its appearance is altogether magnificent. It has nothing of the shining and metallic splendour of the Colcopteræ, for its colours are translucent, and assume their richest hues when they pass before the light. The elytra are purple, variegated with yellow: the wings of a glowing crimson, spotted with black. The abdomen is surrounded with alternate zones of black and yellow, and the legs are throughout of an elegant scarlet, inserior only in brightness to the coral red of the head and thorax. Upon the whole, this species is embellished with such a profusion of various and beautiful colours, that it may be considered as a most splendid example of the Hemipterous order of insects. It is represented on the Iris Chinensis in a slying position.

This is not supposed to be a numerous species in China; on the contrary, it is probably uncommon. Several others of the locust are abundant in that country, and in seasons favourable to their increase, do incredible mischief k. Both the Gryllus tartaricus, and Gryllus migratorius, inhabit Tartary on the northern confines of China, from whence, at certain periods, they descend like an impetuous torrent over the neighbouring countries in quest of food, strip the earth of verdure, and scarcely leave the vestige of vegetation behind them. The Gryllus migratorius, whose myriads are said to darken the face of heaven in their slights, sometimes direct their course westward; cross rivers, sea, and an immense extent of country, till they reach Europe; and though many are lost in these bold migrations, the survivors are in sufficient numbers to commit vast depredations. This species has been known to visit England 1, but not in any abundance. In Little Tartary, and the European provinces of Turkey: in Italy, and in Germany m, they do great mischief in these migrations. The Gryllus flavicornis and nasutus, are two other abundant species in China, and no doubt there are many other common kinds in that country we are at present unacquainted with. The locust is only detrimental when in immense numbers, for in China, as in other eastern countries, they are considered as an article of food, and regularly exposed for sale in the public markets n.

- k "Famines fometimes happen in this part of the province. In fome feafons inundations, produced by torrents from the mountains, and as often the depredations of locusts are causes of this disaster." Vide Sir J. Staunton, Chap. on Tien-sing.
- ¹ The last appearance of this species in England was in 1748. We have a specimen of it from Smyrna, Germany, and China, and deem it too common and general an inhabitant to merit a figure as a Chinese inscet.
- m Roefel fpeaks of this locust infesting the provinces of Wallachia, Moldavia, and Transylvania, in such immense numbers, in the years 1747, 1748, and 1749, that an Imperial and Royal Hungarian edits was iffued, with printed instructions for the best means of exterminating them. Vide Der Heuschrecken-und Grillensammlung, &c. &c. Vol. II. page 193.
- n Sir G. Staunton likewife fpeaks of "a large fpecies of Gryllus" that is kept in cages for amufement in China, and was exposed for sale, with other insects, in the shops of Hai-ten. Neither the species of this, or the locusts noticed in the preceding chapter, are mentioned.



Fulgora Candelaria ().

London Published as the Act directs by E. Donovan, Feb 1.17,98.

FULGORA CANDELARIA.

CHINESE LANTERN-CARRIER.

GENERIC CHARACTER.

The forehead elongated. Antennæ below the eyes: confift of two articulations. The beak, or roftrum, is bent inwards under the body.

SPECIFIC CHARACTER

AND

SYNONYMS.

Trunk curved upwards towards the end. Shells green with yellow marks. Wings yellow, black at the tips.

FULGORA CANDELARIA: fronte roftrata adscendente, elytris viridibus luteo maculatis, alis flavis apice nigris.

Linn, Syft. Nat. 2. 703. 3. Fab. Ent. Syft. t. 4. p. 2. Sp. 4.

Der Fleinere Asiatische oder Chinesische Lanternen-Trager. Roef. Inf. 2.

Gryll. 189. tab. 30.

Acta Holm. 1746. 63. tab. 1. fig. 5, 6.

De Geer Inf. 3. 197. 2.

Edw. Av. tab. 120.

Sulz. Inf. tab. 10. fig. 62.

De Gewapende Cicade. La Cigale armée. Stoll. Cicad.

The phenomena refulting from the properties and effects of light, having engaged the attention of the earlieft philosophers, we must conclude, that phosphorical appearances, and those especially of animated bodies, could not fail to attract their particular notice. Indeed it is evident, from the writings of the accurate observers of nature in remote ages, that they were acquainted with certain insects that have the property of shining in the night. These were known only by general terms, expressive of that property; yet it is probable that some of the Linnæan lampyrides, which are abundant in the south of Europe, as well as in Asia, and some parts of Africa, were the first of the illuminated insects known to them a. Some of the males which are furnished with wings, and are illuminated like the semales, were striking objects of natural history, and could scarcely have escaped their notice. The Greeks included all shining insects under the name lampyris; and the Latins called them cicindela, noctiluca, luciola, lucernata, &c. Whether any of the Fulgoræ were known to the ancients is uncertain: probably they were not, the most remarkable species being peculiar to the warmest parts of America. Asia, once the seat of learning, does indeed produce a few species; but we have no account of these in ancient natural history.

^{*} The lampyris of Pliny is expressly the infect with a shining tail.

The Fulgoræ feem to have been entirely unknown in Europe till the latter end of the laft century; when two writers published descriptions and figures of Fulgora Lanternaria. Madame Merian, of Holland, in her splendid work on the Metamorphosis of the Insects of Surinam, and Dr. Grew, of London, in his Rarities of Gresham College.

Reaumur b is the next author who described the Fulgora Lanternaria, and after him Roesel, in his "Amusing History (or recreation) of Insects c." This brings us to the period in which Fulgora Candelaria, our Chinese species, was first known in Europe: a circumstance of much importance to naturalists at that time, because the first mentioned species was a solitary example of its singular genus. The transactions of the Stockholm academy includes the earliest sigure and description of this extraordinary insect; from these Linnæus described it in his Systema Naturæ; but his arrangement has undergone many alterations since that time. At first he included it among the Coleopterous insects; then he called them Lanternariæ; and again, they were classed with the Cicadæ. These have been since corrected, and a new genus formed of insects with elongated trunks on their foreheads exclusively, under the name Fulgora, a name that has been adopted by later authors, and finally by Fabricius.

Roefel has given three figures and a description of the Fulgora Candelaria; from his account we learn that it was known in England before he was acquainted with it. On its peculiar qualities, he observes he must be entirely filent, because he had been unable to derive any information concerning it; his description is notwithstanding extremely prolix, and occupies four quarto pages; as he notices every trisling particular of the spots, colours, &c. of the infect. We have selected the most interesting passage, because it clearly marks the progressive advancement of the knowledge of natural history in Europe, so late as the middle of the present century.

"According to my promife," fays Roefel d, "I now produce the fecond fort of Lantern-carrier, which I never faw before; and of which I have never read in any work on infects. The feareer however it may be, the more I am indebted to Mr. Beurer, apothecary of this place e, &c. for the permiffion he has granted me to draw and enrich my collection with it. Mr. Collinson has fent it to him from London, under the name Lanternaria Chinensis, for which reason I have called it the Afiatic or Chinese Lantern-carrier." Roesel being a respectable entomological writer of his time, we must infer that Fulgora Candelaria was extremely scarce in Europe when his plate and descriptions were published. The commercial concerns of Europeans with the Chinese having greatly increased fince that period, has facilitated many inquiries concerning the natural productions of China; and amongst a variety of other insects that are now usually brought from that country, specimens of Fulgora Candelaria are not uncommon. In China sew insects are found in greater abundance.

b Memoires pour servir a l'Histoire des Inscetes. 1734.

[·] Infecten Beluftigung.

d Versprochener massen liesere ich nunmehr die zwente Sorte des Lanternen-Tragers, &c. Vol. 1. pl. 30. Locust, page 189.

e Nurenberg.

The authors who have figured or described the Chinese Fulgora, since Roesel, are De Geer, Sebas, Sulzerh, and our countryman Edwardsi. De Geer gives a very concise description, and no figure; and Seba merely observes, in his description of Fulgora Candelaria, "La Chine produit une autre espéce du même genre mais beaucoup plus petite et toute differente." Edwards has given a figure of it in one of the plates in his work of Birds. His description is certainly uninteresting k.

Having noticed the feveral authors who have treated on this infect, we come to confider the peculiar properties of its fingular genus; and among these we find the most association that insects can possess, that of emanating light: not merely a momentary shiving appearance, as is produced by many viscous substances, but a clear and constant resemblance to the element, fire; and capable of disfusing light to surrounding objects, though totally destitute of every principle that can do mischies. To the unphilosophical mind it appears at first impossible, and it cannot fail to associate informed. Indeed, some readers might be inclined to doubt the veracity of travellers in foreign countries, who have seen a vegetable of an animal produce light, if our own country could not supply us with abundant analogous proofs of such phænomenæ. The presence of this animated phosphorus, if we may so express it, is observed on several insects that are natives of England: it is needless to enumerate them, because the most striking example must be recollected by every rural inhabitant, or admirer of poetical simplicity.

THOMSON.

The account which Madame Merian gave of the effect of the light produced by the Fulgora Lanternaria was greatly differedited, though Dr. Grew had related some surprising particulars of a specimen of it

- f De Geer. Histoire des Insectes. 1752.
- E Seba. Thefaurus Naturze. 1765.
- * Kenwzeichen der Insecten (Characteristics of Insects). 1761.
- 1 Edwards's Birds, 4to. London, 1743.
- k "I take it to be a kind of fire-fly, and that part on the head the lantern. The wings are all flat, and the upper ones hard and fmooth, like those of the Beetle or Scarabai kind, of which it is a species." Edwards.
- An inflance of this occurs in the fouth of Europe. An account in the Philosophical Transactions relates of the Dictamnus Albus (Fraxinella), that " in the ftill evenings of dry feasons it emits an inflammable air, or gas, and flashes at the approach of a candle. There are certain inflances of human creatures who have taken fire spontaneously, and been totally consumed."
- m The account which Madame Merian has given of the light of the Fulgora Lanternaria is so surprising, that it will certainly prove acceptable to many readers. It is indeed a digression from the account of Fulgora Candelaria, but will tend to prove, that insects of this genus emit a more vivid light than any of the illuminated kinds hitherto known.
- "Once," fays Madame Merian, "when the Indians brought me a number of these Lantern carriers, I put them into a wooden box, without being aware of their shining at night; but one night, being awakened by an unusual noise, and much frightened, I jumped out of bed, and ordered a light, not knowing whence this noise proceeded. We soon perceived that it originated in the box; we opened with some inquietude, but were still more alarmed, after opening it, and letting it fall on the ground, for a stame appeared to issue from it, which seemed to receive additional lustre, as often as another insect flew out of it. When we observed this some time, we recovered from our terror, and admired the splendour of these little animals." Differt. de generations et metamorphibus Insectorum Surinamensse.

[&]quot; On every hedge

[&]quot; The glow-worm lights his gem, and thro' the dark

[&]quot; A moving radiance twinkles."

from Peru. Her account has, however, been generally believed, fince the miffionaries in countries which produce those insects have confirmed her account. It is admitted, that the *Chinese Fulgora* has an illuminated appearance in the night. "The foreheads of many Fulgoræ (especially those found in China) emit a lively shining light in the night time, which, according to some authors, is sufficient to read by."—Yeats.

The light of the Fulgoræ is generally imagined to iffue from the trunk, or elongated projection of the forehead; but Roefel offers a conjecture on the light of the Fulgora Lanternaria, which, on further inveftigation, may enable naturalists to determine whether the light is entirely produced by an innate property of the trunk, or receives additional splendour from some external cause. He notices a white farinaceous substance on several parts of the wings and body, as well as the trunk, which, he observes, looks like the decayed wood which shines at night. We mention this conjecture of Roesel, though the same occurred to us before we perused his observations. We have invariably found a similar white powder on other insects of this genus, but usually upon the trunk only. The remarks of Roesel were necessarily very limited, two species of the Fulgoræ only being then known. We posses twelve distinct species, and have made dissections and observations on several others; from all which we are inclined to imagine, that the white powder has a phosphoric appearance in the living insect, and increases the light, when the end of the trunk is illuminated.

One of the Fulgoræ of confiderable magnitude, from the interior of India, enabled us to make many observations. The trunk is of the same form as that of the Fulgora Candelaria. The colour is a dark but beautiful purple: the apex scarlet, of a perfectly pellucid appearance, and still retains a reddish glare. The spots of white, sprinkled on the purple colour of the trunk, exhibit also a slight appearance of phosphoric matter. On the trunk of the Fulgora Candelaria these white spots are very conspicuous.

Though the generic name Fulgora feems to imply fome effulgent property in the infects that compose the genus, it is uncertain whether all possess that property. They are indeed furnished with a trunk, but it is smaller in proportion, in several species, than in F. Lanternaria, Candelaria, Flammea, Phospho-

n Cucujus Peruvianus.

[&]quot;That which, befide the figure of the head, is most wonderful in this insect, is the shining property of the same part, whereby it looks in the night like a lantern, so that, two or three of these sastened to a stick, or otherwise conveniently disposed of, will give sufficient light to those who travel or walk in the night." Grew. Museum Regalis Societatis, p. 158.

o Le ver-luifant. Ceux que nous voyons à la campagne dans les nuits d'été ne jettent qu'une foible lueur: mais ils y en a dans les Indes modernes qui répandent un éclat très-vif. Ce font, pour ainfi dire, des phofphores animez. "Les Indiens, dit le favant auteur de la Théologie des infeétes, "ne fe fervoient autrefois dans leurs maifons, et dehors d'aucune autre lumiere. Lorsqu'ils marchent de nuit, ils en attachent deux aux gros doigts du pied, et en portent un à la main. Ces infeétes répandent une fi grande clairté, que par leur moyen on peut lire, écrire, et faire dans une chambre toutes les autres choses nécessaires." Lesser Liv. 2. c. 3. rem. 8. Le trait rapporté par le P. du Tert re dans son Histoire des Antilles, auroit bien dû être cité il dit avoir lû son breviare à la clarté d'un de ces vers-luisans.

P Yeats. Inflitutions of Entomology.

rea, and fome others. It has not been determined whether any of the European Fulgoræ shine in the night time. The genus is very limited, including the discoveries of modern naturalists. Fabricius deferibes only twenty-five species; of these, seven are European, and eighteen extra-European. Two species are found in England.

Chryfanthemum Indicum.-Indian Chryfanthemum.

This is a very lately introduced species in England. It is mentioned by Sir G. Staunton among the plants collected in the provinces of Shan-tung and Kiang-nan. Thunberg describes it as a native of Japan in the Flora Japanica. A variety of it is figured in the Hortus Malabaricus; and in the Herbarium Amboinense is another. We observe a great dissimilarity between the figures of this plant in different works. That figured in the Herb. Amb. has very small flowers, scarcely broader than our large daisy q: the leaves in clusters, some very large, and others small. In the Hort. Malab. the flowers are twice the fize of the former, and the leaves are placed much as afunder. The flowers of our specimen are considerably larger than either of these; yet not of the magnitude represented in the plate of Mr. Curtis's Botanical Magazine.

This Chrysanthemum is not peculiar to China, though it has been long cultivated in that country. It grows spontaneously in some parts of Japan; and from the name it bears in the Arabic, Persic, and other languages, is probably known in most parts of Asia.

9 Chryfanthemum Leucanthemum.

CICADA SANGUINEA.

RED AND BLACK CHINESE FLEA-LOCUST.

GENERIC CHARACTER.

Roffrum bent under the breaft. Antennæ fetaceous. Wings four, membranaceous, declining along the fides of the body.

SPECIFIC CHARACTER

AND

SYNONYMS.

Dark brown. Abdomen fanguine red; two spots of the fame colour on the thorax.

CICADA SANGUINEA alis superioribus fuscis, fronte abdomine thoracisque maculis binis sanguineis.

De Geer Inf. 3. 221. 18. tab. 33. fig. 17.

Tettigonia Sanguinolenta nigra ore thoracis maculis duabus abdomineque fanguineis. Fab. Spec. Inf.
2. 321. 19. Ent. Syft. 4. p. 25. 32.

LA CIGALE CHOINOISE a taches rouge de fang. Stoll. Cicad. tab. 13. fig. 62.

Fabricius refers to the cabinet of Mr. Drury of London for a specimen of this rare insect. The authors who have given sigures of it are De Geer, a French entomologist, and Stoll, in a work lately published on the continent, including Cimices and Cicadæ. The Cicada sanguinolenta of Linnæus being common in Europe, the similarity of names might create confusion, were we not to notice it. Our species is a Cicada of Stoll, as well as of De Geer, which latter author we have followed in the specific name. Fabricius having separated the Linnæan Cicadæ into several genera, this species will be found in his Entomologiæ Systematica under the name Tettigonia sanguinolentä.













Cicada sanguinea.
____ambigua.
___lanata.



Tettigonia splendidula. Cicada abdominalis. _____frontalis.

CICADA AMBIGUA.

SPECIFIC CHARACTER.

Olive colour. Wing cases clear, anterior margin testaceous.

CICADA AMBIGUA: olivacea, elytris hyalinis, marginibus anticis testaceis.

Mr. Drury received this infect from China. It is a species of ambiguous character, but appears to us undescribed. It is not noticed among the Tettigoniæ of Fabricius, in his last systematic arrangement of infects.

CICADA LANATA.

SPECIFIC CHARACTER.

Wing cases black at the ends; spotted with blue. Front and sides of the head red. Abdomen tusted with wool or down.

CICADA LANATA: elytris apice nigris: punctis cœruleis, frontis lateribus rubris, ano lanato. Linn. Syft.

Nat. 2. 711. 42.

Fab. Ent. Syft. T. 4. p. 30. 16.

One of the most beautiful species of the Indian Cicadæ. The wing cases are black, elegantly reticulated, and spotted with bright blue. At the extremity of the abdomen it has a tust of long and very delicate hairs, intermixed with others that are rather convoluted and of a coarser texture. The whole of this insect, but particularly between the abdomen and wings, is sometimes profusely covered with a sine powder of a snowy whiteness, similar to that observed on the Cicada limbata in the impersect state; hence we may conclude it is also one of those insects, which furnish the white wax d so highly esteemed in China.

TETTIGONIA SPLENDIDULA.

SPECIFIC CHARACTER.

Wing cases brown, highly glossed with a golden tinge. Thighs of the fore pair of legs thick; armed with teeth: colour red.

Tettigonia splendidula: elytris fusco aureis, tibiis anticis incrassato dentatis rusis. Fab. Ent. Syst. T. 4. 25. 33.

The infect added under this Fabrician character, is the fame that author described from the cabinet of Mr. Drury, and to which he exclusively refers. We believe the specimen is unique, at least we have not feen it in any other collection; and it has not been figured in any preceding publication. In the Linnæan arrangement it must be placed in the Cicada genus.

CICADA ABDOMINALIS.

SPECIFIC CHARACTER.

Shining black, with two yellowish bands across the wing-cases. Abdomen blood red.

Cicada abdominalis: atra nitida elytris fasciis duabus flavescentibus. Abdomine sanguineo.

An undescribed species; allied both to the Cercopis cruentata and Cercopis versicolor of Fabricius, but clearly distinct from either; the thorax of the former being black; and the elytra of the latter marked at the base with two white spots: the abdomen black. The thorax of our species is black: it has no white spots on the elytra; and the abdomen is of a sanguineous red colour


CICADA FRONTALIS.


SPECIFIC CHARACTER.

Wing-cases red. Five black spots on the head and thorax, and one in front between the eyes.

Cicada frontalis: pallida occipite thoraceque punctis quinque nigris, fronte puncto nigro inter oculos, elytris sanguineis.

We are unable to discover this species among the descriptions of Fabricius, and conclude it must be a new insect. The situation of the black spots on the thorax, and especially that on the front of the head, constitute the decisive criterion of this species.





Cicada Atrata.

London Published as the lot directs by E. Domovan Sept 21798.

CICADA ATRATA.

GREAT BLACK CHINESE FROG-HOPPER.

SPECIFIC CHARACTER.

Black; wings white, black at the base; veins yellowish brown.

TETTIGONIA ATRATA: atra alis albis basi nigris: venis testaceo. Fab. Ent. Syst. 4. 203. 28.

Though the observations that Sir G. Staunton has made on the natural production of China, in his late publication, were necessarily very general, the science of insects appears to have engaged his particular attention; and on that account we must lament, that untoward events, precluded him from observing more minutely, the peculiarities of some kinds, and the economical purposes of others.

In the general histories of unknown countries, (and we can scarcely consider China in any other view) the entomologist must expect to find his favourite science neglected, or treated in a manner more likely to excite curiosity than reward enquiry. The work of this learned writer is an exception to our remark; though a few inaccuracies occur in it. We peruse the following account of an unknown species of Cicada with particular regret, because, it withholds information interesting to the naturalist, and, from its air of novelty, is likely to promote an erroneous opinion concerning that singular tribe of insects.

"The low and fometimes marshy country, through which the river a passes, is favourable to the production of insects; and many of them were troublesome, some principally by their sting; and others by their constant stunning noise. The music emitted by a species of Cicada was not of the vocal kind; but produced by the motion of two slaps or lamellæ which cover the abdomen or belly of the insect. It is the signal of invitation from the male of that species to allure the semale, which latter is quite unprovided with these organs of courtship b."

Again, when describing a town higher up the river, that author says, "The shops of Hai-tien, in addition to necessaries, abounded in toys and trisses, calculated to amuse the rich and idle of both sexes, even to cages containing insects, such as the noisy Cicada, and a large species of the Gryllus s."

The reader may imagine from the first account, that the music of every other species of Cicada is of the

vocal kind, or that it is peculiar to this Chinese insect to be furnished with lamellæ that cause a sound. The latter account confirms such conjecture, by alluding in a specific manner to the noisy Cicada, as to an insect described in a sormer part of the work. We must remark, that not only the males of the species mentioned by that author, are furnished with those lamellæ, but the whole of that section of the Linnæan Cicadæ which Fabricius has called Tettiogonia d. The males of the species included in the other sections of that genus are certainly furnished with them also; though some of them are too minute to be observed without a glass. These lamellæ vary in size in different species; but the accounts we have of them from travellers in foreign countries, and naturalists both ancient and modern, prove they all emit a certain sound to allure the semale. As we are unable to ascertain the Chinese species Sir George mentions, neither sigure or description accompanying his account of it, we must therefore, speak generally of the whole genus, and then confine our remarks to those species we are acquainted with from China. Among these are C. spendidula, sanguinea, and atrata. The latter, we believe, is the largest species of the Chinese Cicadæ known in Europe.

Some species of this tribe were known to the ancients. With them it was the emblem of happiness and eternal youth e; and if we examine the legends of pagan mythology, we find they were deemed a race of creatures beloved by gods and men. The Athenians wore golden Cicadæ in their hair, to denote their national antiquity; or that like those creatures they were the first born of the earth; and the poets feigned that it partook of the perfection of their deities. Anacreon depictures in glowing colours the uninterrupted felicity of this creature: his ode to the Cicada is appropriate to our enquiry b.

- d This includes all the larger species of Linnzan Cicadz, such as C. Orni, Tibicien, reticulata, harmatodes, stridula, &c. &c. &c. for in some editions of the Systema Naturz of Linnzus we find Cicada Noctiluese, which in later editions are the Fulgorse, and also the C. cruciatse, mannifera, spamantes and deflexes included in one genus.
- e Probably because it was supposed to live only a short time. The renewal of youth is illustrated by the story of the Tithonus transformed by Aurora into a Cicada.
- g These pagan deities were without slesh or blood, and composed of aerial and watery humours. Such they imagined the moisture of the Cicada, and perhaps for that reason first affigned it a place among their demi-gods.

h Happy creature! what below

Can more happy live than thou?

Seated on thy leafy throne,

(Summer weaves the verdant crown,)

Sipping o'er the pearly lawn

The fragrant nectar of the dawn;

Little tales thou lov'ft to fing,

Tales of mirth—an infect king:

Thine the treafures of the field,

All thy own the feafons yield;

Nature paints for thee the year,

Songster to the shepherds dear:

Innocent, of placid fame,

What of man can boast the fame?

Thine the lavished voice of praise,

Harbinger of fruitful days;

Darling of the tuneful nine,

f Pharbus is thy fire divine;

Pharbus to thy notes has giv'n

Music from the spheres of heav'n:

Happy most, as first of earth,

All thy hours are peace and mitth;

Cares nor pains to thee belong,

Thou alone art ever young;

Thine the pure immortal vein,

Blood nor stefa thy life sustain;

Rich in spirits—health thy feast,

Thou'rt a demi-god at least.

GREEN'S Transs. Ode +3.

In the infant ftate of mufic, men, feem to have preferred the natural founds of fome animals, to those of their uncouth inftruments. We cannot otherwise account for the extravagant praise, bestowed on the noise of this little creature. It is true, authors agree that the founds of some kinds are exceeding loud and harmonious, and in the early ages of the world these might have a powerful influence on the human mind. It is related that the ancient Locri, a people of Greece, were so charmed with the song of the Cicada, that they erected a statue to its honour i.

The ancients had attentively observed the manners of its life, though they indulged in many poetical fictions concerning it; and particularly, when they affirmed that it subsisted on dew. They have told us that it lives among trees k, which circumstance discountenances the opinion of those moderns, who imagine the grashoppers 1 were the Cicadæ of the ancients.

Neither were they ignorant that the males only were furnished with those instruments which externally appear to produce its found, or the purpose for which that sound was emitted "; though it was reserved for more accurate naturalists to discover the complex organs by which it was caused and modulated. Aldrovandus, near two centuries ago, described the lamellæ, which he compares to the fruit of some herbs, called by modern botanists Thiaspin.

Among later naturalists who have noticed the Cicadæ of foreign countries are Merian°, Margravius p, &c.

Merian fays, its tune refembles the found of a lyre, which is heard at a distance; and that the Dutch in the

- i Some fay that once a certain player of Lori, contesting in the art of music with another, would have lost the victory, by the breaking of two strings of his instrument; but a Cicada slew to his aid, and resting on the broken instrument, sung so well, that the Lorian was declared victor. The Lorians erected a statue to the Cicada as a testimony of their gratitude. It represented the player with the insect on his instrument.
- k Dr. Martyn supposed this refers to the smaller branches in hedges, rather than to the lofty trees in forests: we cannot entirely coincide with that opinion.
- 1 Graftopper. Cicada. They live almost every where in hot countries. Lovel. Hist. Animal. containing the summe of all authors ancient and modern, p. 274, Sc. Sc.
- Circula, a Sauterelle*, or, according to others, a balm cricket.—Non est quod vulgò, a grashopper, vocamus; sed insectum longè diversum, corpore et rotundiore et breviore, qui arbusculis insidet et sonum quadruplo majorum edit. a grashopper, rectè locustum reddideris, Morl ex Ray. Ainstworth.
- = Xenarchus, an old Grecian play-writer, used to say jocosely that "the Cicadæ were very happy because they had filent wives."

 Arishotle also knew the sexual difference of them; he mentions them as a delicious food: he preferred the males when young, but more so
 the sexual difference of them; he mentions them as a delicious food: he preferred the males when young, but more so
 the sexual difference of them;
- Thalafpi parvum Hieraciifolium, five Lunariam luteam Monspel. et Leucoium luteum marinum. Lobel. Stirpium Adversaria nova, p. 74.—Aldrov.
 - · Merian. Infecta Surinamenfia.
 - P Georgi Margravii rer. nat. Brafiliæ. Lib. 7. p. 257.
- * Sauterelle, forte d'insecte. A locust or grashopper. Boyer. Cigale, a stying insect. The Cicada of the ancients, unknown in England. Boyer.

plantations of Surinam (where they are very plenty) call it the Lyre-player 4. Margravius, in his natural history of Brasil, compares it to the sound of a vibrating wire: he says the tune begins with Gir, guir, and continues with Sis, sis, sis. One species is called Kakkerlak in the Indies, perhaps because the sound emitted by it may be likened to the pronunciation of that word. Mr. Abbot, an accurate observer and collector of natural history in North America, has discovered four new species of Cicada, one of them nearly equal in fize to our Cicada Atrata. This, he says, was found in great abundance one season, in some swampy grounds near Susquehanna river, and was remarkable on account of their loud noise, which at a little distance resembled the ringing of horse-bells.

Some naturalists have supposed that the sound of the Cicada, is caused by the flapping of the lamellæ against the abdomen; and others, that it is only a noise occasioned by the rustling of the segments of the body in the contractile motion of that part. Beckman t imagines it is caused by beating the body and legs against the wings: he has endeavoured to explain the meaning of ancient authors, and deduce its etymology from that circumstance u.

Reaumur and Roefel have diffected feveral of the Cicadæ, and difcovered that the lamellæ cannot have that free motion necessary to cause such a sound; but that it is produced by some internal organs of the insect, and only issues through the opening, concealed under the lamellæ, as through the mouth of a mufical instrument *.

The suppositions of these authors seem well founded; we have examined many species that were unknown to them, and find the spine before mentioned, so placed in many insects, as to prevent the motion

- 9 De Lierman.

 F Scopoli carn. Yeats describes the Kakkerlak of the American islands as a species of Blatta, cock-toaches.

 Are there not two insects of that name?—one of them is, we believe, a Blatta.
 - 5 Communicated by Mr. Abbot in North America to Mr. Francillon in London.
- t Roef. Infetten Belluftigung.—Christiani Beckmanni, Bornenfis, manuductionem ad latinam linguam: nec non de originibus latinæ linguæ, &c.
- It is the common opinion that the word Cicada has its origin from quod cito cadat, which, after a general interpretation, implies that the Cicadæ foon vanish, or are short-lived. Beckman maintains that this opinion is abfurd, and proves that its name is derived from finging, because & abruv fignifies a found produced by the motion of a little skin; and that ciccum or cicum is a thin little skin of a pome-granute, that parts the kernels.—Beckman not knowing the infect, or not imagining that the little skin was an appendage to the abdomen, concluded it must mean the transparent wings, and consequently that the sound was produced by beating them against the body: but this interpretation, if applied to the lamelæ instead of the wings, will directly prove the origin of its name, and knowledge of the ancients.
- * For the satisfaction of the curious reader, we detail the most interesting particulars concerning the organization of these parts from Reaumur's Histoire des Insectes, and Roefel's Verschiedene aussaendische forten von Cicaden, Sc.

The music of the Cicada is not caused by the motion of the lamellar, as some have supposed. Reaumur observes, that although the lamellar have a kind of moveable hinge, they have also a stiff and pointed tooth, or spine, that prevents them from being listed far back; and, if strained, are very liable to be broken.

From the anatomical description of Roefel, we find that, within the two hollows that are seen when the lamellæ are listed up, two very smooth skins are visible; these are highly polished, of nearly a semicircular shape, and reflect prismatic colours: there is between these

of the lamelle. We have a specimen from America, which, in addition to the usual organs of sound, have two large hollow protuberances or drums; one on each fide of the abdomen; and must, we imagine, produce a louder sound than any yet discovered: a species very similar to this is also brought from New Holland.

The probofcis of those infects is a hard or horny tube, in which a very acute, slender sucking-pipe is concealed. The horny tube is not unlike a gimlet in form, and is used by those creatures to bore through the bark of trees, to extract the juices, on which it feeds. Linnæus has named the species of one division in his System, Mannifera, because they had been observed to fly among ash trees, bore many holes in them, and when the manna had oozed out, return and carry it off.

With this probofcis they bore holes in the fmall twigs of the extreme branches of trees, and deposit their eggs in them, sometimes to the amount of six or seven hundred. As each cell contains no more than from twelve to twenty eggs, it does great damage to the trees they frequent. Stohl says, "the common one,", which is found at Surinam in the coffee plantations, greatly injures those trees; the semales depositing their eggs in the young shoots, and in holes they bore with their sheath. They live on the juices of the trees."

a hard brown projection, or corner which unites with another piece above them in a longitudinal direction, to the under part of the breaft. This longitudinal piece divides a triangular red space or field into two parts, one on the right fide, and the other on the left. Above these, in a transverse direction, is seen two small yellow skins; the lamellæ in their natural position conceal these organs because they fold exactly over them.

Reaumur, in the exterior appearance of these parts, could discover nothing that could lead to determine the organs of the found; and he was not satisfied that the slight motion of the lamellæ on these parts could produce the loud singing noise of the Cicada. He opened a sew cicadæ on the back part of the body, so that the inner structure of the under side was displayed, and especially the parts connected to the curious organs he had discovered under the lamellæ. At last he discovered two large muscles, which at their point of union formed a space almost square, and were connected with the red triangular fields he had observed on the under side: as he concluded these formed a material part of the organs he wished to discover, he examined them attentively, and found that, by moving them backwards and forwards, he could make a cicada sing that had been dead many months. Although the found was not strong, it tended to prove that he had discovered the instrument that produced it.—In another part he says it is evident the sound is caused by the little skins connected to the muscles, because when they were rubbed with a bit of paper they emitted that kind of sound.

Roefel has discovered two little pieces of horny substance that are connected by a fort of fibre within the skins, in the body, and he supposes when this is in motion, it firikes against the before-mentioned thin skins, and produces a sound, by the same means as a hollow body, or drum, when struck with a stick: and also that this noise may be varied or modulated by a slight motion of the lamellæ, but cannot be produced without the affistance of the internal nerves and muscles connected with the organs first described.

Authors agree that the Cicadæ of hot countries emit the loudeft found. It appears from the papers of Mr. Smeethman (who refided a confiderable time in Africa) published by Mr. Drury, that the found of fome kinds peculiar to that part of the world is fo loud as to be heard at half a mile distance; and that the finging of one within doors, filences a whole company.—

The fame attentive observer says, the open parts of the country are never without their music, some singing in the evening and others only in the day.

y La Cigale Vieilleufe. Cicad. Tibicien.

M. Merian gives a figure and account of the metamorphose of a cicada found in Surinam. She has mistaken the winged insect to be only the pupa of the Fulgora Lanternaria, which is too absurd to deserve contradiction; in other respects her account is interesting, and particularly that part which relates to the pupa state, or chaser, as it is termed. "The pomegranate tree," says Merian, "so well known in all other countries, grows also in the fields of Surinam. On them I have found a species of chaser, which is naturally very lazy, and consequently very easy to be caught. It carries underneath the head a long trunk, with which it easily penetrates the slowers, in order to extract the honey from them. On the 20th of May, when they were laying quite quiet, the skin of the back burst open, and green slies, with transparent wings, issued from them. These slies are found in abundance in Surinam, and have such a rapid slight, that it took me many hours to catch one."

The pupa we received from China with our Cicada atrata, very much refembles that figured by Merian. It has the long fucking trunk or probofcis; but the most formidable of its weapons seem to be the fore feet, which are thick, strong, and armed with spines or teeth; with these it may do more injury to the plants, by tearing off the tender shoots, than by wounding the trunk to extract the mossture.

We have represented the upper and under fide of a male of this interesting species, Cicada atrata, not only to illustrate our preceding remarks, but because we believe no figure has been given of it by any author, unless De Zwarte Chinessche cicade of Stohl is intended for this insect. The Fabrician description has no synonyms.


The general appearance of both fexes of Cicada atrata is very fimilar, except that the female is furnished with a sheath, and the male with lamellæ. The sheath of the female is partly concealed within a valve at the extremity of the abdomen, and is only protruded when the creature lays her eggs. In the figure of the under surface of a male insect, exhibited in the annexed plate, the lamellæ are distinguished by two stars: the single star denotes the situation of the spine, mentioned by Roefel and Reaumur.

Laurus Camphora.—Camphor-tree.

The tree which produces the useful drug camphor is very abundant in Japan and China. Sir G. Staunton says it is the only species of the laurel genus growing in China, where it is a large and valuable timber, and is never cut up for the sake of the drug; but that substance is obtained by decocting the small branches, twigs, and leaves, and subliming the camphor in luted earthen vessels. A purer fort is brought from the island of Borneo and Japan, which is supposed to be a natural exudation from the tree when the bark is wounded. Sir G. Staunton says the Camphor-tree is felled in those countries for the sole purpose of finding the drug in substance among the splinters.

z La Cigale Chinoife noire, pres de Canton en Chine. Cab. de Monf. L. F. Holthufen, pl. 20. fig. 118.

A CONTROL OF THE PARTY OF THE P



Cicada Limbata.

London Published as the Act direct by E. Donovan May 1,750.

CICADA LIMBATA. var.

WHITE-WAX INSECT.

GENERIC CHARACTER.

Roftrum bent under the breaft. Antennæ fetaceous. Wings four, membranaceous, declining along the fides of the body.

SPECIFIC CHARACTER.

Wing cases deflexed, green, margin red. Interior case spotted with black. — Var. whitish, margined with black: a row of black spots on the posterior edge.

CICADA LIMBATA: elytris deflexis viridibus, margine rubro, interiori bafi nigro punctato.

Fab. Spc. Inf. 2. p. 322. 3.

CICADA LIMBATA var. Aliam fimillimam vidi, at totam ex albidam futura bafi nigro punctata margineque elytrorum poftico nigro. Fab. Spe. Inf. ibid.

This fingular infect, and the plant on which it is reprefented, have an equal claim to attention, both as objects of natural curiofity, and importance in domestic economy. The Larva is an elegant and beautiful creature, and China is indebted to its labours for the fine white wax so much esteemed in the East Indies. The plant is not less interesting, as it produces the vegetable tallow, in general use throughout the Chinese empire.

The novelty of these productions could not fail attracting the notice of those learned Europeans who were first permitted to reside in China, and whose object was to promote sciences and arts, as well as the christian knowledge. Both the Wax-insect, and Tallow-tree, are spoken of in their writings, as extraordinary and peculiar advantages to the country. Du Halde, especially in his splendid work L'Histoire de

LA CHINE, treats largely on these productions, in the sections Circ blanche d'Insectes & Parbre qui porte le suif. His relations are, perhaps, too prolix, but they are evidently the result of attentive observation, and serve to illustrate the Natural History, and economical purposes, of the subjects we are noticing: they are briefly these. The white wax of China is called Tchang pe la; it is not the same as the white wax of bees, but is the produce of a fort of insect that feeds on a tree called Tong tein: these insects form a kind of white grease which attaches to the branches of the tree, hardens, and becomes wax. It is scraped from the branches of the trees in autumn, melted on the fire, strained; then poured into cold water, where it coagulates and forms itself into cakes. This wax is very white and glossy; it is mixed with oil and made into candles, and is much superior to the wax of bees for that purpose.

f De la Cire Blanche,

Faite par des insectes, et nommée Tchang pe la, c'est-a-dire, Circ blanche d'insectes.

Ki dit. La Cire blanche dont il s'agit ici, n'eft pas la même que la cire blanche des Abeilles. Ce font de petits infectes qui la forment. Ces infectes fuccent le fue de l'espèce d'arbres nommée Tong tçin, et à la longue ils le changent en une sorte de graisse blanche, qu'ils attachent aux branches de l'arbre.

Il y en a qui disent que c'est la fiente de ces insectes, qui s'attachant à l'arbre, forme cette Cire, mais ils se trompent. On la tire en raclant les branches dans la faison de l'Automne; on la fait sondre sur le seu, et l'ayant passée, on la verse dans l'eau froide où e'le se fige, et se forme en pains. Quand on l'a rompuë, on voit dans les morceaux brisez, des veines comme dans la pierre blanche ou congélation nommée Pe che cao; elle est polie et brilliante: on la mêle avec de l'huile, et on en fait des chandelles. El'e est beaucoup superiéure à celles que sont les Abeilles.

Chi tchin dit. Ce n'est que sous la Dynastie des Yuen qu'on a commencé à connoître la cire formée par ces inscétes. L'usage en est devenu sert commun, soit dans la médecine, soit pour faire des bougies. Il s'en trouve dans les Provinces de Se tehuen, de Hou quang, de Yunnan, de Fo kien, de Tehe kiang, de Kiang nan, et genéralement dans tous les quartiers du Sud-Est. Celle qu'on ramasse dans les Provinces de Se tehuen et d'Yunnan, et dans les territoires de Hen teheou, et de Yung teheou est la meilleure.

L'arbre qui porte cette cire, a les branches, et les feüilles femblables à celles du Tong gin. Il conferve fa verdure durant toutes les faifons : Il pouffe des fleurs-blanches en bouquets durant la cinquième Lune; il porte des fruits en bayes, gros comme le fruit du Kin rampant.

Quand ils ne font pas mûrs, ils font de couleur verte; et ils deviennent noirâtres, lorsqu'ils mûriffent, au lieu que le fruit de Tong sin est rouge. Les insestes que s'y attachent sont fort petits. Quand le soleil parcourt les quinze derniers dégrez des Gemeaux, ils se répandent en grimpant sur les branches de l'arbre; ils en tirent le suc, et jettent par la bouche une certaine bave, qui s'attachant aux branches encore tendres, se changent en une graise blanche, laquelle se durcit, et prend la forme de cire. On diroit que c'est de la gelée blanche que le froid a durcie.

Quand le foleil parcourt les quinze premiers dégrez du Signe de la Vierge, on fait la récolte de la Cire, en l'enlevant de deffus les branches. Si l'on différe à la cuëillir que le Soleil ait entiérement parcouru ce Signe, il est difficile de la detacher, même en la raclant.

Ces infectes font blancs quand ils font jeunes, et c'eft alors qu'ils font leur cire. Quands ils deviennent vieux, ils font d'un châtain qui tire fur le noir, C'est alors que formant de petit pelotons, ils s'attachent aux branches de l'arbre. Ces pelotons font au commencement de la grofficur d'un grain de mil· vers l'entrée du printemps ils commencement a groffic et à s'étendre. Ils font attachez aux branches de l'arbre en forme de grapes, et à les voir, on diroit que l'arbre est chargé de fruits. Quand ils

Chi tchin, a Chinese writer, says, it was not till the dynasty of Yuen that the wax made by these insects began to be known; the custom then became very general to employ it both in medicine and making candles. Its medicinal virtues are highly extolled by the Chinese physicians, and particularly by Tchi hen. It is said to be a drug absolutely necessary to surgeons, that it makes the sless the sless bleeding, appeares pain, unites the nerves, rejoints the bones, and taken in pills, kills the worms that occasion consumptions. It is found in most of the south-east provinces of China; but the best is brought from Sc tchuen and Yunnan, and from the territories of Hen tcheou, and Yung tcheou.

These insects are white when young, and it is then they make their wax. When old, they are of a blackish chesnut colour, and form little pelotons on the branches of trees. These pelotons, at first, are about the fize of a grain of millet; towards the beginning of the spring they increase in bulk and spread; they are attached to the branches like grapes, and at first sight, the trees that bear them, appear loaded with fruit. About the beginning of May, they gather them, and having enveloped them in the leaves of Yo (a species of broad-leaved grass), they are suspended to the trees. At the end of June, and in July, the pelotons open, and the insects come forth, crawl about the leaves, and form their wax.

Sir G. Staunton, in his learned work, has also described the Wax insect; he found it at Turon Bay, in Cochin China, and has caused it to be represented in a vignette plate, with the following description.

"Among other objects of natural curiosity, accident led to the observation of some swarms of uncommon

font fur le point de mettre bas leurs œufs, ils font leur nid de même que les chenilles. Chacun de ce nids ou pelotons contient plufieurs centaines de petits œufs blancs.

Dans le tems que le foleil parcourt la feconde moitié du Taureau on les cüeille, et les ayant enveloppez dans des feüilles de Fa (efpece de fimple à larges feüilles;) on les fuspend à différens arbres. Après que le Soleil est forti du Signe de Gemaux, ces pelotons s'ouvrent, et les œufs produisent des insectes, qui sortant les uns après les autres des seüilles dont ils sont enveloppez, montent sur l'arbre où ils sont ensuite leur cire.

On doit avoir foin d'entretenir le dessous de l'arbre toûjours propre, et de le garentir des sourmis qui mangent ces insectes. On voit deux autres arbres ausquels on peut attacher les insectes, et qui porteront également de la cire; l'un qui se nomme Tien tehu, et l'autre qui est un espéce d'arbre aquatique, dont les seülles resemblent assez à celles du Tilleul.

Qualitez et effects de cette cire.

Elle eft d'une nature qui n'est ni froide ni chaude, et qui n'a aucune qualité nuisible. Elle fait croître les chairs, elle arrête le fang, elle appaise les douleurs, elle retablit les forces, elle unit les nerfs, et rejoint les os, prise en poudre dont on forme de pillules, elles fait mourir les vers qui causent la phtisie.

Tehi hem dit. La Cire blanche est sous la denomination du métal: ses esprits corroborent, sortifient, et sont propres à ramasser et a resserrer. C'est une drogue absolument nécessaire aux chirurgiens: elle a des essects admirables, quand on la fait entrer avec de la peau de Ho hoang dans la composition de l'onguent, qui fait renaitre et croître les chairs. Du Halde, Vol. IV. p. 495. large Folio, 1735.

infects bufily employed upon finall branches of a fhrub, then neither in fruit or flower, but in its general habit bearing fomewhat the appearance of a privet. These insects, each not much exceeding the fize of the domestic fly, were of a curious structure, having pectinated appendages rising in a curve, bending towards the head, not unlike the form of the tail feathers of the common fowl, but in the opposite direction. Every part of the insect was, in colour, of a perfect white, or at least completely covered with a white powder. The particular stem frequented by those insects, was entirely whitened by a substance or powder of that colour, strewed upon it by them. The substance or powder was supposed to form the white wax of the East. This substance is afferted, on the spot, to have the property, by a particular manipulation, of giving in certain proportions, with vegetable oil, such solidity to the composition as to render the whole equally capable of being moulded into candles. The fact is afcertained, indeed, in some degree, by the simple experiment of dissolving one part of this wax in three parts of olive oil made hot. The whole, when cold, will coagulate into a mass, approaching to the firmness of bees wax."

From the accurate description and figures of the latter author, it is evident, the creature that produces the white wax of China, is an impersect insect, or, technically speaking, the pupa of an insect, which, in its mature state, is surnished with wings. This is clearly the fact, for the rudiments of wings are visible in the figures alluded to 5. The metamorphose of insects are so various, and their appearance so changed in passing from one state to another, that the identity of any species in the larva or pupa can only be proved by actual observation; neither do the larva or pupa possess those characteristic differences by which naturalists determine one species from another; and this consideration deters us from deciding on the precise species, to which the pupa before us should be referred.

Stohl, a Dutch author, has been more fortunate; he has afcertained this identical creature to be the pupa of Cicada Limbata, and in his work on Cimices and Cicades, gives a figure of it under the title of De Waldraag ster (Nymphe) or La Cigale Porte Laine, fig. 144, together with the winged insect at fig. 145; and it is on this authority Cicada Limbata is introduced in the annexed plate.

We are ftrongly inclined to credit the accuracy of Stohl in this inftance; there is much fimilarity between the pupa and the cicada, and some striking characteristics are common to both. They agree in the structure of the antennæ, and proboscis, or sucking trunk; the abdomen of the winged insect is also loaded with a fine white powder, and is furnished at the extremity with a tust of down and hairs, similar to that so eminently conspicuous in the pupa state. We have, however, observed the white powder, and tust on the abdomen of Cicada lanata, and have reason to imagine it also forms a white wax, similar to that of the present species.

8 This may account for a paffage in Gordon's description of China, where he says, "In the plains" of Houquang "are vast numbers of little worms that produce wax, in the same manner as bees do honey," if we understand by worms, insects not arrived at maturity; for the larva of Bombyx Mori, is also termed a filk worm, though it belongs to the moth tribe when perfect.

The Cicada limbata is of a light green colour, with a red margin; that which Stohl has figured is of a pale brown, with a black margin. These are the species and variety Fabricius describes, for the specimens he refers to, in the collection of Sir Joseph Banks, agree precisely with our insects. Fabricius notes the habitat Africa. Stohl received the green specimen from the Island of Ceylon; the pale fort from Africa. The larva we have represented is from China; and the Cicada was brought from the East Indies, by the late Mr. Ellis.

Croton Sebiferum—Poplar-leaved Croton, or Tallow-tree.

The Tallow-tree is not the natural food of the Wax infect, but as they mutually illustrate the same inquiry, they are represented in the same plate; and it is further presumed, that a short account of this useful plant, will be deemed a proper sequel to the history of the insect.


Du Halde, when describing the Tallow-tree, says, "Il est de la hauteur d'une grande cerisser. Le fruit est rensermée dans un écorce qu'on appelle Yen Kiou, et qui s'ouvre par le milieu quand il est mûr, comme celle de la châtaigne. Il consiste en des grains blancs de la grosseur d'un noisette, dont la chair a les qualitez du suif; aussi en fait-on des chandelles, après l'avoir fait fondre, en y mêlant souvent un peu d'huile ordinaire, et trempant les chandelles dans la cire qui vient sur l'arbre dont je vais parler: il s'en forme autour du suif une espéce de croûte qui l'empêche de couler b. Page 18. Vol. I.

Sir G. Staunton speaks nearly to the same effect: "From the fruit of the Croton sebiferum, of Linnaus, the Chinese obtain a kind of vegetable sat, with which they make a great proportion of their candles. This fruit, in its external appearance, bears some resemblance to the berries of the ivy. As soon as it is ripe, the capsule opens and divides into two, or, more frequently, three divisions, and falling off, discovers as many kernels, each attached by a separate foot-stalk, and covered with a slessly substance of a snowy whiteness, contrasting beautifully with the leaves of the tree, which, at this season, are of a tint between a purple and a scarlet. The sat, or slessly substance, is separated from the kernels by crushing and boiling them in water. The candles made of this sat are firmer than those of tallow, as well as free

h "It is of the height of a large cherry-tree; the fruit is inclosed in a shell, called Yen Kiou, which, when ripe, opens in the middle like the chesnut. The fruit consists of white kernels of the size of a small, or hazel nut, whose substance has all the qualities of tallow; so that they make candles of it, after having melted and mixed it with a small portion of common oil, and then dipping the candles into the wax before mentioned, it forms a kind of crust round the tallow that prevents its running, or wasting.

from all offensive odour. They are not, however, equal to those of wax or spermaceti." This author further adds, "The wax for candles, is generally the produce of insects, feeding chiefly on the privet, as is mentioned in the chapter of Cochin China. It is naturally white, and so pure as to produce no smoke; but is collected in such small quantities, as to be scarce and dear. Cheap candles are also made of tallow, and even of grease of too little consistence to be used, without the contrivance of being coated with the firmer substance of the tallow tree or of wax." Vide Chapter on Sou-choo foo.

The tallow-tree is now cultivated in the West Indies, where it thrives well, and produces fruit, and by proper attention may hereafter become useful.







Nepa Grandis.

London Published as the Act directs by E. Donovan June 17798.

NEPA GRANDIS. GREAT WATER SCORPION.

GENERIC CHARACTER.

Roftrum bent inwards. Antennæ formed like legs.

SPECIFIC CHARACTER

AND

SYNONYMS.

Brown. Thorax, fcutellum, and elytra, varied with obfcure yellow marks.

Nepa Grandis, fusca flavo maculata. Fab. Ent. Syft. T. 4. p. 61.

Merian Surin. tab. 56.

Die groffe Surinamische, breitleibige Wasserwanze. Roes. Inf. 3. tab. 26.

Le Grand Scorpion-aquatique.

Stoll. Cimic. 2. tab. 7. fig. 4.

De Geer Inf. 3. 379.

M. Merian has given a plate and description of this species in her work on the Insects of Surinam. We learn from that account, that in the larva and pupa state it lives in the water; that it is a voracious creature, and feeds not only on the weaker kinds of aquatic insects, but on some animals much larger than itself. The pupa is represented on the back of a large frog in the water, and is designed to pourtray the manner in which it fastens on those creatures, holds them between its strong curved fore feet in, and extracts the juices of their bodies, through its singularly constructed beak. M. Merian says the winged insect came out of one of these creatures on the twelfth of May 1701.

Every writer on this infect fince M. Merian, appears indebted to her, for their account of these few particulars; for though all the European species of the same genus undergo precisely the same changes in their aquatic dwellings, among decayed vegetables, &c. at the bottom of the water, and quit it only in the

t These are femi-completa: unlike the pupa of the Lepidoptera, &cc. these scarcely differ in appearance or manners of life from the complete insect, but have only the rudiments of the wings.

[&]quot; If these correspond with the antennæ of other insects, the Nepa has only four sect.

winged state v, we are indebted to her for the time of the appearance of this exotic species in that state, as well as for a correct figure of its pupa.

Authors vary in their accounts of its native country. Linnæus, following Merian, makes it Surinam; Margravius, Brafil; and Fabricius, America generally. We observe a slight difference between our Chinese specimen and the figures in preceding works referred to by Fabricius; but in giving it as the Nepa Grandis of that author, we have no hesitation, having compared it with those species referred to by Fabricius in the collection of the Right Hon. Sir Joseph Banks, Bart.

NEPA RUSTICA.

EGG-CARRYING WATER SCORPION.

SPECIFIC CHARACTER

AND

SYNONYMS.

Without tail, brown. Margin of the thorax, and anterior edge of the wing cases, pale.

NEPA RUSTICA: ecaudata fusca thoracis elytrorumque margine antico albido. Fab. Ent. Syst. T. 4. 62. 3. Nepa Plana. Sulz. Hist. Ins. tab. 7. fig 2.

Infects in general discover an extraordinary degree of care, and ingenuity, in depositing their eggs in the most secure situations, or places where the infant brood, when hatched, may be provided with proper fustenance. Those of the aquatic kind usually lay them in recesses in the mud or fand, or under loose stones that lie at the bottom of the water: others, with as much care, and more ingenuity, hollow out the interior substance of the large stalks of water plants, and deposit their eggs in them; or, rising out of the water, lay them in the extreme branches of those plants, to secure them from other aquatic depredators. The Nepa rustica displays even more sagacity, or attachment for its eggs, than those creatures; for it never leaves them. Till they are hatched, it bears them on its back, in a cluster of an oval shape; these eggs are of an oblong form, and are sastened by the narrowest end to a thin film, or plate of cement, that causes them to adhere to the polished surface of the wing cases; when these eggs, about an hundred in number, are hatched, it casts off the exuviæ of the cluster, and differs no longer in general appearance from the male of the same species.

Our figures represent the fituation of the eggs on the back, and the infect also after they are cast off. It is not commonly received with the eggs upon it. Found on the coast of Coromandel, as well as China.

V Nepa Cinerea and Linearis are English species; these live in the water till they have wings, when they occasionally quit it to pursue other winged creatures.—China produces a species which differs very little except in fize from N. Cinerea. Fabricius calls it Nepa Rubba in his Ent. Syst. T. 4. g. 62. sp. 6. We deem a further description of it unnecessary.











+ Nepa rustica.

te Nepa nubra.

London Published as the Act directs by EDenovandoff 1798.







* Cimex Dipar.

: Cimex Hanbuschii.

CIMEX DISPAR.

OCELLATED BUG.

GENERIC CHARACTER.

Beak bent downwards. Antennæ longer than the thorax. Wings folded across each other: the upper ones coriaceous from the base to the middle. Back flat. Thorax margined. Feet formed for running.

SPECIFIC CHARACTER

AND

SYNONYMS.

Efcutcheon extends over the wings and abdomen. Thorax and efcutcheon deep orange with fpots of yellow, and a fmall black point in the middle of each.

CIMEX DISPAR: fcutellaris carneus thorace fcutelloque maculis flavefcentibus: quibufdam puncto ocellari atro. Fab. Spec. Inf. 4. p. 81. 7.

CIMEX OCELLATUS. Thunb. Nov. Spc. tab. fig. 72.

De Ongelyke Schildwantz. La punaise dispar. Stoll. Cimic. tab. 37. fig. 260.

This curious infect is among the number of those lately brought from China. A figure of the upper furface is represented on a leaf of the Camellia Sesanqua, one of the vignette plates of Sir G. Staunton's History of the late Embassy to that country; a coloured figure and a short account of it may therefore prove acceptable to the readers of his volumes.

Professor Thunberg of Upsal discovered this insect in his travels in Japan, and described it among his new species under the name Cimex Ocellatus. Stoll, in a work recently published on Cimices, has also given a figure of it: he has likewise represented another fort, which he considers as the semale (letter A); it has no black points in the yellow spots of the thorax and scutellum: he mentions the Isle of Formosa as the native country of his specimens.

Fabricius has rejected the specific name Ocellatus, which was given by Thunberg, and has called it Dispar: he has added a very minute account of its characteristic marks, &c.

We have reprefented the infect in a flying position, to display the fingular wings, that are concealed under the scutellum when it is at rest. The other figure, distinguished also by a star, exhibits the under side of the infect, which for variety and beauty of colours is not less interesting than the upper surface.

CIMEX STOCKERUS.

BLUE BUG OF STOCKER.

GENERIC CHARACTER.

Beak bent downwards. Antennæ longer than the thorax. Wings folded across each other: the upper ones coriaceous from the base to the middle. Back flat. Thorax margined. Feet formed for running.

SPECIFIC CHARACTER

AND

SYNONYMS.

Oval: green with black fpots: body ferrugineous.

CIMEX STOCKERUS: ovatus corpore viridi: maculis nigris, abdomine ferrugineo. Lin. Syft. Nat. 2. 715. 2. Fab. Ent. Syft. T. 4. 79. 212. 1.

LA PUNAISE BLEU DE STOCKER. Stoll. Cimic. fig. 15. p. 2.

Petiv. Gazoph. 34. tab. 21. fig. 12.

This beautiful infect is not peculiar to China; it is found on the coafts of Bengal and Coromandel, and in the Isle of Java. It feems a very common creature in those parts of the world; for we rarely receive a parcel of the infects of those countries, and from China in particular, that does not include many of them. There are several varieties of this species; some incline very much to green, and vary in the form of their spots, &c. We have a charming miniature variety of it from Africa aguin. about one third of the size of the Chinese specimens, and of a very deep blue colour. The marks both on the upper and under side precisely resemble those in the annexed figures.














· Cimex Stockerus. : Cimex aurantius

:. ____ oruciger.

Reduvius bifidus.

Phasianus



CIMEX AURANTIUS.

SPECIFIC CHARACTER.

Orange. Head, anterior margin of the thorax, and feet, black. Margin of the abdomen marked with black spots.

CIMEX AURANTIUS: aurantius capite thoracis margine antico abdominis maculis marginalibus pedibuíque

atris. Fab. Ent. Syft. T. 4. p. 105, 99.

Sulz. Inf. Tab. 10. Fig. 10.

La Punaise coleur d'orange. Stoll. Cimic. 2. Tab. 6. Fig. 39.

This elegant infect has been figured by Stoll and Sulzer. The four following are not, we believe, in any other work.

CIMEX CRUCIGER.

SPECIFIC CHARACTER.

Oblong: black. Thorax spined, and marked with four longitudinal ferruginous lines. Wing cases black, with a ferruginous cross.

LYGEUS CRUCIGER: thorace acute spinoso oblongus supra niger, thorace lineis elytris cruce serrugineis.

Fab. Ent. Syst. 4. p. 140. 22.

From the collection of Mr. Francillon, who received it from China. Fabricius describes it in his new genus Lygæus as a native of Brazil.

I

CIMEX PHASIANUS.

SPECIFIC CHARACTER.

Brown. Pofterior thighs arched, thick, armed with a fpine. Under fide of the abdomen gibbous at the base.

LYGÆUS PHASIANUS: fuscus femoribus posticis arcuato clavatis unidentatis abdominis basi subtus gibbo.

Fab. Ent. Syst. T. 4. p. 144. 35.

This species is allied to Cimex bellicosus, a native of Africa. Fabricius notes Lygæus Phasianus from Africa also: our specimens were brought from China by the late Mr. Ellis, surgeon.

CIMEX SLANBUSCHII.

SPECIFIC CHARACTER.

Red. Thorax marked with a black band. Scutellum black; and a fpot of the fame in the middle of the wing-cafe.

Lyg. Eus Slanbuschii: fanguineus thorace fascia abbreviata, scutello elytris puncto alisque atris.

Ent. Syst. T. 4. p. 155. 68.

Fabricius describes this insect from the cabinet of Schlanbusch. It is very common in China.

CIMEX BIFIDUS.

SPECIFIC CHARACTER.

Black. A rufous bar across the wing-cases. Scutellum furnished with an erect spine, which is two-cleft or bifid at the apex.

Reduvius Bifidus: ater elytris fascia rufa, scutello spina erecta apice bisida.

Fab. Ent. Syft. T. 4. p. 204. 40.

This very fingular creature is a Cimex of Linnæus. It is a rare species, and has only been described by Fabricius, who places it in the new genus Reduvius a.

a Linnæus comprehends under the general title Cimex, a variety of infects, very opposite in appearance, though not effentially different; and to avoid confusion in the arrangement of these diffimilar species in the same genus, he separates them into eleven distinct samilies, under the names of Apteri, Scutellati, coleoptrati, &c. Fabricius, throughout all his works, endeavours to divide the Linnæan Cimices, and finally, in the Entemologia Systematica, he refers them to the following genera, Acanthia, Cimex, Coreus, Lygeus, Miris, Gerris, and Reduvius.





Landon Published as the Act direct by E. Doner un August 279 b





Landon Published as the Act directs by Donovan June 1.1798

PAPILIO PARIS.

GENERIC CHARACTER.

Antennæ increase in bulk towards the extremity, and usually terminate in a kind of club.

Wings erect, when at rest. Fly in day time.

SPECIFIC CHARACTER.

Posterior wings tailed, black with a large blue spot, and a purple eye near the anterior margin. Seven lunar marks on the under side.

Papilio Paris: alis caudatis nigris: posticis macula cyanea ocelloque purpureo, subtus lunulis septem.

Linn. Syst. Nat. 2. 745. 3.

Fab. Ent. Syft. T. 3. p. 1. 1.

Cram. Inf. 9. tab. 103. A. B.

The fimile proposed by Linnæus for the arrangement of Butterslies, is gleaned from ancient and fabulous history. The species are divided into sections of Trojan and Greek princes, heroes, deities, nymphs, and plebeians: this allegorical system, which is well conducted, seems liable to less objection, than the characters assigned to each section: for many species placed among the Equites, and a more considerable number with the Plebeii, are inconsistent with the essential criterion Linnæus has given. This arrangement has undergone some material alterations in the Entomologia Systematica of Fabricius; alterations, perhaps justified, by the comprehensive view, its author has taken of this pleasing branch of Entomology. The Equites, with many additions, and a few exceptions, are the same as those in the two Linnæan sections: Papilio Priamus, is however, removed from the head of the Equites Trojani, and the precedence given to Papilio Paris.

Papilio Paris is an infect of confiderable beauty. The general colour on the upper furface obscure brown, nearly approaching black, but finely contrasted with brilliant green atoms, that are profusely sprinkled over it. The posterior wings are adorned with a large blue spot, which derives additional lustre from the dusky colour surrounding it. Another species, very similar to Papilio Paris, but without this spot, is also found in China. It is the supposed semale of our species. Fabricius names it Bianor, after Cramer, pl. 103. fig. 6.

PAPILIO CRINO.

SPECIFIC CHARACTER.

Wings tailed, black, entirely covered with resplendent greenish, or golden specks. A band of blue green across the wings. Under side, brown, with lunar marks.

Papilio Crino: alis caudatis nigris: atomis viridi aureis fascia communi cœruleo viridi, posticis subtus lunulis viridibus cœruleis cinereisque. Fab. Ent. Syst. 7. 3. p. 5. 13.

The beauty of the preceding species claims our notice: Papilio Crino, our admiration. Its form is graceful, its colours splendid; and in addition to those attractions, it is extremely rare. We have found an unique specimen of this species in the collection of Mr. Drury, and on that authority we include it as a native of China. It has not been figured by any author. Fabricius describes the same insect under the specific name Crino, erroneously giving its habitat, Africa.

Renealmia exaltata.

Flowered in the fummer of the prefent year, both in the flove of G. Hibberts, Efq. and at Meffrs. Grimwoods and Wyke's, Kenfington. It is a majestic plant, near seven feet in height, and bears a fine pendent group of flowers at the summit.

PAPILIO AGENOR.

SPECIFIC CHARACTER.

Wings indented, black, fanguineous-red at the base: posterior wings white, with black spots.

Papilio Agenor: alis dentatis nigris basi sanguineis: posterio disco albo; maculis nigris.

Lin. Syft. Nat. 2, 747, 14, Fab. Ent. Syft. T. 3, p. 13, 39.

This is one of the largest Chinese Papiliones we are acquainted with. The upper and under surfaces so nearly agree, that we have considered a sigure of the first unnecessary. It is represented with Papilio Coon on the

Plumbago Rofea. Rofe coloured Lead-Wort.

PAPILIO PERANTHUS.

SPECIFIC CHARACTER.

Wings tailed, indented, above black, green at the base: beneath, pale at the ends. Seven lunar marks on the posterior wings.

Papilio Peranthus: alis dentato caudatis nigris fupra bafi viridibus, fubtus apice pallidis: pofticis lunules feptem fulvis. Fab. Ent. Syft. T. 3. p. 1. 44. 15.

Uncommonly scarce, and not hitherto figured. Fabricius has given it as a new species, *Peranthus*, and refers to a specimen in the collection of the Right Hon. Sir J. Banks, Bart. That insect came from Cochin China. We have seen another, which came from Canton: it is represented on a small twig of

Arundo Bambos. Bamboo or Cane.

This well known plant is mentioned by Sir G. Staunton among the most useful productions of China.









Papilio Peranthus.

Lordon, Published or the for direct by Elkenman Japhayay.







Papilio Laomedon .

Landon Published as the Ace directs by E. Denovan April 2.1398.

PAPILIO LAOMEDON.

GENERIC CHARACTER.

Antennæ clubbed at the end. Wings erect when at rest. Fly by day.

SPECIFIC CHARACTER.

Wings indented: anterior pair brown, posterior black: with two red spots on the interior angle.

Papilio Laomedon: alis dentatis: anticis fuscis, posticis nigris: macula duplici rufa anguli ani. Fab. Ent. Syst. 3. p. 1. 12. 35.—Jon. sig. pict. 1. tab. 10.

We have reason to conclude that this superb Butterfly has never been figured by any preceding author. Fabricius described it in his Entomologiæ Systematica from one of the original drawings, in the collection of Mr. Jones, and refers exclusively to that figure, in a synonym to his description. Mr. Jones having obligingly affisted us with his collection of drawings, and given us permission to copy any of them, we cannot withhold enriching our volume with a figure of Papilio Laomedon.

The specimen, from which the drawing was made, was in the collection of Mr. Latham.

PAPILIO COON.

GENERIC CHARACTER.

Antennæ clubbed at the end, Wings erect when at reft. Fly by day.

SPECIFIC CHARACTER.

Anterior wings brown. Posterior wings, and tails, black: white spots at the base; and two yellow spots on the interior angle.

Papilio Coon: alis caudatis concoloribus: anticis fuscis. Posticis nigris basi albo maculaque duplici anguli ani slava. Fab. Ent. Syst. 3. p. 1. 10. 27.—Jon. sig. pict. 1. tab. 36.

This beautiful species is selected from the drawings of Mr. Jones; because we presume that a figure of it has never been published before.

PAPILIO EPIUS.

GENERIC CHARACTER.

Antennæ clubbed at the end. Wings erect when at reft. Fly by day.

SPECIFIC CHARACTER.

Wings indented, brown with yellow fpots: a red fpot on the interior angle of the posterior wings.

Papilio Epius: alis dentatis fuscis flavo maculatis: posticis macula anguli ani rufa. Fab. Ent. Syst. 3.
p. 1. 35. 102.


Papilio Epius and Papilio Demoleus are fo fimilar in their marks and colours, that most authors have confounded one species with the other. Papilio Epius is chiefly distinguished by the red spot in the interior margin of the lower wings, having no blue eye-shaped mark above it.



: Papilio Epius.

* Papilio Demoleus.

London Published as the let direct by Elkenman Feb / 21299.







· Papilio Telamon.

: Papilio Agamemnon.

PAPILIO TELAMON.

TELAMON BUTTERFLY.

GENERIC CHARACTER.

Antennæ increase in bulk towards the extremity, and usually terminate in a kind of club.

Wings, when at rest, erect. Fly in day time.

SPECIFIC CHARACTER.

Wings and tail yellowish, with spots and bands of black. A red streak on the under wings.

Papilio Telamon: alis caudatis concoloribus flavescentibus: maculis fasciis nigris; posticis utrinque striga fanguinea.

The fingular delicacy and beauty of this Papilio is not the only claim it has to the particular attention of Entomologists: it is clearly an undescribed species; and perhaps the only specimen of it yet brought to Europe, is that from which our figure is copied. It was taken near Pekin, by a gentleman in the suite of Earl Macartney, in the late embassy to China; and is at this time in the possession of Mr. Francillon of London, who has kindly permitted us to make drawings and descriptions of this, and every other insect in his magnificent collection that could enhance the value of this publication.

Papilio Telamon bears a diffant refemblance to P. Protefilaus, but a much fironger to P. Ajax: purfuing then the metaphorical method of arranging the butterflies in the Linnæan manner, we have given it the name of the father of Ajax, who was one of the diffinguished Grecian Princes at the fiege of Troy.

PAPILIO AGAMEMNON.

AGAMEMNON BUTTERFLY.

GENERIC CHARACTER.

Antennæ increase in bulk toward the extremity, and usually terminate in a kind of club.

Wings, when at rest, erect. Fly in day time.

SPECIFIC CHARACTER

AND

SYNONYMS.


Upper fide: wings, and tails, black, with green spots. Three lunar red spots on each of the under wings.

Papilio Agamemnon: alis caudatis nigris viridi maculatis: posticis subtus lunulis tribus rufis. Lin. Syst. Nat. 2. 748. 22.

Fab. Ent. Syft. T. 3. p. 1. 33. 98.

Papilio Ægisthus. Cram. Inf. 9. tab. 106. fig. C. D.

Papilio Agamemnon is found in feveral parts of Afia. The under fide is beautifully adorned with a number of bright green fpots of various fizes. The general colour is pale pink, diverfified with shades of chesnut brown. The upper fide is much plainer; the general colour is black, except on the spots, which are green, and precisely agree in shape with those on the under side.





Papilio Menelaus, var.

Papilio_Rhetenor. o.am.

PAPILIO RHETENOR.

SPECIFIC CHARACTER

AND

SYNONYMS.

Wings indented: above shining blue; beneath clouded, and marked with brown spots.

Papilio Rhetenor. Cram. Inf. 6. tab. 15. fig. A. B.

Sulz. Hift. Inf. tab. 13. fig. 1.

Sulz. et Roem. p. 68.

Papilio Menelaus: alis dentatis: fupra cœruleis nitidiffimis, fubtus nebulofis: punctis fufcis.

Linn. Syft. Nat. 2. 748. 20.

Most naturalists are undetermined whether Papilio Menelaus and the supposed variety, Papilio Rhetenor, should not be considered distinct species. The colour of the upper side is nearly the same in both insects; but the under sides are very different. The Linnæan description of the male, P. Menelaus, agrees with Papilio Rhetenor, and probably he considered them the two sexes of one insect. Fabricius would certainly have made a new species of it, if he had not been of the same opinion. We have both insects before us, and should most affuredly make a new species of P. Rhetenor, as Cramer and Sulzer have done, if the authority of the last work of Fabricius were not opposed to us.

Papilio Menelaus is a native of South America. Sulzer's specimen of Papilio Rhetenor came from China. Foreign Entomologists, for this and other reasons, have considered them distinct species. The curious in infects, says Cramer, call this butterfly Le Satiné bleu oblong, to distinguish it from P. Menelaus, or Le Satiné bleu vulgaire.—Cramer has also given a third and much smaller kind of these blue butterflies, Pap. Adonis, which Fabricius also considers a variety of Menelaus.

Whatever effect the artist can produce by a combination of the most brilliant colours employed in painting, he must shrink from comparison of it with this splendid creature. It is impossible to find in any part of the animal creation colours more beautiful or changeable. Pale blue is the principal colour, but new tints meet the eye in every direction, varying from a filvery green to the deepest purple; and the whole surface glittering with the resplendence of highly polished metal.

Thea Laxa.—Bohea, or broad-leaved Tea.

Sir G. Staunton fays the bohea tea is supplied in China from the province of Fochen: the green tea from Kiang-nan. The leaves of these teas vary in some degree in form according to the age of the plant; those of the bohea are the broadest; Thea stricta has much longer leaves, they are lanceolated, and more deeply serrated than those of the bohea. Many authors have considered them varieties of the same species.—Flowers in England in August and September.

PAPILIO OENONE.

GENERIC CHARACTER.

Antennæ clubbed at the end. Wings erect when at reft. Fly by day.

SPECIFIC CHARACTER

AND

SYNONYMS.

Wings indented: yellowish with black margins: the base of the posterior wings bright azure.

Papilio Oenone: alis denticulatis luteis margine nigris: posticis basis cyaneis. Linn. Syst. Nat. 2. 770. 135.

Fab. Ent. Syst. 3. p. 1. 280.


Kleman. Inf. 1. tab. 3. fig. 1, 2.

Found throughout Afia: in China is very common.



· Papilio Oenone. : Papilio Almana.

· Papilio Labentina par







PAPILIO VESTA.

SPECIFIC CHARACTER.

Wings oblong, entire, yellow brown, margined with black fpots.

Papilio Vesta: alis oblongis integerrimis flavescentibus: margine nigro punctato. Fab. Ent. Syst. T. 3. p. 2. 163, 503.

Papilio Vesta is the only insect of the Heliconii division of Butterslies, described by Fabricius as peculiar to China, in his Ent. Syst. It is a rare species, and has not been figured in any work, unless the Papilio Terpsichore of Cramer prove the same species. The Papilio Vesta of that author is a very different insect, being the P. Erato of Fabricius.

PAPILIO PASITHOE.

SPECIFIC CHARACTER.

Wings oblong, entire, black. A central white fpot on each wing, and radiated white marks near the margin. Under fide, posterior pair yellow: a space of blood red colour on the base.

Papilio Pasithoe: alis oblongis nigris albo fubradiatis: puncto centrali albo, posticis subtus luteis basi cruentatis. Linn. Syst. Nat. 2. 755. 53.

Fab. Ent. Syft. T. 3. p. 2. 179. 555.

Papilio Porfenna. Cram. Inf. 4. tab. 43. fig. D. E.

Not so rare as the preceding species; but by no means common. The upper surface is uniformly black, except the radiated white marks, similar, to those seen on the under surface of the anterior wings; and the white central spot. It exhibits no trace of the red, and scarcely a tinge of the yellow colour so conspicuous on the under surface.

F

PAPILIO HYPARETE.

SPECIFIC CHARACTER.

Wings entire, oblong, white, veined with black. Under fide of the posterior pair yellow, with a border of red spots.

Papilio Hyparete: alis oblongis integerrimis albis nigro venofis: pofticis fubtus flavis: margine rubro maculato. Linn. Syst. Nat. 2. 763. 92.

Fab. Ent. Syft. T. 3. p. 2. 178. 554.

Papilio Antonöe. Cram. Inf. 16. tab. 187. fig. C. D.

Papilio Eucharis. Cram. Inf. 17. tab. 201. fig. B. C.

We have two forts of this species; one with the marginal row of red spots on the posterior wings, disposed in a deep border of black; the other has the red spots on a whitish ground. They are certainly the two sexes of Papilio Hyparete. Found near Canton in China.

Sophora Japonica.—Shining-leav'd Sophora.

An elegant and valuable timber tree. Sir G. Staunton speaks of it as very frequent in China. It is noticed in the lifts of plants collected in the journey between Pekin and Zhe-hol in Tartary, and in the provinces of Shan-tung and Kiang-nan, and also in the province of Pe-che-lee.





PAPILIO PRYANTHE.

Antennæ increase in bulk towards the extremity, and usually terminate in a kind of club. Wings, when at rest, erect. Fly in day-time.

SPECIFIC CHARACTER.

Wings roundish, above white, tips black, with a black spot in the middle of the wings. Beneath a pale ash colour, with waved lines and a fulvous spot.

Papilio Pyranthe. Alis rotundatis albis puncto apiceque nigris, fubtus cinereo undatis: puncto fulvo.

Lin. Syft. Nat. 2, 763, 98.

This is a rare species, and has not been figured by any author.

PAPILIO PHILEA.

SPECIFIC CHARACTER.

Wings entire, fomewhat angulated, yellow. A large orange fpot on the anterior pair. Margin of the pofterior pair orange.

Papilio Philea: alis integerrimis angulatis flavis: anticis macula pofticis limbo luteis.

Linn. Syft. Nat. 2. 764. 104.

Roefel has given a figure of this beautiful butterfly in the fourth volume of the Infecten Belluftigung, and calls it die indianische Goldborte; Linnæus also notes it as an Indian species. Our specimen was received from China by the late Mr. Keate: it is represented with the preceding species on the

Melastoma Chinensis,

A plant recently introduced into this country.

PAPILIO GLAUCIPPE.

SPECIFIC CHARACTER.

Wings rounded, entire, white. Anterior pair, black at the ends (and furrounding an orange fpace).

Under fide greyish with brown waves.

Papilio Glaucippe: alis rotundatis integerrimis albis: anticis apice nigris, posticis subtus cinereis susce undatis. Linn. Syst. Nat. 2. 762. 89.

Papilio Glaucippe is an elegant infect: very common in China, and it is faid, in fome adjacent parts of Afia alfo. The Papilio Callirrhoe of Linnæus is considered as the female of this species: few authors deem it more than a variety (β) .

PAPILIO SESIA.

SPECIFIC CHARACTER.

Wings roundish, above yellowish, black at the ends, with a large fulvous orange spot. Beneath, seven white eye-spots.

Papilio Sesia: alis rotundatis albis b apice (medio fulvo) nigris: posticis subtus maculis septem ocellaribus albis. Fab. Ent. Syst. T. 3. p. 1. 203. 636.

Papilio Pyrene. Linn. Syft. Nat. 2. 161. 86.


Fabricius, as well as Linnæus, confiders this as an American infect: Mr. Drury received his fpecimens from China, and that represented in the annexed plate was also brought from the same country by the late Mr. Ellis.

Limodorum Tankervillia,

An elegant and much admired production of China: it was introduced in 1788, and named in compliment to the Earl of Tankerville.

b Alæ anticæ flavissimæ, &c. vide general description.









London Published as the Act directs by E. Dosovan Feb ! 1. 1798.

PAPILIO JAIRUS.

JAIRUS BUTTERFLY.

GENERIC CHARACTER.

Antennæ increase in bulk towards the extremity, and usually terminate in a kind of club.

Wings, when at rest, erect. Fly in day time.

SPECIFIC CHARACTER

AND

SYNONYMS.

Wings entire, brown: middle of the lower wings white; on the upper furface of each, near the posterior margin, an eye-shaped spot; on the under surface two.

Parilio Jairus: F. alis integerrimis fuscis: posticis disco albo supra ocello subtus duobus.—Fab. Ent.

Syst. T. 3. pl. 1. p. 54. 168.

Papilio Jairus. Cram Pap. tab. 6. fig. A. B.—tab. 185. fig. A. B. C. Clerk Icon. tab. 29. fig. 3.

A fpecimen of this extremely rare Butterfly is contained in the collection of Dr. Hunter; a fragment in the British Museum; and one in fine preservation in the collection of Mr. Francillon. Except these, and the specimens from which the figures in the annexed plate are copied, we have never seen it in any cabinet whatever.

It has been figured only by two authors, Clerk in his Icones infectorum rariorum, and Cramer in his Papillons exotique. The figures of Clerk and Cramer do not strictly agree: we observe those of the first much lighter coloured, and the white space on the upper wings considerably larger than in any of the figures in Cramer's plates.

Fabricius says it is a native of the East Indies. One specimen figured by Cramer was brought from the isle of Amboyna. It seems therefore, not peculiar, like some insects, to China.

The copy of this exceedingly fearce work of Clerk, which we had an opportunity of comparing with Cramer's works, is in the invaluable library of the Right Hon. Sir J. Banks, Bart.

PAPILIO ANTIOCHUS.

ANTIOCHUS BUTTERFLY.

GENERIC CHARACTER.

Antennæ increase in bulk towards the extremities, and usually terminate in a kind of club.

Wings, when at rest, erect. Fly in day time.

SPECIFIC CHARACTER

AND

SYNONYMS.

Wings entire, roundish, black: a broad band of bright yellow brown continued across the upper surface of the wings.

Papilio Antiochus: P. F. alis integerimis rotundatis nigris: fupra fascia communi fulva. Linn. Mant.

1. 537.—Fab. Ent. Syst. T. 3. p. 2.—44. 134.

Papilio Eupalemon. Cram. Pap. 12. tab. 143. fig. B. C.

Drury Inf. 3. tab. 7. fig. 3. 4.


Aubent. Mifcell. tab. 68. fig. 3. 4.

Papilio Antiochus is very rare in European cabinets of insects. The specimen figured by Drury came from the Brazils, and Cramer's from Surinam. Fabricius describes it as a native of China. The insect figured in the collection of drawings of Mr. Jones of Chelsea, was a native of China, as well as the specimen in our collection.



: Papilio Jacintha: Papilio Antiochus.

London Published as the let direct by ED ensem December 1174







Papilio Gambriciis.

T. ondon Paklished underliet directly EDenman Harch 2.5709.

PAPILIO JACINTHA.

SPECIFIC CHARACTER.

Wings brown, scalloped. A row of white spots on the anterior pair. Ends of the posterior pair white.

Papilio Jacintha: alis repando dentatis fuscis: anticis striga punctorum alborum, posticis apice albis.

Fab. Ent. Syst. T. 3. p. 1. 60. 187.

This curious butterfly was found in the province of Pe-tche-lee, in China, by the gentleman who difcovered the elegant Papilio, *Telamon*, near Pekin; and has otherwife, by his attention to entomology, added fome interesting species to our catalogue of Chinese insects. P. Jacintha is represented with P. Antiochus, on a leaf of the

Urtica Nivea. White Nettle .

PAPILIO GAMBRISIUS.

SPECIFIC CHARACTER.

Wings indented. Above black, with fpots and ftreaks of green; and a band of white transparent spots on the anterior pair.

Papilio Gambrisius: alis dentatis supra nigris viridi maculatis striatisque: anticis fascia maculari nivea.

Fab. Ent. Syst. T. 3. p. 2. 264.

A specimen of this very rare Papilio, was taken in one of the small islands on the eastern coast of China, and is in the possession of Mr. Francillon. Sir J. Banks, Bart. has a specimen of it from another part of the East Indies.

e Sir G. Staunton speaks of a cloth that the Chinese manusacture from the fibres of a dead nettle. Query, Is this the species employed for that purpose? no other is noticed by that author in the lifts of plants collected in China.

The nettle is of general use in Russian Tartary also; the Kuriles, and other Siberian tribes, make cloth, cordage, thread, &c. of it. Gordon, &c.

PAPILIO LUBENTINA.

SPECIFIC CHARACTER.

Wings dentated, dark, greenish, with spots of white, black, and red.

Papilio Lubentina: alis dentatis obscure virescentibus albo nigro rusoque maculatis.

Fab. Ent. Syst. T. 3. p. 2. 121. 370.

PAPILIO LUBENTINA, is figured only in the works of Cramer: his fpecimen is not precifely like ours, but agrees in all the effential peculiarities, and is unquestionably the same species. The semitransparent spots on the anterior wings are much larger in Cramer's figure than in the infect before us.

PAPILIO LEUCOTHOE.

SPECIFIC CHARACTER.

Wings dentated, above brown, with three white bands across. Beneath yellow brown, with three similar white bands, and black characters.

Papilio Leucothoë: alis dentatis, fupra fufcis: fafciis tribus albis: fubtus luteis: fafciis tribus albis nigro notatis. Linn. Syft. Nat. 2. 780. 179.

This pretty infect is very common about Canton, in China.

PAPILIO POLYXENA.

SPECIFIC CHARACTER.


Wings dentated: above brown, with three white bands across. Beneath yellowish orange, with three white bands: a row of black spots on the lower one.


Papilio Polyxena: alis dentatis, fupra fuscis: fubtus luteis: fasciis utrinque tribus albis, posteriore fubtus nigro-punctata.

This feems to have been confounded with the preceding species both by Linnæus and Fabricius. Linnæus first describes it as the semale, and afterwards as a variety of it; but it certainly is a very distinct species.



London Baklished as the Act directs by EDengram Lonis 1799.







Papilio Bernardus.

I and on Published as the det directs by E. Donesan May 1. 198.

PAPILIO BERNARDUS.

SPECIFIC CHARACTER.

Orange colour. A broad bar of yellow across the anterior pair of wings: ends black. Posterior pair with two tails, and a festoon of black spots, with an eye in the centre of each.

Papilio Bernardus: alis caudatis fulvis: anticis apice atris; fascia flava, posticis striga punctorum ocellatorum. Fab. Ent. Syst. 2. p. 1. 71. 223

This uncommonly rare Chinese buttersly has certainly never been figured in any preceding work. It will also be probably new to the collectors of exotic insects, in this country at least. Fabricius described it only from the accurate drawings of Mr. Jones, as appears by the reference added to the description in his Entomologia Systematica.

Camellia Japonica.—Japan Rofe.

A native of Japan and China. It bears bloffoms from January to May. This is a lofty and magnificent plant, rifing to the height of feveral feet: there is a variety of it with double flowers, perfectly white; and another in which the flowers are variegated with white and red.

PAPILIO ERYMANTHIS. YELLOW JAGGED-BAR BUTTERFLY.

SPECIFIC CHARACTER.

Wings indented, brown: a broad bar of yellowish colour across the anterior pair; exterior ends black.

Lunar marks on the posterior pair, and a row of eye-shaped spots beneath.

Papilio Erymanthis: alis dentatis fulvis: anticis apice nigris; fascia flava, posticis supra lunulis, subtus striga punctorum ocellatorum. Fab. Ent. Syst. 2. part 1. p. 139. 427.

It is the rarity, and not the beauty of this butterfly, which has induced us to add it to our felection. We apprehend it is far from common in China, being very feldom fent to Europe among the infects of that country.

D

PAPILIO ORYTHIA.

VARIABLE BLUE AND BROWN BUTTERFLY.

SPECIFIC CHARACTER

AND

SYNONYMS.

Wings indented, brown. On the upper fide two eyes in each, and another on the under fide of the anterior wings.

Papilio Orythia: alis dentatis fuscis: omnibus supra ocellis duobus, anticis subtus unico. Lin. Syst. Nat. 2. 770. 130.

Fab. Syft. Ent. 2. part 1. p. 91. 284.

The varieties of Papilio Orythia are numerous, and feem to differ according to climate of the countries of which they are natives. It is common in North America, Jamaica, India, &c. The variety from North America is almost wholly brown, and those from Jamaica have less blue in the disk of the lower wings, than those from China.

Papilio Clelia of Cramer, which is found on the coast of Guinea, has been supposed a variety of Papilio Orythia. Fabricius, in the Entomologia Systematica, has made it a distinct species. It greatly resembles P. Orythia, but has no more blue colour on the posterior wings than is concentrated in a large spot near the base.

PAPILIO ALMANA.

DOUBLE-EYE ANGULATED BUTTERFLY.

SPECIFIC CHARACTER

AND


SYNONYMS.

Wings angulated, brown, varied with black marks; a large eye with two pupils in the middle, on the posterior wings. Beneath, entirely brown.

Papilio Almana: alis angulatis fulvis nigro maculatis: pofficis ocello majori; pupilla gemina, omnibus fubtus brunneis. Linn. Syft. Nat. 2. 769.

Fab. Ent. Syft. 3. p. 1. 278.

The angulated form of the wings of this butterfly gives it a remarkable appearance. The eyes on the wings fomewhat refemble those of the Peacock butterfly, to which, in some other respects, it bears no distant similitude. It is common in China; Fabricius gives its habitat Asia.





PAPILIO MÆCENAS.

GENERIC CHARACTER.

Antennæ increase in bulk towards the extremity, and usually terminate in a kind of club.

Wings, when at rest, erect. Fly in day-time.

Plebeii rurales. Linn. c

SPECIFIC CHARACTER.

Upper fide black, difk blue. Two tails to each posterior wing. Under fide clouded with brown.

Hesperia Mæcenas: alis bicaudatis atris: disco cœruleo, subtus brunneo nebulosis.

Fab. Ent. Syst. T. 1. p. l. 271. 45.

We are indebted for this rarity to T. Marsham, Esq. It is described by Fabricius from an original drawing d, and has not been figured by any author.

c Plebeii rurales, alis maculis obscurioribus. Spots on the wings obscure, or not transparent.—The plebeii is the last family of Butterslies in the Linnman system, and is divided into two sections, rurales and urbicolas. Fabricius removes this family from the Papiliones (Butterslies) to a new genus Hesperia. This genus is however divided after the Linnman method into two sections, rurales and urbicolas, and contain nearly the whole of the Linnman plebeii, in addition to the later discovered species. The effential character of the Hesperia is, Palpi two, compressed, hairy, apex cylindrical, naked. Antennae clubbed, oblong, often crooked at the extremity.

4 Jon. fig. pill. 6. tab. 3. fig. 2.

PAPILIO ATYMNUS.

SPECIFIC CHARACTER.

Wings bright brown. Anterior pair black at the tips. Posterior pair furnished with tails.


Hesperia Atymnus: alis caudatis fulvis: anticis apice nigris.

Fab. Ent. Syft. 1. p. 1. 283. 88.

This is also a scarce species. Our specimen is from the collection of the late Duchess Dowager of Portland, who procured it from China. Another specimen in the cabinet of Sir J. Banks, Bart. is from Siam.

Hemerocallis Japonica.

Brought from China by Mr. Slater, and flowered in July 1798 in the green-house of G. Hibberts, Esq. Clapham.





· Sphinx Nechus. Sphinx Polymena. :

London Published as the let directs by EDonovan July 15798.

SPHINX NECHUS.

SPECIFIC CHARACTER.

Wings entire: anterior pair greenish, with testaceous marks. Posterior pair black; a band of yellow spots across the wings, and a single spot near the base.

Sphinx Nechus: alis integris: anticis viridibus; firiga teftacea, posticis nigris: maculis baseos fasciaque flavis. Fab. Ent. Syst. 3. p. 1. 377. 63.

The number of Chinese species of this genus, already described, is very limited: our present insect is the largest of them; but, as this is inferior in size to several kinds found in Europe, we conceive there must remain many larger species of the genus unknown to collectors of foreign insects, and yet very common in China. In the latter part of Sir G. Staunton's work, that author mentions the larva of a Sphinx Moth which furnish an article for the table of the Chinese. We regret that the indefinite expression cannot affist us to determine the species, and scarcely the genus, of the insect alluded to e.

The specimen figured in the annexed plate, is in the collection of Mr. Francillon, who received it from China. A small variety of the same species is found in North America, and sigured by Cramer. Sphinx Batus and Sphinx Gnoma are nearly allied to this insect, particularly the former; both are found in different parts of the East Indies.

e European naturalists are entirely ignorant of the Chinese insects in the state of larva and pupa, if we except a sew species of the Cimices, Cicada, and some altogether uninteresting insects, that have been accidentally brought among others from that country. Hence it must remain undetermined whether they correspond in form with those of other parts of the world. It is, however, highly probable, from their great affinity to those, in the perfect state, that in the state of larva they may also agree. The extensive collection of the larva of sphinges made by Mr. Abbot in North America affords no singularly constructed animal distinct from those sound in Europe; they vary indeed in their colours, but preserve uniformly the characters sound on the same genus in other countries. We noticed among the drawings of the late Mr. Bradshaw the figure of a Chinese sphinx, apparently S. Hylas, together with a larva similar to that of the Sphinx Stellatarum: it was green, and, like all the known larva of the genus, (except the Asserted division) was perfectly free from hairs: it was also surnished with a horn at the posterior part of the body.

SPHINX POLYMENA.

SPECIFIC CHARACTER.

Wings black, with three yellow fpots on the anterior pair, and two on the posterior pair. Abdomen belted with two bright red bands.

SPHINX POLYMENA: Lin. Syft. Nat. 806. 40.

Zygæna Polymena: nigra alis maculis luteis anticarum tribus posticarum duobus. Abdomine cingulis

duobus coccineis. Fab. Ent. Syft. 3. p. 1. 396. 34.

We suspect this beautiful creature is scarce in China; at least it is very rarely found among the insects brought from that country. It is figured on the plate with the

Rofa femperflorens—Ever-blowing China Rofe.

A plant lately introduced from China, but thrives well in this climate, and bears a beautiful deep red flower, throughout most part of the summer season.

SPHINX HYLAS.

SPECIFIC CHARACTER.

Wings transparent. Abdomen green, with a purple belt round the middle. Apex bearded.

SPHINX HYLAS: Linn. Mant. 1. 539.

Sefia Hylas: alis feneftratis, abdomine barbato viridi: cingulo purpureo. Fab. Ent. Syft. 3. p. 1. 379. 3.

The Sphinges with transparent wings may be separated from the others with much propriety: Fabricius includes them in his new genus Sesia; but that genus is not formed exclusively of such insects, his essential characters being taken from the structure of the palpi, tongue, and antennæ. The Sphinx suciformis found in England somewhat resembles this Chinese species.



-:: bifasciata!

SPHINX THALLO.

SPECIFIC CHARACTER

ANO

SYNONYMS.

Wings oblong, entire black, two bars of white on the anterior wings, and a yellow space on the posterior pair.

Papilio Thallo: alis oblongis integerrimis nigris: anticis fafciis duabus, posticis unica flavis. Linn.

Syft. Nat. 2. 756. 62.

Papilio Thallo:

Fab. Ent. Syft. 3. p. 1. 173. 537.

Sphinx pectinicornis: Linn. Syft. Nat. 2. 807. 44?

Fab. Ent. Syft. 3. p. 1. 399. 44?

It is altogether perplexing, and mysterious to us, that Fabricius, throughout all his works, and even in his last enlarged and corrected systematic arrangement, has given a Papilio Thallo in the Heliconii division of that genus, when it is clear no fuch Papilio has existence. Linnæus has, undoubtedly, so named an infeet figured in a plate of Edwards's birds, tab. 226, with a reference to that work; yet it is certain the infect there figured is not a butterfly, but a lepidopterous Infect, to which antennæ of dubious ftructure have been added, to perfect its appearance f. No later author has figured the fame species; and specimens of it being extremely fcarce, it is probable Fabricius has been unable to detect the error. In admitting this, however, we must necessarily notice the synonyms of Sphinx pectinicornis, where he also quotes the same figure in the works of Edwards. Thus we find, throughout all the editions of the Fabrician System, a Papilio Thallo and Sphinx pectinicornis formed of the folitary figure of one infect. Cramer has finally increased the ambiguity of its genera, by making an infect of close affinity, perhaps the same species, a Phalæna. Vide P. Tiberina.

f Coincident observations on perfect specimens of several insects, nearly allied to this species, tend to convince us, that the artift had only a mutilated specimen to copy from. Indeed, though the antennæ in his figure are terminated in a kind of capitulum, like that of the butterflies, the filaments are somewhat jagged, as if intended to appear slightly pectinated. Cramer notices the very close affinity between his infect, P. Tiberina, and that in Edwards's work; which, he observes, differs neither in form nor colour, but only in having clubbed antennæ: those parts of his insect being pectinated or feathered. - "Mr. Edwards a repréfenté un Papillon, qui ne differe de celui-ci, ni pour la coleur, ni pour la deffein, mais qui a des antennes à boutons. Celui que nous donnons ici eft gravé d'apres une Phalêne qui a des antennes plumacées," &c. Cram. C. D. Pl. 32.

Edwards calls this infect the little black and white Butcher fly, because it is figured on a plate with the black and white Butcher bird.

With specimens of both the insects figured by Edwards and Cramer before us, we cannot hesitate to consider them Sphinges of the Adscitæ s family, and of the Zygæna of Fabricius. In removing our insect to the sphinx genus, we have retained the specific name Thallo, as more likely to denote the species than any newly adopted name. The insect figured by Cramer does not precisely agree with Edwards's figure. In the first, the disk of the posterior wings are yellowish, with a deep border of black: in the other, the yellow occupies only a space near the base, and forms a semi-lunar mark near the anterior margin of those wings. We suspect, with Cramer, that they are but the two sexes of one species. Cramer says both his specimens came from China, from whence our insects were also received.

SPHINX RUFICOLLIS.

SPECIFIC CHARACTER.

Wings entire; black purple, a femicircular yellowish band communicating across all the wings; and two
spots of the same colour near the apex. Collar reddish.

Sphinx ruficollis: alis integerrimis nigro-purpurescentibus fascia communi maculisque duabus flavis, thorace antice brunneo.

This and the following species are undoubted nondescripts: both specimens are in the collection of Mr. Francillon, who received them from China.

SPHINX BIFASCIATA.

SPECIFIC CHARACTER.

Wings fulvous or orange, a black bar across the middle of the anterior wings: tips black.

Sphinx bifasciata: alis fulvis fascia apiceque nigris.

Sphinx Hylas, Thallo, ruficollis, and the elegant little species Sphinx bifasciata, are represented on the plate with the

Thuja Orientalis.—China Abor-vitæ Tree,

An ornamental evergreen, much efteemed by the Chinese, and very frequently represented in their landfcapes. Sir G. Staunton remarks in the account of the journey from Pekin to Canton, that great quantities of this plant grew to a prodigious height in the valley in which stands the city of Yen-choo-foo.

E The Adfeitæ differ from the other sphinges, in their general appearance, but have all the characters affigned to that genus.



Phalana Atlas.

PHALÆNA ATLAS.

GENERIC CHARACTER.

Antennæ fetaceous. Wings, in general deflexed when at reft. Fly by night.

Bombyx.

SPECIFIC CHARACTER.

Wings much falcated or hooked. Colour, yellow brown, varied. A transparent spot in the middle of each wing; with a smaller one next that on the anterior pair.

PHALENA ATLAS: alis patentibis falcatis luteo variis: macula fenestrata anticis sesquialtera.

Lin. Syft. Nat. T. 2. 808. 1. Fab. Ent. Syft. T. 3. p. 1. 407. 1.

The Phalænæ are a tribe of infects remarkable for the neatness and simplicity of their colours. Their elegancies consist in the infinite variety, and delicacy of intermingled tints: the contrast of spots, specklings, and lineations, which constitute the minutiæ of infect beauty. Some species are to be excepted in this remark; the larger kinds are often gaudy, and the smallest exhibit a display of the richest colours, fancifully disposed, and most elegantly diversified.

The European species are numerous, and pretty well ascertained: those of remote countries remain in great obscurity. The species inhabiting China are almost unknown; for the latest systematic writer describes not more than twenty species in all the cabinets in Europe. From this scanty number a few are selected to illustrate the genus, and if these appear descient in point of interest or variety, it may stimulate others to collect new species, whenever an opportunity occurs. The Phalæna tribe, not only of China, but every country, except of Europe, are a desideratum of entomology. In Europe the number of this tribe exceeds that of any other: on the contrary, the extra-European species are comparatively the most inconsiderable of our acquisitions. The Papiliones, or Butterslies, are a showy and lively race: they sport in the open fields in day, and attract the traveller's curiosity; hence our cabinets abound with them. But the moth, infinitely more numerous, and not less pleasing, is seldom seen; in the gloomines of its dispo-

[•] Fab. Ent. Syft. These are chiefly described from insects in the collection of Monfon Londini, of which no figures are extant, and the collection unknown.

fitions, it feeks the obscurity of the forest in the day, and only ventures on the wing when the sun is down. In Europe we visit its nocturnal haunts without difficulty or dread; but in hotter climates these are oftentimes impenetrable, or the lurking places of serocious animals; and sew will expose themselves to their attacks, to increase the catalogue of exotic Phalænæ P.

Phalæna Atlas is the first species we have to notice. It is the largest of the moth tribe 9, and is, indeed, a gigantic creature. The species is common, but not peculiar to China, being sound in other parts of Asia, and in America. The influence of climate is easily traced in the varieties from different countries; that from Surinam is the largest, and of the deepest colours. The Chinese kind is the next in size; the colours incline to orange, and the anterior wings are more falcated or hooked at the ends. We have two other Asiatic varieties still smaller, with the wings extremely falcated.

The larva of Phalæna Atlas is figured by M. Merian, in the Infecta Surinamenfia, plate 52: it is about four inches in length, green, with a yellow firipe difposed longitudinally. Upon each segment are four distinct round tubercles of a coral-like orange colour, which are surrounded with very delicate hairs. The pupa is large, and is inclosed in a web of an ochre colour. The filk of this web is of a strong texture, and it has been imagined, if woven, would be superior in durability to that of the common filk worm. Seba has also represented the larva at fig. 1. plate 57. vol. 4. Thesaurus Naturæ. It is nearly fix inches in length, and bulky in proportion; the Phalæna is also larger than that figured by Merian, which is a

P The far greater number of Phalænæ can only be taken in the woods at night. This is termed mothing by collectors. The moths begin to ftir about twilight, and when almost dark, commence their flight. The collector is furnished with a large gauze folding-net, in which the infects are caught indifcriminately, for it is impossible to distinguish one species from another, and often is so dark, that the object itself can barely be discerned. Different species have their favourite haunts, some the lanes, and skirts of woods, but many of them prefer the open breaks in the most retired places. As it would be unsafe, or impossible, to penetrate the woods in many countries, it it better to collect the larva, or caterpillar, for these may be found on the trees in day-time, and if kept in little gauze cages, and carefully fed, will change into chrysalis, and produce the fly. This is certainly tedious, and sew travellers will divert their attention from more important observations; but were they to appropriate their leisure to this branch of science, they would materially improve entomology. Mr. Abbot has investigated a small district of Georgia, in North America, in this manner, and our cabinet is indebted to his labours for several hundred species, altogether new in Europe. The reader may estimate the importance of these discoveries, by referring to the two expensive volumes of North American Insects, lately published; and reflecting, that the originals of all the species included in that work, are but a small selection from those he has surnished us with. Viewing these as the result of one man's research, in an inconsiderable portion of North America, what a variety of new and splendid kinds would be the reward of those, who should explore the more genial regions of Asia, Asia, Asia on an asouth America, with equal diligence and information!

We have hazarded an affertion which may feem inadmiffible, that the Phalænæ are infinitely more numerous than the Papiliones, or any other tribe of infects. Not that we poffers more; but because, in every country that has been investigated, experience justifies such opinion. For instance, in Great Britain, we have only fixty Papiliones, and by mere accident two or three local species have lately been added; of the Phalænæ we have more than 900. The same comparative proportion is observed throughout the countries of the European continent; and it is singularly analogous, that our opinion is confirmed, by the recent discoveries of Mr. Abbot in America also.

9 When Linnæus described it, sew of the very large species of Phalæna were known. We have two species from the interior of Africa, that are larger than the Chinese var. of Atlas, and several others searcely inferior in magnitude.

fmall specimen of the Surinam kind. According to Merian, there are three broods of this insect in a year; they are very common, and feed on the orange trees. Linnæus says, they adhere so tenaciously to the leaves, that they can scarcely be taken off r.

The common filk worm, or Phalæna Mori, is of this family, and merits observation as a native of China. The art of weaving its threads into filk is of the earliest date. The discovery is attributed to the Seres, a people of the East Indies, supposed the Chinese. In the days of Solomon, we are told, a woman named Pamphilia, of the Island of Co, was skilled in the art of making cloth, of the filk brought from the country of the Seres. The most ancient of the Chinese writers ascribe the invention to one of the women of the emperor Hoang ti, named Si ling, and in honour Yuen feit. When Rome degenerated into voluptuousness, Persia, its dependency, furnished this article of luxury; but it is supposed they were indebted to the Chinese for it, and being supplied only in small quantities, it was consequently dear. In Rome it was so searce, as to be worn only by persons of the first distinction.

The Chinese historians affirm, that the discovery was considered at first of such importance, that all the women in the palace of the emperor were engaged in rearing the insect, and weaving its filk. In after times, the filk of China was a principal article of commerce, but latterly its value has been materially lessened, by the culture and sabrication of filk in other countries. As the Chinese know little of the use of linen, the filk is a staple article of their own consumption. The jesuit missionaries mention several forts of it, in use among the Chinese; some admired for beauty, and others for durability. It is generally supposed these are not merely the effect of different manusacture, but are the produce of distinct insects. Sir G. Staunton speaks of the culture of filk worms in China, but only of the common fort. It will gratify curiosity, if not prove advantageous, should suture observers ascertain what kind of

- Larva verticillata verrucis pilofis nec folliculos grandes, tenaces, vix extricandos. Linn. Syft. Nat.
- ² Velleraque ut foliis depectant tenuia Seres. Virg. Georg. II. 122.
- [†] Du Halde, Des Soyeries. Les plus anciens écrivans de cet Empire, en attribuent la découverte à une des femmes de l'Empereur Hoang ti, nommée Si ling, et surnommée par honneur Yuen fei.
- "M. Merian fays, in the defeription of the Surinam variety of Phalæna Atlas: "Telam ducunt fortem, quare bonum fore fericum rata, iftius aliquam collegi copiam & in Belgium transmis, ubi eadem optima judicata est: ut itaque, si quis Erucas istas congregandi laborem non detrectaverit, et bonae notae bombycem, et maximum hine lucrum sibi comparare posset." The thread of which this coccon's web is composed, is so strong, that it has been imagined it would make good silk. I have brought some of it into Holland, which has been esteemed such. So that if any one would take the trouble to collect a number of these caterpillars, they would be found good silk worms, and produce great profit. Merian.—Abbot informs us, the Moths of the Emperor tribe, in general, are called silk worms by the people of Georgia, and in the description of Phalæna eccropia, is still more explicit: for he says, "the caterpillar spins on a twig. The outside web is coarse, the inner covered with silk, like a filk worm's coccon. It is said this silk has been carded, spun, and made into stockings, and that it will wash like linen."

 Abbot's Inst. by Dr. J. E. Smith.—These insects are all of the same natural order, P. Cecropia is rather smaller, but very similar to P. Atlas, and this information at least corroborates the affection of Merian.

An opinion, that the Chinese rear several kinds of infects for the sake of their silk, has long been prevalent. Dr. Lettsom proposes a query on this subject, "Which species of moth or butterfly is it, the caterpillar of which, in China, affords that strong grey kind of silk, and how is it manufactured or wore? How are these silk worms or caterpillars preserved, sed, and managed? The introduction of such a new silk into England would be a useful acquisition, and redeem entomology from the

infects the Chinese appropriate to making filk, and whether P. Atlas is of the number, as has been conjectured.

cenfure it is now branded with, of being a mere curiofity, void of any real utility "." If Leffer and Lyanet are to be relied on, The Theologie des Infeders answers this query. " At this day there are to be found in China, in the province of Canton, filk worms in a wild flate, which, without any care being taken of them, make in the woods a kind of filk which the inhabitants afterwards gather from the trees. It is grey, without luftre, and is used to make a very thick and ftrong cloth, named there Kien Tcheon. It may be washed like linen cloth, and does not stain." A Gentleman resident in the East Indies, speaks of a large Phalæna, producing filk in that country. "We have a beautiful filk worm north-eaft of Bengal, that feeds on the Ricinus, whence I call it Phalæna Ricini: it is fea green, with foft fpines, very large, and voracious, and fpins a coarfe, but ftrong and useful filk. The moth is of great fize, with elegant dark plumage. Is it known to European Naturalists?" - In a collection of papers published by Dr. Anderson, in Madras, 1788, 1789 .- M. Le Bon, Reaumur, Roefel, and feveral others, have attempted to weave the filk of fpiders, as a fubfitute for that of filk worms, but their experiments rather amufe, and point out the ingenuity of the propofers, than promife to be ufeful; for after many trials, it appears that the filk of spiders would be inferior in luftre, and far more expensive than that of filk worms. Sir G. Staunton alludes to these experiments in his description of the Java forefts. "In fome open fpots were found webs of fpiders, woven with threads of fo ftrong a texture, as not eafily to be divided without a cutting inftrument; they feemed to render feafible the idea of him, who, in the fouthern provinces of Europe, proposed a manufacture from spiders' threads; which is so ridiculous to the eyes of those who have only viewed the flimfy webs fuch infects fpin in England."-Many other fubftances of a foft texture have also been wrought into a variety of trifling articles, as gloves, flockings, &c. of the fibres of Asbeftos earth, or mountain flax, beard of the large Pinna shell, &c. &c.

* Naturalist and Traveller's Companion, 1"74.





* Phalana Bubo. Lhalana zonaria: **Dhalana pagaria.



LEPIDOPTERA.



* Phalana militaris. * Phalana lectric.

LEPIDOPTERA.

PHALÆNA MILITARIS.

SPECIFIC CHARACTER.

Wings fpread. Yellow, with violet fpots: ends of the wings violet, with white fpots.

Phalæna militaris: alis patulis concoloribus luteis: apice maculifque violaceis, anticis extus albo maculatis. Linn. Syft. Nat. 2. 811. 12.

Fab. Ent. Syft. 3. p. 1. 416. 33.

This, and the following species, are scarce. Mr. Drury informs us, he never procured but a fingle specimen of Phalæna lectrix, in the course of thirty years collecting infects.

PHALÆNA LECTRIX.

SPECIFIC CHARACTER.

Wings incumbent, black: anterior pair spotted with blue, yellow, and white; posterior pair spotted with red and white.

PHALÆNA LECTRIX: alis incumbentibus nigris: maculis cœruleis flavis albifque, poficis rubro alboque maculatis. Linn. Syft. Nat. 2. 334. 83.

Fab. Ent. Syft. Sp. 1. p. 475. 212.

PHALÆNA BUBO.

Noctua.

SPECIFIC CHARACTER.

Wings spread, dentated brown, with black indulations. A large bright brown eye in the middle of the anterior wings.

Phalæna bubo: alis patulis dentatis fufcis nigro undulatis: anticis macula magna ocellari brunnea.

Fab. Ent. Syft. T. 3. p. 2. 9. 4.

Phalæna Macrops. Linn. Syst. Nat. 3. 225.

This is the largest of the Chinese Noctuæ; some very similar species, but without the orange eye, and of a smaller size, are peculiar to China.

LEPIDOPTERA.

PHALÆNA PAGARIA.

Geometræ.

SPECIFIC CHARACTER.

Antennæ pectinated. Wings roundish, blue black, with a row of white spots next the tips of the wings.

PHALÆNA PAGARIA: pectinicornis alis rotundatis cœruleo nigris: fascia macularia apicis alba.

Fab. Ent. Syst. T. 3. p. 2. 153. 85.


PHALÆNA ZONARIA.

SPECIFIC CHARACTER.

Antennæ fetaceous. Wings green, deeply bordered with pale red. A green fpot on the exterior margin of each wing.

PHALÆNA ZONARIA: alis viridibus margine posteriore lato rufescente, singulis macula marginali viridi.

Not described in any preceding work.







+ Aeshna davata.

: Libellula indica.

Libellula 6 maculata.

London Published as the Actionets by EDoneron July 2 1298.

LIBELLULA CLAVATA.

GENERIC CHARACTER.

Mouth furnished with more than two jaws. Antennæ like a briftle or hair, very short. Tail of the male armed with a pair of forceps.

SPECIFIC CHARACTER.

Abdomen clubbed at the end, gibbous or thick at the base. Body variegated with green, and deep brown stripes.

Aeshna Clavata: abdomine clavato basi gibbo, corpore susce viridique variegato. Fab. Spec. Inf. 1.
p. 526. 4.—Ent. Syst. 2. 385. 4.

This infect is described by Fabricius, and in conformity with his new system, is termed Aestina clavatac. We reject his generic definition, because it clearly belongs to the Linnæan Libellulæ. Libellulæ clavata must be placed with the European L. grandis and forcipata.—It is scarce, and has never been figured before.

LIBELLULA INDICA.

SPECIFIC CHARACTER.

Wings yellow, barred with brown, changeable to bright purple. Apex of the anterior pair white. Pofterior pair blue at the base.

LIBELLULA INDICA: alis flavo fuscoque variis apice albis: posticis macula baseos cyanea. Fab. Ent. Syst.

T. 2. 376. 9.

Libellula variegata. Linn. Syft. Libellula histrio. Mant. Inf.

Another species of Libellulæ, peculiar to India, and found in China, greatly resembles this insect: it is probably a variety of it.

Fabricius divides these two families of Linnæan Libellulæ into three distinct genera; the first retains the Linnæan name, the second and third are called Aeshna and Agrion. Their most effential characters are taken from the form and situation of the mouth, and therefore require a deep magnifier to determine them with accuracy. We have examined those parts in the greater number of the species Fabricius has described, and find his characters agree, except in one instance; but, which alone proves the impracticability of adopting the whole of his system: he describes Libellula Chinessis, and refers to the only figure that has been given of it, in one of the plates of Edwards's Natural History of Birds, 1745. Had he ever seen and examined this rare species, he must have referred it to his genus Agrion, each of the lips being bish, or two-cleft, as in Libellula virgo and puella,—the effential characteristic of the genus Agrion; for the mouths of the Libellulæ of Fabricius differ altogether in structure, and are not notched in the slightest degree, as Libellula clavata, ferruginea, 6 maculata, and the European species Libellula depress, will sufficiently illustrate.

LIBELLULA 6 MACULATA.

SPECIFIC CHARACTER.

Three black marks on the anterior ribs of the first pair of wings, marginal spot white. Posterior wings clouded with yellow.

LIBELLULA 6 MACULATA: alis maculis tribus coftalibus atris: ultima ftigmate niveo, posticis fasciis flavescentibus. Fab. Ent. Syst. T. 2. p. 381. 37.

These delicate insects appear to be male and female; they are almost a miniature resemblance of the two sexes of Libellula depressa, found in Europe; one having the abdomen yellow, and the other blue.

LIBELLULA CHINENSIS.

SPECIFIC CHARACTER.

Anterior wings brownish, without spots. Posterior pair green. Apex brown.

Libellula Chinensis: alis anticis testaceo obsoletis, posticis viridibus apice suscis. Linn. Syst. Nat. 2.

904. 15.

The only two specimens of this species we are acquainted with, were in the collection of the late Duches of Portland; one of those is now in the possession of Mr. Francillon.

LIBELLULA FERRUGINEA.

SPECIFIC CHARACTER.

Wings clear, white, yellow at the base. Body red.

LIBELLULA FERRUGINEA: alis albis basi flavis, corpore rubro. Fab. Ent. Syst. T. 2. p. 380. 33.

Very common in China.

LIBELLULA FULVIA.


SPECIFIC CHARACTER.

Wings yellowish, anterior margins testaceous, with a pellucid spot in the middle. Marginal spot near the end, brown.

LIBELLULA FULVIA: alis flavescentibus marginibus anticis testaceis, macula media subpellucida: stigmate ad apicem suscentibus marginibus anticis testaceis, macula media subpellucida: stigmate ad apicem suscentibus marginibus anticis testaceis, macula media subpellucida: stigmate ad apicem suscentibus marginibus anticis testaceis, macula media subpellucida: stigmate ad apicem suscentibus marginibus anticis testaceis, macula media subpellucida: stigmate ad apicem suscentibus marginibus anticis testaceis, macula media subpellucida: stigmate ad apicem suscentibus marginibus anticis testaceis, macula media subpellucida: stigmate ad apicem suscentibus marginibus anticis testaceis, macula media subpellucida: stigmate ad apicem suscentibus marginibus anticis testaceis, macula media subpellucida: stigmate ad apicem suscentibus marginibus anticis testaceis, macula media subpellucida: stigmate ad apicem subpellucida: stig


Libellula fulvia. Drury Inf. tab. 46. fig.

This infect has been figured, but not described before. We apprehend Fabricius considered it a variety of some other species, not having noticed it.




Libellula Chinensis. : Libellula ferruginea. ::Libellula fulvia.

London Published as the Act directs by Ellionessen No. 2124.







London Published as the Act directs by EDonovan Feb ? 1.1798.

ARANEA MACULATA.

SPOTTED SPIDER.

GENERIC CHARACTER.

Eyes eight. Mouth armed with two hooks or crotchets. Palpi two, confift of feveral joints, headed by the genitalia of the males in that fex. Anus contains teats for fpinning.

SPECIFIC CHARACTER.

Thorax covered with a fattin-like pile, of a filvery colour. Abdomen cylindrical. Legs long and black.

* * * * * * * Oculis. . . : : . .

Aranea Maculata: thorace holosericeo argenteo, abdomine cylindrico, pedibus longissimis atris.

Fab. Ent. Syst. t. 2. p. 425. sp. 66.


This remarkable creature is peculiar to fome parts of the Chinese empire. It is not the largest of the genus, known; yet it is of sufficient magnitude to excite terror and disgust. To an European, who has seen only the indigenous spiders of his own country, a species sive or six inches in length, and nearly the same in breadth, must appear a frightful creature: Aranea Maculata sometimes exceeds that size; but it has not the forbidding aspect of most insects of the same genus. The legs are unusually long, and the body slender. In its general appearance, it resembles some kinds of the *Phalangiae* that are known in England by the vulgar name *Harvest-men*, being generally seen about that time of the year.

It has been observed, that nature oftentimes adorns the most deformed and loathsome of her creatures in the richest display of colours; and this is especially noticed in many forts of snakes, toads, lizards, &c. Spiders seem also of this description: to a form the most hideous we frequently find united a brilliance of colours, and elegance of marking, that is scarcely excelled by any of the butterfly tribe,—the most beautiful of all lepidopterous infects. Our present subject is a striking proof of the latter part of this observation. The three sigures in our plate of Aranea Maculata exhibit a front and a profile view of the insect, together with the front of the head at the third sigure. The head is furnished with two very strong black mandibles, each terminated in an extremely acute point. The fore part of the thorax, which is wholly of a fine silky appearance, and the colour of silver, bending over the mandibles in the form of an arch, or circular head-piece, give it the resemblance of a black head with a crown of silver on the brow. This appearance is heightened in no small degree by three rugged prominences, one in the centre, and another on each side, on the upper part; and by the minute black eyes, which, like those of most spiders, sparkle with the lustre

of small gems. These eyes are eight in number, four are placed immediately in the front of the filvercoloured circular front piece, and on each side are two more at equal distances, but rather in oblique positions.

The body is really beautiful, the chief colour is deep brown, ftrongly tinged with bright purple; a broad ftripe of orange colour paffes down the abdomen from the thorax to the apex: the whole is elegantly marked with a variety of cream-coloured lines and spots intersecting each other. Very little hair is found on any part of this spider except on the thorax, which being rubbed off, discovers a hard testaceous black substance beneath.

Fabricius is the only author who has described this insect: it has never been figured in any work whatever, as appears by the latest works of Fabricius, and our own observation. It is not always the most decisive method to determine the species from a concise description; but this is sometimes unavoidable, and in the present instance the description given by Fabricius accords in every respect with our specimen. The only insect with which it could possibly be consounded is Aranea Pilipes, which also has never been figured; it differs however from Aranea Maculata in the very hairy clothing of the legs, and it has also two silver stripes down the back: a striking specific distinction to separate it from our insect. It is also a native of the East Indies, but not of China, that we are informed.





Cancer mammillaris.

CANCER MAMMILLARIS.

GENERIC CHARACTER.

Feet eight, claws two. Eyes two, diffant from each other, and placed on a kind of footftalk; moveable.

Palpi two large, furnished with claws. Tail articulated, and without sting.

SPECIFIC CHARACTER.

Thorax nearly oval, fpinous. Three fpines on each fide, beak fhort, with three teeth.

Cancer Mammillaris: thorace ovato aculeato utrinque trispinoso, rostro brevi tridentato.

Fab. Ent. Syst. T. 2. p. 465. 91.

In the Entomologia Systematica of Fabricius we find a species of Cancri described, that seems to agree with our specimen. The specific character assigned by that author is however less satisfactory than the general description he has added. The two red spots are very characteristic.

Fabricius met with this creature in the cabinet of Spengler, and notes its habitat China. It is worthy of remark, that this is the only species of the Cancer genus, Fabricius mentions as a native of that country.

CANCER MANTIS.

SPECIFIC CHARACTER,

AND

SYNONYMS.

Hand claws without tarfi, compreffed, hooked, ferrated, with teeth. Body angulated. Tail ferrated, and armed with fpines.

Cancer Mantis: macrourus articularis, manibus adactylis compressis falcatis serrato-dentatis.

Linn. Syst. Nat. 633. 54.—edit. 10.


Squilla Mantis: manibus adactylis compressis falcatis, ferrato dentatis, corpore subangulato, cauda ferrato spinosa. Fab. Ent. Syst. T. 3. p. 511.


The Linnæan Cancri are numerous, and include many species not less singular in appearance than the extraordinary creature before us. Indeed, some species are so extremely different from the rest, both in structure and manners of life, that we cannot hesitate in concluding the Linnæan character of the genus desective and indefinite. This may be observed in several of the species Linnæus himself described, and throughout a more extensive number of those discovered since the time of that author. It is evident Linnæus could never reconcile the subdivisions of the two principal families; the Brachyuri and Macrouri; or crabs with short and long tails; and later naturalists have ventured, with propriety, to alter this part of his arrangement. The Pagarus, Hippa, Scyllaris, Aslacus, Squilla, and Gammarus, are so many new genera, formed of the Linnæan tribes of Cancri in the last System of Fabricius. The genus, to which Fabricius retains the name Cancer, is distinguished from the rest, by having four short filiform antennæ; with the extreme articulation bisid, as we observe in the preceding species. This author follows Degeer in describing the Cancer Mantis, and alters the generic name to Squilla: he observes, that this genus is entirely distinct from the Cancri, and we are of the same opinion, but adhere to the Linnæan genera, because the species, Mantis, was described by Linnæus, and it may be improper therefore to remove it, to a new genus, in any partial survey of his system. The Cancer Mantis is not peculiar to China.



Cancer Mantis.

Landon Published withe Act devote by E.Doneson Josep, at







Scolopendra Morsitans.

London Published is the Act directs by E. Donovan March 1.1798.

SCOLOPENDRA MORSITANS.

VENEMOUS, OR BITING CENTIPEDE.

GENERIC CHARACTER.

As many feet on each fide, as fegments of the body. Body depressed. Antennæ setaceous.

Palpi 2, jointed.

SPECIFIC CHARACTER

AND

SYNONYMS.

Feet 20 on each fide, eyes eight.

Scolopendra Morsitans: pedibus utrinque 20, oculis octo. Linn. Syft. Nat. 2. 1068. 5.—Fab. Ent. Syft. 2. 390. 6.

Scolopendra Mordante,—Scolopendra pedibus utrinque 21, posterioribus spinosis. De Geer. Inf. 7.

563. 1. tab. 43. fig. 1.

Seba. Muf. 1. tab. 81. fig. 3, 4.

2. tab. 25. fig. 3, 4.

Catefb. Car. 3. 2. tab. 2.

Sulz. Hift. Inf. tab. 30. fig. 14.

Bradl. Nat. tab. 25. fig. 4.

Travellers agree that the temperate parts of Asia would be a terrestrial paradise, were it not for the multitude of troublesome insects and reptiles with which they are insected. In a well cultivated country, like China, many of these creatures can scarcely find shelter; but such as harbour in the walls or furniture of human dwellings are as abundant in that, as any other country that lies within or near the tropics. Amongst the latter, none produce more terrible effects than the Centipede, whose poison is as venemous as that of the scorpion, which also is a native of China.

Sir G. Staunton mentions a remarkable circumstance that occurred during the late embassy to China. The ambassador and his suite were accommodated in a temple near the suburbs of Tong-choo-soo. "In some of the apartments the priests had suffered scorpions and scolopendras to harbour through neglect. These noisome creatures were known only by description to some of the gentlemen in the embassy, who had not visited the southern parts of Europe: the sight of such, for the first time, excited a degree of horror in their minds; and it seemed to them to be a sufficient objection to the country, that it produced these animals." Sir George however adds, that no accident happened in that instance.—The species of Scolopendra he alludes to, is probably Morsitans, which is common in many parts of the world, but is particularly found of a frightful size, and in vast abundance, in the two Indies.

Many authors have described this creature. In the voluminous works of Seba we find several specimens of it from different countries, differing materially in fize, and some trifling particulars, but all possessing the distinguishing characters of Scolopendra Morsitans. The largest of these exceed our figure in magnitude, being near fourteen inches in length: this he calls Millepeda major ex nova Hispania. His figure of Millepeda Africanae is about the fize of our Chinese specimen. He has also a third and fourth figure, Millepeda Orientalis and Millepeda Celonica, mas: the latter is the same length as our figure, but the body is very narrow. Millepeda Orientalis is also the same length, but the body is very broad. Some of these infects are not four inches in length.

Authors agree that they vary exceedingly in fize and colour. De Geer describes them to be sometimes deep reddish brown; at others the colour of yellow ochre. The figure in Catesby's Natural History of Carolina is light brown; we have specimens of a livid yellow, and have seen others strongly tinged with red.

The last pair of legs are considerably larger than the others, and are armed with small black spines. The legs terminate in very sharp hooks or nails of a shining black colour. All the other legs are also furnished with a smaller nail of the same shape and colour.

M. Gronovius fays that all its feet are very venemous; but the most formidable of its weapons are the two sharp, hooked instruments that are placed under the mouth, and with which it destroys its prey.

Lewenhoeck having examined these instruments with a microscope, he sound a small opening at the extremity of each, and a channel from them into the body of the creature. Through this channel he supposes the Scolopendra emits the poisonous fluid into the wound it makes with the hooked instrument. That author further remarks, that he has seen a liquor on that part of living scolopendras. A figure of these instruments on the under side of the head, is represented in one of the diffections in our plate.

The fame author, wishing to ascertain the influence of the poison of Scolopendra morsitans, placed a large fly within its reach. The Scolopendra at first took it between a pair of its middle feet, then passed it from one pair of feet to the next, till the fly was brought under the sharp pointed instrument or crotchets at the head, which it plunged into the fly, and it died instantly. Notwithstanding this experiment, De Geer, Catesby, and other authors, affert, that its bite seldom proves fatal to larger animals; but all agree that its poison is as dangerous as that of the scorpion.

This Scolopendra has eight eyes: they are very fmall; four are placed on each fide of the head near the antennæ. In the diffections a figure is given to exhibit the manner in which the four eyes are placed on one fide.

⁵ These creatures differ from most insects in their manner of growth, insomuch that it it impossible to ascertain when they are of their full fize. The segments of the body increase as they advance in age, which circumstance renders it difficult oftentimes to determine the species, without a minute examination of its other characters.

INDEX

ACCORDING TO THE

SYSTEMA NATURÆ

OF

LINNÆUS.

COLEOPTERA.

SCARABÆUS.
Pl. 1, 2, 3.
naficornis
feniculus
Midas
Bucephalus
Moloffus
Sacer

cinctus Cetonia Chinenfis Melolontha viridis

Leci

CURCULIO.
Pl. 4, 5.
longipes
barbiroftris
Chinenfis
pulverulentus
Squamofus

magnified perlatus verrucofus

Pl. 6. rubus farinofus reticulator

Buprestis. Pl. 7. vittata ocellata

TENEBRIO. Pl. 8. femoratus

Meloe.

HEMIPTERA.

Mantis. Pl. 9. oculata flabellicornis

GRYLLUS.
Pl. 10, 11, 12, 13.
nafutus
vittatus
gryllotalpa
acuminatus
perfpicillatus
flavicornis
morbillofus

FULGORA. Pl. 14. Candelaria

CICADA.
Pl. 15, 16, 17.
atrata
ambigua
fanguinea
fplendidula
lanata
fontalis
abdominalis
limbata var.

NEPA. Pl. 18, 19. grandis rubra ruftica

CIMEX.
Pl. 20, 21.
Stockerus
difpar
cruciger
phafianus
aurantius
bifidus
flanbufchii

LEPIDOPTERA.

Papilio.
Equites Troes.
Pl. 22, 23, 24, 25, 26.
Crino
Paris
Coon
Agenor
Laomedon
Peranthus

Equites Achivi.
Pl. 27, 28, 29, 30, 31, 32.
Telamon
Agamemnon
Rhetenor
Demoleus
Epius

Heliconii. Pl. 33. vefta

Danai Candidi.
Hyparete
Pafithoë
Pl. 34, 35
Glaucippe
Sefia
Pyranthe
Philea

Danai Festivi. Jacintha Antiochus Jairus

Nymphales Gemmati.
Pl. 36, 37, 38.
Bernardus
Oenone
Almana
Lubentina
Orythia

Nymphales Phalerati.
Polyxena
Leucothoë
Erymanthis

Gambrifius

Piebeii Rurales. Pl. 39. Atymnus

Mæcenas

Sphinx. Pl. 40, 41. Nechus Thallo ruficollis bifafciata Hylas

Polymena

Phalæna.
Pl. 42, 43, 44.
Atlas
Militaris
lectrix
Bubo
pagaria
zonaria

NEUROPTERA.

LIBELLULA.
Pl. 45, 46.
clavata
indica
6-maculata
ferruginea
fulvia
chinenfis

APTERA.

ARANEA. Pl. 47. maculata

CANCER. Pl. 48, 49. mammillaris mantis

SCOLOPENDRA.
Pl. 50.
morfitans.

INDEX

ACCORDING TO THE

ENTOMOLOGIA SYSTEMATICA

OF

FABRICIUS.

CLASS 1. ELEUTERATA.
Genus 2. Scarabæus
naficornis
feniculus
Midas
Bucephalus
Moloffus
facer
Leei
cinctus
Genus 55. Sagra
femorata
Genus 63. Mylabris
Cichorei
Genus 76. Cetonia Chinenfis
Chinenfis
Genus 77. Melolontha
viridis
Genus 78. Buprestis
vittata
ocellata
Genus 82. Prionus
reticulator
rubus
Genus 85. Stenocorus
farinofus
Genus 102. Curculio
longipes
barbiroftris
Chinenfis
pulverulentus
fquamofus
mag.
perlatus
verrucofus
Cress 2 Hranes
Genus 120. Mantis
flabellicornis

oculata

nafutus

vittatus

Genus 122. Truxalis

```
Genus 123. Acheta
           Gryllotalpa
 Genus 124. Locusta
           perspicillata
           acuminata
 Genus 125. Gryllus
           morbillofus
           flavicornis
 CLASS 5. ODONATA.
 Genus 164. Libellula
           indica
           chinenfis
           ferruginea
           fulvia
        6-maculata
 Genus 165. Aeshna
          elavata
CLASS 6. MITOSATA.
Genus 167. Scolopendra
morfitans
 CLASS 7. UNOGATA.
Genus 171. Aranea
          maculata
 CLASS 8. AGONATA.
  Genus 175. Cancer
          mammillaris
 Genus 184. Squilla
          mantis
 CLASS 9. GLOSSATA.
  Genus 186. Papilio
   Equites Trojani*
     Crino
```

Paris

Coon

Laomedon

Doronthus
Peranthus
Agenor
** Equites Achivi.
Telamon
Agamemnon
Demoleus
Epius
Epius
*** Festivi.
Jacintha
Antiochus
Jairus
vanus
**** Nymphales.
Bernardus
Gambrifius
Rhetenor
Almana
Oenone
Orythia
Lubentina
Leucothoë
Polyxena
Erymanthis
Erymanuns
**** Heliconii.
Vefta
Hyparete
Pafithoe
Tantiloc
***** Danai.
Pyranthe
Glaucippe
Sefia
Philea
Timea
Genus 187. Hesperia
Mæcenas
Atymnus
Genus 188. Sphinx
Nechus
Genus 180. Selia

Pyranthe
Glaucippe
Sefia
Philea
Genus 187. Hesperia
Mæcenas
Atymnus
Genus 188. Sphinx
Nechus
Genus 189. Sefia
Hylas

Genus 190. Zygænæ
bifasciata
ruficollis
Thallo
Genus 191. Bombyx
Atlas
militaris
lectrix
Genus 194. Noctua
Bubo
Genus 196. Phalæna-
pagaria
zonaria
CLASS 10. RYNGOTA.
Genus 201. Fulgora
candelaria
Genus 203. Tettigonia
atrata
ambigua
fanguinolenta
fplendidula
Genus 204. Cicada
limbata, var.
lanata
Genus 205. Cercopis
abdominalis
frontalis
Genus 208. Nepa
grandis
ruftica
rubra
Genus 212. Cimex
Genus 212. Cimex
Stockerus
difpar
aurantius
Genus 214. Lygæus
cruciger
phafianus

Slanbufchii

bifidus

Genus 217. Reduvius

