

A short view of electricity / [Benjamin Wilson].

Contributors

Wilson, Benjamin, 1721-1788.

Publication/Creation

London : For C. Nourse, 1780.

Persistent URL

<https://wellcomecollection.org/works/taanmwen>

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

A
SHORT VIEW
OF
ELECTRICITY

WILSON

1780

56,156/C

72887

A
SHORT VIEW
OF
ELECTRICITY,

By B. WILSON, F. R. S.

And Member of the IMPERIAL ACADEMY of Sciences
at PETERSBURGH, &c. &c.

L O N D O N :

Printed for C. Nourse, opposite *Catherine-street* in the *Strand*.

M. DCC. LXXX.

To Mr. MOULTOU, jun. of Geneva, F. R. S.

AS the following little Tract was really undertaken to bring you acquainted with the subject of Electricity; at least, so far as comes within the compass of my knowledge, I have endeavoured to attain that end in the plainest and simplest manner; without troubling you with the opinions of others, and perplexing your mind with an endless variety of experiments.

The assistance you have since been pleased to favour me with, in consequence of the plan I pursued, by applying your mathematical abilities to these researches, is a considerable proof that my friendly endeavours have afforded you some satisfaction. On which account there appears to be a propriety in addressing this *Short View of Electricity* to you: and the rather, if I might hope that it would encourage you hereafter to treat this part of philosophy in the same manner as Sir Isaac Newton has done the great subjects in his *Principia*.

I am, SIR, with the greatest respect,

Your most obliged humble servant,

B. WILSON.

21st. Mountjoy Row, Covent Garden, W. R. 1781.

The following letter was really undertaken
to let you acquainted with the subject of Elec-
tricity; at least, to let as comes within the compass of
my knowledge. I have endeavoured to strain that end
in the plain and simple manner, without troubling
you with the opinions of others; and perceiving your
taste with an equal variety of experiments.
The of France, which have been related to France
meant, in consequence of the plan I pursued, by ap-
proaching you with such objections to these matters.
I have been informed that they friendly conversations have
aroused you some questions. On which account there
appears to be a necessity in abstracting this letter, as
of Electricity, and the other, if I might have
that it seems necessary to be added to treat the part
of philosophy in the same manner as Sir Isaac Newton
has done the great subjects in his System.

I am, Sir, with the greatest respect,

Your most obliged humble servant,

E. WILSON.

A

S H O R T V I E W

O F

E L E C T R I C I T Y.

General observations deduced from experiments.

1. **E**LECTRICAL effects seem to depend principally upon certain *properties* and *circumstances* of that universal elastic fluid, or *æther*, described by *Sir Isaac Newton*.
2. The earth, and all bodies upon it, as well as the air, (so far as hath yet been experienced) have naturally a certain quantity of this fluid appropriated to them: which, being extremely subtile and elastic, is liable to be disturbed from a variety of causes; and when disturbed, will be continually endeavouring to recover its natural state.
3. Bodies are said to be electrified, in the technical sense, when the natural quantity of this fluid is, by any cause, either augmented or diminished. In the former case they are said to be electrified *plus*: in the latter, *minus*.
4. Two very light bodies of the same material, size, and shape, properly suspended, and in their natural state, (so far as respects the fluid contained in them) do not change their place, but continue at rest.
5. Two similar bodies equally electrified *plus*, that is, having *received* more of the fluid than naturally belongs to them, recede from each other.
6. Two similar bodies equally electrified *minus*, that is, having *lost* part of the fluid which naturally belonged to them, do also recede from each other.
7. Two similar bodies, equally electrified, but in a contrary state, that is, the one having *more* and the other *less* of the fluid than

than what belongs to them naturally, move towards each other.

8. Two similar bodies, the one electrified and the other not, do also move towards each other ; but not so sensibly.
9. All bodies *resist* the passage of this fluid more or less.
10. Glass, amber, silk, &c. resist more than air.
11. A greater quantity of air resists more than a less quantity.
12. Air, in general, resists more than metal, wood, stone, &c.
13. Surfaces, that are smooth and even, resist more than points or acute angles.
14. In space, void of gross matter, vapour, or air, this fluid moves most freely.
15. All those bodies, such as metals, wood, stones, &c. through which this fluid is found to pass freely, have been usually called *conductors*.
16. And all those bodies, such as glass, amber, silk, &c. through which it does not so freely pass, *non-conductors*.

In the following experiments the wood made use of, as a conductor, is of a cylindrical form, and supported by glass that is dry and free from dust, fibres of down, &c.

Balls formed out of the pith of Elder, not larger than one fifteenth of an inch in diameter, and suspended by very fine flaxen threads, (about two inches long) from the cylinder of wood, are the similar light bodies alluded to in the 4th, and following observations.

The *excited* (solid) *glass* cylinder, employed in these experiments, need only be about six or seven inches long, and three eighths of an inch in diameter.

Woollen cloth will excite this glass sufficiently for most of the purposes designed in the following experiments.

EXPERIMENT

EXPERIMENT I.

IN this experiment, the cylinder of wood A is rounded at both ends, and supported by a common wine glass; but to prevent its rolling off from the glass, it should be made a little flat in one part.

FIG. I.



EXCITED GLASS held across, and over the middle of the wood, at a certain distance from it (which distance will depend upon the power of the glass) forces part of the *natural quantity* of the electric fluid, contained in the wood, into the balls; where it will accumulate to a certain degree.

As a proof of this, remove the glass quickly; then oppose it to the balls immediately, and they will recede from the glass. But this effect will continue only for a little time; because the accumulated fluid in the balls, which was part of the natural quantity forced from the wood by the power of the glass in the first instance, returns again into the wood (and thus recovers its natural state) soon after the glass is taken away.

This return of the fluid is somewhat slower in wood than in metal, because the resistance in the wood is somewhat greater.

EXPERIMENT II.


Repeat the experiment, and instead of removing the excited glass as before, *continue* it in its place. In this case the balls will continue to be repelled.

For if the power of the glass be sufficient, at a given distance, to force a certain quantity of the fluid out of the wood into the balls; it must likewise be sufficient to keep it there, that is, to

prevent its return, so long as the glass continues equally electrified, and at the same distance from the wood.

EXPERIMENT III.

In this experiment, the wood B is round at one end, and pointed at the other.



Excited glass, opposed very near to the point A, parts with a certain quantity of its fluid into the wood, threads, and balls. This additional quantity will continue therein for a considerable time, provided no cause interferes to disturb and weaken it. But if such a cause does interfere, and in a certain manner, the whole additional quantity may be discharged either *slowly*, or *suddenly*, according to the nature and form of the body employed to discharge it, the particular circumstances of communication which this discharging body hath with the earth during the experiment, and the quickness or slowness of its approach.

When the same glass is opposed to the balls, and at a proper distance, they will continue to be repelled by it: and whatever additional quantity of fluid may be in the wood, threads, and balls, the point of a *pin* will not discharge it; provided it be fixed on one end of a stick of sealing wax, and the head thereof be buried in the wax, and while the person who makes the experiment holds the wax by the contrary end: for no more of the fluid will escape into the pin than what will electrify it equally with the wood and balls, on account of the resistance exerted by the wax. The quantity of fluid, therefore, which escapes into the pin, can only be proportional to its quantity of matter. But if the person holding the wax touches the pin with his finger, the additional fluid in the wood, &c. will, at that instant nearly, be discharged.

From hence it is manifest, that if a person, standing upon the earth, could bring a very fine point towards a body electrified, (even

(even in the highest degree) *instantaneously*, the greatest discharge would ensue at that moment. Now as this case is, in strictness, impracticable, and as it requires *time* to bring the point sufficiently near, the accumulated quantity must escape gradually in every instant of its approach. But the quicker such approach is made, the more sudden and violent will be the discharge.

Motion, therefore, is a circumstance of great importance in experiments of this kind.

EXPERIMENT IV.

In this experiment the two bodies, A and B, are each of the same form as in experiment the third; and have their pointed ends placed near each other, or in contact.

FIG. 3.



Excited glass, held over A, (as in the first experiment) forces part of the fluid contained (naturally) in A into the balls hanging to it, and part into B: and if the two bodies A and B are immediately removed from each other, B will be found to have *more* fluid in it than belonged to it naturally, which it will retain for a time; and A will have *less*.


When A is separated from B, the balls at A will move towards excited glass; and the balls at B will be repelled by it. Those at A, therefore, as well as the wood, threads and balls, (having lost part of their natural quantity) will be in a *minus* state; and because the balls at B, as likewise the wood and threads, have received an additional quantity, they will be in a *plus* state. This is further manifest by bringing the two ends of A and B together again as they stood at first; for, immediately on doing this, the balls at the two extremities will no longer repel each other: because the over-charge in B will diffuse itself into A, and render the quantities in each equal.

EXPERIMENT

EXPERIMENT V.

In this experiment the two bodies, A and B, are each of the same form as before, but have their round ends opposed to each other,

FIG. 4.




Excited glass, opposed very near to the end A, will part with a quantity of its accumulated fluid into A; and that quantity, by endeavouring to escape from A at the round end F, will force part of the natural quantity, contained in the similar body B, out at the point C into the air: or, in other words, A will be electrified plus and B minus.

Separate those bodies immediately, and the balls belonging to each will continue to be repelled: but the balls at A will recede from glass, and the balls at B will move towards it. Bring the bodies together and into contact, and the repulsion between the balls will cease. Consequently each body is now restored to the state of its natural quantity: and, in the former part of the experiment, as much of the fluid as B lost was equal to the quantity which A gained.

EXPERIMENT VI.

In this experiment the wood is pointed at each end.

FIG. 5.



Excited glass, held over the middle of the body A, forces some part of the natural quantity within the wood into the balls; and some part out, at the two ends, into the air.

During this experiment the balls at A are repelled by glass; and are therefore in a plus state. But after the excited glass is removed,

moved, in a very little time they change to a minus state: because the two ends of the wood, from the nature of their form, had suffered part of the natural quantity to escape into the air, while the glass was held over the wood. But now the glass being removed, the over-charge in the balls will of course return, and equally diffuse itself into the wood again: and because this over-charge, even with the addition of some part of the natural quantity belonging to the balls, is found insufficient to balance the loss sustained, the wood, threads and balls must be in a minus state.

EXPERIMENT VII.

FIG. 6.



Excited glass, held near the end B, parts with a quantity of fluid into it: but that quantity will be less than the quantity parted with into a body differently formed, as in experiment III.


In this case, glass repels the balls, but the balls do not repel one another to so great a distance as in the third experiment: because here the fluid escapes more readily at the two ends, from the nature of their form: whereas, in the other case, (experiment III.) one of the ends, being formed differently, resists more, and consequently suffers less of the fluid to escape by it.

EXPERIMENT VIII.

In this experiment three bodies, A, B, C, are placed in a line near to, or in contact with, each other. The middle body A is pointed at each end; but the extreme bodies B and C are rounded at one end, and pointed at the other; and have their pointed ends opposed to the pointed ends of A.

FIG. 7.

FIG. 7.




Excited glass, held over A, forces out part of the natural quantity of fluid contained in A into B and C.

After the experiment separate A from B and C immediately, and it will be found that glass will cause the balls at A to move towards it, and the balls at B and C to recede from it: when, therefore, those three bodies are brought again into their first situation, (properly) the balls hanging to B, A and C, will cease to repel each other; because the over-charges (or accumulated quantities) in B and C will then be diffused into A; and consequently each of the three bodies will contain no more than its natural quantity.

EXPERIMENT IX.

In this experiment the four pieces of wood, A, B, C, D, are each of them round at both ends, and in contact with each other.

FIG. 8.



Excited glass, held over A, forces out part of the natural quantity contained in A into B; and the quantity received in B will force out part of the natural quantity contained in C into D.

The moment before the excited glass is removed from A, separate B and D from A and C; after which it will be found that A and C will be in a minus state, and B and D in a plus state: for, from the same reason that the excited glass forced the fluid out of A into B, the increase of fluid in B must, (as the excited glass did in the case of A) force the fluid out of C into D.

That

That the respective bodies will be in those states, described above, is manifest from bringing the excited glass towards the pith balls hanging to A B C and D: for those at A and C will move towards the glass, and those at B and D will recede from it.

EXPERIMENT X.

In this experiment the body D is round at each end, as in the first experiment.

FIG. 9.

D




Excited glass held over D, and *very near it*, parts with a small quantity of its accumulated fluid into D, where it will continue for a time.

Because the resistance at the two ends, arising from their form, hinders, in a great measure, any escape of the fluid into the air, while the glass is held over D; and that D has actually received more fluid, in this case, than what belongs to it naturally, appears from the balls being repelled by glass.

EXPERIMENT XI.

FIG. 10.

A



Excited glass held over A, during the time that any proper conductor (C) communicates with it and the earth, forces *part* of the fluid *out* of A into the conductor towards the earth; and renders

C

A

A *more minus* than if no such communication with the earth was introduced.


On removing the excited glass and conductor (C) at the same time, nay, even if the conductor be removed rather the last from A, the balls will be repelled, and move towards the glass when it is exposed to them. If the experiment be made carefully, the distance between the balls may be observed to *increase* for a few moments after such removal; that is, till the remaining part of the fluid in A becomes equally diffused therein.

Now, because the balls are repelled to a greater distance, in this case, than when no communication with the earth interferes, it is a manifest proof, that air resists the passage of this fluid more than grosser matter.

If further proof be required for explaining the several effects produced in the preceding experiments, we need only pursue a little farther the last experiment, in which A was left in a *minus* state; the communication with the earth being withdrawn, and the balls continuing to repel each other: For

EXPERIMENT XII.

FIG. II.



Excited glass, held over A, at 12 or 15 inches from it, and then moved gently nearer and nearer *towards* A, will cause the balls to approach nearer and nearer; and when the glass hath arrived to a certain distance from A, the balls will shew *no signs* of being electrified. In this state they will *continue* so long as the glass continues in that station. But if the glass be moved a little nearer towards A, the balls will begin to recede again from each other; and the nearer the glass is brought towards A, the *more* the balls will be repelled. In this last case, the balls are electrified plus, because

because they recede from glass: and they will continue in the same state so long as the glass *continues* in the *same place*.

Now as these effects are produced by no other change of circumstances, than merely moving the glass continually nearer towards A, the moving it back again, in the same manner, ought to produce the same effects, and leave the balls in the same minus state they were found in at first.

Make the experiment, and it will succeed accordingly.

The minus electricity, observed from the repulsion of the balls, in the first instance, is rendered less, by the approach of the glass, which forces part of the natural quantity in the wood into the threads and balls. The nearer, therefore, the glass approaches to A, the more fluid is forced from A into them; consequently, the repulsion of the balls must be less and less. When, therefore, the balls have received so much fluid from A, as equals the quantity they have lost, their power of repelling ceases, and may, for the present, be considered as in their natural state; the continuance of which state depends upon the glass that causes it, because the glass must continue to be held exactly in the same place: for the moment the glass is moved a little nearer towards A, (which is the second instance) a little more of the fluid is forced from A into the balls; and whatever that quantity may be, it necessarily electrifies them plus. The moving the same glass, therefore, nearer and nearer to A, must, for the same reason, force more and more fluid from A into the balls; and thus the balls, from having their quantity increased, must be repelled more and more.

However trifling the preceding series of experiments may appear to some electricians, the true philosopher may probably view them in a different light; because the reasonings upon their several effects are, it is apprehended, equally applicable to other experiments made upon a larger scale.

As an instance, take one of the experiments that were made at the *Panttheon* with the *great cylinder*, when a lesser one was joined to it, and when *motion* was introduced. By comparing that with the third of these experiments, their coincidence, in general, will

be manifest. For, in the *Pantheon* experiments, a *spark* was produced, at the end of the second cylinder, by a *point* at *nine* inches distance, when it communicated with the earth and was *moved suddenly*. At the same instant (nearly) all the fluid, or, however, the greatest part, was discharged: the spark obtained, therefore, was at a greater distance than when the same point was brought gradually towards the cylinder.

The case is very different with a different termination, though the other circumstances continue the same. For when a *ball* of metal was properly put in the place of the point, and an equal motion given to it, the greatest distance, at which a spark was produced, was only *six* inches. This difference of distance depended, therefore, upon the metal ball resisting the passage of the fluid more than the point; because less of the fluid escaped by the ball than there did by the point, as appeared from the different *residuums* in the cylinders, that were observed immediately after the explosions happened; there being a greater residuum when the ball had caused a spark, than when the point had done the same. So necessary is it, in all these cases, to attend to the circumstance of *motion* as well as that of *resistance*.

Seeing that motion is of so much consequence in experiments of this kind, a method may be contrived which will cause the greatest discharge possible, from a body containing any given quantity of fluid, by interposing a proper substance (for example, glass or wax, sufficiently broad and thick) between the end of the body proposed to be charged, and the point or round end which is to cause the discharge; and that *without moving* either of those ends in the least.

For when those extremities are brought to a convenient or certain distance from each other, and the body is afterwards charged, the removing such interposed substance *suddenly* from between those extremities (by some proper and simple contrivance) will cause the discharge almost instantaneously; and consequently, the relative distances at which a point or round end will be struck, with a given charge, may be more exactly ascertained. On which account it
appears

appears to be an experiment of some moment, and deserves to be tried; as it is yet in dispute whether a point or surface is struck at the greatest distance.*

If the contrivance, alluded to above, for removing the interposed body suddenly, should consist of a *spring* of any kind, no part thereof must interfere with the experiment, so as to reduce the charge in the least, while such spring is producing its effect. And because the motion, which is to be given to the interposing substance, is proposed to be *considerable*, there must be a cushion, or some other substitute, to counteract the force by which the interposed body is impelled.

* If the two extremities are sharply pointed, the distance between them, at which the discharge happens, will, it is apprehended, be considerably greater than when the same ends have a spherical termination, of a given diameter.

THE preceding experiments having been made with *wood*, which is a conducting substance, it is now time to introduce others made with *glass*, which is usually called a *non-conductor*.

But before these experiments are entered upon, it may be proper to observe, that there are some delicate circumstances belonging to them, which, if not attended to, will mislead the observer more than he may be aware of.

1. Whenever an experiment is to be made with glass, *as a conductor*, particular care must be taken that every part of it is in a natural state, and consequently undisturbed by friction, or any other cause, except that of the air, which is unavoidable.
2. The pith balls employed ought to be suspended by fine flaxen threads, three or four inches long; because, the bringing excited glass near them, to examine their plus or minus state, might otherwise interfere with the glass conductor and disturb the experiment.
3. A greater or lesser degree of power in the excited glass, will occasion some differences in the effects produced by it.


4. The

4. The glass conductor ought to have no bubbles or sand-holes in it, particularly near its surface.
5. Breathing upon any part of the conductor ought also to be avoided.

EXPERIMENT XIII.

B, A, C, represents a solid cylinder of glass near three quarters of an inch in diameter, and about eight inches long; one end of which is rounded, and the other pointed.

FIG. 12.




The same *excited glass*, as was employed in the preceding experiments, held at two inches or thereabout from the rounded end C, forces part of the natural quantity of fluid at that end towards the other; and after a time, (the excited glass being removed) the quantity so forced returns.

Oppose the excited glass to the balls at C, and they will move towards it; then oppose it to those at B, and they will be repelled. And, because the fluid meets with more resistance in glass than in metal or wood, the motion of it will be slower; and therefore must take up some time before the accumulated quantity forced towards B can possibly return to the minus part at the end C.

EXPERIMENT

EXPERIMENT XIV.

FIG. 13.




Repeat the last experiment, but with this difference; let there be a communication of metal (F) with the earth from the point B; and, after a few seconds, remove the communication and excited glass at the same time. When this is done the whole glass will be electrified minus.

For while the excited glass at the end C is forcing the natural quantity towards B, the communication (F) with the earth is conveying part of it off: and therefore when that communication and the excited glass are removed at the same time, the whole of the glass A must be more or less in a minus state: that is, according to the quantity of fluid which escaped at the point B, through F, towards the earth during the continuance of the excited glass at C.

EXPERIMENT XV.

In this experiment the glass is pointed at each end.

FIG. 14.




Excited glass, held over the middle of BC, forces that part of the natural quantity which is nearest to A towards the two ends,
and

and some of it into the air at the points B and C : in a little time afterwards the whole of B C, and all the balls, will be electrified minus.

That the excited glafs forces part of the natural quantity from the middle towards the two ends, appears from the balls at B and C being repelled by glafs : and that the middle has lost part of its natural quantity, is manifest from the balls at D moving towards the glafs. In the other case, where more time is necessary, the whole of B C is electrified minus : because all the balls move towards excited glafs ; and therefore some part of the natural quantity must have escaped at the points : otherwise the whole glafs and balls could not, in such circumstances, be in a minus state.

EXPERIMENT XVI.

FIG. 15.




Excited glafs, held over A, during the time that a metal conductor (F) communicates with any part near the middle of B C and the earth, forces part of the natural quantity out of the glafs, near A, into the conductor (F) towards the earth ; so that when the metal and excited glafs are removed at the same time, the whole of B C, in a few seconds, will be electrified minus.

For part of the natural quantity escapes by the metal (F) ; and therefore, when the excited glafs and metal are removed, the balls at the two ends will repel each other a little, and move towards excited glafs. But when the whole of the remaining fluid in B C has had time to diffuse itself equally throughout, the balls will
repel

repel each other to a greater distance, and move towards the glass with more force.

EXPERIMENT XVII.

FIG. 16.



While the glass BC, and the balls hanging to it, continue in their minus state, if the excited glass be again held over the middle for a few seconds, the two ends will be electrified plus, and the middle part will remain minus. But the plus effect at the two ends will continue for a little time only (after the excited glass is removed) and then the whole of BC will return to its minus state again, as was the case in the last experiment.

The excited glass, in this case, forces more of the remaining fluid from the middle towards the two ends, therefore the balls hanging to those ends will be repelled by glass; and the balls in the middle will be moved towards it. The greater quantities of fluid at those ends must consequently, in a little time, (that is, after the excited glass is removed,) diffuse themselves into the minus part again, and put the whole of BC into the same state in which it was at first.

EXPERIMENT XVIII.

FIG. 17.



Excited glass, held at the end B and in contact with it, parts with some of its accumulated fluid into B; and the quantity admitted therein forces out some part of the natural quantity at the remote end C into the air: and when the quantity received at the end B has had time to diffuse itself through the whole of BC, then all the balls will be repelled by glass.

D

That

That, during the experiment, some part of the natural quantity of the fluid in the glass cylinder B C escapes at the end C, appears from the effect which the excited glass hath upon the balls at C : because their motion is not from, but towards, the glass. But when the quantity received at B has had time to diffuse itself through B C, then the balls at both ends will be electrified plus.

EXPERIMENT XIX.

FIG. 18.



The same excited glass, held at about one inch distance from the end B of a solid cylinder of glass (B D) six feet long and about half an inch in diameter, will force part of the natural quantity of the fluid at the end B towards the remote end D. But in doing this, the natural quantity, belonging to the glass, will undergo several alterations ; as appears from the effects which excited glass hath upon a number of balls suspended at equal distances between B and D. In a little time those alterations will be reversed : that is, the parts of the glass that were electrified plus will become minus, and those that were minus will be plus.

EXPERIMENT XX.

When the same excited glass is held at the end B, as before, but in contact with it (for a very little time) the additional fluid received at B will, in going towards D, cause several alterations in the density of the natural fluid contained in B D. But those alterations of density will be the converse to those in the last experiment. And after a little time they will also be reversed, like those observed in the former case.

Whoever may be disposed to repeat these two experiments, must not be surprized if, in a few trials, he should not make them succeed in the manner here described : because it is extremely difficult,
and

and almost impossible, to find the same number of balls plus, and the same number minus, in each experiment. For a difference in the degree of power applied, or a difference in point of time allowed for the changes, as well as to observe the state of each ball, make considerable differences in the effects. These and other delicate circumstances have been the reason for describing the two last experiments in general terms. As to the cause of the alternate states of the fluid in the whole length of B D, and the changes taking place afterwards, it may be comprehended from what has been observed in the preceding experiments.

Because we have seen (by experiment 5, 9, 13, 17, and 18,) that whenever the fluid, contained within a body, becomes suddenly denser in any one part, the fluid in the neighbouring parts, to a certain distance, will be rarer: and *vice versa*, whenever it is made suddenly rarer in any part, the fluid lying next to it becomes denser. Whence it is manifest, that those alternate variations of rarity and density, must, from the nature of an elastic fluid, continue to oscillate many times backward and forward before the fluid can be at rest. But after those oscillations are weakened to a certain degree, they must become imperceptible to the observer.

From these experiments it is also manifest, that the air surrounding any body electrified (when properly circumstanced) must likewise be electrified, but in a contrary state, to a certain distance; and beyond that, must undergo the like alternate changes to imperceptible distances.

EXPERIMENT XXI.

If instead of the glass cylinder, (employed as a conductor) there be put in its place a glass of a different shape; for example, a *plate* eight or ten inches square, both sides thereof, to a certain distance from the center, will be electrified plus, by opposing the excited glass towards the middle of one of them only, and at a little distance from it.

E X P E R I M E N T XXII.

On the contrary, the middle of the glafs, on both fides, will be electrified minus; provided the distance between the excited glafs and the plate of glafs is considerably increased.

N. B. Instead of opposing the excited glafs (in the two last experiments) to the plate of glafs, it will be found more convenient to oppose the *round end* of a metal conductor, when it is properly electrified.

The difference of form, then, between a plate and cylinder of glafs, can make little or no difference in those general effects.

HAVING now produced a sufficient number of facts to shew the nature of this fluid, and the constant uniformity of its action, in glafs as well as wood (they differing only in the circumstance of resistance) we may safely venture to deduce from them some observations that will be of use in explaining the Leyden experiment.

Observation 1. Whenever a given quantity of the electric fluid (which for the future will be expressed by the word *power*) is brought within a certain distance of the surface of a body, whether it be wood, metal, or glafs, it will force out part of the natural quantity of fluid belonging to that body, (and that the more readily if it communicates with the earth) and will also prevent its return, so long as the power continues the same, and at the same distance from the surface.

Obs. 2. The nearer, therefore, this power is brought to the surface of the same body, the greater will be the effect produced by it.

Obs. 3. More time is required for the electric fluid to pass through a given length, or thickness of glafs, than through metal or wood of the same length or thickness.

Obs. 4. This difference of time, in the passage of the fluid through glafs, metal, or wood, can arise from no other cause than the different resistances made by those substances.

Hence

Hence it appears, that if by any contrivance we can bring the power to, and deposit it at, the surface itself of a plate of glass, (by diffusing it over a given quantity of that surface,) the fluid so lodged must, from having so advantageous a situation, act upon the natural quantity contained in the glass with its greatest force. And consequently not only drive out the greatest quantity of the fluid naturally lying at the opposite surface; but also continue to prevent the return of an equal quantity of fluid to supply the loss it had thus sustained; and that so long as the power lodged continues in its place.

But it is to be observed, that no expulsion of the fluid can in any degree take place at the opposite surface, unless the resistance at that surface be sufficiently weakened (or, at least, in some degree) by a proper communication with the earth.

A plate of glass thus circumstanced, is, for the purpose of experiment, the *Leyden phial*.

For the contrivance, alluded to above, means no more than that this plate should have an equal and partial coating of metal laid properly upon each surface. The one serving to conduct the power immediately to the glass, (so far as the coating extends) and collect it there: and the other to carry off as readily such part of the natural quantity belonging to the glass itself (at that surface) towards the earth, or towards any other equivalent substitute, as the power collected is able to drive away.

When a plate is thus charged, and as highly as it will admit, the removing of the coating cannot remove the charge from the glass, because it is deposited at, or near, the surface itself. Neither can the opposite surface, which is in a contrary state at the same time, receive a similar quantity of fluid from any other power (to supply the loss it has sustained) unless at the time there be a removal of *that* in which the charge consists: by reason of the advantageous situation of the diffused charge which originally drove off and prevents the return of such natural quantity to the opposite surface; as is evident from the 1st and 2d of the last observations.

And

And because the resistance, arising from the air, as well as that occasioned by the substance of the glass itself, prevents a free communication between the two surfaces, any contrivance which will sufficiently lessen the resistance between them, must necessarily suffer the charge to escape towards the other side of the glass, and restore the equilibrium. Consequently, if no alteration takes place to lessen the resistance, the glass must continue in the same charged state.

When a wire, or other convenient substance, is employed for this purpose, it is called the *circuit*: because one end communicates with the coating on one surface; and the other with the opposite coating: for when this is properly made, an explosion ensues, and the plate recovers its natural state; though not absolutely so, there being in many instances some little remainder.

A slow and imperceptible discharge may be produced by moist air, vapour of different kinds, and other conducting matter in the form of vapour, when properly applied, and in sufficient quantity, so as to extend itself to both surfaces.

Another contrivance to lessen the resistance, and thereby cause the discharge, is to reduce the thickness of the glass itself in any particular part, where the charge is to be given. Because, if it be made sufficiently thin, the power exerted at the charged surface will, in this case, force its way through the substance of the thinnest part, and break the glass to come at the opposite surface, and supply the natural quantity of fluid it had lost.

Hence we may now understand, why increasing the thickness of the plate increases the resistance: and therefore, why a plate of glass, beyond a certain thickness, cannot be charged at all by the same degree of power.

That all bodies resist the passage of the fluid more or less has been fully proved: but from what cause this resistance arises hath not yet appeared.

ACCORDING to Sir Isaac Newton, and the experiments hitherto made, a certain elastic medium (similar, except in density,

fity,

sity, to the fluid we are treating of) is supposed to be spread over the surface of all bodies, extending to a certain exceedingly small distance from them; which medium is more or less dense, according to the nature of the body it surrounds.

On those accounts it is continually liable to be disturbed from a variety of causes.

The rings of colours produced by pressing the object-glasses of two long telescopes together, taken notice of by the same great observer of nature, and the idea he entertained that such a medium, as described above, does actually surround the surface of all bodies, were sufficient inducements for trying, whether, by the assistance of those coloured rings, some kind of alteration in the medium might not be observed to take place, when the glasses themselves were charged with the electric fluid in a certain manner.

P R E P A R A T I O N.

A well polished plate of glass, (about 12 inches square, properly guarded with a border of sealing wax all round the extremities on both sides, near three inches broad) was well charged like the Leyden phial. After which the two coatings were carefully removed.

Another glass, plain on one side and a little convex on the other, (four inches and three quarters in diameter, and sufficiently thin, being also guarded a little more than half an inch on both sides round the extremity with sealing wax) was likewise charged, the plain side plus and the *convex minus*. After which the two coatings were also carefully removed.

E X P E R I M E N T XXIII.

When the convex surface, thus charged *minus*, was gently laid upon the *minus* surface of the large plate of glass, the weight itself of the plano-convex, without any other pressure, was sufficient to cause a *distinct* appearance of the coloured rings: and the central spot exhibited a *darkish blue colour*. And notwithstanding this general appearance varied, upon any the least change of accidental

dental circumstances, yet the mean diameter of the spot seemed to be about the twentieth part of an inch.

EXPERIMENT XXIV.

After charging those glasses again as before, except that now the convex side was in a *plus* state, the experiment was carefully repeated, but in a contrary way; for the convex side charged *plus* lay upon the *plus* surface of the large plate of glass: and though the weight or pressure of the glass was the same, as in the first case, yet there was a sensible difference; in the general appearance at least: for the coloured rings were *less distinct*, and the central spot appeared of a *yellowish* or *greenish colour*.

Upon repeating this experiment it was observed, in one or two instances, that the spot appeared of a fine red colour: and in other trials, there was no appearance of any spot or coloured rings whatsoever.

Hence it appears, that the glasses were *nearer* to each other when in the minus state in the 21st experiment: and that the surfaces stood at a *greater* distance from each other when in the plus state in the 22d experiment; the pressure in both experiments being the same.

Query, Does it not follow from hence, that the medium on the surfaces of the two glasses had its natural limit extended, or contracted, according to the nature of the charge, agreeable to the representations in fig. 21 and 22? Unless those differences, in the appearance of the rings, should have arisen from any unevenness of the surfaces. For the two surfaces of the square plate of glass, though finely polished, were not ground parallel to each other: which, to have removed all doubt, they ought to have been.

It is material to observe here, that at the end of each of the preceding experiments with the square plane and plano-convex glasses, it was found that they continued to retain the greater part of their respective charges.

EXPERIMENT

EXPERIMENT XXV.

When the glasses were laid on each other, as before, but without any charge in either, the pressure arising from the weight of the plano-convex, was sufficient to produce the rings, but not so distinctly, as in the 21st experiment; yet more sensibly than in the 22d.

IN ORDER to convey a more clear idea of the manner in which this medium is spread over the surfaces; and how it is affected by different circumstances, it has been thought not improper to represent the different states of the medium, in the case of glass particularly, (when it is not, and when it is, disturbed) by the following figures.

E

FIG.

FIG. 19.

A B C D represents the edge, or section of a plate of glass, in its natural or undisturbed state.

ab, ab , the limit of the medium spread on each side of the surface A B.

dc, dc , the limit of the medium spread on each side of the surface C D.

G G G G, the natural state of the fluid which lays between the medium ab below A B, and the medium dc above C D.

This medium is supposed by Sir *Isaac Newton*, in his letter to Mr. *Boyle*, (and since that in his optics) to be rarest at the surface A B (or C D) from whence, on each side, it increases in density continually and regularly to a certain exceedingly small distance. Beyond which the density decreases as regularly to an equal distance; that is, to ab (or dc): so that the densest part of the medium lies exactly in the middle between A B and ab , above and below; and the rarest part exactly in the middle between ab above A B and ab below it: or, as was observed above, at the surface itself A B.

FIG. 20.

A B C D represents the same glass, and in the same state, as in fig. 19.

$t t$, the upper coating of metal.

$k k$, the under coating of metal.

m , the conductor to communicate the fluid to the upper coating $t t$.

n , a metallic communication between the under coating $k k$, and the earth, to conduct the fluid to the earth when driven out from the under surface by the power exerted at the upper surface.

Being thus circumstanced, the glass is properly prepared for making the *Leyden* experiment.

FIG. 21.

Represents the same glass when its medium is disturbed, or charged, with the electric fluid.

$p p$, the part at the surface A B where the charge being lodged or deposited, produces the effect of a plus electricity.

$m m$, the part at the surface C D, which, being deprived of part of its natural quantity of fluid by the power at $p p$, produces the effect of a minus electricity.

FIG. 22.

A B C D represents the same glass, as in fig. 21. in its charged state, but without the coatings.

Fig. 19.




Fig. 20.




Fig. 21.





Fig. 22.



THIS interesting experiment, though so common, hath not as yet, perhaps, been explained to the satisfaction of the philosopher. All the authors who have written upon it having (as it is apprehended) confined their attention chiefly to the meer result of experiments, rather than to the state of the fluid contained within the substance and at the surface of glafs.

A solution, deduced from Sir Isaac Newton's opinion, touching the natural state of the surface of bodies described in fig. 19, and the consequences to be drawn from certain alterations caused therein, will probably be acceptable to those at least, who embrace his doctrines.

FIG. 23.





FIG. 24.



When the surface of glafs A B (fig. 23.) has received a certain quantity of electric fluid, as represented in fig. 24. the medium, which on each side of A B is supposed to be limited by *a b* and *a b*, must be driven on each side farther from the surface A B than it was before, that is to *k k* and *a b*; and therefore must press the natural fluid which lies between the internal limits *a b* and *d c*, against the opposite medium belonging to the surface C D, and supposed to be limited by *d c* and *d c*. But the fluid naturally belonging to the glafs being thus pressed, must push the limit *d c*, and force it nearer to the surface C D, than it was in its undisturbed state. And then the fluid, which naturally belongs to C D, being also pressed, must necessarily, in part at least, be driven out of the glafs. For without such effect taking place, the limit *d c* cannot change its situation, or be brought nearer to the surface C D. But if *d c* be not brought nearer to C D, neither will *a b* be moved farther from the surface A B, which it before was proved to be.

But it is necessary to open a passage for this fluid to escape from one surface while the other is charging : because, if there be no such passage, the glass cannot be charged in the *Leyden* manner at all ; on account of the resistance arising from the natural fluid belonging to the glass, and the medium itself at the under surface.

The discharging of the *Leyden* phial then being (as observed before) only the restoring of the equilibrium in the fluid belonging to the glass, the effect, produced by the shock of the fluid through the conductor which is employed to restore the equilibrium, must be attributed to the resistance of the glass itself : consequently the fluid cannot, on account of that resistance, be accelerated in going through the conductor. Therefore, whatever difference is perceived in the violence of the shock through conductors of different lengths, that difference should be attributed to the resistance the electric fluid meets with in going from one end of such conductor to the other.

AFTER the preceding explanation of the state of the fluid in the *Leyden* experiment, it will be proper to shew the state of it within the substance of a common conductor when it is charged. We have therefore added the following Theorem respecting that state, and the particular part of the conductor which the charge principally occupies.

T H E O R E M.

If a solid cylinder of metal is charged with electric fluid, I say, the fluid contained therein will be denser at the surface than in any part nearer its axis.

If it be supposed otherwise, and the fluid to be equally dense throughout the whole cylinder, then the effect produced, by discharging it, would be proportional to the quantity of matter contained in the cylinder. But by experiment it is found, that a hollow cylinder, of a similar substance, length, and diameter, with the former, and which will contain only the thousandth part of its solid matter, will produce very nearly the same effect as such a solid

solid cylinder; and not, as might be expected, the thousandth part of it only. The particles of the fluid, therefore, contained in the solid one, must be in a state extremely rare, compared with those at the surface.

IT has been remarked, that bodies terminated with points, receive, or part with the electric fluid more readily than rounded ones. This being a fact of some consequence, we shall attempt to give the reason of it upon the preceding principles.

WHEN a body is charged with electric fluid, I say, that if the surface thereof is terminated by a sharp end, it will part with that fluid more easily than if it is terminated by a round surface.

FIG. 25.





FIG. 26.



Let an acute angle *bac* represent the termination of the body in question; *t* a particle of the fluid lying between the surface of the body and its medium; and let *tq* be the direction of the particle. When the particle *t* arrives at *q*, it must be refracted outwards, or reflected inwards, in the direction of *qr*, which makes the angle *rqs* equal to the angle *tqp*. But in the case of the sharp end, the reflection of the particle in the direction *qr* is more perpendicular to *rs* than *tq* is to *qp*; and by consequence must contribute to produce in the particle a fitter disposition to refraction in that state than when it moved in the first direction.

But

But in the case of the round surface, the reflection cannot produce any such advantage, because the particle is always reflected inwards without changing the direction for one more advantageous.

Therefore the sharp end must part with the fluid sooner, because in all the cases of reflections the particles are reflected according to the directions that are always more advantageous to a refraction; which is not the case with the round end.

Upon ACCELERATION.


THE experiments at the *Pantheon*, which were intended to shew the acceleration of the fluid, having been objected to by many who have not sufficiently attended to the known properties of the elastic fluid, it has been thought proper here to establish this very material point upon mathematical principles, with a view to put an end to all farther disputes on the subject.

But before this is done, it may be necessary to mention a material fact that was omitted in the account of those experiments, which is this,

The shock, received at the *middle* of the long wire, was *considerably less* than that which was received at either end,

T H E O R E M.

F I G. 27.



Let A B represent a cylinder of a given diameter, and suppose this cylinder charged with the electric fluid. I say, if all the particles of this fluid are moved at the same instant towards A, the effect produced, by the shock of this fluid, at A, will be nearly proportional to the square of A B.

For the total effect at A is equal to the sum of the effect of each particle contained in the cylinder A B. And the effect of each particle being proportional to its velocity, the total effect at A will

will be proportional to the sum of all the velocities. But since the fluid is supposed nearly perfectly elastic, all the particles will arrive at A nearly at the same instant. Then the velocity of each particle will be proportional to the distance from the place it sets out: and the total effect at A will be proportional to the sum of all those distances.

But all those distances are expressed by the following numbers, 1, 2, 3, 4, 5, &c. N (N , expressing the length AB) in an arithmetical progression (see Fig. 27.) Then the sum of all the distances will be expressed by the sum of the arithmetical progression 1, 2, 3, 4, 5, &c. N , and the effect at A will be proportional to this sum, that is to say, to N^2 . or AB^2 . Q. E. D.

C O R O L L A R Y.

We may demonstrate in a similar manner that the resistance which the fluid meets with in its passage through the cylinder AB , in going to A, is proportional to AB^2 .

C O R. 2.

Then, generally, the total effect produced by the discharge at A (the resistance being allowed for) is proportional to AB^2 .

Concerning LIMITED ACCELERATION.

IN the foregoing Theorem the body receiving the shock hath been considered as in communication with the Earth, and consequently in the situation proper to produce a complete discharge; and therefore a perfect acceleration.

It is now necessary that we consider the case in which the body, being *insulated*, does not receive the whole discharge from the cylinder, but only so much of the fluid as is necessary to restore the equilibrium between the body and the cylinder: this was fully proved by Experiment III. From which it is plain, that the quantity of electric fluid which the body upon the discharge receives, being limited, the distance also, whence the electrical particles composing that quantity sets out, must be limited,

ted; and therefore the acceleration of the particles, together with the effect produced, will likewise be limited.

In the following analysis it is proposed to determine that limit; and at the same time to shew the law observed in the effects produced by the discharge from different cylinders of equal diameters, but indeterminate lengths, upon an insulated body.



LET K represent a cylinder similar to that in the preceding Theorem, that is to say, of a diameter very small in comparison of its length; a wire for example.

Let A be the body insulated which is to receive the shock from the cylinder K. Now as we may not be able immediately to compare the quantities of the fluid which, after the shock, are contained in the body A and the cylinder K; we will suppose a certain cylinder *a* having its diameter equal to that of K, and its capacity, with respect to the electrical fluid, equal to the capacity of A, to be insulated, and in the place of A to receive the discharge from the cylinder K: so that, whatever shall hereafter be proved respecting the cylinder *a* may be with truth applicable to the body A.

First, Suppose *a* equal to *k*, and let *a* receive the shock.

The two cylinders being insulated must, after the shock, be in equilibrio; and being equal, *a* must have received half of the whole of the electrical fluid which, before the shock, was contained in K. The effect therefore produced, by the discharge from the cylinder K upon the insulated body A, will be equal to the effect which would be produced by the discharge of a cylinder

der of half the length of the cylinder K upon the same body A not insulated. Consequently (and by what hath been proved in the preceding Theorem) the effect produced by the cylinder K upon the insulated body A is but the fourth part of the effect produced by the same cylinder K upon the body A not insulated.

This is a particular case; and is only here introduced, as being the most simple, and consequently the most easy for computation. In what follows the proposition is as general as possible.

Secondly. Suppose now the cylinder K to be of any given length, and that length expressed by n times the length of a .

a

K

The two cylinders, as before, being insulated, a will, upon the discharge, receive no more of the fluid than what is necessary to restore the equilibrium; that is, $\frac{1}{n+1}$ th part of the whole. For the whole quantity contained in K is expressed by n ; and consequently the quantity of fluid received by a (or A) upon the shock, is properly expressed by $\frac{1}{n+1}$. And hence it follows, that the effect produced by the discharge from the cylinder K, upon the insulated body A, is equal to the effect produced upon the same body A, not insulated, by the discharge of a cylinder of the same diameter with the cylinder K; but whose length is to the length of the cylinder a as $\frac{1}{n+1}$ is to 1.

E X A M P L E.

The length of the cylinder a being expressed by 1, and the quantity of fluid, which a is capable of receiving, being consequently expressed by 1, also; if now the length of the cylinder K be successively expressed by the numbers 1, 2, 3, 4, 5, n , the quantity of electrical fluid received after the discharge by the

F

insulated

insulated body A, will be respectively expressed by the numbers $\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5} \dots \dots \frac{n}{n+1}$: this last being a general expression.

L I M I T.

If the length of the cylinder K be supposed to become infinite; that is, if the number n be indefinitely great; we shall then have $\frac{n}{n+1}$ equal to 1; and the quantity of electric fluid received by the insulated body A, will consequently be expressed by 1. And so, the effect produced by the discharge of the boundless cylinder K upon the insulated body A, will be equal to the effect produced by the discharge from a cylinder equal to a upon the body A not insulated.

G E N E R A L C O N C L U S I O N.

Whatever may be the length or the diameter of a cylinder, and with whatever quantity of electrical fluid it be charged; the shock received from such a cylinder, by a body insulated, is limited: and that limit depends entirely upon the body which receives the shock.

C O R O L L A R Y.

1. So that a man *insulated* receiving the shock from a wire three miles in length may hardly perceive any effect: whereas the same man *not insulated* may probably be killed by the shock from the same cylinder equally charged.

2. The less the insulated body is, which receives the discharge, the less is the acceleration.

IN the *Leyden* experiment, the surface of glass, which is in a minus state, cannot receive any undetermined quantity of the fluid, but must receive, from the other surface, only half of the excess.


The *Leyden* phial then, is in the same case with the man who is insulated. Consequently (in such circumstances) there can be no acceleration.

Therefore the effect produced by the *Leyden* charge must be proportional to the quantity of the fluid. And consequently, in
glass

glafs of the fame thickness, proportional to the quantity of furface.

And becaufe the fhock produced by the *Leyden* phial depends upon the two fufaces being electrified one plus and the other minus, it is manifefl, that wherever the like plus and minus electricity can be introduced, and to a certain degree, the effect will be the fame without any glafs at all.

FIG. 28.



Instead, therefore, of having those contrary powers so very near together as in glafs, let two fimilar, and fufficiently large, fufaces A and B (fig. 28.) of any metallic fubftance properly guarded at the edges with fealing wax, and infulated, be placed parallel and at a given diftance from each other; whether five, or twenty-five, feet is equally indifferent. And let a perfon, alfo infulated, ftanding between thofe plates, for example at K, communicate by wires from one plate to the other; except at G, where a given interval muft be left for interpofting the proper fubftance alluded to in pages 12 and 13, which is to interrupt fuch communication. If the plates, in thofe circumftances, are equally electrified, one plus and the other minus; and the interpofted body G be removed *fuddenly*, the

F 2

perfon,

person, at K, will receive a shock like that which is received from a charged glass of a certain size.

Hence it appears, that the violence of the shock will be proportional to the size of those metallic surfaces: and therefore, to equal the charge of any *Leyden* phial, or number of phials, those surfaces must be increased accordingly.

WE have already proved, that there is no acceleration in the case of the *Leyden* phial, and consequently, that the shock is proportional to the quantity of surface. Now as it hath been demonstrated (by the Theorem upon acceleration) that the effect produced with a cylinder of a given diameter (very small compared with its length, a wire for example) is proportional to the the square of its length; it follows

That if we express the effect produced from a wire of a certain length, by a certain number of square feet of *Leyden* phial, we shall be able to determine the number of square feet that will produce an equal effect with a wire of any length.

Let n represent the length of a wire which produces the effect of the *Leyden* phial, containing in square feet the number a . I say, that a wire of any length $q \times n$, will produce an effect equal to the effect produced by $q^2 \times a$, square feet of the *Leyden* phial.

D E M O N S T R A T I O N .

The effect produced by the wire n is to the effect produced by the wire $q \times n$, as $n^2 : q^2 n^2$. But the effect produced by n is equal to the effect produced by the number a in square feet of *Leyden* phial, and $n^2 : q^2 n^2 :: a : q^2 \times a$. Therefore the effect produced by the wire $q \times n$ is equal to the effect produced by the number of $q^2 \times a$, of square feet of *Leyden* phial. Q. E. D.

E X A M P L E .

If we suppose the effect produced (in the *Pantheon*) by a wire of one mile in length, is equal to the effect produced by two square

square feet (only) of *Leyden* phial, how many square feet are there necessary to produce the effect of a wire of 18 miles in length?

Then $n = 1$. $a = 2$. $q = 18$.

Consequently $q^2 \times a = 18^2 \times 2$.
 $= 324 \times 2 = 648$.

It is necessary, therefore, to employ 648 square feet of *Leyden* phial to produce the effect of a wire 18 miles long, in which the fluid is accelerated; and not 30 square feet, as was mentioned in a paper lately read in the Royal Society.

SO FAR we have considered this elastic fluid, simply as such, without any regard to other matter that generally accompanies it, and produces a great variety of singular appearances, such as lightning, and other luminous phenomena.

Of whatever kind this matter may be, which experiment shews to be intimately connected with the elastic fluid, there is sufficient reason to conclude, that it consists of particles not near so subtile as those which compose the fluid itself.


But whether it be *phlogiston* or a combination of *phlogiston* with other matter; and whether the property in air of lighting up this matter into sparks, &c. is a separate and distinct principle, must be left to future investigation.



(part of) the ...
 necessary to ...
 from a ...
 ...
 ...

It is ...
 ...
 ...
 ...

20. AR we have ...
 without ...
 and ...
 ...
 ...
 ...
 ...
 ...
 ...
 ...
 ...



F4 blank

