

**The naturalists' miscellany, or, Coloured figures of natural objects; drawn and described immediately from nature.**

**Contributors**

Shaw, George, 1751-1813.  
Nodder, Frederick Polydore  
Nodder, Richard P

**Publication/Creation**

[London] : [Printed for Nodder & co.], [1790-1813]

**Persistent URL**

<https://wellcomecollection.org/works/xgtyy48u>

**License and attribution**


This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.




Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>


































VIRO ORNATISSIMO

THOMÆ MARTYN,

SANCTÆ THEOLOGIÆ BACCALAUREO,

REGIÆ SOCIETATIS SOCIO,

BOTANICES

IN ACADEMIA CANTABRIGIENSI

PROFESSORI DIGNISSIMO,

*DUODECIMUM HUNC*

NATURÆ VIVARII

*FASCICULUM,*

D. D. D.

GEORGIUS SHAW,

FREDERICUS P. NODDER.



TO  
THE REVEREND  
THOMAS MARTYN,

B. D. F. R. S.

PROFESSOR OF BOTANY

IN THE

UNIVERSITY OF CAMBRIDGE,

*THIS TWELFTH VOLUME*

OF THE

NATURALIST'S MISCELLANY

IS

RESPECTFULLY INSCRIBED


BY

GEORGE SHAW,

FREDERICK P. NODDER.







# MERGUS CASTOR.



## CHARACTER GENERICUS.

*Rostrum* denticulatum, subulato-cylindricum, apice aduncum.

*Lin. Syst. Nat.* p. 207.

## CHARACTER SPECIFICUS, &c.

MERGUS cristatus cinereus, capite colloque supremo spadiceis, gula remigibus intermediis abdomineque albis.

*Lath. ind. orn.* p. 829.

MERGUS CASTOR. M. capite cristato cinereo, subtus ferrugineo, gula alba, rostro pedibusque nigricantibus.

*Lin. Syst. Nat.* p. 209.

MERGUS Gulo.

*Scopoll. ann.* 1. No. 88.

*Anas rubricapilla.*

*Brun. No.* 93.

---

Habita non raro est hæc avis Mergi Merganferis femina. Hodiernis tamen physicis visum est ab illo eam omnino sejungere, speciemque revera diversam ducere.



ducere. In Anglia mergo mergansere sæpius con-  
specta, præcipue in partibus septentrionalibus, lacus  
fluviosque frequentat, vesciturque pisciculis et reli-  
quis ejusmodi. Longa est, ut plurimum, uncias  
circiter viginti septem.










THE  
DUN MERGANSER.



GENERIC CHARACTER.

*Bill* narrow, sub-cylindric, toothed at the edges,  
hooked at the tip.

*Nostrils* small, ovate, towards the middle of the  
bill.

*Feet* tetradactyle, palmated, the exterior toe longer  
than the rest.

SPECIFIC CHARACTER, &c.

Ash-coloured crested MERGANSER, with the  
head and upper part of the neck ferruginous,  
the throat, abdomen and middle wing-feathers  
white.

The DUN-DIVER.

*Lath. syn.* 3. p. 420.

*Brit. Zool. edit. fol.* p. 147. pl. N.

Le Harle femelle.

*Pl. enl.* 953.

---

This species, which is now considered as perfectly  
distinct, has been often regarded as the female of  
the *Mergus Merganser*. It is observed to be more  
frequently

frequently seen in England than that species, and is chiefly found in the northern parts of the kingdom, frequenting lakes and rivers, and feeding on fish, &c. Its general length is about twenty-seven inches.



# EXOCOETUS EVOLANS.



## CHARACTER GENERICUS.

*Pinnæ pectorales* longitudine trunci.

## CHARACTER SPECIFICUS, &c.

EXOCOETUS pinnis ventralibus parvis prope pectus.

*Bloch. ichth.* 12. p. 9. t. 398.

ABDOMINALES.

EXOCOETUS EVOLANS. E. abdomine terti.

*Lin. Syst. Nat.* p. 521.

---

Piscium, quos perpauca continet hoc genus, longæ adeo et magnæ sunt pinnæ quæ *pectorales* vocantur, ut harum ope subito interdum ex aquis exiliant, et per aera quasi volitantes ferantur pedes duccenos vel tricenos; in fluctus se iterum committentes cum pinnæ exaruerint. Quæ in tabula depingitur species mare incolit mediterraneum et atlanticum, in quibus, ut verbis utar celeberrimi physici, “ vitam ducit miserrimam, coryphænis et aliis piscibus voracibus perpetuo vexata; quos si eluserit per aera erumpendo, vel a Laris vel Diomedea corripitur; aut rursus se aquis credere coacta, in ipsas fauces hostium detruditur, volantem intentis oculis parique celeritate observantium.”


Notandum

Notandum est non proprium et peculiarem esse  
volatum Exocoetis solis; sed donari illo et alios ali-  
quot pisces; species nempe nonnullas *Scorpææ*,  
*Triglæ*, &c.

Æquat plerumque magnitudine Exocoetus evolans  
Clupeam quæ Harengus dicitur.







*Exocoetidae, flying fish, from the collection of the Smithsonian Institution.*



THE  
FLYING-FISH.



GENERIC CHARACTER.

*Pectoral fins* as long as the body.

SPECIFIC CHARACTER, &c.

FLYING-FISH with small ventral fins, near the breast.

FLYING-FISH.

*Penn. Brit. Zool.* 3. p. 292. pl. 67.

Der Hochflieger.

*Bloch. ichth.* 12. p. 9. pl. 398.

ABDOMINALES.

---

The fish of this genus, which are very few in number, are remarkable for the extreme length and size of their pectoral fins, by which they are enabled to spring occasionally from the water, and to support a kind of flight or continued motion thro' the air, to the distance of two or three hundred feet, when, the fins becoming dry, they are again obliged to commit themselves to their own element. The species here represented is a native of the Mediterranean and Atlantic seas, where, to use the words of  
an



an eminent naturalist, " it leads a most miserable life. In its own element it is perpetually harrassed by the Dorados, and other fish of prey, and if it endeavors to avoid them by having recourse to the air, it either meets its fate from the Gulls or the Albatross, or is forced down again into the mouth of the inhabitants of the water, which keep pace below with its aerial excursion.

It should be observed that this power of flight or temporary skimming thro' the air to a considerable distance, is not confined to this genus only, but takes place in some species of *Scorpæna*, *Trigla*, &c.

The common Flying-Fish is generally about the size of a Herring.



# BULLA ACHATINA.



## CHARACTER GENERICUS.

*Animal* Limax.

*Testa* univalvis, convoluta, inermis.

*Apertum* subcoarctata, oblonga, longitudinalis,  
basi integerrima.

*Columella* obliqua, lævis.

*Lin. Syst. Nat.* p. 1186.

## CHARACTER SPECIFICUS, &c.

BULLA testa ovata, apertura obovata apiceque  
fanguineis, columella truncata.

*Lin. Syst. Nat.* p. 1186.

BULLA Gallica.


*Seb. mus.* 3. t. 71. f. 1. 2. 3. 7. 8:

---

Jampridem in opere hoc nostro Bullam achatinam  
ore purpureo depinximus. Rarior autem longe est  
eximia hæc varietas, et colorum insignem jactat ele-  
gantiam.



THE  
AGATE BULLA.



GENERIC CHARACTER.

*Animal* resembling a Limax or Slug.

*Shell* univalve, convoluted.

*Aperture* somewhat straitened, oblong, longitudinal, entire at the base.

*Column* oblique and smooth.

SPECIFIC CHARACTER, &c.

Ovate, pointed, wide-mouthed BULLA, with broad fasciæ and truncated column.

*Var.*

With blue variegations and crimson mouth.


La Corne de pourpre.

*Knorr. vergn.* 4. p. 44. pl. 24. f. 1.

---

A purple-mouthed variety of the *Bulla achatina* has already been represented in the present work ; but the beautiful kind here figured is much less frequent, and is highly remarkable for the elegance and delicacy of its colors.

447



R.N.

London, Published Sept. 1. 1850 by F.B. Nozder New-mars Street.







# PAPILIO AMPHIMEDON.

VAR.



## CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatae.

*Alæ* (sedentis) erectæ sursumque conniventes (volatu diurno.)

*Lin. Syst. Nat.* p. 744.

## CHARACTER SPECIFICUS, &c.

PAPILIO alis concoloribus fuscis : anterioribus albo radiatis ; posterioribus macula quinque-fida rubra lunulisque albis.

*Fabr. spec. ins.* 2. p. 8.

EQ. TR.

*Var ?*

Alis superioribus fuscis ochraceo radiatis, inferioribus flavis nigro maculatis.

*Seb. mus.* 4. p. 21. t. 16. f. 7. 8.

---

Inter maximos papiliones exoticos numeratur Papilio Amphimedon. In insula Amboina generatur, & magnitudine vera in tabula exprimitur.



# AMPHIMEDON.

VAR.



## GENERIC CHARACTER.

*Antennæ* commonly thickening towards the end into a clavated tip.

*Wings* (when at rest) meeting upwards. (Flight diurnal.)

## SPECIFIC CHARACTER, &c.

Brown Butterfly, with the upper wings radiated with white; the lower with a five-cleft red spot and white marginal crescents.

*Var?*

Upper wings radiated with ochre-color; lower yellow with black variegations.


---

This insect is one of the largest of the exotic Butterflies, and is a native of Amboina. The plate represents it in its natural size.





London: Published by T. D. Sadler, Newman Street.





## COLUMBA ROSEA.



### CHARACTER GENERICUS.

*Rostrum* rectum, versus apicem descendens.

*Nares* oblongæ, membrana molli tumida semitectæ.

*Lingua* integra.

*Lin. Syst. Nat.* p. 279.

### CHARACTER SPECIFICUS, &c.

COLUMBA roseo-sanguinea, cera aurantia, vertice orbitis apicibusque tectricum minorum albis, remigibus caudaque fuscis.

*Cim. Phys.* p. 105. t. 59.

---

Indiam? incolit formosissima hæc avis, magnitudine columbæ vulgari domesticæ fere æqualis. In opere splendidissimo Domini Miller cui titulus *Cimelia Physica* primum depicta fuit.

# COLUMBA ROSA

## CHARACTER GENERALIS


Forma recta, coram apertum descendens.  
Nervi oblique, interstitia non triplicia sed

simplex.  
Ligula magna.  
Lig. 8. A. Nat. p. 270.

Colora et sordida, etc.  
COLUMBA recta, segmenta, coram apertis, ver-  
the oblique, interstitia non triplicia sed  
simplex, ligula magna.  
Civ. 1. 1. p. 105. t. 10.

Forma recta, coram apertum descendens.  
Nervi oblique, interstitia non triplicia sed  
simplex, ligula magna.  
Civ. 1. 1. p. 105. t. 10.







London: Published, Dec. 1<sup>st</sup> 1800 by W. Crisp, Stationer, Newmarket Street.



THE  
CRIMSON PIGEON.



GENERIC CHARACTER.

*Bill* strait, descending towards the tip.

*Nostrils* oblong, half covered by a soft tumid membrane.

*Tongue* entire.

SPECIFIC CHARACTER, &c.

Sanguine rose-coloured PIGEON, with orange cere ; crown, orbits, and tips of the smaller coverts white ; wing-feathers and tail brown.

*Miller's Plates of Nat. Hist. pl. 59.*

---

This most beautiful bird is a native of India? In size it is nearly equal to a common domestic pigeon. It seems to have been first figured in Mr. Miller's splendid plates of Natural History.

# CRIMSON PIGEON

*Columba erythroneura*

This species is found in the  
mountains of the Himalayas  
and is distinguished by its  
bright red plumage.

The male has a red head and  
neck, and a red breast.  
The female has a red head and  
neck, and a red breast.

The male has a red head and  
neck, and a red breast.  
The female has a red head and  
neck, and a red breast.

The male has a red head and  
neck, and a red breast.  
The female has a red head and  
neck, and a red breast.



# TUBULARIA MAGNIFICA.



## CHARACTER GENERICUS.

*Stirps* tubulosa, simplex vel ramosa, basi affixa.  
*Animal* terminale, capite tentaculis cristato.

## CHARACTER SPECIFICUS, &c.

TUBULARIA tubo simplici albido, tentaculis numerosissimis albo rubroque variatis.

## TUBULARIA MAGNIFICA.

*Act. Lin. Soc.* 5. p. 228. t. 9.

---

Omnes quas norunt physici Tubularias superat longe hæc quæ in tabula depingitur tam magnitudine quam venustate. In insulis Americanis rupium cava incolit, et ut solent reliquæ congeneres, sese ad libitum vel extendere vel contrahere potest, et, instante periculo intra latibulum recedere. Genera præterea Tubulariæ et Amphitrites videtur quodammodo connectere; corpus nempe habens annulatum, Tubulariæ autem præcipuos characteras genericos. Ostenditur in tabula dimidiata magnitudine naturali. In quinto volumine actorum Societatis Linnaeanæ exstat icon archetypa quam delineavit ingeniosus Dominus Davies, copiarum militarium præfectus.











THE  
GREAT TUBULARIA.



GENERIC CHARACTER.

*Stem* tubular, simple or branched, fixed by the base.

*Animal* terminal; the head crested with tentacula.

SPECIFIC CHARACTER, &c.

TUBULARIA with a simple whitish tube, and very numerous tentacula variegated with red and white.

*Trans. Lin. Soc.* vol. 5. p. 228. pl. 9.

---

Of all the Tubulariæ yet known the present may be considered as by far the largest and most magnificent in its appearance. It is a native of the seas about the West Indian islands, inhabiting the cavities of rocks, and possessing, like the rest of the genus, the power of withdrawing itself on any appearance of danger into the hole in which it resides. It may be added that it seems in some degree to connect the genera of Tubularia and Amphitrite, having the annulated body of the one, with the peculiar generic characters of the other. It is represented



presented on the plate of half the natural size. It is to the ingenious pencil of General Davies that we are indebted for the original drawing of this animal, which has been engraved and published in the fifth volume of the transactions of the Linnæan Society,



# TRIGLA VOLITANS.



## CHARACTER GENERICUS.

*Digiti liberi ante pinnas pectorales.*

## CHARACTER SPECIFICUS, &c.

TRIGLA digitis vicens membrana connexis.

*Lin. Syst. Nat. p. 498.*

*Pisces Thoracici.*

TRIGLA pinnis pectoralibus longitudine trunci.

*Bloch ichth. 10. p. 93. t. 351.*

---

Nullorum congenerum magis singularis est conformatio seu splendidior color quam piscis qui in tabula depingitur. Longe majores sunt ei pinnae pectorales quam Triglæ Hirundini nec non Triglæ punctatæ, de qua antea in hoc opere disseruimus. Maris Mediterranei et Atlantici incola est Trigla volitans, longa plerumque duodecim vel quindecim uncias.

# TRIPLA VOLTAX

CHARACTER OF THE

TRIPLA VOLTAX

TRIPLA VOLTAX

CHARACTER OF THE

TRIPLA VOLTAX

TRIPLA VOLTAX

TRIPLA VOLTAX

TRIPLA VOLTAX

TRIPLA VOLTAX

TRIPLA VOLTAX

TRIPLA VOLTAX

TRIPLA VOLTAX

TRIPLA VOLTAX

TRIPLA VOLTAX

TRIPLA VOLTAX









THE  
FLYING GURNARD.



GENERIC CHARACTER.

*Finger-shaped processes before the pectoral fins.*

SPECIFIC CHARACTER, &c.

Reddish GURNARD, with numerous fingers connected by a web, and olive-coloured blue-spotted pectoral fins of the length of the body.

The long-finned GURNARD.

Der fliegende Seehahn.

*Bloch ichth. t. 351.*

---

Of all the Triglae this may be considered as the most beautiful in point of color, as well as the most remarkable in form, the size of the pectoral fins far surpassing those of the Trigla Hirundo, as well as of the Trigla punctata before figured in the present work. It is a native of the Mediterranean and Atlantic seas, and is commonly about twelve or fifteen inches in length.

# PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD

PLANTING CUPBOARD



# NAIS DIGITATA.



## CHARACTER GENERICUS.

*Corpus* repens, longum, lineare, pellucidum,  
depressum.

*Pedunculi* setis simplicibus.

*Tentacula* nulla.

*Oculi* duo aut nulli.

*Lin. Syst. Nat. Gmel. p. 3120.*

## CHARACTER SPECIFICUS, &c.

NAIS setis lateralibus solitariis, cauda laciniata.

*Lin. Syst. Nat. Gmel. p. 3121.*

*Müll. von Würm. p. 90. t. 5. f. 1-4. Hist.*

*Verm. I. 2. p. 22. n. 155. Zool. Dan. prodr. 2651.*


---

Naidi digitatæ, quæ e rarioribus est sui generis, insignis est ad speciei distinctionem caudæ conformatio. Eadem fere est magnitudine qua Nais proboscidea, et aquas incolit stagnantes exeunte æstate. Retrahi vel extendi ad libitum possunt caudæ tentacula; quæ cum ad summum expanduntur, parsque corporis anterior cauli plantæ alicujus aquaticæ adhæret, hydræ tenui et gracili simillima Nais digitata, primo visu spectantem falleret.









London Published Oct. 1800 by T. P. Nodder, Newman Street.



## DIGITATED NAIS.



### *GENERIC CHARACTER.*

*Body* repent, long, linear, pellucid, depressed.

*Feet* consisting of simple bristles.

*Tentacula* none.

*Eyes* two or none.

### *SPECIFIC CHARACTER, &c.*

NAIS with single lateral bristles and laciniated tail.

## DIGITATED NAIS.

Finger-tailed Water-Worm.

---

This animal may be considered as one of the rarer species of its genus, and is readily distinguished by the remarkable form of the tail. It is nearly equal in size to the *Nais proboscidea*, and may be found in stagnant waters towards the decline of summer. The processes or tentacula at the end of the tail may be either retracted or extended at pleasure, and when at their utmost extent, while the fore part of the body is attached to the stalk of some water plant or other object, the animal has so much the appearance of a slender *Hydra* or *Polype*, that it might easily, at first view, be mistaken for such.

# GENERAL CHARACTER

Body robust, long, linear, flattened, compressed.

Feet consisting of simple toes.

Tail very long.

Teeth two or none.

## SPECIFIC CHARACTER

Nails with single lateral lobe and terminal

tail.

## UNIDENTIFIED NAILS

Longer than broad.

This animal may be considered as one of the  
 representatives of its genus and is readily distinguish-  
 ed by the remarkable form of the tail. It is nearly  
 equal in size to the *Nail* pattern, and may be  
 found in stagnant waters towards the middle of  
 summer. The processes of the animal at the end of  
 the tail may be either retracted or extended at will.  
 One, and when in their normal extent, when the  
 fore part of the body is attached to the water, or  
 some water plant or other object. The animal has no  
 teeth, the appearance of a slender fly on which  
 that it might readily be mistaken for a fly.



# TRICHODA SOL.



## CHARACTER GENERICUS.

*Vermis* nudo oculo inconspicuus, crinitus.

## CHARACTER SPECIFICUS, &c.

TRICHODA globularis undique radiata.

*Müll. an. inf.* p. 164. t. 23. f. 13. 15.

*Lin. Syst. Nat.* Gmel. p. 3889.

*Joblot. microsc.* 1. p. 2. p. 64. t. 7. f. 15.

*Eichorn Zugabe* f. 1-7,

*Roes. hist. polyp.* p. 500. t. 83. f. 2. 2 2

---

Digna omnino notatu est hujus animalculi indoles et conformatio ; quæ diligentius inspecta non minus mirabitur philosophus quam hydram e divisis sui ipsius partibus plene renatam. Constare videtur totum corpus e massa quasi medullari, homogenea, globosa, radiis seu spinis subpellucidis undique densissime obsita ; oris officio fungente foramine parvulo in centro tuberculi seu papillæ. Torpido et socordi ingenio, dies totos continuos manet sine motu aliquo visibili, nisi quod subobscurè contrahatur, si extrinsecus acciderit aliquod irritamentum. Non tamen obstare videtur summa hæc inertia quo

c

minus




minus monoculos minores prædetur, præcipue parvulum illum cui nomen *Monoculus Pediculus*, qui nonnunquam in ventre repertus est. Verisimile autem est diu manere posse monoculum incolumem, & digestionis (ut loqui solent medici) vi illæsum; e corpore enim *Trichodæ* exemptus, licet multis horis fuerit inclusus, æque celeriter in aquis natare videtur atque solebat antequam in carcerem esset injectus. Si dividatur, vel potius divellatur *Trichoda* Sol acu seu instrumento aliquo idoneo, fragmenta diversa, quamvis primo valde inæqualia, unius vel duarum horarum spatio, formam pristinam sphæricam integri animalculi recuperabunt, radiis juxta veram proportionem sitis, et totidem perfecta animalcula e frustulis istius quod divisum fuerit, monstrantibus. Nihil ad hoc experimentum valere videtur aeris temperies, quod æque probabitur hyberno ac æstivo tempore. *Trichodæ* Solis magnitudo generalis est quasi apicis aciculæ vulgaris. Color subpellucidus alborem habet quasi margaritæ. Aquas plerumque puriores stagnantes incolit, et stipiti plantulæ alicujus aquaticæ affigitur.

Primo descriptum fuisse videtur hoc animalculum ab Eichorno in opere cui titulus "*Zugabe*," &c. A Müllero quoque descriptum et depictum est in præcellenti libro de animalculis quæ *Infusoria* dicuntur. In rarioribus habetur, mensibus Februario & Martio præcipue visum. Mense Februario anni millesimi septingentesimi nonagesimi octavi, speciei de qua nunc agitur plura specimina ipse egomet in frustula divulsi, quæ omnia fragmenta citius unius horæ spatio totidem perfecta fiebant animalcula. Ex

Trichodis




Trichodis integris una monoculum pediculum absorpserat, qui simul atque e carcere liberatus, aquas hilariter circumnatabat, illæsus, ut videbatur, et omnis incommodi expers.











THE  
SUN TRICHODA.



GENERIC CHARACTER.

*Animalcule* inconspicuous to the naked eye,  
beset with hair.

SPECIFIC CHARACTER, &c.

Globular TRICHODA radiated on all sides.

Radiated TRICHODA,

Solar TRICHODA.

---

The animalcule which forms the subject of the present plate is not more remarkable for the singularity of its form than of its nature, and when accurately examined, will be found to exhibit phenomena equally surprising with the reproduction of the polype.

The whole body appears to consist of an uniform medullary substance, beset on all sides with very numerous transparent diverging rays or prickles, while a small hole in the centre of a papillary protuberance forms the mouth.

It is an animalcule of an extremely inactive nature, and occasionally remains for days together in the same situation, and without any visible motion,  
except



except a very obscure contraction on being irritated by any external object.

Notwithstanding this habitual torpidity, it is, apparently, of a predacious nature, and sometimes seizes on and swallows the smaller Monoculi, and especially that very small species the *Monoculus Pediculus* of Linnæus, which is occasionally found in its body. It is probable however that this insect remains a long time uninjured by the digestive powers of the animalcule; since, on opening a *Trichoda* which has thus confined a *Monoculus* for many hours, the insect appears unhurt, and, when liberated, swims about with the same celerity as before its imprisonment.


If the *Trichoda Sol* be divided, or rather torn into several fragments by the point of a needle or other instrument, the several pieces, tho' at first of a very irregular form, will in the space of an hour or two assume the complete spherical figure of the animal before its division; the rays or spines appearing in their true proportion and situation; thus constituting so many distinct animalcules. This experiment seems to be very little influenced by the weather, since in the coldest part of winter it will be found to succeed nearly as well as in the warmer months.

The general size of this curious animalcule is that of a pin's head, and its usual residence is in the clearer kind of stagnant waters, where it commonly attaches itself to the stem or fibres of some of the smaller aquatic plants. Its color is a semitransparent pearly white. It appears to have been first described



scribed by Eichorn in his work entitled "*Zugabe*" &c. Mr. Müller has also described and figured it in his excellent work on the *Animalcula Infusoria*. It is numbered among the rarer animalcules, and is chiefly to be found in the months of February and March.

In February 1798 I divided several of these animalcules in the manner above described; the respective fragments of each of which, in less than an hour's space, assumed their complete form. One specimen, before dilaceration, had enclosed a *Monoculus*, which was no sooner liberated than it swam about, apparently uninjured by its captor.





## APIS LAPIDARIA.

\*\*\*\*\*

### CHARACTER GENERICUS.

Os maxillis atque proboscide inflexa vaginis duabus bivalvibus.

*Alæ planæ.*

*Aculeus feminis et neutris punctorius reconditus.*

### CHARACTER SPECIFICUS, &c.

APIS hirsuta atra, ano fulvo.

*Lin. Syst. Nat. p. 960.*

Bombylius maximus, totus niger, exceptis duobus extremis abdominis annulis rubris.

*Raj. ins. p. 246.*

*Reaum. ins. 6. t. 1. f. 1-4.*

*Frisch. ins. 9. t. 25. f. 2.*

*Scop. carn. 813.*

---


Apum Britannicarum e maximis est Apis lapidaria, cumque colores sortita sit sibi peculiares, a reliquis, primo visu, facillime possit discerni. Magnitudinis multa est diversitas; parvulaque nonnulla specimina (mascula fortasse) sæpe voluerunt auctores speciem ab hac revera separatam constituere. Nidus perelegans, constans e fibris muscorum majorum arcte inter se contextis, forma ovata, efficto  
in



in latere rotundo foramine, in aggeribus sylvarum  
& viarum diverbiis situs, larvas continet nec non  
cibum, mel nempe rude, in frustula fusca et in-  
æqualia huc illuc sparsum; hæc enim species nullas  
cellulas seu favos ex composito ordinatos construit.  
Larvæ in chrysalidas conversæ, singulæ in theca sua  
quasi ovata includuntur.

Notandum est, apes has admodum hirsutas ab  
Anglis vulgari nomine *Humble-Bees* designari; nec  
defuere qui in gravem errorem lapsi, a defectu acu-  
lei inditum fuisse iis hoc nomen opinati sunt.












THE  
ORANGE-TAILED BEE.



GENERIC CHARACTER.

*Mouth* furnished with jaws and inflected proboscis with two bivalve sheaths.

*Wings* flat.

*Sting* concealed and exertile in the females and neuters.

SPECIFIC CHARACTER, &c.

Hairy black BEE, with orange-coloured tail.

The red-tailed Humble-BEE.

Great ORANGE-TAILED Garden-BEE.

---

The *Apis lapidaria* is one of the largest of the British bees, as well as the most remarkable in point of color, being at all times readily distinguishable from the rest of its congeners. It varies however very considerably in magnitude, and some specimens, (perhaps males) are of so comparatively small a size as to have been often considered and described as a distinct species. The nest constructed by this insect is of a very elegant appearance, being of an oval form, and composed of sprigs of the larger mosses, very



very closely and neatly compacted together, a small round hole or entrance being left on one side. These nests are generally situated on dry shady banks in woods, lanes, &c. and contain the larvæ or young animals, together with their food, which consists of a coarse kind of honey of a brownish color, and disposed in somewhat irregular masses or heaps, this species not forming any angular cells or combs. The larvæ at the time of their change to a chrysalis, are each enveloped in an oval case.

It may not be improper to add, that the Bees of this division in the genus, are popularly known by the title of Humble-Bees, and some authors, inconversant in natural history, have most erroneously imagined them, in consequence of the above name; to be destitute of a sting.









# PHALÆNA JUNONIA.



## CHARACTER GENERICUS.

*Antennæ* setacæ, a basi ad apicem sensim attenuatæ.

*Alæ* (sedentis) sæpius deflexæ, (volatu nocturno.)

*Lin. Syst. Nat.* p. 808.

## CHARACTER SPECIFICUS, &c.

Ph. (pavonia) pectinicornis elinguis, alis rotundatis griseo nebulosis subfasciatis, ocello nictitante subfenestrato.

*Lin. Syst. Nat.* p. 810.

*Attaci.*

*Roes. ins.* 4. t. 15. 17.

*List. Goedart.* t. 28.

*Reaum. ins.* 1. t. 47. 49.

*Knorr. delic.* t. c. 2. f. 2.

*Seb. mus.* 4. t. 60. f. 9. 14.

---


Lepidopterorum quæ in Europa generantur longe maxima Phalæna Junonia coloribus eleganter dispositis insignitur. Nullibi in Anglia reperitur; non raro autem in variis Galliæ, Germaniæ, & Italiæ



Italiæ partibus. Suspiciatur sane Linnæus phalænam pavoniam minorem esse hujus de qua jam loquimur varietatem; sed quamvis fateamur simillima inter se esse hæc duo insecta, non possumus tamen nobis persuadere ad unam eandemque speciem debere referri. Ostenditur in tabulis vera phalænæ nec non larvæ pupæque magnitudo.







A.A.

*London, Published Nov. 1<sup>st</sup> 1800, by F. R. Nodder, Newman Street.*



THE  
GREAT PEACOCK MOTH.



GENERIC CHARACTER.

*Antennæ* setaceous, gradually lessening from the base to the tip.

*Wings* (when sitting) generally deflected. (Flight nocturnal.)

SPECIFIC CHARACTER, &c.

GREAT PEACOCK-MOTH, with whitish-grey rounded wings, with dark and pale-brown transverse variegations, and large subfenes-  
trated eye-shaped spot.

GREAT PEACOCK-MOTH.

---


Of all the European Lepidoptera this is by far the largest, as well as one of the most elegant in the disposition of its colors. It is not very uncommon in many parts of France, Germany, Italy, &c. but has never yet been discovered in our own country. Linnæus indeed imagines that the *Phalæna pavonia minor*, or Emperor Moth, is in reality a variety of the same animal; but tho' there exists the most striking



striking similarity in point of appearance, it is impossible to suppose a real identity of species. The great *Phalæna pavonia*, or *Junonia*, is represented on the annexed plates in its natural size, accompanied by its larva and pupa.



# COLUMBA CORONATA.



## CHARACTER GENERICUS.

*Rostrum* rectum, versus apicem descendens.

*Nares* oblongæ, membrana molli tumida semitectæ.

*Lingua* integra.

*Lin. Syst. Nat.* p. 279.

## CHARACTER SPECIFICUS, &c.

COLUMBA orbitis nigris, crista erecta, corpore cærulescente, humeris ferrugineis.

*Lin. Syst. Nat.* p. 282.

Phasianus cristatus Indicus.

*Briss. av.* 1. p. 279. t. 26.

---

Ob molem insolitam et quasi giganteam ad tribum gallinaceam potius quam columbinam referri debere hanc avem nonnullis physicis visum est. Characteres autem manifesti et dubio carentes fere primo visu verum genus evidenter denotant. In insulis Moluccis generatur *Columba coronata*, et in avibus elegantissimis proculdubio meretur numerari. Vocem emittit quasi palumbis, adeo tamen raucam et sonoram, ut subtimuisse dicantur nautæ quibus

præfuit Dominus Bouganvillius, exaudito cantu gemibundo, in viis horridis et sylvestribus, ubi primo insederant; suspicati strepitum seu ululatum esse hostilem indigenarum ferorum, bellum ex insidiis meditantem. In Europam viva sæpius nuper in-  
vecta est hæc avis, et in magno pretio habetur ab iis quibus curæ est rariora animalium genera colligere.







London Published Dec 1<sup>st</sup> 1802 by J. P. Nodder, Newman Street



THE  
CROWNED PIGEON.



GENERIC CHARACTER.

*Bill* strait, descending towards the tip.

*Nostrils* oblong, half covered by a soft tumid membrane.

*Tongue* entire.

SPECIFIC CHARACTER, &c.

Blue-grey PIGEON, with large compressed upright crest, purple-brown shoulders with a white spot, and red eyes.

Great CROWNED PIGEON.

*Edw. pl.* 338.

Le Faisan couronné des Indes.

*Buff. 2. p.* 354. 542.

*Pl. Enl.* 118.

---

The gigantic size of this species, which is not far short of that of a Turkey, has caused some naturalists to place it rather among the gallinaceous tribe than in the genus *Columba*. Its characters are



however so clearly and decisively marked as to declare at once its real and proper genus. It is undoubtedly one of the most elegant of birds, and is a native of some of the Molucca islands. Its voice resembles that of the wood-pigeon, but in so loud and hoarse a tone, that it is recorded of some of Mons. Bougainville's sailors, that they were greatly alarmed on hearing it for the first time, in the wild and unfrequented spots of some islands on which they landed; supposing it to have proceeded from the savage cries of hostile and concealed natives. This bird has frequently been brought alive into Europe, and is considered as one of the greatest ornaments of the Menagerie.



# SPARUS CHRYSURUS.



## *CHARACTER GENERICUS.*

Opercula squamata, inermia.

## *CHARACTER SPECIFICUS, &c.*

SPARUS ruber, linea laterali aurata.

*Bloch. ichth. 8. p. 25. t. 262.*

*Thoracici,*

---

Aquas Brasilienses incolit formosissimus piscis in  
tabula depictus, longitudine, ut plurimum pedali  
vel sesquipedali.

SPARUS CILICUS

SPARUS CILICUS

CILICUS

CILICUS

CILICUS

CILICUS

CILICUS

CILICUS

---

CILICUS


CILICUS

CILICUS

CILICUS







London, Published Dec 1798, by T. & A. Neill, Newman Street.



THE  
YELLOW-STRIPED SPARUS.



*GENERIC CHARACTER.*

Gill-covers scaly, unarmed.

*SPECIFIC CHARACTER.*

Red SPARUS, with yellow lateral stripe and tail.

Der Goldschwantz.

La Queue d'Or.

The Yellow-tailed SPARUS.

---

This highly beautiful fish inhabits the waters of Brasil. Its general length is from twelve to eighteen inches.

# THE YELLOW PERIL

by L. B. WATSON

Illustrated by J. H. M. WATSON

Published by the Yellow Peril Club, New York

Copyright, 1905, by L. B. Watson

Printed by the Yellow Peril Club, New York

10

The Yellow Peril Club

In General

The Yellow Peril Club

This book is intended for the use of the Yellow Peril Club, and is not to be used for any other purpose.



# PAPILIO PHORCAS.



## CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatae.

*Alæ* (sedentis) erectæ sursumque conniventes, (volatu diurno.)

*Lin. Syst. Nat.* p. 744.

## CHARACTER SPECIFICUS, &c.

PAPILIO niger, fascia supra viridi subtus pallidior, alis posterioribus caudatis.

PAPILIO Doreus. P. alis fascia supra viridi, subtus alba.

*Lin. Syst. Nat.* Gmel. p. 2239.

*Fab. sp. ins.* 2. p. 18.

## PAPILIO PHORCAS.

*Cram.* 1. t. 2. f. B. c.

---

In Sierra Leona præcipue conspicitur Papilio Phorcas, inter pulchriores merito numerandus. Magnitudinem naturalem ostendit tabula.

# PAPILIO PHORCAS

Characteristics

General description

Antennae of male and female, and of pupa

Wings

Color of male and female, and of pupa

(Notes on the)

The male and female

Characteristics

Papilio phorcas, Linn. Syst. Nat. p. 1000

Notes on the

Papilio phorcas, Linn. Syst. Nat. p. 1000

(Notes on the)

The male and female

Notes on the

Papilio phorcas

General description

Antennae of male and female, and of pupa

Wings

Color of male and female, and of pupa







459

9



## PHORCAS.



### *GENERIC CHARACTER.*

*Antennæ* gradually thickening towards the end,  
terminating in a clavated tip.

*Wings*, when at rest, upright. Flight diurnal.

### *SPECIFIC CHARACTER.*

Black Butterfly, with green band, paler beneath,  
the lower wings tailed.

---

This species, which may justly be numbered among  
the most elegant of its tribe, is chiefly found in Sierra  
Leona, and is represented on the plate in its natural  
size.

# PHORAS

=====

## Central Characters

Antennae gradually thickening towards the end,  
terminating in a clavate tip.  
Wings, when at rest, slightly tilted.

## General Characters

Black, thorax with green band, paler beneath.  
the lower wings tinted.


---

This species which is little known among  
the most elegant of the tribe is chiefly found in Japan  
I saw, and a very good one, the type in its natural  
size.



# MUREX TRIBULUS.

VAR.



## CHARACTER GENERICUS.

*Animal* Limax.

*Testa* univalvis, spiralis, exasperata suturis membranaceis.

*Apertura* desinens in canalem integrum, rectum seu subascendentem.

*Lin. Syst. Nat.* p. 1213.

## CHARACTER SPECIFICUS, &c.

MUREX TRIBULUS. M. testa ovata spinis setaceis trifariis, cauda elongata subulata recta similiter spinosa.

*Lin. Syst. Nat.* 1214.

VAR. spinis longissimis.

---

Quas genus *Murex* amplectitur numerosissimæ sunt species. His, ut plurimum, insignis est scaritæ superficies, nunc in longas spinas excurrentes, nunc in processus quosdam crispatos seu quasi foliatis. Testam de qua jam agitur rarissimam alit mare Indicum et Atlanticum. Variant specimina quoad colorem, nec non rostri et spinarum longitudinem.

# MUREX TRIBULUS.

WART.

\*\*\*\*\*

## CHARACTER GENERICUS.

Animal 3-line.

Testa nervis, spiralis, exap. ovata subtriangul.

hirsutis.

Apertura desinens in cordato integro, rotunda.

sen. submarginata.

Line. 2 1/2. Vel. 4. 1818.

## CHARACTER SPECIFICUS, &c.

MUREX TRIBULUS. M. testa ovata spinis

setosis tribulis, cordis elongata subulata

totum striatum spinis.

Line. 2 1/2. Vel. 1818.

Var. epula longissima.

Quamquam Murex tribulus, animalis tribulis  
tribulis, testis, spiralis, exap. ovata subtriangul.  
hirsutis, Apertura desinens in cordato integro, rotunda.  
sen. submarginata. Line. 2 1/2. Vel. 4. 1818.  
Murex tribulus, animalis tribulis, testis, spiralis,  
exap. ovata subtriangul. hirsutis, Apertura desinens  
in cordato integro, rotunda. sen. submarginata.  
Line. 2 1/2. Vel. 4. 1818.








P.N.

*London Published Dec 21<sup>st</sup> 1869, by F.P. Nodder, Newman Street.*



THE  
THORNY MUREX.

VAR.



GENERIC CHARACTER.

*Animal* allied to a Slug.

*Shell* univalve, spiral, roughened with membranaceous sutures.

*Aperture* ending in a strait or subascending channel.

SPECIFIC CHARACTER, &c.

Whitish MUREX, with a triple row of setaceous spines, and long beak with similar spines.

VAR. With extremely long spines.

The Thorny Woodcock.

---

The genus Murex, of which the species are extremely numerous, is distinguished, in general, by a peculiar roughness of surface, which either runs out into long spines, or into processes more or less crisped or foliated. The very rare shell exhibited on the present plate is principally found in the Indian and American seas, and varies as to color and length of the beak and spines.

# THORNY MUREX

VAR.

## GENERAL CHARACTER

Animal allied to a *Physa*.  
Shell univalve, apical region with numerous  
bristly points.  
Aperture ending in a short or subascending  
channel.

## Specific Character, &c.

Whitish MUREX, with a triple row of setaceous  
spines, and long beak with similar spines.

VAR. With extremely long spines.

The Thorny Woodcock.

---

The genus *Murex*, to which the species are ex-  
tremely numerous, is distinguished, in general, by  
a peculiar roughness of surface, which either runs  
out in long spines, or in numerous more or less  
crooked or twisted. The spines are shell exhibited  
on the present plate especially found in the in-  
dian and American species, and varies as to color and  
length of the beak and spines.



# BUCCO MAYNANENSIS.



## CHARACTER GENERICUS.

*Rostrum* cultratum, lateraliter compressum,  
apice utrinque emarginato, incurvato; rictu  
infra oculos protenso.

*Nares* pennis recumbentibus obtectæ.

*Pedes* scansorii.

*Lin. Syst. Nat.* p. 168.

## CHARACTER SPECIFICUS, &c.

BUCCO viridis, capite gulaque rubris, cæruleo  
marginatis, jugulo et pectore flavo, abdo-  
minis macula rubra.

*Lath. ind. orn.* p. 203.

## BUCCO MAYNANENSIS.

*Briss.* 4. p. 102.

BUCCO elegans.

*Lin. Syst. Nat. Gmel.* p. 406.

---

In Americæ australis partibus calidioribus gene-  
ratur pulcherrima hæc avis. Nitore & varietate co-  
lorum plerisque antecellit congeneribus, quarum  
nonnullis obtigit non splendidus, sed paulo obscu-  
rior vestitus.

# BUCCO MAYNANENSIS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

BUCCO MAYNANENSIS

BUCCO MAYNANENSIS

BUCCO MAYNANENSIS

BUCCO MAYNANENSIS

BUCCO MAYNANENSIS

BUCCO MAYNANENSIS

BUCCO MAYNANENSIS

BUCCO MAYNANENSIS

BUCCO MAYNANENSIS

BUCCO MAYNANENSIS







R.N.

London Published

Jan. 9, 1801 by L.P. Nodder, Newman Street.



THE  
MAYNA BARBET.



GENERIC CHARACTER.

*Bill* wedge-shaped, laterally compressed, incurvated at the tip and emarginated on each side : the rictus or gape extending beyond the eyes.

*Nostrils* covered with recumbent feathers.

*Feet* scansorial.

SPECIFIC CHARACTER, &c.

Green BARBET, with the head and throat red, bounded by blue ; the breast yellow, and a red spot on the abdomen.

Le beau Tamatia.

*Buff. ois.* 7. p. 98.

Barbu de Maynas.

*Pl. enl.* 330.

Beautiful BARBET.

*Lath. syn.* 2. p. 498.

---

This beautiful bird is a native of the hotter parts of South America, and in splendor and variety of colors excels most of its congeners, some of which are rather obscure in their plumage.





# MACROURUS RUPESTRIS.



## CHARACTER GENERICUS.

*Cauda attenuata.*

*Bloch. ichth. 5. p. 122.*

## CHARACTER SPECIFICUS, &c.

MACROURUS cauda sensim attenuata.

MACROURUS rupestris.

*Bloch. ichth. 5. p. 123. t. 177.*

*Thoracici.*

Coryphæna rupestris. C. dorso dipterygio, pinnæ dorsalis primæ radio primo retro-dentato.

*Lin. Syst. Nat. Gmel. p. 1195.*

Coryphænoides rupestris.

*Gunner. act. nidr. 3. p. 43. t. 3. f. 1.*

---

Novum hoc genus, a celeberrimo Blochio institutum, unicam continet speciem maria arctica incolentem, crescentemque in longitudinem bipedalem seu tripedalem. A Linnæo *Coryphænis* annumeratum est.

# MACRODONTUS RUFESTRIS.

CHARACTER GENERICUS.

Genus altissimum.

Species unica. p. 122.

CHARACTER SPECIFICUS.

MACRODONTUS rufestris. p. 122.

MACRODONTUS rufestris.

Species unica. p. 122. A. 122.

Florus.

Corphus rufestris. C. dorso digitoque pinnis.

dorsalis pinnis radio pinnis rufestris.

V. M. Stat. N. A. Gmel. p. 119.


Corphus rufestris.

Genus. var. unica. p. 122. A. 122.

Notum hoc, quod a characteribus rufestris non  
differt, sed a rufestris rufestris rufestris  
colorem rufestris rufestris rufestris  
rufestris rufestris rufestris rufestris.








London, Published Jan. 8. 1800 by T. D. Volder, New-mart Street.



THE  
ROCK MACROURUS.



GENERIC CHARACTER.

Tail gradually attenuated.

SPECIFIC CHARACTER, &c.

MACROURUS with very large eyes and gradually attenuated tail.

Der Berglachs.

*Bloch. ichth. t. 177.*

---

The genus *Macrourus*, first instituted by Dr. Bloch, contains only one species, which is a native of the Northern seas, and grows to the length of two, and sometimes three feet. In the arrangement of Linnæus it was considered as a species of *Coryphæna*.





# LIBELLULA GRANDIS.



## CHARACTER GENERICUS.

Os maxillosum : maxillis pluribus.

*Antennæ* thorace breviores.

*Alæ* extensæ.

*Cauda* (maris) hamoso-forcipata.

*Lin. Syst. Nat.* p. 901.

## CHARACTER SPECIFICUS, &c.

LIBELLULA thorace lineis quatuor flavis, corpore variegato.

*Lin. Syst. Nat.* Gmel. p. 2675.

LIBELLULA GRANDIS. L. alis glaucescentibus, thorace lineis quatuor flavis.

*Lin. Syst. Nat.* p. 903.

*Reaum. ins.* 6. t. 35. f. 3.

*Roes. ins.* 2. *aquat.* 2. t. 2. 3.

---

Abunde probat pulcherrima et splendidissima hæc Libellula, quam longe inter se discrepant in eodem animalculo larvæ & insecti perfecti forma atque indoles. Ova, quæ dum aquas prætervolat, deponere solet parens, imum petunt, et certo temporis spatio excluduntur larvæ, quæ adultæ repræsentantur in tabula ; pupa seu chrysalide nihil a larva differente,

nisi



nisi quod appareant alarum principia, brevibus the-  
cis super tergum inclusa. Voracissima larvarum  
gula insectorum imbecilliorum plurimis mortem in-  
fert; ad hoc enim donantur instrumento apprime  
idoneo, forcipis vice fungente, geniculato, quod  
quiescentibus super os retrahitur, ad prædã autem  
captandam subito in longinquum projicitur. Cum  
in aquis vixerit larva duos circiter annos, stipite  
plantæ alicujus aquaticæ inscenso, solis calore gra-  
datim diffringitur cutis, et prorepat insectum plene  
formatum; relictis in caule exuviis quasi illæsis et  
vulneris expertibus.

Cum primum e carcere liberatur, alæ teneræ in  
parvum spatium contrahuntur; elapsa autem qua-  
si semihora ad summum expansis, tentat Libellula  
aeris nisus; ex illo tempore si in aquis nuperrime  
relictis diutius fuerit immersa, non minus mortem  
subitura, quam antea esset larva per idem tempus  
aeri exposita.

Libellulæ insecta imbecilliora voraciter prædantur  
volatu rapidissimo. Mira quoque oculi conforma-  
tio, cujus cornea seu tunica exterior in lenticularum  
æque convexarum millia multa dividitur, clarius  
cernitur in hoc insecto quam in aliis.

Species quam depinximus, e maximis est Libellu-  
larum Europæarum, et per Angliam vulgata. Ico-  
nem elegantem et fidelem ex Roeselii opere cui ti-  
tulus *Insecten Belustigung* qua præstantiorem ex-  
primere desperavimus, imitari et in nostram tabel-  
lam transferre non dubitavimus.





463  
459



R. N.

London, Published Dec 1840, by J. P. Nodder, Newman Street.



THE  
GREAT LIBELLULA.



GENERIC CHARACTER.

*Mouth* consisting of several mandibles.

*Antennæ* very slender, filiform, shorter than the thorax.

*Wings spreading.*

*Abdomen* lengthened.

*SPECIFIC CHARACTER, &c.*

Long-Bodied variegated LIBELLULA, with  
four yellow lines on the thorax.

The variegated LIBELLULA.

GREAT LIBELLULA or Dragon-Fly.

This most beautiful and brilliant insect affords a singular instance of the wonderful diversity of form and manners between the larva and the complete state of one and the same animal. The eggs deposited by the parent insect while it hovers over the waters it frequents, sink to the bottom, and after a certain space, hatch into larvæ, which, when arrived at full growth, are of the figure represented on the annexed plate; the pupa or chrysalis itself differing in no other respect from the larva, than in shewing the rudiments of the future wings, which are enveloped in short cases or processes on the back of the animal. The larvæ are remarkably voracious, and destroy great multitudes of the weaker water-insects, being furnished with a formidable apparatus for seizing their prey, so constructed as to fold over the



the face when at rest, and to be suddenly thrown forwards to a considerable extent when in action. After having remained about two years in this state, the animal ascends the stem of some water-plant, and sitting some time in the sunshine, gives birth to the insect in its perfect or ultimate form, which gradually disengages itself from the skin of the chrysalis, leaving it in its former appearance on the stem. At the period of its first exclusion, the wings, which are then very weak and tender, are complicated into a very short compass; but in about the space of half an hour they become expanded to their full dimensions, and the animal at once commences an inhabitant of the air, and would be as effectually destroyed by a continued submersion under water, as the larva would before have been by exposure to the air.

The Libellulæ in their complete state prey on the smaller insects, and are remarkable for the vigour and celerity of their flight. They also exhibit with greater clearness, and on a larger scale than any other insects, the wonderful structure of the eye, the cornea or exterior coat of which is composed of many thousands of hexagonal, double-convex lenses or segments.

The species here represented is one of the largest of the European Libellulæ, and is very common in our own country. It has been so elegantly and accurately exhibited by Roesel in his *Insecten Belustigung* that it would be in vain to attempt a more expressive representation; for which reason the figures on the present plate are copied from that excellent publication.



# CANCER NORVEGICUS.



## CHARACTER GENERICUS.

*Pedes* octo, (rarius sex aut decem;) insuper  
manus duæ chelatæ.

*Palpi* sex inæquales.

*Oculi* duo distantes, plurimis pedunculati; elongati, mobiles.

*Mandibula* cornea, crassa.

*Labium* triplex.

*Cauda* articulata, inermis.

*Lin. Syst. Nat.* Gmel. p. 2963.

## CHARACTER SPECIFICUS, &c.

CANCER thorace antrorsum aculeato, manibus  
prismaticis, angulis spinosis.

*Lin. Syst. Nat.* p. 1053.

*Astacus* Norwegicus.

*Degeer. ins.* 7. p. 398. t. 24. f. 1.

*Seb. mus.* 3. t. 21. f. 3.


---

Astacum vulgarem magnitudine fere æquat Cancer Norwegicus, in mari arctico præcipue repertus.









London, Published by J. & T. Smith, by F. & J. Nodder, Newman Street.



THE  
NORWAY LOBSTER.



*GENERIC CHARACTER.*

*Legs* generally eight, (in some species six or ten,) besides two claspers or chelated arms.

*Feelers* six, unequal.

*Eyes* two, commonly distant; footstalked, moveable.

*Tail* articulated, unarmed.

*SPECIFIC CHARACTER, &c.*

Long-bodied CANCER, with the thorax aculeated forwards, and prismatic arms with the angles spiny.

NORWAY LOBSTER.

*Br. zool.* 4. p. 12. f. 24.

---

This species is nearly equal in size to the common Lobster, and is principally found in the Northern ocean,

# ALCIDO ATRIALIA

-----

## CHARACTER GENERALIS

Posterior trigonum, crassum, rectum, l. 1. 1. 1.

lingua carnea, brevissima, plana, acuta.

Podas gressori plerique.

Tab. Syst. Nat. p. 178.

## CHARACTER SPECIFICUS

ALCIDO (Aluco) carnea, subita alba, capite

service humilis, tenuis, quae apud nigra

collo infertur torpore, abominanda.

Tab. Syst. Nat. p. 178.

## ALCIDO ATRIALIA

Tab. Syst. Nat. Gmel. p. 178.

Later ex carnis praesentibus merito nomen

tor Alcedo nescitis, magis studiis illud in la-

bula expressa.









THE  
BLACK-CAPPED KINGFISHER.



GENERIC CHARACTER.

*Bill* trigonal, thick, strait, long.

*Tongue* fleshy, very short, flat, sharp-pointed.

*Feet*, in most species, gressorial.

SPECIFIC CHARACTER, &c.

Violet-blue KINGFISHER, with white breast  
and collar, rufous abdomen, and black head  
shoulders and wing-tips.

BLACK-CAPPED KINGFISHER.

*Lath. syn.* 1. p. 624.

Martin-pêcheur de la Chine.

*Pl. enl.* 673.


---

The black-capped Kingfisher may justly be considered as one of the most beautiful of the Chinese birds. It is represented on the plate of half the natural size.










London. Published Feb. 9, 1830, by G. P. Nodder, Newman Street.



THE  
NASAL NOTOCANTHUS.



GENERIC CHARACTER.

Short *prickles* along the back.

SPECIFIC CHARACTER, &c.

NOTOCANTHUS with nose-like snout.

Indian NOTOCANTHUS.

Der Stachelrucken.

*Bloch. ichth.* 12. p. 113. pl. 431.

---

The genus *Acanthonotus* or *Notacanthus* is one of the many new genera instituted by Dr. Bloch. It contains but one species, which is a native of the Indian seas? and is represented of half its natural size on the annexed plate. The particular history and manners of this fish are entirely unknown.

# WASAT NOTOCHTUS

WASAT NOTOCHTUS

WASAT NOTOCHTUS

WASAT NOTOCHTUS

WASAT NOTOCHTUS

WASAT NOTOCHTUS

WASAT NOTOCHTUS

WASAT NOTOCHTUS

The genus *Wasatchia* is one of the many new genera established by the author. It contains but one species, *W. wasatchensis*, which is named in honor of the author's friend and colleague, Dr. J. H. Henshaw. The author's history and figures of the genus are given in the accompanying plates.



# BUCCINUM TUBEROSUM.



## CHARACTER GENERICUS.

*Animal* Limax.

*Testa* univalvis, spiralis, gibbosa.

*Apertura* ovata, desinens in canaliculum (s. retusam lacunam) dextrum, cauda retusum.

*Labium* interius explanatum.

*Lin. Syst. Nat.* p. 1196.

## CHARACTER SPECIFICUS, &c.

BUCCINUM testa cingulis duobus tuberculosus,  
cauda recurva.

*Lin. Syst. Nat.* p. 1198.

*Gualt. test.* t. 41. f. A.

*Seb. mus.* 3. t. 73. f. 5——15.

*Knorr. Vergn.* 3. t. 10. f. 1. 2.

---

In Oceano Americano præcipue innascitur Buccinum tuberosum, magnitudine et coloribus varians, et interdum fere in pedalem crescens longitudinem.

# BUCCINUM TUBEROSUM

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

BUCCINUM TUBEROSUM

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS


CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS

CHARACTERISTICS









# TUBEROUS BUCCINUM.



## GENERIC CHARACTER.

*Animal* allied to a Slug.

*Shell* univalve, spiral, gibbous.

*Aperture* ovate, ending in a channel pointing towards the right.

*Interior Lip* expanded.

## SPECIFIC CHARACTER, &c.

Pale BUCCINUM with rufous and dusky variations, two rows of tubercles, and recurved tip.

Knobbed BUCCINUM,

---

The shell here represented is a native of the American seas, and varies greatly both in size and colors, sometimes measuring almost twelve inches in length.

# TUBEROUS BUCCINUM

1841-1842

General Description

Animal shell to 1.5 inches.  
Shell white, with a  
dorsal band of light brown  
towards the right  
interior of the shell.

The shell is covered with


fine black lines, and the  
aperture, two rows of tubercles, and to  
the right.

Knobbed BUCCINUM

The shell has a  
dorsal band, and the  
aperture, two rows of  
tubercles, and to the right.



# PHALÆNA LECTRIX.



## CHARACTER GENERICUS.

*Antennæ* a basi ad apicem sensim attenuatæ.

*Lingua* spiralis.

*Maxillæ* nullæ.

*Clypeus* (plurimis) corneus brevis.

*Lin. Syst. Nat.* Gmel. p. 2400.

## CHARACTER SPECIFICUS, &c.

PHALÆNA (Noctua) spirilinguis, alis nigris,  
maculis cæruleis flavis albisque, inferioribus  
rubro alboque maculatis.

*Lin. Syst. Nat.* p. 834.

## PHALÆNA (Bombyx) LECTRIX.

*Lin. Syst. Nat.* Gmel. p. 2442.

---

Sinam incolit rarissimum insectum Phalæna Lectrix. In Museo Leveriano asservatur eximium specimen unde delineata est hæc nostra figura.








468



9



## LECTRIX.



### GENERIC CHARACTER.

*Antennæ* setaceous, gradually lessening from the base to the tip.

*Wings* (when sitting) commonly deflected.  
(Flight nocturnal.)

### SPECIFIC CHARACTER, &c.

Moth with black wings, the upper variegated with blue, yellow, and white spots, the lower with red and white.

*Edw.* pl. 318.

*Donov. Chin. Ins.*

---

That extremely rare insect the *Phalæna Lectrix* is a native of China. The beautiful specimen here represented is preserved in the Leverian Museum.

# LECTURE

1884-1885

## General Character

that some serious study is necessary for the

base to the tip.

Wings (when sitting) commonly deflected.

(Flight normal.)

## Sexual Characters, &c.

Stock with black wings. The upper winged

with blue. The lower winged the

lower winged the

lower winged the

lower winged the

That extremely rare. The Chinese

is a native of China. The

is presented in the



# MYCTERIA AMERICANA.



## CHARACTER GENERICUS.

*Rostrum* subascendens, acutum, mandibula superiore triquetra.

*Frons* calva.

*Nares* lineares.

*Pedes* tetradactyli.

*Lath. ind. orn.* p. 670.

## CHARACTER SPECIFICUS, &c.

MYCTERIA alba, remigibus rectricibusque nigro-purpurascentibus.

*Lath. ind. orn.* p. 670.

## MYCTERIA AMERICANA.

*Lin. Syst. Nat.* p. 232.

*Ciconia brasiliensis.*

*Briss. av. 5.* p. 371.

---

Mycteria Americana, quam unicam speciem continere genus non ita pridem opinati sunt physici, Americæ australis loca incolit paludosa, et eodem fere vivendi modo utitur quo ardeæ. Magnitudine ciconiam longe exuperat: immo reperta sunt speci-

mina parum ab ipso struthione distantia. Auctum  
nuperrime est genus novis duabus speciebus, quarum  
altera ex Australasia in Angliam, altera ex India  
illata est.









THE  
AMERICAN JABIRU.



GENERIC CHARACTER.

*Bill* subascendant, sharp, with the upper mandible triquetrous.

*Front* bare.

*Nostrils* linear.

*Feet* tetradactylous.

SPECIFIC CHARACTER, &c.

White JABIRU with black wing-feathers and tail.

Great AMERICAN JABIRU.

---

The American Jabiru, till lately the only species known, is a native of South America, frequenting marshy places, and feeding in the general manner of the Heron tribe. Its size considerably exceeds that of a Stork, and indeed in some specimens almost approaches to that of the Ostrich. This genus has been lately increased by a new species from Australasia, as well as by another from India.





# POLYNEMUS PARADISEUS.



## CHARACTER GENERICUS.

Radii filiformes liberi jugulares.

*Bloch. ichth.* 12. p. 15.

PISCES ABDOMINALES.

## CHARACTER SPECIFICUS, &c.

POLYNEMUS radiis liberis septem jugularibus,  
pinna caudæ bifurca.

*Bloch. ichth.* 12. p. 20.

POLYNEMUS digitis septem, cauda bifida.

*Lin. Syst. Nat.* p. 522.

Piscis *Paradisæa*.

*Edw. t.* 208.

---

Maria Americana incolit Polynemus paradiseus,  
magnitudine dimidiata in tabula depictus.

# TOIYVEMIS PARABIBUS

1888

CHURCH OF THE

TOIYVEMIS PARABIBUS

TOIYVEMIS PARABIBUS

TOIYVEMIS PARABIBUS

CHURCH OF THE

TOIYVEMIS PARABIBUS

TOIYVEMIS PARABIBUS

TOIYVEMIS PARABIBUS

TOIYVEMIS PARABIBUS

TOIYVEMIS PARABIBUS

TOIYVEMIS PARABIBUS


TOIYVEMIS PARABIBUS

TOIYVEMIS PARABIBUS

TOIYVEMIS PARABIBUS











## PARADISE JOURNAL

CONTENTS

CHAP. I.

The first of the month of January, 1841.

The second of the month.

The third of the month.

The fourth of the month.

The fifth of the month.

The sixth of the month.

The seventh of the month.

The eighth of the month.

The ninth of the month.

The tenth of the month.

The eleventh of the month.

The twelfth of the month.

The thirteenth of the month.

The fourteenth of the month.

The fifteenth of the month.



# LIBELLULA DEPRESSA.



## CHARACTER GENERICUS.

Os maxillosum : maxillis pluribus.

*Antennæ* thorace breviores.

*Alæ* extensæ.

*Cauda* (maris) forcipata.

*Lin. Syst. Nat.* p. 901.

## CHARACTER SPECIFICUS, &c.

LIBELLULA abdomine depresso, maris cæruleo,  
feminæ flavo.

LIBELLULA alis omnibus basi nigricantibus,  
thorace lineis duabus flavis, abdomine lanceolato, lateribus flavescente.

*Lin. Syst. Nat.* p. 902.

*Roes.* 2. t. 6. 7.

*Edw. av.* t. 333.

---

Formosæ hujus Libellulæ, cui eadem fere est indoles et vivendi ratio ac reliquis congeneribus, discrepant adeo colores in mari et femina, ut ad speciei distinctionem forma præcipue valeat. Mari est corpus læte cæruleum, maculis in marginibus fuscis; feminæ splendide flavum, maculis iisdem quibus mas conspersum. Non aliter quam *Libellula grandis*, rivos locaque aquosa celerrime prætervolat, sub finem æstatis præcipue conspecta.











THE  
DEPRESSED LIBELLULA.



GENERIC CHARACTER.

*Mouth* consisting of several mandibles.

*Antennæ* very slender, filiform, shorter than  
the thorax.

*Wings* spreading.

*Abdomen* lengthened.

SPECIFIC CHARACTER, &c.

Flat-Bodied LIBELLULA; that of the male  
blue, of the female yellow.

The Flat-Bodied Dragon-Fly.

---

This beautiful insect, the general manners or habits of which resemble those of its congeners, is of so different a colour in the different sexes, that the shape is the chief criterion of the species. The body of the male is of a bright blue, with brown marginal variegations, while that of the female, on the contrary, is of a bright yellow, with similar marginal markings. Like the *Libellula grandis*, this species is principally seen towards the decline of summer, and flies with great rapidity about the neighbourhood of brooks and stagnant waters.





# PAPILIO PERSEUS.



## CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatae.

*Alæ* (sedentis) erectæ sursumque conniventes, (volatu diurno.)

*Lin. Syst. Nat.* p. 774.

## CHARACTER SPECIFICUS, &c.

PAPILIO alis pallide cæruleis apice atris ferrugineo maculatis, subtus undatis: ocellis tribus quatuorque.

*Lin. Syst. Nat.* Gmel. p. 2245.

*Fab. sp. ins.* 2. p. 24.

EQ. ARCHIV.

*Seb. mus.* 4. t. 17. f. 13. 14. & t. 18. f. 15. 16.

*Cram. pap.* 6. t. 71. f. A. B.

---

Inter papiliones Americanos eminet P. Perseus, cujus veram magnitudinem ostendit tabula. In Americæ Australis regionibus calidioribus præcipue conspicitur.









London, Published March 1851, by J. P. Walker, Newman, Street.



# PERSEUS.



## GENERIC CHARACTER.

*Antennæ* thickening towards the upper part,  
and generally terminating in a clavated tip.

*Wings* (when sitting) erect, and meeting upwards. (*Flight* diurnal.)

## SPECIFIC CHARACTER, &c.

**BUTTERFLY** with pale-blue glossy wings,  
blackish and spotted with ferruginous towards the edges: beneath undulated and marked with two or three ocellated spots.

*Seb.* 4. t. 17. 18.

*Cram.* 6. t. 71. A. B.


---

The *Papilio Perseus* may be considered as one of the most elegant of the American Butterflies. It is represented on the plate in its natural size, and is principally found in the hotter regions of South America.





# ORIOLOUS MELANOCEPHALUS.



## CHARACTER GENERICUS.

*Rostrum* conicum, convexum, acutissimum,  
rectum: mandibula superiore paulo longiore,  
obsolete emarginata.

*Lingua* bifida, acuta.

*Pedes* ambulatorii.

*Lin. Syst. Nat.* p. 160.

## CHARACTER SPECIFICUS, &c.

ORIOLOUS luteus, capite remigibus apicibusque  
rectricum intermediarum nigris.

*Lin. Syst. Nat.* p. 160.

*Sturnus* luteolus.

*Lin. Syst. Nat. ed.* 10. p. 167.

ORIOLOUS Bengalensis.

*Briss.* 2. p. 329.

---

In India, Sina, variisque partibus orientalibus,  
nec non in Africa innascitur Oriolus melanocephalus.  
In tabula ostenditur magnitudine naturali  
saltem dimidiata.

...the ...  
...the ...  
...the ...  
...the ...


...the ...  
...the ...  
...the ...  
...the ...

...the ...  
...the ...  
...the ...  
...the ...

...the ...  
...the ...  
...the ...  
...the ...

...the ...  
...the ...  
...the ...  
...the ...








R. A.

London Published April 1850 by F. L. Nodder Newman Street.



## BLACK-HEADED ORIOLE.



### GENERIC CHARACTER.

*Bill* conic, convex, very sharp-pointed, strait :  
the upper mandible rather longer than the  
lower, and slightly emarginated.

*Tongue* bifid, sharp-pointed.

*Feet* formed for walking.

### SPECIFIC CHARACTER, &c.

Yellow ORIOLE, with the head, remiges, and  
tips of the middle tail-feathers black.

Le Lorient de la Chine.

*Buff.* 3. p. 262.

BLACK-HEADED Indian Icterus.

*Edw.* pl. 77.

---

The *Oriolus melanocephalus* is a native of India, China, and many other parts of the East Indies, as well as of some parts of Africa. The plate represents it about half the natural size or rather smaller.

# BLACK-LEADED CRAYON

FOR WRITING AND DRAWING

CHARLES C. CHAMBERS

It is a very fine, very smooth, and very  
the paper is of a light color, and the  
the paper is of a light color, and the  
the paper is of a light color, and the  
the paper is of a light color, and the

the paper is of a light color, and the  
the paper is of a light color, and the  
the paper is of a light color, and the  
the paper is of a light color, and the  
the paper is of a light color, and the

the paper is of a light color, and the  
the paper is of a light color, and the  
the paper is of a light color, and the  
the paper is of a light color, and the  
the paper is of a light color, and the



# CHÆTODON TRICOLOR.



## CHARACTER GENERICUS.

Dentes setacei : corpus pictum.

*Bloch. ichth.* 6. p. 35.

## CHARACTER SPECIFICUS, &c.

CHÆTODON antice flavus, postice niger, cauda  
marginibusque pinnarum rubris.

## CHÆTODON TRICOLOR.

*Bloch. ichth.* 12. p. 97. t. 425.

---

Americana est mirabilis hæc species, primusque  
eam descripsisse videtur celeberrimus Blochius.  
Crescere solet in longitudinem plus pedalem.

# CHARTERED COMPANIES

OF THE UNITED KINGDOM

AND OF THE COLONIES

AND OF THE FOREIGN COUNTRIES

TO WHICH THEY HAVE BEEN GRANTED

CHARTERS BY THE PARLIAMENT

OF GREAT BRITAIN AND OF IRELAND

IN THE SEVENTEENTH AND EIGHTEENTH CENTURIES

BY J. H. COLEMAN

ESQ. OF THE BAR AT LINCOLN'S INN


AND OF THE COURT OF COMMONS

IN THE HOUSE OF COMMONS

IN THE YEAR 1843








London: Published April 1840, by F. R. Waller, Newman Street.



# TRICOLOR CHÆTODON.



## GENERIC CHARACTER.

*Teeth* setaceous. Body varied or banded.

## SPECIFIC CHARACTER, &c.

Yellow CHÆTODON, with the hind part black;  
the tail and fins bordered with red.

The TRICOLOR or triple-coloured CHÆTO-  
DON.

---

This highly singular species, which grows to the length of a foot or more, is a native of the American seas, and seems to have been first described in the work of Dr. Bloch.

# TRICOLOR CHARTER

General Instructions

## General Instructions

These instructions are intended to be read in the order in which they are numbered.

## General Instructions

Yellow CHARTER ON, and the back of the

the (all and the back of the)

THE TRICOLOR CHARTER ON, and the back of the

TRICOLOR

This highly important work is the result of a long and arduous task, and is the work of the TRICOLOR CHARTER ON, and the back of the



# ASTERIAS TOREUMA.



## CHARACTER GENERICUS.

*Corpus* depressum, subtus sulcatum; crusta coriacea tentaculis muricata.

Os centrale quinquevalve.

*Lin. Syst. Nat.* Gmel. p. 3160.

## CHARACTER SPECIFICUS, &c.

ASTERIAS granulata, radiis quinque obtusis margine articulato-tuberculatis.

Stella marina Artocreas.

*Seb.* 3. t. 6. f. 5—10.

*Link.* t. 23. f. 37. t. 24. f. 39. t. 37. f. 45.

---

Asteriarum omnium quas adhuc novimus nulla videtur magis esse coelata et quasi artificis manu e ligno elaborata, quam illa quæ in tabula ostenditur. In maribus Indicis generatur, diversa magnitudine, diametro interdum unciali, interdum sexunciali. Color rubeo-pallet, subtus flavescens seu fere albicans.

# ASTERIAS FORBESII

=====

CHARACTER GENERIQUE

Couleur brun-rouge, surface lustrée; cristallin.

base: 1. surface lustrée.

La surface lustrée.

Long. 1.5 cm. Diam. 0.8 cm.

Caractères distinctifs.

ASTERIAS spiculata, dans l'espèce, obtus.

marginé irrégulièrement.

Stella marina, Atrociss.


Long. 1.5 cm. Diam. 0.8 cm.

Long. 1.5 cm. Diam. 0.8 cm.

Asterias communis, dans l'espèce, obtus.  
videtur in parte conica et parte latere  
hinc, lobata, quia illa pars in lobis  
in variis lobis, quia illa pars in lobis  
dividitur in lobos, quia illa pars in lobis  
Color ruber, sed in lobis, quia illa pars in lobis










## CARVED ASTERIAS.



### *GENERIC CHARACTER.*

*Body* depressed, coriaceous, roughened with small tentacula above.

*Mouth* beneath, central, quinquevalvular.

### *SPECIFIC CHARACTER.*

Granulated ASTERIAS, with five obtuse rays articulated with tubercles on the margin.

---

Of all the Asteriæ yet known, the present is one which has most the aspect of an elaborate artificial composition; appearing as if curiously carved on wood. It is a native of the Indian seas, and is found of various sizes, from an inch to six inches in diameter. Its colour is pale red, yellowish or whitish beneath.

# CARVED ASTERIAS

-----

CHARACTER & HABITS.

Body depressed, coriaceous, rounded with

small reticulate spots.

Mouth beneath, central, transverse.

SPERMATOPHYTES.

Translated Asterias, with the oblique ribs

articulated with tubercles on the margin.

Of all the Asterias known, the present is one

which has most the aspect of an elaborate artificial

composition, appearing as if purposely carved on

wood. It is a native of the Indian sea, and is

found at various parts, from six inch to six feet

in diameter. Its colour is pale red, yellowish or



# LIBELLULA VIRGO.



## CHARACTER GENERICUS.

Os maxillosum ; maxillis pluribus.

*Antennæ* thorace breviores.

*Alæ* extensæ.

*Cauda* (maris) hamoso-forcipata.

## CHARACTER SPECIFICUS, &c.

LIBELLULA viridi-cærulea, alis hyalinis fascia  
lata nigra.

LIBELLULA VIRGO. L. alis erectis coloratis.

*Lin. Syst. Nat.* p. 904.

*Roes.* 2. t. 9.


---

Libellula Virgo, qua species nulla Anglicana vulgarior, matutino tempore, per æstatem prætervolat, ut plurimum, rivulos et loca aquosa. Corporis color generalis est cæruleo-viridis saturatior; alis magnam in medio habentibus maculam seu aream ovatam nigro-cærulescentem. Variant tamen hujus speciei colores magis quam aliarum; interdum enim alæ maculis omnino carent, interdum penitus nigræ sunt; corpore quoque vel viridi-aureo, vel lucido-cæruleo, vel etiam fusco.










R.A.

*London, Published April 18, 1851, by F. & J. Widdow, Newman Street.*



# BLACK-WINGED LIBELLULA.



## GENERIC CHARACTER.

*Mouth* consisting of several mandibles.

*Antennæ* very slender, filiform, shorter than thorax.

*Wings* spreading.

*Abdomen* lengthened.

## SPECIFIC CHARACTER, &c.

Blue-Green LIBELLULA, with a broad black band across the wings.

Black-winged Dragon-Fly.

Banded LIBELLULA.

---

The Libellula Virgo, one of the most common species in this country, is generally seen flying, during the morning hours, about the banks of rivulets, or stagnant waters. The general colour of the body is deep blue-green, while the wings are marked in the middle by a very large patch or area of blueish-black. The insect varies however in point of colour more than any other species, and is sometimes seen with the wings perfectly plain or unmarked, and sometimes, on the contrary, entirely blue-black: the tinge of the body also varies in a similar manner, being either bright golden-green, deep lucid blue, or even sometimes brown.



# STARK COUNTY

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891

1890-1891









## CAYENNE OWL.



### GENERIC CHARACTER.

*Bill* crooked, without cere.

*Nostrils* covered with recumbent bristly feathers.

*Head* large ; *Ears* and *Eyes* large.

*Tongue* bifid.

### SPECIFIC CHARACTER, &c.

Rufous OWL, with numerous, brown, transverse, linear undulations, and yellow eyes.

Chat-huant de Cayenne.

*Buff. ois.* 1. p. 391.

*Pl. Enl.* 442.

## CAYENNE OWL.

*Lath. syn.* 1. p. 146.

---


This elegant species appears to have been first described by the Count de Buffon. It is a native of Cayenne, and is about the size of the common white or barn owl. The plate, of course, represents it as considerably diminished from its natural magnitude.







# MUREX TRITONIS.



## CHARACTER GENERICUS.

*Animal* Limax.

*Testa* univalvis, spiralis, exasperata suturis membranaceis.

*Apertura* desinens in canalem integrum, rectum seu subascendentem.

*Lin. Syst. Nat.* p. 1213.

## CHARACTER SPECIFICUS, &c.

MUREX testa ventricosa oblonga lævi, anfractibus rotundatis, apertura dentata, cauda brevi.

*Lin. Syst. Nat.* p. 1222.

Buccinum TRITONIS.

*Rumph. mus.* t. 28. f. B.

---

Magnificam hanc concham, magnitudine interdum fere sesquipedali, generant præcipue maria Indica. Corniformis cum sit, convertere eam dicuntur pastores Asiatici et Africani in tubam pecuariam; barbaræque nonnullæ gentes in bellicam. Mihi sane persuasissimum est hanc ipsam testam in animo habuisse Ovidium, qui, simul atque audita sit grandisona Tritonis buccina, maris iram coercitam et immanes diluvii aquas recedentes cecinerit.


Jupiter



Jupiter ut liquidis stagnare paludibus orbem,  
Et superesse videt de tot modo millibus unum ;  
Et superesse videt de tot modo millibus unam,  
Innocuos ambos, cultores numinis ambos ;  
Nubila disjecit: nimbisque Aquilone remotis,  
Et coelo terras ostendit, et æthera terris.  
Nec maris ira manet: positoque tricuspile telo  
Mulcet aquas rector pelagi: supraque profundum  
Extantem, atque humeros innato murice tectum  
Caeruleum Tritona vocat; conchæque sonaci  
Inspirare jubet; fluctusque et flumina signo  
Jam revocare dato. Cava buccina sumitur illi  
Tortilis, in latum quæ turbine crescit ab imo:  
Buccina, quæ medio concepit ut aëra ponto,  
Littora voce replet sub utroque jacentia Phoebo.  
Tum quoque, ut ora Dei madida rorantia barba  
Contigit, et cecinit jussos inflata receptus,  
Omnibus audita est telluris et æquoris undis:  
Et quibus est undis audita, coercuit omnes.  
Jam mare littus habet: plenos capit alveus amnes:  
Flumina subsidunt: colles exire videntur.  
Surgit humus: crescunt loca decrescentibus undis.”

Color illi generalis est albidus seu albo-flavescens, zonis transversis plus minus fuscis, ferrugineis, seu nigricantibus undulatus. Variant enim hujus nec non plerarumque concharum colores ob ætatem, et aliis multis de causis.









THE  
TRITONIAN MUREX.



*GENERIC CHARACTER.*

*Animal* allied to a Slug.

*Shell* univalve, spiral, roughened with membranaceous sutures.

*Aperture* ending in a strait or subascending channel.

*SPECIFIC CHARACTER, &c.*

MUREX with large, oblong, smooth, whitish shell, with transverse ferruginous undulations, toothed aperture, and short top.

The Sea Trumpet, or great clouded MUREX.

Triton's Trumpet.

---

This superb shell, which is sometimes seen of near a foot and a half in length, is principally found in the Indian seas. From its peculiar form it is well adapted for the purpose of a horn or trumpet, and is said to be often used as such by the African and Asiatic shepherds. It is said to be occasionally used as a military trumpet among barbarous nations. We may also conclude it to have been the shell intended by



by Ovid in his description of the retiring of the waters of the deluge on the sound of the trumpet of Triton.

But when th' all-powerful ruler of the sky  
Saw earth in one wide waste of waters lie,  
While late of all its peopled realms contain'd  
One only pair of human race remain'd,  
Innocuous both, resign'd to Heav'n's decree,  
Celestial pity touch'd the Deity.  
At his command, by winds resistless driv'n,  
Dispersing clouds to earth unveil the heav'n :  
And Neptune bids blue Triton sound amain  
His powerful trump, and bind the waves again.  
The sea-born herald at the call appears;  
High o'er the surge his scaly shoulders rears;  
And while the foaming billows round him swell  
Grasps with his bearded lip the spiral shell.  
With such a blast the sounding conch he blew  
O'er all the globe at once the thrilling signal flew.  
Back to their ancient bounds, from pole to pole  
Th' alarmed seas in reflux circles roll:  
The hills emerge; the woods their branches shew;  
And earth restor'd peeps slowly from below.

This shell is generally of a whitish or yellowish colour, richly variegated with deep and pale ferruginous, brown, and blackish, transverse undulations; varying, like all other shells in the intensity and beauty of its colours, according to age or other circumstances.



# PERCA GUTTATA.

\*\*\*\*\*

## CHARACTER GENERICUS.

*Squamæ duræ, asperæ.*

*Opercula spinosa.*

*Bloch. ichth. 2. p. 56.*

*Thoracici.*

## CHARACTER SPECIFICUS, &c.

PERCA rubra guttis rubris.

*Bloch. ichth. 9. p. 78. t. 312.*

PERCA pinnis dorsalibus unitis, cauda integra,  
corpore punctis sanguineis adperso.

*Lin. Syst. Nat. p. 485.*

Gugupuguacu.

*Catesb. 2. p. 14. t. 14.*

---

Perca guttata, quam satis abundanter alunt maria Americana, in cibis est laudatioribus; perca communi Europea seu fluviatili fere duplo major. Color communis pulcherrime rubet, maculis plurimis obscurioribus conspersus.

# PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA


PERCA CUTTATA

PERCA CUTTATA

PERCA CUTTATA








London Published May 1880 by F. & J. Widdow, Newman Street.



THE  
SANGUINE PERCH.



GENERIC CHARACTER.

*Gill-Covers* spiny.

*Scales* hard and rough.

SPECIFIC CHARACTER, &c.

Red PERCH, with numerous deep-red spots.

The Hind.

*Catesb. Car.* 2. p. 14. pl. 14.

La Sanguinolente.

*Bonaterre Encycl. Ichth.* p. 130.

The Hind, or red spotted Perch.

---

The *Perca guttata*, a very common inhabitant of the West Indian seas, and, in general, much esteemed as an article of food, grows sometimes to near double the size of the common European or river Perch. Its general colour is a most beautiful red, with numerous round spots of a deeper tinge.

# SANGUINE PERCH

\*\*\*\*\*

## GENERAL CHARACTER

Gen. Character: small, slender, hard and tough.


## Scientific Characters

Red PERCH, with numerous deep red spots.  
The Head.  
Length 1.5 ft. 10 in. 14.  
In Sanguine.  
Locality: Florida, Lake p. 100.  
The Head, or red spotted perch.

The Perch is a very common inhabitant of the West Indies, and is found in abundance as an article of food, and is sometimes used as an article of the table. It is a very good fish, its general color is a most beautiful red, with numerous round spots of a darker red.



# LIBELLULA PUELLA.



## CHARACTER GENERICUS.

Os maxillosum : maxillis pluribus.

*Antennæ* thorace breviores.

*Alæ* extensæ.

Cauda maris hamoso-forcipata.

*Lin. Syst. Nat.* p. 901.

## CHARACTER SPECIFICUS, &c.

LIBELLULA alis erectis hyalinis, corpore cæruleo vel rubido.

LIBELLULA PUELLA. L. alis erectis hyalinis.

*Lin. Syst. Nat.* p. 905.

Agrion PUELLA.

*Fab. sp. ins.* p. 527.

*Roes.* 2. t. 10. 11.

*Reaum. ins.* 6. t. 35. f. 6.

*Merian Eur.* 78. t. 156.

---

Loca aquosa sæpissime circumvolitare solet Libellula Puella æstivo tempore. Hanc specie nulla Britannica minor aut vulgatioꝛ. Variat color corporis non minus quam Libellulæ Virginis; vel læte cyaneus, vel rufo-fuscus, vel nigricans. Certus tamen et constans alarum color, scilicet hyalinus; macula fusca parvula prope marginum extremitates ducta.









London Published May 1804 by F. & J. Neider, Newman Street.



## SMALL LIBELLULA.



### GENERIC CHARACTER.

*Mouth* consisting of several mandibles.

*Antennæ* very slender, filiform, shorter than the thorax.

*Wings* spreading.

*Abdomen* lengthened.

### SPECIFIC CHARACTER, &c.

LIBELLULA with upright hyaline wings, and sky-blue or reddish body.

Small Dragon-Fly.

Little slender-bodied LIBELLULA.


---

This is the least of the British Libellulæ, and is a very frequent insect in the neighbourhood of watery places during the summer months. In the colour of its body it varies as much as the L. Virgo, and is either of a bright sky-blue, reddish-brown, or blackish: the wings however are constantly of a similar appearance, or transparent, with a small, dark marginal spot towards their extremities.





# VAGINALIS AUSTRALIS.



## CHARACTER GENERICUS.

*Rostrum* robustum, conico-convexum, compressum, mandibula superiore vagina mobili cornea tecta.

*Facies* nuda, papillosa.

## CHARACTER SPECIFICUS, &c.

VAGINALIS alba, facie papillosa.

VAGINALIS Chionis.

*Lath. ind. orn.* p. 774.

VAGINALIS alba.

*Lin. Syst. Nat. Gmel.* p. 705.

---

Genus *Vaginalis* ex iis est novis ob avium nuperime detectarum abundantiam necessario institutis. Constat ex unica specie cui magnitudo est quasi columbæ vulgaris. Australasiam insulasque maris pacifici præcipue incolit.











THE  
SOUTHERN SHEATHBILL.



GENERIC CHARACTER.

*Bill* strong, conically-convex, compressed; the upper mandible covered at the base by a moveable horny sheath.

*Face* naked, covered with papillæ.

SPECIFIC CHARACTER, &c.

White SHEATHBILL, with scattered papillæ about the face.

The Southern or White SHEATHBILL.

White SHEATHBILL.

*Lath. syn.* 3. p. 268.

---

The genus Vaginalis or Sheathbill is one of the new genera which, in consequence of the late discoveries in ornithology, it has been found necessary to institute. The species here represented is the only one at present known, and is about the size of a common Pigeon. It is a native of New Holland and of the scattered Southern islands.

# EXOTIC FLORA

## General Description

The plant is a small, bushy shrub, growing to a height of about 10 feet. It has a woody stem and is covered with small, dark, glossy leaves. The flowers are small and white, and the fruit is a small, round, red berry. The plant is native to the tropics and is found in the mountains of the East Indies.

## The Fruit of the Plant

The fruit is a small, round, red berry, about the size of a pea. It is covered with a thin, waxy skin and has a sweet, juicy flesh. The seed is small and round, and is covered with a thin, brown skin.

The plant is a small, bushy shrub, growing to a height of about 10 feet. It has a woody stem and is covered with small, dark, glossy leaves. The flowers are small and white, and the fruit is a small, round, red berry. The plant is native to the tropics and is found in the mountains of the East Indies.



# SCARUS CROICENSIS.



## CHARACTER GENERICUS.

*Caput*: *Dentium* loco maxillæ ipsæ eminentes;  
margine dentato-crenatæ osseæ.

*Membrana branchiostega* radiisquinque; Oper-  
culum integerrimum.

*Corpus*: *Linea* lateralis plurimis ramosa.

*Lin. Syst. Nat. Gmel. p. 1280.*

## CHARACTER SPECIFICUS, &c.

SCARUS pinna caudæ rotundata.

*Bloch. ichth. 7. p. 18. t. 221.*

*Pisc. Thoracici.*


---

Maria Indica et Americana incolit Scarus Croi-  
censis, a Blochio primum depictus. Coloribus va-  
riat; interdum totus ruber, interdum fasciis ar-  
genteo-albis insignitus. In pedalem crescit longi-  
tudinem.









London: Published for W. & A. G. & Co. by J. B. Notholt, a Freeman & Son.

Thalassoma



THE  
RED SCARUS.



GENERIC CHARACTER.

Strong *bony processes*, crenated at the edges,  
instead of teeth.

SPECIFIC CHARACTER, &c.

RED SCARUS, with rounded tail, and the body  
sometimes marked by silvery-white stripes.

Der rothe Papagey-Fisch.

*Bloch. ichth. pl. 221.*

---

This elegant species, measuring about twelve inches in length, is a native of the Indian and American seas and was first figured in the work of Dr. Bloch. In color it varies; being sometimes entirely red, and sometimes marked with longitudinal silvery-white stripes.

THE  
RED SCARUS

\*\*\*\*\*

CLARENCE CLARK, JR.

Strong form, slender, rounded at the edges,  
inches of length.

CLARENCE CLARK, JR.

RED SCARUS, with white spots and the body  
somewhat flattened, white stripes.

For to the 10 pages of the  
book, which is 231.

---

The length of the body is about twelve  
inches in length, and the head is about  
one inch and a half long, and the body of Dr.  
Smith. The color is red, and the body is  
red, and the head is red, and the body is  
very thin and



# PINNA MURICATA.

\*\*\*\*\*

## CHARACTER GENERICUS.

*Animal* Limax.

*Testa* subbivalvis fragilis, erecta hians, emittens  
barbam byssinam.

*Cardo* edentulus, coalitis in unam valvis.

*Lin. Syst. Nat. Gmel. p. 3363.*

## CHARACTER SPECIFICUS, &c.

PINNA testa striata, squamis concavis, ovatis,  
acutis.

*Lin. Syst. Nat. p. 1160.*

Concha Pinna.

*Hasselq. itin. 447.*

*Varietates potius quam distinctæ species sunt  
plures hujus generis.*


---

Genus *Pinnæ*, cujus nonnullæ species in peram-  
plam crescunt magnitudinem, præter cæteros cha-  
racteras hoc quoque habet insigne, quod animal  
testam incolens ab apice linguæ tubulatæ sæpissime  
exertæ et retractæ guttulam glutinosam deponat,  
unde efficitur quasi sericus innumerarum fibrarum  
fasciculus, e quo suspensum rupibus aliisque id genus  
tuto

tuto se possit affigere. Inter species minores est Pinna muricata, longa nempe sex vel octo uncias. Color et superficiei scabrities longe variant. Plerumque tamen color exterior est fusco-saturator, interior subrubrus cum nitore quodam margaritaceo. Pinnarum byssus seu sericum aureo-fuscum in chirothecas aliasque parvulas vestes interdum convertitur ab incolis orarum mediterranei maris, Italiæ nempe, Siciliae, &c. &c. Notandum porro est posse se suspendere sericis filamentis, ab animali testam incolente contextis, Pinnæ generi non proprium omnino et peculiare esse, sed et a nonnullis Mytylorum speciebus, parcius tamen et contractius, eandem artem exerceri.







R.V.

*London Published June 1<sup>st</sup> 1801, by T. R. Nodder, Newman Street.*



THE  
MURICATED PINNA.



GENERIC CHARACTER.

*Animal* allied to a Limax.

*Shell* subbivalve, fragile, gaping at the upper part, emitting a beard or tuft of silken filaments.

*Hinge* toothless, the valves coalescing together.

SPECIFIC CHARACTER, &c.

PINNA with striated shell, with acute, ovate, concave scales.

*Gualt. test.* t. 79. f. D.

*Seb.* 3. t. 92.

*Obs.* Several of the Pinnæ are considered by Linnæus as varieties rather than distinct species.

---

The genus Pinna, of which some species grow to a very large size, is distinguished, among other circumstances, by the remarkable power which the inhabiting animal possesses, of affixing itself at pleasure to rocks, or other substances, by a vast number of fine silky threads or filaments, which it forms by extending



extending its tubular trunk or tongue, and discharging from its tip a minute drop of gluten, which, by the retraction of the same organ, is of course formed into a silken filament; and this operation being several thousand times repeated, a thick and beautiful tuft of silky fibres is composed, by the help of which the animal is securely fastened or anchored in such places as it finds convenient. The *Pinna muricata* is among the smaller species of this genus; measuring about six or eight inches in length. In color, as well as in the degree of roughness of its external surface, it varies considerably. It is commonly of a deep brown externally, and of a reddish tinge, varied with a degree of pearly lustre internally. The silk or byssus of the *Pinnæ* is of a rich golden brown color, and is occasionally manufactured into gloves, &c. by the inhabitants of those parts of the Mediterranean coasts where it most abounds, viz. on some of the coasts of Italy and Sicily. It should be added, that the power of adhering by means of silken filaments thus drawn from the inhabiting animal, is not entirely confined to the genus *Pinna*, but takes place, in a smaller degree, in some species of *Mytilus*.



# PAPILIO XUTHUS.



## CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatae.

*Alæ* (sedentis) erectæ sursumque conniventes, (volatu diurno.)

*Lin. Syst. Nat.* p. 774.

## CHARACTER SPECIFICUS, &c.

PAPILIO alis nigris albo striato maculatis; posterioribus subtus ocellis cæruleis fulvisque subfasciatis.

*Fab. sp. ins.* 2. p. 19.

PAPILIO alis caudatis nigris albido striato-maculatis; posticis subtus cæruleo fulvoque subocellatis.

*Lin. Syst. Nat.* p. 751.

---

Inter Lepidoptera rariora numeratur Papilio Xuthus, Indiæ orientalis incola. Tabula magnitudine vera pictum exhibet.










21

184



# XUTHUS.



## GENERIC CHARACTER.

*Antennæ* thickening towards the upper part, and generally terminating in a clavated tip.

*Wings* (when sitting) erect, and meeting upwards. (Flight diurnal.)

## SPECIFIC CHARACTER, &c.

Black subcaudated Butterfly, with yellowish white striated spots; the lower wings marked beneath with a band of blue and orange ocellated spots.

---

The *Papilio Xuthus* is numbered among the rarer species of *Lepidoptera*, and is a native of the East Indies. The plate represents it in its natural size.





# FALCO FORMOSUS.



## CHARACTER GENERICUS.

*Rostrum* aduncum, *cera* instructum.

*Caput* *pennis* arcte tectum.

*Lingua* bifida.

*Lin. Syst. Nat.* p. 124.

## CHARACTER SPECIFICUS, &c.

FALCO *cera* orbitis pedibusque luteis, jugulo  
purpureo, corpore supra cærulescente-rubro,  
abdomine purpureo.

*Lath. ind. orn.* p. 38.

FALCO *aquilinus*.

*Lin. Syst. Nat. Gmel.* p. 280.

---

Sive formæ sive colorum habeatur ratio, merito  
numerari debet hæc avis in pulcherrimis sui generis.  
Americam meridionalem incolit, magnitudine Fal-  
coni cyaneo fere æqualis.









R.V.

London Published by W. Woodley Newman Street



THE  
PURPLE-THROATED FALCON.



GENERIC CHARACTER.

*Bill* hooked; with a cere or naked skin at the base.

*Head* thickly beset with feathers.

*Tongue* generally bifid.

SPECIFIC CHARACTER, &c.

Blue FALCON, with purple throat, and reddish abdomen.

Red-Throated FALCON.

*Lath. syn.* 1. p. 97.

Petit Aigle d'Amerique.

*Buff.* 1. p. 142.

*Pl. Enl.* 417.


---

This bird, whether we consider its shape or colors, must be numbered among the most beautiful of the genus. It is a native of South America, and is nearly equal in size to the *Falco cyaneus* or Hen-Harrier.





# PAPILIO ORONTES.



## CHARACTER GENERICUS.

*Antennæ* apicem versus crassiores, sæpius clavato-capitatae.

*Alæ* (sedentis) erectæ, sursumque conniventes, (Volatu diurno.)

*Lin. Syst. Nat.* p. 744,

## CHARACTER SPECIFICUS, &c.

PAPILIO alis nigris, fasciis duabus virescentibus, caudis albis distantibus.

*Lin. Syst. Nat.* Gmel. p. 2237.

PAPILIO alis caudatis nigris, albido-virescente fasciatus, caudis albis distantibus.

*Lin. Syst. Nat.* p. 750.

---

Inter rariores hujus generis est *Papilio Orontes*, magnitudine vera in tabula depictus. Indiam incolit, coloribus interdum leviter varians.

# TABULAE ORONTES

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes

Tabulae Orontes









# ORONTES.



## GENERIC CHARACTER.

*Antennæ* terminating generally in a clavated tip.  
*Wings* (when sitting) meeting upwards. Flight  
diurnal.

## SPECIFIC CHARACTER.

Black Butterfly, with two greenish bands, and  
two distant white tails or processes.

The Green-Banded tailed Butterfly.

---

The *Papilio Orontes* is one of the rarer species of  
Butterflies, and is represented on the plate in its na-  
tural size. It is a native of India, and sometimes  
varies a little in point of color.

# ONIONTS

-----

ONIONTS

ONIONTS  
ONIONTS (when sitting) meeting  
ONIONTS

ONIONTS  
ONIONTS  
ONIONTS

ONIONTS

ONIONTS  
ONIONTS  
ONIONTS

ONIONTS

ONIONTS

ONIONTS

ONIONTS

ONIONTS

ONIONTS

ONIONTS

ONIONTS

ONIONTS

ONIONTS



# ANTHIAS FORMOSUS.



## CHARACTER GENERICUS.

*Caput* totum squamosum; operculo anteriore serrato.

*Bloch. ichth.* 9. p. 86.

## CHARACTER SPECIFICUS, &c.

ANTHIAS flavus, lineis longitudinalibus cæruleis.

*Perca formosa.*

*Lin. Syst. Nat.* Gmel. p. 1322.

*Perca marina* capite striato.

*Catesb. Car.* 2. t. 6.

---

In maribus Americanis præcipue conspicitur *Anthias formosus*, longitudine, ut plurimum, pedali.

# ANTHUS FORMOSUS

CHARACTER SPECIES

CHARACTER SPECIES

CHARACTER SPECIES

CHARACTER SPECIES

CHARACTER SPECIES

CHARACTER SPECIES

CHARACTER SPECIES

CHARACTER SPECIES

CHARACTER SPECIES

CHARACTER SPECIES

CHARACTER SPECIES


CHARACTER SPECIES

CHARACTER SPECIES

CHARACTER SPECIES







London: Published July 18 1851. by W. R. Nodder, & Newman, Proprs.



THE  
BLUE-STRIPED ANTHIAS.



GENERIC CHARACTER.

*Head* completely scaled: anterior gill-cover  
serrated.

SPECIFIC CHARACTER, &c.

Yellow ANTHIAS with longitudinal blue lines.

The Grunt.

*Catesb. Carol.* 2. p. 6.

L'Ecuriel.

*Bonaterre Encycl. Ichth.* p. 135.

Der Rothmund.

*Bloch. ichth.* pl. 323.

---

This beautiful fish is principally found in the American seas, and is generally about twelve inches in length.





# MUREX HAUSTELLUM.

\*\*\*\*\*

## CHARACTER GENERICUS.

*Animal* Limax.

*Testa* univalvis, spiralis, exasperata suturis  
membranaceis.

*Apertura* desinens in canalem integrum rectum  
s. subascendentem.

## CHARACTER SPECIFICUS, &c.

MUREX testa ovata, tuberculata, cauda elon-  
gata subulata recta muricata.

## HAUSTELLUM.

*Rumph. mus.* t. 26. f. F.

*Argenv. conch.* t. 19. f. B.

---

Muricem Haustellum magnitudine vera depictum  
ostendit tabula. In mari Asiatico præcipue innas-  
citur.

# MURPHY HAUSTORIUM

HAUSTORIUM

HAUSTORIUM

HAUSTORIUM

HAUSTORIUM

HAUSTORIUM

HAUSTORIUM

HAUSTORIUM

HAUSTORIUM

MURPHY HAUSTORIUM

HAUSTORIUM

HAUSTORIUM

HAUSTORIUM

HAUSTORIUM

Murphy Haustorium

ostendit capula. In anti. Apertio

clum.









THE  
PIPE MUREX.



GENERIC CHARACTER.

*Animal* allied to a Slug.

*Shell* univalve, spiral, roughened by membranaceous sutures.

*Aperture* ending in a strait or subascending channel.

SPECIFIC CHARACTER, &c.

MUREX with ovate tuberculated shell, with an elongated tapering muricated tip.

La Becasse.

*Argenv.* p. 257. pl. 16. f. B.

---

This shell is represented on the plate in its natural size. It is chiefly found in the Asiatic seas.





# TROCHILUS MINIMUS.



## CHARACTER GENERICUS.

*Rostrum* subulato-filiforme, apice tubulato, capite longius: Mandibula superior vaginans inferiorem.

*Lingua* filiformis, filis duobus coalitis tubulosa.

*Pedes* ambulatorii.

*Lin. Syst. Nat.* p. 189.

## CHARACTER SPECIFICUS, &c.

TROCHILUS rectirostris fusco-virescens, subtus albidus, rectricibus lateralibus margine exterioriore albis.

TROCHILUS MINIMUS. T. rectirostris, rectricibus lateralibus margine exterioriore albis, corpore viridi nitente subtus albido.

*Lin. Syst. Nat.* p. 193.

Guainumbi minor corpore toto cinereo.

*Raj. syn.* p. 83.

---

Perpusillum hoc Naturæ miraculum irradiat nullus colorum variorum splendor, quo plurimo gaudet fere reliquum genus. Color ei generalis virescit quasi sub-aureo fuscus, dorso humerisque paulo lucidioribus.

M

bus.

bus. Alæ caudaque e purpureo fusca. Corpus inferius fere albet. Minus est saturati nitoris feminae quam mari. Avium omnium minimus est Trochilus de quo jam loquimur, longus scilicet ab apice rostri ad extremum caudæ unciam cum quadrante. In plurimis Americæ australis regionibus, interdum etiam in insula Jamaica conspicitur.








R.N.

London Published August 1801, by Elizabeth Nodder, Sons & Co. Newman Street.



THE  
LEAST HUMMING-BIRD.



GENERIC CHARACTER.

*Bill* slender, tubular, the upper mandible sheathing the lower.

*Tongue* very long, extensile; formed of two conjoined cylindric tubes.

*Toes* three forward, one backward.

SPECIFIC CHARACTER, &c.

Brownish - green strait - billed HUMMING-BIRD, whitish beneath, with the exterior tail-feathers white on the outside.

LEAST HUMMING-BIRD.

*Edw.* pl. 105.

Le plus petit Oiseau-Mouche.

*Pl. Enl.* 276.

---

This little miracle of Nature is not distinguished by the peculiar splendor and variety of colour so conspicuous in many others of the genus, its general tinge being an obscurely gilded greenish brown above, and whitish beneath: the wings and tail have a cast of purplish brown: the colour of the



female is more obscure than that of the male: the usual length of this species is about an inch and quarter from the tip of the bill to the end of the tail, being by far the least of all birds. It is a native of many parts of South America, and is sometimes found in the island of Jamaica.



## PHALÆNA FRAXINI.



### CHARACTER GENERICUS.

*Antennæ* setacæ, a basi ad apicem sensim attenuatæ.

*Alæ* (sedentis) sæpius deflexæ. (Volatu nocturno.)

*Lin. Syst. Nat.* p. 808.

### CHARACTER SPECIFICUS, &c.

PHALÆNA (Noctua) spirilinguis cristata, alis dentatis cinereo-nebulosis, inferioribus supra nigris, fascia cærulescente.

*Lin. Syst. Nat.* p. 843.

*Roes.* 4. t. 28. f. 1.

*Mer. Eur.* t. 46.

---

In Germania aliisque nonnullis Europæis regionibus frequentius nascitur Phalæna fraxini, quam in maximis et rarissimis, insectis quotquot *lepidoptera* vocantur, numerat Britannia. Larva ejus fusco-pallida fraxini et salicis folia præcipue depascitur, et in chrysalidem leviter lanuginosam et fuscam mense Junio convertitur, e qua mense qui proxime insequitur, erumpit Phalæna. A mari illico dignoscitur femina, quod longe major sit.











THE  
CLIFDEN MOTH.



GENERIC CHARACTER.

*Antennæ* setaceous, decreasing in size from the base to the tip.

*Wings*, when at rest, generally deflected.

*Flight* nocturnal.

SPECIFIC CHARACTER, &c.

MOTH with crested thorax and denticulated wings, the upper pair grey clouded with brown, the lower black with a broad transverse blueish band.

The CLIFDEN MOTH. Clifton Beauty. Non-pareil-Moth, &c. &c.


---

The *Phalæna Fraxini*, one of the rarest as well as largest of the British *Lepidoptera*, is much less uncommon in Germany and several other parts of Europe; its larva, which is of a pale brown colour, feeds principally on the ash and willow, and changes into a slightly hairy brown chrysalis in the month of June, the Moth making its appearance in the following month. The male is easily distinguished from the female by its smaller size.





# LUTIANUS CHRYSOPS.



## CHARACTER GENERICUS.

*Opercula* squamata, serrata, inermia.

*Bloch. ichth.* 7. p. 84.

## CHARACTER SPECIFICUS, &c.

LUTIANUS violaceus, oculis aureis.

LUTIANUS radiis tribus spinosis tredecimque  
mollibus in pinna ani.

*Bloch. ichth.* 7. p. 91. t. 248.

PISC. THORACICI.

---


Genus Lutianus, a Blochio primum institutum, varias continet species, quarum illa quæ in tabula depingitur maria incolit Americana? magnitudine, ut plurimum Clupeæ vulgaris.







40



BN



THE  
GOLDEN-EYED LUTIAN.



GENERIC CHARACTER.

*Gill-Covers* scaly, serrated, unarmed.

SPECIFIC CHARACTER, &c.

Violet-coloured LUTIAN with gold-coloured eyes.

Das Goldauge.

*Bloch. ichth. t. 248.*

---

The genus *Lutianus*, instituted by Dr. Bloch, contains several species, of which that represented on the present plate is a native of the American seas, and usually about the size of a common Herring.





# OSTREA PALLIUM.



*Animal* Tethys.

*Testa* bivalvis, (plurimis inæquivalvis,) subaurita.

*Cardo* edentulus, fossula cava ovata striisque-lateralibus transversis.

*Lin. Syst. Nat.*

*CHARACTER SPECIFICUS, &c.*

OSTREA testa æquivalvi, radiis duodecim convexis, striata, scabra, squamis imbricata.

*Lin. Syst. Nat.* p. 1145.

*Pecten* secundus.

*Rumph. mus.* t. 44. B.

PALLIUM Ducale.

*Argenv.* t. 24. f. 1.

---

Inter testas pulcherrimas hujus generis merito numeratur Ostrea Pallium, coloribus plurimum varians. In oris Indicis præcipue conspicitur, et magnitudine vera in tabula exprimitur.

THE  
MANTLE SCALLOP.



GENERIC CHARACTER.

*Animal* resembling a Tethys.

*Shell* bivalve, (in most species unequally,) sub-auriculated.

*Hinge* toothless, with an ovate or subtrigonal fossule and lateral transverse streaks.

SPECIFIC CHARACTER, &c.

Variegated SCALLOP with equal valves, striped by twelve rough, imbricated, scaly rays.

The Ducal MANTLE SCALLOP.


Le Manteau Ducal.

*Argenv. Knorr. &c. &c.*

---

This shell is very justly considered as one of the most beautiful of its tribe: it varies much in colour, and is principally found about the coasts of the Indian seas: it is represented on the plate in its natural size.









## INDEX.

PL.

465. Alcedo atricapilla.  
 487. Anthias formosus.  
 454. Apis lapidaria.  
 475. Asterias Toreuma.  
 447 438. Bulla achatina.  
 461. Bucco Maynanensis.  
 467. Buccinum tuberosum.  
 471. Chætodon tricolor.  
 464. Cancer Norvegicus.  
 449. Columba rosea.  
 457. ——— coronata.  
 446. Exocoetus evolans.  
 485. Falco formosus.  
 459. Libellula grandis 463  
 471. ——— depressa.  
 476. ——— Virgo.  
 480. ——— Puella.  
 491. Lutianus Chrysops.  
 469. Myceteria Americana.  
 445. Mergus Castor.  
 462. Macrourus rupestris.  
 460. Murex Tribulus.  
 479. ——— Tritonis.  
 488. ——— Haustellum.  
 452. Nais digitata.  
 466. Notocanthus Nasus.  
 478. Oriolus melanocephalus.  
 492. Ostrea Pallium.  
 472. Papilio Perseus.  
 486. ——— Orontes.  
 448. ——— Amphimedon.  
 459. ——— Phorcas.  
 484. ——— Xuthus.  
 456, 455. Phalæna Junonia  
 468. ——— Lectrix.  
 490. ——— Fraxini.  
 478. Perca guttata.  
 483. Pinna muricata.  
 470. Polynemus paradiseus.

## INDEX.

PL.

475. Anthias blue-striped.  
 487. Asterias carved.  
 461. Barbet Mayna.  
 454. Bee orange-tailed.  
 467. Buccinum tuberosus.  
 438. Bulla agate. 447  
 472. Butterfly Perseus.  
 459. ——— Phorcas.  
 484. ——— Xuthus.  
 486. ——— Orontes.  
 448. ——— Amphimedon.  
 471. Chætodon tricolor.  
 446. Flying-Fish.  
 485. Falcon purple-throated.  
 451. Gurnard flying.  
 489. Humming-Bird least.  
 469. Jabiru American.  
 465. Kingfisher Black-capped  
 459. Libellula great. 463  
 471. ——— depressed.  
 476. ——— black-winged.  
 480. ——— small.  
 464. Lobster Norway.  
 491. Lutian golden-eyed.  
 462. Macrourus rock.  
 445. Merganser dun.  
 468. Moth Lectrix.  
 455, 456. Moth Peacock.  
 490. Moth Clifton.  
 460. Murex thorny.  
 479. ——— Tritonian.  
 488. ——— pipe.  
 452. Nais digitated.  
 466. Notocanthus nasal.  
 477. Owl Cayenne.  
 478. Oriole black-headed.  
 478. Perch sanguine.  
 449. Pigeon crimson.  
 457. ——— crowned.  
 483. Pinna muricated.

482. *Scarus Croicensis*.  
458. *Sparus chrysurus*.  
477. *Strix cayana*.  
450. *Tubularia magnifica*.  
453. *Trichoda Sol.*  
451. *Trigla volitans*.  
489. *Trochilus minimus*.  
481. *Vaginalis Australis*.


✦ 470. *Polynemus Paradise*.  
✦ 481. *Sheath-Bill Southern*.  
✦ 482. *Scarus red*.  
✦ 492. *Scallop Mantle*.  
✦ 458. *Sparus yellow-striped*.  
✦ 450. *Tubularia great*.  
✦ 453. *Trichoda Sun*.  
✦


















RB  
500.9  
S534



