

The naturalists' miscellany, or, Coloured figures of natural objects; drawn and described immediately from nature.

Contributors

Shaw, George, 1751-1813.
Nodder, Frederick Polydore
Nodder, Richard P

Publication/Creation

[London] : [Printed for Nodder & co.], [1790-1813]

Persistent URL

<https://wellcomecollection.org/works/xgtyy48u>


License and attribution


This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.





Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>




















VIRIS ORNATISSIMIS
REGIÆ INSTITUTIONIS
LONDINENSIS
FUNDATORIBUS ET PATRONIS

HUNC
DECIMUM OCTAVUM
NATURÆ VIVARII

FASCICULUM

D. D. D.

GEORGIUS SHAW,
E. NODDER.

TO THE
ROYAL INSTITUTION
OF
LONDON,
DISTINGUISHED BY ITS
ZEALOUS PROMOTION
OF
USEFUL KNOWLEDGE,
THIS EIGHTEENTH VOLUME
OF THE
NATURALIST'S MISCELLANY
IS
RESPECTFULLY INSCRIBED
BY
GEORGE SHAW,
E. NODDER.

TINAMUS VARIEGATUS.

CHARACTER GENERICUS.

Rostrum longiusculum, obtusum; *nares* in medio rostri.

Rictus amplus.

Latera capitis subdenudata.

Cauda breviuscula.

Pedes tetradactyli, digito postico a terra elevato.

Lath. Ind. Orn. p. 633.

CHARACTER SPECIFICUS, &c.

TINAMUS fuscus, rufo variatus, verlice nigro, pectore ferrugineo.

TINAMUS rufo fusco nigricanteque fasciatus, subtus rufus, pileo nigro, gula abdomineque medio albis.

Lath. Ind. Orn. p. 634.

Tetrao variegatus. T. pedibus rostroque fuscis, capite et cervice nigris, corpore supra lineis transversis spadiceis et nigris vario, subtus rufo, gula medioque abdominis albis.

Lin. Syst. Nat. Gmel. p. 768.

Americam Australem, præcipue Cayanam incolit Tinamus variegatus, magnitudine quasi Tetraonis Perdices Linnæi.

THE
VARIEGATED TINAMOU.

GENERIC CHARACTER.

Bill longish, obtuse ; *nostrils* in the middle of the bill.

Gape wide.

Cheeks slightly bared.

Tail rather short.

Feet tetradactyle, with the hind-toe raised from the ground.

SPECIFIC CHARACTER, &c.

Brown TINAMOU, with rufous variegations, black crown, and ferruginous breast.

VARIEGATED TINAMOU.

Lath. Syn. 2. p. 726.

TINAMOU varié de Cayenne.

Pl. Enl. 828.

The variegated Tinamou is a native of South-America, and particularly of Cayenne. Its size is that of a common partridge.



CYPRÆA TESTUDINARIA.

CHARACTER GENERICUS.

Animal Limax.

Testa univalvis, involuta, subovata, obtusa, lævis.

Apertura utrinque effusa, linearis, utrinque dentata, longitudinalis.

CHARACTER SPECIFICUS, &c.

CYPRÆA subflavescens, nebulis maculisque fuscis atomis albis irroratis.

CYPRÆA testudinaria. C. testa obtusa subcylindrica, extremitatibus depressis.

Lin. Syst. Nat. p. 1173.

TESTUDINARIA.

Rumph. Mus. t. 38. f. c.

In suo genere Cypræa testudinaria est species facile maxima. In maribus Indicis nascitur, coloribus interdum nonnihil varians.

THE
TORTOISE-SHELL COWRY.

GENERIC CHARACTER.

Animal resembling a Limax or Slug.

Shell univalve, involute, obtuse, ovate.

Aperture linear, longitudinal, toothed on both sides.

SPECIFIC CHARACTER, &c.

Yellowish COWRY, with brown clouds and spots freckled with minute white specks.

The Great Indian COWRY.

Porcelaine a ecaille de tortue.

Knorr. Vergn. 4. pl. 27. f. 2.

The tortoise-shell Cowry is the largest of the genus *Cypræa*. It is a native of the Indian seas, and sometimes varies slightly in colour.

Callinectes sapidus



PHALÆNA IMPERATORIA.

CHARACTER GENERICUS.

Antennæ setaceæ, a basi ad apicem sensim attenuatæ.

Alæ (sedentis) sæpius deflexæ, (volatu nocturno.)

CHARACTER SPECIFICUS, &c.


PHALÆNA IMPERATORIA. P. (Bombyx) elinguis, alis reversis flavis fusco irroratis; omnibus macula subocellari strigae undata purpurascente.

Smith. Abbot. Ins. Amer. 2. p. 109.
t. 55.

PHALÆNA imperialis.

Fabr. entom. emend. 4. p. 435.
Drury Ins. 1. t. 9. f. 1, 2.

Iconem quam ostendit tabula suppeditavit nobis eximium opus de insectis quæ in America Septentrionali generantur ab Abboto conscriptum, et Smithii, societatis Linnæanæ præsidis, annotationibus illustratum. Larvam cum chrysalide primus repræsentasse videtur Abbotus. Depascitur larva folia platani Americanæ, et mense Septembri idoneum sibi sub humo nacta latibulum, in chrysalidem convertitur; unde Julio proxime sequente erumpit insectum plene conformatum. In Georgia et Carolina præcipue cernitur *Phalæna imperialis*.



THE
IMPERIAL MOTH.

GENERIC CHARACTER.

Antennæ setaceous, gradually lessening from base to tip.

Wings (when sitting) generally deflex, (flight nocturnal).

SPECIFIC CHARACTER, &c.

MOTH with yellow wings, freckled with brown; each marked by a subocellated spot and waved purplish stripe.

The Great Plane-Tree Moth.

Smith's Abbot. 2. p. 109. pl. 55.

It is from the highly elegant work of Mr. Abbot on North-American Insects, accompanied by the notes of Dr. Smith, president of the Linnæan Society, that the beautiful moth here represented is copied. Mr. Abbot seems to have been the first who has figured its caterpillar and chrysalis: the former feeds on the leaves of the Occidental or American Plane-Tree, and undergoes its change in the month of September, retiring for that purpose under ground; and gives birth to the complete animal in the month of July in the succeeding year. It is principally found in Georgia and Carolina.

MONOCULUS CYPRINACEUS.

CHARACTER GENERICUS.

Pedes natatorii.

Corpus crusta tectum.

Oculi, plerisque, approximati, testæ innati.

CHARACTER SPECIFICUS, &c.

MONOCULUS orbicularis, acetabulis duobus
pectoralibus, cauda biloba.


MONOCULUS Delphinus. M. pedibus octo.

Lin. Gmel. p. 3005.

Mull. entomotr. p. 123. No. 58.

Corporibus, ut plurimum, affigitur bella hæc species diversorum piscium; cyprinorum præcipue; quibus adhæret ope duorum brevium acetabulorum utrinque e thorace sitorum. Ostenditur in tabula tum vera magnitudo tum microscopice aucta. Primus eam descripsisse videtur Bakerus in opere cui titulus "Employment for the Microscope;" qui iconem addidit fidam satis, carentem tamen elegantia. Observavit Dominus Bakerus mori hanc speciem unius aut duorum dierum spatio, nisi in eadem aqua servetur una cum pisce quem morsu infestat. In hoc tamen eum errare ausim asserere; cum monoculos cyprinaceos diu valentes vixisse cognoverim sine visibili ullo nutrimento præter animalcula quæ suppeditarunt aquæ.

In lateribus vasis in quo servantur hæc insecta, deponi solent ovorum ovatorum, alborum, acervi horizontales, oblongi, singulo acervo ova quasi triginta seu quadraginta continente.



THE
CARP MONOCULUS.

GENERIC CHARACTER.

Feet formed for swimming.

Body covered by a crustaceous tegument.

Eyes, in most species, approximated, and imbedded in the shell.

SPECIFIC CHARACTER, &c.

Orbicular MONOCULUS, with two pectoral suckers, and bilobate tail.

Carp-Louse.

Baker Empl. for the Microscope, p. 374.
pl. 14. f. 14.

This elegant species is generally found attached to the bodies of different kinds of fish ; more especially to the carp ; adhering by means of two short, cylindric suckers or acetabula situated on each side the thorax. The plate shews the animal both in its natural size, and as it appears when magnified. It seems to have been first described by Mr. Baker in his work entitled "Employment for the Microscope," where a figure may be found, sufficiently exact to ascertain the species, though not very elegant. Mr. Baker observes that, unless this animal be kept in the same water with the fish it infests, it dies in the space of a day or

two; but this I have found to be a mistake; having preserved several in perfect vigour for a very considerable time, though without any visible nutriment, except the animalcules floating about in the water. The eggs of these insects are whitish, and are deposited in small, horizontal, oblong groupes of thirty or forty in each, on the sides of the glass vessel in which they are kept.

PSITTACUS COCCINEUS.

CHARACTER GENERICUS.

Rostrum aduncum, mandibula superiore mobili,
cera instructa.

Nares in rostri basi.

Lingua carnea, obtusa, integra.

Pedes scansorii.

Lin. Syst. Nat. p. 139.

CHARACTER SPECIFICUS, &c.

PSITTACUS macrourus coccineus, dorso antico, abdomine, fascia capitis, caudaque violaceis, alarum rectricibus fusco-cæruleo marginatis.

PSITTACUS Indicus. P. coccineus, fusco et violaceo variegatus, capite et collo superioribus, pectore et tænia pone oculos violaceis, rectricum majorum apice dilute fusco, minorum rectricibusque ex fusco violaceis.

Lin. Syst. Nat. Gmel. p. 318.

In insulis Indicis, præcipue in Amboina conspicitur
Psittacus coccineus, coloribus interdum leviter varians.
Magnitudo ejus est quasi Psittaci Domicellæ.



THE
INDIAN LORY.

GENERIC CHARACTER.

Bill hooked: upper mandible moveable.
Nostrils round, placed in the base of the bill.
Tongue fleshy, broad, blunt at the end.
Legs short; feet scansorial.

SPECIFIC CHARACTER, &c.

Long-tailed crimson LORY, with violet-blue
variegations; the wing-coverts with dusky-
blue margins.

INDIAN LORY.

Lath. syn. 1. p. 217.

Perruche des Indes orientales.

Pl. Enl. 143.

The splendid species here represented is found in the Indian isles, particularly in Amboyña, and sometimes is observed to vary a little in its colours. Its size is that of the *Psittacus Domicella* or purple-capped Lory.

SPONGIA PALMATA.

CHARACTER GENERICUS.

Animal? fixum, flexile, polymorphum, torpidissimum, contextum vel e fibris reticulatis, vel e spinulis gelatina viva vestitis.

Osculis seu foraminibus superficiei aquam respirantibus.

Soland. et Ellis Zooph. p. 182.

CHARACTER SPECIFICUS, &c.

SPONGIA flavo-ferruginea palmata, poris prominulis.

SPONGIA palmata. S. palmata, digitis apice subdivisis, poris prominulis inordinate dispositis.

Soland. et Ellis Zooph. p. 189. t. 58. f. 6.

Circa litora Britannica conspicitur Spongia palmata, ab Ellisio primum descripta. Tabula magnitudinem veram exprimit.



THE
PALMATED SPONGE.

GENERIC CHARACTER.

Animal? fixed, flexible, torpid, of various forms; composed either of reticulated fibres, or masses of small spines interwoven together; clothed with a gelatinous flesh full of small mouths or foramina on its surface, by which it sucks in and throws out the water.

SPECIFIC CHARACTER, &c.

Yellow-ferruginous, palmated SPONGE, with prominent pores.

PALMATED SPONGE.

Ellis Zooph. p. 189. pl. 58. f. 6.

The palmated Sponge is found about the British coasts, and seems to have been first described by Mr. Ellis. The plate represents it in its natural size.

SPHINX FICUS.

CHARACTER GENERICUS.

Antennæ medio crassiores, seu utraque extremitate attenuatæ, subprismaticæ.

Alæ deflexæ, (volatu graviore vespertino seu matutino.)

Lin. Syst. Nat. p. 796.

CHARACTER SPECIFICUS, &c.

SPHINX alis superioribus olivaceo-fuscis, apicibus pallidis, inferioribus fasciis duabus subnigris.

SPHINX FICUS. S. alis fusco-nebulosis; posticis angulo ani albis.

Lin. Syst. Nat. p. 800.

SPHINX alis fusco-nebulosis, posticis fascia nigra, anguloque ani albo.

Fab. Sp. Ins. 2. p. 145.

Surinamiam incolit Sphinx Ficus. Larva ejus folia Ficus præcipue depascitur, et in tabula nostra cum chrysalide et insecto perfecto magnitudine vera exprimitur.



A. Nodding Del. et Sculp.

THE
FIG-TREE SPHINX.

GENERIC CHARACTER.

Antennæ subprismatic, thickest in the middle,
and attenuated at each extremity.

Wings deflected. Flight generally matutine or
vespertine.

SPECIFIC CHARACTER, &c.

SPHINX with olive-brown upper wings with
pale tips; and pale under-wings with two
blackish bars.

The FIG-TREE SPHINX.

Merian Sur. Ins. pl. 33.

This species is a native of Surinam. The Caterpillar feeds chiefly on the leaves of the Fig-tree, and is represented on the annexed plate, together with the chrysalis and complete Insect, in its natural size.

THE
OF THE
IN THE
AND THE
AND THE
AND THE

THE
OF THE
IN THE
AND THE
AND THE
AND THE

THE
OF THE
IN THE
AND THE
AND THE
AND THE

THE
OF THE
IN THE
AND THE
AND THE
AND THE

THE
OF THE
IN THE
AND THE
AND THE
AND THE

THE
OF THE
IN THE
AND THE
AND THE
AND THE

THE
OF THE
IN THE
AND THE
AND THE
AND THE

THE
OF THE
IN THE
AND THE
AND THE
AND THE

ONCHIDIUM TYPHÆ.

CHARACTER GENERICUS.

Corpus limacinum, subverrucosum.

Caput brachiis duobus brevibus, totidemque tentaculis.

Os infra.



CHARACTER SPECIFICUS, &c.

ONCHIDIUM griseo-olivaceum, subtus flavescens margine nigro.

ONCHIDIUM TYPHÆ.

Buchanan in Trans. Lin. Soc.
vol. 5. p. 132.

Genus *Onchidium*, cui cum genere *Limacis* arcta videtur esse cognatio, instituit Buchananus; qui in quinto volumine actorum societatis Linnæanæ speciem descripsit de qua jam agitur, Bengalæ incolam, et in specie quadam Typhæ, Typhæ scilicet *elephantinæ* præcipue repertam. Magnitudo eadem fere est atque atri limacis communis: color supra leviter olivaceus infra flavo-pallescit.



18. 19. 20. 21. 22.

THE
LIMACINE ONCHIDIUM.

GENERIC CHARACTER.

Body snail-shaped, slightly warted.

Head with two short *lateral arms*, and two *tentacula*.

Mouth beneath.

SPECIFIC CHARACTER, &c.

Greyish-olive ONCHIDIUM, yellowish beneath with black margin.

ONCHIDIUM.

Trans. Lin. Soc. vol. 5. p. 132.

The genus *Onchidium*, extremely nearly allied to that of *Limax*, appears to have been instituted by Dr. Buchanan, who, in the fifth volume of the Transactions of the Linnæan Society, has described the present species, which is a native of Bengal, and is chiefly found on a large species of *Typha*, called *Typha elephantina*. Its size is that of the common black slug, and its colour pale olive above, inclining to yellow beneath.

FALCO COMMUNIS.

CHARACTER GENERICUS.

Rostrum aduncum, basi cera instructum.

Caput pennis arcte tectum.

Lingua bifida.

CHARACTER SPECIFICUS, &c.

FALCO fuscus, rufo undulatus, cauda subnigro fasciata, pectore abdomineque albidis fusco maculatis.

FALCO COMMUNIS. F. rostro cærulescente, cera iridibus pedibusque luteis, corpore fusco, pennarum margine rufo, rectricibus fasciis saturatioribus.

Lath. Ind. Orn. p. 30.

Falconem hunc ad id docilitatis perducere solent aucupes, ut inita cum iis societate, dato signo, in perdices et aves majores volantes irruat, comprehensasque unguibus retinens ad dominos deportet. Exuperat paulum magnitudine falconem *palumbarium* Linnæi, longus nempe, ut plurimum, pedem cum dimidio. Color avis superior infuscatur, marginibus pennarum ferrugineis, caudaque fasciis nigricantibus transversis decorata; inferior albet, non sine maculis fuscis huc illuc dispersis. Notissimum tamen est variare situm et dispositionem colorum. In omni fere Europa reperitur Falco communis.



THE
COMMON FALCON.

GENERIC CHARACTER.

Bill hooked, furnished at the base with a cere.

Head closely covered with feathers.

Tongue bifid.

SPECIFIC CHARACTER, &c.

Brown FALCON with rufous undulations, the tail marked with dusky bars; the breast and belly whitish with dusky spots.

Le FAUCON.

Briss. ornith. 1. p. 321.

This bird, so much esteemed in the diversion of Falconry, in which it is usually flown at partridges and other kind of game, is somewhat larger than the *Falco palumbarius* or Goshawk, measuring about eighteen inches in length. The general colour of its plumage is brown above, with rufous margins, the tail being crossed by several dusky bars: the under parts of the bird are white, with numerous dusky spots. It is known however to vary extremely in the particular distribution of its colours. The Falcon is a native of most parts of the European continent.

that of most birds of the same family
some specimens of the same species
it is known however that the
birds of the same family differ in color
the same species in different parts of the
country is known to have different
colours in different parts of the country
and in different parts of the same
country. It is known that the
birds of the same family differ in color
the same species in different parts of the
country is known to have different
colours in different parts of the country
and in different parts of the same
country.

THE BIRDS

and the same species in different
parts of the country is known to have
different colours in different parts of the
country and in different parts of the same
country. It is known that the
birds of the same family differ in color
the same species in different parts of the
country is known to have different
colours in different parts of the country
and in different parts of the same
country.

THE BIRDS

and the same species in different
parts of the country is known to have
different colours in different parts of the
country and in different parts of the same
country.

CALIGUS PISCINUS.

CHARACTER GENERICUS.

Corpus suborbiculare.

Pedes sex vel decem, posterioribus branchiatis.

Cauda appendiculata.

CHARACTER SPECIFICUS, &c.

CALIGUS corpore brevi, pedibus decem, cauda bifida.

Oniscus lutosus.

Slabb. Micr. t. 16. f. 1. 2. ?

CALIGUS curtus.


Mull. entomostr. p. 180. No. 63.

Monoculus piscinus.

Fabr. syst. ent. 2. p. 189. ?

Infestat hocce animalculum varios pisces eodem modo quo solet monoculi species quam in opere hoc nostro antea descripsimus, marinos præcipue, salmones sæpius, interdum et flesos; celeriter nempe huc illuc super corpora discurrendo, et ad libitum se cuti affigendo. Si certorum quorundum signorum habeatur ratio, ad genus *Monoculi* inter insecta videtur accedere; sunt tamen alia, ob quæ ad genus *Lernææ* inter vermes referendum putes. Exprimatur in tabula tam vera magnitudo quam microscopice aucta.

772



R. W. Paul

THE
FISH CALIGUS.

GENERIC CHARACTER.

Body suborbicular.

Legs six or ten, the lower branchiated.

Tail appendiculated.

SPECIFIC CHARACTER, &c.

CALIGUS with short body, ten legs, and bifid tail.

Calige des poisons.

Latreille ins. et crustac. 4. p. 111.

This animal, like the species of *Monoculus* described in a preceding number of the present work, infests several sorts of fishes, but principally the marine ones. It is found more particularly on the Salmon, and sometimes on the Flounder, wandering about with celerity over various parts of the fish, and occasionally fixing itself tenaciously to the skin. It seems allied in some points to the genus *Monoculus* among Insects, and in others to the genus *Lernæa* among the *Vermes*. It is represented in the plate both in its natural size and magnified.

GRYLLUS VIRIDISSIMUS.

CHARACTER GENERICUS.

Caput inflexum, maxillosum, palpis instructum.

Antennæ setaceæ seu filiformes.

Alæ quatuor, deflexæ, convolutæ: inferiores plicatæ.

Pedes postici saltatorii: ungues ubique bini.

Lin. Syst. Nat. p. 692.

CHARACTER SPECIFICUS, &c.

GRYLLUS viridis, antennis setaceis longissimis, linea thoracica subrufa.


GRYLLUS VIRIDISSIMUS. G. thorace rotundato, alis viridibus immaculatis, antennis setaceis longissimis.


Lin. Syst. Nat. p. 698.

Locusta viridis cantatrix.

Degeer ins. 3. p. 428.

Gryllorum Britannicorum maximus et elegantissimus est qui in tabula depingitur, exeunte æstate non raro peragraræ solitus prata et sepes. Foemina mense autumnæ ova paululum sub terra occulit, e quibus vere insequente erumpunt pulli; iiq; certis temporis intervallis cum exuvias sæpius deposuerint, alis tandem instruuntur, et perfecti evadunt Grylli.





THE
GREEN LOCUST.

GENERIC CHARACTER.

Head inflected, armed with jaws, and furnished with feelers.

Antennæ (in most species) setaceous or filiform.

Wings four, deflected and convoluted; the lower ones pleated.

Hind-Legs formed for leaping: claws on all the feet double.

SPECIFIC CHARACTER, &c.

GREEN LOCUST, with very long setaceous antennæ, and a subferruginous line down the thorax.

The common GREEN LOCUST.

The *Gryllus viridissimus* or Great green Locust, the largest and most elegant of the British species, is not very uncommon in fields and hedges towards the decline of summer. In autumn the female deposits her eggs beneath the surface of the ground, and from these, in the following spring, are hatched the young, which, after casting their skin several times, during the different periods of their growth, at length acquire their wings, and become complete.

PAPILIO ANTIOPA.

CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatæ.

Alæ (sedentis) erectæ sursumque conniventes (volatu diurno.)

Lin. Syst. Nat. p. 744.

CHARACTER SPECIFICUS, &c.

PAPILIO alis angulatis purpureo-nigricantibus limbo albo-flavescente.


PAPILIO ANTIOPA. P. alis angulatis nigris limbo albido.

Lin. Syst. Nat. p. 776.

Roes. ins. belust. 1. *pap. diurn.* t. 1.

Inter insecta Britannica rariora numeratur Papilio Antiopa, cujus larva folia Salicis præcipue depascitur, et mense Julio in chrysalidem convertitur, e qua mense Augusto erumpit papilio.

Antropoda



ANTIOPA.

GENERIC CHARACTER.

Antennæ thickening towards the end, and commonly terminating in a clavated tip.

Wings (when at rest) meeting upwards. Flight diurnal.

SPECIFIC CHARACTER, &c.

Butterfly with purplish-black angular wings, with whitish yellow border.

The black willow Butterfly.

The yellow-edged black Butterfly.

The *Papilio Antiopa* is numbered among the rarer insects of Great Britain. The Caterpillar feeds principally on the leaves of the willow, and changes into chrysalis in the month of July, from which in the month of August emerges the complete insect.

month of August several the country was
suffering in the month of July, most of the
cattle on the farms of the country were
killed or driven to the city. The country was
The people of the country were very poor.

The people of the country were very poor.

The people of the country were very poor.

The people of the country were very poor.

The people of the country were very poor.

The people of the country were very poor.

The people of the country were very poor.

The people of the country were very poor.

The people of the country were very poor.

The people of the country were very poor.

The people of the country were very poor.

The people of the country were very poor.

SCOLOPAX HELIAS.

CHARACTER GENERICUS.

Rostrum teretiusculum, obtusum, capite longius.

Nares lineares.

Facies tecta.

Pedes tetradactyli, postico pluribus articulis insistente.

Lin. Syst. Nat. p. 242.

CHARACTER SPECIFICUS, &c.

SCOLOPAX rufa, albo nigroque transversim fasciata, abdomine albo, cauda fascia nigra.


SCOLOPAX HELIAS. S. corpore nigro fulvo alboque fasciato, abdomine strigisque capitis albis, collo rufo lineis transversis nigris, cauda nebulosa fascia nigra.

Lath. Ind. Orn. p. 725.

ARDEA Helias. A. capite lævi, corpore supra nigro lineis transversis ferrugineis undulato, subtus albido, remigum rectricumque fasciis ferrugineis.

Pall. Nord. beytr. Lin. Gmel. p. 640.

In elegantissime variatis Scolopacini generis habenda est species de qua jam agitur, quæque Scolopacem *rusticolam* Linnæi fere æquat magnitudine. Ali creditur eodem fere modo quo reliquum genus, loca nempe aquosa frequentando, vermesque et insecta exinde extrahendo. Nascitur in calidioribus regionibus Americæ Australis.





APN. Del. 1.

THE
CAURALE SNIPE.

GENERIC CHARACTER.

Bill long, slender.

Nostrils linear, situated in a furrow.

Face covered with feathers.

Feet tetradactylous; back toe jointed.

SPECIFIC CHARACTER, &c.

Rufous SNIPE, with black and white transverse
bands, and a black band across the tail.

The CAURALE SNIPE.

Lath. syn. 3. p. 156.

CAURALE, ou petit Paon des Roses.

Buff. ois. 8. p. 169. pl. 14.

Le CAURAL de Cayenne.

Pl. Enl. 782.

This bird may be considered as the most elegantly variegated species of the genus *Scolopax* hitherto discovered. Its size is nearly that of a Woodcock. In its habits it is supposed to resemble the rest of its tribe, frequenting watery places, and feeding on worms, insects, &c. &c. It is a native of the hotter parts of South America.

MULLUS MACULATUS.

CHARACTER GENERICUS.

Caput compressum, declive, squamosum.

Os cirratum.

Membr. branch. triradiata.

Corpus squamis magnis subdeciduis.

CHARACTER SPECIFICUS, &c.

MULLUS ruber, corpore utrinque maculis tribus rotundatis nigris.

MULLUS MACULATUS. M. maculis tribus nigris ad lineam lateralem.

Bloch. ichth. 10. p. 79. t. 348. f. 1.

PIRAMETARA Brasiliensibus.

Marcgr. Bras. p. 156.

MULLUS Surmuletus. var. β .

Lin. Syst. Gmel. p. 1339.

Maria incolit Americana Mullus maculatus, in pedalem vel sesquipedalem non raro crescens longitudinem.

quoniam

quoniam vel adhibetur non ideo carere possit

Mus musculus Americanus. Mus musculus. 10. 12.

1711. 1712. 1713. 1714.

MUS musculus. 1711. 1712.

1713. 1714. 1715.

1716. 1717. 1718.

1719. 1720. 1721. 1722.

1723. 1724. 1725.

MUS musculus. 1726. 1727.

1728. 1729.

MUS musculus. 1730. 1731.

1732. 1733.

1734. 1735.

1736. 1737.


1740.


1741. 1742. 1743.

1744. 1745.

1746. 1747.

MUS musculus





THE
SPOTTED SURMULLET.

GENERIC CHARACTER.

Head compressed, scaly.

Mouth bearded.

Gill-Membrane three rayed.

Body covered with large subdeciduous scales.

SPECIFIC CHARACTER, &c.

Red SURMULLET, with the body marked on
each side by three rounded black spots.

Der gefleckte Rothbart.

Bloch ichth. p. 348. f. 1.

The spotted Surmullet is a native of the American seas, where it often grows to the length of twelve or eighteen inches.

GRYLLUS MONSTROSUS.

CHARACTER GENERICUS.

Caput inflexum, maxillosum, palpis instructum.

Antennæ setaceæ seu filiformes.

Alæ quatuor, deflexæ, convolutæ: inferiores plicatæ.

Pedes postici saltatorii: ungues ubique bini.

Lin. Syst. Nat. p. 692.

CHARACTER SPECIFICUS, &c.

GRYLLUS subfuscus, alis apice contortuplicatis, pedibus utrinque appendiculatis.

GRYLLUS MONSTROSUS. G. elytris alisque caudato-convolutis.

Lin. Syst. Nat. Gmel. p. 2059.

ACHETA monstrosa.

Fabr. Spec. ins. 1. p. 353.

Drury ins. 2. pl. 43.

E plurimis congeneribus insectis vix aliud insignius reperiri possit Gryllo monstroso, qui in India generatus, et Gryllo Europæo qui *Gryllotalpa* dicitur, contribulis, eodem fere, (ut a forma conjicere licet,) utitur vivendi modo. In tabula depingitur alis clausis, nec non alis plene expansis.



THE
SPIRAL-WINGED LOCUST.

GENERIC CHARACTER.

Head inflected, armed with jaws, and furnished with feelers.

Antennæ (in most species) setaceous, or filiform.

Wings four, deflected and convoluted; the lower ones pleated.

Hind-Legs formed for leaping: claws on all the feet double.

SPECIFIC CHARACTER, &c.

Brownish LOCUST, with the wings twisted at the tips, and the feet appendiculated on each side.

The SPIRAL-WINGED Indian LOCUST.

Among the numerous insects of the genus *Gryllus*, hardly any can be adduced of an appearance more remarkable than the present species, which is a native of India, and belongs to the same tribe with the *Gryllus Gryllotalpa* or Mole-Cricket, to which in its habits of life it is probably similar. The plate represents it both in its general state, and as it appears when the wings are fully expanded.

ALCYONIUM SCHLOSSERI.

CHARACTER GENERICUS.

Animal plantæ forma crescens.

Stirps fixa, carnosæ, gelatinosa, spongiosa, vel coriacea.

Superficies cellulosa, poris seu osculis stellatis pertusa, hydras tentaculatas oviparas exserentibus.

CHARACTER SPECIFICUS, &c.

ALCYONIUM planum subcoriaceum subcæruleum, stellis flavis.

ALCYONIUM carnosum asteriscis obtusis ornatum.

Ellis et Schlosser Act. Angl. 49. p. 451.

ALCYONIUM SCHLOSSERI. A. carnosum lividum, asteriscis luteis radiis obtusis ornatum.

Soland. et Ellis Zooph. p. 177.

Quod in tabula depinximus zoophyton, cuique, cum species sit valde anomala *Alcyonii*, genus distinctum et proprium non immerito possit institui, primus accurate descripsisse et depinxisse videtur Albertus Schlosserus in actis Anglicis philosophicis. Apud oras Britannicas reperitur, fortasse etiam apud oras plerasque Europæas, ut plurimum vel fucorum majorum caulibus circumvolutum, vel super rupes et alia marina frustulatim et multiformiter dispersum.



THE
SCHLOSSERIAN ALCYONIUM.

GENERIC CHARACTER.

Animal growing in the form of a plant.

Body fixed; fleshy, gelatinous, spongy, or coriaceous.

Surface cellular with star-shaped openings, from which are extended polype-shaped oviparous heads.

SPECIFIC CHARACTER, &c.

Flat subcoriaceous blueish ALCYONIUM, with yellow stars.

SCHLOSSER'S ALCYONIUM.

Solander & Ellis's Zoophytes. p. 477.

The curious zoophyte represented on the present plate seems to have been first distinctly described, and figured with accuracy, by Dr. Albert Schlosser in the Philosophical Transactions. It is a native of the British and probably of most other European coasts, and is commonly found either investing the stems of the larger fuci, or spread in irregular patches of various size on rocks, &c. It must be considered as a very anomalous species of Alcyonium, and might not improperly constitute a distinct genus.

FRINGILLA TRISTIS.

CHARACTER GENERICUS.

Rostrum conicum, rectum, acuminatum.

Lin. Syst. Nat. p. 317.

CHARACTER SPECIFICUS, &c.

FRINGILLA flava, fronte alis caudaque nigris.

FRINGILLA tristis. F. flava, fronte nigra,
alis fuscis.

Lin. Syst. Nat. p. 320.

Carduelis Americana.

Catesb. Carol. 1. t. 43.

Depingitur in tabula magnitudo vera Fringillæ
tristis, varias Americæ septentrionalis regiones inco-
lantis.



app. 1827. Del. et Sculp.

THE
GOLDEN FINCH.

GENERIC CHARACTER.

Bill conic, slender towards the end, and sharp-pointed.

SPECIFIC CHARACTER, &c.

Yellow FINCH, with black front, wings, and tail.

The American GOLDFINCH.

Catesb. Carol. 1. pl. 43.

The GOLDEN FINCH.

Penn. Arch. Zool. 2. p. 371.

The *Fringilla tristis* is a native of several parts of North America, and is represented on the plate in its natural size.

THE

THE

THE

THE

THE

THE

THE

THE

THE

THE

THE

SPARUS PAGRUS.

CHARACTER GENERICUS.

Opercula squamosa, mutica.

Dentes validi, Molares (plerisque) convexi, læves, in series dispositi.

CHARACTER SPECIFICUS, &c.

SPARUS roseus, maculis duabus nigris utrinque ad opercula.

Gen. Zool. 5. p. 408.

SPARUS PAGRUS. S. rubescens, cute ad radicem pinnarum dorsi et ani in sinum producta.

Lin. Syst. Nat. p. 469.

Sparus radio ultimo pinnæ dorsi anique cute incluso.

Bloch icht. 8. p. 50. t. 267.

Alitur Sparus Pagrus in mari Mediterraneo, Septentrionali, et Atlantico. Intrat etiam interdum majores fluvios. In eam crescit magnitudinem, ut non raro pondo sit octo vel decem. In cibis habetur lautioribus. Hoc habet commune cum aliis multis piscibus, quod colore nonnihil variet: ut plurimum tamen rubet.

SPARUS PAGRUS

CHARACTER GENERICUS

Opercula squamulosa, mutica.
Dentes validi, Molares (pleurales) convexi,
hæves, in series dispositi.

CHARACTER SPECIFICUS, &c.

SPARUS roseus, maculis duabus nigris utrinque
ad opercula.

Gen. Zool. 5. p. 408.


SPARUS PAGRUS. S. rubescens, cute ad
radicem pinnae dorsi et ani in sinum
producta.

Linn. Syst. Nat. p. 469.

Sparus radio ultimo pinnae dorsi aequæ cute
incluso.

Bloch icht. 8. p. 50. t. 267.

Aliter Sparus Pagrus in mari Mediterraneo, Septen-
trionali, et Atlantico. Latet etiam interdum majores
fluvios. In eam crescit magnitudinem, ut non raro
pondo sit octo vel decem. In cibus habetur lautioribus.
Hoc habet commune cum aliis multis piscibus, quod
colore nonnulli variat: ut plurimum tamen rubet.



750

Thalassoma daniellii

THE
PAGRE SPARUS.

GENERIC CHARACTER.

Gill-Covers scaly, unarmed.

Teeth strong ; in most species convex, smooth,
and disposed in rows.

SPECIFIC CHARACTER, &c.

Rose-red SPARUS, with two black spots on
each side near the gill-covers.

The Red Gilthead.

Penn. Br. Zool. 3. p. 212.

PAGRE.

Rondel. Hist. Poiss. 1. p. 127.

PAGRE SPARUS.

Gen. Zool. 6. p. 408.

The Pagre Sparus is an inhabitant of the Mediterranean, Atlantic, and Northern seas, and sometimes enters the larger rivers. It grows to a considerable size, often weighing eight or ten pounds, and is considered as a delicate fish for the table. In colour, like many other fishes, it occasionally varies, but is generally red.

SCARABÆUS GOLIATHUS.

Var.

CHARACTER GENERICUS.

Antennæ clavatæ, capitulo fissili.

Tibiæ sæpius dentatæ.

Corpus crassum, compactum.

Gen. Zool. 6. p. 19.

CHARACTER SPECIFICUS, &c.

Scarabæus scutellatus, thorace inermi, clypeo
bifurco.

Fabr. sp. ins. 1. p. 14.

SCARABÆUS GOLIATHUS. S. scutellatus,
thorace inermi, capite rostro bifurco.

Lin. Mantiss. 530.

Cetonia Goliathus.

Oliv. ins. 1. fig. 33.

Scarabæi Goliathi in opere hoc nostro antea depicti
varietatem repræsentavimus a priore illo coloribus tan-
tummodo discrepantem. Monstratur in tabula insec-
tum quasi tertia parte imminutum. Generatur in
variis partibus Africanis.

SCARABAEUS GOLIATHUS.

Var.

CHARACTER GENERICUS.

Antennae clavatae, capitulo fissili.

Tibiae serpius dentatae.

Corpus crassum, compactum.

Gen. Zool. 6. p. 19.

CHARACTER SPECIFICUS, &c.

Scutellum scutellatum, thorace normali, clypeo
bifurco.

Fabr. Sp. ian. 1. p. 14.

SCARABAEUS GOLIATHUS, S. scutellatus.

Thorace normali, capite rostro bifurco.

Linn. Mantiss. 530.

Ceteris Goliathus.

Oliv. ian. 1. fig. 33.

Scutellum Goliathus in opere hoc nunc primum descriptum
varietatem repentinam a priore his coloribus ten-
tando descripsit. Monstrum in tabula insect-
um quodammodo parum variatum. Generum in
vniuersis affinis.



THE
GOLIATH BEETLE.

GENERIC CHARACTER.

Antennæ or Horns clavate, with a fissile tip.

Legs generally toothed.

Body thick and compact.

Gen. Zool. 6. p. 19.

SPECIFIC CHARACTER, &c.

Scutellated BEETLE, with unarmed thorax
and forked head.

Fork-Headed BEETLE.

Nat. Miscell. 3. p. 80.

Goliathus.

Drury ins. 1. pl. 31. & 3. pl. 40.

The Scarabæus Goliathus has before been figured in the present work. The variety here represented differs only in colour from the preceding. It is a native of several parts of Africa, and is reduced on the plate to about a third smaller than its natural size.

SPHINX CONVOLVULI.

CHARACTER GENERICUS.

Antennæ medio crassiores, seu utraque extremitate attennatæ, subprismaticæ.

Alæ deflexæ, (volatu quaviore vespertino seu matutino.)

Lin. Syst. Nat. p. 796.

CHARACTER SPECIFICUS, &c.

Sphinx alis integris canis fusco nebulosis, inferioribus nigro fasciatis, abdomine roseo cingulis nigris.

SPHINX CONVOLVULI. S. alis integris: posticis nigro fasciatis margine postico albo punctatis, abdomine rubro cingulis atris.

Lin. Syst. Nat. p. 798.

Merian. ins. Europ. t. 75. f. 2.

Roesel. phal. 1. t. 7.

Arcta admodum affinitate conjungitur formosa hæc species cum *Sphinge ligustri* Linnæi; qua tamen, (in Anglia saltem) multo est rarior. Larva *convolvuli sepii* Linnæani folia præcipue depascitur, et mense Junio sub humo occulta in chrysalidem convertitur, e qua mense Septembri erumpere solet ipsa Sphinx.

SPHINX CONVULSUS

CHARACTERES SPECIFICI

CHARACTERES SPECIFICI

Alae medio crassiores, seu unguis exteriori-
late attenuatae, subtransversae.
Ala dextra, (veluti parvioris vestigio, seu
marginis.)
Lin. Sp. Nat. p. 796.

CHARACTERES SPECIFICI

Sphinx alis integris canis fusco rubris, infu-
rioribus nigro fasciatis, abdomine roseo
cinctis nigra.
SPHINX CONVULSUS. 2. alis integris
posterioribus nigro fasciatis marginibus postico
albo punctatis, abdomine rubro cinctis
albis.

Lin. Sp. Nat. p. 796.

Musca in Europ. p. 79. t. 2.

Reuss. Ital. 1. t. 1.

Alis admodum albis, concoloribus, formae pec-
specie cum Spang. figur. Linnaei, qui tamen, in
Anglia saltem, multo est rarior. Larva nectaris, vel
Jannetii, fons principis depascitur, et tamen Jannetii
sub humo occulta in chrysalidem convertitur, e qua
tamen Septembris erumpit, sed ipse Sphinx.



Adiantum *Lucy*

THE
BINDWEED SPHINX.

GENERIC CHARACTER.

Antennæ subprismatic, thickest in the middle,
and attenuated at each extremity.

Wings deflected. Flight generally matutine or
vespertine.

SPECIFIC CHARACTER, &c.

SPHINX with even, grey wings, clouded with
brown; the lower pair barred with black;
the abdomen rose-coloured with black
bands.

The BINDWEED SPHINX, or *Convolvulus*
Hawk-Moth.

This beautiful species is extremely nearly allied to
the *Sphinx ligustri* or Privet-Sphinx, but is (at least
in England) a much rarer insect. The caterpillar feeds
principally on the leaves of the common bindweed,
and, retiring under ground in June, changes to a
chrysalis, out of which in September emerges the com-
plete insect.

THE

BINDWEED SPHINX

.....

.....

Antennae slightly thickened at the middle
and the joint at the base of the antenna
is very distinct. The body is entirely
black.

.....

SPHINX with very long wings, rounded at
brown; the lower part of the wings black;
the abdomen rose colored with black
bands.

The BINDWEED SPHINX, or Copestolus
Hawk-Moth.

This beautiful species is extremely nearly allied to
the Sphinx ligata or Twisted-Sphinx, but is (at least
in England) a much rarer insect. The female is
principally of the color of the common bindweed,
and, retaining under ground in June, changes to a
chrysalis, one of which is deposited in the com-
mon insect.

PICUS RUFUS.

CHARACTER GENERICUS.

Rostrum polyedrum, rectum, apice cuneato.
Nares pennis setaceis recumbentibus obteetæ.
Lingua teres, lumbriciformis, longissima, mucronata, apice retrorsum aculeata setis.
Pedes scansorii.

CHARACTER SPECIFICUS, &c.

Picus flavo-rufescens, undulis transversis nigris, alis caudaque saturatoribus.

PICUS RUFUS. *P. rufus* nigro-undulatus, alis cauda corporeque subtus saturatoribus.

Lath. ind. orn. p. 241.

PICUS RUFUS. *P. rufus* nigro undulatus.

Lin. Syst. Gmel. 1. p. 432.

Pulcherrimam pici speciem Americæ australis incolam juxta veram magnitudinem repræsentavimus. More congenerum prædatur insecta, et sedem sibi quærit in cavis arborum.

PICUS RUFUS.

CHARACTER GENERICUS.

Rostri polyedri, rectum, apice cuneato.
 Vires pennae setaeque recurvatis obtectae.
 Lingua teres, hamuliformis, longissima, mu-
 cronata, apice retorsum aculeata setis.
 Pedes scanonii.

CHARACTER SPECIFICUS, &c.


Picus flavo-fulvus, undula transversa nigra,
 alis caudaeque saturatioribus.
 PICUS RUFUS. P. rufus nigro-undulatus,
 alis caudae corporeque subius saturatio-
 ribus.

Lat. ind. orn. p. 241.

PICUS RUFUS. P. rufus nigro-undulatus.

Lin. Syst. Genl. 1. p. 482.

Polhemianum pici speciem Americae australis inco-
 tam juxta veterum magnitudinem representativum.
 Motu cognatum praedat insecta, et sedem sibi
 quaerit in cavis arborum.





THE
RUFIOUS WOODPECKER.

GENERIC CHARACTER.

Bill polyedral, strait, wedge-shaped towards the tip.

Nostrils covered by recumbent setaceous feathers.

Tongue worm-shaped, very long, sharp-pointed, and edged towards the tip with reversed bristles.

Feet scansorial.

SPECIFIC CHARACTER, &c.

Yellowish-rufous Woodpecker, transversely undulated with black; the wings and tail deeper than the rest.

Rufous Woodpecker.

Lath. syn. 1. p. 594.

Pic roux.

Buff. ois. 7. p. 36.

Pl. Enl. 694. f. 1.

The present highly elegant species of Woodpecker is a native of South America, and is represented on the plate in its natural size. Like the rest of its tribe it feeds on insects, and inhabits the hollows of trees.

RUFIOUS WOODPECKER.

GENERIC CHARACTER.

Bill polydactyl, straight, wedge-shaped towards the tip.
Nose covered by recumbent setaceous feathers.
Tongue worm-shaped, very long, sharp-pointed, and edged towards the tip with reversed bristles.
Feet scissorial.

SPECIFIC CHARACTER.

Yellowish-rufous Woodpecker, transversely undulated with black, the wings and tail deeper than the rest.

Rufous Woodpecker.

Length, 10.5. 504.

Buff, 10.5. 504.

10.5. 504.

The present highly elegant species of Woodpecker is a native of South America, and is represented on the plate in its natural size. Like the rest of its tribe it feeds on insects, and inhabits the hollows of trees.

MADREPORA SERIATA.

CHARACTER GENERICUS.

Animal Medusa.

Corallium cavitatibus lamelloso-flexuosis.

Lin. Syst. Nat. p. 1272.

CHARACTER SPECIFICUS, &c.

MADREPORA SERIATA. M. ramulosa,
ramis attenuatis acuminatis, stellis longi-
tudinaliter seriatis; margine superiore por-
recto, fornicato, ciliato.

Soland et Ellis Zooph. p. 171.
t. 31. f. 1. 2.

MADREPORA SERIATA.

Pall. Zooph. 336.

In maribus Indicis generatur Madrepora seriata,
crescitque interdum in altitudinem quindecim vel
etiam octodecim unciarum. Color communiter albus
est, substantia dura seu quasi lapidosa.

MADREPORA SERIATA.

CHARACTER GENERIS.

Animal Medusa.

Corallium cavatibus lamelloso-flexuosis.

Lam. S. M. Nat. p. 1272.

CHARACTER SPECIES, &c.

MADREPORA SERIATA. M. lamellosa.

Tam. attenuatis acuminatis, stellis longis.

tubusculis seriatis; margine superiore por-

recto, fornicato; ciliato.

Schm. et Ellis Zool. p. 171.

L. 31. t. 1. 2.

MADREPORA SERIATA.

Pall. Zool. 320.

In maribus laticis generis Madrepore
crescitur interdum in altitudinem pedum vel
etiam octoedrum unciarum. Color communiter albus
est, substantia dura seu quasi lapidea.



Scleractinia *Sp.*

THE
SERIATED MADREPORE.

GENERIC CHARACTER.

Animal allied to a Medusa.

Coral marked with lamellar flexuous or star-shaped cavities.

SPECIFIC CHARACTER, &c.

Madrepore with attenuated, pointed branches; the stars disposed in rows; the superior edge of each star projecting, arched, and ciliated.

Slender-branched Madrepore.

Madrepora seriata.

Soland. et Ellis Zooph. p. 31.

The *Madrepora seriata* is a native of the Indian seas, and occasionally arrives at the height of fifteen or eighteen inches: its general colour is white, and substance hard or strong.

SERIALIZED MADREPORE

GENERIC CHARACTER

Animal allied to a Medusa.

Coel marked with lamellar flexuous or star-

shaped cavities.

Sparsely branched, sometimes subglo-

Madrepore with attenuated, pointed branches;

the stars disposed in rows; the superior

edge of each star projecting, notched, and

crinated.

Slender-branched Madrepore.

Madrepore setina.

Small, slender, and sometimes subglo-

ular in form, with a few small, pointed

branches. The Madrepore setina is a

small, slender, and sometimes subglo-

ular in form, with a few small, pointed

branches.

SPARUS FASCIATUS.

CHARACTER GENERICUS.

Opercula squamosa, mutica.

Dentes validi, Molares (plerisque) convexi,
læves, in series dispositi.

CHARACTER SPECIFICUS, &c.

Sparus elongato-subquadratus albidus, subnigro
transverse fasciatus, pinnis nigro margi-
natis.

SPARUS FASCIATUS. S. corpore fasciato,
linea laterali interrupta.

Bloch. ichth. 8. p. 15. t. 257.

Insignit Sparum fasciatum squamarum magnitudo,
nec non colorum æqualis et concinna dispositio. Ali-
tur in maribus Indicis et Americanis, et interdum fere
bipedali est longitudine. Caro in cibis habetur lautio-
ribus.

SPARUS FASCIATUS.

CHARACTER GENERICUS.

Opercula squamosa, trunca.
Dentes validi, Molares (plerique) convexi,
Javes, in series dispositi.


CHARACTER SPECIFICUS.

Sparus elongato-subquadratus albidus, subnigro
transverse fasciatus, pinna nigro margi-
nata.

SPARUS FASCIATUS. S. corpore fasciato,
linea laterali interrupta.

Bloch. Nat. A. 8. p. 16. t. 287.

Insignis Sparum fasciatum squamarum magnitudo,
nec non colorum squarum et corporis dispositio. Ali-
ter in manibus Indis et Americanis, et interdum fere
bipedali est longitudo. Caro in cibus habetur latio-
ribus.



Thalassoma dactylopterus

THE
FASCIATED SPARUS.

GENERIC CHARACTER.

Gill-covers scaly, unarmed.

Teeth strong ; in most species convex, smooth,
and disposed in rows.

SPECIFIC CHARACTER, &c.

Lengthened subquadrate Sparus, with transverse dusky bands, and fins edged with black.

Gen. Zool. 4. p. 412.

This species, so remarkable for the large size of its scales and the regular distribution of its colours, is an inhabitant of the Indian and American seas, where it grows to the length of nearly two feet. It is said to be in good esteem as a table fish.

PAPILIO PAPHIA.

CHARACTER GENERICUS.

Antennae apicein versus crassiores, apicibus clava-
to-capitatis.
Alae (sedentis) eleganter sursumque connatas
(volandi diurno).

CHARACTER SPECIFICUS, &c.

PAPILIO PAPHIA. P. alis dentatis luteis
nigro-maculatis: subtus lineis argenteis
transversis.


Lin. Syst. Nat. p. 785.

Papilio alis dentatis foliis nigro-maculatis;
subtus fasciis argenteis.

Lin. Syst. Nat. p. 2332.

Nymph. phalaena.

In pulcherrimis lepidopterorum Britannicarum mu-
seo haberi possit Papilio Paphia, prope silvas, ut plu-
rimum, visendus. Larva utique depascitur, et mense
Junio in chrysalidem convertitur, unde insequente
Julio erumpit papilio.



756



PAPHIA.

GENERIC CHARACTER.

Antennæ thickening towards the end, and commonly terminating in a clavated tip.

Wings (when at rest) meeting upwards. Flight diurnal.

SPECIFIC CHARACTER, &c.

Butterfly with denticulated fulvous wings spotted with black, and marked beneath with silvery bands.

The Great Wood Fritillary.

The Silver-banded Fritillary.

The *Papilio Paphia* may be considered as one of the most elegant of the British lepidoptera. It is principally seen in the neighbourhood of woods, where its larva feeds on nettles, changing into chrysalis in the month of June, and the fly appearing in that of July.

PAPHIA.

GENERIC CHARACTER.

Wings (when at rest) meeting towards the end, and commonly terminating in a clavated tip.

Butterfly with distinct blackish tubular wings spotted with black, and marked beneath with silvery bands.

The Great Wood Paphia.

The Silver-banded Paphia.

The Paphia Paphia may be considered as one of the most elegant of the British lepidoptera. It is principally seen in the neighbourhood of woods, where its larva feeds on nettles, changing into chrysalis in the month of June, and the fly appearing in that of July.

NUMIDA CRISTATA.

CHARACTER GENERICUS.

Caput collo vel pennis cristatum.

Palearia carunculata ad latera capitis.

CHARACTER SPECIFICUS, &c.

Numida collo subnudo cærulescente, capite pennis cristato.

Numida nigra albo maculata, vertice cristato, collo cærulescente subtus sanguineo.

Lath. ind. orn. p. 622.

NUMIDA CRISTATA.

Pall. spic. zool. 4. p. 15. t. 2.

Generatur hæc avis in Africa interiori, minor paulo Numida Meleagridi, seu illa quæ dicitur communis *Pintado*. Distinguit a reliquis speciem collum prope nudum cum capite cristato. Quod in tabula ostenditur specimen in Angliam illatum a domino domino Seaforth, perhumaniter nobiscum communicavit vir honoratissimus Carolus F. Greville, regię majestati Britannicę e conciliis secretioribus.

NUMIDA CRISTATA

CHARACTER GENERICUS

Caput capillo vel pennis cristatum.
Parsus concoloratus ad latera capitis.

CHARACTER SPECIFICUS, &c.

Numida collo rubundo extrinseca, capite
pennis cristato.


Numida nigra albis maculata, vertice cristato,
collo extrinseco rubundo sanguineo.

Lat. ind. 14. 15. 16.

NUMIDA CRISTATA

Lat. ind. 14. 15. 16.

Generatio hanc avia in Africa nascitur, nam paulo
Numida Malesia, sed hic est distinctus communis
Nivea. Distinguitur a reliquis speciebus quod prope
caput cum capite cristato. Quod in tabula ostendi-
tur specimen in Angliam hactenus a domibus domini
Stewart, postquam in Angliam commigravit in
Angliam, Caro in F. Greville, regis imperatoris
Britannice a comitibus secretariis.





THE
CRESTED PINTADO.

GENERIC CHARACTER.

Head furnished with a horny or feathery crest.

Wattles, hanging on each side the head.

SPECIFIC CHARACTER, &c.

Pintado with nearly naked blueish neck, and
head crested with feathers.

CRESTED PINTADO.

Lath. syn. 2. p. 688.

The bird here represented is an inhabitant of the interior parts of Africa, and is of somewhat smaller size than the Numida Meleagris of Linnæus or Common Pintado. It is distinguished as a species by having the neck nearly bare, and the head furnished with a feathered crest. The present specimen was brought into England by Lord Seaforth, and was politely communicated by the right honourable Charles Fulke Greville.

CRISTED TINTADO.

=====

CRISTED TINTADO.

Washed furnished with a pair of feathers
Washed, hanging on each side the head.

CRISTED TINTADO.

Pintado with head in center of the head, and

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

CRISTED TINTADO.

SCARABÆUS AURATUS.

CHARACTER GENERICUS.

Antennæ clavatæ, capitulo fissili.

Tibiæ sæpius dentatæ.

Corpus crassum, compactum.

CHARACTER SPECIFICUS, &c.

SCARABÆUS aureo-viridis subdepressus, segmento abdominis primo lateribus unidentato.

SCARABÆUS AURATUS. S. scutellatus muticus auratus, segmento abdominis primo lateribus unidentato, clypeo planiusculo.

Lin. Syst. Nat. p. 557.

Insectum hoc eximiæ pulchritudinis magnitudine fere æquale est scarabæo vulgari, seu *stercorario* Linnæi: formæ tamen est paulo depressioris, colorisque splendidissime aureo-viridis, qui etiam interdum prope inferiorem elytrorum partem lineis aliquot transversis albidis leviter notatur. Non infrequens est apud nos formosa hæc species quamdiu duraverit æstas calidior; quo tempore varias plantas floresque solet supervolare. Larva plerumque degit vel in cavis arborum, vel in terra levi siccaque juxta radices; nonnunquam etiam in tumulis quos congesserint formicæ: nec nisi post exactos circiter tres annos in chrysalidem convertitur; e qua brevi tempore erumpit Scarabæus.

SCARABAEUS AURATUS

SCARABAEUS AURATUS

SCARABAEUS AURATUS

SCARABAEUS AURATUS

SCARABAEUS AURATUS

SCARABAEUS AURATUS

SCARABAEUS AURATUS

SCARABAEUS AURATUS

SCARABAEUS AURATUS

SCARABAEUS AURATUS

SCARABAEUS AURATUS

SCARABAEUS AURATUS

SCARABAEUS AURATUS

SCARABAEUS AURATUS

SCARABAEUS AURATUS

Insectum hoc est in pulcherrima magnitudine
et replete est coloribus vulgaribus, sed maxime
coloribus rufis et paulo depressis, coloribus
hinc inde rufis, qui etiam inter se
inter se clypeum patet hinc alius transversus
albidus lateris rufus. Non infrequens est apud nos
formae hae species quibusdam hinc rufas
quo tempore vides plures hinc rufas
hinc plures rufas et hinc rufas, vel in
terris hinc rufas hinc rufas; nonnullam etiam
in annis quae congregant formae hae non post
excessum hinc rufas in chrysalidem convertunt;
et quae hinc rufas hinc rufas.

758



Scarabaeus sacer

THE
GOLDEN BEETLE.

GENERIC CHARACTER.

Antennæ or horns clavate, with a fissile tip.

Legs generally toothed.

Body thick and compact.

SPECIFIC CHARACTER, &c.

Subdepressed gold-green Beetle, with the first segment of the abdomen furnished on each side with a prominent tooth.

The Green Garden-Beetle.

The *Scarabæus auratus* or Golden Beetle is a species of peculiar beauty, and is about the size of the common black or garden beetle, but of a somewhat flatter shape, and of a most brilliant golden-green colour, sometimes marked towards the lower part of the wing-sheaths by a few transverse whitish streaks. This elegant animal is not uncommon during the hottest part of summer, frequenting various plants and flowers. Its larva or caterpillar is commonly found in the hollows of trees, or among the loose dry soil at their roots; and sometimes in the earth of ant-hills. It remains about three years before it changes to a chrysalis, out of which in a short time afterwards emerges the complete insect.

LABRUS GUTTULATUS.

CHARACTER GENERICUS.

Dentes robusti subacuti: molares, interdum, conferti, convexi: *labia* crassa, duplicata: *pinnæ dorsalis* radii, in certis speciebus, ramento elongati.

Opercula mutica, squamosa.

CHARACTER SPECIFICUS, &c.

Labrus albo confertim guttatus, dorso subfusco, squamis minutis.

Labrus guttatus. L. maculis rotundis, radiis muticis.

Bloch. ichth. 8. t. 287. f. 2.

Maria incolit Indica pulcherrimus hic piscis, et magnitudine vera in tabula depingitur.

LABRUS CUTTULATUS.

CHARACTER GENERICUS.


Dentes robusti subacuti: molares, interdum
conflati, convexi: labia crassa, duplicata:
pinnæ dorsales tres, in certis speciebus, sa-
piens elongata.
Opercula molia, squamosa.

CHARACTER SPECIFICUS, &c.

Labrus albo confertius guttatus, dorso subnaso,
spinae minutis.
Labrus guttatus. L. maculis rotundis, radiis
molis.

Bloch. Nat. G. t. 237. f. 2.

Natus in India paucissima hic piscis, et
magis in tabula depictus.



THE
GUTTULATED LABRUS.

GENERIC CHARACTER.

Teeth strong and subacute : the grinders sometimes convex and crowded : *lips* thick and doubled : rays of the *dorsal fin*, in some species, elongated into soft processes.

Gill-covers unarmed and scaly.

SPECIFIC CHARACTER, &c.

Labrus with numerous round white spots, brownish back, and minute scales.

GUTTULATED LABRUS.

General Zoology, 4. p. 524.

This beautiful fish is an inhabitant of the Indian seas, and is represented on the plate in its natural size.

HYDRACHNA GEOGRAPHICA.

CHARACTER GENERICUS.

Caput, thorax et abdomen unita.

Palpi duo articulati.

Pedes octo.

Lin. Syst. Nat. Gmel.

CHARACTER SPECIFICUS, &c.

HYDRACHNA globosa, glabra, nigerrima,
coccineo maculata.

HYDRACHNA GEOGRAPHICA. H. ni-
gra sphærica, maculis quatuor punctisque
coccineis.

Lin. Syst. Nat. Gmel. p. 2939.

Mult. Zool. dan. prodr. 2270.

Aquas incolit stagnantes puriores Hydrachna geo-
graphica, specierum omnium hætenus detectarum
sine dubio formosissima. In tabula nostra exprimitur
tam vera quam aucta magnitudo. Insecta prædatur
minora aquatica, ut solet reliquum genus, natansque
motu fertur strenue incitato.

HYDRACHNA GEOGRAPHICA.

CHARACTER CIMENTICUS.

Caput, thorax et abdomen unita.

Papilli duo articulat.

Peda octo.

Lin. Syst. Nat. Gmel.

CHARACTER SPECIFICUS, &c.

HYDRACHNA globosa, glabra, nigrissima,

coccineo maculata.

HYDRACHNA GEOGRAPHICA. H. ni-

gra sphaerica, maculis quatuor punctatis

coccineis.

Lin. Syst. Nat. Gmel. p. 2029.

Mull. Zool. dan. prod. 2270.

Adnas incolit stagnantes putiores. Hydrachna geo-
graphica, speciem omnium hactenus detectarum
sic habio formosissimam. In tabula nostra exprimitur
tam vera quam aetna magnitudo. Insecta praedantur
minora aquarum, et solum solum genus, natusque
magis fortis strenue indicato.



THE
VARIEGATED HYDRACHNA.

GENERIC CHARACTER.

Head, thorax, and abdomen united.

Feelers two, articulated.

Legs eight.

SPECIFIC CHARACTER, &c.

Subglobular, smooth, jet-black Hydrachna,
with scarlet spots.

Great variegated HYDRACHNA.

Crimson-spotted Hydrachna.

The Hydrachna geographica, by far the most beautiful of all the species yet discovered, is an inhabitant of the clearer kind of stagnant waters, and is represented on the plate both in its natural size and magnified. Like the rest of its congeners, it preys on the smaller aquatic insects, and swims with a peculiar degree of vigour and celerity.

THE
VARIEGATED HYDRACHNA.

THE HYDRACHNA VARIEGATA.

GENERIC CHARACTER.

Head, thorax, and abdomen united.

Feet two, articulated.

Legs eight.

SPECIFIC CHARACTER.

Subglobular, smooth, jet-black (Hydrachna).

with scarlet spots.

Great variegated HYDRACHNA.

Crimson-spotted Hydrachna.

The Hydrachna geographicus, by far the most beautiful of all the species yet discovered, is an inhabitant of the clearest kind of stagnant waters, and is reported on the plain to be in its natural size and magnified. Like the rest of the congeners, it feeds on the smaller aquatic insects, and swims with a peculiar degree of vigour and celerity.

TANAGRA SILENS.

CHARACTER GENERICUS.

Rostrum conicum, acuminatum, emarginatum,
basi subtrigonum, apice declive.

Lin. Syst. Nat. p. 313.

CHARACTER SPECIFICUS, &c.

Tanagra olivacea, subtus albida, capite utrinque albo fasciato, humeris flavis.

TANAGRA SILENS. T. viridis, capite subtusque incana, superciliis vitta oculari fasciaque jugulari nigris.

Lath. ind. orn. p. 432.

In America australi generatam Tanagram silentem numerosiorem habet Guiana. Discrepant mores a reliquis congeneribus : amans enim solitudinis latibulum sibi quærit in densissimis sylvis. Ab apice rostri ad extremum caudæ quasi septuncialis est mensura.



W. Verelst. Del. et. Sculp.

THE
SILENT TANAGER.

GENERIC CHARACTER.

Bill conic, acuminated, somewhat inclining towards the point; upper mandible slightly ridged, and notched near the tip.

SPECIFIC CHARACTER, &c.

Olive Tanager, whitish beneath, with the head banded on each side with white, and yellow shoulders.

Oiseau silencieux.

Buff. ois. 4. p. 304.

Tangara de la Guiane.

Pl. Enl. 742.

This species is a native of South America, and is particularly found in Guiana. In its manners it differs from the rest of the Tanagers, being a bird of a solitary nature, and inhabiting the deepest recesses of the woods. It usually measures about seven inches from the tip of the bill to the extremity of the tail.

LABRUS BRASILIENSIS.

CHARACTER GENERICUS.

Dentes robusti, subacuti : molares, interdum, conferti, convexi : *labia* crassa, duplicata : *pinnæ dorsalis* radii, in certis speciebus, ramento elongati.

Opercula mutica, squamosa.


CHARACTER SPECIFICUS, &c.

Labrus flavus, dorso lunulis cæruleis, pinna dorsali analique cæruleo striatis.

LABRUS BRASILIENSIS. L. pinna dorsi anique lineata.

Bloch ichth. 8. t. 280.

Maria incolit Americana pulcherrimus hic piscis, in pedalem vel sesquipedalem crescens longitudinem.



THE
BRASILIAN LABRUS.

GENERIC CHARACTER.

Teeth strong and subacute : the grinders sometimes convex and crowded : *lips* thick and doubled : rays of the *dorsal fin*, in some species, elongated into soft processes.

Gill-covers unarmed and scaly.

SPECIFIC CHARACTER, &c.

Yellow Labrus, with the back marked by blue crescents, and the dorsal and anal fin with blue stripes.

BRASILIAN LABRUS.

Gen. Zool. 4. p. 497.

Der brasilianische Lippfisch.

Bloch ichth. 8. pl. 280.

This beautiful fish is an inhabitant of the American seas, and arrives at the length of twelve or eighteen inches.

GRYLLUS MORBILLOSUS.

Var.

CHARACTER GENERICUS.

Caput inflexum, maxillosum, palpis instructum.

Antennæ setaceæ seu filiformes.

Alæ quatuor, deflexæ, convolutæ: inferiores plicatæ.

Pedes postici saltatorii: ungues ubique bini.

Lin. Syst. Nat. p. 692.

CHARACTER SPECIFICUS, &c.

GRYLLUS MORBILLOSUS. Gryllus thorace quadrato rubro verrucoso, elytris fuscis albo punctatis, alis rufis.

Lin. Syst. Nat. p. 700.

Seb. mus. 4. t. 79. f. 7, 8.

Roes. ins. 2. t. 18. f. 6.

Insectum, cujus iconem suppeditavit præclarum Roeselii opus, idem ferme putant auctores atque gryllum morbillosum Linnæi. In hoc tamen ab eo differt quod alæ superiores olivaceo-virent, maculis nullis albis, sed plurimis fuscis aspersæ. Africam incolit inferiorem; nec carere eo dicuntur quædam etiam Americæ australis regiones.



THE
GRANULATED LOCUST.

Var.

GENERIC CHARACTER.

Head inflected, armed with jaws, and furnished with feelers.

Antennæ, in most species, setaceous or filiform.

Wings four, deflected and convoluted; the lower ones pleated.

Hind-Legs formed for leaping: claws on all the feet double.

SPECIFIC CHARACTER, &c.

Locust with square, red, warted thorax, brown wing-cases spotted with white, and red wings.

The Spotted Cape Locust.

The present species, which is copied from the work of the admirable Roesel, is generally quoted by entomological writers as the *Gryllus morbillosus* of Linnaeus. It differs however in the colour of the upper wings, which, instead of being spotted with white, are of an olive-green, with very numerous dusky specks. It is a native of the lower parts of Africa, but is said to occur also in some parts of South-America.

GRANULATED LOCUST

N.Y.

GENERAL CHARACTER

Head reddish, fringed with jaws, and furnished with feelers. Antennae, in most species, setaceous or filiform. Body four, definite, and colored; the lower ones often spotted. First leg formed into a long, sharp, and double.

SPECIES OF THE GENUS

Locust with square, red, white, brown, brown wing-chains spotted with black, and red wings.

The Spotted Cape Locust

The present species, which is found from the wing of the admirable Rose, is generally colored by conological writers as the British specimens of this genus. It differs however in the color of the upper wings, which, instead of being spotted with white, are of an olive green, with very abundant dusky spots. It is a native of the lower parts of Africa, but is said to occur also in some parts of South America.

ACARUS PECTINATUS.

CHARACTER GENERICUS.

Pedes Octo.

Oculi duo, ad latera capitis.


Tentacula duo, articulata, pediformia.

Lin. Syst. Nat. p. 1022.

CHARACTER SPECIFICUS, &c.

Acarus ovatus albidus, capite subacuminato,
tentaculis crassissimis apice pectinatis.

Æqualis fere magnitudine est species de qua jam
agitur Acaro vulgari seu *Sirani* Linnæi; locorum prope
eorundem incola: cum tamen longe sit rarior, eo, ut
opinor, non antea descripta est.



Dr. L. Zelt

8. Müllerer. Sculp.

THE
PECTINATED MITE.

GENERIC CHARACTER.

Legs eight.

Eyes two, situated on each side the head.

Feelers two, jointed, leg-formed.

SPECIFIC CHARACTER, &c.

Ovate whitish Mite, with subacuminated head,
and very thick chelated tentacula with
pectinated tips.

The present species of *Acarus* is about the size of the common domestic Mite or *Acarus Siro* of Linnaeus, and is found in similar situations: it is however a much rarer animal, and seems to have been hitherto undescribed,

THE

PECTINATED MITE

GENERIC CHARACTER

Legs eight.

Legs two, situated on each side the head.

Legs two, jointed, leg jointed.

SPECIFIC CHARACTER

Ovary whitish, with subacuminated head,
and very thick, elliptical, reniform, with
pectinated tips.

The present species of *Acanth* is about the size of
the common domestic *Acanth* of *Acanth* and of *Acanth*.
It is found in similar situations. It is how-
ever a much rarer animal, and seems to have been in-
tentionally introduced.

AMPELIS MAYNANA.

CHARACTER GENERICUS.

Rostrum rectum, convexum : mandibula superiore longiore, subincurvata, utrinque emarginata.

Lingua acuta, cartilaginea, bifida.

Lin. Syst. Nat. p. 297.

CHARACTER SPECIFICUS, &c.

Ampelis nitida cærulea, gula violacea, pennis capitis collique angustatis.

Ampelis Maynana. *A. nitida* cærulea, gula violacea.

Lin. Syst. Nat. p. 298.

Ampelis nitida cærulea, gula violacea, remigibus intus albis.

Lath. ind. orn. p. 365.

Americæ Australis partes calidiores incolit *Ampelis Maynana*. Magnitudine *Turdo iliaco* paulo inferior est.



THE
SILKEN-FEATHERED
CHATTERER.

GENERIC CHARACTER.

Bill strait, a little convex above, bending towards the point: notched on each side near the tip of the upper mandible.

Tongue sharp-pointed, cartilaginous, bifid.

SPECIFIC CHARACTER, &c.

Shining-Blue Chatterer, with violet-purple throat; the head and neck covered with narrow feathers.

The Silky Chatterer.

Lath. synops. 2. p. 96.

Le Cotinga des Maynas.

Briss. av. 2. p. 341.

Le Cotinga à plumes soyeuses.

Buff. ois. 4. p. 447.

This highly elegant species is a native of the hotter parts of South America. Its size is somewhat inferior to that of a Redwing.

TRICHODA TRISETA.

CHARACTER GENERICUS.

Vermis nudo oculo inconspicuus, pellucidus,
altera parte crinitus.

Lin. Syst. Nat. Gmel.

CHARACTER SPECIFICUS, &c.

Trichoda subovata hyalina interaneis fuscis,
seta unica barbata antica, et duabus pos-
ticis.

Animalculum quod depinximus nemo, ut opinor,
antea descripsit. Ut clare et distincte conspici possit
et examinari, microscopio opus est.

Detectum est mense Junio in aqua stagnanti et tur-
bida. Pellis, seu theca externa, conchæ bivalvi non
longe absimilis, pellucida admodum est; intestinorum
autem color fuscus. Movet se animalculum quasi
subsultim, seu impetu quodam subitaneo.

TRICHODA TRISTATA.

Trichoda tristata, n. sp.

CHARACTERES SPECIFICI.

Forma rudo oculo incomposita, pilulosa
alabastrina parte cinerea.


Loc. Agra, N. G. G.

CHARACTERES SPECIFICI.

Trichoda subovata hyalina nitens, fere
seta unica barbata antica, et duabus
tenuis.

Animalculum quod depictum nemo, ut opinor
antea descripsit. Ut clare et distincte compari possit
et examinari, microscopio opus est.
Dorsum est nigrum, fere in aqua saturatum, et ful-
gidum. Pells, seu theca externa, coram pectus non
longe abeuntibus, pellicula admodum est, intermedium
autem color fuscus. Moveri et animalculum quod
subulium, seu corpus quoddam subulium.

766
767



W. J. L. L. L.

U. S. - Dalt

THE
THREE-BRISTLED TRICHODA.

GENERIC CHARACTER.

Animalcule inconspicuous to the naked eye, beset on one part with hairs.

SPECIFIC CHARACTER, &c.

Subovate hyaline Trichoda, with brown intestines, a single bearded bristle in front, and two behind.

The animalcule here figured appears to have been hitherto undescribed. It is distinctly visible by the assistance of the microscope only, and was discovered in turbid stagnant water in the month of June.

The exterior covering, which has some appearance of a bivalve shell, is perfectly transparent, but the intestines are of a brown colour. The motions of this animalcule are generally performed by sudden starts or springs.

MILLEPORA VIOLACEA.

CHARACTER GENERICUS.

Animal Hydra.

Corallium poris turbinatis teretibus.

Lin. Syst. Nat. p. 1282.

CHARACTER SPECIFICUS, &c.

Millepora violacea, in plano ramosa, ramulis tereti-compressis, sutura marginali porosa.

Millepora Violacea. M. in plano ramosa, ramulis ascendentibus flexuosis tereti-compressis, sutura porosa marginem ambiente.

Soland. et Ellis zooph. p. 140.

Sufer rupes maris Indici conspicitur Millepora violacea, magnitudine vera in tabula depicta.

MILLEPORA VIOLOACEA

SYNOPSIS

CHARACTER SPECIFICUS

Ammon, Hydrog.

Conchologia portus turbinata testibus


Vol. 2, p. 122.


CHARACTER SPECIFICUS, etc.

Millepora violacea, in plano tenuis, ramulis
tortis compressis, sicut marginali porosa.

Millepora Violacea. M. in plano tenuis, 12-
radis ascendentibus flexuosis tortis com-
pressis, sicut porosa marginem ambiens.
Soland. et Ellis. voyag. p. 140.

Sicut super maris Indici conspicitur Millepora vi-
olacea, nuperrime vero in tabula depicta.





R. W. Wood Del. et Sculpit

THE
VIOLACEOUS MILLEPORE.

GENERIC CHARACTER.

Animal allied to a Polype.

Coral with very numerous cylindric pores.

SPECIFIC CHARACTER, &c.

Flat, branching, violet-coloured Millepore,
with subcompressed branches margined
with pores.

Violet-Coloured Millepore.

Soland. et Ellis zooph. p. 140.
pl. 26. f. 3.

The Violaceous Millepore is found on the rocks of
the Indian Ocean, and is represented on the plate in
its natural size.

SCORPÆNA VOLITANS.

CHARACTER GENERICUS.

Caput magnum, aculeatum, cirrosum, obtusum, squamis nudum, subcompressum: *Oculi* vicini: *Dentes* maxillis, palato, faucibusque.

Membr. branch. radiis septem.

Corpus torosum. *Pinna dorsalis* unica, radiis anterioribus spinosis.

CHARACTER SPECIFICUS, &c.

Scorpæna fusco flavoque fasciata, pinnis pectoralibus maximis violaceis albo guttatis.

Scorpæna volans. S. pinnis pectoralibus trunco longioribus.

Bloch. ichth. t. 184.

Gasterosteus volitans. G. spinis dorsalibus tredecim, cirris senis, pinnis pectoralibus corpore longioribus.

Lin. Syst. Nat. p. 491.

In maribus Indicis et Americanis generatur *Scorpæna*, forma singulari magis quam pulchra insignis. Attingit interdum longitudinem plus pedalem. Hoc ei commune est cum exocoetis, et alijs nonnullis piscibus, quod pinnis magnis pectoralibus quasi alis supra aquas elata, per breve tempus volitare possit, et hoc modo hostium assultus non raro eludere.

SCORPENA VOLITANS.

SCORPENA VOLITANS.

SCORPENA VOLITANS.

Caput marginatum, aculeatum, ciliatum, obtusum,
spinae nudum, subcompressum: Oculi
virescentes: Dentes maxillares, palato, lacinias
que.

Abdomen breviter radiis septem.
Corpus robustum. Pinnae dorsales, radiis
anterioribus spinosis.

SCORPENA VOLITANS.

Scorpena trisco flavo fasciata, pinnis pec-
toralibus maxime violaceis albis guttatis.
Scorpena volans. 2. pinnis pectoralibus trunco
longioribus.

Abdomen breviter radiis septem.
Gasterostea volitans. G. spinis dorsalibus
tridactylis, ciliis acutis, pinnis pectoralibus
corpore longioribus.

Lin. Syst. Nat. p. 491.

In maribus laticis et rivieris generat Scor-
pena, forma elongata magis quam pinnis longior.
Atingit interdum longioribus pinnis pectoralibus. 1. for-
a communis est cum pectoralibus, et alia ventralibus pinnis
albis, quibus pinnis magis pectoralibus pinnis longioribus
pinnis acutis, pinnis longioribus pinnis longioribus.
hoc non. Scorpena volans. 2. pinnis pectoralibus trunco



THE
FLYING SCORPÆNA.

GENERIC CHARACTER.

Head large, aculeated, cirrhose, obtuse, without scales, subcompressed: *Eyes* placed near each other: *Teeth* in the jaws, palate, and throat.

Gill-Membrane seven-rayed.

Body fleshy: *Dorsal-Fin* single, with the rays of the fore-part spiny.

SPECIFIC CHARACTER, &c.

Scorpæna variegated with brown and yellow transverse bands, with very large violet-coloured pectoral fins spotted with white.

Flying Scorpæna.

Gen. Zool. 4. p. 277.

The Scorpæna volitans, or Flying Scorpæna, more remarkable for its singularity than its beauty, is a native of the Indian and American seas, and has been sometimes seen of the length of more than twelve inches. Like the Exocoeti, and some other fishes, it possesses the power of using its large pectoral fins for the purpose of short occasional flight over the surface of the water, and thus frequently eludes the pursuit of its enemies.

PSITTACUS INCERTUS.

CHARACTER GENERICUS.

Rostrum aduncum, mandibula superiore mobili,
cera instructa.

Nares in rostri basi.

Lingua carnosae, obtusa, integra.

Pedes scansorii.

Lin. Syst. Nat. p. 139.

CHARACTER SPECIFICUS.

Psittacus brachyurus viridi-cæruleus, humeris
subtus scapularibusque rubris, cauda flava.

Obs. Remiges subtus nigræ.

Psittaci speciem, de qua jam agitur, satis ad discriminationem a nullo auctore descriptam fuisse arbitror. Major est psittaco pullario Linnæi, longitudine quasi septunciali. Habitu est robusto magis quam delicato; et specimen (unde ipsa avi quarta parte minor delineata est similitudo) provectioris fuisse ætatis crediderim: quod satis probat mandibula superior plus solito elongata. In India generari creditur. Vixit avis quam descripsimus novenos annos in aviario dominæ Read.



22. 10. 1880. Pelat. Lark.

THE
BLUE-GREEN PARROT.

GENERIC CHARACTER.

Bill hooked : upper mandible moveable.

Nostrils round, placed in the base of the bill.

Tongue fleshy, broad, blunt at the end.

Legs short ; feet scansorial.

SPECIFIC CHARACTER.

Short-tailed blue-green Parrot, with the shoulders beneath, and scapular-feathers red ; the tail yellow.

The present species of Parrot does not appear to have been distinctly described by any ornithological writer. In size it considerably exceeds the *Psittacus pullarius* of Linnæus, measuring about seven inches in length. Its general habit is rather distinguished by an appearance of strength than delicacy, and the specimen from which the figure, (reduced about one fourth in size) is taken, appears to have been of advanced age, as the considerable elongation of the upper mandible seems to prove. This bird is supposed to be a native of India, and was in the possession of Lady Read, with whom it lived during the space of nine years.

BLUE GREEN PARROT.

It is a small, compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

compact, and very

LACERTA FURCIFRONS.

CHARACTER GENERICUS.

Corpus tetrapodum, elongatum, caudatum, nudum.

CHARACTER SPECIFICUS.

Lacerta fusca, fronte bifida, abdominis lateribus flavo maculatis.

Cameleon bifurque.

Brongniart, B. Soc. Phil. No. 36.

Cum in opere quod edidimus zoologico prætermissa fuerit hæc chamæleontis species, iconem ejus e libro Daudinii de amphibiiis desumptam in tabulam nostram transtulimus.

Lacertam furcifrontem primus descripsisse videtur Jacobus Parsonsus in actorum philosophicorum volumine quinquagesimo. Si habitum generalem spectas, summa ei videtur esse affinitas cum chamæleonte communi; nisi quod capitis peculiaris sit conformatio, corpusque non virescat, sed supra fusco-nigricet, infra pallidius. Plantæ aurantiæ sunt; lateraque abdominis variata serie duplici macularum parvarum flavarum et rotundarum, in longitudinem ducta. Insignem hanc lacertam in insulis Indicis generari compertum est.

FACTA FUERUNT

in hunc modum

et cetera

et cetera

et cetera

et cetera

et cetera

et cetera

et cetera

et cetera

et cetera

et cetera

et cetera

et cetera

THE
FORK-HEADED CHAMÆLEON.

GENERIC CHARACTER.

Body four-footed, elongated, tailed ; without any secondary integument.

SPECIFIC CHARACTER.

Brown Chamæleon, with bifid front, and yellow spots on each side the abdomen.

Le Cameleon nez-fourchu.

Daud. Buff. Sonnin. rept. 4 p. 217.

Chameleon, &c. *Pars. Phil. Trans.*
vol. 54.

This remarkable species of Chamæleon having been omitted in the work entitled General Zoology, I here introduce its figure from Mons. Daudin's history of the Amphibia. The first description of this animal seems to have been published by Dr. Parsons, in the fiftieth volume of the Philosophical Transactions. In its general appearance it is allied in the highest degree to the common Chamæleon, but differs in the extraordinary form of its head, and in the colour of the body, which, instead of being green, as in that species, is of a dusky or blackish brown above, and paler beneath ; the soles of the feet are orange-coloured, and the sides of the abdomen are variegated by a double longitudinal row of small, round, yellow spots. It is a native of the Indian islands.

FORK-HEADED CHAMÆLEON

Body four-fifths black, tinged with
any color, irregularly
marked with
Brown. Dorsal spots, and front, and
for spots on sides of the abdomen.
The throat, neck, and
Dorsal spots, and
Chamæleon, the Fork-Headed
vol. 1

This remarkable species of Chamæleon having been
noticed in the late edition of General Zoology, I have
the object in view to give a more detailed history of the
species. The first description of this animal seems
to have been published by Dr. Parson, in the fifth
volume of the Philosophical Transactions. It was
first described as being in the highest degree of the
common Chamæleon, but differs in the extraordinary
form of its head, and in the color of the body, which
instead of being green, as in that species, is of a dusky
or blackish brown above, and paler beneath: the sides
of the feet are orange-colored, and the sides of the
abdomen are variegated by a double longitudinal row
of small round yellow spots. It is a native of the
Indian islands.

VORTICELLA CYATHUS.

CHARACTER GENERICUS.

Corpus contractile, nudum, ciliis rotatoriis.

Lin. Syst. Nat. Gmel. p. 3874.

CHARACTER SPECIFICUS.

Vorticella corpore infundibuliformi subhexagono, ciliis quatuor exsertis.

Quod in tabula exprimitur animalculum, ope microscopii summopere auctum, ad genus Linnæanum Vorticellæ summo jure pertinere arbitror. Dubitari tamen possit annon antea descriptum fuerit. Forma non longe distat a similitudine infundibuli; parsque superior leviter hexagona dividitur intus in sex quasi cellulas seu cavitates. Ab ore protruduntur quatuor tentacula acuta; et in centro visibilia sunt duo breviora. Animalculum pellucide pallens fusco-flavescit: intestinorum longe saturatior est color. Motu fertur alacri validoque; et hoc commune habet cum multis congeneribus, quod ad libitum contrahere possit et occludere hiatum sive corporis partem superiorem: quo in actu depingitur in tabulæ figura secunda, quasi pyriforme

VORTICELLA CYATHUS.

CHARACTER GENERICUS.


Corpus contractile, nudum, ciliis rotatoriis.

Lin. Syst. Nat. Gmel. p. 3874.


CHARACTER SPECIFICUS.

Vorticella corpore infundibuliformi subhexagono, ciliis quatuor exsertis.

Quod in tabula exprimitur animalculum, ope microscopii summopere auctum, ad genus Linnæanum Vorticellæ summo jure pertinere arbitror. Dubitari tamen possit annon antea descriptum fuerit. Forma non longe distat a similitudine infundibuli; parsque superior leviter hexagona dividitur intus in sex quasi cellulas seu cavitates. Ab ore protruduntur quatuor tentacula acuta; et in centro visibilia sunt duo breviora. Animalculum pellucide pallens fusco-flavescit: intestinorum longe saturatior est color. Motu fertur alacri validoque; et hoc commune habet cum multis congeneribus, quod ad libitum contrahere possit et occludere hiatum sive corporis partem superiorem: quo in actu depingitur in tabulæ figura secunda, quasi pyriforme



771



G. P. - Det.

R. P. N. det. P. L.

THE
CUP VORTICELLA.

GENERIC CHARACTER.

Body contractile, naked, furnished with rotatory organs.

SPECIFIC CHARACTER.

Vorticella with subhexagonal cup-shaped body, and four exserted cilia.

The animalcule represented as highly magnified on the present plate belongs most properly to the Linnæan genus Vorticella : it may be doubted however whether it has ever before been described. Its form somewhat resembles a funnel, and the superior part is of a slightly hexagonal shape, divided internally into six cells or cavities. Four sharp tentacula are protruded from the opening, and two smaller are visible in the central part. The colour of this animalcule is a very pale, transparent, yellowish brown, the intestines appearing of a much darker cast. Its motions are swift and vigorous, and it possesses the power, like many others of this genus, of contracting at pleasure the upper part or mouth, in which state it assumes a pear-shaped form, as represented by the second figure.

CUP VORTICELLA.

CHARACTERISTICS.

CHARACTERISTICS.

Body cylindrical, naked, furnished with setae
long, sparse.

CHARACTERISTICS.

Vorticella with subhexagonal cup-shaped body,
and four exerted cilia.

The animalcule represented as highly magnified on
the present plate belongs to the Japanese
genus Vorticella. It may be doubted however whether
it has ever before been described. Its form somewhat
resembles a funnel, and the superior part is of a slightly
hexagonal shape, divided internally into six cells or
lobes. Four sharp processes are protruded from the
opening, and two smaller setae in the central part.
The colour of the animalcule is a very pale, opaque
red, yellowish brown, the interior appearing of a
much darker hue. Its motions are swift and vigorous,
and it possesses the power, like many others of this ge-
nus, of contracting at pleasure the upper part or mouth,
in which state it assumes a pear-shaped form, as repre-
sented by the second figure.

GRYLLUS DUX.

CHARACTER GENERICUS.

Caput inflexum, maxillosum, palpis instructum.

Antennæ quatuor, deflexæ, convolutæ: inferiores plicatæ.

Pedes postici saltatorii: ungues ubique bini.

Lin. Syst. Nat. p. 692.


CHARACTER SPECIFICUS.

Gryllus viridis, thorace carinato scabro, elytris olivaceis, alis rubris nigro maculatis et marginatis.

Gryllus Dux. G. thorace carinato scabro, elytris viridibus, alis rufis fusco maculatis.

Fab. sp. ins. 1 p. 362.

Magnitudine et facie generali simillima est hæc species gryllo *cristato* Linnæi, de quo in opere hoc nostro antea disseruimus. Ab eo tamen differt, quod alæ inferiores pulcherrime rubræ, nigroque marginatæ, maculis plurimis nigris et quasi sagittatis decorantur. Americam incolit australem gryllus dux, nec ulla videtur ejus extitisse descriptio nisi in elegantissimo Druryi opere de insectis exoticis.



22. *Acrida rubra*

THE
IMPERIAL LOCUST.

GENERIC CHARACTER.

Head inflected, armed with jaws, and furnished with feelers.

Antennæ in (most species) setaceous, or filiform.

Wings four, deflected and convoluted; the lower pair pleated.

Hind-Legs formed for leaping: claws on all the feet two.

SPECIFIC CHARACTER.

Green Locust, with rough, keeled thorax, olive-brown wing-sheaths, and red wings spotted and edged with black.

Gryllus Dux.

Drury ins. 2. pl. 44.

This superb insect in size and general appearance is very nearly allied to the *Gryllus cristatus*, or Great Egyptian Locust, before figured in the present work, from which it differs in having the lower wings of a beautiful red colour, edged with black, and marked with numerous arrow-shaped spots of the same colour. It is a native of South-America, and seems to have been first described in the elegant work of Mr. Drury.

THE

IMPERIAL LOCUST.

GENERAL CHARACTER.

Head reddish, armed with jaws, and furnished with locusts.

Antennae in (most species) setaceous, or filiform. Wings four, deflexed and convoluted; the lower pair obsolete.

Hind-legs formed for leaping; claws on all the feet two.

SPECIFIC CHARACTER.

Green Locust, with rough, keeled thorax, olive-brown wing-sheaths, and red wings spotted and edged with black.

Gryllus Dux.

Described by Mr. Dux.

This species inhabits a vast and general appearance is very nearly allied to the Chinese locust, or Great Egyptian Locust, but it differs in the present work from which it differs in having the lower wings of a beautiful red color, edged with black, and marked with numerous brownish spots of the same color. It is a native of South America, and seems to have been first introduced in the present work of Mr. Dux.

TETRAO FRANCOLINUS.

CHARACTER GENERICUS.

Rostrum convexum, breve.

Macula prope oculos nuda, aut papillosa, aut plumis rarius tecta.

CHARACTER SPECIFICUS.

Tetrao ferrugineo nigroque varius, abdomine gulaque atris.


Tetrao Francolinus. T. pedibus nudis calcaratis, abdomine gulaque atris, cauda cuneata.

Lin. Syst. Nat. p. 275.

Perdix Francolinus. P. calcarata, corpore vario, abdomine gulaque atris, cauda cuneata.

Lath. ind. orn. p. 644.

Varias Europæ calidioris regiones incolit Tetrao Francolinus, magnitudine quasi Perdix vulgaris sive cinereæ.





R. N. Adair Del. et Sculp.

THE
FRANCOLIN.

GENERIC CHARACTER.

Bill convex, short.

Spot near the eyes, either naked, granulated,
or but sparingly covered with feathers.

SPECIFIC CHARACTER.

Ferruginous Partridge, with black variegations,
abdomen and throat black, and cuneated
tail.

The Francolin Partridge.

Le Francolin.

Buff. ois. 2. p. 438.

Pl. Enl. 147. 148.

The Francolin.

Edwards pl. 246.

The elegant bird here represented is a native of
many of the warmer parts of Europe. Its size is that
of a common Partridge.

SIMIA ARGENTATA.

CHARACTER GENERICUS.

Dentes primores utrinque quatuor, approximati.

Laniarii solitarii, longiores, hinc remoti.

Molares obtusi.


CHARACTER SPECIFICUS, &c.

Simia alba, facie auriculisque rubris, cauda castanea.

Simia argentata. *S. caudata* imberbis exalbida, facie rubra, cauda fuscescente.

Lin. Syst. Nat. Gmel. p. 41.

Contribulium pulcherrima est species quam depinximus. Magnitudo est quasi communis sciuri. Color albus; partibus corporis superioribus subflavo levissime tinctis. Facies internaue aurium superficies splendide miniata. Cauda castanea. Americam incolit australem *Simia argentata*; unde multis abhinc annis in Europam transtulit Condaminus unicum specimen quod exstat apud Europæos, quodque in Museo asservatur Parisiorum.





THE
SILVER-HAIRED MONKEY.

GENERIC CHARACTER.

Front-Teeth in each jaw four, placed near together.

Canine-Teeth solitary, longer than the others, distant from the grinders.

Grinders obtuse.

SPECIFIC CHARACTER.

White Monkey, with the face and insides of the ears red, and the tail chesnut-coloured.

Le Mico.

Buff. Quadr. 15. p. 121. pl. 18.

Fair Monkey.

Pennant Quadr.

This species may be considered as the most beautiful of its tribe. Its size is that of a common squirrel, and its colour white, with a very slight cast of pale yellow on the upper parts : the face and insides of the ears are of a vermilion red, and the tail of a deep chesnut-colour. It is a native of South America, from whence the only specimen yet seen in Europe, and which is preserved in the Paris Museum, was many years ago imported by Mons. Condamine.



R.H. - Sculpt.

BRACHIONUS GALEATUS.

CHARACTER GENERICUS.

Animalculum nudo oculo inconspicuum, contractile, testa tectum.

CHARACTER SPECIFICUS.

Brachionus pellucidus, corpore elongato, capitis clypeo transverse ovato, cauda apice bifida.

Neminem opinor insigne hoc animalculum antea descripsisse. Aquas incolit stagnantes mense Junio. Pellucidum admodum est, et tantæ velocitatis, ut in microscopio ægre possit examinari, donec guttula incarcerationis evaporata fuerit. Caput subtriangulum est, rostro obtuso, oculisque duobus nigris; super quos sita est seta tenuissima, antennis insecti non absimilis. Totum caput quasi theca contegitur eximie pellucida. Cauda qua corpori inseritur geniculata, deinceps in setas duas seu furcas flexiles dividitur. Ostenditur animalculum in tabula summopere auctum. A Brachiono lamellari Mülleri species revera diversa est, quamvis ad ejus similitudinem proxime videatur accedere.

775



R. P. V. Sculpt.

THE
GALEATED BRACHIONUS.

GENERIC CHARACTER.

Animalcule inconspicuous to the naked eye, with contractile body, covered by a shell.

SPECIFIC CHARACTER.

Pellucid Brachionus, with lengthened body, head with transversely oval shield, and tail bifid at the tip.

This remarkable animalcule, which seems to have been hitherto undescribed, is an inhabitant of stagnant waters in the month of June. It is extremely transparent, and so swift as to be with difficulty kept in view till the small drop in which it lies confined, while under examination by the microscope, begins to evaporate. The head is of a subtriangular shape, with an obtuse snout and two black eyes, above each of which is situated a fine bristle, resembling the antenna of an insect. The whole head seems covered by a kind of shell or case of extreme transparency. The tail is jointed at its insertion to the body, and soon afterwards divides into two branches or flexible forks. This animalcule is represented on the plate as magnified to a very high degree. It seems to approach nearest to the *Brachionus lamellaris* of Müller, but is a distinct species.

THE GALILEATED BRACHIONUS

GENERAL CHARACTER

Animals microscopic to the naked eye, with
conspicuous body, covered by a shell.

SPECIFIC CHARACTER

Galileated Brachionus, with elongated body,
head with transverse oval shield, and tail
with a tip.

The microscopic animal, which seems to have
been first described, as an individual of stagnant
waters in the month of June. It is extremely trans-
parent, and so small as to be with difficulty kept in
view with the small drop in which it has been observed, when
under examination by the microscope. It seems to ex-
hibit a head of a subtriangular form, with an
oblique shield on the back, and above each of which
is situated a fine bristle. Behind the shield is an
inverted T-shaped head shield, covered with fine
setae. The tail is of extreme length, and the tail is
divided at its insertion to the body, and runs inwards
division into two branches or leaf-like lobes. The an-
terior end is represented on the plate as related to a
very small figure. It seems to approach nearest to the
Brachionus latipennis of Lillie, but is a distinct
species.

PHALÆNA ODORA.

CHARACTER GENERICUS.

Antennæ setacæ, a basi ad apicem sensim attenuatæ.

Alæ (sedentis) sæpius deflexæ (*Volatu nocturno.*)

Lin. Syst. Nat. p. 808.

CHARACTER SPECIFICUS.

Phalæna alis crenatis fuscis, subferrugineo undulatis, superioribus ocello nigro luteo marginato.

Phalæna odora. *P. pectinicornis spirilinguis*, alis crenatis nebulosis; superioribus ocello unico; inferioribus sesquialtero.

Lin. Syst. Nat. p. 811.

Americam australem insulasque occiduas incolit insignis hæc phalæna, cujus veram magnitudinem exprimit tabula.



THE
SABLE MOTH.

GENERIC CHARACTER.

Antennæ setaceous, gradually decreasing from base to point.

Wings (when at rest) generally deflected. Flight nocturnal.

SPECIFIC CHARACTER.

Moth with crenated dark-brown wings with subferruginous undulations; the upper pair marked by a black eye-shaped spot with luteous margin.

Drury ins. 1. pl. 3. f. 1.

This magnificent Moth is a native of South America and the West Indian islands. The plate represents it in its natural size.

COLUMBA MELANOCEPHALA.

CHARACTER GENERICUS.

Rostrum rectum, versus apicem descendens.

Nares oblongæ, membrana molli tumida semitectæ.

Lingua integra.

CHARACTER SPECIFICUS.

Columba viridis, capite cano, occipite nigro, gula crissoque flavis, rectricibus utrinque tribus rubris.

Columba melanocephala. *C. viridis*, capite cinerascens, occipite nigro, gula abdomine imo crissoque flavis, rectricibus tribus exterioribus purpureo-coccineis.

Lath. ind. orn. p. 610.

Indiam insulasque Indicas incolit formosa hæc avis, columba vulgari sive domestica paulo minor.

COLLEGE & MUSEUM

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...



W. L. G. Del. & Sculp.

THE
BLACK-CAPPED PIGEON.

GENERIC CHARACTER.

Bill strait, descending towards the tip.

Nostrils oblong, half covered by a soft tumid membrane.

Tongue entire.

SPECIFIC CHARACTER.

Green Pigeon, with grey head black behind, throat and vent yellow, and three of the tail-feathers on each side red.

Turvert.

Buff. ois. 2. p. 555.

Black-capped Pigeon.

Lath. syn. 2. p. 654.

This beautiful bird is a native of India and the Indian islands, and is somewhat smaller than the common domestic pigeon.

BODIANUS APUA.

CHARACTER GENERICUS.

Habitus Percæ.

Opercula squamosa, non serrata, aculeata.

Squamæ (pluribus) læves.

CHARACTER SPECIFICUS.

Bodianus ruber, dorso maculis, corpore punctis
nigris, pinnis nigro marginatis.

Gen. Zool. 4. p. 572.

Bodianus Apua. *B. aculeis* septem in pinna
dorsi, unicoque ad operculum.

Bloch ichth. 7. p. 37. t. 229.

In rivis et circa littora Brasiliæ præcipue conspicitur
Bodianus Apua. In longitudinem plusquam pedalem
crescit, et in cibis habetur lautioribus.


Всего же в 1812 году
всего же в 1812 году
всего же в 1812 году

Всего же в 1812 году
всего же в 1812 году
всего же в 1812 году

Всего же в 1812 году
всего же в 1812 году
всего же в 1812 году

Всего же в 1812 году
всего же в 1812 году
всего же в 1812 году

Всего же в 1812 году
всего же в 1812 году
всего же в 1812 году



THE
APUA BODIAN.

GENERIC CHARACTER.

Habit of the genus *Perca*.

Gill-Covers scaly, not serrated, aculeated.

Scales (in most species) smooth.

SPECIFIC CHARACTER.

Red Bodian, with the back spotted, the body speckled, and the fins edged with black.

Apua Bodian.

Gen. Zool. 4. p. 573.

Pirati Apua.

Marcgr. Bras. p. 157.

The fish here represented is principally found about the coasts and in the rivers of Brasil. It grows to more than twelve inches in length, and is numbered among the most delicate fishes for the table.

LIBELLULA CÆRULATA.

CHARACTER GENERICUS.

Os maxillosum, maxillis pluribus.

Antennæ brevissimæ.

Alæ extensæ.

Cauda maris hamoso-forcipata.


CHARACTER SPECIFICUS.

Libellula corpore gracili longissimo, alis pellucidis, macula prope apicem magna atrocærulea.

Libellula cærulata.

Drury ins. exot. 3. t. 50. f. 1.

In America australi generatur Libellula cærulata, et magnitudine vera in tabula exprimitur.



THE
BLUE-TIPPED DRAGON-FLY.

GENERIC CHARACTER.

Mouth furnished with several jaws.

Antennæ very short.

Wings four, extended.

Tail (in the male) hook-forcipated.

SPECIFIC CHARACTER.

Dragon-Fly with very long slender body, and pellucid wings marked near the tip by a large blackish-blue spot.

Drury ins. 3. pl. 50. f. 1.

The blue-tipped Dragon-Fly is a native of South-America, and is represented on the plate in its natural size.

THE HISTORY OF THE CITY OF LONDON

BY JOHN STOW.

London, the chief city of England, is situated on the north side of the River of Thames, which runs through the middle of it, and divides it into two parts, the one called the Old London, and the other the New London. The Old London is the part which was first inhabited, and is now the most populous part of the city. The New London is the part which was built by Henry the Second, and is now the most beautiful part of the city. The city of London is the most populous city in England, and is the most beautiful city in the world. It is situated on the north side of the River of Thames, which runs through the middle of it, and divides it into two parts, the one called the Old London, and the other the New London. The Old London is the part which was first inhabited, and is now the most populous part of the city. The New London is the part which was built by Henry the Second, and is now the most beautiful part of the city.

The city of London is the most populous city in England, and is the most beautiful city in the world. It is situated on the north side of the River of Thames, which runs through the middle of it, and divides it into two parts, the one called the Old London, and the other the New London. The Old London is the part which was first inhabited, and is now the most populous part of the city. The New London is the part which was built by Henry the Second, and is now the most beautiful part of the city.

The city of London is the most populous city in England, and is the most beautiful city in the world. It is situated on the north side of the River of Thames, which runs through the middle of it, and divides it into two parts, the one called the Old London, and the other the New London. The Old London is the part which was first inhabited, and is now the most populous part of the city. The New London is the part which was built by Henry the Second, and is now the most beautiful part of the city.

SEPIA GRANULATA.

CHARACTER GENERICUS.

Corpus carnosum, vagina excipiente pectus, ad
cujus basin tubus.

Brachia (præter bina tentacula pedunculata in
plerisque) octo, interius adspersa cirris
verrucosis.

Os inter brachia, terminale, corneum.

Lin. Syst. Nat. Gmel. p. 3149.

CHARACTER SPECIFICUS, &c.

Sepia corpore granulato, brachiis longissimis,
dorso oculisque appendiculis cutaneis.

Octopus corpore tuberculis sparsis granulato,
cotyledonibus crebris biserialibus.

Lamarck mem. hist. nat. Par. an. 7. p. 20.

Sepiæ speciem depinximus *Sepiæ octopodiæ* Linnæi
valde affinem, marium Europæorum incolam, descrip-
tamque a Montforto in historia, quam edidit, Mol-
luscorum; quæ scilicet pars est naturalis historiæ a
Buffono conscriptæ, a Sonnino autem continuatæ.
Extat ipsissimum specimen in Museo Principis Arau-
siaci, quod jam penes est Parisios. De vivendi modo,
cui præcipue assuescat animal, nihil pro certo comper-
tum est: crediderim tamen convenire mores moribus
Sepiæ octopodiæ,

THE
GRANULATED CUTTLE.

GENERIC CHARACTER.

Body fleshy, sheathing the breast, which is furnished at its base with a tubular opening.

Arms eight, and, in most species, two pedunculated tentacula, beset with numerous suckers or acetabula.

Head short : *Eyes* large : *Mouth* resembling a parrot's beak.


SPECIFIC CHARACTER.

Cuttle with granulated body, very long arms, and cutaneous appendages at the back and eyes.

Le Poulpe fraisé.

Monfort. Hist. Mollusq. 3. p. 5.

The present species of *Sepia*, which is much allied to the common eight-armed Cuttle-Fish, is a native of the European seas, and is described by Mons. Denys-Montfort in his *History of Mollusca*, (forming a part of the continuation of Mons. Sonnini's edition of Buffon's *Natural History*.) The specimen itself is preserved in the Museum of the Prince of Orange, now at Paris ; but nothing seems to be known relative to the particular history of the animal, which in its modes of life, in all probability, resembles the *Sepia octopodia*.



INDEX.


PL.
 764. *Acarus pectinatus*.
 748. *Alcyonium Schlosserian*.
 765. *Ampelis Maynana*.
 775. *Brachionus galeatus*.
 778. *Bodianus Apua*.
 742. *Caligus piscinus*.
 777. *Columba melanoptera*.
 734. *Cypræa testudinaria*.
 741. *Falco communis*.
 749. *Fringilla tristis*.
 743. *Gryllus viridissimus*.
 747. ——— *monstrosus*.
 763. ——— *morbillosus*.
 772. ——— *Dux*.
 759. *Hydrachna geographica*.
 760. *Labrus guttulatus*.
 762. ——— *Brasiliensis*.
 770. *Lacerta furcifrons*.
 779. *Libellula cærulata*.
 754. *Madrepore seriata*.
 767. *Millepora violacea*.
 736. *Monoculus cyprinaceus*.
 746. *Mullus maculatus*.
 757. *Numida cristata*.
 740. *Onchidium Typhæ*.
 744. *Papilio Antiopa*.
 756. ——— *Paphia*.
 735. *Phalæna imperatoria*.
 776. ——— *odora*.
 753. *Picus rufus*.
 737. *Psittacus coccineus*.
 769. ——— *incertus*.
 758. *Scarabæus auratus*.
 751. ——— *Goliathus*.
 745. *Scolopax Helias*.
 774. *Simia argentata*.
 780. *Sepia granulata*.
 768. *Scorpæna volitans*.
 739. *Sphinx Ficus*.
 752. ——— *Convolvuli*.
 750. *Sparus Pagrus*.
 755. ——— *fasciatus*.
 738. *Spongia palmata*.
 761. *Tanagra silens*.
 733. *Tinamus variegatus*.
 773. *Tetrao Francolinus*.
 766. *Trichoda triseta*.
 771. *Vorticella Cyathus*.


PL.
 748. *Alcyonium Schlosserian*.
 778. *Bodian Apua*.
 775. *Brachionus galeated*.
 758. *Beetle golden*.
 751. ——— *Goliath*.
 744. *Butterfly Antiopa*.
 756. ——— *Paphia*.
 742. *Caligus fish*.
 765. *Chatterer silken-feathered*.
 734. *Cowry tortoise*.
 780. *Cuttle granulated*.
 770. *Chamæleon fork-headed*.
 779. *Dragon-Fly blue-tipped*.
 741. *Falcon common*.
 749. *Finch golden*.
 773. *Francolin*.
 759. *Hydrachna variegated*.
 760. *Labrus guttulatus*.
 762. ——— *Brasilian*.
 743. *Locust green*.
 763. ——— *granulated*.
 747. ——— *spiral-winged*.
 772. ——— *imperial*.
 737. *Lory Indian*.
 754. *Madrepore seriated*.
 767. *Millepore violaceous*.
 764. *Mite pectinated*.
 736. *Monoculus carp*.
 735. *Moth imperial*.
 776. ——— *sable*.
 774. *Monkey silver-haired*.
 740. *Onchidium limacine*.
 777. *Pigeon black-capped*.
 757. *Pintado crested*.
 769. *Parrot blue-green*.
 745. *Snipe Caurale*.
 768. *Scorpæna flying*.
 750. *Sparus Pagre*.
 755. ——— *fasciated*.
 739. *Sphinx Fig-tree*.
 752. ——— *Bindweed*.
 738. *Sponge palmated*.
 746. *Surmullet spotted*.
 733. *Tinamou variegated*.
 761. *Tanager silent*.
 766. *Trichoda three-bristled*.
 771. *Vorticella Cup*.
 753. *Woodpecker rufous*.

MA
570-15
L-33

~~MA
570-15
L-33~~

RB
500-9
553-4





RB
500.9
S534

