HIV: now there is a treatment that may prevent HIV infection after the virus has entered the body: PEP post exposure prophylaxis: treatment only works if started within 72 hours of exposure: 1 month course of anti-HIV medication ... / Terrence Higgins Trust.

### Contributors

Terrence Higgins Trust

## **Publication/Creation**

London: Terrence Higgins Trust, [2004]

## **Persistent URL**

https://wellcomecollection.org/works/axgsa4fr

### License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



# PEP

**POST EXPOSURE PROPHYLAXIS** 

TREATMENT ONLY WORKS IF STARTED WITHIN 72 HOURS OF EXPOSURE

1 MONTH COURSE OF ANTI-HIV MEDICATION

CAUTION: PEP IS NOT A SUBSTITUTE FOR CONDOMS

SIDE EFFECTS CAN INCLUDE DIARRHOEA, NAUSEA AND SEVERE HEADACHES

> Terrence HIGGINS Trust

> > For information you need to know about PEP, visit www.tht.org.uk/pep or call THT DIFECT 0845 12 21 200