

A translation of the pharmacopoeia of the Royal College of Physicians of London, 1836. With notes and illustrations / By Richard Phillips.

Contributors

Royal College of Physicians of London.
Phillips, Richard, 1778-1851.

Publication/Creation

London : S. Highley, 1837.

Persistent URL

<https://wellcomecollection.org/works/txtwervu>

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



40898/B


Last prelim leaf (contents with
corrigenda on verso) and
pp. 1-16 wanting



40898/B

PHARMACOPOEIAS:

London [RCP]



Digitized by the Internet Archive
in 2018 with funding from
Wellcome Library

<https://archive.org/details/b29340998>

A

54800

TRANSLATION
OF THE
PHARMACOPŒIA

OF THE
ROYAL COLLEGE OF PHYSICIANS
OF LONDON,

1836.

WITH NOTES AND ILLUSTRATIONS.

BY
RICHARD PHILLIPS, F.R.S. L. & E. &c.

LECTURER ON CHEMISTRY AT ST. THOMAS'S HOSPITAL.

By Permission.

LONDON :
S. HIGHLEY, 32, FLEET STREET.

1837.



PRINTED BY RICHARD TAYLOR,
RED LION COURT, FLEET STREET.

7/5
Coxd

TO

CLEMENT HUE, M.D.

FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS,
AND SENIOR PHYSICIAN TO ST. BARTHOLOMEW'S HOSPITAL,
&c. &c.

MY DEAR SIR,

INDUCED by various motives I
inscribe this Translation to you. I have witnessed the
unremitting zeal which directed your scientific attain-
ments to the improvement of the original *Work*, and the
candour with which you appreciated the suggestions of
others. I am moreover desirous of acknowledging the
obligation which I am under to you, for the assistance
rendered me on many important occasions connected
with this publication.

I remain,

My dear Sir,

Yours faithfully,

RICHARD PHILLIPS.

ST. THOMAS'S HOSPITAL,
December, 1836.

ADVERTISEMENT.

THE Royal College of Physicians having appointed a Committee to revise their Pharmacopœia, it may perhaps be allowed me to state, that I have, at their request, either conducted or inspected the preparation of almost every medicine which has been introduced, and in many cases repeated the processes which the Pharmacopœia already contained. While thus engaged, I have repeatedly received highly useful information from Mr. Hennell of Apothecaries' Hall, and I am anxious to express my sense of its value.

In preparing the remarks which accompany this Translation my attention has been directed to two classes of persons :—First, those who may have been for some time engaged in the practice of Physic, but who, not having watched the rapid progress which Chemistry has made within a few years, are imperfectly acquainted with the important changes which it has produced by the introduction of new medicines from various and unexpected sources.

The other class to which I allude is the numerous one of Medical Students ; to these I have found, by no inconsiderable experience, that concise descriptions of the chemical changes which occur during the preparation of medicines have been extremely useful. They who know how small the portion of time is which the medical pupil has at his disposal for the acquirement of chemical and pharmaceutical knowledge, will readily admit the propriety of assisting his progress by familiar modes of illustration. With this latter view I have made much use of diagrams, and I shall offer a few words in explanation of the method I have adopted in framing them. It is to be understood, that the new compounds formed during a process, or constituents assuming a fresh state, are denoted by being printed in italics: thus, a solution of sulphate of soda being mixed with one of nitrate of lime, the new compounds formed are *sulphate of lime* and *nitrate of soda*, and supposing one of the resulting substances to be a solid, that is generally placed at the bottom of the diagram. When nitric acid is added to carbonate of lime, the carbonic acid assumes a fresh state, and in this case it is thus described—carbonic acid *gas* ; the only change which it undergoes being from the state of solid combination, to that of an uncombined elastic fluid.

In the alphabetical arrangement now adopted by the College, under the head of *Præparata à Potassio*, Liquor Potassæ correctly precedes Potassæ Carbo-

nas; but my object being that of explaining the nature of the chemical compounds, and of describing the causes upon which their difference depends, I have on this occasion, as well as in some other cases, altered the order, and have treated of carbonate of potash before the solution, because a knowledge of the properties of the carbonate must be acquired before the separation of its carbonic acid by the action of the lime can be understood.

In my translation of the late Pharmacopœia I mentioned the properties which indicated the qualities of various medicines, and described the means which might be adopted for the detection of impurities and adulteration. This very useful appendage to a Pharmacopœia has now been introduced by the College, and to their original directions for this purpose I have added such remarks as it appeared to me might further the purpose of their introduction. The tests which have been proposed are the more necessary, because the College no longer insist that the medicines which they direct, should be prepared exactly in the mode prescribed, provided they will stand the trials of their purity, to which it is proposed they shall be submitted.

I have in many cases given the crystalline forms of various salts; and these, with scarcely any exception, have been derived from the communications of my friend Mr. Brooke to the *Annals of Philosophy*. I have also added the symbols of most chemical compounds, not that I would be under-

stood as admitting their utility, but in compliance with the practice of some of the most eminent chemists of the present day. From numerous plans which have been proposed I have selected the symbols given by Professor Brande in his Manual of Chemistry, and those of Berzelius: the latter are adopted, generally, without any other alterations than the slight ones sanctioned by Professor Turner in his Elements of Chemistry; and I have on many other occasions advantageously consulted both the excellent works now mentioned.

Well knowing how necessary it is that the student should be acquainted with the powers and doses of preparations, I have generally given an account of them; but not being a medical practitioner, the best authorities on the subject have been consulted and quoted, and I need do no more to inspire confidence in this statement, than to observe that I am chiefly indebted for them to Dr. Paris and Dr. Hue; and I have great pleasure in acknowledging their friendly assistance on various occasions connected with this translation.

R. P.

P R E F A C E.

CONSIDERING how much has been effected in the present age towards extending and almost re-creating the arts on which the science of Medicine depends, it will excite no surprise that the Pharmacopœia published twelve years ago, should have become somewhat imperfect and obsolete. The College of Physicians had long ago determined on revising and reforming it, but many circumstances prevented its completion. For not only is it always a difficult undertaking to collect and justly to appreciate the opinions of a number of persons, but it was frequently our duty to examine the most trivial points, and carefully to consider the rejection or improvement of what had become antiquated, and the admission of what was new, and to judge of the best mode of carrying these alterations into effect. On these points we have been much engaged. However, but little progress had been made when a fresh difficulty arose; as we were desirous of compiling, not as before, a London Pharmacopœia, but one which should include Scotland and Ireland with England, it was requisite to consult with the Fellows of both Colleges; and as, on account of the great distance, this was with difficulty accomplished,

we were at length constrained to abandon the negotiation which had been commenced.

This is not a proper occasion for stating by what individuals or by what means the boundaries of Science have been enlarged; though the College of Physicians claim as members, many among those who have greatly promoted its extension. When, however, the knowledge of all Nature had made such rapid advances, it would have been unworthy of Medicine to have remained stationary in that department which relates to remedies.

It will here be proper briefly to explain what we have effected. We have not been satisfied with describing a number of medicines without thoroughly examining them all; we have especially subjected the chemical preparations, which have yearly increased in number and importance, to the test of rigid experiment. If some of these should appear but little approved by experience, we have nevertheless included them in this work, that if any one should think fit to administer them, he might have all accurately prepared. Lest anything further should be wanting to complete the work, or to the security of the sick, we have now for the first time added short notes, by which the purity of the medicines required may be for the most part easily ascertained, sufficiently, at least, for the use of physicians and students in medicine, if not always for the exact analysis of chemists.

We have usually omitted to mention the methods of preparing such medicines as may be had sufficiently pure and without trouble from those who deal in them; though we certainly had rather that some of them should be prepared according to our directions, than trust to the attention or inattention of others. And although more beautiful crystals and of brighter colour are elsewhere produced on account of the more abundant materials

which are employed, these are circumstances which do not at all contribute towards the alleviation of disease.

There is also this novelty in our Pharmacopœia, that we have resolved to adopt an alphabetical arrangement; and if to some this plan should appear unsystematic, we wish to contribute rather to the convenience of the physician, than to consult the opinions of philosophers, or indulge the fancy of theorists. Some chemical substances are excepted from this arrangement, which, having a general connexion, and being united, as it were, by some kind of affinity, could not be separated with propriety.

In the present day it is scarcely necessary to discuss the change of names; for though all admit that it ought not to be done inconsiderately, yet it was to be feared, lest the avoidance of the error likely to arise from the change should lead to a greater one: we are persuaded that the name which the most eminent Professors of the science have bestowed upon a substance will eventually be the most certain and permanent. We have therefore determined to follow in their footsteps, and put up for a short time with inconvenience, if any should arise, or learn to speak barbarously, rather than incur the danger of becoming obscure from employing a doubtful or ill-defined expression. If we have ourselves invented one or two, we have done so in accordance with these principles.

Such are the observations with which we have thought proper to commence. We cannot hope to afford satisfaction to all, or that we have performed everything which we could have wished. Some, perhaps, will think our Work too copious; others too brief and limited. It has at any rate been our endeavour, if we could not please all, at least to benefit many; and we shall be gratified

if studious youths and the compounders of medicine, whether learned or unlearned, gather any fruit from our labour. But we have especially aimed at furnishing the household of the physician and consulting the welfare of the sick, that the former may more certainly and safely contend with diseases, and the latter be sooner restored to the wished-for enjoyment of health.

Scammonium, <i>Scammony.</i>	Convolvulus Scammonea, <i>The Gum-resin.</i>
Scilla, <i>Squill.</i>	Scilla maritima, <i>The fresh Bulb.</i>
Scoparius, <i>Broom.</i>	Cytisus Scoparius (DE CANDOLLE), <i>The fresh Tops.</i>
Senega, <i>Senega.</i>	Polygala Senega, <i>The Root.</i>
Senna, <i>Senna.</i>	Cassia lanceolata (DE CANDOLLE), <i>The Leaves.</i>
	Cassia obovata (DE CANDOLLE), <i>The Leaves.</i>
Serpentaria, <i>Serpentary.</i>	Aristolochia Serpentaria, <i>The Root.</i>
Sevum, <i>Suet.</i>	Ovis Aries, <i>The Suet.</i>
Simaruba, <i>Simaruba.</i>	Simaruba officinalis (DE CANDOLLE), <i>The Bark of the Root.</i>
Sinapis, <i>Mustard.</i>	Sinapis nigra, <i>The Seeds.</i>
Sodæ Acetas, <i>Acetate of Soda.</i>	
Sodæ Carbonas impura, <i>Impure Carbonate of Soda.</i>	
Sodæ Phosphas, <i>Phosphate of Soda.</i>	
Sodii Chloridum, <i>Chloride of Sodium.</i>	
Spigelia, <i>Indian Pink.</i>	Spigelia marilandica, <i>The Root.</i>
Spiritus rectificatus, <i>Rectified Spirit.</i>	Spiritus, <i>Its specific gravity is .838.</i>
Spiritus tenuior, <i>Proof Spirit.</i>	Spiritus, <i>Its specific gravity is .920.</i>
Spiritus Vini Gallici, <i>Spirit of French Wine,</i> <i>(Brandy.)</i>	Spiritus, <i>Distilled from French</i> <i>Wine.</i>

- Stannum,
Tin.
- Staphisagria,
Stavesacre.
- Stramonii Folia,
Leaves of Stramonium.
- Stramonii Semina,
Seeds of Stramonium.
- Styrax,
Storax.
- Succinum,
Amber.
- Sulphur,
Sulphur.
- Tabacum,
Tobacco.
- Tamarindus,
Tamarind.
- Taraxacum,
Dandelion.
- Terebinthina Canadensis,
Canadian Turpentine.
- Terebinthina Chia,
Chio Turpentine.
- Terebinthina vulgaris,
Common Turpentine.
- Terebinthinæ Oleum,
Oil of Turpentine.
- Testæ,
Shells.
- Tiglii Oleum,
Oil of Croton.
- Tormentilla,
Tormentil.
- Toxicodendron,
Sumach.
- Delphinium Staphisagria,
The Seeds.
- Datura Stramonium,
The Leaves.
- The Seeds.*
- Styrax officinale,
The Balsam.
- Sulphur (*sublimatum*).
- Nicotiana Tabacum,
The dried Leaves.
- Tamarindus indica,
The Pulp.
- Leontodon Taraxacum,
The Root.
- Pinus Balsamea,
The liquid Resin.
- Pistachia Terebinthus,
The liquid Resin.
- Pinus sylvestris,
The liquid Resin.
- The Oil distilled from the Resin.*
- Ostrea edulis,
The Shells.
- Croton Tiglium,
The Oil expressed from the Seeds.
- Potentilla Tormentilla
(DE CANDOLLE),
The Root.
- Rhus Toxicodendron,
The Leaves.

Tragacantha,	Astragalus verus (OLIVIER, <i>Voy. dans l'Emp. Ottom.</i>), <i>The concrete Juice.</i>
<i>Tragacanth.</i>	
Tussilago, <i>Coltsfoot.</i>	Tussilago Farfara.
Valeriana, <i>Valerian.</i>	Valeriana officinalis (<i>sylves- tris</i>), <i>The Root.</i>
Veratrum, <i>White Hellebore.</i>	Veratrum album, <i>The Root.</i>
Vinum Xericum, <i>Sherry Wine.</i>	
Ulmus, <i>Elm.</i>	Ulmus campestris, <i>The Bark.</i>
Uva, <i>Raisins.</i>	Vitis vinifera, <i>The dried Berries with the stones taken out.</i>
Uva ursi, <i>Whortleberry.</i>	Arctostaphylos Uva ursi (SPRENGEL, <i>Syst. Veget.</i>), <i>The Leaves.</i>
Zincum, <i>Zinc.</i>	
Zingiber, <i>Ginger.</i>	Zingiber officinalis (ROSCOE, <i>in Act. Soc. Linn.</i>), <i>The Rootstock.</i>

NOTES.

It has been deemed proper to add short notes, relating chiefly to the chemical preparations, by which their purity may, as nearly as possible, be ascertained. This, for the most part, is less necessary with vegetable and animal substances, and is also attended with more difficulty. For although the peculiar character of each plant and animal is sufficiently defined in books on botany and zoology, yet the extracts from them and the weaker preparations are frequently so much altered in taste, colour, and smell, that they cannot be distinguished by any certain sign, or be briefly described.

ACETUM. *Vinegar*.—A yellowish liquor, of a peculiar odour, a fluidounce of which is saturated by a drachm of crystallized carbonate of soda. Solution of chloride of barium being added, the sulphate of barytes precipitated does not exceed 1.14 grain. Hydrosulphuric acid being added, its colour is not altered.

Remarks.—The strongest vinegar contains 5 per cent. of real acetic acid; usually it does not exceed 4.6 per cent.; a fluidounce weighs about 446 grains, saturating 58 grains of carbonate of soda; and two grains of it, making up the drachm, are allowed for saturating the sulphuric acid permitted to be mixed with the vinegar, and for decomposing the sulphates of the water used in vinegar-making: the 1.14 grain of sulphate of barytes is derived from the same sources, and proves that too much sulphuric acid has not been used. The non-action of hydrosulphuric acid demonstrates the absence of most metallic oxides.

ACETUM DESTILLATUM. *Distilled Vinegar*.—Totally vaporized by heat. No precipitate is formed on the addition of acetate of lead, nitrate of silver, nor iodide of potassium. Neither hydrosulphuric acid nor ammonia alters its colour. After the digestion of a plate of silver in it, hydrochloric acid occasions no precipitation. Thirteen grains of the crystals of carbonate of soda are saturated by 100 grains of distilled vinegar.

Remarks.—The total evaporation shows that no solid impurity is dissolved in the vinegar. The non-precipitation by acetate of lead proves the absence of sulphuric acid; by nitrate of silver,

that of hydrochloric acid; by iodide of potassium, that of lead. The non-action of hydrosulphuric acid and of ammonia proves the absence of metallic admixture. If adulterated with nitric acid, silver would be dissolved by digestion in it, and afterwards precipitated as a chloride by hydrochloric acid. The quantity of carbonate of soda to be saturated indicates the presence of 4.6 per cent. of real acetic acid.

ACIDUM ACETICUM. *Acetic Acid.*—The specific gravity of this acid is 1.048. Eighty-seven grains of crystals of carbonate of soda are saturated by 100 grains of this acid. The acid when saturated with carbonate of soda and evaporated, yields crystals of acetate of soda. Other tests agree with those of the preceding preparation.

Remark.—The saturating power of this acid shows that it contains 30.7 per cent. of real acetic acid.

ACIDUM ARSENIOSUM. *Arsenious Acid.*—It is entirely sublimed when heated. Mixed with charcoal and exposed to heat, it emits an alliaceous smell. It is dissolved by boiling water; and hydrosulphuric acid, when added, throws down a yellow precipitate, and lime-water yields a white one.

Remarks.—The precipitate thrown down by hydrosulphuric acid is sesquisulphuret of arsenic, and that by lime-water is arsenite of lime.

ACIDUM BENZOICUM. *Benzoic Acid.*—When cautiously heated it totally evaporates with a peculiar odour. It is sparingly soluble in water, but plentifully in rectified spirit. It is entirely dissolved by solution of potash or by lime-water, and is precipitated by hydrochloric acid.

ACIDUM CITRICUM (crystalli). *Citric Acid (crystals).*—This acid is soluble in water; what is precipitated from the solution by acetate of lead, is dissolved by nitric acid. No salt of potash, except the tartrate, is precipitated by solution of citric acid. It is totally dissipated in the fire.

Remarks.—Any precipitate obtained by acetate of lead, which is insoluble in nitric acid, may be regarded as sulphate of lead, and would denote the presence of sulphuric acid or a sulphate in the citric acid. If the citric acid contained any tartaric acid, that would decompose other salts of potash besides the tartrate, and a crystalline and difficultly soluble precipitate of bitartrate of potash would be formed. As citric acid consists of oxygen, hydrogen and carbon, all of which are dissipated at a red heat, any substance remaining after ignition is an impurity.

ACIDUM HYDROCHLORICUM. *Hydrochloric Acid.*—Colourless ; entirely vaporized by heat. When mixed with distilled water, neither chloride of barium, nor ammonia, nor the sesquicarbonate of ammonia throws down anything. Strips of gold, even when heated in it, are not acted upon by it. It does not destroy the colour of solution of sulphate of indigo. Its specific gravity is 1.16. One hundred and thirty-two grains of crystals of carbonate of soda are saturated by 100 grains of this acid.

Remarks.—The total evaporation by heat proves that no fixed substance is dissolved in the acid. The non-action of chloride of barium proves that no sulphuric acid is present, or sulphate of barytes would be precipitated. Ammonia or the sesquicarbonate of ammonia would detect the presence of most metals and earths by precipitating, and, in some cases, by afterwards redissolving them. Gold, even when heated in hydrochloric acid, is not dissolved unless chlorine be present, and then it is taken up, and may be precipitated from the solution by chloride of tin ; the precipitate is of a purple or dark colour. If chlorine be present it will also destroy the colour of the solution of indigo. When 100 parts of this acid saturate 132 of carbonate of soda, it contains rather more than 33 per cent. of hydrochloric acid gas.

ACIDUM HYDROCYANICUM DILUTUM. *Dilute Hydrocyanic Acid.*—Free from colour ; goes off in vapour by heat, exhaling its peculiar odour. It turns litmus of a slight fugacious red colour ; hydrosulphuric acid, when added, does not discolour it. One hundred grains of this acid, when solution of nitrate of silver is added, precipitate 10 grains of cyanide of silver, which are readily dissolved by boiling nitric acid. If the iodo-cyanide of potassium and mercury when mixed with the hydrocyanic acid is reddened, it contains some other acid. In 100 grains of this diluted acid there are contained 2 grains of real hydrocyanic acid ; and to this standard, in whatever mode it is distilled, we direct it should be reduced.

Remarks.—Its total evaporation shows the absence of fixed impurity. If the hydrocyanic acid redden litmus-paper strongly and permanently, then some other acid is mixed with it : the absence of most metallic salts is denoted by the non-action of hydrosulphuric acid. If the hydrocyanic acid contain hydrochloric acid, then the precipitate formed by nitrate of silver, being chloride of silver, is insoluble in the nitric acid. Any acid mixed with the hydrocyanic acid decomposes the iodo-cyanide of potassium and mercury, and forms biniodide of mercury, which is of a red colour.

ACIDUM NITRICUM. *Nitric Acid.*—By heat it wholly passes off in vapour. When mixed with distilled water, neither nitrate of

silver nor chloride of barium throws down anything. Its specific gravity is 1.50. About 217 grains of the crystals of carbonate of soda are saturated by 100 grains of this acid.

Remarks.—The total evaporation proves that no fixed impurity is held in solution by the acid. Nitrate of silver gives no precipitate when chlorine or its compounds are absent; chloride of barium gives a precipitate of sulphate of barytes when sulphuric acid or a sulphate is present. Two hundred and seventeen grains of carbonate of soda are equivalent to about 81 grains of real nitric acid.

ACIDUM PHOSPHORICUM DILUTUM. *Diluted Phosphoric Acid.*—Chloride of barium or nitrate of silver being added, whatever is thrown down is readily dissolved by nitric acid. Strips of copper and silver are not at all acted upon by it, nor is it coloured when hydrosulphuric acid is added. Its specific gravity is 1.064; 42 grains of carbonate of soda are saturated by 100 grains of this acid, and nothing is thrown down.

Remarks.—The absence of sulphuric acid and of a sulphate is proved by there being no precipitate yielded by chloride of barium, which is insoluble in nitric acid; that no hydrochloric acid nor any chloride is held in solution, is proved by nitrate of silver giving no precipitate which is insoluble in nitric acid. The quantity of carbonate of soda neutralized by 100 grains of this acid shows that it contains $10\frac{1}{2}$ per cent. of real phosphoric acid; and when the carbonate of soda gives no precipitate, no phosphate insoluble in water is dissolved by the phosphoric acid.

ACIDUM SULPHURICUM. *Sulphuric Acid.*—It is free from colour. Its specific gravity is 1.845. What remains after the acid has been distilled to dryness does not exceed the four hundredth part of its weight. Diluted sulphuric acid is scarcely coloured by hydrosulphuric acid.

Remarks.—Its being colourless shows that no carbonaceous matter has fallen into and been decomposed by the acid. The small quantity of matter left after distillation to dryness is sulphate of lead, and generally a little sulphate of potash. The non-production of colour by hydrosulphuric acid, as well as the small quantity of matter left by distillation, show that the acid contains no important metallic impregnation.

ACIDUM TARTARICUM (*crystalli*). *Tartaric Acid (crystals).*—Totally soluble in water. The solution throws down bitartrate of potash from any neutral salt of potash. Whatever is precipitated from this solution by acetate of lead, is dissolved by diluted nitric acid.

Remarks.—The insolubility of any precipitate produced in the

solution of tartaric acid by acetate of lead, in dilute nitric acid would show that the tartaric acid contains either sulphuric acid or a sulphate.

ACONITINA. *Aconitina*.—An alkali prepared from the leaves and roots of Aconite. It is very soluble in sulphuric æther, less in alcohol, and very slightly in water. It is totally consumed in the fire, no salt of lime remaining. This substance possessing strong power, is not to be rashly employed.

Remarks.—As it consists of oxygen, hydrogen, azote and carbon, it is of course entirely destructible by fire, and any remaining substance is an impurity.

ADEPS. *Lard*.—Is not to be used without being carefully washed with water.

ÆRUGO. *Verdigris*.—May be partly dissolved in water, and is almost entirely soluble either in ammonia, or with the assistance of heat, in diluted sulphuric acid.

ÆTHER SULPHURICUS. *Sulphuric Æther*.—Its specific gravity is 0.750. What is sold fluctuates between 0.733 and 0.765. It totally evaporates in the air. It reddens litmus slightly: it combines sparingly with water; for example, in the proportion of a fluidounce to half a pint, and remains limpid.

Remarks.—The specifically lighter it is the sooner it evaporates, and it contains the less alcohol or water. If it reddens litmus strongly, it has been either improperly prepared or too long kept. The more perfect it is the less water or alcohol it contains, and the less soluble it is in water.

ALCOHOL. *Alcohol*.—The specific gravity of this is .815; it is free from colour; when heated it evaporates; it combines with water and with æther; it tastes and smells like wine.

ALUMEN (*crystallinum*).—**ALUM** (*crystalline*).—It is entirely soluble in water. From the solution, ammonia or potash, when added, throws down alumina free from colour; which again dissolves when the potash is added in excess.

Remarks.—The crystallization in octohedrons and perfect solubility in water show that there is no uncombined earthy matter; and the precipitation by ammonia and potash, the solubility of the precipitate in excess of the latter, and its being colourless, show that pure alumina has been precipitated.

AMMONIÆ LIQUOR. *Solution of Ammonia*.—By heat it totally evaporates in evanescent alkaline vapours, as shown by turmeric.

It gives no precipitate with lime-water. When saturated with nitric acid neither sesquicarbonate of ammonia nor nitrate of silver throws down anything. The specific gravity of this solution is 0.960.

Remarks.—Pure water remains after the expulsion of the ammoniacal gas. If lime-water give no precipitate with solution of ammonia, it shows that it contains no carbonic acid; and if neither sesquicarbonate of ammonia nor nitrate of silver give any precipitate when it has been saturated with nitric acid, it proves that no earthy matter, hydrochloric acid, nor any chloride is present.

AMMONIÆ LIQUOR FORTIOR. *Stronger Solution of Ammonia.*—The specific gravity of this is .882. This solution may be reduced to the strength of Liquor Ammoniaë by adding to every fluidounce of it three fluidounces of distilled water.

Its properties are similar to those of Ammoniaë Liquor.

AMMONIÆ ACETATIS LIQUOR. *Solution of Acetate of Ammonia.*—It is not coloured by the addition of hydrosulphuric acid, nor is anything precipitated by nitrate of silver or chloride of barium. The water being evaporated, the residue yields ammonia, and is dissipated by heat.

Remarks.—When the vinegar has been improperly distilled and condensed in a metallic worm, it usually contains some metallic oxide, which is detected by hydrosulphuric acid. Nitrate of silver gives a precipitate of chloride with hydrochloric acid, and chloride of barium yields sulphate of barytes with sulphuric acid, if it be present. Acetate of ammonia is totally decomposed and dissipated by heat, and if there be any residue after its operation, it is an impurity.

AMMONIÆ HYDROCHLORAS (*crystallina*). *Hydrochlorate of Ammonia (crystalline).*—Translucent; it is sublimed by heat, totally dissolved by water. It changes the colour of litmus slightly red. Chloride of barium throws down nothing. Potash or lime being added to it, ammonia is evolved.

AMMONIÆ SESQUICARBONAS (*crystallina*). *Sesquicarbonate of Ammonia (crystalline).*—Translucent, but falls to powder in the air; it is totally dissipated by heat. It is entirely soluble in water; it changes the colour of turmeric. Nitric acid being added to it to saturation, nothing is thrown down either by chloride of barium or nitrate of silver.

Remarks.—When it has lost its transparency it is less pungent to the smell and less active as a medicine, being partially converted into bicarbonate of ammonia: if anything remain after the

application of heat or the action of water, it is an impurity. The non-action of nitrate of silver and chloride of barium proves the absence of hydrochloric and sulphuric acid.

ANTIMONII OXYSULPHURETUM. *Oxysulphuret of Antimony*.—Totally soluble in nitro-hydrochloric acid, emitting hydrosulphuric acid.

Remarks.—By boiling in a solution of bitartrate of potash it loses about 12 per cent., which is protoxide of antimony.

ANTIMONII POTASSIO-TARTRAS (*crystalli*). *Potassio-tartrate of Antimony (crystals)*.—Totally soluble in water, no bitartrate of potash remaining in the vessel, and hydrosulphuric acid being added, a reddish coloured precipitate is obtained. Neither chloride of barium nor nitrate of silver being added to the solution, throws down anything. Nitric acid throws down a precipitate, which is dissolved by an excess of it.

Remarks.—The crystalline form, and solubility in a moderate quantity of water, prove the absence of bitartrate of potash uncombined with oxide of antimony. Hydrosulphuric acid precipitates red hydrated sesquisulphuret of antimony. The non-action of chloride of barium proves the absence of sulphuric acid and sulphates, and that of nitrate of silver, the absence of chlorine and chlorides. Nitric acid precipitates oxide of antimony, which an excess of it redissolves.

ANTIMONII SESQUISULPHURETUM (*striatum*). *Sesquisulphuret of Antimony (striated)*.—With heat it is totally dissolved by hydrochloric acid. From the acid in which it is boiled, a white precipitate is thrown down by distilled water; from the strained liquor hydrosulphuric acid afterwards throws down a reddish-coloured substance.

Remarks.—Hydrochloric acid would dissolve some lead, if any of that metal in the state of sulphuret existed in the sesquisulphuret of antimony: this would remain in solution after the precipitation of the white substance by water, and hydrosulphuric acid would then give a dark-coloured precipitate with the strained solution instead of the reddish one, derived from a small quantity of antimony not thrown down by water. If also any copper had been dissolved, the same appearance would be induced by hydrosulphuric acid as with lead.

ARGENTUM. *Silver*.—It is totally dissolved by diluted nitric acid. This solution on the addition of chloride of sodium throws down a precipitate, which an excess of ammonia dissolves, and it should be free from colour. The chloride of silver being removed, and hydrosulphuric acid added to the solution, it is not

coloured by it, and nothing is thrown down. The specific gravity of silver is 10.4.

Remarks.—If the silver contain gold, which is frequently the case, it remains undissolved as a dark-coloured powder; if lead, it will be dissolved by the nitric acid; and if the quantity be considerable, it will be precipitated with the silver, also in the state of a chloride by chloride of sodium, and this, unlike chloride of silver, would not be dissolved by excess of ammonia. When the chloride of silver is removed, the liquor may contain copper and some lead; with these hydrosulphuric acid would give dark-coloured precipitates.

ARGENTI NITRAS. *Nitrate of Silver.*—It is originally white, but blackens by exposure to light. It is entirely soluble in water. Copper put into the solution precipitates silver; its other properties are as above detailed respecting silver.

Remarks.—If silver containing copper be used, the nitrate is greenish by the presence of subnitrate, or blackish on account of the oxide of copper which it contains. Chloride of sodium should give a white precipitate totally soluble in excess of ammonia, which it will not do if it contain chloride of lead. The solution after precipitation and the removal of the chloride of silver should give no precipitate, nor suffer any discoloration by hydrosulphuric acid; if it should, copper or lead, or both, may be present.

ARGENTI CYANIDUM. *Cyanide of Silver.*—By heat it yields cyanogen, and is reduced to silver.

Remarks.—If pure, the residual silver will be totally dissolved by nitric acid, and the solution will exhibit the properties above described respecting silver.

BARYTÆ CARBONAS. *Carbonate of Barytes.*—Totally soluble in diluted hydrochloric acid. This solution, on the addition of ammonia or hydrosulphuric acid, does not give any precipitate, and it remains colourless: when more sulphuric acid is added than is necessary to saturation, nothing is afterwards thrown down by carbonate of soda.

Remarks.—If totally soluble in hydrochloric acid, it contains no sulphate of barytes; if ammonia occasion no precipitate, it contains no alumina, oxide of lead, nor peroxide of iron; if hydrosulphuric acid occasion no colour, it is free from copper or lead; if carbonate of soda throw down nothing from the solution from which the sulphate of barytes has been precipitated by sulphuric acid, it contains no lime.

BISMUTH. *Bismuth.*—It is dissolved by diluted nitric acid; when subnitrate of bismuth is precipitated from this solution by

ammonia, the liquor is free from colour. Its specific gravity is 9.8.

Remarks.—If the bismuth contain copper, a blue-coloured solution remains after the precipitation of the bismuth by ammonia.

BISMUTHI TRISNITRAS. *Trisnitrate of Bismuth.*—It is soluble in nitric acid without effervescence. Diluted sulphuric acid being added to the solution, nothing is thrown down.

Remarks.—If it possess these properties, it contains no carbonate whatever, nor any oxide of lead.

BORAX (*crystalli*). *Borax* (*crystals*).—Totally soluble in water. Sulphuric acid throws down scaly crystals from the solution. These dissolved in alcohol burn with a green-coloured flame.

Remark.—The crystals are boracic acid, separated from the soda of the borax by its superior affinity for sulphuric acid; sulphate of soda remains in solution.

BROMINIUM. *Bromine.*—Evaporates at a gentle heat, with an acrid smell. It is sparingly soluble in water, more in rectified spirit, and most in æther. Its specific gravity is 3.0.

CALAMINA. *Calamine.*—Almost entirely soluble in diluted sulphuric acid, emitting a few bubbles of carbonic acid, unless it has been previously burnt. The solution, when ammonia or potash is added to it, gives a precipitate, which either of them added in excess dissolves.

Remarks.—If soluble in sulphuric acid, the calamine can contain but little carbonate of lime. The sulphuric solution should be colourless, and remain so, after the addition of the excess of ammonia; if blue, copper is present; if it contain iron, both ammonia and potash throw down the oxide, which neither of them redissolves when added in excess.

CALCII CHLORIDUM. *Chloride of Calcium.*—Free from colour; slightly translucent; hard and friable; totally soluble in water: the solution gives no precipitate on the addition of ammonia or chloride of barium, nor when diluted with much water, with ferrocyanide of potassium.

CALCIS HYDRAS. *Hydrate of Lime.*—Dissolves in dilute hydrochloric acid without effervescence. Ammonia added to the solution throws down nothing.

Remarks.—The solubility in dilute hydrochloric acid without effervescence proves the absence of carbonic acid, that the lime has been well burnt, and that no silica is present. If the

solution gives no precipitate with ammonia, it contains neither oxide of iron nor alumina.

CALX. Lime.—Water being added it cracks and falls to powder. Its other properties are as above mentioned.

Remark.—Such portions as do not slack on the addition of water are insufficiently burnt; and when put into dilute muriatic acid effervesce, on account of the undecomposed carbonate of lime which they contain.

CALX CHLORINATA. Chlorinated Lime.—Dissolves in dilute muriatic acid, emitting chlorine.

Remark.—The chlorine gas is recognised by its colour, peculiar odour, and power of destroying vegetable and animal colour.

CARBO ANIMALIS (*purificatus*). Animal Charcoal (*purified*).—Emits no bubbles on the addition of hydrochloric acid; nor is anything thrown down from the acid either by ammonia or the sesquicarbonate of ammonia.

Remarks.—The purification is effected by means of hydrochloric acid, which should dissolve all the carbonate and phosphate of lime; if any of the former remain, it will dissolve with effervescence on the addition of hydrochloric acid, and the solution give a precipitate with sesquicarbonate of ammonia; if phosphate of lime be dissolved by the hydrochloric acid, the solution gives a precipitate both with ammonia and the sesquicarbonate of ammonia, which is phosphate of lime.

CORNU. Horn.—After it has been well burnt is almost entirely dissolved by nitric acid; then lime is separated by oxalate of ammonia; and phosphoric acid is precipitated by nitrate of lead.

Remarks.—The phosphate of lime, of which the horn after burning almost entirely consists, is dissolved by the nitric acid; oxalate of lime is formed on adding oxalate of ammonia to the solution, which being insoluble in water is precipitated. Phosphate of lead results from the union of the phosphoric acid with the oxide of lead of the nitrate when added to the nitric solution, and this also being insoluble in water, it is precipitated.

CREASOTON. Creasote.—Oleaginous; colourless; its smell peculiar; translucent; boils at 397° . Does not congeal at -50° . Soluble in acetic acid.

CRETA. Chalk.—Totally soluble in dilute hydrochloric acid with effervescence. From this solution, after it has been boiled, when ammonia is dropped in, it throws down nothing.

Remarks.—If totally soluble in hydrochloric acid it contains no silica; and if the solution gives no precipitate with ammonia, it is free from alumina and oxide of iron.

CUPRI SULPHAS (*crystalli*). *Sulphate of Copper (crystals).*—In the air it becomes slightly pulverulent and of a greenish colour. It is totally soluble in water. Whatever ammonia throws down from this solution an excess of ammonia dissolves.

Remarks.—If it become very green on the surface by exposure to the air, it is owing to the presence of sesquioxide of iron: if it contain this oxide, it is precipitated by ammonia, and an excess does not redissolve it; whereas oxide of copper is readily taken up by it.

CUPRI AMMONIO-SULPHAS. *Ammonio-Sulphate of Copper.*—By heat it is converted into oxide of copper, evolving ammonia. Dissolved in water it changes the colour of turmeric, and solution of arsenious acid renders it of a green colour.

Remarks.—If it do not alter the colour of turmeric there is no excess of sesquicarbonate of ammonia; and without this it is not totally soluble in water, but is decomposed by it with precipitation. Arsenious acid unites with the oxide of copper to form green arsenite of copper, which, being insoluble in water, is precipitated.

FERRI PERCYANIDUM. *Percyanide of Iron.*—It is pure if, after being boiled with dilute hydrochloric acid, ammonia throws down nothing from the filtered solution.

Remarks.—If the percyanide of iron contained uncombined sesquioxide of iron or any alumina, they would be precipitated from the solution by ammonia.

FERRI AMMONIO-CHLORIDUM. *Ammonio-Chloride of Iron.*—Totally soluble in proof spirit and water. Potash added to the solution throws down sesquioxide of iron; afterwards, when added in excess, it evolves ammonia.

Remarks.—The iron of the sesquichloride is precipitated, by the action of the potash, in the state of sesquioxide; while another portion of this alkali decomposes the hydrochlorate of ammonia and evolves its ammonia.

FERRI IODIDUM. *Iodide of Iron.*—Emits violet vapours by heat, and sesquioxide of iron remains. When fresh prepared it is totally soluble in water. From this solution, when kept in a badly stopped vessel, sesquioxide of iron is very soon precipitated; but with iron wire immersed in it, it may be kept clear in a well-stopped vessel.

Remarks.—By the action of the oxygen of the air the iron is converted into sesquioxide, which is insoluble in water; by the operation of the metallic iron, without the access of air, the decomposition even of the solution is prevented.

FERRI POTASSIO-TARTRAS. *Potassio-tartrate of Iron.*—Totally soluble in water: the solution does not change either litmus or turmeric; nor is it rendered blue by ferrocyanide of potassium; nor is anything precipitated from it by any acid or alkali. The magnet does not act upon it.

Remarks.—When improperly prepared, as by using bitartrate of potash and iron filings, a large portion is usually insoluble in water; and sometimes it contains metallic iron attracted by the magnet. Other salts of sesquioxide of iron, give a blue precipitate with ferrocyanide of potassium, and the oxide is thrown down by any alkali. If the solution of this preparation act upon litmus-paper, the tartaric acid of the bitartrate of potash is not saturated with sesquioxide of iron.

FERRI SESQUIOXYDUM. *Sesquioxide of Iron.*—Dissolved totally by dilute hydrochloric acid with very slight effervescence, and it is precipitated by ammonia.

Remark.—The effervescence denotes the presence of a small unimportant portion of carbonic acid; after the precipitation of the sesquioxide of iron by ammonia, no other reagent should produce any change in the solution.

FERRI SULPHAS (*crystalli*). *Sulphate of Iron (crystals).*—Colour bluish green, dissolved by water. Iron put into the solution does not precipitate copper.

Remark.—When these crystals have been kept in a badly stopped bottle, or when exposed to the air, especially if moist, the protoxide of iron becomes sesquioxide, and the crystals are first rendered green, and eventually a yellow deposit is formed on their surface, and this is insoluble in water.

HYDRARGYRUM (*purificatum*). *Mercury (purified).*—Totally dissipated in vapour by heat. Dissolved by diluted nitric acid. When boiled in hydrochloric acid, the acid when cold is not coloured, nor is anything precipitated from it, by hydrosulphuric acid. Its specific gravity is 13.5.

Remarks.—If the mercury contain other metals, most of them would remain after its vaporization; the solubility in nitric acid shows that it contains no tin, and hydrosulphuric acid not acting upon the hydrochloric acid after the mercury has been boiled in it, indicates the absence of most other metals.

HYDRARGYRUM CUM CRETA. *Mercury with Chalk.*—Part is evaporated by heat; what remains is colourless, and totally soluble in acetic acid with effervescence: this solution is not coloured by hydrosulphuric acid. These substances can scarcely be so diligently triturated as that no globules shall be visible.

Remarks.—If the mercury be pure, then it is totally evaporated; and the residue is merely chalk or carbonate of lime, which the acetic acid dissolves with the evolution of carbonic acid. If the mercury contained any metal, it would either be left or dissolved by the acetic acid; if the latter, hydrosulphuric acid would, except in few cases, detect it in the solution.

HYDRARGYRI OXYDUM (cinereum). *Oxide of Mercury (grey).*—Digested for a short time with diluted hydrochloric acid and strained, neither solution of potash nor oxalate of ammonia throws down anything. It is totally soluble in acetic acid. By heat it is totally dissipated.

Remarks.—If the oxide of mercury be pure, it is totally converted by hydrochloric acid into protochloride, which remains insoluble. If it contain any binoxide, it will be dissolved by the acid and precipitated from solution of an orange colour by potash; and if during its preparation any carbonate of lime should have been precipitated with the oxide of mercury, it will be dissolved by the hydrochloric acid, and precipitated as an insoluble oxalate by the oxalate of ammonia. If it contain undecomposed chloride of mercury, it will remain after the action of the acetic acid; and anything which is not evaporated by heat is an impurity.

HYDRARGYRI BINOXYDUM (rubrum). *Bin oxide of Mercury (red).*—On the application of heat it yields oxygen, and the mercury either runs into globules, or is totally dissipated. It is entirely soluble in hydrochloric acid.

Remarks.—When it is dissolved in nitric acid, no precipitate is yielded by nitrate of silver; or if there should be any, either it has not been sufficiently washed, or the bichloride of mercury has been imperfectly decomposed, and, consequently, precipitates the silver of the nitrate of silver as a chloride.

HYDRARGYRI NITRICO-OXYDUM. *Nitric-oxide of Mercury.*—On the application of heat no nitric vapour is emitted. Neither lime-water nor hydrosulphuric acid throws down anything from the water in which it has been boiled. In other respects it resembles the preceding preparation.

Remarks.—Nitric vapour, should it arise, results from the decomposition of nitric acid, which should have been previously expelled by heat; if lime-water or hydrosulphuric acid throw

Process.—Acetate of soda is a salt now placed in the *Materia Medica*; it is made almost exclusively for the preparation of acetic acid in the mode above described. An impure acetic acid, called formerly *pyroligneous acid*, is procured by heating and decomposing wood in iron cylinders; it is rendered impure by various substances, but especially by tarry matter, from which it is partially freed by distillation, and then being combined with soda, the resulting acetate is purified by repeated solution and crystallization.

The crystals of acetate of soda are usually very small; the primary form is an *oblique rhombic prism*. It is colourless, inodorous, and rather sweet to the taste.

It is composed of


One equivalent of Acetic Acid.....	=	51	or	37.23
One equivalent of Soda	=	32	„	23.37
Six equivalents of Water.....	9×6	=	54	„ 39.40
<hr/>				
Equivalent.....		137		100.

Symbol,—Berzelius and Turner $\text{Na} \bar{\text{A}} + 6\text{H}$.

Brande..... $\text{S} + \text{ac}' + 6\text{q}$.

Sulphuric acid having greater affinity for soda than acetic acid has, the acetate of soda is decomposed; the acetic acid being volatile rises and is condensed with the water, of both the acid and salt, anhydrous sulphate of soda remaining in the retort.

Acidum Aceticum, P.L.



Properties and Composition.—Acetic acid has never been obtained except in combination with water or with a base; it was formerly, when of the greatest strength, called *radical vinegar*. It is limpid and colourless, its smell is pungent and refreshing, and its taste acrid unless moderately dilute, and then it is agreeably acid, and devoid of the empyreumatic taste of distilled vinegar. Its acid powers are strongly marked by its

action on litmus paper, which it reddens, and by its decomposing the carbonates of potash and soda, &c. with effervescence. It is volatile even at common temperatures, and its vapour, especially when arising from heated acid, is inflammable, burning with a white light. Its salts are termed *acetates*; they are decomposed by most acids except the carbonic, and indeed even this happens in one or two cases.

Anhydrous acetic acid, as it exists in dry acetate of soda or potash, &c., is composed of

Three equivalents of Hydrogen....	$1 \times 3 =$	3	or	6
Four equivalents of Carbon	$6 \times 4 =$	24	,,	47
Three equivalents of Oxygen	$8 \times 3 =$	24	,,	47
		<hr/>		<hr/>
Equivalent		51		100

Symbol,—Berzelius and Turner .. $\bar{A} = H^3 C^4 O^3$.

Brande $ac' = 4car + 3h + 3o$.

It has been mentioned that acetic acid cannot exist without water or a base; and *glacial acetic acid*, as it has been called, on account of its becoming solid and crystalline at a low temperature, about 40° of Fahrenheit, is the strongest acetic acid procurable; it consists of

One equivalent of anhydrous Acetic Acid	$=$	51	or	85
One equivalent of Water	$=$	9	,,	15
		<hr/>		<hr/>
Equivalent		60		100

Symbol,—Berzelius and Turner..... $\bar{A} + H$.

Brande..... $ac' + q$.

One hundred grains of the acetic acid prepared according to the present directions, saturate 87 grains of crystallized carbonate of soda; and as 144 of this salt are equivalent to 51 of real acetic acid, it follows that this acetic acid is composed of

Anhydrous Acetic Acid	30.8
Water	69.2
	<hr/>
	100.

A mixture of 15 parts by weight of this acid and 85 of water is equal in strength to distilled vinegar.

Impurities and Tests.—See Notes : ACIDUM ACETICUM.

Incompatibles.—Alkalis, alkaline carbonates, alkaline earths and their carbonates, and most substances acted upon by other acids.

Pharmacopœia Preparations.—Acetum Cantharidis, Potassæ Acetas, Plumbi Acetas, Oxymel.

Medicinal Uses.—Acetic Acid when diluted is refrigerant, and may be advantageously administered in hæmorrhage; especially in cases where the acetate of lead has been given, since the solubility of this salt is much increased by it. Externally it may be a useful adjunct to lotions containing lead.

ACETUM CANTHARIDIS. (*Epispasticum.*)

Vinegar of Cantharides. (*Epispastic.*)

Take of Cantharides rubbed to powder two ounces,
Acetic Acid a pint;

Macerate the Cantharides with the Acid for eight days, frequently shaking: lastly, press and strain.

This preparation is now first introduced, and is employed as an extemporaneous blister.

ACETUM COLCHICI.

Vinegar of Meadow Saffron.

Acetum Colchici, P.L. 1809, P.L. 1824.

Take of fresh Meadow Saffron cormus, sliced, an
ounce,

Distilled Vinegar sixteen fluidounces,
Proof Spirit a fluidounce;

Macerate the Meadow Saffron cormus with the Vinegar, in a covered glass vessel, for three days; afterwards press and strain [the liquor] and set it by, that the dregs may subside: lastly, add the Spirit to the clear liquor.

Remarks.—It has generally been supposed that the virtue of meadow saffron resides in a peculiar vegetable alkali similar to that contained in white hellebore and called *veratria*; according, however, to the late experiments of Hesse and Geiger, it is

an alkali differing from veratria, and to which they have given the name of *colchicia*.

Colchicia is prepared by digesting the seed of meadow saffron in boiling alcohol, which dissolves an acid salt; this is to be precipitated by magnesia, and what is thrown down again treated with boiling alcohol, which by evaporation deposits colchicia. In this process the magnesia combines with the acid previously united with the colchicia, which is precipitated with the excess of magnesia and dissolved in an uncombined state by the second portion of alcohol. Colchicia crystallizes in slender needles; it is inodorous, its taste is first bitter and afterwards biting, but not so acrid as veratria, from which it also differs in not exciting sneezing, and in being soluble in water.

Hydrate of colchicia is feebly alkaline to tests, but it completely neutralizes acids and forms crystallizable salts with them, which veratria does not. Its salts have a bitter taste. The aqueous solution of this alkali precipitates a solution of chloride of platina. Nitric acid turns it deep violet, blue, and afterwards quickly green and yellow. It is very poisonous; a tenth of a grain given to a cat acted violently upon it, and killed it in about twelve hours. It has not been analyzed, but like other vegetable alkalis it is probably composed of oxygen, hydrogen, azote, and carbon.

Incompatibles.—Alkalis, their carbonates, the alkaline earths and their carbonates, or any substances on which vinegar is capable of exerting any action.

Medicinal Use.—Diuretic. Dose, f3ss to f3j in any bland fluid. It is also employed in the gout.

ACETUM SCILLÆ.

Vinegar of Squill.

Acetum Scilliticum, P.L. 1720, P.L. 1745.

Acetum Scillæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Squill, fresh dried, fifteen ounces,

Distilled Vinegar six pints,

Proof Spirit half a pint;

Macerate the Squill with the Vinegar with a gentle heat, in a covered glass vessel, for twenty-four hours; afterwards press out the [liquor] and set it by, that the dregs may subside: lastly, add the Spirit to the clear liquor.

Remarks.—Squill contains a peculiar vegetable product to which the name of *scillitine* has been given; it is prepared by the action of spirit upon dried squill, evaporating the solution after the alcohol has been distilled from it, and treating the residue again with spirit and with æther. Scillitine is of a bright yellow colour, and is at first obtained in flocks; these soften in hot water and unite into a mass which becomes brown and brittle by drying. Its taste is very acrid and bitter; when heated it fuses, swells and exhales, first an aromatic odour, and afterwards a urinous smell. It is perfectly soluble in alcohol, but not in æther; dilute acids have no action upon it. A single grain of it is sufficient to kill a large dog. It has not been analyzed.

Incompatibles.—The same as the last preparation.

Medicinal Use.—Expectorant and diuretic. Dose f3ss to f3ij in any aromatic distilled water.

ACIDUM BENZOICUM.

Benzoic Acid.

Flores Benzöini, P.L. 1720.

Flores Benzöes, P.L. 1788.

Acidum Benzoicum, P.L. 1809, P.L. 1824.

Take of Benzoin a pound;

Put the Benzoin in a proper vessel placed on sand, and the heat being gradually raised, sublime until nothing more rises; press that which is sublimed, wrapped in bibulous paper, and separate it from the oily part; afterwards again sublime it.

Remarks.—Benzoin is a balsam or resinous exudation from the *Styrax Benzoin* of Sumatra; this is the only plant which yields it in sufficient quantity to be worth collecting from, but there are others which afford it in small quantity. When it first exudes it is soft, but it becomes gradually hard by exposure to the air; benzoin is usually met with in fragments of a yellowish or fawn colour, intermixed with pieces of wood and leaves. That which exhibits most white, translucent, friable pieces when broken is preferred.

The specific gravity of benzoin is about 1.063; it is easily powdered, has an agreeable odour, and its taste is at first sweetish

and afterwards stimulating. It is soluble in alcohol and æther, but not in oils.

According to the analysis of Stolze, two varieties yielded as follows :—

	White Benzoin.	Brown Benzoin.
Yellow resin, soluble in æther	79·83	8·80
Brown resin, insoluble in æther	0·25	69·73
Benzoic acid	19·80	19·70
Extractive	0·00	0·15
Impurities	0·00	1·15
Moisture and loss, and a trace of volatile oil	0·12	0·47
	<hr/> 100·	<hr/> 100·

Process.—Benzoic acid may be obtained from benzoin by several processes; thus, when it is boiled in water with lime, a benzoate of that base is formed, which is soluble in water, and is decomposed by hydrochloric acid, which precipitates the benzoic acid; or it may be boiled, reduced to powder, with rather more than twice its weight of carbonate of soda, by which a benzoate of soda is formed; and this may be decomposed with sulphuric acid, which combining with the soda precipitates the benzoic acid on account of its sparing solubility in water. The simplest and best process is however that by sublimation, as here directed; the benzoic acid is volatilized at a moderate heat, and condenses in the upper and cool part of the apparatus. The oily matter from which the acid is directed to be separated by absorption and pressure, is probably formed by the decomposition of the resin of the benzoin, and a fresh arrangement and combination of its elements.

Properties.—This acid, when pure, is colourless; it crystallizes in soft and rather elastic crystals, which have scarcely any smell, which however is stronger when the acid is prepared by sublimation than by precipitation; its taste is rather aromatic and penetrating than sour; by exposure to the air it suffers no change; it requires two hundred times its weight of cold or twenty-four of boiling water for solution; on cooling a crystalline mass is obtained which has the appearance of fat; alcohol takes it up readily and in large quantity, and by spontaneous evaporation prismatic crystals of the acid are obtained. Although the aqueous solution acts but feebly upon litmus paper, the acid nevertheless combines readily with alkalis and metallic oxides, forming salts called *benzoates*, which are but little employed in scientific, and not at all in medical chemistry.

Benzoic acid fuses and sublimes at a moderate heat; if strongly heated it burns with a bright yellow flame; it dissolves in sulphuric or nitric acid without suffering decomposition.

Composition.—Benzoic acid cannot exist without water or a base; when forming a dry benzoate, as that of silver, it consists of

Fourteen equivalents of Carbon ..	$6 \times 14 = 84$
Five equivalents of Hydrogen ..	$1 \times 5 = 5$
Three equivalents of Oxygen ..	$8 \times 3 = 24$
	<hr/>
The crystals contain	113
One equivalent of Water	$= 9$
	<hr/>
Equivalent	122

Symbol.—Berzelius and Turner. $\ddot{\text{Bz}} = \text{C}^{14} \text{H}^5 \text{O}^3 + \dot{\text{H}}$.
 Brande..... $\text{ben}' + q$.

This water cannot be separated by heat, but may be by combining the acid with oxide of silver.

According to the experiments of Wöhler and Liebig, benzoic acid is to be considered as the oxide of a compound inflammable body, which they call *benzule*, and they regard it as consisting of 14 equivalents of carbon = 84, 5 of hydrogen = 5, and 2 of oxygen = 16; its equivalent is consequently 105, which by the addition of one equivalent of oxygen = 8, yields, as above, 1 equivalent of anhydrous benzoic acid = 113, and this with one equivalent of water = 9, constitutes crystallized benzoic acid as already shown.

Benzule is capable of combining with sulphur, chlorine, and some other elementary bodies; with hydrogen and azote it forms *benzamide*; when benzoic acid is heated with hydrate of lime a peculiar fluid is obtained which is called *benzine*; and another product, composed of carbon, hydrogen, and oxygen, called *benzone*, has been formed by heating benzoic acid with lime.

Impurities and Tests.—See Notes: ACIDUM BENZOICUM.

Incompatibles.—Such substances as neutralize or combine with acids in general, as alkalis, their carbonates, metallic oxides, &c.

Pharmacopœia Preparations.—Tinctura Camphoræ composita. The gum-resin, and of course the benzoic acid which it contains, is also used in the Tinctura Benzoës composita.

Medicinal Uses.—It is supposed to be stimulant and expectorant, but is rarely used except in the Tinctura Camphoræ composita.

ACIDUM CITRICUM.

Citric Acid.

Acidum Citricum, P.L. 1809, P.L. 1824.

Take of the Juice of Lemons four pints,
Prepared Chalk four ounces and a half,
Diluted Sulphuric Acid twenty-seven fluid-
ounces and a half,
Distilled Water two pints ;

Add the Chalk gradually to the Juice of Lemons made hot, and mix. Set by, that the powder may subside ; afterwards pour off the supernatant liquor. Wash the Citrate of Lime frequently with warm water. Then pour upon it the diluted Sulphuric Acid, and the distilled water, and boil for a quarter of an hour. Press the liquor strongly through linen, and strain ; evaporate the strained liquor with a gentle heat, and set it by, that crystals may be formed.

Dissolve the crystals, that they may be pure, again and a third time in water, and strain the solution as often ; evaporate and set it aside.

Remarks.—There are several fruits, as raspberries, gooseberries, &c., which contain citric acid, and in some instances it is associated with malic acid : the juice of lemons may however be considered as an aqueous solution of citric acid, nearly free from any other, but mixed with a considerable quantity of mucilage which prevents the acid from crystallizing, although it may be evaporated sufficiently.


Process.—Chalk consists of carbonic acid and lime, and is termed chemically, carbonate of lime ; when this is added to the lemon-juice, the citric acid, owing to its greater affinity for the lime than of the carbonic acid for it, the latter acid is expelled with effervescence in the gaseous state. The citrate of lime thus formed being but sparingly soluble in water, most of it remains undissolved and subsides in the state of a white powder, whilst the greater part of the mucilage of the lemon-juice is held in solution.

The first steps in the operation are then the formation of citrate of lime, and the separation of the mucilage ; the aqueous part of the lemon-juice holding the mucilage in solution is rejected,


and the citrate of lime is rendered free from any remains of it by repeated washing.

The citrate of lime when heated with the diluted sulphuric acid is decomposed, on account of the greater affinity existing between sulphuric acid and lime than between citric acid and lime; the sulphate of lime thus formed subsides on account of its slight solubility, and the citric acid remains in solution; by evaporation crystals of citric acid are obtained, which are at first of a brownish colour, but rendered at last white by the repeated solution and crystallization directed to be performed.

The operations now described may perhaps be rendered more intelligible by the annexed diagrams :




Solution of Citric Acid.



Properties.—Citric acid is colourless, inodorous, extremely sour; the primary form of the crystal is a *right rhombic prism*, but which usually exhibits the planes described in the annexed figure: M and M' being the lateral primary planes.

M on M'	101° 30'
M on h	129 15
M on g	163 23
g on g'	134 45
a on a'	111 50
a on b	161 30
h on c ¹	139 45
h on c ²	121 15
c ¹ on c ²	161 30
c ² on c' ²	117 30



By exposure to a damp atmosphere the crystals absorb moisture. One hundred parts of citric acid require 75 of cold and 50 parts of boiling water for solution. The solution reddens litmus paper strongly, and like lemon-juice, decomposes and becomes mouldy by keeping. When citric acid is subjected to distillation it yields pyro-citric acid, acetic acid, carbonic acid, empyreumatic oil, carburetted hydrogen gas, and water, and charcoal remains in the retort. When heated with nitric acid it is converted into oxalic acid. One drachm of the crystals of this acid saturates very nearly two drachms of crystallized carbonate of soda. About eleven drachms and a half of citric acid, dissolved in a pint of distilled water, give a solution equal in strength to lemon-juice.

The following table exhibits the quantities, very nearly, of crystallized citric acid, lemon-juice, and solution of citric acid prepared as above, necessary for the decomposition of one scruple of the alkaline salts named :

<i>A Scruple of</i>	<i>Lemon-Juice or Solution of Citric Acid.</i>	<i>Citric Acid.</i>
Bicarbonate of Potash	f3iijss	gr. 14 - - /5
Carbonate of Potash	f3iiij	gr. 17 - - /8
Sesquicarbonate of Ammonia	f3vi	gr. 24 - - 25

It is to be observed that in the above statements the bicarbonate of potash is considered as crystallized; the carbonate as dry, but containing, as it usually does, about 16 per cent. of water; and the sesquicarbonate of ammonia as translucent and moderately hard; if it be opaque and powdery, the change is owing to the loss of carbonate of ammonia, and bicarbonate being formed, its saturating power is less, and to an extent which is uncertain.

Composition.—Citric acid, like the acetic and benzoic acids, is a compound of hydrogen, carbon, and oxygen; when anhydrous it consists of

Two equivalents of Hydrogen..	$1 \times 2 =$	2 or	3.44
Four equivalents of Carbon....	$6 \times 4 =$	24 „	41.38
Four equivalents of Oxygen ..	$8 \times 4 =$	32 „	55.18
Equivalent.....	58	100.	

Symbol,—Berzelius and Turner.... $\bar{C} = H^2 C^4 O^6$.

Brande $cit' = 4car + 2h + 4o$.

The crystals of citric acid which deposit on the cooling of a solution saturated at 212° contain one equivalent of water ; whereas the acid usually met with in commerce consists of 3 equivalents of anhydrous acid and 4 of water ; or each equivalent of acid is combined with $1\frac{1}{3}$ equivalent of water.

Three eqs. of anhydrous Acid $58 \times 3 = 174$ or 1 eq. = 58 or 82.86
 Four eqs. of Water $9 \times 4 = 36$ „ $\frac{4}{3}$ eq. = 12 „ 17.14

Equivalent....	210	70	100.
----------------	-----	----	------

Symbol,—Berzelius and Turner.. $\bar{C}^3 + H^4$.

Brande $3cit' + 4q$.

At 212° these crystals lose half their water, and they then consist of 3 equivalents of acid and 2 of water, but cannot be deprived of more without suffering decomposition.

Incompatibles.—Citric acid is incompatible with all alkaline solutions and substances, converting them into citrates, as ammonia, potash, soda, lime, magnesia, &c. It decomposes the alkaline, earthy and metallic carbonates, most acetates, the alkaline sulphurets and soaps. It is also incompatible with tartrate of potash, which it converts into citrate and bitartrate of potash.

Impurities and Tests.—See Notes : ACIDUM CITRICUM.

Medicinal Uses.—It is employed as a refrigerant, combined with potash or ammonia in the proportions already stated. Half a fluidounce of lemon-juice, or an equal quantity of a solution of citric acid of the same strength so saturated, is generally exhibited as a dose.

ACIDUM HYDROCHLORICUM.

Hydrochloric Acid.

Spiritus Salis, P.L. 1720.

Spiritus Salis marini Glauberi, P.L. 1745.

Acidum Muriaticum, P.L. 1788, 1809, 1824.

Take of Chloride of Sodium, dried, two pounds,

Sulphuric Acid twenty ounces,


Distilled Water twenty-four fluidounces ;

Add the Sulphuric Acid, first mixed with twelve fluid-ounces of the water, to the Chloride of Sodium put into a

glass retort. Pour what remains of the water into a receiver; then, the retort being fitted to it, let the Acid, distilled from a sand-bath, pass over into this water, the heat being gradually increased.

Process.—The Acidum Hydrochloricum of the Pharmacopœia is an aqueous solution of hydrochloric acid gas; this acid may be obtained in its gaseous state in several modes; first, when equal volumes (or measures) of hydrogen gas and chlorine gas are mixed and exposed to daylight, they combine slowly to form this acid gas; in the sunshine or by the taper or the electric spark, the union is effected instantaneously and with explosion, and the combination being effected without any alteration of volume, there remains hydrochloric acid gas equal in volume to the two measures of the elementary gases employed. Another method of procuring the gas is that of decomposing chloride of sodium by sulphuric acid, but not as directed in the Pharmacopœia, diluted with water. The changes which take place are these:

Sixty parts of chloride of sodium (common salt) are composed of 36 of chlorine and 24 of the metal sodium; liquid sulphuric acid, the acidum sulphuricum of the Pharmacopœia, consists of 40 of dry acid and 9 of water, and the water is composed of 1 of hydrogen and 8 of oxygen. When these 60 of chloride of sodium and 49 of sulphuric acid act upon each other, the water and chloride are both decomposed; the 1 of hydrogen uniting with 36 of chlorine constitute 37 of hydrochloric acid gas, and the 8 of oxygen with the 24 of sodium form 32 of oxide of sodium or soda; and the 40 of dry sulphuric acid uniting with these 32 of soda, they form 72 of dry sulphate of soda, which remain in the retort. The use of the water with which the sulphuric acid is diluted will be presently explained.



When this acid is required in its gaseous state it must be received in glass jars filled with and inverted in mercury, on account of its ready solubility in water.

Properties.—Hydrochloric acid gas is colourless, invisible, and permanently elastic under common circumstances; when, however, it is subjected to a pressure of 40 atmospheres at the temperature of 50° it is rendered liquid, but when the pressure is removed it immediately reassumes the gaseous state. Hydrochloric acid gas has a pungent odour, an acid and acrid taste, and is quite irrespirable and unflammable; it reddens litmus paper strongly, and evinces all the other properties of a powerful acid. It has great affinity for water, and when it escapes into the air combines with the moisture of it so as to form a dense white vapour; a few drops of water introduced into a jar of the gas immediately causes its absorption. Water at 40° is capable of dissolving nearly 480 times its bulk of this gas. It is not altered by heat, but by electricity this gas is partially decomposed.

Composition.—It has been already observed that this gas consists of equal volumes of its elementary gases, and

50 cubic inches of Hydrogen gas weigh.....	1.075 grs.
50 cubic inches of Chlorine gas	38.700 —
100 cubic inches of Hydrochloric acid gas weigh..	39.775 grs.

By weight it is composed of

One equivalent of Hydrogen....	=	1	or	2.7
One equivalent of Chlorine	=	36	„	97.3

Equivalent	37.	100.
-----------------	-----	------

Analytical proof of the nature of hydrochloric acid gas is derived from heating binocide of mercury in it; the new products are bichloride of mercury and water, resulting from the double decomposition of the original compounds, and the fresh arrangement and recombination of their elements.

Symbol,—Berzelius and Turner.. $H + C$, or $H Cl$.

Brande..... $h + c$, or hc' , or m' .

Properties of Solution of Hydrochloric Acid, the Acidum Hydrochloricum of the Pharmacopœia.—The water with which the sulphuric acid is diluted, and that into which the hydrochloric acid gas is passed in the receiver, combine with it and form liquid hydrochloric acid, for brevity's sake usually termed merely hydrochloric acid: its properties are, that when perfectly pure it is a limpid, colourless liquid; it emits white suffocating fumes, which turn vegetable blues red, as the gas and liquid acid also do; its taste is strongly sour and acrid: when its specific gra-

vity is 1.16, as directed, it contains about one third of its weight of hydrochloric acid gas.

It acts upon and dissolves several metals with the evolution of hydrogen gas, arising from the decomposition of the acid: this is the case with iron and zinc, and the chlorine of the acid combining with the metals, they remain dissolved as chloride of iron, zinc, &c. Its saline compounds are termed hydrochlorates, as hydrochlorate of ammonia, of morphia, &c.; but when it acts upon a metal, a chloride is formed for the reason just stated; and when upon a metallic oxide, then a chloride also results; but no hydrogen is evolved, for it unites with the oxygen of the oxide to form water. No metallic hydrochlorates therefore exist.

Impurities and Tests.—See Notes: ACIDUM HYDROCHLORICUM.

Incompatibles.—This acid is incompatible with alkalis, most earths, oxides and their carbonates; sulphuret of potassium, tartrate of potash, potassio-tartrate of antimony, nitrate of silver, acetates of lead, &c.

Pharmacopœia Preparations.—Acidum Hydrochlorium Dilutum—Antimonii Potassio-tartras—Tinctura Ferri Sesquichloridi—Ferri Ammonio-chloridum.

Medicinal Use.—According to Dr. Paris, it may be advantageously employed in malignant cases of scarlatina and typhus; and mixed with a strong infusion of quassia, he considers it to be the most efficacious remedy for preventing the generation of worms.—Dose, ℥v—xx, frequently repeated.

ACIDUM HYDROCHLORICUM DILUTUM.

Diluted Hydrochloric Acid.

Take of Hydrochloric Acid four fluidounces,
Distilled water twelve fluidounces;
Mix.

Medicinal use.—See ACIDUM HYDROCHLORICUM.

Dose of Acidum hydrochloricum dilutum ℥xx—lxx.

One fluidrachm of the undermentioned diluted Acids saturates, very nearly, the annexed quantity of crystallized Carbonate of Soda:

Acidum Hydrochloricum dilutum	32 grains.
Acidum Sulphuricum dilutum	28 —
Acidum Nitricum dilutum	19 —

ACIDUM HYDROCYANICUM DILUTUM.

Diluted Hydrocyanic Acid.

Take of Ferrocyanide of Potassium two ounces,
Sulphuric Acid an ounce and a half,
Distilled water a pint and a half;

Mix the Acid with four fluidounces of the water, and to these when cooled and put into a glass retort, add the Ferrocyanide of Potassium first dissolved in half a pint of the water. Pour eight fluidounces of the water into a cooled receiver; then, the retort being fitted on, let six fluidounces of Acid pass into this water distilled with a gentle heat in a sand-bath. Lastly, add six more fluidounces of distilled water, or as much as may be sufficient, that 12·7 grains of nitrate of silver dissolved in distilled water, may be accurately saturated by 100 grains of this Acid.

Diluted Hydrocyanic Acid may be also prepared, when it is more immediately wanted, from forty-eight grains and a half of Cyanide of Silver, added to a fluidounce of distilled water, mixed with thirty-nine grains and a half of Hydrochloric Acid. Shake all these in a well-stopped vial, and after a short interval pour off the clear liquor into another vessel. Keep this for use, the access of light being prevented.

Remarks.—This acid was first obtained by Scheele in 1782, and as it was procured, though intermediately, from Prussian blue, it was originally called Prussic acid; its present appellation was given to it for reasons presently to be stated. It appears extremely probable that this acid exists ready formed in several vegetable products; for when the leaves of the cherry laurel, the peach tree, or when bitter almonds and some other substances are subjected to distillation, the distilled fluid has the peculiar smell of hydrocyanic acid, and produces, after a certain time, Prussian blue when

added to a saturated solution of iron in carbonic acid. It is however possible that the hydrocyanic acid thus obtained is a product and not an educt, and then its elements only, exist in the substances which yield it by distillation; in the mode, however, in which it is here directed to be prepared, it is unquestionably entirely an artificial product. The use of cyanide of silver was, I believe, first proposed by Mr. Everitt.

Ferrocyanide of Potassium is a well-known salt, frequently called prussiate of potash, or prussiate of potash and iron. It is prepared by heating to redness a mixture of animal matter, such as hoofs, horns, or blood, and potash in an iron vessel. By their mutual decomposition and action on the iron, a coaly mass is obtained, which is partially soluble in water, and the solution by evaporation yields large translucent crystals of a fine yellow colour, the primary form of which is an octahedron with a square base. This salt is inodorous; its taste is rather saline; water at 60° dissolves about one third, and at 212° its own weight of this salt: it is insoluble in alcohol; when moderately heated it loses about 13 per cent. of water, and becomes colourless; when heated to redness with access of air it suffers partial decomposition; the residue, when put into water, leaves oxide of iron, and cyanuret of potassium is dissolved; but if the heat be longer continued, then hydrocyanic acid and ammonia are evolved, and the residue consists entirely of carbonate of potash and sesquioxide of iron. When the aqueous solution of ferrocyanide of potassium is mixed with one of a protosalt of iron, a white precipitate is formed, which speedily becomes blue by exposure to the air; and this, as well as the blue precipitate obtained at once from the salts of sesquioxide of iron, is Prussian blue, or the percyanide of iron.

Composition.—This salt may be regarded under two points of view; first, as consisting of ultimate elements; and secondly, as formed of compounds of these elements. According to Berzelius, by ultimate analysis it yields:

Six equivalents of Carbon	6×6	= 36
Three equivalents of Azote	14×3	= 42
One equivalent of Iron		= 28
Two equivalents of Potassium . .	40×2	= 80
Three equivalents of Water	9×3	= 27

Equivalent 213

The experiments of Gay-Lussac have, however, rendered it more than probable that the carbon and azote exist in combination, forming a distinct compound which he terms *cyanogen*, from Greek words signifying to *generate blue*, because it is necessary to the production of Prussian blue. Cyanogen is obtained by heating

cyanide of mercury in a retort : it is gaseous, inflammable, and by cold and pressure is reduced to a liquid. It is composed of

Two equivalents of Carbon.....	6×2	= 12
One equivalent of Azote		= 14
		—
Equivalent.....		26

It is therefore a bicarburet of azote.

Supposing, then, that the carbon and azote exist as cyanogen combined with the metals, the ferrocyanide of potassium consists of

One equivalent of Cyanide of Iron	$26 + 28$	= 54
Two equivalents of Cyanide of Potassium.	$52 + 80$	= 132
Three equivalents of Water.....	9×3	= 27
		—
Equivalent.....		213

Process.—It appears from the experiments of Mr. Everitt (London and Edinburgh Phil. Mag., Feb. 1835), that when 6 equivalents of sulphuric acid are heated with 2 equivalents of ferrocyanide of potassium, the action that ensues is as follows : Two equivalents of ferrocyanide of potassium contain 4 eq. of cyanide of potassium, of which only 3 eq. are decomposed, as are also 3 eq. of water, the 3 eq. of oxygen of which combine with 3 eq. of potassium and form 3 eq. of potash, which unite with the 6 eq. of sulphuric acid to form 3 eq. of bisulphate of potash ; the 3 eq. of hydrogen of the decomposed water unite with the 3 eq. of cyanogen separated from the potassium, and constitute with them 3 eq. of hydrocyanic acid. There remain undecomposed 1 eq. of cyanide of potassium, and the 2 eq. of cyanide of iron, and these combining form, what is called by Mr. Everitt *yellow salt*, the constitution of which, it will be observed, is precisely the converse of that of the ferrocyanide of potassium, with respect to the quantities of the cyanides.

The annexed statement will show how the various constituents of the acid and salt are disposed of, except, indeed, that the quantity of water distilled with the acid is not given :

<i>Submitted to Distillation.</i>		<i>Results of Distillation.</i>	
6 eq. Sulphuric Acid	240	3 eq. Bisulphate of Potash.....	384
4 — Cyanide of Potassium ..	264	3 — Hydrocyanic Acid.....	81
2 — Cyanide of Iron.....	108	1 — Cyanide Potassium 66	$\left\{ \begin{array}{l} \text{Yellow} \\ \text{Salt....} \end{array} \right\} 174$
12 — Water $\left\{ \begin{array}{l} 6 \text{ in the Acid} \\ 6 \text{ in the Salt} \end{array} \right\}$	108	2 — Cyanide Iron.....108	
		9 — Water	81
	—		—
	720		720

It will be noticed, that the proportions of sulphuric acid and ferrocyanide of potassium here mentioned do not differ much from those in the Pharmacopœia; but a large quantity of water is used with them in the latter case to prevent the waste of hydrocyanic acid which would occur, for want of condensation, without it.

Various other processes have been given for preparing hydrocyanic acid, of which only one needs be mentioned, but is that by which it is procured of the greatest strength, and consequently best exhibits its peculiar properties. Gay-Lussac prepared it as follows: he put into a retort some bicyanide of mercury, to which he adapted a tube of about two feet in length and about half an inch in diameter; one third of this tube, and that nearest the retort, had pieces of marble put into it, while the remaining two thirds were occupied with chloride of calcium. On the bicyanide of mercury were poured about two thirds its weight of concentrated hydrochloric acid, and a gentle heat was applied. Any undecomposed hydrochloric acid was stopped by the marble, and all moisture by the chloride of calcium. During this operation the chlorine of the hydrochloric acid combines with the mercury of the bicyanide to form bichloride, which remains in the retort, while the hydrogen of the hydrochloric acid takes the cyanogen of the bicyanide, and these forming hydrocyanic acid, it is vaporized, and afterwards condensed by the cooled water in the receiver. Any portion of the hydrocyanic acid which may be condensed on the fragments of marble are to be volatilized by a gentle heat and sent in vapour to the receiver.

The properties of the acid thus obtained are, that it is a colourless liquid with a strong odour; its taste is first cooling and afterwards burning, and it is very poisonous. Its specific gravity is .6969: it boils at about 80° , becomes solid at 5° of Fahrenheit, and crystallizes in fibres like nitrate of ammonia. It acts feebly as an acid, but reddens litmus paper, the blue colour of which returns as the acid evaporates. It is extremely volatile, and when a drop on paper is exposed to the air, the evaporation of a part of it renders the remainder so cold that it solidifies; and this effect is produced even when the temperature of the air is 68° . The specific gravity of its vapour is to atmospheric air as 0.9476 to 1. It combines with water and alcohol in all proportions.

It forms but few salts on account of the facility with which it decomposes, and is very subject to spontaneous decomposition even when kept in well-stopped bottles, and especially if exposed to light; this is owing to the tendency of its elements to form new combinations. The first appearance of decomposition is indicated by the acid becoming of a brown tint, which gradually becomes deeper, and eventually ammonia is formed and a black

powder subsides; this contains carbon and azote, and it has been supposed to be a peculiar acid composed of carbon, azote, and hydrogen, and has been called azulmic acid. When diluted with water, and especially if a little hydrochloric acid be added, the decomposition is much retarded.

The above-mentioned are the properties of real or anhydrous hydrocyanic acid; that of the Pharmacopœia, which contains only two per cent. of it, possesses them of course in a much lower degree.

Composition.—Regarding it as a ternary compound, hydrocyanic acid consists of

Two equivalents of Carbon	$6 \times 2 = 12$	or	44.4
One equivalent of Azote	$= 14$	„	51.9
One equivalent of Hydrogen ..	$= 1$	„	3.7

Equivalent 27 100.

It is, however, usually considered as a compound of hydrogen and cyanogen, and hence the name of hydrocyanic acid.

One equivalent of Hydrogen	1	or	3.7
One equivalent of Cyanogen	26	„	96.3

Equivalent 27 100.

Considering it in its elastic state, it may be further regarded as composed of

and as these combine without condensing, }	100	50 { cubic inches of cyanogen gas, weighing	27.950 grs.
		50 { cubic inches of hydrogen gas, weighing	1.075 gr.
		100 { cubic inches of hydrocyanic acid vapour weigh	29.025 grs.

I shall give the symbols of all the compounds which I have had occasion to describe at one view:

Ferrocyanide of Potassium:—

Berzelius and Turner.. $2 K Cy + Fe Cy + 3 H$.

Brande. $(fe + cy) + 2 (po + cy) + 3 q$.

Hydrocyanic Acid:—

Berzelius and Turner.. $H + Cy$ or $H + N_2 C$

Brande. $(cy + h)$ or hcy' .

Yellow Salt :—


Berzelius and Turner... $K\ Cy + 2\ Fe\ Cy.$

Brande $(2\ fe + cy) + (po + cy).$

Incompatibles.—When added to salifiable bases it is in most cases decomposed on account of the slight affinity with which its elements are held in combination; so that when mixed with metallic oxides it usually transfers its hydrogen to the oxygen they contain, and the result is not a hydrocyanate, but a metallic cyanide and water.

Impurities and Tests.—See Notes: ACIDUM HYDROCYANICUM.

In the process introduced for preparing hydrocyanic acid extemporaneously the changes which occur are these: the hydrochloric acid is composed of hydrogen and chlorine, and the cyanide of silver, as its name imports, of cyanogen and silver; when these are mixed double decomposition ensues, and the results are chloride of silver which remains insoluble, and hydrocyanic which is held in solution and poured off when clear. The reaction is explained by the annexed diagram.



Medicinal Uses.—Hydrocyanic Acid has been thought by Majendie, who chiefly introduced it into use, to act as a sedative without possessing the stimulating property of opium. It has been used to allay gastric and pulmonary irritation; in the former, when proceeding from dyspepsia or functional disorder of the stomach, as well as in cancerous affections; and in the latter, to allay cough, and the first symptoms of tubercles in the lungs. Great caution should be observed in its use, and the dose at first should not exceed five or six \mathfrak{m} , which may be increased according to the prudence and judgment of the practitioner. It has likewise been employed in convulsions and spasmodic affections, as hooping-cough and asthma.

ACIDUM NITRICUM.

Nitric Acid.

Aqua fortis simplex et duplex, P.L. 1720.*Spiritus Nitri Glauberi*, P.L. 1745.*Acidum Nitrosum*, P.L. 1788.*Acidum Nitricum*, P.L. 1809, P.L. 1824.


Take of Nitrate of Potash, dried,

Sulphuric Acid, each two pounds;

Mix in a glass retort, then let the Acid distil in a sand-bath.

Process.—The quantities of nitrate of potash and *densest* sulphuric acid directed to be used, are nearly in the proportion of one equivalent of the salt to two equivalents of the acid; and if we employ sulphuric acid of specific gravity 1.8433, which is more commonly met with, the equivalent proportions are still more nearly those above stated, or we may consider two equivalents of the salt and four of the acid as submitted to distillation, which will be more convenient in explanation.

Two hundred and four parts of nitrate of potash are composed of 108 of dry nitric acid and 96 of potash; 205 parts of sulphuric acid consist of 160 of dry acid and 45 of water. When these 204 of the salt and 205 of the acid are mixed and heated, double decomposition occurs;—the 96 of potash = 2 equivalents, combine with 160, the 4 equivalents of dry sulphuric acid and 18 = 2 equivalents of water, and form 274 = 2 equivalents of bisulphate of potash, which remain in the retort:—the 108 = 2 equivalents of dry nitric acid rise in vapour and combine with 27 = 3 equivalents of the water, and are condensed in the receiver forming 135 = 2 equivalents of liquid nitric acid.

135 *Liquid Nitric Acid.*

The process is improved by omitting the rectification of the acid, which was directed in the last Pharmacopœia. Nitric acid of sp. gr. 1.5033 to 1.504, as obtained by this method, is, I believe, the strongest procurable; and as two equivalents of the dry acid require three equivalents of water for their condensation, this circumstance will explain the reason for using so much sulphuric acid, on account of the water it contains, when means are not provided for condensing the nitric acid by passing it into water in a Woulfe's apparatus; which is the method practised by those who prepare this acid for manufacturing purposes; and using iron retorts and a high degree of heat, they employ only half the quantity of sulphuric acid.

Properties.—Liquid nitric acid, usually called merely nitric acid, is a dense colourless fluid; it emits white disagreeable fumes; its taste is extremely sour and acrid, and the skin is indelibly tinged of a yellow colour by it. When exposed to the air it attracts water, for which it has considerable affinity; and when they are suddenly mixed, heat is evolved.

Nitric acid has usually a yellowish tint, owing to the presence of a small and unimportant quantity of nitrous acid, formed by the partial decomposition of a little of the nitric acid during its preparation. If the coloured nitric acid be moderately heated in a retort, nitric oxide is expelled, and it is rendered colourless. By exposure to light, and especially to the direct rays of the sun, nitric acid becomes first of a straw colour, and then of a deep orange; this change is owing to the evolution of oxygen, and the consequent formation of nitrous acid. If concentrated, it does not act upon the metals in general, at ordinary temperatures, nor on some of them even when boiled in it; but when a little water is added, most of them decompose a portion of the acid, and the water also; and the metals combining with the oxygen of both, they are either oxidized and become insoluble, or are dissolved and converted into nitrates by the nitric acid remaining undecomposed; during this action nitric oxide gas (nitrous gas) is given out, which uniting with the oxygen of the air, forms red nitrous acid gas. Nitric acid is decomposed by some combustible bodies with great rapidity, as by charcoal, phosphorus, and sugar.

Composition.—Dry nitric acid, as it exists in nitrate of potash and other anhydrous nitrates, is composed of

Five equivalents of Oxygen..	$8 \times 5 = 40$	or 74
One equivalent of Azote	$= 14$	„ 26
	<hr/>	<hr/>
Equivalent	54	100

The elements of nitric acid are incapable of existing in the proportions above stated, unless combined with a salifiable base,

as with potash, forming nitrate of potash, or with water, constituting liquid nitric acid; this, when of the greatest strength, or of sp. gr. 1.5033 to 1.504, is composed of

Two equivalents of Nitric Acid	=	108	or	80
Three equivalents of Water 9×3	=	27	,,	20
		135		100

It is however more convenient in practice to consider liquid nitric acid of this strength as composed of

1 equivalent of Nitric Acid	=	54
1½ equivalent of Water	..	13·5
		<hr/>
Equivalent		67·5

Symbol,—(Dry Acid) Berzelius and Turner.. $\text{N} + 5 \text{O}$ or $\ddot{\text{N}}$.

Brande $n + 50$ or n' .

Symbol,—(Liquid Acid) Berzelius and Turner. $\cdot\ddot{\text{N}} + 3\dot{\text{H}}$.

Brande $2n' + 3q$.

Adulteration.—If pure nitrate of potash be employed in the College process, the nitric acid obtained is perfectly free from all admixture except a little nitrous acid, which, as already noticed, is quite unimportant. The impurities usually occurring in the nitric acid of the shops are the sulphuric and hydrochloric acids.—See Notes: ACIDUM NITRICUM.

Incompatibles.—It has been before observed, that, when moderately diluted, this acid is readily decomposed by most metals; but it has no action upon platina or gold, and they, of course, do not decompose it. When mixed with hydrochloric acid, both suffer decomposition, and chlorine and nitrous acid result. The mixture is called either aqua regia or nitro-hydrochloric acid, and it possesses the power of dissolving both platina and gold. The action of combustible bodies upon this acid has been adverted to. It is incompatible with sulphate of iron, the protoxide of which decomposes it, and combining with its oxygen, becomes sesquioxide, and the colour of the solution of iron changes from bluish green to yellowish red. It acts strongly upon spirit of wine, and by their mutual decomposition nitric æther is formed. Oxides, earths, alkalis, and their carbonates, are incompatible with this acid, and sulphurets are decomposed by it. It decomposes the solution of acetate of lead and acetate of potash, expelling the acetic acid, and forming nitrate of lead and nitrate of potash.

Pharmaceutic Uses.—Nitric acid is employed in several preparations; as, Argenti Nitras, Hydrargyri Nitrico-oxydum, Spiritus Ætheris Nitrici, and Unguentum Hydrargyri nitratis. It is sometimes employed externally as an escharotic.

Medicinal Uses.—See Diluted Nitric Acid.

ACIDUM NITRICUM DILUTUM.

Diluted Nitric Acid.

Acidum Nitrosum dilutum, P.L. 1788.

Acidum Nitricum dilutum, P.L. 1809, P.L. 1824.

Take of Nitric Acid a fluidounce,
Distilled Water nine fluidounces;
Mix.

Composition.—One hundred grains of this diluted acid contain 14·3 of the concentrated acid, and consequently saturate about 31 grains of crystallized carbonate of soda; by weight, therefore, their respective strengths are to each other exactly as 1 to 7: the specific gravity of the diluted acid is 1·080, each fluidrachm contains nearly 9 grains of the concentrated acid, saturating $19\frac{1}{2}$ grains of crystallized carbonate of soda.

Medicinal Use.—This acid is a very powerful antiphlogistic remedy, and is probably serviceable in restraining the progress of syphilis, when an impaired constitution or other circumstances render the exhibition of mercury improper. If sufficiently diluted, it forms an excellent lotion for old indolent ulcers. It is expectorant, and is occasionally used with success in counteracting the consecutive effects of opium. Dose ℞ to xl.

ACIDUM PHOSPHORICUM DILUTUM.

Diluted Phosphoric Acid.

Take of Phosphorus an ounce,
Nitric Acid four fluidounces,
Distilled Water ten fluidounces;
Add the Phosphorus to the Nitric Acid mixed with the


water in a glass retort placed in a sand-bath ; then apply heat, until eight fluidounces are produced. Put these again into the retort that eight fluidounces may distil, which are to be rejected. Evaporate the remaining liquor in a capsule made of platina until only two ounces and six drachms remain. Lastly, add to the Acid, when it is cold, as much distilled water as may be sufficient to make it accurately measure twenty-eight fluidounces.

Remarks.—Phosphorus is a well-known elementary body; combined with oxygen it forms phosphoric acid, and this united with bases constitutes some of the salts of the animal fluids, and united with lime it forms phosphate of lime, almost the whole of the harder portion of bone. Phosphorus is procured from the phosphoric acid of burnt bones, by treating them with dilute sulphuric acid, which combining with the lime separates the phosphoric acid, and this by evaporation and subsequent treatment with charcoal in a retort at a high temperature, loses its oxygen, and the phosphorus being vaporized is condensed in water.

Phosphorus is solid, translucent and nearly colourless ; it is so soft that it may be indented by the nail and very easily cut. It fuses at about 108° , and at 550° it is vaporized ; it is insoluble in water or alcohol, but dissolved by æther and by oils. It is extremely inflammable, and has been known to take fire in the air spontaneously when its temperature was not above 60° . On this account it is always kept in water ; it undergoes slow combustion when exposed to the air, and hence is luminous in the dark, and emits a disagreeable garlic-like smell. It absorbs oxygen with great facility, and is capable of uniting with it in several proportions, forming one oxide and several acids, of which the phosphoric is the best known, and may be obtained by merely burning phosphorus in oxygen gas or in atmospheric air.

Process.—Nitric acid, as has been already explained, is a compound of oxygen and azote, which, when exposed to, and especially if heated with, certain bodies that have a powerful affinity for oxygen, it is decomposed by them. This is particularly the case with phosphorus, which if added to strong nitric acid decomposes it with explosion and rapid combustion. When diluted nitric acid is used, as here directed, the action is slow, the phosphorus gradually melts, separates the acid into oxygen, with which it combines, and nitric oxide gas, which is evolved. A portion of the nitric acid distils before the whole of the phosphorus is acidified, and hence the necessity of returning it into the retort as ordered. The reaction which occurs will be understood by the annexed

diagram, presuming that although nitric acid is decomposed, the decomposition is not total but partial, so that whilst part of the oxygen combines with phosphorus, another portion of it goes off in combination with the azote of the nitric acid, as nitric oxide gas.



Properties.—The phosphoric acid obtained by the above process is a colourless, inodorous, sour fluid; it acts strongly upon litmus paper, and evinces powerful acid properties by the permanent saline compounds which it forms with alkalis, earths and metallic oxides; although evaporated so as to become a very dense fluid it does not destroy animal or vegetable fibre like sulphuric acid. Even when heated to redness it does not lose all its water, but is converted into metaphosphoric acid.

Phosphoric acid combined with soda forms phosphate of soda, a well-known purgative salt. When lime water is added to phosphoric acid an insoluble phosphate of lime is precipitated. The phosphates of barytes, strontia, lead, &c. are insoluble in water, but differ from the sulphates of these bases in being soluble in dilute nitric acid. The phosphates give a yellow precipitate with nitrate of silver, which is a phosphate; but if the phosphate be heated to redness before it is dissolved in water, it gives a white precipitate with nitrate of silver, which is a pyrophosphate of silver.

Composition.—Phosphoric acid is composed of

Five equivalents of Oxygen..	$8 \times 5 = 40$	or $2\frac{1}{2}$ eq. = 20	or 55.5
Two equivalents Phosphorus	$16 \times 2 = 32$	„ 1 eq. = 16	„ 44.4
Equivalent....	72	36	99.9

Symbol,—Berzelius and Turner $2\text{P} + 5\text{O}$ or $\ddot{\text{P}}$.

Brande $(p + 2\frac{1}{2}\text{O})$ or p' .

Impurities and Tests.—See Notes: ACIDUM PHOSPHORICUM.

Medicinal Uses.—Phosphoric Acid possesses the tonic properties of Sulphuric Acid, and is preferable to it in point of flavour. It has also been used with advantage to correct those morbid

states of the system in which a tendency exists to unusual depositions of phosphate of lime, such as in cases of exostosis, or formation of bony tumors, as well as in some forms of urinary concretions.

ACIDUM SULPHURICUM DILUTUM.

Diluted Sulphuric Acid.

Spiritus Vitrioli tenuis, P.L. 1745.

Acidum Vitriolicum dilutum, P.L. 1788.

Acidum Sulphuricum dilutum, P.L. 1809, P.L. 1824.

Take of Sulphuric Acid a fluidounce and a half,
Distilled Water fourteen fluidounces and a half;

Add the Acid to the Water gradually, and mix.

Remarks.—Dry sulphuric acid, sometimes called anhydrous or real sulphuric acid, to distinguish it from the liquid acid which contains water, is composed of

Three equivalents of Oxygen $8 \times 3 = 24$	or Oxygen60
One equivalent of Sulphur ..	= 16 „	Sulphur40
		<hr/>
Equivalent....	40	100

Liquid sulphuric acid, most frequently termed, as in the Pharmacopœia, merely sulphuric acid, and often oil of vitriol, consists of

One equivalent of dry Acid = 40	or Dry Sulphuric Acid	81.6
One equivalent of Water .. = 9 „	Water	18.4
		<hr/>
Equivalent....	49	100.0

Symbol,—(Dry Acid) Berzelius and Turner .. $\text{S} + 3\text{O}$ or S^{III} .
Brande $\text{S} + 3\text{O}$ or S' .

Symbol,—(Liquid Acid) Berzelius and Turner .. $\text{S}^{\text{III}} + \text{H}$.
Brande $\text{S}' + \text{q}$.

The specific gravity of liquid sulphuric acid at 62° of Fahren-

heit is to that of water nearly as 1·845 to 1·000 ; if it exceed this, its usual purity may be questioned ; generally however it is only about 1·8433 and then it is constituted very nearly of

Four equivalents of dry Acid = 160 or Dry Sulphuric Acid 78
 Five equivalents of Water.. = 45 „ Water22

Equivalent....	205	100
----------------	-----	-----

My future observations will in general apply to acid of the last-mentioned strength. Liquid sulphuric acid is colourless, transparent, inodorous and of an oily consistence ; it is highly acrid and corrosive ; its acid properties are extremely strong, so that a single drop gives to a large quantity of water the power of reddening vegetable blue colours ; but when undiluted it has the property of turning vegetable yellow colours brown, as the alkalis do, but it is removed by water. Its boiling point is about 545°, and it solidifies at 15° below zero.

Sulphuric acid has great affinity for water. By exposure to the air, in an open vessel, it imbibes one third of its weight in 24 hours, and more than six times its weight in a twelvemonth. When one part of water is suddenly mixed with four times its weight of concentrated sulphuric acid, both at the temperature of 50°, it is raised to 300° ; but, according to Dr. Ure, the greatest heat is excited by mixing 73 parts of acid with 27 of water. The mixture of sulphuric acid and water always occupies less space than before combination, and the last-mentioned proportions more especially, for the highest temperature is always succeeded by the greatest condensation.

Concentrated sulphuric acid acts very slowly upon the metals at ordinary temperatures ; but at a boiling heat many of them decompose it, and are oxidized by combining with a portion of its oxygen, while sulphurous acid is given out in the gaseous state. When diluted, it rapidly dissolves those metals which decompose water by its agency, as iron and zinc, with the evolution of hydrogen gas, and it dissolves the oxides of most other metals. It readily combines with the alkalis and earths, and forms with them various important salts.

Most vegetable and animal substances, on account of the carbon they contain, decompose and are decomposed by sulphuric acid ; and this renders the acid of a dark colour. Although sulphuric acid ought to be colourless, yet the slight colour which it often acquires from the circumstance just mentioned, does not materially deteriorate it in quality or reduce its strength.

Sulphuric Acid acts upon alcohol ; and the nature of the product depends upon the relative proportions employed. If equal weights of the acid and rectified spirit be heated in a retort, the product is sulphuric æther ; when two of acid and one of spirit

are used, ætherial oil is obtained ; and when the proportions are 7 acid to 1 spirit, olefiant gas is plentifully formed.

Medicinal Uses.—It possesses the refrigerant and antiseptic virtues common to other acids ; and it has astringent properties that render it a most valuable medicine in weakness and relaxation of the digestive organs, in colliquative sweats and in internal hæmorrhage. A fluidrachm of the acidum sulphuricum dilutum contains about ten grains of the strong acid, and will saturate 28 grains of crystallized carbonate of soda. Dose \mathfrak{m} x to xl. See Infusum Rosæ.

Pharmaceutical Uses.—Sulphuric acid enters into the composition of Infusum Rosæ, Potassæ Sulphas and Bisulphas, Sodæ Sulphas, Ferri Sulphas and Zinci Sulphas. It is employed in preparing Acidum Citricum, Hydrochloricum, Nitricum and Tartaricum, Æther Sulphuricus, Antimonii Oxysulphuretum, Hydrargyri Chloridum and Bichloridum, Potassæ Bicarbonas, and Sodæ Sesquicarbonas.

Incompatibles.—All substances that combine with this acid are of course incompatible with it ; such, as already mentioned, are most of the metals, their oxides, some of the earths, their carbonates, and the alkaline carbonates. The solutions of acetate of lead and of chloride of calcium are decomposed by it, white precipitates of sulphate of lead and sulphate of lime being obtained. Its presence is detected by the action of barytic salts, with the base of which it forms sulphate of barytes, soluble only in concentrated sulphuric acid.

Adulteration.—Sulphuric acid always contains sulphate of lead, derived from the chambers in which it is manufactured, and some sulphate of potash : these impurities generally amount to about 1-4th of a grain per cent. When water is added to the acid, the sulphate of lead is precipitated in the state of a white insoluble powder, from which the diluted acid should be poured off for use. If sulphate of potash should be fraudulently mixed with the acid, for the purpose of increasing its specific gravity, the best method of detecting it is to saturate the acid with ammonia, and expel the sulphate of ammonia formed, by putting it into a crucible and subjecting it to a red heat ; the sulphate of potash will remain in the crucible.

ACIDUM TARTARICUM.

Tartaric Acid.

Acidum Tartaricum, P.L. 1824.

Take of Bitartrate of Potash four pounds,
Boiling distilled Water two gallons and a half,
Prepared Chalk twenty five ounces and six
drachms,
Diluted Sulphuric Acid seven pints and seven-
teen fluidounces,
Hydrochloric Acid twenty six fluidounces and
a half, or as much as may be sufficient.

Boil the Bitartrate of Potash with two gallons of distilled water, and add gradually half the prepared chalk; afterwards, the effervescence having ceased, add the remainder of the chalk first dissolved in the Hydrochloric Acid with four pints of distilled water. Lastly, set by, that the Tartrate of Lime may subside; pour off the liquor, and wash the Tartrate of Lime frequently with distilled water, till it is free from taste. Then pour on it the diluted Sulphuric Acid, and boil for a quarter of an hour. Evaporate the strained liquor with a gentle heat, that crystals may be formed.

Dissolve the crystals, that they may be pure, again and a third time in water, and strain it as often, boil down, and set it aside.

Remarks.—Bitartrate of Potash, sometimes called Supertartrate of Potash, Tartar or Cream of Tartar, is a well-known acidulous salt deposited from wine. It occurs in the state of small colourless hard crystals, for an account of which see *Potassæ Tartras*. This salt is very sparingly soluble in water. In its original impure state it is called *argol*, and is of a yellowish or red colour, according to that of the wine which yields it. It contains tartrate of lime, colouring matter and other impurities, from which it is freed by solution in boiling water, and crystallization as the solution cools.

Properties.—Bitartrate of Potash is inodorous, has a sour

taste; when dissolved in water the solution reddens litmus paper; it requires 95 parts of cold and 15 of boiling water to dissolve one part of it; the solution decomposes by exposure to the air, and becomes a weak solution of carbonate of potash. When exposed to distillation it yields a great quantity of carbonic acid, some acetic acid, carburetted hydrogen, carbonate of ammonia, and empyreumatic oil. When calcined in an open fire it leaves carbonate of potash mixed with a little carbonate of lime derived from the tartrate of lime, of which it contains generally about 5 per cent.

The primary form of the crystal of bitartrate of potash is a *right rhombic prism*: the first of the annexed figures represents the planes of its ordinary crystal in a perfect state; M and M' are the lateral primary planes, and the crystals admit of cleavage parallel to those planes, and to the plane *h*, which is parallel to the shorter diagonal of the primary prism; it also cleaves parallel to the longer diagonal. The crystals are not, however, commonly so perfect as this figure, nor indeed is it usual to observe all its planes; for owing to the extraordinary enlargement of certain of them, others are either much diminished, or totally disappear. The common crystals are represented by the second figure; and in observing them, it must be recollected that the plane *h* is constantly striated, as represented in both figures.

Fig. 1.





Fig. 2.



M on M'	107°	30'
— <i>h</i>	126	15
— <i>b</i>	117	2
<i>b</i> on <i>b'''</i>	74	0
<i>b</i> on <i>c</i>	141	25
<i>b</i> on <i>c'</i>	103	18
<i>c</i> on <i>h</i>	125	30
<i>c</i> on <i>c'</i>	109	0

It is composed of


Two equivalents of Tartaric Acid..	$66 \times 2 = 132$	or 70
One equivalent of Potash	$= 48$	„ 25.3
One equivalent of Water	$= 9$	„ 4.7

Equivalent 189 100.


Symbol,—Berzelius and Turner.. $\bar{K} \bar{T}^2 + H$.

Brande. $2 \text{ tar}' + P + q$.

Process.—The tartaric acid of the bitartrate of potash is by two distinct operations converted into tartrate of lime. When half the chalk or carbonate of lime is added, as directed, to the whole of the bitartrate of potash, one half of the acid which it contains, and which constitutes it a bitartrate, acts upon it as a free acid, carbonic acid is expelled, and tartrate of lime formed, which being a salt of little solubility is precipitated; there remains in solution neutral tartrate of potash, as shown by the annexed diagram :




When the remaining half of chalk or carbonate of lime is dissolved in hydrochloric acid, we obtain, as shown under the head of *CALCII CHLORIDUM*, a solution of chloride of calcium; and when this is added to the now *neutral* tartrate of potash, left as described in the first operation, the changes that occur by double elective affinity and decomposition, are, that the chlorine quits the calcium to combine with the potassium of the potash, chloride of potassium is formed and remains in solution; the oxygen separated from the potassium unites with the calcium left by the chlorine, lime is the result, and this combines with the tartaric acid separated from the potash, and tartrate of lime is formed, which, as already explained, is precipitated on account of its insolubility, thus :



When the tartrate of lime thus formed is mixed with the diluted sulphuric acid, it is decomposed owing to the superior affinity of the sulphuric acid for lime, and the sulphate of lime precipitating on account of its insolubility in water, the tartaric acid remains in solution :

Solution of Tartaric Acid.



Sulphate of Lime.

The solution of tartaric acid by evaporation yields crystals, which are purified from their colouring matter by the repeated solution and crystallization directed to be performed.

Properties.—Tartaric acid is colourless, inodorous, and very sour to the taste; it occurs in crystals of considerable size, the primary form of which is *an oblique rhombic prism*.

Fig. 1. exhibits the crystal as usually modified, with the planes symmetrically placed. Fig. 2. exhibits the same modified form, with the planes irregularly disposed, as they appear in most of the crystals, the corresponding planes in both being marked with the same letters. This affords another instance of irregularity, which renders it not easy immediately to perceive the relations of the several planes to each other.

Fig. 1.





Fig 2.



P on M, or M'	97° 10'
M on M'	88 30
P on e or e'	128 15
P on a	134 50
P on h	100 47

Tartaric Acid suffers no change by exposure to the air; water at 60° dissolves about one fifth, and at 212° twice its weight; the solution acts strongly upon vegetable blue colours; it decomposes and becomes mouldy by keeping. It is soluble in

alcohol, but more sparingly so than in water. When the crystals are heated to a little above 212° , they melt into a fluid which boils at 250° , leaving a semitransparent and rather deliquescent mass on cooling. When more strongly heated in a retort, it is decomposed, and yields among other products a peculiar acid, called pyrotartaric acid; when heated in the air, a coaly mass is procured, but which is eventually dissipated.

When treated with sulphuric acid it is decomposed, and acetic acid is obtained; by nitric acid it also suffers decomposition, and a portion of its carbon combines with sufficient oxygen to convert it into oxalic acid. Tartaric acid has a remarkable tendency to form double salts (Sodæ Potassio-tartras).

Tartaric acid when in solution readily acts upon those metals which decompose water, such as iron and zinc; like the bi-salt from which it is obtained, it combines with most metallic oxides, with the alkalis, and most earths to form salts, which are called tartrates.

Composition.—Tartaric acid free from water, in which condition, however, it cannot be obtained by mere exposure to heat, is composed of

Four equivalents of Carbon	$6 \times 4 = 24$	or	36.4
Five equivalents of Oxygen	$8 \times 5 = 40$	„	60.6
Two equivalents of Hydrogen	$1 \times 2 = 2$	„	3.
	<hr/>		<hr/>
Equivalent	66.		100.

In the state of crystals the acid consists of

1 equivalent of Dry Acid	66	or	88
1 equivalent of Water	9	„	12
	<hr/>		<hr/>
	75.		100.

Symbols.—Berzelius and Turner .. $\overline{T} = H^2 C^4 O^5$.

Do. (Crystals) .. $\overline{T} + H$.

Brande $(4 car + 2 h + 5 o)$ or tar' .

Do. (Crystals) .. $(tar' + q)$.

Incompatibles.—Tartaric acid, as already noticed, combines with the alkalis and decomposes their carbonates; its effects are similar upon most earths and oxides and their carbonates, and it is therefore incompatible with them. It decomposes the salts of potash when in solution, converting them into bitartrate, which is quickly precipitated in minute crystals; the solution also gives immediate precipitates with lime-water, the salts of lime and of lead; and when tartrate of potash is heated with a solution of

chloride of platina, metallic platina is precipitated in the state of a black powder.

Impurities and Tests.—See Notes : ACIDUM TARTARICUM.

Medicinal Uses.—This acid being cheaper than citric acid, it is sometimes employed instead of it, especially in preparing what are called sodaic powders, used as substitutes for soda water. It is largely used in calico-printing. Bitartrate of potash is employed in the Pharmacopœia in preparing Potassæ Tartras, Sodæ Potassio-tartras, Antimonii Potassio-tartras, Ferri Potassio-tartras.

PREPARATIONS OF ÆTHER.

ÆTHER SULPHURICUS.

Sulphuric Æther.

Æther vitriolicus, P.L. 1788.

Æther rectificatus, P.L. 1809, P.L. 1824.

Take of Rectified Spirit three pounds,

Sulphuric Acid two pounds,

Carbonate of Potash, previously ignited, an ounce ;

Pour two pounds of the spirit into a glass retort, add the acid to it and mix. Afterwards place it on sand, and raise the heat so that the liquor may quickly boil, and the æther pass into a receiving vessel cooled with ice or water. Let the liquor distil until some heavier portion begins to pass over. To the liquor which remains in the retort, after the heat has subsided, pour the remainder of the spirit, that æther may distil in the same manner.

Mix the distilled liquors, then pour off the supernatant portion, and add to it the carbonate of potash, shaking them frequently during an hour. Lastly, let the æther distil from a large retort, and be kept in a stopped vessel.

Properties.—Sulphuric æther is a colourless, limpid, transparent fluid, of a pleasant smell and a pungent taste; it is extremely volatile, and its vapour when respired mixed with air produces very exhilarating effects. When recently prepared, æther is not at all acid, but after long keeping it reddens litmus paper, owing to the formation of acetic acid by absorbing oxygen from the air. According to Mitscherlich its specific gravity is $\cdot 724$ at 55° Fahrenheit; when its density exceeds this, it contains either alcohol or water, or both. At 30° below zero it begins to assume a foliated appearance, and at 17° lower, it becomes a white, solid, crystalline mass.

Owing to the extreme volatility of æther a considerable quantity evaporates, even while being poured from one vessel to another, and during this evaporation it produces much cold, as may be felt by pouring it on the hand; if a small thin glass tube enveloped in cloth, and containing water, be dipped a few times in æther, allowing it to evaporate after each immersion, the water may be frozen; in a cold atmosphere even mercury may be solidified by its cooling power.

The specific gravity of the vapour of æther is $2\cdot 586$, that of air being 1; so that although it is a very light fluid it yields a dense vapour. Under a mean atmospheric pressure æther boils at about 96° of Fahrenheit, and *in vacuo*, even at 40° below zero; therefore if it were not for the pressure of the air, it would always exist in the aëriform state.

Æther and the vapour which arises from it are very inflammable, and during its combustion under common circumstances, water is formed by the union of its hydrogen with the oxygen of the air, and carbonic acid by the combination of its carbon with the same element. When, however, a coil of platina wire is heated to redness, and then suspended above the surface of æther contained in an open vessel, the wire instantly begins to glow, and continues in that state until the æther is consumed; during this combustion acetic acid is formed instead of the carbonic, and it was at first supposed by Professor Daniell to be a distinct acid; he has however since found that it is acetic acid combined with a peculiar compound of carbon and hydrogen. When the vapour of æther is passed through a red hot porcelain tube it is decomposed, and the products are charcoal, water, carbonic oxide and carburetted hydrogen.

But little water is dissolved by æther, and water takes up only one ninth of its volume of æther. When æther contains alcohol it may be separated by shaking with water, which dissolves the alcohol, and the water dissolved by the æther may be separated by agitation with lime and subsequent distillation. In the formula carbonate of potash is used for the last-mentioned purpose.

Æther dissolves the resins, and even caoutchouc, which is not acted on by alcohol; it takes up a little sulphur and phosphorus; the solution of this last substance becomes luminous in the dark when poured on the hand or hot water. The alkalis, potash and soda are not soluble in æther, which constitutes another marked difference between it and alcohol.

Process.—The nature of the reaction by which æther is produced is a subject which has been much discussed, and some difference of opinion still exists respecting it.

Spirit of wine, or rather the alcohol which it contains, and to which its properties are owing, is composed of

Two equivalents of Carbon	$6 \times 2 = 12$
One equivalent of Oxygen	$= 8$
Three equivalents of Hydrogen $1 \times 3 =$	3
	<hr/>
Equivalent	23

Æther consists of

Four equivalents of Carbon	$6 \times 4 = 24$
One equivalent of Oxygen	$= 8$
Five equivalents of Hydrogen . . $1 \times 5 =$	5
	<hr/>
Equivalent	37

As the equivalent of Æther contains twice as much carbon as that of alcohol, it is evident that at least two equivalents of alcohol are required to produce one of æther, and it is equally apparent that the difference between them depends upon the different proportions of the other elements; thus :

	Carbon.	Oxygen.	Hydrogen.	
Two eq. of Alcohol	$= 24 +$	$16 +$	$6 =$	46
One eq. of Æther	$= 24 +$	$8 +$	$5 =$	37

$$8 + 1 = 9, \text{ one eq. of water.}$$

When then either an equivalent of water, or of each of the elements which constitute it, is separated from two equivalents of alcohol, the result is one equivalent of æther; and it was supposed by Fourcroy and Vauquelin that the sulphuric acid in its action upon alcohol produced this effect in a direct mode, on account of its well-known great affinity for water; this however does not appear to be the case.

According to Mr. Hennell the action of the sulphuric acid is intermediate; he found that when two equivalents of sulphuric acid, and two of alcohol were merely mixed, the acid immediately lost four sevenths of its power of precipitating oxide of lead, and

undergoing great change of properties, was converted into a peculiar acid called the sulphovinic, consisting of

$$\begin{array}{rcl} \text{Two equivalents of Sulphuric Acid..} & 40 \times 2 = & 80 \\ \text{Two equivalents of Alcohol.....} & 23 \times 2 = & 46 \end{array}$$

$$\text{Equivalent.... } 126$$

or, what is the same, of

$$\begin{array}{rcl} \text{Two equivalents of Sulphuric Acid..} & 40 \times 2 = & 80 \\ \text{Two equivalents of Oxygen,} & 8 \times 2 = & 16 \\ \text{Six equivalents of Hydrogen,} & 1 \times 6 = & 6 \\ \text{Four equivalents of Carbon,} & 6 \times 4 = & 24 \end{array}$$

$$\text{Equivalent.... } 126$$

When an equivalent of this acid is heated it is decomposed ; the two equivalents of sulphuric acid, and one equivalent of water, remain in the retort, while the other elements combine to form an equivalent of æther, which is distilled ; thus :—

	Sulphuric Acid.	Oxygen.	Hydrogen.	Carbon.	
1 eq. Sulphovinic Acid =	80	+ 16	+ 6	+ 24	= 126
Remain in the retort	80	+ 8	+ 1		= 89
Distilled....	8	+ 5	+ 24		= 37 æther.

In the mode in which its constitution is here stated, æther is a mere ternary compound of its elements ; it has however with great plausibility been supposed that two of these elements exist as a binary compound, and that it is united with the other to form æther. Thus it may be shown that the elements of æther are equivalent to, and have been supposed to exist as a dihydrate of olefiant gas ; as a hydrate of etherine, that is of a peculiar carburetted hydrogen to which this name has been given ; lastly, it has been proposed by Dr. Kane (Dublin Journal of Medical Science, &c., January 1833) to consider æther as a protoxide of *ethereum*, the name he gives to a supposed compound, consisting of the 4 eq. carbon and 5 eq. of hydrogen which are contained in the æther. A similar view was soon afterwards promulgated by Berzelius, who gave the theoretic carburetted hydrogen the name of *ethule*.

Preferring this to the other theoretical opinions, I shall exhibit the composition of *ethereum*, æther, alcohol and sulphovinic acid in accordance with it, referring to Turner's Chemistry, and Brande's Manual for further information, and for an explanation of the other opinions alluded to.

OLEUM ÆTHEREUM.

Æthereal Oil.

Oleum Vini, P.L. 1788.

Oleum Æthereum, P.L. 1809, P.L. 1824.

Take of Rectified Spirit two pounds,
Sulphuric Acid four pounds,
Solution of Potash,
Distilled Water, of each a fluidounce, or as
much as may be sufficient;

Mix the Acid cautiously with the Spirit. Let the liquor distil until a black froth arises; then immediately remove the retort from the fire. Separate the lighter supernatant liquor from the heavier one, and expose the former to the air for a day. Add to it the Solution of Potash first mixed with water, and shake them together. Lastly, when sufficiently washed, separate the æthereal oil which subsides.

Process.—The proportion of sulphuric acid directed to be used, does not differ materially from that employed by Hennell or Serullas. The products of the distillation are æther, water, sulphurous acid, and a yellow oily fluid which floats upon the water. The formation of the æther and water has been already explained; the sulphurous acid results from the mutual decomposition of a portion of the sulphuric acid and alcohol, the black froth being charcoal deposited from the spirit. The yellow oily fluid by exposure to the air loses the æther with which it is mixed, by evaporation, and the residue after the action of the potash to separate the sulphurous acid, is æthereal oil, sometimes called heavy oil of wine to distinguish it from a lighter oil, which it yields by partial decomposition.

Properties.—Æthereal oil is a yellow fluid, somewhat resembling oil of lavender in appearance; it has a penetrating aromatic odour; its taste is rather sharp and bitter; in water it is insoluble, but dissolved by æther and by alcohol. Its specific gravity according to Hennell is 1.05; while Dumas states it to be 1.133, and he also observes that its composition is rather variable; this from Hennell's statement appears to depend upon the different quantities of a peculiar carburet of hydrogen which it is apt to contain, and which by long keeping separates in prismatic crystals. Chloride of barium when added to this oil gives no pre-

cipitate, but when they are heated together the oil is decomposed and sulphate of barytes is precipitated; the necessity of heat to produce this effect proves that the sulphuric acid is intimately combined with the other constituents of the oil.

When distilled with water, æthereal oil is vaporized and condensed unchanged, but when heated without it, decomposition ensues, and there comes over much thick oily matter, and a considerable quantity of sulphuric acid; in the retort there remain a little sulphuric acid and some charcoal.

Composition.—The analysis of Hennell gives as the composition of æthereal oil,

Sulphuric Acid..	38	or nearly one equivalent ..	40
Carbon	53.7	nine equivalents	54
Hydrogen	8.3	nine equivalents	9
	100.		103

According to Serullas, whose analysis differs but little from that of Liebig, it contains

Sulphuric Acid	55.00
Carbon.....	33.00
Hydrogen	6.18
Oxygen	5.42
	99.60

If this analysis be correct, or even nearly so, the oil examined by Hennell must have contained excess of the peculiar carburated hydrogen in solution, which, as has been noticed, he found on long keeping to be capable of crystallizing from it.

Serullas terms æthereal oil a *sulphate of æther*, and Dumas *sulphatic æther*, observing that if the name of sulphuric æther had not been already in use, it would be the proper appellation of æthereal oil. On this view of the subject the oil must be composed of

One equivalent of Sulphuric Acid	= 40	or	52.00
Four eq. } one eq. of æther or	Carbon 24		31.16
Five — } oxide of ethereum	Hydrogen 5		6.49
One — }	Oxygen 8		10.35
			77
Equivalent of Oleum Æthereum.....			100.

Except in the quantity of oxygen, this statement agrees nearly with the analysis of Serullas; and if it be correct, it appears that when æther is formed in the presence of a great excess of sulphuric acid, one equivalent of each combines to form æthereal oil or sulphate of æther; the symbols of which will be

Berzelius and Turner.. $H^5 C^4 + O + \ddot{S}$ or $\dot{E}l + \ddot{S}$.

Brande. $(4car + 5h + o) + (s')$ or $(eth + s')$.

Pharmacopœia Preparation.—Spiritus Ætheris Sulphurici compositus.

SPIRITUS ÆTHERIS NITRICI.

Spirit of Nitric Æther.

Spiritus Nitri dulcis, P.L. 1745.*Spiritus Ætheris nitrosi*, P.L. 1788.*Spiritus Ætheris nitrici*, P.L. 1809, P.L. 1824.

Take of Rectified Spirit three pounds,

Nitric Acid four ounces;

Add the Acid gradually to the Spirit and mix; then let thirty-two fluidounces distil.

Process.—It has been shown that when sulphuric acid acts upon alcohol, æther is produced without decomposing the acid; when, however, nitric acid is employed, both it and the alcohol suffer decomposition. It has been mentioned that nitric acid is composed of oxygen and azote; during its action upon alcohol it loses a portion of oxygen, and the pure æther yielded by its action, when separated in a mode presently to be described, from the water and undecomposed spirit with which it distils, is composed of, according to Dumas and Boullay,

One equivalent of Æther ..	=	37	or	49·33
Three equivalents of Oxygen	=	24	„	32·00
One equivalent of Azote ...	=	14	„	18·67

Equivalent	75	100·
-----------------	----	------

As an equivalent of nitric acid contains 5 equivalents of oxygen 40, and 1 equivalent of azote 14, it is evident that 1 equivalent by converting alcohol into æther loses 2 equivalents of oxygen 16, and by this it is reduced to hyponitrous acid, which combines with the equivalent of æther formed, constituting hyponitrous æther, or hyponitrite of æther, composed of

One equivalent of Hyponitrous Acid	=	38	or	50·66
One equivalent of Æther	=	37		49·34

Equivalent	75	100·
-----------------	----	------

The oxygen lost by the nitric acid produces various compounds with the elements of those portions of alcohol which are decomposed and yet not converted into æther; thus with its hydrogen it forms water, with the carbon, carbonic acid which is evolved, and oxalic acid which remains in the retort; with portions of hydrogen and carbon it gives rise to acetic and malic acids; and when the operation is long continued a quantity of nitric acid loses more oxygen so as to be reduced to nitric oxide gas, and probably even nitrous oxide and azote are evolved.

Properties.—The Spiritus Ætheris nitrici of the Pharmacopœia is a mixture of hyponitrous æther and alcohol; it is colourless, has a peculiar and rather fragrant æthereal odour; is very inflammable and volatile, producing much cold during evaporation. Its specific gravity should not exceed $\cdot 834$; but when the distillation is carried too far the product is specifically heavier, high-coloured, very acid so as to act strongly on litmus paper and decompose the alkaline carbonates with effervescence. It mixes with water and alcohol in all proportions. By keeping it also becomes more acid than when recently prepared.

Hyponitrous æther may be separated from the alcohol, water and uncombined acid, which the preparation of the Pharmacopœia contains, by digesting lime reduced to powder in it and subjecting the mixture to distillation. The pure hyponitrous æther thus procured is of a pale yellow colour; its smell is æthereal, and when diffused, not unlike that of ripe apples; its specific gravity at about 40° is $\cdot 886$. It is extremely volatile, and it boils at nearly 70° under the usual pressure; it undergoes ebullition even when held in the hand, by evaporation produces a great degree of cold, and yields a very inflammable vapour, which burns with a bright flame. It does not act upon litmus paper until it has suffered partial decomposition, which takes place spontaneously, especially when in contact with water, by which one part is dissolved and another decomposed; when mixed with a solution of potash it yields hyponitrite and a little acetate of potash and alcohol. The density of its vapour is $2\cdot 627$. When passed through a porcelain tube heated to redness hyponitrous æther yields nitric oxide gas, azotic gas, and cyanide of ammonia.

Symbols.—Berzelius and Turner .. $\text{El} + \ddot{\text{N}}$.

Brande $(\text{ETH} + n')$.

Impurities and Tests.—See Notes: SPIRITUS ÆTHERIS NITRICI.

Medicinal Uses.—Refrigerent: Diuretic. Dose, \mathfrak{M} x to \mathfrak{M} xl.

SPIRITUS ÆTHERIS SULPHURICI COMPOSITUS.

Compound Spirit of Sulphuric Æther.

Spiritus Ætheris vitriolici compositus, P.L. 1788.

Spiritus Ætheris compositus, P.L. 1809.

Spiritus Ætheris sulphurici compositus, P.L. 1824.

Take of Sulphuric Æther eight fluidounces,
Rectified Spirit sixteen fluidounces,
Æthereal Oil three fluidrachms ;

Mix.

Remarks.—This preparation is analogous to the *Liquor anodynus mineralis* of Hoffman. By the admixture of spirit with the æther and oil these are rendered miscible with water, and more convenient for exhibition.

Medicinal Uses.—Stimulant, Antispasmodic. Dose f3ss. to f3ij.

ALKALIS.

ACONITINA.

Aconitina.

Take of the Root of Aconite, dried and bruised, two pounds,

Rectified Spirit three gallons,

Diluted Sulphuric Acid,

Solution of Ammonia,

Purified Animal Charcoal, each as much as may be sufficient ;

Boil the Aconite with a gallon of the Spirit for an hour in a retort with a receiver adapted to it. Pour off the

liquor and again boil the residue with another gallon of Spirit, and with the Spirit recently distilled, and pour off the liquor. Let the same be done a third time. Then press the Aconite, and all the liquors being mixed and strained, let the Spirit distil. Evaporate what remains to the proper consistence of an extract. Dissolve this in water and strain. Evaporate the Liquor with a gentle heat, that it may thicken like a syrup. To this add of dilute Sulphuric Acid, mixed with water, as much as may be sufficient to dissolve the Aconitina. Then drop in the solution of Ammonia, and dissolve the Aconitina precipitated, in dilute Sulphuric Acid and water, mixed as before. Afterwards mix the Animal Charcoal, frequently shaking them during a quarter of an hour. Lastly, strain, and solution of Ammonia being again dropped in that the Aconitina may be precipitated, wash and dry it.

Remarks.—Aconitina is one of a numerous class of bodies called vegetable alkalis or alkaloids. It appears to have been first stated by Pallas in 1825, that aconite contains a peculiar alkali; and it has since been more particularly examined by Hesse. Like the other substances of this class it unquestionably exists in the aconite in combination with some vegetable acid, the nature of which has not however been determined. Whatever be the nature of this vegetable salt it is dissolved by alcohol with some portion of the colouring matter.

Properties.—This alkali crystallizes from a spirituous solution in granular crystals, but it is stated that this is not quite pure; indeed, the greater part is incapable of crystallizing; when obtained by evaporation it is a transparent colourless mass, with a glassy lustre: it has no smell; the taste is at first bitter, and afterwards acrid, but this is not permanent, and appears to be derived from another principle (anemonin), from which it may be separated, by repeated solution in and precipitation from acids.

Aconitina is soluble in 150 times its weight of cold, and 50 of boiling water. Alcohol and æther dissolve it in large quantity, and the solutions are decidedly alkaline; it does not change by exposure to the air; when heated moderately it fuses, and when strongly heated it is decomposed, yielding ammonia in the same manner as the other vegetable alkalis, like which it is composed of carbon, oxygen, hydrogen, and azote, but in proportions that have not yet been determined.

Although aconitina combines with acids to form salts, they do not, as far as they have been examined, crystallize, but dry into a gummy mass, their taste is very bitter; the solution of nitrate of aconitina is colourless, that of the sulphate is yellow at first and afterwards becomes of a dark violet; the alkalis decompose them, precipitating the aconitina.

This alkali is in the highest degree poisonous; the 50th part of a grain dissolved in spirit of wine killed a sparrow in a few minutes, and the 20th instantly; applied to the eye it occasions a temporary dilatation of the pupil.

Impurities and Tests.—See Notes: ACONITINA.

Medicinal Uses.—It is too powerful a medicine to be exhibited internally, but has been applied with success in the form of an ointment, in the proportion of one grain to a drachm of lard, in neuralgic affections. Very similar medicinal powers appear to reside in *Delphia*, an alkali obtained much more readily from the *Delphinium Staphisagria*.

LIQUOR AMMONIÆ.

Solution of Ammonia.

Aqua Ammoniæ, P.L. 1788.

Liquor Ammoniæ, P.L. 1809, P.L. 1824.

Take of Hydrochlorate of Ammonia ten ounces,

Lime eight ounces,

Water two pints;


Put the Lime slacked with water into a retort, then add the Hydrochlorate of Ammonia broken into small pieces, and the remainder of the water. Let fifteen fluid-ounces of solution of Ammonia distil.

Process.—Hydrochlorate of ammonia, frequently called muriate of ammonia and sal ammoniac, is a compound of hydrochloric acid and ammonia, or the volatile alkali. When mixed with lime and water and subjected to distillation the changes which occur are these: hydrochloric acid is composed of hydrogen and chlorine and the lime of oxygen and the metal calcium, or it is an oxide of the metal; when the hydrochlorate of ammonia and lime act upon each other, not only is the hydrochlorate decomposed so as to yield its ammonia in the gaseous state, but the hydrochloric acid and lime, or oxide of calcium, also undergo

decomposition; the chlorine of the acid and the calcium of the oxide combine to form chloride of calcium, which remains in the retort, while the hydrogen of the acid and the oxygen of the oxide form water, and the ammonia set at liberty is expelled, and being vaporized with the water, is condensed with it in the receiver.

Liquor Ammoniacæ, P.L.

Solution of Ammonia.



Ammoniacal gas is transparent, colourless, and of course invisible. Its smell is extremely pungent, and its taste acrid. Its sp. gr. compared with atmospheric air is as 0.5893 to 1, and 100 cubic inches weigh nearly 18.27 grains. An animal immersed in it is quickly killed; it extinguishes the flame of a taper, but it is enlarged before extinction. It is very rapidly condensed by water; the solution is colourless and transparent, and like the gas possesses properties which are most strongly alkaline, turning vegetable yellow colours brown, blues green, and by combining with acids it destroys their power of reddening vegetable blue colours. When subjected to a pressure of about 6.5 atmospheres at the temperature of 50°, ammoniacal gas was found by Faraday to become a colourless transparent fluid, having a sp. gr. of 0.760. When ammoniacal gas is mixed with oxygen gas, and fired by the taper, water is formed, and azotic gas left, and by being passed through a red hot tube, it is resolved into hydrogen gas and azotic gas. The aqueous solution decomposes by exposure to the air, and still more readily by heat, the ammonia being dissipated in the elastic or gaseous form.

Composition.—Ammoniacal gas is composed of 3 volumes of hydrogen gas and 1 volume of azotic gas, condensed into 2 volumes, or by weight it is composed of

Three equivalents of Hydrogen	$1 \times 3 =$	3	or	17.64
One equivalent of Azote	$\dots\dots\dots =$	14	,,	82.36

Equivalent	17.	100.
------------	------	-----	------

A solution of sp. gr. 0·960, as directed in the Pharmacopœia, is composed very nearly of

Ammoniacal gas	10
Water	90
	<hr/>
	100

Incompatibles.—Liquor Ammoniaë is incompatible with acids, acidulous and most earthy and metallic salts, but it does not decompose the salts of lime, barytes or strontia, those of magnesia only partially, and the potassio-tartrate of iron is not at all precipitated by it.

Impurities and Tests.—See Notes: AMMONIÆ LIQUOR.

Pharmacopœia Preparations.—Linimentum Ammoniaë, Spiritus Ammoniaë Succinatus, Linimentum Camphoræ compositum, Linimentum Hydrargyri.

Medicinal Uses.—Liquor Ammoniaë is stimulant, rubefaciant and antacid; it may be exhibited in milk, water, or any cold liquid which is not incompatible with it. Dose ℥ x. to ℥ xxx. If it should be swallowed by mistake, the best antidote is vinegar or lemon-juice.

AMMONIÆ SESQUICARBONAS.

Sesquicarbonate of Ammonia.

Sal Volatilis, P.L. 1720.

Sal Volatilis Salis Ammoniaci, P.L. 1745.

Ammonia præparata, P.L. 1788.

Ammonia Carbonas, P.L. 1809.

Ammonia Subcarbonas, P.L. 1824.


Take of Hydrochlorate of Ammonia a pound,

Chalk a pound and a half;

Rub them separately to powder; then mix, and with a heat gradually raised, sublime.

Process.—The reaction in this case is more complicated than that of the last; instead of lime, carbonate of lime is employed, and the product is consequently a carbonate of ammonia.

Ammoniae Sesquicarbonas, P. L.
Hydrated Sesquicarbonate of Ammonia.



Composition.—This salt consists, in its perfect state, of

Three eqs. of Carbonic Acid $22 \times 3 = 66$ or Carbonic Acid 55.93	
Two eqs. of Ammonia $17 \times 2 = 34$ „ Ammonia 28.81	
Two eqs. of Water $9 \times 2 = 18$ „ Water 15.26	
Equivalent....	118 100.

Instead, however, of regarding it as composed of 3 equivalents of acid and 2 equivalents of base, it is more convenient to consider it as in the Pharmacopœia, as constituted of $1\frac{1}{2}$ equivalent of carbonic acid united to 1 equivalent of ammonia and 1 of water. On this view its composition is thus stated :

One and a half equivalent of Carbonic Acid..	= 33
One equivalent of Ammonia.....	= 17
One equivalent of Water	= 9
Equivalent....	59

It is however to be remarked that both the hydrochlorate of ammonia and carbonate of lime are neutral compounds, that is, each consists of 1 equivalent of acid and 1 of base ; the production of sesquicarbonate of ammonia, which is a supersalt, instead of a neutral one, as usually happens when neutral compounds suffer mutual decomposition, is explained by supposing three equivalents of each salt to undergo decomposition, when, if no loss occurred in the operation, the carbonate of ammonia would be neutral and hydrated, consisting of

Three equivalents of Carbonic Acid..	$22 \times 3 = 66$
Three equivalents of Ammonia	$17 \times 3 = 51$
Three equivalents of Water.....	$9 \times 3 = 27$

During sublimation, however, one of the equivalents of the ammonia liberated and one of the water formed are dissipated; whilst the quantity of carbonic acid remaining undiminished, the carbonate actually sublimed consists of three equivalents of carbonic acid and only two of ammonia, which, as just shown, constitute it a sesquicarbonate.

Symbols,—Berzelius and Turner $H^3 N + 1\frac{1}{2} \ddot{C} + \dot{H}$.

Brande $n + 3h$ or $A + 1\frac{1}{2}car' + q$.

Properties.—When recently prepared, sesquicarbonate of ammonia is a colourless translucent mass of a striated crystalline appearance, and it is moderately hard. Its smell is pungent, and its taste sharp and penetrating; turmeric paper when held over it is turned of a reddish brown colour by the carbonate of ammonia which escapes. It is soluble in about four times its weight of cold water, and by hot water it is decomposed with effervescence. When the bottle which contains this salt is frequently opened, or if a small quantity of it be kept in a large bottle, it gradually becomes opaque and friable, and its pungency is much diminished; if it be exposed to the air for some time, it is rendered quite devoid of smell, owing to the volatilization of neutral carbonate of ammonia, bicarbonate being left; and it will be observed that a compound of three equivalents of carbonic acid and two of ammonia, is equal to one equivalent of neutral carbonate, which evaporates, and one of bicarbonate of ammonia, which remains as an inodorous salt combined with water, consisting of

Two eqs. Carbonic Acid . .	$22 \times 2 = 44$	or Carbonic Acid	55.70
One equivalent of Ammonia	= 17	„ Ammonia	21.52
Two equivalents of Water . .	$9 \times 2 = 18$	„ Water	22.78
	<hr/>		<hr/>
Equivalent	= 79		100.

Incompatibles.—Sesquicarbonate of ammonia is decomposed by acids, by potash and soda, and their carbonates; by lime, lime-water, solution of chloride of calcium, magnesia, alum, acidulous salts, as bitartrate and bisulphate of potash, and solutions of iron, except the potassio-tartrate; bichloride of mercury, the acetate and diacetate of lead, sulphate of iron and of zinc, are also incompatible with this salt. With sulphate of magnesia it affords no precipitate.

Official Preparations.—Liquor Ammoniae Sesquicarbonatis, Liquor Ammoniae Acetatis, Cupri Ammonio-sulphas, &c.

Medicinal Uses.—It is stimulant, antispasmodic, diaphoretic, powerfully antacid, and in large doses emetic. In the form of smelling salts it is useful in syncope and hysteria. It must not be kept in powdered mixtures, and although in the form of pill

its properties are longer retained, it is by no means an eligible mode of exhibiting it. Dose, gr. v. to xx. : xxx. grains are emetic.

LIQUOR AMMONIÆ SESQUICARBONATIS.

Solution of Sesquicarbonate of Ammonia.

Spiritus Salis Ammoniaci, P.L. 1720, P.L. 1745.

Aqua Ammoniacæ, P.L. 1788.

Liquor Ammoniacæ Carbonatis, P.L. 1809.

Liquor Ammoniacæ Subcarbonatis, P.L. 1824.

Take of Sesquicarbonate of Ammonia four ounces,

Distilled Water a pint ;

Dissolve the Sesquicarbonate of Ammonia in the Water,
and strain.

This solution ought not to be prepared in large quantities at a time ; for by keeping, or rather by occasional exposure to the air, its pungency and powers suffer diminution. Dose, ℥ xxx. to f3j. in any bland liquid. This solution is of course incompatible with the substances already named as such with the sesquicarbonate of ammonia.

LIQUOR AMMONIÆ ACETATIS.

Solution of Acetate of Ammonia.

Aqua Ammoniacæ Acetataæ, P.L. 1788.


Liquor Ammoniacæ Acetatis, P.L. 1809, P.L. 1824.

Take of Sesquicarbonate of Ammonia four ounces and
a half, or as much as may be sufficient,

Distilled Vinegar four pints ;

Add the Sesquicarbonate of Ammonia to the Vinegar
to saturation.

Process.—This is a case of single elective affinity and decomposition. In preparing this solution, carbonic acid gas is evolved owing to the stronger affinity of the acetic acid for the ammonia with which it was combined, and acetate of ammonia is formed and remains in solution.



If the sesquicarbonate of ammonia have become opake by exposure to the air, a larger quantity will be required, on account of its having been partly converted into bicarbonate of ammonia. It is better that the acid should appear to be in excess than the alkaline salt; for the carbonic acid which remains for some time in solution, and which seems to indicate excess of acetic acid, is eventually dissipated; it is owing to the presence of this acid that solution of acetate of ammonia, when mixed with that of diacetate of lead, often gives a white precipitate of carbonate of lead, and a fallacious appearance of the presence of sulphuric acid in the distilled vinegar used. Vinegar which has been condensed in a metallic worm, affords a dark-coloured precipitate when employed in preparing solution of acetate of ammonia.

Incompatibles.—Acids; potash, soda and their carbonates; lime and lime-water; the acetate and diacetate of lead also, on account of the carbonic acid which usually remains diffused through the *Liquor Ammoniae Acetatis*, are incompatible with it, and they are especially so, if it contain undecomposed sesquicarbonate of ammonia; carbonate of lead being in both cases precipitated.

Tests and Impurities.—See Notes: *AMMONIAE ACETATIS LIQUOR.*

Medicinal Uses.—This preparation is not unfrequently employed as a collyrium, in which case it is especially requisite that there should be no excess of sesquicarbonate of ammonia. When assisted by warmth and plentiful solution, it is an excellent diaphoretic, and in some cases it acts as a diuretic. Dose, fʒiv. to fʒvi. Externally as a lotion it is refrigerant.

MORPHIA.

Morphia.

Take of Hydrochlorate of Morphia an ounce,
 Solution of Ammonia five fluidrachms,
 Distilled Water a pint ;

Add the Hydrochlorate of Morphia first dissolved in the pint of water to the solution of Ammonia with an ounce of water, shaking them together. Wash what is thrown down with distilled water, and dry it with a gentle heat.

Remarks.—Morphia was the first discovered of the vegetable alkalis. It was obtained from opium by Sertuerner in 1803 ; it exists in this substance in combination with a peculiar vegetable acid called the *Meconic Acid*, and probably also with sulphuric acid ; for the process by which the hydrochlorate is obtained, see MORPHIÆ HYDROCHLORAS.

Process.—When ammonia is added to the solution of hydrochlorate of morphia, this alkali having greater affinity for the acid than the morphia has, hydrochlorate of ammonia is formed, and the morphia being quite or nearly insoluble in water is precipitated.

Properties.—Morphia is precipitated by the ammonia in a flocculent state, and on stirring and standing it assumes a crystalline appearance ; it has a bitter taste. According to Berzelius it is insoluble in cold water, and boiling water dissolves rather more than $\frac{1}{100}$ of its weight, and on cooling the solution yields crystals ; the hot solution turns turmeric paper brown, thus evincing its alkaline property. It is soluble in 40 parts of cold anhydrous alcohol and 30 parts when boiling ; in æther it is nearly insoluble ; it is soluble in the volatile and fixed oils. Potash and soda dissolve it in considerable quantity, and ammonia in smaller proportion. When heated strongly in the air it emits a resinous smell, smokes and burns with a lively red flame, and leaves charcoal.

Composition.—Morphia has been repeatedly analysed ; the results do not greatly differ ; it appears to consist of, in its anhydrous state,

Thirty-four equivalents of Carbon	$6 \times 34 = 204$	or 71.83
Eighteen equivalents of Hydrogen	$1 \times 18 = 18$	„ 6.33
Six equivalents of Oxygen.....	$8 \times 6 = 48$	„ 16.90
One equivalent of Azote	14	„ 4.94

Equivalent.... 284 100.

The crystals contain two equivalents of water or

One equivalent of Morphia .. 284 or 94.04


Two equivalents of Water.... 18 „ 5.96

Equivalent.... 302 100.

Symbols,—Berzelius and Turner $C^{34} H^{18} O^6 N^1$.

Brande..... $(34car + 18h + 6o + n)$ or MOR.

The crystals obtained by spontaneous evaporation from alcohol have a pearly lustre, and their primary form is a *right rhombic prism*, only the lateral planes of which appear on the crystals; one cleavage only has been obtained parallel to the plane *h*.



M on M'	127° 20'
M on <i>h</i>	116 20
<i>h</i> on <i>c</i>	132 20
<i>c</i> on <i>c'</i>	95 20

MORPHIÆ ACETAS.

Acetate of Morphia.

Take of Morphia six drachms,

Acetic Acid three fluidrachms,

Distilled Water four fluidounces;

Mix the Acid with the Water and pour them upon the Morphia to saturation. Let the liquor evaporate with a gentle heat that crystals may be formed.

Properties.—Acetate of morphia crystallizes with some difficulty in nearly colourless radiating needles, and during the evaporation of the solution a part of the acid is sometimes dissipated, and a portion of the acetate, suffering at least partial decomposition, becomes insoluble in water, and requires an addition of acetic acid to dissolve it; it is very soluble in water, especially when there is a slight excess of acetic acid, and less so in alcohol. It

is decomposed, like the other salts of morphia, by ammonia, potash, soda, &c., the morphia being precipitated.

Composition.—Acetate of morphia is probably composed of

One equivalent of Acetic Acid.. = 51 or 15.23

One equivalent of Morphia = 284 „ 84.77

Equivalent.... 335 100.

It has not, I believe, been determined whether the crystals contain water or not, and consequently, whether the crystallized salt differs in power from that obtained, as it commonly is, by evaporation to dryness.

Tests and Impurities.—See Notes : MORPHIÆ ACETAS.

Symbols.—Berzelius and Turner $C^{34} H^{18} O^6 N^1 + C^4 H^3 O^3$ or \bar{A} .

Brande (MOR + *ac'*.)

Incompatibles.—The stronger acids, the alkalis and alkaline earths, and many or most earthy and metallic salts.

Medicinal Uses.—Morphia is perhaps the most active principle of opium, but owing to its sparing solubility in water is never used alone medicinally. Its combinations with either the acetic or hydrochloric acids form salts, which being soluble admit of being exhibited in very small doses and with great effect. The advantage which they seem to possess over opium is chiefly ascribed to the absence of narcotina; their use in ordinary cases not being followed by either headache or sickness. The dose is from gr. $\frac{1}{8}$ to gr. $\frac{1}{4}$.

MORPHIÆ HYDROCHLORAS.

Hydrochlorate of Morphia.

Take of Opium, sliced, a pound,

Crystals of Chloride of Lead two ounces, or as much as may be sufficient,

Purified Animal Charcoal three ounces and a half,

Hydrochloric Acid,

Distilled Water,

Solution of Ammonia, each as much as may be sufficient ;

Macerate the Opium in four pints of distilled Water


for thirty hours, and bruise it; afterwards digest it for twenty hours more and press it. Macerate what remains again and a third time in water, that it may become free from taste, and as often bruise and press it. Evaporate the mixed liquors, with a heat of 140° , to the consistence of a syrup. Then add three pints of distilled Water, and when the impurities have subsided pour off the supernatant liquor. Gradually add to this two ounces of Chloride of Lead, or as much as may be sufficient, first dissolved in four pints of boiling distilled Water, till nothing further is precipitated. Pour off the liquor and wash what remains frequently with distilled Water. Then evaporate the mixed liquors as before, with a gentle heat, that crystals may be formed. Press these in a cloth, then dissolve them in a pint of distilled Water, and digest with an ounce and a half of Animal Charcoal, in a heat of 120° , and strain. Lastly, the Charcoal being washed, evaporate the liquors cautiously that pure crystals may be produced. To the liquor poured off from the crystals first separated, previously mixed with a pint of water, gradually drop in as much Solution of Ammonia, frequently shaking it, as may be sufficient to precipitate all the Morphia. To this, washed with distilled Water, add Hydrochloric Acid that it may be saturated: afterwards digest it with two ounces of Animal Charcoal and strain. Lastly, the Animal Charcoal being thoroughly washed, evaporate the liquors cautiously, that pure crystals may be produced.

Remarks.—Opium, the inspissated juice of the *Papaver somniferum*, contains various secondary principles, consisting of acids, alkalis, and neutral compounds; those which are peculiar are

Meconic Acid.	
Morphia.	
Paramorphia or Thebaia.	} Alkalis.
Codeia.	
Narcotina.	
Narceia.	
Meconin.	} Neutral.

Besides these it contains several other secondary principles, some of which are met with in other vegetable products, viz. sulphuric acid, potash, lime, gum, bassorin, caoutchouc, lignin, extract, fixed oil, and a volatile principle. For an account of these I refer to works on chemistry; in a medicinal point of view morphia is the only substance of much importance.

Process.—Morphia exists in opium combined principally with meconic acid, forming meconate of morphia; the first step in the process is to procure an aqueous solution of the soluble portion of opium, and this contains the meconate of morphia. When a solution of chloride of lead is added to it, it decomposes and is decomposed by water, owing to the interference of the meconate; the hydrogen of the water unites with the chlorine to form hydrochloric acid, and its oxygen with the lead to form oxide of lead; the morphia of the meconate combines with the hydrochloric acid, and constitutes with it hydrochlorate of morphia, which remains in solution, and the oxide of lead unites with the meconic acid of the meconate, and forms meconate of lead, which is precipitated.



As the solution contains some sulphuric acid, a little sulphate of lead will also be formed, and precipitated with the meconate of lead.

A quantity of hydrochlorate of morphia remains in solution, after the separation of the crystals, which is directed to be decomposed by ammonia, as explained under the head of Morphia, and this being redissolved in hydrochloric acid yields crystals of the hydrochlorate by evaporation. The animal charcoal is em-

ployed to render the salt colourless, which it effects by its well-known decolorizing power.

Properties.—Hydrochlorate of morphia, commonly called muriate of morphia, is a colourless, inodorous, bitter salt which crystallizes in plumose acicular crystals; it is soluble in 16 to 20 times its weight of water, and when boiling water is saturated with it, a crystalline mass is formed as the solution cools. It is also soluble in alcohol. It is totally decomposed and dissipated by exposure to a red heat. The crystals are anhydrous and consist of

One equivalent of Hydrochloric Acid..	=	37	or	11.52
One equivalent of Morphia.....	=	284	,,	88.48
				<hr/>
Equivalent		321		100.

Symbols.—Berzelius and Turner.. $C^{34} H^{18} N^1 O^6 + HCl$.

Brande ($MOR + mur'$.)

Medicinal Uses.—The hydrochlorate may be justly preferred to the acetate of morphia, in as much as it is more easily obtained in a crystallized form, and is not so subject to decomposition, during its preparation.

QUINÆ DISULPHAS.

Disulphate of Quina.

Take of Heart-leaved Cinchona, bruised, seven pounds,
Sulphuric Acid nine ounces,
Purified Animal Charcoal two ounces,
Hydrated Oxide of Lead,
Solution of Ammonia,
Distilled Water, each as much as may be
sufficient;

Mix four ounces and two drachms of the Sulphuric Acid with six gallons of distilled Water, and add the Cinchona to them; boil for an hour and strain. In the same manner again boil what remains in Water and Acid, mixed in the

same proportions, for an hour. Lastly, boil the Cinchona in eight gallons of distilled water for three hours, and strain. Wash what remains frequently with boiling distilled water. To the mixed liquors add Oxide of Lead while moist, nearly to saturation. Pour off the supernatant liquor, and wash what is thrown down with distilled water. Boil the liquors for a quarter of an hour, and strain; then gradually add Solution of Ammonia to precipitate the Quina. Wash this until nothing alkaline is perceptible. Let what remains be saturated with the rest of the Sulphuric Acid, diluted. Afterwards digest with two ounces of Animal Charcoal, and strain. Lastly, the Charcoal being thoroughly washed, evaporate the liquor cautiously, that crystals may be produced.

Remarks.—The different varieties of Cinchona contain two vegetable alkalis, Quina and Cinchonina: the *Cinchona cordifolia*, or yellow bark, contains chiefly the former; the *Cinchona lancifolia*, or pale bark, the latter; and the *Cinchona oblongifolia*, or red bark, yields both of them. The Quina exists in combination with a peculiar acid called Kinic Acid, forming with it Kinate of Quina, which is soluble to a certain extent in water, and is rendered more so by the sulphuric acid employed in the process, and perhaps by decomposing it. Whatever may be the state of combination, the solution contains sulphuric acid, kinic acid, and quina, mixed with extractive and colouring matter, the latter being got rid of by the animal charcoal. On adding oxide of lead the sulphuric acid combines with it, and the resulting sulphate being insoluble is precipitated, while the kinic acid and quina remain in solution; when ammonia is added after the separation of the sulphate of lead, the kinic acid unites with it and the kinate of ammonia formed is soluble, while the quina is precipitated, and this when afterwards combined with sulphuric acid forms disulphate of quina, which crystallizes.

Sulphuric Acid.
Oxide of Lead.

Kinic Acid.
Ammonia.

Quina.
Sulphuric Acid.

Sulphate of Lead. Kinate of Ammonia. Sulphate of Quina.

Properties and Composition of Quina.—When precipitated from pure sulphate of quina by ammonia it is flaky, colourless, inodo-

rous, and very bitter. It is scarcely soluble in water; to proper tests it is alkaline, and saturates acids. It is soluble in alcohol, and the solution by spontaneous evaporation in a dry place during winter yields small crystals, which contain water. Crystals may also be obtained by dissolving quina in weak alcohol, and adding water till the solution begins to be milky; after some days' exposure to the air a fluid of a resinous appearance separates, which gradually becomes radiated acicular crystals. Quina is also soluble in æther, and slightly in the volatile and fixed oils, when heated.

When quina is dried it is apt to become brown; when more strongly heated it becomes still darker-coloured, fuses, is rendered viscid, decomposes with the formation and evolution of ammonia, and charcoal remains; and this is dissipated when ignited in the air.

Quina is composed of

Twenty equivalents of Carbon..	$6 \times 20 = 120$	or	74
Twelve equivalents of Hydrogen	$1 \times 12 = 12$	„	7.4
Two equivalents of Oxygen....	$8 \times 2 = 16$	„	9.9
One equivalent of Azote.....	$= 14$	„	8.7
	<hr/>		<hr/>
Equivalent	162		100.

Symbol,—Berzelius and Turner .. $C^{20} H^{12} O^2 N^1$.

Brande..... $(20car + 12h + 2o + n.)$

Sulphates of Quina.—The sulphate composed of one equivalent of acid and base, though neutral in composition, is acid to litmus paper, but is not sour to the taste. It may be prepared by crystallizing a solution of sulphate of quina, which has not dissolved as much of the alkali as it is capable of taking up. It effloresces when exposed to the air. It crystallizes in square prisms; it is soluble in 11 times its weight of water at 55° , and in 22 times at 73° . At 212° it fuses in its water of crystallization. It is soluble in alcohol, and totally destroyed by ignition.

It is composed of

One equivalent of Sulphuric Acid	$= 40$	or	14.6
One equivalent of Quina	$= 162$	„	59.1
Eight equivalents of Water 9×8	$= 72$	„	26.3
	<hr/>		<hr/>
	274		100.

This salt is not employed in medicine.

Properties of Disulphate of Quina.—The crystals are colourless, acicular, have a pearly lustre, a bitter taste, and effloresce when exposed to the air. One part of this salt requires for so-

lution about 740 parts of cold, 30 of boiling water, 80 of cold alcohol of specific gravity .850, and much less if boiling. When heated, disulphate of quina fuses and looks like melted wax; it afterwards reddens, begins to decompose, and when the heat is raised to ignition in the air, charcoal is obtained, which is eventually dissipated.

Composition.—This salt consists of

One equivalent of Sulphuric Acid..	=	40	or	9.17
Two equivalents of Quina	$162 \times 2 =$	324	„	74.31
Eight equivalents of Water	$9 \times 8 =$	72	„	16.52
		<hr/>		<hr/>
Equivalent.....		436		100.

By exposure to the air or the temperature of boiling water, half the water is expelled, and when heated to 240° it loses half the remainder, retaining only two equivalents; it is questionable whether more can be expelled without at the same time decomposing the salt.

Symbol,—Berzelius and Turner $2C^{20}H^{12}O^2N^1 + \ddot{S} + 8H.$

Brande $(2QUI + S' + 8q.)$

Impurities and Tests.—See Notes: QUINÆ DISULPHAS.

Incompatibles.—Alkalis, their carbonates and lime-water; these separating the sulphuric acid and precipitating the quina. The soluble salts of barytes, lead, &c. precipitate the sulphuric acid.

Medicinal Uses and Dose.—Quinæ Disulphas is a prominent example of the advantages which Medicine has derived from Chemistry. It possesses all the virtues of the cinchona, unmixed with inert or superfluous substances, such as the woody fibre or resin. Dose, from gr. iij. to gr. x.

STRYCHNIA.

Strychnia.


Take of Nux vomica, bruised, two pounds,
 Rectified Spirit three gallons,
 Diluted Sulphuric Acid,
 Magnesia,
 Solution of Ammonia, each as much as may be
 sufficient;

Boil the bruised *Nux vomica* with a gallon of the Spirit for an hour in a retort, to which a receiver is fitted. Pour off the liquor, and again and a third time boil what remains with another gallon of Spirit, and the Spirit recently distilled, and pour off the liquor. Press the *Nux vomica*, and let the Spirit distil from the mixed and strained liquors. Evaporate what remains to the proper consistence of an extract. Dissolve this in cold Water and strain. Evaporate the liquor with a gentle heat until it has the consistence of a Syrup. To this while yet warm gradually add the Magnesia to saturation, shaking them together. Set it aside for two days, then pour off the supernatant liquor. Press what remain wrapped in cloth. Boil it in Spirit, then strain, and let the Spirit distil. Add to the residue a very little diluted Sulphuric Acid mixed with Water, and macerate with a gentle heat. Set it aside for twenty-four hours that crystals may form. Press and dissolve them. Afterwards to these dissolved in Water add Ammonia, frequently shaking them, that the Strychnia may be thrown down. Lastly, dissolve this in boiling Spirit, and set it aside that crystals may be produced.

Remarks.—The vegetable alkali Strychnia was discovered in 1818 by Pelletier and Caventou, who procured it from the *Strychnos nux vomica*, and hence its name. It is contained also in some other species of *Strychnos*, and exists in them combined with an acid originally, and yet sometimes called, *Igasuric Acid*, but now usually termed *Strychnic Acid*. The *upas* also contains strychnia.

Process.—*Nux vomica* consists of strychnate of strychnia, strychnate of brucia, which is another vegetable alkali, colouring matter, gum, bassorin, starch, wax, fixed oil and lignin; when reduced to powder and digested in spirit, the vegetable salt is dissolved with some admixture. After distilling the alcohol, the magnesia added decomposes the strychnate of strychnia, the base of which remains mixed with the magnesian salt formed, and any excess of its base employed; when this mixture is again digested in spirit, the strychnia is dissolved: and this being left after distillation is taken up by dilute sulphuric acid, and the resulting sulphate of strychnia is crystallized, dissolved in water, and decomposed by ammonia, which, combining with the sulphuric acid,

the strychnia on account of its insolubility precipitates, and this again dissolved in boiling spirit, crystallizes by cooling and spontaneous evaporation.



Properties.—Strychnia is colourless, inodorous, crystalline, unalterable by exposure to the air; it is so extremely bitter as to impart that taste to 600,000 times its weight of water. It requires about 6600 times its weight of cold, and 2500 times its weight of boiling water for solution. It is insoluble in absolute alcohol or in æther; alcohol even of '820 specific gravity scarcely dissolves any when cold, but in diluted alcohol it is more soluble. By rapid evaporation of the alcoholic solution it is deposited in a granular state, but by spontaneous evaporation it is procured in the form of the octahedron and square prisms, terminated by flat four-sided pyramids.

It acts like the alkalis on vegetable colours, and neutralizes acids, and forms salts with them.

It is extremely poisonous; one eighth of a grain is sufficient to kill a dog, and a quarter of a grain produces a decided effect upon a man. When heated it is decomposed, and yields the same products as similarly constituted compounds. As usually obtained it is probably mixed with some *brucia*, another extremely powerful vegetable alkali.

Composition.—It is composed of

Thirty equivalents of Carbon.....	$6 \times 30 = 180$	or
Sixteen equivalents of Hydrogen ..	$1 \times 16 = 16$	
Three equivalents of Oxygen.....	$8 \times 3 = 24$	
One equivalent of Azote.....	$= 14$	

Equivalent.... 234

Symbol,—Berzelius and Turner.. $C^{30}H^{16}O^3N^1$.

Brande $(30car + 16h + 3o + n.)$

Impurities and Tests.—See Notes: STRYCHNIA.

Incompatibles.—Acids and acidulous salts, which neutralize it. It probably decomposes some earthy and metallic salts by combining with their acid.

Medicinal Use.—This alkali, which is one of the most virulent furnished by the vegetable kingdom, has been exhibited with very variable results in paralytic affections. Dose, gr. $\frac{1}{16}$ to gr. $\frac{1}{8}$.

VERATRIA.

Veratria.

Take of Cevadilla, bruised, two pounds,
Rectified Spirit three gallons,
Diluted Sulphuric Acid,
Solution of Ammonia,
Purified Animal Charcoal,
Magnesia, each, as much as may be sufficient;

Boil the Cevadilla with a gallon of the Spirit for an hour in a retort, to which a receiver is fitted. Pour off the liquor, and again boil what remains with another gallon of Spirit and the Spirit recently distilled, and pour off the liquor: and let it be done a third time. Press the Cevadilla and let the Spirit distil from the mixed and strained liquors. Evaporate what remains to the proper consistence of an extract. Boil this three or more times in Water, to which a little diluted Sulphuric Acid is added, and with a gentle heat evaporate the mixed liquors to the proper consistence of a syrup. To this, when cold, put in the Magnesia to saturation, frequently shaking them; then press and wash. Let this be done two or three times; then dry what remains, and digest with a gentle heat in Spirit two or three times, and strain as often. Afterwards let the Spirit distil. Boil the residue in water, to which a little Sulphuric Acid and Animal Charcoal are added, for a quarter of an hour, and strain. Lastly, the Charcoal being thoroughly washed, evaporate the mixed liquors cautiously until they have the consistence of a syrup, and add to them as much Ammonia as may be sufficient to throw down the Veratria. Wash and dry it.

Remarks.—Veratria is a vegetable alkali, which, as well as Strychnia, was discovered by Pelletier and Caventou, in 1819. It was originally procured, as its name imports, from the *Veratrum*

album, or white hellebore; it is now generally and with greater facility obtained from *Cevadilla*, the seeds of the *Helonias officinalis*. In both these substances the alkali is combined with gallic acid, forming gallate of veratria.

Process.—By being boiled in spirit as directed, the cevadilla yields gallate of veratria, colouring matter, and some other compounds. When the residue, after the distillation of the alcohol, is treated with sulphuric acid, sulphate of veratria is formed; this is decomposed by magnesia, and the veratria set free is separated from much of the matter with which it is mixed by digestion and solution in spirit; when this is distilled off the veratria is treated with sulphuric acid and animal charcoal, by which the sulphate of veratria is decolorized, and ammonia then added to it, sulphate of ammonia is formed, and veratria, on account of its insolubility, is precipitated. Substituting veratria for strychnia, &c., the diagram illustrating the preparation of that alkali will serve for this also.

Properties.—This alkali is colourless and pulverulent; it is not crystallizable. It produces violent and dangerous sneezing; the taste is extremely acrid and burning, but free from bitterness. When taken internally it excites nausea and vomiting, and proves fatal to animals in small doses. Authors differ as to its point of fusion, but it becomes on cooling a transparent yellowish mass. When ignited in the air it is totally decomposed and dissipated. In cold water it is nearly insoluble, and boiling water dissolves only 1-1000dth of its weight, and the solution is acrid; alcohol dissolves veratria very readily, but æther takes it up sparingly. Veratria possesses the alkaline property of restoring the blue colour of litmus paper which has been reddened by acids, and also that of saturating and forming salts with acids, which crystallize with great difficulty. According to Couerbe the veratria thus prepared contains at least two other principles, viz. sabadilline and veratrine.

Composition.—Veratria consists of

Thirty-four equivalents of Carbon	$6 \times 34 = 204$	or 70.83
Twenty-two equivalents of Hydrogen..	$1 \times 22 = 22$	„ 7.63
Six equivalents of Oxygen	$8 \times 6 = 48$	„ 16.66
One equivalent of Azote	$= 14$	„ 4.88
	<hr/>	
	288	100.

Symbol,—Berzelius and Turner.. $C^{34}H^{22}O^6N^1$.

Brande $34car + 22h + 6o + n$.

Impurities and Tests.—See NOTES: VERATRIA.

Incompatibles.—See *Strychnia*.

Medicinal Use.—This very powerful alkali in moderate doses increases all the secretions, and has been thought efficient, carefully exhibited, in gout and rheumatism. Dose gr. $\frac{1}{8}$ to gr. $\frac{1}{4}$.

ANIMALIA.

Preparations from Animals.

CARBO ANIMALIS PURIFICATUS.

Purified Animal Charcoal.

Take of Animal Charcoal a pound,
Hydrochloric Acid,
Water, each, twelve fluidounces;

Mix the Hydrochloric Acid with the Water, and pour it gradually upon the Charcoal, then digest for two days with a gentle heat, frequently shaking them. Set by, and pour off the supernatant liquor, then wash the Charcoal very often with water, until nothing acid is perceptible; lastly, dry it.

Remarks.—Bone is principally a compound of phosphate and carbonate of lime and gelatine; when it is perfectly calcined with access of air, the whole of the gelatine is dissipated, and there remains a white mixture of phosphate and carbonate of lime; when, however, the heat is applied to the bone in vessels with small apertures, and it is not too long continued, a portion of the charcoal of the gelatine remains mixed with the phosphate and carbonate of lime, and constitutes animal charcoal, usually termed *ivory black*. This charcoal possesses in a very high degree the power of removing animal and vegetable colouring matter, and is on this account largely used in sugar-refining; with this application of it, the carbonate and phosphate of lime do not interfere, and therefore are not separated. In several of the cases, however, in which it is applied to chemical uses, these earthy compounds would be acted upon by the acids which the solutions to

be decolorized contain, as in preparing Veratria, &c. The hydrochloric acid is therefore employed to dissolve the earthy phosphate and carbonate, and this it does without acting upon the charcoal. The solution contains chloride of calcium and phosphate of lime.

CORNU USTUM.

Burnt Horn.

Burn pieces of Horns in an open vessel until they become perfectly white; then powder and prepare them in the same manner as directed with respect to Chalk.

Remarks.—In this operation the whole of the gelatine of the horn is decomposed and dissipated, and the residue is phosphate of lime very nearly in a state of purity.

Properties.—Phosphate of lime is soluble in most acids without decomposition, and is precipitated from them by ammonia, potash and their carbonates in a gelatinous state, but unaltered in composition. It is however decomposed by sulphuric acid, the results being a precipitate of sulphate of lime, and phosphoric acid which remains in solution. Phosphate of lime is perfectly insoluble in water, and unchangeable by heat, even when carbonaceous matter is present.

Composition.—Bone phosphate, sometimes called bone earth, is a subphosphate of lime, consisting, according to Berzelius, of three equivalents of acid and eight of lime; employing however the equivalent weights generally adopted by English chemists, it is a subsesquiphosphate of lime, containing

One equivalent of Phosphoric Acid..... = 36 or 46.15

One and a half equivalent of Lime $28 + 14 = 42$ „ 53.85

Equivalent.... $\frac{78}{100}$

Symbols.—Berzelius and Turner $\text{Ca}^8 \text{P}_3$.

Brande $1\frac{1}{2}c + p'$.

Medicinal Use.—It forms a part of the Pulvis Antimonii compositus.

TESTÆ PRÆPARATÆ.

Prepared Shells.

Wash the Shells, first freed from impurities, with boiling water; then prepare them in the same manner as directed for Chalk.

Remarks.—Shell consists principally of carbonate of lime, but it is mixed with indurated albumen, which is the cause of the smell of burnt horn experienced when they are put into the fire.

Prepared shells are harder than prepared chalk and less readily acted upon by acids, and probably the animal matter which they contain retards their action. Prepared shells were formerly used in the *Confectio aromatica*, but prepared chalk is now substituted.

AQUÆ DESTILLATÆ.

Distilled Waters.

AQUA DESTILLATA.

Distilled Water.

Take of Water ten gallons;

First let two pints distil, which being thrown away, let eight gallons distil. Keep the distilled Water in a glass bottle.

Remarks.—Most spring and river water contain impurities in solution; these are generally carbonic acid, carbonate of lime, sulphate of lime, and common salt. There are some preparations whose power is much diminished, and whose solutions are rendered turbid by these compounds. Such, more especially, are lime-water, acetate and diacetate of lead; and sulphate of iron is even decomposed by the atmospheric air which water always contains. Water may be nearly deprived of carbonic acid, carbonate of lime, and atmospheric air, by mere ebullition; but at the same time, owing to the evaporation which takes place, the proportion of the other impurities is increased, and therefore water

which has been long boiled, may be more impure even than before ebullition.

The following tests will determine the presence of the usual impurities :—

Lime Water.—If carbonic acid be present, this will cause precipitation of carbonate of lime before ebullition, but not after it.

Chloride of Barium.—If sulphate of lime be present, this will give a precipitate of sulphate of barytes insoluble in nitric acid.

Oxalate of Ammonia.—If this give a precipitate of oxalate of lime before the water is boiled, it may be owing to the presence either of carbonate or of sulphate of lime ; but if only after ebullition, then to the presence of sulphate, provided chloride of barium gives also a precipitate.

Nitrate of Silver.—If common salt or any other chloride be contained in water, this re-agent will afford a precipitate of chloride of silver insoluble in nitric acid.

Properties.—Distilled water is colourless, transparent, inodorous, tasteless, and also vapid on account of the absence of air. No change occurs in its appearance on the addition of lime-water, oxalate of ammonia, nitrate of silver, diacetate of lead, or hydrosulphuric acid. A pint weighs, at 62°, 8750 grains, or one pound six ounces one drachm two scruples and a half, or ten grains less than eighteen ounces and a quarter.

Few chemists are, I believe, in the practice of keeping a still for the purpose of distilling water only ; yet this ought to be done, or the distilled water will have a faint smell and taste of the last herbs which had been subjected to distillation.

AQUA ANETHI.

Dill Water.

Aqua Seminum Anethi, P.L. 1748.

Aqua Anethi, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Dill Seeds, bruised, a pound and a half,

Proof Spirit seven fluidounces,

Water two gallons ;

Let a gallon distil.

Remarks.—The odour and pungency of plants frequently reside in an essential oil, and this has its volatility so much increased

by the vapour of the boiling water, that they rise together in distillation, and a sufficient quantity of the oil is either dissolved by or intimately mixed with the water to impart the peculiar taste and smell of the plant, or the parts of it employed.

Distilled waters in some cases, however, are mixed with other principles besides the volatile oil ; thus cinnamon water contains, according to Soubeiran, cinnamic acid ; valerian and pepper water are not inserted in the Pharmacopœia, but the first contains acetic and valerianic acid ; and the latter ammonia, according to Vauquelin. Distilled waters, it will be observed, are in some cases prepared directly from the oil instead of the plant which yields it.

Waters distilled from herbs are intended merely as vehicles for the exhibition of more important remedies ; when they have been long kept, they undergo a kind of decomposition, and become mucilaginous and sour : this is intended to be prevented by the addition of a small quantity of spirit, which in former Pharmacopœias was directed to be mixed with the distilled product, but is now ordered to be distilled with the water.

AQUA CARUI.

Carraway Water.

Aqua Seminum Carui, P.L. 1745.

Aqua Carui, P.L. 1809, P.L. 1824.

AQUA FŒNICULI.

Fennel Water.

Aqua Fœniculi, P.L. 1788, P.L. 1745, P.L. 1809, P.L. 1824.

These are prepared in the same manner as Dill Water.

AQUA FLORUM AURANTII.

Orange Flower Water.

Take of Orange Flowers ten pounds,
Proof Spirit seven fluidounces,
Water two gallons ;
Let a gallon distil.

AQUA CINNAMOMI.

Cinnamon Water.

Aqua Cinnamomi tenuis, P.L. 1720.

Aqua Cinnamomi simplex, P.L. 1745.

Aqua Cinnamomi, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Cinnamon, bruised, a pound and a half, *or*
Oil of Cinnamon two drachms,
Proof Spirit seven fluidounces,
Water two gallons ;
Let a gallon distil.

AQUÆ MENTHA PIPERITÆ.

Peppermint Water.

Aqua Menthæ Piperitidis simplex, P.L. 1745.

Aqua Menthæ Piperitidis, P.L. 1788.

Aqua Menthæ Piperitæ, P.L. 1809, P.L. 1824.

Take of Peppermint, dried, two pounds, *or*
Oil of Peppermint two drachms,
Proof Spirit seven fluidounces,
Water two gallons ;
Let a gallon distil.

AQUA MENTHÆ PULEGII.

Pennyroyal Water.

Aqua Pulegii simplex, P.L. 1745.

Aqua Pulegii, P.L. 1788, P.L. 1809, P.L. 1824.

AQUA MENTHÆ VIRIDIS.

Spearmint Water.

Aqua Menthæ vulgaris simplex, P.L. 1745.

Aqua Menthæ sativæ, P.L. 1788.

Aqua Menthæ viridis, P.L. 1809, P.L. 1824.

These are prepared in the same manner as Peppermint Water; but when the fresh herb is employed in distilling either that or these, double the weight is to be used.

AQUA PIMENTÆ.

Pimenta Water.

Aqua Piperis Jamaicensis, P.L. 1745.

Aqua Pimento, P.L. 1788.

Aqua Pimentæ, P.L. 1809, P.L. 1824.

Take of Pimenta, bruised, a pound, *or*

Oil of Pimenta two drachms,

Proof Spirit seven fluidounces,

Water two gallons;

Let a gallon distil.

AQUA ROSÆ.

Rose Water.

Aqua Rosarum Damascenarum, P.L. 1745.

Aqua Rosæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Damask Roses ten pounds,

Proof Spirit seven fluidounces,

Water two gallons;

Let a gallon distil.

AQUA SAMBUCL.

Elder Water.

Take of Elder Flowers ten pounds, *or*
Oil of Elder two drachms,
Proof Spirit seven fluidounces,
Water two gallons ;

Let a gallon distil.

Several of the Distilled Waters above mentioned may be prepared in a very short time, when wanted for more immediate use, by carefully triturating a drachm of any distilled Oil with a drachm of Carbonate of Magnesia, and afterwards with four pints of distilled Water. Lastly, let the water be strained.

CATAPLASMATA.Cataplasms.

CATAPLASMA CONIL.

Cataplasim of Hemlock.

Take of Extract of Hemlock two ounces,
Water a pint,

Mix, and add

Linseed, bruised, as much as may be sufficient
to make it of a proper consistence.

Medicinal Uses.—This has been applied to irritable sores, and scrofulous glandular swellings.

CATAPLASMA FERMENTI.

Cataplasm of Yest.

Take of Flour a pound,
Yest of Beer half a pint ;
Mix, and apply a gentle heat until they begin to rise.

This is applied to painful and foul ulcers, and it is stated that it diminishes the foetor of the discharge and hastens the sloughing of the sores. Its efficacy is supposed to depend upon the carbonic acid gas evolved during the fermentation occasioned by the yest.

CATAPLASMA LINI.

Cataplasm of Linseed.

Take of Boiling Water a pint,
Bruised Linseed as much as may be sufficient
to make it of a proper consistence ;
Mix.

CATAPLASMA SINAPIS.

Cataplasm of Mustard.

Take of Linseed,
Mustard Seed, each powdered, half a pound,
Boiling Vinegar as much as may be sufficient
to make them of the consistence of a Cataplasm ;
Mix.

This Cataplasm is stimulant and rubefacient ; applied spread on cloth to the soles of the feet in the low stage of typhus fever, when stupor or delirium is present. It is also used in the same way in apoplexy and coma, and other cases in which there is great determination to the head.

CERATA.Cerates.

CERATUM.

Cerate.

Ceratum, P.L. 1809.*Ceratum simplex*, P.L. 1824.

Take of Olive Oil four fluidounces,

Wax four ounces ;

Add the Oil to the melted Wax, and strain.

This is used as a cooling dressing, and as a basis for more active preparations.

CERATUM CALAMINÆ.

Cerate of Calamine.

Ceratum Epuloticum, P.L. 1745.*Ceratum Lapidis Calaminiaris*, P.L. 1788.*Ceratum Calaminæ*, P.L. 1809, P.L. 1824.

Take of Calamine,

Wax, each half a pound,

Olive Oil sixteen fluidounces ;

Mix the Oil with the melted Wax ; then remove them from the fire, and when first they begin to thicken, add the Calamine, and stir constantly, until they cool.

This cerate, well known by the name of *Turner's Cerate*, is used as a dressing to excoriations and ulcers, and to burns after the inflammation has subsided.

CERATUM CANTHARIDIS.

Cerate of Cantharides.

Ceratum Cantharidis, P.L. 1788.

Ceratum Lyttæ, P.L. 1809.

Ceratum Cantharidis, P.L. 1824.

Take of Cantharides, rubbed to very fine powder, an ounce,
Spermaceti Cerate six ounces;
Add the Cantharides to the Cerate softened by heat, and mix.

This Cerate is employed to promote a discharge from a blistered surface; it generally answers the purpose, without exciting much irritation; but sometimes occasions strangury, and produces swelling of the lymphatics, and general irritation.

CERATUM CETACEI.

Cerate of Spermaceti.

Ceratum Album, P.L. 1745.

Ceratum Spermatis Ceti, P.L. 1788.

Ceratum Cetacei, P.L. 1809. P.L. 1824.

Take of Spermaceti
White Wax two ounces,
Olive Oil four fluidounces;
Add the Oil to the Spermaceti and Wax melted together, and stir them with a spatula until they cool.

This is a soft cooling dressing, and is a convenient basis for more active preparations.

CERATUM HYDRARGYRI COMPOSITUM.

Compound Cerate of Mercury.

Take of the Stronger Ointment of Mercury,
Soap Cerate, each four ounces,
Camphor an ounce ;
Rub them together until they are incorporated.

This is employed to promote the dispersion of indolent tumors.

CERATUM PLUMBI ACETATIS.

Cerate of Acetate of Lead.

Unguentum Cerussæ Acetatae, P.L. 1788.

Ceratum Plumbi Superacetatis, P.L. 1809.

Ceratum Plumbi Acetatis, P.L. 1824.

Take of Acetate of Lead, powdered, two drachms,
White Wax two ounces,
Olive Oil eight fluidounces ;

Dissolve the Wax in seven fluidounces of the Oil, then
to these gradually add the Acetate of Lead separately
rubbed with the remainder of the Oil, and stir with a
spatula until they unite.

A cooling dressing in cases of burns and excoriations.

CERATUM PLUMBI COMPOSITUM.

Compound Cerate of Lead.

Ceratum Lithargyri Acetati, P.L. 1809.

Ceratum Plumbi compositum, P.L. 1824.

Take of Solution of Diacetate of Lead three fluid-
ounces,

Wax four ounces,

Olive Oil half a pint,

Camphor half a drachm ;

Mix the melted Wax with eight fluidounces of the Oil; then remove them from the fire, and, when first they begin to thicken, add gradually the solution of Diacetate of Lead, and stir them constantly with a spatula until they cool ; lastly, mix with them the Camphor dissolved in the remainder of the Oil.

This is commonly known by the name of *Goulard's Cerate*. It is applicable to the same cases as the preceding cerate. It is stated to be particularly serviceable in chronic ophthalmia of the tarsus, and for the increased secretion of tears, which so frequently affects the eyes of persons advanced in years.

CERATUM RESINÆ.

Cerate of Resin.

Ceratum Citrinum, P.L. 1745.

Ceratum Resinæ flavæ, P.L. 1788.

Ceratum Resinæ, P.L. 1809, P.L. 1824.

Take of Resin,

Wax, each a pound,

Olive Oil sixteen fluidounces ;

Melt the Resin and Wax together with a slow fire ; then add the Oil, and press the Cerate while hot, through a linen cloth.

This is commonly called *Yellow Basilicon*. It is employed as an application to foul and indolent ulcers.

CERATUM SABINÆ.

Cerate of Savine.

Ceratum Sabinæ, P.L. 1809, P.L. 1824.

Take of Savine, bruised, a pound,

Wax half a pound,

Lard two pounds;

Boil the Savine in the Lard and Wax melted together;
then press through a linen cloth.

In those cases in which the use of *Ceratum Cantharidis* excites too much irritation, this has been recommended as a substitute.

CERATUM SAPONIS.

Cerate of Soap.

Ceratum Saponis, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Soap ten ounces,

Wax twelve ounces and a half,

Oxide of Lead, powdered, fifteen ounces,

Olive Oil a pint,

Vinegar a gallon;

Boil the Vinegar with the Oxide of Lead over a slow fire, constantly stirring them until they unite; then add the Soap, and boil again in a similar manner, until all the moisture is evaporated; lastly, mix with these the Wax previously dissolved in the Oil.

This cerate is occasionally used as a cooling dressing.

CONFECTIONES.

Confections.

CONFECTIO AMYGDALÆ.

Confection of Almonds.

Take of Sweet Almonds eight ounces,
Acacia, powdered, an ounce,
Sugar four ounces ;

The Almonds being first macerated in Water, and their external coats removed, pound all the ingredients together until thoroughly incorporated.

This Confection may be longer kept unchanged if the Almonds, Acacia and Sugar, separately powdered, are afterwards mixed. Then whenever the Confection is to be used, pound all the ingredients together until thoroughly incorporated.

Remarks.—This Confection being subject to spoil, it is now very advantageously directed that the ingredients should be kept ready mixed in a dry state, and the water added to them when the confection is wanted for use. The confection affords an expeditious mode of preparing Almond Mixture.

CONFECTIO AROMATICA.

Aromatic Confection.

Confectio Raleighana, P.L. 1720.

Confectio Cardiaca, P.L. 1745.

Confectio Aromatica, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Cinnamon,
Nutmegs, each two ounces,

Cloves an ounce,
Cardamoms half an ounce,
Saffron two ounces,
Prepared Chalk sixteen ounces,
Sugar two pounds ;

Rub the dry ingredients together to a very fine powder and keep them in a close vessel. And whenever the Confection is to be used, add water gradually, and mix until they are thoroughly incorporated.

Medicinal Uses.—Stimulant. Cordial. Dose, gr. xx. to 3j. or more. It is incompatible with acids, acidulous and metallic salts, on account of the carbonate of lime which it contains.

As this preparation when mixed with water is subject to ferment, and act upon the carbonate of lime, it is now advantageously directed to be kept dry, and water added when wanted. Another improvement is, probably, the substitution of prepared chalk for prepared shells ; the chalk being much more easily levigated, the Confection is on this account less likely to be gritty.

CONFECTIO AURANTII.

Confection of Orange.

Conserva flavedinis Corticum Aurantiorum, P.L. 1720,
P.L. 1745.

Conserva Corticis exterioris Aurantii Hispalensis,
P.L. 1788.

Confectio Aurantii, P.L. 1809.

Confectio Aurantiorum, P.L. 1824.

Take of the fresh Rind of Orange, separated by a rasp,
a pound,

Sugar three pounds ;

Bruise the Rind with a wooden pestle in a stone mortar ; then, the Sugar being added, again pound them until they are thoroughly incorporated.

CONFECTIO CASSIÆ.

Confection of Cassia.

Electuarium e Cassia, P.L. 1745.

Electuarium Cassiæ, P.L. 1788.

Confectio Cassiæ, P.L. 1809, P.L. 1824.

Take of Cassia [pulp] half a pound,

Manna two ounces,

Tamarind [pulp] an ounce,

Syrup of Rose eight fluidounces ;

Bruise the Manna, then dissolve it in the Syrup ; afterwards mix in the Cassia and Tamarind [pulp], and evaporate the moisture until a proper consistence is attained.

Purgative in doses of ʒij. to ʒj.

CONFECTIO OPII.

Confection of Opium.

Philonium Romanum, P.L. 1720.

Philonium Londinense, P.L. 1745.

Confectio Opiata, P.L. 1788.

Confectio Opii, P.L. 1809, P.L. 1824.

Take of hard Opium, powdered, six drachms,

Long Pepper an ounce,

Ginger two ounces,

Carraway three ounces,

Tragacanth, powdered, two drachms,

Syrup sixteen fluidounces ;

Rub them together to a very fine powder, and keep it

in a close vessel. And whenever the Confection is to be used, add sixteen fluidounces of Syrup made hot, and mix.

Medicinal Properties.—Narcotic. Stimulant. Dose, gr. x. to gr. xxx. One grain of opium is contained in about 36 grains of this confection.

CONFECTIO PIPERIS NIGRI.

Confection of Black Pepper.

Confectio Piperis Nigri, P.L. 1824.

Take of Black Pepper,

Elecampane [Root], each a pound,

Fennel [Seeds] three pounds,

Honey,

Sugar, each two pounds ;

Rub the dry ingredients together, to a very fine powder, and keep them in a covered vessel. And whenever the Confection is to be used, the Honey being added, pound them until thoroughly incorporated.

Medicinal Uses.—This preparation was introduced into the last Pharmacopœia ; it is probably intended as a substitute for Ward's Paste for Piles, &c. Dose, from ʒj. to ʒij. With respect to Ward's Paste, Dr. Paris observes, that " it is principally useful in those cases attended with considerable debility, in leucophlegmatic habits, and when piles arise from a deficient secretion in the rectum ;" in cases attended with inflammation it does harm.

CONFECTIO ROSÆ CANINÆ.

Confection of Dog Rose [Hips.]

Conserva Fructus Cynosbati, P.L. 1720, P.L. 1745.

Conserva Cynosbati, P.L. 1788.

Confectio Rosæ Caninæ, P.L. 1809, P.L. 1824.

Take of Dog Rose [pulp] a pound,

Sugar, powdered, twenty ounces ;

Expose the Pulp of the Rose to a gentle heat in an earthen vessel ; then add the Sugar gradually, and rub together until thoroughly incorporated.

This is principally employed as an agreeable vehicle for making up more active medicines into pills and electuaries.

CONFECTIO ROSÆ GALLICÆ.

Confection of Red Rose.

Conserva Rosarum rubrarum, P.L. 1720, P.L. 1745.

Conserva Rosæ, P.L. 1788.

Conserva Rosæ Gallicæ, P.L. 1809, P.L. 1824.

Take of Red Rose [petals] a pound,

Sugar three pounds ;

Bruise the Petals in a stone mortar ; then, the Sugar being added, pound them again until thoroughly incorporated.

This Confection is employed for the same purposes as the last.

CONFECTIO RUTÆ.

Confection of Rue.

Electarium e Baccis Lauri, P.L. 1720, P.L. 1745.

Confectio Rutæ, P.L. 1809, P.L. 1824.

Take of Rue, dried,

Carraway,

Bay Berries, each an ounce and a half,

Sagapenum half an ounce,

Black Pepper two drachms,

Honey sixteen ounces;

Rub the dry ingredients together to a very fine powder and preserve them. Then, whenever the Confection is to be used, add the Honey to them, and mix them all.

This Confection is used as an antispasmodic in enemata only.

CONFECTIO SCAMMONII.

Confection of Scammony.

Electarium Caryocostinum, P.L. 1720.

Electarium e Scammonio, P.L. 1745.

Electuarium Scammonii, P.L. 1788.

Confectio Scammoneæ, P.L. 1809, P.L. 1824.

Take of Scammony, powdered, an ounce and a half,

Cloves, bruised,

Ginger, powdered, each six drachms,

Oil of Carraway half a fluidrachm,

Syrup of Rose, as much as may be sufficient;

Rub the dry ingredients together to very fine powder, and preserve them; then, whenever the Confection is to

be used, the Syrup being gradually poured in, rub again; lastly, the Oil of Carraway being added, mix them all.

This is a stimulating cathartic, and may be given in the dose of ʒss. to ʒj. It is but seldom used.

CONFECTIO SENNÆ.

Confection of Senna.

Electarium Lenitivum, P.L. 1720, P.L. 1745.

Electuarium Sennæ, P.L. 1788.

Confectio Sennæ, P.L. 1809, P.L. 1824.

Take of Senna eight ounces,

Figs a pound,

Tamarind [pulp],

Cassia [pulp],

Prunes [pulp], each half a pound,

Coriander four ounces,

Liquorice three ounces,

Sugar two pounds and a half,

Water three pints;

Rub the Senna with the Coriander, and by a sieve separate ten ounces of the mixed powder. Boil down the Water, with the Figs and the Liquorice added, to half; then press out [the liquor] and strain it. Evaporate the strained liquor in a water-bath, until of the whole, twenty-four fluidounces remain; then, the Sugar being added, let a Syrup be made. Lastly, rub the pulps gradually with the Syrup, and having thrown in the sifted powder, mix them all.

This is much employed as a laxative, but is generally very badly prepared, containing neither senna nor cassia, and is sold for one third the price which the genuine preparation costs. Dose, ʒij. or more.

DECOCTA.Decoctions.

Decoctions differ from hot infusions only in the application of a longer-continued heat; by this the solvent power of the water is increased, and some substances which are sparingly dissolved by mere infusion in hot water, have their virtues readily extracted by boiling in it.

In some cases, however, infusions contain more of the active principle of medicines than decoctions; thus aromatics and substances which contain essential oils, are diminished in power by their volatilization during the long-continued action of the heat. Another circumstance to be noticed is this, that some of the principles, which are dissolved by hot water, are deposited as the solution cools; this is particularly the case with cinchona, and therefore this decoction should always be exhibited turbid, from the suspension of particles which had become insoluble by cooling. Decoctions ought always to be strained hot, for the reasons which have been just stated, and they should be prepared either with soft or with distilled water; undistilled water which has been long boiled should be especially avoided.

Decoctions suffer decomposition by being kept, in the same manner as infusions, and consequently they ought to be prepared only a very few hours before they are intended for use.

DECOCTUM ALÖES COMPOSITUM.

Compound Decoction of Aloes.

Decoction Aloes Compositum, P.L. 1809, P.L. 1824.

Take of Extract of Liquorice seven drachms,
Carbonate of Potash a drachm,
Aloes, powdered,
Myrrh, powdered,
Saffron, of each a drachm and a half,
Compound Tincture of Cardamom seven fluid-
ounces,
Distilled Water a pint and a half;

Boil down the Liquorice, Carbonate of Potash, Aloes, Myrrh, and Saffron with the Water, to a pint, and strain; then add the Compound Tincture of Cardamom.

Medicinal Uses.—Mildly cathartic. Dose, from $f\text{ʒ} \text{ss.}$ to $f\text{ʒ} \text{j.}$

Incompatibles.—Acids, acidulous salts, earthy and metallic salts and all substances which are decomposed by carbonate of potash, or which decompose it.

The quantities of ingredients ordered in the present Pharmacopœia differ from those of the last, but the proportions, and consequently the strength of the preparation, remain the same.

DECOCTUM AMYLI.

Decoction of Starch.

Mucilago Amyli, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Starch four drachms,

Water a pint;

Rub the Starch with the Water gradually added, then boil for a short time.

Remarks.—This Decoction should be prepared from Starch which is perfectly colourless, and not that which is generally employed for domestic purposes, since it is coloured by smalts, or powdered blue glass. It is used as a demulcent.

DECOCTUM CETRARIÆ.

Decoction of Liverwort.

Decoction Lichenis, P.L. 1809, P.L. 1824.

Take of Liverwort five drachms,

Water a pint and a half;

Boil down to a pint, and strain.

Remarks.—According to Berzelius, Liverwort contains about 44 per cent. of a peculiar starch, which has been called *Lichenin*, 3 per cent. of a bitter principle mixed with bilichenates of potash and lime, phosphate of lime, gum, sugar, extract and amylaceous fibrin.

Medicinal Uses.—This Decoction is mucilaginous and bitter ; it is employed as a remedy in debility, phthisis, and disorders requiring nutritive tonics.

Dose, from f℥j. to f℥iv.

DECOCTUM CHIMAPHILÆ.

Decoction of Winter-green or Pyrola.

Take of Winter-green or Pyrola an ounce,

Distilled Water a pint and a half ;

Boil down to a pint, and strain.

Medicinal Uses.—It has been found useful in cases of dropsy and some affections of the urinary organs ; its action is principally diuretic. Dose, from f℥j. to f℥iss.

DECOCTUM CINCHONÆ CORDIFOLIÆ.

Decoction of Heart-leaved Cinchona.

Take of Heart-leaved Cinchona, bruised, ten drachms,

Distilled Water a pint ;

Boil for ten minutes in a lightly covered vessel, and strain the liquor while hot.

Medicinal Uses.—Tonic in dyspepsia, &c. Dose from f℥j. to f℥iij. two or three times a day. Although cinchona in the form of decoction is less powerful than when exhibited in substance, yet in the former state it may be taken by persons with whom the powder would not agree.

DECOCTUM CINCHONÆ LANCIFOLIÆ.

Decoction of Lance-leaved Cinchona.

Decoctum Cinchonæ, P.L. 1788, P.L. 1809, P.L. 1824.

DECOCTUM CINCHONÆ OBLONGIFOLIÆ.

Decoction of Oblong-leaved Cinchona.

These are prepared in the same manner as the Decoction of Heart-leaved Cinchona.

Remarks. — The three varieties of cinchona contain two vegetable alkalis, namely, *quina* and *cinchonina*, in which their medicinal virtues reside; and it will appear from the annexed comparative statement of their analyses, by Pelletier and Caventou, that the *Cinchona oblongifolia*, or Red Bark, differs materially in composition from the others in containing cinchonina, they being similar to each other, except that the *Cinchona lancifolia*, or Pale Bark, contains gum, which the *Cinchona cordifolia*, or Yellow Bark, does not :

<i>Red Bark.</i>	<i>Pale and Yellow Bark.</i>
Acidulous Kinate of Cinchonina.	Kinate of Quina.
Acidulous Kinate of Quina.	Kinate of Lime.
Kinate of Lime.	Tannic Acid.
Tannic Acid (Tannin).	Red Colouring matter.
Red Colouring matter.	Yellow Colouring matter.
Yellow Colouring matter.	Starch and Gum.
Starch.	Fatty matter.
Fatty matter.	Lignin.
Lignin.	

An account of the properties and composition of quina has already been given; cinchonina is usually prepared from the pale bark which contains it, on the same plan as quina is from yellow bark. Its properties are as follows: When the alcoholic solution is suffered to evaporate slowly the cinchonina separates in slender prismatic crystals; but when the evaporation is rapid it is deposited in colourless, translucent, crystalline plates. It requires 2500 times its weight of boiling water for solution, and in cold

water it is nearly insoluble ; it has a bitter taste, which is slowly developed on account of its slight solubility ; on the addition of an acid it becomes intensely bitter. It suffers no change by exposure to the air. It is very soluble in alcohol, especially when hot, and on cooling crystals are deposited. It restores the colour of litmus which has been reddened, and combines with acids to form neutral and crystallizable salts. When strongly heated it is totally decomposed, yielding ammonia.

Cinchonia is composed of

Twenty equivalents of Carbon	$6 \times 20 = 120$	or 78.
Twelve equivalents of Hydrogen . . .	$1 \times 12 = 12$	7.8
One equivalent of Oxygen	$= 8$	5.2
One equivalent of Azote	$= 14$	9.
	<hr/>	<hr/>
Equivalent	154	100.

Symbol.—Berzelius and Turner $C^{20} H^{12} O^1 N^1$.

Brande $(20car + 12h + o + n, \text{ or } CIN.)$

Medicinal Uses.—Being a much less powerful remedy than quina, it is not separately prepared for medicinal use.

DECOCTUM CYDONIÆ.

Decoction of Quince [Seeds].

Mucilago Seminis Cydonii Mali, P.L. 1788.

Decoction Cydoniæ, P.L. 1809, P.L. 1824.

Take of Quince [Seeds] two drachms,

Distilled Water a pint ;

Boil over a slow fire for ten minutes ; afterwards strain.

Medicinal Uses.—Quince seeds contain a large quantity of inodorous and insipid mucilaginous matter, which is readily dissolved by water. The decoction is viscid and nearly colourless ; it has been recommended as an application to erysipelatous surfaces : it is also employed in aphthous affections and excoriations of the mouth, &c. It very readily suffers decomposition, and on this account should never be kept ready prepared.

Incompatibles.—Alcohol, acids, and most metallic solutions.

DECOCTUM DULCAMARÆ.

Decoction of Woody Nightshade.

Decoctum Dulcamaræ, P.L. 1809, P.L. 1824.

Take of Woody Nightshade, sliced, ten drachms,
Distilled Water a pint and a half;
Boil down to a pint, and strain.

Remarks.—The active principle of the *Solanum dulcamara* is an alkali which has been named *Solania*; it is combined in the plant with malic acid; solania is colourless, pulverulent, and pearly. It is insoluble in cold water, and requires 8000 times its weight when boiling to dissolve it. In oil it is insoluble, sparingly soluble in æther, but readily in alcohol. It has the alkaline properties of restoring reddened litmus paper and saturating acids to form salts.

It is extremely poisonous, and according to Blanchet is composed of

Carbon	62·0
Hydrogen	8·9
Azote	1·6
Oxygen	27·5
	<hr/>
	100.

The accuracy of this analysis is, however, questionable, since it would indicate the combination of a much greater number of equivalents than has hitherto been ascertained to occur.

Medicinal Uses.—Diuretic and narcotic. Dose, from fʒ iv. to fʒ j. three times a day, combined with an aromatic.

DECOCTUM GRANATI.

Decoction of Pomegranate.

Take of Pomegranate [rind] two ounces,
Distilled Water a pint and a half;
Boil down to a pint, and strain.

Remarks.—The rind of the pomegranate is astringent, containing gum, extract and tannin or tannic acid. The decoction is given in doses of fʒ ss. to fʒ j. It has been found useful in cases of tapeworm, as well as in dysentery.

DECOCTUM HORDEI.

Decoction of Barley.

Decoctum Hordei, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Barley [Pearl Barley] two ounces and a half,
Water four pints and a half;

First wash away with cold water the foreign matters from the Barley Seeds; afterwards, half a pint of the Water being poured upon them, boil the Seeds a little while. This Water being thrown away, pour on that which is left, first made hot; then boil down to two pints, and strain.

DECOCTUM HORDEI COMPOSITUM.

Compound Decoction of Barley.

Decoctum Hordei Compositum, P.L. 1788,
P.L. 1809, P.L. 1824.

Take of Decoction of Barley two pints,
Figs, sliced, two ounces and a half,
Liquorice [Root], sliced and bruised, five drachms,
Raisins two ounces and a half,
Water a pint;
Boil down to two pints, and strain.

Medicinal Uses.—This and the simple decoction are useful demulcents in fever, phthisis, gonorrhœa and strangury, given *ad libitum*.

DECOCTUM MALVÆ COMPOSITUM.

Compound Decoction of Mallow.

Decoctum Malvæ Compositum, P.L. 1809, P.L. 1824.

Take of Mallow, dried, an ounce,
Chamomile, dried, half an ounce,
Water a pint;

Boil for a quarter of an hour, and strain.

Medicinal Uses.—Employed in fomentations and enemas.

DECOCTUM PAPAVERIS.

Decoction of Poppy.

Decoctum Papaveris, P.L. 1809, P.L. 1824.

Take of Poppy [Capsules], sliced, four ounces,

Water four pints;

Boil for a quarter of an hour, and strain.

Medicinal Uses.—External as an anodyne fomentation in painful swellings, and in the excoriations produced by the acrid discharge of ulcers.

DECOCTUM QUERCUS.

Decoction of Oak [Bark].

Decoctum Quercus, P.L. 1809, P.L. 1824.

Take of Oak [Bark] ten drachms,

Distilled Water two pints;

Boil down to a pint, and strain.

Remarks.—The well-known astringent property of oak and similar barks has been ascribed to a proximate principle called *tannin*, on account of its power of converting skin into leather, and it has been lately found that they contain besides gallic acid a quantity of a peculiar acid in which the tanning power has been, at least partly, supposed to reside, called *tannic acid*. All vegetable matters which are employed in tanning give an insoluble precipitate with albumen and gelatine, and a dark-coloured one with the salts of iron, which has been called *tanno-gallate of iron*. The exact effect produced by each peculiar principle has not however been satisfactorily ascertained.

Medicinal Uses.—This decoction is principally employed in the form of gargle, injection, or lotion, as a local astringent. It is nearly inodorous and has a very astringent taste.

Incompatibles.—Decoction of cinchona, metallic salts, solution of isinglass, and alkaline solutions destroy its astringency.

DECOCTUM SARZÆ.

Decoction of Sarsaparilla.

Decoction Sarsaparillæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Sarsaparilla, sliced, five ounces,

Boiling Distilled Water four pints ;

Macerate for four hours, in a vessel lightly covered, near the fire, then take out and bruise the Sarsaparilla. When bruised, return it to the liquor, and again macerate in the same manner for two hours ; afterwards boil down to two pints, and strain.

Remarks.—The medicinal power of Sarsaparilla appears to exist in a peculiar neutral vegetable product called *Parillin*.

Medicinal Uses.—Alterative, Demulcent. Dose, from fʒiv. to fʒviii. three or four times a day.

Incompatibles.—Lime-water and acetates of lead and also some solutions of mercury.

DECOCTUM SARZÆ COMPOSITUM.

Compound Decoction of Sarsaparilla.

Decoction Sarsaparillæ Compositum, P.L. 1788,
P.L. 1809, P.L. 1824.

Take of Decoction of Sarsaparilla, boiling, four pints,

Sassafras, sliced,

Guaiacum Wood shavings,

Liquorice, bruised, each ten drachms,

Mezereon three drachms ;

Boil for a quarter of an hour, and strain.

Remarks.—The mezereon is the only very active substance here added to the sarsaparilla ; it contains a neutral vegetable matter called *Daphnin*, from the name of the root, *Daphne Mezereon*.

Medicinal Uses.—Diaphoretic and alterative. It is esteemed to be useful in secondary syphilis and in rheumatism. Dose, fʒiv. to fʒvi. three or four times a day.

DECOCTUM SCOPARII COMPOSITUM.

Compound Decoction of Broom.

Take of Broom,
Juniper Berries,
Dandelion, each half an ounce,
Distilled Water a pint and a half;
Boil down to a pint, and strain.

Medicinal Uses.—This is possessed of diuretic properties, which may be still further increased by the addition of small doses of tartrate or acetate of potash. It has been found efficient in dropsy.

DECOCTUM SENEGÆ.

Decoction of Senega.

Decoction Senegæ, P.L. 1809, P.L. 1824.

Take of Senega Root ten drachms,
Distilled Water two pints;
Boil down to a pint, and strain.

Remarks.—Senega is supposed to contain a peculiar proximate principle, called *Senegin*, which is neutral, possessing neither acid nor alkaline properties.

Medicinal Uses.—Expectorant, diuretic, and diaphoretic. It has been recommended in pneumonic affections attended with accumulation of mucus in the bronchia, and as a diaphoretic in chronic rheumatism. Dose, fʒiss. to fʒiij. two or three times a day.

DECOCTUM TORMENTILLÆ.

Decoction of Tormentil.

Take of Tormentil, bruised, two ounces,
Distilled Water a pint and a half;
Boil down to a pint, and strain.

Medicinal Uses.—It has been found useful as an astringent and tonic, in diarrhoea and alvine fluxes. Dose $f\text{ʒj.}$ to $f\text{ʒiss.}$ three or four times a day.

DECOCTUM ULMI.

Decoction of Elm [Bark].

Decoction Ulmi, P.L. 1788, P.L. 1809, P.L. 1824.

Take of fresh Elm [Bark], bruised, two ounces and a half,
Distilled Water two pints;
Boil down to a pint, and strain.

Remarks.—Elm bark contains several proximate principles, among the rest tannin or tannic acid, but not in so great quantity as oak bark.

Medicinal Uses.—Diuretic, and in herpetic eruptions. Its powers are questionable. Dose $f\text{ʒiv.}$ to $f\text{ʒvj.}$ three or four times a day.

DECOCTUM UVÆ URSI.

Decoction of Whortleberry.

Take of Whortleberry, bruised, an ounce,
Distilled Water a pint and a half;
Boil down to a pint, and strain.

Medicinal Uses.—A good bitter, first recommended by De Haen, and subsequently very much employed in purulent and other affections of the urinary organs.

DECOCTUM VERATRI.

Decoction of White Hellebore.

Decoctum Hellebori Albi, P.L. 1788.

Decoctum Veratri, P.L. 1809, P.L. 1824.

Take of White Hellebore, bruised, ten drachms,

Distilled Water two pints,

Rectified Spirit three fluidounces ;

Boil the Hellebore in the Water down to a pint, and when it has cooled, add the Spirit, then press and strain.

Remarks.—The medicinal power of white Hellebore resides in a peculiar vegetable alkali, for an account of which see VERATRIA.

Medicinal Uses.—It is employed externally as a lotion in scabies, tinea capitis, and other cutaneous eruptions.

EMPLASTRA.

Plasters.

EMPLASTRUM AMMONIACI.

Plaster of Ammoniacum.

Emplastrum Ammoniaci, P.L. 1809, P.L. 1824.

Take of Ammoniacum five ounces,

Distilled Vinegar eight fluidounces ;

Dissolve the Ammoniacum in the Vinegar ; then evaporate the liquor, with a slow fire, constantly stirring, to a proper consistence.

Medicinal Uses.—Stimulant and discutient, applied to white swellings, scrofulous tumours, &c.

EMPLASTRUM AMMONIACI CUM HYDRARGYRO.

Plaster of Ammoniacum with Mercury.

Emplastrum ex Ammoniaco cum Mercurio, P.L. 1745.

Emplastrum Ammoniaci cum Hydrargyro, P.L. 1788,
P.L. 1809, P.L. 1824.

Take of Ammoniacum a pound,
Mercury three ounces,
Olive Oil a fluidrachm,
Sulphur eight grains ;

Add the Sulphur gradually to the heated Oil, stirring constantly with a spatula, until they unite, then rub the Mercury with them, until globules are no longer visible ; lastly, gradually add the Ammoniacum melted, and mix them all.

Medicinal Uses.—Similar to the former, but more powerful, especially in venereal nodes.

EMPLASTRUM BELLADONNÆ.

Plaster of Deadly Nightshade.

Take of Plaster of Resin three ounces,
Extract of Deadly Nightshade an ounce and
a half ;

Add the Extract to the Plaster, melted by the heat of a water-bath, and mix.

Medicinal Uses.—Anodyne and antispasmodic. Applied to the sacrum it relieves pain in dysmenorrhœa.

EMPLASTRUM CANTHARIDIS.

Plaster of Cantharides.

Emplastrum Vesicatorium, P.L. 1745.

Emplastrum Cantharidis, P.L. 1788.

Emplastrum Lyttæ, P.L. 1809.

Emplastrum Cantharidis, P.L. 1824.

Take of Cantharides, rubbed to very fine powder, a pound,

Wax Plaster a pound and a half,

Lard half a pound;

Sprinkle the Cantharides in the Plaster and Lard melted together, and removed from the fire, a little before they concrete, and mix them all.

In spreading this plaster, great care should be taken that heat be not employed, or that it be merely sufficient to soften the plaster; a high temperature decomposes the animal matter, and totally destroys its efficacy.

EMPLASTRUM CERÆ.

Plaster of Wax.

Emplastrum Attrahens, P.L. 1745.

Emplastrum Ceræ, P.L. 1809, P.L. 1824.

Take of Wax,

Suet, each three pounds,

Resin a pound;

Melt them together, and strain.

This plaster is principally used as an ingredient in the preceding.

EMPLASTRUM GALBANI.

Plaster of Galbanum.

Emplastrum Commune cum Gummi, P.L. 1745.

Emplastrum Lithargyri compositum, P.L. 1788.

Emplastrum Galbani compositum, P.L. 1809, P.L. 1824.

Take of Galbanum eight ounces,
Lead Plaster three pounds,
Common Turpentine ten drachms,
Resin of the Spruce Fir, powdered, three
ounces ;

Add first the Resin of the Spruce Fir, then the Lead Plaster melted with a slow fire, to the Galbanum and Turpentine melted together, and mix them all.

Medicinal Uses.—Stimulant. Discutient. It is more powerful than the preceding, and is said to be particularly serviceable in cases of indolent glandular enlargements of a strumous character.

EMPLASTRUM HYDRARGYRI.

Plaster of Mercury.

Emplastrum Commune cum Mercurio, P.L. 1745.

Emplastrum Lithargyri cum Hydrargyro, P.L. 1788.

Emplastrum Hydrargyri, P.L. 1809, P.L. 1824.

Take of Mercury three ounces,
Lead Plaster a pound,
Olive Oil a fluidrachm,
Sulphur eight grains ;

Add the Sulphur gradually to the heated Oil, stirring constantly with a spatula until they unite ; afterwards rub the Mercury with them, until globules are no longer visible ; then gradually add the Lead Plaster melted with a slow fire, and mix them all.

Medicinal Uses.—Alterative. Discutient. It is less powerful than the *Emplastrum Ammoniaci cum Hydrargyro*.

EMPLASTRUM OPII.

Plaster of Opium.

Emplastrum Opii, P.L. 1809, P.L. 1824.

Take of Hard Opium, powdered, half an ounce,
Resin of the Spruce Fir, powdered, three
ounces,
Lead Plaster a pound,
Water eight fluidounces ;

Add the Resin of the Spruce Fir, the Opium, and
the Water to the melted Plaster, and with a slow fire
boil down, until all unite into a proper consistence.

Medicinal Uses.—Anodyne.

EMPLASTRUM PICIS.

Plaster of Pitch.

Emplastrum Cephalicum, P.L. 1745.

Emplastrum Picis Burgundicæ, P.L. 1788.

Emplastrum Picis compositum, P.L. 1809, P.L. 1824.

Take of Burgundy Pitch two pounds,
Resin of the Spruce Fir a pound,
Resin,
Wax, each four ounces,
Expressed Oil of Nutmegs an ounce,
Olive Oil,
Water, each two fluidounces ;

Add first the Resin of the Spruce Fir, then the Oil of
Nutmegs, the Olive Oil, and the Water, to the Pitch,
Resin and Wax melted together. Lastly, mix them all,
and boil down to a proper consistence.

Medicinal Uses.—Stimulant. Rubefacient in pulmonary com-
plaints ; but it frequently produces too great a degree of irrita-
tion.

EMPLASTRUM PLUMBI.

Plaster of Lead.

Diachylon Simplex, P.L. 1720.

Emplastrum Commune, P.L. 1745.

Emplastrum Lithargyri, P.L. 1788.

Emplastrum Plumbi, P.L. 1809, P.L. 1824.

Take of Oxide of Lead, rubbed to very fine powder,
six pounds,
Olive Oil a gallon,
Water two pints ;

Boil them together over a slow fire, constantly stirring, until the Oil and Oxide of Lead unite into the consistence of a plaster ; but it will be proper to add a little boiling Water, if nearly the whole of that which was used in the beginning, should be consumed before the end of the boiling.

Medicinal Uses.—It is largely employed in the bases of many other plasters, and is a common application to excoriations, and for retaining the edges of fresh-cut wounds in a state of apposition, and defending them from the air.

EMPLASTRUM RESINÆ.

Plaster of Resin.

Emplastrum Commune adhæsivum, P.L. 1745.

Emplastrum Lithargyri cum Resina, P.L. 1788.

Emplastrum Resinæ, P.L. 1809, P.L. 1824.

Take of Resin half a pound,
Plaster of Lead three pounds ;

Add the Resin, powdered, to the Plaster of Lead melted over a slow fire, and mix.

Medicinal Uses.—Stimulant. Defensive.

EMPLASTRUM SAPONIS.

Plaster of Soap.

Emplastrum e Sapone, P.L. 1745.

Emplastrum Saponis, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Soap, sliced, half a pound,

Plaster of Lead three pounds ;

Mix the Soap with the melted Plaster ; then boil down to a proper consistence.

Medicinal Use.—Discutient.

ENEMATA.

Enemas.

ENEMA ALOËS.

Enema of Aloes.

Take of Aloes two scruples,

Carbonate of Potash fifteen grains,

Decoction of Barley half a pint ;

Mix, and rub them together.

Medicinal Use.—It is employed for dislodging ascarides from the rectum, and likewise as a stimulant in constipation attendant upon amenorrhœa.

ENEMA COLOCYNTHIDIS.

Enema of Colocynth.

Take of Compound Extract of Colocynth two scruples,
Soft Soap an ounce,
Water a pint;
Mix, and rub them together.

Medicinal Use.—A very efficient enema in cases of obstinate constipation and colic.

ENEMA OPII.

Enema of Opium.

Take of Decoction of Starch four fluidounces,
Tincture of Opium thirty minims;
Mix.

Medicinal Use.—The bulk of the fluid is small for the obvious purpose of causing it to be retained for some time, so that it may act as an anodyne to irritable bowels.

ENEMA TABACI.

Enema of Tobacco.

Take of Tobacco a drachm,
Boiling Water a pint;
Macerate for an hour and strain.

Medicinal Use.—A very drastic enema recommended by some in cases of hernia, but with doubtful success.

ENEMA TEREBINTHINÆ.

Enema of Turpentine.

Take of Oil of Turpentine a fluidounce,
Yelk of Egg as much as may be sufficient ; rub
them together and add
Decoction of Barley nineteen fluidounces ;
Mix.

Medicinal Use.—Employed in cases of intestinal worms, chiefly of tænia, and likewise in some spasmodic affections, as in chorea.

EXTRACTA.

Extracts.

Extracts are those preparations which are obtained when vegetable substances are boiled in water, or have their soluble parts dissolved in proof spirit of wine, or when the expressed juices of recent plants are boiled down to a proper consistence for forming into pills ; and in some cases, the evaporation is carried so far that the extract is reducible to powder.

As the medicinal power of some vegetable substances resides, to a certain extent, in principles which are insoluble in water, but dissolve in spirit of wine, different modes of operating are adopted ; in the first case, that is, when the virtues of the medicines are completely soluble in water, such for example as those of gentian, the extract is termed a watery extract : when the vegetable contains resinous or other matter insoluble in water, it is extracted by spirit, and then termed a spirituous extract ; while the juices of recent plants, when evaporated to a proper degree were formerly called inspissated juices, but they are classed by the College with the extracts.

That part of vegetable bodies which is soluble in water, and reduced by evaporation to the state of extract, has, on this ac-

count, received the name of extractive matter, extract or extractive; it is evident, however, that extracts consist of all the various substances soluble in water, and they must therefore contain very different ingredients; for some are neutral, such as *gentianin*, the peculiar principle of gentian; but others contain neutral salts; for example the extracts of cinchona contain *kinate of cinchonia* and *kinate of quina*, extract of opium contains *meconate of morphia*, &c., &c.; their medicinal powers existing generally, if not entirely, in the alkali.

In preparing all extracts, unless otherwise ordered, evaporate the water, by a water bath, in a pan, as quickly as possible, towards the end stirring constantly with a spatula, until a proper consistence is acquired for forming pills.

Sprinkle upon all softer extracts, a little rectified spirit, that they may not become mouldy.

EXTRACTUM ACONITI.

Extract of Aconite.

Extractum Aconiti, P.L. 1809, P.L. 1824.

Take of Aconite Leaves, fresh, a pound;

Bruise them, sprinkled with a little water, in a stone mortar; then press out the juice, and evaporate it unstrained, to a proper consistence.

Medicinal Uses.—Narcotic; in some cases diuretic. The dose should not at first exceed half a grain; but it may be gradually increased to gr. v. The medicinal power of aconite resides in a peculiar alkali: see ACONITINA. This extract is of a brown colour; it has a disagreeable smell, and an acrid taste, and is not much employed.

The six following Extracts are prepared in the same manner as Extract of Aconite.

EXTRACTUM BELLADONNÆ.

Extract of Deadly Nightshade.

Extractum Belladonnæ, P.L. 1809, P.L. 1824.

Medicinal Uses.—Most useful as an external application to the eyebrows in ophthalmic surgery, to produce dilatation of the pupil.

EXTRACTUM COLCHICI CORMI.

Extract of the Cormus of Meadow Saffron.

Medicinal Uses.—Given in the dose of one grain every four hours, it has been found most efficient in several forms of acute rheumatism, particularly in its earliest stage, and is very generally used by Dr. Hue at St. Bartholomew's Hospital.

EXTRACTUM CONII.

Extract of Hemlock.

Succus Cicutæ Spissatus, P.L. 1788.

Extractum Conii, P.L. 1809, P.L. 1824.

Medicinal Uses.—Anodyne. In doses of five grains every eight hours, and gradually increased to the same quantity every four hours, or until headache or sense of constriction across the forehead supervene, it has been successfully employed in cases of acute rheumatism in a more advanced period than the preceding. It is likewise of service in hooping-cough.

EXTRACTUM DIGITALIS.

Extract of Foxglove.

Medicinal Uses.—The exhibition of Digitalis in this form requires great caution; for the virtues of the extract must vary with its mode of preparation, the quality of the leaf, as affected by the season, and the degree of concentration of the extract itself.

EXTRACTUM HYOSCYAMI.

Extract of Henbane.

Extractum Hyoscyami, P.L. 1809, P.L. 1824.

Medicinal Uses.—This extract is an excellent anodyne and antispasmodic, and possesses the advantage of not materially confining the bowels.

Dose, gr. v. to gr. x. twice or thrice a day.

EXTRACTUM LACTUCÆ.

Extract of Lettuce.

Extractum Lactucæ, P.L. 1824.

Medicinal Uses.—This preparation is much esteemed by some as a mild opiate and narcotic.

Dose, gr. v. to gr. x. twice or thrice a day.

Remarks.—The above extracts contain the respective peculiar principle of the plants from which they are obtained, and I shall briefly state the nature of each.

ACONITUM. See ACONITINA.

BELLADONNA.—The alkali contained in this plant is called *Atropia*; the root furnishes about 3–1000dth of its weight. It is colourless, inodorous, crystallizes in silky transparent prisms, soluble in about 1–500dth of its weight of cold water, and more soluble in hot water. The solution has a disagreeable bitter taste, and the alkaline property of restoring the blue colour of litmus reddened by an acid. The aqueous solution very readily and permanently dilates the pupil of the eye. It gives a white precipitate with infusion of galls, a lemon yellow one with chloride of gold, and an Isabella yellow with chloride of platina. It is soluble in absolute alcohol and in æther.

Atropia combines with acids to form definite salts; the sulphate and acetate are more disposed to crystallize than the nitrate or hydrochlorate.

Atropia is totally decomposed at a high temperature, yielding ammonia. According to Liebig it is composed of

Sixty-eight equivalents of Carbon . . .	$6 \times 68 = 408$ or	75.41
Twenty-three equivalents of Hydrogen $1 \times 23 =$	23 „	4.27
Twelve equivalents of Oxygen	$8 \times 12 = 96$ „	17.74
One equivalent of Azote	$= 14$ „	2.58

Equivalent	541	100.
--------------------	-----	------

Symbol,—Berzelius and Turner . . $C^{68}H^{23}O^{12}N^1$.

Brande $(68car + 23h + 12o + n)$.

COLCHICUM contains the alkali *Colchicia*. See ACETUM COLCHICI.

CONIUM contains the vegetable alkali *Conia*, which differs from all hitherto discovered in being both fluid and volatile. It has the appearance of a colourless volatile oil, is lighter than water, and of a very powerful diffusible, repulsive, odour, somewhat like that of the hemlock itself, and intensely acrid to the taste. It has a strong alkaline action on reddened litmus and on turmeric. It readily combines with and neutralizes acids, but its salts have not yet been obtained in a crystalline form. It is sparingly soluble in water, to which it imparts its odour and its taste. It forms a hydrate with about a fourth of its weight of water; by exposure to the air it quickly becomes of a dark colour, and spontaneously decomposes with the evolution of ammonia. Its boiling point is 370° ; but it distils with water when heated to 212° ; it is partially decomposed during the operation.

According to Dr. Christison few poisons equal Conia in subtilty or swiftness. A single drop put into the eye of a rabbit killed it in nine minutes; three drops, used in the same way, killed a strong cat in a minute and a half. Two grains of conia neutralized with hydrochloric acid and injected into the femoral vein of a young dog, killed it in about three seconds. It is composed, like the other vegetable alkalis, of carbon, hydrogen, oxygen and azote.

DIGITALIS.—The active principle of this plant has been called *Digitalia*: its properties have been but imperfectly examined. When the solution obtained in acetic æther is evaporated, the digitalia remains in the form of an orange-coloured, bitter mass, which is hard when cold, but becomes soft when it is heated. It attracts moisture from the air, is very soluble in water and in alcohol, though but little in common æther. The aqueous solution

is precipitated by diacetate of lead and infusion of galls. It has been called an alkali, but its properties have been imperfectly made out. It is said to crystallize under favourable circumstances: its composition is unknown, and it requires further examination.

HYOSCYAMUS.—*Hyoscyama*, the alkaline and active principle of this plant, crystallizes in colourless stellated prisms, of a silky lustre. They are inodorous and slightly soluble in water, but very soluble in alcohol and æther; their taste is disagreeable and bitter. It gives an abundant white precipitate with tincture of galls, a yellowish white precipitate with chloride of gold, but none with that of platina. This alkali is very poisonous, and when moist it has alkaline properties. If heated it is decomposed with the formation of ammonia; it neutralizes acids, and forms crystalline salts with some of them, and they are as poisonous as the hyoscyama itself. It has not been analyzed, but contains azote like the other vegetable alkalis.

LACTUCA.—The virtues of the Lettuce are stated to reside in a peculiar principle, which Dr. Duncan has called *lactucarium*. Its properties have not been minutely examined, but when in a pure state it is said to possess narcotic power, and even to contain morphia. In the extract, however, this is mixed with a large quantity of comparatively inert matter.

EXTRACTUM ALOËS PURIFICATUM.

Purified Extract of Aloes.

Extractum Aloës, P.L. 1809.

Extractum Aloës Purificatum, P.L. 1824.

Take of Aloes, powdered, fifteen ounces,

Boiling Water a gallon;

Macerate for three days with a gentle heat; afterwards strain, and set by that the dregs may subside. Pour off the clear liquor, and evaporate it to a proper consistence.

Remarks.—Aloes contains a peculiar bitter extractive matter; by treatment with water this medicine is deprived of its resinous matter, and is then said to be less irritating and more purgative in equal doses.

Medicinal Uses.—Purgative. Stomachic. Dose, gr. v. to gr. xv.

EXTRACTUM CINCHONÆ CORDIFOLIÆ.

Extract of Heart-leaved Cinchona.

Take of Heart-leaved Cinchona, bruised, fifteen ounces,
Distilled Water, four gallons ;

Boil down in a gallon of Water to six pints, and strain the liquor while hot. Boil down the bark in an equal measure of Water four times, and strain. Lastly, all the liquors being mixed, evaporate to a proper consistence.

EXTRACTUM CINCHONÆ LANCIFOLIÆ.

Extract of Lance-leaved Cinchona.

Extractum Corticis Peruviani, P.L. 1745.

Extractum Cinchonæ, P.L. 1788, P.L. 1809, P.L. 1824.

EXTRACTUM CINCHONÆ OBLONGIFOLIÆ.

Extract of Oblong-leaved Cinchona.

This and the preceding Extract are prepared in the same manner as the Extract of Heart-leaved Cinchona.

Remarks.—The nature of the substances in which the medicinal power of the different varieties of cinchona reside has been already mentioned. Extract of cinchona is of a dark brown colour, nearly inodorous and of a bitter taste. The active principles of cinchona are more soluble in spirit than in water ; during ebullition, however, a considerable portion of kinate of quina and kinate of cinchonina is dissolved by the water, with a portion of inert soluble matter, and these together constitute extract of bark, which since the discovery of quina has been much less employed than before.

Medicinal Use.—Tonic. Stomachic. Dose, gr. x. to gr. xxx.

EXTRACTUM COLCHICI ACETICUM.

Acetic Extract of Meadow Saffron.

Take of the Fresh Meadow Saffron Cormus a pound,
Acetic Acid three fluidounces ;

Bruise the Cormus, gradually sprinkled with the Acetic Acid, then press out the juice, and evaporate it in an earthen vessel which is not glazed with Lead, to a proper consistence.

Remarks.—The use of the acetic acid is to render the salt of colchicia, which the meadow saffron contains, more soluble.

Medicinal Uses.—It has been much employed in cases of acute rheumatism and gout, in dose of gr. j. to gr. ij. twice or thrice a day.

EXTRACTUM COLOCYNTHIDIS.

Extract of Colocynth.

Extractum Colocynthidis, P.L. 1809, P.L. 1824.

Take of Colocynth, cut in pieces, a pound,
Distilled Water two gallons ;

Mix and boil with a slow fire for six hours, frequently adding distilled Water, that it may always fill the same measure. Strain the liquor while yet hot ; lastly, evaporate it to a proper consistence.

Remarks.—This is a dark-coloured and extremely bitter extract. Colocynth contains a purgative principle to which the name of *colocynthin* has been given. It is a resinlike matter, which is obtained by digesting the colocynth in spirit, and evaporation. It is extremely bitter and purgative ; more soluble in alcohol than in water, and readily dissolved by acids and by alkalis.

Medicinal Uses.—Extract of Colocynth is purgative. Dose, gr. x. to gr. xxx.

EXTRACTUM COLOCYNTHIDIS COMPOSITUM.

Compound Extract of Colocynth.

Extractum Catharticum, P.L. 1745.

Extractum Colocynthidis compositum, P.L. 1788,
P.L. 1809, P.L. 1824.

Take of Colocynth, cut in pieces, six ounces,
Purified Extract of Aloes, twelve ounces,
Scammony, powdered, four ounces,
Cardamom, powdered, an ounce,
Soap three ounces,
Proof Spirit a gallon ;

Macerate the Colocynth in the Spirit, with a gentle heat, for four days. Strain the Spirit, and add to it the Aloes, Scammony, and Soap ; afterwards evaporate to a proper consistence, the Cardamom being mixed towards the end.

Medicinal Uses.—Cathartic. Dose, gr. v. to gr. xxx. It is esteemed to be particularly efficacious when combined with chloride of mercury in relieving habitual costiveness and obstinate visceral obstructions.

EXTRACTUM ELATERII.

Extract of Elaterium.

Elaterium, P.L. 1745, P.L. 1788.

Extractum Elaterii, P.L. 1809, P.L. 1824.

Slice ripe wild Cucumbers, and strain the juice, very gently expressed, through a very fine hair sieve ; then set it aside for some hours, until the thicker part has subsided. The thinner supernatant part being rejected, dry the thicker part with a gentle heat.

Medicinal Uses.—Hydragogue. Cathartic. Dose, from half a grain to two grains. This extract has a greenish colour; its taste is bitter and rather acrid; and when tolerably pure, it is light, pulverulent and inflammable. Its properties have been particularly examined by Dr. Paris, and according to his experiments, they reside in a peculiar substance which he has called *elatin*, and of which the extract contains only about 10 per cent.—*Pharmacologia*, vol. ii. p. 241, 5th edit.

Elaterium has also been examined by Mr. Hennell. He found it to contain earthy matter, starch, and lignin, 21 per cent. of purgative resin [*elatin*?], and 40 per cent. of crystallizable matter, which was bitter, neither acid nor alkaline, slightly soluble in water and æther, but readily dissolved by boiling alcohol. It consisted of carbon 36·9, hydrogen 23·9, and oxygen 39·2 in 100 parts.

EXTRACTUM GENTIANÆ.

Extract of Gentian.

Extractum Gentianæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Gentian, sliced, two pounds and a half,

Boiling distilled Water two gallons;

Macerate for twenty-four hours; then boil down to a gallon, and strain the liquor while yet hot; lastly, evaporate to a proper consistence.

Remarks.—Gentian contains a peculiar neutral vegetable matter which has been called *gentianin*. It is yellow, inodorous, possesses in a great degree the peculiar taste of gentian. It is not very soluble in cold water, but more so in boiling water. Æther and alcohol dissolve it very readily, and it separates from them by spontaneous evaporation in yellow-coloured acicular crystals. It is neither acid nor alkaline. It dissolves in dilute acids, which very much diminish its colour.

Medicinal Uses.—Tonic. Stomachic. Dose, gr. x. to gr. xxx. twice or three times a day. This extract is of a dark brown colour, nearly inodorous and bitter. It is frequently exhibited in combination with chalybeates.

The seven next following are prepared in the same manner as Extract of Gentian :

EXTRACTUM GLYCYRRHIZÆ.

Extract of Liquorice.

Extractum Glycyrrhizæ, P.L. 1745, P.L. 1788,
P.L. 1809, P.L. 1824.

Remarks.—This is a well-known extract, of a dark colour and sweet taste. It is usually imported from Spain, and when it has had a fresh form given to it, it is employed under the name of *refined liquorice* as a demulcent in tickling coughs.

Liquorice contains a peculiar sugar, which is called *glycyrrhizin*; it is a yellow transparent substance, which is extremely sweet, readily dissolves in water and alcohol, and combines also with acids and alkalis, and occasions precipitates in most metallic salts.

EXTRACTUM HÆMATOXYLI.

Extract of Logwood.

Extractum Ligni Campechensis, P.L. 1745, P.L. 1788.
Extractum Hæmatoxyli, P.L. 1809, P.L. 1824.

Remarks.—This extract is of a deep red colour, and has a sweetish astringent taste. It becomes very hard by keeping, so that pills made of it pass through the body unchanged.

Medicinal Use.—Astringent in protracted diarrhœa and dysentery. Dose, gr. x. to gr. xxx. in some aromatic distilled water.

Logwood contains a peculiar colouring principle called *hematin*; it separates from the aqueous solution in small reddish crystals, which have a bitter astringent taste; the aqueous solution is of a fine red when boiling hot, and becomes yellow on cooling. It unites with metallic oxides; alkalis render it first purple, then violet, and eventually brown, apparently by decomposition.

EXTRACTUM LUPULI.

Extract of Hops.

Extractum Humuli, P.L. 1809, P.L. 1824.

Remarks.—This is a dark-coloured, bitter extract, totally devoid of the aromatic principle of the hop. The virtues of the hop are supposed to reside in a peculiar bitter principle, to which the name of *lupulin* has been given.

Medicinal Uses.—Sedative. Dose, gr. v. to gr. xx.

EXTRACTUM PAREIRÆ.

Extract of Pareira.

Medicinal Uses.—See INFUSUM PAREIRÆ.

EXTRACTUM SARZÆ.

Extract of Sarsaparilla.

Extractum Sarsaparillæ, P.L. 1809, P.L. 1824.

Remark.—The peculiar principle of Sarsaparilla has been already noticed.

Medicinal Use.—Alterative. Dose gr. xx. to ʒj. given in pills, or dissolved in the decoction. Even among those who admit the efficacy of other preparations of sarsaparilla, there is great difference of opinion respecting the activity of this.

EXTRACTUM TARAXACI.

Extract of Dandelion.

Extractum Taraxaci, P.L. 1809, P.L. 1824.

Medicinal Uses.—Aperient. Deobstruent. Dose, gr. x. to ʒj. in obstructions of the liver, and in visceral disease.

EXTRACTUM UVÆ URSI.

Extract of Whortleberry.

Medicinal Uses.—See DECOCTUM UVÆ URSI.

EXTRACTUM JALAPÆ.

Extract of Jalap.

Extractum Jalapii, P.L. 1745, P.L. 1788.

Extractum Jalapæ, P.L. 1809, P.L. 1824.

Take of Jalap Root, powdered, two pounds and a half,
Rectified Spirit a gallon,
Distilled Water two gallons;

Macerate the Jalap Root in the Spirit for four days, and pour off the tincture. Boil down the residue in the Water to half a gallon; afterwards strain the tincture and the decoction separately, and let the latter be evaporated, and the former distil, until each thickens. Lastly, mix the extract with the Resin, and evaporate to a proper consistence.

This Extract should be kept *soft*, which may be fit to form pills, and *hard*, which may be rubbed to powder.

Medicinal Use.—Purgative. Dose, gr. x. to gr. xx. This extract is nearly inodorous, has a brown colour and a bitter taste. Jalap has been supposed to contain a peculiar principle called *jalapin*, but some doubt remains on this subject.

EXTRACTUM OPII PURIFICATUM.

Purified Extract of Opium.

Opium Colatum, P.L. 1745.

Opium Purificatum, P.L. 1788.

Extractum Opii, P.L. 1809, P.L. 1824.

Take of Opium, sliced, twenty ounces,

Distilled Water a gallon ;

Add a little Water to the Opium, and macerate for twelve hours, that it may soften ; then, the remaining Water being poured in gradually, rub them until they are very well mixed, and set by that the dregs may subside ; afterwards strain the liquor, and evaporate to a proper consistence.

Properties.—Although the cold infusion of opium possesses the peculiar smell of the drug, yet it is dissipated during evaporation, so that the extract is nearly inodorous. It is of a brown colour, and has a bitter taste. Dose, gr. j. to gr. v. for an adult. The form of extract is to be preferred to that of tincture, when it is intended to continue the operation of the medicine, and not to obtain its full effects at once ; but in cases of accident, or in which the effects of opium are to be called into immediate action, the tincture should be employed.

EXTRACTUM PAPAVERIS.

Extract of Poppy.

Extractum Papaveris albi, P.L. 1788.

Extractum Papaveris, P.L. 1809, P.L. 1824.

Take of Poppy [Capsules,] bruised, the seeds being taken out, fifteen ounces,

Boiling distilled Water, a gallon ;

Macerate for twenty-four hours ; then boil down to four pints, and strain the liquor while hot ; lastly, evaporate to a proper consistence.

Medicinal Uses.—Anodyne. Narcotic. Dose, from gr. ij. to gr. xx. given in the form of pills. This extract is said to be less apt than opium to occasion nausea, headache, and delirium, and therefore to be preferred for procuring sleep in diseases in which the head is much affected.

EXTRACTUM RHEI.

Extract of Rhubarb.

Extractum Rhei, P.L. 1809, P.L. 1824.

Take of Rhubarb, powdered, fifteen ounces,
Proof Spirit a pint,
Distilled Water seven pints ;

Macerate for four days with a gentle heat, afterwards strain, and set by, that the dregs may subside. Pour off the liquor, and evaporate when strained, to a proper consistence.

Medicinal Use.—Purgative. Dose, from gr. x. to gr. xxx. in the form of pills, or dissolved in an aromatic water. The powers of rhubarb residing in a peculiar principle called *rhein*, are said to be much diminished during the process of extraction.

EXTRACTUM STRAMONII.

Extract of Thorn Apple.

Extractum Stramonii, P.L. 1824.

Take of Thorn Apple Seeds fifteen ounces,
Boiling distilled Water a gallon ;

Macerate for four hours in a vessel lightly covered, near the fire ; afterwards take out the Seeds, and bruise them in a stone mortar : return them when bruised to the liquor. Then boil down to four pints, and strain the liquor while hot. Lastly, evaporate to a proper consistence.

Medicinal Use.—Narcotic. Dose, gr. $\frac{3}{4}$ to gr. ij. daily, in maniacal and asthmatic affections.

INFUSA.Infusions.

Infusions are mere solutions of vegetable matter in water, which is sometimes used cold, but in the London Pharmacopœia it is in every instance directed to be boiling: in this state it is poured upon the substance, the active principles of which are intended to be dissolved. The aromatic, bitter, astringent, and mucilaginous properties of vegetable products are, to a considerable extent, soluble in water; while resinous bodies, on the contrary, are totally unacted upon by it.

The substances infused should be only coarsely powdered, or cut into thin slices; for if they are employed in the state of fine powder, not only is the proper action prevented by the proximity of their particles, but the infusion is with difficulty rendered clear.

Hard water should, as much as possible, be avoided, for it not only acts less powerfully as a solvent, but the precipitation which takes place by boiling renders it extremely turbid, and increases the difficulty of procuring a clear infusion. The infusions prepared with cold water are weaker than those in which hot water is employed, unless the digestion be continued for a much longer time.

Dried vegetables are stated to yield their virtues by infusion more readily than when they are in a recent state.

If infusions be long kept, and especially in hot weather, they become turbid, deposit the matter which they had dissolved, and undergo decomposition; they ought, therefore, never to be kept for use longer than a few hours, but prepared for the occasion upon which they are prescribed.

INFUSUM ANTHEMIDIS.

Infusion of Chamomile.

Infusum Anthemidis, P.L. 1809, P.L. 1824.

Take of Chamomile five drachms,

Boiling distilled Water a pint;

Macerate for ten minutes in a vessel lightly covered, and strain.

Medicinal Use.—Stomachic, in dyspepsia; and the infusion prepared with cold water, is said to be more grateful than that made with hot. Dose, $\text{f}\overline{\text{z}}\text{i}$. to $\text{f}\overline{\text{z}}\text{ij}$.

It is employed warm for promoting the operation of emetics.

Incompatibles.—Solutions of the salts of iron, mercury, silver and lead.

INFUSUM ARMORACIÆ COMPOSITUM.

Compound Infusion of Horseradish.

Infusum Armoraciæ compositum, P.L. 1809, P.L. 1824.

Take of Horseradish, sliced,

Mustard, bruised, of each an ounce,

Compound Spirit of Horseradish a fluid-ounce,

Boiling distilled Water a pint;

Macerate the Root and the Seeds in the Water for two hours in a vessel lightly covered, and strain; then add the compound Spirit of Horseradish.

Medicinal Use.—Stimulant in paralysis. Dose, fʒ iss.

Incompatibles.—Solutions of the salts of silver and mercury, and of the alkaline carbonates.

INFUSUM AURANTII COMPOSITUM.

Compound Infusion of Orange Peel.

Infusum Aurantii Compositum, P.L. 1809, P.L. 1824.

Take of Orange Peel, dried, half an ounce,

Lemon Peel, fresh, two drachms,

Cloves, bruised, one drachm,

Boiling distilled Water a pint;

Macerate for a quarter of an hour in a vessel lightly covered, and strain.

Medicinal Use.—Stomachic. Dose, fʒi. to fʒij.

INFUSUM CALUMBÆ.

Infusion of Calumba.

Infusum Calumbæ, P.L. 1809, P.L. 1824.

Take of Calumba, sliced, five drachms,
Boiling distilled Water a pint;
Macerate for two hours in a vessel lightly covered,
and strain.

Medicinal Uses.—Tonic and stomachic. Dose, fʒjss. to fʒij. It very soon spoils; it contains no astringent matter.

Incompatibles.—Solutions of the acetates of lead, bichloride of mercury and lime-water.

INFUSUM CARYOPHYLLI.

Infusion of Cloves.

Infusum Caryophyllorum, P.L. 1809, P.L. 1824.

Take of Cloves, bruised, three drachms,
Boiling distilled Water a pint;
Macerate for two hours in a vessel lightly covered, and
strain.

Medicinal Uses.—Stimulant and stomachic. Dose, fʒi. to fʒij.

It is generally exhibited in combination with other medicines.

Incompatibles.—Lime-water, solutions of the salts of iron, zinc, lead, silver, and antimony.

INFUSUM CASCARILLÆ.

Infusion of Cascarilla.

Infusum Cascarillæ, P.L. 1809, P.L. 1824.

Take of Cascarilla Bark, bruised, an ounce and a half,
Boiling distilled Water a pint;

Macerate for two hours in a vessel lightly covered, and strain.

Medicinal Uses.—Tonic and stomachic. Dose, fʒjss. to fʒij.

Incompatibles similar to those enumerated under the last infusion.

INFUSUM CATECHU COMPOSITUM.

Compound Infusion of Catechu.

Infusum Catechu, P.L. 1809.

Infusum Catechu compositum, P.L. 1824.

Take of Extract of Catechu six drachms,

Cinnamon, bruised, a drachm,

Boiling distilled Water a pint;

Macerate for an hour in a vessel lightly covered, and strain.

Medicinal Use.—Astringent in diarrhoea. Dose, fʒi. to fʒiij. every four hours.

INFUSUM CINCIONÆ.

Infusion of Cinchona.

Infusum Cinchonæ, P.L. 1809, P.L. 1824.

Take of Lance-leaved Cinchona bruised, an ounce,

Boiling distilled Water a pint;

Macerate for six hours in a vessel lightly covered, and strain.

Medicinal Uses.—Tonic in dyspepsia, &c. Dose, fʒi. to fʒiij. three or four times a day.

INFUSUM CUSPARIÆ.

Infusion of Cusparia.

Infusum Cuspariæ, P.L. 1809, P.L. 1824.

Take of Cusparia, bruised, five drachms,
Boiling distilled Water a pint;
Macerate for two hours in a vessel lightly covered, and strain.

Medicinal Uses.—Tonic and stimulant in dyspepsia. Dose, fʒjss. to fʒij.

Incompatibles.—The solutions of the salts of most metals.

INFUSUM DIGITALIS.

Infusion of Foxglove.

Infusum Digitalis, P.L. 1809, P.L. 1824.

Take of Foxglove Leaves, dried, a drachm,
Spirit of Cinnamon a fluidounce,
Boiling distilled Water a pint;
Macerate the Foxglove Leaves in the Water for four hours in a vessel lightly covered, and strain; then add the Spirit.

Medicinal Use.—Diuretic. Dose, fʒss. to fʒj. twice a day.

Incompatibles.—It is decomposed by solutions of the salts of iron, and probably by those of most other metals.

INFUSUM DIOSMÆ.

Infusion of Buchu.

Take of Buchu an ounce,
Boiling distilled Water a pint;

Macerate for four hours in a vessel lightly covered, and strain.

Medicinal Uses.—Diuretic and tonic.

INFUSUM GENTIANÆ COMPOSITUM.

Compound Infusion of Gentian.

Infusum Amarum simplex, P.L. 1720, P.L. 1745.

Infusum Gentianæ compositum, P.L. 1809, P.L. 1824.

Take of Gentian, sliced,

Orange Peel, dried, of each two drachms,

Lemon Peel, fresh, four drachms,

Boiling distilled Water a pint;

Macerate for an hour in a vessel lightly covered, and strain.

Medicinal Uses.—Stomachic and tonic. Dose, fʒjss. to fʒii.

Incompatibles.—Solution of diacetate of lead, sulphate of iron, and analogous salts.

INFUSUM KRAMERIÆ.

Infusion of Rhatany.

Take of Rhatany an ounce,

Boiling distilled Water a pint;

Macerate for four hours in a vessel lightly covered, and strain.

Medicinal Uses.—Tonic and astringent. Dose, fʒiss to fʒij.

INFUSUM LINI COMPOSITUM.

Compound Infusion of Linseed.

Infusum Lini, P.L. 1809.

Infusum Lini compositum, P.L. 1824.

Take of Linseed, bruised, six drachms,
Liquorice, sliced, two drachms,
Boiling distilled Water a pint;
Macerate for four hours near the fire, in a vessel lightly covered, and strain.

Medicinal Uses.—Demulcent in dysuria and catarrh.

Incompatibles.—Preparations of lead and iron, and probably most metallic salts.

INFUSUM LUPULI.

Infusion of Hops.

Take of Hops six drachms,
Boiling distilled Water a pint;
Macerate for four hours in a vessel lightly covered,
and strain.

Medicinal Uses.—Tonic, stomachic and slightly narcotic.
Dose, fʒj. to fʒiss.

INFUSUM PAREIRÆ.

Infusion of Pareira.

Take of Pareira six drachms,
Boiling distilled Water a pint;
Macerate for two hours in a vessel lightly covered,
and strain.

Medicinal Uses.—Employed in cases of irritation of the bladder and *catarrhus vesicæ*. Dose, fʒj. to fʒiss. twice or thrice a day. The activity of the infusion may be increased by the addition of the extract.

INFUSUM QUASSIÆ.

Infusion of Quassia.

Infusum Quassiæ, P.L. 1809, P.L. 1824.

Take of Quassia, sliced, two scruples,
Boiling distilled Water a pint.

Macerate for two hours in a vessel lightly covered, and strain.

Medicinal Uses.—Stomachic and tonic. Dose, fʒjss. to fʒij.

Incompatibles.—There are few substances which produce any effect upon this solution; even the preparations of iron are unchanged by it.

INFUSUM RHEI.

Infusion of Rhubarb.

Infusum Rhei, P.L. 1809, P.L. 1824.

Take of Rhubarb, sliced, three drachms,
Boiling distilled Water a pint;

Macerate for two hours in a vessel lightly covered, and strain.

Medicinal Uses.—Stomachic and tonic. Dose, fʒj. to fʒiij.

Incompatibles.—The stronger acids, metallic solutions, some astringent infusions. The alkalis darken the colour of this infusion, but do not decompose it.

INFUSUM ROSÆ COMPOSITUM.

Compound Infusion of Roses.

Take of Red Rose [Petals,] dried, three drachms,
Diluted Sulphuric Acid a fluidrachm and a half,
Sugar six drachms,
Boiling distilled Water a pint;

Pour the Water upon the Rose Petals in a glass vessel; then mix in the Acid. Macerate for six hours, and strain the liquor; lastly, add the Sugar to it.

Medicinal Uses.—Astringent and refrigerant. Dose, fʒj. to fʒjss. or more.

Incompatibles.—Alkalis and earths, and all substances which combine with sulphuric acid, or are acted upon by small quantities of it; acetate of lead of course throws down a copious precipitate. Sulphate of iron gives it a brown colour, but no precipitate is formed for some hours. It is much employed as a vehicle for the exhibition of cathartic salts.

INFUSUM SCOPARII.

Infusion of Broom.

Take of Broom an ounce,
Boiling distilled Water a pint;
Macerate for four hours in a vessel lightly covered, and strain.

Medicinal Uses.—Aperient. Diuretic. It has been used in dropsy, and it is said with success.

INFUSUM SENNÆ COMPOSITUM.

Compound Infusion of Senna.

Infusum Sennæ, P.L. 1720.

Infusum Sennæ commune, P.L. 1745.

Infusum Sennæ simplex, P.L. 1788.

Infusum Sennæ, P.L. 1809.

Infusum Sennæ compositum, P.L. 1824.

Take of Senna fifteen drachms,
Ginger, sliced, four scruples,
Boiling distilled Water a pint;
Macerate for an hour in a vessel lightly covered, and strain.

Medicinal Use.—Purgative. Dose, $f\text{℥} \text{ij.}$ to $f\text{℥} \text{iv.}$

This infusion spoils quickly; when exposed to the air a yellow precipitate is formed in it, and its purgative qualities are lost.

Incompatibles.—Strong acids, lime-water, and most metallic salts.

INFUSUM SERPENTARIÆ.

Infusion of Serpentry.

Take of Serpentry half an ounce,

Boiling distilled Water a pint;

Macerate for four hours in a vessel lightly covered, and strain.

Medicinal Uses.—Diaphoretic. Tonic. Dose, $f\text{℥} \text{j.}$ to $f\text{℥} \text{ij.}$ two or three times a day.

INFUSUM SIMARUBÆ.

Infusion of Simaruba.

Infusum Simaroubæ, P.L. 1809, P.L. 1824.

Take of Simaruba, bruised, three drachms,

Boiling distilled Water a pint;

Macerate for two hours in a vessel lightly covered, and strain.

Medicinal Uses.—Tonic, in the latter stages of dysentery. Dose, $f\text{℥} \text{ij.}$

Incompatibles.—Lime-water, alkaline carbonates; many metallic salts, especially those of lead, silver and mercury.

INFUSUM VALERIANÆ.

Infusion of Valerian.

Take of Valerian half an ounce,
Boiling distilled Water a pint ;
Macerate for half an hour in a vessel lightly covered,
and strain.

Medicinal Use.—Antispasmodic. Dose, fʒiss. to fʒij. two
or three times a day.

LINIMENTA.

Liniments.

LINIMENTUM ÆRUGINIS.

Liniment of Verdigris.

Unguentum Ægyptiacum, P.L. 1720.

Mel Ægyptiacum, P.L. 1745.

Oxymel Æruginis, P.L. 1788.

Linimentum Æruginis, P.L. 1809, P.L. 1824.

Take of Verdigris, powdered, an ounce,
Vinegar seven fluidounces,
Honey fourteen ounces ;
Dissolve the Verdigris in the Vinegar, and strain
through a linen cloth ; afterwards, the Honey being added,
boil down to a proper consistence.

Detergent and escharotic.

LINIMENTUM AMMONIÆ.

Liniment of Ammonia.

Linimentum Ammoniæ Fortius, P.L. 1788,
P.L. 1809, P.L. 1824.

Take of Solution of Ammonia a fluidounce,
Olive Oil two fluidounces;
Shake them together until they are mixed.

LINIMENTUM AMMONIÆ SESQUICARBONATIS.

Liniment of Sesquicarbonate of Ammonia.

Linimentum Volatile, P.L. 1745.
Linimentum Ammoniæ, P.L. 1788.
Linimentum Ammoniæ Carbonatis, P.L. 1809.
Linimentum Ammoniæ Subcarbonatis, P.L. 1824.

Take of Solution of Sesquicarbonate of Ammonia a
fluidounce,
Olive Oil three fluidounces;
Shake them together until they are mixed.

In the first of these preparations the union between the ammonia and oil is most perfect, on account of the carbonic acid combined with the ammonia of the last; but in both cases a kind of fluid soap is formed. They are used as stimulants in cynanche tonsillaris, spread on flannel, and applied round the throat.

LINIMENTUM CAMPHORÆ.

Liniment of Camphor.

Linimentum Camphoræ, P.L. 1809, P.L. 1824.

Take of Camphor an ounce,
Olive Oil four fluidounces ;
Dissolve the Camphor in the Oil.

This is employed as a stimulant embrocation to sprains and bruises, and in rheumatism.

LINIMENTUM CAMPHORÆ COMPOSITUM.

Compound Liniment of Camphor.

Linimentum Camphoræ compositum, P.L. 1788,
P.L. 1809, P.L. 1824.

Take of Camphor two ounces and a half,
Solution of Ammonia seven fluidounces and
a half,
Spirit of Lavender a pint ;
Mix the Solution of Ammonia with the Spirit ; then let
a pint distil from a glass retort, with a slow fire ; lastly,
dissolve the Camphor in it.

This is used for the same purposes as the former, and is much more powerful on account of the ammonia which it contains.

LINIMENTUM HYDRARGYRI COMPOSITUM.

Compound Liniment of Mercury.

Linimentum Hydrargyri, P.L. 1809.

Linimentum Hydrargyri compositum, P.L. 1824.

Take of stronger Ointment of Mercury,

Lard, each four ounces,

Camphor an ounce,

Rectified Spirit a fluidrachm,

Solution of Ammonia four fluidounces ;

Rub the Camphor, first with the Spirit, then with the Lard and Ointment of Mercury ; lastly, the Solution of Ammonia being gradually poured in, mix them all.

This liniment is stimulant and discutient. One drachm, containing nearly ten grains of mercury, may be rubbed on the affected part night and morning. It is said to salivate sooner than mercurial ointment, when freely employed.

LINIMENTUM OPII.

Liniment of Opium.

Take of Liniment of Soap six fluidounces,

Tincture of Opium two fluidounces ;

Mix.

This is a useful sedative liniment.

LINIMENTUM SAPONIS.

Liniment of Soap.

Linimentum Saponaceum, P.L. 1745.

Linimentum Saponis compositum, P.L. 1788,
P.L. 1809, P.L. 1824.

Take of Soap three ounces,
Camphor an ounce,
Spirit of Rosemary sixteen fluidounces ;

Dissolve the Camphor in the Spirit ; afterwards add the Soap, and macerate with a gentle heat until it is dissolved.

This is a stimulant application ; it is less powerful than the *Linimentum Camphoræ Compositum*, but is used for similar purposes.

LINIMENTUM TEREBINTHINÆ.

Liniment of Turpentine.

Take of Soft Soap two ounces,
Camphor an ounce,
Oil of Turpentine sixteen fluidounces ;
Shake them together until they are mixed.

This is a more powerful stimulant application than the preceding.

MELLITA.

Preparations of Honey.

MEL BORACIS.

Honey of Borax.

Mel Boracis, P.L. 1809, P.L. 1824.

Take of Borax, powdered, a drachm,
Honey an ounce ;

Mix.

Medicinal Uses.—Detergent and cooling in aphthous affections of the tongue and fauces.

MEL ROSÆ.

Honey of Rose.

Mel Rosatum, P.L. 1720.

Mel Rosaceum, P.L. 1745.

Mel Rosæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Red Rose [Petals] dried, four ounces,
Boiling distilled Water two pints and a half,
Honey five pounds ;

Macerate the Rose Petals in the Water for six hours ;
then add the Honey to the strained liquor, and, in a water
bath, boil down to a proper consistence.

Medicinal Use.—As an adjunct to detergent and astringent gargles.

OXYMEL.

Oxymel.

Oxymel Simplex, P.L. 1720, P.L. 1745.

Mel Acetatum, P.L. 1788.

Oxymel, P.L. 1809.

Oxymel Simplex, P.L. 1824.

Take of Honey ten pounds,

Acetic Acid a pint and a half;

Mix the Acid with the Honey made hot.

Medicinal Use.—Detergent; principally used as the basis of gargles and expectorant remedies. Dose, fʒj. to fʒj.

OXYMEL SCILLÆ.

Oxymel of Squill.

Oxymel Scilliticum, P.L. 1720, P.L. 1745.

Oxymel Scillæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Honey three pounds,

Vinegar of Squill a pint and a half;

Boil down in a glass vessel, with a slow fire, to a proper consistence.

Medicinal Use.—Expectorant. Dose, fʒss. to fʒij. in chronic coughs. In large doses it is emetic.

METALLICA.

METALLIC PREPARATIONS.

PRÆPARATA EX ALUMINIO.

Preparations of Aluminium.

ALUMEN EXSICCATUM.

Dried Alum.

Alumen Ustum, P.L. 1788.

Alumen Exsiccatum, P.L. 1809, P.L. 1824.

Let Alum liquefy in an earthen vessel over the fire ; then let the fire be increased, until the ebullition has ceased.

Remarks.—*Aluminium* is a metal obtained with considerable difficulty ; it is of a grey colour, generally in small scales or spangles of a metallic lustre ; alumina is the only known oxide of this metal, it is by Berzelius considered as a sesquioxide, and by Dr. Thomson as a protoxide composed of 1 equivalent of oxygen 8, and 1 equivalent of aluminium 10, the number of alumina being consequently 18.

Symbols,—Berzelius and Turner Al.

Brande AL.

Alum is a well-known double salt which crystallizes in regular octahedrons ; it is styptic and astringent. It is unaltered by exposure to the air ; dissolves in 18 times its weight of water at 60°, and in $\frac{3}{4}$ ths of its weight at 212°. The solution reddens litmus paper.

Composition.—Considered as double salt, alum, according to Dr. Thomson, consists of

Three eqs. of Sulphate of Alumina	$58 \times 3 = 174$	or 35.73
One eq. of Sulphate of Potash ..	88	„ 18.07
Twenty-five equivalents of Water	$9 \times 25 = 225$	„ 46.20
	<hr/>	<hr/>
	487	100.

Symbols.—Berzelius and Turner.. $\dot{\text{K}} \ddot{\text{S}} + \dot{\text{Al}} \ddot{\text{S}}^3 + \dot{\text{H}}^{25}$.

Brande $3(\text{AL} + \text{S}') + (\text{P} + \text{S}') + 25\text{q}$.

Process.—When alum is moderately heated as directed, it fuses in its water of crystallization, and when this is expelled it becomes spongy and opaque; if the heat be too strong, then a portion of sulphuric acid is driven off with the water.

Impurities and Tests.—See Notes: ALUMEN.

Incompatibles.—Alkalis and their carbonates; lime and lime-water, magnesia and its carbonate, tartrate of potash, acetates of lead, &c.

Official Preparations.—Alumen Exsiccatum. Liquor Aluminis Compositus.

Medicinal Uses.—Alum is internally a powerful astringent in hæmorrhages and inordinate fluxes, and is externally useful in repellent astringent lotions and collyria. Dose, gr. x. to gr. xx.

LIQUOR ALUMINIS COMPOSITUS.

Compound Solution of Alum.

Aqua Aluminosa Bateanæ, P.L. 1745.

Aqua Aluminis composita, P.L. 1788.

Liquor Aluminis compositus, P.L. 1809, P.L. 1824.

Take of Alum,

Sulphate of Zinc, of each an ounce,

Boiling Water three pints;

Dissolve the Alum and Sulphate of Zinc together in the water; afterwards strain.

Medicinal Uses.—This solution is powerfully astringent, and is successfully used as a detergent lotion to old ulcers, as a collyrium and as an injection; it will also often remove chilblains, and relieve slight excoriations.

PRÆPARATA EX ANTIMONIO.

Preparations of Antimony.

ANTIMONII OXYSULPHURETUM.

Oxysulphuret of Antimony.

Sulphur Antimonii præcipitatum, P.L. 1745, P.L. 1788.
Antimonii Sulphuretum præcipitatum, P.L. 1809, P.L. 1824.

Take of Sesquisulphuret of Antimony, powdered, seven ounces,

Solution of Potash four pints,

Distilled Water two gallons,

Diluted Sulphuric Acid as much as may be sufficient ;

Mix the Sesquisulphuret of Antimony, Solution of Potash and Water together, and boil with a slow fire for two hours, frequently stirring, distilled Water being often added, that it may fill about the same measure. Strain the liquor, and gradually drop into it as much diluted Sulphuric Acid as may be sufficient to throw down the Oxysulphuret of Antimony ; then wash away the Sulphate of Potash with water and dry what remains with a gentle heat.

Remarks.—Antimony is a white brilliant brittle metal, the specific gravity of which is about 6·7 ; it is often called *regulus of Antimony* to distinguish it from what is termed *crude Antimony*, which is the sesquisulphuret of the Pharmacopœia. Sesquisulphuret of antimony is the most abundant ore of the metal; it is a brittle

grey substance, which has usually a striated crystalline appearance. It is composed of

One and a half equivalent of Sulphur.. = 24
 One equivalent of Antimony = 65

Equivalent 89

Symbols.—Berzelius and Turner... $2\text{Sb} + 3\text{S}$ or $\text{Sb}^2 \text{S}^3$.

Brande $(an + 1\frac{1}{2} s)$.

Process.—When the sesquisulphuret of antimony is boiled as directed in a solution of potash (the oxide of the metal potassium), a portion of each is decomposed, and the results are sulphuret of potassium and sesquioxide of antimony, mixed with a large proportion of the sesquisulphuret dissolved in, but not decomposed by the potash. When the diluted sulphuric acid is added to this solution, it saturates the potash, sulphate of potash is formed, and remains in solution, while the oxide and sulphuret of antimony are precipitated together, forming oxysulphuret of antimony. The sulphuric acid acting at the same time upon the sulphuret of potassium and water, the water is decomposed, its hydrogen combines with the sulphur of the sulphuret and forms hydrosulphuric acid which is expelled in the state of gas, and its oxygen uniting with the potassium converts it into potash, with which the sulphuric acid combines.

Properties.—This preparation is of a bright orange colour, and has a slightly styptic taste. It is insoluble in water, and the greater portion of it is not readily acted upon by dilute acids. When boiled in a solution of twice its weight of bitartrate of potash, I find that about 12 per cent. of sesquioxide of antimony is dissolved.

Composition.—It is generally admitted by chemists that the composition of this preparation is uncertain; the specimen which I examined was a hydrated oxysulphuret, consisting very nearly of

Sesquioxide of Antimony..... 12.
 Sesquisulphuret of Antimony 76.5
 Water 11.5

100.

Impurities and Tests.—See Notes: ANTIMONII OXYSULPHURETUM.

Pharmacopœia Preparations.—*Pilulæ Hydrargyri Chloridi compositæ.*

Medicinal Uses.—It is but seldom employed, except in the above-named preparation, being, on account of its variable composition, less certain in its operation than other antimonials.

Dose.—In herpetic and other eruptions, from gr. j. to gr. iv. twice a day. In larger doses it is emetic.

ANTIMONII POTASSIO-TARTRAS.

Potassio-tartrate of Antimony.

Tartarus Emeticus, P.L. 1720.

Tartarum Emeticum, P.L. 1745.

Antimonium Tartarizatum, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Sesquisulphuret of Antimony, rubbed to powder,

Nitrate of Potash, powdered, each two pounds,

Bitartrate of Potash, powdered, fourteen ounces,

Hydrochloric Acid four fluidounces,

Distilled Water a gallon;

Mix the Sesquisulphuret of Antimony, accurately, with the Nitrate of Potash; the Hydrochloric Acid being then added, and the powder spread upon an iron plate, ignite it. Rub what remains to very fine powder, when it is cold, and wash it with boiling water until it is free from taste. Mix the powder thus prepared with the Bitartrate of Potash, and boil for half an hour in a gallon of distilled water. Strain the liquor while yet hot, and set it aside that crystals may be formed. These being removed and dried, let the liquor again evaporate that it may yield crystals.

Remarks.—In the last Pharmacopœia *glass of antimony* was used in preparing this medicine; it is, however, not only difficult to obtain it, but glass of lead is frequently substituted for, and what is still worse, mixed with it. By the present formula an oxysulphuret of antimony is obtained similar in composition to the *crocus antimonii* of the Pharmacopœia of 1788; in that preparation, however, the mixture was fused, which rendered the sesquioxide less easily soluble in the bitartrate of potash than that obtained by the present process.

Process.—When nitrate of potash is mixed and ignited with sulphuret of antimony, rapid combustion ensues; both are decomposed; a portion of the sulphur of the sulphuret combines with part of the oxygen of the decomposed nitric acid, and the result is sulphuric acid, which forms sulphate of potash, with the potash of the decomposed nitrate; another portion of the oxygen unites with the antimony of the decomposed sulphuret, and sesquioxide of antimony is formed which remains mixed with sulphate of potash, and some sulphuret of antimony; and there would also be free potash and sulphuret of potassium, were it not for the hydrochloric acid employed, which saturates the alkali, and either prevents the formation of the sulphuret or immediately decomposes it when formed. When the residue of the combustion is washed, as directed, sulphate of potash and chloride of potassium are removed by it, and a mixture of sesquioxide and sulphuret of antimony remains, to which perhaps the name of oxysulphuret may be given, as well as to the last preparation, though it is not certain that either of them is a definite compound.

Sesquioxide of Antimony, the base of this preparation, is composed of

One and a half equivalent of Oxygen	12
One equivalent of Antimony	65
	<hr/>
Equivalent	77

Symbols,—Berzelius and Turner . . . Sb

Brande $(an + 1\frac{1}{2}o)$.

Bitartrate of potash is a salt containing two equivalents of tartaric acid and one equivalent of potash; when it is boiled in water with the oxysulphuret of antimony, the equivalent of acid in excess combines with two equivalents of sesquioxide of antimony, and the sulphuret remains unacted upon; the solution therefore consists of two equivalents of tartaric acid, one equivalent of potash, and two equivalents of sesquioxide of antimony, and these combining crystallize together as a double salt, or potassio-tartrate of antimony.

Antimonii Potassio-tartras, or
Tartrate of Potash and Ditartrate of Antimony.

Bitartrate of Potash.	{	2 eqs. Tartaric Acid.	2 eqs. Sesquioxide	} Oxysul- phuret of Antimony.
		1 eq. Potash.	of Antimony.	
		Sulphuret of Antimony.		

Composition.—Antimonii Potassio-tartras is a compound of

One equivalent of Tartrate of Potash.....	66 + 48 =	114
One equivalent of Ditartrate of Antimony	66 + 154 =	220
Three equivalents of Water	9 × 3 =	27
Equivalent.....		361

Symbols,—Berzelius and Turner.. $(\dot{\text{K}}\bar{\text{T}} + \ddot{\text{S}}\bar{\text{b}}\bar{\text{T}}) + 2\text{H}$.

Brande..... $(2\text{tar}' + \text{P} + 2\text{an} + 1\frac{1}{2}\text{o} + 3\text{q})$.

Or it contains

Two equivalents of Tartaric Acid	66 × 2 =	132	or	36·6
One equivalent of Potash.....	=	48	„	13·3
Two eqs. of Sesquioxide of Antimony..	77 × 2 =	154	„	42·6
Three equivalents of Water	9 × 3 =	27	„	7·5
Equivalent.....		361		100·

Properties.—Tartarized antimony crystallizes with great facility, and the general character of the crystals of this compound is that of an *octahedron with a rhombic base*. One distinct cleavage only has been obtained, which is parallel to the plane a of the accompanying figure. The planes z and y are generally striated.

The following are the nearest to coinciding measurements taken on several crystals :



P on P'	108° 16'
P over the edge on the left...	104 15
P on z^1	166 40
P on z^2	165 40 nearly
a on P, or P'	122 00
a on y	90 00

The crystals of this salt are colourless and inodorous, but have a styptic metallic taste: on exposure to the air, they effloresce slightly and become opake. When strongly heated this salt is decomposed, and an alloy of antimony and potassium is obtained. It is soluble in about fifteen times its weight of water at 60°, and twice its weight at 212°. The aqueous solution decomposes spontaneously after it has been some time prepared. It is insoluble in alcohol.

Impurities and Tests.—See Notes : ANTIMONII POTASSIO-TARTRAS.

Adulteration.—This salt should never be purchased in powder, but always in crystals : in the former state it frequently contains a portion of bitartrate of potash uncombined with any oxide of antimony, and which, in preparing the vinum antimonii potassio-tartratis, is precipitated. To judge if the crystals have been properly prepared, drop one or two into a solution of hydro-sulphuric acid, and an orange-coloured deposit will be formed on them.

Incompatibles.—The solution of this salt is decomposed by acids, by alkalis and their carbonates, by some of the earths and metals, and their oxides, by lime water, chloride of calcium, and the acetates of lead. Many vegetable infusions, and especially those which are bitter and astringent, decompose it, such as cinchona, rhubarb, catechu, &c.

Medicinal Uses.—Potassio-tartrate of antimony either sweats, vomits, or purges, according to the quantity exhibited. A *quarter of a grain*, if the skin be kept warm, will promote a diaphoresis; *half a grain* will first prove purgative, and then diaphoretic; and *one grain* will generally vomit, then purge, and lastly sweat the patient. It may be given in solution.

VINUM ANTIMONII POTASSIO-TARTRATIS.

Wine of Potassio-tartrate of Antimony.

Vinum Antimonii Tartarizati, P.L. 1788.

Liquor Antimonii Tartarizati, P.L. 1809, P.L. 1824.

Take of Potassio-tartrate of Antimony two scruples,
Sherry Wine a pint ;

Dissolve the Potassio-tartrate of Antimony in the Wine.

Process.—The College have restored the use of wine in this preparation. When the antimonial salt has been carelessly prepared, and contains bitartrate of potash uncombined with oxide of antimony, a deposit is apt to be formed in this solution; those practitioners therefore, who purchase the potassio-tartrate should insist on having it in the state of crystals, in which there is but little chance of the occurrence of this imperfection. If any deposit be observed in this preparation it ought to be rejected; each fluidounce contains two grains of the potassio-tartrate.

Incompatibles.—See ANTIMONII POTASSIO-TARTRAS.

Medicinal Uses.—In doses of ℥xv. to fʒj. it acts as a diaphoretic when given with saline medicines, warm drinks, &c.

PULVIS ANTIMONII COMPOSITUS.

Compound Powder of Antimony.

Pulvis Antimonialis, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Sesquisulphuret of Antimony, powdered, a pound,

Horns, shaved, two pounds;

Mix and throw them into a crucible red hot in the fire, and stir constantly until vapour no longer arises. Rub that which remains to powder, and put it into a proper crucible. Then apply fire, and increase it gradually that it may be red hot for two hours. Rub the residue to very fine powder.

Process.—Sesquisulphuret of antimony, as already mentioned, consists of sulphur and antimony; horn shavings are composed of phosphate of lime mixed with carbonaceous animal matter. When the sulphuret and horn are heated together, the sulphur is expelled in vapour; and the antimony, combining with the oxygen of the air, is converted into antimonious acid. Part of the animal matter is dissipated by the heat, but the phosphate of lime suffers no change; there remains, therefore, in the crucible, a mixture of antimonious acid, consisting of two equivalents of oxygen 16 and 1 equivalent of antimony 65 = 81, and phosphate of lime, forming *Pulvis Antimonii compositus*.

Qualities.—This preparation is an inodorous insipid powder, of a dull white colour. It is insoluble in water, and only partially soluble in acids; if, however, the antimony it contains were in the state of sesquioxide, as has been stated, then hydrochloric acid, when heated, would entirely dissolve it.

Composition.—In consequence of Dr. Elliotson's statement, that he had exhibited upwards of 100 grains of this medicine without producing any effect, I procured specimens of it from two respectable sources, and subjected them to analysis. I found one of them to consist of

Antimonious Acid	35
Phosphate of Lime	65

The other yielded

Antimonious Acid	38
Phosphate of Lime	62
	<hr/>
	100

I have also analysed James's powder, of which the Pulvis Antimonii Compositus is an imitation, and found it to consist nearly of

Antimonious Acid	56
Phosphate of Lime	44
	<hr/>
	100

These proportions agree almost exactly with the results obtained by Dr. Pearson; and the high state of oxidizement of the antimony will fully account for the inactivity of both preparations.

Adulteration.—No doubt can be entertained that this preparation, like every other, has been sophisticated; but owing to want of power in the genuine article, the practitioner probably has not been disappointed by its adulteration.

Medicinal Uses.—It is stated to be diaphoretic, alterative, emetic, or purgative, according to the extent of the dose and the state of the patient. The doses mentioned are from gr. v. to gr. x. It is worth the consideration of the practitioner, whether the employment of this preparation may not be altogether superseded by that of potassio-tartrate of antimony.

PRÆPARATA EX ARGENTO.

Preparations of Silver.

ARGENTI NITRAS.

Nitrate of Silver.

Causticum Lunare, P.L. 1720, P.L. 1745.

Argentum Nitratum, P.L. 1788.

Argenti Nitras, P.L. 1809, P.L. 1824.


Take of Silver an ounce and a half,

Nitric Acid a fluidounce,

Distilled Water two fluidounces;


Mix the Nitric Acid with the Water, and dissolve the Silver in them in a sand-bath. Afterwards increase the heat gradually, that the Nitrate of Silver may be dried. Liquefy this in a crucible, with a slow fire, until, the Water being expelled, ebullition has ceased; then immediately pour it into proper moulds.

Process.—Nitric acid is composed of oxygen and azote, and when silver is dissolved in it, a portion of the acid is partially decomposed into nitric oxide gas and oxygen; the former escapes into the atmosphere, and separating a portion of its oxygen from admixture with the azotic gas, red nitrous acid gas is formed by their union. The oxygen of the decomposed acid unites with the silver to form oxide of silver, composed of 1 equivalent of oxygen 8 + 1 equivalent of silver 108 = 116, and this is dissolved by the nitric acid undecomposed, and converted into nitrate of silver.



Properties.—Solution of nitrate of silver readily yields transparent colourless crystals, the primary form of which is a *right rhombic prism*.

P on <i>d</i>	116°	36'
M on <i>d</i>	148	0
M on M'	129	31
<i>d</i> on <i>d'</i>	126	48



In some crystals the planes *d* are barely visible, while in others those planes encroach so much on *M* and *M'* as to leave only minute portions of them discernible.

Water, at the temperature of 60° , dissolves its own weight of this salt. It is not deliquescent. By exposure to a strong light it becomes brown, owing to the reduction of a part of the silver to the metallic state. When moderately heated it readily melts, swells, and then remains liquid. On cooling it forms a grey-coloured mass, having a striated and crystalline structure. If subjected to a red heat it is decomposed. It stains the skin black: the crystals contain no water.

Composition.—Nitrate of silver is composed of

One equivalent of Nitric Acid . . .	=	54	or	31.76
One equivalent of Oxide of Silver	=	116	,,	68.24
<hr/>				
Equivalent		170		100.

Symbols,—Berzelius and Turner . . $\text{Ag} + \ddot{\text{N}}$.

Brande $(ag + n')$.

Incompatibles.—Almost all spring and river water, on account of the common salt which they usually contain; chlorides; the alkalis, potash, soda and their carbonates; lime-water. Ammonia added in excess redissolves the precipitate as first formed; the sesquicarbonate throws down carbonate of silver. The sulphuric, hydrochloric and tartaric acids, and the salts which contain them, decompose nitrate of silver. It is decomposed by hydrosulphuric acid and its salts, by the soluble sulphurets and astringent vegetable infusions.

Pharmacopæia Preparations.—Liquor Argenti Nitratis. Argenti Cyanidum.

Medicinal Uses.—It is the most manageable and powerful of all escharotics. Internally it is tonic and antispasmodic, and has been especially exhibited in cases of epilepsy; when it has been long taken it is sometimes deposited in the rete mucosum, so as to give a permanent dark purple hue to the patient. Dose, *one eighth of a grain* gradually increased to *one grain*. But very much larger doses have been given. It should be made into pills with crumb of bread, and mixed with a little sugar to prevent the mass from becoming too hard.

LIQUOR ARGENTI NITRATIS.

Solution of Nitrate of Silver.

Take of Nitrate of Silver a drachm,
Distilled Water a fluidounce;

Dissolve the Nitrate of Silver in the Water and strain; then the access of light being prevented, keep it in a well-closed vessel.

Remarks.—This is employed in many cases to determine the presence of chlorides and hydrochloric acid, with which it gives a white precipitate of chloride of silver insoluble in acids and the fixed alkalis, but readily dissolved by ammonia, and precipitated from it by saturation with an acid.


ARGENTI CYANIDUM.

Cyanide of Silver.

Take of Nitrate of Silver two ounces and two drachms,
Diluted Hydrocyanic Acid,
Distilled Water, each a pint;

Dissolve the Nitrate of Silver in the Water, and add to them the diluted Hydrocyanic Acid, and mix. Wash what is precipitated with distilled Water, and dry it.

Process.—The nature and composition of hydrocyanic acid and nitrate of silver have already been stated. When solutions of them are mixed as directed, the cyanogen of the hydrocyanic acid combines with the silver of the oxide to form cyanide of silver, which is precipitated, and the hydrogen of the acid unites with the oxygen of the oxide to form water, which remains in solution with the nitric acid of the nitrate of silver.



Properties.—Cyanide of silver is precipitated in the form of a white powder. It is insoluble in water, and dissolves in sul-

phuric and nitric acids only when they are concentrated and very hot. Hydrochloric acid, hydrosulphuric acid and the hydrosulphates decompose it readily. The alkalis, potash and soda do not dissolve it, but it easily dissolves in ammonia. It is decomposed by a red heat, and gives, when it has been well dried, silver and cyanogen gas; but when it contains water, it yields hydrocyanic acid and cyanogen, and leaves silver mixed with charcoal.

Composition.—This cyanide is composed of

One equivalent of Cyanogen....	=	26	or	19·4
One equivalent of Silver.....	=	108	„	80·6
		<hr/>		<hr/>
Equivalent....		134		100·

Symbols.—Berzelius and Turner..... Ag + Cy, or Ag Cy.

Brande..... (*ag* + *cy*).

Impurities and Tests.—See Notes: ARGENTI CYANIDUM.

Incompatibles.—Sulphuric and nitric acids, when concentrated and hot. Hydrochloric acid, hydrosulphuric acid and hydrosulphates, and ammonia.

Pharmacopœia Preparation.—It is employed for the extemporaneous preparation of hydrocyanic acid.

PRÆPARATUM EX ARSENICO.

Preparation of Arsenic.

LIQUOR POTASSÆ ARSENITIS.

Solution of Arsenite of Potash.

Liquor Arsenicalis, P.L. 1809, P.L. 1824.

Take of Arsenious Acid, broken into small pieces,
 Carbonate of Potash, each eighty grains,
 Compound Tincture of Lavender five fluidrachms,
 Distilled Water a pint;

Boil the Arsenious Acid and Carbonate of Potash with half a pint of the Water in a glass vessel until they are dissolved. Add the Compound Tincture of Lavender to the cooled liquor. Lastly, add besides, of distilled Water, as much as may be sufficient, that it may accurately fill a pint measure.

Remarks.—Arsenic is a grey brittle metal, which when fresh broken has considerable lustre. Its specific gravity is 5.884. It combines with two different quantities of oxygen, forming *arsenious acid* and *arsenic acid*. Arsenious acid, employed in the Pharmacopœia, is formed of $1\frac{1}{2}$ equivalent of oxygen 12, and 1 equivalent of arsenic $38=50$. Arsenic acid consists of $2\frac{1}{2}$ equivalents of oxygen 20, and 1 equivalent of metal $38=58$.

Properties of Arsenious Acid.—It occurs in large colourless pieces, which externally are usually opaque; but internally, when recently broken, they are frequently transparent, and have the appearance of glass. This substance, called also *white arsenic*, and oxide of arsenic, is moderately hard and brittle; it is inodorous, has hardly any taste, and is extremely poisonous. Its specific gravity, when transparent, I find to be 3.715, and when opaque 3.260; the opacity I believe to be owing to the absorption of water from the atmosphere. Arsenious acid is volatilized at the temperature of about 380° , and when thus vaporized, it is inodorous, although usually stated to possess an alliaceous smell, which belongs only to volatilized metallic arsenic. A thousand parts of water at a mean temperature are said to dissolve 9.6 parts of transparent and 12.5 of opaque arsenious acid in 36 hours; the same quantity of boiling water dissolves 97 parts of the transparent kind, of which 18 are retained on cooling and 79 deposited in the state of small crystals, the form of which is the regular octahedron. The solution of white arsenic reddens litmus paper slightly, and it combines with the alkalis, potash and soda, with great facility; when heated in nitric acid it absorbs more oxygen, and is rendered much more powerfully acid, being converted into arsenic acid.

Process.—This solution of arsenite of potash is very readily prepared; a few minutes' ebullition in a Florence flask is sufficient to dissolve the arsenious acid; during ebullition carbonic acid gas is evolved, owing to the greater affinity existing between the arsenious acid and potash, than between carbonic acid and potash.

In preparing *Liquor Potassæ Arsenitis*, the arsenious acid usually sold in powder should not be employed; it is very com-

monly adulterated with sulphate of lime or gypsum, which being insoluble in the solution of carbonate of potash, the operator supposes that it is difficult to prepare this medicine. This adulteration, and most others likely to occur, may be detected by heating the powder in a crucible; whatever is not volatilized is an impurity, &c.

Incompatibles.—Acids and acidulous salts, hydrosulphuric acid and hydrosulphates, sulphuret of potassium and similar compounds, lime water; earthy salts, such as alum, sulphate of magnesia, and chloride of calcium; metallic salts, as sulphate and sesquichloride of iron, nitrate of silver, and sulphate of copper; decoction of cinchona.

Medicinal Uses.—This solution is a powerful tonic: it is especially employed in intermittent and remittent fevers, periodic headaches, and some diseases of the skin. Dose, *four minims to thirty minims*, twice a day.

PRÆPARATA È BARIO.

Preparations of Barium.

BARII CHLORIDUM.

Chloride of Barium.

Take of Carbonate of Barytes, broken into small pieces,
ten ounces,

Hydrochloric Acid half a pint,

Distilled Water two pints;

Mix the Acid with the Water, and add the Carbonate of Barytes gradually to them. Then, heat being applied, and the effervescence finished, strain and boil down the liquor, that crystals may be formed.


Remarks.—Barium is a metal best known in combination with oxygen, forming the alkaline earth or metallic oxide barytes; this is met with in large quantity in the state of sulphate and of carbonate of barytes.

Barytes or oxide of barium when pure is nearly colourless,

inodorous, soluble in water, and when dissolved in hot water yields crystals on cooling. It has strongly marked alkaline properties, a disagreeable caustic taste, is extremely poisonous and combines readily with acids. It is composed of 1 equivalent of oxygen 8, and 1 equivalent of barium $68=76$.

Carbonate of barytes is usually found in amorphous masses, though sometimes crystallized. It is moderately hard, but easily powdered, nearly or quite colourless, sometimes translucent and very heavy. It is nearly insoluble in water, but readily decomposed by acids with effervescence. It is composed of 1 equivalent of carbonic acid 22, and 1 equivalent of barytes $76=98$.

Process.—When carbonate of barytes is acted upon by hydrochloric acid, not only is the carbonic acid expelled and the barytes dissolved, but owing to the mutual decomposition of the barytes or oxide of barium and hydrochloric acid, water and chloride of barium are formed.



Properties.—Chloride of barium yields a colourless solution which by evaporation gives crystals in rhombic plates, of which 100 parts of water at 60° dissolve about 40 parts; the boiling-point of a saturated solution is 222° , and this consists of 100 parts of water and 78 of crystals. The solution is immediately decomposed by sulphuric acid and sulphates, and by the alkaline carbonates, but ammonia gives no precipitate in it; by exposure to a moderate degree of heat, the crystals lose their water; but the salt is not decomposed even at a very high temperature.

Anhydrous Chloride of Barium consists of

One equivalent of Chlorine.....	= 36 or 34.6
One equivalent of Barium	= 68 „ 65.4

Equivalent....	104	100.
----------------	-----	------

The crystals consist of

One equivalent of Anhydrous Chloride of Barium..	104 or 85.3
Two equivalents of Water.....	18 „ 14.7
	122 100.

Symbols.—Berzelius and Turner $\text{Ba Cl} + \text{H}^2$.

Brande $(ba + c + 2q.)$

Uses.—This salt has been occasionally employed in medicine ; but it is principally used as a reagent for detecting the presence of sulphuric acid and sulphates ; for which purpose the following solution is directed to be prepared.

LIQUOR BARI CHLORIDI.

Solution of Chloride of Barium.

Take of Chloride of Barium a drachm,
 Distilled Water a fluidounce ;
 Dissolve the Chloride of Barium and strain.

PRÆPARATUM È BISMUTHO.

Preparation of Bismuth.

BISMUTHI TRISNITRAS.

Trisnitrate of Bismuth.

Bismuthi Subnitrates, P.L. 1824.

Take of Bismuth an ounce,
 Nitric Acid a fluidounce and a half,
 Distilled Water three pints ;

Mix a fluidounce of the Water with the Nitric Acid, and dissolve the Bismuth in them ; then pour off the solution. To this add the remainder of the Water, and set by that the powder may subside. Afterwards, the super-

natant liquor being poured off, wash the Trisnitrate of Bismuth with Distilled Water, and dry it with a gentle heat.

Remarks.—Bismuth is a metal of a reddish white colour; its structure is usually crystalline, and by cautious cooling after it has been melted, it may be made to assume a cubic form. It is a brittle metal; its specific gravity is 9.882; it melts at the temperature of 476°. By exposure to the air it tarnishes, but does not readily oxidize; but when heated in the air it is converted into a yellow oxide, consisting of

One equivalent of Oxygen	=	8	or	10
One equivalent of Bismuth	=	72	„	90
		<hr/>		<hr/>
Equivalent		80		100.

Symbols,—Berzelius and Turner $\text{Bi} + \text{O}$ or Bi .

Brande ($bi + o$).

Process.—In preparing the nitrate of bismuth, part of the nitric acid is decomposed, with the occurrence of phenomena and effects similar to those which have been described as taking place during the solution of silver in nitric acid. See ARGENTI NITRAS. The oxide of bismuth formed is held in solution by the nitric acid remaining undecomposed.

The solution of nitrate of bismuth is colourless, and when water is added to it, as directed in the formula, it combines with the greater part of the acid, which, however, retains some oxide of bismuth in solution, while a white precipitate is formed, which is a trisnitrate of bismuth, composed of

One equivalent of Nitric Acid	=	54	or	18.36
Three equivs. of Oxide of Bismuth 80×3	=	240	„	81.64
		<hr/>		<hr/>
Equivalent		294		100.

Symbols,—Berzelius and Turner $3\ddot{\text{Bi}} + \ddot{\text{N}}$.

Brande ($3\text{Bi} + n'$).

Peroxide of bismuth is a heavy brown powder, it is a sesquioxide, composed of $1\frac{1}{2}$ equivalent of oxygen $12 + 72$ bismuth $= 84$. It does not form salts with acids, but is decomposed by them with the evolution of oxygen.

Qualities.—This substance was formerly employed as a cosmetic, under the name of magistery of bismuth. It is a white,

inodorous, tasteless powder, insoluble in water. It is rendered black by hydrosulphuric acid.

Medicinal Uses.—This medicine is represented to possess antispasmodic powers, and to be especially serviceable in those forms of dyspepsia which are attended with painful contractions of the stomach. Dose, from gr. v. to gr. xv.

PRÆPARATA È CALCIO.

Preparations of Calcium.

C A L X.

Lime.

Calx Viva, P.L. 1720.

Calx, P.L. 1809, P.L. 1824.

Take of Chalk a pound ;

Break it into small pieces, and burn it in a very strong fire for an hour.

Remarks.—Chalk is composed of carbonic acid and lime, and lime is the oxide of the metal calcium ; it is composed of

One equivalent of Oxygen..... = 8 or 28.58

One equivalent of Calcium = 20 „ 71.42

Equivalent 28 100.

Symbols,—Berzelius and Turner.. $\text{Ca} + \text{O}$ or $\dot{\text{Ca}}$.

Brande (*cal* + *o*) or C.

Carbonate of lime consists of

One equivalent of Carbonic Acid .. = 22 or 44

One equivalent of Lime = 28 „ 56

Equivalent.... 50 100.

Symbols,—Berzelius and Turner .. $\dot{\text{Ca}}$ $\ddot{\text{C}}$.

Brande ($\text{C} + \text{car}'$).

Process.—By the action of heat the carbonic acid is expelled from marble or carbonate of lime; and, as the pure part of chalk, marble, and limestone contains 44 per cent. of carbonic acid, 100 parts should furnish 56 of lime. If the quantity remaining exceed this, the excess must be derived either from earthy impurity, or from a portion of the chalk or limestone remaining undecomposed by the heat. The impurities which limestone contains are insoluble in water, and unimportant in all cases for medicinal uses.

Properties.—Pure lime is colourless, moderately hard, but easily reduced to powder; unlike the limestone from which it is procured, it is sonorous, although but slightly. It is inodorous, and has a burning, alkaline taste, and it corrodes animal substances. Vegetable blue colours are changed to green by lime, and yellow to brown, evincing its alkaline properties. By exposure to the air it imbibes moisture and falls to powder, and is gradually reconverted to the state of carbonate by combining with the carbonic acid of the atmosphere.

When water is poured upon lime it is rendered extremely hot, swells, becomes powdery, and combining with a portion of the water is converted into hydrate of lime. Lime is slightly soluble in water, and the solution possesses alkaline properties. If lime be long exposed to atmospheric air it loses its property of slacking, owing to its having combined with water and carbonic acid, and it is then unfit for use.

Hydrate of lime, or, as it is usually termed, *slacked lime*, is composed of

One equivalent of Lime	=	28	or	75.68
One equivalent of Water	=	9	„	24.32
		<hr/>		<hr/>
Equivalent		37		100.

Incompatibles.—All acids and acidulous salts, alkaline carbonates, ammoniacal salts, metallic salts, borates, and astringent vegetable infusions.

Officinal Preparations.—Liquor Calcis, Liquor Ammoniae, Liquor Potassae, Potassa cum Calce, Calcii Chloridi.

LIQUOR CALCIS.

Lime Water.

Aqua Calcis, P.L. 1720.*Aqua Calcis Simplex*, P.L. 1745.*Aqua Calcis*, P.L. 1788.*Liquor Calcis*, P.L. 1809, P.L. 1824.

Take of Lime half a pound,

Distilled Water twelve pints;

First add a little Water to the Lime, when slacked add the remainder of the Water, and shake them together; then immediately cover the vessel, and set it by for three hours; afterwards keep the Solution with the remaining Lime in stopped glass vessels, and when it is to be used, take from the clear Solution.

Process.—This is a simple solution of lime in water. Unlike most other substances, lime is more soluble in cold water than in hot; and when lime water which has been prepared with cold water is heated, small crystals of lime, probably containing water, are formed and deposited.

I find that

A pint of water at 32° dissolves 13.25 grains of lime.

Ditto	60	11.6	ditto
-------	----	------	-------

Ditto	212	6.7	ditto
-------	-----	-----	-------

It is then evident that water at 32° takes up nearly one seventh more lime than water at 60°, and almost double the quantity dissolved by boiling water.

Properties.—Lime water is colourless and inodorous, but has a disagreeable alkaline taste. It turns vegetable blues green, and yellows brown; and it unites with oil by agitation, forming an imperfect soap. When lime water is exposed to the atmosphere it absorbs carbonic acid, a thin crust of carbonate of lime is rapidly formed on the surface, and eventually the whole of the lime is precipitated from solution: on this account lime water should be preserved from the air as carefully as possible.

Incompatibles.—Lime water is incompatible with the substances already enumerated with respect to lime itself.

Medicinal Uses.—It is antacid, and therefore useful in dyspepsia attended with acidity; it is also astringent in leucorrhœa, in the last stages of dysentery, and in protracted diarrhœa. Dose, in milk, f ʒj. to f ʒvj.

CALCII CHLORIDUM.


Chloride of Calcium.

Calcis Murias, P.L. 1824.

Take of Chalk five ounces,
 Hydrochloric Acid,
 Distilled Water, each half a pint ;

Mix the Acid with the Water ; and to these gradually add the Chalk to saturation. Then the effervescence being finished, strain ; evaporate the liquor until the salt is dried. Put this into a crucible, and having liquefied it in the fire, pour it upon a flat clean stone. Lastly, when it is cold break it into small pieces, and keep it in a well-closed vessel.

Process.—It has been already shown that hydrochloric acid is a compound of chlorine and hydrogen, and that lime is composed of calcium and oxygen ; when carbonate of lime is dissolved in the acid it is converted into chloride of calcium. The changes that occur are, that the carbonic acid is expelled in the state of gas ; the hydrogen of the acid combines with the oxygen of the lime to form water ; and the chlorine and calcium uniting constitute chloride of calcium ; the water used, and that formed, being expelled by evaporation and fusion.



Properties.—Chloride of calcium is colourless, translucent, and inodorous ; its taste is very bitter and pungent. By exposure to the air it deliquesces, and is of course very soluble in water ; at 60°, water dissolves nearly four times its

weight, and hot water a still larger quantity. It is also very soluble in alcohol. By evaporation the solution yields crystals containing a large quantity of water.

Composition.—This salt is composed of

One equivalent of Chlorine.....	= 36	or 64.3
One equivalent of Calcium.....	= 20	„ 35.7
	<hr/>	<hr/>
Equivalent....	56	100.

Symbols.—Berzelius and Turner..... $\text{Ca} + \text{Cl}$, or CaCl .

Brande..... $(\text{cal} + \text{c})$.

Impurities and Tests.—See Notes: *CALCII CHLORIDUM*.

Incompatibles.—This salt is decomposed by sulphuric acid and by sulphates, by the alkalis potash, soda, and their carbonates. If ammonia be added to the solution, no change occurs; but carbonate of ammonia decomposes it, and precipitates carbonate of lime.

For *Medicinal Uses*, see *LIQUOR CALCII CHLORIDI*.

LIQUOR CALCII CHLORIDI.

Solution of Chloride of Calcium.

Liquor Calcis Muriatis, P.L. 1824.

Take of Chloride of Calcium four ounces,

Distilled Water twelve fluidounces;

Dissolve the Chloride of Calcium, and strain.

Remarks.—The solution ordered in the last Pharmacopœia was about double the strength of the present; it was however so concentrated as to crystallize in cold weather.

Medicinal Uses.—Deobstruent and tonic; it is stated to have been advantageously given in bronchocele and scrofula.

Dose, ℥xl. to fʒij., or more.

CALX CHLORINATA.

Chlorinated Lime.

Take of Hydrate of Lime a pound,

Chlorine as much as may be sufficient ;


Pass Chlorine to the Lime, spread in a proper vessel, until it is saturated.

Chlorine is very readily evolved from Hydrochloric Acid added to Binoxide of Manganese, with a gentle heat.

Remarks.—The exact nature of this compound not having been yet determined, the term *chlorinated lime* is provisionally given to it.

This substance is prepared very largely for the use of the bleacher, and is called *bleaching powder*, when so employed. It was formerly termed also Oxymuriate of Lime ; it is now known by the name of Chloride of Lime. Berzelius however considers it as a chlorite, Balard as a hypochlorite of lime ; but in the absence of positive proof I shall consider it, what it has been long termed, a chloride of lime.

Process.—On this view of the subject, when hydrochloric acid acts upon binoxide of manganese, the changes that take place are these : two equivalents of hydrochloric acid are composed of two eqs. of hydrogen = 2, and two eqs. of chlorine = 72 ; one eq. of binoxide of manganese consists of two eqs. of oxygen = 16, and one eq. of manganese = 28 ; when these act upon each other, the two eqs. of hydrogen combine with the two eqs. of oxygen and form two eqs. of water, while one of the eqs. of chlorine unites with the one eq. of manganese to form chloride of manganese, and the other eq. of chlorine is evolved in the gaseous state, and being absorbed by the lime, chloride of lime, the *calx chlorinata* of the Pharmacopœia, is formed.



When chlorine gas ceases to be absorbed, the chloride of lime obtained appears, from the statements of Brande and Grouvelle, and also from my own experiments, to consist of

$$\begin{array}{rcl}
 \text{One equivalent of Chlorine} & \dots\dots\dots & = 36 \\
 \text{Two equivalents of Hydrate of Lime} & \dots\dots & 37 \times 2 = 74 \\
 & & \hline
 & & 110
 \end{array}$$

When water is added to this, the chloride of lime dissolves, leaving nearly all the lime insoluble; it is therefore probably composed of

$$\begin{array}{rcl}
 \text{One eq. of Bihydrated Chloride of Lime} & \dots\dots\dots & 18 + 36 + 28 = 82 \\
 \text{One equivalent of Lime} & \dots\dots\dots & = 28 \\
 & & \hline
 & & 110
 \end{array}$$

It appears extremely probable that the whole of the lime is not converted into chloride, from the deficiency of water; for when the dry chlorine acts upon anhydrous lime, the lime is decomposed, oxygen gas is evolved, and chloride of calcium remains. Dr. Thomson also states that a compound of one equivalent each of chlorine and lime is now formed at Glasgow; such a compound probably results from the intervention of water, as also indicated by the experiments of Houton-Labillardière.

Properties.—Chloride of lime, when pure, is white, but generally has a brownish tint; it emits a weak smell of chlorine, and its taste is strong. It is only partially soluble in water, the lime uncombined with chlorine being comparatively insoluble. It possesses powerful bleaching properties: when exposed to the air it is gradually decomposed, chlorine is given out, and carbonate of lime formed and precipitated. It is also decomposed by heat; some chlorine gas comes over first, and afterwards oxygen derived from the decomposition of the lime, chloride of calcium remaining.

The stronger acids, such as the nitric, hydrochloric and sulphuric, and the alkaline carbonates decompose it; the acids evolve chlorine copiously, and the carbonates precipitate carbonate of lime, while the alkaline chlorides formed remain in solution.

Uses.—It is employed as a disinfectant; when exposed to the air the carbonic acid which it contains evolves chlorine, this powerfully corrects the putrid odour arising either from diseased or decomposing animal matter. It is commonly called Labarracque's Disinfecting Fluid.

CRETA PRÆPARATA.

Prepared Chalk.

Creta, P.L. 1720.

Creta Præparata, P.L. 1745, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Chalk a pound,

Water as much as may be sufficient;

Add a little Water to the Chalk, and rub it to fine powder. Put this in a large vessel with the remainder of the Water; then stir it, and after a short interval, pour off the supernatant Water, still turbid, into another vessel, and set it by that the powder may subside; lastly, the Water being poured off, dry this Powder and keep it for use.

Shells, first freed from impurities and washed with boiling water, are prepared in the same manner.

Process.—This method of preparing this variety of carbonate of lime called chalk, is termed elutriation, and is an effectual method of reducing it to a fine powder.

Properties.—Chalk is a substance so well known that it is hardly requisite to notice its qualities. When pure it is very nearly white. It is dull, opaque, soft, and light, and it always occurs massive. Its sp. gr. is about 2·3; it is sometimes of a greyish tint, and then contains an admixture of foreign matter.

Composition.—By the analysis of Bucholz, chalk is composed of

Carbonic Acid.....	43
Lime	56·5
Water	·5

100·0

The water is an accidental admixture, and, when perfectly pure, carbonate of lime is composed of

One equivalent of Carbonic Acid	= 22	or 44
One equivalent of Lime	= 28	„ 56

Equivalent..... 50 100

Symbols.—Berzelius and Turner.. $\text{Ca} + \ddot{\text{C}}$.

Brande $(\text{C} + \text{car}')$.

Impurities and Tests.—See Notes : CRETA.

Adulteration.—Chalk is so cheap an article that accidental admixture only can be suspected. If, however, what is termed *grey chalk* be used, the prepared chalk will contain some foreign matter, and the colour will be less perfect.

Incompatibles.—Chalk, or carbonate of lime, is incompatible with acids and acidulous salts, for they combine with its base and expel the carbonic acid in the state of gas.

Officinal Preparations.—Ammoniae Sesquicarbonas, Calx, Calcii Chloridum, Hydrargyrum cum Creta, Mistura Cretæ, Confectio Aromatica.

Medicinal Uses.—It is antacid and absorbent, and therefore it is useful in acidities of the primæ viæ, and in diarrhœa, after removing all irritating matters by previous evacuation. It is also a good application to ulcers discharging thin ichorous matter. Dose, gr. x. to gr. xl. or more.

PRÆPARATA È CÜPRO.

Preparations of Copper.

CUPRI AMMONIO-SULPHAS.

Ammonio-Sulphate of Copper.

Cuprum Ammoniatum, P.L. 1809, P.L. 1824.

Take of Sulphate of Copper an ounce,

Sesquicarbonate of Ammonia an ounce and a half;

Rub them together until Carbonic Acid ceases to evolve; then dry the Ammonio-Sulphate of Copper, wrapped in bibulous paper, in the air.

Remarks.—Copper forms two different compounds with oxygen; the first is of a red colour, and is a dinoxide composed of one equivalent of oxygen 8, and two equivalents of copper $64 = 72$; this does not yield salts with acids. The oxide is black, and consists of one equivalent of oxygen 8, and one of copper $32 = 40$. It combines with acids to form salts.

Sulphate of copper is a well-known salt, which is of a fine blue colour, crystallizes in right rhombic prisms, and dissolves readily in water. It consists of

One equivalent of Sulphuric Acid ..	= 40	or 34.48
One equivalent of Oxide of Copper ..	= 40	„ 34.48
Four equivalents of Water	= 36	„ 31.04

Equivalent.....	116	100.
-----------------	-----	------

When this salt is decomposed by sesquicarbonate of ammonia, a portion of the carbonic acid of the latter is expelled with effervescence; there appear to be formed carbonate of copper and sulphate of ammonia, which with the excess of the sesquicarbonate of ammonia employed, form the cupri ammonio-sulphas.

Properties.—This compound, when it has not been too much dried, and retains some excess of sesquicarbonate of ammonia, is of a fine azure blue colour and has an ammoniacal smell; its taste is styptic and metallic.

Composition.—This must be liable to some variation, dependent upon its state of dryness and the excess of the ammoniacal carbonate.

Impurities and Tests.—See Notes: CUPRI AMMONIO-SULPHAS.

Incompatibles.—This preparation is incompatible with acids, the alkalis, potash and soda, and with lime-water.

Officinal Preparations.—Liquor Cupri Ammonio-Sulphatis.

Medicinal Uses.—It is tonic and antispasmodic. It has been employed in chorea, and also advantageously in epilepsy. Dose, *one quarter of a grain*, cautiously increased to *five grains*, twice a day. It is given in the form of pills, made up with crumb of bread.

LIQUOR CUPRI AMMONIO-SULPHATIS.

Solution of Ammonio-Sulphate of Copper.

Aqua Sappharina, P.L. 1720, P.L. 1745.

Aqua Cupri Ammoniati, P.L. 1788.

Liquor Cupri Ammoniati, P.L. 1809, P.L. 1824.

Take of Ammonio-Sulphate of Copper a drachm,

Distilled Water a pint;

Dissolve the Ammonio-Sulphate of Copper in the water, and strain.

Properties.—This solution has a fine blue colour, but unless the ammonio-sulphate of copper retains some excess of sesquicarbonate of ammonia, I have found that it is decomposed, and one half of the oxide of copper is precipitated.

Medicinal Uses.—It is detergent and mildly escharotic. When still more largely diluted, it is employed for removing specks from the cornea.

PRÆPARATA È FERRO.

Preparations of Iron.

FERRI SULPHAS.

Sulphate of Iron.

Sal seu Vitriolum Martis, P.L. 1720.

Sal Martis, P.L. 1745.

Ferrum Vitriolatum, P.L. 1788.

Ferri Sulphas, P.L. 1809, P.L. 1824.


Take of Iron Filings eight ounces,

Sulphuric Acid fourteen ounces,

Water four pints;

Mix the Sulphuric Acid with the Water, and add the Iron to them; then apply heat, and when bubbles have ceased to escape, strain the liquor, and set it aside that crystals may be formed. Evaporate the liquor poured off that it may again yield crystals. Dry them all.


Process.—Concentrated sulphuric acid and iron do not act upon each other at common temperatures, but if the acid be diluted with water, rapid action takes place. Water consists of oxygen and hydrogen, and a portion of it is decomposed by the action of the sulphuric acid and iron. Its oxygen combines with the iron to form oxide or protoxide of iron, and its hydrogen being set at liberty, assumes the elastic form and is evolved in the state of gas. The oxide of iron is dissolved by the sulphuric acid, and sulphate of iron is formed.



The solution of sulphate of iron thus obtained is of a bluish green colour, the iron being in the state of protoxide; if it be long exposed to the air, it first loses its blue tint, owing to the absorption of oxygen, which converts the protoxide into sesquioxide or per-oxide of iron, and eventually it becomes of a red colour.

Properties.—The primary form of sulphate of iron is an *oblique rhombic prism*, MM' and P of the annexed figure being the primary planes: the crystals sometimes exhibit the secondary planes a and e .

P on M, or M'	99° 20'
M on M'	82 20
P on e_1	153 00
P on e_2	123 55
P on a_1	159 00
P on a_2	136 10



The crystals, when recently formed, are of a bluish green colour; by exposure to the air the protoxide of iron which the salt contains attracts oxygen, and the yellow colour of the sesquioxide of iron formed renders the crystals green by admixture with the blue protosulphate of iron. When the exposure has been long continued, the surface of the crystals is encrusted with sesquioxide of iron, and they ought then to be rejected; the solution, as already noticed, attracts oxygen, and it is rendered first green and then red, depositing at the same time a considerable quantity of subsesquisulphate of iron.

Sulphate of iron has a disagreeable styptic taste; it is soluble in about two parts of cold water and 3-4ths of its weight of boiling water. The solution is precipitated of a greenish white by alkalis; the oxide thrown down gradually absorbs oxygen, and becomes red, or sesquioxide of iron; when free from this, the ferrocyanide of potassium occasions a white precipitate, which becomes speedily blue by exposure to the air. When the solution has absorbed oxygen by the action of the air, or by any other means, it then gives immediately a deep blue precipitate

with the same test, and a black one with astringent vegetable infusions and tinctures.

By exposure to a moderate heat the crystals lose 6-7ths of their water, their crystalline form, and become white and powdery; subjected to a strong heat they are decomposed, yielding a peculiar kind of sulphuric acid, and sesquioxide of iron.

Composition of the Oxides of Iron.—There are two well-marked oxides of this metal, the protoxide and the peroxide or sesquioxide; and there is an ore of iron, which is either a peculiar oxide, or is a compound of the protoxide and sesquioxide.

The two oxides just mentioned consist of

Protoxide,	One equivalent of Oxygen ..	8	or	22.2
	One equivalent of Iron	28	,,	77.8
		<hr/>		<hr/>
	Equivalent.....	36		100.

Sesquioxide,	One and a half eq. of Oxygen	12	or	30
	One equivalent of Iron.....	28	,,	70
		<hr/>		<hr/>
	Equivalent.....	40		100.

Composition of Sulphate of Iron.

One equivalent of Sulphuric Acid	40	or	28.8
One equivalent of Protoxide of Iron ..	36	,,	25.9
Seven equivalents of Water	63	,,	45.3
	<hr/>		<hr/>
Equivalent.....	139		100.

Symbols.

<i>Protoxide of Iron.</i>	Berzelius and Turner ..	$\text{Fe} + \text{O}$, or $\dot{\text{Fe}}$.
	Brande	$(fe + o)$, or Fe .
<i>Sesquioxide of Iron.</i>	Berzelius and Turner ..	$2\text{Fe} + 3\text{O}$, or $\ddot{\text{Fe}}$.
	Brande	$(fe + 1\frac{1}{2}o)$.
<i>Sulphate of Iron.</i>	Berzelius and Turner ..	$\dot{\text{Fe}}\ddot{\text{S}} + 7\dot{\text{H}}$.
	Brande	$(\text{Fe} + s' + 7q)$.

Impurities and Tests.—See Notes: FERRI SULPHAS.

Incompatibles.—Ammonia, potash, soda and their carbonates, lime-water, and chloride of calcium, nitrate of silver, the acetates of lead, and soaps. The salts of barytes and strontia, as well as the earths they contain, are incompatible with this salt. It is decomposed also by astringent vegetable bodies.

Officinal Preparations.—This salt is employed in preparing

Ferri Sesquioxylum, Mistura Ferri Composita, and Pilulæ Ferri Compositæ.

Medicinal Uses.—Tonic, astringent, emmenagogue, and anthelmintic; in large doses it occasions griping in the bowels. Dose, gr. j. to v. or more, made into pills with extract of gentian. It should never be given in solution without previously boiling the water, to free it from atmospheric air, the oxygen of which is readily absorbed, and the sulphate of iron is decomposed by it.

FERRI SESQUIOXYDUM.

Sesquioxide of Iron.

Chalybis Rubigo præparata, P.L. 1745.

Ferri Rubigo, P.L. 1788.

Ferri Carbonas, P.L. 1809.

Ferri Subcarbonas, P.L. 1824.

Take of Sulphate of Iron four pounds,


Carbonate of Soda four pounds and two ounces,

Boiling Water six gallons;


Dissolve the Sulphate of Iron and Carbonate of Soda separately, in three gallons of water; then the liquors being mixed together, set by, that the powder may subside. Lastly, the supernatant liquor being poured off, wash what is precipitated in water, and dry it.

Process.—The nature of sulphate of iron has been stated; carbonate of soda is the alkaline salt, formerly called subcarbonate of soda.

When the solutions of these salts are mixed, double decomposition takes place. The carbonic acid of the carbonate combines with the oxide of iron of the sulphate, and the carbonate of iron formed being insoluble in water, it is precipitated; the soda of the carbonate unites with the sulphuric acid of the sulphate of iron, and the sulphate of soda resulting being soluble, remains in solution.



The formation of the carbonate or protocarbonate is the first step in the process ; during the washing which is necessary to get rid of the sulphate of soda, the protoxide of iron by exposure to the air acquires oxygen, and loses carbonic acid, and thus becomes sesquioxide of iron.



This compound, however, generally contains a small quantity of carbonic acid ; it is prepared nearly in the same mode as what was called subcarbonate of iron in the last Pharmacopœia, and which usually was, what indeed it is now called, merely sesquioxide of iron.

Properties.—This preparation is of a reddish brown colour ; it is inodorous and has a disagreeable taste ; insoluble in water, and not readily dissolved by any acid except the hydrochloric acid.

Composition and Symbols have been already given.

Impurities and Tests.—See Notes : FERRI SESQUIOXYDUM.

Incompatibles.—Acids and Acidulous Salts.

Official Preparations.—Ferri Ammonio-Chloridum, Tinctura Ferri Sesquichloridi, Ferri Potassio-Tartras, &c.

Medicinal Uses.—Tonic and emmenagogue. Dose, from gr. v. to xxx. combined with myrrh or aromatics. In doses of half a drachm to a drachm, two or three times a day, it is said to have proved efficacious in tic douloureux. In doses of ʒi. to ʒiv. it is stated to have afforded relief in chorea.

TINCTURA FERRI SESQUICHLORIDI.

Tincture of Sesquichloride of Iron.

Tinctura Martis cum Spiritu Salis, P.L. 1720.*Tinctura Martis in Spiritu Salis*, P.L. 1745.*Tinctura Ferri Muriati*, P.L. 1788.*Tinctura Ferri Muriatis*, P.L. 1809, P.L. 1824.


Take of Sesquioxide of Iron six ounces,

Hydrochloric Acid a pint,

Rectified Spirit three pints ;

Pour the Acid upon the Sesquioxide of Iron in a glass vessel, and digest for three days, frequently shaking. Lastly, add the spirit, and strain.

Process.—When sesquioxide of iron is dissolved in hydrochloric acid they undergo mutual decomposition ; the hydrogen of the acid combines with the oxygen of the oxide to form water, and the chlorine of the acid combining with the iron they form sesquichloride of iron.



Sesquichloride of iron is very soluble both in water and in spirit ; it is composed of

One and a half equivalent of Chlorine = 54 or 65·85

One equivalent of Iron = 28 „ 34·15

Equivalent 82 100·

Symbols.—Berzelius and Turner . . $2F + 3Cl$, or $Fe^2 Cl^3$.

Brande $(fe + 1\frac{1}{2} c)$.

Properties.—This tincture is of a reddish brown colour, and though it is essentially composed of sesquichloride of iron, it contains some hydrochloric acid, without which a deposit is apt to be formed in it. Its taste is acid and extremely styptic, and its smell resembles that of hydrochloric æther. Its specific gravity is about 0.992, and a fluidounce yields nearly 30 grains of sesquioxide of iron.

Incompatibles.—Alkalis and their carbonates, lime-water, carbonate of lime, magnesia, and its carbonate. It is rendered black by astringent vegetable bodies, and is decomposed by a solution of gum arabic.

Medicinal Uses.—When made with proper care, it is one of the most certain and active preparations of iron; and it remains for a very long time without suffering any variation of strength from decomposition. Dose ℥ x. to ℥ xxx. or f 3j.

It is stated to be particularly useful as a tonic in scrofula; in dysuria, ℥ x. given every ten minutes until some sensible effect is produced, afford speedy relief; and it is a powerful styptic in hæmorrhage from the bladder, kidneys, or uterus. It is used externally as a styptic in cancerous and fungous sores, and for the purpose of destroying venereal warts.

FERRI POTASSIO-TARTRAS.

Potassio-tartrate of Iron.


Ferrum Tartarizatum, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Sesquioxide of Iron three ounces,
Hydrochloric Acid half a pint,
Solution of Potash four pints and a half, or as
much as may be sufficient,
Bitartrate of Potash eleven ounces and a half,
Solution of Sesquicarbonate of Ammonia a pint,
or as much as may be sufficient,
Distilled Water three gallons;

Mix the Sesquioxide of Iron with the Acid, and digest for two hours in a sand-bath. Add to these two gallons of the water, and set aside for an hour; then pour off the

supernatant liquor. The solution of Potash being added, wash what is precipitated frequently with water, and while moist boil it with the Bitartrate of Potash, previously mixed with a gallon of the water. If the liquor should be acid when tried by litmus, drop into it solution of Sesquicarbonate of Ammonia until it is saturated. Lastly, strain the liquor, and with a gentle heat let it evaporate, so that the salt may remain dry.

Process.—It has been already explained that, when sesquioxide of iron is dissolved by hydrochloric acid, both are decomposed, and the results are water and sesquichloride of iron; this when mixed with solution of potash is decomposed, and a precipitate is obtained, which is hydrated sesquioxide of iron, and chloride of potassium remains in solution.



When this hydrated sesquioxide of iron is boiled in water with the bitartrate of potash, the excess of acid which this salt contains dissolves the oxide, and a solution is obtained, which if not quite neutral to the litmus test, is to be saturated with ammonia; it consists of tartrate of potash and sesquitartrate of iron, and this evaporated to dryness constitutes the potash tartrate of iron.

Properties.—This preparation is of a brownish colour, with a shade of green; it is inodorous, and has but little of the disagreeable taste of the iron, when properly prepared. It is readily soluble in water, and becomes moist in a damp atmosphere. It gives a dark-coloured precipitate with astringent vegetables, but does not afford a blue precipitate with ferrocyanide of potassium; neither potash nor soda, nor their carbonates, decompose this solution unless heat be applied, and even then ammonia and its carbonate produce no effect upon it. Of all chalybeate prepa-

rations it is the least nauseous, and the solution will remain for a considerable time without suffering decomposition.

Composition.—According to Soubeiran, from whom this much-improved process is taken with slight alterations, this double salt contains 13 per cent. of sesquioxide of iron. I found it to consist of very nearly

One equivalent of Tartrate of Potash.... = 114 or 51.82

One eq. of Tartrate of Sesquioxide of Iron = 106 „ 48.18

Equivalent....	220	100.
----------------	-----	------

This would give rather more than 18 per cent. of sesquioxide of iron, which agrees very nearly with my direct experiment.

Impurities and Tests.—See Notes : FERRI POTASSIO-TARTRAS.

Medicinal Uses.—This preparation is advantageously exhibited in all cases in which chalybeates prove useful. From its slight taste it may be readily given when other preparations of iron prove nauseating. The dose is from gr. x. to 3ss. given either in solution, or in the form of bolus, combined with an aromatic, but should not be long kept in either way ; and in its perfect state it cannot be given in the form of powder, on account of its attracting moisture.

FERRI AMMONIO-CHLORIDUM.

Ammonio-Chloride of Iron.

Ens Veneris, P.L. 1720.

Flores Martiales, P.L. 1745.

Ferrum Ammoniacale, P.L. 1788.

Ferrum Ammoniatum, P.L. 1809, P.L. 1824.

Take of Sesquioxide of Iron three ounces,

Hydrochloric Acid half a pint,

Hydrochlorate of Ammonia two pounds and a half,

Distilled Water three pints ;

Mix the Sesquioxide of Iron with the Hydrochloric Acid in a proper vessel, and digest them in a sand-bath

for two hours; afterwards add the Hydrochlorate of Ammonia first dissolved in the distilled Water; strain and evaporate all the liquor. Lastly, rub what remains to powder.

Remarks.—This preparation was ordered in former Pharmacopœias to be sublimed. These directions were however seldom complied with, and the present process will yield a preparation of uniform strength and appearance.

Process.—It has been explained that when sesquioxide of iron is dissolved in hydrochloric acid, the resulting compound is a sesquichloride of iron, which is here mixed with hydrochlorate of ammonia and evaporated to dryness.

Properties.—The colour of this preparation is an orange red; it becomes moist when exposed to the air, is readily soluble in water, and is partially, at least, soluble in alcohol. It has a sharp saline and metallic taste, but no smell.

Composition.—This is a mixture rather than a definite compound, consisting very nearly of

Sesquichloride of Iron	15
Hydrochlorate of Ammonia	85
	<hr/>
	100.

It yields about 7 per cent. of sesquioxide of iron when decomposed by an alkali.

Impurities and Tests.—See Notes: FERRI AMMONIO-CHLORIDUM.

Incompatibles.—This preparation is decomposed by the alkalis and their carbonates, sesquioxide of iron being precipitated, and ammonia evolved; lime-water produces a similar effect; and like other preparations of iron it is rendered black by astringent vegetable infusions.

Officinal Preparations.—Tinctura Ferri Ammonio-Chloridi.

Medicinal Uses.—It is stated to be tonic, emmenagogue, and aperient. Its dose may be estimated by what I have mentioned respecting its composition.

TINCTURA FERRI AMMONIO-CHLORIDI.

Tincture of Ammonio-Chloride of Iron.

Tinctura Martis Mynsichti, P.L. 1720.

Tinctura Florum Martialium, P.L. 1745.

Tinctura Ferri Ammoniacalis, P.L. 1788.

Tinctura Ferri Ammoniati, P.L. 1809, P.L. 1824.

Take of Ammonio-Chloride of Iron four ounces,

Proof Spirit a pint;

Dissolve the Ammonio-Chloride of Iron in the Spirit,
and strain.

Remarks.—This preparation does not appear to possess any advantage over the *Tinctura Ferri Sesquichloridi*, and from which it differs chiefly in containing hydrochlorate of ammonia.

A fluidounce yields by decomposition gr. 5·8 of sesquioxide of iron.

FERRI IODIDUM.

Iodide of Iron.

Take of Iodine six ounces,

Iron Filings two ounces,

Distilled Water four pints and a half;

Mix the Iodine with four pints of the Water, and to these add the Iron. Heat them in a sand-bath, and when it has acquired a greenish colour, pour off the liquor. Wash what remains with the half pint of Water, boiling. Let the mixed and strained liquors evaporate at a heat not exceeding 212° in an iron vessel, that the salt may be dried. Keep it in a well-stopped vessel, access of light being prevented.

Remarks.—Iodine is a non-metallic elementary solid body, which was discovered in 1812 by M. Courtois of Paris. It exists in combination with potassium or sodium in many springs, and has been detected in the water of the Mediterranean, and in marine molluscous animals and sea-weeds. It has also been met with in the mineral kingdom combined with silver.

Properties.—Iodine is a soft crystalline opaque solid, of a bluish black colour and metallic lustre. Its odour is disagreeable, and resembles that of chlorine; its taste is acrid, and it is poisonous. When heated it is vaporized, and its name is derived from the Greek for the violet colour of the vapour. It is very slightly soluble in water, a pint dissolving but little more than a grain of it. It is readily dissolved by alcohol, and the solution is of a reddish brown colour. Its characteristic property is that of giving a blue colour with a solution of starch. It combines readily with metals, and, like chlorine, forms an acid with oxygen and also with hydrogen.

Process and Properties.—The solution obtained is one of protiodide of iron; it is of a green colour, and by evaporation with as little contact of air as possible, green tabular crystals are formed. By evaporation to dryness and heating moderately, this salt is fused, and on cooling becomes an opaque crystalline mass of an iron-grey colour and metallic lustre. When exposed to the air it attracts moisture, and is very soluble both in water and in alcohol: in order to prevent the deposition of sesquioxide of iron by the absorption of oxygen, the solution should be kept with an iron wire in it.

Composition.—Protiodide of iron is composed of

One equivalent of Iodine	126	or	63·3
One equivalent of Iron	28	,,	14·
Five equivalents of Water	45	,,	22·7
	<hr/>		
Equivalent	199		100·

Symbols.—Berzelius and Turner . . Fe + I, or FeI.

Brande (*fe + i*).

Impurities and Tests.—See NOTES: FERRI IODIDUM.

Incompatibles.—This salt is decomposed by the alkalis, ammonia, potash, soda and their carbonates, by lime-water, and all other substances with which sulphate of iron is incompatible.

Medicinal Use.—Used as an emmenagogue, from one to two grains at a dose.

PRÆPARATA EX HYDRARGYRO.

Preparations of Mercury.

HYDRARGYRUM CUM CRETÂ.

Mercury with Chalk.

Mercurius Alkalizatus, P.L. 1745.

Hydrargyrus cum Cretâ, P.L. 1788.

Hydrargyrum cum Cretâ, P.L. 1809, P.L. 1824.

Take of Mercury three ounces,

Prepared Chalk five ounces ;

Rub them together until globules are no longer visible.

Process.—I have been informed on authority upon which I can rely, that the addition of a small quantity of water greatly accelerates the operation here directed. I have found that a small portion of the mercury is, by the long trituration required, converted into binocide, and this being the case the effects derived from the use of this medicine are readily accounted for.

Impurities and Tests.—See Notes : HYDRARGYRUM CUM CRETÂ.

Incompatibles.—Acids and acidulous salts act upon this preparation, and dissolve the chalk with the effervescence of carbonic acid gas.

Medicinal Uses.—It is one of the mildest of the mercurial preparations. Dose, as an alterative, gr. x. to gr. xxx.

HYDRARGYRI BICHLORIDUM.

Bichloride of Mercury.

Mercurius Corrosivus Sublimatus, P.L. 1720, P.L. 1745.

Hydrargyrus Muriatus, P.L. 1788.

Hydrargyri Oxymurias, P.L. 1809, P.L. 1824.


Take of Mercury two pounds,

Sulphuric Acid three pounds,


Chloride of Sodium a pound and a half;

Boil the Mercury with the Sulphuric Acid in a proper vessel, until the Bipersulphate of Mercury remains dry; rub this when it is cold with the Chloride of Sodium in an earthen mortar; then sublime with a heat gradually raised.


Process.—Supposing the sulphuric acid to be of the greatest density, and the excess of it to be merely evaporated, the changes which occur during the formation of Hydrargyri Bichloridum, commonly called corrosive sublimate, are as follows:—
4 eqs. of liquid sulphuric acid = 196, consist of 36 = 4 eqs. of water, and 160 = 4 eqs. of dry sulphuric acid; during ebullition the 36 of water are evaporated; and 80 = 2 eqs. of dry sulphuric acid are decomposed into 64 = 2 eqs. of sulphurous acid gas, which are evolved, and 16 = 2 eqs. of oxygen, which combine with 202 = 1 eq. of mercury and form 218 = 1 eq. of binoxide of mercury, which uniting with 80 = 2 eqs. of sulphuric acid, remaining undecomposed, there are formed 298 = 1 eq. of bipersulphate of mercury.



Chloride of sodium, or common salt, is then mixed and heated in a subliming vessel with these $298 = 1$ eq. of bipersulphate of mercury, consisting, as already stated, of 1 eq. of mercury $= 202$ 2 eqs. of oxygen $= 16$ and 2 eqs. of sulphuric acid $= 80$; the $120 = 2$ eqs. of chloride of sodium, are composed of 2 eqs. of chlorine $= 72$ and 2 eqs. of sodium $= 48$; during the action of these substances upon each other, $202 = 1$ eq. of mercury, combine with 2 eqs. of chlorine $= 72$, and form $274 = 1$ eq. of bichloride of mercury; $16 = 2$ eqs. of oxygen separated from the mercury, are transferred to the 2 eqs. of sodium $= 48$ and form 2 eqs. of soda $= 64$, which combining with $80 = 2$ eqs. of sulphuric acid, give $144 = 2$ eqs. of sulphate of soda, remaining in the lower part of the subliming vessel.



Properties.—The bichloride of mercury being volatile at the temperature at which it is formed, rises in vapour, and condenses into a white semi-transparent crystalline mass, and perfect crystals are occasionally procurable. The cleavages in the crystals of this substance are parallel to the lateral and to the terminal planes of a *right rhombic prism* of $93^{\circ} 44'$, which therefore may be regarded as the primary form.



P on M, or M'	90° 00'
M on M'	93 44
M on h	133 8

Bichloride of mercury is inodorous; it has an acrid and nauseous taste, which remains long in the mouth. It is a violent poison. Its specific gravity is 5.200 ; water, at 60° Fahr. dissolves rather more than 1-20th, and boiling water one-third of its weight. Although light has no action upon this salt in its solid state, yet it partially decomposes the aqueous solution, and chloride of mercury is precipitated. It is much more soluble in

alcohol, ether, hydrochloric acid, and solution of hydrochlorate of ammonia, than in water. When the alkalis potash and soda, or lime-water are added to a solution of bichloride of mercury, they throw down a reddish yellow precipitate, which is hydrated binoxide of mercury. With ammonia a white precipitate is obtained, as will be presently again noticed.

Composition.—Bichloride of mercury consists of

Two equivalents of Chlorine.. $36 \times 2 = 72$ or 26.27

One equivalent of Mercury .. $= 202$ „ 73.73

Equivalent 274 100.

Symbols,—Berzelius and Turner.. $\text{Hg} + 2 \text{Cl}$, or Hg Cl^2 .

Brande ($hg + 2c$).

Impurities and Tests.—See Notes: HYDRARGYRI BICHLORIDUM.

Incompatibles.—Ammonia, potash, soda, and their carbonates; lime-water, potassio-tartrate of antimony, nitrate of silver, the acetates of lead, sulphuret of potassium, hydrosulphates, soap, many metals, infusion of bitter and astringent vegetables, and some vegetable bodies which possess neither of these qualities.

Officinal Preparations.—Liquor Hydrargyri Bichloridi, Hydrargyri Binoxidum, Hydrargyri Ammonio-Chloridum.

Medicinal Uses.—It is frequently serviceable in secondary syphilis, and in some cutaneous diseases, particularly combined with an antimonial, in lepra. Dose from one eighth to one fourth of a grain, made into a pill with crumb of bread.

LIQUOR HYDRARGYRI BICHLORIDI.

Solution of Bichloride of Mercury.

Liquor Hydrargyri Oxymuriatis, P.L. 1809, P.L. 1824.

Take of Bichloride of Mercury,

Hydrochlorate of Ammonia, each ten grains,

Distilled Water a pint;

Dissolve the Bichloride of Mercury and Hydrochlorate of Ammonia together in the Water.

Remarks.—In the former Pharmacopœias the solvent power of the water was increased by spirit, instead of hydrochlorate of ammonia as now directed. A fluidounce contains half a grain of bichloride of mercury.

Dose, half a fluidrachm to two fluidrachms in infusion of linseed.

HYDRARGYRI CHLORIDUM.

Chloride of Mercury.

Calomelas, P.L. 1720.

Mercurius Dulcis Sublimatus, P.L. 1745.

Calomelas Hydrargyrus Muriatis Mitis, P.L. 1788.

Hydrargyri Submuriatis, P.L. 1809, P.L. 1824.

Take of Mercury four pounds,


Sulphuric Acid three pounds,

Chloride of Sodium a pound and a half,

Distilled Water as much as may be sufficient;


Boil two pounds of the Mercury with the Sulphuric Acid in a proper vessel, until the Bipersulphate of Mercury remains dry; rub this when it is cold with two pounds of Mercury in an earthen mortar, that they may be perfectly mixed. Then add the Chloride of Sodium, and rub them together, until globules are no longer visible; then sublime. Rub the sublimate to very fine powder, and wash it carefully with boiling distilled Water and dry it.

Process.—It has been already mentioned, that when mercury and sulphuric acid are boiled together, the metal is converted into bipersulphate, which when rubbed as directed with a quantity of mercury equal to that which it already contains, we may consider as forming with it protosulphate of mercury, or sulphate of the protoxide; for the first portion of mercury yields half its oxygen to the second portion, and both become protoxide, and combine with the sulphuric acid.




500 = 2 eqs. *Protosulphate of Mercury.*

Now when this protosulphate is mixed and heated with common salt, the changes which occur are, that 500, the 2 eqs. of protosulphate of mercury, composed of $404=2$ eqs. of mercury, $16=2$ eqs. oxygen and $80=2$ eqs. sulphuric acid, decompose $120=2$ eqs. chloride of sodium, consisting of $72=2$ eqs. chlorine and $48=2$ eqs. sodium; the 2 eqs. of mercury= 404 combine with the 2 eqs. of chlorine= 72 , and form $476=2$ eqs. of chloride of mercury; 16 the 2 eqs. of oxygen combine with 48 the 2 eqs. of sodium and form $64=2$ eqs. of soda, which uniting with 80 the 2 eqs. of sulphuric acid, there result $144=2$ eqs. of sulphate of soda.



Properties.—Chloride, or as it is called when a pointed distinction is necessary, *protochloride* of mercury is commonly called calomel. It is a white semitransparent crystalline mass, inodorous, insipid, and insoluble in water. Its specific gravity is 7.175 : by long exposure to light it is rendered of a dark colour, owing to partial decomposition. Occasionally perfect crystals are obtained, in which, although there does not appear to be any distinct cleavage, there are indications of it parallel to all the planes of a *square prism*, and this may be regarded as the primary form.



P on M, or M'	90°	00'
P on a	112	5
P on c	119	50
M on M'	90	00
M on c	150	10

Composition.—Chloride of mercury is composed of

One equivalent of Chlorine.....	=	36	or	16
One equivalent of Mercury.....	=	202	„	84
		<hr/>		<hr/>
Equivalent.....		238		100.

Symbols.—Berzelius and Turner.. Hg + Cl, or Hg Cl.

Brande (hg + c).

Impurities and Tests.—See Notes: HYDRARGYRI CHLORIDUM.

Incompatibles.—Chloride of mercury is immediately decomposed by ammonia, potash, soda, and lime, protoxide of mercury being precipitated; carbonate of ammonia produces the same effect, but the carbonates of potash and soda less rapidly; bicarbonate of potash does not decompose it at all. By nitric acid it is partially converted into bichloride. It is decomposed by iron, copper, and lead, and also by hydrosulphuric acid and its salts.

Officinal Preparations.—Hydrargyri Oxydum, Pilulæ Hydrargyri Chloridi compositæ.

Medicinal Uses.—It is an extremely efficient purgative, and it is alterative, antisyphilitic, and a valuable remedy in obstructions and hepatic affections. It is particularly useful in the diseases of children, and they frequently bear larger doses of it than adults. Dose as an alterative gr. ss. to gr. j. night and morning; as a purgative from gr. ij. to gr. x. or in some cases considerably more. Its insolubility and great specific gravity prevent its being eligibly exhibited in any other form than that of powder or pill.

HYDRARGYRI AMMONIO-CHLORIDUM.

Ammonio-Chloride of Mercury.

Mercurius Præcipitatus Albus, P.L. 1745.

Calx Hydrargyri Alba, P.L. 1788.

Hydrargyrus Præcipitatus Albus, P.L. 1809.

Hydrargyrum Præcipitatum Album, P.L. 1824.

Take of Bichloride of Mercury six ounces,
 Distilled Water six pints,
 Solution of Ammonia, eight fluidounces;

Dissolve the Bichloride of Mercury, with the application of heat, in the Water. To this when it is cold add the Solution of Ammonia, frequently stirring; wash the powder thrown down until it is free from taste; lastly, dry it.

Remarks.—In former Pharmacopœias this preparation was obtained by the addition of solution of carbonate of potash and hydrochlorate of ammonia to that of the bichloride of mercury; the more direct method of employing ammonia is now substituted.


Process.—When ammonia is added to bichloride of mercury, it appears, from the experiments of Dr. Kane, that half the chlorine is removed, and the whole of the mercury thrown down, combined consequently with only so much chlorine as reduces it to the state of protochloride, and this is combined with ammonia.

Composition.—Various statements have been made on this subject. Mr. Hennell considers it as a compound of 1 equivalent of hydrochlorate of ammonia and 1 of binoxide of mercury; Dr. Kane, as constituted of binamide and bichloride of mercury; he admits, however, that it may contain oxygen; and adopting this latter opinion, I am disposed to consider this preparation as containing

One equivalent of Binoxide of Mercury .	= 218
One equivalent of Bichloride of Mercury	= 274
Two equivalents of Ammonia	= 34

526

If this be correct the changes which occur are the following, the figures representing the *number* and not the *weight* of the equivalents: When 2 of bichloride of mercury are dissolved in water and ammonia is added to the solution, 2 of water are decomposed, the 2 of hydrogen of which unite with the 2 of chlorine of one of the equivalents of bichloride of mercury, and form 2 of hydrochloric acid, and these combining with 2 of ammonia give 2 of hydrochlorate of ammonia, which are poured off in the supernatant liquor. The 2 of oxygen of the 2 of decomposed water unite with the 1 of mercury, separated from the 2 of chlorine, and form 1 of binoxide of mercury, which is precipitated with the 1 of bichloride of mercury undecomposed and 2 of ammonia, forming the Hydrargyri Ammonio-Chloridum.



Hydrargyri Ammonio-Chloridum, P.L.

Properties.—This is a light and perfectly white powder. It is inodorous, insipid, and insoluble in water, but dissolved by sulphuric, nitric, and hydrochloric acid. When heated in solution of potash it suffers partial decomposition, becomes yellow, and yields ammonia.

Impurities and Tests.—See Notes: HYDRARGYRI AMMONIO-CHLORIDUM.

Medicinal Use.—It is employed only externally in cutaneous affections.

HYDRARGYRI OXYDUM.

Oxide of Mercury.

Hydrargyri Oxydum Cinereum, P.L. 1809, P.L. 1824.


Take of Chloride of Mercury an ounce,

Lime Water a gallon;

Mix, and frequently shake them. Set by, and when the oxide has subsided, pour off the liquor; lastly, wash it in distilled Water until nothing alkaline can be perceived, and dry it in the air wrapped in bibulous paper.

Process.—When chloride of mercury is acted upon by lime,

both suffer decomposition. The chlorine of the chloride unites with the calcium of the lime, and chloride of calcium is formed, which remains dissolved in the water, and is eventually poured away with it. The oxygen of the lime combining with the mercury forms oxide of mercury, which is precipitated.



Properties.—This preparation is nearly black, insoluble in water and the alkalis, but dissolves readily in nitric acid; it decomposes and is decomposed by hydrochloric acid, which reconverts it to chloride of mercury, with the formation of water. It is totally volatilized by heat.

Composition.—It is composed of

One equivalent of Oxygen	8 or 3·8
One equivalent of Mercury	202 „ 96·2
	210 100·
Equivalent	

Symbols.—Berzelius and Turner .. $\text{Hg} + \text{O}$, or Hg .

Brande ($hg + o$).

Impurities and Tests.—See Notes: HYDRARGYRI OXYDUM.

Incompatibles.—Acids. Acidulous Salts. Hydrosulphuric Acid and Hydrosulphates.

Medicinal Use.—Alterative. Dose, gr. j. to gr. iij. in the form of pill twice a day.

HYDRARGYRI BINOXYDUM.

Binoxide of Mercury.


Mercurius Calcinatus, P.L. 1745.*Hydrargyrus Calcinatus*, P.L. 1788.*Hydrargyri Oxydum rubrum*, P.L. 1809, P.L. 1824.

Take of Bichloride of Mercury four ounces,
 Solution of Potash twenty-eight fluidounces,
 Distilled Water six pints ;

Dissolve the Bichloride in the Water ; strain and add the solution of Potash. Wash the powder thrown down in distilled Water, until nothing alkaline can be perceived, and dry it with a gentle heat.

Remarks.—In former Pharmacopœias, binoxide of mercury was prepared by the slow operation of heat and air, and though similar in composition was very different in appearance from that obtained by the present process.

Process.—When bichloride of mercury is mixed with solution of potash, both are decomposed ; the 2 of chlorine which the bichloride contains take 2 of potassium from the potash, and 2 of chloride of potassium are formed and remain dissolved ; the 2 of oxygen separated from the potassium combine with the 1 of mercury and are precipitated together as binoxide of mercury.



Properties.—Binoxide of mercury as above prepared is an orange-red powder ; it is inodorous, acrid to the taste, and insoluble in water. At a heat below redness, it is decomposed, yielding oxygen

gas, and the mercury returns to the metallic state. It is readily dissolved by the nitric, hydrochloric, and some other acids.

Composition.—Binoxide of mercury is composed of

Two equivalents of Oxygen	=	16	or	7.3
One equivalent of Mercury	=	202	„	92.7
			<hr/>		<hr/>
Equivalent....	218		100.		

Symbols,—Berzelius and Turner.. $\text{Hg} + 2\text{O}$ or $\ddot{\text{Hg}}$.

Brande ($hg + 2o$).

Impurities and Tests.—See Notes: HYDRARGYRI BINOXYDUM.

Incompatibles.—Acids and acidulous salts. Hydrosulphuric acid and hydrosulphates.

Medicinal Uses.—It is a very active medicine; but as it frequently occasions vomiting, purging, and sometimes affects the stomach and bowels violently, it is now but little employed. Dose, gr. j. combined with gr. ss. of opium.

HYDRARGYRI NITRICO-OXYDUM.

Nitric Oxide of Mercury.

Mercurius Præcipitatus Corrosivus, P.L. 1720.

Mercurius Corrosivus Ruber, P.L. 1745.

Hydrargyrus Nitratus Ruber, P.L. 1788.

Hydrargyri Nitrico-Oxydum, P.L. 1809, P.L. 1824.

Take of Mercury three pounds,
Nitric Acid a pound and a half,
Distilled Water two pints;

Mix them in a proper vessel and apply a gentle heat until the Mercury is dissolved. Boil down the liquor, and rub what remains to powder. Put this into another very shallow vessel; then apply a slow fire, and gradually increase it until red vapour ceases to arise.

Process.—The mercury decomposes a part of the nitric acid, and, combining with a portion of its oxygen, is converted first

into oxide and then into nitrate of mercury ; this nitrate of mercury when exposed to a strong heat is decomposed, and so also is the acid which it contains. The red vapour above-mentioned results from the union of the nitric oxide evolved, with the oxygen of the atmosphere, and the consequent formation of nitrous acid gas. (See Argenti Nitras.)

Properties.—This preparation is of a bright red colour, and has a crystalline appearance.

Composition.—Like the last, this preparation is binoxide of mercury : it sometimes contains a little undecomposed nitrate, and has on this account been called, but improperly, Subnitrate of Mercury. Excepting a small and accidental portion of undecomposed nitrate, it consists of the same quantities of oxygen as the last preparation, and its composition, symbols, and incompatibles are similar.

Impurities and Tests.—See Notes : HYDRARGYRI NITRICO-OXYDUM.

Officinal Preparation. — Unguentum Hydrargyri Nitrico-Oxydum.

Medicinal Uses.—It is employed only externally as an escharotic.

HYDRARGYRI BICYANIDUM.

Bicyanide of Mercury.

Take of Percyanide of Iron eight ounces,
Binoxide of Mercury ten ounces,
Distilled Water four pints ;

Boil them together for half an hour and strain. Evaporate the liquor that crystals may be formed. Wash what remains frequently with boiling distilled Water, and again evaporate the mixed liquors that crystals may be formed.

Bicyanide of Mercury may be otherwise prepared by adding as much Binoxide of Mercury as will accurately saturate it, to Hydrocyanic Acid distilled from Ferrocyanide of Potassium with diluted Sulphuric Acid.

Remarks.—Percyanide of Iron, usually called Prussian Blue,


and sometimes Ferrosesquicyanide of Iron, is regarded as a compound of

Nine equivalents of Cyanogen $26 \times 9 = 234$ or 54.4
 Seven equivalents of Iron $28 \times 7 = 196$ „ 45.6


Equivalent 430 100.

It is prepared on the large scale as a pigment by adding the ferrocyanide of potassium already mentioned (see Hydrocyanic Acid) to a solution of sulphate of iron, and it is the result of a very complicated play of affinities, in which the oxygen of the air acts an important part.

Process.—When percyanide of iron and binoxide of mercury are boiled together in water, they act upon each other, though neither of them is soluble in it. The reactions which occur in preparing this compound are, that the cyanogen quits the iron to combine with the mercury, forming bicyanide of mercury, which is dissolved; on the other hand, the iron left by the cyanogen takes the oxygen which the mercury has quitted, and forms with it sesquioxide of iron, which remains insoluble. As the iron is only a sesquioxide, while the mercury is a binoxide, the decomposition is not quite complete.



In the second method mentioned for preparing bicyanide of mercury, the changes are between hydrocyanic acid and binoxide of mercury; two equivalents of the acid decompose one equivalent of the oxide, and the results are bicyanide of mercury and water.



Properties.—The solution of bichyanide of mercury is colourless, and by evaporation yields colourless crystals, the form of which is a right square prism, with numerous modifying planes; this salt has a metallic taste, is poisonous, much more soluble in hot water than in cold, and but little soluble in alcohol. It is decomposed by heat, the results being cyanogen and mercury. It is dissolved by nitric acid without decomposition, but it is decomposed by sulphuric acid, and by hydrochloric acid, which evolves hydrocyanic acid, and bichloride of mercury is formed; the affinity existing between cyanogen and mercury is so strong that the alkalis do not decompose the aqueous solution; but hydrosulphuric acid and the hydrosulphates readily produce this effect.

Composition.—Bichyanide of Mercury consists of

Two equivalents of Cyanogen.....	$26 \times 2 =$	52	or	20.4
One equivalent of Mercury		=202	„	79.6
		<hr/>		<hr/>
Equivalent.....		254		100.

Symbols.—Berzelius and Turner.. Hg + 2Cy, or Hg Cy.

Brande (hg + 2cy).

Impurities and Tests.—See Notes : HYDRARGYRI BICYANIDUM.

Incompatibles.—Sulphuric Acid, Hydrochloric Acid, Hydrosulphuric Acid and Hydrosulphates.

Official Preparation.—Acidum Hydrocyanicum.

HYDRARGYRI IODIDUM.

Iodide of Mercury.

Take of Mercury an ounce,

Iodine five drachms,

Alcohol as much as may be sufficient;

Rub the Mercury and Iodine together, adding the alcohol gradually, until globules are no longer visible. Dry the powder immediately with a gentle heat without the access of light, and keep it in a well-stopped vessel.

Properties.—This compound is a greenish yellow powder, it is

insoluble in water. It should not be exposed to light, as by its action and that of heat it is apt to be resolved into mercury and biniodide; when quickly heated, however, it sublimes nearly or quite unaltered.

Composition.—Iodide of mercury is composed of

One equivalent of Iodine	126	or	38.4
One equivalent of Mercury	202	,,	61.6
	<hr/>		
Equivalent	328		100.

Symbols,—Berzelius and Turner.. Hg + I, or HgI.

Brande (*hg* + *i*).

Impurities and Tests.—See Notes: HYDRARGYRI IODIDUM.

Medicinal Uses.—Has been given internally in scrofulous habits, from gr. j. to gr. iij.; but it is chiefly employed in the form of ointment.

HYDRARGYRI BINIODIDUM.

Biniodide of Mercury.

Take of Mercury an ounce,
Iodine ten drachms,
Alcohol as much as may be sufficient;

Rub the Mercury and Iodine together, adding the Alcohol gradually, until globules are no longer visible. Dry the powder with a gentle heat, and keep it in a well-stopped vessel.

Properties.—The biniodide of mercury is a red colour approaching to scarlet; it fuses readily, and sublimes in rhombic scales, which are yellow at first but become red on cooling. It is not dissolved by water. It is soluble in some acids, and in alcohol also when heated.

Composition.—The salt consists of

Two equivalents of Iodine ..	$126 \times 2 = 252$	or	55.5
One equivalent of Mercury	$= 202$	„	44.5
	<hr/>		
Equivalent	454		100.

Symbols.—Berzelius and Turner.. $\text{Hg} + 2\text{I}$, or HgI^2 .

Brande

Impurities and Tests.—See Notes: HYDRARGYRI BINIODIDUM.

Medicinal Uses.—Like the preceding, it has been tried in scrofulous and syphilitic affections in doses of gr. ss. to j. daily; likewise as an ointment.

HYDRARGYRI BISULPHURETUM.

Bisulphuret of Mercury.

Cinnabaris Factitia, P.L. 1745.

Hydrargyrus Sulphuratus Ruber, P.L. 1788.

Hydrargyri Sulphuretum Rubrum, P.L. 1809, P.L. 1824.

Take of Mercury two pounds,

Sulphur five ounces;

Mix the Mercury with the Sulphur melted over the fire, and, as soon as the mass swells, remove the vessel from the fire, and cover it strongly lest inflammation should occur; then rub to powder and sublime.

Process.—By the action of heat in the first instance, combination takes place between the mercury and a portion of the sulphur; by continuing it, the excess of the latter appears to be expelled, and by sublimation, the red, per- or bi-sulphuret of mercury is obtained.

Properties.—When in mass, this substance is of a dark colour, which, when reduced to a fine powder, is of a brilliant red, and it is often called cinnabar or vermilion. It is inodorous and insipid; unalterable by exposure to the air or moisture. When heated to redness in an open vessel, the sulphur is converted into

sulphurous acid, and the mercury escapes in vapour. It is decomposed when distilled with lime, potash, or soda, and also by several of the metals.

When it is heated with sulphuric acid, sulphurous acid is evolved and a sulphate of mercury is formed. It is insoluble in nitric or hydrochloric acid, but when mixed, the chlorine evolved acts upon and dissolves it, without the assistance of heat.

Composition.—This sulphuret consists of

Two equivalents of Sulphur	$16 \times 2 = 32$	or	13.6
One equivalent of Mercury	$= 202$	„	86.4
	<hr/>	<hr/>	
	234		100.

Symbols,—Berzelius and Turner. . $\text{Hg} + 2\text{S}$, or HgS^2 .

Brande ($hg + 2s$).

Impurities and Tests.—See Notes: HYDRARGYRI BISULPHURETUM.

Medicinal Uses.—It is employed for the purpose of mercurial fumigations by heating ʒss. of it on a red hot iron.

HYDRARGYRI SULPHURETUM CUM SULPHURE.

Sulphuret of Mercury with Sulphur.

Æthiops Mineralis, P.L. 1745.

Hydrargyrus cum Sulphure, P.L. 1788.

Hydrargyri Sulphuretum Nigrum, P.L. 1824.

Take of Mercury,

Sulphur, each a pound ;

Rub them together, until globules are no longer visible.

Process.—The mercury combines with a portion of the sulphur by mere trituration ; according to Mr. Brande, (Manual of Pharmacy, p. 303,) when Hydrargyri Sulphuretum cum Sulphure is boiled in a solution of potash, the excess of sulphur is removed, and a black insoluble powder remains, which when washed and

dried is not acted upon by nitric acid, sublimes at a red heat without decomposition, and assumes the characters of the Hydrargyri Bisulphuretum.

Properties.—This preparation, well known by the name of Æthiops mineral, is a very black, insipid, and inodorous powder.

Composition.—It follows from what has been stated, that Hydrargyri Sulphuretum cum Sulphure is a mixture of

Bisulphuret of Mercury	58
Sulphur	42
	<hr/>
	100

Medicinal Uses.—It is an inefficient preparation. Dose, from gr. v. to gr. xxx. as an alterative.

PRÆPARATA EX MAGNESIO.

Preparations of Magnesium.

MAGNESIA.

Magnesia.

Magnesia Usta, P.L. 1788.

Magnesia, P.L. 1809, P.L. 1824.

Take of Carbonate of Magnesia four ounces;

Burn it for two hours in a very strong fire.

Remarks.—*Magnesium* is a peculiar metal, of which the alkaline earthy substance magnesia, is the only well-known oxide, consisting of

One equivalent of Oxygen..... = 8 or 40

One equivalent of Magnesium..... = 12 „ 60

	<hr/>	<hr/>
Equivalent	20	100.

Process.—The carbonate of magnesia, like the carbonate of lime, parts with its carbonic acid at a high temperature, and the magnesia remains in its pure or caustic state.

Properties.—Colourless, inodorous, and tasteless if pure; it does not, like lime, become hot when mixed with water; it is very nearly insoluble in water, and although the moistened earth exhibits alkaline properties by turning vegetable blues green, and yellows brown, yet water in which it has been agitated does not dissolve enough to produce this effect, as lime-water readily does. By exposure to the air it slowly attracts carbonic acid and is reconverted to carbonate.

Symbols.—Berzelius and Turner $Mg + O$, or $\dot{M}g$.

Brande (*mag + o*).

Impurities and Tests.—See Notes : MAGNESIA.

Incompatibles.—Acids, Acidulous Salts, Metallic Salts, and Hydrochlorate of Ammonia.

Medicinal Uses.—Antacid, and when acidity prevails, purgative; it is preferable to the carbonate whenever the bowels are distended with flatus; in other respects its virtues are the same. Dose, ʒss. to ʒj.

MAGNESIÆ CARBONAS.

Carbonate of Magnesia.

Magnesia Alba, P.L. 1788.

Magnesiæ Carbonas, P.L. 1809.

Magnesiæ Subcarbonas, P.L. 1824.

Take of Sulphate of Magnesia four pounds,
Carbonate of Soda four pounds and eight
ounces,

Distilled Water four gallons;

Dissolve separately the Carbonate of Soda and Sulphate of Magnesia in two gallons of the Water, and strain; then mix the liquors and boil, stirring constantly with a spatula for a quarter of an hour; lastly, the liquor being poured off, wash the precipitated powder with boiling distilled Water, and dry it.

Remarks.—Although sulphate of magnesia is an article of the

Materia Medica, I shall take this opportunity of stating its qualities, crystalline form, and composition. It was originally called Epsom Salt, having been procured from a spring at that place.

Sulphate of magnesia is one of the saline ingredients of sea water, and for a long time it was procured only from the bittern remaining after the preparation of common salt; thus obtained it was usually mixed with so considerable a quantity of chloride of magnesium, that owing to the deliquescent property of this salt, the sulphate was usually damp. It has since been much better prepared from magnesian limestone, by a very ingenious process, invented by the late Dr. Henry, and the salt so formed being unmixed with chloride of magnesium, does not attract moisture from the air.

Sulphate of magnesia crystallizes with great readiness, and although the crystals are usually small, they may be obtained of considerable size by slowly cooling the solution. The primary form of this substance may be regarded as a *right prism* with a *rhombic base*, whose angles are $90^{\circ} 30'$ and $89^{\circ} 30'$.

There is only one cleavage, which is parallel to the short diagonal of the prism, and consequently to the plane *h* of the accompanying figures.

Fig. 1. represents a crystal of a form which frequently occurs, and of which the following are the measurements :

M on M' (primary) ..	$90^{\circ} 30'$
M on <i>h</i>	134 45
M on <i>e</i>	129 00
<i>a</i> on <i>a'</i>	120 nearly

Fig. 1.





Fig. 2. represents a form in which the crystals also frequently appear. In this form only two of the four planes *e* are seen on each summit, and alternating in position as shown in the figure.

On some of the crystals, however, which resemble this figure, the two other planes *e* may be perceived, but they are very minute.

Fig. 2.



Sulphate of magnesia is an extremely bitter salt; it is readily soluble in cold water, and still more so in hot water, the former dissolving an equal weight, and the latter one third more; it is

unalterable by exposure to the air, but when heated it loses its water of crystallization.

Composition.—Sulphate of magnesia is composed of

One equivalent of Sulphuric Acid = 40 or 32.52

One equivalent of Magnesia = 20 „ 16.26

Seven equivalents of Water $9 \times 7 = 63$ „ 51.22

Equivalent	123	100.
--------------------	-----	------

Symbols.—Berzelius and Turner .. $\text{Mg} \ddot{\text{S}} + 7\dot{\text{H}}$.


Brande $(\text{M} + \text{S}' + 7q)$.

Incompatibles.—Sulphate of magnesia is incompatible with the alkalis potash and soda, and their carbonates, but the bicarbonates do not decompose it until part of their carbonic acid is expelled by heat. Ammonia decomposes it but partially, and the sesquicarbonate not at all. Lime-water and chloride of calcium are both incompatible with this salt, and so also are the acetates of lead.

Medicinal Use.—Sulphate of magnesia is extensively employed as a purgative. Dose, from ʒss. to ʒjss.

The process of preparing carbonate of magnesia, from the sulphate, is one in which double decomposition takes place; the carbonic acid of the carbonate combines with the magnesia of the sulphate, and the carbonate of magnesia formed being insoluble in water, it is precipitated; the soda of the carbonate unites with the sulphuric acid of the sulphate of magnesia, and the resulting sulphate of soda remains in solution.

Sulphate of Soda.



Carbonate of Magnesia.

Properties.—Carbonate of magnesia when pure is colourless, inodorous, tasteless, and unalterable in the air; it is insoluble in water; and is decomposed by a strong heat, which expels the carbonic acid.

Composition.—Carbonate of magnesia is composed of

One equivalent of Carbonic Acid = 22 or 52.4

One equivalent of Magnesia = 20 „ 47.6

Equivalent	42	100.
--------------------	----	------

It usually contains some water.

Incompatibles.—Acids and acidulous and metallic salts, hydrochlorate of ammonia, and lime-water.

Medicinal Uses.—Antacid and purgative, and in lithic calculi in doses of ℥j. to ʒj.

PRÆPARATA È PLUMBO.

Preparations of Lead.

PLUMBI ACETAS.

Acetate of Lead.

Saccharum Saturni, P.L. 1720, P.L. 1745.

Cerussa Acetata, P.L. 1788.

Plumbi Superacetas, P.L. 1809.

Plumbi Acetas, P.L. 1824.

Take of Oxide of Lead, rubbed to powder, four pounds
and two ounces,

Acetic Acid,

Distilled Water, each four pints ;

Mix the Acid with the Water, and add the Oxide of Lead to them, and a gentle heat being applied dissolve it : then strain. Lastly, evaporate the liquor that crystals may be formed.

Remarks.—Lead is a well-known, soft, bluish white metal of sp. gr. 11.381. It may be made to combine with four different preparations of oxygen :

1st, Dark gray Din oxide, composed of

1 equivalent oxygen 8 + 2 equivalents lead 208 = 216

2nd, Yellow Oxide, or Protoxide, composed of

1 equivalent oxygen 8 + 1 equivalent lead 104 = 112

3rd, Red Deutoxide, composed of

4 equivalents oxygen 32 + 3 equivalents lead 312 = 344


4th, Brown Binoxide, composed of

2 equivalents oxygen 16 + 1 equivalent lead 104 = 120

Of these the protoxide only combines with acids to form salts; when prepared by heating lead exposed to the air in the process of making red lead, it has a pale yellow colour, and is called in commerce *massicot*; while *litharge*, which is the oxide directed by the College, is obtained during the separation of silver from lead ore, and having undergone partial fusion and crystallization, it has a different appearance, though its composition is the same.

Process.—This is a case of single affinity merely; the solution is colourless, and by evaporation yields crystals of acetate of lead.

Properties.—Acetate of lead is crystalline, colourless, nearly inodorous, of a sweetish astringent taste, and is poisonous; it suffers but little change by exposure to the air. The crystals are usually very small; but if they are suffered to form slowly, they may be obtained of considerable size. Their primary form appears to be a *right oblique-angled prism*; the only modification which it has been as yet observed to present, is exhibited in the annexed figure:—



d on d'	128°	0'
d on M	116	0
d on T	98	30
M on T	109	32

Water at 60° dissolves about one fourth of its weight of this salt, and it is not much more soluble in boiling water. When the solution is exposed to the air, the acetate is partly decomposed by the absorption of carbonic acid, and carbonate of lead is precipitated: water which contains carbonic acid also decomposes acetate of lead to a certain extent; and if a current of carbonic acid gas be passed through the solution, one half of the acetate is converted into carbonate and precipitated, and binacetate of lead remains in solution.

Composition.—Acetate of lead is composed of

One equivalent of Acetic Acid	=	51	or	26.8
One equivalent of Oxide of Lead	=	112	„	58.9
Three equivalents of Water .. 9×3 =		27	„	14.3
		<hr/>		<hr/>
Equivalent		190		100.

Symbols.—Berzelius and Turner, $\text{Pb} + \bar{\text{A}} + 3\text{H}$.
 Brande, $(\text{Pl} + ac' + 3q)$.

Impurities and Tests.—See Notes: PLUMBI ACETAS.

Incompatibles.—It is decomposed by all those acids and their compounds which form, with oxide of lead, salts nearly insoluble in water, as the sulphuric, hydrochloric, carbonic, citric, and tartaric acids. It is decomposed by lime-water; by the alkalis potash and soda; but, if added in excess, they redissolve the precipitate at first formed. Hard water usually contains three ingredients which decompose it, viz. carbonate of lime, sulphate of lime, and common salt; and hence, when dissolved in such water, the solution is always turbid. It is decomposed by hydrosulphuric acid and its salts, which give a black sulphuret: *Liquor Ammoniae Acetatis* also decomposes it, on account of the carbonic acid diffused through it.

Officinal Preparations.—*Ceratum Plumbi Acetatis*, *Liquor Plumbi Diacetatis*.

Medicinal Uses.—It is principally employed externally, in solution in water, as a collyrium in ophthalmia, an astringent in gonorrhœa, and as a wash in external inflammation. Internally it is given cautiously, and combined with opium, in protracted diarrhœa, and in pulmonary and intestinal hæmorrhage. Dose, gr. ss. to gr. j.

LIQUOR PLUMBI DIACETATIS.

Solution of Diacetate of Lead.

Aqua Lithargyri Acetati, P.L. 1788.

Liquor Plumbi Acetatis, P.L. 1809.

Liquor Plumbi Subacetatis, P.L. 1824.

Take of Acetate of Lead two pounds and three ounces,
Oxide of Lead, rubbed to powder, one pound
and four ounces,
Water six pints;

Boil them for half an hour frequently stirring, and when the liquor is cold, add of distilled Water as much as may be sufficient, to measure with it six pints; lastly, strain the solution.

Process.—It has been already mentioned that acetate of lead is a salt, composed of one equivalent each of acid and oxide;

but acetic acid is capable of combining with it an additional equivalent of oxide of lead, by which it becomes a compound of one equivalent of acid and two equivalents of base, forming a subsalt called a *diacetate*.

Properties.—This preparation is colourless; it has an astringent sweetish taste; its specific gravity is 1.260. It is decomposed by spring and other hard water, for reasons which have been already stated with respect to acetate of lead, and the quantity of insoluble salt formed is much larger; distilled water which contains the smallest portion of carbonic acid also decomposes this solution. This preparation is very often improperly prepared with the residue of the distillation of vinegar; it has then a very dark colour and ought to be rejected.

Composition.—This solution, as its name imports, contains diacetate of lead, composed of

One equivalent of Acetic Acid	=	51 or 18.5
Two equivalents of Oxide of Lead..	$112 \times 2 = 224$	„ 81.5
		<hr/>
Equivalent	275	100.

Symbols,—Berzelius and Turner .. $\text{Pb}2\bar{\text{A}}$.

Brande $(2\text{PL} + ac')$.

Incompatibles.—Similar to those which are such with the acetate of lead.

Officinal Preparation.—Liquor Plumbi Diacetatis dilutus.

Medicinal Uses.—External in superficial and phlegmonic inflammations of the skin.

LIQUOR PLUMBI DIACETATIS DILUTUS.

Diluted Solution of Diacetate of Lead.

Aqua Lithargyri Acetati composita, P.L. 1788.

Liquor Plumbi Subacetatis dilutus, P.L. 1809, P.L. 1824.

Take of Solution of Diacetate of Lead a fluidrachm and
a half,

Distilled Water a pint,

Proof Spirit two fluidrachms;

Mix.

Medicinal Use.—Employed as an application in superficial inflammation.


PLUMBI CHLORIDUM.

Chloride of Lead.

Take of Acetate of Lead nineteen ounces,
Boiling distilled Water three pints,
Chloride of Sodium six ounces ;

Dissolve the Acetate of Lead and Chloride of Sodium separately, the former in three pints of distilled Water, and the latter in one pint of distilled Water. The liquors being then mixed together, wash what is precipitated with distilled Water, when it is cold, and dry it.

Process.—When these solutions are mixed, the chloride of sodium and acetate of lead are both decomposed, and the results are acetate of soda, which remains in solution, and chloride of lead, which is precipitated ; and this is accompanied with the transfer of the oxygen of the oxide of lead to the sodium of the chloride :—



Properties.—This compound is colourless, fusible, and on cooling after fusion assumes a horn-like appearance, and hence was formerly called *horn lead*. It dissolves in 30 parts of water

at 60° and in 22 at 212° , separating, as the solution cools, in small flat anhydrous crystals, which have frequently much brilliancy. It is more soluble in water containing a little nitric acid. The solution is decomposed by the alkalis, but potash and soda in excess redissolve the precipitate; the alkaline carbonates throw down carbonate of lead.

Composition.—Chloride of lead consists of

One equivalent of Chlorine.....	36 or 25.7
One equivalent of Lead	104 „ 74.3

Equivalent	140	100.
------------------	-----	------

Symbols.—Berzelius and Turner.. $\text{Pb} + \text{Cl}$.

Brande

(*pl*+*c*).

Uses.—Employed in preparing hydrochlorate of morphia.


PLUMBI IODIDUM.

Iodide of Lead.

Take of Acetate of Lead nine ounces,
Iodide of Potassium seven ounces,
Distilled Water a gallon;

Dissolve the Acetate of Lead in six pints of the Water and strain; add to these the Iodide of Potassium first dissolved in two pints of the Water. Wash what is precipitated, and dry it.

Process.—The properties of iodide of potassium will be presently stated; when it is mixed with a solution of acetate of lead both are decomposed, accompanied with the transfer of the oxygen of the oxide of lead to the potassium of the iodide, and there are formed acetate of potash, which remains dissolved, and iodide of lead, which is precipitated:—



Properties.—Iodide of lead is of a yellow colour; though very sparingly soluble in cold water, it dissolves in it when boiling, and on cooling, shining yellow minute crystalline scales are deposited. It is soluble in solution of potash. It is decomposed and volatilized by heat.

Composition.—Iodide of lead is composed of

One equivalent of Iodine.....	=	126	or	54.78
One equivalent of Lead	=	104	,,	45.22
Equivalent.....		230		100.

Symbols.—Berzelius and Turner.... Pb + I.

Brande (*pl* + *i*).

Impurities and Tests.—See Notes: PLUMBI IODIDUM.

Medicinal Uses.—Used only in the form of ointment.

PLUMBI OXYDUM HYDRATUM.

Hydrated Oxide of Lead.

Take of Solution of Diacetate of Lead six pints,
 Distilled Water three gallons,
 Solution of Potash six pints, or as much as
 may be sufficient to precipitate the oxide;

Mix. Wash what is precipitated with water until nothing alkaline remains.

Process.—This is a case of single elective affinity and decomposition; the oxide of lead combines however with some water, which constitutes it a hydrated oxide; this being insoluble is thrown down, and the acetate of potash formed remains in solution.

Properties.—Hydrate of lead is a perfectly white powder; it is soluble in excess of potash, and therefore in preparing it care should be taken not to employ too much of the alkaline solution. It is also soluble in nitric acid: with hydrochloric acid it forms chloride of lead, and with the sulphuric an insoluble sulphate. It is blackened by hydrosulphuric acid and its salts.

Composition.—It consists of the protoxide of lead combined with water in proportions which have not been determined; its white colour is dependent upon the presence of the water.

Uses.—It is employed in preparing the Disulphate of Quina.

PRÆPARATA È POTASSIO.

Preparations of Potash.

POTASSÆ CARBONAS.

Carbonate of Potash.

Sal Absinthii, Sal Tartari, P.L. 1745.

Kali Præparatum, P.L. 1788.

Potassæ Subcarbonas, P.L. 1809, P.L. 1824.

Take of impure Carbonate of Potash two pounds,
Distilled Water a pint and a half;

Dissolve the impure Carbonate of Potash in the Water, and strain; then pour it off into a proper vessel, and evaporate the Water, that the liquor may thicken; then stir it constantly with a spatula until the salt concretes.

Carbonate of Potash may be prepared more pure from the crystals of Bicarbonate of Potash heated to redness.

Remarks.—Potassium is a metal discovered by Davy in 1807. It is white and bright, readily tarnishes by exposure to the air. The specific gravity is 0·865; its affinity for oxygen is so great, that it not only combines with it at common temperatures, but even decomposes water, and combines with its oxygen so rapidly as to occasion vivid combustion. There are two oxides, of which the oxide or protoxide only, and which is usually called *potash*, combines with acids to form salts.

These oxides are composed of

Oxide, Protoxide, or Potash,	One eq. of Oxygen ..	=	8 or 16·7
	One eq. of Potassium	=	40 „ 83·3

Equivalent	48	100·
------------	------	----	------

Peroxide,	Three equivalents of Oxygen ..	$8 \times 3 =$	24 or 37·5
	One equivalent of Potassium ..	=	40 „ 62·5

Equivalent	64	100·
------------	------	----	------

Process.—Impure Carbonate of Potash is the *Pearlash* of commerce: this consists of the carbonate mixed with various saline and some earthy substances. By solution in water the greater portion of the earthy impurity, at least, is removed, and it is better to employ cold water than the hot, formerly directed by the College, and the quantity is advantageously diminished.

Properties.—This salt is colourless and inodorous; its taste is strong and disagreeable; it does not readily crystallize, and is never kept in crystals; it is deliquescent, attracting in a short time enough water from the atmosphere to become fluid; water dissolves an equal weight of it, and any residue may be considered as impurity. The solution turns vegetable blues green, and yellows brown; it is insoluble in alcohol. When heated to redness it loses about 16 per cent. of water.

Composition.—This salt is composed of

One equivalent of Carbonic Acid ..	=	22	or	31·43
One equivalent of Potash	=	48	„	68·57

Equivalent	70	100·
------------	-------	----	------

It is a sesquihydrate, consisting of

One equivalent of Carbonate of Potash ..	70·	or	84
One and a half equivalent of Water	13·5	„	16

Equivalent	83·5	100
------------	-------	------	-----

As this salt contains one equivalent of its constituent acid and base, its proper appellation is that now given it by the College;

on account of its alkaline property of altering vegetable colours, it has long been called a *subcarbonate*.

Symbols.

Berzelius and Turner $\dot{\text{K}} \ddot{\text{C}} + 1\frac{1}{2}\text{H}$.

Brande $(po + o) + (car + 2o)$, or $(P + car' + 1\frac{1}{2}q)$.

Impurities and Tests.—See Notes : POTASSÆ CARBONAS.

Incompatibles.—Acids and acidulous salts, hydrochlorate of ammonia, acetate of ammonia, lime-water, chloride of calcium, sulphate of magnesia, alum, and most other salts, whether alkaline, earthy, or metallic.

Officinal Preparations and Uses.—Liquor Potassæ Carbonatis, Liquor Potassæ, Potassæ Acetas, Potassæ Sulphas, Potassæ Tartras, Potassii Sulphuretum, Liquor Potassæ Arsenitis, Spiritus Ammoniae, Spiritus Ammoniae Aromaticus, Spiritus Ammoniae Foetidus, Decoctum Aloes compositum, Mistura Ferri composita, Pilulæ Ferri compositæ.

Medicinal Uses.—Antacid and diuretic. Dose from gr. x. to gr. xxx. It is far less pleasant than the bicarbonate of potash; its principal use is in making saline draughts, and bicarbonate of potash.

LIQUOR POTASSÆ CARBONATIS.

Solution of Carbonate of Potash.

Oleum Tartari per deliquum, P.L. 1720.

Lixivium Tartari, P.L. 1745.

Aqua Kali præparati, P.L. 1788.

Liquor Potassæ Subcarbonatis, P.L. 1809, P.L. 1824.

Take of Carbonate of Potash twenty ounces,

Distilled Water a pint;

Dissolve the Carbonate of Potash in the Water, and strain.

Properties.—This solution has a specific gravity of 1.473. It is colourless, inodorous, and possesses the other properties above mentioned. Dose, from ℥x. to fʒj.

POTASSÆ BICARBONAS.

Bicarbonate of Potash.

Potassæ Carbonas, P.L. 1809, P.L. 1824.

Take of Carbonate of Potash six pounds,


Distilled Water a gallon;

Dissolve the Carbonate of Potash in the Water; afterwards pass Carbonic Acid through the solution to saturation. Apply a gentle heat, so that whatever crystals have been formed may be again dissolved. Then set aside [the solution] that crystals may be again produced; the liquor being poured off, dry them.

Carbonic Acid is very easily obtained from Chalk rubbed to powder and mixed with water to the consistence of a syrup, upon which Sulphuric Acid is then poured diluted with an equal weight of Water.

Process.—It has been mentioned that Carbonate of Potash, as indeed its name indicates, is composed of one equivalent each of acid and alkali; the object of the present process is to add another equivalent of carbonic acid, and thus to form Bicarbonate of Potash. Chalk, or carbonate of lime, is composed of one equivalent of carbonic acid and one of lime. When sulphuric acid is added to the carbonate of lime, it is decomposed by the superior affinity of the sulphuric acid for the lime; and the carbonic acid evolved in the gaseous state, being passed into the solution of carbonate of potash, combines with and converts it into bicarbonate, while sulphate of lime remains in the vessel in which the sulphuric acid is poured upon the chalk.

Bicarbonate of Potash = 2 eqs. Acid + 1 eq. Potash.




Sulphate of Lime.

Properties.—This salt is inodorous, colourless, and crystalline. When properly prepared it has scarcely any alkaline taste, and acts but slightly, if at all, upon turmeric paper. It suffers no change by exposure to the air. It requires four times its weight of water at 60° for solution: by boiling water it is partially decomposed, and rendered more soluble by the loss of carbonic acid. It is insoluble in alcohol. When exposed to a low red heat, it loses half its carbonic acid, the whole of its water of crystallization, and returns to the state of carbonate; and this is the method of procuring the latter in a state of purity now adopted by the College, as already seen.


The primary form of this substance is a *right oblique-angled prism*, which is not readily traced in the secondary crystals, but may be derived from cleavage, and is shown in fig. 1. There is also a cleavage parallel to a plane passing through the diagonal marked on the terminal planes.

P on M, or T.....	90°	00'
M on the diagonal plane	53	15
M on T.....	103	25



The planes which appear on the crystals are represented in fig. 2; but the planes *e* are sometimes very disproportionately extended, so as nearly to efface T and *f*, giving to the crystals the character of another primary form.

The planes T do not commonly occur on the crystals, and without these they nearly resemble a secondary form of the *right rhombic prism*; they may, however, be distinguished by the unequal inclination of M on the two adjacent planes. On cleaving or otherwise breaking the crystal, water may be observed between the laminæ, which probably occasions the measurement on the cleavage planes not accurately to agree. This is also the case with many other of the factitious salts.



M on plane parallel to <i>f</i>	127°	35'
M on <i>e</i>	126	45
T on <i>e</i>	156	50
T on <i>f</i>	128	50
<i>e</i> on <i>f</i>	105	40
M on <i>d</i>	111	00
<i>d</i> on <i>d'</i>	138	00

Composition.—This salt consists of

Two equivalents of Carbonic Acid ..	$22 \times 2 = 44$	or 43.56
One equivalent of Potash	$= 48$	„ 47.53
One equivalent of Water	$= 9$	„ 8.91
	<hr/>	<hr/>
Equivalent.....	101	100.

Symbols.—Berzelius and Turner.. $\dot{K} \ddot{C} + \dot{H}$.

Brande

$$(P + 2car + q).$$

Impurities and Tests.—See Notes: POTASSÆ BICARBONAS.

Incompatibles.—These are nearly the same as enumerated when treating of carbonate of potash. It does not, however, produce any precipitate in a solution of sulphate of magnesia; and chloride of mercury is not at all decomposed by it.

Medicinal Uses.—In cases where an alkali is indicated, this preparation offers an agreeable and efficient remedy; and experience has shown that its additional proportion of carbonic acid does not in the least invalidate its alkaline agency. Dose, grs. x. to xxx.

LIQUOR POTASSÆ EFFERVESCENS.

Effervescing Solution of Potash.

Take of Bicarbonate of Potash a drachm,

Distilled Water a pint;

Dissolve the Bicarbonate of Potash in the Water; and pass into it of Carbonic Acid compressed by force, more than sufficient for saturation. Keep the Solution in a well-stopped vessel.

Remarks.—The carbonic acid gas may be procured in the mode already stated. It is intended by the excess of carbonic acid to render the medicine less disagreeable, and this is probably effected without diminishing its power. It is not to be regarded as a definite compound, but as a definite salt mixed with excess of carbonic acid.

LIQUOR POTASSÆ.

Solution of Potash.

Lixivium Saponarium, P.L. 1745.

Aqua Kali puri, P.L. 1788.

Liquor Potassæ, P.L. 1809, P.L. 1824.

Take of Carbonate of Potash fifteen ounces,

Lime eight ounces,

Boiling distilled Water a gallon;

Dissolve the Carbonate of Potash in half a gallon of the water. Sprinkle a little of the water upon the lime in an earthen vessel, and the lime being slacked, add the rest of the water. The liquors being immediately mixed together in a close vessel, shake them frequently until they are cold. Then set aside, that the Carbonate of Lime may subside. Lastly, keep the supernatant liquor when poured off, in a well-stopped green glass bottle.

Process.—This is a case of single elective affinity and decomposition:—the lime has a strong affinity for the carbonic acid which has been expelled from it by heat; and when it is mixed with the carbonate of potash, owing to the greater affinity existing between the earth and the acid than between the alkali and the acid, the carbonate of potash is decomposed, and the potash remaining in solution, the carbonate of lime formed is precipitated.

Solution of Potash.

Solution of
Carbonate
of Potash.

Water.
Potash.

Carbonic Acid.

Lime.

Carbonate of Lime.

Properties.—Solution of potash is limpid, colourless, and inodorous; its taste is extremely acrid and caustic; and, when

rubbed between the fingers, it feels soapy, in consequence of a partial solution of the cuticle. Its specific gravity is 1.063. It should be carefully preserved from contact with the air, in order to prevent the absorption of carbonic acid; and as it is apt to act upon and destroy flint glass, the College have advantageously ordered it to be kept in green glass bottles.

Impurities and Tests.—See Notes: POTASSÆ LIQUOR.

Incompatibles.—Acids, acidulous salts, sesquicarbonate, acetate and hydrochlorate of ammonia, preparations of metals and earths held in solution by acids; chloride and bichloride of mercury.

Officinal Preparations.—Potassæ Hydras, Potassa cum Calce, Antimonii Oxysulphuretum.

Medicinal Uses.—Antacid, diuretic, alterative, and lithontriptic; it has also been found useful in some cutaneous diseases, as in lepra, psoriasis, &c. Dose ℥ x. to f 3 ss. It is recommended to give it in veal broth or in table beer: the latter is said to disguise its nauseous flavour completely. Care, however, ought to be taken that the beer is not sour.

POTASSÆ HYDRAS.

Hydrate of Potash.

Lapis Infernalis sive Septicus, P.L. 1720.

Kali Purum, P.L. 1788.

Potassa Fusa, P.L. 1809, P.L. 1824.

Take of Solution of Potash a gallon;

Evaporate the water in a clean iron vessel over the fire, until, the ebullition being finished, the Hydrate of Potash liquefies; pour this into proper moulds.

Properties.—Hydrate of potash is a compound of potash and water; when pure it is white, hard and brittle, but as usually prepared for medicinal purposes, it contains the various impurities of the solution, and frequently peroxide of iron, acquired during evaporation. It is generally of a brownish and sometimes of a bluish tint, is extremely caustic, and very deliquescent, attracting first water and then carbonic acid from the atmosphere; water dissolves nearly an equal weight of it, and during solution heat is extricated.

Unlike the carbonate and bicarbonate of potash, it dissolves readily in alcohol. It possesses in the strongest degree the properties denominated alkaline.

Hydrate of potash melts when exposed to a low red heat; but so great is the affinity existing between the potash and the water, that although they may be evaporated together at a white heat, the water cannot be separated by it. During the preparation of the hydrate of potash a portion of the potash becomes peroxide of potassium, but the additional oxygen thus acquired, is given out again in the gaseous state, during solution in water.

Composition.—This preparation consists of

One equivalent of Potash = 48 or 84·2

One equivalent of Water = 9 „ 15·8

Equivalent 57 100·

Symbols,—Berzelius and Turner K H.

Brande (P + q).

Medicinal Uses.—Potassæ hydras is used only externally as a caustic; excepting for particular purposes, the argenti nitras or lunar caustic is generally preferred; for, on account of the deliquescent property of the hydrate, it is difficult to confine its action within the requisite limits.

POTASSA CUM CALCE.

Potash with Lime.

Causticum Commune fortius, P.L. 1745.

Calx cum Kali puro, P.L. 1788.

Potassa cum Calce, P.L. 1809, P.L. 1824.

Take of Hydrate of Potash,

Lime, each an ounce;

Rub them together and keep them in a well-stopped vessel.

POTASSÆ ACETAS.

Acetate of Potash.

Sal Diureticus, P.L. 1745.*Kali Acetatum*, P.L. 1788.*Potassæ Acetas*, P.L. 1809, P.L. 1824.

Take of Carbonate of Potash a pound,

Acetic Acid twenty six fluidounces,

Distilled Water twelve fluidounces ;

Add the Carbonate of Potash to the Acid first mixed with the water, to saturation, then strain. Evaporate the liquor in a sand-bath, the heat being cautiously applied, until the salt is dried.

Process.—In this operation the carbonic acid of the carbonate is expelled in the gaseous state on account of the more powerful affinity existing between acetic acid and potash, than between carbonic acid and potash.

Properties.—Acetate of potash thus obtained is colourless pulverulent salt ; it is nearly inodorous, and has a pungent saline taste ; it is extremely deliquescent, very soluble in water, and is dissolved also by alcohol ; it is decomposed by a red heat and converted into carbonate of potash. As usually prepared it has a foliated appearance, which is given to it by fusion and cooling ; in this operation however, unless very carefully conducted, the acetate is apt to be decomposed.

Composition.—Acetate of potash consists of

One equivalent of Acetic Acid = 51 or 51.5

One equivalent of Potash = 48 „ 48.5

99 100.

Symbols.—Berzelius and Turner . . $\dot{\text{K}} \bar{\text{A}}$.

Brande $(\text{P} + \text{ac}')$.

Impurities and Tests.—See Notes : POTASSÆ ACETAS.

Incompatibles.—It is decomposed by the sulphuric, nitric, and hydrochloric acids, the acetic acid being expelled. It is also decomposed by sulphate of soda and sulphate of magnesia, and by several other earthy and metallic salts.

Medicinal Uses.—In small doses it is diuretic, and in larger ones mildly cathartic. Dose as a diuretic from \mathfrak{zj} . to \mathfrak{zj} . ; as a cathartic from \mathfrak{zj} to \mathfrak{zij} . As it is deliquescent it must be exhibited in solution.

POTASSÆ SULPHAS.

Sulphate of Potash.

Tartarum Vitriolatum, P.L. 1720, P.L. 1745.

Kali Vitriolatum, P.L. 1788.

Potassæ Sulphas, P.L. 1809, P.L. 1824.

Take of the Salt which remains after the distillation of
Nitric Acid two pounds,
Boiling Water two gallons;

Ignite the Salt in a crucible until the excess of Sulphuric Acid is entirely expelled, then boil it in the two gallons of Water until a pellicle floats, and the liquor being strained, set it aside that crystals may be formed. The liquor being poured off, dry them.

Process.—It has been already explained that the salt remaining after the distillation of nitric acid is composed of bisulphate of potash and water; the excess of acid is now directed to be expelled by heat instead of, as formerly, saturated by the addition of carbonate of potash.

Properties.—This salt is colourless, inodorous, bitter and rather hard; water at 60° dissolves only one sixteenth of its weight, but boiling water a much larger quantity; it is insoluble in alcohol. It suffers no change by exposure to the air. When exposed to a strong heat it merely decrepitates, losing but little weight, for it contains no water of crystallization.

The primary form of this salt is a *right rhombic prism*; M M' and P are primary planes.

Fig. 1 is a single modified crystal.

M on M'	120° 30'
M on h	120 45
M on e	146 22
h on c	146 10
e on e'	131 12

Fig. 1.




Fig. 2. is the compound crystal, which consists of three single crystals, so united that their upper edges meet at angles of 120° , and consequently the planes of junction incline to each other at the same angle. Hence

M on M''	119° 30'
e on e''	130 24





Fig. 3. is one of the common bi-pyramidal crystals, whose relation to the preceding figures may be perceived from the corresponding letters on the planes.



The union of these three crystals takes place at an angle of 120° .

Composition.—This salt is composed of

One equivalent of Sulphuric Acid . . .	= 40	or 45.45
One equivalent of Potash	= 48	„ 54.55
Equivalent	88	100.

Symbols.—Berzelius and Turner $\overset{\cdot\cdot\cdot}{K} \overset{\cdot\cdot\cdot}{S}$.

Brande $(po + o) + (S + 3o)$, or $(P + S')$.

Adulteration.—This salt is so extremely cheap, and in its crystalline state any mixture would be so obvious, that adulteration is hardly to be suspected. It may, however, be observed, that the solution should produce no change in the colour of litmus or turmeric paper; no precipitate with solution of sulphate of silver, nor any upon the addition of ammonia or its sesquicarbonate.

Incompatibles.—The solution of this salt is decomposed by tartaric acid, which forms crystals of bitartrate of potash; by chloride of barium, barytes water, and chloride of calcium, but not by lime-water as has been asserted; it also decomposes the solutions of acetate and diacetate of lead.

Medicinal Uses.—It should be exhibited in the form of powder, in conjunction with rhubarb or some other purgative medicine. On account of its hardness it is an eligible substance for triturating with other bodies and dividing powders; with this intention it enters into the composition of Pulvis Ipecacuanhæ Compositus. Dose, gr. x. to $\overline{3}$ ss.

POTASSÆ BISULPHAS.

Bisulphate of Potash.

Potassæ Supersulphas, P.L. 1809, P.L. 1824.


Take of the Salt which remains after the distillation of
 Nitric Acid two pounds,
 Sulphuric Acid one pound,
 Boiling Water four pints;

Dissolve the Salt in the Water, and add the Acid to it, and mix. Lastly, boil down, and set aside, that crystals may be formed.

Process.—When bisulphate of potash is dissolved in water, and the solution is allowed to crystallize, it very commonly happens that some sulphate and a large proportion of sesquisulphate are obtained instead of bisulphate, owing to the partition of the excess of sulphuric acid between the water and the sulphate of potash. This inconvenience is remedied by the addition of sulphuric acid now directed to be employed.

Properties.—This salt is extremely acid and bitter; it is very soluble in water, the solution acts strongly upon vegetable blue colours, and decomposes the alkaline, earthy, and metallic carbonates with effervescence. By a red heat the water of crystallization and half the acid are expelled, and common sulphate of potash remains. The primary form of the crystal of this salt has not been ascertained; it may be either a *right rhombic prism*, or an *octahedron with rhombic bases*. There appears to be but one cleavage, namely, parallel to the plane *a*. The crystal is often much flatter than the sketch.

<i>a</i> on <i>c</i> or <i>c'</i>	135° 0'
— <i>o</i> or <i>o'</i>	108 30
<i>c</i> on <i>c'</i>	125 10
<i>o</i> on <i>o'</i>	103 52
<i>o</i> on <i>o''</i>	142 44



Composition.—It is composed of

Two equivalents of Sulphuric Acid ..	$40 \times 2 = 80$	or 54.80
One equivalent of Potash	$= 48$	„ 32.87
Two equivalents of Water	$9 \times 2 = 18$	„ 12.33

Equivalent....	146	100.
----------------	-----	------

Symbols.

Berzelius and Turner $K\ddot{S}^2 + 2\dot{H}$.

Brande $(po + o) + 2(S + 3o)$, or $(P + 2S' + 2q)$.

Incompatibles.—This salt is incompatible with alkalis, earths, and their carbonates; many metals and most oxides are acted upon by the excess of acid which it contains.

Medicinal Uses.—It is exhibited combined with other purgatives, especially with Rhubarb. Dose, gr. x. or more.

POTASSÆ TARTRAS.

Tartrate of Potash.

Tartarum Solubile, P.L. 1745.

Kali Tartarizatum, P.L. 1788.


Potassæ Tartras P.L. 1809, P.L. 1824.

Take of Bitartrate of Potash, powdered, three pounds,
Carbonate of Potash sixteen ounces, or as
much as may be sufficient,

Boiling Water six pints;

Dissolve the Carbonate of Potash in the boiling Water, then add the Bitartrate of Potash, and boil. Strain the liquor, and afterwards boil it down until a pellicle floats, and set it aside that crystals may be formed. The liquor being poured off, dry these, and again evaporate the liquor that crystals may be produced.


Process.—The nature and composition of both the salts employed in this formula have been stated; when they are made to act upon each, the excess of tartaric acid in the bitartrate of potash combines with the potash of the carbonate and expels its carbonic acid in the gaseous state; so that one equivalent of each produces two equivalents of the neutral tartrate.



This salt has a saline bitter taste; it is soluble in less than twice its weight of water, and hence its former name of *soluble tartar*, to distinguish it from the bitartrate of potash: it is nearly insoluble in alcohol. In a damp atmosphere it attracts moisture; by a red heat it is decomposed and converted into carbonate of potash. It is commonly met with in the shops in the state of powder, but it ought always to be crystallized. When this salt has been properly prepared it does not alter either litmus or turmeric paper.

The primary form of tartrate of potash is a *right oblique-angled prism*, with cleavages parallel to the lateral planes.

M on T	89° 30'
M on <i>e</i>	142 13
M on <i>b</i>	107 30
T on <i>e</i>	127 17
T on <i>b'</i>	103 40



Composition.—Tartrate of potash consists of

One equivalent of Tartaric Acid ..	=	66	or	57.9
One equivalent of Potash	=	48	„	42.1
		114		100.
Equivalent....		114		100.

Symbols,—Berzelius and Turner .. $\dot{\text{K}}\bar{\text{T}}$.

Brande

(*tar* + P).

Impurities and Tests.—See Notes: POTASSÆ TARTRAS.

Incompatibles.—Tartrate of potash is decomposed by most acids, and many acidulous salts, for when added to a solution, they occasion the formation and crystallization of bitartrate of potash. It is decomposed by lime-water and chloride of calcium, and by solutions of lead and silver, &c.

Medicinal Uses.—It is a mild and efficient purgative, and when given with resinous purgatives or senna, it corrects their griping properties by accelerating their operation. Dose, ʒj. to ʒj. in solution.

POTASSII BROMIDUM.

Bromide of Potassium.

Take of Bromine two ounces,

Carbonate of Potash two ounces and one
drachm,

Iron Filings an ounce,

Distilled Water three pints;

First add the Iron, and afterwards the Bromine, to a pint and a half of the distilled Water. Set them by for half an hour, frequently stirring them with a spatula. Apply a gentle heat, and when a greenish colour occurs, pour in the Carbonate of Potash dissolved in the remainder of the Water. Strain, and wash what remains in two pints of boiling distilled Water, and again strain. Let the mixed liquors evaporate so that crystals may be formed.


Remarks.—Bromine is an elementary fluid body, which was discovered by M. Balard in 1826. It exists in sea-water and some saline springs, combined probably with magnesium or sodium; but the quantity is so small that 100 pounds of sea-water yielded only $3\frac{1}{3}$ grains of bromine. It is separated by means of the greater affinity of chlorine for the base with which it is combined, and when set free is dissolved by æther.

The properties of bromine are, that at common temperatures it is a deep reddish brown liquid, of a very disagreeable and suffocating odour; hence its name. Its specific gravity is about 3. At a little below 0° it becomes solid. It dissolves sparingly in water, but combines readily with alcohol. It is considerably volatile, for at average temperatures it emits a red-coloured vapour very similar in appearance to nitrous acid; when heated to about 116° it boils; and when passed through red-hot earthen tubes it suffers no alteration of property, not being resolved into any simpler form of matter. It is poisonous; its test is a solution of starch, to which it gives a yellowish tint.

It resembles Oxygen, Chlorine and Iodine in being elicited at the positive pole of the voltaic battery, and like them also it is a powerful supporter of combustion, some metals, as antimony, burning spontaneously when thrown upon it. In smell also it

greatly resembles chlorine and iodine, but is much more powerful than either of them; like them also it combines with oxygen and hydrogen, and forms bromic and hydrobromic acid.

Process for Bromide of Potassium.—Bromine unites with the metals by direct combination; thus the first step in this operation is to obtain bromide of iron; when this is procured, it is decomposed by the carbonate of potash; oxygen is transferred from the potash to the iron, which, combining with it, and the oxide uniting with the carbonic acid, is converted into carbonate of iron and precipitated, while the Bromide of Potassium remains in solution, and is by evaporation crystallized.



Properties.—*Bromide of Potassium* is a colourless inodorous salt; it crystallizes in cubes or quadrangular prisms; they contain no water: it has a penetrating taste, is very soluble in water, and more so in hot than cold; alcohol dissolves a little of it; it decrepitates when heated, and undergoes igneous fusion without suffering decomposition or any alteration of properties. It is decomposed by chlorine, which expels bromine, and chloride of potassium is formed.

Composition.—This salt consists of

One equivalent of Bromine	=	78	or	66.1
One equivalent of Potassium	=	40	,,	33.9
		118		100.
Equivalent		118		100.

Symbols.—Berzelius and Turner . . . K Br.

Brande (*po + b*).

Impurities and Tests.—See Notes: POTASSII BROMIDUM.

Incompatibles.—Acids, acidulous and metallic salts.

Medicinal Uses.—This salt has been successfully employed in cases of enlarged spleen by Dr. Williams, of St. Thomas's Hospital. Dose, from 3 to 10 grs. two or three times a day.

POTASSII IODIDUM.

Iodide of Potassium.

Take of Iodine six ounces,
 Carbonate of Potash four ounces,
 Iron Filings two ounces,
 Distilled Water six pints;

Mix the Iodine with four pints of the Water, and add the Iron, stirring them frequently with a spatula for half an hour. Apply a gentle heat, and when a greenish colour occurs, add the Carbonate of Potash, first dissolved in the two pints of Water, and strain. Wash what remains with two pints of boiling distilled water, and again strain. Let the mixed liquors be evaporated, so that crystals may be formed.

Process.—Although iodine like bromine is capable of immediately uniting with the metals, yet an indirect process is in both cases preferred; and as the principle is the same, substituting iodine for bromine in the last explanation and diagram both will be explained.

Properties.—Iodide of potassium is colourless, inodorous, crystallizes in cubes, which contain no water; it has a penetrating taste. Water at 65° dissolves nearly one and a half time its weight; it is sparingly soluble in absolute alcohol, but more so in that which contains water. It fuses at a moderate temperature, but undergoes no change.

Composition.—This salt is composed of

One equivalent of Iodine	=	126	or	76
One equivalent of Potassium	=	40	,,	24
Equivalent		166		100.

Symbols.—Berzelius and Turner K I.

Brande (*po+i*).

Incompatibles.—Acids, Acidulous and Metallic Salts.

Medicinal Uses.—Is a most valuable medicine, for the introduction of which into this country we are indebted to Dr. Williams of St. Thomas's Hospital. It is but justice due to him to refer to an excellent paper read at one of the evening meetings of the College of Physicians, in which he has illustrated the medicinal properties of this salt in various forms of the secondary symptoms of syphilis. Dose, from gr. v. to gr. x. or more, two or three times a day.

LIQUOR POTASSII IODIDI COMPOSITUS.

Compound Solution of Iodide of Potassium.

Take of Iodide of Potassium ten grains,

Iodine five grains,

Distilled Water a pint ;

Mix, that they may be dissolved.

Remarks.—In this mixture the iodide of potassium, by uniting with an additional portion of iodine, renders it soluble in water ; it has been called *ioduretted iodide of potassium*. It is a brown-coloured solution, and has the peculiar smell and taste of iodine, and its reaction on starch.

Medicinal Uses.—This is another mode of exhibiting iodine, which has been found very serviceable in dispersing some forms of bronchocele. Dose, from ℥v. to ℥x., which may be considerably increased ; but its effects varying on different constitutions, its exhibition requires the exercise of great judgment.

POTASSII SULPHURETUM.

Sulphuret of Potassium.

Kali Sulphuratum, P.L. 1788.

Potassæ Sulphuretum, P.L. 1809, P.L. 1824.

Take of Sulphur an ounce,

Carbonate of Potash four ounces ;


Rub them together, and place them upon the fire in a covered crucible, until they have united.

Remarks.—Sulphur is a well-known elementary or undecomposed body, which sometimes occurs in nature nearly pure, but more commonly in combination with the metals, forming sulphurets. The greater part of that which is used in the arts is the produce of volcanic countries. Its colour is yellow with a shade of green ; it is nearly inodorous and tasteless, insoluble in water, and is with difficulty dissolved by spirit of wine. The sp. gr. of sulphur is about 2 ; at a moderate temperature it melts, and at a higher one is converted into vapour ; it burns readily with a lambent blue flame, and suffocating vapours of sulphurous acid are formed by its combining with the oxygen of the air during combustion. When pure, or crystallized, it is frequently translucent. The primary form of the crystal is *an acute octahedron with a rhombic base*, subject to various modifications.

In commerce, the various kinds of sulphur are distinguished by the names of Crude Sulphur ; Flowers of Sulphur (the Sulphur Sublimatum of the Pharmacopœia) ; and Roll Sulphur, prepared by melting crude sulphur, and pouring it while fluid into moulds.

Process.—When sulphur and carbonate of potash are heated together, the carbonic acid is expelled from the latter, and three-fourths of the potash, or oxide of potassium, are decomposed ; its oxygen combines with sulphur to form sulphuric acid, and this uniting with the one-fourth of undecomposed potash, sulphate of potash results. The potassium of the decomposed potash combines also with sulphur, and sulphuret of potassium is formed ; so that when the operation has been properly conducted the Potassii Sulphuretum of the Pharmacopœia is chiefly a compound of sulphate of potash and sulphuret of po-

tassium. The peculiar properties of the preparation depend upon the sulphuret of potassium.



Properties.—This substance is hard ; it is of a liver-brown colour, and hence its former name of Hepar Sulphuris. It is inodorous while dry, but when moistened it emits a smell of hydrosulphuric acid ; it dissolves readily in water. Its taste is acrid and bitter. By exposure to the air this preparation is soon spoiled, for the sulphur and potassium both attracting oxygen, sulphate of potash is formed ; it then becomes inodorous and white, and is totally unfit for use.

Incompatibles.—This compound is decomposed by acids, they expelling hydrosulphuric acid and precipitating sulphur. It is decomposed also by solutions of most of the metals, which, uniting with the sulphur, are precipitated in the state of sulphuret.

Medicinal Uses.—It is principally used externally in cutaneous diseases, and has been recommended as a lotion for the itch in infants, and is stated to have succeeded after the sulphur ointment has failed. It is rarely used internally.

PRÆPARATA È SODIO.

Preparations of Sodium.

SODÆ CARBONAS.

Carbonate of Soda.

Natron præparatum, P.L. 1788.

Sodæ Subcarbonas, P.L. 1809, P.L. 1824.

Take of Impure Carbonate of Soda two pounds,

Distilled Water four pints ;

Boil the impure Carbonate of Soda in the Water, and strain it while yet hot. Lastly, set it by that crystals may be formed.

Remarks.—Carbonate of Soda is a compound of carbonic acid and the alkali soda ; this substance has been long known as the *fossil alkali* or *natron* ; it occurs in various parts of the earth in the state of a peculiar carbonate. It is obtained by burning certain plants and sea-weed, but is now generally prepared by the decomposition of common salt.

In 1807 it was shown by Davy, that soda is a metallic oxide, very similar in many of its properties to potassium, and it may be procured from soda by processes similar to those used for obtaining potassium from potash.

Sodium is a brilliant white metal ; it is soft, malleable, and it tarnishes rapidly by exposure to the air, owing to oxidizement. Its specific gravity is 0.972. It softens at about 122°, fuses at about 190°, and at a white heat it is volatilized. It burns when heated in contact with air, and is converted to soda or oxide of sodium. Under particular circumstances it decomposes water

with combustion, as potassium does, but in general it is accompanied merely by a hissing noise.

Sodium, like potassium, forms two compounds with oxygen; the protoxide, which is the alkali soda, consists of one equivalent of oxygen 8, and 1 equivalent of sodium $24=32$. The peroxide is a sesquioxide; it is quite unimportant; being decomposed by water, it forms no salts with acids.


Soda resembles potash in appearance and is obtained in the same way; it is best known in the state of hydrate, and procured, as hydrate of potash is, by evaporating a solution and igniting the residue. In medicine, however, it is not employed in a separate state.

Process.—The impure carbonate of soda formerly employed was *barilla*, obtained by the burning of certain plants in Spain; there is, however, now prepared, and with greater economy, a much preferable, though still not quite pure carbonate of soda. When this is dissolved in hot water to saturation, crystals are deposited as the solution cools, which are carbonate of soda.

Properties.—The crystals of this salt are frequently very large, and are obtained for various uses of considerable purity; the primary form of the crystal appears to be an *oblique rhombic prism*.

This figure represents the ordinary shape of the crystals.


Fig. 1.



P on M, or M'	108° 43'
P on e, or e'	128 52
P on h	121 20
M on M'	76 12
M on h	128 6
M on k	141 54
e on e'	79 44
e on k	143 8

The crystals represented by fig. 2. are reduced in height, and so thin as to leave scarcely a vestige of the planes M and h, and several are hemitropes, the plane of imaginary section being parallel to P.

Fig. 2.



Qualities.—Carbonate of soda is colourless and inodorous; its taste is alkaline and disagreeable, but less so than that of carbonate of potash; the crystals contain a large quantity of water, the greater part of which they readily lose by exposure to the air, and at high temperatures the salt becomes fluid and boils. Water at 60° dissolves at least half its weight of carbonate of soda, and boiling water considerably more. The solution possesses the alkaline property of rendering vegetable yellows brown.

Composition.—Carbonate of soda in the crystallized state consists of

One equivalent of Carbonic Acid	= 22	or 15.3
One equivalent of Soda	= 32	„ 22.2
Ten equivalents of Water 9×10	= 90	„ 62.5
	<hr/>	<hr/>
Equivalent	144	100.

Soda. Crystals of Carbonate of Soda.

Symbols.—Berzelius and Turner.. Na . $\text{Na}\ddot{\text{C}} + 10\text{H}$.

Brande (S). $(\text{S} + \text{car}' + 10q)$.

Tests and Impurities.—See Notes : SODÆ CARBONAS.

Incompatibles.—This salt is incompatible with acids, acidulous salts, lime-water, hydrochlorate of ammonia, earthy and metallic salts.

Pharmaceutic Uses.—In preparing Ferri Sesquioxylum, Pilulæ Ferri Compositæ, Magnesiae Carbonas, &c.

Medicinal Uses.—These are similar to those of the carbonate of potash, but this salt is preferable as being more mild and less nauseous. Dose, from gr. x. to 3ss. twice or thrice a day.

SODÆ CARBONAS EXSICCATA.

Dried Carbonate of Soda.

Sodæ Subcarbonas exsiccata, P.L. 1809, P.L. 1824.

Take of Carbonate of Soda a pound ;

Apply heat to the Carbonate of Soda in a proper vessel, until it is dried, and afterwards heat it to redness. Lastly, rub it to powder.

Process.—The greater part of the water which crystallized carbonate of soda contains, is first expelled by a moderate degree of heat, and the total expulsion is effected by ignition ; the first is applied because the fused salt, if strongly heated, would boil over, and the ignition is requisite to render the preparation of uniform strength.

Composition.—It follows from what has been stated that this anhydrous salt consists of

One equivalent of Carbonic Acid	= 22	or 40·7
One equivalent of Soda	= 32	„ 59·3
	<hr/>	<hr/>
Equivalent	54	100·

Symbols.—Berzelius and Turner .. $\text{Na} + \ddot{\text{C}}$.

Brande (S + *car'*).

Medicinal Use.—In this dry state carbonate of soda may be exhibited in the form of powder mixed with other medicines. Dose, gr. v. to gr. xv.

SODÆ SESQUICARBONAS.

Sesquicarbonate of Soda.

Sodæ Carbonas, P.L. 1809, P.L. 1824.

Take of Carbonate of Soda seven pounds,

Distilled Water a gallon ;

Dissolve the Carbonate of Soda in the Water, and strain; then pass Carbonic Acid into the Liquor to saturation that the Salt may subside. Dry this with a gentle heat, wrapped and pressed in cloth.

Process.—In preparing this salt carbonic acid is to be obtained in the same mode as directed for the bicarbonate of potash; in this case, however, instead of forming a bicarbonate, the product contains one fourth less carbonic acid, and is consequently a sesquicarbonate, consisting of

One and a half equivalent of Carbonic Acid ..	33	or 39·76
One equivalent of Soda	32	„ 38·55
Two equivalents of Water	18	„ 21·69
	<hr/>	<hr/>
Equivalent	83	100·

Properties.—This salt is colourless, and in the form of minute indistinct crystals; it is much less soluble in water than the car-

bonate, and hence it crystallizes as it is formed. The solution acts, though slightly, on turmeric paper. Unlike the carbonate it does not precipitate a solution of sulphate of magnesia, and its taste is much less alkaline and disagreeable. When strongly heated it loses half an equivalent of carbonic acid and all its water, and is reduced to dried carbonate of soda. This salt occurs native in Africa and also in South America.

Symbols,—Berzelius and Turner . . $\text{Na}^2 \text{C}^3 + 4\text{H}$.

Brande $(\text{S} + 1\frac{1}{2}\text{C} + 2\text{q})$.

Incompatibles.—The same as the carbonate, except that it does not decompose the salts of magnesia until the mixture is heated.

Impurities and Tests.—See Notes : SODÆ SESQUICARBONAS.

Medicinal Uses.—Similar to those of the carbonate. Dose, gr. x. to gr. xxx. This salt is largely employed for the purpose of making what are called *sodaic powders*, by mixture with tartaric acid, and taken during effervescence; these are sometimes intended as a substitute for *soda water*, from which they differ in yielding tartrate of soda, with a portion of carbonic acid diffused through the solution, instead of consisting of bicarbonate of soda with excess of carbonic acid gas. As they cannot have any alkaline action, they ought not to be employed where such is required.

SODÆ SULPHAS.

Sulphate of Soda.

Sal Catharticus Glauberi, P.L. 1745.

Natron Vitriolatum, P.L. 1788.

Sodæ Sulphas, P.L. 1809, P.L. 1824.

Take of the Salt which remains after the distillation of
Hydrochloric Acid two pounds,
Boiling Water two pints,
Carbonate of Soda as much as is necessary;


Dissolve the Salt in the Water; then gradually add as much Carbonate of Soda as is sufficient to saturate the Acid. Boil down until a pellicle appears, and, the Liquor

being strained, set it aside, that crystals may be formed. The Liquor being poured off, dry them.

Process.—The production of sulphate of soda during the preparation of hydrochloric acid has been explained. The excess of sulphuric acid remaining after the decomposition of the chloride of sodium being comparatively small, the saturation of it by carbonate of soda, instead of expulsion by heat, is of less consequence than in the case of sulphate of potash.

Properties.—Sulphate of soda very readily crystallizes. The primary form of this salt is an *oblique* rhombic prism.

P on M, or M'	101° 20'
P on e, or e'	133 18
P on h	107 44
P on c'	130 45
M on M'	80 24
M on h	130 12
M on l	162 38
M on k	139 48



This salt has a very bitter taste. By exposure to the air it effloresces, and a white powder is left. It is extremely soluble in water, three parts of which, at 60°, dissolve one part of the salt: boiling water dissolves its own weight. It is insoluble in alcohol. When exposed to heat it first undergoes watery fusion by melting in its water of crystallization; when the water has evaporated it becomes white, and at a red heat it melts.

Composition.—Sulphate of soda is composed of

One equivalent of Sulphuric Acid ..	= 40	or 55.55
One equivalent of Soda	= 32	„ 44.45
Equivalent	72	100.

In the crystallized state this salt consists of

One equivalent of Sulphuric Acid...	= 40	or 24.69
One equivalent of Soda	= 32	„ 19.75
Ten equivalents of Water	= 90	„ 55.56
Equivalent	162	100.

Symbols.—Berzelius and Turner .. $\text{Na}\ddot{\text{S}} + 10\text{H}$.

Brande..... $(\text{S} + \text{S}' + 10\text{q})$.

Tests and Impurities.—See Notes: SODÆ SULPHAS.

Incompatibles.—Carbonate of potash, chloride of calcium,

solution of barytes and barytic salts; acetate and diacetate of lead and nitrate of silver, if the solutions be strong.

Medicinal Uses.—A common and efficient purgative. Its nauseous taste may be in a great degree disguised by the addition of a small quantity of lemon-juice, or of bitartrate of potash. Dose, \bar{z} ss. to \bar{z} ij.

SODÆ POTASSIO-TARTRAS.

Potassio-Tartrate of Soda.

Natron Tartarizatum, P.L. 1788.

Soda Tartarizata, P.L. 1809, P.L. 1824.

Take of Bitartrate of Potash, powdered, sixteen ounces,

Carbonate of Soda twelve ounces,

Boiling Water four pints;


Dissolve the Carbonate of Soda in the boiling Water, and add gradually the Bitartrate of Potash. Strain the Liquor; then apply a gentle heat, until a pellicle floats, and set aside, that crystals may be formed. The Liquor being poured off, dry them. Evaporate the Liquor again that it may yield crystals.

Process.—In this preparation the excess of tartaric acid contained in the bitartrate of potash is saturated with soda, by decomposing the carbonate and expelling its carbonic acid in the gaseous state.


Properties.—This salt forms large and beautiful crystals. The form derived from cleavage is a *right rhombic prism*. This is modified in the crystals measured, as shown in fig. 1.

Fig. 1.

P on M, or M'	90°	0'
P on c	138	50
M on M'	100	0
M on g	163	0
M' on g'		



There is a peculiarity in the crystals of this substance. They are produced nearly in halves, and appear to have rested or been formed on planes which would have passed through the middle of the entire crystal. One of these natural segments is shown in fig. 2; but in others of them the front half of fig. 1 is the portion produced, the plane *f* being then uppermost. In some of the segments, however, there is a slight deviation from this exactness of position of the planes *f* or *h*.



This salt, sometimes called *Rochelle Salt* and *Sel de Seignette*, is colourless, inodorous, bitter and saline, very slightly efflorescent when exposed to the air. It is soluble in five parts of water at 60°, and more so in boiling water. It is decomposed by a strong heat; the residuum is a mixture of carbonate of potash and carbonate of soda.

Composition.—This is a double salt, consisting of

One equivalent of Tartrate of Potash	=	114	or	40.
One equivalent of Tartrate of Soda..	=	98	„	34.5
Eight equivalents of Water	=	72	„	25.5
Equivalent		284		100.

Symbols.—

Berzelius } $\dot{K}\bar{T} + \dot{Na}\bar{T} + 8\dot{H}$.
and Turner }

Brande .. $(tar' + P + tar' + S + 8q)$, or $(2tar' + P + S + 8q)$.

Impurities and Tests.—See Notes: SODÆ POTASSIO-TARTRAS.

Incompatibles.—Most acids and acidulous salts, except the bitartrate of potash. By the action of the acids the tartrate of potash is converted into bitartrate. The acetate and diacetate of lead, and the salts of lime, are decomposed by this compound.

Medicinal Uses.—Dose, as a purgative, from 3ij. to 3j.

LIQUOR SODÆ EFFERVESCENS.

Effervescing Solution of Soda.

Take of Sesquicarbonate of Soda a drachm,

Distilled Water a pint;

Dissolve the Carbonate of Soda in the Water; and

pass into it, compressed by force, more Carbonic Acid than is sufficient for saturation. Keep the solution in a well-stopped vessel.

Remarks.—A solution thus prepared is commonly known by the name of soda water; it is a pleasant mode of exhibiting the alkali, and its powers are supposed not to be diminished by the excess of carbonic acid with which it is combined.

Tests.—See Notes: SODÆ CARBONATIS LIQUOR EFFERVESCENS.

LIQUOR SODÆ CHLORINATÆ.

Solution of Chlorinated Soda.

Take of Carbonate of Soda a pound,
Distilled Water forty-eight fluidounces,
Chloride of Sodium four ounces,
Binoxide of Manganese three ounces,
Sulphuric Acid four ounces;

Dissolve the Carbonate of Soda in two pints of Water; then put the Chloride of Sodium and Binoxide of Manganese, rubbed to powder, into a retort; and add to them the Sulphuric Acid, previously mixed with three fluidounces of Water and cooled. Heat the [mixture] and pass the Chlorine first through five fluidounces of Water, and afterwards into the solution of Carbonate of Soda above directed.

Process.—The extrication of chlorine by the action of sulphuric acid upon a mixture of chloride of sodium and binoxide of manganese, has been already explained. In this case it is passed through a small quantity of water, which will combine with any hydrochloric acid that might otherwise pass into the solution of carbonate of soda.

The precise nature of this solution, usually called *Labarraque's Soda disinfecting Liquid*, has not been determined. When the quantity of chlorine gas liberated from the ingredients here directed is not exceeded, no carbonic acid is expelled from the carbonate of soda, and the compound form may be made to cry-

stallize, and consists of chlorine and carbonate of soda ; these crystals when redissolved reproduce the disinfecting liquid.

Properties.—This solution is of a pale yellow colour ; its taste is sharp, saline, and astringent ; it first reddens and then bleaches turmeric paper. When exposed to the air it gradually evolves chlorine, and crystals of carbonate of soda are formed ; its disinfecting property depends upon this gradual escape of the chlorine.

Tests.—See Notes : SODÆ CHLORINATÆ LIQUOR.

PRÆPARATA È ZINCO.

Preparations of Zinc.

ZINCI SULPHAS.

Sulphate of Zinc.

Sal Vitrioli, P.L. 1720, P.L. 1745.

Zincum Vitriolatum, P.L. 1788.

Zinci Sulphas, P.L. 1809, P.L. 1824.

Take of Zinc, in small pieces, five ounces,

Diluted Sulphuric Acid two pints ;

Pour gradually the diluted Sulphuric Acid upon the pieces of Zinc, and the effervescence being finished, strain the Liquor ; then boil it down until a pellicle begins to form. Lastly, set it aside that crystals may be formed.

Remarks.—Zinc is a white metal with a tint of blue ; it has considerable lustre, and its specific gravity is about 7. It is crystalline in its structure, and hard, but when rolled at a temperature between 210° and 300° , it becomes malleable and ductile. It melts at 773° , and when slowly cooled crystallizes in prisms. When strongly heated in close vessels it sublimes,

but when in the air it combines with oxygen and burns rapidly, yielding a white oxide composed of


One equivalent of Oxygen	8	or	80
One equivalent of Zinc	32	,,	20
	<hr/>		
Equivalent	40		100

Symbols,—Berzelius and Turner. $\text{Zn} + \text{O}$, or Zn .

Brande $(\text{zn} + \text{o}, \text{or } \text{Zn})$.

Process.—The phenomena and effects which are produced during the solution of zinc in sulphuric acid are precisely analogous to those which occur during the solution of iron in the same acid.


Concentrated sulphuric acid does not act upon zinc at common temperatures, but when water is added, it is decomposed; the oxygen combines with the zinc to form oxide, similar to that procured by combustion, which is dissolved by the acid, and the hydrogen is evolved in the state of gas.



Properties.—The solution of sulphate of zinc is colourless, and by evaporation it readily yields crystals, which are also devoid of colour; the primary form of this salt is a *right rhombic prism*.

It may be cleaved parallel to the plane *h* of the annexed figure; no distinct cleavages have been observed in any other direction.

M on M'	91°	7'
M on <i>f</i>	135	33
M on <i>h</i>	134	27
M on <i>e</i>	128	58
<i>a</i> on <i>f</i>	120	0
<i>h</i> on <i>c</i>	119	23



Properties.—The crystals of this salt are colourless, usually very small, and not readily by appearance distinguishable from those of sulphate of magnesia; sulphate of zinc has a disagreeable metallic taste; it is not altered by exposure to the air, but if moderately heated loses its water of crystallization, and when it is subjected to a high temperature is entirely decomposed, the acid being expelled, and the oxide only remaining; it is soluble in two and a half times its weight of water at 60° , and much more so in boiling water. The alkalis, ammonia, potash, and soda, decompose the solution, and give a white precipitate; but if they are used in excess, then the precipitate is redissolved; the alkaline carbonates throw down white carbonate of zinc; water impregnated with hydrosulphuric acid decomposes the solution, and forms a white precipitate.

Composition.—Sulphate of zinc is composed of

One equivalent of Sulphuric Acid ..	40	or	28
One equivalent of Oxide of Zinc.....	40	„	28
Seven equivalents of Water.....	63	„	44
	<hr/>		
Equivalent	143		100

Symbols.—Berzelius and Turner .. $\text{Zn}\ddot{\text{S}} + 7\text{H}$.

Brande

$$(\text{Z}_\text{N} + \text{S} + 7\text{q}).$$

Impurities and Tests.—See Notes : ZINCI SULPHAS.

Incompatibles.—Alkalis and their carbonates, lime-water, and astringent vegetable infusions.

Officinal Preparations.—Liquor Aluminis Compositus. Zinci Oxydum.

Medicinal Uses.—Internally as a tonic and astringent. Dose, gr. i. to gr. ij., which may be gradually increased to gr. v. or gr. vi. without exciting nausea. It operates quickly as an emetic, in doses of gr. x. to gr. xxx. Externally it is employed as an astringent, as a substitute for the preparations of lead, in the proportion of gr. x. to eight fluidounces of water.

ZINCI OXYDUM.

Oxide of Zinc.

Zincum Calcinatum, P.L. 1788.

Zinci Oxydum, P.L. 1809, P.L. 1824.

Take of Sulphate of Zinc a pound,

Sesquicarbonate of Ammonia six ounces and
a half,

Distilled Water three gallons;

Dissolve the Sulphate of Zinc and Sesquicarbonate of Ammonia, separately, in twelve pints of the distilled Water and strain; then mix. Wash what is precipitated frequently with water; and lastly, burn it for two hours in a strong fire.

Process.—This is a case of double elective affinity, producing double decomposition; the sulphuric acid of the sulphate of zinc takes the ammonia of the sesquicarbonate, and sulphate of ammonia is formed, which remains in solution; the greater part of the carbonic acid of the sesquicarbonate of ammonia unites with the oxide of zinc, and the resulting carbonate of zinc is precipitated in the state of a white powder. This, when dried and ignited, loses its carbonic acid, and oxide of zinc remains.

Properties.—This oxide is of a yellowish white colour; it is inodorous, insipid, insoluble in water, but readily taken up by acids in general, and the alkalis ammonia, potash, and soda, but not by their carbonates.

Composition, has been already stated.

Tests and Impurities.—See Notes: ZINCI OXYDUM.

Incompatibles.—This oxide is of course incompatible with the alkalis, acids, and acidulous salts.

Official Preparation.—Unguentum Zinci.

Medicinal Use.—Tonic. Dose, gr. j. to gr. vj. twice a day in the form of pill.

CALAMINA PRÆPARATA.

Prepared Calamine.

Calamina Præparata, P.L. 1788, P.L. 1809, P.L. 1824.

Burn the Calamine ; then bruise it. Afterwards let it be made into a very fine powder in the same manner as we have directed Chalk to be prepared.

Remarks.—Calamine is a native carbonate of zinc which occurs plentifully in some parts of England ; it is usually, however, impure, being mixed with sesquioxide of iron and earthy matter. It is sometimes externally applied in excoriations, and is the basis of the Ceratum Calaminæ.

MISTURÆ.

Mixtures.

Remarks.—The term *mixture* was originally employed in pharmacy to denote those preparations, in which a soluble substance forming a viscid solution with water, was used to suspend an insoluble one ; as when gum arabic is dissolved for the purpose of holding chalk in mechanical mixture : there are a few of the preparations now classed as mixtures which are scarcely included in this definition ; and, in prescribing, the word *mixture* is frequently used to signify a compound, all the ingredients of which are in perfect solution.

MISTURA ACACIÆ.

Mixture of Acacia.

Mucilago Arabici Gummi, P.L. 1788.

Mucilago Acaciæ, P.L. 1809, P.L. 1824.

Take of Acacia, powdered, ten ounces,

Boiling Water a pint ;

Rub the Acacia with the Water gradually poured in, and dissolve it.

MISTURA AMMONIACI.

Mixture of Ammoniacum.

Mistura Ammoniaci, P.L. 1809, P.L. 1824.

Take of Ammoniacum five drachms,

Water a pint ;

Rub the Ammoniac with the Water gradually poured in, until they are perfectly mixed.

Medicinal Use.—Expectorant. Dose, fʒss. to fʒj. It may be advantageously combined with tincture of squills, and more so than with the vinegar of the same medicine, for it is slightly curdled by acids. In this mixture the resinous and insoluble matter of the ammoniacum is suspended by the solution of its gummy constituent.

MISTURA AMYGDALÆ.

Mixture of Almond.

Emulsio communis, P.L. 1745.

Lac Amygdalæ, P.L. 1788.

Mistura Amygdalæ, P.L. 1809.

Mistura Amygdalarum, P.L. 1824.

Take of Confection of Almond two ounces and a half,

Distilled Water a pint ;

Add the Water to the Confection of Almond gradually while rubbing them, until they are mixed ; afterwards strain through linen.

Medicinal Use.—Demulcent and diluent. It is generally employed as a vehicle for more active medicines. Acids, spirit of wine, and of course, tinctures, render this preparation turbid, and should not be exhibited with it.

MISTURA ASSAFÆTIDÆ.

Mixture of Assafœtida.

Lac Assafœtidæ, P.L. 1788.

Mistura Assafœtidæ, P.L. 1809, P.L. 1824.

Take of Assafœtida five drachms,

Water a pint ;

Rub the Assafœtida with the Water, gradually poured in, until they are perfectly mixed.

Medicinal Use.—Antispasmodic. Dose, from fʒss. to fʒj. repeated at short intervals in hysteric paroxysms. As it is extremely nauseous, it is rarely used, except as an enema in worms, and the convulsions of infants, arising from irritation of the bowels during dentition.

MISTURA CAMPHORÆ.

Mixture of Camphor.

Julepum e Camphorâ, P.L. 1745.

Mistura Camphorata, P.L. 1788.

Mistura Camphoræ, P.L. 1809, P.L. 1824.

Take of Camphor half a drachm,

Rectified Spirit ten minims,

Water a pint ;

First rub the Camphor with the Spirit, then with the Water gradually poured in, and strain through linen.

Medicinal Use.—Stimulant. Dose, fʒj. to fʒij. every three or four hours. Water dissolves very little camphor ; this mixture is therefore generally used only as a vehicle for more important medicines.

MISTURA CASCARILLÆ COMPOSITA.

Compound Mixture of Cascarilla.

Take of Infusion of Cascarilla seventeen fluidounces,
Vinegar of Squill a fluidounce,
Compound Tincture of Camphor two fluid-
ounces ;

Mix.

Medicinal Uses.—A combination which is of use in chronic affections of the mucous membrane of the lungs. Dose from f ʒj. to f ʒiss. twice or thrice a day.

MISTURA CRETÆ.

Mixture of Chalk.

Julepum e Cretâ, P.L. 1745.

Mistura Cretacea, P.L. 1788.

Mistura Cretæ, P.L. 1809, P.L. 1824.

Take of prepared Chalk half an ounce,
Sugar three drachms,
Mixture of Acacia a fluidounce and a half,
Cinnamon Water eighteen fluidounces ;

Mix.

Medicinal Use.—Antacid in diarrhœa. Dose, f ʒj. to f ʒij. every three or four hours ; its utility is increased when it is combined with opium, catechu, or any other astringent. It is of course incompatible with every acid and acidulous salt.

MISTURA FERRI COMPOSITA.

Compound Mixture of Iron.

Mistura Ferri Composita, P.L. 1809, P.L. 1824.

Take of Myrrh, powdered, two drachms,
Carbonate of Potash a drachm,
Rose Water eighteen fluidounces,
Sulphate of Iron, powdered, two scruples and
a half,
Spirit of Nutmeg a fluidounce,
Sugar two drachms;

Rub together the Myrrh with the Spirit of Nutmeg and the Carbonate of Potash, and to these, while rubbing, add first the Rose Water with the Sugar, then the Sulphate of Iron. Put the mixture immediately into a proper glass vessel, and stop it.

Process.—In this preparation double decomposition takes place, precisely as when sulphate of iron is decomposed in preparing the Ferri Sesquioxylum; except that, carbonate of potash being used in this case, sulphate of potash is formed instead of sulphate of soda.

Qualities.—This preparation contains protocarbonate of iron in a state of suspension. Iron in this form is probably more active than when it has become sesquioxide, being then difficultly soluble. This mixture has at first a greenish colour, but the protocarbonate of iron, to which that is owing, very readily absorbs oxygen from the air, and becomes reddish yellow sesquioxide.

Mistura Ferri Composita should not be made long before it is wanted for use; for not only is its efficacy diminished by keeping, but, from the different appearances which it presents when recently prepared, to those it exhibits when long kept, the patient would naturally suppose that some mistake had occurred in preparing it.

Medicinal Uses.—Astringent. Tonic. Dose, f ʒi. to f ʒij. two or three times a day. It is especially recommended in hysteria and chlorosis, and is unquestionably one of the most efficacious preparations of iron.

Incompatibles.—Acids and acidulous salts, which dissolve the protocarbonate of iron. Vegetable astringents render it black, and are therefore incompatible with it.

MISTURA GENTIANÆ COMPOSITA.

Compound Mixture of Gentian.

Take of Compound Infusion of Gentian twelve fluid-
ounces,

Compound Infusion of Senna six fluidounces,

Compound Tincture of Cardamom two fluid-
ounces ;

Mix.

Medicinal Uses.—Usefully employed in dyspeptic affections accompanied with constipation.

MISTURA GUAIACI.

Mixture of Guaiacum.

Lac Guaiaci, P.L. 1788.

Mistura Guaiaci, P.L. 1809, P.L. 1824.

Take of Resin of Guaiacum three drachms,

Sugar half an ounce,

Mixture of Acacia half a fluidounce,

Cinnamon Water nineteen fluidounces ;

Rub the Guaiacum with the Sugar, then with the Mixture of Acacia, and to these, while rubbing, add gradually the Cinnamon Water.

Medicinal Uses.—Stimulant. Diaphoretic. Dose, f ʒss. to f ʒij. two or three times a day.

MISTURA MOSCHI.

Mixture of Musk.

Julepum e Moschá, P.L. 1745.

Mistura Moschata, P.L. 1788.

Mistura Moschi, P.L. 1809, P.L. 1824.

Take of Musk,

Acacia, powdered,

Sugar of each three drachms,

Rose Water a pint;

Rub the Musk with the Sugar, then with the Acacia,
the Rose Water being gradually poured in.

Medicinal Use.--Antispasmodic. Dose, f ℥i. to f ℥ij.

MISTURA SPIRITÛS VINI GALLICI.

Mixture of Spirit of French Wine [Brandy].

Take of Spirit of French Wine [Brandy],

Cinnamon Water, of each four fluidounces,

The yelks of two Eggs,

Purified Sugar half an ounce,

Oil of Cinnamon two drops;

Mix.

Medicinal Uses.--Stimulant and restorative, and as such employed in the last stage of fever.

OLEA DESTILLATA.

Distilled Oils.

OLEUM ANISI.

Oil of Anise.

Oleum e seminibus Anisi, P.L. 1745.

Oleum essentielle Anisi, P.L. 1788.

Oleum Anisi, P.L. 1809, P.L. 1824.

OLEUM ANTHEMIDIS.

Oil of Chamomile.

Oleum Chamamalinum, P.L. 1720.

Oleum e floribus Chamameli, P.L. 1745.

Oleum Anthemidis, P.L. 1788, P.L. 1809, P.L. 1824.

OLEUM CARUI.

Oil of Carraway.

Oleum essentielle e seminibus Carui, P.L. 1745.

Oleum essentielle Carui, P.L. 1788.

Oleum Carui, P.L. 1809, P.L. 1824.

OLEUM JUNIPERI.

Oil of Juniper.

Oleumessentiale e baccis Juniperi, P.L. 1720, P.L. 1745.

Oleum essentielle Juniperi baccæ, P.L. 1788.

Oleum Juniperi, P.L. 1809, P.L. 1824.

OLEUM LAVANDULÆ.

Oil of Lavender.

Oleum essentielle e floribus Lavendulæ, P.L. 1745.

Oleum essentielle Lavendulæ, P.L. 1788.

Oleum Lavandulæ, P.L. 1809, P.L. 1824.

OLEUM MENTHÆ PIPERITÆ.

Oil of Peppermint.

Oleum essentielle e foliis Menthæ piperitidis, P.L. 1745.

Oleum essentielle Menthæ piperitidis, P.L. 1788.

Oleum Menthæ piperitæ, P.L. 1809, P.L. 1824.

OLEUM MENTHÆ PULEGII.

Oil of Pennyroyal.

Oleum essentielle e foliis Pulegii, P.L. 1745.

Oleum essentielle Pulegii, P.L. 1788.

Oleum Pulegii, P.L. 1809, P.L. 1824.

OLEUM MENTHÆ VIRIDIS.

Oil of Spearmint.

Oleum essenziale e foliis Menthæ vulgaris, P.L. 1745.

Oleum essenziale Menthæ sativæ, P.L. 1788.

Oleum Menthæ viridis, P.L. 1809, P.L. 1824.

OLEUM ORIGANI.

Oil of Marjoram.

Oleum essenziale e foliis Origani, P.L. 1745.

Oleum essenziale Origani, P.L. 1788.

Oleum Origani, P.L. 1809, P.L. 1824.

OLEUM PIMENTÆ.

Oil of Pimenta.

Oleum Pimentæ, P.L. 1809, P.L. 1824.

OLEUM ROSMARINI.

Oil of Rosemary.

Oleum essenziale e foliis Roris marini, P.L. 1745.

Oleum essenziale Roris marini, P.L. 1788.

Oleum Rosmarini, P.L. 1809, P.L. 1824.

OLEUM SAMBUCI.

Oil of Elder Flowers.

The Fruit of Anise, Carraway, and Juniper, the Flowers of Chamomile, Lavender, and Elder, the Berries of Pimenta, the tops of Rosemary, and the entire and fresh Herbs of the rest should be employed.

Put any one of these into an alembic, and add as much Water as is sufficient to cover it, then let the Oil distil into a large cold vessel.

Remarks.—Distilled Oils are frequently called volatile, essential, or æthereal oils. Their chemical characters are nearly the same from whatever vegetables they are procured; but in their sensible qualities they vary considerably, possessing different colours, consistence, smell, and taste; the two latter properties are, of course, derived from that of the plant from which they are obtained; their colours, like those of the fluid fixed oils, are various shades of yellow, green, and brown: they are generally fluid; but some of them, as especially oil of anise, congeal by a very moderate reduction of temperature. They are very sparingly soluble in water, but sufficiently so to impart their smell and flavour to it. They are very readily dissolved by spirit of wine, and they boil at different temperatures. Their volatility is much increased by the presence of water, with the vapour of which they rise in distillation, at a temperature considerably below their boiling point. They are extremely combustible, and much more so than the expressed oils. Most of them are lighter than water, but some sink in that fluid: among the former are the oils of lavender, rosemary, and mint; and of the latter, the oils of cassia, cinnamon, and cloves are examples. They are easily decomposed by sulphuric and by nitric acid, and when suddenly mixed with the latter, some of them inflame.

Like the expressed oils, they are composed of different proportions of oxygen, hydrogen, and carbon.

The volatile oils are capable of dissolving the fixed oils, and hence the latter are sometimes employed in adulterating them: this fraud may be easily detected by dropping some of the suspected oil on paper: if there be any fixed oil mixed with it, it

will remain on the paper after exposure to a moderate heat. Where a cheaper volatile oil has been employed to adulterate a more costly one, the detection can scarcely be made by any other means than by the difference of odour. If spirit of wine be mixed with the oil, then, when it is dropped upon water, a milky fluid is formed, instead of there remaining a transparent film of oil on the surface of the water.

OLEUM SUCCINI.

Oil of Amber.

Oleum Succini Rectificatum, P.L. 1788.

Oleum Succini, P.L. 1720, P.L. 1745, P.L. 1809, P.L. 1824.

Put Amber into an alembic, so that an Acid Liquor, an Oil, and a Salt, contaminated with the Oil, may distil in a sand-bath, with a heat gradually increased. Afterwards, let the Oil distil again and a third time.

OLEUM TEREBINTHINÆ PURIFICATUM.

Purified Oil of Turpentine.

Oleum sive Spiritus Terebinthinæ, P.L. 1720.

Oleum Terebinthinæ Æthereum, P.L. 1745.

Oleum Terebinthinæ Rectificatum, P.L. 1788,
P.L. 1809, P.L. 1824.

Take of Oil of Turpentine a pint,

Water four pints;

Let the Oil carefully distil.

PILULÆ.

Pills.

PILULÆ ALOËS COMPOSITÆ.

Compound Pills of Aloes.

Pilulæ Aloës Compositæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Aloes, powdered, an ounce,
Extract of Gentian half an ounce,
Oil of Carraway forty minims,
Syrup as much as may be sufficient;
Beat them together until incorporated.

Medicinal Use.—Purgative. Stomachic, in habitual costiveness. Dose, gr. x. to gr. xx.

PILULÆ ALOËS CUM MYRRHA.

Pills of Aloes with Myrrh.

Pilulæ Ruffi, P.L. 1720.

Pilulæ Rufi, P.L. 1745.

Pilulæ Aloës cum Myrrha, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Aloes two ounces,
Saffron,
Myrrh, each an ounce,
Syrup as much as may be sufficient;
Rub the Aloes and Myrrh separately to powder; then
beat the whole together until incorporated.

Medicinal Use.—This preparation is yet commonly called *Pilulæ Rufi*, and has been very long in use. Dose, gr. x. to gr. xx. as a stimulant and cathartic.

PILULÆ CAMBOGIÆ COMPOSITÆ.

Compound Pills of Camboge.

Pilulæ Cambogiæ compositæ, P.L. 1809, P.L. 1824.

Take of Camboge, powdered, a drachm,
Aloes, powdered, a drachm and a half,
Ginger, powdered, half a drachm,
Soap two drachms ;

Mix the powders together ; afterwards, the Soap being added, beat the whole together until incorporated.

Medicinal Use.—Cathartic. Dose, gr. x. to gr. xx.

PILULÆ CONII COMPOSITÆ.

Compound Pills of Hemlock.

Take of Extract of Hemlock five drachms,
Ipecacuanha, powdered, a drachm,
Mixture of Acacia as much as may be sufficient ;

Beat them together until incorporated.

Medicinal Use.—Antispasmodic and slightly narcotic. Of use in hooping-cough and incipient stage of phthisis : gr. v. three times a day.

PILULÆ FERRI COMPOSITÆ.

Compound Pills of Iron.

Pilulæ Ferri cum Myrrhâ, P.L. 1809.

Pilulæ Ferri Compositæ, P.L. 1824.

Take of Myrrh, powdered, two drachms,
Carbonate of Soda,
Sulphate of Iron,
Treacle, each a drachm ;

Rub the Myrrh with the Carbonate of Soda ; then, the Sulphate of Iron being added, rub them again ; afterwards beat the whole in a vessel previously warmed until incorporated.

Remarks.—In this preparation the sulphate of iron is decomposed by the carbonate of soda, precisely in the same manner, and in the first instance, with the production of similar compounds as in preparing the *Ferri Sesquioxidum*. While, however, the sulphate of soda is washed away from the sesquioxide of iron, it remains with it in preparing the pills, but the quantity is so extremely small as to be quite unimportant. Nearly the same precautions as those which have been given with respect to the *Mistura Ferri Composita*, will apply to this preparation ; viz. that the pills should be prepared only at the moment in which they are wanted, for the protocarbonate of iron at first formed is very readily converted into sesquioxide by absorbing the oxygen of the atmosphere, by which its solubility and power are diminished. The dose is from gr. x. to gr. xx. two or three times a day, in the same cases as the *Mistura Ferri Composita*.

PILULÆ GALBANI COMPOSITÆ.

Compound Pills of Galbanum.

Pilulæ Gummosæ, P.L. 1720, P.L. 1745.

Pilulæ Galbani Compositæ, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Galbanum an ounce,
Myrrh,
Sagapenum, each an ounce and a half,
Assafoetida half an ounce,
Syrup as much as may be sufficient ;
Beat them together until incorporated.

Medicinal Use.—Antispasmodic and emmenagogue. Dose,
gr. x. to gr. xx.

PILULÆ HYDRARGYRI.

Pills of Mercury.

Pilulæ Mercuriales, P.L. 1745.

Pilulæ Hydrargyri, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Mercury two drachms,
Confection of Red Rose three drachms,
Liquorice, powdered, a drachm ;
Rub the Mercury with the Confection, until globules
can no longer be seen ; then, the Liquorice being added,
beat the whole together until incorporated.

Process.—The mercury in this preparation is probably in the
state of minute division only.

Medicinal Uses.—It is by far the best form for the internal
exhibition of mercury ; when it is intended to act upon the system
as an alterative, it should be administered in doses of from gr. iv.
to gr. vj. Opium may be advantageously given with it, if it
should occasion irritation. In doses, from gr. x. to gr. xx. it
acts as a mild but efficient purgative.

PILULÆ HYDRARGYRI CHLORIDI COM-
POSITÆ.

Compound Pills of Chloride of Mercury.

Pilulæ Hydrargyri Submuriatis compositæ,
P.L. 1809, P.L. 1824.

Take of Chloride of Mercury,

Oxysulphuret of Antimony, each two drachms,
Guaiacum Resin, powdered, half an ounce,
Treacle two drachms;

Rub the Chloride of Mercury with the Oxysulphuret
of Antimony, afterwards with the Guaiacum Resin and
the Treacle until incorporated.

Medicinal Uses.—Alterative. Dose, gr. v. to gr. x. This
pill is much employed in cutaneous eruptions, and in secondary
syphilitic symptoms, particularly when affecting the skin. It is
commonly known by the name of *Plummer's Pill*.

PILULÆ HYDRARGYRI IODIDI.

Pills of Iodide of Mercury.

Take of Iodide of Mercury a drachm,

Confection of Dog Rose three drachms,
Ginger, powdered, a drachm;

Beat them together until incorporated.

Medicinal Uses.—See HYDRARGYRI IODIDUM.

PILULÆ IPECACUANHÆ COMPOSITÆ.

Compound Pills of Ipecacuanha.

Take of Compound Powder of Ipecacuanha three drachms,

Fresh dried Squill,

Ammoniacum, of each a drachm,

Mixture of Acacia as much as may be sufficient ;

Beat them together until incorporated.

Medicinal Uses.—Sudorific and narcotic. Dose, gr. v. three times a day, or gr. x. at night.

PILULÆ RHEI COMPOSITÆ.

Compound Pills of Rhubarb.

Take of Rhubarb, powdered, an ounce,

Aloes, powdered, six drachms,

Myrrh, powdered, half an ounce,

Soap a drachm,

Oil of Carraway half a fluidrachm,

Syrup as much as may be sufficient ;

Mix the Powders, then beat the whole together until incorporated.

Medicinal Use.—Slightly aperient or laxative.

PILULÆ SAGAPENI COMPOSITÆ.

Compound Pills of Sagapenum.

Take of Sagapenum an ounce,
Aloes half a drachm,
Syrup of Ginger as much as may be sufficient;
Beat them together until incorporated.

Medicinal Uses.—Antibilious and laxative. Successfully employed in colic, induced by sedentary occupations. Dose, gr. x.

PILULÆ SAPONIS COMPOSITÆ.

Compound Pills of Soap.

Laudanum, P.L. 1720.
Pilulæ Saponaceæ, P.L. 1745.
Pilulæ Opii, P.L. 1788.
Pilulæ Saponis cum Opio, P.L. 1809, P.L. 1824.

Take of Hard Opium, powdered, half an ounce,
Soap two ounces;
Beat them together until incorporated.

Medicinal Uses.—Anodyne. Narcotic. Dose, gr. iij. to gr. x.
Five grains contain one grain of opium.

PILULÆ SCILLÆ COMPOSITÆ.

Compound Pills of Squill.

Pilulæ Scillæ, P.L. 1788.

Pilulæ Scillæ Compositæ, P.L. 1809, P.L. 1824.

Take of Squill, fresh dried and powdered, a drachm,
Ginger, powdered,
Ammoniacum, powdered, each two drachms,
Soap three drachms,
Syrup as much as may be sufficient;

Mix the Powders together; then beat them with the Soap, and add the Syrup so as to obtain a proper consistence.

Medicinal Uses.—Expectorant. Diuretic. Dose, gr. x. to gr. xx.

PILULÆ STYRACIS COMPOSITÆ.

Compound Pills of Storax.

Pilulæ e Styrace, P.L. 1745.

Take of Purified Storax three drachms,
Hard Opium, powdered,
Saffron, of each a drachm;
Beat them together until incorporated.

Medicinal Uses.—Balsamic and slightly expectorant in chronic affections of the lungs. Dose, gr. iij. to gr. x. Five grains contain one grain of opium.

PULVERES.

Powders.

PULVIS ALOËS COMPOSITUS.

Compound Powder of Aloes

Pilulæ Aromaticæ, P.L. 1745.

Pulvis Aloës cum Guaiaco, P.L. 1788.

Pulvis Aloës Compositus, P.L. 1809, P.L. 1824.

Take of Aloes an ounce and a half,
Guaiacum Resin an ounce,
Compound Powder of Cinnamon half an ounce;
Rub the Aloes and the Guaiacum Resin separately to powder; then mix them with the Compound Powder of Cinnamon.

Medicinal Uses.—This powder is cathartic and sudorific.
Dose, gr. x. to gr. xx.

PULVIS CINNAMOMI COMPOSITUS.

Compound Powder of Cinnamon.

Species Aromaticæ, P.L. 1745.

Pulvis Aromaticus, P.L. 1788.

Pulvis Cinnamomi Compositus, P.L. 1809, P.L. 1824.

Take of Cinnamon two ounces,
Cardamom an ounce and a half,
Ginger an ounce,
Long Pepper half an ounce ;

Rub them together, so that a very fine powder may be made.

Medicinal Uses.—This preparation is stimulant and carminative. Dose, gr. v. to gr. x. in the form of bolus, or mixed with water. It is generally employed to give warmth to more active remedies.

PULVIS CRETÆ COMPOSITUS.

Compound Powder of Chalk.

Diascordium, P.L. 1720.

Pulvis e Bolo compositus sine Opio, Species e Scordio sine Opio, P.L. 1745.

Pulvis Cretæ compositus, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Prepared Chalk half a pound,
Cinnamon four ounces,
Tormentil,
Acacia, each three ounces,
Long Pepper half an ounce ;

Rub them separately to very fine powder ; then mix.

Medicinal Uses.—Astringent and antacid. Dose, gr. v. to gr. xxx.

PULVIS CRETÆ COMPOSITUS CUM OPIO.

Compound Powder of Chalk with Opium.

Pulvis e Bolo compositus cum Opio, Species e Scordio cum Opio, P.L. 1745.

Pulvis Cretæ compositus cum Opio, P.L. 1788,
P.L. 1809, P.L. 1824.

Take of Compound Powder of Chalk six ounces and
a half,

Hard Opium, powdered, four scruples ;

Mix.

Medicinal Uses.—Astringent. Anodyne. Dose, gr. v. to gr. xxx.
Forty grains contain one grain of opium. This and the former
preparation, on account of the carbonate of lime which they
contain, are incompatible with acids and acidulous salts.

PULVIS JALAPÆ COMPOSITUS.

Compound Powder of Jalap.

Take of Jalap three ounces,

Bitartrate of Potash six ounces,

Ginger two drachms ;

Rub them separately to powder ; then mix.

Medicinal Use.—Purgative. Dose, gr. xx. to ʒj.

PULVIS IPECACUANHÆ COMPOSITUS.

Compound Powder of Ipecacuanha.

Pulvis Ipecacuanhæ Compositus, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Ipecacuanha, powdered,
Hard Opium, powdered, each a drachm,
Sulphate of Potash, powdered, an ounce;
Mix.

Remarks.—This powder has been long employed as a sudorific, under the name of Dover's Powder. The sulphate of potash is used merely to divide the more active ingredients. In doses of gr. v. to gr. xx. it acts as a powerful sudorific; it may be given diffused in a mucilaginous fluid, or in the form of bolus. Ten grains contain one grain of opium.

PULVIS KINO COMPOSITUS.

Compound Powder of Kino.

Pulvis Kino compositus, P.L. 1809, P.L. 1824.

Take of Kino fifteen drachms,
Cinnamon half an ounce,
Hard Opium a drachm;
Rub them separately to very fine powder; then mix.

Medicinal Use.—Astringent. Dose, gr. v. to gr. xx. Twenty grains contain one grain of Opium.

PULVIS SCAMMONII COMPOSITUS.

Compound Powder of Scammony.

Pulvis comitis Warwicensis, P.L. 1720.

Pulvis e Scammonio compositus, P.L. 1745.

Pulvis Scammonii compositus, P.L. 1788.

Pulvis Scammoneæ compositus, P.L. 1809, P.L. 1824.

Take of Scammony,
Hard Extract of Jalap, each two ounces,
Ginger half an ounce;
Rub them separately to very fine powder; then mix.

Medicinal Use.—Cathartic. Dose, gr. v. to gr. xx.

PULVIS TRAGACANTHÆ COMPOSITUS.

Compound Powder of Tragacanth.

Species Diatragacanthæ frigidæ, P.L. 1720.

Pulvis e Tragacanthâ compositus, P.L. 1745.

Pulvis Tragacanthæ compositus, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Tragacanth, powdered,

Acacia, powdered,

Starch, each an ounce and a half,

Sugar three ounces ;

Rub the Starch and Sugar together to powder ; then,
the Tragacanth and Acacia being added, mix them all.

Medicinal Use.—Demulcent. Dose, gr. x. to ʒi.

SPIRITUS.

Spirits.

Spirit of wine, or alcohol diluted with water, is employed in pharmacy for various important purposes, and of different degrees of strength, according to circumstances. In its concentrated state it is termed *alcohol* ; when diluted with a small proportion of water it is called *rectified spirit* ; and when more largely diluted, *proof spirit*. The two latter are articles of the *Materia Medica*, and the first is prepared by the process stated below.

Some of the preparations in which Spirit is used in the *Pharmacopœia*, are classed together under the title of *Spiritus* ; it includes spirit of ammonia, and several aromatic distilled spirits : *Tincturæ* and *Æthereæ* are the two other classes.

ALCOHOL.

Alcohol.

Alcohol, P.L. 1788.*Alcohol*, P.L. 1809, P.L. 1824.

Take of Rectified Spirit a gallon,

Chloride of Calcium a pound;

Put the Chloride of Calcium into the Spirit, and when it is dissolved, let seven pints and five fluidounces distil.

Process.—Chloride of calcium is a salt which, as already noticed, has great affinity for water, and is soluble in spirit; when the solution is subjected to distillation, the chloride remains in the retort with nearly the whole of the water.

The strongest spirit which has hitherto been produced is of sp. gr. 0·796, at the temperature of 60°: and it is, probably, alcohol free from water; according to Saussure, it consists of

Three equivalents of Hydrogen ..	$1 \times 3 = 3$	or 13·04
Two equivalents of Carbon	$6 \times 2 = 12$	„ 52·17
One equivalent of Oxygen	$= 8$	„ 34·79

Equivalent....	23	100·
----------------	----	------

Properties.—Alcohol, when pure, is colourless and transparent; its odour is rather pleasant, and its taste is penetrating. It has never been rendered solid by exposure to any degree of cold, either natural or artificial. Alcohol is that part of fermented liquors from which their intoxicating power is derived. It is extremely volatile, producing great cold during its evaporation; and the stronger the alcohol the greater is the cold produced. It is highly inflammable, and during combustion, water and carbonic acid are generated, the quantity of the former exceeding that of the weight of alcohol burned.

Alcohol of sp. gr. 0·800 boils at 174°, or 38° below the boiling point of water, and it is very expansible by heat. When it is mixed with water, heat is evolved, the capacity of the compound being less than that of its ingredients; and the mixture occupies considerably less space than the water and alcohol do when separate.

Alcohol prevents animal substances which are immersed in it from decay; and hence its use in the preservation of anatomical preparations. Its solvent power is very great, and it is on this

account that it is in many cases employed in pharmacy, especially in the preparation of the tinctures of those substances which are resinous, and insoluble in water. It is also largely employed in the preparation of æther.

SPIRITUS AMMONIÆ.

Spirit of Ammonia.

Spiritus Salis Ammoniaci, P.L. 1720.

Spiritus Salis Ammoniaci dulcis, P.L. 1745.

Spiritus Ammoniacæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Hydrochlorate of Ammonia ten ounces,
Carbonate of Potash sixteen ounces,
Rectified Spirit,
Water, each, three pints;
Mix, and let three pints distil.

Process.—In this operation the hydrochlorate of ammonia is decomposed, as already described when treating of the Ammoniacæ Sesquicarbonas; in the present case, however, chloride of potassium remains instead of chloride of calcium, because carbonate of potash is employed instead of carbonate of lime, and the carbonate of ammonia is a neutral, instead of a sesquicarbonate. This is also the case with the Spiritus Ammoniacæ Aromaticus and Spiritus Ammoniacæ Foetidus. It is composed of

One equivalent of Carbonic Acid = 22 or 56.5

One equivalent of Ammonia = 17 „ 43.5

Equivalent	39	100.
--------------------	----	------

As this carbonate contains only two thirds as much carbonic acid as that procured by the use of carbonate of lime, the greater pungency of Spiritus Ammoniacæ and Spiritus Ammoniacæ Aromaticus, than of Liquor Ammoniacæ Sesquicarbonatis, is readily accounted for.

Properties.—Spiritus Ammoniacæ is a transparent colourless fluid; its smell is pungent and its taste acrid; it turns turmeric brown, indicating its alkaline properties. There is usually more carbonate of ammonia formed than the spirit is capable of dissolving, and this remains in the receiver in an imperfectly crystalline state.

SPIRITUS AMMONIÆ AROMATICUS.

Aromatic Spirit of Ammonia.

Spiritus Salis Volatilis oleosus, P.L. 1720.

Spiritus Volatilis Aromaticus, P.L. 1745.

Spiritus Ammoniæ compositus, P.L. 1788.

Spiritus Ammoniæ Aromaticus, P.L. 1809, P.L. 1824.

Take of Hydrochlorate of Ammonia five ounces,
Carbonate of Potash, eight ounces,
Cinnamon, bruised,
Cloves, bruised, each two drachms,
Lemon Peel four ounces,
Rectified Spirit,
Water, each four pints ;

Mix, and let six pints distil.

Properties.—This preparation resembles the last, but is rendered more agreeable by the aromatics, whether applied to the nostrils or internally exhibited.

Incompatibles.—Acids, acidulous salts, earthy and metallic salts, and lime-water.

Officinal Preparations.—Tinctura Guaiaci composita, Tinctura Valerianæ composita.

Medicinal Use.—Stimulant in languors and flatulent colic.
Dose, f 3ss. to f 3j. in water.

SPIRITUS AMMONIÆ FÆTIDUS.

Fetid Spirit of Ammonia.

Spiritus Volatilis fætidus, P.L. 1745.

Spiritus Ammoniæ fætidus, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Hydrochlorate of Ammonia ten ounces,
Carbonate of Potash sixteen ounces,
Rectified Spirit,
Water, each three pints,
Assafoetida five ounces ;
Mix ; then with a slow fire let three pints distil.

Properties.—Colourless, pungent, and, as its name expresses, fetid. By long keeping it acquires a brownish colour.

Incompatibles.—The same as with the last preparations.

Medicinal Uses.—Stimulant. Antispasmodic. Dose, f ʒss. or more in water.

SPIRITUS ANISI.

Spirit of Aniseed.

Aqua seminum Anisi composita, P.L. 1745.

Spiritus Anisi compositus, P.L. 1788.

Spiritus Anisi, P.L. 1809, P.L. 1824.

Take of Aniseed, bruised, ten ounces,
Proof Spirit a gallon,
Water two pints ;
Mix ; then with a slow fire let a gallon distil.

Medicinal Uses.—Stimulant and carminative in flatulent colic, &c. Dose, f ʒij. to f ʒ iv. in water.

SPIRITUS ARMORACIÆ COMPOSITUS.

Compound Spirit of Horseradish.

Aqua Raphani composita, P.L. 1720, P.L. 1745.

Spiritus Raphani compositus, P.L. 1788.

Spiritus Armoraciæ compositus, P.L. 1809, P.L. 1824.

Take of Horseradish, sliced,
Dried Orange Peel, each twenty ounces,
Nutmeg, bruised, five drachms,
Proof Spirit a gallon,
Water two pints;

Mix; then with a slow fire let a gallon distil.

Medicinal Uses.—Stimulant. Dose, f ʒij. to f ʒiv.

SPIRITUS CARUI.

Spirit of Carraway.

Aqua Seminum Carui, P.L. 1745.

Spiritus Carui, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Carraway, bruised, twenty-two ounces,
Proof Spirit a gallon,
Water two pints;

Mix; then with a slow fire let a gallon distil.

Medicinal Uses.—Carminative. Stimulant. Dose, f ʒij. to f ʒiv.

SPIRITUS CINNAMOMI.

Spirit of Cinnamon.

Aqua Cinnamomi fortis, P.L. 1720.

Aqua Cinnamomi spiritiosa, P.L. 1745.

Spiritus Cinnamomi, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Oil of Cinnamon two drachms,
Proof Spirit a gallon,
Water a pint;

Mix; then with a slow fire let a gallon distil.

Medicinal Uses.—Stomachic. Stimulant. Dose, f ʒij. to f ʒiv.

SPIRITUS JUNIPERI COMPOSITUS.

Compound Spirit of Juniper.

Aqua Juniperi composita, P.L. 1745.

Spiritus Juniperi compositus, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Juniper Fruit, bruised, fifteen ounces,
Carraway, bruised,
Fennel, bruised, each two ounces,
Proof Spirit a gallon,
Water two pints ;

Mix ; then with a slow fire let a gallon distil.

Medicinal Uses.—Stimulant. Diuretic. Dose, f ʒiij. to f ʒj.
It is principally exhibited with other diuretics, as foxglove, &c.

SPIRITUS LAVANDULÆ.

Spirit of Lavender.

Spiritus Lavendulæ simplex, P.L. 1745.

Spiritus Lavendulæ, P.L. 1788.

Spiritus Lavandulæ, P.L. 1809, P.L. 1824.

Take of fresh Lavender two pounds and a half,
Rectified Spirit a gallon,
Water two pints ;

Mix ; then with a slow fire let a gallon distil.

Pharmaceutical Uses.—In preparing the Tinctura Lavandulæ Composita and Linimentum Camphoræ Compositum.

SPIRITUS MENTHÆ PIPERITÆ.

Spirit of Peppermint.

Aqua Menthæ piperitidis spirituosæ, P.L. 1745.

Spiritus Menthæ piperitidis, P.L. 1788.

Spiritus Menthæ piperitæ, P.L. 1809, P.L. 1824.

Take of Oil of Peppermint three drachms,

Proof Spirit a gallon,

Water a pint ;

Mix ; then with a slow fire let a gallon distil.

Medicinal Uses.—Stimulant. Carminative. Dose, fʒij. to fʒj.

SPIRITUS MENTHÆ VIRIDIS.

Spirit of Spearmint.

Aqua Menthæ vulgaris spirituosæ, P.L. 1745.

Spiritus Menthæ sativæ, P.L. 1788.

Spiritus Menthæ viridis, P.L. 1809, P.L. 1824.

SPIRITUS MENTHÆ PULEGII.

Spirit of Pennyroyal.

Aqua Pulegii spirituosæ, P.L. 1745.

Spiritus Pulegii, P.L. 1788, P.L. 1809, P.L. 1824.

These are prepared in the same manner as the Spirit of Peppermint.

SPIRITUS MYRISTICÆ.

Spirit of Nutmeg.

Aqua ucis Moschatæ, P.L. 1745.

Spiritus Myristicæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Nutmegs, bruised, two ounces and a half,

Proof Spirit a gallon,

Water a pint ;

Mix ; then with a slow fire let a gallon distil.

SPIRITUS PIMENTÆ.

Spirit of Pimenta.

Spiritus Pimento, P.L. 1788.

Spiritus Pimentæ, P.L. 1809, P.L. 1824.

This is prepared in the same manner as the Spirit of Nutmeg.

The four last preparations are employed in similar doses, and with the same intention as Spirit of Peppermint.

SPIRITUS ROSMARINI.

Spirit of Rosemary.

Spiritus Rosmarini, P.L. 1745, P.L. 1788.

Spiritus Rosmarini, P.L. 1809, P.L. 1824.

Take of Oil of Rosemary two drachms,

Rectified Spirit a gallon,

Water a pint ;

Mix ; then with a slow fire let a gallon distil.

This is employed in the following officinal preparations :—
Tinctura Lavandulæ Composita, and *Linimentum Saponis Compositum*.

SYRUPI.Syrups.

Syrups are strong solutions of sugar in water, generally coloured or flavoured with vegetable matter; and sometimes, but more rarely, they are active medicines; it is particularly requisite that they should be kept in a cool place, or otherwise acetic acid will be generated by fermentation, and this may interfere with medicine, the virtues of which it is employed to increase, or whose disagreeable flavour it is intended to disguise.

Syrups are to be kept in a place where the heat never exceeds 55°.

SYRUPUS.

Syrup.

Syrupus Simplex, P.L. 1745.

Syrupus, P.L. 1788, P.L. 1809.

Syrupus Simplex, P.L. 1824.

Take of Sugar ten pounds,

Water three pints;

Dissolve the Sugar in the Water with a gentle heat.

SYRUPUS ALTHÆÆ.

Syrup of Marshmallow.

Syrupus de Althæâ, P.L. 1720.

Syrupus ex Althæâ, P.L. 1745.

Syrupus Althææ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Marshmallow Root, bruised, eight ounces,
Sugar two pounds and a half,
Water four pints ;

Boil down the Water with the Root to half, and press out the cooled liquor. Set it by for twenty four hours, that the dregs may subside ; then pour off the liquor, and, the Sugar being added, boil down to a proper consistence.

This syrup contains the mucilaginous matter of the marshmallow, and is used as a demulcent. It is apt to spoil by fermentation, and does not possess any active property.

SYRUPUS AURANTII.

Syrup of Orange [Peel].

Syrupus de cortice Aurantiorum, P.L. 1720.

Syrupus e corticibus Aurantiorum, P.L. 1745.

Syrupus corticis Aurantii, P.L. 1788.

Syrupus Aurantii, P.L. 1809.

Syrupus Aurantiorum, P.L. 1824.

Take of fresh Orange Peel two ounces and a half,
Boiling Water a pint,
Sugar three pounds ;

Macerate the Peel in the Water for twelve hours in a vessel lightly covered ; then pour off the liquor, and add the Sugar to it.

This syrup is employed merely on account of its grateful aromatic flavour.

SYRUPUS CROCI.

Syrup of Saffron.

Syrupus Croci, P.L. 1720, P.L. 1745, P.L. 1788.
P.L. 1809, P.L. 1824.

Take of Saffron ten drachms,
Boiling Water a pint,
Sugar three pounds ;

Macerate the Saffron in the Water for twelve hours in a vessel lightly covered ; then strain the liquor, and add the Sugar to it.

It is used merely on account of its fine colour.

SYRUPUS LIMONUM.

Syrup of Lemons.

Syrupus e succo Citrionum, P.L. 1720.
Syrupus e succo Limonum, P.L. 1745.
Syrupus succi Limonum, P.L. 1788.
Syrupus Limonis, P.L. 1809.
Syrupus Limonum, P.L. 1824.

Take of Lemon Juice, strained, a pint,
Sugar two pounds and a half ;

Dissolve the Sugar in the Lemon Juice, with a gentle heat, then set aside for twenty four hours ; afterwards remove the scum, and pour off the clear liquor from the dregs, if there be any.

This is a pleasant syrup ; but it must be remembered that its acidity prevents its being employed in any composition that contains alkalis, alkaline earths, or their carbonates.

SYRUPUS MORI.

Syrup of Mulberry.

Syrupus Mororum, P.L. 1745.

Syrupus Mori, P.L. 1788, P.L. 1809, P.L. 1824.

Take of the Juice of Mulberries, strained, a pint,
Sugar two pounds and a half;

Dissolve the Sugar in the Mulberry Juice with a gentle heat, and proceed in the same manner as directed for Syrup of Lemons.

This is used for the same purposes as the former, and it has the advantage of a fine colour.

SYRUPUS PAPAVERIS.

Syrup of Poppy.

Syrupus de Meconio sive Diacodium, P.L. 1720.

Syrupus de Meconio, P.L. 1745.

Syrupus Papaveris albi, P.L. 1788.

Syrupus Papaveris, P.L. 1809, P.L. 1824.

Take of Poppy [Capsules] three pounds,
Sugar five pounds,
Boiling Water five gallons;

Boil down the Capsules in the Water to two gallons, and press strongly. Boil down the strained liquor again to four pints, and strain while hot. Set it by for twelve hours that the dregs may subside; then boil down the clear liquor to two pints; add the Sugar and dissolve it.

Medicinal Use.—Anodyne. Narcotic. Dose, $f\text{ʒj.}$ to $f\text{ʒj.}$ This syrup is very apt to ferment, and hence the necessity of keeping it cool. It is principally used for children.

SYRUPUS RHAMNI.

Syrup of Buckthorn.

Syrupus de Spina cervina, P.L. 1720.

Syrupus e Spina cervina, P.L. 1745.

Syrupus Spinæ cervinæ, P.L. 1788.

Syrupus Rhamni, P.L. 1809, P.L. 1824.

Take of the fresh Juice of Buckthorn four pints,
Ginger, sliced,
Pimenta, powdered, each six drachms,
Sugar four pounds;

Set by the Juice for three days, that the dregs may subside, and strain. To a pint of the clear Juice add the Ginger and Pimenta; then macerate with a gentle heat for four hours, and strain; boil down that which is left to the measure of a pint and a half; mix the liquors; add the Sugar and dissolve it.

Medicinal Use.—Cathartic. Dose, fʒss. to fʒj. It is an unpleasant remedy both to the taste and in its operation, and is but little used.

SYRUPUS RHŒADOS.

Syrup of Red Poppy.

Syrupus de Papavere erratico, P.L. 1720.

Syrupus Papaveris erratici, P.L. 1745, P.L. 1788.

Syrupus Rhæados, P.L. 1809, P.L. 1824.

Take of Red Poppy [Petals] a pound,
Boiling Water a pint,
Sugar two pounds and a half;

Add the Petals of the Red Poppy gradually to the Water, heated in a water-bath, frequently stirring them;

then, the vessel being removed, macerate for twelve hours; afterwards press out the liquor, and when the dregs have subsided, add the Sugar and dissolve it.

This syrup is of a fine red colour, and is used only on that account.

SYRUPUS ROSÆ.

Syrup of Rose.

Syrupus e Rosis siccis, P.L. 1720.

Syrupus Rosarum solutivus, P.L. 1745.

Syrupus Rosæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Damask Rose [Petals] dried, seven ounces,
Sugar six pounds,
Boiling Water three pints;

Macerate the Rose Petals in the Water for twelve hours, and strain; evaporate the strained liquor in a water-bath to two pints; then add the Sugar, and dissolve it.

Medicinal Properties.—Purgative, but weakly so; it is sometimes given to infants. Dose, fʒij. to fʒj.

SYRUPUS SARZÆ.

Syrup of Sarsaparilla.

Syrupus Sarsaparillæ, P.L. 1824.

Take of Sarsaparilla, sliced, fifteen ounces,
Boiling Water a gallon,
Sugar fifteen ounces;

Macerate the Sarsaparilla in the Water for twenty four

hours; then boil down to four pints, and strain the liquor while hot; afterwards add the Sugar, and evaporate to a proper consistence.

This is employed as an adjunct to the decoction of Sarsaparilla.

SYRUPUS SENNÆ.

Syrup of Senna.

Syrupus Sennæ, P.L. 1809, P.L. 1824.

Take of Senna two ounces and a half,
Fennel, bruised, ten drachms,
Manna three ounces,
Sugar fifteen ounces,
Boiling Water a pint;

Macerate the Senna and Fennel in the Water with a gentle heat for an hour. Mix the Manna and Sugar with the strained liquor; then boil down to a proper consistence.

This is a purgative syrup intended for children. Dose, fʒij. to fʒiv.

SYRUPUS TOLUTANUS.

Syrup of Tolu.

Syrupus Balsamicus, P.L. 1720, P.L. 1745.

Syrupus Tolutanus, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Balsam of Tolu ten drachms,
Boiling Water a pint,
Sugar two pounds and a half;

Boil the Balsam in the Water for half an hour in a vessel lightly covered, frequently stirring, and strain the cooled liquor; then add the Sugar, and dissolve it.

It is employed merely to give a pleasant flavour to draughts and mixtures.

SYRUPUS ZINGIBERIS.

Syrup of Ginger.

Syrupus Zingiberis, P.L. 1745, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Ginger, sliced, two ounces and a half,

Boiling Water a pint,

Sugar two pounds and a half;

Macerate the Ginger in the Water for four hours, and strain: then add the Sugar, and dissolve it.

This syrup is impregnated with the flavour and warmth of the ginger, and is a useful adjunct to bitter infusions and griping purgatives.

TINCTURÆ.

Tinctures.

Tinctures are solutions of various substances in spirit of wine, of different degrees of strength; they are principally prepared from vegetable matters, but in some cases metallic salts are dissolved in it; in others the tinctures contain ammonia, and in one instance animal matter is dissolved by spirit.

The substances which are best adapted for tinctures are those which are active in small doses; for if large ones should be required, they might be in many cases objectionable on account of the quantity of spirit necessarily exhibited with them.

Those substances which are imperfectly soluble in water, or totally insoluble in it, or which spoil unless they are preserved by spirit, are proper for tinctures, provided they can be given in sufficiently large doses; opium, digitalis, &c. are bodies of this class.

Tinctures are frequently useful additions to infusions and decoctions, the spirit preventing the decomposition, which otherwise occurs rapidly. Tinctures which hold resinous matter in

solution, such as that of guaiacum, suffer decomposition on the addition of water.

All Tinctures should be prepared in stopped glass vessels, and frequently shaken during maceration.

TINCTURA ALOËS.

Tincture of Aloes.

Tinctura Aloës, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Aloes, powdered, an ounce,
Extract of Liquorice three ounces,
Water a pint and a half,
Rectified Spirit half a pint;
Macerate for fourteen days, and strain.

Medicinal Uses.—Purgative. Stomachic. Dose, fʒss. to fʒiss.

TINCTURA ALOËS COMPOSITA.

Compound Tincture of Aloes.

Elixir Proprietatis, P.L. 1720.

Elixir Aloës, P.L. 1745.

Tinctura Aloës composita, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Aloes, powdered, four ounces,
Saffron two ounces,
Tincture of Myrrh two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—Purgative. Stomachic. Dose, fʒj. to fʒij.

TINCTURA AMMONIÆ COMPOSITA.

Compound Tincture of Ammonia.

Spiritus Ammoniae Succinatus, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Mastich two drachms,
Rectified Spirit nine fluidrachms,
Oil of Lavender fourteen minims,
Oil of Amber four minims,
Stronger Solution of Ammonia a pint;

Macerate the Mastich in the Spirit, that it may be dissolved, and pour off the clear tincture; then add the other ingredients, and shake them all together.

Qualities.—This preparation has a milky appearance, owing to the separation of the mastich from its solution in spirit by the *Liquor Ammoniae*. It is commonly called *Eau de Luce*, but no oil of amber is contained in the preparation originally so denominated.

Incompatibles.—Acids; acidulous, metallic, and earthy salts.

Medicinal Uses.—Stimulant and antispasmodic. Dose, ℥v. to ℥x. in water.

TINCTURA ASSAFŒTIDÆ.

Tincture of Assafoetida.

Tinctura Fœtida, P.L. 1745.

Tinctura Asæ fœtidæ, P.L. 1788.

Tinctura Assafœtidæ, P.L. 1809, P.L. 1824.

Take of Assafoetida five ounces,
Rectified Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—Stimulant. Antispasmodic. Dose, fʒss. to fʒiiss. This tincture is rendered turbid when mixed with water, owing to the precipitation of the resinous matter of the assafoetida.

TINCTURA AURANTII.

Tincture of Orange [Peel].

Tinctura corticis Aurantii, P.L. 1788.

Tinctura Aurantii, P.L. 1809, P.L. 1824.

Take of dried Orange Peel three ounces and a half,
Proof Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—Tonic. Stomachic. Dose, fʒij. to fʒiij.
It is a useful adjunct to bitter infusions and decoctions.

TINCTURA BALSAMI TOLUTANI.

Tincture of Balsam of Tolu.

Take of Balsam of Tolu two ounces,
Rectified Spirit two pints;
Macerate until the Balsam is dissolved, and strain.

Medicinal Use.—Employed in old coughs and catarrhal affections.

TINCTURA BENZÖINI COMPOSITA.

Compound Tincture of Benzoin.

Balsamum Traumaticum, P.L. 1745.

Tinctura Benzöes composita, P.L. 1788.

Tinctura Benzöini composita, P.L. 1809, P.L. 1824.

Take of Benzoin three ounces and a half,
Storax, strained, two ounces and a half,
Balsam of Tolu ten drachms,
Aloes five drachms,
Rectified Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—Stimulant. Expectorant. Dose, fʒss. to fʒij. In chronic catarrh and confirmed asthma. It is decomposed by water, resinous matter being precipitated, and must therefore be triturated with yolk of egg, or with mucilage. It is more employed externally than internally, as a stimulant to languid ulcers; but its application to fresh wounds, for which it is mostly employed under the name of *Friar's Balsam*, appears to be injurious, by preventing the wound from healing by the first intention.

TINCTURA CALUMBÆ.

Tincture of Calumba.

Tinctura Colombæ, P.L. 1788.

Tinctura Calumbæ, P.L. 1809, P.L. 1824.

Take of Calumba, sliced, three ounces,
Proof Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—Tonic. Stomachic. Dose, fʒj. to fʒiij.

TINCTURA CAMPHORÆ.

Tincture of Camphor.

Spiritus Vini Camphoratus, P.L. 1720.

Spiritus vinosus Camphoratus, P.L. 1745.

Spiritus Camphoratus, P.L. 1788.

Spiritus Camphoræ, P.L. 1809, P.L. 1824.

Take of Camphor five ounces,
Rectified Spirit two pints;
Mix, that the Camphor may be dissolved.

Medicinal Uses.—Stimulant. It is used only externally. It is frequently applied to chilblains, and in cases of chronic rheumatism and numbness.

It is decomposed by water, which, combining with the spirit, precipitates the camphor.

TINCTURA CAMPHORÆ COMPOSITA.

Compound Tincture of Camphor.

Elixir Paregoricum, P.L. 1745.

Tinctura Opii Camphorata, P.L. 1788.

Tinctura Camphoræ composita, P.L. 1809, P.L. 1824.

Take of Camphor two scruples and a half,
Hard Opium, powdered,
Benzoic Acid, each seventy two grains,
Oil of Aniseed a fluidrachm,
Proof Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Use.—Anodyne. Dose, fʒj. to fʒiij. A fluid-ounce contains nearly two grains of opium.

TINCTURA CANTHARIDIS.

Tincture of Cantharides.

Tinctura Cantharidum, P.L. 1720, P.L. 1745.

Tinctura Cantharidis, P.L. 1788.

Tinctura Lyttæ, P.L. 1809.

Tinctura Cantharidis, P.L. 1824.

Take of Cantharides, bruised, four drachms,
Proof Spirit two pints ;
Macerate for fourteen days, and strain.

Medicinal Uses.—Diuretic. Stimulant. Dose, ℥ x. to f ʒj. given in some demulcent infusion. It is useful in gleet, fluor albus, and incontinence of urine. It is likewise employed externally as a stimulating embrocation or rubefacient, in conjunction with camphor liniment, &c.

TINCTURA CAPSICI.

Tincture of Capsicum.

Tinctura Capsici, P.L. 1809, P.L. 1824.

Take of Capsicum, bruised, ten drachms,
Proof Spirit two pints ;
Macerate for fourteen days, and strain.

Medicinal Use.—Stimulant. Dose, ℥ x. to f ʒj. It is employed in the low stage of typhus, and similar cases.

TINCTURA CARDAMOMI.

Tincture of Cardamom.

Tinctura Cardamomi, P.L. 1745, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Cardamom, bruised, three ounces and a half,
Proof Spirit two pints ;
Macerate for fourteen days, and strain.

Medicinal Uses.—Stimulant. Carminative. Dose, f ʒj. to f ʒij. It is generally employed as an adjunct to bitter infusions, but less frequently than the following.

TINCTURA CARDAMOMI COMPOSITA.

Compound Tincture of Cardamom.

Tinctura Stomachica, P.L. 1745.

Tinctura Cardamomi composita, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Cardamom,

Carraway, each, bruised, two drachms and a
half,

Cochineal, powdered, a drachm,

Cinnamon, bruised, five drachms,

Raisins five ounces,

Proof Spirit two pints;

Macerate for fourteen days, and strain.

Medicinal Uses.—As the former, and in similar doses.

TINCTURA CASCARILLÆ.

Tincture of Cascarilla.

Tinctura Cascarillæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Cascarilla, bruised, five ounces,

Proof Spirit two pints;

Macerate for fourteen days, and strain.

Medicinal Uses.—Tonic. Stomachic. Dose, f ʒj. to f ʒij.

TINCTURA CASTOREI.

Tincture of Castor.

Tinctura Castorei, P.L. 1720, P.L. 1745, P.L. 1788,
P.L. 1809, P.L. 1824.

Take of Castor, powdered, two ounces and a half,
Rectified Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—Antispasmodic. Stimulant. Dose, ℥ xx.
to f ʒij.

TINCTURA CATECHU.

Tincture of Catechu.

Tinctura Japonica, P.L. 1745.
Tinctura Catechu, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Catechu three ounces and a half,
Cinnamon, bruised, two ounces and a half,
Proof Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Use.—Astringent. Dose, f ʒj. to f ʒiij. It is a
very useful and grateful adjunct to *Mistura Cretæ* in diarrhœa.

TINCTURA CINCHONÆ.

Tincture of Cinchona.

Tinctura Corticis Peruviani simplex, P.L. 1745.

Tinctura Cinchonæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of heart-leaved Cinchona, bruised, eight ounces,
Proof Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—Tonic. Stomachic. Dose, fʒj. to fʒiij.
It is principally used in mixtures, with the Infusion or Decoction of Bark.

TINCTURA CINCHONÆ COMPOSITA.

Compound Tincture of Cinchona.

Tinctura Cinchonæ composita, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of lance-leaved Cinchona, bruised, four ounces,
Orange Peel, dried, three ounces,
Serpentary, bruised, six drachms,
Saffron two drachms,
Cochineal, powdered, a drachm,
Proof Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—Tonic. Stomachic. Dose, fʒj. to fʒiij.
It contains considerably less cinchona than the simple tincture, but is rendered more grateful by the admixture of the bitters and aromatics.

TINCTURA CINNAMOMI.

Tincture of Cinnamon.

Tinctura Cinnamomi, P.L. 1745, P.L. 1788, P.P. 1809,
P.L. 1824.

Take of Cinnamon, bruised, three ounces and a half,
Proof Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—See *Tinctura Cinnamomi composita*.

TINCTURA CINNAMOMI COMPOSITA.

Compound Tincture of Cinnamon.

Tinctura Aromatica, P.L. 1745.
Tinctura Cinnamomi composita, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Cinnamon, bruised, an ounce,
Cardamom, bruised, half an ounce,
Long Pepper, powdered,
Ginger, sliced, each two drachms and a half,
Proof Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—This and the former are both stomachic and astringent. Dose, f ʒj. to f ʒij.

TINCTURA COLCHICI.

Tincture of Meadow Saffron.

Take of Meadow Saffron Seeds, bruised, five ounces,
Proof Spirit two pints ;
Macerate for fourteen days, and strain.

Medicinal Uses.—In rheumatism and gout.

TINCTURA COLCHICI COMPOSITA.

Compound Tincture of Meadow Saffron.

Spiritus Colchici Ammoniatus, P.L. 1824.

Take of Meadow Saffron Seeds, bruised, five ounces,
Aromatic Spirit of Ammonia two pints ;
Macerate for fourteen days, and strain.

Medicinal Use.—Diuretic. Dose, f ʒss. to f ʒj. in water. The substances enumerated as incompatible with the *Spiritus Ammoniae Aromaticus*, are also such with this preparation.

TINCTURA CONII.

Tincture of Hemlock.

Take of Hemlock Leaves, dried, five ounces,
Cardamom, bruised, an ounce,
Proof Spirit two pints ;
Macerate for fourteen days, and strain.

Medicinal Uses.—Narcotic and antispasmodic. Dose, f ʒss. to f ʒj.

TINCTURA CUBEBÆ.

Tincture of Cubeb.

Take of Cubebs, bruised, five ounces,
Rectified Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—Stimulant. Internally taken in cases of gonorrhœa.

TINCTURA DIGITALIS.

Tincture of Foxglove.

Tinctura Digitalis, P.L. 1809, P.L. 1824.

Take of Foxglove Leaves, dried, four ounces,
Proof Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—Diuretic. Sedative. Dose, ℥ x. to ℥ xl., gradually increased. If it occasion vomiting or purging, its diuretic powers will be lost, which may be prevented by the use of a small quantity of opium.

TINCTURA GALLÆ.

Tincture of Gall.

Take of Galls, bruised, five ounces,
Proof Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Use.—Astringent.

TINCTURA GENTIANÆ COMPOSITA.

Compound Tincture of Gentian.

Tinctura Amara, P.L. 1745.

Tinctura Gentianæ Composita, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Gentian, sliced, two ounces and a half,
Orange Peel, dried, ten drachms,
Cardamom, bruised, five drachms,
Proof Spirit two pints ;
Macerate for fourteen days, and strain.

Medicinal Uses.—Tonic. Stomachic. Dose, fʒj. to fʒij.
It is most advantageously exhibited in combination with the Infusum Gentianæ compositum.

TINCTURA GUAIACI.

Tincture of Guaiacum.

Tinctura Guaiaci, P.L. 1809, P.L. 1824.

Take of Guaiacum Resin, bruised, seven ounces,
Rectified Spirit two pints ;
Macerate for fourteen days, and strain.

Medicinal Uses.—Stimulant. Diaphoretic. Dose, fʒj. to fʒij. When mixed with water the guaiacum is precipitated ; it should therefore be exhibited in mixture with some mucilage, or with yelk of egg.

TINCTURA GUAIACI COMPOSITA.

Compound Tincture of Guaiacum.

Tinctura Guaiacina volatilis, P.L. 1745.

Tinctura Guaiaci Ammoniata, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Guaiacum Resin, bruised, seven ounces,
Aromatic Spirit of Ammonia two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—Stimulant. Diaphoretic. Dose, fʒss. to fʒj. This is a more powerful preparation than the simple tincture, on account of the presence of ammonia. Like the simple tincture it is decomposed by water, and must therefore be exhibited with similar precautions.

It is incompatible with acids, and with acidulous, earthy, and metallic salts.

TINCTURA HELLEBORI.

Tincture of Hellebore.

Tinctura Hellebori, P.L. 1720.

Tinctura Melampodii, P.L. 1745.

Tinctura Hellebori nigri, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Hellebore, bruised, five ounces,
Proof Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Use.—Emmenagogue. Dose, ℥ xxx. to fʒj.

TINCTURA HYOSCYAMI.

Tincture of Henbane.

Tinctura Hyoscyami, P.L. 1809, P.L. 1824.

Take of Henbane Leaves, dried, five ounces,
Proof Spirit two pints ;
Macerate for fourteen days, and strain.

Medicinal Use.—Narcotic. Dose, fʒss. to fʒij. It is stated to procure sleep without affecting the head, or producing the costiveness which opium is apt to do.

TINCTURA JALAPÆ.

Tincture of Jalap.

Tinctura Jalapii, P.L. 1745, P.L. 1788.
Tinctura Jalapæ, P.L. 1809, P.L. 1824.

Take of Jalap, bruised, ten ounces,
Proof Spirit two pints ;
Macerate for fourteen days, and strain.

Medicinal Use.—Cathartic. Dose, fʒj. to fʒss. It is an efficient medicine, but is rarely administered except as an adjunct to cathartic combinations.

TINCTURA IODINII COMPOSITA.

Compound Tincture of Iodine.

Take of Iodine an ounce,
Iodide of Potassium two ounces,
Rectified Spirit two pints ;
Macerate until they are dissolved, and strain.

Medicinal Uses.—See POTASSII IODIDUM.

TINCTURA KINO.

Tincture of Kino.

Tinctura Kino, P.L. 1809, P.L. 1824.

Take of Kino, bruised, three ounces and a half,
Rectified Spirit two pints ;
Macerate for fourteen days, and strain.

Medicinal Uses.—Astringent. Dose, fʒj. to fʒij. It consists chiefly of tannin, and is said to be less efficacious than the *Tinctura Catechu*.

TINCTURA LAVANDULÆ COMPOSITA.

Compound Tincture of Lavender.

Spiritus Lavendulæ compositus Matthiæ, P.L. 1720.

Spiritus Lavendulæ compositus, P.L. 1745.

Tinctura Lavendulæ composita, P.L. 1788.

Spiritus Lavandulæ compositus, P.L. 1809, P.L. 1824.

Take of Spirit of Lavender a pint and a half,
Spirit of Rosemary half a pint,
Cinnamon, bruised,
Nutmeg, bruised, each two drachms and a half,
Red Saunders, sliced, five drachms.

Macerate for fourteen days, and strain.

Medicinal Uses.—Stimulant. Stomachic, in languors, &c. Dose, from fʒss. to fʒij. in water or any convenient liquid.

TINCTURA LUPULI.

Tincture of Hop.

Tinctura Humuli, P.L. 1809, P.L. 1824.

Take of Hops six ounces,
Proof Spirit two pints ;
Macerate for fourteen days, and strain.

Medicinal Uses.—Sedative. Tonic. Dose, from f3ss. to f3ij.
Its powers are questionable as a narcotic, but are stomachic.

TINCTURA MYRRHÆ.

Tincture of Myrrh.

Tinctura Myrrhæ simplex, P.L. 1720.

Tinctura Myrrhæ, P.L. 1745, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Myrrh, bruised, three ounces,
Rectified Spirit two pints ;
Macerate for fourteen days, and strain.

Medicinal Uses.—Tonic. Deobstruent. Dose, f3ss. to f3j.
It is, however, rarely used internally, but as an external application to foul ulcers, and when diluted with water as a lotion for spongy gums. It is decomposed, and its resin precipitated, by mixture with water.

TINCTURA OPII.

Tincture of Opium.

Tinctura Opii, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Hard Opium, powdered, three ounces,
Proof Spirit two pints ;
Macerate for fourteen days, and strain.

Qualities.—This tincture is of a deep brownish red colour, and possesses the peculiar odour and taste of the opium itself. Its specific gravity I find to be about 0·952, when prepared with proof spirit, as directed in the Pharmacopœia; about 19 minims contain one grain of opium; this was proved by boiling down the tincture, and also by determining the quantity of opium left undissolved. It will appear from what has already been stated, that proof spirit is a much better solvent of opium than cold water; for the latter dissolves less than 3-7ths of the opium, whereas proof spirit, as I found in preparing the tincture, dissolves more than 2-3rds of it.

Incompatibles.—This tincture is decomposed by ammonia, potash, and soda, and their carbonates, morphia being precipitated; most metallic salts, and infusion of galls, also decompose it.

Medicinal Use.—Narcotic. As 19 minims contain one grain of opium, the quantity exhibited must depend upon that of the opium which it is intended to give. Its dose is generally stated to be from ℥ x. to ℥ lx. It is given in preference to opium in substance, in cases of accident or of sudden and extreme pain; it is sometimes preferred to solid opium in chronic cases, on account of the facility with which the dose may be apportioned and varied according to circumstances. It is externally employed as an anodyne in lotions.

TINCTURA RHEI COMPOSITA.

Compound Tincture of Rhubarb.

Tinctura Rhabarbari composita, 1788.

Tinctura Rhei composita, P.L. 1809, P.L. 1824.

Take of Rhubarb, sliced, two ounces and a half,
Liquorice, bruised, six drachms,
Ginger, sliced,
Saffron, each three drachms,
Proof Spirit two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—Purgative. Stomachic. Dose, f ʒij. to f jss.

TINCTURA SCILLÆ.

Tincture of Squill.

Tinctura Scillæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Squill, fresh dried, five ounces,

Proof Spirit two pints ;

Macerate for fourteen days, and strain.

Medicinal Uses.—Expectorant. Diuretic. Dose, ℥ x. to
℥ xxx.

TINCTURA SENNÆ COMPOSITA.

Compound Tincture of Senna.

Elixir Salutis, P.L. 1720.

Tinctura Sennæ, P.L. 1745, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Senna three ounces and a half,

Carraway, bruised, three drachms and a half,

Cardamom, bruised, a drachm,

Raisins five ounces,

Proof Spirit two pints ;

Macerate for fourteen days, and strain.

Medicinal Uses.—Stomachic and purgative. Dose, f ʒij. to
f ʒj.

TINCTURA SERPENTARIÆ.

Tincture of Serpentary.

Tinctura Serpentariæ virginianæ, P.L. 1720.

Tinctura Serpentariæ, P.L. 1745, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Serpentary, bruised, three ounces and a half,

Proof Spirit two pints ;

Macerate for fourteen days, and strain.

Medicinal Uses.—Tonic. Diaphoretic. Dose, f ʒi. to f ʒiij.

TINCTURA VALERIANÆ.

Tincture of Valerian.

Tinctura Valerianæ simplex, P.L. 1745.

Tinctura Valerianæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Valerian, bruised, five ounces,

Proof Spirit two pints ;

Macerate for fourteen days, and strain.

Medicinal Use.—Antispasmodic. Dose, from fʒi. to fʒiij. It is seldom employed except as an adjunct to the infusion of valerian.

TINCTURA VALERIANÆ COMPOSITA.

Compound Tincture of Valerian.

Tinctura Valerianæ volatilis, P.L. 1745.

Tinctura Valerianæ Ammoniata, P.L. 1788, P.L. 1809.

Tinctura Valerianæ Ammoniata, P.L. 1824.

Take of Valerian, bruised, five ounces,

Aromatic Spirit of Ammonia two pints ;

Macerate for fourteen days, and strain.

Medicinal Use.—Antispasmodic. Dose, fʒss. to fʒj. It is more powerful than the simple tincture, only on account of the ammonia which it contains. It is incompatible with acids, and with acidulous, metallic, and earthy salts.

TINCTURA ZINGIBERIS.

Tincture of Ginger.

Tinctura Zingiberis, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Ginger, sliced, two ounces and a half,

Rectified Spirit two pints ;

Macerate for fourteen days, and strain.

Medicinal Uses.—Stimulant. Carminative. Dose, fʒi. to fʒij. It is useful in gout when it attacks the stomach, and in flatulent colic, and as a corrigent to griping purgatives.

VEGETABILIA.

Vegetables.

Vegetables are to be collected in dry weather, when wetted neither with showers nor dew; they are to be collected annually, and those which have been kept longer are to be rejected.

Most **Roots** are to be dug up before the stalks or leaves appear.

BARKS ought to be collected at that season in which they can most easily be separated from the wood.

LEAVES are to be gathered after the flowers are blown, and before the seeds ripen.

FLOWERS are to be gathered recently blown.

SEEDS are to be collected when ripe, and are to be kept in their own seed-vessels.

VEGETABILIIUM PRÆPARATIO.

The Preparation of Vegetables.

Vegetables shortly after they have been gathered, those excepted which ought to be fresh, are to be lightly strewed, and dried as quickly as possible, with a gentle heat; keep them afterwards in proper vessels, excluded from the access of light and moisture.

Lay up those **Roots** which we have directed to be kept fresh, in dry sand. Cut the **Cormus** of Meadow Saffron and the **Bulb** of the Squill, before drying, transversely into thin slices, the dry rind being previously peeled off.

Put pulpy fruits, if they are unripe, or if ripe and dry, in a moist place, that they may soften; then press the pulps through a hair sieve; afterwards boil them over a slow fire, frequently stirring; lastly, evaporate the water in a water-bath, until the pulps become of a proper consistence.

Pour boiling water upon the bruised Pods of Cassia, that the pulp may be washed out, which press first through a coarse sieve, and afterwards through a hair one; then evaporate the water in a water-bath, until the pulp acquires a proper consistence.

Press the pulp or juice of ripe and fresh fruits through a sieve, no boiling being used.

GUM-RESINS.

Separate Opium from foreign substances, especially the external, as carefully as possible. Let Opium be kept soft, which may be fit to be formed into pills, and hard, which has been so dried in a water-bath, that it may be rubbed to powder.

Those Gum-Resins are to be reckoned best, which are so perfect, that no purification is necessary. But those which appear to be less pure, are to be boiled in water until they soften, and squeezed with a press through a hempen cloth; then to be set by, that the resinous part may subside. The supernatant liquor being poured off, evaporate it in a water-bath, the resinous part being added towards the end, that it may unite with the gummy part.

The Gum-Resins which melt easily, may be purified by putting them into an ox-bladder, and keeping them in boiling water, until they become so soft, that they may be separated from impurities by a press through a hempen cloth.

Dissolve Storax in rectified spirit, and strain; then let the spirit distil with a gentle heat, until it becomes of a proper consistence.

V I N A.

Wines.

Medicated Wines should be prepared in stopped glass vessels, and frequently shaken during maceration.

VINUM ALOËS.

Wine of Aloes.

Tinctura Hieræ, P.L. 1720.

Tinctura Sacra, P.L. 1745.

Vinum Aloës, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Aloes, rubbed to powder, two ounces,
Canella, powdered, four drachms,
Sherry Wine two pints ;

Macerate for fourteen days, frequently shaking, and strain.

Medicinal Uses.—Stomachic, in doses of f ʒi. to f ʒij. Purgative, f ʒi. to f ʒij.

VINUM COLCHICI.

Wine of Meadow Saffron.

Vinum Colchici, P.L. 1824.

Take of dried Meadow Saffron Cormus, sliced, eight ounces,

Sherry Wine two pints ;

Macerate for fourteen days, and strain.

Medicinal Uses.—Diuretic. Dose, from ℥ xxx. to f ʒi. It is stated to be a specific in the gout, allaying the pain, and cutting short the paroxysm.

VINUM IPECACUANHÆ.

Wine of Ipecacuanha.

Vinum Ipecacoanhæ, P.L. 1745.

Vinum Ipecacuanhæ, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Ipecacuanha, bruised, two ounces and a half,
Sherry Wine two pints;
Macerate for fourteen days, and strain.

Medicinal Uses.—Diaphoretic. Dose, ℥ xx. to ℥ xl. Emetic. Dose, f ʒij. to f ʒiv. It is as efficacious an emetic as *Vinum Antimonii Potassio-Tartratis*, and, being milder in its operation, is better adapted for infants, a tea-spoonful or f ʒss. being administered every ten or fifteen minutes till it operates.

The active power of ipecacuanha resides in a peculiar alkali, to which the name of *Emetina* has been given. The root contains 14 per cent. of it, mixed with woody fibre, starch, gum, &c. Dr. A. T. Thomson (*Dispensatory*, p. 817) states, that a pint of sherry takes up 100 grains of the soluble matter of ipecacuanha. *Emetina* is nearly insoluble in water, but soluble in alcohol. It is insoluble in æther, but is dissolved by most acids. The solutions are not decomposed by potassio-tartrate of antimony, but they are incompatible with salts of lead and mercury, and infusion of galls.

According to the analysis of MM. Dumas and Pelletier, *emetina* consists of

Carbon	64.57
Azote	4.30
Hydrogen	7.77
Oxygen.....	22.95

99.59

VINUM OPII.

Wine of Opium.

Laudanum liquidum Sydenhami, P.L. 1720.

Tinctura Thebaica, P.L. 1745.

Vinum Opii, P.L. 1809, P.L. 1824.

Take of Purified extract of Opium two ounces and a half,

Cinnamon, bruised,

Cloves, bruised, each two drachms and a half,

Sherry Wine two pints ;

Macerate for fourteen days, and strain.

Medicinal Use.—Narcotic. Dose, ℥ x. to f ʒi.

This preparation differs from the *Tinctura Opii*, not only in containing aromatics, but in the purification of the Opium. Various circumstances render it difficult to form an estimate of the comparative powers of these preparations ; they probably differ but little, for respectable authorities agree in representing their doses as similar. The *Vinum Opii* must be less disagreeable to most persons than the tincture, not only on account of the aromatics which it contains, but because the opium during purification loses its peculiar and disagreeable smell and taste.

VINUM VERATRI.

Wine of White Hellebore.

Vinum Veratri, P.L. 1824.

Take of White Hellebore, sliced, eight ounces,

Sherry Wine two pints ;

Macerate for fourteen days, and strain.

Medicinal Uses.—Emetic and cathartic, acting usually with considerable violence. Dose, ℥ v. to ℥ x. See VERATRIA.

UNGUENTA.Ointments.

UNGUENTUM ANTIMONII POTASSIO-
TARTRATIS.

Ointment of Potassio-tartrate of Antimony.

Take of Potassio-Tartrate of Antimony, rubbed to
powder, an ounce,
Lard four ounces ;
Mix.

Medicinal Use.—Employed in chronic swellings of the joints, particularly after rheumatism. The use of this ointment is followed by the appearance of a pustular eruption; and if persisted in for some time, the eruption is not confined to the part upon which it may have been rubbed, but appears diffused over the whole body; a singular effect, dependent no doubt upon its absorption into the system.

UNGUENTUM CANTHARIDIS.

Ointment of Cantharides.

Unguentum Cantharidis, P.L. 1824.

Take of Cantharides, rubbed to very fine powder, an
ounce,
Distilled Water four fluidounces,
Cerate of Resin four ounces ;

Boil down the Water with the Cantharides to half, and strain. Mix the Cerate with the strained liquor; afterwards let it evaporate to a proper consistence.

Medicinal Use.—This is sometimes employed for the same purpose as the Ceratum Cantharidis; it is a milder preparation, and frequently inefficacious.

UNGUENTUM CETACEI.

Ointment of Spermaceti.

Linimentum Album, P.L. 1745.

Unguentum Spermaceti, P.L. 1788.

Unguentum Cetacei, P.L. 1809, P.L. 1824.

Take of Spermaceti six drachms,

White Wax two drachms,

Olive Oil three fluidounces ;

Melt them together with a slow fire, then stir constantly until they become cold.

Medicinal Use.—There is no difference in the properties of this and the Ceratum Cetacei, excepting that the ointment is softer. They are used for similar purposes.

UNGUENTUM CREASOTI.

Ointment of Creasote.

Take of Creasote a fluidrachm,

Lard an ounce ;

Rub and mix them.

Medicinal Use.—Employed in mild cases of ringworm.

UNGUENTUM ELEMI.

Ointment of Elemi.

Unguentum e gummi Elemi sive Linimentum Arcæi,
P.L. 1720.

Unguentum e gummi Elemi, P.L. 1745.

Unguentum Elemi compositum, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Elemi a pound,
Common Turpentine ten ounces,
Suet two pounds,
Olive Oil two fluidounces ;

Melt the Elemi with the Suet ; then remove them from the fire, and immediately mix with them the Turpentine and the Oil ; afterwards press through a linen cloth.

Medicinal Use.—Stimulant and digestive. It is used to keep open setons and issues, and as an application to ulcers which do not admit of the use of adhesive straps.

UNGUENTUM GALLÆ COMPOSITUM.

Compound Ointment of Galls.

Take of Galls rubbed to a very fine powder, two
drachms,
Lard two ounces,
Hard Opium, powdered, half a drachm ;
Mix.

Medicinal Use.—Astringent. Used in hæmorrhoidal affections.

UNGUENTUM HYDRARGYRI FORTIUS.

Stronger Ointment of Mercury.

Unguentum Cæruleum fortius, P.L. 1745.

Unguentum Hydrargyri fortius, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of Mercury two pounds,
Lard twenty-three ounces,
Suet an ounce ;

First rub the Mercury with the Suet and a little of the Lard until globules can no longer be seen ; then add that which is left of the Lard, and mix.

Process.—During trituration with the fatty matter, the mercury is probably reduced to the same state as that in which it exists in the *Pilulæ Hydrargyri*.

Medicinal Uses.—This ointment furnishes a prompt and probably one of the least exceptionable modes of introducing mercury into the system. It is generally applied by rubbing ʒss. to ʒi. on some part of the body where the cuticle is thin, generally, in syphilitic cases, on the inside of the thigh : in chronic hepatitis it is usually applied in the region of the liver.

As the preparation of this ointment is an exceedingly tedious operation, various means, and most of them of an objectionable nature, have been resorted to in order to shorten it. Some employ *Oleum Sulphuratum*, the use of which, on account of the well-known power of sulphur in diminishing the effects of mercury, ought always to be reprobated. By others, turpentine is used on account of its tenacity ; but this is apt to produce pustules, which prevent the continuance of the friction. I have been assured that the admixture of a portion of old ointment greatly facilitates the operation. The ointment contains half its weight of mercury.

UNGUENTUM HYDRARGYRI MITIUS.

Milder Ointment of Mercury.

Unguentum Cæruleum mitius, P.L. 1745.

Unguentum Hydrargyri mitius, P.L. 1788, P.L. 1809,
P.L. 1824.

Take of stronger Ointment of Mercury a pound,
Lard two pounds ;
Mix.

Medicinal Use.—This is used as a dressing, and for those purposes in which the preceding preparation would be too powerful. Six drachms contain one drachm of mercury.

UNGUENTUM HYDRARGYRI NITRATIS.

Ointment of Nitrate of Mercury.

Unguentum Hydrargyri Nitrati, P.L. 1788, P.L. 1809.

Unguentum Hydrargyri Nitratis, P.L. 1824.

Take of Mercury an ounce,
Nitric Acid eleven fluidrachms,
Lard six ounces,
Olive Oil four fluidounces ;

First dissolve the Mercury in the Acid ; then mix the solution, while hot, with the Lard and Oil, melted together.

Process.—By the decomposition of a portion of the nitric acid, the mercury is oxidized, and dissolved by the acid remaining undecomposed.

Medicinal Uses.—Stimulant and detergent. When its strength is diminished by the addition of lard, it is a local remedy of great efficacy in eruptions and various cutaneous diseases.

UNGUENTUM HYDRARGYRI NITRICO-OXYDI.

Ointment of Nitric-oxide of Mercury.

Unguentum Hydrargyri Nitrico-Oxydi, P.L. 1809,
P.L. 1824.

Take of Nitric-oxide of Mercury an ounce,
White Wax two ounces,
Lard six ounces;

Add the Nitric-oxide of Mercury, rubbed to very fine powder, to the Wax and Lard, melted together, and mix.

Medicinal Use.—This is applied in the same manner, and for similar purposes, as the preceding ointment.

UNGUENTUM HYDRARGYRI IODIDI.

Ointment of Iodide of Mercury.

This is prepared in the same manner as the ointment of Nitric-oxide of Mercury.

Medicinal Use.—Used for dressing to scrofulous sores.

UNGUENTUM HYDRARGYRI BINIODIDI.

Ointment of Biniodide of Mercury.

This is prepared in the same manner as the ointment of Nitric-oxide of Mercury.

Medicinal Uses.—A more active preparation than the former, and employed in analogous cases.

UNGUENTUM HYDRARGYRI AMMONIO-CHLORIDI.

Ointment of Ammonio-chloride of Mercury.

Unguentum e Mercurio præcipitato albo, P.L. 1745.

Unguentum calcis Hydrargyri albæ, P.L. 1788.

Unguentum Hydrargyri præcipitati albi, P.L. 1809,
P.L. 1824.

Take of Ammonio-chloride of Mercury a drachm,
Lard an ounce and a half;

Add the Ammonio-chloride of Mercury, to the Lard melted over a slow fire, and mix.

Medicinal Uses.—Stimulant and detergent.

UNGUENTUM IODINII COMPOSITUM.

Compound Ointment of Iodine.

Take of Iodine half a drachm,
Iodide of Potassium a drachm,
Rectified Spirit a fluidrachm,
Lard two ounces;

First rub the Iodine and Iodide of Potassium with the Spirit, then mix with the Lard.

Medicinal Uses.—Used in bronchocele.

UNGUENTUM PICIS LIQUIDÆ.

Ointment of Liquid Pitch [Tar].

Unguentum e Pice, P.L. 1745.

Unguentum Picis, P.L. 1788.

Unguentum Picis Liquidæ, P.L. 1809, P.L. 1824.

Take of Liquid Pitch [Tar],

Suet, each a pound;

Melt them together, and press through a linen cloth.

Medicinal Uses.—This ointment is employed for the removal of tetter, and in tinea capitis.

UNGUENTUM PICIS NIGRÆ.

Ointment of Black Pitch.

Unguentum Basilicum nigrum, vel Tetrapharmacum,
P.L. 1745.

Unguentum Picis Aridæ, P.L. 1809.

Unguentum Picis nigræ, P.L. 1824.

Take of Black Pitch,

Wax,

Resin, each nine ounces,

Olive Oil, sixteen fluidounces;

Melt them together, and press through a linen cloth.

Medicinal Uses.—Digestive and stimulant.

UNGUENTUM PLUMBI COMPOSITUM.

Compound Ointment of Lead.

Take of Prepared Chalk eight ounces,

Distilled Vinegar six fluidounces,

Plaster of Lead three pounds,

Olive Oil a pint;

Melt the Plaster in the Oil with a slow fire; then gradually add the Chalk separately mixed with the Vinegar, the effervescence being finished, and stir constantly until they are cooled.

Medicinal Use.—Used as a dressing to indolent ulcers.

UNGUENTUM PLUMBI IODIDI.

Ointment of Iodide of Lead.

Take of Iodide of Lead an ounce,
Lard eight ounces ;
Rub together, and mix.

Medicinal Use.—Employed in chronic enlargement of joints.

UNGUENTUM SAMBUCL.

Ointment of Elder.

Unguentum Sambucinum, P.L. 1720, P.L. 1745.
Unguentum Sambuci, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Elder [Flowers],
Lard, each two pounds ;
Boil the Elder Flowers in the Lard until they become
crisp ; then press through a linen cloth.

Medicinal Use.—This is used for the same purposes as the
Unguentum Cetacei, over which it possesses no advantage but
a pleasant smell.

UNGUENTUM SULPHURIS.

Ointment of Sulphur.

Unguentum e Sulphure, P.L. 1745.
Unguentum Sulphuris, P.L. 1788, P.L. 1809, P.L. 1824.

Take of Sulphur three ounces,
Lard half a pound,
Oil of Bergamot twenty minims ;
Mix.

UNGUENTUM SULPHURIS COMPOSITUM.

Compound Ointment of Sulphur.

Unguentum Sulphuris Compositum, P.L. 1809, P.L. 1824.

Take of Sulphur half a pound,
White Hellebore, powdered, two ounces,
Nitrate of Potash a drachm,
Soft Soap half a pound,
Lard a pound and a half,
Oil of Bergamot thirty minims;

Mix.

Medicinal Use.—These ointments are both used for the cure of the itch; the latter sometimes excites too much irritation.

UNGUENTUM VERATRI.

Ointment of White Hellebore.

Unguentum Hellebori Albi, P.L. 1788.

Unguentum Veratri, P.L. 1809, P.L. 1824.

Take of White Hellebore, powdered, two ounces,
Lard eight ounces,
Oil of Lemons twenty minims;

Mix.

Medicinal Use.—This is used for the cure of scabies, but is said to be less certain in its effects than the Sulphur ointment.

UNGUENTUM ZINCI.

Ointment of Zinc.

Unguentum Zinci, P.L. 1809, P.L. 1824.

Take of Oxide of Zinc an ounce,
Prepared Lard six ounces;

Mix.

This may be considered as an improvement upon the Ceratum Calaminæ. It is recommended as being very useful in some species of ophthalmia, smeared upon the tarsi every night.

TABLE OF FORMER AND NEW NAMES.

FORMER NAMES.

NEW NAMES.

A.

Acidum aceticum dilutum.

——— muriaticum.

Ammoniae murias.

——— subcarbonas.

Antimonii sulphuretum.

——— sulphuretum præci-
pitatum.

Antimonium tartarizatum.

Arsenicum album.

Acetum destillatum.

Acidum hydrochloricum.

Ammoniae hydrochloras.

——— sesquicarbonas.

Antimonii sesquisulphuretum.

——— oxysulphuretum.

——— potassio-tartras.

Acidum arseniosum.

B.

Bismuthi subnitras.

Bismuthi trisnitras.

C.

Calcis murias.

Ceratum simplex.

Confectio amygdalarum.

——— aurantiorum.

——— scammonæ.

Cuprum ammoniatum.

Calcii chloridum.

Ceratum.

Confectio amygdalæ.

——— aurantii.

——— scammonii.

Cupri ammonio-sulphas.

D.

Decoctum cinchonæ.

——— lichenis.

——— sarsaparillæ.

——— sarsaparillæ compo-
situm.

Decoctum cinchonæ cordifoliæ.

——— cetrariæ.

——— sarzæ.

——— sarzæ compositum.

FORMER NAMES.

NEW NAMES.

E.

Emplastrum picis compositum.	Emplastrum picis.
Extractum cinchonæ.	Extractum cinchonæ cordifoliae.
———— opii.	———— opii purificatum.

F.

Ferri subcarbonas.	Ferri sesquioxylum.
Ferrum ammoniatum.	———— ammonio-chloridum.
———— tartarizatum.	———— potassio-tartras.

H.

Hydrargyri oxydum cinereum.	Hydrargyri oxydum.
———— oxydum rubrum.	———— binoxylum.
———— oxymurias.	———— bichloridum.
———— submurias.	———— chloridum.
———— sulphuretum nigrum.	———— sulphuretum cum sulphure.
———— sulphuretum rubrum.	———— bisulphuretum.
———— præcipitatum album.	———— ammonio-chloridum.

I.

Infusum caryophyllorum.	Infusum caryophylli.
-------------------------	----------------------

L.

Linimentum ammoniæ fortius.	Linimentum ammoniæ.
———— ammoniæ subcarbonatis.	———— ammoniæ sesquicarbonatis.
———— hydrargyri.	———— hydrargyri compositum.
———— saponis compositum.	———— saponis.
Liquor ammoniæ subcarbonatis.	Liquor ammoniæ sesquicarbonatis.
———— arsenicalis.	———— potassæ arsenitis.
———— calcis muriatis.	———— calcii chloridi.
———— cupri ammoniati.	———— cupri ammonio-sulphatis.

FORMER NAMES.

NEW NAMES.

Liquor hydrargyri oxymuriatis. Liquor hydrargyri bichloridi.

———— plumbi subacetatis. ———— plumbi diacetatis.

———— potassæ subcarbonatis. ———— potassæ carbonatis.

M.

Magnesiae subcarbonas. Magnesiae carbonas.

Mistura amygdalarum. Mistura amygdalæ.

Mucilago acaciæ. ———— acaciæ.

———— amyli. Decoctum amyli.

O.

Oleum pulegii. Oleum menthæ pulegii.

———— terebinthinæ rectificatum. ———— terebinthinæ purificatum.

Oxymel simplex. Oxymel.

P.

Pilulæ hydrargyri submuriatis compositæ. Pilulæ hydrargyri chloridi compositæ.

———— saponis cum opio. ———— saponis compositæ.

Plumbi oxydum semivitreum. Plumbi oxydum.

———— subcarbonas. ———— carbonas.

Potassa fusa. Potassæ hydras.

Potassæ carbonas. ———— bicarbonas.

———— subcarbonas. ———— carbonas.

———— supertartras. ———— bitartras.

———— sulphuretum. Potassii sulphuretum.

Pulvis antimonialis. Pulvis antimonii compositus.

S.

Sodæ carbonas. Sodæ sesquicarbonas.

———— murias. Sodii chloridum.

———— subboras. Sodæ biboras.

———— subcarbonas. ———— carbonas.

Soda tartarizata. ———— potassio-tartras.

Spiritus camphoræ. Tinctura camphoræ.

FORMER NAMES.	NEW NAMES.
——— lavandulæ compositus.	——— lavandulæ composita.
Spiritus pulegii.	Spiritus menthæ pulegii.
Syrupus aurantiorum.	Syrupus aurantii.
——— sarsaparillæ.	——— sarzæ.
——— simplex.	Syrupus.

T.

Tinctura ferri ammoniati.	Tinctura ferri ammonio-chloridi.
——— ferri muriatis.	——— ferri sesquichloridi.
——— sennæ.	——— sennæ composita.

V.

Vinum antimonii tartarizati.	Vinum antimonii potassio-tartratis.
------------------------------	-------------------------------------

U.

Unguentum elemi compositum.	Unguentum elemi.
——— hydrargyri præcipitati albi.	——— hydrargyri ammonio-chloridi.

POSOLOGICAL TABLE.

Absinthium	ʒj to	ʒj.
Acacia	ʒss	ʒij.
Acetum Colchici	fʒss	fʒj.
—— Destillatum.....	fʒi	fʒss.
—— Scillæ	fʒss	fʒij.
Acidum Benzoicum	gr. x.	ʒss.
—— Citricum	gr. x	ʒss.
—— Hydrochloricum	℥v	℥xx.
—— Hydrochloricum dilutum	℥xx	℥lxx.
—— Nitricum dilutum	℥x	℥xl.
—— Hydrocyanicum dilutum	℥v.	
—— Phosphoricum dilutum	℥xx	fʒj.
—— Sulphuricum dilutum	℥x	℥xl.
—— Tartaricum	gr. x	ʒss.
Aconiti Folia	gr. j	gr. v.
Æther Sulphuricus	fʒss	fʒij.
Ærugo	gr. $\frac{1}{8}$	gr. j.
Aloës Extractum	gr. v	gr. xv.
Alumen	gr. x	ʒj.
Ammoniacum	gr. x	ʒss.
Ammoniæ Hydrochloras	gr. x	ʒss.
—— Sesquicarbonas	gr. v	ʒj.
Anethi Fructus	gr. x	ʒj.
Anisi Fructus	gr. x	ʒj.
Anthemidis Flores	gr. x	ʒss.
Antimonii Oxysulphuretum	gr. j	gr. iv.
—— Potassio-tartras (Diaphoretic).	gr. $\frac{1}{4}$	gr. ss.
—— (Emetic)	gr. j	gr. ij.
—— Sesquisulphuretum	gr. x	ʒss.

Aqua Anethi	}	f3j to f3iv.
— Carui		
— Cinnamomi		
— Fœniculi		
— Menthæ piperitæ		
— ————— pulegii		
— ————— viridis		
— Pimentæ		
Argenti Nitras	gr. $\frac{1}{8}$	gr. j.
Armoraciæ Radix	3j	3j.
Assafoetida	gr. x	3ss.
Balsamum Peruvianum	gr. x	3ss.
———— Tolutanum	gr. x	3ss.
Belladonnæ Folia	gr. ss	gr. x.
Benzoinum	gr. x	3ss.
Bismuthi trisnitræ	gr. v	gr. xv.
Cajuputi Oleum	℥j	℥v.
Calumbæ Radix	gr. x	3j.
Cambogia	gr. ij	gr. x.
Camphora	gr. iij	3j.
Canellæ Cortex	gr. x	3ss.
Capsici Baccæ	gr. v	gr. x.
Cardamines Flores	3j	3j.
Cardamomi Semina	gr. v	3ss.
Carui Fructus	gr. x	3j.
Caryophylli	gr. v	3ss.
———— Oleum	℥ij	℥v.
Cascarillæ Cortex	gr. x	3j.
Cassiæ Pulpa	3ss	3j.
Castoreum	gr. v	3j.
Catechu	gr. x	3ij.
Centaurii Cacumina	gr. xv	3j.
Cetaceum	3j	3iss.
Cinchonæ cordifoliæ Cortex	gr. x	3iss.
———— lancifoliæ Cortex	gr. x	3iss.
———— oblongifoliæ Cortex	gr. x	3iss.
Cinnamomi Cortex	gr. v	3j.

Cinnamomi Oleum	℥j to	℥iij.
Colchici Cormus	gr. j	gr. v.
Colocynthis Pulpa	gr. j	gr. v.
Confectio Amygdalæ	ʒj	ʒij.
———— Aromatica	gr. xx	ʒj.
———— Aurantii	ʒj	ʒj.
———— Cassiæ	ʒij	ʒj.
———— Opii	gr. x	ʒss.
———— Piperis nigri	ʒj	ʒij.
———— Rosæ caninæ	ʒj	ʒj.
———— Gallicæ	ʒj	ʒj.
———— Scammonii	ʒj	ʒj.
———— Sennæ	ʒj	ʒiij.
Conii Folia	gr. v	ʒj.
Contrajervæ Radix	gr. x	ʒss.
Copaiba	ʒj	ʒj.
Coriandri Fructus	ʒj	ʒj.
Creta præparata	gr. x	gr. xl.
Cubeba	ʒj	ʒij.
Cupri Ammonio-Sulphas	gr. $\frac{1}{4}$	gr. v.
Cupri Sulphas	gr. ss	gr. ij.
———— (Emetic)	gr. ij	gr. x.
Cuspariæ Cortex	gr. x	ʒj.
Cymini Fructus	ʒj	ʒj.
Dauci Fructus	ʒj	ʒj.
Decoctum Aloës compositum	fʒss	fʒj.
———— Cetrariæ	fʒi	fʒiv.
———— Chimaphilæ	fʒj	fʒij.
———— Cinchonæ	fʒj	fʒiij.
———— Dulcamaræ	fʒss	fʒj.
———— Granati	fʒss	fʒj.
———— Sarzæ	fʒiv	fʒviii.
———— compositum	fʒiv	fʒviiij.
———— Scoparii compositum	fʒi	fʒiss.
———— Senegæ	fʒiss	fʒiij.
———— Tormentillæ	fʒi	fʒiss.
———— Ulmi	fʒiv	fʒvj.
———— Uvæ Ursi	fʒj	fʒij.

Digitalis Folia	gr. ss to	gr. iij.
Extractum Aconiti	gr. ss	gr. v.
——— Aloës purificatum	gr. v	gr. xv.
——— Cinchonæ	gr. x	℥ss.
——— Colchici Aceticum	gr. ss	gr. ij.
——— Cormi	gr. ss.	gr. ij.
——— Colocynthis	gr. v	℥ss.
——— compositum.....	gr. v	℥ss.
——— Conii.	gr. v	℥j.
——— Digitalis	gr. ss	gr. j.
——— Elaterii	gr. ss	gr. ij.
——— Gentianæ	gr. x	℥ss.
——— Hæmatoxyli	gr. x	℥ss.
——— Hyoseyami	gr. v	gr. x.
——— Jalapæ	gr. x	℥j.
——— Lactucæ	gr. v	gr. x.
——— Lupuli	gr. v	℥j.
——— Opii purificatum	gr. j	gr. v.
——— Papaveris	gr. ij	gr. xx.
——— Rhei	gr. x	℥ss.
——— Sarzæ	gr. x	℥j.
——— Stramonii	gr. $\frac{3}{4}$	gr. ij.
——— Taraxaci	gr. x	℥j.
——— Uvæ Ursi	gr. x	gr. xx.
Ferri Ammonio-chloridum	gr. iij	gr. xv.
—— Iodidum	gr. i	gr. iij.
—— Potassio-tartras	gr. x	℥ss.
—— Sesquioxylum	gr. v	gr. xxx.
—— Sulphas	gr. j	gr. v.
Fœniculi Fructus	℥j	℥j.
Galbani Gummi-resina	gr. x	℥ss.
Gentianæ Radix	gr. x	℥j.
Granati Cortex	℥j	℥j.
Guaiaci Resina	gr. x	℥ss.
Hydrargyri Bichloridum	gr. $\frac{1}{8}$	gr. $\frac{1}{4}$.

Hydrargyri Biniodidum	gr. ss to	gr. j.
———— Binoxidum	gr. ss	gr. j.
———— Chloridum (Alterative)	gr. ss	gr. j.
———— (Purgative)	gr. v	gr. x.
———— Iodidum	gr. j	gr. iij.
———— Oxydum	gr. j	gr. iij.
———— Sulphuretum cum Sulphure .	gr. v	ʒss.
Hydrargyrum cum Cretâ	gr. x	ʒss.
Hyoscyami Folia	gr. v	gr. x.
Jalapæ Radix	gr. x	ʒss.
Infusum Anthemidis	f ʒj	f ʒij.
———— Armoraciæ compositum	f ʒj	f ʒiss.
———— Aurantii compositum	f ʒj	f ʒij.
———— Calumbæ	f ʒiss	f ʒij.
———— Caryophylli	f ʒj	f ʒij.
———— Cascarillæ	f ʒiss	f ʒij.
———— Catechu compositum	f ʒj	f ʒiij.
———— Cinchonæ	f ʒj	f ʒiij.
———— Cuspariæ	f ʒiss	f ʒij.
———— Digitalis	f ʒss	f ʒj.
———— Diosmæ	f ʒij	f ʒiij.
———— Gentianæ compositum	f ʒiss	f ʒij.
———— Lupuli	f ʒi	f ʒiss.
———— Quassiæ	f ʒiss	f ʒij.
———— Krameriæ	f ʒiss	f ʒij.
———— Pareiræ	f ʒi	f ʒiss.
———— Rhei	f ʒj	f ʒiij.
———— Rosæ compositum	f ʒj	f ʒiss.
———— Scoparii	f ʒj	f ʒij.
———— Sennæ compositum	f ʒiij	f ʒiv.
———— Serpentariæ	f ʒi	f ʒij.
———— Simarubæ	f ʒj	f ʒij.
———— Valerianæ	f ʒiss	f ʒij.
Ipecacuanha (Diaphoretic)	gr. ss	gr. ij.
Ipecacuanha (Emetic)	gr. v	ʒss.
Kino	gr. x	ʒss.
Krameria	gr. x	gr. xx.

Lavandulæ Flores	᠑j	to	᠑j.
Lauri Baccæ et Folia	gr. x		᠑ss.
Lichen	᠑j		᠑j.
Liquor Ammoniaë	℥x		℥xxx.
————— Acetatis	f᠑iv		f᠑vi.
————— Sesquicarbonatis	f᠑ss		f᠑j.
————— Calcis	f᠑j		f᠑vj.
————— Calcii Chloridi	℥xl		f᠑ij.
————— Hydrargyri Bichloridi	f᠑ss		f᠑j.
————— Potassæ	℥x		f᠑ss.
————— Arsenitis	℥iv		℥xxx.
————— Carbonatis	℥x		f᠑j.
————— Potassii Iodidi compositus	℥v		℥x.
Magnesia	᠑ss		᠑j.
————— Carbonas	᠑j		᠑j.
————— Sulphas	᠑ss		᠑iss.
Malva	᠑j		᠑j.
Manna	᠑ss		᠑ij.
Mastiche	gr. x		᠑ss.
Marrubium	᠑j		᠑j.
Mentha piperita	gr. x		᠑j.
————— viridis	gr. x		᠑j.
Menyanthes	᠑ss		᠑j.
Mezerei Cortex	gr. j		gr. x.
Mistura Acaciæ	f᠑j		f᠑j.
————— Ammoniaci	f᠑ss		f᠑j.
————— Assafoetidæ	f᠑ss		f᠑j.
————— Camphoræ	f᠑j		f᠑ij.
————— Cascarillæ Compositæ	f᠑i		f᠑iss.
————— Cretæ	f᠑j		f᠑ij.
————— Ferri composita	f᠑j		f᠑ij.
————— Gentianæ composita	f᠑i		f᠑ij.
————— Guaiaci	f᠑ss		f᠑ij.
————— Moschi	f᠑j		f᠑ij.
————— Spiritûs Vini Gallici	f᠑ss		f᠑iss.
Morphiæ Acetas	gr. $\frac{1}{8}$		gr. $\frac{1}{4}$.
————— Hydrochloras	$\frac{1}{8}$		$\frac{1}{4}$

Moschus	gr. ij	to ʒj.
Myristicæ Nuclei.....	gr. v	gr. x.
Myrrha.....	gr. x	ʒj.
Oleum Anisi.....	℥j	℥v.
—— Anthemidis	℥j	℥v.
—— Carui.....	℥j	℥v.
—— Caryophylli	℥j	℥v.
—— Cinnamomi	℥j	℥iij.
—— Juniperi.....	℥j	℥v.
—— Lavandulæ	℥j	℥v.
—— Menthæ piperitæ	℥j	℥iij.
———— viridis	℥j	℥v.
———— pulegii.....	℥j	℥v.
—— Origani	℥j	℥iij.
—— Pimentæ	℥j	℥iij.
—— Ricini	fʒiv	fʒiss.
—— Rosmarini.....	℥ij	℥v.
—— Succini	℥v	℥x.
—— Terebinthinæ purificatum (diuretic)	℥x	fʒss.
———————— (vermifuge)	fʒij	fʒj.
Olibanum	gr. x	ʒss.
Opium	gr. ss	gr. v.
Opoponax	gr. x	ʒss.
Oxymel	fʒj	fʒj.
—— Scillæ	fʒss	fʒij.
Pilulæ Aloës compositæ	gr. x	ʒj.
———— cum Myrrhâ	gr. x	ʒj.
—— Cambogiæ compositæ	gr. x	ʒj.
—— Conii compositæ	gr. iij	gr. v.
—— Ferri compositæ	gr. x	ʒj.
—— Galbani compositæ	gr. x	ʒj.
—— Hydrargyri (alterative)	gr. iv	gr. vi.
—————— (purgative)	gr. x	ʒj.
———— Chloridi compositæ ..	gr. v	gr. x.
—— Ipecacuanhæ compositæ	gr. v	gr. x.
—— Rhei compositæ	gr. x	gr. xx.
—— Saponis compositæ	gr. iij	gr. x.

Pilulæ Sagapeni compositæ.....	gr. v	to gr. x.
—— Scillæ compositæ	gr. x	ʒj.
—— Styracis compositæ	gr. iij	gr. x.
Pimentæ Baccæ	gr. v	ʒj.
Piperis longi Fructus	gr. v	ʒj.
—— nigri Baccæ	gr. v	ʒj.
Plumbi Acetas	gr. ss	gr. j.
Potassæ Acetas	ʒj	ʒj.
—— Bicarbonas	gr. x	ʒss.
—— Bisulphas	gr. x	ʒj.
—— Bitartras.....	ʒj	ʒss.
—— Carbonas	gr. x	ʒss.
—— Nitras.....	gr. x	ʒss.
—— Sulphas	gr. x	ʒss.
—— Tartras	ʒj	ʒj.
Potassii Bromidum	gr. iij	gr. x.
Potassii Iodidum	gr. v	gr. x.
Pulvis Aloës compositus	gr. x	ʒj.
—— Antimonii compositus	gr. v	gr. x.
—— Cinnamomi compositus.....	gr. v	gr. x.
—— Cretæ compositus.....	gr. v	gr. xxx.
—— Cretæ compositus cum Opio	gr. v	gr. xxx.
—— Jalapæ compositus	gr. xx	ʒij.
—— Ipecacuanhæ compositus	gr. v	ʒj.
—— Kino compositus	gr. v	ʒj.
—— Scammonii compositus	gr. v	ʒj.
—— Tragacanthæ compositus	gr. x	ʒj.
Quassiæ Lignum	gr. v	ʒss.
Quercus Cortex	gr. x	ʒss.
Quinæ Disulphas.....	gr. iij	gr. x.
Rhamni Baccæ.....	ʒj	ʒij.
Rhei Radix	gr. x	ʒss.
Rosæ caninæ Pulpa	ʒj	ʒj.
Rosæ centifoliæ Petala.....	ʒj	ʒj.
—— Gallicæ Petala.....	ʒj	ʒj.
Rosmarini Cacumina	gr. x	ʒss.
Rutæ Folia	gr. xv	ʒij.

Sabinæ Folia	gr. x	to ʒss.
Sagapenum	gr. x	ʒss.
Sapo durus	gr. v	ʒss.
Sarzæ Radix	ʒj	ʒj.
Sassafras Lignum.....	ʒj	ʒj.
Scammonium	gr. v	ʒj.
Scillæ Radix recens.....	gr. v	gr. xv.
———— exsiccata.....	gr. j	gr. iij
Senegæ Radix	ʒj	ʒij.
Sennæ Folia.....	ʒj	ʒj.
Serpentariæ Radix	gr. x	ʒss.
Simarubæ Cortex	gr. x	ʒss.
Sinapis Semina.....	ʒj	ʒss.
Sodæ Biboras	gr. x	ʒss.
———— Carbonas	gr. x	ʒss.
———— exsiccata	gr. v.	gr. xv.
———— Potassio-tartras	ʒij	ʒj.
———— Sesquicarbonas	gr. x	ʒss.
———— Sulphas	ʒss	ʒij.
Spigeliæ Radix	gr. x	ʒij.
Spiritus Ætheris nitrici	℥x	℥xl.
———— sulphurici compositus ..	fʒss	fʒij.
———— Ammoniæ aromaticus.....	fʒss	fʒj.
———— foetidus	℥xv	℥xxx.
———— Anisi	fʒij	fʒss.
———— Armoraciæ compositus	fʒij	fʒss.
———— Carui	fʒij	fʒss.
———— Cinnamomi.....	fʒij	fʒss.
———— Juniperi compositus	fʒij	fʒj.
———— Menthæ piperitæ	fʒij	fʒss.
———— pulegii	fʒij	fʒss.
———— viridis	fʒij	fʒss.
———— Myristicæ	fʒij	fʒss.
———— Pimentæ	fʒij	fʒss.
Stannum	ʒi	ʒss.
Strychnia	gr. $\frac{1}{16}$	gr. $\frac{1}{8}$.
Sulphur	ʒj	ʒij.
Syrupus Papaveris	fʒj	fʒj.
———— Rhamni	fʒss	fʒj.

Syrupus Sarzæ	f3j	to	f3ss.
———— Sennæ	f3ij		f3iv.
Tabaci Folia	gr. ss		gr. v.
Tamarindi Pulpa	3ss		3j.
Terebinthina Canadensis	3j		3j.
———— Chia	3j		3j.
———— vulgaris	3j		3j.
Tiglii Oleum	℥j		℥ij.
Tinctura Aloes	f3ss		f3iss.
———— composita	f3j		f3ij.
———— Ammoniae composita	℥v		℥x.
———— Assafoetidae	f3ss		f3iss.
———— Aurantii	f3ij		f3iij.
———— Balsami Tolutani	f3j		f3ij.
———— Benzoini composita	f3ss		f3ij.
———— Calumbæ	f3j		f3iij.
———— Camphoræ composita	f3j		f3iij.
———— Cantharidis	℥x		f3j.
———— Capsici	℥x		f3j.
———— Cardamomi	f3j		f3ij.
———— composita	f3j		f3ij.
———— Cascarillæ	f3j		f3ij.
———— Castorei	℥xx		f3ij.
———— Catechu	f3j		f3iij.
———— Cinchonæ	f3j		f3iij.
———— composita	f3j		f3iij.
———— Cinnamomi	f3j		f3ij.
———— composita	f3j		f3ij.
———— Colchici	℥xx		℥xxx.
———— Colchici composita	f3ss		f3j.
———— Conii	f3ss		f3j.
———— Cubebæ	f3ss		f3j.
———— Digitalis	℥x		℥xl.
———— Ferri ammonio-chloridi	f3ss		f3ij.
———— sesquichloridi	℥x		f3j.
———— Gentianæ composita	f3j		f3ij.
———— Guaiaci	f3j		f3iij.
———— composita	f3ss		f3j.

Tinctura Hellebori	f3ss	to f3j.
———— Hyoscyami	f3ss	f3ij.
———— Jalapæ	f3j	f3ss.
———— Kino	f3j	f3ij.
———— Lavandulæ composita	f3ss	f3ij.
———— Lupuli	f3ss	f3ij.
———— Myrrhæ	f3ss	f3j.
———— Opii	℥x	f3j.
———— Rhei composita	f3ij	f3iss.
———— Scillæ	℥x	f3ss.
———— Sennæ composita	f3ij	f3j.
———— Serpentariæ	f3j	f3ij.
———— Valerianæ	f3j	f3ij.
———— ————— composita	f3ss	f3j.
———— Zingiberis	f3j	f3ij.
Tormentillæ Radix	gr. x	3ss.
Tragacantha	gr. x	3j.
Valerianæ Radix	3j	3ij.
Veratria	gr. $\frac{1}{8}$	gr. $\frac{1}{4}$.
Veratri Radix	gr. ij	gr. v.
Vinum Aloës	f3j	f3ij.
———— Antimonii potassio-tartratis	℥xv	f3j.
———— Colchici	℥xxx	f3j.
———— Ipecacuanhæ (Diaphoretic)	℥xx	℥xl.
———— ————— (Emetic)	f3ij	f3ss.
———— Opii	℥x	f3j.
———— Veratri	℥v	℥x.
Uva Ursi	gr. x	3j.
Zinci Oxydum	gr. j	gr. vi.
———— Sulphas	gr. j	gr. v.
———— ————— (Emetic)	gr. x	3ss.

INDEX.

	Page		Page
A.		Argenti Cyanidum 198	
Acetum Cantharidis.....	51	— Nitras 195	
— Colchici	51	B.	
— Destillatum.....	47	Barii Chloridum	201
— Scillæ	52	Bismuthi Trisnitræ	203
Acidum Aceticum	48	C.	
— Benzoicum	53	Calamina præparata	290
— Citricum	56	Calcii Chloridum	208
— Hydrochloricum.....	59	Calx	205
— Hydrochloricum dilu-		— Chlorinata	210
lutum	62	Carbo Animalis purificatus ...	114
— Hydrocyanicum dilu-		Cataplasma Conii	121
tum	63	— Fermenti	122
— Nitricum.....	69	— Lini	122
— Nitricum dilutum ...	72	— Sinapis.....	122
— Phosphoricum dilu-		Ceratum	123
tum	72	— Calaminæ	123
— Sulphuricum dilutum	75	— Cantharidis	124
— Tartaricum	78	— Cetacei	124
Aconitina	92	— Hydrargyri C.	125
Æther Sulphuricus	83	— Plumbi Acetatis.....	125
Alcohol.....	315	— — compositum .	126
Alumen Exsiccatum.....	186	— Resinæ	126
Ammoniaë Sesquicarbonas ...	96	— Sabinæ	127
Antimonii Oxysulphuretum... 188		— Saponis	127
— Potassio-tartras ...	190	Confectio Amygdalæ	128
Aqua Anethi	117	— Aromatica	128
— Carui	118	— Aurantii	129
— Cinnamomi.....	119	— Cassiæ	130
— Destillata	116	— Opii	130
— Florum Aurantii.....	118	— Piperis Nigri	131
— Fœniculi.....	118	— Rosæ Caninæ	132
— Menthæ Piperitæ	119	— Rosæ Gallicæ	132
— Menthæ Pulegii.....	119	— Rutæ	133
— Menthæ Viridis	120	— Scammonii	133
— Pimentæ.....	120	— Sennæ	134
— Rosæ	120		
— Sambuci	121		

	Page		Page
Cornu Ustum	115	Extractum Cinchonæ C.	160
Creta Præparata	212	————— L.	160
Cupri Ammonio-sulphas	213	————— O.	160
D.		————— Colchici aceticum .	161
Decoctum Aloës compositum	135	————— Colchici cormi.....	156
————— Amyli	136	————— Colocynthis	161
————— Cetrariæ	136	————— Colocynthis C. ...	162
————— Chimaphilæ	137	————— Conii	156
————— Cinchonæ cordifo-		————— Digitalis	157
————— liæ	137	————— Elaterii	162
————— Cinchonæ lancifo-		————— Gentianæ	163
————— liæ	138	————— Glycyrrhizæ	164
————— Cinchonæ oblongi-		————— Hæmatoxyli	164
————— foliæ	138	————— Hyoscyami.....	157
————— Cydoniæ	139	————— Jalapæ	166
————— Dulcamaræ.....	140	————— Lactucæ	157
————— Granati	140	————— Lupuli.....	165
————— Hordei	141	————— Opii purificatum ..	167
————— Hordei C.	141	————— Papaveris	167
————— Malvæ C.	141	————— Pareiræ	165
————— Papaveris	142	————— Rhei	168
————— Quercus	142	————— Sarzæ	165
————— Sarzæ	143	————— Stramonii	168
————— Sarzæ C.	143	————— Taraxaci.....	165
————— Scoparii C.	144	————— Uvæ Ursi	166
————— Senegæ	144	F.	
————— T tormentillæ	145	Ferri Ammonio-chloridum ...	223
————— Veratri	146	————— Iodidum	225
————— Ulmi	145	————— Potassio-tartras	221
————— Uvæ Ursi	145	————— Sesquioxylum	218
E.		————— Sulphas	215
Emplastrum Ammoniaci.....	146	G.	
————— Ammoniaci cum		Gummi-resinæ.....	352
————— Hydrargyro... ..	147	H.	
————— Belladonnæ.....	147	Hydrargyri Ammonio-chlori-	
————— Cantharidis.....	148	————— dum.....	233
————— Cerae	148	————— Bichloridum ...	228
————— Galbani	149	————— Bicyanidum.....	239
————— Hydrargyri.....	149	————— Biniodidum.....	242
————— Opii.....	150	————— Binoxylum	237
————— Picis	150	————— Bisulphuretum ..	243
————— Plumbi	151	————— Chloridum	231
————— Resinæ	151	————— Iodidum	241
————— Saponis	152	————— Nitrico-oxylum	238
Enema Aloës	152	————— Oxylum	235
————— Colocynthis	153	————— Sulphuretum cum	
————— Opii.....	153	————— sulphure	244
————— Tabaci	153	Hydrargyrum cum Cretâ ...	227
————— Terebinthinæ	154	I.	
Extractum Aconiti	155	Infusum Anthemidis	169
————— Aloës purificatum	159		
————— Belladonnæ	156		

	Page		Page
Infusum Armoraciae C.	170	Liquor Sodae Chlorinatae.....	285
— Aurantii C.....	170	— Sodae Effervescens ...	284
— Calumbae	171		
— Caryophylli.....	171	M.	
— Cascarillae	171	Magnesia	245
— Catechu C.	172	Magnesiæ Carbonas.....	246
— Cinchonæ	172	Mel Boracis	184
— Cuspariæ.....	173	— Rosæ.....	184
— Digitalis	173	Mistura Acaciæ	290
— Diosmæ	173	— Ammoniaci	291
— Gentianæ C.	174	— Amygdalæ	291
— Krameriae	174	— Assafoetidæ.....	292
— Lini C.	175	— Camphoræ	292
— Lupuli	175	— Cascarillæ C.	293
— Pareiræ	175	— Cretæ	293
— Quassiæ	176	— Ferri C.	294
— Rhei	176	— Gentianæ C.	295
— Rosæ C.	176	— Guaiaci	295
— Sennæ C.	177	— Moschi	296
— Scoparii	177	— Spiritus Vini Gallici	296
— Serpentariæ	178	Morphia	101
— Simarubæ	178	Morphiæ Acetas	102
— Valerianæ	179	Morphiæ Hydrochloras	103
L.		O.	
Linimentum Æruginis.....	179	Oleum Æthereum	88
— Ammoniae	180	— Anisi	297
— Ammoniae Ses.	180	— Anthemidis	297
— Camphoræ	181	— Carui	297
— Camphoræ C....	181	— Juniperi	298
— Hydrargyri C.	182	— Lavandulæ	298
— Opii	182	— Menthæ Piperitæ ...	298
— Saponis	183	— Menthæ Pulegii	298
— Terebinthinæ... ..	183	— Menthæ Viridis	299
Liquor Aluminis C.	187	— Origani	299
— Ammoniae	94	— Pimentæ	299
— Ammoniae Acetatis... ..	99	— Rosmarini	299
— Ammoniae Sesquicarb.	99	— Sambuci	300
— Argenti nitratis	197	— Succini.....	301
— Barii Chloridi	203	— Terebinthinæ purifica-	
— Calcii Chloridi.....	209	tum	301
— Calcis	207	Oxymel.....	185
— Cupri Ammonio-sul-		— Scillæ	185
phatis	214		
— Hydrargyri Bichloridi	230	P.	
— Plumbi diacetatis ...	251	Pilulæ Aloës C.	302
— Plumbi diacetatis di-		— Aloës cum Myrrhâ... ..	302
lutus.....	252	— Cambogiæ C.	303
— Potassæ	262	— Conii C.	303
— Potassæ Arsenitis ...	199	— Ferri C.	304
— Potassæ Carbonatis ...	258	— Galbani C.	305
— Potassæ Effervescens	261	— Hydrargyri	305
— Potassii Iodidi C. ...	274	— Hydrargyri Chloridi C.	306

	Page
Tinctura Ferri Sesquichloridi	220
— Gallæ	342
— Gentianæ C.	343
— Guaiaci	343
— Guaiaci C.	344
— Hellebori	344
— Hyoscyami	345
— Jalapæ	345
— Iodinii C.	346
— Kino	346
— Lavandulæ C.	346
— Lupuli	347
— Myrrhæ... ..	347
— Opii	347
— Rhei C.	348
— Scillæ	349
— Sennæ C.	349
— Serpentariæ	349
— Valerianæ	350
— Valerianæ C.	350
— Zingiberis	350

V.

Vegetabilia	351
Veratria	112
Vinum Aloës	353
— Antimonii Potassio-	
Tartratis	193
— Colchici	353
— Ipecacuanhæ	354
— Opii	355
— Veratri	355

U.

Unguentum Antimonii Potas-	
sio-tartratis ...	356

	Page
Unguentum Cantharidis	356
— Cetacei	357
— Creasoti	357
— Elemi	358
— Gallæ C.	358
— Hydrargyri for-	
tius	359
— Hydrargyri mi-	
tius	360
— Hydrargyri Ni-	
tratis	360
— Hydrargyri Ni-	
trico-oxydi ...	361
— Hydrargyri Io-	
didi	361
— Hydrargyri Bin-	
iodidi	361
— Hydrargyri Am-	
monio-chloridi	362
— Iodinii C.	362
— Picis Liquidæ ...	362
— Picis nigræ	363
— Plumbi C.	363
— Plumbi Iodidi ...	364
— Sambuci	364
— Sulphuris	364
— Sulphuris C. ...	365
— Veratri	365
— Zinci	365

Z.

Zinci Oxydum	289
— Sulphas	286

ENGLISH INDEX.

A.		Page			Page
Acetate of Lead		249	Chloride of Lead		253
————— Morphia		102	————— Mercury.....		231
————— Potash.....		265	Chlorinated Lime.....		210
Acetic Acid		48	Cinnamon Water		119
————— Extract of Meadow-			Citric Acid		54
Saffron		161	Compound Cerate of Lead ...		126
Aconitina		92	————— Mercury		125
Æthereal Oil		88	————— Decoction of Aloes		135
Alcohol.....		315	————— Barley		141
Ammonio-chloride of Iron ...		223	————— Broom.		144
————— Mercury		233	————— Mallow		141
————— Sulphate of Copper		213	————— Sarsa-		
Aromatic Confection		128	parilla		143
Aromatic Spirit of Ammonia		317	————— Extract of Colo-		
			cynth		162
			————— Liniment of Cam-		
			phor		181
			————— Mixture of Casca-		
			rilla.....		293
			————— Gentian		295
			————— Iron ...		294
			————— Ointment of Galls		358
			————— Iodine		362
			————— Lead		363
			————— Sulphur		365
			————— Pills of Aloes		302
			————— Camboge ..		308
			————— Chloride of		
			Mercury		306
			————— Galbanum..		305
			————— Hemlock...		803
			————— Iron.....		304
			————— Ipecacuan-		
			ha		307
			————— Rhubarb ...		307
			————— Sagapenum		307
			————— Soap		308
			————— Squill		309
			————— Storax		309
			————— Powder of Aloes ...		310
			————— Antimony		194
			————— Chalk ...		311
			————— with		
			Opium		312
			————— Cinnamon		310
			————— Jalap.....		312
B.					
Benzoic Acid		53			
Bicarbonate of Potash.....		259			
Bichloride of Mercury.....		229			
Bicyanide of Mercury		239			
Biniodide of Mercury		242			
Binoxide of Mercury		237			
Bisulphate of Potash		268			
Bisulphuret of Mercury		243			
Bromide of Potassium.....		271			
Burnt Horn		115			
C.					
Carbonate of Magnesia		246			
————— Potash		256			
————— Soda		277			
Carraway Water		118			
Cataplasm of Hemlock.....		121			
————— Linseed		122			
————— Mustard		122			
————— Yest.....		122			
Cerate		123			
Cerate of Acetate of Lead ...		125			
————— Calamine		123			
————— Cantharides.....		124			
————— Resin		126			
————— Savin		127			
————— Soap.....		127			
————— Spermaceti		124			
Chloride of Barium		201			
————— Calcium		208			

	Page		Page
Compound Powder of Ipecacu-		Decoction of Oak Bark	142
anha ...	312	————— Pomegranate ...	140
————— Kino	313	————— Poppy	142
————— Scammony	313	————— Quina	139
————— Traga-		————— Sarsaparilla	143
canth...	314	————— Senega.....	144
————— Solution of Alum ..	187	————— Starch	136
————— Iodide of		————— Tormetil	145
Potassium	274	————— White Hellebore	146
————— Spirit of Horse-		————— Whortleberry ...	148
radish	318	————— Winter-green ...	137
————— Juniper..	320	————— Woody - Night-	
————— Sulphuric		shade.....	140
Æther...	92	Dill Water	117
————— Tinct. of Aloes	331	Diluted Hydrochloric Acid...	62
————— Ammonia..	332	————— Hydrocyanic Acid ...	63
————— Benzoin ...	333	————— Nitric Acid.....	72
————— Camphor...	335	————— Phosphoric Acid.....	72
————— Cardamom	337	————— Solution of Diacetate	
————— Cinchona...	339	of Lead	252
————— Cinnamon..	340	————— Sulphuric Acid	75
————— Gentian ...	343	Distilled Vinegar.....	47
————— Guaiacum .	344	————— Water	116
————— Iodine	346	Disulphate of Quina	106
————— Lavender...	346	Dried Alum	186
————— Meadow-		Dried Carbonate of Soda.....	279
Saffron...	341		
————— Rhubarb...	348	E.	
————— Senna	349	Effervescing Solution of Pot-	
————— Valerian ...	350	ash	261
Confection, Aromatic	128	————— Solution of Soda	284
————— of Almond	128	Elder Water.....	121
————— Black Pepper ...	131	Enema of Aloes	152
————— Cassia	130	————— Colocynth	153
————— Dog Rose.....	132	————— Opium.....	153
————— Opium	130	————— Tobacco	153
————— Orange.....	129	————— Turpentine	154
————— Red Rose.....	132	Extract of Aconite	155
————— Rue	133	————— Aloes, purified.....	159
————— Scammony	133	————— Cinchona, Heart-	
————— Senna	134	leaved	160
Cyanide of Silver.....	198	—————, Lance-	
		leaved	160
D.		—————, Oblong-	
Decoction of Barley.....	141	leaved	160
————— Cinchona, Heart-		————— Colocynth	161
leaved	138	————— compound	162
—————, Lance-		————— Dandelion	165
leaved	138	————— Deadly Nightshade	156
—————, Oblong-		————— Elaterium	162
leaved	138	————— Foxglove	157
————— Elm Bark	145	————— Gentian	163
————— Liverwort	136	————— Hemlock.....	156

	Page		Page
Extract of Henbane	157	Infusion of Rose, compound...	176
———— Hops	165	———— Senna, compound..	177
———— Jalap	166	———— Serpentry	178
———— Lettuce	157	———— Simaruba	178
———— Liquorice.....	164	———— Valerian	179
———— Logwood	164	Iodide of Iron	225
———— Meadow Saffron ...	156	———— Mercury	241
————, Ace-		———— Lead	254
———— tic	161	———— Potassium	273
———— Opium, purified ...	167		
———— Pareira	165	L:	
———— Poppy	167	Lime	205
———— Rhubarb	168	—— Water	207
———— Sarsaparilla	165	Liniment of Ammonia	180
———— Thorn Apple	168	———— Camphor	181
———— Whortleberry	166	————, com-	
		———— pound	181
F.		———— Mercury	182
Fennel Water	118	———— Opium.....	182
Fetid Spirit of Ammonia.....	317	———— Sesquicarbonate	
		———— of Ammonia..	180
G.		———— Soap	183
Gum-Resins	352	———— Turpentine	183
		———— Verdigris.....	179
H.			
Honey of Borax	184	M.	
Honey of Rose	144	Magnesia	245
Hydrated Oxide of Lead ...	255	Materia Medica	3
Hydrate of Potash	263	Mercury with Chalk	227
Hydrochlorate of Morphia ...	103	Milder Ointment of Mercury	360
Hydrochloric Acid	59	Mixture of Acacia	290
		———— Almond	291
I.		———— Ammoniacum ...	291
Infusion of Broom	177	———— Assafoetida	292
———— Buchu.....	173	———— Camphor.....	292
———— Calumba.....	171	———— Cascarilla, com-	
———— Cascarilla	171	———— pound	293
———— Catechu	172	———— Chalk	293
———— Chamomile	169	———— Gentian, compound	295
———— Cinchona	172	———— Guaiacum	295
———— Cloves.....	171	———— Iron, compound ...	244
———— Cusparia	173	———— Musk	296
———— Foxglove.....	173	———— Spirit of French	
———— Gentian, compound	174	———— Wine	296
———— Hops	175	Morphia	101
———— Horseradish	170		
———— Linseed, compound	175	N.	
———— Orange Peel, com-		Nitrate of Silver	195
———— pound	170	Nitric Acid	69
———— Pareira	175	Nitric Oxide of Mercury.....	238
———— Quassia	176		
———— Rhatany	174	O.	
———— Rhubarb	176	Oil of Amber	301

	Page
Oil of Anise	297
— Carraway	297
— Chamomile.....	297
— Elder Flowers	300
— Juniper	298
— Lavender	298
— Marjoram	299
— Pennyroyal.....	298
— Peppermint	298
— Rosemary	299
— Spearmint	299
— Turpentine, purified ...	301
Ointment of Ammonio-chlo-	
ride of Mercury	362
— Binioidide of Mer-	
cury	361
— Black Pitch.....	363
— Cantharides.....	356
— Creasote	357
— Elder	364
— Elemi	358
— Iodide of Lead... ..	364
— Mercury	361
— Nitrate of Mer-	
cury	360
— Nitric-oxide of	
Mercury	361
— Potassio-tartrate	
of Antimony... ..	356
— Spermaceti	357
— Sulphur	364
— Tar	362
— White Hellebore	365
— Zinc.....	365
Orange Flower Water	118
Oxide of Mercury	237
— Zinc	289
Oxymel.....	185
— of Squill	185
Oxysulphuret of Antimony...	188

P.

Pennyroyal Water	119
Peppermint Water	119
Pills of Aloes with Myrrh ..	302
— Iodide of Mercury ...	306
— Mercury	306
Pimenta Water	120
Plaster of Ammoniacum.....	
—, with	
Mercury	147
— Cantharides.....	148
— Deadly Nightshade	147

	Page
Plaster of Galbanum	149
— Lead	157
— Mercury	149
— Opium.....	150
— Pitch	150
— Resin	151
— Soap.....	152
— Wax	148
Potash with Lime	264
Potassio-tartrate of Antimony	190
— Iron	221
— Soda ...	283
Prepared Calamine.....	290
— Chalk	213
— Shells	116
Preparation of Vegetables ...	
Purified Animal Charcoal ...	114
— Oil of Turpentine ...	301

R.

Rose Water	120
------------------	-----

S.

Sesquicarbonate of Ammonia	96
— Soda	280
Sesquioxide of Iron	218
Solution of Acetate of Ammo-	
nia	99
— Ammonia	94
— Ammonio-sulphate	
of Copper	214
— Arsenite of Potash	199
— Bichloride of Mer-	
cury	230
— Carbonate of Pot-	
ash	258
— Chloride of Barium	203
— Calcium	209
— Chlorinated Soda... ..	285
— Diacetate of Lead.. ..	251
— Nitrate of Silver... ..	197
— Potash.....	262
— Sesquicarbonate of	
Ammonia	99
Spearmint Water	120
Spirit of Ammonia	316
— Anise	318
— Carraway	319
— Cinnamon	319
— Lavender	320
— Nitric Æther	90
— Nutmeg	322
— Pennyroyal.....	321

	Page		Page
Spirit of Peppermint.....	321	Tincture of Cascarella	337
—— Pimenta	322	—— Castor	338
—— Rosemary	322	—— Catechu	338
—— Spearmint	321	—— Cinchona	339
Stronger Ointment of Mercury	359	—— Cinnamon	340
Strychnia	109	—— Cubebs	342
Sulphate of Iron	215	—— Foxglove	342
Sulphate of Potash	266	—— Galls	342
—— Soda	281	—— Ginger	350
—— Zinc	286	—— Guaiacum	343
Sulphuret of Mercury with		—— Hellebore	344
Sulphur	244	—— Hemlock	345
Sulphuret of Potassium	276	—— Henbane	345
Sulphuric Æther	83	—— Hop	347
Syrup	323	—— Jalap	345
—— of Buckthorn	327	—— Kino	346
—— Ginger	330	—— Meadow Saffron... 341	
—— Lemons	325	—— Myrrh..... 347	
—— Marshmallow..... 323		—— Opium	347
—— Mulberry	326	—— Orange	333
—— Orange	324	—— Serpentry	349
—— Poppy	326	—— Sesquichloride of	
—— Red Poppy..... 327		Iron	220
—— Rose	328	—— Squill	349
—— Saffron	325	—— Valerian... 350	
—— Sarsaparilla..... 328		Trisnitrate of Bismuth	203
—— Senna	329		
—— Tolu	329		
T.		V.	
Tartaric Acid	78	Vegetables	351
Tartrate of Potash	269	Veratria	112
Tincture of Aloes	331	Vinegar of Cantharides..... 51	
—— Ammonia-chloride		—— Meadow Saffron ... 51	
of Iron..... 225		—— Squill	52
—— Assafoetida	332		
—— Balsam of Tolu ... 333		W.	
—— Calumba	334	Weights, Measures, &c. 1	
—— Camphor..... 334		Wine of Aloes	353
—— Cantharides	335	—— Ipecacuanha	354
—— Capsicum	336	—— Meadow Saffron..... 353	
—— Cardamom	336	—— Opium	355
		—— White Hellebore ... 353	

THE END.

PRINTED BY RICHARD TAYLOR,
RED LION COURT, FLEET STREET.

VALUABLE STANDARD WORKS,

ON

MEDICINE,
SURGERY,

MIDWIFERY,
CHEMISTRY, &c.

PRINTED FOR

LONGMAN, REES, ORME, BROWN, GREEN, AND LONGMAN,
Paternoster-Row.

A TREATISE ON OPERATIVE SURGERY.

By W. P. COCKS, Surgeon,

Author of "Illustrations of Mr. Samuel Cooper's Surgical Dictionary," comprised
in above 400 Plates, with Anatomical Descriptions.

1 vol. 8vo. illustrated by Twelve Plates.—*In the Press.*

*** This Work will be a Comprehensive Manual of Practical Surgery, embracing illustrated details of all the Operations, Major and Minor, as they are performed on the Living and Dead Subjects by the most eminent Surgeons in this country, on the European Continent, and in America.

ELEMENTS OF MEDICAL JURISPRUDENCE.

By T. R. BECK, M.D.

Professor of the Institutes of Medicine, and Lecturer on Medical Jurisprudence, in the College of Physicians and Surgeons of the Western District of the State of New York, &c. ; and

J. B. BECK, M.D.

Professor of Materia Medica and Medical Jurisprudence in the College of Physicians and Surgeons, New York, one of the Physicians to the New York Hospital, &c. &c.

8vo. 5th Edition, brought down to the present time, including the Notes
of Dr. Dunlop and Dr. Darwall. 2ls.

"Beck's Medical Jurisprudence, in its present enlarged form, has been rendered quite an Encyclopædia on the subject: it is an admirable book of reference, and ought to be in the hands of every member of the profession."—*Med. Gaz.* March 19, 1836.

FIRST LINES OF THE PRACTICE OF SURGERY;

Explaining and illustrating the Doctrines relative to the Principles, Practice, and
Operations of Surgery, as taught

By SAMUEL COOPER,

Surgeon to the North London Hospital, Professor of Surgery and Clinical Surgery in
the University of London, &c.

By the same Author,

Dictionary of Practical Surgery. 1 thick vol. 8vo. 28s.

Illustrations of Ditto, in above 400 Plates, with Anatomical Descriptions.

By W. P. COCKS, Surgeon.

2 vols. 8vo.—Vol. I. 2l. 2s. ; Vol. II. 3l. 7s. 6d.

Mr. Abernethy.

SURGICAL AND PHYSIOLOGICAL WORKS

OF

JOHN ABERNETHY, F.R.S. &c.

4 Vols. 8vo. 2l. 5s. 6d. Consisting of the following Treatises and Lectures, which may also be had *separately* :—

- | | |
|---|---|
| <p>I.—On the CONSTITUTIONAL ORIGIN and TREATMENT of LOCAL DISEASES, and on ANEURISMS, 8s.</p> <p>II.—On DISEASES resembling SYPHILIS, and on DISEASES of the URETHRA, 6s.</p> | <p>III.—On INJURIES of the HEAD, and MISCELLANEOUS SUBJECTS, 7s.</p> <p>IV.—On LUMBAR ABSCESES and TUMORS, 6s.</p> <p>V.—LECTURES ON SURGERY, 8s.</p> <p>VI.—PHYSIOLOGICAL LECTURES, 10s. 6d.</p> |
|---|---|

Dr. Ainslie.

MATERIA INDICA;

Or, some Account of those Articles which are employed by the Hindoos, and other Eastern Nations, in their Medicine, Arts, &c.

By WHITE LAW AINSLIE, M.D. M.R.A.S. In 2 vols. 8vo. 2l.

Dr. Aldis.

INTRODUCTION TO HOSPITAL PRACTICE,

In various Complaints; being a Clinical Report of Fever, Gout, Rheumatism, Cholera, &c. &c.; with Remarks on their Pathology and Treatment.

By C. J. B. ALDIS, M.A. M.B. and L.M. 1 vol. 8vo. 5s.

Dr. Arnott.

ELEMENTS OF PHYSICS,

Or Natural Philosophy, General and Medical, written in plain or non-technical language.

By NEIL ARNOTT, M.D.

8vo. many Wood Engravings, 5th edit. Vol. I. 21s. and Vol. II. Part I. 10s. 6d.

The concluding Part of Vol. II. of the work is in progress.

Dr. Abercrombie.

PATHOLOGICAL AND PRACTICAL

RESEARCHES ON DISEASES OF THE BRAIN AND SPINAL CORD.

By JOHN ABERCROMBIE, M.D. Oxon. and Edin. V.P.R.S.E. &c. &c.

12mo. 3d Edit. greatly enlarged, 5s. 6d.

Dr. Addison & Mr. Morgan.

AN ESSAY ON THE

OPERATION OF POISONOUS AGENTS UPON THE LIVING BODY.

By JOHN MORGAN, F.L.S. Surgeon to Guy's Hospital; and
THOMAS ADDISON, M.D. Assistant Physician to Guy's Hospital.

8vo. 5s.

Mr. Bacot.

TREATISE ON SYPHILIS,

In which the History, Symptoms, and Method of Treating every Form of that Disease, are fully considered.

By JOHN BACOT, Surgeon to the St. George's and St. James's Dispensary, &c. 8vo. 9s.

*Dr. Baillie.***MORBID ANATOMY**

Of some of the most important Parts of the HUMAN BODY.

By MATTHEW BAILLIE, M.D.

With Observations on Diseased Structures, by JAMES WARDROP, Surgeon to the King.
12mo. 7s.

Life and Works of Dr. Baillie. 2 vols. 8vo. with Portrait, 1l. 5s.

*Mr. Balbirnie.***THE SPECULUM APPLIED TO THE DIAGNOSIS AND TREATMENT OF
THE ORGANIC DISEASES OF THE WOMB :**An Inaugural Dissertation, presented to the University of Glasgow,
for the Degree of Doctor in Medicine.

By JOHN BALBIRNIE, A.M. 8vo. 12s. bds.

*Dr. Bateman.***PRACTICAL SYNOPSIS OF CUTANEOUS DISEASES,**

According to the Arrangement of DR. WILLAN ;

Exhibiting a concise View of the Diagnostic Symptoms, and the Method of Treatment.

By T. BATEMAN, M.D. Seventh Edition, edited by A. TODD THOMSON, M.D. F.L.S.
Professor of Materia Medica in the University of London. 8vo. 15s.Delineations of the Cutaneous Diseases comprised in the Classification of the
late DR. WILLAN. By T. BATEMAN, M.D. &c. 4to. with 72 coloured Plates, 12l. 12s.Atlas of Delineations of Cutaneous Eruptions ; illustrative of the Descrip-
tions in the above Synopsis. By A. TODD THOMSON, M.D. &c. Royal 8vo. with 29 col'd
Plates, 3l. 3s.*Sir George Ballingall.***OUTLINES OF THE COURSE OF LECTURES ON MILITARY SURGERY,**

Delivered in the University of Edinburgh.

By SIR GEO. BALLINGALL, M.D. F.R.S.E. &c. &c. 8vo. 18s.

*Sir Charles Bell.***ANATOMY AND PHYSIOLOGY OF THE HUMAN BODY.**

By JOHN and CHARLES BELL.

7th Edition, systematized by SIR C. BELL.

3 Vols. 8vo. numerous Engravings, 2l. 12s. 6d.

Engravings of the Arteries ; illustrative of the above Work.

By SIR C. BELL. Imperial 8vo. 15s. ; coloured, 1l. 1s.

By the same Author,

Illustrations of the Great Operations of Surgery :

Trepan—Hernia—Amputation—Aneurism—Lithotomy.

4to. with 20 Plates, 3l. 15s. plain ; 5l. 5s. col'd. The Parts *separately*, 15s. plain ; 21s. col'd.

Nervous System of the Human Body.

4to. with Plates, 1l. 16s.

Diseases of the Urethra, Prostate, Rectum, &c.

8vo. 13s. 6d.

Engravings from Specimens of Morbid Parts, in the Author's Collection ;

Selected from the divisions Urethra—Vesica—Ren—Morbosa—Læsa, &c.

Folio, 36s.

Mr. G. H. Bell.

TREATISE ON CHOLERA ASPHYXIA, OR EPIDEMIC CHOLERA,

As it appeared in Asia, and more recently in Europe; &c.

By G. H. BELL, F.R.C.S.E. &c. 2d Edit. 6s.

By the same Author,

Treatise on the Diseases of the Liver, and on Bilious Complaints. 6s.

Dr. Bennett.

MANUAL OF ANATOMY;

From the French of BAYLE.

Revised and improved by W. BENNETT, M.D. 18mo. 7s. 6d.

Dr. Blackall.

OBSERVATIONS ON THE NATURE AND CURE OF DROPSIES.

By JOHN BLACKALL, M.D. 8vo. 4th Edit. 10s. 6d.

Dr. Bright.

REPORTS OF MEDICAL CASES;

Selected with a view of illustrating the Symptoms and Cure of Diseases by a reference to Morbid Anatomy.

By RICHARD BRIGHT, M.D. F.R.S. one of the Physicians to Guy's Hospital. 4to.

Vol. I. embraces Dropsy, Inflammation of the Lungs, Phthisis, and Fever; with 16 coloured Plates, 4l. 4s.

Vol. II. contains Diseases of the Brain and Nervous System; with 38 Plates, 9l. 9s. col'd.

Drs. Bright & Addison.

ELEMENTS OF THE PRACTICE OF PHYSIC.

By RICHARD BRIGHT, M.D. F.R.S., and THOMAS ADDISON, M.D., Lecturers on the Practice of Physic at Guy's Hospital.

In 8vo.

* * * The First Fasciculus will embrace the subject of FEVER.

Dr. Burne.

PRACTICAL TREATISE ON THE TYPHUS OR ADYNAMIC FEVER.

By J. BURNE, M.D. &c.

8vo. 7s. 6d.

Sir B. C. Brodie, Bart.

LECTURES ON THE DISEASES OF THE URINARY ORGANS.

By Sir B. C. BRODIE, Bart. V.P.R.S.

Second Edition, with alterations and additions. 8vo. 8s.

By the same Author,

Pathological and Surgical Observations on Diseases of the Joints.

8vo. 3d Edition, revised, altered, and enlarged. 10s. 6d.

Professor Burns.

PRINCIPLES OF MIDWIFERY;

Including the Diseases of Women and Children.

By JOHN BURNS, M.D. Regius Professor of Surgery, Glasgow.

8vo. 8th Edit. with important Additions, 16s.

By the same Author,

Principles of Surgery,

VOL. I. containing Inflammation, Tumors, Aneurisms, &c. 8vo. 14s.

Dr. Carbutt.

CLINICAL LECTURES, DELIVERED IN THE MANCHESTER INFIRMARY.

By EDWARD CARBUTT, M.D. 12s.

Mr. Carmichael.

ESSAY ON VENEREAL DISEASES,
AND THE USES OF MERCURY IN THEIR TREATMENT.

With Drawings of the forms of Venereal Eruptions.

By RICHARD CARMICHAEL, M.R.I.A. 2d Edition, 8vo. 16s.

Dr. Carswell.

ILLUSTRATIONS OF THE ELEMENTARY FORMS OF DISEASE.

Imperial 4to. with col'd Figures of the natural size, and Descriptive Letter-press.

By R. CARSWELL, M.D. Prof. of Path. Anat. London University.

Fas. I. TUBERCLE; Fas. II. and Fas. III. CARCINOMA; Fas. IV. MELANOMA;
Fas. V. SOFTENING; Fas. VI. HÆMORRHAGE; Fas. VII. MORTIFICATION;
Fas. VIII. PUS; Fas. IX. HYPERTROPHY.
15s. each.

Mr. Chitty.

PRACTICAL TREATISE ON MEDICAL JURISPRUDENCE;

With all the Laws relating to Medical Practitioners, and Explanatory Notes.

By J. CHITTY, Esq. Barrister at Law.

Royal 8vo. Part I. 21s.

Dr. Christison.

TREATISE ON POISONS,

In relation to Medical Jurisprudence, Physiology, and the Practice of Physic.

By R. CHRISTISON, M.D. 8vo. 2d Edition, 18s. bds.

Sir C. M. Clarke.

OBSERVATIONS ON THE DISORDERS OF FEMALES.

Illustrated by Plates.

By SIR CHAS. MANSFIELD CLARKE, Bart. M.D. F.R.S. Physician in Ordinary
to the Queen.

3d Edition, 2 Vols. royal 8vo. 14. 16s.

Mr. Clark.

A PRACTICAL TREATISE ON TEETHING,

From Infancy to the completion of the Second Dentition, at about 24 years of Age;
In Four Chapters, each descriptive of an Era in the Development of the Teeth and Jaws.

By J. PATERSON CLARK, M.A.

Author of a "New System of Treating the Human Teeth."

8vo. 3s.

Mr. Le Gros Clark.

THE PRACTICAL ANATOMY AND ELEMENTARY PHYSIOLOGY
OF THE NERVOUS SYSTEM :

Designed for the Use of Students in the Dissecting-Room.

By F. LE GROS CLARK, Demonstrator of Anatomy in St. Thomas's Hospital.

9s. bds.

Dr. Collins.

A PRACTICAL TREATISE ON MIDWIFERY;

Containing the Result of SIXTEEN THOUSAND SIX HUNDRED AND FIFTY-FOUR BIRTHS,
occurring in the Dublin Lying-in Hospital, during a period of Seven
Years, commencing November, 1826.

By ROBERT COLLINS, M.D. late Master of the Institution.
8vo. price 12s. 6d.

Dr. Combe.

OBSERVATIONS ON MENTAL DERANGEMENT.

By A. COMBE, M.D. &c. Post 8vo. 7s. 6d.

Dr. Conquest.

OUTLINES OF MIDWIFERY,

Developing its Principles and Practice.

Intended as a Text-Book for Students. By J. T. CONQUEST, M.D. F.L.S.
5th Edit. with Engravings, 12mo. 7s. 6d.

Sir Astley Cooper.

ANATOMY AND SURGICAL TREATMENT OF ABDOMINAL HERNIA.

In Two Parts.

By SIR ASTLEY COOPER, Bart. F.R.S. &c.

Edited by C. ASTON KEY, Senior Surgeon to Guy's Hospital.

2d Edition, in folio, with Additional Plates, 5l. 5s.

By the same Author,

Observations on the Diseases of the Testis.

4to. with Plates, price 3l. 3s. coloured, or 1l. 11s. 6d. plain.

Anatomy of the Thymus Gland.

4to. with 5 Plates, 14s.

Dr. Copland.

A DICTIONARY OF PRACTICAL MEDICINE.

By J. COPLAND, M.D. F.R.S. &c.

8vo. Parts 1, 2, and 3, price 9s. each.

To form 1 thick volume, uniform with "Cooper's Surgical Dictionary."

By the same Author,

Pestilential Cholera: its Nature, Prevention, and Cure.

12mo. 5s.

Dr. Craigie.

ELEMENTS OF GENERAL AND PATHOLOGICAL ANATOMY.

By DAVID CRAIGIE, M.D. 8vo. 16s.

Dr. Cullen.

FIRST LINES OF THE PRACTICE OF PHYSIC.

By W. CULLEN, M.D. With an Appendix, commenced by the Author, and
continued and completed by J. C. GREGORY, M.D. F.R.S.E. &c.

2 Vols. 8vo. 1l. 4s.

Dr. Cutler.

POPULAR SURGERY :

Being plain Directions offered to the Public at large, for affording Relief in all Cases of Accident, and dangerous Disease, Poisoning, &c.,

In the absence of a regular Practitioner.

From the French of M. MAYOR. With Notes and Additions, by
THOMAS CUTLER, M.D. 4s. bds.

Dr. Thomas Davies.

LECTURES ON DISEASES OF THE LUNGS AND HEART,

Delivered at the London Hospital, by THOMAS DAVIES, M.D.

Mem. Roy. Col. Phys., &c. &c.

8vo. 12s.

Dr. Dowson.

INTRODUCTION TO THE STUDY AND PRACTICE OF MEDICINE.

By JOHN DOWSON, M.D. 12mo. 4s. 6d.

Mr. H. Earle.

**TWO LECTURES ON THE
PRIMARY AND SECONDARY TREATMENT OF BURNS.**

By HENRY EARLE, F.R.S. Surgeon Extraord. to the King, &c.
With 2 Plates, 2s. 6d.

Mr. J. W. Earle.

NEW EXPOSITION OF THE FUNCTIONS OF THE NERVES.

By JAMES WILLIAM EARLE.
8vo. Part I. 7s. 6d.

Edinburgh Medico-Chirurgical Society.

**TRANSACTIONS OF THE MEDICO CHIRURGICAL SOCIETY OF
EDINBURGH.**

8vo. Vol. I. with Plates, 18s. bds.; Vol. II. 14s. bds.; Vol. III. Parts I. and II. 12s. each.

Dr. Elliotson.

HUMAN PHYSIOLOGY.

By JOHN ELLIOTSON, M.D. Cantab. F.R.S.

With which is incorporated, much of the Elementary Part of the INSTITUTIONES
PHYSIOLOGICÆ of J. F. BLUMENBACH,

Professor in the University of Göttingen.

5th Edit. in 8vo. with numerous Woodcuts. Part I. price 10s. 6d.

* * The remaining Part will appear in the course of 1836.

Dr. Farre.

**TREATISE ON SOME
PRACTICAL POINTS RELATING TO THE DISEASES OF THE EYE.**

By the late J. C. SAUNDERS. Edited by J. R. FARRE, M.D.

3d Edition, 14s. plain, and 1l. 5s. coloured.

Journal of Morbid Anatomy, &c.

Edited by J. R. FARRE, M.D.

8vo. with Plates, Part I. 6s.

By the same Author,

Apology for British Anatomy.

4to. with 2 beautiful Lithographic Illustrations, 9s. plain; 12s. coloured.

Mr. D. Fox.

THE SIGNS, DISORDERS, AND MANAGEMENT OF PREGNANCY ;

The Treatment to be adopted during and after Confinement; and the
Management and Disorders of Children.

Written expressly for the Use of Females.

By DOUGLAS FOX, M.R.C.S. and one of the Surgeons to the Derbyshire General Infirmary.
8vo. 6s.

Dr. Fyfe.

ELEMENTS OF CHEMISTRY.

By ANDREW FYFE, M.D. F.R.S.E.

2d Edition, comprising all the recent Discoveries. 8vo. 24s.

By the same Author,

Manual of Chemistry. 18mo. 7s.

Mr. R. T. Gore.

INTRODUCTION TO THE COMPARATIVE ANATOMY OF ANIMALS ;

Compiled with constant reference to Physiology.

By C. G. CARUS, Med. et Phil. Doct. From the German, by R. T. GORE, M.R.C.S.
2 Vols. 8vo. and a 4to. Vol. of 20 Plates, with References, 3*l*.

Dr. Gooch.

PRACTICAL COMPENDIUM OF MIDWIFERY.

By the late ROBERT GOOCH, M.D.

Prepared for publication by GEORGE SKINNER, M.R.C.S.L.
12mo. 7s.

Dr. Mason Good.

THE STUDY OF MEDICINE.

By JOHN MASON GOOD, M.D. F.R.S. &c.

4th Edit. improved from the Author's MSS. and by reference to the latest advances in
Physiology, Pathology, and Practice.

By SAMUEL COOPER, Professor of Surgery in the London University.
4 thick Vols. 8vo. 3*l*. 3s.

Sir. S. L. Hammick.

PRACTICAL REMARKS ON
AMPUTATIONS, FRACTURES, AND STRICTURES OF THE URETHRA.

By SIR S. LOVE HAMMICK, Surgeon Extraordinary to the King, &c.
8vo. 9s.

Dr. Harrison.

THE DUBLIN DISSECTOR ; OR, MANUAL OF ANATOMY.

Comprising a Description of the Bones, Muscles, Vessels, Nerves, and Viscera ; also the
relative Anatomy of the different Regions of the Human Body ;
together with the Elements of Pathology.

By ROBERT HARRISON, A.M. M.B. T.C.D. &c. &c.

12mo. price 9s. in boards, the Fourth Edition, considerably enlarged and improved.

Dr. G. C. Holland.

AN INQUIRY INTO THE PRINCIPLES AND PRACTICE OF MEDICINE,
FOUNDED ON
ORIGINAL PHYSIOLOGICAL INVESTIGATIONS.

By G. CALVERT HOLLAND, M.D.
Physician to the Sheffield General Infirmary.
8vo. 12s.

Dr. Hooper.

MEDICAL DICTIONARY;

Containing an Explanation of the Terms in Anatomy, Botany, Chemistry, Materia Medica
Midwifery, Mineralogy, Pharmacy, Physiology, Practice of Physic, Surgery, &c.

By ROBERT HOOPER, M.D. F.L.S.

Physician to St. Marylebone Infirmary, &c.

1 large vol. 6th Edition, 28s.

By the same Author,

Morbid Anatomy of the Human Brain.

Illustrated by col'd Engravings; imp. 4to. 2l. 12s. 6d.

Morbid Anatomy of the Human Uterus and its Appendages.

Imp. 4to. with 21 col'd Plates, 3l. 3s.

Mr. Ingleby.

PRACTICAL TREATISE ON UTERINE HÆMORRHAGE,

In connexion with Parturition.

By JOHN INGLEBY, M.R.C.S. Lecturer on Midwifery, Birmingham.

8vo. 12s.

Dr. Jewel.

LONDON PRACTICE OF MIDWIFERY:

Including the most important Diseases of Women and Children.

Chiefly designed for the Use of Students and early Practitioners.

With Alterations and Additions, by G. JEWEL, M.D.

Senior Physician-Accoucheur to the St. George's and St. James's Dispensary; Lecturer on
Midwifery, &c.

12mo. 6th edit. 6s. 6d.

Dr. Kay.

THE PHYSIOLOGY, PATHOLOGY, AND TREATMENT OF ASPHYXIA;

Including Suspended Animation in New-born Children, and from Hanging,
Drowning, Wounds of the Chest, &c.

By J. P. KAY, M.D. formerly President of the Royal Med. Soc. Edinb.

8vo. 10s. 6d.

Mr. C. Aston Key.

MEMOIR ON THE
ADVANTAGES AND PRACTICABILITY OF DIVIDING THE STRICTURE
IN STRANGULATED HERNIA ON THE OUTSIDE OF THE SAC.

By C. ASTON KEY, Surgeon to Guy's Hospital, Lecturer on Surgery, &c.

With Cases, and 3 col'd Drawings. 7s. 6d.

Mr. Knox.

THE ANATOMIST'S INSTRUCTOR AND MUSEUM COMPANION:

Being Practical Directions for the Formation and subsequent Management
of Anatomical Museums.

By FREDERICK JOHN KNOX. 4s. 6d.

Dr. Latham.

LECTURES ON SUBJECTS CONNECTED WITH CLINICAL MEDICINE,

Delivered at St. Bartholomew's Hospital, by Dr. LATHAM.

1 vol. 12mo. 6s. 6d.

Mr. Lee.

AN ACCOUNT OF THE
MOST FREQUENTED WATERING PLACES ON THE CONTINENT,
And of the Medicinal Application of their Mineral Springs:
With Tables of Analysis, and an Appendix, on the principal Mineral Waters of England.

By EDWIN LEE, M.R.C.S.,

Author of Observations on the Continental Medical Institutions.

1 vol. post 8vo.

Mr. Liston.

ELEMENTS OF SURGERY.

By ROBERT LISTON, Surgeon to the North London Hospital.

One thick volume 8vo. 27s. The Three Parts may still be had separately, price 9s. each.

London Medical and Chirurgical Society.

TRANSACTIONS OF THE MEDICAL AND CHIRURGICAL SOCIETY OF
LONDON;

Comprising a mass of valuable and important Papers on Medicine and Surgery.

8vo. with Engravings. Vol. XIX. 15s.

Mr. Lucas.

A CONCISE

ANATOMICAL DESCRIPTION OF THE ARTERIES OF THE HUMAN BODY:

Together with full Directions for cutting down upon and securing the
several Arterial Trunks.

For the Use of Students in Anatomy.

By P. BENNETT LUCAS, Member of the Royal College of Surgeons, London.

4s. 6d. bds.

Dr. Macaulay.

DICTIONARY OF MEDICINE,

Designed for Popular Use.

By ALEXANDER MACAULAY, M.D. 8vo. 2d Edition, 14s.

Dr. Macbraire.

A SYSTEM OF MEDICAL NOSOLOGY.

By J. MACBRAIRE, M.D. L.E., &c. &c. 5s.

Dr. Mackenzie.

A PRACTICAL TREATISE ON DISEASES OF THE EYE.

By WM. MACKENZIE, M.D.

Lecturer on the Eye in the University of Glasgow.

2d Edit. in a thick volume 8vo. with Copper-plates and above 100 Woodcuts, 25s.

By the same Author,

Description of the Human Muscles; with a Table of their combined
Actions, &c. 3s.

Dr. Mackintosh.

ELEMENTS OF PATHOLOGY AND PRACTICE OF PHYSIC.

By J. MACKINTOSH, M.D. Acting Surgeon to the Ordnance in North Britain, &c.

2 vols. 8vo. 3d Edit. 31s. 6d.

Mr. Maclean.

ILLUSTRATIONS OF TEETHING ;

Or, a Treatise on the Progress and Shedding of the Human Teeth, to the completion of the permanent set :

Shewing the Causes of the Irregularity and Decay of the Teeth :
and the means most appropriate to their Preservation.

By R. MACLEAN, Dentist.

Royal 8vo. with Plates, 10s. plain, and 12s. col'd.

M. Magendie.

ELEMENTARY COMPENDIUM OF PHYSIOLOGY,

For the Use of Students.

By F. MAGENDIE, M.D. &c.

From the French, with Notes, Tables, &c. by E. MILLIGAN, M.D. &c.

4th Edit. greatly enlarged, 8vo. 21s.

Dr. Maunsell.

THE DUBLIN PRACTICE OF MIDWIFERY.

By HENRY MAUNSELL, M.D. Lecturer on Midwifery, &c. at the Medical School,
Park Street, Dublin. 12mo. 5s.

Dr. M'Cormac.

AN EXPOSITION OF THE
NATURE, TREATMENT, AND PREVENTION OF CONTINUED FEVER.

By HENRY M'CORMAC, M.D. M.R.C.S.E.

In 1 vol. 8vo. 6s.

Mr. Middlemore.

A TREATISE ON THE
DISEASES OF THE EYE AND ITS APPENDAGES.

By RICHARD MIDDLEMORE, M.R.C.S. Surgeon to the Birmingham Eye Infirmary, &c.

In 2 thick vols. 8vo. price 1*l.* 15s.

Dr. Morries & Mr. Graves.

HORTUS MEDICUS ;

Or Figures and Descriptions of the more important Plants, either used in Medicine, or
possessed of Poisonous Qualities, principally confined to those indigenous to
Great Britain : with the Chemical Analyses, &c.

By GEORGE GRAVES, F.L.S. ; and JOHN DAVIE MORRIES, M.D.

4to. with 44 Engravings, and numerous Wood-cuts. 30s. plain ; 42s. correctly coloured.

Dr. Murray.

A DISSERTATION ON THE INFLUENCE OF HEAT AND HUMIDITY ;

With practical Observations on the Inhalation of Iodine and various Vapours,
in Consumption, Catarrh, Croup, Asthma, and other Diseases.

By JAMES MURRAY, M.D., &c.

1 Vol. 10s. 6*d.* cloth.

Dr. Oke.

PRACTICAL OBSERVATIONS ON THE
IMMEDIATE TREATMENT OF THE PRINCIPAL EMERGENCIES

THAT OCCUR IN SURGERY AND MIDWIFERY,

Systematically arranged.

By W. S. OKE, M.D.

Part I. 8vo. 8s. Part II. 6s.

Dr. Palmer.

A DICTIONARY OF TERMS EMPLOYED BY THE FRENCH,
In Anatomy, Physiology, Pathology, Practical Medicine, Surgery, Midwifery,
Pharmacy, Medical Zoology, Botany, and Chemistry;
With their Derivations from the Greek and Latin; &c. &c. &c.
By SHIRLEY PALMER, M.D. formerly of Tamworth.
Parts I. and II. 6s. each. To be completed in 3 Parts.

Mr. Parkes.

THE CHEMICAL CATECHISM.
By the late SAMUEL PARKES, F.L.G. and W.S. &c.
Author of "Chemical Essays," "Rudiments of Chemistry," &c.
Adapted to the present state of Chemical Science, by E. W. BRAYLEY, Jun. A.L.S.,
of the London Institution.
8vo. 13th Edit. 15s.

Mr. Percivall.

ANATOMY OF THE HORSE,
Embracing the Structure of the Foot.
By WILLIAM PERCIVALL, M.R.C.S. Veterinary Surgeon in the 1st Life Guards.
8vo. 20s.
By the same Author,
Hippopathology: a Systematic Treatise on the Disorders and Lameness
of the Horse.
With Woodcuts. 8vo. Vol. I. 10s. 6d.

Pharmacopœia Londinensis.

PHARMACOPŒIA COLLEGII REGALIS MEDICORUM LONDINENSIS,
MDCCCXXIV.
8vo. 8s.; and 18mo. 5s.

Dr. Wilson Philip.

TREATISE ON THE MEANS OF PRESERVING HEALTH,
And particularly the prevention of Organic Diseases.
By A. P. WILSON PHILIP, M.D. F.R.S.L. and E. 8vo. 12s.

Mr. B. Phillips.

TREATISE ON THE URETHRA
Its DISEASES, especially STRICTURE, and their CURE.
By B. PHILLIPS. 8vo. with 2 Plates, 8s.

Mr. Rees.

A TREATISE ON THE ANALYSIS OF BLOOD AND URINE,
In Health and Disease:
With Directions for the Analysis of Urinary Calculi.
By G. O. REES.
Intended as an Introduction to Animal Analysis, for the use of the Medical Profession.
1 vol. 8vo. 5s. 6d.
By the same Author, translated from the French,
The Analysis of Inorganic Bodies. By J. J. BERZELIUS. 12mo. 5s.

M. Richerand.

ELEMENTS OF PHYSIOLOGY.

By A. RICHERAND.

Fifth Edition. Translated from the French, and supplied with Notes and a copious Appendix, by JAMES COPLAND, M.D.
8vo. 2d Edit. 18s.

Mr. Russell.

OBSERVATIONS ON THE TESTICLES.

By JAMES RUSSELL, M.R.C.S. 4s.

Mr. Sandwith.

INTRODUCTION TO ANATOMY AND PHYSIOLOGY:

For General Readers, Young Persons, and Junior Medical Students.

By T. SANDWITH, Surgeon. Plates. 7s. 6d.

Mr. Scott.

SURGICAL OBSERVATIONS ON THE
TREATMENT OF CHRONIC INFLAMMATION

In various Structures;

Particularly as exemplified in the Diseases of the Joints.

By JOHN SCOTT, Surgeon to the London Hospital, &c. 8vo. 7s. 6d.

By the same Author,

Cases of Tic Douloureux, and other Forms of Neuralgia.

8vo. 2s. 6d.

Sir Charles Scudamore.

TREATISE ON THE NATURE AND CURE OF RHEUMATISM.

By Sir C. SCUDAMORE, M.D. F.R.S. 15s.

By the same Author,

Cases illustrating and confirming the Remedial Power of the Inhalation of Iodine and Conium, in Tubercular Phthisis and various Disorders of the Lungs.

8vo. 2d edit. 7s. 6d.

Treatise on the Mineral Waters of Buxton, Tunbridge Wells, Bath, Cheltenham, the Beulah Spa (Norwood), &c.

2d Edition, 8s.

Observations on Laennec's Method of forming a Diagnosis of the Diseases of the Chest by means of the Stethoscope. 5s.

Treatise on the Nature and Cure of Gout and Gravel.

Large 8vo. 4th Edit. 20s. bds.

Further Examination of the Principles of the Treatment of Gout,

With Observations on Colchicum.

8vo. 2d edit. 6s.

Essay on the Blood.

8vo. 6s.

Principles of the Treatment of Gout;

With a further Examination of the Effects of Colchicum as a Remedy;

And some Observations on the Use of Veratria in that Disease.

8vo. 2s. 6d.

Dr. Seymour.

ILLUSTRATIONS OF
SOME OF THE PRINCIPAL DISEASES OF THE OVARIA,
Their TREATMENT, &c.

By EDWARD J. SEYMOUR, M. D.

8vo. with an Atlas of 14 Engravings in folio, 21s.; India Paper, 31s. 6d.

By the same Author,

Observations on the Medical Treatment of Insanity.

8vo. 5s.

Mr. Shaw.

NATURE AND TREATMENT OF THE VARIOUS DISTORTIONS
TO WHICH THE SPINE AND BONES OF THE CHEST ARE SUBJECT.

By JOHN SHAW. 8vo. 10s. 6d.

Dr. Southwood Smith.

SYSTEMATIC TREATISE ON FEVER.

By SOUTHWOOD SMITH, M.D. Physician to the London Fever Hospital.
8vo. 14s.

Mr. Spender.

OBSERVATIONS ON THE
CAUSES AND TREATMENT OF ULCEROUS DISEASES OF THE LEG.

By J. C. SPENDER, M.R.C.S.L.
1 vol. 8vo. 7s. 6d.

Mr. Stafford.

ON PERFORATION AND DIVISION OF
PERMANENT STRICTURE OF THE URETHRA
BY THE LANCETTED STILETTES;

With Observations on the Nature and Treatment of Spasmodic and Inflammatory Stricture,
and on various other Urethral Affections.

By R. A. STAFFORD,
Surgeon to the St. Marylebone Infirmary, Member of the Royal College of Surgeons, &c.
8vo. Third Edition, greatly enlarged and improved, 9s.

Mr. Stanley.

ACCOUNT OF THE
MODE OF PERFORMING THE LATERAL OPERATION OF LITHOTOMY.

By EDWARD STANLEY,
Lecturer on Anatomy and Physiology at St. Bartholomew's Hospital.
Royal 4to. Plates, 15s.

Mr. Swan.

ILLUSTRATIONS OF THE
COMPARATIVE ANATOMY OF THE NERVOUS SYSTEM.

By JOSEPH SWAN.
In 4to. Part I. price 10s. 6d.

By the same Author,

Demonstration of the Nerves of the Human Body.

Imp. folio, with 50 Engravings, half-bd. russia, 14l.

Demonstration of the Nerves of the Human Body.

Being the whole of the foregoing work, on 25 Steel Plates, by FINDEN.
In 4to. price 1l. 11s. 6d.

Treatise on Diseases and Injuries of the Nerves.

8vo. 14s.

Mr. W. H. Thomas.

THE SURGICAL AND DESCRIPTIVE
ANATOMY OF THE BONES, LIGAMENTS, AND JOINTS.

By W. H. THOMAS. 12mo. 6s.

Dr. Thomas.

MODERN PRACTICE OF PHYSIC,

Exhibiting the Character, Causes, Symptoms, Prognostics, Morbid Appearance,
and improved Method of treating the Diseases of all Climates.

By ROBERT THOMAS, M.D.
In 1 thick Vol. 8vo. 9th Edition, revised, 18s.

Dr. A. Todd Thomson.

ELEMENTS OF MATERIA MEDICA AND THERAPEUTICS ;

Including the recent Discoveries and Analyses of Medicines.

By A. TODD THOMSON, M.D. F.L.S. &c. Professor of Mat. Med. and Therapeutics,
and of Medical Jurisprudence, in the University of London.

1 Vol. 8vo. 21s.

By the same Author,

London Dispensatory ;

Containing Translations of the Pharmacopœias, &c. &c. : forming a Practical Synopsis
of Materia Medica, Pharmacy, and Therapeutics.

1 large Vol. 8vo. 8th Edition, with important Additions, 18s.

Atlas of Delineations of Cutaneous Eruptions ;

Illustrative of the Descriptions in Dr. BATEMAN'S Practical Synopsis of
Cutaneous Diseases.

Royal 8vo. 29 col'd Plates, 3*l.* 3s.

Conspectus of the Pharmacopœias.

New Edit. 5s. 6*d.*

Some Observations on the Preparation and Medicinal Employment of
Ioduret and Hydriodate of Iron.

8vo. 2s. 6*d.* sewed.

Dr. Thorburn.

**ELEMENTS OF BEDSIDE MEDICINE AND GENERAL PATHOLOGY,
&c. &c.**

By J. STEWART THORBURN, M.D. 8vo. 14s.

Mr. Travers.

**A FURTHER INQUIRY CONCERNING
CONSTITUTIONAL IRRITATION, AND THE PATHOLOGY OF THE
NERVOUS SYSTEM.**

By BENJ. TRAVERS, F.R.S. Senior Surgeon to St. Thomas's Hospital, &c. &c.
8vo. 14s.

By the same Author,

Observations on the Pathology of Venereal Affections.

8vo. 3s.

Inquiry into the Process of Nature in Repairing Injuries of the Intestines.

8vo. with Plates, 15s.

An Inquiry concerning that Disturbed State of the Vital Functions,
usually denominated Constitutional Irritation.

2d Edition, revised, 8vo. 14s.

Dr. Turnbull.

**ON THE MEDICINAL PROPERTIES OF THE NATURAL ORDER
RANUNCULACEÆ ;**

And more particularly on the Uses of *Sabadilla* Seeds, *Delphinium Staphysagria*, and
Aconitum Napellus : and their Alkaloids—*Veratria*, *Sabadilline*, *Delphinia*, and *Aconitine*.

By A. TURNBULL, M.D. 8vo. 6s.

By the same Author,

An Investigation into the Remarkable Medicinal Effects resulting from the
External Application of *Veratria*.

2d Edit. 8vo. 5s.

On the Preparation and Medicinal Employment of *Aconitine*,
in the Treatment of *Tic Douloureux*, &c.

8vo. 3s. 6*d.*

Sir George L. Tuthill.

NEW PHARMACOPŒIA OF THE ROYAL COLLEGE OF PHYSICIANS
OF LONDON. MDCCCXXIV.

Translated into English.

By SIR GEORGE L. TUTHILL, KNT. M.D. F.R.S.

8vo. 7s.; and 18mo. 4s.

The Translation is printed page for page with the original Latin, and both may
be had done up together.

Mr. Waite.

THE GUMS:

With late Discoveries on their Structure, Growth, Connexions, Diseases, and Sympathies.

By MR. GEORGE WAITE, Surgeon-Dentist, M.R.C.S.

1 vol. 12mo. 6s.

By the same Author,

The Surgeon-Dentist's Manual.

12mo. 5s. 6d. bds.

Mr. Wardrop.

ANEURISM, AND ITS CURE BY A NEW OPERATION.

By JAMES WARDROP, Surgeon to his Majesty.

Royal 8vo. 10s. 6d.

Mr. White.

COMPENDIUM OF THE VETERINARY ART;

Containing Rules for the Treatment of all the Disorders and Accidents to which the
Horse is liable; &c.

By J. WHITE, late Vet. Surgeon of the 1st, or Royal Dragoons.

15th Edit. 8s.

By the same Author,

Treatise on Veterinary Medicine;

Vol. II. containing the Materia Dietetica, the Materia Medica, and the Pharmacopœia.

12mo. 6th Edit. considerably improved, 6s.

Treatise on Veterinary Medicine;

Vol. III. containing Practical Observations on the Structure, Diseases, &c.
of the Digestive Organs of the Horse, &c.

8th Edit. 6s.

Compendium of Cattle Medicine,

Or Practical Observations on the Disorders of Cattle and the other Domestic Animals,
except the Horse.

12mo. 5th Edition, 6s.

THE LONDON MEDICAL GAZETTE:

CONSTITUTING A

COMPLETE AND AUTHENTIC RECORD OF MEDICAL LITERATURE.

Published every Saturday Morning: it is also issued (with the Magazines) in MONTHLY
PARTS, with a full and arranged Table of Contents.

