[Report 1905] / Medical Officer of Health, Leeds City.

Contributors

Leeds (England). City Council.

Publication/Creation

1905

Persistent URL

https://wellcomecollection.org/works/nqtmkmft

License and attribution

You have permission to make copies of this work under a Creative Commons, Attribution license.

This licence permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See the Legal Code for further information.

Image source should be attributed as specified in the full catalogue record. If no source is given the image should be attributed to Wellcome Collection.



LEEDS.

ANNUAL REPORT, 1905.

(PRELIMINARY).

Digitized by the Internet Archive in 2017 with funding from Wellcome Library

41, PARK SQUARE,

LEEDS, February 8th, 1906.

To the Chairman of the Sanitary Committee.

SIR.

The Committee have been already furnished with a general statement of the mortality in Leeds and the eight other largest towns in the United Kingdom during the year 1905. The death-rate in Leeds, calculated upon the figures returned to the Registrar General, and which exclude the deaths of non-municipal paupers, but include those of Leeds persons dying in our own fever hospitals and in the Hunslet workhouse outside the borough boundaries, was 15'3 per thousand per annum.

This is the lowest death-rate I have yet had to record. In 1894 the death-rate fell to 17.9, the lowest on record up to that period; in 1902 it was 17.6, in 1903, 16.6. Not only is the rate 15.3 the lowest for Leeds, but it is the lowest for any of the nine largest towns in the United Kingdom during 1905.

From table 1 it will be seen that during the five years 1885-1889 our Leeds death-rate was 21.16, taking into account all recorded deaths whether inside public institutions or not. In the following five years 1890-1894, containing the influenza period, it was again 21.16, notwithstanding the unusually healthy year which terminated that quinquennium. In the five years 1895-1899, the death-rate fell to 19.77, and in the following five years (1900-1904), to 18.36. The year 1905 had a death-rate, calculated upon the larger number of deaths, i.e., without excluding death of outsiders in public institutions of just under 15.65. The decrease of the general death-rate last year on that of the preceding year was 2.40; on that of the earliest of the quinquennia mentioned (1885-1889), 5.51.

From the same table it will be seen that the death-rate from the seven so called zymotic diseases was 0.97 below the rate in 1904, and 1.18 below that of the five years 1885-1889. This rate is also the lowest that I have had to record. The two years most nearly approaching to it were 1894, when the rate from the seven diseases was 1.98, and 1903 when the rate was 1.74.

The rate from small-pox was below 0.01 (0.006), the year before (1904) the deaths had not reached 0.005. In only four years during the sixteen since 1889 was there no death from small-pox.

From measles the rate of 0.24 last year was the lowest since I came to Leeds in 1889, with the exception of that of the year 1892, when it was 0.20.

From scarlatina last year our rate of 0.09 was the lowest but one in any of these sixteen years; the exceptional year was 1893, when the scarlatina rate was 0.08.

Diphtheria (exclusive of membranous croup) had a rate of 0.08 in 1905. It was 0.08 in 1892, 0.07 in 1890, 0.04 in 1891. If we add to the 0.08, our death-rate last year from diphtheria, 0.03 deaths per thousand from some form of croup, the death-rate from the combined group, including not only diphtheria and membranous croup but other croups not characterized as spasmodic, was o'11. The year 1892, with a rate of 0.08 from diphtheria alone, had a rate of 0.13 from croup, making 0'21 from the combined group. The year 1891, which had the lowest rate from diphtheria alone (0.04), had a rate of 0.08 from "croup," making the combined rate one of 0.12. The year 1890, with a rate of 0.07 from diphtheria alone, had a rate of 0.09 from croup, making 0.16 from the combined group. So that our rate from this group of diseases was lower in 1905 than in any other of the sixteen years, the figures for which are given in table D. part I.

1905

ANNUAL REPORT, LEEDS.

TABLE 1.

Annual deaths per 1,000 of the estimated population.

	All causes.	Seven zymotics.†	Consump-	Bronchitis, pneumonia, pleurisy.	Other lung diseases, without influenza.
Five years, 1885-89 (261 weeks)	21.16	2.78	1.70	3.93	0.52
*Five years, 1890-94 (261 weeks)	21.16	2.25	1.61	4.43	0.31
*Five years, 1895-99 (261 weeks)	19.77	2.74	1.47	3.24	0.55
*Five years, 1900-04 (261 weeks)	18.36	2.48	1.36	3.18	0.10
*Year 1904 (52 weeks)	18.05	2.57	1.40	2.96	0.18
*Year 1905 (52 weeks)	15.65	1.60	1.23	2.76	0.16
1905 increase on 1904					
" decrease " 1904	2.40	0.97	0.12	0.50	0.03
1905 increase on '85-9					
" decrease on '85-9	5.21	1.18	0.47	1.12	0.11

^{*} Estimated of course upon the population calculated from the result of the recent census.

⁺ Exclusive of membranous croup which the Registrar-General did not include in the seven zymotics until 1894.

TABLE 2.

Shewing the death-rates in the nine largest towns of the United Kingdom for the 52 weeks, and each of the thirteen week periods of 1905.

-							
		-	First quarter of 1905.	Second quarter of 1905.	Third quarter of 1905.	Fourth quarter of 1905.	52 Weeks.
	Leeds	-	15.7	14.6	15.3	15.4	15.3
	London	-	16.8	14.3	14.6	16.8	15.6
	Edinburgh	-	17.8	16.1	14.2	16:2	16:2
	Birmingham	-	16.9	15.7	15.6	16.4	16.5
	Sheffield	-	17.8	17.5	18.3	14'4	17.0
	Glasgow	-	19.4	18.2	15.5	18.6	17.9
-	Manchester	-	18.4	16.9	19.4	17:3	18.0
	Liverpool	-	20.4	17.7	19.9	20'4	19.6
-	Dublin	-	24.6	20.4	19.6	20'4	21.3

The rates are calculated upon the populations estimated by the Registrar-General for England from the results of the recent census.

TABLE 3.

Births and deaths registered in the City of Leeds in the four periods of 13 weeks ended respectively April 1st, July 1st, September 30th, and December 30th, 1905. Deaths in age groups.

		Мо	RTALITY	FROM AL	L CAUSES	AT SUBJ	OINED AC	SES.
1	- 2	3	4	5	6	7	8	9
1905. Estimated	Regis- tered Births.	At all Ages.	Under I Year.	and under 5-	and under 15.	and under 25.	and under 60.	60 and upwards.
population at these ages		456,787	11,964	41,144	93,645	91,934	192,143	25,957
I. Quarter	3,255	1,819	397	175	75	94	509	569
II. Quarter	3,214	1,720	373	182	74	106	494	491
III. Quarter	3,081	1,790	668	202	58	70	404	388
IV. Quarter	2,787	1,795	437	260	62	55	488	493
52 weeks	12,337	7,124	1,875	819	269	325	1,895	1,941

TABLE 4.

Birth and death rates in the City of Leeds in the four periods of 13 weeks ended respectively April 1st, July 1st, September 30th, and December 30th, 1905. Death rates in age groups.

			DE.	ATHS PI	er Ann	UM PER	1,000	Living	+	
1905.	2 Birth- rate.	At all Ages.	4 Under I year.	5 and under 5.	5 and under 15.	7 15 and under 25.	8 25 and under 60.	9 60 and upwds.	10 25 10 65	11 Over 65
I. Quarter					3.5	4.1	10.6			111.8
II. Quarter III. Quarter IV. Quarter	27.1	15.7	224	19.7	3·2 2·5 2·7		10 ³ 8 ⁴	60.0		. 78.7
52 weeks			-			3.2	-			

Mortality in Children under one year of age, during the 52 weeks of 1905.

	First Quarter.	Second Quarter.	Third Quarter.	Fourth Quarter.	YEAR.
Calculated per 1,000 of the population under 1, estimated to the middle of 1904, on the supposition that the whole population of the city was that estimated by the Registrar General, and that the number of children under one bore the same proportion to the population as at the 1901 census	133	125	224	147	157
Deaths under I per I,000 births registered in same period	122	116	217	157	152
Deaths per 1,000 registered births, the latter instead of those for the same quarterly period being the average of the same and four preceding quarters. The rate for the year is the mean of the four					
quarterly rates	126	118	213	142	150
Average rate by last method for five preceding years	155	149	234	164	176
Average rate by second method for years 1886-90 (from table 7a 1890 report)	158	147	223	171	175

TABLE D, Part 1.—Shewing death-rates from certain causes for the years 1890 to 1905.

			SEVEN	COMMON		ZYMOTICS.						
	Small-pox.	Measles.	Scarlatina.	Diph- theria.	Whooping cough.	"Fever."	Diarrhœa.	All seven.	Croup (mem- branous and undefined)	Phthisis.	Influenza and diseases of the air-passages other than consumption.	
1890 (53 wks.)	00.0	0.27	0.28	,20.0	0.20	0.30	86.0	2.39	60.0	99.1	29.5	
1891 (52 wks.)	00.00	0.71	0.18	0.04	0.41	0.50	98.0	2.41	80.0	64.1	11.9	
1892 (52 wks.)	0.02	0.50	0.50	0.08	0.45	21.0	01.1	2.18	0.13	1.42	4.56	
1893 (52 wks.)	80.0	06.0	80.0	0.15	0.44	0.30	09.1	3.55	81.0	1.70	4.60	
1894 (52 wks.)	10.0	0.75	0.13	0.15	0.34	0.14	0.45	86.1	0.17	1.49	3.64	
1895 (52 wks.)	:	0.35	0.13	0.10	0.56	0.22	1.58	2.65	0.12	1.55	4.34	
1896 (53 wks.)	00.00	0.48	0.18	0.10	09.0	0.31	69.0	2.27	60.0	05.1	4.02	
1897 (52 wks.)	:	0.40	0.23	0.13	0.24	0.21	1.58	2.79	0.10	1.44	3.60	
1898 (52 wks.)	0.00	0.45	0.50	0.49	0.39	0.23	1.24	3.10	0.12	1.39	3.41	
1899 (52 wks.)	:	0.37	0.15	0.71	0.38	0.17	96.0	2.73	0.12	1.41	3.72	_
1900 (52 wks.)	0.00	0.58	0.12	0.22	0.36	0.50	60.1	2.93	90.0	141	4.19	
1901 (52 wks.)	:	0.58	61.0	0.38	0.33	61.0	1.47	3.14	20.0	141	3.46	
1902 (53 wks.)	10.0	0.43	0.13	0.18	0.46	0.18	19.0	2.00	0.02	1.31	3.52	
1903 (52 wks.)	0.02	0.28	0.25	0.14	0.27	0.13	0.62	1.74	0.03	1.27	3.00	
1904 (52 wks.)	00.0	0.77	0.13	60.0	0.46	0.11	10.1	2.57	0.02	1.40	3.21	
1905 (52 wks.)	0.01	0.24	60.0	80.0	0.27	0.12	0.80	09.1	0.03	1.23	3.07	
				See not	See note to table D. part 2.	D. part 2.						

TABLE D, Part 2. For whole District.

	D 1 .:	Birth	is.	Deaths one year			s at all Total.		Deaths	Deaths	Death ages	s at all net.
Year.	Population estimated to middle of each year.	Number.		Number	regis- tered.	Number		in Public Institu- tions.	of non- resid'nts regis- tered in district.	tered beyond district.	Number	
1	2	3	4	5	6	7	8	9	10	11	12	13
1890	362,768	12,336	33.4	2,128	173	8,370	22.7	. 819	132	No return	8,238	22.4
1891	369,099	12,538	34'I	2,216	177	8,429	22'9	869	144	do.	8,285	22.2
1892	375,540	12,546	33.2	2,114	168	7,403	19.8	789	129	do.	7,274	19'4
1893	382,094	12,348	32'4	2,542	206	8,512	22'4	874	147	24	8,365	22'0
1894	388,761	12,502	32.3	1,945	156	6,935	17.9	772	142	7	6,793	17.5
1895	395,546	12,478	31.7	2,384	191	8,101	20.6	882	167	No return	7,934	20'1
1896	402,449	12,573	30.8	2,120	169	7,682	18.8	908	161	8	7,521	18:4
1897	409,472	12,912	31.6	2,454	190	8,148	20.0	881	175	I	7,973	19.2
1898	416,618	12,971	31.2	2,372	183	7,996	19'3	940	142	9	7,854	18.9
1899	423,889	12,939	31'2	2,222	172	8,105	19.2	1,005	181	26	7,924	18.8
1900	431,287	13,091	30.2	2,397	183	8,619	20'I	1,084	184	90	8,525	19.8
1901	430,630	12,898	30.1	2,429	188	8,283	19.3	1,176	171	92	8,204	19.1
1902	437,341	13,245	29.8	2,113	160	7,814	17.6	1,154	211	96	7,699	17.3
1903	443,559	12,996	29.4	1,992	153	7,334	16.6	1,094	184	113	7,263	16.4
1904	450,142	12,561	28.0	2,207	176	8,096	18.0	1,185	173	116	8,039	17.9
Averages for years 1895- 1904		12,866	30.4	2,269	176	8,018	19.0	1,031	175		7,894	18.6
1905	456,787	12,337	27.1	1,875	152	7,124	15.6	1,225	178	102	7,047	15.5

* Rates calculated per 1,000 of estimated population. The populations given in column 2 for the 11 years, 1890 to 1900, are those estimated by geometrical progression at the rate of increase which obtained between censuses of 1881-1891. That for 1901 has been estimated similarly from the rate which prevailed between the censuses of 1891-1901. To make the populations for 1900 and the previous years correspond they should be decreased respectively by the following percentages: (for 1900) 1.72, (for 1899) 1.53, (for 1898) 1.34, diminishing by about 0.19 each year to (for 1892) 0.21, and (for 1891) 0.05. The rates in columns 4, 8, and 13 must be correspondingly increased by these percentages (see Annual, 1901, and table on p. 58).

The deaths included in column 7 of this table are the whole of those registered during the year as having actually occurred within the City or in the fever hospital at Seacroft. The deaths included in column 12 are the numbers in column 7, corrected by the subtraction of the numbers in column 10 and the addition of the numbers in column 11. Column 11 is however, incomplete in some of the years. Deaths in fever hospitals belonging to but outside the City are, unless those of strangers, included in column 7.

By the term "Non-residents" is meant persons brought into the district on account of illness, and dying in some public institution, and paupers from townships outside the City dying at the Workho se; and by the term "Residents" is meant persons who have been taken out of the district on account of illness, and have died in some public institution, other than our pwn fever hospitals, and inmates of the County Asylums from the City townships.

In the 52 weeks ended 30th December, 1905, the following births and deaths were registered, and cases of infectious diseases reported.

Population estimated by the Registrar General after Census	456,787
Acreage	21,572
Births registered 12,337 Birth-rate per thousand	27.1
Average birth-rate, ten years, 1895-1904 ‡	30.2
Deaths (all ages) 7,124 Death-rate per thousand	15.6
(R.G. 6946=15.26).*	
Average death-rate, ten years, 1895-1904* ‡	19.0
Deaths (under 1 year) 1,875 Ratio to 1,000 births	152

	Total deaths.	Death rate p	per thousand.	Cases
	Total deaths.	1905.	1895-1904.	notified. 1905.
			‡	
I. Smallpox	3	0.01	0,01	129
2. Measles	109	0.24	C'47	304
3. Scarlatina	39	0.00	0.18	935
Diphtheria	38	0.08	0'29	364
4. Memb. croup	6	0.01	0.02	8
Other croups	6	0.01	0.03	6
5. Whooping cough	121	0.27	0.38	
(Typhus	1	0.00	0.00	I T
6. Typhoid	52	0.11	0.18	. 292
Ctd. fever	2	0.00	0.00	
Puer. sept	9	0.03	0.03	28
Erysipelas	5	0.01	0'03	380
7. Diarrhœa	365	0.80	1.00	
Cholera (English)			0.00	
Phthisis	558	1.23	1'41	556§
Other tuberculous diseases	288	0.63	0.78	
Seven commoner zymotics) (including mem. croup) Bronchitis	736	1.62	2.64	
Pneumonia	1,258	2.76	3'34	
Other, lung diseases (without influenza)	72	0.19	0.50	
Malignant growths	444	0.08	0.79	
Violence and accident	273	0.60	0.62	

* The R.G. excludes deaths of non-municipal paupers. The rate is calculated on the larger number.

The average of the published returns were birth-rate 30.4, death-rate 19.0. The higher figures obtained from calculating the births and deaths on the arithmetical mean of populations for 1899 and 1900, obtained by method of interpolation explained in report for 1901, have been printed above. The death-rates for special groups are the uncorrected means of the published returns, from which non-residents were not excluded.

[|] Means not notifiable.

[§] Notification voluntary,

Whooping-cough caused a death-rate last year of 0.27 The rate was the same in 1903. It was 0.24 in 1897, the only occasion on which it was lower than last year, although it was only 0.29 in 1895.

From the combined group of "fever" deaths, constituted principally of deaths from typhoid, the rate last year was 0.12, against 0.11 in 1904, and 0.13 in 1903. Each of the previous thirteen years had a rate higher than the highest of these three, though the year 1894, the especially healthy year, had a fever death-rate of only 0.14. The other years varied in their fever death-rates from 0.17 to 0.30, the average of the ten years, 1895-1904 being 0.18.

From typhoid alone our death-rate was 0'11 last year against an average of 0'18 in the ten preceding years, and 0'21 in the five years immediately preceding them. The progress of typhoid, taking the three quinquennial periods 1890-1894, 1895-1899, and 1900-1904, is interesting.

The death-rate from typhoid fever was 0.21 in 1890-1894, in 1895-1899 it was 0.20, in 1900-1904 it was 0.16. The drop to 0.11 last year, the healthiest year on record, is not of course of equal importance to the gradual lowering that has taken place during the three periods of five years. The great drop from 0.20 to 0.16 has co-incided with the removal of a large number of our old privy middens.

From diarrhoea our death-rate last year was 0.80. This is not the lowest rate during the sixteen years, but it is perceptably below the average of the ten years 1895-1904, which was 1.09. The five years before that had a death-rate of 1.00. Carried to three places of decimals, the rates in the three quinquennia were in 1890-1894, 0.998*, in 1895-1899, 1.210*, and in 1900-1904, 0.960*.

^{*} These figures are averaged from table D, the figures in which were taken from the tables published at the time.

From consumption our death-rate of 1'23 was the lowest in any year of the sixteen. The rate was 1'27 in 1903, 1'31 in 1902, but in no other year lower than 1'39. As compared with the preceding year the rate from consumption was 0'17; as compared with the five years 1885-1889 it was 0'47, lower.

Bronchitis, pneumonia, and pleurisy also had a low deathrate in 1905. The rate of 2.76 was lower by 0.20 than that of the preceding year, and by 1.17 than that of the quinquennium 1885-1889. That quinquennium, however, had not the highest death-rate from this group, for the years 1890-1894, including the especially influenza years, had a rate of 4.43.

Excluding influenza, the "other lung diseases" group showed a decrease of 0.02 upon the rate of 1904, and of 0.11 on the quinquennium, 1885-1889, and of 0.15 upon the influenza quinquennium, 1890-1894.

It has been already mentioned that Leeds had the lowest death-rate of any of the nine largest towns. The rate, however, in London closely approximated that of Leeds; our rate was 15.3, the London rate, 15.6. Then came Edinburgh and Birmingham each 16.2, Sheffield and Glasgow 17.0 and 17.9, Manchester and Liverpool 18.0 and 19.6, and Dublin 21.3. These figures will be found in table 2.

During the first period of five years, for which these figures has been got out, 1890-1894, we stood fourth on the list. During the period 1895-1899 we stood second on the list. During the period 1900-1904 we fell to the third place. Taking the whole period of sixteen years our position was third. The position of the several towns being London 18.7, Edinburgh 18.9, Leeds 19.4, Birmingham 20.1, Sheffield 20.4, followed by Glasgow 21.5, Manchester 23.1, Liverpool 24.1, and Dublin 26.1. Leeds, I believe, was the only one of the provincial towns which did not enlarge its area during these sixteen years.

Infant mortality.—The mortality amongst children in 1905 was very much below the average of the previous fifteen years. During the years 1890-1904, as shown on the chart in the Annual Report for 1904, there was an average of 176 deaths in children under one for every thousand born. Last year the rate fell to 152; a rate lower than in any of the fifteen years. In 1894, the first of our specially healthy years, the rate had been 156. In 1903, the healthiest year until the one we are speaking of, the children's death-rate was 153.

Table 5 shows in the second line of figures the ratios of mortality in children under one year of age to each thousand births registered during each of the quarters of 1905. In the third line the rates are calculated in the same way upon the births not in the same quarter but upon the mean births in that quarter and the four preceding it. The same table gives also in the first line the death-rate of children under one upon the population at that age estimated from the census returns.

The diminution in infantile death-rate, it will be seen, was not entirely due to lessened mortality in the third quarter. The rate was less in the first quarter than the average either of the five preceding years or of the years 1886-1890. So also in the second quarter. It was appreciably but not so markedly less in the third quarter, and it was considerably less again in the fourth quarter.

It is fair to assume that the absence during the greater part of the year of measles and whooping-cough, in epidemic form, along with the usually corresponding diminution in the death-rate from bronchitis and broncho-pneumonia accounted in great part for the diminution of infantile mortality throughout the year, although one would fain hope that some of the work we have now done for several years is beginning to bear fruit. The subject, however, is so important that I propose to return to it in the supplementary portion of the present report.

JAMES SPOTTISWOODE CAMERON.

. . .

ANNUAL REPORT

(With SUPPLEMENT)

- MADE TO THE

URBAN SANITARY AUTHORITY

OF THE

CITY OF LEEDS,

FOR THE YEAR

1905,

AND PARTLY FOR 1906.

BY

J. SPOTTISWOODE CAMERON,

M.D., B.Sc., &c.,

Medical Officer of Health to the City.

CONTENTS.

	PAGE
GENERAL REPORT—Incidence of disease—Infant mortality	I
Supplementary on Infantile Mortality	9
The new table—calculation of infant mortality—mortality under one,	
1890-1904: quarterly variations—age groups under one—age	
groups in districts—mortality from different causes in districts—	
comparisons between South-East Leeds and Chapeltown.	
PART II.—Special diseases	32
Tuberculosis	32
DISEASES CHIEFLY AFFECTING LUNGS AND AIR PASSAGES	35
Phthisis	35
Other lung diseases	37
COMMONER INFECTIVE DISEASES	39
Small-pox-Notes to table 8-Vaccination and mortality, Jan., 1902,	37
to July, 1906	39
Measles	47
Scarlet fever—Relations of scarlatinal death rate and isolation	49
Diphtheria	51
0 1 1 6	
Enversage to two proposed a proposed	52
	52
OTHER DISEASES	57
ILLNESSES NOTIFIED	57
PART III.—HEALTH OF DISTRICTS.	
Wards	58
REGISTRATION SUB-DISTRICTS	59
INTERCEPTS	66
LOCAL FACTORS	66
PART IV.—Administrative work	67
Changes in Staff	67
W	74
T D D D D D D D D D D D D D D D D D D D	78
	10
WOMEN INSPECTORS (also p. 83)—Infectious diseases—outworkers—work- places—infantile mortality—visits to newly-born—midwives' Act—	
other work	88
ASHPIT CLEANSING—DESTRUCTORS—STREET CLEANSING	93
FOOD INSPECTION—Cowkeepers—cowsheds—inspection of cattle—examination	93
of milk for tubercle-bacteriological examination of milk-food and	
drugs-meat and markets	96
Smoke inspection	134
Description and District District Control of the Principle	135
TEMPORARY DWELLINGS INSPECTION—Students' lodgings—conditions found	133
—dangers from inside water closets—the calcium phosphide test	
	137
LIST OF TABLES.	
Births, deaths, and infectious diseases heard of	vi.
In the text.	
TABLE 1Mortality in Leeds in 1905, from all and certain groups of causes,	
and comparison with previous years	3
" 2.—Mortality in nine largest towns in United Kingdom in 1905	6
,, 3.—Deaths in Leeds, 1905, and quarters; age groups—births	7
,, 4.— Do. per 1,000 living—birth-rate	8
- Infantile mortality, 1905, deaths at certain ages from stated causes	10
5.—Death-rates, children under I year, 1905, calculated several ways.	
— Mortality of children under one, each quarter for the fifteen years,	
1890-1904, showing variations above and below average of	
whole period	17
- —Mean temperatures each quarter, 1890-1904	19
— Rainfall each quarter, 1890-1904	19
- Age groups under one-percentage of all deaths under one	20
- Age groups under one, excess at certain ages in districts per cent.	
of city rate at same ages	23
	3

								PAGI
[ABL	E X.—Meat inspection, 1905		***					13:
,,	XI.—Smoke inspection, 1905		***			***		13.
- , ,	XII.—Disinfection, 1905	***	***					130
,,	——Flushing, 1905		***			100	***	13
,,	XIII. and XIV.—Patients admitted	to infe	ectious	s hospit	als, 19	05		136
,,	Other work of removal st	aff				121		135
,,	XV.—Inspection of canal boats,	1905						13
	XVI -Inspection of houses let-in		ngs, I	905		100		13
,,,	XVIIOther work of temporary				ctor, I	905		130
	XVIII.—Results of examination						ents'	
	lodgings		***	111		14.5		46-14
	· · · · · · · · · · · · · · · · · · ·		**					
		ENDI		1 11				
"	A.—Part I.—Causes of death in as separate districts.	regist	ration	sub-dis	tricts;	institut	ions	
,,	A.—Part 2.—Populations, births tion sub-districts, public districts.							
,,,	A.—Parts 3, 4, 5, 6, similar to p part 7 for the first quarte			eral qua	irters o	f 1905,	and	
,,	B.—Part I.—Population, births, in districts, and admis	new ca	ses of					
	arranged for registration							
,,	B.—Part 2.—Similar, except as	to birth	is, to	part I,	but for	wards.		
,,	B.—Parts 3, 4, 5, 6, cases rep							
	during the several quar quarter of 1906.	ters of	1905	, and p	art 7,	during	first	
,,	C.—Deaths classified as to cause allocated. Notes (See				Institu	ation de	aths	
,,	D.—Death-rates in Leeds from the sixteen years, 1890-1	certai			causes	for eac	h of	
,,	D.—Part 2.—Population, births per 1,000 births, deaths	s, birtl						

nett, after allocation for the sixteen years, 1890-1905, averages for decade, 1895-1904.

E.—Part 1.—Births, birth-rate, deaths, death-rate for registration districts, and deaths and death-rate for municipal wards (institution deaths in both cases allocated) for the year 1905.

E.—Parts 2, 3, 4, 5, similar for each quarter of 1905, and 6 for first quarter of 1906.

F.—Births, deaths in Leeds from all and certain groups of causes, sickness, and meteorological data for each week of fourth quarter, 1904, of each quarter of 1905, and of the first quarter of 1906.

The material asked for in the new Local Government tables will be found as follows :-

I.—In table D, part 2.

II. - In table A for current and previous years.

III.—Age groups in new table 18, pp. 54-55; localities in table B.

IV.—Ages in table 17, p. 53 (altered from old table 17 to include ages), including, however, all deaths in institutions; localities in table C, where institution deaths allocated.

V.—Infantile mortality, pp. 10, 23, 24.

ERRATA, 1904 REPORT.

TABLE 25. - Col. 6 (p. 96), in line 22, for 89 read 90; line 24, for 12 read 11; col. 8, line 22, for 68 read 69; line 24, for 7 read 6. In cases "dead" (p. 97), line 22, for 282 read 284. Cases "alive," line 24, for 25 read 23. In total cases "alive," bottom line but one, for 2,036 read 2,034; bottom line, for 2,811 read 2,809. In total cases "dead," last line but one, for 1,460 read 1,462; last line, for 1,861 read 1,863.

TABLE VIII. b.—Two samples of milk from Birstwith, December 7th and 15th, result should have been entered as "incomplete."

In the 52 weeks ended 30th December, 1905, the following births and deaths were registered, and cases of infectious diseases reported.

Population estimated by the Registrar General after Census	456,787
Acreage	21,572
Births registered 12,337 Birth-rate per thousand	27'1
Average birth-rate, ten years, 1895-1904 ‡	30.2
Deaths (all ages) 7,124 Death-rate per thousand	15.6
(R.G. 6946=15.26).*	
Average death-rate, ten years, 1895-1904* ‡	10,0
Deaths (under 1 year) 1,875 Ratio to 1,000 births	152

	Total deaths	Death rate	per thousand.	Cases
	Total deaths	1905.	1895-1904.	notified. 1905.
			†	
1. Smallpox	3	0.01	0.01	129
2. Measles	100	0.24	c'47	304
3. Scarlatina	39	0.00	0.18	935
, Diphtheria	38	0.08	0'29	364
4. Memb. croup	6	0,01	0.02	8
Other croups	6	0,01	0.03	6
5. Whooping cough	121	0.52	0.38	
Typhus	I	0.00	0.00	1
6. Typhoid	52	0.11	0.18	292
Ctd. fever	2	0.00	0.00	
Puer. sept	9	0'02	0.03	28
Erysipelas	5	0,01	0.03	380
7. Diarrhoea	365	0.80	1,00	
Cholera (English)			0.00	
Phthisis	558	1'23	1'41	555\$
Other tuberculous disea	ases 288	0.63	0.78	
Seven commoner zymo (including mem. crou		1.62	2.64	***
Bronchitis Pneumonia Pleurisy	1,258	2.76	3'34	
Other lung diseases (without influenza)	} 72	0.19	0.50	
Malignant growths	444	0.08	0.79	
Violence and accident	273	0.60	0.62	

^{*} The R.G. excludes deaths of non-municipal paupers. The rate is calculated on the larger number.

The average of the published returns were birth-rate 30'4, death-rate 19'0. The higher figures obtained from calculating the births and deaths on the arithmetical mean of populations for 1899 and 1900, obtained by method of interpolation explained in report for 1901, have been printed above. The death-rates for special groups are the uncorrected means of the published returns, from which non-residents were not excluded.

Means not notifiable.

[§] Notification voluntary.

ANNUAL REPORT, 1905.

To the Chairman of the Sanitary Committee.

SIR,

The Committee have been already furnished with a general statement of the mortality in Leeds and the eight other largest towns in the United Kingdom during the year 1905. The death-rate in Leeds, calculated upon the figures returned to the Registrar General, and which exclude the deaths of non-municipal paupers, but include those of Leeds persons dying in our own fever hospitals and in the Hunslet workhouse outside the borough boundaries, was 15'3 per thousand per annum.

This is the lowest death-rate I have yet had to record. In 1894 the death-rate fell to 17.9, the lowest on record up to that period; in 1902 it was 17.6, in 1903, 16.6. Not only is the rate, 15.3, the lowest for Leeds, but it is the lowest for any of the nine largest towns in the United Kingdom during 1905. (Table 2.)

From table 1 it will be seen that during the five years 1885-1889 our Leeds death-rate was 21°16, taking into account all recorded deaths whether inside public institutions or not. In the following five years 1890-1894, containing the influenza period, it was again 21°16, notwithstanding the unusually healthy year which terminated that quinquennium. In the five years 1895-1899, the death-rate fell to 19°77, and in the following five years (1900-1904), to 18°36. The year 1905 had a death-rate, calculated upon the larger number of deaths, *i.e.*, without excluding deaths of outsiders in public institutions, of just under 15°65. The decrease of the general death-rate last year on that of the preceding year was 2°40; on that of the earliest of the quinquennia mentioned (1885-1889), 5°51.

From the same table it will be seen that the death-rate from the seven so called zymotic diseases was 0.97 below the rate in 1904, and 1.18 below that of the five years 1885-1889. This rate is also the lowest that I have had to record. The two years most nearly approaching to it were 1894, when the rate from the seven diseases was 1.98, and 1903 when the rate was 1.74.

The rate from small-pox was below 0.01 (0.006). The year before (1904) the deaths had not reached 0.005. In only four years during the sixteen since 1889 was there no death from small-pox. (Table D, Part 1.)

From measles the rate of 0.24 last year was the lowest since I came to Leeds in 1889, with the exception of that of the year 1892, when it was 0.20. (D 1.)

From scarlatina last year our rate of 0.09 was the lowest but one in any of these sixteen years; the exceptional year was 1893, when the scarlatina rate was 0.08. (D 1).

Diphtheria (exclusive of membranous croup) had a rate of 0.08 in 1905. It was 0.08 in 1892, 0.07 in 1890, 0.04 in 1891. If we add to the 0.08 (our death-rate last year from diphtheria) 0.03 deaths per thousand from some form of croup, the death-rate from the combined group, including not only diphtheria and membranous croup but other croups not characterized as spasmodic, was o'11. The year 1892, with a rate of 0.08 from diphtheria alone, had a rate of 0.13 from croup, making 0'21 from the combined group. The year 1891, which had the lowest rate from diphtheria alone (0.04), had a rate of 0.08 from "croup," making the combined rate one of 0'12. The year 1890, with a rate of 0'07 from diphtheria alone, had a rate of 0.00 from croup, making 0.16 from the . combined group. So that our rate from this group of diseases was lower in 1905 than in any other of the sixteen years, the figures for which are given in table D. part 1.

TABLE 1.

Annual deaths per 1,000 of the estimated population.

	All causes.	Seven zymotics.†	Consumption.	Bronchitis, pneumonia, pleurisy.	Other lung diseases, without influenza.
Five years, 1885-89 (261 weeks)	21.16	2.78	1.70	3.93	0.22
*Five years, 1890-94 (261 weeks)	21.16	2.22	1.61	4.43	0.31
*Five years, 1895-99 (261 weeks)	19.77	2.74	1.47	3.24	0.55
*Five years, 1900-04 (261 weeks)	18.36	2.48	1.36	3.18	0.10
*Year 1904 (52 weeks)	18.05	2.57	1.40	2'96	0.18
*Year 1905 (52 weeks)	15.65	1.60	1.53	2.76	0.16
1905 increase on 1904					
" decrease " 1904	2.40	0.97	0.12	0.50	0.03
1905 increase on '85-9					
" decrease on '85-9	5.21	1.18	0.47	1.12	0.11

^{*} Estimated of course upon the population calculated from the result of the recent census.

Whooping-cough caused a death-rate last year of 0.27. The rate was the same in 1903. It was 0.24 in 1897, the only occasion on which it was lower than last year, although it was only 0.29 in 1895.

From the combined group of "fever" deaths, constituted principally of deaths from typhoid, the rate last year was 0.12,

[†] Exclusive of membranous croup which the Registrar-General did not include in the seven zymotics until 1894.

against 0.11 in 1904, and 0.13 in 1903. Each of the previous thirteen years had a rate higher than the highest of these three, though the year 1894, the especially healthy year, had a fever death-rate of only 0.14. The other years varied in their fever death-rates from 0.17 to 0.30, the average of the ten years, 1895-1904 being 0.18.

From typhoid alone our death-rate was 0.11 last year against an average of 0.18 in the ten preceding years, and 0.21 in the five years immediately preceding them. The progress of typhoid, taking the three quinquennial periods 1890-1894, 1895-1899, and 1900-1904, is interesting.

The death-rate from typhoid fever was 0.21 in 1890-1894, in 1895-1899 it was 0.20, in 1900-1904 it was 0.16. The drop to 0.11 last year, the healthiest year on record, is not of course of equal importance to the gradual lowering that has taken place during the three periods of five years. The great drop from 0.20 to 0.16 has co-incided with the removal of a large number of our old privy middens.

From diarrhœa our death-rate last year was 0.80. This is not the lowest rate during the sixteen years, but it is perceptibly below the average of the ten years 1895-1904, which was 1.09. The five years before that had a death-rate of 1.00. Carried to three places of decimals, the rates in the three quinquennia were in 1890-1894, 0.998*, in 1895-1899, 1.210*, and in 1900-1904, 0.960*.

From consumption our death-rate of 1'23 was the lowest in any year of the sixteen. The rate was 1'27 in 1903, 1'31 in 1902, but in no other year lower than 1'39. As compared with the preceding year the rate from consumption was 0'17; as compared with the five years 1885-1889 it was 0'47, lower.

Bronchitis, pneumonia, and pleurisy also had a low deathrate in 1905. The rate of 2.76 was lower by 0.20 than that

^{*} These figures are averaged from table D, the figures in which were taken from the tables published at the time.

General. 5

of the preceding year, and by 1.17 than that of the quinquennium 1885-1889. That quinquennium, however, had not the highest death-rate from this group, for the years 1890-1894, including the especially influenza years, had a rate of 4.43.

Excluding influenza, the "other lung diseases" group showed a decrease of 0.02 upon the rate of 1904, and of 0.11 on the quinquennium, 1885-1889, and of 0.15 upon the influenza quinquennium, 1890-1894.

It has been already mentioned that Leeds had the lowest death-rate of any of the nine largest towns. The rate, however, in London closely approximated that of Leeds; our rate was 15'3, the London rate, 15'6. Then came Edinburgh and Birmingham each 16'2, Sheffield and Glasgow 17'0 and 17'9, Manchester and Liverpool 18'0 and 19'6, and Dublin 21'3. These figures will be found in table 2.

During the first period of five years, for which these figures has been got out, 1890-1894, we stood fourth on the list. During the period 1895-1899 we stood second on the list. During the period 1900-1904 we fell to the third place. Taking the whole period of sixteen years our position was third. The position of the several towns being London 18.7, Edinburgh 18.9, Leeds 19.4, Birmingham 20.1, Sheffield 20.4, followed by Glasgow 21.5, Manchester 23.1, Liverpool 24.1, and Dublin 26.1. Leeds, I believe, was the only one of the provincial towns which did not enlarge its area during these sixteen years.

Infant mortality.—The mortality amongst children in 1905 was very much below the average of the previous fifteen years. During the years 1890-1904, as shown on the chart in the Annual Report for 1904, there was an average of 176 deaths in children under one for every thousand born. Last year the rate fell to 152; a rate lower than in any of the

TABLE 2.

Shewing the death-rates in the nine largest towns of the United Kingdom for the 52 weeks, and each of the thirteen week periods of 1905.

	First quarter of 1905.	Second quarter of 1905.	Third quarter of 1905.	Fourth quarter of 1905.	52 Weeks.
Leeds -	15.7	14.6	15.3	15'4	15.3
London -	16.8	14.3	14.6	16.8	15.6
Edinburgh -	17.8	16.1	14.5	16.5	16.5
Birmingham -	16.9	15.7	15.6	16.4	16.5
Sheffield -	17.8	17.5	18.3	14'4	17.0
Glasgow -	19.4	18.5	15.2	18.5	17.9
Manchester -	18:4	16.9	19.4	17.3	18.0
Liverpool -	20.4	17.7	19.9	20'4	19.6
Dublin -	24.6	20'4	19.6	20.4	21.3

The rates are calculated upon the populations estimated by the Registrar-General for England since the results of the 1901 census were published.

fifteen years. In 1894, the first of our specially healthy years, the rate had been 156. In 1903, the healthiest year until the one we are speaking of, the children's death-rate was 153.

Table 5 (page 15) shows in the second line of figures the ratios of mortality in children under one year of age to each thousand births registered during each of the quarters of 1905.

In the third line the rates are calculated in the same way upon the births not in the same quarter but upon the mean births in that quarter and the four preceding it. The same table gives also in the first line the death-rate of children under one upon the population at that age estimated from the census returns.

TABLE 3.

Births and deaths registered in the City of Leeds in the four periods of 13 weeks ended respectively April 1st, July 1st, September 30th, and December 30th, 1905. Deaths in age groups.

		Мо	MORTALITY FROM ALL CAUSES AT SUBJOINED AGES.								
1	2	3	4	5	6	7	8	9			
1905. Estimated population at	Regis- tered Births.	At all Ages.	Under I Year.	and under 5.	5 and under 15.	and under 25.	and under 60.	60 and upwards.			
these ages		456,787	11,964	41,144	93,645	91,934	192,143	25,957			
I. Quarter	3,255	1,819	397	175	75	94	509	569			
II. Quarter	3,214	1,720	373	182	74	106	494	491			
III. Quarter	3,081	1,790	668	202	58	70	404	388			
IV. Quarter	2,787	1,795	437	260	62	55	488	493			
52 weeks	12,337	7,124	1,875	819	269	325	1,895	1,941			

The diminution in infantile death-rate, it will be seen, was not entirely due to lessened mortality in the third quarter. The rate was less in the first quarter than the average either of the five preceding years or of the years 1886-1890. So also in the second quarter. It was appreciably but not so markedly less in the third quarter, and it was considerably less again in the fourth quarter.

It is fair to assume that the absence during the greater part of the year of measles and whooping-cough, in epidemic form, along with the usually corresponding diminution in the death-rate from bronchitis and broncho-pneumonia accounted

TABLE 4.

Birth and death rates in the City of Leeds in the four periods of 13 weeks ended respectively April 1st, July 1st, September 30th, and December 30th, 1905. Death rates in age groups.

	DEATHS PER ANNUM PER 1,000 LIVING.									
1	2	3	4	5	6	7	8	9	10	11
1905.	Birth- rate.	At all Ages.	Under I year.	and under 5.	5 and under 15.	and under 25.	and under 60.	60 and upwds.	25 to 65	Over 65
I. Quarter	28.6	16.0	133	17.1	3.5	4.1	10.6	88.0	13.5	111.8
II. Quarter	28.2	15.1	125	17.8	3.5	4.6	10.3	75.9	12.7	92.7
III. Quarter	27·I	15.7	224	19.7	2.2	3.1	8.4	60.0	9.9	78.7
IV. Quarter	24.2	15.8	147	25.4	2.7	2.4	10.5	76:2	12.0	101.7
52 weeks	27.1	15.6	157	20.0	2.9	3.2	9.9	75.0	12.0	96.2

in great part for the diminution of infantile mortality throughout the year, although one would fain hope that some of the work we have now done for several years is beginning to bear fruit. The subject, however, is so important that I propose to return to it in the supplementary portion of the present report.

JAMES SPOTTISWOODE CAMERON.

February 8th, 1906.

INFANTILE MORTALITY.

(ADDENDUM TO PART I.)

The subject of infantile mortality was referred to briefly in the preliminary portion of this report. When the work of preparing the tables was pretty well in hand a suggestion for a new table was received from the Local Government Board, and as we were intending to deal rather more fully with the subject of infantile mortality we availed ourselves of the opportunity of not only getting out this table for the whole City as suggested by the Board, but of making further use of the material collected in the several districts for each of the diseases and for each of the age groups in the full table. The table on the lines asked for will be found on page 10.

The deaths of all children under one year of age registered during the fifty-two weeks of 1905 are accounted for in this table, and they are divided into age periods dealing with every week of age for the first month, and with every month of age for the rest of the year. We have added to the table as suggested by the Local Government Board a few extra lines. Where it has been doubtful whether it was intended that a particular disease should be included in a group we have put it in under a separate heading.

For instance in the first group, that of common infectious diseases, diphtheria and croup are included in one line. We have given three lines to the group; one for diphtheria so called, one for membranous croup, and one for other cases of croup not stated to be membranous or spasmodic.

In the same way in the group of diarrhoal diseases we have given two lines to gastritis and gastro-intestinal catarrh, and have included in the latter three deaths certified as due to gastric catarrh.* We have also added a third line for gastro-enteritis. Cases of gastro-enteritis have not been included in the line for "other causes" at the bottom of the table.

^{*} One aged 3 months, two aged 6 months.

Infantile Mortality during the year (52 weeks ended 30th December) 1905. Deaths from stated causes in weeks and months under one year of age.

	Total deaths under one year.	1857	H H & G L : 6 10	302 26 10 24	279 70 11 07 11 82 114	2 0 0 6 5 8 8	2 29 12 33 134	156	17 56 90	1863
	Nine and under twelve months.	249	: :11: 11: 1	6 :	1111 2 1821	n :n :r	:: 20	31 22 14	9 83:	249
	rr-r2 mths.	%:	::0:+::4	r + : : w	:::: ~:04	*: H: M	:: 0 + 4	12 2 2	:00 m	08
	ro-rr mths.	∞ :	::4:::700	24::4	1111 71110	H : a : w	::000	00 + 1	: 00 01	80
e.	9-10 mths.	35 :	::**:::0	130 H : : 7	:::: >:+6	::0:=	:: + m +	10:2	:	85
or age.	Six and under nine months.	305	::11::16	68 : 23 : 5	13.1.1.25	44417	33.	345	1 18 16	305
year	8-9 mths.	S :	::0:::750	17 :: : :	:=::::	онн ч ;	: 0 0	110	:00	95
one y	1-8 1 hx	56 :	::**:::	3 : : 3	:=:: >:::	a a : : +	1188	16 10 15	:00 m	94
	6-7 mths.	911	::"::::	\$ n : u w	:::::::::0 +-0	: = m : m	: 0 + 0 0	12 17 1	101	911
under	Three and under six months.	374	::".::6	H 6 4 10 10	eм :: 4мой	3 10 10	9 4 17 15	34 29 10	13	374
months	5-6 mths.	112	:::::::	E = = : 0	7 7 1 :	H : M : H	: 8 = 0 =	5 5 : 4	; co a	112
moi	4-5 mths.	120	::"::::	37 : 2	w:::::::	4 4 N : 4	: 10 8 10 10	11 70	H (J) 1/J	120
and	3-4 mths.	142	::::::	1 6 6 1 6	13 6 1 2 1 6 1	: 4 2 7 5	: 01 H 10 =	11 7 : 5	H 60 F	142
weeks	One and under three months.	347	:-::::#	965444	23 14 15 16 16 17	102:	18 : 40	36 21 5	18	347
n we	2-3 mrbs.	147	:*:::::	m 01 : H 04	40 : : 0440	1 = 0 1 0	- w : a g	5 : 5	4 7 6	147
-	1-2 mths.	500	::::::	27 + + : 2	20 2 2 3 2 3 4 5 5	; H 60 ; 61	: 60 50 50	:: 0 33	: 0 10	200
causes	Total under one month.	582	T ::::::	125 : 23	252 555 11 11 145 151 18	:::	16 : :69	13 2 2	10 :12	288
stated	3-4 wks.	75	:::::::	ен::>	\$ s : : V : 0 0	::"::	- + : : 0	97:1	::9	92
	2-3 wks.	8:	*::::::	r:":"	00 nu : : 0 = 10 th	:::	: : : : 2	e, ::	8 : 10	8.
from	1-2 w ks.	95 :	:::::::	::.	00 HH 00:0H	:::::	: - : : 2	m~::	1 : 1	56 :
ths	r week.	322	:::::::	:*:::	174 37 10 26 	:::5:	: 6:: 12	H::H	r:0	327
Deaths	CAUSES OF DEATH.	All causes { Certified	Cricken-pox	Charloga, all forms Enteritis (not tuberculous) Gastritis Gastro-intestinal catarrh Gastro-enteritis	Congenital defects E.S. Vant of breast milk	Tuberculous meningitis Tuberculous peritonitis Tabes mesenterica Hidrocephalus	Erysipelas	Bronchitis Broncko-pneumonia Laryngitis Pneumonia	Suffocation, over-laying	

In the group of wasting diseases we have given three lines to atrophy, debility and marasmus, as they are not grouped together in our death ledger, and we have added a further heading for inanition, malnutrition and privation. These three causes are entered on the same page of our death ledger with "want of breast milk," which cause, however, was specially marked and easily separated. It has been put, as suggested, in the table in a separate line.

For a similar reason tuberculous peritonitis and tabes mesenterica have been separated into two lines and hydrocephalus added in a line to itself. The line "other tuberculous diseases" therefore does not contain the line last named; it does contain, however, three deaths certified as tuberculosis of the lungs, tuberculous broncho-pneumonia and phthisis.

In originally drawing out the table we included a line for scrofula, which has been kept separate in our death ledger, but as there was no death from that cause we have struck it out.

Further down in the table, after "bronchitis" we have introduced broncho-pneumonia which is kept separate in our books. In drafting the table we looked also for cases of pleuro-pneumonia and pleurisy, but there was no death under one year from either of these groups; we have therefore struck the line out of our original draft.

It will be understood therefore, that where a disease is called "pneumonia" or "croupous-pneumonia," or "lobar-pneumonia," it has been entered under the head of pneumonia; but where it is called broncho-pneumonia, or catarrhal-pneumonia, or lobular-pneumonia, it has been entered under the head of broncho-pneumonia. It is probable that a great many of the cases entered as pneumonia ought to be classed along with this group, but it is not quite clear whether the intention of the Board was to include them under bronchitis

or under pneumonia, or to exclude them altogether. We have therefore followed the classification adopted in 1892, and since, in table C.

In each case where we have added a line to the table it has been printed in italics. Where we have simply split up the components of one of the Board's lines the type has not been altered.

In another particular we have also departed from the Board's table. In the form sent out the deaths for each of the first four weeks of life are placed in separate columns, and the total deaths under one month collected together in another column. We have carried this principle of collection a little further, and have added columns for the deaths of those over one and under three months, over three and under six months, over six and under nine months, and over nine and under twelve months. The figures in all these age groups have been printed in heavier type, and are inclosed by heavier vertical lines.

We have taken for granted that in the fifth column "total under one month" are only to be included deaths of children of less than four weeks of age. In all other parts of the table we have interpreted the month as being a calendar month.

The 1,863 deaths recorded in children under one during the 52 weeks were those of infants born in Leeds. In addition to these, 12 children brought into Leeds, after birth, died in the first year of life. In the general estimate of the infantile death-rate elsewhere in the report, these 12 are included, for though their births were not registered in Leeds, it is possible that a nearly corresponding number of children actually born in Leeds may have left the town, and died outside. The age periods of these twelve outsiders who died in Leeds were as follows:—under one week, 2; from two to three weeks, 1;

three months and under four, 4; four to five months, 2; six and under seven months, 2; ten to eleven months, 1; total 12. Two died in the Women and Children's Hospital, ten in the General Infirmary, both institutions in West Leeds.

Adding the 12 to the 1,863 the deaths of children under one were equivalent, as will be seen from tables 4 and 5, to a death rate upon the estimated population under one year of age of 157; the rates for the several periods of thirteen weeks making up the fifty-two, having been 133, 125, 224, and 147. This however, as has been previously pointed out, is not the usual way of calculating infantile mortality, and in the second line of table 5* will be found the rate of mortality as compared with births registered during the same period. The rate for the whole year (52 weeks) was 152 deaths registered for every thousand births registered in the same period, and the rates for the several quarters in the same way 122, 116, 217, and 157 per thousand children whose births were registered during those special quarters. As has been frequently pointed out in these reports, the children under one year whose deaths are recorded in any particular quarter were not necessarily born in that quarter of the year, and in the third line in table 5 the deaths in each quarter (given in table 3) are calculated per thousand of the average number of births registered in the same and the four preceding quarters. This method of calculation gives sufficient prominence to the seasonal birthrate by counting the births registered in the corresponding quarter in two separate years, but it prevents the accidental and abnormal increase or decrease of registrations in any particular quarter from exerting too great an influence upon the rate per thousand. The rates for the several quarters (each of thirteen weeks) of 1905, calculated in this way, were 126, 118, 213, and 142, the average of these numbers being 150 instead of 152 as in the second line.

^{*} Page 15.

It will be noticed that this mode of computation makes the death-rate in the first and second quarters a little higher than when the deaths are calculated upon the births registered during those quarters. In the third or diarrhœa quarter the rate comes out slightly lower, and in the fourth quarter considerably lower, than by the second method. The difference of two per thousand in the whole year, and of varying amounts in the third and fourth quarters, is in part due, doubtless, to the larger number of births in the earlier year.

It will be seen on consulting table 5 that the death-rate in children under one in each quarter of last year, as given in line 3, was considerably lower than the average rates calculated in the same way in the five preceding years. It will also be seen that the average rates in those five preceding years did not differ very greatly from the average calculated by the second method for the five years 1886-1890. While, therefore, each quarterly death-rate last year was considerably below the average of the earlier period named (1886-1890), it was also below the average of the five years preceding 1905. Its lowness in 1905 must be therefore looked upon as to some extent due to special circumstances and not to any very great change in our methods.

In connection with this matter I should like to refer you to the tables and chart of infantile mortality given in the report for 1904. In the table on page 21 of that report the death-rate of infants under one per thousand births registered in the same period is given for each year and for each quarter of the year from the beginning of 1890 to the end of 1904, the rate for that last entire year being 176. According as the difference for any particular quarter and any particular year was above or below that 176, which curiously enough was the average for the whole period of fifteen years embraced in the table, that difference has been expressed as a percentage of the 176 above or below as

Mortality in Children under one year of age, during the 52 weeks of 1905.

	First Quarter.	Second Quarter.	Third Quarter.	Fourth Quarter.	YEAR.
Calculated per 1,000 of the population under 1, estimated to the middle of 1904, on the supposition that the whole population of the city was that estimated by the Registrar General, and that the number of children under					,
one bore the same proportion to the population as at the 1901 census Deaths under 1 per 1,000 births registered in same	133	125	224	147	157
Deaths per 1,000 registered births, the latter instead of those for the same quarterly period being the average of the same and four preceding quarters. The rate for the year is the mean of the four quarterly rates	122	118	217	142	150
Average rate by last method for five preceding years	155	149	234	164	176
Average rate by second method for years 1886-90 (from table 711890 report)	158	147	223	171	175

the case may be. That table showed that during those fifteen years, in both the first and the second quarters, the death-rate was always below that of the whole period:-that in the third quarter, with the exception of the year 1902, the rate was always above that of the whole period, and that in the fourth quarter on three occasions the rate was above the average of the whole period, on one occasion was the same as that average, and in the remaining eleven years below the average. This is another way of saying the third quarter is the quarter in which the rate of mortality was almost always highest. In that third quarter, during those fifteen years, the rate varied from the exceptional year 1902, when it was 2 per cent. below the average of the whole period, to the exceptional years 1901 and 1897, when it was respectively 82 and 81 per cent. in excess of that average. It will also be noticed in this connection that in the three years in which the mortality in the third quarter was more than 75 per cent. above the average of that of the whole period, the mortality of those years taken as a whole was also above that average, but by no means proportionately so. In 1893 the rate in the third quarter was 76 per cent. above the average of the whole period, and the rate for the whole year was 17 per cent. above. But that year was interposed between two in which the year's mortality was in the one case 5 per cent., in the other 11 per cent. below the average. In the year 1901, with the highest third quarter mortality (82 per cent.), the rate for the whole year was only 7 per cent, above that of the whole period, but here the rate of the preceding or second quarter had been as much as 28 per cent. below that average.

The two tables on page 22 in the 1904 report show the corresponding average temperatures for the same quarters of the year, and the total rainfall in inches during the same quarters, distinguishing in the latter case those quarters which consisted of fourteen instead of thirteen weeks. The material in these tables was put in a graphic form on a chart opposite page 26, which was constructed on the principle of showing for

Mortality of Children under One, per 1,000 born in the same period, for each quarter of the 15 years 1890-1904, and for each of those years along with in each case the amount per cent. above or below the average of the whole period.*

7 11 12	1		I	I.	11	I.	1,	v.	YE	AR.
	Deaths per 1,000 Births.	above+ below-	Deaths per 1,000 Births,	above+ helow-	Deaths per 1,000 Births.	above+ below-	Deaths per 1,000 Births.	above+ below-	Deaths per 1,000 Births.	above+ below-
1890	152	- 14	143	- 19	217	+ 23	178	+ 1	173	- 2
1891	156	- 11	173	- 2	205	+17	175	- I	177	+ 1
1892	159	- 10	139	- 21	209	+19	166	- 6	168	- 5
1893	175	- 1	167	- 5	310	+76	171	- 3	206	+17
1894	138	- 22	131	- 26	184	+ 5	173	- 2	156	- 11
1895	163	- 7	133	- 24	296	+68	176	{ + o	191	+ 9
1896	155	-12	164	- 7	207	+ 18	146	- 17	169	- 4
1897	141	- 20	118	- 33	319	+81	187	+ 6	190	+ 8
1898	145	- 18	140	- 21	286	+63	160	- 9	183	+ 4
1899	141	- 20	128	- 27	262	+49	156	- 11	172	- 2
1900	173	- 2	160	- 9	241	+ 37	157	- 11	183	+ 4
1901	143	- 19	127	- 28	320	+82	164	- 7	188	+ 7
1902	152	- 14	142	- 19	172	- 2	172	- 2	160	- 9
1903	131	- 26	116	- 34	177	+ 1	190	+ 8	153	- 13
1904	160	- 9	147	- 17	238	+35	158	- 10	176	{ + o

each quarter the percentage above or below the mean of the whole period, and the percentage by which the temperature exceeded or fell short of the average of the mean temperature of the period 51.59 degrees

^{*} Reprinted along with the two tables on page 19 from the Annual Report for 1904.

The same chart showed the rainfall in each quarter in inches in a graphic method. On page 23 a note explanatory to this chart and drawing attention to some of the variations will be found. For convenience, the tables which are still in type have been inserted here on pages 17 and 19.

Age groups under one.—It will be seen* that of the total 1,863 deaths 588, or 316 per cent., were those of children under four weeks of age, whilst the deaths between the end of the first four weeks and the close of the first three months of life amounted to 186 and the total deaths in the first three months to 502 per cent. of the total deaths under one. During the next three months of age—more than three and less than six months—the percentage was 201. The deaths which occurred in the six earlier months of life were 7026 per cent. of the whole mortality in children under one.

The next two periods of three months contributed 16:37 and 13:37 per cent. respectively. These figures are without taking into account the 12 deaths of new comers into Leeds. Of these three died within the first four weeks of life, and added to the 588 they make 31:5 per cent. of the total 1,875 deaths under one including the 12 outsiders. In the same way when we take the whole three months the mortality is 50 per cent. of the whole, whilst of those under six months of age the percentage is 70:3, the proportion of deaths at these ages not being materially affected by these 12 outsiders. It is noticeable that whether we include or exclude these outsiders half the deaths occurred during the first three months and 31 per cent. within the first month of life.

^{*} See table, page 10.

Mean temperature in degrees Fahrenheit for each quarter and for each year from 1890 to 1904 inclusive.

	I.	II.	III.	IV.	YEAR.
1890	42.9	55.0	61.6	43.5	50.6
1891	40.5	55.0	61.3	43.8	20.1
1892	40°I	54.6	59.7	42.3	49.2
1893	42.3	59°I	63'9	46.3	52.9
1894	42.6	22.1	61.0	47.5	51.6
1895	36.8	57.9	64.0	44.9	50.4
1896	44'I	58.5	61.3	42.9	51.9
1897	41.7	56.0	63.4	47.3	52.1
1898	44°I	55'3	64.3	49'I	53.2
1899	42.6	56.2	64.5	45'9	52.4
1900	40'1	55'9	64.0	48.9	52.2
1901	40'I	58.0	66.0	45'3	52.4
1902	42.6	54.0	60.9	47.2	21.1
1903	45.2	55.8	61.4	45.6	52.1
1904	41.4	56.4	63.5	46.2	21.9
Means	41.8	56.2	62.7	45.8	51.6

Rainfall-inches.

	I.	II.	III.	IV.	YEAR.
1890	5.65	4.86	8.71	3.26*	22.62*
1891	2.36	4.88	7.08	10.35	24.64
1892	4.27	8.02	5.99	9.00	27.28
1893	4.35	2.96	8.51	6.34	21.83
1894	8.78	6.70	5.69	6.81	27.98
1895	6.27	4.86	10.00	8.79	29.92
1896	4.48	5.35	6.24*	8.72	24.76*
1897	7.67	5.68	5.72	6.04	25.11
1898	2.83	7.05	4.10	7.47	21.24
1899	5.67	6.11	6.03	5.29	23.40
1900	8.36	4.88	9.17	8.14	30.22
1901	5.03	3.84	3.25	8.93	21.05
1902	3.12	5.68	6.12	5.95*	21 01*
1903	6.62	5.93	7.86	5.95* 8.83	29.24
1904	7.05	4.40	5.65	3.44	20.24
Means	5.20	5.41	6.66	7:20	24.76

^{*} The fourth quarter of 1890, and third of 1896, and the fourth of 1902 contained 14 weeks. The total rainfall for the quarter has been corrected by deducting a fourteenth part; and a fifty-third part of the total has been deducted from the amount for the year for the same periods.

```
327 under one week ...
                                = 17.6 per cent. of the 1863.
 95 over 1 under 2 weeks...
                                ... 5'1
 90 ,, 2 ,, 3 ,, ...
                                ... 4.8
 76 ,, 3 ,, 4 ,, ...
                                ... 4'I
  - 588 ,, 4
                                   31.6
200 over 4 weeks under 2 months
                                ... 10'7
147 ,, 2 months ,, 3 ,,
                               ... 7'9
--- 935 under 3 months
                                        50'2
142 over 3 months under 4 months
                                ... 7.6
                                ... 6.4
120 ,, 4 ,,
                ,, 5 ,,
                                ... 6.0
112 ,, 5 ,,
                ,, 6 ,,
—— 374 over 3 months under 6 months ——
116 over 6 months under 7 months
 94 ,, 7 ,, 8 ,,
 95 ,, 8 ,, ,, 9 ,,
                                ... 5.1

 305 over 6 months under 9 months — 16.4

 85 over 9 months under 10 months
 84 ,, 10 ,, ,, 11 ,,
                                ... 4'5
 80 ,, 11 ,, ,, 12 ,,
                                ... 4'3

 249 over 9 months under 12 months — 13'4
```

It is noticeable that in the first week of life the mortality was greater than in any single month except the first. It was greater also than in the last period of three months (that is from nine to twelve months). It was also greater than that of the period which preceded it, *i.e.*, six to nine months.

It will be noticed also that the first week mortality accounted for more than half of all the deaths in the first four weeks; and the deaths in these first four weeks were nearly three times as many as in the second calendar month.

These figures deal merely with the deaths at certain short age periods as compared with all deaths in children under one year of age. They may also be compared with the births during the same period. This has been already done for the total deaths under one in table 5 and the accompanying portion of the text of this report. The comparison at the shorter age periods would, however, only give us the several proportions—32 per cent. under four weeks, 50 per cent. under three months, &c.—of the whole infantile death rate of 152 per 1,000 births.

There is evidently a slight error in thus estimating death rates. There were 12,337 births in the period we are considering. It is not very far wrong to say that the 588 deaths in the first month of life might be calculated on these 12,337 births as an approximate population. But evidently in calculating the death rate in the second month of life we ought to deduct at least these 588 deaths from this imaginary population. Similarly, in calculating the mortality at six to twelve months of age, we ought to diminish the births by at least the 1,309 deaths which had occurred in the six earlier months and also by rather more than half the deaths that occurred in the same (or later) six months.

On this assumption we may regard the death rate during the first week of life as approximately 26.5 per 1,000 living, calculated on all births in the year. During the first month it would be 49.3, calculated on the births less the deaths in the first fortnight in life.

During the second half of the first year of life the mortality would be 51.7, instead of 44.9 calculated on the whole births in the year.

Similarly the mortality per 1,000 living of children of less than one year of age would be approximately 169 instead of 152 if calculated on the whole births. This differs also from the 157 per 1,000 given in tables 4 and 5 as the yearly rate calculated upon the population estimated from the

census figures. The estimate 169 is unfortunately more likely to be correct. The probable error from the immigration of babies into Leeds is perhaps negligable.

Age groups in districts.—It is important, however, to consider the age groups in the several districts. In the whole city the mortality under one was 152. In North Leeds it was 169, in West 150, in South-East registration district it was 203. Thus in the township of Leeds in the North part the infantile mortality was 11'2 per cent. above that of the whole city, in the West part it was 1.6 below the city rate, whilst in the district of South-East Leeds, always having a high rate of infantile mortality, the rate was no less than 33.7 per cent. above that of the whole town. In some of the outer districts the difference was not very great. Hunslet was 5 per cent. above, Holbeck and Wortley 3 and 2 per cent. below the town rate, Kirkstall was 20 per cent. below, Bramley 33 per cent., and Chapeltown 27 per cent. below. Evidently the outer and less populous districts had a lower rate of infantile mortality, the highest, South-East Leeds, being 34 per cent. above, Kirkstall, Bramley, and Chapeltown being 20, 33, and 27 per cent. below the town rate.

When we come to consider the incidence of mortality in the first week of life, which was 26.7 on the births in the whole city, it will be found that in Kirkstall, in which the rate in all children under one was 20 per cent. below that of the city, that in the first week of life it was 34 per cent. above the city rate at that period. Bramley and Chapeltown both had rates, the one of 2, the other 8, below that of the city. South-East Leeds had a rate of 12 above that of the city, as compared with 34 excess in the whole year of life. The North district was 18 per cent. below, and the Wortley district 14 per cent. above. The West district, which had a rate of only 2 per cent. below that of the whole city in the whole year, had a rate of 10 per cent. below in regard to the first week of life.

Infant mortality during the year (52 weeks ended December 30th), 1905. Excess or otherwise compared with rate in whole city of district rates at certain age periods.

District.	Births in year.	Under 1 week.	Under 1 month.	1 and under 3 months.	3 and under 6 months.	6 and under 9 months.	9 and under 12 months.	Year.
North	1,792	- 18.4 %	- 4.6 %	+26'1 %	₹ 26.2 %	+ 5.2 %	+12.8 %	+11.5 %
West	1,905	- 9.7 %	- 9.2 %	+ 3.9 %	- 5.2 %	+ 8.8 %	+ 1.0 %	- 1.6%
South-East	1,235	+12.4%	+16.5%	+51.6%	+64.6 %	+ 39.8 %	- 4.4 %	+ 33.7 %
Hunslet	2,140	- 3.7 %	+12.9%	+ 3.9 %	- 6.6 %	+ 10.8 %	- 1.0 %	+ 5.1 %
Holbeck	1,071	- 2.2 %	- 4.6%	- 27.6 %	- 5.2 %	+ 1.3 %	+33.5%	- 3.0 %
Wortley	1,643	+13.9%	- 2.3 %	- 16.3 %	+ 13.7 %	- 9.6 %	+ 4.9 %	- 2.0 %
Kirkstall	981	+ 33.7 %	+ 6.3 %	- 31.4 %	- 29.8 %	- 46.6 %	- 19.7 %	- 20.2 %
Bramley	459	- 2.2 %	- 22.9 %	- 45.9 %	- 64.3 %	- 12.4 %	- 14.3 %	- 32.7 %
Chapeltown	1,016	- 7.9 %	- 7.7 %	- 16.6 %	- 58.0 %	- 32.9 %	- 32.0 %	- 26.9 %
Osmondthorpe	3							
City death rates per 1,000 births registered	12,245*	26.7	48°0	28.3	30.2	24.9	20.3	152.1

If we now turn to the first month, it will be found that South-East Leeds and Hunslet had rates respectively of 17 and 13 per cent. above the rate of the whole city which was 48 per thousand. North Leeds was 5 per cent. and West Leeds 9 per cent. below that rate, Holbeck and Wortley 5 and 2 per cent. below, Kirkstall 6 per cent. above, Bramley on the other hand was 23 per cent. and Chapeltown 8 per cent. below the city rate.

Taking now the next two months of life (one and under three months), the rate in the whole city was 28.3 per thousand. The rate in South-East Leeds was 52 per cent above, that in Bramley was 46 per cent below, North Leeds

^{*} Exclusive of 92 births in workhouses.

Infantile Mortality during the year (52 weeks ended 30th December) 1905, from stated causes in each registration district, with rates per thousand births in the district.

200	ear ooo		N N CO	000000000000000000000000000000000000000			h 2 4 4			V.
City death-	1000	7.00 7.00 7.00 7.00 7.00 7.00 7.00 7.00	247	5.55	250	5.0 7.2 7.2	12.7	7.7	1.251	12,245
OSMOND- THORPE,	Rate per 1,000 births.	:::::::	:::::	1111 111	:::::	:::::	::::	:::	:	23
OSA	Destpe		1::::			:::::	::::	:::		
PEL,	Rate per 1,000 births.	3::::::	8.6	7.01 3.9 5.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7	2.0	2:0	9.8 5.9 	3.0	2.111	910
CHAPEL-	Destps.	:::::::	2 50 : : 50	74:: 7:52	: = 0 : 0	: 01 H :00	10 0 1	ппо	113	1,016
ILEY.	Rate per r,000 births.	::0.0	7:5:5	9.61 2 :: 5		34.5	0.5	:67	t.201	0
BRAMLEY.	Deaths	::"0::::	2 : 1 : 2	0::: 4::1		: : 4 4	10 to : H	: 40	47	459
TALL	Rate per r,000 births.	::::::.6	9.5	27.5 10.2 10.0 10.0 10.0 3.1 5.1	10: 10: 14	12:22	3.1	3.1	1213	-
KIRKSTALL	Deaths.	:::::::::::::::::::::::::::::::::::::::	O. O. H : N	101 - 6:01	4 : H : 4	: + : + >	20:1	m % 0	61.1	981
LEV.	Rate per 1,000 births.	::2:::%	27.0	27. 120 30 : 120 43. 120	\$ 200:	30	8.5	17.2	1.641	43
WORTLEY.	Destps.	::":::0	8::00	£∞ 4 : 2 20 L X	::+	: H 0 10 4	77::	400 7	245	1,643
ECK.	Rate per 1,000 births.	:::::::::::::::::::::::::::::::::::::::	2.87	\$43.3	2 : 5 : 4	5.000	5.6	5.0	5.141	1,1
HOLBECK.	Deaths.	::::::	g m : : >	00 m : 0 : 40	a : w : w	: = 0 0 0	25 1 25	12 0	158	1,071
ET.	Rate per r,000 births.	. : : : : : : : : : : : : : : : : : : :	27.7 000 000 005 005	25 :: 25 :: 25	7:000	20 20 20 20 20 20 20 20 20 20 20 20 20 2	85.9 0.55 7.5	5.6	8.651	0
HUNSLET.	Dearps.	:: > 0 : : :	CH 10 10 00	55 2 : : 2 52	w: 47 4	H 10 H 9 10	34 19 16	1.5	342	2,140
-31.	Rate per r,000 births.	3.5 ::::	5.45	27.5 3.2 1.6 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0	5.00	5.7	4.01 17.8 1.0	404	2.802	,235
South-		:: = ::: 04 +	15 E H : 7	# + 4 : \$: 0 8	0×1-1:	: 2 : 20	4 00 : 01	(U 1/2 (U	251	1,2
7.	Rate per 1,000 births.	2::::2::	377	5000 : 240	03.7 500	4.2 17.6 10.5	89 73	179	9.641	95
WEST.	Destps.	:::::::::::::::::::::::::::::::::::::::	# 1 2 1 8	800 8 : > 1 2 3	8 100 7 10	100 4+ 00 G	17 17 17 17 17 17 17 17 17 17 17 17 17 1	E 0 4	285	1,905
Н.	Rate per 1,000 births.	33	37.9	1337	0.0	33 000	13.4	379	1.691	92
NORTH.	Destps.	H H W : : : : 19	8 4 6 : 7	80 m: ∞: 128	::4	30 0 10 :	27 24	27.7	303	1,792
	CAUSES OF DEATH.	Small-pox Chicken-pox Measles Scarlet fever Diphtheria Ocit Other croup Other croup	B & Enteritis (not tuberculous) E Gastrius Gastro-intestinal catarth Gastro-enteritis	Congenital defects Congenital defects Injury at birth Want of breast milk Vais hamition, madnutrition, privation Atrophy Debility Marsamis	us meni us perit senterica halus erculous	Erysipelas	Bronchitis	Suffocation, over-laying	Totals	Births, exclusive of those in public institutions

was 26 per cent. above, Kirkstall 31 per cent. below, Holbeck Wortley, and Chapeltown all had rates below, whilst Hunslet was 4 per cent. in excess.

In children between three and six months old the rate was 30.5 per thousand in the city. It was 65 per cent. higher in South-East Leeds, 30 per cent. lower in Kirkstall. It was also higher in North Leeds, 26 per cent., and very much lower, 64 per cent. and 58 per cent., in Bramley and Chapeltown.

During the following three months of age, from six months to nine months, the town had a death rate of 25 per thousand born in the whole year. Chapeltown was 33 below that, South-East Leeds was 40 above, Kirkstall 47 below.

Taking now the last three months of the year, nine to twelve months, the city rate was 20'3 per thousand births. In South-East Leeds the deaths were 4 per cent. below that average, in Holbeck they were 34 per cent. above, in Kirkstall, Bramley, and Chapeltown the rates were 20, 14, and 32 per cent. below.

These figures dealing only with a single year are of course not of very great value, but they seem to point to the fact that the mortality in some of the least healthy districts is not so much in excess of that of the town in the earlier weeks as it is in the middle months. The contrast between South-East Leeds and Chapeltown is very marked. The latter district is 8 per cent. below the rate of the town in the first month, 58 per cent. below in the period from three to six months. South-East Leeds is 17 per cent. above in the first month, 65 per cent. above in the period from three to six months. This is a matter which additional figures of the same kind will probably throw greater light upon.

Causes of death in districts.—The new table on page 10 deals with the actual causes of death, amongst children under one, at each age group in the whole city. We have inserted (p. 24) a supplementary table dealing with all the deaths under one, during the year, from the same groups of causes according to the districts in which they occurred. The total deaths from each group of causes will be found in the last column of the earlier table. The rates from these, per 1,000 births in the city, are given in the last column of the new table. For each district, in the newer table, we have given the deaths from each group of causes that occurred in that district and the mortality from that group per 1,000 births in The Local Government Board has also by the district. horizontal lines grouped the groups. The first of these larger groups consists of seven common infectious diseases: smallpox, chicken-pox, measles, scarlet fever, diphtheria, membranous croup and whooping cough, to which we have added other forms of non-spasmodic croup. These must be distinguished from the seven diseases in columns 11 to 17 of the Registrar General's weekly table for the large towns, and to which he now gives the name of "principal epidemic diseases." It will be noticed that in the whole city the rates for this group of infectious diseases add up to 6.8; the number calculated on births and deaths would really be 6.6, several of the rates having been raised to the nearest figure to one place. Taking 6.6 at the rate in the whole city, it will be noticed that in the North district the rate from this group adds up to 7.3, which requires no correction. The difference between 7'3 and 6'6 is 10'6 per cent. of the latter—that is of the rate for the whole town. West district has a rate from these diseases below that of the town. In the South-East district, the one with the greatest mortality amongst children under one, we had only 7 deaths from these diseases, the printed rates adding up to 5.6 but really equivalent on the actual deaths to 5.7 per 1,000 births registered in the district

during the period. The difference between 5.7 and 6.6 is 14.4 per cent. of the latter. The rate from the group is thus 14'4 per cent. lower than in the city. Hunslet, on the other hand, a district in which the death rate of all children under one was only 5 per cent. above that of the whole town, had a rate from these diseases of 9.8 or 481 per cent. above that of the town. The 7 deaths in Holbeck from these diseases run up the rate in that district to close upon the average of the town, whilst the 11 deaths in Wortley-two from measles, nine from whooping cough-bring the mortality of that district quite up to that of the town. Whooping cough alone in Kirkstall gave a rate of 8.2-24 per cent. above that of the city, and that from one single disease of the seven. The high mortality from measles in Bramley, along with a single death from whooping cough, brought the rate of that district, which has a small population, up to 13:1 or nearly double the rate of the city; whilst the rate in Chapeltown is less than half that of the city.

Taking next the deaths from the various intestinal fluxes, it will be noticed that the whole death-rate from this group adds up in the last column to 30, only a decimal point from the calculated rate. In North Leeds the rates add up to 45.7, very considerably in excess of the rate in the town (51 per cent.). In West Leeds, on the other hand, they make 32.5, which is about 8 per cent. above the rate of the city. In South-East Leeds, the district with the highest infantile deathrate, they amount to 48.5 (or more correctly 48.6), that is to say, 61 per cent. more than the rate in the city. In Hunslet, on the other hand, the added rates are only 200, fractionally below the rate of the city. In Holbeck they are 22.4, 25% per cent. below. In Wortley 24.9, also below that of the city by nearly 17 per cent. In Kirkstall the added rates are only 14.2, considerably less than half the rate of the city. In Bramley they are 8.8, 71 per cent. below the rate of the city, whilst in Chapeltown they are 16.7, 45 per cent. below.

Taking individual rates in the rest of the table it will be noticed that from premature birth, whilst the rate was 22.8 in the whole city, it was below that rate in the North, West, Bramley, and Chapeltown districts, and above it slightly in Hunslet and Holbeck, and more considerably and nearly to the same extent in South-East Leeds, Wortley, and Kirkstall.

The very uncertain term "convulsions" accounts for 10'9 deaths per 1,000 births of children under one in the city. The numbers in West Leeds and in Hunslet were 10'5 and 10'7. In South-East and North Leeds 16'2 and 17'9, and in the remaining districts they were under the city rate.

Bronchitis, which is a convenient term to put in a death certificate, accounted for 12.7 deaths of children under one per 1,000 births in the city. This rate was slightly exceeded in the North, and a little more so (15 per cent.) in Wortley. In Hunslet the rate of 15.9 was a quarter more than that of the city, whilst in South-East Leeds the rate, 194, was half as much again as in the city. The rate from this cause was low in Holbeck (5.6) and more or less below that of the city in the other districts. If, however, we take broncho-pneumonia as part of bronchitis, the rate in the whole city becomes 22.7. In North Leeds it is 28.5, that is more than a quarter in excess. In West Leeds it was only 16.3, about as much below. South-East Leeds the combined rate from the two groups of causes was 37'2, more than half as much again. In Hunslet the rate 24.8 was slightly in excess. In Holbeck the rate 19.6 was also lower than in the city. In Wortley 23'1, it was slightly in excess, and in Kirkstall 15'3, a third below the In Bramley the rate was 17:4, between a fourth and a fifth below, whilst in Chapeltown it was 15.7, more than a third below. Overlaying was proportionately most fatal in Kirkstall, more so than in South-East Leeds, though the actual deaths were three in each case, so that if calculated on the whole population of the district instead of the births, the opposite relations would hold.

South-East Leeds and Chapeltown.-The first-named of these districts had a rate per 1,000 births below that of the town from the more infectious maladies, but a much higher rate from the intestinal fluxes. The rate was considerably higher from premature birth, but not so from congenital defects. It had a rate 80 per cent. above that of the city from inanition, malnutrition, privation, and marasmus. From tuberculous diseases other than consumption its rate was about a third more. From the next group of five diseases, including erysipelas, &c., the rate in South-East Leeds was half as much again as that in Leeds generally, whilst from lung diseases other than consumption the rate was 38.9, or 47 per cent. in excess of the rate (26.5) in the city-nearly half as much again. The remaining group, 8:8 against 13'4 in the city, showed in favour of South-East Leeds. So that it is the diarrhoeal diseases, the diseases of nutrition, and the tubercular diseases of childhood, that, along with such nondescript groups as convulsions and the convenient lung disease group, especially sent up the rate in South-East Leeds.

It is interesting now to compare the above with the rates in Chapeltown. We have already seen that in the infectious group and in the diarrhoeal group the rates in this district were very considerably below that of the town. In the third group the rate of 45.3, against 51.9 for the city, was nearly 13 per cent. below the city rate. In the tuberculous group the rate was 4.9 against 6.7, or 27 per cent. below the general city rate. In the lung disease group it was 16.7 against 26.5—37 per cent. below. In the last group the rates were about the same.

The interesting points about South-East Leeds are—(1) that while the death rate under one from all causes is very much in excess of that of the town, it is not so much in excess in the earlier weeks as it is in the middle months (see page 23); (2) that while the more infectious diseases did not cause anything like the havoc in that district that they did in some places, at least at this age, the diseases generally attributed to insufficient

and wrong feeding were greatly in excess, and along with them the diseases of a tuberculous nature. Singularly enough, cases of rickets are absent from the returns in South-East Leeds. Probably some of the deaths that might have been ascribed to that cause have gone to convulsions or bronchitis.

Deaths per 1,000 births in South-East Leeds, Chapeltown, and the city in children under one, from certain groups of causes, 1905:—

	City.	South-East Leeds.	Chapeltown.	Percentage of excess in South-East Leeds, as compared with Chapeltown.
More infectious diseases	6.6	5.7	3.0	90
Diarrhœal diseases	30.1	48.6	16.7	191
Wasting diseases	51.9	66.4	45'3	47
Tuberculous dis- eases	6.7	8.9	4.9	82
Fifth group	17.1	25.9	10.8	140
Respiratory diseases	26.5	38.9	16.4	133
Seventh group	13.3	8.9	13.8	-36
All causes	152.1	203.5	111.5	83

Contrasting South-East Leeds with Chapeltown, as is done in the annexed table, it will be seen that the rate from the more infectious diseases was 90 per cent. in excess in the former of that in the latter; that the diarrhœal diseases were in excess in South-East Leeds to the extent of 191 per cent. of the rate in Chapeltown; that the wasting diseases were 47 per cent. more

fatal in South-East Leeds than in Chapeltown, and the deaths from tubercle 82 per cent.; that in the miscellaneous fifth group the South-East Leeds rates were 140 per cent. in excess of those in Chapeltown; that in the lung disease group the excess was 133 per cent. On the other hand, in the concluding group the rates in South-East Leeds were 36 per cent. below those in Chapeltown. As a whole, the mortality under one per thousand births was in South-East Leeds 83 per cent. in excess of that of Chapeltown. For every 100 deaths in Chapeltown at this age period, 183 occurred in South-East Leeds amongst the same number of children born.

The reason South-East Leeds has been contrasted with Chapeltown rather than Bramley, will be evident from the following table for the ten years, 1894-1903:—

Infant Mortality, 1894-1903.

Chapeltov	vn		118	Sout	h-Eas	st	 214
Kirkstall			138	Holl	oeck		 189
Bramley			145	Osm	ondtl	norpe	 188
North			173	Hun	slet		 183
Wortley			175	West	t		 177
		City			1	74	

PART II. - SPECIAL DISEASES.

TUBERCULOSIS.

The number of deaths from consumption during the year was 558, making, with other forms of tuberculosis, a total of 846 or an annual rate of 1.86 per 1,000. This is the lowest annual rate from tuberculous diseases that I have had to report to you. In the year 1890 the rate was 2.52. From 1891 to 1895 it averaged 2.37. From 1896 to 1900 the average was 2.21. Including the present year the average from 1901 to 1905 has been 2.09.

Taking consumption alone the rate in 1900 was 1.66. In the five years, 1891-95 it was 1.59; in the five years

TABLE 6.

Mortality from tuberculosis, 1905.

Annual Control of the								
1905.	Tuberculosis, general and undefined.	Phthisis.	Hydro- cephalus.	Tuberculous meningitis.	Tuberculous peritonitis.	Tabes mesenterica.	Scrofula.	Total.
I. Quarter	34	178	2	26	4	5	_	249
II. do	27	137	2	2 I	18	17	_	222
III. do	28	117	2	23	8	18	2	196
IV. do	17	126	4	15	9	8	-	179
Year (52 weeks)	106	558	10	85	39	48	_	846
Annual death- rate, 52 weeks of 1905		1.53	0.03	0,10	0.00	0,11	-	1.86

1896-1900 it was 1.45. During the five years 1901-05 it has been 1.32. In both cases there has been a decline in each quinquennial period. These figures will be found in table 6b.

TABLE 6a.

Shewing deaths from each of certain groups of tuberculous diseases in the 52 or 53 week periods of the years 1890-1905, and the populations of those years estimated from the changing ratios of the 1881-91, and 1891-1901 intercensal periods.

YEAR.	Population by interpolation.	General or undefined.	Phthisis.	Hydro- cephalus.	Tuberculous meningitis.	Tuberculous peritonitis.	Tabes mesenterica.	Scrofula.	TOTAL.
1890*	363,018	92	612	20	92	14	92	6	928
1891	369,034	84	657	18	72	25	102	9	967
1892	375,081	84	530	14	65	14	93	9	809
1893	381,157	99	648	17	67	32	86	12	961
1894	387,259	79	576	9	72	18	61	13	828
1895	393,387	114	611	16	77	15	90	2	925
1896*	399,535	66	613	15	74	22	73	10	873
1897	405,716	102	589	14	91	36	99	2	933
1898	411,895	96	576	17	73	26	98	3	889
1899	418,101	86	596	9	102	33	80	I	907
1900	424,322	101	605	I 2	94	48	87	I	948
1901	430,555	107	605	7	75	46	136		976
1902*	436,800	109	584	4	100	36	80	3	916
1903	443,052	100	562	.8	106	37	82	I	896
1904	449,311	137	626	8	88	43	93		995
1905	455,573	106	558	10	85	39	48		846

^{* 1890, 1896,} and 1902 were 53 week years, the others 52.

TABLE 6 b.

Shewing deaths from different forms of tuberculosis since 1890 per 1,000 of the population, estimated at the varying ratios prevailing during 1881-91, and 1891-1901 intercensal periods.

			1000.					
YEAR.	General or undefined.	Phthisis.	Hydro- cephalus.	Tuberculous meningitis.	Tuberculous peritonitis.	Tabes mesenterica.	Scrofula.	TOTAL.
1890*	.25	1.66	.02	.25	.04	*25	.02	2.22
1891	.23	1.79	.02	'20	.07	.28	'02	2.63
1892	'22	1'42	.04	17	.04	.25	.03	2.16
1893	.26	1.41	.04	81.	.08	.53	.03	2.23
1894	'20	1.49	'02	.19	.05	.19	.03	2.12
1895	.59	1.26	.04	'20	.04	.53	.01	2.36
1891 }	.24	1.59	.04	.19	.05	.53	'02	2.37
1896*	.16	1.21	.04	.18	.02	.18	'02	2.12
1897	.25	1.46	.03	.53	.09	.24	.00	2.31
1898	.53	1'40	.04	.18	.06	'24	01	2'17
1899	'2 I	1'43	'02	.24	.08	.19	.00	2.18
1900	'24	1'43	.03	'22	.11	'2 I	.00	2.54
1896)	'22	1.45	.03	'21	.08	.51	.01	2.5 I
1891)	.53	1.2	.04	.50	.07	22	'02	2.59
1901	.52	1.41	'02	17	.1.1	'32		2.27
1902*	.25	1.31	.01	.53	.08	.18	.01	2.06
1903	.53	1 2 7	'02	.24	.08	.19	.00	2.03
1904	31	1.40	.02	.20	.10	.51		2,55
1905	.23	1'23	*02	.19	.09	.11		1.86

^{* 1890, 1896} and 1902 were 53 week years, the others 52. The death rates in each case are calculated on a year containing 52'17747 weeks. The rates for 1905 are calculated on a population of 456,787 in this table, as in the report generally, if calculated upon the interpolation population given in table 6a the rates from undefined tuberculosis and phthisis become 0'24 and 1'26 respectively, and the total death rate from the whole group 1'91 instead of 1'86.

LUNG DISEASES.

PHTHISIS.

In table 6c are given the rates of mortality from phthisis per 1,000 of the estimated population in each intercept of the town. As the rates are for a single year and the numbers in some of the intercepts very small, too much importance must not be attached to the death rate. For instance in the small triangle, bounded by New Briggate, Lowerhead Row and North Street, two deaths occurred, making a mortality of 9 per 1,000, but when the rest of the Central ward in the North registration district of Leeds is added, increasing the total mortality to 36, the death rate falls to 1.89. In this table the mortality is shown according as it occurred under the age of 25, between 25 and 50, and over the age of 50.

Since May 1901, voluntary notification of phthisis has been asked for in Leeds. In the seven concluding months of that year 282 cases of consumption were reported to us. The deaths in that year were 605. I give below the number of cases of consumption reported and the deaths from the same disease in each of the four following years.

YEAR.	DEATHS FROM PHTHISIS.	CASES REPORTED.
1902	584	452
1903	562	586
1904	626	631
1905	558	555

It will be noticed that during the two last of these years the deaths have been 1,184 and the cases reported 1,186. Although the number of phthisis cases living in any year is probably three times the number of deaths, we, of course, only ask to have each case reported once—not once a year. It

CITY OF LEEDS.

Instructions to Assistant Inspectors.

CONSUMPTION

about phthisis cases should be obtained with as little distress to the feelings of the patients or friends as possible, and the latter should be made to understand that it is for the sake of trying to help them that we are asking these questions.

Disinfection should be suggested after any fatal case but, in the meantime, not enforced against the wishes of the relatives. The disinfection recommended should consist of washing the walls, floors, ceilings, etc., with chloride of lime. (The Corporation would compensate to a certain extent for wallpaper damaged by this process.) All carpets, bedding, etc., to be taken to the steam disinfector.

Where the disease is reported during life a copy of the handbill should be given and

commented upon, and, with the consent of the doctor, disinfection carried out occasionally.

In examining the house note air space—give details for each room—number of occupants of each bedroom, etc. Give approximately the size of the windows of each room in the house, and the distance of neighbouring buildings. Especially state whether house opens into a wide or narrow street or court (give dimensions). Make a small plan of the situation of the house in regard to the surrounding buildings, stating approximate height of latter. Any information about the ventilation of the rooms will be of value.

Information should be obtained as to the patient's recent and previous employments, and especially if he has been engaged in any occupation in which there is much dust or which is carried on in an ill-ventilated room.

The question of whether any other member of the family or any near relative has suffered from the same complaint should be enquired into. The patient should have a separate bed, and, if possible, a separate room.

J. S. C.

SANITARY OFFICES,

LEEDS, February 3rd, 1902.

Phthisis. 37

seems therefore probable that we now hear of a very large proportion of the cases that occur in the town.

It may be asked what use do we make of the information. In every case heard of, either by direct notification from a medical man or through the Tuberculosis Association, unless requested not to do so, one of our inspectors visits the house and obtains the information which is usually asked for in cases of zymotic disease. Special instructions were issued to our inspectors in 1901 on a hektographed form. These we reprinted in February, 1902. A copy is given on p. 36.

The handbill which they are instructed to leave at the house, and if necessary explain, is printed the same size as our leaflets for mothers. The size is a convenient one for going into a notebook or the visiting case of a medical man. It will be noticed that the inspector is specially instructed only to ask for disinfection where the medical man in attendance approves.

The work of the Tuberculosis Association still continues. The number of beds at their command are now, at Gateforth, 20 for men and 14 for women, and at Armley, 24 for men, 10 for women, and two cots for children.

OTHER LUNG DISEASES.

The 558 deaths from phthisis form less than a third of the fatal lung diseases. A larger number of persons died from bronchitis than from phthisis, from broncho-pneumonia nearly half as many as from phthisis, from pneumonia, pleuro-pneumonia, pleurisy, and other lung diseases nearly as many as from phthisis.

I append for 1905 the deaths in Leeds from these diseases as well as those from phthisis, grouped, as in table 6c, according as those who died were under the age of 25, between 25 and 50, or over 50. In an earlier part of this report the death rate from lung diseases this year in comparison with earlier

TABLE 6 c.

Table showing deaths from phthisis at all and at certain ages, and death rates at all ages in the intercepts of the wards and townships of Leeds in the year 1905.

			Age.		TOTAL.	Deaths	
		-25.	25-50.	50+.	TOTAL	1,000.	
HOLBECK	Holbeck	10	20	2	42	1.37	
HOLBECK	TIT TT 1	19	6	3	10	2.69	
	West Hunslet			3	771		
HUNSLET	West Hunslet	12	II	- 1 -	24	0.86	
	East Hunslet	9	18	7	34	0.03	
	South		6	3	9	0.93	
SOUTH-EAST	South \ \2.68 [3	5	3	11	2.46	
LEEDS	South }2.68		I	3 6	4	3.24	
	East	11	18	6	35	1.50	
0						60	
OSMONDTHORPE	East	I	I		2	5.60	
Chapeltown	North-East	2	10		12	1.37	
(part of)	North Chap. All. 110	8	10	5	23	4.48	
	North Pottern'tn 1110	1	4	I	6	0.58	
NORTH LEEDS	North	10	7	5	22	1.25	
	North-East	10	23	9	42	1.88	
		I	19	14	34	1.81	
	Central in West } 1.89		2		2	9.08	
WORTLEY	New Wortley	11	11	- 4	26	1'41	
100	{Armley } 0.79{	7	- 11	3	21	0.67	
	\Wortley \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	3	4	4	II	1.19	
	Farnley	I	1		2	0.43	
Bramley	Bramley	9	7	5	21	1.14	
KIRKSTALL	Kirkstall	I	I	I	3	0.69	
	Burley 0.86	6	14	6	26	0.01	
	Headingley)	3	8	***	11	0.82	
CHAPELTOWN	Headingley)		I	1	2	1.21	
(part of)	North-West 1'14			***	***		
	Brunswick		2	I	3	1,00	
WEST	Brunswick	6	15	6	27	1.35	
	North-West	10	17	10	37	1.00	
	Mill Hill	2	8	2	12	1.69	
	West	15	16	9	40	1.69	
Сітч		162	277	115	554	1.55	
	Outsiders	2	2		4	***	

The rates are calculated on populations estimated by the Registrar General's method described at p. 88 of the Annual for 1904.

years has been commented upon (p. 5). Deaths from these disease groups, in children under five, are separated in tables A and C, and in children under one, and between one and five, in table 17.

Deaths from lung diseases, grouped according to nature of disease, and under age groups used for phthisis in certain tables.

1905.	-25.	25-50.	50 +	TOTALS.
Phthisis	. 164	279	115	558
Bronchitis	. 222	54	326	602
Broncho-pneumonia	. 263	6	18	287
Pneumonia and pleuro-pneumonia	124	100	128	352
Pleurisy	. 4	1	12	17
Other lung diseases	. 23	11	38	72
Totals	. 800	451	637	1,888

SEVEN COMMON ZYMOTIC DISEASES.

SMALL-POX.

The subject of small-pox was so fully gone into in the reports for 1903 and 1904 that it is not necessary to do more than give it a passing notice here. Table 8 last year contained a list of all the cases heard of as far as May 10th, 1905, In the corresponding table in this year's report these have only been reprinted from the end of the first quarter. The details about the three following cases have not been previously printed. The table contains also information as to two cases which occurred in the first quarter of 1906.

Small-pox had been somewhat prevalent in the latter part of 1904 and during January, February and March, 1905, it was present in Leeds to a very considerable extent. In the second quarter of 1905 we heard of six, in the third, of none, in the fourth, of three cases, whilst in the first quarter of 1906 we had two cases, in the second, none. The particulars about the renewed outbreak at the beginning of 1905 have been given in the report for 1904 and need not be repeated here.

From the 10th of May, 1905, until October 31st we heard of no case in Leeds. A case of small-pox occurred in Wakefield, and through the courtesy of the Medical Officer of Health we learned, on the 30th of October, that a man from Leeds had visited the same house about a fortnight earlier and was reported to have chicken-pox. The address given was not quite correct; but we succeeded in finding the patient, and ascertained the name of the medical man who was attending him. On communicating with this gentleman to the effect that a person in contact with his patient a fortnight earlier had developed small-pox in Wakefield, he notified the illness to us as one of that disease. patient (7191:8) was a lodger in Leeds, traveller for a firm of dressmakers' furnishers. He had been vaccinated in infancy, and had three marks covering together rather less than half a square inch. He made a good recovery. case was notified on the 31st of October; he was removed the same day. There were three other persons in the house, a man of 29, a woman of 30, and a small boy of four. All had been vaccinated in infancy. The man had four marks, covering nearly three-quarters of a square inch; the woman three, covering a little more than a square inch. The boy had three marks, covering a little less than half a square inch. All three were taken to our cottages on the Manston Hall Estate. All three were revaccinated on October 31st. The re-vaccination was unsuccessful in the man and his wife, and was repeated on the 6th of November, one vesicle afterwards appearing in each case. The little boy, although vaccinated not more than four years ago, responded to the first revaccination in one place. The man (7301:4)

Table 8.-Vaccination and Infectivity.

Showing cases of Small-pox reported from end of first quarter of 1905 to end of first quarter of 1906.

REMARKS AS TO SOURCES OF	INFECTION, ETC.	From 7055:7 From 7030:5 group. Note	77059:6. Note Ilkley Horsforth	Wakefield	From 7191:8	? Oldham From 7373:2
VACCINATION (IF ANY) AFTER CONTACT.		Mar. 26, succ. (E. Apl. 5) From 7055:7 Mar. 26, succ. (E. Apl. 6) From 7030:5		Oct. 31, not suc (Nov. 6, succ. (E. 13th) (Oct. 31, not suc. (E. 13th) (Nov. 6, succ. (E. 13th))		Feb. 22, succ. (E. Mar. 7) From 7373:2
RE-VACCINATION (IF ANY)	BEFORE CONTACT.	11				::
NATTON.	Date.	infancy never	infancy infancy infancy	infancy	infancy	infancy
VACCI	Area.	-101 E-1	x <2x -4+ -60	95(XC 95)-8	-100	Hos op
PRIMARY VACCINATION.	No. of marks.	1 :: 0	1000	ε 4	10	3.6
	Sex.	f. II.	: i - i	i i	ť	ii ii
	Age.	15° 5	28 32 41	29	30	\$48
	Ward.	W.H.* W.H.*	A. & W.	z ×	*.	E.H. *
	Zym. book No.	7104:6	7122:5 7124:7 7124:7 7127:8	7301:4	7301:5	7373:2
		1905. April 6	13 May 3 IO	Oct. 31 Nov. 13	13	1906. Feb. 21 Mar. 8
	Date of eruption.		May 2-3	Oct. 28-29 Nov. 13		1906. Feb. 15-17 March 7

* Developed symptoms when in quarantine. Refe

ne. References to notes on cases up to May 10th, 1905, will be found in the Report for the year 1904.

and his wife (7301:5) developed the eruption of small-pox on November 13th, seven days after the successful, thirteen days after the unsuccessful, revaccination. The child did not take the disease, although carried to the hospital when the father and mother were taken there. The child, therefore, was probably the one most exposed to the infection. It is difficult to say what the relative amounts of exposure to small-pox were before the three contacts went to the cottages, but from the date of their eruption it is probable that the father and mother received their infection from the lodger on the 30th or 31st of October. The child's exposure to infection at home would probably be not as great as theirs. But the child was exposed, in addition, to infection from his parents when they were attacked, and still more so when he went with them to hospital. He did not, however, develop the disease, although actually living for three weeks in a small-pox ward. It is not impossible that had the first revaccination on October 31st in the father and mother been successful they might have escaped the disease. At least the evidence of Leicester as to the immunity conferred by successful vaccination after contact points in this direction.

The measures taken to prevent the spread of the disease were as follows:—A list was obtained from the wholesale merchants of the 24 places in Leeds visited by their traveller as far back as October 19th. They were all business premises of small drapers or dressmakers. They were all visited repeatedly by our lady inspectors. No case occurred at any of them. Four persons had visited at the house on the Sunday after the eruption appeared. They were all kept under observation for upwards of a fortnight.

During the rest of the year no further case of small-pox was discovered in Leeds.

It was on the 21st of February, 1906, that we heard of the next case. A woman of 48 (7373:2), in East Hunslet, vaccinated in infancy, had developed an eruption about the 15th to 17th of February. A medical man was attending, but seemed

to be uncertain as to the nature of the disease. The patient was taken to hospital for observation, and her illness turned out to be a slight attack of small-pox. The following day the rest of the family, consisting of the husband, a grown-up daughter, and the husband of the latter, were removed to the cottages. The patient's husband was a driller at an engineer's in Jack Lane, the son-in-law a shopman in one of the Co-operative Stores, and the daughter a tailoress in a ready-made clothing factory. The patient herself had only domestic employment. All three were re-vaccinated on the 22nd of February. The husband, aged 58, had three marks of vaccination in childhood, covering a little more than a quarter of a square inch of surface. The daughter, in addition to being vaccinated in infancy, had been revaccinated in 1903, along with others employed at the wholesale clothing factory. Her second re-vaccination on the 22nd of February, 1906, was also successful. Her husband had not been vaccinated since infancy, but his re-vaccination on the 22nd of February was successful. The man of 58 (7385:5), notwithstanding his successful re-vaccination on the 22nd of February, developed the disease. His eruption appeared on the 7th of March, pointing to the 21st of February as the probable date of infection—that is, the very day before he was re-vaccinated. It is possible, however, that in this case the development of the small-pox eruption may have been delayed by the vaccinia. It seems unlikely that if the man were very susceptible to smallpox he should have delayed contracting the disease from the 17th to the 21st. In this case also the successful re-vaccination did not prevent the attack of small-pox.

In addition to the contacts removed to the cottages, a woman living next door and a man living on the opposite side of the street were kept on our visiting list for upwards of a fortnight. The former was revaccinated. The latter is said to have had small-pox 25 years ago. The places at which other members of the family had worked were also kept under observation and any absentees visited at home.

TABLE 8 a.

Showing cases of Small-pox and deaths amongst vaccinated and unvaccinated personst during the 235 weeks ended 30th June, 1906.

65+	10	I	6	н	1	:	1	:
-100	:	:	:	:	:	:	:	:
90-	:	:	:	:	:	:	:	:
85-	:	:	:	:	:	:	:	:
85		:	:	:	:	:	:	:
75.	-	1	1	:	:	:	:	:
70-	20	1	4	-	:	:	1	:
65.	4	:	4	:	:	- :	:	
25- 65	452	22	404	13	33	5	15	4
-009	18	64	14	1	3	:	1	1
55-	23	4	19	61	ro.	1	1	1
50-	27	4	23	6	. 10	1	1	I
45.	55	3	53	3	:	:	64	:
45	65	0	62	23	1	:	61	1
35-	79	10	99	и	01	60	3	1
30-	104	-	16	I	6	:	4	:
25.	81	:	94	:	+	:	-	:
15.	138	9	911	33	19	33	60	:
20-	.83	9	89	33	12	33	3	1
15.	55	:	48	1	7	1	:	:
5.	89	:	31	:	33	. :	4	:
10-	40	1	23	:	91	:	-	:
10.	28	:	∞	-:	17	1	m	:
1-5	19	33	1	:	19	m	1	:
4.5	6	1	:	1	6	1	1	1
3.4	ω.	1	:	:		-	:	:
2-3	4	:	:	:	4	:	:	:
1.2	6	-	:	:	m	П	:	:
0-1	00		:	1	00	10	:	:
All Ages.	969	35	260	17	112	14	23	4
	Cases -	* Deaths	Cases .	ed. Deaths	Cases .	nated. Deaths	Cases -	l. Deaths
	Total.		Unvaccinated		Unknown or Doubtful.			

* Including death Z. 6052:3 certified as Meningitis (v. pp. 44 and 68, Annual Report, 1903).

+ To save a line in the table two cases are included amongst the vaccinated who had previously had small-pox, but neither of whom had been successfully vaccinated. These cases were 6276:4, xt. 70, and 6282:0, xt. 46. Both recovered, and are mentioned in the text and in table 8, report 1903.

Fifteen patients, entered as unvaccinated in table 8a in the 1903 report, have been transferred to the unknown or doubtful group. They had no marks.

It was difficult to be quite sure of the origin of the first of these cases. The date of the eruption was not very clearly ascertained. It was stated to be the 17th; but, as the initial symptoms seem to have commenced on the 13th, the date might be as early as the 15th. This would point to infection somewhere between the 31st of January and the 3rd of February. On the 31st of January the other members of the family were at their various employments—the patient at home. On the 1st of February she visited her grocer and was employed at home most of the day cleaning. On the 2nd of February she was also at home, and on the 3rd she was again at the grocer's and also at the butcher's. On the 31st, however, her son-in-law and daughter visited a theatre at which a pantomime was being performed. They seem to have met inside with one of the employees at the theatre, a brother of the young man, and to have been able to go about the place pretty freely. company playing at this theatre had come directly to Leeds from Oldham, playing here from January 15th to February 3rd. During the week ended January 13th, 21 cases of small-pox were reported in Oldham, and the following week 22. Medical Officer of Health for Oldham was not aware of any member of the company having been in contact with small-pox, and no case had been reported in any of the houses where they The epidemic in Oldham was "chiefly amongst children, and in a mild form." At present we are inclined to think the disease had been carried by some member of this company and communicated to our patient.

Table 8a is similar to the tables in the reports for 1903 (p. 49) and 1904 (p. 47), but is brought up to the 30th of June, 1906. The table, therefore, deals with a complete period of four and a half years. The numbers after all are not very large, especially when they come to be dissected into age groups. The interesting point in regard to vaccination is that amongst the vaccinated below the age of five there was no case, and that amongst 79 vaccinated persons between the ages of five and

20 who contracted the disease there was no death. Amongst unvaccinated persons under the age of 20 there occurred 67 cases, six of them fatal. The six fatal cases were all under the age of five years, three of them being infants of less than one year. Between the ages of 20 and 40 301 vaccinated persons were attacked, and six of them died. At the same ages 35 unvaccinated persons were attacked, and six died. Over 40, 180 cases occurred, with 11 deaths, amongst the vaccinated portion of the population; whilst amongst the unvaccinated portion 10 cases occurred, with two deaths.

Put briefly, small-pox seems to have avoided altogether vaccinated children; to have attacked, but without any mortality, a small number of those between 5 and 20; to have attacked with greater freedom vaccinated persons between 20 and 40 with a mortality of less than two per cent.; and to have manifested itself with some amount of frequency amongst vaccinated persons over 40 with a mortality more than three times that of the last group.

Amongst the admittedly unvaccinated the mortality under one year of age was very high. The mortality at this age is, of course, high from any disease, but between the end of the first year and the end of the fifth, 19 cases occurred amongst the unvaccinated, and nearly 16 per cent, of them died. As the unvaccinated portion of the community even yet in Leeds at this age is very much smaller than the vaccinated portion (probably less than a fourth), this fact is not without its significance. Between the ages of five and twenty, although there were 40 cases, none of them were fatal. Between twenty and forty, 35 unvaccinated persons contributed six deaths, a mortality of over 17 per cent. Above forty, the cases were only 10 amongst the unvaccinated population. Two of them were fatal, or 20 per cent.

Taken as a whole, about 80 per cent. of the population contributed 17 deaths from small-pox, and about 20 per cent. contributed 14 deaths, but in those under 20 years of age all the

deaths from small-pox were amongst the 20 per cent, unvaccinated. The cases in which evidence as to vaccination or non-vaccination was insufficient are here omitted. They are given, however, in table 8a.

TABLE 8 b.

S. Oliveration	and in a	VACCINATE	iD.	Unvaccinated.			
Ages.	Cases.	Deaths.	Mortality per cent.	Cases.	Deaths.	Mortality per cent.	
0 I	0	0	nil	8	3	37.5	
1— 5	0	0	nil	19	3	15.8	
5-20	79	0	nil	40	0	nil	
20—40	301	6	2.0	35	6	17'1	
40+	180	11	6.1	10	2	20'0	
All ages.	560	17	3.0	112	14	12.2	

MEASLES.

During the year 109 deaths in Leeds were ascribed to measles. One hundred and two houses were examined in which the patient was already dead, six in which the patient died later, 290 in which the patient was alive and in which no patient died and 7 houses (included in the 108 death houses) were visited where a patient had died but where recovery of others of the family attacked by the disease had taken place. Of the 108 "death houses" 16 were "throughs," 92 "not throughs." Of the 290 measles "case-houses" in which no one died 47 were "throughs," 243 "not throughs."

As measles is not reportable, the number of cases heard of in different districts is not a very clear indication of the relative prevalence of the disease. The disease, however, was the most

TABLE 12.

Showing cases of Scarlet Fever heard of in Leeds during the sixteen years, 1890-1905, with the numbers admitted to the city fever hospital and the deaths in the city and in hospital; showing also for biennial periods the relations to the population of the deaths in the city, and the admissions to hospital.

	Cases. †		Deaths.		Per 1,000 living (Annual rates).	
	Heard of.	Admitted to hospital.	City.	Hospital.	Deaths in city.	Cases in
1890	337	133	103	23	0.53	0:10
1891	328	152	66	18	5 0 23	0.39
1892	812	440	74	19)	0-
1893	316	188	31	6	} 0.14	0.83
1894*	967	453	52	18	1	*14.
1895	874	493	52	29	} 0.13	1,51
1896	1,216	441	72	20	0.51	1.56
1897	1,791	576	95	27	5 0 21	1 20
1898	2,002	532	121	25	0.22	1.43
1899	1,620	649	64	2 I	1 0 22	1 43
1900	1,745	722	52	2 I	0.16	2.07
1901	2,280	1,038	82	39	5 0 10	207
1902	1,962	1,041	56	23	0.10	2:27
1903	2,465	1,063	109	36) 019	2.37
1904	1,295	850	59	34) 0.11	1'7'
1905	935	720	39	30	5 011	1'74

^{*} Notification became compulsory in May, 1894.

t "Cases heard of" does not include one fatal in the third quarter of 1894 (included, however, in the 52 deaths), of which we had no information till we received the Registrar's returns. Similarly one death in 1895, another in 1896, two in 1897, one in 1900, one in 1901, and one in 1903, are included in the death column, but not in that of cases heard of. Before May, 1894, there were, of course, many such.

fatal in North Leeds, where the mortality during the year was 0.46. In Hunslet the mortality was 0.39, in Bramley 0.38, and in South-East Leeds 0.37, after which came Holbeck 0.35. Kirkstall and Osmondthorpe had no death from this cause.

The deaths were most numerous in the third and fourth quarters of the year. The death rate in the town was equal to a rate of 0.24.

There was an error in the record of death houses and recovery houses in table 25 in the Report for 1904 by the transference of two of the former to the lines belonging to the latter. The correction is given elsewhere.

SCARLET FEVER.

The scarlet fever cases reported last year fell to 935. It is difficult to know how many cases existed before 1894. In that year, during which notification was compulsory for eight months, 967 cases were heard of; in the following year 874. Then the number rose in 1896 to 1,216. Taking the ten years (1895-1904) during which notification was compulsory, the average number of cases heard of has been 1,725 each year. Our last year's figure (935) was not much more than half that number. The average number taken to hospital during the same period was 741, or nearly 43 per cent. Last year of the 935 heard of 720 were admitted to hospital, or 77 per cent.

I refer you to the table on page 48 showing the number of cases heard of in Leeds during the last sixteen years, the number admitted to hospital, the deaths in the city and in hospital, the city figures including the hospital ones in both cases. The mortality is also given for each biennium.

It will be noticed that we start with a mortality of 0.23 per thousand, which fell to 0.14, and then to 0.13. The mortality then rose to 0.21 and 0.22, and for the last three biennia it has been 0.16, 0.19, and 0.11.

In the last column is given the proportion per thousand of the population of cases treated in hospital. It must be remembered, however, that, during the years of greater prevalence, cases actually treated in hospital could not always be removed at once for lack of accommodation, thus lessening the amount of good work that might have been done by isolation had the hospital at that time been large enough to take promptly all the cases that should have been sent in. It happened again and again, owing to the lack of beds, that we were bound to leave cases at home sufficiently long to allow them to spread the disease, even although many of them were eventually taken to hospital. Isolation to be effectual must be prompt.

The biennial death rates in the table are only given as far back as 1890-1. Before the year 1894 the cases heard of were chiefly (1) cases sent into hospital, or (2) cases heard of where we were asked to disinfect the houses after recovery. The numbers, therefore, before 1895 give no idea as to the number of cases in Leeds. For instance, the death rate in 1890-1 was 0.23 per thousand, but the number of cases heard of in these two years was only 665. In 1904-5 the cases were 2,230, more than three times as many, and the death rate O'II per thousand, only half. In the two earlier years, with a death rate more than double that in the two later, only four cases in every ten thousand of the population were sent into hospital. In the two later years, with a death rate of only 011, seventeen cases per ten-thousand of the population were sent into hospital. Scarlet fever is, as everyone knows, a disease whose prevalence occurs in cycles. It is probable that the commencement of one of these cycles occurred about two years after notification became compulsory, and that the cycle extended, but with diminishing intensity, over about six years. The formidable appearance in the table of the large number of cases heard of is probably not due so largely to the greater prevalence of scarlet fever as it is to the fact that the cases under compulsory notification were heard of in greater numbers.

Unfortunately, as already said, we have no reliable statistics as to the cases earlier than 1894, but I append a table printed at p. 62 in the Report for 1899 showing the deaths and death rates from scarlet fever during three periods of ten years each, which began 1870 and ended 1899. To this table have been added the figures for the six years 1900-1905, which include the later years of the cycle already spoken of. As was perhaps to be expected, considering the increase of the population and the presence of an epidemic wave, the actual number of deaths is not proportionately a very much smaller one than in the last period: 395 in six years corresponding to 435, six-tenths of those in the decade. In the ten years 1890-99 the deaths averaged 72.5 per annum. During the six recent years they had averaged 65.8, but the death rate has fallen in the six years to 0.15 per thousand, and, as already said in the last biennium, to 0'11.

TABLE 12 a.

Showing deaths from Scarlet Fever in Leeds during three periods of ten years each and one period of six years.

Periods.	Deaths.	Deaths per 1,000.*	Fall per cent. on rate of preceding period.
Ten years 1870-79	 3,090	1,00	
Ten years 1880-89	 2,255	0.68	38
Ten years 1890-99	 725	0.10	72
Six years 1900-05	 395	0.12	

^{*} The rate 1 og is probably somewhat lower than the real rate for the period. It is calculated on an over-estimated population, and should probably be 1 11. As, however, the recent populations are mere estimates, the figure has been allowed to stand. The deaths are in each line those given by the Registrar General, which differ slightly in recent years from our own returns. The rates are recalculated from results of last census, as explained at pp. 69-71 of the Annual Report for 1902, where this matter was discussed.

DIPHTHERIA.

From table 18 on pp. 54-55 it will be seen that 364 cases of diphtheria were reported in the course of the year, and that

a third of them occurred in the first quarter. Of the 364, 110 were those of children under five, 175 of persons from five to fifteen, making altogether 285, out of 364, in the fifteen earlier years of life. To these should be added eight cases of membranous croup, six of them in children under five and two in children under fifteen, after which that disease disappears from the record.

From table 17 it will be seen that the mortality from these two diseases during the year was 0.10 per thousand, and from table D, Part I., it will be noticed that diphtheria itself was less fatal than in any year since 1892. This table, however, does not show the combined death rate from diphtheria and membranous croup. The death rate from all forms of croup (not spasmodic) in 1892 was 0.13; last year it was 0.03. If we combine diphtheria with all forms of croup not spasmodic, the rate was lower in 1905 than in any year since I have had the honour of reporting to you.

ERYSIPELAS AND PUERPERAL FEVERS.

During the year 380 cases of erysipelas were reported to us, and nearly a third of the cases again occurred in the first quarter. While 15 occurred amongst children under one, 8 in those between one and five, and 20 between five and fifteen, the numbers increased to 49 in the next ten year period, whilst the great mass of cases occurred in the period from 25 to 65. The numbers are a little larger than in the previous year, as are also the number of cases of the puerperal fevers. The largest proportion of cases of the latter group, however, occurred in the third quarter, not in the first.

CONTINUED FEVERS.

Continued fevers were reported in 293 cases, one of them, that of a boy of 14, having been regarded by the medical man who certified the death as a case of typhus.

TABLE 17.
Causes of, and ages at, death during year (52 weeks) 1905.

Causes of Death.	Di		IN W			ICT A	т	QUAR	RTERL	SEVE Y PERI . AGES	ODS	DEATH PER
CAUSES OF DEATH.	Allages.	Under 1.	r and under 5.	5 and under 15.	15 and under 25.	25 and under 65.	65 and upwards.	I.	11.	III.	IV.	1,000 FOR VEAR.
Small-pox	3	I				1	1	3				0.01
Measles	100	23	74	12				S	8	19	74	0'24
Scarlet fever	39	2	26	11				14	5	13	7	0.00
Whooping-cough	1	51	67	3				21	27	35	38	0.22
Diphtheria and membranous croup		1	29	13		1		20	7	8	9	0.10
Croup	6	2	4					1	2	2	1	0.01
(Typhus				1			***		1			0.00
Fever Enteric	1		1	5	18	28		16	10			0.11
Other continued	1 0	1				1			1		1	0.00
Epidemic influenza	66	5	2		5	31	23	42	11	2	11	0.14
Chalan												
DI			***									***
Diambana	265	303	47	1		7	7	6		1. 33	1000	0.80
Paradile	65	52	7	1	1	I	3	13	12			0.14
D	0				4	5		4	4			0.05
	1	2				2	DOM:	1	I		2	
Out and the same	7	3			1	3		1	1		4	
TM -1.1.1	1 == 8			27	121	371		1 0	3	1 3	100	
0.1 . 1 . 1 . 1	288	80		48		100000		71	85	100000		
			1	40	3	318					1 3 3	100000000000000000000000000000000000000
Cancer, malignant disease Bronchitis	600	156		3			0000000				1000	
	1000	123		12		100			1000		0.000	
Broncho-pneumonia			1000							9.0	10000	
Pneumonia and pleuro-pneu			1	1	20		1		1		700	
Pleurisy	1	***	,		-	0	5	4	5	3	. 5	0 04
Other diseases of respirator organs	72	8	8	4	3	33	16	24	19	13	16	0.16
Alcoholism	1				3							1
Cirrhosis of liver	58				1	49	8	18	9	16	15	0.13
Venereal diseases	. 33	29	I		1	2		. 9	7	10	7	0.02
Premature birth								60	68	76	75	0.61
Diseases and accidents of partu	-											
rition	1 28	10		***	4			9	10000	13	11	0.08
Heart diseases	. 536	2	4	20	26	319	165		1	119	149	1.18
Accidents	. 223	24	32	24	12	90	41	63	61	42	57	0.49
Suicides	. 50				5	39	6	16	1000			
All other causes	2393	673	168	60	75	673	744	640	628	548	577	5.26
	7124			1.7								1

The deaths of outsiders in the workhouses within the city, which are excluded from the Registrar-General's figures, are included in this table. The diseases from which they died will be found enumerated in table C.

TABLE 18.

Cases of infectious diseases notified during the year 1905.

Cases notified in whole district.

		_				_					_		
Notifiable disease.	All ages.	Under 1 year.	1.2	2-3	3-4	4-5	1.5	5-10	10-15	5-15	15-20	20-25	15-25
Small-pox. I.	120	3		2		3	5	9	9	18	7	13	20
II.	6	1											
IV.	3												
Totals	129	4		2		3	5	9	9	18	7	13	20
Cholera. I.	::		17				::				::		**
III. IV.									**				
Totals													
Diphtheria, I.	122	2	8	8	11	14	41	42	13-	55	8	6	14
II.	75 88	3	2 4	5	5 8	5	16 20	26 37	13	39 45	10	4	8 16
IV.	79	2	2	- 8	9	6	25	28	8	36	5	7	12
Totals	364	8	_16_	24	33	29	102	133	42	175	27	23	50
Memb. croup. I.	4 2	::	1	**	1 2	1	3 2	1	**	1	::	::	::
III. IV.	ż				ï		ï	ï		ï	::		
Totals	8		1		4	1	6	2		2		11	
Erysipelas. I.	116	4	2	1	1		4	1	1 4	2	13	4	17
II.	88 88	4	ï				ï	1 2	3	5	4 3	7	7 10
IV.	88	3	1		1	'n	3	4	4	8	4	11	15
Totals	380	15	4	1	2	1	8	8	12	20	24	25	49
Scarlet fever. I. II.	284 195	8	7 4	16 5	25 15	41 19	89 43	104	44 32	148 112	22 16	7 8	29 24
III.	199	2	6	9	20	28	63	74	35	109	7	6	13
IV	257		6		18	24	50	136	30	168	10	7	17
Totals	257 935	12	23	41	18 78	24 112	59 254	136 394	32 143	168 537	10 55	7 28	17 83
-	935	1	_	11	-	24	59 254		32 143	-	_	-	17
Totals	257 935 1	12	23	41	78	112	59 254	394	32 143 1	537 `i	55	28	83
Totals Typhus fever I. II. III. IV.	257 935 1	12	23	41	78	112	254	394	32 143 1	537 `i	55	28	83
Totals Typhus fever I. III. III. IV. Totals	257 935 1 1	12	23	11 41 	78	112	59 254	394	32 143 1 1	537 i 1	55	28	17 83
Totals Typhus fever I. III. IV. Totals Enteric fever I. II.	257 935 1 1 101 56	1 12 	23	11 41	78	24 112	59 254 	394	32 143 1	537 1 1 34 8	55 15 7	28	17 83
Totals Typhus fever I. III. IV. Totals Enteric fever I. III. III.	257 935 1 1 101	1 12	23	11 41 	78	24 112	59 254 4 5 1	394 	32 143 1 1 21 5 13	537 1 1 34 8 20	55	28	17 83
Totals Typhus fever I. III. IV. Totals Enteric fever I. II.	257 935 1 1 101 56 69	1 12	23	11 41 	78	24 112	59 254 4 5	394	32 143 1 1 21 5	537 1 1 34 8	55 	28 9 13 12	83
Totals Typhus fever I. III. IV. Totals Enteric fever I. III. IV. Totals Totals	257 935 1 1 101 56 69 66	1 12 	23	11 41 	78	24 112 1 4	59 254 4 5 1 2 12	394 	32 143 1 1 21 5 13 8 47	537 1 1 34 8 20 11	55 	28 	17 83 24 20 22 17
Totals Typhus fever I. III. IV. Totals Enteric fever I. III. IV. Totals Relapsing I. fever. II.	257 935 1 1 101 56 69 66 292	1 12 	23	11 41 	78	24 112 3 1 4	59 254 4 5 1 2 12	394 	32 143 1 1 21 5 13 8 47	537 1 1 34 8 20 11 73	55 	28 	17 83 24 20 22 17 83
Totals Typhus fever I. II. IV. Totals Enteric fever I. II. III. IV. Totals Relapsing I. fever. II. III. IV.	257 935 1 1 101 56 69 66 292	1 12 	23	11 41 	78	24 112 1 4	59 254 4 5 1 2 12	394 	32 143 1 1 21 5 13 8 47	537 1 1 34 8 20 11	55 	28 	24 22 22 17 83
Totals Typhus fever I. III. IV. Totals Enteric fever I. III. III. IV. Totals Relapsing I. fever. III. III. IV. Totals Relapsing I. Fever. III. III. IV. Totals	257 935 1 1 101 56 69 66 292	1 12	23	11 41 	78	24 112 3 1 4	59 254 4 5 1 2 12	394 	32 143 1 1 21 5 13 8 47	537 1 1 34 8 20 11 73	55 	28 	17 83 24 20 22 17 83
Totals Typhus fever I. III. IV. Totals Enteric fever I. III. IV. Totals Relapsing I. fever. II. III. IV. Totals Relapsing I. fever. II. III. IV. Totals Continued I.	257 935 1 1 101 56 69 66 292 	1 12 	23	11 41 	78	24 112 3 1 4	59 254 4 5 1 1 2 12	394	32 143 1 1 21 5 13 8 47	537 1 1 34 8 20 11 73 	55 15 7 10 8 40 	28 	17 83 24 20 22 27 17 83
Totals Typhus fever I. II. IV. Totals Enteric fever I. II. IV. Totals Relapsing I. fever. II. III. IV. Totals Continued I. fever II. III. IV.	257 935 1 1 101 56 69 66 292 	1 12	23	11 41 	78	24 112 1 4	59 254 4 5 1 2 12 	394 	32 143 1 21 5 13 8 47	537 i 1 34 8 20 11 73 	55 15 7 10 8 40 	28 	17 83 24 20 22 17 83
Totals Typhus fever I. III. IV. Totals Enteric fever I. III. IV. Totals Relapsing I. fever. II. III. IV. Totals Continued I. fever II.	257 935 1 1 101 56 69 66 292 	1 12	23	11 41 	78	24 112 4	59 254 4 5 1 1 2 12 	394 26 	32 143 1 21 5 13 8 47	537 1 1 34 8 20 11 73 	55 15 7 10 8 40 	28 	17 83 24 20 22 217 83
Totals Typhus fever I. III. IV. Totals Enteric fever I. III. IV. Totals Relapsing I. fever. II. III. IV. Totals Continued I. fever II. III. IV. Totals Continued I. fever II. III. IV. Totals Puerperal I.	257 935 1 1 101 56 69 66 292 	1 12 	23	11 41 	78	24 112 3 1 4	59 254 4 5 1 1 2 12 	394	32 143 1 21 5 13 8 47	537 1 1 34 8 20 11 73 	55 15 7 10 8 40 	28 	17 83 24 20 22 17 83
Totals Typhus fever I. III. IV. Totals Enteric fever I. III. IV. Totals Relapsing I. fever. II. III. IV. Totals Continued I. fever II. III. IV. Totals Puerperal I. fever. II. fever. II. fever. II.	257 935 1 1 101 56 69 66 292 		23	11 41 	78	24 112 	59 254	394	32 143 1 21 5 13 8 47 	537 1 34 8 20 11 73 	55 15 7 10 8 40 	28 	17 83 24 20 22 17 83
Totals Typhus fever I. III. IV. Totals Enteric fever I. III. IV. Totals Relapsing I. fever. II. III. IV. Totals Continued I. fever II. IV. Totals Puerperal I. fever. II. IV. Totals	257 935 1 1 101 56 69 66 292 6 4 13 5	1 12 	23	11 41 	78	24 112 3 1 4	59 254	394	32 143 1 21 5 13 8 47	537 1 1 34 8 20 11 73 	55 15 7 10 8 40 	28 	17 83 24 20 22 27 17 83
Totals Typhus fever I. III. IV. Totals Enteric fever I. III. IV. Totals Relapsing I. fever. II. III. IV. Totals Continued I. fever II. III. IV. Totals Puerperal I. fever. II. fever. II. III. IV.	257 935 1 1 101 56 69 66 292 6 4 4 13	1 12	23 	11 41 	78	24 112 	59 254	394	32 143 1 21 5 13 8 47	537 1 34 8 20 11 73 	55 15 7 10 8 40 	28 	17 83 24 20 22 17 83
Totals Typhus fever I. III. IV. Totals Enteric fever I. III. IV. Totals Relapsing I. fever. II. III. IV. Totals Continued I. fever II. III. IV. Totals Puerperal I. fever. II. III. IV. Totals Plague. I. Plague. I.	257 935 1 1 101 56 69 66 292 6 4 13 5 28	1 12	23	11 41 	78	24 112 	59 254	394	32 143 1 21 5 13 8 47 	537 1 1 34 8 20 11 73 	55 15 7 10 8 40 	28 9 13 12 9 43 	17 83
Totals Typhus fever I. III. IV. Totals Enteric fever I. III. IV. Totals Relapsing I. fever. II. III. IV. Totals Continued I. fever II. III. IV. Totals Puerperal I. fever. II. III. IV. Totals Puerperal I. fever. II. III. IV. Totals Plague. I. III. III. IV.	257 935 1 1 101 56 69 66 292 6 4 13 5 28	1 12	23	11 41 	78	24 112 3 1 4 	59 254	394	32 143 1 21 5 13 8 47 	537 1 34 8 20 11 73 	55 15 7 10 8 40 	28 	17 83
Totals Typhus fever I. III. IV. Totals Enteric fever I. III. IV. Totals Relapsing I. fever. II. III. IV. Totals Continued I. fever II. III. IV. Totals Puerperal I. fever. II. III. IV. Totals Puerperal I. fever. II. III. IV. Totals Puerperal I. fever. II. III. IV. Totals Plague. I. III. IV. Totals	257 935 1 101 56 69 66 292 6 4 13 5 28		23	11 41 	78	24 112 	59 254	394	32 143 1 21 5 13 8 47 	537 1 34 8 20 11 73 	55 15 7 10 8 40 	28 9 13 12 9 43 	17 83

TABLE 18-continued.

Cases of infectious diseases notified during the year 1905. Cases notified in whole district.

Notifiable disease.	25-30	30.35	35.40	40-45	45-50	50-55	55-60	60-65	25-65	65-70	70-75	75-80	80-85	85-90	90-95	95.100	65 up- wds.
Small-pox. I.	10	9	17	18	5	4	7	1	71	1	2						3
II.	1	3	::	1					5				**				
Totals	13	13	17	19	5	4	7	1	79								
Cholera, I.					-	_		-		-				-	-		
II.															1		
IV.																	
Totals Diphtheria, L.	2	3	1	- 3	-	1	-:-		10					-:-	_		
II.	2 4 1	3 3 2	1	ĭ	i	1			11 4		::						
IV.	2	2 2						-:-	4								
Totals	9	10	2	4	1	3		_	29			=					_
Memb. Croup. I.													::				
III.												-:-					**
Totals																	
Erysipelas. I. II.	12 5 5	5	15	6	13	10 10	7 7	9 6	70 60	7 6	7 2 3	1	2 3	2	**	::	19 12
III. IV.	5	10	6	9	5 3	8 7	7 5	4	54 51	11 4	3 2	ż			**		14 8
Totals	33	24	39	32	27	35	26	19	235	28	14	4.	5	2			53
Scarlet fever. I.	6 8	3 5	1 2		**	-:-			10 15	::							
III. IV.	6	4	3	2		ï			12 12								
Totals	26	13	6	3		1			49								1.0
Typhus fever. I.						*					215						
III. IV.																	
Totals						-:-											
Enteric fever I.	12	10	6	6	1	-;		3	38	1							1
II. III.	6 5 10	9 7 12	6 2 4 8	6 5	1	1	i		23 25 36		**	1					i
IV.	33	38	20	21	4	2	1	3	122	1		1					2
Relapsing I.																	
fever. II. III.		**											::				**
Totals																	
Continued I.	-			-:-		-			-	-		-		-			==
fever. II.		* **	::		1.0		**										
Totals					-												
Puerperal I.	2	1	1	1			-	-	5				-			-	
fever. II.		i	2	2					9								**
IV.	1			1	-:-	-:-	-:-		2		::	-:-	_::	-:-			
Totals	7	2	3	4					16								
Plague. I.								**	**								**
III.			::														
Totals																	

TABLE 19.

Table shewing deaths in the four quarters of 1905, for each Municipal Ward, with the estimated population and the death-rate of the ward for the Year.

MUNICIPAL WARDS.	Population, estimated to middle of 1905.	First quarter, 1905.	Second quarter, 1905.	Third quarter, 1905.	Fourth quarter, 1905.	Fifty- two weeks.	Death- rate.
Central	20,216	- 88	76	73	82	319	15.83
North	44,096	149	125	170	129	573	13.04
North-East	31,280	166	125	142	158	591	18.96
East	29,554	154	153	178	153	638	21.66
South	14,161	69	55	83	73	280	19.84
East Hunslet	37,015	113	136	129	156	534	14.48
West Hunslet	31,712	114	113	124	107	458	14:49
Holbeck	30,664	112	137	130	131	510	16.69
Mill Hill	7,136	33	17	28	27	105	14:76
West	23,683	114	113	92	118	437	18.51
North-West	34,015	130	118	100	114	462	13.63
Brunswick	23,034	85	67	83	81	316	13.77
New Wortley	18,516	73	75	75	90	313	16.96
Armley & Wortley	40,561	153	130	161	136	580	14'35
Bramley	23,127	74	80	64	78	296	12.84
Headingley	48,017	161	146	108	119	534	11.16
Outsiders		31	54	50	43	178	
Totals	456,787	1,819	1,720	1,790	1,795	7,124	15.65

The rates are calculated on populations estimated by the Registrar General's method, described at p. 88 of the Annual for 1904. Those for several quarters, given in tables E 2, E 3, E 4, and E 5, are on populations as previously estimated.

OTHER DISEASES.

In Table 17 will be found tabulated certain causes of death, with the age groups at which they occurred. In the same table are also given the number of deaths in each quarter at all ages, and the death rate per thousand per annum for the year.

ILLNESSES NOTIFIED.

The two-paged table dealing with infectious diseases notified has been numbered 18 this year. This number has not been made use of since 1899. It was the number attached to a table in earlier reports, the complement of table 17, and contained the death rates in the year and its several quarters from the diseases enumerated in that table. In 1900 table 17 was altered. The death rates for the whole town were given from a somewhat enlarged group of diseases, along with the deaths at certain age groups and in each quarter from the same causes. The death rate from each of these causes was given for the whole year. Rates from a still more enlarged group of causes are given in table C for the whole year, and from the old smaller group in tables A, parts 3, 4, 5, and 6 for each quarter of the year. In 1900, to avoid any confusion, the new table dealing with infectious diseases was introduced without a number.

Tables of cases notified and deaths registered from the notifiable diseases had been given previously in most of the reports since 1894. In 1900 the more extended form given in table 18 was adopted. In this table the cases notified from each disease are enumerated for all ages and for each quarter of the year. They are enumerated also for the year and its several quarters at certain groups of ages, under one year separately, from one to five years, and for each five years from five upwards. They are also gathered into the groups used in tables 3 and 4 and in table A, part 2. It was the intention, in collecting the information in these minute details, to make it comparable with the returns in our death ledgers.

PART III.-HEALTH OF DISTRICTS.

MUNICIPAL WARDS.

A comparison of mortalities in the different parts of the town is difficult on account of the changes in the population that have taken place since the census. The populations, however, have been estimated by the Registrar General's method for the intercepts of wards and townships, and these populations have been added for the estimated populations for the different wards given in Table 19.

Assuming the comparative accuracy of these figures, it will be noticed that the death rate in the Central Ward, after allocating institution deaths, is given as 15.83; whilst that of the whole town is 15.65. When the 178 deaths of outsiders are deducted, the death rate of the whole town falls to 15:26, so that the death rate of the Central Ward is appreciably higher than that of the whole town when treated in the same manner; that is, when the institution deaths not belonging to Leeds are deducted from both and when those belonging to the Central Ward but occurring in hospitals outside its area are added. In the North Ward the death rate, on the other hand, was 13'04, more than 2 per 1,000 below that of the whole town. The actual difference, 2.22 deaths per thousand, is one of 141 per cent. The North-East, East, and South Wards had much higher rates, 18.96, 21.66, and 19.84. East Hunslet had a rate lower than that of the whole town, 14'48, against 15'26. The rate in West Hunslet was practically the same as in East. Holbeck had a death rate nine per cent, higher than that of the town. Mill Hill, a small district not altogether without slums, had a death rate slightly below that of the town. The West ward had a death rate of 18.51, more than 21 per cent. above that of the town. North-West and Brunswick wards had rates each about 11 deaths per thousand of the population below that of the town, the two rates being respectively 10.7 and 9.7 per cent. lower than the city rate. The rate in New Wortley was about as much above that of the town (111 per cent.). Armley and Wortley had a rate slightly below, Bramley and Headingley both considerably below that of the town—the latter to the extent of 26.8 per cent.

(2) REGISTRATION SUB-DISTRICTS.

The populations given in Table A, Part II., are estimated in a similar manner for the townships and sub-districts, and in Table 21 will be found the rates calculated upon them for these sub-districts from each member of the seven zymotic groups, from croup, phthisis, lung diseases, and from other and all causes. In this case also all institution deaths of persons not belonging to Leeds have been eliminated, and institution deaths of those belonging to the several districts have been allocated to them.

It will be noticed that the North registration district had the highest rate but one from the seven zymotic diseasesmeasles, whooping-cough, and diarrhoa contributed chiefly to this high rate. West Leeds had a smaller death rate from the same group than the North registration district. South-East Leeds, on the other hand, had a rate nearly double that of the whole town. Not only did measles and diarrhœa contribute considerably to this high rate, but diphtheria and fevers had both rates beyond those in the town. In Hunslet, a very large township, the zymotic rate was 1.96, against 1.60 in the town, measles and whooping-cough again contributing largely to this result—the diarrhœa rate being 50 per cent above that of the town, whilst the "fever," scarlet fever and diphtheria rates were below. Holbeck had a zymotic rate slightly above that of the town. Measles and whooping-cough were again chief causes -the rate from diarrhœa being slightly less than in the town generally. Wortley had a rate below that of the town from every one of the seven diseases except scarlet fever and whooping-cough. That from the former was the same as that from the latter, one-third higher than that of the city. Kirkstall—that is the parish of Kirkstall which, with the exception of the small bit of Chapeltown intercepted by the Meanwood Beck corresponds to the Headingley Ward—had a zymotic rate less

TABLE 20.

Shewing the number of deaths from certain specific causes and groups causes in the 52 weeks of 1905 in the Sub-Registration Districts in the Ci of Leeds. All deaths in public institutions within the City of person belonging to the City have been referred to the Sub-District to which the belonged.

	Small-pox.	Measles.	Scarlatina.	Diphtheria.	Whooping-cough	"Fever."	Diarrhœa.	All seven.	Croup.	Pluthisis.	Influence and diseases of the air-passages other than consumption.	All other causes,	All causes
Leeds, North	2	27	4	5	16	12	78	144	3	98	221	574	1040
,, West		15	7	8	11	10	52	103	1	118	216	852	1290
" South-E.	I	13	2	7	8	10	68	109	3	- 50	189	420	77
Hunslet		29	4	3	32	7	71	146	I	67	245	621	1086
Holbeck		12	5	3	13	3	26	62	1	52	129	350	594
Wortley		5	6	2	23	5	45	86	3	60	167	627	94.
Kirkstall			3	4	11	5	10	33		40	83	355	5111
Bramley		7	6		3	: 3	3	22		21	42	161	246
Chapeltown		I	2	6	4		12	25		46	89	305	465
Osmondthorpe										2	2	2	6
City of Leeds	3	109	39	38	121	55	365	730	12	554	1383	4267	6946

Four deaths from phthisis, 13 from influenza and diseases of the air-passages other than consumption, and 1 deaths from other causes, occurred in the City of persons not belonging to Leeds; on the other had 101 deaths occurred during the year of Leeds persons in West Riding Asylums and other publinstitutions, outside the City. Of these 17 were from phthisis, 5 from the lung groups, and 79 from other causes.

Croup includes 6 deaths from membranous croup, 3 in North, 1 in West, 1 in Hunslet, and 1 in Wortley.

than half that of the town. The diphtheria rate was slightly in excess (12½ per cent.), the diarrhœa rate a little more than a quarter that of the town. Owing chiefly to measles and scarlet fever, the Bramley rate, although lower than that of the town as a whole, did not stand nearly so well as the Kirkstall rate. Chapeltown, which consists of the parishes of Potternewton and Chapel-Allerton—the latter including Meanwood, the Headingley intercept already spoken of—had a rate nearly 61 per cent-

TABLE 21.

Shewing the mortality stated in deaths per 1,000 of the population of the Sub-Districts, as estimated to the middle of 1905.

	Small-pox.	Measles	Scarlatina,	Diphtheria	Whooping-cough.	"Fever."	Diarrhœa.	All seven.	Croup.	Phthisis.	Influenza and diseases of the air-passages, other than consumption	All other causes.	All causes.
Leeds, North	0.03	0.46	0.04	0.09	0.52	0.50	1.33	2.45	0.02	1.67	3.76	9.77	17.71
,, West		0.18	0.08	0.00	0.13	0'12	0.01	1.55	0.01	1.39	2.22	10.09	15.53
,, S. E.	0.03	0.32	0.06	0.50	0.53	0.50	1.06	3.14	0.09	1'44	5.45	12.10	22.55
Hunslet		0.39	0.02	0.04	0.43	0.00	0.00	1.06	0.01	0.00	3.30	8.36	14.23
Holbeck		0.32	0.12	0.09	0.38	0.00	0.76	1.80	0.03	1.21	3.75	10.18	17.28
Wortley		0.08	0'09	0.03	0.36	0.08	0.21	1.35	0.02	0.94	2.63	9.87	14.84
Kirkstall			0.09	0.00	0.24	0.11	0.51	0.21		0.86	1.78	7.63	10.98
Bramley		0.38	0.33		0.19	0.19	0.19	1.50		1.14	2.29	8.76	13.39
Chapeitown		0.03	0.02	0.12	0.10		0.30	0.63		1.19	2.22	7.70	11.75
Osmondthrp	***									5.61	5.61	5.61	16.82
City of Leeds	0.01	0.24	0.00	0.08	0.52	0.13	0.80	1.60	0.03	1'22	3.04	9.37	15.26

our deaths from phthisis, 13 from influenza and diseases of the air-passages other than consumption, and 161 deaths from other causes, occurred in the City of persons not belonging to Leeds.

eaths from membranous croup are included under croup, not under diphtheria in this table.

below that of the town. The diphtheria rate was, however, nearly double that of the town. That of the other six diseases was much below that of the city as a whole.

It will be noticed on looking at the last column of this table that the total death rates in Kirkstall and Chapeltown were very much below the rate of the whole town, 41 deaths per thousand in the one case, 31 in the other. Then came Bramley, nearly 2 deaths per thousand below that of the town, followed by Hunslet and Wortley approximately 2 deaths per three thousand and I death per two thousand respectively below that of the town. Then came West Leeds, with its rate nearly the same as that of the town, after which in order, but above the rate of the town, Osmondthorpe, Holbeck, North Leeds, and South-East Leeds, the latter with a rate of 22:22, or nearly 7 deaths per thousand more than in the town generally. The 7 deaths are accounted for, 11 from the zymotic group as already shown, nearly 21 from bronchitis and other lung diseases, nearly 1 from consumption alone, and 23 from other diseases not set out in the table. The rate from the seven zymotic diseases* was much higher than in any other district of the town. That from phthisis (1:44) was higher than in any other district in the town but North Leeds (1.67) and Holbeck (1.51), if we except for reasons to be presently given the small district of Osmond-The rate from other lung diseases (5:45) was higher than in any district except the small district just mentioned. The rate from all other causes (12:10) was higher than in any district whatever.

Osmondthorpe requires a word to itself. The population is so small that a single death in any group raises the rate of that group considerably. The total mortality, giving a rate of 16.82 for the whole district, corresponds to six deaths.

^{*} These are not the same seven diseases spoken of earlier in connection with infant mortality. The latter exclude diarrhoea, which was one of the most fatal diseases amongst infants in South-East Leeds.

TABLE 22.

Table showing deaths from all causes at all ages, and death rates at all ages in the intercepts of the wards and townships of Leeds for the year 1905.

		Deaths, 1905.	Average death rate per thousand.
Ногвеск	Holbeck West Hunslet	510	16.69
HUNSLET	West Hunslet East Hunslet South	375 534 172	13.45 14.51 17.84
SOUTH-EAST LEEDS	South } 24.81 Central }	108 31 632	24.15 27.44 21.72
OSMONDTHORFE	East) 19.84	6	16.81
CHAPELTOWN (part of)	North-east North C. A. Pott.	138 63 204	15.72 12.26 9.59
NORTH LEEDS	North North-east Central Central in W. } 15.14	306 453 281 7	17.45 20.23 14.95 31.76
WORTLEY	New Wortley Armley } 14.35 Farnley	313 426 154 50	16.96 13.67 16.62 10.69
Bramley	Bramley	246	13:39
KIRKSTALL	Kirkstall Burley Headingley	65 306 140	14.88 10.67 10.39
Chapeltown (part of)	Headingley North-west Brunswick	23 3 34	17:33 37:66 11:36
West	Brunswick North-west Mill Hill West	282 459 105 437	14.13 13.57 14.76 18.52
Сіту		6,946	15.26
	Outsiders	178	

The rates are calculated on populations estimated by the Registrar General's method, described at p. 88 of the Annual for 1904.

Returns for 1903 and 1904 are given in that report, p. 89.

In Annual for 1891, tables 22, 23, and 24 give deaths and rates in sections of insanitary area No. 1, i.e., the triangular area in the Norh-East ward. The numbers have not been used for any tables since; but in 1894 two tables dealing with Camp Field and West ward, and in 1905 two tables dealing with Camp Field and the York Street area, were inserted unnumbered between tables 21 and 25.

Shewing case-houses examined on account of certain diseases heard of during 1905, and some of the conditions found as to drainage and closet arrangements.

	1	2	3	4	5	6	7	8	9	10	
		-	-	D	rains	severe	d.				
			Water	-close	t.						52 weeks, 1905.
	F	Ins	ide.	F.V.	Out	side.	Т.	w.c.	M.	or P.	
	def.	not	def.	not	def.	not	def.	not	def.	not.	
Through Back-to-back		14				11 25	0.000	1 20		3 5	Smallpox
Through Back-to-back	3	126	 I	5 4	2	127		16 70		18	Scarlet fever
Through Back-to-back	13	30 15	 I	3	13	43 64		4 45		3	Diphtheria
Through Back-to-back		***				1 2		3			Membranous croup
Through Back-to-back			7			4					"Croup"
Through Back-to-back											Typhus fever
Through Back-to-back	7	16 9	2	I	3 16	29 89	1 4	5 28	 I	4 3	Typhoid fever
Through Back-to-back	***		***				***				Continued fever
Through Back-to-back		29 15		1 2		38 108		50		7 4	Erysipelas
Through Back-to-back		I			2 I	1 6		4		 I	Puerperal fever
Through Back-to-back		3 2				6 34		2 27			Measles: death-houses
Through Back-to-back		6				25 104		11 68			Measles: recovery houses
Through Back-to-back						4		I I			Measles: recoveries in death- houses
Through Back-to-back		13 9		 I		34 124		13 78	P.:	2 2	Diarrhœa
Through Back-to-back		23 22		2 I		58 85		5 40		8 7	Phthisis
Through Back-to-back		14		3 2		34 145		12 100			Phthisis: recovery houses
Through Back-to-back		15 12				24 93		4 45		6	Broncho-pneumonia
Through Back-to-back		18		2.		30 113		4 51		3	Pneumonia
Through Back-to-back		4 2				3		2			Pleuro-pneumonia
Through Back-to-back		I				3		2			Pleurisy
Through Back-to-back		 I									Laryngitis
Through Back-to-back		7 6		 I		7 8		5		:	Influenza
Through Back-to-back Both		316 168 484	2 2 4	15 15 30		470 1288 1758		86 639 725	 I I	51 72 123	

		1	2	3	4	5	6	7	8	9	10	11			
					1	Drains	not se	vered.			-		Cas	es.	Total
			V	ater-	loset.					-					deaths
		F.V	Insid	le. not F	.v.	Outs	ide.	T.W	.c.	М. о	r P.	No drain	Alive.	Dead.	City.
	1	def.	not	def.	not	def.	not	def.	not	def.	not	- 1			
Through Back-to-back							8		1 26				31 84		3
Through Back-to-back			7 23		 I	1 2	21 80		8 36		7 4		340 572		39
Through . Back-to-back .	-1	4		3		4 12	18	 I I	I 22	 I	2 I	1	120 236	3	38
Dools to book							2						1 6	 1	6
Deals to beal.							2	 I	2				6	3	6
Doals to book							 I							 I	I
Deals to beals						6	10	7	5				91	I	52
Deals to beals															2
Deals to beals			I 2				15 31		1 28		3		97 242	I	5
Doub to book		1				 I	3		3				6 .	I 2	9
Dool to book			I			I	16		2 10		I			16 92	109
Deale to beale			S				6 28		2 28			 I	47 243		
Deals to book							 I						6		
Back-to-back							30		5 45		4			70 295	365
Back-to-back			1 5				5 33		1 28			2		105 222	558
Back-to-back			2				12 56		3 47		1 2	 I	82 372		***
T) - 1 - 1 - 1			2				2 39		4 32	I	I	 I		53 231	287
Dools to book			2			1	30		3 25		5	1		64 239	340
Back-to-back														5 7	12
Back-to-back							1 4							4 10	17
Back-to-back										1				1 4	8
71 1 1			I I				5		5	1	I			21 33	66
Back-to-back		4	17 47 64	3 3	1 2 3	13 29 42	86 398 484	1 21 22	37 348 385	3 1 4	14 20 34	5 5 10	816 1977 2793	345 1145 1490	1923

It will be noticed that the populations already alluded to as given in table A are given differently in different parts of that table. In the quarterly tables A3, 4, 5, and 6, and in the lower line in Part I. the estimates are given as originally calculated by geometrical progression, the numbers being toned down proportionately to correspond with the Registrar-General's total for the whole town. In the upper line in Part I. and in the figures given in Part II. the numbers are those obtained by adopting the Registrar-General's method for the large towns, and applying it to the several districts. The differences in some cases are considerable, and it is difficult to say which method gives the least inaccurate result.

(3) INTERCEPTS.

The populations for the intercepts have been calculated by the Registrar-General's method for the large towns. Table 22 gives the deaths in each intercept of wards and districts, with the rate upon the thus estimated population. I went so fully into the difficulties of the estimates of population in small districts in last year's report (pp. 90-95), that I need not any further allude to it here. In that report the deaths in each intercept for 1903 and 1904 were given and the assumed death rates (Table 22, p. 89).

Table 22 thus gives the deaths and rates in these intercepts for a third year. The case rates for notifiable diseases were given for the five years 1900-4 in last year's report (p. 99). Those for a single year would have so little value that they are not given.

(4) OTHER LOCAL FACTORS.

I hoped to have had time this year to have discussed table 25, with its predecessors under the same number. On the same lines as this table a record has been kept of the University lodging-houses examined by Mr. Carter, and in table XVIII., and in the text preceding it in part IV., will be found a discussion of the results found.

PART IV.-ADMINISTRATIVE WORK.

CHANGES IN STAFF.

Scavenging superintendent.—To the great regret of the whole department, Mr. George Darley, who had been scavenging superintendent, so far as the removal of house refuse was concerned, since 1890, and, so far as both street cleansing and ashpit cleansing were concerned, since 1897, died after a short illness on the 19th of June, 1905. Mr. Darley was originally an engineer, and had made himself thoroughly acquainted with the working of both stationary and locomotive engines. He worked for the Corporation on the destructors when they were novelties, and for several years made an efficient smoke inspector. When in 1890 the Corporation came to the conclusion that it would be better to take the cleansing of ashpits into their own hands, Mr. Darley was appointed to superintend this work, and he at once made the thing a success. On the death of Mr. Hanford, who had for many years superintended the street cleansing and watering department, it was considered advisable to combine these two formerly distinct sections of scavenging work under one manager. Mr. Darley thus undertook the superintendence of the whole of the cleansing. He was ably assisted by Mr. Mann, who became his deputy, and the Council never had occasion to regret the changes made by taking the whole of the scavenging into their own hands. Mr. Darley had a good deal of originality, and though during the last year or two he had suffered from time to time in his health, he had succeeded in training a good staff of assistants to carry out the work.

Mr. Darley was one of the earliest amongst our staff to take the inspector's diploma of the Sanitary Institute, of which Institute he became an Associate in 1888. He always took an interest in the work of the Inspectors' Association, and was President in 1899-1900 of the Association of Cleansing Superintendents. On the death of Mr. Darley Mr. Mann was appointed temporarily for twelve months to carry on the work, which has in no way suffered at his hands.

Divisionals.—Mr. John Wm. Mills, the senior amongst our divisional inspectors, died on the 17th of May, 1905. Mr. Mills had been on our staff from 1875, when he was appointed foreman joiner and works inspector in the scavenging department. Between 1879 and 1880 he was inspector over the scavenging done by the contractor, and for a portion of that time superintended the street sweeping directly. In the last named year he became assistant nuisance inspector, having charge of the East and Kirkgate wards. On my coming to Leeds I found him inspector of the Mill Hill district, a position he had occupied for a few months. He had certificates of the Science and Art department in building construction, geometry, and freehand drawing, and of carpentry and joinery, brickwork and masonry, from the City and Guilds. In 1891 he was chosen as one of two divisional inspectors on the institution of that office, and was told off to what we then called the western division of the town. The following year he took the diploma of the Sanitary Institute as a nuisance inspector. When in 1902 the number of divisional inspectors was increased to four Mr. Mills retained the northern part of the western division, including the Headingley with Burley ward, North-west ward and the Brunswick ward, to which was added in lieu of the West ward the part of the Central ward north of Lady Lane, in order that the Jewish population might be as far as possible under the oversight of the same divisional inspector.

On January 11th of the present year Mr. Carter, our canal boats and temporary dwellings inspector, was appointed to the division, retaining in addition the superintendence and general control of his old work. On this condition his salary was made fifty shillings a week, the sum which Mr. Mills had been receiving for some years at the time of his death.

Assistant inspectors.—No change took place amongst the works inspectors.

On March 25th, as mentioned in my report for 1904, Mr. Charles Firth left us to take up his duties as inspector of nuisances to the Borough of Ossett. Mr. Firth was one of what we might call the new school of inspectors. Born in 1880, he received a Board School education in the neighbouring village of Drighlington, attending classes at the Higher Grade, and later at the Grammar Schools in Bradford. He became a registered plumber by examination, taking a firstclass certificate and the silver medal of the City and Guilds, receiving also the diploma in plumbing of the Bradford Technical College. He held certificates in mathematics and chemistry, geometrical, freehand, and model drawing, and took the nuisance inspectors' diploma of the Sanitary Institute in 1902. In 1903 he applied for and was appointed a probationer, and was so useful, that although no vacancy occurred at the end of six months, he was allowed to continue on probation for about a year, and in 1904 I had great pleasure in recommending him for a vacancy amongst our assistant inspectors. He was appointed officially to the care of the South ward, in which he had been working as a probationer, and later in the same year he took charge of East Hunslet on the resignation of Mr. Spetch.

On the 4th May, Mr. Henry McCaragher died. Mr. McCaragher, as mentioned in previous reports, had been ailing for a considerable length of time. His work in the South ward had latterly been done by Mr. Firth as just mentioned. Originally a pit sinker for coal, he was for some nine years night foreman and superintendent of the re-laying of old drains. Then for a time he was an inspector of drainage under the Streets and Sewerage Committee. He was one of those selected by the Sanitary Committee in 1891, from whom to chose the two divisional inspectors at the time that office

was a new one. His health was, however, for several years very indifferent, and he was practically blind some months.

On July 15th Mr. Ellis, inspector for the time being in Mill Hill and a portion of the Central ward, died. Mr. Ellis was one of the men who had risen from the ranks. An Irishman by birth, he came upon the staff of the City Engineer in connection with drainage work somewhere in the eighties. In 1890 he was made night foreman on Mr. Darley's staff of ashpit cleansers. In 1894 he was appointed an assistant nuisance inspector to the East ward on condition that he obtained before his second advance the diploma of the Sanitary Institute. He got the ordinary advance to 33s. in 1895, took the diploma of the Sanitary Institute in 1896, getting his second advance the following year. He remained in the East ward until April, 1902, when it was considered advisable on account of his health to give him rather lighter work. Mr. Ellis was one of our inspectors who had a good deal of personal character. Amongst his other good qualities he had a great love for old books.

At the close of the year four men were on probation— Hutchinson, Pollard, Saxton and Broadbelt. Since the beginning of the present year they have all been made assistant inspectors.

Other inspectors.—Mr. Petty, as mentioned in my last report, our senior meat inspector, died in 1904. Temporary arrangements were made for carrying on his work by Mr. Marsden, our junior meat inspector, and Mr. Bentley, one of our ward inspectors, trained as a butcher, who had taken the Food Inspection diploma of the Sanitary Institute. In June, 1905, the Committee appointed Mr. James A. Dixon consulting meat inspector as well as to certain duties connected with the inspection of cowsheds and cattle. Mr. Dixon is a veterinary surgeon of experience, having filled for some time a similar office as meat inspector and inspector of cows in the borough

of St. Helens. His appointment is that of "Veterinary Assistant to the Medical Officer of Health." In addition to acting as referee in all difficult cases of diseased meat and diseased animals brought into town for food, he has thrown himself very thoroughly into the work of examining the cows in the various cowsheds within the borough for tuberculosis. As the Committee are aware this work had been done by a veterinary surgeon practising in the town with great advantage.

Table VIIIa. in my reports for the last few years has contained a list of the number of cows examined in this way for disease of the udder. There was of course the difficulty that is apt to occur where a man in practice acts as a sort of referee in regard to the condition of cattle in the sheds of the various farms, and though there was every reason to believe that Mr. Bowman was thoroughly conscientious in carrying out the work entrusted to him by the Corporation, he was himself the first to confess that it would be an advantage to the department to have a veterinary surgeon not in practice to carry out this work. There is another advantage about the new arrangement. Our own veterinary surgeon is able to adapt his time of visiting to the circumstances of the case more easily than a gentleman in private practice, especially in the summer time when the cattle are out at grass. Mr. Bowman's habit was to devote a whole afternoon to the examination of cows in neighbouring farms, but to discontinue these examinations during the summer months. Mr. Dixon has been able to continue even in greater numbers the examination of cattle at farms during the winter, and during the summer has been able to arrange his work in such a way as to examine cattle in certain cowsheds at the milking time.

Women inspectors.—Miss Turner and Miss Hunter left us in the latter part of 1904 (Annual, p. 107).

On the 20th March, 1905, Miss Agnes Helena Bhose, who had been a probationer since August of the previous

year, was made a sub-inspector. At the time of her becoming a probationer Miss Bhose held the nuisance inspector's certificate of the Royal Sanitary Institute. She had received nursing training or acted as a nurse at the Royal Southern Hospital, Liverpool, at the Royal Devon and Exeter Hospital, and at the Isolation Hospital, Dartford, Kent. Miss Bhose had the certificates of the College of Preceptors, second and third class, and had generally a good secondary education.

On the same day (March 20th) Miss Enid Orange was made a sub-inspector. Miss Orange was selected as a probationer at the same time as Miss Bhose, but came to us a few weeks later. She had been educated at the Kensington High School, the Guildhall School of Music, at University College and Bedford College, London. She had a first class certificate from the College of Preceptors, had passed the London Matriculation and Preliminary Scientific examination, the Royal Sanitary Institute inspector's examination in 1903, and that of the Sanitary Inspector's Board in 1904. She had had some experience in rent collecting in London with Miss Mary Lumsden.

In March, also, Miss Florence Isabel Lansdown came to us as a probationer. She held the higher certificate of the Oxford and Cambridge Joint Board, the National Health Society's certificate in hygiene, and the certificate of the Sanitary Inspector's Examination Board. She had also worked in the sanitary department of the City of London for six months, and was advised to come to us for further training by Dr. Collingridge. She was made a sub-inspector in December.

Miss C. Frank-Smith came to us as a probationer in August, 1905. She held both senior and junior Cambridge Local certificates, and those of the II. and III. class of the College of Preceptors. She had taken in January, 1903, the certificate of the Sanitary Inspector's Board, and in April, 1905, the diploma of the National Health Society.

Table of Ward Inspectors' Work. Year 1905.

		ı			1000						-	
Quarters.	rters.	rters.		>	Total 1305.			-	Quarters.	il. I	IV.	Total 1905.
					-			1		-	1	T
NSPECTION.	-		-	1			nued).	0.0	good	000	-	0
Houses and premises Infective disease	940 1025	250	250 15	3 10	811	33.	diams by tests	0000	865	042 1	1241 208	2082
House to bound more than 1000 1000	2000 1500	1590	2000		r			000		2000		1100
(House-to-nouse work 1233 1233 12	5753 1373	1373	37.5 34	+0		200	are control	200				000
mises Occupants 53 93 91	03	16		5			Touries allu letters serveu	-	:	:	5	67
Buildings and Offices 202 409 337	409 337	337		230			Dwelling houses until for human habitation closed	:	:			
305 332 300	332 300	390	300	300			Dwelling houses rendered in 10r numan natitation					: .
Number of houses wholly or partly examined 3118 3199 3050 5583	3199 3050	3050	1050 55	10	3 15550		Houses cleansed	144	169x	139	153	000
Total number of above houses where sanitary defects						42.	Overcrowded houses dealt with	62	16	94	101	340
1229 1102	1102		285 16	0		43.	Defective spouting, &c., repaired	156	169	245	243	SII
found in above houses 1654 (480)	1760 1760	1760	760 24	3	7 305							:
							ed					
, or 33,	0			-		i v						:
	170 155	133		00	245		DO. PEDMIN		:			:
011 85 98 110	98 110	110		6			converted into		:		:	:
damp or dilapidated r32	117 143	143		~	003	48	Do. do. ordinary water closets	350	351	280	1001	188
with defective eave-gutters or			-			40.	closets erected	14			96	TOL
000 100 000	000 700	999		- 2			also	000	-	Sec		1001
tan pipes	434	200		71.5	000		a bandle	433			260	5
s ,, hadly drained 751 054 042	054 022	0.62		6				:	:	:	:	:
found in ,, without sink drain 12 7	7		:	*	4 23		Fall closets converted into water closets		:	:	I	-
: :			-	~	1.1	33	Trough closets altered into water closets	*	11		10	100
badly ventilated 5	IO	_	IOI	24	36	54.		245	230		328	-002
						100	Drains in course of construction inspected	266		278		1017
closet accommodation 842 1172 810	1172 810	810		+	2577			24.7				Tro
ore rigo	010	200					increased when connection made to cause	200			1	
+/2 642 402	t/z 64z	+/=		÷.	1133	57.	or to sewer	5			7.5	543
				4		_	iouse drains effected	1001	261	095 IC		2500
417 389	389 432	435		ö	1 IJOI	39	Cesspools filled up			10	-	0
	705		876 9	-	3144	90	Public or private wells abolished	:	:	100		CH
od in houses 3855 3872 2768 4	200 4 2		758 42	-	- 344	61.	Houses supplied with town's water		1	-		-
despess wars formed 5007 5545	0000		180 of	15	1.00			200	0.40	1027	84.4	Sant
C+CC +CCC DIMOI SAN CONTROL STORY	2242		200	6	0.00			100	240			1/4
200 310 410 3	310 410	410		0				000	1003	1259 1		4757
42 50 79	50 79	79		8	245			2950	2950 2971 3100	3100	3225 12	12252
240 200	240 200	200		oò.	040		Houses in which all defects found have been remedied a	2653	2653 2665	2830 3	3113 11	11261
1020 A107 AEED A	4407 4550	4550	-		5		Street online cleansed	24.0	000	444		1000
+600+ /+++ on60	4447 43394	40004	4		1,000			144				3
Complaints unfounded 12 47 15 1	_	_	15	-	99	07.	Offensive accumulations removed	61	300	40	44	141
OTHER WORK DONE.						89	Pollutions of river or streams remedied	0.0	7	CH	4	122
Infanting diseases			.h.	- 73	.000	_		1				0
Injective disease 20715 5110	SILO		504 13	+				20	25			340
-	2237		057 25	ĕ	٠,	70.	o inspect work in progress	2492	2222	2515 2		680
	180		310 1	125		71.	Total nuisances abated	3661	3700 4400	7 007	. 14	бкоо
Come only door	See orga		2000	0	-					-		-
10r Cuner causes 1990 1049 1720 2	6tor		7202	0	7552							Ī
		Not love									-	

The figures for the four quarters and year in the above table have in some cases been altered from those given in the quarterly reports, which contain some errors.

In December, Miss Lucy Heath Pearson left us to take up her duties as lady inspector to the borough of Lambeth Educated at the South Hampstead High School and the Girls' Grammar School, Birkhamsted, as well as in Berlin and Cörlitz, Miss Pearson held the higher certificate of the Joint Oxford and Cambridge Board, a diploma of the National Health Society, and the diploma of the Sanitary Inspector's Examination Board, the latter dated 1903. In June of that year she came to us as a probationer, and in May of the following year was appointed as assistant inspector, serving in the latter capacity for about 19 months before she left to take up the more highly remunerated position already mentioned at Lambeth.

WARD INSPECTION.

The number of houses examined completely on account of infectious disease, alleged nuisance, or in connection with house-to-house work, was increased by 266, or from 12,439 in 1904 to 12,705 in 1905. On the other hand the number of houses partially examined fell from 2,865 to 2,851 in the same period. Taken together the houses examined wholly or partially increased from 15,304 to 15,556, an increase of 252 more or less complete examinations during the year.

From line 8 in tables I. and II. it will be seen that the number of houses in which sanitary defects were found was rather smaller than in the previous year, 5,309 against 5,434. On the other hand line 9, the number of defects found, was 7,305 against 7,220 in the preceding year. The special examinations of drains by tests also slightly increased from 10,264 in 1904 to 10,383 in 1905, and the number of defects revealed by this examination from 3,831 to 3,935. The total number of nuisances found, however, fell from 19,115 to 17,853, although the fall amongst those in which the abatement was secured was only from 16,932 to 16,509.

The number of extra visits paid to houses on account of infective disease, nuisances found, completion of reports, and

other causes, was 10,000 greater last year than in the previous year. The total number from all those causes in 1904 was 35,213, and in 1905 45,667. The principal increase was in the first of these groups (line 29).

A large portion of this increase was in the first quarter of 1905, when no fewer than 20,715 secondary visits on account of infective disease were paid, the number in the corresponding quarter of 1904 having been 1,648, instead of 20,715. In that quarter the number of cases of small-pox was 120, as compared with 6 in the corresponding quarter of the preceding year, and a very determined effort was made to clear the town of that disease.

Water closets erected either in lieu of privies or as altogether new amounted to 1,289, against 1,317 the previous year. Drains in course of construction inspected were 1,017, against 1,708 the previous year, whilst 1,941 were reconstructed against 2,361; together 2,958 against 4,069.

The number of disconnections of house drains effected fell from 4,052 to 3,506. It has of course to be remembered that the more work we do of this kind the less work there is to do in the future.

The same remark applies to the falling off of the number of extra visits paid to the work inspected, which fell from 11,230 to 9,689, corresponding to the smaller number of drainage operations required.

A good deal of the Inspectors' time has been occupied in searching for and revisiting the comparatively small number of privies now left unconverted, the difficulty with the last few thousands being naturally greater than at the commencement of our operations.

Women's work in wards.—No separate tables have been kept as last year of this work. What information we have about it will be found in the general report under the head of women inspectors. The work already mentioned is simply that of the ordinary ward and works inspectors.

76

Analysis of work done by District Inspectors in the several Wards, 1905.

1					- 1							
	"AI	а	1066	1753 1753 541	4296	1638		244 148 164	428 438 128 128 128 128 128 128 128 128 128 12	1023	491 985 4301 3805	129 237 237 4866
	tral. th of Lane).	C.	88	178	253	129		11 471	842 ::	12	36 57 270 217	14 17 322 ::
Division on-122,322.	Central. (North of Lady Lane)	1	H	25 364	171	213		144	886 :	179	152 218 1196 1182	222 1243 1243
D. IV. Divisio	wick.	1	4 4	: : 10 :	45	11		HE03	٠:::	6.00	∞r-26	: :19 :
No. IV.	Brunswick.	C	:33	235	1204	514		24 52 52	135	319	145 302 1092 927	41 35 67 1235
No.	. West.	N	300	28 e8	721	235		17	141	139	556 556	32,22,21
	m.jeh.	а	199	367	732	888		1629	5: 458	913	555137	\$888b
1	-guibe		109	285	169	316		Boon	98 : :°°	159	549 549 331	628.38
ion.	noisivie .III	п		2757	4	1020		130 91 149	425 225 25 25 25 25 25 25 25 25 25 25 25	1287	355 416 3256 3104	247 19 132 3654
is :06,8	samley.	В	249	888 888	1031	354		17	127	288	888288	44 25 799 199
i i	mley and Vortley.	W I	526	1042	932	88		888	159	621	3885	523333
No. III Division. Population—105,887.	New Vortley.		11,	882	1064	149		383	35 1 1 1	321	714 698 1	128 178 179 179
N S	West.			910	12	306		37	53 113 2	120	148 107 682 663	21 8 767 767
1905.					Number of houses wholly or partly examined	lefects			utters or	dation	", with drains, &C., temporarily stopped in stopped " s found in houses in which above nuisances were found	
61			1	se work	bed .	ary d			1 : : : : : : :	latio	emp	:::::
st,			. 98	se we	: imi	. : fill		: : 79	9	A	C., t	:::::
31st,			INSPECTION.	Alleged nusances House-to-house work Occupants Buildings and Offices	tly e.	here	&c.	dirty overcrowded damp or dilapidated	ed in	500	is: 11.7	11111
			ECT	se-to se-to ipan ding	par par	es w		wde r dil	fall pipes y drained out sink d y lighted y ventilat	defective loset accor lirty closet	drams, stopped other nu houses	
De	WARDS.		Infe	Hou Docci Buil	lly o	hous in al	CES	dirty overcrowded damp or dilap	fall pipes fall pipes badly drained without sink o badly lighted badly lighted	with di	with of with of d in ho	***
nded Dec.	WAI		- 2	-	who	of above houses where sand	UISANCES,		Y Z.Y ZZ.	× .×.	wi md ii hich	artion ance and and
			HOUSEs and premises	competery examined on account of Houses and premises examined only	uses	nd a		Houses	1111	2 2	,, with drain stopped ,, with other i. Total nuisances found in houses No. of houses in which above nu	Street gulfies stopped Offensive accumulations Other outside nuisances Total nuisances found Complaints unfounded
.s e.			HOO pun	on account uses and prexamined or	r of bo	Total number of were found Sanitary defect	Z	_	5		isanc	Illies racci rside isanc
weeks	1		ses a	on account	uper '	wer wer			Nuisances found in above	houses	d nu	nsive er ou d nu uplain
	:		Hon	Hou	Nun	Tota			a a	hor	Tota No.	Stre Offe Oth Com
5.9			Ti.	d in 4 10 1	6 4	∞ ಕ		0 1 6	÷ + 50 0 10	5 6		2 2 2 2 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
No. II Division. Population-113,552.	noisivi0 ,11	1	1237		l k3	2350	+	154 97 212	434 1320 2 2 3 14	386	486 990 4964 3761	352 2512 5512 50
Vo. II Division Population-113,552	Iolbeck.	I	276		prod.	814	1100	25 69	1935	287	272 418 1725 1195	141 16 56 1938 4
O U	West funslet.	I	35.	986	2	974		35	2588 : :	130	73 102 1479 1051	119 10 68 68 1676
= bulat	East Hunslet.	1	426	328	- 00	180		14.5	148	251	2846 9778	33.08.08.52 2.00.08.08.52
2 %	South		182		- 00	218		45.84	138	88	822 735 735	82880
	noisiviC I	I	1881		14.3	9 1534		2 115 8 77	9238 678 678 25 46 25	7 237	2546	440 440 440 440 440 440
026.	East.		88		1 .4	0 136		0 1 6 2 8 4 8 8	5 47 11 11 11 11 11 11 11 11 11 11 11	3 87	9 190 2 255 4 768	7 182 4 225 0 1007 4
Division.	V. East.	Į	448			909		1113	0 137 9 375 1 1	194	282 282 9 1135 884	77 1470 184 184 1870 1870
D tion-	North.	1	146	245	1219	446		1181	682 :	41	626538	144 13 86 912 3
No. I Divisior Population—115,026.	South of dy Lane).	9	. 85	iα :==	29	88		212	87 · · ·	9.7	8782	7 16 102
				44 5 5	245	38		4 :00	1: 188	192	52525	33073
-	Hill Hill.	X	46	118	1 20						118	. н
	CITY TOTAL	X	4,411 811 46	-	1	5,309		643 386 602	2,884 23 23 11 36	3,577	1,701 3,144 15,450 13,216	1,218 245 940 17,853 3

	g
	2
	25
	8
16	3
	5
	8
	4
	1
	1
ь	÷
•	=
	_
	_
1	
1	
1	
1	
1	ABLE
1	ABLE
1	ABLE
1	

	noisivi ,VI	D!		6567 2011 472 2041	2372 1248 467 7 :: 241 141	301 301 11 15 344 344	889 222 767 1079 3648 3648	3405 243 64 64 54 5948	4647	
	tral. h of ane).	C.		273 138 52 52	F 525 : : : 1 4	13 : : : : : : : : : : : : : : : : : : :	~848 : : : 88€	188	289	
	Central. (North of Lady Lane).	-		273 079 136	521 45 33 143 85	2 : : : : : : : : : : : : : : : : : : :	15 35 102 1589 1589	1378 122 1 123	968	-
	Brunswick.	-		57	ФЮН : : : 10100	8.::::::::	1 : : 6 : : : : 2488	\$00::::	73	
	Bruns	C.		774 494 6 127	172 325 132 132 132 8	65 : : : : : : :	23 152 153 772 772	282 :88	1191	
	West.	'N		1550 650 444	194 194 101 101 179	10 1 10 10 11 11 11 11 11 11 11 11 11 11	25. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	550 97 112 636	872	
!	rley.	Bn		3247 152 413 1165	2853	8 . : : : 8 7 9 : : : 6	159 159 159 251	215 40 8 8 : 1	822	
	-Snibs (part),	ley Hes		587 507 41 116	\$485 :: 500	8::::825::::21	833.21.1 :8888	88 82 82 84 84 85 85 85 85 85 85 85 85 85 85 85 85 85	504	I.
	noisivi III.			3907 1843 78 2403	3177 481 260 : 1 : 1 : 127 : 81	160 845 49 519 519	571 185 45 348 : : 539 614 5190	221 221 15 37 37 2415	3512	See note to table
	amiley.	Bu		910	593 133 68 17 17	45 102 14	37. 38. 17.1 450	419 42 3 23 567	654	te to
	ortley, and	A		868 169 43 493	1451	371 371 371 371 371 371 371 371 371 371	52883 : : 58855	1143 45 45 1149	675	e no
	New ortley.	M		590	888 : : : 888	25 : : : : : : : 52 : : : 52 : : : 52	115 33 47 1120 120 120 5161	109 109 111 111 1111	566 1675	Se
	Vest.	1		539 409 549	897 155 51 37 15 15	8 ::::: ₀ 8 :::2	8884 : : : 8888	262 : 58 583 : 58	617	or.
				::::	1111111111	241111111111	11::::::::	been	-	pect
	:			::::	 n closed shitation	:::::::		have be	1	Works Inspectors and Yiddish Inspector
ontininea.	-		шi	::::	0 11	closets er closet	do. of re-construction inspected do. of re-construction do. inspected when connection made to nections of house drains effected als filled up or private wells abolished and water closets repaired and water closets repaired ouse nuisances remedied ouses for which above work done	H : : :		Yiddi
07110			DOM	: : orts	y te	water of y water () It ter closets , &c.)	inspinstruction on mer freet	defects found removed ams remedied noces removed inspect work in		and
Ĭ				Sep Rep	ins b t for ith	aire b wr ary v ary v arter clos ed, 8	tion ruction ins espect shed swa was was was c wo	fects loves s ret es re		tors
	SDS		WORK	foun of ses	dra ed fi	repaired uilt rough water ordinary wat bs cts built into water ch water-closets water-closets	struc const const dra tholi own;	de :: : : : : : : : : : : : : : : : : :		sbec
4	WARDS.		×	ve d nces etior Cau	servitto ser	&c. s bu repa	con re-c hen ouse close close close rich i	illi sed tions or str nui id to	peri	In s
AB			HER	Infective disease Nuisances found Completion of Reports Other Causes	by di by di tters tters s un s rei d	outing, &c., repaired n privies built privies repaired rebuilt retted into trough water clos do. ordinary water clos the erected hpits or tubs n water closets built converted into water closets treed into water-closets need (lime-washed, &c.)	ourse of construction is or, of re-construction ons of house drains effe lided up rivate wells abolished plied with town's water plied with town's water water closets repaired and which above work s for which above work	which all defects found s cleansed cumulations removed river or streams remedied onestic nuisances removed isits paid to inspect work in	nces abated	Vork
-	-		-	-	amin und ents nd le hous hous hous	spou rel pr rel nvert sets eshpi th w s cou	cour do. spect tions filled priva pplie d wa	ed	ance	1
	:		0	iona paid uses	ts for the same of	ective middle middle Do. jes cot Do. jes cot Closer clos v dry a v troug closet cles ets cle cts cle cts cle cts cle cts cle cts cle cts cle	ins innunec ools ools ools ools ools ools ools ool	remediecet gullie fensive ac llutions o her non-d	nuis	by for
				Additional visits paid to houses for	Special examinations of drains by tests Defects found by ditto Appointments Notices and letters served Dwelling houses unfit for human habitati Houses cleansed Overcrowded houses dealt with	Defective spouting, &c., repaired New midden privies built Old midden privies repaired Do. rebuilt Privies converted into trough water Do. do. ordinary wat Water closets erected New dry ashpirs or tubs New trough water closets built Pail closets converted into water closets T. W. C.'s altered into water closets Closets cleansed (lime-washed, &c.)	Drains in course of construction in Do. do. of re-construction Do. inspected when connections Disconnections of house drains effecesspools filled up. Public or private wells abolished Houses supplied with town's water Trough and water closets repaired Other house nuisances remedied Total houses for which above work	Houses in which all defects Street gullies cleansed Offensive accumulations removed Pollutions of river or streams rem Other non-domestic nuisances ren Additional visits paid to inspect v	Total nuisar	one l
				32. V	5 1 4 3 3 3 3 4 4 4 4 4 4 4 4 4 4 4 4 4 4	55555555555555555555555555555555555555	100 THP CD	65. H 66. S 70. A	71. T	This table includes work done by
	.II			8166 3183 54 1554	2252 2252 256 256 3 5 125 80	211 1183 41 1155 1155 328	132 781 82 909 362 1551 3092	258 258 228 22 22 22 22 22 24 1792	4297	des w
	noisiv			TO THE RESIDENCE	283 : : 283	137 : : : : : : : : : : : : : : : : : : :	225 225 226 226 226 226 226 226 226 226	1109 30 117 2 12 2 501 17		inclu
	inslet,			229 2118 309 1101 22 27 303 283	1010 PO 00101	37	H00H0 00000	553 553 553 553 553 553 553	981 1003 1468	ple
	Vest.	٨		great are 6		8 10 to to 10			1 100	is ta
	1sa2			2802 2017 92 681 305 663			6 41 5 222 6 222 6 222 7 59 7 733	832 48 3 48 17 2 17 2 68 3 68		E
	-dane	S			2 499 170 170 12 12 21 2 12 2 2 2 2 2 2 2 2 2 2 2 2 2	8 : : : : 8	51123 : : : 2008	382 : 358	845	
	noisiv	Die		2 9595 7 2038 7 201 1554	31 2582 31 212 31 212 	1 139 1 57 1 188 1 188 1 248	7 425 2 661 100 1482 1 100 5 5 5 613 2 322 2 322 2 322	285 285 40 40 189 189 2554	4053	
	.isa.	E		5122 57 57 7 7 8 350	43	11 : : : : : : : : : : : : : : : : : :	72222 : : : 4468 282 : : : 4482	321 10 10 57 4	591	
	East.	N.		892 57 892 57 91 7 48 350	227	10 : : : :	126 175 175 176 1069	936 11 752 752	1360 1476	
	orth.	N		834 515 76 342	2239 2239 57 13 15	22. : : : 38	130 239 249 249 277 277 507	643 79 716	1360	
	ntral orth of	(Mo		40 199 162	1 1 200 33	7::::9::::7	13 17 42 51 103 103	1937: 65	171	
	Hill:	Mil		107 375 24 652	273 67 33 4	28 : : : : : : : : : : : : : : : : : : :	79 38 173 173 112 261	138 28 4 198 498	456	
	Cirv	TOTAL		28,235 9,075 805 7,552	10,383 3,985 1,195 1,3 605 348	811 1,188 1,188 1,104 1,104 1,067	1,017 1,941 2,949 3,506 2 2 1,476 4,757	11,261 1,007 141 12 348 9,689	16,509	

INSPECTION OF WORKPLACES.

Bakehouses.—At the end of 1904 there were 856 bakehouses on our list in the town. At the end of 1905 the number had increased to 997; 223 new ones having been put upon our list, and 82 existing ones removed. The total number of visits paid by the ward inspectors was 2,542. In addition to these the senior workshops inspector (Mr. Lonsdale) made 35 visits, and my laboratory clerk (Mr. Ferguson) 70, nine of them with Mr. Lonsdale, eight with myself, 53 alone.

In the Central ward, which is in the care of three different inspectors, the 10 overground bakehouses in the northern part visited by the ordinary inspector increased to 16; nine new bakehouses having been added to our list and three old ones discontinued. The visits he paid were 45. Underground the five existing bakehouses were reduced to one, four having been discontinued; the visits paid were six. In the same part of the Central ward, but visited by the Jewish inspector, the overground bakehouses had been 10. They increased by the addition of a new one to 11, and to them 75 visits were paid. The underground ones, which had been three, were reduced to two by the closing of one; 16 visits were paid. In the southern part of the Central ward (south of Lady Lane), the seven overground bakehouses on our list were increased to II by the addition of four; 25 visits were paid. The two underground bakehouses were increased to three by the addition of one; six visits were paid. The total for the whole ward was therefore 27 overground bakehouses at the beginning; 38 at the end of the year; 14 new ones put on our list and three old ones discontinued. The visits made by the ward inspectors during the year were 145. The underground bakehouses, which were 10 at the beginning of the year, were six at the end; 28 visits were paid by the ward inspectors. Mr. Lonsdale paid one to an overground bakehouse in this ward, and Mr. Ferguson three to underground ones.

TABLE III.

Shewing the number of drains or sources of pollution diverted from the River Aire and its tributaries and connected to the town sewers.

Date.	Mills, factory, house drains, stables, and pigstyes.	Water closets.	Privies.	Trade pollu- tions.	Total.
Previous to 31st Dec., 1904	5,840	494	233	78	6,645
During the 52 weeks ended 30th Dec., 1905	31	12			43
Totals	5,871	506	233	78	6,688

In the North Ward the inspector reports 49 at the beginning, 45 at the end of the year; two new, six discontinued; 92 visits. These do not include underground bakehouses, which were five all throughout the year, 11 visits having been paid. Mr. Lonsdale made four visits to bakehouses above ground, one to an underground; Mr. Ferguson 10, all to undergrounds, one with Mr. Lonsdale, eight alone, and one with me.

North-East.—Overground 28, increasing to 66; 43 new, five discontinued; ward inspector's visits, 154. Underground four throughout the year; six visits paid. Three new underground bakehouses were opened but closed within the year. Mr. Lonsdale's visits were one to an overground, three to undergrounds. Mr. Ferguson's 10, six alone, three with Mr. Lonsdale, one with myself, all to undergrounds.

East ward.—Overground, 28, increasing to 31; three new, 67 visits. Underground, two at the beginning, which were both discontinued; two visits by ward inspector. Mr. Ferguson also visited one of these alone.

South ward; 60, increasing to 65 by the addition of five new ones; ward inspector made 264 visits. Underground, seven throughout the year; 36 visits. Mr. Lonsdale made two visits to two overground bakehouses; Mr. Ferguson two alone and a third with me to the underground variety.

East Hunslet.—88, increasing to 89 overground; seven new, six discontinued; 219 visits. Underground, three throughout the year; six visits. In addition to these visits by the ward inspector, Mr. Lonsdale made three to overground and three to underground, Mr. Ferguson two with Mr. Lonsdale to the latter.

West Hunslet.—66, increasing to 116 by the addition of 50 to our list; 243 visits. Underground, six throughout the year; 15 visits by ward inspector. Mr. Lonsdale visited three, one to overground, two to underground. Mr. Ferguson three more to undergrounds, one of them with myself.

Holbeck.—106, falling to 105; nine new, 10 disused; ward inspector's visits, 249. Underground four, falling to three by closure of one; 10 visits. Additional visits to underground, two by Mr. Lonsdale, on both occasions with Mr. Ferguson.

Mill Hill.—Visited by the same inspector as the southern part of the Central ward; overground 14, increasing to 15 by the addition of one new; 64 visits. Underground five throughout the year; 20 visits. Mr. Lonsdale made two overground and one overground visit. Mr. Ferguson made one visit to an underground with Mr. Lonsdale, and three alone.

West.—Overground 47, increasing to 59; 18 new, six discontinued; 104 visits. Underground, three throughout the year; five visits. Additional visits by Mr. Lonsdale to overground, three, and six by Mr. Ferguson to underground alone.

North-West.—Overground 57, reduced to 56; one new, two discontinued; 161 visits by ward inspector. Below ground

nine, reduced to seven by disuse of two; 23 visits. Mr. Lons-dale visited one overground, Mr. Ferguson 14 underground, one of the latter with me.

Brunswick.—Overground 36, increased to 44 by the addition of eight to our list; 105 visits by ward inspector. Underground eight, reduced to three by the disuse of five; 13 visits. This ward is visited by the same inspector who inspects the non-Yiddish part of the northern Central ward. Mr. Lonsdale also visited one aboveground, and Mr. Ferguson made three underground visits.

Armley and Wortley.—Aboveground 36, rising to 56, one discontinued, 21 new; 113 visits. Underground eight, falling to four by the discontinuance of four; six visits by ward inspector. Mr. Ferguson visited one underground bakehouse with me. Mr. Lonsdale visited one overground.

New Wortley.—Overground 56, increasing to 57 by the addition of one new bakehouse; 141 visits. No underground bakehouse. Mr. Lonsdale made an additional visit to an overground.

Bramley.—Aboveground 13, increasing to 22; 11 new; two disused; 75 visits. Underground, two throughout the year; ward inspector's visits, 7. Mr. Lonsdale made one underground visit.

Headingley (without Burley).—Aboveground 10, increasing to 19; 10 new, one discontinued; 20 visits paid. Underground, two at the beginning, one at the end, by the disuse of one; two visits by ward inspector. Mr. Lonsdale also made three overground visits.

Burley (part of Headingley ward).—Overground, 52 at the beginning, 45 at the end of the year; 10 new, 17 disused; 112 visits. Underground five, increasing to 10 by the addition of five newly-discovered. 24 visits by ward inspector. Mr. Ferguson made eight visits underground.

At the foregoing, 928 bakehouses above ground, 268 persons not members of the family of the occupier found work in 132. In 796, only members of the family were employed. In the 69 underground bakehouses, 48 persons outside the occupier's family were employed in 30 of them. In the remaining 39, only members of the occupier's family were employed. Baking, it is evident in Leeds, is still chiefly carried out in private families, supplemented by a few domestic bakeshops.

Shewing the sanitary conditions at time of visit of workshops on register and occupied, during the 52 weeks ended December 30th, 1905.

TABLE IV.

	Employees S & AT TIME OF VI							CONDITION OF PREMISES.			
DATE.	DATE. Visits to		4:		- 1		rou	ms.	closets.		rksho d clo
1905.	Vi	male.	female.	total.	good.	defect- ive.	clean.	dirty.	clean.	dirty.	Workshops found closed.
4 weeks ended Jan. 28	160	344	722	1,066	158	2	158	2	147	13	16
4 weeks ended Feb. 25	111	166	283	449	110	I	98	13	105	6	2
5 weeks ended Apl. I	41	260	223	483	36	5	26	15	20	21	9
4 weeks ended ,, 29	51	207	349	556	46	5	37	14	35	16	18
5 weeks ended June 3	250	1,427	1,446	2,873	241	9	222	28	184	66	14
4 weeks ended July I	61	273	346	619	61		53	8	49	12	5
4 weeks ended ,, 29	185	713	753	1,466	183	2	175	10	163	22	18
5 weeks ended Sept. 2	73	153	281	434	72	1	69	4	66	7	8
4 weeks ended ,, 30	74	211	429	640	73	1	70	4	66	8	6
4 weeks ended Oct. 28	125	497	441	938	121	4	113	12	107	18	19
5 weeks ended Dec. 2	192	953	861	1,814	189	3	177	15	177	15	22
4 weeks ended ,, 30	139	989	703	1,692	138	I	133	6	121	18	15
Totals	1,462	6,193	6,837	13,030	1428	34	1331	131	1240	222	152

Workshops.—The visits paid to workshops already on the register were 1,462, against 1,489 in the previous year. The conditions tound are shown in table IV. On the whole, they are more favourable than last year. As explained in my last

TABLE IVa.

Factories, Workshops, Laundries, Workplaces, and Homework. 1.—INSPECTION.

Including Visits and Inspections made by Sanitary Inspectors or Inspectors of Nuisances.

		Number of					
Premises.	Inspections.	Written Notices.	Prosecutions.				
Factories (Including Factory Laundries.)		694	64				
Workshops (Including Workshop Laundries.)		3,613	426				
Workplaces		277	3				
Homeworkers' Premises		1,026	41				
Total		5,610	534				

2.- DEFECTS FOUND.

	N	umber of Defe	ects.	Number
Particulars.	Found.	Remedied.	Referred to H.M. Inspector.	of Prosecu- tions.
Nuisances under the Public Health Acts:-*				
Want of cleanliness	273	294		
Want of ventilation	35	33		
Overcrowding	6	6	***	
Want of drainage of floors	I	I		
Other nuisances	180	161		
Sanitary accom- (insufficient	36	39	***	
modation. unsuitable or defective	313	288		
Sec. 22 in force. not separate for sexes	21	16	***	***
Offences under the Factory and Workshop Act:				
Illegal occupation of underground bake-				
house (S. 101)	68	28		
Breach of special sanitary requirements for				
bakehouses (SS. 97 to 100)	I	2	***	***
Failure as regards lists of outworkers (S. 107)	129	129		
Giving out work to be (unwholesome (S. 108)	***		***	
done in premises				
which are (infected (S. 110)		***		
Allowing wearing apparel to be made in				
premises infected by scarlet fever or				
sniallpox (S. 109)		***		
Other offences	***	***		
Total	1,063	997		

^{*} Including those specified in Sections 2, 3, 7, and 8, of the Factory Act as remediable under the Public Health Acts.

3.-OTHER MATTERS.

Class.	Num	ber.
Matters notified to H.M. Inspectors of Factories: Failure to affix Abstract of the Factory and Workshop Act (S. 133) Action taken in matters referred by H.M. Inspectors as remediable under the Public Health Acts, but not under the Factory Act (S. 5). Other Other Motified by H.M. Inspector Reports (of action taken) sent to H.M. Inspectors	6	6 3 7
Underground Bakehouses (S. 101):— Certificatés granted during the year In use at the end of 1905	6	8
Homework:—	Numb	per of
List of Outworkers (S. 107):—	Lists.	Outworkers
Lists received	390	1,040
Addresses of outworkers { forwarded to other Authorities received from other Authorities	3	
Homework in unwholesome or infected premises:—	Wearing Apparel;	Other.
Notices prohibiting homework in unwholesome premises (S. 108) Cases of infectious disease notified in homeworkers' premises Orders prohibiting homework in infected premises (S. 110)		
Workshops on the Register (S. 131) at the end of 1905 (Including 162 workshop bakehouses, not including 835 domestic bakehouses). Total number of workshops on Register		199

report, these are complete inspections, not mere visits, and in table IVa. it will be seen that our visits to workshops total up to 3,613, exclusive of 277 visits paid to workplaces and 694 to factories. In table IV., only complete inspections are counted. In table IVa. every official visit is considered as an inspection.

In table V. will be found the work done in visiting workplaces coming on our register for the first time. During the year these numbered 179 against 192 in the preceding year. The general description of the drainage as to disconnection was on the whole better in new workshops than the previous year. Although the number of inside closets was actually fewer, the proportion of those with inside closets in

TABLE V.

Shewing workshops not previously visited to which visits have been paid during the year, and the sanitary arrangements at time of visit.

			RIPTIC		Sit	UATION	OF CLOS	ETS.	
	to to er.	-	AINA	GE.	ins	ide.	out	tside.	hops
DATE. 1905.	Workshops added to register.	cut off.	not cut off.	none.	w.c. soil-pipe carried up.	w.c. soil-pipe not car- ried up.	Water Closets.	Privies.	Workshops found closed,
4 weeks ended Jan. 28	9	5	1	3	4		5		
4 weeks ended Feb. 25	6	4	I	I	3		3		
5 weeks ended April 1	21	9	2	10	9		12		
4 weeks ended ,, 29	12	8	2	2	4		8		
5 weeks ended June 3	21	14	5	2	7		14		I
4 weeks ended July I	13	5	3	5	I		11	1	
4 weeks ended ,, 29	12	9	I	2	5		7		I
5 weeks ended Sept. 2	14	8	3	3	5	I	6	2	I
4 weeks ended ,, 30	13	9	2	2	3	,	10		
4 weeks ended Oct. 28	20	15	2	3	5		15		4
5 weeks ended Dec. 2	26	12	6	8	9		17		I
4 weeks ended ,, 30	12	6	***	6	4		8		3
Totals	179	104	28	47	59	I	116	3	11

which the soil pipe was carried up the same diameter above the eaves of the building was larger. The number with outside water closets was larger both actually and proportionately, and the number of privies smaller, also both actually and proportionately. In tables IV. and V., and also in table IVa., the visits paid by women inspectors are included. The work shown in table Va. (outworkers) is entirely work done by the ladies.

TABLE Va. (Outworkers).

Showing visits paid by Women Inspectors to the houses of Homeworkers during the 52 weeks ended 30th December, 1905.

w	Complete inspections	Addit	ional visits	about	T1
Wards.	of houses on first visit.	work ordered.	infectious disease.	other causes.	Total.
				- 12	1
Central	30	14		29	73
North	2 I	6		29	56
North East	57	25	2	37	121
East	73	9		41	123
South	4	2		1	7
East Hunslet	11	1		11	23
West Hunslet	19	***	3	20	42
Holbeck	22	4		33	59
Mill Hill	6			11	17
West	136	24		113	273
North West	20	10		16	46
Brunswick	2 I	17		9	47
New Wortley	32			27	59
Armley and Wortley	18	3		16	37
Bramley	1				1
Headingley	22			13	35
Totals	493	115	5	406	1,019

Turning for a moment to table IVa. (the Home Office table) it will be noticed that the number of inspections and visits paid to factories was larger than last year, although the number of written notices was fewer. On the other hand the

TABLE Vb.
Visits paid by Senior Workshops Inspector on account of

				Factories.	Workshop
Non-abatements			 	200	512
Drain inspection			 	7	I
Disease enquiries			 	65	14
River pollution			 	28	15
Complaints			 	134	200
Measurement of	workro	oms	 		20
Wage enquiries			 	. 14	11
Bakehouses	underg	ground	 		13
Dakenouses	above	ground	 		2 2
Other causes		***	 	96	220
Appointments			 	66	51
	<u> </u>	TOTAL		610	1,079

Work done.

		Factories.	Workshops
Drains tested	 	 I 2	19
River pollutions abated	 	 28	15
Nuisances abated	 	 122	223

inspections of workshops (3,613) was fewer than last year, though the number of written notices (426) was nearly double that of the previous year. The cause of this great increase is probably to be found in the encouragement of the written notices as against the verbal. In many cases now both are

employed where formerly only a verbal notice was given. The number of visits to workplaces, all part of the women's work, was 277 against 477; the notices three against 24. The home workers received 1,026 visits and 41 notices, against 2,453 and 96 last year. The reason for this apparent diminution in the work carried out by our women inspectors (who made 1,019 of the 1,026 visits) is that their time has been very largely taken up with the investigation of the conditions prevailing in the homes of the new-born in the South-East Registration District, and in the following up and watching the progress of those children. This work is an exceedingly valuable one, but as our staff has not been increased some of it had to be done at the expense of inspections of workplaces and home workers' premises. Correspondingly there is a diminution in the number of defects found and dealt with, enumerated in part 2 of the Home Office table. The number of out-workers does not differ very much from last year.

The matter of underground bakehouses has been already referred to.

Other work.—The special work of the senior inspector, not included in tables IV. and V., is given in table Vb., which shows a slight increase on the figures of last year.

WOMEN INSPECTORS.

Infectious disease.—On account of puerperal fever 29 houses have been completely examined and 103 other visits paid. Thirteen additional visits were paid to nurses and midwives who were connected with the cases, and 17 visits were paid by midwives to the disinfecting station for the purpose of being disinfected personally, and as to their wearing apparel, under the immediate oversight of our inspectors. This number does not include the relations and neighbours who nursed the patients, and whose clothing was sent from the house where the illness occurred to be disinfected at the station.

On account of small-pox 108 visits were made to factories, 260 to other workplaces, and 328 to schools; 675 visits were paid to contacts at home because absent from school or workplace.

Only two visits were paid in consequence of measles, the Medical Officer of Health considering it inadvisable that the ladies should visit such cases on account of the important work they were doing at the time amongst infants.

On account of infectious disease amongst persons employed, 32 visits and enquiries were made in factories, 10 in workshops, 3 in other workplaces, and one in a private house. Five houses of absent employees were visited in connection with this work.

In the girls' and infants' departments of elementary schools 224 inspections were made on account of infectious illness, not including small-pox. To the homes of absentees 247 visits were paid, 107 return visits to schools, and 45 other visits were made in connection with this branch of work. Notices to the number of 37 were sent to the officers of the Education Committee warning them of cases in which it was considered inadvisable that children should return to school until medical advice had been obtained. It was necessary to give 6 notices for cleansing and limewashing of schools, all of which were attended to. There were also 5 notices in respect of other nuisances all of which were complied with.

Outworkers.—Table Va. gives, as in previous years, the work of the women inspectors in visiting at their homes persons employed as outworkers. In addition to the figures given in table Va. as to outworkers, II visits were paid to their employers.

Fourteen structural defects, rendering insanitary a much larger number of the houses of home workers required notices which were all complied with. In 39 houses notices for

cleansing or limewashing were carried out. Here as many were for the cleansing of closets or yards used by several occupiers, the actual number of notices was in excess of the nuisances themselves.

Workplaces.—There were 1,092 inspections made of workshops and 1,213 return visits were paid. Five notices were given for over-crowding and complied with, 12 for defective ventilation and complied with, 177 for limewashing and cleansing, of these two have lapsed because occupiers have left. In the case of 136 notices for other nuisances 4 lapsed on account of the workshops having been closed and 9 are still unremedied. Of these notices 142 were verbal.

In workshop laundries 19 inspections were made and 40 return visits paid. One notice for nuisance was sent and complied with.

Of domestic laundries there were 50 inspections and 14 other visits were made. No notices were sent.

Of restaurant kitchens 170 inspections were made and 102 other visits paid. There were 10 notices sent on account of defects, one of which in respect of defective ventilation remains unremedied. There were also 42 limewashing and cleansing orders all carried out.

In connection with factories 163 visits were paid (32 as already stated being in respect of infectious disease). For cleansing and limewashing 8 notices were given and all complied with. Two notices for other nuisances were given, one was attended to and the other not. All these notices were verbal.

Infant mortality.—During the year in South-east Leeds 295 investigations were made of the deaths of infants and 270 return visits paid.

In connection with this work 34 notices were given and 32 were complied with, 1 lapsed owing to the house being

closed, and I was not attended to. Limewashing notices to the number of 20 were given, all of which were complied with.

Visits to newly born.—In the beginning of 1905 the visiting of houses in this district, where a birth had occurred, was commenced. Such visits were made in 1,236 cases, detailed information collected and recorded. In connection with the work of the Committee for the supply of pure milk in this district a systematic revisiting of infants receiving this milk was carried out during the latter part of the year. In the course of this work 6,083 visits have been made to the mothers and guardians of the children under supervision. This matter is alluded to elsewhere.

Midwives' Act.— In addition to the work already given, a considerable amount of time has been occupied in the carrying out of the Midwives' Act, 1902. Our Chief Lady Inspector early in the year obtained the certificate of the London Obstetrical Society. She reports:—

"In the early part of the year every effort was made to "ensure that all women practising as midwives in Leeds should be informed about the existence and provisions of the Midwives' "Act, 1902. In spite of this, several midwives who ought to have registered did not apply till after March 31st, 1905. Of those who notified their intention to practise during the year, "75 did so by virtue of having been bona fide midwives in July, "1901, and five as holding recognised certificates, namely, three from the London Obstetrical Society, one from the Manchester "Maternity Hospital, and one from the Glasgow Maternity "Hospital. Of the bona fide midwives, two have left the town, one has moved outside the borough boundary, and one is dead. "Several of those who have notified their intention to practise have not so far taken cases on their own responsibility, but have acted as monthly nurses.

"A very large proportion of the midwives are illiterate; some, however, can read, though they cannot write. Those who can read the clinical thermometer are few. Six of those on the register are between 30 and 40 years of age, 29 between 40 and 50, 22 between 50 and 60, and 19 between 60 and 70 years of age. The total number of midwives now practising is 76.

"A series of six lectures was arranged for those midwives "who had notified their intention to practise. There was an "average of a little over 44 out of 77 possible attendances. "This, considering the nature of their work and the distance "from some of their homes, may be looked upon as good. The "lectures were also attended by a few district nurses and others "interested. The midwives showed that they had benefitted by "the lectures, and expressed their gratitude to the lecturers.

"Notifications of 56 still births have been received from midwives as having occurred during the year 1905. There were 102 notifications of requests for medical assistance. Eight deaths of infants were notified where there was no medical practitioner in attendance. It is to be expected that the number of notifications will increase as the midwives come to realise their obligations more completely.

"The proportion of cases attended by midwives in the "South-east Registration District, one of the poorest in the "town, was 554 out of 1,243 births registered during the year. "There were 44 births in this district where the attendance is "unknown.

"During the year 193 inspections were made of bags and case books, and in each case the cleanliness of the person and house of the midwife was reported upon, and a certain amount of instruction on the rules of the Central Midwives' Board was given. Other visits to the houses of patients and to doctors to investigate reported cases of neglect were paid to the number of 58, and 152 interviews with midwives took place at the

"Public Health Offices. Seventeen personal disinfections of midwives were superintended by our inspectors. In each case the inspector arranged that all facilities for obtaining a bath at the disinfecting station should be given. In 16 cases the inspector was present. Not only were the midwife's clothes disinfected by steam, and not only did she pass through the bath to the uninfected side of the station, but the inspector personally ascertained that she actually had the bath, examined ther hands and finger nails carefully after her bath, and made ther place her hands in a disinfecting solution for a few minutes."

Other work.—In addition to the above work 44 visits were paid to lavatories used by women, 7 nuisance notices were given and complied with, and 2 notices for cleansing and limewashing, both complied with. Notice was also sent for the cleansing and limewashing of a midwife's house and complied with. In consequence of complaints 18 notices were given for ordinary nuisances, 17 were remedied and 1 was not. Fifteen other cleansing and limewashing notices were sent, 11 were complied with, 3 were not attended to, and 1 lapsed.

CLEANSING.

Under the head of changes in staff mention has been made of this subject.

I print the usual tables and statistical material.

Ashpits inspected by Nuisance Inspectors during the 52 weeks ended 30th December, 1905.

Number of inspections of ashpits.	Requisitions to cleanse sent to Cleansing Sub- Department from Sanitary Office.	Number of latter returned as carried out.	Ashpits not cleansed within four days of requisition.	Condition of ashpits generally.
22,820	5,337	5,334	12	Good.

TABLE VII.

Refuse Removal (same period).

No. of	No. of	No. of		LOA	DS REMO	VED.	
No. of midden emptyings.	dry ashpit or tub emptyings.	box or pail emptyings.	Total.	Night- soil.	Dry ashes.	Rubbish.	Total.
3,328	1,288,551	14,490	1,306,369				191,814

Destructors.—The following work has been done at the four destructors:—

At Armley Road 29,530 loads of rubbish, weighing 25,477'8 tons (0'86 tons or 17'26 cwts. per load) were destroyed during 309 working days. On an average of 12 cells, the work per cell per day would be 6'87 tons. The highest observed temperature was 1,500° Fahr.,* the lowest 500°; average, 1,477°. There were 7,416 observations taken. The firemen employed averaged 14'6 a day, and the amount turned over by each averaged 5'68 tons daily.

At Beckett Street, 23,312 loads of rubbish, weighing 18,804.2 tons (0.81 tons or 16.13 cwts. per load) were destroyed in 8 cells during 309 working days, being an average of 7.61 tons per cell per day. The highest observed temperature was 1,500° Fahr.,* the lowest 1,260°; average 1,497°. There were 7,416 observations taken. The average number of firemen employed was 9.94, and their daily turnover 6.1 tons.

At Kidacre Street, 25,955 loads of rubbish, weighing 21,382'9 tons (0'82 tons, or 16'48 cwts. per load) were destroyed in 12 cells during 309 working days, being an average of 5'77 tons per cell per day. There were 7,416 observations of temperature taken, the readings averaged 1,459° Fahr., the highest was 1,500°,* the lowest 560°. The average number of firemen was 11'85, and they turned over an average of 5'84 tons a man daily.

^{*} The pyrometer does not register above 1,500° Fahr.

At Meanwood Road, 19,826 loads of rubbish, weighing 16,902'4 tons (0'85 tons or 17'05 cwts. per load) were destroyed in 8 cells during 309 working days, being an average of 6'84 tons per cell per day. There were 7,416 observations of temperature taken, the readings averaged 1,461° Fahr., the highest was 1,500°,* the lowest 600°. The average number of firemen was 6'75, and they turned over an average of 8'1 tons each daily.

Street cleansing.—This heading is intended to include the work done in the ordinary sweeping and gully cleansing to the paved streets, in the watering of streets and roads, in the cleansing of certain Macadam roads and their gullies and in the removal of snow.

Mr. Mann reports that during 311 working days, 314,249 street cleansings were effected, an average of 1,010 a day. The length of paved streeting cleansable by us is 252 miles.

The cleansing of gullies was equivalent to cleansing 225,595 or an average of 725 per day; 225,595 charges of disinfectant were applied to gullies. In addition to this, 67,015 loads of street refuse were carted away (215 a day), and 124,436 cleansings of courts and yards were effected, an average of 400 a day.

The number of horse-days for street cleansing was 24,083 and for watering 5,969, being an average of 77 and (during 111 days) 54 horses a day respectively.

During the III street watering days 120,109 barrels of water were used.

The work done in connection with the unpaved streets was equivalent to the cleansing of 29,250 roads and 11,277 gullies; whilst 20,354 loads of refuse were removed, averaging for 300 working days, respectively 98 roads, 38 gullies, and 68 loads per day. The number of horse-days was 5,171, an average of 17 horses a day.

^{*} The pyrometer does not register above 1,500° Fahr.

During the 52 weeks, the sub-department removed 6,240 loads of snow at a cost to the city of £495.

The cost of street cleansing was £23,030, of street watering, £3,827, and of cleansing highways £7,494, making a total cost (including snow removal) of £34,846.

FOOD INSPECTION.

FOOD, DRUGS, AND DAIRIES.

Cowkeepers.—Table VIII. contains, as in recent years, the ordinary statistical information about the visits of our ordinary inspectors, and in addition the number of visits paid by the Veterinary Surgeon. It contains also the usual information as to the number of cowkeepers in the city, the number of milk vendors on the register and the alterations in the latter during the year.

Condition of sheds.—Table VIIIa. contains, as usual, the conditions of the sheds found when the Veterinary Surgeon made his inspection of the cattle, when examining the latter principally on account of the condition of their udders.

TABLE VIII.

Work done in connection with the Cowsheds and Dairies Order during the 52 weeks ended 30th Dec., 1905.

Cowkeepers on the register .			149
Milk-retailers " .			510
Visits to cowsheds			886
" " (Veterinary	7)		294
" milkshops			791
" railway stations .			210
Cowkeepers discontinued busin	ness		7
New cowkeepers registered .			13
New cowsheds built			
Cowsheds reconstructed and			
additional light and ventilat			13
Farms or milkshops visited of	on accour	nt of	
infectious disease			16

TABLE VIIIa. Veterinary inspection of Cattle, year ended 30th Dec., 1905.

			Cattle	and Cond	ition.	
Date of	Register					Condition of Shed.
Inspection	No.	Ward.	No.	Udder diseased.	General	Condition of Shed.
			15 Administra	discused.	Condition	
1905.						
Jan. 3.	41	N.E.	4		healthy	Dirty and unfit for use.
,,	478	,,	12		,,	Satisfactory.
	IOI	E.	19		,,	Do.
	351	,,	10		,,	Do.
	172	,,	28		,,	Do.
	145	,,	22		,,	Do.
	145a	,,	16		,,	Do.
	964	,,	16		,,	Do.
,,,	19	,,	16		,,	Wants cleaning.
Jan. 13.	835	S.	9		, ,,	Satisfactory.
,,	198	E.	13	I	others healthy.	Do.
	90		26	(mammitis)	healthy	Do.
"	329	,,	13			Do.
"	1011	27	6		"	Do.
Jan. 20.	520	N.E.	15		"	Do.
1	1108	,,	27		"	Do.
,,	220		25		,,	Do.
",	136	"	15			Do.
"	138	"	46	1	others healthy.	Do.
",		"		(mammitis)		
Jan. 26.	553	Bnk.	13		healthy	Do.
"	67	N.	22	***	"	Do.
,,	170	"	22		"	Do.
,,	4	,,	12		"	Do.
,,	665););	18		12	Do.
,,	295	N.E.	11		others	Do.
"	34	E.	12	(mammitis)	others healthy.	Do.
Feb. 8.	45	N.	17		healthy	Do.
,,	602	,,	15		,,	Do.
,,	343	,,	15		,,	Do.
,,	196	,,	2 I		"	Do.
,,	896	,,	9		,,	Do.
,,	343a	,,	26		22	Do.
Feb. 21.	104	A. &W.	22		"	Do.
,,	38	,,	24		12	Do.
,,	99	,,	23	1	others healthy.	Do.
1	,			(tuber- culous)		
,,	89	Brmy.	3		healthy	Do.
,,	126	,,	13		"	Do.
,,	63	,,	7		,,	Do.
,,	360	,,	18		,,,	Wants cleaning.
,,	337	*,,	2 I	1	others healthy.	Satisfactory.
March 1.	553	Bnk.	9	(mammitis)	healthy	Do.
	39	N.	46			Do.
",	66	C.	12		"	Do.
	00	0.	1.6		,,	170.

			Cattle	e and Conc	lition.	
Date of Inspection.	Register No.	Ward.	No. Examined	Udder diseased.	General Condition	Condition of Shed.
1905.						
March 1.	7	C.	5		healthy	Satisfactory.
,,	153	,,	10		23	Clean but badly ventilated.
"	151	;;	8		"	Satisfactory.
,,	88	N.W.	26		,,,	Do.
"	154	,,	4		,,	Fairly clean, very old build- ings, and close to houses.
March 9.	49	. ,,	24		,,	Satisfactory.
,,	393	Hdy.	34		,,	Do.
,,	347	,,	8		,,	-Do.
,,	195	,,	18		,,	Do.
,,	833	N.	24		,,	Do.
,,	75	Hdy.	9		,,	Do.
-,,	338	N.	2		,,	Do.
,,	714	Hdy.	8		,,	Do.
March 16.	94	,,	8		,,	Do.
,,	213	,,	5		,,	Do.
,,	609	,,	16		,,	Do.
,,	550	,,	11		,,	Do.
,,	676	"	7	***	,,	Do.
,,	353	"	21		1,	Do.
,,	384	,,	I 2		,,	Do.
,,	470	,,	7))	Do.
"	146	,,	26	2 (mammitis)	others healthy.	Do.
,,	146a	,,	8	***	healthy	Do.
March 29.	932	,,	26		,,	Do.
,,	912	,,	24		,,	Do.
,,	167	,,	12		,,	Do.
,,	389	,,	11		,,	Do.
,,	431	A. & W.	29		,,	Do.
"	813	,,	26	I	others' healthy but	Very dirty.
April 3.	36	Brmy.	2	(mammitis)	healthy	Satisfactory.
	243	,,	8	9	,,	Do.
"	682	- ,,	20		,,	Dirty.
,,	201	,,	4		,,	Satisfactory.
,,	822	,,	8		"	Do.
,,	21	,,	9		,,	Do.
,,	352	,,	6		,,	Do.
,,	110	,,	4	I (tuber-	others healthy.	Do.
	227	-	8	culous)	healthy	Wants whitewashing.
,,	574	,,	14			Satisfactory.
April 7.	206	A. &W.	4		"	Do.
	392	W.H.	16		"	Do.
,,	390		19	I	others	Do.
,,	390	"		(mammitis)	healthy.	

Г		1		Cattle	and Cond	ition. 1	
	Date of Inspection.	Register No.	Ward.	No. Examined	Udder	General Condition	Condition of Shed.
	1905. April 7.	135	W.H.	16		healthy	Satisfactory.
1	"	85	"	7		"	Do. Do.
1	"	359 266	E.H.	2		. "	Do.
1	"			3 4		"	Walls dirty, ventilation bad.
1	April 12	557 408	Brmy.	14		"	Dirty.
1	,,	148	,,	9		"	Do.
1	,,	70	,,	4		,,	Satisfactory.
ı	,,	147	"	5		,,	Do.
1	,,	322	,,	4		,,	Do.
	April 14	351	E.	9		,,,	Do.
	May 17.	321	Hdy.	I 2	I	others healthy.	Do.
	,,	812		7	(mammitis)	healthy	Do.
L	,,	589	N.W.	5		,,	Do.
ı	"	543	N.	19	***	,,	Do.
П	"	170	,,	23		,,	Do.
1	June 15.	67	,,	26		٠,	Do.
П	July 1.	909	Brmy.	2	I	others healthy.	Do.
	July 7.	909	,,	13	I	,,	Do.
	July 26.	136	N.E.	14	I	,,	Do.
1	,,	222	,,	19	(mammitis	healthy	Do.
1	,,	1108	,,	26		,,	Do.
1	July 28.	285	,,	9		,,	Do.
П	,,	170	N.	25		,,	Do.
1	,,	543	,,	21		,,	Do.
1	August 4.	67	,,	7		,,	Do.
1	,,	909	Brmy.	I	I		Do.
1					(tuber- culous)		
1	August 10.	332	W.H.	55	***	,,	Do.
	"	43	,,	37		,,	Do.
1	Amount xx	392	N.E.	17	***	,,	Do. Do.
1	August 11.	136		1 28	1	others	Do. Do.
	",	138	"	38	(mammitis		
	,,,	45	N.	17		healthy	
1	August 15.	609	Hdy.			"	Satisfactory.
	,,	213	,,	I		"	Do.
15	"	247	"	7		,,	Do. Do.
-), August 16	393	,, E	32		"	Do.
1	August 16.	145	E.	16		"	Dirty.
	"	964	"	3.0		"	Satisfactory.
	"	100000000000000000000000000000000000000	"	15		"	Do.
1	August 18.	153	ď.	7		"	Satisfactory as to cleanli-
1	2050000	.33	0.	'		,,,	ness but structurally bad.
			1	1		1	

TABLE VIII a .- Continued.

7			Cattle	e and Cone	lition.	
Date of Inspection.	Register No.	Ward.	No. Examined	Udder diseased.	General Condition	Condition of Shed.
1905.						
August 18.	17	C.	6		healthy	Satisfactory.
,,	66	,,	II		,,	Do.
,,	39	N.	46		,,	Do.
August 23.	67	,,	13		,,	Do.
August 29.	390	W.H.	22		,,	Dirty.
,,	135	,,	10		,,	Satisfactory.
,,	64	,,	10		,,	Do.
,,	150	Hol.	13		,,	Do.
August 31.	146	Hdy.	23		,,	Satisfactory. Watersupp
						required.
,,	146a	,,	17		,,	Satisfactory.
,,	912	,,	2 I		"	Do.
Sept. 1.	353	.,,	19		,,	Do.
,,	67	N.	16		"	Do.
,,,	138	N.E.	38		"	Do.
Sept. 4.	38	A.&W.	25		"	Do.
"	114	,,	22		others	Do.
"	431	"	30 cows	I	healthy.	Do.
Comb 6	813);	out.		1	Dirty.
Sept. 6.	67	N.	16		healthy	Satisfactory.
Sont =	45	LId.	17		"	Do. Do.
Sept. 7.	146	Hdy. A.& W.	18		"	
Sept. 11	813		000755	I	"	Dirty. Satisfactory.
,,	431	,,	I	suspicious		Satisfactory.
Sept. 12.	360	Brmy.	18		healthy	Do.
,,	63	,,	cows out.			Do.
,,	243	,,	,,			Do.
,,	201	,,	,,			Do.
,,	682	,,	,,			Do.
,,	822	,,	,,			Do.
,,,	21	,,,	"			Do.
Sept. 13	67	N.	16		healthy	Do.
Cont.	390	W.H.	22		>>	Do.
Sept. 14.	4	N.	14 cows		,,	Do.
Cont .0	665	A 9-117	out.		1 - 141 -	Do.
Sept. 18.	445	A. &W.			healthy	Do.
,,	10	"	4		,,,	Do. Do.
"	256 246a	"	I 2 cows		"	Do. Do.
, ,,	246b		out.			Do.
"	708		"			Do.
,,	402	"	"			Do.
Sept. 20.	67	Ň.	19		healthy	Do.
Sept. 21.	36	Brmy.	2	133		Do.
20pm 211	30	Ziny.	-		"	

70			Cattle	e and Cone	dition.	
Date of Inspection.	Register No.	Ward.	No, Examined	Udder diseased.	General Condition	Condition of Shed.
1905.		A 9-117			boolthu	Catiafastam
Sept. 21.	99	A. &W.	2 2 cows			Satisfactory. Do.
,,	708	"	out.			Do.
Sept. 22.	544	Ň.	17		healthy	Dirty.
35	553		21		,,	Satisfactory.
"	404	"				No cows kept.
Sept. 26.	351	É.	11		healthy but dirty.	Dirty.
,,,	101	,,	16		healthy	
,,	478	N.E.	17		,,	Do.
,,	520	,,	15		,,	Do.
Sept. 27.	67	N.	19		,,	Do.
Sept. 29.	405	A. &W.	27		healthy	Do.
,,	1014	W.H.	18		but dirty.	Do.
17	96	, ,,	6 cows		healthy	Do.
.,,	206	A. & W.	out.		1 1.1	Do.
Oct. 3.	159	E.H.	25		healthy	Do.
Oct. 4.	67	N.	19		. 22	Do. Do.
Oct. 11. Oct. 12.	67	Druss.	19		"	Do.
Oct. 12.	201	Brmy.	5		"	Do.
,,	559 88	"	5 26	1	others healthy.	Do.
				(mammitis)		
Oct. 18.	67	N.	19		healthy	Do. Do.
Oct. 20.	896	"	8 28		others	Do. Do.
Oct. 23.	343	"	10.39	I	healthy.	Do.
Oct. 25.	67 88	N.W.	19			Do.
	343	N.	I	1	,,,	Do.
Oct. 30.	343	,,	I	I (tuber-		Do.
Nov. 2.	321	Hdy.	14	culous)	healthy	Do.
,,	812	,,	8		,,	Do.
Nov. 4.	67	Ň.	19		,,	Do.
Nov. 7.	78	Brmy.	19		,,	Do.
Nov. 9	67	N.	19	***	,,	Do.
Nov. 13.	553	Bnk.	12		others healthy.	Do.
Nov. 14.	909	Brmy.	14		healthy	Fairly clean.
Nov. 15.	35	W.	I I		,,	Satisfactory.
"	167	Hdy.	14		"	Do. Do.
Nov. 16.	389	N.W.	II	***	"	Dilapidated.
	154		15		"	Satisfactory.
23	164	Hdy.	3		"	Do.
"	30		2		"	Do.
,,	67	Ň.	21		"	Do.
Nov. 17.	550	Hdy.	11		"	Do.

TABLE VIII a .- Continued.

			Cattle	and Cond	ition.	
Date of Inspection.	Register No.	Ward.	No.	Udder	General	Condition of Shed.
	2.01			diseased.	Condition	
1905.						
Nov. 17.	384	Hdy.	12		healthy	
,,	47	,,	6		,,	Do.
,,	714	"	8		others	Do.
,,	75	"	13	I (mammitis)	healthy.	Do.
,,	338	N.	3		healthy	Do.
,,	833	,,,	26		,,	Do.
,,,	94	Hdy.	8		"	Do.
Nov. 20.	557	E.H.	5		healthy	Do.
"	226	,,,	2		but dirty.	Dirty.
37	835	S.	10		healthy	Satisfactory.
"	353	Hdy.	17	***	"	Do. Do.
,,	676	,,	7		"	Do.
Nov. 21.	75 378	W. H.	9	I	healthy	Satisfactory, but gas lime
1407. 21.	3/0	**. 11.	9		nearing	used for bedding.
	663	E.H.	19			Satisfactory.
,,	85	Hol.	7		"	Do.
Nov. 22.	67	N.	21		",	Do.
Nov. 23.	173	Brmy.	10		,,	Do.
,,	801	,,,	6		,,	Do.
,,	77	,,	34		,,	Do.
,,	635	,,	8		,,	Do.
,,	400	,,	27	1	others healthy.	Do.
,,	73		22	(mammitis)	healthy	Do.
i ",	249	"	7		,,	Do.
,,	92	,,	T		,,	Do.
,,	406	,,	7		,,	Do.
,,	107	,,	4		,,	Do.
,,	765	,,	II		,,	Do.
,,	398	,,	15		,,	Do.
,,	318	,,	24	***	,,	Do.
,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	141	, ,,	19		,,	Do.
Nov. 24.	708	A.& W			,,	Do.
,,	402	"	10		,,	Do.
"	2461		6		"	Do. Do.
Nov. 27.	246A		10	***	"	170.
Nov. 27.	75	Hdy. E.H.	1 25	1	healthy	Satisfactory, but water
1101. 20.	493	13.11.	35	***	neartify	supply not yet improved.
1	204		6			Dirty and dark.
Nov. 30.	430	Brmy.			"	Satisfactory.
,,,	67	N.	21		1000	Do.
Dec. 1.	199	Brmy.		I	others healthy.	Do.
				(mammitis	healthy	Do.
"	127	"	25			Do.
,,,	, 120	,,,	7		"	. 170.

			Cattle	e and Conc	lition.	
Date of Inspection.	Register No.	Ward.	No. Examined	Udder diseased.	General Condition	Condition of Shed.
1905. Dec. 1.	515	Brmy.	11		healthy	Satisfactory,
,,	120	,,	22		,,	Do.
,,,	126	,,	14		,,	Do.
,,	337	,,	23		,,	Do.
,,	310	,,,	5		,,	Do. but rather dark.
,,	227	,,	8		,,	Do.
Dec. 5	. 90	E.	26		. 22	Do.
,,	172	,,	30		healthy except one.	Do.
,,	329	,,	14		healthy	Do.
,,,,,	1011	,,	6		,,	Do.
Dec. 6		Brmy.	5		,,	Do.
,,	243	12	9		,,	Do.
,,	682	,,	20		,,	Do.
,,	822	,,	9		,,	Do.
,,	2 I	,,	II		,,	Do.
,,	70	"	5		,,	Do.
,,,	410	"	7		,,	Do.
,,	147	,,	6		.,,	Do.
, "	887	,,	5		, 22	Do.
,,	408	,,	3		"	Do.
,,	574	,,	14		,,,	Do.
,,	148	"	10		21	Dirty walls.
D 11	310	,,,	5		"	Satisfactory.
Dec. 8.		"	8		others	Walls dirty.
"	352	,,	6	I mammitis	healthy.	Satisfactory.
,,	372	A.& W.	3		healthy	Do.
,,	445	N.	17		,,	Do.
,,	67	,,	21		,,	Satisfactory, but ventila-
						tion requires improvem't.
Dec. 1	1 49	N.W.	22		,,	Satisfactory.
,,	213	Hdy.	5		,,	Do.
,,	644	,,	3		,,	Do.
,,	196	N.	2 I		13	Do.
Dec. 1	3. 321	Hdy.	1	I		Do.
,,	195	N.	19	(mammitis	healthy	One shed satisfactory,
						other requires recon-
Dec. 2	1. 67	",,	21		,,	structing. Satisfactory. Ventilation improved.
Dec. 2:	2. 543		24		,,	Satisfactory.
Dec. 20		Č.	8		,,	Clean (see August 18).
The state of the state of	151		2		,,	Satisfactory.
,,,	7	, "	5		,,	Do.
,,	66	"	12		,,	Do.
,,	67	Ň.	21		"	Do.
,,	-1				,,,	

Cowshed No. 41 has been a recurring nuisance. In the report for 1903, page 152, it was stated that some alterations were being made with a view of improving it. In 1904, page 137, it was pointed out that a new window had been put in, the stands repaired, and the wooden obstructions at the head of the cows removed; it was added: "the shed is not an ideal one even yet, and the occupier's notions of cleanliness are somewhat primitive." I am happy to think that this is probably the last occasion I shall have to report on this shed to you, as it has now been closed in consequence of the pressure we have put upon the owner.

Farm No. 19 was reported upon not only on the 3rd of January, but also again on the 16th of August. The owner was written to and the cowshed has since been visited and found in a satisfactory condition.

On February 21st the sheds at farm 360 required cleaning. They were found clean on the inspector's visit on June 8th, and also on the Veterinary Surgeon's visit on September 12th. The occupier, however, requires looking after, as his shed was reported for want of ventilation in the previous year.

On March 1st, cowshed No. 153 was found badly ventilated. The occupier was written to on the following day and the shed was visited by the ordinary inspector on March 12th, May 2nd, June 15th. The ventilation was better on these occasions, but the shed was overcrowded on May 2nd. On August 18th, Mr. Dixon visited with Mr. Walker and again on December 29th. On the former date he reported the shed as satisfactory as to cleanliness, but structurally bad, and on the latter date again as clean, but referred to the previous report. In 1903 this shed was reported upon and more ventilation was procured by opening a window into the yard. It is referred to on page 153 in the report for that year. The shed is of an old type in the insanitary area and although improved by the new window, really ought to be re-constructed

or demolished. As the whole of this district is intended to be pulled down, it is difficult to get owners to make any reconstructions, and it may be necessary to ask the Committee to require its disuse.

Cowshed 154 was visited on March 1st. It has been a source of trouble to me ever since I came to Leeds. place had, I am told, been personally visited by the Sanitary Committee before my advent, but they resolved to allow it to be continued in spite of official representations to the contrary. A good deal has since been done to get the occupier to keep the place cleaner, but it is situated in quite an improper place for a cowshed. The occupier was a very old man and one scarcely expected him to be able to continue his work so long as he did. He died towards the end of the year, and his friends I fear, looked upon me as responsible for his demise. We treated him as gently as we could consistent with justice to the ratepayers. He was visited by the ordinary inspector on February 20th and March 1st, and received letters on the 21st of September and the 5th of October, seeing me personally after the latter date. Since his death, the cows have been gradually got rid of and are now all gone.

On April 3rd the Veterinary Surgeon reported the shed No. 682 as dirty. A letter was sent to the occupier the same day. The place was visited again on the 10th and 11th of April and on the 2nd and 8th of August, and found in better condition. It was also visited again and found satisfactory by the Veterinary Surgeon on September 12th.

On the same date, April 12th, shed No. 227 was reported as requiring lime-washing. Notice was given, and on the 14th it was found that this had been carried out.

The following week on April 7th, at shed No. 557, the walls were dirty and ventilation bad. Only four cows were kept. It was visited on the 15th and found clean and the ventilation in proper order. It was visited again later in the year and found all right.

On April 12th, at farm 408, which contains two cowsheds, they were found dirty. A letter was written on April 17th. At subsequent visits on the 21st and 29th of June, they were found clean. This was also the case on December 6th when Mr. Dixon visited.

On the same date, April 12th, shed 148 in Bramley was found dirty. The occupier was a new tenant, the shed having been disused for cow-keeping for some years. Mr. Walker had written to the tenant on March 15th as to certain structural alterations, and had visited on the 8th of April about these. On the 12th the place was still dirty, but on the 20th it was in a more satisfactory condition. The cows were out in the fields. On the 6th of December, at Mr. Dixon's first visit, the man had again dirty walls. During the present year it has been visited again, and the conditions found better. It, however, requires to be watched.

On August 11th, farm 45 was visited and found rather dirty. The occupier was spoken to at the time, and was written to on August 14th. At the Veterinary Surgeon's visit on September 6th the farm was satisfactorily clean, and this has been the case at subsequent visits.

On August 29th, farm 390 was found dirty. The cows were out in the field, returning to be milked. The occupier was written to on August 31st. On September 13th, Mr. Dixon again visited and found the shed satisfactory.

On August 31st, Mr. Dixon reported that farm 146 had a deficient water-supply. The owner was written to, and a better supply procured. It has now got Corporation water.

On September 4th, at farm 813, the shed was found dirty, though the cows were out in the field. Letters about cleanliness have been sent to the occupier of this farm on March 27th, April 8th, September 5th, and visits paid January

18th, April 3rd, 5th, 11th, July 27th, and September 4th. The Veterinary Surgeon visited again on the 11th and examined the cows. The occupier has since discontinued cow-keeping, greatly to our satisfaction.

On September 22nd, at farm 544, both sheds were found dirty. The occupier, being a new tenant, was given to understand the requirements of the Authority, both verbally and in writing, and the sheds, during the rest of the year, have been found satisfactory, but he had again to be written to this spring.

On September 26th, cowshed 351 was found dirty. The occupier was written to, and the sheds have since been found in a better condition.

On November 20th, No. 226, a shed with only two cows, was found dirty. At subsequent visits it has been found in better condition.

On November 21st, it was found that at the cowshed at farm 378, gas-lime was being used for bedding. A letter was written, and the practice has been discontinued.

On November 28th, Mr. Dixon reports that at farm 493 the sheds were moderately satisfactory, but the bad water-supply not yet improved. I see nothing for it but to ask the Committee to request the occupier of this farm to discontinue keeping cows. It is impossible to keep them healthy and clean in the present condition of affairs. The matter has been reported to the Committee by Dr. Porter in connection with the water supply.

On the same day Mr. Dixon reported that at farm 204 the shed was dirty and dark. I have had to report about this place before. In 1903 it was reported to you that the occupier had his cows dirty, his sheds dirty, and his drains stopped. In April, 1904, these conditions had been removed.

In October he had again to be written to by Mr. Walker for very dirty sheds. On November 23rd, 1905, the place was again dirty, and in December Mr. Walker wrote asking him to remove the wooden slides from the windows and to substitute glass. This has been done. The place has been somewhat cleaner since.

On December 1st the cowshed at farm 310 was found dark. One of the windows had been partly blocked up, and the remainder were dirty. On December 6th this had been remedied.

On December 8th cowshed 209 was found with dirty walls. The owner was spoken to. At subsequent visits the place was found satisfactory.

On the same day at farm No. 67 the ventilation was unfavourably reported upon. An improvement was made by the removal of the boarding above the cows' heads.

On the 13th of December, Mr. Dixon reported that farm 195 had one shed satisfactory and one that required to be reconstructed. On the 20th, Mr. Dixon and Mr. Walker met the occupier and explained what was required, and the reconstruction has since been carried out to their satisfaction.

Inspection of cattle.—On January 13th an inflamed udder was found in one of the cows at farm 198. The cattle were shortly afterwards sold and the dairy farm given up.

On January 20th a case of mammitis was found at farm 138. Mr. Dixon examined the udder of a cow at this farm on August 11th and September 1st, and came to the conclusion that the disease was old standing suppurative mammitis.

On January 26th another case of mammitis occurred at cowshed 34. All the cows were healthy at this farm on Mr. Dixon's visit on August 16th.

On February 21st a diseased udder was found at farm 99. As the condition was suspicious, the cow was re-visited on March 2nd and May 29th, the cow in the meantime having been taken out of the milk herd and the milk destroyed. The Corporation paid £4 to the owner in compensation. On April 1st she was slaughtered at the public abattoir and her carcase destroyed.

On the same day at farm 337 one cow out of a herd of twenty-one was found to have mammitis; the others were healthy.

On March 16th, of 26 cows at farm 146 two were suffering from mammitis. This farm was frequently visited, and has already been mentioned on the question of water supply.

On March 29th, at the dirty farm 813, a case of mammitis occurred, and Mr. Bowman remarked that "the other 25 cows were healthy but dirty." The cow was isolated for a few weeks and then sent away.

On April 3rd, at shed 110, one of four cows had a tuberculous udder. A few days later an auction sale occurred at the farm, and we were unable to ascertain what became of the cow. She did not reappear in Leeds.

On April 7th, at farm 390, one of 19 cows was suffering from mammitis. This cow was also sent away, and we did not hear what became of her. The farm has already been referred to in the previous section.

On May 17th a cow suffering from mammitis was found at shed 321. She was isolated from the herd and sold a few weeks later. We heard nothing more about her.

On July 1st, at farm 909, Bramley, one of two cows had a diseased udder suspected to be tuberculous. The cow was isolated and Mr. Bowman's suspicions confirmed on July 7th, and on August 4th Mr. Dixon certified tuberculous mammitis. In this case also the owner got rid of the cow, on which account I refused to certify for loss of milk.

On July 26th, at farm 136, a case of mammitis was found in one of 14 cows. On re-examining on August 11th the cow had recovered.

On September 4th, of 30 cows at farm 431, the udder was suspicious of tubercle in 1. The cow was isolated and seen again on September 11th, when the suspicion of tubercle was confirmed. The cow was sold and we lost sight of her.

On October 17th mammitis was diagnosed in one of 36 cows at shed 88. She was re-visited on the 26th, when the inflammation had subsided.

On October 23rd, at farm 343, one of 28 cows had diseased udder. On a suspicion of tubercle the cow was isolated and seen again on the 26th. Mr. Dixon considered that the animal had lung tubercle, and that the udder was also tuberculous. He offered to test with tuberculin. The owner preferred a consultation. Mr. Dixon met his veterinary surgeon on October 30th, and the two agreed that the animal was tuberculous. She was afterwards sold and we lost sight of her. No compensation of course was made.

On November 17th, at farm 75, one of 13 cows had an inflamed udder. She was visited again on November 20th and November 27th. The animal lost the quarter.

On November 23rd one of 27 cows at farm 400 had an inflamed udder. She had been purchased under a warranty, and was sent back to the vendor.

On December 1st, at farm 199, one of 15 cows was found with an inflamed udder. The milk was kept out of the supply, and the animal, which was quite fat, was sent to the butcher.

On December 8th, at farm 352, one of six cows had an inflamed udder. She was fat, and went to the butcher.

On December 13th, at owner's request, a cow suffering from mammitis at shed 321 was examined. The milk was kept out of the supply, and the cow recovered. Milk. 111

It will therefore be seen that 22 out of 3,975 animals examined were found to have some disease of the udder. In 4 of these the disease turned out definitely to be tuberculosis; in 1 tubercle was suspected, but the animal was removed too soon for certainty; in the remaining 17 the disease was probably in most cases mammitis.

Relying upon the liberality of the Committee, I have advised Mr. Dixon to persuade cow-keepers to keep all animals suffering from suppurative mammitis out of the milking herd until the udder is well, and have advised compensation where the owner was evidently not at fault. Compensation was given during the year to one farmer to the extent of £4 4s.

Bacteriological examination of milk.—In addition to the table (VIIIb.) given for the first time last year showing the result in samples of milk tested at the University laboratory for tuberculosis, I present you this year with a further series of observations showing the bacteriological contents of various samples of milk. The table (VIIIc.) is carried out beyond the end of the year, but some of the following remarks apply only to the part concerned in the year 1905. The table shows the part of the town in which the various samples of milk were taken. The letter "a" following the name means that the milk was cooled, generally speaking at the dairy of the Pure Milk Supply Association, before being sent out. In some of these cases the milk was that of the same morning on which the examination was made; in others it had been collected on the afternoon or the morning of the previous day.

The letter "c" indicates that the sample was taken either from the can or the milk cart, generally speaking in the street, but in a good many cases the milk so indicated was sampled at Potternewton dairy from the milk sent from the farm in Harehills Lane to be cooled at the dairy at Potternewton Park. In all these latter cases the letter "H," instead of the

number of the milkman, signifies that the milk was from the same farm as that cooled and sent out by the Association. The letter "d" shows that the milk was purchased from a dairy; "f" indicates that the sample of milk was taken at the farm or immediately on leaving the farm. The letter "h" indicates that it was procured in a house, generally the house of a milk seller; "s" indicates that the milk was taken from the counter of a shop; the letter "t" shows that the milk was taken directly from the teat of a cow; "t.j" indicates the first milk drawn from the teat; "t2" that it was the later milk from the teat.

The reasons for taking the samples from the same farm milked at the same time (1) from the teat of the cow, (2) from the can sent to the dairy, (3) after the milk had been over the cooler, was in order to find out if possible whether any contamination resulted in the process of cooling. In all these cases the milk, if not cooled before being bottled, was cooled immediately afterwards, the bottle being placed in ice; but in a few samples during the winter, the milk after having been thoroughly cooled was not kept in ice up to the time of examination. Usually, however, the Association milk had been kept cool up to the time of examination.

The observations were made by several different men. Those to which a dagger (†) is prefixed were made at the Pathological Department of the University; those with an asterisk (*) in our own laboratory at the Municipal Buildings. In the latter case the examination was generally made as soon as the milk reached the laboratory, arrangements having been made to carry out the examination either morning or afternoon on receipt of the milk. In the former case the examinations were generally made in the forenoon, and where the milk was collected the previous day the conditions of temperature at the Philosophical Hall have been entered for both days. It would have been interesting to add the conditions as to rainfall

previous to the collection of the samples; but to do so would have involved too large a table.

It may perhaps be necessary for the sake of those not familiar with the process to indicate briefly the way in which the counting was done. A small measured quantity of milk was removed from the sample bottle (which had in each case been previously sterilised) by a sterilised pipette. This was diluted with a known quantity of sterilised water and thoroughly mixed. Measured quantities of diluted milk were mixed with sterilised nutrient gelatine, placed in flat covered dishes and exposed to room temperature. The number of colonies in each specimen multiplied by the number of the dilution and fraction of cubic centimetre gives the number of colonies in the last column but one; the results generally being averages of one or two cultures. Each "colony" represents a micro-organism present in the diluted milk under examination. The last column shows the growths which a cubic centimetre of certain dilutions was capable of producing in broths treated with bile-salts. certain classes of bacteria grow in this medium, principally those associated with animal excretions. No attempt was of course made to count these, but according as the characteristic growth resulted from I cubic centimetre of a dilution of I in 10, I in a 100, or I in 1,000, the result was entered as more than 10, more than 100, or more than 1,000 organisms of this class to the cubic centimetre. If no growth occurred with a 1 in 10 dilution, it was entered in the last column as nil.

Similarly in the last column but one a few results are given as <1,000. In these cases the observer having been accustomed to deal with large numbers, had not made a sufficiently strong dilution to prove the absence of bacteria, and <1,000 in these cases means that the dilution of 1 in a 1,000 gave no growth on the nutrient gelatine plate.

Any conclusions from individual cases would be dangerous, for it is always possible in a somewhat dense fluid like milk to take up in a pipette a single particle which, broken up, might afford a large number of colony-producing bacteria in the sample taken. Dealing, however, with large numbers, and restricting ourselves in the meantime to the results obtained during the calendar year, it will be found that of the milks examined directly from the farm, 7 of which were examined on the day of collection, and 3 the following day, the average bacterial content was 30,660. Of the 10 samples bacteria appeared in the taurocholic nutrient broth medium in the strongest dilution in 2, and in the weaker dilutions not at all, the remaining 8 showing no growth. In the 43 samples which had been through the cooler and bottled and, after that, examined, 31 of them the same day and 12 the next day, the average bacterial content was 44,851. In the bile salts ten of them are considered as having more than 10 colonies to the cubic centimetre, I of them as having a 100, the rest none. Comparing these with samples of milk taken from shops and carts in the East part of the town, leaving out certain samples in which an accident had occurred to the gelatine, in the remaining 20 the average number of colonies was 4,039,235 per cubic centimetre. Of the 23 samples in which a result was obtained with the bile-salt broth, 6 showed more than a 1,000, 12 more than a 100, and 5 more than 10. Samples in other parts of the town were taken and showed a varying average, but very much above that of the Association milk. Comparative experiments were also made by keeping the bottled milk in bottle unopened, and under the conditions recommended as to coolness, and also by allowing the bottle to be opened and part of the milk used. To detail all these experiments would take rather too long in this part of the report, but possibly they may be spoken of elsewhere. Experiments of this kind are marked x in the table.

Food and drugs.—The tables in regard to food and drugs are on the same lines as last year.

TABLE VIII b.

Samples of Milk sent to the Bacteriological Laboratory for Examination for Tubercle during 1905.

Date.	Where farm is situate.	No. in register.	Result of the test.
1905.			
Jan. 10.	Guiseley	 368	Pseudo-tuberculosis.
,,	Weeton	 530	Not tuberculous.
,,	Thorner	 14	Tuberculous.
,,	Weeton	 548	Not tuberculous.
Jan. 13.	Skipton	 684	Do.
,,	Elslack	 158	Incomplete.
Jan. 18.	Thornton	 B.	Not tuberculous.
,,	Barnoldswick	 530	Do.
,,	Rawdon	 72	Do.
,,	Do	 30	Incomplete.
Jan. 25.	Horsforth	 812	Not tuberculous.
",	Do	 498	Do.
,,	Do	 936	Do.
Jan. 31.	Rawdon	 268	Do.
,,	Do	 268	Do.
,,	Thornton	 257	Do.
,,	Horsforth	 268	Tuberculous.
Feb. 8.	Barnoldswick	 В.	Not tuberculous.
,,	Rylston	 530	Do.
,,	Do	 530	Do.
,,	Bell Busk	 530	Do.
Feb. 15.	Thornton	 530	Do.
,,	Newbiggin	 240	Tuberculous.
,,	Rylston	 367	Not tuberculous.
Feb. 22.	Darley	 530	Do.
,,	Goldsboro'	 530	Do.
,,	Leeds	 215	Tuberculous.
,,	Do	 530	Not tuberculous.
Feb. 28.	Thorner	 14	Do.
,,	Bardsey	 530	Do.
,,	Leeds	 99	Do.
March 1.	Arthington	 336	Do.
,,	Leeds	 515	Do.
March 8.	Bramhope	 88	Do.
,,	Do	 44	Do.
,,	Do	 529	Do.
,,	Cookridge	 530	Do.
March 15.	Leeds	 290	Do.

TABLE VIII b .- Continued.

Date.	Where farm is situate.	No. in register.	Result of the test.
1905.			
March 15.	Leeds	121	Not tuberculous.
,,	Swillington	303	Do.
,,	Seacroft	747	Do.
March 21.	Leeds	456	Do.
,,	Crossgates	456	Incomplete.
,,	Whinmoor	456	Not tuberculous.
,,	Whitebeck	456	Do.
March 29.	Silsden	74	Do.
,,	Husthorpe	259	Do.
,,	Bingley	245	Do.
,,	Leeds	445	Incomplete.
April 3.	Menston	530	Do.
,,	Burley-in-Wharfedale	530	Do.
,,	Guiseley	579	Not tuberculous.
,,,	Menston	530	Incomplete.
April 12.	Middleton	395	Not tuberculous.
,,	Do	859	Do.
"	Rothwell	450	Tuberculous.
, ,,	Do	239	Not tuberculous.
April 18.	Leeds	290	Do.
	Horsforth	684	Do.
April 26.	Shadwell	421	Do.
. 27	Eccup	335	Incomplete.
April 27.	Bardsey	530	Not tuberculous.
"	Collingham	530	Incomplete.
May 2.	Whinmoor	857	Not tuberculous.
" "	Newsome Green	745	Do.
May 8.	Leeds	110	Do.
"	Scarcroft	113	Incomplete.
"	Ben Rhydding	0.0	Do. Do.
Man, 16	Halton		Not tuberculous.
May 16.	Shropshire		Probably tuberculous.
"	Apperley Bridge Rawdon		
"	Ct 1		Incomplete. Do.
May 24	D. 11 D. 1	0.	Not tuberculous.
May 24.	D 1	00	Incomplete.
,,	Ben Rhydding	530 640	Not tuberculous.
,,	Thornton	504	Tuberculous.
May 31.	Bramhope	643	Incomplete.
	Leeds	321	Do.
,,	Do	725	Do.
",	Do	367	Not tuberculous.
,,,	200	37	

TABLE VIII b .- Continued.

Date.	Where farm is situate. No. in register. Result of the tes		Result of the test.	
1905.				
June 7.	Cookridge		826	Not tuberculous.
,,	Horsforth		812	Incomplete.
,,	Rigton		1072	Not tuberculous.
,,	Bramhope			Incomplete.
June 14.	Apperley Bridge		100	Do.
June 17.	Pool			Do.
,,	Weeton		148	Do.
June 22.	Bramhope			Not tuberculous.
,,	Leeds		2.	Do.
,,	Steeton			Incomplete.
June 28.	Bell Busk			Not tuberculous.
,,	Earby			Do.
	Do			Do.
"	Thornton			Incomplete.
July" 5.	Do		222	Do.
	Pannal			Not tuberculous.
July 12.	Sherburn		1000	Do.
	Long Preston			Pseudo-tuberculosis.
"	Grassington			Incomplete.
"	Horsforth		0 -	Do.
July" 19.	Tanda			Do.
	De			Do.
",	110			Not tuberculous.
,,	De		0	Do.
July 26.	Early.		0	Incomplete.
	Elalask	•••		Do.
"	Consum		1 00	Not tuberculous.
Δυσ ο	Looda	•••	0	
Aug. 2.	De	• • • •		Incomplete. Do.
"	Control of the Contro			Tuberculous.
"	Do Do		0	Incomplete.
A.,	Earby			Do.
Aug. 11.		• • • •	00	Not tuberculous.
"	Ben Rhydding			
"	Grassington		00	Do. Do.
111	Earby		0.	Take the second
Aug. 16.	Menston	•••		Do.
,,	Horsforth			Incomplete.
,,,	Long Preston		00	Do.
Aug. 23.	Sherburn		00	Not tuberculous.
"	Leeds			Incomplete.
"	Elslack		00	Do.
,))	Thornton		00	Do.
Aug. 30.	Leeds		120	Do.

TABLE VIII b .- Continued.

Date.	Where farm is situate.	No. in register.	Result of the test.
1905.		1	
Aug. 30.	Leeds	63	Incomplete.
,,	Do	10.	Do.
,,	Elslack	417	Do.
Sept. 6.	Leeds	6-	Not tuberculous.
,,	Bramhope	-	Incomplete.
,,	Leeds	66	Not tuberculous.
Sept. 20.	Goldsboro'		Incomplete.
,,	Birstwith	200	Do.
	Ben Rhydding	1	Do.
Sept. 27.	Austhorpe		Do.
	Leeds		Do.
,,,	Steeton		Do.
"	Colton	0	Do.
Oct." 5.	Rylston		Not tuberculous.
	Menston		Incomplete.
,,	Leeds		Do.
"	Do		Do.
Oct." 10.	De		Do.
	Association Dailes		Not tuberculous.
"	Dimelon		Incomplete.
,,	T - J-	1 6	Do.
Oct." 18.	D-		Not tuberculous.
Oct. 25.	Danadhan	0	Incomplete.
	Tanda		Do.
,,	De	1	Do.
,,	Ctanles	1 .	Do.
Nov. 1.	Consideration	00	Do.
	De	0	Do.
,,,	Do		Do.
,,,	Cattle		Do.
Nov. 15.	I am a Ducaton		Do.
	Early.		Do.
,,	Dall Duals	00	Do.
,,	Clarkan	00	Do.
Nov. 22.	C-11-1-1	00	Do.
	Hamanata		Do.
,,	Daulou	. 00	Do.
"	Dissessith	00	Do.
Dec. 6.	Commission	0	Do.
	D-		Do.
,,	Da:	100000000000000000000000000000000000000	Do.
"	Do	0	Pseudo-tuberculosis.
,,	Do	100	1 seddo-tuberediosis.

TABLE VIII c.
Bacteriological examination of milk.

				Ten	peratu	re in de	egrees Fahre	enheit.				
	Place of taking sample.			Milk- man.			C	of air.			No. of	Growths
			Time.			Philoso	phical	Hall.	Time and	Of milk	1 c.c. All kinds,	McConkey broth
						Min. previous night.	10 a. m.	4 p.m.	place of sampling.	sampling.	All kinds.	per c.c.
1†	Kirkgate (s)		1905 July 7 10.50 a.m.	21	1	52°	66°	73°			1,460,000	100
2†	Potternewton (d)		10. 0 a.m.	3	9	52°	66°	73°			60,000	100
3†	Sheepscar (d)		10.25 a.m.	5	7	52°	66°	73°			610,000	100
4†	York Street (s)		10.55 a.m.	65	8	52°	66°	73°			1,380,000	1,000
5†	Harehills Lane (d)		July 14 10.40 a.m.	6	7	60°	73°	80°		77°	4,800,000	10
6†	Richmond Road (s)		11.25 a.m.	18	0	60°	73°	80°		71°	2,400,000	1,000
7†	Greenfield Road (s)		11.35 a.m.	6	8	60°	73°	80°		72°	8,000,000	1,000
8†	Moor Road (d)		9.30 a.m.	7	9	60°	73°	80°		79°	960,000	100
9†	Cottage Road (c)		9.45 a.m.	5	0	60°	73°	80°		81°	1,440,000	100
10†	Roundhay (f)		July 28 6.55 a.m.	6	7	50°	65°	67°	60°	81°	110,000	10
11†	Roundhay (f)		6.55 a.m.	6	7	50°	65°	67°	60°	80°	550,000	10
12†	Potternewton (h)		10.15 a.m.	9	1	50°	65°	67°	71°	72°	850,000	1,000
13†	Chapeltown (h)		10. 5 a.m.	18	5	50°	65°	67°	67°	71°	1,640,000	1,000
14†	Harehills Lane (t j)	{	Aug. 3 3. 0 p.m. Aug. 4	} +	ı. {	54° 53°	56° 63°	59° 66°	60°	89°	} 7,500	nil
15†	Potternewton (a)	{	Aug. 3 3. 0 p.m. Aug. 4	} +	I. {	54° 53°	56° 63°	59° 66°	60°	::	} 5,000	nil
16†	Harehills Lane (f)	{	Aug. 3 3. 0 p.m. Aug. 4	}	ı. {	54° 53°	56° 63°	59° 66°	60°	89°	} 3,333	nil
17†	Richmond Street (h)	{	Aug. 3 4.15 p.m. Aug. 4	} 56	4 {	54° 53°	56° 63°	59° 66°	63°	59°	}12,160,000	1,000
18†	Potternewton (x)	{	Aug. 8 p.m. Aug. 11 a.m.	}	1.	54° 52°	66° 58°	67° 62°			29,600,000	1,000
19†	Potternewton (a)		a.m.	ŀ	1.	52°	58°	62°			23,000	nil
20†	Harehills Lane (t j)	{	Aug. 10 3. 0 p.m. Aug. 11 a.m.	} 1	1.	51° 52°	62° 58°	63° 62°			3,300	nil
21†	Dyer Street (s)		11. 5 a.m.	5	3	52°	58°	62°	58°	54°	820,000	1,000
22†	High Street (s)		11.15 a.m.	25	9	52°	58°	62°	57°	58°	90,000	100
23†	Potternewton (a)	{	Aug 17 3. 0 p.m. Aug. 18 9. 0 a.m.	}	ł	49° 57°	58°	70° 64°			410,000	nil

TABLE VIII c .- Continued.

_				Ten	peratu	re in de	grees Fahre	enheit.		
					0	f air.			No. of	Growth
	Place of taking sample.	Time.	Milk- man.	Philosophical Hall.				Of milk	colonies in 1 c c.	McConhi
				Min. previous night.	10 a m.	4 p.m.	Time and place of sampling.	at time of sampling.	All kinds.	per c.c.
24†	Potternewton (a)	Aug. 18 7. 0 a.m.	H.	57°	58°	64°			< 10,000	nil
25†	Potternewton (d)	7.30 a.m.	39	57°	58°	64°	58°	54°	3,720,000	1,0000
26†	Metcalfe Street (h)	10.20 a.m.	321	57°	58°	64°	58°	55°	3,150,000	1,0000
27†	Park Lane (d)	Aug. 26 9.30 a.m.	2	56°	58°	61°	64°	70°	153,000	1000
28†	Harper Street (s)	9.45 a.m.	992	56°	58°	61°	62°	70°	1,950,000	1000
29†	Harchills Lane (t)	5.45 a.m.	H.	56°	58°	61°	60°	(100°)	10,000	nil
30†	Potternewton (a)	7.15 a.m.	H.	56°	58°	61°	62°	56°	116,000	nil
31†	Harehills Lane (t)	Sep. 1 5.45 a.m.	Н.	54°	58°	60°	58°	(100°)	< 10,000	nil I
32†	Potternewton (a)	6.0 a.m.	н.	54°	58°	60°	62°	56°	143,000	nil
33†	Hunslet Road (s)	10.15 a.m.	321	54°	58°	60°	65°	62°	133,000	1,0000
34†	Hunslet (c)	10.40 a.m.	458	54°	58°	60°	62°	75°	1,093,000	100
35†	Harehills Lane (t)	Sep. 8 5.45 a.m.	H.	51°	60°	61°	60°	(100°)	10,000	nil
36†	Potternewton (a)	7. 0 a.m.	H.	51°	60°	61°	62°	50°	23,000	nil
37†	Harehills Lane (c)	6.55 a.m.	1,120	51°	60°	61°	56°	78°	105,000	100
38†	Dyer Street (s)	10.15 a.m.	53	51°	60°	61°	63°	74°	90,000	1000
39†	Harehills Lane (t)	Sep. 15 5.35 a.m.	Н.	41°	53°	57°	63°	(100°)	none developed	nil
40†	Harehills Lane (c)	6.45 a.m.	137	41°	53°	57°	65°	72°	10,000	1000
41†	Potternewton (a)	7. 0 a.m.	H.	41°	53°	57°	58°	50°	10,000	nil
42†	Wheeler Street (s)	10.40 a.m.	87	41°	53°	57°	60°	56°	none developed	100
43*	Potternewton (a)	Sep. 19 6.50 a.m.	Н.	52°	56°	60°	54°	49°	9,000	nil
44*	Harehills Lane (t)	2.20 p.m.	H.	52°	56°	60°	63°		< 1,000	nil
45*	Potternewton (a)	2.45 p.m.	Н.	52°	56°	60°	56°	50°	6,000	nil
46*	Quarry Hill (s)	10.40 a.m.	301	52°	56°	60°	63°	72°	3,000	nil
47*	Kirkgate Depot	Sep. 19 3. 0 p.m. Sept. 20 10. 0 a.m.	} H. {	52° 51°	56° 59°	60°	 58°	 46°	11,000	nil
48*	Kirkgate (s)	Sep. 20 10.10 a.m.	211	51°	59°	60°	58°	73°	1,255,000	100
49†	Harehills Lane (t)	Sep. 22 5.15 a.m.	H.	49°	57°	59°	63°		, 10,000	10
50†	Potternewton (a)	6. 0 a.m.	н.	49°	57°	59°	53°	50°	85,000	10 13
					- 1					

TABLE VIII c .- Continued.

				X 611	Temperature in degrees Fahrenheit.						
			24:11		0	f air.	No. of colonies in	Growths in			
	Place of taking sample.	Time.	Milk- man.	Philosophical Hall. Time and				Of milk	1 c.c.	McConkey broth	
				Min. previous night.	10 a. m.	4 p.m.	place of sampling.	sampling.	ZII KIIIVISI	per c.c.	
51†	Richmond Road (h) .	. 10.30 a.m.	696	49°	57°	59°	62°	78°	42,000	100	
52†	Wheeler Street (s) .	a.m.	583	49°	57°	59°	65°	69°	430,000	1,000	
53*	Potternewton (f.c) .	Sep. 25 3. 0 p.m.	H.	49°	55°	56°	57°	92°	2,000	nil	
54*	Potternewton (a) .	3.10 p.m.	H.	49°	55°	56°	57°	54°	14,000	nil	
55*	York Road (a)	Sep. 26 10.55 a.m.	н.	48°	54°	56°	59°	53°	2,800	nil	
56*	Freehold Street (h)	11. 5 a.m.	279	48°	54°	56°	58°	62°	92,500	100	
57†	Potternewton (x)	Sep. 27 3.30 p.m. Sep. 29	} H. {	49°	54°	58°	58°	50°	23,300	10	
	,	a.m.) (52°	55°	55°		**)		
58†	Harehills Lane (x) .	Sep. 29 6.30 a.m.	H.	52°	55°	55°	59°	72°	40,000	10	
59†	Potternewton (a) .	7.15 a.m.	Н.	52°	55°	55°	54°	48°	16,600	nil	
60†	Hunslet (d)	10. 0 a.m.	240	52°	55°	55°	56°	53°	2,570,000	100	
61*	Potternewton (f.c) .	Oct. 2 2.50 p.m.	н.	43°	48°	54°	51°	92°	700	nil	
62*	Potternewton (a) .	3.10 p.m.	H.	43°	48°	54°	51°	56°	3,600	nil	
63*	Duke Street (s)	Oct. 3 11.35 a.m.	229	40°	49°	51°	57°	59°	11,000	100	
64*	Potternewton (a)	Oct. 2 2.50 p.m. Oct. 3	} H. {	43°	48°	54°	51°	56°	10,800	nil	
		11.57 a.m.) (40°	49°	51°	59°	59°) 10,000		
65†	Potternewton (a) .	Oct. 6 7.15 a.m.	н.	38°	47°	50°	44°	16°	complete liquefaction	100	
66†	Cross Green Lane (x)	Oct. 5 7. 0 a.m. Oct. 6 9. 0 a.m.	} H. {	38°	47°	50°			complete liquefaction	nil	
67†	Spring Close Street (s) .	Oct. 6 9.25 a.m.	255	38°	47°	50°	60°	63°	complete liquefaction	10	
68†	Cross Green Lane (s) .		375	38°	47°	50°	54°	68°	complete liquefaction	100	
69*	Potternewton (a.j) .	Oct. 9 6.50 a.m.	н.	48°	56°	62°	52°	50°	24,000	nil	
70*	Potternewton (a.ij) .	7.35 a.m.	H.	48°	56°	62°	52°	52°	3,600	nil	
71*	Potternewton (x)	Oct. 9 7.15 a.m. Oct. 10 10.15 a.m.	} H. {	48° 47°	56° 52°	62° 57°	52° 59°	50°	4,500,000	1,000	

Food inspection.

TABLE VIII c .- Continued.

				Tem	peratu	e in de	grees Fahre	nheit.		
					0	f air.			No. of	Growthi
	Place of taking sample.	Time.	Milk- man.	Philoso	phical	Hall.	Time and	Of milk at time of	colonies in 1 c c. All kinds.	McConki broth
				Min. previous night.	10 a m.	4 p.m.	place of sampling.	sampling.	All killos.	per c.cc
	(Oct. 9) (1	
72*	N. E. Station (x)	9.15 a.m. Oct. 10	1,003	48°	56°	62°	55°	60°	11,500,000	1,0000
		10.25 a.m. Oct. 13) (47°	52°	57°	59°	60°)	
73†	Harehills Lane (f.x)	Carl Philasens	Н.	44°	47°	48°	58°	(100°)	180,000	nil
74†	Harehills Lane (f.c)	6.30 a.m.	Н.	44°	47°	48°	58°		6,660,000	100
75†	Potternewton (a)	7.15 a.m.	H.	44°	47°	48°	54°	52°	91,300	nil
76†	Hunslet (c)	Oct. 16	216	44°	47°	48°	50°	56°	16,000,000	1,0000
77*	Potternewton (a.x)	6.50 a.m. 7.15 a.m.	} н.	35°	41°	46°	40°	49°	14,000	100
78*	Midland Station	3.40 p.m.	1,013	35°	41°	46°	47°	78°	6,000	nil
79*	York Road (x), v. 77	Oct. 16 6.50 a.m.) H.	35°	41°	46°	40°	49°)	
19-	York Road (x), v. 77	Oct. 17 10.45 a.m.	J)	33°	40°	47°	58°	57°	J 240,000	nil
	(Oct. 16) (1	
80*	Midland Station (x),v.78	3.40 p.m. Oct. 17	1,013	35°	41°	46°	47°	78°	42,500	100
		10.45 a.m.)	33°	40°	47°	58°	57°	,	
81†	Potternewton (a)	Oct. 19 3.15 p.m.	Н. {	39°	43°	45°	46°	46°	48,500	nil
		Oct. 20 Oct. 19		36°	42°	47°			,	
82†	Potternewton (d.x)	3.15p.m. Oct. 20	} н. {	39° 36°	43° 42°	45° 47°	46°	87°	33,000	nil
	,	Oct. 20	,	30	42	**		**	,	
83†	Richmond Road (s)	9.40 a.m.	180	36°	42°	47°	54°	62°	25,000,000	1000
84†	Wentworth Terrace (h)	Oct. 20 9,50 a.m.	63	36°	42°	47°	54°	60°	4,920,000	100
	Went worth restant (ii) 11	Oct. 23		00	4.2	31		00	1,020,000	10
85*	Potternewton (a.x)	6 40 a.m.	H.	36°	43°	47°	41°	46"	5,000	nil
86*	Mid. Station	4.15 p m.	1.009	36°	43°	47°	47°	72°	111,500	1000
074	Pottomo to (Oct. 23 6.40 a.m.) (36°	43°	47°	41°	46°	1	
87*	Potternewton (a.x) v. 85	Oct. 24 10.15 a.m.) H. {	36°	44°	47°	66°	66°	\$02,000	1000
	(Oct. 23) (1		1	
88*	Mid. Station (x). v. 86	4.15 p.m. Oct. 24	1,009	36°	43°	47°	47°	72°	2,050,000	1,000
1		10.15 p.m.	,	36°	44°	47°	66°	66		
89†	Harehills Lane (t.j)	Oct. 26 2.15p.m.	} н. {	39°	45°	47°	57°	.,	< 1,000	nil
		Oct. 27	,	44°	47°	52°			,	1
90†	Harehills Lane (t.ij)	Oct. 26 2.55 p.m.	} н. {	39°	45°	470	58°		{[1,000	nil
		Oct. 27		44°	47°	52°				

TABLE VIII c .- Continued.

-										
				Ten	peratu	re in de	grees Fahre	enheit.		
			Milk-		0	f air.			No. of colonies in	Growths
	Place of taking sample.	Time.	man.	Philoso	phical	Hall.	Time and	Of milk at time of	1 c c. All kinds	McConkey broth
				Min. previous night.	10 a m.	4 p.m.	place of sampling.	sampling.	111 111103	per c.c.
91†	Potternewton (a) {	Oct. 26] 3.10 p.m. Oct. 27	} н. {	39° 44°	45° 47°	47° 52°	49°	45°	} 85,000	nil
92†	Bridge Street (s)	Oct. 26 3.45p.m. Oct. 27	} 953 {	39° 44°	45° 47°	47° 52°	60°	53°	330,000	10
93*	Potternewton (a)	Oct. 30 6.45 a.m.	Н.	41°	48°	51°	49°	48°	93,000	10
94*	Park Square (x)	Oct. 30 even. Oct. 31 9.35 a.m.		41° 41°	48° 44°	51° 48°	 54°	 54°	26,600,000	1,000
95*	Kirkgate (x) v. 93	Oct. 30 6.45 a.m. Oct. 31 9.15 a.m.	} H. {	41° 41°	48° 44°	51° 48°	49° 59°	48° 59°	} 1,117,000	10
96†	Potternewton (a)	Nov. 2 3.15 p.m. Nov. 3 morn.	} H. {	46° 43°	49° 46°	51° 49°	54° 44°	48°	10,650	10
97†	Potternewton (a)	Nov. 3 morn.	н.	43°	46°	49°_	44°	46°	10,500	nil
98†	Upper Cross Street (s)	Nov. 3 9.30 a.m.	187	43°	46°	49°	50°	65°	206,000	10
99†	York Street (s)	9.45 a.m.	199	43°	46°	49°	52°	54°	20,000,000	100
100†	Potternewton (a)	Nov. 10 7.15 a.m.	Н.	38°	40°	44°	44°	46°	24,300	10
101†	Elland Road (s)	9.15 a.m.	545	38°	40°	44°	55°	66°	226,700	10
102†	New Princess Street (s)	9.30 a.m.	208	38°	40°	44°	57°	62°	73,300	100
103†	Sweet Street (d) Lower Wortley (c)	Nov. 14	791	38° 40°	40°	44° 45°	48°	60° 72°	3,330,000 15,500	100
105†	Potternewton (a.j)	Nov. 17	н.	34°	37°	39°	38 ⁴	40°	1,766	nil
106†	Potternewton (a.ij)	8. 0 a.m.	H.	34°	37°	39°	38°	40°	700	nil
107†	St. James' Place (s)	9.30 a.m.	14	34°	37°	39°	48°	54°	40,000	10
108†	St. James' Street (c)		74	34°	37°	39°	48°	64°	60,000	10
109*	Harehills Lane (f)	Nov. 22 6.35 a.m.	H.	37°	42°	46°	32°	92°	3,950	nil
110*	Potternewton (a)	6.45 a.m.	H.	37°	42°	46°	32°	42°	7,000	10
111*	Midland Station	4.45 p.m.	1,232	37°	42°	46°	44°	72°	206,000	10
						1 3				

TABLE VIII c .- Continued.

	The state of the s			Tem	peratu	re in de	grees Fahre	enheit.		
					0	f air.	-		No. of	Growth
	Place of taking sample.	Time.	Milk- man.	Philoso	phical	Hall.	Time and	Of milk	colonies in 1 c.c. All kinds.	McConker broth
				Min. previous night.	10 a. m.	4 p.m.	place of sampling.	sampling.	All Killus.	per c.c.
112†	Station, Headingley	Nov. 24 3.25p.m. Nov. 25 morn.	}1,012 {	42° 41°	44° 43°	45° 46°	48°	85°	12,000	10
113†	Station, Headingley {	Nov. 24 3.27p.m. Nov. 25	}1,025 {	42° 41°	44° 43°	45° 46°	48°	76°	{ 16,000	10
114†	Station, Headingley {	Nov. 24 3.30 p.m. Nov. 25	}1,028 {	42° 41°	44° 43°	45° 46°	48°	52°	} 63,000	nil
115†	Potternewton (a)	Nov. 24 3. 0 p.m. Nov. 25 10. 0 a.m.	} H. {	42°	44° 43°	45°	48°	42°	103,000	nil
116*	Potternewton (a) .	Nov. 28 6.45 a.m.	н.	40°	44°	46°	42°	46°	3,000	nil
117*	Midland Station	Nov. 29 10.10 a.m.	1,198	37°	40°	44°	45°	67°	142,000	100
118*	Kirkgate (a.x)	Nov. 28 6.45 a.m. Nov. 29 11.15 a.m.	} H. {	40° 37°	44° 40°	46° 44°	42° 67°	46° 62°	100,000	nil
119†	Station, Headingley	Nov. 30 3.25p.m. Dec. 1	}1,029 {	39°	41° 41°	45° 46°	46°	72°	} 183,000	nil
120†	Station, Headingley	Nov. 30 3.27p.m. Dec. 1	}1,030 {	39°	41° 41°	45° 46°	46°	75°	} 50,000	nil
121†	Midland Station	Nov. 30 4. 0 p.m Dec. 1	}1,031	39°	41° 41°	45° 46°	44°	72°	} 37,000	nil
122†	Potternewton (a)	Nov. 30 4. 0 p.m Dec. 1	. } н	39°	41° 41°	45° 46°	44°	46°	} 19,000	nil
123*	Potternewton (a) .	Dec. 4 7.15 a.m.	Н.	39°	42°	45°	44°	42°	10,250	10
124*	Kirkgate (s)	. 4.30 p.m.	211	39°	420	45°	63°	66°	150,000	100
125*	Kirkgate (v. 123)	Dec. 4 11.30 a.m.	Н.	39°	42°	45°	52°	43°	35,000	10
126†	Moor Rd., Headingley (c)	Dec. 7 4.15 p.m. Dec. 8	} 79	45° 45°	52° 48°		50°	66°	} 59,000	10
127†	Potternewton (a)	Dec. 7 4.10 p.m Dec. 8	} н.	45° 45°	52° 48°			46°	} 13,000	nil
128†	Harehills Lane (f)	Dec. 8 7.15 a.m	Н.	45°	48°	49°	52°	78°	1,189,000	10

TABLE VIII c .- Continued.

				Ten	peratu	re in de	grees Fahre	enheit.		
					C	of air.			No. of colonies in	Growths
	Place of taking sample.	Time.	Milk- man.	Philoso	phical	Hall.	Time and	Of milk at time of	1 c.c.	McConkey broth
				Min. previous night.	10 a.m.	4 p.m.	place of sampling.	sampling.	All kinds.	per c.c.
129†	Potternewton (a)	Dec. 8 7.30 a.m.	н.	45°	48°	49°	52°	48°	11,000	nil
130*	Potternewton (a)	Dec. 11 6.45 7.15	} H.	39°	40°	43°	34°	38°	6,050	nil
131*	Midland Station	. 10.55 a.m.	1033	39°	40°	43°	41°	60°	13,500	nil
132†	Potternewton (d)	Dec. 16 3.45 p.m. Dec 17	} н.	43	45°.	46°	46°	80°	} 7,000	nil
133†	Potternewton (a)	Dec. 16 4.10 p.m. Dec. 18	} н.	43°	45°	46°	46°	48°	} 7,000	nil
134†	Harehills Lane (c)	. 4.55 p.m.	1034	43°	45°	46°	56°	76°	45,000	nil
135†	Gipton (c)	. 4.30 p.m.	137	43°	45°	46°	56°	85°	13,000	nil
136*	Potternewton (c)	Dec. 18 2.40 p.m.	н.	40°	42°	43°	43°	86°	11,000	10
137*	Potternewton (a)	. [3.40 p.m.	Н.	40°	42°	43°	43°	44°	15,000	10
138*	Brownhill Crescent (h) .	. 4.32 p.m.	496	40°	42°	43°	58°	68°	27,500	nil
139†	Potternewton (f.c)	Dec. 18 3.40 p.m. Dec. 19	{ н.	40° 38°	42° 43°	43° 45°	43°	86°	} 5,000	nil
140†	Potternewton (a)	Dec. 18 3.45 p.m. Dec. 19	{ н.	40° 38°	42° 43°	43° 45°	43°	44°	} 31,000	10
141†	Brownhill Crescent (h)	Dec. 18 4.50 p.m. Dec. 19	} 496	40° 38°	42° 43°	43° 45°	58°	68°	} 31500	nil
142†	Potternewton (c) .	1906. Jan. 5 6.50 a.m.	н.	42°	46°	47°	42°	87°	3,000	nil
143†		. 7. 0 a.m.	H.	42°	46°	47°	42°	39°	2,000	nil
144†	Park Lane (d)	. 11.15 a.m.	2	42°	46°	47°	54°	63°	10,000	nil
145*	Potternewton (c) .	. 6.50 a.m.	H.	420	46°	47°	42°	87°	8,600	pil
146*	Potternewton (a) .	. 7. 0 a.m.	Н.	42°	46°	47°	42°	39°	58,200	10
147*	Park Lane (d)	. 11.15 a.m.	2	7 42°1	46°	47°	54°	63°	69,100	10
148*	Potternewton (c)	Jan. 12 7.0 a.m.	H.	₩ 40°	47°	48°	43°	88°	10,300	nil
149*	Potternewton (a) .	. 7.15 a.m.	H.	₹ 40°₽	47°	48°	43°	40°	5,900	nil
150†	Potternewton (c) .	. 7. 0 a.m.	Н.	3 40°7	47°	48°	43°	88°	16,400	nil
151†	Potternewton (a) .	7.15 a.m.	H.	40°F	47°	48°	43°	40°	18,200	nil
152†	Potternewton (c) .	. 7.25 a.m.	H.	400]	47°	48°	43°	40°	8,200	nil

TABLE VIII c .- Continued.

				IADLE			ontinue				
					Tem	peratu	re in de	grees Fahre	mheit.		
						0	f air.			No. of	Grows
	Place of taking sample	e.	Time.	Milk- man.	Philoso	phical	Hall.	Time and	Of milk	I. C. C.	McCord
					Min. previous night.	10 a.m.	4 p.m.	place of sampling.	at time of sampling.	All kinds.	per cor
153†	Park Lane (s)		9.45 a.m.	994	40°	47°	48°	55°	65°	15,700	nidi
154*	Potternewton (c)		Jan. 19 6.45 a.m.	Н.	32°	39°	41°	-32°	86°	6,000	ni
155*	Potternewton (a)		7. 0 a.m.	H.	32°	39°	41°	43°	39°	500	11
156†	Potternewton (a)	{	Jan. 18 3. 0 p.m. Jan. 19 10.30 a.m.	} H. {	37° 32°	40° 39°	42° 41°	44°	42°	28,000	nili
157†	Potternewton (a)		Jan. 19 7. 0 a.m.	67	32°	39°	41°	43°	39°	44,000	nili
158†	Mill Hill (d)		10. 0 a.m.	776	32°	39°	41°	45°	63°	1,600,000	100
159†	Midland Station		10.20 a.m.	1,018	32°	39°	41°	42°	68°	610,000	100
160†	Potternewton (a)		Jan. 26 7. 0 a.m.	Н.	43°	50°	51°	47°	42°	10,000	nilii
161†	Potternewton (d)		8.10 a.m.	39 -	43°	50°	51°	88°	80°	145,000	100
162†	Ellerby Terrace (d)		10.20 a.m.	34	43°	50°	51°	63°	68°	67,000	10
163†	Easy Road (h)		10.30 a.m.	917	43°	50°	51°	52°	50°	300 000	nili
164†	Potternewton (a)		Feb. 2 7.30 a.m.	Н.	38°	42°	46°	44°	42°	20,000	nili
165†	Richmond Street (s)		10. 0 a.m.	564	38°	42°	46°	59°	68°	1,000,000	nili
166†	Richmond Street (s)		10.10 a.m.	750	38°	42°	46°	59°	72°	151,000	nil
167†	Ellerby Lane (kitchen)		10.20 a.m.	226	38°	42°	46°	59°	60°	137,000	nil
168*	Potternewton (a)		Feb. 7 6.50 a.m.	H.	33°	39°	44°	36°	42°	1,900	nill
169*	Park Lane (s)		10.45 a.m.	994	33°	39°	44°	54°	55°	35,000	10
170†	Potternewton (a)		Feb. 9 6.30 a.m.	Н.	30°	31°	39°	38°	42°	70,000	16
171†	York Street (s)		10.10 a.m.	199	30°	31°	39°	51°	52°	350,000	nili
172†	York Road (s)		9.50 a.m.	413	30°	31°	39°	41°	55°	1,000,000	1,000
173†	York Road (s.g)		10. 5 a.m.	414	30°	31°	39°	48°	38°	92,000	nili
174*	Potternewton (a)		Feb. 15 6.45 a.m.	H.	32°	36°	41°	34°	42°	4,650	nili
175*	Richmond Street (s)		10.55 a.m.	750	32°	36°	41°	55°	60°	6,300	nil
176*	Harehills Lane (t.j)		Feb. 22 2.30 p.m.	Н.	33°	35°	45°	55°	(100°)	< 1,000	nil
177*	Harehills Lane (f)		2.33 p.m.	Н.	33°	35°	45°	55°	89°	4,050	nil
178*	Potternewton (a)		3. 5 p.m.	Н.	33°	35°	45°	42°	42°	4,250	nil

TABLE VIII c .- Continued.

				Теп	peratu	re in de	grees Fahre	enheit.		
			Nem.		0	f air.			No. of colonies in	Growths in
	Place of taking sample.	Time.	Milk- man.	Philoso	phical	Hall.	Time and	Of milk at time of	1 c.c. All kinds.	McConkey broth
				Min. previous night.	10 a.m.	4 p.m.	place of sampling.	sampling.	All Killusi	per c.c.
179*	Harehills Lane (c) .	March 9 3. 0 p.m.	Н.	38°	40°	42°	45°	84°	8,600	nil
180*	Potternewton (x) .	3.15 p.m.	Н.	38°	40°	42°	45°	82°	2,850	nil
181*	Kirkgate Depot (x) .	4.45 p.m.	H.	38°	40°	42°	54°	54°	404,500	10]
182*	Kirkgate Depot	March 15 3. 0 p.m. March 16 10.40 a.m.	} H. {	35° 33°	51° 49°	53° 56°	 43°	 44°	16,650	nil
183*	Kirkgate	March 10.45 a.m.	н.	33°	49°	56°	43°	42°	41,350	10
184*	Kirkgate (s)	. 10.50 a.m.	422	33°	49°	56°	44°	45°	5,700	nil
185†	Potternewton (d.c) .	March 30 7. 0 a.m.	Н.	36°	45°	48°	40°	88°	26,000	nil
186†	Harehills Road (d.c.) .	. 7.20 a.m.	543	36°	45°	48°	44°	83°	43,000	nil
187*	Potternewton (a) .	. 7.20 a.m.	H.	36°	45°	48°	45°	42°	7,500	nil
188*	Potternewton Dairy	March 29 3, 0 p.m. March 30 7.30 a.m.	} H· {	37° 36°	41° 45°	46° 48°	 45°	 44°	63,400	nil
189*	Park Lane (d)	March 30 10.30 a.m.	2	36°	45°	48°	50°	60°	74,200	10
190*	Harehills Lane (t.j) .	April 24 6.15 a.m.	Н.	37°	43°	44°	59°	(100°)	600	nil
191*	Potternewton (a) .	. 6.35 a.m.	H.	37°	43°	44°	48°	46°	2,800	nil
192*	Gipton Wood (a) .	. 6.45 a.m.	543	37°	43°	44°			10,400	10
193*	Potternewton (d.c)	May 2 7. 5 a.m.	Н.	36°	47°	48°	43°	90°	4,000	nil
194*	Potternewton (a)	. 7.15 a.m.	Н.	36°	47°	48°	43°	46°	14,100	nil
195*	Templenewsam (b)	. 7.47 a.m.	1,116	36°	47°	48°	45°	71°	37,200	10
196*	Kirkgate Depot (a)	May 26 10.30 a.m.	н.	48°	53°	56°	50°	48°	2,550	nil
197*	Kirkgate Depot (a)	May 25 3. 0 p.m May 26 10.35 a.m.	I II.	46° 48°	55°	56°	 50°	 44°	3,000	nil
198*	Park Lane (d)	May 26 11.10 a.m.		48°	53°	56°	54°	58°	6,050	100
							100			

Samples of food sent to the City Analyst for examination during the 52 weeks ended 30th December, 1905.

	Article.		Genuine.	Poor in quality.	Adul- terated.	Total.	Sum- moned.	Con- victed.	Dis- missed or With- drawn.
Milk		 	430	70	22	522	20	17	3
Butter		 	20		8	28	8	6	2
Jam		 			4	4			
Lard		 	4			4			
Mace		 	1			I			
Tinned I	Peas	 	I			I			
Vinegar		 	24		4	28	I	I	
Cordial		 			5	5			
Tomato S	Sauce	 			1	I			
Castor C	il	 	2			2			
Seidlitz	Powder	 	3		1	4			
	Total	 	485	70	45	600	29	24	5

TABLE IXa.

Summonses issued during the 52 weeks of 1905 under the Sale of Food and Drugs Acts, 1875, 1879 and 1899*, for articles other than butter.

No. of sample.	Article	Percentage adulteration	of n.		£	ines		Remarks.
4	Milk	 $8\frac{1}{2}\%$ added water			3	0	0	and costs
13	Do.	 7% do.			2	0	0	do.
49	Do.	 23% fat removed				_		withdrawn
50	Do.	 40% do.				_		do.
165	Do.	 17% added water			5	0	0	and costs
166	Do.	 31% do.			3	0	0	
167	Do.	 8½% do,				_		bound for six
237	Do.	 11% do.			2	0	0	and costs
244	Do.	 9% do.			2	0	0	do.
300	Do.	 10% do.			3	0	0	do.
370	Do.	 9% do.			0	10	0	
509	Do.	 17% do.			5	0	0	
510	Do.	 9½% do.				_		to pay costs
515	Do.	 9% do.				_		do.
540	Do.	 3½% do. 12% f	at remo	ved		_		discharged on warranty
543	Do.	 14% fat removed			I	0	0	and costs
571	Do.	 17% added water			I	0	0	- 1
581	Do.	 29% do.				_		to pay costs
593	Do.	 4% do. 6½%	fat remo	ved		_		do.
598	Do.	 32% added water			10	0	0	
	-			£	37	10	0	

^{*} For cases under Margarine Act see table IXb.

TABLE IXb.

Other Summonses taken out under the Margarine Act, 1887, and the Food and Drugs Acts, 1875, 1879 and 1899, during the 52 weeks ended 30th December, 1905.

No. of	Article.			centage ulteration		4	1	Fine	S.	Remarks.
ampie.			au	uiteratioi	le .		£	s.	d.	
160	Butter	86%	foreign	fat			1	0	0	and costs
172	Do	88%	do.					-		to pay costs
250	Do	89%	do.				1	0	0	and costs
311	Do	87%	do.					_		dismissed on invoice
320	Do	85%	do.				1	0	0	and costs
321	Do	86%	do.				0	2	6	and costs
354	Do	88%	do.							dismissed on invoice
596	Do	88%	do.				I	5	0	and costs
440	Vinegar			_			1	0	0	and costs
-	For givin	g fals	e warr:	anty in	writin	g	3	0	0	and costs
_	For refusi	ng to	sell m	ilk to l	Inspecto	or	2	0	0	and costs
						£	10	7	6	

TABLE IXc. Adulterated Samples where no proceedings were taken.

No. of Sample.	Article		Adultera	ation.			Remarks.
2	Milk		48% fat removed				unofficial sample
3 8	Do.		3% water added	***			purchased officially
	Do.		1% do.				do.
10	Do. Do.		5% do.		***		do.
12 16	Do.	***	4% do. 2½% do.				do.
58	Do.		2½% do. 6% fat removed				do.
61	Do.		2% water added				do.
62	Do.		9% fat removed				do.
63	Do.		1% do.	***			do.
71	Do.		11/2% water added		***		do.
78	Do.		4% do.				do.
87 97	Do. Do.		1½% do. 1½% do.		***		do.
104	Do.		$\frac{1\frac{1}{2}\%}{2\frac{1}{2}\%}$ do.				do.
108	Do.		4½% do.				do.
IIO	Do.		1½% do.				do.
III	Do.		3% do.				do.
113	Do.		2% do.				do.
114	Do.		1% do.				do.
116	Do. Do.		4½% do.		***		do. do.
121	Do.		$\frac{3\frac{1}{2}\%}{1\%}$ do.	***			do.
131	Do.	::	3% water added				- do
133	Do.		2% do.				do.
135	Do.		1% do.				do.
169	Do.		3½% fat removed				do.
177	Do.		2% water added	***			do.
192	Do.		1% do.		***		do. do.
203	Do. Do.	***	1½% do. 3% do.			***	do.
207	Do.		3% do. 3% do.				do.
222	Do.		2% do.				do.
227	Do.		3½% fat removed				do.
233	Do.		I% do.				do.
235	Do.		2% do.				do.
238 268	Do. Do.		1½% do.		***		do. do.
273	Do.		3½% water added 1% do.				do.
339	Do.		1% do. 1½% do.				do.
351	Do.		4% do.				do.
374	Do.		4½% do.				do.
377	Do.		4% do.	3% fat	remov	ved	do.
379	Do.		2% water added			***	do.
380 386	Do. Do.		2% do. 4% do.		***	****	do. do.
387	Do.		4% do. 4% do.				do.
388	Do.		$\frac{4}{5\frac{1}{2}}$ % do.	6% fat	remo		do.
393	Do.		4% water added				do.
433	Do.		5% do.				do.
465	Do.		3% do.				do.
477	Do.		3% do.				do. do.
482 484	Do. Do.		3% do. 4% do.	***		***	unofficial sample
487	Do.		4% do. 2% do.				purchased officially
490	Do.		1% fat removed				do.
491	Do.		$4\frac{1}{2}\%$ water added,		remo		do.
494	Do.		4% water added	131			do.
500	Do.		4% fat removed			and in	do.
501	Do.		21% water added,				unofficial sample
505 506	Do. Do.	***	3½% water added do.			***	purchased officially do.
527	Do.		$\frac{3\frac{1}{2}\%}{3\frac{1}{2}\%}$ do.				do.
540	Do.			2% fat			do.
541	Do.		$4\frac{1}{2}\%$ do.				do.
569	Do.		4% fat removed				do.
572	Do.		$3\frac{1}{2}\%$ do.				do.
	Do.		1½% do.		***		do.
575							
575 592 594	Do. Do.		2% water added $3\frac{1}{2}\%$ do.			***	do. do.

K

Meat and markets.—The inspectors paid 5,189 visits to 96 private slaughter-houses, two public, and two knackers' yards. They have also visited the general market, the wholesale meat market, and the cattle market, the shops in the town in which meat, fish and fruit are sold, and two knackers' yards. Their visits to these number 5,059.

Slaughter House and Meat Inspection, 52 weeks ended 30th December, 1905.

Class of meat seized and des- troyed.	Weight in stones of 14 lb.	No. of seizures.	No. of persons sum- moned.	No. of convic- tions.	Penalties.
Beef	63	3	3	2	Imprisonment, 6 weeks. Do. 3 months.

It will be seen from table X that three official seizures were made. Four summonses were issued; two of them, however, on the same person.

A screw butcher who had been previously convicted and sent to prison for an offence at Middlesbrough, and who had since been living in a neighbouring district of the West Riding, had settled in Leeds. On October 25th, the inspectors learnt that there had been slaughtered for him a tuberculous animal, which they had seen in the neighbourhood of the market and on which they were keeping their eyes. The butcher claimed the meat in my presence. It was condemned the following day by the Stipendiary, and afterwards a summons was issued against the butcher as owner of the meat, and another against the occupier of the private slaughter-house in which it had been killed.

Before these summonses, however, were issued, on the 3rd of November we seized the four quarters of another cow, also tuberculous and unfit for food, in the shop of the man who had slaughtered the previous animal for the butcher, and the same day at the shop of the butcher himself, three quarters of another cow also unfit for food. A third summons was therefore issued against the man who had killed the previous animal for having unsound meat upon his premises, and a fourth against the butcher, the owner as is believed of all three carcases. When the summonses were returnable the slaughterer and the slaughter-house keeper in the first case attended. The owner of the meat did not and a warrant was issued. The case against the other men was partly heard, but not completed. Later, the slaughterer of the first animal was sent to prison for six weeks without the option of a fine, for having unsound meat in his shop.

For some time the screw butcher managed to evade the detectives, but he was eventually brought to Leeds from Glasgow, and early this year sentenced to three months' imprisonment without the option of a fine—the maximum penalty. As the slaughter-house keeper appeared to have been merely guilty of carelessness, the summons was withdrawn on the completion of the case against the owner.

In addition to the articles seized officially, and included in table X., 55,254 lbs. of beef, 5,073 of mutton, 7,105 of veal, 4,459 of pork, and 546 of fish, together 72,437 lbs., or 5,174 stones were destroyed by the owners with our consent. English offal, consisting of tripe, livers, plucks, &c., weighing 2,019 lbs. was destroyed. What our inspectors describe as foreign offal, to the extent of 5,333 lbs. was also destroyed. German sausages, foreign lard and other animal matter to the extent of 2,128 lbs., 1,100 eggs, and 24 lbs. of bacon were also destroyed, and vegetables to the weight of 5,908 lbs. There were also destroyed, 2,387 lbs. of fruit.

Ten Sunday morning visits were made. Prospect Row, the resort of hawkers, shops in High Street, St. Peter's Street, York Street, and Marsh Lane in the North-East Ward, and others in the Holbeck and Hunslet districts were visited. Occasionally a piece of frozen meat which had been kept rather too long was found and the owner allowed to destroy it. But on the whole the meat examined was not exactly unfit for food.

The meat destroyed privately includes the carcases and organs of three animals affected with anthrax. One case occurred in March, at Beeston, within the borough. The animal had been bought at the Otley auction some weeks earlier and it is uncertain how it became infected with the disease. There was no attempt to sell the carcase, which was destroyed by us.

In August a case occurred at Stanningley. We destroyed the carcase and organs and the Contagious Diseases officials took charge of the disinfection of the premises.

TABLE XI. Smoke, 1905 (52 weeks).

Complaints received					25
Furnaces inspected			1		8,305
Observations taken of c					
Total number of minutes	dense sn	noke			3,026
Average minutes durat each observation of					onds)
Smoke prevention applia	nces adap	ted to fu	rnaces		83
Chimneys newly erected					11
Furnaces in connection v	with new o	chimneys			43
Notices served upon mar	nufacturer	s			7
Do. do. stol	cers				2 I
Persons summoned before	re the mag	gistrates			2
Do. convicted					2
Total amount in fines		£1	os.	od.	
Costs			8s.	od.	

A third case occurred in October, this time outside the borough. The carcase and organs had been consigned to a merchant in the Leeds wholesale meat market. He drew the inspector's attention to them. They were at once sent to the destructor and the West Riding Constabulary informed of the facts. A fine of £5 and costs was obtained by the County police from the owner for not reporting.

REMOVAL AND DISINFECTING WORK.

Removals.—In addition to the work shown in tables XII. and XIII., our staff also removed 353 contacts to Manston Hall Cottages, Seacroft. They also took five persons to the Gildersome Convalescent Home, four from the town and one from Manston Hall. They took one from Manston Hall to the General Infirmary and one to the Women and Children's hospital, one from the Killingbeck hospital to the Bramley Union Workhouse, one from home to the Leeds Union Workhouse, and one home from the General Infirmary. A case of smallpox was removed from Halton to the Waterloo Cottage Hospital for the District Authority, making a total of 363 removals. There were also 41 other ambulance journeys where removal did not take place.

Disinfection.—It is not mentioned in table XII. that 489 persons went or were taken to one of our disinfecting stations. Each had a bath and his or her clothes disinfected.

The work of the disinfecting staff included some disinfections on account of phthisis, which it may be interesting to separate from the details given in table XII. The houses disinfected on account of this disease were 186, whilst 399 rooms were stoved with sulphur, and in 2 the paper was stripped. Beds and mattresses to the number of 514, articles of bed clothing to the number of 919, articles of wearing apparel and miscellaneous articles to the number of 493 and 306 respectively were removed and disinfected at the station.

TABLE XII.

Work done by Disinfecting Staff, 1905 (52 weeks).

Houses disinfected			2,362	
Rooms disinfected (stripped 310, limewa	ashed	278)	5,900	
Beds and mattresses disinfected			11,001	
Articles of bed clothing disinfected			18,623	
Articles of wearing apparel disinfected			52,608	
Miscellaneous articles disinfected			43,853	

TABLE XIII.

Cases removed to hospital by our own staff.

Classified according to diseases certified

Small- pox.	Scarlet fever.	Diph- theria.	Typhus fever.	Typhoid fever.	Other diseases.	Total, 1905.
132	730	165	I	140	41	1,209

(52 weeks).

TABLE XIV.

Return for the 52 weeks ended 30th December, 1905, of patients in hospital.

	Small-pox	Scarlet Rever.	Diphtheria. co	Typhus 4 fever.	Enteric, or typhoid on fever	Other or doubtful ocases.	Total. 2
No. in Hospital on Saturday, 31st December, 1904	7	125	15		19	13	179
No. since admitted	126	720	153		185	207	1,391
No. discharged	130	721	135		154	182	1,322
No. died	3	29	14		28	12	86
No. remaining in Hospital, 30th December, 1905		95	19		22	26	162

We are now well off for disinfecting stations. We have a new one at Kidacre Street with a large and small steam chest of the Lyon's type, and a drying chamber. There are also two bath rooms and two waiting rooms. The whole is worked by steam from the Kidacre Street destructor.

We have at our newly re-constructed station at Burmantofts two old but efficient steam disinfectors,—a Warner which
we have had some years, and a Washington Lyons, which was
removed from the Beckett Street Hospital when the latter was
pulled down. We have convenience for drying the clothes
which have been through the steam chest, and two good bathrooms, with waiting-rooms attached, to both sides. We have
also a laundry divided into two parts, one on each side of the
disinfecting station,—a receiving place for the steeping and
preliminary washing of soiled linen before disinfection, and a
drying, ironing, and getting-up room for the treatment of the
linen which has been through the preliminary washing and then
through the disinfector.

Flushing.—During the 52 weeks of the year, 9 carts, each with 2 attendants, have been employed flushing drains. In this period 57,461 house drains, 34,092 water closets, and 36,234 gullies have been flushed, chiefly on account of illness. The above-mentioned figures include the flushing of drains in connection with 334 schools. In addition to this, 2 men with a horse and cart have been engaged putting an iron solution into tanks connected with the sewers, and in this manner 18,630 gallons of disinfecting solution have been allowed to trickle into certain of the sewers. This last shows a considerable increase of this part of our work in the year.

CANAL BOATS AND TEMPORARY DWELLINGS.

It is not necessary to deal in detail with the whole of the matters contained in tables XV., XVI., and XVII., which are on the lines of those in last year's report. I propose,

TABLE XV.
Canal Boats (52 weeks).

Registered during the year 1905				8
Transferred to fresh owners				_
Struck off register				I
On register, 30th December, 1905			4	101
Visits of inspection to wharves and	lock	s	Ι,	102
Boats completely inspected				381

TABLE XVI.
Houses Let in Lodgings.

				Houses.	Rooms.
Registered during the year	; let a	is furnis	hed		
rooms				4	16
Struck off register				12	24
On register, 30th December	er, 190	5		210	394
Houses let in lodgings	visite	d, but	not		
registered				245	507
Visits for registration purp	oses			4	16
Visits for additional inspec	ction		3	,268	
Nuisances found and abat	ed-				
Overcrowded rooms				Found.	Abated.
Dirty rooms				100	
Dirty and bad bedding				24	24
Dilapidated dwellings				85	83
Defective drains				75	75
Dirty closets				10	10
				1000	
		Total	133	333	331

TABLE XVII.

Other work of Temporary Dwellings Inspector.

	o common lodging-houses			1,30
"	" " " as			44
,,	furnished rooms as to sm	all pox		2,18
,,	vans			21
,,	tents			9
,,,	cellar dwellings			7
,,	" " closed			I
,,	overcrowded houses			
,,	infectious diseases			5
,,	test drains			3
,,	as to other causes			55
Univer	sity lodging houses—			
Hous	ses inspected, 29; containi	ng 97 r	ooms.	
Drai	n testings, 29 (defects foun	d 29, ir	12 ho	uses).
	sting of drains of houses pr (1904) 45 (defects found 4,			ined
Tota	l visits to these houses			9

however, to say a little about the last addition to the work of the lodging house inspectors, that of investigating the lodging houses on the University list.

Students' lodgings.—Mr. Carter has compiled this year, for the first time, a table dealing with the houses examined since we commenced our systematic inspection of students' lodgings in 1901. We have omitted the addresses of the houses, distinguishing them only by the number in our register. Most of the houses examined are in the North West ward.

In 1901, when the thing was new, we examined, by smell test, 61 houses, and we got a positive result from the test,

showing that something was wrong with the drainage in 31 of them—more than half. Before the house was put upon the University list the defects found were in every case remedied. In one case, No. 56, the house was not put on the list at all, as the defects were not remedied in time. This house shortly after became unoccupied. It remained unoccupied for some time. Work has been done since which would probably prevent the smell test from revealing any defect, but the actual requirements of our department in regard to students' lodging houses have not been complied with. Where we grant a certificate we insist upon conditions, some of which we might not be able to enforce under an ordinary nuisance prosecution.

The figures at the top of the columns in table XVIII. refer to the years in which the examinations were made. The sign + means that the examination of the houses by the smell test gave a positive result indicating a defect in the drainage. The sign - indicates that the smell of the test was not discovered anywhere either inside the house or in its immediate proximity. A thick vertical line in the table after the result of the smell test in any year shows that application was not made after that year for inclusion in the list.

In Nos. 2, 9, 21, 32, 38, and 91, although the result of the test was negative and is so entered in the table, there were structural conditions which in our opinion required to be altered. Some of them were such as would probably have led to a positive result at some later time. In No. 2 the soil-pipe acted as a rain-water pipe, and there were damp walls in the basement. In No. 9 the water closet was situated in the middle of the house. In No. 21, which had a water closet in the basement, the soil-pipe acted as a rain water pipe. In No. 32 the bath room was dirty. In No. 38 the water closet pan was offensive and the cistern defective. All these occurred amongst the 30 which gave a negative result

to the test made the first year (1901). In No. 91 the result had been negative when the house was first submitted for examination in 1904. The fall pipes, however, were still connected to the drains. These and some other matters were put right and next year the house again gave a negative result.

At first it had not been thought necessary to examine the following year a house which had given a negative result, or where the defect found had been remedied and a further test made with a negative result. It was found, however, on re-examining houses in 1903 which had been examined with a negative result in 1901, that two of these (No. 2 and No. 33) gave positive results in 1903, whilst Nos. 4, 19, and 24, which had been examined in 1901, and the defects put right, showed other defects in 1903. Amongst the houses submitted in 1903 which had been examined the previous year, one (No. 65) gave a negative result in 1902 and a positive one in 1903. It was therefore thought desirable to make it a general rule-first, that no house should be regarded as fit to be put on the list unless two separate testings of the drainage revealed no defect; and second, that where a defect was found, and thought to be remedied, two tests with negative results should be made before registering the house. Since 1902, therefore, where the result of the first test was negative, a second has been made with a like negative result before the house is entered on the list of lodgings. When the result of a test is positive, or when there is a condition likely to lead soon to a positive result, a house is not now entered upon the list of lodgings until the defects have been remedied, and two further tests made, both with a negative result.

It will be remembered that none of the 61 cases examined in 1901 were re-tested in 1902. The drainage of all the houses, however, whose occupiers applied for registration in 1903 was tested whether it had been tested before or not. In 1903, as

already said, 5 of the 34 houses which had been found right, or put right in 1901, again yielded positive results on our testing.

Of the 12 houses examined or placed on the list for the first time in 1902, seven were found defective and the defects remedied. Five of these 12 houses were not presented for registration the following year, but of the seven which were one showed a defect.

Of the 15 houses offered for registration for the first time in 1903, nine showed defects which were remedied before the houses were certified. Ten of the fifteen were offered again in 1904. One (No. 85) does not seem to have been re-examined, but the nine which were all gave negative results, and in the eight of them which we examined again (i.e., for the third time) in 1905 the result was again negative in all.

In 1904 30 new houses were offered for registration, and 13 gave a positive result to one or other of the two tests. Of the 30 only 13 were put upon the list for report in 1905, and of these 13 in two (both of which had given double negative results in the previous year) the result was positive.

In 1905 29 new houses were put on the list. Twelve of them gave positive re-action to the test, 17 did not, although tested a second time.

It will be thus seen that during the five years 147 houses have been thus examined for the first time, and 72, or 49 per cent., were found to have defective drainage. The defects were, with the exception of the one already mentioned, in every case remedied, and the drains tested with a negative result. Sixty-four houses, which had been either without a detected defect at the first examination, or where the defect had been remedied, were examined later a second time. Of these 35 were examined two years after they were first certified as sound, and 5 of them gave positive results.

Twenty-nine were examined a second time the year after they had satisfied us, and 3 of the 29 gave positive results. Of the whole 64, 8, or 12½ per cent., responded to our test on the second examination. The numbers are small, but it is perhaps worth noticing that of the 35, where a two years' interval had occurred, the percentage found faulty was 14'3; of the 29 examined with only a year's interval the percentage was 10'3. Thirty-eight were examined in this way a third time, and two were found defective (5'3 per cent.). Twenty-four were tested a fourth year, one failing (4 per cent.).

It will be noticed in regard to No. 19 which was on our list for four years, in three of which we tested its drainage, that this was found faulty on each of these three occasions. On the first occasion the joint of the water closet in the basement allowed the smell of the test to escape into the house, and the yard drain similarly allowed escape into the yard. The joints of the water closets were made good and several pipes in the yard taken up, until the result of a new test was negative. There was no disconnecting chamber and no means of applying a water test. No test was made in 1902. In 1903 the smell of the test in the watercloset of the basement again indicated a leakage. It was found that rats in the two years' interval had made burrows into the house from the neighbourhood of the sewer, and that leaky joints in the water closet drain outside had allowed the smell of the test to come up these burrows into the house. The faulty drain was laid bare for about three yards. It was found that the bend pipe and the pipes of the drain leading to the sewer had been laid with clay joints. The joint at the bend and about five other joints were remade with concrete. The owner, refusing to carry the relaying of the drain to the sewer as we wished, contented himself with stopping his repairs as soon as the negative result was obtained by the test. negative result was obtained twice. In the following year, 1904, the test again gave a positive result in this house, but

in a different place. On this occasion the defect revealed by the smell test was between the joint of the water closet and the soil-pipe on the first floor. This joint, which had been made with putty, was remade with white lead and again two tests applied, both with negative results. We have not had any further application to register this house, which is at present standing empty. The occupier has moved into the house numbered 126 in the table.

Dangers from inside water closets.—Of the 147 houses examined, 125 had inside water closets (93 F.V., 32 not F.V.), and 22 were houses with water closets outside. Of the 125 inside water closet houses, the first examination revealed defects in 65, or nearly 52 per cent. Of the 22 houses, with water closets only outside, defects were found in seven, or less than 32 per cent. This indicates, as has been pointed out in previous reports, that the existence of a water closet in the house adds enormously to the chance that the drainage will be found defective on testing. It will be remembered that in diphtheria houses and in typhoid houses, and in house-to-house work throughout the borough, the smell test has generally revealed defects in about one-third of the houses examined. In the non-inside water closet portion of these lodging houses the proportion found defective is thus about the same as throughout the town. The excessive proportion of defects found in the whole of these houses as compared with the town generally is evidently due to the more frequent existence of an inside water closet.

In 1897, when the Congress of the Sanitary Institute was held in Leeds, a large number of lodging houses were examined by our inspectors, and we were struck by the great number of defects found, especially amongst those with inside water closets. The present figures confirm these observations, and point to the general conclusion that, convenient as an inside water closet may be, its presence adds greatly to the risk incurred by the inmates of breathing an atmosphere impregnated with sewer gas.

It is not satisfactory to deal with very small numbers, but it is perhaps worth mentioning that of the 93 F.V. houses—that is those in which the soil pipe was carried full bore above the eaves—42 (or 45 per cent.) showed a defect on the first testing. Of the 32 not F.V. houses—that is those in which the soil pipe was used as a rain water pipe, or in which the ventilator had a smaller diameter than that of the soil pipe—the first test gave a positive result in 23, or 72 per cent.

I think the general inference from these and the figures put before you in previous years must be that an inside water closet is a source of considerable danger, and that to limit the amount of danger from these conveniences the following points should be insisted upon:—

(1) No inside water closet ought to open directly into a sleeping or living room, or into a passage communicating generally with the rooms of the house.

An ante room ventilated to the outside air, if possible with a through draught, ought to be regarded as an essential accompaniment of an inside water closet. Where the joints of the soil-pipe are leaky, as is probably the case some time or other in nearly half the water closets of the town, even the presence of a window in the apartment is not sufficient to dilute the gases continually given off from leaky joints, and the passages of the house should not be polluted by the atmosphere from an apartment containing such a soil-pipe. Where the structure of an old house does not permit of such an ante room the closet itself ought to be provided with permanent, unclosable ventilation.

(2) In all cases where there is an inside water closet the efficiency of the joints ought to be tested at least every second year. Manufacturers allege that there are joints which will not leak. Doubtless certain forms of joints resist the processes of decay longer than others, but none are immortal. Variations

TABLE XVIII.

	Snev	ving	resul	ts of	exa	mina	tion	of nts	stı	udelo	dgin	gs si	nce '	1901.	
	1901.	1902.	1903.	1904.	1905.				1901.	1902.	1903.	1904.	190 5.		
1	+							38	7						
2	-		+	-	-			39	-						
3	_	1.	-	- 1				40	-		-	-	-		
4	+		+	-			1	41	-						
5	+		-	-	-			42	+						
6	-		-	-				43	+		-	-			
7	-							44	+		-	-	-		
8	+		-	-	-			45	+		-				
9	-	1						46	+		-				
10	+		-	-				47	+		1	-	1-1		
11	-							48	+						
12	+		-	-	-		14	49	+		-	-	1-		
13	-							50	+						
14	+	20		-				51	-				1.30		
15	-							52	+						
16	-		-	_	-			53	+		_	_	-		
17	-	1						54	-		-	-	-		
18	-							55	+		-				
19	+		+	+				56*	+			1 18			
20	+						-	57	-		-	+	-		
21	-							58	+		-	-	-		
22	+		-	-	-			59	-		-	-	-		
23	-	1						60	-		-		13		
24	+		+	-	-			61	+		-	-	-		
25	- "							62		+	-				
26	-							63		-					
27	+		-	-	-			64		+	-	-	_		
28	+		_					65		-	1+				
29	+							66		+					
30	+		-	-	-			67		+	-				
31	-		-	-	-			68		-					
32	-					1		69		+	-	-	-		
33	-		+		1012			70		-			4		
34	+		-					71		+	-	-			
35	-							72		+	-	-	+		
36	-		-	-	-			73		-					
37	-							74			-	-	-		

4 See text.

					Т	ABLE	. XV	111.—C	ontinue	ed.			-		
	1901.	1902.	1903.	1904.	1905.				1901.	1902.	1903.	1904.	1905.		
75			+	-	-			112				-			
76			+	-	-			113				-			
77			+					114				+			
78			+	-	-			115				+	-		
79			-					116				+			
80			-					117				+			
81			+	-				118				-			
82			+	-				119					+		
83			-	-	-			120					-		5
84			+	-	-			121					-		
85			-		-			122					-		1
86			-	-	-			123					+		
87			+					124					-		
88			+					125					-		N.
89				-			1	126					-		
90				-	-			127					+		
91				-	-			128					+		
92				+				129					-		
93				+	-			13C					-		
94				-				131					+		
95				+				132					+		
96				+				133					+		
97				+				134					-		
98				-				135					-		
99				-	+			136					-		
100				-				137					+		
101				-				138					+		4.0
102				-				139					+		
103				+	-			149					+		
104				, -	-			141					-		
105				+				142					-		
106				-	-			143					-		
107				+	-			144					+		
108				-	+			145					-		
109				-	-			146					-		
110				-	-			147					-		
111				+											

in heat and cold—the different co-efficients of expansion of the materials used, all point to this inference, and even the hydraulic test of a joint proves nothing except that the joint was tight at the time the testing was made.

It should be remembered that a flaw in a soil-pipe joint means often more than the mere entrance of sewer gas. It frequently means the deposit by evaporation of matters suspended in the fluids from the closet, and the retention thus of solid and offensive substances within the dwelling.

The calcium phosphide test.—A word may perhaps be necessary about our continued use of the smell test. With proper precautions the calcium phosphide will cause so strong an odour in a moist drain as to reveal even a very slight leakage in the immediate neighbourhood of the house. Properly used and under favourable weather conditions it may reveal defects in joints many yards away from the house. The conditions of the weather, however, require to be kept in mind, and a single negative test should never in any case be relied upon. Where there is a disconnecting chamber it is well, when permission can be obtained, to test all recent drains with the water test. The water test, however, only reveals faults in the drain below the surface, and does not indicate the failure of joints in the vertical lengths of a soil-pipe, and between the soil-pipe and the basement. The water test is frequently considered to be too severe a test to be applied to old drains. It has even been alleged that it is capable of producing a leakage. It is therefore objected to by many owners. It has occasionally happened to us that the smell test has not revealed some defect in pipes underground. Such cases are, however, rare, and are generally to be explained by the mode in which the test was applied.

APPENDIX.

TABLE A, Part I.

Table shewing deaths recorded in the City of Leeds during the fifty-two weeks ended 30th December, 1905, classified according to cause, age, and the registration sub-districts in which they occurred.

	ring ring	500	::::::	: ::	::	: : :	1:::	ंद	8 11	8	120	160	90	
S OF	Out- siders occurring	nud 5	111111	1 1 1	::]	: : :	:::	1 ::	et :=	1 01	16	18	178	
DEATHS			111111		7.1		:::	12:	9 9	1 0	9		1	
DE	Leeds persons securring outside	0				* * *		1	-	-	1	9	101	1
	1 5 5 5 C	Ping C	1:::::	111	::	: :	111	::	1 1	1 1	-	-		
	Annual rate per 1,000		000000	0.00	0.80	10.0	0.02	1-23	2.76	7-49	8-16	15-65		
-	iliy	all		121	365	4 00	Nr 5	558	1258	3410	3714	7196		15-6
Total	Mortality in City.	5.00	131122	5 12	15	. 4 0	10 et 0	542	712	2113	2281	0270	3	11-0
-	N	pun 2	-288800	118	350	: :	C4 F0	1.91	949	0.0	14332	0,000	5 1	50-91
	1 8	0v. 5	::::::		::		1::	.4	r- 13	: t2	34	58	5	ID.
	Bramley	- American						1		. 00	100	5 5	13	
		pun .	11111		::			1 : :	1 1	1				
SES.	Holbeck	5.00	111111	1 11	:::		::::	:00	9 10	: 2	133	18	37	
HOUS	H	pun	:= ::::	1 11	::	1 1	: :::	1:4	- : - :	: -	1:	-		
WORKHOUSES.	slet	200	::::::	1 11	: :	1 1	121 4	1:0	-10 W-	15	30	45	00	
M	Hunslet	nud 5	::::::	: ::	::	: :	:::	1::	: :	: :	100	10	4	
		5.00		: :00	::	1.1	11:	:83	5 3	207	200	101		
	Leeds.	Sandration.				2 30 1				1 03			436	
-		pung		1 11	:-	1 :		1			22	83		
	Fever Hospitals	50	23 :00 ++ ::	: :8	::	1 : :	-::	1::	00	8	-	22	88	
	Hos	pun 2	1 :88 : :	: : :	::	: : :	111	::	: :*	35		123		
	uy,	5.5	111111	: :19	::	: : 6	; PO +-1	: 13	61 83	141	304	445	10	
	In- firmary, &c.	pun	:::=::		:0	: :	:::	: 10	n ng		99	801	553	
								-03	FI .	. 100	-	4 10		
	Osmond- thorpe.	379 d ov.		: ::	::			1	-		-	03	9	16.8
		o mid		- 11	::			1 :::						
1	Chapel- town.	44,378 nd ov. 5 5			0			1 35	5 46	7 128	149	1 277	418	9-01
		13		* * *		1 1		1.	64 .	4	韦	#		
	aley 435	17,874 ind ov. 5 5	1 . : : : :	: :=	:-	1 1	::-	:83	61 83	3 8	82	162	225	12.2
	Brar 18,	17, und 5	15 : : : :	19 ::	:03	1. 1	11.1	::	15	4 8	34	13	93	17
	3 2	1.00	: : :00 : :	: ::0	:-	: :	- : -	:55	# 8	141	172	515	23	0
	irks 16,68	45,495 und ov 5	· : HO1 : :	# ::	00	1 1		1:-	98 :		97	150 3	463	10.0
-	У. ч	1.		: :01	: =	: 04 -		:13	E 25		1	1		
	ortle 1,769	62,925 and ov. 5 5						1	_	1 64	4 253	9 482	821	12.9
-	. W.	1		53 ::	:4			1 3 3	8 :	-	194	339		
	beck 88	34,720 and ov. 5 5	:07 :::	: : : =			: :::	:5	88 82	168	133	301	520	15-1
	Holl 34.	and St.	10 10 1H	13 : .	:83	::	: :03	:-	45 :	01	118	213	43	7
	let.	1 .	;∾ ; ⊢ ; ;	: :01	:10	: : *	:	57:	103	1 - 1	216	459	_	00
	Hunslet, Holbeck, Wortley, Kirkstall Bramley 74,576 34,439 63,769 46,685 18,435	75,482 und ov.	:2021	25 ::	: 93	: :	:	1.50	121	1	1083	492	951	12-8
-		1 22	·0H-	100	:10			: 23	77 11		1286	282		
	South E.	A A						1					627	18.1
1.		13	11 :1 :2	°° : :	: 6		111		8 :		159	335		
LEEDS	West.	84,714 5 ov.	;co := : :	- :0	:~	: : 4		: 8	011	2 88	376	702	1,062	12.5
CEE	≥ 8	SE Pus	:5000	2 ::	:5	: :	. 7:	: 4	8 01	0 4	216	360	1,	12
	ė o	1	1 .0 01-	03 1414	:00	: : 0	3 : : H3	1:5	28 27		181	394		-
	North.	58,344 und ov 5	:22 :00 00 :	1:	:22:	::		1	93	1.00	136	412 3	806	13-7
-			:::::@g		· ·	:	-::-		05 .	:			1	h d
	TOWNSHIP &c. Estimated Population 8	70	Small-pox Measles Scarlatina Diphtheria Croup (memb)	Whooping- cough Typhus	doubtr doubtr urrhorn	Cholera Rheumatic Fever Acute & Sub-	acute Rheu- matism Erysipelas Pyæmia Puerperal	Ague	Bronchitis Pneumonia Pleurisy Heart Disease	Total of Above causes	All other	eran	Total	Mortality per 1,000 per an,

† Including 12 deaths in the Consumptive Hospital, Armley (see quarters' returns).

* There were 85 deaths at Manston, and 3 deaths at Killingbeck (outside the City boundary).

TABLE A, Part 2.

Table of populations, registered births, and mortality at certain ages, in the registration sub-districts.

(Public institutions regarded as sub-districts.)

-	pulatio	n estimated at each age		of 1905		456,787	11,964	41,144	93,645	91,934	192,143	11,273	14,684	
DECL					10.									
REGIS	STRA	TION		ation at ges.§	red		Mo	rtality fro	m all cau	ses, at su	ıbjoined a	iges.		Death-
	OF OF EDS C		Census 1901	Estimated to middle of 1905.	Registered Births.	At all ages.	Under 1 year.	and under 5	5 and under 15	15 and under 25	25 and under 60	60 and under 65	65 and upwards	rate per t,000 for each district.
Leeds To	ownship	-North	59,281	58,932	1,792	806	290	122	31	30	177	59	97	13.72
	do	West	84,340	85,005	1,905	1,062	266	94	31	46	273	75	277	12:54
Do.	do.	South	34,299	34,817	1,235	627	244	91	25	22	125	32	88	18:07
Hunslet			69,064	74,576	2,140	951	337	155	31	32	192	61	143	12:80
Holbeck			31,572	34,489	1,071	520	153	66	18	27	120	36	100	15-13
Wortley			59,328	63,769	1,643	821	238	101	34	35	203	47	163	12-92
Kirkstall		11 75	41,561	46,686	981	463	113	37	13	15	110	53	122	9-95
Bramley	**		17,299	18,435	459	225	47	16	10	18	59	24	51	12:25
Chapelto	wn		31,845	39,720	1,016	418	109	32	10	16	123	27	101	10.56
Osmondt	horpe		379	358	3	6		2	1	1	1		1	16.82
Infirmary						541	55	53	45	51	259	41	37	
*Fever He	ospitals			**		88	1	34	16	13	21	1	2	
	- (Leeds	***		78	436	17	12	2	12	183	56	154	
WORKHO	USES	Hunslet				48	2	1		2	15	3	25	
		Holbeck	**		4	37		1			10	8	18	
		Bramley	**	4.4	10	63	3	2	2	1	16	10	29	
Consump	tive H	ospital				12		**		4	8			
For the w	For the whole City 428,968 456,787 12,33					7,124	1,875	819	269	325	1,895	533	1,408	15-65

^{*} There were 85 deaths at Manston Hospital, and 3 deaths at Killingbeck Hospital (outside the city boundary). See note to table B.

The rates are calculated for the year on populations estimated by the Registrar General's method, described at p. 88 of the Annual for 1904. Those for several quarters are on populations as previously estimated.

[§] For explanation of discrepancies between district populations for 1905 in this table and those which immediately follow (A 3, 4 5, 6), see p. 66, part III.

rable shewing Deaths recorded in the City of Leeds during the thirteen weeks ended 1st April, 1905, classified according to cause, age, and the registration sub-districts in which they occurred.

The Registry The											_					-
No.	ths	ring ring	04.	:::::	: :	: ::	: :		::::	1::	-	HE	2	23	31	
No. No.	Dea	O de la company	5 5	:::::	: :	: ::	: :	:	-111	1::	:	: :	:	100		
Carrier Carr		Annual ate per 1,000	i.	0-03 0-12 0-14 0-04 0-01	0-18	0.09	: :	0.07	0.01	12:	3.03	0.69	7-31	8-67	15.98	
Control Cont			All	5841	21 :1	9: 9	: :	00	4	178	344	123	832	987	1819	16.0
The Registration of the Property North, No	TAI.	in in ty.	S A	01-000 H ;	: :9	: =	: :	00	4	174	221	127	627	620		12-4
Care	To	Mor	o pd	128821	22 :	cu:	: :	:	:::	:4	123	19	202	367		52
Charge C	_	o			: ::	::	: :	:	:::	1 :0	:	::	et .		4	
Charge C	Coni	Hos- pital.	o pu				: :	:	:::	1::	:	::	:	:		
North-Hamiltonian	-		1	::::::	: ::	::	: :	:	:::	:00	1	ø :	7	-	9	
North-Hamiltonian		ramb		::::::	: : :	1::	: :	:	::::	::	:	::		69	-	
North-Hamiltonian	of.	eck B	ov.	::::::	: : :	::	: :	:		::	-	:3	-	2	00	
North-Hamiltonian	OUSE	Holb	5 5	::::::	: : :	::	: :	:		::	:	.::	:	:		
Recommendation Companies	окки		ov.	::::::	: : :	::	: :	:	:::	:-	-	::	63		6	
North. West, South E. Hussler, Hollacd Wortley, Kirkstall Bransley, Chingel, Osmond, Alexanded, St. 1	W	Hun	and 5	::::::	: : :	:::	: :	:	:::		:					
Charge C		- de		:::::	: :-	::	: :	:			12				152	
Charlest C	_			:::::			: :	-:		1 ::	-					
North Reserved North Nor		ever pital	l ov.							1::					33	
No. No. Still No. th. West, South B. Hunslet, Holbeck No. tless Still Still No. tless Still No. tless Still No. tless Still Still			-	- :=- : :			1	03		- 10	4					
North Nort		In- mary, etc.								1::	-				129	
Charge C			_				1 : :	-	1::	-	:	_	:			
Particle North. West. South E. Hunslet. Holbeck Wortley. Kirkstall Bramley. Chaptellower Se. 34 St. 74 St. 475 T. 24.82 St. 720 62.925 45.495 T. 277 44.378 St. 74 St. 778 St. 77		norpe 379	o pu				: :	:	:::	::	:	::	:	:	:	:
Charles North. West. South E. Hunslet. Holbeck Worrley. Kirkstall Bramley. Townships. Charles South E. Hunslet. Holbeck Worrley. Kirkstall Bramley. Townships. Charles South E. Hunslet. Holbeck Worrley. Kirkstall Bramley. Townships. Charles South E. Hunslet. Holbeck Worrley. Kirkstall Bramley. Townships. Charles South E. Hunslet. Holbeck Holbeck			,		::		: :	ce	:::	19:	19	90			0	9
Preserved Court		Shape town 44.37			H :	: ::	: :	:	:::	1::	10	:-			12	10
Delta Delt		9	. io	:::::	: :	: ::	: :	:	:::	1000	10	36	19		53	\$.5
Delta Delt		framil 17.8	5 S	: : : : : :	- :	: :-	: :	:	:::	0 10 10 10 10	64	;=	10			2
Delta Delt		11 Se		:::=::	: :	: :-	: :	:		12:	13			-	143	9-2
Delta Delt		Kirks 45,4	nmd 5	::: : : :	7 :	: :-	: :	:	:::	1	5					
Delta Delt		rley.	9.50	;- ;- ; ;	: :-		: :	:	-:-						203	12.9
Delta Delt		W or	nud 5	:4::4:			: :	_	100	-	_					
Delta Delt		lbeck ,720						:						*****	116	13.4
Delta Delt		. H.								-			-			-
Delta Delt		inslet 5,482	di ov.												210	11.5
December County										-	-	-				
Cheer and cover 5.		nth J	o pu									:-		one more	143	16.6
Cheer and over 5. Section	. S.							04			-	128	100			M)
Cownship Ac. North. Retimated open and over 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5.	ED	West 84,714	o pu				: :	:							231	12.
OWNSHIP &c Nor Retinated 156.787 156.787 Under and over 5. Sandhina Over 5. Sandhina Over 5. Sandhina Opplitheria Oppl	LB						::	C/4	::10	12:	23	12 6	19	8	8	60
TOWNSHIP &c. Estimated Population 156.787 Under and over 5. Snathpox Nansles Scarlatina Coup (memby n undefined Whooping Coup (memby n undefined Whooping Coup (fermit Coup (fermit Coup (fermit Coupler or Ague Phthisis Prever Ague Total of All other Coupler or Total of All other Total Total Mortality per Lioup per au,		Nort 58.34	pung 2		::	: : : =	1 : :	:	::::	1 .:	22		B	69	33	12
	Washington Com		Under and over 5.	Smallpox Mersles Scarlatina Diphtheria Croup (memb)	Whooping Cough	Other or doubtfi	TCholera Rheumatic Fever	Acute & Sub-	Eryspelas Pyzenia Puerperal	Ague	Bronchitis	Heart Disease Injury, &c	Total of above causes			Mortality per 1,000 per an.

* There were twenty ween deaths at Manston Hospital, and three deaths at Killingbeck Hospital during this quarter. * Finglish.

A. 4.

Table shewing Deaths recorded in the City of Leeds during the thirteen weeks ended 1st July, 1905, classified according to cause, age, and the registration sub-districts in which they occurred.

Part North Nation	_	~									warmh's and		_	-	-
Control Cont	eaths	Out- iders in in	d ov.	1:::::			: :	:		:04		· · · · · ·	durin.	3	
Converged Conv	A		E G	11111	:::	::	::	:	:::	::	: ::	1:	63		
The Property Prope		Annus rate po 1,000	pop.	:00000 :00000	0.24 0.01 0.09	0.01	0.05	0.04	0.01	1.38	2.84	02.9	8-42	15-11	
The Property Prope	.2	ty	Ages	:000000	27		: 0	4	4 11	137	323 110 81	762	1 %	1720	15.1
Convergence	Cora	in in City.	over.	:======================================	1 16		: 03	Ф	:- ·	133	139	299	610		11.6
Charles Parimeter Parime			und.	; t~010000	% :F	-2-	: :	:	⊢ ::	:4	139	502	348		41.9
Part		ptive os- tal.	5.	::::::	:::	::	: :	:	:::	:03	: ::	03	:		
Part	_ (1		111111	: ::	::	: :	:	:::	::	: ::	:	:		
Part		amley	d ov.		: ::		: :	:			······			19	
Part		ck Bra					: :	:	2000						
Part	USES	Tolbe	o pu											13	
Part	RKHO	let.			: ::		.: :	:	:::			10	10		
Part	Wo	Huns	pun 2	::::::	: ::	::	: :	:	:::	::	: ::	:		15	
County C			5.0	:::::	: :-	::	: :	:	:::	:00 :	12 2	29	29	03	
Comparison			-		: ::	::	: :	:	:::	: = 0	.:		100	F	
Comparison		Fever iospital	and ov.		: ::	::	: :	:		: : *			:	12	
Comparison		arry, I		:::::	: ::	::	: :	-	::-	:03 0	20 00 25	4	8	69	
Formation Form		firm	onu 5	::::::	: ::	::	: :	:	:::	: 64:	9 :19	o	16	-	
Formation Form		Samond- thorpe.	5 5				: :	: :					; ;	1	9.01
Computation				:::::	: ::	::	: -	1	:: -	:o e	- 6000	83		1	do.
Choker and cover & Secretarian Cover & S			5 5	:::::	: ::	::	: :	:	:::	:: 4	٠ : :	9	1	9	00
Choker and cover & Secretarian Cover & S		mley.	1 ov.	::::::	: ::	::	: :	:	:: -	: + 4	0 -10			63	4.1
Choker and cover & Secretarian Cover & S		II Bra	1		- ::		: :	:							
Choker and covered by the control of covered by the covered by t		irksta 45,496	o pu											123	10-9
Choker and covered by the control of covered by the covered by t		S 63.		::::::	: :=		: -	-	:::					0	1
Choker and covered by the control of covered by the covered by t		Wortl	5 5	:::::	9 ::	::	: :	:	:::			interest	diame.	31	12
Choker and covered by the control of covered by the covered by t		lbeck 720	5.0	:-:::	: ::	-:	: :	:	:::	:21		47	12	40	63
Choker and covered by the control of covered by the covered by t		Но				-	: :	:	:::						16
Choker and covered by the control of covered by the covered by t		umslet 3,482	o po				4 1		:" :					223	12.5
Estimated															_
Estimated		South 34,47	5 5					;						146	17.0
Estimated Population— Estimated Page, Test of Cough C	DS.			::::::	- ::	; :	: :	1	:::			87	16	03	6
Estimated Population— Estimated Page, Test of Cough C	EE	We 84,7	5 5	: := := :	04 ::	::	::	:	:::	:: £	:-	0.00000001		255	11
Estimated Population— 156,787 Under and over 5. Smallpox Smallpox Croup (memb) ,, undefined Whooping Cough Cough Diphtheria Croup (memb) ,, undefined Whooping Cough Cough Diphtheria Croup (memb) Lever Croup (memb) Cough Cough Population Esteric Cough Cough Cough Cough Mortalism Fever Fever Phinsis Pherperal Pever Phinsis Pheniss Pheniss Pheniss Phoenochitis Pheniss Prever Total of Total of Total Total Total Total Total Total Total	I	orth. 344	50	:::-::	:		: :	:	:::					297	1.2
Estimated Population— \$456,787 Under and over 5. Smallpox Measles Scarlatina Diphtheria Croup (mem., undefin) Whooping Cough Cough Cough Cough Cough Cholera Fever			5		02 :-		: :	:	11.1	. : 8		83	4		
	Control Control	Estimated Population— 456,787	Under and over 5.	Smallpox Measles Scarlatina Diphtheria Croup (memb	evers.	Diar	Acute & Sub.	acute Rheu-	Erysipelas Pysemia Puerperal Fever	Ague Phthisis Bronchitis	Pleurisy Heart Disease Injury, &c	Total of above causes	All other		Mortality per 1,000 per an

* There were twelve deaths at Manston Hospital during this quarter. ¶ English.

A. 5.

1905.—THIRD QUARTER.

Table shewing Deaths recorded in the City of Leeds during the thirteen weeks ended 30th September, 1905, classified according to cause, age, and the registration sub-districts in which they occurred.

_	-														
1 de la constitución de la const	of Out- siders securring	City.	1:::::	: : :	: ::	::	:	:::	1::	-	1 9	00	18		
1		and City		: : :	: ::	::	:	:::	::	-	:-	63	15	28	
	Annual rate per	pop.	0.011	0.09	5.99	10.0	0.03	0.01	1.03	1-67	1-06	7.86	7.87	15-73	
	^	All	:0100 :00	35 :01	325	: "	140		117	190	611	168	988	1790	15-7
	Mortality in City.	und. over.	:01D : : :	10: 2		: -	ю	- : -	114	106	38	410	510	920 1	9.1
1	Terai. Mortalit in City.		:5000 :01	12 ::	315	: :	:	;	:10	88	72	484	388	870	
-		5.0		1 : ::	::	: :	:		:4		::	4	:	1 00	65.7
	Con- sumptive Hos- pital.	nnd o		1 : : :	::			::::	1::		::	:	:	4	
-		5.0	::::::	:::	::	: :	:	11:	:	-	r :	ю	6	1	
	Bram	end 5	:::::	:::	::	: :	:	:::	1::	:	::	:	-	13	
05	ec,	5.0	::::::	:::	::	: :	:	:::	::	:	٦:	н	-		
tonos	Holl	nug 2	:::::	: ::	::	: :	:	:::	::-	:	::	:	:	00	
WORKHOUSES	Hunslet. Holbeck Bramley.	ov.	:::::	:::	::	: :	:	:::	:-	:	::	-	9	п	
=		5 S	::::::	: ::	::	: :	:	::::	::	:	::	:	:	-	
	Leeds.	. o.	:::::	: ::	::	: :	:	:::	17	00	22	42	\$	85	
-		pun 2	111111	: ::	:	: :	:	:::	::	04	::	ю	7	0,	
	In. Fever firmary, Hospitals	d ov.	::4:::	: :0	::	: :	: '	- : :	:::	03	::	12	:	21	
-	. Но	o S	:: co+ ::	1 : : :	::	: :	:	:::	: :		03	6	:		
	In- mary etc.	d ov.			::				:-		3 14 2	20	69 /	122	
-		o S	:::=		.6	: :	-	111	::			1 16	. 17		
	Osmond- thorpe.	und ov.			::	: :		::::	:-	······	::	:		-	10-6
-		1 . 1	::::::			: :	:		-10	40 1	-01	19	35		
	Chapel- town.		::::	H ::	10:	: :	:		:			16	21 2	99	8.1
-		1 2		: ::	:	: :	: -	:::	: 47	- 0		41	22		
	3ramley 17.874	bun 5	:::::	: : :	111	: :	:	:::	::	-	::	-	6	45	10-1
	Illan 88	5.0	::::::	: :04	::	: :	:	:::	:4	4 1	2-	52	40	87	9
	Kirkstal 45,495	5 5	::-:::	⊣ ::	-1:	: :	:	:::	::	co.	:-	15	82	6	9.8
	Vortley. 62,925	5.0	:: : :: :	: ::	:-	: -	:	: : :	:23	= =	0.0	22	54	122	14.1
	Word	pun 2	:∾ : : : -	œ ::	:00	: :	:	:::	::	-	.03	28	8	0	14
	Holbeck 34,720	0,0	:7::::	: ::	::	: :	: :	: :				53	*	131	15.1
	. Н.о.	o S	:00 : . : :	۳ : :	-	: :	: :	- :			::	28	32		H
	South E. Hunslet. Holbeck Wortley. Kirkstall Bramley. 34,476 73,482 34,720 62,925 45,495 17,874	d ov.	:-:::	: ::	:03		-	: -				68	49	260	11:2
	H E	5 5	: : = = = : :	9 ::			: :				-	8	99	-	_
	34,476	und ov.	.03			: :	: :			10		32	88	181	21.4
s.	4			: : 01	:02	: : 0		- 1			- 1	8	44	1	_
LEEDS.	West. 84,714	und ov.	: : : : : :	4 ; ;	:4		: :	: :	1 21				59 83	261	12.4
LE			::::::	04 ::	-61		: :				-		02	-	-
	North. 58.344	5 5		~ ::	. 59		: ;	: :				*******	63 3	83	15.8
0			- 10000	: : :				:				-1-	-	:	an'.
TOWNSHIP	&c. Estimated Population- 456.787	Under and over 5.	Smallpox Neasles Scanlan Diphtheria Croup (memb) undefined	Cough Cough Typhus Enteric	Diarrhoga	Rheumatic Fever Acute & Sub-	natism Erysipelas	Pysemia Puerperal Fever	Ague Phthisis Bronchitis	Plenrisy J	njury, &c.	above causes	Chuses		1,000 per an.
TOW	Fopul 15	Con	Scarl Diph Cross	GVETS.	Diarrho	Rhe Fe	natism Erysipel	Pyre Per	Ague Phthisis Bronchiti	Plem	Injur	bove caus	5	Total	1,0
-				,niino.	2	-						6			_'

* There were twenty-one deaths at Manston Hospital during this quarter. * English.

1905.—FOURTH QUARTER.

Table shewing Deaths recorded in the City of Leeds during the thirteen weeks ended 30th December, 1905, classified according to cause, age, and the registration sub-districts in which they occurred.

-	No.	-		-											-
Deaths	of Out- siders occurring in City.	and ov.	111111	1 : : :	: :	1::	:	:::	:04	C/I	4:	00	63	43	
A			:::::	: ::	::	::	:	:::	::	:	::	1:	9	4	
	Annual rate per	dod.	0.06	0.33	0.01	10-0	0.03	0.05	rin	3.52	1:31	8.10	7.67	15-77	0
		All	:47.6 :1	16: 38	1.8	: -	100	014 :	126	401	149	922	873	796	15.8
TOTAL	Mortality in City.	over.	:∞⊶∞::	: :9	:10	: ⊣	10	H02 :	121	202	568	292	1 5	1098 1795	10-9
5	I No		:900:1	23 ::	23.	::	:	H01 :	:10	66	24	365	332	169	52.7
To	200		::::::	: ::	::	::	:	:::	:01	:	::	64	1 :		165
	Con- sumptive Hos- pital.	and ov.	::::::	: ::	::	: :	:	:::	1::	:	::	:	:	0.1	
		ov.	::::::	: ::	::	: :	:	:::	:-	ю	ı٥ :	0	9		
	Bram	end 5	:::::	: ::	::	::	:	:::	::	:	::	:	:	15	
SES.	Hunslet, Holbeck Bramley	o.5.	:::::	:::	::	::	:	:::	:03	02	∾ :	9	62	00	
HOUS	Hol	enn 5	:::::	: ::	::	: :	:	:::	::	;	::	:	:	"	
WORKHOUSES	islet.	0.0	:::::	: ::	::	::	:	:::	:10	03		7	0	13	
		and 5	::::::	: ::	::	::	:	:::	::	:	::	:	-		
	Leeds.	d ov.		: ::	::	::	:	:::	:19	9	10	43	33	(8)	
-		end 5	111111	: : :	::	::	:	:::	::	-	::	-	ю		
	Fever Hospitals	und ov.	::=01::	: :9		:::	:		1::		:-	14	-	13	
-		-	::n4::	01	::	1	:	-03 -	1 :::	4	36	6	-		
	In- firmary, etc.	und ov.							-03		1 9 1	12 32	15 82	141	
i-				1 11	::		÷	111		-		2	-	-	
9	Osmond thorpe. 379	und ov.							1::		::	:	-	4	42.4
-		ov. u	::::::	: ::	;=	::	:	::::	:0	12	910	該	42		0
	Chapel- town, 44.378	pun 2	:- : : : :	ca ::	::	: :	:	:::	::	6	::	12	23	111	10-0
-		5.0	::::::	: :-	::	: :	:	:::	: 1-	00	0100	22	17	03	0
	3ramley 17.874	5 5	:-::::	- ::	:-	::	:	:::	1::	9	::	6	6	62	13-9
	tall E	_	:::=::	: :: ::		: :	:	- ::	:00	6 2 :	21	18	0	90	
	South E. Hunslet. Holbeck Wortley. Kirkstall Bramley. 34,476 73,482 34,720 62,925 45,495 17,874	und ov.	:::-::	: ::	e :	::	:	:::	::	7	:-	10	23	100	60
	Vortley. 62,925		:::::	: ::	::	: :	:	:::	13:	83	55	88	9	202	13.2
1	Wor	und ov.	:-:::	ca ::	:40	: :	:	:- :	::	8	::	81	47	03	12
	Holbeck 34,720	o. 5	::::::	: ::	:-	; =	:	:::	:00	23	000	47	Sign 1	133	15.4
	Hol 34,	e s	:0::::	4 ::	:10	: :	:	:≓ :	::	12	::	8	88	-	11
1	Tunslet. 73,482	o. 5	:= : : : :	: :-	:-	::	63	:::	12		13	61	48	258	11:1
_	Hu 73	5 5	:2::::	13 ::	:-	: :	:	:::	;	4	-:	88	13		-
	outh E 34,476	d ov.	:01 : : : :	: :-	::	: :	:				99	8	31	154	17-9
o's		bun 9	;o ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ;	ю ::	;10	1 :	:	:::		45	:	8	83		_
LEEDS	West. 84,714	5 5.00	:014	.: :	::	: :	-	- : :			11 28	58	1 87	268	12.7
LE	1.	5. 5 5	· A · · · ·		.		-			CA		24	9 54		_
	North. 58,344	5 5	17: ::: 3:	9 ::		::	:	: : : : : :		83	.2 18	64 62	39 49	214	14.7
-		T		~ : :		:	-	::-	1::-	_	0 1	0	70	:	± ii
	Estimated Population— 456.787	Under and over 5.	Smallpox Measles Scarlatina Diphtheria , undefined	Whooping Cough ever Typhus ever Cherric	Diarrhoea .	Cholera Rheumatic Fever Acute & Sub-	natism	Erystpelas Pyæmia Puerperal Fever	Agne Pluthisis Bronchitis	Pneumonia Pleurisy	Heart Diseas Injury, &c.	Total of above causes	All other causes	Total	Mortality per 1,000 per an:

* There were twenty-five deaths at Manston Hospital during this quarter. ¶ English.

A.

Table shewing Deaths recorded in the City of Leeds during the thirteen weeks ended 31st March, 1906, classified according to cause, age, and the registration sub-districts in which they occurred.

Deaths	out- siders curring in City.	0.00	:::::	: ::	::	: :	:	:::	::	- 22	00	37	99	
Death of Out- siders occurrin in Ciry.		und ov.	:::::	: ::	::	: :	:	:::	::	: ::	:	5	(Z)	
	Annual rate per 1,000	dod	1.26 0.03 0.03 0.01	0.36	0.00	10-0	90.0	0.02	1.46	3.32 1.57 0.55	9-30	8-59	17.79	
,		Ages	146 31 31 1	12: 42	10	: =	-	m : 0	169	383	1062	286	2064	17.8
Terai. Mortality in City.		over.	:6121 ::	: :=	:10	: □	-	۰: ۳	162	203	159	612	1263	12.4
		und.:	137	5 : 다		: :	:,	01 : :		. 9	411	38)	791 1	6-89
	9	o	::::::	: ::	::	: :	:	:::	;=	: ::	1	: [1	(2)
Con- sumptive Hos- pital.		nind o	::::::	: ::	::	: :	:	:::	::	: ::	:	:	-	
		5.	:::::	: ::	::	: :	:	:::	:-	4 4:	6	9		
	Holbeck Bramley	5 5	:::::	: ::	::	: :	:	:::	:: :	: ::	:	;	15	
68.	beck	o. 55.	111111	1 14	::	: :	:	:::-	-:-	1 000	9	2		
WORKHOUSES.		5	:::::	: ::	::	: :	:	:::	::	: ::	:	:	=	
ORKI	Hunslet.	ov. 5	:::::	: ::	::	: :	:	:::	:"	: -:	03	9	11	
*	Har	2 5	:03 : : : :	:::	::	: :	:	:::	::	: ::	63	-		
	Leeds.	i ov.	::::::	: :-	::	: :	:	♂ ::	:23:	¥ %4	76	46	130	
		end 5		: ::	::	: :	:	:::	:"	٠: ١	4	17	H	
	Fever Hospitals	d ov.	:===	; ;t-	::	: :	:	::-	::	: :0	19	4	27	
		bun 5	:10 : :	: ::	::	1010	:	:::	::	: ::	4	:		_
	firmary, etc.	d ov.	111111	: :01	::	: :		:::		7 7 5 19	4	88	152	
		bin 5	::::::	: ::	:-		-	1::		. H .	00	1 18		
	Osmond- thorpe.	und ov.			::	: :	:	113	1 1				04	22.7
							1		-01	12 8	:	8 9	-	
	Chapel- town. 41.594	und ov.	.00 .03 .11							9	98 100	17 4	121	11.7
		5. ti		: ::	::	::	-	111	-00	900	16 1	13 1		
	tramley 18.709	o pun	:::=::		::			111	::	ታ ::	5 1	6	43	9
	HE I		;= ;m : :	1 : : :	::	::	-	::-		2 22	18	49		
	Cirkstal 47,911	o pun	:4H4 : :	: ::	:01	: :	:	1:::		4 :-	16	24	125	10.5
	57 K		:01 : : :	111	::	: :	:	::::	12:	62 25	9		00	7
	Vortley 64,837	o pun	:8: -: :	9 ::	::	: :	:	- : :		9 ::	13	52 64	238	14.7
	eck N		:::::	: ::	::	::	:	:::	10	0 00 1	83	8	CA	_
	Holbeck 35,189	und ov.	:10 : : :	r ::	:=	: :	:	:: .	:01	1: 14	31	43	132	15-1
	Huuslet, Holbeck Wortley, Kirkstall Bramley, 75,901 35,189 64,837 47,911 18,709	500	:0 :□ : :	: ::	:-	: :	:	:: ~	24:	13 33	Ħ	젊	82	ю
	Hunslet 75,901	und: o	:%: -0 :	# ::	:-	: :	:	- : :	:-	g ::	88	19	308	16.3
	outh E. 34,949		::::::	: ::	:-	: -	6/3	:::	7.	% Sr	8	32	179	9
	South E. 34,949	und ov.	: 17: 21:	r ::	:=	: :	:	:::	::	F :- S	8	188	7	9.00
LEEDS	West. 85,186	0.00	:01:10:	: :-	:-	: :	:	::-	:83	8 4,	115	88	333	15.7
CE	₩ 85,	-	:846 : :	- ::	:-	: :	:	:::	:-	ន ::	8	8	8	15
-	North. 58.866	d ov.	111111	: ::	::	: :	64	:::	19	2 84	19	49	226	15.4
		ping 2	:8 :m ::	64 :	::	1 : :	:	:::	::	8 :10	99	51		
	Rc Estimated Population—163,495	Under and over 5.	Smallpox Measles Scarlatina Diplutheria Croup (memb)	Whooping Cough	Diarrhoea .	¶Cholera Rheumatic Fever	Acute & Sub-	Erysipelas Pysemia Fuerperal	Ague Phthisis	Preumonia Pleurisy Heart Disease Injury, &c	Total of above causes	All other)	Total	Mortality per 1,000 per an.
Contin														

TABLE B, Part I. (SUB-DISTRICTS.)

8. .

TABLE OF POPULATION, BIRTHS, AND OF NEW CASES OF INFECTIOUS SICKNESS, coming to the knowledge of the Medical Officer of Health, during the 52 weeks of 1905, in the Urban Sanitary District of Leeds; classified according to Diseases, Ages, and Localities.

_	-				_
the als.	13	латоТ	138848881889918894488488884888848888	293	297
Cases Removed from their Homes in the for Treatment in the Isolation Hospitals	12	Other.	5x5vc-Zvo-vo-24 :4xxx+v :-4 :4 :4xv : : :	539 39	110
Hom on H	11	Erysipelas.	: :0 : := : :0 : :0 : : : : : : : : : :	: :2	12
their	10	Cholera.		:::	:
from t	6	Puerperal.	1:01:::::::::::::::::::::::::::::::::::	::0	0
ed fr	00	Relapsing.	11111111111111111111111111111111111	:::	:
Cases Removed for Treatment in	7	Continued.		1:::	:
es Re Freat	9	Enteric or	:4:200-11-08028 '00 :00 :00 :00 :40 : : :	985	182
	10	Lyphus.	**********************	:::	1
Number of such several Localities	4	Membranous Croup.		:::	:
Loca	00	Diphtheria.	00001240014004014001000010000	282	152
Number of everal Loca	61	Scarlatina.	860888414088410884188881216886865	95	7101
sev N	-	.xod-llem2	2008 : 10148461 : : 22 : 21 : : 4 : : 1 : : 8 : : :	858	8
	00	TVIOT	F388484854858888888888888888888888888888	566 955 949	-
the	-	.латоТ		ವಾಹಹ	2470
coming to the Health.	12	Other,	28487847 9888338 83987 : 100817 : : :	128	347
Meal	=	Erysipelas.	2000 00 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2	882	88
	10	Сројета.	:::::::::::::::::::::::::::::::::::::	:::	:
Locality, Officer of	6	Puerperal.	1:12:14:10:10:14:14:10:11:10:11	: :8	88
100000000000000000000000000000000000000	00	Relapsing.		:::	:
N.B.	7	Continued.	::::::	:::	:
	9	Enteric or Typhoid.	48840E28E85E :40 :60 :47 :60 :67 : : :	2525	289
	10	Typhus.	1 ^{rt}	: □ :	-
es of	4	Membranous Croup.	04 14 1111104 11114 11111111111111111	903 :	00
Cases	00	Diphtheria.	38-58835.245150.44-05150.40580 : : :	110	363
New	61	Scarlatina.	::: 1336.4415282838538524146.8851	266 537 122	925
	-	-xod-liem2	30% :upua8apul : 100 (3 : 14 : 11 ' 18 : 1 :	982	129
	_ (c)	rd.5.	Under 5, 5 under 15, 15 upwards.	Under 5, 5 under 15, 15 upwards.	vi.
	Aged under 5	5 under 15, 15 upwards.	Under 5, 5 under 15, 15 upwards. Under 5, 15 under 15, 15 upwards. Under 5, 5 under 15, 15 upwards.	nder	All ages.
4		15.0	1000 1000 1000 1000 1000 1000 1000 100	15	V
	'st	E Rich	1,870 1,905 1,235 2,140 2,140 2,140 459 459 459	12,337	:
	bered	Regist			
r at	'906I	Estimated of middle of	58,932 85,006 34,817 74,576 65,769 46,686 46,686 39,720	456,787	
all ages.				_	
Population at all ages.		Census, 1901.	59,281 84,340 84,340 69,064 69,064 41,561 17,299 31,845 31,845	428,968	
-		ರೆ"	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 24	
	ties	nese			
Names of Localities adopted for the purpose of these Statistics.		of th	: : : : : : : : : : : : e	:	I tot
		istics istics	East let cck cck cck let rall let let let let let let let let let l	Totals	Grand total
ames adoy pury Stat			North West South-East Hunslet Holbeck Wortley Kirkstall Chapeltown	T	9
	Acres .		X > 0 H H > X H O O		

Notification has been compulsory since the first of May, 1894. The Small-pox Hospital is at Killing'eck, outside the town. New wards for general fever cases were opened on the Small-pox Hospital is a further extensions were made and completed in October, 1904. The use of the Fever Hospital in Beckett Street (the old House of Recovery) was discontinued in November, 1904, the fever cases being treated entirely in the new hospital at Manston, except that, for a short time during the furnishing, the hitherto unused Small-pox hospital at Killingbeck was utilised for the purpose. The wards of the new Small-pox hospital at Killingbeck are a quarter of a mile away from any of the buildings on the Manston estate. The whole of the fever accommodation for Leeds is now in the township of Searcoft, only a few hundred yards from the boundary of the buildings on the buildings on the Manston during 1905. Three cases of small-pox, admitted to Killingbeck Hospital, two cases of tiphtheria, one case of typhoid fever, and one case of "other" disease, admitted to Manston Hospital, cane from outside the city, and are not included in this table. Also fourteen members of the Manston Hospital, and three for typhoid fever, all over 15. None of these fourteen are counted, as they could not be classified under any district of the city. One case of erysipelas, included in the 2,470 notified in 1905, is included in the 1,297 removed to hospital, although the actual removal did not take place until 1906.

TABLE B, Part 2. (Wards).

		lation at ages.		Γ	Nev	Car kno	ses of	Sick ge of	ness the	in ea	ch L	ocali Office:	ty, co	omin Iealt	g to t	he
Names of Localities		283	Aged	1	2	3	4	5	6	7	8	9	10	11	12	13
adopted for the purpose of these	Census,	Estimated to middle of 1905.	under 5, 5 under 15, 15 upwards.	эх.	131	ia.	sno			ever	-		1	· S		
Statistics.	1901.	stima	ao apwards.	Small-pox.	Scarlatina.	Diphtheria.	Membranous Croup.	bus.	Enteric or Typhoid.	Continued	sing	Puerperal.	Cholera.	Erysipelas.	Other.	TOTAL.
(a)	(6)	(c)	(d)	Smi	Sca	Dip	Men	Typhus.	Iyph	onti	Relapsing	nerp	S	Erys	0	T
(4)	(0)	(c)	Under 5,		13		2		1		1			-	4	25
Central	20,996	20,216	5 under 15, 15 upwards.	22	19				20			3		iò	14	33 51
North	38,762	44,096	Under 5, 5 under 15, 15 upwards.		30 62 12	6	::	i	 8 19	::		5	::	2 2 41	12 11 3	60 111 95
North-East	29,084	31,290	Under 5, 5 under 15, 15 upwards.	2 6 20	31 44 10		i		9 7	::		ï		2 22 22	18 10	51 90 73
East	28,297	29,554	Under 5, 5 under 15, 15 upwards.	1 2 13	13 37 9	14			8 29			2	::	2 24	6 6	31 69 85
South	15,047	14,161	Under 5, 5 under 15, 15 upwards.	1 5 13	11 11 4	5 8 2			1 4 3			ï		20	10 3 4	28 31 47
East Hunslet	33,450	37,015	Under 5, 5 under 15, 15 upwards.	1 2	12 26 6	11 19 5	1		3 9 30			i		5 2 26	12 15 3	44 72 73
West Hunslet	29,267	31,712	Under 5, 5 under 15, 15 upwards.	3 3 6	22 34 9	11 22 11	1		2 4 12			··· i·		3 24	18 14 6	60 78 69
Holbeck	27,871	30,664	Under 5, 5 under 15, 15 upwards.	5	18 31 14	7 12 3		::	4 9	::	::	 4		i 17	10 10 2	35 58 54
Mill Hill	7,736	7,136	Under 5, 5 under 15, 15 upwards.		6	252			2			::	::	1 25	5 +	8 17 33
West	23,914	23,683	Under 5, 5 under 15, 15 upwards.	i	15 36 4	4 1 13	1		3 8			3	::	1 2 16	8 7 5	29 50 49
North-West	32,239	34,015	Under 5, 5 under 15, 15 upwards.	4	10 25 14	5 8 5			1 4 14					20 20	2 2 7	20 41 62
Brunswick	22,893	23,034	Under 5, 5 under 15, 15 upwards.	3	12 17 6	9 13 7			3 2 7			i	::	3 i1	5 3 5	32 35 40
New Wortley	18,734	18,516	Under 5, 5 under 15, 15 upwards.	::	12 19 6	1 1 1			2 2			2		 10	622	19 24 23
Armley and Wortley	36,243	40,561	Under 5, 5 under 15, 15 upwards.	2 11	17 28 6	1 1 1	1		3 13			ï		1 2 15	3 7	25 37 54
Bramley	21,650	23,127	Under 5, 5 under 15, 15 upwards.	2	24 83 5	8 2			5 12			ï		1 3 20	12 11 3	41 110 45
Headingley	42,785	48,017	Under 5, 5 under 15, 15 upwards.	4	23 59 17	11 18 13		::	2 17			2		3 1 36	21 19 7	58 99 96
Totals	428,968	456,787	Under 5, 5 under 15, 15 upwards.	9 18 102	266 1 537 1 122	175	6 2	i	12 73 204			28		23 20 337	140 129 78	566 955 949
Grand total				129	925 3	563	8	1	289			28		380	347	2470

TABLE B, Part 2.-Wards (continued).

		lation at ages.		N	umbe seve	r of	such ocal	Cas	ses R for T	emo	ved f	rom in Ise	their olatio	Hor n He	mes i	in the
Names of Localities		999	Aged	1	2	3	4	5	6	7	8	9	10	11	12	13
adopted for the purpose of these Statistics.	Census, 1901.	Estimated to niddle of 1905	under 5, 5 under 15, 15 upwards. (d)	Small-pox.	Scarlatina.	Diphtheria.	Membranous Croup.	Typhus.	Enteric or Typhoid.	Continued a	Relapsing.	Puerperal.	Cholera.	Erysipelas.	Other.	TOTAL
Central	20,996	20,216	Under 5, 5 under 15, 15 upwards.	9	10 18 2			.,	6 14			 i	::		4	17 28 30
North	38,762	44,096	Under 5, 5 under 15, 15 upwards.	9	22 45 8	6 12 3		**	6 12						5 2 3	33 65 35
North-East	29,084	31,280	Under 5, 5 under 15, 15 upwards.		23 37 9	4 6 2			7 6					2	2 8 7	31 64 46
East	28,297	29,554	Under 5, 5 under 15, 15 upwards.		10 36 8	10			 8 19				::	2	5 6	23 61 48
South	15,047	14,161	Under 5, 5 under 15, 15 upwards.		9 11 4	3 1	::	::	2 3	::		::		ï	2 3	16 21 24
East Hunslet	33,450	37,015	Under 5, 5 under 15, 15 upwards.	1 2 3	9 17 4	2			1 6 19			ï		i	9	13 35 27 20
West Hunslet	29,267	31,712	Under 5, 5 under 15, 15 upwards. Under 5,	6	12 27 8 17	5 3 4			3 9	::		::		4	1 3	39 33 21
Holbeck	27,871	30,664	5 under 15, 15 upwards. Under 5,	4	25 9 3	10 2	1000		2 6			i			4 1 3	41 23 8
Mill Hill	7,736	7,136	5 under 15, 15 upwards. Under 5,		1	3	::				::	::			33	7 4 11
West	23,914	23,683	5 under 15, 15 upwards. Under 5,	i	31 3	1 1 4			1 3 1					ï	5	34 13
North-West	32,239	34,015	5 under 15, 15 upwards. Under 5,	4	18 9 7	2 1 5			4 4 2						1 2 2	25 20 16
Brunswick	22,893	23,034	5 under 15, 15 upwards. Under 5,		9 4	7 1 1			1 4						3	20 14 12
New Wortley	18,734	18,516	5 under 15, 15 upwards. Under 5,	2	15 6 10	1			2			ï			1 1	17 9
Armley and Wortley		40,561	5 under 15, 15 upwards. Under 5,	ii 	18 4 22 67	4			8						2	20 25 27 81
Bramley	21,650	23,127	5 under 15, 15 upwards. Under 5,	2	5 19	7 1 2 2		7	5	**		i 		`i	2	15 21
Headingley	42,785	48,017	5 under 15, 15 upwards. Under 5,	4 9	39 12 201	7 3 51		::	9	::					1 4 26	48 32 293
Totals	428,968	456,787	5 under 15, 15 upwards.	18 99	414 95	80 21	::	::	55 121			5	::	12	39 45	606 398
Grand total				126	710	152			182			5		12	110	1297

B. 3.

New cases of Infectious Sickness heard of in the several Sub-districts and Wards of the City of Leeds during the thirteen weeks ended 1st April, 1905.

	tilliteen	Where	Small-	Scar-	Diph-	Mem-	Ty-	Ty-	Con-	Puer-	Erysi-	Cho-		
		treated.	pox.	let fever.	theria.	bran- ous croup.	phus fever.	phoid lever.	tinued fever.		peias.	lera.	Other.	TOTALS.
	North	(Hosp. (Home	38 I	20 2	6	2		22					7 12	93 137
	West	Hosp.	7	40 12	8			8			34		13	81 146
	South-East	Hosp.	26	16	8			12		2	7		3	64 86
TS.	Hunslet	Hosp.	20	31	6			15					5 27	77 180
RIC	Holbeck	(Hosp.	5	13	30			3			24		I	36) 6.
SUB-DISTRICTS.	Wortley	Home Hosp.	12	30	3			4			7		13 I	50 75
JB-I	Kirkstall	(Home (Hosp.	4	25	2 I	1		I		1	9		5 2	23)
SI		(Home (Hosp.	 I	8	3			4			10		19	52) 5
	Bramley	(Home		4				2			I		I	8) 24
	Chapeitown	(Home	4	25 13	10			2 I			7		9	33 73
	Osmondthorpe	Home												:::}
_		CII												
01	Central	{Hosp. Home	9 I	2	2 2	1		13		2	3		I	10 37
	North	Hosp. Home	5	27 13	5 7			7 4			12		3	47 94
	North-East	Hosp.	28	14 1	I	 I		5			8		3	5; 72
	East	Hosp. Home	16	13	7 3			11		2	7		2 2	49 68
	South	Hosp.	18	14	3			3			***			38)
	Fact Hunslet	(Hosp.	3	12	- 4 I			10			7		4	30) 70
	West Hunslet	Home Hosp.	10	13	4			7 3			10		_ 5 _ I	31 83
S.		Home Hosp.	4	9	8	I		3			8		20 I	52 5 03
WARDS.	TIOIOCCK	Home Hosp.	1	- 1 I	3			I			6		13	25) 34
W,	Militaria	Home Hosp.		2					7.1		10		1 I	4 18
	11 Cat	Home		14	7			3			8		5	20 3 43
	Trottii- trest	Hosp. Home	4	18	3			4 5			8		1 4	²⁹ ₂₅ 54
	Dittionated	Hosp. Home	2 I	8	6								3	12 36
	New Wortley	Hosp. Home		9	1		***	I			2		2	11 16
	Armley & Wortley		II	8			***	2					I	22) 25
	Bramley	(Home	2	21	5			5			4		2	33) 48
	Headingley	Hosp.	4	7 26	- 1			1		***	4		2	15)
		Home		II	II						10		19	34 87
		Hosp.	117	213	46			71					20	467 870
	CITY	Home	3	67	76	4		29		6	116		102	
		Cases	120	280	122	4		100		6	116	***	122	870

Three nurses and one maid at Manston Hall contracted scarlet fever, and one nurse contracted typhoid fever, and all were treated in the hospital. They are not included in the 213 and 71 given above. Manston Hall is outside the city.

New cases of Infectious Sickness heard of in the several Sub-districts and Wards of the City of Leeds during the thirteen weeks ended 1st July, 1905.

		Where treated.	Small- pox.	Scar- let fever.	Diph- theria.		Ty- phus fever.		Con- tinued fever.	peral	Erysi- pelas.	Cho- lera.	Other.	TOTALS.
	North	Hosp. Home		14	4 8			3 6			12		3 2	²⁴ 37) 61
	West	Hosp.		15	10	 I		3 7		···	12		6	34 85
	South-East	Hosp.	1	13	 I		***	6			6		5 3	25 41 16 41
TS.	Hunslet	Hosp.	2	23	2 9			9 5			I 19		3	40 92 52 92
TRIC	Holbeck	Hosp.		18	6	***		1 1						25 \ 17 \ 42
SUB-DISTRICTS.	Wortley	∫ Hosp.	2	20 20	I					1	5		3	26) 57
UB-1	Kirkstall	Hosp.		26	2			2			10	***	9	31) (0
S	Bramley	Home Hosp.		2 I	6			I		I	10		18	38) 69
		(Home		17	6						3		3	0)
	Chapeltown	Home		11	8			2			9		_ 4	35)
-	Osmondthorpe	Home									1			} 1
-	((Hosp.		-	2									6)
	Central	(Home	-	1	3	1		3			3			11) 17
	North	(Home	1	6	7 8		1	4			12		3	$\binom{26}{35}$ 61
	North-East	{Hosp. Home		14	3			I			6		3	18 40
	East	{Hosp. Home		7 2				5 4			6		4 2	16 31
	South	{Hosp. Home	1	8	I			I			7		I	12 21
1	East-Hunslet	/ T.T.		8	8			6 3			7		2 4	16 42
	West-Hunslet			13	5			3			1	***	1	25 51
DS.	Holbeck	Hosp.		18	2			I			6			21 \ 21
WARDS	Mill Hill	∫Hosp.			- 2 I			I		1	5		I	2) 11
1	West	∫ Home ∫ Hosp.		3			***	I	***		6	***	2	9)
	North-West	(Home	1	8	1	1		2 2		I	2		9 2	13 28
		(Home		3	3 8			2 I			4		3	15)
	Brunswick	(Home		6	4			2			***		2	14 28
	New Wortley	Home		4				2			3		1 2	5 13
	Armley & Wortley	(Frome		11 6				2			6		6	16 36
	Bramley	{Home		6 3							4		1 2	7 16
1	Headingley	{Hosp. Home		27	7			2		 I	10		18	3 ² 39) 71
					-									377
	Стту	∫Hosp. Home	6	147 47	31 44	2		24 30		4	87		25 69	234) 518
	0111 111211111	Cases	6	194	75	2	I	54		4	53		94	518

A nurse contracted scarlet fever, and a nurse and a maid contracted t phoid fever, and all were treated in the hospital. They are not included in the 147 and 24 given above. Manston Hall is outside the City.

New cases of Infectious Sickness heard of in the several Sub-districts and Wards of the City of Leeds during the thirteen weeks ended 30th September, 1905.

		Where treated.	Small- pox.	Scar- let fever.	Diph- theria.	Mem- bran- ous croup.	Ty- phus fever.	I y- phoid fever.	Con- tinued fever.	Puer- peral fever.	Erysi- pelas.	Cho- lera.	Other.	TOTALS.
	North	Hosp.		18	6			1 I 2		I 2	I 15		12 1	49 75
	West	Hosp.		20	4			4 6		2	I 22		5 2	34 85
	South-East	Hosp.		16 1	9			6 2			5		3 5	36 so
TS.	Hunslet	Hosp.		15	3			9 2		2	2 14		4	33 77
FRIC	Holbeck	Hosp.		8	3 2			2 3		I _ 1	2		I	15 25
DIST	Wortley	Hosp.		38	2 I			4		- I			2 3	47 71
SUB-DISTRICTS.	Kirkstall	Hosp.		6	4			5			8		2	14 42
S	Bramley	∫ Hosp.		17	3			5 2						22 \ 28
	Chapeltown	(Home ∫Hosp.		13	9			I			5		2	25 \ 45
	Osmondthorpe	Home Hosp.		8	3			-1			7			}
-		(Home											***)
	Central	Hosp.		7	I			3		I	3		4	16 21
	North	Hosp.		13	6			6		2			3	28 56
	North-East	Hosp.		II	8			3			13		7	30 42
	East	Hosp.		16	8			6		I	I		3	34 42
	South	∫ Hosp.		2	I			2			1 I			4) 14
	East-Hunslet	\ Home ∫ Hosp.		4	1			4		I	3 1		5	14) 25
	West-Hunslet	(Home ∫Hosp.		9	7			5	-:-		8			17) 20
S.	Holbeck	∫ Hosp.		5 8	3			2		I	5		I	$\begin{bmatrix} 22 & 39 \\ \hline 15 \\ 7 & 22 \end{bmatrix}$
WARDS.	Mill Hill	∫ Home ∫ Hosp.		2	2					I	I			2) 12
=	West	∫ Home ∫ Hosp.		12	I						8		1	15) 20
	North-West	∫ Home ∫ Hosp.		5	4 1			1 2		I	6			15) 30
	Brunswick	\ Home ∫ Hosp.	-	3	3			3			6		4	16) 24
	New Wortley	∫ Home ∫ Hosp.	-:	13	4 I			2 I		I	2			11 5 20
	Armley & Wortley	Home		6	1			2		I			2	8)
	Bramley	(Home		7 36	4			4		 I	3		1 2	15/ 23
		∫ Home ∫ Hosp.		6	4			4			6			95 33
-	Headingley	(Home		7	5			5		1	8		2	28 42
_		(Hosp.		151	43			43		3	6		29	275)
	Стту	Home		47	44			26	:	10	82		14	275 498
_		Cases		198	87			69		13	88		43	498

A nurse contracted scarlet fever, and another nurse contracted diphtheria. Both were treated in the hospital. They are not included in the 151 and 43 given above. Manston Hall is outside the City. In addition to the 275 cases hospitalled (out of 498 reported during the quarter) I case of scarlet fever, which had been reported in the previous quarter, was also taken to hospital.

B. 6.

New cases of Infectious Sickness heard of in the several Sub-districts and Wards of the City of Leeds during the thirteen weeks ended 30th December, 1905.

		Where treated.	Small- pox.	Scar- let fever.	Diph- theria.	Mem- bran- ous croup.	Ty- phus fever.	Ty- phoid fever.	Cen- tinued fever.	Puer- peral fever.	Erysi- pelas.	Cho- lera.	Other.	TOTALS.
	North	Hosp.		38	6 5			5 3		 I			6 5	55 82
	West	Hosp.	***	29	6	***		6			13		12	53 93
	South-East	Hosp.		18	4 3			6 3			7	***	6	35 48
TS.	Hunslet	Hosp.		18	3 9	2		9 7		1	2 12		5	38 85
SUB-DISTRICTS	Holbeck	Hosp.		12	3 3			2 I			7		4 3	21 23 44
DIS	Wortley	Hosp.		15	I			5		I	13		23	22 42 64
SUB-	Kirkstall	{Hosp. Home		11	5			3			12		2	21 32 53
0,1	Bramley	Hosp.		28	7 2 1			2			1 6		3	33 51
	Chapeltown	{Hosp. Home	3	29	2 8			6			5		I 3	41 64
	Osmondthorpe	{Hosp. Home												}
)
	Central	{Home		19	2 2			4 3						²⁶ ₈ 34
	North	{Hosp. Home	3	21 4	3 7			3		 I	8		3	$\binom{3^2}{23}$ 55
	North-East	{Hosp. Home		30	3			4			5		4 5	41 60
	East	{Hosp. Home		18	4 2			5 2			7		5	33 44
	South	Hosp. Home			3			2			2		3 4	7 16
	East-Hunslet	{Hosp. Home		6 I	4	 I		6 5		I	7		9	15 42
	West-Hunslet	{ Hosp. Home		12 1	1 5			2 2			2 4		2 2	19 34
SDS.	Holbeck	Hosp. Home		12	3			2 I		 I	6		3	20 40
WARDS	Mill Hill	Hosp. Home		3 2	2						2		8	11 6 17
	West	Hosp. Home		16	I 5			3		 I	2		1	18 34
	North-West	Hosp. Home		2 3	3 2			I						6 17
	Brunswick	Hosp. Home		5	2 2			5					3	15 23
	New Wortley	Hosp. Home		6 I						1	5			7 16
	Armley & Wortley	C		7 4							5			12 22
	Diamie,	(Home		30 8	3			2 4			9			$\binom{36}{41}$ 77
	Headingley	Hosp. Home		9	5 7			3			12		2 3	21 32 53
		/ II		***										
	Сіту	Hosp. Home	3	198 55	32 47	2		44 22		3	84	:::	36 52	$319 \atop 265$ 584
		Cases	3	253	79	2		66		5	88		88	584

Four nurses at Manston contracted scarlet fever and were treated in the Hospital They are not included in the 198 given above. Manston Hospital is outside the city.

B. 7.

New cases of Infectious Sickness heard of in the several Sub-districts and Wards of the City of Leeds during the thirteen weeks ended 31st March, 1906.

		Where treated.	Small- pox.	Scar- let fever.	Diph- theria.	Mem- bran- ous croup.	Ty- phus fever.	Ty- phoid fever.	Con- tinued fever.	Puer- peral lever.	Erysi- pelas.	Cho- lera.	Other.	TOTALS.
	North	Hosp.		21	15			9		2	14		7 6	5 ² 33) 85
	West	Hosp. Home		25 17	9	 I		3 7		2	2 23	***	12	51 115
	South-East	Hosp. Home		6 2	2 4	2		3					4	15 35
CTS.	Hunslet	Hosp.	2	18	10	2		15		2	3 14		5 5	53 108
SUB-DISTRICTS	Holbeck	Hosp. Home		13	1 4			4			6		6 8	24 18) 42
-DIS	Wortley	{Hosp. Home		2I IO	6			5 1			I 14		4 I	33 65
SUB	Kirkstall	{ Hosp. Home		14 7	29 20			2		1	7		11 5	55 96
	Bramley	{Hosp. Home		19	I			I 2			4		3	24) 10) 34
	Chapeltown	{Home		10	3			1 4		 I	8		2 I	16 28 44
	Osmondthorpe	Hosp.												}
		(Hosp.												
	Central	Home (Hosp.		12	4			I			3		5	15) 36
	North	Home (Hosp.		5 2	9 7			3			9		I	23 40
	North-East	Home Hosp.		3	7			7 3		2	9		4	30 50
	East	(Home		6 2	4	2		3			9		4	15 35
	South	Hosp. Home		1	4			3						8 12
	East-Hunslet	Hosp. Home	2	10	13			6			10		2	$\frac{24}{34}$ 58
1.6	West-Hunslet	Home	-	2	5 3	1		6		2	4		3	26 48
WARDS.	Holbeck	Hosp. Home			4			3			5_		5	19 32
WA	Mill Hill	(Hosp.		2	2 I			1			5_		6	9) 19
	West	Hosp. Home Hosp.		7	3	 I		3			9		3	18 44
	North-West	Hosp. Hosp.		8 2	4			2			6	·	I	12 27
	Brunswick	Hosp.		6	1 1			2 I		2	4		2 I	10 25
		Home Hosp.		4	I						7		I	6) 16
	Armley & Wortley	Home Hosp.		9	4						6		3	19 43
		Home Hosp.		3	2			3			5		3	13 40
	Headingley	Home		7	29			2			7		5	56 99
		Hosp.	2	147	72			41		I	6		54	222)
	CITY	Home		53	76	5		30		7	99		31	323 624 301 624
		Cases	2	200	148	5		71		8	105		85	624

A nurse at Manston Hall contracted scarlet fever, and was treated in the hospital. The case is not included in the 147 given above. Manston Hall is outside the city. In addition to the 323 cases hospitalled (out of the 624 reported during the quarter) 1 case of erysipelas which had been reported in previous quarter was also taken to hospital.

o TABLE

Table shewing deaths recorded in the City of Leeds during the fifty-two weeks ended 30th December, 1905, classified according to cause, age, and the registration sub-districts to which the patients belonged. Deaths in institutions allocated districts to which patients belonged. to . ..

Annua 00,00 0,00 rate 0.00 0.30 0.08 0.54 0,05 0,05 0.03 \$0.0 0.02 10,0 10,0 10,0 S 17 10 14 52 30 6 4 3 198 333 W 10 38 12 85 2 23 FOTAL City. over 2 4 4 5 100 10 0 0 04 H 20 E under 1 100 1 350 8 " Ĉŧ. : 00 > 500 71 occurring in over . H outsiders : : : 5 Deaths City. ander 5 over Osmondthorpe. 358 under over Chapeltown : 61 + 68 : 100 . 10 39,720 under w ; 04 : : 09 0. 4 4 : : over 100 Bramley. 18,435 under ,E9 : : : : : * : 00 over : 01 × Kirkstall. 46,685 under + : 0 over = : 04 m 9 Wortley. 63,769 under н 15 1 :: 2 H 60 H : # -Holbeck. 34,489 0 . 04 : 10 .0 -01 Hunslet. 74,576 20 : : : : : : : : : : 00 . 00 19 00 South E. 34,817 2: : 99 over . 0 H + 60 OE. - 101 : 0 - : 0 LEEDS. West. 85,005 under : : 2 10 : 8 :::00 69 - 60 O O : C : H O North. 58,932 under 00 : : : 0 : # # 10 : 92 2000 Rheumatic fever ... urethra Estim. Pop., 456,787 Zoogenous diseases TOWNSHIPS, Cholera (English) Diarrhœa, &c. .. Under and over 5 Erysipelas
Pyæmia ...
Phagedena ...
Septiczemia Phlebitis ...
Puerperal fever Parasitic diseases Whooping-cough Diphtheria rheumatism Starvation, &c. Small-pox Chicken-pox Measles ... Scarlet fever Other or doubtful Syphilis ... Gonorrhœa Stricture of u Rheumatism Influenza Rickets Ague Gout Contin. Fevers.

			LEEDS.	DS.		-		-		-		-		-		-		-		- D	Deaths of	-	Torra		-
TOWNSHIPS,	North.	÷	West.	St.	South E.	점	Hunslet.	let.	Holbeck.	ock.	Wortley.	sy.	Kirkstall.	all.	Bramley.		Chapeltown.		Osmond- thorpe.		outsiders occurring in City,		mortality in City.	ty	Annum
Under and over 5	under 5	over 5	under	over 5	under 5	over 5	under	over 1	under 5	over to	under o	over tu	under	over ur	under	over u	under o	over un	under over 5	er under	er over	under	r over	all	per 1,000
9		11148418110	-5:05:::::::	:::+00:0:0:=	100 10 4 11111	11181411110	02:04:14:11	:: 10 + 8 : 8 : + +	-00 : : W : : F : F :	1117119119	(2 = NW)	11461010110	+ w : a w : : : : . w :	1::0:4:0::=	:0:HH::::::	:::-:::-::-	10116111111		1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -	1000070000		10 8 - 4 00 + ; m ; m ;	:: 040 # + 4 + 0 8	200000000000000000000000000000000000000	10,00 11,00 10,00
Dis. of layer Dis. of bowels	: : :	- 00	; - -	2 7 :	:- :	m m =	: "	- 0 :	: + :	in +	: m :	n h m	11 1	- m +	111	; a =	. 01	m = :	11 1	: 07 -	mog +	: 2 0	0.00	0,00 9	0.09
Nephritis Kidney disease Albuminuria Other urinary	26 1 1 1	I C : 2	::::	77 2 2 2	m : : :	00 + ; ;	1171	±∞ :-	::::	4 m ; =	-:::	==:-	:::::	00 g : m	1111	**!!	1111	0 m : N	1111	1111	++ :+	9:::	80 5 2 ±	20 a ±	0.21
Dis. of generative Organs	:	71	1	13	:	64	1.5	6	:	-	:	+		-	:	-	3	н			7	:	35	35	80.0
Childbirth Disease of bones (15), 1	60	1	Ĉ4	7	64	.01	:	~	-	0	-	10	-	-		-	:	-		:		01	60	80 (%)	50.0
Joints (o), arthritis (8) J Ulcer: Phlegmon (o) Skin disease	1 1 1	m + ;	: -	0 8:	1 11	m ; ;	: 17	m ::	2 2 3	- ::	: 1.4	1 11	: ::	+	1 11	- ::	: ::	: ::	1 11	3 :-	N ::	: : 5	2 4 H	₩ ₩ 400	0.00
Injury Lead poisoning	= :	**	r ;	60 N	7	91	0 :	00 :	M3 +	60 :	0 :	E :	m :	0 =	-:	2 :	N ;	9 :		-:	17	98 :	217	273	09.0
Mortification Debility Marasmus, Atrophy Tumour Abscess	0 D 0 : 1	40:04	1: 20.11	+ 10 ; + 10	:28::	H 4 : 4H	: 50 H : 0	HH : 40	: +:: : "	a = ; = 0		+ + ; m n	10 m ; ;	H::N:	::10::	H = : 10 10	: 57 : :	; ; ; m a	11111	1:9::	H ; ; # N	E 4 1 : 0	27.88	16 141 33 28	0.04
	1	:	1	:	:		;	:	1	:	:	1	-	1	:	:	:	1	:	:	;	**		-	00,0
Total under 5 and over 5	440	9009	399	168	350	421	500	125	237	357	364	579	159	352	69	177	148 3	317	4	17	191	2,694	4,430	7,124	15.65
	oto,I	07	I,S	1,290	1	771	1,080	00	594		943		511	м	245	10	465		9		178		7,124		3
Mortality per 1,000 l	2.21	1	15.5	.01	22,5	2	5,11	10	17.3		0°.+1		0.11		13,4		11.0		8-94		:	6.05	0,11	15.6	:

This table is printed for the first time in the report for 1892. In it the causes of death are more detailed than in the other tables. They are classified as in Table A, part 2, according as the deaths were those of persons under or over the age of five. All the deaths which occurred in the following public institutions: the Infirmary, the Women and Children's Hospital, the Borough Fever and Small-pox Hospitals, the Leeds, Hunslet, Holbeck, and Bramley Workhouses, have been classified under the districts to which the patients belonged. The 178 deaths of persons who belonged to no district in the city have been separated in two columns by themselves, as deaths of outsiders; these deaths are, however, included in the total mortality of the city. As far as possible, the order of the Registrar-General has been followed in the arrangement of this table. The horizontal lines correspond with the groups in the Registrar-General's annual report.

Septicamia includes deaths from pyæmia (4), phlebitis (0), phagedæna (0), septicæmia (not puerperal) (3). Parasitic diseases include thrush (5). Starvation includes purpura hæmorrhagica (4), scurvy (1), privation and want of breast milk (0), malnutrition (18), and inanition (50 deaths). Rheumatic fever in reports previous to 1892 had only the deaths ascribed in those terms to this disease by the medical attendant. Deaths from acute and sub-acute rheumatism had previously been classed under "rheumatism." A separate line has been given, both in Table C and Table A, to prevent confusion and enable comparison. "Rheumatism" includes chronic rheumatism and disease simply described as "rheumatism" (see Report, 1893, page 144). Anæmia includes chlorosis (2), hoemophilia (1), but not leucocythemia; deaths from the latter have been referred to diseases of the ductless glands. In malformations are included cyanosis (2), patent foramen ovale (5), spina bifida (7), atelectasis (22), imperforate anus (4), cleft palate (4), harelip (2), and (28) other congenital defects.

Brain disease includes deaths registered from such causes as cerebral congestion, cerebral homorrhage, and softening of the brain. Meningitis includes diseases classified as meningitis (94), and spinal (1) meningitis, but not tuberculous. Apoplexy includes all apoplexies not otherwise defined. Paralysis includes hemiplegia, paraplegia, and "paralysis." General paralysis (8) is included under insanity, and does not include deaths from "softening of the brain." Convulsions includes diseases so certified, and deaths (0) due to "fits." Fits of apoplexy, &c., come under other headings. We have tried, as far as possible, to keep to the old headings.

Endocarditis, &c., includes valvular disease of the heart. "Heart disease" includes such diseases as hypertrophy, atrophy, fatty degeneration, weak heart, cardiac disease or degeneration and "disease of the heart." Angina pectoris includes only those deaths in which the symptom but no disease is stated. Aneurism includes all the aneurisms so stated. Other diseases of the circulatory system includes atheroma. Other respiratory diseases includes asthma (18), emphysema (18), empyæma (12), pulmonary congestion (5), "lung disease" (0), and others (11). Tonsillitis, &c. includes pharyngitis (0), parotiditis (0). Diseases of the stomach includes dyspepsia, hoematemesis, gastritis. Disease of liver includes hepatitis (2). Diseases of the bowels includes melœna (0), ulcer of intestines, obstruction of bowels, strangulation not due to hernia, intussusception, appendicitis. Kidney disease includes deaths from granular kidney (6), Bright's disease (47), other kidney diseases (18), and uræmia (5). Albuminuria includes only deaths in which the symptom without any pathological cause was registered. Diseases of the urinary system includes calculus, hoematuria, cystitis and other diseases of the bladder. Disease of the generative organs includes uterine disease (1), ovarian disease (6), and "other diseases" of the generative organs, male (25), female (3). Childbirth includes all the accidents of parturition, except puerperal fever.

Diseases of the bones and joints includes disease of the spinal column, but not, of course, such diseases as spinal sclerosis, which are now referred to disease of the nervous system. A comparison of mortality previous to 1890 is difficult, as the term spinal disease was used to include both diseases of the spinal column and of the spinal marrow. Abscess includes cellulitis (3), carbuncle (4). Injury includes deaths from accident or negligence, homicide, misadventure, suicide, and execution. The death from "other causes" was that of a newly-born male infant found dead on the 8th May, at Cross Cliff Road, Headingley. The Coroner's jury had not sufficient evidence to show whether the death was due to natural causes or no.

TABLE D, Part 1.—Shewing death-rates from certain causes for the years 1890 to 1905.

			SEVEN	COMMON		ZYMOTICS.					
	Small-pox.	Measles.	Scarlatina.	Diph- theria.	Whooping cough.	"Fever."	Diarrhœa.	All seven.	Croup (mem- branous and undefined)	Phthisis.	Influenza and diseases of the air-passages other than
1890 (53 wks.)	00.0	0.27	0.28	20.0	0.20	0.30	86.0	2.39	60.0	99.1	2.62
1891 (52 wks.)	0.00	0.71	0.18	0.04	0.41	0.50	98.0	2.41	80.0	62.1	11.9
1892 (52 wks.)	0.05	0.50	0.50	80.0	0.45	0.17	01.1	2.18	0.13	1.42	4.56
1893 (52 wks.)	0.08	06.0	80.0	0.15	0.44	0.30	09.1	3.55	0.18	1.70	4.60
1894 (52 wks.)	10.0	0.75	0.13	0.15	0.34	0.14	0.45	86.1	21.0	1.49	3.64
1895 (52 wks.)		0.35	0.13	0.10	0.56	0.22	1.58	2.65	0.12	1.55	4.34
1896 (53 wks.)	00.0	0.48	0.18	01.0	09.0	0.51	69.0	2.27	60.0	1.50	4.02
1897 (52 wks.)	:	0.40	0.23	0.13	0.24	0.51	1.58	2.79	0.10	1.44	3.60
1898 (52 wks.)	0.00	0.45	0.56	0.46	0.39	0.23	1.24	3.10	0.12	1.39	3.41
1899 (52 wks.)		0.37	0.15	12.0	0.38	0.17	96.0	2.73	0.12	1.41	3.72
1900 (52 wks.)	00.00	0.58	0.12	0.55	0.39	0.50	60.1	2.93	90.0	141	4.19
1901 (52 wks.)	:	0.58	61.0	0.38	0.33	61.0	1.47	3.14	20.0	1.41	3.46
1902 (53 wks.)	10.0	0.43	0.13	0.18	0.46	0.18	19.0	2.00	0.02	1.31	3.52
1903 (52 wks.)	90.0	0.28	0.25	0.14	0.57	0.13	0.62	1.74	0.03	1.27	3.00
1904 (52 wks.)	00.0	22.0	0.13	60.0	0.46	0.11	10.1	2.57	0.02	1.40	3.21
1905 (52 wks.)	10.0	0.24	60.0	80.0	0.27	0.12	08.0	09.1	0.03	1.23	3.07
				See note	to table	D. part 2.					

See note to table D, part 2.

TABLE D, Part 2. For whole District.

	0	Birth	is.		under of age.		s at all Total.		Deaths	Deaths	Death ages	s at all net.
Year.	Population estimated to middle of each year.	Number.	Rate.*	Number	regis- tered.	Number		in Public Institu- tions.	of non- resid'nts regis- tered in district.	tered beyond district.		
1	2	3	4	5	6	7	8	9	10	11	12	13
1890	362,768	12,336	33'4	2,128	173	8,370	22.7	819	132	No return	8,238	22'4
1891	369,099	12,538	34.1	2,216	177	8,429	22'9	869	144	do.	8,285	22.2
1892	375,540	12,546	33.2	2,114	168	7,403	19.8	789	129	do.	7,274	19.4
1893	382,094	12,348	32.4	2,542	206	8,512	22'4	874	147	24	8,365	22.0
1894	388,761	12,502	32.3	1,945	156	6,935	17.9	772	142	7	6,793	17.5
1895	395,546	12,478	31.7	2,384	191	8,101	20.6	882	167	No return	7,934	20'I
1896	402,449	12,573	30.8	2,120	169	7,682	18.8	908	161	8	7,521	18.4
1897	409,472	12,912	31.6	2,454	190	8,148	20.0	881	175	1	7,973	19.5
1898	416,618	12,971	31.5	2,372	183	7,996	19.3	940	142	9	7,854	18.9
1899	423,889	12,939	31.5	2,222	172	8,105	19.2	1,005	181	26	7,924	18.8
1900	431,287	13,091	30.2	2,397	183	8,619	20°I	1,084	184	90	8,525	19.8
1901	430,630	12,898	30.1	2,429	188	8,283	19.3	1,176	171	92	8,204	19.1
1902	437,341	13,245	29.8	2,113	160	7,814	17.6	1,154	211	96	7,699	17:3
1903	443,559	12,996	29.4	1,992	153	7,334	16.6	1,094	184	113	7,263	16.4
1904	450,142	12,561	28.0	2,207	176	8,096	18.0	1,185	173	116	8,039	17.9
Averages for years 1895- 1904		12,866	30'4	2,269	176	8,018	19.0	1,031	175		7,894	18.6
1905	456,787	12,337	27.1	1,875	152	7,124	15.6	1,225	178	102	7,047	15.2

^{*} Rates calculated per 1,000 of estimated population. The populations given in column 2 for the 11 years, 1890 to 1900, are those estimated by geometrical progression at the rate of increase which obtained between censuses of 1881-1891. That for 1901 has been estimated similarly from the rate which prevailed between the censuses of 1891-1901. To make the populations for 1900 and the previous years correspond they should be decreased respectively by the following percentages: (for 1900) 1'72, (for 1899) 1'53, (for 1898) 1'34, diminishing by about 0'19 each year to (for 1892) 0'21, and (for 1891) 0'05. The rates in columns 4, 8, and 13 must be correspondingly increased by these percentages (see Annual, 1901, and table on p. 58).

The deaths included in column 7 of this table are the whole of those registered during the year as having actually occurred within the City or in the fever hospital at Seacroft. The deaths included in column 12 are the numbers in column 7, corrected by the subtraction of the numbers in column 10 and the addition of the numbers in column 11. Column 11 is however, incomplete in some of the years. Deaths in fever hospitals belonging to but outside the City are, unless those of strangers, included in column 7.

By the term "Non-residents" is meant persons brought into the district on account of illness, and dying in some public institution, and paupers from townships outside the City dying at the Workhouse; and by the term "Residents" is meant persons who have been taken out of the district on account of illness, and have died in some public institution, other than our own fever hospitals, and inmates of the County Asylums from the City townships.

TABLE E. VITAL STATISTICS FOR 1905.

The following Births and Deaths were recorded in the several Sub-Registration Districts of the City of Leeds during the fifty-two weeks ended 30th December, 1905. The figures in italics after the Births and Deaths give the proportion per annum per 1,000 of the estimated population.

Districts.	Births.	Birth Rate.	Deaths.	Death All causes.	Rate. 7 Zymotics.
S (North	 1,870	31.84	1,040	17.71	2.49
West	 1,905	22.49	1,290	15.23	1.24
South-East	 1,235	35.59	771	22.22	3.14
Hunslet	 2,140	28.79	1,080	14:53	1.98
Holbeck	 1,075	31.28	594	17.28	1.80
Wortley	 1,653	26.01	943	14.84	1.37
Kirkstall	 981	21.08	511	10.08	0.69
Bramley	 459	24.08	246	13:39	1.25
Chapeltown	 1,016	25.67	465	11.75	0.63
Osmondthorpe	 3	8.41	6	16.82	
Outsiders	 		178		
Totals	 12,337	27.10	7,124	15.65	1.62

Considered as occurring in the Municipal Wards, the foregoing Deaths are classed as follows:—

Wards.	I	Deaths.	Death	Wards.		Deaths.	Death
Eastern Div	ision.		Rate.	Western Di	vision.		Rate.
Central		319	15.83	Mill Hill		105	14.76
North		573	13.04	West		437	18.51
North-East		591	18.96	North-West		462	13.63
East		638	21.66	Brunswick		316	13.77
South		280	19.84	New Wortley		313	16.96
East Hunslet		534	14.48	Armley		580	14'35
West Hunslet		458	14'49	Bramley		296	12.84
Holbeck		510	16.69	Headingley		534	11.16

In both these tables deaths occurring in public institutions have been referred to the districts to which the patients belonged. The births in workhouses are included in those of the districts in which these institutions are situated. There were eighty-five deaths at Manston Hospital, and three deaths at Killingbeck Hospital, during the year.

The rates are calculated for the year on populations estimated by the Registrar General's method, described at p. 88 of the Annual for 1904. Those for several quarters are on populations as previously estimated.

The following Births and Deaths were recorded in the several Sub-Registration Districts of the City of Leeds during the thirteen weeks ended 1st April, 1905. The figures in italics after the Births and Deaths give the proportion per annum per 1,000 of the estimated population.

Districts.	Births.	Birth Rate:	Deaths.	Death Rate. All causes. 7 Zymotics				
o (North	 505	34.7	282	19:4	1.0			
West South-East	 517	24.5	352	16.7	0.4			
South-East	 312	36.3	194	22.6	1.2			
Hunslet	 549	30.0	241	13.2	0.7			
Holbeck	 285	32.9	134	15.5	0.0			
Wortley	 434	27.7	240	15:3	0.8			
Kirkstall	 276	24'3	154	13.6	1.0			
Bramley	 113	25.4	60	13.5	2.0			
Chapeltown	 263	23.8	131	11.8	0.3			
Osmondthorpe	 I	10.6						
Outsiders	 		31					
Totals	 3,255	28.6	1,819	16.0	0.8			

Considered as occurring in the Municipal Wards, the foregoing Deaths are classed as follows:—

Wards.		Deaths.	Death	Wards.	Deaths.	Death
Eastern Division.			Rate.	Western Div	Rate.	
Central		89	17.8	Mill Hill	 33	18.4
North		149	13.8	West	 114	19.6
North-East		166	19.6	North-West	 130	15.3
East		154	21.3	Brunswick	 85	15.0
South		69	19.5	New Wortley	 73	16.0
East Hunslet		113	12.5	Armley	 153	15.4
West Hunslet		114	147	Bramley	 74	13.1
Holbeck		112	14'5	Headingley	 160	13.0

In both these tables deaths occurring in public institutions have been referred to the districts to which the patients belonged. The births in workhouses are included in those of the districts in which these institutions are situated. There were twenty-seven deaths at Manston Hospital and three deaths at Killing beck Hospital during this quarter.

Central ward deaths read 88, Headingley 161; death-rates 17'6 and 13'1.

The following Births and Deaths were recorded in the several Sub-Registration Districts of the City of Leeds during the thirteen weeks ended 1st July, 1905. The figures in italics after the Births and Deaths give the proportion per annum per 1,000 of the estimated population.

Districts.	Districts.		Birth Rate.	Deaths.	Death Rate. All causes. 7 Zymotics				
North		498	34'3	219	15.1	0.8			
West		491	23.3	308	14.6	0.4			
South-East		335	39.0	178	20.7	0.5			
Hunslet		536	29:3	264	144	0.6			
Holbeck		258	29.8	161	18.6	1.5			
Wortley		441	28.1	217	13.8	0.7			
Kirkstall		251	22·I	140	12.4	0.5			
Bramley		136	30.5	68	15:3	0.5			
Chapeltown		267	24'I	110	9.9	0.1			
Osmondthorpe		1	10.6	I	10.6				
Outsiders				54					
Totals		3,214	28.2	1,720	15.1	0:6			

Considered as occurring in the Municipal Wards, the foregoing Deaths are classed as follows:—

Wards.		Deaths.	Death	Wards.		Deaths.	Death
Eastern Divi	Eastern Division.		Rate.	Western Div	ision.		Rate.
Central		76	15.2	Mill Hill		17	9.5
North		125	11.6	West		113	19.4
North-East		125	148	North-West		118	13.8
East		153	21.1	Brunswick		67	11.8
South		55	15.6	New Wortley		75	16.5
East Hunslet		136	15.0	Armley		130	13.1
West Hunslet		113	145	Bramley		80	142
Holbeck		137	17.7	Headingley		146	11.0

In both these tables deaths occurring in public institutions have been referred to the districts to which the patients belonged. The births in workhouses are included in those of the districts in which these institutions are situated. There were twelve deaths at Manston Hospital during this quarter,

The following Births and Deaths were recorded in the several Sub-Registration Districts of the City of Leeds during the thirteen weeks ended 30th September, 1905. The figures in italics after the Births and Deaths give the proportion per annum per 1,000 of the estimated population.

Districts.	Births.	Birth Rate.	Deaths.	Death Rate. All causes. 7 Zymotic					
oi (North	 460	31.6	281	19:3	5.9				
West South-East	 472	22.4	299	14.2	2.8				
South-East	 315	36.7	216	25°I	8.4				
Hunslet	 538	29.4	287	15.7	4.3				
Holbeck	 269	31.1	147	17.0	3.1				
Wortley	 396	25.3	248	15.8	34				
Kirkstall	 262	23.I	106	9.4	I'I				
Bramley	 96	21.6	52	11.7	I.I				
Chapeltown	 273	24.7	103	9.3	1.4				
Osmondthorpe	 		I	10.6					
Outsiders	 		50						
Totals	 3,081	27.1	1,790	15.7	3.6				

Considered as occurring in the Municipal Wards, the foregoing Deaths are classed as follows:—

Wards.	Wards. Deaths.		Death	Wards.	Wards. Deaths.				
Eastern Division.			Rate.	Western Divi	Rate.				
Central		73	14.6	Mill Hill		28	15.6		
North		170	15.8	West		92	15.8		
North-East		142	16.8	North-West		100	11.7		
East		178	24.6	Brunswick		83	14.6		
South		83	23.5	New Wortley		75	16.5		
East Hunslet		129	14.3	Armley		161	16.2		
West Hunslet		124	15.9	Bramley		64	11.3		
Holbeck		130	16.8	Headingley		108	8.8		

In both these tables deaths occurring in public institutions have been referred to the districts to which the patients belonged. The births in workhouses are included in those of the districts in which these institutions are situated. There were twenty-one deaths at Manston Hospital during this quarter.

E. 5.

The following Births and Deaths were recorded in the several Sub-Registration Districts of the City of Leeds during the thirteen weeks ended 30th December, 1905. The figures in italics after the Births and Deaths give the proportion per annum per 1,000 of the estimated population.

Districts.	Births.	Birth Rate.	Deaths.	Death Rate. All causes. 7 Zymotics				
Si (North	 407	28.0	258	17.7	2.4			
West South-East	 425	20.I	331	15.7	1.4			
South-East	 273	31.8	183	21.3	2.7			
Hunslet	 517	28.2	288	15.7	2.4			
Holbeck	 263	30.4	152	17.6	1.6			
Wortley	 382	24.4	238	15.2	0.7			
Kirkstall	 192	16.9	III	9.8	0.3			
Bramley	 114	25.6	66	14.8	1.6			
Chapeltown	 213	19:3	121	10.0	0.5			
Osmondthorpe	 1	10.6	4	42.4				
Outsiders	 ***		43					
Totals	 2,787	24.5	1,795	15.8	1.5			

Considered as occurring in the Municipal Wards, the foregoing Deaths are classed as follows:—

Wards.	Deaths.	Death	Wards.	Death	Death
Eastern Div	ision.	Rate.	Western Divi	sion.	Rate.
Central	82	16.4	Mill Hill	27	15.0
North	129	12.0	West	118	20.3
North-East	158	18.7	North-West	114	13.4
East	153	21.1	Brunswick	81	14:3
South	73	20.7	New Wortley	90	19.8
East Hunslet	156	17:3	Armley	136	13.7
West Hunslet	107	13.8	Bramley	78	13.8
Holbeck	131	16.0	Headingley	110	9.7

In both these tables deaths occurring in public institutions have been referred to the districts to which the patients belonged. The births in workhouses are included in those or the districts in which these institutions are situated. There were twenty-five deaths at Manston Hospital during this quarter.

The following Births and Deaths were recorded in the several Sub-Registration Districts of the City of Leeds during the thirteen weeks ended 31st March, 1906. The figures in italics after the Births and Deaths give the proportion per annum per 1,000 of the estimated population.

Districts.	Births.	Birth Rate.	Deaths.	Death Rate. All causes. 7 Zymotics				
oi (North	 480	32.7	290	19.8	2.6			
West South-East	 445	21.0	405	19.1	2.0			
South-East	 334	38.4	229	26.3	2.9			
Hunslet	 564	29.8	340	18.0	3.0			
Holbeck	 259	29.5	146	16.7	13			
Wortley	 392	24.3	265	16.4	2.3			
Kirkstall	 286	24.0	148	12.4	1.9			
Bramley	 109	23.4	45	9.7	0.4			
Chapeltown	 241	23.3	134	12.0	1.3			
Osmondthorpe	 2	22.7	2	22.7				
Outsiders	 		50					
Totals	 3,112	26.9	2,054	17.8	2.1			

Considered as occurring in the Municipal Wards, the foregoing Deaths are classed as follows:—

Wards.	Dea	ths.	Death	Wards.		Deaths.	Death
Eastern Division.			Rate.	Western Div		Rate.	
Central		90	18.0	Mill Hill		36	20.7
North	І	56	13.8	West		137	23.3
North-East	I	72	21.7	North-West		144	16.8
East	1	90	25.5	Brunswick		94	16.4
South		90	25.9	New Wortley		91	19.8
East Hunslet	I	77	18.8	Armley		163	15.7
West Hunslet	1	39	17:3	Bramley		56	9.6
Holbeck	1	15	14.7	Headingley		154	12.5

In both these tables deaths occurring in public institutions have been referred to the districts to which the patients belonged. The births in workhouses are included in those of the districts in which these institutions are situated. There were twenty-seven deaths at Manston Hospital during this quarter.

TABLE F (1).

Shewing Births, Deaths, from all and certain causes, Home Patients of the Dispensary, admissions to the Fever Hospitals, and some of the Meteorological conditions and the Death-rates from certain causes in Leeds: with the Birth and Death-rates from all causes in the 76 large English towns for each of the thirteen weeks ended 31st December, 1904.

	1	1	OCT	THE RESERVE AND ADDRESS OF THE PERSON NAMED IN		I		VEMI		ember		**	MADEL			
1 2 17 5									1		_	ECE	-	1	OR BS.	
1904.		Oct. 8th.	Oct. 15th.	Oct. 22nd.	Oct. 29th.	Nov. 5th.	Nov. 12th.	Nov. 19th.	Nov. 26th.	Dec. 3rd.	Dec. roth.	Dec. 17th.	Dec. 24th.	Dec. 31st.	TOTALS OR AVERAGES.	YEAR.
Total Births Total Deaths	1 2	241 121	266 136	250 125	224 140	237 128	225 150	219 137	214	244 147	218 129	251 113	216 171	216 201	3,021 1,828	12,561 8,096
Under 1 year	3 4 5 6 7	43 8 3 40 27	48 6 7 52 23	36 9 4 45 31	41 16 7 40 36	4 ² 9 5 4 ² 30	40 12 7 62 29	32 11 6 48 40	34 6 4 54 32	40 10 5 53 39	29 14 6 42 38	21 9 3 38 42	35 8 4 68 56	36 11 7 80 67	477 129 68 664 490	2,207 735 546 2,614 1,994
Deaths: Small-pox Measles Scarlet Fever *Diphtheria Whooping-cough { Typhus Fever Typhoid Fever Other or doubtful Diarrhoea or Dysent.	_	 I I	 1 2 14	3 1 1	 2 2 2 1 	1 2 1 1	1 1 2 2 2	 1 1 3	 I 	 I I 2 	 2 2 3 	 I 2 I 	I 2 1 2 2	 1 2 	2 2 7 15 15 1 15 	2 344 59 50 208 3 47
All seven	17	10	17	6	10	6	8	6	3	4	8	5	6	3	92	1,164
Croup	21 22	7 9 17 4 10 23	117 117 15 10 5 11 28	35 1 9 10 3 9	33 7 14 3 8 16	35 8 10 3 10 25	36 1 16 11 5 12 30	38 1 14 6 9 15 27	22 12 16 6 13 26	36 1 17 8 10 16	38 5 15 3 9 23	 20 5 7 6 16 24	 42 1 14 19 8 16 28	1 64 7 15 20 4 18 20	5 423 13 146 163 69 163 301	10 1,442 32 626 677 245 630 1,185
Dispensary: visits pd.	27	231	270	241	357	310	362	293	332	294	297	287	314	234	3,822	15,639
Cases admitted to our own hospitals	28	28	27	22	47	22	38	44	28	34	22	22	20	28	382	1,263
Attached Ther. F Dry bulb	30 31 32 33 34 35 36 37 38 39 40	54.77 52.46 48.69 76.15 57.00 44.71 12.29 0.14 NW sw 2	53.92 52.39 49.69 82.77 56.14 35.86 20.28 0.06 E.SE 2 	57.39' 57.31 55.08 86.08 61.14 48.29 12.85 0.16 w se 2	57:46 49:46 46:85 82:38 56:14 41:71 14:43 0:08 SE NW 2	55.08 51.77 48.77 80.23 54.71 46.57 8.14 0.03 SW NE 2	54.15 50.23 47.77 83.15 54.43 42.00 12.43 0.91 w sw 3 26.1	50.46 45.92 44.00 86.08 50.57 36.71 13.86 w 2 	41 '92 36 '23 34 '85 87 '00 40 '00 32 '86 7 '14 0 '61 NW 2 	46.00 44.08 42.46 87.38 46.86 37.86 9.00 0.01 sw.w.2 	46.85 41.15 39.77 84.77 46.00 36.86 9.14 0.58 sw 2	46 15 44 69 42 69 85 23 46 86 38 14 8 72 0 74 w sw 2	41°54 33°15 31°92 84°23 36°00 30°29 5 71 sw 2 25°0	43'31'41'69'40'00'85'92'44'43'35'00'9'43'0'12'W NW 3'''25'0	46°20 44°04 83°95 50°02 38°99 11°03 3°44 2	29'79 53'33 51'88 47'72 75'10 56'21 43'07 13'14 20'54 2
Death-rate (Leeds) Death-rate (76 towns) Birth-rate (76 towns)	43 44	14.0 12.1	15.8	14.5 15.9 28.5	29.6 12.0	14'9 30'0	15.7 28.2	16.6	17°3	17.0 21.0	15'0	13.1 19.5 28.8	19.8 19.4 25.7	23°3 22°4 26°0	16·3 17·5 28·1	18.0 17.3 29.1
D.R. lung dis. (Leeds) D.R. 7 Zymotics		0.8 1.5	2.0	4.1 0.2	3.8	4°1 0°7	4.5 0.0	4.4 0.2	0.3 0.9	4.5 0.2	4'4 0'9	0.6 5.3	4'9 0'7	7.4	3.8	3.5

The Dispensary returns are furnished me by the kindness of the resident staff, and have regard to a week ended in each case a day earlier than that given in the heading.

The meteorological data are compiled from returns sent us by Mr. Crowther. They are uncorrected readings, made at 10 a.m. and 4 p.m. The humidity each week is the average of the humidities calculated on each of the thirteen observations of the wet and dry bulbs.

^{*} Includes membranous croup. Line 19 includes non-spasmodic croup not returned as membranous. † Line 20 includes line 21.

TABLE F (2).

Shewing Births, Deaths, from all and certain causes, Home Patients of the Dispensary, admissions to the Fever Hospitals, and some of the Meteorological conditions and the Death-rates from certain causes in Leeds: with the Birth and Death-rates from all causes in the 76 large English towns for each of the thirteen weeks ended 1st April, 1905.

	1		JANI	UARY	7.	F	EBRU	JARY	7.		М	ARC	H.		
1905.		Jan. 7th.	Jan. 14th.	Jan. 21st.	Jan. 28th.	Feb. 4th.	Feb. 11th.	Feb. 18th.	Feb. 25th.	Mar. 4th.	Mar. 11th.	Mar. 18th.	Mar. 25th.	Apr. 1st.	TOTALS OR AVERAGES.
Total Births Total Deaths	1 2	243 181	278 179	258 140	278 141	25I 122	261 143	237 139	222 116	245 129	23I 142	260 141	243 133	248 113	3,255 1,819
Under I year I to 2 years 2 to 5 years 5 to 60 years 60 yrs. and upwards	3 4 5 6 7	33 9 10 66 63	31 8 3 69 68	33 6 2 52 47	27 4 9 55 46	25 8 3 43 43	25 9 7 51 51	33 7 4 50 45	20 3 11 49 33	24 11 3 64 27	37 8 6 50 41	41 11 6 48 35	37 8 7 44 37	31 4 8 37 33	397 96 79 678 569
Deaths: Small-pox Measles Scarlet Fever *Diphtheria Whooping-cough Typhus Fever Typhoid Fever Other or doubtful Diarrhœa or Dysent.	8 9 10 11 12 13 14 15 16	 2 1 2	1 2 1 I	 I 1	3 1 I	 I 	 I I 4 I 2	 1 2 1 1 3	 4 1	1 1 4 2 I	 1 3 3	3 	3 3 	 1 3 1	3 8 14 20 21 16
All seven	17	6	5	3	8	3	9	9	7	9	8	7	7	7	88
Cholera (English) Croup Dis. of Resp. System Influenza† Phthisis Dis. of Circul. System Violent Deaths Inquest cases Deaths in Pub. Inst.	18 19 20 21 22 28 24 25 26	62 9 13 14 12 29 25	45 9 16 15 11 19 32	 1 31 1 19 10 6 8 37	33 4 9 12 4 13 23	25 12 10 4 14 32	28 5 14 8 10 21 30	34 4 14 6 3 6 23	 17 2 18 9 7 15 26	26 3 10 14 9 18 27	27 1 15 12 3 16 31	29 13 7 5 15 23	27 1 17 8 3 15 23	26 3 8 10 2 8 16	1 410 42 178 135 79 197 348
Dispensary: visits pd.	27	305	359	386	334	316	277	226	249	291	267	255	263	240	3,768
Cases admitted to our own hospitals	28	42	55	33	33	63	36	42	33	29	33	36	39	32	506
Wet bulb	30 31 32 33 34 35 36 37 38 39 40 41 42 43	47 '31' 45 '77' 43 '85' 62' 47 '86' 39 '71' 8 '15' 0 '02' w 2 27 '8 20 '7	46.62 41.92 40.23 86.92 45.14 36.57 0.35 w 3 31.8 20.4 17.5	40'00 33'15 32'23 88'31 36'71 28'29 8'42 0'10 SE E 2 29'5 16'0 18'1	44.08 41.77 40.54 90.31 45.43 35.86 9.57 0.05 sw 2 31.8 16.1 18.4	47 '38 45 '38 42 '77 80 '85 48 '57 39 '00 9 '57 0 '36 W 3 28 '7 13 '9 18 '0	48.08 43.38 40.77 80.77 48.71 38.29 10.42 0.02 w.sw 2 29.8 16.3 17.4	48'31'47'08'44'31'80'77'50'00'39'71'10'29'0'35'w'2'	45'00 39'77 36'62 75'46 42'71 34'29 8'42 0'24 NE NW 2 25'4 13'3 15'7	45.00 42.23 39.77 81.62 45.57 35.29 10.28 0.50 w sw 3 28.0 14.7 17.1	47.69 44.85 41.92 79.31 49.71 38.14 11.57 0.58 sw 3 26.4 16.2 17.2	48.69 48.15 45.00 78.54 52.43 41.29 11.14 0.65 sw 3 29.7 16.1 15.8	50°38 48°92 46°08 81°00 54°71 39°71 15°00 0°18 SE E 2 27°8 15°2 15°9	49.75 48.92 45.92 80.08 52.29 41.86	46'77 43'92 41'51 82'29 47'68 37'54
D.R.lung dis. (Leeds) D.R. 7 Zymotics	45 46	7°1 0°7	0.4 2.1	3.2	3.8	2.0	3.5	3.0	0.8	3.0	0.8 3.1	3.3	3.1	3.0	3.6

The Dispensary returns are furnished me by the kindness of the resident staff, and have regard to a week ended in each case a day earlier than that given in the heading.

The meteorological data are compiled from returns sent us by Mr. Crowther. They are uncorrected readings, made at to a.m. and 4 p.m. The humidity each week is the average of the humidities calculated on each of the thirteen observations of the wet and dry bulbs.

^{*} Includes membranous croup. Line 19 includes non-spasmodic croup not returned as membranous. † Line 20 includes line 21.

TABLE F (3).

Shewing Births, Deaths, from all and certain causes, Home Patients of the Dispensary, admissions to the Fever Hospitals, and some of the Meteorological conditions and the Death-rates from certain causes in Leeds: with the Birth and Death-rates from all causes in the 76 large English towns for each of the thirteen weeks ended 1st July, 1905.

			AP	RIL.			N	MAY.		1		1			
1905.		April 8th.	April 15th.	April 22nd	April 29th.	May 6th.	May 13th.	May 20th.	May 27th.	June 3rd.	June 10th.	June 17th.	June 24th.	July 1st.	TOTALS OR AVERAGES,
Total Births Total Deaths	1 2	229 142	247 135	260 147	233 136	264 153	272 150	266 147	244 143	228 122	253 110	211 105	273 111	234 119	3,214 1,720
Under 1 year	3 4 5 6 7	27 13 7 62 33	32 7 11 48 37	33 11 5 62 36	33 7 5 48 43	34 11 5 60 43	33 8 5 44 60	39 4 6 55 43	19 7 12 62 43	28 5 5 49 35	24 6 6 48 26	17 4 4 48 32	31 6 45 28	23 11 10 43 32	373 95 87 674 491
Deaths: Small-pox Measles Scarlet Fever *Diphtheria Whooping-cough Typhus Fever Typhoid Fever Other or doubtful Diarrhœa or Dysent.	8 9 10 11 12 13 14 15 16	 I I	1 4 I	I I I I 	 	I 2 I	 I 2 I 	 I 1 	 2 3 1	 I 6 I 	2 I 1 2	 I I 2	 3 2 3	I I I 2	8 5 7 27 1 10 1 8
All seven	17	2	7	4	2	5	5	3	6	8	6	4	8	7	67
Cholera (English) Croup Dis. of Kesp. System Influenza† Phthisis Dis. of Circul. System Violent Deaths Inquest cases Deaths in Pub. Inst.	21 22	38 2 13 7 2 6 22	36 I 10 I4 10 I9 26	35 3 14 12 12 17 16	25 I II 9 3 I2 24	30 11 8 6 21 28	33 11 16 11 22 26	 30 1 6 15 5 14 27	 27 1 13 18 7 13 35	 21 1 11 10 5 10 22	 18 1 12 8 5 12 27	 14 8 9 3 7 26	 21 8 7 7 7 12 22	 25 9 13 5 7 21	353 11 137 146 81 172 322
Dispensary: visits pd.	27	302	238	298	197	270	200	252	202	278	223	197	238	201	3,176
Cases admitted to our own hospitals	28	23	35	20	18	26	15	16	21	24	14	9	25	18	264
Mn. of highest reading ,, lowest ,, daily range Total rainfall (inches) Wind {Direction Force 0-6 Amount of Cloud Birth-rate (Leeds)	30 31 32 33 34 35 36 37 38 39 40	48.69 46.15 43.23 79.69 51.00 37.29 13.71 0.55 N W 3 26.2	50.08 50.54 47.77 81.92 54.29 42.14 12.15 0.31 8E.SW 3 	47.67 44.58 42.83 87.08 47.71 38.29 9.42 0.28 NE N 3 	49°54 51°69 48°85 81°23 54°43 42°00 12°43 0°66 w 2 	50.92 50.15 46.23 74.46 53.86 41.00 12.86 1.03 SW NE 3	53.85 56.92 50.46 63.85 61.29 44.71 16.58 w nw 3	56·15 56·77 51·31 69·69 03·29 46·71 16·58 0·07 NE 2 	53.69 56.15 50.62 68.54 66.14 44.71 21.43 sw nw 2	60.62 64.15 56.77 62.00 68.71 52.86 15.85 0.35 sw 2	58.00 58.46 51.85 63.85 62.71 46.86 15.85 0.15 NE 3	61'31'64'85'57'46'62'92'70'86'49'29'21'57'0'86'E NE 2	61.85 67.08 60.46 66.31 73.57 53.29 20.28 1.23 SW.E.3 	65°38 67°54 60°69 67°23 74°29 53°86 20°43 0°35 NE 2	55°26 56°61 51°48 71°35 61°70 45°62 16.08 5°84 3
Death-rate (Leeds) Death-rate (76 towns) Birth-rate (76 towns)	42 43 44	16.2 15.7 27.7	15:4 16:6 28:6	16.8 15.6 26.5	15.2 16.2 28.3	17.5 15.4 31.9	30.2 12.0	16·8 14·4 29·5	16.3 14.6 28.9	13.0 14.7 27.2	13.5 28.5	13.4 24.9	30.3 13.0	13.6 12.7 28.4	15°1 14°7 28°5
D.R. lung dis. (Leeds) D.R. 7 Zymotics		4'3 0'2	0.8 1.1	0.2	0.5	3.4	3.8	3'4 0'3	3.1 0.2	0.9	0.7	0.2	2.4 0.0	0.8 0.8	3.1 0.9

The Dispensary returns are furnished me by the kindness of the resident staff, and have regard to a week ended in each case a day earlier than that given in the heading.

The meteorological data are compiled from returns sent us by Mr. Crowther. They are uncorrected readings, made at to a.m. and 4 p.m. The humidity each week is the average of the humidities calculated on each of the thirteen observations of the wet and dry bulbs.

^{*} Includes membranous croup. Line 19 includes non-spasmodic croup not returned as membranous. † Line 20 includes line 21.

TABLE F (4).

Shewing Births, Deaths, from all and certain causes, Home Patients of the Dispensary, admissions to the Fever Hospitals, and some of the Méteorological conditions and the Death-rates from certain causes in Leeds: with the Birth and Death-rates from all causes in the 76 large English towns for each of the thirteen weeks ended 30th September, 1905.

			IU	ILY.			A	UGUS	ST.						
1905.		July 8th.	July 15th.	July 22nd	July 29th.	Aug. 5th.	Aug. 12th.	Aug. 19th.	Aug. 26th.	Sept. 2nd.	Sept. 9th.	Sept. 16th.	Sept. 23rd.	S pt. 3oth.	TOTALS OR AVERAGES.
Total Births Total Deaths	1 2	231 81	248 122	242 158	245 173	282 152	191 165	262 145	250 156	240 134	236 134	222 128	216 131	216 111	3,081 1,790
Under 1 year	3 4 5 6 7	20 5 5 33 18	32 9 7 43 31	70 4 6 44 34	71 16 6 45 35	77 7 8 33 27	77 14 5 40 29	58 11 4 46 26	55 15 8 44 34	48 -9 -8 45 -24	51 5 6 36 36	42 10 4 40 32	37 7 10 45 32	30 6 7 38 30	668 118 84 532 388
Typhus Fever Typhoid Fever Other or doubtful	8 9 10 11 12 18 14 15	1 2	2 2 6 I	3	 I I 5	1	 2 2 	5	 4 4 1	 2 3 2	 I 2 2 2	 I 4 I	3 3 4 	 1 2 3 	 19 13 8 35 10
Diarrhœa or Dysent. All seven	16	5	16	37 41	49	50	55	37 42	41	32	25	21	18	12	325 410
Cholera (English) Croup Dis. of Kesp. System Influenza† Phthisis Dis. of Circul. System Violent Deaths Inquest cases Deaths in Pub. Inst.	18 19 20 21 22 23 24 25 26	13 7 7 7 2 6 16	10 12 9 4 11	13 12 14 6 9	23 10 10 2 7 25	 17 6 10 1 10 14	23 12 7 6 10 23	 15 7 12 5 9 28	 15 9 12 6 14 20	 17 11 6 3 7 22	 19 10 8 3 6 19	 12 8 10 4 10 18	 1 13 2 9 10 4 6 34	 15 4 9 3 10 14	 2 - 205 2 117 124 49 115 271
Dispensary: visits pd.		241	252	294	280	258	283	317	290	272	238	294	249	299	3,567
Wind Direction Force 0-6	29 30 31 32 33 34 35 36 37 38 39 40 41 42 43	63.54 67.08 57.85 56.00 73.00 52.14 20.86 0.10 NW W 2 26.4 9.3 11.6	68·77 71·08 64·92 69·46 78·43 59·57 18·86 0·13 NW I 28·3 13·9 12·9	66 23 68 85 60 92 61 62 73 43 57 00 16 43 0 34 NW 2 27 6 18 0 13 4	65.31 65.92 59.85 68.62 70.14 56.43 13.71 0.61 8.8 W.W. 2 28.0 19.8 15.0	61.69 62.69 56.85 69.62 66.86 54.00 12.86 0.91 sw 2 32.2 17.4 17.0	60.46 61.77 55.92 68.38 66.86 51.57 15.29 0.30 sw w 2 21.8 18.8	61.69 63.69 57.38 67.54 69.29 53.71 15.58 0.92 sw 2	60.46 61.46 56.08 71.15 67.29 53.00	60.77 60.38 56.08 75.00 65.43 53.43 12.00 0.53 NE 2 27.4 15.3 17.3	60.31 61.15 57.46 78.46 64.86 54.14 10.72	56.15 57.46 52.38 70.54 62.43 47.43 15.00	56.23 57.92 54.00 76.92 62.00 48.86 13.14	55.62 54.54 52.92 89.15 58.57 48.57	62.62 57.12 70.06 67.58 53.07 14.51
D. R. lung dis. (Leeds)	_	1.2	0.6	1.2	2.6	1.9	2.6 5.8	1.7 4.5	1.7	1.9	5.1 5.2	1 '4 1 '7	1.2	1.7	1.8

The Dispensary returns are furnished me by the kindness of the resident staff, and have regard to a week ended in each case a day earlier than that given in the heading.

The meteorological data are compiled from returns sent us by Mr. Crowther. They are uncorrected readings, made at 10 a.m. and 4 p.m. The humidity each week is the average of the humidities calculated on each of the thirteen observations of the wet and dry bulbs.

^{*} Includes membranous croup. Line 19 includes non-spasmodic croup not returned as membranous. † Line 20 includes line 21.

TABLE F (5).

Shewing Births, Deaths, from all and certain causes, Home Patients of the Dispensary, admissions to the Fever Hospitals, and some of the Meteorological conditions and the Death-rates from certain causes in Leeds: with the Birth and Death-rates from all causes in the 76 large English towns for each of the thirteen weeks ended 30th December, 1905.

1	17/1		OCTO	BER			NOV	VEMI	BER.		D	ECE	MBEI	R.	e	
1905.		Oct. 7th.	Oct. 14th.	Oct. 21st.	Oct. 28th.	Nov. 4th.	Nov. 11th.	Nov. 18tn.	Nov. 25th.	Dec. 2nd.	Dec. 9th.	Dec. 16th.	Dec. 23rd.	Dec. 30th.	TOTALS OR AVERAGES.	YEAR.
Total Births Total Deaths	1 2	218	214 105	220 99	244 123	203 142	223 142	192 146	214 157	216 139	227 170	216 155	234 150	166 157	2,787 1,795	12,337 7,124
Under I year	3 4 5 6 7	26 11 6 35 32	26 6 3 47 23	25 5 41 27	31 9 7 46 30	37 14 5 56 30	30 7 8 53 44	38 5 12 49 42	43 10 13 53 38	30 9 11 44 45	49 11 16 36 58	34 20 15 41 45	15 50 41	34 13 18 54 38	437 130 130 605 493	1,875 439 380 2,489 1,941
Deaths: Small-pox Measles Scarlet Fever *Diphtheria Whooping-cough Typhus Fever Typhoid Fever Other or doubtful Diarrhoea or Dysent.	11 12 13 14 15	3 3 1 	3 1	 1 2 3	 2 2 1	 2 5 1 3	7 1 2 2 1	 4 1 2 1	 4 2 1 6 3	7 4 1	3 3 1	16 1 1 2 1 	 8 5 2 	15 4 1	74 7 9 38 16 1 26	3 109 39 44 121 1 52 2 365
All seven	17	12	10	6	5	11	14	9	16	13	13	24	16	22	171	736
Cholera (English) Croup Dis. of Resp. System Influenza† Phthisis Dis. of Circul. System Violent Deaths Inquest cases Deaths in Pub. Inst.	21 22	14 1 6 13 7 14 22	1 24 1 9 11 5 14 18	 20 1 10 6 4 24	22 1 8 10 8 11 23	34 1 13 11 1 9	35 2 7 11 5 17 22	39 1 9 16 6 18 24	 41 14 17 9 20 20	32 1 12 11 2 8 20	56 1 11 13 5 12 31	10 10 7 20 22	34 1 8 14 3 7 23	37 9 15 6 16	1 428 11 126 158 64 170 284	 6 1,396 66 558 563 273 654 1,225
Dispensary: visits pd.		269	342	375	410	343	371	486	431	510	416	527	501	292	5,273	15,784
Cases admitted to our own hospitals		32	20	20	28	29	21	32	34	25	21	21	25	26	335	1,391
Attached Ther. °F Dry bulb Wet bulb Humidity Mn. of highest reading , lowest ,, daily range Total rainfall (inches) Wind {Direction Force 0-6 Amount of Cloud	30 31 32 33 34 35 36 37 38 39 40	53.08 51.62 50.23 90.38 57.57 43.14 14.43 0.38 NW 3	53°38 52°08 50°69 90°46 58°29 45°57 12°72 0°57 NW 2	44'00 42'62 89'54 52'43 36'86 15'57 0'34 NW 2	46.62 46.38 44.77 88.23 50.57 38.20 12.28 0.05 NW I	49°08 47°00 45°77 91°23 53°86 42°29 11°57 1°28 sw sb 1	48.00 46.08 44.46 88.38 50.29 40.57 9.72 6.78 8W NW I	45'31'41'31'39'46'85'31'46'00'37'71'8'29'0'34	44 '00 42 '77 40 '69 84 '08 45 '43 35 '86 9 '57 0 '33 Nw sw 2	46.69 44.54 42.77 86.85 48.29 39.57 8.72 0.81 sw se 2	48.62 46.69 44.15 82.15 49.86 42.00 7.86 0.12 sw 2	47.15 44.62 42.77 86.23 47.14 40.71 6.43 NW 2	47.15 44.92 43.15 86.85 47.86 39.86 8.00 se.sw 2	42 '92' 39 '08' 37 '46' 86 '54' 42 '57' 34 '00' 8 '57' 0 '14' NE SE 2	45'47 43'77 87'40 50'01 39'73 10'28 5'14 2	29.85 52.69 52.16 48.48 78.01 56.74 43.99 12.75 20.75
Birth-rate (Leeds) Death-rate (Leeds) Death-rate (76 towns) Birth-rate (76 towns)	42 43	13.2	13.7	11'3	14.0 14.8	16 2	16.5	16.7		15.9	16.1 16.1		26.7 17.1 17.8 27.8	19'0 17'9 17'4 21'4	24.5 15.8 15.8 26.6	27°1 15°6 15°7 28°2
O.R.lung dis. (Leeds) O.R. 7 Zymotics		1.6	2.2 1.1	2.3	2.2	3.9	4.0	4.2	4.7 1.8	3.7	6.4	4.6	3.9	4'3 2'5	3.8	3.1

The Dispensary returns are furnished me by the kindness of the resident staff, and have regard to a week ended in each case a day earlier than that given in the heading.

The meteorological data are compiled from returns sent us by Mr. Crowther. They are uncorrected readings, made at 10 a.m. and 4 p.m. The humidity each week is the average of the humidities calculated on each of the thirteen observations of the wet and dry bulbs.

^{*} Includes membranous croup. Line 19 includes non-spasmodic croup not returned as membranous. † Line 20 includes line 21.

TABLE F. (6).

Shewing Births, Deaths, from all and certain causes, Home Patients of the Dispensary, admissions to the Fever Hospitals, and some of the Meteorological conditions and the Death-rates from certain causes in Leeds; with the Birth and Death-rates from all causes in the 76 large English towns for each of the thirteen weeks ended 31st March, 1906.

Total Births	EDINGUISM MAICH, 1900.
Total Births 1 233 231 210 257 244 256 227 244 221 281 220 248 240 3,1 Total Deaths 2 194 172 154 156 142 154 180 155 157 165 14c 145 140 2,0 Under I year 3 40 32 33 44 33 37 31 36 22 32 30 44 33 4 1 to 2 years 4 23 21 17 17 17 11 19 12 19 13 10 10 14 2 2 to 5 years 5 19 12 9 9 8 10 14 10 14 15 5 6 10 1 5 to 60 years 6 62 66 59 53 46 52 64 53 62 52 60 51 47 7 60 yrs, and upwards 7 50 41 36 33 38 44 52 44 40 53 35 34 36 5	FEBRUARY. MARCH.
Total Deaths 2 194 172 154 156 142 154 180 155 157 165 146 145 140 2,0 Under I year 3 40 32 33 44 33 37 31 36 22 32 30 44 33 4 1 to 2 years 423 21 17 17 11 19 12 19 13 10 10 14 2 2 to 5 years 5 19 12 9 9 8 10 14 10 14 15 5 6 10 1 5 10 1 5 6 60 66 59 53 46 52 64 53 62 52 60 51 47 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 </td <td></td>	
I to 2 years	
Measles 9 15 16 9 15 8 11 14 7 12 14 7 11 7 1 Scarlet Fever 10 1 1 1 <	17 11 19 12 19 13 10 10 14 203 8 10 14 10 14 15 5 6 10 141 8 46 52 64 53 62 52 60 51 47 727
Diarrhoea or Dysent. 16 I I I I 2 2 I I	8 II I4 7 I2 I4 7 II 7 I46 I I 4 6 2 3 I 2 4 3 3 3 34 3 2 3 2 4 5 2 3 2 2 42 2 I I I I 2 I I 12
	1 16 19 20 15 21 20 13 17 14 248
Influenza	I I 5 28 19 33 36 24 22 29 29 32 412 3 2 1 2 12 2 7 10 18 10 19 9 17 11 16 169 0 10 15 19 15 13 22 23 6 6 187 4 4 4 11 4 7 4 5 4 1 63 0 12 18 19 9 12 12 15 13 9 168
Dispensary: visits pd. 27 363 365 359 352 332 260 264 258 278 252 295 288 240 3,9	2 332 260 264 258 278 252 295 288 240 3,906
Cases admitted to our own hospitals 28 20 30 27 18 27 26 22 31 34 24 24 26 36 3	3 27 26 22 31 34 24 24 26 36 345
Wind {Direction 38 SE SW SW NW SW NW SW NW SW NW SW NW SW SE NE W SW SW SW NW SW NW SW NW SW NW SW NW SW NW SW SE NE NW NE	9 55 23 51 69 50 60 52 23 52 08 57 08 54 85 57 77 57 69 52 35 24 57 38 39 31 38 23 38 08 40 50 47 77 44 31 42 15 43 69 42 14 043 31 37 23 36 23 36 15 38 25 44 54 41 69 38 08 38 92 39 62 84 62 52 62 82 92 84 31 82 58 78 31 81 69 71 08 68 62 81 32 64 9 00 43 29 45 00 44 71 44 43 52 71 47 29 46 71 47 00 46 36 40 71 33 00 33 43 32 71 34 71 42 14 33 86 35 14 35 57 35 92 8 29 10 29 11 57 12 00 9 72 10 57 13 43 11 57 11 43 10 44 3 0 08 0 49 0 18 0 46 0 75 0 42 0 31 0 36 0 19 6 06 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
Birth-rate (76 towns) 44 29 9 30 3 26 7 29 3 28 2 28 4 29 4 28 4 28 1 29 2 28 0 28 1 29 5 2	28.5 58.4 56.4 58.4 58.1 56.5 58.0 58.1 56.2 58.2

The Dispensary returns are furnished me by the kindness of the resident staff, and have regard to a week ended in each case a day earlier than that given in the heading.

The meteorological data are compiled from returns sent us by Mr. Crowther. They are uncorrected readings, made at 10 a.m. and 4 p.m. The humidity each week is the average of the humidities calculated on each of the thirteen observations of the wet and dry bulbs. On January 17th, 1906, the barometer was removed into an inner room in which there was a fire.

^{*} Includes membranous croup. Line 19 includes non-spasmodic croup not returned as membranous the Line 20 includes line 21.



