[Report 1936] / Medical Officer of Health, Newcastle-upon-Tyne City & County.

Contributors

Newcastle upon Tyne (England). City & County Council.

Publication/Creation

1936

Persistent URL

https://wellcomecollection.org/works/fs3wbrpj

License and attribution

You have permission to make copies of this work under a Creative Commons, Attribution license.

This licence permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See the Legal Code for further information.

Image source should be attributed as specified in the full catalogue record. If no source is given the image should be attributed to Wellcome Collection.





CITY AND COUNTY OF NEWCASTLE UPON TYNE

ANNUAL REPORT

OF THE

MEDICAL OFFICER OF HEALTH

ON THE

Sanitary Condition of the City

DURING THE YEAR

1936.

PRINTED BY
CHRISTIE, MALCOLM, LTD.,
CHARLOTTE SQUARE, NEWCASTLE UPON TYNE.

Digitized by the Internet Archive in 2017 with funding from Wellcome Library

CONTENTS.

NTRODUCTION AND SUMMARY—	
	PAGE.
HEALTH COMMITTEE	8
MATERNITY AND CHILD WELFARE COMMITTEE	9
Organisation of Health Department	9A
Staff	10-12
LETTER TO CHAIRMAN OF HEALTH COMMITTEE—	
Vital Statistics	13-21
Infectious Diseases	21-22
Nutrition	22-23
General Hospital Services.	24
Reorganisation of the Municipal and Voluntary Hospitals in	~.
the City	24-25
Whitton Tower Convalescent Home for Children	25
Slum Clearance	25-26
Conclusion Summary of Statistics (1936)	27 28
Summary of Statistics (1990)	20
REPORT—	
I.—GENERAL STATISTICS—	
(Including Population and Vital Statistics, Mortality Tables as prescribed by Ministry of Health, Social Conditions, Geology, Climatology, Water Supply, Disposal of Refuse, and Adoptive and Local Acts in Force)	31-40
Hospitals in the City.	41
II.—THE CHILD—	
Births and Deaths.	45-46A
Report of the Maternity and Child Welfare Medical Officer—	10-101
Diphtheria Immunisation	47-48
Dental, Aural, and Nasal Treatment	48
Whalton Rest Home	48
Orthopædia	48-49
Home Helps	49
Maternity Hospital	49 50-51
Midwives' Fees	51-52
Maternal Mortality	52
Puerperal Septicæmia and Puerperal Pyrexia	52-53
Consultants' Services	53
Midwives Acts	53-54
Ultra-Violet Ray Therapy	54 54
Nursery Schools	54
Dietho	51 55

CONTENTS—Continued.

TI	TH	E C	HIL	D (C	onting	red)_
-1.1		100	7 A A A A A A		C/ / 2 4 4 1 1 4 4 1	00.00

Report of the Maternity and Child Welfare Medical Officer (Continued)—

(Community)	
Death	PAGE 55
Deaths	55-56
Toddlers	56
Children Acts, 1908–1933.	56-57
Municipal Training Course for Health Visitors	57
Welfare Centres—Attendances, etc.	57-60
Dried Milk	60-61
Notification of Live and Still Births	61
Work of the Health Visitors—Details, etc	62-63
Voluntary Workers	63
Maternity and Nursing Homes (Report of the Board	
of Inspection)	64-68
III.—INFECTIOUS DISEASE—	
Deaths—Ward Distribution	71
Ages and Deaths of Notified Cases	72
Cases—Ward Distribution	73
Households Affected	74
Schools	75
Public Institutions	75
Scarlet Fever	76
Diphtheria	76
Measles and Rubella	76-78
Whooping Cough	78
Food Poisoning	78-79
Enteric Group of Fevers	79-80
Diarrhœa	80
Smallpox	81
Vaccination	81
Chickenpox	81
Erysipelas	81
Puerperal Septicæmia and Puerperal Pyrexia	81
Influenza and Pneumonia	81-83
Encephalitis Lethargica	83
Acute Poliomyelitis	83
Cerebro-Spinal Meningitis	83-84
Bacillary Dysentery	85-86
Venereal Diseases	86
CITY HOSPITALS FOR INFECTIOUS DISEASES—	
Report of the Deputy Medical Superintendent—	
Accommodation, Admissions, Average Stay, Per-	
centages, and present Death Rates compared with	
previous years, etc	87-91
Diphtheria	91-94
Scarlet Fever	94-96
Otorrhœa and Rhinorrhœa	96-97

CONTENTS—Continued.

CITY HOSPITAL FOR INFECTIOUS DISEASES (cont	inued)—
Report of the Deputy Medical Superintendent (:ontinued)-	PAGE.
" Return " Cases	97-98
Erysipelas	
Mixed Infections	
Cross Infections	
Staff Sickness	
Smallpox and Isolation Hospitals	. 101
Disinfection—Disinfectants Distributed, etc	
Bacteriological Investigations	.102-110
IV.—TUBERCULOSIS—	
Report of the Tuberculosis Medical Officer-	
Tuberculosis Dispensary Clinics	. 113
Attendances at the Dispensary	
Contacts	
Domiciliary Visits	. 117
Relations with other Departments and Hospitals	. 117
Voluntary Tuberculosis Care Council	. 117
Nurses' Visits to Patients	. 118
X-Ray Examinations.	. 118
Sputum Examinations	. 118
Work of the Sanitary Inspector	
Notifications and Deaths	.118-121
Duration of Illness	
Family History	
House Accommodation	
Tuberculosis and Pregnancy	. 123
Institutional Treatment—Barrasford, Stannington,	
City Hospital and Newcastle General Hospital	
BARRASFORD SANATORIUM—Report of the Medica	1
Superintendent—	
General	
X-Ray Plant	
Dental Clinic	
Occupational Therapy	
Admissions	
Discharges	
Status and Occupation	
Diagnosis	.134-133
Treatment—Artificial Pneumothorax, Gold Salts,	195 197
Ultra-Violet Ray Radiation	
	107-109
V.—GENERAL DISEASES—HOME AND HOSPITAL—	
Domiciliary Medical Services	143
NEWCASTLE GENERAL HOSPITAL—Report of the	
Medical Superintendent—	111111111111111111111111111111111111111
Lectures	
Tuberculosis	144

CONTENTS—Continued.

V.—GENERAL DISEASES (continued)—	PAGE
Maternity Ward	145
Mental Cases	145
Infectious Diseases	145
Operations	146
Dental Treatment	146
Convalescent Home for Children, Rothbury14	6A-147
Paying Patients	
Staff Sickness1	
Statistics1	
Report of Department of Neurosurgery1	
,, ,, ,, Fever Therapy1	
,, ,, ,, Thoracic Surgery1	
" " Diabetic Clinic	
Maintenance in other Institutions	
VI.—FOOD—	
BOVINE TUBERCULOSIS, AND THE INSPECTION OF MEAT AND	
Provisions and Food and Drugs-	
Tuberculous Milk	181
Report of the Veterinary Officer, Inspector of Meat, etc	
Bovine Tuberculosis	82-183
Milk and Dairies Order, 1926.	
The Milk (Special Designations) Order of 19361	
Cow-keepers	
Anthrax	
Meat, Provisions, etc., Inspection of	
Animals Slaughtered	
Animals Condemned	
Public Health (Meat) Regulations, 1924	
Imported Foodstuffs	
Caseous Lymphadenitis	
Foreign Meat, etc., arriving by Vessel	
Exported Foodstuffs	191
Food Destroyed as Unfit for Consumption1	
Microscopical Examinations	195
Slaughterhouses	95-196
Merchantise Marks Act, 1926	
Rats and Mice (Destruction) Act, 1919	
Contraventions	198
Report of the Chief Sanitary Inspector—	
Food and Drugs Adulteration	SA-200
Offences other than Adulteration	201
Condensed Milk	202
Artificial Cream Act, 1929.	202
Milk and Water—Bacterial Examinations	202
Milk Churns	202
Premises on which Food is prepared2	
Ice Cream	203
Milk and Dairies (Amendment) Act, 1922	204
Milk and Dairies Order, 1926.	204

CONTENTS-Continued.

VII.—THE HOME AND THE WORKSHOP—

Nuisances, Housing, Factories and Workshops, etc.— Report of the Chief Sanitary Inspector—

	PAGE.
Introduction	207
Nuisance Abatement.	208
Notices served	208
Magisterial Proceedings	208
Rent and Mortgage Interest Restrictions Acts	208
Conversion of Dry Closets to Water-Closets	8-209
Atmospheric Pollution, Smoke Abatement	210
,, Observation Station Records2	11-214
Sunshine	215
Sanitary Inspection of Theatres, Cinemas, etc21	15-216
Offensive Trades	216
Summary of Nuisances dealt with	7-218
Visits and Inspections in respect of Nuisances	219
Summary of Legal Proceedings	220
Housing—	
Housing Act, 1935 (Overcrowding)	21-224
Housing Acts, 1925 to 1935 (Inspections)	224
Ministry of Health Table 22	
Housing Act, 1930, Section 17, 19, 20, 55, 5622	27-228
Clearance Areas22	
Houses Demolished	230
Houses Built	230
Tents, Vans, etc.	230
Tenemented Houses	231
Tenement Bye-laws	231
New Buildings and Sanitary Alterations	231
Common Lodging Houses	31-232
Factories and Workshops-	
Inspections	32-233
Outworkers	233
Defects found and dealt with	234
Trades carried on in Workshops of City	235
Council and other Schools.	235
Rag Flock Acts 23	35-236
Exhumations and Re-Interments.	236
Fertilisers and Feeding Stuffs Act, 1926.	236
Merchandise Marks Act, 1926.	236
Agricultural Produce (Grading and Marking) Act, 1928	237
Pharmacy and Poisons Act, 1933	237
Shops Act, 1934	237
Staff Changes and Conclusion	238

APPENDICES "A and B"-

- "A" A Study of the Diets of Sixty-nine Working Class Families.
- "B" Domiciliary Medical Services—Joint Medical Relief District, Period 1st March, 1935—29th February, 1936.

Members of Council who served on the

HEALTH COMMITTEE.

The Lord Mayor (Alderman JOHN GRANTHAM, J.P.)

Councillor Walter Thompson, J.P., Chairman.

Alderman David Adams, J.P., M.P., Vice-Chairman.

Alderman J. Chapman.

,, J. Moore, J.P.

" W. Locke, J.P.

Councillor Catherine A. Auld, J.P. Councillor May Newton.

,, Н. Моат.

R. M. Rowe.

J. Pearson, J.P.

T. M. TAYLOR, J.P.


A. E. Bedson.

CATH. A. LOCKE, J.P.

J. E. Scanlan, O.B.E., J.P.

E. F. WEIDNER.

Table showing the various Sections of the Health Committee's work which is under the direct charge of the Medical Officer of Health.



MATERNITY AND CHILD WELFARE COMMITTEE.

```
*Alderman John Chapman, Chairman.
```

*Alderman David Adams, J.P., M.P.

*Councillor Walter Thompson, J.P. †Dr. R. P. Ranken Lyle, J.P.

* ,, J. E. SCANLAN, O.B.E., †Dr. J. C. SPENCE. I.P.

* ,, MAY NEWTON. †Mrs. E. I. LEACH.

* ,, J. Pearson, J.P. ‡Councillor Jeanie L. Gibbin, O.B.E., J.P.

* ,, Cath. A. Locke, J.P. ‡ ,, Frances E. Taylor, J.P.

† Co-opted member.

‡ Appointed by City Council.

^{*}Councillor Catherine A. Auld, J.P., Vice-Chairman.

^{*} Member of the Health Committee.

STAFF.

- J. A. CHARLES, M.D., B.S., F.R.C.P., D.P.H., Medical Officer of Health and Medical Superintendent of the City Hospitals for Infectious Diseases.
- E. F. DAWSON-WALKER, M.D., B.S., B.Hy., D.P.H., Deputy Medical Officer of Health.
- WM. GRAY, Senior Sanitary Inspector.
 - JAS. McNichol, Chief Assistant Inspector and Assistant Workshops Inspector.
 - ISAAC CLARK, Assistant Workshops Inspector (retired July).
 - W. E. Perkins, Assistant Workshops Inspector (Appointed July).
 - A. FLOCKHART and A. KIRSOP, Assistant Inspectors under the Food and Drugs Acts.
 - L. W. Johnson, J. Brown, L. Wade, M. Swales (resigned 31st Dec.), H. W. Grieves (resigned July), R. G. Suddick (resigned Feb.), A. Ibbitson, W. M. Pettigrew, R. S. Cooper, E. Banks, T. Sayer, D. Wilkinson, District Inspectors.
 - W. Stewart, F. Galton, F. James (April), F. Thompson (Temp. Aug.), Slum Clearance Inspectors.
 - N. MAYNE, Assistant Inspector of Common Lodging Houses.
 - WM. BEAN, R. CHAPMAN (Temp.), J. SHIPLEY (April), Infectious Diseases Inspectors.
 - Jas. Robson, Jas. Bruce, Jno. R. Cragie, J. W. Robson, Thos. Moore-J. Robson, Jun., Ambulance Drivers and Disinfectors.
- *ALFRED HEDLEY, M.S.M., *GEO. CUTHBERTSON, *ALEC. M. WALKER, JOS. GILHESPY, H. G. OLIVER, *D. H. MACPHERSON, *R. DOBBIN, H. G. COATES, *F. PELLATT, *L. SMALLEY, A. CAMPBELL, L. WHITEMAN (appointed June), R. A. RIDLEY (Temp.), Clerks in the Health Department. ALICE FENWICK, E. STOBART (Temp.), A. E. BLAIR (Temp.), (Typists),

Those marked * hold the Sanitary Inspector's Certificate of the Royal Sanitary Institute.

- THOS. PARKER, F.R.C.V.S., Veterinary Officer and Inspector of Provisions, H. Thornton, M.R.C.V.S., B.V.Sc., D.V.H., Assistant Veterinary Inspector, Jas. M. Anderson, W. Cockburn, Geo. Phillips, Assistant Inspectors of Provisions. *Norman Dickson, Clerk.
- A. F. G. SPINKS, M.D., Maternity and Child Welfare Medical Officer.
 - E. G. Brewis, M.D., B.S., M.R.C.P., D.P.H., Assistant Maternity and Child Welfare Medical Officer (appointed Jan.).
 - a Georgina B. Cameron, M.B.E.*, Chief Health Visitor and Supt. of Midwives.
 - t Catherine M. Thextont, b Marion Moody*, c Lizzie Isa Pritchard, c Louise Shell, d Florence Martha Hatfield*, d Norah B. Willson*, b E. Hisco*, b E. Johnson*, b N. E. Carr*, b T. Mason*, b E. M. Hastie*, b N. Lewis*, b M. A. Simpson*, b N. Thompson*, g C. N. Phillips, b D. A. Atkinson, b M. Batty, b A. Craggs, b P. E. Pearce, b R. Roxby, b M. Scorer, b E. G. Sayer, b L. Youell, b A. Bradley, b C. Barron* Health Visitors. Edith Rodgers, Marion S. Batt, Blanche Dickson, A. Dougal, Clerks.
- (Qualifications of those marked a C.M.B., General and Fever Nursing and R.S.I. Certificates.

 b C.M.B., General Nursing and R.S.I. c C.M.B. and R.S.I. d C.M.B. and General Nursing.

 1 C.M.B., Fever Nursing and R.S.I. g C.M.B.).
 - * State Registered Nurse. † State Registered Fever Nurse.

IRENE COOK, S.O.N.A., Orthopædic Nurse.

Annie G. Bainbridge, Superintendent of Welfare Centres (retired March).

GLADYS PATTISON (resigned Sept.), IRENE GAWMAN, CATH. BARNES, MARY E. MUSE, EDITH TAYLOR (Temp. appointed Aug.), VIOLET SIMPSON (Temp. appointed Aug.), Clerks.

H. GLEN DAVISON, M.D.

L. MABEL R. CAMPBELL, M.B., Ch.B.

H. HARVEY EVERS, M.B., F.R.C.S., (resigned Oct.).

GERTRUDE H. G. HICKLING, M.D., Ch.B., B.Sc., D.P.H.

C. N. Armstrong, M.B., B.S., M.R.C.P., B.Hy., D.P.H.

A. G. OGILVIE, M.B., B.S., M.R.C.P.
ANNE FAIRWEATHER, M.D., B.S.,
B.Hy., D.P.H., D.P.M. (Psych, and
Ment. Def.).

F. E. STABLER, M.D., B.S., F.R.C.S.

C. C. Ungley, M.D., M.B., B.S., F.R.C.P., M.R.C.S.

DOROTHY HOPKINSON, M.B., B.S.

NORA LONG, M.B., B.S.

ELSIE B. WRIGHT, M.D., B.S., M.R.C.S.,

Assistant Medical Officers (part time) Welfare Centres.

G. HURRELL, M.D., B.S., B.Hy., D.P.H., Tuberculosis Medical Officer.

N. V. Hepple, M.D., M.B., B.S., D.P.H. B.Hy., Assistant Tuberculosis Medical Officer (resigned Aug.).

John Stokoe, м.в., в.s., Assistant Tuberculosis Medical Officer (appointed February.

WM. H. DICKINSON, O.B.E., M.D., Ch.B., M.R.C.P.(Ed.), D.P.H., Tuberculosis Medical Officer (part time).

c Constance M. Bayne, d Annie Booth, a W. E. Dale*, b J. P. Kenmir*, e M. Young, Tuberculosis Visiting Nurses.

(Qualifications of those marked a General Nursing. b General Nursing, C.M.B. and R.S.I. c General Nursing and Health Visitors and School Nurses Certificates of R.S.I. d Fever Nursing. e Fever Nursing and C.M.B.

* State Registered Nurse.

C. Sandilands, Assistant Inspector (appointed August).
Geo. Magnay, Gertrude Gillender, M. Pringle, Clerks.

BARRASFORD SANATORIUM.

C. G. R. GOODWIN, M.R.C.S., L.R.C.P., Medical Superintendent.

HAZEL I. ASHFORD, M.B., Ch.B., Assistant Medical Superintendent (resigned February).

Doris Cohen, M.B., B.S., Assistant Medical Superintendent (appointed Feb.)
Frances Baguley, A.R.R.C., Matron. Sisters, Nurses, Domestic Staff.

CITY HOSPITAL FOR INFECTIOUS DISEASES.

E. F. DAWSON-WALKER, M.D., B.S., B.Hy., D.P.H., Deputy Medical Superintendent.

J. F. Caithness, M.B., Ch.B., D.P.H., Senior Resident Medical Assistant.

L. H. Murray M.B., B.S., Resident Medical Assistant (appointed Aug.)

W. FRANK WILSON, M.B., B.S., Consulting Oto-Rhinologist.

J. L. WATT, Matron.

H. PHILLIPS, Steward.

Jessie Laing, Assistant Matron. Sisters, Nurses, Clerks, Domestic Staff. Maud B. Elliott, Dispenser.

GEO. COCKBURN, Engineer.

Lodge Keepers, Firemen, Porters, Gardeners, Joiner and Handyman.

SMALLPOX AND ISOLATION HOSPITALS.

MATTHEW and ISABELLA ROBSON, Caretakers.

NEWCASTLE GENERAL HOSPITAL.

- G. P. HARLAN, M.D., Ch.B., B.Hy., D.P.H., Medical Superintendent.
- G. F. Duggan, M.B., B.Ch., M.A.O., F.R.C.S. (Edin), Deputy Medical Superintendent.

JUNIOR RESIDENT HOUSE PHYSICIANS AND SURGEONS (6).

A. BARON, Matron.

- S. Lake and A. Lunt, (appointed May), Assistant Matrons. Sisters, Nurses, Domestic Staff.
- N. H. Herdman, Dispenser (resigned Aug.) G. H. Darling, Dispenser (appointed Aug.)
- JAMES MATTHEWS, Steward. Ambulance Drivers, Porters, Male Nurses, Clerks.

CONSULTING STAFF, Etc.

THOMAS BEATTIE, M.D., B.S., F.R.C.P., Medical Director.

F. J. NATTRASS, M.D., B.S., F.R.C.P., Physician.

ELSIE B. WRIGHT, M.D., B.S., M.R.C.S., L.R.C.P., Medical Registrar.

W. G. A. Swan, M.B., B.S., M.R.C.P., Medical Registrar.

JOHN CLAY, C.B.E., M.B., B.S., F.R.C.S., Surgical Director.

J. C. Stewart, M.S., F.R.C.S., Surgeon.

- W. E. WARDILL, M.B., B.S., F.R.C.S., Plastic and Genito Urinary Surgeon.
- G. S. Clark-Maxwell, M.A., M.B., B.Sc., Surgical Registrar.
- A. Logan, M.B., Ch.B., F.R.C.S. (Eng. and Ed.), Surgical Registrar.
- G. A. Mason, M.B., B.S., F.R.C.S., Thoracic Surgeon (appointed April)
- A. R. D. Pattison, M.B., B.S., F.R.C.S., L.R.C.P., Neurological Surgeon.
- S. W. DAVIDSON, M.D., B.S., M.R.C.P., Radiologist.
- A. MacRae, M.A., M.D., Ch.B., D.O.M.S., Ophthalmic Surgeon.
- D. R. MACGREGOR, B.Sc., M.B., Ch.B., D.L.O. (R.C.P.S.), Oto-Rhinologist.

PHILIP AYRE, M.R.C.S., L.R.C.P., Anaesthetist.

- W. J. PHILLIPS, M.B., B.S., Anaesthetist.
- E. Joan Thompson, M.B., B.S., D.A., Anaesthetist (appointed Feb.)
- S. F. Evans, Ph.D., M.Sc., Radiotherapist.

DISTRICT MEDICAL OFFICERS.

Dr. W. Simpson (resigned Feb.), Dr. R. W. Nevin, Dr. T. J. Ryan.

PUBLIC VACCINATORS.

Drs. J. MacRae (died November, A. Sutcliffe appointed Temp. November), Richard Dagger, T. J. Ryan, H. R. Smith (died March, L. S. Davison appointed Temp. March, Terminated July, W. A. Slater appointed June), A. M. Paterson, J. A. Brand, G. P. Harlan (Newcastle General Hospital). H. L. Taylor, S. Fullerton, H. R. Kendal.

VACCINATION OFFICERS.

EASTERN DISTRICT—W. H. F. GARRETT. WESTERN DISTRICT—W. W. CUMMINGS.

To Councillor WALTER THOMPSON, J.P., Chairman of the Health Committee of the Corporation of Newcastle upon Tyne.

SIR,

I have the honour to present the sixty-fourth Annual Report of the Medical Officer of Health on the sanitary conditions of the City.

The practice which has been followed in previous years whereby the detailed reports of the work of the several sections of the Health Department are submitted by the officers responsible has not been varied, and these documents constitute the major portion of the text.

In this letter, which is not intended either to summarise or to consolidate the reports of the sub-departments—for these of themselves are important contributions to our health records some comment will be made on the outstanding features of the year 1936, and the more urgent of our many continuing problems will be reviewed.

Vital Statistics-Marriage and Birth Rates.

The vital statistics of 1936 were in several ways remarkable and the more noteworthy details will now be referred to. Marriages were more numerous both actually and relatively than in any year since 1921. The total of 2,474 is only seven more than the record for 1935, but it exceeds the average for the three year period 1933—1935 by 124, and is equivalent to a marriage rate of 17.0 persons per 1,000 of the population.

The associations of prosperity and depression with the fluctuations in the marriage rate have been discussed in these pages on previous occasions. For the year 1936 it is notable that the matrimonial increase goes hand in hand with a more than welcome reduction of 6,387 persons in the number of the male and female unemployed. This return to employment was undoubtedly the most significant feature in the life of the City during the year.

The marriage rate has been increasing since 1931, but curiously enough the city birth rate does not yet reflect this fact. The birth rate for 1936 was 15.6 per 1,000, a figure considerably lower

than any hitherto recorded, though we may derive what little consolation we may from the knowledge that it is still higher than the rates for England and Wales as a whole, and for the 122 great towns which were 14.8 and 14.9 respectively.

Thirty years ago, in 1906, the Newcastle birth rate was 31.9, or rather more than twice the present rate. The effects of this change in the fertility of the population are met with in every phase of our social and economic life.

Maternal and Infantile Mortality.

Before dealing with the general death rate for the City something must be said of those special death rates which are concerned with the mortality of childbirth and of infancy. In neither case are the records comforting or complimentary. The local maternal mortality rate was equivalent to 5.92 deaths of mothers per 1,000 live and still births. This rate is the highest ever recorded in the City, and is disappointing in the extreme. It is 50 per cent, higher than the national rate, and even when we make every possible allowance for the greater completeness in statistical recording in recent years, for the evils of overcrowding and poverty which are still so prevalent, for the somewhat haphazard character of the midwives service, now happily to be co-ordinated, there still remains a doubt, not easily to be dismissed, that something is lacking either from the planning or in the execution of our midwifery services. Whether that deficiency is to be found in the hospital arrangements, or in the work of the medical and midwife personnel, or in the central direction of the scheme, is a matter which calls for careful enquiry and investigation.

The incidence of infantile mortality similarly gives us no cause for self-congratulation. Following upon a rate of 83 in 1934, and 86 in 1935, the infantile mortality rate in 1936 reached the high figure of 90 per 1,000 births. This should be contrasted with the national rate of 59, and the rate recorded for the 122 great towns which was 63.

Of the 408 deaths which occurred under the age of one year, 208 were attributable to the so-called neo-natal conditions of the first month of life, while the infective conditions of infancy, measles, whooping cough, bronchitis, broncho-pneumonia, diarrhœa and enteritis were responsible for 117 of the remaining 200. Actually the bad record of 1936 is due to a number of outbreaks, both

domiciliary and institutional, of infective enteritis. Such epidemics can be prevented if the simple rules of personal cleanliness are regarded by all those entrusted with the care and feeding of infants. In many cases observance of these elementary principles is rendered difficult by insanitary home conditions, or by the overcrowding and consequent relative understaffing which is so much a feature of our local hospitals.

The whole of the accommodation for the treatment of sick children which is provided in the various institutions of the City has long been overdue for reconstruction, and though their respective managing bodies are aware of this fact, the financial difficulties have not been appreciated or resolved by the general public.

In the Annual Report for 1935, reference was made to the investigation which had been carried out into the comparative frequency of deaths in the first month of life; *i.e.*, of the neo-natal mortality—amongst infants born of normal labours in hospital and under home surroundings respectively.

It was stated that the chances of survival of the home-born infant were greater at any rate in Newcastle, than those of the child born to a mother confined in hospital. Our enquiries along these lines have been continued and the evidence which has been collected in the local hospitals supports the general statement already made.

Even though the enquiry was limited to the subsequent history of children born of normal labours, it is appreciated that there may have been conditions in the mother necessitating her admission to hospital, which might in turn have been detrimental to the newly born child. It would be unwise, therefore, without wider knowledge of all the circumstances, to stress unduly the fact that a normal confinement in hospital may be prejudicial to the future welfare of the infant. But the evidence does suggest and suggest strongly, that overcrowded maternity wards and inferior maternity hospital accommodation, do play some part in the causation of neo-natal mortality, and that just as our children's hospitals are in need of reconstruction, so also are those institutions which for many years have given at all times refuge and succour to women in labour.

It is pleasing therefore to report that plans have been prepared for the erection of a new Princess Mary Maternity Hospital of 140 beds on the Castle Leazes site; and that the existing accommodation at the Newcastle General Hospital is to be replaced by a modern unit to comprise 30 beds in the first instance.

The General Death Rate.

The general death rate for the City showed a slight increase and rose from 12.6 per 1,000 population in 1935 to 13.1, the highest rate since 1931. This crude rate represents the occurrence of mortality in the Newcastle population, which of course is not constituted in the same proportions as regards the age and sex of its members as is the standard population of England and Wales. By means of comparability factors, which have been prepared by the Registrar General, it is now possible to adjust these differences in the population of individual towns and counties, and to place them all on a common basis of comparison.

In Table I. are recorded the crude and adjusted death rates for all cities in England with a population of 250,000 and over, together with similar information for certain local towns and counties.

TABLE I.

Name of Town.	Population as estimated by Registrar General, Mid. 1936.	General Death Rate.	Death Rate adjusted by Comparability Factor.
Portsmouth	251,400	11.8	11.7
Leicester		11.6	11.8
Bristol	413,900	12.3	12.0
Birmingham		11.3	12.4
London County	4,141,100	12.4	12.6
West Ham		11.6	13.3
Nottingham		13.2	13.6
Sheffield	518,200	12.2	13.8
Hull	321,500	12.7	14.0
Leeds	489,862	13.6	14.6
NEWCASTLE UPON TYNE		13.1	14.8
Liverpool	846,400	12.9	14.8
Bradford	291,085	14.9	14.9
Stoke-on-Trent	273,100	12.4	15.1
Manchester		13.5	15.4
Northumberland County	406,550	12.0	12.7
Tynemouth	66,800	11.6	12.8
Durham County	897,287	11.9	13.6
Sunderland	184,179	12.8	14.3
Middlesbrough	140,000	12.2	14.4
South Shields	111,800	13.2	14.8
Gateshead	119,034	13.2	14.9

As regards the actual order of merit it will be noted that our position is in the lower half of the national table, while amongst the local counties and county boroughs we are superior only to Gateshead.

Mortality from certain special causes.

Turning now to the individual causes of death the six most important of these are set out in order below (Table II.) The figures and positions for 1935 are reprinted for the purpose of comparison.

TABLE II.
INDIVIDUAL CAUSES OF DEATH.

		1936.		
No.	Cause of Death.	Number.	Percentage of Total Deaths.	
1	Diseases of the Heart	871	22.5	
2	Cancer	413	10.6	
3	Diseases of Veins and Arteries	412	10.6	
4	Bronchitis and Pneumonia	366	9.5	
5	Diseases of Nervous System	276	7.1	
6	Tuberculosis—Pulmonary	265	6.8	

		1935.			
No.	Cause of Death,	Number.	Percentage of Total Deaths.		
1	Diseases of the Heart	701	19.1		
2	Cancer	433	11.8		
3	Diseases of Veins and Arteries	406	11.1		
4	Bronchitis and Pneumonia	358	9.7		
5	Diseases of Nervous System	243	6.6		
6	Tuberculosis—Pulmonary	240	6.5		
		Li selia	Janes		

It will be seen that the relative ranking of these major causes of death is the same as in 1935. Nevertheless, there are at least three points upon which some emphasis must be laid.

In the first place the steady increase in deaths from diseases of the cardio-vascular system (i.e., heart, veins and arteries) still continues. The lives of 1,283 persons ended as the result of some form of cardio-vascular disease or failure. In 1934 and 1935 the comparable totals were 935 and 1,107 respectively. In the past three years the proportion of these deaths to deaths from all causes has risen from 25.6% (1934) through 30.2% (1935), until in 1936 it was 33.1% or practically one-third of the total mortality.

Secondly, the number of deaths from cancer has fallen from 442 in 1934 and 433 in 1935 to 413 in 1936. But these figures of themselves signify nothing as there have been many years in which even smaller totals of cancer deaths have been recorded.

For the first time, however, a detailed enquiry has been made into the deaths from cancer occurring in Newcastle citizens, and an attempt has been made to obtain and to correlate information on the following, amongst other, points:—

- (a) whether the patient delayed unduly in seeking his doctor's advice,
- (b) whether the doctor diagnosed the patient's condition promptly and took steps thereafter to obtain appropriate treatment.
- (c) whether there was institutional delay in the provision of treatment,
- (d) whether every possible form of treatment was available, and
- (e) whether in certain patients the disease progressed to an unfavourable conclusion despite everything that could be done.

Generally speaking the answers to these questions obtained from an investigation of 202 cases were as follows:—

(a) There was frequently reluctance on the part of the patient to consult the doctor until the disease was well established. No fewer than 11 per cent. of patients fell into this category.

- (b) In nine cases out of ten the diagnosis of the patient's condition was made promptly by the medical practitioner. In the remaining tenth case, owing to a more prolonged investigation, or the indefinite nature of the symptoms, diagnosis was somewhat delayed. Nevertheless, immediately diagnosis had been established, steps were invariably taken to obtain adequate treatment.
- (c) Admission to hospital, provided the patient acquiesced in this course, followed almost immediately, with at the most, only a few days' delay.
- (d) Adequate treatment, whether by surgery or by radium, was available for all City cases. Deep X-ray therapy was also available for the majority of patients requiring it, but here the need for further facilities could not be denied.
- (e) A large number of cases, amounting to one-third of the whole, proved to be completely resistant to every form of treatment, even when these were applied promptly and efficiently after early diagnosis.

To one result of this enquiry attention must be gratefully directed. The Schools and Charities Committee of the Corporation, having been informed of the trend of these investigations, offered to establish and maintain at the Newcastle General Hospital a fully equipped and staffed Deep X-ray Therapy Department. It is unnecessary to say that this offer, conceived in a spirit of the truest service to the community, was wholeheartedly welcomed and accepted by the Health Committee. The department will be completed in 1938, and both the Health Committee and the Schools and Charities Committee will be associated in its administration.

Thirdly, as regards pulmonary tuberculosis, the total number of tuberculous deaths of which 265 were recorded is 25 higher than the total of the previous year, and the percentage of the total death roll which is due to this disease is also slightly greater.

This is a reminder, if any were required, that in the City as on Tyneside generally the conquest of tuberculosis is not yet complete, and that unremitting attention and vigilance are necessary to keep it within such measure of control as we now possess.

TABLE III.

Average Death Rates per 100,000 in England and Wales and Newcastle upon Tyne during the Nine Year Period 1927-1935. (Based upon the Registrar-General's Abridged List of Causes of Death.)

			1	Newcastle
		England	Newcastle	as a per-
No.	Causes of Death.	and	upon	centage of
		Wales.	Tyne.	England
				and Wales
(1)	(2)	(3)	(4)	(5)
1-7	(-/	(-)	(-/	
	All causes	1,211	1,288	106.3
	*Infantile Mortality	64.4	82.7	128.3
1.	Typhoid and paratyphoid fevers	0.73	0.78	106.8
2.	Measles	8.2	17.6	214.6
3.	Scarlet Fever	1.7	3.0	176.5
4.	Whooping Cough	7.4	10.7	144.6
5.	Diphtheria	7.8	4.5	57.7
6.	Influenza	35.6	27.5	77.2
7.	Encephalitis lethargica	2.3	3.1	134.8
8.	Cerebro-spinal fever	2.0	5.2	260.0
9.	Tuberculosis of respiratory system	71.6	100.7	140.6
10.		20.6	24.6	119.4
11.	Other tuberculous diseases			200.0
	Syphilis	3.4	6.8	200.0
12.	General paralysis of the insane,	10	0.0	444.7
10	tabes dorsalis	4.8	6.6	141.7
13.	Cancer, malignant disease	148.5	142.2	95.8
14.	Diabetes	14.6	14.6	100.0
15.	Cerebral haemorrhage, etc	66.1	54.1	81.8
16.	Heart disease	241.9	226.7	93.7
17.	Aneurysm	3.3	3.6	109.1
18.	Other circulatory diseases	60.2	92.6	153.8
19.	Bronchitis	58.1	58.1	100.0
20.	Pneumonia (all forms)	80.2	94.3	117.6
21.	Other respiratory diseases	12.9	13.4	103.9
22.	Peptic ulcer	10.3	11.1	107.7
23.	Diarrhoea, etc. (under 2 years)	10.2	18.7	183.3
24.	Appendicitis	7.3	6.0	82.2
25.	Cirrhosis of liver	3.9	2.9	74.4
26.	Other diseases of liver, etc	6.4	7.6	118.7
27.	Other digestive diseases	†	+	†
28.	Acute and chronic nephritis	38.7	44.6	115.2
29.	‡Puerperal sepsis	1.7	1.9	111.8
30.	Other puerperal causes	2.5	2.7	108.0
31.	*Congenital debility, premature	2.0		100.0
180000	birth, etc.	32.4	34.9	107.7
32.	Senility	46.8	28.5	60.9
33.	Suicide	13.1	11.5	87.8
34.	Other violence	41.7	39.2	94.0
35.	Other defined causes	†	+	4
36.	Causes ill-defined or unknown	4	+	+
00.	oauses in-defined of unknown			

^{*} The rates for these headings are per 1,000 live-births.

[†] Not extracted.

[†] The rates for these headings are per 1,000 live-births for 1927 and per 1,000 live and still births 1928 and onwards.

Death rates which are equivalent to 125 per cent. or more of the similar rates for England and Wales are indicated in heavier type.

Table III. is of particular interest, as it indicates at a glance the average death rates for England and Wales and for the City, from the 36 causes set out in the Registrar General's Abridged List. The period over which the rates have been averaged is now one of nine years, and the fifth column shows the significant differences between our own and the national record.

The following are the causes of deaths of which our experience is 25 per cent. greater than the average of the country as a whole—infantile mortality, measles, scarlet fever, whooping cough, the epidemic diseases of the central nervous system, tuberculosis of respiratory system, syphilis, general paralysis of the insane and tabes dorsalis, certain diseases of the circulatory system, and diarrheea in children under two years.

Infectious Diseases.

The notifiable infectious diseases, other than tuberculosis, and those minor infectious conditions of which whooping cough and diarrhœa are the outstanding examples, were responsible for 308 deaths as compared with 303 in 1935. The distribution of the deaths amongst the individual diseases showed considerable variation, there being a marked reduction in the mortality from pneumonia and from whooping cough. These improvements, however, were more than offset by the deaths in the outbreaks of diarrhœa and enteritis to which reference has already been made. No fewer than 105 children under the age of 2 years died from these conditions. During the previous year there were 65 deaths in this age group from the same cause.

Of the other infectious diseases scarlet fever showed the most marked decline and the total of 937 notified cases was the smallest since 1930. The case mortality rate, based upon eight recorded deaths, was 0.80 per cent., which is only slightly higher than the average for the past 10 years.

As regards diphtheria, the epidemic manifestation of this disease, which was the subject of special comment in the last Annual Report, still continues. The number of cases increased from 675 to 693, which constitutes a record for the City. 35 deaths occurred and the case mortality rate of 5.1 per cent. is approximately the same as in the previous year. Other cities in the North of England, notably Leeds and Hull, have experienced similar epidemics in recent years, and we are probably fortunate

that both the incidence of the disease and the fatality have not been greater. Undoubtedly the immunisation clinics which have been in operation since 1934, have contributed in protecting large numbers of children from this serious infection.

Measles has again happily failed to appear in anything like a major epidemic. Although 4,022 cases were notified, the case mortality was very low, namely 0.42 per cent. But even this low mortality represented the loss of 17 young lives, or more than twice as many as died from scarlet fever. Nor must we forget the complications, the "running ears," and "delicate chests," which will be a continually recurring cause of invalidity in later years. The reduction in the mortality and morbidity rates of measles is almost entirely a question of the avoidance of over-crowding, and the improvement of the home environment. Even if our great housing schemes pay no other dividend than the relegation of measles and whooping cough to the position of truly minor infectious disorders, they will have proved worth while.

Nutrition.

In December, 1936, the report on the dietary survey of 1934 was published under the title of "A Study of the Diets of Sixty-nine Working Class Families in Newcastle upon Tyne," and a copy of this document is included as Appendix A. of the present report.

The reception given to the survey report by the medical and lay press alike was exceedingly favourable, and the many anonymous individuals, who had been concerned in its preparation, found therein a gratifying acknowledgment of their labours. More particularly is it necessary to stress the extent of our debt to the Medical Department of the Ministry of Health not only for information which was freely placed at our disposal, but also for much unofficial assistance.

One criticism of the report was to the effect that a knowledge of the economic and dietary conditions of the families investigated in 1934, was of little value in 1936. But such a criticism neglects altogether the importance of having the results of a carefully organised survey for use as a starting point and datum line for future enquiries.

In point of fact, a re-survey of the financial circumstances, though not of the diets of 64 of the 69 families has been made during the course of the present year, and has yielded most interesting results. Many of the features which were the subject of comment in the 1934 investigation are still in existence to-day. In other directions changes have undoubtedly occurred.

For example, no fewer than 16 of 35 traced unemployed families are now in the employed group, with larger incomes and greater spending powers. But the conditions of 24 families who had been employed in 1934 deserve some special attention. Three of them were found to be unemployed, and the remaining 21 though still employed were not, on the whole, as well off economically as they had been in 1934. Such minor increments as had been added to the incomes of these latter families were more than counterbalanced by (a) higher rentals and (b) the increased size of the family.

When these two factors have been taken into account, the average family in this group (i.e. those who were employed both in 1934 and 1937) has less money available now than it had in 1934, for the other outgoings of the family budget. The family manager has then to decide whether she will spend less on food or whether she will maintain the dietary expenditure at the expense of other sections of the budget. When it is realised that the present cost of foodstuffs in Newcastle is 16 per cent. higher than in 1934, it would appear probable that either less food is being consumed by the families under discussion, or that the same quantities of nutrients are being obtained from cheaper sources. What precisely happens could only be determined by another dietary survey. At any rate, it is obvious that we are only at the beginning of our understanding of this linked problem of family incomes and family dietaries.

Domiciliary Medical and Nursing Services.

A special report describing the working of the "open choice" method of providing domiciliary medical services which has been adopted in eight of the ten medical relief districts of the City was published in October, 1936, and is included in the present Annual Report as Appendix B.

There is nothing to add to the information therein set out, and to one point only is further reference indicated. Since the period covered by the special report, namely, from March 1st, 1935, to February 29th, 1936, full co-operation with the nursing associations has been achieved, and the value of the work of these organizations on behalf of the sick poor of the City can be gratefully acknowledged.

General Hospital Services.

The steady progress and continuous development of the work of the Newcastle General Hospital is seen in the columns of Table IV., in which are recorded the main statistical data since the transfer of the hospital to the City Council.

TABLE IV.

Year.	Admissions.	Discharges.	Operations.	Maternity Cases
1930	3,048	3,099	596	97
1931	3,598	3,574	1,125	99
1932	4,522	4,447	1,428	161
1933	4,776	4,763	1,560	194
1934	5,544	5,555	2,076	225
1935	6,245	6.215	2,722	273
1936	6,707	6,695	2,722	388

More detailed information is given in the pages of the hospital report itself, which now incorporates special sections relating to the activities of the important departments which have been established for neuro-surgery, fever therapy, diabetes and thoracic surgery. These latter departments are an outstanding tribute to the energy and industry of the consulting staff of the hospital, and to the foresight of the Health Committee which has encouraged and fostered these newer enterprises.

The Re-organisation of the Municipal and Voluntary Hospitals in the City.

Towards the end of the year, schemes for the complete reorganization of the City Hospital for Infectious Diseases and the Newcastle General Hospital, were submitted to the Commissioner for the Special Areas, with a request that they should be considered as being eligible for grant.

The schemes submitted in respect of the City Hospital for Infectious Diseases provided for the re-development of the Walker Gate site utilising the areas east and west of Benfield Road.

As regards the Newcastle General Hospital, the proposals put forward were based on the assumption that the Elswick Grange Institution would be removed to Benton, and that the site vacated would be available for hospital purposes. A comprehensive scheme, extending the hospital to 944 beds, with admission block, out-patient department, tuberculosis dispensary and maternity and child welfare centre, has been prepared by the City Architect, Mr. R. G. Roberts, F.R.I.B.A., and irrespective of the receipt of a grant from the Commissioner's funds, should form the ground plan for the future development of the hospital.

Applications of a similar character have been laid before the Commissioner by the governing bodies of a number of the voluntary hospitals in the City. These proposals, when considered alongside the schemes for the re-development of the City Hospital for Infectious Diseases and the Newcastle General Hospital, constitute a notable contribution to the problem of the re-organization and rationalisation of the hospital services of the City and of the adjacent areas. So far, although certain of these schemes were presented to the Commissioner upwards of two years ago, there has been no indication as to whether he will regard them as ranking for grant and if so, what proportion of the total cost will be defrayed from the funds at his disposal. It is to be hoped that the difficulties which the Commissioner is apparently experiencing in formulating his policy on these matters, will be presently resolved.

Whitton Tower Convalescent Home for Children.

One other item of re-organization may be referred to here-As a result of the generosity of Councillor Angus Watson, J.P., the City now possesses in Whitton Tower, Rothbury, one of the most beautifully situated and splendidly equipped convalescent homes for children. Suitable patients are transferred there from both municipal hospitals, and its facilities are also available for City children from the voluntary hospitals.

Through its possession we have been enabled to fill one more of the gaps in our defences, and the success of Councillor Watson's inspiration is visible in the rapid restoration to health of children with long histories of ill-health in their ordinary surroundings.

(A photograph of the Home is inserted opposite page 147).

Slum Clearance and Overcrowding.

During the year two public enquiries under the Housing Act of 1930 were held and 11 Clearance Orders and three Compulsory Purchase Orders were submitted for confirmation to the Minister of Health.

These orders comprised a total of 545 houses in which were accommodated 1,141 families, with a total population of 3,930 persons.

The actual amount of slum clearance accomplished during the years 1935 and 1936 has been on the whole rather less than that effected in the years immediately succeeding the introduction of the 1930 Act, but the slowing down of the set programme is only temporary. As soon as certain difficulties which have been experienced by the Housing Committee have been removed, progress should be continuous in every direction.

The following is a summary of the work which has been carried out in the campaign between 1931 and 1936.

Ninety Clearance Orders and six Compulsory Purchase Orders have been made, and have necessitated the holding of eleven public enquiries.

These Orders as confirmed by the Minister have dealt with 2,332 premises in which were housed 5,315 separate families, with a total population of 19,477. The activities of the Health Committee so far, as represented by the ninety-six Orders referred to, and by action taken against individual unfit houses will result eventually in the re-housing of no fewer than 19,436 persons or 6.7 per cent. of the total population of the City. This total will have been exceeded to a very considerable extent before the full programme has been completed.

During the first six weeks of the year the Census of Over-crowding in accordance with the requirements of the Housing Act of 1935, was taken by a band of specially appointed enumerators and clerks. The results of the Census are of historical interest, and are summarised in the report of the Chief Sanitary Inspector. The percentage of the population who were ascertained to be living in conditions of overcrowding as defined on the admittedly low standards of the Act, was no less than 10.7 per cent.

Newcastle is therefore amongst the most densely overcrowded areas in the country. Amongst the greater cities, namely, those with a population of one-quarter of a million and upwards, only West Ham and Liverpool with overcrowding percentages of 8.4 and 7.4 respectively, have records which are in any way comparable to our own.

Conclusion.

There remains, Sir, the very pleasant privilege of thanking yourself, for the first time in your capacity as Chairman, your predecessor, who is now the occupant of the Vice-Chair, and the members of the Committee for the unfailing kindness and helpfulness which have been shown to me during a year when encouragement was exceedingly welcome.

The staff of the Department, and Dr. E. F. Dawson-Walker and Mr. A. Hedley in particular, have placed me more than ever in their debt for their willingness at all times, and for their cooperation and enthusiasm which together have lightened so many labours.

I am, Sir,

Your obedient servant,

J. A. CHARLES,

Medical Officer of Health.

Health Department, Town Hall,

> Newcastle upon Tyne, July, 1937.

SUMMARY OF STATISTICS, 1936.

Population (estimated mid. 193	6)	****	****	10.00	290,400
Area of City (acres)	****	****		****	11,401
Estimated number of houses	****				77,758
Rateable value	2				£2,543,478
Sum produced by 1d. rate	****				£9,919
Births				4419	4,537
Birth rate (per 1,000 populatio	n) 🤐	****			15.6
Marriages		****			2,474
Deaths					3,878
Death rate (per 1,000 population					13.1
,, ,, ,, adjust	ed by co	mparal	oility fa	ctor	14.77
Infantile Mortality (deaths unde	er one y	ear per	1,000	live	
births)	****				90
Natural increase in population (e	xcess of	births	over de	aths	
in the year)					659

CHIEF CAUSES OF DEATH.

Cause.				Number.	Percentage of total deaths.
Diseases of the Heart				871	22.5
Cancer			****	413	10.6
Diseases of the veins and arte	ries			412	10.6
Bronchitis and pneumonia				366	9.5
Tuberculosis (all forms)	****			308	7.9
Do. (Pulmonary)				265	6.8
Diseases of the nervous system	1	1	****	276	7.1
Diseases of the genito urinary	system		1117	164	4.2
Diseases of early infancy, and	congeni	tal ma	lfor-		
mations under 1 year		****		191	4.9

INFECTIOUS DISEASES.

Disease.				Cases notified.	Number of deaths.	Death rate per 1,000 population
Scarlet fever		****	****	937	8	0.028
Diphtheria				693	35	0.117
Enteric fever	,		****	8	-2	0.007
Erysipelas		****	****	176	12	0.041
Cerebro-spinal fever		****	****	8	7	0.024
Measles	****			4,022	17	0.059
Tuberculosis (all forms)	1111			584	308	1.040
				(new cas	es)	

Whooping cough, which is not notifiable, caused 7 deaths. Influenza, which is not notifiable, caused 38 deaths.

Health Report, 1936.


I.—GENERAL.

MORTALITY TABLES, SOCIAL CONDITIONS, CLIMATOLOGY, WATER SUPPLY, DISPOSAL OF REFUSE. ARTE ROLL SATEADWER TO TENDES, GHA TEN

Health Report. 1936

SPIRE TORRESTORS, CLIMATO

												Beaute			-			Street S	area.				None					an from				Barre			Hone					70-86									
										-		-	110	-		-120	13	1			-	=	-		100	, the	王	100	無	700 100 100 100	E .	1								100							王		
01.20 (01.00 (01	0.00 mm m		100 (100 (100 (100 (100 (100 (100 (100	77 2 2 2 3 4 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	1		ADD ADD ADD ADD	CONTROL CONT	141 140 140 140 140 140 140 140 140	THE PRESENTATION OF	2	11 18 182222 11 18 11 11 11 11 11 11 11	9.71	8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	** diffe * thin in a little of the same of	0 11 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		014 015 013 013 013 014 015 016 017 017 017 017 017 017 017 017 017 017	0.00 0.00	0.00 0.00 0.00 0.00 0.00 0.00 0.00	100 100		10.0 m (10.0 m) (10.0	200 200 200 200 200 200 200 200 200 200	PAMP PAMP	#33 #33 #34 #35 #35 #35 #35 #35 #35 #35 #35 #35 #35		42 44 43 44 44 44 44 44 44 44 44 44 44 44	4.0 4.0 4.0 4.0 4.0 4.0 4.0 4.0 4.0 4.0	10 4 4 1 3 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	100 147 150	15. 10. 10. 10. 10. 10. 10. 10. 10. 10. 10	 ### ### ### ### ### ### ### ### #### ####	10 10 10 10 10 10 10 10		10	에 가 에 에 에 에 에 에 에 에 에 에 에 에 에 에 에 에 에 에	100 100	### ### ### ### ### ### ### ### ### ##	1	10 10 10 10 10 10 10 10	100 100		0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	100 110 110 110 110 110 110 110 110 110	Cast	100 mm of 100 mm	110 110 110 110 110 110 110 110 110 110	



GENERAL STATISTICS.

POPULATION.—As estimated by the Registrar General at the middle of the year 1936—290,400.

(Col. 8). Population per acre (exclusive of River Areas). 30 Gross (Col. 3 & RETURN SHEWING THE ESTIMATED POPULATION OF THE DIFFERENT WARDS IN THE CITY, 4) 6) 26 Spaces and (exclusive Net Area of Public River Open Areas) acres. 126 2,092 145 232 232 89 1,428 1,428 1,465 172 197 197 1139 1139 1173 577 9,546 Spaces (exclusive of Area of Public River Open Area). acres. 1,102 1,550 Acreage, Population per Acre, etc. and River clusive of Area (in-Present Added Areas Areas). 3,194 181 249 213 250 142 608 1,508 174 174 443 1,585 139 11,401 acres. 143 90 River 305 Areas. 51 acres (5) River area (1-4-1935 1,550 Added Areas. 318 2,638 764 Boundary Extension included) prior to (3) acres. 1,644 1,644 1,644 1,644 1,644 1,190 1,190 1,190 1,190 1,143 1,143 1,149 1 8,458 Area not Population (estimated). 1,825 115,259 111,884 117,721 14,383 13,570 9,317 22,339 25,701 11,107 18,655 11,742 17,742 30,022 290,400 (2) Byker ... St. Lawrence .. St. Anthony's ... Walker Arthur's Hill St. Andrew's St. Thomas'. St. John's ... Stephenson... Benwell Armstrong .. St. Nicholas' esmond Elswick WARD. Fenham .. All Saints' Dene..... Westgate Heaton...

INHABITED HOUSES.—77,758 inhabited houses, which, on the estimated population, shows an average of 3.73 persons per dwelling.

RATEABLE VALUE.—£2,543,478. A penny rate produced £9,919.

SOCIAL CONDITIONS.—The principal **Trades and Occupations** are of a healthy nature, being generally engineering and machine making; conveyance of men, goods, and messages; building and works of construction, e.g., ship building; and connected with ships and boats, sea-faring and harbour work; food, tobacco, drink, and lodging; coal and shale mines; and commercial or business occupations.

The amount of **Public Assistance** granted during the year ended 31st March, 1936, was £389,891 for out-door relief, and £40,070 for indoor maintenance, making a total of £429,961, as compared with £423,607 in the previous year.

The number of registered male unemployed was 22,935 at the beginning of the year, and 17,193 at its close, whilst the figures for females were 2,804 and 2,159 respectively.

The City contains many **Hospitals** and other medical charities, but since wide surrounding districts are also served by them, figures as to patients treated are not of local value. A list of municipal and voluntary hospitals serving the city is given on page 41.

MARRIAGES.—2,474 marriages took place during the year, as compared with 2,467 in 1935, and 2,312 in 1934.

BIRTHS.—4,537, equivalent to a rate of 15.6 per 1,000 population.

DEATHS.—(All causes)—5,148, (53 weeks) equivalent to a gross rate of 17.3 per 1,000 population, and, after deduction of the deaths of 1,421 non-citizens and addition of 151 Newcastle residents who died elsewhere, to a net rate of 13.1 per 1,000 population. In 1935 the death rate was 12.6.

12 Orders for Burial (Newcastle upon Tyne Improvement Act, 1882, Sec. 47) were made, 3 being in respect of bodies lying in inhabited rooms, and 9 being cases from hospital.

Cremation Act, 1902.—The Crematorium, West Road, was opened on the 22nd October, 1934. The following table shows the number of cremations up to the 31st December, 1936:—

	Newcastle Residents.	From Outside of the City.	Total.
*1934	11 84 109	15 104 161	26 188 270
TOTAL	204	280	484

^{* 22}nd Oct.—31st Dec., 1934.

Total Deaths during recent years from certain classes of Disease. Classification in Table III. of Ministry of Health.

	Nervous System.	Circu- latory.	Respira- tory.	Digestive.	Externa Causes.
1912	410	435	603	204	152
1913	457	453	722	332	114
1914	448	505	863	465	142
1915	470	635	873	361	163
1916	477	448	856	281	117
1917	497	478	864	268	135
1918	498	503	957	252	135
1919	439	497	1,040	272	133
1920	384	534	861	275	124
1921 .	347	581	726	297	113
1922	363	689	913	181	92
1923	363	623	623	219	112
1924	376	667	749	206	110
1925	359	696	681	248	131
1926	335	742	596	220	158
1927	328	751	615	204	123
1928	331	796	480	247	153
1929	311	893	577	226	148
1930	256	874	469	227	137
1931	250	991	509	195	158
1932	232	976	413	201	161
1933	237	1,003	362	213	151
1934	266	935	405	215	134
1935	243	1,107	391	223	130
1936	276	1,283	408	266	154

CANCER DEATHS IN AGES (MALE AND FEMALE), 1936.

Site.	Sex.	Under 1 Year.				15—25 Years.			Years and over.	Total.
Cancer of the buccal						1				
cavity and pharynx	M.							15	11	26
curry and panely	F.					****		2	2	4
Cancer of the diges-								- 27		
tive system	M.				****	****	12	51	64	127
	F.			****			7	46	55	108
Cancer of the respi-	**							40	_	40
ratory organs	M.	1111	1101	****	****		4	10	5	19
Concer of the uterms	F.			****		****		5	3 9	8 27
Cancer of the uterus Cancer of other fe-	P.		****		****	4118	4	14	9	21
male genital organs	F.						2	3	4	9
Cancer of the breast	M.									
	F.						6	20	10	36
Cancer of the male genito-urinary										
organs	M.				****		3	4	8	15
Cancer of the skin		1117	2111	****		****		1	3	4
	F.		****		****	****		1	1	2
Cancer of other or	M							7	=	10
unspecified organs	M. F.	****		4449	****			7 8	5	12
	F.	2000		****	****		****	0	0	10
	M.						19	88	96	203
	F.						19	99	92	210
Total						3	38	187	188	413

The average age at death for males was 62 and females 62.

INFANTILE MORTALITY.—408 infants died before completing the first year of life, representing a rate of 90 deaths per 1,000 live births.

ZYMOTIC DEATH RATE.—There were 195 deaths from the "Chief Zymotic Diseases"—smallpox, measles, scarlet fever, diphtheria, whooping cough, fever (typhus, simple continued, and enteric) and diarrhoea (all ages)—equivalent to 0.66 deaths per 1,000 population.

TUBERCULOSIS.—308 persons died from various forms of tuberculosis, 265 being from pulmonary, and 43 from non-pulmonary. The equivalent death rates are: all forms 1.04 pulmonary 0.90, and non-pulmonary 0.14, per 1,000 population.

For comparison of death rates with previous years see large table, page 31A.

For particulars of deaths as to site of disease, age, etc., see table page 37A.

GEOLOGY.—The geological formation of the area consists of heavy clay on the top of hard sandstone, which overlies coal seams.

CLIMATOLOGY.—The following is a brief summary of the main features of the weather in 1936, as recorded on the "Newcastle Chronicle's" instruments:—

The mean barometer reading was 29.7 inches. The mean maximum and minimum temperatures were 74.5 F. and 39.8 F. respectively.

The rainfall for the year was 24.80 inches, 0.80 inches less than that of 1935 (25.60).

The following table shows the frequency of the directions of the wind:—

W.	on	40	days
N.W.	on	117	,,
N.E.	on	54	,,
E.	on	8	,,
S.E.	on	62	,,
S.W.	on	75	**
S.	on	8	
N.	on	2	

Sunshine.

Sunshine records have been available by the courtesy of Professors G. W. Todd and J. A. Hanley, of Armstrong College. The observations are taken at Cockle Park Farm (fifteen miles north of the City, and in a rural area), and at the College itself. During the year 1,155 hours of sunshine were registered in the City, as compared with 1,428 at Cockle Park.

WATER SUPPLY.—The City is served by the Newcastle and Gateshead Water Company with a plentiful supply of pure upland surface water, collected from large catchment areas at Catcleugh, close to the Cheviots, and in lower Northumberland. It is stored in large impounding reservoirs at Catcleugh, Hallington, and Whittle Dene, and passes through filters at Whittle Dene and Throckley. It was found, however, that filtration did not secure the degree of freedom from bacteria which was desirable, and during the last few years it has been supplemented by chlorination, with marked improvement.

In the vast majority of cases the household taps are served directly from the mains without intervening cisterns. A separate trade supply is piped to some of the great riverside works from a point above the filters.

The bacteriological reports upon the water are given on page 105.

SEWERAGE.—There are 373 miles of sewers in the City (before extension) and 16 miles (approximate) in added areas discharging directly into the Tyne, which is tidal, at various points along the 81 miles of river frontage.

CLEANSING AND SCAVENGING .- A weekly collection of refuse is made from 75 per cent. of premises and twice weekly from the remainder.

There are 80,787 dry ashtubs and galvanised iron bins, 127 dry ashpits, and 128 conservancy system closets in the City. Conversions are proceeding steadily and, during 1936, 21 pailclosets, 13 combined privies and ashpits and four "cell" privies were removed and water closets substituted. Seven dry ashpits were also removed and dustbins substituted. One school (in the area added in 1935) is served by "chemical" closets, there being no sewers available. With this exception, all the schools are served by the water-carriage system.

ADOPTIVE AND LOCAL ACTS IN FORCE.

Adopted Acts.—Infectious Disease (Prevention) Act, 1890. Section 4.

Public Health Acts Amendment Act, 1890.—Part III.— Whole of; Part IV.—Whole of.

Public Health Acts Amendment Act, 1907.—Part II.—Sections 20, 22, 23, 26, 27, 28, 29, 30, 31 and 33; Part III.—Sections 34, 35, 36, 37, 38, 43, 45, 48, 49, 50 and 51; Part IV.—Sections 52, 53, 56, 58, 59, 61, 62, 63, 64, 65 and 68; Part X.—Whole of.

Public Health Act, 1925.—Part II., Sections 15, 21, 22, 23, 24, 25, 26, 27, 28, 30, 31, 32, 33 and 35; Part III.—Whole of; Part IV.—Whole of; Part V.—Whole of.

Local Acts.—Newcastle-upon-Tyne Improvement Act, 1837.

,,		"	1846.
,,		,,	1853.
,,		,,	1865.
,,		,,	1870.
,,		,,	1882.
,,		,,	1892.
Newcastle-upon-Tyne	Tramways and In	aprovement Act,	1899.
Newcastle-upon-Tyne	Corporation Act		1911.

Newcastle-upon-Tyne Corporation Act 1926. Newcastle-upon-Tyne Corporation (General Powers) Act, 1935.

			REI	UK	N O	F D	EAT		FROM			CA	USE	3	DUI	KIN(o Ti	HE 5	3 W.	BEK	SE	NDE	D 2	in J.				DEA	THE.							1	TRAN	19-1	37A
CATRE OF DEXTS.	Under Lyear	1 year and under 2.	2 years and under 8.	5 years and under 15.	15 years and 5 under 28.	ander 45.	under 68.	and above.	Torat	Under 1 year	1 year and under 2.	2 years and under 5.	8 years and under 15.	under 25. N	25 years and under 45.	under 65.	and above.	Torat (Ner)	St. Nicholas'.	St. Thomas',	St. John's.	Strphemion.	Armstrong	Westgate.	Arthur's Hill.	Bennett,	Fesham,	All Salety'.	St. Andrew's.	Jesucet.	Dene.	Heaton.	Byker.	St. Lawrence.	St. Anthony's.	Walker,	DEATH DEATH	Outward. F	Tarelitations in the City of "Ecoldents or "New Residents,
I.—EPIDEMIC, ENDEMIC AND INFECTIOUS DISEASES.																																							
Enterio Fever Mendes Scarfer Fever Whooping Cough Diphtheria Influenza Dycentery Erysipolas Policencephalitis Lethangica Gerebro-Spinal Fever Tetamis	8 1 5 4 2 2 6 3	4 5 2 5 =	7 2 2 17 5	16 21 4 4	2 1 2 3 3	1 8 1 3 4 1	9 1 3 1 1	15	3 20 11 9 43 39 4 16 	6 4 3 2 1 5 3	14000-	7 1 12 12 11 11 11	1 15 2 	2 2 3 1	1 7 1 3 4 -	9 3	15	17 8 7 35 38 2 12 10 7	11111-111-1	111111-111111	1 2 5	1 1 1 1 1 1 1 1 1 1		la language	11111-11111	21 19 3 11 11	1 6 4 1	5 1	4 1 2 1 1 1	1 2 4	11-1-111111		222	1 4 22 1 1 22 1 1	1 2 1 3 1 4 1	2 2 2	1	3 3 2 8 2 2 4 9 5	4 12 11 4 41 7 4 10 5 16 6
Tuberculosis of the Respiratory System Tuberculosis of the Central Nervous System Tuberculosis of the Peritoricum and Intestines Tuberculosis of the Peritoricum and Intestines Tuberculosis of the Events and Joints Tuberculosis of Benato Urinary System Tuberculosis of Other Organ Disseminated Tuberculosis TOTAL TUBERCULOSIS	2 2 4	3 8 1	3 8 1	11 15 3 1 	74 1 9 5 1 1 1 - 3	3 6 1 1 1 1 2	1 4	2	286 44 20 3 4 1 1 15	1	1 4 1 6	1 6 1	8 4 1 1 1 15	71 2 2 1 2 78	3 1 1 1 1 1 1	3	1	265 19 10 3 2 1 1 7 308		1 1 1	10	27		7 12	-	16 3 1 2 22	20 2 2 1 1	14 2 2 1	12	2	1	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	6 1	13 4	1 1 1 48	27 3 1 1	- 1	25 10 2	141 39 19 3 4 1 1 13 221
Syphilis. Pyaesia. Septicamia Other Diseases due to Protorox Hydatid Cyats. Other Diseases due to Helminths Other Infectious or Parasitic Diseases	3	1 3		1411111	1 = 1 = 1	5 3 3	22 6 1	1	32 18 4 1 1 6	2 2					4 1 1 1 1 1	20	1	27 5 1 -1 3				11111	1	3 3 1 1 1	4 11111	1 1 1	1			2			5 1	2	2 1	41		5 13 3 1	22 17 4 1
II.—GANCER AND OTHER TUMOURS. Cancer of the Duccal Cavity Cancer of the Depetive Organs and Peritoneum Cancer of the Respiratory Organs. Cancer of the Terns Cancer of the Female Genital Organs Cancer of the Female Genital Organs Cancer of the Breast Cancer of Male Genito Urinary System Cancer of other or unspecified Organs Tumours (not madignant) Tumours of andetermined nature		111111111		111111111111111111111111111111111111111	12	26 6 5 3	171 1 22 18 7 26 8 5	15 67 12 10 5 11 10 4 17 17	36 366 40 33 15 44 21 10 48 23 39						19 4 4 2	97 1 15 14 3 20 4 2	13 (19 8 9 4 10 8 4 13 13	30 235 27 27 27 9 36 15 6 28 9		2 13 1 1 4	1 1 1 1	2 18 1 2 1 1	1 9 1	3 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 1	2 14 3 3 4 2 3	3 23 3 2 2 1 1 2	2 12 1 4	[6 m 1 1 1 1 1 1 1 1 1	7 3 1 2 2	1 12 2 1 1 1 1 3 1 3	3 17 3 5 2	1731221111	2 1 2	1 15 1 5 3 1 3 1	3 21 5 2 2 5 2 3 2 1	11111	6	19 214 28 13 10 17 15 7 30 23 32
III.—RHEUMATISM, DISEASES OF NUTRI- TION AND OF ENDOCRINE GLANDS AND OTHER GENERAL DISEASES.				7	7	10	4		28 10				4	4	9	2 1	6	19 8		1	2		1	-		1	1	1	1		1	1		1	2	5 1		9 2	14 6
Chronic Rheumatism. Diabetes: Rickets. Discases of the Pituitary Ghand Discases of the Thyroid and Parathyroid Ghands Exophthalmic Gottre Discases of the Africails Other General Discases.				1111111	1	6 1 23	2 22 3 2 1	6 22 2	51 1 1 3 8 3 5	1					3 1 2 2	8 21	17 2 1	28 1 1 2 4 2 2		CHILLIE.		1	1	1	i lini	4 1	1	111111	2	-111111	- 111111		111111	2	*	1	THE LITTLE	23 1 4 1 3	33 1 2 7 3 2
IV.—DISEASES OF THE BLOOD AND BLOOD-FORMING ORGANS. Harmorrhagic conditions Anaesis, Caloronis Leukemia, Aleukemia Diseases of the Spiece	3				1 5	1 3 1 2	[77]	5 4	4 17 19 2					1 3		7 4	44	1 12 12 12		1111	12	1	-1	1 1		2 1	- 17	1		111	3			1	1 2 1	-2	100	3 7 7 1	3 12 14 2
V.—CHRONIC POISONING. Alcoholism							3		3							3		3		1									1							1			-1
VI.—DISEASES OF THE NERVOUS SYSTEM AND SENSE ORGANS. Encephalitis Meningitis Locomotor Ataxy Other Diseases of the Spinal Cord Cerebral Haemorthage, Apoplexy Hemphers of the Haesee Other John State Other Johnses of the Nervous System Diseases of the Ear and of the Mastoid Sinus	160 2 8		1 1 1 1 2 4	2	3 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 12 2 17 27	1 5 3 46 6 2 2 1 6 4	987	2 10 6 7 159 16 4 3 10 20 22 46	15 16	11111111111	1 2	3 2	3 1 5	1 1 5 1 5 1 9 5 5	1 4 3 47 5 5 3 2 5	2 101 7 9	1 2 5 6 155 14 10 4 17 18 21 23		1 5 2 1 3		4 1 1 1 1 4 1	2 1 1 2 2 2	1 5 1 2		1 2 1 2 1 1 2 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 1 2 17 4 1 2 1 2 1 3 3 3 6		8 1 2		11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 8 3 1 1 1 1 1 1	8 1 2 3	1 12 1 1 1 1 1 1	10 to	2 19 1		1 8 1 1 17 2 1 1 2 5 5 23	2 9 4 4 57 7 4 3 3 3 10 44
VII. DISEASES OF THE CIRCULATORY SYSTEM. Pericanlitis					1 2	4 2 32	1 4 57	1	6 9					3	2 3	1 2	57 416	4 8		1		1	1		2		1 -		72	IST.	1	1	1.5	1 24	1	1111	2 10	2 3	5 6 54
Acute Endocarditis Chronic Endocarditis, Valvular Disease Diseases of the Myocardium Angina Pectoris Other Diseases of the Heart Anseuryan Atterio Scienosis Gangariu Diseases of the Arterio Diseases of the Lymphatic System Abnormalities of Blood Pressure		1	1		2 10	14 5 4 6 2 1	131 82 11 12 58	56 419 66 20 3 296 12	566 153 35 23 356 10 2 9	- 1		111111111	1	8	3 2 3	8	18 1 286 6	28 13 340 8 2 7 2	1 5	17 8 1	1	43	31 6 4 19	1	13 36 7 5 12 1 1	2 2	6 22	11	1 10	18 8 22	29 14 1 22 1 1 1	1	2	2		23	17 4 1	20 33 23 8 10 27 11 1 2 6	269 48 13 18 128 16 1
VIII. DISEASES OF THE RESPIRATORY SYSTEM. Diseases of the Nasal Fosse and Annexa					- 1	1			1						1		-	7					11	5.1.	17	1	1		-		13	1.54	1 39	1 24	10	13	11 14	1	1 1
Diseases of the Laryns Brouchies Brencho-Pieumenia Lobar Preumenia Pieumenia Pieumenia Congestion and Haemoerhagic Infanct of Lung Authura Pulmonary Emphysema Other Diseases of the Rospitzatory System.		7 8 1	2 3		1 3 3 4 8 4 8 4 2	10	3	17 6 2 3	179 109 21 16 8	41 000	1	3 3	22 1 1 1 1 1	2 9 9	1822112	40	16 5 1 3 7	110 144 96 16 9 5 18 3 6	111111	1 3	12	8 10 5 2 3 1 1	81	11 10 1 1 1	4 5 6 3	3 4 1	7 1	6 39 11 00 42 11 11 11 11		1000	5 1	22 11	7 3 1 1 2	7 10 3	10 10 5 1	The same of		8 7 1 3	24 122 75 9 9 2 2 1 4
Carried forward.	16	3 6	7 7	8 10	9 181	420	1068	8 149	9 3615	111	3 4	9 51	0 50	133	3 1319	874	138	2984	1	4 100	129	193	142	199	160 1	01 2	52 25	1 14	13 10	5 113	2 167	147	130	161	215	258	102	733	1893

		RE	IUK	N U	F D	EA	HS	FR	OM "			USE	5	DU	RIN	5 T	HE.	53 W	EEN	SE	NDE	D 2	ND J	ANI	7777	10000	2000	ERT D		20000	-	-	-	-	-		e Tax		374
CAUSE OF DEATH.	oder 1 year	year and oder 2.	years and sdor 8.	years and oder 15.	Syvars and South of the Care and the Care an	Synamand Pader 45.	years and oter 65.	years d above.	Total Gross).	der 1 year	year and	years and	years and oler 18.	years and N	vestra and "	years and ider 68.	years d above.	Torse. (Next).	Nicholas'.	Thomas'.	John's.	ophenica.	metrong.	rwick		thur's Hill.	DWGB.	Saints.	Andread	Andrew A.	100	aton.	Seer.	Lawrence.	Anthony's.	aller.	Das	SEC.	Parities in the of Engineer.
Brought forward	163	- 9	78	109	181	420	1998	1499	3615	113	49	50	37	133	319	\$7.6	1389	2984	24	106	128	93 1	42 1	99 1	(i) (ii)	1 25	2 25	1 143	105	1112	167	9H	130	161	215	258	102	One	A Car
IXDISEASES OF THE DIGESTIVE																																							
SYSTEM. Discuss of the Buccal Cavity, Pharynx, etc	2	1	1	1	1	1 27	*	2 13	17							7	.1.	8				1		1		2	1		1			1							13
Other Diseases of the Stomach Diarrhoes and Enteritis (under 2 years) Diarrhoes and Enteritis (over 2 years)	133	11	-	3	-	1	49 5	12	90 11 144 19	3 94	11	1 1 19	1130		9 7	18	6	33 8 103		7	6		1 11	1 5	8	2 1	3 3	0 0		3	2	100	2	-+	9	9		57 3 39	5
Ciceration of the Intestines	-		3	10	5 3	10 2	1 17 34	1 10 28	4 55 78	119		I.E.	1 2	27	222	8 0	3 9	18 3 17 22		22.1		1 0	1	2			2 1	2 2		-	1	1	100	-	-	1 2 2	2	3 38	4 3 51 76 12
Other Diseases of the Intestines						1 2	8 7 6	1 1	9 9						1	4 5 4	1 1	7 7 6		-	-	3	1		-				-	1	1	1	-	2	1 1	1		58 5 77 7	12 8
Boliary Calcoli Other Diseases of the Gall Bladder and Ducts Diseases of the Paucress Peritonitis without stated cause				-		3 1	21 8 4 4	21 8 1	45 19 6 7						3	5 1 1	7 6	15 13 1 3			1	2	11.00	1 2	1111	1	1	1	1	2	1 1 1	1	1	1		1		30 7 5 4	39 14 6
X. NON-VENEREAL DISEASES OF THE GENITO-URINARY SYSTEM AND ANNEXA.																																						1	
Acute Nephritis Chronic Nephritis Nephritis (not stated to be acute or chronic)	1		1	2	5 2	16	1 48 4	5 57 10	14 127 17				1	3 7	8	31 4	4 48 10	8 90 17	1	2 7	1	3	1 5	7	8 3	1	1 -	1	17.	9	5	5	1	3	1 5	7	2	8 39	72
Other Diseases of the Kidney and Annexa Calculi of the Urinary Passages. Cystitis	2 -				2	2	9 2 3	5 1 7	20 3 13	2				1		1 1	3	8 1 9				11	-		1	3	1		2	2	1	1	1	-	1	1	-	12 2	72 3 16 3
Diseases of the Urethra, Urinary Abocess, etc. Diseases of the Prostate Diseases of the Male Genital Organs Diseases of the Female Genital Organs	1				1111	1 12	7 14 5	45	9 59 1 8						-	3 2	21	24		1	7		2		1 .	3	2 4		1	3	2	1	3	1111		1	1 3	4 6 36 1	7 8 50 1
XI. DISEASES OF PREGNANCY, CHILD- BIRTH AND THE PUERPERAL STATE.																		4													1							5	8
Post-abortive Sepsis. Abortion, not returned as Septic					1	11 8			12 8					1	3			4.3				1		1	-						1			1	7	1		8 5	11
Ectopic Gestation Other Accidents of Pregnancy Pureperal Hamorrhage Discreased Sensis					1 6	1 9			1 10					-	1			1																				2 9	7 3 9
Puerperal Sepsis Puerperal Albuminuria and Convulsions Other Toxemias of Pregnancy Other Accidents of Childberth					3 2	25 13 3 4	1		31 17 3 6					1	6			7		1		2		1	1	1					14		1			1	1	25 10 3	31 16 3
Other or unspecified conditions of the Puerperal state					2	4			6					2	1			3					1								1					1		3	6
XII.—DISEASES OF THE SKIN AND CELLULAR TISSUE. Carbuacie, Boil						-14	2																																
Cellulitis, Acute Alucrus Other Diseases of the Skin and Annexa	3	1	i		1	1	6	1	12 2	1		1		1		1	1	4. 2				1		1 .			-	7	1	1		1		-		-	,1	4 8 1	7 12 2
XIII. DISEASES OF THE BONES AND ORGANS OF LOCOMOTION.																																						1	
Acute Infective Osteomyelitis and Periostitis Other Diseases of the Bones . Diseases of the Joints				141		1 2	1 1	7	1 6 5				- 1		1	1	-	1 4								-		- 2			177	-1			9	191	8	1 5 1	6 3
XIV.—CONGENITAL MALFORMATIONS. Congenital Malformations	50	3			1	2			36	27	1					,		29		1						1.										1			
XV. DISEASES OF EARLY INFANCY.										-0.5											1					4	2	1	1		1	4	1	4	2	5	1 2	8	43
Congruital Debility	36 160								36 160	26 106								26 106		1 4	2 2	1 10	5	1	6 2	1 9	2	2 5	3	2 5	1.3	-3	4	2 7	4	2 9	3 5	0	22
Injury at Birth Atelectasis Icterus Neonatorum	17 6								23 17 6	8 12 6								8 12 6				4	1 1	1	2	1	4	1				1	-		1 3	1		5	20 6 6
Other Diseases peculiar to Early Infancy	10								10	6								6		1	1	1	1 -			2												4	6
Senile Dementia								56	36								3 64	3 64		3	1	2	2	3	7.0	1 5	1 3	3	3	4	7	-	6	4	1 5	-6	3 .	9	
XVII.—DEATHS FROM VIOLENCE.																																							
Suicide by Poison Possonous Gas Hanging or Strangulation						10	1 16 1	1 3 2	2 29 4						10	15	3 2	2 28 3		2	-	3	2	1	2 1	4	1 5	17	2	2			-	1	2	2	1 -	2	3
Drowning						1 1	2 4	6	3 8						-	100		2 4		1	1					1					12		-					1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1
Jumping from a High Place Crushing Accidental Absorption of Poisonous Gas					2	1	1	1	1 2					2	1	1		3 1 2				1			1	=	1				-		-	1	-			1	3 1
Harns (confligrations excepted)		1	3	1	2	1 2	3	6	19 3 1		1	2	-	1	2 2 1		2	8 2 2		2			2 -			1		1		-	2					1	2 1	3 1 1	18
Injury by Fitnarms Injury by Fall, Crushing, etc. Electricity (Lightning excepted)		2	8	13	17	34	41	29	141		1	3	II	9	19	24	16	83	1	4	1	10			2 2	3	12	3	2	2	2	2	2	7	3	18 1	12 7	0 1	35
Unstated Forms of Accidental Violence Violent Deaths of Unstated Nature Wounds of War	1			2	1	2	- 6	4	14	1			11.	1	1	2 5 2	7	3 9 2		1		1	1 -	-		3		1				1		2		1	1 1 2	6 1	341
XVIII. ILL-DEFINED DISEASES.																		100																					
Sudden Death Cause of Douth unstated or sli-defined	1					1	1	2	3						ī	1	3	5								1	1	1				1		ï		1	1	1	7
Total	622	86	97 1	53 2	146 6	44 2	458 1	842,5	148	408	63	59	78 1	65 4	11 1	0071	627 3	878	16	150 1	82 24	64 19	14 24	1 200	128	329	349	182	140	153	210	182	61 2	13 2	74 3	40 15	51 14	21 30	35

VITAL STATISTICS, YEAR 1936, AND INFECTIOUS DISEASES.

COMPARISON WITH OTHER DISTRICTS.

ATTACK RATE PER 1,000 BIRTHS. (live & still)	Puerperal Fever.	3.27	2.12	1.85	20.00	200	8,93	3.39	200	2.6	6.21	5.66	18.17	1.95	1000	78.7	21 -	4.1	0.19	0.40	20.02	1.60
	Ery- sipelas.	0.40	0.01	0.35	0.09	0.68	0.48	0.42	0.82	0.00	0.32	0.62	0.35	0.46	0.24	0.43	0.60	0.00	0.34	17.0	0.00	0.56
PULATION	Enteric Fever and Con- tinued Fever.	0.06	20.0	0.09	0.01	0.00		0.01	0.02	0.00	0.02	0.03	0.04	0.02	0.04	00.00	0.01	0.01	0.21	0.00	0.01	0.05
1,000 Pc	Diph- theria.	1.39	2.30	50.00	202	98 86	1.56	2.86	2.51	1.01	1.41	1.10	1.52	1.07	0.97	1.07	977	0.74	0.09	1.00	1.02	3.57
CATE PER	Scarlet Fever.	0101 e	3.22	20.00	10.0	4.85	3.12	5.64	1.80	10.2	2.19	3.84	1.60	2.00	010	2000	2.99	0000	1.00	11.0	*****	5.26
ATTACK RATE PER 1,000 POPULATION	Typhus,	1.1	:	:	:				0.00		0.0			0.00					0.00	0.00	:	: :
	Small- pox.	0.0		:					0.00		0.0			00.00			:		0.00	00.00		
	Tubercu- losis (all forms) Death Rate.	++ 2	1.04	0.96	0.00	0.68	1.01	1.14	0.97	0.90	0.85	0.78	1.05	0.81	0.81	0770	1.12	0.00	0.00	1.07	12.0	0.77
Death Rate per 1,000 from Enteric	Smallpox, Scarlet Fever, Measles, Whooping Cough, and	0.21	47.0	19.0	0.20	0.34	0.35	0.53	0.49	0.07	0.28	0.21	0.17	0.19	0.11	0.20	0.43	0.16	0.10	20.0	0.10	0.43
	Infantile Mortality Rate.	65 65 65 65 65 65 65 65 65 65 65 65 65 6	200	88	3 22	29	11	81	0.0	0 00	7.4	62	000	40	64	88	200	657	82.	100	20	12
Death	adjusted by Compara- bility Factor.	:: 41	20.	14.0	14.93	13.79	15.39	16.5	19.00	11.80	15.1	12.4	13.3	12.02	19.60	12.00	200	19.61	14.9	11.20	19.01	13.6
	General Death Rate.	123 12		12.7	14.9	19.5	13.5	14.0	10.0	11.6	12.4	11.3	12.6	200	8.11.8	10.01	200	11.8	10.8	10.0	10.01	11.9
.00	Birth Rate.	14.8	2:	18.4	13.4	15.2	14.7	15.0	15.0	14.5	16.8	15.8	1.0.1	77.4	900	0.61	16.0	18.1	10.6	18.5	15.9	17.3
	DISTRICT.	England and Wales 122 Great Towns (including London) NEWCASTLE UPON TYNE		Hall	Bradford	Sheffield	Manchester	Salford	Liverpool	Leicester	Stoke-on-Trent	Birmingham	Cardiff	Dristol	London (County)	Cottoband	Catterness Shields	Tenemonth		Middleshrough	County of Northumberland	*County of Durham

† Not available.

* Administrative County.

Vital Statistics of Whole District during 1936 and previous Years.

			BIRTHS.		TOTAL I REGISTE THE DE	RED IN		TERABLE THS.	NET		BELONGIN	G TC
YEAR.	Population estimated to Middle		Ne	t.			of Non- resi-	of Resi-	Under of A	ige.	At all	Age
	of each Year.	Uncor- rected Number	Number	Rate.	Number	Rate.	dents regis- tered in the District	dents not reg- istered in the District	Number	Rate per 1,000 Nett Births.	Number	Ra
_ 1	2	3	4	5	6	7	8	9	- 10	11	12	1
1011	007.001	7.089	7,082	26.5	4.667	17.5	448	165	973	137	4.384	1
1911 1912	267,261	7,089	7,194	26.7	4,221	15.7	529	146	727	101	3,838	1
1913	269,193 271,295	7,480	7,154	27.5	4,611	17.0	560	141	908	122	4,192	1
1914	271,523	7,564	7,538	27.8	5,069	18.7	546	138	1,029	137	4,660	1
1914	278,107	7,575	7,545	27.8	5,257	18.9	693	207	1,007	133	4,771	1
1916	278,107	7,332	7,343	26.2	4,875	17.5	680	232	899	123	4,427	1
1917	278,107	6,548	6,495	23.4	4,646	16.7	718	246	732	113	4.174	1
1918	278,107	6,555	6,468	23.3	5,380	19.3	872	308	692	107	4.816	1
1919	275,099	6,793	6,674	23.3	5,358	19.5	737	234	806	120	4,855	1
1920	286.061	8,433	8,070	28.0	4,609	16.1	779	195	817	101	4.025	1
1921	278,400	7,720	7,284	26.2	4,602	16.5	817	142	699	96	3,927	1
1922	281,600	7,432	6,987	24.8	4,698	16.7	831	145	646	92	4.012	1
1923	283,800	6,961	6,367	22.4	4,298	15.1	789	150	623	98	3,659	1
1924	285,900	7.029	6,335	22.2	4,607	16.1	929	172	632	100	3,850	1
1925	286,300	7.031	6,215	21.6	4,732	16.5	989	165	550	88	3,908	î
1926	284,700	6,728	6,007	21.0	4,460	15.7	979	161	530	88	3,642	i
1927	288,500	6,215	5,395	18.7	4,468	15.5	1,058	178	474	88	3,588	1
1928	281,500	6,360	5,429	19.2*	4,683	16.6	1,178	179	447	82	3,684	1
1929	283,400	6.120	5.126	18.1	5.040	17.8	1,313	172	438	85	3,899	î
1930	283,400	6,190	5,223	18.4	4,665	16.5	1,232	133	384	74	3,566	î
1931	283,600	6,058	5,056	17.8	4,911	17.3	1,251	145	467	92	3,805	1
1932	285,100	6,006	4,883	17.1	4,579	16.0	1,174	134	370	76	3,539	1
1933	286,500	5,770	4,712	16.4	4,695	16.4	1,182	127	359	76	3,640	î
1934	287,050	5,848	4.695	16.4	4,823	16.8	1,322	145	389	83	3,646	1
1935	292,700†	5,895	4,666	16.0	5,040	17.3	1,489	121	400	86	3,672	1
1936	290,400	5,709	4,537	15.6	5,148	17.4	1,421	151	408	90	3,878	1

^{*} Calculated on a population of 282,200.

Corrected Death Rates in different Wards, 1936.

St. Nicholas'.	St. Thomas'.	St. John's.	Stephenson.	Armstrong.	Elswick.	Westgate.	Arthur's Hill.	Benwell,	Fenham,	All Saints'.	St. Andrew's.	Jesmond.	Dene.	Heaton.	Byker.	St. Lawrence,	St. Anthony's.	Walker,
8.8	9.8	12.8	14.9	13.5	18.2	14.7	13.7	14.7	13.6	13.2	13.3	13.8	11.3	13.7	11.1	12.0	17.6	11.3

[†] Rates calculated on a population of 291,025.

CAUSES OF DEATH AT DIFFERENT PERIODS OF LIFE FOR 1936.

(REGISTRAR GENERAL'S RETURN).

Causes of Death.	Sex	All Ages	0-	1-	2-	5-	15-	25-	35-	45-	55-	65-	75
all Causes	M. F.	2013 1813	238 167	24 39	29 29	37 43	75 85	81 90	128 105	235 151		465 417	308 415
1—Typhoid and para- typhoid fevers	M. F.	1					1	1			****	****	
2—Measles	M. F.	7 11	4 2	1 3	2 6								
3—Scarlet Fever	M. F.	6		5	2	1				****			
4—Whooping Cough	M. F.	3 5	2 3	1	1								
5—Diphtheria	M. F.	17 18	3	3 2	5 7	6 9					****		
6—Influenza	M. F.	20 18	2	1		1	1 1	2	1 4	5	2 2	3 4	4 5
7—Encephalitis lethargica	M. F.	4 8					2	4	1	2	1 1		
8—Cerebro-spinal fever	M. F.	2 3	1 1			1 2							
9—Tuberculosis of respiratory system	M. F.	153 112	1	1	1	3 6	27 43	25 24	33 16	31 11	24 6	8 4	,
0—Other tuberculous diseases	M. F.	15 24		3 2	2 5	4 3	4 2	2 4	3	1	3	1	
11—Syphilis	M. F.		3			1	1	ï	2 1	4 4	5 3		
12—General paralysis of the insane, tabes dorsalis	M. F.								4	6	. 0		
13—Cancer, malignant disease	M F.						1	5	15 12			69 71	2 2
14—Diabetes	M F.							2	9	• 4		3 13	
15—Cerebral hæmorrhage, etc.	M F.							1 1	2		31 18	36 31	1 2
16—Heart disease	M F.				i	2	6					175 156	
17—Aneurysm	M F.					1	1				5 1	3	
18—Other circulatory diseases	M F			ļ			1	1			20 21	47 47	

40

Causes of Death at different periods of life for 1936—continued.

		A 22		10 33									
Causes of Death.	Sex	All Ages	0-	1-	2-	5-	15-	25-	35-	45-	55-	65-	75
19—Bronchitis	M. F.	64 46	1	2	1	2	1 1			12 1	13 5	19 12	16 23
20—Pneumonia (all forms)	M. F.	154 93	28 24	7 10	8 3	1 3	8 4	7 4	12 4	28 3	30 11	18 16	7
21—Other respiratory diseases	M. F.	15 8		1	1	1		1	2 2	2 2	3	3	1
22—Peptic ulcer	M. F.	29 7				****	****	3	7	6 3	7	5 2	1
23—Diarrhœa, etc.	M. F.	59 56	48 38	3 8	1 2	2	****	1	2	1	 1	1	1 3
24—Appendicitis	M. F.	12 4				2	1		1	2 2	5	1 1	1
25—Cirrhosis of liver	M. F.	9							1	3	3	1	1
26—Other diseases of liver, etc.	M. F.	7 21							1	1 3	4 6	1 7	1
27—Other digestive diseases	M. F.	25 27	3 2	1	****	1	1	2	1 2	6 3	4 8	6 5	3
28—Acute and chronic nephritis	M. F.	56 63				1	3 3	2 3	5	6 7	12 15	16 11	11 24
29—Puerperal sepsis	F.	11			****		3	7	1				
30—Other puerperal causes	F.	17		****			5	5	7				***
31—Congenital debility, premature birth, malformation, etc.	M. F.	115 73	115 72					1					
32—Senility	M. F.	21 41										5 7	10
33—Suicide	M. F.	28 14					2	6	2 3	6 4	8 5	2 2	
34—Other violence	M. F.	81 37	2	2	4	10 4	10	9 3	10 5	16 7	6 4	8 4	
35—Other defined diseases	M. F.	153 138	25 23	2 4	4	6	9	7 11	8 19	12 19	24 14	34 22	2
36—Causes ill-defined, or unknown	M. F.	3 1									1	1	

Resident Population 290,400.

UNDER 1 YEAR.

	L	egitimate.	Illegitimate.
M.		232	6
F.		157	10

HOSPITALS.

Name.	Purpose.	No. of Beds.	For Newcastle Cases.	For Cases from outside City.
35				
MUNICIPAL. City Hospital for	Infectious			
Infectious Diseases	Diseases,	200		
Smallpox and Isolation	Tuberculosis Smallpox and	338	338	
Hospitals	Isolation	172	172	
Newcastle General Hospital	Medical, Surgical and Maternity	Men 271 Women 319		
Hospital		Children 205)	****
Barrasford Sanatorium, Barrasford	Tuberculosis	95	75	20
Newcastle Mental	Mental	1,067	1,067	
Hospital, Gosforth Shotley Bridge Colony,	Mental Defec-	473	473	
Shotley Bridge Colony,	tives	473	4/3	
St. Mary Magdalene	Chronic Sick	96	96	
Hospital, Newcastle				
VOLUNTARY.	Conoral Medical	708	208	500
Royal Victoria In- firmary, Newcastle	General, Medical and Surgical,	708	200	300
	Venereal Discourse etc			
Do.	Diseases, etc. Convalescents	50		
Fleming Memorial	Children	88	30	58
Hospital, Newcastle Princess Mary Maternity	Maternity	73	30	43
Hospital, Newcastle Eye Infirmary,	Eyes	35	11	24
Newcastle	Lyes	33	11	24
Throat, Nose and Ear	Throat, Nose and	35	****	
Hospital, Newcastle Hospital for Diseases of	Ear Diseases of the	Out patients	only.	
the Chest, Newcastle	Chest	20		
Catherine House, Newcastle	Maternity	20		
Babies' Hospital and	Children	26	13	13
Mothercraft Centre, Newcastle				
Stannington Sana-	Tuberculosis	310	30	280
torium, Stannington Dental Hospital,	(Children) Dental	Out patients	only.	
Newcastle				
Walker Accident Hospital	Shipyard Accidents	21	21	
Newcastle Dispensary	General, Medical			
Hospital for Diseases of the Skin	Skin Diseases	6		
Hospital for Women	Women			07
Sanderson's Home for Crippled Children,	Children	134	67	67
Gosforth				



REPORT OF THE MATERNITY AND CHILD WELFARE MEDICAL OFFICER.

II.—THE CHILD.

INFANTILE MORTALITY, MATERNITY AND CHILD WELFARE, NURSING HOMES.

BRAYJEN GILLS SEE THE STANK

CHIE CHIE

ONIO WELLARD HURSING HORES

INFANTILE MORTALITY.

SUMMARY OF BIRTHS AND DEATHS, 1936.

	LF	GITIMA	TE.	ILLE	Cuand		
Their same learn bear	M.	F.	Total.	M.	F.	Total.	Grand Total.
Total Births in the year	2,838	2,617	5,455	126	128	254	5,709
Net ,, ,, ,, ,,	2,257	2,110	4,367	86	84	170	4,537
Net Deaths under 1 year	231	156	387	7	14	21	408
Death Rate per 1,000 births	102	74	89	80	167	124	90

BIRTHS AND DEATHS (NET), 1936.

BIF	CTHS AND	DEATHS (N.	ET), 1936.	
Ward.	Births.	Deaths under 1 year of age.	Children under I year of age— Death rate per 1,000 births.	Birth rate per 1,000 population.
St. Nicholas'	2	_	-	1.1
St. Thomas'	153	14	92	10.0
St. John's	150	18	120	12.6
Stephenson	304	35	115	17.2
Armstrong	290	31	107	20.2
Elswick	195	19	97	14.7
Westgate	205	21	102	15.1
Arthur's Hill	91	7	77	9.8
Benwell	523	45	86	23.4
Fenham	462	39	84	18.0
All Saints'	210	22	105	15.2
St. Andrew's	137	15	109	13.0
Jesmond	94	9	96	8.5
Dene	201	8	40	10.8
Heaton	117	- 11	94	8.8
Byker	211	13	62	14.6
St. Lawrence	293	25	85	16.5
St. Anthony's	414	45	109	26.6
Walker	485	31	64	16.2
CITY	4,537	408	90	15.6

All births and deaths occurring in Public Institutions have been allotted to the Wards to which they properly belong.

ANALYSIS OF INFANTILE MORTALITY.

	1927	1928	1929	1930	1931	1932	1933	1934	1935	1936
Death-rate of Infants under 1 year per 1,000 births	88	82	85	74	92	76	76	83	86	90
Death-rate of Infants under 3 months per 1,000 births	55.6	50.8	52.5	46.7	48.1	51.2	45.0	57.2	57.4	58.4
Death-rate of Infants from Premature Birth, per 1,000 births	22.6	20.6	24.5	17.8	20.2	20.7	20.4	21.5	21.9	23.4
Death-rate of Infants under 1 year per 1,000 births from Premature Birth, plus all Congenital Causes*	38.6	35.4	38.8	33.7	34.2	37.3	36.7	38.6	43,3	42.5
Death-rate of Infants under 1 year per 1,000 births, from Diarrhoea and all other Digestive Diseases†	9.3	13.4	15.0	11.3	12.5	9.2	12.9	13.4	13.7	22.2
Death-rate of Infants under 1 year per 1,000 births, from Infantile Atrophy, Debility and Marasmus Death-rate of Infants under	6.5	4.4	3.7	4.6	2.2	4.9	4.7	4.5	5.6	5.7
1 year per 1,000 births, from Measles	0.6	2.2	3.7	0.5	5.7	0.8	1.5	1.9	0.6	1.3
Death-rate of Infants under 1 year per 1,000 births, from Whooping Cough	1.3	3.9	1.4	2.5	2.6	2.5	2.3	1.9	2.4	0.9
Death-rate of Infants under 1 year per 1,000 births, from Respiratory Diseases	27.1	16.6	16.4	16.8	24.7	16.0	12.9	15.5	15.9	12.1
Death-rate of Infants under 1 year per 1,000 births, from Tuberculosis (all forms)	2.4	1.3	1.0	1.1	2.0	0.8	1.3	0.9	0,9	0.2

For particulars of deaths, as to causes, etc., see Table on page 46A.

^{* &}quot;All Congenital Causes" includes Syphilis, Congenital Defects and Diseases of Early Infancy.

^{† &}quot;Diarrhoea and all other Digestive Diseases" includes Diarrhoea, Dysentery, Epidemic or Zymotic Enteritis, Rickets, Diseases of the Stomach, Enteritis, Obstruction of Intestine, Peritonitis and other Diseases of the Digestive System.

									1	AGE I	PERIO	DS.									.g:
					GROS	s.				-		Ni	er (a	fter a	llowir	g for	tran	sfers)			
Cause of Death.	Under I Week.	1-2 Weeks.	2-3 Weeks.	3-4 Weeks.	Total under I Month.	1-3 Months.	3-6 Months.	6-9 Months.	9-12 Months.	Total under I Year of Age.	Under 1 Week.	1-2 Weeks.	2-3 Weeks.	3-4 Weeks.	Total under 1 Month.	1-3 Months.	3-6 Months.	6-9 Months.	9-12 Months.	Total under 1 Year of Age.	Deaths in Institutions the City of " Residents or " Non-Residents."
EPIDEMIC, ENDEMIC AND INFECTIOUS DISEASES.																					
Measles Scarlet Fever Whooping Cough Diphtheria Influenza Dysentery Erysipelas Cerebro-Spinal Fever						2 1 1 1 1	2 	3 1 1 1 2	1 3 1	8 1 5 4 2 2 6 3						2 1 1 1 1	2 3 2 2	1 1 2	1 3 1	6 4 3 2 1 5 3	7 1 1 4 2 2 6 3
Tuberculosis of the Respiratory System Disseminated Tuberculosis						1	1 1		1	2 2						1				1	2 2
Total Tuberculosis	-					1	2		1	4						1				1	4
Syphilis. Pyæmia Septicæmia Other Infectious or Parasitic Diseases			2		2	2 1	2	3		4 3 3					ï	1	1	2		2 1 2	4 3 3
RHEUMATISM, DISEASES OF NUTRITION AND OF ENDOCRINE GLANDS AND OTHER GENERAL DISEASES. Chronic Rheumatism Rickets Other General Diseases.					1		 1		ï	1 1 1 1							 "ï			 1 1	1 1
DISEASES OF THE BLOOD AND BLOOD- FORMING ORGANS.																1					
DISEASES OF THE NERVOUS SYSTEM AND SENSE ORGANS.	2	1	****	****	3		****			3		1			1			****	****	1	2
Meningitis Infantile Convulsions Diseases of the Ear and Mastoid Sinus Other Diseases of the Nervous System	****	2 1	2		7	3 1 1	1 4 2 	2 5	2	3 16 8 2	3	2	ī		6	3 1 1	4	2 5	1	1 15 6 1	2 2 8 1
Diseases of the Lymphatic System						1			1	2 1						1	- 1			1 1	2
Myocardium	****		****		****		1			1		****							****		****
Bronchitis		1 1	2 3 1 	ī 	3 5 1 	3 8 2 	19 3 2	1 10 2 3	16 1 1	7 58 9 6			1 3 1	ïi	1 4 1 	1 6 1 	13 2 2	1 8 1 3	10 1 1	3 41 6 5	5 38 6 1
Carried forward	6	6	10	1	23	29	49	34	28	163	3	3	7	1	14	21	33	27	18	113	109

Return of Deaths under One Year of Age during the 53 Weeks ended 2nd January, 1937.

										Age	PERI	_									I gr
					GR	oss.		-		-	-	1	NET (after	allow	ving f	or tr	ansfer	rs).	-	lons
Cause of Death.	Under 1 Week	1-2 Weeks.	2-3 Weeks.	3-4 Weeks.	Total under I Month.	1-3 Months.	3-6 Months.	6-9 Months.	9-12 Months.	Total under 1	Under 1 Week	1-2 Weeks.	2-3 Weeks.	3-4 Weeks.	Total under 1 Month.	1-3 Months.	3-6 Months.	6-9 Months.	9-12 Months.	Total under 1	Deaths in Institutions the City of "Residents or "Non-Residents"
Brought forward	6	6	10	1	23	29	49	34	28	163	3	3	7	1	14	21	33	27	18	113	109
DISEASES OF THE DIGESTIVE SYSTEM. Diseases of the Buccal Cavity, Pharynx, etc. Other Diseases of the Stomach. Diarrhoea and Enteritis. Hernia, Intestinal Obstruction NON-VENEREAL DISEASES OF GENITO-	1	7	1 25	10	1 43	1 27 	2 1 36 3	17 5	10	2 3 133 9	1	3	1 15	7	1 26 	1 19	1 26 1	14 1	9	3 94 2	2 114
Other Diseases of the Kidney and Annexa Cystitis Diseases of the Male Genital Organs DISEASES OF SKIN AND CELLULAR		****				1 1	1	1	1	1 2 1 1					****	1		1	1	2 1	1 1 1
TISSUE. Cellulitis, Acute Abscess						1	1		1	3							1	****		1	3
Congenital Malformations. Congenital Malformations	19	3	4	2	28	10	8	3	1	50	12	2	3	1	18	5	4			27	36
Congenital Debility Premature Birth. Injury at Birth Atelectasis Icterus Neonatorum Other Diseases peculiar to Early Infancy	94 16	3 28 5 2 2	10 20 1 1	5 11 1 1 1	24 153 22 17 5 8	6 6 1 1 2	5 1		1	36 160 23 17 6 10	5 59 6 12 1 3	2 18 1 2	7 15 1 1	3 10 1	17 102 8 12 5	4 4	4		1	26 106 8 12 6 6	20 106 20 6 6 6
DEATHS BY VIOLENCE.							1									-	****	****	****	0	0
Violent Deaths of Unstated Nature		****				****	****	1		1	****										
ILL-DEFINED DISEASES.											****	****	****	****	****	****	****	1	****	1	5311
Causes of Death Unstated or Ill-defined	1				1				****	1			****								1
Total	164	56	73	32	325	86	107	61	43	622	102	31	51	24	208	57	70	44	29	408	432

Report of the

Maternity and Child Welfare Medical Officer.

TO THE MEDICAL OFFICER OF HEALTH.

SIR.

I have the honour to submit to you my seventeenth annual report.

Innovations.

The following additional services and altered conditions were provided during the year :—

Ante-Natal Sessions were established in Scotswood, St. Lawrence and Heaton Centres, and sessions for children were added at Scotswood and Cowgate Centres. The Scotswood Centre, previously conducted in St. Margaret's Church Hall, Denton Road, was moved to the new hall in Norland Road, in which improved accommodation is provided.

A new Centre was opened in St. Francis Church Hall, Cragside, for the use of residents in the new estate which is rapidly developing in that area.

Diphtheria Immunisation.

No. of Clinics held:

Diana Street, Monday	42
Scotswood, Wednesday	24
Byker, Wednesday	14
,, Thursday	16
Walker, ,,	
Diana Street, Saturday	36
Byker, ,,	7
Denton School, Wednesday	5
Atkinson Road School, Tuesday	. 5
	177

TOTALS FOR 1936.

Total No. of New Children sent for.	Total Attendances for 1st Treatment.	Total Attendances for 2nd Treatment.	Total Attendances for 3rd Treatment.	Total Atten- dances.
2,398	1,745	1,592	1,466	4,803

Dental, Aural and Nasal Treatment.

Under the arrangements made with the Education Authority 243 nursing or expectant mothers and 359 children were referred for dental treatment. Of these 190 women and 285 children were treated.

Similarly 204 children were sent for aural and nasal treatment and of these 133 were treated.

Whalton Rest Home.

Twenty-two mothers accompanied by 22 children were sent through the Centres for two weeks' holiday at the Rest Home during the year. All were enthusiastic in their praise of the Home and of the pleasure and benefit which they had derived from their holiday.

Orthopædia.

179 new cases were visited during the year. Of these 113 have attended for examination by the Orthopædic Surgeon at the Education School Clinic and a total of 280 examinations and re-examinations were made. Of cases notified to attend for examination four parents refused.

Average number attending twice weekly for treatment —Massage and Exercises	25
Admitted to the W. J. Sanderson Orthopædic Hospital	
School	8
Waiting for admission to the W. J. Sanderson Ortho- pædic Hospital School	4
Refused institutional treatment	4
X-Ray Examinations.	12
Photographic Records	3

40 cases attended other institutions.

SURGICAL APPLIANCES :-

Plaster of Paris splints made	in 3	cases	(9	plasters).
New splints supplied in	28	***		
New boots supplied in	64	,,,		
Alterations to boots in	42	**		
Splints repairs in	17	,,		

During the year 406 children were visited. This includes the 179 new cases; of these

Parents refused any kind of treatment in	14	cases.
	11	
Died		
	73	

leaving 333 on visiting list at end of 1936.

Classification of 179 New Cases Visited.

Congenital Deformities :	
Congenital Club Feet	7
Congenital Flat Feet	3
Congenital Abnormalities—Toes	2
Spastic Paralysis.	
Poliomyelitis	5
Ataxia	
Mentally Defective and M.D. Paralysis	1
Old injury to lower limb	2 2
Disparity in Growth of Legs	2
Torticollis	1
Muscular Dystrophy	1
Tubercular Spine ,	1
Tubercular Knee.	1
Flat Feet	36
Rickets—Bow Legs	42
Knock Knees and Flat Feet	48
General	8
Talipes Calcaneus	1
Metatarsal Varus	1
Renal Dwarfism	1
Pecularity of Cait	1
Pecularity of Gait	1
Periostitis	î
Œdema of Feet	1
Total	179

Home Helps.

During the year 16 mothers were supplied with the services of a Home Help. Of these six contributed towards the cost.

Maternity Hospital.

Free beds in the Princess Mary Maternity Hospital were placed at the disposal of 119 women and 41 others were admitted to the Newcastle General Hospital for confinement during the year. In addition free outdoor attendance on the various districts by the Professional Staff of the Hospital was provided for 150 women. The necessary orders were given by the Medical Officers at the Centres.

Ante-Natal Centres.

During the year 1,847 women attended the Municipal Clinics, this being a decrease of 189 over the corresponding figure for the previous year.

The following table shows the attendances at the ante- and post-natal clinics:—

CENTRE.	ANTE-	NATAL	Post-Natal.			
CENTRE.	Attendance.	Individuals.	Attendance.	Individuals		
Benwell	1,089	290	121	39		
Byker	1,679	388	14	4		
Fenham	569	135	1	1		
Diana Street	807	212	2	2		
Elswick	901	274	2	2		
Heaton	75	20	1	1		
Scotswood	305	68	33	22		
St. Lawrence	200	40	2	2		
Walker	585	231				
Wharncliffe St	995	189	4	4		
	7,205	1.847	180	77		

Women Attending Ante-Natal Centres.

The following details refer to the confinements of 1,709 expectant mothers who attended the municipal ante-natal centres during 1936, and whose children were born during that year.

Mothers were sent to the ante-natal centres by the following :-

	Cases.	Percentage.
Doctors		14.2
Midwives	284	16.6
Health Visitors on Districts	123	7.2
Welfare Centres and Voluntarily	1,059	62.0
	1,709	
	Management .	

The result of the subsequent confinements were :-

Tune of	Number	Resulting in					
Type of Confinement.	of Cases.	Living Children.	Still-born Children.	Sets of Twins.			
Normal	1,338	1,301	27	10			
Instrumenta!	215	170	38	7			
Cæsarian Section	16	14	2				
Induction	3	3	100				
Abortion	19			****			
	1,591	1,488	67	17			
Not Pregnant	64	****	****	****			
Left City	54						
Total	1.709						

Abnormalities were found in 101 or 5.9 per cent. of the cases, and the ultimate results were as follows:—

Abnormality.	No.	Normal Confinements.		Instrumental Confinements.			Cæsarian Sections.		Induced Labour.				
		No.	Living Children.	Still-born Children,	No.	Living Children.	Still-born Children.	No.	Living Children.	Still-born Children.	No.	Living Children.	Still-born Children.
Breach Presentation	59	37	34	3	20	13	7	2	2				
Deformed Pelvis	8	2	2					6	6		·		
Albuminuria	21	17	15	2	3	3					1	1	
A.P. & P.P. Hæmor- rhage	13	6	5	1	7	2 *	5						

7 mothers subsequently died: --Sepsis, 3; Cardiac Disease, 1; Toxæmia, 2; Pneumonia, 1.

Notices for medical help sent to Local Authority by the midwives:—

FOR THE MOTHER.		During Puerperium-	
During Pregnancy— Ante Partum Hæmorrhage Abortions	21 3 7	Rise of Temperature Eclampsia Undefined Illness of Mother	14 1 21 36
Albuminuria & Puffiness of hands and feet	14	Total calls for mother	338
During Labour— Uterine Inertia	79 37 3 9 129 257	FOR CHILD. Prematurity Discharging Eyes Congenital Defects Illness of Baby Still-births Rashes Total calls for mother and child	26 36 6 25 3 5 101 439

In 24.3 per cent. of the midwives' cases the services of a doctor were requisitioned.

Claims from doctors for fees in respect of calls from midwives :-

	Cases.
For forceps delivery	102
For post partum hæmorrhage	6
For ante partum hæmorrhage	10
For illness of mother	36
For illness of child	56
For premature birth	9
For discharging eyes	17
Other	104
Specialists called in	6
Total cases	346

20 claims for payment of midwives' fees were received.

Maternal Mortality.

4,537 live and 190 still-born births belonging to the City occurred during the year. 28 women died as a result of childbirth, a mortality rate of 5.92 per 1,000 live and still births as compared with 5.13 in the previous year.

Causes.	1936	1935	1934	1933	1932
Abortions (Septic)	4		1	1	1
Abortions (Not Septic)	3	1	1	4	1
Accidents of Pregnancy	1		1		
Puerperal Hæmorrhage	1	3	3	4	3
Other Accidents of Childbirth	1	2	2	2	4
Puerperal Fever	6	12	7	7	9
Other Toxæmias of Pregnancy	1	1	1	2	2
Puerperal Phlegmasia		4	1		1
Puerperal Insanity					
Puerperal Disease of Breast		****		200	
Ectopic Gestation	1		1		1
puerperal state	3		2	2	
Convulsions	7	2	6		
	28	25	26	22	22

Puerperal Septicæmia and Puerperal Pyrexia.

One hundred and thirty cases were notified during the year—35 puerperal fever and 95 pyrexia. Details of these are embodied in the following table:—

	Total Cases Notified.	Newcastle Cases.	Extra Mural Cases.	Admitted to Hospital.	Total Deaths.	
) (Newcastle	10
Puerperal Septicæmia	35	9	26	100	Extra Mural	33
				123	Newcastle	0
Puerperal Pyrexia	95	19	76		Extra Mural	0

Of the 28 City cases 26 were visited and the attendants at the confinements are indicated in the following table:—

	Puerperal Septicæmia.	
Doctors		
Doctors and Midwives	1	3
Midwives	3	2
Princess Mary Maternity Hospital Staff		10
Newcastle General Hospital		2
Gables Maternity Home		
		177
	9	17
	-	Minne

Consultants' Services.

The services of Obstetrical Specialists were asked for and provided on six occasions and the Midwifery Emergency Service was sent to two cases during the year.

Midwives Acts.

Fifty-three midwives notified their intention to practise midwifery in the City. Two of these were registered as being in practice prior to the 1902 Act, and the remaining fifty-one qualified by examination and possessed the Certificate of the Central Midwives Board. Nine of them only did temporary work at Maternity Homes.

Midwives practising in the City attended approximately two-fifths of the total births taking place in the City. Midwifery bags, appliances, and records kept by practising midwives were examined as a routine, and all cases of ophthalmia neonatorum, puerperal pyrexia and septicæmia were carefully investigated and supervised. Clothing and appliances thought to be infected were disinfected—the owners being suspended from practice so long as was deemed necessary for the safety of others.

For these various purposes 622 visits were made; 112 visits to septic patients, and 302 to cases of ophthalmia neonatorum were also made.

Births attended by Midwives.—1,711 (net) living births (a decrease of 107 on the previous year) and 49 (net) still-births (9 less than in 1935) were attended by midwives during the year. Midwives attended 38.0 per cent. of the net births in the City. In addition midwives attended in the capacity of maternity nurses with doctors in 347 cases, as compared with 298 in 1935.

Ultra-Violet Ray Therapy.

Those children who are brought to the Centres, and who are considered to be in need of artificial sunlight, are referred for such treatment to the Light Department of the Newcastle General Hospital, or to the Brinkburn Street Sun-Ray Clinic. 367 children were so referred during the year, and they received a total of 6,202 treatments.

Health Talks.

A lecture lasting about ten minutes and dealing with an appropriate subject—such as digestive disorders among children in the spring and summer, and the respirating diseases in the autumn and winter—is given by the Centre Health Visitor at every Centre. There is a complete syllabus of the subjects of these talks, and this is closely adhered to. It embraces everything conducive to maintaining good health in mothers and children, and the talks are listened to with interest.

Nursery Schools.

These were conducted by members of the Voluntary Association in Diana Street and Wharncliffe Street Centres, and were much appreciated and enjoyed by the parents and scholars respectively. It is a pleasure to pay public tribute to the ladies who are kind and interested enough to give their time and services every week for this praiseworthy object.

Births.

Of the 4,537 infants born alive in 1936, and belonging to Newcastle residents, 2,343 were boys and 2,194 were girls. Of the former 102 per 1,000, and of the latter 77 per 1,000 died during their first year.

26.1 per cent. of the City's births occurred in institutions, as shown in the following table:—

104
573
136
373
1,186

The Birth Rate in the City for the year 1936 was 15.6 per 1,000 which is 0.4 less than the rate for 1935.

Illegitimate Births.

One hundred and seventy illegitimate children were born—this being 28 less than in the previous year. The death rate among these children is high everywhere, and in Newcastle in 1936 it was 124 per 1,000 compared with 89 per 1,000 legitimate children. Every effort is made to save these children, and when it is possible to get the mothers to bring the children regularly to the Centres the children's lives are practically secure. In all instances the single girl is provided with free milk for her infant when this is suitable and necessary.

Deaths of Children.

	1932	1933	1934	1935	1936
Deaths of children during first week of life	133	126	112	109	102
Deaths of children during first month	175	177	197	117	106
Deaths from prematurity	101	96	101	104	106
Deaths of twins and triplets	32	39	33	29	23

Toddlers.

As in previous years care and attention was bestowed on the children of toddling age, among whom health deteriorates rapidly unless it is closely watched. For the last seventeen years special efforts have been made in Newcastle to encourage mothers to bring toddlers to the Centres, and it is gratifying to report that of the 104,954 attendances at the Centres last year, more than half were made by children of 1–5 years of age.

TODDLERS ATTENDING THE CENTRES.

Year.	Number of Children.
1931	. 4,257
1932	. 4,422
1933	
1934	. 4,198
1935	. 4,288
1936	. 4,315

Ophthalmia Neonatorum.

The number of cases notified was 54, of which 51 were City cases. 50 of these were visited. The confinements were attended by:—

Doctors	8
Midwives	20
Princess Mary Maternity Hospital	16
Doctor and Midwife	3
Newcastle General Hospital	2
Gables Maternity Home	1
	=0
	50

301 calls were made on the 50 visited cases in the City, and the ultimate results were :—

Recovere	ed complet	ely	46
Died			3
Slightly	defective		1
			-
			50
			-

The ophthalmia incidence per 1,000 births for the last five years has been as follows:—

1932	 9.2
1933	 11.0
1934	 11.5
1935	 12.3
1936	 11.2

Children Acts, 1908-1933.

At the beginning of the year there were 118 nursed-out children in the City, and 112 at the close of the year, with foster mothers (private) 51, Institutions 61.

All these children were regularly supervised and kept under observation. 3 children died, all in Hospital.

Municipal Training Course for Health Visitors.

The sixth Training Course for Health Visitors commenced in September, 1935, and ended with the examination at the College of Medicine in April, 1936. Of the ten students enrolled eight qualified in April, one in July and one in December.

The seventh Training Course commenced with 16 students in October, 1936.

Welfare Centres.

The following table shows the geographical position of the Centres in the City, together with details of Centre days :—

Ante-Natal Sessions.	Friday, 2 p.m. Mr. F. E. Stabler. Friday, 2 p.m. Dr. Mabel Campbell. Thursday, 2 p.m. Mr. Harvey Evers. Wednesday, 10 a.m. Mr. Harvey Evers. until 22/7/36. Mr. E. Stabler from 29/7/36. Mr. Harvey Evers. until 28/7/36. Mr. Harvey Evers. Mr. W. Hunter from 4/8/36. Mr. W. Hunter from 67/36. Mr. W. Hunter from 7/8/36. Mr. W. Harvey Evers. Friday, 10 a.m. Mr. Harvey Evers. Friday, 10 a.m. Dr. Mabel Campbell. Friday, 10 a.m. Or Wharncliffe Street (see below). Thursday, 2 p.m. Dr. Mabel Campbell. Thursday, 10 a.m. Wr. F. E. Stabler. Thursday, 10 a.m. Tuesday, 10 a.m. (see Walker above). Tuesday, 10 a.m. (see Walker above). Tuesday, 10 a.m. (see Walker above). Tuesday, 10 a.m. Dr. Mabel Campbell.	
Health Visitor.	Miss Batty Miss Johnson Miss Pritchard Miss Pritchard Miss Hastie Miss Hastie Miss Lewis Miss Carr Miss Scorer Miss Scorer Miss Storel Miss Phillips Miss Shell	Miss Pearse
Medical Officer.	Dr. E. G. Brewis Dr. Dorothy Hopkinson Dr. Anne Fairweather Dr. C. H. Armstrong Dr. C. C. Ungley, p.m. Dr. Elsie Wright, a.m. Dr. Elsie Wright, a.m. Dr. Elsie Wright Dr. A. G. Ogilvie Dr. A. G. Spinks Dr. A. F. G. Spinks Dr. A. F. G. Spinks Dr. A. F. G. Spinks Dr. Anne Fairweather Dr. A. F. G. Spinks Dr. Anne Fairweather Dr. A. F. G. Spinks Dr. A. F. G. Spinks Dr. A. F. G. Spinks Dr. Anne Hackling Dr. A. Gogilvie Dr. Glen Davison Dr. Gertrude Hickling Dr. Gertrude Hickling Dr. A. G. Ogilvie. Dr. Gertrude Hickling Dr. Mabel Campbell Dr. Dorothy Hopkinson, since 2/7/36	Dr. A. F. G. Spinks
Women and Children.	Monday Monday Tuesday Wednesday Tuesday Friday Wednesday Tuesday Wednesday Tuesday Wonday Tuesday Wonday Thursday Thursday Thursday Wednesday Thursday Thursday	Thursday
Address.	Y.W.C.A. Club, Buddle Road Corner of Dalton Street and Shipley Street Princess Mary Maternity Hospital, Jubilee Road Methodist Church Hall, Stamfordham Road 25, Diana Street Malvern Street Church Hall, Grange Road St. Gabriel's Parish Hall, Chillingham Road St. Margaret's Church Hall, Norland Road St. Jude's Parish Hall, Dinsdale Road Benson Memorial Chapel, Ancrum Street Harbottle Street Church Street Conservative Assoc. Hall, Walpole Street, Walkergate 18, Wharncliffe Street	St. Francis Church Hall
Centre.	Benwell Byker City Cowgate Diana Street, Westgate Elswick Heaton Scotswood Scotswood Shieldfield Shieldfield Walker St. Lawrence Walker	Benton (September 1936) St. Francis Church Hall

Na	Ante- Natal.	Post- Natal.	st-	New	New Children.	ren.	Inc	Individuals.	als.	At	Attendances.	ses.	Med	Medical Sessions.	mate.	Pud ng	Individ- uals.
-busttend-	ances.	-bivibnI .slau	Attend- ances.	Under 12 months	Over 12 months	Total.	Under 12 months	Over 12 months	Total.	Under 12 months	Over 12 months	Total.	Number.	Average Attend'ce	itigəllI	Boys.	Girls.
- 1		1	-	26	12	38	24	20	44	306	307	613	16	38.3		21	23
290 1089	6	39	121	263	45	308	404	421	825	4541	7359	11900	192	62.0	13	396	429
388 1679	6	4	14	256	39	295	375	329	704	3424	4635	8059	188	42.9	24	366	338
1		1	1	108	17	125	154	210	364	1678	3183	4861	94	51.7	00	184	180
1		1	1	41	7	48	78	107	185	879	1072	1921	99	29.6	1	84	101
212 807		01	61	227	46	273	354	374	728	4061	4693	8754	190	46.1	6	396	332
274 901		01	01	308	49	357	465	443	806	5132	5635	10767	192	56.1	32	452	456
135 569		-	-	219	17	236	315	234	549	3528	2250	5778	96	60.2	4	271	278
20 75		-	-	183	31	214	268	211	479	2768	2034	4802	96	50.2	00	245	234
68 305		22	33	173	28	201	237	272	509	2275	3424	5699	110	51.8	11	264	245
1		1	:	211	26	237	340	375	715	3775	5856	9631	192	50.2	21	370	345
. !		-	1	64	12	76	66	73	172	878	587	1565	47	31.9	4	98	86
40 200	0	61	61	290	38	328	463	449	912	5007	6255	11262	194	58.1	18	494	418
231 585	10	1	1	215	23	238	361	366	727	3402	5127	8529	192	44.4	22	372	355
-		1		137	30	167	221	226	447	2291	2664	4955	86	50.6	7	233	214
189 995	10	4	4	207	29	236	321	205	526	3437	2391	5828	86	59.5	25	268	258
1847 7205		77	180 2	2928	449	3377	4479	4315	8794	47482	57472	104954	2061	6.06	206	4502	4292

Attendances at Maternity and Child Welfare Centres.

(CHILDREN ONLY.)

YEAR.	No. of Attendances.	No. of Individuals.	Average Attendance per Individual.	Average Attendance at each Session.
1920	22,596	3,751	6.0	44.2
1921	32,538	4,734	6.8	40.7
1922	36,020	4,835	7.4	44.9
1923	42,515	5,153	8.2	46.5
1924	45,766	5,587	8.2	45.5
1925	45,476	5,744	7.9	43.6
1926	50,697	6,467	7.8	46.2
1927	46,672	6,522	7.1	42.4
1928	53,960	6,532	8.3	49.3
1929	52,460	6,574	7.9	48.2
1930	67,626	7,776	8.7	44.2
1931	83,561	8,927	9.4	43.1
1932	100,658	9,251	10.9	51.5
1933	99,103	8,955	11.1	50.9
1934	107,717	8,872	12.1	54.6
1935	104,174	8,952	11.6	52.2
1936	104,954	8,794	11.9	50.9

Dried Milk.

During the year 91,278 lb. cartons of dried milk were given gratis, and vouchers for 15,116 were given for cost price milk, the latter being distributed by the chemists as formerly. Approximately one-third of the children attending the Centres received free milk, and expectant mothers to the number of 469 also received it gratis.

The following table shows the quantity of dried milk distributed each month during the year 1936:—

Month.	FREE.	AT COST PRICE
	lbs.	lbs.
January	6,487	1,170
February	6,755	1,760
March	8,796	1,158
April	6,860	2,368
May	7,120	1.276
June	9,112	959
July	5,784	1,354
August	6,256	828
September	8,638	1,404
October	7,181	1,582
November	9,495	557
December	8,794	700
	91.278	15,116

Children attending Centres	8,794
Children given free milk	3,603
Percentage	41.0
Expectant mothers given milk	469
Free milk given to children (lbs.)	${88,346 \atop 2,932} = 91,278$
Free milk given to expectant mothers (lbs.)	2,932 = 51,276

NOTIFICATION OF BIRTHS ACTS.

Of the 5,709 live, and 371 still-births (gross) which were registered in the City in 1936, 5,395 or 88.7 per cent. were notified as follows:—

Notified by.	Gross Living Births.		Gross Still Births
Medical Practitioners	267	5000	1 i
Medical Practitioners and Midwives Midwives	328 1,750		19 51
Princess Mary Maternity Hospital	2,065		224
Newcastle General Hospital	351 285		30
Parents	5	****	
	5,051		344
	-		-

Still-Births.—Of the total net notifications of births received, 169 were of still-births, which gives a rate of 42.0 per 1,000 of net live and still-births.

Still-births Registered (net)		190
Still-births Notified.		169
Percentage Notified		88.9
Still-births Visited.		172
Duration of Pregnancy.	No.	Percentage to Total.
At or under 7 months	13	7.6
At 7–8 months.	61	35.5
At full time	98	56.9
uggested causes of the still-births :-		
		Cases.
(a) Ill-health of the mother(b) Foetal deformities and malpresentat	ions and	14
uterine inertia		49
etc.	norrhage,	20
(d) Other causes, including albuminuria and	accidents	

Syphilis was returned as the cause of death of two children below the age of 1 year.

Health Visiting.

With the exception of the so-called residential districts such as parts of Jesmond, every district in the City was visited regularly by members of the Health Visiting Staff.

For all purposes the Health Visiting Staff during the year 1936 made a grand total of 87,245 visits.

4,104 births were visited, and 21,094 re-visits were paid, an average of about 5 re-visits per child. These gave a total of 25,198 visits to children under 1 year.

SUMMARY OF VISITS.

	Primary.	Subsequent.	Total.
Births.	4,104	21,094	25,198
Measles ,	3,625	4,251	7,876
Pneumonia	478	552	1,030
Diarrhœa	91	148	239
Children over 1 year			40,669
Hospital Cases			428
Expectant Mothers			1,438
Special Visits			1.093
Visits to Boarded-out or Nursed-out Children Unsuccessful Visits (Outs and			596
Removals)			7,308
Orthopædic Work			1,058
Tuberculosis Contacts			312
			87.245

Infants on Visiting List.

Of 4,285 children born in the City in 1935, 3,358 completed their first year in 1936 and of the remainder:

370 died,

261 left the City,

269 could not be traced.

27 were visited only once.

The following figures are therefore based on the 3,358 who completed the first year, *plus* the 370 who died, making in all a total of 3,728, and of that total 2,447, or 65.6 per cent., attended the Welfare Centres.

Of the number (2,447) attending the Centres 123 died, a rate of 50.3 per 1,000, as compared with 90 per 1,000 for the City.

Illnesses.—Among the children visited 287, or 7.7 per cent., contracted measles; 72, or 1.9 per cent., contracted whooping cough; 134, or 3.6 per cent., contracted diarrhœa; 323, or 8.7 per cent., contracted bronchitis or pneumonia.

Details as to the stated **Feeding** of the 3,728 children under supervision during the year are given in the following table:—

School of the second	FEEDING.						
. Tollands	Breast.		MIXED.		ARTIFICIAL.		
	No.	Per- centage.	No.	Per- centage.	No.	Per- centage.	
At First Visit	3,284	88.1	121	3.2	323	8.7	
above ChildrenAt time of Death	274 143	8.3 4.4	18 30	14.9 24.8	78 197	24.1 61.0	
Surviving Children(3,358) at 9 months	1,278	38.1	280	8.3	1,800	53.6	

Details as to children who should have attained the age of 5 years during 1936:—

Well and attending school		
		Left City or failed to trace
Died in 2nd year	1,113	
Died in 3rd year	33	
Died in 4th year	23	
Died in 5th year	19	
Total surviving	2.837	
Total deaths.		
Total reported on	4,113	

The addresses of 250 children who left the City were sent to the Medical Officers of Health for the districts to which they had gone.

Voluntary Workers.

As in other years the lady members of the Voluntary Association, under the presidency of Mrs. Leach, rendered valuable services, not only at the Centres, but also in the districts.

I am, Sir,

Your obedient servant,

A. F. G. SPINKS, M.D.,

Maternity and Child Welfare Medical Officer.

Maternity and Child Welfare Department,

10, Bigg Market,

Newcastle upon Tyne,

May, 1937.

MATERNITY AND NURSING HOMES.

REPORT OF THE BOARD OF INSPECTION.

1.—Introductory.

The Annual Inspection of the Maternity and Nursing Homes in the City, under the Nursing Homes Registration Act, 1927, has again been carried out by the Board specifically appointed for the purpose.

2.-Record of Inspection.

Nineteen homes in all were inspected, sixteen of these being homes which were inspected and reported upon in considerable detail last year. The three remaining homes comprised one which was not inspected last year owing to extensive alterations then in progress, and two homes which had applied for exemption from registration under sub-section 1, Section 6, of the Nursing Homes Registration Act. The position of these two homes is the subject of special reference in paragraph 4 of this report.

One home which was inspected and placed on the register last year has never been started as a nursing home, and it is recommended that the registration be withdrawn.

3.-Assessment of Nursing Homes.

In the report for the year 1935, an attempt was made to assess in general terms, the efficiency of the nursing homes inspected, and to place them in the following categories:—

- (a) Satisfactory. These are nursing homes concerning which little or no criticism is offered. At most, certain minor additions to staff or equipment are necessary.
- (b) Moderately Efficient. These are nursing homes which are are open to criticism on a number of points. They all require fairly considerable additions to the nursing or domestic staffs and/or, in some cases, additional equipment.
- (c) Unsatisfactory. These are homes which possess marked defects in staff, equipment and/or management.

Under this scheme of assessment, in 1935, only four homes were adjudged to be "satisfactory." Eight fell into the group of "moderately efficient," and four were classified as "unsatisfactory."

Using the same classification, as a result of the inspection for 1936, the findings were :—

Three of the four homes which were counted as "satisfactory" last year continue to hold their place, but one was not entirely satisfactory when visited this year and must be placed in the second category of "moderately efficient" homes. To this group must be added five homes which were ranked in the second group last year but have so far improved as to allow them to be classified as "satisfactory," and one home which was not inspected last year. In short, nine nursing homes in the City can now be counted as "satisfactory."

Eight homes were last year classed as "moderately efficient." Two remain in this category, and to these must be added the one already mentioned which has lapsed from the "satisfactory" list. Two others, which last year were considered to be definitely "unsatisfactory" have sufficiently improved to justify their inclusion in the second group, thus making a total of five homes ranking as "moderately efficient." One home, which was placed in this second group last year, could not be classified this year, owing to extensive alterations which were in progress.

Of the four homes classed as "unsatisfactory" last year one must still remain in this category. Two of the four, as has been stated, have improved sufficiently to be ranked as "moderately efficient," while the fourth was undergoing extensive alterations and could not be classified.

Of the 17 homes referred to in this paragraph the following is a summary of classification:—

Satisfactory	 	9
Moderately Efficient	 	5
Unsatisfactory	 	1
Unclassified	 	2

Application for Exemption under Section 6 (Sub-Section I.) of the Nursing Homes Registration Act, 1927.

Two homes have made application for exemption from inspection under Section 6 of the Nursing Homes Registration Act, 1927. This Section states:—

- "(1) A local supervising authority may grant exemption from the operation of this Act in respect of any hospital or institution not carried on for profit.
 - (2) Any exemption granted under this section in respect of any hospital or institution shall continue in force for one year only from the date on which it is granted, but without prejudice to the power of the local supervising authority to grant any further exemption in respect of that hospital or institution."

These two homes were inspected and found to be "moderately efficient." On certain points both institutions fell short of the standard which has been laid down for nursing homes by the Health Committee. In one case it was evident that several of the wards were overcrowded and in both cases the number of the nursing staff was not entirely satisfactory. In addition to these major points, certain minor points were noted where improvements should be made.

Observations.

It can be said that, as a result of the exhaustive survey made last year, there has been some improvement in the general conditions of the nursing homes in the City. This has been most marked in two respects, firstly, in the number and qualifications of the nursing staffs employed, and secondly, in the standard of accommodation provided both for the nursing and domestic staffs. While, however, there is evidence of this improvement there yet remains room for further advance along these lines, since in many cases both the number of the qualified staff, and the standard of accommodation provided for nursing and domestic staffs, fall considerably below what is now provided in up-to-date General Hospitals.

6.—Recommendations.

It is suggested that the findings and recommendations of the Board be communicated to the keepers of the homes which have been classed as "satisfactory" and "moderately efficient," and that in the case of the home which has on both occasions been classed as "unsatisfactory," the keeper again be interviewed.

As regards the two homes applying for exemption under Section 6 of the Nursing Homes Registration Act, 1927, it is recommended that such exemption be granted, but conditional upon the rectification of the matters referred to in paragraph 4 above.

With regard to the nursing home which, though inspected and placed on the register last year, has never yet functioned as a nursing home, it is recommended that the registration be withdrawn.

E. F. Dawson-Walker,

Deputy Medical Officer of Health.

A. F. G. Spinks,

Maternity and Child Welfare Medical Officer.

J. L. WATT, Matron, City Hospital for Infectious Diseases.

Health Department,

Town Hall,

Newcastle upon Tyne, 1.

APPENDIX 1.

CITY AND COUNTY OF NEWCASTLE UPON TYNE.

Nursing Homes Registration Act, 1927.

LIST OF NURSING HOMES REGISTERED.

Ref. No.	Address.	Registered in the Name of
1.	26, Archbold Terrace	Miss M. M. Anderson
2.	10, Eslington Terrace	Miss M. Toyne
3.	1a, Clayton Road	Miss C. E. Balfour
4.	1, Park Terrace	Mr. R. J. Willan, F.R.C.S.
5.	5, Saville Place	Dr. H. Drummond
6.	4, Bentinck Terrace	Mrs. L. Newton
7.	10, Fernwood Road	Miss N. P. Hunter
8.	"The Minories," Jesmond Rd.	Mother Prioress
9.	"The Gables," Elswick Road.	The Matron
10.	24, Grosvenor Road	Miss Kirby and Miss Rooney
11.	6, Osborne Road	Miss M. H. Robertson
12.	"Elswick Lodge," Park Road	Chairman of Management Committee
13.	5. Osborne Terrace	Mrs. H. W. B. Gordon
14.	9, Windsor Terrace	Miss I. M. Middleton
15.	"Catherine House," 63, Osborne Road	Salvation Army
16.	"The Cheviot," Bowland Lodge, Western Avenue	Mrs. M. I. Behn
17.	10, Collingwood Terrace	Miss E. M. Myers
18.	"Cairney House," 10, Osborne Villas	Mr. J. Gilmour, F.R.C.S.
19.	Walker Accident Hospital, Airey Terrace, Walker	The Honorary Secretary
20.	*Northern Women's Hospital, 1a, Osborne Avenue	The Honorary Secretary

^{*} Exempted from Registration under Section 6 of the Act.

INCLUDING REPORTS OF THE
DEPUTY MEDICAL SUPERINTENDENT OF THE
INFECTIOUS DISEASES HOSPITAL
AND THE BACTERIOLOGIST.

III.—INFECTIOUS DISEASE.

FEVERS, FOOD POISONING,
CITY HOSPITALS FOR INFECTIOUS DISEASES,
DISINFECTION, BACTERIOLOGY.

THE TO STROTER ASSESSMENT OF THE PARTY OF TH

MEADERO EUOITOMENI-.III

PEVERS, FROD POISONING, CITY HOSPITALS FOR INFECTIOUS SHEARS! SIMULFECTION, BACTERIOLOGY

DEATHS (CORRECTED) FROM NOTIFIABLE INFECTIOUS DISEASES AND NON-NOTIFIABLE ZYMOTIC DISEASES.

berculos's.	V nL	:	o :	07	17	1 1	10	10	000	100	70	10	0 0	10	10	01	100	er er	400	32	308
Dysen- tery.			*****			:	:			*****						-		-			5
Diarr- hoea (under 2		-	-	9 0	12	= "	00	00	10	200	01	0 -	4		4		1 -	4	500	5	105
Whoop- ing	- Sono	-		-		-		-		-			: 0	7					-		7
Small- pox.		1					:	*****	:			-	****	****	::				:		1
Puer-		:			-		7				-		****		-	****		-	*****	61	10
Measles, and	Nuocina	-	:			-		-		.1	-	-	4	-				4	01		17
Polio- myelitis		-	:				*****		****	****	:::			*****		****	****	*****	*****	:	1
oenceppu-	Poli	-	:	****						****		:		****	*****		*****	*****	****	:	1
Enceph- alitis Lethar-	esta.	-				-				-	-	-	_					-	-	C1	10
Cere- bro- Spinal	rever.	-		1	_	_		-			:	1	_		****	:	*****	-		-	7
Pneu- monia.		-	4	21	17	23	22	11	9	24	21	16	0	6	6	4	11	13	16	24	256
Enteric Fever.		:	:		1010							****		****	****	::		_	-	:	61
Scarlet Fever.		:			-				****	-	1		*****	-	1		*****	5	-	-	os .
Ery- sipelas.		:	:		-	:		-	*****	-	*****	-	-	*****		-		61	4	-	12
Diph- theria.			61	61	3	-	****	2	****	6	9	****	-	*****	-	:	27	-	3	61	35
WARD.		St. Nicholas	St. Thomas'	St. John's	Stephenson	Armstrong	Elswick	Westgate	Arthur's Hill	Benwell	Fenham	All Saints'	St. Andrew's	Jesmend	Dene	Heaton	Byker	St. Lawrence	St. Anthony's	Walker	CLTY

Note: -All deaths in Public Institutions have been allotted to the Wards to which they properly belong.

NOTIFIED CASES OF INFECTIOUS DISEASE AND DEATHS (GROSS).

EXCLUSIVE OF TUBERCULOSIS.

AGES OF CASES OF INFECTIOUS DISEASE NOTIFIED AND DEATHS REGISTERED DURING THE YEAR 1936.

	org) lsti	Cases to Hosp	877 877 877 9688 331 32 1138 138 138 138	2118
T.	6.	Deaths.	35 10 10 17 17 17 17 17 17	359
NET TOTAL.	1936	*,sess.	693 176 937 8 8 8 3 3 51 546 1942 1441 1144	7650
	16	Desths.	20 6 6 17 17 10 10 10 10 10 10 10 10 10 10 10 10 10	490
GROSS TOTAL	1935	Cases.	688 257 1304 10 10 10 10 60 117 62 62 734 734 1897	8587
oss	936.	Desths.	43 110 110 110 110 110 110 110 110 110 11	475
GR	19	Cases.	705 193 955 8 10 5 35 95 95 95 1144	7852
	es it wn.	Deaths.		-
	Ages not known	Cases.		6
1 3	and by ds.	Desths.	9	69
	65 and up- wards.	Cases.	35	59
	o .	Deaths.		98
	45 to 65.	Cases.	9 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	165
só	9.	Deaths.	36 1 1 3 3 3 3 4 3 4 5 5 5 5 5 5 5 5 5	78
AGES—YEARS	25 to 45.	Cases.	85 57 58 58 4 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	345
ES	o .	Deaths.	1 2 2 1 1 1 1 1 1	28
Ar AG	15 to 25.	Cases.	250 20 20 38 38 38 38 38 38 38	313
A	15.	Desths.	9 2 4	33
	5 to 15	Cases.	476 9 9 582 3 2 2 2 1110 1110	3348
	9.	Deaths.	23 2 3 3 4 5 7 5 7 5 7 7 7 7 7	84
	1 to 5.	Cases.	110 110 1 1 155 155 2341 285	3144
	ler	Destps.	491 8	97
1 9	Under 1.	Cases.	7 4 4 7 1 2 2 4 4 7 2 3 3 0 0 3 3 6 3 8 6 9 8 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	469
	NOTIFIABLE DISEASE.		Diphtheria (including Membranous Croup) Erysipelas Scarlet Fever Enteric Fever Cerebro-Spinal Fever Acute Poliomyelitis Acute Poliomyelitis Acute Poliomyelitis Acute Poliomyelitis Puerperal Fever Puerperal Pyrexia Ophthalmia Neonatorum Pneumonia Malaria Dysentery † Chickenpox	

* Cases from outside the City excluded for the purpose of calculating NET Death Rates. ‡ Temporarily notifiable. † Ministry of Health Regulations, 1920.

 $\begin{array}{c} 7 \\ 333 \\ 2296 \\ 2296 \\ 2298 \\ 2298 \\ 2298 \\ 2209 \\$

TOTAL. Tuberculosis (all forms), 584 34 Dysentery. 1001 10-1 : 10 4 10 Malaria. 1144 Chickenpox, Smallpox. WARD DISTRIBUTION OF INFECTIOUS DISEASES (NET). Acute Influenzal Pneumonia. 39 0100-10-01-410 3101010-03-01: Acute Primary Pneumonia, 507 Neonatorum. 1010-00-010 51 Ophthalmia Puerperal Pyrexia. 13 Puerperal Fever, 0 :01 --01-36173712363333 97 Rubella 3925 32 66 120 365 291 266 117 241 159 279 338 523 Measles. Encephalitis Lethargica, Acute Polio-encephalitis. 3 Poliomyelitis. Cerebro-Spinal Fever. 00 - 01 937 Scarlet Fever. 00 Enteric Fever, - 1 2 2 3 2 3 2 3 4 4 9 4 9 8 176 Erysipelas, 693 Diphtheria. CITY ... Anthony's St. Lawrence. St. Anthony's Arthur's Hill All Saints' St. Andrew's St. Nicholas'
*St. Thomas...
St. John's ...
Stephenson ...
Armstrong ...
Elswick Nicholas' Thomas. Benwell.... Byker WARD Westgate Walker ... Heaton .. lesmond Fenham Dene ...

Royal Victoria Infirmary and Fleming Memorial Hospital for Sick Children. Elswick Grange and Newcastle General Hospital. City Hospital for Infectious Diseases, Walker Gate. Includes

: :

HOUSEHOLDS AFFECTED WITH INFECTIOUS DISEASES EXCLUSIVE OF TUBERCULOSIS, MEASLES AND CHICKENPOX.

			HOUSEHOLDS WITH	LDS WITH			Mili-			Cases.	
DISEASES.	Single Cases.	Cases each.	3 Cases each.	4 Cases each.	5 Cases each.	6 Cases and over	tary or Naval Cases.	Insti- tutions.	TOTAL CASES (Gross).	from outside of City.	CASES.
Diphtheria (including											
Membranous Croup)	550	40	00	2	1	1	*****	38	705	12	693
Erysipelas	149			****	*****	****		44	193	17	176
Scarlet Fever	723	89	90	3	1			55	955	18	937
Enteric (or Typhoid Fever)	7			****	****			1	00	-	8
Cerebro-Spinal Fever	9	****	****		****	*****	****	4	10	5	00
Poliomyelitis	61	****		***				3	9	2	3
Encephalitis Lethargica					:			*****			-
Polioencephalitis		****			****	-		****		****	***
Puerperal Fever	9	****	****	****	****			29	35	26	6.
Puerperal Pyrexia	10	-	-				****	85	95	76	19
Ophthalmia Neonatorum	43		****			-	-	11	54	3	51
Pneumonia	533	5	1	****	****			27	573	27	546
Malaria		****	****			****		****	*****	*****	****
Dysentery	20	-			::	:		19	41	7	34
-	2,049	114	17	5	2		1	316	2.674	190	2,484

* See next page.

INFECTIOUS DISEASES.

Schools and Infectious Disease.—It was not found necessary to close any school on account of infectious disease during the year.

PUBLIC INSTITUTIONS AND INFECTIOUS DISEASE.

The following notifications were received during the year:-

.лутоТ	252 48 48 48 10 10 10 10 12 12 12 12 12 12 12 12 12 12 12 12 12	357°
Dysentery.	797	19
.einelel/		- 1
Smallpox,		
Cerebro-Spinal Fever,	00	4
Polio- encepbalitis,	111 1 111111 111111	:
Poliomyelitis.		60
Ophthalmia Neonatorum,		=
Chickenpox.	1 2 1 2 1 3	21
Paeumonia.	2-: : -::::: ::::::	27
Pyrexia.	1 61 82 1 1 1 1 1 1	85
Puerperal Fever.	1 2 3	29
Measles and Rubella,	8 10 - - 20 -	20
Encephalitis Lethargica.	111 1 111111 111111	1
Enteric Fever.	1-1 1 111111 11111	-
Scarlet Fever,	20	55
Erysipelas.	848 2 1 4	44
Diphtheria.	21 1 1 1 1 1 3 2 3 3 3 3 3 3 3 3 3 3 3 3	38
Institutions, &c.	Royal Victoria Infirmary Fleming Memorial Hospital Newcastle General Hospital City Hospital for Infectious Diseases Princess Mary Maternity Hospital Military Barracks Eye Infirmary National Children's Home Throat, Nose and Ear Hospital Common Lodging Houses Royal Victoria School for the Blind Babies' Hospital, West Parade Northern Counties Orphanage Gables Maternity Home Nursing Homes Hostels, etc.	Total

* Does not include any cases belonging to the City which could properly be assigned to their homes.

SCARLET FEVER.

Notifications of 937 cases were received during the year, and there were 8 deaths, equivalent to a case mortality of 0.8 per cent.

DIPHTHERIA.

693 cases were notified during the year, and 35 died, a case mortality of 5.1 per cent., as compared with 5.2 in 1935.

MEASLES AND RUBELLA.

4,022 cases (including 97 of rubella) were notified, and there were 17 deaths (net) in 1936, representing a death rate of 0.06 per 1,000 population, as compared with 0.06 in 1935, and a case mortality of 0.4 per cent. of notified cases (net).

The following table shows the deaths in the various wards, and at different age periods:—

WARD.	Under 3 months.	3 and under 6 months.	6 and under 9 months.	9 and under 12 months.	1 and under 2 years.	2 and under 3 years.	3 and under 4 years.	4 and under 5 years.	5 and under 10 years.	Cver 10 years.	Totals
St. Nicholas'		****			****						
St. Thomas'											
St. John's					1						1
Stephenson			1111	1		****			****	****	1
Armstrong		1									1
Elswick				4444							
Westgate					****			1			1
Arthur's Hill								1			1
Benwell		1							****		1
Fenham				1111		1117	1111	1			1
All Saints'			****			****					
St. Andrew's					2	1	1				4
Jesmond											
Dene		4440				****					
Heaton											
Byker											
St. Lawrence	1		1			1		1			4
St. Anthony's	1		300		1						2
Walker							****	****		****	
TOTAL	2	2	1	1	4	2	1	4			17

Each Health Visitor visited and re-visited selected cases occurring in her district. By this arrangement each case is seen immediately on receipt of the notification, and advice is given regarding the nursing and isolation of the patient. The cases are kept under supervision until they recover, and should subsequent cases occur in the family they are recorded.

Measles Cases, including Rubella, notified during 1936.

Cases notified by Medical Practitioners Cases found by Health Visitors Cases notified by Parents and others Cases found from Returns of Deaths	3,347 668 16 3	
Less 12 cases from outside the City:—		gross. net.

Of the total number of measles cases notified, 3,625 in 2,916 households (or 89.9 per cent.) were visited by the Health Visitors, and 4,251 revisits were paid, a total of 7,876 visits.

The following particulars refer to the cases visited :-

Algorithm and the state of the		Dw	ELLINGS	oF.		Total.
	1 room	2 rooms	3 rooms	4 rooms	More than 4 rooms	Total houses visited.
Families	170	890	969	736	151	2,916
Children	333	2,050	2,237	1,923	349	6,892
Cases	203	1,147	1,203	897	175	3,625
Percentage of Cases to						
Children	61.0	56.0	53.8	46.6	50.1	52.6
Cases developing Pneumonia Percentage of Cases develop-	10	24	24	18	1	77
ing Pneumonia		2.1	2.0	2.0	0.6	2.1
Deaths from Measles	7	2	3	5		17
Deaths certified Pneumonia	1	2				3
Case Mortality per cent	3.9	0.3	0.2	0.6		0.6

Total unvisited cases 409, including 389 better-class houses in which no deaths occurred, and 19 in institutions, with 3 deaths.

Medical Attendance.—In 97.9 per cent. of the cases visited a doctor was in attendance.

Condition of Patient.—In 89.8 per cent. of the cases visited the disease ran a normal course, but bronchitis, pneumonia or other complications developed in the remainder.

Attendance at Schools.—1,101, or 30.4 per cent., of the affected children had previously attended school, and 2,524, or 69.6 per cent., had never attended school. In connection with 1,298 of the latter cases, however, other children in the infected houses were scholars, equivalent to 35.8 per cent. of the total cases.

The following were the ages of visited children who were suffering from measles:—

Under 1	year	 290
1-2	years	 501
2-3	years	 537
3-4	vears	 594
4-5	years	 594
5-6	vears	 718
Over 6	years	 391
		3,625
		-

WHOOPING COUGH.

7 deaths occurred from whooping cough. The particulars are as follows:—

Winn			YEARS	of Age			Total
WARD.	0-1.	1-2.	2-3.	3-4.	4-5.	5-10.	Total
St. Nicholas'							
St. Thomas'	1111	****					
St. John's	1						1
Stephenson							
Armstrong	1						1
Elswick							
Westgate			1				1
Arthur's Hill							
Benwell	1						1
Fenham							
All Saints'		101					
St. Andrew's						****	
Jesmond	1	1					2
Dene			****		****	****	
Heaton							
Byker							
St. Lawrence							
St. Anthony's		1		****			1
Walker				1111		3334	****
CITY	4	2	1		-		7

The death rate in 1936 was equivalent to 0.02 per 1,000 population, as compared with 0.08 in 1935.

FOOD POISONING.

Eleven cases of illness due to organisms of the food poisoning group were notified during the year. Two of these, a man and his wife, were found to have been infected with the para typhoid C. bacillus. The history was that they had attended a garden party held in a village in the County of Durham, where they had eaten some salmon sandwiches. A number of other persons who also partook of these sandwiches were taken ill, and were found to have been infected with this organism.

The remaining nine cases were due to infection with the bacillus ærtrycke.

Six were notified from the Fleming Memorial Hospital, three being Newcastle cases and three extra mural cases.

Two were notified from the Newcastle General Hospital, and one from the Royal Victoria Infirmary, the latter being a Newcastle case. All of these cases were transferred to the City Hospital, with the exception of one extra mural case, which died at the Fleming Memorial Hospital.

One case died in the City Hospital. No source of infection was traced in these nine cases.

ENTERIC GROUP OF FEVERS

During the year 1936, 15 cases of the enteric group of infections were brought to notice. The distribution of these cases, according to the months in which they were notified, the type of infection (typhoid or para-typhoid), and their place of origin, is recorded in the following table:—

Distribution of Enteric Group Infections for 1936.

	EXTRA-	MURAL.	Newo	ASTLE.
	Typhoid.	Para- typhoid B.	Typhoid.	Para- typhoid E
January		1		
February				****
March			1	1
April				
May	****			
June		1	****	****
July		1		1
August	1		1	
September	****		2(2)	****
October	****		1	
November	1	1	****	
December	1		1	
Totals	3	4 .	6 (2)	2

Figures in parentheses indicate deaths.

It will be seen that seven patients came from without the City's boundaries, the remaining eight being Newcastle cases proper. Of the seven extra mural cases, four were notified from the Royal Victoria Infirmary, two from the Fleming Memorial Hospital, and one from a Nursing Home in the City. All these cases were admitted to the City Hospital, at the request of the Local Authority concerned.

The eight City cases, which were all admitted to hospital, were made up of 6 cases of typhoid and two para typhoid B. infections. Two of the typhoid cases proved fatal. The cases were all of the sporadic type, and no connection could be traced between them.

In all there were 25 admissions to the City Hospital. Apart from the 15 patients mentioned above, these cases were all notified in the areas of neighbouring Local Authorities, and were admitted to the City Hospital at the request of the Local Authority concerned. Seven of these cases were suffering from typhoid fever, one having been infected in the Bournemouth epidemic which occurred during the summer. Another was a chronic carrier of the disease who was sent in for special investigation, in order to see whether it was possible to eradicate the infection.

The remaining five cases, four of whom were members of the same family, were infected in a localised outbreak which occurred in a mining village in Northumberland. The remaining three cases were para typhoid B. infections.

Among the 25 cases admitted to hospital, there were four deaths, all due to typhoid infection, equivalent to a case mortality of 16 per cent.

In six cases anti-typhoid serum, introduced by Felix & Pitt, was given, three of these cases proving fatal.

DIARRHOEA.

There were in all 126 deaths from the disease, equal to a death rate of 0.43 per 1,000 population, and this number included 105 deaths of children under two years of age.

SMALLPOX.

No case of this disease occurred in the City during the year.

The following are the particulars of **Vaccination** during the last thirty-two years:—

	rar. Registered Vaccinations Vaccinations 005 7,958 7,264 2 006 7,721 6,733 2 007 7,610 6,702 1 08-12 35,265 27,240 11 3-17 34,296 21,251 3 8-22 34,372 19,011 9 928 5,780 4,320 1 929 5,638 3,555 3 930 †6,195 3,897 3 931 6,059 3,754 3 932 6,009 3,600 2 933 5,770 3,479 1	Unsuccessful	Exemptio	n Certificates.	Deaths, Removals	
Year.		Vaccinations	Number.	Percentage to Total Births.		
1905	7,958	7,264	27	65	0.8	
1906	7,721	6,733	28	92	1.2	
1907	7,610	6,702	16	94	1.2	****
*1908-12	35,265	27,240	114	3,398	9.6	
1913-17	34,296	21,251	33	7,144	20.8	****
1918-22	34,372	19,011	95	9,262	26.9	
1923-27	31,290	19,658	30	5,542	17.7	
1928	5,780	4,320	19	912	15.8	****
1929	5,638	3,555	33	1.092	19.4	
±1930	†6,195	3,897	31	1,264	20.4	1,003
1931	6,059	3,754	39	1,343	22.2	923
1932	6,009	3,600	27	1,395	23.2	889
1933			18	1,377	23.9	809
1934	5,890	3,467	27	1,449	24.6	874
1935	5,899	3,474	32	1,401	23.7	901
§1936	5,713	2,798	26	1,305	22.8	

- * Vaccination Act, 1907, came into force.
- † Walker District included.
- ‡ Supervision of Vaccination transferred from Guardians to Health Committee on 1st April, 1930.
- § Provisional figures only.

CHICKENPOX.

1,144 cases were notified. There were no deaths.

ERYSIPELAS.

176 cases of this disease were notified and there were 12 deaths.

PUERPERAL SEPTICÆMIA AND PUERPERAL PYREXIA.

28 cases were notified, with 10 deaths. Inquiries were made concerning 26 of these.

INFLUENZA AND PNEUMONIA.

These diseases accounted for 294 deaths as against 327 last year.

Total deaths at age periods.

Under 5 years.	5-15.	15-25.	25-45.	45-65.	65 and over.	Total.
84	6	13	34	85	72	294

As will be seen from the above figures, 84, or 28.6 per cent., of the deaths occurred below the age of 5 years.

546 cases of pneumonia, including influenzal-pneumonia, were notified. For the ages and ward distribution, see pages 72 and 73.

Of that number 478, or 87.5 per cent., were visited by Health Visitors. It was found that 325, or 68.0 per cent., were primary pneumonia, 54, or 11.3 per cent., were cases of influenzal-pneumonia, and 99, or 20.7 per cent., were cases of pneumonia following other diseases.

Sex.—60.3 per cent. of the cases were males.

Ages.—The ages of the 478 cases visited were as follows:—

Under 1	year	45
	years	145
	years	99
	years	46
	years	59
	years	62
	years	22
		478

Housing.—24 cases occurred in 1 roomed dwellings, 144 cases occurred in 2 roomed dwellings, 146 cases occurred in 3 roomed dwellings, and 164 cases occurred in more than 3 roomed dwellings.

Type of House.—194 cases occurred in flats, 154 cases in tenements, and 130 in self-contained houses.

Previous History .-

There wa	s a previous	history of	Measles	in	207	cases.
,,	**	,,	Whooping Cough	in	149	cases.
,,	,,	,,	Influenza	in	94	cases.
,,		,,	Frequent winter			
			Coughs and Colds	in	376	cases.
.,,	**	,,	Pneumonia	in	134	cases.
	T.,		Tuberculosis	in	9	cases.

Hospital Treatment.—140 cases of pneumonia were treated in the Infectious Diseases Hospital. The majority of these were from houses where there was over-crowding or other unsuitable home conditions. 34 of these patients died, giving a case mortality of 24.4 per cent.

Deaths.—104, or 21.8 per cent., of the visited cases of pneumonia died.

ENCEPHALITIS LETHARGICA.

One case of encephalitis lethargica was admitted to the City Hospital during the year.

ACUTE POLIOMYELITIS.

One proved case of poliomyelitis was admitted to the City Hospital during the year, and was later transferred to the Newcastle General Hospital.

CEREBRO-SPINAL MENINGITIS.

During the last three years the incidence of cerebro-spinal fever, which for some years previous to 1932 had been steadily increasing, has somewhat declined, and during 1936 17 cases were notified in Newcastle. The figures for 1934 and 1935 were 31 and 27. Twenty-six cases of this disease were nursed in the City Hospital during the year. Eight of these were Newcastle cases, while the remainder were admitted either direct, or through one of the hospitals in the City, from surrounding areas.

There were 10 deaths among these 26 cases, equivalent to a case mortality rate of 38.4 per cent. The corresponding figures for 1934 and 1935 were 52.9 per cent. and 44.4 per cent.

The distribution of these cases, according to the months in which they were admitted, and their places of origin, is recorded in the following table:—

the state of the s	Newcastle.	Extra-Mural.	Totals.
January	2(1)	1(1)	3 (2)
February	1	2	3
March		2(1)	2(1)
April	1	2 (1)	3(1)
May	1 (1)	1	2(1)
June		2	2
July	1(1)	1	2(1)
August	1	1(1)	2(1)
September		1	
October		1(1)	1 (1)
November	1	2(1)	3(1)
December	****	3 (1)	3 (1)
Totals	8 (3)	18 (7)	26 (10)

The figures in parentheses, which are included in the numbers alongside which they stand, indicate fatal cases.

The circumstances of all the Newcastle cases have been carefully investigated, but in no case has it been possible to trace the source of the infection.

More than two-thirds of the cases of cerebro-spinal fever admitted to hospital came from extra-mural authorities in the neighbourhood, and wherever possible, it has been the policy of the Health Department to give assistance to authorities whose hospital accommodation is of such a character as to prevent them from giving adequate treatment to patients suffering from this extremely dangerous disease. The following table shows the age and sex distribution of the 26 cases admitted to hospital:—

Ages.	0-1.	1–2.	2-5.	5–15.	15-25.	25-45.	45 and up- wards.	Totals.
Male Female	3 (1)	2 (1)	3 (3)	5 (2) 2 (2)	7	1	1 (1)	21 (7) 5 (3)
Totals	4 (1)	2 (1)	3 (3)	7 (4)	8	1	1 (1)	26 (10)

Figures in parentheses indicate deaths.

It will be noted that the mortality rate is particularly high under the age of 5, and that young persons and adults from 15 to 25 have the best chance of recovery from the disease.

BACILLARY DYSENTERY.

Bacillary dysentery has been prevalent in the City since 1928, and during the past year 94 cases were notified. In 53 of these cases the diagnosis was confirmed bacteriologically, six of these being extra-mural cases, which had been admitted to one of the City's hospitals suffering from the disease.

Forty-six proved cases of bacillary dysentery were admitted to the City Hospital. Among these cases there were six deaths.

During the year 17 cases were notified from the Newcastle General Hospital. There was one outbreak in this Institution, which occurred in the autumn. During the month of October ten cases were notified, four being due to the Sonne organism, and six to the Flexner type. The remainder were sporadic cases.

Nine cases occurred in the Fleming Memorial Hospital during the year, all being of the sporadic type. Of the total nine cases, five were due to the Sonne organism, three to the Flexner organism, and one to the Newcastle type. There were two deaths in the Sonne group, and one in the Flexner group. Three of the cases were extra mural in origin.

Five cases were notified from the Royal Victoria Infirmary during the year.

There was one outbreak which occurred in the City Hospital on a diphtheria ward. It was caused by the Flexner type of organism, and was due to careless nursing.

The circumstances and history of all cases were carefully investigated with a view to obtaining information as to the probable sources of infection. The age, sex, and mortality incidence of the series of 53 cases are given in the following table:—

Ages.	0-1.	1-2.	2-5.	5-15.	15-25.	25-45.	45 and up- wards.	Total.
Males Females	2 (1) 3 (2)	6 5 (2)	6 7	9 (1)	3 1 (1)	4	1	31 (2) 22 (5)
Total	5 (3)	11 (2)	13	14 (1)	4 (1)	5	1	53 (7)

The figures in parentheses indicate fatal cases.

The distribution of these organisms among the cases is as follows:—

	1 5 1 1	ER.		Sonne Bacil-	astle	Atypical	Not	Totals.		
train a la	w.	X.	Y.	Z.	XZ.	lus.	Newcastle	Atyl	Tyl	Totals.
Total No. of Cases	1	5	1	11	3	22	7	1	2	53
Fatal Cases			1	1		3	1		1	7
Non-Fatal Cases	1	5		10	3	19	6	1	1	46

VENEREAL DISEASES.

Syphilis was certified as the cause of death in 27 cases.

The work of the treatment clinic has been continued successfully. 1,451 old and new cases attended 23,807 times as outpatients. 10 cases accounted for 149 in-patient days. Of the 769 new cases, 132 were syphilis, 410 gonorrhea, 3 soft chancre, and 224 were conditions other than venereal. 68 per cent. were males.

2,555 doses of salvarsan substitutes, 3,431 doses of mercury and 1,085 doses of bisoxyl were administered to out-patients and in-patients.

1,170 Wasserman reactions were carried out by the College of Medicine, and 1,350 microscopical examinations of pathological material were made at the treatment clinic. The irrigation stations for males and for females in connection with the clinic have been in full use during the year.

56 medical practitioners in the City are qualified to receive free supplies of arseno-benzol compounds. 15 made application for these supplies during the year and 847 doses were given.

Newcastle Residents Notified as Attending other Centres.

Cases.—Syphilis, 2; gonorrhoea, 7; conditions other than venereal, 7.

Attendances. -362.

Doses of salvarsan substitute given to out-patients and inpatients, 53.

In-patients.—In-patient days, 84.

Information as to ophthalmia neonatorum will be found in Section II. (The Child).

CITY HOSPITALS FOR INFECTIOUS DISEASES.

Report of the Deputy Medical Superintendent.

Accommodation.

NAMES AND SITUATION OF HOSPITALS.	TOTAL AVAILABLE BEDS.
City Hospital for Infectious Diseases, Walker Gate— Beds. Fever Pavilions	
Tuberculosis Pavilions	338 172

City Hospital, Walker Gate.

YEAR.	Population of the City.	Number of Beds at Hospital for Fever Cases.	Total Admissions (exclusive of Pulmonary Tuberculosis and Smallpox).	Percentage of Scarlet Fever, Diphtheria and Enteric Fever Cases Admitted to Cases Notified.
1890	182,866	104	219	21.3
1900	213,039	104	290	38.6
1910	265,077	172	912	83.0
1920	286,061	232	1,710	86.4
1921	278,400	232	1,683	82.4
1922	281,600	232	1,032	86.3
1923	283,800	232	991	92.6
1924	285,900	232	1,502	90.5
1925	286,300	*232	1,711	86.4
1926	284,700	*232	1,397	89.1
1927	288,500	*232	1,493	89.7
1928	281,500	*232	1,294	92.9
1929	283,400	*232	1,713	89.1
1930	283,400	*232	1,649	96.4
1931	283,600	*232	2,347	95.6
1932	285,100	*232	2,143	96.4
1933	286,500	*232	3,040	96.3
1934	287,050	*232	3,292	95.3
1935	292,700	*232	2,881	97.2
1936	290,400	*232	2,471	97.0

^{* 30} of these beds temporarily appropriated for Tuberculosis patients.

CITY HOSPITAL, WALKER GATE.

(Fever Pavilions.)

Admissions during the year-2,471.

The average daily number of patients in the hospital was 213, exclusive of 98 cases of Tuberculosis.

RATE PER CENT. OF CASES REMOVED TO HOSPITAL TO CASES NOTIFIED.

co	-		0	
1936	96.0	98.4	100.0	96.7
1935	96.3	98.7	0.001	6.96
1934	94.5	98.2	0.001	95.0
1933	96.1	100.0	0.001	0.96
1931 1932	96.3	8.96	0.001	96.3
1931	95.2	1.66	92.3	92.6
1930	95.9	97.5	97.6	96.1
1925	85.0	94.1	96.4	86.0
1920	85.7	89.1	0.06	86.4
1915	91.3	89.1	87.0	90.5
1910	84.5	80.1	90.5	83.0
1905	50.1	36.8	52.0	47.8
	35.0	40.0	54.5	38.6
1890 1895 1900	18.4 33.0	28.7	48.0	34.6
1890	18.4	8.3	38.9 48.0	21.3
	Scarlet Fever	Diphtheria	Enteric Fever	All cases of the above, together with Continued and Typhus Fever and Cerebro-Spinal Fever, etc.

CITY HOSPITAL FOR INFECTIOUS DISEASES, WALKER GATE.

Diseases Admitted-1936.

													1	Prov.	ED T	O BE	:											
SENT IN AS	Number.	Scarlet Fever.	Diphtheria.	Diphtheria Carriers.	Enteric Group of Fevers.	Dysentery.	Measles.	Rubella,	Varicella.	Mumps.	Pertussis.	Epidemic Cerebro- Spinal Meningitis,	Other forms of Meningitis.	Poliomyelitis.	Encephalitis Lethargica,	Pneumonia.	Bronchitis.	Influenza.	Other Respiratory Diseases.	Erysipelas,	Skin and Septic Conditions,	Puerperal Pyrexia.	Tonsillitis,	Gastro- Intestinal Diseases.	Ophthalmia Neonatorum,	General Diseases,	Injuries,	Inchesition
Scarlet Fever	968	915	2		1		3	6			1				l	4	2		l		3		9	1,	1			23
Diphtheria	679	12	526		1		5		****		****		****			1			12		1		101	6		****	****	1
piphtheria Carriers	222		1	220	-					1775	****	****														****	****	1 1
nteric Group of Fevers	31		100		23	2							1			1		1						1				
ysentery	74				1	44	1			****		****		****		î			****			****		25				
leasles	93	2	1				87						****			2			****		****		****					
Rubella	2		****					2											****									
aricella	3								2							1		****										
lumps	3									2											****		1		****			1
nthrax	1												****								1			****	****			1
ertussis	8				****					****	5					1	1							1				
Epidemic Cerebro-Spinal																10000												
Meningitis	32	****	****	****		****		****	****			23	5	+++4		2 5				****	****	****	****	****				
Other forms of Meningitis	21	****	****				1					2	9			5	1	1		****	****	****	****	1				
Poliomyelitis	1	****				****			****					1	1111	1000					****		****	****	****	1110	3141	
Encephalitis Lethargica	2		****	****	****										1	****		****		****	****	****				****		
neumonia	138	****	****	****	****	****	1		****		2	1	****	****	****	121	5		5	****	****	****	****	1	****			3
Bronchitis			****	****	****						****			****	****		****		****	****	****						****	1
nfluenza	9		****	****	****	****	****	****	****	****	****	****		****	****	****	****	9	****		****		****	****	****	****	****	
Other Respiratory Diseases			****		****				****	****					****		****	1444				****	****		****	****	****	
Erysipelas	87					****	****		++++	****				****	****	1	****	****	****	80	5	****	****	****	****	1		
kin and Septic Conditions	30												****		****				****	****	29		****	****			****	
Puerperal Pyrexia	2		****	****					****			****	****	****	****	****	****	1448	****	****	****	2		****	****	****	****	
Consillitis	19		****	1						****	****	****	****	****		****	1-14	****	****	****	****	****	17	1		****		+-
Gastro-Intestinal			100																					10				
Diseases	19	****	****										1	****	****	****	****	****	4111	****	****	****	****	18			****	**
phthalmia Neonatorum	6	****				****							****	****	****	1111	****	****	21.44	****	****	****	****	1	5			
General Diseases			****				****			****		****	*****	****	****	****	2752	2111	****	****	****	****	****	****	****			
njuries	6 15	****	12.55	****		****	****			****				****	****	****	*****	****	1	****	7.00	****	****	1	****	****	6	1
Unclassified	15	****	****	****		****	****								****		****			****	****	****	****	1	****	****		1:
Totals	2,471	929	530	221	25	46	98	8	2	2	7	26	16	1	1	140	9	11	18	80	39	2	128	57	5	1	6	6



Present Death Rates compared with those of Previous Years.

RETURN SHOWING THE NUMBER OF CASES OF
SCARLET FEVER, DIPHTHERIA AND ENTERIC FEVER ADMITTED TO HOSPITAL
AND MORTALITY RATES PER CENT.
1891-1900.

YEARS.		BER OF C		NUMB	NUMBER OF DEATHS.			E MORTA PER CENT	
TEAKS.	Scarlet Fever.	Diph- theria.	Enteric Fever.	Scarlet Fever.	Diph- theria.	Enteric Fever.	Searlet Fever.	Diph- theria.	Enterio Fever.
1891-1895	1,105	92	277	34	26	51	3.1	28.3	18.4
1896-1900	1,087	103	442	41	21	86	3.8	20.6	19.5
			19	15–193	4.				
1915-1919	3,402	998	194	99	89	21	2.9	9.0	10.8
1920-1924	3,919	1,037	78	37.	73	9	0.9	7.5	11.6
1925-1929	3,612	908	123	43	62	23	1.2	6.8	18.7
1930-1934	6,296	860	220	76	53	15	1.2	6.1	6.8
			19	935-193	6.				
1935	1,236	549	23	9	41		0.7	7.4	
1936	929	530	25	12	40	4	1.2	7.5	16.0

	Totals.	20 4 9 8 1 1 1 1 1 1 1 1	152
	December.	0101 01	14
	November.		00
	October.	2	13
	September.	0 -	01
HS.	August.	199	7
DEATHS	July.	- -	-
П	June.	01	7
	May.	0 -0 1 -0 4 -	91
	April.	010 01 1 -0 4 01 1 - 5	50
	Матсh.		12
	February.	-0 -	16
-	January.		18
	Totals.	0.00040 01 4 1180 0 10 0 1	2471
	December.	103 24 25 27 27 28 11 11 11 11 12 13 14 14 14 15 16 17 17 18 18 18 18 18 18 18 18 18 18 18 18 18	230
	November.	99 34 17 17 18 19 19 19 19 19 19 19 19 19 19 19 19 19	236
	October.	83 22 23 27 20 20 20 20 20 20 20 20 20 20 20 20 20	220
	September.	2220428	143
NS.	August.	310 4 6 1	136
DMISSION	July.	25° 1 2 2 1 1 2 1 1 2 1 1 2 1 1 2 1 1 2 1 1 2 1	119
ADM	June.	208 91 102 1 1 2 1 1 1 1 1 1 1 1 1 1 2 1 1 2 1 1 2 1 1 2 1 1 2 1 1 2 1	121
	May.	474 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	231
	.lingA	55 12 12 13 13 13 13 13 13 13 13 13 13 13 13 13	961
	Матсh.		272
	February.	467 1 1 1 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	260
	January.		307
	DISEASE.	Scarlet Fever Diphtheria Diphtheria Carriers Enteric Group of Fevers. Dysentery Measles Rubella Numps Pertussis Epidemic Cerebro-Spinal Meningitis Other forms of Meningitis Poliomyelitis Poliomyelitis Encephalitis Lethargica Bronchitis Influenza Other Respiratory Diseases Erysipelas Skin and Septic Conditions Puerperal Pyrexia Tonsillitis Gastro-Intestinal Diseases Ophthalmia Neonatorum General Diseases Injuries Unclassified	Totals

Length of Stay in Hospital of Early Fatal Cases.—The following cases died within a short period after their admission to hospital:—

		in urs.	Within 48 hours	
Scarlet Fever	3		1	
Diphtheria	14		3	
Dysentery			1	
Measles	1			
Encephalitic Lethangian	1		****	
Encephalitis Lethargica	. 1			
Epidemic Cerebro-spinal Meningitis	2		2	
Other Forms of Meningitis	. 3		1	
Pneumonia			4	
Other Respiratory Diseases				

Erysipelas			1	
Puerperal Pyrexia			1	
Gastro-Intestinal Conditions	2			
General Diseases			1	
Series as to the series and the series are the series are the series and the series are the seri				
	-		-	
Total	28		15	
	Name of Street		*****	

Average Stay in Hospital during the last Twenty-nine Years.

All Cases.		Scarlet Fever.		Diphtheria (inc'uding carriers).		Enteric Fever		Other Diseases.		
TEARS.	Average No.	Average Stay in Days.	Average No.	Average Stay in Days.	No.	Average Stay in Days,	No.	Average Stay in Days.	Average No.	Average Stay in Days.
1908-12. 1913-17. 1918-22. 1923-27. 1928. 1929. 1930. 1931. 1932. 1933. 1934. 1935. 1936.	1,538 1,408 1,419 1,294 1,713 1,649 2,347 2,143 3,040 3,292 2,881	46.7 39.6 31.2 31.9 22.5 21.7 23.9 27.3 30.3 27.6 30.1 31.2 31.3	599 929 758 751 452 543 584 989 1,120 1,934 1,669 1,236 929	51.7 45.6 37.1 35.2 29.3 29.7 32.5 36.5 35.2 32.7 35.0 33.8 33.8	326 220 215 185 205 247 194 113 162 114 503 736	41.3 39.9 43.2 44.3 33.6 29.6 34.7 46.3 57.5 61.6 41.1 45.5 43.4	68 70 15 21 25 38 66 21 33 34 66 23 25	46.3 47.4 46.6 54.0 44.5 42.2 44.3 50.2 47.0 41.2 40.0 37.2 43.8	61 318 420 462 612 885 805 1,224 828 958 1,054 886 766	29.6 20.6 16.8 21.1 12.9 13.6 13.5 17.8 17.7 12.7 15.5 16.6

DIPHTHERIA.

The marked increase in the incidence of diphtheria in Newcastle and the surrounding area, which first became evident during the autumn of 1934, persisted through 1936, as is shown by the number of cases and carriers admitted to hospital. As the above table shows the combined figures for these cases, which since 1929 had never reached 200, in 1934 rose to 503, and in 1936 reached 751. While the number of cases and carriers combined in 1936 exceeded that of 1935 by 15, there were actually 19 fewer cases and 34 more carriers than in the preceding year.

530 true cases were admitted to hospital, and 221 carriers. 404 of these cases were of the simple faucial or tonsillar type, and in this group there were four deaths. Of these fatal cases, one died suddenly two hours after admission, another died within 24 hours of admission, a third died after operation for intestinal obstruction, and the fourth developed a severe toxic myocarditis.

In 21 cases the infection was limited to the nose, and there were two cases of aural diphtheria, one requiring a mastoidectomy.

In a group of 75 faucio-pharyngeal cases, with varying degrees of nasal involvement, there were 23 deaths—equivalent to a case mortality of 30.6 per cent.

There were 28 cases of laryngeal or tracheal diphtheria of whom thirteen, or 46.4 per cent. died. In sixteen of these cases the obstruction was so considerable that tracheotomy was performed shortly after admission to hospital, and of these nine died. In two cases tracheotomy had been performed before admission to hospital, and both of these died.

The case mortality of the whole series of 530 cases was 7.5 per cent., compared with 7.4 per cent. in 1935, and 7.6 per cent. in 1934.

The death rate for diphtheria has been practically constant for the last three years. This covers the period during which the disease has become more prevalent in this district, and since the rates for the two preceding quinquennial periods were 6.8 per cent. and 6.1 per cent. respectively, it can be said that, although a proportion of severe cases are being met with, the virulence of the infection does not seem to have increased with the incidence of the disease in this area to the same extent as it would appear to have done in certain other areas.

In one hundred cases, where virulent organisms persistently remained in the throat after recovery from the disease, including also a number of healthy carriers, tonsillectomy was performed. In the majority of cases this procedure rendered the patient free from infection after a short period. Mastoidectomy was carried out on two cases, one of which was a case of aural diphtheria.

The increased number of cases necessitated the use of one spare ward throughout the year, and for the first six months of the year it was found necessary to requisition for diphtheria cases the pavilion at Walker Gate which usually accommodates thirty of the tuberculosis patients. It was also necessary for a period to utilise a ward at the Moor Hospital for the accommodation of convalescent cases and carriers.

The scheme, inaugurated in June, 1934, whereby free immunisation against diphtheria of all children under the age of five was offered to parents, has been continued throughout 1936. The work has been carried out by a part-time officer, and four clinics have been held each week at certain Maternity and Child Welfare Centres. A number of children of school age, whose parents were anxious that they should be immunised, have been treated at clinics held on Saturday mornings.

In the autumn, in co-operation with the Education Authority, it was decided to circularise the parents of the children in a number of the schools of the City, emphasizing the advantages of prophylactic inoculation, and offering free facilities.

The response was most encouraging and the numbers were so large that it was decided, with the consent of the Education Authority, to hold the clinics in the schools, thus minimising as far as possible the amount of time during which the children would be absent from their classes. These clinics commenced in the last week in November, and ten had been held before the end of the year.

A total of 177 clinics were held during the year, 124 being for infants and 53 for school children.

During the year 1,114 infants and 352 school children were immunised, a total of 1,466.

An investigation, commenced in 1934, was continued during 1936, into the incidence and clinical significance of the three types of diphtheria bacilli, *Gravis*, *Intermedius and Mitis*. The following facts for 1936 have been supplied by Dr. J. F. Caithness. Wherever possible the organisms from each patient were isolated and classified by the Public Health Laboratory, and an intra-dermal virulence test performed.

In 589 patients—cases and carriers which were proven bacteriologically—the diphtheria bacilli isolated were subjected to further examination in order to classify them into their respective types—Gravis, Intermedius, and Mitis.

Of these 42, or 7.1 per cent., whilst exhibiting virulence as determined by the intra-dermal guinea pig inoculation test, failed in their characteristics to conform to any of the three types.

A further 15, or 2.5 per cent., were capable of classification but were non-virulent to the intra-dermal test; 3 were Gravis, and 12 were Mitis in type. The following table shews the distribution between those types of the remaining 532 bacilli, which were virulent and which could be classified into their respective types. The clinical severity of the cases from which they were isolated is also indicated

	No. of Cases.	Per-		ditta I	CL	INICAL CLAS	SIFIC	ATION		
		centage of Total.	ntage Severe.		Moderate.			Mild. Ca		arriers.
		or rotal.	No.	Percentage	No.	Percentage	No.	Percentage	No.	Percentage
Gravis	179	33,64	44	24,58	55	30.73	44	24.58	36	20.11
Intermedius	98	18,42	21	21.42	36	36.73	28	28.57	13	13,26
Mitis	255	47.93	14	5.49	63	24.70	91	35,68	87	34.11

The number of deaths occurring in each group were :-

Gravis ... 13 deaths.

Intermedius ... 4 deaths.

MITIS ... 4 deaths, 1 due to broncho-pneumonia.

SCARLET FEVER.

The incidence of scarlet fever in the City which, as shown by the notification and admissions to hospital, had been gradually increasing since 1928, until it reached its peak in 1933, now appears to be on the wane, and in 1936 929 cases of scarlet fever were admitted to hospital, as against 1,669 in 1934 and 1,236 in 1935. This figure is, however, somewhat in excess of what would be termed a normal year.

The Isolation Hospital on the Town Moor was in use until the end of May to accommodate convalescent cases.

The prevailing type of scarlet fever remained mild, and the mortality rate was 1.3 per cent., as compared with 1.8 per cent. in 1934, and 0.7 per cent. in 1935.

Of the twelve fatal cases six had received antitoxin. One was suffering from severe burns and died three hours after admission, another died from an acute cellulitis which spread from a septic finger, and a third was admitted with an appendix abscess and died shortly after operation. Of the other three in this group, one was an acute case which died four hours after admission, another developed a streptococcal pharyngitis, and the third, a child of one year, had a double mastoidectomy performed, and subsequently developed a streptococcal tracheitis which necessitated a tracheotomy.

The six cases which did not receive antitoxin consisted of one case suffering from burns, one from scalds, a case desquamating on admission which had developed a mediastinal abscess, and one which was admitted as mastoiditis following scarlet fever from an extra mural area. This child died from secondary hæmorrhage in the operating theatre before the operation was commenced. The infection, spreading from the mastoid area, had eroded the jugular vein, and the subsequent hæmorrhage had formed a retro-pharyngeal swelling which suddenly burst.

Of the remaining two cases, one developed broncho-pneumonia, and one died very suddenly from cardiac failure.

Seven deaths were, therefore, due to scarlet fever and its complications, without any additional factor, giving a more accurate death rate of 0.75 per cent.

Scarlet fever antitoxin has been used to a slightly greater extent than last year, but to a lesser extent than in the six previous years. The numbers and relative proportions of patients receiving this form of treatment for the period 1926-1936, are as follows:—

	1926	1927	1928	1929	1930	1931	1932	1933	1934	1935	1936
Scarlet Fever Cases											
admitted Number treated	831	741	452	543	584	989	1,120	1,934	1,669	1,236	929
with Antitoxin Percentage treated	78	172	177	169	249	483	380	436	331	260	243
with Antitoxin	9.5	20.3	39.2	31.1	42.6	48.8	33.9	22.5	19.8	22.0	26.1

Scarlet fever antitoxin has now been in use for eleven years, and opinions as to its value and its limitations are becoming more definite.

It seems clear that while it is very efficacious in the treatment of severe cases showing signs of toxæmia, and of considerable value in true "toxic" cases, its power to prevent the onset of complications is negligible, and furthermore, it is of little value in the treatment of these complications or the septic sequelæ of scarlet fever.

The general consensus of opinion seems to be that it should be reserved for the treatment of the more severe cases, and that its routine use in mild cases of scarlet fever is not advisable. In the following table is summarised the statistical information regarding all cases of scarlet fever treated during the year under review:—

Cornena	Num	Per- centage	Per-			Average stay in Days in Hospital.			
Scarlet Fever.	Num- ber.	treated with Anti- toxin.	centage with Compli- cations.	Mor- tality Rate.	Return - Case Rate.	All Cases.	Complicated Cases.	Non- compli- cated Cases.	
All Cases	929	26.1	33.8	1.2%	3.6%	33.3	43.8	27.7	
Antitoxin Cases	243	100	37.8	2.47%	4.9%	34.8	45.8	28.1	
Non-Anti- toxin Cases	686	Nil.	32.3	0.87%	2.7%	32.7	42.9	27.6	

PERCENTAGE INCIDENCE OF COMPLICATIONS.

- 0	Rhin- orrhœa.	Ot- orrhœa.	Adenitis	Rheu- matism.	Album- inuria.	Neph- ritis.	Cardiac.	Other Compli- cations.
All Cases	12.8	7.1	7.4	0.4	1.3	0.6	1.7	2.4
Antitoxin Cases	15.2	4.9	9.5	0.8	1.6	0.4	2.0	3.3
Non-Anti- toxin Cases	11.9	7.9	6.7	0.3	1.2	0.7	1.6	2.0

Oto-Rhinologist to the Hospital (Mr. W. Frank Wilson), in the treatment and supervision of scarlet fever cases complicated by otorrhœa or rhinorrhœa has been continued along lines developed in recent years.

The incidence of these complications was high and showed a slight increase on last year. 314 cases occurred in 929 admissions—a complication rate of 33.8 per cent., as contrasted with 24.1 per cent. in the previous year.

The distribution of these cases according to whether or not they were treated with scarlet fever antitoxin, and their respective

stay in hospital, are shown in the following table :-

posymbolini clinins qui ni si Maniquina se bad su dui s	day andrasign day been wha	Number of Cases.	Average Stay in Hospital (days).
Non-Antitoxin Cases	Rhinorrhœa	82	43.6
	Otorrhœa	54 37	52.9
Antitoxin Cases	Rhinorrhœa		50.6
	Otorrhœa	12	75.5
Company of the contract of the	Total	185	49.6

The average stay per patient of cases in this group was 49.6 days, as contrasted with the figure given for 1935, namely, 48.7 days.

In the treatment of these patients it was found necessary to perform sixty-one operations—thirty-two for the removal of tonsils and adenoids, and twenty-nine for mastoidectomy.

Subsequent Progress.—As in previous years, supervision of cases of rhinorrhœa and otorrhœa has been maintained wherever possible after their discharge from hospital, and 245 cases of this type have been visited at varying intervals. The result of these visits showed that amongst 119 cases of rhinorrhœa, three, or 2.5 per cent., still had slight nasal discharge, whilst ten or 15.1 per cent., of 66 cases of otorrhoea had slight persisting deafness or discharge from the ear.

All the cases in which the nasal or aural discharge has persisted have been kept under observation by Mr. Frank Wilson at the Out-Patient Department of the Royal Victoria Infirmary.

"Return" Cases.—The year's total admissions of scarlet fever cases, which numbered 928, produced 34 "Return" cases, a percentage of 3.6. These arose from 31 "Infecting" cases, a percentage of 3.3.

SEASONAL OCCURRENCE.

Overson	Total Scarlet	11	Infecting '' Cases.	" Return " Cases.		
Quarter.	Fever Admissions.	No.	Percentage.	No.	Percentage.	
January to March	312	13	4.1	14	4.5	
April to June	166	3	1.8	4	2.4	
July to September	166	3	1.8	3	1.8	
October to December	284	12	4.2	13	4.6	

Of the 31 "Infecting" cases (a) 13 had no complications or discharges whilst in hospital, and remained "clean" after reaching home, (b) 1 had no complications whilst in hospital, but developed discharges after reaching home, while (c) 17 had complications whilst in hospital, but were "clean" on discharge.

The figure of 3.3 for the percentage of "Infecting" cases is slightly lower than that for last year, and is a distinct improvement on the figures for the three preceding years. It was, however, to be expected that, with the lighter incidence of scarlet fever in the area, the return case rate would fall.

V	Total Scarlet Fever	"	Infecting '' Cases.	" Return " Cases.		
YEARS.	Admitted.	No.	Percentage.	No.	Percentage.	
1906-10	2,203	63	2.8	82	3.7	
1911-15	5,185	217	4.2	251	4.8	
1916-20	3.202	104	3.2	112	3.5	
1921-25	3,850	93	2.4	105	2.7	
1926-30	3,160	111	3.5	110	3.5	
1931	989	37	3.7	39	3.9	
1932	1,120	49	4.4	56	5.0	
1933	1,934	96	5.0	107	5.5	
1934	1,669	86	5.1	94	5.6	
1935	1,236	48	3.8	52	4.2	
1936	928	31	3.3	34	3.6	

"RETURN" CASES FOR YEARS 1906-1936.

ERYSIPELAS.

Of recent years erysipelas has shown a tendency to become one of the commoner and severer infectious diseases prevailing in the City. Its incidence and mortality approximate roughly to those of diphtheria, with the notable exception that while the latter is a disease of children and young people, erysipelas principally attacks the middle-aged and elderly.

In the following table the number of notifications of erysipelas, the deaths caused by the disease, and the case mortality rate are detailed for the years 1926-1936. In addition, similar information is given for such of these cases as were admitted to the City Hospital, together with the duration of their stay in hospital.

		4 1600			CITY H	OSPITAL.	
YEAR.	Total Notifica- tions.	Deaths.	Mor- tality Rate. Per cent.	Admis- sions.	Deaths.	Mor- tality Rate. Per cent.	Dura- tion of stay in Hospital (days).
1936	176	12	6.8	80	9	11.2	18.3
1935	239	15	6.3	127	20	15.7	13.1
1934	240	16	6.6	126	23	18.2	14.2
1933	264	12	4.5	116	15	12.9	17.4
1932	205	13	6.4	100	11	11.0	14.6
1931	218	11	5.0	91	4	4.4	14.0
1930	208	12	5.8	107	11	10.3	11.3
1929	220	11	5.0	85	8	9.4	13.0
1928	234	19	8.1	49	6	12.2	12.6
1927	212	12	5.7	51	2	3.9	14.5
1926	172	5	2.9	31	2	6.5	25.6

The mortality rate for these cases remains high. In 1936 this was 11.2 per cent. for all cases of erysipelas treated in hospital—as contrasted with 15.7 per cent. in 1935 and 18.2 per cent. in 1934.

Of the 80 cases admitted to hospital, 7 were given antitoxin, and of these one died, giving a mortality rate of 14.3 per cent. Fifty-five cases received local treatment only, and of these seven died, equivalent to a mortality rate of 12.7 per cent.

The remaining eighteen cases were treated with the new pharmacological preparation "Prontosil" which has its origin from Germany. Of these 18 cases one died, a mortality rate of 5.5 per cent. This fatal case was a two-months old baby.

This new remedy has only been used during the last four months of the year, and the number of cases treated as yet is small. From clinical observations alone, however, as has been stated by other observers, it would appear that the use of "Prontosil" or its derivatives will mark an important advance in the treatment of erysipelas.

Mixed Infections.

33 patients, or 1.3 per cent., of those sent into hospital were found, on or shortly after admission, to be suffering from or

incubating two distinct infectious diseases, as follows:-

Scarlet Fever with Diphtheria	6
Scarlet Fever with Measles	2
Scarlet Fever with Rubella	3
Scarlet Fever with Varicella	4
Scarlet Fever with Pertussis	1
Scarlet Fever with Mumps	1
Scarlet Fever with Erysipelas	2
Diphtheria with Scarlet Fever	5
Diphtheria with Pertussis.	1
Typhoid Fever with Diphtheria Carrier	1
Dysentery with Varicella	1
Rubella with Varicella	1
Measles with Varicella	2
Measles with Scarlet Fever	1
Mumps with Scarlet Fever	1
Pneumonia with Diphtheria	1
	33
	17.7

Cross Infections.

During the year 48 patients, or 1.9 per cent. of the total admissions, contracted a second infection in the wards of the hospital. The details are as follows, the primary infection being stated first:—

Scarlet Fever with Diphtheria	5
Scarlet Fever with Varicella	11
Scarlet Fever with Pertussis	5
Diphtheria with Dysentery	5
Diphtheria with Scarlet Fever	17
Diphtheria with Measles	1
Diphtheria with Rubella	2
Diphtheria with Pertussis	1
Diphtheria with Varicella	1
Pneumonia with Scarlet Fever	1
Total	48

There were no deaths.

Staff Sickness.

The amount of sickness among the Nursing Staff showed an increase on the previous year.

Nursing Staff.—71 of the nursing staff were off duty owing to sickness for a total of 1,699 days. One nurse contracted tuberculosis, one diphtheria, and one developed tonsillitis and was found to be a diphtheria carrier. Six suffered from influenza, fifteen from tonsillitis, fifteen from various skin and septic conditions, and three from minor accidents. Four suffered from general diseases, such as rheumatism, and one developed appendicitis. The remainder were nursed in their own homes.

Domestic Staff.—78 were off duty through sickness for a total of 1,336 days. One developed tuberculosis, five suffered from influenza, eleven from tonsillitis, and nine from skin and septic conditions. Two developed scarlet fever, three became temporary diphtheria carriers, one developed appendicitis, two suffered from general diseases, and five from minor accidents. The remainder were nursed in their own homes.

During the year the practice of immunising the staff against scarlet fever, diphtheria, and the enteric group of fevers has been carried out as previously.

The nurse who contracted diphtheria had previously been immunised, and gave a negative Schick test. She was an example of the occasional case of diphtheria occurring in a Schick negative person. Both she and the nurse who was found to be a carrier required to have their tonsils removed before they could be rendered free from infection.

The nurse who developed tuberculosis was transferred to Barrasford Sanatorium, where she made good progress.

SMALLPOX AND ISOLATION HOSPITALS, TOWN MOOR.

Owing to the disappearance of smallpox from the neighbourhood of Newcastle upon Tyne, it was not found necessary to bring the wards of the smallpox hospital into use for that disease at any time throughout the year.

The wards of both hospitals, however, were again utilised to accommodate convalescent cases of scarlet fever and diphtheria. The wards were in use from January to May, temporary nurses and domestics being engaged to staff the hospital.

E. F. DAWSON-WALKER, M.D.,

Deputy Medical Superintendent.

City Hospital for Infectious Diseases,

Newcastle upon Tyne,

8th May, 1937.

DISINFECTION, Etc.

8,068 cases of notifiable infectious disease were inquired into by the Infectious Disease Inspectors, Health Visitors and Tuberculosis Nurses and, with the exception of measles and chickenpox, the houses or rooms connected therewith disinfected by spraying with formalin. In connection with cases of tuberculosis, 713 houses, including 950 rooms, were similarly disinfected. 804 visits were made, and disinfection was also carried out in 329 special cases.

INFECTED ARTICLES TREATED IN THE DISINFECTING APPARATUS AT THE CITY HOSPITAL FOR INFECTIOUS DISEASES, WALKER GATE.

ARTICLES F	ARTICLES FROM CITY.		ARTICLES—HOSPITAL PROPERTY		
1936.	1935.	1936.	1935.		
27,247	29,153	12,150	15,554		

5,354 articles were also disinfected at the Smallpox Hospital. The staff have thus dealt with 44,751 articles during the year.

Fluid disinfectant, in half-pint tins, was given out free on the order of the special inspectors, for home use in connection with infectious disease. Every precaution was taken to ensure that the disinfectant was properly and economically used.

DISINFECTANTS DISTRIBUTED-1936.

France	FOR INFECTIOUS DISEASES.	FOR PHTHISIS.
From	FLUID (½ pint tins.)	FLUID (½ pints.)
Health Department. Tuberculosis Dispensary Corporation Yard, Benwell	69	640
Total	70	640

BACTERIOLOGICAL EXAMINATIONS, 1936.

The following is a report of the bacteriological examinations carried out on behalf of the Health Department of the Newcastle Corporation, at the Public Health Laboratory (University of Durham College of Medicine), at Armstrong College, Newcastle upon Tyne.

A total of 12,579 examinations were made during the year. This is a slight increase on the previous year, when 12,071 was the total, while in 1934 the total was 9,757.

The nature of the investigations and the results obtained are given under the various sections and are as follows:—

BACTERIOLOGICAL EXAMINATIONS :-

	Swab for Diphtheria.		Sputum for Tubercle bacilli (microscopically).		Swabs for Hæmolytic Streptococci.	
Taylor !	Total.	Positive.	Total.	Positive.	Total.	Positive
Number of Examinations	5,341	552	595	77	55	33
Percentage Positive	10.33		12.94		60.0	

The number of swabs examined for diphtheria bacilli has continued to be heavy and shows a slight increase even on 1935. The number was especially heavy in March and even in the summer months the numbers examined were in marked excess of previous years.

AGGLUTINATION REACTIONS :-

i. Enteric Fever.

A total of 66 bloods were examined for Widal Reactions against B. typhosus, B. paratyphosus—A. and B. paratyphosus—B., 11 were positive and 3 were suspicious with B. typhosus "H" emulsions,

7 were positive and one was suspicious with B. paratyphosus—B. "H" emulsions,

1 gave a doubtful reaction with B. paratyphosus—A. "H" emulsion,

38 were negative to all "H" emulsions and 5 negative to B. typhosus "H" only,

1 was positive, 1 suspicious and 2 were negative to B. typhosus "O" emulsions.

ii. Abortus Fever.

3 bloods were examined for agglutination to Brucella abortus and Brucella melitensis, all were negative.

MILK EXAMINATIONS :-

i. For tubercle bacilli by animal inoculation :-

A total of 376 milks were inoculated and 10 positives were obtained = 2.66%.

ii. Routine milks for examination for B. coli and total count.

When B, coli is taken as indicator the results were :-

B. coli not found in B. coli found in B. coli ,, B. coli ,,				$ \begin{bmatrix} 59 \\ 34 \\ 34 \\ 13 \end{bmatrix} $ 140
B. coli found in B. coli ,, B. coli ,,	0.001 ml. but not in less			11) 17 11) 39
		Tota	1	179

Thus, 140 were considered satisfactory and 39 unsatisfactory. With these samples a total count of the number of organisms able to grow on agar medium at 37°C. was also made. Taking 200,000 organisms per ml. as the standard (as in Grade "A" milk) it was found that,

119 of the samples gave totals below 200,000 and were thus *satisfactory* and 60 of the samples were above 200,000 and were *unsatisfactory*.

iii. Graded Milks.

142 samples of "Graded Milks" were examined up to the end of May in accordance with the scheme of the Ministry of Health under the Milk and Dairies (Amendment) Act, 1922, and Milk (Special Designations) Order, 1923.

The following results were obtained from January to the end of May:—

Certified Milk			Failed to satisfy test.
Grade "A" (T.T.)	71	67	4
Grade "A" Pasteurised Milk		14	3
Totals	142	133	9
	Marie Service	NAMES AND ADDRESS OF	Person mining

After 1st June, 1936, the new Milk (Special Designations) Order, 1936, started to operate, although the new method of testing did not come into force until 1st January, 1937.

Under the new Order 202 samples were examined from 1st June to the end of December, the results being as follows:—

	Total.		Failed to satisfy test.
Ministry samples, T.T	. 41	36	5
Tuberculin Tested		75	24
Accredited	. 36	31	5
Pasteurised		15	11
Totals	. 202	157	45
	-	and the same of	-

WATER EXAMINATIONS :-

 Routine samples gave the following results in a total of 184 examinations:—

Class I. Class II.				ml. or less but not in		
Class III. Class IV.	 "	10	ml. ml.	"		20 6
					7 172 11	184

ii. During the months of February and September, 31 samples of water from various Swimming Baths in the City were examined. Detailed reports were sent and the following is a summary of the results, taking only the presence of B. coli as the indicator:—

Class I.	B. coli	not foun	d in	100	ml. or less		10
Class II.		found in	100	ml.	but not in l	ess ,	10
Class III.	,,	**	10	ml.	,,		5
Class IV.	11	,,	1	ml.	,,		4
Class V.		,,,	0.1	ml.	,,	*******	2
						-	
							31
						-	-

iii. In addition some special waters were examined and detailed reports were returned:—

During August samples of water from taps	in		
Sutton's Dwellings			2
During October samples of water from the			
Northern Counties' Orphanage			2
Grammar School Baths			2
and samples of water (before and after chloris	nation)	from	
Barrasford Sanatorium			2
			-
Total	Los mil		8

VENEREAL DISEASES :-

	Total.	Serological reactions.	Microscopical examinations.
From Treatment Centres	1,170	1,170	
From Institutions	1,384	1,374	10
From Private Practitioners	563	382	181
Total	3,117	2,926	191

OTHER EXAMINATIONS :-

(a) Diphtheria.—The typing of diphtheria bacilli according to the types of Anderson, McLeod and others and intra-dermal virulence tests have been continued. Subcutaneous virulence tests have also been done on a number of cultures mostly from convalescents. The following results were obtained during the year:—

VARIETY OF ORGANISMS ISOLATED.

	1	Diphthe	ria types		В.	Saccharose	No diphtheria
Total.	Gravis.	Mitis.	Inter- mediate	Atypi- cal.	Hof- manni.	fermenters.	bacilli isolated.
795	206	303	110	74	36	8	58
		Tota	1693				Mala

VIRULENCE TESTS.

Intra	dermal.	Subcut	aneous.
Positive.	Negative.	Positive.	Negative
641	56	140	27
Total	697	Total	167

The total of 56 negative intradermal tests were obtained with the following types:—Gravis 3, Mitis 20, Intermediate 1, Atypical 26, B. hofmanni and Saccharose fermenters 3 each.

(b) Enteric Fevers.—The following specimens of fæces were received and examined for organisms of the enteric group:—

From the City Infectious Diseases Hospital	pecimens.	Positive.
From the City Health Department From the Newcastle General Hospital		5 1
Total	82	34

From this total of 82 specimens, 34 positive results were obtained:—

- B. typhosus being isolated ... 12 times
- B. paratyphosus—B being isolated 22 ,,

Various non-pathogenic non-lactose fermenting organisms such as B. paracolon, B. Morgan No. 1, B. proteus and B. pyocyaneus were also isolated a number of times.

Specimens of Urine were also similarly examined: -

From the City Infectious Diseases Hospital. 26.

From these B. paratyphosus—B. was isolated once, the remaining 25 specimens were negative.

(c) Bacillary Dysentery and Food Poisoning.—The examination of fæces for dysentery bacilli and organisms of the food poisoning group has been continued, a total of 325 specimens being submitted as follows:—

From the City Infectious Diseases Hospital From the City Health Department From the Newcastle General Hospital	12	Positive. 45 6 23
Totals	325	74
	Transport Contract	Street Section

From the 74 giving positive results the following organisms were isolated:—

B. dysenteriæ Flexner	23	times.	
B. dysenteriæ Sonne	18	33	
B. dysenteriæ Newcastle	6	.,,	
Salmonella ærtrycke		22	
Group phase salmonella		11	
Salmonella newport		,,	
Atypical non-lactose fermenters	5	- 22	

The Flexner dysentery bacilli isolated were of the following types:—

Type V. 0, W. 0, X. 10, XZ. 2, Y. 0, Z. 11. Total = 23.

The numbers of dysentery bacilli isolated this year are much less than in previous years, but the number of Salmonella group organisms is increased. The group phase Salmonella was isolated from patients connected with an outbreak of "Food Poisoning" which occurred at Sedgefield, Co. Durham, at the end of August, 1936. The organism isolated was in group phase and it was later identified as Salmonella Thompson (Scott), although it had shown best agglutination with a B. paratyphosus—C serum.

(d) Cerebro-spinal Fluids.

(i) A total of 28 cerebro-spinal fluids were received for bacteriological examination; detailed reports were returned and the following is a summary of the results:—

(1) Tuberculous	4
(2) Meningitis with inconclusive indication of in-	
fecting organism	12
(3) Streptococcal	1
(4) No definite indication of meningitis	11
Total	28
	Name of Street

(ii) 21 specimens of cerebro-spinal fluids from cases of suspected C.S. Meningitis were received for cultivation and typing, Results were as follows:—

(1) Group I.	6
(2) Group II.	0
(3) Atypical Gram-negative cocci (not agglutinating with Group I. or II. serum)	3
(4) No meningococci grown (sterile)	4
(5) No meningococci isolated (cultures contaminated)	8
Total	21

(e) Typing of Pneumococci.

This was started in February, typing being done with peritoneal exudate after mouse inoculation by (1) rapid method on a slide and also by (2) macroscopic method with a young broth culture, only Type I., II. and III. sera being employed.

A total of 18 sputa were examined and practically none of these were from Acute Lobar Pneumonia during the first 3 days. Mostly, both methods gave similar results.

> 3 were Type II. by both methods, 10 belonged to Group IV. by both methods, 2 were negative or very scanty Pneumococci present but 3 gave variations:—

Rapid.	Macroscopic.
1. Group IV.	Type II. atypical.
2. Type I. and II.	Type II. ,,
3. Type I. and II.	Type I.

(f) Miscellaneous Examinations. Direct examinations of throat swabs for diphtheria bacilli or Vincent's angina 13 From the City Health Department. (a) P.M. specimens from a suspected case of Foodpoisoning (small intestine, colon, spleen) (b) Tomato Sauce? Food-poisoning organisms 1 (c) Blood agglutination test for dysentery 1 From the City Hospital, Walker Gate. Urines for bacteriological examination Urines for guinea-pig inoculation Blood cultures for organisms..... Blood for agglutination test for dysentery 2 Smear for B. anthracis 1 Culture from appendix wound? diphtheria bacilli 1 Total 33

- (g) Measles Serum.—One batch was put up during March.
- (h) Newcastle General Hospital.—A number of bacteriological examinations for the Hospital have been carried out and the following is a summary:—

CHARACTER OF EXAMINATION.	
Autogenous Vaccines	7
Blood cultures for organisms	
Blood agglutination test for dysentery	
Catgut for sterility	1
Cerebro-spinal fluids for bacteriological examinations	8
Fæces for tubercle bacilli	9
,, parasites	9
" general bacteriological examination	3
Fluid from liver abscess and scrapings from cyst wall for amœbæ	3
Leptospira:—Bile	1
Blood	1
Urine	2
Pleural fluid or empyema pus for organisms	16
Pus from various sources	12
Sputum for organisms	3
Swabs (throat) for organisms	2
Urines for bacteriological examination	14
" microscopical "	4
,, animal inoculation for tubercle bacilli	5
Total	108

The following table gives a complete summary of the various examinations, including the year 1935 for comparison:—

Nature of Investigation.	1935.	1936.
Throat swabs for B. Diphtheriae	5,114	5,341
Sputa for Tubercle bacilli	571	595
Swabs for Hæmolytic Streptococci	45	35
Against the Enteric Fevers	65	66
Against Brucella Abortus	3	3
For the Tubercle Bacillus	347	376
For Bacillus Coli and Count	154	179
Graded Milk	401	1142
Water Examinations :—	401	(202
For Bacillus Coli	196	184
For complete examination	40	39
Venereal Diseases	2,919	3,117
Other Examinations :—	-,0-10	
(a) Diphtheria—Type of bacilli	692	795
Virulence tests—(i) intradermal	700	697
(ii) subcutaneous	136	167
(b) Enteric Fevers—(i) Fæces	98	82
(ii) Urine	25	26
(c) Bacillary Dysentery and Food-poisoning		
group	425	325
(d) Meningitis (various) C.S. Fluid	41	49
(e) Pneumococcal Typing	_	18
(f) Miscellaneous	51	33
(g) Newcastle General Hospital	48	108
Total	12,071	12,579

(Signed) S. H. WARREN, M.R.C.S. (Eng.), D.P.H. (Lon.),

Director, Public Health Laboratory.

University of Durham College of Medicine, Newcastle upon Tyne, 4th March, 1937. REPORTS OF THE
TUBERCULOSIS MEDICAL OFFICER
AND
MEDICAL SUPERINTENDENT,
BARRASFORD SANATORIUM.

IV.—TUBERCULOSIS.

TUBERCULOSIS DISPENSARY, BARRASFORD SANATORIUM. PIRO HIDRIRITE VI

TURESCULDSIS DISPENSANT, BARRASTORD BARATORIUM.

TUBERCULOSIS.

Report of the Tuberculosis Medical Officer.

TO THE MEDICAL OFFICER OF HEALTH.

SIR,

I beg to submit, herewith, my report for the year 1936.

The general work of the department has been on similar lines to that of 1935 and there have been no changes in the Dispensary staff or weekly number of clinics held. Certain improvements have been made, however, with regard to the examination of contacts and more time has been devoted to this work and the consideration of the home and family of the patient, the details of which are set out in the appropriate paragraphs in this report.

The co-operation between the Tuberculosis Dispensary and the doctors in the City has always been very good, but has been further improved by more frequent correspondence with regard to patients and contacts. Doctors have also been loaned x-ray films of their patients in cases where there was difficulty in diagnosis or some point of special interest. This service has been greatly appreciated.

The X-ray facilities in the City are good; patients living in the West being X-rayed at Newcastle General Hospital and in the East at the Sanatorium Pavilions, City Hospital for Infectious Diseases, Walker Gate, but these arrangements mean additional journeys and inconvenience for patients. To make the Tuberculosis Scheme thoroughly efficient an X-ray apparatus should be installed in the Tuberculosis Dispensary. Difficulties exist in the present building, which is old and too small. During the year the Dispensary was visited by several members of Committee and plans for a new building were drawn up by the City Architect, from which an approximate cost was estimated. This has been sent to the Commissioner for Special Areas to ascertain if a grant towards the cost can be obtained.

I am indebted to the Health Committee for sending me away for one week during the year to visit five different Local Authorities to see certain points in their dispensary organisations and sanatoria.

The Tuberculosis Dispensary. 447 clinics have been held at the Dispensary at each of which an average of 19 patients was seen, of these 5 were stripped and carefully physically examined, and in addition many minor examinations were made and recorded.

Attendances at the Dispensary.—2,949 persons attended the Dispensary during the year or were visited in their homes, registering 8,623 attendances, and 2,334 complete physical examinations were made. 1,710 cases attended for the first time; of these 781 were sent by general practitioners, 282 by the Dispensary Visiting Nurses, 104 by the Newcastle-upon-Tyne Dispensary, 23 by the Royal Victoria Infirmary, 23 by the School Medical Officer, 76 by the staff of the Newcastle General Hospital, 30 by the Maternity and Child Welfare Centres, 39 came of their own accord under special circumstances, 313 were sent by the Tuberculosis Dispensary Medical Staff and smaller numbers from other sources.

Of the 1,710 new cases, 446 had lived with patients known to have tubercle bacilli in their sputum, 232 with tuberculosis patients who had not tubercle bacilli in their sputum, and 47 were contacts of persons who had died from pulmonary tuberculosis.

Table 1 gives details of the New Cases examined (excluding contacts), and Table 2 gives details of the recommended contacts as per Memo. 37/T (revised). I am pleased to report that the latter figures have increased from 447 in 1935.

New Cases Examined (excluding Contacts), during the Year 1936. (First Schedule, Part A., Memo. 37/T., Revised).

Diagnosis	Ma	les.	Fem	ales.	Totals
Diagnosis.	Over 15 yrs.	Under 15 yrs.	Over 15 yrs.	Under 15 yrs.	Totals
Pulmonary Tuberculosis Non-Pulmonary Tuberculosis Diagnosis not completed Non-Tuberculosis	169 22 19 212	12 25 9 110	128 18 10 261	19 18 4 76	328 83 42 659
TOTALS	422	156	417	117	1,112

Contacts.—A complete list of names and ages of the contacts of each tuberculous case is kept in the patient's dossier and every effort is made by the medical and nursing staff of the Dispensary to have all these people examined and X-rayed. The names of any that are children below the age of 5 years are sent at weekly intervals to the Maternity and Child Welfare Medical Officer and those between the ages of 5 and 14 years to the School Medical Officer. These two Medical Officers are co-operating in this

114A

MEMO, 37/T. REVISED. SCHEDULE III. PULMONARY TUBERCULOSIS.

ANNELA RETURE SHOWING IN SUMMARY FORE (a) THE CONDITION AT THE RED OF 1938 OF ALL PAULETT REMAINING ON THE DISPERSANY.

REGISTER: AND (6) THE SEASON FOR THE REMOVAL OF ALL CARM WRITTERS OF THE REMOVAL PROPRIES ACCORDING TO
THE YEARS IN WHICH THE PATISHIN WERE FERST ENTERED ON THE DISPERSANY REGISTER AS DEPOSITE CARES OF PULMONARY
TERRECULORS, AND THEIR CAMPAGEARY REGISTER AS DEPOSITE CARES OF PULMONARY

	Pre	vious.	to 192	8.		19	26.			19	V27.			- 1	1928.		1		1929.				1930.				1931.		-1-		1932.				1933.				1934.		1		1935				1936	
Condition at the time of the	4	Class T	i.B. pl	MR.		Class	T.B.	plus.	1	Class	T.B.	plus.	1	Cla	es T.I	i, plus.	1	C	ass T.I	B. plus		, a	lass T.	B. plus		Cla	os T.B	plus.	1	Ca	iss T.II	i, plus.	1	Cla	as T.I	, plus.	- 6	Ch	us T.1	3. plus		Ch	MA T.	B. plu		5 0	lass T	.13. p
last Record made during the year to which the Return relates,	Cles T.R. selv	Crosp J.	Group 3.	Total (Class T.R. plos).	One T.S. sule	Group I.	Comp 3.	Total Klass T.R. plest	Clas T.R. sale	Group 1.	George 2.	Total (Class T.B. ples)	On T.B. nie	Greep 1.	Group #.	Group L. Total (Clear	Cles T.R. selve	Group 3.	Comp 2.	Group 3.	T.B. plot.	Group J.	Comp 2	Georg 3. Total (Class	T.R. plot.	Group L.	Group 2.	Googs A. Total (Class T.R. class	One T.B. mo	Greep I.	Group Z.	Group S. Thush (Class	One T.B. min	Group 1.	Greep Z.	Group 3. Tetal (Class	Cles T.R. 160	Greep I.	Group 2s.	Group St. Total (Class	Clear T.R. ed.	Group 1.	Group 1.	Group 3.	T.R. plot.	Grap L	Seept	Comp &
Disease Arrested— Adults—Mass Female Children	4 3	3 6	2	9 11 1	-		2 -	2 1 1			1			1	6 2 1	1	7 1		3 4 1	1	3 5 1	3	4 3	1	8 3 2	1	3 2	1 3	18 11 2		1		7 3 3				000-											
Disease not Arrested— Adults—Male . Female Children	2	8 14 5 2 3	1	111		3	2 -	2 2		1	3	3 7	1	1	7		8 9	2 2	12 7	3	12	4 1	9 14 2		15 5 16 7 3 7	1	14 6 4	4 15 3 5 - 4	8		20 7 1	12 30 7 14 1 1	20	1	20 13 2	20 4	1 31 3 31 2 30			29 4 12 3			36 18 2	32 18 2	70 7 87 8 4 3	4 1 2	39 32 1	34 29 3
Total on Dispensary Register a 31st December	15	19 37	18	74	2	4	5	. 9	1	1	5	4 10	2	3	17	1 2	1 18	8 2	27	8 :	4 1	5 4	32	9 4	15 28	2	29	8 39	61		29	20 4	69	1	35	30 6	1 98		41	44 8	8 127	3	56	52 1	11 13	6 3	72	66
Discharged as Recovered— Adults—Male : Female :	00 28 22	10 24	2	98 62 4	3 5 4		6 -	6 2	7 6 6		1 2 1		9 -3		3		3 3	2 2	1	2 1	8 2	67.8	3		3 1	111		1 1					=															
Lost eight of, or otherwise removed from Dispensary Register.	31	12 2	7	134	16	3 :	20	5 28	27	2	14	6 21	17	2	16	3 2	1 30	1	21	6 :	38 4	0 1	15	9 2	18 32	1	11	8 20	28		11	16 22	14	1	6	8 1	31	1	8	5 1	1 11		6	12	18	5	1	4
DEAD—Adults—Male Female Children	23 13 3		12		9 7 4	4	50 40 34 5 2	0 93 1 89 8 10	13 13	3		50 91 80 58 7 7	7 2 2	1 2	68 27 6	34 10 26 5 2	3 15 5 11 8 4	2	42 47 9	51 1 39 3 3 1	(3 1) (8 1 12	9 1 1 1 1		79 13 37 6 5	0 22 3 12 7 3	E	24	97 131 82 76 3 8	10.		18 24 1	56. 80			17 20 2	64 8 54 7 6 8	1 11 8		14	70 7 41 5	5 11			31 3 33 4 3		7 -	8 2	22 16 1
Total written off Dispensary Register	180	83 153	60	730	48	10 1	14 10	4 228	72	6	83 1	3 182	41	8	121	65 19	1 81	3	124	102 2	1 9	2 4	97	130 23	1 79	1	75 1	61 237	82		54 1	41 190	27	1	45 1	12 171	5 55	1	31 1	20 18	2 40		22	79 11	1 2		-9	43
GRAND TOTALS	195 1	92 190	78	804	50	14 1	19 10	4 237	73	7	88 9	7 193	43	8	138	66 21	2 96	7	151	107 2	10	7 8	129	139 27	6 107	3	104 1	10 276	116	-	83 1	61 244	95	2	80 1	92 244	133	1	72 1	64, 23	7 173	3	78	131 2	2 18	1 3	81	109

MEMO. 37/T. REVISED. SCHEDULE III.—Continued. NON-PULMONARY TUBERCULOSIS.

Annual Return servings of numberay from (a) the condition at the ring of 1006 of all Pathents remaining on the Despensary Register; and (a) the reasons for the benoval of all cases written opt the Register.

		Pre	vious 1	to 192	6.		16	k26.	_	1		1927.				1928.		-1		1921				193	0.			190	11.			193	2.			190	3.			190	34.				1935.			1	1906	
	Condition at the time of the last Record made during the year to which the Return relates.	Bone std. Juste,	Abdessites Other Organia	Projeteral Clauds	Total	Bear and Joint,	Abdominal	Other Organic	Charle	Boos and later,	Althoused	Other Organi.	Charle of the Charles	Bon sat	Abdominal	Other Copania	Periphend Glands.	Total	Just.	Other Organic	Peripheral Clearly	Total	Rose int	Abdressinal Other Organs,	Perplema	Total	Bears and Joseph.	Abdominal Other Option	Perphend	Total	Boars and Justs,	Other Organi.	Prophent Clash,	Total.	Brees and Joseph and	Other Owner	Pertphenal Classics,	Total	Peace and Joseph	Abbasing	Postpheni	Gleeds, Treat.	Rese tod Jon a	Abdominal	Other Organs,	Complete	Brass and	Abd minet	Other Organic	Fresh
on Dispense	Disease Aerested— Adults—Male Fernale. Children	1 2	1 1	111	1 4		-			-	1		2 1						2 1	-	2 1	2 5	1 - 2	1 =	1	2 3	3	2 1	8	15	2 1 2	-	1 3	3 1 6	1 -2	1 1	7	1 1 11	2	2 -	1 1000	4 2 5		1		3	1	-		
rmaining y Register	Disease not Arrested— Adults—Male Female Children	2 .	- 2 - 1 - 4	1 3	5 1 16	- 3		1 -	1 4	1 3			1 1	3 4	-		2	3 -	1 8 1		111	1 9	1 2 4	1 1 1 2	1 4	2 5 10	2 1 9	1 -	1 7	3 2 22	2 2 10	3	1 2	6 4 16	5 2 14	2 -	1 1 5	8 5 31	237	3 1	2 2	6 6 35	7 6 12	2 2 15	127	1 1 5 1 25 5	7 11	9 2 11	1 2 2	20 44
3 3	Total on Dispensary Register at 31st December.	14	1 8	4	27	3	1	2 _	- 6	3	1		3 5	7	1	1	2	11 1	1 3	-	3	17	10	4 2	6	22	15	8 3	16	42	19	7 3	7	35	24 1	5 4	14	57	15	7 3	3 21	58	28	20	4 3	14 S	25	23	4 30	85.
	Transferred to Pulmonary	4	1 _	5	10		1		1 2	1	1		3 8	2	1	1	1	8 -	_ 3	-		3	2	2 3		7	2	1 3	1	7		2	1	3		2 2	1	4						1		1 3				
wensary na for	Discharged as Received— Adults—Male Female. Children	2 3 8	1 1 1 8 2	3 3 18	22 7	2 8	2 6	1 -	5 4 4 6 21	1 1 7	5		3 4 3 4 7 20	1 -6	1	1	1 8	4 1 15	1 2 2 3 4		2	- 6	4 - 6 1		4	6		1 1	2 1 5	2 3 10	1 2	2 -	1 2 10	3 5 15			1 22	1 1 4												
on Dis	Lost sight of, or otherwise removed from Dispensary Register	22	1 3	20	46	16	2	2 2	2 42	8	8	2 1	6 44	11	5	3	26	45 1	5 8	3	18	44	12	9 3	15	39	7	7 6	14	34	13 1	7	12	42	12	7 3	9	31	3	6	9	19	2	3	1	4 1	2			
Not now gister a	DEAD—Adults—Male Female Children	1 -4 :	1 2	3	16 7 8	3	20.02	1 -	1 5 2 5	2	1 1	3 1 1	2 8000	-3	2 3	2		2 4 6	1 3 2 1	1	1	3 6 3	3 1 5	1 1 1 5 1	1	6 4 12	3	1 1 4 1 3 3	2	5 6 8	5 2 2	1 1	- 2	8 3 9	3	1 4	1 1	10 7 6	=	1 1		2 2	4	1 2 2	2 -4	1 1	3	1	3 3	1 2 5
(8) (8)	Total written off Dispensary Register.	40 1	1 10	49	188	31	14	6 3	3 84	19	16	8 4	1 84	21	15	6	35	77 2	5 18	5	28	76	33 2	6 9	28	96	15	17 12	24	68	25 2	1 12	27	85	23 1	8	16	60	4	7 3	9	23	7	7	7	6 23	3	2	3 2	35
	Gramp Toraza of (a) and (b) (excluding those transferred to Pulmonary)	54 1	2 18	53	215	34	15	8 3	3 90	24	17	8 4	14 90	28	16	7	37 1	88 3	6 21	5	31	93	43 3	0 11	34	118	30	25 15	40	110	44 2	8 15	34	21	47 2	8 12	30	117	19	24 2	5 30	81	33	27	11 4	10 111	28	25	2 35	95

work. The Maternity and Child Welfare Medical Officer keeps a separate card index of these children and supplies them with emulsion. The School Medical Officer pays special attention to them during routine examinations, and in both Departments if any of them are thought to be delicate they are referred to the Tuberculosis Dispensary for expert opinion.

In every case where a death from tuberculosis has occurred in the City the consent of the General Practitioner, who attended the deceased, is first obtained and then the remainder of the family are seen or written to and urged to be examined in case any of them may be developing tuberculosis. In this way a number of very early cases of tuberculosis have been found and many contacts seen.

As in previous years contacts have been examined and x-rayed by the Tuberculosis Medical Officer during visiting hours on Wednesday afternoons at the Sanatorium Pavilions, City Hospital for Infectious Diseases, Walker Gate.

The details of the contacts seen, which number 598, are set out in the following table.

169 were seen during domiciliary visiting and 55 at the Sanatorium Pavilions, City Hospital, Walker Gate, on relatives' Visiting Afternoons.

Contacts Examined during the Year 1936. (First Schedule, Part A., Memo. 37/T., Revised).

Diagnosis.	Ma	les.	Fem	ales.	7P-4-1-
Diagnosis.	Over 15 yrs.	Under 15 yrs.	Over 15 yrs.	Under 15 yrs.	Totals
Pulmonary Tuberculosis Non-Pulmonary Tuberculosis Diagnosis not completed Non-Tuberculosis	18 2 73	2 1 7 130	14 4 6 199	5 4 6 127	39 9 21 529
TOTALS	93	140	223	142	598

221 cases, who had been seen previously and discontinued, returned for re-examination. The details are given below:—

Cases Discontinued in Previous Years, and Returned during the Year 1936 (Included in Previous Tables 1 and 2.)

Diagnosia	Ma	les.	Fem	ales.	Totals
Diagnosis.	Over 15 yrs.	Under 15 yrs.	Over 15 yrs.	Under 15 yrs.	Totals
Pulmonary Tuberculosis Non-Pulmonary Tuberculosis Diagnosis not completed Non-Tuberculosis	19 2 5 39	32	15 2 2 2 78	2 1 24	36 4 8 173
TOTALS	65	32	97	27	221

The following table gives further details of patients and cases who attended the Dispensary or were visited in their homes:—

Type of the transfer over	the Dis	spensary (or	nd Cases wh were visited the Year 19	in their
	Total.	Males.	Females.	Under 15 years of age.
"Sputum Positive Cases"	743	435	294	14
"Negative Cases"	2,206	643	758	805
Totals	2,949	1,078	1,052	819

Cases and patients written off the Dispensary register during the year.

Cases and Patients written off the Dispensary Register during the Year 1936.
(First Schedule, Part A., Memo. 37/T., Revised).

Drugwegg	Mai	ES.	FEMA	LES.	Tomiro
Diagnosis.	Over 15 yrs.	Under 15 yrs.	Over 15 yrs.	Under 15 yrs.	TOTALS
Pulmonary Tuberculosis, Recovered	22	1	16		39
Non-Pulm. Tuberculosis, Recovered	16	5	18	4	43
Non-Tuberculosis	313	244	474	219	1,250
Left district, lost sight of, or will not attend Dispensary	80	20	67	11	178
Totals	431	270	575	234	1,510

At the end of the year the number of patients and cases on the Dispensary register was 1,797. These are tabulated below:—

Number of Cases and Patients on Dispensary Register at end of Year 1936. (First Schedule, Part A., Memo. 37/T., Revised).

Diagnosis.	MA	LES.	FEMA	LES.	TOTAL
DIAGNOSIS.	Over 15 yrs.	Under 15 yrs.	Over 15 yrs.	Under 15 yrs.	TOTAL
Pulmonary Tuberculosis (T.B. in Sputum)	404	4	267	9	684
T.B. in Sputum)	268	61 130	206 105	60 106	595 454
Diagnosis not completed	21	16	16	11	64
Totals	806	211	594	186	1,797

The two tables (pages 114A and 114B) are self-explanatory and are required by the Minister of Health under Memo. 37/T (revised).

Domiciliary Visits.—During the year I made 382 domiciliary visits to 213 patients and 169 contacts. These are included in "Attendances at Dispensary."

Relations with other Departments and Hospitals.—The closest co-operation has existed between the Tuberculosis Dispensary and all the various departments of the Health Department. Constant interchange of information and patients has taken place between the School Medical Service and the Tuberculosis Dispensary. The relation between it and the other Medical Services in the City has, as formerly, been most satisfactory.

Voluntary Tuberculosis Care Council.—In November, 1936, Mr. Girling, Joint Hon. Sec., resigned. The work of the Council has gone on with as little interruption as possible, but at the time of writing its reconstruction is under consideration. During 1936, 136 patients were referred to the Voluntary Tuberculosis Care Council for consideration. The assistance given was as follows:—

	Number of Patients.
Loan of bed and bedding	37
Total number of beds and bedding on loan	147
Milk	
Outfits of clothing	
Eggs	
Surgical Appliances	15
Other assistance	
Referred for adoption under "Jubilee Scheme"	71

Nurses' Visits to Patients.—793 new patients were seen and 10,104 subsequent visits made. A total of 1,708 were upon their lists on December 31st, 1936, comprising of 762 males, 566 females and 380 children. As there are four nurses constantly visiting in the City, each has approximately 427 patients upon her books. The infectious cases of pulmonary tuberculosis were visited once a month. The others at longer intervals.

X-ray Examinations.—If the diagnosis of a case is uncertain an X-ray examination gives valuable assistance. This is carried out either at the Sanatorium Pavilions, City Hospital, Walker Gate, or at Newcastle General Hospital, and now with the improved facilities that exist, every new patient is X-rayed except where peculiar circumstances are present. During the year a total of 1,637 films were taken in connection with Dispensary patients, 991 at the City Hospital, Walker Gate, and 646 at Newcastle General Hospital.

Sputum Examinations.—An effort is made to examine the sputum from every possible case and in many instances repeated examinations are necessary. 1,223 specimens of sputum were examined at the Dispensary, of these 240 contained tubercle bacilli. In addition 595 sputum examinations were carried out at the University of Durham Bacteriological Laboratory in the City, to which medical practitioners may send specimens. 77 of these were positive.

The Sanitary Inspector.—Disinfection has been carried out by the Sanitary Inspector as necessary in houses after a death, or change of address of a person suffering from pulmonary tuberculosis. Bedding and clothing have been removed and disinfected also and attention has been given to overcrowding and sanitary defects.

Notifications.—622 notifications were received during the year, but some were duplicates, so that the total number of new cases was 584, of whom 449 were certified to be suffering from "pulmonary" and 135 from "non-pulmonary" tuberculosis. This is the lowest number of notifications ever received in a year in the City.

The details as regards sex and age are given in the accompanying table:—

Summary of Notifications during the Period, 1st January to 31st December, 1936.

(The Public Health (Tuberculosis) Regulations, 1930.)

					P	rimary	Notifi	ication	s.				Total Notifications (including
Age Periods.	0 to 1.	1 to 5.	5 to 10.	10 to 15.	15 to 20.	20 to 25.	25 to 35.	35 to 45.	45 to 55.	55 to 65.	65 and up- wards.	TOTAL.	Cases previously notified by other doctors).
Pulmonary—													
Males	****	8	9	8	27	29	51	40	36	31	8	247	263
Females		8	12	8	30	45	42	34	11	8	4	202	214
Non-Pulmonary													
Males	1	17	11	11	9	5	5	4	****	2	2	67	71
Females		14	10	11	5	5	6	7	2	5	3	68	74
TOTALS	1	47	42	38	71	84	104	85	49	46	17	584	622

As far as possible every notified case is visited by the nurses and urged to visit the Dispensary for examination and classification with a view to treatment.

Of the 584 cases notified, 497 attended the Dispensary and 32 others were visited in their homes by the Health Visitors in the course of the year. The names of the patients certified to have died from tuberculosis, but not previously notified, are entered in the notification register, so that if the 29 patients in this category, and 21 who died within one week of notification and were not known to the Dispensary be deducted, it will be seen that the Dispensary gets into touch with nearly all of the known cases of tuberculosis. The only cases not known to the Dispensary were five who were living in institutions or refused to be visited.

A table has been prepared to illustrate these points and also to show the nature of the institutional treatment afforded to the cases notified during 1936. 313 of the 449 patients notified as suffering from pulmonary tuberculosis were treated in beds belonging to, or controlled by the City Council, and 66 out of a total of 135 patients notified as suffering from forms of tuberculosis other than pulmonary were treated in such beds.

The number of patients dying in the year of notification is also given, and it will be seen that 146 (equal to 25 per cent.) of all the new cases died in the same year as they were notified.

NOTIFICATIONS OF TUBERCULOSIS DURING 1936.

		ry d	by ut ded ry.			ed Institution			
Part Affected.	Notifi- cations.	Attended	Visited by Nurse but not attended Dispensary.	Barras- ford Sana- torium.	Sanat- orium Pav. Walker Gate.	Stann- ington Sana- torium.	New- castle Gen. Hosp,	Totals.	Died during the Year.
Pulmonary— Male	247	218	9	52	116	4		172	64
Female Non-Pulmonary	202	173	17	39	96	6		141	46
Male	67	52	4		****	5	33	38	15
Female	68	54	4 2			3	25	28	21
TOTALS	584	497	32	91	212	18	58	379	146

Cases re-admitted to the Sanatorium Pavilions, Walker Gate, and those transferred to Barrasford Sanatorium during the year are counted as only receiving treatment on one occasion.

During the year 205 cases (35.1 per cent. of the total) were notified by the Dispensary Medical Staff.

Practitioners were written to by the Medical Officer of Health when notification appeared to have been neglected.

Public Health (Tuberculosis) Regulations, 1930.

Number of Cases of Tuberculosis remaining on the Notification Register at the end of Year.

Veen	I	ULMONARY	7.	Non-Pulmonary.			Total		
Year.	Males.	Females.	Total.	Males.	Females.	Total.	Cases		
1925	855	608	1,463	340	312	652	2,115		
1926	744	515	1,259	297	263	560	1,819		
1927	644	441	1,085	236	204	440	1,525		
1928	720	443	1,163	294	254	548	1,711		
1929	744	501	1,245	319	270	589	1,834		
1930	737	495	1,232	316	264	580	1,812		
1931	767	501	1,268	298	251	549	1,817		
1932	801	513	1,314	292	240	532	1,846		
1933	795	531	1,326	294	270	564	1,890		
1934	792	538	1,330	292	237	529	1,859		
1935	799	569	1,368	283	236	519	1,887		
1936	776	598	1,374	267	217	484	1,858		

Deaths.—There were 308 deaths from tuberculosis of New-castle-upon-Tyne residents. 265 pulmonary and 43 non-pulmonary, giving a death rate per 1,000 population—

	Number of Deaths.	Death Rate per 1,000 Population.
Pulmonary Tuberculosis	265	0.90
Non-Pulmonary	43	0.14
All forms of Tuberculosis	308	1.04

Page 37A in the Report of the Medical Officer of Health gives further particulars of deaths from tuberculosis.

Of the 265 persons who died from pulmonary tuberculosis 90.0 per cent, were known to the Dispensary staff, 220 having visited the Dispensary and an additional 18 having been attended in their homes by the visiting nurses.

41.9 per cent. of the persons who died from "non-pulmonary" tuberculosis were attended at or from the Dispensary. This is much lower than the pulmonary cases; the main reason being that 30.2 per cent. of the non-pulmonary cases were not notified before death.

Of 265 deaths from pulmonary tuberculosis the diagnosis was verified bacteriologically in 189 instances, i.e., 71.3 per cent.

Six other Dispensary patients who were known to be suffering from pulmonary tuberculosis, and in whose sputum tubercle bacilli had been found, died during the year. The cause of death being registered as: bronchitis and myocardial degeneration, 1; broncho pneumonia, 1; renal calculus, 1; miliary tuberculosis, 1; cardiac disease, 2.

Duration of Illness.—Whenever possible, in pulmonary cases, enquiry was made as to the length of time the deceased had been ill, and the average duration of illness was found to be 46.5 months. As in previous years, important differences were discovered when age and sex were considered, the figures being 54.7 months for adult males, 37.6 months for adult females and 21.4 months for those below 15 years of age (both sexes).

The period between notification and death was, as one would expect, longer in the adult males than in the adult females and children, but averaged 33.6 months for all cases. 29.4 per cent. of the patients had either not been notified prior to death (5.6 per cent.) or died within 3 months of notification (23.8 per cent.)

Further details and comparative figures for previous years are submitted in the following table:—

RETURN OF DEATHS FROM PULMONARY TUBERCULOSIS OCCURRING IN :-

	Deaths which occurred in these years,											
	Average	Average	Average	Average for	1000	1094	1935	1936.				
	1913-17.	1918-22.	1923-27.	1928-32.	1800	1934	1300	м.	F.	CHD	Tota	
Persons not notified ,, notified under 1 mth. ,, between 1 and 3 ,, ,, between 3 and 6 ,,	43 35 94 53	51 47 48 30	33 50 44 38	23 38 45 36	17 29 24 26	25 33 22 36	8 23 27 30	12 16 15 15	3 10 16 6	3 4 1	15 29 35 22	
Total under 6 months	226	183	166	140	96	116	88	58	35	8	101	
Persons notified between— 6 and 12 months 12 and 18 ,, 18 and 24 ,, 2 and 3 years	47 28 15 20 21	46 21 15 18 47	40 25 17 22 53	36 22 17 21 59	31 27 17 25 66	32 18 15 24 75	22 24 16 25 65	13 8 8 7 56	15 11 6 10 26	1 1 2	29 19 14 18 84	
Totals	357	331	324	296	262	280	240	150	103	12	265	

The figures for non-pulmonary forms of tuberculosis show that in 13 instances out of 43 deaths, the disease had not been notified prior to death; 9 of the 15 fatal unnotified cases of pulmonary tuberculosis, and 12 of the 13 fatal unnotified cases of non-pulmonary tuberculosis, died in hospitals; included in the 12 "other forms" were 6 cases of tuberculosis meningitis.

Family History.—In 98 instances amongst the 247 cases of pulmonary tuberculosis known to the Dispensary who had died during the year, *i.e.*, in 39.7 per cent., there was a history that some near relation was suffering from, or had died of pulmonary tuberculosis. The figures were 39.3 per cent. for males and 40.2 cent. for females.

House Accommodation.—The home conditions of the people are intimately associated with tuberculosis. The numbers of rooms in the dwellings occupied by the above 247 persons were as follows:—

Rooms in Dwelling.	1	2	3	4	More than 4	Common Lodging Houses.	Not Known.	Total.
Deaths	16	55	72	63	28	5	8	247

As regards the type of house occupied, 112 were flats, 55 tenements, 67 self-contained, 5 were common lodging houses, and in 8 cases the particulars were not known.

It is noteworthy that of the 229 patients suffering from pulmonary tuberculosis who attended the Dispensary and died in 1936, 210, or 91 per cent., had received institutional treatment, on one or more occasions. This is a high percentage and shows what a large proportion of the cases visiting the Dispensary avail themselves of the accommodation provided.

Tuberculosis and Pregnancy.—From the 1st January, 1924, to the 31st December, 1936, 88 women, who had had tubercle bacilli in their sputum and were attending the Dispensary, became pregnant and each gave birth to a child. Their classification, in the terms of the Ministry of Health grouping, at the time they were entered upon the Dispensary Register was:—T.B. positive, Group 1. 12; Group 2. 56; and Group 3. 20. At the time of confinement 37 were "disease arrested," none of whom have relapsed or died as a result of the confinement. 36 of the women had a family history of tuberculosis.

2 of the children were premature and died at birth and 1 was stillborn. 16 are now known to be dead and 5 were certified to be due to some form of tuberculosis.

14 of these patients cannot now be traced because they have moved into other districts. This can be understood because the investigation dates back 12 years and on this account no further useful statistics about the matter can be obtained.

From this it will be seen that there has been only a very small number of tuberculous women known to the Dispensary, who have given birth to children. In 1935, out of 4,666 births in the City only 4 were known to be complicated by tuberculosis. This figure appears to be very low, but it compares with that of the London County Council Hospitals where, out of 13,000 annual confinements, only 20 or 30 are those of tuberculous mothers (British Medical Journal, 17th October, 1936, page 766).

INSTITUTIONAL TREATMENT.

Approximately 76 beds were provided at Barrasford Sanatorium for Newcastle-upon-Tyne patients suffering from pulmonary tuberculosis and 136 hospital beds at the Sanatorium Pavilions, City Hospital for Infectious Diseases, Walker Gate; 60 beds at Newcastle General Hospital for the treatment of non-pulmonary tuberculosis, and 30 beds at Stannington Sanatorium for children for both surgical and medical cases.

Barrasford Sanatorium.—The report of the Medical Superintendent of Barrasford Sanatorium, which will be found under a separate heading, contains details and statistics of Newcastle patients treated in that Institution.

Sanatorium Pavilions, City Hospital, Walker Gate.—459 patients were admitted (274 males and 185 females) and included 57 transferred from Newcastle General Hospital who were found to be suffering from pulmonary tuberculosis.

Details of the number of patients admitted and discharged are given in the accompanying table :—

Patients who received Treatment in the Sanatorium Pavilions, Walker Gate, during the Year 1936.

		Sex	In Institu- tion on 1st January, 1936.	Ad- mitted during the Year.	Discharged during the Year.	Died in Institu- tion during the Year.	In Institu- tion on 31st Dec., 1936.
Number of Patients.	Adults Do Children Do	F. M.	53 21 3 9	210 124 8 9	176 93 10 15	54 27 1 3	33 25
Observation Cases.	Adults Do Children Do	F. M.	3 3 2	44 39 12 13	33 34 12 8	5 5 1	9 3 6
TOTALS			94	459	381	96	76

N.B.—37 patients were re-admitted and are counted as 74 admissions.

Of the 98 patients discharged who had been under observation 48 were found to be suffering from tuberculosis. The total number of days of those who received treatment was 39,562 giving an average length of stay as 89 days.

96 patients died in the Institution; the conditions of the other patients on discharge is given in the table below:—

	Males,	Females.	Total.
Improved	192	112	304
Without Improvement	39 60	38 36	304 77 96
Died in Hospital	60	36	96
Totals	191	186	477

Many of those discharged "improved" were fit for light work; 29 were transferred to Barrasford Sanatorium and 1 to Stannington Sanatorium. 13 patients were sent to the Newcastle General Hospital for surgical treatment.

Treatment has been on Sanatorium lines, modified to some extent in view of the type of patient; the essentials are the same, however, namely, rest and good food under satisfactory hygienic conditions, with exercise graduated to the patient's tolerance.

X-ray Examinations.—During the year 1,659 thoracic films were taken. These included 991 Dispensary patients, 285 inmates of the Sanatorium Pavilions, 92 patients from the City Hospital for Infectious Diseases, 228 in connection with artificial pneumothorax treatment and 63 nurses and maids belonging to the staff of the Hospital. In addition, 1,450 routine screen examinations were made, 44 to out-patients and 1,406 in connection with the same number of artificial pneumothorux refills.

Artificial Pneumothorax.—There were 43 initial inductions of artificial pneumothorax and 1,406 refills performed at the Sanatorium Pavilions, City Hospital, Walker Gate, during the year. Since the year 1922, 400 patients have received this form of treatment.

Sanocrysin.—This was administered in 1 case.

Lipiodol has been used to assist with diagnosis when necessary.

NEWCASTLE GENERAL HOSPITAL.

117 patients were admitted (61 males and 56 females). Details are given in the following table:—

Patients who received Treatment in Newcastle General Hospital during the Year 1936.

Emporter ogrande	Sex	In Institu- tion on 1st Jan., 1936.	Ad- mitted.	Dis- charged.	Died in Institu- tion.	In Institu- tion on 31st Dec., 1936.
Pulmonary Adults	M.	4	3 4	6	1	
Do Do.	F.	****	4	6 2 29 31	200	2 13 7
Non-Pulmonary Do.	M. F.	18	33 36 25	29	9	13
Do Do.	F.	10	36	31	9 8 7	7
Do. Children	M.	19	25	20		17
Do. Do.	F.	8	16	15	5	4
Totals		59	117	103	30	43

5 patients were re-admitted and are counted as 10 admissions.

1 patient was ,, twice ,, is ,, 3 ,,

The results of the treatment received are given in the table below :—

	Males.	Females.	Children.	Totals.
Improved Without Improvement Died in Hospital	33 2 10	19 14 8	32 3 12	84 19 30
Totals	45	41	47	133

The total number of days of those who received treatment was 30,825, giving an average length of stay of 232 days.

Thoracic Surgery.—The Thoracic Surgeon at Newcastle General Hospital has continued to operate upon suitable cases. Whenever necessary he has visited the Sanatorium Pavilions, City Hospital, Walker Gate, and seen patients, in consultation with the Tuberculosis Medical Officer, that were thought likely to benefit by operative procedure.

Two cases of artificial pneumothorax had cauterisation of adhesions performed; three cases of bronchiectasis had a lobectomy carried out, one case of pulmonary tuberculosis had a thoracoplasty done, and two phrenic evulsions have been carried out.

STANNINGTON CHILDREN'S SANATORIUM.

The 30 beds maintained in this Institution for the treatment of Newcastle-upon-Tyne patients were kept fully occupied throughout the year and 55 children completed treatment. The details appear below:—

CHILDREN WHO RECEIVED TREATMENT IN STANNINGTON SANATORIUM DURING YEAR 1936.

	In Sana- torium	Ad- mitted		Persons who completed Treatment during the year.				
	on 1st Jan., 1936.	during the Year.	Number	Total Number of Days.	Average length of stay in Days.	torium on 31st Dec., 1936.		
Pulmonary Males		12	12	3,247	271	8		
Do. Females Non-Pulm, Males	5	13 16	9 15	1,679 1,764	186 118	11		
Do. Females		14	19	4,410	232	6 5		
Totals	30	55	55	11,100	202	30		

In every case except four benefit accrued to the patient, as is shown in the following returns:—

	Males.	Females.	Total.
Disease quiescent	7 18 2	9 17 2	16 35 4
Totals	- 27	28	55

No action has been taken under the Public Health Act of 1925 (compulsory removal of patients to hospital) or under the Public Health Prevention of Tuberculosis Regulations, 1925, dealing with milk.

I wish to acknowledge the loyal support and interest of my staff.

Yours faithfully, GEORGE HURRELL, M.D., D.P.H., Tuberculosis Medical Officer.

Tuberculosis Dispensary, 91, New Bridge Street, Newcastle upon Tyne, 2, 19th April, 1937.

BARRASFORD SANATORIUM.

Report of the Medical Superintendent.

TO THE MEDICAL OFFICER OF HEALTH.

SIR,

I beg to submit a report on the work at Barrasford Sanatorium during the year 1936.

General.—The optimum number of patients accommodated remains at 95, with which number a few single rooms are available on the ground floor for sick patients.

This number of beds is in excess of those required for Newcastle alone, and some of the surplus beds are rented to neighbouring local authorities as follows:—

Gateshead Corporation	10	beds for male cases.	
West Hartlepool Corporation	6	beds-3 for each sex.	
Tynemouth Corporation	2	beds for female cases.	

Early in the year Dr. Hazel Ashford resigned from the post of assistant medical officer, after approximately a year's service, and was succeeded by Dr. D. Cohen.

During the year the usual yearly sum was expended on painting—this time most of it was devoted to internal decoration, and a smaller amount to roofs.

Further development of the grounds was carried out, and the improved appearances reflect considerable credit on the work of the gardener (Mr. J. Henderson) and his staff.

Another length of road was macadamised, and progress in this direction is being made, though there is still much road work to be done

In the laundry the old ironing callender was replaced by a new machine, which was urgently needed. It is essential now to consider the replacement of the other two machines—the washing machine and the hydro extractor, both of which go back to the first days of the Sanatorium, that is 1907. X-ray Plant.—Each patient has had a radiograph taken on admission to the Sanatorium and during treatment when it seemed that useful information would be produced by further films. During the year 270 films were completed and the interpretations entered up in the patients' notes. All artificial pneumothorax cases are "screened" as a routine, and the screenings numbered 685, and the appearances were reproduced by diagram in the patients' records.

Dental Clinic.—The majority of patients have some carious teeth, and a smaller number have such a number decayed, together with infected gums, as to be incompatible with good health. An attempt is made to improve the state of the mouth in all cases in which the above conditions are present. Clinics are held each fortnight, when treatment is in the hands of Mr. G. Hutchinson, L.D.S.

During 1936 the following work was completed:-

Extractions	247
Fillings	68
Temporary fillings	19
Scalings	37
Attention to dentures	4
Examinations	10

The total attendances numbered 308.

Occupational Therapy.—No change has been made in this phase of sanatorium routine, which is of the greatest value to the person undergoing a treatment which is often prolonged, and from the very nature of the surroundings apt to be tedious and monotonous, especially to town dwellers. Occupational therapy offers to patients something practical and constructive, to occupy some of their time. It is applicable both to the more robust type of patient who has completed the long walks, and those who through their disability are unable to walk far and on whose hands time is likely to hang heavily.

It is divided into two main types, the one is the following of crafts, and the other is woodwork combined with suitable sorts of estate work. The former is carried out in workshops where the patients are instructed by a handicrafts instructor (Mr. J. A. Caughey). The crafts are leather, raffia and cane work, rug and basket making, poker work, and the making of other articles requiring concentration and care.

The standard of the finished articles is remarkably good, and the bulk of the goods made are sold and there is very little wastage of material. The attendances numbered 5,460; the women worked 5,274 hours and the men 5,646.

The woodworking section is housed in a separate workshop under the control of the joiner (Mr. F. C. Gerdes). Men only are employed, and as far as possible patients who normally earn their livings by the use of tools are given the opportunity of keeping in practice whilst under treatment. Here suitable woodwork is undertaken, and patients also are able to help in repairs and improvements about the estate.

In addition, men patients assist the gardener in suitable weather, in the maintenance and development of the garden and grounds.

Admissions.—The total number of cases admitted to the Sanatorium during the year was 192, three less than in the previous year. The number of Newcastle admissions was 144, as against 145 in 1935. Gateshead Corporation had 28, Tynemouth Corporation had 4, and West Hartlepool Corporation had 16. There were no admissions of private cases.

So far as Newcastle cases are concerned, there is rarely any delay in admission; a patient is usually in the Sanatorium within a week after the receipt of his application form. In the cases of the outside authorities who have a fixed number of beds and therefore often a waiting list, there is occasionally delay in admission.

Of the 192 admitted cases, 20 had been in the Sanatorium previously, and were disposed as follows:—

10	case had l	been adr	nitted t	twice pr	eviou	sly	in 1929 and 1	
1		23	,,	,,		**	in 1933 and 1	934
10	of the re-a	admittee	cases	was firs	tadm	nitted i	in1	909
1	11	.,		.,				920
1			"				1	923
1	**	277	3.5	**	**	.,,		
1	**	- 33	11	2.7	3.3	2.2		924
2	,,	,,		were	**	**	1	927
1	**	,,	**	was	**	.,,	1	931
4	**			were				933
5					33.0		1	934
0	,,	,,	33	11	**		1	935
4		22	33	33	**	**		200

Of these 20 re-admitted cases, 17 had had at some time or other tubercle bacilli demonstrated in the sputum, and in three cases tubercle bacilli had never been seen.

131

ADMISSIONS TO THE SANATORIUM DURING 1936.

Authority.	Male.	Female.	Total.
Newcastle Corporation	90	54	144
Gateshead Corporation	28		28
Tynemouth Corporation	1	3 5	4
West Hartlepool Corporation	11	5	16
	130	62	192
During 1935	123	72	195
During 1934	104	54	158
During 1933	108	51	159
During 1932	114	54	168
During 1931	125	60	185
During 1930	121	65	186
During 1929	124	54	178

Note.—Figures relating to the years 1921-1928 are given in the Report for the year 1932.

Discharges.—There were 190 discharges during 1936, as compared with 205 in 1935. No case died in the Sanatorium during the year. There were no summary dismissals during the year, and the total of these is only seven since 1921 when the Corporation acquired the Sanatorium.

DISCHARGES FROM THE SANATORIUM DURING 1936.

Authority.	Male.	Female.	Total.
Newcastle Corporation	85	59	144
Gateshead Corporation	27		27
Tynemouth Corporation	1	3	4
West Hartlepool Corporation	11	4	15
	124	66	190
During 1935	137	68	205
During 1934	97	47	144
During 1933	108	57	165
During 1932	111	60	171
During 1931	124	60	184
During 1930	131	59	190
During 1929	115	54	169

Note.—Figures relating to the years 1921-1928 are given in the Report for the year 1932.

SUMMARY OF MOVEMENTS OF PATIENTS DURING 1936.

Authority.	In residence night of Dec. 31st, 1935.	Admitted during 1936.	Dis- charged during 1936.	In residence night of Dec. 31st, 1936.
Newcastle Corporation Gateshead Corporation Tynemouth Corporation West Hartlepool Corporation	59 9 2 5	144 28 4 16	144 27 4 15	59 10 2 6
	75	192	190	77

Details in connection with Discharged Cases.

The particulars of patients and the results of their treatment, which are set out later, are based on the completed cases discharged. Of these 190, 17 exhibited no definite signs or symptoms of clinical tuberculosis, and were discharged as soon as this fact was established, and are excluded from the particulars and results of treatment which follow. The details (c to f) are, therefore, based on the 173 cases of definite tuberculosis.

(a) Length of stay-

The average duration of treatment of all cases was 149.53 days. Excluding the 17 non-tuberculous cases, 157.18 days. The 144 Newcastle cases alone averaged 168.82 days. The longest stay was 408 days, the shortest 5 days.

(b) Beds occupied and patient days-

Average number of beds occupied, 85.63. 52.06 by males, and 33.57 by females.

Total number of patient days was 31,340. 19,053 male, and 12,287 female.

Below is given an analysis of the average number of beds occupied, and the number of patient days.

Authority.	Average Beds occupied daily.	Patient Days.
Newcastle Corporation	67.98 9.94	24,882
Gateshead Corporation	1.97	3,641 724
West Hartlepool Corporation	5.71	2,093

(c) Social Status-

	Male.	Female.	Total.
Single	51	45	96
Married	57	16	
Widowers	2		73 2
Widows		2	2
TOTAL	110	63	173

(d) Age-

	Years.	Male.	Female.	Total.
16-20		20	9	29
20 - 25		18	19	37
25 - 30		17	13	30
30-35		19	11	30
35-40		10	4	14
40-45		10	2	12
45-50		6	4	10
50-55		6 5 2 3	1	6
55-60		2		2
60–65		3		3
	Total	110	63	173

(e) Occupations of 110 Male Patients-

Labourers Engineering and metal workers	19 7
Clerks	7
Motor drivers and mechanics	5
Salesmen	5
Joiners	4
Blacksmiths and Blacksmiths' strikers	3
Post office workers (outside)	3
Grocers	2
Railway workers (outside)	2
Railway workers (inside)	1 2
Electricians	2
Barmen	2
Rivetters	2
Ex-Naval ratings	2
Warehousemen	2

and one each of the following:—seaman, draughtsman, machinist, despatch hand, watch repairer, hawker, holder-up, message boy, caterer, hotel porter, butcher, rigger, gas fitter, schoolboy, cabinet

maker, mason, mineral water worker, hospital porter, painter, paper maker, insurance official, aircraft fitter, moulder, fruiterer, bus conductor, bricklayer, cook, boot repairer, van boy, office boy, upholsterer, paint mixer, window dresser, typewriter mechanic, electric welder, and one had no occupation. Total 110.

(f) Occupations of 63 Female Patients-

Housewives	16
Shop assistants	7
Housework at home	7
Clerks and typists	6
Domestic servants	5
Machinists	3
Factory workers	2
Nurses	2
Dressmakers	2

and one each of the following:—tailoress, tobacco stripper, cashier, laundress, theatre attendant, waitress, box maker, bakeress, paper bag maker, barmaid, office cleaner, dancing pupil, saleswoman. Total 63.

Diagnosis.

The diagnosis of pulmonary tuberculosis was confirmed bacteriologically either before admission or during residence in 117 cases (74 males and 43 females). 45 patients (34 males and 11 females) were apparently without tubercle bacilli in the sputum, and 11 patients (2 males and 9 females) said they had no expectoration; making 56 cases of tuberculosis in whose sputa tubercle bacilli had never been demonstrated. The clinical findings in all sputum negative cases can be divided as follows:—

Not suffering from clinical tuberculosis	17
Definite pleural tuberculosis without evidence of lung tuberculosis	18
Definite physical signs and X-ray evidence of lung tuberculosis without demonstrable bacilli	38

In the cases of the 38 patients in the last group, the radiographs all showed appearances suggesting the presence of deposit in the pulmonary situation for which tuberculosis shows a predilection. 237 sputum examinations were made in connection with these 38 cases, and as 9 had no sputum the average examinations in those that had was 8 each. 1,504 sputum examinations were made at the Sanatorium during the year; of these 474 were positive as

regards the presence of tubercle bacilli, and 1,030 were negative. 959 complete examinations of the chest were made during the year, together with routine examinations of the larynx and urine on admission of the patient, and subsequently when necessary.

During the year 17 cases were discharged as not suffering from pulmonary tuberculosis, and the diagnosis in these cases were as follows:—

Bronchiectasis	4
No pathological condition detected	4
Chronic inflammatory lesion.	3
Spontaneous pneumothorax	3
Chronic nephritis.	1
Hernia of stomach into pleural space	1
Diagnosis not established—sent to Hospital	

These 17 non-tuberculous cases were included in the 20 patients sent for observation for the purpose of making a diagnosis. Three were found to be suffering from pleural tuberculosis.

The period of observation for the purpose of diagnosis is set out below:—

Marakan	Under 1 week.				2 to 4 weeks.		More than 4 weeks.	
	М.	F.	М.	F.	М.	F.	M.	F.
Tuberculous Non-tuberculous	 5		1 3		1 4	1	2	

Lipiodol was employed when necessary in diagnosis, and 4 lipiodol bronchograms were produced, all of which showed the characteristic appearances of bronchiectasis.

Treatment.

Rest, diet, graduated exercise going on to occupation, are the points regarded as most important in routine treatment. It is essential, however, on admission and subsequently, to be certain that the bodily temperature is ranging normally. Unless there is any cause other than lung tuberculosis, a raised temperature is usually an indication of active disease, and rest in bed is essential until it has been reduced by bedrest alone or in conjunction with some form of special treatment.

93 of the 173 definite cases of tuberculosis were found to have normal temperatures during the whole of their residence, whilst 80 patients were feverish at some time or other of their treatment in the Sanatorium.

Afebrile throughout Treatment.	Febrile on Admission, Afrebile on Discharge.	Febrile Intermittently.	Febrile throughout Treatment.	Afebrile on Admission, Febrile on Discharge.
93	40	11	24	5

Lung collapse or artificial pneumothorax was used to a considerable extent as in previous years. It continues to give effective results in a large group of cases, which would probably not be attained by any other means. It controls symptoms and restores working capacity in successful cases, and is by far the most useful and widely used form of special treatment at the command of the tuberculosis worker. The rest provided by successful lung collapse allows stable healing to occur if the treatment is kept up sufficiently long.

39 of the 173 cases discharged in 1936 were considered to be suitable for treatment by lung collapse, but in 10 of them changes in the chest in the course of the disease on the chosen side, prevented the treatment from being carried out.

Of the 29 discharged cases treated, 10 were right sided and 18 were left and one was a simultaneous bilateral case. In addition to these, 14 cases had had an artificial pneumothorax induced before admission, bringing the number treated during the year to 43 (17 right, 25 left and 1 bilateral).

In connection with the above cases, all of whom were discharged during the year, 553 insufflations of air were performed, whilst during the year the total number of such operations was 685.

In 26 cases the induction of lung collapse seemed to be effective in controlling symptoms, 16 of them losing tubercle bacilli previously present in the sputum. In the remaining 17 the procedure was ineffective and was abandoned—in 3 cases only on account of effusion of fluid.

On discharge, the lung collapse is maintained by the Tuberculosis Medical Officer in most cases.

Since 1922, 446 cases have been treated by lung collapse at Barrasford, exclusive of those cases where it was induced before admission, which total 82.

Gold Salts.—Following on the unfavourable or valueless results in the use of gold salts in previous years, its employment has now been discontinued at the Sanatorium.

Ultra Violet Radiation.-No case was treated by ultra violet radiation during the year.

Results of Treatment.

The problem in the treatment of pulmonary tuberculosis is the case with tubercle bacilli in the sputum. The case of tuberculosis whose lung damage is represented by a pleurisy with fluid, or minimal lung involvement without bacilli in the sputum, almost invariably does well, provided an adequate period of treatment is accepted and commonsense management of activities for 1 to 2 years afterwards.

But the sputum T.B. positive case is different, and the most significant criterion of improvement is the sustained abolition

of sputum.

The case which, better in every other aspect, still has tubercle bacilli in the sputum at the end of treatment, is still a potential source of danger to others and himself, and it is probably true to say that relapse occurs sooner or later.

31 of the 117 discharged patients lost their bacilli during treatment. A case is regarded as T.B. negative who has had absence of sputum for a month, or where serial examinations of

sputum have been negative for at least this period.

During the year the estimation of the blood sedimentation rate was carried out as a routine, being done monthly in every case. The Westergren method, employing a 200 m.m. tube, has been used, and during the year 1,112 estimations were made and recorded.

The following are the weight records of the 173 definite cases of tuberculosis and the 17 non-tuberculous cases.

		Gained up to 7 lbs.	Gained 7 to 14 lbs.	Gained over 14 lbs.	Remained station- ary.	Lost up to 7 lbs.	Lost over 7 lbs.	Not weighed on dis- charge.	TOTAL
- (Gained weight	55	62	33		12			150 17
173 definite	Lost weight Stationary		****		4				4
cases. Not weighed on discharge							2	2	
То	tal	55	62	33	4	12	5	2	173
	Gained weight	7	6	2					15
17 non-									1
tuber- culous cases. Stationary Not weighed on discharge				1				1	
							1	1	
T	otal	7	6	2	1			1	17

Under the classification of cases introduced by the Ministry of Health, patients suffering from pulmonary tuberculosis are divided into:—

Class T.B. Minus, or those cases in which tubercle bacilli have never been demonstrated in the sputum, and,

Class T.B. Plus, viz., cases in which tubercle bacilli have at any time been found.

The latter class is further divided into three groups :-

Group 1.—Cases with slight constitutional disturbance, if any, and in which the obvious physical signs are of very limited extent.

Group 3.—Cases with profound systemic disturbance or constitutional deterioration, with marked impairment of function, and with little or no prospect of recovery.

Group 2.—All cases which cannot be placed in Groups 1 or 3.

To indicate results of treatment, the following terms are laid down:—

- "Quiescent."—Cases which have no symptoms of tuberculosis and no signs of tuberculous disease, except such as are compatible with a completely healed lesion, and in which the sputum, if present, is free from tubercle bacilli.
- "Arrested."—In pulmonary cases the term should be applied only to cases which have been "quiescent" for a period of at least 2 years.
- "Improved."—Cases short of "quiescent," in which the general health is fair and the symptoms of tuberculosis have materially diminished.
- "No Material Improvement."—All other patients who are alive.

When considered in these terms, the results of treatment of the 173 cases of lung or pleural tuberculosis can be set out as follows:—

	T.B. Minus			
		M.	F.	Total.
	Quiescent	19	7	36
	Împroved	12	8	20
	No Material Improvement	5	7 8 5	10
	T.B. Plus.			
		M.	F. 1 1	Total.
	(Quiescent	1	1	2
G.1	Improved	1	1	2
	Quiescent	1		2 2 1
	(Quiescent	1		1
G.2	Improved	31	29	60
	Quiescent	17	8	25
	(Quiescent			
G.3	Improved	3		3
	Quiescent	19	4	23

The number of T.B. minus cases which improved to the stage of quiescence is made up of cases of pleural tuberculosis which had no evidence of disease in the lung itself.

It is pleasing to record the great assistance rendered by Dr. Cohen in the clinical work, and to acknowledge the administrative efficiency and help of the Matron (Miss F. Baguley, A.R.R.C.) The loyalty and work of the rest of the staff, both nursing and lay, is very much appreciated.

Yours faithfully,

CECIL G. R. GOODWIN, L.R.C.P., M.R.C.S.,

Medical Superintendent.

Barrasford Sanatorium,

Northumberland,

25th March, 1937.

the locally and worth of the rest of the eight field hapsing and

REPORT OF THE MEDICAL SUPERINTENDENT, NEWCASTLE GENERAL HOSPITAL.

V.—GENERAL DISEASES HOME AND HOSPITAL.

DOMICILIARY MEDICAL SERVICE, NEWCASTLE GENERAL HOSPITAL. MEDICAL DESCRIPTION HOSPITAL

HOME AND HOSPITAL

DOMESTIANY MEDICAL BERVICE,

DOMICILIARY MEDICAL SERVICES.

This work was originally carried on by District Medical Officers, each of whom was in charge of a specified district in the City, and gave both medical attendance and medicines. These officers were remunerated by the payment of a salary and bonus.

By resolution of the City Council dated 20th September, 1933, an "open choice" method for the provision of Domiciliary Medical Services was introduced into six of the Medical Relief Districts as from 8th November, 1933.

During the year a further district was added to the seven already in the scheme, and the eight districts are now designated the Joint Medical Relief District.

It is proposed to add to the Joint Medical Relief District any other districts which may become vacant.

Domiciliary Medical Services in the Joint Medical Relief District are given by a panel of medical practitioners who have contracted with the City Council to provide the required services. Medicines, etc., for patients in the area of the Joint Medical Relief District are supplied from two municipal dispensaries which have been established at the Newcastle General Hospital and the Newcastle Dispensary, New Bridge Street.

A report on the working of the Scheme during the period 1st March 1935—29th February 1936, is included in this report (Appendix B).

The following table gives particulars of the work carried out during 1936 of the remaining District Medical Officers whose areas are not included in the Joint Medical Relief District.

District No.	District Medical Officer.	Number of Cases Treated.	Attendances by the M.O. at the Homes of the Patients.	Attend- ances by the Patients at the M.O.'s Surgery.
7	Dr. W. Simpson	*173	233	125
8	Dr. R. W. Nevin	2,591	2,863	3,705
10	Dr. T. J. Ryan	1,379	3,988	6,156

^{*}Resigned, figures relate to the period 1st January-1st February, 1936.

NEWCASTLE GENERAL HOSPITAL.

TO THE MEDICAL OFFICER OF HEALTH.

SIR.

It affords me much pleasure to submit for your consideration this report on the year's work in the Newcastle General Hospital.

Again I have to report an increased activity in all departments of the Hospital, the admissions having increased by 462 and the discharges by 480. It will be again noted that the increase has been due to the larger number of female patients and children admitted, the number of male patients admitted being only slightly greater than during the previous year. This has necessitated putting up extra beds in every ward. It clearly indicates the absolute necessity of increased accommodation. I have found it necessary to refuse immediate admission in the case of paying patients and it has also been found necessary to discontinue the admission of patients requiring dental treatment.

During the year 2,908 letters were sent out to doctors, of these 1,043 referred to medical cases and 1,865 to surgical. This service is much appreciated by medical practitioners.

The Sunday Morning Lectures were continued during the Spring and Summer months, the attendance at these being fairly well maintained. It was decided that during the Winter months when the general practitioners were likely to be busy and the weather stormy it might be advisable to discontinue them.

The course of lectures and demonstrations for undergraduates started in 1935 was continued, the average attendance being approximately 25. I consider that something of a more ambitious nature as regards clinical teaching could well be undertaken, with, I feel sure, benefit to the undergraduate.

We have had to deal with an increased number of cases of pulmonary tuberculosis, the large proportion of them having been admitted as suffering from some other disease. I think it is hardly realised how frequently the condition occurs in infants. The ready co-operation of the Tuberculosis Medical Officer has greatly facilitated the arrangements necessary in dealing with these patients.

A number of patients suffering from tuberculous phthisis have been treated by operation, these cases having been admitted from the City Hospital and the various surrounding Sanatoria, on the recommendation of the Tuberculosis Medical Officers. There has been a great extension in the thoracic surgery work, but this the subject of a special report by Mr. Mason, so requires no further reference from me.

This year's work in the Maternity Ward has again been a record one, 388 cases having been dealt with, compared with 273 in 1935. The number of children born was 397. We were fortunate in having no untoward happening, and the medical and nursing staff dealing with this branch of the work are to be commended on the excellent results obtained. The number of beds available in the present Maternity Ward is quite inadequate, but extra accommodation has been found in D. Block, and, although far from being an ideal arrangement, has definitely eased the situation. If the number of cases this year (1937) is maintained at the present level 1936 will be easily exceeded. These facts all go to prove the urgent necessity of expediting the building of a new Maternity Ward.

The Mental Wards, formerly administered by the Public Assistance Committee, were taken over by the Health Committee, the change taking place on 1st November, 1936. I have advocated this step for a number of years and am delighted that it has been accomplished. No difference should be made in dealing with the mentally sick and the physically sick. The new arrangement is working smoothly, and all concerned, patients, relatives and staff, appreciate the change.

The number of patients admitted was again high, only being 6 below 1935, which was a record year. It was found necessary to certify 30.4% of the males and 35.9% of the females, the remainder being discharged or transferred to other departments. A few cases were sent to the City Mental Hospital as voluntary patients.

Of the admissions to the Mental Wards 15% belonged to districts outside the City and were chiefly accounted for by admissions from the Royal Victoria Infirmary and to a smaller extent from the Princess Mary Maternity Hospital.

The incidence of infectious diseases was much lower than in the previous year, both as regards adults and children. Although sporadic cases of dysentery occurred it caused considerably less worry than it did in the previous year. The staff also was remarkably free from any trouble of an infectious nature. The number of operations performed in the theatres was the same as in the previous year, viz.:—2,722, but there was an increase of 84 in the number of major operations, and a similar decrease in the number of dental cases.

I would like to make mention of the excellency of the work of the anæsthetists and the important part they play in the good results obtained in the serious surgical operations which are undertaken. The new methods of anæsthesia have made these operations possible, and, what is of the greatest importance, have reduced to a considerable extent the apprehension and dread felt by all patients to a greater or lesser degree. It has also made much easier the period immediately following the operation. Anæsthesia has really become a very specialised part of medicine. The appointment of a full time anæsthetist has proved of the greatest possible advantage and has been very much appreciated by the surgeons.

The number of patients admitted for treatment following miscarriage continues to increase, being 38 higher than last year.

The Shadowless Lamp installed in the main theatre has proved a great improvement on the old lighting.

The work performed by Mr. Pattison, the Neurological Surgeon, has shown a marked increase, but this is the subject of special report so need not be further mentioned.


The arrangements for dealing with dental cases remains very unsatisfactory and this problem will have to be dealt with sooner or later. The system of admitting a limited number each week on Friday night for operation on the Saturday morning had practically to be discontinued from September on account of the lack of accommodation.

As a special report is being given by Dr. Nattrass and Mr. Evans on the work done in the radio-therm department it need not be referred to by me.

The admission of children again increased to the extent of 200 during the year. The incidence of specific infectious diseases was very low, but a large number of cases of gastro enteritis of a very severe type were admitted. No specific organism was isolated.

The new Children's Block in process of completion should prove a most valuable asset in the treatment of the children.







The Convalescent Home at Louisville was closed in July, up to which time 70 children had passed through. The new Home at Rothbury, Whitton Tower, was opened in July, children being admitted the following month. Up to the end of the year 36 cases had passed through, but it is fully expected that it will be possible to make full use of the accommodation during the summer months. A photograph of this Home is shewn on page 146A. The transport has caused a certain difficulty, but this has been overcome now, it is hoped.

The number of paying patients admitted for treatment continues to increase, as will be seen from the following returns for the last six years:—

1931	 63	1934	 170
1932	 105	1935	 218
1933	 124	1936	 308

This number could be very readily increased provided the accommodation were available, and again I would recommend considering the provision of accommodation of a private nature at an increased fee.

The Diabetic and Pernicious Anæmia Clinics continue to perform a useful part in the work of the Hospital. A special report is given by Dr. Swan on the work performed during the year.

As in other departments so has there been a great increase in the work done in the X-ray, Massage and Electrical departments. Treatments by massage have practically been doubled, this having been made possible by the appointment of a second masseuse.

The number of prescriptions dispensed for Domiciliary Medical Service patients was 28,746, an increase of 2,675, and the number of prescriptions dispensed for patients from the Diabetic Clinic was 744.

The number of cases admitted belonging to other districts, continues to increase, and, on the whole, less difficulty has been experienced in making the necessary arrangements, although even yet the procedure might be facilitated and expedited.

The number of nurses admitted to the sick wards was 46, one less than the previous year, and of these 7 were under treatment on more than one occasion. As is usually experienced, the most prevalent sickness was septic throat, and again it occurred usually in probationers during the early months of their training. Two nurses had to be transferred to Barrasford Sanatorium, one suffering from tuberculosis phthisis, and the other from pleurisy.

The former is now a patient in her home Sanatorium (Aberdeen), and appears to be making good progress. The other has made a good recovery and is now on duty. The following are the outstanding illnesses:—

Septic Throats	19	Septic Fingers	15
Quinsy	1	Tuberculous Phthisis	1
Acute Rheumatism	1	Tubercular Knee	1
Appendicitis	2	Pleurisy	1
Jaundice	2	Minor Injuries	5

The amount of special nursing involved, particularly in the case of surgical cases, has been a considerable tax on the Nursing staff, but the duties have been carried out without any question or complaint. An increase in the staff is urgently required and this necessitates the provision of additional accommodation.

The accommodation provided for the Resident Medical Staff calls for improvement, particularly as regards sitting rooms, these being very bare and uninteresting.

The Hospital has passed through a very strenuous year, still under difficulties, and I would like to place on record my appreciation of the good work done by all branches of the staff.

In concluding I would also express to you my thanks for the constant interest you have shown and the help you have given in all pertaining to the administration of Newcastle General Hospital.

ADMISSIONS AND DISCHARGES, ETC., FOR THE YEAR ENDED 31st DECEMBER, 1936.

	Males.	Females.	Children.	Total.
Admissions	2,281	2,928	1,498	6,707
Discharges	2,304	2,912	1,479	6,695
Of the Disc	harges—C	ured	1,649	
	R	Relieved	4,001	
	Γ	Died	1,045	
		Total	6,695	
(There were	also 28 de	eaths in the El	swick Grange.)	

TABLE OF AGES OF PATIENTS TREATED. Men over 60. 753 Women over 60. 597 Men under 60. 1,551 Women under 60. 2,315 Boys, 3-16. 348 Girls, 3-16. 325 Children under 3. 806 6,695

TRANSFERS FROM OTHER HOSPITALS, HOMES AND AUTHORITIES.

Royal Victoria Infirmary	49 + 6*
Gateshead County Borough	7
Gateshead P.A.C	14 + 1†
Gateshead Tuberculosis Care Committee	9
Shotley Bridge Colony	4
South Shields	4
Prudhoe Hall Colony	1
Northumberland County Council	46 + 1‡
North Riding of Yorkshire	2
Durham County Council	60 + 2‡
Tynemouth County Borough	2 + 1†
Tynemouth P.A.C.	2
Sunderland P.A.C.	1
Stoke-on-Trent	1
West Hartlepool	4
Northallerton & District Hospital Assn	1
PRIVATE CASES ADMITTED	308
Inquests held :	
Hospital cases (36); Elswick Grange cases (7)	43

^{*} Observation ward cases, 3 of which are chargeable to other Institutions.

^{,, ,,} case.

^{* ,,} cases (included in Royal Victoria Infirmary Nos.)

OPERATIONS			
FOR YEAR ENDED 31st DE		1936.	
Abdominal			
Gynæcological			
Thoracic			
Genito-Urinary			
Nose, Throat, Ear and Eye		176	
Blood Vessels			
Rectum Brain and Special Cases			
Skin and Subcutaneous Tissues.			
Examination under anæsthetic.		122	
Amputations		17	
Teeth			
Radium Cases		2	
Teeth extracted under local and	æsthetic	12	
Total		2,722	
Major Operations		786	
Teeth		and the same	
		2,722	
RETURN OF CASES TREATED IN A	MASSAGE	DEPAR	TMENT
Medica	ıl		
Massage. Electric Treatments 5,029 2,621		ight. 48	Total. 10,198
X-RAY DEPARTM	ENT.		
	sures		
,, ,, T.B. Dispensary 647 ,, Babies' Hospital 66	, T.B.	Dispensa es' Hospi	ry 647 tal 66
Barium Meals	Pari	um Meal	
Screened 205	,		ened 205
Total cases X-rayed 3,125	Total exp	osures	4,477
	zour en		
		36.	
		Vomen.	Total.
Under treatment, January 1st, 1936 Admitted during 1935	7 216	6 183	13 399
Admitted during 1900	210		
	223	189	412
Discharged during 1935:			-
Discharged during 1935 :— Cured	15	7	22
Improved	53	41	94
I.S.Q.	7	10	17
Transferred to :— Mental Hospital	68	68	136
General Hospital	40	37	77
A. and I. Wards	4	7	11
House (Able-bodied)	6	7	13
,, (Chronic) ,, Epileptic	10	1 2	11
Deaths	12	5	17
Under treatment, December 31st,			
1936	4	4	8

PATHOLOGICAL DEPARTMENT. SPECIMENS EXAMINED :-510 Sputa Fæces 479 Blood Sugars..... 111 Blood Ureas..... 376 Blood Cell Counts..... 264 Blood Vandenbergh..... Gastric Analyses 1,322 77 G.C. Smears Pus and Effusions 124 Cerebro-Spinal Fluid 150 Complete Urinary examinations...... 1,266 17 Urea Concentration tests..... Blood Sedimentation tests..... 26 Total...... 4,723 Diabetic Clinic Urines...... 1,049 Blood Examinations (Pernicious Anæmia Clinics) G.C. Smears 73 Total..... 1,276 TOTAL REPORTS FOR HOSPITAL LABORATORY 5,999. By arrangement with other Laboratories :-Wassermann Reactions 688 Bacteriological Examinations 461 Histological Sections 347 Total...... 1,496 ADULTS .- CLASSIFIED LIST OF DISEASES TREATED. MEDICAL. RESPIRATORY. Bronchitis 197 Broncho pneumonia 54 Lobar pneumonia..... Asthma Pleurisy..... Hypostatic pneumonia 48 Bronchiectasis..... 21 Others 13 17 Bronchitis and Emphysema DIGESTIVE. 74 Gastritis Dyspepsia..... 6 Gastric Ulcer Gastro-Enteritis 55 12 Cirrhosis of Liver 12 Constipation Duodenal Ulcer..... 33 Jaundice 7 Colitis 7 Others 16 11 Alcoholism NERVOUS. 86 Neuritis Cerebral Hæmorrhage..... 10 Cerebral Thrombosis Neurasthenia 43 45 General Paralysis 36 22 Functional Disseminated Sclerosis..... 18 Mental..... 83 44 Neuralgia 9 Epilepsy Locomotor Ataxy Others

Paralysis Agitans

4

DEFICIENCY DISEASE. Scurvy 37

INFEC	TIOUS	DISEASES.	
Encephalitis Lethargica	21	Diphtheria	2
Influenza		Paratyphoid	1
Erysipelas		Cerebro-Spinal Meningitis	1
Dysentery	3	Others	3
Scarlet Fever	1		
C	RCULA	TORY.	
Valvular Disease of Heart		Aneurism	11
Myocarditis		Pernicious Anæmia	
Pericarditis		Leukæmia	
Arterio-sclerosis		Secondary Anæmia	
Acute Endocarditis	5	Senility	
Hyperpyesia		Debility	32
Angina Pectoris		Others	6
Coronary Thrombosis	10		
1	RHEUM	ATIC.	
Acute Rheumatism	29	Sciatica	24
Chronic Rheumatism		Acute Arthritis	
Rheumatoid Arthritis		Chorea	
Lumbago		Others	
Gout			
F	XCRET	ORY.	
Acute Nephritis		Uraemia	17
Chronic Nephritis		Cystitis	
Pyelitis		Others	
		ECRETORY.	
Myxœdema		Diabetes Mellitus	48
Goitre	75	Others	8
		RVATION.	00
Cases for Observation			30
Cases for Observation	BERCU	JLOSIS.	
Cases for Observation	BERCU		30 67
Cases for Observation* TU Pulmonary	BERCU 80	JLOSIS	
Cases for Observation* TU Pulmonary	BERCU 80	JLOSIS. Non-Pulmonary OF DISEASES TREATED.	
Pulmonary	BERCU 80 D LIST SURGIO	JLOSIS. Non-Pulmonary OF DISEASES TREATED. CAL.	67
Cases for Observation	BERCU 80 D LIST SURGIO 225	JLOSIS. Non-Pulmonary OF DISEASES TREATED. CAL. Septic conditions	67
Pulmonary	D LIST SURGIO 225 5	JLOSIS. Non-Pulmonary OF DISEASES TREATED. CAL.	67 27 27
Cases for Observation	BERCU 80 D LIST SURGIO 225 5 139	JLOSIS. Non-Pulmonary. OF DISEASES TREATED. CAL. Septic conditions. Ulcers.	27 27 27 27 66
Cases for Observation	BERCU 80 D LIST SURGI 225 5 139 174 81	JLOSIS. Non-Pulmonary	27 27 27 27 66 22
Cases for Observation	D LIST SURGIO 225 5 139 174 81 17	JLOSIS. Non-Pulmonary OF DISEASES TREATED. CAL. Septic conditions Ulcers Cellulitis Abscess Gangrene Mastitis.	27 27 27 27 66 22 20
Cases for Observation	D LIST SURGIO 225 5 139 174 81 17 51	JLOSIS. Non-Pulmonary	27 27 27 66 22 20 16
Cases for Observation TU Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction	D LIST SURGIO 225 5 139 174 81 17 51 28	JLOSIS. Non-Pulmonary	27 27 27 66 22 20 16 4
Cases for Observation TU Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis	D LIST SURGIO 225 5 139 174 81 17 51 28 8	JLOSIS. Non-Pulmonary	27 27 27 26 22 20 16 4 9
Cases for Observation "TU Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70	JLOSIS. Non-Pulmonary	27 27 27 27 66 22 20 16 4 9 59
Cases for Observation "TU Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord Gastroptosis	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70 6	JLOSIS. Non-Pulmonary	27 27 27 27 66 22 20 16 4 9 59
Cases for Observation TU Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord Gastroptosis Peritonitis.	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70 6 4	JLOSIS. Non-Pulmonary	277 277 277 266 222 200 166 4 9 59 8 12
Cases for Observation TU Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord Gastroptosis Peritonitis. Plastic cases	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70 6 4 15	JLOSIS. Non-Pulmonary	277 277 277 666 222 200 166 4 9 59 8 122 37
Cases for Observation "TU Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord Gastroptosis Peritonitis Plastic cases Renal Calculus	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70 6 4 15 30	JLOSIS. Non-Pulmonary	277 277 277 666 222 200 166 4 9 59 8 12 377 18
Cases for Observation "TU Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord Gastroptosis Peritonitis Plastic cases Renal Calculus Kidney	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70 6 4 15	JLOSIS. Non-Pulmonary	277 277 277 666 222 200 166 4 9 59 8 122 377 188 222
Cases for Observation Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord Gastroptosis Peritonitis Plastic cases Renal Calculus Kidney Perinephritic Abscess	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70 6 4 15 30 8	JLOSIS. Non-Pulmonary OF DISEASES TREATED. CAL. Septic conditions Ulcers Cellulitis Abscess Gangrene Mastitis Carbuncle Varix Phlebitis Hæmorrhoids Empyema Diseases of Bone Diseases of Rectum Cystitis Displaced Cartilage Bursitis	277 277 277 666 222 200 166 4 9 59 8 12 377 18
Cases for Observation "TU Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord Gastroptosis Peritonitis Plastic cases Renal Calculus Kidney	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70 6 4 15 30 8 6 5	JLOSIS. Non-Pulmonary	277 277 277 666 222 200 166 4 9 59 8 12 377 18 222 266 25 5
Cases for Observation Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord Gastroptosis Peritonitis Plastic cases Renal Calculus Kidney Perinephritic Abscess Lung Diseases Fractures Dislocations	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70 6 4 15 30 8 6 5 104 5	JLOSIS. Non-Pulmonary	277 277 277 666 222 200 166 4 9 9 8 122 377 188 222 266 25 5 15
Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord Gastroptosis Peritonitis Plastic cases Renal Calculus Kidney Perinephritic Abscess Lung Diseases Fractures Dislocations Injuries, Wounds, etc.	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70 6 4 15 30 8 6 5 104 5 55	JLOSIS. Non-Pulmonary. OF DISEASES TREATED. CAL. Septic conditions. Ulcers. Cellulitis. Abscess Gangrene Mastitis. Carbuncle Varix. Phlebitis Hæmorrhoids. Empyema. Diseases of Bone Diseases of Rectum Cystitis. Displaced Cartilage. Bursitis. Deformities Post operative Simple Tumours Teeth	277 277 277 666 222 200 166 44 99 599 8 122 266 255 5 151 174
Cases for Observation Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord Gastroptosis Peritonitis Plastic cases Renal Calculus Kidney Perinephritic Abscess Lung Diseases Fractures Dislocations Injuries, Wounds, etc. Burns	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70 6 4 15 30 8 6 5 104 5 55 13	JLOSIS. Non-Pulmonary. OF DISEASES TREATED. CAL. Septic conditions. Ulcers. Cellulitis. Abscess Gangrene Mastitis. Carbuncle Varix. Phlebitis Hæmorrhoids Empyema. Diseases of Bone Diseases of Rectum Cystitis. Displaced Cartilage. Bursitis. Deformities Post operative Simple Tumours Teeth Toxic Goitre	277 277 277 666 222 200 166 4 9 9 59 8 12 266 25 5 15 174 9
Cases for Observation "TU Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord Gastroptosis Peritonitis Plastic cases Renal Calculus Kidney Perinephritic Abscess Lung Diseases Fractures Dislocations Injuries, Wounds, etc. Burns Prostate	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70 6 4 15 30 8 6 5 104 5 55 13 31	JLOSIS. Non-Pulmonary OF DISEASES TREATED. CAL. Septic conditions Ulcers Cellulitis Abscess Gangrene Mastitis Carbuncle Varix Phlebitis Hæmorrhoids Empyema Diseases of Bone Diseases of Rectum Cystitis Displaced Cartilage Bursitis Deformities Post operative Simple Tumours Teeth Toxic Goitre Osteo Arthritis	277 277 277 666 222 200 166 44 99 599 8 122 266 255 5 151 174 9 18
Cases for Observation "TU Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord Gastroptosis Peritonitis Plastic cases Renal Calculus Kidney Perinephritic Abscess Lung Diseases Fractures Dislocations Injuries, Wounds, etc. Burns Prostate Hydrocele	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70 6 4 15 30 8 6 5 104 5 55 13 31 14	JLOSIS. Non-Pulmonary OF DISEASES TREATED. CAL. Septic conditions. Ulcers. Cellulitis. Abscess Gangrene Mastitis. Carbuncle Varix. Phlebitis Hæmorrhoids Empyema. Diseases of Bone Diseases of Rectum Cystitis Displaced Cartilage. Bursitis. Deformities Post operative Simple Tumours Teeth Toxic Goitre Osteo Arthritis Intestinal Colic	277 277 277 666 222 200 166 44 99 599 8 122 266 255 5 151 174 9 18 13
Cases for Observation "TU Pulmonary ADULTS.—CLASSIFIE Carcinoma Sarcoma Hernia Appendicitis Cholecystitis Gastric Ulcer Duodenal Ulcer Intestinal Obstruction Diverticulitis Brain and Spinal Cord Gastroptosis Peritonitis Plastic cases Renal Calculus Kidney Perinephritic Abscess Lung Diseases Fractures Dislocations Injuries, Wounds, etc. Burns Prostate	D LIST SURGIO 225 5 139 174 81 17 51 28 8 70 6 4 15 30 8 6 5 104 5 55 13 31	JLOSIS. Non-Pulmonary OF DISEASES TREATED. CAL. Septic conditions Ulcers Cellulitis Abscess Gangrene Mastitis Carbuncle Varix Phlebitis Hæmorrhoids Empyema Diseases of Bone Diseases of Rectum Cystitis Displaced Cartilage Bursitis Deformities Post operative Simple Tumours Teeth Toxic Goitre Osteo Arthritis	277 277 277 666 222 200 166 44 99 599 8 122 266 255 5 151 174 9 18

PREGNANCY AND DISEASES OF WOMEN.

Albuminuria of Pregnancy Hyperemesis Gravidarum Pyelitis of Pregnancy Placenta Praevia Disorders of Pregnancy Miscarriage Puerperal Pyrexia Ectopic Pregnancy	391 4 13 7 4 21 159 1 3	Puerperal Sepsis Ovarian Cyst Salpingitis Uterine Fibroid Pelvic Cellulitis Diseases of Uterus Disorders of Menstruation Menopause Others	3 16 25 16 1 82 34 1 23
DISEASI Dermatitis	40 4 5 1	THE SKIN. Scabies	5 6 8 8
Conjunctivitis	3 13 3 ES OF	DISEASES. Gon. Rheumatism Late Syphilis Mixed Infection THE EYE. Others	3 6 1
DISEASES OF T Tonsillitis Tonsils and Adenoids Otitis Media Deflected Septum	19 67 18	Γ, NOSE AND EAR. Laryngitis Mastoid Others	2 5 13
CHILDREN.—CLASSIFIE	ED LIS	T OF DISEASES TREATED.	,
Bronchitis Pleurisy. Lobar Pneumonia. Broncho Pneumonia Bronchiectasis Circulatory Acute Nephritis Acute Rheumatism Chorea Excretory	63 4 35 47 14 24 10 13 30 9	Digestive Diabetes Mellitus Epilepsy Prematurity Marasmus Rickets Nursing For Observation Others	3 5 20 4 8
Impetigo	SKI 77 46 11 3 6	N. Tinea Verminous Congenital Syphilis Others	1 2 6 8
Conjunctivitis		THE EYE. Blepharitis	1

TH	ROAT, NOSE AND EAR.	
12 8 5	Tonsils and Adenoids Others	52 2
SURG	ICAL.	
38 16 2 1 8 12 16 13 12 21	Sarcoma Abscess Septic conditions Cellulitis Deformities Osteomyelitis Phimosis Teeth Others	1 36 5 4 3 6 9 11 25
NFEC	TIOUS.	
6 1 5 15 1	Varicella Cerebro-Spinal Meningitis Erysipelas Dysentery Diphtheria	1 2 3 10 7
BERG	CULOSIS.	
		46
	12 8 5 8 16 2 1 8 12 16 13 12 21 NFEC 6 1 5 15 1 1	8 Others

GEO. P. HARLAN, M.I.

Medical Superintendent.

Newcastle General Hospital, 14th March, 1937.

Report of Department of Neurosurgery.

In presenting a review of the work of the neurosurgical department for the year 1936, it is necessary that the work of previous years should be referred to, since this is the first time that the work of any year has been subjected to an analysis. By this means one is provided with a standard of comparison by which the work of 1936 may be judged and which may provide some indication of the lines for future developments.

Until 1933 there was no provision for the treatment of neurosurgical cases in the north-east of England. A few patients who were in the position to afford it were sent to the neurosurgical clinics in Edinburgh or Manchester, but as far as the majority were concerned, patients were seldom referred by the physician to the surgeon on account of the prohibitive mortality which attended brain operations when performed with the use of general surgical technique. Moreover, there was an increasing disinclination on the part of general surgeons to undertake cerebral surgery, due to their recognition of the totally different operative technique which was called for.

There is a common impression that the surgery of the nervous system consists almost exclusively of the surgery of intracranial tumours, and that this field is one in which the results achieved are a small reward for the great expenditure of time and labour involved. If it were possible to place a monetary value upon human life there would be some justification for regarding neuro-surgery as uneconomic, for apart from the high mortality of many operative procedures, many of the cases which the neurosurgeon justly regards as his most brilliant successes are scarcely able to return to a state of full economic usefulness.

Nevertheless, the surgery of intracranial tumours is and doubtless will continue to be the most important activity of a neurosurgical clinic, being invariably the neurosurgeon's dominant interest. In this field there is an unsurpassed opportunity for scientific observation, and it is my firm belief that to regard neurosurgery from a material and utilitarian viewpoint is a mistaken and shortsighted policy, for any line of scientific inquiry, if resolutely followed, will sooner or later and often in some unexpected manner contribute to the common welfare.

The exacting demands of the surgery of intercranial tumours have raised the standard of technique in the performance of other neurosurgical operations to such a level that the success which attends these operations could never have been achieved had the surgery of such conditions as epilepsy, intracranial vascular disease and trigeminal neuralgia developed along their own lines. Moreover, the highly disciplined team work which is an essential attribute of neurosurgery is reflected to an increasing extent in other fields of surgery.

In reviewing the in-patient material which has passed through the department the cases of the tumour group will be considered first. The "verified" cases are those in which the presence of an intracranial tumour has been confirmed by the histological examination of tissue obtained at operation or autopsy. If such confirmation is lacking the case is classed in the group of "unverified" tumours until the diagnosis is finally settled. This group for any given period of time diminishes, for most of the cases belonging to it will either be re-operated upon or will come to autopsy when the diagnosis will be histologically confirmed or refuted. The only cases which remain permanently in this group are those in which no autopsy is carried out. The group of "suspect" cases for a given period also diminishes with the passing of time, for with continued observation a case will either declare itself as a definite case of intracranial tumour, or the possibility of the presence of a tumour will be excluded by arriving at an alternative diagnosis.

It is of some interest to note the extent to which the services of the department have been given to cases from outside the Newcastle boundaries. This applies equally to cases of all types admitted to the clinic, but a consideration of the domicile of the cases of the tumour group is sufficient illustration of this point.

VERIFIED TUMOURS (INCLUDING SPINAL CORD).

Year.	Number.	Newcastle.	Tees-side.	Other districts.
1933	25	9	3	13
1934	33	16	2	15
1935	36	15	8	13
1936 (to Augu	ist) 25	7	2	16
	-			11 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1
Total	119	47	15	57
	-	-	-	Printer.

Cases of other types.

ARACHNOIDITIS-SEROUS AND CISTERNAL.

1933		 2
1934	****	 9
1935		 7
1936		 4

This condition is one of considerable importance since it is a frequent cause of blindness. Four patients of the above series were already blind when admitted to hospital, but in the others good results have been obtained following simple decompressive operations. There is no striking increase in the number of patients of this class for reasons which are probably related to some obscure epidemiological factor. It is noteworthy that the great majority of these cases have been operated upon during the winter months.

EPILEPSY.					
1933			3		
1934		****	5		
1935			7		
1936		****	13		

The recent understanding of the essentially focal nature of the origin of the convulsive attack in epilepsy has resulted in the successful surgical treatment of many cases which have previously been regarded as "idiopathic." This is reflected in the increasing number of epileptic patients who are admitted to the clinic. These cases have been selected from a rapidly increasing number of epileptics who are referred by their local doctors. The majority of these cases, even in the light of modern knowledge, are obviously unsuitable for surgical treatment and their number is already proving a difficult problem in the absence of adequate facilities for dealing with out-patients. Moreover, the clinic is often besieged by epileptic patients who attend for no other reason than the monthly supply of prominal, for which the National Health Insurance makes no provision. No detailed records of these out-patient attendances have so far been kept.

```
VERIFIED INTRACRANIAL TUMOURS.
              .... 25 (20 operated)
       1933
              .... 33 (25 operated)
.... 33 (21 operated)
.... 28 (27 operated)
       1934
       1935
       1936
 VERIFIED SPINAL CORD TUMOURS.
       1933
             .... 0
       1934
                          0
       1935
                          2
       1936
UNVERIFIED INTERCRANIAL TUMOURS.
       1933
       1934
       1935
                          6
       1936
UNVERIFIED SPINAL CORD TUMOUR.
       1935
              ....
 INTRACRANIAL TUMOUR SUSPECTS.
       1933
             .... 0
                          2
       1934
                         12
       1935
             ....
                     .... 9
       1936
```

The large number of verified cases of intracranial tumour during the year 1933 calls for some comment. Regular operative work did not start until April of that year, and all work in the clinic was interrupted for four months in the later part of the year. The fact that 25 cases were dealt with in the course of six months is a sufficient indication of the urgent need which existed in the district for the provision of a neurosurgical clinic; it indicates also the ready response of the profession to the provision of facilities for treating a class of case hitherto unprovided for.

The majority of cases in the "suspect" group are those sent directly to the hospital by practitioners with the diagnosis of an intracranial tumour in which subsequent investigations reveal no positive evidence of a tumour or other grounds for operative intervention. The number of these cases is very small in proportion to the number of verified tumours since most cases of the latter are seen by me in the hospitals throughout the district which co-operate with this clinic. It should be mentioned that the number of cases of all types seen in other hospitals with the view to their transfer to the neurosurgical clinic amounts to at least twice the number of cases handled in this clinic although no record of their actual number is available.

Intracranial Vascular Disease (including Aneurisms).

1000			
1933	****	****	4
1934			6
1935			10
1936	****		13

This class is represented almost entirely by cases from the tumour suspect group in which investigations have excluded the presence of an intracranial tumour. The increase in the number of cases in this group would appear to be proportional to the number of cases admitted for investigation.

TRIGEMINAL NEURALGIA AND ALLIED CASES.

1933		 5
1934		 8
1935	****	 0
1936	1000	3

The large incidence of these cases in the early period of activity of the clinic is perhaps explained by the fact that in many of them the symptoms were of many years' duration, no surgical treatment having been undertaken in view of the prohibitive risk which attends the section of the trigeminal sensory root when performed in the absence of proper facilities. Trigeminal neuralgia is a rare disease and it is improbable that more than four or five cases will be seen each year.

UN	CLASSI	FIED.	
1933			5
1934			7
1935			12
1936			16

The foregoing group does not include many cases of head injury which have been supervised at the request of the general surgeons. Unless these cases have displayed features of some special neurological interest no record has been kept of them outside the general surgical records. The supervision of cases of head injury has frequently been undertaken in other hospitals cases by the invitation of the surgeons concerned.

OPERATIVE '	W	O	RK	
-------------	---	---	----	--

Major operations.	1933.	1934.	1935.	1936.
Osteoplastic explorations	21	20	23	40
Suboccipital explorations	5	16	8	5
Subtemporal decompressions.	9	8	11	6
Trigeminal nerve sections etc.	3	14	0	5
Auditory nerve sections	2	0	4	1
Ventriculograms	16	23	34	43
Laminectomies	0	2	4	3
Unclassified	12	10	5	13
	-	-	_	
Total	68	93	89	116
	Total Control	-		-

The foregoing statistics of the work of the neurosurgical department during its four years' existence show a steady increase in all branches of neurosurgery. The only possible exception is that of intracranial tumours if one is to judge the work of the clinic by the number of patients admitted, but the number of operations for intracranial tumour performed during 1936 is more than in any previous year. An increasing number of patients are seen and supervised in other hospitals in the district, and when a patient's prospects are seen to be hopeless he is not admitted to this hospital. In previous years all patients with the diagnosis or even the suspicion of an intracranial tumour were admitted to this hospital, but with the existence of better facilities for the selection of cases only those cases are admitted in which there are some hopes of ultimate improvement. This policy has been attended by an increasing reduction in operative mortality, but it is important that it should not be pursued too far for a certain amount of clinical material of scientific interest will otherwise fail to reach the department.

In reviewing the work of the department one is faced with the question of how far the existing facilities satisfy the needs of the population of this district. The department has constantly, though at times with difficulty, kept pace with the demands made upon it. How far the neurosurgical requirements of the population are met is another matter. It is an established principle in the United States of America that a population of one million will keep a neurosurgeon fully occupied. If the incidence of neurosurgical diseases in this district is equal to that of America one is forced to the conclusion that a large amount of clinical material in this district remains untouched. Moreover, in considering the increase in the number of patients who are admitted to the department, one must not lose sight of the fact that an increasing number of patients are being admitted to the department from areas as far afield as the West Riding of Yorkshire and the North Midlands.

The absence of any official association of the department with the Newcastle College of Medicine has contributed to the formation of close ties elsewhere. Since October, 1934, all routine pathological work has been carried out in the laboratory of the Royal College of Physicians of Edinburgh, a large amount of this having been done gratuitiously when the pathological material has had some features of special interest. With the combined material of the Edinburgh and Newcastle neurosurgical clinics an Intracranial Tumour Registry has been established by the Royal College of Physicians of Edinburgh. This institution, which is the only one of its kind in this country, provides an unrivalled opportunity for the study and correlation of information regarding the less common types of intracranial tumour. A close interchange of material and opinion exists between this and the Manchester clinics. Certain extensive studies have been carried out on our material in the departments of anatomy and surgery of Manchester University and papers are shortly to be published on the combined experience of the two clinics on certain subjects.

In concluding my report it is the source of the greatest pleasure for me to be able to express my appreciation to my colleagues and assistants, to the operating and ward nursing staffs, and to Dr. Harlan, for the help and encouragement which they have so constantly given.

A. R. D. PATTISON, F.R.C.S.,

Surgeon in Charge of the Department of Neurosurgery.

Newcastle General Hospital, June, 1937.

FEVER THERAPY DEPARTMENT.

It has long been recognized that fever in disease is part of a bodily reaction of defence, usually against micro-organisms. From time to time attempts have been made to use artificially induced fever as a means of treatment, for example, the practice of bathing in very hot natural springs in Japan, for the subjects of certain diseases, goes back for centuries. Occasionally the accidental occurrence of an infection with high fever in the course of some chronic disease has been followed by prompt and unexpected recovery of the latter.

Such isolated observations culminated in the great discovery by Wagner-Jauregg of Vienna, in 1918, of the curative effect of artificially induced malaria in general paralysis of the insane (G.P.I.) To inoculate with malaria patients already seriously ill was in the first instance a bold undertaking, and indeed Wagner-Jauregg planned the method for many years before he adopted it. In the meantime he tried various other methods of inducing fever, methods which he hoped would be less severe than malaria, but though considerable effects were produced they fell far short of those obtained when at length he employed malaria. In fact, the outlook in this disease, formerly associated with progressive insanity and uniformly fatal within five years, has been enormously improved by Wagner-Jauregg's discovery, though it is still a very serious malady.

Naturally enough, this work led to renewed interest in the possibilities of fever treatment, and research has proceeded chiefly in two directions. Firstly, workers have sought to discover what other diseases, if any, would respond to fever; secondly, numerous other methods for producing fever, besides malaria, have been investigated.

In the Fever Therapy Department of the Newcastle General Hospital we are endeavouring to take a share in both aspects of this work. Before the Department opened in 1933 much preliminary research had been done by one of us (S.F.E.) in the Physics Department of Armstrong College under the supervision of Professor W. E. Curtis, F.R.S., and with the support of the Medical Research Council. The work was inspired by the observation of a great American engineer, Dr. W. R. Whitney, of a rise in body temperature of persons working in the neighbourhood of short-wave radio transmitters. Subsequently, Carpenter and Page in Whitney's laboratory devised an apparatus whereby short waves could be utilized to produce fever in man. The work at Armstrong College

proceeded on similar lines, and so soon as technical perfection appeared to have been reached, a number of students and others submitted themselves to practical trials and permitted their temperatures to be raised to high levels. Finally, the apparatus was installed in the Newcastle General Hospital, where it has been used to produce fever nearly 500 times. It may be said at once that experience has amply justified our confidence in its reliability and safety.

The chief claim for the electrical method, as opposed to the inoculation of malaria, is that it is much safer because of its precise control and because it does not weaken the patient as does an attack of a severe infective illness. Most workers are also convinced that it produces a higher percentage of cures. Further, it has become manifest that in an important group of conditions, namely, gonococcal infections, in which this treatment is of great value, fever is required of a range and duration which can only be obtained by physical methods.

No finality has been reached in regard to the precise electrical method, and at the present time, with the further help of the Medical Research Council, we are engaged in constructing additional apparatus. The original machine is the property of the Medical Research Council, who have loaned it to the Hospital and have throughout watched our work with sympathetic interest. We feel that this example of co-operation between the national research organization and a municipal hospital is specially worthy of comment.

Malarial treatment originated in Austria, and electrical fever in the United States of America, where it has now been studied on a large scale with characteristic energy and enthusiasm. In the spring of this year the First International Conference on Fever Therapy was held in New York, and this we were privileged to attend. For the first time workers on this subject were able to discuss their mutual problems both in formal conference and privately. We were part of a very small British contingent. In addition we paid personal visits to fever therapy centres in New York, Rochester and elsewhere. The experience so gained was invaluable, and indeed without it we should have been unable to institute advances in treatment which we believe to be important and upon which we intend to start work during the coming winter.

F. J. NATTRASS. S. F. EVANS.

Newcastle General Hospital, July, 1937.

Report of Department of Thoracic Surgery.

It is my privilege to submit this, the first report on the work done in the Department of Thoracic Surgery at the Newcastle General Hospital, from its commencement until the end of December, 1936—a period of just over two and a half years.

A pressing need for accommodation as a result of the recent and rapid developments being made in Thoracic Surgery led to the establishment—at first on an experimental basis—of this Department in May, 1934. Since then in the period under review 174 cases of thoracic diseases have been undertaken and 340 operations performed. The steady way in which the volume of the work has increased is illustrated by these figures:—

During May—Dec., 1934, 17 cases were admitted and 27 operations performed.

During 1935, 65 cases were admitted and 118 operations performed.
,, 1936, 92 ,, ,, 195 ,, ,,

Some indication is also given by these figures, that several "operations" were required by many of these cases. Indeed, not only are many of the remedial operative interventions of necessity carried out in stages, but a large proportion of these cases require "major" investigations ranking for all practical purposes as operations. Such an investigation is usually some form of "endoscopic" examination, such as bronchoscopy, oesophagoscopy, or thoracoscopy, where the bronchus, gullet, or the interior of the chest respectively are inspected with the aid of specially designed electrically illuminated instruments; in this series, 116 of these examinations were made and are included in the figures just given. Direct estimations of the blood pressure within the great veins are being made with increasing frequency in connection with the work now in progress on cardio-vascular diseases.

Although only a relatively small proportion of those who suffer from chest diseases are suitable for surgical treatment at the present stage of its development, this does not detract from the value of your Committee's action in making it available for the citizens of this district. Their decision was a necessary one, because such facilities were not previously available in any of

the "voluntary" or other hospitals in the vicinity. Actually, this is one of the first hospitals in the country, certainly the first in the provinces, to make such special provision for all types of surgical thoracic work.

The Public Authorities have been responsible for many years for the treatment of tuberculosis, and in this connection they maintain an elaborate system of Dispensaries to which cases of many types of chest diseases are sent for investigation, etc. Although, as the statistics given later show, thoracic surgery is far from being principally concerned with phthisis, the value of a close association with this Dispensary system is obvious, and such is most easily effected, when the surgical clinic is established in a Public Authority's Hospital. Indeed, under ordinary circumstances, the almost complete absence of any private practice, whereby he may make a living consistent with the nature of his work, makes it impossible for a surgeon specialising in this difficult branch of surgery, at any rate in the provinces, to do so as part of the work of a "voluntary" institution. This position is almost unique amongst the various specialities into which modern medicine is divided, therefore thoracic surgery requires special consideration. The following factors are probably responsible; an unusually high proportion of those requiring treatment occurs apparently amongst the poorer classes of the community; very prolonged periods of hospitalisation are required—the average stay in hospital of our cases has been 66.3 days, this figure includes those who have been admitted for investigation for a few days only, and others which have been in hospital for more than 200 daysand the actual stay of those subjected to any major thoracic operation has rarely been shorter than three or four months; finally, all these cases require repeated radiological examinations and other expensive investigations, on such a scale, that only a wealthy patient would be able to bear the cost personally. Accordingly, it seems obvious that the Thoracic Surgeon must be subsidised, not, as are his colleagues in general surgery, by private practice, but directly by the Public Authorities of the district in which he works, the services of each individual Thoracic Surgeon being utilised by several such Authorities. Certain cases of tuberculosis may and should be operated upon in the Sanatoria, if these are large enough to carry the necessary equipment, otherwise, as many cases as possible requiring thoracic surgery should be concentrated in one centre, so that the most economical use may be made of both personnel and equipment, and needless and wasteful reduplication avoided. Likewise, academically and scientifically greater benefit would probably be derived from the experience at a "clinic" in which the bulk of the work from the area is centralised than from a number of small and—by nature of the expensive equipment required—relatively inefficient clinics.

The most useful contribution which the Thoracic Surgeon would apparently be able to make towards the work of the voluntary hospitals of his district, will probably prove to be of a consultative nature, cases being seen by him, when requested, and transferred if suitable for further special investigation and operation to the Public Authority's Hospital—an arrangement already obtained elsewhere and likely to prove successful. Actually, this has been the unofficial practice in this city ever since our Department was established, and in an attempt to have this arrangement made "official," an appeal was made some time ago to the Staff of the Royal Victoria Infirmary asking for their co-operation; their answer has not yet been received, but it is hoped that it will be favourable, and an arrangement, beneficial to the district and facilitating the working of this Department, will result.

The Authorities of the surrounding districts have co-operated where possible in arranging for the admission of patients living in their areas, and it is likely that in the near future this co-operation will become closer. At present it is necessary to negotiate each such admission separately, an arrangement which sometimes entails the loss of much valuable time; the institution of some "automatic" mechanism whereby such cases could be dealt with without undue delay is desirable. "Unification of Tyneside" in whatever form it eventually takes should be of help in this way.

Although the majority—102—of the cases treated in the Department since its beginning were naturally citizens of Newcastle-upon-Tyne, a considerable number—actually 73—were from the area of some other Local Authority through whose co-operation their admission was facilitated. These latter included:—

22 cases from Gateshead.

21 ,, ,, Durham County.

20 ,, ,, Northumberland.

5 ,, Middlesbrough, Hartlepool, and Billingham.

3 ,, ,, South Shields.

2 ,, ,, Darlington.

Scope of Thoracle Surgery.

It may not be out of place in this first report, to outline some of the conditions with which we have had to deal, and to explain the measures adopted for their relief. The impression, widely held, that this branch is principally concerned with the surgical relief of phthisis is erroneous. Admittedly, this forms a very important but by no means the principal part of the work, as a glance at the following analysis of the 174 cases dealt with in the Department shows :-

Of these, 50 were cases of pleural or pulmonary tuberculosis;

67 were cases of other inflammatory conditions of the lungs, such as abscess, gangrene, pneumonitis, infected lung cysts and bronchiectasis;

27 were cases of other inflammatory conditions of the pleura, chest wall, and sub-diaphragmatic spaces;

12 were cases of malignant disease of the lungs and bronchi;

mediastinal growths and abscesses;

oesophageal obstruction;

n n diseases of the heart and great vessels.

(a) Tuberculosis is usually treated under the direction of a physician in a Sanatorium. At first it is customary to employ only constitutional measures, the patient being kept under close observation during this period. The disease may resolve completely in favourable cases with this treatment, but it not uncommonly fails to do so, and then tends to run one of two courses-one on the whole favourable, in which there is a more or less marked tendency to form scar tissue whereby the condition may heal; the other, and usually unfavourable type, is characterised by the occurrence of pneumonic consolidations in the affected portion of the lung. In the former some form of "collapse" therapy is likely to prove beneficial, but this is not the case with the latter type in which constitutional measures are usually relied upon in the hope that as the general condition improves so will the type of the local disease change. Efforts are being made to ascertain the possibility of so influencing by operative measures the local condition in the "pneumonic" types of the disease that a more "fibrous" reaction will result.

"Collapse" therapy includes a number of measures designed to facilitate the collapse of a diseased lung so that any scar tissue which has formed therein may contract and so healing is encouraged. The best and simplest

form of such collapse available is that attained by the establishment of an artificial pneumothorax; this may be inadequate because of the presence of adhesions which hold the diseased lung to the chest wall and so prevent its collapse. If these adhesions are not too dense, they may be detached from the chest wall by means of an electric cautery operated through a small puncture with the aid of the thoracoscope.

If the nature or distribution of the adhesions is such as to preclude this relatively simple operation, then it becomes necessary either to separate the lung from the chest wall by blunt dissection and to maintain the resulting collapse by inserting a plug of paraffin wax—"plombage" or to remove portions of the ribs from the overlying chest wall and collapse this in with the adherent lung; (Fig. 1, page 175A) this operation of "thoracoplasty" formerly one of great danger and severity has now by improvements in technique, based largely upon the work of Sauerbruch, and by carrying it out in several stages, been made comparatively safe. It is important to realise, that so long as a "focus" of tuberculosis remains uncontrolled and unhealed, its products-sputum-are a menace not only to the community but to the rest of the patient's lung tissue. Statistics are notoriously difficult to assess, and our experience is too recent to permit of any attempt to do so, but it is indisputable that the expectation of life of a patient with such an uncontrolled focus is exceedingly short. It is equally indisputable that there are some thousands of people in Central Europe, Scandinavia, and America, alive, cured of the disease, and working to-day, as a direct result of such operations, who would otherwise now be dead.

The development of septic fluid—pyopneumothorax—in the pleural cavity is one of the most serious complications of pulmonary tuberculosis; in such cases an operation to drain the fluid away is urgently required, and if the patient survives this serious illness, a thoracoplastic operation is necessary to obliterate the resultant cavity.

(b) Suppurative diseases of the lungs are frequently seen by the Thoracic Surgeon, but the majority recover without operation—occasionally after a very prolonged period of observation. Spontaneous resolution of suppurative pneumonitis, of lung abscess, and even of lung gangrene may occur, and the symptoms of infected lung cysts and bronchiectasis—even in the inevitable persistence of gross anatomical deformity—may completely disappear. Hence active surgical intervention is withheld until it becomes obvious that either the maximum spontaneous recovery has occurred, or that the patient is in the optimum condition for it. A patient is in danger of death from septic poisoning, super-added pneumonia, or from abscesses in the brain so long as there is an active focus of lung sepsis. Some of these patients, who are regarded as free from sputum, are often found on examination to be swallowing relatively large quantities. Further, patients with such diseases as bronchiectasis, even when free from symptoms, are always in danger of relapse; the imminence-or what is of more importance for purposes of prognosis, the likelihood-of such a relapse cannot be foretold.

Lung abscesses are drained as a rule only when it is obvious that spontaneous recovery will not occur, when the acute phase is over and the lesion well encapsulated, and when precautions have been taken to ensure that the lung is so adherent to the chest wall that the drainage can be carried out without risk of spreading the infection to the pleura. (Fig. 2, page 175B.)

Removal of the involved portion of the lung is advised for bronchiectasis when the disease is limited to the lower part (Fig. 3, page 175c) or even the whole of one lung; it is now also known that it is possible to remove the lower portions of both lungs under certain circumstances. Collapse therapy has little practised application in this condition, being of no curative value and not without danger. Extirpation, although a serious operation, is followed in the great majority of cases by recovery, restoration to normal health, and freedom from expectoration.

Lung cysts of congenital origin occasionally become infected and may give rise either to the expectoration of septic sputum or to symptoms resembling empyemata for which operations may be radically performed; the symptoms are only permanently relieved when its true nature is recognised and the cyst removed. (Fig. 4, page 175D).

(c) Empyema—or septic pleurisy—is almost always a complication of acute pneumonia and was formerly treated by drainage as soon as the presence of the pus was detected. This is now known to be unwise, especially in the case of infants and young children, and in others in whom the empyema forms early in the course of the pneumonia; it is now customary to temporise by aspirating the pus—repeatedly, if need be—until the tissues have stiffened sufficiently to permit of the chest being opened without undue danger of displacement of the heart and other important organs.

Adequate drainage having been established its maintenance is imperative until there is incontrovertible radiological evidence that the lung has completely re-expanded and is everywhere in contact with the chest wall, otherwise recurrence and consequent chronic invalidism follows. The rate of this re-expansion varies enormously in different cases but it can usually be facilitated by suitable breathing exercises carried out under the supervision of a masseuse familiar with the object in view. The majority of the chronic empyemata referred to the Clinic have been directly attributable to premature removal of the drainage tube; simple restoration of the drainage for an adequate period sufficed to restore many of these cases to health, but in others an extensive thoracoplastic operation was required to terminate what was probably an unnecessarily prolonged illness.

(d) Growths of the lungs and bronchi are being recognised with increasing frequency as a result of the improved methods of diagnosis now available. Unfortunately the vast majority of these growths have already progressed beyond the bounds of legitimate surgical enterprise when first seen by the Thoracic Surgeon. This has been the experience in this Department; indeed, of the 12 cases admitted to the Department, not one was found suitable for removal, although in several instances exploratory operations were carried out to ascertain the possibility of this being done. Experience elsewhere, notably in America, has shown that it is possible to detect the presence of some of these growths at a sufficiently early stage to permit of their removal. When the occurrence of abnormal expectoration

—notably blood—is regarded as a symptom urgently requiring investigation, just as urgently as either the vomiting, of blood or its passage in the urine, more cases will be diagnosed during a phase amenable to surgical treatment.

The provision of facilities for treatment by high-powered X-radiations within the near future is likely to be of benefit to these cases to whom at present little or no relief can be given.

(e) The mediastinum is a region situated vertically in the centre of the chest between the two lungs, and about its diseases, and the possibility of their surgical relief, knowledge is slowly being accumulated. In foreign clinics the most frequent indication for opening the mediastinum is for goitres deeply situated behind the sternum and giving rise to asphyxial symptoms; strangely enough, no cases of this type have presented themselves for treatment here. When infections occur in such a deeply situated and relatively inaccessible region, they are naturally likely to have made considerable progress before a diagnosis, sufficiently accurate to permit of operation, can be made; even so, it is occasionally possible to bring such cases to a successful conclusion. (Fig. 5, page 175E.)

Asphyxial disturbances and congestion of the great veins due to compression of the structures within the mediastinum either by malignant growths or large aortic aneurysms may be afforded symptomatic relief by splitting the upper part of the sternum.

Dermoid cysts in the mediastinum may attain comparatively large dimensions without attention being drawn to their presence; indeed, they are usually discovered either as a consequence of some complication, such as erosion into a bronchus, or accidently. Their removal is now a recognised surgical procedure. (Fig. 6, page 175F.)

(f) The œsophagus, when there is any appreciable disturbance in the function of swallowing, requires a complete investigation in which radiological methods and direct endoscopic inspection are indispensable. Only too often, when a patient complains of dysphagia, an automatic diagnosis of "cancer of the gullet" is made and the patient literally abandoned. In many cases no doubt this diagnosis is correct, but in no case should it be made until the investigation referred to has been carried out, as it is only thus that certain benign lesions amenable to surgery may be discovered.

Cardiospasm—or spasmodic dysphagia—may be cured in many instances by psychological measures and in others by some form of dilatation; a certain number can only be relieved by anastomosing the lower end of the oesophagus to the stomach, an operation short-circuiting the region affected by the spasm or stricture and carried out most satisfactorily through the chest.

Several cases are now on record where it has been possible to remove the portion of the gullet affected by a cancerous growth, but speaking generally the results are extremely disappointing; it is hoped that technical developments will improve this position. In conjunction with Mr. Thurgar of the Newcastle Radium Centre a case of this type was treated by applying radium directly to the growth after bi-lateral exposure of the oesophagus; the result, though not in the long run successful, gave us much valuable information which we feel may be applied with some confidence of success to future cases.

(g) The surgery of the heart and great vessels is probably the most recently opened field for surgical enterprise. It has of course been known for long that cases have survived wounds of the heart after their repair; similarly, recovery has followed removal of fragments of shot and other missiles from the heart, but such are rare in this country, at any rate in peace-time. Empyema of the pericardium is an infrequent but lethal complication of sepsis elsewhere; when it is the principal disease, it appears to kill, rather by compressing and so embarrassing the action of the heart than by its virulence; accordingly, it is well worth attempting drainage when the condition is diagnosed, this having been followed by recovery.

Decompression of the heart by incising or by removing the pericardium is now a recognised surgical procedure for those cases in which it is apparent that the pericardium by reason of its tightness or its adhesion to the heart is interfering unduly with the return of the blood along the great veins to that organ. In these cases, ascites, cyanosis, and distension of the veins of the neck are prominent features; the heart may or may not be grossly enlarged, and it may or it may not be affected with valvular lesions; the venous blood pressure however is constantly found to be elevated—often markedly. It is important, when operating on these cases, to remember that the operation should be primarily a decompression, as we have found that extensive stripping of an adherent pericardium from the heart may deprive that organ of an important part of its blood supply, with serious consequences.

In certain cases, where the heart muscle receives an inadequate blood supply due to changes in the coronary arteries by which it is normally supplied, either heart failure or severe attacks of pain and syncope-angina pectorisoccur and ultimately kill the patient. O'Shaughnessy has shown that certain of these cases, if otherwise suitable, may be greatly benefited by grafting the omentum on to the surface of the heart; this operation is neither severe nor unduly dangerous, if the appropriate precautions are taken. The results of the operation are distinctly encouraging. Natural cure of saccular aneurysms of the aorta is by clotting and subsequent organisation of the blood within the sac; this process may be initiated and hastened by the introduction of gold wire (Colt) into the sac; in one such case of ours the aneurysm was pointing through the front of the chest wall; the patient, a bedridden invalid, was completely relieved after treatment by this method and now reports occasionally as an Outpatient.

Accommodation.

Naturally at first, when only a few cases were undertaken and admissions were sporadic, very little demand was made upon the accommodation; later, however, the position changed so rapidly that during the earlier part of 1936 about 25 beds were constantly occupied by these cases. Of late it has been necessary to increase this accommodation so that by the end of the year 30 beds were in use and it seems not unlikely that even this figure

will be exceeded in the near future in view of the frequency with which cases are presenting themselves for investigation and treatment. As the scope of thoracic surgery becomes more widely appreciated and as the arrangements for the admission of cases from surrounding areas become more automatic, the demand for accommodation in all probability will be such that it will be possible to keep between 70 and 80 beds more or less in constant use. The most suitable type of accommodation for these cases is that in which the majority of the patients are housed in rooms of about 4 beds each, together with an adequate number of single-bedded rooms for dangerously ill and other cases requiring such segregation. It is hoped that when the Department is re-housed in the proposed new block these requirements will be met. Accommodation is also required for the private cases which occasionally come for treatment and for whom there is no suitable accommodation available elsewhere in the City.

Academic Activities.

At present there is no systematic instruction in thoracic surgery given at the University of Durham College of Medicine, but during the past two years demonstrations have been given as part of the voluntary classes which the students may attend in this Hospital during the summer term. Similarly, some of the students have attended the operations in the Department. Your Thoracic Surgeon has also participated for the past three years in the various Postgraduate classes and intensive courses arranged by the College, including the Sunday morning Ward Demonstrations for general practitioners. Systematic instruction in the fundamental principles and scope of thoracic surgery is manifestly required by medical students, and it is desirable that this should if possible be given in a Clinic such as ours. Students from the College already have to attend affiliated institutions for instruction in infectious diseases, psychiatry, children's diseases, vaccination, and midwifery, so that no new precedent would be established if such an arrangement were implemented.

Numerous surgeons, not only from other centres in this country but from abroad, have visited the Clinic and seen the work in progress therein. Contributions to the literature relevent to the work of the Department include the following:—

- "Surgical Aspects of Phthisis," Newcastle Medical Journal, Vol. XIV., October, 1934.
- "The Post-Operative Management of Acute Empyema Thoracis," *British Medical Journal*, 29th December, 1934, Vol. II., p. 1197.
- "Some Impressions of Continental Thoracic Surgery,"

 University of Durham College of Medicine Gazette, February,
 1935.
- "Pulmonectomy" (A preliminary report on two cases in which an entire lung has been successfully removed), British Medical Journal, February 16th, 1935, Vol. I., p. 299.
- "Surgical Treatment of Pulmonary Tuberculosis," Clinical Journal, June, 1935.
- "Extirpation of the Lung," Lancet, May 9th, 1936, p. 1047.
- "The Treatment of Pulmonary Suppuration," Post-Graduate Medical Journal, November, 1936.

Every effort has been made to keep the work of the Department abreast of that in progress in other Clinics; accordingly, whenever and as frequently as possible these have been visited, discussions participated in, and conferences both in this country and on the continent attended. By the courtesy of the Council of the Royal College of Surgeons of England it has been possible to carry out a certain amount of instructive and necessary experimental work at their Research Farm at Downe, some of which is still in progress. As a Hunterian Professor, in February, I delivered a lecture before that College on "Extirpation of the Lung"—based largely upon experience derived from the work in the Department.

A cinematographic film illustrating the underlying principles and execution of the operation of Thoracoplasty was made in conjunction with Dr. James Whillis, of the University of Durham College of Medicine Anatomy Department. It received very flattering criticisms from the Film Institute, and has been shown at numerous medical meetings in this country.

Conclusion.

The work outlined in this report, although arduous, exacting, and often worrying, is of the greatest possible interest. Many of the operations performed are admittedly serious undertakings

but almost without exception the diseases for the relief of which their aid is invoked are such that the victims have little or no prospect of either prolonged existence or reasonable health if the operations are withheld.

It is impossible to acknowledge individually all those—especially my physician colleagues and the Radiological Department—upon whose co-operation the success of the work has largely depended. It is necessary, however, to record my appreciation of the highly skilled assistance rendered by my Anæsthetist, Dr. Philip Ayre, and the loyal way in which Mr. Andrew Logan, Surgical Registrar at the Hospital, my House Surgeons, and the Nursing Staff have done everything in their power to facilitate the work.

George A. Mason, M.B., B.S. (Dunelm), F.R.C.S. (Eng.),


Surgeon in Charge of the

Department of Thoracic Surgery.

Newcastle General Hospital, 31st March, 1937.

Fig. 1.

Total Paravertebral Thoracoplasty for Pulmonary Tuberculosis.




A young woman who had suffered from tuberculosis for several years, and who had received several courses of sanatorium treatment, was referred for surgical treatment. X-ray examination (a) showed that the left lung was extensively affected, but that the disease was of a fibrous type, as the lung, as a result of its shrinkage, had pulled the heart and trachea far over into the left side of the chest. Following thoracoplasty the patient has returned to normal life in good health and free from sputum. Subsequently X-ray examination shows that the lung has been satisfactorily collapsed (b).



Fig. 2.

Enormous Lung Abscess: Drainage: Recovery.




A man had suffered for several weeks from cough, expectoration of large quantities of foul sputum like anchovy sauce, wasting, anæmia and pyrexia. An X-ray picture obtained, when he was transferred to the Clinic, showed a large collection of fluid occupying the lower part of the left side of the chest (a). At operation this was found to be an unusually large abscess within the lung. Complete recovery followed drainage (b).



Fig. 3.


Lobectomy for Unilobar Bronchiectasis.



A young miner had suffered from occasional attacks of hæmoptysis and on several occasions there had been periods during which foul sputum was expectorated. He had been for a while in a sanatorium, where it was decided that he was not suffering from tuberculosis. Examination with the aid of Lipiodol demonstrated that there was bronchiectasis present and that this was confined to the lower lobe of the left lung (b). The affected lobe was removed at operation. Some months later the patient was able to return to work in good health.



Fig. 4. Congenital Lung Cyst.



A young woman had during the previous few years been operated upon on several occasions in different parts of the country for a supposed empyema. On admission to hospital, although the wounds of the previous operations were soundly healed, she was coughing up about half a pint of foul sputum each day. X-ray examination revealed the presence of a cyst (Fig. 4) in the lower and back part of the right lung. This was removed and the defect in the chest wall closed in several stages, the patient eventually returning home soundly healed and in good health.





Fig. 5.
Mediastinal Abscess: Drainage: Recovery.




A heavily built young man sustained a fracture in the vicinity of the second left costo-chondral junction. About ten days later he was admitted to hospital with a high temperature and a swelling on the left side of the chest. X-ray examination showed a large shadow continuous with that of the heart occupying the middle of the left side of the chest (a). At operation a large mediastinal abscess was drained and a complete recovery followed (b).



Fig. 6.

Dermoid Tumour of the Mediastinum.



A young woman had suffered from coughing and the expectoration of slimy sputum for a few months prior to admission. X-ray examination showed a large shadow bulging from the mediastinum into the right side of the chest (a). At operation this mass was found to be a dermoid tumour of the mediastinum and was removed with complete success, the patient subsequently obtaining work as a canteen worker. The condition of the chest after operation is seen in (b).



Report of Diabetic Clinic.

SIR.

I have pleasure in submitting a report on the activities of the Out-patient Diabetic Clinic at the Newcastle General Hospital for the years 1936 and 1937 up to 1st March.

For this period 85 patients are on the books of the Clinic, of whom females number 60 and males 25.

75 of these patients are regular attenders, the usual interval between visits being three weeks. 66 patients are on insulin and with a very few exceptions are supplied with it by the Hospital. There is very little trouble with the injections and only one sore arm has occurred during the period. The diets are as simple as possible, scales being supplied for weighing bread, etc.

Five patients have pulmonary tuberculosis but with one exception this was present before they started to attend. Only one case of peripheral neuritis has occurred.

The age groups are rather unusual as will be seen from the following table; the largest number of cases fall between the ages of 50 and 70. Nearly all of these are women.

AGE GROUPS.

AGE.	0-10.	11-20.	21–30.	31-40.	41-50.	51-60.	61-70.	71–80.	Total.
No. of patients	0	5	8	. 7	14	25	22	4	85

Ten clinic patients died during the period and are not included in the above figures.

The facts are as follows :-

Sex.	AGE.	Cause of Death.
M.	66	Not known.
F.	71	Carcinoma of gall bladder.
F.	65	Heart failure, hypertension.
F.	53	Diabetic coma, insulin stopped at home.
F.	57	Diabetic coma, cellulitis of leg.
M.	57 27	Diabetic coma, died at home.
F.	14	Diabetic coma, insulin stopped at home.
F. M.	61	Gangrene of leg and hemiplegia.
F.	70	Gangrene of leg.
F.	67	Arterio sclerosis, hypertension.

W. G. A. SWAN,

Medical Registrar and Physician in charge of the Diabetic Clinic.

Newcastle General Hospital, 15th April, 1937.

MAINTENANCE IN OTHER INSTITUTIONS.

Nine persons were maintained in various special institutions in different parts of the country during the year. The details are as follows:—

Institution.	Μ.	F.	Type of Case.
Home for Epileptics, Maghull St. Elizabeth's School for	2	1	Epileptic.
Epileptics, Much Hadham St. Vincent's Hospital for the		1	Epileptic.
Dying, Liverpool	****	1	Advanced Phthisis.
and Dumb, Boston Spa		1	Blind, Deaf and Dumb.
St. John's Home, Birmingham Hospital of St. John of God,		1	Deformity.
Scorton	*2		Cripples.
Total	4	5	

^{* 1} Case died 10th May, 1936.

REPORTS OF THE VETERINARY OFFICER
AND INSPECTOR OF PROVISIONS,
AND OF THE INSPECTOR UNDER THE FOOD AND
DRUGS ACTS (CHIEF SANITARY INSPECTOR).

VI.-FOOD.

BOVINE TUBERCULOSIS.

INSPECTION OF MEAT AND PROVISIONS.

INSPECTION OF FOOD AND DRUGS.

ANDISTO THANSATES, THE TO STOURS OF A COURSE TO HOLOTON THAT TO CHA

0007-IV

INSPECTION OF MEAT AND PROVISIONS
INSPECTION OF MEAT AND PROVISIONS
INSPECTION OF FROD AND DRUCK.

BOVINE TUBERCULOSIS, AND THE INSPECTION OF MEAT AND PROVISIONS AND FOOD AND DRUGS.

TUBERCULOUS MILK.

During the year 376 samples of milk were taken for bacteriological examination, 10 of which were reported by the Bacteriologist to contain tubercle bacilli. The positive samples were from the supplies of seven different farms, six of which were situated in the County of Northumberland, and one in the County of Durham.

The following table shows the percentage of milk samples found to contain tubercle bacilli during the past 17 years:—

	Percentage of Samples found Tuberculous.
1920	6.3
1921	5.5
1922	7.0
1923	4.5
1924	3.2
1925	8.0
1926	4.0
1927	3.7
1928	3.7
1929	8.7
1930	4.2
1931	3.7
1932	1.8
1933	2.0
1934	2.6
1935,	3.4
1936	2.7

Note:—Figures relating to the years 1907-1919 are given in the Annual Report for the year 1932.

Report of the

Veterinary Officer, Inspector of Meat, etc.

TO THE MEDICAL OFFICER OF HEALTH.

I have pleasure in submitting the following report which includes the work of inspection under the Public Health Acts during the year 1936.

Tuberculosis.

During the year, six animals, housed in registered cowsheds within the City, were found affected with one of the forms of the disease which required them to be dealt with under the Tuberculosis Order of 1925.

The animals, five of which were giving tuberculous milk, were subsequently slaughtered and the owners compensated according to the valuation before slaughter, as agreed upon by the Veterinary Officer, on behalf of the Corporation, and the owners, in accordance with the Order. Upon examination of the carcasses and internal organs, the disease in one case was found to come within the category of "tuberculosis not advanced," as defined by the Order, whilst in the remaining five, the animals were found to have been suffering from "tuberculosis advanced," necessitating the condemnation and destruction of the entire carcass and organs of each as unfit for human consumption.

When the amounts paid, comprising compensation to the owners, costs for cartage and slaughtering, commission on sales, etc., were deducted from the total obtained through the disposal of the carcasses, hides, offals, together with the percentage recoverable from the Ministry of Agriculture, etc., there remained a balance of £2 18s. 0d. in favour of the Corporation on the administration of the Tuberculosis Order during the year.

In the course of milk and meat inspection within the City during the year, 1,291 animals were found on slaughter to be affected with the disease, this being an increase of 12.55 per cent. as compared with the number found diseased during the previous year. In 984 cases some part of the carcass or internal organs of each was condemned and destroyed as diseased, whilst in the case of each of the remaining 307 animals it was found necessary, owing to the extent and distribution of the disease, to destroy the entire carcass and internal organs.

The Milk and Dairies Order of 1926.

Within the City there are 22 cow-keepers, registered as occupying 22 premises; and on the registered premises there is a total of 35 cowsheds in which are housed 515 milch cows.

During the year, 210 visits were made for the purpose of inspecting the animals, buildings, conditions as to cleanliness, etc.

In addition to the six tuberculous animals previously referred to, four milch cows within the registered sheds were found suffering from mastitis and other illnesses. As with the tuberculous animals, it was found necessary in one case, immediately the presence of disease was detected clinically, to adopt precautionary measures by excluding the milk in question from the public supply. When a sample of milk from this animal was examined microscopically, the mastitis was found to be due to streptococcal infection. In each of the three remaining cases the animal, although not affected with any condition likely to contaminate the milk supply, was, however, suffering from traumatic or other disease, necessitating its disposal by slaughter.

The Milk (Special Designations) Order of 1936.

The above-named Order, which came into operation on 1st June, 1936, revoked the Milk (Special Designations) Orders of 1923 and 1934. The designations now prescribed for the various grades of milk are generally recognised as an improvement on those prescribed under the revoked Orders. The latter comprised "Certified," "Grade A (Tuberculin tested)" and "Grade A;" and it was "Grade A," although inferior to the other two in respect of possible freedom from tuberculosis infection, which was commonly believed to be the best milk of all.

The special designations which may now be used in relation to milk are :—

- i. Tuberculin Tested;
- ii. Accredited; and
- iii. Pasteurised.

Tuberculin Tested Milk is milk from cows which have passed a veterinary examination and a tuberculin test; it is bottled on the farm or elsewhere and it may be raw or pasteurised. If bottled on the farm, it may be described on the bottle caps or cartons as Tuberculin Tested Milk (Certified). If pasteurised it is described as Tuberculin Tested Milk (Pasteurised). It must satisfy certain bacteriological tests.

ACCREDITED MILK is raw milk from cows which have passed a veterinary examination; it is bottled on the farm or elsewhere. It must satisfy the same bacteriological tests as raw tuberculin tested milk, and must not be treated by heat.

PASTEURISED MILK is milk which has been retained at a temperature of 145° to 150° F. for at least thirty minutes; and does not contain more than 100,000 bacteria per millilitre.

It is unlawful for any person to use any of the above designations unless he holds a licence from the appropriate Licensing Authority authorising him to do so.

Before granting Producers' Licences for "Tuberculin Tested" and "Accredited" milks, the Local Authority requires to be satisfied that the structure and arrangement of cowsheds and dairies, so far as lighting, ventilation, cleansing, drainage and water supply are concerned; the precautionary methods during milking; the cooling, storing and handling of milk; the apparatus for the sterilising of utensils; the condition of the herd, ascertained by clinical (veterinary) examination and, where necessary, the application of the tuberculin test and the results of the bacteriological examination of one or more samples of the milk, are such as to make it reasonably probable that the milk will regularly comply with the prescribed tests at all stages and at all times of the year.

DISEASED COWS FOUND IN REGISTERED PREMISES WITHIN THE CITY.

No. of Registered Premises.	₽.	of tered heds.	of Milch in City.	No. of Diseased Cows.						
	of tere			Tuber	culosis.	Other Diseases,		Destroyed		
	Regis Prem	No. of Registered Cowsheds.	No. of Cows in	Of Udder.	Other than Udder.	Udder.	Other than Udder.	(under the Tuberculosis Order, 1925)*		
†1920	26	26	40	565						
1921	25	25	38	575	1921	****				
1922	25	25	39	489	****					
1923	25	25	39	484	2		8		1	
1924	22	22	34	436	3	2	2		4	
1925	21	21	33	337	9	****	1		4*	
1926	20	20	31	410	5	2	1	3	5*	
1927	18	18	29	334	9 5 2 3	4	2	3	6*	
1928	19	19	31	308	3	1	1	3	4*	
1929	19	19	30	258	4 2	1	1	2	4*	
1930	17	17	28	251	2	3	1	4	4*	
1931	16	16	27	243	4	7	1	3	9*	
1932	16	16	27	246	4	2	7	3	6*	
1933	16	16	27	243	1		5	4	1*	
1934	14	14	22	223		2	6	4	5*	
1935	23	23	38	504	3	3	3	2	6*	
1936	22	22	35	515	5	1	1	3	6*	

[†] Figures relating to the years 1907-1919 are given in the Annual Report for the year 1932.

Anthrax.

During the past thirty years, the City has been fortunate in not being visited by this disease on more than eight occasions. In each of the years 1907 and 1908, two outbreaks occurred and in the years 1919 and 1925, one outbreak, involving only one animal in each case, occurred. Then after a period of nearly ten years' freedom, the disease occurred in 1935, and again during the year under report which is briefly referred to as follows:—

On 18th July, 1936, a veterinary practitioner notified suspected anthrax as existing in a cow found dead on a farm within the district. Within two hours of receiving the notification, the farm in the Kenton area was visited when blood smears, taken from the dead animal and examined microscopically, disclosed the presence of anthrax bacilli, thus confirming the original report. The usual procedure under the Anthrax Order, 1928, was then followed, including the declaration of the premises as an infected place; notifying the Medical Officer of Health and also the Ministry of Agriculture and Fisheries; segregation of non-infected animals; arranging for the disposal of the carcass; cleansing and disinfection, besides making the necessary inquiries as to the origin of the outbreak.

Within forty-eight hours of the death of the cow referred to, a horse which had been grazing in the same field was found dead. An examination of material, microscopically, disclosed the fact that the horse also had died from anthrax. The carcass of this animal was dealt with in the same manner as the cow carcase.

Fortunately, the milch cows on the farm were far removed from the field in which the six non-lactating cows and one horse involved in the outbreak—were grazing.

On making inquiries as to the possible source of origin, it was found that whilst no previous sudden or unexplained deaths had occurred on the farm in the past and that the animals involved had not received any artificial feeding stuff or any foreign litter, the field to which these animals were confined had been dressed with artificial manure of animal origin. Having regard to all the circumstances, the artificial manure used for dressing the field in question was suspected as a possible source of infection.

In the Ministry of Agriculture and Fisheries' Report, by the Chief Veterinary Officer, of Proceedings under the Diseases of Animals Acts for the year 1935, it is stated that the 320 outbreaks of anthrax in Great Britain unconnected with any previous case may have originated from the following sources:—

Effluent from tanyards, etc.	, caus	sing con	tamin	ation	
of streams					9
Feeding infected carcasses or	offals	to anin	nals		2
Use of imported feeding stuff	s				208
Use of artificial manure of a	nimal	origin o	on the	land	25
Simultaneous use of both imp	ported	l feeding	g stuff	s and	
artificial manures					49
No explanation available					27

Within Great Britain, 468 outbreaks of the disease were confirmed, in which 549 animals were attacked, as compared with 386 outbreaks during the previous year, in which 443 animals were attacked.

INSPECTION OF MEAT AND OTHER FOODS.

The number of animals slaughtered within the City for food purposes during the year was 257,343. Whilst there were 1,013

more cattle, 2,206 more pigs, there were 120 fewer calves and 637 fewer sheep slaughtered than during the previous year, as indicated in the following table:—

Animals Slaughtered on Licensed Premises within the City.

YEAR 1936.	1935.	1934.	1933.	1932.
Horses 1,565	1,610	982	950	1,266
Cows 3,005 Heifers. 15,760 Bulls 600 Bullocks 9,134	27,486	21,623	20,278	18,895
Calves 7,358	7,478	4,705	3,475	2,976
Sheep 172,844	173,481	163,556	167,653	186,662
Pigs 47,077	44,871	37,737	41,281	48,642
Total Animals 257,343	254,926	228,603	233,637	258,441

Five hundred and seventy-one animal carcasses, together with 2,168 lbs. of meat (excluding offal, etc.) were condemned and destroyed as being unfit for human consumption, as compared with 417½ animal carcasses and 3,170 lbs. of meat condemned and destroyed during the previous year. Of the 571 carcasses, 307½ were condemned on account of tuberculosis, as compared with 219½ carcasses condemned on account of that disease out of the previous year's total of 417½ carcasses. It is of interest to note that of beef alone, the total number of carcasses condemned for tuberculosis during the year under report is greater than that of the previous year by 51.57%, and is the highest since the year 1919.

Of the 47,077 pigs slaughtered, 725 were found affected with tuberculosis, which proportion—approximately 15 per 1,000—is practically the same as that of the pigs found tuberculous amongst those slaughtered during the previous year. As occurred during the previous year, the percentage of cases in which lesions of the disease were found confined to the region of the throat was high, for, out of the 725 found affected, in 606, or 83.59 per cent., were the lesions so localised.

Cattle, Calves and Pigs Slaughtered within the City.	Diseased, otherwise	nimals found Unsound or unfit for ensumption.	Number of Animals found Tuberculous.				
(See also previous Table).	Whole Carcasses Condemned,	Parts or Organs Condemned.	Whole Carcasses Condemned.	Parts or Organs Condemned			
Year 1936.		Year	1936.				
Cows 3,005	163	144	156	126			
Heifers15,760	47	104	44	80			
Bulls 600	4	17	3	14			
Bullocks 9,134	41	66	38	42			
Sex not known	****	264	****	54			
Totals28,499	255	595	241	316			
Calves 7,358	36	7	8	1			
Pigs 47,077	129	1,166	58	667			

CARCASSES OF BEEF CONDEMNED WITHIN THE CITY DURING THE PAST SEVENTEEN YEARS.

Total o	condemned.	Numbers condemned on account of Tuberculosis.	Percentage Tuberculous.			
Year.	Carcasses.	Carcasses.	Per Cent.			
*1920	198	171	86.36			
1921	90	78	86.66			
1922	85	79	92.94			
1923	69	58	84.05			
1924	66	61	92.42			
1925	157	130	82.80			
1926	126	102	80.95			
1927	123	107	86.99			
1928	115	109	94.78			
1929	124	118	95.16			
1930	147	124	84.35			
1931	117	94	80.34			
1932	135	120	88.89			
1933	128	116	90.62			
1934	186	158	84.94			
1935	182	159	87.35			
1936	255	241	94.51			

Note.—The above refers to whole carcasses and quarters, but does not indicate the total number of animals found tuberculous, and therefore does not include those carcasses in which only the organs or parts were found diseased and condemned. See preceding table.

*Figures relating to the years 1907-1919 are given in the Annual Report for the year 1932.

Public Health (Meat) Regulations of 1924.

For the purposes of these Regulations, 6,781 visits were made to meat and provision shops, restaurants, stalls, vehicles, etc., and as a result, 28 contraventions were found requiring to be dealt with. In two instances only were further proceedings necessary, the results of which appear on page 198, under the heading "Contraventions."

Inspection of Carcasses sent into the City from Outside Districts during the Year 1936, including the Carcasses of Animals taken under the Tuberculosis Order, 1925, by other Local Authorities and Slaughtered within the City.

Material Examined.	Condition	Found.	How Dealt with.
*4 Cow carcasses & Organs *1 Cow Carcass and	Tuberculos	is	Carcasses and organs condemned.
Organs	**		Lungs and udders condemned.
*1 ,, ,,			Head, lungs, mesenteric fat and udders condemned.
*1 ,, ,,	,,		Head, lungs and mesenteric fat condemned.
*1 ,, ,,	,,		Lungs and udders condemned.
1 Juliock Carcass and	Normal		Passed.
Organs 1 Heifer Carcass and			**
Organs	Congestion		Liver condemned.

^{*} Slaughtered under the Tuberculosis Order, 1925, certificate of examination in each case being sent to the local authority concerned.

Imported Foodstuffs.

During the year, 210 vessels carrying meat and other foods from Denmark, Holland, America, Canada, Australia, Russia and Madagascar, arrived at the Quayside, this being a decrease of three as compared with the number of arrivals during the previous year.

Four hundred and sixty-four visits were made to the wharves and vessels alongside, 2,401 packages, containing meat, etc., being opened and examined. Regarding these visits, seven were in response to official notices received from the Customs House concerning foodstuffs detained for our inspection and certification. Both before and while being discharged from the vessel, it is not practicable to make more than a general survey or superficial inspection of frozen beef quarters and carcasses of mutton, but these, as well as imported meat arriving by rail within the City, are subjected to supervision and inspection within the cold storage depots and wholesale meat shops.

Caseous Lymphadenitis.

During the year, of 26,314 carcasses of mutton, comprising seven separate consignments arriving at the Quayside direct from Australia, 479 were examined, none being found affected with the disease, as compared with the arrival of 17,225 Australian carcasses, comprising eight separate consignments, during the previous year, of which 505 were examined, two being found diseased.

Foreign Meat, etc., arriving by Vessel.

Salted Meat, Offal, etc.

Pork.—30 barrels.

Offal (barrels).—864 feet, 1,249 maws, 493 heads, 31 cheeks and 7 casings.

Frozen Meat.

BEEF.-15,835 fore and hind quarters and 16,269 crops.

(Packages).-1,358 boneless and 15 shin beef.

Offal (Packages).—25 tails, 4,017 kidneys, 3,153 livers, 15 hearts, 10 skirts, 160 (mixed) offals, 164 tripes, 5 heads, and 18 tongues.

Veal.—400 hind quarters and 100 crops and 172 packages.

Offal (Packages).—5 kidneys, 3 hearts, 871 livers, 3 tongues and 15 (mixed) offals.

Mutton and Lamb.—26,314 carcasses.

Offal (Packages).—60 livers and 5 tongues.

PORK.—3,844 carcasses and 455 sides. (Packages).—316 cuts.

Offal (Packages).—15 kidneys.

Other Goods.

561,754 sides Danish, Dutch and Canadian bacon. (Cases).—
1,335 American bacon and hams, 31,132 tinned meats and 5 sausages.

NUMBER AND ORIGIN OF VESSELS ARRIVING WITH FOOD.

Denmark.	Holland.	America.	Canada.	Australia.	Russia.	Mada- gascar.
106	71	8	14	8	2	1

Exported Foodstuffs.

The number of horses slaughtered within the City, for the purpose of the carcasses being exported for consumption on the Continent, was 1,565 or 45 fewer than during the previous year.

Number of Visits and Inspections of Premises during the Year 1936.

		Centr Iarke		Me Sho		Fis Shop		Provision Shops,						
Slaughterhouses.	Meat and Provisions.	Fruit and Vegetables.	Fish.	Wholesale.	Retail.	Wholesale.	Retail.	Wholesale.	Wharves and Vessels.	Cold Stores.	Restaurants.	Stalls, Carts, etc.	Food Preparing Factories.	Bakeries.
17,588	450	349	336	3450	720	27	2	4	464	3	2	2052	103	2

Total Carcasses, &c., Destroyed as Being Unfit for

		Carcas	ses, etc.		Lu	ings.		Kie	ineys.	L
	Beef.	Veal.	Mutton.	Pork.	Sets Ox.	Sets Pig.	Ox Hearts.	Ox.	Pig.	Ox.
Tuberculosis		8		58	312	5	41		****	133
Pyrexia	2 qrs.	****	****	26+ 20 lbs.		****	1			1
Pyæmia	1	9	4							
Septic Conditions	2+204	3	18+2	5+1	4		2	1		47
septic conditions	lbs.		qrs. +	qr.+ 57 lbs.						
Uræmia				2						
Jaundice				2				833		
Fatty Degeneration	****	****								2
Hyaline Degeneration	39 lbs.							88		
Pneumonia	****	1	2		5	293				
Pleurisy	****	2	1+1 qr.		31	40				
Pleurisy and Pneumonia		****	2			****				
Peritonitis		****	3	5	****	****	10.00			1
Pleurisy and Peritonitis Mastitis		1111	-	11	****	****				****
Cirrhosis		4417	44.17	****		1111	****		****	57
Cavernous Hæmangioma										6
Edema and/or Emaciation Parasites (distomatosis,		1	46	****		****				
cysts, etc.)		****	****		5	****			2	****
Bleeding, Congestion, etc.	2	4	38+1 qr.	20	3			3		3
Immaturity Γraumatism	2 qrs.+	Q lbe	3+2 qrs.	Q The	+***	****				****
	617 lbs.		+84 lbs.							
Decomposition		14+1		45½lbs.	5	1111	25		12	31-
	+7831bs.	qr.	qrs.+							10 lbs.
Unmarketable (including			298½ lbs.							105
animals from centres of										
infection of scheduled										
disease)		****	****	****		****				

Human Consumption during the Year 1936.

-	Heads.			Plucks		ts,	hs and tines.	F	at.	Udo	lers.				eads.
						Sets,	Stomachs an Intestines.					ails.	Ox Tongues,	ins.	Sheep Sweetbreads.
Calf.	Sheep.	Pig.	Calf.	Sheep.	Pig.	Ox.	Pig.	Ox.	Pig.	Cow.	Pig.	Ox Tails.	Ox I	Griskins.	Sheep
		564+42 halves	1		69	5	2	41		2			****		
				2	1										
				****	****			****		1111					
***	****		****			****	****		****		1111	****		****	
***			****			****	****			****		****		4411	
***	****		1		50									****	
			î	2	34										
	****	****		1000	2			****						****	
			****			2									
	****			1111	13			****	****					,	
***		****			****	****	****	****	****	5	1	****		****	
	4419		****	****	****	0.00	11/4	****	****	****		****		****	
	****	****		1111	****	0.00		****	****		****			****	****
***	****	****	****	****	****	2015	1111	****	****	****	****	****		****	****
				13	4										
	****	****		4	28			****						****	
	****			****	****	****				****		****	****		
***	****			1217	****	****	****	****	****	****		****		0.044	
2	48	24 cwts.		153	17				265 lbs.			25+ 15 lbs.	12 lbs.	1 cwt	4 lbs.
		14					688								
		****	****	****	****		000	****	****		****		****	****	

POULTRY, GAME, FISH, FRUIT AND VEGETABLES, PROVISIONS, &C., DESTROYED AS BEING UNFIT FOR HUMAN CONSUMPTION

		8 4 3 3 3	15	lbs.	11	11	2.486	1 968	1,523	13	11	11	100	9	11	100	1 1	1,146		
		::		tins	- 00	4-	- 1	- 1	95	152	12	00 00	850	11:	400	22	110	1-		
	Provisions, etc.	Bacon Biscuits	Raisins Rice, 81 bags	TINNED GOODS.	Applies	. 0	Catfish Fillets	Crab	Loganberries Lunch Tongue	Milk	Oranges	Pears Pineapole Cubes	Plums	Pork Brawn	Prawns Roast Duck and Ham	Salmon	Sausage	Veal and Ham		
		lbs.	1	4,368	1	1	1	1	1	1		1	1							
DURING THE YEAR 1936.	Fruit and Vegetables.	Bilberries, 381 chips		Carrots, 29 packages, +	Cauliflowers, 18 crates	Grape Fruit, 166 cases	Lemons, 5 cases	Lettuce, 119 crates	Oranges, 759 cases	Pears, 1 box	Potatoes, 50 packages +	10 tons 6 cwts.	Tomatoes, 236 packages							
DURING TH	Fish.	Codlings 532	Haddocks 35	Halibut, 1 + 101	Ling 17	Plaice, 6 boxes + 360	Salmon 30	Skate 123	Trout 195		SHELL	Craw Fish 120	Shrimps, 2 bags							
	Poultry and Game.	Chickens 10 Codlings	Ducks 1	Partridges 2	Pheasants 7	Rabbits182	Turkeys 2													
	Cause of Unfitness.								Dusound	out o	Unwholesome.									

Total Weight of Meat and Other Foodstuffs Condemned.

The approximate total weight of meat and other foodstuffs condemned during the year was 102 tons 10 cwts. 3 qrs. 17 lbs., comprising:—

	tons.	cwts.	qrs.	lbs.
Beef, Mutton, Veal, Pork	83	4	3	5
Offal and Provisions	19	6	0	12
end rezer site andar to	102	10	3	17

Microscopical Examinations.

During the year, microscopical examinations were carried out as an aid to, or confirmation of, diagnosis in connection with 36 cases under investigation.

The material examined comprised specimens of milk, blood and swabs taken from the throats of cows. Of the samples of milk examined for tuberculosis, five were found positive and twelve negative; and of the throat swabs examined for the same disease, one was positive and one negative. Of the specimens of blood and other tissues examined for anthrax two were found positive and fourteen negative. A specimen of milk examined for organisms other than tuberculosis, was found positive.

MICROSCOPICAL EXAMINATIONS.

	Specimens Examined.	Result of Examination.				
Consider the more and the sense to	Examined.	Positive.	Negative			
Samples of Milk examined for Tuberculosis	17 2	5	12			
Throat Swabs examined for Tuberculosis Blood, etc., examined for Anthrax Milk examined for Organisms other than	16	2	14			
Tuberculosis	1	1				
	36	9	27			

Slaughterhouses.

During the year, 75 separate premises were licensed for slaughtering purposes, this being one more than during the previous year.

Of the 75 licensed premises, 14 were vacant during the whole or part of the year and 22 were occupied by wholesale firms, the remainder being occupied by retail butchers.

Of the total animals dealt with within the City, 183,645, or 71.36 per cent., were slaughtered by wholesalers, the remaining 73,698, or 28.64 per cent., being slaughtered by retailers.

Licensed Slaughtermen.

Under the Slaughter of Animals Act, 1933, slaughtermen's licences were granted during the year to 14 persons whose applications were approved by the Health Committee.

The Merchandise Marks Act, 1926.

In accordance with the procedure outlined in section 2 (1) of the above-named Act, Orders have been made—by His Majesty in Council—known as the Merchandise Marks (Imported Goods) No. 3 Order, 1934; No. 5 Order, 1934; No. 7 Order, 1934; and No. 8 Order, 1931, relative to bacon and ham; dead poultry; certain classes of chilled, frozen, boneless and salted meat and edible offals; and salmon and sea trout, respectively, each of which, subject to the provisions contained in certain Merchandise Marks (Imported Goods) Exemption Directions, makes it unlawful to import into or sell or expose for sale within the United Kingdom any of the first three mentioned classes of foodstuffs or to sell or expose for sale within the United Kingdom imported foodstuffs of the last mentioned variety, unless they bear an indication of origin as described in the respective Orders referred to.

One of the objects of the Orders is to ensure that when imported meat, bacon or ham, poultry or fish is exposed for sale (whether whole, jointed, in pieces or cuttings on trays or in packages) it shall be so readily identifiable by means of marks, labels, tickets or notices, as the case may be, that a would-be purchaser could scarcely fail to gain a correct impression as to its origin.

As the Order relative to meat is somewhat complicated, because of the considerable variety of carcasses, joints and offals involved, an explanatory letter was prepared as a guide and sent to the wholesale and retail meat traders within the City.

Rats and Mice (Destruction) Act, 1919.

During the year, 173 visits were made to premises in respect of 135 complaints received, 231 premises, including others than those complained of, being inspected and dealt with. Of the 231 separate premises, rats were found infesting 221, the remaining 10 being found free from any evidence of infestation. As will be seen in the following table, the premises most frequently invaded by the pests were dwellings, allotments and shops, these accounting for a little more than 84 per cent. of the whole.

As pointed out in previous reports, many occupiers have the impression that it is only necessary to lay poison bait to solve the problem, no attention being given to the question of prevention. The rat problem, so far as buildings are concerned, is one that invariably involves the question of construction and repair. With the view of obtaining successful results and efficient administration, it has always been the practice—within the City, at least—to place before the public, so far as possible, the best known means of prevention, besides educating the individual occupiers as to the advantages of rendering premises independent, structurally, one from the other, and of operating simultaneously.

RATS AND MICE (DESTRUCTION) ACT, 1919.

Complaints received	135
Number of premises inspected and dealt with in connection with	001
the above	231
Number of premises infested with rats	221 173
KIND OF PREMISES DEALT WITH.	
Allotments	51
Bakehouses	2
Builders' Yards	2
Dwellings.	105
Factories	2
Garages	3
Halls	3
Hospitals	2
Hostels	3 2 2 2
Hotels	3
Housing Estate	1
Offices	2
Poultry Runs	1
Restaurants	â
Shops (food)	24
Shops (other than food)	14
Warehouses	3
	4
Waste ground Wharfs	2
	1
Workrooms	1
Total	921

CONTRAVENTIONS.

Offence.	No. of Cases.	Action taken, etc.
The Public Health Act, 1875. Deposited for the purpose of preparation for sale within a slaughterhouse, one carcass of cow beef which was diseased	2	Two persons were charged, one being fined £10 and £2-2-0 costs, the case against the other being dismissed.
Failing to give notice to the Local Authority of disease in the carcass and internal organs of a cow	2	In one case offender fined £5 and £1-1-0 costs, the other being dismissed.
Meat conveyed in dirty vehicles Meat, unprotected, lying on floor Dirty butchers' shops Using for the preparation and storage of food a room with which a w.c.	8 6 4	Offenders cautioned. Offenders cautioned. Offenders cautioned.
directly communicated	3 2	Offenders cautioned. Offenders cautioned.
Premises used for domestic washing where food is prepared for sale	1	Offender cautioned. Offender cautioned.
Blowing mutton carcasses with the breath	1	Offender cautioned.
Bye Laws. Dirty slaughterhouses	3	Offenders cautioned.
Imported meat, not labelled as such, exposed for sale	3	Offenders cautioned.

Thomas Parker, f.r.c.v.s., Veterinary Officer.

Town Hall,

Newcastle upon Tyne, 31st March, 1937.

Samples taken for Analysis during the Year 1936.

		of Sam btained			ilt of lysis.	A	ction	Tak	en.	
ARTICLE,	Formal.	Informal.	Total.	Genuine.	Not Genuine.	Prosecutions	Convictions.	Cases D sm ssed.	Cases With- drawn,	Remarks.
New Milk	724	4	728	669	59	24	3	20	1	In 29 of the remaining 35 cases (of the 59 samples "not genuine") the vendors were cautioned by order of the Health Committee, and in 6 no action was taken, these being "appeal to cow" samples, etc.—not for
Skimmed Milk	9		9	8	1	1		****	1	proceedings.
Cream (Tinned and Fresh)		10	10	10						
Butter	****	17	17	17	****		****			
Margarine		17	17	17						
Lard		1 2	1 2	1 2		****	****		****	
Tea		5	5	5		****				
Coffee	****	6	6	6			***		****	
Baking Powder		1	1	1	1111		****		****	
Custard Powder	1000	1	1	1						
Jams, Jellies and Marmalade Lemon Curd		14	14	14		****			1111	
Rice, Ground Rice, Tapioca, Semo-			10		2000		****	****		
lina, Sago and Corn Flour		6	6	6						
Flour, Wholemeal and Oatmeal Split Peas, Barley, Lentils and		3	3	3		****	****	****	****	
Haricot Beans		4	4	4		400		1000		
Herbs (Sage)		9	9	9				****	****	
Dried Fruits (including Currants	****		9		****			210.0	20.00	
and Raisins)		13	13	13	****					
Candied Peel, Glacé Cherries, Almonds (Ground and Whole)	****	18	18	18		4171		****		
Mincemeat		5	5	5	****	****		****		
Soups (Tinned)	****	4 2	2	4 2	****			****	****	
Vinegar, Mustard and Pepper	****	6	6	6		****		1118	****	
Meat Pastes		1	1	1			****		****	
Tinned Fish, (Salmon and Brisling).		2	2	2		1119		****		I sample of Tinned Brisling contained a negligible quantity of tin compounds.
Biscuits (Assorted)		15	17	15	2	****			****	The vendors of the 2 samples "not genuine" were cautioned. (For further information see p. 200).
Bacon	****	1	1	1			****	1110	****	
Tripe	7	3	3	3	4111	10000	****			
Sausage		30	37	26	11	7	4	2	1	The samples "not genuine" all contained sulphur dioxide, either undeclared or in excess of the limit allowed. The remaining 4 samples (of those "not genuine") were taken informally (not for proceedings) (See also p. 200).
Table Jellies (and Gelatine)		2	2	2	and and	****		****	****	
Ice Cream		12	12	12	****					Most of the samples were poor in their fat content. No
Household Drugs, etc. :— (Including Glauber Salt, Gregory Powder, Compound Liquorice Powder, Cascara Sagrada, Syrup										action was taken, there being no fixed standard for "ice cream."
of Figs, Castor Oil, Camphorated Oil, Syrup of Rhubarb, Tincture of Rhubarb, Paregoric, Syrup of Squills, Glycerine, Cream of Tartar, Tartaric Acid, Liquid Paraffin, Chemical Food, Crushed Linseed, Eucalyptus Oil, Composition Essence, Sulphur and Zinc Ointments,	4100	53	53	53				3130		2 samples of Glauber Salt had effloresced and were not in accordance with the requirements of the British Pharmacopæa (1932), indicating storage in a place where the temperature had been too high. The attention of the vendors was drawn to the matter.
Cod Liver Oil (and "extracts")	1	0	0	0						
Spirits (Whisky, Rum and Gin)	1	8 8	8 9	8 8	1		****	****		The sample "not genuine" was one of Gin containing 8.76% excess water. This was taken informally, and a subsequent (formal) sample being genuine, the case was met by a caution.
Totals	743	295	1,038*	964	74	32	7	22	3	Amount of Penalties:—£19 10s. 0d.

^{*} Includes 158 samples taken in course of delivery (at railway stations, hospitals, etc.).

1.

FOOD AND DRUGS ADULTERATION, Etc.

Total Samples.—The number of samples of foods and drugs obtained for analysis during the year was 1,038, as against 1,057 in 1935. They were of a varied nature, and included most articles in common use in the household. All of these were submitted to the Public Analyst, who certified that 964 were genuine and 74 not genuine.

Informal Samples.—295 informal samples were taken, as against 294 last year. Legal proceedings cannot be taken if these samples are found not genuine; this method is, however, a useful guide to the general quality of foodstuffs sold in any particular district. Any adulterated samples are followed up by "formal" or "official" samples, so that legal proceedings may be taken if necessary.

Milk Samples.—As usual, the greatest number of samples obtained has been of milk, one of the most important articles of food, and which unfortunately readily lends itself to fraudulent practices. 728 samples were taken, and of these 59 were certified to be below the minimal limits fixed by the "Sale of Milk Regulations, 1901." Of this number 19 were deficient in non-fatty solids, 36 in milk-fat and 4 in both. The percentage of deficiency in fat varied from 0.3 to 33.3 (the average being 7.33), and of solids not fat from 0.7 to 29.7 (average 7.82).

"Appeal to Cow" Samples.—Four farms were visited and, after witnessing the milking operations, 22 samples were taken, and submitted to the Public Analyst for analysis. 18 of these were up to the standard, whilst the remaining 4 fell below. No action could be taken in respect of these 4 samples, as the milk was "as it came from the cow." The deficiencies amounted to 12.0 and 12.6 per cent. in milk-fat and 2.3 and 3.0 per cent. in non-fatty solids, respectively.

Notwithstanding the large number of samples (1,038) of over 90 different articles of foods and drugs, it was only necessary to institute legal proceedings in 32 cases.

Samples not Genuine, etc.—The percentage of all samples not genuine to the total number taken was 7.13 (compared with 2.65 for the previous year). The percentage of non-genuine

milk samples to the total number of milk samples obtained was 8.10 (as against 1.72 in 1935). The total number of samples taken was at the rate of 3.57 per 1,000 of the population (estimated) of the City for the year 1936. This is in excess of the number suggested by the Ministry of Agriculture (viz., 3 per 1,000 of the population).

Margarine.—17 samples of margarine were purchased and analysed. All were genuine, free from preservatives, and in compliance with the requirements of the Act in all other respects.

Margarine Warehouses.—48 visits were made to the registered margarine warehouses in the City. The packages were examined as regards proper marking, and all found to comply with the Act.

Preservatives in Food.—Of the total samples obtained for analysis (1,038), only 44 contained preservative, the quantity in most instances being well within the limits allowed.

26 samples of sausage contained preservative (sulphur dioxide), the quantity in 22 instances being within the permissible limit. In the case of the remaining 4 samples (informal and formal, respectively, from each of two retailers), the vendors were summoned and fined, in each case, 20/- in respect of the excess quantity and 10/- for failing to declare the presence of the preservative. 5 vendors were summoned in respect of failure to give the necessary declaration; one fined 20/-, one 10/-, 2 cases dismissed, and one withdrawn, the case being met by a caution.

2 samples of biscuits ("rum wafers") taken at a warehouse in the City, were found to contain 200 and 100 parts, respectively, of boric acid, contrary to the Regulations. The firm concerned, immediately on becoming aware of the presence of the preservative, withdrew from sale and surrendered their stock of these goods for destruction. The offence was met by a caution.

OFFENCES OTHER THAN ADULTERATION.

Offence.	No. of Cases.	Action Taken, etc.
Milk and Dairies (Consolidation) Act,	ya-tere	
1915; Section 6.— Names and addresses of vendors not inscribed upon milk vessels and/or vehicles.	2	In 1 case offender sum- moned; case dismissed with a caution. The other case was met by a caution.
Milk and Dairies Order, 1926; Section 6 (3).—		
Selling milk without being registered for the purpose.	3	In 2 cases offenders cautioned; in the other summoned, case dis- missed with a caution.
Section 28.— Milk vessels not properly cleansed before being returned.	1	Offender cautioned.
Section 29 (1).— Churn conveying milk in course of delivery to retailer bearing the name and address of a different farmer (on lid and body of churn).	1	Offender cautioned.
Section 29 (2).— Milk churns in a condition contravening the Order. Section 30.—	3	Offenders cautioned.
Selling skimmed milk from churn and vehicle not marked with the words "skimmed milk."	1	Offender summoned; case withdrawn.
Section 33.— Vehicle (barrow) used for conveyance of milk in an offensive condition (from previous conveyance of fish).	1	Offender cautioned.
Milk and Dairies (Amendment) Act, 1922; Sections 2-3.— Milk and Dairies Order, 1926; Section 6.		
The Milk (Special Designations) Order, 1923; Section 3.— Selling "Pasteurised" milk without being registered and licensed for the purpose.	2	Offenders cautioned.
Food and Drugs (Adulteration) Act, 1928; Section 6 (3) (d).— Margarine labelled with unauthor-	2	Offenders cautioned.
ised descriptive names. Section 8.—	-	Offenders cautioned.
Wholesale margarine warehouse not registered.	1	Offender cautioned.
The Milk (Special Designations) Order, 1936; Third Schedule, Part 1, B 3.—		
Churns containing "Tuberculin Tested" milk forwarded unsealed.	1	Offender cautioned.
Total	18	State Halder - State of State

The Public Health (Condensed Milk) Regulations, 1923-1927.

Six samples of condensed milk were obtained. All were genuine and in compliance with the Regulations.

Artificial Cream Act, 1929.

There are two premises (retail shops) on the register and, during the year, no further applications have been received for registration.

BACTERIAL IMPURITY OF MILK AND WATER.

For details of examinations under this heading see pages 104, 105 and 181.

Cleanliness of Milk Churns.—During the year 18,910 churns awaiting return to the farmers were examined at the various railway stations in the City. Of this number, 2 (from 1 dealer) were found in an uncleansed condition. The case was met by a caution.

In addition, 2,958 churns in course of transit through the City were examined; all were found to have been rinsed as required.

Water.—Samples were collected from all parts of the City and at the water works, and examined for the presence of bacillus coli.

The results are described on page 105.

PREMISES ON WHICH FOOD IS PREPARED.

Bakehouses.—There are in the City 268 bakehouses, of which 45 are factories (i.e., places in which mechanical power is used), and 223 are workshops.

The number of "domestic" bakehouses, or private dwelling houses in which the occupier makes bread for sale amongst the neighbours, is 66, a decrease of 12 as compared with 1935. Domestic bakehouses are under the same supervision as when the business is carried on in an ordinary bakehouse, and, generally speaking, are kept in a cleanly state. A few minor contraventions were found, which were immediately remedied on verbal request.

Restaurant Kitchens (including hotels, cafés and dining rooms).

—Regular inspection and strict supervision are exercised over these places, in order to ensure the handling and preparation of food under hygienic conditions. The number on the register is 111 (a decrease of 4 during the year). 9 notices were served on occupiers in regard to lime-washing and general cleansing. In each case the notice was immediately complied with.

Fried Fish Shops.—The number of these is 158 (as against 161 in the previous year). For comments see "Offensive Trades" (Section VII.).

Manufacture and Sale of Ice Cream.—Under Section 4 of the Newcastle upon Tyne Corporation (General Powers) Act, 1935, which came into operation in August last, compulsory registration is now required of all ice cream manufacturers, vendors, dealers, and premises. Applicants who are refused registration are afforded an opportunity of appearing before the Health Committee to show cause why registration should not be refused. If aggrieved by refusal, appeal may be made to a court of summary jurisdiction.

251 applications for registration were received, of which 242 were granted by the Committee, and 9 refused owing to the unsuitability of the premises.

Of those granted, the applications were made as :-

(a)	manufacturers and vendors and as owner-occupiers of the premises	40
(b)	manufacturers and vendors and as occupiers	36
(c)	vendors and as owner-occupiers	46
(d)	,, ,, occupiers	113
(e)	street vendors	6
	owners of premises	

The powers under Section 4 are strictly enforced, ensuring sanitary conditions for the preparation, storage, and handling of this delectable foodstuff. Each year consumption is increasing and, as no legal standard is fixed for the milk-fat content, it is permissible to market this commodity devoid of cream. The time is now overdue for such a standard to be fixed. 12 samples submitted for analysis were found to vary in their milk-fat content between 1.7 and 14 per cent.

The Milk and Dairies (Amendment) Act, 1922, Sec. 2, and The Milk and Dairies Order, 1926, Sec. 6.—During the year 40 applications were received for permission to retail milk, 39 being granted and 1 refused on sanitary grounds. At the close of the year there were 684 retail milk-shops in the City, including 69 belonging to the 10 larger dairy companies. Of the total, 68 were shops in which only dairy products and like commodities were retailed, 576 were shops selling other articles, and 40 were hawkers. All milk-shops and dairies were regularly inspected, and the conditions generally found to be satisfactory.

W. GRAY,

Inspector under the Food and Drugs Acts, etc.

Health Department,

Town Hall,

15th May, 1937.

REPORT OF THE CHIEF SANITARY INSPECTOR.

VII.—THE HOME AND THE WORKSHOP.

NUISANCES, HOUSING, FACTORIES AND WORKSHOPS, Etc.

BUT TO THOSE HIM

VII. THE HOME AND THE

ROISANCES, HOUSING, FASTORIES AND WORKSHOPS, Esc.

NUISANCES, HOUSING, FACTORIES AND WORKSHOPS, ETC.

The following is the Report of the Chief Sanitary Inspector.

TO THE MEDICAL OFFICER OF HEALTH.

SIR,

I have the honour to submit the following report on the work carried out by my section of the Department during the year 1936.

There is no outstanding feature to record. The work, by its very nature, is not spectacular, yet much solid work has been accomplished, with ensuing benefit and contentment to the community.

Inspectorial time, released in the rapid demolition of worn-out and unfit houses under Slum Clearance Orders, has been profitably directed into other channels, the result of which is reflected in the various tabular records.

The attention of the appropriate Committees has been drawn by special report to certain matters arising in connection with the holding of the Temperance Festival on the Town Moor during Race Week.

Suggestions have been made with a view to the improvement of existing arrangements with respect to the sale of foodstuffs and the matters of water supply, sanitary conveniences, and nuisances, which will in no wise interfere with this very popular Festival, which attracted over half-a-million persons during the eight days on which it was held.

New legislation in the *Pharmacy and Poisons Act*, 1933— Listed Sellers of Part II. Poisons came into operation on the 1st May. Briefly, the provisions affecting a local authority are: registration of premises, and the control of storage, transport, labelling and sale of certain specified poisons.

NUISANCES.

The number of nuisances reported upon and dealt with during the year was 21,773, which is an increase of 2,663 upon the previous year.

Analysis reveals, however, that the variation is much in accord with previous years. It is satisfactory to note that less action has had to be taken against the dirty class of tenant, whose actions and lack of action create so much trouble to all.

It is particularly noted, in the complaints received at the Department, the increasing number of tenants of poor type property who desire Council houses.

Notices Served.

The following are the numbers of notices and letters issued during the year:—

Number of notices served :-		
Informal	4,365	
Statutory	340	
•		4,705
Number of special letters sent		2,664
Number of circular letters sent		2,128
Total		9,497

Magisterial Proceedings.

It was necessary to report 75 cases of statutory "Notices not complied with" to the Health Committee, who ordered legal proceedings to be taken, but in only 2 cases were summonses issued. The work in the other instances was carried out before making application for summonses.

The details of this part of the work are given on page 220.

The Rent and Mortgage Interest Restrictions Acts.

No application was received from tenants for a certificate under the above Acts certifying that the house was not reasonably fit for habitation.

Conversion of "Dry" Closets to Water-closets.

The conversion of these conveniences to water-closets, under Section 36, Public Health Act, 1875, has proceeded in a satisfactory manner. During the year, 21 pail-closets have been converted, also 4 "cell" privies, and 13 combined privies and ashpits. In addition, 17 pail-closets were abolished in the demolition of Slum Areas. Ten years ago upwards of 1,800 privies were in existence, whereas to-day only 9 remain within the boundaries of the City as before the recent extension. These remaining closets are generally in good structural condition and, in certain instances, no sewer is available to enable conversion to be effected.

In addition, 25 "waste-water" closets have been abolished and replaced by up-to-date conveniences. These "waste-water" closets consist simply of a trapped stoneware shaft of 12 in. to 18 in. diameter in the w.c., and from 4 ft. to 9 ft. deep. The solids in the trap are flushed away by slop- and rain-water from the yard drains, the shafts (usually caked with excreta) being only cleansable by scraping. Unfortunately, this type of closet, which is never free from effluvia, is deemed to be, under existing law, on the "water-carriage" system. The conversion of these to proper water-closets must of necessity be slow.

Seven dry ashpits were also removed and, in three instances, were replaced by regulation dustbins (the first of which was supplied free of charge). 29 free bins were supplied during the year.

RETURN OF PRIVIES, ETC., IN THE VARIOUS WARDS OF THE CITY.

Wards.	Total No. Privies.	Pail- Closets.	" Cell " Privies.	Combined Privies and Ashpits.
St. Nicholas'				
St. Thomas'		9		46
St. John's				
Stephenson	2444		****	
Armstrong	****	1111	****	****
Elswick		****		****
Westgate	****		****	
Arthur's Hill				****
Benwell	4	1	3	****
Fenham	49	5	****	44
All Saints'		****		
St. Andrew's		****	****	****
Jesmond		****	22	
Dene	19		17	2
Heaton		****		****
Byker	****	****		
St. Lawrence			****	
St. Anthony's		****		
Walker	1	****		1
Total in City	128	15	20	93

Smoke Abatement.

Under the provisions of the *Public Health* (Smoke Abatement) Act, 1926, a byelaw was made and came into operation on the 1st June, allowing 3 minutes' emission of black smoke during a period of 30 minutes, anything above this being deemed to be an offence and a nuisance. Whenever this permissible amount is found to be exceeded, and also where a heavy emission of "medium" smoke is observed, the cause is inquired into and advice given, wherever possible, to remedy the fault.

It is pleasing to record increased activity in the number of observations, together with a decreased number of offences, as compared with the previous year.

595 observations were made of 100 factory and other chimneys. In addition there were 104 observations of the chimney at the City Hospital for Infectious Diseases, Walker Gate.

15 informal notices were served during the year. Drivers of 2 steam wagons were verbally cautioned about the quantity of black smoke given off whilst proceeding through the City.

The following table gives details as to smoke inspection:-

No. of chimneys watched.	No. of observations made.	No, of chimneys from which black smoke issued in such quantity as to	No. of times when smoke issued so as to be a nuisance.	No. of served abatement nuisa	No. of Prosecu- tions.	
wateried.	made.	be a nuisance.	be a nuisance.	Informal.	Statutory.	tions.
100	595	11	27	15		

Note.—Not including 104 observations and 39 excessive emissions at the City Hospital, Walker Gate.

ATMOSPHERIC POLLUTION.—Newcastle Records, 1936.
TOWN MOOR GAUGE.

211

	es).			METR	IC TON KILO	s of D metre	EPOSIT PER MO	PER SO	UARE		
Month.	(Millimetres).	Insol	uble Ma	atter.		uble tter.	DS.		Includ Soluble		
	RAIN (M	Tar.	Other Car- bonaceous.	Ash.	Loss on Ignition.	Ash.	TOTAL SOLIDS.	Sulphate as S.O3.	Chlorine as Cl.	Ammonia as N.H3.	Lime as Ca.O.
January February March April May June July August Sept October Nov Dec	79.5 99.4 54.4 71.0 99.4 78.1 71.0 83.8 42.6 102.2	0.14 0.11 0.20 0.04 0.12 0.13 0.10 0.11 0.10 0.23 0.03 0.64	1.85 1.42 1.94 1.48 1.89 1.96 1.62 1.18 1.14 0.87 1.02 1.62	1.96 2.39 2.30 1.86 2.43 1.52 1.43 1.35 1.03 1.23 1.01 1.93	0.91 2.07 2.39 2.12 1.28 1.39 1.87 1.70 2.19 1.96 2.46 2.04	3.63 3.98 3.18 2.54 3.83 3.18 1.72 1.99 2.17 2.30 3.26 1.79	8.49 9.97 10.01 8.04 9.55 8.18 6.74 6.33 6.63 6.59 7.78 8.02	1.48 1.80 1.70 0.80 1.02 0.75 0.75 0.72 1.09 0.85 1.41 1.02	0.64 0.80 0.92 0.67 0.50 0.60 0.28 0.26 0.48 0.94 1.05 0.37	0.09 0.09 0.10 0.06 0.06 0.09 0.10 0.07 0.13 0.07 0.26 0.23	0.18 0.20 0.17 0.14 0.06 0.33 0.07 0.11 0.34 0.28 0.34
Total, 12 months	937.6	1.95	17.99	20.44	22.38	33.57	96.33	13.39	7.51	1.35	2.46
Average per month	78.1	0.16	1.50	1.70	1.87	2.80	8.03	1.20	0.63	0.11	0.2

An average of 8.03 metric tons per square kilometre per month=7.7 cwts. per acre per annum, or 247 tons per square mile per annum, as compared with 7.1 cwts. per acre, or 229 tons per square mile in 1935.

WESTGATE CEMETERY GAUGE.

212

	es)			METR			PER MO		UARE		
MONTH.	llimetr	Insol	uble M	atter.		uble tter.	DS.		Include Soluble	led in Matter.	
	Rain (Millimetres)	Tar.	Other Car- bonaceous.	Ash.	Loss on Ignition.	Ash.	TOTAL SOLIDS.	Sulphate as S.O3.	Chlorine as Cl.	Ammonia as N.H3.	Lime as Ca.O.
January February March April May June July August Sept October Nov Dec	60.1 70.8 38.7 53.4 76.2 63.7 60.1 65.5 42.7	0.12 0.19 0.28 0.05 0.08 0.17 0.08 0.33 0.11 0.24 0.08 0.13	5.22 2.63 3.89 2.29 2.38 2.25 2.26 1.43 2.02 2.15 2.10	4.44 3.15 4.22 3.30 3.43 2.74 2.91 2.02 1.49 2.75 2.74 2.38	3.09 1.44 1.42 0.85 1.07 0.91 0.96 0.96 1.31 1.71 1.92 0.48	6.73 3.25 3.39 1.94 2.46 2.28 1.93 1.32 1.96 2.31 1.76 2.08	19.60 10.66 13.20 8.43 9.42 8.35 8.14 6.06 6.30 9.03 8.65 7.17	3.07 1.15 1.46 0.69 0.99 0.73 0.65 0.87 0.86 0.88 1.10 1.01	0.90 0.64 0.65 0.52 0.41 0.43 0.31 0.17 0.47 1.03 0.73 0.40	0.17 0.09 0.08 0.04 0.05 0.01 0.04 0.01 0.07 0.08 0.24 0.08	0.47 0.29 0.41 0.33 0.48 0.37 0.07 0.15 0.27 0.38 0.48
Total, 12 months	791.8	1.86	30.05	35.57	16.12	31.41	115.01	13.46	6.66	0.96	4.28
Average per month	66.0	0.16	2.50	2.96	1.34	2.62	9.58	1.12	0.56	0.08	0.3

An average of 9.58 metric tons per square kilometre per month=9.2 cwts. per acre per annum, or 294 tons per square mile per annum, as compared with 9.2 cwts. per acre, or 295 tons per square mile in 1935.

213
WELBECK RESERVOIR GAUGE.

Month.	res).			MEIK			PER Me	PER SO	OAKE		
	üllimet	Insol	luble M	atter.		uble tter.	TOTAL SOLIDS.		Includ Soluble	led in Matter	
	RAIN (Millimetres).	Tar.	Other Car- bonaceous.	Ash.	Loss on Ignition.	Ash.		Sulphate as S.O3.	Chlorine as Cl.	Ammonia as N.H3.	Lime as Ca.O.
January February March April May June July August Sept October Nov Dec.	39.1 50.8 75.5 63.8 62.5 63.8 23.4 78.1	0.04 0.09 0.19 0.08 0.14 0.15 0.12 0.14 0.21 0.21 0.23 0.26	1.56 1.25 1.41 1.39 1.85 1.21 1.82 1.41 1.52 0.86 1.81 0.99	2.02 2.92 2.80 3.29 3.48 3.57 2.35 2.20 2.07 2.21 2.79 2.26	1.38 1.82 1.64 1.17 1.31 1.07 1.41 0.87 1.02 0.79 2.03 1.73	3.03 3.28 3.05 2.34 3.15 2.11 1.65 1.38 2.04 1.26 3.59 3.55	8.03 9.36 9.09 8.27 9.93 8.11 7.35 6.00 6.86 5.33 10.45 8.79	1.28 1.38 1.29 0.73 0.98 0.62 0.82 0.77 0.74 0.62 1.29 1.17	0.64 1.00 0.90 0.78 0.72 0.64 0.30 0.17 0.49 0.42 1.38 0.61	0.10 0.09 0.10 0.07 0.08 0.04 0.03 0.03 0.03 0.04 0.20 0.09	0.17 0.34 0.34 0.29 0.34 0.38 0.31 0.16 0.27 0.39 0.53
Total, 12 months	668.0	1.86	17.08	31.96	16.24	30.43	97.57	11.69	8.05	0.90	3.68
Average per month	55.7	0.16	1.42	2.66	1.35	2.54	8.13	0.99	0.67	0.80	0.31

An average of 8.13 metric tons per square kilometre per month=7.8 cwts. per acre per annum, or 250 tons per square mile, per annum, as compared with 6.5 cwts. per acre, or 207 tons per square mile in 1935.

214
TOTAL IN THREE GAUGES IN THE CITY, 1936.

	tres)				Kilo	METRE	PER MO	ONTH.			
MONTH.	illime	Inso	luble M	atter.		uble tter.	TOTAL SOLIDS.		Includ Soluble		
	RAIN (Millimetres).	Tar.	Other Car- bonaceous.	Ash.	Loss on Ignition.	Ash.		Sulphate as S.O3.	Chlorine as Cl.	Ammonia as N.H3.	Lime as Ca.O.
Total, 12 months	2397.4	5.67	65.12	87.97	54.74	95.41	308.91	38.54	22.22	3.21	10.42
Average per month	199.8	0.47	5.43	7.33	4.56	7.95	25.74	3.21	1.85	0.27	0.87
Average per gauge 12 mths.	799.1	1.89	21.71	29.32	18.25	31.80	102.97	12.85	7.41	1.07	3.47
Average per gauge per month	66.6	0.16	1.81	2.44	1.52	2.65	8.58	1.07	0.62	0.09	0.29

An average of 8.58 metric tons per square kilometre per month=8.3 cwts. per acre per annum, or 264 tons per square mile per annum, as compared with 7.6 cwts. per acre, or 244 tons per square mile in 1935.

For comparison with the foregoing, the following returns of sunshine recorded at the Armstrong College, Newcastle, and at Cockle Park, near Morpeth (about 15 miles from the City), are given:—

Month.	Armstrong College. Sunshine (hours).	Cockle Park. Sunshine (hours).
January	45.4	56.5
February	50.3	65.4
March.	46.5	51.2
April	154.2	203.5
May	122.4	174.6
June		209.8
July		157.3
August		186.3
September	74.8	85.8
October	86.5	105.0
November	48.1	70.0
December	47.2	62.3
Total for year	1154.9	1427.7
Average per month	96.2	119.0

CINEMAS, THEATRES, AND OTHER PLACES OF PUBLIC ENTERTAINMENT.

By a Ministry of Health Circular issued in 1920, Sanitary Authorities are required to give particular attention to premises holding a licence for music, dancing, etc., special regard having to be given to sanitary conveniences, dressing rooms, ventilation, and means of escape in case of fire.

In pursuance of this order, 6 applications were received for certificates of sanitation which must be submitted to the Licensing Justices before a music or dancing licence is granted. After a careful inspection of the premises all were granted.

The number of places so certified is now 4 theatres and music halls, 38 cinemas, and 119 concert halls, billiard rooms, cafés, etc., 173 visits were made both during the day and night time to inspect the sanitary arrangements, dressing rooms, etc., which were generally found to be in order.

Testing (with the "Kata" thermometer) of the air and ventilating system of all theatres, music halls and cinemas has been carried out. In 14 cinemas, conditions were found to be unsatisfactory. This state was immediately brought to the attention of the reponsible persons, who rectified matters. The trouble was due to inattention to the varying climatic conditions (alternating hot and cold spells) and the proper use of efficient appliances.

In all other cases the conditions were satisfactory.

In addition, tests for demonstration purposes were made at the Durham University College of Medicine and at the Rutherford College.

OFFENSIVE TRADES.

14 applications for permission to establish the trade of a fish fryer were received during the year. Of that number 5 were granted and 7 refused, the proposed premises not being suitable for the purpose, whilst 2 applications were withdrawn.

This class of "offensive trade" still predominates, there being now 158 on the register (against 161 last year). As compared with the previous year there is an increase of 1 gut scraper, and a decrease of 3 fish fryers, 1 rag and bone dealer, 1 tripe boiler, and 1 bone boiler.

The fried fish shops are regularly inspected by day and occasionally at night-time. It was necessary to serve notices for the abatement of nuisances in only 3 cases.

With the exception of a few minor contraventions, all the offensive trades have been conducted in a satisfactory manner.

In consequence of receipt of complaints regarding the untidy condition of streets in the vicinity of fried fish shops, due to the deposit thereon of greasy wrapping papers, etc., "Litter" notices were sent to and are displayed in all these shops, directing the attention of the public to the byelaw in respect of litter in streets. A satisfactory improvement followed the issue of these notices in the affected districts.

The number of offensive trades now on the register is :-

Fish fryers	158
Rag and bone dealers	8
Tripe boilers	6
Gut scrapers	7
Dealers in hides and skins	
Bone boilers	2
Fat melters and extractors	1
Glue and size makers	2
Soap boiler	1

All are systematically inspected, 1,402 such visits being made during the year.

Summary of Nuisances, etc., for the Abatement of which Notices were Served during 1936.

Foul pail-closets (to replace with water-closets). Foul privies and ashpits (to replace with water-closets). Defective waste-water closets (to replace with fresh-water closets with flushing cisterns, etc.). Foul or defective ashpits not connected with privies (to remove and provide dust-bins). Insufficient water-closet accommodation (additional water-closets ordered) Defective or insufficient dust-bins (for houses). """ (for business premises). Defective water-closets Water-closets without adequate water supply. Choked water-closets (mostly served on tenants). Dirty water-closets (all served on tenants). Defective drains (to repair, or construct new drains). Insufficient means of drainage. Rainwater fall-spouts connected directly with drains. Sunsufficient ventilation to drains. Choked drains, etc Defective, want of, or choked sinks, waste-pipes, etc No sink provided. Defective or choked soil-pipes, vent shafts, etc 215 Sink waste-pipes not trapped. Want of or defective pavement in yards, passages, etc Boirty padding. Damp rooms. Dirty yards, passages, stairs, etc Dirty yards, passages, stairs, etc Animals, pigeons, and fowls improperly kept Offensive accumulations. Accumulations of manure Want of or defective manure pits Broken roofs and want of or defective or choked spouting. Parken roofs and want of or defective or choked spouting. Parken roofs and want of or defective or choked spouting. Parken roofs and want of or defective or choked spouting. Parken roofs and want of or defective or choked spouting. Parken roofs and want of or defective or choked spouting. Parken roofs and want of or defective or choked spouting. Parken roofs and want of or defective or choked spouting. Parken roofs and want of or defective or choked spouting. Parken roofs and want of or defective or choked spouting. Parken roofs and want of or defective or choked spouting. Parken roofs and want of or defective or choked spouting. Parken roofs and want of or defective or choked spou
Foul privies and ashpits (to replace with water-closets) Defective waste-water closets (to replace with fresh-water closets with flushing cisterns, etc.). Foul or defective ashpits not connected with privies (to remove and provide dust-bins). Insufficient water-closet accommodation (additional water-closets ordered) Defective or insufficient dust-bins (for houses) Defective water-closets Water-closets Water-closets (mostly served on tenants) Dirty water-closets (mostly served on tenants) Dirty water-closets (all served on tenants) Defective drains (to repair, or construct new drains) Ensufficient means of drainage Rainwater fall-spouts connected directly with drains Insufficient ventilation to drains Choked drains, etc. Defective, want of, or choked sinks, waste-pipes, etc. No sink provided Defective or choked soil-pipes, vent shafts, etc. 21 Sink waste-pipes not trapped Want of or defective pavement in yards, passages, etc. Birty yards, passages, stairs, etc. Dirty yords. Dirty yards, passages, stairs, etc. Dirty yards, passages, and fowls improperly kept Offensive accumulations Accumulations of manure Want of or defective manure pits Broken roofs and want of or defective or choked spouting Smoke nuisances Want of proper ventilation to rooms (including to floor space), broken window cords, etc. Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) Cisterns supplying water to sinks, etc. dirty or defective
Defective waste-water closets (to replace with fresh-water closets with flushing cisterns, etc.). Foul or defective ashpits not connected with privies (to remove and provide dust-bins). Insufficient water-closet accommodation (additional water-closets ordered) Defective or insufficient dust-bins (for houses) Defective water-closets Water-closets without adequate water supply Choked water-closets (mostly served on tenants) Dirty water-closets (all served on tenants) Defective drains (to repair, or construct new drains) Ensufficient means of drainage Rainwater fall-spouts connected directly with drains Insufficient ventilation to drains Choked drains, etc. Defective, want of, or choked sinks, waste-pipes, etc. No sink provided Defective or choked soil-pipes, vent shafts, etc. 21 Sink waste-pipes not trapped Want of or defective pavement in yards, passages, etc. Dirty bedding Damp rooms Dirty yards, passages, stairs, etc. Animals, pigeons, and fowls improperly kept Offensive accumulations Accumulations of manure Want of or defective manure pits Broken roofs and want of or defective or choked spouting Want of water (other than in tenements—see below). Smoke nuisances Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) Cisterns supplying water to sinks, etc. dirty or defective 39 Page of the move and of the common of the defective or defective 9989 Cisterns supplying water to sinks, etc. dirty or defective
with flushing cisterns, etc.). Foul or defective ashpits not connected with privies (to remove and provide dust-bins). Insufficient water-closet accommodation (additional water-closets ordered). Defective or insufficient dust-bins (for houses). , , , , (for business premises). Defective water-closets. Water-closets without adequate water supply. Choked water-closets (mostly served on tenants). Dirty water-closets (all served on tenants). Dirty water-closets (all served on tenants). Defective drains (to repair, or construct new drains). Rainwater fall-spouts connected directly with drains. Insufficient means of drainage. Rainwater fall-spouts connected directly with drains. Insufficient ventilation to drains. Choked drains, etc
and provide dust-bins). Insufficient water-closet accommodation (additional water-closets ordered) Defective or insufficient dust-bins (for houses), (for business premises), (for business premises, (for business premises), (for business premises), (for bu
and provide dust-bins). Insufficient water-closet accommodation (additional water-closets ordered) Defective or insufficient dust-bins (for houses), (for business premises), (for business premises, (for business premises), (for business premises), (for bu
and provide dust-bins). Insufficient water-closet accommodation (additional water-closets ordered) Defective or insufficient dust-bins (for houses), (for business premises), (for business premises, (for business premises), (for business premises), (for bu
Insufficient water-closet accommodation (additional water-closets ordered) Defective or insufficient dust-bins (for houses) , , , , , , (for business premises) Defective water-closets , , , , , , , , (for business premises) Defective water-closets without adequate water supply 44 Choked water-closets (mostly served on tenants) Dirty water-closets (all served on tenants) Defective drains (to repair, or construct new drains) Defective drains (to repair, or construct new drains) Examinater fall-spouts connected directly with drains Insufficient means of drainage Rainwater fall-spouts connected directly with drains Insufficient ventilation to drains Choked drains, etc. Defective, want of, or choked sinks, waste-pipes, etc. No sink provided Defective or choked soil-pipes, vent shafts, etc. Sink waste-pipes not trapped Want of or defective pavement in yards, passages, etc. Dirty rooms. Dirty bedding Damp rooms Damp rooms Page Animals, pigeons, and fowls improperly kept Offensive accumulations Accumulations of manure Want of or defective manure pits Broken roofs and want of or defective or choked spouting Want of water (other than in tenements—see below). Smoke nuisances Want of proper ventilation to rooms (including to floor space), broken window cords, etc. 939 Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) Cisterns supplying water to sinks, etc., dirty or defective.
closets ordered) Defective or insufficient dust-bins (for houses) """, (for business premises) Defective water-closets Water-closets without adequate water supply Choked water-closets (mostly served on tenants) Dirty water-closets (all served on tenants) Defective drains (to repair, or construct new drains) Insufficient means of drainage Rainwater fall-spouts connected directly with drains Insufficient ventilation to drains Choked drains, etc Defective, want of, or choked sinks, waste-pipes, etc. Sink provided Defective or choked soil-pipes, vent shafts, etc. Sink waste-pipes not trapped Want of or defective pavement in yards, passages, etc. Dirty bedding Damp rooms Dirty yards, passages, stairs, etc. Animals, pigeons, and fowls improperly kept Offensive accumulations Accumulations of manure Want of or defective manure pits Broken roofs and want of or defective or choked spouting Want of poper ventilation to rooms (including to floor space), broken window cords, etc. Six yards, passay, stairs, walls, fireplaces, etc.) Scieterns supplying water to sinks, etc. dirty or defective
Defective or insufficient dust-bins (for houses) 1,235 , , , , , , , , , , , , , , , , , , ,
Defective water-closets
Water-closets without adequate water supply Choked water-closets (mostly served on tenants) Dirty water-closets (all served on tenants) Defective drains (to repair, or construct new drains) Rainwater fall-spouts connected directly with drains Insufficient ventilation to drains Choked drains, etc. Defective, want of, or choked sinks, waste-pipes, etc. No sink provided Defective or choked soil-pipes, vent shafts, etc. Sink waste-pipes not trapped Want of or defective pavement in yards, passages, etc. Birty bedding Damp rooms Dirty bedding Damp rooms Dirty yards, passages, stairs, etc. Animals, pigeons, and fowls improperly kept Offensive accumulations Accumulations of manure Want of or defective manure pits Broken roofs and want of or defective or choked spouting Want of water (other than in tenements—see below) Smoke nuisances Want of proper ventilation to rooms (including to floor space), broken window cords, etc. Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) Sink provided Same roofs and wanter to sinks, etc., dirty or defective 344 445 446 447 448 448 449 449 449 440 440 440
Water-closets without adequate water supply Choked water-closets (mostly served on tenants) Dirty water-closets (all served on tenants) Defective drains (to repair, or construct new drains) Rainwater fall-spouts connected directly with drains Insufficient ventilation to drains Choked drains, etc. Defective, want of, or choked sinks, waste-pipes, etc. No sink provided Defective or choked soil-pipes, vent shafts, etc. Sink waste-pipes not trapped Want of or defective pavement in yards, passages, etc. Birty bedding Damp rooms Dirty bedding Damp rooms Dirty yards, passages, stairs, etc. Animals, pigeons, and fowls improperly kept Offensive accumulations Accumulations of manure Want of or defective manure pits Broken roofs and want of or defective or choked spouting Want of water (other than in tenements—see below) Smoke nuisances Want of proper ventilation to rooms (including to floor space), broken window cords, etc. Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) Sink provided Same roofs and wanter to sinks, etc., dirty or defective 344 445 446 447 448 448 449 449 449 440 440 440
Choked water-closets (mostly served on tenants) Dirty water-closets (all served on tenants) Defective drains (to repair, or construct new drains) Insufficient means of drainage Rainwater fall-spouts connected directly with drains Insufficient ventilation to drains Choked drains, etc. Defective, want of, or choked sinks, waste-pipes, etc. No sink provided Defective or choked soil-pipes, vent shafts, etc. Sink waste-pipes not trapped Want of or defective pavement in yards, passages, etc. Dirty rooms Dirty bedding Damp rooms Dirty yards, passages, stairs, etc. Animals, pigeons, and fowls improperly kept Offensive accumulations Accumulations of manure Want of or defective manure pits Broken roofs and want of or defective or choked spouting Smoke nuisances Want of proper ventilation to rooms (including to floor space), broken window cords, etc. Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) Cisterns supplying water to sinks, etc., dirty or defective 13 Particular defective 14 15 16 17 18 18 19 19 19 19 19 19 19 19
Dirty water-closets (all served on tenants) Defective drains (to repair, or construct new drains) Insufficient means of drainage Rainwater fall-spouts connected directly with drains Insufficient ventilation to drains Choked drains, etc. Defective, want of, or choked sinks, waste-pipes, etc. No sink provided Defective or choked soil-pipes, vent shafts, etc. Sink waste-pipes not trapped Want of or defective pavement in yards, passages, etc. Dirty rooms Dirty bedding Damp rooms Total Dirty yards, passages, stairs, etc. Animals, pigeons, and fowls improperly kept Toffensive accumulations Accumulations of manure Want of or defective manure pits Broken roofs and want of or defective or choked spouting Smoke nuisances Want of proper ventilation to rooms (including to floor space), broken window cords, etc. Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) Cisterns supplying water to sinks, etc., dirty or defective
Defective drains (to repair, or construct new drains) Insufficient means of drainage Rainwater fall-spouts connected directly with drains Insufficient ventilation to drains Choked drains, etc. Defective, want of, or choked sinks, waste-pipes, etc. No sink provided Defective or choked soil-pipes, vent shafts, etc. Sink waste-pipes not trapped Want of or defective pavement in yards, passages, etc. Dirty bedding Damp rooms Dirty yards, passages, stairs, etc. Animals, pigeons, and fowls improperly kept Offensive accumulations Accumulations of manure Want of or defective manure pits Broken roofs and want of or defective or choked spouting Want of water (other than in tenements—see below). Smoke nuisances Vant of proper ventilation to rooms (including to floor space), broken window cords, etc. Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) Cisterns supplying water to sinks, etc., dirty or defective.
Insufficient means of drainage Rainwater fall-spouts connected directly with drains Insufficient ventilation to drains Choked drains, etc. Defective, want of, or choked sinks, waste-pipes, etc. No sink provided Defective or choked soil-pipes, vent shafts, etc. Sink waste-pipes not trapped Want of or defective pavement in yards, passages, etc. Birty rooms Dirty bedding Damp rooms Dirty yards, passages, stairs, etc. Animals, pigeons, and fowls improperly kept Offensive accumulations Accumulations of manure Want of or defective manure pits Broken roofs and want of or defective or choked spouting Want of water (other than in tenements—see below). Smoke nuisances Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) Cisterns supplying water to sinks, etc., dirty or defective. 38 38 38 38 38 38 38 38 38 38 38 38 38
Rainwater fall-spouts connected directly with drains Insufficient ventilation to drains Choked drains, etc. Defective, want of, or choked sinks, waste-pipes, etc. No sink provided Idefective or choked soil-pipes, vent shafts, etc. Sink waste-pipes not trapped Want of or defective pavement in yards, passages, etc. Dirty rooms Dirty bedding Bamp rooms Dirty yards, passages, stairs, etc. Animals, pigeons, and fowls improperly kept Offensive accumulations Accumulations of manure Want of or defective manure pits Broken roofs and want of or defective or choked spouting Smoke nuisances Want of proper ventilation to rooms (including to floor space), broken window cords, etc. Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) Cisterns supplying water to sinks, etc., dirty or defective. 38 38 38 38 30 38 30 30 30 30 30 30 30 30 30 30 30 30 30
Insufficient ventilation to drains
Choked drains, etc
Defective, want of, or choked sinks, waste-pipes, etc. No sink provided
No sink provided Defective or choked soil-pipes, vent shafts, etc
No sink provided Defective or choked soil-pipes, vent shafts, etc
Sink waste-pipes not trapped
Sink waste-pipes not trapped
Want of or defective pavement in yards, passages, etc. Dirty rooms
Dirty rooms 26 Dirty bedding 8 Damp rooms 762 Dirty yards, passages, stairs, etc. 52 Animals, pigeons, and fowls improperly kept 77 Offensive accumulations 48 Accumulations of manure 13 Want of or defective manure pits 9 Broken roofs and want of or defective or choked spouting 2,164 Want of water (other than in tenements—see below) 132 Smoke nuisances 11 Want of proper ventilation to rooms (including to floor space), broken window cords, etc. 939 Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) 9,982 Cisterns supplying water to sinks, etc., dirty or defective 3
Dirty bedding 8 Damp rooms 762 Dirty yards, passages, stairs, etc. 52 Animals, pigeons, and fowls improperly kept 77 Offensive accumulations 48 Accumulations of manure 13 Want of or defective manure pits 9 Broken roofs and want of or defective or choked spouting 2,164 Want of water (other than in tenements—see below) 132 Smoke nuisances 11 Want of proper ventilation to rooms (including to floor space), broken window cords, etc. 939 Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) 9,982 Cisterns supplying water to sinks, etc., dirty or defective 3
Damp rooms 762 Dirty yards, passages, stairs, etc. 52 Animals, pigeons, and fowls improperly kept 77 Offensive accumulations 48 Accumulations of manure 13 Want of or defective manure pits 9 Broken roofs and want of or defective or choked spouting 2,164 Want of water (other than in tenements—see below) 132 Smoke nuisances 11 Want of proper ventilation to rooms (including to floor space), broken window cords, etc. 939 Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) 9,982 Cisterns supplying water to sinks, etc., dirty or defective 3
Dirty yards, passages, stairs, etc. 52 Animals, pigeons, and fowls improperly kept 77 Offensive accumulations 48 Accumulations of manure 13 Want of or defective manure pits 99 Broken roofs and want of or defective or choked spouting 2,164 Want of water (other than in tenements—see below) 132 Smoke nuisances 11 Want of proper ventilation to rooms (including to floor space), broken window cords, etc. 939 Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) 9,982 Cisterns supplying water to sinks, etc., dirty or defective 3
Animals, pigeons, and fowls improperly kept 7 Offensive accumulations 48 Accumulations of manure 13 Want of or defective manure pits 9 Broken roofs and want of or defective or choked spouting 2,164 Want of water (other than in tenements—see below) 132 Smoke nuisances 11 Want of proper ventilation to rooms (including to floor space), broken window cords, etc. 939 Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) 9,982 Cisterns supplying water to sinks, etc., dirty or defective 3
Offensive accumulations
Accumulations of manure
Want of or defective manure pits 9 Broken roofs and want of or defective or choked spouting 2,164 Want of water (other than in tenements—see below) 132 Smoke nuisances 11 Want of proper ventilation to rooms (including to floor space), broken window cords, etc. 939 Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) 9,982 Cisterns supplying water to sinks, etc., dirty or defective 3
Broken roofs and want of or defective or choked spouting
Want of water (other than in tenements—see below). Smoke nuisances
Smoke nuisances
Want of proper ventilation to rooms (including to floor space), broken window cords, etc. Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.)
broken window cords, etc. 939 Structural defects—internal and external—(broken plaster, floors, stairs, walls, fireplaces, etc.) 9,982 Cisterns supplying water to sinks, etc., dirty or defective 3
floors, stairs, walls, fireplaces, etc.) 9,982 Cisterns supplying water to sinks, etc., dirty or defective 3
floors, stairs, walls, fireplaces, etc.) 9,982 Cisterns supplying water to sinks, etc., dirty or defective 3
Cisterns supplying water to sinks, etc., dirty or defective
Cisterns supplying water to sinks, etc., dirty or defective
Stables (unsuitable, defective, etc.)
Piggeries (,, ,,)
Food manufactured or stored for sale under improper conditions 10
Bakehouses—Dirty, etc. 90
Fried fish shops—(Plaster and pointing defective, and dust-bins
required) 3
Schools (Council and other) :-
Foul privies (2) and ashpit
Urinal insufficiently drained
Dust-bins required
Sink waste-pipe choked and defective
Water tap defective
rater tap detective
Carried forward 19,97
Carried forward

SUMMARY OF NUISANCES, ETC.—Continued.

Brought forward	19,971
Cellar dwellings illegally occupied.	3
Inadequate accommodation for :-	
* Storage of food (other than in tenements)	432
Cooking of food (,, ,,)	105
Washing of clothes (,, ,,)	82
Water supply and sink not conveniently accessible	
(other than in tenements)	139
Rooms, staircases, etc., insufficiently lighted (other	
than in tenements)	27
Tenements—Limewashing not done .	18
No adequate accommodation for washing of clothes	21
,, storage of food	67
,, ,, preparation and	
cooking of food	11
Water supply and sinks not adequate, conveniently	
accessible, etc	104
Water supply (only) not adequate, conveniently	
accessible, etc.	16
Insufficient number of water-closets provided	5
Inadequate lighting of common staircases—	
Natural20	
Artificial59	79
Staircases without proper handrails, etc	4
Cinemas—(Dust-bins required)	1
Temperature excessive	14
Tents, vans, sheds, and similar structures—(Sanitary arrange-	1.1
ments unsatisfactory)	1
Ice Cream (Floor of manufactory broken)	î
Premises unregistered	2
Dairy (floor dirty)	1
Shops Act, 1934; Sec. 10—(Defects and contraventions)	84
"Offensive trade" established without consent	1
Sites of demolished houses not properly cleared	246
Derelict buildings (unsafe)	4
Unclassified minor nuisances	334
Chomostica millor nuisances r	334
TOTAL	21.775
	,,,,,

Details Relating to certain Works carried out in the Abatement of Nuisances and to Inspections made during 1936.

Length (in yards) of old drains removed	1,707
Length (in yards) of new drains constructed	2,510
New trapped gullies provided to drains	466
Combined Privies and ashpits removed—Privies	13
Ashpits	11
"Cell" privies removed	4
Pail-closets removed	38
Defective water-closets removed	155
Water-closets provided (in place of the foregoing privies and	
defective water-closets removed, also in 25 cases where the	
accommodation was previously insufficient)	213
"Chemical" closets provided (no sewer being available)	2
Dry ash-pits removed and replaced by galvanised iron dust-bins	7
Dust-bins substituted for dry ash-pits where water-closets	
existed, and provided in cases where privies have been	2200
replaced by water-closets	*29
No. of drains tested	1,087
No. of tests of above drains made by smoke and water	1,111
No. of inspections from complaints made at office (verbally or	
by letter)	5,085
No. of tenement inspections made.	7,696
No. of contraventions of Tenement Bye-laws for which notices	500
have been served to obtain remedy	568
Inspections of houses made from complaints received outdoors	
or nuisances discovered in the districts, including a large	
number of minor nuisances, such as choked drains and	
dirty yards, the abatement of which was accomplished at	2,436
the time of visit, and without legal notice	14,384
Supervisions of work in progress	8,354
Common yards and courts in the worst localities specially	0,004
visited on Friday afternoons and Saturday mornings to	
obtain weekly cleansing	16,551
Inspections after infectious disease	1,446
Inspections of milk shops and ice creameries (including retail	1,110
shops)	1,628
,, bakehouses	†1,537
offensive trades	1,402
wholesale margarine warehouses	48
,, as to limewashing of tenements	1,295
,, schools	181
,, shops (re Shops Act, 1934)	452
public houses	74
under Housing Acts	11,368
", re overcrowding (Preliminary Survey)	80,660
Inspection of cinemas, etc. (day visits, 124; night visits, 49)	173
,, Tents, vans, sheds and similar structures	57
Miscellaneous visits	3,082

^{*} Dust bins supplied free by Corporation.
† Including 1,316 inspections made under the Factory and Workshop Acts by the Assistant Inspectors of Workshops.

Summary of Legal Proceedings ordered to be taken before the Magistrates for the Abatement of Nuisances, etc., during the Year 1936.

		Nuisances the Sum- being ap-		Summonses issued.
NATURE OF COMPLAINT.	No. of Cases.	Work done and Ne abated without the monses ordered be plied for.	Work done and Summonses withdrawn.	Other Results.
Public Health Acts :-		la di santa		
Roofs and/or spouting defec-	00	10	0	
Dampness in rooms, etc	20	18	2	
Rooms inadequately venti-				
lated (broken sash-cords,etc.) Fireplaces, ovens, etc., defec- tive (causing emission of	4	4		
smoke into rooms)	5	5		
rotted, etc	1	1		
untrapped, etc	1	1		
Basement room dilapidated, causing room above to be	1	1	****	
damp, cold, draughty, dirty,	,	,		
Drains defective, imperfectly	1	1		
trapped, etc	5 8	5 8		
Public Health Act, 1875, Sec. 36, and Newcastle upon Tyne Improvement Act, 1892, Sec. 53:—				
Water-closets defective	14	14		
Newcastle upon Tyne Cor- poration Act, 1911, Sec. 55: Want of or defective dustbins		par sea a		
for house refuse	6	6		
Newcastle upon Tyne Cor- poration Act, 1926, Sec. 14: Newcastle upon Tyne Cor-				
poration (General Powers) Act, 1935, Sec. 9:—				
Houses not provided with sufficient and properly ven-			-	
tilated larder or other food storage accommodation	1	1	****	
Total	75	73	2	2008

HOUSING.

Housing Act, 1935.-Overcrowding.

PRELIMINARY SURVEY.

Section 1 of the above Act imposes on Local Authorities the duty of ascertaining the number of dwelling-houses in their area which are overcrowded.

The survey for this purpose, authorised by the City Council on the report of the Health Committee of 4th December, 1935, has now been completed.

The enumeration of the houses in the City, other than those on the Corporation estates, has been carried out by a temporary staff of 20 men and 2 clerks, who were appointed for this purpose.

The houses on the Corporation estates were dealt with by the City Treasurer's staff.

The overcrowding provisions of the Act are set out in two tables. Table I. states the maximum number of persons permitted to occupy a house in accordance with the number of rooms. Table II. states the maximum number in accordance with the floor areas of the rooms. Further, it is not permissible for persons of opposite sexes who are over 10 years of age, and not living together as husband and wife, to sleep in the same bedroom.

A circular (No. 1,507) issued by the Minister of Health on the 19th November, 1935, directs that :—

- (a) The inspection of the houses be completed by the 1st April, 1936.
- (b) The report of the inspections, showing the result of the inspections and the number of new houses required to abate the overcrowding be submitted by 1st June, 1936.
- (c) The proposals for the provision of the new houses required be submitted by 1st August, 1936.

In certain other towns the survey has been confined to houses of the working classes, but in view of the special local conditions prevailing in Newcastle where the development of tenemented dwellings in residential areas has been carried out on a large scale, and also in view of the marked degree of overcrowding which was revealed by the 1931 census, it was thought advisable to make the scope of the survey as wide as possible.

HOUSING ACT, 1935.—OVERCROWDING SURVEY.

TABLE 1.

SEPARATE DWELLING-HOUSES.

Private. Council. 3,650 44 48 2,191 32 1,509 460 111 9,64 6.8 2,463 1,620 76 4,232 1,509 460 111 9,64 6.8 3,580 919 239 2,939 2,430 117 39 17,49 6.8 4,185 919 239 2,5182 907 39 17,49 6.8 8.9 4.6 6.8 8.9 4.6 6.8 8.9 4.6 6.8 8.9 4.6 6.8 8.9 4.6 6.8 8.9 4.6 6.8 8.9 4.6 6.8 8.9 4.6 6.8 8.9 4.6 8.9 4.6 8.9 4.6 8.9 4.6 8.9 4.6 8.9 4.6	MUNICIPAL WARDS.	Tota	Total Nos. Recorded.	Em	Empty.	Not Ove	Not Overcrowded.	Overci	Overcrowded.	Q.	Percentage of Overcrowding	je oje
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		Private.	Council.	Private.	Council.	Private.	Council.	Private.	Council.	Private.	-	Total.
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$, , , , o m	0.000		07		101.0	00	044	01	00 00	0 10	00 00
2,560 99 2,939 2,939 14,66 14,66 14,66 14,66 14,66 14,66 14,66 14,66 14,66 14,66 14,66 14,66 14,66 14,66 14,66 17,49 17,23 16,7 17,23 16,7 17,23 16,7 17,23 16,7 17,23 16,7 17,23 16,7 17,23 16,7 17,23 16,7 17,23 16,7 17,23 16,7 17,23 16,7 17,23 16,7 17,23 16,7 17,23 16,7 17,23 16,7 17,23 17,2 17,2	II Saints	3,053	44	848		2,191	32	814	12	26.66	27.3	26.06
2,463 1,620 76 2,463 4,232 1,509 460 111 9.64 6.8 3,636 919 228 2,972 1,509 460 111 9.64 6.8 4,185 919 228 2,182 907 59 12 1.749 1.3 4,185 168 3,642 907 59 12 1.04 6.8 4,185 168 3,642 907 59 12 1.04 6.8 6,523 2,567 193 6,129 2,450 201 117 8.96 1.3 3,874 73 1948 1,948 1,43 17.28 8.96 4.6 11.3 1.3 1.3 1.3 1.3 1.3 1.3 1.3 1.3 1.3 1.3 1.3 1.3 1.3 1.3 1.3 1.3 1.3 1.3 1.5 1.5 1.1 1.3 1.5 1.5 1.1 1.5 1.5	rmstrong	3,560	****	66		2,939	4	522	****	14.66	****	14.66
4,768 1,620 76 4,232 1,509 460 111 9.64 6.8 3,636 2,8 2,972 1,509 460 111 9.64 6.8 5,480 919 239 2,972 907 59 12 1.07 1.3 4,185 168 3,678 2,450 201 117 8.96 4.6 4,185 193 6,129 2,450 201 117 3.08 4.6 3,874 73 2,67 3,494 7 0.18 3.69 4.6 2,413 2,68 3,494 7 0.18 3.69 4.6 117 3.08 4.6 2,413 4,9 1,948 1,948 1,143 3.69 4.6 11.7 3.69 4.6 11.7 3.69 4.6 11.5 3.69 4.6 11.5 3.69 4.6 11.3 3.69 4.6 11.3 4.6 11.3 4.6 11.5 <td< td=""><td>rthur's Hill</td><td>2,463</td><td>4444</td><td>24</td><td></td><td>2,407</td><td></td><td>32</td><td>*****</td><td>1.29</td><td></td><td>1.29</td></td<>	rthur's Hill	2,463	4444	24		2,407		32	*****	1.29		1.29
3,636 919 28 2,972 907 636 12 17.49 1.3 5,480 919 239 5,182 907 59 12 17.49 1.3 4,185 1,68 3,642 2,450 201 117 8.96 4.6 3,874 7 3,658 2,450 201 117 3.08 4.6 3,707 206 3,494 7 0.18 3.69 1.723 4.6 2,413 49 1,948 416 7 0.18 8.9 2,530 41 1,948 1,948 416 17.23 8.9 2,530 44 1,853 3,011 2,181 651 214 17.58 8.9 3,651 436 18 3,709 92 206 12.33 11.5 4,096 104 181 3,719 2,179 641 242 11.99 9.96 5,345 2,431 105 <td>enwell</td> <td>4.768</td> <td>1,620</td> <td>76</td> <td>::</td> <td>4,232</td> <td>1,509</td> <td>460</td> <td>111</td> <td>9.64</td> <td>6.8</td> <td>8.94</td>	enwell	4.768	1,620	76	::	4,232	1,509	460	111	9.64	6.8	8.94
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	vker	3,636	*****	28		2.972		636	*****	17.49		17.49
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	ene	5.480	616	239		5,182	907	59	12	1.07	1.3	1.1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	lswick	4.185		168		3,642		375	****	8.96		8.96
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	enham	6,523	2,567	193		6,129	2,450	201	117	3.08	4.6	3.5
3,707 206 3,494 7 0.18 2,413 49 1,948 416 17.23 8.9 2,413 49 41,948 1,948 416 17.23 8.9 2,530 44 1,853 3,011 2,181 651 214 17.58 8.9 2,530 44 1,853 363 628 73 17.2 16.7 4,096 104 181 3,709 92 206 12 5.02 11.5 4,327 184 3,310 833 19.25 11.5 5,345 2,431 105 10 4,599 2,179 641 242 11.99 9.96 5,345 2,431 105 1 4,599 2,179 641 242 11.99 9.96 3,484 43 10 61,514 9,713 7,759 793 10.91 7.54 81,614 1,825 10 61,514 9,713 7,759 793 10.91 7.54 81,614 1,825 71,27 8,552 71,27 8,552	eaton	3,874		73		3,658		143	:	3.69	:	3.69
2,413 2,413 49 1,948 1,948 416 17.23 8.9 2,530 4,3651 44 1,853 3,011 2,181 651 214 17.58 8.9 2,530 444 1,853 3,011 2,181 628 73 17.2 16.7 3,651 4,66 104 181 3,005 363 628 73 17.2 16.7 4,096 104 181 3,709 92 206 12 5.02 11.5 4,327 184 3,310 833 19.25 11.5 5,345 2,431 105 10 4,599 2,179 641 242 11.99 9.96 5,345 2,431 1,825 10 61,514 9,713 7,759 793 10.91 7.54 81,614 1,525 71,227 8,552 8,552 10.991 7.54	puomse	3,707		206	-	3,494	4011	7	-	0.18	****	0.18
3,703 2,395 41 3,011 2,181 651 214 17.58 8.9 2,530 44 1,853 3,005 363 628 73 17.2 16.7 3,651 436 18 3,005 363 628 73 17.2 16.7 4,096 104 181 3,709 92 206 12 5.02 11.5 4,327 184 3,310 4,599 2,179 641 242 11.99 9.96 5,345 2,431 105 10 4,599 2,179 641 242 11.99 9.96 3,484 43 2,976 2,976 2,976 13.34 13.34 13.34 71,098 10,516 1,825 10 61,514 9,713 7,759 793 10.91 7.54	t. Andrew's	2,413		49	:	1,948		416		17.23		17.23
2,530 44 1,853 633 73 25.01 16.7 3,651 436 18 3,005 363 628 73 17.2 16.7 4,096 104 181 3,709 92 206 12 5.02 11.5 5,345 2,431 105 10 4,599 2,179 641 242 11.99 9.96 3,484 43 2,976 2,976 465 13.34 13.34 71,098 10,516 1,825 10 61,514 9,713 7,759 793 10.91 7.54 81,614 1,835 71,227 8,552		3,703	2,395	41	:	3,011	2,181	651	214	17.58	8.9	14.18
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		2,530		44	****	1,853	-	633	-	25.01		25.01
300 6 3,709 92 37 12.33 11.5 4,096 104 181 3,709 92 206 12 5.02 11.5 5,345 2,431 105 10 4,599 2,179 641 242 11.99 9.96 3,484 43 2,976 2,976 13.34 13.34 71,098 10,516 1,825 10 61,514 9,713 7,759 793 10.91 7.54 81,614 1,835 71,227 8,552	t. Lawrence	3,651	436	18		3,005	363	628	73	17.2	16.7	17.15
4,096 104 181 3,709 92 206 12 5.02 11.5 4,327 184 3,310 4,599 2,179 641 242 19.25 11.99 9.96 5,345 2,431 105 10 4,599 2,179 641 242 11.99 9.96 3,484 3,484 1,825 10 61,514 9,713 7,759 793 10.91 7.54 81,614 1,835 71,227 8,552		300	****	9	****	257	1	37	4111	12.33		12.33
4,327 184 10 3,310 2,179 833 19.25 19.25 5,345 2,431 105 10 4,599 2,179 641 242 11.99 9.96 3,484 43 2,976 2,976 13.34 13.34 71,098 10,516 1,825 10 61,514 9,713 7,759 793 10.91 7.54 81,614 1,835 71,227 8,552		4,096	104	181	-	3,709	92	206	12	5.02	11.5	5.19
5,345 2,431 105 10 4,599 2,179 641 242 11.99 9.96 3,484 43 2,976 465 13.34 71,098 10,516 1,825 10 61,514 9,713 7,759 793 10.91 7.54 81,614 1,835 71,227 8,552	cephenson	4.327		184		3,310		833	:	19.25		19.25
3,484 43 2,976 465 13.34 71,098 10,516 1,825 10 61,514 9,713 7,759 793 10.91 7.54 81,614 1,835 71,227 8,552	/aiker	5,345	2,431	105	10	4,599	2,179	641	242	11.99	96.6	11.35
71,098 10,516 1,825 10 61,514 9,713 7,759 793 10.91 7.54 81,614 1,835 71,227 8,552	'estgate	3,484	***	43		2,976	1	465	:	13.34		13.34
1,835 71,227	Totals	71,098	10,516	1,825	10	61,514	9,713	7,759	793	10.91	7.54	10.48
1,835 71,227)	1)	1)	1)	1			
		81	614	8,1	35	71,	227	8,5	52			

TABLE 2

7 71 8 81 9 91 10 101 111 111 12 13 141 16 171 19 201 22 24 26 29 301 32 4 1 11 2 21 3 31 (a) (b) 7,593 3538 126 1759 40 __ 218 1 __ 81 1 __ 54 __ _ 17 17 9 1 __ _ _ 2 37 67 15 44 __ 16 (a) 130 (b) 18,918 911 4144 694 5281 1 3972 195 3 4 1975 327 48 (a) 495 (b) 5,820 __ 7 8 1 __ 1 __ ... 86 49 1550 2 1 452 20 26 1940 8 408 ___ 1065 198 (a) 541 (b) 15,549 257 134 59 21 21 2 2 25 1091 161 - 111 4,472 21 515 25 12 814 169 3 738 1 9 407 (a) 1,481 (b) 8,738 230 121 74 26 _ 113 11 1097 243 17 138 1822 3 2454 1054 11 1497 216 1 603 (a) 592 (b) 2,072 13 39 37 528 1 31 17 12 1 1 ... 2,664 65 __ 40 2 447 51 4 524 444 7 251 64 1 86, (a) 832 (b) 4,755 5,587 53 104 758 1 155 93 54 29 28 4 10 1 2 13 1240 818 16 756 191 26 9 13 3 1 ... 271 22 1 301 17 105 3 336 388 8 15 138 54 2,488 __ 1 83 53 38 12 31 5 6 1 2 14 1 254 13 1 388 14 128 49 126 438 47 83 291 96 3 191 1 38 22 73 201 46 3 _ 12 5 7 1 2 2 1 1 ... 896 .. 131 8 12 26 74 32 1 165 45 27 16 7 9 2 4 4 3 1,090 33 15 52 228 14 5 82 48 130 67 1 2 141 5 3 1 1 1 38 35 6 26 1 1 14 10 91 6 22 9 5 6 6 1 1 1 ... 1 453 36 1 5 (a) 102 (b) 48 150 6 29 26 25 144 21 47 1 1 ... - - - 1 - 1 - 1



		۶	

HOUSING ACT, 1935.—OVERCROWDING SURVEY.

TABLE 3.

HOUSES OTHER THAN THOSE ON CORPORATION ESTATES.)

																			-1.	ABLE	3.											(Ho	USES C	THER:	THAN	THOSE	ON C	ORPORATION ES	IMILO
umber Persons							Nun	nber o	f Famil	ies con	taining	the Nu	imber e	of Pers	ons in	the Fire	st Colu	mm occ	upying	Dwelli	ngs wit	h the P	ermitte	d Num	ber sho	wn at t	he He	ad of th	is Colu	20.0.								FAMILIES.	Tot
Family.	1	10	2	21	3	3)	4	41	5	51	6	6}	7	71	8	81	9	91	10	101	11	111	12	13	14]	16	171	19	20}	22	24	26	29	301	32	35	361	(a) Overcrowded (b) Uncrowded.	d Fami
1	13	3458	126	1689	40		991		4		666				218			81	1		54			17	17	9	1											(a) (b) 7,385	7,38
14		2	37	67	15		44				16				1			4			1			1														(a) (b) 188	18
2	93	36	883	4138	291		5281	2	47	1	3972	3	3	4	1975		2	588			327			165	70	21	9	5	2	1	1							(a) 129 (b) 17,791	17,90
21		418	76	49	1445	1	1940	8	25		1065	1	2	1	452			86			26			7	5	1		1										(a) 494 (b) 5,115	5,60
3		371	78	89	1972	6	3997	25	150	1	3646	3	5	5	2010			716	1		482			257	134	59	21	21	2	2	1							(a) 538 (b) 13,516	14,05
31		109	10	814	114	57	940	21	113	3	738	3	1	9	407	1		109			68			25	12	- 6		1										(a) 1,047 (b) 2,514	3,5
4		113	11	1097	190	17	62	138	1405	3	2454	10	7	11	1497	1	1	595	1		363			230	121	74	26	37	8	5	2	2		1				(a) 1,428 (b) 7,054	8,48
4)		40	2	447	32	13	37	37	360	4	524	7		7	251	1	1	85			65		1	31	17	12	1	1										(a) 571 (b) 1,405	1,9
5		20	4	585	40	7	45	104	617	13	1240	20	7	16	756	1		337		1	232		1	155	93	54	29	28	4	10	1	2	1	1	1			(a) 805 (b) 3,620	4,4
51		20		271	9	1	301	17	36	3	336	8	5	15	138			61			45			26	9	13	3			1								(a) 655 (b) 663	1,3
6		14	1	254	8	1	388	14	73	49	126	75	47	83	291	5	3	187	2		116		1	83	53	38	12	31	5	6	1	2			1			(a) 802 (b) 1,168	1,9
61		2		131	4	1	165	1	11	22	66	24	12	26	74	2		45	3		30			12	5	7	1	2	2	1	1							(a) 403 (b) 247	6
7		3		86		1	141	2	15	15	52	44	14	37	130	2	3	76	1	2	60			45	27	16	7	9	2	4	4	3			1		1	(a) 359 (b) 446	8
71				34	1	1	58	3	7	10	21	4	6	20	38	2	6	24	1	1	14			5	3	1					1	1						(a) 145 (b) 117	2
5				26	1		59		8	16	68	8	5	8	7	6	9	33	8		29			22	9	5	6	6	1	1	1		1					(a) 196 (b) 141	3
81				13			20		2	8	29	3	1		2	6	3	8			8			6	1	5												(a) 78 (b) 37	1
				2	3		17			4	21			-1	4	2	2	24	1		8	2		12	5	2	2	1		1						1		(a) 54 (b) 61	1
9)							4			1	8	1			2	2	3	2	1	1	4		1	3	1				1									(a) 21 (b) 14	
10				1			3				6				1		4	2	1		3			2	5	1	2		2	1	3		1					(a) 17 (b) 21	
104							1				2				1			2	-									2										(a) 6 (b) 2	
11											2			-1	2			1	1	-					1			2		1								(a) 7 (b) 4	
115																					1																	(a) 1 (b)	
12											1									1		1		3				1					1					(a) 3 (b) 5	

.

HOUSING ACT, 1935.—OVERCROWDING SURVEY.

TABLE 4.

(Corporation Estates only.)

umber Persons -	Nun	nber of	Familie	es conta	ining t	he Num	ber of	Persons	s in the	First C	olumn	оссиру	ing Dw	ellings v	vith the	Permit	ted Nu	mber sh	own at	the Hea	ad of th	is Colu	nn.	FAMI		Total
Family.	1	11	2	21	3	31/2	4	41	5	5 1	6	61	7	71	8	81	9	91	10	101	11	111	12		rcrowded.	Familie
. 1		80	****	70		****			47	2112		10				1								(a) (b)	208	208
11	****					***	****		16	****		2							2010			1010		(a) (b)	18	18
2		1	28	6	403			****	450	****		192				48	- Anna		1-14					(a) (b) 1	1,127	1,128
21/2			1	2111	105				383			197	****			20								(a) (b)	705	706
3			3		147				941		****	781				161	44.00	3				****		(a) (b) 2	3 2,033	2,036
31	1444		****	****	55				402		2004	404			-	48		2						(a) (b)	55 856	911
4			-	1414	53			***	417	-	2201	1044				215		8						(a) (b) 1	53 1,684	1,737
41				44.00	19	1448	2		166			437	4117			63		1	2100	222				(a) (b)	21 667	688
5					19		8		141			798			200	190	- 100	6		****				(a) (b) 1	27 1,135	1,162
51			1-		13	-			69			390			***	54	41.49	2						(a) (b)	82 446	528
6	-				5				55			363				91	****	4						(a) (b)	60 458	518
61					4			-	27		7	177				30	-1-1-	1				****		(a) (b)	38 208	246
7	2107				1				18	6497		184		11	+++00	65		6			11.11			(a) (b)	203 82	285
71		****							13			87			1117	33		2						(a) (b)	100 35	135
8		2024			2	100			4			77	1014			30	2000	3			4114			(a) (b)	83 33	116
	****			1					1	1010		23				10		1						(a) (b)	24 11	35
8½ 9	-	9100	****	-			-	-	3			16			4010	3	6	1						(a) (b)	22 7	29
	22,00	2000	1111	1000	1117				4000			6		-			6						11/10	(a) (b)	12	12
91	27-15		****						-			3					3						***	(a) (b)	6	6
	4144											44.00					1			4444		****	****	(a) (b)	1	1
101		2010.								2.10				0100			1111	2116		****				(a) (b)		
11	11.04	+114	200	200									1000									-	***	(a) (b)		
111	7710	(0000)	100	100									-				1			ins	200	0.00		(a) (b)	1	1



10,516 houses on the Corporation estates and 71,098 ascertained other dwelling-houses have been recorded, making a grand total of 81,614 separate dwelling-houses.

The following table summarises the conditions found in :-

- (A) Council houses.
- (B) Other dwelling-houses.

A. B.
Overcrowded ... 7.54 per cent. 10.91 per cent.
Not overcrowded... 92.37 ,, 86.52 ,,
Empty 0.09 ,, 2.57 ,,

The overcrowding throughout the City, i.e., the houses on the Corporation estates and other ascertained dwelling-houses is 10.48 per cent.

The actual percentage of overcrowding existing in the Wards of the City of all dwelling-houses is set out in Table 1.

The survey of the 81,614 separate dwelling-houses has necessitated 88,113 visits by the enumerators and the measurement of 32,624 rooms.

The City Treasurer's staff has made 13,058 visits and measured 7,560 rooms of the foregoing totals.

It is of interest to note that in 211 dwelling-houses overcrowding exists solely on account of the necessity for sex separation, and that within the next two years 584 dwelling-houses may possibly become overcrowded due to the growth of families in age and numbers.

The work of the survey has been carried out very efficiently by the temporary staff and, throughout the whole of the survey, it has not been necessary in any instance to have recourse to legal action to obtain information or entry to houses.

Details of the survey are set out in Tables 1, 2, 3, 4 (pages 222-222c).

The Minister of Health has fixed the 1st January, 1937, as the "appointed day" with respect to overcrowding and, in consequence, the 1st July, 1937, is the date on and after which overcrowding offences will commence. Provisions are embodied in the Act safeguarding certain specified cases of overcrowding.

Generally, the offences which may commence on the 1st July, 1937, are in the cases of empty houses being let to families whose numbers are in excess of the permitted number to occupy the house.

It is the duty of the local authority, on request from the owner or occupier, to provide the "permitted number" in relation to a house, and all "permitted numbers" must be inserted in the rent books or similar documents by the owners on or before the 1st July, 1937.

To obtain the "permitted numbers" it is necessary to measure the rooms of the house and, for this purpose, a temporary staff has again been appointed, and commenced their task in January, 1937.

The City Treasurer has already commenced to abate overcrowding in Council houses. During the year, 299 families, affecting 1,581 persons, have been re-housed in suitable dwellings.

The Housing Acts, 1925 to 1935.

The number of inspections under the Housing Acts was 11,368, an increase of 866 over last year's total.

MINISTRY OF HEALTH TABLE.

1.—Inspection of Dwelling Houses during the Year:—	
(1) (a) Total number of dwelling houses inspected for housing defects (under Public Health or Housing Acts)	7,621
(b) Number of inspections made for the purpose	21,300
(2) (a) Number of dwelling houses (included under sub-head (1) above) which were inspected and recorded under the Housing Consolidated Regulations, 1925	
(3) Number of dwelling houses found to be in a state so dangerous or injurious to health as to be unfit for human habitation	*1,202
(4) Number of dwelling houses (exclusive of those referred to under the preceding sub-head) found not to be in all respects reasonably fit for human habitation	4,084
* (Dealt with as Clearance Areas or as Individual Unfit Houses	5).
 Remedy of Defects during the year without Service of formal Notices Number of defective dwelling houses rendered fit in consequence of informal action by the Local Authority or their 	:
officers	1,417
3.—Action under Statutory Powers during the Year:—	
(a).—Proceedings under Sections 17, 18 and 23, of the Housing Act, 1930—	
(1) Number of dwelling houses in respect of which notices were served requiring repairs	524
(a) By owners	503
(b) By Local Authority in default of owners	
(1) Number of dwelling houses in respect of which notices	2,143
(2) Number of dwelling houses in which defects were remedied after service of formal notices:—	
(a) By owners(b) By Local Authority in default of owners	2,128
(c).—Proceedings under Sections 19 and 21 of the Housing Act, 193	30 :
(1) Number of dwelling houses in respect of which Demolition Orders were made	19
(2) Number of dwelling houses demolished in pursuance of Demolition Orders.	† 20
† (16 of these dealt with in previous years.)	
(d).—Proceedings under Section 20 of the Housing Act, 1930 :-	
(1) Number of separate tenements or underground rooms in respect of which Closing Orders were made	31
(2) Number of separate tenements or underground rooms in respect of which Closing Orders were determined, the tenement or room having been rendered fit	

MINISTRY OF HEALTH TABLE-Cont nued.

(a).—(i) Number of dwellings overcrowded at the end of the	
year	8,55
(ii) Number of families dwelling therein	26,9
(b).—Number of new cases of overcrowding reported during the year	***
(c). (i) Number of cases of overcrowding relieved during the year	29
(ii) Number of persons concerned in such cases	1,58
(d). Particulars of any cases in which dwelling-houses have again become overcrowded after the Local Authority have taken steps for the abatement of overcrowding	
(e). Any other particulars with respect to overcrowding conditions upon which the Medical Officer of Health may consider it desirable to report	****
†This number is calculated in accordance with the Ministry of He formula. The number of persons, irrespective of ages, is 46,563	

Housing Act, 1930; Section 17.

The systematic visits under this Section have been increased by 343 in comparison with last year. 1,941 houses (selfcontained and flat type) have been inspected within the 10 sanitary districts. Considerable difficulty has been experienced at times with a certain type of owner regarding the requirements of notices. However, obstacles were overcome without recourse to report for statutory action.

Under this valuable Section, owners are required by notice to carry out works or repairs, making houses reasonably fit for habitation.

The nature and number of the defects so dealt with are shown in the following summary:—

HOUSING ACT, 1930; SECTION 17.

DETAILS OF WORKS CARRIED OUT UNDER NOTICE.

No. of houses involved	
Roofs repaired (including chimney stacks)	1,107
Spouting repaired, renewed, etc.	793
External walls repaired, re-pointed, etc.	1,336
Under-floor ventilation provided.	238
Yard pavements repaired or renewed	822
Dampness remedied (from causes other than those defined above)	870
Drains repaired, reconstructed, etc.	651
Water-closets repaired	1,237
,, ; additional conveniences provided	39
Dustbins provided	349
Accommodation for washing clothes, provided, repaired, etc	423
Coalhouses repaired	541
Ceiling- and wall-plaster repaired or renewed	1,724
Floors repaired or renewed	586
Window sash-cords renewed or repaired	664
Windows repaired or renewed	469
Doors repaired or renewed	947
Cooking accommodation provided	76
Fireplaces repaired or renewed.	815
Ventilated food stores provided	548
Water supply and/or sinks provided, waste-pipes repaired, etc	1,088
Staircases.—Stairs, handrails, etc., repaired or renewed	516
Light (natural and/or artificial) provided	126
Ventilation of rooms, etc., improved	84
Rooms, staircases, etc., cleansed	49
Minor repairs (not included in the above)	
Total	

Housing Act, 1930; Sections 19 and 20. Individual Unfit Houses.

These sections give a local authority power to order the demolition (Sec. 19) or closure as dwellings (Sec. 20) of insanitary houses. The owner has the right to appear before the Health Committee and, if aggrieved by their decision after hearing his case, may appeal to the County Court.

Houses are reported monthly. The numbers dealt with and the results are given in the following table:—

	Nu	mber of	
	Houses.	Separate Holdings (or Families).	Popula- tion.
Demolition Orders made	19	47	147
Closing Orders made	17	31	87
Premises retained for business purposes (owners finding alternative accommo- dation for tenants displaced)	5	10	20
Otherwise dealt with	10	19	51
Totals	51	107	305

Housing Act, 1935; Sections 55 and 56.

Section 55 affords the owner of a working class dwelling the opportunity of submitting to the local authority proposals for the improvement of the house (other than decoration or repair).

When the proposals are agreed and the works completed, a certificate is issued to the owner, covering a period of from five to ten years, exempting the house from any action under "slum clearance," as an unfit house.

Under Section 56, any proposals submitted under Section 55 in regard to a house scheduled in a "slum" area may be transmitted to the Minister of Health for his consideration. No proposals may be made in respect of houses confirmed as "unfit" houses by the Minister.

No application was received under these Sections.

Clearance Areas.

In February and September the Minister of Health held Inquiries into 14 areas. The February Inquiry, in respect of 10 clearance areas and 1 compulsory purchase order, concerned 136 houses, 295 separate dwellings, and a population of 967 persons. The September Inquiry comprised 1 clearance order and 2 compulsory purchase orders, of 415 houses, 857 separate dwellings, and a population of 3,000 persons.

13 areas were fully confirmed by the Minister and 1 with slight modifications.

The first Inquiry only lasted 1 day, the second 6 days. Both were largely attended by the owners and others acting for them, the opposition to the action of the Corporation being particularly strong with respect to the second Inquiry; no objection was offered in regard to 2 areas.

Details of the confirmations are appended in the following tables:—

Ann	(As origi	(a) nally repr	resented).		(b) nfirmed t ter of He	
Area.	Houses.	Dwell- ings.	Popu- lation.	Houses.	Dwell- ings.	Popu- lation
Miller's Hill	12	63	215	12	63	215
Long Row		50	134	31	50	134
Vine Lane		27	78	9	27	78
Fawdon Square (Compulsory						
Purchase Order)	25	25	100	25	25	100
Denton Square		15	57	15	15	57
Mary's Place		18	70	5	18	70
White Street, No. 1		29	78	11	29	78
" ,, No. 2		43	166	13	43	166
awdon Old Pit Cottages	5	6	20	5	6	20
Blythe Nook	2	11	21	2	11	21
Bridge Row		8	28	2 8	8	28
Barker Street	8	24	64	8	24	64
Wesley Street (Compulsory						1
Purchase Order)	266	529	1,799	260	518	1,762
Bentinck (Compulsory Purchase						
Order)	141	304	1,137	141	304	1,137
TOTAL	551	1,152	3,967	545	1,141	3,930

EXCLUSIONS.

dening Structures	Houses.	Dwellings.	Population.
1 Compulsory Purchase Order. Properties excluded from Order and Area, and to be dealt with under a new order	6	11	37

The houses in all the areas were very old, damp, dilapidated, congested, and beyond repair or reconstruction.

Amongst the outstanding defects may be mentioned :-

Structures.—Defective brick- or stone-work, cracked and bulging walls, damp-proof courses either defective or non-existent, chimney stacks burst and in many cases in danger of falling.

Roofs.—Slates and tiles broken, loose and missing; timbers sagging and broken, flashings and spouts defective.

Floors.—Broken, rotted, worn and out of level.

Staircases.—Treads broken, worn, out of level, handrails loose, broken and missing, dark, badly ventilated, and difficult of access.

Windows.—Rotting and perished frames and sashes, broken sash-cords, etc.

Doors.—Dilapidated, badly fitting, warped.

Grates and Stoves.—Badly set, defective, fire-bars missing, ovens out of order.

Sanitary accommodation.—W.C.'s used in common by several tenants in bad structural condition.

Water supplies and Sinks.—Insufficient and not conveniently accessible to all the tenants.

Overcrowding.—Both in the houses and on space, prevalent in every area.

Houses Demolished, etc.—Apart from action by the Health Committee, 14 self-contained houses, 4 flats and 1 tenemented house (of 2 holdings) have been demolished, or have ceased to be used as dwellings, for various reasons (conversion to business premises, estate development, dilapidations, etc.)

Houses built during the year 1936.—The City Engineer reports that there were 1,056 self-contained houses, and 569 flats (tenancies), built privately during the year under report. In addition, 643 dwellings were provided under housing schemes.

Tents, Vans, Sheds and Similar Structures.

There now remain only 2 vans in the City occupied as dwellings. These are on isolated plots of land, are in a clean state, and comply with the bye-laws.

Tenemented Houses.

The demolition of houses in slum areas has again reduced the number of tenemented houses. The conversion of large selfcontained houses into the maximal number of separate dwellings, which was a feature of the preceding few years, has noticeably decreased.

The plans submitted with respect to these houses now show generally a desirable dwelling.

The number of tenemented houses in the City at the end of the year was 2,796, consisting of :—

> 1,708 One-room holdings. 4,720 Two-room holdings. 857 Three-room holdings. 103 Four-room holdings. 2 Five-room holdings.

A total of 7,390 separate holdings. During the year 7,696 inspections have been made of this type of dwelling.

Tenement Bye-laws.

In addition to the Clearance Areas already reported upon, 54 tenemented houses, comprising 247 separate holdings, have been inspected and reported upon in detail during the year, with a view to the bye-laws being put in force. In no case was it found necessary to report for legal proceedings.

New Buildings and Sanitary Alterations.

431 plans were examined by the Medical Officer of Health before their submission to the Town Improvement and Streets Committee and, where necessary, suggestions forwarded to the City Engineer for his consideration, as compared with 397 during the previous year.

Common Lodging Houses.

At the end of the year there were on the Register 17 common lodging houses, as compared with 18 in 1935, one house having been closed and removed from the register, owing to lack of lodgers, etc.

The total number of lodgers for which the houses are registered is 595, showing a decrease of 19 from last year, due to the removal above-mentioned. 2,201 inspections during the day-time and 140 at night-time have been made, and it is satisfactory to note that it was unnecessary to resort to legal proceedings to remedy contraventions of the bye-laws governing the management of the houses.

The average number of lodgers per night was 387, the highest number being 403, and the lowest 353.

The following summary shows in detail the accommodation as at the end of the year :—

Description of		No. of			Accomm	odation.	
Lodgers.	Houses.	Single Beds.	Double Beds.	Married Couples.	Single Women.	Single Men.	Total.
Married couples and single women Women only Men only	1	60 18 497	10	10	60 18	497	80 18 497
TOTAL	17	575	10	10 (20 persons)	78	497	595

Summary of inspections, contraventions found, etc. :-

Number of houses on the register at the end of the year	1
Sec. 63); 18 granted, 1 refused	1
Houses ceased to be occupied as common lodging houses	
Inspections made in the day-time	
Inspections made in the night-time	14
Notices served re washing of bed-clothes, 70, re lime-washing of houses, 35;	10
Defects and Contraventions of Bye-laws, etc. :—	
Structural defects in houses	
Defective water-closets.	
Defective roofs and defective or choked spouting	1
Choked w.c.s and drains	1
Sink waste-pipes obstructed	
Dust-bins defective or insufficient.	
Lack of efficient ventilation (broken sash-cords, etc.) Lavatory wash-basin defective	
Yard pavement defective	
Unclassified minor nuisances (burst water-pipes, etc.) Beds not "aired"	1
Slops not emptied	
Floors not swept	
Lodger occupying an unregistered room	
Deaths reported.	
Cases of infectious disease reported (measles 1, tuberculosis 3)	

Factories and Workshops.

The inspection of these has been maintained during the year, the total number of inspections being 9,655, as against 9,056 for the previous year. These included visits to workshops, domestic workshops, workplaces, laundries and bakehouses, also to factories on receipt of complaint from H.M. Inspector. Generally speaking, their condition as regards sanitary accommodation, ventilation, cleanliness, water supply, and other matters of a hygienic nature, was found satisfactory.

During the year 31 lists of outworkers were received, 8 employers having sent in their lists in February and August, as required by the Factory and Workshop Act, 1901, and 15 employers only once. Included in the lists were the names and addresses of three outworkers residing in other towns, and these, in accordance with the requirements of the Act, were forwarded to the Local Authorities of the districts concerned. No contravention of the Act was found in any of the 74 outworkers' premises inspected.

39 notices as to insanitary conditions in factories and workshops were received from H.M. Inspector of Factories, 30 of which related to factories (which are visited by the Health Department staff only on receipt of a complaint from H.M. Inspector), and 9 to workshops. Some of the latter had, however, been found and dealt with by the District Inspectors prior to receipt of the complaint. The others received due attention and the necessary works were carried out without having to resort to legal proceedings.

Administration of the Factory and Workshop Act, 1901, in connection with Factories, Workshops and Workplaces, during the Year 1936.

Home Office Tables.

INSPECTION OF FACTORIES, WORKSHOPS AND WORKPLACES.
 INCLUDING INSPECTIONS MADE BY SANITARY INSPECTORS.

	Number of			
Premises. (1)	Inspections. (2)	Written Notices. (3)	Occupiers Prosecuted (4)	
Factories (Including Factory Laundries.) Workshops (Including Workshop Laundries.) Workplaces. (Other than Outworkers' premises.)	394 7,618 1,643	214		
TOTAL	9,655	214		

2.—DEFECTS FOUND IN FACTORIES, WORKSHOPS AND WORKPLACES.

	Numb	Number of Offences		
Particulars.	Found.	Re- medied.	Referred to H.M. In- spector.	in respect
*Nuisances under the Public Health Acts:— Want of cleanliness Want of ventilation Overcrowding Want of drainage of floors Other nuisances Sanitary accommodation insufficient unsuitable or defective not separate for sexes	194 8 1 76 47 116 12	194 8 1 76 44 115 12		
Offences under the Factory and Workshop Acts:— Illegal occupation of underground bakehouse (s. 101) Other offences (Excluding offences relating to out-work and offences under the Sections mentioned in the Schedule to the Ministry of Health (Factories and Work- shops Transfer of Powers Order, 1921.)	1	1	ïï	
TOTAL	455	451	11	

^{*} Including those specified in sections 2, 3, 7 and 8 of the Factory and Workshop Act, 1901, as remediable under the Public Health Acts.

OUTWORK IN UNWHOLESOME PREMISES, Section 108.

Nature of Work. (1)	Instances.	Notices served. (3)	Prosecutions.
As per Home Office List	None.	None.	None.

TRADES.

Particulars as to the number and nature of the various trades carried on in the workshops of the City :—

Trades.	Work- shops.	Domestic Work- shops.	Work- places.
Athletic Outfitters, etc.	12	100	****
Bacon Curing, Pickles, etc.	50	L	200
Bags, Waterproofs, etc. (making and repairing)	19	2	3
*Bakehouses	268		
Blacksmiths, Plumbers, etc.	123		4
Bouquets and Wreaths (making, etc.)	14		0.00
Boots, etc. (making and repairing)	137	20	****
Dressmaking, Underclothing, etc.	253	15	
Drysalters, Cleaning & Packing Fruit, Tea, etc.	29	1	94
Eurniture Making Leiners etc.	226	11	0.1
Furniture Making, Joiners, etc.		11	****
Harness, etc. (making and repairing)	20		4554
Jewellery, Watches, etc. (making & repairing)	79	2	
Laundries	10	****	****
Machines and Tools (making and repairing)	149	2000	5
Painters, Engravers, Photographers, etc.	82	6	15
Restaurant Kitchens, etc.			111
Tailoring, Shirts, etc.	243	13	
Miscellaneous	112		117
Totals	1,826	71	349

^{*} Includes 45 "Factory" and 66 "Domestic" Bakehouses.

Inspection of Council and other Schools.

During the year 181 inspections were made. In three cases defects were found. These were brought to the attention of the Education Authorities and subsequently remedied.

At one school (acquired under the Northumberland Review Order, 1935), privies and ashpits have been replaced by "chemical" closets. Water-closets could not be provided owing to the absence of sewers.

Rag Flock Acts, 1911 and 1928.

There are no manufacturers of rag flock in the City, the principal users being upholsterers and bedding makers. The number of these who use (or are likely to use) rag flock is 26, in 15 workshops and 11 factories. To the former (which are also inspected under the Factory and Workshop Acts) 74 visits were made, factory premises being only visited on receipt of complaint from H.M. Inspector.

Eight samples of rag flock were purchased and submitted for analysis. All were found to conform to the standard of cleanliness prescribed by the Regulations. Eight samples of feathers (used in the manufacture of bedding and cushions) were purchased and submitted for analysis under the "rag flock" test. Of these, only one conformed to the prescribed standard of cleanliness. No legal action could be taken, as the law relates only to rag flock. There is definitely a need for the amendment of the law in this respect, fixing a standard of cleanliness or purity and the materials to be used in this class of work.

Exhumations.

Two exhumations were carried out under the supervision of the Department during the year, both being authorised by Home Office Licence. The operations were carried out in the early morning in a sanitary and reverent manner and with due regard to the conditions set out in the Licence.

Fertilisers and Feeding Stuffs Act, 1926.

In pursuance of this Act, 24 visits were made to factories, warehouses, and retail shops where fertilisers or feeding stuffs were prepared or stored for sale, for the purpose of seeing that the requirements were carried out as to the marking of packages, inspection of registers, etc.

Six samples of fertilisers and 1 of feeding stuff were obtained (mostly informally) and submitted for analysis to the Agricultural Analyst. In the case of one sample of fertiliser the amount of nitrogen was found to be below that stated on the sample. This was the subject of correspondence with the Ministry of Agriculture and Fisheries, who advised that the article did not come within the scope of the Act, and no further action was taken. The remaining samples all complied with the requirements.

Merchandise Marks Act, 1926.

In the administration of this Act, 335 inspections and personal visits were made to shopkeepers, stall-holders, hawkers, etc., in order to ascertain whether imported goods were properly marked with the "indication of origin" required by the Act and the Orders made thereunder. Attention was drawn to the requirements where necessary, in 93 instances there was left a copy of a printed notice to traders (setting out the principal provisions of the Act), and in 65 cases cautions were administered (55 verbally and 10 by special letter).

Agricultural Produce (Grading and Marking) Act, 1928.

143 inspections of markets, shops and stores, were made as to the grading and marking of eggs. No contravention of the Regulations was found.

Pharmacy and Poisons Act, 1933.

LISTED SELLERS OF PART II. POISONS.

102 applications were received for registration as listed sellers. The applicants consisted of :—

Grocery, Provisions and General Stores						50
Hairdressers			****	****		16
Druggists					****	14
Hardwaremen,	etc.		5.0			9
Seed and Agric	cultural	Mer	chants	****		8
Chemical Disir	fectant	Man	ufactur	ers		3
Electrical Sup	plies		21.01	****		1
Manufacturing	Chemi	sts				1

Of these, 101 applications were granted by the Health Commottee, and 1 refused, the premises being considered unsuitable.

One dealer, after several applications being made to him, defaulted regarding payment of the statutory fee, and his registration was accordingly cancelled.

Shops Act, 1934; Section 10.

The duties which fall upon the Health Department under this Section are in respect to ventilation, temperature, sanitary accommodation, lighting, washing facilities, and accommodation for the taking of meals, for persons employed in or about the business of the shop.

Routine inspection is now in progress and, during the year, 452 visits have been made. 84 contraventions of the Act and Section were dealt with.

In 3 cases, where premises were not provided with suitable and sufficient sanitary conveniences and such accommodation was otherwise conveniently available, certificates of exemption were issued under the provisions of the particular sub-section.

Staff Changes.

I have again to record a number of Staff changes. Mr. Isaac Clark, Assistant Inspector of Factories and Workshops, retired on superannuation in June, after 34 years' service with the Corporation, whilst District Inspectors Ralph W. Suddick, Herbert W. Grieves and Maurice W. Swales resigned on taking up service as Sanitary Inspectors to the Urban District Councils of Longbenton, Edmonton and Spennymoor, respectively.

Conclusion.

In conclusion, Sir, I would express to you my sincere appreciation of your unfailing support and guidance afforded to me at all times, and to the whole of the Staff I tender thanks for assiduous attention to their various duties.

I am, Sir,

Your obedient servant,

W. GRAY,

Chief Sanitary Inspector, Inspector of Common Lodging Houses, etc.

Health Department,

Town Hall,

15th May, 1937.

CITY AND COUNTY OF NEWCASTLE UPON TYNE.

A STUDY OF THE DIETS OF SIXTY-NINE WORKING CLASS FAMILIES IN NEWCASTLE UPON TYNE.

1.—INTRODUCTORY.

The dietary survey, of which this report is the record, was not undertaken as an academic exercise, but as a practical enquiry into the food consumption and expenditure of certain working class families in Newcastle during the autumn of 1934. It may be helpful, therefore, to describe briefly the historical and social setting of the investigation.

During the early years of the present decade, Newcastle in common with the other Tyneside towns, lay under a heavy burden of unemployment. The measure of this can be gauged from the facts that between 1931 and 1934, out of a population of approximately 285,000 the average number of persons in receipt of Public Assistance was 19,222. With such a large proportion of the population in circumstances which could only be described as impoverished, special watchfulness was necessary to detect or, if possible, to anticipate the presence of untoward developments.

The first indication of alarm was given towards the end of 1932 in the Annual Report of the Newcastle Dispensary for the previous year. It was stated that "the Committee are gravely concerned about the great increase in poverty, sickness and malnutrition amongst the poorest classes in the City."

Largely as a result of this pronouncement, the City Health Committee requested Dr. J. C. Spence, F.R.C.P., to carry out an investigation into the health and nutrition of certain of the city children between the ages of one and five years. The record of this investigation was published in March, 1934. Dr. Spence's primary conclusion which was to the effect that at least 36% of the children from the poor districts of the city which he had examined, were unhealthy or physically unfit and appeared malnourished, concentrated attention on the urgency of the situation.

A dietary survey carried out by Miss Doris G. Rogers, of the Newcastle Dispensary, during the later months of 1933, emphasised the disparity between the circumstances of employed and unemployed families and the inadequacy of the food supply of the latter when compared with the recently issued recommendations of the Nutrition Committee of the British Medical Association. A similar enquiry into the diets of 15 unemployed and 9 employed

families conducted by the staff of the Maternity and Child Welfare Section of the City Health Department, only served to confirm these observations.

Both these surveys were of a relatively simple character, being based in the main on personal enquiries as to the purchases made by housewives over definite periods, a strictly quantitive study being out of the question in the circumstances. Nevertheless, the value of these enquiries as indicators of the trend of events should not be under-estimated.

There had thus been accumulated a body of evidence, some of real significance, as in the case of Dr. Spence's findings, the remainder largely suggestive, which, re-inforced as it was by a number of unconfirmed impressions and opinions, justified the City Health Committee in setting on foot an enquiry of a more ambitious character. It was accordingly decided to undertake a comprehensive dietary survey of a number of working class families in different parts of the city.

The study was carried out in September, 1934, and the families were selected so as to be representative as far as possible of the working class population. Some measure of special selection was inevitable, as families who were likely to be non-co-operative were obviously avoided. The main objective of the investigation was to compare the diets of unemployed and employed families whether domiciled in "old houses" or living upon "new housing estates." For this purpose, a number in each of the employed and unemployed groups was taken from the old dwellings and the rest from houses on new estates. The number of families studied and their distribution amongst the various sub-groups are tabulated below. Attention is directed to the identifying initials by which these various categories will be referred to subsequently.

	Total No. of families studied.	No. of families in old houses.	No. of families on New Housing Estates.	Identifying Initials.
Unemployed	38	24	14	U.O.H. Unemployed Old Houses U.N.E. , New Estates.
Employed	28	18	10	E.O.H. Employed Old Houses. E.N.E. New Estates.
Widows	3	2	1	W.O.H. Widows Old Houses. W.N.E. ,, New Estates.
	69	44	25	

2.—OCCUPATION OF WAGE-EARNERS.

The occupations of the employed wage-earners are classified as follows:—

labourers, including building trade labourers	
and one chimney sweep	9
metal workers, including plumbers and	
fitters	4
indoor workers, including shop attendants	
and a caretaker	4
transport workers, including 'bus drivers and	
carters	4
outdoor agents, including one advertising	
agent	3
railway workers 2; joiners 2.	

Some children of the employed families were also wage-earners, namely:—

boys, 3 (pit boy, butcher's boy and office boy); girls, 4 (3 maids and 1 dentist's assistant).

Amongst the unemployed families 3 juveniles were employed as follows:—

boys, 2 (labourer and mine firer); girl, 1 (packer).

The heads of all the unemployed families except one had been out of work for periods ranging from six months up to nine years preceding the enquiry.

3.—PERSONNEL EMPLOYED IN THE STUDY.

The practical part of the investigation was conducted by three specially selected Health Visitors working under the personal direction and instruction of Miss M. L. Clark, who has had long experience in the conduct of dietary investigations. The Health Visitors selected had previously assisted in the dietary investigation under the auspices of the Newcastle Health Department to which reference has already been made. On that occasion they gave convincing evidence of their ability to conduct the present more exacting enquiry.

4.—METHOD OF INVESTIGATION.

The method employed was essentially the same as that of Paton, Cathcart and their colleagues. It consists in weighing, measuring and recording all foodstuffs in the house on the first and on the last day of study and also food purchased during the

investigation. The daily out-goings of waste are also recorded. The period of study for each family lasted a week, and the investigator spent some time daily in each house carrying out the weighings and measurings and obtaining the information required. Care was taken throughout to check the amounts purchased by weight, including canned and packet foods. The housewife had been previously instructed to collect all the inedible and edible wastage, such as peelings, meat and fish bones, bread and potato waste, etc., into separate parcels, paper being provided for the purpose. Food supplied to dogs and cats was also ascertained as far as possible and reckoned as waste. From the data thus obtained the net amounts of different foods eaten by the family over the period were determined. Purchases of sweets, ice cream, etc., by children, or from the "fish and chip" shops, were ascertained by enquiry and checked so far as possible. Account was taken of all meals taken in the house by each member of the family and of meals which did not come from the family larder. The latter were approximately assessed in terms of dietman-values and deducted from the total diet-man-value of the family. The commonest evaluations used were breakfast 1rd. dinner $\frac{1}{3}$ rd, tea and supper combined $\frac{1}{3}$ rd, i.e., of the appropriate man-value, but this proportion was occasionally altered according to variations in the relative sizes of meals in different families. Meals eaten in the houses by visitors were similarly assessed and added to the total diet-man-value of the family.

Most of the members of each family were weighed during the survey. Blood hæmoglobin was also determined in the majority soon after the research was finished.

5.—ANALYSIS OF FOODS.

Special chemical analysis had to be made of some foods bearing local names. Analyses for protein, fat and carbohydrate were carried out by the City Public Analyst or in the Ministry of Health Foods Laboratory; mineral analyses were made by Dr. R. A. McCance and Miss Widdowson, of King's College Hospital, London, who also supplied figures for some other foods not listed in analytical tables.

6.—CALCULATION OF RESULTS.

The family co-efficients suggested by Cathcart and Murray and the tables of food values of Plimmer, Sherman, McCance and Lawrence, McCance and Shipp, and Hutchison and Mottram were employed. Account was taken of the loss of nutrients from some foods through cooking, a precaution made possible by the use of unpublished data kindly supplied by Dr. McCance and Miss Widdowson. Losses from cooking occur when food-stuffs are boiled and the water thrown away. "Cabbage water" for example contains on the average 61.6 per cent. of the total calcium and 49.3 per cent. of the total protein present in the raw vegetable.

7.—STANDARDS OF COMPARISON.

The standards we have adopted for comparison are: Calories 3,000, total protein (animal and vegetable) 80 gm., animal protein 37 gm., calcium 0.68 gm., phosphorus 1.32 gm. and iron 15 mg. per man daily. These standards although not all based on indisputable evidence have been assented to by most workers on nutrition as adequate for the average man of a community weighing 70 kg. The community would of course include men engaged in sedentary occupations, light work and hard work. The standards are averages obtained from numerous laboratory and dietary studies and are therefore only comparable with average and not with individual diets. Table I. sets out the standards adopted, together with the daily intake of nutrients per man value for the various groups and sub-groups investigated. Certain other relevant information as to rent, food expenditure and income is also included.

The outstanding feature of the present enquiry was the occurrence of wide variations from the mean values. It therefore seemed more appropriate to compare the diets and their constituents with ranges or zones of values which were regarded as standard, rather than with the usual recognised standards themselves. Accordingly, ranges of the more important nutrients have been chosen so that diets providing amounts inside these limits may be regarded as sufficient.

The ranges adopted which are also indicated by horizontal lines in Table II. are as follows:—

 Calories
 ...
 2,500
 — 3,499.

 Protein
 ...
 70
 — 109 gms.

 Animal Protein
 ...
 30
 — 59 gms.

 Calcium
 ...
 0.5
 — 0.999 gms.

The selection of these ranges is based on the recommendations of experienced investigators in nutrition, but it does not necessarily follow that individual diets containing lesser or greater amounts of nutrients than those inside the ranges are to be judged insufficient or excessive. To pronounce with certainty on the precise requirements of every individual in a community would be to presume a more exact knowledge of human nutrition than we possess. It is a fundamental biological fact that variations exist among all living organisms. This applies to food requirements as well as to body functions, but, in the present state of knowledge, it is impossible to state the extent of variations in food requirements for different individuals. For this reason and to avoid misunderstanding the following terms are used in appraising individual diets in relation to these ranges: "within the range," "above the range "and" below the range." In the case of gross deviations from the range the terms "deficient" or "excessive" are employed.

RESULTS.

8.—AVERAGE COMPOSITION OF DIETS AND FREQUENCY DISTRIBUTION.

The composition of the family diets in terms of pure nutrients is summarised in Table I. The average for the whole 69 families taken at its face value suggests that the diets as a whole were fairly adequate when judged by the quoted standards. Calories, total protein and animal protein may be considered as practically sufficient. The quantity of iron is exactly standard but calcium and phosphorus are deficient to the extent of about 8 per cent.

The general import of the averages in Table I. should not be unduly stressed. That their presentation of the facts requires further amplification is evident from Table II., which shows the frequency distribution of representative constituents of the diet, Calories, total protein, animal protein and calcium. The values extend over a very wide range, the scatter being more marked in employed than in unemployed families. On account of the wide variation in consumption of nutrients, which is the most conspicuous feature of the results, the means lose a good deal of their apparent significance. The range of intake per man value as regards Calories extends from 1,846 to 5,261 and of calcium from 0.24 gms. to 1.41 gms. In these circumstances and in view of the small number of data the means of the two major groups (i.e., unemployed and employed) and four sub-groups (Table I.) give

an imperfect picture of the relative nutritional value of the diets in these groupings. It will be found more informative to study the *range of variation* in the consumption of nutrients as experienced in the various groups and sub-groups.

All families. In the 69 families, the intake of Calories was below the selected range in 25 per cent., calcium in 35 per cent., total protein in 32 per cent. and of animal protein in 42 per cent. The intakes were above the range for Calories in 20 per cent., calcium in 10 per cent., total protein in 12 per cent. and animal protein in 12 per cent. The Calorie, calcium, total protein and animal protein contents in the remaining diets were within the range.

Unemployed and Employed. Considering the unemployed and employed groups separately the Calorie content was below the range in 31 per cent. of the former and in 18 per cent. of the employed, and above the range in 16 per cent. of the unemployed and in 25 per cent. of the employed.

In 2 widows' families the Calorie content was within the range and in one above the range. Four employed families consumed 4,500—5,499 Calories, which was almost certainly an excessive amount, whereas the three highest intakes amongst the unemployed were between 4,000 and 4,249 Calories.

The distribution of the calcium intake shows no appreciable difference between unemployed and employed except that the maximum intake and percentage above the range was higher in the latter (i.e., 1.4—1.49 gm., 2 families) than in the unemployed (i.e., 1.1—1.19 gm., 2 families). Calcium intake below the range was as common amongst the employed as the unemployed, being found in about 35% of the families in each group. It was less than 0.4 gm. and therefore deficient in 6 unemployed, 3 employed families and one widow's. The intake of calcium in two widows' families was within the range and in one below the range.

The intake of total protein was below the range in 29 per cent. of the unemployed and in 36 per cent. of the employed families and of animal protein in 45 per cent. and 32 per cent. respectively. The difference regarding total protein is too small to be important but that between the figures for animal protein is too large to be ignored and suggests, when taken together with the fact that 21 per cent. of the employed but only 5 per cent.

of the unemployed consumed animal protein above the range, that the unemployed as a whole consumed less animal protein than the employed families. Three unemployed and 4 employed families consumed less than 60 gm. of protein which was definitely below the standard range. One unemployed and 3 employed families had more than 130 gm. of protein daily which appears to be an excessive amount. It would be illogical to speak of a deficiency of animal protein because it is not strictly essential for life. It is, however, extremely probable that amounts above 70 gm. are excessive. One unemployed and 2 employed families were in this category. Two of the widows' families consumed amounts of total protein within the range and one below the range. The intake of animal protein was below the range in all three.

The general feature of the frequency distribution is a larger proportion of unemployed than of employed at the lower level of consumption of Calories, calcium and animal protein, and correspondingly smaller proportions at the higher levels. Regarding total protein, the difference of 16 per cent. in favour of the employed above the range is offset to some extent by one of 7 per cent. in favour of the unemployed below the range.

If attention is confined to the mean values in the unemployed, employed, O.H. and N.E. groupings (Table I.) it would seem that on the average the diet of the unemployed was poorer than that of the employed and the diet of the N.E. families poorer than that of the O.H. families. These differences would seem to be attributable in the main to the U.N.E. families since the average diet of this sub-group is lower than that of any of the others and so must have affected the means in both the unemployed and N.E. groups. The wide variation round the means throws some doubt on the validity of this inference and statistical analysis reveals only the following significant differences between them, the term significant being interpreted in its formal statistical sense, i.e., not likely to be due to chance:—

- All unemployed families consumed significantly less fat and animal protein than employed.
- 2. Unemployed families on new housing estates consumed significantly less Calories, fat and animal protein than employed families resident on new housing estates.

These differences are confirmed for Calories and animal protein by the frequency distributions in Table II. Old Houses and New Houses. Comparisons between O.H. and N.E., between U.O.H. and U.N.E. and between E.O.H. and E.N.E. did not reveal any significant differences. On the other hand, the means are uniformly lower in the unemployed than in the employed whether the comparisons are made with or without reference to housing. This consistency suggests that all the differences are real. It would, therefore, seem that the average diet of the unemployed families was on the whole poorer than that of the employed, but as Table II. shows there was very considerable overlap in the diets of individual families.

9.—DIETS BELOW THE RANGE IN CALORIES, TOTAL PROTEIN, ANIMAL PROTEIN OR CALCIUM.

Inspection of those diets falling below the range in either Calories, total protein, animal protein or calcium (Table II.) disclosed that most of them were also below the range in other constituents. To illustrate this point the diets were grouped according to the intake of Calories and averaged in groups (Table IIIa). Total protein, fat, carbohydrate, phosphorus and iron are seen to increase with Calories, whereas animal protein rises and then falls and calcium rises somewhat irregularly. The changes in man-value, rent and food expenditure are not consistent. In two only of the 41 diets listed in Table IIIa, did the calcium intake reach the standard level, 0.68 gm. One of these diets contained 0.75 gm. calcium and 4,098 Calories and the other 0.78 gm. calcium and 2,693 Calories. Some other diets in this grouping are noteworthy on account of extreme variations in their constituents. One had 3,509 Calories but only 17.9 gm. animal protein, another 3,190 Calories but only 11.3 gm. animal protein and 0.24 gm. calcium, and a third 2,646 Calories, 40.6 gm. of animal protein and 0.65 of calcium. The lowest intake of iron was 7.2 mg, in a diet of 2,099 Calories. This was an E.O.H. diet; and its content of other nutrients was, total protein 52 gm., animal protein 17.5 gm., calcium 0.26 gm., and phosphorus 0.61 gm. Another very low diet, also in an E.O.H. family contained 1,846 Calories, 53 gm. total protein, 20 of animal protein, 0.41 of calcium, 0.83 of phosphorus and 10.9 mg. of iron. These 2 diets were doubtless deficient in every respect. The highest iron intake within this grouping of diets was 20.9 mg. in a diet of 2,413 Calories.

The same group of diets was classified according to size of family and the constituents averaged in groups (Table IIIb).

All nutrients except animal protein, which changed but slightly, fell in general with increase in man-value up to the 7.0 man-value level. Food expenditure also fell, but rent showed no constant change. The mean intake of nutrients of families in the 2–2.9 man-value group exceeded that of families in the 5–5.9 group by the following percentages: Calories 11, protein 21, animal protein 13, fat 9, carbohydrate 11, calcium 51, phosphorus 27 and iron 12.

Summarising this analysis it would seem-

- (a) that the Calorie content of the diets is a measure of the total protein, fat, carbohydrate, phosphorus and iron, but not of the calcium or animal protein contents, and
- (b) that the larger the family the lower are the food expenditure and intake of all nutrients per man except animal protein.

10.—DIETS WITHIN THE RANGE IN CALORIES, TOTAL PROTEIN AND ANIMAL PROTEIN.

The diets in this category have been grouped in the same way as the others. The averages are shown in Table IV. (a and b). Seven diets fell into the 2,500—2,999 Calories group and seven into the 3,000—3,499 group. All other nutrients and food expenditure rise with increase in Calories while rent falls slightly and man-value remains almost unchanged. The decrease in rent and increase in food expenditure are consistent with one another and with the increase in consumption of nutrients, for it might ordinarily be expected that the lower the rent the greater would be the food expenditure and, consequently, the consumption of food.

The man-value classification shows that all nutrients and food expenditure fell while rent rose with increase in man-value.

11.—DIETS ABOVE THE RANGE IN CALORIES, CALCIUM, TOTAL PROTEIN OR ANIMAL PROTEIN.

The diets above the range were grouped according to Calorie content and man-value and averaged with the results shown in Table V. (a and b). Total protein, carbohydrate, calcium and iron run parallel with Calories. The general trend of fat and phosphorus is in the same direction but animal protein shows little correspondence with Calories, and man-value, rent and food expenditure none.

In the man-value classification total protein, animal protein, fat and iron fall as man-value rises. The changes in other constituents are irregular, but the larger families, with man-values above 3, are below the smaller, where the man-values are less than 3, in the consumption of every food constituent. They are lower also in food expenditure but higher in rent.

In general, it would appear from this analysis that-

- (a) the Calorie contents of these diets may be regarded as an approximate indication of the amounts of the other nutrients considered except animal protein, and
- (b) the larger families had on the whole poorer diets and spent less money per man-value weekly on food than the smaller families.

It will be observed that the total of families included in Tables III., IV. and V. is 71, *i.e.*, two more than the number studied, 69. The reason for the increase is that two diets which were below the range in animal protein and were therefore put in Table III. were above the range in Calories and so were included in Table V. The details of these diets are noteworthy. They are:—

Total Animal Calories. Protein. Protein. Calcium. Phosphorus. Iron.
Diet A ... 3,509 79.5 gm. 17.9 gm. 0.58 gm. 1,24 gm. 16.3 mg.
,, B ... 4,098 106.6 gm. 29.2 gm. 0.75 gm. 1,96 gm. 20.5 mg.

These examples serve to show to what extent diets may be poor in some respects although they appear adequate or even excessive in others. Thus, Diet A is deficient in animal protein and low also, but to a less extent, in calcium and phosphorus whereas the other nutrients are present in adequate amounts; Diet B is probably too high in Calories while animal protein is on the low side, the other nutrients being quite adequate.

12.—COMPARISON OF TABLES III., IV. AND V.

To illustrate the general characteristics of these diets in the several Tables and to compare the differences, the figures were averaged and the differences calculated on a percentage basis. The results are shown in Table VI.

The dietary constituents increase uniformly from Table III. through IV. to V. The increase is least marked for carbohydrate, the cheapest food-stuff, and most marked for calcium and animal protein which are relatively expensive nutrients. Food expenditure per man-value per week also increases from Table III., through

IV. to V. but much more markedly than any of the food constituents. This indicates that the costs of the diets within the range and above the range were higher on the average than those below the range. The man-value per family moves in the opposite direction to the food constituents and food expenditure. The larger families would appear, therefore, to have had poorer diets than the smaller and to have spent less on food. Neither the dietary constituents nor food expenditure would appear to have been substantially influenced by the rent in the families studied, for this was lowest for the families in Table III. and highest for those in IV. Had diet been appreciably affected by rent, this would have been highest for the families in Table III.

13.—COST OF NUTRIENTS.

The average number of Calories and grammes of protein and calcium obtained per penny of total money spent on food are shown in Table VII. The values were obtained by dividing the total amounts of these nutrients in each diet, exclusive of all foods supplied free of cost, by the amount of money spent on food. Garden produce was reckoned at wholesale prices. The figures do not indicate any appreciable differences between the O.H. and N.E. families whether unemployed or employed or between all unemployed and all employed in regard to calcium. unemployed, however, obtained per penny expended, 22 per cent. more protein and 23 per cent, more Calories on the average than the employed; these differences were found to be statistically significant. The most probable reason for the higher cost of Calories and protein to the employed than to the unemployed is the fact that the former consumed more animal protein and fat which are expensive sources of protein and Calories respectively.

14.—RENT, MAN-VALUES AND FOOD EXPENDITURE PER MAN-VALUE.

The average rents, man-values and expenditure on food per man weekly are stated in Table I. and the family distribution of these is given in Table II. under the headings of the several sub-groups.

The average rent was lowest in the U.O.H. group, 7.86s. per week. The differences between the other groups are slight, but the relatively high rent in the U.N.E. group, 10.27s. per week, is striking. On the average, rent was 21 per cent. higher for

the N.E. than for the O.H. families. The rent did not appear to be related to intake of food or expenditure on food per head. For example, unemployed living in "old houses" with an average rental of 7.86s. spent 53.5d. per man-value per week, while similar families on "new estates" paying on an average 10.26s. weekly for rent, spent 54.3d. per head on food.

The differences between the average man-values are so small and the distribution so wide especially in the two employed groups that no importance can be attached to them.

The distribution of food expenditure per man-value in the different groups is very wide particularly in the employed subgroups (O.H. and N.E.) The average expenditure under this heading was practically identical in the two unemployed, 53.5d. and 54.3d., and the two employed sub-groups, 72.5d. and 75.6d., respectively. The averages of all the unemployed, 53.94d., and all the employed, 73.62d., may therefore be compared. These show that the expenditure of the employed exceeded that of the unemployed by 36.5 per cent. This difference is in all probability due, as pointed out below, to the purchase of (a) more expensive foods and (b) greater quantities of fat and animal protein and probably of other foods, by the employed than by the unemployed.

15.—QUALITATIVE ANALYSIS OF CONSUMPTION OF FOOD STUFFS.

The various kinds of foods and beverages, including different cuts of meat, brands of cereals and vegetables amounted to 151. The list of these foods has been abbreviated so as to illustrate dietary and economic habits, and is set out in the Appendix to the report. The Appendix shows the number and percentage of families who consumed each article of food and also the average intake per man of those families who ate the food. Statistical tests were applied to the most outstanding differences between the percentages of unemployed and employed families consuming important foodstuffs, disregarding housing conditions. Statistical analysis of the average quantities of foods consumed was not justifiable because the number of families involved was too small. It will be convenient to discuss in the first place the differences found to be statistically significant.

These are dealt with in sub-paragraph (a). Those differences which were not found to be capable of statistical analysis, or were not statistically significant, are discussed in the subsequent subparagraphs.

Statistically Significant Differences.

(a) Cakes, fresh tomatoes, butter, fresh milk, bacon and boiled ham.

More employed families consumed cakes, fresh tomatoes, butter, fresh milk, bacon and boiled ham than unemployed families, the differences between the numbers of families in the two groups being greater than is likely to have arisen by chance.

For instance, 76 per cent. of the unemployed families consumed butter and 66 per cent. fresh milk compared with 96 per cent. and 100 per cent. respectively of the employed. The average consumption of fresh milk was low throughout, according to the commonly accepted standards. For the four main sub-groups it was respectively:—

Unemployed—Old Houses	 0.13 p	ints	Per man-
Unemployed—New Estates	 0.26	,,	and the second second
Employed—Old Houses	 0.19	,,	value per
Employed—New Estates	 0.26	,,	diem.

The relative costliness of cakes, tomatoes, fresh milk, butter, bacon and ham is probably the main explanation for these differences in consumption.

Fifty-three per cent. of unemployed and only 7 per cent. of employed took a proprietary brand of dried milk which many of the unemployed received free from the City Health Department. It has been observed in Newcastle that many of the families who receive dried milk free of cost buy only enough liquid milk for tea; some indeed consume no fresh milk at all, but use instead condensed skimmed milk all of which in the present study came from countries outside Great Britain. Skimmed milk, as is well-known, is little inferior in nutritive value to whole milk, the only difference between them being that the bulk of the fat and A and D vitamins are absent from skimmed milk, but these nutrients can easily be supplied by other suitable fats. Skimmed milk, like whole milk, is an excellent source of protein of high biological value, of calcium and of other inorganic elements. In the present study only three families, two unemployed and one employed, consumed liquid skimmed milk. There is no doubt, as was verified by enquiry, that more would have bought skimmed milk in liquid form if it had been easily obtainable; but little or none is placed on the market in this country. It is known that much of the skimmed milk produced is used for purposes other than human food, mainly the feeding of pigs and poultry, and some may be run to waste.

It is difficult to see how the lower consumption of butter by the unemployed than by the employed can be attributed entirely to economic causes, because butter at that time could be obtained almost as cheaply as margarine. Possibly a margarine-eating habit established when butter was more expensive was also a factor.

Differences which were not found to be capable of Statistical Analysis or which were not Statistically Significant.

(b) Cereals.

Bread and flour have been combined because of the very common practice of home baking in the North of England. The great bulk of the bread weighed in the houses at the beginning and end of the study was home-made. The consumption of white flour and bread was almost identical in the 4 sub-groups, about 10 ozs. or 1,000 Calories per man-value daily. The great bulk of the bread consumed was white; only about 45 per cent. of all the families consumed brown bread at all. One U.O.H. family consumed only brown bread.

Biscuits, tapioca and sago and rice show no clear differences between the sub-groups in numbers of families or quantities consumed. About 30 per cent. of all families used oatmeal.

(c) Jams and Sugar.

The fact that only 36 per cent. of the U.N.E. families purchased jam is probably fortuitous; in the other 3 sub-groups the proportions were 71, 70 and 67 per cent. There was little difference in the quantities consumed per man-value in the 4 sub-groups, about 0.6 oz. daily. The quantities of marmalade consumed were small in all groups. The average sugar consumption by all the sub-groups together was about 2.4 ozs., 280 Calories per man daily.

(d) Fruit.

Apples were used fairly extensively by all groups, the proportions of families who used them being 75, 64, 78 and 100 in the 4 sub-groups in the order in which they appear in the table. The percentages for bananas were 46, 43, 50 and 70, thus suggesting slightly higher consumption by the employed families. Other fruits consumed by a minority of the families included oranges, pears, plums, grapes, bilberries, blackberries, grape-fruit and dried fruits.

Potatoes were consumed to an almost equal extent by the 4 sub-groups, about 5.9 ozs., 146 Calories, per man daily. It is noteworthy that nearly twice this amount of energy was derived

from sugar and nearly seven times from white bread. The popularity of the remaining vegetables was in the following descending order: onions, fresh tomatoes, carrots, cabbage, dried peas, turnips, leeks, beetroot, cauliflower, lettuce. The low consumption of lettuce is surprising in view of the season, September. The data suggest that cauliflower was used more extensively by the E.O.H. families, 44 per cent. against 10—14 per cent. of the others. The reason was probably a question of taste or habit, it could scarcely be price, because this vegetable was least popular with the E.N.E. families.

Other vegetables consumed by a minority of the families were cucumber, rhubarb, parsnips, green peas, french beans, radish and celery.

(e) Dairy Products.

Consumption of dairy products as regards those where statistical differences were found to occur has been dealt with in sub-paragraph (a) above, but the following additional information is of interest. Only one unemployed family but three employed families purchased cream. A relatively large number of families, about 80 per cent., consumed cheese, which is one of the best and cheapest sources of animal protein and calcium. The average amount consumed per man-value of the employed exceeded that of the unemployed by about 44 per cent. This difference is probably due to the fact that cheese sandwiches feature largely in the workman's mid-day lunch.

Eggs were cheap in Newcastle at the time of the study, hence their popularity which was not far from equal in the 4 sub-groups.

The great majority of families in all groups purchased condensed skimmed milk, the percentages varying from 67 per cent. in the E.O.H. to 92 per cent. in the U.O.H. The amounts consumed per man were almost equal in the U.N.E. and in the two employed sub-groups and these amounts were exceeded by the U.O.H. families by about 35 per cent. About one-fifth of the employed families purchased condensed full cream milk, but the number of unemployed families who used it was almost negligible. It would, therefore, appear that condensed skimmed milk was more popular with the unemployed and condensed whole milk with the employed. The reason for this qualitative difference is most probably one of price; for condensed skimmed milk at the time generally cost about 3d. and condensed whole milk about 6d. per lb. tin. Although these data only suggest the existence of the above difference the case for its reality is greatly strengthened

by the fact that the unemployed purchased all their food at a lower rate than the employed.

The great bulk of the dried milk was consumed by the unemployed, particularly the U.O.H. families. Most of it was a proprietary brand, supplied gratis from the Welfare Centres.

(f) Fish.

The low consumption of fish is surprising but the paucity of data makes it impossible to say whether it was more popular with one or other group or sub-group. There is no indication that expensive fishes were purchased instead of equally, or more, nutritious fish of the cheaper varieties such as haddock and herring. Herrings were cheap in Newcastle at the time of the survey and the small amounts consumed suggest that this class of fish is not popular with the working classes of the city.

(g) Meat.

The consumption of bacon has already been considered in sub-paragraph (a). Stewed beef was by far the most popular meat. This would also account for the popularity of onions and carrots, one or both of which was always cooked along with the meat. Stewing is a very economical method of cooking as there is little or no loss from either the meat or the vegetables. Stewing was more popular than the amount of stewed beef consumed would indicate because it was not always possible to be certain whether the mutton purchased was intended for stewing, roasting or frying. Mutton appeared to be slightly more popular with the employed than the unemployed.

The relatively small consumption of the expensive pork and chicken as well as the preponderance of beef over pork sausages are noteworthy. Evidently rabbit is not popular in Newcastle, in spite of its low cost. Liver was fairly widely used by unemployed and employed alike. Other offal and meat products included black and white puddings (10), kidney (9), head and tripe (4), pig's cheek, cows' heel and heart (3). The figures in brackets are the numbers of families using these foods and are an indication of their popularity.

(h) Canned and Cooked Food.

The consumption of boiled ham has already been discussed in sub-paragraph (a).

Fried potato chips were the only food in this category used by a bare majority of the families, 35 (including one widow's family) out of 69. The next most popular of these foods was fried fish which was purchased by about one-third of all the families. Baked beans, corned beef and meat pies were consumed by not more than one-fourth of the families. Pineapple was the most popular canned fruit and it was apparently favoured more by the employed than by the unemployed. The other more expensive canned fruits, peaches, pears and fruit salad, were also used more by the employed than by the unemployed. The same applies to canned peas, salmon and boiled tongue. Other foods in this category purchased by a small minority of families were apple tart, meat and fish pastes, polony, pickles, sauces, sardines (used by three unemployed families only), wheat flakes and similar cereal products (seven families), fish cakes, and veal and ham. The numbers of families using these foods varied from ten for sauces to two for yeal and ham pies. Proprietary meat extracts and prepared soups were consumed by six and two families, respectively.

In this qualitative analysis it was impossible owing to the small number of families and to the variations in numbers consuming different foods and in quantities consumed to make comparisons between the four sub-groups. There did not appear, however, from the inspection of the detailed results to be any evidence of a marked difference in consumption either qualitative or quantitative as between the O.H. families and N.E. families. Except in the case of fish and chips the relatively expensive and prepared canned foods were bought by a minority of the families, and in regard to many of the foods the number of consumers was extremely small. There is, therefore, little evidence of generally extravagant buying or of serious decline in home cooking by the families studied. Indeed, practically all the bread consumed was home baked. Further evidence of the prevalence of home cooking can be seen by comparing the total consumption by all the families of prepared meat and fish with that of meat and fish which required cooking in the home. Prepared meat and fish, including meat pies and fish cakes, amounted to 97.37 lbs.; while the purchases of meat and fish, including all offal, which required cooking amounted to 470 lbs., i.e., about 5 times the amount of meat and fish purchased in the cooked or prepared state.

Fresh meat was consumed by all 69 families and fresh vegetables by all but one. All the employed and most of the unemployed families had fresh fruit, but more of the employed had fish, canned and cooked meats and canned fruit and vegetables. Two of the 3 widows' families had no fresh fruit or fresh fish, two had canned vegetables but none of them canned fruit.

A partial summary of the data set out in the Appendix is given in Table VIII.

16.—HÆMOGLOBIN CONTENT OF BLOOD.

The estimations were made by Dr. J. B. Tilley, of the Health Department, Newcastle, using the most recent type of Dare hæmoglobinometer, which is graduated so that a reading of 100 per cent. is equivalent to a concentration of 15.9 gm. of hæmoglobin per 100 cc. Dr. Tilley had previously been trained in the use of the instrument and the accuracy of his results with it was checked against readings taken on the same hospital patients by the Haldane instrument. This is standardised so that a reading of 100 per cent. is equivalent to 13.8 gm. hæmoglobin. Readings of the same sample on the Haldane should therefore be somewhat higher than on the Dare instrument. The results obtained were as follows:—

Haldane.	Dare.	Difference
Hb.%	Hb.%	Haldane—Dare.
72	69	3
48	41	7
67	65	2
87	86	1
89	81	8
57	49	8
67	58	9

On the average the Dare readings are about 5 per cent. below the Haldane readings but differences as high as 9 per cent. were obtained. In collaboration with others, many more comparisons of the Dare with the Haldane and photo-electric cell methods have been made. The Dare nearly always gave lower results but the maximum difference was never greater than 9 per cent. of hæmoglobin. On account of these experiences it was decided to regard Dare readings as understating by approximately 10% the values which would be obtained with the Haldane instrument on the same bloods. The result of this decision, therefore, is that the lower limit of normality of hæmoglobin percentage on the Dare instrument is placed 10% lower than on the Haldane. Since the latter type of hæmoglobinometer is most frequently

employed in this country it is desirable, for purposes of comparison, to fix as far as possible the critical level at which normality ends and anæmia begins with reference to the Haldane instrument. There is, however, some difference of opinion as to the precise position of these critical levels on the Haldane scale for people of different ages of both sexes. It is clear from consideration of the literature on the subject that further research on healthy people of different ages will be necessary before the critical levels can be definitely established, but it would appear that the levels which would meet with the greatest measure of assent are 80% for males over 12 and 75% for women and children below 12 years. It is probable that the hæmoglobin values of the great majority of cases of anæmia would fall below these levels, which, in accordance with the comparisons already quoted, correspond with 70% and 65% respectively on the Dare instrument. It was, therefore, decided that all hæmoglobin values falling below 70% in males over 12 and below 65% in women and children, should be regarded as indicative of anæmia.

The hæmoglobin values of the employed and unemployed groups are compared in Table IX. The mean hæmoglobin percentages are practically identical in both groups for males, females and children respectively.

As regards the distribution of the values, this was approximately the same for the males of both categories and it may be inferred that there was no difference between the hæmoglobin content of the blood of employed and unemployed males.

The distribution of values shown by the women and children in the employed and unemployed groups show some differences.

The numbers and percentages of **anæmic** persons, as previously defined, were as follows:—

Women.

Employed Group (4 out of 39) = 11%Unemployed Group (12 out of 37) = 32%

Children (under 12).

Employed Group (10 out of 59) = 17%Unemployed Group (17 out of 79) = 22%

As regards the percentage of anæmic women the difference between the employed and unemployed groups is rather more than would be expected to occur by chance. On the other hand there are relatively more women in the employed group who have hæmoglobin percentages of the moderately low order of 65—69%, and rather fewer at the high level of 80% hæmoglobin or over

(Table IX.) There is therefore no evidence of a general shift downwards in the scale of values shown by the unemployed women and the mean values in the two groups, are as previously indicated, nearly identical.

The children in the two groups show the same type of differences, namely, slightly more persons in the unemployed group falling below the critical anæmia level, i.e., 65% hæmoglobin; proportionately fewer at the moderately low level of 65-69% hæmoglobin and more in the high group of 80% or over.

The differences are not more than might have arisen by chance, though they confirm those recorded for the women in showing more individuals in the unemployed group at the two ends of the scale.

Adopting the criterion of anæmia as defined in the preceding paragraphs, this analysis then suggests that there is a somewhat higher proportion of anæmic women amongst the unemployed.

17.—HÆMOGLOBIN VALUES IN RELATION TO INTAKE OF IRON.

In Table X. is shown the distribution of hæmoglobin values in relation to the iron intake per man-value daily. The hæmoglobin values are classified according to whether they fell above or below the suggested anæmia level, 70 per cent. for men and 65 per cent. for women and children. Of the 2 anæmic men, one consumed 5-10 mg. and one 15-20 mg. of iron daily. Of the 16 anæmic women, there were 9 in the 10-15 mg. group, 3 in each of the 15-20 and 20-25 mg. groups, one in the 25-30 mg. group, and none in the 5-10 mg. group. The 27 anæmic children fell into the following groups:-17 in the 10-15 mg. group, 3 in each of the 5-10 and 15-20 mg. groups, 4 in the 20-25 mg. group and none in the 25-30 mg. group. It is, therefore, clear that there is no relation between the hæmoglobin of the blood and the amount of iron in the food consumed during the week of the study. It should not, however, be inferred that there is no relation between diet and the hæmoglobin content of the blood as other nutritional factors have also to be taken into account. Thus the bulk of the hæmoglobin molecule consists of a protein complex, the metabolism of iron is affected by the calcium content of the diet, and anæmia may be caused by a deficiency of vitamin C. It is therefore conceivable that some of the deficiencies in hæmoglobin were caused by deficiency of protein, calcium or vitamin C in the diet.

18.—WEIGHTS AND HEIGHTS OF INDIVIDUALS.

The weights and heights of members of the families studied were recorded in the Welfare Centres, the number of refusals being low. Babies were weighed and measured naked, the height being taken by means of a special measuring apparatus. The school children were measured without shoes and upper garments, while adults were measured in ordinary indoor attire with shoes removed. The averages are shown in Table XI.

In view of the fact that the numbers of children at different ages were too small to allow comparison of weights and heights being made between employed and unemployed families, the data for all the children were grouped together according to the age groups. Every person aged 20 years and over was reckoned as an adult and the measurements of the small number between 14 and 20 years were excluded from the averages. In the unemployed group there were 33 men and 39 women, in the employed group 28 men and 28 women, so that the numbers were large enough to justify averaging them separately. The only noteworthy difference between the means is that between the unemployed and employed men, the latter being the heavier by 13 lbs. 4 ozs., a difference which was found to be significant on statistical analysis. The cause of this difference is not clear. If the lower weight of the unemployed was due to their diet being less adequate than that of the employed, similar differences would be expected in the women and children. The other conceivable causes are:-

- (a) that these men lost their employment in the first instance wholly or partly because of poor physique and
- (b) that the loss of weight was due to wasting of muscular tissue which is a physiological result of lack of exercise.

It is impossible to say to what extent (a) was a contributory cause of the lower weight of the unemployed men. The fact that no material differences were found in the weights of women and children would suggest that (b) was an important factor but not necessarily the only one.

23

SUMMARY AND CONCLUSIONS.

- 1. The intake of food of 69 families in Newcastle selected so as to be representative as far as possible of the working class community was determined quantitatively for a week in September, 1934. The bread-winner in 38 families had been unemployed for from six months up to nine years; 24 of these families lived in old houses and 14 on new estates. The bread-winner was employed in 28 of the families of whom 18 resided in old houses and 10 on new estates. There were also 3 widows' families.
- 2. The average diet of the 69 families supplied, per manvalue daily, 2,960 Calories, 82 gms. of protein (36 from animal sources), 94 of fat, 445 of carbohydrate, 0.63 of calcium, 1.23 of phosphorus and 15 mgs. of iron. The average man-value was 3.6, the expenditure on food per man-value per week being 62.4d. As judged by commonly accepted standards this diet is just about sufficient in all these nutrients, except calcium and phosphorus which are both deficient by about 8 per cent.
- 3. Wide variations were found in the consumption of nutrients of different families. The ranges in the 69 families were:—Calories, 1846—5261; protein 51—161 gm.; calcium 0.24—1.41 gm.; phosphorus 0.61—2.44 gm.; iron 7.2—28.9 mg. per manvalue daily. The variations were more marked in the employed than in the unemployed families.

In the frequency distribution of diets according to Calories, total protein, animal protein and calcium contents, more unemployed than employed fell below a range of intake within which sufficiency is considered to lie, for Calories, animal protein and calcium; more employed than unemployed fell above the ranges in Calories, total protein, animal protein and calcium. It is not suggested that all diets below the ranges are necessarily inadequate for the families concerned, because the variations in nutritional needs of individuals are known to be considerable but it is probable that such diets are within the danger zone in regard to the genesis of malnutrition.

Two employed families had less than 1,900 Calories; 6 unemployed, 3 employed and one widow's family less than 0.4 gm. of calcium; 3 unemployed and 4 employed families less than 60 gm. of protein per man daily. These diets were therefore deficient in these nutrients. Four employed families had 4,500 or more Calories; 3 employed and one unemployed family had 130 gm. or more protein and one employed family over 90 gm. animal protein daily. These intakes were probably excessive.

- 4. The average diet of the unemployed was lower than that of the employed in fat and animal protein; the average diet of the unemployed, new estate, families was lower than that of the employed, new estate, families in Calories, fat and animal protein. These differences were shown on statistical analyses to be greater than would be likely to arise by chance. The differences between the other means could have been due to chance; but, since the frequency distribution and the averages showed the unemployed diets to be consistently lower in every particular than the employed diets, irrespective of housing conditions, it is very probable that the diets of the unemployed as a whole were inferior to those of the employed.
- 5. The amounts of energy and protein obtained per penny of money spent on food were 23 per cent. higher in the unemployed than the employed families. Statistical analysis showed that these differences were unlikely to be due to chance. The unemployed therefore purchased these nutrients at lower cost than the employed. The cost of calcium was the same in both groups of families.

The employed spent 36 per cent. more money on the purchase of food than the unemployed, but there was no indication of markedly excessive purchase of expensive foods to the exclusion of equally nutritious cheaper foods by either group of families.

6. There were no important differences between the average man-values in the different groups.

Rent was 21 per cent. higher in the new housing estates than in the old houses but rent did not appear to be closely related to intake of food.

7. The larger families on the whole had less adequate diets and spent less money on food than the smaller families. The Calorie content of the diets gave a rough indication of their composition in the other essential nutrients considered, except animal

protein and calcium, the amounts of which did not always vary in the same direction as Calories.

8. Significantly more employed than unemployed families consumed cakes, fresh tomatoes, butter, fresh milk, bacon and boiled ham. The differences between the numbers in the two groups is most probably related to the greater outlay on food by the employed families. The new estate families consumed on the average 0.26 pint of liquid milk per man daily, the employed, old houses, families 0.19 pint per day, and the unemployed, old houses, families 0.13 pint per day. Seventy-six per cent. of the unemployed consumed butter and 66 per cent. fresh milk; the figures for the employed were 96 and 100 per cent. respectively.

Only three families consumed liquid skimmed milk, but the majority consumed condensed skimmed milk. One-fifth of the employed but only a negligible number of the unemployed used condensed full cream milk.

9. Nearly all the bread consumed was baked at home. Brown bread was used by 45 per cent. of all families. The average total consumption of bread per man-value daily was 10 ozs. (1,000 Calories). Thirty per cent. of families used oatmeal.

The average consumption of sugar by all families, excluding that present naturally in foods or added to prepared foods and dishes such as jams, biscuits and sweets, was 2.4 ozs. (280 Calories) per man-value daily; the figure for potatoes was 5.9 ozs. (146 Calories).

- 10. Ninety-nine per cent. of families consumed fresh vegetables and 88 per cent. fresh fruit. Thirty per cent. of all families consumed canned vegetables and 17 per cent. canned fruits.
- 11. All 69 families consumed fresh meat, 71 per cent. fresh fish, and 68 per cent. canned or cooked meat and fish.
 - Stewing was the most popular method of cooking meats.
- 13. The consumption of canned, prepared and cooked foods was higher among the employed than the unemployed.
- 14. The hæmoglobin content of the blood was determined in 38 of the men, 76 of the women and 138 of the children included

in the investigation. Using the Dare hæmoglobinometer, readings below 70 per cent. in the men or 65 per cent. in women and in children below 12 years, were considered to be indicative of anæmia. The average hæmoglobin values were: in the men 83%, in the women 73% and in the children 73%.

Adopting the definition of anæmia as previously stated, 2 men (5 per cent. of the total), 16 women (21 per cent.) and 27 children (20 per cent.) were found to be anæmic.

Of the 37 women in the unemployed group, 12 or 32% were anæmic, as compared with only 4, i.e., 10% of the 39 women in the employed group.

There was, however, no general shift downwards in the hæmoglobin level amongst the unemployed group, slightly more in the unemployed group than in the employed having high hæmoglobin values.

No relation could be found between the hæmoglobin levels and the content of iron in the family diets.

15. No difference was found between the heights and weights of children or women in any of the groupings, but the employed men were heavier on the average than the unemployed men by 13.25 lbs. The possible causes of this difference are discussed; it was probably not due to difference in diet, since a similar difference in weight was not found between the women of the employed and unemployed groups.

ACKNOWLEDGMENTS.

From the collection of the basic data to their final analysis and presentation, this report has been the work of many hands.

We would express our most grateful thanks to the Medical Department of the Ministry of Health, not only for information which has been freely placed at our disposal, but also for much unofficial assistance.

In no less measure our best thanks are also due to Dr. Bradford Hill, of the London School of Hygiene and Tropical Medicine, who carried out the statistical work of the report, and to Dr. R. A. McCance and Miss E. M. Widdowson, of King's College Hospital, who undertook a number of highly specialised food analyses and made available for our use certain unpublished results of their own.

To the many other members of the team whose assistance may not have been recognised by name elsewhere, our appreciation and thanks are also extended.

J. A. CHARLES,

Medical Officer of Health.

Health Department,

Town Hall,

Newcastle upon Tyne,

December, 1936.

TABLE I.

Average	e daily l	Average daily Intake of Nutrients per Man-value	Nutrier	its per I	fan-valı	ue; average	ge :	1-value,	Rent, 1	ood Exp	enditure	Man-value, Rent, Food Expenditure and Income.	me.
STANDARDS.	No. of Families	Calories No. 3,000	Protein Gms.	Animal Protein Gms.	Fat Gms.	Carbo- hydrate Gms.	Calcium Gms.	Phos- phorus Gms.	Iron Mgms,	Man- value per Family.	Rent per Family per week Shillings.	Food ex- penditure per man- value per week. Pence.	Income per head per week. Shillings.
All families	69	2,960	81.95	36.31	94.55	445	0.627	1.232	15.3	3,60	8.91	62.40	8.07
Unemployed	38	2,847	79.27	33,03	88.70	433	0.591	1.187	15.0	3,43	8.75	53,94	89.9
Employed	28	3,094	86.24	42.08	103.73	454	0.691	1.307	15.6	3.86	9,33	73.62	66'6
Widows	3	3,143	75.78	23.98	82.94	523	0.497	1.106	15,3	1	1	1	1
Old Houses	44	3,018	84.23	37.55	90.76	452	0.638	1,278	15,3	3,50	8.36	62.15	7.96
New Estate	25	2,857	77.93	34.13	90,12	434	0.608	1.151	15.3	3.80	10.11	63.89	8.25
Unemployed (Old Houses)	24	2,958	83.22	35.72	93.20	446	0.626	1.277	15.4	3,34	7.86	53,5	6.44
Unemployed (New Estate)	14	2,656	72.50	28.43	80.99	411	0.530	1.033	14.4	3.57	10.26	54.3	7.08
Employed (Old Houses)	18	3,100	87.10	41.69	104.12	453	0.674	1.308	15.4	3.71	9.02	72.5	10.10
Employed (New Estate)	10	3,085	84.69	42.79	103.01	454	0.721	1.304	16.1	4.12	68.6	75.6	9.80
Widows(Old Houses)	2	3,012	70.54	22.90	79.84	503	0.463	1.025	13.5	1	1	1	1
Widows (New Estate)	-	3,404	86,26	27.36	89.15	564	0.566	1,269	19,0	1	1	1	ı

TABLE II.

Distribution of Families according to Intake per Man-value daily of Calories, Calcium, Protein and Animal Protein; and Distribution according to Man-value, Rent and Food Expenditure.

					100000000000000000000000000000000000000	Expen				
				Nos.	of Far	MILIES.	,			
	Uı	nemploy		E	mploye		Wid	ows.	3	% of Tetal.
	Old Houses.	New Estate,	% of Total Unem- ployed.	Old Houses,	New Estate,	% of Total Em- ployed,	Old Houses,	New Estate,	Total.	
Calories. 1,750–1,999 2,000–2,249 2,250–2,499		5	31	2 1 2		18	=	-	2 3 12	25
2,500-2,749 2,750-2,999 3,000-3,249 3,250-3,499	4 3 3 1	4 3 2	53	4 2 1 1	4 1 2 1	57	1		17 9 8 4	55
3,500-3,749 3,750-3,999 4,000-4,249 4,250-4,499 4,750-4,749 4,750-4,999 5,000-5,249 5,250-5,499	3 - 3 - - -		16	2 - - 1 1 1	1 - - 1 - -	25	1	11111111	$ \begin{array}{c} 7 \\ \hline 3 \\ \hline 2 \\ \hline 1 \\ 1 \end{array} $	20
Totals	24	14	100	18	10	100	2	1	69	100
Calcium (Gms.) .200299. .300399. .400499.	1 2 5	1 2 2	34	1 1 5		36		=	3 7 14	35
.500599	6 2 4 1 1	5 2 1 1	61	1 3 2 1 1	1 2 - 2	46	1 _ _ _	1 - - -	15 9 7 5 2	55
1.00-1.09 1.10-1.19 1.20-1.29 1.30-1.39 1.40-1.49			5	1 1 -1 -1	1 - 1	18		11111	- 4 1 - 2	10
TOTALS	24	14	100	18	10	100	2	1	69	100
Protein (Gms.) 50–59	2 4	1 4	29	4 2	4	36		=	7 15	32
70–79 80–89 90–99 100–109	7 3 3 3	7 1 1	66	4 1 2 1	$\frac{2}{2}$	43	1 - -	- 1 -	21 6 8 4	56
110-119 120-129 130-139 140-149 150-159 160-169	1 - 1 -		5	1 1 - 1 -	1 1 - -	21			3 1 1 2 -	12
TOTALS	24	14	100	18	10	100	2	1	69	100

Sufficiency of each nutrient is considered to lie between the heavy horizontal lines.

Total Tota				TABLE	11.	contini	ecu.				
Total Tota					N	os. of	FAMILI	ES.			
Animal Protein Gims. New Fotal Unem. Houses. Estate. Ployed. Houses. Estate. Ployed. Houses. Ployed. Ployed. Houses. Ployed. Ployed. Houses. Ployed. Ployed. Ployed. Houses. Ployed. Ployed. Ployed. Houses. Ployed. Ployed. Ployed. Houses. Ployed. Ployed.		Ur	employ	red.	E	mploye	ed.	Wid	ows.		% of
C(Gms.)				Unem-			Em-			Total.	Total.
40-49	(Gms.) 10-19	9	2 6	45	2 5	1 1	32	1 1	<u></u>	6 23	42
Totals	40-49			50	1	1	47	=	=	8	46
Rent. s. d. s. d. 3 0-4 11 1	70-79 80-89	1	Ξ	5	_	1 —	21	=	_	2	12
S. d. s. d. 3 0-4 11	Totals	24	14	100	18	10	100	2	1	69	100
Man-value.	s. d. s. d. 3 0—4 11 5 0—6 11 7 0—8 11 9 0–10 11 11 0–12 11 13 0–14 11	6 12 4	8	1003	4 2 8 1	1			1	10 23 26 3 1	
1.0-1.9	TOTALS	24	14		18	10		2	1	69	
Food Expenditure per Manvalue per Week s. d. s. d. 2 0-2 11 1	1.0-1.9. 2.0-2.9. 3.0-3.9. 4.0-4.9. 5.0-5.9. 6.0-6.9.	5	6		4 1 3 1	1 1 1		1 	_	26 19 11 6 3	
ture per Manvalue per Week s. d. s. d. 2 0-2 11 1 3 0-3 11 5 5 5 2 4 0-4 11 13 6 5 0-5 11 2 2 0-2 11 1 2 0-2 11 2 2 0-2 11 2 3 0-3 11 2 4 0-4 11 1 2 0-5 11 2 2 0-6 0-6 11 2 2 0-7 11 1 3 0-8 11 1 4 1 - 9 0-9 11 - 10 0-10 11 - 11 0-11 11 - 12 0-12 11 - 13 0-13 11 - 14 0-14 11 - 15 0-15 11 -	Totals	24	14		18	10		2	1	69	
TOTALS 24 14 18 10 2 1 69	ture per Man- value per Week s. d. s. d. 2 0—2 11 3 0—3 11 4 0—4 11 5 0—5 11 6 0—6 11 7 0—7 11 8 0—8 11 9 0—9 11 10 0—10 11 11 0—11 11 12 0—12 11 13 0—13 11 14 0—14 11 15 0—15 11	1 5 13 2 2 1 1 — — — — — — — — — — — — — — — —	- - - - - - - - - - - - - - -		1 4 1 - 1 - 1	1 2 1 1 1 		1		15 26 7 4 3 2 5 2 - 2 - 1	

Sufficiency of each nutrient is considered to lie between the heavy horizontal lines.

Comparison of Averages of Diets listed in Tables III., IV. and V. (inset opposite). TABLE VI.

gm. gm. pl 6.67 9 0.97 40 102	Carbo-	Phos-		Man-	Sent per	Rent ner Food Expendi-
2.613 70.0 27.2 78.3 408 0.48 2.983 82.1 38.9 103.9 430 0.67 3,934 113.9 55.8 129.4 579 0.97 14 17 43 33 5 40	hydrate. Calcius	m. phorus.	Iron.		week.	ture per Man
2.613 70.0 27.2 78.3 408 0.48 2,983 82.1 38.9 103.9 430 0.67 3,934 113.9 55.8 129.4 579 0.97 14 17 43 33 5 40 50 63 105 65 42 102		gm.	mg. Fa	Family.	s. d.	value per week pence.
2,983 82.1 38.9 103.9 430 0.67 3,934 113.9 55.8 129.4 579 0.97 14 17 43 33 5 40 50 63 105 65 42 102		1.02	13.2	4.25	4 6	, 27.8
3,934 113.9 55.8 129.4 57.9 0.97 14 17 43 33 5 40 50 63 105 65 42 102		1.25	16.9	2.91	9 5	71.6
14 17 43 33 5 40 50 63 105 65 42 102	-	1.80	19.8	2.48	8 1	102.6
50 63 105 65 42 102		22	801	- 46	109	157
	42 102	7.1	20	- 71	80	269

TABLE III.

Diets failing below the Range in either Energy, Protein, Animal Protein or Calcium (see Table II.):
Average Intake of Nutrients per Man-value Daily and Food Expenditure per Man-value Weekly in Groupings
according to Level of Calories and Man-value.

No. of Families	Calories No.	Protein Gms.	Animal Protein Gms.	Fat Gms.	Carbo- hydrate Gms.	Calcium Gms.	Phos- phorus Gms.	Iron Mgms.	Man- value per Family.	Rent per Family per week	penditure per Man- value per week. Pence.
01	1,847	54,53	22,23	58.71	275	0,356	0.805	9.90	4.58	3. d.	29.6d.
15	2,301	64.48	27.87	68.36	357	0,442	0.937	12.20	4.35	4 3	28.5d.
18	2,689	71.29	28.03	81.06	420	0,508	1.042	13.30	4.49	4 10	25.7d.
+	3,224	80.76	25.04	99.34	502	0,456	1.079	15,20	3,04	4 1	33,3d.
01	3,804	93.06	23,55	105.27	621	0,666	1.603	18.40	3,45	01	39.6d.
1	2,767	78.24	27.94	83.76	428	0.599	1.179	13.80	2,49	60	42.2d.
12	2,722	72,18	28.21	81.91	424	0.467	1.051	13.70	3,54	4 10	30,9d.
11	2,576	68.59	27.41	73.28	411	0,502	1.025	13.50	4.30	4 9	23,3d.
9	2,487	64.48	24.68	76.76	385	0,398	0.930	12,30	5.64	C1	19,4d.
83	2,275	61.19	26.73	77.11	334	0.369	0.818	10.60	6.26	3 7	19,8d.
0.0	2,497	65.01	25,21	71.39	388	0.412	0.879	12.00	7.30	4 3	22.0d.

TABLE IV.

Diets falling within the Range in either Energy, Protein, Animal Protein or Calcium (see Table II.):

Average Intake of Nutrients per Man-value Daily and Food Expenditure per Man-value Weekly in Groupings
according to Level of Calories and Man-value.

	No. of Families	No. of Calories Families No.	Protein Gms.	Animal Protein Gms.	Fat Gms.	Carbo- hydrate Gms.	Calcium Gms.	Phos- phorus Gms.	Iron Mgms.	Man- value per Family.	Rent per Family per week	penditure per Man- value per week. Pence.
a. Calories. 2,500—2,999	7	2,822	78.98	36.60	97.00	408	0.651	1.199	16.5	2.99	s. d. 10 3	. 64.0d.
3,000—3,499	7	3,144	85.27	41.30	110.76	451	969'0	1.298	17.2	2.82	8 6	79.2d.
b. Man-value.	6	3,058	82.55	40.12	109.06	437	689.0	1.269	17.3	2.67	4	74.5d.
3-3.9	20	2,847	81,36	36,85	94.55	418	0,645	1.212	16.0	3,31	11 3	66.2d.

TABLE V.

Diets falling above the Range in either Energy, Protein, Animal Protein or Calcium (see Table II.):

Average Intake of Nutrients per Man-value Daily and Food Expenditure per Man-value Weekly in Groupings according to Level of Calories and Man-value.

	No. of Families	No. of Calories Families No.	Protein Gms.	Animal Protein Gms.	Fat Gms.	Carbo- hydrate Gms.	Calcium Gms.	Phos- phorus Gms.	Iron Mgms.	Man- value Per Family.	Rent per Family per week	penditure per Man- value per week. Pence.
a. Calories. Below 3,500	es.	2,980	92,73	61.95	120.07	382	906'0	1.483	16.3	1.98	s. d. 7 10	119.6d.
3,500—3,999	7	3,602	101.78	45,98	111.46	548	0.840	1.599	17.7	2.69	8 11	83.4d.
4,000—4,499	8	4,138	121.41	56.65	145,51	585	1.022	2.046	20.1	2,66	7 9	72.6d.
4,500 4,999	61	4,525	128.26	64,83	144.50	878	1.159	1.931	24.1	2,53	6 7	132,8d.
Above 5,000	67	5,156	151.64	73,83	162,10	773	1.231	2,333	26.2	1.93 *One	*7 3 Rent Fre	167,5d.
b. Man-value. 1-1.9	01	3,972	126.06	77.37	155,25	517	0.946	1.804	20.6	1.83	7 0	135,9d.
2-2.9	11	4,035	116.39	57.97	136,06	586	1.039	1.865	20.3	2,33	7 6	108,4d.
3—3.9	C S	3,554	104.97	41,38	88.40	584	0.821	1.731	18,4	3.18	8 10	55.7d.
4-4.9	1	3,509	79,55	17.91	86.24	603	0.579	1.240	16.3	4.06	0 8	60.50

		4				
					3 5 5 .	

TABLE VII.

Average Amounts of Nutrients purchased per Penny.

	UNEMP	LOYED (38 Fa	milies).	EMPL	OYED (28 Fam	illies).
	Old Houses	New Estate	All	Old Houses	New Estate	All
	(24)	(14)	Unemployed.	(18)	(10)	Employed.
Protein (Gms.)		9.34	9.43	7.64	7.91	7.74
Range		4:61–12.74	4.61-12.74	4.45-11.25	5.42-12.89	4.45–12.89
Calcium(Gms.) Range		.061	.058 .037099	.054 .040084	.062	.057 .040094
Calories	346.7	353.0	349.0	277.5	293.5	283.2
Range	274.8–467.4	185.2-543.2	185.2-543.2	155.2-451.5	193.8-467.5	155.2-467.5

Table VIII.

Numbers and Percentages of Families who consumed the following Categories of Foods.

GROUP.	Total number of Families.	Fresh Fruit.	Fresh Vege- tables,	Fish (includ- ing Fried).	Fresh Meat.	Canned or Cooked Meat and Fish	Canned Vege- tables.	Canned Fruit.
Unemployed								
(Old Houses)	25	21	25	19	25	15	6	2
Percentage		84	100	76	100	60	24	2 8
Unemployed								
(New Estate)	14	12	14	9	14	6	- 2	2
Percentage		86	100	64	100	43	14	14
Employed								
(Old Houses)	17	17	17	13	17	13	6	4
Percentage		100	100	76	100	76	35	21
Employed								
(New Estate)	10	10	10	7	10	10	5	4
Percentage		100	100	70	100	100	50	40
Widows	3	1	2	1	3	3	2	
Percentage	-	33	67	33	100	100	67	-
TOTAL	69	88%	99%	71%	100%	68%	30%	17%

TABLE IX.

Distribution of Hæmoglobin Values.

(Percentages stated are on the Dare scale.)

Percentage below suggested anaemia level, f.e., 70% in males over 12. 65% in females and children under 12.	9	5	10	32	17	22
Per- centage.	I	1	10	32	17	22
No. below 65% Hb.	1	1	4	12	10	17
Per-	9	5	24	14	27	15
No. 65%- 69% Hb.	-	-	6	5	16	12
Per-	22	15	38	19	34	28
No. 70%- 79% Hb.	4	3	15	7	20	22
Per-	72	80	28	35	22	35
No. 80% Hb. & over.	13	16	11	13	13	28
Mean Hb.%	82	84	74	73	72	74
Number examined.	18	20	39	37	59	79
	Employed	Unemployed	Employed	Unemployed	Employed	Unemployed
Mean Hb.%	63	6	7.3	01	7.3	2
Number examined.	30	00	7.6	0/	136	001
Persons examined.	Molec	Males	Famalas	remaies	Children	12 years)

TABLE X.

Distribution of Hæmoglobin Values according to Iron Intake.

	w.		* .		
50	Belo 70%	0	Belo 65%	1	0
25—30 mg.	At and above 70%	0	At and above 65%	1	ю.
60	Below 70%	0	Below 65%	3	4
20—25 mg.	At and above 70%	3	At and above 65%	4	6
	Below 70%	1	Below 65%	3	8
15—20 mg.	At and above 70%	7	At and above 65%	13	21
5.0	Below 70%	0	Below 65%	6	17
10—15 mg.	At and above 70%	25	At and above 65%	38	71
ò	Below 70%	1	Below 65%	0	8
5—10 mg.	Number of At and above Below Persons. 70% 70% 70% 70% 70% 70% 70% 70% 70%	1	At and above Below 65% 65% 65% 65% 65% 65% 65%	4	7
f Iron.	Number of Persons.	38		76	138
Intake of Iron.	GROUP.	Men		Women	Children be- low 12 years

TOTALS.

... 61

::

ADULTS.

Males ...

Females

TOTALS.

... 94

CHILDREN.
Males ...
Females

Distribution of Weights and Heights of Certain Members of the Families Studied. Weights (lbs.) and Heights (inches). TABLE XI.

A. Aleston	(i.e. over 20 years).	Unemployed.	lb. oz. 130 3 (33 persons)	Employed.	143 7 (28 persons)	Unemployed.	lb. oz. 123 7 (39 persons)	Employed.	125 13 (28 persons)	Unemployed.	66.50 (33 persons)	Employed.	66.54 (28 persons)	Unemployed.	61.58 (39 persons)	Employed.	60.33 (28 persons)
	(i.e. o	All.	lb. oz.		19	All.	lb. oz.		29	All.	66.52		19	AII.	61.06		67
Total persons	20 years.				16				88				56				88
	14-20.	lb. oz.	96 13		01		102 0		1		60.25		01		60.00		-
	9-10. 10-11. 11-12. 12-13. 13-14. 14-20.	lb. oz.	80 15		00		83 14 102 0		21		57.00		00		58.35		01
	12-13.	lb. oz.	72 12		00		63 12		10		55.71		00		52.83		20
	11-12.	lb. oz.	63 3		2		61 8		01		51.33		10		50.94		01
	10-11.	lb. oz.	55 5		10		63 5		9		51.38		52		51.67		9
	9-10.	lb. oz.	53 9		1-		53 15		00		48.48		1-		48.51		00
	8-9.	Ib. oz.	51 4		80		0 19		П		47.89		00		50.00		-
YEARS.	7-8.	Ib. oz.	4		9		43.1		9		44.04		9		43.73		9
AGE IN	6-7.	. Ib. oz.	40 12		ro		41 8		6		42.93		10		42.84		6
	5-6.	. Ib. oz.	40 12		9		37 14		150		41.61	7	9		40.70		10
	4-5.	. Ib. oz.	35 4		=		37 3		Ξ		38.31		11		38.35		Π
	9-	. Ib. oz.	33.4		10		3 31 10		00		36.88		2		36.50		00
	2-3	. Ib. oz.	28 6		00		24 13		4		33.36		00		32.50		4
	1-2.	Ib. oz.	22 9		Π		22 10		00		30.93		11		31.10		00
	6/12 to 11/12.	lb. oz.	19 7		1-		17 15		7		27.14		-1		25.88		4
	Birth to 5/12.	Ib. oz.	15 12		01		12 7		1		26.5		01		25.13		1
			Mean Weight (lbs and ozs.)	7	Persons		Mean Weight (Ibs. and ozs.)	Number of	Persons	Mean Height		Number of	Persons	Mann Holobt		Number of	Persons
	Sex.		Males				Females			Moloc	comm			Pomolo	Lemans		

Appendix.

Principal Foods consumed; Numbers of Families and Average Amounts per Man-value Weekly (Widows' Families excluded).

APPENDIX.

Principal Foods consumed: Numbers of Families and Average Amounts per Man-value Weekly (Widows' Families excluded).

			UNEMPLO	UNEMPLOYED (38).					EMPLOY	EMPLOYED (28).		
	I PIO	old Houses	es (24).	New	Esta	New Estate (14).	PIO	Hous	Old Houses (18).	New	Esta	New Estate (10).
	No. of Families	%	Average amount per man- value per week.	No. of Families	%	Average amount per man- value per week.	No. of Families	%	Average amount per man- value per week.	No. of Families	%	Average amount per man- value per week
CEREALS.			lbs. ozs.			lbs. ozs.			lbs. ozs.			lbs. ozs.
Biscuits	12	20	- 2.85	6	64	- 3.57	11	61	- 3.44	7	20	- 3.22
Cakes	4	17		4	59	- 5,33	7	39	- 8.57	9	09	- 5.51
Flour, white	23	96	4 12.88	14	100	4 9.79	18	100	4 6.31	10	100	4 3.62
Flour, brown	10	42		9	36	- 6.89	12	67	1 0.49	89	30	1 2.44
Oatmeal and Oats	00	33	- 8.46	9	43	- 1.11	8	17	- 2,33	4	40	- 2.99
Rice	20	83	- 2.74	11	79	- 2.73	10	99	- 2.64	6	90	- 2.60
Fapioca and Sago	7	53	- 1.68	C1	14	- 3.61	3	17	- 1.48	3	30	- 0.83
SUGAR AND JAMS. Jams and Jellies Sugar Black Treacle	17 24 4	171 100 17	lbs. ozs. - 4.14 - 15.38 - 1.95	5 1 1	36	lbs. ozs. - 3.83 - 15.24 - 4.74	18 4	100	lbs. ozs. - 4.37 - 15.36 - 2.67	10	100	lbs. ozs. - 4.51 1 4.96 - 2.88

			UNEMPLO	UNEMPLOYED (38).						EMPLOY	EMPLOYED (28).		
	Old Houses	Hous	es (24).	New	New Estate (14).	te (1.	4).	Old	Hous	Old Houses (18).	New	Esta	New Estate (10).
	No. of Families	%	Average amount per man- value per week.	No. of Families	%	Ave am per va va por	Average amount per man- value por week.	No. of Families	%	Average amount per man- value per week.	No. of Families	%	Average amount per man- value per week
FATS AND DAIRY PRODUCTS.			Ibs. ozs.			lbs.	ozs.			lbs. ozs.			lbs. ozs.
Butter	18	75	- 5.38	111	79	1	4.90	17	94	- 6.75	10	100	- 8.25
Margarine	19	79	- 7.47	11	79	1	5,65	13	72	- 5.16	00	80	- 3.06
Cream	1	1	1	1	1	1	1.39	2	11	- 1.28	1	10	- 1.13
Cheese	20	83	- 1.87	10	71	1	1.62	15	83	- 2.61	00	80	- 2.38
Eggs	23	96	- 4.24	12	86	1	5.25	18	100	- 4.22	10	100	- 6.71
Milk, fresh	16	67	.90 pt.	6	64	1.81	pts.	18	100	1.33 pts.	10	100	1.80 pts.
Milk, skimmed	1	4	.24 pt.		7	1.00	pt.	1	1	٠.	1	10	0.17 pt.
Milk, condensed			lbs. ozs.			lbs.	ozs.			lbs. ozs.			lbs. ozs.
*Sk. Sw.	22	92	- 6.36	. 11	79	1	4.54	12	67	- 4.92	6	06	- 4.66
*F.C.Sw.	1	7	- 7.86	1	7	1	5.06	3	17	- 5.41	01	20	- 5.39
Milk, Ev.F.C.		-		c	-		0 10	c	1		o	30	0 40
Dried Milk Pro-	1	+	00.6 -	4	*	1	0.72	0	17	00.6	0	20	1.7
prietary Brand	15	62	- 6.47	20	36	1	3.90	1	9	- 2.86	-	10	- 10.76
Other Dried Milks	1	1	1	1	1	1	1	4	22	- 4.75	-	10	5.37

* Sk. Sw. = Skimmed, sweetened.
F.C. Sw. = Full cream, sweetened.
Ev. F.C. Unsw. = Evaporated, full cream, unsweetened.

			UNEMPLO	UNEMPLOYED (38).					EMPLOY	EMPLOYED (28).		
	PIO	Old Houses	es (24).	New	Esta	New Estate (14).	Old	Hous	Old Houses (18).	New	Esta	New Estate (10).
	No. of Families	%	Average amount per man- value per week.	No. of Families	000	Average amount per man- value per week	No. of Families	%	Average amount per man- value per week.	No. of Families	%	Average amount per man- value per week.
FRUIT AND			lbs. ozs.			lbs. ozs.			lbs. ozs.			lbs. ozs.
Apples.	18	75	5.58	6	64	8.23	11	78	7.45	10	100	4 95
Bananas	111	46	- 3.08	9	43	- 3.00		20	- 4.17	7	70	- 3.34
	80	33	- 2,84	2	14	5.92	ıc	28	4.75	00	30	- 9.13
Cabbage	6	37	- 12.16	œ	57	8.17		44	- 7.60	9	09	7.92
Carrots	14	58	- 3.45	6	64	- 3.04		72	- 3,30	7	70	5.61
Cauliflower	8	12	- 4.34	2	14	7.82		44	- 5.94	-	10	- 1.24
Leeks	00	33	- 2,55	7	50	- 5.60		17		1	10	- 2.82
Lettuce	3	12	- 2.04	4	29	- 3,10		11	- 3.07	7	40	96.0 -
Onions	24	100	- 4.80	13	93	- 6.72		94		10	100	- 4.23
Peas (Dried)	13	54	- 3.76	00	36	- 2.40		33	- 2.64	10	20	- 2.13
Potatoes	24	100	2 12,69	14	100	2 7.01	-11	100	200	10	100	2 3.73
Tomatoes, fresh	19	79	- 7.30	00	57	- 5.62	16	68	- 6.43	10	100	
Turnips	=======================================	46	- 5,85	20	36	- 7.73		39	86.9 -	4	40	66.9 -
FISH.			lbs. ozs.			lbs. ozs.			lbs. ozs.			lbs ozs
Cod	01	00		1	1	1	4	22	- 10.92	-	10	
Haddock	4	17	- 5.01	1	1	- 3.07	4	22	- 4.68	01	20	- 11.40
Herring, fresh	9	21	- 10.24	03	21	- 3.75	_	9	- 11.33	1	10	- 7.36
Kippers	00	33	- 3.78	4	53	- 3.05		22	- 1.72	1	10	- 1.84
Sole	1	1	1	1	7	- 1.11	1	1	1		1	1

			UNEMPLO	UNEMPLOYED (38).					EMPLOY	EMPLOYED (28).		
	PIO	Hous	Old Houses (24).	New	Esta	New Estate (14).	PIO	Hous	Old Houses (18).	New	Esta	New Estate (10).
	No. of Families	%	Average amount per man- value per week.	No. of Families	%	Average amount per man- value per week.	No. of Families	%	Average amount per man- value per weck.	No. of Families	%	Average amount per man- value per week
MEAT.			lbs. ozs.			lbs. ozs.			lbs. ozs.			lbs. ozs.
Bacon	16	67	- 4.19	10	71		16	68	- 5.56	10	100	
Beef, roast	14	58	- 10.27	7	50	- 7.57	1	39	- 10,34	9	09	- 7.35
Beef, stewed	21	87	- 5.65	14	100	- 7.42	18	100	- 7.23	10	100	- 4.41
Liver	7	59	- 2.29	3	21	99'1 -	1	9	- 1.43	4	40	- 1.65
Mutton, stewed	-	,										
Mutton, roast and fried	12	20	- 6.87	9	43	- 4.25	12	67	- 8.39	9	09	- 4.65
Pork	4	17	- 4.09	1	7	- 6.68	5	28	- 2.76	1	1	1
Rabbit	2 .	00	- 2.87	1	1	1	9	33	- 8.13	1.	10	- 11.80
Chicken	1	4	- 8.36	1	1	1	1	1	1	1	1	1
Sausage, beef	19	79	- 6.03	12	98	- 5,48	13	72	66.9 -	6	06	- 5.52
Sausage nork	-	4	- 1.90	1	7	- 6.32	3	17	- 6.73	1	1	1

			UNEMPLO	UNEMPLOYED (38).					EMPLOY	EMPLOYED (28).			
	PIO	Old Houses	ses (24).	New	Esta	New Estate (14).	DIO	Hous	Old Houses (18).	New	Esta	New Estate (10).	
	No. of Families	0,	Average amount per man- value per week.	No. of Families	%	Average amount per man- value per week.	No. of Families	%	Average amount per man- value per week.	No. of Families	%	Average amount per man- value per week	
CANNED AND			lbs. ozs.			lbs. ozs.			lbs. ozs.			lbs. ozs.	
Baked Beans	4	17	- 2.91	5	14	- 5.01	3	17	- 4.26	+	40	- 2.45	25
Corned Beef	7	59	- 2.21	57	14	- 2.37	5	Ξ	- 2.50	9	09	- 3.59	
	7	29	- 6.76	4	59	- 5.73	ic	28	- 4.41	9	09	- 5.13	
Fried Chips	15	62	- 10.46	4	59	- 7.61	6	50	6.79	9	09	- 7.09	
Boiled Ham	67	00	- 2.44	00	21	- 1,19	80	44	- 1.44	20	50	- 2.10	
Ice Cream	1	1	1	1	7	- 1.67	1	9	- 2.01	2	20	- 0.55	
Meat Pies	3	12	- 3.13	4	58	- 13,18	57	11	- 2.06	4	40	- 2,98	
Pineapple, tinned	73	00	- 10,07	-	7		2	Ξ		3	30	- 4.45	
Peaches, tinned	1	1	1	1	7	- 4.46	1	9	- 7.44	61	20	- 7.82	
Peas, tinned	1	4	- 8.00	1	1	1	67	11	- 5.11	1	10	- 0.26	
Salmon, tinned	3	12	- 5.12	1	1	1	5	=	- 5.02	01	20	- 3.41	
Sweets	5	21	- 1.50	03	21	- 1.46	4	22	- 1.12	10	20	- 2.37	200
Tomatoes, tinned	2	8	- 5.06	1	1	1	-	9	- 6.83	1	10	- 4.78	_
Tongue, boiled	1	1	1	-	7	- 0.56	01	11	- 1.67	1.	10	- 0.52	
Pears, tinned	1	1	1	1	1	1	1	9		1	1	1	
	1	1	1	1	1	1	1	9	- 3.20	1	I	1	



CITY AND COUNTY OF NEWCASTLE UPON TYNE.

DOMICILIARY MEDICAL SERVICES.

JOINT MEDICAL RELIEF DISTRICT.

REPORT OF THE MEDICAL OFFICER OF HEALTH ON THE WORKING OF THE DOMICILIARY MEDICAL SERVICES IN THE JOINT MEDICAL RELIEF DISTRICT DURING THE PERIOD 1ST MARCH, 1935—29TH FEBRUARY, 1936.

1. **Introductory.** A previous report on this subject was presented to the City Council on the 20th March, 1935. It recorded the establishment of the "open choice" method of providing domiciliary medical services in the City, and described the working of the scheme during the initial period of approximately ten months, *i.e.*, from November 8th, 1933, to August 31st, 1934.

Certain features of the scheme may be briefly summarised as follows:—

- (a) It applied to six of the ten medical relief districts of the City, which were designated the Joint Medical Relief District.
- (b) Drugs and dressings were provided from municipal dispensaries and not at the cost of the medical officers.
- (c) The medical practitioners on the panel, by which the scheme was operated, agreed that during the trial period of twelve months their remuneration, apart from certain special fees, should be derived from a pool of £1,200. This pool when divided amongst the participating practitioners in proportion to the total number of services rendered to their patients, yielded an amount of 5.656 pence per unit service.
- 2. Remuneration of Panel Practitioners. In order to put this obviously inadequate rate of remuneration upon a more equitable basis, discussions took place between representatives of the Health Committee and of the panel practitioners, with the result that a further report, embodying terms which had been agreed by both parties was submitted to the City Council and approved by that body on April 3rd, 1935.

The newly agreed terms provided :-

- (a) That while no further payment could be made to the practitioners in respect of the first year period (November 8th, 1933—November 7th, 1934) during which it had been agreed to operate the scheme on the "pool" basis, a sum of £875 would be added to the amount normally payable on the pool scale for the period November 8th, 1934— February 28th, 1935.
- (b) That as from March 1st, 1935, payments to practitioners would be at the rate of 5/- per quarter per patient treated, but that where a patient had received treatment during two consecutive periods of three months, and was still found to require further medical attention at the end of the second quarter, a payment of £1 should be made to cover treatment for a period of twelve months.
- (c) That the special fees payable in respect of certificates required by Relieving Officers, confinements and fractures, etc., should be continued. The fee for a certificate had previously been fixed at 1/-. Confinements, fractures, etc., were paid for on the recognised scale.
- (d) That the new financial arrangements should be embodied in the official contract of service, and that the whole position should be reviewed as soon as possible after February 29th, 1936.

Owing to the retirement of one of the four remaining salaried District Medical Officers in September, 1934, an additional Medical Relief District was included in the Joint Medical Relief District from that date.

The present report, therefore, deals with the operation of the scheme under the new conditions of service in seven of the ten original medical relief districts during the period of twelve months from March 1st, 1935, to February 29th, 1936.

 Administrative Procedure. The administrative procedure under the new arrangements, and the method of assessing the payments due to members of the panel will now be described in general terms.

A patient requiring treatment presents to the panel practitioner of his choice the relieving officer's order, together with a medical record card which is valid for a period of three months from the date of issue. At the end of that period the record card is forwarded to the Health Department, and if the treatment of the patient has been completed, the practitioner is credited with a fee of 5/-. If, in the opinion of the doctor, treatment is still necessary, a note to that effect is made on the record card, and the Public Assistance Officer is instructed by the Health Department to issue a renewal of treatment card, valid for a further period of three months.

All cards returned to the Health Department are carefully scrutinised by a medical member of the staff, and in certain cases where it is found that a recommendation for further treatment has been made in error, the issue of an additional record card is not authorised. Where the necessity for the continuance of treatment would appear to be open to question, the case is reviewed by the Medical Referee (i.e., Dr. G. P. Harlan, Medical Superintendent of the Newcastle General Hospital) in the presence of the panel practitioner, if the latter so desires. In actual practice recommendations of the panel practitioners as to continuance of the treatment are usually confirmed by the Medical Referee.

The amounts payable to members of the panel are based primarily on the number of three-monthly record cards which they send to the Health Department, each card being valued at 5/-. Practitioners are also credited with one shilling in respect of every medical certificate issued by them at the request of a Relieving Officer, with the fees payable for confinements and other special services, and with the actual cost of emergency drugs dispensed by them to patients.

The initial financial settlement was in respect of the halfyear ended 31st August, 1935, but subsequent payments have been made quarterly.

4. Number of persons in receipt of Public Assistance. The average number of persons in the City who were in receipt of Public Assistance during the twelve-month period March 2nd, 1935—February 29th, 1936, is stated in Table I., where the corresponding numbers for the ten-month period reviewed in the previous report are also given.

TABLE I.

Penson	Average nun	nber of Persons	receiving Publ	ic Assistan
Period.	Men.	Women.	Children.	Total.
30th Oct., 1933 25th Aug., 1934	6,274	6,611	8,510	21,395
2nd March, 1935) 29th Feb., 1936 }	6,379	6,758	7,549	20,687

(It will be appreciated that these numbers indicate the average number of persons receiving public assistance in each of the quoted periods. The total number of individuals who received public assistance at one or another time during the period is obviously greater.)

The 20,687 persons who on an average were in receipt of relief between March 2nd, 1935, and February 29th, 1936, were not evenly distributed throughout the medical relief districts of the City. Two of the three districts excluded from the scheme and still worked by part-time salaried officers had an unduly high proportion of their population on relief. In the remaining districts the proportion of persons on relief was probably about the average for the City.

Taking these facts into consideration it is estimated that the average number of persons in receipt of relief in the area comprised by the seven amalgamated medical relief districts—the Joint Medical Relief District—was approximately 13,500 during the period under consideration.

5. Record of patients treated, etc. Between March 1st, 1935, and February 29th, 1936, in the area of the Joint Medical Relief District, 8,193 persons were attended by medical practitioners under the Domiciliary Medical Scheme.

Of these 611 were "long period" cases receiving treatment for three consecutive quarters or for the whole year. From the medical standpoint these could be regarded as "chronic" patients. The remaining 7,582 were "short period" cases where treatment was required for periods of varying duration, but never throughout three consecutive quarters. Medically speaking the great majority of these cases were acute conditions of short duration, but a number were patients suffering from sub-chronic or recurrent disorders.

More complete details of the groups and sub-groups to which the various types of case were allocated are given in the following tabular statement (Table II.) which also includes information as to the number of medical record cards issued for each type of case, and the cost to the local authority.

TABLE II.

Designation and Sub-group of case.	Medical Grouping (approxi- mate).	Duration of Treatment.	Number of three month- ly Record Cards issued.	Cost of Fatient to Local Authority.
Long Period	Chronic	More than six months con- secutively.	3 or 4	£1
Short Period- Sub-group A.	Acute	One period not exceeding three months.	ī	5/-
" " В.	Acute	Two separate periods, each not exceeding three months.	2	10/-
,, ,, C.	Acute	Three separate periods, each not exceeding three months.	3	15/-
,, ,, D.	Sub- Chronic	Two consecutive periods of three months, not ex- ceeding six months.	2	10/-
,, ,, E.	Sub- Chronic	One period not exceeding three months, followed or preceded by a separate period not exceeding six months.	3	15/-

In Table III. the 8,193 patients treated in the Joint Medical Relief District are distributed amongst the several groups and sub-groups already referred to. The average number of medical services rendered to each patient is also recorded, together with the average remuneration received by the medical practitioner per unit service.

TABLE III.

Designation and Sub- group of case.	Number Treated.	Total No. of services rendered.	Cost to Local Authority.	Average No. of Unit services per patient.	Average remunera- tion per unit service.
Long Period	611	16,145	£ s. d. 611 0 0	26.4	9.08d.
Short Period- Sub-group A. ,, B. ,, C. ,, D. ,, E.	835 82	26,568 7,761 1,120 8,148 3,371	£ s. d. 1,476 15 0 417 10 0 61 10 0 286 10 0 138 15 0	4.5 9.3 13.7 14.2 18.2	s. d. 1 1.34 1 0.91 1 1.18 - 8.44 - 9.88
Total of Short Period Cases	7,582	46,968	2,381 0 0	6.2	s. d. 1 0.17
Total of All Cases. Long and Short Periods	8,193	63,113	2,992 0 0	7.7	- 11.38d.

For a complete understanding of these data one further item of information is required, namely, the actual character of the services rendered to the various types of patient. This is set out for "Long Period" and "Short Period" cases in Table IV.

TABLE IV.

Designation.	No. of Patients.	Attend- ances at Doctor's surgery.	Visits at patient's home.	Total Units of Medical Service.
Long Period Cases.	611	11,126	5,019	16,145
Average per Patient.	-	18,2	8.2	26.4
Short Period Cases.	7,582	29,025	17,943	46,968
Average per Patient.	-	3,8	2.4	6,2

Ratio of Attendances to Visits—Long Period Cases, 2.2 to 1.
Short Period Cases, 1.6 to 1.
All Cases, 1.7 to 1.

- 6. Analysis of services rendered by Panel Practitioners. A number of deductions can doubtless be made from these statistical data. For the present it is sufficient to submit:—
 - (a) That the remuneration per unit service, which amounted to 11.38d. per unit is twice the previous rate of 5.656d.
 - (b) That the rate for all "Short Period" cases (1s. 0.17d.) is 25% higher than the rate for "Long Period" cases (9.08d.)
 - (c) That the highest rate of remuneration (1s. 1.34d.) is obtained from the true "Short Period" case where treatment is given for a period not exceeding three months (Sub-Group A.)
 - (d) That the services rendered to both Long and Short Period cases appear to be reasonable in number. The typical "Short Period" case in which treatment is complete within three months (and frequently in a much shorter period) has 4.5 visits or attendances. The typical chronic case received 26.4 services during the nine months or year during which it was under medical attention. This can be regarded as equivalent to a fortnightly visit or attendance.
 - (e) That the ratio of attendances to visits is a valuable index of the services rendered by the doctors, for it can be assumed as a general rule that a patient attending the surgery takes up less of the practitioner's time than does a visit to the patient's home.

During the first review period (8th November, 1933, to 31st August, 1934) the total numbers of attendances and visits were 18,338 and 23,318 respectively, yielding an attendance to visit ratio of 0.8 to 1., in other words, domiciliary visiting played a more important part in the scheme than the surgery consultation.

The present review period shows a reversal of this state of affairs. The total numbers of attendances and visits were 39,333 and 22,962 respectively, the attendance to visit ratio being 1.7 to 1.

For "Long Period" patients the ratio was 2.1 attendances to 1 visit, as contrasted with 1.6 attendances to 1 visit for the "Short Period" patients. This difference between the two types of case is understandable as many of the long period patients are suffering from chronic conditions which do not prevent them from getting about.

Individual doctors vary in their methods of practice some appear to do the greater part of their work in the surgery, others visit their patients with commendable frequency. The tendency to replace domiciliary visiting by the easier process of surgery consultation cannot be regarded favourably, and its further extension will be to the detriment of the service as a whole.

7. **Expenditure.**—Expenditure upon the Domiciliary Medical Services falls under three heads—Medical Practitioner Services, Dispensary Services and Administrative Charges.

Details of these for the twelve months ending February 29th, 1936, will be given in the following paragraphs, together with any additional information which may appear relevant.

Medical Practitioners' Services.

- (a) Remuneration of Medical Practitioners upon the panel of the Joint Medical Relief District. The method of calculating this is described in paragraph 2 (b) above. The cost for the twelve months under consideration was £2,992.0.0, equivalent to 7s. 3½d. per patient treated.
- (b) Fees for Certificates. 5,634 certificates were completed by Medical Practitioners at the request of Relieving Officers. The cost of these at 1/- per certificate amounted to £281.14.0.

It will be noted that this expenditure is equal to 9.4% of the ordinary remuneration of the Panel Practitioners. If this £281.14.0 is spread over the 63,113 unit medical services rendered to patients it is equivalent to an additional 1.07d. per unit service. Added to the average fee per unit service it increases this from 11.38d. to 1s. 0.45d.

(c) Confinement and other Special Fees. Under this heading £50.11.6 was disbursed during the year. This amount represents payment for 36 confinements.

Owing to the miserable housing conditions of many of the families in receipt of public assistance, the majority of confinements take place at the Princess Mary Maternity Hospital and the Newcastle General Hospital.

No payments in respect of special fees, e.g., fractures, etc., were made during the period under review.

(d) Emergency Medicines. Claims for the cost of emergency medicines were received to a total of £4.16.0.

The total expenditure under heads (b), (c) and (d) was £337.1.6, equivalent to 10d. per patient treated.

Summary of Expenditure on Medical Practitioners' Services :--

Remuneration of Medica	l Prac	titioners	on	£	s.	d.
the Panel				2,992	0	0
Fees for Certificates				281	14	0
Fees for Confinements				50	11	6
Emergency Medicines				4	16	0
				£3,329	1	6

The average expenditure under these various sub-heads amounted to 8s. 1½d. per patient treated.

Dispensary Services. No alteration has been made in the method of providing dispensary services as described in the previous report. There has been, however, a very definite increase in the number of prescriptions dispensed from the two dispensaries. Between November 8th, 1933, and August 31st, 1934, the total number of prescriptions dealt with was 30,219. For the twelvemonth period now under review, 61,357 prescriptions were dispensed. Even when allowances have been made for the extended duration of the second review period and the larger number of patients treated, there is an actual increase in the number of prescriptions dispensed of at least 40%. No satisfactory explanation of this increase can be offered, but it is obviously a matter which will require close supervision.

As a result of this increased work the staffs at both dispensaries have been augmented and this is reflected in the salary costs for the Municipal Dispensary at the Newcastle Dispensary, as set out in the summary statement on page 10.

At the Newcastle General Hospital no definite amount is appropriated for dispensers' salaries. The sum transferred from the Hospital account to the Domiciliary Medical Service account includes both the cost of materials and an allowance for the services of the Dispensary and its staff.

Summary of Dispensary Services Costs :-

(a) Municipal Dispensary, Newcastle Dispensary, New Bridge Street.

	£	s.	d.
Rent, Cleaning, Light, Water, etc.	 200	0	0
Dispensers' Salaries	 296	0	0
Special Saturday afternoon duty	 26	0	0
Drugs, Dressings, etc	 364	0	0
	£886	0	0

Prescriptions dispensed, 33,872. Average cost per prescription, 6.3d.

(b) Newcastle General Hospital. Transfer from Hospital Account	for	£	S.	d.
Drugs and Dispensary Services		562	0	0
Special Saturday afternoon duty		26	0	0
		£588	0	0

Prescriptions dispensed, 27,485. Average cost per prescription, 5.1d.

(c) General.

Total cost of Dispensary Services = £1,474.

Total number of prescriptions dispensed, 61,357.

Average cost per prescription, 5.7d.

Average cost per patient treated = 3s. 7d.

This average cost per prescription dispensed 5.7d. contrasts very favourably with the average cost per prescription under the National Health Insurance Scheme, which is 7.8d.,* more especially when it is remembered that the National Formulary is used in both schemes. The fact that only two dispensaries are available under the Domiciliary Medical Scheme has not been proved to be a serious inconvenience either to patients or their relatives, but with any additional extension of the scheme some decentralisation of the dispensary arrangements would appear advisable.

Administrative Charges.

(a) Clerical Services. At the time when the Domiciliary Medical Services were provided by part-time salaried officers the clerical supervision of their work took up

^{*} This figure has been kindly supplied by the Clerk to the Newcastle upon Tyne Insurance Committee.

approximately one-seventh, i.e., 5 hours weekly, of the time of one intermediate grade clerk (salary £210) at a cost of £30. Following the adoption of the "open choice "method operated on the basis of a financial "pool," with from 45 to 50 doctors participating it was found necessary to employ an additional full-time clerk (salary £182) and to allocate two-fifths, i.e., 14 hours weekly, of the time of the intermediate grade clerk to this purpose. replacement of the " pool " by the new system of remuneration has not resulted in any saving in clerical services. The expenditure on these services chargeable against the Joint Medical Relief District remains at approximately £250 per annum. As the cost of the Domiciliary Medical Services was £5,129 in the twelve months under review the clerical services represent 5% of that total. It has been suggested that this is an excessive proportion, but it is doubtful whether such a contention can be maintained when the clerical administrative costs of certain other sections of the Health Services are considered. Apart from any question of comparative cost the clerical section of the Domiciliary Medical Services is valuable in itself. It is, in fact, a key point of the scheme for through it is maintained a close liaison between the panel practitioner on the one hand and the municipal hospitals, the statutory health services, the domiciliary nursing services and the public assistance department on the other. The liaison work is in addition to its primary statistical and financial functions.

It is possible that by certain modifications in the present method of assessing the remuneration of the medical practitioners, notably by the substitution of an **annual** payment and medical record card for the present **quarterly** payment and card, a reduction could be effected in the work of the clerical section. By such a procedure the number of record cards handled could be reduced by 30%. But it should be remembered that practically 75% of patients are under treatment for a maximum period of three months, and a twelve-month record card issued at the beginning of the incapacity of such a patient would be in use for three months and dormant for nine months. Incidentally, during the dormant period the patient might cease to be a person in receipt of Public Assistance.

The present arrangement whereby record cards are returned quarterly undoubtedly assists the central office in exercising a continuous supervision of the scheme. For "Long Period" or "Chronic" cases the twelve-month record card would have certain advantages.

(b) Printing, Postages. Apart from clerical salaries the only item chargeable as administrative expenditure is printing and postages, totalling £76.

Summary of Administrative Charges.

Salaries (additional)	 	 	£250
Printing, Postages	 	 	76
			£326
			Accessed to the

Average cost per patient treated=9.5d.

Table V. states the total expenditure on the Domiciliary Medical Services during the period 1st March, 1935—29th February, 1936.

TABLE V.

ITEM.	ACTUAL COST YEAR ENDED 31/3/36.	
Salaries	2,992	s. d. 7 3,5
Special Fees (Certificates, Confinements,		, 0.0
Emergency Medicines, etc.)	0.07	- 10
Dispensing Services		3 7
Administrative Charges:		
Printing, Postages	76	1
Additional Clerical Assistance £182 Proportion of Intermediate Grade Clerk's Salary £68	250	- 9.5
Total	€5,129	12 6

8. Number of Practitioners on the Panel of the Joint Medical Relief District. The number of medical practitioners on the panel was 56 on March 1st, 1935, and had increased to 61 by February 29th, 1936. Practically 60% of the National Health Insurance practitioners in the area of the Joint Medical Relief District serve on the Domiciliary Medical Services panel.

- 9. Medical Records. The medical record cards are kept with a degree of accuracy and completeness not inferior to the standard of record-keeping under the National Health Insurance Scheme. Some administrative inconvenience is caused occasionally by doctors failing to return expired record cards to the Health Department promptly.
- 10. Complaints against Practitioners. No complaints were preferred by patients against practitioners on the panel during the period under review. The right to change their doctor is rarely exercised by patients on the grounds of dissatisfaction with the services provided.

These two facts only serve to confirm what has been the invariable experience of all who have been concerned with the working of the "open choice" method. No other scheme of domiciliary medical service has ever achieved so large a measure of acceptance and popularity amongst those members of the community whom it is intended to serve.

- 11. Relations with Relieving Officers. The relations between Medical Practitioners and Relieving Officers have continued excellent throughout the year. There is now a very complete and mutual understanding of their respective rôles in the scheme.
- 12. Certification of Fitness. Here again, experience has given the panel practitioners an increased knowledge of his duties in this matter. Complaints as to laxity in certifying fitness have been relatively few, and it is the considered opinion of the Public Assistance Department that the members of the "open choice" panel discharge this duty even more conscientiously than did their predecessors, the salaried part-time district medical officers. In the few doubtful cases the services of the Medical Referee (Medical Superintendent of the Newcastle General Hospital) are called upon.
- 13. Recommendations for Medical Extras. The view expressed in the previous report that the dangers of excessive recommendations for medical extras had been exaggerated has been substantiated during the past year.

Table VI. states the expenditure on medical extras, and the average cost per relieving officers' order for the three periods:—

a. 1st November, 1932, to 31st August, 1933 (10 months period operated by part-time salaried district medical officers).

- b. 1st November, 1933, to 31st August, 1934 (10 months period operated by panel of practitioners serving on the panel of the Joint Medical Relief District).
- c. 1st March, 1935, to 29th February, 1936 (12 months period under review).

TABLE VI.

Period.	Relieving Officers' Orders.	Milk and Eggs.	Other extras- Cod Liver Oil, etc.	Total Cost.	Cost per Relievin Officers Order.	g
a. { 1st Nov., 1932 } 31st Aug., 1933 }	5,096	297	£ 42	339	s. d. 1 3,9)
b. { 1st Nov., 1933 } 31st Aug., 1934 }	8,836	513	56	569	1 3.4	1
c. { 1st Mar., 1935 } 29th Feb., 1936 }	13,396	657	72	729	1 1,1	1

14. Association with the Newcastle General Hospital. There has been a marked increase in the work of the Newcastle General Hospital since 1930 (Table VII.). To this the success of the Domiciliary Medical Services has certainly contributed but the main cause is the steady progress which the hospital has made in the regard of the inhabitants of the City. There is no evidence that panel practitioners refer cases to hospital unnecessarily.

The hospital provides a general consultative service for outpatients and has established two important clinics for diabetic and anæmia cases. These latter are particularly useful in the case of Public Assistance patients. For example, diabetic patients attend the hospital for blood sugar examinations and the regulation of their insulin dosage. The necessary information on these points is communicated to the panel practitioners together with any suggestions as to special dietetic requirements, which can be recommended for issue as medical extras.

TABLE VII.

Year.	Admissions.	Operations.	Maternity Cases
1930	3,048	596	97
1931	3,598	1,125	99
1932	4,522	1,428	161
1933	4,776	1,560	194
1934	5,544	2,076	225
1935	6,245	2,722	273

- 15. Relations with Statutory Health Services. The association of the Domiciliary Medical Services with the other organised health services of the Local Authority, particularly the Maternity and Child Welfare and Tuberculosis services is being steadily developed.
- 16. Relations with District Nursing Associations. During the course of the past year an agreement was reached whereby the local Nursing Associations provide domiciliary nursing services for any public assistance patient on request by the medical practitioner. This agreement applies to all the Nursing Associations except one which is organised on lines somewhat different from those of the ordinary district association.

The importance of this arrangement cannot be overstated, and the co-operation of the associations has been very welcome.

17. Incidence of Sickness, etc. It has frequently been observed that while the statistics of birth and mortality are now adequately recorded by the Registrar General, there is little information regarding the incidence of sickness apart from notifiable infectious disease.

Details concerning one section of the population are now forthcoming from the records of the National Health Insurance Scheme but these obviously relate to insured persons over the age of 16. The Domiciliary Medical Scheme provides an opportunity for studying sickness and invalidity amongst one large section of the population including males and females of all ages.

The importance of such data is only now beginning to be understood. With proper statistical methods it should be possible to correlate the effect of bad housing conditions, low income levels, etc., upon the health of the public assistance community. The epidemiology of many of the minor infectious diseases could also be studied as well as the incidence of those common causes of incapacity—tonsillitis, rheumatism and gastritis.

Certain elementary facts have already been extracted. Table VIII. (page 16) shows the age and sex distribution of the patients treated under the scheme. There are a number of interesting features in this table, but the only one to which attention is particularly directed is the fact that 41.4 per cent of the clientele of the scheme are children under the age of 15 years.

Table IX. (page 17) gives the classified distribution of the diseases for which patients were treated. This table is based on a similar table in the Annual Report of the Chief Medical Officer of the Ministry of Health for 1933, which records the incidence of disease amongst insured persons.

TABLE VIII.

AGE AND SEX DISTRIBUTION OF PATIENTS TREATED.

(1ST MARCH, 1935—29TH FEBRUARY, 1936).

	9	CHRC	NIC	CHRONIC PATIENTS.	NTS.			ACI	UTE P	ACUTE PATIENTS	TS.		CHR	CHRONIC AND ACUTE PATIENTS.	AND AC	UTE	PATIE	NTS.
	Males.		Fem	Females.	Total Males & Females.	tal Females.	Ma	Males.	Females.	ales.	Total Males & Females.	Total	Males.	les.	Females.	ales.	Total Males & Females.	Total & Females.
2.00	No. tag	Percen- tage of Total.	No.	Percen- tage of Total.	No.	Percen- tage of Total.	No.	Percen- tage of Total.	No.	Percen- tage of Total.	No.	Percen- tage of Total.	No.	Percen- tage of Total.	No.	Percen- tage of Total.	No.	Percen- tage of Total.
-	11	5.2	16	4.0	27	4.4	851	28.2	688	19.5	1,740	22.9	862	26.7	905	18.2	1,767	21.6
-	6	4.2	15	3.8	24	3.9	745	24.7	850	18.6	1,595	21.0	754	23.3	865	17.5	1,619	19.8
1	8	1.4	17	4.3	20	3.3	163	5.4	327	7.2	490	6.5	166	5.1	344	6.9	510	6.2
71		33,3	87	21.8	158	25.9	693	22.9	1,290	28.3	1,983	26.2	764	23.7	1,377	27.8	2,141	26.1
66		46.5	136	34.2	235	38.5	447	14.8	692	16.8	1,216	16.0	546	16.9	902	18.2	1,451	17.7
20		9.4	127	31.9	147	24.0	120	4.0	438	9.6	558	7.4	140	4.3	565	11.4	705	8.6
213		100	398	100	611	100	3,019	100	4,563	100	7,582	100	3,232	100	4,961	100	8,193	100

TABLE IX.

information for insured persons as set out in the Annual Report of the Chief Medical Officer of the Ministry of Proportional distribution of disease amongst acute and chronic Public Assistance Patients. Health for 1933 is given for comparison.

DISEASE. Paper Part 1935 Part 1935 Part 1936 Part 1936 Part 1936 Part 1936 Part 1937 Part 19		Popuoconto	thing Arone	ALE	WOACTE	TIDON TV	NE
Triple DISEASE Curo Per 1,000 Pe		Represents England a	and Wales.	(1st Mar	ch, 1935—2	9th Februar	ту, 1936.)
Influenza Infl		Re ₁ (Year	oort. 1933.)	Ас Ратт	UTE ENTS.	Спр	CONIC ENTS.
Influenza 14,905 118.6 373 34.7 17 17 17 178 12.6 19.0 9.3 24 15.89 12.6 12.6 19.5 10.0 9.3 24 15.89 12.6 10.0 9.3 24 15.89 12.6 10.0 9.3 24 10.5 10.0 10.5 10.0 9.3 24 10.5 10.0 10.5 10.0 10.5 10.0 10.5 10.0 10.5 10.0 10.5 10.0 10.5 10.0 10.5 10.0 10.5 10.0 10.5 10.0 10.5 10.0 10.5 10.0 10.5 10.0 10.5 10.0 10.5 10.0 10.5 10.0 1.0 10.5 10.0 1.0 10.5 10.0 1.0		No.	Per 1,000 of Total.	No.	Per 1,000 of Total.	No.	Per 1,000 of Total.
Tuberculosis, all forms Tuberculosis, all forms 1,589 12.6 513 47.7 165 Organic Heart Disease 1,578 12.6 513 47.7 165 Anæmia 1,589 12.6 495 46.1 61 Bronchitis, tonsillitis, nasal catarrh, cold, etc. 29,698 236.4 2,773 258.2 244 Bronchitis, tonsillitis, nasal catarrh, cold, etc. 1,698 236.4 2,773 258.2 244 Diseases of the digestive system 1,698 10.2 28 64 10.2 131 Diseases of the genito-urinary system 13,915 110.7 1,098 102.2 131 Diseases of the genito-urinary system 3,854 30.7 394 36.7 55 Diseases of the genito-urinary system 3,854 30.7 394 36.7 55 Skin Diseases 10,808 86.0 236 22.1 14 Abscess, boils, and other septic conditions 11,329 90.2 566 52.8 94 <	1	14,905	118.6	373	34.7	17	13.9
Organic Heart Disease 1,589 12.6 513 47.7 165 Anæmia Anæmia 1,578 12.6 495 46.1 61 Bronchitis, tonsilitis, nasal catarrh, cold, etc. 29,698 236.4 2,773 258.2 244 Pheumonia and other diseases of the digestive system 1,699 113.6 283 26.4 64 Diseases of the genito-urinary system 13,915 110.7 1,098 102.2 131 Diseases of the genito-urinary system 3,854 30.7 394 36.7 55 Diseases of the genito-urinary system 3,854 30.7 394 36.7 55 Diseases of the genito-urinary system 3,854 30.7 394 36.7 55 Skin Diseases 6,112 48.6 43.1 40.1 24 Abscess, bolls, and other septic conditions 8,808 70.1 429 39.9 19 Lumbago, relumnatism, etc. 6,115 48.7 783 52.8 10 Malignant Disease		902	7.2	100	9.3	24	19.6
Anæmia Anæmia and other diseases of the respiratory system Piseases of the digestive system (1,699 13,6 2,773 258.2 244 Preumonia and other diseases of the respiratory system (1,699 13,6 2,873 258.2 244 Diseases of the digestive system (1,699 13,6 2,83 102.2 131 Diseases of the digestive system (1,699 13,915 110,7 1,098 102.2 131 Diseases of the nervous system and special senses. (2,046 56.1 48.6 43.3 112 Nimitaries and accidents (1,000 10,739 10,809 10,809 10,809 10,809 Lumbago, rheumatism, etc. (2,046 2,046 2,046 2,049 1,000 1,222† Debility, neuralgia and headache (2,046 2,044 13 Malignant Disease (2,046 2,046 2,044 1,000 1,222† Total (1,000 10,739* 1,000 1,222†	_	1,589	12.6	513	47.7	165	135.0
Bronchitis, tonsillitis, nasal catarrh, cold, etc. 29,698 236.4 2,773 258.2 244 Pneumonia and other diseases of the respiratory system 1,699 13.6 283 26.4 64 Diseases of the digestive system 13,915 110.7 1,098 102.2 131 Diseases of the genito-urinary system 3,854 30.7 394 36.7 55 Diseases of the genito-urinary system 3,854 30.7 465 43.1 112 Diseases of the nervous system and special senses. 7,046 56.1 465 43.3 112 Skin Diseases of the nervous system and special senses. 6,112 48.6 40.1 22.1 14 Injuries and accidents 8,60 236 22.1 14 40.1 14 Abscess, boils, and other septic conditions 8,808 70.1 429 39.9 19 Lumbago, rheumatism, etc. 8,808 70.1 48.7 78.3 72.9 80 Malignant Disease 8,808 70.1 48.7 78.3	. Anæmia	1,578	12.6	495	46.1	61	49.9
Pneumonia and other diseases of the respiratory system 1,699 13.6 283 26.4 64 Diseases of the digestive system 13,915 110.7 1,098 102.2 131 Diseases of the digestive system 3854 30.7 394 36.7 55 Diseases of the nervous system and special senses. 7,046 56.1 465 43.3 112 Skin Diseases of the nervous system and special senses. 10,809 86.0 236 43.3 112 Skin Diseases of the nervous system and special senses. 10,809 86.0 236 22.1 14 Abscases, boils, and other septic conditions. 11,329 86.0 236 52.8 94 Lumbago, theumatism, etc. 8,808 70.1 48.7 783 72.9 80 Malignant Disease 11,329 90.2 566 52.8 94 Other Diseases 7,063 56.1 598 55.7 82 Infectious Diseases 24.4 13 Total 1,000 10,739*	Bronchitis, tonsillitis, nasal catarrh, cold, etc	29,698	236.4	2,773	258.2	244	200.0
Diseases of the digestive system 13,915 110.7 1,098 102.2 131 Diseases of the genito-urinary system 3,854 30.7 394 36.7 55 Diseases of the nervous system and special senses. 7,046 56.1 465 43.3 112 Skin Diseases 48.6 48.6 43.1 40.1 24 Injuries and accidents 8,808 70.1 429 39.9 19 Abscess, boils, and other septic conditions 8,808 70.1 429 39.9 19 Lumbago, rheumatism, etc. 11,329 90.2 566 52.8 94 Debility, neuralgia and headache 6,115 48.7 783 72.9 80 Malignant Disease 7,063 56.1 598 55.7 82 Other Diseases 7,063 56.1 598 55.7 82 Infectious Diseases 7,063 56.1 24.4 13 Puerperal state 20.2 1,000 10,739* 1,000 1,222†	. Pneumonia and other diseases of the respiratory system	1,699	13.6	283	26.4	64	52,4
Diseases of the genito-urinary system 3,854 30.7 394 36.7 55 Diseases of the nervous system and special senses. 7,046 56.1 465 43.3 112 Skin Diseases 10,809 86.0 236 43.1 24 Injuries and accidents 10,809 86.0 236 22.1 14 Abscess, boils, and other septic conditions 11,329 90.2 566 52.8 94 Lumbago, rheumatism, etc. 11,329 90.2 566 52.8 94 Debility, neuralgia and headache 6,115 48.7 783 72.9 80 Malignant Disease 221 1.8 39 3.6 17 Other Diseases - 901 83.9 6 Infectious Diseases - 901 83.9 6 Puerperal state - - 901 83.9 6 Puerperal state - - 262 24.4 13 Total 125,646 1,000 10,739* 1,000 1,222‡	_	13,915	110.7	1,098	102.2	131	107.2
Diseases of the nervous system and special senses. 7,046 56.1 465 43.3 112 Skin Diseases Skin Diseases 48.6 431 40.1 24 Injuries and accidents 10,809 86.0 236 22.1 14 Abscess, boils, and other septic conditions 11,329 90.2 566 52.8 94 Lumbago, rheumatism, etc. 11,329 90.2 566 52.8 94 Debility, neuralgia and headache 6,115 48.7 783 72.9 80 Malignant Disease 7,063 56.1 598 55.7 82 Infectious Diseases - 901 83.9 6 Puerperal state - - 262 24.4 13 Total 125,646 1,000 10,739* 1,000 1,222†		3,854	30.7	394	36.7	55	45.0
Skin Diseases 48.6 48.6 48.1 40.1 24 Injuries and accidents 10,809 86.0 236 22.1 14 Abscess, boils, and other septic conditions 8.808 70.1 429 39.9 19 Lumbago, rheumatism, etc. 11,329 90.2 566 52.8 94 Debility, neuralgia and headache 6,115 48.7 783 72.9 80 Malignant Disease 1.8 39 3.6 17 Other Diseases 56.1 598 55.7 82 Infectious Diseases - 901 83.9 6 Puerperal state - 262 24.4 13 TOTAL 125,646 1,000 10,739* 1,000 1,222†		7,046	56.1	465	43.3	112	91.6
Injuries and accidents 10,809 86.0 236 22.1 14 Abscess, boils, and other septic conditions 8.808 70.1 429 39.9 19 Lumbago, rheumatism, etc. 11,329 90.2 566 52.8 94 Lumbago, rheumatism, etc. 11,329 90.2 566 52.8 94 Debility, neuralgia and headache 6,115 48.7 783 72.9 80 Malignant Disease 221 1.8 39 3.6 17 Other Diseases 56.1 598 55.7 82 Infectious Diseases 56.2 24.4 13 Puerperal state 262 24.4 13 TOTAL 125,646 1,000 10,739* 1,000 1,222†		6,112	48.6	431	40.1	24	19.6
Abscess, boils, and other septic conditions 8,808 70.1 429 39.9 19 Lumbago, rheumatism, etc. 566 52.8 94 Lumbago, rheumatism, etc. 521 11,329 90.2 566 52.8 94 Debility, neuralgia and headache 6,115 48.7 783 72.9 80 Malignant Disease 72.0 39 3.6 17 Other Diseases 6 52.8 94 Infectious Disease 72.9 80 Infectious Diseases 6 6 Puerperal state 70.00 10,739* 1,000 1,222†		10,809	86.0	236	22.1	14	11.5
Lumbago, rheumatism, etc. 11,329 99.2 566 52.8 94 Debility, neuralgia and headache 6,115 48.7 783 72.9 80 Malignant Disease 1.8 39 3.6 17 Other Diseases 56.1 598 55.7 82 Infectious Diseases 901 83.9 6 Puerperal state 262 24.4 13 Total 125,646 1,000 10,739* 1,000 1,222†		8,808	70.1	429	39.9	19	15.5
Debility, neuralgia and headache 6,115 48.7 783 72.9 80 Malignant Disease 39 3.6 17 Other Diseases 56.1 598 55.7 82 Infectious Diseases 901 83.9 6 Puerperal state 262 24.4 13 TOTAL 125,646 1,000 10,739* 1,000 1,222†		11,329	90.2	566	52.8	94	76.9
Malignant Disease 1.8 39 3.6 17 Other Diseases 7,063 56.1 598 55.7 82 Infectious Diseases - - 901 83.9 6 Puerperal state - - 262 24.4 13 TOTAL 125,646 1,000 10,739* 1,000 1,222†		6,115	48.7	783	72.9	80	65.4
Other Diseases 7,063 56.1 598 55.7 82 Infectious Diseases — — 901 83.9 6 Puerperal state — — 262 24.4 13 TOTAL 125,646 1,000 10,739* 1,000 1,222†		221	1.8	39	3.6	17	13,9
Infectious Diseases — — 901 83.9 6 Puerperal state — 262 24.4 13 TOTAL 125,646 1,000 10,739* 1,000 1,222†		7,063	56.1	598	55.7	82	67.1
Puerperal state — — 262 24,4 13 Total		1	1	106	83.9	9	4.9
125,646 1,000 10,739* 1,000 1,222†		1	1	262	24.4	13	10.6
	TOTAL	125,646	1,000	10,739*	1,000	1,222+	1,000

(See Table VIII.). *This figure represents illnesses suffered by 7,582 acute patients. ., 611 chronic patients. **

A further extension of these statistical investigations should yield interesting and helpful results.

- 18. **Conclusions.** The essential features of a model Domiciliary Medical Scheme can be variously stated but the following are probably its main requirements:—
 - (a) The Scheme should supply to public assistance patients a complete range of medical, dispensing and ancillary services not inferior to those provided under the National Health Insurance Acts.
 - (b) Its administration should be identified as little as possible with the Poor Law, and wherever feasible it should allow free choice of doctor by the patient.
 - (c) From the administrative standpoint it should be efficient, convenient, elastic and not unnecessarily expensive.
 - (d) It should work in close co-ordination with the statutory health and hospital services of the local authority, and should co-operate effectively with the voluntary organizations which are concerned with the nursing and medical treatment of the sick poor.
 - (e) It should offer a reasonable remuneration to those medical practitioners engaged in its service.
 - (f) Its medical records should be capable of throwing light upon the problems of sickness and invalidity in the community.

It would be premature to suggest that at the present stage of its development the Newcastle Scheme complies with all these requirements, but it is submitted that during the period covered by this report it has been steadily approximating to the ideal which was present in the minds of its originators.

> J. A. Charles, Medical Officer of Health.

Health Department,
Town Hall,
Newcastle upon Tyne,
October, 1936.