The anatomy of the brain. Containing its mechanism and physiology; together with some new discoveries and corrections of ancient and modern authors upon that subject to which is annex'd a particular account of animal functions and muscular motion ... / [Humphrey Ridley].

Contributors

Ridley, Humphrey, 1653-1708. Cowper, William, 1666-1709

Publication/Creation

London: Sam. Smith and Benj. Walford, 1695.

Persistent URL

https://wellcomecollection.org/works/z4kezdbc


License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org











V b

63087/3


The Library of the Wellcome Institute for the History of Medicine


OF
LONDON
DEPOSIT


Accession Number


Press Mark

RIDLEY, H.









The distance of the Brain Digmon Judicamas qui Imprimanas, Tanna Dayman, P.

Tractatum hunc cui titulus

The Anatomy of the BRAIN;

Dignum Judicamus qui Imprimatur,

Dat. ex Ædibus Collegii in Comiriis Cenfor. Sept. 7. 1694 Thomas Burwell, Præses.

Samuel Collins,
Fred. Slare,
William Dawes,
Tancred Robinson

In white

ANATOMY BRAIN.

SCAL SO

Containing its

Mechanism and Physiology;

Together with fome

New Discoveries and Corrections

Ancient and Modern Authors Upon that SUBJECT.

To which is annex'd a particular Account of


ANIMAL FUNCTIONS AND Muscular Motion.

The Whole illustrated with Elegant Sculptures after the life.

By H. RID LET, Coll. Med. Lond. Soc.

LONDON:

Printed for Sam. Smith and Benj. Walford, Printers to the Royal Society, at the Princes Arms in St. Paul's Church-yard, 1695.



Spectatissimo Doctissimoque Viro D. D. JOHANNI LAWSON

Collegii Regalis Medicorum London.

Presidi Dignissimo

SOCIIS,

Et inter eos speciatim

CENSORIBUS

Vel eo nomine Clarissimis

SAMUELI COLLINS, RICHARDO TORLESS, EDVARDO TYSON, MARTINO LISTER.

NECNON

D. D. Electoribus Meritissimis Omnibus & Singulis.

Tam præ Universali Exquisita sua eruditione, quam Artis Apollineæ Praxi fælicissima longe Celeberrimis

CÆTERIS Denique,

Egregiis Viris

Inclytissimæ bujus Societatis ascriptis

Paginas has eorum jussu in lucem prodeuntes,

Honoris & Obsequii Ergo

quam Humillime Offert,

H. R.

Section (Said office Page 1700) Collegii Pegalis Medicorum Londori, THE PRINCIPLE OF THE PARTY OF T Time over Delection of section for emilising

PREFACE

TO THE

Reader.

HAT Reason which, upon first thoughts, seemed of most force to disswade me from engaging my felf upon the Subject I have made choice of in these few following Sheets, (which was, its having been already undertaken by two fo eminent Persons, as the late Willis, and the present Vieussenius) upon second became the greatest motives to it. Seeing that even after the best Proofs they have either of them been able to give of Skill or

A 4

The PREFACE

or Industry upon this Subject, there hath yet escap'd undiscover'd both a great deal of the Materials which Nature is wont to furnish for the framing of Parts, and Contrivance too in ranging of them, in order to bring about that great design of making them all contribute their share to the conservation of the whole.

The cruth of this becoming still more evident whilst I became more conversant in Diffection, after some time, put me upon an endeavour, by a deeper Scrutiny, to discover something more than what as yet had come to light: and this I undertook so much the more vigorously, as by how much I reckon'd it more preferrable to contribute my Mite towards the perfecting of a Work already so happily be-

gun and successfully carried on, than to break the Ice only (the common fate of the sirst attempt) of another. With what success I have done it the Reader must be Judge.

Through the whole description of Parts I have offer'd nothing but Matter of Fact, and have taken all possible care to avoid being impos'd upon my felf, by making Experiments in proportion to my Doubts. Some of them have been upon Subjects in their natural, some in their morbid estate, some upon those of Untimely Death; and on those last fometimes whilst the natural Fluids remained in their proper Vessels, though after a preternatural manner occasion'd by Strangulation; sometimes when in the room thereof, other Bodies have been introduc'd

To the READER.

by Injection, as Tinged Wax and Mercury, the first of which by its consistence chiefly, the other by its permanent nature and colour, contribute mightily towards bringing to view the most minute ramifications of Vessels, and secretest recesses of Nature.

By this various disposition of the Subject it is that so great Dissiculties are overcome in search after Truth, many things appearing oftentimes very plain in one state, which either lay concealed, or seemed otherwise modified in any of the other.

The Figures were delineated by the hand of that Compleat Anatomist Mr. Comper the Surgeon, whose great Skill in Dissection renders that Talent so fortunate both to himself and his Friends: and how exactly that

that Work is performed, I submit to the Severest Censure of any who will be at the pains to compare any of the Cuts to the life.

What I have faid upon the Physiologia, in relation to Nutrition and Muscular Motion, depends on Microscopical Observations; and as to the Postulatum on which they both depend, though at first fight it may appear surprizing, yet I am confident it will become far less so to those who have been acquainted with what hath been faid of the Vascular Compages of Plants by Malpighius and Grew, and of several other Subjects by Lewenboeck.

And to conclude, I must confess I have been the better satisfied with it my self, since I met with some Passages in the Works

The PREFACE

Works of those learned Micrographists Dr. Power, and of Mr. Hooke, relating to this Subject, in which last, the medium made use of for solution of that famous Phanomenon of that Plants contraction at the first appulse of Touch from external Objects, as well as the manner of its acting, is the same with that made use of here as a Postulatum, upon which the whole of what is faid about Muscular Motion is built : Altho' at the same time I am sensible 'tis not so apply'd in that place by the aforefaid Author, whose opinion in reference to Muscular Motion (being the same with that of Dr. Mayow already taken notice of in the following Sheets) is exprefly otherwise in the account he gives of those natural Hygrometers the Beards of Wild Oats, of all the forts of Cranes Bills and

and Cats Guts, conformable to the manner of Nature's acting on which, in order to make them proper Indexes of the various Changes of Weather (viz., by wreathing and unwreathing) he supposes that to be of Muscular Motion.

I have quoted Authors, not out of oftentation, but both for their Truth and Errors, to the end that at the same time we may see it reasonable and convenient to read all they say, we may be render'd cautious how we believe; and to put us in mind, that as we find something done to our hands by those who have gone before, there is reason we should do something for those who are to come after.

THE

INTRODUCTION.

Owsoever the Controversie may Stand amongst Learned Men, about the Method and Order which Nature makes use of in the framing the different Parts of Animals, especially as to precedency of Time, some of them supposing a rudimentary delineation, or pre-existence of the whole, which, as the Ingenious Bruner hath rightly observed, must necessarily imply an actual existence of the whole Race of Mankind at once, either in the Testicle or Ovarium of Eve, according to the Learned Harvey, Malpighius, Swammardam, &c. or in that of Adam, according to Lewenhoeck, Dr. Garden, and Several others, and consequently must needs also infer an extinction of the same Progeny, as soon as the number of those humane Germens or Animalcles shall be exhausted; others a gradual formation of parts,

0136

The Introduction.

one after another, by an intestine motion begun and carried on from the time of coition, by the Subtle matter in the Cicatricula of the Egg: I see no reason to make my self a Party on either side at this time, seeing the fineness of structure and dignity of functions are sufficient to give preference to one above another, and to render it more worthy of a particular consideration. And this part I take to be the Brain, the delicacy of whose stru-Eture is such, that with no little resemblance to its divine Author, whilft it gives us the greatest and clearest discoveries of other things, lies most concealed it self.

And seeing all that Mystick Know-ledge, which in ancient times, in the eyes especially of the Vulgar, appeared meer Necromancy or Witchcraft, as well as all the Curious Discoveries of more modern Ages upon the whole subject of Nature, now going under the more familiar and proper term of Refined Sence, or Philosophy, hath been meerly owing to a more acurate knowledge of the parts and modification of Matter, I see not any more likely way of conquering the difficulties yet

The Introduction.

behind upon any particular subject, than the endeavouring after a further and more nice scrutiny into it by such means and experiments as serve to bring its most minute parts and texture under the test of Sence. which so assisted, doth the same office to the discerning faculty as good artificial Glasses do to it, bringing the Object and Judgment to such a nearness, that even the first Link of the Chain becomes discernable, and the mechanical proceedings of Nature so highly instructive to the Understanding, in its finding out and affiguing proper Causes to Effects much more obvious and intelligible.

I shall therefore treat this Noble Part after the aforesaid manner, with all the Justice I can, leaving those invisible, and almost divine things called Animal Spirits, to be treated of more at large, by those more illuminated Philosophers, who see best when their Eyes are shut, and content my self with making an inquiry into, and giving a description of, what so ever upon this Subject, by Dissection, shall offer it self as an Object of our

Senles.

THE

THE

ANATOMY

OFTHE

BRAIN.

CHAP. I.

Of the Anatomy of the Brain.

H E topmost part or Olla of the Cranium being removed, the first part of the Brain that comes in view is the

Dura Mater, which, with the subjacent Pia Mater, is accounted only an improper part of the Brain, strictly so called, however of great use in many respects to it.

Tis by Spigelius and other Anatomists reckon'd, and I think not undefervedly, the thickest and hardest Membrane of the whole Body, enclosing the whole Brain, properly so called, somewhat losely, sticking almost

most inseparably to the Basis of the Cranium, and to the top and sides, under the Coronal, Sagittal, and Lamdoeid Sutures, very fast by the Sinus's whose description will come in ano-

ther place.

In some places of the upper part of the Cranium, which on each fide of the Sagittal Suture or Vertex are called Offa Bregmatis, it adheres not to the Bone, not with francing the politive Opinion of Van Roonhuyse, in his Let-Roonh. ter to Du Foy, to the contrary, who P. 149. for that very reason would fain take away in a great measure the use of the Trepan and Trefoyne, and altogether the use of the Instrument called Decufforium, which skilful Surgeons do often make use of to make room for the discharge of subfided matter below the fractur'd place in many Accidents of the Brain. To boul a mi

Tis very discernably double, as Columbus and several others formerly, col. p. 348 and Vieussenius lately, have observed, having very strong and large Fibres on the inside, but very small, and hardly visible, on that side next the Skull; as appeared to me, after having sirst let it lye a little time in boiling or at least very scalding Water.

But

But as to the distribution of the double fort of Fibres on each side this Membrane, I could not by any means find them agreeing with the description Vieusenius hath given of them, as running in an oblique semicircular manner, externally from before backwards, and in the same figure internally from behind forwards; but far otherwife, on the infide, where they are very strong, they feem manifestly to have three originals from the top part of the Processus Falcatus, before, behind, and in its middle; those before running in a curved manner backwards, half the length, and a great width of the Dura Mater, and those behind running after the same manner forwardly with this difference, that a great number of them bend foon after their rife from that process in a kind of a femilunary way to it again a little on this fide the rife of the middle Series of Fibres, others of them making a bigger arch after having stretched themselves wider upon the Dura Mater, bend back again to, and terminate in the Falia little beyond the rife of the aforesaid middle Series of Fibres.

Those from the middle part of the Falx run backwardly, but less curved than the rest, terminating as the Fibres which arise backwardly do, at some distance from the Process in the inward Superficies of the Dura Mater.

As to those belonging to the exernal side or second Lamina of the Dura Mater, they are extream small and obscure, running from behind forwards.

Besides these, there are no less remarkable ones belonging to the Falx it self, of two sorts of Orders, the one running streight about half the length of it, on its upper part, from before backwards, the other transverse, from the inseriour or fifth Sinus to the superiour or third, on the hinder part of the Process, and are most conspicuous there, as the other are towards its foremost part.

As to the Use of these Fibres, it may be remembred that this Membrane consists of two Lamina's, between which the Veins which reduce the Bloud from the Arteries, which furnish the whole Brain with it, run sol some space after the manner of the Ureters in the Bladder, in large

Trunks,

Trunks, before they enter the Sinus; fo that the Fibrous Constitution of this Membrane here, where the Bloodvessels are largest (together with the curved entrance of them into the Sinus, especially in an erect position of the Body) do the office of Valves, support the weight, and promote the afcent of the Blood. But that which is most considerable, is this, That if the inward Lamina of this part, which makes the inferiour and lateral part of the Sinus, was not in some meafure furnish'd with additional strength on this fide suitable to that which it hath on the other, by reason of its cohesion to the Skull, the Blood which is continually running through it with no small rapidity, especially in great plenitude of the Vessels or preternatural Ebulitions, would frequently burst out, or at least cause fuch distentions as could not but be very injurious to a part so very exquifitely fensible; yet notwithstanding, tho' Nature seems plainly to have made a double provision against fuch Accidents, by the transverse Ligaments within the Sinus, and thele ftrong and numerous Fibres without, I have rarely open'd any strangled Body,

Body, where some such Rupture, or at least Distention, hath not hapned.

This Membrane hath plenty of Nerves from the foremost Branch of the fifth Pair, and is thereby made very fensible, so that from any molestation given it by the ill Crass or undue motion of the Blood, it becomes accordingly affected. And as the various distribution of Fibres before described serve in a natural estate to give a kind of springiness to the Veffels, whose Coats are extended by the Blood as they run between the Laminæ of this Membrane, to the end the same may be the more readily circulated through them; fo in a preternatural estate, no doubt, they are subject to Spasms, which may retard the course of the Blood in such fort, that in some kind of violent Headachs, where the Membrane is affected through overfulness of Blood, and particularly in those which are wont to proceed from Vapours (fo called) or Convulsive Motions of Nervous parts, we often observe a fixed ruddiness in the Face, attended with a kind of stiffness and foreness in the Eyes, proceeding doubtless from a flagnation in some measure of the

Humours in those parts, through the too slow passage of them into the reductory Vessels or Sinus's. And to this preternatural assection of the Sinus's may certainly many other ill Symptoms of the Brain be imputed, and not to any irregular Systole and Diastole of the Membrane it self, occasion'd through any convulsive or paralytical state thereof, as that curious Speculatist Dr. Mayow hath affirmed, mayow, med, seeing not any living Dissection Tr. 4. p.49 hath ever been found to give Authority to any such Hypothesis.

The First Process of the Dura Mater.

It hath two Processes, the first of which arises from that part of the Os Ethmoeides, called Crista Galli, and is extended from thence backwards, as far as the concourse of the four greater Sinus's, commonly called Torcular Herophili, in the figure of a Sicle, whence it hath that denomination of Falx, and by reason of the strict connexion it hath by certain Membranous Fibres with the Cranium in those places which are immediately under the Sutures, and with the Brain it felf, by the intervention of the Pia Mater, (to which it is joyned both by the intervention of large Blood-veffels, propagated thence to the longitudinal B 4 and

and lateral Sinus's, and certain carnous Adnascencies, as it descends down betwixt the two Hemispheres of the Brain, and afterwards at its approach to the back of the Corpus Callofum, (over which that Membrane is loofely expanded) both by continuity of its Membranous Substance and Rami-, fications of Blood-veffels, terminating in the fifth Sinus, at the bottom of the Process. so that in a Diseased Brain I once law it drawn up the length of an Inch from the faid Corpus Callefum, in the exact form of a membranous thin Production, continued to the fifth Sinus running at the bottom of this Process,) it keeps the Brain fuspended in such a natural conformation, that it needs not, to that internal part by the Ancients call'd Fornix, nor that by Vieussenius of late fubstituted in the room of it, call'd Corpus Callosum, for its support.

Another Use it hath is, partly to desend the Cerebellum from Compression, to which, by its connexion with the Galli Crista, it doth not a little contribute, but chiefly the two Hemispheres of the Brain from the like Injury from each other, upon its various position in Sleep or otherwise;

and

and therefore is wanting in many other Creatures, as Calves, Sheep, &c. which not only Sleep less, but for the most part in a less injurious posture.

The Second Process of the Dura Mater.

The fecond is that which arising fo forwardly as from the hindermost Process of the Wedglike Bone, which composes the back and uppermost part only of the Sella Equina; it passes up betwixt the Cerebrum and Cerebellum, all the way adhering to the internal Eminencies of the Offa Petrosa to the lateral Sinus's, by which means not only the Cerebellum immediately, as is commonly observed, but consequently all the Parts from the beginning of the fourth Sinus, or the Glandula Pinealis, to the last Foramen of the Skull, (viz.) the Caudex Medullaris, with its Appendices the Nates and Testes, (which being placed upon the upper part of the Medulla Oblongata, make a fort of an Istbmus betwixt the Cerebrum and Cerebellum) together with the Nerves proceeding out of it, are defended from the injurious pressure of the hinder Limbs of the Brain.

lujury from each other, upon its vari-

AHAD m sleep or otherwife

CHAP. II.

Of the Pia Mater.

H E Second Integument of the Brain, commonly called Pia or Tenuis Mater, by Galen and many others, Choroeides, from its likeness in substance and ramification of Bloodveffels to that Membrane of the Secondines call'd Chorion, with much more reason than Vesalius, on behalf of the Plexus Choroeides it self advances against it; was by all the Ancients look'd upon as its only other Integument, being a very thin and pellucid Membrane, co-extended with the Brain it felf, not only in its outward but inward structure too, as likewise through all its Plicatures, Intersfices, and Cavities, even over the Corpus Callosum it self, the' loosely, as hath been already observ'd, notwithstanding the great Vefalius af- Vefal. firms the contrary: Which Membrane P. 778. also a chance cut in pareing the toppart of the Brain down to the lateral Ventricles with a Razor, in a Body I lately had, gave me an opportunity

of

of showing as fair in those Ventricles as the largest Membrane of the whole Body, to several who stood by, not-withstanding Molinetti, who laughs Mol. p.78. at all that pretend to have found any

fuch thing, affirms the contrary. But this is to be enquired for either in recent Bodies, or fuch who have before death been, thro' some Diseafes, fill'd with extravalated Serum, as Dropfies, Stoppage of Vrine, some fort of Apoplexies, or the like: That way which in want of the other opportunities discovers it best, is the separating the Septum Lucidum near to its rife, which is just from the Fornix, where it arises from its two Roots, near to which place the Medulla of the Brain begins to advance into the Corpora Striata; for from thence for above half way of its passage backward toward the hinder limbs of the Brain, it continues hollow, and, I am apt to think, is but a Duplicature of this part, tho' it may be somewhat medullary, and therefore, by reason of its transparency, hath the Name of Septum Lucidum.

This Opinion of the Ancients, of its being the only other, and that a fingle Integument of the Brain, was equally

equally receiv'd for Truth by the late two learned and curious Anatomists Willis and Vieussenius, together with all the other modern Writers, except Bidloo and Bohn, both which affirm, Bid. Tab. 8 they have found another diffinct Bohn p. 333 membranous Integument of the Brain coming betwixt the other outward Dura, and inward Pia Mater, the one three hours, the other fifteen days after death; and by them both reckoned the original of the second proper Integument of the Spinal Marrow which Tulpius first discovered, Tulp.cent. 1 and Vieussenius supposes to be a Dupli- Vieussen. cature of the Pia Mater in that part P. 143. only.

Now, that there was a middle Membrane in some parts of the Brain, and particularly at the Basis of the Cerebellum, from whence it's continued down to the Spinal Marrow, constituting the second proper Integument of that part as afore-mentioned, I had long since observed; but whether it be another absolute distinct Membrane from that other subjacent one, by the aforesaid Authors properly named the Pia Mater, and common to the Spinal Marrow with the Brain it self, like as is this other

ic-

fecond middle one too, or only one and the fame Membrane double, as confisting of two Lamina's, may well be doubted of.

Wherefore, for fatisfaction concerning this difficulty, I have lately made the strictest enquiry possible, and that in a subject most likely to afford a decision in such a Controversie, and this was an Human Brain extreamly hydropical, where there was no Cavity or Interstice, without abundance of Water extravalated, infomuch that where ever, according to the natural construction of Parts, there was any larger than ordinary duplicature of this Membrane, as there are at the end of the Calamus Scriptorius, betwixt the superincumbent Cerebellum and Medulla Spinalis, in the Isthmus or space betwixt the Cerebrum and Cerebellum, upon the Processes called Nates and Testes, in the depressed part also of the Brain, between the beginning of the Annular Process, and the first appearance or coming out of the O!factory Nerves, by Vefalius taken notice of and called a Process of the Vesal.p.794 Pia Mater, there was found a great deal of Water distending this Duplicature much beyond its natural li-

mits; fo that by way of confequence, if these Cavities were only Interstices of two different Membranes distinctly investing the Brain, and not a Duplicature only of one and the same, the Water would then probably have infinuated it felf betwixt them, and made them to have appear'd far different from what they did, agreeable to what it hath often been found to do in some Dropfies of the Belly, where the Water hath been found to to have divided or parted the double Membrane of that Region call'd Peritoneum, as to have render'd it capable of containing the quantity of fifteen Gallons of Water, and upon a difcharge of the same after death, by cutting the external Lamina of that Membrane, the other inward one being yet (unknown to the Dissector) left whole, to have imposed upon the Spectators, and those very sagacious ones, fo as that at first fight, till after having recollected themselves, and 70b Meetr. divided the other second Lamina too, Obf. 52. they thought the Bowels of this part to have been wanting; but contrary to this Event, in this Subject I found this Membrane entire, and free from any divultion throughout its whole CIT-

circumference, excepting the places afore taken notice of. However, suppoling the like conformation here in this with the Membranes of the other parts, I attempted to divide it, and did to fuccessfully in many parts of it. but most readily in the beginning of the superficial Plicatures of the cortical part of the Brain, where there are naturally small Interstices, betwint which many of the Blood-veffels creep into and immerge themselves in the cortical and medullary parts thereof: So that I think there cannot remain any further scruple of its being only a double, and not two diffinet Membranes of the Brain.

Bidleo very truly observes this sirst or middle Membrane, by him so called, by me only the sirst, or one Lamina of a double Membrane, to be thinner than the Dura Mater above it, and thicker than the other Membrane or Lamina under it; which last most properly it is that infinuates it self through all the close Plicatures of the Brain, and that, as by frequent inspection I have often observed, not in a continuous, but rather retiform contexture, and so, by such as love hard words, or terms of Art, may be call'd

called after the same name of that Membrane investing the crystalline Humour of the Eye, Arachnoeides.

The Advantages accrueing to the whole through such a disposition of this part, as hath already been observed, are very considerable, inasmuch as that thereby first of all it becomes not only an Integument of inclosure, on behalf of the Brain, and the Bloodvessels belonging to it in general, but of expansion for Strength too, where the peculiar structure of Parts, in such places as were before mentioned, require it.

As to the first, the Brain is not only kept more warm, close, and compact, and better desended on its depending part from the asperity of the Bone it lies upon, but the Vessels hereby more strongly supported, and it self secured from being broken or torn, whilst between its duplicature they climb up into the Brain, whose delicate tender Fibres must otherwise of necessity have suffer'd violence by the largeness and pulsation of the Arteries, together with the weight of them,

and the other reductory Vessels, from

which the Sinus's meet them.

Nextly,

Nextly, as it is an Integument of Expansion in the places before mention'd, that tender small part the Infundibulum, where it quits the Brain, in order to its passage into the Glandula Pituitaria, by the circumtension of this outward Lamina, is fortified upon any violent Accident from difruption, and the Brain and Medulla Oblongata, in those places where they are only loofely contiguous, are better preserved in their natural due connexion; all which Advantages, inalmuch as they may more reasonably be ascribed to one double Membrane than two fingle ones, tho' of the like strength when joyned fast together, may not unreasonably be thought to argue for the duplicature of this Membrane exclusively, to the introduction of a third or new one.

Laftly, as to what concerns the Glandes and Plexus's which Dr. Will. Will. p. 26. lis affirms to be scatter'd all over this col. 1. Membrane; as to the former, I could never fee them, but I have feen the external Superficies of the cortical part of the Brain, in strangled Bodies, appear glandulous very plainly, through this transparent Integument, which upon bare inspection,

The Anatomy of the Brain. 18

> without further enquiry, might eafily impose upon the less cautious Spectator.

> As to the latter, the Plexus's, and distribution of Blood-vessels from them, after a separation of the serous gross part of the Blood in the aforementioned supposed Glandules, (according to that learned person's conjecture) into the substance of the Brain, in order to produce the finer Animal Spirits; I cannot but look upon it altogether conjectural, till fuch time as not only the Glandes, but their excretory Ducts also, together with the Emunctories where the fuppoled excrementitious Juice is eliminated, (lymphatick or reductory Glandes (if they could be found) never having been by Nature defigned to any such use) be first discovered.

This Membrane hath Blood-veffels

of two forts.

Of the first are these properly belonging to the Brain it felf, which, as it hath already been observ'd, it doth as it were conduct through its Duplicature, in their passage allowing them thereby the opportunity of growing extreamly fine, after many serpentine twinings towards their capillary Ex-

tremities.

Blood-vesfels of the Pia Mater. tremities, before they are protended Bid. Tab. into the Brain it felf, and those are 8. f. 5. l-M chiefly spread all-along upon the under or second Lamina of this Mem- Ib. l. G. brane.

The second are those which belong to this part it self, for its own nourishment, and these I sound upon diligent inspection, whilst I separated its second Lamina spread plentifully upon the inside of the outermost or first Lamina, and both these you will find very well delineated in the

places quoted in Bidloo.

This Duplicature is also very plainly communicated to all the Nerves both within and without the Cranium, making by its outward Lamina a fecond Integument under the first from the Dura Mater to the whole Fasciculus of Nerves, and a third by its inward Lamina, which yields an involucram or covering to each fingle Fibrilla, which collectively make up the whole Nervous Body it felf, thro' the admirable fineness of which Membrane invefting those medullary Fibrils, altogether insensible of themfelves, it happens there is such a nimble confent betwixt part and part, and betwixt all and the Brain it felf.

C 2 CHAP.

CHAP. III.

Of the Vessels belonging to the Brain in general.

part in common with the rest of the Body, though in reality but one continued Canal variously modified, yet, through the diversity of Fluids they contain, go commonly under the denomination of Arteries, Veins, Sinus's, and Lymphædusts, and not without good reason, perhaps, the Nerves may be in some sence of the same kind too.

The two first of these may, with relation to their different distribution, be deservedly consider'd in a two-sold respect, either as they belong to the first Integument of the Brain, or the Brain, properly so called,

it felf.

The Arteries therefore belonging to this part called Dura Mater, or first Integument, are three fair Branches on each side.

The

The first and foremost of which are fent out from the Carotid Artery, whilst it remains in the fourth Fig. 2. hh. hole of the Cranium, and are propagated chiefly through the foremost part of the bottom of the Dura Mater, as in the Figure delineated, but greatly mistaken by Dr. Willis p. 2. perhaps taking it upon trust from col. 2. Wepfer, equally with himself there- wepf.p.105 in mistaken; who describes it for a small branch of the Carotid Artery, that runs betwixt the two first Lobes of the Brain, which instead of coming out of the Bone of the Forehead, as he would have it, goes into it without lending any branches to this Membrane at all, being truly delineated and described by the aforemention'd accurate Vieussenius. Vieus. Tab.

And that this Artery was not on- 17. dd,bb. ly mistaken by, but unknown to the par. 4. aforesaid Wepfer, is plain, seeing he Wipf. p. 101 fays, that from the very styliform Process, where the Carotid Artery does indeed enter the long Canal, to the place where it perforates the Dura Mater to enter the Brain. there is not one Branch fent out from it; which Error, by injecting with Wax, which keeps longer in, and .

and shews the Vessels much better than small tinged Liquors, had very

easily been avoided.

Fig. 2. ii. The second Branch of Arteries ascend into the Dura Mater by the fixth hole of the Cranium, together with a Branch of the internal Jugular Vein, and are dispersed laterally all over the fore-part of this Membrane, as far as the very Sinus Longitudinalis, (which nevertheless it enters not, as there will be occafion to take notice of hereafter) as in the Figure delineated.

The third Branch of Arteries climb into the Dura Mater by the eighth hole of the Calvaria, together with a small reductory Branch of the Vertebral Vein, where the F16.2. kk. lateral Sinus's enter the internal

Jugular (which occasion'd the Ingenious Highmore erroneously to believe it enter'd the very lateral Si. Highmore, nus it felf) and the eighth pair of par. 1. Nerves pass out of the Cranium, which passage of this Artery is not hitherto described by any that I know of; neither have I ever feen it figured, but in Vieusenius's first vieus.tab. " Cut, and there but very faintly.

It arises from the external Bran-Vieuss. tabe ches of the Vertebral Artery, accor8.f. 1. c. Barthol.
ding to Vieussenius, but Bartholine p. 431.
makes it to be a slip of the Carotid Par. ult.
Artery, calling it the lesser Branch thereof; wherein he is mistaken.

The Veins As to the Veins, Riolane, and af-Riol.p.252 of the Dura ter him Willis, seems to say this Will. p. 2. Mater. Membrane hath none; for the the col. 2.

latter hath this obscure expression of par. 6.
them, Tam crebris Venarum propaginibus quam Arteriarum nusquam consitu est; speaking of the Crassa Meninx, by which we might guess he win. p.22.
thought it had some, yet in another col. 1.
place he plainly substitutes the Sinus's for the reductory Vessels, as
well on behalf of this Membrane as
the Brain it self; as appears plain
enough in the Page noted.

Vieusenius indeed allows Veins Vieus.p.32
to this part, and says, they all-along par. 3.
accompany the Arteries, and afterward terminate, according to Vestin-Vesting.
gus, in the internal Jugular; yet in Vieus. p. 4.
another place he says, some of the par. 2.
Venal Branches discharge the Blood into the Sinus Longitudinalis. Which last is a flat contradiction to the place foregoing, inasmuch as in that he says, they accompany the Arteries

C4 all

all-along after the same manner of distribution or ramification; which, if so, who sees not that they must needs grow capillary towards the Sinus, and consequently be uncapable of reducing the Blood into them, all reductory Vessels being always capillary in the place from which, and not to which, they bring that

which they contain.

Now therefore, neither what the one nor the other fays can possibly be true; for, as to the former the learned Dr. Willis, if his Affertion was good, it must of necessity follow, that all the Arteries dispersed thro' this Membrane must terminate in some of the Sinus's, otherwise there will want a reductory Veffel: the first of which is contrary to ocular demonstration, the last to common reason. To describe stores

As to Vieusenius the latter, befides what hath been already faid against him, if what he fays in the place aforecited be true, that the Veins of the Dura Mater run concomitantly along with the Arteries, then they must of necessity answer the ends of other Veins throughout the whole Body, in reducing the Blood

Blood adduced by the Arteries, unless the Arteries they accompany discharge their Blood into the Sinusses, (which, as hereafter shall be shown, they plainly do not) for otherwise, seeing they both grow capillary in their ascent from the Basis of the Cranium, they must necessarily be both adductory Vessels, than which, by the Laws of Circulation, there can be no greater an Absurdity.

Wepfer not knowing of these Veins, was forced to think, and consequently to affirm, That the Arteries leave the Dura Mater in their extremities, and terminate in the Pia Mater, and so have their Blood reduced by the Veins there; but this is evidently not so to the Eye of any who heedfully separates this Membrane from the

other.

Before therefore I proceed to the description of the Blood-vessels belonging to the Brain it self, which by the exactness of method I ought to do, I hope it may be pardonable, if I make a short enquiry after the unaccustom'd distribution of Blood-vessels Nature hath furnish'd the Brain in general with, and the Reasons of its procedure therein.

The Truth then concerning this affair, is, That contrary to what hath hitherto been observed, the Bloodvessels belonging to this part in general, as hath already been observed, are of two forts, the one belonging to the Brain it self, the other to its out-

most Integuments.

Now, as to the first, 'tis observable, that the Veins enter not the Brain, nor run concomitantly, like as in other parts of the Body, with the Arteries, (the carotid entring at the fourth hole in the Basis of the Skull, and the internal Jugular at the eighth; the Vertebral Artery at the last and largest hole of the Skull, and the Vertebral Vein at the ninth (which Vieussenius mistakenly calls Vieussen. the tenth) thro' which it runs into par. 3. the internal Jugular, at that Veins entrance into the round hole at the bottom of the Skull, under the Styliform Process, where the Sinus Laterahis meets it) where after having advanc'd into certain venous productions called Sinus's, they descend from thence in large Trunks, growing capillary all-along in their passage till they meet the Extremities of the Arteries, and are indeed no other than

meer

meer Branches of the Sinus's, and consequently I look upon the Sinus's themselves no other than large Veins.

The common reason all modern Authors give for this different diffribution of Blood-vessels belonging to the Brain, from the other parts of the Body, is, that it may receive an equal warmth at the top as at the bottom, as being thereby very much affifted in the production of Animal Spirits in an equal proportion all over; and that it is so may very well be granted: but, that Nature had yet another provident Intention, will be as evident, if we confider, that if the Veins had ascended with the Arteries thro' the holes in the bortom of the Cranium, upon all great Ebulitions of the Blood, the pullation of the Arteries would in that Stricture of the Veffels made by the Bone, of necessity hinder the freedom of its return by the Veins, and confequently oceasion a stagnation of Blood through the whole Brain, to the utter subversion of all its faculties, nothing being more certain, than that upon any confiderable abatement of circulation there presently happens by

by way of restagnation, a secession of the watery and thin from the more gross and red part of the Blood.

The other way of the Veins entring the Brain (viz. those appertaining to its outward Integument, one at the fixth hole of the Basis of the Cranium, the other at the eighth, as aforelaid) is, their ascent with the Arteries after a quite different manner from the former, even to their capillary Extremities; a manifest indication that they serve for the reduction of fo much Blood from the Dura Mater as the aforesaid fort of Vessels, the Arteries, have brought thither; and although by reason of their smallness Nature seems not to have been so sollicitous in avoiding the Inconvenience supposed to have follow'd, upon the Artery's entring the same hole with the Veins, taken notice of in the preceding Cafe, where they are very large, and consequently the Effect might prove much more injurious, yet Nature hath not been wanting in providing a Remedy against it; as will plainly appear in the following Pages.

gad adiable oreds her From

From this manner of their entring the Brain at the same inlet of the Skull with the Arteries, may, for ought I know, be very rationally accounted for that violent troublesome Noise which many, in Distempers arifing from the turgescency of the Blood, caufing a preternatural beating of the Arteries, do fo much complain of; a Symptom happening from the Stricture before mention'd which the unyielding circumference of the Bone occasions upon the different Blood-vessels entring at one and the same Foramen, to which effect also the nearness of the Os Petrofum, through which the Hearing Nerves do país to this hole, which is in that part of the Wedglike Bone that joyns to, or is conterminous with it, does not a little contribute.

To the same cause, in some measure doubtless, may be ascribed the frequent Headachs happening in Feavers, the Artery then so swelling and compressing the Vein against the edges of the Bone, that the Blood cannot be returned back through it in a due proportion, and consequently by its stagnation the Membrane becomes instanced and painful.

So

So that conformable to what hath already been taken notice of concerning the wife contrivance of Nature, in ordering the different distribution of the Blood-vessels, so as to avoid the Inconveniencies which might accrew to the Brain by compression of the reductory Vessels, occasion'd through their entrance at one and the same hole with the Arteries; it feems very much worth our observing, that befides the Veins of the Dura Mater, which enter the Cranium together with the Arteries, as hath before been mention'd, there are also several others belonging to this Membrane, having their rife at, and their descent after a very remarkable manner, from a Veinhereafter to be describ'd on each side of the Longitudinal Sinus, as you may fee in dd,nn,&c. grow capillary in their descent down ner to the other; and these do visibly inosculate with some of the Extremities of the aforesaid capillary Arteries, after the same manner as those larger Veins belonging to the Pia Mater do with the Arteries be-

belonging to the Brain and it, by which means it so falls out, that a considerable part of that Blood brought up by the Meninx Arteries, is carried back by these Veins, to the end that, especially in all preternatural swelling of the Blood, the inconvenience of Compression and all its ill consequences happening, by reason of an overfulness of these Vessels, may be in a great measure avoided.

sales a more de normana

read on each indesection ongs with a bright and sent may fee in the highest and confequently must grove capillary in their defociat down

oly inolegiate with long or the Ex-

activate uniprobe arreries

Trom the dier a quite contrary man! CHAP.

CHAP. IV.

Of the Veins belonging to the Brain it self.

A FTER this short digression, by order of Method, the Blood-vessels belonging properly to the Brain it self, fall under consideration.

The curious Anatomist Malpighius, Malp. de in his Letter to Fracassatus, says, they par. 2. bear a third proportion to those of De Cort. the whole Body; and for what rea- cereb. p 81 par. 2. son, seeing the part it self bears not the same proportion to the whole, it is so, it will be worth our while to enquire hereaster.

These are either Arteries or Veins. The former go under the name of

Carotid and Vertebral.

The first of which, after a curved passage (which is very well expressed in a Fig. of Dr. Wills) from the place Willis p. 29. where it begins to enter the Basis of Fig. 1. the Cranium (which is from the Styliform Process of the Os Petrosum) to the place where, on the inside, they pass through the Dura Mater, and ascend into the Brain, (which is at the fore-

foremost internal Process of the Os Cuneiforme) there is very near an inch and an half distance. I say, after this crooked passage into the Brain, they are propagated quite through its substance, having first divested themfelves of that thick Coat borrowed of the Dura Mater during their stay in the passage aforementioned; but not without the mediation or intervention of the Pia Mater, which Mem-. brane all the Branches of the aforesaid. as well as the Vertebral Artery, more or less first prop themselves upon before they enter on and disperse themfelves' through the substance of the Brain it felf, and is very finely expresfed in a Cut of Placentinus, at the end Spig.p. 179 of Spigelius; insomuch that Molinetti Mol. p. 77. (with whom also agrees Marchetti)p. 191. looks upon it as only a production par. s. of those numerous Vessels; whereas all those little ramifications both of the Carotid and Vertebral Arteries, viz. those from the carotid Artery, which as foon as it gets through the Dura Mater, and parts with its bor- Vieuffen. rowed Coat, are fent to the a Infun- p.35.par.r. Merves, together with those other e. of the Vertebral Artery which accom- . 16. gg.

pany

pany the d third, e fourth, f fifth, & fixth, d Vieuffen. h seventh, i eighth, k ninth, and I tenth P.35. par. I. pairs of Nerves, inafmuch as they en- 1 Tab. 17. P ter not the Brain it felf, are altogether P.35.par. 1. exempt from that Membrane; any of Tab. 17. which now-mention'd Blood-veffelsh h. p. 35. you either find delineated in Vieuse-h Tab 4. hb. nius's 17th Table, or mention'd in 'Tab. 17. some other place of his Book, by Fig. 2. hh. those Directions here placed in the 16. Fig. 2. margin; all which, tho' existent in hard. 4. Nature, are nevertheless there painted too stiff and formal (I am afraid by guess) inasmuch as that without an injection of Mercury (except those two which belong to the Olfactory and Optick Nerves) they do rarely come to fight in any form at all, Wax being over gross a body to enter fuch minute Vessels as those are; whereas by an injection with Mercury I find scarce any Nerves but what hath fome fuch small ramifications of Blood-veffels in them.

To go about to describe distinctly the whole ramification of Arteries through this part, which as was before noted, is here more remarkable for number and fize than in any other part of the Body, would not only be to do what in a great measure hath

been

been already done by Vieusenius, in his fixth Chapter, but feem to have also in it much more of oftentation than use.

I shall therefore only take notice of such propagations of them, as are either remarkable for magnitude, some curiosity of Structure, or useful

design of Nature.

And of this fort may well be efteemed the Vertebral Artery, next after the Carotid, which hath already been described, as entering the Brain at the last and largest Foramen of the Skull, contrary to what Dr. Willis, Willis, p.29. and before him Wepfer, affirms, col. 1. coming thither on each fide out of par. 2. the hole in the transverse Process of parx. ibid. the first Vertebra of the Neck, after Wepf.p.112. a very remarkable curved manner, as Low. Tab. 4.

Fig. 1. EE you fee in the Figure, (and by no means like to the delineation and description given by Dr. Lower and Dr. Willis,) ascending laterally upon the Medulla Oblongata as far as the beginning of the Processus Annularis, where they meet together in one fingle Trunk continuing fo the length thereof, by Vieusenius call'd vieusen. Arteria Cervicalis, after which they Tab. 4. either send forth two Branches, or receive two from the carotid Artery,by

D 2

means

means whereof there is a communication betwixt these two large Bloodvessels, and that of great use and benesit to the Brain, for by this means it happens, that if even three of the sour great Arteries which surnish this part with Blood, were totally obstructed, there would yet be a way lest for a competent supply from the other unobstructed fourth. These I call the Communicant branches, very ill painted in Bidloo's ninth Table, but very well in Vieussenius's fourth; as may plain-Vieussen.

Fig. 1. 4d ly appear here in the Figure taken tab. 4. b b.

exactly from Nature it self.

The structure and smallness of these Arteries seem to suggest two, yet surther, provident Intentions of Nature.

The first is the same it hath expressed in several other places, as in the ascent of the Blood by the Carotid Arteries, both which enter the Brain in a crooked line, the first at the sourth hole of the Basis of the Skull, the second from the hole in the transverse process of the first Vertebra of the Neck, after the manner already in both places described. So in the like manner here, by the narrowness of these Branches, the Blood is in a great measure retarded in its motion

motion to the carotid Artery, and by consequence to the Brain it self, which, for Reasons hereaster to be given in describing the Sinus's, would otherwise be in great danger of being over-flowed with extravasated and restagnant Blood.

The second is, a forcing the Blood more plentifully into the Spinal Artery, with which, tho' through the conical structure of the Arteries in common it cannot be altogether unfurnish'd, yet by its perfectly-reflexed position, would have it very scantily, were it not that by reason of the narrowness of the aforesaid Communicant-branches betwixt the two great Arteries, the Blood was driven back in a fort of a retrograde motion.

'Tis true, there is a conformation of Arteries something like this, tho' not altogether in the mammary and epigastrick Branches; but 'tis worth noting, that in both these places the main Artery from which these Branches spring is much more taper or conical, and the succeeding exporting Vessels far less both in number and size than those of the carotid Artery here, whose foremost and hinder lateral ramifications between the

D 3

Ibid. p. c

Lobes

Lobes of the Brain, bear an overproportion to the Trunks from whence they come, and confequently must, according to the aforefaid observation of Malpighius, in his Letter to Fracassatus, receive the blood brought thither far more freely and

plentifully.

Besides, the Cervical Artery here is fo far from being Conical, that being made up of two vertebral Arteries joyning together, it is much wider than either of them fingle, as appears plain-Fig. 1. g. ly in the Figure, and consequently would have carried away the Blood forwardly from the Spinal Artery more freely, had not Nature order'd the Structure of Vessels after another manner here than it does in other parts of the Body, where there is not the same necessity of contrivance.

One more Branch I take leave to mention only upon the score of its never hitherto having been taken notice of by any, and that's a small Artery attended with a Vein paffing through the lateral part of the Os Cuneiforme, (which constitutes the back part of the Orbite of the Eye, just under a very little Process of that Bone, (which either by reason of its

fize hath escaped being seen, or inconsiderable use, was never before, as far as I know, thought worth the mentioning;) and this, upon raising the fore Lobes of the Brain, offers it felf to the Eye of any heedful Obferver.

CHAP. V.

Of the Sinus's belonging to the Brain.

Third fort of Vessels offer themselves next to our consideration, under the general name of Sinus's.

These formerly were reckon'd only four, to which Vefalius added a fifth Vefal. p. 758 at the bottom of the Falx, by him Fig. 3 F. only call'd a Vein, which tho' frequently found, yet in some Subjects is wanting. Bourdon mentions two Bourd. more at the bottom of each fide the par. 2. fecond Process of the Dura Mater, D 4 under

under the lateral ones, which I never faw but once, and I am apt to think with Vieussenius, are most commonly

wanting.

Vieussenius describes four more, Vieus. which I find long before taken notice P. 6. par. 5. of, and exactly describ'd by Falloppius, p. 114. and after him, tho' but rudely, by Vid. Vid. that laborious Collector Vidus Vi- cap. 10. dius.

I think I can shew one more, but be their number what it will, I judge it reasonable to look upon them no other than Veins, whether we confider them in respect to either Office or Structure. All the business is, to confider and shew for what end they appear as fuch large Channels into which all the Veins of the Brain, like fo many small Rivulets after an unufual manner do empty themselves; and that I will endeavour to do after having first shown their several respective situations.

The first two are called Laterales, F164. BB. which run within a strong duplicature of the hinder Process of the Dura Mater, down upon the Os Occipitale over the Cerebellum, till in their further defcent, after a tortuous manner, upon the lower production of the Offa Pe-

trola

Fig.2.GC. trosa they wind under them in order to their passage out of the Cranium at the eighth hole, common to the

third Branch of Arteries belonging to the Dura Mater, and the internal

Jugular coming in, which is through two round bony Cells in the Os Petrofa, just under the Styloeid Processes into the internal Jugular Vein, into which, together with the Vertebral, all the rest of the Veins and Sinus's belonging to the Brain discharge the restuent Blood.

FIG. 4. AA, &c. The next is called the third or longitudinal one, from its rife at the bony Process called *Crista Galli*, and progress the whole length of the Brain to the hinder and somewhat declining part of the occipital Bone, where it seems to be cleft into the two lateral ones.

Into this third Sinus not only the internal Veins of the Brain it self are inserted, but also some of those belonging to its outward Integuments, which Falloppius sirst, one of the Lu-Fallop. minaries of Anatomy, observed; and tom 1. p.82. par. 3 after him Vieusenius, which are by Vieus. p.19. Wepfer mistakenly taken for Arteries, par. 2. Wepfer mistakenly taken for Arteries, Wepf. p. 42. who nevertheless, for ought I know, par 2.

may

may be in the right, in affigning the overcloseness of the Pores of the Cranium (by what Accident soever happening) thro' which the refluent Blood is transmitted to the Sinus. for a frequent cause of inveterate obstinate Headachs.

The fourth, which from its situa-Fig. 4. C. tion may not improperly be called the Internal Sinus, comes from the under part of the falcated Process, at that point where it becomes continuous to the second Process of the

Ibid. II. Dura Mater, and a large double Vein belonging to the Plexus Chorocides.

together with the fifth Sinus, (when Toid K. there is one) enters it at an Interstice made between the end of the Corpus Callosum, the Nates, Testes and Cere. bellum, from whence having first passed over the Gerebellum, it at last arrives with the other three at that place of union, which from its Author hath ever fince retain'd the Name of Tor-

cular Herophili.

The four others of Falloppius and Vidus Vidius, or Vieussenius, by this last called Superiores and Inferiores, the dd first two of which being longer and narrower, are call'd Superiores, are on the Basis of the Brain *, arise,

Ibid. EE

Ibid. A.

Ibid. cc.

according to him, from the Receptacula Sellæ Æquinæ, by the same Author so named, (hereaster to be described, though more truly, from the eccircular Sinus, as I hope in its place to make appear, running down from thence upon the internal Process of the Os Petrosum, and terminating in the Sinus Laterales, where they begin to be declive and tortuous in their passage to the internal Jugular.

The other two, called co Inferiores, which are much shorter and wider than the others, descend from the same place as the former, between the Os Petrosum and Occipitale, down to the aforesaid eighth hole of the Cranium, where the Jugulars come up into the Brain, and end there.

Another I discover'd by having first injected the Veins with Wax running round the Pituitary Gland on its upper side forwardly within a duplicature of the Dura Mater, backwardly between the Dura Mater and Pia Mater, there somewhat loosely stretched over the subjacent Gland is self, and laterally in a sort of a Canal made up of the Dura Mater above, and the carotid Artery on each outside of the Gland, which by being fasten'd

fasten'd to the Dura Mater above, and below at the Basis of the Skull too; seaves only a little Interstice betwixt it self and the Gland, thereby constituting a Cavity communicating with the two foremention'd forward and backward ones, from whence the abovemention'd four small Sinus's do descend, by a visible continuity, on each side from a little beneath the hinder Process of the Sella Turcica:

Fig. 2.EE and this from its Figure may not unfitly be called the Circular Sinus.

Vieusenius, it may be, saw some part of this Sinus where the other four small ones enter it, which is at the hindermost part of his Receptacula Sellæ Equinæ lateribus adjacentia, so called, and from thence thought those Receptacles to communicate with and to be capable of performing the office he assigns them, (viz.) of bringing back Blood from the nourishment of the subjacent Bone call'd Guneiforme, together with the Water separated stom the Pituitary Gland, into these sour inferiour Sinus's.

Now, as concerning these Receptacles of his, 'tis certain that they are not any where existent in Human Brains, (according to the description he gives of them in the place here noted) seeing both the third, fourth, Pag. 16. two foremost Branches of the fifth, as well as its third hindermost one, together with the fixth pair of Nerves, do not only run out of the Brain enclosed in so many distinct little Capfula's or Coverings made of the Dura Mater, during their passage through that part of the Basis of the Cranium by him call'd Receptacula, &c. but even the whole Dura Mater, together with its Membranous Productions constituting the aforesaid Coverings of those Nerves, in that place sticks close to the Basis of the subjacent Bone, (viz.) the External Process of the Os Cuneiforme, on its under side, and to the Carotid Artery (which also both above and below (as was before noted) by its borrow'd coat sticks close to the Dura Mater,) on that fide towards the aforesaid Gland. leaving no room at all for either Blood or Serum to be contain'd there, as he would have it; tho' in the same place which he describes for his Receptacles I have in several injected Bodies observ'd two very fair and large Veins, one coming into the Cranium at the second Foramen from the

the Orbit of the Eye, (and possibly may be a Reductory Vessel to that part) and so climbs up on the fide of the lateral Process of the Wedglike Bone, almost up to the Circular Sinus: the other at the fifth Foramen, which climbs up upon the same Bone till it meet and joyns with the other, from whence they make one short Branch. which enters the Circular Sinus very near the place where the two other inferiour ones on each fide descend down from it; which if they should chance to be cut by accident in any enquiry made into that part, might cause an appearance of Blood, and thereby become an occasion of the aforesaid erroneous Hypothesis.

Neither is it possible (granting there were any such Receptacles as he mentions) they should serve to the end he assigns, seeing the Glandu-Vieussp. 1a Pituitaria is on all sides enclosed by both the Dura and Pia Mater; which sirst (notwithstanding what he says to the contrary) is on all sides of this Gland of a very strong and equal thickness; yea, in that very part where (as hath been before taken notice of) there is a kind of a Chase made by a certain duplicature

of

of the Dura Mater, constituting the foremost part of the Circular Sinus.

And if this also was granted, yet would the manner he describes of the Serum or Water getting into these Receptacles (which is by transcolation) render his Supposition very unprobable, feeing 'tis by no means conformable to the Custom of Nature in all other parts of the Body that Arteries should depose a Serum, or any thing elfe but Blood, (except what goes for Nourishment to the Part it felf) in any Part, without being furnish'd either with its Excretory or Secretory Ductus, neither of which was ever pretended to have been found here.

And as a thorow confirmation of all this, faid in opposition to the afore-faid Hypothesis, I shall only add this, and conclude, that in several Injections made use of in order to find out the use of Parts, I never sound one drop of the tinged Liquors on that side of the Carotid Artery, where he hath made the situation of these Receptacles.

The use of this Circular Sinus is in common with the rest to reduce Blood returning from all the adjacent parts, as the Pituitary Gland, the

Wedg-

Wedglike Bone also, and it may be from the Rete Mirabile, which in Brutes is very large, and therefore feems to require the Service of this Sinus, either mediately or immediately, for reducing a share of its Blood, seeing the Glandula Pituitaria appears no where furnish'd with Veins terminating any where elle sufficient to carry off the refluent Blood from this Plexus, notwithstanding Vieusenius faith on the contrary it hath no Veins, and therefore is forc'd to have recourse to those small Branches of Veins which accompany the Branches fent out by the carotid Artery, before it perforate the Dura Mater, with the Optick Nerves, or those which go to the Gangliforme Plexus of the fifth Nerve, or those coming out of the Wedglike Bone, for reductory Vessels to this Part; but with what probability I know not.

CHAP

CHAP. VI.

Of the Motion of the Brain and Sinus's.

O these Sinus's, especially the Longitudinalis, and by way of consequence to the Lateralis also, most if not all the Ancients, as well as Moderns too, particularly Willis and Vieusenius, have unanimously Viensfen. ascrib'd Pulsation, after the manner of P. 14. Arteries, by reason of some Arteries (as they thought) from the Dura Mater terminating in them: of the truth whereof being somewhat doubtful, I resolv'd to make use of such an Experiment as might remove all future Scruples, and most satisfactorily put an end to the Controversie; which was as follows,

I took off the upper part of the Skull of a Dog alive, by which means the Dura Mater with its third Longitudinal Sinus lay bare to the Eye and Touch, to neither of which Senses, at first, either any beating of E

the Membrane in general, or of the Sinus, was the least discernable. After some pause, by chance the Sinus it felf, which I defign'd to have open'd with a Lancet, being touch'd with a cauterizing Iron (which in making the Experiment there was occasion to make use of) pour'd out the Blood very violently, and at first without any very remarkable pulfation, but after some time discernable enough, both as to the Blood and Membrane too.

I cut this Sinus through almost the length of it, to fee whether any Arteries (whereof many, according to Vieussenius, which was also long afore affirm'd, and that upon Experience too, by the learned Wepfer, did terminate in it, and so occasion its beat- Wepf p 116 ing.) would discover themselves by par. 1. throwing out their falient Blood, but

no fuch Sign appear'd.

After all which 'tis manifest the Sinus's themselves have no pullation, other than what is communicated to them from the fubjacent Brain, which contrary to what Bourdon affirms, hath an evident pulsation through the Bourd. multitude of Arteries dispersed thro' par. 2.

it so forcible as to create a sensible Systole and Diastole in its outward

coverings.

'Tis worth noting, that while the Blood-veffels are all full, so as to keep the Dura Mater upon its full stretch, the pulsation is not visible at all, or at least very faintly; but after a depletion of the Vessels, so, as that grows somewhat more lax, the beating becomes very visible, equally in the Sinus and Membrane too.

After having made this Experiment I found one Author of the fame opinion, and that is Falloppius, who in vindication of Galen against Vefalius, his Contemporary, fays, all I have faid upon the foregoing Experiment, and all the great Vefalius was able to answer in his own vindication in his ingenious Book call'd Anatomicam Gabr. Falloppi Observat. Examen,

falls very short of its aim.

As to the Transverse Ligaments * Fig. 4 r. which are in fome places * round, cordal, and in others if broad or membranous, in the Longitudinal Sinus chiefly, both ferving for Strength and (in concurrence with the cruciform ligamentous Fibres, taken no-E 2

tice

tice of by Vieusenius, on the under and outside of this Sinus, from whence the Fibres belonging to the falcated Process asoremention'd seem to have their original,) Elasticity to this part for its more vigorous reduction of the Blood passing through it, together with its blind Cavities or Diverticulums serving to moderate the overswift or violent motion of the Blood; seeing I find them so exactly describ'd by Vieusenius, to whom the Reader may have recourse, I think their description need take up no room here.

But as to the manner of the Veins entring this Sinus, I find it far different from that which is describ'd by Lower first, and afterwards by Vieus. Low. fig. 4. h h. senius, both whom make them enter vieus trab.2 with their Orifices from behind for DD, &c. wards, (two or three only excepted by Vieusenius) and that for some other useful purposes than what have hitherto been taken notice of.

And this is as follows, (viz.) About Fig. 4. one half of them (tho' intermixedly) dd, &c. (but all, after having first upon their arival at the Sinus infinuated themselves for some space after the manner of

the

the Pancreatick Duct or Ureters first 16. dd, &c. taken notice of by Lower, betwixt the Duplicature of the Dura Mater) from behind forwards, the other half from before backwards, as in the Figure.

> Now, by this contrivance 'tis plain, that first of all there are made two contrary Torrents in one and the fame Channel, by which means the refluent Blood, made poor by the vast quantity of its richest parts drawn off as it were into Animal Spirits, thro' a collision of Parts, which by this contrivance must needs fall out, is preferv'd in its due mixture, which when at any time lost through the languishing of its intestine motion or elasticity, retards even its circular or progressive motion, which when it happens but in some degree, is the cause of many Distempers; and when altogether, of Death it felf.

In the next place the circulation is at all times not only somewhat retarded, and the Blood hinder'd, (together with the help of the bony Cell at which the internal Jugular Veins enter the Sinus's) especially in an erect posture, from descending with that rapidness and weight it would

E 3 others

otherwise have done upon the descending Cava to the Heart; but also much more so retarded in a supine position of the Head, a posture most natural and ordinary for Mankind to take their rest in, through which contrivance, in concurrence with that of the Lateral Sinus's, (whose structure is such, that in the aforesaid posture the Blood is forced to climb upwards before it can arrive at the place of its descent into the Jugular Vein) there is made a more plentiful generation of Animal Spirits, one chief Cause of the great refreshment and vigorous disposition of the whole Body we find after Sleeping.

As to the other manner of the Veins entring this Sinus, (viz. from before backwards) it from thence happens, that in a prone Position of the Brain, a posture not uncommon amongst Men, the Blood is help'd forward in its circulation through the Sinus; the truth and design whereof are at once both evident and pointed at by Nature from the Structure of this part (and which therefore shews the great usefulness of Comparative Anatomy) in Brutes, who by reason of

fuch

fuch a Position, which the necessity of Feeding almost always keeps them in, have always such a disposition of this Part, to assist the Blood in

its heavy circulation.

The defign of Nature in making these Channels so wide on a sudden, in respect to the Branches of Veins lately treated of terminating in them, feems to correspond with the conformation of the Parts just now treated of, and with that it had in making the Ramifications of Arteries afore taken notice of so large and unproportionable to the Trunks from which they spring, which is a flower than ordinary circulation of Blood through the Brain, in order to make a still more copious production of the Animal Spirits so called. Which profitable Defign and End of Nature had nevertheless been attended with a very great Inconvenience, (viz.) an extravalation of too much Serum, the usual effect or consequence of a flacken'd Circulation, had it not been for another provident Contrivance of Nature in the two Communicantbranches, betwixt the Carotid and Vertebral Arteries aforemention'd, p. 36. E 4

p. 36. by the narrowness of whose Channel the influent Blood is in some measure represt in its motion, and an overcharging the Vessels with Blood

prevented.

These Sinus's differ in structure one from another, the Longitudinal and Lateral ones having many transverle Ligaments which the other have not, and the Longitudinal having many finall Cavities or bland Diverticulums, as aforesaid, which the Lateral have not; the use of them all being for strengthening and defending them from giving way to the violent irruption of Blood into them, against which sometimes notwithstanding they are not able to defend themfelves; as I have feen in many Skulls ni which the Blood hath burst open the fides of the Sinus's, and found its way between the Duplicature of it, so as even to have made a Fovea or Cavity in the Cranium it felf, as was before noted, one of which I have now by me.

CHAP. VII.

Of the Plexus Choroeides.

Body made up of Arteries, Veins, Membrane, and Glands, double on each fide, (which hath not before been taken notice of) and confequently having two Originals.

The first Original is from the foremost Branch of the Communicant
Artery, which running backward up
betwixt the hinder Lobes of the
Brain, (in which for some part of the
way it is immerged, and to which it
gives many large Branches) and the
Medulla Oblongata at length arrives
one part of the Plexus on each
side.

The second Original is from the hindermost Branch of that Communicant Artery, which running more backwardly, ascends betwixt the hinder Limbs of the Brain and the Cerebellum, till it comes to the Isthmus, where

Fig. I. fecond ce-

where communicating with the first Branch abovemention'd, they make a reticular broad Expansion, which covers both Nates, Testes, and Glandula Fig. 5. GG Pinealis, and constitutes the second or other part of the Plexus Choroeides.

> The first Branch begins to divide it self into divers Network Fouldings, interspersed with Glands somewhat before it enters'the Ventricles, and continues such to its Extremity on each fide, where they both under the Fornix wind crossthe third Ventricle into a mutual inosculation.

> The fecond begins to assume the same shape or contexture as soon as it begins to enter the Isthmus, continuing such throughout its entire abovemention'd Expansion.

> These two on each side are joined together by a twofold connexion, the first is by an Artery running under the Bombyces, intervening betwixt them, which could not be here inferted fo as to come in view.

> The second is by a production of the Pia Mater, which is extended all over these parts of the Lateral Ventricles, and the third Ventricle which

lyes

Ibid. 5.

Ives betwixt the first two parts of the Plexus forwardly, and down to the other two hinder parts of the Plexus backwardly under the Fornix and Septum Lucidum; so that what soever Water is transmitted out of these Ventricles, must slip down not only under the Fornix, but that Membranous Production it felf; from which kind of structure and position of this Membrane may probably be understood how there might happen such an Hydrocephalus as the learned Tul- Tulp. lib. L. pius mentions, in which there was found above two pounds of Water in one Ventricle, without any at all in the other: and fuch another as Wepfer mentions, where the Water Wepf. p.69 caufing the Hydrocephalus in an Heifer, was found contain'd in a Cystis, and that only in the left Ventricle too: for, supposing this membranous production of the Pia Mater to be double here, as it certainly is in all other places, 'tis not difficult to conceive, that the Water which is extravasated must needs infinuate it self betwixt the two Lamina's, till by a continual encrease it extends them into the shape of a large Bladder, fuch

fuch a one as the latter found there and drew out with his Fingers; and that which feems to put out of all Controversie that it was so, is, that in those places, both above towards the Corpus Callosum, and below on the Basis of the Ventricle, he found some fort of Asperities as though the Bladder fill'd with Water had been covered with some small Protuberances not much unlike to White Poppyfeed, in those places where it was contiguous to them; which Protuberances doubtless were the small Glands interspersed quite through this Plexus.

How this Distemper came to be on one fide only, though fometimes it is on both, as you may fee in another place of the aforesaid Tulpius, may likely enough be from an Ad- Willia p. 10 nascency of both the Lamina's of this par. 1. Membranous Production, in that place where the Septum Lucidum finks down from the Fornix, occafion'd by some small fort of pressure of the superincumbent Brain. fides these Veins, which are very truly describ'd by Willis, I have always found two more meeting the tore-

foremost Extremities of this Plexus, from between the two first Lobes of the Brain, where it seems to end under the foremost part of the Corpora Striata, by which it is there fixed and as it were kept in its due situation: and from these Branches are on each side sent forth many more little ones to the Corpora Striata, and several other parts adjacent.

To this Plexus belong also Veins, which from the Extremities of that part of it in the Lateral Ventricles part of it in the Lateral Ventricles ty large Trunks, running down thro' the middle of the third Ventricle, as far as the fourth Sinus, and there receiving some Branches from the other hinder part of the Plexus spread over the Isthmus, discharge the restuent Blood into that Sinus.

But besides this sort of Reductory Vessels, it hath also another, (viz.) Lymphæducts, which I first discover'd in the Brain of a strangled Body, and shew'd to several then present, running in different ramifications amongst the reticulated Vessels and Glands of this part: Which Observation being added to that of the great Anatomist

Ibid. 99.

An-

Anthony Nuck, who in that curious Piece call'd Adenogrophia fays, he Nuc.p. 150. faw one coming from the Glandula Pinealis, and that his Friend another Anatomist, whose Name he mentions not, (but I know it was one Bodivol, whom I had the Happiness to be very well acquainted withal, now dead) fent him word, he saw another not far from the aforefaid place; may be of fufficient authority to evince the real Existence of these Vessels hitherto so much enquir'd after, in the Brain as well as in other parts of the Body.

The Glands belonging to this Plexus are very many, but very small, and their Use, according to all the Moderns, especially Willis, Duncan, and Vieusenius, to carry off the redundant watery part of the Blood, but that without ever shewing by what rational contrivance of Structure it can be done, feeing none of them ascribe a Secretory Duct, which must always be in readiness when any unprofitable part is to be discharg'd.

Since therefore this part is found furnishid with Lymphæducts, 'twill be no hard matter to conceive the genuine use of the Glands, which is, to

lepa-

feparate a rich nutritious Juice from the influent Blood, and by the Lymphaduets to refund it to the refluent, after the loss of its noblest parts lest behind in the Brain, in its passage to

the Heart again.

It may also, for ought I know, according to the Opinion of Willis, ferve to warm its neighbouring parts the Internal Superficies of the Brain, which being purely medullary, hath not fo plentiful a share of Blood-vesfels dispersed through it as the rest, and consequently, to maintain an equality of warmth conducing fo much to the conferving the Spirits in their due vigour and exercise, must borrow an additional supply from hence. It is fituated upon the middle of the Thalami Nervorum Opticorum, all-along them length way, and, contrary to what Willis fays, is, by vertue of feveral Blood-vessels, join'd to that medullary part of the Brain fo call'd, immediately lying under it.

CHAP.

Of the Rete Mirabile.

7Otwithstanding the Opinions of the late Wepfer, Willis, and Vieussenius too, (which two last indeed, tho' but now and then, are willing to allow it an existence only in Men, (who nevertheless, if the Suppofition of Willis be true, viz. That such cannot but be Fools) had better be Willis p. 27. without it,) together with almost all col. 2. the Ancients, as Vefalius, Columbus, &c. to the contrary, I have never found this Rete wanting, or with any difficulty discoverable in Men, springing from and lying on the infide of each Carotid Artery, in that place of the Circular Sinus chiefly which looks into the four abovemention'd inferiour and superiour Sinus's in the Basis of the Brain, and in some measurealso the whole length of the Sella Turcica, on each fide, between the Gland and the Carotid Artery.

And that it is so small in them with respect to what it is in Brutes of several kinds, is no way surprizing, when consideration is had to the Use and Service of it in those Creatures,

who,

who, by reason of their prone Position, would otherwise be in danger of having their Brains deluged as it were with an over-great quantity of the Influent Blood, and of a Rupture of the Vessels, by its violent ingress, and this Danger so much the more threatned by how much the same Cause which brings it into the Brain with that force is equally as great and effectual to hinder its proportionable return; for the relief of which Inconveniency Nature hath contriv'd a means of its more easie and safe defcent into the Brain, by turning that one large Stream of Blood, (which through its being penn'd in one Channel, becomes to rapid) into many more, (by which means the Carotid Trunk above the Dura Mater in those Creatures is very small to what it is beneath, whereas that Artery in Men, &c. hath the same bigness on both sides that Membrane,) and they not only reticulated and contorted for the more flow and laborious (which Contrivance the Ancients thought was only for a more exact preparation of the Blood for Animal Spirits) descent of the Blood, but also many of them by their infertion into the Glandula Pi-

tuitaria

tuitaria, attended with small Veins issuing thence, to take off some part of the burden too, and leavest as snow

This last contrivance of Nature methinks may be sufficient to render Vieus. p. 46 that Controversie of Vieussenius with par. 2. Willis (which, before them, was betwixt Waleus and Rolfincius) the two latter on each side denying this Rete to have any Veins, very needless; feeing that if the Pituitary Gland have any, which I am confident it hath, (notwithstanding the positive Asser. Diemerbr. tion of Diemerbroeke, in order to serve par. 3. his own most unprobable Hypothesis, to the contrary) as having feen them plain injected with Wax; then this part of the Blood in some of the Branches of the faid Rete, which are plainly inferted into the Gland, is equally capable of being reduced by thole Veins without any necessity of having recourse to those remote Branches Vieussenius hath been forced to seek for, as if it had had par. 2. them of its own. The true at 1

And that to the aforefaid Position of different Creatures ought chiefly to be ascrib'd the variety of Magnitude of this Rete in feveral of them, its fize in Dogs feems highly to evince;

111

in which, by reason of their Horizontal Position, being neither so prone as several Brutes who seed on Grass, nor so creek as Man, that Rete is found smaller than in the first,

and larger than in the last.

Another Use it hath been thought to have, is, to carry off a confiderable quantity of a dull watery part of the Blood, in order to the production of the finer Animal Spirits; and this it is thought to effect by means and help of the Pituitary Gland, betwixt which and it felf there is constantly observ'd a great affinity, the one being either greater or leffer in proportion as the other is fo, and betwixt which there are in all Creatures, but more remarkably in those where they are both large, a diffribution of feveral Branches coming from the aforesaid Rete. And this is look'd upon by Vieussenius so considerable an office of the Glandula Pituitaria, that in those Creatures where it is but small, as in Men, Horses, Dogs, &c. he hath sub-views.p.102 flituted many, but particularly par. 3. two Cavities, for that use in the Wedglike Bone, just under the Sella. Turcica, in which he supposes that

part of the aforesaid Serum, which by the smallness of the Rete cannot be return'd that way, is remitted by several little Arteries slipt off from the Carotid, whilst under the Sella Turcica, terminating in the two abovenamed Cavities, there either deposing a part of the Serum to be carried off by a strange way he there mentions, (viz.) by two holes, into the Nostrils, and thence into the Fauces; or else by certain Veins meeting them in that place, as their views. p.9. Proper Reductory Vessels, to the par. 2. Heart.

Now, as to this office of the Glandula Pituitaria, I cannot easily be perswaded it is either design'd for, or capable of it, till such time the Abettors of this Opinion can be able to show me it furnish'd with an Excre-

tory Duct for this purpole.

And if they offer, that the Veins are such, I reply, That (besides its being very unprobable that so vast a quantity of Blood as continually is brought by the Carotid Arteries to the Brain, should be able to get rid of any considerable quantity of its Serosity, by so small a part as the Glandula Pituitaria is;) 'tis not the usual

usual way of Nature to part with any Share of its Juices out of its Veffels, when fo unactive and unprofitable as this is, and immediately to receive it in again, seeing it is provided of Emunctories enough to con-

vey it away by.

Moreover, granting (which by no reasonable means is to be granted) it were so as they would have it, yet nevertheless, in conformity to Nature's proceedings in all fuch-like cases, there ought to be an intermediate passage by way of a Secretory Duct, which none hath been able hitherto to discover.

And so far as Vieussenius seems to Vieus. p. 102 be of this opinion, which in one place Par. 3. he plainly is, making it of fo gross and viscid a nature, as is only fit to be discharg'd at the Emunctory of the Nose; the same Reply is satisfactory: But when by way of flat contradiction to himself he comes to make the same gross Humour a perfect fine Lympha, the Answer is then, Vieng. P. 54 That there is no need of parting with it beforehand, feeing we find that Liquor only separated by the Lymphæducts of the Brain afterwards.

Seeing therefore there is such an affinity as before mention'd, between the Rete Mirabile and Glandula Pituitaria, and taking it for granted, that the office of the Glandula Pituitaria is not what it hath generally hitherto been believ'd, to the end we may attain a more exact knowledge of what it really is; it seemeth not altogether immethodical to take that part into confideration in the next place, together with the Infundibulum, which last hath not only as near a relation to the Gland as the Gland hath to the Rete, but such a close communication with it, that it feems in a manner almost impossible to treat of one independently on the other.

CHAP.

therefore there is such

d taking it for granted, that

GHAP. IX.

Of the Glandula Pituitaria, and Infundibulum.

HIS Gland is seated in and fills up in a manner all that space contain'd within the Sella Tur-

cica (Vessels only excepted).

Tis cover'd on all fides with the Pia and Dura Mater, excepting that part on its upper Superficies, in which there is a little round hole, by which the Infundibulum descends slopingly into it, being at its entrance inviron'd with a Production of the Pia Mater, for its more firm connexion with that part, as was before noted.

But as to the Dura Mater, it encompasses it after a far different manner than what Vieusenius hath de-Vieus. P.51 scrib'd, not suspending it in Man as pare 5 it doth in Brutes, so as to hinder it from touching the bottom of the Sella, and that forasmuch as there is not the same reason for its so doing in one as there is in the other, for in Brutes

Brutes the Rete Mirabile is not only fituate on each fide this Gland, but runs quite under its hinder part, by which one fide of the Rete communicates with the other, a Disposition of this Part which Vieusfenius was altogether unacquainted with; whereas in Man, inafmuch as there is not that fort of Structure in the one (i. e. the Rete) 'tis not necessary it should

be requir'd in the other.

However, in neither one nor the other is the Reason which Vieusenius gives for Nature's contrivance of this affair of any weight, feeing neither the Rete Mirabile, much less the few vieuff.p. 50 small Veins belonging to the Bone par 1. beneath, could possibly any way be compressed by this Gland, though superincumbent, because it is so firmly knit to the Dura Mater, lying above and upon it, which is supported by the two foremost and hindermost Processes of the Sella Turcica. in fuch a manner as is fufficient to fustain and keep from pressing upon any subjacent part ten times a greater weight than the Glandula Pituitaria is. All of the state would w

Moreover, the Dura Mater is fo far from fulpending it from that Bone, Bone, that it is, together with the Gland, fixed to that very Bone it self.

The substance of this Gland is far differing from that of all the rest, which I have often upon this account particularly examin'd; in consistence indeed 'tis the same with most of the Conglobate kind, if not somewhat harder, but then being pressed or squeezed, it emits much more Water than any of them.

As to the Conglomerate fort, it hath not the least resemblance to any of them, and consequently cannot be supposed, as it hath hitherto been by all, to carry off any excrementatious or unprofitable part of the Blood.

Now, if we consider this part, together with the appended Infundibulum, we shall certainly find a conformation far different from any other
part in the whole Body of Man, inassumed as that which this Gland
receives by the Infundibulum, or which
is the same, what this Infundibulum
conveys to it, is not separated from
the mass of Fluids by any visible
Secretory Duct, which in its ordinary method Nature is observed constantly to make use of, whensoever

it parts with any part of the Blood, whether excrementitious or reductitious, throughout the whole compages of the Body.

Nor hath the manner of Nature in transmitting a certain Liquor to the Gland been less abstruse in carrying it off from that part again, the reductory Vessels from the Gland being equally conceal'd, as the adductory to the Infundibulum; that way of Transudation, according to the invention of Vieusenius, being to the greatest degree improbable, as having no resemblance to the course of Nature throughout the whole Body.

Nay, even a possibility it self seems hardly allowable, if we take but notice of that part in Brutes in whom its Integuments are extraordinary denle, the Dura Mater, as he truly observes, investing it close on every fide, (and which he perceiving, and confequently foreseeing what might from thence unanswerably be objected against him) was forced to make them much more than in Men; in which last indeed there is feemingly Views P.52 fome reason for its being so, inas- par. 2. much as the Rete lies in a Duplicature as it were of the Dura Mater, on

each

each fide of the hindermost part of the Sella Turcica, as tho' one Lamina of it was spread upon the subjacent Bone, and the other over the Pituitary Gland, (a disposition contrary to that in Brutes, as hath already been taken notice of) but nevertheless there is no necessity that it should be so divided in this place, nor doth the faid Author ever offer a Reason for its being so, (which looks as though his Affertion was only a Guess) seeing this Membrane can fend out new Productions as well double as fingle, as we find in its two eminent Processes before describ'd, and Sinus's; agreeable to what it also therefore may and does do here, where the Integuments of this part appear plainly to be of too thick a confistence to admit of his imaginary way of transudation, which is manifest not only by fight and section, but in that by the greatest force made use of in compressing and squeezing it between ones Fingers, we find it impossible to force out the least appearance of Humidity through its aforesaid Inclosure or Integuments,

Being therefore very inquisitive after the true use of this part, and despai-

despairing of ever attaining to such a Knowledge without first knowing the exact Structure thereof, besides all other means commonly made use of in all Anatomical Enquiries, I made ule of all forts of Injections serviceable to fuch an end, as of tinged Liquors, Wax, and Mercury, but all with little, if any, success according to my expectation, the Wax not penetrating its Texture at all, the tinged Liquors but very superficially, and the Mercury, (where my chief Hopes were) always by its weight (do what I could to the contrary) either breaking through the fides of the Infundibulum, where it leaves the Brain, or elfe falling down in greater Globuli than the extream narrow Panages were capable of admitting, and by this means became altogether ufelefs...

Being compelled therefore for the present to leave off a little while a further enquiry into the Structure of this part, by reason of the great mist it is involved in, and to gain a little more Light for our Guidance in searching after Truth, (which like many other things of greatest value lyes deep, and is with

with great difficulty accessible) it may not be amiss to see what Affiftance can be had, by making diligent Scrutiny into the Structure of its Appendix the Infundibu-

dibulum.

The Infun- This is a thin medullary Duck. covered with the Pia Mater, descending from the internal Concave Superficies of the Brain, to which, by reason of its wideness towards one end, and narrowness towards the other, in resemblance to a Tunnel, as well as by reason also of the parity of their Uses, the Ancients gave the Name of Infundibulum.

> In Man it is closely invested with the Pia Mater at its very entrance into the Gland, and from that place hath not any manifest Cavity I could discover by blast or style, but is altogether of a medullary substance, contrary to what it is in Sheep or Calves, in which last Creature, where the Parts are larger, by inferting a Blow-pipe into that part of the Infundibulum, next to the Gland, I have feen its further Tract or Passage on the upper part thereof a little puffed up, and a confiderable

siderable quantity of Water regurgitate, as though it had lain contain'd either in some Pipes or Porulous Substance of that Gland.

This Difference is not taken notice of by Vieussenius, and therefore what he says of this part seems chiesly in this respect, if not altogether, applicable to the Structure it hath in Men.

Those two Divisions or Ramissications of this part the faid Author men- Vienffen. tions, one forwardly, and the other P. 49. backwardly, in Sheep, Calves, &c. I have always found correspondent to the Descriptions he there gives of them; but whether the first be protended so, and terminate after the manner he there describes I somewhat scruple, seeing I have always observ'd the Extremity of that part in Brutes, towards the foremost part of the Gland, finking as it were into the very Substance thereof, and afterwards becoming presently altogether imperceptible, and in Man the termination thereof just after the same manner, save only that in the last it happens forthwith upon its approach to the Gland,

without being protended either back-

wardly or forwardly.

The Use of this part is certainly to convey some fort of Humidity from that great concamerated Cavity within the Brain, resulting from its inward complication of parts, to the Pituitary Gland, and the office of it is to receive and carry off this transmitted Humidity; but as to how either this Humidity is collected in the aforesaid Cavity, or how, when convey'd into the Gland, it is carried off, we are still as much in the dark as ever.

I know very well there is nothing more easie with the Visionary Philosophers than such a Knack as this; and now I think on't, the great Williamakes nothing of turning a Vein Will. p. 46. into a Lymphædust in the Glandula Pinealis and Plexus Chorocides, no less than which does also the accurate Vieusenius, in the Plexus belong. Vieusento ing to the fourth Ventricle; but how par. 3. consonant this is to the rational structure or mechanism of parts, neither the one or the other have been so kind as to explain.

Sits of Masongers 231

Now

Now, as to the Plexus and Glands before mention'd, 'tis evident by what hath been already discover'd and accordingly given an account of in the preceding Pages, they are furnish'd with Lymphæducts, as proper reductory Vessels; so that so far the Pro-

phecy is vanish'd.

But as to the remaining Gland, I am not so fond of guessing to say it hath any, and consequently all I can say is, that as I look upon the Infundibulum to be no more than a large Lymphæduct variously ramified through the Glandula Pituitaria, discharging its Liquor by those many small Branches into the Veins dispersed through that part to be reduced after the manner 'tis in all other Secretory Glands back to the Blood again.

And that which seems most to favour this Conjecture, is the extraordinary humidity of this Gland, especially in Brutes, above the rest of the whole Body, as serving not only to export what Lympha is separated from several Arteries dispersed thro it, but that also which it is charged

with from the Brain it felf.

And

And to this twofold manner or double office of Secretion is owing the two distinct Substances it seems to consist of, the one being accommodated to that part of the Lympha coming from the Brain, and is therefore whitish, the other to that separated immediately out of the Blood, and is therefore reddish.

Lastly, As to the manner how the Lympha passes down thro' the Infundibulum from the Brain to the Glandula Pituitaria, I look upon it to be in the form of condenled Vapours arifing from the Arteries of the Plexus Choroeides, emitted thence for the keeping moist and in good order that inward Production of the Pia Mater; spread all over its Parietes, which being a membranous dry part of it felf, might otherwise become injurious to that fine medullary part lying under and being contiguous to it; in which there is a continual motion of Animal Spirits, whose Tracts, and consequently they themselves, through any the least intemperance of this Membrane, would be in great danger of either some obstruction or disorder.

And that this Lympha is only the result of the aforesaid Vapours, I am the more readily enclin'd to believe, because I never saw Water in that part of any sound Brain, nor unsound neither, where the Plexus Choroeides was firm; and there was no reasonable ground, by the extravalation of Serum in some other remote parts of the Brain, to believe it had its rise from thence.

CHAP.

CHAP. X.

Of the Glandula Pinealis.

THE Gland call'd Pinealis, from its Figure, is about the bigness of an ordinary Pea, prefix'd to the two Prominencies call'd Nates, hereafter to be describ'd, at the end of the third Ventricle, immediately under the broad and hinder part of the Fornix, (with which nevertheless it hath no connexion, as Vieusenius vieuse.71. saith it hath) and over that part of the Rima in the third Ventricle call'd Anus.

'Tis joyn'd to the Nates by several Fibrous Roots, and becomes a support to that part of the Plexus Choroeides there situate.

In an hydropical Brain of a strumous Boy, I have seen it swelled to a fize of three times its ordinary magnitude, and by reason of the abundance of stagnate gelatinous Lympha contain'd in it, perfectly transparent.

G 2 Hence

Hence it most plainly appears that this part is a meer Gland, and, by what was faid before conformable to what hath been observ'd in this hydropick Brain, of the Conglobate or Lymphatick kind, and by confequence a very unfit part to be made a Receptacle for Animal Spirits, as Vieuse- Vieus. p.71. nius makes it, and much more a place of refidence for the Soul, according

to Des Chartes.

'Tis true, there are two fair medullary Tracts arifing feemingly from the two Roots of the Fornix, stretching length-way upon the Thalami Nervorum Opticorum, as far back as this Gland, (by Vieussenius called vieus, p. 64 Tractatus Medullaris Nervorum opti-par. 3. corum Thalamis interjectus, as though it was only one, and accordingly is fe delineated by him, Tab. 7. GG, but indeed is two, one on each fide) about which place they turn in, and by a transverse bending kind of a Process (by the same Author call'd Tractus medullaris natibus antepositus) unite, as he hath exactly observ'd: And willis, p. 9. this, doubtless, gave occasion to the col. 1. Error of Des Chartes, as Willis tru. par. 1. ly thought, (whose sublime and most

deservedly-admir'd Philosophy had doubtless been much more useful, had he convers'd more with Diffections, and less with Invisibility) and Vieusenius too, (with whom in p. 508. the same Mistake doth agree Mu will p. 100 raltus and Willis) for upon a col 1. par. 3 more heedful inspection (as was most evident in the Brain afore. mention'd) it will be found that no part of the Process aforesaid. however near it comes to this Gland, does in any wife become continuous to it.

Whart.

Dr. Wharton also stum'led upon P. 141. these medullary Tracts, placing them amongst the Nerves themfelves, and ascribes the same unreasonable use to them as he does to the Nerves in many other Parts of the Body, (viz.) of separating a fuperfluous Humour from the Cruça Medulla Oblongata, or Thalami Nervorum Opticorum, (being the same Part, and only on the other fide or upper part of the Brain, under another denomination) which he supposes to be the Commune Sensorium.

CHAP.

CHAP. XI.

Of the Brain in general.

HAT part of this Treatife relating to the Vessels, being dispatch'd, I shall in the next place proceed to an account of the Brain it self, under which term are generally comprehended the Cerebrum and Cerebellum, and Medulla Oblongata, which Parts being in many respects so different one from another, may justly challenge a distinct and orderly description.

The Brain then, in the first place, as distinct from the other two, is that large and almost spherical Body which comes first to fight in the old way of Diffection, filling the greatest part of all that space contain'd in the Cranium, confisting of two different Substances (first taken notice of by Archangelus Piccolominius) Piccolom. both in Colour, Confistence, and Office, the one being more com-G4 pact,

pact, white, medullary, or fibrous, the other foster, greyish, and glandulous.

The utmost Malpighius (by vertue of his Microscopes) could do, Malp. de was to discover the Cortical part to p. 78, 81. confift of Glands of an oval depref- par. 3. fed Figure, and in his Opinion, of the Conglomerate kind, (but that how properly, as also his calling the Nerves their Excretory Ducts, I leave to the Judgment of others) and the Medullary part to confift of various Fibres immerged in and having their original from the aforesaid Glands, deriving from them a certain Liquor call'd Nervous Juice, concerning the Existence of which, in the usual sence 'tis taken in, as a fluid body, contain'd and running continually in the Channel of the Nerves, as Water in Wooden or Leaden Pipes, for either Nutrition or Censation, is a thing somewhat improbable, it being not only possible, but very easie to resolve those two Phænomena's, the first from the Blood, and the other from the Natural Tenseness of Sensible Parts maintain'd by the supply of a proper Liquor from the Blood.

Blood, both in their Originals and continued or elongated Productions; inasmuch as it doth as certainly circulate in them as in any other parts of the Body. And as to the manner how this is done, it will appear very plain and intelligible, after the innate Structure of the Part hath been more accurately enquired into.

The Curious Lewenboeck made a Lewenh. far deeper scrutiny into these two Cirib. p.37-Parts, being very probably affifted by better Glasses, and from what occurr'd to his view, called the cortical part a pellucid Vitrious Oily Substance, (the seeming oiliness of which Substance I attribute only to the stagnating of the pure Liquor, growing cold after death of the Creature,) from such a close and regular Position of the Globuli swimming therein, as allows the Rays of Light to pass them without refraction, contrary to what they do in the other or medullary part of the Brain, in which they are so dispos'd that the Light cannot pals them in right lines, and consequently being a little distorted, makes them appear white,

white, notwithstanding Malpighius Malpig. de on the contrary neither allows the Cereb. p. 2. Parts of the Brain to be diaphanous, nor the Animal Spirits to be any

thing a-kin to Light.

'Tis true, even by his own confession, that his most nice and disigent Inspections could not free him from many Scruples about what he saw; yet some things to our purpose were plain enough, as Reticular Bodies of a red colour, which being larger in the Cortical Parts than Medullary, helps to give it that greyish or subrunneous colour, as he calls it.

Nextly, a transparent Vitrous-colour'd Substance contain'd in most minute Vessels; whence 'tis plain there are two forts of Liquors in this Cortical Part, one of a red colour, or Blood, contain'd in larger Vessels, whose Globuli, which give it its redness, either by reason of their size or figure, cannot enter those small Vessels which with the Fluid contained in them constitute this transparent, cineritious, or cortical part of the Brain.

The other a transparent Liquor, contained in most minute Vessels, as aforemention'd; from whence I am induced to believe this Cortical part to be only an Aggregate of different Vessels, (as also I do of all the rest of the Parts of the Body) containing different sorts of Fluids.

Of these Vessels some contain a more compound Liquor, commonly call'd Blood, which whilst in that state, by reason of the Globuli swimming on it, looks red, and by reason of a tubulous Pore of a proper size and sigure so continued to the Vessel we call a Vein, that it undergoes a continual quick circulation.

Another fort of Vessels there is which receive and contain a more simple sluid body, of a thin transparent nature, which when in some parts of the Body, gives the name of Lymphæducts to the Vessels that it runs in; but when in these Vessels, which are discover'd to make up the great Substance of the Brain, whether Cortical or Medullary, may be allow'd the name of Fluidum Animale.

And

And this last sort of Vessels I look upon to be either a certain Protension of an Artery, by its smallness render'd capable of holding such a sort of Liquor only as the last spoken of, or else such a tubulous production of the Artery as by its Orifice or Pore answers to the figure and size of the Fluid it is by

Nature intended to receive.

Upon the same exact

Upon the same exact Enquiry made by a Microscope, the medullary part of the Brain appears to be of the very same constitutive parts, ranged only after a somewhat different manner, which makes this part appear more white, as was before observ'd. But over-and-above (if it may be allowable to make a Conjecture) I am enclin'd to think the Whitenels of this part may be owing in some, if not the greatest part, to such a narrowness of the Vessels discover'd here, containing the pellucid Substance aforemention'd as will not entertain any Fluid whatfoever, without its being first reduc'd into very minute Particles, or Septometry fo called: Which last Vessels I therefore suppose to be only

only yet more Capillary Productions of the aforesaid Cortical Vesfels, as they are of the red or Blood-veffels indu'd with fuch a Pore as fits them only for the reception of a most subtile, fine, fost Liquor, which I esteem the true Medullary and Nervous Juice, which being contained in its proper Capfula, and many of them collected into one Fasciculus, at its egress out of the Brain, being there wrapped up in more thick and strong Coverings made of the two outward Membranes of the Brain, do constitute that part we call a Nerve, which having all its Integuments or Membranous Inclosures always kept turvid and tense by its contain'd Fluid, after a flow and leifurely manner continually dispensed from the Fountain, and by its growing more taper towards the place of its termination, by which means it acquires a greater streightness or narrowness of its Pores ordinarily call'd Fibrilla, it so falls out that all inward Impressions, upon all occasions, are the more easily and speedily transmitted through it.

The

The very fame notion also concerning Nutrition (which in the truest sence is only an apposition of Parts nourishing to Parts pre-existent to be nourished) in the rest of the Parts of the Body, I have thought reasonable to entertain ever since, by affiftance of the Microscope I have plainly discern'd the Veins to be only continuations of Arteries, and the Blood to run in the fame Channel variously modified, without the least suspicion of Extravalation, (viz.) a continual transmission of Nutritive Juice out of the Pores of Arteries, after many windings like Tindrils of Vines (Analogus to which the red Reticular Bodies of Lewenboeck seem to be in the Brain,) grown very capillary into certain Tubuli's or Pores of a corresponding bigness and figure, making up the whole fleshy part of the Body, whose Substance, when 'tis freed by washing or injection of Water, we fee to confift only of large and small Blood-vessels and Fibres; which last, whether Nervous or Membranous, or fuch as relate to Muscular Motion, commonly called Carnous, I tupsuppose to be full of minute distinct Vessels for the communicating and receiving their proper Liquors or Fluids after the manner already expres'd, which as contain'd in the faid Tubuli or Pores, whilst they retain their Natural Constitution and Proportion, I presume it is which keeps the Habit of the Body plump and vigorous, the more thin and languid being perpetually carried back by the Lymphatick Vessels, and a great part wholly exterminated by meer simple Transpiration; which I adventure to think is not only superficial from the Sudorifick Glands in the Skin, but also through the whole Substance of inward parts. through small Canaliculi's or Meatus's in even the Viscera themselves: by which, not unlikely, we may guels at the Meaning of Hippocrates, when he said, All things were conspirable and transpirable.

The minuteness of Vessels is that which hath so embroil'd the Thoughts of Naturalists upon this Subject, and set Realities so remote from the Understanding, otherwise 'tis no Paradox to affirm the Existence of Vasa

Vaso.

Vasorum almost to Infinitum, some containing Liquids in a continual more nimble circulation, others in a gentle protrusion only: Which will appear altogether unfurprizing, if it be consider'd that the aforemention'd Ingenious Author hath computed, that even the 64th part Lewenh. of a Miriad (i.e.) of a Ten hundred p. 46. thousandth part of any Substance but as big as a small grain of Sand, cannot, especially if of a rigid or inflexible nature, enter those little Vesfels, which are feen in a retiform manner distributed amongst, and fixed to the aforesaid pellucid Globules, which swimming in those little Vesfels, are discover'd to make up both the Cortical and Medullary part of the Brain. As also further, that even the tender Coats of the smallest of those Vessels which contain the aforesaid most minute Globular Fluid Bo- Lewent. ib. dies, are also full of yet far more minute Vessels than they themselves are.

Nay, I am fo far from being furpriz'd at this kind of Vascular Constitution of Parts, that I apprehend not how Nature could otherwise have

acted

acted without the confequence of a boundless Accretion, inatmuch as that when any parts of a Fluid become extravalate, they necessarily lose much of their progressive motion, and if of a groß consistence, are either proscrib'd by the wider passages, or of a finer, through those more straight and elaborate (viz.) by Transpiration; so that what Particles of Matter soever continually arrive, for either the augmentation or reparation of the Parts, must (unless the ruine of the Subject do not first happen, as we see it often does in Diseases proceeding from fuch Causes) needs (if not confin'd in Vessels) advance into a monstrous preternatural accumulation, as being, by reason of their gross consistence, altogether uncapable of being carried off proportionably to the measure of their aggestion, in the form of subtile Steams or Exhalations.

Besides a rational explication of the natural Functions which this Hypothesis furnisheth us with it also, seems to clear a great many Difficulties which have hitherto puzzel'd the most refined Physiologists relating to

fation and Muscular Motion; of which last here in the next place, the other being reserved for the last Chapter, which treats of Sensation and Motion in general.

O recite the Opinion of others upon this Subject would be a thing altogether useless here, seeing an Abstract of them is already extant in the Philosophia vetus & nove by Mr. Colbert; and befides, the most correct of them are not only very unprobable, but ablolutely repugnant to plain Reafon and Matter Fart too; an Inflance whereof Phave in Dr. Willia's Tendi- Wills at nous Refervatories of Animal Spirits, p. 35 in Dr. Mayon's Twisting or Fiddle Mayow de -3]8m .20m thring Fibres, with whom of Mr. Regu agrees, by which the Musicie must needs lose a great deal of its thickness, than which nothing is more contrary to Experiment; in Duncan's first and second Element Dane p. 90 of Des Chartes, which he makes the Animal Spirits to confult of, contrary even to the very Principles of that great Man's Philotophy, which 31-

the Animal Faculty, Juch as are Senfation and Muscular Motion; of which

Of Muscular Motion.

O recite the Opinion of others upon this Subject would be a thing altogether useless here, seeing an Abstract of them is already extant in the Philosophia vetus & nova by Mr. Colbert; and besides, the most correct of them are not only very unprobable, but absolutely repugnant to plain Reason and Matter of Fact too; an Instance whereof you may have in Dr. Willis's Tendi- Willis de Mot. Musc. nous Reservatories of Animal Spirits, p. 35. in Dr. Mayow's Twifting or Fiddle- Mayow de flring Fibres, with whom of late p. 73. Mr. Regis agrees, by which the Muscle must needs lose a great deal of its thickness, than which nothing is more contrary to Experiment; in Duncan's first and second Element of Des Chartes, which he makes the Animal Spirits to confift of, contrary even to the very Principles of that great Man's Philosophy, which al-H 2 lows

lows no Elasticity to those Bodies themselves, though the Authors of it in all others; likewise in Dr. Croon's croone, making the Blood it self, as well as p. 23, 24, the Animal Spirits, to be mov'd by 25, 33. Philos. cell. the power of the Soul to any p. 23. Muscles; as likewise the extravasation of those two Liquors first into the spaces betwixt the Fibres, and then their introvasation into the Fibres themselves again, in order to make inslation, an Error incident to the Immortal Borellus also, whose ima-Borel de ginary Discourse upon this Subject mot. Anim. p.ult. prop. seems of a very different Thread 23. & plusom the rest of his Excellent ribus alus locis.

dy said about the Structure of Parts be remembred, (viz.) That the Medullary Part of the Brain is only a Contexture of Vessels; that its Nervous Propagation or Nerve is also a Compages of Vessels, formerly call'd Filaments, much more narrow than those of the Brain it self; and, that these Nerves produce, or at least terminate in the Fibres of all forts of sensible Parts whatsoever, though of a different texture, as well

well as those carnous ones of Muscles, which last are tubulous, 'twill not be in the least unreasonable to infern, That thefe Botered dies being kept continually turgid with the contained Fluid; are equally capable of transmitting or receiving Impressions of the Object, as if they were Aretched longitudinally like a Bow-string from each Extremity, according as Borellus hath obsermake inflation, an Error incidenbay.

Croones

And as to Muscular Motion, allowmink som ing only what may directly be inquiq sla ferred from what hath previously been said, (viz.) That the Nervous and Carnous Fibres are only a congeries of Fluids contained in certain Vessels communicating with each other, that by reason of a Plenitude in the aforefaid Fibres, the whole Machine is in a constant Equilibrium, it will necessarily follow, upon the common Postulatum, (to which all Mankind must be beholden upon all fuch Explications as these to the World's end) viz. that the Sensative or Rational Soul can command the Animal Spirits (which I call only a Nervous Fluid) into a equivamen

H 3

Primus Impetus, or local motion, that a part of that Liquor, whenever a Mufele contracted is transmitted through the Vessels which contain it from the great Refervatory thereof, the Brain, to its Car nous Fibres, into whose Vessels, being so much narrower than those of the Nerves, even by vertue of the fame force which moves it from the Brain, that Liquor is driven after a most rapid manner, (which Effect, to any acquainted with the nature of Fluids and mechanical Laws of Motion by Projection, needs not any demonstration) causing the Intumescence or Inflation of the Muscle, the same Liquor at the same time being driven back again with an equal speed from the Antagonist Muscle into the room of the first, which was transmitted from the Brain to the contracted one, in order to maintain the same Plenitude or (which is the same thing in the fence of the old Philosophers) to avoid a Vacuum. And if any object the wideness of the Passage it is to come back by from the reflexed relaxed Muscle, as an impediment to an it comes to pais by

equivalent speed in that Liquors retrocession, I have to answer, that the Emptiness being made first, is a sufficient recompence for that.

And here I cannot but take notice, that all they who contend for Animal Spirits, analogous to those we fee produc'd from various Subjects by Fire, as the only adequate medium for all forts of Muscular Motion, have been forced to have recourse either to certain Tracts or Interstices betwixt the Filaments of the Nerves continued from the Brain, or the Original of the Nerves through their whole Productions to the Muscle, of which fort are the Cartefians, or else to a certain Neryous Juice, for their place of refidence, of which fort are most of the Moderns, and particularly Vieuf-Senius, by which Passages, or out of which Juice these fine invisible things are either voluntarily, by the command of the Soul, or inadvertently, from feveral either inward or outward impressions, transmitted, in order to produce Motion: which if true, and the only ways of producing Muscular Motion, I beg leave to ask, how it comes to pais, by H 4 either

either of these ways, that when another person bends my Arm, and that against my Will too, the bending Muscles of the Arm become as tumid as when voluntarily or inadvertently contracted at any other time; which hath been truly observ'd the not latisfactorily accounted for, by Dr. Croone, or any other I know of. Croone, p.7.

But how this or any other fort of contraction of a Mulcle happens, does by the other afore mention'd Hypothesis become explicable, with-Yout any manner of difficulty at all: For when the Cause of Contraction is from the Command of the Soul, the pressure is first from the Fluid in the Brain, by which all the interfacent or continued Fluid flows towards the Part to be moved, the fame proportion of Fluid being at the same instant transferred into its room from the relaxed Muscle; and when the contraction of the Muscle is from the above-mention'd external of force bending the Arm against my will, then the Liquor contained in the relaxed carnous Fibres or Vascula is transmitted through the whole concinuity of Fluids, to that which is contracted, and all this without being beholden to the wild Conceits of a dry and moist part of the Nervous Juice, blind Passages, invisible Tubuli betwixt the Antagonist Muscles on Valves in the Nerves, by a meer Equilibrium of the Fluids contained in the Vessels the Parts consist of

vd. At the same time I am not infenfible of the Solution some have given this Instance of Involuntary Motion upon another Hypothesis, (viz.) by supposing an equality of Tension or Elasticity in all the Muscles of the whole Body; by which means it falls out, that when any new additional force (though never to small) is added to the Fibres of any Muscle, as in voluntary motion, or the power of Elasticity of in the Antagonist Muscle, overcome by outward force, as in the aforemention'd Instance of Involuntary Motion, the other Muscle then becomes contracted anno addition

Now, that this is one concurrent Cause in both forts of Instances, as being confirm'd by the Experiment of cutting a Muscle through, either towards the Extreams or in the middle, by which the Fibres, by their

-sncontracted, and all this without being

natural Elasticity, are found to contract either to one or the other, or to both Extreams, is allow'd to be true; but to be the only Cause, is altogether as falfe. Tobash bon raffiff

For, in the first place, as to the ease of voluntary Contractions, it is allow'd to proceed from a transmission of Spirits from the Brain into the carnous Fibres, (that Hypothesis of Steno to the contrary having been convicted long fince by Borellus, in his Book De Motu Animalium) though not without the concurrence or fympraxis of the natural Elasticity of the Fibres belonging to the Muscle to be contracted, remaind the regues a mort

bo So likewife, without the transmillion of Animal Spirits from fome force or another, I deny even the posfibility of that stiffness or hardness which is eafily preferved in all contracted Muscles, feeling and seeming as though they were indurated and fwelled out, as really they are, whether it be in the case of voluntary or involudtary motion; in confirmation of which, I affirm, that though by the cutting of the carnous Fibres of any Muscle through, which way fo-

ever

ever it be, the contracted part may, and doubtless does, grow thicker by the fhortning of its Fibres, yet by that means only it does not become stiffer and harder, so as we find Mulcles do when contracted by any natural Cause, nor is there any neceffity it should do so, according to any Rules of Mechanism, seeing the Fibres fhortning only by their own elastick force, when they find the circumambient space give way have no necessity of subintration of parts, which is always requifite to procure a stiffness or hardness to a part altering its dimensions as Muscles do, from a longer and thinner to a shorter and thicker circumference; and upon this it must needs follow, that in a Muscle contracted by involuntary force (in which Action the Brain is altogether unconcern'd) that stiffness or hardness then perceivable in it, must needs be owing to the Fluid or Spirits in the antagonist Muscle, after the manner already explained, transmitted to it. no item or is bullowni

of which is afternot that though by

CVOY

the cutting partition carnous Tibros of

Now, to define what fort of thing this Animal Fluid (fo called) is, I fee no occasion to trame any other Idea of it than what we ordinarily bave of the pureft Liquors, feeing the Nerves are a Substance which (to the Senses of either Smell or Taste difcovers very little elfe than what is infipid) are always reckon'd amongst the least hot parts of the Body, and doubtless far less warm in Fishes than us, who yet have as great a flock of Animal Spirits as any other Creatures. And this Confideration may be it was that occasion'd an Author to give the Animal Spirits the Epithire of Frigidiusculi. by before Ido Du Ham.

Tis plain enough, that the Vessels T. 1. p.753 which contains this Fluid are extream minute, and consequently the Content must needs be of a very fine and depurate consistence, though without much resemblance to either the aforesaid nimble, saline, or sulphurous Productions of the Fire.

Tis in a continual, gentle, direct motion, though perhaps contained in curved or reticulated Vessels, from its original source to the ends of the carnous Fibres, from whence

pens

it

it is convey'd into the Membranous or Tendinous Productions, according as the Fibres terminate, and it may be by filtration only; in which, as in other and particularly in Glandulous Parts not Subservient to Muscular Motion, where Nervous Ramifications are very copious, whether it be of any other use than to keep the Parts in their proper tone, in order to their regular discharge of the office of Secretion, must still remain a Controversie, notwithstanding all that hath been yet advanced against it, inafmuch as wastings and numbnesses of Parts, the common Symptoms of obstructed or divided Nerves, (which doubtless by their hastening through happening fuch Causes to Muscular Parts, gave the first rise to that Conjecture about the Existence and Use of that Juice throughout the whole Body) are equally explicable by the want of Tone, as of that supposed Liquor.

To the proof of all this an Experiment frequently made does not a little contribute, and that is the injecting the Arteries of a Dog, or any fuch Creature, when dead, upon which there immediately hap-

pens

pens a contraction of the Muscles, according to the different strength of them, (viz.) of the Extenders in the hinder Legs, and of the Benders in the fore Legs, though the Injection be only of cold Water, the reason of which effect in particular, if it be remembred what hath been before observed, (viz.) that the Blood-vessels do most certainly enter the composition of the Nerves themselves, will not only become very easily explicable, but the whole Hypothesis at least very highly probable.

If it be faid, That this speedy instantaneous ressure of the Animal Fluid is opposed by the aforementioned constant direct motion it hath from its Source to the parts to be moved, 'tis case to reply, That its slow direct motion that way is easily overcome and repelled by the violent impulse of the forcibly-relaxed Muscle the other way.

If further it be demanded, by what means it so happens that in the Instance before us of an Arm bent by force, that the refluent Ani-

mal

mal Fluid is rather towards the Muscle, which by that means then proves contracted, than towards any other whatfoever, to all which it may indifferently have access, I think the Solution feems not difficult, if it be consider'd, that at the same time that the one Muscle is forced from. the other is forced into a contraction; from whence it lo falls out, that the carnous tubulous Fibres of the last, which by being extended under the state of relaxation, did loe their cavity, must needs by their natural elasticity, when freed from the preponderant force of its Antagonist, acquire it again, by which means a space being made, the repelled Fluid, by the Laws of Libration, (to lay nothing of the habitual motion of the Animal Spirits, or Liquor, by most Authors, especially Borellus, urged as a Reason for this effect) must needs be driven this ther.

In fine, though I am not averse to think most of the Phænomena relating to Sensation and Motion may be solved by this Theory, tho of so small an apparatus, yet I am so

far

112 The Anatomy of the Brain.

far from being fond of it, that I have referved a far greater share of Friendship for any other that may seem but of never so little more a kin to Truth, and submitting all I have said on this Subject to the candid Sentiments of the more judicious Proceedee in describing the other parts of the Brain as they offer themselves in the usual modern way of Dissection.

MARKET OF STOCKER TOOKER

ding to as ourward, and next to its

Oscwardly its convex and corri-

Diese Mater called Fals, from the body Frocels called Crifes Gall for were landerment pare of the Crammar where there the two Dienisons are frontliked over the care from factor which pare all the care fedity to a test the factor following the factor of the factor of the factor of the factor following the factor of the fact

and to descript on CHAP.

in eard appearance.

and or CHAP. XIII.

be distributed the little filmed

Of the Brain in particular.

ferib'd and consider'd in general, as consisting of two different Substances commonly called its Similar Parts, and the Source of all Sense and Motion, comes now to be taken notice of in a more particular manner, with respect to its dissimilar parts or conformation; and this I think may best be done first according to its outward, and next to its inward appearance.

Outwardly 'tis convex and cortical, exactly divided into two Hemisipheres by the first Process of the Dura Mater called Falx, from the bony Process called Crista Galli forwardly to the very hindermost part of the Cranium, where these two Divisions are stretched over the Cerebellum, from which part also 'tis perfectly separated by the second Process of the Dura Mater, to the end it

ma

The Anatomy of the Brain.

14

may not cause any prejudicial compression upon that part, either by its weight or pulsation was (alaste)

The foremost Division is made only as deep as the Corpus Callosum, the latter to the very Medulla Oblongata it self-manual visitabin bus even

Tis further imperfectly divided into four Lobes, two whereof (which being the less) are forwardly, and two (which are much bigger) backwardly.

These Divisions appear best in the inverted or Varolian Dissection, being marked out as it were by four Branches of the Carotid Artery, two before, and one on each side.

These I call Impersect Divisions of the Brain, because though the Pia Mater runs betwixt them, together with the aforesaid Branches of the great Artery, yet they adhere by several Fibres, both of that Membrane and the Blood-vessels them-selves.

Tis also imperfectly divided thro' all its external cortical part by the Ria Mater, though not so prosoundly, to the end the Blood-vessels may penetrate this part in more fine and recicular Ramifications; and that by the

the pullation of the Arteries the interjacent cortical Glands, (or rather Vessels) may more freely make their Proper Secretions, I flomesof ad I

Nextly, it may be confider'd in its inward appearance, which is concave and medullary, taking its original from the Extremities or Apices of the Medulla Oblongata, (or rather a little more forwardly from the foremost part of Vieussenius's oval Center) commonly called Processus Lentiformes, or according to Dr. Willis, Corpora Striata. W 11 28 100 bestrem

From hence 'tis presently reflected back on each fide in the form of a Vault, very near as far as the Nates and Testes, a little below which on each fide 'tis joyn'd with the Crura Medulla Oblongata on their under fide, being continuous there to those Parts commonly call'd the Crura Forbrane, and the Blood veffels taisin

The middle and uppermost part of this Medullary Substance, by the Ancients always called Corpus Callofum, is therefore by Vieusenius cal- vieus per led Fornix Vera, in his Opinion par. 1. fustaining that Office (though I see not that it does, or for the Reasons I 2 before

116 The Anatomy of the Brain.

before given in the description of the

Dura Mater and its Processes, needs

to do any fuch thing.)

This is that part which, as was before noted, was thought (but mi-flakenly) by Vefalius and others to escape the covering of the Pia Mater and in it are not visible any bloody Specks, as in most other parts of the Medulla Cerebri.

Tis the medium uniting the medulary part of each Hemisphere or Division of the Brain, samous for the transverse Stria running through it from each side of the aforesaid Hemispheres, the Septum Lucidum

only coming between.

In this large or principal Cavity are contained the three Ventricles, the Fornix, the Septum Lucidum, Corpora Striata, Thalami Nervorum Opticorum, the Roots of the Fornix, the Tractus Intermedius of the Corpora Striata, the Tractus Medullaris Thalamis Nervorum Opticorum Interjectus, (which last has bin already described) the Vulva, Anus, and Rima or Passage to the Glandula Pituitaria by the Infundibulum, and Glandula Pinealis, (which also hath already been described) of all which briefly in their order. The

The three Ventricles, by cutting FIG. 5, AA asunder the Fornix near to its Roots, and turning it backwards over the The three Nates, Testes, and Glandula Pinealis, Ventricles, appear to be but one, those on each fide it being called the Laterales, in which are the Corpora Striata Thalami Nervorum Opticorum and Crura Medulla Oblongata, that Rima, fo far as 'tis covered with the Fornix and parts the Crura Medulla Oblongata, being the third.

From the extream Limits of these viens.T.10 two fide Ventricles, from before to AA, &c. Centrum behind, does arise that medullary space called by Vieussenius, Centrum

Ovale, in his Opinion the great Dispensatory of Animal Spirits, the

Ovale.

fore part whereof Willis calls Limbus Will. de An.

anterior corporis striati. T. 8. E.

Tot Fornix. The Fornix is a medullary part Fig. 5. AA, arifing from two Roots in the foremost part of the Basis of the Brain, lying betwixt and upon the uppermost parts of the Thalami Nervorum Opticorum, which Roots come out of the foremost part of the Geminum Centrum semicirculari, so called by Vieusenius, like two large I 3 Nerves,

Nerves, and afterwards joyn toge-A ther, constituting a broadish medullary Body, which after having first projected it felf for some space for wardly betwixt the Corpora Striata, and afterwards run the length of the third Ventricle, growing all the way broader and broader, and towards its edges (by Vieussenius called Fimbra) Vieussen. thinner; and being reflected backward Tab. 6. D towards the hinder part of the lateral) Ventricles, like two Arms, commonly called Crura Fornicis, the beginnings whereof on each fide are by Auran- Aurant. tius called Hippocampi and Bombyces, Anat. Obs. (from whence, I know, he had P. 45. chiefly observ'd this part in Brutes, in which, by vertue of the hinder part of the Fornix, in that place growing fomewhat thicker, and running over the hinder and upper parts of the Th. Nerv. Opticorum, which are more prominent in them, as in Sheep, Calves, Ge. than in Men) it is made to appear on each fide like the bending Creft of the Sea-borfe, and is in colour much like the Silk-worm, certain minute Malp de Stria's, which Malpighius calls Fi- cereb. p. s. bra, croffing them like Rings obliquely, contrary to what the fame

Au-

Author's Account is of them, who fays those Fibra or Stria run upon them otherwise, wiz. as they do on the Septum Lucidum (i.e. longitudinally) and embracing the Th. Ner. Open on their upper part on both fides, but adhering close to them as one continued Substance on their under part, (in which place they are called, by Wieusenius, Posteriores veri fornicis Vieus p.61. (viz. Corporis Callos) Columna) becomesthere continuous with the hinder part of the Corpus Callosum, where it winds down upon the fides of the Crura Medulla Oblongata, and Toid. makes up that undermost space or cavity of the two fide Ventricles, by the faid Aurantius called Ventriculi Hippocampi or Bombycini, and Vieussenius called the hinder part of the Centrum Ovale, which by that kind of curved passage loses something of its oval figure and in manim thad in Men) it is made to appear on

The Septum Lucidum. The Septum Lucidum some of the Moderns think to arise from the Fornix, thence ascending to the internal Superficies of the Corpus Callosum; others from this last descending down to the Fornix, but most likely

likely from this last, where towards its foremost part I have always found it double, (first taken notice of by Sylvius de le Boe) and as Vieuffe- Sylv. dele nius truly fays, often with Water in Boe Diff. its duplicature. Os part mag floorish Med. p.19.

Tis a very thin, medullary, Thef. 13. transparent Body, intermediate to the Corpus Callosum and Subjacent Fornix, by means whereof the two lateral Ventricles are in that place

separated one from another.

The Corpos I 1, 600.

The Corpora Striata, or Processus Fig. 5. Lentiformes, are two Prominencies fituated fomething higher than, and in Men a great part of them on each fide (though Dr. Willis fays, where musos the Corpora Striata ends the Thalami Nervorum Opticorum begins, which is only fo in Brutes) of the Thalami Nervorum Opticorum, or Juga Crurum Medulla Oblongata, and are so called from the many white Streaks appearing in them, descending obliquely to the Medulla Oblongata, with Cineritious Substance coming betwist them when they are cut horizontally. He paragraphs to nound

They run down on each fide the Thalami Nervorum Opticorum as far as till the Corpus Callosum be-. gins to wind back upon the Crura Medulla Oblongata, towards the hindermost part thereof.

have got them delineated here exactly true, (tho' by neglect without the Striæ) finding all the Cuts of them in Willis to be from Brutes, except one, which is done very ill, and those in Vieussenius very false, unless in Figure the 8th, which also wants the Strice or Strice or Strice

38cd. p. 19

AL-TEG

entiformes, are two Promunencies The Thala- The Thalami Nerverum Opticorum mi Nervo- are two prominent Bodies, more corum. purely medullary on their outward Superficies than within, which meeting together like the two topmost Aroaks of a Y inverted, constitute the uppermost part only of the Crura Medulla Oblongata in that form, the other or undermost side being quite of another figure; and feeing they are the immediate continued Productions of the Medulla Globofa Cerebri, (which contrary to the old Opinion of Praxagoras and Philotimus, afferting the Brain to be only a Germination

mination of the Dorsal Marrow, of late reviv'd by Bartholine, (if any prece. Caufab. in dency of Parts as to time may be al p. 137. low'd) I look upon to be rather the original than the production of the Medulla Oblongata and Spinalis (00) and may more properly be called Capita than Crura of the Medulla foremost part meeting toget. John and

Wun

The Tops or Juga do, as already F16.5. CC. observed, encline close, yea, joyn together, as Vieusenius hath rightly observed contrary to Willis, (whose Figures of that part are utterly false) unless where the Rima ad Infundibulum parts them, leaving like the Corpora Striata an obtuse angle their external medulla med ninds

> Betwixt these two last mention'd Bodies there is a medullary space on each fide, which in a bending manner encompasses the Thalami themfelves, and receive the Extremities HALL COL of the Striæ in the Corpora Striata, and del as they descend from the aforemention'd Centrum Ovale, and is therefore by Vieussenius called Gemi- Vieus. p.67. num Centrum Semicirculare, by Willis par. 2. Willis de Limbi Posteriores Corporum Stria- An. Brut. P. 42, T.8. torum

> > The

The reason why they are called of dalami Nervorum Opticorum, is from certain Fibres supposed to be in them, arifing both from their true medullary Superficies (by Vieussenius call'd a Medullary Membrane) and fome from within their own Substance, which at last, towards their foremost part meeting together, make up the Bodies of the Optick Nerves.

Willis says nothing of these Fibres, though in his Opinion Galen did not improperly give them that name. Vieusenius paints them very falle) unless where the Rima gnorth

lary Trast.

As for my part, I never could find any Fibres at all appearing in their external medullary part, those within are very small at best, and fcarce discernable a si ayod soiboll

On the outside of these I have always found and often showed a very A Medul- fair medullary Tract, here descri-Tab.5 mm bed, running all-along betwixt the Corpora Striata, & from the very hindermost extent of the Corpora Striata forwardly, down to the very Roots of the Fornix, to which they feem

With-

The Paffage Within this Cavity of the Brain into the In- are likewise two passages into the fundibulum Infundibulum, and so on to the Glandula Pituitaria, the foremost of which is called by the odd Name of The

hima, or just under the Roots of the

The Vulva, of Vulva, and the hindermost of The Anus, from their fituation, which with the Rima betwixt them, is called, as was before noted, the third

Ventricle.

ramina.

The places whence all this Water iffues are commonly by the latter Anatomists described under the name of Tria Foramina, situated so as to give passage from all the eminent Regions of the Brain, from whence there can be access had to them for the Water (or rather the Lympha, properly to called) to fall into the aforefaid Infundibulum, the first whereof is behind the Testes, under the Valvula major, (hereafter to be described) the other just under the Pineal Gland, or the beginning of the Rima, which two meet in an Aperture, under the Nates and Testes, by Vieus- Vieus.p.73. Senius call'd Aqua Emissarium, having par. 3. a steep descent into the Infundibu-

lum; and the last at the end of the

Rima.

Rima, or just under the Roots of the Fornix, and all ending at length (tho' by two different passages) in the Infundibulum. of bus mulndihan

The Nates It may not be unseasonable in the and Testes next place to take notice of two remarkable very fair Processes, called Nates and Testes, by former Anatomists so named from the resemblance they had to those parts; but it is plain from thence they were only used to dissect Brutes, in which they have fuch a proportion as is betwixt them; whereas in Men tis plain they are very near of the fame fize, and not very different in form, being oblong and accuminated towards their Extremities; but in Sheep, Calves, and most other Creatures the Nates are round and large, and the Testes oblong, somewhat accuminated, and very small.

> Before these Natiform Procesfes, under the Glandula Pinealis, runs a transverse Process before taken notice of Pag. 84, by Vieu Jenius called Vieuffen-Processus Natibus Antepositns, and Tab. 8 f Nervuli Æmulus, which upon further enquiry, by drawing the Thala-

mi Nervorum Opticorum still wider, appears to be rather Nervi than Nervuli Æmulus, being as thick as that behind the Roots of the Fornix, to which in situation 'tis just opposite, and feems to joyn the Thalami Nerverum Opticorum together, as that

does the Corpora Striata.

In what rank to place them 'tis hard to fay, as being neither proper Appendices to either the Brain or Cerebellum, properly fo called, and being divided from the Medulla Oblongata in some measure by an Interstice commonly called Ductus ad Infundibulum by the Moderns, but by the Ancients a Passage for the Animal Spirits to the fourth or noble Ventricle.

The Ifthmus.

They are fituated upon that part of the Medulla Oblongata which is between the Cerebrum and Cerebellum, which space was before called Isthmus, opposite to that part called from its Author Pons Varolii, and by many Authors, as Bartholine, Spigelius, Highmore, &c. thought to be the two hindermost Roots of the Spinalis Medulla, which much more likely Riolanus makes the Processes of the Cerebellum to be, and with him Vefalius the great Vesalius, who paints them p.766,767. From this intermediate fituation fig. 11. GG.

Dr. Willis thought fit to make them as it were an Intelligence Office betwixt the Cerebrum and Cerebellum, how rightly, I refer to the Judgment

of others.

'Tis certain they are medullary Bodies, and contribute to the making the Animal Fluid or Spirits fo called after the same manner as the rest of the Brain does; for in cutting them through, (after having taken the reticular expansion of Bloodvessels off from them, which is very large here, and eminently conspicuous in injected Brains) I find them of the very same substance with the Processus Annularis and the Thalami Nervorum optici, partly cineritious, and partly medullary, and in fresh Brains somewhat, but very faintly, striated.

I know not of any part within the Brain, properly so called, that is not already described, except a certain Medullary Chord at the end of the third Ventricle, and the Valoula The

major.

The first of these is a Medullary willis p.43 fura Craf- Process, which joyns the Corpora col. 2. wi E nula Striata together, according to Dr. of Vieuge- Willis, by Vieugenius called Comiffura Crassioris Nervi æmula; and according to him is the Medium or Commissura by which his Geminum centrum semicirculare intervening between the two Corpora Striata Superiora anteriora & posteriora, and his Tractus medullaris transversus & obliquus intervening between his two Corpora striata inferiora anteriora and posteriora, have a communication with each other.

Dr. Willis places this Chord or willi, p.6. Commissure under the Roots of the col 1. Fornix, but it is always behind it, tho' contiguous to it.

The Valvu-

The second is the Valvula mala major. jor, so called by Vieussenius, but vieus. p. 76. plainly enough discovered by Dr. Wil- Willis, p.49 lis long before, and its proper use col.2.par.2 described.

> It is a thick (especially in Men) medullary Membrane, adhering forwardly to the inferiour part of the Testiforme Process, a little behind that transverse medullary Process from whence the pathe-

tick or fourth Pair of Nerves arise, laterally to the Process ascending from the Nates to the Cerebellum, on its hindermost Expansion, to the foremost Vermicular Process of the Cerebellum, and no where that I know of to any part of the Pous Varolii, as Vieussenius will have it, (who vieusen. seems to have mistaken another part p. 76. for that Process) unless just where Id. p. 73. the fecond Process of the Cerebellum comes out from thence, which jointly with its fellow Process on the other fide, when they meet together, after their transverse descent on the backpart of the Medulla oblongata, do really make up that part which by Willis is call'd (and that no doubt from Varolius) Protuberantia Annularis, and by others, from its true Author, Pons Varolii.

By raising up the foremost abovemention'd Vermicular Process of the Cerebellum with the Finger, it rarely fails to come in sight; but if not so, 'tis easily shown, by blowing into the Foramen situated under the Pineal

Gland.

Its use, according to Vieusenius, vieus.p. 110 is to hinder any part of that Water par. 2. which falls into the hindermost Fo-

K

ramen

ramen behind the Testes, from running into the fourth Ventricle, or Vice versa from the fourth Ventricle into it, or from getting out on each side of the Medulla oblongata, over the afore-mention'd Processes, so as to fall down upon the Nerves arifing thereabouts below from the Medulla oblongata: Which last use is evidently most true, (whether it be understood of Water preternaturally or accidentally collected there, for I must needs confess I could never find any there, any more than I could in the third Ventricle in Subjects free from those Diseases incident to that part, as hath before already been remarked p.82) but as to that relating to the passage from the Cerebellum to the last or third Foramen, I much doubt the Truth of it, for many Reafons, of which this is one, viz.

That the Plexas Chorocides in the fourth Ventricle, together with the adjacent Parts, being of the same Texture as the other are in and about the two lateral ones of the Brain, renders it as reasonable to suppose that Water may be collected there as in other parts of the Brain, (nay, that it is so, he himself also allows as Matter of Fact) and consequently as necessary

ceffary to have a place of vent for the Water whenever it happens to gather there, as it was for that which was at any time got into the other Ventricles. And confequently,

In the next place, I do not fee how this tender Film can be able to intercept a passage of so searching a body as Water at any time forced against it (not withstanding the supposed declivity of this Part, which in Man, by reason of the largeness of the subjacent prominent annular Procels, is very inconfiderable) which by Pulsation must needs happen whenever we suppose that Cavity filled with it.

And, in the last place, notwithstanding all the Contrivance the aforefaid Author hath shewn in conveying the gross part of the Water (which, as was before noted, he grants may be, nay, constantly is deposed there from the Glands of the Plexus Chorocides here situate) by the Extremities of Views p. 111 Veins, out of this Ventricle, I am suspicious, if there was no spedier reductory passage found out, there would frequently happen very great Mischiefs to the Medulla Spinalis it felf, and the Nerves springing from it, seeing the Extremity of that Ventricle called the Calamus Scriptorius is there

parted from the Spinal Marrow behind it, but only by the Pia Mater, which notwithstanding it is there double, as it is also quite down the whole Spine, lest perhaps the Water should fall down upon the Nerves which arise from it teo readily, yet upon such an occasion may be casily supposed subject to violation. Not to say any thing of the high improbability of any such Conveyance at all by the Veins, seeing that in a natural state they are always, as hath been already observed, conti-

nuations only of Arteries.

'Tis true, this may hinder the fall of Water into the fourth Ventricle, by realon of a Passage under the Nates before mention'd, by Viensfenius call'd Aque Emissarium, to near at hand to receive it when it finds its further palfage that way obstructed by the interpolition and relillance of this Valve. And for the same reason doubtless it was, that in Vieussenius's wall at Experiment which he brings for a Proof of his Opinion, no Water was view p. 110 found in the fourth Ventricle, it has par. 2. ving got a passage immediately, upon its non-admittance by that Valve, to convey it another way, which by reason of the steepness thereof, is done much more readily.

CHAP.

bind it but only by the Pia Mater. which not with flanding it is there dou-

bluodi CHAP. XIV.

tall down upon the Nerves which arrie na of the Cerebellum. moul occasion may be cally supposed subject

to violation. Not to lay any thing of

HE Cerebellum falls next in order to our confideration, in describing of which I hope a great deal of pains may reasonably be spared, seeing all that hath been already spoken of the cortical or cineritious part of the Brain, as also of its medullary part, is equally applicable to the Cerebellum. Nor is what hath been faid already of the Plexus Choroeides in the Ventricles of the one part less applicable to that Plexus in this. And for the fame realing

The Plexus This Plexus Chorocides in the Chorocides fourth Ventricle begins to be glanbellum. dulous just under the Eighth Pair of Nerves, from whence it runs up on the fide of the Caudex Medullaris to the chordal or third Process of the Cerebellum, and from it enters the fourth Ventricle, by Aurantius called Aurant. K 3 Cifterna obs. p. 48. Cisterna Spirituum, (which Ventricle, conformably to what that Author hath in the aforesaid place obferved, I always find broader than long, and double, though not divided by any intervening Body, as the two lateral ones of the Brain are;) not lying loofe therein, nor at the bottom of it, as the Plexus does in the Ventricles of the Brain, but quite contrariwife, (and which hath not heretofore, as I know of, been taken notice of) adhering close to the top of this Ventricle, or the bottom of the superincumbent Cerebellum, then running transverse just at the end of the Calamus Scriptorius, there becomes continuous to the Plexus of the other fide; as hath been observed of the Plexus in the lateral Ventricles of the Brain.

This Plexus arises from a ramisication of the second or backwardest Branch of the Cervical Artery, as one part of the other Plexus of the Brain mention'd in that Chapter where the said Plexus is treated of, doth) and another smaller Branch of the said Artery about the place where it ascends from the Vertebrals,

to a reticular Expansion first, and then a little space further meeting with the other constitutes this Plexus.

This part differs from the Brain in its cortical structure, inasmuch as its Interstices are here eliptical or pieces of imperfect Circles, growing shorter towards those two Productions of the Cerebellum, before and behind, (which by reason of certain annular depressions occasion'd by Bloodvessels there embracing them, seem as tho they were wrinkled like Worms, and therefore called Processus Vermiculares) as Parallels upon the Globe do towards each Pole.

The three Processes of the Cerebellum.

It hath three Processes, which joyned together on each side, make up as it were two sair Roots, according to the Ancients called the hinder Roots of the Oblongata Medulla, by the Moderns Peduncles or Stalks, by which this part grows to the Medulla Oblongata.

Fig. 7. gg The first of these ascend from the Cerebellum to the Nates, the se-

Medulla Oblongata, which meeting together on the under fide thereof,

K4

as was before noted, make up that large Protuberance by Willis called Var. Anat. Processus Annularis, by others from the first Author Pons Varolii.

This I find full of Stria's or medullary Tracks, much fronger and Fig. 6. cc larger than those of the Corpora Striata, running transverse on each fide the length of the whole Procels, and terminating in a medullary long Tract, dividing that Procels into two equal parts, as you Ib. ce fee in the Figure, the use whereof, as having never been before observed, will be hereafter taken notice of.

The third descends from this part backwards, upon the upper fide of Fig. 7. hh the Medulla Oblongata, like two longish thick Chords on each side, making the Medulla look fomewhat thicker and broader in that place, and not unfirly fliled the Chordal Process. and of Parson ravo this ste

These Stalks, when they joyn together at the other end, make up the Meditallium or Corpus Callosum of the Cerebellum.

The tranfwerfe Procef of the fourth Ven-Bricke.

p. 26.

ned are alittle above the original of the There are two or three fair medullary Processes close to, and sometimes

Medulla

riding one over another, a little on this fide the fourth Ventricle, or about the beginning of the Calamus Scriptorius, which joyn the two ceffes together that descend from the Cerebellum to the Medulla Oblongata; and there are two more deteending length-way from that other transverse Process behind the Testes, down to thefe, guitanimiat bus

inside of the Medulla

Thefe long medullary Processes I ceffes on the never find wanting, though in different numbers, fometimes having feen Oblongata: three, sometimes two, and once I oould find but one, (though larger than ordinary) and constantly, in what number foever, ending in the transverse Processes at the afore-mention'd beginning of the fourth Ventricle.

lary long Traft, dividing that Pro-

These long descending Processes are just over-against the Corpora Pyramidalia, on the other or under fide of the Medulla Oblongata, and the transverse Processes at the beginning of the fourth Ventricle last mentioned, are alittle above the original of the Eighth Pair of Nerves, infomuch that without being very circumspect one

may mistake them for the original of that Nerve, whereas in reality I find them to be the original of the loft or hindermost Branch of the Seventh, as will be more particularly taken notice of hereafter, in the description of those Nerves; and therefore cannot but wonder how Dr. Willis (who Wills Ce speaks in one place as though he p 12. col 2 had feen them) came to affign them

for the Root of the ninth Pair, beneath which and this Process I have always observed the space of already been looken chan in alleh in the next place that we take no-

rice of it on as other fide, where are

led, which Crura are only the un-

der nert of the Thatems Nergaran

Crura Me most considerable its Cruras so caldulla Ob SOURATE.

Optioning before described, which in their Extremities becoming continuous to the under fide of the medullary hinder nort of the Brain,

anuti or ameiona ent b CHAP.

the Brain there, as it had its, him dermost from the Properties of the Oxedellam tout room is more diligent enquiry it appears; that mofe

dulla Oblongata nad use

CHAP. XV.

nay anticke them for the original of that Morvey whereas in reality I find

Of the Medulla Oblongata. confer the west wandernereiore cane

nor inniwonily how this Walle (who

HE third part of the Brain, in its general acceptation, according to the foregoing method, is called the Medulla Oblongata, all whose parts on its foreside having already been spoken of, it remains in the next place that we take notice of it on its other fide, where are Crura Me- most considerable its Crura, so called, which Crura are only the under part of the Thalami Nervorum Opticorum before described, which in their Extremities becoming continuous to the under fide of the medullary hinder part of the Brain, Ancients to think occasion'd the the Medulla Oblongata had its foremost Roots immediately from the Brain there, as it had its hindermost from the Processes of the Cerebellum; but upon a more diligent enquiry it appears, that these

dulla Oblongata.

Crura are more deeply immerged in and knit to the Medulla Globofa of and anog the Brain forwardly, by vertue of the Corpora Striata, as also by the very medullary part of the Brain it felf, which there, from the back or undermost winding part of the Corpus Callosum is perfectly mingled I Pair of Nerves begin ti diw

Where these two Crura begin to come close together, the Protuberan-Fig 6. BB tia Annularis, or Pons Varolii, made up of the second Process of the Cerebellum aforemention'd, begins to cover the Medulla Oblongata for about the space of an inch and an half, after which this Medulla Oblongata in one large Trunk is continued to the first Vertebra of the Spine, and so quite down to the end thereof. Werling to the thereof.

with the Corpora Pyramidalia and Whilst the Brain is in this position The two white Brdies behind it may not be unseasonable to take the Infun- notice of two fair white Bodies on dibulum. Fig. 1. bb this side of the Infundibulum, in that depressed part of the Brain, where the Pia Mater (as hath before been taken notice of) is fo remarkably double.

There

pora Oli

Waria. Ebid: 0

There are also two white long pora Pyra- medullary Processes called Corpora Fig. 1. n. Pyramidalia both by Willis and Vieussenius, which arise just at the ending of the Annular Process running down upon the Med. Oblongata the space of an inch, ending a good space below the place where the Eighth Pair of Nerves begin, which have their original between the Corpora Olivaria and the Chordal Processes partly on the other fide thereof, contrary to the account we have of them by Dr. Willis, who describes them as Willis, p. 13. ending in pointed Extremities, just col.1.par.1 p.61. col.2,

after which this Medulla Ob-

where those Nerves have their original, par. 3.

The Corpora Olivaria. Ibid. o.

On each fide of these appear plainly the Corpora Olivaria, fo called from their Figure, as the former were by Vieusenius, which with the Corpora Pyramidalia and two white Bodies behind the Infundibulum, he calls Conceptacula Spirituum Animalium, or places containing Animal Spirits upon feveral occasions of use to the Brain, both in its natural and intellectual Faculties, to some notice and selection

THAP double.

CHAP. XVI

Crura Medulla Oblongata, where they advance on each fide into the Globous

whence running concealed berwixt to foremost the Nerves ob-

N the same position of the Brain we also have a fit time of taking a view of the Nerves, which are still medullary Productions of the Brain dispersed to all the parts of the Body, which have need of either Sense or Motion, and these are in number ten Pairs or Conjugations, having their Names and Originals as follows.

The first is the Olfastory Pair, which after they leave the former Lobes of the Brain, and begin to run to the Bone called Ethmoides, take the name of Processus Mammillares; but this is chiefly in Brutes, where through their largeness they have that appearance, and are manifestly hollow.

By the utmost Scrutiny I have been able to make, they have but one Original, and that is from the undermost and soremost part of the

Crura

advance on each fide into the Globous medullary part of the Brain, from whence running concealed betwixt its foremost and hinder Lobes obliquely, for a good space, at last they come in fight, as you see them in the Figure: And by what means Vieusenius comes to find such disfused Originals for them as he speaks of Iknow not.

Their Use is known to most, and a particular account thereof, as of the rest, together with the manner of Sensation, with relation to the external Organs of Sense, is much more sit for a Physiological Tract than one of this kind.

I shall therefore only at this time give a general description of the Nerves belonging to the Brain, how and where they arise, the difference or variety whereof serve very well to inform us, according to several late Theories, concerning the different Reservatories of the Animal Fluid or Spirits, and the different dispensation of the same to several parts of the Body.

The

Pair.

The fecond Pair are called the The Second Optick or Seeing Nerves, of which Ibid. 2 2. I find no more Originals than of the former, and that is from those medullary parts called Thalami Nervorum Opticorum, tho' Vieussenius says they are from feveral parts; and Willis in general terms from the afores faid Thilami Nervorum Opticorum, behind the Corpora Striata: which description is more exact in Quadrupeds, where the Thalami Nervorum Opticorum are altogether in situation behind the Corpora Striata, than in Men, where a great part of the Corpora Striata are fituated on the outfides of the Thalami Nervorum Opticorum, and only their Heads or Extremities before them.

> The Blood vessels mention'd both by Willis and Vieussenius belonging to these Nerves, I have seen to run not only upon or with them, but also in injected Bodies exactly quite thro' the medullary substance of them, into the rericular Coat of the Eye, wherein they end in an infinite number of the most capillary Ramifications, which by an injection of that Artery

> > made

made with Mercury, become very delightfully conspicuous to the Eye.

The Nervous Fibres also, from the fifth and third Pair of Nerves, do twine about the Bodies of these Nerves, as the two above-mention'd Authors do truly affirm, but how rightly they both assign to them the office of dilating and contracting them subserviently to the visory faculty, and preternaturally in Convulsions of the Eye, as though these Fibres were truly Muscles, or of the carnous kind, I refer to the Judgment of others.

These go out of the Skull at its first Foramen.

The Third

The third Pair arise out of the Fig. 1833 forward and upper part of the Annular Process, where 'tis contiguous to, and covered with the under part of the Thalami Nervorum Opticorum, coming out into fight from between them, just where that Process terminates forwardly, which is where the Crura Medulla Oblongata come together into one body, constituting the Caudex Medulla Oblongata.

by an Intellign of the Amer Thele These running through a duplicature of the Dura Mater, on the outside of the Circular Sinus, go out of the second hole of the Skull to the Eyes, and are therefore called Par Oculorum Motorium, to the voluntary motion of which only they are granted to be subservient, which, seeing they have their original from the Cerebellum, afford us no weak Argument against the Hypothesis of Dr. Willis, who hath reserved that part in Nerves subservient to involuntary motions only.

The Fourth Pair. Ibid. 44.

The fourth Pair is very small, coming from the transverse Process on the forefide of the Medulla Oblongata behind the Testes, first coming in fight between the undermost part of the hinder Lobe of the Brain and the Cerebellum laterally, croffing that part where the Annular Process ends towards the Crura Medulla Oblongata, from whence they pass into a duplicature of the Dura Mater, and afterwards, a little more outwardly than the former, goes through the same fecond hole to the Trochlear Muscle of the Eye, and are called from their momoving of that according to the paffions of the Mind, the Pathetick Pair.

The Fifth Pair. Ibid. 5 5.

The fifth Pair is broad and large, made up of many thick Fibres continuous to each other, some softer than others, arifing from the uppermost part of the Processus Annularis, which is backward laterally, where 'tis broadest, by reason of the second Process of the Cerebellum there entering it.

The several the fifth

Pair.

This Nerve, after having first Branches of climb'd over the inner Process of the Os Petrosum into a kind of a Cavity made of a duplicature of the Dura Mater in that place, immediately fwells into a kind of a thickness, cal-Fig. 3. B led a Ganglion, from whence several Branches are propagated, lying betwixt the Dura Mater and the Cranium, on each fide the Sella Turcica. without any Fovea or Cavity at all, going out of the Skull at three several places, its superiour small Branch at the second hole with the third and fourth Pair of Nerves, its inferiour smaller Branch at the third

hole, and its posteriour or largest

Branch at the fifth.

FIG. 3. C, D, E

> L 2 From

From the infide of the foremost FIG. 2. y. Branch two little ones turn back, and meeting with another small Branch a little lower turned back also from the fixth Pair, where that Nerve is fasten'd to the outmost or borrowed Coat of the Carotid Artery, make up a small Trunk of a reddish or fleshy colour, like to that which 'tis of when passed out of the Cranium, (as Veslingius hath truly observed, who calls it The Intercostal Pair. The Internal Branch of the Sixth pair) x. z z. ' which descending obliquely, and creeping under that Artery, betwixt its external, proper, and borrowed Coat, goes out with the Carotid Artery at the fourth hole of the Skull, which is in a manner double between the Os Petrosum and Cuneiforme, and from its passage through the Thorax, near

The fixth Pair. Fig. 1st. 66. The fixth are about the bigness of the third, arising from the hinder part of the Annular Process overagainst, and not far off from the beginning or head of the Corpora Py-

which, it receives a Branch from the Intercostal Nerves) is call'd, The

Intercostal Pair.

pyramidalia. It sends out sometimes one (in this Subject very short) sometimes two slips, as was afore said, for the making up the Trunk of the Intercostal Nerve, and after that (with the foremost Branch of the sistent Pair, in one and the same duplicature of the Dura Mater, together with the preceding third and fourth Pair of Nerves) goes out at the second hole of the Skull, and terminates in the abductory Muscles of the Eye.

The seventh Pair. Fig. 1st, n

Nerve is large, and comes out almost just over-against the original of the fifth Pair, on the lower or under side of the second Process of the Cerebellum, where it first appears coming out from the Cerebellum to make the aforesaid Protuberantia Annularis between the Corpus Olivare and that Protuberance, as though it crept out betwixt them, and had (as it really hath) a more remote extraction.

It confifts of two distinct Processes, the first of which is more round, hard, and less than the second, that being for Motion,

this for Sense, but tho' they seem as though they had the same original, being feemingly continuous at their rise from the Brain, (which Dr. Willis affirms they have, tho willis, p. 12 sometimes he makes it in one place, & p. 56. and fometimes in another) yet upon col.1-par.4. a further enquiry it does appear otherwife, the first or hardest having its original from the Caudex Medullaris, not far from the place where it comes first in view; the second very remote from the transverse Process or Procesfes in the passage to the fourth Ven-Fig. 7.11 tricle before described, (which in

another place the same Author seems col. I-pat. plainly to have observ'd, taking it for the Original of the other Procels of this Nerve;) from whence it ascends all-along on the sides of the Medulla Oblongata till it arrives at the afore-mention'd place, where it first, together with the other Branch, leaves the Medulla, to pais out of it at the seventh hole in the Bone called Petrofum.

The eighth The eighth, or Par Vagum, arises Fig. 1688,2 very little beneath the feventh, but yet not from any part of the Annular Protuberance, but exactly in that somewhat hollow place betwixt the Corpus Olivare and third or Chordal Process, having numerous (I have counted ten or twelve) Fibres, but all continuous at their first rise, for its original.

This in a multitude of Ramifications is spent upon the Bowels, and goes out at the eighth hole with the Spinal Accessory Nerve, where the great lateral and the inferiour little Sinus's in the Basis of the Skull go out into the Internal Jugular.

To this eighth Pair about half an inch from its first rise, whilst it climbs upon or sticks to the Pia Mater upon the Basis of the Cere
Ibid. * * bellum, ascends a Nerve called Spinalis Accessorius by Willis, but long before him taken notice of, nay, painted and described, by Vidus Vidus Vidius, the original whereof I find to dius, p.93. be as far as the seventh Vertebral T. 18.

Pair, from the foremost and hinder most beginnings of that Nerve, not-withstanding Vieussenius confines its

L 4 original

part only.

This Nerve runs under the Vertebral Artery near half an inch on the fide of the Medulla Oblongata, at length, about half an inch from the beginning of the eighth Pair, leaves the aforesaid Medulla Oblongata, running obliquely upon the Pia Mater of the Cerebellum, to joyn with the aforesaid Pair, which it really does in that very place, though it part with it afterwards again.

The ninth Pair. Ibid. 9 9.

The ninth hath several (in one Body I counted seven or eight) pretty large Fibres for its original, very distant one from another, the first of them coming higher, from the very top of the Corp. Olivare; the next, and several others, are much less, a quarter of an inch lower; and the last much lower yet, about the ending of the Corpus Olivare, or beginning of the tenth Pair, with several others between the Pia Mater and subjacent Medulla Oblongata; but after all, its Trunk is very little, about the bigness of the Accessory Pair.

Thro' the Fibres of this Nerve there runs commonly a small but very visible Branch of the Verrebral Artery, at its original; as you fee in the Figure expressed by the Letter k on the right fide, going out at the ninth hole, together with this Nerve and the Vertebral Vein, which Vein Vieus.p. 163 Vieussenius mistakenly makes to go out at a tenth hole, for almuch as that is never found in Nature, neither need be, seeing the tenth Pair goes out at the last or great Foramen, by which the Medulla Oblongata passes into the Spine.

The tenth Pair.

FIG.I. k.

The tenth Pair, (which had it a Ibid. 10 10 double Original from each fide of the Spinal Marrow, (as all the rest of the SpinalNerves have) might much more properly be called the first Vertebral. inasimuch as that both a great part of its rife and egress is quite out of the bounds of the Granium) serving chiefly the Muscles of the Neck, it begins with three, and fometimes more, small Fibres lower a great deal, out of the Medulla Oblongata, almost an inch below the Trunk of the ninth Pair, and is about the fize thereof.

It

It goes out of the Cranium betwixt the first and second Vertebra of the Neck, making its passage through the Dura Mater from the Medulla Oblongata, about half an inch below the place where the said Arte-

ry comes in.

The Structure of these Nerves is consistent of many Fibrilla's or Stria's, a certain number whereof being first enclosed in a production of that delicate inward Lamina of the Pia Mater afore described and spoken of, makes up a Fasciculus or Bundle, and many of these collective-

ly the Body of a Nerve.

In these Fibrilla's or Stria's (betubulous and always turgid, as in so many Rivulets springing from the main Fountain the Brain, and from thence distributed to every respective part of the Body) is contain'd the Animal Fluid, by means whereof there is maintain'd a constant intercourse betwixt it and the Soul, and reciprocal acts of Friendship betwixt one part and another.

This Animal Fluid I look upon only as a Body confisting of very minute and flexile Particles, contain'd

tain'd in such a space as allows them a capacity of being agitated on all fides by vertue of the fubtile matter, or Æthereal Globuli they fwim in, by which means they are render'd capable of pervading the narrowest Channels of the whole Machine, provided its Orifice or Pore be adapt thereto, in contradistinction to those other fort of groffer Particles of Matter, which by reason of the narrowness and figure of the space they are to enter, do approximate so close, as to become contiguous in all their Superficies, whereby they become deprived of their former expansive agitation, which is always necessary to make a Body fluid, and like fo many small Filaments orderly disposed, do constitute the Inclosures or Coats of those Vessels the Fluids are contained in.

This Animal Fluid I conceive to be in a continual state of Transpiration, proportionable to the measure of its leisurely production, seeing no more necessity of ascribing any further Uses to it, besides those afore-mention'd, than I do to the

156 The Anatomy of the Brain.

the watery Humour of the Eye, besides its service to Vision, which is always in a state of fresh production, as by the Excellent Nuck's Nuck de Experiment is plainly manifest; and Duc. Aqu. yet, by vertue of Transpiration, Oculor. some way or other, though to us P. 109. not visible, without any inconveniency to that noble Organ.

CHAP.

CHAP. XVII.

Of Sensation and Motion in general.

HE Nerves thus constituted, become accommodated for Use in relation to their several and distinct Functions, in some consisting of Sence only, fuch as are those appertaining to the particular Sensories, (viz.) the Smelling and Seeing Nerves, as also the soft Process of the Hearing Nerve, some Branches of the fifth, and it may be of the ninth Pair, for Tasting; in short, all the Nerves belonging to those external Sensories, by way of eminency, and in a less eminent or general way all the Nerves of the whole Body, which are distributed to fuch Parts as by reason of their structure are capable of Sensation only, any of which, as furnish'd with the Nervous Fibrils, but more eminently the Cuticula, may properly be call'd

an Organ or Sensory of Feeling; in others of Motion chiefly, such as are all the whole System of Nerves, (excepting them only afore-mention'd) ses, which though in a less eminent manner, are nevertheless sensitive Nerves also: In others of both, in all respects (viz.) either in a more eminent or less eminent Sensation, and Motion too, with relation to the different Fibres they consist of in their Originals, as the fifth and ninth Pairs.

These two different Functions of Sensation and Motion are executed after two as different manners.

The first of which, being occasion'd from external Objects, is discharged by a pressure thereof made
on the Instrument of Sense, so that
the Motion is backward irom one
Extream of the Organ to the other,
where it terminates in the Commune
Sensorium, commonly so called, and
is therefore stiled Perception, Passion,
or Assertion.

The other is discharged by some manner of impulse upon the Organ from within outwardly, with a tendency either to acquire some Good, or avoid some Evil; by which Impulse, when carried on so far, either in a natural or moral sence, as to terminate in, or to be executed upon its proper Object; the Object then may be said to suffer as before in the other case it might be said to act, and the perceptive Faculty now to act as before it might be said to suffer, and this Action is commonly called Local Motion.

For whose sake, seeing 'tis of different kinds, learned Men have thought fit to organize or divide the Brain into two distinct Provinces invested with feveral Rights and Jurisdictions abating the Power of the Sensitive Soul, which before was looked upon universal over the whole Brain, allowing it only a principal, but no absolute Empire there: And this they have done upon no weak or unreafonable grounds, seeing that Local Motion is not only in many respects performed without its affiftance, but even against its power of resistance; as in the Pulsation of the Heart, vermicular Motion of the Bowels, and in a great measure the Act of Respiration.

Now,

Now, that which hath been taken from the Brain hath been conferr'd on the Cerebellum, to which, though some Power in this Affair may justly be allowed, as was before observed, yet possibly not altogether so

much as there hath been.

Dr. Willis, who is Chief in this Cause, having distinguish'd Motion into voluntary and involuntary only, hath made the Cerebrum accountable for the one, and the Cerebellum chiefly for the other; and to that end hath furnish'd it with the like number of Nerves, as in his own words is expressed, Ot divisum willis c.18. cum ipso (i. e.) Cerebro, imperium Cerebellum habeat; nay, considering the Intercostal Pair, derived from the sistent and sixth Pair, which belong to the Cerebellum, he hath made it exceed.

I am apt to think that Learned Person too soon fell in love with his first Thoughts, the ordinary reason of either ones seeing falle, or not far enough.

Retional Creatures, the one purely matural, fuch as is Pulistion of the Heart, and various contraction of the Proceeding from a certain por-

I I

STO13

Nothing being more apparent, than that most of those Actions or Animal Motions he calls Involuntary, and of which he gives so many Instances, are equally found in Brutes and rational Creatures too, whilst in the state of Infancy, as well as when grownup, with this only difference, that all of them in the last are under the controuling power of the Soul, and confequently may be suspended upon a reflex'd Act of the Understanding; whereas in Brutes and Infants they are necessary, and do as naturally ensue upon the impulse of the Object, as Water, when unconfin'd, runs towards a Plain.

Now, if all these were supposed to be under the power of the Gerebellum only in Brutes and Infants, the Brain it self must necessarily be thought altogether useless in them.

It will be necessary therefore to take notice, that there are two sorts of Animal Motion in Brutes, as in Rational Creatures, the one purely natural, such as is Pulsation of the Heart, and various contraction of the Viscera, proceeding from a certain portion

tion of the Animal Fluid continually dispensed to the Nerves in an
equal proportion, and so may be
said to have their cause originally co-existent with the Creature,
and always present: And this kind
we find by a most convincing Experiment hereafter to be mentioned,
to be from the Cerebellum, and absolutely free from the dominion of
the Brain, in its ordinary way of acting or influx.

The other is that of Instinct, relating to the Sensative Soul, or an aptitude of the Nervous Structure, to act according to the Impressions made upon the Nerves, either from within, or from without, and so may be said to depend on the presence of such Causes as are supervenient and extraneous to Nature, suitable to the impressions whereof the Animal either pursues or avoids the Object, obeys, or results the Im-

Now, I take it for granted, that no body will deny but that the Nerves (by vertue whereof these last actions of Instinct are performed) whether they arise from the Cerebrum or Cere-

bellum,

bellum, are equally under the command of the Soul; or elfe, as I faid before, the Brain in those Creatures is to no purpole the nave the soquuq on ot

And of this fort I reckon all those actions in rational creatures of Instinct before they have attain'd to the use of their Understanding, from any fort of Impressions, or inadvertent and inconfulted, when he hath the controuling power of Reason allow'd him and makes no use of it, such as are called Habitual, which at first were produced by command of the Rational Part only, but through frequent repetitions at last, without any command from that, out of a blind obedience to a bare impulse from the Object; or lastly, such as happen when he hath altogether lost the use of it, as in Sleep or Distraction; in which last Cases 'twill be very difficult to diftinguish him from a meer Machine or Automaton.

Now, from what hath been faid, I cannot but think it plain, that many of the Actions before spoken of in Dr. Willi's sence, by him called Involuntary, as proceeding M2 from

bellum.

from the dominion of the Cerebellum only, such as he calls the various Configuration of the Face, from some Impulse or Provocations in the Viscera or elsewhere, erecting the Ears, turning the Neck and Eyes about, Judden Shrieks and Outcries upon some extraordinary frightful Object furprizingly affecting one Sense or another, furnished with either fuch Nerves as he fuppoles to be altogether under the command of the Cerebellum, as the fifth and feventh, or elfe to have a very near correspondence with that part by vertue of Vicinity, as the ninth, do more truly progeed from that perceptive faculty, or (to use his own words) that part of the Soul, he hath confin'd to that part of the Medullary System called the Gerebrum, inalmuch as in realog nable Creatures they may and commonly are suspended, as well as the Nerves they flow from, fometimes made use of as Instruments of Voluntary Motion by it also; and to think the contrary, is as much Involunt M 45 procelam

as to fay, that when any body happens to express any of the aforemention'd involuntary Acts, or but hit his Bedfellow a box of the Ear, whilst asleep, all these must be allow'd to proceed only from the Organ of Involuntary Motions called the Cerebellum.

And of this kind also in a great meafure I reckon Respiration, concerning which I cannot easily be brought to think it fatisfactorily explain'd by Dr. Willis, from the Energy of those Animal Spirits which flow only from the Cerebellum in the Par Vagum, after the same manner they do to the Heart by the Intercostal and that Pair for its pullation, and as only under the command of the Soul, to be stopt now and then, as it pleases, by vertue of some Nerves communicated to the Intercoftal Muscles and Diaphragm. the chief Infruments of breathing, From the of . 1700 birde of the old sand Voluntary Motion by it allo; and

M 3

to think the contrary, is as much

Iam

1660 The Anatomy of the Brain IT

I am therefore rather enclin'd to bel think this Motion is of the other or different kind before spoken of, une 28 der the Title of Instinct, proceed in ing from ban extraneous superve-biv nient Gause, acting conformably wo to the course of Nature cimement other Cases of the same kindulas in to Hunger and Thirft, and theotike puo where the obtaining the defigned of Endoer Effect renders the part from W whence comes the Motion for viorne time inlenfible of the impression da and where, after the ceafing of the Effect or Motion, the fense of the impression revives again, whence there happens an equal reciprocasso tion between the Sense and Fruition, 18 or Sense and Motions mort noillargui

manner and reason of the Spirits acting upon the Stomach land Pa-os late in relation to Hunger and Thirst, to that of the Systole and Dia-stole of the Lungs or Respiration, twill be needful to take notice, that in an Insant unborn there is no Respiration, but yet there is a Cerebellum; and that if this sort of Motion cal-

318

led Instinct, which I make to differ from purely Natural Motions, fuching as are contemporary with even the first living Rudiments of the Indi- 196 vidual, was altogether and folely got owing to the Cerebellam, after the manner of that of the Heart; then of of necessity the Child in the Womb ought too respire. But being fatisauH fied of the contrary, it remains that W we account for its respiration another ther way, which is as afore noted. through the presence or absence of the first moving Cause or Impulseons which I make or suppose to be any thing impressing the Nerves, product pagated through the Organs of Breathing, for as to transmit the impression from within to the perceptive Faculty, prefiding both over the Cerebrum and Cerebellum too. to the tend the Spirits may from thence forthwith be commanded into fuch other Nerves as act those Muscles which serve for enlarging the whole Cavity of the Thorax, in order to let the Air into the Lungs more plentifully which was the thing aimed at by Nature; and thefe M4 are

168 The Anatomy of the Brain.

phragm. I and danger to Blood to

Now 'tis easie to conceive, that whilft the Child is enclosed in its Mothers Belly, there is not that occasion for Respiration as when 'tis born, the main Stream of Blood all that while finding no passage thro' them, and thats which does by the Ruy shian Artery made of Juices much more mild and cooler, the vo native wheat being dittle, and the Aliment meer Chyle or Milks from whence it falls out that the Polmonick Nerves go altogether unprovoked, which after birth are continually otherwise impressed or provoked by the hot Effluorums of Blood, new bred of ftronger Food, no and by a ftronger native hear, and wholly flowing through them swhich heat continually, as the Child acquires a greater maturity a Tencreavisting, may for lought disknow, anot a little contribute, by way of natural documpuliento its exclusionals ous as

clearly appear to any who will the more take the pains to confiden well of the flructure of Parts in Children

the

un-

- inunbornation who morthen usual circuit of Blood through the Lungs, which and blis wante for the same and beneather and bereits but hedeldhed mixture of Bladel and and Chale, is denoted; vals dalforthrough menthe Liveon terving chiefly for fepabootating athat gross Excrement the Gall, not bred brack deaft in any and proportion) in and Infant . unborn, and in lieu of thele, other Paffages, od (which become altogether unnecefad fary after birth) sprovided by Nature after a Thorter and more compendious way, (viz) by the Foramen Ovale betwirt the Vena Cava and Vena Pulmon and Tubulus Arteriofus between the Art. Pulm. and Aorta in the Lungs, and the Tubulus Venolus between the Sinus of the Porta and the Cava in the Liver 3 as hath been most no fagaciously observ'd by the late Learos ned Dr. Walter Needham oo seed

Tis true, That in several Creatures there are some Nerves very much depending on the Cerebellum, as are they which minister (though in a different manner, as hath already been taken notice of, and will be hereafter further explained) to the Natural and Vital Functions, (viz.)

170 The Anatomy of the Brain.

the Par Vagum and Intercostal Pairs, and therefore the aforefaid Author, pol who is in this as in many other of his Discoveries very fortunate, and highly commendable, maderalad very good guess when he brought in these Faculties into Subjection brown that part, inafmuch as obyoife my veral others, as well as by my own is Experience upon living Bodies, we up find, that notwithstanding most parcog of the Brain be pared off with a all Razor, yea, even after the Medulla no Oblingata be divided betwixt the Cerebrum and Cerebellum, and taken !! wholly out of the Cranium, wthem Heart will beat, whenat the same time if the Cerebellum it felf be but cut in pieces, though all the rest of the Brain be kept entire, the Creature expires prefently it to notification bus

Yea, I have feen Respiration (which is only in part depends on the Cerebel-ing lum) totally to cease upon only a sudden violent compression of that part by a blow, and, after its being wounded, the Heart to cease beating immediately.

ediately. The supervented to use the word

Bustlet

All

All which must of natural confequence fall out upon the Hypothefis to That those Functions of Nature do depend on the Cerebellum for their fource and influence, which is constant, uninterrupted, and out of the arbitrary jurisdiction of the Brain; yet with this wdifference, that in Motions purely natural, and either contemporary with the Embrio, as the first figus of its vitality, such as is Pulfation of the Heart, during its enclosure within the Mother, or Supervenient upon sits further growth and more visible organisation of Parts, w as the natural contraction of the other Viscera subservient to the offices of Protrusion of the Chyle, feparation of the Glandular Juices, and profcription of the Excrements, the Animal Fluid or Spirits do altogether flow from the Cerebellum, the Nerves there both descending from the Cerebellum, and terminating in those parts afore-mentioned ; whereas in Respiration, which I call a Motion of Supervenient Instinct (if I may be allowed to use the word Instinct 11A

Instinct in that sence) the Nerves descending from the Cerebellum, and propagated through the Lungs from the Par Vagum, serve only to convey the first impulse or impression of the Object to those parts which are by Nature framed and qualified to produce Respiratory Motion, and those are the Nerves of the Spinel Marrow. receiving the impression from the Cerebellum, sceing that by the aforesaid Experiment it appears plain, that as ter the whole Cerebrum was divided from the Cerebellum and Medulla Oblengata, the act of Respiration continued for a confiderable time entire, which Motion is dependent on the Sensitive Faculty presiding in the Cerebellum, transmitting the first Impulse produced by the eighth Pair or Par Vagum (as before observ'd) and communicated thence to those Spinal Nerves which act the Intercostal Muscles and Diaphragin.

Vagum, which is propagated thro' the Lungs, is to convey the Impression from thence to the Cerebellum, which by vertue of its con-

nexion

nexion with the Caudex Medullaria (from whence the Ancients rightly thought that part had its hindermost Roots from the Cerebellum, as before taken notice of) it is able to transmit it further, as the Sensative Faculty presiding there shall direct, and that too by the common way, the Medulla Oblong et a and Spinal Nerves.

And further; That this part is as capable thereof as the Cerebrum, and is not wholly and only deputed for the service of such Nerves or Organs as are employed by the involuntary part or portion of the Soul, (as Dr. Willis would have it) appears in that the third Pair of Nerves, by him allowed to be 2mongst the number of the other kind of Nerves, (viz,) those commanded by the Will, from hence (as hath been already shewn) hath its original. And here also furthermore give me leave to add, by way of conjecture, that the reason why the Soul hath not an equal command over those afore mention'd Nerves dedicated to the vital and natural Motions, is, the early date

or commencement of the office of those Nerves, by which means they contract an habitual irreliftible Influx, much less fo in those belonging to the Respiratory Functions, the mexercife whereof is of a later date : and laftly, the Influx is not in the least of habitual in those other subservient to the Organical Functions of the Limbs, inafmuch as they are not capable of being exercifed till a much longer time after, and then not fo uninterruptedly as either the first or the fecond, but gradually, and with intermissions.

So that the only reason why upon cutting the Cerebellum Respiration ceases, is, that by that means its structure is discomposed, and render'd unfit either to receive or transmit the impression further to the aforefaid Nerves, which are fubfervient to the Instruments of Reall spiration solution and garvom to

Tis true, there are reciprocal communications betwixt the Nerves of the Intercostal Pair, Vertebræ, and Diaphragm, yet seeing they terminate not immediately in the Parts of each others particular distinct jurisdictions, there is no interchangeable act or office from thence produced betwixt them.

For as, notwithstanding there are fo many Branches of Nerves communicated from the Spinal Nerves fublervient to voluntary motion to the Intercostal Pair, on their descent to the Viscera, and yet by reason of their not ferminating in those parts, they are not in the least able to bring these Nerves under the commands of the Rational Soul, by which provident Care of Nature it fo falls out, that 'tis not in the power of any, by misguided Reason, to act injuriously to themselves : So by vertue of feveral Branches reciprocally communicated from the Intercostal Pair in its passage down to the Viscera, to the Spinal Nerves, there is no power given to them of moving the Mulcles to which they are subservient uninterruptedly, after the meer manner of the of the Intervolut Pair, Verrellas, and

Diaphragm, yet feeing they termi-

But now, to return to where we lest off, in some Creatures it's very plain, that Nature hath extended this imperial residence of the Soul beyond the Cerebellum, even as far as the Spinalis Medulla, having not only put this last motion, but that of Pulsation too, under the jurisdiction of that elongation of the Brain; as appears in the samous Experiment of the Industrious Caldest upon the Tortoise, which after the Head was cut off lived, and carried its Shell about, the space of six Months.

Besides which, 'tis remarkable, (by way of digression) according to another Experiment by the aforesaid Author made upon that Creature, that after even the Heart and all the Viscera besides, were taken out, except the Lungs, that Creature (to use his own Expression) was found so to resist Death, as to turn it self from the inverted or supine position it had been placed in, in order to make the Experiment, to its prone or natural one, and to live and move six hours after. From whence

whence it appears, that Muscular Motion is capable of being performed by the Animal Fluid alone, without the concurrence of the Blood, by most Authors constantly hitherto Caldess. made to go a share therewith in the

performance of that action.

So that we find Nature hath not stinted it felf to one place for the Seat of the Sensative Soul, or Refervatories of the Animal Spirits fo called, in order to the discharge of the afore-mention d Functions. no more than it is at a los about the maintaining them in their Integrity by other ways, when it hath fo tallen out that the natural Aructure of the Organs, destin'd by Nature to that end, have utterly been defirov'd. of which we have many Inftances in the Anatomical History, those Functions in Several Creatures remaining perfect, where after death there have been found neither any Cerebrum or Cerebellum at all, or at least such as by their constitution was utterly render'd useless to any such end.

N OF

The truth of which is full more

mort

Of the first is an Instance of the Learned Wepfer, in a Child living fixteen hours after it was born, and discharging all the Duties of Nature that one of its age was capable of, and by the by (which all the patrons of a nutritious Juice by the Nerves may do well to take no- Mifc. Curiof. tice of) of a very strong and good habit of Body, whose Brain, after death, was found to be only an heap of Watery Bladders on Hydatides, except a small part at the bottom of the Skull, lying din a Sinus made in the Wedglike Bone, where the Pituitary Gland is commonly found confifting only of three Medullary Bodies, two of which being each of the bigness of a Kidney Bean, and the third behind them of a Pea only, from which indeed there did proceed fome, but very inconfiderable Nerves, de 100 nor Nervous Fibrils, but fuch as none can judge of a due proportion requifite to fatisfie the Exigencies of the common anatural, and dvital theless after death fourmoiningly

beilinghe

The truth of which is still more plain, and without exception, in another Instance in the Miscell. Med. Misc. Med. Physic. Gallic. of a Child living five Phys. Gall. days after it was born, whose Head An.3.P.54-had nothing but Water contained within the inclosures of the Dura and Pia Mater, without the least footsteps of any medullary part at all.

Parallel to which two last Instances, I had one communicated to me by that curious Anatomist and learned Person Dr. Tyson, in a Child born alive, with no more Brain in the Skull than what might lye in a Filbird-shell, the Medulla Spinalis being much larger than ordinary, as though part of the absent Brain had been squeez'd down thisther.

of the last (viz. where the natual ral conformation hath been depraved) there is extant an Instance in two several places of the Mission obs. 26.

Scell. Curios. in a fat Ox, in obs. 26.

which while living there were obs. & 130.

serv'd but very little signs of any fuch thing, whose Brain was nevertheless after death found wholly petrissed.

N 2 From

From all these 'tis manifest the Sensative Faculty is able to answer its internal or external Impressions, by one part as well as another, and that the Medullary System of the Spinalis Medulla may become as adequate a Sensory, in relation to the aforesaid Functions sometimes, as either Cerebrum or Cerebellum.

And as to the power or influence the Soul in general exercises over the Nerves, howfoever different in their original, feeing we have already observed what a provident care Nature hath taken for the preserving Creatures from their own violence, in that it hath not only constituted the chief Fountain from whence the great current of Spirits is derived, for the fervice of the vital and natural parts, by the Eighth and Intercostal Nerves, which is the Cerebellum, fo as to be free from the commands of the Rational Will in its ordinary way of acting, but bath alfo taken care that not any of those Branches which have their originals from Trunks, which are under the power of voluntary dictates of the Soul, should terminate in such Organs by which those Functions are discharg'd, (abare communication between Nerves of different Provinces not being sufficient to such ends or offices, as hath been observed in those afore-mention'd additional subsidiary smaller Streams of Spirits slowing to the parts confecrate to the natural and vital Functions by Branches propagated from the Spinal Marrow, to the Intercostal Nerve, all the way of its descent to the lower Venter.)

So we may further also remark, that as there are some manner of Impressions made upon the perceptive Faculty, after such fort of a manner as that it even lofes its power over its own Subjects, (viz.) the Nerves, which are subservient to its voluntary commands, as in Laughing, Sneezing, and libidinous Erections, the Organs by which these Actions are produc'd, being altogether under the power of those Nerves subservient to the voluntary dictates of the Soul, and acted after the very same manner as those of Respiration, as often as proportionable objects present, and (notwithstanding the assertion of Dr. Wil-

N 3

listo the contrary, who makes Laughing proper to Man only, and, by the authority of Aristotle, Sneezing an Affection proper but to few, it any other Creature, befides Man) might also produce the fame effects in Brutes, provided their Rupid Souls were capable of being equally impressed by such Objects as are proper for exciting a rational Laughter, as we fee they are by those producing the aforemention'd venereous actions, feeing the want of the Plexus Cervicalis, of the Intercostal Nerves, and two or three fmall Branches propagated from thence to the Nerve of the Diaphragm (which he calls a Disposition peculiar to Man, and consequently in his opinion the cause of that Affection in him) might be in a great measure suplied not only by that nervous Branch we find propagated from the inferiour Plexus of the Par Vagum (which Nerve is equally dependent on the Cerebellum, as the Intercoffal) to the third Brachial Nerve, from which the Nerve of the Diaphragm hath one of its originals, but also by that other propagated from the Thoracick Plexus of the Intercostal

Will pro6

costal Nerve it self, to the same aforesaid Brachial Nerve, into which the Nerve of the Diaphragm is inserted.

So, on the contrary, there are fome Impressions made upon the Soul sometimes, through which it acquires a power over those Nerves at other times in no wife subject to it, and those are the impressions either of great Joy or great Grief, suitable to which the Vital and Natural Faculties are made either much more or else so much less vigorous than ordinary, as even quite to languish.

How this comes to pais, according to Dr. Willis in favour of his own Hypothesis, and particularly in relation to the first, (which allows of no Involuntary Motions, but what come from the Province of the Cerebellum) is explained by suppofing an undulating or rowling motion of the first impression upon the Brain out of it again, through the Natiform Processes into the Cerebellum, and from thence by the Annular Process into the Intercostal Pair of Nerves, and so to the Nerve of the Diaphragm, (and he should, to make this way of explication en-Deoffal

entire, have taken in also all these Vertebral Branches inserted into the Intercostal Nerve, in order to the moving of the Intercostal Muscles, without which that action cannot be performed) by a correspondence between which Nerves and those of the Face, being all of one family, the aforesaid Gesture of Laughing is performed.

Now, befides the needlefness of bringing the Conceptions or Impressions of the Brain under a neceffity of being executed by the in-Feriour Province of the Cerebellum, till fuch time as 'tis proved, that fuch motions of the Spirits, upon extraordinary occasions, may rationally be granted, without supposing a regular motion of the same through fuch supposed Passages leading from one Part to the other at all other times, (the allowing whereof does necessarily imply a capacity of the Soul to alter the course of the Spirits influencing the vital and natural Organs, at least in fome measure, at its pleasure, which is plainly contrary to Experience;) I shall hardly look upon that Hypothefis -Sat13

pothetis to be any more than meerly precarious.

And further, to shew, that such Effects or Alterations of the Vital Organs happening upon violent Paffions of the Mind, are no way owing to such a transmission of the Animal Fluid from the Cerebrum to. the Cerebellum as the aforesaid Author supposeth, I ask, how it should come to pass that in the contrary Passion of Grief, especially when occasion'd by surprizing frightful Accidents, the Heart should so languish, as fometimes wholly to cease beating, feeing in the aforefaid Experiment we find that Motion felffufficient, by vertue of a constant irradiation or influence of the Cerebellum only, and consequently could not be thought so to languish upon fuch occasions for want of those Spirits it never stood in need of.

Without therefore being forc'd to have recourse to that other Hypothesis clogg'd with so many difficulties, I think the aforesaid case may admit of another manner of explication, consistent with what I have all-along advanc'd upon this Subject

rela-

relating to the true fource of voluntary and involuntary Actions if we suppose, that from such Impressions upon the Soul as are either extreamly more or less welcome to it, (in which case the Object is faid to act unproportionably upon the Subject) it may not only act accordingly, above its usual irradiation and force over the Cerebellum, and by that means, as fending the Spirits either more or less copioufly to the Vital Organs, particularly the Heart, the nearest way, (viz.) by the Par Vagum and Intercostal Pair, for that time render them more vigorous, or more languid in their operations, in proportion to the difference of the Passions, just after the manner it happens in cales of Mienation of Mind or Diftraction, where by the Strength of the Impression, or Idea upon the Mind, it drives the Spirits with such an impetus into the Limbs, as makes them act with a vast greater force than what they were wont to do, even above the refistance of Chains or Bars of Iron; but also it may transmit the Spirits more or less cowhooiqueal actions of the Fifera

piously, to the Vital and Natural Faculties, the other way freed from the subsidiary Nerves of the Spina aforementioned, to the Intercostal Pair, which sends forth ramifications to the Heart (in Men especially) equally with, if not more plentifully than the Par Vagum, and from the Vertebral and Brachial to the Nerve of the Diaphragm and Intercostal Muscles, by which means it so falls out, upon such impressions, that the Organs of Respiration to the sight, and that of Pulsation to the touch, are very remarkably affected.

By this means I have endeavour'd to restore the Brain to a capacity of putting its own Conceptions or Impressions made upon it into execution, without being beholden to its neighbour the Cerebellum, and that either in relation to its voluntary, inadvertent, or involuntary Acts; where, note, I make a distinction between Acts involuntary and those of inadvertency, inalmuch as these last, though they are not with, yet they are not contrary to the actual confent of the Will, after the manner of the natural actions of the Viscera,

fuch

fuch as are out of the power of the Will to hinder; besides which, I look upon no other in Rational Creatures (in a strict sence consider'd) to be involuntary, foralmuch as 'tis a contradiction to fay a Voluntary Agent does any thing against his Rational Will (though it may be against his Approbation) by which he is only distinguish'd from a Brute: Though Dr. Willis hath all-along used the word involuntario in another sence, confounding it with acts of meer Ignorance under the term of Inscie, and those also done only inadvertently, or without confideration, under the term of Inconsulto; and doubtless upon this notion of Involuntary Motions built his Hypothefis, which makes all those Actions which are perform'd at any time without the notice of the Intelle-Atual Faculty, notwithstanding at other times they are altogether under its command, equally depending on the Cerebellum as those purely natural, which are always free from the power of the first, and also absolutely subject to the last. Thefe

These Actions I have therefore called by the term of Supervenient Instinct, and being the meer Effect of external or internal Impressions upon Sensative Bodies, as Ecchoes are to those upon such as are only natural, are equally competent to Rational and Irrational Creatures, and capable of being exerted by the influence of the very fame Nerves which minister to the Senfative Faculty, whether it act advertently or inadvertently in the one, or foont aneously in the other, (where, by the way, it may not be altogether unworthy of our taking notice, the genuine fence of that word in Actions performed by those Creatures, is much nearer a-kin to the term Inconfulto than Involuntario in Men) without the supposed rambling Motions of Impressions made upon it, (through Passages only at some times or upon extraordinary occasions made use of) out of the Cerebrum into the Cerebellum.

Now, as to the organisation of this Part, made to consist of various Medullary Prominencies, Appendixes, and Treets, by Nature contrived for aud adjusted to the various functions

ot

of the Soul, and dispensation of the Animal Spirits thro' the whole System of the Nerves, which first are confin'd to, or made to refide in such and such places as so many distinct apartments, viz. the Commune Sensorium in one place, the Imagination and Judgment in another, and the Memory in a third; of which there is such a large and formal apparatus and description (tho' with great discrepancy of opinion) in Willis and Vieussenius, the one placing the Commune Sensorium in his Corpora Striata only, the other in the Superiour and middle Corpora Striata, jointly with the Centrum Ovale; from both whom Des Cartes and feveral others, and with much more shew of Reason, particularly Malpighius, differ, placing it in the ex- Malpig. de tream limits of the medullary part par. 2. of the Brain, where 'tis continuous with the cineritious circumaffuled Part; I must confess, that as I have not been able, by the best enquiry I could make either into Brains diffected whilft fresh, or when boiled in Oyl, to discover any such actual configuration or disposition of Parts, as we find to formally delineated by either

ther of them, but especially the

So neither do I see any necessity thereof, feeing we may much more eafily, and to the lelf-tame ends and advantages, look upon the Soul as one internal principal Sensative Faculty. and the whole medullary part of the Brain, as confifting of fuch Fibrils or Vascula's as in some places more nearly in others more remotely communicate with the Nerves propagated thence to all the external Senfories. one adequate Common Sensory, by which that principal Faculty both receives all its impressions, and accordingly, as by so many gradations of one and the same power, executes or performs those different Functions commonly going under the aforefaid Names of The Common Sense, or Simple Apprehension, Imagination, Judgment, and Memory.

Malplg. de Cerco.p. 116

Par. 2.

And as to the second, (viz.) the Medullary Tracts, by which the Animal Fluid, as by so many Rivulets, is derived from the great Pond or Magazine into many Rivers, surnishing the whole Body therewith, all I could find by the most diligent search,

search, were only those which have already in the preceding Sheets been remark'd, of which, in the first place, are those in the Corpora Striata, very

large and difcernable. We you stugist

Superficies of the Corpus Callosum running transversely by the Septum Lucidum into the Fornix, and from that longitudinally into its hinder Thighs or Pillars formerly called Bombyces, over which they run in a wreathed manner, as was before observed, terminating in the back part of the Lateral Ventricles, enclosed in the hinder Limbs of the Brain, which Ventricles at length terminate in, and are continuous to the subjacent fore-part of the Crura Medulla Oblongata.

Those in the Thalami Nervorum
Opticorum running obliquely down
to part of the subjacent Crura and

Caudex Medullaris.

AME.

Those of the Nates and Testes running after the same manner, and terminating so too, only something lower.

the Messelle Spendis, (vie.) that at tishoffe time the Nerves on one fide

Those in the Annulary Process, which forafmuch as they have ne ver before been taken notice of, I have caused to be engraved in a Figure by themselves, whose Medullary Tracts or Strie, furnished with Spirits both from the continuous medullary Caudex, and Productions of the Cerebellum too, of which the Annular Process is made, (by means whereof the Nerves appertaining thereto may be rationally supposed to be under the influence of both those Parts, conformable to what hath all-along been afferted;) are as visible, being more thick, and of a far harder confistence, than that of the Corpora Striata themselves, (tho upon every attempt of cutting that Process, they may not appear so) and most of them terminating in a middle Medullary Tract, by means whereof there is the same inconveniency prevented, at least in some measure, as there is by that sepimentum of the Pia Mater, continued from the joyning together of the Crura Medulla Oblongata, down quite thro' the Medulla Spinalis, (viz.) that at the same time the Nerves on one side may

may, (as Molinetti, tho' in another Mol. p. 104 place of the Brain, hath truly observed) by any morbid cause, be injured, those on the other may efcape. Is not be as your or may effect the second by the second by the other may effect the second by the se

Concerning these, seeing they seem to have a particular aspect or relation to those Nerves, whose originals we find nearest them, it may not be unreasonable to think they are particular Conduits, from whence the said Nerves are surnished with Animal Fluid, though at the same time we must allow a very free communication betwixt them all.

And consequently, we may suppose the suff of those to convey Spirits from the globous medullary part of the Brain next to it by Vieussenius called the Superious Part of the Centrum Ovale, down to the subjacent medullary part of the Brain, to augment those which are produced lower, and particularly for the service of the Olfactory and Visory Nerves, which last hath more eminently its Supply from the Thalami Nervorum Opticorum.

The

The second fort, or the transverse Striæ's of the Corpus Callosum, to convey an additional Supplement by way of the wreathed
Tracts in the hinder Columns of the Fornix, to the Crura Medulla
Oblongata, where they become continuous to the reflex'd part of the
Lateral Ventricles backwardly, for the service also of the aforesaid two Pair of Nerves, but more particularly to those arising lower either on the Annular Process or Caudex Medullaris.

Those of the Thalami Nervorum Opticorum and Natisorm Processes, the sirst of which lies upon, and is continuous to the subjacent medullary part of the Crura Medulla Oblangata, the other to the Caudex Medullaris, may be supposed to derive Spirits on the behalf of those Nerves which spring from any adjacent parts, whether on this or the other side of the Annular Process or Caudex Medullaris.

O 2 And

tucersum:

cuberance is made, and this fee-And of this fort are the Optick Nerves, which are supplied immediately from the first of those Me. dullary Prominencies, and not unlikely from those fair Medullary Tracts afore-mentioned, running from the Root of the Fornix, extending themselves all the way between the Corpora Striata and Thalami Nervorum Opticorum, in which last at length they are obliterate. The Third, Fifth, Sixth, and First or hard Branch of the Auditory Nerves, mediately by continuity of them with the Annular Protuberance, to all which the other or lesser Medullary Prominencies called Nates. by vertue of their continuity with the subjacent parts, may be suppoled to contribute fomething alfo : and these seems to be better provided for than the rest of the Nerves, inalmuch as belides this way of being supplied from the Cerebrum, they have also another very visible, and much larger, from the Second Process of the Cerebellum, of which the Annular Protuberance

tuberance is made, and this feemingly not without a provident Defign of Nature, seeing the Nerves which are derived thence are much larger, and have a greater Task of fervice layed upon them than any others of the whole Brain, as hath also the Par Vagum, or eighth Pair, which therefore, by vertue of its infertion between the Chordal or third Process of the Cerebellum and Corpus Olivare (and not according to Dr. Willis, from the points or extremities of the Corpora Pyramidalia) hath a double tribute of Spirits, one from the Caudex Medullaris or Cerebrum, the other from the Cerebeldullary Prominencies called M.mui

And to this End or great Service it looks as though this Process was furnished with such a Texture as it appears to have, of strong, large, medullary Striæ's, capable of receiving and containing a Supply from both Fountains.

very visible and much larger, from the Second Process of the Cere-

tuberance

eshed Wich the Annular Pro-

veyed a confiderable Portion of the Whence it may not be unfeafo nable to remark, That not without fhew of good Reason I have allalong afferted the Propriety of the Brain to those Nerves in part, allowed by Dr. Willis to be no further affected by any Impressions of the Brain, than as first conveved from it into the Province of the Cerebellum, and confequently to depend immediately on this last for influence entirely in order to convey Animal Spirits to those parts wherein they are inferted, noque snooth Brain may truly be supposed to be

Upon the Caudex Medullaris, on its under side contiguous to the hinder Extremities of the Annular Process, are situate the Corpora Pyramidalia and Olivaria, overagainst which are the two long Medullary Tracts lately taken notice of, seeming to come from the transverse Medullary Process behind the Testes, and terminating in those other transverse Medullary Process before the entrance into the Fourth Ventricle on the other side,

fide, by which there may be conveyed a confiderable Portion of the Animal Fluid to the Pathetick Nerve, which hath its rife from the first transverse Process, and to the soft or second Branch of the Auditory Nerve, which hath its rife from the second on that side, and also to the Ninth and Tenth Pair on the other side.


veyed from it into the Province


And to conclude, From all these taken together, with the rest of the whole medullary part of the Brain, the Overplus of what is not fpent upon the inmate Nerves of the Brain may truly be supposed to be promiscuously dispensed to all those other extraneous ones produced from the elongation of the Brain, call'd the Spinal Marrow. In which last there is this conformation or disposition of Parts differing from that of the Brain. that whereas in that the cineritions part is external, 'tis here internal; and this for very good reason, and by a provident contrivance of Nature, feeing that not only the cineritious part of the Brain forves for and Fourth Ventricle on the other


fide,


supplying those Nerves which have their original thence, as well as all the rest of the Spinal Marrow; and consequently ought to have the largest space and dimensions possible, which without this situation could not have been; but also without this contrivance the Nerves of this part must of necessity have had their originals from the cineritious part of the aforesaid Marrow, contrary to both the custom and convenience of Nature too.


THE


















Fig: IV.



M. Vander Gucht Sculp:

bh The

Fig:11.



M. Vander Gucht. Sculp:

topplying shole feerves, which have their original thence, as well as all the

reft of the Spine I M.D. Iva and com-

Exhibits the Basis of the Brain, with part of the Medulla Oblongara, the Blood-vessels being injected with Wax.

A A The fore Lobes of the Brain.

BB The hinder Lobes.

CC The Cerebellum,

D D The lateral Sinus's.

E E The Vertebral Arteries as they pass between the first
Vertebra and the Bone of the Occiput.

F The Vertebral Sinus.

G,&c The Dura Mater on the right fide taken off from the Spinal Marrow, and remaining on the left.

1,2,3 The ten pair of Nerves belonging to the Brain,

4,50. with seven of the Spinal Marrow.

The Foramen that opens into the Pituitary Gland from the Infundibulum.

b b The two white Protuberances behind the Infundi-

bulum.

c & The two Trunks of the Carotid Artery cut off where they begin to run betwixt the fore and hinder Lobes of the Brain.

d The two Arteries joyning the Carotids with the Cervical Artery, called the Communicant Branches.

times seeming as the came from the Communicant Branch on each side, from the first of which the Plexus Chorocides hath its original in chief, and from the last the Plexus Chorocides of the 4th Ventricle.

Several little Branches arising from the Carotid

Artery

g The Cervical Artery composed of the two Trunks of the Vertebral Artery within the Cranium.

D

bh The

h h The two Trunks of the Vertebral Artery.

iii The Spinal Artery.

k A small Branch of an Artery running through the 9th pair, broken off from its other part thro' inadvertency of the Graver.

I The Crura of the Medulla Oblongata.

m m The Annular Protuberance, or Pons Varolii.

n That part of the Caudex Medullaris on the right fide called by Willis and Vieussenius Corpora Pyramidalia.

. o That part on the fame fide called Corpus Olivare.

P The foremost Branch of the Carotid Artery, dividing the fore Lobes of the Brain, consisting of two Branches, one of them only appearing here.

q q Little Branches of Arteries helping to make the

Artery upon and thro' the Annular Protuberance.

s s Part of the 2d Process, or Podunculi, of the Cerebellum.

mamail by Proceedings and all of the State o

THE SOUTH AND MANY MANY PARTY AND ASSESSMENT

place a is diese et o its chree B

* * The Spinal Accessory Nerve.

FIG. II.

Shewing the internal Basis of the Cranium, the Sinus's being injected with Wax.

AA	The Edges of the Skull.
BB	The Dura Mater upon the bottom of the Skull.
CC	The lateral Sinus's
d d	The superiour, longer and narrower Sinus's.
ee	The inferiour, shorter and wider Sinus's.
f	The Process of the Bone Cribriforme, called Crista
	Galli.
88	Some small descending Branches of Veins upon the
	bottom of the Dura Mater.
h h	The first Branch of Arteries proper to the D. Mater.
ii.	The fecond Branch of Arteries belonging to the
	Dura Mater.
k	The third Branch belonging to the Dura Mater.
L	The last hole of the Skull.
mn	Several Veins communicating with the inferiour
	short Sinus's.
n	Part of the Os Jugale.
00	The Os Ethmoeid, where the first pair of Nerves or
1	mammillary Processes go forth.
PP	The Optick Nerves cut off.
qq	The Carotid Arteries cut off.
r	The third pair of Nerves visible only on one fide.

The fourth pair of Nerves turned up.

out at the second hole of the Skull.

The fixth pair of Nerves.

The fifth pair of Nerves on one fide expanded before it is divided into its three Branches, on the other fide whole; which Nerves, with its three

Branches, are expressed in the third Figure. Its foremost superiour Branch on the left side, going

X The

SS

The Intercostal Nerve, in this subject proceeding from two Branches of the fifth Nerve, joyning

with the body of the fixth Nerve.

Two Branches of the fifth pair of Nerves, in this subject running almost close to the 6th pair, being partly the Roots of the Intercostal Nerve, which creeps out of the Skull under and between the Coats of the Carotid Artery.

The Body of the Carotid Artery, after it has entred

the Cranium.

- The Glandula Pituitaria.
- The Circular Sinus.
- The Infundibulum. 3

The Frontal Arteries.

- The place where the Lateral Sinus's begin to be 5 declive and tortuous.
- The Dura Mater raised and reclined to shew the Subjacent Nerves.
- The feventh or Auditory Nerves.

The eighth pair, or Par Vagum.
The ninth pair.

I wo large Veilas, III ere B of Taners, the found

were on web file go out mee the mored Turns

Being the Fifth Nerve, with its Branches, whilft within the Cranium. Tomal on a mage

A

Its Trunk.
Its Ganglion, Its first or superiour Branch, going out at the second hole of the Cranium.

Its fecond or midle Branch, going out at the fecond hole. Its third or hindermost Branch, going out at the fifth

sers total Veins on the Pure Maren, which and ore choic that true but the fales of the found.

display of pathypas a consider

hole:

FIG. IV.

Shews the Superiour and lateral Sinus's of the Dura Mater, opened after they had been injected with Wax.

A A. The third or longitudinal Sinus.

BB The first and second, or lateral Sinus's.

C The fourth Sinus.

ddd A Vein running on each fide of the third Sinus.

Sinus of the Dura Mater, after a contrary manner one to the other.

f f The fifth Sinus at the bottom of the Falx.

g The Torcular, where all the superiour and lateral Sinus's meet.

hh The tortuous part of the lateral Sinus running under the Cerebellum.

i i The Veins entering the fourth Sinus from the Plexus Choroeides.

k The place where the fourth Sinus arises.

** The Specus or round hole at which the lateral Sinus's on each fide go out into the internal Jugular Vein.

11 Two large Veins, whereof one enters the fourth Sinus upon the second Process of the Dura Mater, so as to resist the course of the Blood in that Sinus, in its ascent to the Torcular; the other upon the same Process, so as to hinder its descent to the Internal Jugular, contrary to a conformation of Vessels which Vieussenius mentions in his third Table, HH.

mmm Transverse Chordal Ligaments in the longitudinal and lateral Sinus's.

n n Part of the Dura Mater on each fide of the longitu-

o o Portions of the Pia Mater.

PPEC Divers small Veins on the Dura Mater, which enter those that run on the sides of the longitudinal Sinus, according to its length.

99, Sc.

qq& The Veins of the Cerebrum as they appear under the Pia Mater, before they enter the longitudinal Sinus.

R R The falcated Process, with its Veins which enter the fifth Sinus.

S S The second Process of the Dura Mater.

† † The beginnings of the Jugular Veins.

FIG. V.

Representing the Brain in a middle section, the Blood vessels being first injected with Wax.

A A The Fornix cut off at its Roots and turned back.

b b Its Roots at the beginning of the Thalami Nervorum Opeicorum.

cc.Sc. The Thalami Nervorum Opticorum.

d d That part of the Crura Fornicis which growing fomewhat thicker as it turns off towards the Lateral Ventricles, runs over the Crura Medulla Oblongata, which being very prominent in Sheep, and Calves, helps to thrult it up into such a Protuberance as the Ancients called Bombyces or Hyppocampi.

of the first Branch of the Cervical Artery, sometimes seeming as thô it came from the Communicant Branch, in the Lateral Ventricles.

The place where those two Plexus's on each fide

meet under the Fornix.

g g That other part of the Plexus which is made of the fecond Branch of the Cervical Artery joyned with the first by a Communicant Branch not to be seen here, lying under the Crura Fornicis, which is expanded all over the Isthmus, becoming glandulous near to, and especially under the Glandula Pinealis covered here with the Fornix.

hh Two

h h Two large Veins coming from the top of the upper part of the Plexus down to the other Branch of the Plexus, all the length of the third Ventricle, and then terminates in the fourth Sinus.

i i The Trunks of several Arteries, appearing as they were cut off in dividing the Medullary † and

Cineritious * part of the Brain.

k k A Venous Branch on each fide entring the Plexus
Choroeides, from whence there are many slips
branched upon the Corpora Striata.

△ △ The Corpora Striata whole.

The Rima of the third Ventricle.

m m A long Medullary Tract between the Thalami Nervorum Opticorum and Corpora Striata.

on, &c. The Centrum Ovale of Vieussenius.

O The fourth Sinus of the Dur. Mater.

P The Torcular, where the four, and fometimes five, Sinus's meet.

Q Q The Lateral Sinus's.

R A large Vein entering the Lat. Sinus's on one fide. SS, &c. The Gerebellum covered with the second Process of the Dura Mater on its uppermost part,

T T The Vertebral Arteries. V V The Vertebral Sinus's.

W The Medulla Spinalis, with its integuments.

x x The Style supporting the large Veins of the Plexus
Choroeides in the third Ventricle.

q q The Lymphæduets of the Plexus Choroeides.

Y Y Two of the Cervical Nerves springing from the Medulla Oblongata.

tt, &c. The Medullary part of the Brain.

**, &c. The Cineritious part.

South spiem of a poly to the and

The state of the s

FIG. VI.

Being a draught of the Annular Protuberance, Med. Spinalis, Sc. cut through the middle lengthway.

A A The Crura Medulla Oblongata.

BB The Annular Process, or Pons Varolii divided.

cc The Transverse Strie.

ce The intervening Medullary Tract in which the

ff The third or chordal Process of Dr. Willis.

h The Spinal Marrow.

i i Some part of the Cerebellum.

k k The second Processes of the Cerebellum, which com-

1 1 The cineritious part of the Medulla Oblongata.

FIG. VII.

Being the Cerebellum cut through on its hinder part, and reclined laterally.

A A The Cerebellum.

B B The arboreous ramification of the Meditallium of the Cerebellum appearing, being cut right downwards.

CC The Pathetick Nerves.

c c The Nates.

d d The Testes.

e The transverse Process whence the Pathetick Pair have their original.

f The Glandula Pinealis.

g g The first Process of the Cerebellum, running from it to the Nates here extended laterally.

h h The third or Chordal Processes.

i i The transverse medullary Process in the 4 Vent. from whence the soft Branch of the 7 N. has it original.

k k The Medullary Process descending from the Transverse Process behind the Testes, down to the aforemention'd other Medullary Transverse Process.

11 The Originals of that Process a little too low.

m m The eighth pair of Nerves.

n The Calamus Script.or Extremity of the 4th Ventricle

o The Spinal Marrow.

P P The Accessory Nerves.

q q The tenth pair of Nerves,

THE

TABLE.

A

A Rtery Carotid, and the manner no	ite ou
A Rtery Carotid, and the manner of trance into and distribution throi	unh the
Brain. Page	22 22
Where it parts with its borrowed Coat.	n 23
Artery Vertebral, and its manner of or	atriana d
and distribution.	D 26
Artery Cervical	ihid
Why not Conical	D. 28
and distribution. Artery Cervical Why not Conical The Communicant Branches of Arteries Their 71se and Benefit	D.26
Their Use and Benefit ibid.	& 37
Their Use and Benefit ibid. A small Branch of Arteries not before	obler-
oca .	n 18
The Thirty of the Brush Chief M	or the
Cranium with the Veins	D. 27
Artery Spinal, how the Blood is forced i	nto it,
Animal Spirits, how more plentifully pro-	duced,
Augustin Committee of the contraction of the contra	P. 54
Anatomy Comparative, its use	2.54
Arteries, why their Ramifications are ove	r-pro-
portionable to the Trunks in the Brain	
Q and a second	l he

The TABLEAT

The Use of the narrowness of the Communica	MA.
branches, min's odt ni hob Pros	6
How the Carotid Artery in Brutes comes to	be
Smaller above the Dura Mater than under	t,
eget of its firudure, ibid & &	35
Animal Fluid, what, Amonp of not 8.8.1	55T
How it passes out of the Carnous Fibres, P. I.	9
Its effect in glandulous and other parts not fe	167
viceable to Muscular Motion, warthis alb	id.
Its production, barydp. I	1618
Anus, a lufpended, pain Brain	44
Its production, Anus, Aquæ emissarium of Vieussenius, P. I.	3.2
be Cerebellum, B thid, beyond be cerebellum, B thid, thid, this medullary part, part	125
be Cerebellum, B thid	1 :
anal of its medullary part. D. 116	3729
Brain, its Vessels in general, A ship 299 Per Their kinds,	20
Their kinds,	id.
Blood vessels, their different distribution in	re-
lation to the Brain it self, and its Inte	BHTI
ments, bidi the Brain to Nerves often estinam	26
The reason of this different distribution, P.	27
Blood-vessels belonging to the Brain it self on	ely,
p.	32
What proportion the Blood vessels of the Br	arn
bear to the rest of those of the whole Bo	dy,
BUNE P. 32, C	38
The remarkable propagations of its Arter	ies,
Quical Arteries, their Ole, post se	35
Blood-vessels belonging to the Nerves, p.33	,34
Blood in the Sinus's hath two contrary torres	
Adultary part, ibid.	53
The effects thereof, muloli o	old.
The Quantity	Tow

	200
How the circulation of the Blood comes to be	N.
retarded in the Brain, p. 53	
The Bony cell where the Sinus's go out into the	Y
internal Tuontar IV BILL add acode 18 man	
The effect of its structure, ibid. & 54	
The Brain, how distinguish d, p. 87, 112	A
Its two Substances, and add to mo called ibid.	7
Their structure, sale and molubnal p. 80.00	1
Why of a different colour, p. 90, & 92	
Blood, why red	Y
How the Brain is suspended, p. 7, 8.	×
Its Lobes and particular description, p. 113, 114	-
Its hinder Lobes stretched backwardly beyond	
the Cerebellum, ibid.	
Original of its medullary part, p. 115	
Bombyces, vide Hippocampi, Blood-vellels of	1
Watery Bladders instead of Brain, p. 178, 179	1
Watery Bladders instead of Brain, p.178,179	X
Brain petrified, an , to let , and of por 179	
Propriety of the Brain to Nerves often thought	13
under the power of the Cerebellum, p. 198	1
Blood-vessels belonging to the Brain it self only,	A
There were never the Plant nelled water Porge	3
Confent of Parts ! See bool and normodorp. 19	
Carotid Artery, vid. Artery, of the san or inso	
The Circular Sinus, p. 43	2
Communicant Branches between the Carotid and	*
Cervical Arteries, their Use, p. 55, 56.	1
The Cortical or cineratious part of the Brain,	一天
	-
Corpus Callosum, ibid.	-
Q 2 Its	
113	

Its Striæ,	p. 116
Centrum Ovale of Vieussenius,	p. 117
Crura Fornicis,	p. 818
Commissura Crassioris Nervi Æmula	of Willis
and Vieussenius, a gassa sous to as	P.126
Corpus Callofum, and and an anatal	p. 115
Centrum Oyale,	p. 117
Crura Fornicis	p. 110
Corpora Striata,	p. 120
The Cerebellum, a seddo we to the	p.133
Its d fference from the Brain,	p. 135
Corpora duo alba pone Infundibulu	m, p. 140
Corpora Pyramidalia,	p. 141
Olivaria, a sadar to mortisque va san	ibid.
Power of the Cerebellum, p.	170,173
D	
The Dura Mater.	p. 1
The Dura Mater, Its manner of adhesion to the Cranius	n, p. 2
Its manner of adhesion to the Cranius	n, p. 2
Its manner of adhesion to the Cranius	n, p. 2
Its manner of adhesion to the Cranius Is double. What sort of Fibres, and their distri	n, p. 2
Its manner of adhesion to the Cranius Is double. What sort of Fibres, and their distributer Use,	p. 2 ibid. bution,p 3 p. 4, 5, 6
Its manner of adhesion to the Cranius Is double. What sort of Fibres, and their distriction Their Use, How affected in some Distempers, p	bution, p 3 p. 4, 5, 6 articularly
Its manner of adhesion to the Cranius Is double. What sort of Fibres, and their distributer Use,	bution, p 3 p. 4, 5, 6 articularly
Its manner of adhesion to the Cranium Is double, What sort of Fibres, and their distriction Their Use, How affected in some Distempers, pain Vapours, Its Nerves, Its Processes,	bution, p 3 p. 4, 5, 6 articularly p. 6 ibid. p. 7, 9
Its manner of adhesion to the Cranium Is double. What sort of Fibres, and their distriction of Their Use, How affected in some Distempers, pain Vapours, Its Nerves, Its Processes, Their Uses,	p. 2 ibid. bution,p 3 p. 4, 5, 6 articularly p. 6 ibid. p. 7, 9 p. 7, 8, 9
Its manner of adhesion to the Cranius Is double. What sort of Fibres, and their distriction Their Use, How affected in some Distempers, p in Vapours, Its Nerves, Its Processes, Its Processes, Its Blood vessels apart from the B	p. 2, ibid. bution, p 3 p. 4, 5, 6 articularly p. 6 ibid. p. 7, 9 p. 7, 8, 9 rain, their
Its manner of adhesion to the Cranium Is double. What sort of Fibres, and their distribution, when the Cranium Is double. What sort of Fibres, and their distribution, and their distribution, when I is double. Its Nerves, Its Processes, Its Blood vessels apart from the Bunumber and distribution,	p. 2 ibid. bution,p3 p. 4, 5, 6 articularly p. 6 ibid. p. 7, 9 p. 7, 8, 9 rain, their p. 20, 22
Its manner of adhesion to the Cranium Is double. What sort of Fibres, and their distriction of Their Use, How affected in some Distempers, pain Vapours, Its Nerves, Its Processes, Its Processes, Its Blood vessels apart from the Bunmber and distribution, Two sorts of Dropsies of the Brain of	p. 2. ibid. bution, p. 3. p. 4, 5, 6 articularly p. 6 ibid. p. 7, 9 p. 7, 8, 9 rain, their p. 20, 22 bierved by
Its manner of adhesion to the Cranium Is double. What sort of Fibres, and their distriction of Their Use, How affected in some Distempers, pain Vapours, Its Nerves, Its Processes, Its Processes, Its Blood vessels apart from the Bunmber and distribution, Two sorts of Dropsies of the Brain of Tulpius and Wepler,	p. 2, 2, 6, 6 articularly p. 6 ibid. p. 7, 9 p. 7, 8, 9 rain, their p. 20, 22 biferved by p. 59
Its manner of adhesion to the Cranium Is double. What sort of Fibres, and their distriction of Their Use, How affected in some Distempers, pain Vapours, Its Nerves, Its Processes, Its Processes, Its Blood vessels apart from the Bunmber and distribution, Two sorts of Dropsies of the Brain of	p. 2. ibid. bution, p. 3. p. 4, 5, 6 articularly p. 6 ibid. p. 7, 9 p. 7, 8, 9 rain, their p. 20, 22 bierved by p. 59 ibid. & 60
Its manner of adhesion to the Cranium Is double. What sort of Fibres, and their distriction of Their Use, How affected in some Distempers, pain Vapours, Its Nerves, Its Processes, Its Processes, Its Blood vessels apart from the Bunmber and distribution, Two sorts of Dropsies of the Brain of Tulpius and Wepler,	p. 2, 2, 6, 6 articularly p. 6 ibid. p. 7, 9 p. 7, 8, 9 rain, their p. 20, 22 biferved by p. 59
Its manner of adhesion to the Cranium Is double. What sort of Fibres, and their distriction of Their Use, How affected in some Distempers, pain Vapours, Its Nerves, Its Processes, Its Processes, Its Blood vessels apart from the Bunmber and distribution, Two sorts of Dropsies of the Brain of Tulpius and Wepler, Their Solution,	p. 2. ibid. bution, p. 3. p. 4, 5, 6 articularly p. 6 ibid. p. 7, 9 p. 7, 8, 9 rain, their p. 20, 22 bierved by p. 59 ibid. & 60

vale of Vieussenius, p. 117	Centrum C
E giol	Crura Fore
Elasticity of the Blood, wolfer	Examillar
The Effects of that being weakened	ani V bishid
Extragalation of the Nutrition	EL Land
Extravasation of the Nutritious	Finias, its
effect, color	o murpios
Elasticity not competent to the fir	It principles
o of Bodies, Bien	IC STON KOO
The Occasion of it in other Bodies,	Kidi Cerebe
Elasticity its æquilibrium in the	phole compa-
pages of Muscles, lang adla o	The reaction do
The Effect thereof, Milebimer	p. 105
Enteriments by Ling: C. 1	1010. X 106
Experiments by Injection, of what	use in Mus-
cular Motion, mulledone	p. 110
F	
Mater. D. E	The Dura !
The Falx and its particular uses,	D 780
Is wanting in several Creatures, a	wd why 50
Fibres of the Dura Mason A	na way, p.9
Fibres of the Dura Mater do	A CONTRACTOR OF THE PARTY OF TH
Valves,	18 p. 5
Fleshy part of the Body, what,	P. 94
Fluids of the Body, their different	motions, p.96
	p. 117
	8ri.qoceffe
The Fornix of Vieussenius,	Tally mall
The Fornix commonly to catted	p. 115
THE TOTAL COMMINGULATOR CALLED	THE PERSON OF TH

Its Fimbria, according to Vieusienius, p. 118
Its Crura, ibid.
How natural and vital Functions relate to the
Cerebellum, and with what difference,

P. 170, 171

Q3

How

How they come to be not under the power of the rational Soul, p. 174
How performed when the Brain is utterly incapable of acting, p. 177

G

Injection with Mercury makes Blood-vessels ap-The Glandula Pituitaria enclosed in Strong muludibnaini p. 46 Membranes No Serum can get through its Integument, ibid. & 75 8 Glands of the Plexus Chorocides, p. 62,63 The Glandula Pituitaria not capable of carrying an Excrementitious humour, p. 69, 73 The Gland. Pit. its situation, & snimero p. 71 Is not suspended in Men as in Brutes, ibid. & 73 In substance it differs from all other Glands, ib. Is of two forts, and why, with guiratte p. 81 In what manner the Lympha gets into it, ibid. The Glandula Pincalis, its situation and con-8 nexion, Is of the Conglobate kind, 39 9130 330 13 19 19.84 Errors of DesChartes, Lower and others about it, 2848819 al Seafer, their feats, Geminum Centrum Semicirculare of Vieuslenius,

regaments of the Manse

Fleadach, how happening in Feavers, p. 29
From the closeness of the Pores in the Crenium,
P. 42
Two

Two forts of Hydrocephalus, amos vada p. 59 Hippocampi Arantii, 81 11 g rational Souls Their Strie, niars adt nada bamrofrag aibid. pable of alling. Injection with Mercury makes Blood-vessels ap-4 kg Glandula Pituitaria enclosed The Infundibulum, 23mbrames p. 77 The difference between it in Men and Brutes, Its two Ducts in Brutes, suxal 9 ods to shap. 78 Its Office, dan bor va mattaria Rolling 379, 80 Infundibulum, the passage into it by three Foramina's, mostantil eti Jil bris Dp. 124 21 sqt suspended in When as in Brutes, sumdfil. Instinct, what was mort erost to sope 162, 163 The differing effects of some Impressions upon the 18 1 18 18 red the Lympha gets inches iside Transmission of Impressions, according to Doctor Willis, improbable, Pru 847 185 Difference between involuntary and inadver-78 ingres DesChartes, Lower and the trestoutie Internal Seuses, their feats, P. 190 Geminum Centrum Semicirculare of Vieufles D. IZZ Lsgaments of the Sinus's, Their Uses P, 52,56 Lateral Sinus's, we governo and salahab. 140 Longitudinal Sinus's, to to Sono old sup. 11 Lymphæduets of the Brain, p. 61, 62 Lym-DUL 8

Lympha, how generated within the Ventricles,
p. 81,82
To what end, ibid. Laughter, how made, p. 181
Laughter, how made, p. 181
Why peculiar to Man, p. 182
Libidinous Actions, how caused, p. 181, 182
therwest fusce, bon generates bid.
Nerver its firment M P. 93
TO ESS DICE TESTS FREE PRICE OF SOLE
Membranes of the Brain, vid. Dura and Pia
Mater.
Muscular Motion, p.99
Divers Opinions about it, ibid.
Muscle, its inflation or contraction, p.102,106
Muscle, how contracted or swelled by force,
No Glas the Green Coll : 1: 1: 104
Muscles, the effect of their being out through,
Muscles their bending and see fine p. 105
Muscles, their hardness and swelling in con-
Muscular Motion Come particular Phonomera
Muscular Motion, some particular Phænomena about it solved, p. 110, 111
Muscular Motion made by the Nervous Fluid
alone without concurrence of the blood, p.177
A Medullary Tract not before observed, p.123
Medulla Oblongata, p. 139
Hs Crura, mumaro sa midian asura ibid.
Animal Motion, p. 158
Motion voluntary and involuntary, p.159, 160
Two sorts of Motions in Brutes, P. 161
Medullary Tracts of the Brain, p. 192
The state of the s
37

The TABLE. Lympha, how generated within the Ventricles, V. P. 81,82

N P. 51, 52	
To what end, ibid.	admin 8
Noise in the Head, how occasion d,	p. 29
Nervous Juice, walk of tallange of p.	88. 92
Nutrition q below could recitop. 88	,89,94
Nervous Juice, how generated	ibid.
Nerve; its structure	P. 93
The effect of its being taper, ibid.	& 102
Nates of the Brain, . vid. Dara and to 250	p. 125
Nerves Olfactory,	p. 143
Optick Moitold	P. 144
Motorium or third Pair	p. 145
Patheticum or fourth pair	p. 146
The fifth pairs son betauting word	p. 147
The fixth pair	p. 148
The Intercostal and west to south sat	ibid.
The seventh or auditory pair	P. 149
The eighth or Par Vagum	p. 150
The Accessory Spinal Nerve	P.151
The ninth pair when sind smot mostakly	D. 142
The tenth pair	p. 153
The structure of the Nerves	p. 154
Nerves their different functions	p. 157
The effects of Communications betwixt	Nerves,
P. quongata, p. 130	75,181
The Nerves within the Cranium, how	upplied
with Spirits from various medullar	Tracts
of the brain, p. 194,1	95, Gc.
of Mocions in Boutes, p. 161	57.101 and
Variets of the Brote, P. 192	19 11 19 37.63

W

Hath two different Originals

bidi 1	ent Originals	b two differ	Elat
bidi	0	t they are	Wha
to be glandulous.	part begins	विभागी अंपर अंग	24.18
Optick Nerve i	ts original	9 3 33 3	p. 123
External Object	s, how they	act is and a	11158
8 Improportionabl	e actings of t	ne Object	100
Organization of	Animal Flui	1 of the Cer	ehrum
Overplus of the	Antimus I ves	sumos slovos	D TOO
with its use	and to make	20,00	1
art of the Pickus	a Red Pla 2	Feins joynin	Ino.
D.61	ity .	its extrem	ni
The Pia Mater, Why called Cho	ity. Keins en	draw reducte	p. 10
Why called Cho	roeides	, sun	bid.
Its particular o	listribution	mandin Le	toid.
How to Ind it	in the Ventr	icles shape	p. 11
Is double every	where, but	where most	visibly,
	2/10		
Its inward Lam			
Its Ofes quo	richtly baral	China China W	P.10
Its Blood-vessel. How it invests			
ON Fibrils 11 9	110,000	erebellum	D. TO
How the Arteri	es belonging	to the Brain	it felf
are ramified	through it	ellus Ivaribi	p. 33
How Some brane	ches of the C	carotid and	Verte-
8s1 bral Arteries	s are exempt	from it,	P. 34
The Processes o	f the Dura I	Mater, with	their
distribution	ind use	dina a cuit	
Pulsation of the	Sinus s, who	nce Day	P.50
The Plexus Che	proeices of the	ne Drain	p. 57
It is double			ibid. Hath
347			77010

	- I - A - Company of the Company of
Hath two different Originals	ibid.
What they are	ibid.
Where the first part begins to be gi	
Nerve its original biss	Marie Control of the
Where the first part of the Plexus tern	
ortionable allings of the Objection 186	
Where the second part begins to be	
is of the Animal Fluid of the Cerebrum	algravibid.
The double connexion of the two par	ts of the
Plexus	ibid.
Two Veins joyning the first part of the	e Plexus
in its extremity	p.61
Its large reductory Veins entering to	be fourth
bidi Sinus Didi Sinus	ibid.
Mrs Lymphæducts and reliable reliable	
Its Glands relatives out at it but	
li every where, but where All rightly,	
Its situation and use	Married Street, Street
The Pituitary Gland hath Veins	P. 66
Processus Lentiformes, vid. Corpor	a Striata.
Passage into the Infundibulum	P.123
Procellus Annularis, or second Proc	ess of the
Cerebellum p. 129,	135, 140
Procellus Natibus antepolitus	p. 130
Processus Nervi Amulus	Po 125
Processus Nervuli Æmulus	10 por 12 9
The Plexus Choroeides of the Cerebe	H PT24
Processus Vermiculares han wound	p.134
The first Process of the Cerebellum	hididarra
The third or Chosdal Process of the C	erebelling
	ob p. 136
Fark	The

The medullary transverse Processes of the for Ventricle p.	urth
Ventricle p.	136
The medullary Processes descending to	thoje
transverse ones posterolah rolling p.	137
transverse ones The Plexus Choroeides of the Cerebel	lum,
"quim can be separated but by wroner se."	133
Perception or Passion, what,	158
Sinus's bave no gullation of thereis wer	200
Sinus's have no pullation of themselves, P. 500 Pullation is from the Brain, ibid.	
r pullation is from the Brain, bid.	Thes
The Receptacula Sellæ Aquinæ, &c. of V	ieus-
Coming not existent in Men	200
Their Use impossible	0.46
Their Use impossible The Rete Mirabile Always existent in Men	. 64
Always existent in Men	ibid.
Its Intuation	idia.
Why smaller in them than Brutes ib. 6	x 65
The effect of its being so large in Brutes,	p. 65
Is differently situated in Men and Brutes,	P.72
Hindermost Roots of the Medulla Spil	nalis,
what formerly a viscous 20116 3/2 p.	126
Respiration, how performed P.	165
Why a Child unborn respires not, p.	168
Why Respiration ceases upon cutting the	ere-
nobellum, Jaluatev mard sat to sup.	174
rion, bon made, p. 91, 92	1,330
tion, bone explaine pass, 101	
made, p.158	TOEF
Sinus's of the Brain	. 39
Their number mubiou.I min	
	p. 40
The third or longitudinal	D. 41
	The

The fourth or internal one	ibid.
Four other smaller Sinus's, and their	
othor,	P.42, 43
7 0: 1 0:	p. 43
Its particular description ib	id. & 44
Its use Coorceider of the Cerebaju atl	p. 47, 48
No Serum can be separated but by	proper le-
	P.47
The Sinus's have no pulsation of t	hemselves.
	p. 50
Their pulsation is from the Brain,	ibid.
The different Ligaments of the Sinus's	D. 57
The use thereof	ibid
THE DITTOR I ACTIVITED OF LINGUISTING	33C AP 4/4A
Sinus's	515 p. 52
Sinus's Their Uses The Etructure of the Sinus's	wibid.
The structure of the Sinus's	P.53
Effects thereof	ibid.
Animal Spirits, how made more plenti	fully, p.54
Why the Sinus's grow so wide on a su	
Their difference of structure	
The longest Sinus commonly burst in	Strangled
bodies bentro trag orod w	bidi
How the blood passes the Lateral	Sinus's in
different positions of the Brain	P. 54
	mulp. 91
Secretion, bow made,	p. 91, 92
Sensation, how explained	0.88, 101
How made,	p. 158
Corpora Striata	115,120
Septum Lucidum	P. 119
Its Striæ	ibid.
or bengitudinal	The

THE RESIDENCE OF THE PARTY OF T	
The Spinalis Medulla in a	Tortoife officiating
instead of the Brain	p. 176
Sneezing, why peculiar to M	languistra april 82
Commune Senforium, what	191 q the Dura N
Striæ of the Annular Pi	
p. 26, 27, 28,	p. 193, 197
Why terminating in a middle	le medullary Tract,
the Sinus's C. Cz. Cz.	261.d Keins enter
Conformation of the Spinal	Marrow differing
from that of the Brain, as	nd why and parent
from what they have in	Sums a genter.
T. P.54	Men
Bon String	the eins of the Corpor
Torcular Herophili	12023H124-36 b. 45, F
Tractus Medullaris Th.Ner	
of Vicustenius gmv I and	a Poluspuos shada
Tract. Med. Natibus anter	ontus of vicunent
bidi serrerici	Vaidi how continuon
Transpiration, what,	sas gararamo progra
Conspiration of Hippocrates	Digital and Alexand
Tone of Parts 2911911A to 28	Later due bioners
Testes Thalami Nervorum Optica	Valete of Vellalmura
Tria Foramina relating to	the Infundibulum
	ide Infundibulum.
P. 128,129,139,139	ies henarion and who
of the Broin	The shree Venericles
	Vulva Cerebra
psiq L G colled	2 1 1 1 1 1 1 1 1 1 1 1
Vapours, commonly so called	June Moter of
fecting the Fibres of the I	ma mater, P.O

Some Veins of the D. Mater entering the third
IN gad of the Brain sunis
Sinus Vertebral Artery, and or washing vide Artery.
Veins of the Dura Mater enter the Brain with
the Arteries contrary to those of the Brain
Two Veins entering the Circular Sinus, p. 45, 46
How the Veins enter the Sinus's p. 52, 53
The effects of their different entrance, p. 54
The Veins have a different disposition in the
Sinus's of Brutes, from what they have in
Men Veins of the Corpora Striata P. 54 p.61
The large reductory Veins of the Plexus Cho-
bidius Medullans Th. Nerv. Opt. in rebisons
Vapours condensed into Lymphamop. 81
Vascular constitution of Parts 1811 by p. 91
Veins, how continuous to Arteries ibid.
Vessels containing the Animal Fluid are capillary
productions of Arteries of P. 93
Veins only productions of Arteries p. 94 Vessels their minuteness p. 95,96
Vestels of VesselsmurositqO murovre Mibid.
Ventricles of the Brain The Brain p. 117
Valvula major p. 128, 129
its situation and use p. 128,129,130,132
The three Ventricles of the Brain p. 117
Vulva Cerebri p.124

Fapours, commonly so called, how sometimes affecting the Fizesy Waraflater, p. 6

Some

ERRATA.

PAGE 9. 1. 14. for to read towards; p. 16. 1. ult. for from which r. which from; p. 32. in the title of the Chapter, for Veins r. Veisels; p. 32. 1. 13. after Veins insert which last have already been treated of; p. 64. 1. 5. dele only; p. 89. 1. 16. Vitrious r. Vitrous; p. 92. 1. 29. for Septometry r. Leptometry; p. 102. 1.3. for contrasted r. contrasts; lbid. 1.29. for reflexed r. relaxed; p. 109. 1. 18. for hastening r. happening; p. 117. 1. 28. for Semicirculari r. Semicirculare; p. 119. 1. 12. for becomes r. become; p. 138. from And therefore in the 7th line to the end of that Paragraph, leave it out: p. 137. 1. 7. for above r. below; p. 168. 1. 8. after passage add at lease but very little.



