An essay on the uterine haemorrhage, which precedes the delivery of the full grown foetus. Illustrated with cases / [Edward Rigby].

Contributors

Rigby, Edward, 1747-1821.

Publication/Creation

London: Printed for Joseph Johnson ..., 1775.

Persistent URL

https://wellcomecollection.org/works/rupdey5y


License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.




Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



TV.a.713. -4044/8

Carran



T3429.

E S S A Y

ON THE

UTERINE HÆMORRHAGE,

WHICH PRECEDES THE

DELIVERY

OF THE

FULL GROWN FOETUS:


ILLUSTRATED WITH CASES.

BY EDWARD RIGBY.

LONDON: 110

PRINTED FOR JOSEPH JOHNSON, No. 72, ST. PAUL'S CHURCH-YARD.

MDCCLXXV.



E S A

HRT MO

MA

UTERINE HEMORRHAGE,

WHICH PRESENTS WHE

DERIVERRY

HISTORIOAL AD JAUT

ILLUSTRATED WITH CASE

EST, EDWARD KIGEY.

PRINTED FOR JOHNEY JOHNSON, No. 72, Sr.

ADVERTISEMENT.

MOST of the cases on which the doctrine contained in the following pages, is founded, sell under my notice in consequence of being appointed to attend all the poor women in a large and populous city, who have difficult and dangerous labors.

I thought it right to premise this, as were it not known what circumstance gave me an opportunity of collecting them, the number of the cases, when compared with the short space of time in which they occurred must appear so extraordinary, (exceeding, in so great a proportion, the number usually met with even in the most extensive private practice) as possibly, with some readers, to render their authenticity doubtful, and consequently to invalidate the reasoning deduced from them.

Norwich, Sept. 30, 1775. E. R.

MADVERTISEMENT.

Ages is founded in the following fages in the following fages is founded in the following confequence of being attained to attend all the poor voomen in a large and populous city; who have difficult and dangerous labors.

Thought it right de premie this, as were constitued have gave an apportunity of collecting them, the number of the coper, when command with the port face of their face that they are the face for the face of their face that they are considered with the face for the face of the face that they are considered with the face for the face of the f

A T

Sept. goliggs.

A N

Remoived viewed Smit, exh al

stod of pressincy, are delicen

E S S A Y

ON THE

UTERINE HÆMORRHAGE.

turition, exposes women to so much danger, as prosuse Hæmorrhages from the Uterus, towards the latter end of pregnancy, and in the time of labor; the art of midwisery is likewise, in no instance, more at a loss in the use of means for the relief of the patient; an enquiry into the causes of them, and an attempt to improve the practice in such cases, cannot, therefore, be useless.

B

THE

THE treatment of floodings, that come on before the Uterus has acquired any confiderable fize, must be very obvious, and the consequences of them, at that early period of pregnancy, are feldom to be dreaded, as, if the patient lofe blood from the arm, be kept cool, and in an horizontal posture, and such mild, astringent, and anodyne medicines be administered to her, as have been found, by experience, to restrain discharges of blood, they will very frequently stop entirely, and the woman go on to her full time: and if this should not be the case, but the Hæmorrhage should still increase, it will feldom increase to a degree that will endanger the life of the mother, without the small fœtus and fecundines being feparated, and thrown off by it, after which the Uterus will foon contract, and thereby closing the mouths of the bleeding veffels, the difcharge will gradually diminish, till it be entirely stopped: the furgeon has, therefore, in these cases, nothing manual to do, for, notwithstanding it has been recommended commended by * Mauriceau and Deventer, and it is faid to be the practice of fome, to endeavour to bring away the fœtus by art, even in the earliest months, I am persuaded, from experience, that it is never necessary, and were it even necessary, I cannot conceive it possible to do it with the hand.

But floodings that precede the delivery of the full-grown fœtus, when the Uterus is arrived at its greatest stretch, and the vessels have acquired their utmost magnitude, must be ever highly dangerous, being more profuse, and more difficult to suppress, in proportion to the increased size of the vessels; insomuch, that the number of instances in which they have unhappily proved fatal, is very considerable.

Most of the authors whom I have read on this subject, describe these cases as particularly embarrassing, and seem

^{*} Van Swieten's Commentaries, Sect. 1308---9.

alike to acknowledge, that they have always been at a loss when such have occurred to them, to determine, with any degree of certainty and fatisfaction, which of the two methods of practice hitherto recommended, it has been most proper to adopt; whether to endeavour to restrain the discharge, by the means before mentioned for that purpose, and leave nature, by her own efforts, to expell the child, as is the case in floodings of the early months, or at once to introduce the hand into the Uterus, and bring it away by art.—This doubt about the propriety of waiting, or the necesfity of removing the contents of the womb, they fay, is ever owing to the uncertainty of knowing the quantity of blood that has been loft, and, if it were known, to the impossibility of ascertaining the degree of loss, that a woman might fustain, without manifest rifque of life.

They all however agree, that when the discharge becomes very profuse, and

and fuch a confiderable quantity of blood has been loft, as threatens the immediate death of the patient, that nothing but a fpeedy delivery will give any chance of preventing it; and have given us cases, wherein the bringing away the child by art has been attended with fuccess; they likewise inform us of others, in which waiting and purfuing the palliating means has been justified, by the natural pains having come on, and the child having been timely expelled by them; moreover, where both methods have been used, a great number are related which, nevertheless, proved fatal: but no particular reasons have been given, why the different methods of practice were used, why the fame methods in some cases have succeeded, and in others, apparently fimilar, have failed; nor have any hints been fuggested to us, which might lead us, at the beginning of the complaint, even to a probable conjecture, whether the Hæmorrhage be of that kind, which requires the turning the child, or not.

WE need not be furprized, then, to find, that upon a fubject of fuch acknowledged uncertainty, there should be fome writers who give the most opposite advice; for as it is reasonable to fuppose that the furgeon, who has lost a patient by too long waiting for the natural pains, will, in all future cases, think it right to turn the child upon the first attack of the complaint, so it is equally natural to suppose that another, who has had feveral that have terminated fafely without turning, will think it feldom necessary: thus, to instance but two, * Chapman invariably recommends the delivery by art, upon the first coming on of the complaint, and † Puzos advises always to wait for the natural pains, which he believes will rarely fail of putting a fafe end to it.

It is faid that an eminent lecturer in midwifery, in London, directs his pu-

^{*} Essay on the Improvement of Midwifery, chiesly with Regard to the Operation. 1733.

⁺ Memoire sur Pertes de Sang.

pils not to be too hasty in checking the discharge, as he imagines some cases that have fallen under his notice, have, by suffering the vessels to unload a little, turned out better than others have done, in which means were used to restrain it upon the first attack.

ANOTHER, (who is lately deceased,) not less capable of judging upon the subject, acknowledged himself totally at a loss what to advise, and said, that surgeons must, in a great measure, be lest to their own discretion, when such cases occur; but speaking in general terms, he thought it right at first to endeavour to check the Hæmorrhage, and wait for nature's assistance by pains, and if they should not come on, the slooding should increase, and the woman grow weaker, it was then right to have recourse to delivery by art.

It is hardly necessary to observe, that contradictory as these directions are one to another, they must all in their turns be be improper, as they are guided by no fixed rules; and if no information be, therefore, to be had, than what can be collected from books, and no other directions are to influence our practice, than the vague ones we have mentioned, it will ever be uncertain, it must frequently be unfuccessful; for we must either wait undetermined what to do, till the discharge becomes very profuse, and fo much blood is loft as renders what we then do probably useless, or we must do it before much loss has been fustained, at a time when the patient appears to be in no danger, and when we cannot have the fatisfaction of knowing that nature will not be able herfelf to expell the child, and that the turning is absolutely necessary; the timid practitioner, encouraged by no certain guide, and cautiously afraid of giving his patient unnecessary pain, we may reasonably fuppose, will, for the most part, be guilty of the first error; while another, who is more precipitate, through a defire of preventing the danger of delay, will

will as often make use of painful means, when the efforts of nature, assisted by more gentle methods, would probably be equally successful; and, at the same time, he will likewise unnecessarily expose his patient to the danger, which a too early delivery may perhaps occasion.

viomo ottoretsu abl

To remove the uncertainty and embarrassiment, which have hitherto attended the practice in these cases, and determine on more fixed and rational principles, when it is safe to wait for nature's endeavours to expell the child, and when it is absolutely necessary to bring it away by art, would, therefore, certainly be an important improvement in midwifery.

For this purpose two things appear to be indispensably necessary; first, to know the reason why, in cases that have began exactly alike, where the discharge, pulse, and faintness have appeared the same, and there has been no remarkable disference

ference in the age, strength, and constitution of the patient, and the same treatment, has, likewise, been made use of, they have, nevertheless, in the end turned out quite differently, why in fome, the discharge is restrained by using the common palliating means, and the labor terminates fafely by waiting for nature to empty the womb, and in others, notwithstanding the use of the very fame methods, it increases to a degree that exposes the woman's life to the most immediate danger, and thereby renders the turning of the child neceffary: and, fecondly, to be able to procure this information as early as poffible after the coming on of the Hæmorrhage, fo as to enable us to determine with certainty, before too much blood has been loft, whether it be right to endeavour to restrain it by the means before-mentioned, or to proceed at once to delivery.

A KNOWLEDGE of the true causes that produce floodings, will give us all the

UTERINE HÆMORRHAGE. 11

the information, which I have confidered as the first requisite towards an improvement in the practice; for though it has been little noticed by those who have written upon the subject, they certainly arise from two very different causes, which are very different in the danger they produce, and which require a very opposite method of treatment.

FLOODINGS have, indeed, heretofore been confidered as arifing from two different causes, one alone of which was supposed dangerous; a distinction having been made, by fome authors, between the discharge which came from the Vagina, and that which proceeded from the Uterus; and when it came from the Uterus, they also diffinguished whether it came from the bottom or the orifice of the womb, by which was only meant, whether it was occasioned by a feparation of the Placenta, or whether it was owing merely to a rupture of the vessels of the Vagina or Os Uteri, produced

duced by the distension of labor: this distinction, to those who are the least conversant with practice, must appear trifling, as no bleeding of confequence enough to deferve confideration, ever comes from the latter; and that, which is the object of the present enquiry, always proceeds from the Uterus.

THE feparation of the Placenta from the Uterus, before the delivery of the child, and the consequent opening of its vessels, must, therefore, be looked upon as the proximate cause of every confiderable discharge of blood from the womb at that time: but this premature separation of it may be produced by very different causes, and it is a knowledge of this difference that will, in my opinion, remove the difficulty of ascertaining the reason, why the same apparent complaint should, very often, fo widely differ in its termination, and at the fame time remove also the uncertainty of treating it.

THERE is no particular part of the Uterus, to which nature feems constantly and uniformly to fix the Placenta, it is, nevertheless, for the most part, so situated, that if the woman be healthy, and no accident befall her, it does not feparate till the full term of pregnancy, nor then before the entire expulsion of the child, after which it becomes difengaged from the Uterus, and is thrown off, making room for its entire contraction, which shutting up the mouths of the veffels, effectually prevents any confiderable lofs of blood; for which purpose, it is plain, it must be fixed to some part of the womb which does not dilate during labor, namely, to the fundus or fides of it.

In this case, then, when a flooding comes on before the delivery of the child, it is obvious, that the separation of the Placenta must be owing to some accidental circumstance, to violence done to the Uterus by blows or falls, to some peculiar laxity of the uterine vessels

from badness of habit, or fever, or to the influence of the passions of the mind suddenly excited, such as fear, anger, &c.

But from the uncertainty, with which. (as before observed) nature fixes the Placenta to the Uterus, it may happen to be fo fituated, that when the full term of pregnancy is arrived, and labor begins, a flooding necessarily accompanies it, and without the intervention of any of the above accidental circumstances; that is, when it is fixed to that part of the womb which always dilates as labor advances, namely, the Collum and Os Uteri, in which case, it is very certain that the Placenta cannot, as before described, remain secure till the expulsion of the child, but must, of neceffity, be separated from it, in proportion as the Uterus opens, and, by that means, an Hæmorrhage must unavoidably be produced.

UTERINE HÆMORRHAGE. 15

THAT floodings, which arise from these two different causes, which I will distinguish by the names of accidental and unavoidable, though they may appear exactly fimilar in their first fymptoms, should terminate very differently, if left to nature, affifted only by the palliating means before mentioned, cannot feem strange; nor can it be a doubt that of these two kinds of floodings, only one of them, namely, that which is produced by an accidental separation of the Placenta, can be relieved by the use of these palliatives; and that the other, in which the Placenta is fixed to the Os Uteri, and the flooding is therefore unavoidable, cannot possibly be suppressed by any other method whatever, than the timely removal of the contents of the womb: for supposing the discharge to be for a while restrained by bleeding, medicine, cool air, &c. it will inevitably return when nature is fo far recovered as again to bring on labor: in the first case, if the Hæmorrhage have been checked by the use of the above means,

it is not impossible but labor may come on, and the child be fafely expelled by the natural pains, before it returns; or, if it should return, it may not increase in quantity; as in this case, very probably, the feparated part of the Placenta, which occasions the discharge, remains nearly the fame; whereas, in the other case, in which the dilatation of the Os Uteri produces the feparation of the Placenta, every return of pain must be a return of the bleeding, and it must become greater and greater, as the Uterus opens more and more, and the Placenta is in proportion detached, till it increases to a degree that exhausts the patient, and she dies before nature has been able to expell the child. That fuch must, inevitably, be the progress and event of floodings arifing from fuch a cause, if left to nature, is too obvious to be further infifted on.

THAT this attachment of the Placenta to the Os Uteri, is much oftener a cause of floodings than authors and practitioners

UTERINE HÆMORRHAGE. 17

tioners are aware of, I am from experience fully fatisfied; and fo far am I convinced of its frequent occurrence, that I am ready to believe that most, if not all, of those cases which require turning the child, are produced by this unfortunate original situation of it; and, moreover, (which is perhaps of as much practical importance to know,) when the Placenta is not so situated, the events of the annexed cases authorise me to say, that if the patient be properly managed, nature will, for the most part, terminate the labor safely, without any manual assistance of the surgeon*. And,

* I HAVE the satisfaction of adding two very respectable authorities in further confirmation of what I have just said, the one is of Mr. Charles White of Manchester, and the other of Mr. John Aikin, of Warrington, gentlemen well known both as surgeons, and as writers; Mr. White, who has had the most extensive practice in midwifery, as there was a time when almost all the difficult cases not only in Manchester, but in a very populous neighbourhood, through a large circuit of many miles, fell under his care, and who is therefore well qualified to judge upon the subject, tells me, "That the distinction I make between sloodings which are accidental and those which are unavaidable, perfectly agrees with his

independent of the proofs which experience gives, it feems reasonable that in the latter case it should be so; for those who are much conversant with the difficult part of midwifery, must have obferved, how much more nature is able to do for her own relief than is commonly imagined, and how, unexpectedly, she will sometimes effect, what art has been a long time in vain attempting; if we add to this, that when any dangerous circumstance affects the Uterus, nature ever makes some effort to remove it, need we be furprized, that in these cases, when the Placenta is not at the mouth of the womb, and there is, therefore, no impediment to its dilatation, and the expulsion of the child, she should, for the most part, safely effect both?

experience in fuch cases, and that he is very clear that sew, if any, of the former require turning and delivery by art." And Mr. Aikin, whose practice is also considerable, says, "That he has never had occasion to use forcible dilatation and turning except where the Placenta has been found at the mouth of the Uterus."

UTERINE HÆMORRHAGE. 19

There are not, indeed, wanting relations of cases, in which the Placenta has been found at the mouth of the Uterus, but it was usually supposed to have been separated from some other part of it, and pushed down into that situation by its own gravity, and the force of the natural pains, and some have even denied the possibility of its ever being originally fixed there.

Mauriceau* has a chapter on this subject, and La Motte† relates three cases of it; but they both suppose that when the Placenta presents it is ever wholly detached, and considering it therefore as a foreign body, direct that it should be always immediately brought away.

Dionis ‡ fays, "That the afterbirth fometimes loofens before the mem-

^{*} Diseases of Women with Child and in Child Bed, translated by Chamberlen, 1752.

[†] General Treatife of Midwifery, translated by Tom-kyns, 1746.

[†] Treatise of Midwisery, translated from the French, 1719.

branes, which contains the waters, are broke, and when the infant turns itself, it is to be found at the internal orifice of the womb."

Ruysch || says, "It is well known, that the Placenta Uterina sometimes prolapses or subsides before the sætus in the time of parturition."

Deventer † relates, that when the Placenta is detached from the Uterus, it is usually found at the orifice, to which it descends by its weight, "ou fon poids l'entraine;" and in another place he calls it, "la chute," the falling down of the Placenta.

Giffard ‡ has more than twenty cases, where the Placenta was found at the Os Uteri, but he plainly supposes that it

[|] Practical Observations in Surgery and Midwifery, English Translation, 1751.

[†] Observations importantes sur le Manuel des Accouchemens, traduites du Latin, 1734.

¹ Cases in Midwifery, 1734.

had not been originally fixed there, for he fays, "It is cuftomary in floodings to find the Placenta funk down to the mouth of the womb."

Smellie, in his first volume of Midwifery, more than once mentions the possibility of the Placenta being fixed to this part of the Uterus, and in his third volume, describes several cases in which it was there fituated; but there are no practical inferences drawn from them, nor, in his directions about the management of floodings, are there any rules given relative to this fituation of it.

THERE is a fimilar case related by Dr. D'Urban, in his elegant Latin Differtation on the Hæmorrhagia Uterina, which he evidently confiders as a most unufual one; for, speaking of the Placenta being there fituated, and thereby producing the flooding, he fays, "fingularem Hæmorrhagiæ hujus caufam fuiffe."

Dr. Hunter, in his beautiful engravings of different views of the gravid Uterus, lately published, has one, in which the Placenta was found at the Os Uteri, and had been the cause of a fatal flooding *.

More authorities might still be produced, to prove that the Placenta has been often found in this fituation, but these are sufficient; and I have not the least doubt, but in all of them it was originally fixed to the Os Uteri; it is possible, indeed, if the womb open with unufual facility and quickness, and the

* Since I wrote the above, I have feen Levret's Treatise on Midwifery, written in French, and published at Paris a few years ago, and on the subject of stoodings, he has a chapter to prove that the Placenta may be fituated on the Os Uteri without having been previously separated from fome other part, and pushed down there, he illustrates what he fays by four cases, two of which were under his own care, another was communicated by a friend, and the last was an account of a dissection of a gravid Uterus published in the History of the Royal Academy of Sciences at Paris in 1723, in which the Placenta was found there fituated, and had been the cause of an Hæmorrhage which proved mortal.

woman, through a peculiar constitutional strength, be able to support the loss of blood which must necessarily be produced by it, that the Placenta may become wholly detached; and its having been fometimes found lying loofe there, is, undoubtedly, the reason why it has been supposed to have been separated from some other part of the womb, and to have fallen down into that fituaation: the impossibility of fuch a circumstance, will, however, be very evident, if we confider the anatomy of the gravid Uterus; for the Spongy Chorion*, which, by an universal adhesion, connects the membranes to the Uterus, and which is an expansion of the surface of the Placenta, must effectually prevent the latter from changing its place, whilst the former remains unseparated, which

^{*} This fine cellular substance, which is the connecting medium between the Uterus and the Chorion, and through which an infinite number of vessels ramify, was described by the late Dr. M' Kenzie, under the name I have used, but it is sometimes called the Membrana Cribriformis, and I find Dr. Hunter, in his anatomical plates of the gravid Uterus, gives it the name of Decidua.

I am convinced from feveral diffections it always does till the expulsion of the child; indeed, there must be a partial separation of this membrane, in the space between the Placenta and the Os Uteri, to allow of the discharge of blood into the Vagina, but there must be an entire separation of it, above as well as below the Placenta, to admit of its falling down, which, I should suppose, could never take place before the delivery of the child.

It may appear extraordinary, that a circumstance, attended with so much danger, and which seems to be so frequent a cause of the Uterine Hæmorrhage should have hitherto been so little noticed, for though, in the cases which have just been alluded to, the Placenta was found at the Os Uteri, yet it was, in very sew of them, supposed to have been originally sixed there, and I make no doubt but it has often happened when it has not been known at all to the surgeon, as I am induced to believe that

UTERINE HÆMORRHAGE. 25

that in the greatest number of those instances, in which the women have died undelivered, the floodings have been produced by this attachment of it: but this is eafily accounted for, when we confider, that it is very rarely that a furgeon has an opportunity of opening the gravid Uterus after death, that there are very few fymptoms in the course of the complaint, which might lead a perfon unapprized of its frequent occurrence, to fuch a conjecture, and that in the early part of labor, when the Uterus is high in the Pelvis, and the Os Tincæ is very little open, it is not discoverable by the ufual mode of examination: we may also add, that in those few cases where there has been fufficient strength to admit of its being completely open, the Placenta must have been found loofe; and, moreover, which is perhaps the principal reason, that the number of floodings which happen, when compared to the number of labors, is fo fmall, that very few must come under the notice of those who are engaged only in private

private practice, not enow probably in their whole lives to draw their attention fufficiently to the fubject, or to make them competent judges of it.

ADMITTING, then, that floodings are produced by these two different caufes *, and that they require a treatment

* La Motte relates a case of flooding, in his 214th obfervation, produced by a cause different to either of those I have mentioned, and which, probably, would be more dangerous than that which arises from the accidental cause, as it would continue as much during the prefence as the absence of pain, nor would it cease till the child and Placenta were removed, which, unless the progress of the labor was quick, might produce a loss of blood sufficient to destroy the woman; the case I allude to is a rupture of one of the umbilical veffels, owing to the cord being feveral times twifted round the child's neck, and to its texture being remarkably weak and knotty: but as this is fo very extraordinary an inftance, that it, perhaps, may never occur again, and as it would be, moreover, impoffible to know the cause of it till the labor was finished, I think it deferves not to be confidered amongst the general causes of the Uterine Hæmorrhage.

Another circumstance may likewise happen to prevent the expulsion of the child by nature, even when the flooding is not produced by the attachment of the Placenta to the Os Uteri, I mean, when the Pelvis is fo badly formed that the head cannot pass in the usual time: Mr. Aikin has favoured

fo widely different, we cannot be at a loss when such occur to us, and we have discovered the particular cause from which they arise, how to act; as, in the one case, we shall be encouraged to wait, and make use of such means to restrain the discharge, as will be more particularly mentioned hereafter, and in the other, we shall not hesitate to have recourse to delivery by art: for it is very obvious, that the want of fuccess which has fo often attended the turning the child, when fuch has been thought necessary, is to be attributed to the operation having been too long delayed, rather than to any real danger that at-

me with a case somewhat like this; "he was sent for to a woman who had slooded pretty much, the membranes were broken and the Funis was coming down into the Vagina, he immediately introduced his hand to turn, when he sound the head in the passage, which by a pain was forced pretty low, and he thought the labor would soon be over, but after waiting two or three hours the head continued where it was, and the slooding returned at times; the woman being now very weak and the child certainly dead, from the obstruction of the navel-string, which was pushed down, he opened the head, delivered, and the patient did well."

tends the cautious performance of it; as if it be not attempted, as usually happens, till the woman be well night exhausted, it must certainly be a doubtful matter whether she lives through the operation, or if she survive that, whether the debilitated Uterus will be able to contract itself when its contents are removed, fo far as to put an entire stop to the discharge: the chances under fuch circumstances being, then, so unfavourable, no wonder that the most cautious and skilful turning of the child has fo feldom been attended with fuccefs.

THE fuccess of turning depending, therefore, entirely upon its being done before the patient has loft too much blood, it is a matter of the utmost importance to obtain an early knowledge of the necessity there is of doing it, or in other words, to know at the beginning of the discharge, whether it be produced by the Placenta being fituated on the Os Uteri or not; which is the fecond

fecond circumstance I before considered as essentially necessary to enable the surgeon to practice with certainty in these cases, and concerning which I shall now endeavour to give some directions.

THERE is, perhaps, fome difference to be observed in the time and manner that floodings, produced by these different causes, come on; probably, that which is occasioned by the Placenta being fixed to the Os Uteri, will, for the most part, not come on till the full term of parturition, when the Uterus begins to dilate from the approach of labor; the other, which is owing to some accidental separation of the Placenta, may, on the other hand, come on before labor begins, and indeed at any time during pregnancy, and possibly, were we to be very nice in out enquiries, it might be accounted for by the patient's having received fome external injury, having fuffered by a fever, or undergone fome fudden and confiderable fright,

fright, &c.—but as these, with other symptoms that might, very likely, be enumerated, are, at best, but vague and equivocal: and as also, though the Placenta be situated on the mouth of the womb, it may, nevertheless, sometimes be separated by the same accidental means which detach it, when otherwise situated, the only certain knowledge respecting its situation, is to be derived from an examination of the Uterus by the touch.

For this purpose, however, the usual method with one singer will not always suffice, but the hand must be introduced into the Vagina, and one singer infinuated into the Uterus*; for in several of the sollowing cases it will appear, that

^{*} I have lately had an opportunity of seeing an accurate copy of Dr. Young's very excellent Lectures on the Theory and Practice of Midwisery, and though he takes no notice of this singular situation of the Placenta, he advises in flooding always to examine the state of the Uterus, by introducing the hand into the Vagina; the reason he gives for it is, that the coagulated blood which is usually found in the passage, renders it impossible to

though the women were frequently examined in the ufual way, the Placenta was not discovered till the hand was admitted for the purpose of turning the child: if this be done flowly and cautiously, and the hand be properly lubricated, it will feldom give the patient much pain, but if it should give some pain, as it is to obtain information fo effentially necessary to her fafety, that ought never to induce us to omit doing it, or to do it imperfectly: if the Placenta be at the mouth of the Uterus, it will be immediately felt by the finger, and may be distinguished from the membranes, by its greater thickness, and from coagula of blood, by the irregularity and roughness of its interior furface, which will then prefent to the finger.

IT must be acknowledged, indeed, that it may sometimes happen, that, at

feel the Os Uteri with fufficient distinctness by the finger alone: if, then, it be right when the situation of the Placenta is not an object of enquiry, the propriety of my recommending it above must be very obvious.

the

the very first coming on of the complaint, if the discharge be small, and more especially if it be the patient's first child, and the parts be close and unyielding, the admission of the hand into the Vagina, as I have directed, will be attended with the utmost difficulty, and, perhaps, be almost impracticable:-in this case let us wait (but let it be with the patient) till the discharge increases, or has continued long enough to relax the parts, for certainly, if the woman be able to bear losing a little blood, which at first she may fafely do, the examination will be thereby rendered more eafy, and the turning the child, if necessary, be more practicable and fafe.

Supposing, then, that the Placenta should, from this enquiry, be found at the mouth of the womb, the surgeon will be at once convinced of the danger that must unavoidably attend delay, from the impossibility there will be of affording the woman relief by any other

means than the timely removal of the child, and will, on that account, not hefitate to deliver before too great a loss has been sustained.

In recommending early delivery, I think it right, however, to express a caution against the premature introduction of the hand, and the too forcible dilatation of the Os Uteri, before it is fufficiently relaxed by pain or discharge; for it is, undoubtedly, very certain, that the turning may be performed too foon as well as too late, and that the confequences of the one may be as destructive to the patient as the other. I am particularly led to observe this, as I have lately been informed, from very good authority, (namely, a gentleman to whom one of the cases occurred,) of three unhappy instances of an error of this fort, which happened, fome years ago, to three furgeons of established reputation, who, from the fuccess they had met with in delivering feveral who

were reduced to the last extremity, were encouraged to attempt it where but very little blood had been loft, in hopes that their patient's constitutions would suffer less injury, and their recovery be more fpeedy; which, till the experiment was made, was a very reasonable supposition; the women died, and they feemed convinced that their deaths were owing to the violence of being delivered too foon, and not to the loss of blood, or any other cause.

IT becomes, then, necessary to endeavour to ascertain, with a degree of accuracy, the precise time when we may proceed to deliver, without fear of incurring the ill effects either of precipitancy or delay.

Ir has been advised, never to introduce the hand till nature has shewn fome disposition to relieve herself, by the dilatation of the Os Uteri to the fize of a shilling, or a half crown, and this rule is certainly founded on a rational principle

principle, for when it is fo much dilated, there is no doubt but the turning may be easily and fafely effected; but from fome of the annexed cases it appears, that a dilatation to this degree fometimes does not take place at all, and that even when the woman is dying from the great loss of blood, the Uterus is very little open; the reason for which feems to be, that when the discharge has been confiderable, and more particularly when much blood has been fuddenly loft, fuch a faintness is brought on, that though the Uterus be totally relaxed, and might therefore be opened by the most gentle efforts, yet nature is unable to make use of those efforts: and, moreover, if there be flight pains, the adhesion of the Placenta to the internal furface of the mouth of the womb, counteracts their influence, and thereby hinders its giving way to a power, which would otherwife, probably, very eafily open it.

Ir

IT appears, then, that this rule, if invariably adhered to, would, in some cases, be attended with danger, as we might wait for the opening of the Uterus, till it was too late to relieve the woman by turning the child; and for this reason it feems right, that we should fometimes be as much influenced by the Os Uteri being in a state capable of dilatation without violence, as by its being really open; when this is the case, therefore, if the woman's fituation demand speedy affistance, we should not hesitate to attempt delivery, even though to the touch the Uterus feem quite shut, more especially as in making the attempt, we shall know, before we can possibly have injured the Uterus, whether it be fafe to proceed; -if the womb readily give way, and the hand pass with eafe, we may be certain no harm will follow, and may, on that account, confidently profecute the turning; but if, on the contrary, there immediately come on a contraction of the Os Uteri, that, in a purse-like manner, tightly furrounds

rounds the fingers, it will prove difficult, and we ought therefore to defift, and wait till the part be more relaxed by pain or discharge, as difficulty, in these circumstances, is the truest criterion of danger.

As an encouragement that we may fafely fuffer a woman, under fuch circumstances, to lose more blood, the contraction may certainly be looked upon as a proof, that there still remains a confiderable portion of animal strength, and that she has not been so much affected by the loss, as we before imagined; and if we can fo far moderate the discharge, as to prevent the blood from being too fuddenly loft, which, in fuch a case, it must be our endeavour to do, a very confiderable quantity may come away without endangering the life of the patient. But in waiting for a further relaxation, we ought by no means to leave the woman, not even if the flooding, from the means we have used to moderate it, be totally suppressed; as when

when the Placenta is here fituated, the Hæmorrhage will fometimes return fo fuddenly and profusely, that if the furgeon be not at hand immediately to bring away the child, the woman perishes in a very little time *.

* THE case of the King's coachman's wife, related by Smellie in his answer to Douglas, is a striking proof of the danger of leaving a patient in fuch a fituation. woman had flooded feveral times from the middle of the feventh month to within a fortnight of her full reckoning, at which time it increased much, she had slight pains, and the Os Uteri was found open to the fize of a fixpence, beyond which was a foft fubstance that felt like the Placenta; as the dilatation was, he thought, infufficient to admit of delivery, he determined to wait; the advice of another physician was taken, who concurred with him that it was proper to wait till "those pains fhould bring on right labour," they therefore left the patient; but in a few hours after he was again fent for, when he found her in fuch extreme faintness, that she expired foon after his arrival: the body was opened, and the Placenta was found at the Os Uteri.--- It is observed, indeed, that a trial was then made to open the mouth of the womb, but it was not effected without much difficulty and a laceration, fuch an accident happening however after death, (when every strong membranous part is equally incapable of contraction and extension,) is no proof that if the most favourable opportunity had been watched for, and a gradual and repeated endeavour to

To steer fafely, then, between the two dangerous extremes, it appears necesfary that, on the one hand, we should never deliver till the dilatation of the womb can be effected by eafy means, and, on the other hand, when it has been fufficiently relaxed by discharge, if the woman have fuffered much by it, that we should no longer defer it, notwithstanding, from the absence or inefficacy of pain, the Os Uteri should remain unopened; yet, after all, as turning feems to be only necessary when the Placenta is fixed to the mouth of the womb, and that circumstance can feldom be known till the hand be introduced into the Vagina, and one finger be infinuated into the Uterus, I should

open it had been before made, it would not have fucceeded: their determining to wait "till right labor should come on," and leaving their patient without apprehending its bringing on an increase of the discharge, proves their not having thought about the Placenta, and their not being aware of the unavoidable consequence of such a situation of it; and I have related the case as much to prove this, as to exemplify the danger of leaving a patient under such circumstances. imagine it not very likely that we should often be in danger of injuring the patient by premature delivery, as when the hand passes easily into the Vagina, I should suppose there will be seldom much difficulty in its admission into the Uterus.

In introducing the hand for the purpose of turning, when the Os Uteri has been carefully dilated, if the separated part of the Placenta be immediately prefenting, it is best to endeavour to pass the finger through the fubstance of it, and by degrees with other fingers to enlarge the opening, till the hand can get through it into the cavity of the Uterus: the obvious reason for this is, that by this means not more of the Placenta may be feparated, than is necessary for the introduction of the hand, and confequently that as little increase of bleeding as possible may be produced by the operation; but if it be impracticable, as I have more than once found it, and it must ever be when the middle

of the Placenta prefents to the hand, from the thickness of it near the Funis, it must be carefully separated from the Uterus on one fide, and the hand paffed till it gets to the membranes, which being eafily broken, it is admitted into the bag, the floating fœtus is turned, and the delivery finished, as in preternatural positions of the child; except, that in this case the extraction should be more flow, that the Uterus may not be unable to contract, by being too fuddenly emptied: a moderate pressure from the hand of an affistant, upon the Abdomen, as the child is coming away, will likewife be of use to affift the contraction. The Placenta being at the Os Uteri, and being ufually feparated more by the introduction of the hand, commonly comes away immediately, but if a part of it should remain adhering, and the difcharge continue, it should be carefully removed, and as it is fo near, it may very eafily be done.

IF, on the contrary, it be clear from this careful examination of the Uterus, that the Placenta is not at the mouth of it, and that the coming on, or increase of labor, will not of necessity increase the discharge, provided it be not very profuse, (for let it be remembered, that I am fuppofing the examination to be made early, and before any confiderable quantity of blood has been loft,) it certainly will be proper to wait for the natural pains, and, in the mean time, to use fuch methods as are likely to restrain the flooding, which are, the admitting a free circulation of cool air into the room, keeping the patient in an horizontal pofture, giving her anodyne, with Tinctur: Rofar: &c. and fupplying her frequently with fuch cool and fimple nutritious drinks as will support her without quickening the circulation; from pursuing this method it will often happen, that the discharge goes off entirely, and if the woman be not arrived at her full term, and she be kept very still and calm, that it does not return before labor comes on. But if it should still continue, or return frequently, it will be right

right, if possible, to bring the Uterus into a state of contraction, by exciting fome pain, which may often be done by gently irritating the Os Uteri with the finger; if this fucceed, and the mouth of the Uterus be thereby fo far dilated, that the diftended membranes may be felt, they must be immediately pierced by paffing a probe along the finger, as upon the discharge of water thus produced, the womb necessarily contracts to a certain degree, and the flooding proportionably abates; this is, for the most part, foon fucceeded by flight pains, which if the child prefent fair, have very foon an effect upon it, and push it down *.

In the then relaxed and inelastic state of the Uterus, it is astonishing how

* This is the method of practice recommended by Puzos in his Memoire fur Pertes de Sang, which if confidered as relating only to cases produced by an accidental separation of the Placenta, is certainly an excellent one, and these are the only ones that seem to have occurred to him, for he appears not to be aware of the Placenta being sometimes fixed to the Os Uteri, in which case, it

much it is influenced by a trifling degree of pain, dilating and giving way to the most gentle throws; infomuch that, in these cases, the child usually passes with half of the ordinary efforts of nature: it is likewise remarkable, that the discharge commonly abates upon the coming on of pain *; which

is plain, his advice must be dangerous. The success that attended the management of his cases, which were certainly produced by accidental causes, may, I think, serve to strengthen what I have ventured to declare as my opinion, that when fuch is the case, it will for the most part, terminate fafely by the fole affiftance of nature.

* The Fundus and sides of the Uterus being in a flate of contraction during the presence of pain, press upon the Placenta, and lessen the flux of blood into the womb; moreover, when the water is escaped, the child's body comes in contact with the Uterus, and the Placenta may likewise be pressed upon by it, so as to have its veffels stopped, and these are, without doubt, the reasons why it is observed that the flooding usually abates whilst the pain continues, but this must obviously be only when the Placenta is fixed to any part but the Collum and Os Uteri, in which case the reverse must happen, as those parts are dilated during pain: it may be of use to attend to this circumstance, when we cannot so soon as we could wish, make a manual enquiry into the cause of the flooding.

proves the propriety of endeavouring to excite it by the means before-mentioned, when the other methods used to restrain the slooding do not succeed, and from this circumstance, too, the early examination with the hand in the Vagina, and one singer in the Os Uteri, is not only useful when the Placenta is there situated, but from the stimulus it excites, is of service to bring on pain, and facilitate labor when it is not so situated.

But if, notwithstanding this mode of treatment, the discharge should not leffen, if the evacuating the waters should not abate it, and if, moreover, labor pains, sufficient for expelling the child, should not succeed, and the flooding should still increase, so as to endanger the life of the patient, I should imagine it hardly necessary to say, that even in this case, as well as when the Placenta is sixed to the Os Uteri, the only certain method of stopping it should be used, namely, the delivery of the child by turning; for though

though I have never yet met with a case that under fuch circumstances has required it, and believe fuch very rarely happens, yet I would not be supposed to fay fuch a one cannot occur, as the feparation of the Placenta may, for instance, be produced by fuch violence done to the Abdomen, and the Hæmorrhage may be fo profuse, that nothing but a fpeedy delivery by art will put a flop to it. I only mean, that when we are called in early to flooding cases, if we judge only by the quantity of blood that has been loft, which may be fmall, and the present strength of the woman, which may be confiderable, we must frequently be deceived in our judgment of the cases, and be in danger of using a wrong method of treatment, but that the knowledge of the causes which produce them, will in the one case, for the most part, justify our waiting; and in the other, will invariably prove the propriety of turning the child.

THE want of fuccess which has so often attended turning in flooding cases, has, however, induced some people to believe there is danger in the operation itself, and that, independent of the time and manner of its being performed, the mischief in part arises from that: among others, Puzos raifes objections that have a tendency to discourage it; he draws. indeed, a very nice comparison between the influence that natural, and artificial labor have upon the Uterus, and feems thence to infer, that the injury done to it by the latter, is very often the reason why it is unable to contract itself after the child and Placenta have been removed: there can be no doubt but that the womb ever fuffers more from art than from nature, as the latter is more gentle, flow, and regular in her efforts to expell the child, than the former is to bring it away; but he certainly goes too far in attributing fo much mifchief to the operation by art, as if the want of contractile power in the Uterus were owing to the mere mode of delivery, it would very often turn out so in preternatural labors, where the improper position of the children renders the introduction of the hand to turn them necessary, in which too the Uterus being more rigid than in slooding cases, more force is requisite to effect it, and consequently more violence is offered to the womb; but every one's practice in these cases contradicts it; furely the obvious reason then for the want of success is, in most cases, what has been before observed, that the delivery has been too long deferred, and the woman too much exhausted by the great loss of blood.

It has been likewise urged by some, as an additional objection to turning, that in these cases there is, for the most part, such an insensibility of the Uterus, that as nature is, on that account, unable to expell the child, she will, for the same reason, be unable to contract the womb, if delivery be effected by art, and, independent of the injury which turning may do to the Uterus, that all attempts

attempts to stop the discharge, will, for this reason, be ineffectual; but I should fuppose this want of fensibility to be owing to the same cause, the loss of blood; for when the Placenta prefents to the Os Uteri, (which is the case we consider as requiring turning,) no wonder nature is unable to expell the child, as every effort she uses to dilate the womb for that purpose, must separate the Placenta, produce an increase of bleeding, and proportionably leffen the vital power; fuch an idea, therefore, which feems to be an unjust one, ought never to induce us to omit using the only certain method of stopping the slooding, and thereby of preventing that infenfibility, which a further loss of blood alone occasions.

To many practitioners, moreover, the introduction of the hand to turn the child, is a very difagreeable operation, and if they have not been much used to it in cases where the Uterus is but little open, it appears a very difficult and formi-

formidable one: it were to be wished, that even this circumstance had never an improper influence upon surgeons, especially those who are young in practice, and that they were never induced to omit, or too long delay this operation, because they feel unwilling to do it.

But it is not fo difficult as many imagine, for even in preternatural cases, where the Uterus is strong and rigid, and gives way reluctantly, if the hand be flowly and gradually introduced, it will feldom be found impracticable, provided the Pelvis be not badly formed; and in floodings it is effected with peculiar eafe, which should be a further encouragement to attempt it in fuch cases; for as the Uterus necessarily becomes fo relaxed after a confiderable lofs of blood, it very readily gives way to the admission of the hand, that tight contraction of its neck, which in other cases is such an impediment to the introduction of it, being here feldom to be met with; and it may be added, that

in proportion as nature, from the loss she has sustained, is less able to bear violence, happily, a proportionable less force is requisite.

If, therefore, the operation may be performed without much difficulty to the furgeon, if the cautious performance of it be attended with no danger to the patient, and if the becoming early acquainted with the necessity there is for it, give us an opportunity of doing it before the woman has lost too much blood, and before the Uterus has thereby been too much deprived of its fenfibility and power of contraction, if, likewise, that early knowledge may be obtained by purfuing the directions I have given, the turning the child, in the cases I have mentioned, cannot, furely, be too much urged to practitioners; as it is highly reasonable to expect more frequent fuccess, when it is done under more favourable circumstances, if it be certain, that fuccess sometimes attends it, when the patient is in appearance at the last extremity. IT

IT will, however, frequently happen, that we shall not be called in till late in these cases, when the woman will probably be in appearance dying, and indeed, fometimes the fluor is fo rapid, that in a very little time fo much is loft, that the patient finks immediately; but, as was just observed, unexpected success having sometimes attended turning. even under the most unpromising circumflances, it is certainly always our duty to do it, and by that means give the woman the only possible chance. I know there are arguments to be used, which may feem to justify a furgeon's relinquishing his patient under fuch melancholy circumstances; that as people fo frequently judge by the event only, he may incur blame, and his reputation undefervedly fuffer, if it terminate badly, as it is most likely to do; these may be tolerable arguments in trade, but they are very unjustifiable ones in morals, which direct us always to do what is in itself right, independent of the opinion of the world, and the confequences

fequences that may follow it. But I cannot fee much reason to apprehend much injury to our reputation, if, previous to the performing a dangerous operation, the uncertainty of the event be but properly represented to the friends of the patient, or if, before our doing it, we fend for some furgeon of established reputation, to justify our opinion, and to be prefent, and perhaps affift, whilst we do it; which is of all others the most effectual method of preventing any injury to our character; and in places where the gentlemen of the profession behave liberally to each other, there can be no difficulty in procuring fuch affistance.

Thus, I have ventured to place one of the most important subjects in midwifery in a new light, and have endeavoured to establish a hitherto uncertain practice upon principles that are more fixed and constant, by ascertaining when we may with propriety leave nature to do her own work, and when it will be requisite

requifite to proceed to immediate delivery, by turning the fœtus: I have also endeavoured to fix the precise time when it may, with most fafety, be done, and, in order to promote the practice of turning, when fuch becomes necesfary for the woman's fafety, have attempted to obviate the objections which have been made to this operation, from a supposition of its being either difficult, dangerous, or useless. And, from what has been faid, it appears, that the Placenta is fixed to the Os Uteri much more frequently than has hitherto been fupposed; that when it is so situated, nothing but turning the child will put a stop to the flooding; that when it is not fo fituated, nature will, for the most part, expell it fafely herfelf; that an early knowledge of this circumstance is of the utmost importance; that it may be obtained with ease and safety; and that, therefore, it should, in every case, be enquired into before much blood has been lost; that the information, procured by making fuch an enquiry, should govern

govern our management of the case; if we find the Placenta at the mouth of the womb, that we should proceed to delivery; that, if it be not fo fituated, if the discharge be not very profuse, and a very large quantity of blood have not been already loft, we should endeavor to restrain it by the means commonly directed for that purpose, and wait for nature's affiftance in the expulfion of the child: and it is thence evident, that this practice will have an advantage over the uncertain one hitherto adopted, because our determination about what we do, will ever be more fafe and fatisfactory, for if, on the one hand, we wait, we shall have the satisfaction of knowing that, in all probability, nature will be able to expell the fœtus; and if, on the other hand, we immediately turn the child, we shall, also, have the fatisfaction of knowing that nothing but turning can relieve the woman, and that, therefore, we do not give her unnecessary pain: and, finally, that our doing it before the patient has fustained too

too great a loss of blood, will make the chance of success more probable, and thereby be the means of preventing, in some measure, the fatality which has hitherto so frequently attended these cases, and which has, perhaps, been more owing to a rational method of treatment not being known, than is commonly imagined.

allopted, becaute due determination a-

description of the property of the party of

production and inguistion latisfactions

and if on the other bank, ave into al-

Talest ted type world to not finish

named of the site of the same

trattent ban : atantant

CASES.

THE subjects of the following cases were most of them poor women, under the care of midwives when I was sent for to them, and had been slooding a considerable time before I saw them. As they may, on that account, be justly considered as labouring under every disadvantage, none, I think, could better exemplify what I have ventured to advance in the foregoing essay.

CASE I.

FCEMBER 29, 1769, I was fent for, in the afternoon, to the wife of —— Balls. She was at the latter end of the eighth month of pregnancy, and had been feized the preceding evening, with a discharge of blood from the Uterus;

Uterus; it began without pain, and in fmall quantity, but increased by degrees, and was confiderable when I faw her; fhe had now, however, fmall pains, by which the Os Uteri was already fomewhat dilated: I admitted as much cold air as I could into the room. fupplied her frequently with cool and nourishing drinks, and as the pains still continued, waited till the membranes were fo far distended and pushed down, that I could fcratch them with my nail, by which means I broke them, and let the water escape; the discharge immediately leffened, the pains increased, and, in a little while, I felt one foot of the fœtus presenting; I brought it down, and with great ease drew forth a small dead child. The Placenta came away in about a quarter of an hour, the flooding became less and less, and the poor woman, though much reduced by the loss she had fustained, recovered in the ufual time.

CASEII.

FEBRUARY 6, 1770, — Stannard. She was a finall delicate woman, of a fickly relaxed habit, and had born feveral children. About the end of the eighth month, a flooding came on, without any previous pain, or fymptoms of labor. I faw her in the evening, after the discharge had been several hours, though as yet it had not been very profuse; she was, however, very faint and languid: by keeping her upon the bed with but few cloaths upon it, and admitting cool air into the room, it evidently leffened: I found the Os Tincæ relaxed, and a little open; after examining feveral times, (and, probably by the flight irritation, occasioned by the frequent touch) it opened fomewhat more, and the membranes protruded fo far, as to be felt by my finger; I immediately broke them, the discharge abated still more, and some slight pains succeeding, she was, in about half an hour after

after the breaking of them, delivered, with remarkable ease, of a small living child; the Placenta was removed without trouble, the discharge was moderate, stopped at its proper time, and the woman perfectly recovered.

CASE III.

IN the morning of March 12, 1772, - Coufins, a fickly relaxed woman, who had born many children, was feized with a flooding in the latter end of the last month of her pregnancy. I was fent for upon the first attack of the complaint, and living near the patient, was with her before much loss had been fustained by it, though the Hæmorrhage was then confiderable. She was without pain, and I found, upon examination, that the Uterus was very little open: the room being very fmall, and the air in it too warm and impure, I immediately opened the door and windows, drew back the curtains of the bed, took off fome of the cloaths, and did

did every thing to render her cool, and to admit fresh air into the room, by which means the discharge considerably lessened: I gave her an anodyne, directed the coolest drinks, and lest her, desiring to be called upon return of either pain or slooding.

In the evening I was fent for again, when I found the latter had returned, and in an increased quantity, infomuch that the woman was extremely faint and languid; the Uterus was, however, now rather more open, and some slight pains were coming on, and upon examining whilst she had one, I was just able to perceive the membranes pressing against the mouth of the Uterus; I introduced the sharpest end of a probe along my finger, and broke them, the flooding became less immediately, and some pains following foon after, she was fafely, and with fingular eafe, delivered by them of a living child: the Funis being small and tender, broke upon the first gentle effort to draw the Placenta by it, but by waiting about half an hour, it descended far enough into the Vagina, for the fingers to get hold of it, and bring it away. The woman was very much weakened by the loss of blood she had sustained, but in a few weeks she perfectly recovered.

C A S E IV.

IN the evening of August 12, 1772, I was fent for to the wife of —— Leman, a pauper, belonging to the town; she had a midwife with her, who informed me her patient had been flooding in fome degree during the day, and that it had, in the last hour, very much increafed. I examined, found the difcharge confiderable, the Uterus was fcarce at all open, and fhe was without pain; she was, moreover, extremely faint, and feemed to have fuffered much more than any of the former patients: admitting cool air into the room, &c. as in the other cases, for awhile abated the discharge, but as it returned very foon,

foon, and the woman feemed in the most immediate danger, I was very defirous of attempting to relieve her by turning the child, but judging it right to have the opinion of another furgeon, I fent for one who has had confiderable practice in this city. He feemed to think my patient too far gone to receive relief from any attempt whatever to stop the flooding, and as he added, that it was his opinion, she would fink during the operation, as one or two had before done on whom he had made the like attempt, he advised me not to turn the child, His advice prevented my doing it immediately, though before his arrival it was my defign to attempt it, if he justified it: I was determined, however, not to leave the bed-fide, that if there came on the least degree of pain, fo as to allow me to feel the membranes, I might, as I had before done. pierce them with a probe, or if the flooding increased, and I found it practicable to introduce my hand, I still refolved

resolved to attempt the removal of the

By carefully attending to keep the room very cool, by preventing my patient from being the least stirred, and being myself her nurse, in giving her every few minutes small quantities of the coolest drinks, I prevented the difcharge from increasing, and at the same time supplied, as far as I could, the waste of what she did lose, by the drinks she took, being as nutritious as I could venture to give them, without their being irritating: after attending her in this manner about two hours, frequently examining and gently stimulating the Os Internum, there came on at length a flight pain, and foon after, I could just feel the membranes with the end of my finger; I immediately introduced a probe, in the manner I had before done, and broke them; it had the same good effect as before, for the discharge immediately stopped, and pain coming on, the Uterus opened, the head of the child

child was pushed down, and, notwithstanding the very alarming state she had, but just before, been in, she was soon easily, and safely delivered, by the natural pains, of a dead child.

CASEV.

NOVEMBER 5, 1772, about ten o'clock in the evening, I was fent for to - Middleton, a poor woman, who had been flooding a little the greatest part of the day; the discharge was about this time rather increased, which induced the midwife, who attended her, to defire my affistance; I found her without pain, and the Os Tincæ not the least opened; but the discharge was not so great as in the last case: I directed as in the former cases, gave her an anodyne, and left her with orders to be fent for again if the flooding increased, or if pain came I was fent for again about fix in the morning, she had slept in the night, and the discharge had been but little, but it was now very confiderable: na-

F

ture, however, feemed disposed to affist her, for there soon came on a sew pains which opened the Uterus, and distended the membranes so far as to enable me to break them; it had again the good effect of suppressing the slooding, and facilitating the labor, which terminated safely in less than half an hour; the child was dead.

C A S E VI.

Or not have I work to

DECEMBER 1, 1772, about midnight I was fent for to — Welden, another pauper. She had a midwife with her, who informed me, the woman had been flooding a confiderable time, and had loft a large quantity of blood, which feemed to be true, from the state the poor creature was in; for her faintness was extreme, and she had every symptom of the most immediate danger.

Upon examination, I found the Os
Uteri more dilated than in any of the
former cases, and the Placenta evidently
presenting:

prefenting: as no possible relief could, in this case, be expected from waiting, I at once resolved to give her the chance of an immediate delivery; which I effected by introducing my hand into the Uterus, turning and bringing away the Fœtus; and this I did with much greater ease than I could have imagined, as the resistance from the Uterus was very trisling; I endeavoured to pass my hand through the substance of the Placenta, but not being able to do it, I separated it on one side, till there was room for my hand to pass.

THE woman remained very faint and weak a long time after delivery, but being carefully nursed, she recovered by degrees, and was able to go out before the end of the month. This was likewise a dead child.

C A S E VII.

DECEMBER 29, 1772, about fix o'clock in the morning, I was called to F 2 — Free-

— Freeman, a poor woman, who was under the care of a midwife; she had been flooding many hours, and had lost an immoderate quantity of blood, was greatly sunk, and appeared to be almost dying; on examination, I found the mouth of the Uterus as open as in the last case, and the Placenta situated in like manner, which determined me to pursue the same method I had so successfully used in that.

The Pelvis was narrow and difforted, but I introduced my hand into the Uterus, and turned the child with all defirable eafe; the feet, body, and arms of the Fœtus I brought down in the ufual manner, and with no more than ufual difficulty; but when I came to the head, it remained fo fast betwixt the bones of the Pelvis, that, though I got one of my fingers into the mouth, (the face being towards the Sacrum,) and pulled the body, at the same time, with considerable force, I could not move it in the least degree, infomuch that the

Vertebræ of the neck began to give way; which made me defist from pulling fo forcibly, and induced me to send for the affistance of another surgeon.

He made feveral fimilar but unfuccefsful attempts; we, therefore, concluded, that nothing but lessening the fize of the head, by evacuating the brain, would allow it to pass; but to effect this was no easy matter; he thought it possible to pass the scissars through the Os Palati into the head, and attempted it; when the scissars had pierced the bones, I endeavoured to enlarge the opening, but could not do it; in tracing with my finger, round the head, as far as I was able, I thought there was a possibility of pushing in fome curved instrument behind the ear, at the lower end of the temporal bone; but the scissars being strait, I could not use them; however, from the looseness of the scalp, (for it ought to be observed, that,

that the child was dead, and almost putrid, which was certainly the reason why the Vertebræ of the neck separated so eafily, when I attempted to pull the head,) I thought I could push in the curved end of a blunt hook, which, with a good deal of difficulty, I effected, and by degrees infinuated it under the temporal bone; the opening I eafily enlarged by my finger, and with one blade of the forceps, fo that at length fome of the brain came away, the head was thereby compressed into a smaller compass, and she was delivered: but the extreme fatigue she had undergone by this unlucky difficulty, joined to the immoderate loss of blood she had previously sustained, was more than she was able to support, and she died the following morning.

SINCE the above case happened, I have procured a pair of sciffars of Cargill, in Lombard-street, curved at the points, (fomewhat like Tonfil fciffars) which may

may easily be used where it is found necessary to open the head, after delivering the rest of the child.

C A S E VIII.

JANUARY 4, 1773, —— Bygrave, a pauper, fell in labor, and a flooding came on, after having had some pain; I was sent for about an hour after the discharge began, and sound it very considerable; but there had been pain sufficient to dilate the Uterus, and to enable me to break the membranes, when I sound a hand and soot presenting; I immediately brought down both seet, and delivered her, safely, of a living child.

CASE IX.

FEBRUARY 3, 1773. —— Clarke. She had flooded feveral times, and in confiderable quantities, in the last month of her pregnancy; but it was every time suppressed

suppressed by the means I have before related, and she went to her full time.

HER labor was a natural one, and at that time there was very little discharge, though it had been considerable a few days before. She was delivered by the natural pains, of a large, living child; but immediately upon the expulsion of it, there came on a most profuse fluor; I instantly introduced my hand into the Uterus, and was fortunate enough to remove the Placenta, with but little loss of time; the womb immediately contracted, and the discharge abated.

CASEX.

FEBRUARY 12, 1773, I was fent for to — Marshall, a poor woman in the workhouse, who was in her last month of pregnancy, and had been flooding about two hours; she had, in that time, lost a very great quantity of blood, and was so much sunk by it, that she died soon after I came into the room.

I HAD an opportunity of opening the body, the following morning: the membranes adhered univerfally to the Uterus, by the Spongy Chorion; I carefully meafured the water contained in the bag, and there were three pints of it: the child laid, with the head obliquely to the right fide of the Fundus Uteri, and the face towards the Spine; the hands were turned upon the face, holding each one of the feet, fo that the Podex would have presented; the Placenta was situated upon the Os Uteri, and a partial feparation of it, not bigger than a crown piece, was the cause of this fatal Hæmorrhage. Before the died, I examined with my finger, found the Uterus very little open, and did not feel the Placenta.

REMARK.

This case proves, that the Os Uteri sometimes does not at all dilate to the size that has been usually thought necessary for safe delivery, and that it is

not, therefore, always right to defer turning the child, in expectation of it: had I been with the woman fooner, I certainly should have attempted it, and as from the first, there had been a considerable discharge, in all probability the Uterus was so relaxed, that it might have been easily and safely effected.

C A S E XI.

MAY 15, 1773, I was fent for, in the evening, to — Maltward, who had fpoken to me to attend her; she had gone her full time, and some symptoms of labor came on, accompanied with a slight discharge of blood; I gave her an anodyne, directed her to be kept cool, &c. and left her, with orders to be sent for again if the complaint increased.

ABOUT four o'clock in the morning I was called to her again, and found the discharge was increased, but she had had but little pain; however, after frequently

quently examining, in about half an hour, the membranes began to fwell, and I was able to break them; the flooding abated, more pains came on, and the child defcended into the Vagina, with the breech prefenting; but, as the woman was well formed, it paffed eafily, and fine was foon, fafely, and almost entirely by the natural pains, delivered of a dead child.

C A S E XII.

JUNE 25, 1773, I was called to ——
Sherwood, a pauper, under the care of a midwife. She had been flooding feveral hours, had loft a large quantity of blood, and was extremely faint. I treated her as I had done most of the former ones, gently irritated the Os Uteri, and, as soon as I was able, broke the membranes, and she was, in like manner, safely delivered, by the natural pains, of a dead child.

C A S E XIII.

JUNE 27, 1773. — Playford. She was attended by a midwife, at the time I was fent for, and had been flooding very much feveral hours; the discharge was still profuse, the Os Uteri quite shut, and from the faintness she was in, she was altogether without pain, though the complaint came on with labor pains: she had the most threatening appearance, and I very much feared I could be of no service to her, and intimated it to the midwife and the affistants, but added, that if there were any possible chance, it must be from immediately delivering her.

As they were defirous of another furgeon's opinion, I fent for a gentleman, who confirmed what I had faid respecting the danger the woman was in, and agreed with me, that the only chance she could have must be from a speedy delivery; the practicability of which, however he rather doubted, as the Os

Tincæ was fo little open: I apprehended great difficulty in doing it, and feared likewise, that if I succeeded in bringing away the child, the woman would hardly furvive the loss she had fustained; but having succeeded before, much beyond my expectation, I thought it right to attempt it; I introduced my hand into the Vagina, for this purpofe, and first one, then more fingers, into the Uterus, when, (and not before,) I found the Placenta fixed to the Os Uteri; I endeavoured to pass my finger through the fubstance of it, but was not able, though I tried fome time; I, therefore, separated it on one side, and got my hand compleatly into the Uterus; the head of the child presented, but I foon got hold of the feet, brought them down, and delivered with the same ease as in the last case of turning: the woman remained very languid a long while, and feemed hardly alive for many hours; but by supplying her frequently with cool and nutritious drinks, and carefully managing

managing her in other respects, she recovered entirely.

REMARK.

This case appears much to have refembled that of Marshall, who died undelivered; as the discharge was very profuse, and the Uterus very little open, the difference in the event being produced merely by my being fortunately called fooner to this woman. The eafe with which the turning was effected, and the fuccess which attended it, confirm the remark made to that case, that it is fometimes justifiable to deliver where the Os Uteri is not dilated to the fize of a shilling, or a half crown.

C A S E XIV.

JANUARY 1, 1774, I was fent for, about noon, to --- King, a poor woman, who was at the full term of her fourth pregnancy: without any previous accident, or complaint, she was seized with

with a trifling pain, like the beginning of labor-pains, which was accompanied with a little appearance of blood; when I first saw her, she was not at all faint, had lost a very small quantity of blood, and was sitting by the fire side; the Os Tincæ was a little relaxed, but not open, and she had no pain. I defired her to be laid upon the bed, to be kept very cool and quiet, and ordered the midwife to send for me again, if she found the flooding increase.

FROM this treatment it abated, and the woman got some rest in the afternoon; but in the evening her pain returned, and with it so prosuse an Hæmorrhage, that before I could get to her, she had lost an assonishing quantity of blood, and had the most threatening appearance; she, before, had a good pulse, and a slorid, healthy countenance, but nowherpulse were scarce perceptible, her countenance was pale, her lips livid, &c. from the extreme faintness she was now in, the discharge and pain were abated,

but as the Uterus was very little open, as she had no pain, and was so much sunk, there seemed to be no reason to expect assistance from nature, and no chance but from an immediate delivery.

I SENT for another furgeon, to justify what I did; he agreed with me, that it was the only chance of relieving her, though a very poor one; however, as he approved it, and the friends of the patient confented to it, I proceeded to turn the child. I found, as foon as I had introduced one finger into the Os Tincæ, that the Placenta was fituated as in the last case, and I was now able to infinuate my finger and hand through the fubstance of it, which, as I before observed, is an advantage; the child was in a natural posture, but I passed the head, got hold of the feet, and by them brought it away with all defirable facility: the woman immediately, and for fome time after delivery, appeared rather better, as she took nutriment, and

and was rather revived; but nature was unable to recover from the great loss she had sustained; and she died about six hours after.

C A S E XV.

-JANUARY 21, 1774, ——Bond. This was a patient of the same gentleman who was with me in the last case. He sent for me in the evening, and the woman was then, in the same dreadful state that my last patient was in, when I sent for him.

He informed me, that he had feen her the preceding evening, that she was then, to all appearance, in perfect health, but that she had been alarmed by a small discharge of blood from the Uterus; he bled her, gave her an anodyne, ordered her to be kept cool and quiet, and saw her the following morning: she had had a good night, and the appearance of blood was not more considerable, but in the evening it had suddenly increased to a

G

most

most violent degree, infomuch as to have reduced her, in a short space of time, to the deplorable state we found her in.

SHE seemed to be dying, but as we thought it justifiable to give her the only possible chance, by turning the child; by his leave, I introduced my hand, the Uterus was shut, though loofe and relaxed, and as foon as I got one of my fingers into it, I found the Placenta fixed to the mouth of it: in this case I could not pass through the substance of it, but separated it on one side, got my hand within the membranes, turned, and brought away the child, with the same remarkable ease as in the other case; but this was attended with no better fuccess than the former case, for she died in half an hour after delivery.

REMARK.

The unhappy events of the two last related cases, seem, at first, to contradict a remark made but a sew pages before; as it may, perhaps, be thought, that it would have been safer to have deferred the delivery longer, because the Os Uteri was so little open in either of them.

But fo far from the turning having been prematurely done, I am convinced its want of fuccess was owing folely to its having been too long delayed; for when I proceeded to deliver, the women had every fymptom of approaching death, and though my principal motive in turning was to give them a chance of recovery, as no other means could possibly do it, yet I was, likewife, further induced to attempt it, that I might be fatisfied of the fituation of the Placenta, which, about this time, began to excite my attention; and I well remember urging this to the gentleman who

was with me, as an additional reason why I wished to do it, as from their being so much reduced by the great loss of blood, he, at first, rather discouraged it.

The fingular ease with which it was effected in both cases, is, likewise, in my opinion, a positive proof that no mischief was brought on by turning; for, with respect to that operation, it has been before observed, that difficulty in doing it is the true mark of danger, and ease is therefore that of safety.

If I had not delivered, very probably I should not have had an opportunity of opening the bodies after death, and as, consequently, I should have known nothing of the Placenta, I could not possibly, with so much certainty, have made the distinction I have since done, between sloodings which are accidental, and those which are unavoidable.

IT may not be improper further to remark here, that in the management of the cases which have hitherto been related, I was influenced by no other circumstances than those which are usually attended to in floodings, namely, the quantity of blood which had been loft, and the feeming strength of the patient; for, though the Placenta being found at the mouth of the Uterus, was, in the cases of Welden and Freeman, an additional inducement to my turning, yet the idea of the frequent occurrence of fuch a circumstance, did not strike me till those of Playford, Marshall, King and Bond had fallen under my notice; nor, indeed, was I fenfible of the importance of an early enquiry, whether the Placenta was, or was not, fo fituated, till the two last cases happened, in both which it appeared fo evident, that, had its fituation been fooner known, the children might have been brought away before fo great a quantity of blood had been loft, and, very probably, the lives both of the mothers and the children have been faved. CASE

C A S E XVI.

JANUARY 24, 1774, I was fent for on the evening of this day, by the same gentleman with whom I had been in the last case, to --- Flood, who was at the latter end of the ninth month of pregnancy, and had, just before he saw her, been feized with a Uterine Hæmorrhage, but much blood had not been yet loft; I advised a careful and immediate examination of the Uterus, to know whether the Placenta was fituated at the mouth of it, and that the management of the case should be influenced by the information thence gained; this examination was accordingly made, and we were convinced that the Placenta was not there; we, therefore, directed an anodyne, ordered her to be kept still and cool, and left her.

FROM this treatment, the discharge for fome hours abated, but it returned in the morning, accompanied, however, with flight pains, which were increased

by gently irritating the Os Uteri with the finger, till the membranes fwelled, and gave us an opportunity of piercing them; more pains fucceeded, and the patient was at length fafely delivered by their fole affiftance.

C A S E XVII.

APPIL 7, 1774, I was called in the morning to — Howlett, a poor woman, who was feized with a flooding at the end of her ninth month: at the time I faw her, she had a midwife with her; I examined her in the same manner I had done the last patient, and being fully satisfied that the Placenta was not in the way, I gave the same directions as were observed in the last case; the discharge, in like manner, abated, and towards the evening, she was safely delivered by the natural pains, having no other assistant than the midwife, who was with her at first.

C A S E XVIII.

MAY 18, 1774, a very delicate relaxed lady, in the beginning of the ninth month of her first pregnancy, soon after getting her breakfast, had suddenly a slight discharge of blood from the Uterus, occasioned, as she imagined, by a smart shock of sneezing; she was immediately carried to bed, and I was sent for.

THE discharge continued, but was not increased in quantity, and, in about half an hour, I had an opportunity of examining, and sound the Os Uteri quite close and unyielding. As the Vagina was very strict, and there had not been discharge of any kind to relax it sufficiently for the admission of the hand, without considerable pain, and greatly alarming the patient, who had no apprehension of the danger of her situation, I ventured to defer the enquiry, and made use at sirst only of the methods commonly used to restrain it; but I thought

it right to flay with her: the Hæmorrhage was, thereby, kept from increasing many hours; but before the next morning, it became more formidable, fo that I thought it imprudent to remain any longer ignorant of the cause of it; I therefore examined in the way before directed, which was now done with tolerable ease; and finding that the Placenta was not at the mouth of the Uterus, I still waited for the natural pains, and endeavoured to excite them by the gentle stimulus of the finger on the Os Tincæ; in about two hours they came on, and pushed down the bag, which I foon opened; the discharge abated, and the head descending by repeated and gentle pains, she was at length fafely delivered of a dead child.

C A S E XIX.

JULY 14, 1774, about eleven o'clock in the forenoon, I was fent for to—— Fearman, with whom a midwife had been feveral hours; she was in her ninth ninth month, had a small but increasing discharge from the Uterus, was faint, and without labor pain: I examined carefully, without losing any time, and being convinced the Placenta was not at the Os Uteri, I gave her hopes of being released by the natural pains; to effect which I rendered the air as cool as I could, lessened the number of bed-cloaths, and endeavoured to excite pain, by gentle irritation on the OsTincæ; the discharge became less, but no pain coming on, I lest her, with directions to be called again on return of slooding, or coming on of pain.

In a few hours after, I was fent for, on the former account; I then repeated my efforts to make the Uterus contract, and at last so much pain came on as protruded the membranes, which, as before, I instantly broke; more pains succeeded, the Hæmorrhage stopped, and, to the surprize of the patient and her attendants, a few very trisling pains expelled a living child.

CASE

C A S E XX.

NOVEMBER 8, 1774, I was fent for to a poor woman at Ringland, about fix miles from this city: the messenger's account of her was, that she had been slooding, more or less, for several days; that the discharge had that morning increased immoderately; and, that the midwife who was with her, was greatly alarmed, and thought herself unable to relieve her. From this description of her, I told him, there was very little probability of finding her alive; and my conjecture proved true, for she had been dead about an hour before we got there.

The midwife who attended her, being an intelligent and communicative woman, I learned from her, that her patient, who was in her last month of pregnancy, had been as well as usual during the former months; but six days before, without any previous complaint, a slight

a flight discharge came on, accompanied with fome pain, which, however, stopped of itself, and the pain went off; that it had returned feveral times, and at each time in an increased quantity; that she had examined her at several different times, but found the mouth of the womb shut; and with the last attack. which was the morning of that day, and was more violent than any of the preceding ones, she had very considerable pains, and expressed such strong figns of bearing down, that she thought her near delivery; in that state she examined her, and then found the Os Uteri confiderably open; she removed feveral coagula of blood, which were in the Vagina, and at the mouth of the womb, and expected to feel the head of the child, but in its stead she found (to use her own words) a strange lump of stringy substance, unlike any thing she had ever before felt; the flooding still continued, the blood pouring forth with every pain, till the poor creature fell

fell into a fainting, from whence she did not recover.

I was defirous of opening the body, to be convinced of the fituation of the Placenta, but could not obtain the confent of the by-standers; there can, however, I think, be little reason to doubt its being at the Os Uteri, as nothing elfe could answer the midwife's description of a lump of stringy substance, or could hinder the descent of the child, when nature was affifted by fuch strong pains, and when the mouth of the womb was fo much open. I asked her, if she had ever before found the Placenta in fuch a fituation; she told me she had never, till now, been with a patient who flooded before delivery, though fhe had attended feveral hundred women; but she believed this could be nothing elfe.

C A S E XXI.

NOVEMBER 20, 1774, a midwife, with whom I have before frequently been, called upon me for my advice about a patient she was then going to, whose name was Bailey, and with whom she had been the day before, at which time there was a slight flooding, which had been then checked by keeping her still and cool, but she found it was at this time returned.

As I could not conveniently go with her, I defired her to be very careful in examining the patient, and to be fatiffied, whether there was any thing unufual at the Os Uteri, in which cafe I defired her to fend for me; otherwise, she might fafely wait the coming on of pains, in the mean time keeping her cool, and in bed.

SHE called on me the next day, to inform me she had pursued my advice, that

that the mouth of the womb was opening, but she found nothing there like what I had described to her; she, therefore, agreeable to my directions, waited, and, notwithstanding the continuance of the discharge, she was safely and easily delivered, by the natural pains, before the following morning.

C A S E XXII.

DECEMBER 16, 1774, —— Smith, a healthy, strong young woman, in the last month of pregnancy, sent for me in the evening, having had, for several hours, a discharge of blood from the Uterus. I immediately introduced my hand into the Vagina, and with one singer in the Os Uteri, which was soft and yielding, I imagined I found the Placenta, but, upon further examination, was satisfied that it was nothing more than a coagulum of blood, as I very evidently selt the membranes, with the head of the child behind them. The dilatation produced by this examination,

was fufficient to have rendered the introduction of the hand into the Uterus, to turn the fœtus, if it had been necesfary, very practicable; but being confident of the great probability, not to fay certainty, there was of nature's being able to expell the child in due time, I thought it right to endeavour to restrain the present discharge, and wait for pain. This was foon effected, by the usual means, and there feemed a probability of her going to her full time, if kept still and cool, for it stopped entirely for two days.

IT returned, however, on the third day from the first attack, when it was probably occasioned by some imprudent exercise in the woman, as she had not yet gone her full time; but being now accompanied with pains, the Uterus opened, the child was pushed down and expelled with remarkable eafe. The Hæmorrhage, in this cafe, rather increased after the delivery of the child, and did not abate till the Placenta was removed,

removed, which did not descend so soon as it usually does.

C A S E XXIII.

DECEMBER 21, 1774, I visited Mrs. L—, she was in her last month of pregnancy, and had had a slight discharge of blood from the Uterus the preceding day, but as it was stopped when I was with her, and she had no pain, I had not an opportunity of examining the state of the womb; I, therefore, only recommended stillness and rest, and defired to be again sent for, if it returned in the least degree.

On the 25th, I was sent for early in the forenoon, the discharge having returned, and being then considerable; I immediately examined, and, as in the last case, imagined I selt one edge of the Placenta on the posterior side of the Os Tincæ, but on tracing my singer compleatly round it, I distinctly selt the membranes, and the child's head be-

hind them; I ventured, therefore, to wait, and gave my patient hopes of being fafely delivered, without any other than the usual affistance; from cooling the air, &c. it again abated, nor was there any return of it during the remainder of the day, and the following night. About nine o'clock the next morning, the 26th, she was again alarmed by a return of it, and with increafed violence; upon examination, I again thought there was an edge of the Placenta in the mouth of the Uterus, but as it could be but a very small portion of it, and as there was fufficient paffage for the child, and, moreover, during the pains, which now came on, the discharge did not increase, but stopped; I thought it still safe to wait, and endeavoured immediately to break the membranes, which, with some difficulty, I effected; the discharge became less, but the pains went off, and by keeping her very cool and free from motion, the Hæmorrhage was inconfiderable all the day, and the following night.

On the 27th, about fix o'clock, it came on again, and in a larger stream than ever, and as her pains were still but trifling, and, from the frequent return of the flooding, she had lost a large quantity of blood, I began to apprehend danger, and almost to think I had done wrong in fo long trufting to nature; I was therefore, for fome time, embarraffed to determine what was now best to do, but it being again checked by increasing the cold air in the room, which had not been fufficiently attended to in my absence, I was once more encouraged to wait, and fortunately from this time, there was no great return of it; in about two hours the pains increased, the Uterus opened, and the head came forward, and though from its being large, and the Pelvis not a good one, the progress of the labor was much flower than I had ever before found it in fimilar circumstances, it terminated fafely by nature's fole affiftance before one o'clock; the Hæmorrhage was inconfiderable after delivery, and H 2

and my patient, though much reduced, and being in other respects a sickly woman, recovered entirely.

THERE was a peculiarity in the form and texture of the Placenta in this case that deferves notice, as, probably, the Hæmorrhage was, in some measure, occasioned by it; instead of the usual circumscribed and circular cake, thick in the middle, and becoming less and less towards the edges, it was an uneven mass, thinly, and, in some places, almost fuperficially spread over near one fide of the Uterus; the edges of it terminated in a broken manner, forming fomewhat like the lines of a very irregular island on a map, and one edge, making almost a detached lobe, hung down on one fide of the Os Tincæ, and was, I was now convinced, what I had before felt, and what had probably produced some of the flooding, but the principal difcharge feemed, by the discoloration of the Placenta, to have arisen from a separation of it higher up in the Uterus.

CASE

C A S E XXIV.

JUNE 19, 1775, I was fent for to -Hoole, a poor woman, under the care of a midwife: she had been flooding several hours, and, in the last half hour, the discharge had considerably increased. I immediately examined with my hand in the Vagina (for with the finger only I could but just touch the outside of the Os Uteri,) and found, by introducing one finger into the Uterus, that the Placenta was at the mouth of it: she had lost a considerable quantity of blood, and was very faint, but did not appear to have fuffered fo much as to have induced me, had the Placenta not been there, or had I made no enquiry to find it, to have turned the child; but being convinced of the danger of delay, I determined to deliver, and previous to my doing it, fent for a furgeon who had been before with me in some of the foregoing cases.

WHEN

When he came, I told him the woman's fituation, and defired him to examine in the common way, with one finger only, which he did, but could find nothing unufual at the mouth of the womb; I then defired him to introduce his hand, as I had done; he did this, and immediately discovered the Placenta; he, therefore, agreed with me in the propriety of immediate delivery.

I INTRODUCED my hand into the Uterus, and found the child lying in the natural posture, I passed the head, and with tolerable ease got hold of the feet, brought them down, and extracted a dead child.

The woman remained extremely languid for some time after delivery, but, notwithstanding this, and that she laboured under every disadvantage produced by extreme poverty, and a remarkable ignorance in her assistants, having neither pure air, clean linen, and

and hardly common nutriment for feveral hours, yet she perfectly recovered.

REMARK.

As this woman had not gone her full time, being, according to her own account, in the middle of the eighth month, which feemed true from the smallness of the child, I expected to have had some difficulty in introducing the hand, the Uterus not being compleatly distended, but, notwithstanding the mouth of it was but little open, it gave way with the same ease I have ever observed in these cases, and the delivery was effected with no more than usual trouble.

The necessity of introducing the hand into the Vagina, and admitting one finger into the Uterus, in order to distinguish with certainty whether the Placenta be at the mouth of it or not, has been before observed, and several of the former cases have proved the propriety

priety of it: in Playford, King, and Bond, it was not found till the hand was admitted for the purpose of turning, and in Marshall, though the usual examination was made before she died, it was not known till the Uterus was opened: in the present case, its not being discoverable by two different persons who examined at first in the usual way, is a very remarkable proof of the necessity there is for doing it; I, therefore, thought it right again to take notice of it in this place, as it may, perhaps, more strongly impress the reader.

C A S E XXV.

JULY 1, 1775, — Sherringham. In the course of the preceding week she had several times had a slight Hæmorrhage from the Uterus, which returning more considerably this morning, made her send for her midwise: it continued during the day, accompanied, however, with slight pains; but as they did not increase, the slooding continued, and she

she became faint, I was sent for about fix o'clock in the evening.

I IMMEDIATELY examined, as in the last case, and was satisfied that the Placenta was not at the mouth of the womb, as I very distinctly felt the head of the child presenting: I, therefore, waited, and endeavoured to excite pain, and, as foon as I was able, broke the membranes; her pains became stronger, the Os Uteri dilated, and the flooding entirely ceased, and I expected she would have been delivered with the eafe and quickness peculiar to these cases, but in this I was disappointed, for it proved very laborious, and the head descended into the hollow of the Sacrum fo flowly, that she was not delivered till one o'clock in the morning: it was a large living child, the Placenta was carefully removed, the discharge was trifling, and the woman perfectly recovered.

C A S E XXVI.

MRS. B-, the subject of the following case, is patient to the same gentleman who has favoured me with his affistance in several of the former cases: fhe is a woman of a very tender and delicate constitution, has been mother to feven children, and has had very bad health for feveral years past.

AUGUST 20, 1775, being at her full reckoning, she had the preceding day, and all the night, fmall pains in her back, which she considered as the forerunners of labor; about nine o'clock in the morning the pains fuddenly became stronger, and she felt a pressure upon the lower part of the Uterus, which was followed by a discharge; it was in a confiderable quantity, and she imagined it to be water, till, upon examination, fhe found it blood: fhe immediately fent for her furgeon, who came to her about ten o'clock; the pains had then entirely

entirely left her, but the discharge continued slowly trickling from the Uterus; as soon as he had an opportunity, he examined with the hand in the Vagina, and distinctly felt the Placenta at the mouth of the womb; he soon after sent for me, and I was with the patient about eleven o'clock.

SHE was still without pain, and the discharge was less in quantity, and of a paler colour: I examined in the same manner he had done, and found the Placenta in the before-mentioned fituation; the Os Uteri was dilated to about the fize of a shilling, but upon my attempting to introduce three fingers into it, it contracted fo closely round them, and was fo rigid, that I concluded the delivery would as yet be attended with too much difficulty to render it adviseable, and as there had not yet been bleeding fufficient to injure her, and it was now almost entirely stopped, we thought it right to wait till the parts were more relaxed by discharge, or dilated

lated by pain; but we staid with the patient, faw every cloth which had been used, and examined the Uterus alternately once in an hour.

ABOUT three o'clock in the afternoon there came on fuch pains as she before had in the night, but they were not fufficient to open the Uterus more, and therefore did not increase the bleeding: about five o'clock the pain fuddenly became more violent, the Uterus was more widely dilated by it, which producing a further separation of the Placenta, a fresh discharge was occasioned, and it was now rapid and confiderable; there was, therefore, every reason to justify immediate delivery, the Os Uteri was more dilated, it was more relaxed, and more yielding, and the Hæmorrhage was fo confiderable, that a further delay might have been attended with the utmost danger; he, therefore, proceeded to turn, which he did flowly, but with great ease, and extracted a living child.

THERE was no discharge of consequence immediately after delivery, nor whilst we staid with her, which was more than half an hour, but upon my calling again in the evening I found the room too warm, the bleeding had increased, and she was much more faint than she had before been; but from cooling the air, and adding a little pressure to the Abdomen, it abated, she had a tolerable night, and was nearly as well the next morning as she usually found herself after labor.

REMARK.

The happy event of the above-recited case, has afforded me peculiar satisfaction, as the management of it, from the beginning of the complaint, was in exact conformity to the directions I have given, and it appears to be a full proof of the propriety of them; for supposing it had been treated in the way commonly adopted, the situation of the Placenta would not have been known so early;

early; and supposing that circumstance not to have been thought of, as after the first discharge there was no pain, and the bleeding was inconfiderable, no one, I am perfuaded, would have fcrupled leaving the patient in the forenoon. The furgeon, in fuch case, probably would not have been fent for again till the evening, when the pain and Hæmorrhage returned fo fuddenly and profufely, and as some time must necessarily have been loft before he could have been with her, it is not unlikely but at his arrival he might have found her in the fame fituation, that, under fimilar circumstances, the women mentioned in cases No. 14 and 15, were found in, and might, therefore, have been unable to fave her.

It may not be improper, likewise, to observe, that the precise time for turning the child seems very happily to have been hit upon; for had we proceeded to deliver before the Uterus was sufficiently relaxed, there certainly would

would have been great difficulty in effecting it, and the woman might, very probably, have fuffered materially from the violence that must in that case have been used; and, on the other hand, if we had waited for a further dilatation of the womb, as that could not have taken place without an increase of the discharge, she, very likely, would have been unable to sustain the loss of blood, more especially as the extreme weakness of her constitution, and the bad health she had for some time past been in, rendered her a very improper subject for either extreme.

C A S E XXVII.

SEPTEMBER 16, 1775, about five o'clock in the afternoon, I vifited — Olley, a poor woman, who was near her full reckoning; she had a discharge of blood from the Uterus, which first came on about a month before, but it had never been profuse, as it abated by rest, and an horizontal posture, and returned only

upon any confiderable exertion, in walking, stooping, &c. it was this day, however, a good deal alarming, being much increased in quantity, and unaccompanied with pain.

I INTRODUCED my hand into the Vagina, and from a careful examination of the womb, was fatisfied that the Placenta was not at the mouth of it: in making this enquiry, I was able to break the membranes, a confiderable deal of water escaped, and the Hæmorrhage instantly ceased; and, though sufficient pains did not immediately come on, she was, about feven hours after, fafely delivered, by their fole affistance, of a large living child.

C A S E XXVIII.

SEPTEMBER 18, 1775, I was called about midnight to the wife of --- Baxter, another poor woman, under the care of a midwife; I was informed, that in the course of the preceding three weeks weeks she had frequently been alarmed with a discharge of blood from the womb, and that for about two hours before my arrival, she had been slooding most profusely: though she was very near her full time, there were not any symptoms of labor, nor had she yet had the most trisling pain: she was extremely faint, her pulse was hardly perceptible, and she appeared so much reduced as to be in the most imminent danger.

delivered by the netural pains, and the

Vagina, and found it full of coagulated blood, and with my finger carefully examined the mouth of the Uterus, which, though very little open, was, from the long continuance of the Hæmorrhage, loofe and dilatable: being convinced that the Placenta was not in the way, I endeavoured to break the membranes, but I could not do it fo foon as in the last case; after several repeated attempts, however, I at length succeeded, and a very large quantity of I water

water poured forth, by which means the Hæmorrhage was immediately fuppressed.

I CONTINUED a little stimulus to the Os Uteri, and it evidently brought on pain, which gradually increasing, the part became more open, and, in two hours from my first seeing the patient, without there being any return of bleeding, not even fo much as to occasion the least stain in examining, she was fafely delivered by the natural pains, and the child, notwithstanding the great loss which had been fustained, was born alive and vigorous. The Placenta came away without trouble, and no material discharge accompanied it; she remained very faint for some time after delivery, and was very feeble for feveral days; but at the time of writing this, which is a fortnight fince she was brought to bed, fhe has no complaint but weakness.

REMARK.

REMARK.

As the flooding, in this case, was so very profuse, as the patient had not the least pain, and was likewise extremely reduced, it is not unreasonable to suppose, that I should have been induced to have turned the child upon my first feeing her, if I had not before experienced fuch extraordinary proofs of nature's ability to expell the child, when the Placenta was not in the way to prevent the opening of the womb, and when ever fo little pain could be excited by stimulating the Os Tincæ: at the same time it appears very likely, that debilitated as she was, she would have been unable to support any other than the gradual and gentle dilatation of the womb, which nature effected, and that therefore, fuch a method of treatment would, probably, in this case, have been unfuccessful.

CONCLUSION.

OF the foregoing cases of slooding, nine were produced by a separation of the Placenta, occasioned by its being situated on the Os Uteri, and which was therefore, in every one of them, unavoidable; and nineteen were owing to a separation of it, arising from some accidental cause.

Or the latter number it appears, that though many were very alarming cases, as the patients lost large quantities of blood, and were extremely faint, not one proved fatal, not one but terminated safely, by waiting for the efforts of nature to expell the contents of the womb *; whilst, in all the former num-

* In two or three of the cases included in the latter number, it happened, indeed, that the children came into the world footling, and consequently, that more manual assistance was used than in natural presentations; but as this circumstance was totally accidental, and independent

ber, that no means whatever which nature could use, were able to suppress the difcharge, and that notwithstanding the complaint began in most of them in a manner but little alarming, yet nothing but the removal of the fœtus by art could fave the patients lives: in four out of the number, its being timely done, it had manifestly that happy effect; and in the others, where the turning was unfuccessfully used, it seemed to be clearly owing to its having been too long delayed; for, in the cases of King and Bond, where it was most evidently fo, the flooding began by no means in a threatening manner, nor did either of those women appear at first in so much danger as most of the nineteen other patients did, in whom the Hæmorrhage

of the flooding, as even in them, too, the dilatation of the womb was effected folely by nature; and as it is likewise very probable, that, if no affistance had been given (the children being small) that nature would at length have expelled them, I have considered the safe termination of the labors as effected by nature.

was produced by the accidental separation of the Placenta.

FROM hence, then, it is evident, how very frequently the Placenta is fixed to the Os Uteri, and that notwithstanding fo little notice has been taken of it, by those who have written on the subject of Uterine Hæmorrhages, how necessary it is, in every case, to make an enquiry for it: but the inferences which arise from the cases are so obvious, and the method of practice which they point out, has been before fo fully confidered, that it need not now be repeated; especially as the happy events of all the cases which were treated agreeable to the method recommended in the effay, of themselves, so fully speak its superiority over that which governed the mat nagement of the first related cases.

BEFORE I entirely quit the subject, it may not, perhaps, be totally foreign to it, to confider what would probably have been my method of treating the cases which

which succeeded those of King and Bond, if I had not then established some criterion, by which I could judge determinately of the propriety of trusting to nature, or of applying to art.

Ir is very natural to believe, that when my mind had been a good deal affected by the disagreeable events of those two cases, that I should not have hefitated to have had immediate recourse to delivery by art, in every following case in which I found the discharge at all confiderable; from a conclusion, (which, under fuch circumstances, would not have been an unreasonable one,) that as there feemed to be fome latent undifcoverable cause, which sometimes unexpectedly produced the most fatal mifchief, when at the beginning of the complaint there was no appearance of danger, it was justifiable to run the risque of unnecessarily turning the child, in fome cases, rather than be liable to omit doing it, in a fingle instance, where it might be abfolutely necessary for the woman's woman's fafety: and, indeed, could it be admitted that the indifcriminate use of this operation was attended with no danger, this would certainly be the only secure method of treating sloodings, were it not in our power to discover those particular cases in which the Placenta was situated on the mouth of the womb.

For these reasons, therefore, I doubt not but this would have been the method of practice which I should have adopted; and it is upon the same principles, I am persuaded, that the invariable use of turning has been recommended by some authors, and put in practice by some surgeons.

THE objections to fuch a method of treatment, are, however, felf-evident, as it must be particularly inconvenient and irksome to the surgeon, always painful, and sometimes dangerous to the patient.

This confideration, then, still further illustrates the advantage of knowing the true causes from whence these Hæmorrhages proceed; and if, therefore, by what has been said, I have in the least degree added to the knowledge of them, and shall be, on that account, but in a single instance, the means of saving the life of a sellow-creature, the little trouble I have had in throwing my thoughts together upon the subject, will not be lost labor, nor these pages, sew as they are, be written in vain.

THE END.

ERRATA.

DONCLUSION.

Page 12, line 9, for rom, read from.

Page 28, line 21, for of, read for.

Page 29, line 21, for out, read our.

Page 30, line 5, in note, for flooding, read floodings.

Page 39, line 5, in note, for and, read or.

Page 82, line 8, after delivered put a comma instead of the femicolon.

